



W
D916d
1860



NATIONAL LIBRARY OF MEDICINE

Bethesda, Maryland

From QUEEN CITY
BOOK CO.
7 W. Seventh St.
Cincinnati, - Ohio



DUNGLISON'S
MEDICAL DICTIONARY

DUNGLISON'S
MEDICAL DICTIONARY

WORKS BY THE SAME AUTHOR.

I. (JUST ISSUED.)

HUMAN PHYSIOLOGY:

With Five Hundred and Thirty-two Handsome Illustrations on Wood.

EIGHTH EDITION, REVISED, MODIFIED, AND ENLARGED.

In two large and handsome octavo volumes, of about 1500 pages; leather, price, \$7.

In preparing the present edition, "no pains have been spared to make the work a complete expression of the science of the day." This statement our own examination of the work enables us to confirm; every page of it testifying to the author's industry in culling from various quarters and sources all that was valuable in the physiological contributions to science of the last few years. The careful and scrutinizing spirit exhibited by the writer when investigating mooted questions, the extensive information he possesses

of general science in almost every department, and the clear and happy style in which he presents his views, render his Physiology one of the most reliable and attractive works in our language. To the practitioner and general reader, we can heartily recommend it as an excellent résumé of the present state of physiological science. As a text-book for the student, we think it has no superior in our language, and for this object we presume it was chiefly, if not expressly written.—*Med. Examiner, Oct., 1856.*

II. (NOW READY, JULY, 1857.)

GENERAL THERAPEUTICS AND MATERIA MEDICA:

ADAPTED FOR A MEDICAL TEXT-BOOK.

WITH ABOUT TWO HUNDRED ILLUSTRATIONS.

SIXTH EDITION, REVISED AND IMPROVED.

In two very handsome octavo volumes, of about 1100 pages; leather, price \$6.

The most complete and satisfactory exponent of the existing state of Therapeutical Science, within the moderate limits of a text-book, of any hitherto published. What gives the work a superior value, in our judgment, is the happy blending of Therapeutics and Materia Medica, as they are or ought to be taught in all our Medical schools; going no further into the nature and commercial history of drugs than is indispensable

for the medical student. This gives to the treatise a clinical and practical character, calculated to benefit, in the highest degree, both students and practitioners. We shall adopt it as a text-book for our classes, while pursuing this branch of medicine, and shall be happy to learn that it has been adopted as such in all of our medical institutions.—*The N. Y. Journal of Medicine.*

III. (JUST ISSUED.)

NEW REMEDIES:

WITH FORMULÆ FOR THEIR PREPARATION AND ADMINISTRATION.

Seventh Edition, with extensive Additions.

In one very large octavo volume, of 770 pages; leather, \$3.75.

It may be considered almost a work of supererogation to enter into an elaborate criticism of a work which has reached its *seventh edition*. The public has pronounced, in the most authoritative manner, its verdict, and we are certainly not disposed in the present instance to dispute its decision. In truth, such books as this will always be favorably received by the Profession of our country. They are labor-saving productions, which, at the expense of much research and reading to the author, condense in a convenient

space the novelties and discoveries of the age. The present edition of this work is considerably enlarged and improved. The author, with his accustomed accuracy, has elaborated and amplified many of the articles but casually or imperfectly treated of in the former editions; and he has also added considerably to the list of new remedies. About thirty new agents, or novel applications of old remedies, are introduced to the notice of the reader in this edition.—*Va. Med. and Surg. Jour., Sept., 1856.*

IV.

THE PRACTICE OF MEDICINE.

A TREATISE ON SPECIAL PATHOLOGY AND THERAPEUTICS.

THIRD AND REVISED EDITION.

In two large octavo volumes, of about 1500 pages; leather, \$6.25.

The student of medicine will find in these two elegant volumes a mine of facts, a gathering of precepts and advice from the world of experience,

that will nerve him with courage, and faithfully direct him in his efforts to relieve the physical sufferings of the race.—*Boston Med. and Surg. Jour.*

BLANCHARD AND LEA, PHILADELPHIA.

Leup.
MEDICAL LEXICON.

A
DICTIONARY
OF
MEDICAL SCIENCE;

CONTAINING

A CONCISE EXPLANATION OF THE VARIOUS SUBJECTS AND TERMS OF ANATOMY, PHYSIOLOGY,
PATHOLOGY, HYGIENE, THERAPEUTICS, PHARMACOLOGY, PHARMACY, SURGERY, OB-
STETRICS, MEDICAL JURISPRUDENCE, DENTISTRY, Etc.; NOTICES OF CLI-
MATE AND OF MINERAL WATERS; FORMULÆ FOR OFFICIAL,
EMPIRICAL, AND DIETETIC PREPARATIONS, Etc.;

WITH


FRENCH AND OTHER SYNONYMES.

BY

ROBLEY DUNGLISON, M.D., LL.D.,

PROFESSOR OF THE INSTITUTES OF MEDICINE, ETC.,
IN THE JEFFERSON MEDICAL COLLEGE
OF PHILADELPHIA.

REVISED AND VERY GREATLY ENLARGED.



PHILADELPHIA:
BLANCHARD AND LEA.

1860.

W
D 9162
1860

Entered, according to Act of Congress, in the year 1857, by

BLANCHARD AND LEA

in the Clerk's Office of the District Court of the United States for the Eastern District of Pennsylvania.

COLLINS, PRINTER.

P R E F A C E .

ON this new edition of his "*Medical Lexicon*," the author has bestowed more than usual labour. It is not more than four years since a friendly reviewer, in one of the most learned of European medical periodicals, (the "*British and Foreign Medico-Chirurgical Review*" for July, 1853, p. 205,) in noticing a former edition, remarked, that the labour which had been bestowed upon the work had "been something prodigious;" adding—"the work, however, has been now done, and we are happy in the thought, that no human being will have again to undertake the same gigantic task. Revised and corrected from time to time, Dr. Dunglison's '*Medical Lexicon*' will last for centuries."

Yet, in the present edition, not only has the work been "revised and corrected," but about six thousand subjects and terms have been added, which are not to be found in the one noticed by the "*British and Foreign*." Many of these have been introduced into medical terminology in consequence of the progress of the science; whilst others had escaped the author in the previous editions. These additions have necessarily required a great amount of labour, which—as on every former occasion of the kind—has been cheerfully bestowed, in order that the work might be rendered still more worthy of the vast favour which it has experienced. It has ever been the anxious desire of the author to make it a satisfactory and desirable—if not indispensable—lexicon, in which the student may search without disappointment for every term that has been legitimated in the nomenclature of the science; and the present very carefully revised, greatly enlarged, and accurately printed edition cannot fail to be more extensively useful, and to offer stronger claims to the attention of the practitioner and student, than any of its predecessors.

The author is grateful for the opportunity again afforded him of expressing his acknowledgments for the reception which the Dictionary has met with everywhere from the profession.

ROBLEY DUNGLISON

PHILADELPHIA, 1116 GIRARD ST.

JULY, 1857.

PREFACE

TO THE SECOND EDITION.

THE present undertaking was suggested by the frequent complaints, made by the author's pupils, that they were unable to meet with information on numerous topics of professional inquiry,—especially of recent introduction,—in the medical dictionaries accessible to them.

It may, indeed, be correctly affirmed, that we have no dictionary of medical subjects and terms which can be looked upon as adapted to the state of the science. In proof of this, the author need but remark, that he has found occasion to add several thousand medical terms, which are not to be met with in the only medical lexicon at this time in circulation in the country.

The present edition will be found to contain many hundred more terms than the first, and to have experienced numerous additions and modifications.

The author's object has not been to make the work a mere lexicon or dictionary of terms, but to afford, under each, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of medical science. In its preparation, he has freely availed himself of the English, French, and German works of the same nature, and has endeavored to add every subject and term of recent introduction, which has fallen under his notice; yet, with all his care, it will doubtless be found that subjects have been omitted. The numerous additions, however, which he has made, and his strong desire to be useful, "by removing rubbish and clearing obstructions from the paths through which learning and genius press forward to conquest and glory," will, he trusts, extenuate these and other objections that might be urged against the work; especially when the toil, which every compiler of a dictionary must endure, is taken into consideration; a toil which has been so forcibly depicted by the great English Lexicographer, as well as by the distinguished SCALIGER:

"Si quelqu'un a commis quelque crime odieux,
S'il a tué son père, ou blasphémé les Dieux,
Qu'il fasse un Lexicon: s'il est supplice au monde
Qui le punisse mieux, je veux que l'on me tonde."

EXPLANATION.

IF the simple synonymy of any term be needed, a mere reference to the term may be sufficient; but if farther information be desired, it may be obtained under the term referred to. For example, the French word *Tronc* is said to be synonymous with *Trunk*. This may be sufficient for the inquirer: should it not, the requisite information may be found by turning to *Trunk*.

ABBREVIATIONS ARBITRARILY EMPLOYED.

Arab.	Arabic.	P.	Portuguese.
Ch.	Chaussier.	Ph. D.	Pharmacopœia of Dublin.
D.	Dutch.	Ph. E.	“ Edinburgh.
Da.	Danish.	Ph. L.	“ London.
E.	English.	Ph. P.	“ Paris.
F.	French.	Ph. U. S.	“ of the United States of America.
F. or Fah.	Fahrenheit.		
Fam.	Family.	Pl.	Plural.
G.	German.	Prov.	Provincial.
Heb.	Hebrew.	R.	Réaumur.
I.	Italian.	S.	Spanish.
Imp. meas.	Imperial measure.	S. g.	Specific Gravity.
Ir.	Irish.	Sax.	Anglo-Saxon.
L.	Latin.	Sc.	Scottish.
Linn.	Linnæus.	Sex. Syst.	Sexual System.
Nat. Ord.	Natural Order.	Sw.	Swedish.
Old Eng.	Old English.	Vulg.	Vulgarly.
Ord.	Order.		



A

DICTIONARY

OF

MEDICAL SCIENCE.

A.

A

A, before a consonant; *An* before a vowel, *a*, *av*, have, in the compound medical terms, a privative or debasing signification, like that of the particles *in*, *im*, *un*, *ir*, in English. Thus: *Stheni'a* means strength; *Astheni'a*, want of strength; *Anæmia*, want of blood, etc. Occasionally, in compound words, they have an intensive meaning.

AACHEN, Aix-la-Chapelle.

A, or **AA**. See Abbreviation.

AANDE, Breath.

AARZIL, MINERAL WATERS OF. **A** is in the canton of Berne, in Switzerland. The chief spring contains chlorides of calcium and sodium, sulphates of lime and soda, oxyd of iron, and sulphohydric acid gas.

AASMUS, Anhelatio.

ABACH, MINERAL WATERS OF. A hydro-sulphuretted saline spring not far from Ratisbon or Regensburg, in Bavaria.

ABAISSÉMENT, Depression; *a. de la Cataracte*, see Cataract—*a. de la Matrice*, Prolapsus uteri.

ABAISSÉUR DE L'AILE DU NEZ, Depressor alæ nasi—*a. de l'angle des lèvres*, Depressor anguli oris—*a. de la lèvre inférieure*, Depressor labii inferioris—*a. de la langue*, Glossoctochus—*a. de la mâchoire inférieure*, Digastricus—*a. de l'œil*, Rectus inferior oculi.

ABALIENATIO MENTIS, Insanity.

ABALIENATUS, *Corruptus*, Corrupted; from *ab*, and *alienus*, 'different.' *Membra abalienata*. Limbs dead or benumbed.—Celsus, Scribonius Largus.

ABANGA. Name given by the inhabitants of St. Thomas to the fruit of a palm tree, the seeds of which they consider very useful in diseases of the chest, in the dose of three or four, two or three times a day.

ABAPTIST'A, *Abaptis'ton* or *Abaptis'tum*, from *a*, privative, and *βαπτίζω*, 'to plunge.' A term applied to the old trepan, the conical shape of which prevented it from plunging suddenly into the cavity of the cranium.

ABAPTISTON, Abaptista.

ABAPTISTUM, Abaptista.

ABAREMO-TEMO. A Brazilian tree, which grows in the mountains, and appears to be a mimosa. Piso relates that the decoction of its bark, which is bitter and astringent, was applied in that country to ulcers of a bad character.

ABARNAHAS, Magnesia.

ABARTICULATIO, Diarthrosis, and Synarthrosis.

ABÁTARDISSEMENT, Degeneration.

ABATTEMENT, Prostration.

ABBREVIATION

ABATTIS, Giblets.

ABBÉCOURT, MINERAL WATERS OF. A chalybeate spring, six leagues from Paris, and one from Poissy. It was once much frequented, but is now abandoned.

ABBEVILLE, MINERAL WATERS OF. An acidulous chalybeate at Abbeville, in the department of Somme, France.

ABBREVIATION, *Abbreviatio*, *Brachyn'sis*, *Brachys'mos*, *Abbreviatu'ra*. (F.) *Abréviation*, from *brevis*, 'short.' Abbreviations are chiefly used in medicinal formulæ. They are by no means as frequently employed now as of old, when every article had its appropriate symbol. The following are some of the abbreviations which have been or are employed:

R. *Recipe*, Take.

A, **AA**, **ANA**, (*ava*) *utrinque*, of each.

ABDOM. *Abdomen*.

ABS. **FEBR**. *Absente febre*, In the absence of fever.

AD. or **ADD**. *Addé* or *addotur*.

AD. **LIB**. *Ad libitum*, At pleasure.

ADMOV. *Admovere*, Let it be applied.

ALTERN. **HOR**. *Alternis horis*. Every other hour.

ALV. **ADSTRACT**. *Alvo adstrictâ*, The bowels being confined.

AQ. *Aqua*, Water.

AQ. **BULL**. *Aqua bulliens*, Boiling water.

AQ. **COMM**. *Aqua communis*, Common water.

AQ. **FERV**. *Aqua fervens*, Hot water.

AQ. **FONT**. *Aqua fontis*, Spring water.

AQ. **MARIN**. *Aqua marina*, Sea water.

B. **A**. *Balneum arenæ*, A sand-bath.

BALS. *Balsamum*, Balsam.

BB. **BDSD**. *Barbadensis*, Barbadoes.

BIB. *Bibe*, Drink.

BIS IND. *Bis indies*, Twice daily.

B. **M**. *Balneum mariæ*, (F.) *Bain marie*, A water bath.

BOL. *Bolus*.

BULL. *Bulliat*, Let it boil.

BUT. *Butyrum*, Butter.

B. **V**. *Balneum vaporis*, A vapour bath.

C. *Congius*, A gallon.

CÆRUL. *Cæruleus*, Blue.

CAP. *Capiat*, Let him take.

C. **C**. *Cornu cervi*, Hartshorn.

C. **C**. **U**. *Cornu cervi ustum*, Burnt hartshorn

C. **M**. *Cras manâ*, To-morrow morning.

C. **N**. *Cras nocte*, To-morrow night.

C. **V**. *Cras vespere*, To-morrow evening.

COCHL. *Cochleæ*, A spoonful.

COCHL. **AMPL**. *Cochleare amplum*, A large spoonful.

COCHL. INF. *Cochleare infantum*, A child's spoonful.
 COCHL. MOD. or MED. *Cochleare modicum* or *medium*, A dessert-spoonful.
 COCHL. PARV. *Cochleare parvum*, A tea-spoonful.
 COL., *Cola*, and *Colatura*, Strain, and to the strained.
 COMP. *Compositus*, Compound.
 CONF. *Confectio*, Confection.
 CONS. *Conserva*, Conserve.
 CONT. *Continetur*, Let it be continued.
 COQ. *Coque*, Boil.
 CORT. *Cortex*, Bark.
 CRAST. *Crastinus*, For to-morrow.
 CUI. *Cujus*, Of which.
 CUJUSL. *Cujuslibet*, Of any.
 CYATH. *Cyathus*, A glassful.
 CYATH. THEÆ, A cup of tea.
 D. *Dosis*, A dose.
 D. et S. *Detur et signetur* (placed at the end of a prescription).
 D. D. *Detur ad*, Let it be given in or to.
 DEAUR. PIL. *Deauretur pilula*, Let the pill be gilded.
 DEB. SPISS. *Debita spissitudo*, A due consistence.
 DEC. *Decanta*, Pour off.
 DECUB. *Decubitus*, Lying down, going to bed.
 DE D. IN D. *De die in diem*, From day to day.
 DEJ. ALV. *Dejectiones alvi*, Alvine evacuations.
 DEP. *Depuratus*, Purified.
 DEST. *Destilla*, Distil.
 DET. *Detur*, Let it be given.
 DIEB. ALTERN. *Diebus alternis*, Every other day.
 DIEB. TERT. *Diebus tertiis*, Every third day.
 DIG. *Digeratur*. Let it be digested.
 DIL. *Dilutus*, Dilute.
 DIM. *Dimidius*, One-half.
 DIV. *Divide*, Divide.
 DONEC ALV. SOLUT. FUER. *Donec alvus soluta fuerit*, Until the bowels are opened.
 DRACH. *Drachma*, A drachm.
 EJUSD. *Ejusdem*, Of the same.
 ENEM. *Enema*, A clyster.
 EXHIB. *Exhibeatur*, Let it be exhibited.
 EXT. SUPER ALUT. *Extende super alutum*, Spread upon leather.
 F. *Fiat*, Let it be made.
 F. PIL. *Fiat pilula*, Make into a pill.
 F. VENÆ. or F. VS. *Fiat venæsectio*, Let bleeding be performed.
 FEB. DUR. *Febre durante*, The fever continuing.
 FEM. INTERN. *Femoribus internis*, To the inside of the thighs.
 FIST. ARMAT. *Fistula armata*, A bag and pipe—a clyster pipe and bladder fitted for use.
 FL. *Fluidus*, and *Flores*, Fluid, and Flowers.
 FRUST. *Frustillatim*, In small pieces.
 GEL. QUAVIS. *Gelatinâ quavis*, In any kind of jelly.
 G. G. G. *Gummi guttæ Gambiæ*, Gamboge.
 GR. *Granum*, A grain.
 GT. *Gutta*, A drop. GTT. *Guttæ*, Drops.
 GTT. or GUTT. QUIBUSD. *Guttis quibusdam*, With some drops.
 GUM. *Gummi*, Gum.
 GUTTAT. *Guttatim*, By drops.
 HOR. DECUB. *Horâ decubiti*, At bed-time.
 HOR. INTERM. *Horâ intermedii*, At intermediate hours.
 H. S. *Horâ somni*, At bed-time.
 INF. *Infunde*, Infuse.
 IND. *Indies*, Daily.
 INJ. ENEM. *Iniciatur enema*, Let a clyster be given.
 IN PULM. *In pulmento*, In gruel.
 JUL. *Julepus*, A julep.
 LAT. DOL. *Lateri dolenti*, To the pained side.
 LB. and LIB. *Libra*, A pound weight.

LIB. LIB. *Libræ*, Pounds.
 LIQ. *Liquor*.
 M. *Misce*, Mix.
 MAC. *Macera*, Macerate.
 MAN. *Manipulus*, A handful.
 MAN. PRIM. & ante primo, Early in the morning.
 MIC. PAN. *Mica panis*, Crumb of bread.
 MIN. *Minimum*, The 60th part of a drachm by measure.
 MITT. *Mitte*, Send.
 MITT. SANG. *Mittatur sanguis*, Let blood be drawn.
 MOD. PRÆSCRIPT. *Modo præscripto*, In the manner directed.
 MOR. SOL. *More solito*, In the usual manner.
 MUC. *Mucilago*, Mucilage.
 N. M. *Nux moschata*, Nutmeg.
 O. *Octarius*, A pint.
 OL. *Oleum*, Oil.
 OL. LINI. S. I. *Oleum lini sine igne*, Cold-drawn linseed oil.
 OMN. BID. *Omni biduo*, Every two days.
 OMN. BIH. *Omni bihorio*, Every two hours.
 OMN. HOR. *Omni horâ*, Every hour.
 OMN. MAN. *Omni manâ*, Every morning.
 OMN. NOCTE, Every night.
 OMN. QUADR. HOR. *Omni quadrante horâ*, Every quarter of an hour.
 O. O. O. *Oleum olivæ optimum*, Best olive oil.
 OV. *Ovum*, An egg.
 OX. *Oxymel*.
 OZ. *Oncia*, An ounce.
 P. *Pondere*, By weight, and *Pars*, A part.
 P. and PUG. *Pugillus*, A pugil.
 P. Æ. *Partes æquales*, Equal parts.
 PART. VIC. *Partitis vicibus*, In divided doses.
 PERACT. OP. EMET. *Peractâ operatione emeticâ*, The operation of the emetic being over.
 PIL. *Pilula*, A pill. PIL. or PILL. *Pilulæ*, Pills.
 POST. SING. SED. LIQ. *Post singulas sedes liquidas*, After every liquid evacuation.
 POT. *Potio*, A potion.
 P. P. *Pulvis patrum*, Jesuits' bark.
 P. RAT. ÆTAT. *Pro ratione ætatis*, According to the age.
 P. R. N. *Pro re natâ*, As occasion may be.
 PULV. *Pulvis*, A powder.
 Q. P. *Quantum placeat*, As much as may please.
 Q. S. *Quantum sufficit*, As much as is sufficient.
 QUOR. *Quorum*, Of which.
 Q. V. *Quantum volueris*, As much as you wish.
 RAD. *Radix*, Root.
 RAS. *Rasura*, Shavings.
 RECT. *Rectificatus*, Rectified.
 RED. or REDIG. IN PULV. *Redactus in pulverem*, or *Redigatur in pulverem*, Powdered, or Let it be powdered.
 REG. UMBIL. *Regio umbilici*, The umbilical region.
 REPET. *Repetatur*, Let it be repeated.
 S. A. *Secundum artem*, According to art.
 SEM. *Semen*, Seed.
 SEMI-DR. *Semi-drachma*, half a drachm.
 SEMI-H. *Semi-hora*, Half an hour.
 SERV. *Serva*, Keep, preserve.
 SESQUIH. *Sesquihora*, An hour and a half.
 SESUNC. *Sesuncia*, An ounce and a half.
 SI NON VAL. *Si non valet*, If it does not answer.
 SI OP. SIT. *Si opus sit*, If there be need.
 SI VIR. PERM. *Si vires permittant*, If the strength will permit.
 SING. *Singulorum*, Of each.
 SOLV. *Solve*, Dissolve.
 SP. and SPIR. *Spiritus*, Spirit.
 SS. *Semi*, One half.
 ST. *Stet*, Let it stand.
 SUB FIN. COCT. *Sub finem coctionis*, Towards the end of the boiling.

SUM. *Sumat*, Let him take; also, *Summitates*, The tops.
S. V. *Spiritus vini*, Spirit of wine.
S. V. R. *Spiritus vini rectificatus*, Rectified spirit of wine.
S. V. T. *Spiritus vini tenuior*, Proof spirit of wine.
SYR. *Syrupus*, Syrup.
TEMP. DEXT. *Tempori dextro*, To the right temple.

T. O. *Tinctura opii*, Tincture of opium.
TR., TRA. and TINCT. *Tinctura*, Tincture.
TRIT. *Tritura*, Triturate.
V. O. S. or VIR. OV. SOL. *Vitello ovi solutus*, Dissolved in the yolk of an egg.
VS. *Venæsectio*, Venesection.
Z. Z. *Anciently myrrh*: now *zinziber* or ginger.
lb. *Libra*, A pound. **lb.** *Pounds*.
℥. *Uncia*, An ounce.
℥℥. *Fluiduncia*, A fluidounce.
℥. *Drachma*, A drachm.
℥℥. *Fluidrachma*, A fluidrachm.
℥. *Scrupulum*, A scruple.
℥. *Minimum*, A minim.
ss. *Semis*, or half; **iss.**, one and a half.
j., one; **ij.**, two; **ijj.**, three; **iv.**, four, &c. See Symbol.

The same system is not always followed in abbreviating. The subjoined will exhibit the usual mode:

R
Infus. Colomb. f℥ iss
Tinct. Gent. comp. f℥ i
Syr. Cort. Aurant. f℥ ss
Tinct. caps. gtt. xl. M.

Capt. coch. ij. p. r. n.
 This, written at length, is as follows:

Recipe
Infusi Colombæ sesqui-fluidunciam.
Tincturæ Gentianæ Compositæ fluidrachmam.
Syrupi Corticis Aurantiorum semi-fluidrachmam.
Tincturæ Capsici guttas quadraginta.
 Misce.

Capiat cochlearia duo pro re natâ.
 In the United States the directions are always written in English.

ABCS. Abscess—a. *Aigu*, see Abscess.

ABCS EN BOUTON DE CHEMISE. *Shirt-button* or *Shirt-stud Abscess*. A deep-seated mammary abscess, following the septa of the breast, and appearing under the skin, so as eventually to give rise to one or more subcutaneous abscesses, without losing the original character of submammary suppuration. A cavity exists between the integuments and the gland—another, larger, between the mamma and the chest—the two communicating by a passage, which is generally narrow—the whole having the exact appearance of a shirt-stud.—*Velpeau*.

ABCS CHAUD. see Abscess—a. *Chronique*, see Abscess—a. *par Congestion*, see Abscess—a. *Consécutif*, Abscess, metastatic—a. *Diathésique*, see Abscess—a. *Froid*, see Abscess—a. *Métastatique*, Abscess, metastatic—a. *Rétro-utérine*, see Retro-uterine—a. *Scrofuleux*, see Abscess—a. *Soudain*, see Abscess.

ABDO'MEN, from *abdere*, 'to conceal,' because it conceals the viscera; by some, fancifully presumed to be from *abdere*, 'to conceal,' and *omentum*, 'the caul.' From *abdo* is formed *abdomen*, as from *lego*, *legumen*. *Etron*, *Hypogastrium*, *Physce*, *Hypocæ'lium*, *Epis'chion*, *Lap'ara*, *Hypocho'lion*, *Gaster*, *Hypog'trion*, *Nedys*, *Neig'ra*, *Nei'ra*, *Abdu'men*, *Venter*, *Venter imus*, *Venter infimus*, *Alvus*, *Uterus*, *The belly*, (Sc.) *Kyte*, *Penche*, *Weam*, (Prov.) *Baggie*, *Hag*, *Wem*, *Pook*, (F.) *Ventre*, *V. inférieur*, *Bas ventre*. The largest of the three splanchnic cavities, bounded

above, by the diaphragm; below, by the pelvis; behind, by the lumbar vertebræ; and at the sides and fore part, by muscular expansions. It is distinguished into three anterior regions, from above to below; viz. the epigastric, umbilical, and hypogastric, each of which is itself divided into three others, one middle, and two lateral: thus, the epigastric region comprises the *epigastrium* and *hypochondria*; the umbilical, the *umbilicus* and *flanks* or *lumbar regions*; and the hypogastric, the *hypogastrium* and *iliac regions*. None of these regions has its limits well defined. The chief viscera contained in the cavity of the abdomen, *Cœ'lia*, *Cavum Abdom'inis*, are the stomach, intestines, liver, spleen, pancreas, kidneys, &c. It is lined by the peritoneum.

ABDOMEN, PENDULOUS, Physconia.

ABDOM'INAL, *Abdominalis*, *Ventra'lis*, *Ventral*. That which belongs to the Abdomen, as *abdominal muscles*, *abdominal viscera*, &c.

ABDOMINIS EXPLORATIO, *Abdominoscopia*.

ABDOMINOSCOPIA, *Gastrosco'pia*, *Laparosco'pia*, *Abdom'inis Exploratio*. *Abdominoscopy*. A hybrid word, from *abdomen*, the lower belly, and *σκοπεω*, 'I view.' Examination of the lower belly as a means of diagnosis. See Auscultation.

ABDOM'INOUS, *Ventrio'sus*, *Ventr'us*, from *abdomen*, 'the belly,' *Big-bellied*, *Big-paunched*, *Ventrip'otent*. Having a large abdomen.

ABDUCENS LABIORUM, *Levator anguli oris*.

ABDUCENTES, *Motor oculi externus*.

ABDUCTEUR DE L'ŒIL, *Rectus externus oculi*—*a. de l'oreille*, *Abductor auris*—*a. du gros orteil*, *Abductor pollicis pedis*—*a. du petit orteil*, *Abductor minimi digiti pedis*—*a. court du ponce*, *Abductor pollicis brevis*—*a. long du ponce*, *Abductor longus pollicis*.

ABDUCTION, *Abduc'tio*, from *abducere*, to separate, (*ab* and *ducere*, 'to lead.') The movement which separates a limb or other part from the axis of the body.

The word has also been used synonymously with *Abruptio*, *Apag'ma*, *Apoclas'ma*, a fracture near the articular extremity of a bone, with separation of the fragments.

ABDUCTOR, same etymon. (F.) *Abducteur*. A muscle which moves certain parts by separating them from the axis of the body.

ABDUCTOR ARTICULARIS, *Abductor auris*—*a. Indicis pedis*, *Prior indicis pedis*, *Posterior indicis pedis*—*a. Medii digiti pedis*, *Prior medii digiti pedis*—*a. Minimi digiti*, *Flexor parvus minimi digiti*—*a. Minimi digiti*, *Prior minimi digiti*—*a. Oculi*, *Rectus externus oculi*—*a. Pollicis manus*, and *a. Brevis alter*, *Abductor pollicis brevis*.

ABDUCTOR AURIS, *Abductor auriculæ'ris*. (F.) *Abducteur de l'oreille*. A portion of the *posterior auris*, whose existence is not constant, which passes from the mastoid process to the concha.

ABDUCTOR IN'DICIS, *Semi-interos'seus in'dicis*. A muscle which arises from the os trapezium and metacarpal bone of the thumb, and is inserted into the first bone of the forefinger. Its use is to bring the forefinger towards the thumb.

ABDUCTOR MIN'IMI DIG'ITII, *Carpo-phalan'geus minimi digiti*, *Carpo-phalangien du petit doigt*, *Extensor tertii interno'dii minimi digiti*—(Douglas.) *Hypothenar minor metacarpeus*. See *Flexor parvus*. It originates, fleshy, from the os pisiforme, and from the annular ligament near it; and is inserted, tendinous, into the inner side of the base of the first bone of the little finger. Use, to draw the little finger from the rest.

ABDUCTOR MINIMI DIGITI PEMS, *Calco-subphalangeus minimi digiti*, *Calcaneo-phalangien du petit orteil*, *Parath'nar major*—(By Winslow, the muscle is divided into two portions,—

Parathenar major and metatarsus.) *Calcaneos-phalangien du petit orteil*—(Ch.) (F.) *Abducteur du petit orteil*. This muscle forms the outer margin of the sole of the foot, and is immediately beneath the plantar aponeurosis. It arises, tendinous and fleshy, from the outer side of the protuberance of the os calcis, and from the root of the metatarsal bone of the little toe, and is inserted into the outer part of the root of the first bone of the little toe. *Use*, to draw the little toe outwards.

ABDUCTOR POL'LICIS BREVIS, *Abductor Pollicis Manûs*, *Scapho-carpo-super-phalangeus Pollicis*, *Sus-phalangien du ponce*, *A. pollicis manus* and *A. brevis alter*—(Albinus.) (F.) *Abducteur court du ponce*, *Carpo-sus-phalangien du ponce*—(Ch.) A short, flat, triangular muscle, which arises from the anterior surface of the os scaphoides and the annular ligament of the carpus, and terminates at the outside of the upper extremity of the first phalanx of the thumb. A particular portion, on the inner side of this muscle, is called, by Albinus, *Abductor brevis alter*.

ABDUCTOR LONGUS POLLICIS, *A. l. P. Manûs*, *Extensor ossis metacarpi pollicis manûs*, *Extensor primi internodii*—(Douglas,) *Extensor primus Pollicis*, *Cubito-radi-sus-metacarpium du ponce*, *Cubito-sus-metacarpium du ponce*,—(Ch.) (F.) *Abducteur long du ponce*. A long, thin muscle, arising from the posterior surface of the ulna, radius, and interosseous ligament, and inserted at the outer side of the upper extremity of the first metacarpal bone.

ABDUCTOR POLLICIS PEDIS, *Calco-sub-phalangeus Pollicis*. (F.) *Abducteur du gros orteil*. This muscle arises, fleshy, from the anterior and inner part of the protuberance of the os calcis, and tendinous from the same bone where it joins with the os naviculare. It is inserted, tendinous, into the internal os sesamoideum and root of the first bone of the great toe. *Use*, to pull the great toe from the rest.

The name *Abductor* has been given also to all those interosseous muscles of the hand and foot, which perform the motion of abduction on the fingers or toes, and to muscles which execute the same function on other parts of the body.

ABDUMEN, Abdomen.

ABEBÆ'OS, from *a*, neg., and *βεβαιος*, 'firm,' *Infirmus*, *Deb'ilis*. Weak, infirm, unsteady.

ABELLE, Bee.

ABELMELUCH. One of the names of the Ricinus, according to some authors.—Prosper Alpinus says that a tree, which grows about Mecca, is so called. Its seeds, which are black and oblong, are said to be a most violent cathartic.

ABELMOSCHUS, *Hibiscus abelmoschus*—*a*. *Moschatus*, *Hibiscus abelmoschus*.

ABELMUSK, *Hibiscus abelmoschus*.

ABENSBERG, MINERAL WATERS OF. *A.* is a city of Bavaria, where there is a cold, sulphureous spring.

ABERRATIO, Aberration—*a*. *Lactis*, *Galactorrhea*—*a*. *Mensium*, *Menstruation*, vicarious—*a*. *Menstruorum*, *Menstruation*, vicarious.

ABERRA'TION, *Aberra'tio*, from *aberrare*, (*ab* and *errare*), 'to stray,' 'to wander from.' This word has several meanings.

1. The passage of a fluid of the living body into an order of vessels not destined for it. In this sense it is synonymous with the *Error Loci* of Boerhaave.

2. The flow of a fluid towards an organ different from that to which it is ordinarily directed; as in cases of vicarious hemorrhage. *Aberrations of sense or judgment* are certain errors in the perceptions, or certain derangements of the intellectual faculties.

The word is used in optics to designate the

dispersion of the rays of light in passing through a lens.

ABERRATION, CHROMATIC, Aberration of Refrangibility.

ABERRATION OF REFRANGIBIL'ITY, *Chromat'ic aberration*, (F.) *Aberration de Réfrangibilité*, *Chromatisme*, exists, when, as in a common lens, the rays that pass near the circumference of the lens are decomposed, so that a coloured image is observed. This aberration in the human eye is corrected by the iris, which does not permit the rays to fall near the circumference of the lens, and also by the crystalline lens itself, which, owing to its structure, serves the purposes of an achromatic glass.

ABERRATION, SPHERICAL, Aberration of sphericity.

ABERRATION OF SPHERIC'ITY or spher'ical aberration takes place, when the rays, as in a common lens, which pass through the centre of the lens, and those which pass near the circumference, are unequally refracted, so that they do not meet at a common focus.

This aberration of sphericity in the human eye is corrected by the iris and lens.

ABESSI, Realgar.

ABEVACUA'TIO, *Apoceno'sis*, from *ab*, and *evacuare*, 'to empty.' An evacuation. A partial or imperfect evacuation. By some it is applied to an immoderate evacuation.—Kraus.

ABHAL. A fruit well known in India, and obtained from a species of cypress. It passes for an emmenagogue.

ABIES, *Pinus picea*—*a*. *Balsamea*, *Pinus balsamea*.

ABIES BALSAMIFERA, *Pinus Balsamea*—*a*. *Canadensis*, *Pinus Canadensis*—*a*. *Excelsa*, see *Pinus abies*—*a*. *Gallica*, *Pinus picea*—*a*. *Larix*, *Pinus larix*—*a*. *Pectinata*, *Pinus picea*—*a*. *Picea*, *Pinus picea*—*a*. *Rubra*, *Pinus rubra*.

ABIGA, *Teucrium Chamapitys*.

ABIOSIS, Death.

ABIOTOS, *Conium maculatum*.

ABIRRITA'TION, *Abirritatio*, from *ab*, privative, and *irritatio*, 'irritation.' This word strictly means absence or defect of irritation. The disciples of Broussais used it to indicate a pathological condition, opposite to that of irritation. It may be considered as synonymous with debility, asthenia, &c.

ABLACTATIO, Weaning.

ABLASTES, Sterile.

ABLATIO, Extirpation.

ABLEPH'ARUS, from *a*, privative, and *βλεφαρον*, 'eyelid.' One who has no eyelids.

ABLEPSIA, *Cæcitas*.

ABLUTENTIA, Detergents.

ABLUTENTS, Detergents.

ABLUTION, *Ablu'tio*, *Aponip'sis*, *Cataclysmus*, from *ablueri*, (*ab* and *luere*), 'to wash.' A name given to legal ceremonies in which the body is subjected to particular affusions. Ablution (especially of the extremities) with cold or tepid water is employed, therapeutically, to reduce febrile heat. Also, the washing by which medicines are separated from the extraneous matters mixed with them.

ABNORMAL, Abnormous.

ABNORMITY, Anomalia.

ABNORMOUS, *Abnor'mis*, *Enor'mis*, *Abnor'mal*, (F.) *Anormal*, from *ab*, 'from,' and *norma*, 'rule.' Not conformable to rule; irregular.

ABOLI'TION, *Aboli'tio*, destruction or suppression, from *ab* and *luere* (?) 'to wash.' A word, often employed, especially by the French, to express the complete suspension of any symptom or function. *Abolition of the sight*, e.g. is the complete loss of sight.

ABOMA'SUS, *Aboma'sum*, *Enys'tron*, *Ventriculus*.

ulus intestinalis, Rennet, (Sc.) Roddikin, (F.) Caillette. The lowermost or fourth stomach of ruminating animals.

ABOMINATIO, Disgust.

ABONDANCE, see *Plethora*.

ABORSIO, Abortion.

ABORSUS, Abortion.

ABORTICIDIUM, Fœticide.

ABORTIF, Abortive.

ABORTIFACIENS, Abortive.

ABORTION, *Abor'tus, Abor'sus, Abor'sio, Dys-to'cia abor'ti'va, Omoto'cia, Paracy'e'sis abortus, Amblo'sis, Amblo'ma, Amblos'mus, Ec'bole, Embryo'to'cia, Diaph'thora, Etro'r'sis, Examblo'ma, Examblo'sis, Ectros'mos, Apopal'e'sis, Apopal'sis, Apoph'thora, Phthora, Convul'sio u'teri, Depen'dit'io.* (F.) Abortement, *Blessure, Miscarriage*; from *ab* and *oriri*, 'to rise,' applied to that which has arisen out of season. The expulsion of the fœtus before the seventh month of utero-gestation, or before it is *viable*. The causes of this accident are referable either to the mother, and particularly to the uterus; or to the fœtus and its dependencies. The causes, in the mother, may be:—extreme nervous susceptibility, great debility, plethora, faulty conformation, &c.; and it is frequently induced immediately by intense mental emotion, violent exercise, &c. The causes seated in the fœtus are its death, rupture of the membranes, &c. It most frequently occurs between the 8th and 12th weeks of gestation. The symptoms of abortion are:—uterine hemorrhage with or without flakes of decidua, with intermitting pain. When abortion has once taken place, it is extremely apt to recur in subsequent pregnancies about the same period. Some writers have called abortion, when it occurs prior to three months, *Effluxion*. The treatment must vary according to the constitution of the patient and the causes giving rise to it. In all cases, the horizontal posture and perfect quietude are indispensable.

ABORTION is likewise applied to the product of an untimely birth,—*Abor'tus*, *Abor'sus*, *Apoble'ma*, *Apob'ole*, *Eeblo'ma*, *Amblothr'id'ion*, *Eetro'ma*, *Fruc'tus immatu'rus*, *Abortment*, (F.) *Avorton*, *Avortin*.

TO ABORT, *Abori'ri*. To miscarry. (F.)
Avorter.

ABORTIVE, *Abortivus*, *Ecbolicus*, *Ambloiticus*, *Amblothruidium*, *Ambolicus*, *Phthorivus*, *Apophthoricus*, *Ectroficus*, *Abortifaciens*, *Acyterius*, *Expel lens*, *Phthoriconotus*, *Phthoriconotus*, *Ecbolicus*, *Contractor uteri*, *Accelera tor Partus*, *Parturient*, *Parturificans*, *Ecbolic*. (F.) *Abortif.* A medicine to which is attributed the property of causing abortion. There is probably no direct agent of the kind.

ABORTMENT, Abortion.

ABORTUS. Abortion.

ABOUCHEMENT, Anastomosis.

ABOULAZA, a tree of Madagascar, used, according to Flacourt, in the practice of the country, in diseases of the heart.

ABOUTISSEMENT. Suppuration.

ABOYEUSES, see *Convulsionnaire*.

ABRABAX, *Abraax*, *Abraaxas*. A mystic term, expressing the number 365, to which the Cabalists attributed miraculous properties.

ABRACADA'BRA, *Abracada'bra*, the name of a Syrian idol, according to Selden. This word, when pronounced and repeated in a certain form and a certain number of times, was supposed to have the power of curing fevers and preventing many diseases. It was figured on amulets and worn suspended around the neck.

א
ב
ג
ד
ה
ו
ז
ח
ט
י

ABRACALAN, A cabalistic term to which the Jews attributed the same virtue as to the word ABRACADABRA.

ABRASADABRA, Abracadabra.

ABRASAX, Abrabax.

ABRA'SION, (Prov.) *Fleck, Abrasão, Apoxysma, Apoxysmus*, from *abrader*, (ab and radere) 'to rasp.' A superficial excoriation, with loss or substance, under the form of small *shreds*, in the mucous membranes of the intestines,—(F.) *Rachures des Boyaux*. Also an ulceration of the skin, possessing similar characters. According to Vieq d'Azyr, the word has been used for the absorption of the molecules composing the various organs.

ABRATHAN, *Artemisia abrotanum*.

ABRAXAS, Ábrabax.

ABRE, *Abrus precatorius*.

ABRÉVIATION, Abbreviation.

ABRICOT, see *Prunus Armeniaca*—*a. Sauvage*, *Mammea Americana*.

ABRICOTIER, Prunus Armeniaca.

ABROSIA, Abstinence.

ABROTANUM, *Artemisia abrotanum* — a.
Cathsum, *Artemisia abrotanum* — a. Mas, *Artemisia abrotanum*.

ABROTONE, *Artemisia abrotanum*.

ABROTONI'TES, (*οινος*, 'wine,' understood.) Wine impregnated with *Artemisia Abrotanum* or Southernwood.

ABROTONUM, *Artemisia Abrotanum*.

ABRUPTIO. Abduction.

ABRUS PRECATORIUS, (from *abros*, 'elegant.') *Liq'orice Bush, Red Bean, Love pea.* (F.) *Abre, Liane à réglisse.* A small ornamental shrub, found from Florida to Brazil, as well as in Egypt and the West Indies; *Nat. Ord.* Leguminosæ. *Sec. Syst.* Monadelphæa Enneandria; having beautiful scarlet seeds with a black spot. The roots and leaves are sweet mucilaginous demulcents. The seeds of the American kind are considered to be purgative and poisonous. They are employed to form rosaries, and hence called, in mockery, *Jumble beads*.

ABSCCESS, from *abscedo*, (*abs*, and *cedere*), 'I depart,' or 'separate from.' *Absces'sus*, *Absces'sio*, *Aphiste'sis*, *Apost'e'ma*, *Ecy'e'ma*, *Ecy'e'sis*, *Reces'sus*, *Impos'thume*, *Gathering*. (Old Eng.) *Apostemacion*, *Apost'hume*. (Sc.) *Hattrel*. (Prov.) *Coul*, *Numpost*, *Postime*. (F.) *Abcès*, *Depôt*. A collection of pus in a cavity, the result of a morbid process. See *Pyogenia*, and *Suppuration*.

The French have various distinctive terms for Abscesses.

Abcès Chaud, Aigu, Soudain, is one which follows violent inflammation.

Abscès Froid, Chronique, Scrofuleux, cold, chronic, or scrofulous abscess, one which is the result of chronic or scrofulous inflammation.

Abscès par Congestion, A. diathésique, a symptomatic abscess; one which occurs in a part at a distance from the inflammation by which it is occasioned: e. g. a *lumbar abscess*; in which the inflammation may be in the lumbar vertebrae, whilst the pus exhibits itself at the groin.

ABSCCESS, ALVEOLAR, Parulis—a. Cold, see Abscess.

ABSCESS, METASTATIC. *Abscessus metastaticus*, (F.) *Abcès métastatique*, *A. consécutif*, an abscess which forms suddenly, and sometimes without any precursory signs of inflammation, in a part of the body remote from one in a state of suppuration, and without presenting a sufficient reason for its development in the place which it occupies. It is a consequence of phlebitis.

ABSCESS, PERFORATING OF THE LUNG, see Lung, perforating abscess of the—*a.* Perilaryngeal, see Perilaryngitis—*a.* Psosas, Lumbar abscess—*a.* Retropharyngeal, see Retropharyngeal—*a.* Shirtsud, *Abcès en bouton de chemise*.

ABSCESSUS CAPITIS SANGUINEUS NEONATORUM, Cephalæmatoma—*a.* Cerebri, Encephalopyosis—*a.* Gangrænescens, Anthrax—*a.* Gangranosus, Anthrax—*a.* Lacteus, Mastodynia apostematosa—*a.* Lumborum, Lumbar abscess—*a.* Mammæ, Mastodynia apostematosa—*a.* Metastaticus, Abscess, metastatic—*a.* Nucleatus, Furunculus—*a.* Oculi, Hypopyon—*a.* Pectoris, Empyema—*a.* Pulmonum, Pneumapostema—*a.* Renalis, Nephropostasis—*a.* Spirituosus, Aneurism—*a.* Thoracis, Empyema—*a.* Urinosus, Urapostema.

ABSCISSIO PRÆPUTII, Circumcision.

ABSCISSIO, Abscis'sio, Abscis'sio, from *abs-cidere* or *abscindere*, 'to cut off,' *Apoc'opë*, *Apothrausis*, *Diac'opë*. Excision or extirpation of a part, especially of a soft part.—Fabricius Hildanus.

Fracture or injury of soft parts, with loss of substance.—Hippocrates.

Diminution, or loss of voice.—Celsus.

Sudden and premature termination of a disease.—Galen.

ABSCONSO, Sinus.

ABSENCE DU BRUIT RESPIRATOIRE, see Murmur, respiratory.

ABSINTHE, Artemisia absinthium.

ABSINTHITES, αψινθίτης, Apsinth'ites, Wine impregnated with Absinthium or Wormwood.—Dioscorides.

ABSINTHIUM, (Ph. U. S.,) *Artemisia absinthium*—*a.* Marinum, *Artemisia maritima*—*a.* Maritimum, *Artemisia maritima*—*a.* Ponticum, *Artemisia pontica*—*a.* Romanum, *Artemisia pontica*—*a.* Santonicum, *Artemisia santonica*—*a.* Vulgare, *Artemisia absinthium*.

ABSORBANT, Absorbent.

ABSORBENT, Absorb'ens, from *absorbere* (*ab* and *sorbere*), 'to drink, to suck up.' (F.) *Absorbant*. That which absorbs.

ABSORBENT SYSTEM is the collection of vessels, *Vasa absorbentia* seu *resorbentia*, and glands, which concur in the exercise of absorption.

A medicine used for absorbing acidity in the stomach and bowels, as magnesia, chalk, &c. *Incor'tens, Resorb'ens, Sal'urans, Præcip'itans*.

Also, any substance, such as cobweb, sponge, &c., which, when applied to a bleeding surface, retains the blood, and forms with it a solid and adhesive compound, which arrests the hemorrhage.

ABSORPTIO, Absorption—*a.* Sanguinis, Hæmorrhæphesis.

ABSORPTION, Absorp'tio, Resorp'tio, Inhal'tio, Imbibit'io, Anar'rhophe, Anarrhophe'sis, Catapino'sis, Rhoëdë'sis, Catar'rhophe'sis, Catar'rhopê; same etymon. The function of absorbent vessels, by virtue of which they take up substances from without or within the body. Two great divisions have been made of this function.

1. *External absorption, or the absorption of composition*, which obtains, from without the organs, the materials intended for their composition; and, 2. *Internal absorption, or the absorption of decomposition*, which takes up from the organs the materials that have to be replaced by the exhalants.

By *external absorption* is meant not only that which takes place at the external surface of the body, but also that of the mucous membranes of the digestive and respiratory passages. Hence, again, the division of external absorption into *cutaneous—resorp'tio cuti'ne seu cutis, inhal'tio cutis,—intestinal or digestive, and pulmonary or respiratory*.

Internal absorption is also subdivided into, 1. *Molecular or interstitial, nutritive, organic, or decomposing*, which takes up from each organ the materials that constitute it, so that the decomposition is always in equilibrio with the deposition. 2. *The absorption of excremential secreted fluids*, such as the fluid of serous membranes, synovia, &c. As these are constantly exhaled on surfaces which have no external outlet, they would augment indefinitely, if absorption did not remove them in the same proportion as that in which they are deposited. 3. *The absorption of a part of the excremential fluids*, as they pass over the excretory passages.

Absorption does not effect the decomposition of the body immediately. It merely prepares the fluid which has to be eliminated by the secretory organs.

The great agents of external absorption are the veins and chyliiferous vessels; of internal absorption, probably the lymphatics. In the chyliiferous vessels and lymphatics the fluid is always found to possess the same general properties. In them, therefore, an action of elaboration or selection must have taken place. The veins, on the other hand, seem to exert no selection. Any fluid, possessing the necessary tenuity, passes through the coats of the vessel readily by imbibition, and proceeds along with the torrent of the circulation. Watery fluids in this manner enter the blood when they are taken into the stomach. Substances that require digestion, on the other hand, must pass through the chyliiferous vessels and thoracic duct.

ABSORPTION OF COMPOSITION, see Absorption—*a.* Cutaneous, see Absorption—*a.* of Decomposition, see Absorption—*a.* Digestive, see Absorption—*a.* External, see Absorption—*a.* of Excremential Secreted Fluids, see Absorption—*a.* Internal, see Absorption—*a.* Intestinal, see Absorption—*a.* Interstitial, see Absorption—*a.* Molecular, see Absorption—*a.* Nutritive, see Absorption—*a.* Organic, see Absorption—*a.* Pulmonary, see Absorption—*a.* of Excremential Secreted Fluids, see Absorption—*a.* Respiratory, see Absorption.

ABSTÈME, Abstemious.

ABSTÈMIOS, Abste'mius, Aoi'nos, from *abs*, 'without,' and *temetum*, 'wine.' (F.) *Abstème*. Used by the ancient writers, as well as by the French, in the sense only of its roots; one who abstains from wine or fermented liquors in general.

ABSTERGENTIA, Detergents.

ABSTERSIVA, Detergents.

ABSTERSORIA, Detergents.

ABSTINENCE, Abstin'en'tia, from *abs*, 'from,' and *tenerè*, 'to hold,' *Abros'ia, Asit'ia, Liman'chia, Limocet'ia*, Fasting. Privation, usually voluntary, as when we speak of *abstinence from pleasure, abstinence from drink*, &c. It is more particularly used to signify voluntary privation of certain articles of food. Fasting is a useful remedial agent in certain diseases, particularly in those of an inflammatory character.

ABSUS, a kind of cassia, C. Absus, which grows in Egypt and in India, the seeds of which, pulverized and mixed with powdered sugar, have been employed in form of a dry collyrium, in the endemic ophthalmia of Egypt.

ABULIA; from *a*, 'privative,' and *βουλη*, 'will.' Loss of the will, or of volition.

ABULICUS; same etymon. One who has lost the power of will or of volition.

ABUS DE SOI-MÊME, Masturbation.

ABUTA (a Guiana name), Pareira brava.

ABUTILON AVICENNÆ, *A. Cordatum*, *Sida Abutylon*, Indian or Yellow Mallova, Velvet leaf; Order, Malvaceæ, (from *a*, privative, 'bovs,' 'an ox,' and *τιλος*, 'diarrhœa,' that is, 'a remedy for the diarrhœa of cattle;') is naturalized, and common in most parts of the United States. It resembles common mallow in its properties, being mucilaginous and demulcent.

ABVACUATIO, an excessive or colliquative evacuation of any kind.

ACACIA, (Ph. U. S.) *Acaciæ gummi*—*a. Catechu*, *Catechu*—*a. False*, *Robinia pseudo-acacia*—*a. Germanica*, see *Prunus spinosa*—*a. Giraffæ*, see *Acaciæ gummi*—*a. Horrida*, see *Acaciæ gummi*—*a. Indica*, *Tamarindus*—*a. Nilotica*, see *Acaciæ gummi*—*a. Nostras*, see *Prunus spinosa*—*a. Senegal*, see *Senegal, gum*—*a. Vera*, see *Acaciæ gummi*—*a. Zeylonica*, *Hæmatoxylon Campechianum*.

ACACIE GUMMI, *Acacia*, from *ακν*, 'a point,' so called in consequence of its spines, *G. Acaciæ Arabiæ*, *G. Mimosæ*, *G. Arabicum*, *G. Acanthinum*, *G. Leucum*, *G. Thebaicum*, *G. Serapioidis*, *G. Lamac*, *G. Senega* or *Seneca*, (see *Senegal, gum*), *Gum Arabic*. (F.) *Gomme Arabique*. The gum of the *Acacia* seu *Mimosæ Nilotica*, *Acacia vera*, *Spina Egyptiaca*, of Upper Egypt, *Nat. Ord. Mimoseæ. Sex. Syst. Polygamma Monœcia*. It is in irregular pieces, colourless, or of a pale yellow colour, hard, brittle, of a shining fracture, transparent, soluble in water, and insoluble in alcohol, s. g. 1.4317.

It is mucilaginous; but is rarely used, except in pharmacy. Sometimes it is administered alone as a demulcent.

Acacia Horrida and *A. Giraffæ*, of South Africa, yield a good gum.

ACAJOU, (of Indian origin) *Anacardium occidentale*.

ACAJUBA OFFICINALIS, *Anacardium occidentale*.

ACALYPHA BETULINA, *Cauda felis agrestis*, (*a*, 'privative,' *καλος*, 'beautiful,' and *δφν*, 'touch') is employed by the natives of India as a stomachic and in cholera. It is given in the form of infusion of the leaves.

ACALYPHA HISPIDA, *Caturus spiciflorus*.

ACALYPHA INDICA, 'disagreeable to the touch.' *Cupameti*, Order *Euphorbiaceæ*. A common annual in the gardens of India. An infusion of the root, and a decoction of the leaves are cathartic.

ACALYPHA VIRGINICA. Three-seeded mer'cury, indigenous, flowering in August, is said to have expectorant and diuretic properties.

ACAMATUS, from *a*, priv., and *καμνω*, 'I labour.' This word has been sometimes used for a good constitution of the body. According to Galen, it means that position in which a limb is intermediate between flexion and extension; a position which may be long maintained without fatigue.

ACAMPسيا, *Contractura*.

ACANOS, *Onopordium acanthium*—*a. Spina*, *Onopordium acanthium*.

ACANTHA, Vertebral column. Also, Spinous process of a vertebra.

ACANTHABOLUS, *Acanthulus*, *Volseella*, from *ακανθα*, 'a spine,' and *βαλλω*, 'I cast out.' A kind of forceps for removing extraneous substances from wounds.—Paulus of Ægina, Fabricius ab Aquapendente, Scultetus, &c.

ACANTHALZUCA, *Echinops*.

ACANTHE FAUSSE, *Heracleum spondylium*.

ACANTHIA LECTULARIA, *Cimex lectularius*.

ACANTHIUM, *Onopordium acanthium*.

ACANTHULUS, *Acanthabolus*.

ACANTHUS MOLLIS, (*ακανθα*, 'a spine or thorn;') same etymon as *Acacia*. *Melampyllum*, *Branca ursina* seu *vera*, *Brankur'sine*, *Bear's Breech*. (F.) *Pied d'ours*. This plant is mucilaginous like *Althæa*, and is used as a demulcent.

ACAPATLI, *Piper longum*.

ACAR/DIA, from *a*, priv., and *καρδια*, 'the heart.' The state of a fœtus without a heart.

ACARDIOHÆMIA. (F.) *Acardiémie*, from *a*, priv., *καρδια*, 'heart,' and *δυνα*, 'blood.' Want of blood in the heart.—Piorry.

ACARDIONERVIA, (F.) *Acardionervie*: from *a*, priv., *καρδια*, 'heart;' and *νευρον*, 'nerve.' Want of nervous action in the heart as indicated by the sounds ceasing to be audible.

ACARDIOTROPHIA, Heart, atrophy of the.

ACARE, *Acarus*.

ACARICIDE, from *acarus*, and *cadere*, 'to kill.' A destroyer of acari,—as of the *acarus scabiei*.

ACARICO/BA. The Brazilian name for *Hydrocotyle umbellatum*, used by the Indians as an aromatic, alexipharmic, and emetic.

ACARON, *Myrica gale*.

ACARP/Æ, from *a*, 'privative,' and *καρπος*, 'fruit.' A division of the family of cutaneous diseases by Fuchs, in which there is no "fruit," (Germ. *Frucht*), or production from the cutaneous surface—tubercles, vesicles or pustules. Lentigo, Chloasma, Argyria, and Pityriasis belong to it.

ACARUS, from *a*, privative, and *καρπος*, 'divisible.' (F.) *Acare*. A minute insect, one species of which has been noticed, by several observers, in the itch. The *Acarus Scabiei*, see *Psora*.

ACARUS CIRO, see *Psora*—*a. Comedonum*, *Acarus Folliculorum*.

ACARUS CROSSEI, *Crosse mite*. An insect supposed by Mr. Crosse, of England, to have been developed in a solution of silicate of potassa when submitted to slow galvanic action, for the purpose of obtaining crystals of silex. It did not, however, prove to be a new formation.

ACARUS FOLLICULORUM, *A. Comedonum*, *Entozoon seu De'modex seu Sino'nea seu Steatozoon folliculorum*, *Macrogastrer platypus*. An articulated animalcule, discovered in the sebaceous substance of the cutaneous follicles. According to Professor Owen, it belongs to the Arachnida.

ACARUS SCABIEL, *Acarus*, see *Psora*.

ACATALEP/SIA, from *a*, privative, and *καταλαμβάνω*, 'I comprehend.' Uncertainty in diagnosis. Its opposite is *Catalepsia*.—Galen.

ACATAP/OSIS, from *a*, privative, and *καταπιωσις*, 'deglutition.' Incapacity of swallowing. Vogel has given this name to difficulty of deglutition.

ACATASTAT/IC, *Acatastat'icus*, from *a*, priv., and *καταστημι*, 'to determine.' An epithet given to fevers, &c., when irregular in their periods or symptoms.—Hippocrates.

ACATHAR/SIA, from *a*, priv., and *καθαρίζω*, 'I purge;' *Sordes*, Impurities. Omission of a purgative.—Fœsius.

ACATSAVAL/LI, a Malabar plant, which is astringent and aromatic. A bath of it is used in that country in cases of hemicrania. It is supposed to be the *Cassipoua filiformis* of Linnæus.

ACAWERIA, *Ophioxylum serpentinum*.

ACCABLEMENT, *Torpor*.

ACCELÉRATEUR DE L'URINE, *Accelerator urinæ*.

ACCELERATOR PARTUS, Abortive.

ACCELERATOR URINÆ, *Bulbo-cavernosus*, *Bulbo-urethral*—(Ch.), *Ejaculator Seminis*, *Bulbo-symphico-cavernæus*. (F.) *Accélérateur de l'urine*, *Bulbo-caverneux*, *Ano-caverneux*, from *ad* and *celer*, 'quick.' A muscle of the penis, which arises, fleshy, from the sphincter ani and membranous part of the urethra, and tendinous, from the crus and beginning of the corpus cavernosum penis. In its course it forms a thin, fleshy layer, the inferior fibres of which run more transversely than the superior, which descend in an oblique direction; the muscles of both sides completely enclosing the bulb of the urethra. It is inserted into its fellow by a tendinous line running longitudinally on the middle of the bulb. Its use is to propel the urine or semen forwards.

ACCENT, *Sonus vocis*, from *ad* and *canere*, *cantum*, to sing. Inflection or modification of the voice, which consists in raising or dropping it on certain syllables.

The accent exhibits various alterations in disease.

ACCÈS, Paroxysm.

ACCES'SION, *Accessio*, from *accedo*, (*ad* and *cedere*), 'I approach.' The invasion, approach, or commencement of a disease.

ACCESSOIRE, Accessory—*a. du long Fléchi-seur commun des orteils*: see *Flexor longus digitorum pedis profundus perforans* (accessorius)—*a. de l'Obturator interne*, *Ischio-trochanterianus*—*a. du Pied d'Hippocampe*: see *Cornu ammonis*—*a. du Sacro-lombaire*: see *Sacro-lumbalis*.

ACCESSORIUS FLEXOR LONGUS DIGITORUM PEDIS; see *Flexor longus digitorum pedis profundus perforans* (accessorius)—*a. Pedis hippocampi*;—see *Cornu ammonis*.

ACCESSORY, *Accessorius*, (F.) *Accessoire*, *Annexe*, same etymon. A consequence or dependence on any thing; as *accessory ligament*, *muscle*, *nerve*, &c.

ACCESSORY OF THE PAROTID is a name given by Haller to a small gland, which accompanies the parotid duct, and is commonly a mere prolongation of the parotid itself. See *Parotid*.

ACCESSORY SCIENCES TO MEDICINE are those which do not relate directly to the science of man in a state of health or disease; as physics, chemistry, &c.

ACCESSORY OF THE PAR VAGUM, Spinal nerve. The term *accessory* is also given to several muscles.

ACCESSUS, Coition.

ACCIDENS, Symptom—*a. Consecutifs*, Consecutive phenomena.

ACCIDENT, *Accidens*, from *accidere*, (*ad* and *cadere*), 'to happen.' A casualty; an unforeseen event. The French—like older English writers—use the term in nearly the same sense as *symptom*. It means also an unexpected symptom.

ACCIDENTAL, *Adventitious*. That which happens unexpectedly.

The French give the name *Tissus accidentels* to those adventitious textures, that are the result of a morbid process.

ACCIPITER, *Hæraz*, *lepaç*, 'the hawk,' from *accipere* (*ad* and *capio*), 'to take.' *Meneçratis* *Accipiter*, (F.) *Épervier*. A bandage applied over the nose, so called from its likeness to the claw of a hawk.

ACCLIMATED, *Clima'ti assue'tus*, (from *ad* and *clima*.) A word of recent introduction from the French, which means 'accustomed to a climate.'

ACCLIMATATION, Acclimation.

ACCLIMATEMENT, Acclimation.

ACCLIMATION, *Seasoning*. (F.) *Acclimate-*

ment, *Acclimation*. The act of becoming acclimated or accustomed to a climate.

The constitution of a person, who goes to live in another and a very different climate, usually experiences changes, which are frequently of an unfavourable character, and the study of which is of considerable importance in medicine.

ACCOMPAGNEMENT DE LA CATARACTE, Accompaniment of the cataract.

ACCOMPANIMENT, *Adjun'ction*. (F.) *Accompagnement*, (*compagnon*, 'an associate'). That which is joined to any thing.

Accompaniment to the cataract, (F.) *Accompagnement de la cataracte*, is a whitish, viscid substance, which sometimes surrounds the opaque crystalline lens, and remains after the operation for cataract, causing a secondary cataract.

ACCOUCHÉE, Puerpera.

ACCOUCHEMENT, Parturition—*a. Laborious*, *Dystocia*—*a. contre Nature*, see *Presentation*, *præternatural*—*a. Laborieux*, *Laborious labour*,—*a. Provoqué*, see *Parturition*.

ACCOUCHEUR, (F.) *Adju'tor Partûs*, *Obstet'ricans*, *Obstetri'cius*, *Maieu'ter*, *Maietu'tes*. He who practices the art of midwifery. A physician-Accoucheur, a Surgeon-Accoucheur, a Man-midwife, &c.

ACCOUCHEUSE, Midwife.

ACCOUPLEMENT, Coition.

ACCOUTUMANCE, Habit.

ACCRETION, *Accre'tio*, from *ad*, 'to,' and *crecere*, 'to increase.' Augmentation; (F.) *Accroissement*; also, increase by juxtaposition.

ACCRUISSEMENT, Accretion, Increase.

ACCUSATIO, Indication.

ACEDIA, *Incu'ria*, from *a*, privative, and *κηδος*, 'care.' Want of care, neglect. Also, fatigue.—*Hippocrates*.

ACELLA, Axilla.

ACBOGNOSIA, Pharmacognosia.

ACEOLOGIA, Materia Medica.

ACÉPHALE, Acephalous.

ACEPHALHÆMIA, (F.) *Acephalhémie*, from *a*, priv., *κεφαλη*, 'head,' and *αίμα*, 'blood.' Want of blood in the encephalon.—*Piorry*.

ACEPHALIA, see *Acephalous*.

ACÉPHALIE, see *Acephalous*.

ACEPHALOBRA'CHUS, from *a*, privative, *κεφαλη*, 'head,' and *βραχιων*, 'arm.' A foetus without head or arms.

ACEPHALOCAR'DIA, from *a*, priv., *κεφαλη*, 'head,' and *καρδια*, 'heart.' Monstrosity in which there is absence of head and heart.

ACEPHALOCHIRUS, from *a*, privative, *κεφαλη*, 'head,' and *χειρ*, 'hand.' A foetus without head or hands.

ACEPHALOCYST, *Acephalocystis*, (F.) *Acéphalocyste*, from *a*, privative, *κεφαλη*, 'head,' and *κυστις*, 'bladder.' A hydatiform vesicle, without head or visible organs, ranked amongst the Entozoa, although possessed of few animated characteristics. In no organ of the body are acephalocysts so frequently found as in the liver. Generally it is the 'multiple acephalocyst,' *A. socialis* seu *prolifera*, which is met with. At times, however, it is the 'solitary acephalocyst,' *A. eremita* seu *sterilis*.

The *acephalocystis endog'ena* has a firm coat, and is composed of different layers, which have numbers of smaller hydatids within them, and are thrown off from the interior of the parent cyst. This species has hence been termed *endogena*, to distinguish it from the *A. exog'ena* of ruminant animals, in which the young vesicles are developed from the exterior of the parent vesicles.—See *Hydatid*.

ACEPHALOGAS'TER, *Athoraceceph'alius*, from a privative, *κεφαλη*, 'head,' and *γαστηρ*, 'the

belly.' A name given to monsters devoid of head, chest, and abdomen; or to those which have an abdomen, but no chest or head.

ACEPHALOP'ODUS, from *a*, priv., *κεφαλη*, 'head,' and *πους*, *podos*, 'foot.' A monster devoid of head and feet. The condition is called *Accephalopod'ia*.

ACEPHALORA'CHIA, from *a*, priv., *κεφαλη*, 'head,' and *ραχis*, 'spine.' Monstrosity in which there is absence of head and spine.

ACEPHALOS'TOMA, from *a*, privative, *κεφαλη*, 'head,' and *στομα*, 'mouth.' An acephalous foetus, at the upper part of which there is an opening resembling a mouth.

ACEPHALOTHORA'CIA, from *a*, privative, *κεφαλη*, 'head,' and *θωραξ*, *thorax*, 'thorax.' Monstrosity in which there is absence of head and chest.

ACEPHALOTHO'RUS, from *a*, privative, *κεφαλη*, 'head,' and *θωραξ*, 'chest,' *Apctocephalus*. A monster devoid of head or chest.

ACEPH'ALUS, from *a*, privative, and *κεφαλη*, 'head.' (F.) *Acéphale*. A monster born devoid of head. The condition is called *Accephalia*. (F.) *Acéphalie*.

ACER, Acid.

ACER PALMIFOLIUM, A. Saccharinum.

ACER PENNSYLVANICUM, *Striped Maple*, *Striped Dogwood*. A decoction of the bark has been used internally and externally in cutaneous affections; and a decoction of the leaves and twigs is said to relieve nausea and vomiting.

ACER RUBRUM, *Red Maple*; Indigenous. The inner bark is a mild astringent. The Indians use a decoction of it for sore eyes.

ACER SACCHARINUM, A. *palmifolium*. *Maple*, *Sugar Maple*. (F.) *Érable*, Ord., *Aceraceæ*. This tree contains a large amount of sweet sap, whence a considerable quantity of sugar may be extracted. When purified, this sugar can scarcely be distinguished from that obtained from the cane.—See *Saccharum*.

ACERATES LONGIFOLIA, *Long-leaved green Milkweed*; Order, *Asclepiadaceæ*; indigenous, flowering in June and July; has the properties of the order. See *Asclepias*.

ACERATO'SIS, from *a*, privative, and *κερας*, *keras*, 'horn.' Defective development of the corneous tissue.

ACERB', *Acerbus*, *Stryphnos*, from *acer*, 'sharp.' A savour, or taste, compounded of the acid, bitter, and astringent; such as is met with in unripe fruits, &c.

ACER'CUS, from *a*, privative, and *κεκος* 'a tail.' A monster devoid of tail.—Gurlt.

ACE'RIDES, *Acerodes*, from *a*, privative, and *κηρος*, 'wax.' Plasters devoid of wax.—Galen.

ACERODES, *Acerides*.

ACERO'SUS, *Achyrodes*, *Pithyrinus*, from *αχυρος*, 'chaff,' *Purfura'ceus*. An epithet used by Hippocrates for the coarsest bread, made of flour not separated from the chaff.—Fœsius.

ACERVULUS CEREBRI. See *Pineal Gland*—a. *Glandulus Pinealis*, see *Pineal Gland*.

ACES'CENCY, *Acrescentia*, from *acrescere*, 'to grow sour,' (*akis*, 'a point,' *acer*, 'sharp.')

A disposition to acidity. The humourists believed that the animal humours are susceptible of this change.

ACESIA, Cure.

ACESINOSUS, Curative.

ACESIS, Curation, Cure, Medicament.

ACESMA, Medicament.

ACESMIUS, Curable.

ACESMUS, Cure.

ACESODYNES, Anodyne.

ACESOPHORUS, Curative.

ACESTER, Physician.

ACESTIS, Medicament.

ACESTOR, Physician.

ACESTORIA, Medicine.

ACESTORIS, Midwife.

ACESTOS, Curable.

ACESTRA, Needle.

ACESTRIA, Midwife.

ACESTRIS, Midwife.

ACESTRUM, Medicament.

ACETA MEDICATA, *Acetica*.

ACETABULA UTERINA, *Cotyledons*.

ACETAB'ULUM, (F.) *Acétabule*, from *acetum*, 'vinegar,' because it resembles the old vinegar vessel *oxybaph'ion*. A measure capable of containing the eighth part of a modern pint.—Athenæus. Galen. See *Cotyloid*. According to Castelli, the lobes or cotyledons of the placenta of ruminating animals have been so called.

ACETABULUM, *Cotyle*, see *Cotyloid*—a. *Humeri*, see *Glenoid*—a. *Marinum*, *Umbilicus marinus*.

ACETARIA, same etymon. A salad or pickle.

ACETAS, *Acetate*.

AC'ETATE, *Acetas*. (F.) *Acétate*. A salt formed by the union of the acetic acid with an alkaline, earthy, or metallic base. The acetates chiefly used in medicine are the acetates of ammonia, lead, potash, and zinc.

ACE'TICA, *Aceta Medica'ta*. (F.) *Vinaigres Médicinaux*. Pharmaceutical preparations of vinegar.

ACE'TICUM AC'IDUM, *Acidum Aceticum fortius seu forte purum seu glacialē, Acetum radicalē, Oxos, Acetic Acid, Strong Acetous Acid, Acidum Acetosum forte, Radical Vinegar, Spiritus Ven'eriis (when made from verdigris), Spirit of Verdigris*. Concentrated acetic acid, prepared by decomposing an acetate and receiving the acetic acid by distillation, has a very pungent and grateful odour, and an acid and acid taste. Its s. g. is about 1.046, and it is very volatile.

It is stimulant, rubefacient, and escharotic, and is applied to the nostrils in syncope, asphyxia, headache, &c. It destroys warts.

An *Aromatic Spirit of Vinegar*, *Ac'idum Aceticum Camphoratum*, A. *acetosum camphoratum*, is formed of this strong acid, $\frac{1}{3}$ x; *Camphor*, $\frac{3}{4}$ j; *Alcohol*, $\frac{1}{2}$ j.

A strong Acetic Acid was ordered by the London pharmacopœia, prepared from wood. It was called *Vinegar of wood*, *Improved distilled Vinegar*, *Pyroligneous Acid*, *Acetum Ligno'rum*, and its strength was such, that 87 gr. of crystallized subcarbonate of soda should saturate 100 grains of the acid.

Ac'idum Aceticum Dilut'um, A. A. *ten'ue seu debillius*, *Acetum destillatum*, *Acidum aceticum*, *Acidum acetosum destillatum*, *Distil'ed vinegar*, (F.) *Acide Acétique faible*, *Vinaigre distillé*, is prepared by distilling vinegar, until seven-eighths have passed over. An *Acidum aceticum dilutum*, *Diluted acetic acid*, is made by mixing a pint of the strong acetic acid with seven pints of distilled water.—Ph. U. S. Its properties are like those of vinegar.

ACETICUM MARTIALE, *Ferri Acetas*.

ACÉTOLAT, (F.) from *acetum*, *vinegar*. A liquid medicine resulting from the distillation of vinegar on one or more aromatic vegetable substances, and which are formed of vinegar and essential oils, or other volatile principles.—Béral.

ACÉTOLATURE, same etymon. A tincture, formed by digesting a vegetable substance or substances in vinegar.—Béral.

ACÉTOLÉ, same etymon, *Oxetolē*. A medicine, formed of distilled vinegar and medicinal principles dissolved in it.—Béral.

ACÉTOMEL, *Oxymel*.

AC'ETONE, *Ac'eton*, from *acetum*, 'vinegar.'

Epir'itus pyro-acet'icus lign'osus, *Pyro-acet'ic spirit*, *Pyro-acet'ic Ether*, *Mesit'ic Al'cohol*, *Bihydrate of Mesit'ylene*, (F.) *Acetone*; erroneously called *Naphtha* and *Wood Naphtha*. By others, however, the terms are applied to *Pyroxylic Spirit*. A limpid, colourless liquid, having a peculiarly penetrating and slightly empyreumatic odour. Its density in the liquid state, is almost the same as that of alcohol, 0.7921. Its taste is disagreeable, and analogous to that of peppermint. It is miscible in all proportions with water, alcohol, and ether. It may be prepared by distilling a mixture of two parts of crystallized acetate of lead and one part of quicklime in a salt-glaze jar (gray-beard,) the lower part of the jar being coated with fire-clay; and a bent glass tube, half an inch in diameter, adapted to the mouth by a cork, so as to form a distillatory apparatus. The jar is supported on the mouth of a small furnace, by which the lower part only is heated to redness, and the vapours are conducted into a Liebig's condenser. The product is repeatedly redistilled from quicklime, until its boiling point is constant at 132°.

It has been brought forward as a remedy in phthisis pulmonalis; but evidently with unfounded pretensions. It is an excitant, and may be serviceable in chronic bronchitis. The dose is ten to forty drops three times a day, diluted with water.

ACETOSA ALPINA, *Rumex alpinus*—a. *Nostras*, *Rumex acetosa*—a. *Pratensis*, *Rumex acetosa*—a. *Romana*, *Rumex scutatus*—a. *Rotundifolia*, *Rumex scutatus*—a. *Scutata*, *Rumex scutatus*—a. *Vulgaris*, *Rumex acetosa*.

ACETOSELLA, *Oxalis acetosella*.

ACETUM, *oξos*, *Oxos*, *Acetum Vini* seu *Bitar'nicum* seu *Gall'icum*, *Common Vinegar*, *Acidum aceto'sum*, *A'legar*, *Acetum Cerevis'iae*, (F.) *Vinaigre*; from *axis*, 'a point,' *acer*, 'sharp.' A liquor obtained by the acetous fermentation. Vinegar has a pungent odour, and a pleasant acid taste. One fluid ounce of the Acetum of the United States Pharmacopœia is saturated by about 35 grains of crystallized bicarbonate of soda. It is refrigerant in fevers; antiseptic, and anti-narcotic; and externally is stimulant and discutient.

Vinegar Whey is made by stirring a small wineglassful of vinegar, sweetened with a dessert spoonful of sugar, in a pint of milk; boiling for fifteen minutes, and straining. Like tamarind whey it is an agreeable drink in febrile affections.

ACETUM AROMATICUM, *A. Theriacale* seu *quatuor furum*, *Acidum Acet'icum Aromaticum*, *Thieves' Vinegar*, *Vinegar of the four Thieves*, *Marseilles Vinegar*, (F.) *Vinaigre Aromatique*, *V. des quatre voleurs*, (*Rorismarin*, *cacum*, *sicc.*, *Fol. Salvia* sing. $\frac{3j$. *Lavand. flor. sicc.* $\frac{3iv$. *Caryoph. cont.* $\frac{3ss$. *Acid. Acet.* $\frac{Oij$. Macerate 7 days, and filter.—Ph. E.) Odour, pungent and aromatic. Used as a perfume.

ACETUM BRITANNICUM, *Acetum*.

ACETUM CANTHARIDIS, *Vinegar of Cantharides*, (*Cantharid.* in pulv. $\frac{3ij$. *Acid. acet.* $\frac{f3v$. *Acid. pyrolign.* $\frac{f3xv$: *Euphorb.* in pulv. *crass.* $\frac{3ss$. Mix the acids; add the powders; macerate for seven days; strain; express strongly, and filter the liquor.—Ph. E. The London College macerates *cantharid.* $\frac{3ij$ in *acid. acet.* $\frac{Oj$. (Imp. meas.) for eight days; expresses and strains.) It is used as a prompt vesicant.

ACETUM COLCHICI, *Vinegar of meadow saffron*. (*Colchic. rad. contus.* $\frac{3ij$; *Acid. acet.* *dilut.* vel *Acet. destillat.* $\frac{Oij$; Ph. U. S. 1851. It may also be made by displacement.) It is used as a diuretic, and also in gout. Dose $\frac{f3ss$. to $\frac{f3iss$.

ACETUM DESTILLATUM; see *Acetium acidum*—a. *Gall'icum*, *Acetum*—a. *Lignorum*: see *Ac-*

ticum acidum—a. *Mulsum dulce*, *Oxyglyceus*—a. *Opil*, *Guttæ Nigræ*—a. *Plumbicum* seu *Saturninum*, *Liquor Plumbi subacetatis*—a. *Quatuor furum*, *Acetum Aromaticum*—a. *Radiale*, *Acetium Acidum*—a. *Rosatum*, *Oxyrrhodinum*.

ACETUM SCILLÆ, *Acidum Acet'icum Scillit'icum*, *Vinegar of Squills*, (F.) *Vinaigre scillitique*, (*Scilla contus.* $\frac{3iv$; *Acet. destillat.* $\frac{Oij$; Ph. U. S. It may also be made by displacement.) Diuretic, expectorant, and emetic. Dose $\frac{f3ss$ to $\frac{3ij$ as a diuretic and expectorant. It may also be made by the process of displacement.

ACETUM THERIACALE, *Acetum aromaticum*.

ACEYTE DE SAL. A remedy for bronchocele used in S. America. Roulin found it to contain a portion of iodine.

ACHACANA. A species of cactus, in the province of Potosi in Peru. Its root is thick and fleshy, and of a conical shape. It is a good edible, and is sold in the markets of the country.

ACHALYBHÆMIE. See *Chlorosis*.

ACHANACA. A plant of the kingdom of Mely in Africa. It is used by the natives as an antisiphilitic.

ACHAOVAN, a species of Egyptian chamomile.—Prosper Alpinus.

ACHAOVAN-ABIAT. The Egyptian name of *Cineraria maritima*, used in female diseases.

ACHAR, *Atchar*.

ACHE, *Pain*.

ACHE, *Apium graveolens*—a. *des Montagnes*, *Ligusticum levisticum*.

ACHEILIA, *Achi'lia*, from *α*, priv., and *χελος*, 'lip.' A malformation, consisting in a deficiency of a lip or lips.

ACHEILUS, *Achi'lus*: same etymon. One who is without lips.

ACHEIR, *Achir*, *De'manus*, from *α*, privative, and *χερ*, 'hand.' One devoid of hands.—Galen.

ACHEIRIA, *Achi'ria*: same etymon. The state of being devoid of hands.

ACHEROIS, *Populus*.

ACHIA, *Achiar*. A name given in India to the pickled shoots of the bamboo.

ACHIA, *Atchar*.

ACHIAR, *Achia*.

ACHIC'OLUM, *Achit'olus*, *Hidrote'rion*, *Su'da'rium*, *Fornix*, *Tholus*, *Sudato'rium*, *Su'datory Sweating-house*, *Sweating-bath*. The sweating-room in the ancient bagnios.

ACHILIA, *Acheilia*.

ACHILLE, TENDON D', *Achillis tendo*.

ACHILLEA AGERATUM, *A. visco'sa*, *Balsam'ita femin'ea*, *Eupato'rium mes'ues*, *Ageratum*, *Cos'tus horto'rum minor*, *Maudlin*, *Maudlin Tansey*; (F.) *Achillee Visqueuse*; *Ord. Compositæ*; *Sex. Syst.* Syngenesia Polygamia Superflua,—has the same properties as tansey, bitter and aromatic, and is used in like affections.

ACHILLEA ATRATA, *Herba Gen'ipi veri*, (F.) *Achillee Noire*, has similar virtues.

ACHILLEA MILLEFOLIUM, *Achillea Myriophyl'lon*, *Chrysoc'oma*, *Millefol'ium*, *Chilophyl'lon*, *Lumbus Ven'erie*, *Common Yarrow* or *Mi'foil*. (F.) *Millefeuille*, *Herbe aux charpentiers*. The leaves and flowers have an aromatic smell, and a rough, bitterish, somewhat pungent taste. They have been used in dyspepsia, flatulence, hemorrhage, &c. An extract of the plant, made with proof spirit, has been called *Achille'sum*; and is used by the Italians in intermittent fever.

ACHILLEA MOSCHATA, an Alpine plant, has been esteemed diaphoretic and vulnerary (?) A volatile oil called *Espirit d'Iva*, is obtained from it in Switzerland, which is much prized for its musky odour. It is said, by Mérat and De Lens, to be the *Genipi* of the Savoyards, which is usually referred to *Artemisia rupestris*.

ACHILLEA PTARMICA, *Ptar'mica*, *P. Vulga'*.

via, *Pseudo py'rethrum*, *Pyrethrum sylvestre*, *Draco sylvestris*, *Tarchon sylvestris*, *Sternutamentaria*, *Dracunculus Praten'sis*, *Sneeze-wort*, *Bastard Pel'titory*. (F.) *Herbe à éternuer*. The roots and flowers have a hot, biting taste, approaching that of pyrethrum. Their principal use is as a masticatory and sialogogue.

ACHILLEA VISCOSA, A. Ageratum.

ACHILLÉE NOIRE, *Achillea atrata* — a. *Viaqueuse*, *Achillea aegeratum*.

ACHILLEINUM, see *Achillea Millefolium*.

ACHILLE'IS. A beautiful species of barley, mentioned by Theophrastus and Galen, called after Achilles, a labourer. The decoction was used in fevers and jaundice. — Hippocrates.

ACHILLEUM, *Telephium*.

ACHIL'IS TENDO, *Funis Hippoc'ratidis*, *Corda seu Chorda Hippoc'ratidis*, *Corda magna*, *Nervus latus*, (F.) *Tendon d'Achille*. The strong tendon of the gastrocnemii muscles above the heel: so called, because it was the only vulnerable part of ACHILLES, or because of its strength. See Tendon.

ACHILUS, *Acheilus*.

ACHIMBASSI. An archiater or chief of physicians. A name given, at Grand Cairo, to a magistrate who licenses physicians.

ACHIR, *Acheir*.

ACHIRIA, *Acheiria*.

ACHITOLUS, *Achicolum*.

ACHLYS, *Caligo*.

ACHMELLA, *Spilanthus acmella*.

ACHNE. Lint. See *Lintum*. Also, small mucous flocculi seen in front of the cornea. — Hippocrates.

ACHOL'IA, from *α*, privative, and *χολη*, 'bile.' Deficiency or want of bile. Also, Asiatic cholera. A'CHOLUS: same etymon. One deficient in bile.

ACHOR, *Porrigio larvalis*.

ACHO'RES. A term often employed by the ancients to designate both *crusta lac'tea*, and small superficial ulcerations on the skin of the face and head. See *Porrigio Larvalis*.

ACHORES CAPITIS, *Porrigio scutellata*.

ACHORION SCHÖNLEINI. See *Porrigio favosa*.

ACHORIS'TUS, from *α*, priv., and *χωρίζω*, 'I separate.' Any sign which necessarily accompanies a state of health or disease.

ACHOUROU. The Carabæ name for a species of myrtle used in dropsy.

ACHRAS AUSTRALIS, *Sapota* — a. *Sapota*, *Sapota* — a. *Zapota*, *Sapota*.

ACHROI, *Achromatis'ti*, *Achro'mati*, *Achro'mi*, from *α*, privative, and *χρώμα*, 'colour.' Pale individuals. — Hippocrates. It is nearly synonymous with *λεῖψαιμι*, *leiphæ'mi*, persons without colour; bloodless.

ACHRO'MA, *Chloasma album*, *Leucopathi'a partialis acquis'ita*. Partial privation of colour of the skin.

ACHROMASIA, *Decoloration*.

ACHROMATI, *Achroi*.

ACHROMAT'IC, *Achromat'icus*: same etymon. A lens, so constructed as to correct the aberration of refrangibility of common lenses, is so termed. The *Crystalline* is an achromatic lens.

ACHROMATISTI, *Achroi*.

ACHROMATOPSIA, *Achromatop'sy*, *Chromatopseudop'sia*, *Chromatometablep'sia*, *Dyschromatop'sia*, *Parachro'ma*, *Pseudochro'mia*, *Parora'sia*, *Visus de'color*, *Colour blindness*, *Idiot'cy*, *Dal'tonism*, from *α*, privative, *χρώμα*, 'colour,' and *ορρομαι*, 'I see.' Incapability of distinguishing colours; a defect situate in the cerebral part of the visual organ. Persons so circumstanced have been termed, by Mr. Whewell, *Idiot's*. See *Acyanoblepsia* and *Anerythropsia*.

ACHROMI, *Achroi*.

ACHROMODERMIE, see *Albino*.

ACHROMOTRICHOMIE, see *Albino*.

ACHRYSTALLODIAPHANIE, see *Cata-ract*.

ACHYLO'SIS, from *α*, privative, and *χυλος*, 'juice, chyle.' Defective chylosis or formation of chyle.

ACHYMO'SIS, from *α*, privative, and *χυμος*, 'juice, chyme.' Defective chymification.

ACHYRANTHES REPENS, *Illice'brum polygonoides*, *Forty Knot*; indigenous. Order, *Amaranthaceæ*. A decoction of the plant is drunk as a diuretic in dropsy, ischuria, &c.

ACHYRODES, *Acerous*.

ACHYRON, *Furfur*.

A'CIA, from *akis*, a point. A word used by Celsus, which has puzzled commentators, — some believing it to have meant a needle; others the thread; and others, again, the kind of suture. "*Acia mollis, non nimis torta*." — Celsus, Galen. (Chifflet thinks it meant the thread. — Antwerp, 1638.)

ACID, *Ac'idus*, *Oxy*. (F.) *Acide*, *Aigre*, from *akis*, *akidos*, 'a point;' sharp; sour; especially as applied to odorous or sapid substances. The French also use the term *aigre*, when referring to the voice, in the sense of sharp and shrill: — *as une voix aigre, vox aspera*.

ACID, ACETIC, *Aceticum acidum* — a. *Acetic*, of commerce, *Pyroligneous acid* — a. *Acetic*, dilute, see *Aceticum acidum* — a. *Acetous*, strong, *Aceticum acidum* — a. *Aerial*, *Carbonic acid* — a. *Antimonious*, *Antimonium diaphoreticum* — a. *Arsenious*, *Arsenicum album* — a. *Auric*, see *Gold* — a. *Azotic*, *Nitric acid* — a. *Benzoic*, *Benjamin*, flowers of — a. *Benzuric*, *Hippuric acid* — a. *Bezoardic*, *Uric acid* — a. *Boric*, *Boracic acid* — a. *Calcareous*, *Carbonic acid* — a. *Calculus*, *Uric acid*.

ACID, CARBAZOT'IC, *Ac'idum Carbazot'icum*, *Carbonit'ric* or *Picric acid*, *Welter's Bitter*, (F.) *Acide Carbazotique*. This acid is frequently formed by the action of concentrated nitric acid on animal and vegetable substances. It, as well as the carbazotates, is tonic and astringent.

ACID, CARBONACEOUS, *Carbonic acid* — a. *Carbonitric*, *Acid Carbazotic* — a. *Carbonous*, *Oxalic acid* — a. *Caseic*, *Lactic acid* — a. *Chromic*, see *Chromic acid* — a. *Citric*, *Citric acid* — a. *Crotonic*, see *Croton tiglium* — a. *Cyanhydric*, *Hydrocyanic acid* — a. *Cyanohydric*, *Hydrocyanic acid* — a. *Ga-lactic*, *Lactic acid* — a. *Gastric*, *Gastric juice*.

ACID, GALLIC, *Ac'idum Gall'icum*. (F.) *Acide Gallique*. This acid is found in most of the astringent plants that contain tannic acid of the kind obtained from galls. It is in delicate silky needles, usually somewhat yellowish, inodorous, and of a harsh, somewhat astringent taste. It dissolves in one hundred parts of cold and three parts of boiling water. It is very soluble in alcohol, and but slightly so in ether.

It has been highly extolled in internal hemorrhage, especially from the urinary organs and uterus. Dose from ten to twenty grains.

The last Pharmacopœia of the United States (1851) directs it to be made by exposing a thin paste of powdered galls and distilled water for a month, adding the water from time to time to preserve the consistence; expressing the paste; boiling the residuum in distilled water; filtering through animal charcoal, and crystallizing.

ACID, HIPPU'RIC, *Ac'idum Hippu'ricum*, *Uro-benzoic*, *Urino-benzoic* or *Benzu'ric acid*. An acid found in the urine of graminivorous animals. It is contained in human urine, especially after benzoic acid has been taken. See *Hippuria*.

ACID, HYDRIO'IC, *Ac'idum Hydrio'dicum*. This acid is made by mixing solutions of iodide of potassium and tartaric acid; filtering the liquor

to separate the bitartrate of potassa, and adding water to make the resulting hydriodic acid of definite strength.

It has been used in the same cases as the preparations of iodine in general, but is rarely employed.

ACID, HYDROCHLORONITRIC, Nitro-muriatic acid—a. Hydrocyanic, Hydrocyanic acid—a. Hydrocyanic, dilute, see Hydrocyanic acid—a. Hydro-sulphuric, Hydrogen, sulphuretted—a. Hydrothionie, Hydrogen, sulphuretted—a. Igarisur: see *Jatropha curcas*.

ACID, IODIC, *Ac'idum Iod'icum*, (F.) *Acide Iodique*. This is obtained by boiling iodine with nitric acid; or by decomposing iodate of baryta by dilute sulphuric acid. It is a white, transparent solid, slightly deliquescent, and very soluble in water. It has been given with sulphate of quinia in hoarseness, scrofula, incipient phthisis, chronic inflammation, syphilis, etc. Dose three to six grains, or more.

ACID, LACTIC, see Lactic acid.

ACID OF LEMONS, Citric acid—a. Lithenic, Uric acid—a. Lithiasic, Uric acid—a. Lithic, Uric acid—a. Marine deplogisticated, Chlorine—a. Lactic, Lactic acid—a. Marine dulcified, Spiritus ætheris muriatici—a. Mephitic, Carbonic acid—a. of Milk, Lactic acid—a. Muriatic, see Muriaticum acidum—a. Muriatic, dilute, Muriaticum acidum—a. Nanceic, Lactic acid—a. Nitric, see Nitric acid—a. Nitric, dilute, see Nitric acid—a. Nitro-hydrochloric, Nitro-muriatic acid—a. Nitro-muriatic, see Nitro-muriatic acid—a. Nitrous, deplogisticated, Nitric acid—a. Oxypsetonic, Nitric acid—a. Picric, Acid, carbazotic—a. Polygalic, see Polygala senega—a. Prussic, Hydrocyanic acid—a. Pyroligneous, see Aceticum acidum—a. Pyrolignic, Pyroligneous acid—a. Rosacic, see Porphyria—a. of Sorrel, Oxalic acid—a. of Sugar, Oxalic acid—a. Sulphhydric, Hydrogen sulphuretted—a. Sulpho-hydric, Hydrogen sulphuretted—a. Sulphuric, see Sulphuric acid—a. Tannic, Tannin—a. Uric, Uric acid—a. Urino-benzoic, Acid, hippuric—a. Urobenzoic, A. Hippuric—a. Urous, Uric oxide—a. Urylic, Uric acid.

ACIDE ACÉTIQUE FAIBLE, see Aceticum acidum—a. Azotique, Nitric acid—a. Boracique, Boracic acid—a. Chromique, Chromic acid—a. Gallique, Acid, gallic—a. Hydrocyanique, Hydrocyanic acid—a. Hydrosulphurique, Hydrogen, sulphuretted—a. Iodique, Acid, iodie—a. Lactique, Lactic acid—a. Nitrique, Nitric acid—a. Phosphorique, Phosphoric acid—a. Prussique, Hydrocyanic acid—a. Pyroacétique, Pyroligneous acid—a. Pyrolignique, Pyroligneous acid—a. Sulfureux, Sulphurous acid—a. Sulfurique, Sulphuric acid—a. Sulfurique delayé, Sulphuricum acidum dilutum—a. Tannique, Tannin—a. Tartrique, Tartaric acid—a. Urique, Uric acid.

ACIDITATIO, Acidities.

ACID'ITIES, *Ac'o'ies, Acidita'tio, Ac'idum morbo'sum seu prim'um vi'rum, Or'ytes, Sordes ac'idæ*, (F.) *Aigreurs*. Sourness of the stomach, the result of indigestion, indicated by acid eructations, etc. The affection is very common in children, and must be obviated by absorbents, as magnesia, chalk, etc., and by regulated diet.

ACIDOLOG'IA, from *akis, akios*, 'a point, a sharp instrument,' and *logos*, 'a description.' A description of surgical instruments.

ACIDOM'ETER, (F.) *Acidomètre, Pèse-acide*, from *acid*, and *μετρον*, 'measure.' A hydrometer for determining the density of acids.

AC'IDS, *Ac'ida, Ac'o'ies*, are liquid, solid, or gaseous bodies, possessed of a sour, more or less caustic taste, and the principal character of which is the capability of saturating wholly or in part, the alkaline properties of bases.

Acids, in general, are refrigerant and antiseptic. Their particular uses are pointed out under the individual articles.

To **ACID'ULATE**, (F.) *Aiguiser, Aciduler*. To render acidulous, or slightly acid.

ACID'ULOUS, *Ac'id'ulus, Suba'cidus, Oxo'des, Oxo'i'des*, (F.) *Acidule, Aigrelet*. Substances are so called which possess a sourish taste, as tamarinds, cream of tartar, etc.

ACIDULOUS FRUITS. Oranges, gooseberries, etc.

ACIDULOUS WATERS, *Aqua Acidula*. Mineral waters containing carbonic acid gas sufficient to render them sourish. See Waters, mineral.

ACIDULOUS WATER, SIMPLE, *Aqua Ac'idi Carbon'ici* (Ph. U. S.) *Aqua aeris fixi seu acid'ula simplex. Liqueur seu Aqua Sodæ efferves'cens, Aqua Carbonat'is Sodæ acid'ula, Soda water, Mineral water*, (F.) *Eau acidule simple*, is water impregnated with fixed air.

Water, so impregnated, is cooling and slightly stimulating. It is used beneficially in dyspepsia, and in cases of vomiting, etc.

ACIDUM ACETICUM, Aceticum acidum—a. Aceticum aromaticum, Acetum aromaticum—a. Aceticum camphoratum, see Aceticum acidum—a. Aceticum dilutum, see Aceticum acidum—a. Aceticum e Ligno venale, Pyroligneous acid—a. Aceticum empyreumaticum, Pyroligneous acid—a. Aceticum glaciale, Aceticum acidum—a. Aceticum Scilliticum, Acetum scillæ—a. Acetosellæ, Oxalic acid—a. Acetosum, Acetum—a. Allantonicum, Allantoic acid—a. Amnicum, Amniotic acid—a. Arsenicosum, Arsenious acid—a. Arseniosum (Ph. U. S.), Arsenicum album—a. Azoticum, Nitric acid—a. Benzoicum, Benjamin, Flowers of—a. Benzoylicum, Benzoin, Flowers of—a. Boracicum, Boracic acid—a. Borussicum, Hydrocyanic acid—a. Carbazoticum, Acid, carbazotic—a. Carbonicum, Carbonic acid—a. Citricum, Citric acid—a. Gallicum, Acid, gallic—a. Hydriodicum, Acid, hydriodic—a. Hydrocarbonicum, Oxalic acid—a. Hydrochloricum, Muriaticum acidum—a. Hydrocyanicum, Hydrocyanic acid—a. Hydrocyanicum dilutum, see Hydrocyanic acid—a. Hydrothionicum liquidum, see Hydrosulphuretted water—a. Iodicum, Acid, iodie—a. Jatrophiicum, see *Jatropha curcas*—a. Lacticum, Lactic acid—a. Ligneum, Pyroligneous acid—a. Ligni pyroleosum, Pyroligneous acid—a. Ligorum empyreumaticum, Pyroligneous acid—a. Limonis, Citric acid—a. Lithicum, Uric acid—a. Marinum concentratum, Muriaticum acidum—a. Morbosum, Acidities—a. Muriaticum, Muriaticum acidum—a. Muriaticum dilutum, Muriaticum acidum—a. Muriaticum nitroso-oxygenatum, Nitro-muriatic acid—a. Nitri, Nitric acid—a. Nitri duleificatum, Spiritus ætheris nitrici—a. Nitricum, Nitric acid—a. Nitricum dilutum, Nitric acid—a. Nitro-muriaticum, Nitro-muriatic acid—a. Oxalinum, Oxalic acid—a. Phosphoricum, Phosphoric acid—a. Picricum, Acid, carbazotic—a. Primarum viarum, Acidities—a. Prussicum, Hydrocyanic acid—a. Pyroaceticum, Pyroligneous acid—a. Pyrolignum, Pyroligneous acid—a. Pyroxyllicum, Pyroligneous acid—a. Quercitannicum, Tannin—a. Sacchari, Oxalic acid—a. Saccharinum, Oxalic acid—a. Salis, Muriaticum acidum—a. Salis culinaris, Muriaticum acidum—a. Salis marini, Muriaticum acidum—a. Scytodephicum, Tannin—a. Septicum, Nitric acid—a. Succinicum, Succinic acid—a. Sulphuricum, Sulphuric acid—a. Sulphuricum alcoolisatum, Elixir acidum Halleri—a. Sulphuricum aromaticum, Sulphuric acid, aromatic—a. Sulphuricum dilutum, Sulphuric acid, diluted—a. Sulphuris volatile, Sulphurous acid—a. Sulphurosicum, Sulphurous acid—a. Tannicum, Tannin—a. Tartari essentialis, Tartaric acid—a. Tartaricum, Tartaric acid—a. Tartarosum, Tartaric acid—a. Uricum,

Uric acid—a. Urolithicum, Uric acid—a. Vitriolicum, Sulphuric acid—a. Vitriolicum aromaticum, Sulphuricum acidum aromaticum—a. Vitriolicum alcohole aromaticum, Sulphuricum acidum aromaticum—a. Vitriolicum vinosum, Elixir acidum Halleri—a. Zooticum, Hydrocyanic acid—a. Zootinicum, Hydrocyanic acid.

ACIDURGIA, Surgery (operative).

ACIER, Chalybs.

ACIES, Chalybs—a. Digitum manus, Phalanges of the fingers—a. Diurna, Hemeralopia.

ACINE, Acinus.

ACINE'SIA, *Acine'sis*, *Akine'sia*, *Immobilitas*, *Quies*, *Requies*, *Requie'tio*, *Esycheia*, *Erem'ia*, from *a*, privative, and *κίνησις*, 'motion,' *κίωω*, 'I move.' Rest. Immobility. Also, the interval between the systole and diastole of the heart—*Parasy'stolē*.

Under the term *Acineses*, Romberg includes the paralytic neuroses, or those that are characterized by defect of motive power.

ACINI OF MALPIGHI, Corpora Malpighiana.

ACINIFORMIS (TUNICA), Choroid, Uvea.

AC'INUS, *A. glandulosus*, from *ac'inus*, 'a grape-stone,' (F.) *Acine*. *A. glandiform corpuscle*, in which secretion was supposed to take place, and the excretory radicle to arise. Acini are the *globuli arteriarum termini* of Nichols. The term *ac'ini glandulosi* has also been given to glands which, like the pancreas, are arranged, as it were, in clusters. Glands thus formed have been called *glandulæ acino'se*. See Lobule.

ACIPENSER, see Ichthyocola.

ACIURGIA, Surgery (operative).

ACLEITROCARDIA, Cyanopathy.

ACMAS'TICUS, from *ακμή*, 'the top,' and *σῶω*, 'I remain.' A fever which preserves an equal degree of intensity throughout its course. It is also called *Homot'onos*. The Greeks gave it the name of *Epacmas'ticos*, and *Syn'ochos*, when it went on increasing, and *Paracmas'ticos*, when it decreased.—Galen.

ACMÉ, *Vigor*, *Cor'yphe*, *Culmina'tio*, *Status*, *Fastig'ium*. The period of a disease at which the symptoms are most violent. *Arche*, *Ἀρχή*, is 'the commencement;' *anab'asis*, *ἀναβ'asis*, 'the period of increase;' and *acmé*, *ακμή*, 'the height,' (F.) *État*.

ACMELLA, *Spilanthus acemella*—a. Mauritiانا, *Spilanthus acemella*.

ACMON, Incus.

ACNÉ, *Acne*, *Ion'thus varus*, *Varus*, *Psyr'ia*, *Acne*, *Stone Pock*, *Whelk*, *Bubule*, (F.) *Acné*, *Dartre pustuleuse disséminée*. A small pimple or tubercle on the face.—Goræus. Fœsius thinks the word ought to be *Acmé*; and, according to Cassius, it is, at all events, derived from *ακμή*, 'vigour;' the disease affecting those in the vigour of life especially.

Willan and Bateman have adopted the term in their Nosology of cutaneous diseases, and placed it in the order TUBERCULA. *Acne*, with them, is an eruption of distinct, hard, inflamed tubercles, sometimes continuing for a considerable length of time, and sometimes suppurating slowly and partially. They usually appear on the forehead, temples, and chin, and are common to both sexes; but the most severe forms are seen in young men. They require but little management, and consist of four varieties: *Acne indur'ata*, *A. simplex*, (*Haploëné*), *A. punct'ata* (*Ion'thus varus punct'atus*, *Punctæ mucosæ*, *Crino'nes*, *Comedo'nes* or *Maggot Pimple*, *Worms*, *Grubs*), and *A. rosacea*.—See Gutta Rosea.

ACNE MENTAGRA, Sycosis—a. Rosacea, Gutta Rosea—a. of the Throat, Pharyngitis, follicular.

ACNE'MIA, *Akne'mia*, from *a*, privative, and *κνήμη*, 'the leg.' Absence of legs.

ACNÉ, *Acne*—a. *Molluscoïde*, *Molluscum*.

ACNES'TIS, from *a*, privative, and *σκαῖν*, 'to scratch.' The part of the spine which extends, in quadrupeds, from between the shoulders to the loins. According to Pollux, the middle of the loins. The vertebral column.

ACNESTOS, Cneorum tricocum.

ACOE, Audition, Ear.

ACE'LIOS, from *a*, privative, and *κοιλία*, 'belly.' Devoid of belly. One who is so emaciated as to appear to have no belly.—Galen.

ACOËMETER, Acoumeter.

ACOËMETRUM, Acoumeter.

ACOËNOSI, Aconusi.

ACOËSIS, Audition.

ACOGNOSIA, Pharmacognosia.

ACOLASIA, Intemperance.

ACOLOGY, Materia Medica.

ACONE, Mortar.

ACONIT À GRANDS FLEURS, Aconitum cammarum—a. *Salutaire*, Aconitum anthora.

ACONITA, see Aconitum napellus.

ACONITE, Aconitum.

ACONITI FOLIA, see Aconitum—a. Radix, see Aconitum.

ACONITIA, see Aconitum napellus.

ACONITIN, see Aconitum napellus.

ACONITINE, see Aconitum napellus.

ACONITIUM, see Aconitum napellus.

ACONITUM, from *Ac'onā*, a place in Bithynia, where it is common. *Cynoc'oton*, *Pardalian'ches*, *Pardalian'chum*, *Canic'ida*, *Ac'onite*, *Wolfsbane*, *Monkshood*. Ord. Ranunculaceæ. *Sex. Syst.* Polyandria Trigynia.

ACONITUM, Aconite, in the Pharmacopœia of the United States, 1842, is the leaves of Aconitum napellus, and A. paniculatum. In the last edition, 1851, *Aconiti folia* is the official name for the leaves, *Aconiti radix* for that of the root.

ACONITUM AN'THORA, *A. Salutiferum* seu *Nemoro'sum* seu *Candol'lei* seu *Jacquini* seu *Eu'tophum* seu *Anthoroidium*, *Anth'ora vulg'ria*, *An'th'ora*, *Anth'ora*, *Salutary Monkshood*, *Wholesome Wolfsbane*, *Yellow helmet flower*, (F.) *Aconit salutaire*. The root of this variety, as of all the rest, is poisonous. It is used as a cathartic and anthelmintic. Dose ℥ss to ℥j.

ACONITUM ANTHOROIDEUM, A. anthora.

ACONITUM CAM'MARUM, *A. paniculatum*, *A. macran'thum*, *A. Kuneszov'vii*, (F.) *Aconit à grands fleurs*, resembles Aconitum Napellus in properties.

ACONITUM CANDOLLEI, A. anthora—a. Eulophum, A. anthora.

ACONITUM FEROX, a most violent ranunculaceous poison, which grows in Nepal, and is the *Bish* or *Bikh* poison of that country. Its effects are like those of the Aconitum Napellus.

ACONITUM JACQUINI, A. anthora—a. Kusnezovii, A. cammarum—a. Macranthum, A. cammarum.

ACONITUM NAPELL'US, *Aconitum Napel'lus verus*, *Common Monkshood* or *Wolfsbane*, *A. Nemonta'num*, (F.) *Chaperon de Moine*. The leaves are narcotic, sudorific, and deobstruent (?) They have been used in chronic rheumatism, scrofula, scirrhus, paralysis, amaurosis, etc. The active principle is called *Aconit'ia*, *Aconit'ina*, *Aconit'a*, *Aconit'ium* or *Aconitine*. A form for its preparation is contained in the Ph. U. S. (1851). It is made by treating an alcoholic extract of the root with dilute sulphuric acid; precipitating by solution of ammonia; dissolving the precipitate in dilute sulphuric acid; treating with animal charcoal; again precipitating with solution of ammo-

mia; washing with water, and drying. It requires 150 parts of cold and 50 of boiling water to dissolve it, but is readily dissolved by alcohol and ether. It neutralizes the acids, and forms with them uncrystallizable salts. It has been used internally, and especially applied externally, in neuralgic cases, iatroleptically and endermically. Dose of Aconitum, gr. j. to gr. iij.

ACONITUM NEMOROSUM, *A. anthora*—*a. Neomontanum*, *A. napellus*—*a. Paniculatum*, *A. cammarum*—*a. Racemosum*, *Actaea spicata*—*a. Salutiferum*, *A. anthora*.

ACONITUM UNCINATUM, *Wild Monkshood*, indigenous, is said to have similar virtues with *Aconitum napellus*.

ACONUSI, *Acœn'osi*, *Acœn'osi*, from *akon*, 'audition,' and *vosos*, 'disease.' *Morbi aurium et auditus*. Diseases of the ear and audition.

ACONOSI, *Aconusi*.

ACOPIS. Same etymon as the next. Pliny gives this name to a precious stone, which was boiled in oil and used against weariness.

ACOPON, from *a*, privative, and *kopos*, 'weariness.' A remedy against weariness—*Fœsius*, *Goræus*, &c. *Ac'opum*,—*Celsus*, *Pliny*. See *Anagryris*.

ACOPRIA, Constipation.

ACOPROSIS, Constipation.

ACOR BENZOÏNUS, *Benjamin*—*a. Boracicus*, Boracic acid—*a. Succineus*, Succinic acid—*a. Sulphuris*, Sulphuric acid—*a. Tartaricus*, Tartaric acid.

ACORE BÂTARD, *Iris pseudacorus*—*a. Fauca*, *Iris pseudacorus*—*a. Odorant*, *Acorus calamus*.

ACORES, Acids, and Acidities.

ACOR'IA, from *a*, privative, and *kopew*, 'I satiate.' An inordinate or canine appetite.—*Hippocrates*.

ACOR'ITES. A wine made of Acorns.—*Dioscorides*.

ACOR'MUS, from *a*, privative, and *kopmos*, 'trunk.' A monster devoid of a trunk.—*Gurlt*.

ACORN, JUPITER'S, *Fagus castanea*—*a. Oily*, *Guilandina moringa*—*a. Sardinian*, *Fagus castanea*.

ACORNS. See *Quercus alba*.

ACORUS ADULTERINUS, *Iris pseudacorus*.

ACORUS CALAMUS, *A. Versu* seu *Brasiliensis*, *Callamus Aromaticus* seu *Odoratus* seu *Vulgaris*, *Typha Aromatica*, *Clava Rugs'sa*, *Sweetflag* or *Acorus*, *Flagroot*, *Sweet cane*, *Myrtle Flag*, *Sweet grass*, *Sweet root*, *Sweet rush*. (F.) *Jonc roseau* ou *Canne aromatique*, *Acore odorant*. *Ord. Aroideæ*; *Acoraceæ*. (Lindley.) *Sex. Syst. Hexandria Monogynia*. The rhizome—*Callamus* (Ph. U. S.)—is stomachic and carminative, but is rarely used. It is regarded as a good adjuvant to bark and quinia in intermittents.

ACORUS PALUSTRIS, *Iris pseudacorus*—*a. Vulgaris*, *Iris pseudacorus*.

ACOS, Medicament.

ACOS'MIA, from *a*, privative, and *kosmos*, 'order, ornament.' Disorder, irregularity in the critical days, according to *Galen*, who uses the word *kosmos* for regularity in those days. Others, and particularly *Pollux*, call bald persons *acosmoi*, because they are deprived of one of their most beautiful ornaments.

ACOU'METER, *Acœu'm'eter*, *Acœm'eter*, *Acœm'etrum*, *Acu'meter*, *Acusim'eter*, (F.) *Acoumètre*, from *akouo*, 'I hear,' and *metron*, 'measure.' An instrument designed by *M. Itard* for measuring the degree of hearing.

ACOUNÈTRE, *Acoumeter*.

ACOUOPHO'NIA, *Cophonia*; from *akouo*, 'I hear,' and *φωνη*, 'voice.' "Auditory Percussion." A mode of auscultation, in which the

observer places his ear on the chest, and analyzes the sound produced by percussion.—*Donné*.

ACOUS'MA, an imaginary noise. Depraved sense of hearing.

ACOUS'TIC, *Acus'ticus*, (F.) *Acoustique*. That which belongs to the ear; as *Acoustic nerve*, *Acoustic trumpet*.

Acoustic MEDICINE is one used in diseased audition.

ACOUSTICO-MALLÉEN, *Laxator tympani*. *Acous'tics*, *Acus'tica*. (F.) *Acoustique*. The part of physics which treats of the theory of sounds. It is also called *Phonics*.

ACOUSTIQUE, *Acoustic*, *Acoustics*.

ACQUA BINELLI, *Aqua Binelli*—*a. Brocchieri*, *Aqua Brocchieri*—*a. Monterossi*, *Aqua Binelli*—*a. di Napoli*, *Liquor arsenicalis*—*a. della Toffana*, *Liquor arsenicalis*.

ACQUETTA, *Liquor Arsenicalis*.

ACQUI, MINERAL WATERS OF. These thermal sulphureous springs are in *Piémont*. Their temperature is 167° Fahr., and they contain sulphohydric acid and chloride of sodium.

ACQUIRED DISEASES, *Morbi acquis'iti*, *M. advent'iti*, *M. epict'iti*. *Adventitious diseases*. (F.) *Maladies acquises*. Diseases which occur after birth, and which are not dependent upon hereditary predisposition.

ACRAI'PALA, from *a*, privative, and *κραταιη*, 'drunkenness.' Remedies against the effects of a debauch.—*Gornæus*.

ACRA'LEA, from *akros*, 'extremity.' The extreme parts of the body, as the head, hands, feet, nose, ears, &c.—*Hippocrates* and *Galen*. See *Acra*.

ACRA'NIA, from *a*, privative, and *κρανιον*, 'the cranium.' Want of cranium, wholly or in part.

ACRA'SIA, from *a*, privative, or 'bad,' and *κρασις*, 'mixture.' Intemperance. Excess of any kind. Surfeit.

It has been employed to denote debility, synonymously with *Acratia*; but this may have been a typographical inaccuracy.

ACRAT'IA, from *a*, privative, and *κρατος*, 'strength.' Impotence; weakness, fainting.

ACRATIS'MA, from *a*, privative, and *κραννυμι*, 'to mix.' A breakfast, consisting of bread steeped in wine, not mixed with water.—*Galen*, *Athenæus*.

ACRATOM'ELI, from *akparov*, 'pure wine,' and *μελι*, 'honey.' Wine mixed with honey.

ACRATOPE'GÆ, *Akratope'gæ*, from *a*, privative, *κρατος*, 'strength,' and *πνυγ*, 'a spring.' Mineral waters having no marked chemical qualities.

ACRATOPOS'IA, from *acratum*, and *ποσις*, 'drink.' The drinking of pure or unmixed wine.

A'CRATUM, *akparov*, from *a*, privative, and *κρατος*, 'strength.' Unmixed wine, *Acetum vinum*, *Vinum merum*.

ACRATURE'SIS, from *acratia*, 'weakness,' and *ουρησις*, from *ουρον*, 'urine,' 'the act of discharging the urine.' Inability to void the urine, from paralysis of the bladder.

ACRE. The extremity or tip of the nose.

A'CREA, *Acrote'ria*, from *akros*, 'the summit.' The extreme parts of the body, as the feet, hands, ears, &c.

Also the extreme parts of animals that are used as food, *Acroco'lia*.

ACRID, from *akros*, 'pointed,' or from *ακρῖς*, 'a point,' *Acer*. An epithet for substances which occasion a disagreeable sense of irritation or of constriction at the top of the throat.

Acrid heat, (F.) *Chaleur âcre*, is one that causes a hot tingling sensation at the extremities of the fingers.

ACRID POISON, See *Poison*.

ACRIDS, in Pathology, are certain imaginary

substances, supposed by the humourists to exist in the humours, and to cause various diseases. See Acrimony.

ACRIDOPH'AGI, from *ακρις*, *ακριδος*, 'a locust,' and *φαγω*, 'I eat.' Locust-eaters. Acridophagous tribes are said to exist in Africa.—Strabo.

ACRIMONY, *Αει'ιτας*, *Acrimo'nia*, from *αερ*, 'acid,' *ακis*, 'a point.' Acrimony of the humours. An imaginary acid change of the blood, lymph, &c., which, by the humourists, was conceived to cause many diseases.

ACRIN'IA, from *α*, privative, and *κρινω*, 'I separate.' A diminution in the quantity, or a total suspension, of the secretions.

ACRIS, a sharp bony prominence. Also, the locust.

ACRISIA, *Ακρι'σις*, from *α*, privative, and *κρισις*, 'judgment.' A condition of disease, in which no judgment can be formed; or in which an unfavourable opinion must be given.—Hipp. and Galen.

ACRISIS, *Acrisia*.

ACRITICAL, *Ακρι'τος*, (F.) *Acritique*, from *α*, privative, and *κρισις*, 'judgment.' That which takes place without any crisis, or which does not foretell a crisis; as a *critical symptom*, *abscess*, &c.

ACRITOS, *Acritical*.

ACRIVIOLA, *Tropeolum majus*.

ACROBYTIA, *Acropos'thia*, from *ακρος*, 'top,' and *βυω*, 'I cover.' The extremity of the prepuce.—Hippocrates, Rufus.

ACROBYSTITIS, *Posthitis*.

ACROCHEIR', *Acrocheir'*, *Acrocheir'on*, from *ακρος*, 'extremity,' and *χειρ*, 'the hand.' The forearm and hand.—Goræus. Also, the hand.

ACROCHORDON, from *ακρος*, 'extremity,' and *χορδον*, 'a string.' A tumour which hangs by a pedicle. A kind of hard wart, *Verruca penicillis*.—Aëtius, Celsus.

ACROCHORISMUS, from *ακρος*, 'extremity,' and *χορευω*, 'I dance.' A kind of dance, with the ancients, in which the arms and legs were violently agitated.

ACROCOLIA, *Acrea*.

ACROCOLIUM, *Acromion*.

ACROD'RYA, from *ακρος*, 'extremity,' and *δρυς*, 'a tree.' Autumnal fruits, as nuts, apples, &c.

ACRODYN'IA, *Erythema acro'dynum*, *E. acro'dyn'ia*, *Chiro-podalgia*, (F.) *Acro'dynia*, *Mal des pieds et des mains*, from *ακρος*, 'extremity,' and *δυνω*, 'pain.' A painful affection of the wrists and ankles especially, which appeared in Paris as an epidemic, in 1828 and 1829. It was supposed by some to be rheumatic, by others to be owing to spinal irritation. It appears to have been the same as Dengue.

ACROLENION, *Olecranon*.

ACROMASTIUM, *Nipple*.

ACROMIA, *Acromion*.

ACROMIAL, *Acromia'lis*. Relating to the Acromion.

ACROMIAL ARTERY, *External Scapular*, *A. Arter'ia Thorac'ica humera'lis*, *Artère troisième des Thoraciques*,—(Ch.) *A. Thoracique humérale*, arises from the anterior part of the axillary artery, opposite the upper edge of the pectoralis minor. It divides into two branches: one, *superior*; the other, *inferior*,—the branches of which are distributed to the subclavians, serratus major anticus, first intercostal, deltoid, and pectoralis major muscles, as well as to the shoulder joint, &c. They anastomose with the superior scapular, thoracic, and circumflex arteries.

ACROMIAL NERVES, *Nervi acromia'les*. Branches of the fourth cervical nerve, which are distributed to the acromial region.

ACROMIAL VEIN has the same arrangement as the artery.

ACRO'MIO-CORACOIDEUS. Belonging to the acromion and coracoid process.

The triangular ligament between the acromion and coracoid process of the scapula is so called.

ACRO'MION, *Acro'mium*, *Acro'mia*, *Acro'mis*, from *ακρος*, 'the top,' and *ωμος*, 'the shoulder.' Or *Acro'mii*, *Humerus summus*, *Arnus summus*, *Mucro hu'meri*, *Rostrum porci'nium*, *Caput Scap'ulae*, *Acro'mium*. The process which terminates the spine of the scapula, and is articulated with the clavicle.

ACROMIS, *Acromion*.

ACROMPHALIUM, *Acromphalon*.

ACROMPHALON, *Acromphal'ium*, from *ακρος*, 'the top,' and *ομφαλος*, 'the navel.' The extremity of the umbilical cord, which remains attached to the foetus after birth.

ACROMYLE, *Patella*.

ACRO-NARCOTIC, See *Poison*.

ACRONYX, from *ακρος*, 'the summit,' and *ονυξ*, 'the nail.' Growing in of the nail.

ACROPARALYSIS, from *ακρος*, 'extremity,' and *παρالىσις*, 'palsy'; *Paral'ysis extremitat'um*, Palsy of the extremities.—Fuchs.

ACROPOSTHIA, *Acrobystia*.

ACROPOSTHITIS, *Posthitis*.

ACROPSI'ON, from *ακρος*, 'extremity,' and *ψιλος*, 'naked.' The extremity of the glans penis.

ACROBIA, *Vertex*.

ACROBRIHEUMA, *Rheumatis'mus extremitat'um*, from *ακρος*, 'extremity,' and *ρευμα*, 'defluxion, rheumatism.' Rheumatism of the extremities.

ACROS, *ακρος*, 'extremity, top.' The strength of the Athletæ, and of diseases; the prominences of bones; the extremities of the fingers, &c. See *Acrocheir*, *Acromion*, &c.

ACROSTICHUM POLYPODIOIDES, *Polypodium incanum*.

ACROTERIA, *Acrea*. See *Extremity*.

ACROTERIASIS, *Acroteriasmus*.

ACROTERIASMUS, *Acroteri'asis*, from *ακροτηρια*, 'the extremities'; hence *ακροτηριαζω*, 'to mutilate.' Amputation of the extremities.

ACROTHYMTON, from *ακρος*, 'top,' and *θυμον*, 'thyme.' A kind of conical, rugous, bloody wart, a 'moist wart,' compared by Celsus to the flower of thyme.

ACROTICA, from *ακρος*, 'summit.' Diseases affecting the excrent functions of the external surface of the body.

Pravity of the fluids or emunctories that open on the external surface; without fever or other internal affection as a necessary accompaniment.

The 3d order of the class *Eccrit'ica* of Good.

ACROTISMUS, *Asphyxia*.

ACT, *Actus*, from *actum*, past participle of *agere*, 'to do,' 'a thing done.' The effective exercise of a power or faculty. The action of an agent. *Acte* is used by the French, to signify the public discussion, which occurs in supporting a thesis;—thus, *soutenir un Acte aux Ecoles de Médecine*, is, 'to defend a Thesis in the Schools of Medicine.'

ACTA, *Gesta*.

ACTÆA CIMICIFUGA, *A. racemosa*.

ACTÆA RACEMO'SA, *A. Cimicif'uga*, *Cimicif'uga racemo'sa*, *Macro'trys racemo'sa*, *Bot'rophis Serpenta'ria* (?), *Serpenta'ria nigra*, *Black snake-root*, *Richweed*, *Cohosh*, *Squaw root*, *Rattleweed*, *Black Cohosh*. (F.) *Actée à grappes*, *Serpentaire noire*. *Ord. Ranunculaceæ*. *Sax. Syst. Polyanthia Pentagynia*. A common plant in the United States. The root, *Cimicif'uga* (Ph. U. S.), is astringent; and, according to Barton, has been successfully used, in the form of a decoction, as a gargle in putrid sore throat. A decoction of the root cures the itch. It is acro-narcotic, and

has been used in rheumatism, acute and chronic; chorea, &c.

ACTÆA SPICA'TA, *Christophoria'na spica'ta*, *Aconitum racemo'sum*, Baneberry, Herb *Christophæ*. (F.) *Herbe St. Christophe*. A perennial herbaceous European plant, the root of which resembles that of the black hellebore. The root is cathartic, and sometimes emetic, and in overdoses may produce dangerous consequences.

Actæa America'na, of which there are two varieties, *A. alba* and *A. rubra*,—white and red *cohosh*,—is indigenous in the United States. It has the same properties as *A. spicata*.

ACTE, Sambucus.

ACTE, Act.

ACTÉE à GRAPPES, *Actæa racemosa*.

ACTIF, Active.

ACTIO, Action, Function.

ACTION, *Act'io*, *Opera'tio*, *Energi'a*, *Praxis*: from *agere*, *actum*, 'to act.' Mode in which one object influences another.

The *animal actions* are those that occur in the animal body: the *vital*, those that are essential to life: the *physiological*, those of a healthy character: the *pathological* or *morbid*, those that occur in disease, &c. The ancients divided the *physiological actions* into *vital*, *animal*, *natural*, *sexual*, *particular*, *general*, &c. See *Function*.

ACTIONES NATURALES, see *Function*.

ACTIVE, same etymon. *Dras'ticus*, *Act'ivus*, *Sthen'icus*, *Hypersthen'icus*. (F.) *Actif*. This adjective is used, in *Pathology*, to convey the idea of superabundant energy or strength. *Active symptoms*, e. g. are those of excitement. In *Therapeutics*, it signifies *energetic*:—as, an *active treatment*. The French use the expression *Médecine agissante*, in contradistinction to *Médecine expectante*. In *Physiology*, *active* has a similar signification, many of the functions being divided into active and passive.

ACTON. A village near London, at which there is a purgative mineral spring, like that at Epsom.

ACTUAL. Same etymon as *active*. That which acts immediately. A term usually restricted to the red-hot iron, or to heat in any form; in contradistinction to the *potential* or *virtual*, which is applied to caustics or escharotics.

ACTUARIUS. Originally a title of dignity given to the Byzantine physicians.

ACTUS PARTURITIONIS, Parturition.

ACUITAS, Acrimony.

ACULEUS LIGNEUS, Splinter.

ACUMETER, Acoumeter.

ACUPUNCTURE, *Acupunctu'ra*, *Acupuncta'tion*, *Acupunctura'tion*, from *acus*, 'a needle,' and *punctura*, 'a puncture,' A surgical operation, much in use amongst the Chinese and Japanese, which consists in puncturing parts with a very fine needle. It has been employed, of late years, in obstinate rheumatic affections, &c., and apparently with success. Acupuncture is likewise a mode of infanticide in some countries; the needle being forced into the brain through the fontanelles, or into the spinal marrow, &c.

ACURGIA, Surgery (operative).

ACUS, Needle—a. *Capitata*, Pin—a. *Invaginata*, see *Needle*—a. *Ophthalmica*, see *Needle*—a. *Paracenta*, Trocar—a. *Paracenetica*, Trocar—a. *Triquetra vulgaris*, Trocar—a. *Veneris*, *Eryngium campestre*.

ACUSIMETER, Acoumeter.

ACUSIS, Audition.

ACUSTICA, Acoustics.

ACUSTICUS, Auditory.

ACUTE, *Acu'tus*, *Oxy's*, *ἄκς*, (*akis*, 'a point.')(F.) *Aigu*. A disease which, with a certain degree of severity, has a rapid progress, and short

duration, is said to be "acute."—*Oxy'nose'ma*, *Oxy'nosus*, *Oxynu'sus*.

Diseases were formerly subdivided into *Morbi acutis'simi*, very acute, or those which last only three or four days: *M. subacutis'simi*, which continue seven days: and *M. subacu'ti*, or those which last from twenty to forty days.

The antithesis to *acute* is *chronic*. *Acute*, when applied to pain, sound, cries, &c., means *sharp*.

ACUTENACULUM, *Porte-aiguille*.

ACYANOBLEP'SIA, from *a*, privative, *κυανος*, 'blue,' and *βλεπω*, 'I see.' Defective vision, which consists in incapability of distinguishing blue.—Göthe. See *Achromatopsia*.

ACYESIS, Sterilitas.

ACYRUS, *Arnica montana*.

ACYSTINERVIA, Cystoparalysis.

ACYSTURONERVIA, Cystoparalysis.

ACYSTUROTROPHIA, Cystatrophia.

ACYTERIUS, Abortive.

ADACA. The *Sphæranthus Ind'icus*, a Malabar plant, which is acrid and aromatic.

ADAC'RYA, from *a*, privative, and *δακρυω*, 'I weep.' Defective secretion of tears.

ADEMONIA, Anxiety.

ADAKO'DIEN. A Malabar plant of the family *Apocynæ*, used in that country in diseases of the eyes.

AD'ALI, *Lip'pia*. A Malabar plant, which the Orientals regard as an antidote to the bite of the *naja*.

ADAMANTINE SUBSTANCE, Enamel of the teeth.

ADAMAS, Diamond.

ADAMI'TA, *Adami'tum*. A very hard, white calculus.—*Paracelsus*.

The first word has been used for stone in the bladder: the second for lithiasis or the calculous condition.

ADAM'S APPLE, *Pomum Adami*.

ADANSONIA DIGITATA, Baobab.

ADAPTER, from *ad* and *apto*, 'I fit.' A tube employed in pharmaceutical operations for lengthening the neck of a retort; or in cases where the opening of the receiver is not large enough to admit the beak of the retort.

ADAR'CE, *Adar'cion*, *Adar'cis*. A concretion found about the reeds and grass in the marshy regions of Galatia, and hiding them, as it were: hence the name, from a privative, and *δεκω*, 'I see.' It was formerly in repute for cleansing the skin from freckles, &c.

ADARIGO, Orpiment.

ADARNECH, Orpiment.

ADARTICULATIO, Arthrodia.

AD CUTEM ABDOMINIS (ARTERIA). The *superficial artery of the abdomen*,—a branch of the crural or femoral, which arises at the lower part of Poupart's ligament, and ascends towards the umbilicus, being distributed to the integuments.

ADDAD. A Numidian plant; bitter and poisonous.

ADDEPHAG'IA, *Adephag'ia*, from *αδδν*, 'much,' and *φάγειν*, 'to eat.' *Voraciousness*. Galen and Hoffmann have given this name to voracious appetite in children affected with worms. Sauvages refers it to *Bulimia*. Also, the goddess of gluttony.

ADDER'S TONGUE, *Ophioglossum vulgatum*.

ADDISON, DISEASE OF, see *Capsule*, renal.

ADDITAMEN'TUM. A term once used synonymously with *Epiphysis*. It is now restricted to the prolongation of two cranial sutures, the lambdoidal and squamous.

ADDITAMEN'TUM COLI, Appendix vermiformis *cæci*—a. *Necatum*, *Olecranon*—a. ad *Sacro-lumbalem*, see *Sacro-lumbalis*—a. *Uncatum* *ulnæ*, *Olecranon*—a. *Ulnæ*, *Radius*.

ADDUCENS OCULI, *Rectus internus oculi*.

ADDUCTEUR DE L'ŒIL, *Rectus internus oculi*—*a. du Gros orteil*, *Adductor pollicis pedis*—*a. Premier ou moyen*, *Adductor longus femoris*—*a. du Pouce*, *Adductor pollicis manûs*—*a. Second ou petit*, *Adductor brevis*—*a. Troisième ou grand*, *Adductor magnus*.

ADUCTION, *Adductio*, from *ad*, 'to,' and *ducere*, 'to draw.' *Paragôgê*. The action by which parts are drawn towards the axis of the body.

The muscles which execute this function are called *Adductors*.

ADDUCTOR MEDII DIGITI PEDIS, *Posterior medii digiti pedis*—*a. Oculi*, *Rectus internus oculi*.

ADDUCTOR METACARPÏ MINIMI DIG'ITI, *Metacarpæus*, *Car'po-metacarpæus minimi dig'iti*, is situate between the adductor and flexor, next to the metacarpal bone. It arises, fleshy, from the unciform process of the os unciforme, and from the contiguous part of the annular ligament of the wrist, and is inserted, tendinous and fleshy, into the fore-part of the metacarpal bone of the little finger, from its base to its head.

ADDUCTOR POL'ICIS MANÛS, *A. Pollicis*, *A. ad minimum dig'itum*, *Metacarpophalangæus pollicis*—(Ch.) (F.) *Adducteur du pouce*. A muscle which arises, fleshy, from almost the whole length of the metacarpal bone of the middle finger, and is inserted into the inner part of the root of the first bone of the thumb.

ADDUCTOR POL'ICIS PEDIS, *Antihænar*, *Metatarsophalangæus pollicis*—(Ch.) *Tarso-metatarsi-phalangium du pouce*. (F.) *Adducteur du gros orteil*. Arises by a long, thin tendon, from the under part of the os calcis, from the os cuboides, os cuneiforme externum, and from the root of the metatarsal bone of the second toe. It is divided into two fleshy portions, and is inserted into the external sesamoid bone, and root of the metatarsal bone of the great toe.

Bichat has given the general name, *Adductors*, to those of the interosseous muscles of the hand or foot, which perform the action of adduction.

ADDUCTOR TERTII DIGITI PEDIS, *Prior tertii digiti pedis*.

ADDUCTORS OF THE THIGH. These are three in number, which have, by some anatomists, been united into one muscle—the *Triceps Adductor Femoris*.

1. *Adductor longus femoris*, *Adductor femoris primus*, *Triceps minor*, *Pubio-femoralis*—(Ch.) (F.) *Premier ou moyen adducteur*. Arises by a strong tendon from the upper and fore part of the os pubis and ligament of the symphysis, at the inner side of the pectinialis. It runs downwards and outwards, and is inserted by a broad, flat tendon, into the middle of the linea aspera.

2. *Adductor brevis*, *A. femoris secundus*, *Triceps secundus*, *Sub-pubio-femoralis*—(Ch.) (F.) *Second ou petit Adducteur*. Arises tendinous from the os pubis, at the side of its symphysis, below and behind the last muscle. It runs obliquely outwards, and is inserted by a short, flat tendon into the inner and upper part of the linea aspera, from a little below the trochanter minor to the beginning of the insertion of the adductor longus.

3. *Adductor magnus*, *Adductor femoris tertius et quartus*, *Triceps magnus*, *Ischio-femoralis*—(Ch.) (F.) *Troisième ou grand adducteur*, is much larger than either of the others. It arises from the ramus of the pubis, from that of the ischium, and from the tuber ischii, and is inserted into the whole length of the linea aspera. Near the lower part of the linea aspera it is pierced by a kind of oblique, fibrous canal, through which the crural artery and vein pass.

ADEC. The inner man.—Paracelsus.

ADECTA, Sedatives.

ADELIPARIA, Polysarcia.

ADELODAGAM. A bitter Malabar plant, used in asthma, catarrh, and gout.

ADELPHIA, see *Adelphixia*.

ADELPHIX'IA, *Adelphix'is*; from *adelphos*, 'brother.' Consanguinity of parts in health or disease; *Frater'nitas*, *Fratrat'io*. Hippocrates used the word *Adel'phia* for diseases that resemble each other.

ADELPHIXIS, Sympathy.

ADEMONIA, Depression, Nostalgia.

ADEMOSYNE, Depression, Nostalgia.

ADEN, *adên*, 'a gland;' hence *Adenalgia*, *Adeniform*, &c.—see *Gland*.

ADENAL'GIA, *Adenodynia*, from *adên*, 'a gland,' and *algos*, 'pain.' Glandular pain.

ADENECTOP'IA, from *adên*, 'a gland,' and *εκτοπος*, 'removed from its place.' Dislocation of a gland.

ADENEMPHRAX'IS, from *adên*, 'a gland,' and *εμψραξις*, 'obstruction.' Glandular obstruction.

ADÉNIES, *Adénopathies angibromiques*. Diseases of the glands connected with the digestive tube.—Piorry.

ADEN'IFORM, *Adeniform'is*, *Adenoides*, *Adenoid*, from *Aden*, 'a gland,' and *Forma*, 'form or resemblance.' *Glandiform*, or resembling a gland.

ADÉNITE, *Adenitis*—*a. Lymphatique*, *Lymphadenitis*.

ADENITIS, from *adên*, 'a gland,' and *itis*, a termination denoting inflammation. *Phlegma'sia adeno'sa* seu *glandulo'sa*. (F.) *Adénite*. Glandular inflammation.

ADENITIS LYMPHATICA, *Lymphadenitis*.

ADENITIS MESENTERICA, *Mesenteric Ganglion'itis*. Inflammation of the mesenteric glands.

ADENITIS PALPEBRARUM CONTAGIOSA, see *Ophthalmia*.

ADENOCE'LE, from *adên*, 'a gland,' and *κηλη*, 'a rupture,' 'a tumour.' Under this term Mr. Birkett includes tumours of the mammae, which contain structures similar to, if not identical with, the normal secreting tissue of the gland, and more especially the cysto-sarcoma, and chronic mammary tumours of surgical authors.

ADENOCHIRAPSOLOG'IA, from *adên*, 'a gland,' *χειρ*, 'the hand,' *απτο*, 'I lay hold of,' and *λογος*, 'a description.' The doctrine of curing scrofula or the king's evil by the royal touch.

ADENOCHON'DRIUS, from *adên*, 'a gland,' and *χονδρος*, 'a cartilage.' Relating to gland and cartilage,—for example, *Arthrophyma adeno-chon'drium*, a tumefaction of the glands and cartilages of joints.

ADÉNODERMIES SYPHILITTIQUES, *Asymphysiques*, from *adên*, 'a gland,' and *δερμα*, 'skin.' Syphilitic affections of the glands of the skin.

ADENODYNIA, *Adenalgia*.

ADENOG'RAPHY, *Adenog'raphia*, from *adên*, 'a gland,' and *γραφω*, 'I describe.' That part of anatomy which describes the glands.

ADENOID, *Adeniform*.

ADENOIDES, *Adeniform*.

ADENOL'OGY, *Adenolog'ia*, from *adên*, 'a gland,' and *λογος*, 'a description.' A treatise on the glands.

ADENOMALA'CIA, from *adên*, 'a gland,' and *μαλακία*, 'softening.' Mollescence or softening of a gland.

ADENO-MENINGEAL, see *Fever, adenomeningeal*.

ADENONCOSIS, *Adenophyma*.

ADENONCUS, Adenophyma.

ADENO-NERVOUS FEVER, Plague.

ADĒNOPATHIES ANGI-BROMIQUES, Adénies.

ADĒNO-PHARYNGEUS, from *adn̄n*, 'a gland,' and *φάρυγξ*, *φάρυγος*, 'the pharynx.' Some fleshy fibres, which pass from the constrictor pharyngis inferior to the thyroid gland, have received this name. Their existence is not constant.

ADĒNO-PHARYNGITIS. Same etymon. Inflammation of the tonsils and pharynx.

ADENOPHTHALMIA, Ophthalmia tarsi.

ADENOPHY'MA, *Adenoncus*, *Adenoncosis*, from *adn̄n*, 'a gland,' and *φύμα*, 'a swelling,' Swelling of a gland, or glandiform ganglion. (F.) *Glandage*. *Adenophyma* is used by some to signify a soft glandular swelling;—*Adenoncus*, one of a harder character.—Kraus.

ADENOPHYMA INGUINALIS. Bubo.

ADENOSCIRRHUS, *Adenosclerosis*, from *adn̄n*, 'a gland,' and *σκίρρως*, 'induration.' Scirrhous induration of a gland.

ADENOSCLEROSIS, Adenoseirrhus.

ADENOSIS SCROPHULOSA, Scrofula.

ADENO'SUS, (*Abcesus*). A hard, glandular abscess, which suppurates slowly.—M. A. Severinus.

ADENOSYNCHITONITIS, Ophthalmia Neotorum.

ADENOT'OMY, *Adenotomia*, from *adn̄n*, 'a gland,' and *τομή*, 'I cut.' Dissection of the glands.

ADEPHAGIA, Addephagia, Boulimia.

ADEPS, genitive *Adipis*, *Adeps Sullus*, *Oryngium*, *Pinguedo*, *Lardidum*, *Lardum*, *Lard*. (Prov.) *Mort*, *Sallie*, *Swinesame*. Pig's flare. The fat of the hog. In the Ph. U. S., the prepared fat of *Sus scrofa*, free from saline matter.

ADEPS ANSERINUS, *Adeps anseris* or *Goose grease*, (F.) *Graisse d'Oie*, is emollient. It has been used as an emetic.

ADEPS CANTHARIDIBUS MEDICATUS, Unguentum lyttæ medicatum—a. Cortice Daphnes gnidii medicatum, Unguentum epispasticum de Daphne gnidio—a. Humanum, Liqueumunia—a. Hydrargyro medicatus, Unguentum Hydrargyri—a. ex Hydrargyro mitius dictum cinereum, Unguentum oxidi hydrargyri cinereum—a. Hydrargyri muriate oxygenato medicatus, Unguentum muratis hydrargyri oxygenati medicatum—a. Hydrargyri nitrate medicatus, Unguentum hydrargyri nitratis—a. Hydrargyri oxido rubro et plumbi acetate medicatus, Unguentum ophthalmicum—a. Lauro medicatus, Unguentum laurinum—a. Myristicæ, see Myristica—a. Ovilli, Seum—a. Papavere, hyoscyamo, et belladonnâ medicatus, Unguentum populeum.

ADEPS PRÆPARATUS, A. *Sullus Præparatus*, *Hog's lard*, *Barrow's grease*, *Lard*, *Ac'unge*, *Axungia*, *Axungia porcina*, (F.) *Graisse de Porc*, *Saindoux*, *Azonge*, is prepared by melting pig's flare, and straining it. This is called *rendering* the lard. Lard is emollient, but is chiefly used for forming ointments and plasters.

ADEPS SULLUS PRÆPARATUS, *Adeps Præparatus*—a. Sulfure et ammoniæ muriate medicatus, Unguentum sulphuratum ad scabiem—a. Sulfure et carbonate potassæ medicatus, Unguentum sulphuratum alcalinum ad scabiem—a. Tartaro stibii medicatus, Unguentum antimonii tartarizati—a. Oxido zinci medicatus, Unguentum oxidi zinci impuri.

ADEPT, Alehymist.

ADEPTA MEDICINA. Medicine, which treated of diseases contracted by celestial operations, or communicated from heaven.

ADEPTA PHILOSOPHIA, Alchymy.

ADERMONERVIA, Dermataneuria.

ADERMOTROPH'IA, from *a*, priv., *δέρμα*, 'skin,' and *τροφή*, 'nourishment.' Extenuation or atrophy of the skin.

ADFLATUS, Afflatus.

ADHERENT'IA, Adherence.

ADHESIO, Adherence.

ADHATO'DA, *Justicia adhato'da*. The *Malabar Nut Tree*. (F.) *Noyer de Ceylan*. Used in India for expelling the dead fetus in abortion. The word is said to convey this meaning in the Ceylones.

ADHĒRENCE, *Adhesion*, *Adherentia*, *Concretio*, *Atrésia*, *Prosp'physis*, *Proscollésis*, *Adhésio*, from *adhærere*, (*ad* and *hærere*,) 'to stick to.' These words are usually employed synonymously. The French often use *adherence* for the state of union, and *adhesion* for the act of adhering.

ADHESION, Adherence.

ADHĒSIVE INFLAMMA'TION is that inflammation which terminates by an adhesion between inflamed and separated surfaces, and which was, at one time, supposed to be necessary for such adhesion.

Adhesive is also an epithet for certain plasters which stick closely to the skin.

ADIANTHUM, Adiantum.

ADIANTUM, A. pedatum.

ADIANTUM ÆTHIOP'ICUM. A South African plant, *Ord. Filices*, an infusion of which is sometimes used as an emollient in coughs, and in diseases of the chest.

ADIANTUM ALBUM, *Asplenium ruta muraria*—a. *Aureum*, *Polytrichum*.

ADIANTUM CAPILLUS VEN'ERIS, A. *Coriandrifolium* seu *Nigrum*, *Capillus Ven'eri*, from *a*, privative, and *δαίρω*, 'to grow wet,' from the leaves not being easily moistened. *Maiden hair*. (F.) *Capillaire de Montpellier*. A European plant, of feeble, aromatic and demulcent properties. It is used for forming the *Sirap de Capillaire* or *Capillaire*.

ADIANTUM CORIANDRIFOLIUM, A. *Capillus Veneris*—a. *Nigrum*, A. *Capillus Veneris*.

ADIANTUM PEDA'TUM, A. *Canadense* seu *Patens*, *Adiantum*, *Capillus Veneris Canadensis*, *Herba Veneris*, *Filix Veneris*, *Canada Maiden hair*, *American Maiden hair*, *Rockfern*, *Sweetfern*, (F.) *Capillaire du Canada*, has the same properties. *Capillaire* was once made from this. See *Adiantum*.

ADIANTUM RUBRUM, *Asplenium trichomanoides*.

ADIAPHANIE PÉRICHRYSAITIQUE, see Cataract.

ADIAPHORO'SIS, *Adiaphoresis*, from *a*, privative, *ἔω*, 'through,' and *πόρος*, 'a pore.' Defect or suppression of perspiration, *Adiapneustia*.

ADIAPH'OROUS, *Adiaph'orus*, *Indifferens*, *Neutral*. A medicine which will neither do harm nor good.

ADIAPNEUSTIA, *Adiaphorosis*.

ADIARRHŒ'A, from *a*, privative, and *διάρρην*, 'to flow.' Retention of any excretion.—Hippocrates.

ADIATH'ESIC, *Adiathesicus*, from *a*, privative, and *ἰσθῆσις*, 'diathesis.' In the system of controstimulism, *adiathesic diseases* are such as occur without any antecedent diathesis.

ADICE, Urtica.

ADIPATUS, Fatty.

ADIPEUX, Adipose.

ADIPOCERA, *Adipocire*—a. *Cetosa*, *Ceteum*.

ADIPOCIRE, *Adipocera*, from *adeps*, 'fat,' and *cera*, 'wax.' The base of biliary calculi, called also *Chol'esterine*. Also, a sort of soap, formed from animal matter under certain circum-

stances. (F.) *Gras des Cadavres, Gras des Cimetières*. The human body, when it has been for some weeks in water, assumes this appearance; and it has been a subject of legal inquiry, what length of time is necessary to produce it. This must, of course, depend upon various circumstances, as climate, season, &c.

ADIPOCIRE DE BALEINE, Cetaceum.

ADIPOSE, *Ad'ipous*, from *adeps*, 'fat.' (F.) *Adipeux*. That which relates to fat—

as *Adipose membrane*, *A. vessels*, &c. See *Fatty*.
ADIPOSE SARCO'MA of *AB'ERNETHY*, *Emphy'ma sarco'ma adiposum*, is suetty throughout, and enclosed in a thin capsule of condensed areolar substance, connected by means of minute vessels. It is chiefly found on the fore and back parts of the trunk. See *Sarcoma*.

ADIPOSIS. See *Polysarcia*.

ADIPO'SIS HEPAT'ICA, *Pineol'sis seu steato'sis seu malar'sis hepatic'a*, *Hepar adiposum*, *Fatty liver*, *Fatty degeneration of the liver*, (F.) *Dégénérescence graisseuse du Foie*. Fatty disease of the liver.

ADIPOSUS, Fatty.

ADIPOUS, Fatty.

ADIP'SIA, *Dipso'sis exsers*. Absence of thirst.

ADIP'SON, *Adip'sum*, from *a*, privative, and *δρῖν*, 'thirst.' Any substance which relieves thirst. Applied to a decoction of barley to which oxymel was added.—*Hippocrates*.

ADIPSOS, *Glycyrhiza*.

ADITUS, 'an entrance,' 'an approach,' from *adere*, *aditum*, 'to go to.' *Pros'odos*. The entrance to a canal or duct, as *Aditus ad Aquæductum Fallopii*.

ADITUS AD INFUNDIBULUM, Vulva.

ADIULIS'TOS, from *a*, privative, and *δινλίζω*, 'I strain.' Unstrained wine for pharmaceutical purposes.—*Gorreaus*.

ADJUNCTUM, Accompaniment.

ADJUTOR PARTÛS, Accoucheur.

ADJUVANT, *Ad'juvans*, from *adjuvare*, 'to aid.' A medicine, introduced into a prescription to aid the operation of the principal ingredient or basis. Also, whatever assists in the removal or prevention of disease.

ADJINICULE, *Adminic'ulum*, (*ad* and *manus*, 'a hand'). A prop, a support. Anything that aids the action of a remedy.

ADNASCENTIA, *Prospheysis*.

ADNATA (TUNICA,) *Conjunctiva*.

ADNÉE (MEMBRANE,) *Conjunctiva*.

ADOLESC'ENCE, *Adolescen'tia*, *Juven'ta*, *Juven'tas*, *Juven'tus*, *Ætas bona*, *Youth*; from *adolescere*, (*ad* and *ollescere*,) 'to grow.' (F.) *Jeunesse*. The period between puberty and that at which the body acquires its full development; being, in man, between the 14th and 25th years; and, in woman, between the 12th and 21st.

ADOLESC'ENS, *Ju'venis*, *Hebe'tes*, *Hebe'tor*, *Hypene'tes*. A youth. A young man in the period of adolescence.

ADO'LIA. A Malabar plant, whose leaves, put in oil, form a liniment, used in facilitating labour.

ADOR, *Zea mays*.

ADORION, *Daucus carota*.

ADOUC'SSANT, *Demulcent*.

AD PONDUS OM'NIUM. The weight of the whole. In a prescription it means, that any particular ingredient shall equal in weight the whole of the others.

ADRAGANT, *Tragacantha*.

ADRAGANTHE, *Tragacantha*.

ADRA RIZA, *Aristolochia clematitis*.

ADROBO'LON, from *adros*, 'great,' and *βῶλος*, 'mass.' The bdellium of India, which is in larger pieces than that of Arabia.

ADROS, *adros*, 'plump and full.' Applied to the habit of body, and also to the pulse.—*Hippocrates*.

ADSARIA PALA, *Dolichos pruriens*.

ADSPIRATIO, *Aspiration*, *Inspiration*.

ADSTANS, *Prostate*.

ADSTITES GLANDULOSI, *Prostate*.

ADSTRICTIO, *Astriction*, *Constipation*.

ADSTRICTORIA, *Astringents*.

ADSTRINGENTIA, *Astringents*.

ADULAS'SO. The *Justitia bivalvis*. A small shrub, used in India as a local application in gout.

ADULT, see *Adult age*.

ADULT AGE, *Andri'a*, from *adolescere*, 'to grow to,' (*ad* and *olere*, *olitum*, 'to grow.') *Virility*, *manhood*. The age succeeding adolescence, and preceding old age. In the civil law, an adult is one, who, if a boy, has attained the age of fourteen years; and, if a girl, of twelve. In the common law, one of full age. *Adult*, *Adultus*, is also used for one in the adult age.

ADULTERATIO, *Falsification*.

ADULTUS, see *Adult age*.

ADUNCATIO UNGUIUM, *Onychogryphosis*.

ADURENS, *Caustic*.

ADURION, *Rhus coriaria*.

ADUST, *Adus'tus*, from *adurere*, (*ad* and *urere*,) 'to burn.' The blood and fluids were formerly said to be adust, when there was much heat in the constitution, and but little serum in the blood.

ADUSTIO, *Adustion*, *Burn*.

ADUS'TION, *Adus'tio*. State of the body described under *Adust*. In surgery, it signifies *cauterization*.

ADVENTITIOUS DISEASES, *Acquired diseases*.

ADVENTITIUS, *Accidental*.

ADYNA'MIA, *Impotentia*; from *a*, privative, and *δυναμις*, 'strength,' *Adyna'sia*, *Adyna'tia*. Considerable debility of the vital powers; as in typhus fever. Some Nosologists have a class of diseases under the name *Adynamia*, *Ectlyses*, *Morbi asthen'ici*.

ADYNAMIA VIRILIS, *Impotence*.

ADYNAM'IC, *Adynam'icus*, *Hypodynam'ic*, *Hypodynam'icus*; same etymon. Appertaining to debility of the vital powers.

ADYNASIA, *Adynamia*.

ADYNATIA, *Adynamia*.

ADYNATOCOMIUM, *Hospital*.

ADYNATODOCHIUM, *Hospital*.

ADYNATOS, *Sickly*.

ÆDEA, *Genital Organs*.

ÆDÆAGRA, from *aidia*, 'genital organs,' and *αγρα*, 'seizure.' Gout in the genitals.

ÆDÆAG'RAPHY, *Ædæagraph'ia*, *Ædæog'raphy*, from *aidia*, 'organs of generation,' and *γραφω*, 'I describe.' A description of the organs of generation.

ÆDÆAL'OGY, *Ædæalog'ia*, *Ædæol'ogy*, from *aidia*, 'organs of generation,' and *λογος*, 'a description.' A treatise on the organs of generation.

ÆDÆAT'OMY, *Ædæatom'ia*, *Ædæotom'ia*, *Ædæot'omē*, *Ædæot'omy*, from *aidia*, 'organs of generation,' and *τεμνω*, 'I cut.' Dissection of the parts of generation.

ÆDÆI'TIS, *Ædæoti'tis*, *Mede'i'tis*; from *aidia*, 'organs of generation,' and *itis*, denoting inflammation. Inflammation of the genital organs.

ÆDÆOBLENORR'HÆA, *Leucorrhœa*.

ÆDÆODYN'IA, from *aidia*, 'organs of generation,' and *δύω*, 'pain.' Pain in the genitals. *Pudendagra*.

ÆDÆOGARGALUS, *Masturbation*, *Nymphomania*.

ÆDÆOGARGARISMUS, *Masturbation*, *Nymphomania*.

ÆDEOGRAPHY, *Ædœography*.
ÆDEOLOGY, *Ædœology*.
ÆDEOMANIA, *Nymphomania*.
ÆDEOMYCODERMITIS, from *αἰδοῖα*, 'organs of generation,' *μυκός*, 'mucus,' and *δέρμα*, 'skin.' Inflammation of the mucous membrane of the genital organs.
ÆDEON, *Inguen*.
ÆDEOPSOPHESIS, *Ædeopsophia*.
ÆDEOPSOPHIA, *Ædeopsophēsis*, from *αἰδοῖα*, 'organs of generation,' and *ὥσπερ*, 'to make a noise.' Emission of wind by the urethra in man, by the vagina in woman.—Sauvages and Sagar.
ÆDEOPSOPHIA UTERINA, *Physometra*.
ÆDEOSCOPY, *Ædœoscopy*, from *αἰδοῖα*, 'organs of generation,' and *σκοπεῖν*, 'to view.' Exploration of the organs of generation.
ÆDEOTITIS, *Ædœitis*—æ. *Gangrænosa*, *Colpocæ*—æ. *Gangrænosa puellarum*, *Colpocæ infantilis*—æ. *Gangrænosa puerperarum*, *Colpocæ puerperarum*.
ÆDEOTOME, *Ædœatomy*.
ÆDEOTOMIA, *Ædœatomy*.
ÆDEOTOMY, *Ædœatomy*.
ÆDOPTOSIS, *Hysteroptosis*—æ. *Uteri*, *Prolapsus uteri*—æ. *Uteri inversa*, *Uterus*, *inversion of the*—æ. *Uteri retroversa*, *Retroversio uteri*—æ. *Vaginæ*, *Prolapsus V.*—æ. *Vesicæ*, *Exocyste*.
ÆEIG'LUCES, *Æeig'luces*, from *αἰ*, 'always,' and *γλυκός*, 'sweet.' A kind of sweet wine or must.—Gorræus.
ÆGAGROPILA, *Ægagropilus*, from *αἰγας*, 'the rock goat,' and *πίλος*, 'hair,' *Bézoar d'Allemagne*, *Pila Dama'rum* seu *Rupicapra'rum*. A ball composed of hairs, found in the stomach of the goat: once used medicinally.
ÆGEIROS, *Populus*.
ÆGER, *Sick*.
ÆGIAS, *Ægis*, *Ægilia*, *Ægides*, from *αἰ*, *αἰγός*, 'the goat,' *why*, is not known. (F.) *Aige* or *Aigle*. There is obscurity regarding the precise meaning of this word. It was used to designate an ulcer, or speck on the transparent cornea.—Hippocrates.
Ægla, *Mafre Jean* uses it for a calcareous deposit between the conjunctiva and sclerótica.
ÆGIDES, *Ægias*.
ÆGILOPS, *An'chilops*, *An'kylops*, from *αἰ*, *αἰγός*, 'goat,' and *ὤψ*, 'the eye.' An ulcer at the greater angle of the eye, which sometimes does not penetrate to the lachrymal sac, but at others does, and constitutes fistula lachrymalis.—Galen, Celsus, Oribasius, Aëtius, Paulus of Ægina, &c.
ÆGTRINON. An ointment of which the fruit or flower of the poplar was an ingredient; from *αἰγείοος*, 'the black poplar.'
ÆGLA, *Ægias*.
ÆGOCERAS, *Trigonella fœnum*.
ÆGOLETHRON, *Ranunculus flammula*.
ÆGONYCHON, *Lithospermum officinale*.
ÆGOPHONIA, *Egophony*.
ÆGOPHONICUS, *Egophonic*.
ÆGOPIDIUM, *PODAGRARIA*, *Ligusticum podagraria*.
ÆGRIPPA, *Agrippa*.
ÆGRITUDO, Disease—æ. *Ventriculi*, *Vomiting*.
ÆGROTATIO, Disease.
ÆGRÛTUS, Sick.
ÆGYPTIA. An epithet for several medicines, mentioned by Galen, Paulus of Ægina, and Myrseus.
ÆGYPTIA MOSCHATA, *Hibiscus abelmoschus*.
ÆGYPTIA STYPTERIA, *Αἰγυπτια στυπτηρια*, *Ægyptian alum*. Recommended by Hippocrates.
ÆGYPTIA ULCERA; *Ægyptian ulcers*. Ulcers of the fauces and tonsils, described by Aretæus, as common in Egypt and Syria.

ÆGYPTIACUM, *Ægyption*, *Mende'sion*, *Mel Ægyptiacum*, *Phar'macum Ægyptiacum*. A preparation of vinegar, honey, and verdigris, scarcely used now, except by veterinary surgeons as a detergent. See *Linimentum Ærginis*.
ÆGYPTION, *Ægyptiacum*.
ÆGYPTIUM MEDICAMENTUM AD AURES, *Pharmacum ad aures*.
ÆGYPTIUS PESSUS; *Ægyptian pessary*. A pessary, composed of honey, turpentine, butter, oil of lily or of rose, saffron, each one part; with sometimes a small quantity of verdigris.
ÆICHRYSON, *Sedum*.
ÆIPATHEIA, see *Continent (Disease)*.
ÆIPATHIA, see *Continent (Disease)*.
ÆMOPTOICA PASSIO, *Hæmoptysis*.
ÆNEA, *Catheter*.
ÆOLECTHYMA, *Variola*.
ÆOLLION, *Varicella*.
ÆOLLIUM, *Varicella*.
ÆON, *atav*. The entire age of a man from birth till death.—Hippocrates, Galen. Also, the spinal marrow. See *Medulla Spinalis*.
ÆONESIS, *Fomentation*.
ÆONION, *Sedum*.
ÆO'RA, from *αἰωρεῖα*, 'I suspend.' *Gestation*, *swinging*.—Aëtius, Celsus, &c.
ÆQUALIS, *Equal*.
ÆQUATOR OCULI. The line formed by the union of the upper and under eyelid, when they are closed. It is below the middle of the globe.
ÆQUIVOCUS, *Equivocal*.
ÆR, *Air*.
ÆRATION OF THE BLOOD, *Hæmatosis*.
ÆRATUS, *Carbonated*.
ÆERENDOCARDIA, (F.) *Æerendocardie*, from *anp*, 'air,' and *endocardium*. Presence of air in the endocardium or lining membrane of the heart.—Piorry.
ÆERË, *Carbonated*.
ÆERENTERECTASIA, *Tympanites*.
ÆEROLUM, *Æreolus*, *Chalcus*. The sixth part of an obolus by weight, consequently about 2 grains.
ÆERESIS, *αἰρεσις*, 'the removal of any thing.' A suffix denoting a removal or separation, as *Aphæresis*, *Diæresis*, &c.
ÆERETHMIE, *Emphysema*.
ÆERETHMOPNEUMONIE, see *Emphysema of the Lungs*.
ÆERGIA, *Torpor*.
ÆERHÆMATOXIA, (F.) *Æerhæmatoxic*; from *anp*, 'air,' *αἷμα*, 'blood,' and *τοξικόν*, 'poison.' Poisoning by the reception of air into the blood-vessels.
ÆERIFEROUS, *Æerifer*, (F.) *Æerifère*, from *aer*, 'air,' and *ferre*, 'to carry.' An epithet for tubes which convey air, as the larynx, trachea, and bronchia.
ÆERIFLUX'US. The discharge of gas, and the fetid emanations from the sick. *Flatulence*.—Sauvages.
ÆERODERMECTASIA, *Emphysema*.
ÆERODIAPH'THORA, from *anp*, 'air,' and *διάθροα*, 'corruption.' A corrupt state of the air.
ÆERO-ENTERECTASIA, *Tympanites*.
ÆEROHYDROPATHY. *Æerohydropathia*, from *anp*, 'air,' *ὕδωρ*, 'water,' and *πάθος*, 'affection.' A mode of treating disease, in which air and water are the chief agents employed.
ÆEROL'OGY; *Æerolog'ia*, *Æerolog'icē*, from *anp*, 'air,' and *λογος*, 'a description.' That part of physics which treats of the air, its qualities, uses, and action on the animal economy.
ÆEROMANCY, *Æeromanti'a*, from *anp*, 'air,' and *μαντεία*, 'divination.' An art in judicial as-

trology, which consists in the foretelling, by means of the air, or substances found in the atmosphere.

AEROMELI, *Fraxinus ornus*.

AEROPÉRITONIE, see Tympanites.

AEROPHOBIA, from *anp*, 'air,' and *φοβος*, 'fear.' Dread of the air. This symptom often accompanies hydrophobia, and sometimes hysteria and other affections.

AEROPHOBICUS, *Aëroph'obus*; same etymon. One affected with aerophobia.

AEROPHOBUS, Aerophobicus.

AEROPHTHORA, Aerodiaphthora.

AÉROPLEURIE, Pneumothorax.

AÉROPNEUMONASIE, see Emphysema of the Lungs.

AÉROPNEUMONECTASIE, see Emphysema of the Lungs.

AÉRORA'CHIA, from *anp*, 'air,' and *παχis*, 'spine.' Accumulation of air in the spine.

AEROSIS, Pneumatosiis, Tympanites.

AÉROTHORAX, Pneumothorax.

ÆRUCA, *Cupri subacetos*.

ÆRUGINOUS, *Ærugine'us*, *Io'des*, from *ærugo*, 'verdigris.' (F.) *Ærugineus*. Resembling verdigris in colour; as the bile when discharged at times from the stomach.

ÆRUGO, *ios*, from *æs*, 'copper.' The rust of any metal, properly of brass. See *Cupri Subacetos*.

ÆRUGO FERRI, Ferri subcarbonas—æ. Plumbi, Plumbi subcarbonas.

ÆS, *Cuprum*.

ÆSCHOS, *αισχος*. Deformity of the body generally, or of some part.—Hippocrates.

ÆSCULUS HIPPOCASTANUM, from *esca*, 'food,' [?] *Casta'nea equi'na*, *Pav'ina*, *Horse-chestnut*, *Buck-eye*. (F.) *Marronnier* ou *Marronnier d'Inde*. Ord. Hippocastaneæ. Sex. Syst. Heptandria Monogynia. The bark has been advised as a substitute for cinchona. Both bark and fruit are astringent. Externally, it has been employed, like cinchona, in gangrene.

ÆSECAVUM, Brass.

ÆSTATES, Ephelides.

ÆSTHE'MA, *αισθημα*, *αισθηματος*, 'a sensation, a perception.' See Sensation and Sensibility. In the plural, *æsthe'mata*, the apparatuses of the senses.

ÆSTHEMATOLOGY, *Æsthematolog'ia*; from *αισθημα*, *αισθηματος* and *λογος*, 'a discourse.' The doctrine of, or a treatise on, the senses, or on the apparatus of the senses.

ÆSTHEMATONU'SI, *Æsthematorganonu'si*, from *αισθημα*, *αισθηματος*, and *νοσος*, 'diseases.' Diseases affecting sensation.

ÆSTHEMATORGANONUSI, *Æsthematonusi*.

ÆSTHESIOG'RAPHY, *Æsthesiogra'phia*, from *αισθησις*, 'perception, sensation,' and *γραφη*, 'a description.' A description of the sensorial apparatuses.

ÆSTHESIOLOGY, *Æsthesiolog'ia*, from *αισθησις*, 'perception, sensation,' and *λογος*, 'a discourse.' The doctrine of, or a treatise on, sensation or the senses.

ÆSTHESIS, *Aisthe'sis*, from *αισθανομαι*, 'I feel.' The faculty of being affected by a sensation. Perception. Sensibility, as well as the senses themselves. See Sense.

ÆSTHETERION, Sensorium.

ÆSTHETICA, from *αισθανομαι*, 'I feel.' Diseases affecting the sensations. Dullness, depravation or abolition of one or more of the external organs of sense. The 2d order, class *N'urotica*, of Good. Also, agents that affect sensation.—Pereira.

ÆSTIVUS, Estival.

ÆSTUARIUM, Stove.

ÆSTUATIO, Ardor, Ebullition, Fermentation.

ÆSTUS, Ardor.

ÆSTUS VOLAT'ICUS. Sudden heat, scorching or flushing of the face.—Vogel.

ÆTAS, Age—æ. Bona, Adolescence—æ. Deceprota, Deceperitudo—æ. Mala, Senectus—æ. Profecta, Senectus—æ. Senilis, Senectus.

ÆTHER, *Ether*, from *αιθρ*, 'air,' or from *αιθεω*, 'I burn.' *Naphtha*. *Liquor æthe'reus*. A volatile liquor obtained by distillation from a mixture of alcohol and a concentrated acid. See *Æther sulphuricus*, and *Ether*.

ÆTHER ACETICUS, see *Æther Sulphuricus*—æ. Chloricus, Chloroform, *Ether*, chloric.

ÆTHER HYDRID'ICUS, *Hydriod'ic* or *Iodhy'dric Ether*, *Iodide of Ethyle*. (F.) *Éther iodhydrique*: obtained by mixing alcohol, parts v.; iodine, parts x.; and phosphorus, one part, and distilling. It is used for iodine inhalation.

ÆTHER HYDROCHLORICUS; *Æther muriaticus*.

ÆTHER HYDROCYAN'ICUS, *Æ. Prus'icus*, *Hydrocyan'ic* or *Prussic Ether*, *Hydrocyan'ate of Eth'erine*, *Cyan'uret of Eth'yle*, (F.) *Éther Hydrocyanique*, has been advised in whooping-cough, and where the hydrocyanic acid is indicated. Dose, 6 drops.

ÆTHER LIGNOSUS, Acetone—æ. Martialis, *Tinctura seu Alcohol sulfurico-æthereus ferri*.

ÆTHER MURIAT'ICUS, *Æ. Hydrochlor'icus*, *Muriat'ic*, *Chlorohy'dric* or *Marine Ether*, *Muriate of Etherine*, *Chloride of Ethyle*. This ether, on account of its volatility, can only be kept in cool places. It has the properties of the other ethers, and when used is generally mixed with an equal bulk of alcohol. It has been employed as an anæsthetic.

A Chlorinated Chlorohydric Ether, (F.) *Éther Chlorhydrique chloré*, formed by the action of Chlorine on Chlorohydric Ether, has been introduced into practice as a local anæsthetic.

ÆTHER MURIATICUS ALCOHOLICUS, *Spiritus ætheris muriaticus*—æ. Nitricus, see *Æther Sulphuricus*—æ. Nitricus Alcolisatus, *Spiritus ætheris nitrici*—æ. Pyro-aceticus, Acetone—æ. Nitrosus, see *Æther sulphuricus*.

ÆTHER SULPH'RICUS, *Æther* (Ph. U. S.), *Æ. Vitriol'icus*, *Naphtha Vitrioli seu Sulph'rica*, *O'leum vitriolæ dulcè*, *Sulph'uric Ether*, (F.) *Éther*, *Éther sulfurique*, *vitriolique* ou *hydrati-que*. Ether prepared from sulphuric acid and alcohol.

Rectified Ether, *Æther rectific'us*, prepared by distilling 12 oz. from a mixture of sulphuric ether, f^{ss}xiv, fused potass, f^{ss}. and distilled water, f^{ss}ij, is a limpid, colourless, very inflammable, volatile liquor; of a penetrating and fragrant odour, and hot pungent taste. Its s. g. is 0.732.

Æther Sulphuricus, *Sulphuric Ether* of the Pharmacopœia of the United States (1842), *Æther* of that of 1851, is formed from alcohol, Oiv; sulphuric acid, Oj; potassa, f^{ss}xj; distilled water, f^{ss}ij; distilling and redistilling according to the process there laid down. The specific gravity of this ether is 0.750.

It is a diffusible stimulant, narcotic and antispasmodic, and is externally refrigerant. Dose, grt. xxx to f^{ss}ss. When ether is inhaled, it is found to be a valuable anæsthetic agent; and is employed with advantage in spasmodic affections, and in surgical operations. See Anæsthetic.

The Parisian Codex has an *Æther aceticus*, *Naphtha Aceti*, (F.) *Éther actique*; an *Æther muriat'icus seu hydrochlor'icus*, (F.) *Éther chlorhydrique*; an *Æther nitricus seu nitrosus*, (F.)

Æther nitrique ou *nitreux* ou *azoteux*; and an *Æther phosphoratus*. They all possess similar virtues. See *Anæsthetic*.

Æther Sulphuricus Acidus, *Elixir acidum Halleri*—æ. *Sulphuricus* cum *alcohol*, *Spiritus ætheris sulphurici*—æ. *Sulphuricus* cum *alcohol* aromaticus, *Spiritus ætheris aromaticus*.

Æther Terebinthinatus, *Terebinthinated ether*, made by mixing gradually two pounds of *alcohol*, and half a pound of *spirit of turpentine*, with two pounds of *concentrated nitric acid*, and distilling one-half the mixture with a gentle heat. Employed externally and internally in biliary calculi, rheumatism, &c. Dose 20 to 40 drops, in honey or yolk of egg.

ÆTHEREA HERBA, *Eryngium maritimum*.

ÆTHEREAL, *Æthereus*, *Ætheral*, *Æthereous*, (F.) *Éthérée*. An ethereal tincture, (F.) *Tincture éthérée*, is one formed by the action of sulphuric ether, at the ordinary temperature, on medicinal substances. An ethereal oil is a volatile oil. See *Olea Volatilia*.

ÆTHERISMUS, *Etherism*.

ÆTHERISATIO, *Etherization*.

ÆTHERISATUS, *Etherized*.

ÆTHEREO-OLEOSA (Remedia), from *Æthereoleum*, 'a volatile oil.' Remedies, whose properties are dependent upon the volatile oil they contain.

ÆTHEROLEA, *Olea volatilia*.

ÆTHIOPICUS LAPIS, *Ethiopian stone*. A stone formerly supposed to be possessed of considerable virtue.—*Oribasius*.

ÆTHIOPIFICATIO, *Æthiopopoeisis*, *Æthiopismus*, *Æthiopiosis*, from *Æthiops*, and *facere*, 'to make.' The mummy-like colouring of the skin, induced at times by the use of mercurial ointment; and seen in bodies poisoned by arsenic.

ÆTHIOPIOSIS, *Æthiopification*.

ÆTHIOPIS, *Salvia sclarea*.

ÆTHIOPISMUS, *Æthiopification*.

ÆTHIOPOPOESIS, *Æthiopification*.

ÆTHIOPS, from *αἶω*, 'I burn,' and *ὤψ*, 'countenance.' A black or burnt countenance. The ancients gave this name to certain oxides and sulphurets of metals, which were of a black colour.

ÆTHIOPS ALBUS, *Albino*—æ. *Alcalisatus*, *Hydrargyrum cum creta*—æ. *Animal*, see *Choroid*.

ÆTHIOPS MARTIALIS, *Martial Ethiops*, *Ferri Deutoxydum nigrum*. The black deutoxide of iron: once in repute as a tonic. See *Oxydum Ferri nigrum*.

ÆTHIOPS MINERALIS, *Hydrargyri sulphuretum nigrum*—æ. *Narcoticus*, *Hydrargyri sulphuretum nigrum*—æ. *per se*, *Hydrargyri oxydum cinereum*—æ. *Saccharatus*, *Hydrargyrum saccharatum*—æ. *Vegetabilis*, see *Fucus vesiculosus*.

ÆTHOLICES, from *αἶω*, 'I burn.' Fiery pustules on the skin. Some have considered them to have been boils.

ÆTHUSA AMMI, *Sison ammi*.

ÆTHUSA CYNAPIUM, *Fool's Parsley*, (F.) *Faux Persil*, *Petite Ciguë*. Family, *Umbelliferae*. Sex. Syst. *Pentandria Digynia*. A poisonous plant, which has been mistaken for true parsley, producing nausea, vomiting, headache, giddiness, sopor, and at times, fatal results. It resembles conium in its action.

ÆTHUSA MEUM, *Meum*, *M. Athamancticum*, seu *Anthipitium*, *Athamantha Meum*, *Ligusticum Capillaceum* seu *Meum*, *Seoeli Meum*, *Meu*, *Spiguel*, *Baldmney*. (F.) *Éthuse*, *Méum*. The root has been advised as carminative, stomachic, &c.

ÆTIOLGY, *Ætiologia*, *Ætiology*, *Ætiology*, from *αἷτια*, 'cause,' and *λογος*, 'a discourse.' The doctrine of the causes of disease.

ÆTITES, or AÉTITES, from *aeros*, 'an eagle.' *Eagle-stone*, *Pierre d'Aigle*, *Hydrate de trioxide de fer*. This stone was formerly supposed to facilitate delivery, if bound on the thigh; and to prevent abortion, if bound on the arm. It was also called *Lapis Collymus*.

ÆTOI, or AËTOI PHLEBES, Temporal veins.

ÆTOLION, *Cnidia grana*.

AFFADIL, *Narcissus pseudo-narcissus*.

AFFADISSEMENT, (F.) from *fade*, 'insipid.' That condition of the digestive function in which the appetite is diminished, the sense of taste blunted, and the action of the stomach enfeebled; a state usually accompanied by general languor.

AFFABLISSEMENT, *Asthenia*.

AFFAIRES, *Menses*.

AFFAISSEMENT, *Collapsus*.

AFFECTIO, *Affection*—a. *Arthritica Cordis*, *Cardiagra*—a. *Hypochondriaca*, *Hypochondriasis*—a. *Hysterica*, *Hysteria*—a. *Sarmatica*, *Plica*—a. *Tympanica*, *Tympanites*.

AFFECTION, *Affection*, from *afficio* or *affectare* (*ad* and *facere*) 'to move or influence.' Any mode in which the mind or body is affected or modified.

AFFECTIONE TYPHOÏDE, see *Typhus*—a. *Vaporeuse*, *Hypochondriasis*.

AFFECTIONES ANIMI, *Affections of the mind*.

AFFECTIONS OF THE MIND, *Affectus seu Passiones seu Affectiones seu Congruantiones seu Confusiones seu Turbationes seu Perturbationes animi*, (F.) *Affections de l'âme* include not only the different passions, as love, hatred, jealousy, &c., but every condition of the mind that is accompanied by an agreeable or disagreeable feeling, as pleasure, fear, sorrow, &c.

In Pathology, *Affection*, *Pathos*, *Pathema*, is synonymous with disease: thus we speak of a pulmonary affection, a calculous affection, &c.

AFFECTIONS DE L'ÂME, *Affections of the mind*.

AFFECTIVE, That which affects, touches, &c. Gall gives the term *affective faculties*, (F.) *Facultés affectives*, to functions dependent upon the organization of the brain, comprising the sentiments, affections, &c.

AFFECTUS, *Passion*—a. *Faucium pestilens*, *Cynanche maligna*—a. *Hyderodes*, *Hydrops*—a. *Spasmodico-convulsivus labiorum*, *Neuralgia faciei*.

AFFERENT, *Afferens*, *Centripetal*, *Eodidic*, from *affero*, (*ad* and *fero*, 'to carry,') 'I bring.' Conveying inwards, as from the periphery to the centre. The vessels which convey the lymph or chyle to the lymphatic glands, are called *afferent*, *visæ afferentia* seu *inferentia*. Also, nerves that convey impressions towards the nervous centres—*nervi entobanontes*.

AFFION, *Affium*, *Opium*. The Bantamese thus designate an electuary of which opium is the basis, and which they use as an excitant.

AFFLATUS, *Adflatus*, *Epipnoi'a*, from *ad*, 'to,' and *flare*, 'to blow.' Any air that strikes the body and produces disease.

AFFLUENCE, *Affluens*, from *affluere*, (*ad* and *fluere*, 'to flow,') 'to flow to.' A flow or determination of humours, and particularly of blood, towards any part.

AFFLUXUS, *Fluxion*.

AFFRODILE, *Narcissus pseudo-narcissus*.

AFFUSIO, *Affusion*—a. *Frigida*, see *Affusion*—a. *Orbicularis*, *Placenta*.

AFFUSION, *Affusio*, *Pros'chysis*, *Epich'ysis*, from *ad*, 'to,' and *fundere*, *fusum*, 'to pour.' The action of pouring a liquid on any body. *Affusions*, *Rhyptolusis*, cold and warm, are used in

different diseases. The cold affusion, *Affusio seu Perfusio frigida*, is said to have been beneficial in cutting short typhus fever and scarlatina, if used during the first days. It consists in placing the patient in a tub, and pouring cold water over him; then wiping him dry, and putting him to bed. The only precaution necessary is, to use it in the state of greatest heat and exacerbation; not when chilliness, or topical inflammation, is present.

AFIUM, Opium.

AFTER-BIRTH, Secundines.

AFTER-PAINS, See Pains, labour.

AGACEMENT, (F.) from *ακάζειν*, 'to sharpen.' The setting on edge.

AGACEMENT DES DENTS. A disagreeable sensation experienced when acids are placed in contact with the teeth. *Tooth-edge*. Setting the teeth on edge.

AGACEMENT DES NERFS. A slight irritation of the system, and particularly of the organs of sense and locomotion, corresponding nearly to the English *Fidgets*.

AGAH, Intermittent Fever.

AGALACTATIO, Agalactia.

AGALACTIA, *Agalax'ia*, *Agalac'tio*, *Agalac'tio*, *Defectus lac'tis*, *Oligogal'ia*, *Oligogalac'tia*, from *a*, privative, and *γαλα*, 'milk.' Absence of milk in the mammae.

AGALAXIA, Agalactia.

AGAL'LOCIUM, from *αγאלλω*, 'I decorate,' *Calambac*, *Calambouk*, *Liq'mum Agal'lochi veri* seu *Al'oës* seu *Aspal'athi*, *Xyloölöës*, *Aloes wood*. A resinous and very aromatic wood of the East Indies, from *Eccæc'ria Agal'locha*, *Cynome'tra Agal'lochum*, *Alöës'ylon Agal'lochum*. Used in making pastils, &c. — Dioscorides, Oribasius, Paulus.

AGAMOUS, See Cryptogamous.

AGARIC, *Agar'icum*, *Agar'icus*. A genus of plants in the Linnaean system, some of which are edible, others poisonous. It is supposed to have been so called from *Agaria*, a region of Sarmatia. — Dioscorides. Among the edible varieties of the Boletus, the following are the chief:—1. The *Agar'icus edulis* seu *Aren'sis* seu *Sylvat'icus* seu *Campest'ris*, (F.) *Agaric comestible et champignon de couche*. 2. The *Agar'icus odoratus*, (F.) *Mousseron*. The most common poisonous varieties are the *Agar'icus necat'or*, (F.) *Agaric meurtrier*; and, 2. The *Agar'icus acris*, (F.) *Agaric âcre*; besides the *Auranite*, a sub-genus, which includes several species. One of the most delicate is the *Agaricus Aurantiacus*, but care must be taken not to confound it with the *A. Pseudo-aurantiacus*, which is very poisonous. The *A. aurantiacus* is called, in French, *Orange*. See Poisons, Table of.

AGARIC, see Boletus ignarius—*a. Amadouvier*, Boletus laticis—*a. Blanc*, Boletus laticis—*a. de Chêne*, Boletus ignarius—*a. Female*, Boletus ignarius—*a. of the Oak*, Boletus ignarius—*a. Odorant*, *Dædalea suaveolens*—*a. White*, Boletus laticis.

AGARICUM, Boletus ignarius.

AGARICUS, Boletus ignarius—*a. Albus*, Boletus laticis—*a. Arvensis*, see *Agaric*—*a. Aurantiacus*, *Amanitæ*, *Bolites*—*a. Auriculiformis*, *Peziza auriculæ*—*a. Campestris*, see *Agaric*—*a. Chirurgerum*, Boletus ignarius—*a. Ignarius*, Boletus ignarius—*a. Laticis*, Boletus laticis—*a. Pseudo-aurantiacus*, *Amanitæ*—*a. Quercus*, Boletus ignarius—*a. Sylvaticus*, see *Agaric*.

AGARIFIED, See Feverish.

AGASTRONERVIA, (F.) *Agastronervie*, from *a*, privative, *γαστήρ*, 'stomach,' and *νεῦρον*, 'nerve.' Want of nervous action in the stomach.

AGASYLLIS GALBANUM, Bubon galbanum.

AGATHIS DAMARRA, Pinus damarra.

AGATHOSMA CRENATUM, Diosma crenata.

AGATHOTES CHIRAYITA, Gentiana chirayita.

AGA'VE AMERICA'NA, *A. Ramo'sa*, *American Agave*, *American aloe*, *Mayney*, from *αγανος*, 'admirable.' *Nat. Ord.* Bromeliaceæ. *Sex. Syst.* Hexandria Monogynia. This plant has been considered diuretic and antisymphilitic. The favourite drink of the Mexicans—*Pulque*—is the fermented juice of this plant.

AGAVE RAMOSA, A. Americana.

AGAVE VIRGINICA, *Rattlesnake's master*, *False aloe*,—grows in the Southern States. The root is very bitter. It has been used in tincture as a carminative in colic; and as a remedy for bites of serpents.

AGE, *ἡλικία*, *Heli'kia*, *Ætas*;—Of uncertain etymon. Period of life. Time that has elapsed since birth, &c. Five ages are often designated in the life of man: 1. First infancy (*Infant'ia*;) 2. Second infancy (*Pueri'tia*;) 3. Adolescence (*Adolescen'tia*;) 4. The adult age (*Viril'itas*;) 5. Old age, (Old Eng.) *Ethl*, (*Senectus*.)

AGENEIOS, Imberbis.

AGEN'ESIS, from *a*, privative, and *γενεσις*, 'generation,' (F.) *Agénésie*. Imperfect development of any part of the body; as *cerebral agenesia*, i. e. imperfect development of the brain in the fetus. See Monster.

AGENNESIA, Impotence, Sterilitas.

AGENNESIS, Impotence.

AGENOSO'MUS, from *a*, privative, *γεννω*, 'I generate,' and *σῶμα*, 'body.' A malformation in which the fissure and evagination are chiefly in the lower part of the abdomen; the urinary or sexual apparatus absent, or very rudimentary.

AGENT, *Agens*, from *agere*, 'to act.' Any power which produces, or tends to produce, an effect on the human body. Morbific agents, (F.) *Agens morbosiques*, are the causes of disease;—therapeutical agents, (F.) *Agens thérapeutiques*, the means of treating it.

AGER NATURE, Uterus.

AGERASIA, *Insensescencia*, from *a*, privative, and *γῆρας*, 'old age.' A vigorous and green old age.

AGERATUM, *Achillea ageratum*—*a. Altissimum*, *Eupatorium ageratoides*.

AGERATUS LAPIS, *Asymptos*. A stone used by cobblers to polish shoes. It was formerly esteemed discutient and astringent.—Galen, Oribasius, Paulus.

AGES, Palm.

AGEUSIA, Ageusia.

AGEUS'TIA, *Ageus'tia*, *Ageusia*, *Apogeu'stia*, *Apogeu'is*, *Dysæsthesia gustato'ria*, *Anæsthesia gustato'ria*, *Parageu'is*, from *a*, priv., and *γεῦσις*, 'taste.' Diminution or loss of taste, *Anæsthesia lingua*.—Savages, Cullen.

AGGLOMERATE, *Agglomeratus*, from *agglomerare*, (ad and *glomerare*, 'to wind up yarn in a ball,') 'to collect together.' Applied to tumours or glands in aggregation.

AGGLUTINANT, *Agglutinans*, *Collet'icus*, *Glut'inans*, from *gluten*, 'glue,' (F.) *Agglutinant*, *Agglutinatif*, *Colletique*. Remedies were formerly so called, which were considered capable of uniting divided parts.—Paulus.

Plasters are called *agglutinants*, (F.) *agglutinatifs*, which adhere strongly to the skin. Certain bandages are likewise so termed; (F.) *Ban-dellettes agglutinatives*.

TO AGGLUTINATE. The French use the word *agglutiner* in the sense of 'to reunite,' as *agglutiner les lèvres d'une plaie*, 'to reunite the lips of a wound.'

AGGLUTINATIF, Agglutinant.

AGGLUTINATIO, Coition.

AGGLUTINATION, *Colle'sis*, *Epicolle'sis*, *Proscolle'sis*, *Glutinat'io*, from *agglutinare*, 'to glue together.' The first degree of adhesion. Also, the action of agglutinants.

AGGLUTINER, To agglutinate.

AGGREGATE, *Aggrega'tus*, from *aggregare*, (ad and *gregare*,) 'to flock together,' 'to assemble together.' Glands are called aggregate which are in clusters. See Peyer's Glandule. *Aggregate pills*, (F.) *Pilules agrégatives*, signified, formerly, those which were believed to contain the properties of a considerable number of medicines, and to be able to supply their place.

AGHEUSTIA, Ageustia.

AGHOUL, Agul.

AGIAHA'LID or AGIHA'LID or AGRAHA'LID, An Egyptian and Ethiopian shrub, similar to *Ximenia*. The Ethiopians use it as a vermifuge. The fruit is purgative.

AGIHALID, Agiaholid.

AGISSANT, Active.

AGITATION, *Agita'tio*, *Done'sis*, from *agere*, 'to act.' Constant and fatiguing motion of the body, *Tyrbē*, *Tyrba'sia*, *In'quies*,—or distressing mental inquietude,—*An'imi Agita'tio*.

AGITATORIUS, Convulsive.

AGLOS'SIA, from *a*, privative, and *γλωσσα*, 'the tongue.' A malformation, which consists in the want of a tongue.

AGLOSSOTOMA, from *Aglossia*, and *τομα*, 'mouth.' A mouth without a tongue.

AGLOSSOSTOMOG'RAPHY, *Aglossostomogra'phia*, from *a*, priv., *γλωσσα*, 'the tongue,' *στομα*, 'the mouth,' and *γραφω*, 'I describe.' Description of a mouth without a tongue.—Rohland, (of Saumur.)

AGLUTI'TION, *Agluti'tio*, from *a*, priv., and *glutire*, 'to swallow.' A hybrid term, designating impossibility of swallowing.—Linnaeus.

AGMA, Fracture.

AGMATOLOG'IA, from *αγμα*, *αγματος*, fracture, and *λογος*, 'a description.' The doctrine of fractures. A treatise on fractures.

AGME, Fractures.

AGMINA DIGITORUM MANUS, Phalanges of the fingers.—*a*. Membrana, Amnios.

AGMINATED GLANDS, Peyer's glands.

AGNAIL, Hangnail.

AGNA'THIA, from *a*, priv., and *γναθος*, 'jaw.' A malformation, which consists in the want of the jaw, especially of the lower.

AGNINA MEMBRANA, Amnios.

AGNO'IA, *Agne'a*, from *a*, priv., and *γινωσκω*, 'I know.' State of a patient who does not recognise individuals.—Hippocrates, Galen, Foësius.

AGNUS CASTUS, Vitex.

AGO'GE, *αγωγή*. The order or condition of a disease.—Hippoc., Galen. Likewise the state of the air.—Hippoc., Galen, Gorraeus, Foësius.

AGOGUE, *αγωγή*, 'a leader,' from *αγω*, 'I lead or expel.' Hence *Cholagogue*, an expeller of bile; *Hydragogue*, &c.

AGOMPHIASIS, *Agompho'sis*, from *a*, privative, and *γομφω*, 'I nail.' Looseness of the teeth.—Gorraeus. See Gomphiasis.

AGOMPHOSIS, Agomphiasis.

AGON, Agony.

AGONE, Hyoscyamus.

AGONIA, Sterilitas.

AGONISMA, Agony.

AGONISMUS, Agony.

AGONIS'TICA, from *αγών*, 'a combat.' The part of ancient gymnastics, which had reference to the combats of the Athlete.

Also, very cold water, given internally to calm febrile heat.—Paulus of Ægina.

AGONIZANS, Psychorages.

AGONOS, Sterile.

AG'ONY, *Agon'ia*, *Agon*, *Agonis'ma*, *Agonis'-mus*, *Muchthos*, *Mogus*, *Psychorag'ia*, *Psychorhag'ia*, *Angor*, *Throe*, *Throe*, (Sc.) *Patient of death*, from *αγών*, 'a combat.' The last struggle of life.—Galen, Gorraeus, &c. The agony, which is of longer or shorter duration, is characterized by great change in the features, gradual abolition of sensation and motion, loss of voice, dryness or lividity of the tongue and lips, rattling in the throat, small and intermittent pulse, and coldness of the extremities. This state is not present in those who die suddenly. See *Facies Hippocratica*.

AGOS'TUS, from *αγω*, 'I lead.' The fore arm from the elbow to the fingers. Also, the palm of the hand.—Gorraeus. See Palm.

AGRA, *αγρα*, from *αγρω*, 'I seize hold of.' A seizure, as *Odontagra*, a tooth seizure, toothache; *Chiragra*, *Podagra*, &c.

AGRAFE DE VALENTIN. A kind of forceps with parallel branches, employed by Valentin in the operation for hare lip, to effect the approximation of the edges of the wound.

AGRAHALID, Agiaholid.

AGRÉGATIVES PILULES. See Aggregate.

AGRÉMIE, from *agrie*, 'gout,' (Piorry,) and *αίμα*, 'blood.' The altered condition of the blood in gout.—Piorry.

AGRIA, Herpes exedens, Lichen agrius.

AGRIAMPELOS, Bryonia alba.

AGRICOCCEMELEA, Prunus Spinosa.

AGRIE, Gout.

AGRIFOLIUM, Ilex aquifolium.

AGRIMONIA, Agrimony—a. Eupatoria, Agrimony—a. Odorata, Agrimony—a. Officinalis, Agrimony.

AGRIMONIA PARVIFLO'RA, Small-flowered or Sweet-scented Agrimony; indigenous; has similar properties to A. Eupatoria. See Agrimony.

AGR'IMONY, *Agrimo'nia*, *A. Eupatoria* seu *odorata* seu *officinalis*, *Caj'al*, *Lacip'ula hepatica*, *Cockle-bur*, *Stickwort*. (F.) *Aigremoire*. Ord. Rosaceæ. Sex. Syst. Icosandria Digynia. A mild astringent and stomachic. Dose, in powder, from ℥j to ʒj.

AGRIMONY, HEMP, *Eupatorium cannabinum*—a. Small-flowered, *Agrimonia parviflora*—a. Sweet-scented, *Agrimonia parviflora*.

AGRIOCASTANUM, *Bunium bulbocastanum*, *Lycoperdon tuber*.

AGRIOCINARA, *Sempervivum tectorum*.

AGRIORIGANUM, *Origanum majorana*.

AGRIOSELINUM, *Smyrnum olusatrum*.

AGRIOTHYM'IA, from *αγριος*, 'ferocious,' and *θυμος*, 'disposition.' Ferocious insanity.—Sauvages.

AGRIPALMA GALLIS, *Leonurus cardiaca*.

AGRIP'PA, *Ægrip'pa*, from *ager partus*, 'difficult birth' or perhaps from *αγπα*, 'taking or seizure,' and *πους*, 'the foot.' This term has been given to those born by the feet. It is pretended that the family of Agrippa obtained their name from this circumstance. Parturition, where the feet present, is called *Agrippæ partus*, *Agrippinus partus*.

AGRIPPINUS PARTUS, see Agrippa.

AGRO DI CEDRO, see Citrus medica.

AGROPYRUM LÆVISSIMUM, *Triticum repens*.

AGROSTEMMA GITHAGO, *Lychnis githago*.

AGROSTIS, *Bryonia alba*—a. Verticillata, *Andropogon muricatus*.

AGRUNA, *Prunus spinosa*.

AGRYPNIA, Insomnia.

AGRYPNOCOMA, *Coma vigil*.

AGRYPNO'DES, from *αγρυπνος*, 'sleepless.' Characterized by sleeplessness, as *Febris Agrypnodes*, a fever accompanied with sleeplessness.

AGRYPNOTICUS, Anthypnotic.
AGRYPNUS, *aycynos*. Sleepless, vigilant.
AGUA DE VERUGA, see Verugas.
AGUARDIENTE, Brandy. See also Spirit.
a. de Italia, see Spirit.
A'GUE, from Gothic, *agis*, 'trembling.' (?) Intermittent fever; often used in the same sense as chill or rigor.
AGUE AND FEVER, Intermittent fever.
AGUE, BROW, Neuralgia frontalis.
AGUE CAKE, *Placenta febrilis seu sple'nica*, *Physconia sple'nica*, *Splenis Tumor*; (F.) *Gâteau fébrile*. A visceral obstruction—generally in the spleen—which follows agues, and is distinctly felt by external examination. To a greater or less degree, it is not uncommon.
AGUE, DEAD, see Fever, masked.
AGUE DROP, Tasteless, Liquor arsenicalis.
AGUE, DUMB, see Fever, masked—a. Face, Neuralgia, facial.
AGUE, FREE, *Laurus sassafras*.
AGUE, LEAPING, see Leaping ague—a. Quartan, Quartan—a. Tertian, Tertian fever.
AGUE TREE, *Laurus sassafras*.
AGUE, THIRD-DAY, Tertian.
AGUE WEED, *Eupatorium perfoliatum*.
AGU'IOS, from *a*, priv., and *yucov*, 'limb.' Mutilated or wanting limbs.—Hippocr. Weak, feeble.—Galen.
AGUL, *Aghoul*, *Alha'gi*, the *Hedysarum seu Hedysarum alhagi*. A thorny shrub of Persia and Mesopotamia, which affords manna. The leaves are purgative.
AGYR'IAS, from *ayyos*, 'a collection.' Opacity of the crystalline.—Aëtius, Paré.
AGYR'TA, from *ayyos*, 'a crowd.' Formerly, a stroller who pretended to supernatural powers. Subsequently, a quack or illiterate pretender.—See Charlatan.
AGYRTIA, Charlatanry.
AHO'RA, from *a*, privative, and *spa*, 'youth.' Tardy development of the organs: the opposite to *Hyperho'ra*.
AHOUI, Thevetia ahouai.
AHUSAL, Orpiment.
AHYPNIA, Insomnia.
AIAULT, *Narcissus pseudo-narcissus*.
AIDE (F.), *Ad'jutor min'ister*. An assistant to a surgeon in his operations.
AIDOIOMANIA, Nymphomania.
AIERSA, Iris Germanica.
AIGE, Ægias.
AIGLE, MINERAL WATERS OF. Near the city of this name, in Normandy, is the chalybeate spring of St. Xantin, much used in the 16th and 17th centuries.
AIGLE, Ægias.
AIGRE, Acidulous—a. *Voiz*. See Acid.
AIGRELET, Acidulous.
AIGRETTE, *Rumex acetosa*, see Typha latifolia.
AIGREMOINE, Agrimony.
AIGREURS, Acidities.
AIGU, Acute.
AIGUES-CAUDES, MINERAL WATERS OF, *Aignes-chaudes*, *Eaux chaudes*, *Aqua calida*. Sulphurous springs, about six miles from Bonnes, Basses Pyrénées, in France: five in number, the temperature of which is, respectively, about 70°, 81°, 92°, 93°, and 95°.
AIGUILLE, Needle—a. à *Acupuncture*, see Needle—a. à *Appareil*, see Needle—a. à *Bec de Lièvre*, see Needle—a. à *Cataracte*, see Needle—a. de *Deschamps*, see Needle—a. *Engainée*, see Needle—a. à *Fistule*, see Needle—a. à *Gaine*, see Needle—a. à *Ligature*, see Needle—a. à *Manche*, see Needle—a. à *Séton*, see Needle—a. à *Suture*, see Needle.
AIGUILLON (F.), *Spina Helmontii*. A

term used since the time of Van Helmont to designate the proximate cause of inflammation. According to him, an inflamed part is in the same condition as if an *aiguillon* or thorn were thrust into it.
AIGUISER, To acidulate.
AIL, Allium.
AILE, Ala, *Aileron*.
AILERON (F.), *Extrema Ala seu Pin'nula*, diminutive of (F.) *Aile*, a wing. The extremity of the wing of a bird, to which the great feathers are attached.
AILERONS DE LA MATRICE. Three folds at the base of the broad ligaments of the uterus, which are occupied by the ovary and its ligament, the Fallopian tube, and the round ligament.
AILING, Sick, Sickly.
AILEMENT, Disease.
AIMA, 'aipa, see Hæma.
AIMANT, Magnet.
AIMATERA, Hepatirrhœa.
AIMORRHEA, Hæmorrhagia.
AIMORRHOIS, Hæmorrhoids.
AINE, Inguen.
AIPATHIA, Continent disease.
AIPI, *Jatropha manihot*.
AIPIMA COXERA, *Jatropha manihot*.
AIPIPOCA, *Jatropha manihot*.
AIR, *Aër*, *Pneuma*, from *aw*, 'I breathe.' Common Air, Atmospheric air, (F.) *Air atmosphérique*, is an invisible, transparent, inodorous, insipid, ponderable, compressible, and elastic fluid, which, under the form of the atmosphere, surrounds the earth to the height of 15 or 16 leagues.
Air is essentially composed of two gases, oxygen and nitrogen, in the proportion of 20 of the former to 80 of the latter. Oxygen is the vital portion, but the nitrogen is necessary to dilute it. Air also contains a small portion of carbonic acid gas, and has always floating in it aqueous vapour, different terrestrial emanations, etc. Its effects upon the human body vary according to its greater or less density, temperature, moisture, etc.; hence, change of air is found extremely serviceable in the prevention and cure of certain morbid conditions. See Climate, and Respiration.
AIR ACIDE VITRIOLIQUE, Sulphurous acid—a. *Alcalin*, Ammonia—a. *Atmosphérique*, Air.
AIR BLADDER, *Swim-bladder*, *Swimming-bladder*, *The Swim*, (F.) *Vessie natatoire*. An abdominal organ in many fishes, sometimes communicating by means of a duct, *Ductus pneumaticus*, with the alimentary canal, at others, not, which is considered by some to belong to the respiratory system. Its contents are the elements of atmospheric air, but in different proportions; and its chief and general function appears to be to regulate the specific gravity of the fish.
AIR CELLS OF THE LUNGS, Bronchial cells; see Cellule—a. Chamber, Folliculus æris.
AIR, DEPHLOGISTICATED, Oxygen—a. Empty, Oxygen.
AIR DU FEU, Oxygen.
AIR, FACTITIOUS, Carbonic acid—a. Fixed, Carbonic acid—a. *Gaté*, Azote—a. *Hépatique*, Hydrogen sulphuretted—a. Inflammable, Hydrogen, Hydrogen carburetted—a. Mephitic, Carbonic acid.
AIR PASSAGES, (F.) *Voies aériennes*, *V. aérières*. The larynx, trachea, bronchia, etc.
AIR PUANT, Hydrogen, sulphuretted.
AIR, PURE, Oxygen—a. Solid, of Hales, Carbonic acid—a. *Vicié*, Azote—a. Vital, Oxygen.
AIRAIN, Bell-metal, Brass.
AIRE, Areola.
AIRELLE ANGULEUSE, *Vaccinium myrtillus*—a. *Ponctué*, *Vaccinium vitis idæa*.
AIRIGNE, Hook.

AIRTHREY, MINERAL WATERS OF.

Airthrey is situate about two miles north of Stirling, Scotland. The waters are saline cathartics; containing chloride of sodium, chloride of calcium, sulphate of zinc, and chloride of magnesium.

AISELLE, Axilla.

AISTHESIS, *Æsthesis*.

ATHOMO'MA, from *αἶσος*, 'black.' A black condition of all the humours of the eye. A. Paré.

ATIA, Cause.

ATIIOLOGY, *Ætiologia*.

ATION, Cause.

AIX-LA-CHAPELLE, MINERAL WATERS OF. Called by the Germans, Aachen. A thermal, sulphureous mineral water, which contains, in 1000 grammes, 28.54 cubic inches of sulphuric acid gas, 18.05 cubic inches of carbonic acid gas, 0.1304 grammes of carbonate of lime, 0.0440 grammes of carbonate of magnesia, 0.5444 grammes of carbonate of soda, 2.3697 grammes of chloride of sodium, 0.2637 of sulphate of soda, and 0.0705 of silica. The temperature is 134° Fahrenheit.

The factitious water of *Aix-la-Chapelle*, *A'qua Aquigranensis*, (F.) *Eau d'Aix-la-Chapelle*, is made by adding pure water f³ xvijss, to hydro-sulphuretted water f³ iv., carbonate of soda gr. xx, chloride of sodium gr. ix.—Ph. P.

There are thermal sulphureous springs at Aix in Savoy (98°), and some thermal springs at Aix in Provence (91°).

AIZOON, *Sempervivum tectorum*.

AJUGA, *A. pyramidalis*, *Consolida me'dia*, *Bu'gula*, *B. pyramidalis*, *Tro'nerium pyramidalis*, Upright Bugloss, Middle Consound, (F.) *Bugle pyramidale*. This plant is subastringent and bitter.

AJUGA CHAMÆPITYS, *Teucrium chamæpitys*.

AJUGA REPTANS, *Bu'gula*, *B. reptans*, Common Bugle, (F.) *Bugle rampante*, has similar properties.

AKATALIS, *Juniperus communis*.

AKATERA, *Juniperus communis*.

AKERATODIAPHANIE, see *Caligo*.

AKINESIA, *Acinesia*.

AKNEMIA, *Acnemia*.

AKOLOGY, *Materia Medica*.

AKRATOPEGE, *Acratopegae*.

AKSIS, *Intermittent fever*.

ALA, *Pinna*, *Pteryx*, 'a wing,' (F.) *Aile*. A term often used by anatomists for parts which project like a wing from the median line; as the *Ala nasi*, *Ala of the uterus*, etc. See *Axilla*, and *Pavilion of the Ear*. Also, *Pterygium*.

ALA EXTREMA, see *Aileron*.

ALABAMA, MINERAL WATERS OF. *Tallahatta springs* contain sulphur, and salts of iron, lime, and magnesia. But the most noted are those of *Bladen*. See *Bladen*, Mineral Waters of. *Bailey's spring* is an acidulo-sulphureous chalybeate. It is fourteen miles from *Tuscumbia*.

ALABASTER, *Alabastrum*, (F.) *Albâtre*, *Alabastrites*. A variety of compact gypsum; of which an ointment was once made—the *unguentum alabastrinum*; used as a discutient. Alabaster likewise entered into several dentifrices.

ALABASTRITES, *Alabaster*.

ALÆ INTERNÆ MINORES CLITORIDIS, *Nymphæ*—a. *Magnæ*, see *Sphenoid bone*—a. *Majores*, *Labia pudendi*—a. *Minores*, *Nymphæ*, see *Sphenoid bone*—a. *Muliebres minores*, *Nymphæ*—a. *Nasi*, see *Nasus*—a. *Pudendi* *Muliebres*, *Labia pudendi*—a. *Pulmonum*, see *Pulmo*—a. of the *Uterus*, see *Ala*—a. *Vespertilionis*, see *Uterus*.

ALAISE, *Alææ*.

ALAITER, from (F.) *lait*, 'milk.' To suckle.

ALALIA, *Mutitas*.

ALAMBIC, *Alembic*.

ALANFUTA. A name given by the Arabians to a vein, situate between the chin and lower lip, which they were in the habit of opening in cases of factor of the breath.—*Avicenna*.

ALAUQUECA. The Hindoostanee name of a stone, found in small, polished fragments, which is considered efficacious in arresting hæmorrhage when applied externally. It is a sulphuret of iron.

ALARES MUSCULI, *Pterygoid muscles*.

ALA'RES VENÆ. The superficial veins at the fold of the arm.

ALARIA OSSA. The wing-like processes of the sphenoid bone.

ALA'RIS, *Ala'tus*, *Aliform'is*; from *ala*, 'a wing.' Wing-shaped, winged.

ALATERNUS, *COMMON*, *Rhamnus alaternus*—a. *Latifolius*, *Rhamnus alaternus*.

ALAT'US, *Pterygo'des*, *Homo ala'tus*. One whose scapulae project backwards like wings.

ALBA'DARAN, *Aladaban*. The sesamoid bone of the metatarso-phalangeal joint of the great toe. The Rabbis and Magicians attributed extraordinary virtues to it.—*Arabians*.

ALBAGIAZI, *Sacrum*.

ALBAMENTUM, *Albumen ovi*.

ALBAN, SAINT, MINERAL WATERS OF. A French acidulous chalybeate, in the department of the Loire.

ALBARAS ALBA, *Lepra alphoides*—a. *Nigra*, *Lepra nigricans*.

ALBARES, *Lepra alphoides*.

ALBAROS, *Lepra alphoides*.

ALBÂTRE, *Alabaster*.

ALBEDO UNGUIUM, see *Nail*.

ALBIN B'CEUF, *Albumen ovi*.

ALBINISM, see *Albino*.

ALBINISMUS, see *Albino*.

ALBINO, 'White.' *Leuce'thiops*, *Æthiops albus*, *Dondo*, from *albus*, 'white,' (F.) *Blafard*, *Nègre-blanc*. A Spanish word applied to individuals of the human race who have the skin and hair white (*Achromodermie* and *Achromotrichomie*.—*Piorry*); the iris very pale, bordering on red; and the eyes so sensible that they cannot bear the light of day. This condition, which has been called *Leucæthiopia*, *Alpho'sis*, *Æthiopia*, *Albino'smus*, *Albinis'mus*, *Al'binism*, *Kakerlakism*, *Leucopathia*, *Leuco'sis*, is seen more frequently in the Negro. Both sexes are exposed to it. It does not seem to be true that there are tribes of Albinos in the interior of Africa.

ALBINOISMUS, see *Albino*.

ALBOR OVI, *Albumen ovi*.

AL'BORA. A kind of itch or complicated leprosy.—*Paracelsus*.

ALBOT, *Crucible*.

ALBOTIM, *Terebinthina*.

ALBUGINÉ, *Albuginea*, *Albugineous*.

ALBUGIN'EA, *A. Testis*, *Tu'nica albugin'ea*, *Perites'tis*, *Dura mater testis*, *Membra'na capsula'ris testis*. (F.) *Albuginé*, *Tunique albuginée*. A strong, fibrous, and resisting membrane, which immediately envelopes the testicle, and has, at its upper part, an enlargement, called *corpus Highmorianum*. From its inner surface it sends off a number of flat, filiform prolongations or septa, between which are contained the seminiferous vessels. Externally, it is covered by the tunica vaginalis testis.

ALBUGIN'EOUS, *Albugin'eus*, 'white,' from *albus*, (F.) *Albuginé*. A term applied to textures, humours, &c., which are perfectly white.

ALBUGIN'EOUS FIBRE, (F.) *Fibre albuginée*. A name given by *Chaussier* to what he considers one of the four elementary fibres.

The albugineous fibre is linear, cylindrical, tenacious, elastic, but little extensible, and of a shining, satiny appearance. It forms fasciæ cr

fascioli, which constitute the tendons, articular ligaments, and aponeuroses; hence the name *Albugineous membranes*, given by Chaussier to the fibrous membranes.

Gauthier considered, that the rete mucosum consists of four layers, to two of which he gives the names *membrana albuginea profunda* and *membrana albuginea superficialis*, respectively.

ALBUGINITIS, (F.) *Albuginitis*. A term employed by some authors for inflammation of the albugineous tissue. Thus, gout and rheumatism are regarded as species of the genus albuginitis.

ALBUGO OCULORUM, Leucoma—a. Ovi, Albumen ovi.

ALBULA, Leucoma.

ALBUM CANIS, Album græcum—a. Ceti, Cetaceum.

ALBUM GRÆCUM, *Album Canis*, *Cynocœprus*, *Spodium Græcorum*, *Stercus Caninum Album*. The white dung of the dog. It consists almost wholly of phosphate and carbonate of lime, from the bones used as food. It was formerly applied as a discutient gargle in quinsies, and as an anti-epileptic; but is now justly banished from practice.

ALBUM NIGRUM. The excrement of the mouse.

ALBUM OCULI, see Sclerotic.

ALBUM RHAZIS. A white ointment made of ceruse and lard, prescribed by the Arabian physician Rhazes.

ALBUMEN, *Leuco-ma*, *Ooni-nē*, *Ozemun*, from *albus*, 'white.' (F.) *Albumine*. An immediate principle of animals and vegetables, which constitutes the chief part of the white of egg. It is found in the serum, chyle, synovia, serous fluids, &c. There is not much difference in chemical composition between animal and vegetable albumen, fibrin and casein. Also, the white of the eye. See Sclerotic.

ALBUMEN OVI, *Ovalbumen*, *Albu'men*, *Albu'mor*, *Albu'go* seu *Albur* seu *Can'didum Ovi*, *Clare'ta*, *Ovi albus liquor*, *Albumen'tum*, *Lac avis*, white of egg, (F.) *Blanc d'œuf*, (Old F.) *Albin d'œuf*, is used in pharmacy for suspending oils, &c., in water. See Ovum.

ALBUMEN, SALIVARY, see Saliva.

ALBUMINE, Albumen—a. *Cérébrale*, *Neurine*—a. *du Cerveau*, *Neurine*.

ALBUMINOID, *Albuminoides*, from *albumen*, and *idos*, 'resemblance.' Resembling albumen, as fibrin, casein, &c. It is often applied to aliments in the same sense as proteinous.

ALBUMINOSE. The soluble principle of fibrin, identical with the dominant principle of the white of egg.—Bouchardat. According to others, the ultimate product of the transformation, in the stomach, of albuminoid matters, which renders them fit for assimilation and nutrition. Albuminose is not coagulable by heat, and incompletely so by acids; an excess of acid dissolving the precipitate. Robin and Verdel regard it as synonymous with the casein of the small intestine of Tiedemann and Gmelin, the *gelatiniform matter of the intestine* of Prévost and Morin, and the casein of the blood of Dumas. See Peptone.

ALBUMINOSIS. A condition of the blood, in which the ratio of albumen is increased, as in abdominal typhus, variola, rubéola, &c. Moser and Strahl.

ALBUMINURIA. A hybrid term from '*albumen*,' and *urupa*, 'the urine.' A condition of the urine in which it contains albumen, the presence of which is indicated by its coagulation on the application of adequate heat. See Kidney, Bright's disease of the.

ALBUMINURIC, *Albuminuricus*, (F.) *Albuminurique*. Relating or appertaining to albuminuria. One labouring under Albuminuria.

ALBUMINURORRHÉE, Kidney, Bright's disease of the.

ALBUMOR, Albumen ova.

AL'CAEST, *Al'cahest*, *Al'chaest*, perhaps from (G.) all, 'all,' and geist, 'spirit.' A word invented by Paracelsus to designate a liquor, which, according to him, was capable of removing every kind of swelling.

The same word was used by Van Helmont for a fancied universal solvent, capable of reducing every body to its elements.

ALCAEST OF GLAUBER is a thick liquor obtained by detonating nitrate of potassa on hot coals, which transforms it into subcarbonate of potassa.

ALCAEST OF RESPOUR is a mixture of potassa and oxyd of zinc.

ALCAHEST, Alcaest.

ALCAHOL, Alcohol.

ALCALES'CENCE, *Alkales'cence*, *Alcalescen'tia*. The condition in which a fluid becomes alkaline.

ALCALESCENCE OF THE HUMOURS was an old notion of the humourists. It can only occur during the putrid fermentation of animal matters, which contain azote, and produce ammonia. **ALCALINITY** is the quality of being alkaline.

AL'CALI or *Alca'li*, *Al'kali*, from *al* (Arab.,) 'the,' and *kali*, the name of the *Salsola Soda*, a plant which contains a large quantity of one of the principal alkalis—*soda*. The alkalis are substances soluble in water, possessing generally a urinous, acrid, and caustic taste, turning the syrup of violets green, and restoring to blue infusion of litmus, which has been reddened by acids; reddening the yellow of turmeric, and having the greatest tendency to unite with acids, whose character they modify, and form salts with them. In medicine we understand by this term *Potassa*, *Soda*, or *Ammonia*.

ALCALI, CAESTIC, *Al'kali Causticum*. A pure alkali. One deprived of its carbonic acid.

ALCALIS, Fixed, Soda and potassa; **VOLATILE ALCALI**, Ammonia.

ALCALI AMMONIACUM ACETATUM, Liquor ammoniac acetatis—a. Ammoniacum fluidum, Liquor ammoniac—a. Fixum tartarizatum, Potassæ tartaras—a. Minerale sulphuricum, Soda, sulphate of—a. Tartari aceto saturatum, Potassæ acetas—a. Vegetabile salito dephlogisticatum, Potassæ murias hyperoxygenatus—a. Vegetabile tartarizatum, Potassæ tartaras—a. Vegetabile vitriolatum, Potassæ sulphas—a. Volatile acetatum, Liquor ammoniac acetatis—a. Volatile aeratum, Ammoniac carbonas—a. Volatile ex sale ammoniaco, Ammoniac carbonas.

ALCALIGENE, Azote.

ALCALINITY. See Alcalescence.

AL'CALOID; from *alca'li* and *idos*, 'form.' A name given to the organic alcalis to distinguish them from the mineral, from which they differ as regards composition and general properties;—having nothing in common but their basic properties. Brucia, Emetia, Morphia, Strychnia, &c., are alcaloids.

ALCANA, *Anchusa officinalis*.

ALCANA MAJOR LATIFOLIA DENTATA, Prinos—a. Orientalis, Lawsonia inermis—a. Spuria, *Anchusa tinctoria*—a. Vera, *Lawsonia inermis*.

ALCEA, *Hibiscus abelmoschus*—a. *Ægyptiaca*, *Hibiscus abelmoschus*—a. Indica, *Hibiscus abelmoschus*.

AL'CEA Ro'SEA, *Common hollyhock*, *Hoc, Hoc'ce*. Emollient, like Althæa.

ALCIACHENGE, Physalis.

ALCHACHIL, Rosmarinus.

ALCAEST, Alcahest.

ALCHEMIL'LA, said to have been celebrated

with the Alchemists [?], supposed by some to be of Arabic derivation. *A. vulgaris*, *Common Ladies' Mantle*, *Pes Leonis*, *Leontopodium*, (F.) *Pied de Lion*. Formerly in great repute as an astringent in hemorrhage.

ALCHEMY, Alchymy.

ALCHITRAM, see *Pinus Sylvestris*.

ALCHITURA, see *Pinus Sylvestris*.

ALCOHOL, Alcohol.

ALCHORNEA LATIFOLIA, see *Alcornouque*.

AL'CHYMY, *Al'chemy*, *Alchemi'a*, *Alchimi'a*, *Adepta Philosophia*, from *al*, an Arabic particle, signifying, 'superiority, excellence,' and *Chimia*, 'Chymistry.' This word was formerly synonymous with Chymistry; but, from the 7th century, it has been applied to the mysterious art of endeavouring to discover a universal remedy, and a mode of transmuting the baser metals into gold: an operation to which they gave the name *Opus magnum*, and *Philosopher's stone*.

Alchymy has also been called *Scien'tia vel Philosophia Hermetica*, from an idea that Hermes or Mercury was its inventor.

Harris has well defined this chimerical art: '*Ars sine arte, cujus principium est mentiri, medium laborare, et finis mendicare.*'

AL'CHYMYST, (Old Eng.) *Alkymistre*, *Flatuarius*, *Adept'*. One pretending to alchymy.

ALCOCALUM, *Cynara scolymus*.

AL'COHOL, *Al'cohol*, *alcohol*, *Alkol*, *Alcol*, *Al'cool*, *Al'kool*. An Arabic word, formerly used for an impalpable powder, and signifying 'very subtle, much divided.' At the present day it is applied to highly rectified spirit of wine:—see *Spiritus rectificatus* or *rectified spirit*, distilled from dried subcarbonate of potassa. In the Ph. U. S., Alcohol is rectified spirit of the specific gravity 0.835. The Dublin college has a *spiritus fortior*, used in the preparation of certain essences, whose specific gravity is .818. *Alcohol, absolute alcohol*, of the Edinburgh and Dublin Pharmacopœias, is of specific gravity .797.

Alcohol is an inflammable liquor, lighter than water, of a warm, acrid taste, colourless, transparent, and of a pungent, aromatic smell. It is the product of the distillation of vinous liquors; is miscible with water in all proportions, and is the direct solvent of resins, balsams, &c. Various other vegetable principles are soluble in it, and hence it is used, in different states of concentration, in the preparation of *elcirs*, *tinctures*, *essences*, &c.

Alcohol acts on the animal body as a powerful stimulus: as such, in a dilute form, it is used in the prevention and cure of disease. Its habitual and inordinate use is the cause of many serious affections, of a chronic character especially, as visceral obstructions, dropsy, &c.

ALCOHOL, ABSOLUTE, see *Alcohol*—a. *Ethereus Ferratus*, *A. Sulfurico-æthereus ferri*—a. *cum Aloe perfoliata*, *Tinctura aloes*—a. *Ammonie et guaiaci*, *Tinctura guaiaci ammoniata*—a. *Ammoniatum*, *Spiritus ammonie*—a. *Ammoniatum aromaticum*, *Spiritus ammonie aromaticus*—a. *Ammoniatum fetidum*, *Spiritus ammonie fetidus*—a. *Amylicum*, *Oil, fusel*—a. *cum Aromatibus sulphuricatis*, *Sulphuricum acidum aromaticum*—a. *cum Aromatibus compositis*, *Tinctura cinnamomi composita*—a. *Castoriatum*, *Tinctura castorei*—a. *cum Crotonæ cascariâ*, *Tinctura cascariæ*—a. *Dilutum*, *Spiritus tenuior*—a. *Ferratus*, *Tinctura ferri muriatis*—a. *cum Sulphate ferri tartarisatus*, see *Ferrum tartarisatum*—a. *cum Guaiaco officinale ammoniatus*, *Tinctura guaiaci ammoniata*—a. *Iodii*, *Tinctura Iodini*—a. *cum Opio*, *Tinctura opii*—a. *Methylic*, *Spiritus Pyroxylicus*—a. *Pyroxylic*, *Spiritus Pyroxylicus*—a. *Sulphuricatum*, *Elixir acidum Halleri*—a. *Sulphuricum*, *Elixir acidum*

Halleri—a. *Sulphuris*, *Carbonis sulphuretum*—a. *Vini*, *Spiritus rectificatus*—a. *Wood*, *Spiritus pyroxylicus*.

ALCOHOL'IC, *Alcohol'icus*, *Spirituosus*, *Spir'ituosus*. (F.) *Alcoholic*. Relating to or containing alcohol—as an *alcoholic drink* or *remedy*.

ALCOHOLISM'US, *Al'coholism*, (F.) *Alcöolisme*. The series of morbid phenomena produced by the use of alcoholic liquors.

ALCOL, Alcohol.

ALCOLÆ, Apthæ.

ALCOLAT, *Alcolat*.

ALCOOL, Alcohol—*a. Camphoræ*, *Spiritus camphoræ*—*a. de Soufre*, *Carbonis sulphuretum*.

ALCOOLAT, Spirit. It now generally signifies alcohol, charged, by distillation, with the volatile principles of certain substances, or a distilled spirit,—*Alcolat* (*Codex med.* of Paris).

ALCOOLATUM, Tincture—*a. Antiseorbuticum*, *Tinctura de Cochleariis*—*a. Carminativum Sylvi*, *Tinctura de Cochleariis*—*a. de Croco compositum*, *Tinctura de Croco composita*.

ALCOOLATURE, *Teinture avec les plantes fraîches*. A name given by M. Bérard to certain preparations introduced by him, which consist of alcohol, charged, by maceration, with the soluble principles of fresh plants. See *Tincture*.

ALCOOLÉ. A name given, in the new French pharmaceutical nomenclature, to medicines resulting from the solvent action of alcohol on different substances, or from its admixture with liquids which unite with it in all proportions. See *Tincture*.

ALCOOLÉ SUCRÉ, *Ratafia*.

ALCOOLIQUE, *Alcoholic*.

ALCOOLISER (F.) Formerly, 'to reduce into an impalpable powder.' No longer used.

ALCÖOLISME, *Alcoholismus*.

ALCOOLOMETER, *Areometer*.

ALCOOMETER, *Areometer*.

ALCORNOCQUE (F.), *Cortex Alcornoco*. The bark of *Alchor'nea latifolia*, of Jamaica, which has been considered capable of curing phtisis. It is bitter, tonic, and slightly astringent. Dose of the powder \mathfrak{z} i to \mathfrak{z} ss.

AL'CYON, *Hal'cyon*. A swallow of Cochin China, whose nest is gelatinous and very nutritious. It has been proposed in medicine as anæsthetic and aphrodisiac.

ALCYONIUM, *Bastard sponge*. The ashes were formerly employed as dentrifices: they were believed proper for favoring the growth of the hair and beard, and were used in Alopecia.

ALDABARAN, *Albadaran*.

ALDEHYDE, see *Anæsthetic*.

ALDER, AMERICAN, *Alnus serrulata*—*a. Black*, *Prinos*, *Rhamnus frangula*—*a. European*, *Alnus glutinosa*—*a. White*, *Clethra alnifolia*.

ALE, *Cerevisia*.

ALE, WHITE. A drink, used in Devonshire, England, amongst the ingredients of which are milk, spice, and spirit.

ALEACAS, *Glycyrrhiza*.

ALEBERRY. A beverage, formerly made by boiling *ale* with *spice* and *sugar*, and *sops of bread*. It was given to invalids.—*Palsgrave*.

ALECOST, *Tanacetum balsamita*.

ALECTORIUS LAPIS, *Alecto'ria*; from *alektrōp*, 'a cock.' The name of a stone, supposed to exist in the stomach of the cock, or, according to others, in that of the capon, four years old. Many marvellous properties were formerly attributed to it, which are as groundless as its existence. There are no stones in the stomach, except what have been swallowed.

ALEGAR, *Acetum*.

ALEHOOF, *Glechoma hederacea*.

ALEIMMA, *Liniment*.

ALEIPHA, Liniment.

ALEIPTERIUM, from ἀλείφω, 'I anoint.' The place in the ancient gymnasium where the combatants anointed themselves.

ALEIP'TRON. Same etymon. A box for containing ointments.

ALEMA, Farina.

ALEM'BIC (Arab.), *Moorshead, Capitellum, Capitulum, Ambicus*, (F.) *Alambic*. A utensil made of glass, metal, or earthen ware, adapted for distillation. A still. It consists of a body or cucurbit, (F.) *cucurbita, chaudière*, to which is attached a head or capital, (F.) *chapiteau*, and out of this a beak descends laterally to be inserted into the receiver, worm, condenser, or refrigerator, (F.) *serpentin, réfrigérant*, as the case may be.

ALEM'BROTH (Salt.) *Sal Alembroth*. The alchemists designated by this name, and by those of *Sal sapientie, Sal artis, Sal vite* and *S. Scientia*, the product resulting from the sublimation of a mixture of corrosive sublimate and sal ammoniac. It is stimulant, but not employed.

ALÈNE (F.), *Alêne, Alaise, Lin'teum*, from αλέω, 'I preserve.' A guard. A cloth arranged in several folds, and placed upon a bed, so as to guard it from the lochia or other discharges.

ALETON, Farina.

ALETRIS, a. farinosa.

ALETRIS (Ph. U. S.). *A. Farinosa seu alba, Stargrass, Starwort, Blazing star, Aloe-root, Bitter-grass, Black root, Unicorn root, Ayue root, Ayue grass, Colic root, Devil's bit, Mealy starwort*, (F.) *Alétria*. *Nat. Ord. Asphodelææ. Sex. Syst. Hexandria Monogynia*. This plant is an intense and permanent bitter, and is used as a tonic and stomachic. It is common in the United States.

ALETRIS ALBA, *Aletris—a. Aurea*; indigenous—has similar virtues.

ALEURON, Farina.

ALEUROTESIS, see Cribration.

ALEXANDERS, *Smyrnum olusatrum*.

ALEXANDRI ANTIDOTUS AUREA. See Alexandrine.

ALEXAN'DRINE, *Emplas'trum Alexan'dri*. A garlic plaster, invented by Alexander, contemporary of Mesuë. Other ancient preparations were called 'Alexandrine,' as the *Alexan'dri antidotus aurea*, used in apoplexy; the *Collyrium siccum Alexandrinum*, or 'Collyrium of King Alexander,' mentioned by Aëtius.

ALEXICACUM, Amuletum, Alexipharmic.

ALEXIPHAR'MIC, *Alexiphar'macum, Anti-phar'macum, Alexic'cus, Caco-alcite'ria, Leri-phar'macum*, (F.) *Alexipharmaque*, from ἀλέξω, 'to repel,' and φάρμακον, 'poison.' A term formerly used for medicines which were considered proper for expelling from the body various morbid principles, or for preventing the bad effects of poisons taken inwardly.

ALEXPYRETICUS, Febrifuge.

ALEXIR, Elixir.

ALEXITE'RIA, *Cacalexite'ria*, from ἀλεξάσθαι, 'to assist.' Originally, *alexiterium* was used synonymously with remedy. In more modern times it has been applied to a class of medicines, that counteract poisons placed in contact with the exterior of the body, in contradistinction to alexipharmic.

ALEXITERIUM CHLORICUM, see Disinfection—a. Nitricum, see Disinfection.

ALÈZE, Aïse.

ALFUSA, Tutia.

ALTA MARINA, Pila marina.

ALGALIC, Catheter.

ALGALIE, Catheter.

AL'GAROTI, *Al'garot, Algaro'thi Pulvis, Pulvis Angel'icus, Oc'idum seu Submuri'as Stib'i*

præcipitan'do parat'um, Antimo'nii Oc'y'dum, Oc'idum antimo'nii Nitro-muriat'icum, Oc'idum Stib'i *Al'idum Muriat'ico oxyg'enat'o parat'um, Mercur'ius Vita, Mercur'ius Mortis, Flowers of Antimony*, (F.) *Oxyde d'Antimoine*, so called from Victor Algarothi, a Veronese physician; separated from the terchloride of antimony by pouring the terchloride into a large quantity of water. When this, an oxychloride, is washed with abundance of water to separate the muriatic acid, and afterwards by a solution of alkali to remove the terchloride, the *teroxide of antimony, Antimo'nii Oxidum* of the Edinburgh Pharmacopœia is left. It was formerly much used as an emetic, purgative, and diaphoretic.

ALGE'DO, from αλγος, 'pain.' Violent pain about the neck of the bladder, occasionally occurring in gonorrhœa.—Cockburn.

ALGEDON, Pain.

ALGEMA, Pain.

ALGESIS, Pain.

ALGETICUS, See Algos.

AL'GIDUS, from *algor*, 'cold.' (F.) *Algide*. That which is accompanied by coldness.

AL'GIDA FEBRIS, *Febris horrif'ica seu horrida seu quer'quera seu crym'des, Bry'cet'us, Bry'chet'us*. (F.) *Fèvre algide, Algid Fever*. A pernicious intermittent, accompanied by icy coldness, which is often fatal in the second or third paroxysm.

ALGOR, Rigor.

ALGOS, αλγος, 'pain.' See Pain. Hence, *Alget'icus*, 'painful,' as *Epilep'sia alget'ica*. The suffix *algia* has the same signification—as in *Cephalalgia, Pleuralgia, Neuralgia*, &c.

ALGOSPASMUS, from αλγος, 'pain,' and σπασμος, 'spasm.' Painful spasm or cramp of the muscles.

ALGUE COMMUNE, *Pila marina—a. Marine, Pila marina—a. des Verriers, Pila marina*.

ALHAGI, Agul.

ALHANDAL, see Cucumis colocynthis.

ALHASEF, Sudamina.

ALHAUNE, *Lawsonia inermis*.

ALIBILIS, Nutritious.

AL'ICA, *Hal'ica, Farina'rium, Chondrue*, from αλερε, 'to nourish.' A grain from which the ancients made their tisanes; supposed, by some, to have been the *Triticum spelta*. At times, it seems to have meant the tisanic itself.

AL'ICES, from αλέω, 'I sprinkle.' Spots which precede the eruption of small pox.

ALIENATIO, Anomalia—a. Mentis, Insanity.

ALIENATION, MENTAL, Insanity.

ALIENATUS, Insane.

ALIÉNÉ, Insane.

ALIÉNISTE, Psychiater.

ALIENUS, Delirious.

ALIFORMES MUSCULI, Pterygoid muscles.

ALIFORMIS, Alaris, Pterygoid.

ALIGULUS, Confection.

ALIMA, Aliment.

ALIMELLÆ, Parotid.

AL'IMENT, *Aliment'um, Al'ima, Harma'lia, Nutri'men, Nut'riens, Sustain'aculum, Ciba'rium, Broma, Comis'te, Cibus, Esca, Nutri'tus, Nutri'ment'um, Sitos, Trophê*. (Prov.) *Belly-timber*. (F.) *Aliment, Nourriture*, from αλερε, 'to nourish.' Food. Any substance which, if introduced into the system, is capable of nourishing it and repairing its losses.

The study of aliments forms one of the most important branches of hygiene. They are confined to the organized kingdom,—the mineral affording none.

As regards the immediate principles which

predominate in their composition, they have been classed, but imperfectly, as follows:—

TABLE OF ALIMENTS.

1. <i>Feculaceous.</i>	Wheat, barley, oats, rye, rice, Indian corn, potato, sago, peas, beans, &c.
2. <i>Mucilaginous.</i>	Carrot, salsify, beet, turnip, asparagus, cabbage, lettuce, artichoke, melon, &c.
3. <i>Saccharine.</i>	Sugar, fig, date, raisin, apricot, &c. Orange, currant, gooseberry, cherry, peach, strawberry, raspberry, mulberry, prune, pear, apple, sorrel, &c.
4. <i>Acidulous.</i>	Cocoa, olive, sweet almond, nut, walnut, animal fat, oil, butter, &c.
5. <i>Oleaginous and Fatty.</i>	Different kinds of milk, cheese.
6. <i>Cuscons.</i>	Tendon, aponeurosis, true skin.
7. <i>Gelatinous.</i>	cellular texture; very young animals.
8. <i>Albuminous.</i>	Brain, nerve, eggs, &c.
9. <i>Fibrous.</i>	Flesh and blood.

Dr. Prout has four great classes—the *aqueous*, *saccharine*, *oleaginous*, and *albuminous*:—Dr. Pereira, twelve—the *aqueous*, *mucilaginous* or *gummy*, *saccharine*, *amylaceous*, *lignous*, *pectinaceous*, *acidulous*, *alcoholic*, *oily* or *fatty*, *protinaceous*, *gelatinous*, and *saline*.

Liebig divides them into two classes:—the NITROGENIZED or PLASTIC ELEMENTS OF NUTRITION, in which he comprises *vegetable fibrin*, *vegetable albumen*, *vegetable casein*, *flesh* and *blood*; and the NON-NITROGENIZED ELEMENTS OF RESPIRATION, in which he comprises *fat*, *starch*, *gum*, *cane sugar*, *grape sugar*, *sugar of milk*, *pectin*, *bassorin*, *wine*, *beer* and *spirits*. The former alone, in his view, are inservient to the nutrition of organized tissue; the latter are burnt in respiration, and furnish heat.

The following simple arrangement is, perhaps, as little objectionable as any:—

- | | |
|-------------------------------|---|
| 1. Nitrogenized Aliments, | { Fibrinous (Glutinous?)
Albuminous.
Caseinuous.
Amylaceous.
Saccharine.
Oleaginous. |
| (Albuminous, of Prout.) | |
| | |
| 2. Non-nitrogenized Aliments, | |

The second division might be still farther simplified, inasmuch as amylaceous aliments are convertible into sugar during the digestive process; and, from both, oleaginous matter may be formed.

ALIMENTAL, Nutrimental.

ALIMENTARY, Nutrimental—a. Tube, Canal, alimentary.

ALIMENTATION, *Alimenta'tio*. The act of nourishing.

ALIMENTEUX, Nutrimental.

ALIMENTUM, Aliment, Pabulum.

ALIMOS, Glycyrrhiza.

ALINDE'SIS, from *αλινδομαι*, 'to be turned about.' A species of exercise, which consisted in rolling in the dust, after having been anointed with oil.—Hippocrates.

ALIPÆ'NOS, *Alipæ'num*, *Alipæ'nos*, from *α*, priv., and *λειπεναι*, 'to be fat.' An epithet formerly given to every external remedy, devoid of fat or moisture: such as powders.—Galen.

ALIPANTOS, Alipenos.

ALIP'TA, *Alip'tes*, from *αλειψω*, 'I anoint.' He who anointed the Athlete after bathing. The place where this was done was called *Alip'terium*.

ALIPTERIUM, see Alip'ta.

ALIP'TICA, same etymon. The part of ancient medicine, which treated ofunction as a means of preserving health.

ALISIER BLANC, *Cratægus aria*.

ALISMA, *A. plantago*—*a. lanceolata*, *A.*

plantago.

ALIS'MA PLANTA'GO, *Alisma*, *A. lanceolata* seu *graminifolia*, *Plantago aquatica*, *Water Plantain*, (F.) *Plantain d'Eau*. Ord. Alismaceæ. See, *Syst. Hexandria Polygynia*. The fresh root is acrid, and the dried leaves will vesiccate. The leaves have been proposed as substitutes for *Uva Ursi*.

ALITURA, Nutrition.

AL'KALE, *O'leum Gall'næ*. An ancient pharmaceutical name for pullets' fat.

ALKALESCENCE, Alcalescence.

ALKALI, see Alkali—a. *Ammoniacum causticum*, *Ammonia*—*a. Ammoniacum spirituosum*, *Spiritus ammoniac*—*a. Minerale nitratum*, *Soda*, nitrate of—*a. Minerale phosphoratum*, *Soda*, phosphate of—*a. Minerale salinum*, *Soda*, muriate of—*a. Vegetable*, *Potash*—*a. Vegetabile cum aceto*, *Potassa acetata*—*a. Vegetabile fixum causticum*, *Potassa fusa*—*a. Volatile*, *Ammonia*—*a. Volatile causticum*, *Ammonia*—*a. Volatile concrete*, *Ammonia carbonas*—*a. Volatile mild*, *Ammonia carbonas*—*a. Volatile nitratum*, *Ammonia nitrata*—*a. Volatile tartarizatum*, *Ammonia tartaras*—*a. Volatile vitriolatum*, *Ammonia sulphas*.

ALKANET, BASTARD, *Lithospermum officinale*—*a. Dyer's*, *Anchusa tinctoria*—*a. Garden*, *Anchusa officinalis*—*a. Officinal*, *Anchusa officinalis*.

ALKAR, Medicament.

ALKEKENGI, *Physalis*.

ALKER'MES, *Confectio Alker'mes*, *Alcher'mes*. A celebrated electuary, composed of a multitude of substances. It was so called from the grains of kermes contained in it. It was used as a stimulant. Also, kermes.

ALKERVA, see *Ricinus communis*.

ALKITRAN, *Cedria*.

ALKOL, Alcohol.

ALKOOL, Alcohol.

ALKYMISTRE, Alchymist.

ALLA, *Cerevisia*.

ALLAITEMENT, Lactation.

ALLAMAN'DA, *A. Cathartica* seu *grandiflora*, *Orelia grandiflora*, *Gall'aripa*, *Echlinus scandens*, *Apoc'ynum scandens*. A shrub, native of Guiana, the infusion of whose leaves is said by Linnæus to be useful in *Colica Pictorum*.

ALLANTODES, Allantois.

ALLAN'TOIC ACID, *Ac'idum allanto'icum*. A peculiar acid, found in the liquor of the allantois of the cow.

ALLANTOIDES, Allantois.

ALLAN'TOIS, *Allanto'id'es*, *Allanto'd'es*, *Membrana urina'ria*, *M. seu Tunica Farcimin'tis*, *M. Intestina'lis*, the *Allantoid Vesicle*, from *αλλας*, *αλλαντος*, 'a sausage,' and *αιδος*, 'shape.' A sort of elongated bladder, between the chorion and amnion of the fetus, which is thrown out from the caudal extremity of the embryo, and communicates with the bladder by the urachus. It is very apparent in quadrupeds, but not in the human species. As the allantois is developed, its walls become very vascular, and contain the ramifications of what become the umbilical artery and vein, which, by the elongation of the allantois, are brought through the villi of the chorion, into indirect communication with the vessels of the mother.

ALLANTOTOXICUM, from *αλλας*, *αλλαντος*, 'a sausage,' and *τοξικον*, 'a poison.' Sausage poison (G.) Wurstgift. The Germans have given this name to a poison developed in sausages formed of blood and liver.

ALLECTUARY, Electuarium.

ALLELUIA, *Oxalis acetosella*.

ALLEN'THESIS, from *αλλος*, 'another,' and *εισθεσις*, 'introduction.' The entrance into, or

presence of extraneous bodies in, the organism.—Walther.

ALLEVIATOR, from *ad*, 'to,' and *levare*, 'to raise;' 'a soother.' An instrument for raising invalids, invented by Mr. Jenks, of Rhode Island. It consists of two upright posts, about six feet high, each supported by a pedestal; of two horizontal bars at the top, rather longer than a common bedstead; of a windlass of the same length, placed six inches below the upper bar; of a cog-wheel and handle; of linen belts from six to twelve inches wide; of straps secured at one end of the windlass; and at the other having hooks attached to corresponding eyes in the linen belts, and of a head-piece made of netting. The patient lying on his mattress, the surgeon passes the linen belts beneath his body, attaching them to the hooks on the ends of the straps, and adjusting the whole at the proper distance and length, so as to balance the body exactly, and then raises it from the mattress by turning the handle of the windlass. To lower the patient again, and replace him on the mattress, the windlass must be reversed.

ALLGOOD, *Chenopodium bonus Henricus*.

ALLHEAL, *Heracleum spondylium*.

ALLIA'CEOUS, *Allia'ceus*, (F.) *Alliacé*, from *allium*, 'garlic.' Belonging to garlic, as *alliaceus* odour.

ALLI'IRE, *Alliaria*.

ALLIARIA, from *allium*, its smell resembling garlic. *A. officinalis*, *Erysimum alliaria* seu *cordifolium*, *Sisymbrium alliaria*, *Jack-in-the-hedge*, *Stinking hedge Mustard*, *Hedge Garlic*, *Sauce-alone*, *Hesperis alliaria*, (F.) *Alliaire*. *Ord.* Cruciferae. This plant has been sometimes given in humid asthma and dyspnoea. It is reputed to be diaphoretic, diuretic, and antiscorbutic.

The Parisian Codex has a compound syrup of *alliaria*, *Sirup d'érysimum composé*, which is used in hoarseness.

ALLIGATURA, *Fascia*, *Ligature*.

ALLIOTICUS, *Alterative*.

ALLIUM, (Ph. U. S.) from *oleo*, 'I smell.' *A. sativum*, *Theriaca rusticorum*, *Ampeloprasum*, *Scorodion*, *Scorodon*, *Garlic*, (F.) *Ail.* *Ord.* Liliaceae. *Sex. Syst.* Hexandria Monogynia. A native of Sicily, but cultivated for use. The bulbs or cloves, *Ag'lithes*, have a strong, offensive, and penetrating odour, and a sweetish, biting, and caustic taste. Internally, garlic is stimulant, diuretic, expectorant, emmenagogue(?), diaphoretic, and anthelmintic. Externally, it is rubefacient, maturative, and repellent; and a *garlic ointment* is made by digesting, at a moderate heat, for half an hour, two or three cloves of fresh garlic in an ounce of lard, and straining. It is applied as a rubefacient to the chest in croup.

Dose, one to six cloves, swallowed whole, or from fʒss to fʒij of the juice.

Taylor's Remedy for Deafness, a nostrum, appears to consist of garlic, infused in oil of almonds, and coloured by *alkanet root*.

ALLIUM ASCALONICUM, *Échalotte*.

ALLIUM CANADENSE, *Meadow garlic*; indigenous; has the same properties as *allium*.

ALLIUM CEPA, *Cepa Vulgaris*, *Common Onion*, (*Cepula*, *Crom'myon*, (F.) *Oignon*. Acid and stimulating, and possessing very little nutriment. Onions have been used as stimulants, diuretics, and anthelmintics. The boiled or roasted onion, as a cataplasm, is emollient and maturating. The fresh root is rubefacient. The expressed juice is sometimes used in otalgia and in rheumatism.

ALLIUM GALLICUM, *Portulaca*—a. *Plantaginum*, *A. Victoriale*.

ALLIUM PORRUM, *Porrum*, *P. sativum*, *Prasum*, the *Leek* or *Porret*; (F.) *Poireau*, *Porreau*. It possesses the same property as the onion.

The virtues of the genus *Allium* depend upon an acid principle, soluble in water, alcohol, acids, and alkalies.

ALLIUM REDOLENS, *Teucrium scordium*.

ALLIUM VICTORIA'LE, *A. plantagin'eum*, *Cepa victorialis*, *Victoria'lis longa*. The root, which, when dried, loses its alliaceous smell and taste, is said to be efficacious in allaying the abdominal spasms of pregnant women (?).

ALLOCHET'IA, *Allotrichet'ia*, from *allos*, 'another,' and *χῆσαν*, 'to go to stool.' The discharge of extraneous matters from the bowels. The discharge of faeces by an abnormal opening.

ALLOCHOSOS, *Delirious*.

ALLOCHROMA'SIA, from *allos*, 'another,' and *χρῶμα*, 'colour.' A change of colour.

ALLOCEPATHIA, *Allopathy*.

ALLOCEPATHIC, *Allopathic*.

ALLOEOSIS, *Alteration*.

ALLOEOTICUS, *Alterative*.

ALLOIOSIS, *Alteration*.

ALLOIOTICUS, *Alterative*.

ALLONGEMENT, *Elongation*.

ALLOPATH, *Allopathist*.

ALLOPATHES, *Allopathic*.

ALLOPATH'IC, *Allopath'icus*, *Allocepath'ic*, *Allocepath'icus*, *Allopathes*, *Heteropath'ic*, from *allos*, 'another,' and *παθος*, 'affection.' Relating to the ordinary method of medical practice, in contradistinction to the homœopathic. See *Allopathy*.

ALLOPATHIE, *Allopathy*.

ALLOP'ATHIST, *Al'lopath*, same etymon. One who follows allopathy.

ALLOP'ATHY, *Allopath'ia*, *Allocepath'ia*, *Hyperantio'sia*, *Hyperantio'ma*, *Cura'tio contrarior'um per contra'ria*, (F.) *Allopathie*; same etymon. The opposite to homœopathy; or, according to Hahnemann, a method of treatment in which remedial agents are employed, the action of which, on healthy man, produces morbid phenomena different from those that are observed in the sick person. The ordinary medical practice is so designated by the homœopathist.

ALLOPHASIS, *Delirium*.

ALLOTRIOCHET'IA, *Allochetia*.

ALLOTRIODON'TIA, from *allos*, 'foreign,' and *οδους*, 'a tooth.' Transplantation of teeth.

ALLOTRIOEC'CRISIS, from *allos*, 'foreign,' and *εκκρισις*, 'separation.' The separation of extraneous matters from the body in disease.

ALLOTRIOGEUSTIA, *Parageusia*.

ALLOTRIOPHAGIA, *Malacia*.

ALLOTRIOTEX'IS, from *allos*, 'foreign,' and *τεξις*, 'parturition.' The bringing forth of an abnormal foetus.

ALLOTRIU'RIA, from *allos*, 'foreign,' and *ουρον*, 'urine.' Admixture of foreign matters with the urine.

ALLOTROPISM; (F.) *Allotropie*, from *allos*, 'another,' and *τροπος*, 'a turn or change.' A term recently introduced into chemistry, the object of which is to express the property, possessed by certain simple bodies, of assuming different qualities on being subjected to certain modes of treatment. Carbon, for example, furnishes three forms—plumbago, charcoal, and diamond.

ALL-OVERISH, see *Indisposition*.

ALLSPICE, see *Myrtus pimenta*—a. *Bush*, *Laurus Benzoin*—a. *Carolina*, *Calycanthus*—a. *Wild*, *Laurus Benzoin*.

ALLUCINATIO, *Hallucination*.

ALLURE, *Influenza*.

ALLUS POLLEX, *Pollex*, see *Digitus*.

ALMA, Water.

ALMARIAB, see *Plumbi oxydum semivitreum*.

ALMEZERION, *Cnecorum tricoceum*.

ALMOND, *Amygdala*.

ALMOND BLOOM. A liquid cosmetic, formed of *Brazil dust* $\mathfrak{z}\text{ij}$, water Oij ; boil and strain; and add *isinglass* $\mathfrak{z}\text{vj}$, *grana sylvestria* $\mathfrak{z}\text{ij}$, or *cinnamon* $\mathfrak{z}\text{ij}$, alum $\mathfrak{z}\text{ij}$, borax $\mathfrak{z}\text{ij}$; boil again, and strain through a fine cloth.

ALMOND CAKE, see *Amygdala*—a. of the Ear, Tonsil—a. Earth, *Arachis hypogaea*—a. Paste, see *Amygdala*—a. Powder, see *Amygdala*—a. of the Throat, Tonsil.

ALNUS, *A. glutinosa*—a. *Communis*, *A. glutinosa*.

ALNUS GLUTINOSA, *Alnus*, *A. communis*, *Betula glutinosa* seu *emarginata*, *Europea* an *Alder*, (Sc.) *Arn*, *Eller*, *Ord*. *Betulaceæ*. A tree which grows in Europe, in moist places. The bark and leaves are astringent and bitter; and hence are employed in intermittents, and as a tonic and astringent.

ALNUS SERRATULA, *American Alder*, has similar properties.

ALNUS NIGRA, *Rhamnus frangula*.

ALOCHIA, from *a*, privative, and *λοχεια*, 'lochia.' Absence of the lochial discharge.

ALOEDARIUM. A compound medicine, containing aloes.—*Gorræus*.

ALOE. Aloes—a. *False*, *Agave Virginica*.

ALOE ROOT, *Aletris farinosa*.

AL'OE'S, *Al'oë*, *Fel Natu'ra*. The inspissated juice of the *Aloe*. *Ord*. *Asphodelææ*. *Sex. Syst.* *Hexandria Monogynia*.

ALOES BARBADENSIS, *A. hepatica*—a. *Bombay*, *A. hepatica*—a. *des Barbades*, *A. hepatica*.

ALOES CABBALLINA, *A. Guinien'sis*, *Horse-aloes*. Used chiefly for horses. It is collected in Spain and Portugal, and is very coarse.

ALOES EN CALÉBASSES, *A. hepatica*.

ALOES, CAPE, *Shining Aloes*; a cheap and excellent form of aloes, collected at the Cape of Good Hope, from *Aloe ferax*, *A. Africana*, *A. spicata*, and other species.

ALOES, EAST INDIA, *A. Succotorina*—a. *Guinensis*, *A. Caballina*.

ALOES HEPATICA, *A. vulgaria*, *A. Barbaden'sis*, *Hepatic aloes*, *Bombay aloes*, *Barbadoes aloes*, *A. vulgaria extractum*, (F.) *Aloes en calébasses*, *A. des Barbades*. This species has a very disagreeable odour, and an intensely bitter and nauseous taste. Properties the same as the last.

ALOES, HORSE, *A. Caballina*—a. *Lucida*, *A. Succotorina*—a. *Socetrine*, *A. Succotorina*—a. *Spicata extractum*, *A. Succotorina*.

ALOES SUCCOTORINA, *Socetrine aloes*, *Turkey aloes*, *East India aloes*, *Al'oës lu'cida*, *A. Zoctorina*, *A. spicata extractum*, *An'ima Aloës* is the best species. Its odour is not unpleasant; taste very bitter, and slightly aromatic; colour reddish-brown, with a shade of purple; mass hard, friable; fracture conchoidal and glossy; soluble in dilute alcohol. Powder of a bright cinnamon-yellow colour. It is cathartic, warm, and stimulating; emmenagogue, antelmintic, and stomachic. As a cathartic, it affects the rectum chiefly. *Dose*, as a cathartic, gr. v to $\mathfrak{z}\text{ij}$ in pill.

ALOES, TURKEY, *A. Succotorina*—a. *Vulgaris*, *A. hepatica*—a. *Wood*, *Agallochum*—a. *Zoctorina*, *A. Succotorina*.

ALOËTIC, *Al'oëticus*. A preparation which contains aloes.

ALOEXYLON, *Agallochum*.

ALOGOTROPHIA, from *alogos*, 'disproportionate,' and *τροφή*, 'nutrition.' Irregular nutrition. Used particularly to designate the irregular manner in which the nutrition of bones is effected in rickety individuals.

ALOPECES, *Psoræ*.

ALOPEC'IA, from *αλωπηξ*, 'a fox;' (this animal being said to be subject to the affection.) *Capilla'ria deflu'vium*, *Athria dep'ilis*, *Phalacro'tis*, *Depilatio*, *Triph'o'sis* *Atheris*, *Gangra'na Alopec'ia*, *Atrich'ia*, *Deflu'vium seu Lapsus seu Fluxus Pilo'rum*, *Lipotrich'ia*, *Vulpis morbus*, *Pela'da*, *Baldness*, (F.) *Pelade*. Falling off of the hair; loss of the hair. When this is confined to the crown of the head, it is called *calvities*, although the terms are often used synonymously. When congenital, it is called *Atrich'ia seu Alopec'ia adna'ta*. The falling off of the hair in old age is termed *Alopec'ia seni'lis*.

ALOPECIA ADNATA, see *Alopecia*—a. *Areata*, *Porrigi decalvans*—a. *Circumscripta*, *Porrigi decalvans*—a. *Partialis*, *Porrigi decalvans*—a. *Senilis*, see *Alopecia*.

ALOUCHE, *Cratægus aria*.

ALOUCHI. The name of a gum procured from the *canella alba* tree.

ALOUCHIER, *Cratægus aria*.

ALPAM. A shrub which grows on the coast of Malabar. Certain parts of this, infused in oil, form an antispore ointment. The juice of the leaves, mixed with that of *calamus*, is employed against the bites of serpents.

ALPHENIC, *Saccharum candidum*.

ALPHITEDON, see *Fracture*.

ALPHITON, *αλφίτον*, *Polenta*, *Fari'na*. Any kind of meal. Toasted barley-meal.—*Hippocrates*. *Polenta* means, also, a food composed of Indian meal, cheese, &c. See *Farina*.

ALPHON'SIN, *Alphon'sinum*. A kind of bullet forceps, similar to a *Porte-crayon*, so-called from the inventor, *Alphonso Ferri*, of Naples.—*Scultetus*.

ALPHOS, *Lepra alphoides*.

ALPHOSIS ÆTHIOPICA, see *Albino*.

ALPINIA CARDAMOMUM, *Amomum cardamomum*—a. *Exaltata*, *Renealmia exaltata*—a. *Tubulata*, *Renealmia exaltata*—a. *Galanga*, *Maranta galanga*.

ALPISTE, *Phalaris Canadensis*.

ALSANDERS, *Smyrnum olusatrum*.

ALSINÉ ME'DIA, *A. avicula'rum* seu *vulgaris*, from *αλος*, 'a grove,' because growing abundantly in the woods. [?] *Morsus Galli'na*, *Holostæum Alsin'e*, *Stella'ria me'dia*, *Mouse-ear*, *Chickweed*, *Chickenswort*, (F.) *Mouron des Oiseaux*, *Morgeline*. *Ord*. *Caryophyllacææ*. This plant, if boiled tender, may be eaten like spinach, and forms an excellent emollient poultice. It was formerly regarded as a vulnerary and detergent.

ALTAFOR, *Camphor*.

ALTER SEXUS, *Sex*, *female*.

ALTERANS, *Alterative*.

ALTÉRANT, *Alterative*; see, also, *Alteration*.

ALTERATION, *Alteratio*, from *alter*, 'other,' *Alloio'sis*, *Alloio'sis*. This word is used in France to express any change which supervenes; for example, in the expression of the countenance (*altération de la face*), or in the structure of an organ (*altération organique*), or in the nature of fluids excreted (*altération de l'urine*, *des larmes*, *du lait*, &c.)

Alteration is also used in an entirely different sense, to express intense thirst in disease. In this case its etymology is different. It comes from *haléter*, and was formerly written *halteration*.

ALTERATIVE. *Alterans*, *Alloio'icus*, *Alloio'icus*, *Immutans*. An agent considered to be capable of producing a salutary change in a disease, but without exciting any sensible evacuation. As medicine improves, this uncertain class of remedies becomes, of necessity, diminished in number. See *Eutrophie*.

(F.) *Altérant*. The French term likewise

means, that which causes thirst, — *Siticulo'sus*, *Dipsæticus*, as *alter* means both to change, and to cause thirst. *S'alter* is to experience a change for the worse, — (*corrupti*.)

ALTERCANGENON, Hyoscyamus.

ALTERCUM, Hyoscyamus.

ALTERED, see Castratus.

ALTHÆA, from *αλθεω*, 'to heal.' *A. officinalis*, *Malvæ'scum*, *Aristalthæa*, *Hibis'cus*, *Hibiscum*, *Hibiscus*, *Hibis'cha* *mismal'ca*, *Bismal'ca*, *Marsh mallon*, (Prov.) *Malice*, (F.) *Gummauve*. Ord. *Malvacæ*. Sex. Syst. *Monadelphia Polyandria*. The leaves, *Althæa fo'lia*, and root, *Althæa rudi's*, contain much mucilage. They are emollient and demulcent, and are employed wherever medicines, possessing such properties, are required. In the Ph. U. S., *Althæa* is the root of *Althæa officinalis*.

ALTHANATHA, Orpiment.

ALTHEUS, Physician.

ALTHAXIS, Curation.

ALTHOS, Medicament.

ALTILIBAT, Terebinthina.

ALU'DEL, *Alu'tel*, *Vitrum sublimato'rium*. A hollow sphere of stone, glass, or earthen ware, with a short neck projecting at each end, by means of which one glass might be set upon the other. The uppermost had no aperture at the top. Aludels were formerly used in the sublimation of various substances.

A'LULA, diminutive of *ala*, 'a wing.' A little wing.

ALUM, Symphytum — a. Cataplasm, Coagulum aluminosum — a. Egyptian, *Egyptia stypteria*.

ALUM, Roche, *Alu'men de Rochi*, (F.) *Alun de Roche*. So called from *Roccha* in Syria, where there was a manufactory of it. It is in pieces of the size of an almond, covered with a reddish efflorescence.

Common Roche Alum, *A. Rochi Gallis*. Fragments of common alum moistened and shaken with prepared bole. It is white when broken.

ALUM ROOT, *Geranium maculatum*, *Heuchera cortusa*.

ALUM, SOLUTION OF, COMPOUND, *Liquor aluminis compositus*.

ALUM SPRINGS OF VIRGINIA, see Virginia, mineral waters of.

ALUMEN, (an Arabic term, *alum*.) *Alum*, *Hypersul'phas alu'mina et Potas'sæ*, *Potas'sæ alu'mino-sulphas*, *Sul'phas Aluminæ Acid'ulus cum Potas'sæ*, *Sulphas Alu'minae*, *Sul'phas Kal'ico-alumin'icum*, *Sulphas aluminæ'ris*, *Supersul'phas alu'minae et potas'sæ*, *Argill'a sulphu'rica alu'al'ita*, *A. vitriola'ta*, *Sypte'ria*, *Sulpersul'phas Argill'æ alcalisa'tum*, *Argilla Kali'sulphu'rica*. (F.) *Alun*.

ALUMEN CATINUM, Potash of commerce — a. Fixum, see Potash — a. Kinosatum, *Polvis sulphatis aluminæ compositus*.

ALUMEN COMMUNE, Common alum, *English alum*, *Rock alum*, *Alumen facti'tium*, *A. crystal'linum*, *A. ru'peum*, (F.) *Alun d'Angleterre*, is the variety usually employed. It is in octahedral crystals, but generally in large, white, semitransparent masses; has a sweetish, styptic taste; effloresces in the air, and is soluble in 16 parts of water at 60°. It is tonic and astringent, and as such is used internally and externally. Dose, gr. v to xv.

ALUMEN EXSICCATUM, *Alu'men ustum* seu *sicca'tum* seu *calcina'tum*, *Sulphas alu'minae fusus*, *Argill'a sulphu'rica usta*, *Burnt alum*, *dried alum*. (F.) *Alun calciné*, (*Alum* melted in an earthen vessel until ebullition ceases.) Escharotic.

ALUMEN ROMANUM, *Roman alum*, *A. Ru'tinum* seu *Rubrum*. (F.) *Alun de Rome*. In crystals, which are of a pale red when broken, and covered with a reddish efflorescence.

ALUMEN SICCATUM, *Alumen exsiccatum*.

ALUMINA, ACETATE OF, *Aluminæ Acetas* — a. *Depurata*, *Argilla pura* — a. *Pura*, *Argilla pura* — a. *Sulphate of Aluminæ Sulphas*.

ALUMINÆ ACETAS, *Argill'æ Ac'etas*, *Ac'etate of Alu'mina*. A deliquescent salt, obtained by the addition of acetate of lead to sulphate of alumina and potassa. It possesses the same properties as the sulphate of alumina.

ALUMINÆ ET POTASSÆ HYPERSULPHAS, *Alumen* — a. et *Potassæ supersulphas*, *Alumen* — a. *Sulphas*, *Alumen*.

ALUMINÆ SULPHAS, *Argillæ Sulphas*, *Sulphate of Alu'mina*. Simple sulphate of alumina may be made by the direct combination of alumina and sulphuric acid, and contains 30 per cent. of the former, to 70 per cent. of the latter. It is a deliquescent salt; and is an excellent antiseptic and detergent for ulcers. It is chiefly used to preserve dead bodies — a strong solution being injected into the arteries.

Gannal's Solution for preserving animal substances is made by dissolving an ounce of acetate of alumina in twenty ounces of water.

ALUMINÆ SULPHAS ACIDULUS cum POTASSÆ, *Alumen* — a. *Sulphas fusus*, *Alumen exsiccatum*.

ALUMINE FACTICE, *Argilla pura*.

ALUMINII OXIDUM, *Argilla pura*.

ALUN, *Alumen*.

ALUNSEL, *Gutta*.

ALUS, *Symphytum*.

ALUSIA, Hallucination — a. *Hypochondriasis*, *Hypochondriasis*.

ALUTEM, *Aludel*.

ALVAQUILLA, *Psoralea glandulosa*.

ALVARAS NIGRA, *Ichthyosis*.

ALVEARIUM, *Auditory canal*, external.

ALVEOLAR, *Alveola'ris*, from *alveus*, 'a cavity.' (F.) *Alvéolaire*. That which relates to the alveoli.

ALVEOLAR ABSCESS, *Parulis*.

ALVEOLAR ARCHES, (F.) *Arcades alvéolaires*, are formed by the margins or borders, *Juga alveola'ria*, of the two jaws, which are hollowed by the Alveoli.

ALVEOLAR ARTERY, *Supra-maxillary A.*, *Ar'tère sus-maxillaire* of *Chaussier*, arises from the internal maxillary, descends behind the tuberosity of the upper jaw, and gives branches to the upper molar teeth, gums, periosteum, membrane of the maxillary sinus, and buccinator muscle.

ALVEOLAR BORDER, *Limbus alveola'ris*. The part of the jaws that is hollowed by the alveoli.

ALVEOLAR MEMBRANES are very fine membranes, situate between the teeth and alveoli, and formed by a portion of the sac or follicle which enclosed the tooth before it pierced the gum. By some this membrane has been called the *alveolo-dental periosteum*.

ALVEOLAR VEIN. This has a similar distribution with the artery.

ALVÉOLE, *Alveolus*.

ALVEOLI DENTIS, see *Alveolus*.

ALVEOLO-DENTAL PERIOSTEUM, see *Alveolar membranes*.

ALVÉOLO-LABIAL, *Buccinator*.

ALVEOLUS, same etymon. *Bo'trion*, *Bo'thrion*, *Odontoboth'rium*, *Odontopha'nè*, *Frena* (*Alve'oli*), *Mortariolum*, *Ho'micos*, *Præsepium*, *Phatnè*, *Phat'nion*, *Præsep'ium*, *Patnè*, *Pathnè*, (F.) *Alvéole*. The alveoli are the sockets of the teeth, *Alve'oli dentis*, *Ma'nia* seu *Caver'nae dent'ium*, into which they are, as it were, driven. Their size and shape are determined by the teeth which they receive, and they are pierced at the apex by small holes, which give passage to the dental vessels and nerves.

ALVEUS, Auge — a. *Ampullosus*, *Receptaculum chyli* — a. *Ampullesens*, *Thoracic duct* — a.

Communis: see Semicircular canals—a. Utriculus: see Semicircular canals.

ALVI EXCRETIO, Defecation—a. Fluxus aquosus, Diarrhoea—a. Laxitas, Diarrhoea—a. Profluvium, Diarrhoea.

ALVIDUCUS, Laxative.

ALVINE, *Alvīnus*, from *alvus*, 'the abdomen.' That which relates to the lower belly, as *alvine dejections*, *alvine flux*, *alvine obstructions*, &c.

ALVUM EVACUANS, Cathartic.

ALVUS, Abdomen, Uterus—a. Adstricta seu Astricta, Constipation—a. Cita, Diarrhoea—a. Dura, Constipation—a. Renum, Pelvis of the kidney—a. Tarda, Constipation—a. Viridis, Dejection.

ALYCE, Anxiety.

ALYMPH'IA, from *a*, priv., and *lymphā*, 'lymph.' Want of lymph.

AL'YPON, from *a*, priv., and *λυπη*, 'pain.' An acrid, purging plant, described by Matthioli. By some it has been supposed to be the *Globularia alypum* of botanists.

ALYSIS, Anxiety.

ALYSMUS, Anxiety.

ALYSSUM PLINII, Galium mollugo.

ALYSSUS, Antihydrophobic.

ALYX'IA STELLA'TA, *A. aromatica*, (F.)

Alyce aromatique. A plant, of the family Apocynaceæ, the bark of which resembles canella alba, and is used in Batavia in pernicious fevers.

AL'ZILAT. In some of the Arabian writers, a weight of three grains.—Ruland and Johnson.

AMABLE, Lacuna Labii Superioris.

AMADOU, Boletus igniarius.

AMADOUVIER, Boletus igniarius.

AMAGRIR, Emaciate.

AMAGRISSEMENT, Emaciation.

AMAND (SAINT), MINERAL WATERS OF, Saint-Amand-les-Eaux. These springs, in the department Nord, France, are thermal (77° Fahr.), carbonated chalybeates; and are much used internally, or in the form of *boves*, in chronic affections of the digestive organs and liver, and in rheumatism.

AMANDEN, see Amygdala.

AMANI'TÆ, from *a*, privative, and *mania*, 'madness' i. e. 'not poisonous.' (?) A name given, by the Greeks and Romans, to the edible *champignons*. *Amanita* forms, at the present day, a genus, some of which are edible, others poisonous. Amongst others, it contains the *Agaricus aurantiacus* and *A. pseudo-aurantiacus*.

AMARA DULCIS, Solanum dulcamara.

AMARAC'NUM. An ancient and esteemed plaster, containing several aromatics, the marjoram, *aparakos*, in particular.

AMARACUS, Origanum majorana—a. Tomentosus, Origanum dictamnus.

AMARITIES, Bitterness.

AMARITUDO, Bitterness.

AMAROR, Bitterness.

AMARUCACHU, Polyanthus tuberosa.

AMAR'US, *Picros*, 'bitter.' (F.) *Amēr*. The bitter principle of vegetables is the great natural tonic, and hence *bitters*, as they are termed collectively, belong to the class of tonics. Several are used in medicine; the chief are, gentian, quassia, cinchona, calumba, dog-wood, &c.

AMASE'SIS, *Amasse'sis*, from *a*, privative, and *μασσειν*, 'mastication.' Mastication when impeded or impracticable.

AMATORIUM, Lacuna labii superioris.

AMATORII, Oblique muscles of the eye.

AMATORIUM VENEFICUM, Philiter.

AMATORIUS MUSCULUS, Obliquus superior oculi.

AMAURO'SIS, *Obfusca'tio*, *Offusca'tio*, from *αυρος*, 'obscure,' Drop serena, Gutta serena, Cataracta nigra, Parop'sis amauro'sis, Immo-

itas pupill'æ. Suffusio nigra, Black cat'aract, Nervous blindness, (F.) Goutte-serene, Cataracte noire, Amauotocervie (Piorry). Diminution, or complete loss of sight, without any perceptible alteration in the organization of the eye: generally, perhaps, owing to loss of power of the optic nerve or retina—*Optic anæsthesia*. Counter-irritants are the most successful remedial agents, although the disease is always very difficult of removal, and generally totally incurable.

AMAUROSIS DIMIDIATA, Hemipopia—a. Diurnal, Nyctalopia—a. Nocturnal, Hemeralopia—a. Imperfecta, Hypo-amaurosis.

AMAUROTIC, *Amauroticus*, (F.) *Amauroticus*; same etymon. Affected with amaurosis.

AMAUROTIC CAT'S EYE, *Galeamauro'sis*. A name given by Beer to an amaurotic affection, accompanied by a remarkable change of colour in the pupil, which presents, apparently in the fundus of the eye, a lighter tint, yellowish or brownish yellow, instead of its natural clear black.

AMA'ZIA, from *a*, privative, and *μαστος*, 'breast.' A monstrosity, in which there is absence of one or both breasts.

AMBARUM, Ambergis—a. Cineritium, Ambergis.

AMBE, from *αμβαινω*, 'I ascend,' *Ambi*. A superficial eminence on a bone. Also, an old surgical machine for reducing dislocations of the shoulder; the invention of which is ascribed to Hippocrates. It is no longer used.—Hippocrates, Scultetus. See Crista.

AMBER, Succinum—a. Liquid: see Liquid-amber styraciflua.

AMBERGRIS, 'gray amber,' *Ambor*, *Ambra*, *Ambra gri'æa* seu *cineara'cea* seu *ambrosina*, *Ambarum*, *Succinum cine'reum* seu *gri'seum*, *Ambarum cineri'tium*. A concrete substance, of the consistence of wax, cineritious colour, studded with yellow and blackish spots, and exhaling a very pleasant odour. It seems highly probable that ambergris is formed in the intestines of the whale, and voided with its excrement. Like all aromatic substances, ambergris is slightly anti-spasmodic and excitant; but it is oftener employed as a perfume than as a medicine.

AMBIA. A liquid, yellow bitumen, the smell and virtues of which are similar to those of the resin *tacamahaca*. It is obtained from a spring in India.

AMBICUS, Alembic.

AMBIDEXTER, *Amphidex'ius*, from *ambo*, 'both,' and *dexter*, 'right.' One who uses both hands with equal facility. Celsus says the surgeon ought to be '*non minus sinistra quam dextra promptus*.' One of the aphorisms of Hippocrates says, that a woman is never ambidexter. This is a mistake.

AMBILÆVUS, Ampharisteros.

AMBIPIA, Diplopia.

AMBITUS GENITALIS MULIEBRIS, Ves-tibulum.

AMELOMA, Abortion.

AMBLOSIS, Abortion.

AMBLOSISMUS, Abortion.

AMBLOTHRIDION, see Abortion.

AMBLOTHRIDIUM, Abortive.

AMBLOTICUS, Abortive.

AMBLUS, *αμβλυσ*, 'obscure.' Hence, AMBLYAPI'IA, from *αμβλυσ*, 'obscure,' and *αἴη*, 'feeling.' Dulness of the sense of touch.

AMBLYOGMOS, Amblyopia.

AMBLYOPIA, from *αμβλυσ*, 'obscure,' and *οψ*, 'the eye.' *Amblyop'mos*, *Amblyog'mos*, *Amblyopia* (so called by some, according to Castelli, ob ignorantiam Græcæ linguæ), *Hebetudo visus*,

Feebleness of sight, (F.) Amblyopie, Vue faible.
First degree of Amaurosis.—Hippocrates.

AMBLYOPIA CREPUSCULARIS, Hemeralopia—a. Dissitorum, Myopia—a. Meridiana, Nyctalopia—a. Proximorum, Presbytia.

AMBLYOSMOS, Amblyopia.

AMBOLICUS, Abortive.

AMBON, ἀμβων, 'the raised rim of a shield or dish,' from ἀμβαινω, 'I ascend.' The fibro-cartilaginous rings or *bourrelets*, which surround the articular cavities, as the glenoid cavity of the scapula, the acetabulum, &c., have been so called—Galen. See Crista.

AMBOR, Ambergris.

AMBRA, Succinum—a. Ambrosiaca, Ambergris—a. Cineracea, Ambergris.

AMBAGRISEA, Ambergris.

AMBRE BLANC, Succinum (album)—a. Jaune, Succinum.

AMBRETTE, Hibiscus abelmoschus.

AMBRINA AMBROSIOIDES, Chenopodium ambrosioides—a. Anthelmintica, Chenopodium anthelminticum—a. Botrys, Chenopodium botrys.

AMEROISIE DU MEXIQUE, Chenopodium ambrosioides.

AMBROSIA, from α, privative, and βροτος, 'mortal.' Food which makes immortal, or the food of immortals. Food of the gods—Homcr. See also, Chenopodium botrys.

AMBROSIA ELATIOR, see A. Trifida.

AMBROSIA MARITIMA, Ord. Compositæ. A plant which grows on the shores of the Levant, and has a pleasant, bitter and aromatic taste. It is given in infusion, as a tonic and antispasmodic.

AMBROSIA TRIFIDA, *Horseweed, Richweed, Horsemint, Horsecane, Bitterweed, Great or Tall Ragweed, Wild Hemp.* This indigenous plant is found in low grounds and along streams, from Canada to Georgia, and west to Louisiana and Arkansas. It is an annual, and flowers in August and September. An infusion has been recommended locally in mercurial salivation.

Ambrosia elatior, Ragweed, is said by Dr. R. E. Griffith to have much more developed sensible properties.

AMBROSIE DU MEXIQUE, Chenopodium ambrosioides.

AMBULANCE (F.), from *ambulare*, 'to walk.' A military hospital attached to an army, and moving, along with it. Also called *Hôpital ambulant*.

AMBULATIO, Walking.

AMBULATORY, *Ambulans, Ambulativus, Ambulative*, (F.) *Ambulant*. A morbid affection is said to be 'ambulatory,' (F.) *ambulant*, when it skips from one part to another; as *Erysipèles ambulans*, &c. When blisters are applied successively on different parts of the body, they are called *Vésicatoires ambulants*.

AMBULEIA, Cichorium intybus.

AM'BULI. The Brachmanic name for an Indian aquatic herb, which appears to belong to the family *Lysimachia*. The whole plant has a sweet smell. Its decoction has a very bitter taste, and is an excellent febrifuge. It is also taken in milk in cases of vertigo.

AMBULO FLATULENTUS ET FURIOSUS, *Flatus furiosus, Vareni*. Painful, mobile, and periodical tumours affecting different parts, which were once considered as the effect of very subtle vapours—Michaelis. Their nature is by no means clear.

AMBUSTIO, Burn.

AMBTUA, Pareira brava.

AMBUYA-EMBO. A very beautiful, creeping aristolochia of Brazil, the decoction of which is exhibited successfully in obstructions. It is also used in fumigation and in baths as a tonic.

AME, Anima.

AMELI. A Malabar shrub, belonging to a genus unknown. The decoction of its leaves is said to relieve colic. Its roots, boiled in oil, are used to repel tumours.

AMELIA, Apathy.

AMENIA, Amenorrhœa, Emmenagogues.

AMENOMA'NIA. A hybrid word, formed from the Latin *amœnus*, 'agreeable,' and *mania*, 'mania.' A gay form of insanity.

AMENORRHÆA, *Parame'nia obstructio'nis, Menoceryph'ia, Menosta'sia, Apophras'is, Archa'ia, Defectus seu Reman'sio seu Cessatio men'sium, Menstruatio impedi'ta, Ischome'nia, Ame'nia, Metrohæ'mia*, from α, privative, *μην*, 'a month,' and *ρην*, 'I flow.' *Suppression of the menses*, (F.) *Suppression du flux menstruel*. This suppression is most commonly symptomatic, and hence the chief attention must be paid to the cause. Usually, there is an atonic state of the system generally, and hence chalybeates and other tonics are advisable.

Two great varieties of Amenorrhœa are commonly reckoned. 1. *A. Emansio'nis, Emansio men'sium, Menis'chesis, Menos'chesis, Menstruatio retenti'a, Men'sium retenti'o, Retention of the menses*, when the menses do not appear at the usual age: and, 2. *Suppres'sio Men'sium seu Menstruatio'nis, Amenorrhæ'a Suppres'sio'nis, Interrup'tio menstruatio'nis, Menstruatio'nis suppres'sa*, in which the catamenia are obstructed in their regular periods of recurrence. See *Emansio Mensium*, and *Menses*.

AMENORRHÆA DIFFICILIS, Dysmenorrhœa—a. *Emansio*, see Amenorrhœa—a. *Hymenica*, see *Hymenicus*—a. *Partialis*, Dysmenorrhœa—a. *Suppressionis*, see Amenorrhœa.

AMENTIA, Dementia: see, also, *Fatuitas*, and *Idiotism*—a. *Senilis*, Dementia of the aged.

AMER, *Amarus*—a. *de Bœuf*, Bile of the ox.

AMERICAN, see *Homo*.

AMERICAN TUBEROSUM, *Solanum tuberosum*.

AMERTUME, Bitterness.

AM'ETHYST, *Amethys'tus*, from α, privative, and *μεθω*, 'I am drunk.' A precious stone, to which the ancients attributed the property of preventing drunkenness. It was also used as an anti-diarrhœic and absorbent.—Pliny, Albertus Magnus.

AMETH'YSUM, *Amethys'tum, (remedium.)* Same etymon as the last. A remedy for drunkenness.

AMETRIA, Intemperance, from α, privative, and *μετρον*, 'measure.' Also, absence of the uterus; from α, privative, and *μητρα*, 'the uterus.'

AMETROHÆMIA, Amenorrhœa.

AMANTHUM MUSCÆTOX'ICUM, (*αμαντος*, 'pure,' and *αλδος*, 'a flower,') *Melan'thium muscætox'icum, Helo'nias erythrospërma*, 'Fly-poison, Fall-poison,' indigenous. Ord. *Melanthaceæ*. A narcotic poison, not used, however.

AMICULUM, Amnion.

AMIDOLIQUE, from *amidum*, 'starch.' Appertaining or belonging to starch. M. Bérall calls *médicaments amidoliques* those, which owe their general properties to starch.

AMIDON, Amylum—a. *Iodure d'*, Starch, Iodide of.

AMIDONNIÈRE, *Ammi maculatum*.

AMIDUM, Amylum.

AMINÆA, Anime.

AMINÆUM VINUM, *Amine'an wine*, highly esteemed as a stomachic. Virgil distinguishes it from the Falernian.—Pliny, Macrobius, &c.

AMMA, Truss.

AMMI, *Ammi majus seu cicutæfolium seu vulgare* seu *Bolbers*, *Am'mios murica'ta, A'pium*

ammi, *Bishop's weed*. The seeds of this plant are aromatic and pungent. They are said to be carminative and diuretic, and are tonic and stomachic.

AMMI BOLBERI, *Ammi*—*a. des Botaniques*, see *Sison ammi*—*a. Cicutaefolium*, *Ammi*—*a. Majus*, *Ammi*—*a. Verum*, see *Sison ammi*—*a. Vulgare*, *Ammi*.

AMMIION, *Hydrargyri sulphuretum rubrum*.

AMMIOS MURICATA, *Ammi*.

AMMISMUS, *Psammismus*.

AMMOCHO'SIA, *Ammocho'sis*, from *ampos*, 'sand,' and *xew*, 'I pour.' *Arenatio*. Putting the human body in hot sand, for the cure of disease.

AMMO'NIA, *Ammonia* or *Ammoniacal gas*, *Volatile alkali*, *Alkali ammoniacum causticum*, *A. volatile causticum*, *Ammonia caustica seu pura*, *Ammoniacum*, *A. causticum*, *Gas ammoniacale*, *Nephritis urinosæ*, (F.) *Ammoniaque*, *Air alcalin*, *Gaz ammoniacal*. An alkali, so called, because obtained principally by decomposing sal ammoniac (*muriate of ammonia*) by lime. This gas is colourless, transparent, elastic, of a pungent, characteristic odour, and an acrid, urinous taste. It turns the syrup of violets green, and its specific gravity is 0.596. When inhaled, largely diluted with common air, it is a powerful irritant. When unmixed, it instantly induces suffocation.

AMMONIA, ACETATE OF, *SOLUTION OF, Liquor ammoniæ acetatis*—*a. Arseniate of*, *Arseniate of ammonia*—*a. Benzoate of*, *Ammoniac benzoas*—*a. Bicarbonate of*, see *Ammoniac carbonas*—*a. Caustica liquida*, *Liquor ammoniæ*—*a. Chlorohydrate of*, *Ammoniac murias*—*a. Citrate of*, *Ammoniac citras*—*a. Hydriodate of*, *Ammonium, iodide of*—*a. Hydrochlorate of*, *Ammoniac murias*—*a. Hydrosulphuret of*, *Ammoniac sulphuretum*—*a. Iodide of*, see *Iodine*—*a. Liniment of*, strong, *Linimentum ammoniæ fortius*—*a. Liquid*, *Liquor Ammoniac*—*a. Muristica*, *Ammoniac murias*—*a. Nitrata*, *Ammoniac nitras*—*a. Phosphate of*, *Ammoniac phosphas*—*a. Præparata*, *Ammoniac carbonas*—*a. Pura liquida*, *Liquor ammoniæ*—*a. Solution of*, *Liquor ammoniæ*—*a. Solution of*, stronger, *Liquor ammoniæ fortior*—*a. Tartrate of*, *Ammoniac tartas*.

AMMO'NIAC, GUM, *Ammoniacum* (Ph. U. S.), *Gummi Ammoniacum*, *Armoni'acum*, *Mato'rium*, (F.) *Ammoniac*, *Gomme ammoniacque*, so called from *Ammonia* in *Lybia*, whence it is brought. A gum-resin, the concrete juice of *Jore'ma ammoniacum*, of *Persia*: a species of a genus allied to *Ferula*; and also the gum-resin of *Ferula tingitana*. The latter is the ammoniacum of commerce. It is in irregular, dry masses and tears, yellow externally, whitish within. Its odour is peculiar, and not ungrateful: taste nauseous, sweet, and bitter. It forms a white emulsion with water: is soluble in vinegar; partially so in alcohol, ether, and solutions of the alkalis.

Gum ammoniacum is expectorant, deobstruent (?), antispasmodic, discutient, and resolvent. It is chiefly used, however, in the first capacity, and in the formation of certain plasters.

Two varieties are met with in the market, *Guttæ ammoniaci*, the best; and *Lapis ammoniaci*, the more impure.

AMMONIACÆ NITRAS, *Ammoniac nitras*—*a. Sulphas*, *Ammoniac sulphas*.

AMMONIACUM, *Ammonia*, *Ammoniac gum*—*a. Hydrochloratum ferratum*, *Ferrum ammoniatum*—*a. Succinatum*, *Spiritus ammoniæ foetidus*—*a. Volatile mite*, *Ammoniac carbonas*.

AMMONIÆ ACETAS, *Liquor ammoniæ acetatis*—*a. Arsenias*, *Arseniate of Ammonia*.

AMMONIÆ BENZOAS, *Benzoate of Ammonia*. A salt formed by the union of benzoic acid and ammonia, which has been prescribed for the removal of gouty depositions of urate of soda in the joints. It is regarded as a good diuretic.

AMMONIÆ BICARBONAS, *Bicarbonate of Ammonia*, is official in the Dublin Pharmacopœia. It has the same properties as the next, and is more palatable. Dose, six to twenty-four grains.

AMMONIÆ CARBONAS, *A. Subcarbonas seu Sesquicarbonas*, *Salt of bones*, *Sal Osium*, *Sal Fuliginis*, *Salt of wood-soot*, *Salt of urine*, *Volatile Sal Ammoniac*, *Baker's salt*, *Alkali volatile æratum*, *A. volatile ammoniacale*, *A. volatile ex sale ammoniaco*, *Ammoniacum volatile mite*, *Ammonium carbonicum seu subcarbonicum*, *Carbonas ammoniaci alkalinus seu incompletus seu superammoniacus*, *Hypocarbonas ammoniacæ*, *Flores salis ammoniaci*, *Sal cornu cervi volatile*, *Sal volatile salis ammoniaci*, *Concrete or mild volatile alkali*, *Carbonate or Subcarbonate of ammonia*, *Ammonia præparata*, *Sal volatile*, *Smelling salt*, (F.) *Carbonate d'ammoniaque*, *Sel volatil d'Angleterre*, (*Ammon. muriat. bij*; *Oretæ Biss. Sublime*—Ph. U. S.) A white, striated, crystalline mass; odour and taste pungent and ammoniacal: soluble in two parts of water: insoluble in alcohol: effloresces in the air. It is stimulant, antacid, diaphoretic, and antispasmodic. Dose, gr. v to xv.

Carbonate of ammonia is at times used to form effervescing draughts. One scruple saturates six fluidrachms of lemon juice, twenty-six grains of crystallized tartaric acid, and twenty-six grains of crystallized citric acid.

Carbonate of ammonia is the usual smelling salts. When a few drops of *Liquor Ammoniac fortior* are added to it coarsely powdered, and some volatile oil, it forms *Preston Salts*.

AMMONIÆ CITRAS, *Citrate of Ammonia*. Made by saturating lemon or lime juice, or a solution of citric acid, with carbonate of ammonia. Dose, fʒss. It may be made extemporaneously, and taken in an effervescing state. Seventeen grains of citric acid, or half a fluidounce of lemon juice, will be sufficient for thirteen grains of carbonate of ammonia.

AMMONIÆ CUPRO-SULPHAS, *Cuprum ammoniatum*.

AMMONIÆ ET FERRI MURIAS, *Ferrum ammoniatum*—*a. Ferro-citras*, *Ferri ammonio-citras*—*a. Hydriodas*, *Ammonium, iodide of*—*a. Hydrochloras*, *Ammoniac murias*—*a. Hydrosulphuretum*, *Liquor fumans Boylii*—*a. Hypocarbonas*, *Ammoniac Carbonas*.

AMMONIÆ MU'RIAS (Ph. U. S.), *Muriate of Chlorohydrate of Ammonia*, *Hydrochlorate of Ammonia*, *Chlorohydrate of Ammonia* (Ph. U. S.), *Sal Ammoniacum*, *Sal Ammoniac*, *Sal Ammoniacus*, *Ammoniac Muriatica*, *Ammonium Muriatum*, *Ammonia Hydrochloras*, *Chloruretum Ammoniacum*, *Sal Armeniacum*, *Salmiac*, *Fuligo Al'ba Philosophorum*, *Misadir*, (F.) *Muriate d'Ammoniaque*, *Sel Ammoniac ou Armeniac*. A saline concrete, formed by the combination of muriatic acid with ammonia. In Egypt it is manufactured in large quantities by subliming the soot formed by burning camel's dung—26 pounds of the soot yielding 6 pounds. It is also prepared, in great quantities, by adding sulphuric acid to the volatile alkali obtained from soot, bones, &c., mixing this with common salt, and subliming.

Muriate of ammonia is inodorous, but has an acrid, pungent, bitterish, and urinous taste. Three parts of cold water dissolve one. Soluble also in 4.5 parts of alcohol. It is aperient and diuretic, but seldom used internally. Ex-

ternally, it is employed, producing cold during its solution, in inflammations, &c.

AMMO'NIÆ NITRAS, *Nitrate of Ammonia*, *Al'kali volat'ilē nitratum*, *Sal ammoni'acus nitro'sus*, *Ammo'nia nitra'ta*, *Nitras ammoni'acæ*, *Nitrum flammans*, (F.) *Nitrate d'Ammoniaque*. A salt composed of nitric acid and ammonia. It is diuretic and deobstruent. (?) Externally, it is discutient and sialogogue.

AMMO'NIÆ PHOSPHAS, *Ammonium phosphoricum*, *Phosphate of Ammonia*, (F.) *Phosphate d'Ammoniaque*. This salt has been recommended as an excitant, diaphoretic, and discutient. More recently, it has been proposed as a new remedy for gout and rheumatism, as a solvent of uric acid calculus, and for diseases, acute and chronic, connected directly with the lithic acid diathesis.

AMMONIÆ SESQUICARBONAS, *A. carbonas*.

AMMO'NIÆ SULPHAS, *Sulphate of Ammonia*, *Sulphas ammoni'acæ*, *Ammonium sulphuricum*, *Al'kali volat'ilē vitriolatum*, *Sal Ammoniacum secrete* GLAUBERI, *Sal secrete* GLAUBERI, *Vitriolum ammoniacalē*, (F.) *Sulphate d'Ammoniaque*. Formed by adding sulphuric acid either to sal ammoniac or to ammoniacal liquor. Its properties are like those of the muriate of ammonia.

AMMO'NIÆ SULPHURETUM, *Sulphuret of Ammonia*, *Hydrosulphuret of Ammonia*, *Ammonium Sulphhydratum*, *Hydrosulphas Ammoniacæ*, *Spiritus BEGU'NI*, *Sp. fumans BEGU'NI*, *Sulphuretum ammoni'acæ*, *Sp. salis ammoni'aci sulphuratus*, *Liquor ammoni'aci hydrothi'odis*, *Hydrosulphuretum Ammonicum*, *Hydrosulph. ammoniacalē aquosum*, *Hydrog'eno-sulphuretum ammoni'acæ liquidum*, *Spiritus sulphuris volatilis*, *Hepar sulphuris volat'ilē*, BOYLE'S or BEGUINE'S *fuming spirit*, (F.) *Hydrosulphate sulfuré d'Ammoniaque*, *Liquetur fumante de BOYLE*, *Sulfure hydrogéné d'Ammoniaque*, *Hydrosulfure d'Ammoniaque*. Odour very fetid; taste nauseous and styptic; colour dark yellowish green. It is reputed to be sedative, nauseating, emetic, disoxygenizing (?) and has been given in diabetes and diseases of increased excitement. Dose, grt. viij to grt. xx.

AMMO'NIÆ TARTRAS, *Al'kali volat'ilē tartarizatum*, *Sal Ammoniacum tartaricum*, *Tartarus ammoniacæ*, *Tartrate of Ammonia*, (F.) *Tartrate d'Ammoniaque*. A salt composed of tartaric acid and ammonia. It is diaphoretic and diuretic; but not much used.

AMMONIAQUE, *Ammonia*—*a. Arséniate d'*, *Arseniate of ammonia*—*a. Hydrosulfure d'*, *Ammoniac sulphuretum*—*a. Hydrosulfate sulfuré d'*, *Ammoniac sulphuretum*—*a. Liquide*, *Liquor ammonia*—*a. Phosphate d'*, *Ammoniac phosphas*—*a. Sulfure hydrogéné d'*, *Ammoniac sulphuretum*.

AMMONIÆ IODIDUM, *Ammonium*, *iodide of*

AMMONIO-CUPRICUS SUBSULPHAS, *Cuprum ammoniatum*.

AMMO'NION, from *ampos*, 'sand.' An ancient collyrium of great virtue in many diseases of the eye, and which was said to remove sand from that organ.

AMMONIUM ARSENICICUM, *Arseniate of ammonia*—*a. Carbonicum*, *Ammoniac carbonas*—*a. Hydroiodicum*, *Ammonium*, *iodide of*—*a. Iodatum*, *Ammonium*, *iodide of*.

AMMO'NIUM, IODIDE OF, *Iod'idum seu Ioduretum ammonii*, *Ammonium Iodatum seu Hydroiodicum*, *Hydri'odas ammoniacæ*, *Hydri'odate of ammoniac*. This salt is formed by saturating liquid hydroiodic acid with caustic ammonia, and evaporating the solution. It is applied in the form of ointment (3j ad adipis 3j) in lepra, psoriasis, &c.

AMMONIUM MURIATICUM MARTIATUM SEU MARTIALE, *Ferrum ammoniatum*—*a. Muriatum*, *Am-*

moniac murias—*a. Phosphoricum*, *Ammoniac phosphas*—*a. Subcarbonicum*, *Ammoniac carbonas*—*a. Sulphhydratum*, *Ammoniac sulphuretum*—*a. Sulphuricum*, *Ammoniac sulphas*.

AMNA ALCALIZATA, *Water*, *mineral*, *saline*.

AMNEMOSYNE, *Amnesia*.

AMNE'SIA, *Amnest'ia*, *Amnemos'ynē*, from *a*, *privative*, and *μνησις*, 'memory.' *Mo'ria imbec'ilis amne'sia*, *Obliv'io*, *Recollectio'nis jactura*, *Dysæsthe'sia inter'na*, *Debil'itas memo'riae*, *Memoria dele'ta*, (F.) *Perte de Mémoire*, 'loss of memory.' By some nosologists, amnesia constitutes a genus of diseases. By most, it is considered only as a symptom, which may occur in many diseases.

AMNESTIA, *Amnesia*.

AMNIC, *Amniotic*.

AMNIC ACID, *Amniotic acid*.

AMNIITIS, *Amnitis*.

AMNIOLEP'SIS, from *amnios*, and *ελεπω*, 'I steal or take away clandestinely.' Premature escape of the liquor amnii.

AMNIORRHŒ'A, from *amnios*, and *ρρω*, 'I flow.' A premature discharge of the liquor amnii.

AM'NION, *Am'nios*, *Annium*, *Hym'nium*, *Charta virgin'ea*, *Armata'ra*, *Agni'na membra'na*, *Pellu'cida membra'na*, *Galea*, *Scepar'num*, *Indu'sium*, *Amic'ulum*, *Membra'na factum involvens*. The innermost of the enveloping membranes of the fetus:—so called because first observed in the sheep (?), (*αμνος*, 'a sheep.') It is thin, transparent, perspirable, and possesses many delicate colourless vessels, which have not been injected. It is generally considered to be produced by a fold of the external layer of the germinal membrane, rising up, and gradually enveloping the embryo. Its external surface is feebly united to the chorion by areolar and vascular filaments. Its inner surface is polished, and is in contact with the body of the fœtus and the liquor amnii.

AMNIOT'IC, *Amniot'icus*, *Amnic*, *Am'nicus*, (F.) *Amniotique ou Annique*. Relating, or appertaining, to the amnios.

AMNIOT'IC ACID, *Ac'idum am'nicum seu amniot'icum*. A peculiar acid, found by Vauquelin and Buviva in the liquor amnii of the cow.

AMNIQUE, *Amniotic*.

AMNI'TIS, *Amni'tis*, from *amnion* and *itis*, 'inflammation.' Inflammation of the amnion.

AMENOMANIA, from *amēnus*, 'agreeable,' and *mania*. A form of mania in which the hallucinations are of an agreeable character.

AMOME FAUX, *Sison amomum*.

AMO'MUM CARDAMO'MUM (Ph. D.), *A. verum*, *Alpin'ia cardamomum*, *Caro'pi*, *Mato'nia*, *Cardamo'mum*, *Elettaria Cardamo'mum*, *Cardamo'mum Minus*, *Lesser*, *True*, or *Official Cardamom*, (F.) *Cardamome de la Côte de Malabar*, *Cardamome*. Ord. Zingiberaceæ. The fruit of this East India plant—*Cardamonum*, (Ph. U. S.), *C. Malabaren'sē*—has an agreeable, aromatic odour, and a pungent, grateful taste. It is carminative and stomachic: but is chiefly used to give warmth to other remedies. It is called *Amomis*. Dose, gr. v to ʒj.

The *Amo'mum Cardamo'mum* of Linnæus, *Cluster* or *Round Cardamom* of Sumatra, Java, and other islands eastward of the Bay of Bengal, yields the round *Cardamom*, *Cardamo'mum rotundum*, of the shops. The fruits, in their native clusters or spikes, constituting the *Amo'mum racemosum*, are rarely met with.

AMOMUM CURCUMA, *Curcuma longa*.

AMOMUM GALANGA, *Maranta galanga*.

AMOMUM GRANUM PARADISI, *A. max'timum*, *Cardamo'mum majus seu pipera'tum*, *Meleguet'ta*, *Maniguet'ta*. Greater cardamom seeds—*Grana Paradisi*, *Grains of Paradise*, (F.) *Graines de*

Paradis—resemble *A. cardamomum* in properties. They are extremely hot, and not much used.

AMOMUM, GREAT-WINGED, *Amomum maximum*—*a. Hirsutum, Costus.*

AMOMUM MAXIMUM, *Great-winged Amomum.* Yields the fruit known in commerce by the names *Jana* or *Nipal Cardamoms, Bengul Cardamoms* of the Calcutta market. &c.—Percira.

AMOMUM MONTANUM, see Cassuminiar—*a. Pimenta, see Myrtus pimenta*—*a. Racemosum, see A. cardamomum*—*a. Sylvestre, see Cassuminiar*—*a. Zedoaria, Kaempferia rotunda*—*a. Zerumbet, see Cassuminiar.*

AMOMUM ZINGIBER, *Zin'giber officinale* seu *album* seu *nigrum* seu *commune, Zin'ziber, Ginger, (F.) Gengembre.* The black and white ginger, *Zin'ziber fuscum* et *album, Zin'giber* (Ph. U. S.), are the rhizoma of the same plant, *Zin'giber officinale*, the difference depending upon the mode of preparing them.

The odour of ginger is aromatic; taste warm, aromatic, and acrid. It yields its virtues to alcohol, and in a great degree to water. It is carminative, stimulant, and sialogogue.

Preserved Ginger, Zingiberis Radix Conditâ, Radix Zingiberis condita ex Indrâ allatâ, is a condiment which possesses all the virtues of ginger.

Ginger-Beer Powders may be formed of *white sugar* ʒi and ʒij, *ginger gr. v, subcarbonate of soda gr. xxxvj,* in each blue paper: *acid of tartar* ʒiss, in each white paper—for half a pint of water.

Oxley's Concentrated Essence of Jamaica Ginger is a solution of ginger in rectified spirit.

AMOR, Love.

AMORGE, Amurca.

AMORPHUS, Anhistous, Anideus.

AMOSTEUS, Osteocolla.

AMOUR, Love—*a. Physique, Appetite, venereal.*

AMOUREUX, (muscle.) Obliquus superior oculi.

AMPAC, *Amp'acus.* An East India tree, the leaves of which have a strong odour, and are used in baths as detergents. A very odoriferous resin is obtained from it.

AMPAR, Succinum.

AMPELOCAEPUS, Galium aparine.

AMPELOPRASUM, Allium.

AMPELOPSIS QUINQUEFOLIA, (αμπελος, 'the vine,' and οψis, 'appearance,') Virginian Creeper, American Ivy, Five-leaved Ivy, Woody Climber. An indigenous climbing plant. *Ord. Vitaceæ;* which flowers in July. It has been advised as an expectorant.

AMPELOS, Vitis vinifera—*a. Agria, Bryonia alba*—*a. Idea, Vaccinium Vitis Idea*—*a. Melæna, Tannus communis*—*a. Oinophorus, Vitis vinifera.*

AMPHAMPHOTERODIPSIA, Diplopia.

AMPHARIS'TEROS, Ambla'tus, 'awkward,' from ἀμφι, and ἀριστερος, 'the left.' Opposed to ambidexter.

AMPHEMERINOS, Quotidian.

AMPHEMERUS, Quotidian.

AMPHI, αμφι, 'both, around, on all sides.' Hence, a prefix in many of the following terms.

AMPHIAM, Opium.

AMPHIARTHRO'SIS, from αμφι, 'both,' and αρθρσις, 'articulation.' A mixed articulation, in which the corresponding surfaces of bones are united in an intimate manner by an intermediate body, which allows, however, of some slight motion. Such is the junction of the bodies of the vertebrae by means of the intervertebral cartilages. This articulation has also been called *Diarthrose de Continuité.* The motion it permits is but slight.

AMPHIBLESTROIDITIS, Retinitis.

AMPHIBLESTROIDES, Reticular.

AMPHIBLESTROMALACIA, from αμφιβλεστροιδες (membrana), 'the retina, and μαλακία, 'softening.' Mollescence or softening of the retina.

AMPHIBRAN'CHIA, from αμφι, 'around,' and βραχχια, 'the throat.' Amphibron'chia. The tonsils and neighbouring parts.—Hippocrates.

AMPHICAUSTIS, Vulva.

AMHID'EUM, from αμφι, 'around,' and ιδω, 'I bind.' The outermost margin of the cervix uteri: the *Labium uteri.*

AMPHIDEXIUS, Ambidexter.

AMPHIDIARTHRO'SIS, from αμφι, 'about,' and αρθρσις, 'moveable joint.' A name given by Winslow to the temporo-maxillary articulation, because, according to that anatomist, it partakes both of ginglymus and arthrodia.

AMPHIESMA CORDIS, Pericardium.

AMPHIMERINA, Pertussis—*a. Hectica, Hectic fever.*

AMPHIMERINOS, Quotidian.

AMPHION, Maslach.

AMPHIPLIX, Perinæum.

AMPHIPNEUMA, Dyspnœa.

AMPHISMELA, Knife, double-edged.

AMPHISMILE, Knife, double-edged.

AMPHISPHAL'SIS, Circumactio, Circumductio, from αμφι, 'around,' and σφαλλα, 'I wander.' The movement of circumduction used in reducing luxations.—Hippocrates.

AMPHODIPLOPIA, see Diplopia.

AM'PHORA, per syncope, for αμφιφορεus, from αμφι, 'on both sides,' and φεω, 'I bear,' because it had two handles. A liquid measure among the ancients, containing above seven gallons. Also called *Quadrantal, Cera'mium, Ceram'nium, Cudus.*

AMPHORIC RESPIRATION, see Cavernous Respiration.

AMPHOTERODIPLOPIA, see Diplopia.

AMPHRODIPLOPIA, Diplopia.

AMPLEXATIO, Coition.

AMPLEXUS, Coition.

AMPLIFICATIO, Platynosis.

AMPLIOPIA, Amblyopia.

AMPOSIS, Anaposis.

AMPOULES, Essera.

AMPUL'LA, (L.) 'A bottle.' A membranous bag, shaped like a leathern bottle. See *Cavitas Elliptica.* In pharmacy, a receiver.

AMPELA CHYLIFERA SEU CHYLI, Receptaculum chylli.

AMPULLÆ, Phlyctenæ.

AMPUTATION, *Amputa'tio, from amputare, amputatum, (am, 'around,' and putare, 'to cut off,') Apot'omē, Apotom'ia.* The operation of separating, by means of a cutting instrument, a limb or a part of a limb, or a projecting part, as the mamma, penis, &c., from the rest of the body. In the case of a tumour, the term *excision, removal, or extirpation, (F.) Resection,* is more commonly used.

AMPUTATION, CIRCULAR, is that in which the integuments and muscles are divided circularly.

AMPUTATION, FLAP, (F.) *A. à lambeau,* is when one or two flaps are left so as to cover the stump when the limb has been removed.

AMPUTATION, JOINT, *Exarticulatio, (F.) A. dans l'article ou dans la contiguité des membres,* is when the limb is removed at an articulation.

Each amputation requires a different process, which is described in works on operative surgery.

AMPUTATION, SPONTANEOUS, see Spontaneous.

AMULET, Amuletum.

AMULETTE, Amuletum.

AMULE'TUM, from amoliri, 'to remove.' An Amulet, Perian'ma, Apotropa'um, Periap'ton, Phylact'ion, Apoteles'ma, Leucite'ma, Alexica-

eum, Præservativum, Probæscan'ium, Probæscan'tium, (F.) *Amulette*. Any image or substance worn about the person for the purpose of preventing disease or danger.

AMUR'CA, *Amur'ga*, ἀμύργα, from ἀμύρω, 'I press out.' The marc or grounds remaining after olives have been crushed and deprived of their oil. It has been used as an application to ulcers.

AMUR'GA, *Amurca*.

AMUSA, *Musa Paradisiaca*.

A'MYCE, *Amycha*, Ἀμύχ'ις. Excoriation, Scarification.

AMYCHA, *Amyce*.

AMYCTICA, from ἀμύσσω, 'I lacerate.' Medicines which stimulate and vellicate the skin.—*Cælius Aurelianus*.

AMYDOLÉ, *Fecula*.

AMYDRIASIS, *Mydriasis*.

AMEL'IA, from α, privative, and μέλος, 'marrow.' A monstrous formation, in which there is an absence of spinal marrow.

AMELONER'VIA; from α, privative, μέλος, 'marrow,' and νεύρον, 'nerve.' Paralysis or deficient action of the spinal marrow.

AMELOTROPH'IA, from α, privative, μέλος, 'marrow,' and τροφή, 'nourishment.' Atrophy of the spinal marrow.

AMYG'DALA, same etymon as *Amyctica*; because there seem to be fissures in the shell. The *Almond*, of which there are two kinds; *Amygdalæ amara* and *A. dulces*, (F.) *Amygdalus amara*, and *A. douces*, obtained from two varieties of *Amygdalus communis* or *A. sativa*, *Almond tree*, (Old Eng.) *Amyllier*, a native of Barbary. *Ord. Amygdalææ. Sex. Syst. Icosandria Monogynia*.

The taste of *Amygdala dulcis* is soft and sweet; that of *A. amara*, bitter. Both yield, by expression, a sweet, bland oil. The bitter almond contains Prussic acid. They are chiefly used for forming emulsions.

AMYG'DALÆ PASTA, *Almond Paste*, a cosmetic for softening the skin and preventing chaps, is made of bitter almonds, blanched, ʒiv, white of one egg; rose water, and rectified spirit, equal parts, or as much as is sufficient.

AMYG'DALÆ PLACEN'TA, *Almond Cake*, is the cake left after the expression of the oil. The ground *Almond Cake*, *Almond Powder*, *Fari'na Amygdala'rum*, is used instead of soap for washing the hands.

AMYG'DALA, *Tonsil*. Also, a lobule or prominence of the cerebellum, so called from its resemblance to an enlarged tonsil. This and its fellow of the opposite side form the lateral boundaries of the anterior extremity of the *valley*, and are in great part covered by the medulla oblongata. The *Amygdalæ* are seated on either side of the uvula, in the fourth ventricle.

AMYG'DALATOME, *Amygdalatomus*, from ἀμύδαλν, 'tonsil,' and τομή, 'incision,' improperly *Tonsillitome*.—*Tonsil-guillotine*, (F.) *Sc'ateur des Amygdalæ*. An instrument for surgery and excising a portion of the tonsil. See *Kioto-me*.

AMYG'DALATUM, *Emulsio amygdalæ*.

AMYG'DALÉ, *Tonsil*.

AMYG'DALIN, *Amygdali'num*, *Amygdali'na*, *Amygdaline*. A principle contained in bitter almonds, which is prepared by pressing the bruised almonds between heated plates to separate the fat oil; boiling the residue in alcohol; evaporating, and treating with ether, which precipitates the amygdalin in a crystalline powder. A weak solution of it, under the influence of a small quantity of *emulsin* or *synaptase*, which constitutes the larger portion of the pulp of almonds, yields at once oil of bitter almonds and hydrocyanic acid.

AMYG'DALITIS, *Cynanche tonsillaris*.

AMYG'DALUS, see *Amygdala*.

AMYG'DALUS COMMUNIS, see *Amygdala*.

AMYG'DALUS PER'SICA, *Per'sica vulgaris*. The common peach-tree, (F.) *Pêcher*. The leaves and flowers have been considered laxative. They are bitter and aromatic, and have been given in hæmaturia, nephritis, &c. The fruit is one of the pleasant and wholesome summer fruits, when ripe. The kernels, *Amygdalæ Per'sicæ*, as well as the flowers, contain prussic acid.

Peach Brandy is distilled from the fruit, and is much used in the United States.

AMYG'MOS, *Scarification*.

AMYL, *Amylum*.

AMYL'ACEOUS, *Amyla'ceus*, (F.) *Amylacé*, from *amylum*, 'starch.' Having the nature of, or containing starch. Starchy, starch-like.

AMYLACEOUS BODIES, *Corpora amylacea*.

AMYLE, *HYDRATED OXIDE OF*, Oil, fusel.

AMYLENE, see *Potato oil*.

AMYLEON, *Amylum*.

AMYL'I IODIDUM, *Starch, iodide of*—*a. Ioduretum*, *Starch, iodide of*.

AMYL'LIER, see *Amygdala*.

AMYLOID BODIES, *Corpora amylacea*.

A'MYLUM, *A'midum*, *Fec'ula*, *Amyl'leon*, *Amyl'ion*, *Amyl'on*, from α, priv., and μύλη, 'a mill,' because made without a mill. *Starch*, (Old Eng.) *Amyl*, (F.) *Amidon*, *Amylon*. *Starch of Wheat*, *Fari'na*, *Tritic' fari'na*, *Amylum tritic'eum* seu *Tritic'i*, *Fec'ula Amyla'cea*, is inodorous and insipid, white and friable. It is insoluble in cold water and alcohol, but forms with boiling water a strong, semi-transparent jelly. It is demulcent, and is used as an emollient glyster, and as the vehicle for opium, when given *per anum*. Starch is met with abundantly in all the cereal grains, in the stalks of many of the palms, in some lichens, and in many tuberous roots, particularly in the bulbs of the orchis.

AMYLUM AMERICANUM, see *Arrow-root*—*a. Cannaceum*, *Tous-les-mois*—*a. Iodatium*, *Starch, iodide of*—*a. Manihoticum*, see *Jatropha manihot*—*a. Marantaceum*, *Arrow-root*—*a. Palmaceum*, *Sago*—*a. Quercneum*, *Rachahout*.

A'MYON, from α priv., and μυων, 'a muscle,' *Ennusculation*. Without muscle. Applied to the limbs, when so extenuated that the muscles cannot be distinguished.

AMYOSIS, *Synchezis*.

AMYOSTHEN'IA, (F.) *Amyosthénie*, from α, priv., μυων, 'a muscle,' and σθένος, 'strength.' Defect of muscular contraction.

AMYOSTHÉNIE CYSTURIQUE, *Paralysis of the bladder*.—*Piorry*.

AMYRIS COMMIPHORA, see *Bdellium*.

AM'YRIS ELEMIF'ERA, (a. intensive, and μύρον, 'an odoriferous ointment,' because it enters into the composition of such.) (F.) *Balsamier* ou *Baumier Élémifère*. *Ord. Terebinthaceæ. Sex. Syst. Octandria Monogynia*. The plant whence it has been supposed *GUM EL'EMI* is obtained. This gum or resin is brought from the Spanish East and West Indies. *Brazilian Elemi*, according to Dr. Royle, is produced by *Leica leicarbiba*; *Mexican Elemi*, by *Ela'phrium elemiferum*; and *Manilla Elemi*, by *Canarium commune*. It is softish, transparent, of a pale whitish colour, inclining a little to green, and of a strong, though not unpleasant smell. It is only used in ointments and plasters, and is a digestive.

AMYRIS GILEADENSIS, see *A. opobalsamum*.

AM'YRIS OPOBAL'SAMUM, (F.) *Balsamier* ou *Baumier de la Mecque*, *Bal'sem*, *Bal'samum*. The plant from which is obtained the *BALSAM OF MECCA*, *Bal'samum genuinum antiquum* seu *Asiat'icum* seu *Juda'icum* seu *Syriacum* seu *e*

Meerl seu Alpi'ni, Balsamel'on, Egyptiacum
Balsamum, Cuccubalsamum, Oleum Balsami,
Opobalsamum, Xylobalsamum, Balm or *Balm*
of Gilead, (F.) Baume Blanc, B. de Constantino-
ple blanc, B. de Galaad, B. du Grand Caïre, B.
Vrai, Terebinthine de Gilead, T. d'Egypte, T. du
Grand Kaïre, T. de Judée. A resinous juice
obtained by making incisions into Amyris opobal-
samum and A. Gileaden'sis of Linnaeus, Balsa-
maiden'dron Gileaden'se of Kunth. The juice
of the fruit is called Caryobalsamum; that of the
wood and branches Xylobalsamum. It has the
general properties of the milder Terebinthinales.

AMYRIS TOMENTOSUM, Fagara octandra.

AMYRON, Carthamus tinctorius.

AMYUS, from *a*, privative, and *μῦς*, 'a mouse,
 a mouse.' Weak or poor in muscle.

AMYXIA, from *a*, privative, and *μῦξα*, 'mu-
 cus.' Deficiency of mucus.

AMYXIS, Amyce, Scarification.

ANA, *ava*, a word which signifies 'of each.'
 It is used in prescriptions as well as *ā* and *āā*, its
 abbreviations. As a prefix to words, it means
 'in,' 'through,' 'upwards,' 'above,' in opposition
 to *cata*;—also, 'repetition,' like the English *re*.
 Hence,—

ANAB'ASIS, from *αναβαίνω*, 'I ascend.' The
 first period of a disease, or that of increase.—
 Galen. See Augmentation.

ANABEXIS, Expectoration.

ANABLE'SIS, from *ana*, 'again,' and *βλεπω*,
 'I see.' Restoration to sight.

ANABOL'E'ON, *Anabole'us*, from *αναβαλλω*, 'I
 cast up.' An ointment for extracting darts or
 other extraneous bodies.

ANAB'OLE, from *ana*, 'upwards,' and *βαλλω*,
 'I cast.' *Anago'ge, Anaph'ora, Anacine'ma,*
Anacine'sis. An evacuation upwards. An act
 by which certain matters are ejected by the
 month. In common acceptance it includes, *ex-*
spuition, expectoration, regurgitation, and vomit-
ing.

ANABROCHIS'MUS, *Anabron'chismus*, from
ava, 'with,' and *βροχος*, 'a running knot.' An
 operation for removing the eye-lashes, for exam-
 ple, when they irritate the eye, by means of a
 hair knotted around them.—Hippocrates, Galen,
 Celsus, &c.

ANABRONCHISMUS, Anabrochismus.

ANABROSIS, Corrosion, Erosion.

ANACAMPSELOS, Sedum telephium.

ANACARDIUM OCCIDENTA'LE, (*ava* and
καρδία, 'heart,' from the resemblance of the fruit
 to a dried heart,) *Acaju'ba occidenta'lis, Cassi-*
nvium pomiferum, Cashew (W. Indies.) (F.) Ac-
ajou. Ord. Terebinthaceæ. Sex. Syst. Ennean-
dria Monogynia. The Oil of the Cashew Nut,
O'leum Anacar'dii, (F.) Huile d'Acajou, is an
 active caustic, and used as such in the countries
 where it grows, especially for destroying warts,
 &c.

ANACARDIUM ORIENTALE, Avicennia tomentosa.

ANACATHAR'SIS, from *ava*, 'upwards,' and
καθαίρειν, 'to purge.' Purgation upwards. Ex-
 pectoration. See, also, Repurgatio.

ANACATHARSIS CATARRHALIS SIMPLEX, Ca-
 tarrh.

ANACATHARTICUS, Expectoant.

ANACESTOS, Incurable.

ANACHREMPsis, Exspuition.

ANACHRON, Soda.

ANACINEMA, Anabole, Exspuition.

ANACINESIS, Anabole, Exspuition.

ANACLASIS, Repercussion.

ANACLINTE'RIUM, *Anacelin'trum, Recubi-*
to'rium, from *ανακλινω*, 'I recline.' A long chair
 or seat, so formed that the person can rest in a
 reclining posture.

ANACLINTRUM, Anaclinerium.

ANACOLLE'MA, from *ava*, 'together, and
κόλλω, 'I glue.' A healing medicine.

ANACOLLE'MATA, Frontal bandages.

ANACOLUP'PA. A creeping plant of Mala-
 bar, the juice of which, mixed with powdered
 pepper, passes in India as a cure for epilepsy,
 and as the only remedy for the bite of the naja.
 It is supposed to be *Zapala'na nudiplo'ra*.

ANACOLU'TTIE, Incoherence.

ANACOMIDE, Restauratio.

ANACONCHYLISMUS, Gargarism.

ANACONCHYLISMUS, Gargarism.

ANACTESIS, Restauratio.

ANACTIRION, Artemisia.

ANACYCLEON, Charlatan.

ANACYCLUS OFFICINARUM, see Anthe-
 mis Pyrethrum—a. Pyrethrum, Anthemis pyre-
 thrum.

ANADESMUS, Fascia.

ANADIPL'O'SIS, from *ava*, 'again,' and *διπλω*,
 'I double.' *Epanadiplo'sis, Epanalep'sis, Redu-*
plicat'io. The redoubling which occurs in a
 paroxysm of an intermittent, when its type is
 double.—Galen, Alexander of Tralles.

ANADORA, Ecdora.

ANAD'OSIS, from *αναδίδωμι*, 'I give up.'
 Purgation upwards, as by vomiting. Congestion
 of blood towards the upper parts of the body.
Anadosis seems also to have occasionally meant
 chylification, whilst *diadosis* meant capillary nu-
 trition.—Hippocrates, Galen.

ANAD'ROME, from *ava*, 'upwards,' and *δρομω*,
 'I run.' The transport of a humour or pain from
 a lower to an upper part.—Hippocrates. Also,
 the globus hystericus. See Angone.

ANEDGE'US, from *av*, privative, and *αἷσις*,
 'organs of generation.' A monster devoid of
 sexual organs.

ANÆMATOPOE'SIS, from *a*, *av*, privative,
 'αἷμα,' 'blood,' and *ποιεω*, 'I make.' Impeded or
 obstructed hæmatosis.

ANÆMATO'SIS, *Anhæmato'sis*, from *a*, *av*,
 privative, and 'αἷμα,' 'blood.' Defective hæma-
 tosis or preparation of the blood. Anæmia.

ANÆMIA, *Ess'emia, Anæ'masis, Anhæ'mia,*
Anhæmato'sis, Polyganhæ'mia, Anæmo'sis, Oligæ-

dræ'mia, Anæ'mia, (F.) Anémie, Anhémic, Anhé-
matotic, Polyanhémic, Hydrehémic, Essanguinité,
Blood'lessness: from *a*, *priv*, and 'αἷμα,' 'blood.'
 Privation of blood;—the opposite to plethora. It
 is characterized by every sign of debility. Also,
 diminished quantity of fluids in the capillary ves-
 sels:—the opposite to *Hyperæmia*.—The essential
 character of the blood in anæmia is diminution
 in the ratio of red corpuscles.

ANÆMIC, *Anem'ic, Anæ'micus;* same ety-
 mon. Appertaining or relating to anæmia,—as
 an "anæmic person." See Exsanguious.

ANÆMOCH'ROUS, from *a*, *av*, privative, 'αἷμα,'
 'blood,' and *χρῶς*, 'colour.' Devoid of colour,
 pale.

ANÆMOSIS, Anæmia.

ANÆMOTROPHY, *Anæmotroph'ia:* from *av*,
 privative, 'αἷμα,' 'blood,' and *τροφή*, 'nourish-
 ment.' A deficiency of sanguineous nourishment.
 —Prout.

ANÆMYDRIA, Anhydræmia.

ANÆSTHESIA, *Anæsthes'is, Insensibili'tas,*
Analges'ia, Parop'sis æpers, (F.) Anæsthésie;
 from *a*, privative, and *αἰσθάνομαι*, 'I feel.' Pri-
 vation of sensation, and especially of that of
 touch, according to some. It may be general or
 partial, and is almost always symptomatic.

ANÆSTHESIA GUSTATORIA, Ageusia—a. Lin-
 guæ, Ageusia—a. Olfactoria, Anosmia—a. Optic,
 Amaurosis.

ANÆSTHESIS, Anæsthesia.

ANÆSTHET'IC, *Anæsthet'ic, Anæsthet'icus,*

(F.) *Anesthésique*; same etymon. as *Anæsthesia*. Relating to privation of feeling, as an "anæsthetic agent;" one that prevents feeling. The term is, now, almost restricted to agents, which produce such effect by being received into the lungs in the form of vapours or gases, and passing with the blood to the nervous centres on which their action is exerted. Perhaps, as a general rule, the intellectual faculties first feel their influence, — a sort of intoxication supervening, with imperfect power of regulating the movements; the sensory ganglia become afterwards or simultaneously affected, sensation and motion are suspended, and ultimately, if the quantity inhaled be sufficient, the medulla oblongata has its actions suspended or destroyed, respiration ceases, and death is the consequence. Different agents have been used as anæsthetics by way of inhalation — sulphuric ether, chloroform, chloric ether, compound ether, chlorohydric and nitric ethers, bisulphuret of carbon, chloride of olefant gas, benzoin, aldehyde, light coal-tar naphtha, &c.; but the first four are alone employed. They have been, and are, greatly used in serious surgical operations, and during parturition; and in such cases, as well as in many diseases, especially of a painful nature, produce the most beneficial results.

ANÆSTHETIZATION, (F.) *Anesthésisation*; same etymon. The condition of the nervous system induced by anæsthetics.

ANAGALLIS, from *ava*, and *γαλα*, 'milk,' from its power of coagulating milk. *A. arvensis* A. Phœnic'ea, Red Pimpernel, Scarlet Pimpernel, Shepherd's Sun-dial. Nat. Ord. Primulacææ. *See*. Syst. Pentandria Monogynia. (F.) *Mouron rouge*. A common European plant; a reputed antispasmodic and stomachic.

Another species — *Anagallis carulea* is a mere variety of the above.

ANAGALLIS AQUATICA, Veronica Beccabunga. ANAGARGALICTON, Gargarism. ANAGARGARISMUS, Gargarism. ANAGARGARISTON, Gargarism. ANAGLYPHE, Calamus scriptorius. ANAGNOSTAKIS, see Ophthalmoscope. ANAGOGE, Anabole, Rejection. ANAGRAPHÉ, Prescription.

ANAGYRIS, *Anagyris foetida*, *Anagyris*, *Ac'oon*, Stinking Bean Trefoil. (F.) *Anagyris*, from *ava*, 'I lead upwards' [?]. Native of Italy. The leaves are powerfully purgative. The juice is said to be diuretic, and the seeds emetic. — Dioscorides, Paulus.

ANAGYRUS, Anagyris.

ANAL, *Analis*. That which refers to the anus; — as *Anal region*, &c.

ANAL'DIA, (F.) *Analdie*; from *a*, *av*, privative, and *αλδεν*, 'to grow.' Defective nutrition.

ANALEMSIA, Analepsia.

ANALENTIA, Analepsia.

ANALEP'SIA, *Analep'sis*, *Analen'tia*, *Analen'sia*, from *ava*, 'fresh,' and *λαμβάνειν*, (future *ληψομαι*), 'to take.' Restoration to strength after disease. — Galen. A kind of sympathetic epilepsy, originating from gastric disorder. See Epilepsy. Also, the support given to a fractured extremity; — *Appen'sio*. — Hippocrates.

ANALEPSIS, Convalescence, Restauratio.

ANALEP'TICA, *Anapeyc'tica*, *Psychot'ica*, *Refect'iva*, *Reficien'tia*, *Restauran'tia*, *Analep'tics*, same etymon. Restorative medicines or food; such as are adapted to recruit the strength during convalescence; — as sago, salep, tapioca, jelly, &c.

ANALEPTIC PILLS, JAMES'S, consist of *James's Powder*, *Gum Ammoniacum*, and *Pills of Aloës and Myrrh*, equal parts, with *Tincture of Castor*, sufficient to form a mass.

ANALGE'SIA, *Analg'ia*, from *a*, *av*, priv., and *αλγος*, 'pain.' Absence of pain both in health and disease. See Anæsthesia.

ANALGIA, Analgesia.

AN'ALOGUE, *Anal'ogus*; from *ava*, 'again,' and *λογος*, 'a description.' A part in one organized being which has the same function as another part in another organized being.

ANALOGOUS TISSUES, see Tissues.

ANALOSIS, Atrophy.

ANALTESIS, Restauratio.

ANALTHES, Incurable.

ANAMIRTA COCCULUS, *Menispermum cocculus* — a. *Peniculata*, *Menispermum cocculus*.

ANAMNES'TIC, *Anamnesticum*, from *ava*, 'again,' and *μνησκω*, 'I remember.' A medicine for improving the memory. See, also, Commemorative.

ANANAS, *Bromelia ananas* — a. *Aculeata*, *Bromelia ananas* — a. *Americana*, *Bromelia pinguin* — a. *Ovata*, *Bromelia ananas* — Wild, broad-leaved, *Bromelia pinguin*.

ANANAZIP'TA. A word formerly scrawled on amulets to charm away disease.

ANANDRI'A, from *a*, *av*, privative, and *ανηρ*, 'a man.' Want of manliness. Impotence in the male. The state and act of emasculation.

ANANEO'SIS, *Renova'tio*; from *ava*, 'again,' and *νεος*, 'new.' Renovation or renewal, — as of the blood by the chyliiferous vessels and lymphatics.

ANAPET'TA, *Expan'sio meatuum*, from *ava*, and *παρα*, 'I dilate.' A state opposite to the closure of vessels. — Galen.

ANAPHALANTIASIS, *Anaphalanto'ma*, from *αναφαλαντιος*, 'bald.' Loss of the hair of the eyebrows. Also, baldness in general.

ANAPHALANTOMA, Anaphalantiasis.

ANAPHE, Anaphia.

ANAPH'IA, *Anaph'ia*, *An'aphē*, from *a*, *av*, priv., and *αφν*, 'touch.' Diminution or privation of the sense of touch.

ANAPHLASMUS, Masturbation.

ANAPHONE'SIS, from *ava*, 'high,' and *φωνη*, 'voice.' Exercise of the voice: vociferation: — the act of crying out. *Vocifera'tio*, *Clamor*.

ANAPHORA, Anabole.

ANAPHRODIS'TA, from *a*, priv., and *Αφροδιτη*, 'Venus,' *Defect'us Ven'eri*. Absence of the venereal appetite. Sometimes used for *Impotence* and *Sterility*.

ANAPHRODISIAC, Antaphrodisiac.

ANAPHROMELI, *Mel despumatum*.

ANAPHLASIS, *Anaplas'mus*, from *αναπλασσω*, 'I restore.' *Confirma'tio*, *Reposi'tio*. Restoration. Union or consolidation of a fractured bone. — Hippocrates.

ANAPLASMATIC, Anaplastic.

ANAPLASMUS, Anaplasia.

ANAPLAS'TIC, *Anaplas'ticus*; same etymon. An epithet applied to the art of restoring lost parts or the normal shape, — as '*Anaplastic Surgery*.' See Morioplastice. Also an agent, that increases the amount of plastic matter — fibrin — in the blood; *Anaplasmat'ic*.

ANAPLERO'SIS, from *αναπληρω*, (*ava*, and *πληρω*), 'I fill up.' Repletion. That part of surgical therapeutics whose object is to supply parts that are wanting. Also, *Apposition* or *Prosthesis*.

ANAPLEROTICUS, Incarnans.

ANAPLEU'SIS, *Fluetua'tio*, *Innata'tio*, from *αναπλεω*, (*ava* and *πλεω*), 'to swim above.' The looseness or shaking of an exfoliated bone; or of a carious or other tooth, &c. — Hippocrates Paulus.

ANAPLOSIS, Growth.

ANAPNEUSIS, Respiration.

ANAPNOE, Respiration.

ANAPNOËNUSI, from *anapnoë*, 'respiration,' and *nosos*, 'disease.' Diseases of the respiratory organs.

ANAPNOMETER, Spirometer.

ANAPODISIS UTERI, Retroversio Uteri.

ANAPODISMUS UTERI, Retroversio Uteri.

ANAPODOPHYLLUM CANADENSE, Podophyllum peltatum.

ANAP'OSIS, *An'posis*, from *ava*, 'again,' and *posis*, 'drink.' A recession of humours from the circumference to the centre of the body.—Hippocrates.

ANAPSE, Auante.

ANAPSIA, Cæcitas.

ANAPSYCTICA, Analeptica.

ANAPTYSIS, Expectoration.

ANAPTXYXIS, Growth.

ANARCOTINA, Narcotine.

ANARRHEGNU' MINA, from *anarrhēgnun*, 'I break out again.' Fractures are so called when they become disunited; as well as ulcers when they break out afresh.

ANARRHINON, from *ava*, 'upwards,' and *arrhē*, 'the nose.' That which returns by the nose.—Gorrieus.

According to others, that which issues by the skin; from *ava*, and *paros*, 'the skin.'

ANARRHINUM, Sternutatory.

ANARRHĒ'A, *Anar'rhoë*, *Anarrho'pia*, *Anar'tasis*, from *ava*, 'upwards,' and *rhoë*, 'I flow.' Afflux of fluid towards the upper part of the body.

ANARRHOPHE, Absorption.

ANARRHIOPHIENUSI, from *anarrhophē*, 'absorption,' and *nosos*, 'disease.' Diseases of the absorbents.

ANARRHIOPHESIS, Absorption.

ANARRHOPIA, Anarrhœa.

ANAR'THRUS, from *av*, priv., and *arthron*, 'a joint.' Without a joint. One who is so that his joints are scarcely perceptible.—Hippocrates.

ANASAR'CA, from *ava*, 'through,' and *σαρξ*, *sarkos*, 'the flesh.' *Anasar'ca*, *Catar'ca*, *Aqua intercus seu inter cutem*, *Hyposar'ca*, *Hydrops cellul'ris toti'us cor'poris*, *H. Anasar'ca seu inter'cus seu subcut'a'neus seu cellul'us seu cut'a'neus seu tela cellul'œ*, *Catar'ca*, *Episarc'id'ium*, *Hy'draton'cus*, *Hy'draton'cus*, *Hy'dron'cus*, *Hydrosar'ca*, *Hydroder'ma*, *Hydroph'is vera*, *Sar'cites*, *Polytym'phia*, *Hyposarcid'ius*, *Lencophlegmat'ia*, *General dropsy*, *Dropsy of the cellular membrane*, (F.) *Anasarque*. Commonly, it begins to manifest itself by swelling around the ankles; and is characterized by tumefaction of the limbs and of the soft parts covering the abdomen, thorax, and even the face, with paleness and dryness of the skin, and pitting when any of these (especially the ankles) are pressed upon. Like dropsy in general, *Anasarca* may be *active* or *passive*; and its treatment must be regulated by the rules that are applicable to general dropsy. At times, the symptoms are of an acute character, and the effusion sudden, constituting *Dermatoc'h'ysis*, *Hydrops Anasar'ca acut'us*, *Ede'ma cal'idum*, *Æ. acut'um*, *Æ. febri'le* of some. See *Hydrops*.

ANASARCA HYSTERICUM, *Anathymiasis*—a. Pulmonum *Hydropneumonia*, *Edema of the Lungs*—a. *Serosa*, *Pneumatia dolens*.

ANASARCHA, *Anasarca*.

ANASARQUE, *Anasarca*.

ANASISMUS, Concussion.

ANASPADIA, see *Anaspadæus*.

ANASPA'DIAS, *Epispad'ias*, from *ava*, 'upwards,' and *σπαω*, 'I draw.' One whose urethra opens on the upper surface of the penis.

ANASPADISIS, see *Anaspadæus*.

ANASPADISMUS, see *Anaspadæus*.

ANAS'PASIS, *Anaspassm'us*, from *ανασπασω*, 'I contract.' *Retrac'tio*. Contraction, especially of

the bowels. The condition is called *Anaspad'ia*, *Anaspad'isis*, and *Anaspad'ismus*.—Hippocrates.

ANASPASMUS, *Anaspassus*.

ANASSA, *Bromelia ananas*.

ANASTALTICA, *Styptics*.

ANASTASIS, *Anarrhœa*. Also, restoration from sickness. Convalescence.

ANASTŒCHEIO'SIS, from *ava*, 'again,' and *σπαινω*, 'element.' *Re'element'io*. Resolution of a body or its parts into their elements,—Galen.

ANASTOMO'SIS, from *ava*, 'with,' and *σπασω*, 'a mouth.' *Inosculat'io seu Reu'nio vaso'rum*, *Exanastomosis*, *Concur'sus*, (F.) *Abouchement*. Communication between two vessels. By considering the nerves to be channels, in which a nervous fluid circulates, their communication likewise has been called *Anastomosis*. By means of anastomoses, if the course of a fluid be arrested in one vessel, it can proceed along others.

ANASTOMOSIS ANEURISMATICA, *Telangiectasia*—a. *Jacobson's*, see *Petrosal ganglion*.

ANASTOMOT'IC, *Anastomot'icus*, (F.) *Anastomotique*. Same etymon as *anastomosis*. Belonging or relating to *anastomosis*.

ANASTOMOT'ICS, *Anastomot'ica*. Same etymon. Certain medicines were formerly so called, which were believed to be capable of opening the mouths of vessels, as asperients, diuretics, etc.

ANASTOMOT'ICUS MAGNUS (RAMUS), (F.) *Artère collatérale interne*, *A. collatérale du coude*, is a branch of the brachial artery, which comes off a little above the elbow, and bestows branches to the brachialis internus, to the under edge of the triceps, and to the muscles, ligaments, etc., about the elbow joint. See, also, *Articular arteries of the knee*.

ANASTROPHE UTERI, *Inversio uteri*.

ANATASIS, Extension.

ANATHEMA, *Tabula votiva*.

ANATHERIUM MURICATUM, *Andropogon muricatus*.

ANATHOMIA, *Anatomy*.

ANATHYMIAMA, *Anathymiasis*.

ANATHYMI'ASIS, *Anathymiamia*, from *ava*, 'upwards,' and *θυμα*, 'fumigation.' *Ede'ma fugax*, *Ede'ma spas'ticum*, *Ede'ma hyster'icum*, *Anasar'ca hyster'icum*. An uncertain and transient swelling or inflation, said to have been observed at times in nervous and hysterical persons. It also means *Exhalation*, *Fumigation*, and *Hypochochriasis*.

ANATOLE UNGUIUM, see *Nail*.

ANATOME, *Anatomy*—a. *Animata*, *Physiology*.

ANATOMIA, *Anatomy*—a. *Animalis*, *Zootomy*—a. *Comparata*, *Zootomy*—a. *Comparativa*, *Zootomy*—a. *Viva*, *Physiology*.

ANATOMIE, *Anatomy*—a. *Chirurgicale*, see *Anatomy*—a. *des Régions*, see *Anatomy*.

ANAT'OMISM, *Anatomism'us*. Same etymon as *Anatomy*. The doctrine of those who look into the arrangement of parts, to explain all the phenomena of the organism.

ANAT'OMIST, *Anatom'icus*. One who occupies himself with *anatomy*. One versed in *anatomy*.

ANAT'OMY, *Anat'omē*, *Anatom'ia*, *Anathom'ia*, *Prosect'io*, from *ava*, and *τεμνω*, 'to cut,' (F.) *Anatomic*. The word *Anatomy* properly signifies *dissection*; but it has been appropriated to the study and knowledge of the number, shape, situation, structure, and connection—in a word, of all the apparent properties of organized bodies. *Anatomy* is the science of organization. Some have given the term a still more extended acceptation, applying it to every mechanical decomposition, even of inorganic bodies. Thus, *Cryt'allography* has been termed the *Anatomy* of

crystallized minerals. Anatomy has also been called *Morphol'ogy*, *Somatol'ogy*, *Somatol'omy*, *Organol'ogy*, etc. It assumes different names, according as the study is confined to one organized being, or to a species or class of beings. Thus, *Androt'omy*, or *Anthropot'omy*, or *Anthrop'ography*, or *Anthroposomatol'ogy*, is the *Anatomy of Man*; *Zoölogy*, that of the other species of the animal kingdom; and *Vet'inary Anat'omy* is the anatomy of domestic animals; but when the word is used abstractly, it means *Human Anatomy*, and particularly the study of the organs in a physiological or healthy state. *Physiolog'ical Anatomy* is occasionally used to signify the kind of anatomy which investigates structure with a special view to function. The Anatomy of the diseased human body is called *Patholog'ical* or *Morbid Anatomy*, and when applied to Medical Jurisprudence, *Forensic Anat'omy*. Several of the organs possessing a similarity of structure, and being formed of the same tissues, they have been grouped into Systems or Genera of Organs; and the study of, or acquaintance with, such systems, has been called *General Anat'omy*, *Histol'ogy*, or *Morphol'omy*, whilst the study of each organ in particular has been termed *Descriptive Anatomy*, *Anthropomorphol'ogy*. *Histology* is, however, more frequently applied to the *Anatomy of the Tissues*, which is called, also, *Textural* and *Microscop'ic Anatomy*, *Micranatom'ia*, see *Histology*. *Descriptive Anatomy* has been divided into *Skeletol'ogy*, which comprises *Osteol'ogy* and *Syndesmol'ogy*; and idto *Sarc'ol'ogy*, which is subdivided into *Myol'ogy*, *Neur'ol'ogy*, *Angiol'ogy*, *Adenol'ogy*, *Splanchnol'ogy*, and *Dermol'ogy*. *Sur'gical Anat'omy*, *Medico-Chirurgical Anat'omy*, *Topograph'ical Anat'omy*, *Re'gional Anat'omy*, (F.) *Anatomie Chirurgicale*, *A. des Régions*, is the particular and relative study of the bones, muscles, nerves, vessels, etc., with which it is indispensable to be acquainted before performing operations. *Compar'ative Anat'omy* is the comparative study of each organ, with a view to an acquaintance with the modifications of its structure in different animals or in the different classes of animals. *Transcendent'al* or *Philosoph'ical Anatomy* inquires into the mode, plan, or model upon which the animal frame or organs are formed; and *Artif'icial Anat'omy* is the art of modelling and representing, in wax or other substance, the different organs or different parts of the human body, in the sound or diseased state. *Phytol'omy* is the anatomy of vegetables, and *Pictor'ial Anatomy*, anatomy artistically illustrated.

ANATOMY, see *Skeleton*—*a. Artificial*, see *Anatomy*—*a. Comparative*, see *Anatomy*, *Zootomy*—*a. Descriptive*, see *Anatomy*—*a. Forensic*, see *Anatomy*—*a. General*, see *Anatomy*—*a. Human*, see *Anatomy*—*a. of Man*, see *Anatomy*—*a. Medico-Chirurgical*, see *Anatomy*—*a. Microscopic*, see *Anatomy*—*a. Morbid*, see *Anatomy*—*a. Pathological*, see *Anatomy*—*a. Pathological, microscopic*, see *Histology*—*a. Philosophical*, see *Anatomy*—*a. Physiological*, see *Anatomy*—*a. Pictorial*, see *Anatomy*—*a. Practical*, see *Dissection*—*a. Regional*, see *Anatomy*—*a. Surgical*, see *Anatomy*—*a. Textural*, see *Anatomy*—*a. Topographical*, see *Anatomy*—*a. Transcendental*, see *Anatomy*—*a. Veterinary*, see *Anatomy*.

ANATON, Soda.

ANATREPSIS, *Restauratio*.

ANATRESIS, *Perforation*, *Trepanning*.

ANATRIBE, *Friction*.

ANATRIPSIS, *Friction*.

ANATRIPSOL'OGY, *Anatrispolog'ia*, *Anatrispolog'ia*, from *ανατριψις*, 'friction,' and *λογος*, 'a discourse.' A treatise on friction as a remedy.

ANATRIPTOLOGIA, *Anatrispology*.

ANATRON, *Natrum*, *Soda*.

ANAT'ROPE, from *ανα*, 'upwards,' and *τροπε*, 'I turn.' Subversion. A turning or subversion or inverted action of the stomach, characterized by nausea, vomiting, etc.—Galen. We still speak of the stomach turning against any thing.

ANAUDIA, *Catalepsy*, *Mutitas*.

ANAXYRIS, *Rumex acetosa*.

ANAZESIS, *Ebullition*.

ANAZOTURIA, see *Urine*.

ANCHA, *Haunch*.

ANCHELOPS, *Ægilops*.

ANCHONE, *Angone*.

ANCHORALIS PROCESSUS, *Coracoid*.

ANCHUSA ANGUSTIFOLIA, *A. Officialis* —*a. Incarnata*, *A. Officialis*—*a. Lycopsoides*, *A. Officialis*.

ANCHU'SA OFFICINALIS, *A. Angustifolia* seu *Incarnata* seu *Lycopsoïdes*, *Alechna*, *Lingua Bovis*, *Buglossum sylvestre*, *Offic'inal* or *Garden Al'kanet* or *Bugloss*, (Old Eng.) *Langdebeß*; *Ord. Boraginæ*. *Sex. Syst. Pentandria Monogynia*. (F.) *Buglose*. A native of Great Britain. The herb was formerly esteemed as a cordial in melancholia and hypochondriasis; but it is now rarely used. It is also called *Buglossa*, *Buglossum angustifolium majus*, *B. vulgarē majus*, *L. sativum*.

ANCHU'SA TINCTORIA, *Alechna spuria*, *Dyer's Bugloss*, *Anelimum*, *Buglossum Tinctorum*, *Lithospermum villosum*, *Dyer's Al'kanet*, (F.) *Orcanette*. A European plant. The medical properties are equivocal. It is used to give a beautiful red colour to ointments.

ANCHYLOSIS, *Ankylosis*.

ANGISTRON, *Hamulus*.

ANCLE, *Astragalus*, *Malleolus*.

ANCLER, *Malleolus*.

ANCL'ET, *Malleolus*.

ANCLIFF, *Malleolus*.

ANCLOWE, *Malleolus*.

ANCOLIE, *Aquilegia vulgaris*.

ANCON, *Elbow*, *Olecranon*.

ANCONAD, see *Anconal Aspect*.

ANCONAGRA, *Pechyagra*.

ANCON'AL, from *αγκων*, 'the elbow.' Relating, or appertaining to the elbow or the olecranon. **ANCONAL ASPECT**. An aspect towards the side on which the ancon or elbow is situated.—Barclay. *Anconad* is used by the same writer adverbially, to signify 'towards the anconal aspect.'

ANCONÉ, *Anconeus*.

ANCONÉ'US, from *αγκων*, 'the elbow.' A term once applied to every muscle attached to the olecranon. Winslow distinguished four—the *great*, *external*, *internal*, and *small*; the first three being portions of the same muscle, the *triceps brachialis*. The last has, alone, retained the name. It is the *Anconeus minor* of Winslow, the *Anconeus* seu *Cubitalis* *RIOLANT* of Douglas, the *Epicondylo-Cubitalis* of Chaussier, the *Brevis Cubiti*, (F.) *Anconé*, and is situate at the upper and back part of the fore-arm. It arises from the external condyle of the os humeri, and is inserted into the posterior edge of the upper third of the ulna. Its use is to aid in the extension of the fore-arm.

ANCONÉUS EXTERNUS, see *Triceps extensor cubiti*—*a. Internus*, see *Triceps extensor cubiti*—*a. Major*, see *Triceps extensor cubiti*.

ANCTERES. *Fibulae* or *Clasps* by which the lips of wounds were formerly kept together.—Celsus, Galen.

ANCTERIASMUS, *Infibulation*.

ANCU'BITUS, *Petrifac'tio*. An affection of the eye, in which there is a sensation as if sand were irritating the organ.

ANCUNNUENTÆ. A name formerly given to menstruating females.

ANCUS, *Ancus*, from *αγκων*, 'the elbow.' One who cannot extend his arms completely.

Also, the deformity resulting from a luxation of the humerus or fore-arm.—Hippocrates.

ANCYLE. Ankylosis.

ANCYLOBLEPHARON, *Ankyloblepharon*.

ANCYLOCHEILIA, *Ankylocheilia*.

ANCYLOCOLPUS, *Colpactesia*.

ANCYLOGORE, *Ankylogore*.

ANCYLODERE, *Torticollis*.

ANCYLODERIS, *Torticollis*.

ANCYLODONTIA, *Ankylodontia*.

ANCYLOGLOSSIA, *Ankyloglossia*.

ANCYLOMELE, *Ankylomele*.

ANCYLOMERISMUS, *Ankylomerismus*.

ANCYLORRHINIA, *Ankyloorrhinia*.

ANCYLOSIS, *Ankylosis*.

ANCYLOTIA, *Ankylotia*.

ANCYLOTOMUS, *Ankylotomus*.

ANCYRA. Hook.

ANCYROID CAVITY, Digital cavity.

ANCYROIDES PROCESSUS, *Coracoid*.

AND. Breath.

ANDA. Native name of a tree of Brazil—*Anda Come'sii* seu *Brasilien'sis* seu de *Pison*, *Andaessu*, *Anda-açu*, *Joanne'sia princeps*. Ord. *Euphorbiaceæ*. *Sex. Syst.* *Monœcia* *Monadelphia*. An oil—Oil of *Anda*—is obtained from the seeds by pressure, 50 to 60 drops of which act as a cathartic. The fruit is an oval nut, containing two seeds. These have the taste of the chestnut; but are strongly cathartic, and even emetic. The shell is astringent, and is used as such in diarrhœa, etc.

ANDE. Breath.

ANDELY, MINERAL WATERS OF. *Andely* is in France, near Gisors, and eight leagues from Rouen. The water is cold, and a weak chalybeate. It is used in chlorosis and abdominal obstructions.

ANDERSON'S PILLS, see *Pilulæ Aloes* et *Jalapæ*.

ANDIRA IBAI, *Geoffræa Vermifuga*—a. *Inermis*, *Geoffræa inermis*—a. *Racemosa*, *Geoffræa inermis*—a. *Surinamensis*, *Geoffræa Surinamensis*.

ANDRACHAHARA, *Sempervivum tectorum*.

ANDRACHNE, *Arbutus unedo*, *Portulaca*.

ANDRALOGOMÈLE, from *ανρ*, *ανρὸς*, 'a man,' a, priv., *λογος*, 'reason,' and *μυλον*, 'a domestic animal.' A name given by Malacarne to a monster in which he supposed the presence of the body of a man with the limbs of a brute.

ANDRANATOM'IA, *Andranat'omē*, *Androtom'ia*, *Androt'omē*, *Anthropot'omy*, from *ανρ*, genitive *ανρὸς*, 'a man,' and *τεμνειν*, 'to cut.' The anatomy of man.

ANDRI'A. Adult age. Manhood.

ANDRI'A M'ULIER, *Mulier Hermaphrodit'ica*. A female hermaphrodite.

ANDROGEN'IA, from *ανρ*, *ανρὸς*, 'a man,' and *γενειν*, 'generation.' The procreation of males.—Hippocrates.

ANDROG'YNUS, from *ανρ*, *ανρὸς*, 'a man,' and *γυν*, 'a woman.' A hermaphrodite. An effeminate person.—Hippocrates.

ANDROLEPSIA, *Conception*.

ANDROMANIA, *Nymphomania*.

ANDROM'EDA ARBO'REA, *Sorrel Tree*, *Sour Tree*, *Sour Wood*, *Elk Tree*, *Elk Wood*, *Sorrel Wood*, *Sour Leaf*, (F.) *Andromède*, *Andromédier*. A small indigenous tree; Ord. *Ericaceæ*, *Sex. Syst.* *Decandria* *Monogynia*; found in the Alleghany mountains and the hills and valleys diverging from them, as far as the southern limits of Georgia and Alabama; but seldom north of Virginia. The sour leaves are refrigerant and

astringent, and have been used to make a kind of lemonade, which has been given in fevers.

ANDROMEDA MARIA'NA, *Beard-leaved Moorwort*, *Kill-lamb*, *Lamb Killer*, *Staggerbush*. A decoction of this American plant is said to have been successfully employed as a wash, in a disagreeable affection—not uncommon amongst the slaves in the southern parts of the United States—called the *Toe Itch*, and *Ground Itch*.—Barton.

ANDROMEDA NIT'IDA, *Sourwood*, *Sorrel tree*, indigenous; has properties similar to those of *A. arborea*.

ANDROPOGON BICORNIS, *Juncus odoratus*—a. *Calamus aromaticus*, see *Oleum graminis Indici*—a. *Citratus*, *Juncus odoratus*—a. *Citriodorus*, *Juncus odoratus*, *Nardus Indica*.

ANDROPOGON MURICA'TUS, (*ανρ*, *ανρὸς*, 'a man,' and *μυρον*, 'a beard.') *A. squarro'sis*, *Phal'aris zizanioides*, *Agrostis verticillat'a*, *Anathe'rum murica'tum*, *Vetiveria odorat'a*, *Vittiveray*, *Cuscuta*, *Khus-Khus*; *Nat. Ord.* *Graminæ*. The root of this plant, from Bombay, is aromatic and bitterish. It is used as a perfume. It has been used as an excitant and diaphoretic.

ANDROPOGON NARDUS, *Calamus Alexandrinus*, *Nardus Indica*—a. *Schœnanthus*, *Juncus odoratus*—a. *Squarrosus*, *A. Muricatus*.

ANDROSACE, *Umbilicus marinus*—a. *Matthioli*, *Umbilicus marinus*.

ANDROSÆMUM, *Hypericum perforatum*.

ANDROTOMY, *Andranatomia*.

ANDRUM. An East India word, latinized by Kæmpler, signifying a kind of elephantiasis of the scrotum, endemic in southern Asia.

ANÉANTISSEMENT (F.), *Virium extinctio*. This word is often employed hyperbolically, by patients in France, to signify excessive fatigue, debility or syncope.

ANEBIUM, *Ancusa tinctoria*.

ANEBUS, *Impuber*.

ANECPYETUS, from *αν*, for *ανευ*, 'without,' and *πυα*, 'I promote suppuration.' That which does not suppurate, or is not likely to suppurate.

ANEGERTICA, from *ανεγειρω* (*ava* and *γειρω*), 'I awaken.' The art of resuscitating the apparently dead.

ANEILE'MA, *Aneile'sis*, from *ανειλεω* (*ava* and *ειλεω*), 'I roll up.' Applied particularly to the motion of air in the intestines and the tormina accompanying it.—Hippocrates.

ANEILESIS, *Aneilema*.

ANEMIA, *Anæmia*.

ANEMO'NĒ, *Wind Flower*: from *ανemos*, 'the wind,' because it does not open its flowers until blown upon by the wind.

ANÉMONE DES BOIS, *Anemone nemorosa*.

ANEMONE COLLINA, *A. Pulsatilla*—a. *Hepatica*, *Hepatica triloba*—a. *Intermedia*, *A. Pulsatilla*.

ANEMONE LUDOVICIANA, *A. patens*.

ANEMO'NĒ NEMORO'SA, *Ranunculus albus* seu *nemoro'sis*, *Wood Anemone*, (F.) *Anémone des bois*. Ord. *Ranunculacææ*. The herb and flowers are poisonous, acrid, and corrosive. They have been used as rubefaciants.

ANEMO'NĒ PATENS, *A. Ludoviciana*, *Pulsatilla patens*; indigenous; from Illinois and Wisconsin to the Rocky Mountains, is supposed to possess similar properties.

ANEMO'NĒ PRATEN'SIS, *A. Sylvæ'tris*, *Pulsatilla nigricans* seu *pratensis*. This plant has similar properties with the last. It is also called *Meadow Anemone*, (F.) *Pulsatille noire*, *F. des prés*.

ANEMO'NĒ PULSATIL'LA, *A. Collina* seu *Intermedia* seu *Pratensis* seu *Rubra*, *Pulsatilla vulgaris*, *Herba ventis*, *Nola culina'ria*, *Pasque flower*, (F.) *Coquelourde*, possesses like properties.

ANEMO'NĒ RUBRA, *A. Praten'sis*—a. *Rue-leaved*,

Thalictrum anemonoides—a. *Sylvestris*, A. *Pratensis*.

ANEMONY, *Anemone hepatica*—a. Meadow, *Anemone pratensis*—a. Wood, *Anemone nemorosa*.

ANEMOS, Wind.

ANENCEPHALIA, see *Anencephalus*.

ANENCEPHALOHÆMIA, from *av*, privative, *εγκεφαλος*, 'encephalon,' and *αἷμα*, 'blood.' Defect of blood in the brain. Syncope.

ANENCEPHALONEURIA, *Anencephaloneuria*, from *av*, priv., *εγκεφαλος*, 'encephalon,' and *νευρον*, 'nerve.' Want of nervous action in the encephalon.

ANENCEPHALOTROPHIA, from *av*, privative, *εγκεφαλος*, 'the encephalon,' and *τροφή*, 'nourishment.' Atrophy of the encephalon.

ANENCEPHALUS, from *av*, privative, and *εγκεφαλος*, 'brain.' A monster devoid of brain.—Bonetus, G. St. Hilaire. Also, one that has a part only of the brain;—*Paracephalus*. The condition has been called *Anencephalia*. A weak, silly person.—Hippocrates.

ANENERGESIA, Debility.

ANENERGIA, Debility.

ANENTÉRONERVIE SATURNINE, see Palsy, lead.

ANEPISCHEISIS, Incontinentia.

ANEPITHY'IA, from *av*, priv., and *ἐπιθυμία*, 'desire.' Many nosologists have used this word for a loss of the appetites; as of those of hunger, thirst, venery, &c.

ANEPITHY'IA CHLOROSIS, Chlorosis.

ANER, *avv*, genitive *avpos*. A man.

ANERETHIS'IA, *Inirritabil'itas*, from *av*, priv., and *ερεθισ*, 'irritability.' Defect of irritability.—Swediaur.

ANERVISMIE, Paralysis.

ANERYTHROP'SIA, from *av*, priv., *ερυθρος*, 'red,' and *οφία*, 'vision.' Defective vision, which consists in an incapability of distinguishing red.

ANESIS, Remission.

ANESTHÉSIE, Anæsthesia.

ANESTHÉSIE EXTATIQUE. The aggregate of phenomena of impaired feeling produced especially by the manipulations of the animal magnetizer.—Andral.

ANESTHETIC, Anæsthetic.

ANESTHÉSIQUE, Anæsthetic.

ANESTHETIZATION, Anæsthetization.

ANESON, Anethum.

ANESUM, *Pimpinella anisum*.

ANET, Anethum.

ANETH, Anethum graveolens.

ANETHUM, *Aneson*, *Anethon*, *Anethum Fœniculum* seu *Segetum* seu *Piperitum*, *Fœniculum*, *F. Dulce* seu *Officinale* seu *vulgarè*, *Ligusticum fœniculum*, *Faniculum*, *Fennel* or *Finckle*, *Mar'athrum*, *Anet*, *Sweet Fennel*, (Prov.) *Spingel*, (F.) *Fenouil* or *Anis doux*. Ord. Umbelliferae. *Sex. Syst.* Pentandria Digynia. The fruit, *Fœniculum* (Ph. U. S.), has an aromatic odour, and warm, sweetish taste. It is carminative. The oil—*Oleum Fœniculi*—is official in the Ph. U. S. The root is said to be pectoral and diuretic.

ANETHUM FœNICULUM, Anethum.

ANETHUM GRAVEOLENS, *Anethum*, A. *horten'se*, *Pastinaca Anethum* seu *Graveolens*, *Fer'ula Graveolens*, *Dill*, (F.) *Aneth*, *Fenouil puant*. A native of the south of Europe. The seeds are stimulant and carminative. A distilled water—*Aqua anethi*, *Dill-water*—is official in the London and Edinburgh Pharmacopœias. Dose, gr. xv to ʒi.

Oleum Anethi, *Oil of Dill*, (F.) *Huile d'Aneth*, possesses the carminative properties of the plant. ANETHUM PASTINACA, *Pastinaca Sativa*—a. *Piperitum*, *Anethum*—a. *Segetum*, *Anethum*.

ANETICUS, Anodyne.

ANETON, Anethum.

ANETUS, Intermittent fever—a. *Quartanus*, *Quartan*—a. *Quotidianus*, *Quotidian*—a. *Tertianus*, *Tertian* fever.

ANEURALGICON, from *a*, privative, *νευρον*, 'nerve,' and *αλγος*, 'pain.' A name given by Dr. C. T. Downing to an instrument used by him to allay pain in nerves. It is a kind of fumigating apparatus, in which dried narcotic and other herbs are burnt, the heated vapour being directed to any part of the body.

ANEURIA, Paralysis.

ANEURISM, *Aneurys'ma*, *Aneurys'mus*, *Aneuris'ma*, *Cedma*, from *ανευρειν*, (*ava*, and *ερευειν*), 'to dilate or distend.' *Dilatatio Arteriarum*, *Ectasia*, *Embryos'ma*, *Exangi'a aneuris'ma*, *Arterieurys'ma*, *Arterieurys'ma*, *Hæmatocoele arterio'sa*, *Abcessus spirituosus*, *Arterioec'tasis*, (F.) *Anévrysm*, *Aneurisme*. Properly, Aneurism signifies a tumour, produced by the dilatation of an artery; but it has been extended to various lesions of arteries, as well as to dilatations of the heart.

There are various kinds of aneurism. The following are the chief:

I. When the blood, which forms the tumour, is enclosed within the dilated coats of the artery. This is the TRUE ANEURISM, *Aneurys'ma verum*, *Hernia Arteriarum*, (F.) *Anévrysm vrai*.

II. When the blood has escaped from the opened artery, it is called SPURIOUS or FALSE ANEURISM, *Aneuris'ma spurium*, *Ruptura Arteriae*, *Arteriorrhex'is*, *Arteriodial'ysis*, *Echymoma arteriosum*, (F.) *Anévrysm faux*. The latter is divided into three varieties:

1. *Diffused False Aneurism*, (F.) *Anévrysm faux*, *primitif*, *diffus*, *noncirconscrit* ou *par infiltration*, which occurs immediately after the division or rupture of an artery, and consists of an extravasation of blood into the areolar texture of the part.

2. *Circumscribed False Aneurism*, (F.) *Anévrysm faux circonscrit*, *circonscrit* ou *par épanchement*, *enkysté* ou *sacciforme*, *tumeur hémorrhagiale circonscrite*, in which the blood issues from the vessel some time after the receipt of the wound, and forms itself a sac in the neighbouring areolar membrane.

3. *Aneurism by Anastomosis*, *Varicose* or *Circoid Aneurism*, *Phlebarteriodial'ysis*, *Aneurys'ma venoso-arteriosum*, A. *varicosum*, (F.) *Anévrysm par anastomose* ou *variqueux*, A. *par érosion*, A. *de Pott*, A. *des plus petites artères*, which arises from the simultaneous wounding of an artery and vein;—the arterial blood passing into the vein, and producing a varicose state of it.

III. MIXED ANEURISM, (F.) *Anévrysm mixte*, is that which arises from the dilatation of one or two of the coats, with division or rupture of the other. Some authors have made two varieties of this:

1. *Mixed external Aneurism*, where the internal and middle coats are ruptured, and the areolar is dilated.

2. *Mixed internal Aneurism*, in which the internal coat is dilated, and protrudes, like a hernial sac, through the ruptured middle and outer coats. This variety has been called *Aneurys'ma Herniam Arteriae sistens*.

Aneurisms have been termed *traumatic* or *exogenous*, and *spontaneous*, according as they may have been caused by a wound, or have originated spontaneously. The latter, when originating from lesions of the inner coats of arteries, have been termed *endogenous*. They have also been divided into *internal* and *external*.

The internal aneurisms are situate in the great splanchnic cavities, and occur in the heart and great vessels of the chest, abdomen, &c. Their diagnosis is difficult, and they are often inaccessible to surgical treatment.

The *external aneurisms* are situate at the exterior of the head, neck, and limbs, and are distinctly pulsatory.

Aneurisms, especially the internal, may be combated by a debilitating treatment, on the plan of Valsalva, which consists in repeated blood-letting, with food enough merely to support life. In external aneurism, the artery can be obliterated. This is usually done by applying a ligature above the aneurismal tumour.

ANEURISM, DISSECTING, is one in which, owing to rupture of the inner and middle coats of an artery, the blood makes itself a channel between these coats and the outer coat.

In many cases, the lesion appears to consist in a separation of the laminae of the middle coat, between which the blood forms itself a channel.

ANEURISMS OF THE HEART, *Cardiœchi*, *Cardiœurysma*, (F.) *Anévrysmes du cœur*, have been divided into *active* and *passive*. The former can scarcely be esteemed aneurisms, as they most commonly consist of increased thickness of the parietes of the heart, which diminishes its cavity instead of increasing it. The term *Hypertrophy of the heart* better indicates their character. *Passive aneurism*, *Cardiœctosis*, on the contrary, is attended with extenuation of the parietes of the organ, and enlargement of the cavities. The physical signs of *dilatation of the heart* are the following:—The action of the heart is not visible, and no impulse is conveyed to the hand. On percussion, there is a loss of resonance over a larger surface than usual, but the dulness is much less intense than that which accompanies hypertrophy. On auscultation, the action of the heart is only slightly felt, and communicates at once the impression of its diminished power. The impulse is feebler than usual. Both sounds are widely transmitted over the thorax, and are not much fainter at a distance from their point of origin.

Partial or true aneurism of the Heart—*Cardiœctasis partialis*, *Aneurysma consecutivum cordis*—is sometimes seen; rarely, however.

The name *Aneurism of the Valves of the Heart* has been given to pouch-like projections of the valves into the auricles.

ANEURISM BY ANASTOMOSIS, see Aneurism—a. Brador's operation for, see Brador—a. Circoid, see Aneurism—a. Endogenous, see Aneurism, and Endogenous—a. Exogenous, see Aneurism, and Exogenous—a. External, see Aneurism—a. False, see Aneurism—a. False, circumscribed, see Aneurism—a. False, diffused, see Aneurism—a. Internal, see Aneurism—a. Mixed, see Aneurism—a. Mixed, external, see Aneurism—a. Mixed, internal, see Aneurism—a. Spontaneous, see Aneurism—a. Spurious, see Aneurism—a. Traumatic, see Aneurism—a. True, see Aneurism—a. Valsalva's method of treating, see Aneurism—a. Varicose, see Aneurism.

ANEURISMA, Aneurism.

ANEURISMAL, *Aneurysmal*, *Aneurismat'ic*, *Aneurysmat'icus*, *Aneurismal'is*. That which belongs to Aneurism.

ANEURISMAL SAC or CYST, (F.) *Sac ou Kyste anévrysmal*, is a sort of pouch, formed by the dilatation of the coats of an artery, in which the blood, forming the aneurismal tumour, is contained.

ANEURISMATIC, Aneurismal.

ANEURYSM, Aneurism.

ANEURYSMA, Aneurism—a. Cordis activum, Heart, hypertrophy of the—a. Herniam arterie sistens, see Aneurism—a. Spurium, see Aneurism—a. Varicosum, see Aneurism—a. Venoso-arteriosum, see Aneurism—a. Verum, see Aneurism.

ANEURYSME, Aneurism.

ANEURYSMUS, Aneurism, Dilatation.

ANEVRIA, Paralysis.

ANÉVRYSMÉ. Aneurism—a. de l'Arterie, Aorteurysma—a. de Poit, see Aneurism—a. des plus petites Artères, see Aneurism—a. Circoscrit, see Aneurism—a. Diffus, see Aneurism—a. Énhysté, see Aneurism—a. Fanc, see Aneurism—a. Fanc consensitif, see Aneurism—a. Miète, see Aneurism—a. par Anastomose, see Aneurism—a. par Épanchement, see Aneurism—a. par Érosion, see Aneurism—a. par Infiltration, see Aneurism—a. Primitif, see Aneurism—a. Sacciforme, see Aneurism—a. Variqueux, see Aneurism—a. Vrai, see Aneurism.

ANEYS, see Pimpinella anisum.

ANFION, Maslach.

ANFRACUOSITÉS CÉRÉBRALES, Anfractuosités, cerebral—a. Ethmoïdales, see Anfractuosity.

ANFRACUOSITY, *Anfractus*, *Sulcus*, from *am*, 'around,' and *frangere*, *fractus*, 'to break.' A groove or furrow. Used in anatomy to signify sinuous depressions or *sulci*, of greater or less depth, like those which separate the convolutions of the brain from each other. These

ANFRACUOSITIES, CEREBRAL, *Anfractus Cerebri*, (F.) *Anfractuosités Cérébrales*, are always narrow, and deeper at the upper surface of the brain than at its base; and are lined by a prolongation of the pia mater.

The Ethmoid Cells are, sometimes, called *Anfractuosités ethmoïdales*.

ANFRACTUS, Anfractuosity—a. Cerebri, Anfractuosités, (cerebral.)

ANGECTASIA, Angiectasis.

ANGEIAL, Vascular.

ANGIECTASTIA, Angiectasis.

ANGIECTASTIS, Angiectasis.

ANGIECTOMA, Angiectasis.

ANGIEOGRAPHY, *Angiographia*, *Angiograph'ia*, from *αγγειον*, 'a vessel,' and *γραφω*, 'a description.' The anatomy of the vessels.

ANGIEOHYDROGRAPHY, *Angiohydrographia*, *Angieondrographia*, *Angieohydrograph'ia*, *Hydrangiograph'ia*, from *αγγειον*, 'a vessel,' *ὕδωρ*, 'water,' and *γραφω*, 'I describe.' A treatise on the lymphatics.

ANGIEOHYDROTOMY, *Angiohydrotomy*, *Angieondrotomy*, *Angieohydrotom'ia*, *Hydrangioto'm'ia*, from *αγγειον*, 'a vessel,' *ὕδωρ*, 'water,' and *τομω*, 'to cut.' Dissection of the lymphatics.

ANGIOLEUCITIS, *Angioleucit'is*, *Angiolymphit'is*, *Lymphangit'is*, *Lymphangit'is*, *Lymphangit'is*, *Hydrangit'is*, *Lymphit'is*, *Lymphit'is*, *Inflammat'io vasorum lymphaticorum*, from *αγγειον*, 'a vessel,' *λευκος*, 'white,' and *itis*, inflammation. (F.) *Inflammation des vaisseaux lymphatiques ou des tissus blancs*. Inflammation of the lymphatics; lymphatic or scrofulous inflammation.

ANGIEOLOGY, *Angiolog'ia*, *Angieolog'ia*, from *αγγειον*, 'a vessel,' and *λογος*, 'a discourse,' A discourse on the vessels. The anatomy of the vessels. It includes *Arteriolog'ia*, *Phlebolog'ia*, and *Angieohydrolog'ia*.

ANGIEOMALACIA, *Angiomalacia*, from *αγγειον*, 'a vessel,' and *μαλακία*, 'softening.' Mollescence or softening of vessels.

ANGIOMYCES, *Hæmatodes fungus*.

ANGION, Vessel.

ANGIEONDROGRAPHY, *Angieohydrograph'ia*.

ANGIENDROTOMY, *Angieohydrotomy*.

ANGIONOSUS, Angiopathia.

ANGIONUSUS, Angiopathia.

ANGIEOPATHIA, *Angiopath'ia*, *Angion'osis*, *Angionus'us*, *Angio'sis*, from *αγγειον*, 'a vessel,' and *παθος*, 'a disease,' Disease of the vessels.

ANGEIOPLANIA, *Angioplania*, from *αγγειον*,

ANGIDIOSPONGUS, Hæmatodes fungus.

ANGINA PECTORIS, *A. cordis*, *Sternalgia*,
Asthma spastico-arthritis incoⁿsistens, *Asthma*
diaphragmaticum, *Arthritis diaphragmatica*,
Orthopne^a cardiaca, *Sternodynia syncope^{tica}*
et palpitans, *S. syncope^{alis}*, *Cardingmus cordis*
sinistri, *Asthenia pectoralis*, *Angor pectoris*,
Stenocardia, *Diaphragmatic^e gout*, *Asthma con-*
vulsivum, *Asthma arthriticum*, *Cardioneuralgia*,
Neuralgia brachiothoracica, *Hyperæsthesia*
plicæ cardi^{ac}i, *A. dolorifica*, *Syncope angio-*
ca seu angens, *Cardiodyne spasmodica inter-*
mittens, *Pnigophobia*, *Prunella*, *Suspirium*
cardiacum, *Pneumonalgia*, *Suffocative Breast-*
pang, (F.) *Angine de Poitrine*, *Névrose du Cœur*.
 A disease, the precise pathology of which is not
 known. The principal symptoms are, violent
 pain about the sternum, extending towards the
 arms, anxiety, dyspnoea, and sense of suffoca-
 tion. It is an affection of great danger, and is
 often connected with ossification, or other morbid
 condition of the heart. It appears to be neurop-
 athic, and has been termed *Neuralgia of the Heart*.
 Some, however, employ this last term for an
 acutely painful intermittent affection of the heart,
 which seems to differ from angina pectoris more

in regard to the small number of parts which are drawn into morbid consent with the affected cardiac nerves, than in regard either to its nature or appropriate treatment. The most powerful stimulating and narcotic antispasmodics are required during the prooxysm.

ANGINA PELLICULARIS. *A. maligna*, *Diphtheritis of the throat*. A name given to those inflammations about the throat, in which exudations or false membranes are thrown out during the phlogosis of the mucous membranes. *Aphthæ*, *Tracheitis*, when accompanied with the membraniform exudation, are, with some, examples of diphtheritic inflammation.

ANGINA PERNICIOSA. *Cynanche trachealis*—*a. Pestilentialis*, *Pharyngitis*, *diphtheritic*—*a. Polyposa*, *Cynanche trachealis*—*a. Polyposa* seu *Membranacea*, *Cynanche trachealis*—*a. Pseudo-membranosa*, *Pharyngitis*, *diphtheritic*—*a. Pulposa*, *Cynanche trachealis*—*a. Sanguinea*, *Cynanche tonsillaris*.

ANGINA SICCA, (F.) *Angine Sèche*, is a chronic inflammation of the pharynx, with a distressing sense of dryness and heat, in chronic diseases of the stomach and lungs. See *Padauchoue*.

ANGINA SIMPLEX. *Isthmitis*.

ANGINA SQUIRROSA, (F.) *Angine squirreuse*, consists in difficulty of deglutition, caused by scirrhus disorganization of the pharynx or œsophagus, or by enlarged tonsils.

ANGINA STRANGULATORIA, *Cynanche trachealis*—*a. Strepitosa*, *Cynanche trachealis*—*a. Suffocatoria*, *Cynanche trachealis*—*a. Synochalis*, *Cynanche tonsillaris*—*a. Thyreoidea*, *Thyreoiditis*—*a. Tonsillaris*, *Cynanche tonsillaris*—*a. Trachealis*, *Cynanche trachealis*—*a. Ulcerosa*, *Cynanche maligna*—*a. Uvularis*, *Staphyloedema*, *Uvulitis*—*a. Vera et Legitima*, *Cynanche tonsillaris*.

ANGINE GUTTURALE, *Cynanche tonsillaris*—*a. Laryngea*, *Laryngitis*—*a. Laryngea et tracheale*, *Cynanche trachealis*—*a. Laryngea adnata*, *Edema of the glottis*—*a. Esophagica*, *Esophagitis*—*a. Pharyngea*, *Cynanche parotidea*—*a. de Poitrine*, *Angina pectoris*—*a. Sèche*, *Angina sicca*—*a. Simplex*, *Isthmitis*—*a. Squirreuse*, *Angina Squirrosa*—*a. Tonsillaire*, *Cynanche tonsillaris*.

ANGINEUX, *Anginose*.

ANGINOSE, *Anginosus*, (F.) *Angineux*. Relating or appertaining to angina; as *Scarlatina anginosa*.

ANGIOCARDITIS, from *αγγειον*, 'a vessel,' and *carditis*, 'inflammation of the heart.' Inflammation of the heart and great vessels.

ANGIODIASTASIS, *Angiectasis*.

ANGIOGRAPHY, *Angiography*.

ANGIOHÉMIE, *Hyperæmia*.

ANGIOHYDROGRAPHY, *Angiohydrography*.

ANGIOHYDROTOMY, *Angiohydrotomy*.

ANGIOITIS, *Angitis*.

ANGIOLEUCITIS, *Angioleucitis*.

ANGIOLOGY, *Angiology*.

ANGIOLYMPHITIS, *Angioleucitis*.

ANGIOMALACIA, *Angiomalacia*.

ANGIOMYCES, *Hæmatodes fungus*.

ANGIONOSUS, *Angiopathia*.

ANGIONCUSUS, *Angiopathia*.

ANGIOPATHIA, *Angiopathia*.

ANGIOPLANIA, *Angioplania*.

ANGIOPLEROSIS, *Plethora*.

ANGIOPYRA, *Synocha*.

ANGIORRHAGIA, *Hæmorrhagia activa*.

ANGIORRHŒA, *Angiorrhœa*.

ANGIOSIS, *Angiopathia*.

ANGIOSTEGNOSIS, *Angiemprraxis*.

ANGIOSTENOSIS, *Angiemprraxis*.

ANGIOSTOSIS, *Angiostosis*.

ANGIOTROPHE, see *Torsion*.

ANGIOTELECTASIA, *Telangiectasia*.

ANGIOTENIC, *Angiotenic*, *Angiotenicus* seu *Angiotenicus*, from *αγγειον*, 'a vessel,' and *τενειν*, 'to extend.' An epithet given to inflammatory fever, owing to its action seeming to be chiefly exerted on the vascular system.

ANGIOTOMY, *Angiotomy*.

ANGLE, *ἄνγλος*, from *αγκυλος*, 'a hook.' The space between two lines which meet in a point.

ANGLE, FA'CIAL, pointed out by Camper, is formed by the union of two lines, one of which is drawn from the most prominent part of the forehead to the alveolar edge of the upper jaw, opposite the incisor teeth—the *facial line*—and the other from the meatus auditorius externus to the same point of the jaw. According to the size of the angle it has been attempted to appreciate the respective proportions of the cranium and face, and, to a certain extent, the degree of intelligence of individuals and of animals. In the white varieties of the species, this angle is generally 80°; in the negro not more than 70°, and sometimes only 65°. As we descend the scale of animals, the angle becomes less and less; until, in fishes, it nearly or entirely disappears. Animals which have the snout long, and facial angle small, such as the snipe, crane, stork, &c., are proverbially foolish, at least they are so esteemed; whilst intelligence is ascribed to those in which the angle is more largely developed, as the elephant and the owl. In these last animals, however, the large facial angle is caused by the size of the frontal sinuses:—so that this mode of appreciating the size of the brain is very inexact, and cannot be depended upon.

The following is a table of the angle in man and certain animals:

FACIAL ANGLES.

Man.....	from 68° to 83° and more.
Sapajou.....	65
Orang-Utang.....	56 to 58
Guenon.....	57
Maydrill.....	30 to 42
Coati.....	28
Pole-Cat.....	31
Pug-Dog.....	35
Mastiff.....	41
Hare.....	30
Ram.....	30
Horse.....	23

ANGLE, OCCIPITAL, OF DAUBENTON, is formed by a line drawn from the posterior margin of the foramen magnum to the inferior margin of the orbit, and another drawn from the top of the head to the space between the occipital condyles. In man, these condyles, as well as the foramen magnum, are so situate, that a line drawn perpendicular to them would be a continuation of the spine; but in animals they are placed more or less obliquely; and the perpendicular is necessarily thrown farther forward, and the angle rendered more acute.

ANGLE, OPTIC, (F.) *Angle optique*, is the angle formed by two lines, which shave the extremities of an object, and meet at the centre of the pupil.

ANGNAIL, *Hangnail*.

ANGOISSE, *Angor*.

ANGOLAM. A very tall Malabar tree, which possesses vermifuge properties.

ANGONE, properly *Ἀνχονή*, from *αγχω*, 'choke,' *Anadromæ*, *Hysteroptomæ*, *Hysteroptomæ*, *Profectio* *Falcium* seu *Uteri* *na* seu *Matricis*, *Strangulatio uterina*, *Suffocatio uterina* seu *hysterica*, *Globus* seu *Nodus hystericus*, *Orthopnœa hysterica*, *Dysphagia globosa* seu *hysterica*, *Nervous Quinsy*. A feeling of strangulation, with dread of suffocation. It is common

in hysterical females, and is accompanied with a sensation as if a ball arose from the abdomen to the throat.

ANGOR, Anguish, (F.) Angoisse. Extreme anxiety, accompanied with painful constriction at the epigastrium, and often with palpitation and oppression. It is frequently an unfavourable symptom.

ANGOR, Agony, Orthopnoea—a. Faucium, Isthmitis—*a.* Pectoris, Angina pectoris.

ANGOS, Bubo, Uterus, Vessel.

ANGOSTURE VRAIE, Cusparia febrifuga.

ANGOURION, Cucumis sativus.

ANGUIS, Serpent.

ANGUISH, Angor.

ANGUISH, FEBRILE, Angor Febrilis. The combination of weariness, pain, anxiety, and weakness affecting the head and neck, which is so generally observed at the commencement of fever.

ANGULAIRE DE L'OMOPATE, Levator scapulae.

ANGULAR, Angula'ris, from angulus, 'an angle,' (F.) Angulaire. That which relates to an angle.

ANGULAR ARTERY AND VEIN. A name given, 1. to the termination of the facial artery and vein, because they pass by the greater angle of the eye; and, 2. to the facial artery and vein themselves, because they pass under the angle of the jaw. See Facial.

ANGULAR NERVE is a filament furnished by the inferior maxillary, which passes near the greater angle of the eye.

ANGULAR PROCESSES of the frontal bone are seated near the angles of the eyes. See Orbital.

ANGULARIS, Levator scapulae.

ANGULI-SCAPULO-HUMÉRAL, Teres major.

ANGULUS COSTÆ, see Costa—*a.* Ocularis, Canthus.

ANGURIA, Cucurbita citrullus.

ANGUSTATIO, Arctatio—*a.* Cordis, Systole—*n.* Intestini recti vel Ani, Stricture of the rectum.

ANGUSTIA, Angusta'tio, Stenochoria. Anxiety, narrowness, strait, constriction.

ANGUSTIA ABDOMINALIS, Pelvis, (Brim)—*a.* Perinaealis, Pelvis, (Outlet.)

ANGUSTURA, Cusparia febrifuga—*a.* False, Brucea antidysenterica, and Strychnos nux vomica—*a.* Spuria, Brucea antidysenterica, and Strychnos.

ANGUSTURE, FAUSSE, Brucea antidysenterica—*a.* Ferrugineuse, Brucea antidysenterica—*a.* Vraie, Cusparia febrifuga.

ANGUSTURINE, Brucine.

ANHÆMATOSIA, Asphyxia, Anæmia.

ANHÆMIA, Anæmia.

ANHAPHIA, Anaphia.

ANHELATIO, from anhelō, 'I pant.' Anhelitus, Aæmus, Panting, Gasping, Anhelation, (Sc.) Hech, Hegh, (Prov.) Gien, Kaving, Peching, (F.) Essoufflement. Short and rapid breathing. See Dyspnœa.

Anhelatio is sometimes employed synonymously with asthma.

ANHELITUS, Breath.

ANHÆMATOSIE, Anæmia, Asphyxia.

ANHÉMIE, Anæmia.

ANHIS'TOUS, from a, av, privative, and 'istos, 'organic texture,' 'Anorganic.' Amor'phus. The tunica decidua uteri is termed by Velpeau the *anhistous membrane*.

ANHUIBA, Laurus sassafras.

ANHYDRÆMIA, Anæmyd'ria, from av, privative, 'vōwp, 'water,' and 'asma, 'blood.' A condition of the blood in which there is a diminution in the quantity of the serum.

ANHYDROMYEL'IA, from av, priv., 'vōwp,

'water,' and μυελος, 'marrow.' Deficiency or absence, in the spinal cavity, of the cephalo-spinal fluid.

ANICE'TON, Anice'tum, Mesia'mum, from a, privative, and νικη, 'victory,' 'invincible.' A plaster much extolled by the ancients in cases of achores. It was formed of litharge, cerusse, thus, alum, turpentine, white pepper, and oil.

ANID'EUS, from av, privative, and εἶδος, 'shape.' Amor'phus. A monster devoid of shape.—J. G. St. Hilaire.

ANIDRO'SIS, from a, privative, and 'ιδρως, 'sweat.' Sudor'is null'itas vel priva'tio. Absence of sweat. Deficiency of perspiration.—Hippocrates.

ANILEMA, Borborygmus, Tormina.

ANILESIS, Borborygmus, Tormina.

ANILITAS, see Dementia.

AN'IMA, An'imus, Mens, Psychē. The mind, breath, &c., from αἶμος, 'wind or breath.' (F.) **Âme.** The principle of the intellectual and moral manifestations. Also, the principle of life—the life of plants being termed *An'ima vegetati'va*, (F.) *Âme végétative*; that of man, *An'ima sensi'ti'va*, (F.) *Âme sensitive*.

The *Anima* of Stahl, *An'ima Stahlia'na*, was a fancied intelligent principle, which he supposed to preside over the phenomena of life,—like the *Archæus* of Van Helmont.

Under the term *Anima mundi*, the ancient philosophers meant a universal Spirit, which they supposed spread over every part of the universe.

The precise seat of the mind in the brain has given rise to many speculations. The point is unsettled.

With the ancient chemists, *Anima* meant the active principle of a drug separated by some chemical management.

ANIMA ALOES: see Aloes Succotorina—*a.* Articulorum, Hermodactylus—*a.* Hepatis, Ferri sulphas—*a.* Pulmonum, Crocus—*a.* Rhei, Infusum rhei—*a.* Stahlianæ, see Anima—a. Vegetativa, Plastic force.

AN'IMAL, Zoön. A name given to every animated being. Most animals have the power of locomotion; some can merely execute partial movements, such as contraction and dilatation. In other respects it is often a matter of difficulty to determine what is an animal characteristic. The study of animals is called *Zoöl'ogy*.

AN'IMAL, Anima'lis. That which concerns, or belongs to, an animal.

ANIMAL HEAT, Calor anima'lis seu nati'vus, Cal'idum anima'le seu innat'um, Biolychnion, Flam'mula vita'lis, Thërma em'phytum, Thërma em'phytum, Ignis anima'lis seu natura'lis seu vita'lis, (F.) Chaleur animale, is the caloric constantly formed by the body of a living animal, by virtue of which it preserves nearly the same temperature, whatever may be that of the medium in which it is placed. This formation seems to take place over the whole of the body, and to be connected with the action of nutrition.

The following are the natural temperatures of certain animals; that of man being 98° or 100°

ANIMALS.	Temperature.
Arctic Fox	101
Arctic Wolf.....	105
Squirrel.....	
Hare	104
Whale	103
Arctomys citillus, zibul—in summer.....	80 to 84
Do. when torpid.....	103
Goat	102
Bat. in summer.....	101 or 102
Musk	101
Marmota bobac—Bobac.....	
House mouse.....	

ANIMALS.	Temperatur.
Arctomys marmota, <i>marmot</i> ,—in summer.....	101 or 102
Do. when torpid.....	43
Rabbit.....	100 to 104
Polar Bear.....	100
Dog.....	100 to 103
Cat.....	
Swine.....	
Sheep.....	
Ox.....	
Guinea-pig.....	100 to 102
Arctomys glis.....	99
Shrew.....	98
Young wolf.....	96
Fringilla arctica, <i>Arctic finch</i>	111
Rubecola, <i>redbreast</i>	
Fringilla linaria, <i>lesser red poll</i>	110 or 111
Falco palumbarius, <i>goshawk</i>	100
Caprimulgus Europeanus, <i>European goat-sucker</i>	
Emberiza nivalis, <i>snow-bunting</i>	109 to 110
Falco lanarius, <i>lanner</i>	
Fringilla carduelis, <i>goldfinch</i>	109
Corvus corax, <i>raven</i>	
Turdus, <i>thrush</i> , (of Ceylon,).....	
Tetrao perdix, <i>partridge</i>	108
Anas clypeata, <i>shoveler</i>	
Fringilla pugnax, <i>ruff</i>	
Scelopax limosa, <i>lesser godwit</i>	
Tetrao tetrix, <i>grouse</i>	
Fringilla brumalis, <i>winterfinch</i>	107
Loxia pyrrhula.....	
Falco nisus, <i>sparrowhawk</i>	107 to 111
Vultur barbatus.....	
Anser pilechroclis.....	106
Columbus arritus, <i>dusky grebe</i>	
Tringa vanellus, <i>lapwing</i> , wounded.....	105
Tetrao lagopus, <i>ptarmigan</i>	
Fringilla domestica, <i>house sparrow</i>	104
Strix passerina, <i>little owl</i>	
Illeotopus ostraceus, <i>sea-pie</i>	103 to 107
Anas penelope, <i>widgeon</i>	
Anas strepera, <i>gadwall</i>	103
Pelecanus carbo.....	
Falco ossifragus, <i>sea-eagle</i>	89 to 91
Fulica atra, <i>coot</i>	
Anas acuta, <i>pin-tail duck</i>	83
Falco nilvus, <i>kite</i> , (wounded,).....	
Merops apiaster, <i>bee-eater</i>	74
Goose.....	
Hen.....	103
Dove.....	
Duck.....	89 to 91
Ardea stellaris.....	
Falco albicollis.....	83
Picus major.....	
Cossus ligniperda.....	74
Shark.....	
Torpedo marmorata.....	

ANIMAL KINGDOM, (F.) *Règne Animal*, comprises all animated beings.

ANIMAL LAYER, see *Tache embryonnaire*—a. Magnetism, see Magnetism, animal.

ANIMALCULA SEMINALIA, Spermatozoa—a. Spermatica, Spermatozoa.

ANIMAL/CULE, *Animalculum*; diminutive of *animal*. A small animal. An animal well seen only by means of the microscope.

ANIMALCULES, SEMINAL, Spermatozoa—a. Spermatic, Spermatozoa.

ANIMALCULISM, Spermatism.

ANIMAL/CULIST, *Animalist*. One who attempts to explain different physiological or pathological phenomena by means of animalcules. One who believes that the embryo is formed from the so-called spermatic animalcules—*sperm'atist*, *sem'inist*.

ANIMAL/CULOVISM, *Animalculovism*—us, from *animalculum* and *ovum*. The doctrine, now universal, which maintains that the new being is formed by the concurrence of the spermatic animalcule or spermatozoid with the ovum.

ANIMALCULUM, Animalcule.

ANIMALIST, Animalculist.

ANIMALITY, *Animalitas*, (F.) *Animalité*. The aggregate of qualities which distinguish that which is animated. That which constitutes the animal.

ANIMALIZATION, *Animalisatio*. The transformation of the nutritive parts of food into the living substance of the body to be nourished. Assimilation.

To ANIMATE, *Animarē*. To unite the living principle with an organized body. The French use it in the sense of,—to excite or render active; as, *animer un vésicatoire*: to excite a blister to suppurate.

ANIMATION, Animation—a. *Fœtus*, see Quickening.

ANIMATION, *Zoôsis*, *Animalisatio*, from *anima*, 'the soul or mind.' The act of animating. The state of being enlivened.

ANIMATION, SUSPENDED, Asphyxia.

AN'IME, *Gum an'imē*, *Amin'a'a*, *Gummi an'imē*, *Can'camum*, *Can'camp*. A resin obtained from the trunk of *Hymen'œa coum'baril*. Ord. Fabaceæ. It has been given as a cephalic and uterine. It is not used. The plant is also called *Cour'baril*.

AN'IME, (F.) An epithet applied to the countenance, when florid, in health or disease.

ANIMELLÆ, Parotid.

ANIMI CASUS SUBITUS, Syncope—a. De-lirium, Syncope—a. Pathemata, Passions.

ANIMISM, see *Animist*.

AN'IMIST, from *anima*, 'the soul.' One who, following the example of Stahl, refers all the phenomena of the animal economy to the soul, *An'imism*.

The soul, according to Stahl, is the immediate and intelligent agent of every movement, and of every material change in the body. Stahl therefore concluded, that disease is nothing more than a disturbance or disorder in the government of the economy, or an effort by which the soul, attentive to every morbid cause, endeavours to expel whatever may be deranging the habitual order of health. See *Stahlianism*.

ANIMUS, Anima, Breath.

ANIRID'IA, from *an*, priv., and *iris*, *ἰρις* 'the iris.' Absence of the iris.

A'IN, Pimpinella anisum—a. *Aigre*, Cuminum Cyminum—a. *de la Chine*, Illicium anisatum—a. *Doux*, Anethum—a. *Étoilé*, Illicium anisatum.

ANISA'TUM, from *Anisum*, 'Anise.' A sort of medicated wine, formerly prepared with honey, wine of Ascalon, and aniseed.

ANISCALPTOR, Latissimus dorsi.

ANISCHURIA, Enuresis.

ANISE, Pimpinella anisum—a. Star, Illicium anisatum, I. Floridanum—a. Tree, Florida, Illicium Floridanum—a. Tree, yellow-flowered, Illicium anisatum.

ANISEED, see Pimpinella anisum.

ANISEMA DRACONTIUM, *Arum draconium*.

ANISI SEMINA, see Pimpinella anisum.

ANISO'DUS L'URIDUS, *Nican'dra anom'ala*, *Phy'alis stramo'nium*, *Whitley'a stramo'nifolia*. A plant of Nepal, possessed of narcotic properties, and resembling belladonna and tobacco. It dilates the pupil, and is used in diseases of the eye like belladonna. It is given in alcoholic tincture (*dried leaves* 3j to alcohol f3viij). Dose, 20 drops internally in the 24 hours.

ANISOPHYLLUM IPECACUANHA, Euphorbia Ipecacuanha.

ANISOS'THENES, *Inaqua'li rob'orē pollens*. That which is unequal in strength: from *ai*, priv., *ἴσος*, 'equal,' and *σθένος*, 'strength.' An epithet applied particularly to the muscular contractility which, in the sick, is sometimes augmented in certain muscles only,—in the flexors, for example.

ANISOT'ACHYS, from *an*, priv., *ἴσος*, 'equal,

and ταχως, 'quick.' An epithet for the pulse, when quick and unequal.—Gorraus.

ANISUM, Pimpinella anisum—a. Africanum frutescens, Bubon Galbanum—a. Fruticosum galbanum, Bubon Galbanum—a. Officinale, Pimpinella anisum—a. Sinense, Illicium anisatum—a. Stellatum, Illicium anisatum—a. Vulgare, Pimpinella anisum.

ANKLE, Astragalus, Malleolus.

ANKLEY, Malleolus.

ANKUS, Ancus.

ANKYLOBLEPH'ARON, *Ancylobleph'aron*, *Palpebra'rum coal'itus*, from *αγκυλη*, 'a bridle,' and *βλεφαρον*, 'eyelid.' A preternatural union between the free edges of the eyelids. Likewise called *Synbleph'aron*, *Symbblepharo'is*, and *Pro'physis*.

Also, union between the eyelids and globe of the eye.—Aëtius.

ANKYLOCHET'LIA, *Ancylochei'tia*, (F.) *Ancylochei'tie*, from *αγκυλος*, 'crooked,' and *χειλος*, 'lip.' Accidental union of the lips.

ANKYLOCHÉLIE, *Ancylocheilia*.

ANKYLOCOLPUS, Colpatresia.

ANKYLOCORE, Coreelisis.

ANKYLODON'TIA, from *αγκυλος*, 'crooked,' and *οδους*, *οδοντος*, 'a tooth.' An irregular position of the teeth in the jaws.

ANKYLOGLOS'SIA, *Ancyloglos'sia*, *Comere'tio lingue*, from *αγκυλη*, 'a bridle,' and *γλωσσα*, 'the tongue.' Impeded motion of the tongue in consequence of adhesion between its margins and the gums; or in consequence of the shortness of the frænum: the later affection constituting *Tongue-tie*, *Olopho'nia lin'guæ frænata*. It merely requires the frænum to be divided with a pair of scissors.

ANKYLOGLOSSOT'OMUM, from *an'kyloglos'sia*, 'tongue-tie,' and *τομη*, 'incision.' An instrument used in the operation for tongue-tie.

ANKYLOME'LÊ, *Ancylome'lê*, from *αγκυλος*, 'crooked,' and *μηλη*, 'a probe.' A curved probe.—Galen.

ANKYLOMERIS'MUS, *Ancylomeris'mus*, from *αγκυλη*, 'a bridle,' and *μερος*, 'a part.' Morbid adhesion between parts.

ANKYLOPS, Ægilops.

ANKYLORRHIN'IA, *Ancylo'rrhin'ia*, from *αγκυλη*, 'a bridle,' and *ρην*, 'the nose.' Morbid union of the parietes of the nose.

ANKYLOSED, see Ankylosis.

ANKYLO'SIS, *Ancylo'sis*, *Anchylo'sis*, *An'cylê*, *Stiff Joint*, from *αγκυλος*, 'crooked.' An affection, in which there is great difficulty or even impossibility of moving a diarthrodial articulation. It is so called, because the limb commonly remains in a constant state of flexion, and a joint thus affected is said to be *ankylosed*. Anchylosis is said to be *complete* or *true*, when there is an intimate adhesion between the synovial surfaces, with union of the articular extremities of the bones. In the *incomplete* or *false* ankylosis, there is obscure motion, but the fibrous parts around the joint are more or less stiff and thickened. In the treatment of this last state, the joint must be gently and gradually exercised; and oily, relaxing applications be assiduously employed.

ANKYLOSIS SPURIA, *Rigiditas articularum*.

ANKYLO'TIA, *Ancylo'tia*, from *αγκυλη*, 'a bridle,' and *ους*, *ωρος*, 'the ear.' Morbid union of the parietes of the ear.

ANKYLOT'OMUS, *Ancylo'tomus*, from *αγκυλος*, 'crooked,' and *τεμνειν*, 'to cut.' Any kind of curved knife.—Paulus. An instrument for dividing the frænum linguae.—Sculetus.

ANNEAU, Ring—a. *Crural*, *Crural canal*—a. *Diaphragmatique*, *Diaphragmatic ring*—a. *Fémo-*

ral, *Crural canal*—a. *Inguinal*, *Inguinal ring*—a. *Ombilical*, *Umbilical ring*.

ANNEXE, Accessory, Appendix.

ANNI CRITICI, Climacterici (anni)—a. Decretorii, Climacterici (anni)—a. Fatales, Climacterici (anni)—a. Genethliaci, Climacterici (anni)—a. Gradarii, Climacterici (anni)—a. Hebdomadici, Climacterici (anni)—a. Heroici, Climacterici (anni)—a. Natalitii, Climacterici (anni)—a. Scallares, Climacterici (anni)—a. Scansiles, Climacterici (anni).

ANNOTATIO, *Episma'sia*. Under this term some have included the preludes to an attack of intermittent fever—as yawning, stretching, somnolency, chilliness, &c.

ANNOTTO, see Terra Orleana.

ANNUAL DISEASES, *Morbi an'nui*, *M. an'niuersarii*, (F.) *Maladies annuelles*. A name given, by some, to diseases which recur every year about the same period. *Febris annua*, (F.) *Fièvre annuelle*, is a term used for a fancied intermittent of this type.

ANNUENS, *Rectus capitis internus minor*.

ANNUIT'IO, *Nodding*, from *ad*, 'to,' and *nutus*, 'a nod.' A gesture, denoting assent in most countries. Also, the state of somnolency, when the individual is in the erect or sitting posture, with the head unsupported, in which the power of volition over the extensor muscles of the head is lost, and the head drops forward.

ANNULAR, *Annula'ris*, *Anula'ris*, *Cricoi'des*, (F.) *Annulaire*, (*annus*, 'a circle.') Any thing relating to a ring, or which has the shape or fulfils the functions of a ring; from *annulus*, 'a ring.'

ANNULAR FINGER, *Ring Finger*, *Ring man*, *Dig'itus annula'ris*, *Param'esos*. The fourth finger, so called from the wedding ring being worn thereon. See Digitus. Of old, it was believed, that there is a medium of direct communication between the ring finger and the heart.

ANNULAR GANGLION, see Ciliary ligament.

ANNULAR LIG'AMENT, *Transverse ligament*, *Cru'cial ligament*. A strong ligamentous band, which arches across the area of the ring of the atlas, from a rough tubercle upon the inner surface of one articular process, to a similar tubercle on the other. It serves to retain the odontoid process of the axis in connexion with the anterior arch of the atlas.

AN'NULAR LIG'AMENT OF THE RA'DIUS, is a very strong fibro-cartilaginous band, which forms, with the lesser sigmoid cavity of the cubitus, a kind of ring, in which the head of the radius turns with facility.

AN'NULAR LIG'AMENTS OF THE CARPUS, *Ar'mill'æ manus membrano'sæ*, are two in number.

The one, *anterior*, is a broad, fibrous, quadrilateral band, extending transversely before the carpus, and forming the gutter, made by the wrist, into a canal. It is attached, externally, to the trapezium and scaphoides; and internally to the os pisiforme and process of the unciform. It keeps the tendons of the flexor muscles, median nerve, &c., applied against the carpus.

The *posterior* ligament is situate transversely behind the joint of the hand, and covers the sheaths of the tendons, which pass to the back of the hand. Its fibres are white and shining, and are attached, externally, to the inferior and outer part of the radius; internally to the ulna and os pisiforme.

AN'NULAR LIG'AMENTS OF THE TARSUS are two in number. The *anterior* is quadrilateral, and extends transversely above the instep. It is attached to the superior depression of the os calcis, and to the malleolus internus. It embraces the tendons of the extensor muscles of the toes, the *tibialis anticus*, and *peroneus anticus*. The inter-

al is broader than the last. It descends from the malleolus internus to the posterior and inner part of the os calcis, with which it forms a kind of canal, enclosing the sheaths of the tendons of the *tibialis posterior*, *flecor longus digitorum pedis*, and *F. longus pollicis pedis*, as well as the plantar vessels and nerves.

ANNULAR VEIN, *Vena annularis*, is situated between the annular finger and the little finger. Aëtius recommends it to be opened in diseases of the spleen.

ANNULARIS, Cricoid : see *Digitus*—*a. Ani*, *Sphincter ani*.

ANNULI CARTILAGINEI, see *Trachea*—*a. Cartilaginosa Tracheæ*, see *Trachea*.

ANNULI FIBRO-CARTILAGINEI, "*Fibro-cartilaginous or festooned rings*." Tough tendinous tracts in the heart, two of which, anterior, are situated on the right and left of the aortal opening; and one posterior, which runs backwards from the aorta to the border of the auriculo-ventricular septum, where it splits into two slender crura.

ANNULI-TENDINO-PHALANGIENS, *Lumbricales manus*.

ANNULUS, *Dactylus*, *Vulva*—*a. Abdominis*, *Inguinal ring*—*a. Albidus*, see *Ciliary (ligament)*—*a. Cellulosus*, *Ciliary ligament*—*a. Ciliaris*, *Ciliary ligament*—*a. Fossæ ovalis*: see *Ovalis fossa*—*a. Gangliiformis*, see *Ciliary (ligament)*—*a. Repens*, *Herpes circinatus*—*a. Umbilicalis*, *Umbilical ring*—*a. Ventriculi*, *Pylorus*—*a. Viuesenii*, see *Ovalis fossa*.

ANO, *anw*. A prefix denoting 'above, up.'

ANO-CAVERNEUX, *Accelerator urinæ*.

ANOCHILUS, from *anw*, 'above,' and *χειλος*, 'lip.' The upper lip. Also, one who has a large upper lip.

ANOCCELIA, *Stomach*.

ANOCCELADELPHUS, *Cœliadelphus*.

ANO'DIA, from *av*, priv., and *ὠδή*, 'song.' An unconnected or dissonant mode of speech.

ANOD'IC, *Anod'icus*, from *anw*, 'above, up,' and *ὁδός*, 'a way.' Tending upwards. An epithet applied by Dr. Marshall Hall to an ascending course of nervous action.

ANODIN, *Anodyne*.

ANODIN'IA, from *a*, priv., and *ὠδιν*, 'a labour pain.' Absence of labour pains.

ANODMIA, *Anosmia*.

ANODUS, *Edentulus*.

AN'ODYNE, *Anod'ynus*, *Antod'ynus*, *Antid'ynus* (improperly), *Paregor'icus*, *Anet'icus*, *Antal'gicus*, *Acesod'ynes*, (F.) *Anodin* ou *Anodyn*, from *av*, privative, and *ὀδυν*, 'pain.' *Anodynes* are those medicines which relieve pain, or cause it to cease; as opium, belladonna, &c. They act by blunting the sensibility of the encephalon, so that it does not appreciate the morbid sensation.

ANODYN'IA, *Indolen'ia*. Cessation or absence of pain. Vogel has given this name to a genus of diseases, characterized by a cessation of pain, and the exasperation of other symptoms; as we see in gangrene.

ANODYNUM MINERALE, *Potassæ nitratis sulphatis paucillo mixtus*.

ANOE'IA, *Anoi'a*, from *a*, privative, and *νοος*, 'mind.' Delirium, imbecility. See *Dementia* and *Idiotism*.

ANOSIA, *Dementia*—*a. Abstricta*, *Melancholy*.

ANOIA, *Anœa*.

ANOMAL, *Anomalous*.

ANOMALES, *Anomalous*.

ANOMAL'IA, from *av*, privative, and *ομαλος*, 'regular.' *Abnorm'itas*, *Aliena'tio*. Anomaly, abnormality, irregularity. In Pathology, anomaly means something unusual in the symptoms pro-

per to a disease, or in the morbid appearances presented by it.

ANOMALIA NERVORUM, *Nervous diathesis*.

ANOMALIE PAR CLOISONNEMENT, see *Cloisonnement*.

ANOMALOTROPHIES, from *av*, privative, *ομαλος*, 'regular,' and *τροφή*, 'nourishment.' A class of diseases, which consist in modifications in the nutrition of organs.—Gendrin.

ANOM'ALOUS, *Anom'alus*, *Anom'alis*; the same etymon. Irregular; contrary to rule. (F.) *Anomal*. In Medicine, a disease is called *anomalous*, in whose symptoms or progress there is something unusual. Affections are also called *anomalous*, which cannot be referred to any known species.

ANOMALOUS, *Irregular*.

ANOMMATUS, *Anomphthalmus*.

ANOMOCEPH'ALUS, from *a*, priv., *νομος*, 'rule,' and *κεφαλή*, 'head.' One whose head is deformed.—Geoffroi Saint-Hilaire.

ANOMOSPLÉNOTOPIE, *Splenectopia*.

ANOMOTOPIE, *Ectopia*.

ANOM'PHALUS, from *av*, priv., and *ομφαλος*, 'the navel.' One devoid of navel. Many writers have endeavoured to show that Adam and Eve must have been *ανομφαλοι*, as they could not have had umbilical vessels.

ANONA TRILOBA, see *Carica papaya*.

ANO'NA TRIPET'ALA. A tree of the family *Anonæ* or *Anonacæ*; *Sec. Syst.* *Polyandria Polygynia*, from fifteen to twenty feet high, native of South America, which bears a delicious fruit called *Chirimoya*. Both the fruit and flowers emit a fine fragrance, which, when the tree is covered with blossom, is almost overpowering.—Tschudi.

ANONIS, *Ononis*.

ANONYCH'IA, from *av*, privative, and *ονυχ*, *ονυχος*, 'a nail.' Want of nails,—a rare congenital defect.

ANONYME, *Innominatum*.

ANONYMOUS, *Anon'ymus*, *Innomina'tus*, (F.) *Anonyme*, from *av*, privative, and *ονομα*, 'name.' That which has no name.

The word has been applied to many parts of the body:—to the *Anonymous bone* or *Os innominatum*:—the *Anonymous foramen* or *Foramen innominatum*, &c.

ANOPHRESIA, *Anosmia*.

ANOPHTHAL'MUS, *Anom'thalmus*, from *av*, privative, and *οφθαλμος*, 'an eye.' One who is devoid of eyes.

ANOPS'IA, from *av*, priv., and *ωψ*, 'the eye.' A case of monstrosity in which the eye and orbit are wanting.

ANOPTICONERVIE, *Amaurosis*.

ANOR'CHIDES, from *av*, priv., and *ορχις*, 'a testicle.' They who are without testicles.—*Fortunatus Fidelis*.

ANOREX'IA, from *av*, priv., and *ορεξις*, 'appetite.' *Inappetent'ia*, *Limo'sis expert*, (F.) *Peris d'appetit*. Absence of appetite, without loathing. Anorexia or want of appetite is symptomatic of most diseases. Also, *Indigestion*, *Dyspepsia*.

ANOREXIA EXHAUSTORUM, *Frigidity of the stomach*—*a. Mirabilis*, *Fasting*.

ANORGANIC, see *Anihistous* and *Inorganic*.

ANORMAL, *Abnormal*.

ANORTHOP'IA, from *a*, priv., *ορθος*, 'straight,' and *ὄψις*, 'vision.' Vision in which persons are unable to judge when objects are not parallel or not symmetrical.

ANOS'IA, from *a*, priv., and *νοσος*, 'disease.' Health. Freedom from disease.

ANOS'MIA, from *av*, privative, and *οσμη*, 'odour.' Loss of smell. Diminution of the sense of smell. Called, also, *Anosphre'sia*, *Anosphra'sia*, *Anosphre'sia*, *Paros'mia*, *Anod'mia*, *Anosmo'sia*.

cia, Olfactûs amis'sio seu defic'iens, Dysœsthe'sia olfacto'ria, Anœsthe'sia olfacto'ria, Odora'tus deper'ditus, (F.) Perte de l'Odorat.

ANOSMOSIA, Anosmia.

ANOSPHRASIA, Anosmia.

ANOSPHERISIA, Anosmia.

ANO'TUS, from *av*, privative, and *ous*, *oros*, 'the ear.' A monster having no ears.

ANOXÉMIE, Asphyxia.

ANSA INTESTINALIS, *Anse (intestinalle).*

ANSE (F.), *Ansa* (L.), signifies, properly, the handle of certain vessels, usually of an arched form. By analogy, it has been applied to that which is curved in the form of such handle. Thus, the French speak of *Anse intestinale*—*Ansa* seu *Gyrus intestinalis*—to signify a portion of intestine, supported by its mesentery, and describing a curved line:—also, of *Anse nerveuse, Anse anastomotique, &c.*

Anse de fil is used, in Surgery, to designate a thread, curved in the form of an *Anse*.

ANSERINA, *Potentilla anserina*.

ANSÉRINE, *Chenopodium ambrosioides*—*a. Anthelmintique, Chenopodium anthelminticum*—*a. Bon Henri, Chenopodium Bonus Henrius*—*a. Botrys, Chenopodium Botrys*—*a. Féide, Chenopodium vulvaria*—*a. Vermifuge, Chenopodium anthelminticum*.

ANTAC'IDS, *Anti-acids, Antiac'ida, Invertent'ia*, from *anti*, 'against,' and *acida*, 'acids.' Remedies which obviate acidity in the stomach. They are chemical agents, and act by neutralizing the acid. Those chiefly used are ammonia, calcei carbonas, calx, magnesia, magnesie carbonas, potassa, potasse bicarbonas, p. carbonas, sodæ bicarbonas, and s. carbonas. They are, of course, only palliatives, removing that which exists, not preventing the formation of more.

ANTAG'ONISM, *Antagonis'mus, Antis'tasis*, from *avri*, 'against,' and *aywivōqat*, 'to strive.' Action in an opposite direction. It applies to the action of muscles that act in a contrary direction to others. In estimating the force of muscles, this antagonism must be attended to.

ANTAG'ONIST, *Antagonis'ta*. A muscle whose action produces an effect contrary to that of another muscle. Every muscle has its antagonist, because there is no motion in one direction without a capability of it in another.

ANTALGICUS, *Anodyne*.

ANTAPHRODIS'IA, *Antaphrodit'ic, Antaphrodisiacus, Anaphrodisiacus, Anaphrodis'iac, Anterot'icus*, from *avri*, 'against,' and *αφροδισιακος*, 'aphrodisiac.' A substance capable of blunting the venereal appetite.

ANTAPHRODITIC, *Antaphrodisiac*.

ANTAPOD'OSIS, from *avrapodidwmi*, 'I return in exchange.' The succession and return of the febrile periods.—Hippocrates.

ANTAPOLECTICUS, *Antiapoplectic*.

ANTARTHITIC, *Antiarthritic*.

ANTASTHENICUS, *Tonic*.

ANTASTHMATICUS, *Antiasthmatic*.

ANTATROPH'IC, *Antatroph'icus, Antatrophus, Antatroph'icus*, from *avri*, 'against,' and *ατροφια*, 'atrophy.' A remedy opposed to atrophy or consumption.

ANTEBRACHIAL, see *Antibrachial*.

ANTECEDEN'TIA. The precursory or warning symptoms of a disease.

ANTEFLEXIO UTERI, see *Anteversio*.

ANTELA'BIA, *Proche'la*, from *ante*, 'before,' and *labia*, 'the lips.' The extremity of the lips.

ANTELOPE, *Antilopus*.

ANTEBALLOLUMEN, *Succedaneum*.

ANTEN'BASIS, from *avri*, and *εμβασις*, 'I enter.' *Mutuus ingres'sus*. The mutual reception of bones.—Galen.

ANTEMETIC, *Antiemetic*.

ANTEDEXIS, *Counter-indication*.

ANTENDIXIS, *Counter-indication*.

ANTENEAS'MUS, from *avri*, 'against,' and *veav*, 'audacious;' or rather, perhaps, from *avri*, and *τεivo*, 'I extend or stretch.' One furious against himself. Mania, in which the patient attempts his own life.—Zacchias.

ANTENNA'RIA DIOFCA, *Gnaphalium Dioicum, Hispid'ula, Pes cati, Eli'chrysium montanum, Dico'cious Everlast'ing, Catfoot, (F.) Pied de chat. Order, Compositæ*. A common European plant, which has been advised in hemorrhage, diarrhoea, &c.

ANTENNARIA MARGARITACEA, *Gnaphalium margaritaceum*.

ANTENNARIA PLANTAGINIFOLIA, *Gnaphalium plantaginifolium* seu *plantagin'eum* seu *dioicum* variety *plantaginifolium*, *Plantain Life-everlasting, Cudweed*, indigenous, has similar properties.

ANTEPHIALTIC, *Antiephialtic*.

ANTEPILEPTIC, *Antiepileptic*.

ANTEPONENS, *Anticipating*.

ANTERE'ISIS, from *avri*, 'against,' and *επειδω*, 'I support.' The resistance—the solidity—of bones.—Hippocrates.

ANTÉRIEUR DU MARTEAU, *Laxator tympani*—*a. de l'Oreille, Anterior auris*.

ANTE'RIOR, *Ant'icus*, from *ante*, 'before.' Situate before. Great confusion has prevailed with anatomists in the use of the terms *before*, *behind*, etc. Generally, the word *anterior* is applied to parts situate before the median line, the body being in the erect posture, with the face and palms of the hands turned forwards, and the feet applied longitudinally together.

ANTE'RIOR AU'RIS (Muscle), *Auricula'ris antérieur, At'trahens auric'ulam, (F.) Auriculaire antérieur, Antérieur de l'oreille, Zygomato-oriculaire*. A small muscle, passing from the posterior part of the zygoma to the helix. Use, to draw the ear forwards and upwards.

ANTERIOR MALLEI, *Laxator tympani*.

ANTEROTICUS, *Antaphrodisiac*.

ANTEUPHORBIIUM, *Cacalia antieuphorbium*.

ANTEVER'SION, *Antever'sio, Antrover'sio*, from *ante*, 'before,' and *vertere, versum*, 'to turn.' Displacement of the uterus, in which the fundus is turned towards the pubes, whilst its orifice is towards the sacrum. It may be caused by extraordinary size of the pelvis, pressure of the viscera on the uterus, etc.; and is recognised by examination *per vaginam*. A simple forward inclination of the body of the uterus, without the os uteri being carried much backwards, is termed *Anteflex'ion of the uterus, Anteflex'io u'teri*. Not unfrequently, however, it is used synonymously with *Retroversion of the Uterus*. See *Retroversio uteri*.

ANTHÆMPTY'ICUS, *Antihæmpty'icus*, from *avri*, 'against,' and *hæmptyia*, 'spitting of blood.' Against spitting of blood. A remedy for spitting of blood—*antihæmptygium (remedium)*.

ANTHÆMORRHAGICUS, *Antihæmorrhagic*.

ANTHETICUS, *Antihetic*.

ANTHELI'T'RAGUS, (F.) *Anthélitragien*. One of the proper muscles of the pavilion of the ear.

ANT'HELIX, *Anti-helix*, from *avri*, 'before,' and *ελix*, 'the helix.' An eminence on the cartilage of the ear, in front of the helix, and extending from the concha to the groove of the helix, where it bifurcates.

ANTHELMINT'IC, *Antihelmin'ticus, Antiecol'icus, Anthelmin'thicus, Antiscol'eticus, Helmin'thicus, Helminthago'gus, Antivermino'sus, Vermifugus, Vermifuge*, from *avri*, 'against,' and *ελμινς*, 'a worm.' A remedy which de-

stroy or expels worms, or prevents their formation and development. The chief anthelmintics are, *Chenopodium*, *Mucuna*, *Oleum animale Dippelii*, *Oleum Terebinthinæ*, *Sodii Chloridum*, *Spigelia*, and *Pulvis Stanni*. See Worms.

ANTHEMA ERUPTIO, Eranthem.

ANTHEMIS COTULA, from *antheō*, 'I flower.' *A. foetida* seu *Nasturaceæ* *sis*, *Cotula*, *C. foetida*, *Cotu*, *Cyanthemis*, *Chamæelum foetidum* seu *Chamomilla spuria* seu *foetida*, *Marrubia Cotula*, *Mayflower*, *Mayweed*, *Stinking Chamomile*, *Wild Chamomile*, *Dog's fennel*, *Dilly*, *Dibweed*, *Field-weed*, *Pissweed*. Ord. *Compositæ Corymbiferae*. *Sex. Syst.* Syngenesia Superflua. (F.) *Maroute*, *Camomille fétide*, *Camomille puante*.

This plant *Cotula*, (Ph. U. S.), has a very disagreeable smell, and the leaves have a strong, acrid, bitterish taste. It is reputed to have been useful in hysterical affections.

ANTHEMIS FETIDA, A. cotula.

ANTHEMIS NOBILIS, *Anthemis*, *A. odorata*, *Chamæelum*, *Ch. Nobilis* seu *odoratum*, *Chamomilla Romana*, *Eucanthemum*, *Leucanthemum*, (F.) *Camomille Romaine*. The leaves and flowers. — *Anthemis*, Ph. U. S., have a strong smell, and bitter, nauseous taste. The flowers are chiefly used. They possess tonic and stomachic properties, and are much given as a pleasant and cheap bitter. A simple infusion is taken to produce, or to assist vomiting. Externally, they are often used in fomentations.

The *Oleum Anthemidis* possesses the aromatic properties of the plant, but not the bitter and tonic. Consequently, the '*Chamomile Drops*,' as sold by the druggists, must be devoid of the latter qualities. They are made by adding *Ol. anthem.* f3j. to *Sp. vini rectif.* Oj.

ANTHEMIS NOVEBORACENSIS, A. cotula—a. Odorata, A. cotula.

ANTHEMIS PYRETHRUM, *Pyrethrum*, *P. verum*, *Anacælus pyrethrum*, *Buphthalmum Creticum*, *Dentaria*, *Herba salicaria*, *Pes Alexandrinus*, *Stinking Chamomile*, *Pellitory of Spain*, (F.) *Pyrethre*, *Racine salivaire*, *Salivaire*, *Pied d'Alexandre*. The root, *Pyrethrum* (Ph. U. S.), is hot and acrid, its acrimony residing in a resinous principle. It is never used except as a masticatory in toothache, rheumatism of the face, paralysis of the tongue, etc. It acts as a powerful sialogogue.

The *Pellitory of the shops* in Germany is said to be derived from *Anacælus officinarum*; a plant cultivated in Thuringia for medicinal purposes.

ANTHEMIS TINCTORIA, *Buphthalmi Herba*, *Dyer's Chamomile*, a European plant, has a bitter and astringent taste, and has been regarded as stomachic and vulnerary. (F.) *Camomille des Teinturiers*, *Œil de Bœuf*.

ANTHEMIS VULGARIS, *Matricaria Chamomilla*.

ANTHERA, from *antheros*, so called from its 'florid' colour. A remedy compounded of several substances—myrrh, sandarac, alum, saffron, etc. It was used under the form of liniment, collyrium, electuary, and powder.—Celsus, Galen.

ANTHEREON, Mentum.

ANTHORA, *Aconitum anthora*—a. *Vulgaris*, *Aconitum anthora*.

ANTHORISMA, from *avti*, 'against,' and *eotepa*, 'boundary.' *Tumor diffusus*. A tumour without any defined margin.

ANTHOS, see *Rosmarinus*—a. *Sylvestris*, *Ledum sylvestre*.

ANTHRACIA, from *ανθραξ*, *ανθρακος*, 'coal.' *Carbuncular Eranthem*. An eruption of tumors, imperfectly suppurating, with indurated edges, and, for the most part, a sordid and sanious core. A genus in the order *Eranthematica*, class *Hæmatia* of Good, and including *Plague* and *Yaws*.

ANTHRACIA, *Anthracois*—a. *Pestis*, *Plague*—a. *Rubula*, *Framboesia*.

ANTHRACION, see *Anthrax*.

ANTHRACOID, *Anthracoides*, from *ανθραξ*, *ανθρακος*, 'coal,' and *αδως*, 'resemblance.' (F.) *Charbonneau*. As black as coal. Accompanied by or resembling anthrax.

ANTHRACOMA, *Anthrax*.

ANTHRACONECROSIS, see *Sphacelus*.

ANTHRACOPHYLTIS, see *Anthrax*.

ANTHRACOSIA, *Anthrax*.

ANTHRACOSIS, *Anthracia*, *Carbo Palpebrarum*, from *ανθραξ*, *ανθρακος*, 'a coal.' A species of carbuncle, which attacks the eyelids and globe of the eye.—Paulus of Ægina. Also, a carbuncle of any kind. It has been used for the "black lung of coal miners," which is induced by carbonaceous accumulation in the lungs; *Pseudo-melanotic formation*, (Carswell.) *Matière noire des Pouxons*, *Charbon pulmonaire*. When ulceration results from this cause, *black phthisis*, (F.) *Phthisie avec Mélanose*, exists. See *Melanosis*.

ANTHRACOSIS PULMONUM, see *Melanosis*.

ANTHRACOTYPHUS, *Plague*.

ANTHRAKOKALI, *Lithanthrakokali*, from *ανθραξ*, *ανθρακος*, 'coal,' and *kali*, 'potassa.' An article introduced as a remedy in cutaneous diseases. It is formed by dissolving carbonate of potassa in 10 or 12 parts of boiling water, and adding as much slacked lime as will separate the potassa. The filtered liquor is placed on the fire in an iron vessel, and suffered to evaporate, until neither froth nor effervescence occurs, and the liquid presents a smooth surface like oil. To this, levigated coal is added in the proportion of 160 grammes to 192 grammes of potassa. The mixture is stirred, and removed from the fire, and the stirring is continued, until a black homogeneous powder results. A sulphuretted anthrakokali is made by mixing accurately 16 grammes of sulphur with the coal, and dissolving the mixture in the potassa, as directed above. The dose of the simple and sulphuretted preparations is about two grains three times a day.

ANTHRAX, *ανθραξ*, 'a coal,' *Anthrax*, *Carbo*, *Rubinus verus*, *Codæella*, *Erythema gangranosum*, *Gravitraxum*, *Pruna*, *Persiculus Ignis*, *Pyra*, *Gravitraxum*, *Phyma Anthrax*, *Erythema anthrax*, *Carbunculus*, *Anthracois*, *Anthracois*, *Abcessus gangranosus* seu *gangranosus*, *Furunculus malignus*, *F. gangranosus*, *Carbuncle*, (F.) *Charbon*. An inflammation, essentially gangrenous, of the cellular membrane and skin, which may arise from an internal or external cause. In the latter case it is called *Anthracion*, *Vesicula gangranosus*, *Anthracophlyctis*, (F.) *Pustule maligne*; *Bouton d'Alep*, *Feu Persique* (Persian fire), *Mabat*, *Bouton malin*, *Puce maligne*, and is characterized at the outset by a vesication or bleb filled with a sero-sanguinolent fluid, under which a small induration is formed, surrounded by an areolar inflammation, which becomes gangrenous. It has been thought by some to be induced altogether by contact with the matter of the carbuncle of animals, or of the exuviae of the bodies of such as had died of the disease, but it is now known to arise primarily in the human subject. This form of carbuncle has received different names, many of them from the places where it has prevailed:—*Carbunculus contagiosus* seu *Gallienus* seu *Hungaricus* seu *Polonicus* seu *Septentrionalis*, *Lobus pustulosus Finnicus*, *Pustula gangranosa* seu *Licida Esthoniae*, *Pemphigus Hungaricus*, *Puce de Bourgogne*.

Anthrax is a malignant boil, and its treatment is similar to that which is required in case of gangrene attacking a part.

ANTHRAX PULMONUM, Necropneumonia.
ANTHRISCUS CEREFOLIUM, Scandix cerefolium—a. Humilis, Chærophyllym Sylvestre—a. Procerus, Chærophyllym Sylvestre.
ANTHROPE, Cutis.

ANTHROPIATRICA (MEDICINA), from *an*, 'man,' and *iatros*, 'a physician.' Medicine applied to man, in contradistinction to animals.

ANTHROPOCHEMIA, Chymistry (human).
ANTHROPOCHYMY, Chymistry (human).
ANTHROPOGEN'IA, *Anthropogen'esis*, *Anthropog'eny*, *Geneal' therapy*, from *an* *θρῶπος*, 'man,' and *γενεσις*, 'generation.' The knowledge, or study, or phenomena of human generation.

ANTHROPOGRAPHY, *Anthropograph'ia*, from *an* *θρῶπος*, 'man,' and *γραφῶν*, 'a description.' Anthropology. A description of the human body.
ANTHROPOID, *Anthropoi'des*, from *an* *θρῶπος*, 'man,' and *ειδός*, 'resemblance.' Resembling man:—as the ape.

ANTHROPOLITHUS, from *an* *θρῶπος*, 'man,' and *λίθος*, 'a stone.' The petrification of the human body or of any of its parts. Morbid concretions in the human body.

ANTHROPOL'OGY, *Anthropolog'ia*, from *an* *θρῶπος*, 'man,' and *λογος*, 'a discourse.' A treatise on man. By some, this word is used for the science of the structure and functions of the human body. Frequently, it is employed synonymously with *Natural History and Physiology of Man*.

ANTHROPOMAGNETISMUS, Magnetism, animal.

ANTHROPOMANCY, *Anthropomanti'a*, from *an* *θρῶπος*, 'a man,' and *μαντεία*, 'divination.' Divination by inspecting the entrails of a dead man.

ANTHROPOM'ETRY, from *an* *θρῶπος*, 'a man,' and *μετρον*, 'measure.' Measurement of the dimensions of the different parts of the human body.

ANTHROPOMORPHOLOGY, Anatomy (descriptive).

ANTHROPOMORPHUS, *Atropa mandragora*.
ANTHROPONOMY, *Anthroponomi'a*, from *an* *θρῶπος*, 'man,' and *νομος*, 'law.' A knowledge of the special laws which preside over the functions of the human body in action.

ANTHROPOPHAGUS, (F.) *Anthrophage*, from *an* *θρῶπος*, 'a man,' and *φαγῶν*, 'I eat.' A name given to one who eats his own species.

ANTHROPOPH'AGY, *Anthrophoph'gia*, same etymon. The custom of eating human flesh. A disease in which there is great desire to eat it.

ANTHROPOS, Homo.

ANTHROPOSCOPIA, Physiognomy.

ANTHROPOTOMY, Andranatomia.

ANTHUS, Flos.

ANTHYPNOTIC, *Anthypnot'icus*, *Antihypnot'ic*, *Agrypnot'ic*, from *an*, 'against,' and *ὑπνῶτικός*, 'stupefying.' A remedy for stupor.

ANTHYPOCHONDRIAC, *Anthypochondri'acis*, from *an*, 'against,' and *ὑποχονδριακός*, 'hypochondriac.' A remedy for hypochondriasis.

ANTHYSTER'IC, *Antihyster'ic*, *Antihyster'icis*, from *an*, 'against,' and *ὑστερα*, 'the uterus.' A remedy for hysteria.

ANTI, *avri*, as a prefix, in composition, generally means 'opposition.'

ANTIADÉS, Tonsils.

ANTIADITIS, Cynanche tonsillarlis.

ANTIADON'CUS, from *avri* *adēs*, 'the tonsils,' and *ογκος*, 'tumour.' A swelling of the tonsils.—Swedmar. *Anti'ager* has a similar meaning.

ANTIADONCUS INFLAMMATORIUS, Cynanche tonsillarlis.

ANTIAPOPLEC'TIC, *Antiapoplec'ticus*, *Antapoplec'ticus*, *Apoplec'ticus*, from *an*, 'against,' and *ἀποπληξία*, 'apoplexy.' A remedy for apoplexy.

ANTIARIS TOXICARIA, see Upas.
ANTIARTHRIT'IC, *Antarthrit'ic*, *Antiarthrit'icis*, *Antipodag'ric*, from *avri*, 'against,' and *ἀρθρις*, 'the gout,' (F.) *Antigoutteux*. A remedy for gout.

ANTIASTHEN'IC, *Antiasthén'icis*, from *avri*, 'against,' and *ασθενία*, 'debility.' A remedy for debility.

ANTIASTHMATIC, *Antiasthmat'icis*, from *avri*, 'against,' and *ασθμα*, 'asthma.' A remedy for asthma.

ANTIATROPHICUS, Antatrophic.

ANTIBALLOMENUM, Succedaneum.

ANTIBDELLA, *Antlia sanguisuga*.

ANTIBECHICUS, Expectorant.

ANTIBRACH'IAL, *Antibrach'ialis*. That which concerns the fore-arm.—Bichat. J. Cloquet suggests that the word should be written *antebrachial*, from *ante*, 'before,' and *brachium*, 'the arm'—as *antebrachial region*, *antebrachial aponeurosis*, &c.

ANTEBRACH'IAL APONEURO'SIS, (F.) *Aponévrose antébrachiale*, is a portion of the aponeurotic sheath which envelops the whole of the upper limb. It arises from the brachial aponeurosis, from a fibrous expansion of the tendon of the biceps muscle, from the epicondyle, epitrochlea, and, behind, from the tendon of the triceps brachialis. Within, it is inserted into the cubitus, etc.; and, below, is confounded with the two annular ligaments of the carpus. It is covered by the skin, by veins, lymphatics, and by filaments of superficial nerves; it covers the muscles of the fore-arm, adheres to them, and sends between them several fibrous septa, which serve them for points of insertion.

ANTIBRACHIUM, Fore-arm.

ANTIBRO'MIC, *Antibrom'icis*, from *avri*, 'against,' and *βρωμος*, 'foetor.' A *Deodorizer*. An agent that destroys offensive odours—as chloride of zinc, simple sulphate of alumina, &c.

ANTICACHEC'TIC, *Anticachec'ticus*, *Anticacchym'ic*, from *avri*, 'against,' and *καχέξια*, 'cachexy.' A remedy against cachexy.

ANTICACOCHEMIC, Anticachectic.

ANTICANCEROUS, *Anticancero'sus*, *Anticancero'sus*, *Anticarcinom'atous*, *Antischir'ous*, from *avri*, 'against,' and *καρκινωμα*, 'cancer,' carcinoma. Opposed to cancer.

ANTICANCROSUS, Anticancerous.

ANTICARCINOMATOUS, Anticancerous.

ANTICARDIUM, *Fosselle du cœur*, Scrobiculus cordis.

ANTICATAR'RHAL, *Anticatarrrh'alis*, *Anticatarrrho'icis*, from *avri*, 'against,' and *καταρρῶς*, 'catarrh.' A remedy for catarrh.

ANTICAUSOD'IC, *Anticausod'ic*, *Anticausod'icis*, from *avri*, 'against,' and *καυσος*, 'a burning fever.' A remedy for *causos* or inflammatory fever.

ANTICAUSOTIC, Anticausodie.

ANTICHEIR, Pollex, see Digitus.

ANTICHÉRADICUS, Antiscrofulous.

ANTICHOL'ERIC, *Anticholer'icis*, from *avri*, 'against,' and *cholera*. A remedy against cholera.

ANTICHOLERICA, *Sophora heptaphylla*.

ANTIC'IPATING, *Antic'ipane*, *Antepo'neno*, *Prolept'icis*, (F.) *Anticipant*. An epithet for a periodical phenomenon, recurring at progressively shorter intervals. An *anticipating intermittent* is one in which the intervals between the paroxysms become progressively less.

ANTICNEMIUM, Shin.

ANTICŒUR, Scrobiculus cordis.

ANTICOL'IC, *Anticol'icis*, from *avri*, 'against,' and *κολικός*, 'the colic.' That which is opposed to colic.

ANTICOMMA, *Contre-coup*.

ANTICOPE, *Contre-coup*.

ANTICOPOSCOPIUM, Pleximeter.

ANTICRUSIS, *Contre-coup*.

ANTICRUSMA, *Contre-coup*.

ANTICUS, Anterior.

ANTIDARTREUX, Antierpetic.

ANTIDEIXIS, Counter-indication.

ANTIDIARRHEIC, *Antidiarrhœicus*. A remedy for diarrhœa. Opposed to diarrhœa.

ANTIDINIC, *Antidinicus*, *Din'ic*, *Din'icus*, from *avti*, 'against,' and *δινος*, 'vertigo.' Opposed to vertigo.

ANTIDOTAL, *Antidota'lis*, same etymon as *antidote*. Relating to an antidote; possessed of the powers of an antidote.

ANTIDOTA'RIUM, from *αντιδοτον*, 'an antidote.' A dispensatory. A pharmacopeia or formula. ANTIDOTARY was used formerly for any composition that had the properties of an antidote.

ANTIDOTARY, see Antidotarium.

ANTIDOTE, *Antid'otum*, from *avti*, 'against,' and *δοτω*, 'I give.' Originally, this word signified an *internal remedy*. It is now used synonymously with *counter-poison*, *antiphar'macum*, (F.) *Contre-poison*, and signifies any remedy capable of combating the effect of poison.

A List of reputed Antidotes.

1. METALS.	Chalk.
Iron Filings.	Soap.
Zinc Filings.	
2. ACIDS.	5. SULPHURETS.
Tannic Acid.	Sulphuretted Hydrogen,
Acetic or Citric Acid.	dissolved in water.
	Sulphuret of Potassium.
3. SALTS.	6. HALOIDS.
Alkaline or Earthy Sulphates.	Chlorine.
Chloride of Sodium.	7. METALLIC OXIDES.
Hypochlorite of Soda or of Lime.	Hydrated Sesqui-oxide of Iron.
	Mixed Oxides of Iron.
4. ALKALINES.	8. ORGANIC SUBSTANCES.
Ammonia.	Albuminous Substances.
Carbonates of Ammonia.	(Albumen, Casein, and Gluten.)
Carbonates of Soda.	Starch.
Magnesia.	Oil.
Carbonate of Magnesia.	Animal Charcoal.
Lime Water.	

ANTIDOTUM HERACLIDIS, Enneapharmacos—a. Mithridatium, Mithridate.

ANTIDYNAMICA, Debilitants.

ANTIDYNOUS, Anodyne.

ANTIDYSENTERIC, *Antidysenter'icus*, from *avti*, 'against,' *δυσ*, 'with difficulty,' and *εντερων*, 'intestine.' Opposed to dysentery.

ANTIEMETIC, *Antemet'ic*, *Antiemet'icus*, from *avti*, 'against,' and *εμετικος*, 'emetic.' A remedy for vomiting.

ANTIEPHIALTIC or ANTEPHIALTIC, *Antiephial'ticus*, from *avti*, 'against,' and *εφιαλτης*, 'nightmare.' A remedy for nightmare.

ANTIEPILEPTIC or ANTEPILEPTIC, *Antiepilep'ticus*, from *avti*, 'against,' and *επιληψια*, 'epilepsy.' A remedy for epilepsy.

ANTIFEBRILIS, Febrifuge.

ANTIGALACTIC, *Antigalac'ticus*, *Antilac'teus*, from *avti*, 'against,' and *γαλα*, *γαλακτος*, 'milk.' (F.) *Antilacteuz*. Opposed to the secretion of milk or to diseases caused by the milk.

ANTIGONI COLLYRIUM NIGRUM, *Black collyrium* of ANTIGONUS. It was composed of cadmia, antimony, pepper, verdigris, gum Arabic, and water.

ANTIQUA, see West Indies.

ANTIHÆMOPTICUS, *Anthæmopt'icus*.

ANTIEC'TIC, *Antiech'ticus*, *Anthech'ticus*, from *avti*, 'against,' and *εξis*, 'habit of body.'

The *Antiech'ticum* POTERII is the white oxyd of antimony; also called *Diaphoreticum Jovial'e*.

ANTHELIX, Anthelix.

ANTHELMINTICUS, Anthelmintic.

ANTHEMORRHAG'IC, *Antihæmorrhag'icus*, from *avti*, 'against,' and *αιμορραγια*, 'hemorrhage.' That which is against hemorrhage; an antihemorrhagic remedy.

ANTHEMORRHOID'AL, *Antihæmorrhoida'lis*, from *avti*, 'against,' and *αιμορροιδες*, 'hemorrhoids.' A remedy for hemorrhoids.

ANTHERPET'IC, *Antierpet'icus*, from *avti*, 'against,' and *επρες*, 'herpes.' (F.) *Antidartreux*. A remedy for herpes.

ANTHYDROPHOB'IC, *Antihydrophob'icus*, *Antily'seus*, *Alys'eus*, from *avti*, 'against,' *υδωρ*, 'water,' and *φοβος*, 'dread.' A remedy for hydrophobia.

ANTHYDROP'IC, *Antihydrop'icus*, *Hydrop'icus*, from *avti*, 'against,' and *υδρωψ*, 'dropsy.' A remedy for dropsy.

ANTIHPNOTIC, Anthypnotic.

ANTHYSTERIC, Anthysterie.

ANTI-IC'TERIC, *Anti-icter'icus*, *Icter'icus*, from *avti*, 'against,' and *ικτερος*, 'jaundice.' A remedy for jaundice.

ANTI-IMPETIGENES, SOLOMON'S, see Liquor Hydrargyri oxy muriatis.

ANTILABIUM, Prolabium, see Antelabia.

ANTILACTEUS, Antilactetic.

ANTILAITHEUX, Antilactetic.

ANTILEP'SIS, *Apprehen'sio*, from *avtilap'bas*, 'I take hold of.' The mode of attaching a bandage over a diseased part, by fixing it upon the sound parts.—Hippocrates. The mode of securing bandages, &c., from slipping. Treatment by revulsion or derivation.

ANTILETHAR'GIC, *Antilethar'gicus*, from *avti*, 'against,' and *ληθαργικος*, 'affected with lethargy.' A remedy for lethargy.

ANTILITH'ICS, *Antilith'ica*, *Lith'ica*, from *avti*, 'against,' and *λιθος*, 'a stone.' A substance that prevents the formation of calculi in the urinary organs.

The chief antilithics—according as the calculi are lithic acid or phosphatic—are alkalies or acids; with revellents, especially change of air; tonics, as *diosma crenata* [?], and *uva ursi* [?].

ANTILLY, MINERAL WATERS OF. A celebrated French medicinal spring, near Meaux, in France. The waters have not been analyzed; but astonishing and chimerical effects have been ascribed to them.

ANTIOBIUM, Antitragus, Tragus.

ANTIOI'MIC, *Antioi'micus*, *Antio'mic*, *Antipestilential'is*, from *avti*, 'against,' and *λοιμος*, 'the plague.' A remedy for the plague.

ANTIOPOUS. The *Ant'elope*. (F.) *Gazelle*, from *avbos*, 'a flower,' and *ωψ*, 'the eye'—so called from its beautiful eye. An African animal, whose hoofs and horns were formerly given in hysteric and epileptic cases.

ANTILYSSUS, Antihydrophobic.

ANTIMELANCHOLIC, *Antimelanchol'icus*, from *avti*, 'against,' and *μελαγχολια*, 'melancholy.' A remedy for melancholy.

ANTIMEPHIT'IC, *Antimephit'icus*, from *avti*, 'against,' and *μεφhitic*. A remedy against mephitic or deleterious gases.

ANTIMIASMATIC, *Antimiasmatic'us*, from *avti*, 'against,' and *μιασμα*, *μιαματος*, 'miasma.' A remedy against miasmatic affections.

ANTIMOINE, Antimonium—a. *Beurre d'*, Antimonium muriatum—a. *Chlorure d'*, Antimonium muriatum—a. *Oxide d'*, Algaroth—a. *Oxide blanc d'*, Antimonium diaphoreticum—a. *Soufre doré d'*, Antimonii sulphuretum precipitatum—a. *Sulfure d'*, Antimonium—a. *Sulfuré*, hydrosul-

phure rouge d', Antimonii sulphuretum rubrum—*a. Ferre d'*, Antimonii vitrum.

ANTIMO'NIAL, *Antinomia'lis*, *Stibia'lis*, from *antimonium*, 'antimony.' A composition into which antimony enters. A preparation of antimony.

ANTIMO'NIAL POWDER, *Pulvis antimonialis*, *Ox'idum antimo'ni cum phos'phatē calcis*, *Phosphas calcis stibia'tus seu calc'icum stibia'tum*, *Pulvis Jame'sii seu stibia'tus seu de phos'phatē calcis et stib'ii compositus*, *Facitious JAMES'S Powder*, *SCHWANBERG'S Fever Powder*, *CHENEVIX'S Antimonial Powder*, (F.) *Poudre antimoniale composée ou de JAMES*. A peroxide of antimony combined with phosphate of lime. (*Take of common sulphuret of antimony*, ℥ij; *hartshorn shavings*, ℔ij. Roast in an iron pot, until they form a gray powder. Put this into a long pot, with a small hole in the cover. Keep it in a red heat for two hours, and grind to a fine powder.) This preparation has long been esteemed as a febrifuge: but it is extremely uncertain in its action. The ordinary dose is 6 or 8 grains.

ANTIMONIALE CAUSTICUM, Antimonium muriatum.

ANTIMONIATUM SULPHUR, Antimonii sulphuretum præcipitatum—*a. Tartar*; Antimonium tartarizatum.

ANTIMONII BUTYRUM, Antimonium muriatum—*a. Calx*, Antimonium diaphoreticum—*a. Cerrusa*, Antimonium diaphoreticum—*a. et Potassæ tartaras*, Antimonium tartarizatum—*a. Murias*, Antimonium muriatum—*a. Oleum*, Antimonium muriatum—*a. Oxydulum hydrosulphuratum aurantiacum*, Antimonii sulphuretum præcipitatum—*a. Oxydum*, see *Algaroth*—*a. Oxydum auratum*, Antimonii sulphuratum præcipitatum—*a. Oxydum nitro-muriaticum*, *Algaroth*—*a. Oxydum cum sulphure vitrificatum*, Antimonii vitrum—*a. Oxydum sulphuretum vitrificatum*, Antimonii vitrum—*a. Oxysulphuretum*, *A. sulphuretum præcipitatum*—*a. Potassio-tartaras*, Antimonium tartarizatum—*a. Regulus medicinalis*, Antimonium medicinale—*a. Sal*, Antimonium tartarizatum—*a. Sulphur auratum*, Antimonii sulphuretum præcipitatum—*a. Sulphur præcipitatum*, Antimonii sulphuretum præcipitatum—*a. Sulphuretum*, Antimonium.

ANTIMO'NII SULPHURE'TUM PRÆCIPITA'TUM, *Sulphur antimoniatum*, *Hydrosulphuretum stibio'sum cum sulphurē*, *Ox'o'des stib'ii sulphuratum*, *Oxydulum antimo'ni hydrosulphuratum aurantiacum*, *Oxydum auratum antimo'ni*, *Sulphuretum stib'ii oxydula'ti*, *Hydro-sulfuretum luteum ox'ydi stib'ii sulfurati*, *Sulphur antimo'ni præcipitatum*, *Sulphur auratum antimo'ni*, *Golden Sulphur of Antimony*.

Antimo'ni Sulphure'tum Præcipitatum, *A. Oxysulphuretum*, (F.) *Soufre doré d'Antimoine*, of the London Pharmacopœia, is nearly the same as the old *Kermes* or *Chermes Mineral*. It is a powder of an orange colour, of a metallic, styptic taste. It is emetic, diaphoretic, and cathartic, according to the dose; and has been chiefly used in chronic rheumatism, and in cutaneous affections. Dose, gr. j to gr. iv.

Antimonii Sulphuretum Præcipitatum, of the United States Pharmacopœia, is made by boiling together Sulphuret of Antimony, in fine powder, Solution of Potassa, and distilled water; straining the liquor while hot, and dropping into it Diluted Sulphuric Acid so long as it produces a precipitate.

ANTIMO'NII SULPHURE'TUM RUBRUM, *Red Sulphuret of Antimony*, *Hydrosulphuretum stib'ii rubrum*, *Sub-hydrosulfas stib'ii*, *Hydro-sulphuretum rubrum stib'ii sulfurati*, *Pulvis Carthusianorum*, *Kermes* or *Chermes mineral*, (F.) *Hydro-sulfure rouge d'Antimoine sulfuré*, *Vermillon de*

Provence. Properties the same as the last. Dose, gr. j to gr. iv.

ANTIMONII TARTRAS, Antimonium tartarizatum—*a. Tartaras et Potassæ*, Antimonium tartarizatum—*a. Vitrum hyacinthinum*, Antimonii vitrum—*a. Tersulphuretum*, Antimonium.

ANTIMO'NII VITRUM, *Glass of Antimony*, *Antimo'ni ox'ydum sulphuretum vitrificatum*, *Ox'ydum stib'ii semivitreum*, *Antimo'ni vitrificatum*, *Ox'ydum antimo'ni cum sulphure vitrificatum*, *Vitrum stib'ii*, *Antimo'ni vitrum hyacinthinum*, *Oxydulum stib'ii vitreatum*, (F.) *Verre d'Antimoine*. (Formed by roasting powdered common antimony in a shallow vessel, over a gentle fire, till it is of a whitish gray colour, and emits no fumes in a red heat; then melting it, on a quick fire, into a clean, brownish red glass.) It has been used for preparing the tartarized antimony and antimonial wine.

ANTIMONIOUS ACID, Antimonium diaphoreticum.

ANTIMO'NIUM, of unknown derivation; and by some supposed to be from *anti*, 'against,' and *monos*, 'alone,' i. e. not found alone: or, according to others, from *anti*, 'against,' and *moine*, 'a monk;' because, it is asserted, certain monks suffered much from it. *Stibi*, *Stib'ium*, *Reg'ulus Antimo'ni*, *Mineralium*, *Gynac'um*, *Myne'sia Satur'ni*, *Marcas'ita plum'bea*, *Platyophthal'mon*, *Stin'mi*, *Aurum lepro'sum*, *Antimo'niium erudum*, *Antimo'ni sulphure'tum* (Ph. U. S.), *A. Tersulphure'tum*, *Sulphure'tum stib'ii nigrum*, *Common or Crude Antimony*, *Sulphuret of Antimony*, (F.) *Antimoine*, *Sulfure d'Antimoine*. Sulphuret of antimony is the ore from which all the preparations of antimony are formed. In Pharmacy, it is the native sesquisulphuret of antimony, purified by fusion. When prepared for medical use, by trituration and levigation, it forms a powder, *Antimo'ni sulphure'tum præparatum*, (Ph. D.) of a black, or bluish gray colour, which is insoluble. It is slightly diaphoretic and alterative, and has been used in chronic rheumatism, cutaneous diseases, &c.

ANTIMONIUM ALBUM, Bismuth—a. *Calcinatum*, Antimonium diaphoreticum.

ANTIMO'NIUM DIAPHORETICUM, *Diaphoret'io Antimony*, *Antimo'nius Acid*, *Mineral Bez'oard*, *Antimo'niium Calcinatum*, *Mineral Diaphoret'ic*, *Matière perlée de KERKING*, *Peroxide of Antimony*, *Calx Antimo'ni*, *Antimo'niium diaphoreticium lotum*, *Cerrusa Antimo'ni*, *Calx Antimo'ni elota*, *Ox'o'des stib'ii album*, *Oxydum stibio'sum*, *Oxydum stib'ii album median'te nitro confectum*, *Potassæ biantimo'niæ*, *Deutoxide of Antimony*, (F.) *Oxide blanc d'Antimoine préparé par la moyen du nitre*. (Common antimony, ℔j; purified nitre, ℔ij.—Throw it by spoonfuls into a red-hot crucible; powder and wash. The flowers that stick to the side of the crucible must be carefully separated, otherwise they render it emetic.) Dose, gr. x. to xxx.

ANTIMONIUM EMETICUM, *A. tartarizatum*.

ANTIMO'NIUM MEDICINA'LE, *Reg'ulus Antimo'ni Medicinalis*, *Medic'inal Reg'ulus of Antimony*. (*Antimon. sulphur. 3v Potass. subcarb. 3i Sodii chlorid. 3iv*. Powder, mix, and melt. When cold, separate the scoriae at top, powder the mass, and wash it well.) It is conceived to be more active than common antimony.

ANTIMO'NIUM MURIA'TUM, *Antimo'ni Murias*, *Chloruret of Antimony*, *Chloruretum stib'ii*, *Spuma trium draco'um*, *Deuto-murias stib'ii sublimatus*, *Butter of Antimony*, *Muriate of Antimony*, *Chloride of Antimony*, *Butyrum Antimo'ni seu stib'ii*, *O'leum Antimo'ni*, *Caus'ticum antimoniale*, *Antimonium salitum*, (F.) *Chlorure d'Antimoine*, *Beurre d'Antimoine*. (Common an-

timony and corrosive sublimate, of each equal parts: grind together, and distil in a wide-necked retort, and let the butyraceous matter, that comes over, run in a moist place, to a liquid oil.) A caustic, but not much used as such. Sometimes taken as poison.

ANTIMONIUM SALUTUM, Antimonium muriatum.

ANTIMONIUM TARTARIZATUM, *Tartaras antimoniæ*, *Tartaras Antimoniæ et Potassæ*, *Antimoniæ et Potassæ Tartarus* (Ph. U. S.), *Antimoniæ potassio-tartaras*, *Tartarus Antimoniæ*, *Tartar Antimoniatum*, *Sul Antimoniæ*, *Tartaras Potassæ stibiosus* seu *stibialis*, *Tartarus lixivialis stibialis*, *Deuto-tartaras potassæ et stibii*, *Tartarus emeticus* seu *stibialis*, *Tartarus emeticus*, *Antimonium emeticum*, *Tartarized Antimony*, *Tartrate of Antimony and potassa*, *Potassio-tartrate of Antimony*, *Emetic Tartar*, *Tartar Emetic*, (F.) *Tartre stibie*, *Tartre Émétique*, *Émétique*; in some parts of the United States, vulgarly and improperly called *Tartar*: (Made by digesting sulphuret of antimony in a mixture of nitric and muriatic acids with the aid of heat; filtering the liquor, and pouring it into water: freeing the precipitate from acid, by washing and drying it; adding this powder to bitartrate of potassa in boiling distilled water; boiling for an hour, and after filtering the liquor while hot, setting it aside to crystallize.—Ph. U. S.) Tartarized antimony is emetic, sometimes cathartic and diaphoretic. Externally, it is rubefacient, and in the form of ointment, more especially *unguentum antimoniæ* is used to cause pustulation. Dose, as an emetic, gr. j to gr. iv in solution: as a diaphoretic, gr. one-sixteenth to gr. one-quarter.

The empirical preparation, called *Nonnus's Drops*, consists of a solution of tartarized antimony in rectified spirit, disguised by the addition of some vegetable colouring matter.

ANTIMONIUM VITRIFICATUM, *Antimonium vitrum*.

ANTIMONY, *Antimonium*—a. Butter of, *Antimonium muriatum*—a. Chloride of, *Antimonium muriatum*—a. Chloruret of, *Antimonium muriatum*—a. Common, *Antimonium*—a. Crude, *Antimonium*—a. Deutoxide of, *Antimonium diaphoreticum*—a. Flowers of, *Algaroth*—a. Glass of, *Antimonium vitrum*—a. Golden sulphur of, *Antimonii sulphuretum præcipitatum*—a. Medicinal regulus of, *Antimonium medicinale*—a. Muriate of, *Antimonium muriatum*—a. Peroxide of, *Antimonium diaphoreticum*—a. Potassio-tartrate of, *Antimonium tartarizatum*—a. Sulphuret of, red, *Antimonii sulphuretum rubrum*—a. Tartarized, *Antimonium tartarizatum*—a. Terechloride of, see *Algaroth*—a. Teroxide of, see *Algaroth*—a. Vegetable, *Eupatorium perfoliatum*.

ANTIMONY AND POTASSA, *Tartrate of Antimonium tartarizatum*.

ANTINEPHRITIC, *Antinephretic*, *Antinephreticus*, from *avri*, 'against,' and *νεφρις*, 'nephritis.' A remedy for inflammation of the kidney.

ANTINEUROPATHIC, *Nervine*.

ANTINEUROTIC, *Nervine*.

ANTINIAD, see *Antinial*.

ANTINIAL, from *avri*, 'against,' and *ινω*, 'the ridge of the occiput.' An epithet for an aspect towards the side opposite to the *inion*, or ridge of the occiput.—Barclay. *Antiniad* is used adverbially by the same writer, to signify 'towards the antinial aspect.'

ANTIOCHI HYDRA. A preparation extolled by the ancients in melancholy, hydrophobia, epilepsy, &c. It was formed of gerdander, agaric, pulp of colocynth, Arabian stœchas, opoponax, sagapenum, parsley, aristolochia, white pepper, cinnamon, lavender, myrrh, honey, &c.

ANTIOCHI THERIACA. A theriac employed by Antiochus against every kind of poison. It was

composed of thymæ, opoponax, millet, trefoll, fennel, aniseed, nigella sativa, &c.

ANTIODONTAL'GIC, *Antiodontal'gic*, *Antiodontal'gicus*, *Odontal'gic*, *Odont'ic*, *Antiodontal'gicus*, from *avri* 'against,' and *ὀδονταλγία*, 'tooth-ache.' A remedy for toothache.

ANTIORGAS'TIC, *Antioargas'ticus*, from *avri*, 'against,' and *οργαω*, 'I desire vehemently.' A remedy for orgasm or erethism, and for irritation in general.

ANTIPARALYTIC, *Antiparalyticus*, from *avri*, 'against,' and *παλυσίς*, 'palsy.' Opposed to palsy.

ANTIPARASITIC, *Antiparasiticus*, *Antipheiricus*, *Pigh'icus*, *Parasit'icide*, from *avri*, 'against,' and *παράσιτος*, 'a parasite.' An agent that destroys parasites, as the different vermin that infest the body. The chief antiparasitics are *Cocculus*, *Staphysagria*, *Veratrum album*, and certain of the mercurial preparations.

ANTIPARASTATITIS, from *avri*, 'opposite,' and *παράστασις*, 'the epididymis,' also, 'the prostate,' and *itis*, denoting inflammation. Inflammation of Cowper's glands.

ANTIPATHIA, from *avri*, 'against,' and *πάθος*, 'passion, affection.' Aversion. A natural repugnance to any person or thing.

ANTIPATHIC, *Antipathicus*, (F.) *Antipathique*. Belonging to antipathy. Opposite, contrary,—as *humours antipathiques*; humours opposed to each other. Also, palliatives.

ANTI'PATRI THERIACA, *Theriac of Anti'ATER*. A farrago of more than 40 articles: used as an antidote against the bites of serpents.

ANTIPERIODIC, *Antiperiodicus*, *Antityp'icus*, from *avri*, 'against,' and *περίοδος*, 'period.' A remedy which possesses the power of arresting morbid periodical movements;—e. g. the sulphate of quinia in intermittents.

ANTIPERISTAL'TIC, *Antiperistalticus*, *Antiperist'ular*, from *avri*, 'against,' and *περιστάλω*, 'I contract.' An inverted action of the intestinal tube.

ANTIPERIS'TASIS, from *avri*, 'against,' and *περιστάσις*, 'reunion, aggregation.' A union of opposite circumstances: the action of two contrary qualities, one of which augments the force of the other. The peripateticians asserted, that it is by Antiperistasis, that fire is hotter in winter than in summer. Theophrastus attributes the cause, which renders man more vigorous, and makes him digest more readily in winter, to the augmentation of heat caused by Antiperistasis.

ANTIPERNIUS, from *avri*, 'against,' and *Pernio*, 'a chilblain.' A remedy against chilblains:—as *Unguentum antipernium*, an ointment for chilblains.

ANTIPERTUSSIS, see *Zinci sulphas*.

ANTIPESTILENTIALIS, *Antioleimic*.

ANTIPIARMACUS, *Alexipharmic*.

ANTIPHLOGIS'TIC, *Antiphlogisticus*, *Phlog'gus*, from *avri*, 'against,' and *φλέγω*, 'I burn.' Opposed to inflammation; as *Antiphlogistic remedies*, *A regimen*, &c.

ANTIPH'THERIACA, *Antiphthiriaca*, from *avri*, 'against,' and *φθίριον*, 'I am lousy.' A remedy used to destroy lice.

ANTIPH'THIS'ICAL, *Antiphthis'icus*, from *avri*, 'against,' and *φθίσις*, 'consumption.' Opposed to phthisis.

ANTIPHYS'ETICS, *Carninatives*.

ANTIPHYSICA, *Carninatives*.

ANTIPHYSICAL, *Antiphys'icus*, from *avri*, 'against,' and *φύσας*, 'I blow.' An expeller of wind: a *carninative*.

It has also been used for any thing preternatural; here, the derivation is from *avri*, 'against,' and *φύσις*, 'nature.' The French sometimes say, *Un goût antiphysique*, 'an unnatural taste.'

ANTIPLASTIC, *Antiplast'icus*, *Plastilytic*, *Plastilyticus*, from *αντι*, 'against,' and *πλαστικός*, 'formative.' Antiformative. An agent that diminishes the quantity of plastic matter—fibrin—in the blood.

ANTIPLEURITIC, *Antipleuret'icus*, *Antipleuret'icus*, from *αντι*, 'against,' and *πλευρις*, 'pleurisy.' Opposed to pleurisy.

ANTIPNEUMONIC, *Antipneumon'icus*, from *αντι*, 'against,' and *πνευμονία*, 'disease or inflammation of the lungs.' A remedy for disease or inflammation of the lungs.

ANTIPODAGRIC, *Antiarthritic*.

ANTIPRAXIS, from *αντι*, 'against,' and *πραξις*, 'I act.' A contrary state of different parts in the same patient: e.g. an increase of heat in one organ, and diminution in another.

ANTIPSORIC, *Antipsor'icus*, *Antiscab'ious*, from *αντι*, 'against,' and *ψωρα*, 'the itch.' (F.) *Antigalea*. Opposed to the itch.

ANTIPUTRID, *Antiseptic*.

ANTIPYIC, *Antipy'icus*, from *αντι*, 'against,' and *πυρ*, 'pus.' Opposed to suppuration.

ANTIPYRETIC, *Febrifuge*.

ANTIPYROTIC, *Antipyrot'icus*, from *αντι*, 'against,' and *πυρ*, 'fire.' Opposed to burns or to pyrosis.

ANTIQUARTANA'RUM, *Antiquar'tium*. A remedy formerly used against quartan fever.

ANTIQUOUS, *Chronic*.

ANTIRHACHITIC, *Antirhachit'icus*, from *αντι*, 'against,' and *rachitis*. Opposed to rachitis, or rickets.

ANTIRHEUMATIC, *Antirheumat'icus*; from *αντι*, 'against,' and *ρευμα*, 'rheumatism.' A remedy for rheumatism.

ANTIRHINUM ACUTANGULUM, A. *Linar'ia*—a. *Auriculatum*, A. *Elatine*.

ANTIRHINUM ELATINE, A. *auriculatum*, *Elatine*, *E. hastata*, *Linar'ia elatine*, *Cymbalaria elatine*, *Fluellen* or *Female Speedwell*, was formerly used against scurvy and old ulcerations.

ANTIRHINUM HEDERACEUM, A. *Linar'ia*—a. *Hederasfolium*, A. *Linar'ia*.

ANTIRHINUM LINARIA, A. *hederaceum* seu *hederasfolium* seu *acutangulum*, *Linar'ia*, *L. vulgaris* seu *cymbalaria*, *Elatine* *cymbalaria*, *Cymbalaria muralis*, *Oxyris*, *Urtica*, *Common Tansy*, (F.) *Linar'ia*. Ord. *Scrophulariaceae*. The leaves have a bitterish taste. They are reputed to be diuretic and cathartic. An ointment made from them has been extolled in hemorrhoids.

ANTISCABIOUS, *Antipsoric*.

ANTISCIRRHOSUS, *Anticancerous*.

ANTISCOLETICUS, *Anthelmintic*.

ANTISCOLICUS, *Anthelmintic*.

ANTISCORBUTIC, *Antiscorb'uticus*, from *αντι*, 'against,' and *scorbutus*, 'the scurvy.' Opposed to scurvy.

ANTISCROFULOUS, *Antiscroph'ulous*, *Antiscroful'ous*, *Antistrum'ous*, *Antichærad'icus*. Opposed to scrofula.

ANTISEPTIC, *Antisept'icus*, *Antiputrid*, from *αντι*, 'against,' and *σептос*, 'putrid.' *Antiputredin'ous*. Opposed to putrefaction. The chief antiseptics, internally or externally employed, are *Acidum Muraticum*, *Acidum Nitricum*, *Acidum Sulphuricum*, *Alumina sulphur*, *Carbo Ligni*, *Calx Chlorinata*, *Chlorinum*, *Cinchona* and its active principles, *Cressate*, *Danci Radix*, *Fermentum Cerevisiæ*, *Soda Chlorinata*, and *Zinci Chloridum*.

ANTISIALAGOGUE, *Antisialago'gus*, *Antisial'us*, from *αντι*, 'against,' and *σάλις*, 'saliva.' A remedy against pyalism.

ANTISPASIS, *Derivation*, *Revulsion*.

ANTISPASMODIC, *Antispasmod'icus*, *Antispas'us*, from *αντι*, 'against,' and *σπασμ*, 'I con-

tract.' Opposed to spasm. The whole operation of antispasmodics is probably revulsive. The following are the chief reputed antispasmodics. *Ether Sulphuricus*, *Asafetida*, *Castoreum*, *Dracontium*, *Moschus*, *Oleum Animale Dippelii*, and *Valeriana*—with the mental antispasmodics, abstraction, powerful emotions, fear, &c. Of direct antispasmodics, we have no example.

ANTISPASTICUS, *Antispasmodic*, *Derivative*.

ANTISQUAMIC, *Antisqua'micus*; from *anti*, 'against,' and *squama*, 'a scale.' A name given by Mr. Headland to a medicine, which, by acting on the blood, removes cutaneous affections. The name is inappropriate, as the true scaly diseases—lepra and psoriasis—are not the only ones that require the use of emetropics.

ANTISTASIS, *Antagonism*.

ANTISTERGYMA, from *αντι*, 'against,' and *στηνυγμα*, 'a support.' A fulcrum, support, crutch.—*Hippocrates*.

ANTISTERNUM, from *αντι*, 'against,' and *στενον*, 'the sternum.' The back.—*Rufus*.

ANTISTRUMOUS, *Antiscrofulous*.

ANTISYPHILITIC, *Antisyphilit'icus*, from *αντι*, 'against,' and *σφιλίς*, 'the venereal disease.' Opposed to the venereal disease.

ANTITASIS, *Counter-extension*.

ANTITHENAR, *Opponens pollicis*, *Adductor pollicis pedis*.

ANTITHERMA, *Refrigerants*.

ANTITHORA, *Aconitum anthora*.

ANTITRAGICUS, *Antitra'geus*, (F.) *Muscle de l'Antitragus*, *M. antitragien*.—(Ch.) Belonging to the antitragus. A small muscle is so called, the existence of which is not constant. It occupies the space between the antitragus and anthelix.

ANTITRAGIEN, *Antitragicus*.

ANTITRAGUS, from *αντι*, 'opposite to,' and *τραγος*, 'the tragus,' *Antilo'bium*, *Oblo'bium*. A conical eminence on the pavilion of the ear, opposite the tragus.

ANTITYPIA, from *αντι*, 'against,' and *τυπη*, 'I strike.' Resistance. Hardness. Repercussion.

ANTITYPICUS, *Antiperiodic*.

ANTIVENEREAL, *Antivene'reus*, from *αντι*, 'against,' and *venus*, 'Venus.' The same as *Antisyphilitic*. Formerly it was used synonymously with *Antaphrodisiac*.

ANTIVERMICULAR, *Antiperistaltic*.

ANTIVERMINOSUS, *Anthelmintic*.

ANT'LIA or **ANTLI'A**, from *αντλειν*, 'to pump out.' A syringe; a pump. Hence, *Antlia lac'tea*, *Lactisug'ium*, a breast-pump; and *Antlia sanguisug'a*, *Antibell'la*, *Hiru'do artificia'lis*, the exhausting syringe used in cupping.

ANTLIA GASTRICA, *Stomach-pump*.

ANTODONTALGIC, *Antiodontalgic*.

ANTODYNES, *Anodyne*.

ANTRAX, *Anthrax*.

ANTRE, *Antrum*—a. d'Hyghmore, *Antrum of Highmore*.

ANTROVERSI, *Anteversio*.

ANTRUM, 'a cavern,' *Cavern'a*, *Bar'athrum*, (F.) *Antre*. A name given to certain cavities in bones, the entrance to which is smaller than the bottom.

ANTRUM AURIS, *Tympanum*—a. *Buccinosum*, *Cochlea*, *Labyrinth*—a. *Dentale*, see *Tooth*—a. *Pylori*, see *Stomach*.

ANTRUM OF HIGHMORE, *Antrum Highmore'i*. *Antrum seu Gena seu maxilla're seu maxil'la superio'ris*, *Genyan'trum*, *Gnathan'trum*, *Maxillary Sinus*, *Sinus Gena' pituita'rius*, (F.) *Antre d'Hyghmore*, *Sinus Maxillaire*. A deep cavity in the substance of the superior maxillary bone communicating with the middle meatus of the nose.

It is lined by a prolongation of the Schneiderian membrane.

ANTSJAR, Upas.

ANULARIS, Annular.

ANULUS, *Fossette*.

ANURESIS, Ischuria.

ANURIA, Ischuria.

ANUROCRINIE, see Ischuria.

ANURORRHÉE, see Ischuria.

ANUS, 'a circle,' *Podex, Potex, Mol'ynē, Mol'yn'is, Dactyl'ios, Cath'edra, Cyr'ceon, Cys'saros, Cynthos, Aph'edra, Aph'edron, Hedra, Proctos, Archos, Sedes, Culus, Cu'leon*. The circular opening situate at the inferior extremity of the rectum, by which the excrement is expelled. The *fundament*. The *body*. The *seat*, (F.) *Siège, Fondement*.

ANUS also signifies the anterior orifice of the *Aqueduct* of Sylvius. By some, this *Anus*, called, also, *Fora'men commu'nē posterius*, has been supposed to form a communication between the back part of the third ventricle and the lateral ventricles. It is closed up, however, by the tela chorioidea, and also by the fornix, which is intimately connected with this. The foramen is situate between the commissura mollis of the optic thalami and the pineal gland.

ANUS, ARTIFICIAL. An opening made artificially, to supply the place of the natural anus. The term is often used to include preternatural anus.

ANUS CEREBRI, see *Aqueductus Sylvii*.

ANUS, CONTRACT'ED, (F.) *Anus rétréci*. A state of the anus when, from some cause, it is contracted.

ANUS, IMPER'FORATE. A malformation, in which there is no natural anus. See *Atresia* and *adnata*.

ANUS, PRETERNAT'URAL, (F.) *Anus contre nature, A. anormal*. An accidental opening which gives issue to the whole or to a part of the fæces. It may be owing to a wound, or, which is more common, to gangrene attacking the intestine in a hernial sac.

This term is also employed, as well as *Anus devic, de'vius anus*, to the case where the anus, instead of being in its natural situation, is in some neighbouring cavity, as the bladder, vagina, etc.

ANXIETAS, Anxiety—a. *Præcordiorum*, see *Anxiety*.

ANXI'ETY, *Anxi'etas, Anxi'etude, Adæmo'nia, Dyspho'ria anxi'etas, Alys'mus, Al'yce, Al'ysie, Asē*, (F.) *Anxiété*, from *angere, Gr. αγγειν*, 'to strangle, to suffocate.' A state of restlessness and agitation, with general indisposition, and a distressing sense of oppression at the epigastrium, *Anxi'etas præcordio'rum*. *Inquietude, anxiety, and anguish*, represent degrees of the same condition.

ANYPNIA, Insomnia.

AOCHLE'SIA, from *a, priv.*, and *οχλος*, 'disturbance.' *Tranquillity, calmness*.

AOR'TA, *Arte'ria magna seu crassa seu max'ima, Hæmal Axis*, of Owen, (F.) *Aorte*. This name was given by Aristotle to the chief artery of the body. It may have been derived from *αορτισμα*, 'I am suspended,' as it seems to be suspended from the heart; or from *αορ*, 'air,' and *τροω*, 'I keep,' because it was supposed to contain air. It is probable that Hippocrates meant by *aorai* the bronchia and their ramifications. The aorta is the common trunk of the arteries of the body. It arises from the left ventricle of the heart, about opposite to the fifth dorsal vertebra, passes upwards (*ascending Aorta*), forms the *great arch of the Aorta*, and descends along the left of the spine (*descending Aorta*), until it reaches the middle of the fourth or fifth lumbar vertebra, where it bifurcates, to give origin to the common iliacs. The aorta is sometimes divided into the

Thoracic or Pectoral, and the *Abdominal*. For the arteries which arise from it, etc., see *Artery*.

AORTASIE, Aorteurysma.

AORTECTASIE, Aorteurysma.

AORTEURYS'MA, from *αορτη*, 'the aorta,' and *ευρυς*, 'dilated.' *Aneurism of the Aorta*, (F.) *Aneurysme de l'Aorte, Aortectasie, Aortasie*.—*Piurisy*. By carefully auscultating over the dorsal vertebra, a bellows' sound, with a deep and not always perceptible impulse, may be detected.

AORT'IC, *Aorticus*. Relating to the Aorta. The *Aortic ventricle*, (F.) *Ventricule Aortique*, is the left ventricle. The *Aortic valves* are the sigmoid valves at the origin of the Aorta, etc.

AORTITIS, *Inflammat'io Aor'tæ*, from *Aorta*, and *itis*, denoting inflammation. Inflammation of the aorta.

AORTA, *Aortron*. A lobe of the lungs.—*Hippocrates*.

AO'TUS, from *a*, privative, and *οὖς, ωτος*, 'an ear.' A monster devoid of ears.—*Gurli*.

APAG'MA, *Apoclas'ma, Apocœaulis'menon*, from *απο*, 'from,' and *αἶμα*, 'I remove.' Separation, abduction. Separation of a fractured bone.—*Galenus, Fœsius*.

APAGOGÉ, Defecation, Inductio.

APALACHINE, *Ilex vomitoria*—a. *à Feuilles de Prunier*, *Prinos*—a. *Gallis, Ilex vomitoria*.

APAL'LAGÉ, *Apallax'is*, from *απαλλαττω*, 'I change.' Mutation, change. It is generally taken in a good sense, and means the change from disease to health.—*Hippocrates*.

APALLAXIS, Apallage.

APALOT'ICA, from *απαλότης*, 'softness, tenderness.' Fortuitous lesions or deformities affecting the soft parts. The first order in the class *Tychica*, of Good.

APANTHESIS, Apanthismus.

APANTHIS'MUS, *Apanthē'sis*, from *απο*, 'from,' and *ανθεω*, 'I flower.' The obliteration of parts previously inservient to useful purposes, as of the ductus venosus and ductus arteriosus, which are essential to foetal existence, but are subsequently unnecessary. See also *Stuprum*.

APANTHRO'PIA, from *απο*, 'from,' and *ανθρωπος*, 'man.' Detestation of man; desire for solitude.—*Hippocrates*. One of the symptoms of hypochondriasis.

APAPHRISMOS, Despumation.

APARACH'TUM VINUM, from *a*, priv., and *παρυχω*, 'I pour over.' The purest wine: that which has not been mixed with sea-water.—*Galen*.

APARINE, Gallum aparine—a. *Hispidæ, Gallium aparine*.

APARTHROSIS, Diarthrosis.

AP'ATHY, *Apathi'a, Ameli'a*, from *a*, privative, and *παθος*, 'affection,' (F.) *Apathie*. Accidental suspension of the moral feelings. It takes place in very severe diseases, particularly in malignant fevers.

APECHE'MA, from *απο*, 'from,' and *ηχος*, 'sound.' Properly the action of reflecting sound. In medicine, it is synonymous with the Latin *Contrafissura*, a counter-fissure, a counter-blow.—*Gorrieus, Celsus*.

APECTOCEPHALUS, Acephalothorus.

APEL'LA, *Appel'la, Leipoder'mos, Recuti'tus*, from *a*, priv., and *pellis*, 'skin.' One whose prepuce does not cover the glans.—*Galenus, Linnaeus, Vogel*. Retraction or smallness of any other soft appendage.—*Sagar*. One who is circumcised.

APEPSIA, Dyspepsia.

APERIENT, *Aperiens, Aperiti'vus*, from *aperire* (ab and *pario*), 'to open.' *Res'crans*. A laxative. (F.) *Apéritif*. A medicine which gently opens the bowels. The term had formerly a much more extensive signification, and,

like *Catalyticum*, was given to a substance supposed to have the power of opening any of the passages, and even the blood-vessels.

APERIS'TATON, *Aperis'tatum*, from *a*, privative, and *περιστημι*, 'I surround.' An epithet for an ulcer not dangerous nor considerable nor surrounded by inflammation.

APÉRITIF, Aperient.

APERITIVUS, Aperient.

APERTOR OCULI, Levator palpebræ superioris.

APERTO'RIMUM, from *aperio*, 'I open.' An instrument for dilating the os uteri during labour.

APERTURA, Mouth.—*a*. Anterior ventriculi tertii cerebri, Vulva (cerebri).—*a*. Pelvis superior, see Pelvis.

APEUTHYSMENOS, Rectum.

APEX, *Mucro*. The point or extremity of a part,—as the apex of the tongue, nose, etc.

APEX LINGUÆ, Proglottis.

APHERESIS, Apheresis, Extirpation.

APHALANG'IASIS, from *a*, 'intensive,' and *φαλαγγ*, 'phalanx.' The fourth stage of Oriental leprosy, which is recognised chiefly by a gangrenous condition of the fingers.

APHASSOM'ENOS, from *αφασσω*, 'I touch, I feel.' The touching of the parts of generation of the female as a means of diagnosis.—Hippocrates. See Esaphe.

APHEDRA, Anus.

APHEDRIA, Menses.

APHEDRON, Anus.

APHELI'A, *αφελος*, 'simple.' Simplicity. The simple manners of the sect of Methodists in teaching and practising medicine.

APHELX'IA, from *αφελω*, 'I abstract.' Voluntary inactivity of the whole or the greater part of the external senses to the impressions of surrounding objects, during wakefulness. *Revery*, (F.) *Réverie*. Dr. Good has introduced this into his Nosology, as well as *Aphela'ia* *socors* or absence of mind—*A. intentæ* or abstraction of mind: and *A. otio'sa*, *Stu'dium* *ina'næ*, *Cenospul'dia*, *brozen study* or listless musing.

APHEPSEMA, Decection.

APHEPSIS, Decection.

APHE'RESIS, *Apha'resia*, from *αφαίρω*, 'I take away.' An operation by which any part of the body is separated from the other. Hippocrates, according to Foesius, uses the expression *Apha'resis San'guinis* for excessive hemorrhage; and Sennertus, to express the condition of an animal deprived both of the faculties of the mind and of the mind itself.

APHE'SIS, from *αφηνμι*, 'I relax.' A remission. This word expresses sometimes the diminution or cessation of a disease; at others, languor and debility of the lower extremities. See Languor, and Remission.

APHILANTHROPY, *Aphilanthro'pia*, from *a*, privative, *φιλω*, 'I love,' and *ανθρωπος*, 'a man.' Dislike to man. Love of solitude. Vogel has given this name to the first degree of melancholy.

APHISTESIS, Abscess.

APHODEUMA, Excrement.

APHODUS, Excrement.

APHONETUS, Aphonus.

APHON'IA, *Liga'tio* *linguæ*, *Loque'la* *abol'ita*, *Defectus* *loque'læ*, *Dyspho'nia*, (of some,) *Aph'ony*, (F.) *Aphonie*, *Perte de la Voix*, from *a*, privative, and *φωνη*, 'voice.' Privation of voice, or of the sounds that ought to be produced in the glottis. When aphonia forms part of catarrh or of 'cold,' it is commonly but of little consequence; but when produced by causes acting on the nervous system, as by some powerful emotion, or without any appreciable lesion of the vocal apparatus, (*Laryngo-paralysis*), it frequently resists all remedies.

APHONIA, Catalepsy—a. *Surdorum*, *Mutitas* *Surdorum*.

APHONICUS, Aphonus.

APHO'NUS, *Apho'nicus*, *Apho'netus*; same ety.mon. Relating to aphonia.

APHONYA, Aphonia.

APHORIA, Sterilitas.

APHORICUS, Sterile.

APHORUS, Sterile.

APHOR'ME, *αφορμη*, 'occasion.' The external and manifest cause of any thing. The occasional cause of a disease.—Hippocrates.

APHRO'DES, 'frothy,' from *aphros*, 'foam,' and *eidos*, 'resemblance.' Applied to the blood and the excrements.—Hippocrates.

APHRODISIA. Coition, Puberty.

APHRODIS'ACUS, *Aphrodisiacus*, from *Aphroditē*, 'Venus,' (F.) *Aphrodisiaque*. Medicine or food believed to be capable of exciting to the pleasures of love; as ginger, cantharides, &c. They are generally stimulants.

APHRODISIACUS, Venereal.

APHRODISIASMUS, Coition.

APHRODISIOGRAPHY, from *Aφροδιτη*, 'Venus,' and *γραφω*, 'I describe.' Etymologically, this term means a description of the pleasures of love, but it has been placed at the head of a work describing the venereal disease.

APHRODITIC, Venereal.

APHROG'ALA, from *aphros*, 'foam,' and *γαλα*, 'milk.' *Lac spurcolum*. A name formerly given to milk rendered frothy by agitation.

APHRONIA, Apoplexy.

APHRONITRUM, Natrum, Soda.

APHROSYNE, Delirium, Insapity.

APIITHÆ, *Aphthæ*, *Aphthæ*, from *απτω*, 'I inflame.' Thrush or sore mouth, *Aphtha lacte'cimen* seu *Infan'tum*, *Lacte'cimen*, *Lactucim'ina*, *Lactud'mina*, *Al'cola*, *Em'phylisis* *aphtha*, *Ulcera serpentina* *oris*, *Pustula* *oris*, *Febris aphtho'sa*, *Angi'na aphtho'sa*, *Vesicula* *gingiva'rum*, *Stomatit'is exsudat'iva* seu *vesiculosa* *infan'tum*, *Stomac'yra*, *S. aphtha*, *Prunel'la*, *White Thrush*, *Milk Thrush*. Aphthæ consist of roundish, pearl-coloured vesicles, confined to the lips, mouth, and intestinal canal, and generally terminating in curd-like sloughs. In France, the Aphthæ of children, *Aphthes des Enfants*, is called *Muguet*, *Millet*, *Blanchet*, *Catarrhe buccal* and *Stomatite crèmeuse* *pultacee*, *Pultaceous inflammation of the Mouth*; and generally receives two divisions—the *mild* or *discreet*, (F.) *Muguet bénin* ou *discreet*, and the *malignant*, (F.) *Muguet malin* ou *confluent*, the *Black Thrush*. Common Thrush is a disease of no consequence, requiring merely the use of absorbent laxatives. The malignant variety, which is rare, is of a more serious character, and is accompanied with typhoid symptoms.—*Typhus aphthoides*.

APHTHÆ ADULTORUM, Stomatitis, aphthous—a. Præputii, Herpes præputii—a. Serpentes, Cancer aquaticus.

APH'THAPHYTE, *Aphthaph'yton*: *Ofid'ium albicans*, *Champignon du Muguet*, from *αφθα*, 'aphthæ,' and *φυτον*, 'a vegetable.' A parasitic vegetable growth observed in aphthæ.

APHTHÆ GANGRÊNEUX, Cancer aquaticus.

APHTHES DES ENFANS, Aphthæ.

APHTHEUX, Aphthous.

APHTHO'DES, *Aphthoides*, *Aphthoideus*, from *aphthæ*, and *eidos*, 'resemblance.' Aphthous-like. Resembling aphthæ.

APHTHOUS, *Aphtho'sus*, (F.) *Aphtheux*. Belonging to aphthæ; complicated with aphthæ; as *Aphthous Fever*.

APIASTRUM, Melissa.

APICES CRURUM MEDULLÆ OBLON-
GATÆ, Corpora striata—a. Digitorum, Pu-
pule.

APILEPSIA, Apoplexy.

APIOL, *Apiolum*, Parsley oil, from *apium*,
'parsley,' and *oleum*, 'oil.' A yellowish oily
liquid, obtained from *apium petroselinum* or
parsley, which, in the dose of 15 grains, has
been used as an antiperiodic.

APIONTA, see Excretion.

APIOS, *Pyrus communis*.

APIS, Bee.

APITES, from *απιον*, 'a pear.' Perry.—Gor-
raus.

APIUM, A. graveolens—a. Ammi, Ammi—a.
Anisum, Pimpinella anisum—a. Carvi, Carum.

APIUM GRAVEOLENS, *Apium*, *Paludo'pium*,
Bellinum, *Seseli graveolens*, *Sium graveolens* seu
apium, *Snallage*, (F.) *Ache*. Ord. Umbelliferae.
Sex. Syst. Pentandria Digynia. The plants,
roots, and seeds are aperient and carminative.
Sclery is a variety of this.

APIUM HORTENSE, A. graveolens—a. Monta-
num, *Athamanta rupestrinum*—a. *Paludapium*,
A. *Graveolens*—a. *Petraeum*, *Bubon Macedonicum*.

APIUM PETROSELINUM, *Apium Hortense* seu
vulgarē, *Eteoselinum* [?], *Grielum*, *Petroseli-*
nium, *Common Parsley*, (F.) *Persil*. The root—
Petroselinum, (Ph. U. S.)—and seeds are diuretic
and aperient. See *Apiol*.

APIUM SIUM, *Sium nodiflorum*—a. *Vulgare*, A.
graveolens.

APLASTIC, *Aplasticus*, from *a*, privative,
and *πασσω*, 'I form.' That which is not capable
of forming; or is not organizable.

APLASTIC ELEMENT: one which is unsuscep-
tible of any farther amount of organization.—
Gerber.

APLESTIA, Inguivies, Intemperance.

APLEUROS, from *a*, privative, and *πλευρον*,
'a rib.' One without ribs, or pleura.—Hippo-
crates, Galen.

APLOTOMY, *Aplotomia*, from *απλος*, 'sim-
ple,' and *τομω*, 'I cut.' A simple incision.

APNÉE, Apnoea.

APNEUMATOSIS, see Atelectasis.

APNEUMIA, from *a*, priv., and *πνευμων*, 'lung.'
A monstrosity in which there is absence of lungs.

APNEUMONERVIA, *Apneumonēria*, from
a, priv., *πνευμων*, 'lung,' and *νευρον*, 'nerve.'
Want of nervous action in the lungs.

APNEUSTIA, Apnoea, Asphyxia.

APNŒA, from *a*, privative, and *πνέω*, 'I re-
spire.' (F.) *Apnée*, Absence of respiration, *Re-*
spiratio abolita; or insensible respiration. Also,
Orthopnoea. See Asphyxia.

APNŒA INFANTUM, Asthma Thymicum.

APNŒASPHYXIA, Asphyxia.

APNUS, *apnoos*, same etymon. One devoid of
respiration. An epithet applied by authors to
cases in which the respiration is so small and
slow, that it seems suspended.—Castelli. It is
probable, however, that the word was always ap-
plied to the patient, not to the disease.

APO, *apo*, a prefix denoting 'from, of, off, out.'
Hence—

APOBAMMA, Embamma.

APOBAINON, Eventus.

APOBESOMENON, Eventus.

APOBIOSIS, Death.

APOBLEMA, Abortion.

APOBOLE, Abortion.

APOBRASMA, Furfur.

APOCAPNISMUS, Fumigation.

APOCATASTASIS, Considentia, Restauratio.

APOCATHARSIS, Catharsis.

APOCATHARTICUS, Cathartic.

APOCECAULISMENON, Apagma.

APOCENO'SIS, *Apoceno'sis*, from *apo*, 'out,'
and *κένωσις*, 'evacuation.' A partial evacuation
according to some, in opposition to *Cenosis*, which
signifies a general evacuation.—Cullen and Swe-
diar apply it to morbid fluxes.

APOCENOSIS, Abvacuation—a. *Diabetes melli-*
tus, *Diabetes*—a. *Ptyalismus mellitus*, see *Saliva-*
tion—a. *Vomitus pyrosis*, *Pyrosis*.

APOCHOREON, Excrement.

APOCHREMMMA, Sputum.

APOCHREMPSIS, Exspuition.

APOCHYMA, from *αποχω*, 'I pour out.' A
sort of tar, obtained from old ships, which is im-
pregnated with chloride of sodium. It was used
as a discutient of tumours.—Aëtius, Paulus,
Gorraus.

APOCIN GOBE-MOUCHE, *Apocynum an-*
drosamifolium.

APOCLASMA, Abduction, Apagna.

APOCLEISIS, Asitia, Disgust.

APOC'OPE, from *apo*, and *κοπεῖν*, 'to cut.'
Abcession. A wound with loss of substance.
Fracture with loss of part of a bone. Amputation.

APOCOPUS, Castratus.

APOCRISIS, Contagion, Excrement, Secre-
tion.

APOCROUS'TIC, *Apocroustica* seu *Apocrus-*
tica, (remed'ia), from *apo*, 'out,' and *κρούω*, 'I
push.' An astringent and repellent.—Galenus.

APOCRUSTICA, Apocroustic.

APOCYESIS, Parturition.

APOCYN, see *Apocynum Cannabinum*.

APOCYNIN, see *Apocynum Cannabinum*.

APOC'YNUM ANDROSÆMIFOLIUM, (Ph.
U. S.) from *apo*, and *κύνω*, 'a dog,' because es-
teemed, of old, to be fatal to dogs. *Dog's Bone*,
Bitter Dog's Bone, *Milkweed*, *Bitterroot*, *Honey-*
bloom, *Catchfly*, *Flytrap*, *Ip'ceas*, *American Ip'-*
ceas, (F.) *Apocin gobe-mouche*, A. *amar.* Ord.
Apocynaceae. *Sex. Syst.* Pentandria Digynia.
The root of this plant is found from Canada to
Carolina. Thirty grains evacuate the stomach
as effectually as two-thirds of the amount of
Ipecacuanha, by which name it is known in var-
ious parts of the Eastern States. It is in the
secondary list of the Pharmacopœia of the United
States.

APOCYNUM CANNABINUM, (Ph. U. S.) *Indian*
Hemp. This American plant possesses emetic,
cathartic, diaphoretic and diuretic properties, has
been strongly recommended in dropsy, and
has been given in decoction, —3ij of the root
boiled in three pints of water to two. A wine-
glassful for a dose. An active principle, *Apocyn*
or *Apocynin*, has been extracted from the root.

APOCYNUM HYPERICEFOLIUM, a variety of A.
cannabinum, which, as well as A. *androsamifol-*
ium, abounds in a milky juice. This, when ap-
plied to the skin, produces a troublesome erup-
tion resembling flea-bites: hence, the plants have
been termed, by the voyageurs in the Hudson's
Bay territory, *Herbes à la puce*.

APOCYNUM NOVE ANGLIÆ HIRSUTUM, *Ascle-*
pias tuberosa—a. Orange, *Asclepias tuberosa*—a
Scandens, *Allamanda*.

APODACRYTICUS, *Delachrymaticus*, from
apo, 'from,' and *δακρῶω*, 'I weep.' A substance,
supposed to occasion a flow of the tears, and then
to arrest them.—Columella, Pliny, Galenus.

APODEMIALGIA, Nostalgia.

APOD'IA, from *a*, privative, and *πους*, *ποδος*,
'a foot.' Want of feet; hence *Apous* or *Apus*,
one who has no feet.

APODYTERIUM, *Conistarium*, *Spoliatē-*
rium, *Spoliatium*, from *αποδυω*, 'I strip off.' The
ante-room, where the bathers stripped themselves
in the ancient gymnasia.

APOGALACTISMUS, Weaning.

APOGALACTOS, Exuber.

APOGEUSIS, Ageusia.

APOGEUSTIA, Ageusia.

APOGLAUCOSIS, Glaucosis.

APOGON, Imberbis.

APOG'ONUM, from *apo*, and *γίνομαι*, 'I exist.'

A living fetus in utero.—Hippocrates.

APOLEPISIS, Desquamation.

APOLEPISMUS, Desquamation.

APOLEP'SIS, *Apolep'sis*, *Apolip'sis*, from *απολαμβάνω*, 'I retain.' Retention, suppression.—Hippocrates. Asphyxia.

APOLEX'IS, from *απολγω*, 'I cease.' Old age, decrepitude.

APOLINO'SIS, from *apo*, and *λινον*, 'a flaxen thread.' The mode of operating for fistula in ano, by means of a thread of *Hemolinon* or *Linum crutum*.—Hippocrates, Paulus.

APOLIPSIS, Apolepsis.

APOLLINARIS ALTERCUM, Hyosecyamus.

APOLUTICA, Cicatrisantia.

APOLYS'IA, *Apol'ysis*, from *απολύνω*, 'I loosen.'

Solution. Relaxation. Debility of the limbs or looseness of bandages.—Erolian. Expulsion of the fetus and its dependencies. Termination of a disease.—Hippocrates, Galen.

APOMATHE'MA, *Aponathé'sis*, from *apo*, and *μαθάνω*, 'I learn.' Forgetfulness of things taught.—Hippocrates.

APOM'ELI, from *apo*, 'of,' and *μελι*, 'honey.' An oxymel or decoction made of honey.—Galen, Aëtius, Paulus, &c.

APOMEXIS, Munctio.

APOMYLE'NAS, from *απομύλλαινω*, 'I make a wry mouth.' One who pushes his lips forwards, pressing them against each other. Occasionally a symptom of nervous fever.—Galen, Erolian.

APOMYTHO'SIS, *Apomytho'sis*, from *απορυσσω*, 'I snore.' A disease in which there is stertor.—Sauvages, Sagar.

APOMYXIA, Nasal mucus.

APONEUROGRAPHY, *Aponeurogra'phia*, from *απονευρωσις*, an 'aponeurosis,' and *γραφη*, 'a description.' A description of the Aponeuroses.

APONEUROLOG'Y, *Aponeurolog'ia*, from *απονευρωσις*, 'an aponeurosis,' and *λογος*, 'a discourse.' *Aponeurosiology*. The anatomy of the aponeuroses.

APONEUROSIOLOGY, Aponeurology.

APONEURO'SIS, *Aponeuro'sis*, from *apo*, 'from,' and *νευρον*, 'a nerve.' *Pronerva'tio*, *Denerva'tio*, *Enerva'tio*, *Expan'sio nervo'sa*, (F.) *Aponeurose*, *Aponévrose*. The ancients called every white part *νευρον*, and regarded the Aponeurosis as a nervous expansion. The Aponeuroses are white, shining membranes, very resisting, and composed of fibres interlaced. Some are continuous with the muscular fibres, and differ only from tendons by their flat form. They are called *Aponeuroses of insertion*, (F.) *Aponévroses d'insertion*, when they are at the extremities of muscles, and attach them to the bone;—*Aponeuroses of intersection*, (F.) *Aponévroses d'intersection*, if they interrupt the continuity of the muscle, and are continuous on both sides with muscular fibres. Others surround the muscle, and prevent its displacement: they are called *enveloping Aponeuroses*, (F.) *Aponévroses d'enveloppe*.

APONEUROSIS, Fascia.—a. Crural, Fascia lata.—a. Femoral, Fascia lata.—a. Iliac, Fascia iliaca.

APONEUROSIT'IS, from *aponeurosis*, and *itis*, 'denoting inflammation.' Inflammation of an aponeurosis.

APONEUROT'IC, *Aponeurot'icus*. What relates to Aponeuroses:—thus, we say *Aponeurotic expansion*, *Aponeurotic muscle*, &c.

APONEUROTOMY, *Aponeurotom'ia*, from *απονευρωσις*, 'aponeurosis,' and *τέμνω*, 'I cut.' Anatomy of aponeuroses.

Aponeurotomy has, also, been proposed for the division, (*débridement*) of filaments, &c., in aponeurotic openings, and for the section of fasciae.

APONÉVROSE PÉDIEUSE, see Pedal Aponeurosis.—a. *Superficielle de l'Abdomen et de la Cuisse*, Fascia superficialis.

APONÉVROSIS, Aponeurosis.

APON'IA, from *a*, privative, and *παιος*, 'pain.' Freedom from pain.

APONIP'SIS, Ablution.

APOPALLE'SIS, *Apopal'sis*, from *αποπαλλω*, 'I throw off.' Expulsion. Protrusion.—Hippocrates. Also, Abortion.

APOPATE'MA, *Apop'athos*, *Apop'atus*. The excrement, and the place where it is deposited.—Dioscorides, Erolian.

APOPEDasIS, Luxation.

APOPHLEGMATISANS PER NARES, Er rhine.—a. per Os, Sialogogue.

APOPHLEGMATISANT'IA, *Apophlegmatison'ta*, *Apophlegmatism'i*, from *apo*, 'out,' and *φλεγμα*, 'phlegm.' Medicines which facilitate the upward expulsion of mucus from the mucous membrane of the digestive or air passages; as gargles, masticatories, &c.

APOPHLEG'MATISM, *Apophlegmatism'us*. The action of Apophlegmatisantia.—Galen.

APOPHLEGMATISMI, Apophlegmatisantia.

APOPH'RADES, from *αποφρας*, 'unlucky.'

An epithet applied to unlucky days, (*dies nefandi*.) Days on which a favourable change is not expected to occur in a disease.—A. Laurentius.

APOPH'RAxis, Amenorrhœa.

APOPHTHAR'MA, *Apoph'thora*, from *apo*, and *φθίρω*, 'I corrupt.' Abortion, as well as a medicine to procure abortion.

APOPHTHORA, Abortion.

APOPHTHORIUS, Abortive.

APOPHY'ADES, from *apo*, 'from,' and *φύω*, 'I spring.' The ramifications of veins and arteries.—Hippocrates.

APOPHYSE BASILAIRE, Basillary process.—a. *Engainante ou vaginale*, Vaginal process.—a. *Pyramidale*, see Temporal Bone.—a. *Pétrée*, see Temporal Bone.

APOPHYSES ÉPINEUSES, Spinous processes of the vertebrae.

APOPH'YSIS, from *apo*, 'from,' and *φύω*, 'I rise,' *Ec'physis*, *Proces'sus*, *Appendix*, *Prominentia ossis contin'ua*, *A process of a bone*. When the apophysis is yet separated from the body of the bone by intervening cartilage, it is called *Epiph'ysis*. The apophyses or processes are, at times, distinguished by epithets, expressive of their form: as *A. styloid*, *A. coracoid*, &c. Others are not preceded by the word apophysis; as *Trochanter*, *Tuberosity*, &c.

APOPH'YSIS OF INGRAS'SIAS is a term applied to the lesser ala of the sphenoid bone.

APOPHYSIS MAMMILLARIS, Mastoid process.

APOPHYSIS OF RAU, Grêle *apophyse du Mar-teau*: see Malleus.

APOPHYSIS ZYGOMATICA, Zygomatic process.

APOPIES'MA, from *αποπιεζω*, 'I compress.' Hippocrates uses the term to signify a fancied expression or forcing out of humours by the application of bandages in wounds and fractures.

APOPLANESIS, Error loci.

APOPLEC'TIC, *Apoplectic'us*. Referring to Apoplexy. This word has various significations. It is applied, 1. To individuals laboring under apoplexy: 2. To remedies proper for combating apoplexy: 3. To the constitution, temperament, or make, *Architectu'ra apoplectic'a*, *Hab'itus apoplectic'us* seu *quadra'tus* seu *turs'us*, which predisposes to it, and, 4. To the symptoms which characterize apoplexy; as *Apoplectic sleep*, *A.*

stroke, A. stertor, &c. The jugular veins have also, by some, been called *Apoplectic veins*, *Venæ apoplecticae*.

APOPLECTIC CELL. *Focus apoplecticus.* A cavity remaining in the encephalon, after the effusion of blood and its subsequent absorption.

APOPLECTICUS, Antiapoplectic, Apoplectic.

APOPLEXIA, *Apoplexy*.—*a. Catalepsia*, *Catalepsia*.—*a. Cerebralis*, see *Apoplexy*.—*a. Cerebri*, see *Apoplexy*.—*a. Cordis*, *Hæmocardiorrhagia*.—*a. Hepatica*, *Hepatorrhagia*.—*a. Hydrocephalica*, *Hydrocephalus internus*.—*a. Interarachnoidalis*, *Apoplexy*, *meningeal*.—*a. Intermeningealis*, *Apoplexy*, *meningeal*.—*a. Medullaris*, *Apoplexia myelitica*.—*a. Meningea*, *Apoplexy*, *meningeal*.

APOPLEXIA MYELITICA, *A. Medullaris* seu *Spinæ* seu *Rachidis*, *Hæmorrhachis*, *Myelorhagia*, *Myelapoplexia*, (F.) *Apoplexie de la Moëlle épinière*, *Hæmorrhagie de la Moëlle épinière*, *Hémato-myélie*, *Hémo-myélorrhagie*, *Hématorrhachie*. Hemorrhage into the spinal marrow.

APOPLEXIA NERVOSA, *Apoplexy*, *nervous*.—*a. Nervosa traumatica*, *Concussion of the brain*.—*a. Oculi* *Hæmophthalmia*.—*a. Pituitosa*, see *Apoplexy*.—*a. Pulmonalis*, see *Hæmoptysis*.—*a. Pulmonum*, see *Hæmoptysis*.—*a. Renalis*, *Apoplexy*, *renal*.—*a. Rachialis*, *A. myelitica*.—*a. Sanguinea*, see *Apoplexy*.—*a. Serosa*, see *Apoplexy*.—*a. Simplex*, *Apoplexy*, *nervous*.—*a. Spasmodica*, *Apoplexy*, *nervous*.—*a. Spinalis*, *Apoplexia myelitica*.—*a. Temulenta*, see *Temulentia*.

APOPLEXIE CAPILLAIRE, *Mollities cerebri*.—*a. Cérébrale*, *Apoplexy*, *Hæmorrhagie cérébrale*.—*a. du Cœur*, *Hæmocardiorrhagia*.

APOPLEXIE FOUROYANTE, 'Thundering Apoplexy.' A form of apoplexy, which is intense and rapidly fatal.

APOPLEXIE MENINGÉE, *Apoplexy*, *meningeal*.—*a. de la Moëlle Épinière*, *Apoplexy*, *spinal*.

APOPLEXY, *Apoplexia*, (Sc.) *Poplesy*, from ἀποπληξιν, 'to strike with violence.' At the present day, the term apoplexy is employed by many writers to signify interstitial hemorrhage, (F.) *Hæmorrhagie interstitielle*, or every effusion of blood, which occurs suddenly into the substance of an organ or tissue. Hence, we speak of cerebral apoplexy, pulmonary apoplexy, &c. &c. Formerly it was always—and still is by many—used in a restricted sense, to signify, in other words, the train of phenomena, which characterize cerebral apoplexy. This disease, *Hæmorrhagia Cerebri*, *Aphro'nia*, *Carus Apoplexia*, *Coma Apoplexia*, *Apoplexia cerebri sanguinea* seu *cerebra'lis*, *Encephalorrhagia*, *Sanguinis ietus*, *Hæmatencephalum*, *Pulplexia*, *Sideratio*, *Apilepsia*, *Morbus atton'itus*, *Gutta*, *Theophlegia*, *Theoplexia*, (F.) *Apoplexie*, *A. cérébrale*, *Hæmatencephalie*, *Comp de sang*, is characterized by diminution, or loss of sensation and mental manifestation; by the cessation, more or less complete, of motion; and by a comatose state,—circulation and respiration continuing. It generally consists in pressure upon the brain; either from turgescence of vessels, or from extravasation of blood: hence the terms *Hæmencephalus*, *Hæmorrhagie cérébrale*, and *Hæmoencephalorrhagie*, applied to it by some. The general prognosis is unfavourable; especially when it occurs after the age of 35. When Apoplexy is accompanied with a hard, full pulse, and flushed countenance, it is called *Apoplexia sanguinea*, *Cataphora Coma*; when with a feeble pulse and pale countenance, and evidences of serous effusion, *Apoplexia serosa*, *A. pituitosa*, *Serous Apoplexy*, *Cataphora hydrocephalica*, *Encephaloch'ysis semi'lis*, *Hydrocephalus acutus senum*, *Hydroencephalorrhée*,

(Piorry), *Hydropsie cérébrale suraiguë*, *Hydro-rhagie*.

In *Nervous Apoplexy*, *Apoplexia nervosa* seu *spasmodica*, *A. simplex*, *Simple apoplexy*, no lesion whatever may be perceptible on dissection; although the patient may have died under all the phenomena that are characteristic of apoplexy.

APOPLEXY, HEART, see *Coup-de-Soleil*.—*a. of the Heart*, *Hæmocardiorrhagia*.

APOPLEXY, MENINGEAL, *Apoplexia meningea* seu *intermeningea'lis* seu *interarachnoida'lis*, (F.) *Apoplexie méningée*, *Hæmorrhagie méningée*. Hemorrhage from the meninges of the brain or spinal marrow, generally into the great cavity of the arachnoid.

APOPLEXY, NERVOUS, see *Apoplexy*.—*a. Pulmonary*, see *Hæmoptysis*.—*a. Simple*, *A. Nervous*.

APOPLEXY, RENAL, *Apoplexia renal'is*. A condition of the kidney, characterized by knotty, irregular, tuberculated eminences, some of a deep black colour. Effusion of blood into the substance of the kidney.

APOPLEXY, SEROUS, see *Apoplexy*.—*a. Spinal*, *Apoplexia myelitica*.

APOPNEUSIS, *Exhalatio*.

APOPNIxis, *Suffocatio*.

APOPNOE, *Exspiratio*.

APOPNOEA, *Exspiratio*.

APOPSYCHIA, *Syncope*.

APOPTOSIS, from ἀποπίπτω, 'I fall down.' A relaxation of bandages.—*Erotian*.

APORRHŒ, *Aporrhœa*.

APORRHŒA, *Apor'rhœe*, *Apor'rhyis*, *Deflu-vium*, from ἀπορρῶ, 'I flow from.' An emanation, effluvia, contagion.—*Moschion*. A falling off of the hair, according to some.

APORRHYSIS, *Aporrhœa*.

APOSCEN'MA, *Aposcen'sis*, from ἀποσκειν, 'I lie down, I direct myself towards.' Afflux of fluids towards a part. Metastasis. The first word has been applied to the excrements.—*Hippocrates*, *Galen*.

APOSCENOSIS, *Apocenosus*.

APOSCENAPARNIS'MUS, *Deascia'tio*, from ἀπο and σκεπαρῶν, 'a hatchet.' Wound of the cranium, by a cutting instrument, in which a piece of the bone has been cut out, as with a hatchet.—*Gorræus*.

APOSCEPSIS, *Aposcemma*.

APOSCHASIS, *Aposchas'mus*, from ἀποσχάζω, 'I scarify.' *Scarification*. A slight superficial incision in the skin. Also, blood-letting.—*Ilippoc.*

APOSTA, *Sitis defectus*, from ἀ, privative, and ποσις, 'drink.' Want of thirst; absence of desire for liquids.

APOSITIA, from ἀπο, 'from,' and σιτος, 'food.' Aversion for food.—*Galen*. See *Disgust*.

APOSITIC, *Aposit'icus*; the same etymology. Any substance which destroys the appetite, or suspends hunger.

APOSAS'MA, from ἀποσπῶ, 'I tear or lacerate.' (F.) *Arrachement*. A solution of continuity, especially of a ligament; *Rhegma ligamenta're*, *Lacera'tio ligamenta'ria*.

APOSPHACEL'ISIS, *Aposphacelis'mus*, from ἀπο, and σφάκελος, 'mortification.' Gangrene in wounds and fractures, owing to the bandages being too tight.—*Hippocrates*.

APOSPHINX'IS, *ασπινγ'ις*, constriction, compression. The action of a tight bandage.—*Hippocrates*.

APOSPONGIS'MUS, the act of sponging for any purpose.—*Gorræus*.

APOSTALAG'MA, *Apostag'ma*, from ἀπο, 'from,' and σταλάζω, 'I drop.' The ancient name for the saccharine liquor which flows from grapes when not yet pressed.

APOSTASIS, from ἀπο, and ἵστημι, 'I stop.'

The ancients had different significations for this word. It was most commonly used for an abscess. The separation of a fragment of bone by fracture. Removal of disease by some excretion, &c.

APOSTAX'IS, from ἀποσῆζω 'I distil from.' *Stasis*. The defluxion of any humour, as of blood from the nose.—Hippocrates.

APOSTEMA, from ἀπο, 'from,' and ἵστημι, 'I settle,' or from ἀφίστημι, 'I recede from.' This word is used by the ancients somewhat vaguely. It meant an affection in which parts, previously in contact, are separated from each other by a fluid collected between them. The moderns regard it as synonymous with *Abscess*. Some, even of the moderns, have applied it to any watery tumour, and even to tumours in general.

APOSTEMA CEREBRI, Encephalopoyosis—a. Empyema, Empyema—a. Parulis, Parulis—a. Phalangum, *Fourche*—a. Psosaticum, Lumbar abscess.

APOSTEMACION, Abscess.

APOSTERIG'MA, from ἀποστρίζω, 'I support.' Anything that supports a diseased part, as a cushion, a pillow, &c.—Galen. A deep-seated and inveterate disease of the intestines.—Hippocrates.

APOS'THIA, *Leipoder'mia*, from α, privative, and ποσις, 'prepuce.' Want of prepuce.

APOSTHUME, Abscess.

APONTOLÉ, Extractum.

APOSTOLO'RUM UNGUENTUM, *Dodecapharmacum*, Ointment of the Apostles. So called, because as many solid ingredients entered into its composition as there were apostles. It contained several resins and gum-resins, yellow wax, oil, vinegar, verdigris, &c., and was formerly employed as a vulnerary.

APOSTROPHE, from ἀπο, and στρέφω, 'I turn.' An aversion or disgust for food.—Paulus. Also, the direction of humours towards other parts.

APOSYRMA, Abrasion, Desquamation.

APOTELES'MA, from ἀπο, and τέλεσμα, 'completion.' The result or termination of a disease. See, also, Amuletum.

APOTHANASIA, see Death.

APOTHE'CA, *Pharmace'um*, *Pharmacopol'ium*, from ἀπο, and τίθημι, 'to place.' Any place where things are kept, and therefore 'a shop,' and particularly a wine cellar. A place or vessel wherein medicines are kept. See *Pharmacopolium*.

APOTHECARIES' HALL. The Hall of the Corporation or Society of Apothecaries of London, where medicines are prepared and sold under their direction, &c. This Company obtained a charter of incorporation in the 15th year of James the First. No general practitioner can establish himself in England or Wales, without having obtained a license from the Court of Examiners of the Company.

APOTHECARY, *Apotheca'rius*, *Dispensator*, *Pharmacop'olla*, *Pharmacop'us*, *Pharmac'us*, *Pharmacut'us*, *Pimenta'rius*, *Rhizot'omus*, *Myrop'oles*, *Myrop'olus*, *Pharmacter*, *Pharmacur'gicus*, *Pharmacur'gus*, *Pharmacut'ist*, same derivation. (Prov.) *Pot'ecary*, (Sc.) *Pottingar*, (F.) *Apothicaire*, *Pharmacien*, *Pharmacopole*. In every country except Great Britain, it means one who sells drugs, makes up prescriptions, &c. In addition to these offices, which, indeed, they rarely exercise, except in the case of their own patients, the Apothecaries in England form a privileged class of practitioners—a kind of sub-physician. See Surgeon-apothecary.

APOTHERAPE'IA, *Apothērap'ia*, *Apothēra'gia*, from ἀποθεραπεύω, (ἀπο and θεραπεύω), 'I cure.' A perfect cure.—Hippoc. In the ancient Gymnastics, it meant the last part of the exercises:—the friction, inunction, and bathing, for

the purpose of obviating fatigue, or curing disease.—Galen, Gorræus.

APOTHERAPEUSIS, *Apothērapeia*.

APOTHER'EUMUM, from ἀπο, and θερμη, 'heat.' A pickle made of mustard, oil, and vinegar.—Galen.

APOTH'ESIS, from ἀποτίθημι, 'I replace.' The position proper to be given to a fractured limb, after reduction.

APOTHECAIRE, Apothecary.

APOTHECAIRERIE (F.), from αποθήκη, 'a warehouse, shop.' The same as *Apotheca*; also, a gallipot. See *Pharmacopolium*.

APOTILIM'MA, from ἀπο, and θλιβω, 'I press from.' Anciently, the dregs, and sometimes the expressed juice, *Succus expressus*, of plants.—Gorræus.

APOTHRAU'SIS, from αποθραω, 'I break.' Fracture of a bone, with spicula remaining. Extraction of a spiculum of bone.—Gorræus. Also, Abscission.

APOTILMOS, Evulsion.

APOT'OKOS, from ἀπο, and τικω, 'I bring forth.' An abortive foetus.—Hippocrates.

APOTOME, Amputation.

APOTOMIA, Amputation.

APOTROPÆUM, Amuletum.

APOTROPE, Aversion. Also, deviation—as of a limb—*Paratropé*.

APOXYSMUS, Abrasion.

APOZEM, Decoction.

APZESIS, Decoction.

APPARA'TUS, *Parascenz*, from ad and parare, 'to prepare.' This word signifies a collection of instruments, &c., for any operation whatever. (F.) *Appareil*.

In Surgery, it means the methodical arrangement of all the instruments and objects necessary for an operation or dressing. By extension, the French give the name *Appareil*, *Capas chirurgica*, to the case or drawers in which the apparatus is arranged.

Apparatus has likewise been applied to the different modes of operating for the stone. See *Lithotomy*.

In *Physiology*, *Apparatus*, (F.) *Appareil*, is applied to a collection of organs, all of which work towards the same end. A system of organs comprehends all those formed of a similar texture. An *apparatus* often comprehends organs of very different nature. In the former, there is analogy of structure; in the latter, analogy of function.

APPARATUS ALTUS, see *Lithotomy*.

APPARATUS IMMOV'ABLE (F.), *Appareil immobile*, *Immovable Bandage*, *Permanent Bandage*. An apparatus for fractures, which is generally formed by wetting the bandages in some substance, as starch or dextrin, which becomes solid, and retains the parts *in situ*.

APPARATUS LATERALIS, see *Lithotomy*—a. Major, see *Lithotomy*—a. Minor, see *Lithotomy*.

APPAREIL, *Apparatus*, *Boîtier*—a. *Diapno'gène*, see *Perspiration*—a. *Grand*, see *Lithotomy*—a. *Haut*, see *Lithotomy*—a. *Immobile*, *Apparatus*, *immovable*—a. *Lacrymal*, see *Lachrymal passages*—a. *Lateralisé*, see *Lithotomy*—a. *Névrothèle*, see *Névrothèle*—a. *Petit*, see *Lithotomy*—a. *Pigmental*, *Pigmental apparatus*.

APPAREILS DE FORMATION (F.). Gall admits, in the brain, two kinds of fibres; the one, divergent, proceeding from the cerebral peduncles to the convolutions, and constituting what he calls *appareils de formation*: the other, convergent, and proceeding from the convolutions to the centre of the organ, constituting what he calls *appareils de réunion*. The first, as a whole, form the organs of the mental faculties: the latter are commissures, which unite parts of the organ that are double and in pairs.

APPAUVRI, Impoverished.

APPENDICE, Appendix — *a. Cæcal*, Appendix vermiformis cæci — *a. Digital*, Appendix vermiformis cæci — *a. Sous-sternale*, Xiphoid cartilage — *a. Sus-sphénoïdale du cerveau*, Pituitary gland — *a. Xiphoidé*, Xiphoid cartilage.

APPENDICES COLI ADIPOSEÆ, Appendiculæ epiploicæ — *a. Épiploïques*, Appendiculæ epiploicæ. APPENDICULA CEREBRI, Pituitary gland — *a. Vermiformis cæci*, see Appendix — *a. Epiploica*, Epiploic appendage.

APPENDICULÆ PINGUEDINOSÆ, *A. Epiploica*, *Epiploicæ appendices*, *Appendices colidi adiposæ*, *Fimbriæ carnosæ colidi*, *Supplementa epiploica*, *Omentula*, (F.) *Appendices Épiploïques*. Prolongations of the peritoneum beyond the surface of the great intestine, which are analogous in texture and arrangement to omenta.

APPENDIX, *Epiph'ysis*, from *appendere*, (ad and *pendere*, 'to hang,') 'to hang from.' Any part that adheres to an organ or is continuous with it:—seeming as if added to it. An *appendage*; an apophysis, (F.) *Appendice*, *Anexæ*.

APPENDIX AURICULÆ, see Auricles of the Heart — *a. Cerebri*, Pituitary gland — *a. ad Cerebrum*, Cerebellum — *a. Cutanea Septi Narium*, *Statica Septi Narium* — *a. to the Epididymis*, *Vasculum aberrans* — *a. Ventriculi*, Duodenum.

APPENDIX VERMIFORMIS, *Appendicula Vermiformis Cæci*, *Tubus Vermicularis Cæci*, *Ectophyas*, *Additamentum Coli*, *Appendix Cæci*, (F.) *Appendice vermiforme*, *A. cæcal* or *digital*. A vermicular process, the size of a goose-quill, which hangs from the intestine cæcum. Its functions are unknown.

APPENDIX VESICÆ, see Bladder, sacculated.

APPENSIO, see Analeptia.

APPETENCE, *Appeten'tia*, from *appetere*, (ad and *petere*,) 'to desire.' An ardent, passionate desire for any object.

APPETITE, *PERTE D'*, Anorexia.

APPETITE, *Appeti'tus*, *Appeten'tia*, *Appeti'tia*, (ad and *petere*,) 'to seek,' *Cupi'do*, *Orex'is*, *Ormê*: same etymology as the last. An internal sensation, which warns us of the necessity of exerting certain functions, especially those of digestion and generation. In the latter case it is called *venereal appetite*, (F.) *Appetit vénérien*: in the former, simply *appetite*, (F.) *Appetit* ou *Appétition*. If the desire for food, occasioned by a real want, be carried to a certain extent, it is called *hunger*, when solid food is concerned; *thirst*, when liquid. *Appetite* and *hunger* ought not, however, to be employed synonymously: they are different degrees of the same want. Hunger is an imperious desire: it cannot be provoked, like the appetite. It is always allayed by eating: but not so the appetite; for, at times, it may be excited in this manner. They are very generally, however, used synonymously.

APPETITE, MORBID, *Limosis*.

APPETITE, VENE'REAL, Venereal desire, (F.) *Le génésique*, *Amour physique*, *Sens génital*. The instinctive feeling that attracts the sexes towards each other to effect the work of reproduction.

APPETITUS CANINUS, *Boulimia* — *a. Deficiens*, *Dysorexia*.

APPLE, ADAM'S, *Pomum Adami* — *a. Bitter*, *Cucumis colocynthis* — *a. Curassoa*, *Aurantium curassaventium* — *a. Dead Sea*, see *Quercus infectoria* — *a. Eye*, see *Melon* — *a. of the Eye*, *Pupil* — *a. Mad*, see *Quercus infectoria* — *a. May*, *Podophyllum peltatum* — *a. of Peru*, *Datura stramonium* — *a. Root*, *Euphorbia corollata* — *a. of Sodom*, see *Quercus infectoria*.

APPLE TEA, *Apple water*. Slice two large, not over-ripe apples, and pour over a pint of boiling

water. After an hour, pour off the fluid, and, if necessary, sweeten with sugar.

APPLE TREE, *Pyrus malus*.

APPLICATA, from *applicare*, (ad and *plicare*, 'to fold,') 'to apply.' A word, unnecessarily introduced into medical language, to express the objects which are applied immediately to the surface of the body, as clothes, cosmetics, baths, &c. — Hallé.

APPLICA'TION, *Applica'tio*, (same etymon,) in a moral signification, is synonymous with attention. Also, the act of applying one thing to another; as the application of an apparatus, of a bandage, blister, &c.

APPREHENSIO, from *ad* and *prehendere*, 'to take.' This word is employed in various senses. It means catlepsy or catoche. — Paul Zacchias. A kind of bandage for securing any part. Also, a therapeutical indication.

APPROCHE, Coition.

APPROXIMA'TION, *Approxima'tio*, from *ad* and *proximus*, 'nearest.' Ettmüller gave this name to a pretended method of curing disease, by making it pass from man into some animal or vegetable, by the aid of immediate contact.

APRAC'TA, from *a. priv.*, and *πρασσω*, 'I act.' Without action. An epithet for the parts of generation, when unfit for copulation or generation.

APRICATIO, Insolation.

APRICOT, *Prunus Armeniaca*.

APROCTUS, see Atretus.

APRONIA, *Tamus communis*.

APROSOP'IA, *Tricephalia*, from *a. priv.*, and *προσωπον*, 'the face.' A malformation, which consists in the face being deficient.

APROSOPUS, *Microprosopus*.

APSINTHIA'TUM, from *αψινθιον*, 'wormwood.' A sort of drink made of wormwood. — Aëtius.

APSINTHITES, Absinthites.

APSYCHIA, Syncope.

APSYCHISME, Idiotism.

APSYCHY, Syncope.

APSYXIA, Syncope.

APTHÆ, Aphthæ.

APTYS'TOS, from *a. priv.*, and *πτωω*, 'I spit.' Devoid of expectoration. An epithet given to certain pleurisies, in which there is no expectoration. — Hippocrates.

APUS, see Apodia.

APY'ETOS, from *a. priv.*, and *πυνω*, 'pus.' An external affection, which does not end in suppuration.

APYIQUE, *Apyros*.

A'PYQOS, from *a. priv.*, and *πυνω*, 'pus.' (F.) *Apyique*. That which does not afford pus.

APYRECTIC, *Apyretic*.

APYREMELE, *Apyromele*.

APYRETIC, *Apyret'icus*, *Apyrect'ic*, *Apyrect'icus*, *Apyr'etus*, from *a. priv.*, and *πυρ*, 'fire, fever.' Without fever. This epithet is given to days in which there is no paroxysm of a disease, as in the case of an intermittent, as well as to some local affections which do not induce fever. Urticaria is sometimes called an *aporetic exanthem*.

APYREX'IA, *A'pyrezy*. The same etymology. Absence of fever; *Dialen'ma*, *Dialeip'sia*, *Dinlip'sis*, *Tempus intercalare*, *Interval'lum*, *Intermissio*. *Apyrexia* is the condition of an intermittent fever between the paroxysms: the duration of the *appyrexia* consequently depends on the type of the intermittent. Occasionally, the term has been applied to the cessation of the febrile condition in acute diseases.

APYROMELE, *Apyrome'le*, from *a. priv.*, *πυρ*, 'a nut,' and *μύλη*, 'a sound.' A sound or probe, without a button or nut. It is the *Melo'tis*,

Specillum auricularium or *Auricular sound* of Galen.

AQUA, Urine, Water.—a. Acidi carbonici, Acidulous water.—a. Acidula hydrosulphurata, Naples water (factitious)—a. Aeris fixi, Acidulous water (simple)—a. Alkalina oxymuriatica, *Eau de Javelle*—a. Aluminis composita, Liqueur aluminis compositus.—a. Aluminosa Bateana, Liqueur aluminis compositus.—a. Ammoniae, Liqueur ammoniae.—a. Acetatis ammoniae, Liqueur ammoniae acetatis.—a. Ammoniae carbonatis, Liqueur ammoniae subcarbonatis.—a. Ammoniae caustica, Liqueur ammoniae.—a. Amnii, Liqueur Amnii.

AQUA AMYGDALARUM CONCENTRATA, A. amygdalarum amara-rum, (F.) Eau d'Amandes amères, Water of bitter almonds. Made by bruising well two pounds of bitter almonds; adding, whilst triturating, ten pounds of spring water, and four pounds of alcohol; letting the mixture rest in a well-closed vessel, and then distilling two pounds. Used instead of the Aqua Laurocerasi, and the Hydrocyanic acid. It must be given with great caution.

An *Aqua Amygdalæ amaræ, Bitter Almond water*, has been introduced into the last edition of the Ph. U. S., 1851, (*Ol. amygdal. amar. m. xvj; Magnes. Carbon. ʒj; Aqua Oij.*) Dose, f ʒss.

AQUA ANETHI, see *Anethum graveolens*—a. Anisi, see *Pimpinella anisum*—a. Anisi Fortis, Spiritus anisi.—a. Aquisgranensis, see *Aix-la-Chapelle*—a. Auditoria, Cotunnus, Liqueur of— a. Aurantii, see *Citrus aurantium*—a. Azotica oxygenata, Aqua nitrogenii protoxydi.—a. Balsamica arterialis, Aqua Binelli.—a. Bareginensis, Baréges water.—a. Barytæ Muriatæ, see *Baryta, muriate of*—a. Bellilucana, Balaruc waters.—a. Benedicta, Liqueur calcis.—a. Benedicta composita, Liqueur calcis compositus.—a. Benedicta Rulandi, Vinum antimonii tartarizati.

AQUA BINELLI, Aqua Binelli, Aqua Balsamica arterialis, Binelli's styptic, (F.) Eau de Binelli. A celebrated Italian hæmostatic, invented by one Binelli. Its composition is unknown, but its virtues have been ascribed to creasote; although there is reason for believing it to possess no more activity than cold water. Aqua Monterossi, (F.) *Eau de Monterossi*, appears to be of analogous composition.

AQUA BORYONENSIS, Bourbonne-les-Bains, mineral waters of—a. Bristolensis, Bristol water.

AQUA BROCCHEARI, Aqua Broccchieri, Broccchieri or Broccchiar water, Broccchieri's styptic, (F.) Eau de Broccchieri, Eau styptique de Broccchieri. A supposed styptic, which made much noise at Paris at one time. It is devoid of efficacy. Dr. Paris found nothing in it but water perfumed by some vegetable essence.

AQUA CALCARIÆ USTÆ, Liqueur calcis.—a. Calcis, Liqueur Calcis.—a. Calcis composita, Liqueur calcis compositus.—a. Camphoræ, Mistura camphoræ.—a. Camphorata, Bates's, see *Cupri sulphas*—a. Carbonatis sodæ acidula, Acidulous water, simple.—a. Carui, see *Carum Carui*—a. Cassiæ, see *Laurus cassia*—a. Catapultarum, *Arquebuseade, eau d'*—a. Chlorini, see *Chlorine*.

AQUA CINNAMOMI, Cinnamon water, (Sc.) Cannel water, Distilled water of Cinnamon Bark. Prepared also in the following manner. *Ol. Cinnam. f ʒss; Magnes. Carbon. ʒj; Aq. destillat. Oij.* Rub the oil and carbonate of magnesia; add the water gradually, and filter. (Ph. U. S.)

AQUA CINNAMOMI FORTIS, Spiritus Cinnamomi—a. Colcestrensis, Colchester, mineral waters of.

AQUA COLORATA, 'coloured water.' A name given to a prescription in which simple coloured water is contained. Used in hospital cases, more especially, where a *placebo* is demanded.

AQUA CUPRI AMMONIATA, Liqueur c. a.—a. Cu-

pri vitriolati composita, Liqueur cupri sulphatis composita.—a. inter Cutem, Anasarca.—a. Destillata, Water, distilled.

AQUA FABRO'RUM, 'Water of Smiths.' Water in which hot iron has been quenched. A feeble chalybeate.

AQUA FLORUM AURANTII, see *Citrus aurantium*—a. Fluvialis, Water, river.

AQUA FENIC'ULI, Fennel water. The distilled water of fennel seed. It may be prepared also like the aqua cinnamomi.

AQUA FONTANA, Water, spring—a. Fortis, Nitric acid.—a. Goulardi, Liqueur Plumbi subacetatis dilutus.—a. Hepatica, Hydrosulphuretted water.—a. Hordeata, Decoctum hordei.—a. Imbrium, Water, rain.—a. Infernalis, see *Toddy*—a. Intercus, Anasarca.—a. Inter Cutem, Anasarca.—a. Juniperi composita, Spiritus juniperi compositus.—a. Kali, Liqueur potassæ subcarbonatis.—a. Kali caustici, Liqueur potassæ.—a. Kali præparati, Liqueur potassæ subcarbonatis.—a. Kali puri, Liqueur potassæ.—a. Kali subcarbonatis, Liqueur potassæ subcarbonatis.—a. Labyrinthi, Cotunnus, liqueur of—a. Lactis, Serum lactis.—a. ex Lacu, Water, lake.—a. Lauro-cerasi, see *Prunus Lauro-cerasus*—a. Lithargyri acetati composita, Liqueur plumbi subacetatis dilutus.—a. Lucis, Spiritus ammoniæ succinatus.—a. Marina, Water, sea.—a. Medicata, Water, mineral.

AQUA MENTHÆ PIPERITÆ, Peppermint Water. The distilled water of peppermint. It may be prepared like the aqua cinnamomi.

AQUA MENTHÆ PIPERITIDIS SPIRITUOSA, Spiritus menthæ piperitæ—a. Menthæ pulegii, see *Mentha pulegium*—a. Menthæ viridis, Spearmint water, see *Aqua menthæ piperitæ*—a. Menthæ vulgaris spirituosa, Spiritus menthæ viridis.—a. Mineralis, Water, mineral.—a. Mirabilis, Spiritus pimentæ.—a. Monterossi, see *Aqua Binelli*—a. Mulsæ, Hydromeli.—a. Natri Oxymuriatici, Liqueur sodæ chlorinatæ.—a. Neapolitana, Naples water, (factitious)—a. Nephritica, Spiritus myristica.

AQUA NITROGENII PROTOXYDI, Protoxide of Nitrogen Water, Aqua azotica oxygenata, Searle's patent oxygenous aerated water. A patent solution of protoxide of nitrogen, said to contain five times its own bulk of gas. It has been recommended as a nervine, and excitant in nervous conditions, dyspepsia, &c. It has also been used in cholera, and to counteract the evil consequences of drunkenness. The dose is f ʒvj, or ʒviii, two or three times a day; or, in dyspepsia, as a beverage between meals.

AQUA NIVATA, Water, snow—a. Nucis moschatæ, Spiritus myristicæ.—a. Ophthalmica, Liqueur zinci sulphatis cum camphorâ.—a. Paludosa, Water, marsh.—a. Pedom, Urine.—a. Pericardii, see *Pericardium*—a. Phagedenica, see *Hydrargyri Oxymurias*—a. Piceæ, see *Pinus sylvestris*—a. Picis, see *Pinus sylvestris*—a. Pimentæ, see *Myrtus Pimenta*—a. Plumbi, Liqueur Plumbi, subacetatis dilutus.—a. Pluvialis, Water, rain.—a. Potassæ, Liqueur potassæ.—a. Pulegii, see *Mentha pulegium*—a. Pulegii spirituosa, Spiritus pulegii.—a. Putealis, Water, well.—a. ex Puteo, Water, well.—a. Rabelli, Elixir acidum Halleri.—a. Raphani composita, Spiritus armoraciæ compositus.—a. Regia, Nitromuriatic acid.

AQUA ROSÆ, Rose Water, Rhodostag'ma, (Ros. centifol. lbvij.) : Aquæ cong. ij. M. Distil a gallon—Ph. U. S.)

AQUA SALUBRIS, Water, mineral—a. Sâmbuci, see *Sambucus*—a. Sappharina, Liqueur cupri ammoniata.—a. Saturni, Liqueur plumbi subacetatis dilutus.—a. Sclopetaria, *Arquebuseade eau d'*—a. Seminum anisi composita, Spiritus anisi.—a. Seminum carui fortis, Spiritus carui.—a. Sodæ efferves-cens, Acidulous water, simple.—a. Soteria, Water,

mineral.—a. Stygia, Nitro-muriatic acid.—a. Styp-tica, Liquor cupri sulphatis composita.—a. Sulphurata simplex, Hydrosulphuretted water.—a. Sulphureti ammoniæ, Liquor fumans Boylei.—a. Thediana, *Arquebusade eau d'*—a. Theriacalis Bezoardica, Chylostagma diaphoreticum Minde-ri.—a. Tofana, Liquor arsenicalis.—a. Tosti panis, Toast water.—a. Traumatica Thedenii, *Arquebusade eau d'*—a. Vegeto-mineralis, Liquor plumbi subacetatis dilutus.—a. Vicienis, Vichy water.—a. Vitriolica camphorata, Liquor zinci sulphatis cum camphorâ.—a. Vitriolica cærulea, Solutio sulphatis cupri composita.—a. Vulneraria, *Arquebusade eau d'*—a. Zinci vitriolati cum camphorâ, Liquor zinci sulphatis cum camphorâ.

AQUE ACIDULÆ, Acidulous waters.—a. Badigæ, Bath, Mineral waters of.—a. Badizæ, Bath, Mineral waters of.—a. Bathoniæ, Bath, Mineral waters of.—a. Buxtoniensis, Buxton, Mineral waters of.—a. Calidæ, *Aigues chaudes*—a. Cantuarienses, Canterbury, waters of.—a. Chalybeate, Waters, mineral, chalybeate.

AQUE DESTILLATÆ, *Hydrolat'a*, Distilled Waters, (F.) *Hydrolate*. These are made by putting vegetable substances, as roses, mint, pennyroyal, &c., into a still with water, and drawing off as much as is found to possess the aromatic properties of the plant. To every gallon of the distilled water, 5 oz. of spirit should be added to preserve it. The simple distilled waters are sometimes called *Aquæ stillatit'ia simplices*: the *spirituous*, *Aquæ stillatit'ia spirituosæ*, but more commonly *Spirit'ia*.

AQUE FERROSÆ, Waters, mineral, chalybeate.

AQUE MARTIALES, Waters, mineral, chalybeate.

AQUE MEDICATÆ, Medicated waters include, in the Pharmacopœia of the United States, preparations consisting of waters impregnated with some medicinal substance, which are not arranged in any other class. Among these are the "Waters," and "Distilled waters" of the British pharmacopœias.

AQUE METUS, Hydrophobia.—a. *Minerales acide*—a. Waters, mineral, gaseous.—a. *Minerales ferruginosæ*, Waters, mineral, chalybeate.—a. *Minerales sulphuræ*, Waters, mineral, sulphureous.—a. *Stillatitiæ*, *Aquæ destillatæ*—a. *Solis*, Bath, mineral waters of.

AQUEDUCTUS, *Aqueduct*, from *aqua*, 'water,' and *ducere*, *ductum*, 'to lead.' (F.) *Aqueduc*. Properly, a canal for conducting water from one place to another. Anatomists have used it to designate certain canals.

AQUEDUCTUS CEREBRI, Infundibulum of the brain.—a. Cotunnii, Aqueductus vestibuli.

AQUEDUCTUS COCHLEÆ, (F.) *Aqueduc du Limaçon*:—a very narrow canal which proceeds from the tympanic scala of the cochlea to the posterior edge of the *pars petrosa*.

AQUEDUCTUS FALLOPII, *Canalis Fallopii*, *Canal spirôide de l'os temporal* of Chaussier, (F.) *Aqueduc de Fallope*. A canal in the *pars petrosa* of the temporal bone, which extends from the meatus auditorius internus to the foramen stylo-mastoideum, and gives passage to the facial nerve. The opening into this aqueduct is called *Hiat'us Fallopii*, *H. Canalis Fallopii*, *Fissu'ra Canalis Fallopii*, *Foramen Tari'ni*, *F. anony'mum Ferrein'ii*.

AQUEDUCTUS SYLVII, *Canalis eminentiæ quadrigeminæ*, *Iter ad quartum ventricul'um seu à tertio ad quartum ventricul'um*, *Canalis medius*, (F.) *Aqueduc de Sylvius*, *Canal intermédiaire des ventricules* of Chaussier. A canal forming a communication between the third and fourth ventricles of the brain. The ventricular opening is termed *anus cerebri*.

AQUEDUCTUS VESTIBULI, *Aqueductus Cotun'rii*, *Canal of Cotunn'ius*, (F.) *Aqueduc du vestibule* ou *Aqueduc de Cotugno*. This begins in the ves-

tibule, near the common orifice of the two semi-circular canals, and opens at the posterior surface of the *pars petrosa*.

AQUALICULUS, from *aqualis*, 'a water-pot.' That part of the abdomen which extends from the umbilicus to the pubes. See Hypogastrium. It has also been applied to the stomach or intestinal canal.

AQUASTER. A word used, by Paracelsus, to express the visions or hallucinations of patients.

AQUEDUC, Aqueduct.—a. *de Cotugno*, Aqueductus vestibuli.—a. *de Fallope*, Aqueductus Fallopii.—a. *du Limaçon*, Aqueductus cochleæ.—a. *de Sylvius*, Aqueductus Sylvii.—a. *du Vestibule*, Aqueductus vestibuli.

AQUEDUCT, Aqueductus.

A'QUEOUS, *A'queus*, *Aquo'sus*, *Hydato's*, *Hydro's*, from *aqua*, 'water,' (F.) *Aqueur*, Watery. The absorbents or lymphatics are sometimes called, in France, *Conduits* ou *Canaux aqueux*.

AQUEOUS HUMOUR OF THE EYE, *Humor aquo'sus*, *Oœi'des*, *Oo'des*, *Hydato'i'des*, *Hydato's*, *Ovatus* seu *Oviformis* humour, *Albugin'eus* humour, (F.) *Humeur aqueuse*. The limpid fluid which fills the two chambers of the eye, from the cornea to the crystalline, and which is, consequently, in contact with the two surfaces of the iris. Quantity, 5 or 6 grains: s. g. 1.0003. It contains albumen, chloride of sodium, and phosphate of lime in small quantity; and is enveloped in a fine membrane:—the membrane of the aqueous humour, *Tunica propria* seu *Vagi'na* seu *Membrana* seu *Cap'sula humo'ris a'quei* seu *Membrana* *Demuria* seu *Descemet'ii*, *Membrane of Demours* or of *Descemet*; although these last terms are by some appropriated to a third layer of the cornea.

AQUEUS, Aqueous.

AQUIDUCA, Hydragogues.

AQUIFOLIUM, *Ilex aquifolium*—a. *Folius deciduis*, *Prinos*.

AQUILA, Hydrargyri submuriæ, Sulphur.

The alchymists used this word for sublimed sal ammoniac, precipitated mercury, arsenic, sulphur, and the philosopher's stone. See Hydrargyri Submuriæ, and Sulphur.

AQUILA CÆLESTIS; a sort of panacea, of which mercury was a constituent.

AQUILA LACHRYMÆ; a liquor prepared from several ingredients, especially from calomel.

AQUILA PHILOSOPHORUM. The alchymists, whose terms were always mysterious, called mercury thus, when reduced to its original form.

AQUILA VENERIS; an ancient preparation, made by subliming verdigris and sal ammoniac.

AQUILÆ VENÆ, Temporal veins.

AQUILEGIA, *A. vulg'ris* seu *sylvestris* seu *Alpi'na*, Common Columbine or Columbine, (F.) *Ancolie*. Order, Ranunculacææ. From *aquila*, 'the eagle,' owing to some fancied resemblance of the spurs to talons. The seeds, herb, and flowers were formerly used in jaundice and cutaneous diseases. They are still retained in many of the Pharmacopœias of continental Europe.

AQUILEGIA ALPINA, *Aquilegia*.

AQUILEGIA CANADENSIS, *Wild Columbine*, is indigenous, and flowers in April and June. The seeds are said to be tonic.

AQUILEGIA SYLVESTIS, *Aquilegia*—a. *Vulgaris*, *Aquilegia*.

AQUILIC'IA SAMBU'CINA, from *aqua*, 'water,' and *clucire*, 'to attract,' because it grows in moist places. [?] An East Indian plant, the decoction of whose root is used in indigestion to allay heartburn.

AQUIPAROUS, (F.) *Aquipare*, from *aqua*, 'water,' and *pario*, 'I bring forth.' An epithet

for glands which, like the parotid, secrete much water, in contradistinction to the submaxillary glands, which are muciparous.

AQUO-CAPSULITIS, Aquo-membranitis.

AQUO-MEMBRANITIS, *Keratôir'itis*, *Aquo-capsul'itis*, *Hydromening'itis*. Inflammation of the anterior chamber of the eye. A badly compounded term, denoting inflammation of the capsule or membrane of the aqueous humour.

AQUULA, Ceratocela, Hydatid, Hydroa—a. Acustica, Cotunnus, liquor of.

AQUULA seu AQUA MORGAGNI. The minute portion of water which escapes when an opening is made into the capsule of the crystalline.

AQUULA VITREA AUDITIVA, *Vitrine auditivæ*.

ARA PARVA, a small altar;—a kind of bandage invented by Sostratus, which represents the corners of an altar.—Galen.

AR'ABE; a wound, a blow.—Erolian.

ARABICA ANTIDOTUS HEPATICA, *Arabic Hepat'ic Antidote*. A powder composed of myrrh, costus, white pepper, &c. It was administered in new wine.

ARABICUS LAPIS. A sort of white marble, analogous to alabaster, found in Arabia. It was regarded as absorbent and desiccative, and was employed in hemorrhoids.

ARABIS BARBAREA, *Erysimum barbarea*.

ARABIS MALAG'MA. An antiscrofulous medicine, composed of myrrh, olibanum, wax, sal ammoniac, iron pyrites, &c.—Celsus.

ARABISTS, see Arabs, medicine of the.

AR'ABS, MEDICINE OF THE. The Arabians kept the torch of medical science illuminated during a dark period of the middle ages. Before the year of the Hegira, they had schools of medicine; but these were most flourishing during the 10th, 11th, and 12th centuries. The chief additions made by them to medical science were in the departments of pharmacy and in the description of diseases. Their principal writers were Avicenna, Serapion, Averrhoes, Hali Abbas, Moses Maimonides, Avenzoar, Rhazes, Albucasis, &c. The disciples of the Arab school were called *Ar'abists*.

ARACACHA, *Conium mosechatum*.

ARACHIS AFRICANA, *A. hypogea*—a. Americana, *A. Hypogea*.

AR'ACHIS HYPOGE'A, *A. America'na* seu *Africa'na*, *Arachni'da hypogea*, Ground nut, Pea nut, Earth almond, (S.) *Mane*; erroneously called *Pistachio nut*, in the South; *Pindars* of the West Indies. *Ord.* Leguminosæ. Cultivated in the Southern States. The seeds are oily, and are eaten. A kind of inferior chocolate may be made of them.

ARACH'NE, *apayn*, 'a spider,' 'a cobweb.' Hence, *Arachnitis*, &c.

ARACHNIDA HYPOGEA, *Arachis hypogea*.

ARACHNI'TIS, more properly *Arachnoid'itis*, *Arachnoid'itis*, *Inflammation of the Arachnoid*. A variety of phrenitis.

ARACHNODETIS, *Arachnitis*.

ARACH'NOID, *Arachnoideus*, *Arachno'des*, from *apayn*, 'a cobweb,' and *idos*, 'form, resemblance.' Resembling a spider's web.

ARACHNOID CANAL, see Canal, arachnoid.

ARACHNOID OF THE EYE. The lining membrane of a cavity, supposed by some to exist between the sclerotic and choroid.

ARACH'NOID MEMBRANE, *Meninx media* seu *serosa*, *Tunica aranea* seu *crystallina*, *Menin'gion*. A name given to several membranes, which, by their extreme thinness, resemble spider-webs. Celsus and Galen called thus the membrane of the vitreous humour,—the *tunica hyaloidea*. The moderns use it now for one of the membranes of the brain, situate between the

dura mater and pia mater. It is a serous membrane, and composed of two layers; the external being confounded, in the greater part of its extent, with the dura mater, and, like it, lining the interior of the cranium and spinal canal; the other being extended over the brain, from which it is separated by the pia mater, without passing into the sinuosities between the convolutions, and penetrating into the interior of the brain by an opening at its posterior part under the corpus callosum. It forms a part of the investing sheath of the nerves, as they pass from the encephalic cavities. Its chief uses seem to be:—to envelop, and, in some measure, protect the brain, and to secrete a fluid for the purpose of keeping it in a state best adapted for the proper performance of its functions.

ARACHNOIDITIS, *Arachnitis*.

ARACK', *Arrack*; (East Indian.) A spirituous liquor made in India in various ways, often from rice, sometimes from sugar fermented along with the juice of the cocoa nut; frequently from toddy, the juice which flows from the cocoa-nut tree by incision, and from other substances. It is a strong, heating spirit.

ARACK, Mock, is made by adding $\frac{1}{2}$ ij of Benzoic acid to a quart of rum. The celebrated Vauxhall punch is made with such arack.

ARACOUCHINI, *Icica aracouchini*.

ARACUS AROMATICUS, *Vanilla*.

AR'ADOS, *Apados*. The agitation excited in the stomach by the coction of aliments of different nature.—Hippocrates. Likewise, the motion produced by cathartics.

ARÆOMA, *Interstices*.

ARÆOMETER, *Areometer*.

ARÆOTICA, from *apaous*, 'I rarefy.' Medicines supposed to have the quality of rarefying the humours. See *Rarefaciens*.

ARAKI, see Spirit.

ARALIA CANADENSIS, *Panax quinquefolium*.

AR'ALIA HISPIDA, *Dwarf Elder*, *Ord.* *Araliaceæ*, is said to be diuretic, and has been recommended, in decoction, in dropsy.

AR'ALIA NUDICAULIS, *Nardus America'nus*, *Small Spikenard*, *Wild Liquorice*, *Sweet root*, *False* or *Wild Sassaaparilla*, (F.) *Petit nard*. This American plant is said to be a mild stimulant and diaphoretic, and has been recommended as a substitute for sassaaparilla. It is used, also, as a tonic. It is in the secondary list of the Pharmacopœia of the United States.

AR'ALIA RACEMOSA, *American Spikenard*, has the same properties as *A. Nudicaulis*.

AR'ALIA SPINOSA, *Angelica Tree*, *Prickly Ash*, *Toothach Tree*, *Spikenard Tree*, *Prickly Elder*, *Shotbush*, *Pigeon Tree*. Its properties are not clear. The berries, and a tincture of them, have been employed, it is said, successfully in toothach. A spirituous infusion has also been used in colic. The bark is official in the Pharmacopœia of the United States.

ARANEÆ, *Araneæ* Tela—a. *Tarentula*, see *Tarentula*.

AR'ANEÆ TELA, *Aranea*, *Araneum*, *Cobweb*, (F.) *Toile d'Araignée*. Formerly, this substance was much employed, and supposed to possess extraordinary virtues, especially when applied to the wrists. It has been recently used again in intermittents. The spider itself, softened into a plaster and applied to the forehead and temples, is said by Dioscorides to prevent ague. Cobweb is a mechanical styptic, and is so applied at times.

ARANEOSA URINA. A term applied to the urine when loaded with filaments, like cobwebs.

ARANEOSUS (PULSUS); a term employed

to express extreme weakness of pulse; when the movements resemble those of a delicate net raised by the wind.

ARANEUM, *Aranee Telo*.

ARANEUM ULCUS, *Astakillo*. A name given by Paracelsus to a malignant, gangrenous ulcer, extending from the feet to the legs.

ARARA, *Myrobalmus citrina*.

ARASCON, *Nymphomani*, *Satyriasis*.

ARATRUM, *Vomer*.

ARAUCHARIA DOMBEYI, *Dombeya excelsa*.

ARBOISE, *Arbutus unedo*.

ARBOL DE LECHE, *Galactodendron utile*.

ARBOR BENIVI, Benjamin—a. *Indica*, *Laurus cassia*—a. *Maris*, *Coral*—a. *Thurifera*, *Juniperus Lycia*—a. *Uteri Vivificans*, *Palmae uteri plicate*.

ARBOR VITÆ, (F.) *Arbre de vie*. A name given to an arborescent appearance, observed on cutting the cerebellum longitudinally; and which results from the particular arrangement of the white substance with the cineritious. Also, the *Thuya occidentalis*.

ARBOR VITÆ, AMERICAN, *Thuya occidentalis*—a. *Vitæ Uterinus*, *Palmae uteri plicate*—a. *Vitæ of the Uterus*, *Palmae uteri plicate*.

ARBORES. A morbid alteration of the skin, which precedes its ulceration. Ruland.

ARBORIZATION, *Arborisatio*, (F.) *Arborisation*, from *arbor*, *arboris*, 'a tree.' The figure or appearance of a tree or plant. The capillary vessels, when injected, as in inflammation, frequently appear under the form of Arborizations.

ARBOUSIER, *Arbutus unedo*.

ARBRE DE VIE, *Arbor Vitæ*.

ARBUSCULA GUMMIFERA BRAZILIENSIS, *Hypericum bacciferum*.

ARBUTUS, A. *Unedo*—a. *Trailing*, A. *Uva ursi*, *Epigæa repens*.

ARBUTUS UVA URSI, *Arctostaphylos Uva ursi*, *Mairan'nia uva ursi*. Ord. *Ericaceæ*. *Sex Syst.* Decandria Monogynia. (F.) *Busserolle* ou *Raisin d'Ours*. The leaves—(*Uva Ursi*, Ph. U. S.)—of this plant are tonic and astringent, and have been employed, chiefly, in diseases of the urinary organs, and also as a parturifacient. Dose of the powder from gr. xv to ʒss. The English names are *Trailing Arbutus*, *Bear's Whortleberry* or *Bearberry*, *Mountain-box*, *Redberry*, *Upland Cranberry*, *Foxberry*, *Checkerberry*, (Sc.) *Brawlin*.

ARBUTUS UNE'DO, *Arbutus*, *Andrach'nē*, *Une'do*, *U. papyracea*, *κορπος*, (F.) *Arbousier*, *Arboise*. A decoction of the leaves is astringent, and has been used in diarrhœa.

ARC, *Arch*, *Arcus*, (F.) *Arc*, *Arcade* (diminutive). Any part of the body resembling an arch in form; as the *Arch of the colon*, (F.) *Arc du colon*,—the transverse portion of that intestine:—*Arch of the Aorta*, *Arcus aort'æ*, (F.) *Crosse de l'Aorte*, &c., the turn which the aorta takes in the thorax.

ARCA ARCANORUM, *Hydrargyrum*—a. *Cor-dis*, *Pericardium*.

ARCADE, see *Arc*—a. *Anastomotique*, *Arch*, *anastomotique*—a. *Crurale*, *Crural arch*—a. *Inguinale*, *Crural arch*—a. *Orbitaire*, *Orbital arch*—a. *Pubienne*, *Pubic arch*—a. *Zygomatique*, *Zygomatich arch*.

ARCADES DENTAIRES, *Dental arches*—a. *Palmaires*, *Palmar arches*.

ARCDI-TEMPORO-MAXILLAIRE, *Temporals*.

ARCE'US or ARCE'US, *BALSAM OF*, (F.) *Baume d'Arceus*. A kind of soft ointment used in sores, contusions, &c. It is made by melting two parts of mutton suet, one part of hog's lard:

turpentine and rosin, each one part and a half: straining and agitating till cold.

ARCANNO, *Colophonia*.

ARCANUM, from *arca*, 'a chest.' A secret, a *nostrum*, a *quack* or *empirical medicine*, (F.) *Arcane*. A remedy-whose composition is kept secret; but which is reputed to possess great efficacy.

ARCANUM CORALLINUM, *Hydrargyri nitricoxydum*—a. *Duplicatum*, *Potassæ sulphas*—a. *Tartari*, *Potassæ acetas*.

ARCEAU, *Arculus*, *Cradle*.

ARCEUTHOS, *Juniperus communis*.

ARCH, ANASTOMOTIC, (F.) *Arcade Anastomotique*, is the union of two vessels, which anastomose by describing a curved line. The vessels of the mesentery anastomose in this manner.

ARCH of the AORTA, see *Aorta*—a. *Crural*, see *Crural arch*—a. *Femoral*, see *Crural arch*—a. *Gluteal*, see *Gluteal aponeurosis*—a. *Hæmal*, see *Hæmal arch*—a. *Inguinal*, see *Crural arch*—a. *Orbital*, see *Orbital arch*—a. of the *Palate*, see *Palate bone*—a. of the *Pubis*, see *Pubic arch*—a. *Subpubic*, see *Subpubic arch*—a. *Superciliary*, see *Superciliary arches*—a. of a *Vertebra*, see *Vertebra*—a. *Zygomatich*, see *Zygomatich arch*.

ARCHES of the PALATE. These are two in number on each side of the throat, one of which is termed *anterior*, the other *posterior*.

The *anterior arch* arises from the middle of the *velum palati*, at the side of the *uvula*, and is fixed to the edge of the base of the tongue.

The *posterior arch* has its origin, likewise, from the side of the *uvula*, and passes downwards to be inserted into the side of the pharynx. The anterior arch contains the *circumflexus palati*, and forms the *isthmus faucium*. The posterior arch has within it the *levator palati*, and between the arches are the *tonsils*.

ARCHÆ'US, *Arche'us*, from *αρχη*, 'commencement,' (F.) *Archée*. A word invented by Basil Valentine, and afterwards adopted by Paracelsus and Van Helmont. The latter used it for the internal principle of our motions and actions. This archæus, according to Van Helmont, is an immaterial principle, existing in the seed prior to fecundation, and presiding over the development of the body, and over all organic phenomena. Besides this chief archæus, whose seat Van Helmont placed in the upper orifice of the stomach, he admitted several of a subordinate character, which had to execute its orders; one, for instance, in each organ, to preside over its functions; each of them being subject to anger, caprice, terror, and every human feeling.

ARCHANGEL, NEW, MINERAL SPRINGS. About twenty miles to the north of New Archangel, Sitka Island, on the N. W. coast of North America, are some thermal sulphureous waters, the temperature of one of which is upwards of 153° of Fahr. They are much celebrated.—Sir Geo. Simpson.

ARCHANGELICA, *Lamium album*—a. *Officinalis*, *Angelica*.

ARCHE, *αρχη*, *Init'ium*, *Princip'ium*, *Primordium*, *Ori'go*, *Inva'sio*. The first attack of a disease.

ARCHECPTOMA, *Proctocœle*.

ARCHÉE, *Archeus*.

ARCHELL, CANARY, *Lichen roccella*.

ARCHELOG'IA, from *αρχη*, 'beginning,' and *λογος*, 'a discourse.' A treatise on fundamental principles—of medicine, for example.

ARCHEN'DA. A powder of the leaves of the *ligustrum*, used by the Ægyptians after bathing, to obviate the unpleasant odour of the feet.—Prosper Alpinus.

ARCHIATER, *Archiat'rus*, *Protomed'icus*,

Protiatros, from *αρχα*, 'I am first,' and *ιατρος*, 'physician.' The original signification of this word is a matter of dispute. Some consider, with Mercurialis, that it meant physician to a prince, king, emperor, &c.; others, with C. Hoffmann, apply it to every physician who, by his situation, is raised above his colleagues. The former opinion seems to have prevailed — *Archiatre des Rois de France* being applied to the chief physician to the kings of France.

ARCHIG'ENI MORBI. Acute diseases; because they hold the first rank: from *αρχη*, 'beginning,' and *γινωσκει*, 'I am.'

ARCHIMAGIA, Chymistry.

ARCHINGEAY, MINERAL WATERS OF. Archingeay is situate in France, three leagues from St. Jean d'Angely. The waters are prized in all diseases. They seem to contain carbonate of lime, a little chloride of sodium, carbonate of iron, and some bitumen.

ARCHITECTURA APOPLECTICA, Apoplectic make.

ARCHITIS, Proctitis, Rectitis.

ARCHOCELE, Proctocele.

ARCHOPTOMA, Proctocele.

ARCHOPTOSIS, Proctocele.

ARCHORRHA'GIA, from *αρχος*, 'the anus,' and *ρεια*, 'I flow.' *Archorrhæa*. Hemorrhage from the anus.

ARCHORRHŒA, Archorrhagia.

ARCHOS, Arcus, Rectum.

ARCHOSTEGNOMA, Stricture of the Rectum.

ARCHOSTEGNOSIS, Stricture of the Rectum.

ARCHOSTENOSIS, Stricture of the Rectum.

ARCHOSYRINX, Fistula in ano.

ARCIFORM, *Arciformis*, from *arx*, *arcis*, 'a top or ridge,' and *forma*, 'shape.' An epithet given to certain fibres, *Fibræ arciformes*, of the anterior pyramids of the medulla oblongata, which take a curved course around the inferior extremity of each corpus olivare and ascend towards the cerebellum.

ARCTA'TIO, *Arctitudo*, from *arcto*, 'I make narrow;' *Angustatio*, *Coarctatio*, (F.) *Rétrécissement*. Contraction of a natural opening of a canal, and especially of the vulva, of the orifice of the uterus, or of the intestinal canal. Constipation, (see Stegnesis.) Reunion by suture or infubulation. — Scribonius Largus, Paul Zacchias, &c.

ARCTITUDO, Arctatio.

ARCTIUM, A. lappa.—a. Bardana, A. lappa.

ARCTIUM LAPPA. The root and seed of the *Clit'bur*, *Bardana*, *Αρκτιον*, *Arctium*, *A. bardana* seu *major* seu *minus* seu *tomentosum*, *Plaphis*, *Lappa glabra*, *Lappa major* seu *persona'ta*, *Persoll'ata*, *Persoll'ata*, *Persoll'ata*, *Burdock*, (Old Eng.) *Clithe*, *Cuckold*, (F.) *Bardane*, *Glouteron*, *Herbe aux teigneux*. Ord. Compositæ. Sex. Syst. Syngenesia æqualis. Root, *Lappa* (Ph. U. S.), diuretic; seed, cathartic. It has been used in decoction in diseases of the skin and in syphilis.

ARCTIUM MAJUS, A. lappa.—a. Minus, A. lappa.—a. *Tomentosum*, A. lappa.

ARCTOPUS ECHINATUS. A South African plant, Ord. Umbelliferae, which is demulcent and diuretic, somewhat approaching sarsaparilla. The decoction of the root is employed in syphilis, lepra, and chronic cutaneous affections of all kinds.

ARCTOSTAPHYLOS UVA URSI, *Arbutus uva ursi*.

ARCTU'RA, from *arcto*, 'I straighten.' The effects of a nail grown into the flesh, *Arctu'ra unguis*. See Onychogryphosis.

ARCTURA UNGUIUM. The growing in or inversion of the nails. See Onychogryphosis.

ARCUA'TIO, *Concavatio*. An anterior gibbosity or projection of the sternum.

ARCUEIL, MINERAL WATERS OF. Arcueil is about one league south of Paris. The water contains carbonic acid, carbonate of lime, sulphate of lime, chloride of sodium, and some deliquescent salts.

A celebrated society held its meetings at this village, of which Berthollet, Humboldt, La Place, &c., were members.

ARCULA CORDIS, Pericardium.

ARCULÆ. The Orbital Fossæ: *ποσιδεις*. — Rufus of Ephesus.

ARC'ULUS, diminutive of *arcus*, 'an arch.' A small arch; a cradle, (F.) *Arceau*, *Archet*. A semicircular box or basket used for preventing the bed-clothes from coming in contact with injured or diseased parts. An ordinance of the Grand Duke of Tuscany forbade mothers to sleep with an infant near them, unless it was put under a solid cradle.

ARCUS MEDULLARIS, Fornix.—a. Senilis, Gerotoxon.—a. Subpubicus, Subpubic arch.—a. Superciliaris, Superciliary arches.—a. Unguium, see Nail.—a. Zygomaticus, Zygomatic arch.

ARDALOS, Excrement.

ARDAS, Excrement.

ARDENT, *Ardens*, from *ardere*, 'to burn.'

ARDENT FEVER, (F.) *Fièvre ardente*. *Causus*, *Synocha*, or inflammatory fever.

ARDENT or INFLAMED EYES, (F.) *Yeux ardents*. The eyes are so called when injected red.

ARDENT URINE, (F.) *Urine ardente*. Urine of a deep red.

ARDESIA HIBERNICA, Hibernicus lapis.

ARDEUR, *Ardor*—a. du Cœur, Cardialgia—*a. d'Estomac*, *Ardor ventriculi*, Pyrosis—*a. de la Fièvre*, *Ardor Febrilis*—*a. d'Urine*, *Ardor Urinae*.

AR'DOR, (F.) *Ardur*. Heat. A feeling of burning, of violent heat; *Ætus*, *Æstus*, *Æstus*, *Æstus*.

ARDOR FEBRI'LIS, (F.) *Ardur de la Fièvre*. The hot period of fever.

ARDOR STOMACHI, Pyrosis.

ARDOR URINÆ, (F.) *Ardur d'Urine*. A scalding sensation occasioned by the urine in passing over the inflamed mucous membrane of the urethra, or over the neck of the bladder.

ARDOR VENEREUS, Heat.

ARDOR VENTRIC'ULI, *Ebullit'io Stomachi*, *Heartburn*, (Sc.) *Heartacis*, *Heartcald*, *Hersket*, (F.) *Ardur d'Estomac*. See Cardialgia and Pyrosis.

AREA, 'a void place,' 'an open surface.' A Latin word used by some authors to designate a variety of Alopecia, in which the hair changes colour, but does not fall off; also, *Porrigio decalvans*.

AREA GERMINATIVA, *Tache embryonnaire*.

AREA PELLUCIDA. An elliptical depression in the ovum, filled with a pellucid fluid, in the centre of which is the germ.

AREA VASCULO'SA, see *Circulus venosus*.

ARE'CA. The fruit—*Are'ca nut*, *Betel nut*—of *Are'ca Cat'echu* seu *Faufel*, *Cain'ga*; Ord. Palmæ; Sex. Syst. Monœcia Monadelphica; (F.) *Arec*, is astringent and tonic, and enters into the composition of the *Betel*, the great masticatory of the Orientals.

ARECA CATECHU, see *Areca*—a. *Faufel*, see *Areca*.

AREFAC'TION, *Arefactio*, *Xerun'is*, *ξηραν'is*, from *arefacere*, 'to make dry,' (*arere*, 'to dry,' and *facere*, 'to make.') The process of drying substances, prior to pulverization.

ARENA, see Gravel.

ARENAMEN, Bole Armenian.

ARENA'TIO, *Inhumatio*, *Chosis*, *Sand or*

Earth Bath; from *arena*, 'sand,' *Saburra'tio*. The application of hot sand to the body. *Pedi-lu'via* [?] of sand were formerly used in Asclepias.

ARENGA SACCHARIFERA, see Toddy.

ARENOSA URINA, *Sandy Urine*. Urine when it deposits a sandy sediment.

ARENOSUS, *Sabulous*. Also, one who passes sandy urine.

ARENULA, see Gravel.

AREOLA. A diminutive of *Area*, (F.) *Aire*. Anatomists understand by *Areola* the interstices between the fibres composing organs, or those existing between laminæ, or between vessels which interlace with each other.

Areola is also applied to the coloured circle *Halo*, *Halos*, which surrounds the nipple, *Areola papillaris*, and which becomes much darker during pregnancy; as well as to the circle surrounding certain vesicles, pustules, &c., as the pustules of the small-pox, the vaccine vesicle, &c. Chaussier, in such cases, recommends the word *Auricola*, (F.) *Aurêole*.

AREOLA PAPILLARIS, see *Areola*—a. Tubercles of the, see *Mamma*.

AREOLAR, *Areola'ris*. Appertaining to an areola.


AREOLAR EXHALATIONS are those recremen-tital secretions which are effected within the organs of sense, or in parenchymatous structures—as the aqueous, crystalline, and vitreous humours, &c.

AREOLAR TISSUE, Cellular Tissue.

AREOMETER, *Araom'eter*, *Gravim'eter*, *Al-coölom'eter*, *Alcoöm'eter*, *Ärostat'ic Balance*, from *apaus*, 'light,' and *μετρον*, 'measure,' i. e. 'measure of lightness.' An instrument

so called because first em-ployed to take the specific gra-vity of fluids lighter than water. The *Areometer* of *Baumé*, which is the most used in Pharmacy, particularly in France, consists of a tube of glass, largely expanded towards its inferior extremity, and terminating be-low by a small ball, containing mercury or lead, which serves it as a balance, so that it may remain upright in the fluid. This tube is furnished with a graduated scale. If the fluid into which the *Areometer* is plunged be heavier than water, the instrument rises; if lighter, it sinks. There are various *Areometers*, as those of the Dutch, of *Fahrenheit*, *Nichol-son*, &c. The *Areometer* is also called *Hydrom'eter*, (F.) *Aréo-mètre*, *Pèse-liqueur*.

There are some hydrometers which have a general applica-tion for determining the spe-cific gravities of liquids—as *Fahrenheit's*, *Nicholson's*, *Guy-ton de Morveau's*, and the com-mon glass hydrometers, inclu-ding *Baumé's*, *Cartier's*, *Twad-dle's*, *Zanetti's*, and the spe-cific gravity beads; others in-tended for special application—as for estimating the com-parative strength of spirits; the comparative densities of syr-ups, oils, &c.—as *Gay Lus-sac's*, *Sikes's*, and *Dicat's* hy-drometers, and the saccharometer, urinometer, and elæometer.



Baumé's Hydro-meter.

drometers, and the saccharometer, urinometer, and elæometer.

SCALE OF BAUMÉ'S HYDROMETER, WITH CORRE-SPONDING SPECIFIC GRAVITIES.

1. Ascending Scale for light liquids.

Scale of Baumé.	Specific Gravities.	Substances.
70	700	Pure hydrocyanic acid.— <i>Gay Lus-sac</i> .
66	715	Very pure sulphuric ether.
60	742	The same concentrated.
50	782	
48	792	Equal parts of alcohol and ether.
42	819	Very pure alcohol for pharmaceu-tical purposes.
40	827	
36	847	Pure alcohol. Naphtha.
33	863	Alcohol of commerce.
32	868	Essential oil of turpentine.
30	878	
26	900	Hydrocyanic acid of Scheele and pure hydrocyanic acid, mixed with an equal portion of water.— <i>Robiquet</i> .
25	906	
23	915	Acetic ether.
Id.	Id.	Nitric ether.
22	923	Muriatic ether.
Id.	Id.	Liquid ammonia.
20	935	Olive oil.
18	948	
13	980	Brandy.
12	986	
11	993	Burgundy wine.
10	1000	Bordeaux wine.
		Distilled water.

2. Descending Scale for heavy liquids.

Scale of Baumé.	Specific Gravities.	Substances.
0	1000	Common distilled water.
1	1007	
	1009	Distilled vinegar.
2	1014	Common vinegar.
3	1032	
4	1075	Cow's milk.
10	1075	
12	1091	Concentrated acetic acid.
20	1161	
21	1180	
25	1210	Liquid hydrochloric acid.
30	1261	Boiling syrup.
35	1321	Cold Syrup.
40	1384	Common nitric acid.
41	1398	
45	1454	Concentrated nitric acid.
Id.	Id.	
50	1532	Phosphoric acid for medical use.
60	1714	
66	1847	Very concentrated sulphuric acid.
70	1946	Very concentrated phosphoric acid

ARES. A term invented by *Paracelsus* to de-signate the principle on which depends the form of mercury, sulphur, and salt. These the alchymists regarded as the three bodies that give birth to every other.

AR'ETÉ, *apern*, 'virtue.' Mental or corporeal vigour.—*Hippocrates*.

ARETHU'SA, *A. bulbosa*; indigenous. *Order*, *Orchidaceæ*. The bruised bulbs are used in toothache; and as cataplasms to tumours.

A'REUS. A pessary mentioned by *Paulus of Ægina*.

ARGEL, *Cynanchum oleæfolium*.

AR'GEMA, *Argemon*, *Argemum*, from *apyos*, 'white.' *Fos'sula*, (F.) *Encavure*. A white spot or ulceration of the eye.—*Hippocrates*. See *Leucoma*.

ARGEMONÉ, MEXICANA, *Thorn Poppy*, *Prickly Poppy*, *Yellow Thistle*. A native of Mexico, but naturalized in most parts of the world. *Ord.* *Papaveraceæ*. *Sex. Syst.* *Polyan-dria Monogynia*. The juice resembles gamboge,

and has been used as a hydragogue. The seeds are employed in the West Indies as a substitute for ipecacuanha. They are also used as a cathartic.

ARGENSON, MINERAL WATERS OF. A chalybeate situate at Argenson in Dauphiny: used in cases of obstruction, jaundice, &c.

ARGENT, Argentum—*a. Chlorure d'*, see Argentum—*a. et d'Ammoniaque*, *chlorure d'*, see Argentum—*a. Iodure d'*, see Argentum—*a. Oxide d'*, see Argentum—*a. Petit*, Platinum—*a. Vice*, Hydragryum.

ARGENTERIA, *Potentilla anserina*.

ARGENTI CHLORIDUM, see Argentum—*a. et Ammoniae chloridum*, see Argentum—*a. et Ammoniae chloruretum*, see Argentum—*a. Cyanidum*, see Argentum—*a. Cyanuretum*, see Argentum—*a. Iodidum*, see Argentum—*a. Ioduretum*, see Argentum.

ARGENTI NITRAS, *Argentum Nitratum* seu *Nitricum*, *Sal argenti*, (F.) *Nitrate* ou *Azotate d'Argent*, *Nitrate of Silver*. This preparation is sometimes kept in crystals, the *Nitras Argenti* in *crystallos concretus*, *Nitrate d'Argent cristallisé* of the Codex of Paris, *Luna potabilis*, *Crystalli Lunæ*, *Argentum nitricum crystallisatum*, *Nitras argenti crystallinus*, *Nitrum lunare*, *Hydragrym Boylæ*. Generally, however, it is in the fused state: and it is this which is admitted into most Pharmacopœias, and which, besides the name *Nitras Argenti*, is called *Nitras argenti fusus* seu *fusum*, *Cauticum lunare*, *Lapis infernalis*, *Argentum nitricum fusum*, and *lunar caustic*, (F.) *Nitrate d'argent fondu*, *Pierre infernale*.

In the Pharmacopœia of the United States, it is directed to be prepared as follows:—Take of silver, in small pieces, 3j; nitric acid, f3vj, distilled water, f3ij. Mix the acid with the water, and dissolve the silver in the mixture in a sand bath; then crystallize, or gradually increase the heat, so that the resulting salt may be dried. Melt this in a crucible over a gentle fire, and continue the heat until ebullition ceases; then immediately pour it into suitable moulds.

The virtues of nitrate of silver are tonic, and escharotic. It is given in chorea, epilepsy, &c.: locally, it is used in various cases as an escharotic. Dose, gr. 1-8 to gr. 1-4 in pill, three times a day.

When silver is combined with iodine, it is said to have the same effect as the nitrate, and not to produce the slate colour of the surface, which is apt to follow the protracted use of the latter.

ARGENTI OXIDUM, see Argentum.

ARGENTILLA VULGARIS, *Potentilla anserina*.

ARGENTINE, *Argento'eus*, same etymon as the next. Pertaining to silver; as an 'argentine solution,' or solution of a salt of silver.

ARGENTINE, *Potentilla anserina*.

ARGENTUM, *Argentum*, from *argos*, 'white,' *Silver*, *Luna*, *Dia'na*. (F.) *Argent*. A solid metal of a shining white appearance; insipid; inodorous; highly sonorous; malleable and ductile; somewhat hard; crystallizable in triangular pyramids; fusible a little above a red heat, and volatilizable; s. g. 10.4. Not used in medicine, unless in some places for silvering pills. SILVER LEAF, *Argentum foliatum*, is the state in which it is used for this purpose.

ARGENTUM DIVISUM, *metallic silver*, in very fine powder, has been recommended internally in syphilis.

THE CHLORIDE (*Argenti chloridum*, *Argentum muria'tum* seu *chlora'tum* seu *salu'tum*. *Chloruretum Argenti*, *Chloruret* or *Muriate of Silver*, (F.) *Chlorure d'Argent*); the CYANURET;

the IODIDE (*Argenti Iodidum*, *Argentum Ioda'tum*, *Ioduretum Argenti*, *Iod'uret of Silver*, (F.) *Iodure d'Argent*); the OXIDE (*Argenti ox'idum*, *Argentum oxyda'tum*, (F.) *Oxide d'Argent*), and the CHLORIDE of AMMONIA and SILVER (*Argenti et Ammoniae chloridum*, *Argentum muria'ticum ammonia'tum*, *Chloruretum Argenti et Ammoniae*, *Chloruret of Silver and Ammonia*, *Ammonio-chloride of Silver*, (F.) *Chlorure d'Argent et d'Ammoniaque*), have been used in syphilis. At first, these different preparations were administered iatroleptically on the gums; the chloride, the cyanide and the iodide in the dose of 1-12th of a grain; the chloride of silver and ammonia in the dose of 1-14th of a grain, and the oxide of silver and divided silver in the dose of 1-8th and 1-4th of a grain. M. Serre, of Montpellier, who made many trials with them, soon found that these doses were too small; he therefore raised that of the chloride to 1-10th, and of the iodide to 1-8th of a grain, without any inconvenience resulting. The dose of the other preparations was likewise increased in a similar ratio. M. Serre extols the preparations of silver—used internally as well as iatroleptically—as antisyphilitics, but they are not to be depended upon.

The Cyanuret or Cyanide of Silver, *Argenti Cyanuretum* seu *Cyanidum*, *Argentum cyanogenatum*, (F.) *Cyanure d'argent*, is thus directed to be prepared in the Ph. U. S. (1842.) *Argent. Nit. 3xv*, *Acid Hydrocyan.*, *Aq. destillat. 3ā Oj*. Having dissolved the nitrate of silver in the water, add the hydrocyanic acid, and mix them. Wash the precipitate with distilled water and dry it. In the last edition of the Pharmacopœia, (1851,) it is directed to be prepared as follows:—*Nitrate of Silver*, dissolved in *distilled water*, is put into a tubulated glass receiver; *Ferrocyanuret of Potassium*, dissolved in *distilled water*, is put into a tubulated retort, previously adapted to the receiver. *Dilute Sulphuric Acid* is added to the solution in the retort; and, by means of a sand-bath and a moderate heat, distillation is carried on until the liquid that passes over no longer produces a precipitate in the receiver. The precipitate is then washed with distilled water, and dried.

The Oxide of Silver, *Argenti Ox'idum*, has been introduced into the last edition of the Ph. U. S. (1851). It is made by precipitating a solution of the *Nitrate of Silver* by solution of *Potassa*, drying the precipitate.

ARGENTUM CHLORATUM, see Argentum—*a. Cyanogenatum*, see Argentum—*a. Fugitivum*, *Hydragryum*—*a. Fusum*, *Hydragryum*—*a. Iodatam*, see Argentum—*a. Liquidum*, *Hydragryum*—*a. Mobile*, *Hydragryum*—*a. Muriaticum*, see Argentum—*a. Muriaticum Ammoniatum*, see Argentum—*a. Oxydatum*, see Argentum—*a. Salitum*, see Argentum—*a. Vivum*, *Hydragryum*.

ARGIL, PURE, *Argilla pura*.

ARGILE OCHREUSE PALE, *Bolus Alba*.

ARGILLA BOLUS FLAVA, *Terra Lemnia*—*a. Bolus rubra*, *Bole Armenian*—*a. Ferruginea rubra*, *Bole Armenian*—*a. Kalisulphurica*, *Alumen*—*a. Pallida*, *Bolus alba*.

ARGILLA PURA, *Terra Alu'minis* seu *bola'ris*, seu *argilla'cea pura* seu *depu'ra'ta* seu *hydra'ta*, *Alu'mina pura* seu *depu'ra'ta*, *Ox'idum alumin'i*, *pure Argil* or *Alumina*, (F.) *Alumine factice*. This substance, which is prepared by drying alum and exposing it, for twenty or twenty-five minutes, to a red heat, until the sulphuric acid is driven off, has been recommended in indigestion as antacid, as well as in vomiting and diarrhœa accompanied with acidity. The dose to a very young child is from 3ss to 3j; to older children from 3j to 3ij.

ARGILLA SULPHURICA ALCALISATA. Alumen—*a. Sulphurica usta*, Alumen exsiccatum—*a. Superphurica alcalisatum*, Alumen—*a. Vitriolata*, Alumen.

ARGILLÆ ACETAS. Aluminæ acetæ—*a. Sulphas*, Aluminæ sulphas.

ARGOL. RED, Potassæ supertartaras impurus—*a. White*, Potassæ supertartaras impurus.

ARGUMENTUM INTEGRITATIS. Hymen.

ARGYRIA, from *argyros*, 'silver.' The discoloration of the skin occasioned by the internal use of nitrate of silver.

ARGYROCHÆTA, Matricaria.

ARGYROPH'ORA, from *argyros*, 'silver,' and *φωρ*, 'I bear.' A name given, by Myrepsus, to an antidote which he regarded as extremely precious.

ARGYROTROPHEMA, *Blanchemanger*.

ARGYRUS, Argentum.

ARHEUMATIC, *Arheumat'icus*, from *a*, privative, and *ρευμα*, 'fluxion or rheumatism.' One without fluxion or rheumatism.

ARIA, *Cratægus aria*—*a. Cattiva*, see Miasma.

ARIC'INA, *Aric'ia*, *Cus'conin*, *Cusco-Cincho'nia*, so called from Arica in South America, the place where it is shipped. An alkaloid found in Cusco Bark, which is very similar in many of its properties to Cinchonia. Cusco was the ancient residence of the Incas.

ARIC'YMON, from *api*, an intensive particle, and *κρυειν*, 'to conceive.' A name given to a female who conceives readily.—Hippocrates.

ARIDE'NA. A Latin word employed to designate the leanness of any part.—Ettmuller, Sauvages.

ARID'ITY, *Arid'itas*. (F.) *Aridité*, from *arere*, 'to grow dry.' The French use the word *Aridité* to express the dryness of any organ, and particularly of the skin and tongue, when such dryness is so great as to render the organ rough to the touch. *Aridité* also means the lanuginous appearance of the hair in some diseases in which it seems covered with dust.

ARIDU'RA. Wasting or emaciation of the whole or of any part of the body; Marasmus, Atrophy.

ARIDURA CORDIS, Heart, atrophy of the—*a. Hepatis*, Hepatophria.

ARIKA, see Siro.

ARISÆMA ATRO-RUBENS, *Arum triphyl-lum*.

ARISTALTHÆA, *Althæa*.

ARISTOLOCH'IA, from *apistos*, 'very good,' and *λοχια*, 'parturition,' so called, because the different varieties were supposed to aid parturition. *Birthwort*, (F.) *Aristolochie*. *Ord. Aristolochiaceæ*. Several varieties were once in use.

ARISTOLochia CAVA, *Fumaria bulbosa*.

ARISTOLOCH'IA CLEMAT'IS, *Aristoloch'ia Vulgaris* seu *Cre'tica seu ten'uis*, *Adra Riza* (F.) *Aristolochie ordinaire*, *Upright Birthwort*. The root has been considered stimulant and emmenagogue, and as such has been used in amenorrhœa, chlorosis, and cachexia.

ARISTOLOCHIA CRETICA, *A. Clematidis*—*a. Fabacea*, *Fumaria bulbosa*—*a. Hastata*, see *A. serpentaria*—*a. Hirsuta*, see *A. serpentaria*.

ARISTOLOCH'IA LONGA, and *A. Rotun'da*, (F.) *Aristolochie longue et ronde*, *Long and Round Birthwort*. Virtues the same as the preceding.

ARISTOLOCH'IA PISTOLOCH'IA, *Pistoloch'ia Aristoloch'ia*, *Polyrrhi'za*. This variety has an aromatic odour, and an acrid and bitter taste. (F.) *Aristolochie crénelée*.

ARISTOLOCHIA RETICULATA, see *A. serpentaria*. ARISTOLOCH'IA SERPENTARIA, *Serpentaria*, *S. Virginia'na*, *Viper'a*, *Viper'ia*, *Viper'ina* *Virginia'na*, *Culubri'na* *Virginia'na*, *Contrayer'va* *Virginia'na*, (F.) *Serpentaire et Aristolochie serpentaire de*

Virginie, *Culverine* ou *Couleuvrée* ou *Viperine de Virginie*, *Virginia Snakeroot*, *Snakeroot Birthwort*, *Snakeweed*, *Snagrel*. Virtues of the root, *Serpentaria*, tonic, stimulant; and as such, employed in debility, intermittents, &c. ARISTOLOCH'IA HIRSU'TA, *A. HASTA'TA* and *A. RETICULA'TA*; indigenous, are employed indiscriminately with *A. Serpentaria* under the name *Serpentaria* or *Virginia Snakeroot*.

ARISTOLOCH'IA SIPHO, *Dutchman's pipe*, and *A. TOMENTO'SA*; indigenous, have virtues like those of *A. Serpentaria*.

ARISTOLOCH'IA TENUIS, *A. Clematidis*—*a. Tomentosa*, see *A. sipo*—*a. Trifida*, *A. Trilobata*.

ARISTOLOCH'IA TRILOBA'TA, *A. trif'ida*, (F.) *Aristolochie trilobée*. A plant of Surinam and Jamaica; possessing the general virtues of the *Aristolochiaceæ*. The other varieties of *Aristolochia* have similar properties.

ARISTOLOCH'IA VULGARIS ROTUNDA, *Fumaria bulbosa*.

ARISTOLOCH'IC, *Aristoloch'icus*. Same etymology. An old term for remedies supposed to have the property of promoting the flow of the lochia.—Hippocrates, Theophrastus, Dioscorides, &c.

ARIS'TON MAGNUM, and ARISTON PARVUM. These names were formerly given to pharmaceutical preparations, used in phthisis, tormina, and fever.—Avicenna.

ARISTOPHANE'ON. A sort of emollient plaster, prepared with four pounds of pitch, two of apoc'hyma, one of wax, an ounce of opoponax, and half a pint of vinegar.—Gorræus. Not used.

ARKAN'SAS, MINERAL WATERS OF. About 5 miles from the Washita river, and about a quarter of a degree north of the Louisiana line, there are about 70 of those springs;—*Quachita* or *Washita Springs*. They are thermal, varying from 138° to 150° Fahrenheit, and are employed in rheumatism, cutaneous affections, &c.

ARK-BEIN, Pubis os.

ARKEION, *Arctium Lappa*.

ARLADA, *Realgar*.

ARLES, MINERAL WATERS OF. Thermal sulphureous springs in the department of Pyrénées Orientales, France. Their temperature is 103° to 145° of Fahr., and they contain sulphohydric acid.

ARM, Brachium.

ARMA, Penis—*a. Ventr*, Penis.

ARMAMENTARIUM, Arsenal—*a. Chirurgicum*, see Arsenal—*a. Portabile*, see Case.

ARMATORY UNGUENT, *Hoplochrysma*.

ARMATURA, Amnios.

ARME, from *apw*, 'I adapt.' Any physiological or mechanical junction or union of parts.—Hesychius. *A. suture*, as of the cranium.—Galen.

ARMENIACA EPIROTICA, *Prunus Armeniaca*—*a. Malus*, *Apricot*, see *Prunus*—*a. Vulgaris*, *Prunus Armeniaca*.

ARMENIAN STONE, *Melochites*.

ARMENITES, *Melochites*.

ARMILLÆ MANUS MEMBRANOSÆ, Annular ligaments of the carpus.

ARMOISE BLANCHE, *Artemisia rupestris*—*a. Commune*, *Artemisia vulgaris*—*a. Estragon*, *Artemisia dracunculus*—*a. Ordinaire*, *Artemisia vulgaris*.

ARMONIACUM, *Ammoniac*, gum.

ARMORA'CIA, from *Armorica*, where it was largely cultivated. In the Pharmacopœia of the United States, the fresh root of *Cochlearia armoracia*.

ARMORACIA RUSTICANA, *Cochlearia armoracia*—*a. Sativa*, *Cochlearia armoracia*.

ARMOUR, Condom.

ARMURE DES JAMBES, see Cornu am-

nis.

ARMUS, Humerus—a. Summus, Acromion.

ARN. *Alnus glutinosa*.

ARNALDIA, *Arnaldie*. A disease, which appears to have prevailed in England at one time, but whose origin and nature are unknown. It was accompanied with loss of hair; and, by some, is supposed to have been Syphilis. "Deinde uterque Rex incidit in egritudinem, quam *Arnaldium* vocant, in qua ipsi usque ad mortem laborantes, capillos suos deposuerunt."—Rogerus Hovedenus, in Ricardo I., cited by Du Cange.

ARNICA MONTANA. Derivation uncertain. *Arnica*, *Leopard's Bane*, *Doronicum Germanicum* seu *Oppositifolium* seu *Plantaginifolium*, *Arnica Plauensis*, *Alisma*, *Aclyrus*, *Diuretica*, *Punicea lapsidum*, *Plar'nica montana*, *Caltha* seu *Calceolula Alpina*, *Nardus Celtica altera*, (F.) *Arnique*, *Bétoine* ou *Tabac des Montagnes*, *Tabac des Vosges*, *Tabac* ou *Bétoine des Savoyards*, *Doronic d'Allemagne*, *Plantain* ou *Sauai des Alpes*. *Sex. Syst.* Syngenesia Polygamia superflua. *Ord.* Compositæ. The plant and flowers (*Arnica*, Ph. U. S.) are considered, or have been considered, narcotic, stimulant, emmenagogue, &c.; and, as such, have been given in amaurosis, paralysis, all nervous affections, rheumatism, gout, chlorosis, &c. Dose, gr. v to x, in powder. In large doses it is deleterious.

ARNICA NUDICAULIS and A. MOLLIS are supposed to have medicinal virtues like the last.

ARNICA PLAUENSIS, *Arnica montana*—d. Spuria, *Inula dysenterica*—a. *Suedensis*, *Inula dysenterica*.

ARNIQUE, *Arnica montana*.

ARNOGLOSSUM, *Plantago*.

ARNOTT'S DILATOR, see Dilator, *Arnott's*.

ARNUT, *Bunium bulbocastanum*.

AROMA, *Ar'tyma*, 'perfume': (*apt*, intensive, and *osm* or *odm*, 'odour,' or from *apa*, 'to make fitting or agreeable.') *Spiritus Rector*, (F.) *Arôme*. The odorous part of plants. An emanation—frequently imponderable—from bodies, which acts on the organ of smell, and varies with the body exhaling it.

AROMATE, *Aromatic*.

AROMATIC, *Aromaticus*, (F.) *Aromate*. Any odoriferous substance obtained from the vegetable kingdom which contains much volatile oil, or a light and expansible resin. Aromatics are used in perfumes, in seasoning, and embalming. In medicine they are employed as stimulants. Ginger, cinnamon, cardamoms, mint, &c., belong to this class.

AROMATIZE, (F.) *Aromatiser*. To add to a mixture or potion some aromatic, to mask its taste or render it more agreeable. Such mixture is then said to be aromatized.

AROMATOPLA, from *aroma*, 'an odour,' and *πωλω*, 'I sell.' An apothecary or druggist. One who sells spices.

ARON, *Arum*.

AROPH. A barbarous word, which has various significations with the ancients. Paracelsus employed it to designate a lithontripctic remedy. The mandragora, according to some. Also, a mixture of bread, saffron, and wine.—Van Helmont.

AROPH PARACELSI, *Ferrum ammoniacum*.

ARQUEBUSADE, *EAU D'*, *Aqua traumatica Thedæni* seu *Thedæna seu scolopetaria seu vulneraria seu catapulta'rium*, *Mistura vulneraria ac'idia*. A sort of vulnerary water, distilled from a farrago of aromatic plants. *Rosemary*, *hiss: millefoil*, *thyme*, each *hss*: *Proof spirit*, 2 gallons—distil a gallon. This is one form.

ARR, *Cicatrix*.

ARRABON, *Arraphon*.

ARRACHEMENT (F.), from *arracher*, 'to tear out,' *Apospas'ma*, *Abruptio*, *Avulsio*. Act of separating a part of the body by tearing it from the bonds connecting it with others. *Eviscion*. *Laceration*.

Arrachement is applied to certain operations, as to the extraction of a tooth, the extirpation of a *polypos*, &c.

ARRACK, *Arack*. See *Spirit*.

ARRAPHON, *Ar'rabon*, from *a*, *priv.*, and *παφν*, 'a suture,'—'without suture.' A term applied to the cranium when it presents no sutures.

ARRECTIO, *Erection*.

ARRED, see *Scar*.

ARREPTIO, *Insanity*.

ARRESTA BOVIS, *Ononis spinosa*.

ARRÊT D'HILDAN, *Remora Hildani*.

ARRÊTE-BŒUF, *Ononis spinosa*.

ARRHŒA, from *a*, *priv.*, and *ρρω*, 'I flow.' The suppression of any flux. *Amenorrhœa*.

ARRHOSTEMA, *Disease*.

ARRHOSTENTIA, *Disease*.

ARRHOSTIA, *Disease*, *Infirmity*.

ARRHYTHMUS, *Caecorhythmus*.

ARRIBA, *Geoffræa vermicifuga*.

ARRIÈRE-BOUCHE, *Pharynx*—a. *-Dent*, see *Dentition*—a. *-Faix*, *Secundines*.

ARRIÈRE-GOUT (F.), 'after taste.' The taste left by certain bodies in the mouth for some time after they have been swallowed, owing perhaps to the papillæ of the mouth having imbibed the savoury substance.

ARRIÈRES NARINES, *Nares*, posterior.

ARROCHE, *Atriplex hortensis*—a. *Puant*, *Chenopodium vulvaria*.

ARROSEMENT, *Aspersio*.

ARROWHEAD, *Sagittaria variabilis*.

ARROW LEAF, *Sagittaria variabilis*.

ARROW POISON. This differs with different tribes of Indians. By some, the poison capsicum, and infusions of a strong kind of tobacco, and of euphorbiacæ are mixed together, with the poisonous emmet, and the teeth of the formidable serpent, called, by the Peruvian Indians, *Miuamaru* or *Jergon*—*Lachesis picta* of *Tschudi*.

ARROW ROOT, *Fecula Maran'ta*, *Am'ylum marantaceum* seu *America'num*. The fecula of the rhizoma of *Maran'ta Arundinacea*, *Maranta* (Ph. U. S.), *Ord.* *Marantaceæ*, which, like all feculæ, is emollient and nutritive, when prepared with water, milk, &c.

Dr. Carson has shown, that *Florida arrow root* is derived from *Za'mia integrifolia* or *Z. pu'mila*, *Sugar pine*, *Ord.* *Cycadaceæ*; *Bermuda arrow root* being obtained from *Maranta arundinacea*. *Florida arrow root*, as well as the farina, is known in the Southern States under the name *Coonti* or *Coontie*.

According to Dr. Ainslie, an excellent kind of arrow root is prepared in Travancore from the root of *Curcuma angustifolia*, *Ord.* *Zingiberaceæ*.

Arrow root mucilage is made by rubbing *arrow root powder* with a little cold water, in a basin, by means of the back of a spoon, until it is completely mixed with the water; then pouring boiling water over it, stirring assiduously until a soft, gelatinous, tenacious mucilage is formed; and, lastly, boiling for five minutes. A tablespoonful of arrow root powder is sufficient to make a pint of mucilage. It may be moderately sweetened; and wine or lemon juice may be added.

With milk also it forms a bland and nutritious article of diet.

ARROW ROOT, BRAZILIAN. The fecula of *Jatropha Manihot*.

ARROW ROOT, COMMON, see *Solanum tuberosum*.

ARROW ROOT, EAST INDIAN. The fecula of the tubers of *Curcuma angustifolia*, or narrow-leaved Turmeric.

ARROW ROOT, ENGLISH, Arrow root, common.

ARROW ROOT, TAHITI, *Tacca stearch*, *Otaheiti Salep*. The fecula of *Tacca Oceanica*.

ARROW WEED, *Sagittaria variabilis*.

ARROW WOOD, *Euonymus*, *Viburnum dentatum*.

ARS CABALISTICA, Cabal—a. Chymiatrica, Chymistria—a. Glysmatia nova, Infusion of medicines—a. Coquinaria, Culinary art—a. Cosmetics, Cosmetics—a. Culinaria, Culinary art—a. Empirica, Empiricism—a. Formulas medicas concinandi, see Prescription—a. Hermetica, Chymistry—a. Homoeopathica, Homoeopathy—a. Hydriatrica, Hydrosudotherapeia—a. Infusoria, Infusion of medicines—a. Machaonia, Medicina—a. Majorum, Chymistry—a. Medica, Medicina—a. Obstetrica, Obstetrics—a. Sanandi, Art, healing—a. Separatoria, Chymistry—a. Signata, Cabal—a. Spagyrica, Chymistry—a. Sphygmica, Sphygmologia—a. Veterinaria, Veterinary Art—a. Zoiatrica, Veterinary Art.

ARSALTOS, Asphaltum.

ARSATUM, Nymphomania.

ARSECOCKLE (Se.), synonymous with Teut. aers-bleyne, *tuberculus in ano*. Originally, a hot pimple on the hips. Now, in Scotland, a similar pimple on the face or any part of the body.—Jamieson.

ARSENAL (F.), *Chirapotheca*, *Armamentarium*, *A. Chirurgicalium*. A collection of surgical instruments. A work containing a description of surgical instruments. *Armamentarium obstetricum* has the same significations in regard to obstetrics.

ARSENATE, *Arsen'ias*, (F.) *Arse'niat*, from *αἰσν*, 'robust.' A salt formed by a combination of arsenic acid with a salifiable base.

ARSENATE OF AMMONIA, *Arsen'ias Ammo'niæ*, *Ammonium Arsenicum* seu *Arsenicicum*, (F.) *Arse'niat d'Ammoniaque*. This preparation is highly extolled in cutaneous diseases. A grain of the salt may be dissolved in an ounce of distilled water, and 20 to 25 drops be commenced with as a dose.

ARSENATE OF IRON, *Arsen'ias Ferri*, *Ferrum Arseniatum* seu *Arsenicicum oxydulatum*, (F.) *Arse'niat de Fer*. This preparation has been applied externally to cancerous ulcers. An ointment may be made of 3ss of the arseniate, 3ij of the phosphate of iron, and 3vj of spermaceti ointment. The arseniate has also been given internally in cancerous affections, in the dose of one-sixteenth of a grain.

ARSENATE OF POTASSIUM, *POTASSIUM, Proto-arsen'iate of Potas'sium*, *Arsen'iate of Potassa*, *Arsen'ias Potassæ* seu *Kali*, (F.) *Arse'niat de Potasse*, *Sel arsenical de Macquer*. Properties the same as those of arsenious acid.

ARSENATE OF QUINA, *Quinæ Arsenias*.

ARSENATE OF SODA, *Sodæ arsen'ias*, (F.) *Arse'niat de Soude*. Made, according to the Codex of Paris, by mixing and heating to redness arsenious acid and nitrate of soda; dissolving the salt in water; adding carbonate of soda until there is an alkaline reaction; filtering, evaporating, and crystallizing. It is the basis of *Pearson's Solution*, which is formed of one grain of the crystallized arseniate to one fluidounce of distilled water. Dose, grt. xx. *Heinecke's Solution* consists of the arseniate, gr. vj; peppermint water, f 3iiss; wine of cinnamon, f 3ss; laudanum, f 3j. Dose, forty to fifty drops.

ARSENATE OF SODA, *Arseniate of Soda*.

ARSENIC, *Arsenicum*, (Ph. U. S.) A solid

metal; of a steel-gray colour; granular texture; very brittle; volatilizing before melting; very combustible and acidifiable. It is not dangerous of itself, and only becomes so by virtue of the facility with which it absorbs oxygen.

ARSENIC BLANC, *Arsenicum album*.

ARSENIC, IODIDE OF, *Arsen'ici Iod'idum* seu *Teriod'idum* seu *Iodure'tum*, *Arsenicum Iodatum*, *Hydri'odæ arsen'ici*, formed by the combination of arsenic and iodine. This preparation, applied externally, has been highly extolled in various cutaneous affections. An ointment may be made of three grains of iodide to 3j of lard. It has also been given internally in the dose of a tenth of a grain in similar affections. It is official in the Ph. U. S. 1851.

ARSENIC, OXIDE OF, *Arsenicum album*—a. Oxide of, White, *Arsenicum album*—a. White, *Arsenicum album*.

ARSENIC AND MERCURY, IODIDE OF, *Hydrargyri et Arsen'ici Iod'idum*, *Double Iodide of Mercury and Arsenic*, *Iodo-arsenite of Mercury*. A compound, which has been proposed as more efficacious than either the iodide of arsenic or the iodide of mercury. It is made by triturating 6·08 grains of arsenic; 14·82 grains of mercury; 49 of iodine, with a fluidrachm of alcohol, until the mass has become dry, and from being deep brown has become pale red. Eight ounces of distilled water are poured on, and, after trituration for a few moments, the whole is transferred to a flask; half a drachm of hydriodic acid, prepared by the acidification of two grains of iodine, is added, and the mixture is boiled for a few moments. When the solution is cold, make the mixture up to f 3viij with distilled water. This is called by Mr. Donovan, the proposer, *Liquor Arsen'ici et Hydrargyri Iodidi*, each drachm of which, by measure, consists of water, 3j; arsenious acid, gr. 1·8th; peroxide of mercury, gr. 1·4th; iodine, converted into hydriodic acid, gr. 3·4ths. In the last edition of the Ph. U. S. it is directed to be made of *Arsenicum Iodidum* and *Hydrargyri Iodidum rubrum*, each gr. xxxv; and *Aqua destillata*, Oss; dissolving by rubbing, heating to the boiling point, and filtering.

The dose of *Donovan's Solution* is from ℥x to f 3ss, two or three times a day.

It has been used successfully in inveterate cutaneous diseases.

ARSENICAL, *Arsenicalis*, (F.) *Arse'nic*. Relating, or appertaining to, or containing, arsenic.

ARSENICAL PASTE, (F.) *Pâte Arsenicale*. This application to cancers is formed of 70 parts of cinnabar, 22 of dragon's blood, and 8 of arsenious acid; made into a paste with saliva, when about to be applied.

ARSENICAL POWDERS OF FRÈRE CÔME, AND OF JUSTAMOND, see Powder, Arsenical.

ARSENICAL SOLUTION, see Arseniate of Soda, and Liquor Arsenicalis.

ARSENICI HYDRIODAS, Arsenic, Iodide of—a. Iodidum, Arsenic, Iodide of—a. Ioduretum, Arsenic, Iodide of—a. Teriodidum, Arsenic, Iodide of.

ARSENICISM'US, *Intoxicatio Arsenicalis*. Poisoning by arsenic.

ARSENICUM, Arsenic.

ARSENICUM ALBUM, White Arsenic, Oxide of Arsenic, *Ratsbane*, *Arsen'ici oxydum album*, *Calx Arsen'ici alba*, *Ac'idum Arsenico'sum* seu *Arsenio'sum* (Ph. U. S.), *Arsenious acid*, *White oxide of arsenic*, (F.) *Arsenie blanc*. An acid which is met with in commerce, in compact, white, heavy, fragile masses; of a vitreous aspect, opaque, and covered with a white dust; of an acrid and nauseous taste; without smell when cold; volatilizable by heat, and exhaling the odour of gar-

lic; soluble in water, alcohol, and oil; crystallizable in regular octahedrons. It is this that is meant by the name arsenic, as commonly used.

ARSENICUM ALBUM SUBLIMATUM, *Sublimed Oxide of Arsenic*, is the one employed in medicine. It is tonic and escharotic, and is the most virulent of mineral poisons. It is used in intermittents, periodical headaches, neuroses, &c. Dose, gr. one-tenth to one-eighth, in pill. See Poisons, *Table of*.

ARSENICUM TOPATUM, Arsenic, Iodide of—*a. Rubrum factitium*, Realgar.

ARSENIS POTASSÆ, Arsenite of protoxide of potassium—*a. Potassæ aqueus*, Liqueur arsenicalis—*a. Potassæ liquidus*, Liqueur arsenicalis.

ARSENITE, *Ar'senite*. A salt, formed by a combination of the arsenious acid with a salifiable base.

ARSENITE OF PROTOXIDE OF POTASSIUM, *Proto-arsenite of Potassium*, *Ar'senite of Potassa*, *Ar'senite Potassæ*. An uncrystallizable and colourless salt, which forms the basis of the *liqueur arsenicalis*, which see.

ARSENITE OF QUINA, Quinine arsenic.

ARSE-SMART, BITING, Polygonum hydropiper.

ART, HEALING, *Are Sanan'di*, *Medici'na*. The appropriate application of the precepts of the best physicians, and of the results of experience to the treatment of disease.

ART, VETERINARY, Veterinary art.

ARTABĒ, ἀραβή. Name of a Persian measure for dry substances, in use with the ancients, equal at times, to 5 modii: at others, to 3; and at others, again, to 7.—Galen.

ARTANTHE ELONGATA, see Matico.

ARTELSHEIM, MINERAL WATERS OF. These German waters have been much recommended in hysteria, gout, palsy, &c. Their physical or chemical properties have not been described.

ARTEMISIA, *Anact'ri'on*. Ord. Compositæ. Called after a queen of the name, who first employed it; or from *Apræus*, 'Diana,' because it was formerly used in diseases of women, over whom she presided. The Gauls called it *Bricunum*.

ARTEMISIA ABROTANUM, *Abrot'anum*, *Abrot'anum*, *Abrot'anum Cathum* seu *Mus*, *Abathan*, *South'ernwood*, *Oldman*, *Slovenwood*, (F.) *Abrotone*, *Aurone*, *Aurone mâle*, *Aurone des jardins*, *Garderobe*, *Citronelle*. Supposed to be possessed of stimulant properties.

Oil of Southernwood, *O'leum Abrot'ani*, (F.) *Huile d'Aurone*, possesses the aromatic properties of the plant.

ARTEMISIA ABSINTHIUM, *Absin'thium*, *A. vulgârè*, *Apsin'thium*, *Barypi'eron*, *Common Wormwood*, *Wormit*, (F.) *Absinthe*. Properties:—tonic and anthelmintic. The Oil of Wormwood, *O'leum Absin'thi*, (F.) *Huile d'Absinthe*, contains the aromatic virtues of the plant.

ARTEMISIA AFRA, a South African species, is tonic, antispasmodic and anthelmintic; and has been used in debility of the stomach, visceral obstructions, jaundice and hypochondriasis. It is taken in infusion, decoction and tincture. A strong infusion is used by the Cape Colonists as a collyrium in weakness of the eyes; and the pounded leaves and stalks are employed as discutients in oedema and sugillations.

ARTEMISIA ALBA, *A. Santonica*—*a. Balsamita*, *A. Pontica*.

ARTEMISIA BIEN'NIS, *Biennial Wormwood*; indigenous.

ARTEMISIA BOTRYS, *Chenopodium ambrosioides*.

ARTEMISIA CAMPES'TRIS, *Field Southernwood*,

(F.) *Aurone des Champs*. This possesses the same properties as *A. Abrot'anum*.

ARTEMISIA CANADENSIS, Canada Wormwood—*a. Caudata*: indigenous.

ARTEMISIA CHENOPODIUM, *Chenopodium botrys*.

ARTEMISIA CHINENSIS, *A. Indica*, *A. Moxa*. From this the Chinese form their moxas.

ARTEMISIA CONTRA grows in Persia, Asia Minor, and other parts of the East.

ARTEMISIA DRACUNCULUS, *Tar'agon*, (F.) *Armoise estragon*, *Estragon*. Virtues:—the same as the last. The leaves are much used to season food, especially in the form of *Es'tragon vin'egar*.

ARTEMISIA GLACIALIS, *Silky Wormwood*;

ARTEMISIA INDICA, *Artemisia Chinensis*, *A. Santonica*;

ARTEMISIA JUDICA, grows in Palestine, Arabia, China, Cochinchina, and Northern Africa.

ARTEMISIA LEPTOPHYLLA, *A. Pontica*;

ARTEMISIA MARITIMA, *Absin'thium Mari'num* seu *Mari'tinum*, *Sea Wormwood*, *Maritime South'ernwood*;

ARTEMISIA MOXA, *A. Chinensis*;

and

ARTEMISIA PONTICA, *A. Roma'na* seu *Tenuifolia* seu *Balsamita* seu *Leptophylla*, *Absinthium Ponticum* seu *Roma'num*, *Roman Wormwood*, *Lesser Wormwood*, possess like virtues;—as well as

ARTEMISIA ROMANA, *A. Pontica*;

ARTEMISIA RUBRA, *A. Santonica*; and

ARTEMISIA RUPES'TRIS, *Creeping Wormwood*, *Gen'ipi album*, (F.) *Armoise blanc*, *Génipi blanc*. This variety has aromatic virtues, and is used in intermittents, and in amenorrhœa.

ARTEMISIA SANTONICA, *Santon'icum*, *Canni Herba*, *Chamæcedris*, *Chamæcyparissus*, *Absinthium Santonicum*, *Sementi'na*, *Xantoli'na*, *Scheba Ar'abum*, *Sina* seu *Cina* *Levan'tica*, *Tartarian South'ernwood*, (F.) *Barbotine*. From this and other species of absinthium is obtained the so-called *Semen contra vermes* seu *Contra* seu *Zedaira'æ* seu *Santon'ici* seu *Cina*, *Hagiogermum*, *Sanctum Semen*, *Wormseed*, which do not consist of seeds, but of broken peduncles. Virtues:—anthelmintic and stimulant. Dose, gr. x. to ʒi in powder. Its active principle is *San'tonin*, the dose of which is from 2 to 5 grains in the day, for a child six or eight years of age.

ARTEMISIA TENUIFOLIA, *A. Pontica*.

ARTEMISIA VULGARIS, *Artemisia rubra et alba*, *Cin'gulum Sancti Joan'nis*, *Mater Herbal'rum*, *Berenise'cum*, *Busbastecor'dium*, *Canapa'cia*, *Mugwort*, (F.) *Armoise ordinaire*, *A. Commune*, *Herbe de Saint Jean*, *Ceinture ou Couronne de Saint Jean*. This, as well as some other varieties, possesses the general tonic virtues of the *Artemisia*. *Artemisia vulgaris* has been highly extolled by the Germans in cases of epilepsy. Dose of the powder, in the 24 hours, from ʒss to ʒi.

ARTÈRE, Artery—*a. Brachiale*, *Brachial artery*—*a. Brachio-céphalique*, *Innominate arteria*—*a. Bronchique*, *Bronchial artery*—*a. Ciliaire*, *Ciliary artery*—*a. Clitorienne*: see *Clitoris*—*a. Cœcale*: see *Colic arteries*—*a. Collatérale du coude*, *Anastomoticus magnus ramus*—*a. Collatérale externe*, *Arteria profunda humeri*—*a. Collatérale interne*, *Anastomoticus magnus ramus*—*a. Coronaire des lèvres*, *Labial artery*—*a. Coronaire Stomachique*, *Coronary artery*—*a. Crurale*, *Crural artery*—*a. Deuxième des thoraciques*, *Arteria thoracica externa inferior*—*a. Épineuse*, *Meningeal artery*, middle—*a. Femoro-poplitée*, *Ischiatic artery*—*a. Fessière*, *Gluteal artery*—*a. Gastrique droite*, *petite*, *Pyloric artery*—*a. Gutturo-maxillaire*, *Maxillary artery*, internal—*a. Honteuse externe*, *Pudic*, external, artery—*a. Honteuse interne*, *Pu-*

dic, internal, artery—*a. Humérale profonde*, Arteria profunda humeri—*a. Iliacæ primitivæ*, Iliac artery—*a. Innominatæ*, Innominata arteria—*a. Irienne*, Ciliary artery—*a. Ischio-pœnicæ*: see Pudic, internal, artery—*a. Mœdiane antérieure*, Spinal artery, anterior—*a. Mœdiane postérieure du rachis*, Spinal artery, posterior—*a. Mœningée moyenne*, Meningeal artery, middle—*a. Mœningée postérieure*, Prevertebral artery—*a. Montonière*, Mental foramen—*a. Mœsœcephalique*, Basillary artery—*a. Mœsœcalique*: see Colic artery—*a. Musculaire du bras*, Arteria profunda humeri—*a. Musculaire du bras, grande*: see Collateral arteries of the arm—*a. Musculaire grande de la cuisse*, Arteria profunda femoris—*a. Opisthogastrique*, Celiac artery—*a. Orbitaire*, Ophthalmic artery—*a. de l'Oraire*, Spermatic artery—*a. Pelvi-crurale*, Crural artery—*a. Pelvi-crurale*, Iliac artery—*a. Pelvienne*, Hypogastric artery—*a. Première des thoraciques*, Arteria thoracica externa superior—*a. Prévertébrale*, Prevertebral artery—*a. Radio-carpienne transversale palmaire*, Radio-carpal artery—*a. Scrotale*, Pudic, external, artery—*a. Sous-clavière*, Subclavian artery—*a. Sous-pubio-fœmorale*, Obturator artery—*a. Sous-pubienne*, Pudic, internal, artery—*a. Sous-sternal*, Mammary, internal—*a. Spino-épineuse*, Meningeal artery, middle—*a. Stomogastrique*, Coronary artery—*a. Sus-carpienne*: see *Sus-carpien*—*a. Sus-maxillaire*, Alveolar artery—*a. Sus-maxillaire*, Buccal artery—*n. Sus-metatarsienne*, Metatarsal artery—*a. Sus-pubienne*, Epigastric artery—*a. Testiculaire*, Spermatic artery—*a. Thoracique humérale*, Acromial artery—*a. Trachto-cerebral*: see Cerebral arteries—*a. Trochanterienne*, Circumflex artery of the thigh—*a. Troisième des thoraciques*, Acromial artery—*a. Tympanique*, Auditory artery, external—*a. Urticale*: see Ciliary artery—*a. Vulvaire*, Pudic, external, artery.

ARTERIA, Artery—*a. Ad Cutem Abdominis*, see *Ad Cutem abdominis*, (arteria)—*a. Anonyma*, Innominata artery—*a. Aspera*, Trachea—*a. Cerebralis*, Carotid, internal—*a. Cervicalis*, Basillary artery—*a. Coronaria dextra*, Pyloric artery—*a. Crassa*, Aorta—*a. Cruralis*, Crural artery—*a. Externa Cubiti*, Radial artery—*a. Dorsalis metacarpi*, Metacarpal artery—*a. Dura matris media maxima*, Meningeal artery, middle—*a. Encephalica*, Carotid, internal—*a. Gastrica superior*, Coronary artery—*a. Ileo-colica*: see Colic arteries—*a. Iliaca interna*, Hypogastric artery—*a. Iliaca posterior*, Hypogastric artery—*a. Magna*, Aorta—*a. Magna pollicis*, Princeps pollicis—*a. Malleolaris externa*: see Tibial arteries—*a. Malleolaris interna*: see Tibial arteries—*a. Mammaria externa*, A. Thoracica externa inferior—*a. Maxima*, Aorta—*a. Media anastomotica*: see Colic arteries—*a. Meningeæ-media*, Meningeal artery, middle—*a. Musculacis femoris*, A. Profunda femoris—*a. Pharyngea suprema*, Pterygoid artery—*a. Profunda cerebri*: see Cerebral arteries—*a. Pudenda communis*, Pudic, internal, artery—*a. Pudica*, Pudic, internal, artery—*a. Ramulus ductus Pterygoidei*, Pterygoid artery—*a. Speno-epinosa*, Meningeal artery, middle—*a. Spinales*, Spinal arteries—*a. Spinalis*, A. Profunda humeri—*a. Sternalis*, Mammary, internal—*a. Supra-orbitalis*, Frontal artery—*a. Sylviana*: see Cerebral arteries—*a. Thoracica axillaris vel alaris*, Scapular artery, inferior—*a. Thoracica humeralis*, Acromial artery—*a. Transversalis colli*: see Cerebral arteries—*a. Transversalis humeri*, Scapular artery, superior—*a. Ulnaris*, Cubital artery—*a. Uterina hypogastrica*, Uterine artery—*a. Vasta posterior*, A. Profunda femoris.

ARTERIAL, Arteri'acus. A medicine prescribed in diseases of the windpipe. Also arterial.

ARTERIE ADIPO'SÆ. The arteries which secrete the fat about the kidneys are sometimes so called. They are ramifications of the capsular, diaphragmatic, renal, and spermatic arteries.

ARTERIE APOPLECTICÆ, Carotids—*a. Capitales*, Carotids—*a. Ciliares*, Ciliary arteries—*a. Corporis callosi cerebri*, Mesolobar arteries—*a. Jugulares*, Carotids—*a. Lethargicæ*, Carotids—*a. Mesolobica*, Mesolobar arteries—*a. Præparantes*, Spermatic arteries—*a. Somniferæ*, Carotids—*a. Soporales*, Carotids—*a. Soporarie*, Carotids—*a. Venosæ*, Pulmonary veins.

ARTERIAL, Arteri'acus, Arteri'ous. Belonging to arteries.

ARTERIAL BLOOD, (F.) Sang artériel. Red blood is so called because contained in the arteries. The pulmonary veins, however, also contain red blood: hence the name arterial veins, (F.) Veines artérielles, applied to them.

ARTERIAL CONSTITUTION, Plethora arteriosa.

ARTERIAL DUCT, Canalis arteri'ous, Ductus arteri'ous seu Botalli, (F.) Canal artériel, C. Pulmo-aortique, is the portion of the pulmonary artery which terminates in the aorta in the fœtus. When this duct is obliterated after birth, it is called Arteri'al Ligament, Ligamentum arteri'osum, (F.) Ligament artériel.

ARTERIAL SYSTEM includes all the arteries, from their origin in the heart to their termination in the organs. See Vascular System.

ARTERIALIZATION OF THE BLOOD, Hæmatisis.

ARTÉRIARCTIE, from ἀρτηρία, 'artery,' and αρκτο, 'I straiten.' Contraction of an artery.

ARTERIECTASIS, Aneurism.

ARTERIECTOP'IA, from ἀρτηρία, 'artery,' and εκτοπος, 'out of place.' Dislocation of an artery.

ARTERIEURYSMA, Aneurism.

ARTERITIS, Arteritis.

ARTERIODIALYSIS, see Aneurism.

ARTERIOG'RAPHY, Arteriogra'phia: from ἀρτηρία, 'artery,' and γραφή, 'a description.' A description of the arteries.

ARTERIOLOLA. A small artery.

ARTERIOLOGY, Arteriolog'ia; from ἀρτηρία, 'artery,' and λογος, 'a discourse.' A treatise on the arteries.

ARTERIO-PHLEBOTOMY, see Bloodletting.

ARTERIO-PITU'ITOUS. An epithet applied to vessels which creep along the interior of the nostrils.—Ruysh.

ARTERIORRHEXIS, see Aneurism.

ARTERIOS'ITAS, Arterios'ity, from arteria, 'an artery.' A condition of the blood in which it preserves in the veins the arterial character.—The opposite to Venositas.

ARTERIOSTAS SANGUINIS, Prædominium sanguinis arteriosi.

ARTÉRIOSTEIE, from ἀρτηρία, 'artery,' and οσσειον, 'a bone.' Ossification of an artery.—Piorry.

ARTERIOSTENOS'IS, from ἀρτηρία, 'artery,' and στενω, 'contracted.' Contraction or obliteration of an artery.

ARTERIOT'OMY, Arteriotom'ia, from ἀρτηρία, 'an artery,' and τομή, 'I cut.' This word has been used for the dissection of arteries. Most commonly, however, it means a surgical operation, which consists in opening an artery, to draw blood from it. Arteriotomy is chiefly used in inflammatory affections of the head, when the blood is generally obtained from the temporal artery. See Bloodletting.

ARTERITIS, Arteri'itis, Inflammat'io Arteri'arum, (F.) Artérite, Inflammation des artères; from ἀρτηρία, 'an artery,' and itis, a termination denoting inflammation. Inflammation of an ar-

tery. Inflammation of the inner coat of an artery is termed *Endo-arteritis*, or *Endonarteritis*; of the outer, *Exo-arteritis* or *Exarteritis*.

ARTERY, Arte'ria, (F.) Artère, from *anp*, 'air,' and *ῥησναι*, 'to preserve,' quasi, 'receptacle of air,' because the ancients believed that it contained air. They, at first, gave the name Artery to the trachea, *ἀρτηρία ρπαχσια*, because it is filled with air: and afterwards they used the same term for the arteries, properly so called, probably because they commonly found them empty in the dead body. We find, also, *ἀρτῆρες* to designate the arteries, called by the Latins *Venæ mican'tes pulsat'iles*. Arteries, with the moderns, signify the order of vessels, which arise from the two ventricles of the heart, and have valves only at their origin. They are cylindrical, firm, and elastic canals; of a yellowish white colour; little dilatable; easily lacerable; and formed, 1. Of an external, laminated, or areolar membrane, *Tu'nica externa seu adventit'ia seu cellula'ris, Vagina cellula'ris*, of a dense and close character. 2. Of a middle coat, *Tu'nica me'dia seu elastica*, composed of fibres, which do not, however, contract on the application of the galvanic stimulus, formed chiefly of elastic tissue, and also of smooth muscular fibres, and eminently elastic; and, 3. Of an inner coat, *Tu'nica in'tima seu glom'ra*, which is thin, diaphanous, reddish, and polished.

The use of the arteries is to carry the blood from the heart to the various parts of the system. It will be obvious, however, that they cannot all convey arterial blood. The pulmonary artery, for example, is destined to convey the venous blood to the lungs, there to be converted into arterial; whilst the pulmonary veins convey arterial blood back to the heart.

TABLE OF THE PRINCIPAL ARTERIES OF THE BODY.

All the arteries take their rise from the Pulmonary Artery, or the Aorta; and the names generally indicate the parts to which they are distributed.

I. ARTERIA PULMONALIS.

The Pulmonary Artery arises from the right ventricle, and soon divides into a right and left branch, one of which is distributed to each lung.

II. ARTERIA AORTA.

The Aorta arises from the left ventricle. It is the common trunk of the arteries of the body, and may be divided into five portions.

a. Arteries furnished by the Aorta at its origin.

1. A. Cardiaca or coronaria anterior.
2. A. Cardiaca or coronaria posterior.

b. Arteries furnished by the Aorta at its arch.

The arch of the Aorta gives off, to the left, two considerable trunks—the *Arteria carotidea primitiva*, and *A. subclavia*; and, to the right, a single trunk, which is larger—the *A. innominata*, or *Brachio-cephalica*, which divides into the *primitivæ carotid et subclavianæ*.

A. ARTERIA CAROTIDEA PRIMITIVA. Divides into A. Carotidea externa, and A. Carotidea interna.

Furnishes, 1. A. Thyroidea superior.

2. A. Lingualis, which gives off the A. dorsalis lingue and A. sublingualis.

3. A. facialis vel A. Maxillaris externa, which furnishes the A. palatina inferior, the A. submental, and A. coronaria superior and inferior.

4. A. occipitalis, which gives off the A. mastoidea posterior.

5. A. auricularis posterior, which gives off A. stylo-mastoidea.

6. A. pharyngea inferior.

a. A. Carotidea externa.

The external carotid ultimately divides into the temporal artery and internal maxillary.

1. A. Temporalis.

Furnishes A. transversalis faciei, A. auricularis anterior, and A. temporalis media.

2. A. Maxillaris interna.

Furnishes 13 branches, viz.—A. meningea media, A. dentaris inferior, A. temporalis profunda posterior, A. masseterina, A. pterygoidea, A. buccalis, A. temporalis profunda anterior, A. obovatus, A. suborbitalis, A. vidiana, A. pterygoidea or pharyngea superior, A. palatina superior, and A. sphenopalatina.

b. A. Carotidea interna.

Furnishes, 1. A. ophthalmica, which gives off A. lacrymalis, A. centralis retina, A. supraorbitalia vel superciliaris, A. ciliare posterior, A. ciliare longæ, A. muscularis superior et inferior, A. ethmoidalis posterior et anterior, A. palpebralis superior et inferior, A. nasalis, and A. frontalis. 2. A. communicans Willestii. 3. A. choroidæ. 4. A. cerebialis anterior. 5. A. cerebialis media.

B. ARTERIA SUBCLAVIA.

Furnishes, 1. A. vertebralis, which gives off A. spinalis anterior et posterior, A. cerebellosa inferior, and forms—by uniting itself with that of the opposite side—the A. basillaris, divided into A. cerebellosa superior and A. cerebellalis posterior. 2. A. thyroidea inferior, which gives off A. cervicalis ascendens. 3. A. mammaria interna, which gives off the A. mediastina anterior and A. diaphragmatica superior. 4. A. intercostalis superior. 5. A. cervicalis transversa. 6. A. scapularis superior. 7. A. cervicalis posterior vel profunda. Further on, the subclavian artery continues its progress under the name A. axillaris.

A. Axillaris.

Furnishes, 1. A. axillaris. 2. A. thoracica superior. 3. A. thoracica inferior vel longa vel mammaria externa. 4. A. scapularis inferior vel communis. 5. A. circumflexa posterior. 6. A. circumflexa anterior. Further on, the axillary artery continues under the name A. brachialis.

A. Brachialis.

Furnishes A. humeralis profunda vel collateralis externa. 2. A. collateralis interna. It afterwards divides into the radial and cubital arteries.

1. A. Radialis.

Gives off A. recurrens radialis, A. dorsalis carpi, A. dorsalis metacarpi, A. dorsalis pollicis, and terminates in forming the Arcus palmaris profundus.

2. A. Cubitalis.

Gives off A. recurrens cubitalis anterior and posterior; A. interossea anterior and posterior, which latter furnishes A. recurrens radialis posterior. It terminates in forming the superficial palmar arch, which gives off A. collaterales digitorum.

c. Arteries given off by the Aorta in the Thorax.

These arteries are,

1. A. Branchica, dextra et sinistra.
2. A. œsophagea (to the number of four, five, or six).
3. A. mediastina posteriores.
4. A. intercostales inferiores vel aortice (to the number of eight, nine, or ten).

d. Arteries furnished by the Aorta in the Abdomen.

These branches are,

1. The A. diaphragmatica vel phrenica, dextra et sinistra.

2. A. Cœliaca.

Which divides into three branches, 1. A. coronaria ventriculi. 2. A. Hepatica, which gives off A. pylorica, A. gastro-epiploica dextra, and A. cystica; and, lastly, the A. splenica, which gives off A. gastro-epiploica sinistra and Vasa brevia.

3. A. Mesenterica superior.

Which gives off at its convexity the A. colica dextra superior, media et inferior, and at its convex part from 15 to 20 Rami intestinales.

4. A. Mesenterica inferior.

Which gives off A. colica superior media, and inferior, and divides into A. hæmorrhoidales superiores.

5. The A. Capsulares media (to the number of two on each side).

6. A. Renales vel Emulgentes.

7. A. Spermaticæ.

8. A. Lumbares (to the number of four and five on each side).

e. Arteries resulting from the Bifurcation of the Aorta.

The Aorta, a little above its Bifurcation, gives off the *A. sacra media*, and divides into *A. iliaca primitiva*.

- | | | |
|-------------------------------|---|--|
| <i>A. Iliaca primitiva</i> | { | Divides into <i>A. Iliaca interna</i> and <i>A. Iliaca externa</i> . |
| | | Furnishes, 1. <i>A. ilio-lumbaris</i> . 2. <i>A. sacra lateralis</i> . 3. <i>A. glutea vel iliaca posterior</i> . 4. <i>A. umbilicalis</i> . 5. <i>A. vesicalis</i> . 6. <i>A. obturatoria</i> . 7. <i>A. hæmorrhoidalis media</i> . 8. <i>A. uterina</i> . 9. <i>A. vaginalis</i> . 10. <i>A. ischiatica</i> . 11. <i>A. pudenda interna</i> , which gives off the <i>A. hæmorrhoidales inferiores</i> . 12. <i>A. of the septum</i> . <i>A. transversa perinei</i> . <i>A. corporis cavernosi</i> , and <i>A. dorsalis penis</i> . |
| a. <i>A. Iliaca interna</i> | | Furnishes, 1. <i>A. epigastrica</i> , 2. <i>A. iliaca anterior vel circumflexa ili.</i> and is continued afterwards under the name of <i>Crural Artery</i> . |
| | | Furnishes, 1. <i>A. subcutanea abdominalis</i> . 2. <i>A. pubica superficialis et profunda</i> . 3. <i>A. muscularis superficialis</i> . 4. <i>A. muscularis profunda</i> , which gives off the <i>A. circumflexa externa</i> and <i>interna</i> , and the three Perforantes, distinguished into superior, middle, and inferior. Farther on, the crural artery continues under the name <i>A. Poplitea</i> . |
| b. <i>A. Iliaca externa</i> | { | Furnishes, 1. <i>A. Articulæres superiores, interni, medii, et externi</i> . 2. <i>A. Genicula</i> . 3. <i>A. Articulæres inferiores, interni et externi</i> . 4. <i>A. tibialis antica</i> , which, at the foot, takes the name <i>A. dorsalis tarsi</i> , and gives off the tarsal and metatarsal arteries. In the leg, the popliteal artery divides into the peroneal and posterior tibial. |
| A. Cruralis. | | |
| A. Poplitea. | { | Divides into <i>A. peronæa antica</i> and <i>A. peronæa postica</i> . |
| | | Divides into <i>A. plantaris interna</i> and <i>A. plantaris externa</i> . The latter, by anastomosing with the <i>A. dorsalis tarsi</i> , forms the plantar arch, whence arise <i>Rami superiores vel perforantes postici</i> , <i>R. Inferiores postici et antici</i> , which give off <i>Rami perforantes antici</i> . |
| 1. <i>A. Peronæa</i> | { | |
| 2. <i>A. Tibialis postica</i> | | |

ARTERY, ANGULAR, Facial artery.—a. Articular. Circumflex artery.—a. Brachiocephalic. Innominata arteria.—a. Central of the retina, Central artery of the retina.—a. Central of Zinn, Central artery of the retina.—a. Cephalic, Carotid.—a. Cerebral posterior, Vertebral.—a. Cervico-scapular, see Cervical arteries.—a. Coronary of the lips, Labial artery.—a. Crotaphitic, Temporal artery.—a. Fibular, Peroneal artery.—a. Gastric inferior, Gastro-epiploic artery.—a. Gastro-hepatic, see Gastro-epiploic artery.—a. Genital, Pudic (internal) artery.—a. Guttural inferior, Thyroideale A. inferior.—a. Guttural superior, Thyroideale A. superior.—a. Humeral, Brachial artery.—a. Iliac posterior, Gluteal artery.—a. Iliaco-muscular, Ileo-lumbar artery.—a. Labial, Facial artery.—a. Laryngeal superior, Thyroideale artery, superior.—a. Maxillary internal, Facial artery.—a. Median of the sacrum, Sacral artery, anterior.—a. Nasal, lateral, large, Spheno-palatine artery.—a. Palato-labial, Facial artery.—a. Periocephalic, Carotid (external).—a. Pharyngeal, superior, Pterygo-palatine artery.—a. Phrenic, Diaphragmatic artery.—a. Posterior, of the brain, see Cerebral arteries.—a. External scapular, Acromial artery.—a. Spinal, Meningeal artery, middle.—a. Subclavian right, Innominata arteria.—a. Subscapular, Scapular artery, inferior.—a. Superficial of the abdomen, Ad cutem abdominis (arteria).—a. Supramaxillary, Alveolar artery.—a. Suprarenal, Capsular artery.—a. Thoracic, internal, Mammary internal.—a. Urethro-bulbar, Transverse perineal artery.—a. Vesico-prostatic, Vesical artery.—a. Vidian, Pterygoid artery.

ARTETISCUS, from *artus*, 'a limb.' One who has lost a limb.

ARTEURYSMA, Aneurism.

ARTHANITA, from *artos*, 'bread,' the *Cyc-*

lamen or Sourdread. It was formerly made into ointment, *Unguentum Arthanitæ*, with many other substances, and was employed as a purgative, being rubbed on the abdomen.

ARTHANITA CYCLAMEN, Cyclamen.

ARTHETICA, Teucrium chamæpitys.

ARTHRAGRA, Gout.—a. Anomala, Gout, anomalous.—a. Genuina, Gout, regular.—a. Legitima, Gout, regular.—a. Normalis, Gout, regular.—a. Vera, Gout, regular.

ARTHRALGIA, Arthrodynia, Gout. See Lead rheumatism.

ARTHRELCO'SIS, from *αρθρον*, 'a joint,' and *ἐλκωσις*, 'ulceration.' Ulceration of a joint.

ARTHEMBOLE'SIS, same etymon as the next. The reduction of a fracture or luxation.

ARTHEMBOLUS, from *αρθρον*, 'a joint,' *εν*, 'in,' and *βαλλω*, 'I cast.' An ancient instrument used in the reduction of dislocations.

ARTHRETICA, Teucrium chamæpitys.

ARTHRITIC, *Arthrit'icus*, Gouty, from *αρθρον*, 'a joint.' (F.) *Arthritique*, *Goutteux*. That which relates to gout or arthritis, as *arthritic symptoms*, &c. See Podagra.

ARTHRITICUS VERUS, Gout.

ARTHRITIFUGUM, from *arthritis*, 'gout,' and *fugare*, 'to drive away.' A remedy that drives away gout. Heyden terms cold water, internally, the *arthritifugum magnum*.

ARTHRITIS, Gout, Arthrophlogosis, Arthrosis.—a. Aberrans, Gout (wandering).—a. Acuta, Gout (regular).—a. Arthrodynia, Rheumatism, chronic.—a. Asthenica, Gout (atonic).—a. Atonica, Gout (atonic).—a. Diaphragmatica, Angina Pectoris.—a. Erratica, Gout (wandering).—a. Hydrarthros, Hydrarthrus.—a. Inflammatoria, Gout (regular).—a. Juvenilis, see Rheumatism, acute.—a. Maxillaris, Siagonagra.—a. Nodosa, Gout (with nodosities).—a. Planetica, Gout (wandering).—a. Podagra, Gout.—a. Rheumatica, see Rheumatism, acute.—a. Rheumatismus, Rheumatism, acute.—a. Retrograda, Gout (retrograde).

ARTHRO'ACE, from *αρθρον*, 'a joint,' and *κακος*, 'bad.' Disease of the joints; and especially caries of the articular surfaces. *Spina ventosa*.

ARTHROCAE COXARUM, Coxarum morbus.

ARTHROCACOLOG'IA, from *arthroca-* according to Rust, a chronic disease of the joints; and *logos*, 'a description.' A treatise on chronic diseases of the joints.

ARTHROCARCINOMA, from *αρθρον*, 'a joint,' and *καρκινομα*, 'cancer.' Cancer of the joints.

ARTHROCHONDRI'TIS, from *αρθρον*, 'a joint,' *χονδρος*, 'a cartilage,' and *itis*, denoting inflammation. Inflammation of the cartilages and joints.

ARTHRO'DIA, from *αρθρον*, 'a joint.' *Adarticula'tio*, (F.) *Enfoncure*, *Enfonceure*. A moveable joint, formed by the head of a bone applied to the surface of a shallow socket, so that it can execute movements in every direction. *Arthrodium* is 'a small joint,' diminutive of *Arthrodia*.

ARTHRODYN'IA, *Arthronal'gia*, *Arthral'gia*, from *αρθρον*, 'a joint,' and *οδυνη*, 'pain.' Articular pain. Pain in the joints. See Rheumatism, chronic.

ARTHRODYNIA PODAGRICA, Gout.

ARTHROGRAPHY, *Arthrogaph'ia*, from *αρθρον*, 'a joint,' and *γραφω*, 'I write.' A description of the joints.

ARTHROGRYPO'SIS, from *αρθρον*, 'a joint,' and *γρυπω*, 'I incurvate.' Malposition of the joints, especially of the hands and feet.

ARTHROHYDRINE, Synovia.

ARTHROLOGY, *Arthrol'ogy*, from *αρθρον*, 'a joint,' and *λογος*, 'a description.' A description of the joints. The anatomy of the joints.

ARTHRORHOMBOLÉ, from *αρθρον*, and *βαλλω*, 'I cast.' Coaptation, reduction. Reduction of a luxated or fractured bone.

ARTHRORHOMENINGITIS, Meningarthrocace.

ARTHRON, 'a joint.' The ancients used the word *Arthron* for the articulation of bones with motion, in opposition to *Symphysis*, or articulation without motion.

ARTHRONALGIA, Arthrodynia.

ARTHRON'CUS, *Arthrophyma*, from *αρθρον*, 'a joint,' and *αγκος*, 'a swelling.' Tumefaction of a joint.

ARTHRONEMPYESIS, Arthropyosis.

ARTHROP'ATHY, *Arthropathi'a*, from *αρθρον*, 'a joint,' and *παθος*, 'affection.' A disease of the joints.

ARTHROPHLOGO'SIS, from *αρθρον*, 'a joint,' and *φλεγω*, 'I burn.' *Arthritis*, *Ostearthro'sis*. Inflammation of the joints.

ARTHROPHLOGOSIS SYNOVIALIS, Synovitis.

ARTHROPHYMA **ADENOCNDRUM**, see *Adenochondrius*.

ARTHROPYO'SIS, *Arthronempye'sis*, from *αρθρον*, 'a joint,' and *πυω*, 'pus.' Suppuration or abscess of the joints.

ARTHRO-RHEUMATISMUS, Rheumatism (acute).

ARTHRO'SIA, from *αρθρως*, 'I articulate.' *Arthritis*, (of some). Inflammation, mostly confined to the joints; severely painful; occasionally extending to the surrounding muscles. A genus of diseases in the Nosology of Good, including *Rheumatism*, *Gout*, *Articular inflammation*, *Joint-ache*, &c.

ARTHROSIA ACUTA, Rheumatism, acute—a. *Chronica*, Rheumatism, chronic—a. *Lumborum*, *Lumbago*—a. *Podagra*, *Gout*—a. *Podagra complicata*, *Gout* (retrograde)—a. *Podagra larvata*, *Gout* (atonic)—a. *Podagra regularis*, *Gout* (regular).

ARTHIROSIS, Articulation.

ARTHIROSPON'GUS, from *αρθρον*, 'a joint,' and *σπγγος*, 'a sponge.' A white, fungous tumour of the joints.

ARTHIROTRAU'MA, from *αρθρον*, 'a joint,' and *τραυμα*, 'a wound.' A wound of a joint.

ART'IA. According to some, this word is synonymous with *aprrpia*; others use it synonymously with *Trachia*.

ARTICHAUT, *Cynara scolymus*.

ARTICHOKE, *Cynara scolymus*.

ARTICLE, Articulation—a. of Death, see *Psychorages*.

ARTICOALUS, *Cynara scolymus*.

ARTICULAR, *Articula'ris*, (F.) *Articulaire*, from *artus*, 'a joint,' *articulus*, 'a small joint.' That which relates to the articulations—as the *articular capsules*, &c.

ARTICULAR ARTERIES OF THE ARM, Circumflex arteries of the arm.

ARTICULAR ARTERIES OF THE KNEE arise from the popliteal artery, and surround the tibio-femoral articulation. Although of a small size, they are important, as they furnish blood to the lower extremity after the operation for popliteal aneurism. They are distinguished into *superior* and *inferior*. The *superior articular arteries*, *popliteal articular arteries*, are commonly three in number; one of which is *internal*, another *external*, and another *middle*, the *æzygous articular*. The first, *Ramus anastomoticus magnus*, anastomoses by one branch with the external circumflex, and by another with the external superior articular. The second anastomoses with the external circumflex, the superior internal articular, and the inferior external articular; and the third is distributed within the joint. The *inferior articular arteries* are two in number: an

internal and *external*. The former anastomoses with the internal superior articular and the external inferior articular. The latter anastomoses with the recurrent branch of the anterior tibial, and the external superior articular. To each articular artery there is an *articular nerve*.

ARTICULAR FACETTES are the contiguous surfaces, by means of which the bones are articulated.

ARTICULAR PROCESSES, see *Vertebrae*.

ARTICULAR VEINS of the knee follow the same course as the arteries.

ARTICULATED, *Articula'tus*, (F.) *Articulé*, same etymon as *Articulation*. Provided with, or united by, articulations; as an 'articulated skeleton.' See *Skeleton*, articulated; and *Voice*, articulated.

ARTICULATIO, Articulation—a. *Artificialis*, *Pseudarthrosis*—a. *Notha*, *Pseudarthrosis*.

ARTICULA'TION, *Joint*, *Articula'tio*, *Arthro'sis*, *Asarthro'sis*, *Artic'ulus*, *Junctu'ra*, *Cola*, *Conjunctio*, *Nodus*, *Commissu'ra*, *Comp'ages*, *Syntaxis*, *Har'mus*, *Verticula*, *Vertic'ulus*, *Vertic'ulum*, (F.) *Articulation*, *Article*. By some of the French surgeons and anatomists, *article* is restricted more particularly to a movable articulation. Same etymon. The union of bones with each other, as well as the kind of union.

TABLE OF ARTICULATIONS.

Articulations are generally divided into *Diarthroses* or movable articulations, and *Synarthroses* or immovable.

Diarthroses.	{	1. Amphiarthrosis.
		2. Diarthrosis, orbicu- { Enarthrosis,
		lar vague. { Arthro'dia
Synarthroses.	{	3. Alternative or Ginglymus, which admits of varieties.
		1. Suture.
		2. Harmony.
		3. Gomphosis.
		4. Schindylesis.

The articulations are subject to a number of diseases, which are generally somewhat severe. These may be physical, as wounds, sprains, luxations, &c.; or they may be organic, as ankylosis, extraneous bodies, caries, rheumatism, gout, hyarthroses, arthropyosis, &c.

ARTICULATION means also the combination of letters which constitute words. See *Voice*.

ARTICULATION, FALSE, *Pseudarthro'sis*, *Artic'ulus falsus*, (F.) *A. fausse*, *A. accidentelle*, *A. contre nature*, *A. anormale*. A false joint, formed between fragments of bone, that have remained ununited; or between a luxated bone and the surrounding parts.

ARTICULATION SURNUMÉRAIRE, *Pseudarthrosis*—a. *en Chamrière*, *Ginglymus*—a. *de la Hanche*, *Coxo-femoral articulation*.

ARTICULI DIGITORUM MANUS, *Phalanges* of the fingers—a. *Digitum pedis*, *Phalanges* of the toes.

ARTICULO MORTIS, see *Psychorages*—a. *Spinalis*, *Semispinalis colli*.

ARTICULUS NOVUS, *Pseudarthrosis*—a. *Præternaturalis*, *Pseudarthrosis*—a. *Spurius*, *Pseudarthrosis*.

ARTIFICI'IAL, *Artificial'is*, (F.) *Artificiel*, from *ars*, *artis*, 'art,' and *facere*, 'to make.' That which is formed by art.

ARTIFICIAL EYES are usually made of enamel, and represent a sort of hollow hemisphere, which is applied beneath the eyelids, when the eye is lost.

ARTIFICIAL TEETH are made of ivory, porcelain, &c.

Pièces d'Anatomie Artificielles, are preparations of anatomy, modelled in wax, plaster, paper, &c.

ARTISCOCCUS LÆVIS, *Cynara scolymus*.

ARTIS'CUS, from *artos*, 'bread.' See Trochiscus. A troch of the shape of a small loaf. Also, and especially, a troch made of vipers.

ARTOCARPUS. The Bread-fruit Tree, (F.) *Jacquier*, from *artos*, 'bread,' and *karpos*, 'fruit.' A Polynesian tree, so called because the fruit, which is milky, and juicy, supplies the place of bread to the inhabitants. It grows to the height of 40 feet.

ARTOCARPUS INTEGRIFOLIA, Caoutchouc.

ARTOC'REAS, from *artos*, 'bread,' and *κρεας*, 'flesh.' A kind of nourishing food made of various aliments boiled together.—Galen.

ARTOG'ALA, from *artos*, 'bread,' and *γαλα*, 'milk.' An alimentary preparation of bread and milk. A ponicice.

ARTOM'ELLI, from *artos*, 'bread,' and *μελι*, 'honey.' A cataplasm of bread and honey.—Galen.

ARTUS, Membrum.

ARTYMA, Aroma, Condiment.

ARUM, A. maculatum, and A. triphyllum—a. Americanum betæ foliis, Dracontium foetidum.

ARUM DRACON'TIUM, *Arisæma draconitium*, *Green Dragon*; indigenous, has probably the same medical properties as A. triphyllum.

ARUM DRACON'CVLES, *A. polyphyllum*, *Draconculus polyphyllus*, *Colubrina Dracontia*, *Erva de Sancta Maria*, *Giflarus serpentina*, *Serpentina Gaillosum*, Family, Araceæ. See *Syst. Monoclea Polyandria*. The roots and leaves are very acrimonious. The plant resembles the A. maculatum in its properties.

ARUM ESCULENTUM, *Cala'dium esculentum*, *Taro*, *Kalo*. The foliage and roots possess acid qualities, which are dissipated by baking or boiling; in which form it is used as food by the people of Madeira, the Polynesians, &c.

ARUM MACULA'TUM, *Aron*, *Arum* (of the older writers), *A. vulgare*, *Cuckoo Pint*, (Prov.) *Cocky-baby*, *Barba Arvensis*, *Serpentina minor*, *Zin-giber Germanicum*, *Succedatis penis*, *Wake Robin*, *Priest's pintle*, (F.) *Gouet*, *Pied de Veau*, *Pied de lièvre*, *Picotin*, *Giron*, *Amidonnière*. The fresh root is stimulant internally. Dose, ℥j. of the dried root. Externally, it is very acrid. From the root of this Arum a starch is prepared, which is called *Portland Island Sago*, *Gerva serpentina*, *Cerusæa serpentina*, *Fecula ari maculati*.

ARUM, THREE-LEAVED, *Arum triphyllum*.

ARUM TRIPHYLLUM, *Arisæma atrorubens*, *Three-leaved arum*, (F.) *Pied de Veau triphyllus*, *Indian Turnip*, *Dragon Root*, *Dragon Turnip*, *Pepper Turnip*. This plant grows all over the United States, and is received into the Pharmacopœia under the title *Arum*. The recent root, or *Cormus*—*Arum*, (Ph. U. S.)—is very acrimonious, and has been employed in asthma, croup, and whooping-cough. Boiled in lard, it has been used in tinea capitis, and in milk in consumption.

ARUM VIRGINICUM, *Peltandra Virginia*—a. *Vulgare*, A. maculatum—a. Water, *Calla palustris*.

ARUMARI, Caramata.

ARUNDO BAMBOO, Bamboo—a. *Brachii major*, *Ulna*—a. *Brachii minor*, *Radius*—a. *Indica*, *Sagittarium alexipharmacum*—a. *Major*, *Tibia*—a. *Minor*, *Fibula*—a. *Saccharifera*, see *Saccharum*.

ARVA, Ava.

ARVUM, Vulva—a. Naturæ, Uterus.

ARY-ARYTENOIDÆUS, *Arytenoidæus*—a. Epiglotticus, *Aryteno-epiglotticus*.

ARYT'ENA, *apraiva*, 'a ladle.' Hence,

ARYT'ENO-EPIGLOT'TICUS, *Aryt'eno-epiglottidæus*, *Ary-epiglotticus*. That which belongs to the arytenoid cartilages and epiglottis.

Winslow gives this name to small, fleshy fasciculi, which are attached, at one extremity, to the arytenoid cartilages, and, by the other, to the free edge of the epiglottis. These fibres do not always exist. They form part of the arytenoid muscle of modern anatomists.

ARYT'ENOID, *Aryt'enoïdes*, *Arytenoidæus*, from *apraiva*, 'a ladle,' and *eidos*, 'shape.' Ladle-shaped.

ARYTENOID CAR'TILAGES, *Cartilag'ines arytenoïdes* seu *gutturales* seu *gutturinæ* seu *gutturiformes* seu *triq'uetræ*, *Guttur'ina*, are two cartilages of the larynx, situate posteriorly above the cricoid, which, by approximation, diminish the aperture of the glottis. Their upper extremities or cornua are turned towards each other, and are now and then found loose, in the form of appendices, which are considered, by some, as distinct cartilages, and termed *coniform* or *tuberculated Cartilages*, *Cornic'ula Laryngis*.

ARYTENOID GLANDS, *Gland'ula Arytenoidæa*, are small, glandular, whitish bodies, situate anterior to the A. cartilages. They pour out a mucous fluid to lubricate the larynx.

ARYTENOIDÆUS, (F.) *Arytenoïdien*. A small muscle, which passes from one arytenoid cartilage to the other, by its contraction brings them together, and diminishes the aperture of the glottis. Winslow divided the muscle into three portions;—the *Arytenoidæus transversus*, or *Ary-arytenoidæus*, and two *Arytenoidæi obliqui*.

ARYTHM, *Aryth'mia*, from *a*, privative, and *ρυθμος*, 'rhythm,' 'measure.' Irregularity. This word is applied chiefly to the pulse.

ASA, Asafetida. See Assa—a. Dulcis, Benjamin.

ASAFÆTI'DA, *Assafetida* (Ph. U. S.), *Assafetida*, *Stercus diaboli*, *Cibus Deorum*, *Asa*, *Devil's dung*, *Food of the Gods*. A gum-resin—the concrete juice of *Ferula Assafetida*, *Narthex Assafetida*. Order, Umbellifera. It is in small masses of a whitish, reddish, and violet hue, adhering together. Taste bitter and subacid; smell insupportably alliaceous. The Asiatics use it regularly as a condiment.

Its medical properties are antispasmodic, stimulant, and anthelmintic. Dose, gr. v to xx, in pill.

ASAGRÆA OFFICINALIS, see *Veratrina*.

AS'APES, *Asep'ton*, properly, not putrid, from *a*, privative, and *σᾶω*, or *σῴω*, 'I corrupt.' A term applied by Hippocrates to the spruæ, or to other matters evacuated, which do not give signs of coction.

ASAPH'ATUM, from *a*, privative, and *σαφης*, 'clear.' This term has been applied to collections in the sebaceous follicles of the skin, which may be pressed out like little worms, with a black head. See *Aene*.

ASAPHI'A, from *a*, privative, and *σαφης*, 'clear.' *Dyspho'nia immodulata palati'na*, *Parapho'nia gutturalis seu palati'na*. Defective articulation, dependent upon diseased palate.—Hippocrates, Vogel.

ASARABACCA, *Asarum*—a. Broad-leaved, *Asarum Canadense*.

ASAR'CON, from *a*, priv., and *σαρξ*, 'flesh.' Devoid of flesh. Aristotle uses the term for the head when it is but little fleshy, compared with the chest and abdomen.

ASARET, *Asarum*—a. *du Canada*, *Asarum Canadense*.

ASARITES, from *ασαρων*, 'the asarum.' A diuretic wine, of which asarum was an ingredient.—Dioscorides.

AS'ARUM, from *a*, privative, and *σᾶειν*, 'to adorn,' because not admitted into the ancient coronal wreaths [?], *As'arum Europa'um* seu

officina'lis, *Nardus Monta'na* seu *Rust'ica*, *As'arum*, (F.) *Asaret* ou *Cabaret*, *Azarum Cabaret*, *Oreille d'homme*, *Oreillette*, *Givard-Rousin*, *Nard Sauvage*. Order, *Aristolochiaceae*, *Sex. Syst.* *Dodecandria Monogynia*. The plant, used in medicine, is the *As'arum Europa'um*, *Asarabac'ca*, and of this the leaves. They are emetic, cathartic, and errhine, but are hardly ever employed, except for the last purpose.

ASARUM CANADENSE, *A. Carolinia'num*, *Canada Snake-root*, *Wild Ginger*, *Colt's Foot*, *Broad-leaf Asarabacca*, *Indian Ginger*, *Heart Snake-root*, (F.) *Asaret du Canada*. The root *As'arum*, (Ph. U. S.), is used as a substitute for ginger, and is said to act as a warm stimulant and diaphoretic.

ASARUM CAROLINIANUM, *A. Canadense*—*a. Europæum*, see *Asarum*—*a. Hypocistis*, *Cytinus hypocistis*—*a. Officinale*, see *Asarum*.

ASBESTOS SCALL, see *Eczema* of the hairy scalp.

ASCAIN, MINERAL WATERS OF. *Ascain* is a village, situate about a league from St. Jean-de-Luz, in France. The water is a cold chalybeate.

ASCARDAMYC'TES, from *a*, privative, and *καρδάμυξ*, 'I twinkle the eyes.' One who stares with fixed eyes, without moving the eyelids.—*Hippocrates*.

ASCARICIDA ANTHELMINTICA, *Vernonia anthelmintica*.

ASCARIDE LUMBRICOÏDE, *Ascaris lumbricoides*—*a. Vermiculairis*, *Ascaris vermicularis*.

ASCARIS, pl. *ASCARIDES*, from *ασκαρίζω*, 'I leap.' A genus of intestinal worms, characterized by a long, cylindrical body, extenuated at the extremities; and having a mouth furnished with three tubercles, from which a very short tube is sometimes seen issuing. Formerly, there were reckoned two varieties of the *Ascaris*—the *As'caris lumbricoïdes* seu *gigas hom'inis*, *Lumbrici'ens*, *L. teres hom'inis*, *Scolex*, (F.) *Lombricoïde*, *Ascaride lombricoïde*, *Lombric*, or long round worm; and the *As'caris Vermiculairis*—the *Ascaris* proper—the *thread worm* or *maw worm*. The former is alone included under the genus, at present—a new genus having been formed of the *A. vermicularis*, under the name *Oxyuris*. It is the *Oxyu'ris* seu *Fusa'ria vermiculairis*, (F.) *Ascaride*, *A. vermiculairis*, *Oxyure vermiculairis*.

A new species of entozoa has been found by Dr. Bellingham, the *As'caris ala'ta*.

ASCARIS ALATA, see *Ascaris*—*a. Gigas hominis*, see *Ascaris*—*a. Lumbricoides*, see *Ascaris*—*a. Renalis*, *Strongylus gigas*—*a. Trichuria*, *Trichocephalus*—*a. Vermicularis*, see *Ascaris*—*a. Visceralis*, *Strongylus gigas*.

AS'CELES, *As'celes*, *Carens cru'ribus*, from *a*, privative, and *ακλος*, 'a leg.' One who has no legs.

ASCELLA, *Axilla*.

ASCEN'DENS, from *ascendere*, (*ad* and *soan-dere*), 'to ascend.' (F.) *Ascendant*. Parts are thus called, which are supposed to arise in a region lower than that where they terminate. Thus, *Aorta ascendens* is the aorta from its origin to the arch: *Vena cava ascendens*, the large vein which carries the blood from the inferior parts to the heart: *Obliquus ascendens* (*muscle*), the lesser oblique muscle of the abdomen, &c.

ASCEN'SUS MORBI. The period of increase of a disease.

ASCHERSONIAN VESICLES, see *Vesicles*, *Aschersonian*.

ASCESIS, *Exercise*.

ASCHILL, *Scilla*.

ASCHISTODACTYLUS, *Syndactylus*, from *a*, privative, *σχιστος*, 'cleft,' and *δακτυλος*, 'a

finger.' A monster whose fingers are not separated from one another.—*Gurif*.

AS'CIA (an axe), *Azinè*, *Scepar'nos*, *Dol'abra*, *Fas'cia spir'al'is*. Name of a bandage mentioned by the ancients, and figured by Scultetus, in the shape of an axe or hatchet.—*Galen*. See *Doloure*.

ASCILLA, *Axilla*.

ASCITE, *Ascites*—*a. Péritonéale*, *Ascites*.

ASCIT'ES, from *ακος*, 'a leathern sack, a large belly'—*Aski'tes*, *Hydroc'elè Péritonéale*, *Hydrops Abdom'inis* seu *Ascites*, *Hydrogaster*, *Hydroperitone'um*, *Hydroc'elia*, *Hydret'rum*, *Asci'tes*, *Cutlich'ysis*, *Dropsy of the lower belly*, *Dropsy of the Péritoné'um*, (F.) *Ascite*, *A. péritonéale*, *Hydro-péritonie*, *Hydropisie du Bas-ventre ou du péritoné*. A collection of serous fluid in the abdomen. *Ascites* proper is dropsy of the peritoneum; and is characterized by increased size of the abdomen, by fluctuation and the general signs of dropsy. It is rarely a primary disease; but is always dangerous, and but little susceptible of cure. Most generally, it is owing to obstructed circulation in some of the viscera, or to excitement of the vessels of the abdominal organs. The treatment is essentially the same as that of other dropsies. Paracentesis, when had recourse to, can only be regarded as a palliative.

Dropsy of the peritoneum may also be sacculated or in cysts, and occasionally the fluid accumulates exterior to the peritoneum, *Hydropigast'rium*. When in cysts it is termed *Hydrocysts'is*, *Hydrops abdom'inis sacca'tus* seu *cyst'icus* and *Asci'tes sacca'tus*.

ASCITES HEPATO-CYSTICUS, *Turgescencia vesicular' felica*—*a. Ovarii*, *Hydrops ovarii*—*a. Purulentus*, *Pyoc'elia*—*a. Saccatus*, see *Ascites*, *Hydroarion*, and *Hydrops ovarii*.

ASCIT'IC, *Asci'ticus*, (F.) *Ascitique*. Relating or appertaining to, or affected with, *ascites*.

ASCLEPIADÆ, *Asclepi'ades*, from *ασκληπιος*, 'Esculapius.' The priest physicians, who served in the ancient temples of Esculapius, and who took their name from being his descendants.

ASCLEPIADE, *Asclepias vincetoxicum*.

ASCLEPIAS ALBA, *A. vincetoxicum*—*a. Apocynum*, *A. Syriaca*—*a. Asthmatica*, *Tylophora asthmatica*—*a. Cornuti*, *A. Syriaca*—*a. Crispa*, *Gomphocarpus crispus*.

ASCLEPIAS CURASSAV'ICA, *Bastard Ipecacuanha*, *Redhead*, *Bloodweed*. The leaves are emetic in the dose of one or two scruples. It is the *Ipecacuanha blanc* of St. Domingo.

ASCLEPIAS DECUM'BENS; the root. Escharotic, cathartic, sudorific, diuretic.

ASCLEPIAS, FLESH-COLOURED, *A. Incarnata*—*a. Gigantea*, *Mudar*.

ASCLE'PIAS INCARNA'TA, *A. pulchra*, *Flesh-coloured* or *swamp asclepias* or *silkweed*. The root of this plant, which grows in all parts of the United States, and is official in the Ph. U. S., has the same virtues as *A. Syriaca*, and *A. tuberosa*.

ASCLEPIAS OBOVATA, *A. Syriaca*.

ASCLEPIAS PROC'ERA [?], *Beidelossar*, *Beidelcar*. An Egyptian plant, the leaves of which are made into a plaster, and applied to indolent tumours. The milky juice is caustic, and is used as such.

ASCLEPIAS PSEUDOSARSA, *Hemidesmus Indicus*—*a. Pubescens*, *A. Syriaca*—*a. Pulchra*, *A. incarnata*.

ASCLEPIAS SYRIACA, *A. pubes'cens* seu *apoc'y'num* seu *obovata* seu *tomento'sa* seu *Cornu'ti*, *Common Silkweed*, *Milk Weed*, (F.) *Herbe à la houette*. The cortical part of the root, official in the Ph. U. S., has been given, in powder, in

asthmatic and pulmonic affections in general, and, it is said, with success.

ASCLEPIAS SULLIVANII, *Smooth Milkweed*, *Silkweed*: indigenous, possesses the same virtues as the next.

ASCLEPIAS TOMENTOSA, *A. Syriaca*.

ASCLEPIAS TUBEROSA, *Butterfly Weed*, *Pleury Root*, *Flux Root*, *Wind Root*, *Wind Weed*, *White Root*, *Orange Swallow Root*, *Silk Weed*, *Canada Root*, *Orange Apocynum*, *Tuberous Rooted Swallow Root*. Said to have been first recommended by the Asclepiades. In Virginia and the Carolinas, the root of this plant—official in the Ph. U. S.—has been long celebrated as a remedy in pneumonic affections. It is sudorific, and the powder acts as a mild purgative. Its chief powers are said to be expectorant, diaphoretic, and febrifuge. It is occasionally given to relieve pains of the stomach from flatulency and indigestion.

ASCLEPIAS VINETOXICUM, *A. Alba*, *Cynanchum Vincetoxicum*, *Vincetoxicum*, *V. Officinale*, *Hirundinaria*, *Apocynum Novæ Angliæ hirsutum*, &c., *Swallow-Wort*, *White Swallow-Wort*, (*F.*) *Asclepiade*, *Dompthe-venin*.

The root is said to be stimulant, diuretic, and emmenagogue, but is hardly ever used.

ASCLEPIASMUS, *Hæmorrhoids*.

ASCLITES, *Ascites*.

ASCO'MA, from *ασκος*, 'a bottle.' The eminence of the pubes at the period of puberty in females.—Rufus of Ephesus.

ASE, *Anxiety*.

ASELLI, *Onisci aselli*.

ASELLUS, *Oniscus*.

ASE'MA CRISIS, *κρίσις ασμα*; from *a*, privative, and *σμα*, 'a sign.' A crisis occurring unexpectedly and without the ordinary precursory signs.

ASEPTON, *Asapes*.

ASH, BITTER, *Quassia*—*a. Blue*, *Fraxinus quadrangulata*—*a. Mountain*, *Sorbus acuparia*—*a. Mountain*, *American*, *Sorbus Americana*—*a. Prickly*, *Aralia spinosa*, *Xanthoxylum clava Herculis*—*a. Prickly*, shrubby, *Xanthoxylum fraxineum*—*a. Stinking*, *Ptelea trifoliata*—*a. Tree*, *Fraxinus excelsior*—*a. White*, *Fraxinus Americana*.

ASIALORRHŒ'A, (*F.*) *Asialorrhæe*, from *a*, priv., *σialon*, 'saliva,' and *ρρω*, 'to flow.' Diminution in the flow of saliva.

ASIMINA TRILOBA, see *Carica papaya*.

ASIT'IA, from *a*, privative, and *στρος*, 'food.' Abstinence from food. Want of appetite,—*Fastingium ciborum*, *Apocleitæ*.

ASIUS LAPIS, *Assius Lapis*.

ASJAGAN, *As'jogan*. An Indian tree, the juice of whose leaves, mixed with powdered cumin seeds, is employed in India in colic.

ASJOGAM, *Asjagan*.

ASKELES, *Asceles*.

ASKER, *Eschar*.

ASKITES, *Ascites*.

ASO'DES, *Asso'des*, from *ασν*, 'disgust,' 'satiety.' A fever accompanied with anxiety and nausea; *Febris aso'des* vel *aso'des*.

ASPALASOMUS, from *ασπαλαξ*, 'a mole,' and *σμα*, 'a body.' A genus of monsters in which there is imperfect development of the eyes.—I. G. St. Hilaire. Also, a malformation, in which the fissure and eventration extend chiefly upon the lower part of the abdomen; the urinary apparatus, genitals and rectum opening externally by three distinct orifices.—Vogel.

ASPALTUM, *Asphaltum*.

ASPARAGINE, see *Asparagus*.

ASPARAGUS, *Asπαργος*, *Asparagus officinalis*, *Common Asparagus*, *Sparagus*, *Speragus*, *Sparrow Grass*, *Grass*, *Sperage*. Ord., *Aspho-*

delem. Ser. Syst. Hexandria Monogynia. *Asparagi officinalis Turionnes*, (*F.*) *Asperge*. The fresh roots are diuretic, perhaps owing to the immediate crystallizable principle, *Asparagine*, which is said to be sedative in the dose of a few grains. The young shoots are a well known and esteemed vegetable diet. They communicate a peculiar odour to the urine. A syrup made of the young shoots and an extract of the roots has been recommended as a sedative in heart affections.

ASPA'SIA. A ball of wood soaked in an infusion of galls, and used by females for constricting the vagina.

ASPEN, AMERICAN, *Populus tremuloides*—*a. European*, *Populus tremula*.

ASPERS ARTERIA, *Trachea*.

ASPERGE, *Asparagus*.

ASPERITAS ARTERIÆ ASPERÆ, *Rau-cedo*.

ASPERITÉ DES PAUPIÈRES, *Trachoma*.

ASPERITUDU, *Trachoma*.

ASPERITY, *Asperities*, roughness. *Asperities* are inequalities on the surfaces of bones, which often serve for the insertion of fibrous organs.

ASPERMATIA, *Aspermatismus*.

ASPERMATISMUS, *Aspermia*, *Aspermia*, from *a*, privative, and *σπερμα*, 'sperm.' Reflux of sperm from the urethra into the bladder, during the venereal orgasm.

ASPERMIA, *Aspermatismus*.

ASERSIO, *Catapasma*, *Fomentation*.

ASPERSION, *Aspercio*, from *aspergere* (ad and *spargere*), 'to sprinkle,' (*F.*) *Arrosionem*. Act of sprinkling or pouring a liquid *guttatim* over a wound, ulcer, &c.

ASPERULA, *Galium aparine*.

ASPER'ULA ODORA'TA, (diminutive of *asper*, 'rough,' the seeds being rough), *Galium odoratum*, *Matrieyl'va*, *Hepatica stellata*, (*F.*) *Asperule odorante* ou *Muguet des bois*, *Hépatique étoilée*. Ord., *Rubiaceæ*. Ser. Syst. *Tetrandria Monogynia*. Sweet-scented Woodroof. Said to be diuretic, deobstruent, tonic, and vulnerary.

ASPERULE ODORANTE, *Asperula odorata*.

ASPHALTI'TES, *Nephrit'es*, *Nephrit'is*, *Prima Vertebra lumbal'ria*, same etymon as asphaltum. A name given by some to the last lumbar vertebra.—Gorræus.

ASPHAL'TUM, *Nep'ta*, *Arsal'tos*, *Asphaltum*, from *ασφαλτεω*, 'to strengthen,' being used as a cement. [?] With the Greeks, this word signified any kind of bitumen. It is now restricted chiefly to the BITUMEN OF JUDE'A, *B. Judæicum*, *A. sol'idum*, *Jews' Pitch*, *Karabé* of Sodom, (*F.*) *Asphalte*, *Bitume de Judée*, *Pois minérale scorifique*, *Baume de momie* ou *des funérailles*. It is solid, friable, vitreous, black, shining, inflammable, and of a fetid smell. An oil is obtained from it by distillation. It enters into the composition of certain ointments and plasters.

It is collected on the surface of the water of the Dead Sea or Lake Asphaltites, in Judæa.

ASPHARINE, *Galium aparine*.

ASPHOD'ELUS, *Asphodelos*, *A. Ramosus* seu *Albus* seu *Maris*, *Has'tula Regis*, (*F.*) *Lis asphodelæ*. The bulbs of this southern European plant have an acrimony which they lose in boiling water. They contain a fecula with which bread has been made, and have been considered diuretic. They have been used as a succedaneum for the squill.

ASPHYCTOS, *Pulseless*.

ASPHYX'IA, from *a*, priv., and *σφξίς*, 'pulse,' *Defectus Pulsus*, *Acrotismus*, *Sidera'tio*, *Sydera'tio*, (*F.*) *Anémotosis*. For a long time, *Asphyxia* was confined to the sense of 'suspension of circulation or syncope,' *Pulselessness*. It now

generally means *suspended animation*, produced by the nonconversion of the venous blood of the lungs into arterial.—*Apnœa*, *Apnœus'tia*, *Apnœophy'ria*, *Anhæmato'sia*, *Ecl'ysis pneumo-cardi'ca*, *Anozœmie*, *Hypoœmie*,—*Piorry*. Owing to the supply of air being cut off, the unchanged venous blood of the pulmonary artery passes into the minute radicles of the pulmonary veins, and their peculiar excitability requiring arterial blood to excite them, stagnation takes place in the pulmonary radicles, and death occurs chiefly from this cause,—not owing to venous blood being distributed through the system, and 'poisoning' it, as was the idea of Bichat. *Curus asphy'cia*, *Mors appa'rens*, *Mors putati'va*, *Pseudothan'atos*, *Apparent death*, (F.) *Mort appa'rente*, is characterized by suspension of respiration, of the cerebral functions, &c. Several varieties of asphyxia have been designated.

1. ASPHYX'IA OF THE NEW-BORN, *A. neonato'rum*. This is often dependent upon the feeble condition of the infant, not permitting respiration to be established.

2. ASPHYX'IA BY NOXIOUS INHALATION, or inhalation of gases, some of which cause death by producing a spasmodic closure of the glottis; others by the want of oxygen; and others are positively deleterious or poisonous.

3. ASPHYX'IA BY STRANGULATION, or *Suffoca'tion*; produced by mechanical impediment to respiration, as in hanging.

4. ASPHYX'IA BY SUBMERSION, *A. by drowning*, *A. immerso'rum*, *Angiairhydrie* (Piorry), as occurs in the drowned, who perish in consequence of the medium in which they are plunged, being unfit for respiration. See Submersion.

Mr. Chevalier has used the term *Asphy'cia Idiopath'ica*, for fatal syncope owing to relaxation of the heart. See Suffocation.

ASPHYX'IA IMMERSORUM, *A. by submersion*—*a. Local*:—see *Grangrene*—*a. Neonatorum*, *A. of the new-born*—*a. Pestilenta*:—see *Cholera*—*a. Pestilential*:—see *Cholera*.

ASPHYX'IAL, *Asphy'ctic*. Relating to asphyxia—as '*asphyxial* phenomena.'

ASPHYXIE DES PARTIES, *Gangrene*—*a. Lente des nouveau-nés*, Induration of the cellular tissue.

ASPHYX'IED, *Asphy'ctiated*, *Asphyxia'tus*, same etymon. In a state of asphyxia.

ASPIC, *Aspis*, *Lavandula spica*.

ASPIDISCOS, *Sphincter ani externus*.

ASPID'IUM ATHAMANTICUM, from *ασπίδιον*, 'a small shield,' diminutive of *ασπίς*, 'a shield.' A South African fern, *Ord.* Filices, which is possessed of anthelmintic properties. Its caudex, in the form of powder, infusion, or electuary, has been found excellent in helminthiasis, and especially in tapeworm.

ASPIDIUM CORIACEUM, *Calagualæ radix*—*a. Depastum*, *Polypodium filix mas*—*a. Discolor*, see *Calagualæ radix*—*a. Erosum*, *Polypodium filix mas*—*a. Filix fœmina*, *Asplenium filix fœmina*—*a. Ferrugineum*, see *Calagualæ radix*—*a. Filix mas*, *Polypodium filix mas*.

ASPIRATIO, *Inspiration*.

ASPIR'ATION, *Aspira'tio*, *Aspira'tio*, from *aspirare* (*ad* and *spirare*), 'to breathe.' The French sometimes use the term synonymously with inspiration. It also means the act of attracting or sucking like a pump. Imbibition. Also, the pronunciation of a vowel with a full breath.

ASPIS, *aspis*. A name given by the ancients to a venomous serpent—the *Egyptian viper* of Læcæpède, (F.) *Aspis*. Its bite is very dangerous, and it is supposed to have been the reptile which Cleopatra used for her destruction.

ASPLENIUM, *Asple'num*, from *a. priv.*, and *σπλην*, 'the spleen.' *Spleenwort*, *Miltwaste*.

ASPLENIUM ACREUM, *A. ceterach*.

ASPLENIUM CETERACH, *A. ac'reum* seu *latifolium*, *Gymnogram'mè ceterach*, *Doradillo*, *Blechnum squamosum*, *Scolopend'ria*, *Athy'rium*, *Ceterach officina'rum* seu *Canarien'sis*, *Grammitis ceterach* seu *au'rea*, *Gymnopteris ceterach*, *Vittaria ceterach*, (F.) *Doradille*. *Ord.* Filices. Supposed to be subastringent and mucilaginous, and has been recommended as a pectoral. It has also been given in calculous cases.

ASPLENIUM FILIX FÆMINA, *Polypod'rium filix fœmina* seu *mollè* seu *dentalium* seu *incisum* seu *trif'idum*, *Aspidium filix fœmina*, *Athy'rium filix fœmina* seu *mollè* seu *ovatum* seu *trif'idum*, *Pteris palustris*, *Female fern*, *Spleenwort*, (F.) *Fougère femelle*. The root of this plant resembles that of the male fern, and is said to possess similar anthelmintic virtues. The name *female fern* is also given to *Pteris aquilina*.

ASPLENIUM LATIFOLIUM, *A. ceterach*—*a. Murale*, *A. ruta*—*a. Obtusum*, *A. ruta muraria*.

ASPLENIUM RUTA MURARIA, *A. murale* seu *obtusum*, *Paronychia*, *Phyllitis ruta muraria*, *Scolopend'rium ruta muraria*, *Wallrue*, *White Maidenhair*, *Tentwort*, *Adiantum album*, *Ruta muraria*, *Salvia Vitæ*, (F.) *Rue des murailles*, *Sauve-vie*. Used in the same cases as the last.

ASPLENIUM SCOLOPENDRIUM, *Scolopend'rium officina'rum* seu *lingua* seu *phyllitis* seu *vulgarè*, *Scolopend'ra*, *Scolopend'ria*, *Hart's Tongue*, *Spleenwort*, *Phyllitis*, *Lingua cervina*, *Blechnum lignifolium*, (F.) *Scolopendre*, *Langue de cerf*. Properties like the last.

ASPLENIUM TRICHOMANOIDES, *A. Trichomanes*, *Phyllitis rotundifolia*, *Calyphyl'ium*, *Trichomanes*, *T. crenata*, *Adiantum rubrum*, *Common Maidenhair*, *Polytrichum commune*, (F.) *Polytric*. Properties like the last.

ASPLENIUM, *Asplenium*.

ASPREDO, *Trachoma*—*a. Miliacea*, *Miliary fever*.

ASPRELE, *Hippuris vulgaris*.

ASPRITUDO, *Trachoma*.

ASSABA. A Guinea shrub, whose leaves are considered capable of dispersing buboes.

ASSACOU, *Hura Brasiliensis*.

ASSA DOUX, *Benjamin*—*a. Dulcis*, *Benjamin*—*a. Odorata*, *Benjamin*.

ASSAFETIDA, *Asnafetida*.

ASSAFETIDA, *Asnafetida*.

ASSAIERET. A compound of bitter, stomachic, and purgative medicines in the form of pill.—*Avicenna*.

ASSAINISEMENT (F.), from *ad*, 'to,' and *sanare*, 'to make healthy.' The act of rendering healthy, as by the draining of marshes, the disinfection of the air, &c.

ASSAISONNEMENT, *Condiment*.

ASSAKUR, *Saccharum*.

ASSALA, see *Myristica moschata*.

ASSARTHROSIS, *Articulation*.

ASSA'TIO, *Opt'e'sis*. The boiling of food or medicines in their own juice, without the addition of any liquid. Various kinds of cooking by heat.—*Galen*.

ASELLA, *Axilla*.

AS'SERAC, *Assis*. A preparation of opium, or of some narcotic, used by the Turks as an excitant.

ASSERECULUM, *Splint*.

ASSERVATION, *Conservation*.

ASSES' MILK, see *Milk*, *asses'*.

ASSES' MILK, *ARTIFICIAL*, see *Milk*, *asses'*.

AS'SIDENS, from *ad*, 'to,' and *sedere*, 'to be seated.' That which accompanies or is concomitant. An epithet applied to the accessory symptoms, *Assidentia signa*, and general phenomena of disease.

ASSIDENTIA SIGNA, see *Assidens*.

ASSIMILABLE, *Assimilabilis*. Same etymon as Assimilation. That which is susceptible of assimilation.

ASSIMILATING, *Assimilatory*, *Assimilatorius*, (F.) *Assimilateur*. Same etymon as the next. That which effects assimilation, as an *assimilating* or *assimilatory* organ.

ASSIMILATION, *Assimilatio*, *Similatio*, *Appropriatio*, *Exomoiosis*, *Homoiosis*, *Threpsis*, *Threpticis*; from *assimilare*, (ad and similare,) 'to render similar.' The act by which living bodies appropriate and transform into their own substance matters with which they may be placed in contact. The reverse action—that of separating and eliminating matters already assimilated, in the act of nutrition—is termed, by the French, *Déassimilation*.

ASSIS, *Asserac*.

ASSIUS LAPIS, *Asius Lapis*. A sort of stone or earth, found near the town of Assa in the Troad, which had the property of destroying proud flesh.

ASSOCIATED MOVEMENTS, see *Instinctive*.

ASSODES, *Asodes*.

ASSOLPINNANT, *Somniferous*.

ASSOUPISSEMENT, *Somnolency*.

ASSOURON, see *Myrtus Pimenta*.

ASSUETUDO, *Habit*.

ASSULA, *Splint*.

ASSULTUS, *Attack*.

ASSUMPTIO, *Prehension*.

ASTACI FLUVIATILIS CONCREMENTA, *Cancrorum chelæ*.

ASTACUS FLUVIATILIS, *Crab*.

ASTAKILLOS, *Araneum ulcus*.

ASTARZOF. An ointment, composed of litharge, frog's spawn, &c. Also, camphor, dissolved in rose water.—*Paracelsus*.

ASTASIA, *Dysphoria*.

ASTER ATTICUS, *Bubonium*.

ASTER CORDIFOLIUS, *Heart-leaved Aster*, *A. Punicæus*, *Rough-stemmed Aster*, and other indigenous species, *Order Compositæ*, possess aromatic properties.

ASTER DYSENTERICUS, *Inula dysenterica*—a. *Heart-leaved*, *A. cordifolius*—a. *Helenium*, *Inula Helenium*—a. *Lingualis*, *Eryngium campestre*—a. *Officinalis*, *Inula Helenium*—a. *Rough-stemmed*, *A. Punicæus*—a. *Tortifolius*, *Sericocarpus tortifolius*—a. *Undulatus*, *Inula dysenterica*.

ASTERIA GEMMA, *Asterius*, *Astroites*, *Astrios*, *Astrobolus*. The ancients attributed imaginary virtues to this stone,—that of dispersing *Nævi Materni*, for example.

ASTERIAS LUTEA, *Gentiana lutea*.

ASTER'NIA, from *a*, privative, and *στερνον*, 'the sternum.' A monstrosity, in which there is absence of the sternum.

ASTEROCEPHALUS SUCCISA, *Scabiosa succisa*.

ASTHENES, *Infirm*.

ASTHENIA, *Vis imminuta*, from *a*, privative, and *σθένος*, 'force,' 'strength.' Want of strength, debility. (F.) *Affaiblissement*. Infirmary. A word used in this sense by Galen, and employed, especially by Brown, to designate debility of the whole economy, or diminution of the vital forces. He distinguished it into *direct* and *indirect*: the former proceeding from diminution of stimuli; the latter from exhaustion of incitability by the abuse of stimuli.

ASTHENIA DECLUTINONIS, *Pharyngoplegia*—a. *Pectoralis*, *Angina Pectoris*.

ASTHEN'IC, *Asthenicus*, (F.) *Asthénique*. Same etymon as the last. Relating or belonging to *asthenia*.

ASTHENICOPYRA, *Fever, adynamic*.

ASTHENICOPYRETUS, *Fever, adynamic*.

ASTHÉNIQUE, *Asthenic*.

ASTHENO'PIA, *Debitus seu Hebetudo visus*, (F.) *Affaiblissement de la Vue*, from *a*, privative, and *σθένος*, 'strength,' and *ὤψ*, 'the eye.' Weakness of sight; *Weak-sightedness*.

ASTHENOPYRA, *Fever, adynamic*.

ASTHENOPYRETUS, *Fever, adynamic*.

ASTHMA, from *ασπμα*, 'laborious breathing,' from *ασ*, 'I respire.' *A. spasticum adultorum* seu *bronchia'le seu senio'rum seu conculsivum seu spasticum intermittens*, *Spasmus bronchia'lis*, *Dyspnæa et orthopnæa conculsiva*, *Matum Caducum pulmonum*, *Broken-windedness*, *Nervous asthma*, (F.) *Asthme*, *A. nervæ*. Difficulty of breathing, recurring at intervals, accompanied with a wheezing sound and sense of constriction in the chest; cough and expectoration.

Asthma is a chronic disease, and not curable with facility. Excitant and narcotic antispasmodics are required.

There are no pathognomonic physical signs of asthma. In some cases, the respiration is universally puerile during the attack. In the spasmodic form, the respiratory murmur is very feeble or absent during the fit; and in all forms percussion elicits a clear pulmonary sound. The disease generally consists in some source of irritation, and occasionally, perhaps, in paralysis of the pneumogastric nerves, *Bronchoparalysis*, *Paralysis nervi vagi in parte thoracica*, more frequently of the former—all the phenomena indicating constriction of the smaller bronchial ramifications. The treatment is one that relieves spasmodic action—narcotics, counter-irritants, change of air, &c.

ASTHMA ACUTUM, of Millar, *A. spasticum infantum*, *Cynanchæ Trachealis spasmodica*, (F.) *Asthme aiguë*. Probably, spasmodic croup [?] See *Asthma Thymicum*.

ASTHMA AERIUM, *Pneumothorax*—a. *Aërium* ab *Emphysemate Pulmonum*, *Emphysema of the Lungs*—a. *Arthriticum*, *Angina Pectoris*—a. *Bronchiale*, *Asthma*.

ASTHMA CARDIAC, (F.) *Dyspnée ou Asthme Cardiaque*, *Dyspnoea dependent upon disease of the heart*.

ASTHMA CONVULSIVUM, *Angina pectoris*—a. *Diaphragmaticum*, *Angina Pectoris*—a. *Dolorificum*, *Angina pectoris*—a. *Emphysematicum*, *Pneumothorax*.

ASTHMA GRINDERS', *Grinders' Rot*. The aggregate of functional phenomena, induced by the inhalation of particles thrown off during the operation of grinding metallic instruments, &c. The structural changes induced are enlargement of the bronchial tubes, expansion of the pulmonary tissue, and phthisis.

ASTHMA GYPSEUM, *A. pulverulentum*—a. *Hay, Fever*, hay.

ASTHMA HUMIDUM, *Humid, Common*, or *Spitting asthma*, is when the disease is accompanied with expectoration. It is also called *a. humorale*, *A. flatulentum*, *A. pneumonicum*, *Blennorrhæa chronica*, &c.

ASTHMA INFANTUM, *Cynanche trachealis*—a. *Infantum Spasmodicum*, *A. Thymicum*—a. *Koplian*, *A. Thymicum*—a. *Laryngeum Infantum*, *A. Thymicum*—a. *Montanum*, *A. pulverulentum*—a. *Nervous*, *Asthma*—a. *Nocturnum*, *Incebus*.

ASTHMA PULVERULENTUM, *A. gypseum*, *A. montanum*. The variety of asthma to which millers, bakers, grinders, and others are subject.

ASTHMA SICCUM, so called when the paroxysm is sudden, violent, and of short duration; cough slight, and expectoration scanty; spasmodic constriction.

ASTHMA SPASTICO-ARTHRITICUM INCONSTANS, *Angina pectoris*—a. *Spasticum Infantum*, *A. Thymicum*.

ASTHMA THYMICUM, *A. T. Kop'pii*, *Asthma spasmodicum infantum* seu *infantum spasmodicum* seu *largum* seu *infantum* seu *intermittens* *infantum* seu *dentium* seu *periodicum* *acutum*, *Thymasthma*, *Cynanchê trachealis spasmodica*, *Spasmus glottidis*, *Koppian Asthma*, *Thymic Asthma*, *Laryngismus stridulus*, *Laryngo-spasmus*, *Apnoea infantum*, *Spasm of the larynx* or *glottis*, *Croup-like inspiration of infants*, *Child-crouping*, *Spasmodic croup*, *Pseudo-croup*, *Spirious* or *Cerebral croup*, *Phreno-glottism*, *Suffocating nervous catarrh*, (F.) *Laryngitis stridulosa*, *Faux Croup*, *Pseudo-croup nerveux*, *Spasme de la Glotte et du Thorax*. A disease of infants, characterized by suspension of respiration at intervals; great difficulty of breathing, especially on waking, swallowing, or crying; ending often in a fit of suffocation, with convulsions. The pathology of the disease has been supposed to consist in an enlargement of the thymus gland, or of the glands of the neck pressing on the pneumogastric nerves [?]. The ear, by auscultation, at a distance from the chest, detects an incomplete, acute, hissing inspiration, or rather cry; whilst the expiration and voice are croupal, both at the accession and termination of the paroxysm. The heart's action has been observed to be distinct and feeble.

These symptoms are often accompanied by rigidity of the fingers and toes; the thumb being frequently drawn forcibly into the palm of the clenched hand, whence the name *Carpo-pedal spasm*, applied, at times, to the disease.

ASTHMA TRIPICUM. *Asthma* characterized by periodicity.

ASTHMA UTERI, *Hysteria*—*a. Weed*, *Lobelia inflata*.

ASTHMATIC, *Asthmaticus*, *Pneucolyticus*, (F.) *Asthmatique*. Affected with asthma. Relating to asthma.

ASTHME AIGU, *Asthma acutum*—*a. Cardi-aque*, *Asthma cardiacæ*—*a. Nervens*, *Asthma*.

ASTIGMATISM, *Astigmatismus*, from *a. privative*, and *στριγμα*, *striptos*, 'a point'. A state of irregular refraction in the eye, in which the rays are not brought to one focus, but converge at different distances, so as to form two linear images at right angles to each other.

ASTOMUS, from *a. privative*, and *στρομα*, 'a mouth'. (F.) *Astome*. One without a mouth. Pliny speaks of a people in India without mouths, who live *anhelati et odore!*

ASTRAGALE COL D', *Collum astragali*.

ASTRAGALCIDES SYPHILITICA, *Astragalus excapus*.

ASTRAGALUS, *Talus*, the *Ankle*, *Quat'rio*, *Quar'tio*, *Quater'nio*, *Diab'ebos*, *Peza*, *Cavic'ula*, *Cavill'a*, *Tetro'ros*, *Astr'ion*, *Os Ballist'æ* seu *Tes'sera*, from *αστραγᾶλος*, 'a die,' which it has been considered to resemble[?]. A short bone, situate at the superior and middle part of the tarsus, where it is articulated with the tibia. It is the *ankle bone*, *sliding bone*, or *first bone of the foot*. The anterior surface is convex, and has a well-marked prominence, supported by a kind of neck, and hence has been called the *head of the astragalus*. The astragalus is developed by two points of ossification.

Also, the atlas or first vertebra of the neck.

ASTRAGALUS EXSCAPUS, *Astragali'ides syphilitica*, *Stemless Milk-vetch*, (F.) *Astragale à poussees velus*. *Ord. Leguminosæ*. *Sex. Syst. Dialdelphia Decandria*. The root is said to have cured confirmed syphilis.

ASTRAGALUS TRAGACANTHUS, see *Tragacanth*.

ASTRAGALUS VERUS, *Spina hirci*, *Astrag'alus aculeatus*, *Goat's thorn*, *Milk-vetch*. The plant which affords *Gum Tragacanth*. See *Tragacantha*.

ASTRANTIA, *Imperatoria*—*a. Diapensia*, *Sanicula*.

ASTRAPE, *Coruscat'io*, *Fulgar*, *Fulmen*, *Lightning*. Galen reckons it amongst the remote causes of epilepsy.

ASTRICT'ION, *Astrict'io*, *Stypsis*, *Adstrict'io*, *Constrict'io*, from *astringere*, (*ad* and *stringere*,) 'to constringe.' Action of an astringent substance on the animal economy.

ASTRICTORIA, *Astringents*.

ASTRINGENT ROOT, *Comptonia asplenifolia*.

ASTRIN'GENTS, *Astringent'ia*, *Adstricto'ria*, *Adstringent'ia*, *Stryphna*, *Catastalt'ica*, *Constringent'ia*, *Contrahent'ia*, *Stegno'tica*, *Syncret'ica*, *Astricto'ria*. Same etymon. Medicines which have the property of constringing the organic textures. External astringents are called *Styp-tics*.

The following are the chief astringents: *Acidum Sulphuricum*, *A. Tannicum*, *Alumen*, *Argent' Nitras*, *Catechu*, *Crescoton*, *Cupri Sulphas*, *Tinct. Ferri Chloridi*, *Liquor Ferri Nitratis*, *Ferri Sulphas*, *Galla*, *Hæmatoxylin*, *Kino*, *Krameria*, *Liquor Calcis*, *Plumbi Acetas*, *Quercus Alba*, *Quercus Tinctoria*, *Zinci Sulphas*.

ASTRION, *Astragalus*.

ASTRIOS, *Asteria gemma*.

ASTROBLES, from *αστρον*, 'a star,' and *βαλλω*, 'I strike.' One struck by the stars, (*sideratus*.) One who is in a state of sideration—in an apopleptic state.—*Gorræus*.

ASTROBLESTIA, *Astrobolismus*.

ASTROBOLIA, *Astrobolismus*.

ASTROBOLIS'MUS, *Astrobol'ia*, *Astroble'sia*, *Heli'asis*, *Heli'o'sia*. Same etymology. *Sidera'tion*, or action of the stars on a person. *Apoplexy*.—*Theophrastus*, *Gorræus*.

ASTROBOLUS, *Asteria gemma*.

ASTROITIS, *Asteria gemma*.

ASTROLOGER, see *Astronomy*.

ASTROLOG'Y, *Astrolog'ia*, from *αστρον*, 'a star,' and *λογος*, 'a discourse.' The art of divining by inspecting the stars. This was formerly considered to be a part of medicine; and was called *Judicial Astrology*; to distinguish it from astronomy.

ASTRONOM'Y, *Astronom'ia*, from *αστρον*, 'a star,' and *νομος*, 'a law,' 'rule.' A science which makes known the heavenly phenomena, and the laws that govern them. Hippocrates places this and astrology amongst the necessary studies of a physician. In the earlier English writers, *Astronomer* is often used in the sense of *Astrologer*.

ASYSTOLIE, from *a. privative*, and *συστολη*, 'systole. A name given by M. Beau to a period, in diseases of the heart, in which the systole is incomplete; or, at least, is insufficient to free the ventricles from the blood in them.

ASTRUTHIUM, *Imperatoria*.

ASTYPHIA, *Impotence*.

ASTYSIA, *Impotence*.

ASUAR, *Myrobalanus Indica*.

ASULCI, *Lapis lazuli*.

ASYNODIA, *Impotence*.

ATACTOS, *Erratic*.

ATARACTAPOIE'SIA, *Ataractopœ'sia*, from *a. privative*, *ταραχος*, 'troubled,' and *ποιειν*, 'to make.' Intrepidity, firmness; a quality of which, according to Hippocrates, the physician ought to be possessed in the highest degree.

ATARAX'IA, from *a. privative*, and *ταραξις*, 'trouble,' 'emotion.' Moral tranquillity, peace of mind.

ATAVISM, *Atavism'us*, from *átavis*, 'an old grandsire or ancestor indefinitely.' The case in which an anomaly or disease, existing in a family,

is lost in one generation and reappears in the following.

ATAX'IA, from *a*, privative, and *ταξις*, 'order.' Disorder, irregularity. Hippocrates employs the word in its most extensive acceptation. Galen applies it, especially, to irregularity of pulse; and Sydenham speaks of *Ataxia Spirituum* for disorder of the nervous system. Ataxia, now, usually means the state of disorder that characterizes nervous fevers, and the nervous condition.

ATAXIA SPIRITUUM, Nervous diathesis. See Ataxia.

ATAX'IC, *Atax'icus*; same etymon. Having the characters of ataxia.

ATCHAR, *A'chia*, *Achar*. A condiment used in India. It is formed of green fruits of various kinds—garlic, ginger, mustard, and pimento—pickled in vinegar.

ATECNIA, Sterilitas.

ATELEC'TASIS, *Atelek'tasis*, from *ατελής*, 'imperfect,' 'defective,' and *εκτασις*, 'dilatation.' Imperfect expansion or dilatation; as in

ATELEC'TASIS PULMO'NUM, *Pneumotelektasis*, *Pneumotelektasis*. Imperfect expansion of the lungs at birth, giving rise to *Cyano'sis pulmonalis*. A similar condition is observed in lungs which have received air, and in a distinctly lobular form. This is regarded by Dr. W. T. Gairdner as, in all probability, a secondary lesion, and dependent, in the majority of instances, on a catarrhal condition of the bronchial tubes. It is called, by him, *pulmonary collapse*, or *collapse of the lung*, and has received the name *apneumato'sis* from Fuchs.

ATELES, *ατελής*, 'imperfect,' 'defective.' Hence

ATEL'IA, (F.) *Atélie*. A monstrosity in which there is a want of some members.

ATELOCHE'LIA, from *ατελής*, 'imperfect,' and *χελος*, 'lip.' A malformation which consists in an imperfect development of the lip.

ATELOENCEPHAL'IA, from *ατελής*, 'imperfect,' and *εγκεφαλον*, 'the encephalon.' State of imperfect development of the brain.—Andral.

ATELOGLOS'SIA, from *ατελής*, 'imperfect,' and *γλωσσα*, 'tongue.' A malformation which consists in an imperfect development of the tongue.

ATELOGNA'THIA, from *ατελής*, 'imperfect,' and *γναθος*, 'the jaw.' A malformation which consists in an imperfect development of the jaw.

ATELOMYEL'IA, from *ατελής*, 'imperfect,' and *μυελος*, 'marrow.' State of imperfect development of the spinal marrow.—Béclard.

ATELOPROSO'PIA, from *ατελής*, 'imperfect,' and *προσωπον*, 'the face.' A malformation which consists in imperfect development of the face.

ATELORACHIDIA, *Hydrorachis*.

ATELOSTOM'IA, from *ατελής*, 'imperfect,' and *στομα*, 'mouth.' One whose mouth is imperfectly developed.

ATER SUCCUS, *Atrabilis*.

ATHAMANT'A, from Athamas, a place in Thessaly. A genus of plants.

ATHAMANTA ANNUA, *A. Cretensis*.

ATHAMANTA AUREOBELI'NUM, *Oreoseli'num*, *O. legitimum* seu *nigrum*, *Selin'num oreoseli'num*, *Peuced'anum oreoseli'num*, *Apium montanum*, *Black Mountain Parsley*, (F.) *Persil de Montagne*; *Ord. Umbelliferae*. The plant, seed, and roots, are aromatic. It has been considered attenuant, aperient, deobstruent, and lithontriptic. The distilled oil has been used in toothache.

ATHAMANTA CRETEN'SIS seu CRE'TICA, *A. annua*, *Libanotis annua* seu *Creten'sis* seu *hiruta*, *Daucus Creticus* seu *Candia'nis*, *Myrrhis annua*, *Candy Carrot*. The seeds of this plant are acid

and aromatic. They have been used as carminatives and diuretics.

ATHAMANTA MACEDONICA, *Bubon Macedonicum*—*a*. Meum. *Ethusa meum*.

ATHANASIA, *Tanacetum*.

ATHANA'SIA, from *a*, privative, and *θανος*, 'death.' An antidote for diseases of the liver, jaundice, gravel, &c. It consisted of saffron, cinnamon, lavender, cassia, myrrh, juncus odoratus, honey, &c., and was esteemed to be sudorific.

ATHARA, *Athera*.

ATHELASMUS, from *a*, privative, and *θηλη*, 'a breast or nipple.' Impracticability of giving suck; from want of nipple or otherwise.

ATHELXIS, Sucking.

ATHE'NA. Name of a plaster, recommended by Asclepiades, and composed of oxide of copper, sublimed oxide of zinc, sal ammoniac, verdigris, gall nuts, and a variety of resinous and other ingredients.—Oribasius, Aëtius, and P. Egineta.

ATHENIO'NIS CATAPOTIUM. A pill, composed of myrrh, pepper, castor, and opium; used to allay coughing.—Celsus.

ATHE'RA, *Atha'ra*, from *αθηρ*, 'an ear of corn.' A kind of pap for children; also, a kind of liniment.—Dioscorides, Pliny.

ATHERAPEUTUS, Incurable.

ATHERO'MA, from *αθηρ*, 'pap or pulp,' *Emphy'ma encystis atheroma*, *Mollus'cum*, *Pultatio*. A tumour formed by a cyst containing matter like pap or *Bouillie*, or plaster.

ATHEROM'ATOUS, *Atheromato'des*. Having the nature of Atheroma.

ATHE'LETA, from *αθλος*, 'combat.' *Athlete*, (F.) *Athlète*. *Athletæ* were men who exercised themselves in combat at the public festivals.—Vitruvius.

ATHELET'IC, *Athlet'icus*, (F.) *Athlétique*; concerning *Athletæ*. Strong in muscular powers.—Fœsius.

ATHOL PORRIDGE, see Mahogany.

ATHORACOCEPHALUS, *Acephalogaster*.

ATHIRIX, *At'richus*, *Culex*, (F.) *Chauve*; from *a*, privative, and *θριξ*, *τριχος*, 'hair.' *Bald*, (Sc.) *Beld*, *Bellit*. One who has lost his hair.

ATHRIX DEPILIS, *Alopecia*.

ATHYM'IA, *An'imi defectus* et *anxi'etas* seu *demissio*, *Tristitia*, *Mæror*, *Lyp'.* from *a*, priv., and *θυμος*, 'heart,' 'courage.' Despondency. The prostration of spirits often observable in the sick.—Hippocrates. Melancholy.—Swediaur. See Panophobia.

ATHYMIA PLEONECTICA, see Pleonectica.

ATHYRION, *Asplenium ceterach*.

ATHYRIUM FILIX FEMINA. *Asplenium filix femina*—*a*. *Filix* mas, *Polypodium filix mas*—*a*. *Molle*, *Asplenium filix femina*—*a*. *Ovatum*, *Asplenium filix femina*—*a*. *Trifidum*, *Asplenium filix femina*.

ATLANTAD, see Atlantal.

ATLANTAL; same etymon as *Atlas*. Relating or appertaining to the atlas.

ATLANTAL ASPECT. An aspect towards the region where the atlas is situated.—Barclay. *Atlantal* is used by the same writer to signify 'towards the atlantal aspect.'

ATLANTAL EXTREMITIES. The upper limbs.

ATLANTION, *Atlas*.

ATLAS, *Atlantion*, *Astragalus*, from *ατλαω*, 'I sustain.' The first cervical vertebra; so called from its supporting the whole weight of the head, as Atlas is said to have supported the globe on his shoulders. Chaussier calls it *Atloïde*. This vertebra in no respect resembles the others. It is a kind of irregular ring, into which, anteriorly, the *processus dentatus* of the second vertebra is received. Posteriorly, it gives passage to the medulla spinalis.

ATLODID'YMUS, *Atlod'y-mus*, (F.) *Atlodyme*: from Atlas, 'the first bone of the neck,' and *ἰδυμος*, 'a twin.' A monster which is simple below and double above. M. Isid. Geoffroy Saint-Hilaire applies this term to a monster, which has a single body, with two distinct heads supported on one neck.

ATLOÏDE, Atlas.

ATLOÏD'O-AXOID, (F.) *Atloïdo-axoïdien*. Relating to both the Atlas and the Axis or Vertebra Dentata.

ATLOÏDO-AXOID ARTICULATION. The articulation between the first two cervical vertebrae.

ATLOÏDO-AXOID LIG'AMENTS. These are two in number; one anterior and another posterior, passing between the two vertebrae.

ATLOÏD'O-OCCIP'ITAL. Relating to the atlas and occiput. The *Atloïdo-occipital Articulation* is formed by the condyles of the occipital bone and the superior articular surfaces of the Atlas. The *Atloïdo-occipital muscle* is the Rectus capitis posticus minor.

ATLOÏDO-SOUS-MASTOÏDIEN, Obliquus superior oculi—a. *Sous-occipitale*, Rectus capitis lateralis.

ATMIATRIA, *Atmidiat'ricē*, from *ατμος*, 'vapour,' and *ιατρεία*, 'treatment.' Treatment of diseases by fumigation.

ATMIDIATRICE, Atmiatria.

ATMISTERION, Vaporarium.

ATMOS, Breath.

AT'MOSPHERE, *Atmosphæ'ra*, from *ατμος*, 'vapour,' and *σφαῖρα*, 'a sphere;'—as it were, *Sphere of vapours*. The atmosphere is a spherical mass of air, surrounding the earth in every part; the height of which is estimated at 15 or 16 leagues. It presses on the surface of the earth, and this pressure has, necessarily, sensible effects on organized bodies. The surface of the human body being reckoned at 15 square feet, it is computed that a pressure of 33,000 pounds or more exists under ordinary circumstances; and this pressure cannot be increased or diminished materially, without modifying the circulation and all the functions.

ATMOSPHERIZATION, Hæmatisis.

ATOCIA, Sterilitas.

ATOL'MIA, from *a*, priv., and *τολμα*, 'confidence.' Want of confidence; discouragement. A state of mind unfavorable to health, and injurious in disease. It is the antithesis of *Eutol'mia*.

ATOMY, Skeleton.

ATONIA, Atony—a. Ventriculi, Gasterasthenia.

AT'ONIC, *Aton'icus*, (F.) *Atonique*. Wanting tone. Same etymon as the next. Also, a medicine capable of allaying organic excitement or irritation.—Schwlgué.

AT'ONY, *At'on'ia*, *Infr'mitas et Remissio vi-rum*, *Langor*, *Lax'itas*, from *a*, priv., and *τονος*, 'tone,' 'force.' Want of tone. Weakness of every organ, and particularly of those that are contractile. Violent gastritis has been described by Scribonius Largus under a similar name, *Ατωνα*, *At'onon*.

ATRABIL'IARY, *Atrabil'ious*, *Atrabilia'ris*, *Atrabil'ious*, (F.) *Atrabilaire*, *Atrabileux*, from *ater*, 'black,' and *bilis*, 'bile.' An epithet given by the ancients to the melancholic and hypochondriac, because they believed the Atrabilis to predominate in such.

ATRABILARY CAPSULES, ARTERIES and VEINS. The renal capsules, arteries and veins; the formation of Atrabilis having been attributed to them.

ATRABI'LIS, same etymon, *Ater succus*, *Black Bile* or *melancholy*, (F.) *Atrabile*. Ac-

cording to the ancients, a thick, black, acrid humour, secreted, in the opinion of some, by the pancreas; in that of others, by the supra-renal capsules. Hippocrates, Galen, Aëtius, and others, ascribe great influence to the Atrabilis in the production of hypochondriasis, melancholy, and mania. There is really no such humour. It was an imaginary creation.—Aretæus, Rufus of Ephesus, &c.

ATRACHE'LOCEPH'ALUS, from *a*, priv., *τραχηλος*, 'neck,' and *κεφαλή*, 'head.' A monster whose neck is partially or wholly deficient.

ATRACHE'LUS. Same etymon. One who is, very short-necked.—Galen.

ATRAC'TYLIS GUMMIF'ERA, *Carduus pi'neus*, *Ixi'ne*, *Gummy-rooted Attractylis*, *Pine Thistle*. Ord. Compositæ. The root, when wounded, yields a milky, viscid juice, which concretes into tenacious masses, and is said to be chewed with the same views as mastic.

ATRAGENE, *Clematis vitalba*.

ATRAMEN'TUM, *A. Sutor'ianum*, *Ink*, *Chaleant'hum*, (F.) *Encre*. It has been advised as an astringent, and as an external application in herpetic affections.

ATRAMENTUM SUTORIUM, Ferri sulphas.

ATRE'SIA, (F.) *Atresie*, Adherence, Imperforation. Same etymon as Atrētus. See Monster.

ATRE'SIA ANI ADNA'TA, *Anus Imperforatus*, *Imperforatio ani*, *Atrētocy'sia*, (F.) *Imperforation de l'anus*. Congenital imperforation of the intestinal canal.

ATRETELYTRIA, Colpatresia.

ATRETISMUS, Imperforation.

ATRETOCEPH'ALUS, from *ατρητος*, 'imperfect,' and *κεφαλή*, 'head.' A monster, in which some of the natural apertures of the head are wanting.—Gurlt.

ATRETOCOR'MUS, from *ατρητος*, 'imperfect,' and *κορμος*, 'trunk.' A monster in which the natural apertures of the trunk are wanting.—Gurlt.

ATRETOCY'SIA, Atresia ani.

ATRETOMETRIA, Hysteratresia.

ATRETOPSIA, Coreclisis.

ATRETOSTOM'IA, from *ατρητος*, 'imperfect,' and *στομα*, 'mouth.' Imperforation of the mouth.

ATRETURE'THRIA, from *ατρητος*, 'imperfect,' and *ουρηθρα*, 'urethra.' Imperforation of the urethra.

ATRE'TUS, from *a*, priv., and *τραπος*, 'I perforate.' *Imperforatus*, *Imperforate*. One whose anus, or parts of generation, are imperforate (*apoc'itus*).

ATRIA, Auricles of the heart—a. Mortis, see Death.

AT'RICES. Small tumours, which appear occasionally around the anus. Some commentators consider the word to be synonymous with condylomata.—Forestus.

ATRICHIA, Alopecia—a. Adnata, see Alopecia—a. Senilis, see Alopecia.

ATRICHUS, Athrix.

ATRICI. Small sinuses in the vicinity of the anus, not penetrating the rectum.

ATRIO-VENTRICULAR, Auriculo-ventricular.

ATRIPLEX FÆTIDA, *Chenopodium vulvaria*. **ATRIPLEX HORTEN'SIS**, *A. Sati'va*, (F.) *Ar-roche*, *Bonne Dame*, *Follette*. Ord. Chenopodiaceæ. The herb and seed of this plant have been exhibited as antiscorbutics.

Atriplex al'imus, *A. Portulacoi'des*, and *A. Pat'ula*, are used as pickles, and have similar properties.

ATRIPLEX LACINIATA, *A. patula*—a. Mexicana, *Chenopodium ambrosioides*—a. Odorata, *Chenopodium botrys*—a. Olida, *Chenopodium vulvaria*.

ATRIPLEX PAT'UEA, *A. lacinia'ta* seu *Pur-ehin'na*, *Spreading Orache*; indigenous; *Family*, Chenopodiaceæ. The expressed juice is said to be cathartic. It has been used, in place of gamboge, in dropsy and asthma.

ATRIPLEX PURSHIANA, *A. patula*.

ATRIUM CORDIS DEXTER, Sinus dexter cordis—a. Cordis sinistrum, Sinus pulmonalis—a. Vagina. Vestibulum.

AP'ROPA, from *Ἀπορος*, 'immutable,' 'the goddess of destiny,' so called from its fatal effects.

ATROPA BELLADON'NA, *Belladon'na*, *B. baccifera* seu *trichot'oma*, *Deadly Nightshade*, *Common Dwole*, (Sc.) *Mekilhoort*, *Sola'num lethale* seu *horten'se nigrum*, *Sola'num mac'num* seu *furi'um* seu *melanocer'aeus*, (F.) *Belladone*, *Mocelle furieuse*, *Belle Dame*. *Ord.* Solanaceæ. *See Syst.* Tetrandria Monogynia. The leaves—*Belladonna* (Ph. U. S.)—are powerfully narcotic, and also diaphoretic, and diuretic. They are occasionally used where narcotics are indicated. Sprinkling the powdered leaves over cancerous sores has been found to allay the pain; and the leaves form a good poultice. Dose, gr. $\frac{1}{2}$ to gr. j of the powdered leaves.

ATROPA MANDRAG'ORA, *Mandrag'ora*, *M. ver-nalis* seu *offici'alis* seu *acaul'is*, *Circæ'a*, *Anthro-pomorph'us*, *Malum terres'tre*, *Mandrake*. The boiled root has been used in the form of poultice to indolent swellings.

ATROPHIA, *Atrophy*, *Tabes*—a. Ablactat-urum, *Brash*, weaning—a. Cerebri, *Phrenatrophia*—a. Cordis, *Heart*, atrophy of the—a. Glandularis, *Tabes mesenterica*—a. Hepatis, *Hepatatrophia*—a. Infantum, *Pædatrophia*, *Tabes mesenterica*—a. Intestinorum, *Enteratrophia*.

ATROPHIA LACTAN'TIUM, *Tabes nutri'cum* seu *lactea*. The atrophy of nursing women.

ATROPHIA LIENIS, *Splenatrophia*—a. Mesen-terica, *Tabes mesenterica*—a. Testiculi, *Orchidatrophia*—a. Unguium, *Onychatrophia*.

ATROPHIE, *Atrophy*—a. *Mésentérique*, *Tabes mesenterica*.

ATROPHIE MUSCULAIRE PROGRES-SIVE, *Paralytic musculaire atrophique*, *Atrophie musculaire primitive ou idiopathique ou avec transformation graisseuse*. A rare malady, in which the muscles become so greatly atrophied that they cannot perform their functions.

ATROPHIED, see *Atrophy*.

ATROPHY, *Marasmus Atro'phia*, *Atro'phia Maras'mus*, *Maras'sis*, *Ma'cies*, *Contabescen'tia*, *Tabes*, *Marco'res*, *Analo'sis*, from *a*, privative, and τροφή, 'nourishment.' (F.) *Atrophie*, *Des-èchement*. Defective nutrition, *Hypotrophy*. Progressive and morbid diminution in the bulk of the whole body or of a part. Atrophy is generally symptomatic. Any tissue or organ thus affected is said to be *atrophied*.

ATROPHY OF THE HEART, see *Heart*, atrophy of the.

ATROPIA, *Atropine*—a. Sulphate of, *Atropiæ sulphas*.

ATROP'IE SULPHAS, *Sulphate of Atrop'ia* (Ph. L.). Formed by dissolving *atropia* in dilute sulphuric acid; evaporating, and crystallizing. Virtues, the same as those of *Atropia*.

ATROPINE, *Atropi'na*, *Atrop'ia*, *Atrop'ium*, *Atrop'ium*, *Belladon'nin*, (F.) *Atropine*. The active principle of *Atropa Belladonna*, separated by Brandes, by a process similar to that for procuring morphia. A single drop of a solution, of one grain in fʒiv of distilled water with a few drops of acetic acid, applied to the inner surface of the lower eyelid, causes dilatation of the pupil, in fifteen or twenty minutes. A grain to a drachm of lard is an application in neuralgia.

ATTACHE, Insertion.

ATTACK, *Insul'tus*, *Assul'tus*, *Irrep'tio*, *Inva'sio*, *Eis'bolé*, *Lapsis*, (Prov.) *Take*, (F.) *Attaque*. A sudden attack, invasion or onset of a disease. A seizure. One attacked or affected with severe disease is often said, in the United States, to be "taken down," or to be "down" with it.

ATTAGAS, *Attagen*.

ATTAGEN, *Attagas*, the *Fran'colin*. Celebrated with the ancients both as food and medicine.—*Martial*, *Aristophanes*.

ATTANCOURT, MINERAL WATERS OF. A mineral water in France, at Attancourt, in Champagne; about three leagues north of Joinville. The water is a chalybeate, and contains sulphate of lime. In large doses it is purgative.

ATTAQUE, *Attack*—a. *des Nerfs*, *Nervous attack*.

ATTAR OF ROSES, see *Rosa centifolia*.

ATTELE, Splint.

ATTENOTING, MINERAL WATERS OF, in Bavaria. The water contains carbonic acid, carbonates of lime and soda, sulphates of lime and magnesia, chloride of sodium, iron and alum. It is much used in skin diseases, fistula, old ulcers, calculi, and hæmorrhoids.

ATTENUANTS, *Attenuan'tia*, *Leptun'tica*, (F.) *Atténuans*, *Leptontiques*, from *tenuis*, 'thin.' Medicines which augment the fluidity of the humours.

ATTENUA'TION, *Attenua'tio*; same etymon. Thinness, emaciation. A term used by the homœopaths, in the sense of dilution or division of remedies into infinitesimal doses.

ATTRANT, *Attrahent*.

ATTITUDE, *Situs Cor'poris*. Low Latin, *optitudo*; from Latin *optare*, 'to fit.' Situation, position of the body. The attitudes are the different postures which man is capable of assuming. In *General Pathology*, the attitude will often enable the physician to pronounce at once upon the character of a disease, or it will aid him materially in his judgment. In St. Vitus's dance, in fractures, luxations, &c., it is the great index. It will also indicate the degree of nervous or cerebral power; hence sinking down in bed is an evidence of great cerebral debility in fever. The position of a patient during an operation is also an interesting subject of attention to the surgeon.

ATTOL'ENS AUREM, *A. Auric'ulæ*, *Leva'tor Auris*, *Superior Auris*, *Attol'ens Auric'ulum*, *Auricula'ris super'ior*, (F.) *Auriculaire supérieur*, *Temporo-auriculaire*. A muscle of the ear, which arises, thin, broad, and tendinous, from the tendon of the occipito-frontalis, and is inserted into the upper part of the ear, opposite to the anti-helix. It raises the ear.

ATTOLLENS OCULI, *Rectus superior oculi*—a. *Oculum*, *Rectus superior oculi*.

ATTOUCHEMENT, *Masturbation*.

ATTRACTION OF AGGREGATION, *Cohesion*, force of.

ATTRACTIVUM, see *Magnet*.

ATTRACTIVUS, *Attrahent*.

ATTRACTORIUS, *Attrahent*.

ATTRAHENS AURICULAM, *Anterior auris*.

AT'TRAHENT, *At'trahens*, *Attract'ivus*, *Attracto'rius*, from *ad*, 'to,' and *traho*, 'I draw.' (F.) *Attractif*, *Attrayant*. Remedies are so called, which attract fluids to the parts to which they are applied, as blisters, rubefacients, &c.

ATTRAPE-LOURDAUT (F.). A bistoury invented by a French surgeon, called Biennaise, and used in the operation for hernia. See *Bistouri caché*.

ATTRITA, Chafing.

ATTRITIO, *Attrition*, Chafing.

ATTRIT'ION, *Attrit'io*, *Eethlim'ma*, from

ad, and *terere*, 'to bruise.' Friction or bruising. Chafing.—Galen. Also, a kind of cardialgia.—Sennertus. Likewise, a violent contusion.

ATTRITUS, Chafing.

ATYP'IC, *Atypicus*, *At'ypos*, (F.) *Atypique*, from *a*, privative, and *typos*, 'type.' That which has no type. Irregular. Chiefly applied to an irregular intermittent.—*Febris atypica*.

ATYPOS, Erratic.

AUANSIS, Drying.

AUANTE, *Auap'sē*, from *avaris*, 'desiccation.' Hippocrates gave this name to a disease, the principal symptom of which was emaciation. Atrophy.

AUBE-VIGNE, *Clematis vitalba*.

AUDÉPINE, *Mespilus oxyacantha*.

AUBERGINE, *Solanum Melongena*.

AUBIFOIN, *Centaurea cyanus*, *Cyanus segetum*.

AUCHEN, Collum.

AUCHENORRHEUMA, *Torticollis*.

AUCHE'TICUS, from *αυχν*, 'the neck.' One affected with stiff neck or torticollis.

AUCTUMUS, Autumn.

AUDE, Voice.

AUDINAC, MINERAL WATERS OF. Audinac is situate in the department of Arriège, France. The water contains a small quantity of sulphohydric acid, carbonic acid, sulphates of lime and magnesia, carbonates of lime and iron, and a bituminous substance. Temp. 67° Fahr. It is much used in chronic rheumatism, herpes, scrofulous diseases, &c.

AUDITIF, Auditory.

AUDIT'ION, from *audire*, *audire*, 'to hear.' *Audit'io*, *Audit'us*, *Acōē*, *Acōē'sis*, *Acu'sis*, (F.) *Ouïe*, Hearing. The act of hearing. The sensation arising from an impression made on the auditory nerves by the vibrations of the air, produced by a sonorous body. The physiology of Audition is obscure. It probably takes place:—
1. By the vibrations being communicated from the membrana tympani along the chain of small bones to the membrane of the foramen ovale.
2. By means of the air in the cavity of the tympanum, the membrane of the foramen rotundum is agitated. 3. The transmission may be made by means of the bony parietes. In these three ways the vibrations produced by a sonorous body may reach the auditory nerve. Audition may be active or passive: hence the difference between listening and simply hearing.

AUDITORY, *Audit'orius*, *Audit'ivus*, *Acus'ticus*, (F.) *Auditif*. That which relates to audition.

AUDITORY ARTERIES AND VEINS, are vessels which enter the auditory canals, and are, like them, distinguished into internal and external. The external auditory artery, *A. Tympanique*—(Ch.) is given off by the styloid, a branch of the external carotid: the internal is a branch of the basilar artery, which accompanies the auditory nerve, and is distributed to it. The Auditory Veins empty into the internal and external jugulars.

AUDITORY CANAL, EXTERNAL, *Meatus audit'orius exter'nus*, *Alvea'rium*, *Scapha*, *Scaphus*, (F.) *Conduit auditif externe*, *Conduit auriculaire*, commences at the bottom of the concha, at the *Fora'men audit'ivum exter'num*, passes inwards, forwards, and a little downwards, and terminates at the membrana tympani. It is partly cartilaginous, partly osseous, and partly fibrous.

AUDITORY CANAL, INTERNAL, *Meatus audit'orius inter'nus*, *Porus* seu *Sinus acus'ticus*, *Cyar*, (F.) *Conduit auditif interne*, *C. labyrinthique*, is situate on the posterior surface of the pars petrosa of the temporal bone. From the *Fora'men audit'ivum inter'num*, where it commences, it

passes forwards and outwards, and terminates by a kind of *cul-de-sac*, *mac'ula cribro'sa*, perforated by many holes, one of which is the orifice of the *Aquæductus Fallopi*; and the others communicate with the labyrinth.

AUDITORY NERVE, *Nervus audit'orius* seu *acus'ticus*, *Nerf labyrinthique*—(Ch.), is the *Portio Mollis* of the seventh pair. It arises from the corpus testiforme, from the floor of the fourth ventricle, and by means of white striæ from the sides of the calamus scriptorius. As it leaves the encephalon, it forms a flattened cord, and proceeds with the facial nerve through the foramen auditivum internum, and as far as the bottom of the meatus, where it separates from the facial, and divides into two branches, one going to the cochlea, the *cochlear*; the other to the vestibule and semi-circular canals, the *vestibular*.

AUGÉ, *Al'veus*. Some of the older anatomists gave this name to a reservoir, into which liquids flow in an interrupted manner, so that it is alternately full and empty. Such are the ventricles and auricles of the heart.

AUGGERE, Intermittent Fever.

AUGMENT'ION, from *augere*, *auctum*, 'to increase.' *Augment'um*, *Increment'um*, *Anal'asis*, *Aug'tio*, *Auris*, *Progres'sio*, *Progres'sus*, *Acu'e'sis*, (F.) *Augment*. The stage of a disease in which the symptoms go on increasing.

AULISCUS, Canula. See *Fistula*.

AULOS, Canula, *Fistula*. See *Vagina*, and *Foramen*.

AUMALE, MINERAL WATERS OF. Aumale is a town of Upper Normandy, in the country of Caux. Several springs of ferruginous mineral waters are found there, whose odour i penetrating, and taste rough and astringent. They are tonic, and employed in debility of the viscera, &c.

AUMURE, *Parietaria*.

AUNE NOIRE, *Rhamnus frangula*.

AC'NÉE, *Inula helenium*—*a. Dysentérique*, *Inula dysenterica*—*a. Officinale*, *Inula helenium*.

AURA, *Pnoë*. A vapour or emanation from any body, surrounding it like an atmosphere. Van Helmont regarded the vital principle as a gas and volatile spirit, which he called *Aura vitalis*.

In *Pathology*, *Aura* means the sensation of a light vapour, which, in some diseases, appears to set out from the trunk or limbs; and to rise towards the head. This feeling has been found to precede attacks of epilepsy and hysteria, and hence it has been called *Aura Epileptica*, and *A. hyster'ica*.

AURA EPILEPTICA, see *Aura*—*a. Hysterica*, see *Aura*.

AURA SAN'GUINIS. The odour exhaled by blood newly drawn. See *Gaz Sanguinis*.

AURA SEM'INIS, *A. semina'lis*, *Spir'itus genitalis*:—A volatile principle fancied to exist in the sperm, and regarded by some as the fecundating agent. Such is not the case.

AURA VITALIS, Vital principle.

AURAL MEDICINE AND SURGERY, *Otiatria*.

AURANCUM, see *Ovum*.

AURANITE, see *Agaric*.

AURANTIA CURASSAVENT'IA, *A. Curassav'ica*, *Curasso's apple* or *oranges*. Immature oranges, checked, by accident, in their growth. They are a grateful, aromatic bitter, devoid of acidity. Infused in wine or brandy they make a good stomachic. They are also used for *issus peas*. See *Citrus aurantium*.

AURANTII CORTEX, see *Citrus aurantium*—*a. Floris Aqua*, see *Citrus aurantium*.

AURANTIUM, *Citrus aurantium*.

AURELIANA CANADENSIS, *Panax quinquefolium*.

AUREOLA, *Areola*.

AURI CHLORETUM CUM CHLORETO NATRII, see Gold—a. Chloridum, Gold, muriate of—a. Chloretum, Gold, muriate of—a. Cyanidum, see Gold—a. Cyanuretum, see Gold—a. Iodidum, see Gold—a. Ioduretum, see Gold—a. et Natri chloruretum, see Gold—a. Murias, Gold, muriate of—a. Nitro-murias, see Gold—a. Oxidum, see Gold—a. Terchloridum, see Gold—a. Tereyanidum, see Gold—a. Teroxidum, see Gold.

AURICHALCUM, Brass.

AURICLE, *Auricula*, (F.) *Auricle*, *Oreicle*. Diminutive of *auris*, an ear. The pavilion of the ear. See Pavilion.

AURICLES OF THE HEART, *Auricula*, *Atria*, *Cavities innominatae*, (F.) *Oreillettes*, are two cavities, one right, the other left, each communicating with the ventricle of its side. These two cavities receive the blood from every part of the body. Into the *right auricle*, the two *venae cavae* and coronary vein open: into the *left*, the four pulmonary veins. Chaussier calls the former the *Sinus of the Venae Cavae*:—the latter, the *Sinus of the Pulmonary Veins*. The foliated or dog's ear portion of each auricle is called *Appendix auriculæ*. See *Sinus*.

AURICULA JUDÆ, *Peziza auricula*—a. *Muris*, *Hieracium Pilosella*—a. *Muris major*, *Hieracium murorum*.

AURICULARE, *Auricular*, see *Digitus*—a. *Postérieur*, *Retrahens auris*—a. *Supérieur*, *Attollens aurum*.

AURICULAR, *Auricularis*, *Oreicular*, (F.) *Auriculaire*, from *auricula*, 'the ear.' That which belongs to the ear, especially to the external ear.

AURICULAR ARTERIES AND VEINS, *Oreiculaires*—(Ch.), are divided into *anterior* and *posterior*. The *anterior* are of indeterminate number. They arise from the temporal artery, and are distributed to the *meatus auditorius externus*, and to the pavilion of the ear. The *posterior auricular* is given off by the external carotid, from which it separates in the substance of the parotid gland. When it reaches the inferior part of the pavilion of the ear it bifurcates; one of its branches being distributed to the inner surface of the pavilion, the other passing over the mastoid process, and being distributed to the temporal and posterior auris muscles, &c. Before its bifurcation it gives off the *stylo-mastoid artery*. The *Anterior and Posterior Auricular Veins* open into the temporal and external jugular.

AURICULAR FINGER, (F.) *Doigt auriculaire*, is the little finger, so called because, owing to its size, it can be more readily introduced into the *meatus auditorius*.

AURICULAR NERVES are several. 1. The *auricular branch*, *Zygomato-auricular*, is one of the ascending branches of the cervical plexus. It ramifies and spreads over the two surfaces of the pavilion. 2. The *auricular or superficial temporal*, *Temporal-cutaneous*—(Ch.), is given off from the inferior maxillary. It ascends between the condyle of the jaw and the *meatus auditorius externus*, sends numerous filaments to the *meatus* and pavilion, and divides into two twigs, which accompany the branches of the temporal artery, and are distributed to the integuments of the head. There is also a *posterior auricular* furnished by the facial.

AURICULARIA SAMBUCCI, *Peziza auricula*.

AURICULARIS ANTERIOR, *Anterior auris*—a. *Superior*, *Attollens aurum*.

AURICULE, *Auricle*, *Pavilion of the ear*.

AURICULO-VENTRICULAR, *Atrio-ventricular*, *Atrio-ventriculæ*, *Auriculo-ventri-*

culæ, (F.) *Auriculo-ventriculæ*. That which belongs to the auricles and ventricles of the heart. The communications between the auricles and ventricles are so called. The *Tricuspid and Mitral Valves* are auriculo-ventricular valves.

AURIGA. A species of bandage for the ribs, described by Galen. See, also, *Liver*.

AURIGO, *Icterus*—a. *Neophytorum*, *Icterus Infantum*.

AURIPIGMENTUM, *Orpiment*—a. *Rubrum*, *Realgar*.

AURIS, *Ear*.

AURISCALPIUM, *Earpick*.

AURISCOPEIUM, *Auriscope*, from *auris*, 'the ear,' and *σκοπεω*, 'I view.' An instrument for exploring the ear.

AURIST, *Otiater*, *Otiatrus*, *Ear-doctor*, *Ear-surgeon*; from *auris*, 'the ear.' One who occupies himself chiefly with the diseases of the ear and their treatment.

AURIUM FLUCTUATIO, *Bombus*—a. *Marmorata*, *Cerumen*—a. *Sibulus*, *Bombus*—a. *Sonitus*, *Bombus*—a. *Sordes*, *Cerumen*—a. *Susurrus*, *Bombus*.

AURONE, *Artemisia abrotanum*—a. *des Champs*, *Artemisia campestris*—a. *des Jurdins*, *Artemisia abrotanum*—a. *Mâle*, *Artemisia abrotanum*.

AURUGO, *Icterus*.

AURUM, *Gold*—a. *Chloratum*, *Gold*, *muriate of*—a. *Chloratum natronatum*, see *Gold*—a. *Foliatum*, *Gold leaf*—a. in *Libellis*, *Gold leaf*—a. *Leprosum*, *Antimonium*—a. *Limatum*, see *Gold*—a. *Muriaticum*, see *Gold*—a. *Muriaticum natronatum*, see *Gold*.

AURUM MUSIVUM, *Aurum Mosaicum*, *Sulphuret of Tin*, *Deutosulphuret* or *Persulphuret of tin*. (*Quicksilver*, *tin*, *sulphur*, *sal ammoniac*, $\bar{a}\bar{a}$, equal parts. The tin being first melted, the quicksilver is poured into it, and then the whole are ground together, and sublimed in a bolt-head. The *aurum musivum* lies at the bottom.) It is used in some empirical preparations.

AURUM OXYDATUM, see *Gold*—a. *Oxydulatum muriaticum*, *Gold*, *muriate of*—a. *Nitro-muriaticum*, see *Gold*—a. *Salitum*, *Gold*, *muriate of*.

AUSCULTATE, from *auscultare*, 'to listen.' To practise auscultation. '*To auscult*' is at times used with the same signification.

AUSCULTATION, *Auscultatio*, *Echoscopé*, act of listening. Buisson has used it synonymously with *listening*. Laënnec introduced *auscultation* to appreciate the different sounds which can be heard in the chest; and in the diagnosis of diseases of the heart, lungs, &c. This may be done by the aid of an instrument called a *stethoscope*, one extremity of which is applied to the ear, the other to the chest of the patient. This mode of examination is called *Mediate Auscultation*, (F.) *Auscultation médiate*,—the application of the ear to the chest being *immediate auscultation*.

The act of exploring the chest is called *Stethoscopy*, *Thoracoscopy*; of the abdomen, *Abdominoscopy*.

AUSCULTATORY, *Auscultatorius*; *Auscultatory*, *Auscultic*, (with some.) Belonging or having relation to auscultation.

AUSCULTATORY PERCUSSION, see *Acouphonia*.

AUSTERE, *Austerus*. Substances which produce a high degree of acerb impression on the organs of taste.

AUSTRUCHE, *Imperatoria*.

AUTALGIA DOLOROSA, *Neuralgia*, *facial*, *Pleurodynia*—a. *Pruriginosa*, *Itching*—a. *Vertigo*, *Vertigo*.

AUTARCTIA, from *avros*, 'himself,' and *αρκτω*, 'I am satisfied.' Moral tranquillity.—Galen.

AUTEMESIA, from *avros*, 'self,' and *εμεσις*,

'vomiting.' Spontaneous or idiopathic vomiting.—Alibert.

AUTEMPRESMUS, Combustion, human.

AUTHEMERON. A medicine which cures on the day of its exhibition; from *avros*, 'the same,' and *ἡμερα*, 'day.'

AUTHYGIANSIS, Vis medicatrix naturæ.

AUTOCHIR, *Autochirus*, *Suicida*, from *avros*, 'himself,' and *χειρ*, 'hand.' One who has committed suicide. A self-murderer or suicide.

AUTOCHIRIA, Suicide.

AUTOCINE'SIS, *Motus voluntarius*, from *avros*, 'self,' and *κίνησις*, 'motion.' Voluntary motion.

AUTOCRASY, *Autocrati'a*, *Autocrato'ria*, from *avros*, 'himself,' and *κρατος*, 'strength.' Independent force. Action of the vital principle, or of the instinctive powers, towards the preservation of the individual. See Vis Medicatrix Naturæ. Also, the vital principle.

AUTOCRATIA, Autocracy, Vis Medicatrix Naturæ.

AUTOCRATORIA, Autocracy—a. Physiatrix, Vis medicatrix naturæ.

AUTOCTONIA, Suicide.

AUTOGENOUS, from *avros*, 'self,' and *γεννᾶω*, 'I generate.' A term applied by Mr. Owen to parts or elements that are usually developed from distinct and independent centres; as in the case of the different parts or elements that form a vertebra.

AUTOGENIA, Generation, equivocal.

AUTOLITHOTOMUS, from *avros*, 'himself,' *λίθος*, 'a stone,' and *τομεῖν*, 'to cut.' One who operates upon himself for the stone.

AUTOMAT'IC, *Automat'icus*, *Autom'atus*, (F.) *Automatique*, from *avtomatos*, 'spontaneous.' That which acts of itself. Those movements are called automatic which are executed without volition:— involuntary motions, *motus automat'ici* seu *auton'ati* seu *involuntarii*.

AUTOMNAL, Autumnal.

AUTONOM'IA, *Vis medica'trix natu'ræ*. The word *Autonomia* is occasionally employed by the French and Germans for the peculiar mechanism of an organized body. Thus, although individuals of the same species may differ in outward conformation, their mechanism or instinctive laws (*Autonomia*) may be the same.

AUTONYCTOBATIA, Somnambulism.

AUTOPEP'SIA, from *avros*, 'self,' and *πεπῶ*, 'I concoct.' Self-digestion,—as of the stomach after death.

AUTOPHIA, Autopsia.

AUTOPH'ONY, *Autopho'nia*, (F.) *Autophonie*, *Retentissement autophonique*, from *avros*, 'self,' and *φωνή*, 'voice.' An auscultatory sign pointed out by M. Hourmann, which consists in noting the character of the observer's own voice, while he speaks with his head placed close to the patient's chest. The voice, it is alleged, will be modified by the condition of the subjacent organs. The resonance, thus heard, he terms *retentissement autophonique*. This diagnostic agency Dr. K. G. Latham proposes to term *heautophon'ica*.

AUTOPHONIA, Suicide.

AUTOPOSPHORUS, Phosphorus.

AUTOPLAS'TIC, *Autoplas'ticus*, from *avros*, 'self,' and *πλαστικός*, 'formative.' Relating to autoplasty or plastic surgery.

AUTOPLASTICE, Morioplastice.

AUTOPLASTY, Morioplastice.

AUTOP'SIA, *Autopsy*, from *avros*, 'himself,' and *opsis*, 'vision.' *Autoph'ia*, *Autoscop'ia*. Inspection; examination by one's self; self-inspection. Often improperly used for the following:

AUTOP'SIA CADAVER'ICA, (F.) *Autopsie* ou *Ouverture cadavérique*. Attentive examination after death,—*Examination post mortem*, *Sec'tio Cadav'*

eris, *Dissection*, *Nec'roscopy*, *Nec'ropepy*, *Necroscop'ia*, *Necrop'sia*, *Necrop'sis*,—practised for the purpose of investigating the causes and seat of an affection of which a person may have died, &c.

AUTOP'SIA CADAVER'ICA LEGA'LIS, *Sec'tio* seu *Obduc'tio lega'lis*, is the examination after death for medico-legal purposes.

AUTOPYROS, Synecismos.

AUTOSCOPIA, Autopsia.

AUTOSITE, from *avros*, 'self,' and *σιτος*, 'nourishment.' A single monster, capable of deriving nourishment from its own proper organs, in contradistinction to *Omphalosite*.

AUTOTHERAPIA, Vis medicatrix naturæ.

AUTUMN, *Autumn'us*, *Auctum'nus*, from *augere*, *auctum*, 'to increase' [?], *Phthiro'pon*, (F.) *Automne*. One of the seasons of the year, between the 23d of September and the 21st of December. In all climates, the Autumn or Fall is liable to disease; a combination of local and atmospheric causes being then present, favourable to its production.

AUTUM'NAL, *Autumna'lis*, (F.) *Autumnal*. Relating to Autumn; as *Autumnal Fruits*, *Autumnal Fevers*, &c.

AUTUMNAL FEVER generally assumes a bilious aspect. Those of the intermittent kind are much more obstinate than when they appear in the spring.

AUXESIS, Augmentation, Increase.

AUXILIARY, *Auzilia'ris*, from *auxilium*, 'aid.' (F.) *Auxiliaire*. That which assists, or from which assistance is obtained.

AUXILIARY MEDICINE is one which assists the principal medicine or basis. It is synonymous with *Adjuvant*.

AUXILIARY MUSCLES are those which concur in the same movement. Some anatomists have applied the term to several ligaments, as well as to the fleshy fibres, which hang from the *sacrospinalis* muscle.

AUXILIUM, Juvans, Medicament.

AUXIS, Augmentation, Increase.

AVA, *Arva*, *Kava*. An intoxicating narcotic drink, made by chewing the Piper methisticum. It is much used by the Polynesians.

AVAILLES, WATERS OF. A small village in France, 13 leagues S. S. E. of Poitiers, at which there is a cold saline chalybeate. It contains chlorides of sodium and calcium, sulphate and subcarbonate of soda, iron, &c.

AVANT-BOUCHE (F.), *Os anti'cum*. This name has been applied by some to the mouth, properly so called—in contradistinction to the *Arrière bouche* or *Pharynx*.

AVANT-BRAS, Fore-arm.

AVANT-CŒUR, Scrobiculus cordis.

AVANT-GOUT (F.), *Prægusta'tio*. A fore-taste; prægustation.

AVANT-MAIN (F.), *Adver'sa Manus*. The inside of the hand, when extended.

AVANT-PIED (F.). The most advanced part of the foot.

AVANT-POIGNET (F.). The anterior part of the wrist.

AVELINE, *Corylus avellana*, (nut.)

AVELLANA, *Corylus avellana*—a. Cathartica, *Jatropha curcas*.

AVE'NA, *Bromos*, *Oats*, (Prov., North of England) *Aits*. The seeds of *Ave'na sativ'a*. Ord. Gramineæ. *Sex. Syst.* Triandria Digynia. (F.) *Avoine*. Oats are used as food for man in some parts, particularly in the North of England and Scotland. When deprived of the husks, they form *Groats*. Reduced to meal—*Avene Fari'na*, *Oatmeal*, (Ph. U. S.)—they are applied as cataplasms to promote suppuration. The dry meal is sprinkled over erysipelatous parts.

Oatmeal gruel. *Water gruel*, is prepared as follows:—Take of *oatmeal*, 5j; *soft water*, Oiss. Rub the meal in a basin, with the back of a spoon, in a moderate quantity of the water, pouring off the fluid after the grosser particles have subsided, but whilst the milkiness continues; and repeat the operation until no more milkiness is communicated to the water. Put the washings in a pan, after having stirred them well, in order to suspend any fecula which may have subsided; and boil until a soft, thick mucilage is formed.

It is a good demulcent, and is used also as a vehicle for clysters.

AVENA EXCORTICATA, Groats.

AVENÆ FARINÆ, see *Avena*.

AVENHEIM, MINERAL WATERS OF. Avenheim is three leagues from Strasburg: near it is an aperient mineral water.

AVENNES, MINERAL WATERS OF. Avennes is a village in the department of Hérault, in France: near it is a saline spring, the temperature of which rises to 84° Fahrenheit.

AVENS, COMMON, *Geum urbanum*—a. Purple, *Geum rivale*—a. Water, *Geum rivale*—a. White, *Geum Virginianum*—a. Yellow, *Geum urbanum*.

AVERICH, Sulphur.

AVERRHO'A BILIM'BI, *Bilim'bi*, *Bilimbing teren*. Ord. Oxalidæ: called after Averrhoës. An Indian tree, which has a fruit that is too acid to be eaten alone. It is used as a condiment, and in the form of syrup as a refrigerant.

AVERRHO'A CARAM'BOLA, *Malum Cœn'sœ*, *Prunum stellatum*, *Tamarca*, *Conga*, *Caram'bola*. An Indian tree, whose fruits are agreeably acid. The bark, bruised, is employed as a cataplasm, and its fruit is used as a refrigerant in bilious fever and dysentery.

AVERSION, *Aver'sio*, *Apo'tropē*, from *averte*, (*a* and *verte*), 'to turn from.' Extreme repugnance for anything whatever.

AVERSION (F.) also means, in therapeutics, the action of medicines which turn the afflux of fluids from one organ, and direct them to others; being synonymous with *counter-irritation*, or rather, *revulsion* or *derivation*.

AVERTIN (F.). A disease of the mind, which, according to Lavoisier, renders the patient obstinate and furious.

AVEUGLE, Cæcus.

AVEUGLEMENT, Cæcitas—*a. de Jour*, *Nyctalopia*—*a. de Nuit*, *Hemeralopia*.

AVICEN'NIA TOMENTO'SA, *A. Afric'na* seu *resinif'era* seu *nit'ida*, *Bont'ia ger'minans*: called after Avicenna. Ord. Verbenacæ. The plant which affords the *Malac'ca Beun* or *Anacard'ium Orient'al'e* of the Pharmacopœias, *Semecarpus Anacard'ium*. The oil drawn from the bark of the fruit is a corrosive and active vesicatory, but it is not used.

AVICULA CIPRIA, *Patill*—*a. Margaritifera*, see *Pearl*.

AVOIN, *Avena*.

AVON, MINERAL WATERS OF. Avon Springs are in the State of New York, on the eastern branch of the Genesee river, 18 miles from Rochester. There are three sulphureo-saline and one iodine spring. They are applicable to the same set of cases as the Sharon Springs.

AVORTEMENT, Abortion—*a. Provoqué*, see *Parturition*.

AVORTER, to Abort.

AVORTIN, Abortion.

AVORTON, Abortion.

AVULSIO, *Arrachement*.

AVULSION, Evulsion.

AX, MINERAL WATERS OF. Ax is a small town in the department of Arriège, France; where there are several sulphurous springs, the

temperature of which varies from 77° to 162° of Fahrenheit.

AXE, Axis—*a. de l'Œil*, Axis of the eye.

AXE'A COMMISSU'RA, *Trochoides*. A pivot-joint. See *Trochoid*.

AXEN Körper, *Corpuscula tactūs*.

AXES, Intermittent, *Paroxysm*.

AXILE BODIES, *Corpuscula tactūs*.

AXIL'LA, *Ala*, *Assel'la*, *Assell'la*, *Assill'la*, *Acel'la*, *Cordis emuncto'rium*, *Malē*, *Hypom'nia*, *Fol'lea axilla'ris*, *Maschale*, *Maschalis*. (Sc. and Prov.) *Oxtar*, *Oxter*, (F.) *Aisselle*. The cavity beneath the junction of the arm with the shoulder; the *armpit*; (F.) *Crence de l'Aisselle*. It is bounded, anteriorly, by a portion of the pectoralis major; posteriorly, by the latissimus dorsi. It is covered with hair, contains much areolar membrane, lymphatic ganglions, important vessels and nerves, and numerous sebaceous follicles, furnishing an odorous secretion. In consequence of such secretion, the ancients called it *emuncto'rium cordis*.

AXILLARY, *Maschali'us*, (F.) *Aillaire*, from *axilla*, 'the armpit.' Belonging to the armpit.

AXILLARY ARTERY, *Artē'ria axilla'ris*; a continuation of the subclavian, extending from the passage of the latter between the scaleni muscles as far as the insertion of the pectoralis major, when it takes the name of *Brachial*.

AXILLARY GLANDS are lymphatic glands seated in the armpit; into which the lymphatic glands of the upper extremity open.

AXILLARY NERVE, *Cir'conflex N. Scapulo-hum'eral* (CH.), *Nerf circonflexe*, *Articular nerve*; arises from the posterior part of the brachial plexus, particularly from the last two cervical pairs and the first dorsal. It is chiefly distributed to the posterior margin of the deltoid.

AXILLARY VEIN, *Vena Axilla'ris*, *Vena Subala'ris*. This vein corresponds with the artery: anterior to which it is situate. It is a continuation of the *brachial veins*; and, at its termination, assumes the name *Subclavian*.

AXINE, *Ascia*.

AXIRNACH. An Arabic word, used by Albucasis to designate a fatty tumour of the upper eyelid, observed particularly in children.

AXIS, *Axon*, (F.) *Axe*. A right line which passes through the centre of a body.

AXIS. CEREBRO-SPINAL, see *Encephalon*—*a. of the Cochlea*, *Modiolus*—*a. Cranio-Spinal*, see *Encephalon*—*a. Cylinder of Nerve*, see *Nerve fibre*—*a. Cœliac*, *Cœliac artery*.

AXIS OF THE EYE, (F.) *Axe de l'œil*, called also, *Vis'ual Axis* and *Optic Axis*, is a right line, which falls perpendicularly on the eye, and passes through the centre of the pupil.

AXIS, HÆMAL, *Aorta*—*a. Neural*, see *Encephalon*.

AXIS, is also the second vertebra of the neck, *Axon*, *Epistroph'us*, *Epistrophus*, *Maschali'us*: *Ver'tebra Denta'ta*, (F.) *Essieu*. So called, because it forms a kind of axis on which the head moves. Chaussier calls it *Axoïde*, from *axōn*, 'axis,' and *eidōs*, 'shape.'

AXLETEETH, *Molar teeth*.

AXOIDE, Axis—*a. Occipitale*, *Rectus capitis posticus major*.

AXOIDO-ATLOIDEUS. What refers to both the axis and atlas, as *Axoido-atloidean* articulation.

The lesions of the Axoido-atloidean, are, 1. Fracture of the *Processus Denta'tus*. 2. Rupture of the odontoid ligament, and consequently passage and pressure of the process behind the transverse ligament: and 3. The simultaneous rupture of the odontoid and transverse ligaments. These different accidents are fatal.

AXOIDO-ATLOIDIEN, *Obliquus inferior capitis*.

AXON. Axis.

AXONGE, Adeps præparatus.

AXUNGE, Adeps præparatus.

AXUNGIA, Pinguedo—a. Gadi, Oleum Jecoris Aselli—a. de Mumiâ, Marrow—a. Articularis, Synovia—a. Piscina Marina, Oleum Jecoris Aselli—a. Porcina, Adeps præparatus.

AYPNIA, Insomnia.

AZALEA PROCUMBENS, Loiseleuria procumbens.

AZARNET, Orpiment.

AZARUM, Asarum—a. Cabaret, Asarum.

AZEDARACH, Melia Azedarach.

AZEDARACHA AMENA, Melia Azedarach.

AZO'IC, *Azo'icus*, same etymon as Azote. Devoid of life. The "*Azoic period*" of the geologist is that before any living being appeared.

AZOODYNA'MIA, from *a*, priv., *ζωη*, 'life,' and *δυναμις*, 'strength.' Privation or diminution of the vital powers.

AZORES, CLIMATE OF. The Azores or Western Islands are said to afford one of the best examples of a mild, humid, equable climate, to be met with in the northern hemisphere. It is slightly colder and moister than that of Madeira, but even more equable. Sir James Clark thinks, that a change from the Azores to Madeira, and thence to Teneriffe—one of the Canaries—would prove more beneficial to the phthisical valetudinarian than a residence during the whole winter in any one of those islands.

AZOTAS HYDRARGYRICUS LIQUIDUS, see Hydrargyri nitras—a. Hydrargyros—ammonic, see Hydrargyri oxydum cinereum—a. Hydrargyrosus, Hydrargyri nitras—a. Potassicus, Potassæ nitras.

AZOTATE, Nitrate—a. d'Argent, Argenti nitras—a. de Mercure, Hydrargyri nitras—a. de Mercure et d'ammoniaque, see Hydrargyri oxydum cinereum—a. de Plomb, Plumbi nitras—a. de Potasse, Potassæ nitras.

A'ZOTE, *Azo'tum*, from *a*, priv., and *ζωη*, 'life.' Nitrogen, Alcaligene, Gas azo'ticum, Nitrogen'tium, Seyton, (F.) Azote, Nitrogène, Mufette, Air gâté, Air vicié, is a gas which is unfit for respiration. It is not positively deleterious, but proves fatal owing to the want of oxygen. It is one of the constituents of atmospheric air, and a distinguishing principle of animals. Vegetables have it not generally diffused, whilst it is met with in most animal substances. It has been variously called, *phlogistic air*, *vitiated air*, &c.; has been looked upon as sedative, and recommended to be respired, when properly diluted, in diseases of the chest.

AZOTE, PROTOXIDE OF, Nitrogen, gaseous oxide of.

AZOTÉ, Nitrogenized.

AZOTED, Nitrogenized.

AZOTENÈNES, from *azote*, and *νεσος*, 'disease.' Diseases fancied to be occasioned by the predominance of azote in the body.—Baumes.

AZOTIC ACID, Nitric acid.

AZOTIZED, Nitrogenized.

AZOTURIA, see Urine.

AZTEC CHILDREN, see Nanus.

AZUR, Coral, Smalt.

AZU'RRIUM. A compound of two parts of mercury, one-third of sulphur, and one-fourth of sal ammoniac.—Albertus Magnus.

AZ'YGES, *Az'ygus*, *Az'ygous*, *sine pari*, from *a*, priv., and *ζυγος*, 'a yoke', not paired. The *sphenoid bone*, because it has no fellow. Also, a process, *Processus Az'ygus*, *Rostrum sphenoidale*, projecting from under the middle and forepart of this bone.

AZYGOS GANGLION, see Trisplanchnic Nerve.

AZYGOS ARTICULAR ARTERY, see Articular arteries of the skull.

AZYGOS MUSCLE, *Az'ygus Uvula*, is the small muscle which occupies the substance of the uvula.—Morgagni. The name is, however, inappropriate, as there are two distinct fasciculi, placed along-side each other, forming the *Palato-staphylini*, *Staphylini* or *Epistaphylini* muscles, *Staphylini medii* of Winslow.

AZYGOS PROCESS, OF THE SPHENOID, see Azyges.

AZYGOS VEIN, *Vena Azygos*, *Veine Prélobthoracique*—(Ch.), *Vena sine pari*, *Vena paricaris*, (F.) *Veine sans Paire*. This vein was so called by Galen. It forms a communication between the *V. cava inferior* and *V. cava superior*, permitting the blood to pass freely between the two. It rises from the vena cava inferior, or from one of the lumbar or renal veins, passes through the diaphragm, ascends along the spine to the right of the aorta and thoracic duct, and opens into the *V. cava superior*, where it penetrates 'the pericardium. On the left side, the *SEMI-AZ'YGOS*, *Left bronchial* or *left superior intercostal vein*, *Vena demi-azygos*, *V. hemi-az'ygæ*, *Veine petite prélobthoracique*—(Ch.), presents, in miniature, nearly the same arrangement.

AZYMIA HUMORUM, Crudity of the humours.

AZYMUS, from *a*, priv., and *ζυμη*, 'leaven.' Azymous bread is unfmented, unleavened bread.—Galen.

AZZLE-TEETH, Molar teeth.

B.

BABEURRE, Buttermilk.

BABILLEMENT, Loquacity.

BABUZICARIUS, Incubus.

BABY, Infans.

BAC'ARIS, *Bach'aris*. A name given by the ancients to an ointment, described by Galen under the name *Ointment of Lydia*. It was sometimes employed in diseases of the womb.—Hippocrates.

BACCÆ BERMUDENSES, *Sapindus saponaria*—h. seu *Grana actes*, see *Sambucus ebulus*—b. *Jujubæ*, *Jujube*—b. *Myrtillorum*, see *Vaccinium myrtillus*—b. *Norlandicæ*, *Rubus arcticus*

—b. *Piperis Glabri*, see *Piper Cubeba*—b. *Piscatoricæ*, see *Menispermum cocculus*—b. *Zizyphi*, see *Jujube*.

BACCAR, *Bac'caris*, *Bac'charis*. An herb used by the ancients in their garlands, to destroy enchantment. Perhaps, the *Digitalis purpurea*: Some authors have erroneously thought it to be the *Asarum*.

BACCHARIS, *Baccar*.

BAC'CHARIS HALIMIFOLIA, *Groundsel tree*; *Order*, *Compositæ*; indigenous; is used as a demulcent to allay cough, in the form of decoction.

BACCH'IA, from *Bacchus*, 'the god of wine.'

A name applied to the red or pimpled face of the drunkard. See *Gutta rosea*.

BACCHICA, *Hedera helix*.

BACCIVOROUS, *Baccivorus*, (F.) *Baccivore*, from *bacca*, 'a berry,' and *voro*, 'I devour.' Living on berries.

BACHARIS, *Bacaris*.

BACHELOR'S BUTTONS, see *Strychnos nuxvomica*.

BACHER'S TONIC PILLS, *Pilule ex Hellebore et Myrrha*.

BACILE, *Crithmum maritimum*.

BACILLAR LAYER OF THE RETINA, *Tunica Jacobi*.

BACILLUM, *Bacillus*, *Baculus*, *Baculus*: 'a stick.' This name has been applied to a kind of troch, composed of expectorants, and having the shape of a stick. Also, a suppository. *Bacillum* was used by the ancient chemists for several instruments of iron.

BACK, *Dorsum*.

BACK-ACH ROOT, *Liatris*.

BACKBONE, Vertebral column.

BACKSIDE, *Nates*.

BACKSPRENT, Vertebral column.

BACKSTROKE OF THE HEART, Impulse, diastolic.

BACOBOTRYS PICTA, *Saoria*.

BACOVE, *Musa sapientum*.

BACTYRLOBIUM FISTULA, *Cassia fistula*.

BACULUS, *Bacillum*.

BAD, Sick.

BADEN, MINERAL WATERS OF. Baden is a town six miles from Vienna. Here are 12 springs, containing carbonates of lime and magnesia; sulphates of lime, and magnesia, and soda; and chlorides of sodium and aluminum. The water is used in diseases of the skin, rheumatism, &c.

There are two other towns of the same name; one in Suabia, and the other in Switzerland, about 12 miles from Zürich, where are mineral springs. The waters of the last two are thermal sulphureous.

BADEN-BADEN, MINERAL WATERS OF. Celebrated thermal springs, situate about a league from the high road to Basle and Frankfort. Their temperature varies from 130° to 154° Fahrenheit. Their situation is beautiful, and they are much frequented.

BADENWEILER, MINERAL WATERS OF. B. is in Baden, about a league to the westward of Müllheim. The waters are thermal (87° of Fahr.), and contain carbonate of lime, sulphuret and chloride of calcium, and chloride of magnesium.

BADER, Bather.

BADIAGA. A kind of sponge, sold in Russia, the powder of which is said to take away the livid marks from blows and bruises in a few hours. Its nature is not understood.

BADIANE, *Illicium anisatum*.

BADISIS, Walking.

BADLY, Sick.

BADUKKA, *Capparis badukka*.

BAG, DUSTING, see *Dusting-bag* — b. of Waters, see *Liquor Amnii*.

BAGEDIA, Pound.

BAGGIE, Abdomen.

BAGNÈRES-ADOUR, MINERAL WATERS OF. Bagnères-Adour is a small town in the department of *Hautes Pyrénées*, having a great number of mineral springs; some, cold chalybeates; others, thermal salines; but the greatest part sulphureous and warm.

BAGNÈRES DU LUCHON is a small town in the department of *Haute Garonne*, on the frontiers of Spain. It has been for a long time

famous for its numerous sulphureous springs, the temperature of which is from 69° to 148° of Fahrenheit.

BAGNIGGE WELLS. A saline mineral spring of London, resembling the Epsom.

BAGNIO, *Baignoire*.

BAGNOLES, MINERAL WATERS OF. Bagnoles is a village in the department of Orne. The water resembles that of *Baignères de Luchon*.

BAGNOLS, MINERAL WATERS OF. Bagnols is a village, two leagues from Meude, in the department of Lozère. The waters are hydrosulphurous and thermal: 109° Fahrenheit.

BAGOAS, *Castratus*.

BAGUENAUDIER, *Colutea arborescens*.

BAHA'MA ISLANDS, CLIMATE OF. The climate of the Bahamas is not considered to be well adapted for consumptive patients, on account of the rapid alternations of temperature, and the prevalence of winds, often of a dry, cold character. Still, the phthisical valetudinarians from most portions of the United States might derive advantage from a residence there during the winter months. The accommodations are not, however, good, or numerous.

BAHEL, *Columnea longifolia*. A labiated plant of Malabar, whose leaves, bruised, are applied as cataplasms to suppurating tumours.

BAHEL SCHULLI, *Genista spinosa Indica*.

BAIGNEUR, Bather.

BAIGNOIRE, (F.) *Baptistarium*, a Bathing tub, *Bagnio*, *Solium*, *Piscina*. The vessel or place in which bathing is performed. *Baignoire oculaire*, an eye-bath, — a small vessel for bathing the eyes. See *Scaphium oculare*.

BAILEY'S SPRING, see Alabama, Mineral Waters of.

BAILLEMENT, Yawning.

BAILLON, Speculum oris.

BAIN, Bath — b. *Chaud*, Bath, hot — b. *Électrique*, Bath, electric, see *Electricity* — b. *Entier*, Bath, general — b. *de Fauteuil*, Bath, hip — b. *Frais*, Bath, tepid — b. *Froid*, Bath, cold — b. *Hygiénique*, see Bath — b. *Marie*, Bath, water — b. *Médicinal*, Bath, medicated — b. *de Mer*, Bath, sea — b. *de Pied*, Bath, foot, Pediluvium — b. *de Sable*, Bath, sand — b. *de Siège*, Bath, hip — b. *Simple*, see Bath — b. *Tempéré*, Bath, tepid, B. Temperate — b. *de Tête*, Bath, head — b. *Tide*, Bath, tepid — b. *Très froid*, Bath, cold — b. *de Vapeur*, Bath, vapour.

BAINE, Bath.

BAINS, MINERAL WATERS OF. These are situated at Plombières, department of the Vosges. They are said to be saline and thermal by some; others deny them any medical properties.

BAIRN, Infans.

BAIRNWORTS, *Bellis*.

BALAMPULLI, *Tamarindus*.

BALANCE, AEROSTATIC, *Areometer*.

BALANCEMENT, *Compensation*, from (F.) *balance*, 'a balance,' itself from *bis*, 'twice,' and *lanx*, 'a dish.' A law of teratogeny, as maintained by Geoffroy St. Hilaire, by which exuberance of nutrition in one organ is supposed to involve, to a greater or less extent, the total or partial atrophy of some other, and conversely.

BALANDA, *Fagus Sylvatica*.

BALANEUM, Bath.

BALANISMUS, Suppository.

BALANITE CONTAGIEUSE, see Gonorrhoea.

BALANITIS, Gonorrhoea spuria.

BALANOBLENNORRHOEA, Gonorrhoea spuria.

BALANOCASTANUM, *Bunium Bulbocastanum*.

BALANORRHAGIA, see Gonorrhoea.

BALANORRHŒA, *Gonorrhœa spuria*.

BALANOS PHENICOS, Date.

BALANUS, *βαλανός*, 'glands,' 'an acorn.' The glands penis. Hence, *Balanoblenorrhœa*, *Blennorrhœa* of the glands; and *Balanitis*, Inflammation of the glands. Suppositories and pessaries were called *Balani*.

BALANUS, Glands, Suppository—b. Myrepica, *Guilandina moringa*.

BALARUC, MINERAL WATERS OF. *Balaruc* is a town in the department of Hérault, in France. The waters are saline and thermal. They contain carbonic acid, carbonate of lime, carbonate of magnesia, chlorides of sodium, calcium, and magnesium, sulphate of lime, and a little iron. They are considered tonic, and are largely used. Their temperature is about 118° Fahrenheit.

BALARUC WATER, *FACITIVIOUS*, (F.) *Eau de Balaruc*; *Aqua Bellilucana* is made of simple acidulous water (containing twice its bulk of carbonic acid) f 3ssss; chloride of sodium, 5iss; chloride of calcium, gr. xvij; chloride of magnesium, gr. lvi; carbonate of magnesia, gr. j.

BALATRO, *Bambatio*.

BALAUSTINE FLOWERS, see *Punica granatum*.

BALBIS, *βαλβίς*, 'a foundation.' Any oblong cavity.—Galen. Hippocrates, in his treatise on the joints, gives the name *Balbittodes* to the olecranon cavity of the humerus.

BALBUS, (F.) *Begue*. One habitually affected with stammering. A stammerer.

BALBUTTEMENT, *Bulbuties*.

BALBUTIES, *Paellismus*, *Paellotes*, *Blæstias*, *Bargylusis*, *Dysphasia*, *Mogilasia*, *Ischophonia*, *Battariusmus*, *Bombasia*, *Hesitatio*, *Loquelu blasia*, *Timbultio Lingue*, (F.) *Bulbutement*, *Bégaiement*, *Bégagement*, Stuttering, Stammering, Hammering, (Prov.) *Tutt'ering*, *S. Vitis's Dance of the Voice*, (Sc.) *Hab'bering*, *Halbling*. Also, vicious and incomplete pronunciation, in which almost all the consonants are replaced by the letters B and L; *Trantismus*.

BALCHUS, *Bellium*.

BALD, *Athrix*.

BALDMONEY, *Æthusa meum*.

BALDNESS, *Alpecia*, *Calvities*—b. Limited, *Porrigo decalvans*—b. Partial, *Porrigo decalvans*.

BALENAS, *Leviathan* penis.

BALIMBAGO, *Hibiscus populeus*.

BALINEATOR, *Bather*.

BALINEUM, *Bath*.

BALL, *Pila*—b. of the Eye, *Pupil*.

BALLISMUS, *Chorea*.

BALLISTA, *Astragalus*.

BALLOCK-GRASS, *Orchis mascula*.

BALLON, *Receiver*.

BALLONNEMENT, *Tympanites*.

BALLOTA FETIDA, *B. vulgaris* seu *nigra*, *Marrubium nigrum*, *Black Horehound*, *Stinking H.*, (F.) *Marrube noir*. *Ord. Labiate. Sez. Syst. Didynamia Gymnospermia*. This plant is esteemed to be antispasmodic, resolvent, and detensive. (?)

BALLOTA LANA'TA, *Leonurus lanatus*, (F.) *Ballote cotonneuse*. A plant which grows in Siberia. The whole plant, with the exception of the root, has been recommended in dropsy, and in rheumatism and gout, as a diuretic. It is usually given in decoction (℥ss to ℥ij to ℥viij of water).

BALLOTE COTONNEUSE, *Ballota lanata*.

BALLOTTEMENT, (F.) *Mouvement de Ballotement*, *Agitation*, *Succussion*, *Repercussion*, means the motion impressed on the fœtus in utero, by alternately pressing the uterus by means of the index finger of one hand introduced into the vagina; the other hand being

applied on the abdomen. It is one of the least equivocal signs of pregnancy.

BALLS, *DEER*, *Elaphomyces granulatus*.

BALLSTON SPA. This village is situate in Saratoga County, New York. The spring Sans Souci belongs to the class of Acidulous Chalybeates. It contains iodide of sodium. There is also a sulphur spring.

BALM, *Melissa*—b. Apple, *Momordica balsamina*—b. Bastard, *Melitis Melissophyllum*—b. of Gilead, *Solomon's*, see *Tinctura cardamomi*—b. of Gilead, *Poplar*, *Populus canadensis*—b. of Gilead tree, *Dracocephalum Canariense*—b. Indian, *Trillium latifolium*—b. Mountain, *Monarda coccinea*—b. Red, *Monarda coccinea*—b. Scarlet rose, *Monarda coccinea*—b. Stinking, *Hedeoma*.

BALMONY, *Chelone glabra*.

BALNEA CŒNOSA, *Boue des eaux*.

BALNEARIUM, *Hypocaustum*.

BALNEARIUS, *Bather*.

BALNEATOR, *Bather*.

BALNEOGRAPHY, *Balneographia*, from *βαλανεον*, 'a bath,' and *γραφη*, 'a description.' A description of baths.

BALNEOLLOGY, *Balneology*, from *βαλανεον*, 'a bath,' and *λογος*, 'a description.' A treatise on baths.

BALNEOTHERAPIA, from *βαλανεον*, 'a bath,' and *θεραπεια*, 'treatment.' Treatment of disease by baths.

BALNEUM, *Bath*—b. Acidum, *Bath*, acid—b. Alkalium, *Bath*, alkaline—b. Animale, *Bath*, animal—b. Antipsoicum, *Bath*, antipsoic—b. Anti-syphiliticum, *Bath*, antisiphilitic—b. Arenæ, *Bath*, sand—b. Gelatinosum, *Bath*, gelatinous—b. Mariæ, *Bath*, water—b. Marinum, *Bath*, sea—b. Mariæ, *Bath*, water—b. Medicatum, *Bath*, medicated—b. Sulphuris, *Bath*, sulphur.

BALSAM, *Balsamum*, *Balsamus*, *Bol'son*, *Del'son*, (F.) *Baume*. This name is given to natural vegetable substances, concrete or liquid, but very odorous, bitter, and piquant: composed of resin, benzoic acid, and sometimes of an essential oil—which allow benzoic acid to be disengaged by the action of heat; readily dissolved in volatile oil, alcohol, and ether; and, when treated with alkalies, afford a soluble benzoate, and throw down resin. We know of only five balsams—those of Peru, and Tolu, Benzoin, solid *Styrax* or *Storax*, and liquid *Styrax*. (See those different words.) There are, however, many pharmaceutical preparations and resinous substances, possessed of a balsamic smell, to which the name *balsam* has been given; but they differ essentially in composition and properties: hence the distinction of balsams into *natural* and *artificial*. The *natural balsams* include the five before mentioned; the *artificial* the remainder.

BALSAM, *ACOUSTIC*, *Balsamum Acousticum*, (F.) *Baume acoustique*. A mixture of fixed and essential oils, sulphur, and tinctures of fetid gums. Used in cases of atonic deafness, dropped into the ear. The *acoustic balsam* of Dr. Hugh Smith is made by mixing three drachms of *oz-gall*, with one drachm of *balsam of Peru*.

BALSAM, *AMERICAN*, see *Myroxylon Peruvianum*—b. Anodyne, *Bates's*, *Linimentum saponis et opii*.

BALSAM, *APOPLECTIC*, *Balsamum Apoplecticum*, *Balsamus Apoplecticus*, (F.) *Baume Apoplectique*. A medicine composed of several *balsams* properly so called, resins, and volatile oils. It is of a stiff consistence, is worn in ivory boxes about the person, and is smelled at in headaches, &c.

BALSAM APPLE, *Momordica balsamina*.

BALSAM OF ARCEUS, *Balsamum Arcei*, *Unguentum E'lemi*, (F.) *Baume d'Arcæus*. A soft ointment; sometimes employed in wounds, ul-

cers, &c. It is made by melting, with a gentle heat, two parts of mutton suet, one of lard, one and a half of turpentine, and as much resin.

BALSAM, CALABA, see *Fagara octandra*.

BALSAM, CANADA, see *Pinus balsamea*—b. Canary, *Draacocephalum Canariense*—b. Capivi, *Copaiba*.

BALSAM OF CARPA'THIA, *B. Carpathian*, *Bal'samum Carpathicum*, (F.) *Baume de Carpathie*. The resin of the *Pinus Cembra*, a tree, which grows in Switzerland, Libya, and the Krapac mountains in Hungary.

BALSAM, CHALYBEATE, *Bal'samum Chalybeatum*, (F.) *Baume d'aier ou d'aiguilles*. A mixture of nitrate of iron, alcohol, and oil, prepared by dissolving needles in nitric acid. It was formerly employed in friction in pains of the joints.

BALSAM, COMMANDER'S, *Tinctura benzoini composita*—b. for Cuts, *Tinctura benzoini composita*.

BALSAM, COR'DIAL, of SENNER'TUS, *Bal'samum Cordiale Senner'ti*, (F.) *Baume cordiale de Senner't*. A stimulant medicine, composed of the essential oils of citron, cloves, and cinnamon, of musk, and ambergris. Dose, 6 to 15 drops.

BALSAM OF FIERABRAS. A celebrated Spanish vulnerary balsam, mentioned by Cervantes; the composition of which was oil, rosemary, salt, and wine. (?)

BALSAM, SPIRITUOUS, OF FIORAVENTI, *Bal'samum Fioraventi spirituosum*, (F.) *Baume de Fioraventi spiritueux*. Different products of the distillation of resinous and balsamic substances, and of a number of aromatic substances, previously macerated in alcohol, have been thus called. The *Spirituous Balsam of Fioraventi*, the only one now used in friction, in chronic rheumatism, is the first product of the distillation from a sand-bath. It is entirely alcoholic. The *Oily Balsam of Fioraventi* is obtained by removing the residue, and distilling it in an iron vessel, at a white heat. It has the appearance of a citrine-coloured oil. The *Black Balsam of Fioraventi* is the black oil, obtained when the temperature is sufficient to carbonize the substances in the cucurbit.

BALSAM OF FIR, see *Pinus balsamea*.

BALSAM OF FOURCROY or of LABORDE, (F.) *Baume de Fourcroy ou de Laborde*. A kind of liniment composed of aromatic plants, balsams, resins, aloes, turpentine, theriac, and olive oil. Used in chaps of the skin and nipples.

BALSAM, FRIAR'S, *Tinctura benzoini composita*.

BALSAM OF GENEVIÈVE, (F.) *Baume de Geneviève*. An ointment composed of wax, turpentine, oil, red saunders, and camphor. Used in contused wounds, gangrene, &c.

BALSAM OF HONEY (HILL'S). A tincture made of *tolu*, *honey* (3*ss* ℥ij) and *spirit* (a gallon). A pectoral, used in coughs. The committee of the New York College of Pharmacy recommend the following formula—(*Gum Benzoin*. ℥v. *Bals. Tolut.* ℥j, *Mellis* ℥viij. *Alcohol.* Oij—digest for 10 days and filter). See *Mel*.

BALSAM OF HOREHOUND (FORD'S). A tincture of *horehound*, *liquorice-root*, *camphor*, *opium*, *benzoin*, *dried squills*, *oil of aniseed*, and *honey*. It has the same properties as the above. See *Marrubium*.

BALSAM, HUNGARIAN, see *Pinus mughos*.

BALSAM, HYPNOTIC, *Bal'samum Hypnoticum*, (E.) *Baume Hypnotique*. A preparation of which opium, hyoscyamus, camphor, and some other sedative substances, form the basis. It is used externally in friction, to provoke sleep.

BALSAM, HYSTERIC, *Bal'samum Hystericum*, (F.) *Baume Hystérique*. A preparation made of opium, aloes, asafetida, castor, distilled oils of *rae*, amber, &c. It is held to the nose, applied

to the navel, or rubbed on the hypogastrium in hysterical cases.

BALSAM, INDIAN, see *Myroxylon peruvianum*.

BALSAM, INDURET, *Bal'samum induratum*, (F.) *Baume hydrinduré*, *B. induré*, *Gelle contre le goître*. A balsam used in the way of friction, in goitre, at Lausanne, in Switzerland. It may be made as follows—*animal soap*, 60; *infuse of potassum*, 42; *alcohol* at 85°, 500; *essence of lemon*, 4 parts. The iodide is dissolved in the alcohol, which is added to the soap melted in a water-bath. The whole is then filtered, and put into bottles.

BALSAM OF LEICTOURE, OF CONDOM or VINGUÈRE, *Bal'samum Lectorale*. A strongly stimulant and aromatic mixture of camphor, saffron, musk, and ambergris, dissolved in essential oils. The ancients burnt it for the purpose of purifying the air of a chamber, when infected with a disagreeable odour.

BALSAM OF LIFE OF HOFFMANN, *Bal'samum Vita Hoffmanni*, (F.) *Baume de Vie d'Hoffmann*. A tincture, composed of essential oils and ambergris, employed internally and externally as a stimulant. A mixture of essential oils without alcohol constitutes the *Saxon Balsam*, *Bal'samum apoplecticum*, *B. aromat'icum*, *B. cephal'icum*, *B. Saxon'icum*, *B. nervi'um*, *B. SCHERZERI*, *B. Stomach'icum*. Employed in friction as a stimulant.

BALSAM OF LIFE, Decoction aloes compositum—b. of Life, Turlington's, see *Tinctura benzoini composita*.

BALSAM OF LOCATELI or LUCATELI, *Bal'samum Lucatel'i*, (F.) *Baume de Lucatel*. A sort of ointment, composed of wax, oil, turpentine, sherry, and balsam of Peru, coloured with red saunders. It was once administered in pulmonary consumption.

BALSAM OF MECCA, see *Amyris opobalsamum*—b. Mexican, see *Myroxylon Peruiferum*—b. Natural, see *Myroxylon Peruiferum*.

BALSAM, GREEN, OF METZ, *Bal'samum Viride Meten'sium*, *B. Viride*, (F.) *Baume vert de Metz*, *Baume de Feuillet*, *Huile verte*, *O'huile verte cupri viride*. This is composed of several fixed oils, holding, in solution, subcarbonate of copper, sulphate of zinc, turpentine, aloes, and the essential oils of cloves and juniper. It is green and caustic, and is employed to hasten the cicatrization of atonic ulcers.

BALSAM, NEPHRITIC, OF FULLER, *Bal'samum Nephriticum Fulleri*. A liquid medicine, composed of oils, resins, and balsams, which have experienced an incipient state of carbonization from concentrated sulphuric acid. It was given in the dose of 15 to 30 drops in certain affections of the kidneys.

BALSAM, NERVOUS, *Bal'samum Nervinum*, (F.) *Baume nervin ou nerveal*. A kind of ointment, composed of fatty bodies, volatile oils, balsam of Peru, camphor, &c. It is employed in friction in cases of sprains and rheumatic pains.

BALSAM, PARALYTIC, OF MYNSIGHT. A sort of liniment or soft mixture of the essential oils of different aromatic plants, oils of turpentine and amber.—Lémery.

BALSAM OF PAREIRA BRAVA, *Bal'samum Pareira bravae*. A soft mixture of balsam, resin, muriate of ammonia, and powder of the root of *Pareira brava*. It is given internally, to excite the urinary secretion.

BALSAM, PERUVIAN, see *Myroxylon Peruiferum*—b. of Peru, red, see *Toluifera balsamum*—b. of Peru, white, see *Myroxylon Peruiferum*.

BALSAM OF RACKASTRA or of RAKASTRI. This substance is of a yellowish-brown colour; semitransparent; fragile, when dry, but softening by heat; adhering to the teeth, when chewed. It

has a smell similar to that of the Balsam of Tolu, and is slightly bitter. It is brought from India in gourd shells, and has been employed in diseases of the urinary and genital organs, especially in gonorrhœa.

BALSAM, RIGA. Prepared from the shoots of the Scotch Fir, macerated in spirit of wine. *Internally*, stimulant and diuretic; *externally*, a vulnerary. See *Pinus Cembra*.

BALSAM OF SATURN, Balsamum Saturni. A solution of acetate of lead in spirit of turpentine, concentrated by evaporation; to which camphor has been added. This balsam was applied to hasten the cicatrization of wounds.

BALSAM OF THE SAMARITAN, (F.) Baume du Samaritain. A sort of liniment, prepared by boiling together, at a gentle heat, equal parts of wine and oil. It is said to have been the ointment used by the Samaritan of the Gospel to cure a patient covered with ulcers.

BALSAM, SAXON, Balsam of Life of Hoffmann.

BALSAM OF SULPHUR, Balsamum Sulphuris, (F.) Baume de Soufre. A solution of sulphur in oil.—*B. sulph. anisatum, (F.) B. de Soufre anisé.* A solution of sulphur in essential oil of aniseed; given as a carminative.—*B. Sulph. succinatam, (F.) B. de Soufre succiné.* A solution of sulphur in oil of amber.—*B. Sulphuris terebinthinæ, Common Dutch Drops, (F.) B. de Soufre térébinthine.* A solution of sulphur in essential oil of turpentine, administered as a diuretic.—The *Balsam of Sulphur of RULAND* is a solution of sulphur in linseed oil or nut oil.

BALSAM OF SYMPATHY, Balsamum Sympathicum, (F.) Baume de Sympathie. A balsam, used in the days when sympathetic influence was strongly believed in. It was composed of the raspings of a human skull, blood, and human fat, and was applied to the instrument which had inflicted the wound.

BALSAM, THIBAUT'S. A tincture of myrrh, aloes, dragon's blood, flowers of St. John's wort, and Chio turpentine. *Internally*, diuretic; *externally*, vulnerary.

BALSAM OF TOLU, see Toluifera Balsamum.

BALSAM, TRANQUIL, Balsamum tranquillum seu tranquillans, (F.) B. tranquille. A liquid medicine employed, externally, in the shape of friction: it is prepared by macerating and boiling, in olive oil, narcotic and poisonous plants,—belladonna, mandragora, hyoscyamus, &c.—and afterwards infusing, in the filtered decoction, different aromatic plants. It was employed as an anodyne.

BALSAM, TURKEY, Dracocephalum Canariense—*b. Turlington's*, see *Tinctura Benzoini composita*.

BALSAM OF TURPENTINE, Dutch Drops, Balsamum Terebinthinæ. Obtained by distilling oil of turpentine in a glass retort, until a red balsam is left. It possesses the properties of the turpentine.

BALSAM, VERRAIN'S, Tinctura Benzoini composita.

BALSAM, VULNERARY, OF MINDERER'S, Balsamum vulnerarium Mindereri, (F.) B. vulnéraire de MINDERER. A kind of liniment, composed of turpentine, resin elemi, oil of St. John's wort, and wax. Employed in friction, and as a dressing to wounds.

BALSAM WOOD, Impatiens fulva.

BALSAM WOUND, Tinctura Benzoini composita.

BALSAMADENDRON GILEADENSE, Amyris Gileadensis—*b. Myrrha*, see *Myrrha*.

BALSAMARIA INOPHYLLUM, see Fagara octandra.

BALSAMELÉON, Myroxylon Peruiferum.

BALSAMIC, Balsamicus, (F.) Balsamique, from βαλσαμον, 'balsam.' Possessing the quali-

ties of balsams. *Balsamic odour*:—a sweet, faint, and slightly nauseous smell. *Balsamic substance*:—one resembling the balsams in property.

BALSAMIER ÉLÉMIÈRE, Amyris elemifera—*b. de la Mecque, Amyris opobalsamum*.

BALSAMINA, Momordica balsamina.

BALSAMINE, Momordica balsamina.

BALSAMIQUE, Balsamic.

BALSAMITA FEMINEA, Achillea ageratum—*b. Major, Tanacetum balsamita*—*b. Mas, Tanacetum balsamita*.

BALSAMITA SUAVEOLENS, B. odorata seu maris, Mentha Saracenicæ seu Romana. Ord. Compositæ. Ser. Syst. Syngenesia Polygamia superflua. A plant, common in the south of France, and cultivated in the gardens; where it bears the names of *Menthe coq ou romaine, Grand baume, Baume coq ou Costus des Jardins*. Its smell is strong and aromatic, and taste hot. It is used for the same purposes as tansey, i. e. as a stimulant, vermifuge, &c.

BALSAMITA SUAVEOLENS, Tanacetum balsamita—*b. Vulgaris, Tanacetum balsamita*.

BALSAMO-SACCHARUM, Elæo-Saccharum.

BALSAMUM, see Balsam, Amyris opobalsamum—*b. Ægyptiacum, see Amyris opobalsamum*—*b. Album, see Myroxylon Peruiferum*—*b. Alpini, Dracocephalum Canariense*—*b. Alpini, see Amyris opobalsamum*—*b. Anodynum, Linimentum saponis et opii*—*b. Apoplecticum, Balsam of life of Hoffmann*—*b. Aromaticum, Balsam of life of Hoffmann*—*b. Asiaticum, see Amyris opobalsamum*—*b. Braziliense, Copaiba*—*b. Calaba, see Fagara octandra*—*b. Canadense, see Pinus balsamea*—*b. Catholicum, Tinctura benzoini composita*—*b. Cephalicum, Balsam of life of Hoffmann*—*b. Copaibæ, Copaiba*—*b. Genuinum antiquorum, see Amyris opobalsamum*—*b. Hyperici simplex, see Hypericum perforatum*—*b. Ioduretum, Balsam, ioduretted*—*b. Judaicum, see Amyris opobalsamum*—*b. Libani, see Pinus cembra*—*b. Maria, see Fagara octandra*—*b. c. Meccæ, see Amyris opobalsamum*—*b. Mercuriale, Unguentum hydrargyri nitratis*—*b. Nervinum, Balsam of life of Hoffmann*—*b. Nucistæ, see Myristica*—*b. Opodeldæ, Linimentum saponis camphoratum*—*b. Ophthalmicum rubrum, Unguentum hydrargyri nitrico-oxydi*—*b. Persicum, Tinctura benzoini composita*—*b. Peruvianum, see Myroxylon Peruiferum*—*b. Saturninum, Unguentum plumbi superacetatis*—*b. Scherzeri, Balsam of life of Hoffmann*—*b. Stomachicum, Balsam of life of Hoffmann*—*b. Styracis, Styrax*—*b. Styracis benzoini, Benjamin*—*b. Succini, see Succinum*—*b. Sulphuris Barbadense, Petroleum sulphuratum*—*b. Sulphuris simplex, Oleum sulphuratum*—*b. Syriacum, see Amyris opobalsamum*—*b. Tolutanum, see Toluifera balsamum*—*b. Tranquillans seu Tranquillum, Balsam, tranquil*—*b. Traumaticum, Tinctura benzoini composita*—*b. Universale, Unguentum plumbi superacetatis*—*b. Viride, Balsam, green, of Metz; see Fagara octandra*.

BALSAMUS, Balsam—*b. Palustris, Mentha aquatica*.

BALSEM, Amyris opobalsamum.

BAMBA, Bamboo.

BAMBALIA, Balbuties.

BAMBALIO, Bam'balo, Bala'tro, from βαλῶ, 'I speak inarticulately.' One who stammers or lisps, or utters inarticulate sounds. According to KRAUSE, one who speaks as if he had pap in his mouth, or as if his tongue were paralyzed.

BAMBOO, (F.) Bambou, Bambu. Fam. Graminæ. Ser. Syst. Hexandria Monogynia. The young shoots of *Bambos arundinacea, Arundo bambos, Bambusa arundinacea*, and of *Bambos verticillata*, contain a saccharine pith, of which

the people of both the Indies are very fond. They are sometimes made into a pickle.

BAMBOS ARUNDINACEA, Bamboo—b. Verticillata, Bamboo.

BAMBUSA ARUNDINACEA, Bamboo.

BAMIX MOSCHATA, Hibiscus abelmoschus.

BAMMA, from βαμν, 'I plunge,' 'a paint: a dye.' Anciently, liquids were so called, in which certain bodies were plunged, to moisten or soften them. In the case of tea, for instance, into which bread is dipped, the tea would be the *bamma*.

BANANA, Musa sapientum.

BANANIER, Musa sapientum.

BANAUSIA, Charlatanry.

BANC D'HIPPOCRATE, Bathron.

BANCAL, (F.) One who has deformed legs. It includes the *valgus*, *compennis*, and *varus*, which see.

BANCROCHE, (F.) A vulgar epithet for a rickety individual.

BAND, PRIMITIVE, see Nerve Fibre.

BAND'AGE, *Desma*, *Syndes'mus*, *Hypodes'mia*, *Hypodes'ma*, *Hypodes'mus*, (the last three signify properly an under bandage.) A *binder*, from Sax. *bindan*, 'to bind.' This word, with the French, is generally used to express the methodical application of rollers, compresses, &c., *Band'aging*, *Syn'desis*, to fix an apparatus upon any part,—corresponding to the words *deligatio*, *fasciatio*, *fasciolum applicatio*, *epid'esia*. With us the noun is usually applied to the result of the application, or to the bandage itself;—a sense in which the French employ the word *Bande*. Bandages are *simple* or *compound*. The simple bandage is *equal*, if the turns are applied circularly above each other; *unequal*, if the turns are not accurately applied upon each other. If each turn of the bandage be only covered one-third, it forms the *doloire* of the French; if the edges touch only slightly, it is the *moussé*; if the turns are very oblique and separated, it is the *spiral* or *creeping*, (F.) *rampant*; if folded upon each other, it is termed the *reversed*, (F.) *renversé*. By uniting various kinds of bandaging, we have the *compound*; and these compound bandages have received various names expressive of their figure, or of the parts to which they are applied, as *capistrum*, *epica*, &c. Bandages are divided, also, as regards their uses, into *uniting*, *dividing*, *retaining*, *expelling*, *compressing*, &c.

BANDAGE OR ROLLER, *Fas'cia*, *Tæ'nia*, *Epides'mos*, *Vin'culum*, the *Bande* of the French. It may be made of linen, flannel, or other stuff capable of offering a certain resistance. The two extremities of a bandage are called *tails*, (F.) *chefs*, and the rolled part is termed its *head*, (F.) *globe*. If rolled at both extremities, it is called a *double-headed roller* or *bandage*, (F.) *Bande à deux globes*.

BANDAGE À BANDELETTES SÉPARÉES, Bandage of separate strips.

BANDAGE, BODY, *Manti'le*, (F.) *Bandage de Corps*, is used for fixing dressings, &c., to the trunk. It is formed of a towel, napkin, or some large compress, folded three or four times; the extremities of which are fastened by pins. This is again fixed by means of the *scapulary bandage*, which is nothing more than an ordinary bandage, stitched to the anterior and middle part of the napkin, passing over the clavicles and behind the head, to be attached to the back part of the napkin.

BANDAGE, COMPOUND, see Bandages.

BANDAGE, COMPRESSING, or ROLLER, *Fascia compressiva* seu *convoluta*, (F.) *Bandage compressive* ou *roulé*, is the simple roller with one head; and is employed in cases of ulcers, varices,

&c. of the limbs. Whenever this roller is applied to the lower part of the limbs, it is carried upwards by the *doloire* and *reversed* methods above described.

BANDAGE, COMPRESSIVE, see Bandage, compressing—b. *de Corps*, see Bandage—b. *Dividing*, see Bandage—b. *Divisif*, *Dividing bandage*—b. *à dix-huit Chefs*, Bandage, eighteen-tailed—b. *en Doloire*, *Doloire*—b. *Double-headed*, see Bandage.

BANDAGE, EIGHTEEN-TAILED, *Fas'cia octoid'ecim capit'ibus*, (F.) *Bandage à dix-huit chefs*. This bandage is made of a longitudinal portion of a common roller; and with a sufficient number of transverse pieces or tails, to cover as much of the part as is requisite. It is a very useful bandage, inasmuch as it can be undone without disturbing the part.

BANDAGE, EXPELLING, see Bandage.

BANDAGE, GALEN'S, *B. for the Poor*, *Fas'cia Gale'ni* seu *Pau'perum*, (F.) *Bandage de Galien ou des Pauvres*, *Galen*, is a kind of *cuneus* or hood, (F.) *Couvetche*, divided into three parts on each side; of which GALEN has given a description. See Cancer Galeni.

BANDAGE OF GENGIA, Bandage of Theden—b. *Hernial*, see *Truss*—b. *Immovable*, *Apparatus*, *immovable*.

BANDAGE, IN'GUINAL, *Fas'cia inguinal'is*. A bandage for keeping dressings applied to the groin. It consists of a cincture, to which is attached a triangular compress, adapted for covering the groin. To the lower extremity of this, one or two bandages are attached, which pass under the thigh, and are fixed to the posterior part of the cincture. This bandage may be either simple or double.

Other bandages will be found described under their various names.

BANDAGE DES PAUVRES, Bandage, Galen's—b. *Permanent*, *Apparatus*, *immovable*—b. of the Poor, see Bandage, Galen's, and Cancer Galeni—b. *Rampant*, see Bandage—b. *Reversed*, see Bandage—b. *Roulé*, see Bandage—b. of *Scul'tetus*, Bandage of separate strips.

BANDAGE OF SEPARATE STRIPS, or B. OF SCUL'TETUS, *Fas'cia fasci'olis separati'im disposit'is* seu *Scul'teti*, (F.) *Bandage à bandelettes séparées* ou *de Scul'tet*. This is formed of linen strips, each capable of surrounding once and a half the part to which they have to be applied, and placed upon each other, so as to cover successively one-third of their width. It is used chiefly for fractures, requiring frequent dressing.

BANDAGE, SIMPLE, see Bandage.

BANDAGE OF THEDEN, *B. of Gengia*. A bandage employed, at times, in brachial aneurism, which commences at the fingers, and extends to the axilla.

BANDAGE, UNDER, *Hypodes'mis*—b. *Unequal*, see Bandage—b. *Unissant*, *Uniting bandage*—b. *Uniting*, see Bandage, and *Uniting bandage*.

BANDAGING, see Bandage—b. *Doctrine*, of, *Desmaturgia*.

BANDAGIST, (F.) *Bandagiste*. One whose business it is to make bandages, and especially those for hernia.

BANDE, Bandage. The word *Bande*, in anatomy, is used by the French for various narrow, flat, and elongated expansions.

BANDE À DEUX GLOBES, see Bandage.

BANDE D'HÉLIODORE, a kind of bandage for supporting the mamme.

BANDEAU, (F.) A kind of simple bandage, which consists of a piece of cloth, folded four times, and applied round the head. There is also the *Bandeau ou Mouchoir en triangle* or

triangular bandage, a kind of *couvrechef*, made of a square piece of cloth, or of a handkerchief, folded diagonally, and applied round the head.

BANDELETTE, (F.) Diminutive of *Bande Fasciola*, *Tanola*, *Vitta*: a narrow bandage, strip, or fillet. Also *Tenia semicircularis*.

BANDETTES AGGLUTINATIVES. Small strips, covered with a glutinous plaster. *Vitta agglutinantes*. See *Agglutinant*.

BANDETTES DÉCOUPÉS, are strips of linen, notched on one edge, and covered, on one side, with ointment. They are applied to wounds to prevent the lint from sticking, and the laceration of the cicatrix.

BANDELETTE SEMICIRCULAIRE, *Tænia semicircularis*—*b. des Cornes d'amon*, *Corpus fibriatum*—*b. des Eminences pyriformes*, *Tænia semicircularis*—*b. de l'Hippocampe*, *Corpora fibriata*.

BANDURA, *Nepentha destillatoria*.

BANDY-LEGGED, *Cnemosecoliosis*.

BANEBERRY, *Actæa spicata*.

BANGUE, Bhang, Bang, Bangi or Beng, Sed-hee, Subjee. Adanson believes this to be the *Nepenthes* of the ancients. The largest leaves and capsules without the stalks of *Can'nabis Indica*, (F.) *Chanvre Indien*, Indian hemp, probably identical with *C. sativa*. Family, *Urticæ*. See *Syst. Dicoica Pentandria*. The leaves and flowers of *Cannabis* are narcotic and astringent. They are chewed and smoked. The seeds, mixed with opium, areca, and sugar, produce a kind of intoxication, and are used for this purpose by the people of India. An alcoholic extract of the plant, *Churrus—Extractum Can'nabie* (Ph. U. S.)—has been used in India, and since then in Europe and in this country, as a narcotic and anti-convulsive, in the dose of from half a grain to ten or more. It requires, however, great caution in its administration. The pure resin—*Cannabine, Haschischine*—is active in the dose of two-thirds of a grain.

The dried plant, which has flowered, and from which the resin has not been removed, called *Ganjah, Ganjah, Guaza, and Guazab, Haschisch, Haschich, Hachisch, Hashish, or Haschisch*, of the Arabs, consists of the tops and tender parts only of the plant, collected immediately after inflorescence, and simply dried.

BANICA, *Pastinaca sativa*.

BANILAS, *Vanilla*.

BANILLA, *Vanilla*.

BANILLOES, *Vanilla*.

BANISTERIA ANGULO'SA. This plant, in Brazil and the Antilles, passes for a powerful sudorific, and an antidote to the poison of serpents.

BANKSIA ABYSSINICA, *Hagenia Abyssinica*—*b. Speciosa, Costus*.

BANNIÈRES, MINERAL WATERS OF. Bannières is a village in Quercy, diocese of Cahors, France. The waters are probably chalybeate. They are celebrated in amenorrhœa, cachexia, jaundice, &c.

BA'OBAB, Adansonia digitata, of Africa. *Nat. Ord. Bombacæ*. One of the largest productions of the vegetable kingdom. Its fruit is called, in the country, *Pain de singe*. The pulp is sourish, and agreeable to eat; and a refreshing drink is made from it, which is used in fevers. Prospero Alpini and Dr. L. Frank think that the *Terra Lemnia* was prepared, in Egypt, from the pulp. All the parts of the Baobab abound in mucilage. The bark has been given as a substitute for cinchona.

BAPTISIA LEUCANTHA, see *Sophora tinctoria*—*b. Tinctoria, Sophora tinctoria*.

BAPTISTERIUM, Baignoire.

BAPTORRHŒA, see *Gonorrhœa*.

BARAQUETTE (F.) A name given by Raisons, physician at Nîmes, in France, to a catarrhal epidemic, which occurred there in 1761. See *Influenza*.

BARATHRON, *Juniperus sabina*.

BARATHRUM, *Antrum*.

BARBA, Beard—*b. Aaronis, Arum maculatum*—*b. Capræ, Spiræa ulmaria*—*b. Hirci, Tragopogon*—*b. Jovis, Sempervivum tectorum*.

BARBADOS, see West Indies—*b. Leg*, see *Elephantiasis*.

BARBAREA, Erysimum barbarca—*b. Stricta, Erysimum barbarca*.

BARBAROS'SE PIL'ULÆ, Barbaros'se's Pills. An ancient composition of quicksilver, rhubarb, diagridium, musk, &c. It was the first internal mercurial medicine which obtained any real credit.

BARBE, Beard—*b. de Bouc, Tragopogon*.

BARBEAU, Centaurea cyanus, Cyanus segetum.

BARBER-CHIRURGEONS. A Corporation of London, instituted by King Edward IV. The barbers were separated from the surgeons, by 18 Geo. II., c. 15; and the latter were erected into a *Royal College of Surgeons* at the commencement of the present century.

BARBERS, ARMY, see *Bathers*.

BARBERIE, MINERAL WATERS OF. These mineral waters are half a league from Nantes. They contain carbonic acid, chlorides of magnesium and sodium, sulphate of magnesia, carbonates of magnesia, lime, and iron. They are used as chalybeates.

BARBERRY, Oxycantha Galeni—*b. American, see Oxycantha Galeni*.

BARBIERS. A variety of paralysis chiefly prevalent in India; and by many considered to be the same as *Beriberi*. *Beriberi* is commonly an acute disease. *Barbiers* is generally chronic.

BARBITIUM, Beard.

BAR-BONE, Pubis, os.

BARBOTINE, Artemisia Santonica.

BARBULA CAPRINA, Spiræa ulmaria.

BARCLAY'S ANTIBILIOUS PILLS, Pilulæ antibiliosæ.

BARDADIA, Pound.

BARDANA, Arctium lappa—*b. Minor, Xanthium*.

BARDANE PETITE, Xanthium.

BARÈGES, MINERAL WATERS OF. Barèges is a village in the department of Hautes Pyrénées, near which are several springs. They are sulphureous and thermal, the heat varying from 85° to 112° Fahrenheit. They contain chlorides of magnesium and sodium, sulphates of magnesia and lime, carbonate of lime, sulphur, &c. These springs have long enjoyed a high reputation, and are daily advised in cutaneous and serofulous affections, &c. A nitrogenized matter was first found in these waters, and afterwards in other sulphureous springs, to which Longchamp gave the name *Barégine*.

FACTITIOUS BARÈGES WATER, Aqua Fæciginensis seu Baretginensis, (F.) Eau de Barèges, is made by adding hydrosulphuretted water, $\frac{1}{5}$ iv, to pure water, $\frac{1}{5}$ xvijss, carbonate of soda, gr. xvj, chloride of sodium, gr. ss. Bottle closely.

BARÉGINE, see *Barèges*.

BARGADA, Convolvulus pes capræ.

BARGOU. An alimentary preparation formed of ground oats, boiled to a proper consistence with water.

BARIGLIA, Soda.

BARI CHLORIDUM, Baryta, muriate of—*b. Iodidum, Baryta, hydriodate of*.

BARILLA, Soda.—b. Alicant, Soda.—b. Carthagena, Soda.—b. Turkey, Soda.

BARILLOR, Soda.

BARIUM, *Baryum*, *Barytium*, *Pluto'nium*, from *βαρυς*, 'heavy.' The metallic base of baryta, so called from the great density of its compounds.

BARIUM, CHLORIDE OF, Baryta, muriate of—b. *Chlorure de*, Baryta, muriate of—b. Iodatum, and Iodide of, Baryta, hydriodate of—b. Protoxide of, Baryta.

BAR, Cinchona.—b. Arica, see Cinchona cordifolia cortex.—b. Ash, see Cinchona.—b. Bitter, Pinckneya pubens.—b. Bogota, see Cinchona.—b. Calisaya, Cinchona cordifolia cortex.—b. Calisaya, spurious, see Cinchona cordifolia cortex.—b. Carabaya, see Cinchona cordifolia cortex.—b. Caribæan, Cinchona Caribæa cortex.—b. Carthagena, see Cinchona.—b. Coquette, see Cinchona.—b. Crown, Cinchona lanceifolia cortex.—b. Cusco, see Cinchona cordifolia cortex.—b. Doon, Sassy-bark.—b. Elk, Magnolia glauca.—b. Essential salt of, see Cinchona.—b. Florida, Pinckneya pubens.—b. Fusagasuga, see Cinchona.—b. Georgia, Pinckneya pubens.—b. Gray, see Cinchona.—b. Huamiles, see Cinchona.—b. Huancu, see Cinchona.—b. Indian, Magnolia glauca.—b. Iron, see Kino.—b. Jésuit's, Cinchona.—b. Jaen, see Cinchona.—b. Lima, see Cinchona.—b. Loxa, Cinchona lanceifolia cortex.—b. Pale, Cinchona lanceifolia cortex.—b. Maracaybo, see Cinchona.—b. Ordeal, Sassy-bark.—b. Peruvian, Cinchona.—b. Pitaya, Cinchona Caribæa cortex, see Cinchona.—b. Red, Cinchona oblongifolia cortex.—b. Royal, Cinchona cordifolia cortex.—b. of St. Ann, see Cinchona cordifolia cortex.—b. Saint Lucia, Cinchona Caribæa cortex.—b. Santa Martha, see Cinchona.—b. Sassy, Sassy-bark.—b. Seven, Hydrangea arborescens.—b. Silver, see Cinchona.—b. Yellow, Cinchona cordifolia cortex.

BARLERIA BUXIFOLIA, Cara schulli.

BARLEY-BREE, Cerevisia.

BARLEY-CORN, Cerevisia.

BARLEY, PEARL, see Hordeum.—b. Scotch, Hordeum

BARLEY-WATER, Decoctum hordei.

BARM, Yest.

BARN, Infans.

BARNET, MINERAL WATERS OF, Barnet is not far from London. The water is of a purging quality, like that of Epsom, and about half the strength.

BAROMACROMETER, *Pædobaromacrometer*, *Pædometer*, from *βαρος*, 'weight,' *μετρος*, 'long,' and *μετρον*, 'measure.' An instrument invented by Stein to indicate the length and weight of a new-born infant.

BAROMETER, *Baroscopium*, *Ba'roscope*, from *βαρος*, 'weight,' and *μετρον*, 'measure.' (F.) *Baromètre*. An instrument which measures the weight of the air. A certain degree of density in this medium is necessary for health. When we ascend high mountains, great inconvenience is experienced, owing to the diminished density. Changes of this character are indicated by the Barometer or weather-glass.

BA'ROS, *βαρος*, 'heaviness.' Employed by the Greek physicians to designate the feeling of lassitude and heaviness observable in many diseases.—Hippocrates, Galen.

BAROSCOPE, Barometer.

BAROSMA CRENATA, Diosma crenata.

BAROTE, Baryta.

BAROTES SALITUS, Baryta, muriate of.

BARRAS, see Pinus sylvestris.

BARRE, MINERAL WATERS OF, Barre is a small town, six leagues from Strasburg. The waters are thermal, and contain much iron, calcareous salt, &c. They are diuretic and tonic.

BARRE (F.), *Barrure*, *Tara*, 'a bar.' A projection or prolongation of the symphysis pubis, —a deformity rendering delivery difficult.

BARREÉ (F.). A term applied, in France, to a female whose pelvis has the deformity described under *Barre*.

BARRÉS (DENTS), *Barred Teeth*. The molar teeth, when the roots are spread or tortuous, so that they cannot be extracted without being broken; or without a portion of the alveolar arch being removed.

BARREL OF THE EAR, Tympanum.

BARREN, Sterile.

BARRENESS, Sterilitas.

BARROS, Terra Portugallica.

BARRURE, *Barre*.

BARTON'S FRACTURE, see Fracture of the Radius, Barton's.

BARYCOCCALON, *Datura stramonium*.

BARYCOITA, *Baryeccia*.

BARYCOI'A, *Barycoi'ta*, *Bradyecoi'a*, *Paracoi'sa obtusa*, *Discoi'a*, *Dysceoi'a*, *Audi'tus diffic'ilis*, *A. gravis*, *A. immin'utus*, *Obaudi'tus*, *Obaudi'tus*, *Hypocoph'o'is*, *Hypocoph'o'is*, (F.) *Durété d'Oreille*, *Dyscote*, from *βαρυς*, 'heavy,' and *ακον*, 'hearing.' Hardness of hearing, incomplete deafness. See *Cophosis*, and *Deafness*.

BARYGLOSSIA, Balbuties, Baryphonia.

BARYI HYDRAS IODATI, Baryta, hydriodate of.

BARYLALIA, Baryphonia.

BARYOD'YNE, from *βαρος*, 'heavy,' and *δυνω*, 'pain.' A dull, heavy pain.

BARYPHO'NIA, *Baryglos'sia*, *Baryla'lia*, *Loque'ta impedi'ta*, from *βαρυς*, 'heavy,' and *φωνη*, 'voice.' Difficulty of voice or speech.

BARYPIRON, *Artemisia abrotanum*.

BARYSOMATIA, *Polysarcia adiposa*.

BARYSOMATICA, *Polysarcia adiposa*.

BAR'Y'TA, from *βαρος*, 'heavy,' *βαρυνω*, 'weight.' *Terra ponderosa*, *Bary'tes*, *Ponderide of Bar'ium*, *Heavy Earth*, *Ponderous Earth*, (F.) *Baryte*, *Barote*, *Terre pesante*. This earth and its soluble salts are all highly corrosive poisons. It is never employed in medicine in the pure state. When externally applied, it is caustic, like potassa and soda.

BAR'YTA, CARBONATE OF, *Baryta Carbonas*, (Ph. U.S.), (F.) *Carbonate de Baryte*, *Craie barotique*, is only used officially to obtain the muriate.

BARYTA, HYDRIODATE OF, *Baryta Hydriodas*, *Baryta Hydriod'ica*, *Hydras Baryi Iodati*—(in the dry state, *Iodide of Bar'ium*, *Baryi Iod'idum*, *Bar'ium Iod'atum*.) has been given in serofulous and similar morbid conditions. It may be administered internally in the dose of one-eighth of a grain three or four times a day, and be applied externally to serofulous swellings, in the form of ointment, (gr. iv to ʒj of lard.)

BARYTA HYDRIODICA, Baryta, hydriodate of.

BARYTA, MURIATE OF, HYDROCHLORATE OF, *Bary'ta mur'ias*, *Chlo'ride of Bar'ium*, *Bar'ii Chlo'ridum* (Ph. U.S.), *Chlo'ruret of Bar'ium*, *Terra ponderosa sal'ta seu mur'iat'a*, *Sal mur'iat'icum barot'icum*, *Baro'tes sal'itus*, (F.) *Chlo'rure de barium*, is the combination chiefly used. The Muriate of Baryta may be formed as follows: *Baryt. Carbon.* in frustulis, lbj; *Acid. Muriat.* fʒxij; *Aqua* Oijj. Mix the acid with the water, and gradually add the Carbonate of Baryta. Toward the close of the effervescence, apply a gentle heat, and, when the action has ceased, filter the liquor, and boil it down so that crystals may form as it cools.—Ph. U. S.

It is given in the form of the *Solutio Muriat'is Bary'tæ*, *Liquor Bar'ii Chlo'ridi*, Ph. U. S. *Aqua bary'ta mur'iat'a*, (F.) *Solution de Muriate de*

Baryte, (*Muriate of Baryta*, one part; *distilled water*, three parts.) and is employed in scrofulous cases, worms, and cutaneous diseases. Externally, to fungous ulcers and to specks on the cornea.

BARYTA, SULPHATE OF, *Baryta sulphas*, (F.) *Sulfate de Baryte*, is used in pharmacy, like the carbonate, to obtain the chloride of barium.

BARYTES (CARBONAS, Baryta, carbonate of—b. Hydriodas, *Baryta, hydriodate of—b.* Murias, *Baryta, muriate of—b.* sulphas, *Baryta, sulphate of.*

BARYTE, *Baryta—b.* Carbonate de, *Baryta, carbonate of—b.* sulfate de, *Baryta, sulphate of.*

BARYTHMIA, Melancholy.

BARYTIUM, Barium.

BARYUM, Barium.

BAS-FOND, see Urinary Bladder.

BAS-LASSÉ, Stocking, laced.

BAS-VENTRE, Abdomen.

BASAAL. The name of an Indian tree, the decoction of whose leaves, in water, with ginger, is used as a gargle in diseases of the fauces. The kernels of the fruit are vermifuge.

BASANASTRA'GALA, from *basavos*, 'torture,' and *αστραγαλος*, 'the astragalus.' Pain in the ankle joint; gout in the foot.

BASANIS'MOS, from *basanizein*, 'to explore.' 'A touch-stone.' Investigation or examination.—Hippocrates, Galen.

BASE, *Basis*, from *βαινω*, 'I proceed,' 'I rest,' 'I support myself.' That which serves as a foundation or support. That which enters, as a principal matter, into a mixture or combination. In anatomy, it is employed in the former sense, as *Base of the Cranium*, *Base of the Brain—Basis seu Pons* *Anterior cerebri*; *Base of a process*, &c., *Base of the heart—Basis vel cornua cordis*. In dentistry, it means a metallic, ivory, or hippopotamus plate, which is used as a support for artificial teeth. In the art of prescribing, *Basis* is the chief substance which enters into a compound formula.

BASE-BORN, Illegitimate.

BASEMENT MEMBRANE, see Membrane, basement.

BASIATTO, Coition.

BASIATOR, Orbicularis oris.

BASIL BUSH, *Ocimum caryophyllatum—b.* Citron, *Ocimum basilicum—b.* Common, *Ocimum basilicum—b.* Small, *Ocimum caryophyllatum—b.* Wild, *Chenopodium vulgare—b.* Wild, *Cunila mariana—b.* Wild, *Pycnanthemum incanum*.

BASILAD, see Basilar Aspect.

BASILAR, *Basila'ris, Bas'ilary*, (F.) *Basilaire*. That which belongs to the base, from *basis*, 'base.' This name has been given to several parts, which seem to serve as basis to others. The sacrum and sphenoid have been so called.

BASILAR ARTERY, *A. basila'ris seu cervica'lis*, (F.) *Artère ou Tronc basilairre, A. mésocephalique* (Ch.) The union of the two vertebral arteries. It ascends along the middle groove on the inferior surface of the tuber, and is supported, beneath, by the *Fossa basilaris*. It terminates in the posterior cerebral arteries.

BASILAR ASPECT. An aspect towards the base of the head.—Barclay. *Basilad* is used adverbially by the same writer to signify 'towards the basilar aspect.'

BASILAR FOSSA, (F.) *Gouttière ou Fosse basilairre*, is the upper surface of the basilar process, —so called because it is channelled like a *Fossa* or *Gutter*. The *Tuber annulare* rests upon it.

BASILAR PROCESS, *Processus basila'ris ossis occip'itis seu cuneiformis ossis occip'itis*, (F.) *Apophyse basilairre, Prolongement sous-occipital, Cuneiform Process*, is the bony projection, formed

by the inferior angle of the os occipitis, which is articulated with the sphenoid.

BASILAR SINUS, *Sinus transversus*.

BASILAR SURFACE, (F.) *Surface basilairre*, is the inferior surface of the process. It is covered by the mucous membrane of the pharynx.

BASILAR VERTEBRA. The last vertebra of the loins.

BASIL'IC, Basil'icis, (F.) *Basilique*, from *βασιλικος*, 'royal.' This name was given, by the ancients, to parts which they conceived to play an important part in the animal economy.

BASILIC VEIN, *Vena basil'ica seu cubiti inte'rior*, (F.) *Veine Basilique, Veine cubitale cutanée of Chaussier*. This vein is one of those on which the operation of bloodletting is performed. It is situate at the internal part of the fold of the elbow, in front of the humeral artery, and is formed by the anterior and posterior cubital veins, and by the median basilic. It terminates, in the arm-pit, in the axillary vein. The ancients thought, that the basilic of the right arm had some connexion with the liver, and hence they called it *hepatic*. The vein of the left arm, for a similar reason, they called *splenic*. The *Median Basilic Vein*, (F.) *Veine médiane basilique*, is one of the branches of the preceding vein. It joins the median cephalic at an acute angle, or rather by a transverse branch, and receives some branches of the deep radial and cubital veins, and a considerable subcutaneous vein—the common median.

BASILIC COMMUN, *Ocimum basilicum—b.* *Sauvage*, *grand*. *Chenopodium vulgare*.

BASIL'ICUM, Basil'icum. 'Royal,' or of great virtue. An ointment, composed of yellow wax, black pitch, and resin, of each one part, olive oil, four parts. Hence it was called *Unguentum Tetrapharmacum*, (*τετραφαρμακα*, 'four drugs'.)—Celsus. Scribonius Largus.

BASILICON, Basilicum, of the Parisian Codex, is the *Onguent de Poix et de Cire*. In most Pharmacopœias, it is represented by the *Unguentum* or *Ceratum Resinæ*. It is used as a stimulating ointment. See *Ceratum Resinæ*; and *Unguentum Resinæ Nigræ*.

BASILICON, Basilicon, Ocimum Basilicum—b. *Citratum, Ocimum basilicum—b.* *Majus, Ocimum basilicum*.

BASILIQUE, *Basilie*.

BASILISCUS, Syphilis.

BASIO-CERATO-CHONDRO-GLOSSUS, Hyoglossus.

BASIO-CERATO-GLOSSUS, from *basis*, 'base,' *κερας*, 'cornu,' and *γλωσσα*, 'tongue.' A name given to a part of the hyoglossus, which is inserted into the cornu of the os hyoides and base of the tongue.

BASIOCES'TRUM, from *basis*, 'the base,' and *κεστρα*, 'a dart.' An instrument for opening the head of the foetus in utero, invented by Mesler, a German.

BASIO-GLOS'SUS, Hypoglossus, *Hypobasiloglossus, Ypsiloglossus*, from *basis*, 'base,' and *γλωσσα*, 'the tongue.' A name formerly given to the portion of the hyoglossus which is inserted into the base of the os hyoides.—Riolan, Thomas Bartholine. See *Lingual Musc*.

BASIO PHARYNGÆ'US, from *basis*, 'base,' and *φαρυγξ*, 'the pharynx.' A name given to some fibres of the constrictor pharyngis medius.—Winslow.

BASIS, see Prescription—*b.* *Cerebri*, *Base of the Brain—b.* *Cordis*, *Radix cordis—b.* *Corporis*, *Sole*.

BASSI-COL'ICA. Name of a medicine composed of aromatics and honey.—Scribonius Largus.

BASSIA BUTYRACEA, see Spirit, (Arrack.)

BASSIN, Pelvis—*b. Oculaire*, *Scaphium oculare*.

BASSINER, *Fove're*, to foment.

BASSINET, Pelvis of the kidney, *Ranunculus bulbosus*.

BASSORA, GUM. A gum, obtained from a plant unknown, which came originally from the neighbourhood of Bassora, on the Gulf of Persia, whence its name. It is in irregularly shaped pieces, white or yellow, and intermediate in its transparency between gum Arabic and gum tragacanth. Only a small portion is soluble in water. The insoluble portion is a peculiar principle, called *Bassorin*. It is not used in medicine; but *Bassorin* enters into the composition of several substances.

BASSORIN, see *Bassora* gum.

BASSWOOD, *Tilia Americana*.

BASTARD, Illegitimate.

BATA, *Musa Paradisiaca*.

BATATA DE PURGA. The Brazilian and commercial name of the purgative, feculent, and gum-resinous roots of two plants of the family *Convolvulaceæ*, the one called *Joticoua*, *Meehounca*, *Convolvulus meehoucanus*, the *Piptostegia Pissalis* of Von Martius; the other *Convolvulus operculatus*, the *Ipomoea* seu *Piptostegia operculata* of Von Martius.

BATATAS. The inhabitants of Peru gave this appellation to several tuberous roots, especially to *Convolvulus Batatas* or *Sweet Potato*. Our word, *Potato*, comes from this. See *Solanum tuberosum*.

BATEMAN'S PECTORAL DROPS, see *Pectoral Drops*, *Bateman's*.

BATERION, *Bathron*.

BATES'S ALUM WATER, *Liquor aluminis compositus*—*b. Anodyne Balsam*, *Linimentum saponis et opii*—*b. Camphorated Water*, *Lotion*, camphorated, of *Bates*—*b. Collyrium*, see *Lotion*, camphorated, of *Bates*.

BATH, Anglo-Saxon *bað*, *Bal'neum*, *Balan'eum*, *Baline'um*, *Loutron*, (Old Eng.) *Baine*, (F.) *Bain*. Immersion, or stay, for a longer or shorter duration, of the whole or a part of the body, in some medium, as water. Act of plunging into a liquid, sand, or other substance, in which it is the custom to bathe, *Plunge Bath*. Also, the vessel in which the water is put for bathing. Also, a public or private establishment for bathing. The common water-bath, used for hygienic as well as for therapeutical purposes, is the *Bain simple* ou *hygiénique* of the French.

In *Pharmacy*, a vessel, placed over a fire, and filled with any substance, into which another vessel is placed, containing matters for digestion, evaporation, or distillation.

BATH ACID, *Bal'neum ac'idum* (*Acid. muriat.* *℞ij*; *Aqua* cong. *℥xvi*. One half, one third, or one fourth the quantity of acid is more frequently employed.)

BATH ACID, *Scott's*, see *Scott's Acid Bath*—*b. Air*, hot, see *Bath*, hot—*b. Air*, warm, see *Bath*, hot.

BATH ALKALINE, *Bal'neum alkali'num*. This may be made of half a pound or a pound of *pearl-ash* or of *carbonate of soda*, to sixty-six gallons of water.

BATH ANIMAL, *Bal'neum animal'e*, consists in wrapping an animal recently killed, or its skin, around the body, or some part of it.

BATH ANTIPSORIC, *Bal'neum antipso'ricum*. Recommended in cases of itch and other cutaneous diseases. (*Potass. sulphuret.* *℥iv*, *Aqua* cong. *℥x*.)

BATH ANTISYPHILITIC, *Bal'neum antisyp'hilit'icum*, *Mercurial bath*. Made by dissolving from two drachms to an ounce of the corrosive chloride of mercury in sixty gallons of water.

BATH ARM, *Brachil'vium*. A bath for the arm.

BATH COLD, see *Bath*, hot—*b. Cool*, see *Bath*, hot.

BATH DRY, is one made of ashes, salt, sand, &c. The ancients used these frequently for the rapaeutical purposes.

BATH EARTH, *Arenatio*.

BATH ELECTRIC, (F.) *Bain électrique*, consists in placing the person upon an insulated stool, communicating by a metallic wire, with the principal conductor of the electrical machine in action. The Electric Bath produces general excitement of all the functions, and especially of the circulation and secretions.

BATH FOOT, *Pedil'vium*, (F.) *Bain de Pied*, a bath for the feet.

BATH GELATINOUS, *Bal'neum gelatino'sum*. Made by dissolving two pounds of *gelatin* or *glue* in a gallon of water. A *gelatino-sulphurous bath* may be made by adding a pound of *glue*, previously dissolved in water, to the sulphur bath.

BATH GENERAL, (F.) *Bain Entier*, is one in which the whole body is plunged, except the head; in contradistinction to the *partial bath*, *Merobalan'eum*, *Merobal'neum*.

BATH HALF, *Semien'pium*, *Eccathisma*, *Inces'sio*, *Inces'sus*, is one adapted for half the body. One, for receiving only the hips or extremities, is also so called.

The *Sitz-bath*, (G.) *Sitzbad*, of the hydropathists is a tub of cold water, in which the patient sits for a variable period.

BATH HAND, *Manub'vium*, (F.) *Bain de Main* ou *Manicule*, is a bath for the hands.

BATH HEAD, *Capitul'vium*, (F.) *Bain de Tête* ou *Capitulu'e*, a bath for the head.

BATH HIP, *Coxal'vium*, (F.) *Bain de Fau-teuil*, *Bain de Siège*, is one in which the lower part of the trunk and upper part of the thighs are immersed.

BATH HOT, *Bal'neum Cal'idum*, *Zestolu'ria*, (F.) *Bain chaud*, is a bath, the temperature of which is 98° and upwards; the *WARM BATH* from 92° to 98°; the *TEMPID BATH*, (F.) *Bain Tiède*, *Bal'neum tep'idum*, from 85° to 92°; the *TEMPERATE BATH*, (F.) *Bain tempéré*, from 75° to 85°; the *COOL BATH*, (F.) *Bain frais*, from 60° to 75°; the *COLD BATH*, *Bal'neum frigidum*, *Frigida'rium*, (F.) *Bain froid*, *Bain très froid*, (of some,) from 30° to 60°; and the *VAPOUR BATH*, *Bal'neum vapo'ris*, (F.) *Bain de Vapeur*, *Étuve Humide*, from 100° to 130°, and upwards. See *Vaporarium*. A *WARM AIR BATH*, or *HOT AIR BATH*, consists of air the temperature of which is raised.

BATH MEDICATED, *Bal'neum Medica'tum*, (F.) *Bain médicinal*, is a bath, formed of decoctions or infusions of vegetable substances, or of any ingredient, introduced into the water for therapeutical purposes.

BATH MERCURIAL, *Bath*, antisyp'hilitic—*b. Nitro-muriatic acid*, *Scott's acid bath*—*b. Plunge*, see *Bath*.

BATH SAND, *Bal'neum Are'næ*, (F.) *Bain de Sable*, consists of a vessel filled with sand, and placed over the fire. Into this vessel, the one is put which contains the substance to be evaporated. See *Psammismus*.

BATH SEA, *Bal'neum mari'num*, (F.) *Bain de mer*. A sea-water bath.

BATH SHOWER, *Implu'vium*, *Hydroco'nium*, is one in which the water is made to fall like a shower on the body. See *Douche*.

BATH SITZ, see *Bath*, half.

BATH STEAM, may be formed by introducing steam into a properly closed vessel in place of water, as in the water bath.

BATH SUCCESSION, *Transit'ion bath*. A term

applied to the rapid succession or transition from a cold to a warm or hot bath, or conversely.—Bell.

BATH, SULPHUR, *Bal'neum Sulphuris*, *Sulphuretted bath*. A bath much used in psora, and other chronic cutaneous affections. It may be composed of two ounces of diluted sulphuric acid, and eight ounces of sulphuret of potassium added to each bath; or simply of the sulphuret of potassium.

BATH, SULPHUR VAPOUR, see Sulphurous Acid.

BATH, TAN. An astringent bath, prepared, at times, by boiling two or three handfuls of ground oak-bark,—such as is used by tanners—in two or three quarts of water, for half an hour, and then adding the decoction to the water of the bath.

BATH, TEMPERATE, see Bath, hot—b. Tepid, see Bath, hot—b. Transition, Bath, succession—b. Vapour, see Bath, hot, and Vaporarium—b. Warm, see Bath, hot.

BATH, WATER, *Balneum Mariæ* seu *Mariæ*, (F.) *Bain Marie*, in chemistry, consists of a vessel filled with boiling water, or salt water, in which the vessel is placed, that contains the substance to be evaporated.

Bathing is much employed in the treatment of disease. The cold bath, especially the cold sea bath, is a sedative and indirect tonic: the warm bath a relaxant; and the hot bath a stimulant.

The regular use of the bath is extremely conducive to health; but if too much indulged in, it is apt to produce injurious effects.

BATH ALUM SPRINGS, see Virginia, Mineral Waters of.

BATH, MINERAL WATERS OF, *Aquæ Bathoniæ* vel *Bad'niæ*, *Aquæ Solis*, *Aquæ Bad'niæ*. Celebrated thermal springs at Bath, in England. They contain but little impregnation, and are chiefly indebted to their temperature, from 112° to 117° Fahrenheit, for their utility. The main ingredients are sulphate of lime, chloride of sodium, sulphate of soda, carbonate of lime, protoxide of iron, free carbonic acid and azote.

These waters are employed in the most heterogeneous cases; and are serviceable where the simple thermal springs are indicated, as in rheumatism, paralysis, &c.

BATH SPRINGS, in Berkley County, Virginia, is a mild carbonated water: temperature 73° Fahr. It is a useful bath in cutaneous affections, and is said to contain some of the salts of lime and magnesia.

BA'THER, same etymon; *Balnea'rius*, *Balinea'tor*, (F.) *Baigneur*. One who bathes. Anciently, the name was given to those that administered baths to the diseased,—the *Ætiastes* of the French. At the present day, in remote districts in Germany, the country people call their medical practitioners *Bäder* or 'bathmen,' and *Feldscheeren* or 'army barbers.'

BATHING, see Bath.

BATHMIS, *Bathmus*, 'base, support.' The cavity of a bone, which receives the eminence of another; and especially the two *Fossettes* at the inferior extremity of the humerus into which the processes of the ulna are received, during the flexion and extension of the fore-arm.

BATHRON, *Bathrum Hippocratis*, *Scamnum Hippocratis*, *Bate'rión*, 'a step, a ladder.' (F.) *Banc d'Hippocrate*. An instrument, used for the extension of a limb, in cases of fracture or luxation. The description of it is found in Galen, Orisbanus, and Scultetus, with a figure.

BATHRUM HIPPOCRATIS, *Bathron*.

BATIA, Retort.

BATISSE, MINERAL WATERS OF. Ba-

tisse is three leagues from Clermont, in France. The water is tepid, and contains subcarbonate and sulphate of soda, sulphates of lime and iron, muriate of magnesia, and carbonate of lime.

BATOS, *Robus Idaeus*.

BATRACHUS, *Ranula*.

BATTALISM'US, *Battaris'mus*, from *Serrapiscus*, 'to stammer.' *Balbuties*. Stammering with incapacity to pronounce the R.

BATT'ALUS, *Bat'tarus*, same etymon. A stammerer, a stutterer.

BATTARISMUS, *Battalismus*.

BATTARUS, *Battalus*.

BATTATA VIRGINIANA, *Solanum tuberosum*.

BATTEMENS DOUBLES, see *Bruit du Cœur fictil*.

BATTEMENT, *Pulsation*.

BAUDRICOURT, MINERAL WATERS OF. Baudricourt is a town of France, two leagues and a half from Mirecourt. The waters are sulphureous.

BAUDRUCHES, *Condom*.

BAUHIN, VALVE OF, *Valve of TUL'PIUS*, *V. of FALL'OPPIUS*, *V. of VAROL'US*, *Hæmocolic Valve*, *Ileo-colic Valve*, *Valvula Ilei*, *Valvula Coli*, *V. Cæci*, *Operculum Ilii*, *Sphincter Ilei*. This name is given to the valve situate transversely at the place where the Ileum opens into the cæcum, and which Bauhin says he discovered at Paris, in 1759. It had, however, been previously described by several anatomists; as by Vidus Vidius, Postius, &c.

BAUME, *Balsam*—*b. d'Acier*, *Balsam*, *chalybeate*—*b. d'Aiguilles*, *Balsam*, *chalybeate*—*b. d'Amérique*, see *Toluifera balsamum*—*b. Apoplectique*, *Balsam*, *apoplectic*—*b. d'Arcæus*, *Arceus*, *balsam of*; see, also, *Balsam of Arcæus*—*b. d'Arcæus*, *Unguentum elemi compositum*—*b. Aromaticum*, *Balsam*, *aromatic*—*b. Benjoin*, *Benjamin*—*b. Blanc*, see *Amyris Opobalsamum*—*b. du Brésil*, *Copaiba*—*b. de Canada*, see *Pinus balsamea*—*b. de Cannelle*, *Laurus cinnamomum*—*b. de Carpathie*, *Balsam of Carpathia*—*b. de Carthagène*, see *Toluifera balsamum*—*b. de Constantinople blanc*, see *Amyris opobalsamum*—*b. de Copahu*, *Copaiba*—*b. Cordiale de Sennerte*, *Balsam*, *cordial*, of *Sennertus*—*b. d'Eau à feuilles ridées*, *Mentha crispæ*—*b. de Feuille*, *Balsam*, *green*, of *Metz*—*b. de Fioraventi spiritueuse*, *Balsam*, *spirituous*, of *Fioraventi*—*b. de Fourcroy ou de Laborde*, *Balsam of Fourcroy or Laborde*—*b. des Funérailles*, *Asphaltum*—*b. de Galaad*, see *Amyris opobalsamum*—*b. de Geneviève*, *Balsam of Geneviève*—*b. Grand*, *Tanacetum balsamita*—*b. du Grand Caire*, see *Amyris opobalsamum*—*b. Hydriodaté*, *Balsam*, *ioduretted*—*b. Hypnotique*, *Balsam*, *Hypnotic*—*b. Hystérique*, *Balsam*, *hysteric*—*b. des Indes*, see *Myroxylon Peruiferum*—*b. Ioduré*, *Balsam*, *ioduretted*—*b. des Jardins*, *Mentha viridis*—*b. de Lucatel*, *Balsam*, *Lucatelli's*—*b. Marie*, see *Fagara octandra*—*b. de Monie*, *Asphaltum*—*b. Nervi*, *Balsam*, *nervous*—*b. de Perou*, see *Myroxylon Peruiferum*—*b. de Saint-Thomas*, see *Toluifera balsamum*—*b. du Samaritain*, *Balsam of the Samaritan*—*b. Saxon*, *Balsam*, *Saxon*—*b. de Soufre*, *Balsam of sulphur*—*b. de Sympathie*, *Balsam of sympathy*—*b. Tranquille*, *Balsam*, *tranquil*—*b. de Tolu*, see *Toluifera balsamum*—*b. de Vanille*, *Vanilla*—*b. Vert*, see *Fagara octandra*—*b. Vert de Metz*, *Balsam*, *green*, of *Metz*—*b. de Vie d'Hoffmann*, *Balsam of Life*, of *Hoffmann*—*b. de Vie de Lelièvre*, *Tinctura aloes composita*—*b. Vrai*, see *Amyris opobalsamum*—*b. Vulnèraire de Minderer*, *Balsam*, *vulnerary*, of *Mindererus*.

BAUMIER, *Balsamier*.

BAURAC, (*Arab.*) *Nitre*, or salt in general. From this word comes *Borax*.

BAURIN, MINERAL WATERS OF. Baurin is a village four leagues from Roye, department of Somme. The waters are strongly chalybeate.

BAVE (F.), *Saliva ex ore fluens, Spuma, Humor Salivæ.* Frothy, thick, viscid saliva, issuing from the mouth. This *drivelling* or *slavering*, we see in children, old people, &c. The term is, also, applied to the frothy liquid, which flows from the mouth of rabid animals. Sauvages uses it synonymously with salivation.

BAVEUX, *Bacense*, an epithet, occasionally applied by the French to the spongy flesh of a wound, which suppurates, and exhibits but little tendency to heal.

BAY, CASTOR, *Magnolia glauca*—b. Rose, *Rhododendron chrysanthemum*—b. Rose, American, *Rhododendron maximum*—b. Sweet, Laurus—b. White, *Magnolia glauca* and *M. macrophylla*.

BDALSIS, Sucking.

BDELLA, Hirudo.

BDELLIUM, *Myrrha imperfecta, Bilehon, Madeleon, Balchus.* A gum-resin, brought from the Levant and India, and supposed to be obtained from a species of *Amgris*, little known. It is solid, brittle, of a deep brown colour, of an acrid and bitter taste, and sweet odour. It was much vaunted by the ancients, but is now little employed. Two different gum-resins have been in the shops, distinguished by the names *Indian* and *African bdellium*. Dr. Royle was informed that the former was obtained from *Amgris Commiphora*, growing in India and Madagascar. The latter is said to be from *Hendelotia Africana*, which grows in Senegal.

BDELLOMETER, from *βδέλλα*, 'a leech,' (*βδέλλα*, 'I suck,' and *μετρον*, 'measure.' An instrument, proposed as a substitute for the leech; inasmuch as we can tell the quantity of blood obtained by it, whilst we cannot by the leech. It consists of a cupping-glass, to which a scarificator and exhausting syringe are attached.

BDELLUS, Fart.

BDELYGMIA, Fart.

BDELYGMUS, Fart.

BDESMA, Flatulence.

BDOLUS, Fart.

BEAD TREE, *Melia Azedarach.*

BEAL, Phlegmon, to Suppurate.

BEALING, Pregnant.

BEAN, CARTHAGENA, *Habilla de Carthagena*—b. Egyptian, *Nymphaea nelumbo*—b. French, *Phaseolus vulgaris*—b. Garden, common, *Vicia faba*—b. Indian, Catalpa—b. Kidney, *Phaseolus vulgaris*—b. Malacca, *Avicennia tomentosa*—b. Pontic, *Nymphaea nelumbo*—b. Red, *Abrus precatorius*—b. Sacred, *Nelumbium luteum*—b. St. Ignatius's, *Ignatia amara*—b. Tonka, *Dipterix odorata*—b. Tree, Catalpa—b. Tree, white, *Cratogeomys aria*.

BEAN TREFOIL TREE, see Cytisine.

BEARBERRY, *Arbutus uva ursi*.

BEARD, (Sax.) beard, (G.) Bart., *Barba, Pogon, Genion, Barbittium*, (F.) *Barbe*. The hair which covers a part of the cheek, the lips, and chin of the male sex, at the age of puberty.

BEAR'S BREECH, *Acanthus mollis*—b. Foot, *Helleborus fetidus*—b. Fright, *Heptallon graveolens*—b. Grass, *Yucca filamentosa*—b. Whorlberry, *Arbutus uva ursi*.

BEARWEED, *Veratrum viride*.

BEASTINGS, Colostrum.

BEASTLINGS, Colostrum.

BEATING OF THE HEART, see Heart.

BEAUGENCY, MINERAL WATERS OF. Beaugency is a quarter of a league from Orleans.

The waters contain subcarbonate of soda, iron, magnesia, and lime. They are tonic and aperient.

BEAUMONT ROOT, *Gillenla trifoliata*.

BEAUVAIS, MINERAL WATERS OF. These waters are chalybeate. Beauvais is in Picardie, France.

BEAVER, Castor fiber—b. Wood, *Celtis occidentalis*. *Magnolia glauca*—b. Tree, *Magnolia glauca*, *M. macrophylla*.

BEBEERIA, see Bebeeru.

BEBEERINE, see Bebeeru.

BEBEERU, *Sipeeri*. A tree of British Guiana, which yields two alkalies—*Bebeerine*, *Bebeerine*, *Bebeerine* or *Biberine*, and *Sipeerine*; and in its properties resembles the *Cinchona*. It has been referred to *Nectandra Rodiei*. Ord. Laurineæ. The timber of the tree is known to ship-builders by the name *libicu* or *green-heart*. The *Sulphate of Bebeeria* has been employed in intermittents. *Warburg's Fever Drops*, *Tinctura antifebrilis Warburgi*, an empirical antiperiodic preparation, have by some been considered to be a tincture of the seeds of the Bebeeru, but this is questionable.

BEC, (F.) Rostrum, Beak. This name has been applied to various parts.

BEC CORACODIEN, (F.) Cor'acoid beak, is the end of the coracoid process.

BEC DE CUILLE, Hamulus. An instrument used for the extraction of balls. It consists of an iron rod, 7 or 8 inches long, having at one extremity a small cavity, into which the ball is received to be drawn outwards. See *Cochleariformis*.

BEC DE GRUE MUSQUÉ, *Geranium Moschatum*—b. *de Grue Robertin*, *Geranium Robertianum*—b. *de Lièvre*, *Harelip*.

BEC DE LA PLUME À ÉCRIRE (F.), *Beak of the Calamus Scriptorius*, is a small cavity at the superior part of the medulla oblongata, which forms part of the 4th ventricle.

BEC, (LE.) MINERAL WATERS OF. Bec is six leagues from Rouen, in Normandy. The water is strongly chalybeate.

BECCABUNGA, *Veronica Beccabunga*.

BECHÆSTHESIS, from *βήξ*, *βήξος*, 'cough,' and *αίσθησις*, 'sensation.' The excitement or desire to cough.

BECHIA, Tussis.

BECHIAS, Tussis.

BE'CHICS, *Be'chica, Becha, Bee'chica, Be'chita*, from *βήξ*, *βήξος*, 'cough,' (F.) *Béchiques*. Medicines adapted for allaying cough.

BECHITA, Bechie.

BECHUM, Tussilago.

BECHORTHOPNEA, Pertussis.

BECUIBA, Ibicniba.

BED, HYDROSTATIC, Water bed—b. Water, Arnott's, see Water bed.

BED'EGAR, Bedeguar, Bedeguard, Spon'gia Cynosbati, Fungus Rosarum, F. Cynosbati, (F.) Pomme mousseuse, Éponge d'eglantier. An excrescence, which makes its appearance on different species of wild roses, and which is produced by the puncture of a small insect.—*Cynips Roseæ*. It was formerly employed as a lithontriptic and vermifuge, but is not now used. It was slightly astringent.

BEDFORD, MINERAL WATERS OF. Bedford is a village, situate on the great Western Turnpike road from Philadelphia to Pittsburg, a few miles east of the chief elevation of the Alleghany mountains. The "Springs" are about a mile and a half from the village. They are saline and sulphureous. The most celebrated—the "Mineral Spring" or "Anderson's Spring"—contains carbonic acid, sulphate of magnesia, chlorides of sodium and calcium, and carbonate of iron, but none of these articles in great quantity; hence, the main action of the waters is dia-

retic. At some distance from the Springs, there is a chalybeate water; and about 10 miles S. W. of Bedford, at Milliken's Cove, a strong sulphureous spring. The climate of Bedford Springs is agreeable. During the heat of summer, the nights are generally cool. In the morning of July 14, 1853, the author marked the thermometer, at five o'clock, A. M., at 45°. Three days before, however, it was at 70° at the same hour.

BEDLAM, Morotrophium.

BEDLAMITE, Insane.

BEDSTRAW, *Galium verum*—b. Ladies, greater, *Galium, mollugo, Galium verum*—b. Rough, *Galium asprellum*—b. Ladies, rough, *Galium asprellum*.

BEE, Sax. beo. *Apis, A. mellifica* seu domestica, *Melissa, Melitta*, (F.) *Abille*. This insect was formerly exhibited, when dried and powdered, as a diuretic.

BEE IN THE BONNET, see Insanity.

BEEBREAD, Propolis.

BEECH, *Fagus sylvatica*—b. Drop, *Orobanche Virginiana*—b. Drops, false, *Hypopitys lanuginosa*.

BEECH, ALBANY, *Pterospora Andromedea*.

BEECH, MAST, see *Fagus sylvatica*.

BEDDY'S EYES, *Viola tricolor*.

BEEF ESSENCE, see Beef tea.

BEEF TEA, *Jus horum*. An infusion of beef, much used in debilitating maladies, and in convalescence. It may be made as follows: Take two pounds and a half of lean beef: cut it, in small pieces, into three parts of water in an earthen pipkin: let this simmer, but never boil, until the liquor is consumed to a pint and a half: then strain carefully. It ought to be entirely free from fat or grease.—Dr. E. J. Seymour.

Essence of beef—as it has been called—may be made by putting a pound of good beef, freed from fat, and cut into small pieces, into a porter-bottle, corking lightly. The bottle must be put into boiling water, and kept there until the water has been boiling at least half an hour. As the boiling goes on, the cork may be inserted a little more tightly, to retain the contents of the bottle. The juices of the beef are thus separated, and constitute the 'essence,' which may be seasoned to the taste. It contains much nutriment.

BEEH, *Centaurea behen*.

BEER, *Cerevisia*—b. Black, see Falltrank—b. Chowder, see Chowder—b. Pipsissewa, see *Pyrola umbellata*.

BEER, JEWS', see *Pinus sylvestris*—b. Springs, see Water, Mineral (giseous.)

BEER, SPRUCE, *Cerevisia Abietia*, may be prepared as follows. Take of *Essence of Spruce*, half a pint: *Pimento*, bruised; *Ginger*, bruised; *Hops*, of each, four ounces; water, three gallons. Boil for five or ten minutes; strain, and add of warm water eleven gallons; yeast, a pint; molasses, six pints. Mix and allow to ferment for twenty hours. Used as an agreeable drink in summer.

BEER, TAR, see *Pinus sylvestris*.

BEET, Beta.

BEEST, Colostrum.

BEESTINGS, Colostrum.

BEETLEWEED, *Galax aphylla*.

BÉGALEMENT, Balbuties.

BEGMA,—according to some, *Bregma*,—from *βρεχω* or *βρεχωμαι*, 'to expectorate after coughing.' Coughing; also, the sputum or expectorated matter.—Hippocrates.

BEGONIA. The *Begonia grandiflora* and *B. tomentosa* have astringent roots, which are used in Peru in cases of hemorrhage, scurvy, low fevers, &c.

BÈGUE, Balbus.

BEHEN ABIAD, *Centaurea behen*—b. Album, *Centaurea behen*—b. Officinatum, *Cucubalus behen*—b. Range, *Statice limonium*—b. Vulgaris, *Cucubalus behen*.

BEHMEN ACKMAR, *Statice limonium*.

BEHAHALALEN, *Sempervivum testorum*.

BEIDELASAR, *Asclepias procera*.

BEIST, Colostrum.

BEISTINGS, Colostrum.

BEISTYN, Colostrum.

BEJUIO, *Habilla de Carthagera*.

BELA-AYE or BE-LAHE, A tonic and astringent bark of a Madagascar tree. Du-petit-Thouars and Sonnerat think it may be substituted for the Sinarouba.

BELADAMBOC. A species of convolvulus of the Malabar coast, which contains an acid milky juice. From this a liniment is formed with oil and ginger, which is used against the bites of rabid animals.

BE-LAHE, Bela-aye.

BELA-MODAGAM. A kind of *Scævola* of the Malabar coast, the leaves of which are considered diuretic and emmenagogue.

BELANDRE, (F.) A litter, surrounded with curtains, in which patients are sometimes carried to hospitals.

BELCHING, Eructation.

BELD, Athrix.

BELEMNOID, Belenoid.

BELEMNOIDES PROCESSUS, Styloid processes.

BEL'ENOID, BEL'ONOID, BEL'EMNOID or BEL'OID, *Belenoides* or *Belenoides Processus*, from *belos*, 'an arrow,' and *eidos*, 'shape.' This name has been given to styloid processes in general.—*Processus belenoides*.

BELESME, see Bellesme.

BELESON, Balsam, *Mussaenda frondosa*.

BELILLA, *Mussaenda frondosa*.

BELINUM, *Apium Graveolens*.

BELI OCLUS, Belloculus.

BELL, CANTERBURY, *Campanula trachelium*.

BELLADONE, *Atropa belladonna*.

BELLADON'NA, in the Pharmacopœia of the United States, is the official name of the leaves of *Atropa Belladonna*.

BELLADONNA BACCIFERA, *Atropa belladonna*—b. Trichotoma, *Atropa belladonna*.

BELLADONNIN, Atropine.

BELLE-BLOME, *Narcissus pseudo-narcissus*.

BELLE DAME, *Atropa belladonna*.

BELLEGU, *Nyrobalanus*.

BELLEREGI, *Nyrobalanus*.

BELLESME, MINERAL WATERS OF.

Bellesme is about three leagues from Montagne, in France. The waters are chalybeate.

BELLEY, MINERAL WATERS OF. The waters at Belley, department of Ain, in France, are saline aperients.

BELLFLOWER, *Narcissus pseudo-narcissus*.

BELLIDOIDES, *Chrysanthemum leucanthemum*.

BELLIS, *Bellus*, ('pretty,') *Bellis perennis* seu minor seu hortensis, *Synphytum minus*, *Brickwort*, Common Daisy, (Sc.) *Ewe-gowan*, (Prov.) *Bairnswort*, (F.) *Paquerette vivace*, *petite Marguerite*. Ord. Compositæ. The leaves and flowers are rather acid. They were, at one time, considered to cure different species of wounds. See *Osmitopsis asteriscoides*.

BELLIS HORTENSIS, *Bellis*—b. Major, *Chrysanthemum leucanthemum*—b. Minor, *Bellis*—b. Perennis, *Bellis*—b. Pratensis, *Chrysanthemum leucanthemum*.

BELLIT, Athrix.

BELL METAL, *Cal'cocos*, (F.) *Airain*, *Métal des cloches*. An alloy of copper, zinc, tin, and a

small quantity of antimony, used for making bells. The mortars of the apothecary are often formed of this material. They require to be kept clean, to avoid the formation of verdigris.

BELLOCULUS, *Beli Oculus*. A kind of gem, which the Assyrians considered efficacious in the cure of many diseases. They imagined that the figure of an eye could be seen in it, and hence its name. *Beli's Eye*, from the god Bel.

BELLON, Colic, metallic.

BELLOTAS, see *Ilex major*.

BELLOWS' SOUND, *Bruit de soufflet* — b. s. Encephalic, see *Bruit de soufflet*.

BELLOWS' SOUND, *Funic*, a single murmur of the bellows kind, synchronous with the first sound of the heart; heard by some observers, and referred by them to diminished calibre of the umbilical arteries, either by pressure or stretching of the funis, or both.

BELLOWS' SOUND, *PLACENTAL*, *Bruit placentaire*.

BELLWORT, *SMALLER*, *Uvularia perfoliata*.

BELLY, *Venter*, from *Ir. bolg*, 'the belly, a bag or pouch' [?]. At the present day, the abdomen. Formerly, all the splanchnic cavities were called *bellies*; — the *lower belly*, *venter inferior*, being the abdomen; the *middle belly*, *venter medius*, the thorax; and the *upper belly*, *venter superior*, the head. Also, the womb. See *Venter*.

BELLY-ACH, *Colica*.

BELLY-ACH, *DRY*, *Colica metallica*.

BELLY-ACH, *ROOT*, *Angelica lucida*.

BELLY-BAND, *Belt*, *Russian*.

BELLY-BOUND, *Constipated*.

BELLY-GOD, *Glutton*.

BELLY-HARM, *Colic*.

BELLY, *POT*, *Physconia*.

BELLY-THIRA, *Colic*.

BELLY-TIMBER, *Aliment*.

BELLY-WARK, *Colic*.

DELMUSCHUS, *Hibiscus abelmoschus*.

DELNILEG, *Myrobalanus*.

BELOID, *Belenoid*.

BELOIDES PROCESSUS, *Styloid processes*.

BELONE, *Needle*.

BELONODES, *Styloid*.

BELONOID, *Belenoid*.

BEL'S EYE, *Belloculus*.

BELT, *RUSSIAN*, *Ventra'le*, — vulgarly, *Belly-band*, — *Abdominal supporter*. A broad bandage applied to the abdomen, so as to support, and make methodical pressure upon it. Different forms have been termed *obstetric binders*, *utero-abdominal supporters*, &c.

BELUL'CUM, from *βελος*, 'a dart,' and 'ελκω', 'I draw out.' An instrument used for extracting darts or arrows. Many instruments of this kind have been noticed by surgeons. — Ambrose Paré, Fabricius ab Aquapendente.

BELZOE, *Benjamin*.

BELZOIM, *Benjamin*.

BELZOINUM, *Benjamin*.

BEN, *Guilandina moringa* — b. of Judæa, *Benjamin* — b. *Nut*, *Guilandina moringa*.

BENATH, *Pustule*.

BENEDICTA SYLVESTRIS, *Geum rivale*.

BENEDICTUM LAXATIVUM, *Confectio senne*.

BÉNÉFICE DE LA NATURE, *Beneficium natura* — b. *de Ventre*, see *Beneficium nature*.

BENEFIC'IUM NATU'RÆ, (F.) *Bénéfice de la nature*. This term is used by the French pathologists for cases in which diseases have got well without medical treatment. With them, *Bénéfice de nature*, or *B. de ventre*, is synonymous also with *Alvi Proflu'vium*; — a spontaneous diarrhoea, often acting favourably either in the prevention or cure of disease.

BENEL, *Croton racemosus*.

BENEOLENS, from *bene*, 'well,' and *olere*, 'to smell.' *Eudotis*, *Succedans*. A sweet-scented medicine, as gums, &c.

BENCI, *Bangue*.

BENGALÉ INDORUM, *Cassumuniar*.

BENGAL ROOT, *Cassumuniar*.

BENGI, *Hyoseyamus*.

BENIGN', *Benignus*, *Euth'es*, (F.) *Bénin*, *Bénigne*. Diseases of a mild character are so called: as well as medicines whose action is not violent, as a *Benign Fever*, *Febri benigna impetris*, &c.

BÉNIN, *Benign*.

BENJAMIN, corruption of *Ben'zoin*, *Benzo'inum* (Ph. U. S.), *B. ceruum*, *Assu adon'tu seu dulcis*, *Benjai*, *Benjoin*, *Assa* or *Ben'joyn*, *Benjo'inum*, *Belzoe*, *Belzoim*, *Ben'zoe*, *Styl'acis Benzoini Balsamum*, *Liquor Cyped'iacus*, *Canton Benzoë*, *Ben of Judæ'a*, *Acor Benzo'inus*, *Sal Ac'idum seu essential'e seu volat'il'e Benzoë*, (F.) *Benjoin*, *Baume Benjoin*, *Assa doux*. A resinous, dry, brittle substance, the concrete juice of *Styrax Benjoin*, *Arbor Benivi*, *Laurus Benjoin*, of *Sumatra*. The odour is extremely fragrant, and taste slightly aromatic. It is principally used for the preparation of the acid which it contains. It is also employed in some vulnerary tinctures, and as an expectorant. *Benzoic Acid*, *Ac'idum Benzo'icum*, is obtained from it by sublimation. The purest Benjamin is in *amygdaloid masses*: hence called (F.) *Benjoin amygdaloïde*.

BENJAMIN BUSH, *Laurus benzoïn*.

BENJAMIN, *FLOWERS OF*, *Ben'zoic Acid*, *Ac'idum Benzo'icum seu Benzoylicum*, *Flores Benzoë* seu *Benzo'ini*, *Ac'idum Benzo'icum per sublimatio'nem*, (F.) *Acide Benzoïque*, *Fleurs de Benjoin*. This acid exists in all the balsams, but chiefly in *Benjoin*, from which it is obtained by sublimation. It is in vanilla, canella, the urine of infants, and of herbivorous animals. Its odour is aromatic and fragrant; taste hot, slightly acidulous, and agreeable. The crystals consist of white, satiny flakes, slightly ductile. It is probably stimulant; and has been used, as such, in chronic catarrh; but it has little efficacy.

BENJAOY, *Benjamin*.

BENJOIN, *Benjamin*.

BENJOINUM, *Benjamin*.

BENJUI, *Benjamin*.

BEN MOENJA. A Malabar tree. An alexipharmic decoction is made of its roots, in the country, which is much praised in cases of malignant fever. Its bark, boiled with *Calamus aromaticus* and salt, forms a decoction used in bites of poisonous serpents.

BENNE, *Sesamum orientale*.

BENNET, *HERB*, *Geum urbanum*, and *Q Virginianum*.

BENOÎTE, *Geum urbanum* — b. *Aquatique*, *Geum rivale* — b. *des Ruiss'eaux*, *Geum rivale* — b. *de Virginie*, *Geum Virginianum*.

BENZENE, *Benzole*.

BENZIN, *Benzole*, see *Anæsthetic*.

BENZOATE OF AMMONIA, *Ammonia benzoas*.

BENZOE, *Benjamin*.

BENZOENIL, *Vanilla*.

BENZOIN, *Benjamin* — b. *Odoriferum*, *Laurus Benjoin*.

BEN'ZOLE, *Benzin*, *Benzene*, *Phene*, *Hy'dru-ret of Phen'yle*. A constituent of coal-gas tar, obtained by distilling coal naphtha. It is a colourless, oleaginous liquid, of agreeable odour. Sp. gr. 0.85. The vapour, when inhaled, acts as an anæsthetic.

BERBERINE, see *Oxyacantha Galeni*.

BERBERIS, *Oxyacantha Galeni* — b. *Canadensis*, see *Oxyacantha Galeni*.

BERRINE, *Verbena officinalis*.

BERCE, *Hieracium spondylium*.

BERENDAROS, *Ocimum basilicum*.

BERENICE, *Succinum*.

BERENICIUM, *Potassæ nitras*.

BERENISECUM, *Artemisia vulgaris*.

BERGAMOT, *Bergamote*, *Bergamot'ta*, (F.) *Bergamotte*. A small orange, of a very agreeable taste, and peculiar odour. From its rind an oil, *Oleum Bergamiti* (Ph. U. S.), is obtained, which is much employed as a perfume, and sometimes in medicine. See *Citrus mella rosa*.

BERIBERI, *Beriberia*, *Synclonus Beriberia*, *Indosynclonus*, *Paralysis Beriberi*, from *beri* in the Singalese language, which signifies 'weakness;' therefore, *beriberi*, 'great weakness.' This word is also said to be Hindustanee, and to mean a *sheep*.—Bontius. Beriberi is an Indian disease, little known in Europe. It consists in debility and tremors of the limbs,—sometimes, indeed, of the whole body; with painful numbness of the affected parts, &c.—the patient walking doubled, and imitating the movements of sheep! Some authors have esteemed it rheumatic; others, paralytic; others, to be a kind of chorea. It is, almost always, incurable; is rarely fatal; and is treated by exercise, stimulant friction, sudorifics, &c. It is sometimes called *Bambiers*, but this would seem to be a different disease.

BERICOCCO, *Prunus armeniaca*.

BERKLEY SPRINGS, see Bath Springs.

BERLE NODIFLORE, *Sium*.

BERLUE, *Metamorphosis*.

BERMUDAS, CLIMATE OF. Pulmonary invalids are occasionally sent to Bermuda, but the principal objection to a winter residence there, is the prevalence of strong winds; especially of the dry, sharp, and cold north-west winds, during the winter and spring. Still, it affords a good winter retreat for the phthisical, from any part of the United States, provided due care be selected in choosing a suitable locality. The neighbourhood of Hamilton has been strongly recommended with this view.

BERNARD THE HERMIT, *Cancellus*.

BERRIES, INDIAN, see *Menispermum cocculus*—b. Turkey, yellow, see *Piper cubeba*.

BERS. A sort of electuary, composed of pepper, seed of the white hyoscyamus, opium, euphorbia, saffron, &c. The Egyptians used it as an excitant.—Prospero Alpini.

BERU, MINERAL WATERS OF. Beru is in Champagne, France. The waters are slightly chalybeate.

BERULA, *Sium nodiflorum*—b. *Angustifolia*, *Sium nodiflorum*.

BESASA, *Ruta*.

BESICLES, Spectacles.

BESOIN, Want—b. *de Respirer*, see Want—b. *de la Vie*, Necessary of life.

BESSANEM. A word used by Avicenna, for redness of the skin, limbs, and face, produced by the action of cold.

BENNON,

BENNONNE, } See Gemellus.

BETA. The Beet, *Sicula*, (F.) *Bette*, *Betterave*. Family, Chenopodæ. Sex. Syst. Pentandria Digynia. A genus of plants, of which the following are the chief varieties:

BETA HYBRIDA, *Root of Scarcity*. Root red, outside; white, within. Very nutritive; yields sugar.

BETA VULGARIS ALBA, *White Beet*. The root yields sugar, and the leaves are eaten as a substitute for spinach.

BETA VULGARIS RUBRA, *Red Beet*. Root red and nutritive; yields a small quantity of sugar.

BETAYNE, *Betonica officinalis*.

BETEL, *Piper Betel*, *Chavica betle*. A species of pepper, cultivated in several parts of India. The East Indians are in the habit of chewing the leaves with lime and areca; and they give the name *Betel* to this preparation. It is used in all the equatorial countries of Asia. Betel is said to be tonic and astringent. It is also called *Bette*, *Bètre*, *Bette*. See *Areca*.

BETIROOT, *Trillium latifolium*—b. Broad-leaf, *Trillium latifolium*.

BÉTISE, *Dementia*.

BÉTOINE, *Betonica officinalis*—b. *des Montagnes*, *Arnica montana*—b. *des Savoyards*, *Arnica montana*.

BÉTON, *Colostrum*.

BETONICA AQUATICA, *Scrophularia aquatica*.

BETONICA OFFICINALIS, *B. purpurea*, *Vettonica*, *Cestron*, *Bet'ony*, *Wood Betony*, (Old Eng.) *Betayne*, *Psychotrophum*, *Veronica purpurea*, (F.) *Bétoine*. Order, Labiatae. Sex. Syst. Didynamia Gymnospermia. Betony was in much esteem amongst the ancients, who employed the flowers and leaves, in decoction, in gout, sciatica, cephalalgia, &c. It was so called, according to Pliny, from being in great repute among the Vettones or Bettones, an ancient people of Spain. Antonius Musa is said to have written a volume in praise of it; recommending it in no less than 47 different diseases. It has, however, little or no virtue. The leaves are said to be aperient, and the root emetic.

BETONICA PAULI, *Veronica*.

BETONY, *Betonica officinalis*—b. Paul's, *Lycopus sinuatus*, *Lycopus Virginicus*, *Veronica*—b. Water, *Scrophularia aquatica*—b. Wood, *Betonica officinalis*.

BÈTRE, *Betel*.

BETTE, *Beth*.

BETTERAVE, *Beta*.

BETULA ALBA, *Эрвва*. The Birch, (Sc.) *Birk*, (F.) *Bouleau commun*. Order, Betulaceæ. The young leaves are slightly odorous, astringent, and bitter. They are applied to wounds and ulcers. They have been regarded as antiscorbutic and anthelmintic. The tree furnishes a saccharine juice, which is considered antiscorbutic and diuretic.

BETULA EMARGINATA, *Alnus glutinosa*—b. *Glutinosa*, *Alnus glutinosa*.

BETULA LENTA, *Sweet Birch*, *Black Birch*, *Cherry Birch*, *Mountain Mahogany*, is an American species, the bark and leaves of which have the smell and taste of *Gaultheria procumbens*. An infusion is sometimes made of them, and used as an excitant and diaphoretic. The volatile oil is nearly, if not wholly, identical with that of *Gaultheria*.

BEURRE, Butter—b. *de Bambouc*, Butter of bambouc—b. *de Cacao*, Butter of cacao—b. *de Coco*, Butter of cocoa—b. *Végétale*, *Persea gratissima*.

BEUVRIGNY, MINERAL WATERS OF. Beuvrigny is in the vicinity of Bayeux, in Normandy. The water is chalybeate.

BEVERAGE, Drink.

BÉVUE, *Diplopia*.

BEX, Tussis—b. *Convulsiva*, *Pertussis*—b. *Humida*, *Expectoration*—b. *Theriodes*, *Pertussis*.

BEXIS, Tussis.

BEXUGO. Under this name, a purgative root was formerly introduced into Europe from Peru. It is supposed to have been the root of a Hippocraten.

BEZ' OAR, *Bez' oar*, *Bez' ehard*, *Pa' zahar*, from Persian *Pa*, 'against,' and *zahar*, poison. *Lapis Bezoar'dicus*, *Calculus Bez' oar*, *Enterolithus Bezoar'dus*, *Bezoard*. A calculus concretion, found

in the stomach, intestines, and bladder of animals. Wonderful virtues were formerly attributed to these Bezoars. There were two great varieties: the *Bezoar orientalis*, *Animal Bezoar-ticum orientale*, formed in the fourth stomach of the gazelle of India (*Gazella Indica*, or rather *Antilopæ cervicapra*;) and the *Bezoar occidentalis*, *Animal Bezoar-ticum occidentale*, found in the fourth stomach of the wild goat or *chamois* of Peru. These substances were esteemed to be powerful alexipharmics; but the former was the more valued. It was believed that no poison, and no eruptive, pestilential, or putrid disease, could resist its influence. As so many virtues were ascribed to it, other animal concretions were substituted for it; and factitious Bezoards were made of crabs' eyes and claws, bruised and mixed with musk, ambergris, &c.

BEZOAR BONTINUM, (F.) *Bézoard de Boeuf*, *Bezoard of the beef*. A concretion formed in the fourth stomach of beeves; also, a biliary calculus found in the gall-bladder.

BEZOARD OF THE DEER, *B. of the Luchrymal Fossa of the Deer*, *Deer's Tears*. A moist, highly odorous, fatty matter, found below the anterior caecum of the orbit of the red deer—*Cervus elaphus*. It has been used, like castor, as an antispasmodic, in the dose of from 5 to 15 grains, two or three times a day.

BEZOAR EQUINUM, *Bezoard of the horse*—b. *Hystriels*, *Bezoard of the Indian porcupine*.

BEZOARD OF CAYMAN. This was once much prized. It is now unknown.

BÉZOARD D'ALLEMAGNE, *Ægagropila*.

BEZOARD OF THE CHAMOIS, and **B. OF THE HONSE**, *Bezoar equinum*, *Hippolithus*, &c., exhibit their origin in the name.

BEZOARD OF THE INDIAN PORCUPINE, *Bezoar Hystriels*, *Lapis Porcinus seu Malucensis*, *Petro del Porco*, (F.) *Bézoard de Porc-Épie*, was formerly the dearest of all the Bezoards, and was sold at an enormous price in Spain and Portugal.

BEZOARD, MINERAL, *Antimonium diaphoreticum*—b. *Vegetable*, see *Calappite*.

BEZOARDIC, *Bezoar'dicus*, (F.) *Bézoardique*; concerning the bezoard. Bezoardic medicines are those supposed to possess the same properties with the bezoard; as antidotes, alexiteria, alexipharmics, cordials.

BEZOARDICA RADIX, *Dorstenia contrajerva*.

BEZOARDICUM SATURNI. A pharmaceutical preparation, regarded by the ancients as antihysteric. It was formed of protoxide of lead, butter of antimony, and nitric acid.

BEZOARDICUM HUMANUM. Urinary calculi were formerly employed under this name as powerful alexipharmics.

BEZOARDICUM JUVIALÈ. A sort of greenish powder, used as a diaphoretic, and formed of antimony, tin, mercury, and nitric acid.

BEZOARDICUM LUNARÈ. A medicine formerly regarded as a specific in epilepsy, convulsions, megrim, &c. It was prepared of nitrate of silver, and butter of antimony.

BEZOARDICUM MARTIALÈ. A tonic medicine, used by the ancients in diarrhoea. It was prepared from the tritoxide of iron and butter of antimony.

BEZOARDICUM MERCURIALÈ. A medicine, formerly vaunted as an antisiphilitic, and prepared from the mild chloride of mercury, butter of antimony, and nitric acid.

BEZOARDICUM MINERALÈ; the deutoxide of antimony; so called because its properties were supposed to resemble those of animal bezoard.

BEZOARDICUM SOLARÈ. A diaphoretic medi-

cine, prepared of gold filings, nitric acid, and butter of antimony.

BEZOARDICUM VENERIS. A pharmaceutical preparation, formerly employed in lepra, diseases of the brain, &c.; which was made from filings of copper, butter of antimony, and nitric acid.

BHANG, *Bangue*.

BI, as a prefix to words, has the same signification as *Di*.

BIAOTHANATI, *Biathanati*.

BIBERINE, see *Bebeera*.

BIBIRU, *Bebeeru*.

BIBITORIUS, *Rectus internus oculi*.

BIBLIOGRAPHY, MEDICAL, from *βιβλίου*, 'a book,' and *γραφω*, 'I describe.' A description or history, or skill in the knowledge, of medical books. The most distinguished medical bibliographers have been: J. A. VAN DER LINDE, Amsterdam, 1662, octavo, (L.) M. LIPENIES, France, ad Mœn. 1679, fol. (L.) G. A. MERCKLEIN, Norimb. 1686, (L.) J. J. MANGET, Genex, 1695 to 1731, (L.) TARIN (anatomical), Paris, 1753, (F.) A. VON HALLER, Zürich, 1774, &c., (L.) VIGILIUS VON CRUETZENTELD (surgical), Vindob. 1781, (L.) C. G. KUHN, Lips. 1794, (L.) C. L. SCHWEIKARD (anat., phys., and legal medicine), Stuttgart, 1796 to 1800, (L.) G. G. PLOEQUET, Tubing. 1808 to 1814, (L.) C. F. BURDACH, Gotha, 1810 to 1821, (G.) J. S. ERSCH. (since 1750), Leipz. 1822, (G.) TH. CH. FR. ENSLIS, (of Germany, since 1750.) Berlin, 1826, (G.) J. B. MONTFALCON, Paris, 1827, (F.) J. FORBES, M.D., F.R.S., London, 1835. A. C. P. CALLSEN, Copenhagen, 1845, (G.) E. MORWITZ, Leipz. 1849, (G.)

BICAUDALIS, *Retrahens auris*.

BICAUDATUS, *Caudatus*, from *bi*, and *cauda*, 'a tail;' 'double-tailed.' A monster having two tails.

BICEPHALUM, *Dicephalum*. A hybrid word, from *bi* and *κεφαλή*, 'head.' Sauvages applies this epithet to a very large sarcoma on the head, which seems to form a double head.

BICEPHALUS, *Dicephalus*.

BICEPS, from *bi*, and *caput*, 'head.' That which has two heads. This name has been particularly given to two muscles; one belonging to the arm, the other to the thigh.

BICEPS EXTERNUS MUSCULUS. The long portion of the *Triceps Brachialis*.—Douglas.

BICEPS FLEXOR CUBITI, *B. Cruris*, *Biceps*, (F.) *Biceps Cruralis*, *Biceps Femoris*, *Ischio-femoro-pæronius*—(Ch.)—A muscle on the posterior part of the thigh; one head arising from the tuberosity of the ischium, and the other from a great part of the linea aspera. It is inserted into the top of the fibula. It serves to bend the leg on the thigh.

BICEPS FLEXOR CUBITI, *Biceps*, *B. Brachii seu manus seu internus seu internus humeri*, *Coraco-radialis*, (F.) *Scapulo-radialis*, (Ch.)—*Biceps Brachialis*. A muscle, situate at the anterior and internal part of the arm; extending from the edge of the glenoid cavity and from the top of the coracoid process to the tuberosity of the radius. It bends the fore-arm upon the arm.

BICHE DE MER, *Sea Slug*. A molluscan animal, belonging to the genus *Holothuria*, which is caught amongst the islands of the Feejee group, New Guinea, &c., and when prepared finds a ready sale in China, where it is used as an ingredient in rich soups.

BICHET, *Terra Orleansa*.

BICHICULE, probably corrupted from *βήχας*, 'cough.' Pectoral medicines, composed of liquorice juice, sugar, blanch almonds, &c.—*Rhazes*.

BICHIOS, *Dracunculus*.

BICHO, *Dracunculus*—b. *di Culo*, *Proctocæc*.

BICHOS. A Portuguese name for the worms that penetrate the toes of people in the Indies; and which are destroyed by the oil of the cashew nut.

BICIP'ITAL, from *biceps* (*bi* and *caput*), 'two headed.' Relating to the biceps.

BICIP'ITAL GROOVE, (F.) *Couliasse ou Gouttière bicipitale*, *Couliasse humérale*, (Ch.) is a longitudinal groove, situate between the tuberosities of the os humeri, which lodges the long head of the biceps.

BICIP'ITAL Tu'BERCLE, *Bicipital tuberosity*, (F.) *Tubérosité bicipitale*; a prominence near the upper extremity of the radius, to which the tendon of the biceps is attached.

BICORNE RU'DE, *Ditrachyceros*.

BICUSPID, *Bicuspidatus*, from *bi* and *cuspis*, 'a spear.' That which has two points or tubercles.

BICUSPID TEETH, *Dentes Bicuspidati*, (F.) *Dents bicuspidées*. The small molars. See Molar.

BIDDY, *Podiculus*.

BIDENS ACME'LLA, *Spilanthes acmella*.

BIDET, (F.) *Bidet*; pronounced *beeday*. A small horse formerly allowed to each trooper for carrying his baggage. Hence, perhaps, applied to a chamber bathing apparatus, which has to be bestridden. It is a useful arrangement, in case of hemorrhoids, prolapsus ani, affections of the sexual organs, &c.

BIECHO, *Bische*.

BIÈRE, *Cerevisia*—*b. Médicinale*, see *Cerevisia*.

BIESTINGS, *Colostrum*.

BIFÉMORO-CALCANIEN, *Gastrocnemii*.

BIFURCATION, *Bifurcatio*, from *bi* and *furca*, 'a fork.' Division of a trunk into two branches; as the *bifurcation of the trachea, aorta*, &c.

BIGARADE, see *Citrus aurantium*.

BIGASTER, *Digastricus*.

BIGBELLIED, *Abdominus*.

BIG BLOOM, *Magnolia macrophylla*.

BIGBONE, see *Kentucky*, *Mineral Waters of*.

BIGPAUNCHED, *Abdominus*.

BIGEMINAL BODIES, *Quadrigenina tubercula*.

BIGGAR. A disease of Bengal, remarkable for the intensity and danger of the cerebral symptoms. — Twining.

BIG-LEAF, *Magnolia macrophylla*.

BIGLES, see *Strabismus*.

BIGNONIA CAPREOLA'TA, *B. crucig'era*; indigenous. The root and plant are employed, like sarsaparilla, in infusion and decoction, in syphilis, chronic rheumatism, and in cachexia in general.

BIGNONIA CATALPA, *Catalpa*—*b. Crucigera*, *B. capreolata*.

BIGNONIA IN'DICA. The leaves are employed in India, as emollients, to ulcers.

BIGNONIA LEUCOX'YLON, *Trumpet tree*. A South American tree, the sap of which is the surest antidote against the manchineel.—Schleiden.

BIGNONIA RADICANS, *Tecoma radicans*—*b. Sem-pervirens*, *Gelseminum nitidum*.

BIJON, see *Pinus sylvestris*.

BIKH, *Aconitum ferox*.

BILATE OF SODA, *Picromel*.

BILATERAL SYMMETRY, see *Symmetry*.

BILAZAY, MINERAL WATERS OF. Bilazay is a town in France, two leagues from Thouar, department of Deux Sèvres, near which is a thermal sulphureous spring. Temperature about 77° Fahrenheit.

BILBERRY, *Vaccinium myrtillus*—*b. Red*, *Vaccinium vitis idæa*.

BILE, *Bilis*, *Fel*, *Chol'oe*, *Cholâ*, *Choler*, (F.)

Bile, *Fiel*. A yellow, greenish, viscid, bitter, nauseous fluid, secreted by the liver. It is distinguished into *hepatic* and *cystic*; according as it flows immediately into the duodenum from the liver or from the gall-bladder. It contains, according to Muratori, water; a peculiar fatty matter; colouring matter (*Cholepyrrhin* or *Bilipha'in*, *Matière Jaune de la Bile*; cholesterolin, combined with soda; picromel or *bilin*; extract of flesh, mucus; soda, phosphate of soda; phosphate of lime, and chloride of sodium.

The use of the bile is to remove from the body superfluous hydro-carbon; and it is probably inservient to useful purposes in digestion.

BILE, *Furunculus*—*b. Black*, *Atrabilis*—*b. de Boaf*, see *Bile*—*b. Repandue*, *Icterus*.

BILE OF THE BEAR, *Gall of the Bear*, *Fel Ursi*, was thought to be anti-epileptic; and that of the *Eel*, *Fel anguill'ar*, to facilitate labour.

BILE OF THE OX, *Gall of the Ox*, *Ox Gall*, *Fel Tauri* seu *Bovis* seu *Bovinum*, (F.) *Bile* ou *Fiel* ou *Amer de Boaf*, was once reputed cosmetic and detergent, anti-otalgie and emmenagogue; as well as to possess the power of facilitating labour. It has also been given as a bitter stomachic and antelmintic; and as a tonic and laxative, in cases of deficiency of the biliary secretion.

BILE OF THE SWINE, *Bilis porcina*, inspissated, has been used in the same cases as the bile of the ox.

BIL'IARY, *Bilia'ris*, *Bilia'rius*, *Fel'leus*, (F.) *Biliaire*. That which relates to bile.

BIL'IARY APPARA'TUS, *B. organs*, *B. passages*. The collection of parts that concur in the secretion and excretion of bile—viz., the liver, pori biliiari or tubuli biliferi; hepatic, cystic, and choledoch ducts, and gall-bladder.

BIL'IARY CONCRE'TIONS are concretions found in some parts of the biliary apparatus.

BILIARY DUCTS, *Pori biliiari*.

BILIEUX, *Bilious*.

BILIA'TION, *Bilia'tio*; from *bilis*, 'bile.' The excretion of bile; as *urination* means the excretion of urine.

BILIFULVINE; from *bilis*, 'bile,' and *fulvus*, 'reddish yellow.' The colouring matter of the bile of the ox.—Berzelius.

BILIMBI, *Averrhoa bilimbi*.

BILIMBING TERES, *Averrhoa bilimbi*.

BILIN, *Picromel*.

BILIOUS, *Bilio'us*, *Chol'icus*, *Chol'ius*, *Fel'in'us*, *Epich'olos*, *Pteroch'olos*, *Fel'leus*, (F.) *Bili'eux*; from *bilis*, 'bile.' That which relates to bile, contains bile, or is produced by bile. An epithet given to certain constitutions and diseases, which are believed to be the effect of superabundance of the biliary secretion; as *Bilious temperament*, *B. symptoms*, *B. fever*.

BILIPHÆIN, see *Bile*.

BILIS FLUXIO, *Cholera morbus*.

BILITICUS, *Cholagogue*.

BILIVERD'IN, from *bilis*, 'bile,' and *viridis*, 'green.' On adding an acid to a solution of the yellow colouring matter of bile, a precipitate of green flocculi takes place, which possesses all the properties of chlorophyll, or the green colouring matter of leaves. This is the *biliverdin* of Berzelius.

BILNA. MINERAL WATERS OF, see *Püllna*.

BILOCULAR, see *Unilocular*.

BILUMBI BITING-BING, *Malus Indica*.

BI'MANUS, from *bi* and *manus*, 'a hand.' One that has two hands. A term applied only to man, because he is the sole mammiferous animal that possesses two perfect hands.

BINDER, *Bandage*.

BINDERS. OBSTETRIC, see *Belt*, *Russian*.

BINDWEED, *Polygonum aviculare*—*b. Fiddle-leaved*, *Convolvulus panduratus*—*b. Great*,

Convolvulus sepium — b. Hedge. *Convolvulus sepium* — b. Lavender-leaved. *Convolvulus Cantabrigia* — b. Sea. *Convolvulus soldanella* — b. Virginian. *Convolvulus panduratus*.

BINDWOOD, *Hedera helix*.

BINKHUMBA, *Phyllanthus urinaria*.

BINOCTULAR, *Binoctularis*: same etymon as the next. Relating to or affecting both eyes — as '*binoctular vision*' — vision with both eyes; or from impressions made upon both retinas, which are amalgamated into *single vision*.

BINOCTULUS, *Binocle*, *Diophtalmica Falcia*, *Oculus duplex*, from *bi* and *oculus*, 'an eye.' (F.) *Oeil double*. A bandage applied over both eyes. It was also formerly called *Diophtalmus*.

BIN'SICA. A Rabbinical term for disorder of the mind. According to VAN HELMONT, an atrophy of the organ of imagination.

BIOCHYMIA, Chymistry, vital.

BIOID, *Vis vitalis*.

BIODYNAMICS, *Biodynamica*, *Biodynamica*, *Biochymia*, from *bios*, 'life,' and *dynamis*, 'power,' 'force.' The doctrine of the vital activity or forces.

BIOGAMIA, Magnetism, animal.

BIOLOGY, from *bios*, 'life,' and *logos*, 'a description.' The doctrine of life or of living bodies. — Physiology. "*Biological phenomena*" are those presented by living bodies.

BIOLYCH'NION, *Biolych'nium*, from *bios*, 'life,' and *lych'nium*, 'a lamp.' Innate heat, vital heat, animal heat. *Lych'nium*, *Lychnid'ium*, *Thermum caphyllum*, *Flamma* seu *Flammula vitalis* seu cordis. Also, a secret preparation of which BEERIS and RESERVE make mention.

BIOLYSIS, see *Biolysis*.

BIOLYTIC, *Biolyticus*; from *bios*, 'life,' and *lysis*, 'solution.' Relating to the destruction of life. A '*biolytic agent*' is one that tends to *biolysis*, or destruction of life. A depressing agent. — Schultz.

BIO-MAGNETISMUS, Magnetism, animal.

BIONOMY, Physiology.

BIOPHENOMENOLOGIA, Physiology.

BIOS, *bios*. Life. Also, what is necessary for the preservation of life.

BIOSPHIA, Biodynamics.

BIOSTATICS, Statistics, medical.

BIOTAX'Y, *Taxon'omy*, from *bios*, 'life,' and *taxis*, 'order, arrangement.' The arrangement or co-ordination of living organisms, according to their external characters.

BIOTE, Life.

BIOTHAN'ATI, *Biothanth'ati*, from *bios*, 'life,' and *thanas*, 'death.' Those who die of a violent death very suddenly, or as if there was no space between life and death.

BIOTIC, Vital.

BIOTICS, Physiology.

BIOTOMIA, Vivisection.

BIOTOS, Life.

BIPARIETAL, *Biparietalis*, (F.) *Biparietal*, from *bi* and *parietale* (os), 'the parietal bone.' An epithet for the diameter of the cranium from one parietal fossa to the other.

BIPARIETAL SUTURE, Sagittal suture.

BIPIN'NA, from *bi* and *pinnæ*, 'a wing-femur.' A term used by the ancients for a diminutive penis, not exceeding in size two quills.

BIR, Thorax.

BIRA, Cerevisia.

BIRCH, *Betula alba* — b. Black. *Betula lenta* —

b. Cherry, *Betula lenta* — b. Sweet, *Betula lenta*.

BIRDS' BREAD, *Sedum acre*.

BIRDS' NEST, *Hypopitys lanuginosa*.

BIRK, *Betula alba*.

BIRTH, Parturition — b. Cross, Presentation,

preternatural — b. Live, see *Born alive* — b. Plural, see *Multiparous*.

BIRTHWORT, *Aristolochia* — b. Snakeroot, *Aristolochia serpentaria*.

BISCHE, *Biecho*. A malignant kind of dysentery, which often prevails in the island of Trinidad.

BISCUIT, *Biscuitus*; *bis*, 'twice,' and *coctus*, 'baked,' (F.) *bis* and *cuit*, 'twice baked.' A kind of dry, hard bread, or cake, which is variously made; and, when without eggs or butter, is easy of digestion. It was formerly called *Dipyrrites*, and *Dipyrus*. MEDIC'INAL OR MEDICATED BISCUITS, (F.) *Biscuits Médicinaux*, *Massepains*, *Maccarons*, are occasionally made by adding to biscuit paste a medicinal solution or powder, and baking the mass.

BISCUIT, MEAT. An alimentary preparation, proposed by Mr. G. Borden, Jr., of Texas, which consists in combining the matters, extracted from meat by boiling, with flour, so as to form biscuits; which keep well, and are of course nutritive.

BISCUITS MÉDICINAUX, see *Biscuit*.

BISERMAS, *Salvia sclarea*.

BISERIENS, *Dicrotus*.

BISH, *Aconitum ferox*.

BISHOP'S WEED, *Ammi*.

BISLINGUA, *Ruscus hypoglossum*.

BISMALVA, *Althæa*.

BISMUTH, *Antimonium album*, *Chalcitas*, *Luna imperfecta*, *Stannum glaciæ* seu *cine'reum*, *Bismuthum*, *Wismuthum*, *Regulus of Bis'muth*, *Marcasita*, *Tin glass*, (F.) *Étain gris*, *É. de Glace*. A metal, in spicular plates, of a yellowish-white colour; s. gr. 9.822; fusible at 400° Fahrenheit, and volatilizable at a high temperature. It is used only in the preparation of the subnitrate.

BISMUTH, NITRATE OF, *Bismuth*, *Subnitrate of* — b. Oxide of, *Bismuth*, *Subnitrate of* — b. Oxide of, white, *Bismuth*, *Subnitrate of* — b. *Regulus of Bismuth*.

BISMUTH, SUBNITRATE OF, *Bismuthi subnitras* seu *Nitras*, *Marcasita alba*, *Plumbum cine'reum*, *Magisterium Marcasita* seu *Bismuthi*, *Bismuthum Nitricum* seu *Subnitricum*, *Nitras Subbismuthicum*, *Cale Bismuthi*, *Bismuthum oxygallatum album*, *Bismuthum album*, *Subazotus bismuthicus*, *Oxide of Bismuth*, *Nitrate of Bismuth*, *White Bismuth*, *Trisnitrate of Bismuth*, *White Oxide of Bismuth*, *Magisterium of Bismuth*, *Pearl White*, *Spanish White*. (F.) *Sousnitrate ou sousazotate de bismuth*, *Oxide blanc de B.*, *Blanc de fard*, *Blanc de perle*. (*Bismuth*, in frustulis, $\mathfrak{z}\mathfrak{i}$. *Acid nitric*, f $\mathfrak{z}\mathfrak{i}\mathfrak{j}$. *Aq. destill.* q. s. Mix a fluid-ounce of distilled water with the nitric acid, and dissolve the bismuth in the mixture. When the solution is complete, pour the clear liquor into three pints of distilled water, and set the mixture by, that the powder may subside. Lastly, having poured off the supernatant fluid, wash the subnitrate of bismuth with distilled water, wrap it in bibulous paper, and dry with a gentle heat. Ph. U. S.) It is considered to be tonic and antispasmodic, and has been chiefly used in gastrodynia.

BISMUTH, TRISNITRATE OF, *Bismuth*, subnitrate of.

BISMUTH, VALERIANATE OF, *Bismuthi valerianæ*, *Bismuthum valerianicum*. Prepared by mixing a neutral solution of oxide of bismuth in nitric acid, with valerianate of soda; washing, and drying the precipitate. Used in gastrodynia, chronic gastralgia, neuralgia, and chronic palpitation, as a nerveine. Dose, $\frac{1}{2}$ a grain to 2 grains, three or four times a day, in pill.

BISMUTH, WHITE, *Bismuth*, subnitrate of.

BISMUTHI NITRAS, Bismuth, subnitrate of—b. Valerianæ, Bismuth, valerianate of.

BISMUTHUM, Bismuth—b. Album, Bismuthi subnitræ—b. Nitricum, Bismuth, subnitrate of—b. Oxydulatum album, Bismuth, subnitrate of—b. Subnitricum, Bismuth, subnitrate of—b. Valerianicum, Bismuth, valerianate of.

BISPRUS, Dipnoos.

BISSUM, Hydrangen arboreus.

BISTORT, OFFICINAL, Polygonum bistorta—b. Virginian, Polygonum virginianum.

BISTORTA, Polygonum bistorta.

BISTORTE, Polygonum bistorta.

BISTORTIER, (F.) A name given by the *Pharmacien* to a long wooden pestle used for reducing soft substances to powder, and in the preparation of electuaries.

BISTOURI, (F.) *Pistorien'sis gladius, Scalpellus, Scalpium, Bistoury*. A small cutting-knife, used in surgery.—so called, according to Iluet, from the town of Pistori, which was formerly celebrated for the manufacture of those instruments. A bistoury has the form of a small knife, and is composed of a blade and a handle. The blade, which is most commonly movable in the handle, may be fixed by a button, spring, &c. When fixed in the handle, the bistouri is called by the French, *B. à lame fixe ou dormante*.

The chief bistouries are:—1. The STRAIGHT B. (F.) *B. droit*, in which the blade and cutting edge are straight, the point being fine, round, or square. 2. The CONVEX B. (F.) *B. convexe*; the blade of which is convex at the cutting edge, concave at the back. 3. The CONCAVE B. (F.) *B. concave*; the blade of which is concave at its edge, and convex at the back. 4. BLUNT-POINTED B. (F.) *B. boutonné*; the blade of which has a button at its extremity. 5. The BLUNT OR PROBE-POINTED BISTOURY of POTT; concave at its cutting edge, and its point blunt; so that it can be carried on the palmar surface of the index finger, to divide the stricture, in strangulated hernia. Sir Astley Cooper has recommended a useful modification of this, to avoid wounding the intestine, should it come in contact with the edge of the knife. His Bistoury has an edge of not more than eight lines in length, situate about five lines from the point. 6. BISTOURI À LA LIME, (F.) is a straight bistoury; the blade fixed in the handle, the extremity with a button, and the edge made with a file. It is chiefly used for dilating parts. 7. BISTOURI ROYAL, (F.) A Bistoury used in operating upon Louis XIV. for fistula in ano. 8. BISTOURI GASTRIQUE, (F.) A complicated instrument, invented by Morand, for dilating wounds of the abdomen. 9. BISTOURI CACHÉ, *B. herniaire, ou Attrape-lourdaud de Biennaise, Forceps de cepto'ria*. A curved bistouri, the blade of which is placed in a canula, whence it issues on pressing a spring.

The word *Bistouri* is used by the French, at times, where we would employ knife.

BITCH-DAUGHTER, Incubus.

BITNOBEN, *Salt of Bitu'men, Padnoon, Soucherloon, Khala mimuc*. A white, saline substance, which is a Hindoo preparation of great antiquity, and has been supposed to be the *Sal asphaltites* and *Sal Sodome'nis* of the ancients. It is used by the Hindoo in the prevention or cure of almost all diseases.

BITHNIMAL'CA, *Gas'teranax*. Two unmeaning words, used by Doleus, to designate an active principle supposed to have its seat in the stomach, and to preside over chymification, &c.

BITOS DE KIS, Proctoocæ.

BITTER, Amarus—b. Bark, Pinckneya pubens—b. Bloom, Chironia angularis—b. Holy, *Hiera piera*—b. Redberry, Cornus Florida—b. Root, Apocynum androsæmifolium, Gentiana

Catesbæi, Menyanthes verna—b. Sweet nightshade, Solanum Dulcamara—b. Sweet vine, Solanum Dulcamara—b. Welter's, Acid, carbazotic.

BITTERNESS, *Amaritudo, Amarit'ies, Amaror, P'cria*, (F.) *Amertume*. A particular taste, which belongs to many substances. In some diseases there is a sense of bitterness felt in the mouth.

BITTERS, COLUMBO, Tinctura Calumbæ—b. Spirit, Tinctura gentianæ composita—b. Wine, Vinum gentianæ compositum.

BITTERSWEET, Solanum dulcamara.

BITTERWEED, Ambrosia trifida.

BITTERWOOD TREE, Quassia.

BITTOS. A disease, in which the chief symptom is an acute pain in the anus.—Chomel.

BITUME DE JUDEE, Asphaltum.

BITUMEN, GLUTINOUS, Pissasphaltum—b. Judaicum, Asphaltum—b. of Judeæ, Digasticus. Asphaltum—b. Petroleum, Petroleum—b. Malta, Pissasphaltum—b. Salt of, Bitnoben—b. Solidum, Asphaltum.

BIVENTER, Digasticus—b. Cervicis, Complexus musculus—b. Maxille, Digasticus.

BIVENTRAL LOBE OF THE CEREBELLUM, see Lobe, biventral.

BIXA AMERICANA, see Terra Orleana—b. Orleana, see Terra Orleana—b. Orellana, see Terra Orleana.

BLABE, Wound.

BLACCIE, Rubcola.

BLACIA, Debility.

BLACKBERRY, AMERICAN, see Rubus fruticosus—b. High or Standing, see Rubus fruticosus—b. Low, Rubus trivialis.

BLACK BOTTLE, see Infusum Sennæ.

BLACK DOSE, see Infusum Sennæ.

BLACK DRAUGHT, see Infusum Sennæ.

BLACK DROP, Guttae nigrae.

BLACK LION. A term given to a sloughing syphilitic ulcer, under which the British soldiers suffered greatly in Portugal.

BLACK ROOT, Aletris farinosa, Pterocaulon pycnostachyum, Leptandria purpurea.

BLACKWATER, Pyrosis.

BLACTIE, Rubcola.

BLADDA, Buttermilk.

BLADDER, Urinary Bladder.

BLADDER, FASCIC'ULATED. A condition of the bladder in which the interior of the organ is marked by strong reticulated ridges or columns.

BLADDER, GALL, see Gall Bladder—b. Irritable, Cysterethismus.

BLADDER, SAC'ULATED. A condition of the bladder, in which the mucous coat protrudes through the muscular, so as to produce a hernia, which may go on increasing, so as to form a vesical sacculus or appen'dix vesicæ.

BLADDER, SWIM, Air bladder—b. Urinary, see Urinary Bladder.

BLADEBONE, Scapula.

BLADOCH, Buttermilk.

BLADON SPRINGS. Mineral springs in Alabama, about 85 miles from Mobile, which are said to be similar in composition to those of Seltzer, Spa, and Aix-la-Chapelle, of Europe.

BLADUM, Blé.

BLAEBERRY, Vaccinium myrtillus.

BLÆ'SITAS, *Blæsa lingua*. Some authors have used this word as synonymous with stammering. See Balbuties. Sauvages understands by it a defect in pronunciation, which consists in substituting soft consonants for those that are hard; as the z for s, the d for t, the s for g and j, &c. Also, Lisper, *Traul'simus, Trau'lotes*, (F.) *Blé-sité, Blé (parler)*.

BLÆSOPODES, see Kylosis.

BLESOPUS, see Kyllosis.

BLÆSUS. A distortion; especially the outward distortion of the legs. Also, a stammerer.

BLAFARD, (F.) *Pallidus*, *Pallidulus*. This epithet is sometimes given to the skin, when pale and dull; but, most frequently, to the flesh of a wound, when it has lost its colour, and become white. The word is, also, sometimes used synonymously with Albino.

BLAIN, (Sax.) *blegne*, *Bleine*. A pustule; a both; a blister. In Scotland, a mark left by a wound; the discolouring of the skin after a sore.—Jamieson.

BLAK OF THE EYE, Pupil.

BLANC D'ARGENT, Plumbi subcarbonas—*b. de Baleine*, Cetaceum—*b. de Céruse*, Plumbi subcarbonas—*b. de Crems*, Plumbi subcarbonas—*b. de Fard*, Bismuth, subnitrate of—*b. de Kremnitz*, Plumbi subcarbonas—*b. de Krems*, Plumbi subcarbonas—*b. de l'Œil*, Sclerotic—*b. d'Œuf*, Albumen ovi—*b. de Perle*, Bismuth, subnitrate of—*b. de Plomb*, Plumbi subcarbonas—*b. de Zinc*, Zinci oxydum.

BLANC-MANGER (F.), *Cibus albus*, *Leucophagium*, *Leucophagum*, *Argyrotrophe* ma. An animal jelly, so called on account of its colour, combined with an emulsion of sweet almonds, to which sugar has been added, and some aromatic. It is sometimes prescribed as a nutriment in convalescence and chronic diseases.

BLANC-RAISIN, Blanc Rhazis.

BLANC RHAZIS, *Blanc-raisin*. An ointment composed of cerusa, white wax, and olive oil.

BLANCA, Plumbi subcarbonas.

BLANCH, from (F.) *blanchir*, 'to whiten, to bleach.' To whiten by depriving of the outer rind; as 'to bleach almonds;' i. e. to peel them.

BLANCHET, (F.) A blanket. A term given by the French Pharmacists to the woollen strainer through which they filter syrup and other thick fluids. See, also, Aphthæ.

BLANCHING, Etiolation—*b. of the hair*, Canities.

BLANCNON ORIBASII, Polypodium filix mas.

BLAS' An unmeaning term invented by Van Helmont to designate a kind of movement in the body; at times, local,—at others, under extraneous influence. Thus, he speaks of the *Blas meteoros* of the heavenly bodies, and the *Blas humanum*, that which operates in man.

BLAS ALTERNATIVUM, Plastic force.

BLASÉ, (F.) 'cloyed.' An epithet given to one whom the abuse of enjoyment has prevented from any longer deriving satisfaction or pleasure from it.

BLAST, *Contagio*, (Sax.) *blart*, (G.) *blasen*, 'to blow.' The infection of any thing pestilential. The stroke of some sudden plague.

BLASTE'MA, *Blaste'sis*, (F.) *Blastème*, from *blasarav*, 'I bud.' A germ. The sense of this word, which is often used by Hippocrates, is obscure. Castelli thinks it means the eruption of some morbid principle at the surface of the body. Also, the matrix or general formative element of tissues.

BLASTEMA PILI, see Hair.

BLASTEMAL, *Blastemal'is*. Relating or appertaining to a blastema,—as 'blastemal formations,' those that are formed from a blastema.

BLASTOCARDIA, see Molecule.

BLASTODERMA, see Molecule.

BLATTA BYZANTIA, *Unguis odoratus*, (F.) *Blatte de Byzance*. This name seems, formerly, to have been given to a marine production from some of the Conchylia. It had an agreeable smell, a reddish tint, and the shape of a nail. It was prescribed in epilepsy, hysteria, and hepatic obstructions. Rondellet affirms, that it was the

production of the shell-fish *murex* or *purpura*; and that the name *Blatta* is derived from the Greek *βλαττος*, 'purple.'

BLAVELLE, *Centaurea cyanus*.

BLAVÉOLE, *Centaurea cyanus*.

BLAVEROLLE, *Centaurea cyanus*.

BLAWING, Swelling.

BLAWORT, *Centaurea cyanus*.

BLAZING-STAR, *Chamaelirium luteum*, *Liatris*.

BLÉ, *Bladum*. This word answers, in France, to the word *Corn* in England; i. e. any kind of grain employed for making bread. Wheat being most commonly used for this purpose. *Blé* is sometimes restricted to this. *Blé mêlé* is a mixture of wheat and rye.

BLÉ 'CORNT', Ergot—*b. d'Espagne*, Zea mays—*b. d'Italie*, Zea mays—*b. Météil*, see *Blé*—*b. Noir*, Polygonum fagopyrum—*b. de Turquie*, Zea mays.

BLÉ (PARLER,) *Blæstias*.

BLEABERRY, *Vaccinium myrtillus*.

BLEACHING, Dealcation—*b. Liquid*, *Eau de javelle*.

BLEAREDNES, Lippitudo.

BLEAR-EYE, Lippitudo.

BLEAR-EYED, Lippus.

BLEAR-EYEDNESS, Lippitudo.

BLEB, Bulla.

BLECHNON, Polypodium filix mas.

BLECHNUM LIGNIFOLIUM, *Asplenium Scolopendrium*—*b. Squamosum*, *Asplenium ceterach*.

BLECHROPYRA, see Blechros.

BLECHROPYRUS, Typhus mitior.

BLECHROS, *βλεχρος*, 'weak, feeble, slow.' An epithet applied to different affections, and particularly to fevers. Hence *Blechropyra*, 'a slow fever.' *Blechrophyg'mia*, 'a slow pulse.'

BLECHROSPHYGMIA, see Blechros.

BLÉD, Corn—*b. Avorté*, Ergot—*b. Farouche*, Ergot.

BLEDOCH, Buttermilk.

BLEEDING, Bloodletting, Hæmorrhagia.

BLEEDING FROM THE NOSE, Epistaxis.

BLEEDING HEART, *Cypripedium luteum*—*b. Boist*, Cupping glass.

BLEIB, Bulla.

BLEINE, Blain.

BLÈME, (F.) This word has nearly the same signification as *Blafard*. Generally, however, it includes, also, emaciation of the countenance.

BLENCYSTORRHÉE, Cystirrhœa.

BLENNA, Mucus—*b. Narium*, Nasal mucus.

BLENNADENITIS, from *βλεννα*, 'mucus,' *αδην*, 'a gland,' and *itis*, denoting inflammation. Inflammation of mucous follicles.

BLENNELYT'RIA, from *βλεννα*, 'mucus,' and *ελυτρον*, 'a sheath.' A discharge of mucus from the vagina. Leucorrhœa.—Alibert.

BLENNEMESIS, *Βλεννοειδ'εσις*, *Vomitus pituitosus*, from *βλεννα*, 'mucus,' and *εμεσις*, 'vomiting.' Vomiting of mucus.

BLENNENTERIA, Dysentery.

BLENNISTH'MIA, from *βλεννα*, 'mucus,' and *ισθμος*, 'the gullet.' Increased flow of mucus from the pharynx and larynx.—Alibert.

BLENNOCHEZIA, Diarrhœa, mucosa.

BLENNOCYSTIDES, Bursæ mucosæ.

BLENNOCYSTITE, Cystirrhœa.

BLENNODES, Muciform.

BLENNOMESIS, Blennemesis.

BLENNOG'ENOUS, *Blennog'ennus*, *Mucif'ie*, *Mucif'icus*, from *βλεννα*, 'mucus,' and *γενναω*, 'I form.' Forming or generating mucus. Bireschet and Roussel de Vauzème describe an apparatus

of this kind for the secretion of the mucous matter that constitutes the cuticle, composed of a glandular parenchyma or organ of secretion situated in the substance of the true skin, and of excretory ducts, which issue from the organ, and deposite the mucous matter between the papillae.

BLENNOIDES, Muciform.

BLENNOIDEUS, Muciform.

BLENNOMETRIS, see Leucorrhœa.

BLENNOPHTHALMIA, Ophthalmia, (purulent.)

BLENNOPTYSIS, from *βλεννα*, and *πτωω*, 'I spit.' Expectoration of mucus. Catarrh.

BLENNOPYRA, *Blennopy'ria*, from *βλεννα*, and *πυρ*, 'fire.' Alibert has classed, under this head, various fevers with mucous complications; as *Mesenteric fever*, *Adeno-meningeal fever*, &c.

BLENNORRHAGIA, Gonorrhœa—b. Genitulum, Leucorrhœa—b. Notha, Gonorrhœa spuria—b. Spuria, Gonorrhœa spuria.

BLENNORRHAG'IC, *Blennorrhag'icus*, (F.) *Blennorrhagique*, from *βλεννα*, 'mucus,' and *ρρῑναι*, 'I break forth.' Relating or appertaining to blennorrhagia or gonorrhœa.

BLENNORRHAGIC EPIDIDYMITIS, Hernia humoralis.

BLENNORRHAGIE FAUSSE, Gonorrhœa spuria—b. *du Gland*, Gonorrhœa spuria.

BLENNORRHÉE, Blennorrhœa.

BLENNORRHINIA, Coryza.

BLENNORRHË'A, *Blennor'rhœ*, *Blennorrhag'ia*, *Phlegmorrhœ'a*, *Phlegmorrag'ia*, (F.) *Blennorrhée*, from *βλεννα*, 'mucus,' and *ρεω*, 'I flow.' Inordinate secretion and discharge of mucus. Also, Gonorrhœa.

BLENNORRHËA CHRONICA, (gleet,) see Gonorrhœa—b. Genitulum, Leucorrhœa—b. Luodes, Gonorrhœa impura—b. Nasalis, Coryza—b. Oculi, see Ophthalmia—b. Oculi gonorrhœa, see Ophthalmia—b. Oculi neonatorum, see Ophthalmia—b. Oculi purulenta, see Ophthalmia—b. Urethralis, Gonorrhœa, Cystorrhœa—b. Ventriculi, Gastrorrhœa—b. Vesicæ, Cystorrhœa.

BLENNORSES, from *βλεννα*, 'mucus.' Affections of the mucous membranes.—Alibert.

BLENNOTHORAX, Catarrh, Peripneumonia notha—b. Chronicus, Asthma humidum.

BLENNOTORRHËA, Otirrhœa.

BLENNURETHRIA, Gonorrhœa.

BLENNURIA, Cystorrhœa.

BLEPHARADENITIS, Ophthalmia Tarsi.

BLEPHARANTHRACOS'IS, *Bléphari'tis gangræno'sa*, *Carbuncula'tio Oculi*. Gangrenous inflammation of the eyelids.

BLEPHARELOSIS, Entropion.

BLEPHARIDES, Cilia.

BLEPHARIDOPLASTICE, Blepharoplastice.

BLEPHARISMUS, Nictation.

BLEPHARITIS, (inflammation of the eyelids,) Ophthalmia tarsi—b. Gangrænosa, Blepharanthracosis.

BLEPHAROBLENNORRHËA, Ophthalmia, purulent—b. Neonatorum, see Ophthalmia (purulenta infantum.)

BLEPH'ARO-CONJUNCTIVITIS, *Blepharconjunctivitis*, from *βλεφαρον*, 'an eyelid,' *conjunctiva*, and *itis*, denoting inflammation. Ophthalmia affecting the conjunctiva and eyelids.

BLEPHARODYSCHRË'A, from *βλεφαρον*, the 'eyelid,' *δυσ*, 'with difficulty,' and *χρoα*, 'colour.' Discoloration of the eyelid. Nævus of the eyelid.—Von Ammon.

BLEPHARCEDEMA AQUOSUM, Hydroblepharon.

BLEPHARON, Palpebra—b. Atoniaton, Blepharoptosis.

BLEPHARONCO'SIS, *Blepharon'cus*, *Blepharophy'ma*, *Palpebra'rum Tumor*, from *βλεφα-*

ρον, 'eyelid,' and *oγκος*, 'tumour.' A tumour of the eyelid.

BLEPHARONCUS, Blepharonecosis.

BLEPHAROPHIMO'SIS, from *βλεφαρον*, 'the eyelid,' and *φίμος*, 'a cord.' Congenital diminution of the space between the eyelids.—Ammon. See Blepharostenosis.

BLEPHAROPHTALMIA, Ophthalmia tarsi—b. Neonatorum, see Ophthalmia—b. Purulenta, Blepharopyorrhœa.

BLEPHAROPHTHALMITIS GLANDULOSA, Ophthalmia, purulent, of infants.

BLEPHAROPHYMA, Blepharonecosis.

BLEPHAROPLAS'TICE, *Blepharidoplas'ticê*, *Insi'tio Cilio'rum*, *Bleph'aroplasty*, from *βλεφαρον*, 'the eyelid,' and *πλαστικός*, 'forming,' 'formative.' The formation of a new eyelid.

BLEPHAROPLEGIA, Blepharoptosis.

BLEPHAROPTO'SIS, *Blepharople'gia*, *Casus pal'pebræ superioris*, *Delap'us* seu *Prolap'sus* seu *Proptosis* seu *Pto'sis pal'pebræ*, *Atoniaton blepharon*, from *βλεφαρον*, 'the eyelid,' and *πτωσις*, 'fall.' A falling down of the upper eyelid over the eye, caused by a paralysis of the *Levator palpebræ superioris* muscle. This paralysis is an unfavorable symptom, as it is generally connected with a state of the brain favouring apoplexy or palsy.

BLEPHAROPTOSIS ECTROPIUM, Ectropium—b. Entropion, Entropion.

BLEPHAROPYORRHË'A, *Blepharophtal'mia purulen'ta*, *Pyorrhœ'a pal'pebræ*, from *βλεφαρον*, 'eyelid,' *πυρ*, 'pus,' and *ρεω*, 'I flow.' Secretion of pus from the eyelids.

BLEPHAROPYORRHËA NEONATORUM, see Ophthalmia (purulenta infantum.)

BLEPHARORRHË'A, from *βλεφαρον*, 'eyelid,' and *ρεω*, 'I flow.' A discharge of mucus from the eyelids.

BLEPHAROSPAS'MUS, from *βλεφαρον*, 'eyelid,' and *σπασμος*, 'spasm.' A spasmodic action or tic of the orbicularis palpebrarum muscle.

BLEPHAROSTENO'SIS, from *βλεφαρον*, 'the eyelid,' and *στενωσις*, 'contraction.' Accidental diminution of the space between the eyelids.—Ammon. See Blepharophimosi.

BLEPHAROSYNDESMITIS, Blepharoconjunctivitis.

BLEPHAROTIS, Ophthalmia tarsi—b. Glandularis contagiosa, see Ophthalmia.

BLEPHAROTITIS, Ophthalmia tarsi.

BLEPHAROTOSIS, Ectropium.

BLEPHAROXYS'TUM, *Blepharoxys'trum*, from *βλεφαρον*, 'eyelid,' and *ξύω*, 'I scrape.' An instrument used, by the ancients, for removing callosities, which made their appearance in the affection called, by the Greeks, *τραχώμα*.—Paulus of Ægina, Goræus.

BLEPHIL'IA HIRSU'TA, *Ohio Horsemint*, *Hairy Horsemint*, from *βλεφαρίς*, 'an eyelash,' in reference to its hairy fringed bracts. An indigenous plant of the Mint family, Labiateæ, which has the aromatic properties of the Mints.

BLËSITE, Blasitas.

BLESSED HERB, Geum urbanum.

BLESSURE, Abortion, Wound.

BLESTRIS'MUS. Restlessness of the sick.—Hippocrates.

BLETA. A word, used by Paracelsus for white or milky urine, arising from diseased kidneys. *Biota alba* has the same meaning.

BLEU DE BERLIN, Prussian blue—b. *de Prusse*, Prussian blue.

BLEUET DES MOISSONS, Cyanus segetum.

BLÉVILLE, MINERAL WATERS OF. Bléville is a village about two miles from Havrê. The waters are acidulous chalybeate.

BLIGHT IN THE EYE, Ophthalmia, catar-

chal. Palsy of the orbicularis palpebrarum and muscles of the eyebrow is also vulgarly called *blight*.

BLIGHTS, see *Lichen urticatus*.

BLINDNESS, *Cæcitas*—b. Colour, *Achromatopsia*—b. Day, *Nyctalopia*—b. Nervous, *Amaurosis*—b. Night, *Hemeralopia*—b. Snow, see *Snow blindness*.

BLISTER, *Vesicatorium*, *Emplastrum Vesicatorium* seu *Lytta*, *Epispasticum*, *Vesicant*, *Blister plaster*, (F.) *Vésicatoire*, *Vésicant*. Any substance which, when applied to the skin, irritates it and occasions a serous secretion, raising the epidermis, and inducing a vesicle. Various articles produce this effect, as *cantharides*, *mustard*, *garon*, *euphorbium*, *garlic*, *annamonia*, &c. Blisters are used as counter-irritants. By exciting a disease artificially on the surface, we can often remove another which may be at the time existing internally. A *perpetual blister* is one that is kept open for a longer or a shorter time by means of appropriate dressings.

BLISTER of vesication, (Prov.) *Ercle*, also means the *vesicle* produced by vesicatories or other causes.

BLISTER, FEVER, see *Fever blister*.

BLISTER, MAGISTRAL, (F.) *Vésicatoire magistral*. A prompt means of producing vesication recommended by M. Valleiæ. It is prepared as follows—Take powdered *cantharides* and *wheat-flour*, of each equal parts; *vinegar*, a sufficient quantity to form a soft paste.

BLISTER BEETLE, *Cantharis*—b. Fly, *Cantharis*—b. Plaster, *Blister*.

BLISTERWEED, *Ranunculus acris*.

BLISTERING FLY, *Cantharis*—b. Paper, see *Sparadrapum vesicatorium*—b. Tissue, *Sparadrapum vesicatorium*.

BLISTERS, WHITE, *Rupia escharotica*.

BLITUM AMERICANUM, *Phytolacca decandra*.

BLOOD, Anglo-Saxon, *bloo*, from *blevan*, 'to bleed.' *Sanguis*, *Cruor*, *Lapis animalis*, *Hæma*, *æma*, (Old Eng.) *Blede*, (F.) *Sang*. An animal fluid formed chiefly from the chyle; acquiring important properties during respiration; entering every organ through the circulation; distributing the nutritive principles to every texture, and the source of every secretion. The blood is white in the molluscous and inferior animals, which have been, hence, called *white-blooded*, to distinguish them from the *red-blooded*, which class includes the mammalia, birds, reptiles, and fishes. Human blood is composed of water, albumen, fibrin, an animal colouring substance, a little fatty matter—*hæmateleum*—and different salts; as chlorides of potassium and sodium, phosphate of lime, subcarbonate of soda, lime, magnesia, oxide of iron, and lactate of soda, united with an animal matter. *Arterial blood* is of a florid red colour, strong smell; s. g. 1.049. *Venous blood* is of a brownish red; s. g. 1.051. The difference in colour has given occasion to the first being called *red blood*; the latter, *black*. The former, which is distributed from the heart, is nearly the same through its whole extent: the latter is the remains of the arterial blood after the different elements have been taken from it in nutrition, and probably differs in composition. It likewise contains different substances absorbed. Venous blood, taken from a vessel and left to itself, becomes solid, and separates into two distinct parts,—the *serum* or watery supernatant fluid; and the *cruor*, *coagulum*, *crassamentum*, *hepar* seu *placenta sanguinis*, *placenta cruoris*, *insula*, *thrombus*, or clot. The serum is chiefly water, holding albumen in solution and the salts of the blood. The clot contains the fibrin, colouring matter—*hæmatoxin*, a little serum, and a

small quantity of salts. M. Le Canu found the blood to be composed—in 1000 parts—of water, 785.590; albumen, 69.415; fibrin, 3.565; colouring matter, 119.626; crystallizable fatty matter, 4.300; oily matter, 2.270; extractive matter soluble in alcohol and water, 1.920; albumen combined with soda, 2.010; chlorides of sodium and potassium; alkaline phosphates, sulphates, and subcarbonates, 7.304; subcarbonate of lime and magnesia, phosphate of lime, magnesia and iron, peroxide of iron, 1.414; loss, 2.586. The four principal components of the blood are fibrin, albumen, corpuscles, and saline matter. In the circulating blood they are thus combined—

Fibrin, } In solution forming *Liquor Sanguinis*.
Albumen, }
Salts, }

Red Corpuscles—suspended in the *Liquor Sanguinis*.

In *coagulated blood* they are thus combined:

Fibrin, } Forming the *crassamentum* or
Red Corpuscles, } clot.
Albumen, } Remaining in solution, forming
Salts, } serum.

The following table exhibits the computations of different physiologists regarding the weight of the circulating fluid—arterial and venous.

Harvey,	1lb.
Lister,	8
Moulin,	8
Abildgaard,	10
Blumenbach,	10
Lobb,	10
Lower,	10
Sprengel,	10 to 15
Günther	15 to 20
Blake	164 to 194
Müller and Burdach	20
Wagner	20
Quenai	20 to 25
F. Hoffmann	23
Haller	28 to 30
Young	40
Hamberger	80
Keill	100

The proportion of arterial blood to venous is about 4 to 9.

Much attention has been paid to the varying condition of the blood in disease. The average proportion of each of the organic elements in 1000 parts of healthy blood is as follows, according to Le Canu, and MM. Andral and Gavarret:—fibrin, 3; red corpuscles, 127; solid matter of the serum, 80; water, 790.

Dried human blood was, at one time, considered to be anti-epileptic; that of the goat, dried, *Sanguis hirci siccat*, sudorific and antileptic. The dried blood of the ox—*Extractum Sanguinis Bovini*—and the dried red corpuscles have been given as analeptics, especially where there was a deficiency of red corpuscles.

BLOOD, ARTERIAL, see *Blood*—b. Black, see *Blood*—b. Black, vascular system of, see *Vascular*.

BLOOD, CASEIN, Globulin—b. Corpuscles, Globules of the blood—b. Disease, *Hæmatonosis*—b. Disks, Globules of the blood.

BLOOD, DRIED, see *Blood*—b. Loss of, *Hæmorrhagia*—b. Menstrual, see *Menses*—b. Red, see *Blood*—b. Red, system of, see *Vascular*—b. Spitting of, *Hæmoptysis*—b. Venous, see *Blood*—b. Vomiting of, *Hæmatemesia*—b. White, *Lymph*.

BLOOD CRYSTALS, *Hæmatoidin*.

BLOODING, *Blood-letting*.

BLOODLESSNESS, *Anæmia*.

BLOOD-LETTING, *Missio* seu *Detractio Sanguinis*, *Hæmat*, *Catascasmus*, *Bleeding*,

Bleeding, (F.) Saignée, Emission sanguine. A discharge of a certain quantity of blood produced by art: an operation which consists in making an opening into a vessel to draw or let blood from it. When practised on an artery, it is called *Arteriotomy*; on a vein, *Phlebotomy*, *Venæsectio*, *Venesection*; and on the capillary vessels, *local* or *capillary*, in contradistinction to the former, which is termed *general*. Blood-letting is used both during the existence of a disease, as in inflammation, and in the way of prophylaxis. It is employed to fulfil various indications. 1. To diminish the actual mass of blood;—when it is termed, by the French pathologists, *Saignée évacuative*. In such case, fluids ought not to be allowed too freely afterwards. 2. To diminish the turgescence in any particular organ—(F.) *Saignée révulsive*, *Revulsive blood-letting* or *bleeding*, *Venæsectio revulsoria*, when performed far from the part affected; and *Saignée dérivative*, when near. 3. To diminish the consistence of the blood, (F.) *Saignée spoliative*. The immediate effects of blood-letting are: diminution of the mass of blood and of heat; retardation of the pulse, and sometimes syncope. Blood-letting from the veins—*phlebotomy*—is practised on the subcutaneous veins of the neck, the face, the fore-arm, and the leg; sometimes on those of the hand or foot. The necessary apparatus consists of a bandage or riband, a compress of rag, and a lancet or fleam.

The veins selected for the operation, are, 1. *In the fold of the arm*, five;—the cephalic, basilic, the two median, and the anterior cubital. 2. *In the hand*, the cephalic and salvatella. 3. *In the foot*, the great and little saphena. 4. *In the neck*, the external jugular. 5. *In the forehead*, the frontal. 6. *In the mouth*, the ranine. The operation of phlebotomy in the limbs is performed by tying a circular bandage round the limb, in order that the subcutaneous veins may become turgid by the course of the blood being obstructed: the bandage not being so tight, however, as to compress the arteries of the limb. A puncture is made into the vein, and the desired quantity allowed to flow. The ligature is now removed, and a compress and retaining bandage applied. *Capillary or local blood-letting*, *arterio-phlebotomy*, is practised on the skin or mucuous membranes, by means of leeches, the lancet, or cupping.

BLOOD-LETTING, CAPILLARY, see Blood-letting—b. Derivative, see Blood-letting—b. Evacuative, see Blood-letting—b. General, see Blood-letting—b. Local, see Blood-letting—b. Revulsive, see Blood-letting—b. Spoliative, see Bloodletting.

BLOODLIKE, Sanguine.

BLOOD LIQUOR, Liquor Sanguinis.

BLOODROOT, Sanguinaria Canadensis.

BLOODSHOT, Hyperæmia.

BLOODSTONE, Hematites.

BLOODSTROKE, Coup-de-sang.

BLOOD VESICLE, Globule of the blood.

BLOOD VESSEL, (F.) *Vaisseau sanguin*. A vessel destined to contain and convey blood.

BLOOD VESSEL, BREAKING, BURSTING, RUPTURE OF A. Hemorrhagia.

BLOODWEED, Asclepias curassavica.

BLOODWORT, Hieracium venosum, Sanguinaria Canadensis.

BLOODY, *Sanguin'eus*, *Cruent'us*, *Sanguin'eus*, (F.) *Sanguin*. Having the character of blood. Relating to blood. See Sanguine.

BLOODY FLUX, Dysentery.

BLOOM, HONEY, Apocynum androsæmifolium.

BLOSSBURG MINERAL SPRINGS. The town of Blossburg is in Tioga County, 133 miles

north of Harrisburg. They contain pure sulphuric acid; and are astringent and tonic.

BLOSSOM, see Grog-blossom.

BLOTA ALBA, Bleta.

BLOTCH, (Prov.) *Splavin*, *Sprote*. A pustule; more commonly, an eruption of a large kind.

BLOW, *Ictus*, *Plegè*, (F.) *Coup*. Effect produced by one body striking another. The impression made by any body which strikes us, or against which we strike;—a common cause of wounds, contusions, fractures, &c.

BLOWING SOUND, *Bruit de Souffle*—b. s. Diffused, see Murmur, respiratory.

BLUE-BELLS, Gentiana catesbæi.

BLUE-BERRY, Caulophyllum thalictroides, Lantana—b. Low, Vaccinium Pennsylvanicum.

BLUE-BONNETS, Cetaurea cyanus.

BLUE-BOTTLE, Cetaurea cyanus, Cyanus segetum.

BLUE LICKS, see Kentucky, Mineral Waters of.

BLUE STONE, Cupri sulphas.

BLUET, Cetaurea cyanus.

BLUET DES MOISSONS, Cyanus segetum.

BLUTER, Borborygmus.

BLUSH, see Flush.

BLUSH, CUTANEOUS, see Efflorescence—b. Inflammatory, Erythema.

BLUTER, Borborygmus.

BOA, *Boia*. An eruption of red, ichorous pimples.—Pliny. See, also, Hidroa and Sudamina.

BOA UPAS, Upas.

BOÆ, Syphilis.

BOALA, see Scherliev.

BOBERRI, Curcuma longa.

BOCHE, Swelling.

BOCCHOE, Diosma crenata.

BOCHET, Bochetum.

BOCHETUM, (F.) *Bochet*. A term formerly applied to the second decoction of the sudorific woods. The French word has been lately revived.

BOCIUM, Bronchocele.

BOCHO, Diosma crenata.

BOCIUM, Bronchocele.

BOCK, Vomiturition.

BOCKING, Eructation, Vomiting.

BOCKLET, MINERAL WATERS OF. The springs of Bocklet, in Bavaria, are acidulous chalybeates.

BODY, *Corpus*, *Soma*, (F.) *Corps*. From Teutonic *boden*, the 'fundus or bottom.' [?] The human body, (F.) *Corps humain*, (Old E.) *Liche*, is the collection of organs which compose the frame. At times, however, body is used synonymously with *trunk*. We say, also, *body of the femur*, of the *sphenoid*, &c., to designate the shaft or middle portion of those bones; *body of the uterus*, &c. Likewise, any distinct portion of matter perceived by the senses. Also, the rectum.

BODY, COMING DOWN OF THE, Proctoele.

BODY-SEARCHER, *Searcher*. One who formerly examined the bodies of the dead to report as to the cause of death. In the time of the plague this was done by "the chirurgeons," who were allowed "twelve pence," to be paid out of the goods of the party searched, if able; otherwise, by the parish. Until of late, this office was executed in England by two old women—called 'searchers'; and hence the imperfections of the Bills of Mortality. In plague times, the office was an important one; and it is stated, that a noted searcher, named Snacks, finding his business increase so much, that he could not transact it alone, offered to any one who would join him in its hazards, half the profits; and they who joined him were said to "go with Snacks." Hence, the saying "to go Snacks," or to divide the spoils.—Wadd.

BODY-SNATCHER, Resurrectionist.

BOE, Cry.

BOELLI, Intestines.

BOETHEMA, Medicament.

BOG-BEAN, *Menyanthes trifoliata*.

BOHON UPAS, Upas.

BOIA, Boa.

BOIL, *Furunculus*—*b. Blind*, see *Furunculus*—*b. Gun*, *Parulis*—*b. Malignant*, see *Furunculus*—*b. Wasp's nest*, see *Furunculus*.

BOIS AMER, *Quassia*—*b. de Campêche*, *Hæmatoxylum Campechianum*—*b. de Cerf*, *Cornu cervi*—*b. de Chypre*, *Rhodium lignum*—*b. de Couleuvre*, see *Strychnos*—*b. Doux*, *Glycyrrhiza*—*b. de Marais*, *Cephalanthus occidentalis*—*b. de Plomb*, *Dorca palustris*—*b. Puant*, *Prunus padus*—*b. de Rose*, *Rhodium lignum*—*b. de Sappan*, *Cæsalpinia sappan*—*b. Sudorifique*, *Wood*, sudorific.

BOISSE, MINERAL WATERS OF. These waters are situate about half a league from Fontenay-le-Compte, in France. They are purgative, and seem to contain carbonate and sulphate of lime and chloride of sodium.

BOISSON, Drink.

BOIST, Swelling.

BOÎTE, (F.) A box or case, *Capsa*, *Pyxis*. An apparatus for the reception of any matters which it may be desirable to preserve. In *Surgery* and *Anatomy*, *Boîtes à dissection*, *B. à amputation*, *B. à trépan*, *B. à cataracte*, &c., mean the cases containing these various instruments. *Boîte du Crane* is the bony case which receives the brain. *Boîte* is, also, the portion of the stem of the trephine which receives the pyramid or centre-pin. *Boîte de Petit* is a machine, invented by M. Petit, to retain the fractured portions of bone in apposition, when the leg has been fractured in a complicated manner. *Boîte* is, also, a kind of case put before an artificial anus to receive the fæces, which are continually being discharged. The vulgar, in France, give the name *Boîte*, to various articulations,—*B. de genou*, *B. de la hanche*; "knee-joint, hip-joint."

BOITEMENT, Claudication.

BOITERIE, Claudication.

BOÏTER, (F.) *Appareil*, *Cap'sula unguentaria*, *Capad'rium*. A dressing-case. A box, containing salves and different apparatus, used more particularly by the dressers in hospitals.

BOK, Vomiturition.

BOKKING, Eructation, Vomiting.

BOL, *Bolus*—*b. d'Arménie*, *Bole*, *Armenian*—*b. Oriental*, *Bole*, *Armenian*—*b. Blanc*, *Bolus alba*—*b. Rouge*, *Bole*, *Armenian*.

BOLA, Myrrha.

BOLCHON, *Bdellium*.

BOLE, *Bolus*, (F.) *Bol*, *Terre bolaire*, meant, with the older writers, argillaceous earth, used as an absorbent and alexipharmic. The various boles had different forms given to them, and were stamped, or sealed, hence called *Terræ sigillatæ*; as in the following:

BOLE ARME'NIAN, *Bole Arme'niac*, *B. Ar'menic*, *Argilla ferrugin'ea rubra*, *A. Bolus rubra*, *Sinapis*, *Arena'men*, *Bolus Orientalis* seu *Armeniaca* seu *Arme'niæ* seu *rubra*, (F.) *Bol d'Arménie* ou *oriental* ou *rouge*. A red, clayey earth, found not only in Armenia, but in several countries of Europe,—in Tuscany, Silesia, France, &c. It was once esteemed a tonic and astringent, and was applied as a styptic. It is now, scarcely, if ever, used. It consists of argil, mixed with lime and iron.

BOLESIS, Coral.

BOLESON, Balsam.

BOLET ODORANT, *Dædalea suaveolens*.

BOLETUS AGARICUS, *B. Laricis*—*b. Albus*, *Boletus laricis*—*b. Cervinus*, *Elaphomyces granulatus*—*b. Discoidens*, *Dædalea suaveolens*.

BOLETUS ESCULENTUS, (F.) *Morelle*. *Ord.* Fungi. An eatable mushroom, found in the woods in Europe, and much admired by *Gastronomes*. It was formerly esteemed to be aphrodisiac.

BOLETUS FULVUS, *B. ignarius*—*b. Hippocrepis*, *B. ignarius*.

BOLETUS IGNARIUS. The systematic name for the *Agaric*, *Agar'icus*, *Agar'icus* seu *Pharmacopœias*, *Agar'icus Chirurgo'rum* seu *Quercus seu ignarius*, *Polyporus ignarius*, *Is'ca*, *Boletus unguilatus seu fulvus* seu *hippocrepis* seu *obtus*, *Spunk*, *Am'adou*, *Punk*, (Prov.) *Funk*, *Fungus Ignarius*, *Fungus Quercinus*, *Agaric of the Oak*, *Touchwood*, *Touchwood Boletus*, *Female Agaric*, *Tinder*, (F.) *Agaric de chêne*, *Amadouvier*. It was formerly much used by surgeons as a styptic.

BOLETUS LARICIS, *B. Laric'icus* seu *purgans* seu *albus* seu *agar'icus* seu *officina'lis*, *Fungus Laric'is*, *Polyporus officina'lis*, *Agar'icus albus* seu *Laric'is*, *A. Albus optimus*, *White Agaric*, (F.) *Agaric blanc*, *A. Amadouvier*. On the continent of Europe it has been given as a cathartic and emetic, as well as to moderate the sweats in phthisis.—*De Haen*. Externally, styptic.

BOLETUS OBTVS, *B. ignarius*—*b. Officinalis*, *B. laricis*—*b. Purgans*, *Boletus laricis*—*b. Salicis*, *Dædalea suaveolens*—*b. Suaveolens*, *Dædalea suaveolens*—*b. Touchwood*, *Boletus ignarius*.

BOLI MARTIS, *Ferrum tartarisatum*.

BOLISMOS, *Boulimia*.

BOLI'TES. The mushroom; perhaps the *Agar'icus Aurantiacus*.—*Pliny*, *Martial*, *Seutonius*, *Galen*. It was so called, in consequence of its shape,—from *Bolus*.

BOLLYNGE, Swelling.

BOLNING, Swelling.

BOLT. Used, at times, for to swallow without chewing,—as to "bolt one's food."

BOLUS, *βωλος*, *Buccella*, a morsel, a mouthful, a bole, (F.) *Bol*. A pharmaceutical preparation, having a pilular shape, but larger; capable, however, of being swallowed as a pill.

BOLUS ALBA, *Terra Sigilla'ta*, *Argilla pallidior*—called *sigilla'ta*, from being commonly made into small cakes or flat masses, and stamped or sealed with certain impressions. (F.) *Bol blanc*, *Terre Sigillée*, *Argile ochreuse pâle*. It was used like *Bole Armenian*, and was brought from Etruria. See *Terra*.

BOLUS, ALIMEN'TARY, *Bolus Alimentarius*. The bole formed by the food, after it has undergone mastication and insalivation in the mouth; and been collected upon the tongue prior to deglutition.

BOLUS ORIENTA'LIS. A kind of bolar earth, only distinguished from *Bole Armenian* in being brought from Constantinople. See *Bole*, *Armenian*.

BOLUS RUBRA, *Bole Armenian*.

BOMA'REA SALSIL'LA. The inhabitants of Chili use this plant as a sudorific. It is given in infusion in cutaneous diseases.

BOMBAX, *Gossypium*.

BOMBEMENT, *Bombus*.

BOMBITATIO, *Tinnitus aurium*.

BOMBUS, *Aurium fluctuatio*, *A. Sib'itus*, *A. Son'itus*, *A. Suen'rus*, (F.) *Bombement*. A kind of ringing or buzzing in the ears;—characterized, according to SAUVAGES, by the perception of blows or beating repeated at certain intervals. Also, *Borborygmus*. See *Fart*, *Flatulence*, and *Tinnitus Aurium*.

BOMBXY MORI, see *Sericum*.

BON, *Coffea Arabica*.

BONA, *Phaseolus vulgaris*.

BONANNTA OFFICINALIS, *Sinapis alba*.

BONA FEVER, see *Fever*, *Bona*.

BONDUE, *Gymnocladus Canadensis*.

BONE, Os, Os'teon, Os'teum, (F.) Os, (Sax.) ban. The bones are the solid and hard parts, which form the basis of the bodies of animals of the superior classes; and the union of which constitutes the *skeleton*. The human body has, at the adult age, 208 bones, without including the 32 teeth, the ossa Wormiana, and the sesamoid bones. Anatomists divide them, from their shape, into 1. *Long bones*, which form part of the limbs, and represent columns for supporting the weight of the body, or levers of different kinds for the muscles to act upon. 2. *Flat bones*, which form the parietes of splanchnic cavities; and, 3. *Short bones*, met with in parts of the body where solidity and some mobility are necessary. Bones are formed of two different textures; *spongy* and *compact*. They afford, on analysis, much phosphate and carbonate of lime, a little phosphate of magnesia, phosphate of ammonia, oxides of iron and manganese, some traces of alumina and silica, gelatin, fat, and water. The uses of the bones are mentioned under each bone. They give shape to the body, contain and defend the viscera, and act as levers to the muscles.

TABLE OF THE BONES.

BONES OF THE HEAD.	Bones of the Cranium or Skull.	Frontal.....	1
		Parietal.....	2
		Occipital.....	1
		Temporal.....	2
		Ethmoid.....	1
		Sphenoid.....	1
	Bones of the Face.	Superior Maxillary.....	2
		Inferior Maxillary.....	2
		Jugal or Cheek.....	2
		Nasal.....	2
		Lachrymal.....	2
		Palatine.....	2
	Dentes, or Teeth.	Inferior Spongy.....	2
		Vomer.....	1
		Inferior Maxillary.....	1
		Incisores.....	8
		Cuspidati.....	4
		Molares.....	20
BONES OF THE TRUNK.	Bone of the Tongue.	Hyoïd.....	1
		Malleus.....	2
	Bones of the Ear.	Incus.....	2
		Orbicular.....	2
		Stapes.....	2
		Cervical.....	7
	Vertebrae.	Dorsal.....	12
		Lumbar.....	5
	Sacrum.....		1
	Os Coccygis.....		1
	The Thorax.	Sternum.....	1
		Ribs.....	24
	The Pelvis.	Innominate.....	2
		Clavicle.....	2
BONES OF THE UPPER EXTREMITY.	The Shoulder.	Scapula.....	2
		Humerus.....	2
	The Arm.	Ulna.....	2
		Radius.....	2
	Fore-arm.	Navicular.....	2
		Lunare.....	2
	The Hand.	Cuneiforme.....	2
		Orbicular.....	2
	Carpus, or Wrist.	Trapezium.....	2
		Trapezoides.....	2
	Metacarpus.	Magnus.....	2
		Unciforme.....	2
BONES OF THE LOWER EXTREMITY.	The Thigh.	Phalanges.....	10
		Femur.....	28
	The Leg.	Tibia.....	2
		Fibula.....	2
	Tarsus, or Instep.	Calcis Os.....	2
		Astragalus.....	2
	The Foot.	Cuboides.....	2
		Navicular.....	2
	Metatarsus.	Cuneiforme.....	10
		Phalanges.....	10
	Total.....		240

The bones of the ox, *Bos Taurus*, are employed in pharmacy.

BONE-ACH, Osteocopus — b. Ague, Osteocopus.

BONE, BACK, Vertebral column — b. Bar, Pubis, os. **BONE BLACK**, see Carbo animalis.

BONE, BLADE, Scapula — b. Bont-like, Os scaphoides — b. Breast, Sternum — b. Crupper, Coccyx.

BONE EARTH, see Cornu ustum — b. Fever, see Inflammation.

BONE, HAUNCH, Ilion — b. Interparietal, Interparietal bone.

BONE PHOSPHATE OF LIME, see Cornu ustum.

BONE, RUMP, Coccyx — b. Share, Pubis — b. Splinter, Fibula.

BONEBINDER, Osteocolia.

BONE-DOCTOR, Renoueur.

BONE-NIPPERS, Osteul'cum, Tenac'ula, from *teneo*, 'I hold.' (F.) *Tenaille incisive*. An instrument used for cutting off splinters and cartilages. It is a kind of forceps, the handles of which are strong, and the edges, which touch each other, cutting.

BONEST, Eupatorium perfoliatum — b. Rough, Eupatorium teucrifolium — b. Upland, Eupatorium sessifolium.

BONE-SETTER, Renoueur.

BONE-SHAVE, Neuralgia femoro-poplitea.

BONES, BRITTLINESS OF THE, Fragilitas ossium — b. Friability of the, Fragilitas ossium — b. Salt of, Ammoniac carbonas — b. Softening of the, Mollities ossium.

BONHOMME, Verbascum thapsus.

BONICLATTER, Bonnyclabber.

BONIFACIA, Ruscus hypoglossum.

BONITHLOBBER, Bonnyclabber.

BONNE DAME, Atriplex hortensis.

BONNES, MINERAL WATERS OF. Bonnes is a village six leagues from Pau, in the department Basses Pyrénées, France. Here are several thermal springs. They were celebrated as early as the time of Francis I., under the name *Eau d'Arquebusade*. They contain chlorides of sodium and magnesium, sulphates of magnesia and lime, sulphur, and silica. The temperature is from 78° to 90° Fahrenheit.

The *fictitious* EAU DE BONNES is made of Hydrosulphuretted water, f $\frac{3}{4}$ iv; pure water, Oj. and f $\frac{3}{4}$ ss; chloride of sodium, gr. xxx; sulphate of magnesia, gr. i.

BONNET, Reticulum.

BONNET À DEUX GLOBES, Bonnet d'Hippocrate.

BONNET D'HIPPOCRATE, Cap of Hippocrates, Mitra Hippocrat'ica, Fascia capit'alis, Pi'leus Hippocrat'icus. A kind of bandage, the invention of which is ascribed to Hippocrates. It consists of a double-headed roller, passed over the head, so as to envelop it like a cap. The French also name it *Bonnet à deux globes, Capeline de la tête*.

BONNYCLABBER, Clabber, from Irish, *baine*, 'milk,' and *clabar*, 'mire.' In Ireland, sour buttermilk. In this country, the thick part of sour milk. *Boniclatter* and *Bonithlobber*, also, meant cream or "good milk gone thick." — Halliwell.

BONPLANDIA ANGUSTURA, Cusparia febrifuga — b. Trifoliata, Cusparia febrifuga.

BONTA GERMINANS, Avicennia tomentosa.

BONUS GENIUS, Peucedanum — b. Henricus, Chenopodium bonus Henricus.

BONY, Osseous.

BOOCHO, Diosma crenata.

BOON UPAS, Upas.

BOONA, Phaseolus vulgaris.

BOOTIA VULGARIS, Saponaria.

BOOTIKIN. A glove with a partition for the thumb, but no separate ones for the fingers — like an infant's glove — made of oiled silk. — Dr. E. J. Seymour. Horace Walpole speaks in raptures of the benefit he derived from bootikins in gout.

BORAC'IC ACID, Ac'idum Borac'icum, Sal sedati'vus HOMBER'GI, Boric Acid, (F.) Acide

boracique. An acid obtained from borax, which was once looked upon as sedative. It was also called *Acor Borac'icus*, *Sal vitrioli narcot'icum*, *Sal volat'ile Borac'is*, and *Flores Borac'is*.

BORAGE, *Borago officinalis*.

BORAGO OFFICINALIS, *Buglossum verum* seu *latifolium*, *Borra'go*, *Corra'go*, *Borago hortens'is*, *Borago*, *Burrago*, (F.) *Bourrache*. Ord. Boraginæ. Sex. Syst. Pentandria Monogynia. The leaves and flowers have been considered aperient.

BORAS SUPERSODICUS, Borax.

BORATHRON, *Juniperus Sabina*.

BORAX, *Borax Soda* seu *supersaturus soda* seu *alcales'cens* seu *aleali'num* seu *supersod'icus*, *Soda Bibo'ras*, *Subboras Soda* seu *nat'ricum*, *Soda Borax'ia*, *Chrysocol'ia*, *Capi'strum auri*, *Subborate* of *protoz'ide* of *Sodium*, *Subprotoborate* of *Sodium*, *Borax Ven'etus*, *Borax'trion*, *Nitrum facti'tium*, &c., *Subbo'rate* or *Biborate* of *Soda*, *Borate* of *Soda*, (F.) *Borate* ou *Sous-borate* de *Soude*, *Borate saturé* de *soude*. It is found in an impure state in Thibet and Persia. It is inodorous; taste cool, and somewhat alkaline; soluble in 12 parts of water. Borax is seldom used except as a lotion in aphthæ.

BORATE OF MERCURY has been recommended as an antisyphilitic.

BORAXTRION, Borax.

BORBONIA RUSCIFO'LIA. A small South African shrub, used in asthma and hydrothorax. In decoction, it is given as a diuretic.—Pappe.

BORBORUS, *Fimus*.

BORBORYG'MUS, from βορβορῆς, 'I make a dull noise.' *Murmur* seu *Bombus* seu *Motus Intestini'rum*, *Anile'ma*, *Anile'sis*, *Oeliosoph'ia*, *Intona'tio intestinal'is*, *Murmur ventris* seu *intestinal'is*, *Bor'borygm*, (Sc.) *Bluter*, *Blutter*, (Prov.) *Crolling*, *Crouling*, (F.) *Gargouillement*, *Grouillement d'Entrailles*. The noise made by flatus in the intestines. This happens often in health, especially in nervous individuals.

BORD (F.), *Margo*, *Edge*, *Margin*. Anatomists have so named the boundaries of an organ. Thus, the bones, muscles, &c. have *bords* as well as bodies. The 'free edge,' *bord libre*, is one not connected with any part; the 'adhering edge,' *bord adhérent*, one that is connected; and the *bord articulaire*, or 'articular margin or edge,' that which is joined to another bone.

BORD CILIAIRE, Ciliary margin.

BORDEAUX, MINERAL WATERS OF. Near this great city, in the south-west of France, is a saline, chalybeate spring. It contains oxide of iron, carbonate and sulphate of lime, chlorides of sodium and calcium, subcarbonate of soda, and sulphate of magnesia.

BORE, Boron.

BORE-TREE, *Sambucus*.

BORGNE (F.), *Cocles*, *Unoc'ulus*, *Luscus*, *Luscio'cus*. One who has only one eye, or sees only with one. The word has been used, figuratively, for *blind*, in surgery and anatomy. See *Cæcus*.

BORIUM, Boron.

BORKHAUSENIA CAVA, *Fumaria bulbosa*.

BORN, past participle of *bear*, (F.) *né*. Brought forth from the womb.

BORN ALIVE. It has been decided by English judges, that 'to be born alive,' means, that acts of life must have been manifested after the whole body has been extruded; and that respiration *in transitu* is not evidence that a child was born alive. It must be 'wholly born alive;' hence respiration may be a sign of *life*, but not of *live birth*.

BORON, *Bo'rium*, *Borum*, (F.) *Bore*. A simple substance, the basis of boracic acid; obtained, by heating potassium with boracic acid, as a dark,

olive-coloured powder, devoid of taste and smell. Heated in the air or in oxygen, it is converted into boracic acid.

BOR'OSAIL, *Zael*. Ethiopian names for a disease, very common there, which attacks the organs of generation, and appears to have considerable analogy with syphilis.

BORRAGO, *Borago officinalis*.

BORRI, *Curcuma longa*.

BORRIBERRI, *Curcuma longa*.

BORSE, MINERAL WATERS OF. Borse is a village in Béarn. The waters are chalybeate.

BORUM, Boron.

BOSA. An Egyptian name for a mass, made of the meal of darnel, hemp-seed, and water. It is inebrating.—Prospero Alpini. At the present day, it is applied to a kind of acidulated and, sometimes, fermented scherbet. In its commonest form, it is made by fermenting an infusion of millet-seed. A more agreeable Bosa is prepared by the Egyptians from tamarinds and honey; and by the Greeks from the fruit of *Solanum Lycopersicum*. This last is called, by them, *Αποστρεκον*, *Drosi'sticon*.

BOSCHESJESMANSTHEE, *Methystophyllum glaucum*.

BOSOM, see *Mamma*.

BOSSA, Plague token.

BOSSÉ, Hump, Protuberance — *b. Nasale*, Nasal protuberance.

BOSSU, see *Hump*.

BOSWELLIA SERRATA, see *Juniperus lycia* — *b. Thurifera*, see *Juniperus lycia*.

BOTAL FORA'MEN, *Fora'men Botal'is* seu *Botal'lii*, *Fora'men Oval'is*, (F.) *Trou de Botal*, *Trou ovale*. A large opening which exists in the fetus in the partition between the two auricles of the heart; and by means of which the blood passes from one to the other. Its discovery is generally attributed to Leonard Botalus, Botal, or Botalli, who wrote in 1562. It was spoken of, however, by Vesalius, and even by Galen.

BOTANE, Herb.

BOTANICAL DOCTOR, Herb-doctor.

BOTANIQUE MÉDICALE, Botany, medical.

BOTANY, MEDICAL, *Botan'ica Med'ica*, *Medic'na Botan'ica*, *Phytolog'ia Med'ica*, from *βοταν*, 'an herb,' (F.) *Botanique Médicale*. The knowledge of the properties, characters, &c. of those vegetables which are used in medicine.

BOTARGO, (F.) *Botargue*. A preparation made in Italy and the south of France, with the eggs and blood of the *Mugilceph'alus* or *Mullet*, strongly salted, after it has become putrescent. It is used as a condiment.

BOTARGUE, Botargo.

BOTCH, (I.) *Bozza*. Of old, a swelling of the skin. A phlegmon. A large ulcerous affection.

BOTHOR. An Arabic term for abscess in the nares. It means, also, a tumour in general; especially those which are without solution of continuity.

BOTHRIOCEPH'ALUS, *B. latus*, *Bothrioceph'alum*, from βοθριον, 'a small pit,' and κεφαλή, 'head.' *Bothrioceph'alus*, *Diboth'rium latum*, *Tænia lata* seu *vulgar'is* seu *os'culis lateral'ibus gem'inis* seu *grisea* seu *membrana'cea* seu *tene'lis* seu *denta'ta* seu *humana* *in'er'mis* seu *prima* seu *os'culis lateral'ibus solita'riis* seu *aceph'ala* seu *os'culis superficial'ibus*, *Lumbri'cus latus*, *Plat'a*, *Haly'sis membrana'cea* seu *luta*, (F.) *Ténia à anneaux courts* ou *non armé*, *Ver solitaire*, *Broad Tape-worm*. Common in Switzerland, Russia, and some parts of France. It inhabits the intestines of man, and extends to an enormous length. A broken specimen has been obtained 60 yards long.—Goëze.

BOTH'RION, *Both'rium*, from βοθριον, 'a small

pit, cavity,' &c. An *alveolus*, or small fossa. A small, deep ulcer on the cornea.—Galen, Paulus of Aegina. See *Fossette*.

BOTHRIUM, Bothrion, *Fossette*.

BOTHROS, Fovea.

BOTIN, Terebinthina.

BOTIUM, Bronchocele.

BOTOTHINUM. An obscure term, used by Paracelsus to denote the most striking symptom of a disease:—the *Flos morbi*.

BOTOU, Pareira brava.

BOTRIOCEPHALUS, Bothriocephalus.

BOTRION, Alveolus.

BOTROPHIS SERPENTARIA, *Actæa racemosa*.

BOTRYS, *Chenopodium botrys*, see *Vitis vinifera*—*b. Ambrosioides*, *Chenopodium ambrosioides*—*b. Americana*, *Chenopodium ambrosioides*—*b. Anthelmintica*, *Chenopodium anthelminticum*—*b. Mexicana*, *Chenopodium ambrosioides*.

BOTTINE (F.). A thin boot or buskin, *O'crea le'vior*. An instrument which resembles a small boot, furnished with springs, straps, buckles, &c., and used to obviate distortions of the lower extremities in children.

BOTTLE-NOSE, *Gutta rosea*.

BOTTLE-STOOP. In Pharmacy, an arrangement for giving the proper inclination to a bottle containing a powder, so as to admit of the contents being readily removed by the knife, in dispensing medicines. It consists of a block of wood with a groove in the upper surface, to receive the bottle in an oblique position.

BOTTOM, Nates.

BOTTRY-TREE, *Sambucus*.

BOUBALIOS, *Momordica elaterium*, *Vulva*.

BOUBON, *Bubo*.

BOUCAGE ANIS, *Pimpinella anisum*—*b. Majeur*, *Pimpinella magna*—*b. Mineur*, *Pimpinella saxifraga*—*b. Petit*, *Pimpinella saxifraga*.

BOUCHE, Mouth—*b. Pâteuse*, Clammy mouth.

BOUCLEMENT, Infibulation.

BOUE, see Corruption.

BOUE SPLÉNIQUE. The altered blood extravasated into the splenic cells has been so called.

BOUES DES EAUX (F.), *Boues Minérales*, *Bul'nea Cæno'ea*. The mud or swamp, formed near mineral springs, impregnated with the substances contained in such springs, and consequently possessing similar properties. The *Boues* are applied generally and topically, in France, at the springs of St. Amand, Bagnères de Luchon, Bagnols, and Barèges; in the United States, at the White Sulphur in Virginia, &c.

BOUES MINÉRALES, *Boues des eaux*.

BOUFFE, (F.). The small eminence, formed by the junction of the two lips.—Delaurens.

BOUFFE, Eructation.

BOUFFÉE DE CHALEUR, Flush.

BOUFFISSURE, Puffiness.

BOUGIE, (F.) A wax candle: *Candel'ula*, *Cande'la*, *C. cæ'ra seu medica'ta*, *Ce'reum medicatum*, *Cere'olus Chirurgo'rum*, *Da'dion*, *Speci'ulum ce'reum*, *Virga ce'rea*, *Cère'olus*. A flexible cylinder, variable in size, to be introduced into the urethra, œsophagus, rectum, &c., for the purpose of dilating these canals, when contracted. A *Simple Bougie* is composed of solid and insoluble substances; as plaster, elastic gum, gutta percha, catgut—(*Bougie de Corde à boyau*,) &c. It acts of course only mechanically.

BOUGIE, MED'ICATED, (F.) *B. Médicamenteuse*, has the addition of some escharotic or other substance to destroy the obstacle; as in the *Cautic Bougie*, (F.) *B. cautérisante*, *B. armée*, *B. emplastique*, which has a small portion of *Lunar Caus-*

tic or *Common Caustic* inserted in its extremity. Ducamp has recommended a *Bougie*, which swells out near its extremity, for the better dilating of the urethra. This he calls *B. à ventre*. The *metallic Bougie*, invented by Smyth, is a composition of metal, allowing of great flexibility; and a *hollow Bougie* is one with a channel running through it, to be used in the same manner as the catheter, or otherwise.

BOUGIE, UTERINE, Sound, uterine.

BOUILLIE, (F.) *Pulvic'ula*, *Pap*, from (F.) *bouillir*, 'to boil' Flour, beaten and boiled with milk. It is a common food for infants.

BOUILLON, (F.) from *bouillir*, 'to boil,' *Jus*, *Sorbi'tio*. A liquid food, made by boiling the flesh of animals in water. The osmazome, gelatin, and soluble salts dissolve; the fat melts, and the albumen coagulates. Bouillon is nourishing, owing to the gelatin and osmazome. The *Jus de Viande* is a very concentrated Bouillon, prepared of beef, mutton, veal, &c.

BOUILLON, in common language, in France, means a round fleshy excrescence, sometimes seen in the centre of a venereal ulcer.

BOUILLON BLANC, *Verbascum nigrum*.

BOULLONS MÉDICINAUX ou *PHARMACEUTIQUES*, *Medicinal* or *Pharmaceutic Bouillons*, contain infusions or decoctions of medicinal herbs. The *Bouillon aux herbes* is generally composed of sorrel or beet.

BOUILLON d'OS, (F.) *Bouillon from bones*, is obtained by treating bones with muriatic acid, in order to dissolve the earthy parts. The gelatin, which remains, is then boiled with a little meat and vegetables.—D'Arceet. Bouillon, however, can be easily obtained from the bones of roast meat by simple coction.

BOULLONNEMENT, Ebullition.

BOUIS, *Buxus*.

BOULE D'ACIER, *Ferrum tartarizatum—b. de Mars*, *Ferrum tartarizatum—b. de Molsheim*, *Ferrum tartarizatum—b. de Nancy*, *Ferrum tartarizatum*.

BOULEAU COMMUN, *Betula alba*.

BOULESIS, *Voluntas*.

BOULIM'IA, *Bulim'ia*, *Bulim'ius*, *Bul'timus*, *Boul'timos*, *Bulim'ia'sia*, *Bolis'mos*, *Echim'ia*, *Fames can'ina seu bovi'na seu lupi'na*, *Appeti'tus caninus*, *Appeten'tia can'ina*, *Alepha'gia*, *Cynorex'ia*, *Orex'is cyno'des*, *Bupi'na*, *Bupe'i'na*, *Phage'na*, *Phagedæ'na*, from *βovs*, 'an ox,' and *λῆμος*, 'hunger'; or from *βov*, augmentative particle, and *λῆμος*, 'hunger,' (F.) *Boulimie*, *Faim canine*, *F. dévorante*, *Polyphagie*. An almost insatiable hunger. A canine appetite. It is sometimes seen in hysteria and pregnancy; rarely under other circumstances.

BOULIMIE, *Boulimia*.

BOULOGNE, MINERAL WATERS OF. Boulogne is in the department of Pas-de-Calais, France. The waters are chalybeate.

BOUNCING BET, *Saponaria*.

BOUND, *Costive*.

BOUNTREE, *Sambucus*.

BOUQUET ANATOMIQUE DE RIOLAN, (F.) from *bouquet*, a collection of flowers or other substances tied together. A name given, by some anatomists, to the collection of ligaments and muscles, inserted into the styloid process of the temporal bone.

BOUQUET FEVER, *Dengue*.

BOURBILLON, see *Furunculus* (oore).

BOURBON-LANCY, MINERAL WATERS OF. Bourbon-Lancy is a small village in the department of Saône-et-Loire, France; where there are thermal saline springs, containing carbonic acid, chloride of sodium, and sulphate of soda, chloride of calcium, carbonate of lime, iron,

and silica. Their heat is from 106° to 135° Fahrenheit.

BOURBON L'ARCHAMBAUT, MINERAL WATERS OF. This town is in the department of Alier, six leagues west from Moulins, and has been long celebrated for its thermal chalybeate waters. They contain sulphohydric acid, sulphate of soda, magnesia, and lime, carbonate of iron, and silica. Their temperature varies between 136° and 145° Fahrenheit.

BOURBONNE-LES-BAINS, MINERAL WATERS OF. These springs are seven leagues from Langres, department of Haute-Marne, France. They are thermal and saline, and have been long celebrated. Temperature from 106° to 135° Fahrenheit. The *Facitiosus water*, (F.) *Eau de Bourbonne-les-Bains, Aqua Borronensis*, is composed of water, containing twice its bulk of carbonic acid, f3xxss; chloride of sodium, f5j, chloride of calcium, gr. x, &c.

BOURBOULE, MINERAL WATERS OF. A village near Mont d'Or, where there are two thermal saline springs.

BOURDAINE, *Rhamnus frangula*.

BOURDONNEMENT, *Tinnitus aurium*.

BOURDONNET, *Pulvis*, *P. e linamentis confectus seu rotundus, Dossil*. A term in French surgery for charpie rolled into a small mass of an olive shape, which is used for plugging wounds, absorbing the discharge, and preventing the union of their edges. In cases of deep and penetrating wounds, as of the abdomen or chest, a thread is attached to them by which they may be readily withdrawn, and be prevented from passing altogether into those cavities.

BOURGÈNE, *Rhamnus frangula*.

BOURGEON, Granulation, Papula—*b. Chariv.*, Granulation.

BOURGEONNEMENT, see Generation.

BOURGEONS, *Gutta rosea*.

BOURGUÉPINE, *Rhamnus*.

BOURRACHE, *Borago officinalis*.

BOURRELET (F.) A *Pad*, a *Border*. A fibro-cartilaginous border, which surrounds certain articular cavities, such as the glenoid cavity of the scapula and the acetabulum; by which the depth of those cavities is augmented. See *Corpus Callosum*.

BOURRELET ROULÉ, *Cornu ammonis*.

BOURSE à BERGER, *Thlaspi bursa*—*b. à Pasteur*, *Thlaspi bursa*.

BOURSE (LES), *Scrotum*—*b. Mucilagineses*, *Bursæ mucosæ*—*b. Muqueuses*, *Bursæ mucosæ*.

BOURSOUFFLURE, Puffiness.

BOURTREE, *Sambucus*.

BOUSSOLE, *Glossanthrax*.

BOUT DE SEIN (F.), 'End of the breast,' 'nipple.' An instrument of caoutchouc or ivory to place over the nipple when the infant sucks, soon after delivery, or where the nipple is in a morbid condition.

BOUTON, Papula—*b. d'Alep*, see Anthrax—*b. Malin*, see Anthrax—*b. d'Or*, *Ranunculus acris*.

BOUTONNIÈRE (F.), *Fissura, Incisio*. A long incision made into the urethra to extract a calculus from the canal, when it is too large to be discharged.

Also, a small incision or puncture, made in the peritoneum, or above the pubis, to penetrate the bladder in certain cases of retention of urine.

BOVACHEVO, *Datura sanguinea*.

BOVILLÆ, *Rubeola*.

BOVISTA, *Lycoperdon*.

BOWEL, Intestine.

BOWLEGGED, see *Cnemosciosis*.

BOWMAN'S CAPSULE, see *Kidney*.

BOWMAN'S ROOT, *Euphorbia corollata*, *Gil. lenia trifoliata*, *Leptandria purpurea*.

BOXBERRY, *Gaultheria*.

BOX, MOUNTAIN, *Arbutus uva ursi*.

BOX TREE, *Buxus*, *Cornus Florida*.

BOXWOOD, *Cornus Florida*.

BOYAU, Intestine—*b. Pollinique*, see *Pollen*.

BRABYRON, *Prunum Damascenum*.

BRACHERIOLUM, Truss.

BRACHERIUM, Truss.

BRACHIA COPULATIVA, see *Peduncles of the Cerebellum*.

BRACHIA PONTIS, see *Peduncles of the Cerebellum*.

BRACHILEUS, *Brachial*—*b. Internus*, *Brachialis anterior*.

BRACHIAL, *Brachialis*, *Brachius*, from *Brachium*, 'the arm.' What belongs to the arm.

BRACHIAL APONEUROSIS. An aponeurosis, formed particularly by expansions of the tendons of the latissimus dorsi, pectoralis major, and deltoides muscles, and which completely envelops the muscles of the arm.

BRACHIAL ARTERY, *Arteria brachialis*, *Humeral Artery*, (F.) *Artère ou Tronc brachial*. The artery, which extends from the axilla to the bend of the elbow; where it divides into *A. cubitalis* and *A. radialis*. It passes along the internal edge of the biceps, behind the median nerve and between the accompanying veins. Under the name *Brachial Artery*, Chaussier includes the subclavian, axillary, and humeral, the last being the *brachial proper*.

BRACHIAL MUSCLE, ANTERIOR, *Musculus Brachialis Anterior*, *Brachialis internus* seu *antiscus*, *Brachius*, *B. internus*, (F.) *Muscle brachial interne*, *Huméro-cubital*—(Ch.) This muscle is situate at the anterior and inferior part of the arm, and before the elbow-joint. It arises, fleshy, from the middle of the os humeri, and is inserted into the coronoid process of the ulna. *Use*, To bend the fore-arm.

BRACHIAL PLEXUS, *Plexus Brachialis*, is a nervous plexus, formed by the interlacing of the anterior branches of the last four cervical pairs and the first dorsal. It is deeply seated in the hollow of the axilla, and extends as far as the inferior and lateral part of the neck. It gives off the *thoracic nerves*, *supra* and *infra scapular*, and the *brachial* (which are six in number), the *axillary*, *cutaneous*, *musculo-cutaneous*, *radial*, *cubital*, and *median*.

BRACHIAL VEINS are two in number, and accompany the artery, frequently anastomosing with each other: they terminate in the axillary. Under the term *Brachial Veins*, Chaussier includes the humeral, axillary, and subclavian.

BRACHIALE, *Carpus*.

BRACHIAL'GIA, *Neural'gia Brachialis*, from *βραχίων*, 'the arm,' and *algos*, 'pain.' Pain in the arm, neuralgia of the arm.

BRACHIALIS, *Brachial*—*b. Anticus*, *Brachial muscle*—*b. Externus*, see *Triceps extensor cubiti*—*b. Internus*, *Brachial muscle*.

BRACHERIUM, Truss.

BRACHILE, Truss.

BRACHILUVIUM, Bath, arm.

BRACHIO-CEPHALIC ARTERY, *Innominate arteria*—*b. Veins*, *Innominate venæ*.

BRACHIO-CUBITAL, *Brachio-cubitalis*. That which belongs both to the arm and cubitus. This name has been given to the internal lateral ligament of the elbow-joint; because it is attached to the os brachii or os humeri and to the cubitus or ulna.

BRACHIOCYCLOSIS, from *βραχίων*, 'the arm,' and *κυλλωσις*, 'the act of making crooked.'

Curvature of the arm inwards. Paralysis or loss of power from curvature of the arm.

BRACHION, Brachium.

BRACHIONŌUS, from βραχιον, 'the arm,' and οζος, 'a swelling.' A tumour of the arm.

BRACHIO-RA'DIAL, Brachio-radialis. That which belongs to the brachium and radius. This name has been applied to the external lateral ligament of the elbow-joint, because it is attached to the humerus and to the radius. See *Supinator radii longus*.

BRACHIORRHEU'MA, Rheumatismus brachii, from βραχιον, 'the arm,' and ρευμα, 'defluxion, rheumatism.' Rheumatism of the arm.

BRACHIOTOMY, Brachiotomia, from βραχιον, 'arm,' and τομη, 'incision.' Amputation of the arm.

BRACHIOLOM, Truss.

BRACHIUM, Brachion, Laceratus, (F.) Bras, 'the arm.' The arm from the shoulder to the wrist, or the part between the shoulder and elbow. See *Humeri Os*.

BRACHIUM ANTERIUS. A rounded process, which passes from the anterior pair of the corpora quadrigemina (nates) obliquely outwards into the thalamus opticus.

BRACHIUM MOVENS QUARTUS, Latissimus dorsi.

BRACHIUM POSTERIUS. A rounded process, which passes from the posterior pair of the quadrigemina (testes) obliquely outwards into the optic thalamus.

BRACHUNA, Nymphomania, Satyriasis.

BRACHYAU'CHEN, from βραχυς, 'short,' and αυχην, 'neck.' One who has a short neck.

BRACHYCEPH'ALÆ (Gentes), 'short heads,' from βραχυς, 'short,' and κεφαλη, 'head.' In the classification of Retzius, those nations of men whose cerebral lobes do not completely cover the cerebellum—as the Slavonians, Fins, Persians, Turks, Tartars, &c.

BRACHYCHRONIUS, from βραχυς, 'short,' and χρονος, 'time.' That which continues but a short time. A term applied to diseases which are of short duration.—Galen.

BRACHYGNA'THUS, from βραχυς, 'short,' and γναθος, 'the under jaw.' A monster with too short an under jaw.—Gurlt.

BRACHYNSIS, Abbreviation.

BRACHYPNŌEA, Dyspnœa.

BRACHYPOTI, from βραχυς, 'short,' and ποτος, 'drinker.' They who drink little, or who drink rarely. Hippocrates, Galen, Foësius.

BRACHYRHYN'CHUS, from βραχυς, 'short,' and ρυγχος, 'snout.' A monster with too short a nose.

BRACHYSMOS, Abbreviation.

BRACING, Corroborant.

BRACKEN, Pteris aquilina.

BRADYŒSTHE'SIA, from βραδύς, 'difficult,' and αἰσθησις, 'sensation.' Impaired sensation.

BRADYBOLISMUS, Bradyspermatismus.

BRADYECŌIA, Deafness.

BRADYLOG'IA, Dyslogia; from βραδύς, 'difficult,' and λογος, 'a discourse.' Difficulty of speech.

BRADYMASE'SIS, Bradymasse'sis, improperly Bradymaste'sis, Masticatio difficilis, from βραδύς, 'difficult,' and μασσις, 'mastication.' Difficult mastication. See *Dysmasesis*.

BRADYMASTESIS, Bradymasesis.

BRADYPEP'SIA, Tarda ciborum concoctio, from βραδύς, 'slow,' and πεπω, 'I digest.' Slow digestion.—Galen. See *Dyspepsia*.

BRADYSPERMATIS'MUS, Bradybolis'mus, Ejaculatio seminis impedita, Dyspermatis'mus, from βραδύς, 'slow,' and σπερμα, 'sperm.' A slow emission of sperm.

BRADYSUR'IA, Tenes'mus vesicæ, (F.) Ténue vesical, from βραδύς, 'difficult,' and ουρην,

'to pass the urine.' Painful evacuation of the urine, with perpetual desire to void it. Dysuria. **BRADYTŌCIA, Dystocia.**

BRAG'GET, Braggart, Bragwort. A name formerly applied to a tisan of honey and water. See *Hydromeli*.

BRAI LIQUIDE, see Pinus sylvestris—b. Sec. Colophonia.

BRAIN, Cerebrum—b. Fag, see Nervous diathesis.

BRAIN, LITTLE, Cerebellum.

BRAIN PAN, Cranium—b. Sand, see Pineal gland.

BRAINE, MINERAL WATERS OF. Braine is a small village, three leagues from Soissons, France, which has purgative waters similar to those of Passy.

BRAIRETTE, Primula veris.

BRAKE, COMMON, Pteris Aquilina—b. Rock, Polypodium vulgare, Polypodium ineanum.

BRAKE Root, Polypodium vulgare.

BRAGING, Vomiting, Vomiturition.

BRAMBLE, AMERICAN HAIRY, see Rubus fruticosus—b. Common, Rubus fruticosus.

BRAMBLE-BERRIES, see Rubus fruticosus.

BRAN, Furfur.

BRANC-URSINE BÂTARDE, Heracleum spondylium.

BRANCA GERMANICA, Heracleum spondylium—b. Ursina, Acanthus mollis—b. Vera, Acanthus mollis.

BRANCH, from (F.) Branche, originally, probably, from βραχιον, 'an arm,' [?] because branches of trees, &c., go off like arms. A term applied, generally, to the principal division of an artery or nerve. The word is commonly used synonymously with *Ramus*; but often, with the *Rameau*, *Branche* signifies the great division;—*Rameau*, Lat. *Ramus*, the division of the branches; and *Ramuscules*, Lat. *Ramusculi*, the divisions of these last.

The French, also, speak of the *branches* of the pubis for the *Rami* of that bone, *branches* of the ischium for the *rami* of the ischium, &c.

BRANCHES DE LA MOËLLE ALLONGÉE (PETITES), Corpora restiformia.

BRANCHI, Branchæ. Swellings of the tonsils, or parotid, according to some;—of the thyroid gland, according to others.

BRAN'CHIA, (Gr.) βραγchia. The gills or respiratory organs of fishes, corresponding to the lungs of terrestrial animals.

BRANCHUS, βραγχος, Branc's do. A catarrhal affection of the mucous membranes of the fauces, trachea, &c.—Galen. Hoarseness.

BRANCI, Cynanche tonsillaris.

BRANCIA, Vitrum.

BRANDWINE, Brandy.

BRANDY, (G.) Brannntwein, Dutch, Brandwijn, 'burnt wine,' (Old Eng.) Brandywine. Vinum adustum seu crematum, Aqua Vitæ, (F.) Spiritus vini Gallici, (Ph. U. S.) Spiritus Gallicus, Eau de vie, (S.) Aguardiente. The first liquid product obtained by distilling wine. It is composed of water, alcohol, and an aromatic oily matter, which gives it its flavour. Brandy is a powerful and diffusible stimulant, and as such is used in medicine. It has been also called *Liquor Aquile'gius*. See *Spirit*.

BRANDY, APPLE, see Pyrus malus—b. Egg, see Ovum.

BRANDY-BOTTLES, see Nymphæa lutea.

BRANDY-FACE, Gutta rosea.

BRANKS, Cynanche parotidæa.

BRANKURSINE, Acanthus mollis.

BRANNTWEIN, Brandy.

BRAS. See Oryza.

BRAS, Brachium—b. du Cervelet, Corpora restiformia.

BRASDOR'S OPERATION FOR ANEURISM. An operation by ligature, proposed by Brasdor, which consists in the application of the ligature on the distal side of the tumour.

BRASÉGUR, MINERAL WATERS OF. Brasegur is a place in the diocese of Rhodéz, where there are cathartic waters.

BRASENIA, B. Hydropeltis.

BRASENIA HYDROPELTIS, Brase'nia, B. Pel-tu'ta, Hydropel'tis purpu'rea, Gelatina aquat'ica, Frogleaf, Little Water Lily, Water Jelly, Water shield, Water target, Deerfood. An indigenous plant, *Nat. Ord. Ranunculaceæ, Sex. Syst. Poly-andria Polygynia*, flourishing from Kentucky to Carolina and Florida; and covering the surface of ponds, marshes, &c. The fresh leaves are mucilaginous, and have been used in pulmonary complaints, dysentery, &c., like Cetraria.

BRASENIA PELTATA, B. Hydropeltis.

BRASH, (Sc.), of uncertain etymology. A transient attack of sickness. A bodily indisposition.—Jamieson. A rash or eruption. (Prov. West of England.)

BRASH, WATER, Pyrosis,

BRASH, WEANING, (Sc.) Speaning brash, Atroph'ia Ablactatorum. A severe form of diarrhœa, which supervenes at times on weaning. The *Maladie de Crueithier* appears to be a similar affection.

BRASILETTO, see Cæsalpinia.

BRASIUM, Malt.

BRASMOS, Fermentation.

BRASS, Sax. bray, Welsh, prés. Aurichal-cum, properly, Orichal'cum, ('mountain brass,' from *ops*, 'a mountain,' and *chalcos*, 'brass.') *Chrysochal'cos, Chalcos, Escavum, (F.) Airain.* A yellow metal, formed by mixing copper with calamine. The same general remarks apply to it as to copper. See Cuprum.

BRAS'SICA, Crambè, B. olera'cea seu capita'ta seu cuna'na of the old Romans. The Cabbage, (F.) *Chou potager. Order, Cruciferae. Sex. Syst. Tetradynamia Siliquosa.* Cato wrote a book on its virtues. It is a vegetable by no means easy of digestion when boiled; when raw, it appears to be more digestible. When forming a solid globular mass, like a head, it is the *B. Capita'ta, (F.) Chou-Cabus, Chou Pommé.*

BRASSICA CANINA, Mercurialis perennis—b. Capitata, Brassica—b. Cumana, Brassica.

BRAS'SICA ERU'CA, B. his'pida, Eru'ca, E. fe'tida seu sati'va, Sina'pis eru'ca, Sisym'brium erucas'trum, Garden Rocket, Roman Rocket, Shinlock, &c., (F.) Chou Roquette, Roquette. This was considered by the Romans an aphrodisiac.—Columella. The seeds were ordinarily used.

BRAS'SICA FLOR'IDA, —Bras'sica Pompeia'na of the ancients—the Cauliflower, *Caulis Flor'ida, (F.) Chou-fleur*, is a more tender and digestible variety.

The *Broc'coli, B. Sabel'lica* of the Romans, *B. Ital'ica*, belongs to this variety.

BRASSICA HISPIDA, B. eruca—b. Italica, B. Florida—b. Marina, Convolvulus soldanella.

BRAS'SICA NAPUS, Napus Sylvestris, Bu'nias, Rape, (F.) Navette. The seed yields a quantity of oil.

BRASSICA NIGRA, Sinapis nigra—b. Oblonga, B. rapa—b. Oleracea, Brassica—b. Pompeiana, B. Florida.

BRAS'SICA RAPA, Rapa rotun'da seu oblon'ga seu napus, Rapum majus, Sina'pis tuberosa, Turnip, (F.) Chou navet, Navet, Rave. The turnip is liable to the same objection (but to a less extent) as the cabbage.

BRASSICA SABELLICA, B. Florida.

BRATHU, Juniperus sabina.

BRATHYS, Juniperus sabina.

BRAWLINS, Arbutus uva ursi, Vaccinium vitis idæa.

BRAWN, Muscle.

BRAYER, Truss.

BRAYERA ANTHELMINTICA, Hagenia Abyssinica.

BRAZIL WOOD, Cæsalpinia echinata.

BREAD, see Triticum.

BREAD, GLUTEN. Bread made of wheat dough deprived of the chief portion of its starch by washing. Bread, made of gluten only, cannot be eaten, on account of its hardness and toughness; hence one-fifth of the normal quantity of starch is allowed to remain, and in this form the bread is said to be tolerably light, eatable, and moderately agreeable.

BREAD, HOUSEHOLD, Syncomistos.

BREADBERRY, Pap.

BREAD-FRUIT TREE, Artocarpus.

BREAKBONE FEVER, Dengue.

BREAST, Thorax, Mamma—b. Abscess of the, Mastodynia apostematosa—b. Pigeon, see Lardosis.

BREAST-GLASS, Milk-glass. A glass applied to the nipple to receive the milk when secreted copiously by the mamma.

BREAST, IRRITABLE, Neuralgia mammae.

BREAST-PANG, SUFFOCATIVE, Angina pectoris.

BREAST-PUMP, Antlia Lactea.

BREASTWEED, Saururus cernuus.

BREATH, Sax. bræse, (Old Eng.) Aande, and Ande, Hal'itus, Anhel'itus, An'imus, Spir'itus, A'mos, (F.) Haleine. The air expelled from the chest at each expiration. It requires to be studied in the diagnosis of thoracic diseases especially. See Respiration.

BREATH, OFFENSIVE, Fætor Oris, Catostomatophre'sia, Hal'itus oris fæ'tidus, Ozæ. An offensive condition, which is usually dependent upon carious teeth, or some faulty state of the secretions of the air passages. The internal use of the chlorides may be advantageous.

BREATH, SATURNINE, see Saturnine—b. Short, Dyspnœa.

BREATHING AIR, see Respiration.

BREATHING, DIFFICULTY OF, Dyspnœa.

BRECHET, (F.) The Brisket. This name is given in some parts of France to the *cartilago ensiformis*, and sometimes to the sternum itself.

BRECHMA, Bregma.

BRECHMUS, Bregma.

BRÉDISSURE (F.), Trismus Capistratus. Incapacity of opening the mouth, in consequence of preternatural adhesion between the internal part of the cheek and gums; often occasioned by the abuse of mercury.

BREDOUILLEMENT (F.), Tituban'tia. A precipitate and indistinct mode of utterance, in which a part only of the words is pronounced, and several of the syllables viciously changed. This defect is analogous to stuttering, but differs from it in being dependent on too great rapidity of speech; whilst stuttering is characterized by continual hesitation, and frequent repetition of the same syllables.

BREE, Supercilium.

BREED, Race.

BREEDING, Generation, Pregnant.

BREEDING, CROSS. The act of raising or breeding from different stocks or families.

BREEDING-IN-AND-IN. The act of raising or breeding from the same stock or family.

BREELLS, Spectacles.

BREGMA, Breehma, Brechmus, from βρεχμα, 'to sprinkle; Fontanel'la, Sina'ciput. The top of the head was thus called, because it was believed to be humid in infants; and, according to

some, because it was conceived to correspond to the most humid part of the brain.

BREGMATODYMIA, see *Cephalodymia*.

BRENNING, Burning.

BREPHOTONON, *Conyza squarrosa*.

BREPHOTROPHE'UM, *Echthobrephotrophe'um*, from *βρεφος*, 'a new-born child,' and *τροφειν*, 'to nourish.' A founding hospital.

BREUILLET, *Cesalpinia sappan*.

BRE'VIA VASA, *Short Vessels*. This name has been given to several branches of the splenic arteries and veins, which are distributed to the great *cul-de-sac* of the stomach.

BREVIS CUBITI, see *Anconens*.

BRICK, (F.) *Brique*. Hot bricks are sometimes used to apply heat to a part, as to the abdomen in colic, or after the operation for popliteal aneurism: or, reduced to very fine powder, and mixed with fat, as an application to herpetic and psoric affections.

BRICKS, *Fornaceæ Testæ* or *Tiles* were formerly bruised in vinegar, and the liquid was used as a specific in cutaneous affections. They entered, also, into a cerate used for scrofulous humours, &c. To the *Terra Fornacea*, or *Brick earth*, the same virtues were assigned.

BRICKLIKE SEDIMENT, see *Lateritious*.

BRICUMUM, *Artemisia*.

BRIDE (F.), 'a bride.' *Fræ'nulum*, *Retinaculum*. This term is given, in the plural, to membranous filaments, which are found within abscesses or deep-seated wounds, and which prevent the exit of pus. The term is, also, applied to preternatural adhesions, which occur in cicatrices of the skin, in the urethra, or in inflamed serous or synovial membranes.

BRIEF, *Rife*.

BRIER, *WILD*, *Rosa Canina*.

BRIGHT'S DISEASE OF THE KIDNEY, see *Kidney*, *Bright's disease* of the.

BRIGHTON, CLIMATE OF. The air of this fashionable watering-place, on the south coast of England, is dry, elastic, and bracing. Its climate appears to the greatest advantage in the autumn and early part of the winter; when it is somewhat milder and more steady than that of Hastings. Accordingly, it is adapted for all cases in which a dry and mild air at this season of the year proves beneficial. In the spring months, owing to the prevalence of, and its exposure to, north-east winds, the climate is cold, harsh, and exciting to the delicate. It is well adapted for convalescents, and for all who require a dry and bracing sea air.

BRIMSTONE, Sulphur.

BRINE, *Muria*.

BRINTON ROOT, *Leptandria purpurea*.

BRION, *Corallina*.

BRIQUE, *Brick*.

BRIQUEBEC, MINERAL WATERS OF. This town is three leagues from Cherbourg, in France. The water contains chloride of iron.

BRIQUETÉ, *Lateritious*.

BRISE-COQUE (F.), from *briser*, 'to break to pieces,' and *coque*, 'a shell.' An instrument designed by Heurteloup for breaking to pieces the shell of a vesical calculus, after it has been hollowed by his *mandrin à virgule*.

BRISE-PIERRE ARTICULÉ (F.), from *briser*, 'to break to pieces,' and *piierre*, 'a stone.' An instrument invented by Jacobson for crushing the stone in the bladder.

BRISTOL HOT WELL, *Bristolien'sis Aqua*. Bristol is about thirteen miles from Bath, in England. The water is an almost pure thermal; slightly acidulated. It contains chlorides of magnesium and sodium, sulphate of soda, sulphate of lime, carbonate of lime, carbonic acid,

oxygen and azote. Temperature, 74° Fah. The *Hot Well* has been long celebrated. Its action is like that of thermal waters in general. The climate of Bristol is mild, and hence the water has been celebrated for the cure of incipient pulmonary consumption. See *Clifton*.

BRIZOCERAS, *Ergot*.

BROAD, *Sac. βρᾶς, Latius*, (F.) *Large*. Any body is so termed whose transverse extent is considerable compared with its length. The *Broad Bones*, such as the frontal, parietal, occipital, iliac, aid in forming the parietes of splanchnic cavities. *Broad Muscles* generally occupy the parietes of cavities, and especially those of the chest and abdomen. The epithet has also been applied to other parts—as to the *broad ligaments* of the womb, &c.

BROCCOLI, *Brassica sabellica*.

BROCHIOS, *βροχος, La'queus*. A bandage.

BROCHT, Vomiting.

BROCH'THUS, *βροχ'θος, Gula*. The throat. Also, a kind of small drinking vessel.—*Hipp*.

BROCHUS, *βροχος*. This name has been given to one who has a very prominent upper lip. According to others it means one whose teeth project in front of the mouth.

BRO'DIUM. A synonym of *Jus* or *Jusculum*. Broth, or the liquor in which any thing is boiled. *Bro'dium salis*—a decoction of salt.

BROIEMENT, see *Cataract*, *Laceration*—*b. de la Pierre*, *Lithotrixy*.

BROKELEAK, *Rumex hydrolapathum*.

BROKEN DOSES, see *Doses*, *broken*.

BROKEN-WINEDNESS, *Asthma*.

BROMA, *Aliment*, *Bromine*.

BROMATOECRISIS, *Lientery*.

BROMATOGRAPHY, *Bromatograph'ia*, *Bromography*, *Bromograph'ia*, from *βρωμα*, 'food,' and *γραφειν*, 'a description.' A description of aliments.

BROMATOL'OGY, *Bromatolog'ia*, *Sitiol'ogy*, from *βρωμα*, 'food,' and *λογος*, 'a discourse.' A treatise on food.

BROME, *Bromine*.

BROMEGRASS, *Bromus ciliatus*—*b. Soft*, *Bromus ciliatus*.

BROME'LIA ANA'NAS, called after *Olaus Bromel*, a Swede. *Carduus Brasilia'nus*, *Ana'nas ova'ta* seu *aculea'ta*, *Anas'sa*, *Capa-Isiak'ka*, *Ana'nas* or *Pine Apple*, *Soursop*. Order, *Bromeliaceæ*. A West India tree, which produces the most delicious of fruits.

BROME'LIA PINGVIN, *Ana'nas America'na*, *Pinguin*, *Broad-leaved wild Ana'nas*, &c. The West India plant, which affords the *Pinguin* fruit. The fruit is refrigerant, and the juice, when ripe, very austere. It is used to acidulate punch. A wine is made from the *Pinguin*, which is very intoxicating, and has a good flavour.

BROMIC, *Bro'miens*; same eymon as *Bromine*. Containing bromine.

BROMIDE OF IRON, see *Bromine*—*b. of Mercury*, see *Bromine*—*b. of Potassium*, see *Bromine*.

BROMIDRO'SIS, from *βρωμος*, 'stench,' and *ἰδρως*, 'sweat.' Offensive sweat.

BROMINE, *Bro'minum* (Ph. U. S., 1842), *Bro-min'um* (1851), *Broma*, *Bromin'eum*, *Bro'mium*, *Bro'mina*, *Bromum*, *Mu'rina*, *Muride*, *Brome*. A simple body, of a very volatile nature, and highly offensive and suffocating odour, whence its name, from *βρωμος*, 'a stench.' It is met with chiefly in sea-water, and in many animal and vegetable bodies that live therein. It has likewise been found in many mineral waters of this and other countries. In its chemical relations, it may be placed between chlorine and iodine. With oxygen it forms an acid—the *Bromic*, and with hydrogen another—the *Hydrobromic*.

PURE BROMINE; BROMIDE OF IRON, *Ferri Bro-midum*, *Verum bromat'um*, (F.) *Bromure de Fer*, in solution, Hydrobromate of Iron, *Ferri hydro-bromas*, *Ferrum hydrobromicum oxydatum*, (dose, gr. i or ij.); and BROMIDE OF POTASSIUM, *Potas-sii bromidum*, (F.) *Bromure de potassium*, have been used medicinally, and chiefly in scrofulosis—internally, as well as applied externally. Bromine may be dissolved in forty parts of distilled water, and six drops be commenced with as a dose. BROMIDES OF MERCURY (*Hydrargyri Bro-mida*, (F.) *Bromures de Mercure*), have been given in syphilis. The *protobromide* and the *bibromide* are analogous in composition and medicinal properties to the corresponding iodides of mercury.

CHLORIDE OF BROMINE, *Bromin'ii chlo-ridum*—made by passing chlorine through bromine, and condensing the resulting vapours by cold, has been prescribed internally, as well as externally, in cancer; but chiefly in the latter mode, in the form of a caustic paste, either alone or with other chlorides—as those of zinc, antimony, and gold.

BROMIUM, Bromine.

BROMOGRAPHY, Bromatography.

BROMOS, *βρωμος*. One of the cerealia, sup-posed, by some, to be oats. See *Avena*.

BROMOSUS, Fetid.

BROMUM, Bromine.

BROMURE DE FER, see Bromine—*b. de Mercure*, see Bromine—*b. de Potassium*, see Bromine.

BROMUS CILIATUS, *B. purgans*, *Brome grass*; indigenous: Order, Gramineæ; is said to be emetic, and anthelmintic [?], cathartic and diuretic. It purges cattle.

BROMUS GLABER, *Triticum repens*.

BROMUS MOLLIS, *Soft Brome Grass*. The seeds are said to cause giddiness in man, and to be fatal to poultry.

BROMUS PURGANS, *B. ciliatus*—*b. Temulentus*, *Lolium temulentum*.

BRONCHARCTIA, Bronchiostenosis.

BRONCHECTASIS, Bronchia, dilatation of the.

BRONCHES, Bronchia—*b. Ganglions lym-phatiques des*, Bronchial glands.

BRONCHI, Bronchia.

BRONCHIA, *Bronchiæ*, *Bronchi*, from *βρογχος*, 'the throat.' The Latins used the term *Bronchus* for the whole of the trachea; whilst they called its ramifications *Bronchia*. *Bronchia*, *Bronchiæ*, and *Bronchi*, (F.) *Bronches*, now mean the two tubes, with their ramifications, which arise from the bifurcation of the trachea, and carry air into the lungs—*Canalæ pulmo-num*, *Syringes seu Canales aëri'eri*.

BRONCHIA, DILATATION OF THE, *Dilated Bronchia*, *Bronchectasis*, *Bronchiectasis*, *Dilatatio bronchio-rum*. The physical signs of this condition are the following: *Percussion* usually clear, but not unfrequently less so than natural, although very seldom quite dull. *Auscultation* detects coarse mucous or gurgling rhonchi, increased by the cough, combined with, or replaced by, bronchial or cavernous respiration, which is often effected as if by a sudden puff or whiff. The resonance of the voice is increased, but it seldom amounts to perfect pectoriloquy. The most common situations for dilated bronchia are the scapular, mammary, or lateral regions. They are almost always confined to one side.

BRONCHIA, OBLITERATION OR COMPRESSION OF THE. The inspiratory murmur on auscultation is weaker or wholly suppressed over a limited portion of the chest; the expiration is generally more distinct and prolonged; all the other conditions are natural.

BRONCHIE, see Bronchia.

BRONCHIAL, *Bronchic*, *Bronchia'lia*, *Bron-chicus*, (F.) *Bronchial*, *Bronchique*. That which relates to the bronchia.

BRONCHIAL ARTERIES, (F.) *Artères Bronchiques*. These are generally two in number, one going to each lung. They arise from the thoracic aorta, and accompany the bronchia in all their ramifications.

BRONCHIAL CELLS, (F.) *Cellules Bronchiques*. The *Air-cells*; the terminations of the bronchia.

BRONCHIAL COUGH, (F.) *Toux bronchique*, *T. tubaire*. This generally accompanies bronchial respiration. They both indicate obstruction to the entrance of air into the air-cells.

BRONCHIAL GLANDS, *Glandulæ Vesicula'nae*, *Glandæ Vesicula'nae*, (F.) *Glandes bronchiques*, *Ganglions lymphatiques des bronches*, are numerous glands of an ovoid shape; of a reddish hue in the infant, and subsequently brown and black, seated in the course of the bronchia. Their functions are unknown. The bronchial glands may be presumed to be affected by scrofulosis, when, in addition to the existence of tumours in the neck, percussion gives a dull sound under the upper and central part of the sternum, whilst there is no appreciable lesion of the lungs.

BRONCHIAL NERVES, (F.) *Nervi bronchiques*, are furnished by the two pulmonary plexuses.

BRONCHIAL PHTHISIS, see Phtthisis bronchial—*b. Respiration*, see Murmur, respiratory.

BRONCHIAL VEINS arise from the last divisions of the arteries of the same name, and pass, on the right side, into the vena azygos; on the left, into the superior intercostal.

BRONCHIC, Bronchial.

BRONCHIECTASIS, Bronchia, dilatation of the.

BRONCHITIS, Bronchitis.

BRONCHIOLE, *Bronchiolum*, *Bronchiolus*; diminutive of *Bronchium* or *Bronchus*. A minute bronchial tube.

BRONCHIOPEUMONIA, Bronchopneumonia.

BRONCHIOSTENOSIS, *Bronchiarectia*, from *βρογχος*, 'a bronchus,' and *στενωσις*, 'contraction.' Contraction or narrowness of the bronchi.

BRONCHITE CONVULSIVE, Pertussis—*b. Pseudomembraneæ*, Polypus bronchialis.

BRONCHITIS, *Bronchiti'is*, *Inflammatio bronchio-rum*, *Catar'rhus Pulmo-num seu bronchio-rum*, *Pleuriti'is humida seu bronchiti'is*, *Broncho'tasis*, *Angi'na bronchia'lia*, *Pul'monary Catarrh*, (F.) *Inflammation des Bronches*, *Inflammation of the lining membrane of the bronchial tubes*. This is always more or less present in cases of pulmonary catarrh; and is accompanied by cough, mucous expectoration, dyspnoea, and more or less uneasiness in breathing. The acute form is accompanied with all the signs of internal inflammation, and requires the employment of antiphlogistics followed by revulsives. The chronic form, *Tussis seni'lis*, *Catar'rhus seni'lis*, *Rheuma catarrha'te*, *Peripneumonia nobilis*, *Bronchorrha'ia acuta*, *Winter Cough*, *Chronic Catarrh*, may be confounded with phtthisis; from which it must be distinguished mainly by the absence of hectic fever and of the physical signs that are characteristic of the latter, as well as by the nature of the expectoration, which is generally mucous, although at times mucopurulent. When the expectoration is little or none, the bronchitis is said to be dry, dry catarrh, (F.) *Catarrhe Sec.*

When bronchitis affects the smaller tubes, it is termed *capillary bronchiti'is*, *bronchi'tis capilla'ris*, *broncho'ac'c' infanti'lis* [?], and is often fatal to children. *Vesicular bronchitis* is the term proposed by M.M. Rilliet and Barthez for the vesicular pneumonia of children.

BRONCHITIS, Catarrh—b. Asthenica, Peripneumonia notha—b. Capillary, see Bronchitis—b. Chronic, see Bronchitis—b. Crouposa, Polypus bronchialis—b. Convulsiva, Pertussis—b. Exsudativa, Polypus bronchialis.

BRONCHITIS, MECHANICAL. Inflammation of the lining membrane of the air-tubes, induced by the inhalation of irritating particles.

BRONCHITIS MEMBRANACEA, Polypus bronchialis—b. Plastic, Polypus bronchialis—b. Pseudomembranous, Polypus bronchialis—b. Summer, Fever, hay—b. Vesicular, see Bronchitis.

BRONCHIUS, Sterno-thyroides.

BRONCHLEMMITIS, Polypus bronchialis.

BRONCHOCACE, Peripneumonia notha—b. Infantilis, see Bronchitis.

BRONCHO-CATARRHUS, Catarrh.

BRONCHOCE/LE, from *βρογχος*, 'a bronchus,' and *κηλη*, 'tumour.' An inaccurate name for the affection which is called, also, *Bo'chium*, *Botium*, *Bocium*, *Hernia gutturis seu gutturalis seu bronchialis*, *Guttur uterium seu globosum*, *Tracheolophy'ma*, *Thyreoc'le*, *Thyreoc'le*, *Tracheoc'le*, *Tracheloc'le*, *Thyremphez'is*, *Thyreophez'is*, *Thyrophraz'is*, *Thyreopraz'is*, *Thyron'cus*, *Thyron'cus*, *Deiron'cus*, *Deron'cus*, *Gossam*, *Gottum*, *Ecechebron'chus*, *Gongro'na*, *Struma*, *Glands*, *Tuber gutturo'sum*, *Gut'teria*, &c., the Derbyshire neck, *Snelled neck*, *Wen*, *Gaitre*, &c., (F.) *Gaitre*, *Gouëtre*, *Hypertrophie du Corps Thyroïde*, *Grosse Gorge*, *Gros Cou*. This is no rupture, but consists of an enlargement of the thyroid gland. It is common at the base of lofty mountains in every part of the world; and has been supposed to be owing to the drinking of snow-water, but it occurs where there is no snow. The tumour is sometimes very extensive. Iodine has great power over it, and will generally occasion its absorption, when the case has not been of such duration as to have ended in a cartilaginous condition.

BRONCHOCEPHALITIS, Pertussis.

BRONCHOPARALYSIS, Asthma.

BRONCHOPHONY, Resonance—b. Pectoriloquous, Pectoriloquy—b. Strong, Pectoriloquy.

BRONCHOPLASTIC, *Bronchoplasticus*, from *βρογχος*, 'a bronchus,' and *πλασσω*, 'I form.' An epithet given to the operation for closing fistule in the trachea: *Bronchoplasty*.

BRONCHOPNEUMONIA, *Bronchiopneumonia*, from *βρογχος*, 'a bronchus,' and *Pneumonia*. Inflammation of the bronchia and lungs.

BRONCHORRHŒA, (F.) *Bronchorrhée*, *Catarrhe pituitæ*, *Phlegmorragie pulmonaire*, *Flux bronchique*, from *βρογχος*, 'bronchus,' and *ρρω*, 'I flow.' An increased secretion of mucus from the air passages, accompanied or not by inflammation, — a gleet, as it were, of the pulmonary mucous membrane. When excessive, it may constitute *Phthisis pituitosa seu mucosa*, *Mycophthisis*.

BRONCHORRŒA ACUTA, Bronchitis (chronic).

BRONCHOSTASIS, Bronchitis.

BRONCHOTOME, *Bronchotomus*, from *βρογχος*, and *τεμνω* 'to cut.' A kind of lancet, with a blunt and rounded point, mounted on a handle, and fitted to a canula, which passes in along with it, and is allowed to remain in the opening made in the trachea.

BRONCHOTOMY, *Bronchotomia*, (F.) *Bronchotomie*. Same etymology. A surgical operation, which consists in making an opening either into the trachea, (*Tracheotomy*;) into the larynx, (*Laryngotomy*;) or into both, (*Trachea-laryngotomy*;) to extract foreign bodies or to permit the passage of air to the lungs. These different parts are divided transversely or vertically, according to circumstances.

BRONCHO-VESICULAR RESPIRATION, see Respiration.

BRONCHUS, see Bronchia, Trachea.

BROOKLINE, *Veronica beccabunga*.

BROOM, *Sophora tinctoria*, *Spartium scoparium*—b. Butcher's, *Ruscus*—b. Clover, *Sophora tinctoria*—b. Indigo, *Sophora tinctoria*—b. Rape, of Virginia, *Orobanche Virginiana*—b. Spanish, *Spartium junceum*—b. Yellow, *Sophora tinctoria*.

BROSSARDIÈRE, MINERAL WATERS OF. Brossardière is a chateau in Bas-Poitou, France. The waters contain carbonates of iron and lime, chloride of sodium, and sulphate of lime. They are aperient.

BROSSE, Brush.

BROTH, CHICKEN, see Chicken Broth.

BROTH, VEGETABLE. Take two *potatoes*, a *carrot*, and an *onion*, all cut fine; boil in a quart of water for an hour, adding more water from time to time, so as to keep the original quantity; flavour with *salt*, and a small quantity of *potterbs*; strain. A little mushroom catchup improves the flavour.

BROTHER, UTERINE, see Uterine.

BROUILLARD, Caligo.

BROUSSAIST. One who is a believer in, and professor of, the physiological and pathological opinions of Broussais. The system itself was called Broussaism, or the *Physiological Doctrine*.

BROW, Front—b. Ague, Neuralgia frontalis.

BROWN RED, Colcothar.

BROWNIAN, *Brownian*, *Brownian*. Relating to the system or opinions of John Brown.

BROWNIISM, *Brownism*, *Brownianism*. The doctrines of Brown.

BROWNIST, *Brownian*, *Brownian*. A follower of the system of Brown.

BRUCEA ANTI-DYSENTERICA. Called after Bruce, the Abyssinian traveller. *B. ferruginea*, *Angustura spuria*, (F.) *Fausse Angusture*, *A. Ferrugineuse*. The systematic name of the plant whence was obtained—it was supposed—false *Angustura* or false *Cusparia Bark*. It is really the bark of *Strychnos nux vomica*.

BRUCIA, *Brucine*.

BRUCINE, *Brucia*, *Bruci'na*, *Bruci'num*, *Brucium*, *Pseudangusturi'num*, *Caniram'mum*, *Caniram'mum*, *Vom'icine*, *Angusturine*. An organic, salifiable base, discovered in the false *angustura*—*Brucea anti-dysenterica*, and obtained from *Strychnos nux vomica*. It is of a pearly white; crystallizes in oblique prisms with a parallelogrammatic base; is very bitter, slightly acid and styptic, and soluble in water, but more so in alcohol. *Brucia* is a less active poison than *strychnia*. It resembles it, however, and may be used as a substitute for it and for the extract of *nux vomica*. Dose, half a grain.

BRUCKENAU, MINERAL WATERS OF. These springs are in Bavaria, and contain carbonic acid and iron.

BRUCOURT, MINERAL WATERS OF. Brucourt is three leagues and a half from Caen, in Normandy. The waters contain carbonic acid, chloride of sodium, and sulphate of soda, much sulphate of lime, &c.

BRUICK, *Furunculus*.

BRUISE, Contusion.

BRUISE-ROOT, *Stylophorum diphyllum*.

BRUISEWORT, *Bellis saponaria*.

BRUISSEMENT (F.), *Frem'itus*. This word has much the same signification as *Bourdonnement*, as well as *Bruit*.

BRUIT (F.), 'sound.' A French term, applied to various sounds heard on percussion and auscultation, viz.:

BRUIT DE CRAQUEMENT, *B. de Tiraillement*, *Bruit de cuir neuf*, *Cri de cuir*, 'sound of crackling, or bursting, or of new leather.' A sound produced by the friction of the *pericardium*, when dried and roughened by inflammation.

BRUIT DU CŒUR FŒTAL, *Battemens doubles*. *Double bruit du Cœur du Fœtus*. The pulsations of the foetal heart heard in auscultation in the latter half of utero-gestation.

BRUIT DE OUIR NEUF, *Bruit de craquement*.

BRUIT DE DIABLE, *Roulement du Diable*, *Bruit de soufflé à double courant*, 'noise of the diable, or humming-top.' *Venous hum*. A high degree of *Bruit de soufflet*, heard on auscultating the arteries or veins—probably the latter—of the neck, in chlorosis. It denotes an impoverished state of the blood.

BRUIT DOUBLE DU CŒUR DU FŒTUS, *Bruit du Cœur fœtal*—*b. d'Expansion pulmonaire*, see *Murmur*, respiratory—*b. de Frôlement*, see *Frôlement*—*b. de Froissement pulmonaire*, see *Froissement pulmonaire*—*b. de Frôlement péricardique*, see *Frôlement péricardique*.

BRUIT DE FROTTEMENT ASCENDANT ET DESCENDANT, 'sound of friction of ascent and descent.' Sounds produced by the rubbing of the lung against the parietes of the chest, as it rises and falls during inspiration and expiration. They are distinctly heard in pleuritis, when the pleura has become roughened by the disease. *Friction sounds*, *Rubbing sounds*, *To-and-fro sounds* are also heard in pericarditis and peritonitis.

BRUIT HUMORIQUE, *B. Hydropneumatique*. The sound afforded on percussion when organs are filled with liquid and air.

BRUIT HYDROPNEUMATIQUE, *Bruit humorique*—*b. de Jappement*, see *Sifflement modulé*—*b. de Lime à bois*, see *Bruit de Scie*.

BRUIT DE MOUCHE (F.), 'fly sound.' A sound analogous to the *Bruit de diable*—so called from its likeness to the buzzing of a fly—heard on auscultating the neck in chlorotic cases.

BRUIT MUSCULAIRE. The sound accompanying the first sound of the heart, referred by some to muscular contraction. Called, also, *Bruit rotatoire*, in consequence of its having been thought to resemble the rumbling of distant wheels.

BRUIT MUSCAL, *Sifflement modulé*.

BRUIT DE PARCHEMIN, 'parchment tone.' A sound as if produced by two sheets of parchment applied to each other. It is said to be produced by thickening and rigidity of the valves of the heart.

BRUIT DE PAULEMENT, see *Sifflement modulé*.

BRUIT PLACENTAIRE, *B. de souffle placentaire* ou *utérin*, *Souffle utérin* ou *placentaire*, *Placental bellows' sound*, *Utero-placental murmur*, *Uterine murmur*. The bellows' sound heard on auscultating over the site of the placenta in a pregnant female. It does not appear to be owing to the placental vessels; but to the uterine tumour pressing upon the large vessels of the mother.

BRUIT DE POT FÉLÉ, 'sound of a cracked vessel.' *Cracked pot sound*. A sound heard on percussion, when a cavern in the lungs is filled with air, and has a narrow outlet. It is not diagnostic, however.

BRUIT DE RACLEMENT, 'sound of scraping.' A sound produced by the scraping of hard, solid membranes, as the pericardium, against each other, very analogous to *Bruit de craquement*.

BRUIT DE RAPE, 'sound of a rasp.' A sound heard during the contraction of either the auricles or ventricles. It is constant; and the contraction of the cavity is more prolonged than natural, and emits a hard, rough, and—as it were—stiff sound.

It indicates contraction of the valvular orifices by cartilaginous deposits, or ossification, and is better heard near the apex of the heart, if the auriculo-ventricular valves be concerned,—near the base, if the semilunar valves be the seat of the disease.

BRUIT RESPIRATOIRE, *Murmur*, respiratory.

BRUIT ROTATOIRE, *Bruit musculaire*.

BRUIT DE SCIE, or 'saw sound,' and **BRUIT DE LIME À BOIS**, or 'file sound,' resemble the *Bruit de Râpe*.

BRUIT DE SOUFFLE À DOUBLE COURANT, *Bruit de Diable*.

BRUIT DE SOUFFLET, *Bruit de Souffle*, 'bellows' sound,' 'blowing sound.' A sound like that of a bellows, heard occasionally by the ear applied to the chest during the contraction of the ventricles, auricles, or large arteries. It coexists with affections of the heart, but is heard, also, without any disease of that organ,—whenever, indeed, an artery is compressed. An *Encephalic bellows' sound*, (F.) *Bruit de soufflé céphalique*, has been described by Drs. Fisher and Whitney. It is heard on applying the ear to the occiput or to the top of the head; and is considered to indicate turgescence of vessels, or inflammation. When such turgescence exists, the vessels are compressed, and the compression gives rise to the sound in question. [?]

BRUIT DE SOUFFLE CÉPHALIQUE, see *Bruit de soufflet*—*b. de Souffle placentaire*, *Bruit placentaire*—*b. de Souffle utérin*, *Bruit placentaire*—*b. de Tiraillement*, *Bruit de craquement*.

BRUIT DE SOUPAPE, 'valvular or flapping noise.' A sound heard in respiration, when a foreign body is in the air passages. It somewhat resembles the flapping of a valve; hence its French name.

BRUIT DE TAFFETAS, 'sound of taf-feta.' '*Sarcenet sound*.' A respiratory sound, so named, by M. Grisolles, from its resembling the sound caused by the tearing of a piece of taffeta; and which he considers to indicate hepatization of the lung, limited to the surface, in pneumonia.

BRUIT TYMPANIQUE, 'tympanic sound.' The clear sound afforded by percussing the stomach and intestines when containing air.

BRUIT UTÉRIN, *B. placentaire*.

BRUITS DU CŒUR, see *Heart*.

BRUK, *Furunculus*.

BRULURE, *Burn*.

BRUNELLE, *Prunella*.

BRUNNER'S GLANDS, *Brunneri Glan'dulae*, *G. Brunneri* *næ seu solitariae*, *Solitary glands or follicles*, *Second pancreaticas*. Compound muciparous follicles, seated between the mucous and muscular coats of the stomach, along the two curvatures of that organ, and in the duodenum; so called from their discovery having been generally attributed to Brunner. The solitary intestinal follicles are often known, at the present day, as the glands of Brunner, although Brunner restricted the latter term to the glands of the duodenum.

BRUNONIAN, *Brownian*.

BRUNONIANISM, *Brownism*.

BRUNUS, *Erysipelas*.

BRUCUS, *Ruscus*.

BRUSH, *Scopula*, (F.) *Brosse*. A well-known instrument, used in medicine chiefly for the following purposes:—1. To clean the teeth. 2. To remove the saw-dust which adheres to the teeth of the trephine, during the operation of trephining. 3. To rub the surface of the body, for the purpose of exciting the skin, and favouring transpiration. Westring, a Swedish physician, has

recommended metallic brushes for the purpose of conveying galvanism to a part. These brushes consist of a plate of ebony fitted to another of gold, in which threads of the same metal are fixed;—the brush being connected with one of the poles of the galvanic pile.

BRUSH, STOMACH, *Excusia ventriculi*.

BRUTA, *Juniperus sabina*.

BRUTIA. A sort of thick pitch, obtained from Brutia, in Italy. From *Piz Brutia* was obtained the *Oleum Picinum*.

BRUTIA, *Instinet*.

BRUTINO, *Terebinthina*.

BRUTOLÉ, see *Cerevisia*.

BRUXANELLI. A Malabar tree, the bark and leaves of which have a strong smell, and are astringent. On the coast of Malabar, its juice, mixed with butter, is applied to boils. Its bark is esteemed to be diuretic, and its roots antihirritic.

BRUYÈRE VULGAIRE, *Erica vulgaris*.

BRUYÈRES, MINERAL WATERS OF. Bruyères is a small village. $7\frac{1}{2}$ leagues from Luneville. The waters are acidulous and chalybeate.

BRYCETOS, see *Algidus*.

BRYCHETHMOS, *Rugitus*.

BRYCHETOS, see *Algidus*.

BRYGMA, *Brygnus*, *Trisita*, *Prisis*, *Prismus*, *Odontopri'sis*, *Stridor Dentium*, (F.) *Grincement des Dents*. Grinding of the teeth. A common symptom, in children, of gastric or other derangement, but often present when there is no reason to suspect any.

BRYONE, *Bryonia*—*b. d'Amérique*, *Convolvulus Mechoacan*.

BRYONIA AFRICANA, *African Bryony*, from *βρω*, 'I bud forth.' A South African plant, common amongst the Hottentots, which, in the form of decoction, acts simultaneously as an emetic, cathartic, and diuretic. It is used by the natives in cutaneous diseases, dropsy, and syphilis. The tincture is a powerful emetic and cathartic.—Thunberg.

BRYONIA ALBA, *White Bryony*, *Vitis alba sylvestris*, *Agrastis*, *Agriampelos*, *Am'pelos d'grin*, *Echetro'sis*, *Bryonia aspera* seu *Dioica*, *Cedros'tis*, *Chelido'nium*, *Labrus'ca*, *Mol'thrum*, *Ophrostaph'ylon*, *Pailo'thrum*. Ord. *Cueurbitaceæ*. Sex. Syst. Monoclea Monadelphia. (F.) *Conteu'rée*, *Vigne vierge*, *V. blanche*, *Navet du diable* ou *galant*. The root is large and succulent, and has an acrid, bitter, and disagreeable taste. It is a drastic cathartic. Externally, it has been applied, in form of cataplasm, in gout. When repeatedly washed, a good starch is obtained from it. The active principle has been separated from it, and called *Bry'online*.

BRYONIA MECOACANNA NIGRICANS, *Convolvulus jalapa*—*b. Nigra*, *Tamus communis*—*b. Peruviana*, *Convolvulus jalapa*.

BRYONINE, see *Bryonia alba*.

BRYONY, BLACK, *Tamus communis*—*b. White*, *Bryonia alba*—*b. Wild*, *Sycios angulatus*.

BRYTIA, *Marc* of grapes.

BRYTOLATURE, see *Cerevisia*.

DRYTOLÉ, see *Cerevisia*.

BRYTON, *Cerevisia*.

BU, *βου*, abbreviation of *βους*, 'an ox,' in composition expresses 'excess, greatness.' Hence *Bulimus*, *Buphthalmia*, &c.

BUBASTECORDIUM, *Artemisia vulgaris*.

BUBE, *Pustule*.

BUBO, *βουβων*, *Pano'chia*, *Panus inguinalis*, *Atenoph'nia inguinalis*, *Bubonopa'nus*, *Bubon'oncus*, *Bubon'cus*, *Cambu'ca*, *Angus*, *Boubon*, *Culuce'le*, *Culose'le*, (F.) *Bubon*, *Poulain*. In the works of Hippocrates and Galen, this word

sometimes signifies the groin—*Inguen*; at others, the inguinal glands; and at others, again, swelling or inflammation of these parts. The moderns apply the term to an inflammatory tumour seated in the groin or axilla, and they generally distinguish—1. *Simple* or *Sympathetic Bubo*, which is independent of any virus in the economy. 2. *Venereal Bubo*, (F.) *Bubon vénérien*, which is occasioned by the venereal virus. 3. *Pestilential Bubo*, or *B. symptomatic of the Plague*. The last two have by some been called malignant *Bubo*, (F.) *Bubon malin*.

Primary *Bubo*, (F.) *Bubon primitif*, shows itself with the first symptoms of syphilis: the consecutive not till afterwards.

BUBON, *Bubo*, *Inguen*—*b. Gummiiferum*, see *Ammoniac gum*.

BUBON D'ENBLÉE (F.). An enlargement and suppuration of one or more of the inguinal glands, not preceded by any other of the more common forms of venereal disease, nor by any other syphilitic symptom.

BUBON GALBANUM. The systematic name of a plant which has been supposed to afford galbanum; *Meo'pion*, *Mato'rium*. The plant is also called *Fer'ula Africa'na*; *Oreoseli'num Africa'num*, *Ani'sum frutico'sum galbanif'erum* seu *Africa'num frutes'cens*, *Seli'num Galbanum*, *Agayl'lis galbanum*, *The long-leaved* or *longue-leaved Galbanum*. Ord. *Umbelliferae*. The plant can scarcely, however, be considered to be determined. By the Dublin College, it is referred to *Opo'dia Galbanif'era*, Ord. *Umbelliferae*. Galbanum is the gummi-resinous juice. Its odour is fetid, and taste bitter and acrid: the agglutinated tears are of a white colour, on a ground of reddish-brown. It forms an emulsion when triturated with water, and is soluble in proof spirits of wine, and vinegar: sp. gr. 1.212. It has been given as an antispasmodic and expectorant, in pill or emulsion. Dose, from gr. 10 to 60. Externally, it is applied as a cataplasm.

Bubon galbanum is a South African plant; and is reputed to be an excellent diuretic, under the name of *Wild Celery*. A decoction of the leaves is given in dropsy and gravel. According to Pappe, the resinous matter, which exudes from the stem, differs in appearance, smell, and in every respect, from *Gummi Galbanum*.

BUBON MACEDONICUM, *Athaman'ta Macedonica*, *Petroseli'num Macedonicum*, *A'pium petr'a'num*, *Petra'pium*, (F.) *Pera'il de Macédoine*, *Macedonian Parsley*. Its properties are similar to those of common parsley, but weaker and less grateful. The seeds are an ingredient in the celebrated compounds, *Mithridate* and *Theriac*.

BUBONA, *Nipple*.

BUBONALGIA, from *βουβων*, 'the groin,' and *algos*, 'pain.' Pain in the groin.

BUBONCUS, *Bubo*.

BUBONIUM, *Aster At'ticus*, *Golden Starwort*. A plant anciently supposed to be efficacious in diseases of the groin, from *βουβων*, 'the groin.'

BUBONOCELÉ, from *βουβων*, 'the groin,' and *κηλη*, 'tumour,' 'rupture.' *Hernia inguinalis*, *In'guinal Hernia*, *Rupture of the Groin*. (F.) *Hernie inguinale*. Some surgeons have confined this term to hernia when limited to the groin, and have called the same affection, when it has descended to the scrotum, *Oscuto'cele*, *Scrotal Hernia*. The rupture passes through the abdominal ring: and, in consequence of the greater size of the opening in the male, it is more frequent in the male sex.

BUBONONCUS, *Bubo*.

BUBONOPANUS, *Bubo*.

BUBONOREX'IS, from *βουβων*, 'the groin,' and *ρηξις*, 'a rupture.' A name given to *bubonocèle* when accompanied with a division of the pe-

ritoneum, or when, in other words, it is devoid of a sac.

BUBON'ULUS, *Bubon'culus*. A diminutive of *Bubo*. A painful swelling of the lymphatics of the penis, extending along the dorsum of that organ to the groin. It is an occasional accompaniment of gonorrhœa.

BUBUKLE. A word used by Shakspeare for a red pimple on the nose.

BUBUNCULUS, *Bubonulus*.

BUCAROS, Terra Portugallica.

BUCCA, *Gnathos*. The mouth. The cheek and hollow of the cheek. Also, the vulva.

BUCCAC'RATON, from *Bucca*, and *κρω*, 'I mix.' A morsel of bread sopped in wine, which served of old for a breakfast.—Linden.

BUCCAL, *Bucca'lis*, from *Bucca*, 'the mouth,' or rather 'the cheek.' That which concerns the mouth, and especially the cheek.

BUCCAL ARTERY, *Artère Sus-maxillaire*, (Ch.) arises from the internal maxillary or from some of its branches, as the *Temporalis profunda antica*, or the *Alveolar*. It distributes its branches to the buccinator muscle, and to the buccal membrane.

BUCCAL GLANDS, *Molar Glands*. Mucous follicles, seated in the buccal membrano, opposite the molar teeth. They secrete a viscid humour, which mixes with the saliva, and lubricates the mouth.

BUCCAL MEMBRANE, (F.) *Membrane Buccale*. The mucous membrane, which lines the interior of the mouth.

BUCCAL NERVE, *Buccinator Nerve*, *Buccolabial*—(Ch.), is given off by the inferior maxillary. It sends its branches to the cheek, and especially to the buccinator muscle.

BUCCAL VEIN follows the artery.

BUC'CEA, *Buccel'la*. The fleshy excrescence of nasal polypus, so called because it was believed to proceed from the mouth.—Paracelsus. Also, a mouthful.

BUCCELA'TON, *Buccela'tus*. A loaf-shaped cathartic medicine; made chiefly of scammony. —Aëti, Paulus of Ægina.

BUCCELLA, *Bolus*, *Buccæa*.

BUCCELLA'TIO. A mode of arresting hemorrhage, by applying a pledge of lint to the bleeding vessel.—Avicenna, Fallopius.

BUCCINA, Turbinated bones.

BUCCINA'TOR, from *buccinare*, 'to sound the trumpet.' The *Buccinator Muscle*, *Retractor Anguli Oris*, *Bucco-Alvéolo-maxillare*, *Alvéolo-labial*—(Ch.), *Masoe'rius*, is situate in the substance of the cheeks. It extends between the posterior portions of the alveolar arches of the two jaws and the commissure of the lips, which it draws backward. It assists in mastication, by pushing the food back towards the teeth; and, if the cheeks be distended by air, its contraction forces it out.

BUCCO. One who is blub-cheeked, or wide-mouthed. In composition, the mouth.

BUCCO-ALVÉOLO-MAXILLAIRE, *Buccinator*.

BUCCO-LABIAL NERVE, *Buccal nervé*.

BUCCO-PHARYNGE'AL, *Bucco-Pharynge'us*, (F.) *Bucco-Pharyngien*. Belonging to the mouth and pharynx. The *Bucco-pharynge'al Aponeuro'sis* or *Intermaxillary Ligament*, extends from the internal ala of the pterygoid process to the posterior part of the lower alveolar arch, and affords attachment, anteriorly, to the buccinator, and, posteriorly, to the constrictor pharyngis superior.

BUC'ULA, from *Bucca*, 'the mouth.' A small mouth. The fleshy part beneath the chin.—Bartholine.

BU CERAS, *Trigonella fœnum*—b. *Fœnum Græcum*, *Trigonella fœnum Græcum*.

BUCHU, *Diosma crenata*—b. *Leaves*, *Diosma crenata*.

BUCKBEAN, *Menyanthes trifoliata*—b. *American*, *Menyanthes verna*.

BUCKBERRY, *Vaccinium stamineum*.

BUCKET FEVER, *Dengue*.

BUCKEYE, *Æsculus hippocastanum*.

BUCKHO, *Diosma crenata*.

BUCKTHORN, *PURGING*, *Rhamnus*.

BUCKU, *Diosma crenata*.

BUCKWHEAT, *Polygonum fagopyrum*—b. *Plant*, eastern, *Polygonum divaricatum*.

BU CNEMIA. see *Elephantiasis*—b. *Tropica*, see *Elephantiasis*.

BUCTION, *Hymen*.

BUFF, *INFLAMMATORY*, *Corium phlogisticum*.

BUFFY COAT, *Corium phlogisticum*.

BUG, (BED,) *Cimex*.

BUGANTIA, *Chilblain*.

BUG'GERY, *Sod'omy*, *Sodom'ia*, *Co'itus Sodomiticus*; (I.) *Bugarone*. Said to have been introduced by the Bulgarians; hence the name. A carnal copulation against nature, as of a man or woman with any animal; or of a man with a man, or a man unaturally with a woman. The unnatural crime.

BUGLE, *Prunella*—b. *Common*, *Ajuga reptans*—b. *Pyramidalis*, *Ajuga*—b. *Reptans*, *Ajuga reptans*—b. *Water*, *Lycopus Virginicus*—b. *Weed*, *Lycopus*.

BUGLOSE, *Anchusa officinalis*.

BUGLOSS, *DYER'S*, *Anchusa tinctoria*—b. *Garden*, *Anchusa officinalis*—b. *Upright*, *Ajuga*.

BUGLOSSA, *Anchusa officinalis*.

BUGLOSSUM ANGUSTIFOLIUM MAJUS, *Anchusa officinalis*—b. *Latifolium*, *Borago officinalis*—b. *Sativum*, *Anchusa officinalis*—b. *Sylvestris*, *Anchusa officinalis*—b. *Tincturum*, *Anchusa tinctoria*—b. *Verum*, *Boracic acid*—b. *Vulgar majus*, *Anchusa officinalis*.

BUGRANDE ÉPINEUSE, *Ononis spinosa*.

BUGRANE, *Ononis spinosa*—b. *des Champs*, *Ononis arvensis*.

BUGLA, *Ajuga*—b. *Chamæpitys*, *Teucrium chamæpitys*—b. *Pyramidalis*, *Ajuga*—b. *Reptans*, *Ajuga reptans*.

BUIS, *Buxus*.

BUISARD, *MINERAL WATERS OF*. Baisard is two leagues from Chateau-Thierry, in France. The water contains chloride of calcium and carbonate of lime.

BULB, *Bulbus*, (F.) *Bulbe*. A name, given by anatomists to different parts which resemble, in shape, certain bulbous roots. The *Bulb of the Aorta* is the great sinus of the Aorta. *Bulb of a Tooth*; the vascular and nervous papilla contained in the cavity of a tooth. The *Bulb* or *Root of the Hair* is the part whence the hair originates. The *Bulb of the Urethra* is the dilated portion formed by the commencement of the *Corpus spongiosum* towards the root of the penis. We say, also, *Bulb*, for *Globe*, of the eye.

BULB OF THE EYE, see *Eye*—b. of the Female, *Bulbus vestibuli*—b. *Rachidian*, see *Medulla oblongata*.

BULBE, *Bulb*—b. *Rachidian*, see *Medulla oblongata*—b. *du Vagin*, *Bulbus vestibuli*—b. *de la Voûte à trois Piliers*, Mamillary tubercles.

BULBI FORNICIS, Mamillary tubercles—b. *Priorum Crurum Fornicis*, Mamillary tubercles.

BULBOCASTANEUM, *Bunium bulbocastanum*.

BULBO-CAVERNEUX, *Accelerator urinæ*.

BULBO-CAVERNOSUS, *Accelerator urinæ*—b. *Synoesmo-cavernæ*, *Accelerator urinæ*—b. *Uréthral*, *Accelerator urinæ*.

BULBOCODIUM, *Narcissus pseudonarcissus*.

BULBONACH, *Lunaria rediviva*.

BULBUS, Bulb.

BULBUS ESCULENTUS. The *Es'culent Bulb*: a particular kind, so denominated by the ancients. It is supposed to have been the *Cepa Ascalonica*.—Dioscorides, Celsus, Pliny, &c.

BULBUS GLANDULOSUS, Proventriculus—b. Medullæ spinalis, Medulla oblongata—b. Oculi, see Eye—b. Olfactorius, see Olfactory Nerves—b. Pili, see Hair—b. Rachidicus, see Medulla oblongata—b. Vaginae, B. vestibuli.

BULBUS VESTIBULI, *B. Vag'inæ*, *Plexus retiformis* seu *reticularis* seu *cavernosus*, *Crura clitoridis interna*, Bulb or Semi-bulb of the Female, (F.) *Bulbe du Vagin*. A close-packed plexus of intricately anastomosing veins, inclosed in a fibrous investment,—being an immediate continuation and extension of the *pars intermedia*, and occupying the space between the beginning or vestibule of the vagina and the rami of the pubic arch on each side. It is regarded by Lauth, Taylor, Morgagni and Kobelt as the analogue of the male bulb.

BULBUS VOMITORIIUS. A plant, said by Dioscorides to be emetic and diuretic. It is the *Musk-grape flower*, according to Ray,—the *Hycinchus Muscari*.

BULESIS, Voluntas.

BULGA, Vulva.

BULIMIA, Boulimia.

BULLITHOS, from *bovs*, 'an ox,' and *lithos*, 'a stone.' A bezoar or stone, found in the kidneys, gall-bladder, or urinary bladder of an ox or cow.

BULLA, (F.) *Bulle*. A *Bleb*, (Sc.) *Bleib*. A portion of the cuticle, detached from the skin by the interposition of a transparent, watery fluid. It forms the 4th order in Willan's and Bateman's arrangement of cutaneous diseases, and includes erysipelas, pemphigus, and pompholyx. By some, Bulla has been used synonymously with *Pemphigus*. See, also, Hydatid.

BULLACE PLUM, Prunus in Sitia.

BULLÆ ROTUNDÆ CERVICIS UTERI, Nabothi glandulæ.

BULLDOG, Tabanus.

BULL-FISTS, Lycoperdon.

BULLOUS, *Bulla'sus*, (F.) *Bulleux*. Having relation to a bulla or bleb, as a 'bullous eruption.' Pemphigus has been designated *Maladie bulleuse*, and *Fievre bulleuse*, when accompanied by fever.

BULLSEGG, *Typha latifolia*.

BUMBLEKITES, see *Rubus fruticosus*.

BUMELLIA, *Fraxinus excelsior*.

BUM-GUT, Rectum.

BUNA, *Coffea Arabica*.

BUNCOMBE, WARM AND HOT-SPRINGS OF. See North Carolina, Mineral Waters of.

BUNDURH, *Corylus avellana*.

BUNEWAND, *Heracleum spondylium*.

BUNIAS, *Brassica napus*.

BUNIOID, *Napiform*.

BUNION, Bunyon.

BUNITES VINUM. A wine, made by infusing the *Bunium* in must. It is stomachic, but scarcely ever used.

BUNION CARVI, Carum.

BUNION BULBOCASTANUM, *Bouvion*, so called, it has been supposed, from growing on hills, from *bovos*, 'a hill,' or from the tuberosity of its root. *B. minus*, *Bulbocastanum*, *Sium bulbocastanum*, *Scander bulbocastanum*, *Carum bulbocastanum*. Order, Umbellifere. The systematic name of a plant, whose root is called *Pig-nut*, *Agriocastanum*, *Nucula terrestris*, *Bulbocastanum majus et minus*, *Earth-nut*, *Hawk-nut*, *Kipper-nut*, (Sc.) *Arnut*, (F.) *Terre-noix*. The root is tuberosus, and is eaten raw or roasted. It has been sup-

posed to be of use in strangury. It is not employed in medicine.

BUNNIAN, Bunyon.

BUNWEED, Senecio Jacobæa.

BUN'YON, *Bun'ion*, *Bun'nica*, from *bovos*, 'an eminence.' [?] An enlargement and inflammation of the bursa mucosa at the inside of the ball of the great toe.

BUOPHTHALMIA, Buphthalmia.

BUPEINA, Boulimia.

BUPHTHALMI HERBA, *Anthemis tinctoria*.

BUPHTHAL'MIA, *Buphthal'mia*, *Buphthal'mos*, *Elephantom'ma*, from *bovs*, 'an ox,' and *ophthalmos*, 'an eye.' *Ox-eye*. Under this name, the generality of authors have designated the first stage of hydrophthalmia. Others, with Sabatier, mean by it, turgescence of the vitreous humour, which, by pushing the iris forwards, forms around the crystalline a sort of border.

BUPHTHALMUM CRETICUM, *Anthemis Pyrethrum*—b. *Majus*, *Chrysanthemum leucanthemum*.

BUPHTHALMUS, Hydrophthalmia, *Semper-vivum tectorum*.

BUPINA, Boulimia.

BUPLEUROIDES, *Bupleurum rotundifolium*.

BUPLEURUM ROTUNDIFOLIUM, *Bupleu'ron*, *Bupleuroi'des*, from *bov*, augmentative, and *πλευρον*, 'side,' (F.) *Buplèvre*, *Percefeuille*, *Round-leaved Hare's Ear*, *Thorowax*. Order, Umbellifere. The herb and seeds are slightly aromatic. It was formerly celebrated for curing ruptures, being made into a cataplasm with wine and oatmeal.

BUPLÈVRE, *Bupleurum rotundifolium*.

BURAC, Borax. Also, any kind of salt. (Arabic.)

BURBOT, see *Oleum Jacoris Aselli*.

BURDOCK, *Arctium lappa*—b. Lesser, *Xanthium*—b. Prairie, *Silphium terebinthaceum*.

BURIAL ALIVE, Zoothapsis.

BURIS, Hernia, accompanied by scirrhus tumefaction; or, perhaps, a scirrhus tumour only.—*Avicenna*.

BURN. Sax. *beþnan* or *þýpan*, 'to burn or burn.' *Us'tio*, *Ambus'tio*, *Adu's'tio*, *Tres's Causis*, *Eryth'ma Ambus'tio*, *Causis*, *Encau'sis*, *Pyri-caus'tum*, *Combustu'ra*, *Catacau'ma*, *Combustio*, (F.) *Brûlure*. An injury produced by the action of too great heat on the body. Burns are of greater or less extent, from the simple irritation of the integument to the complete destruction of the part. The consequences are more or less severe, according to the extent of injury, and the part affected. Burns of the abdomen, when apparently doing well, are sometimes followed by fatal results. Their treatment varies,—at times, the antiphlogistic being required; at others, one more stimulating.

BURNEA, see *Pinus Sylvestris*.

BURNET, CANADA, *Sanguisorba Canadensis*.

BURNETT'S DISINFECTING LIQUID or FLUID. A solution of chloride of zinc, first used by Sir William Burnett for preserving timber, canvass, &c., from dry rot, mildew, &c., and afterwards as an antiseptic and antiseptic, especially in the case of dead bodies. The Dublin Pharmacopœia has a *Zinci Chloridi Liquor*, *Solution of Chloride of Zinc*, which is not near so strong as Sir William's preparation.

BURNING, *Brenning*. A disease mentioned by old historians, from which authors have unsuccessfully endeavoured to demonstrate the antiquity of syphilis.—*Parr*.

BURNING OF THE FEET, see *Feet*, burning of the.

BURNT HOLES. A variety of *rupia*, popularly known in Ireland under this name; and not

unfrequent there amongst the ill-fed children of the poor. See *Rupia escharotica*.

BURR, Rotacismus.

BURRAGE, *Borago officinalis*.

BUR-REED, GREAT, *Sparganium ramosum*.

BURRH, Rotacismus. •

BURRHI SPIRITUS MATRICALIS. The Spirit of *Burhus* for diseases of the Womb. It is prepared by digesting, in alcohol, equal parts of myrrh, oilibanum, and mastic. Boerhaave frequently prescribed it.

BURSA CORDIS, Pericardium—b. Omentalis, see Epiploon, gastro-hepatic—b. Pastoris, Thlaspi bursa—b. Testium, Scrotum—b. Virilis, Scrotum.

BURSÆ MUCILAGINOSÆ, B. mucosæ.

BURSÆ MUCOSÆ, B. mucosæ vesiculæ res, Bursæ seu Capsulæ synoviales, *Blennocystides*, *Sacci mucosi*, *Vesicæ unguinalæ tendinum*, *Vaginæ Synoviales* seu *mucilaginosæ*, *Synovial Crypts* or *Follicles*, (F.) *Bourses Synoviales* ou *muqueuses* ou *mucilagineuses*. Small membranous sacs, (from *bursa*, 'a bag, a purse,') situate about the joints, particularly about the large ones of the upper and lower extremities, and, for the most part, lying under the tendons. They are naturally filled with an oily kind of fluid, the use of which is to lubricate surfaces over which the tendons play. In consequence of bruises or sprains, this fluid sometimes collects to a great extent. The bursæ are, generally, either of a roundish or oval form, and they have been arranged under two classes, the spherical and the vaginal.

BURSÆ SYNOVIALES, Bursæ mucosæ.

BURSAL, *Bursalis*. Relating or appertaining to bursæ—as a 'bursal tumour.'

BURSALIS, Obturator internus.

BURSERIA ACUMINATA, B. gummifera.

BURSERIA GUMMIFERA, B. acuminata, *Terebinthus gummifera*, *Jamaica Bark Tree*. Order, *Terebinthaceæ*. A resin exudes from this tree, which, as met with in the shops, is solid externally; softish internally; of a vitreous fracture; transparent; of a pale yellow colour; turpentine smell, and sweet, perfumed taste. It has been used like balsams and turpentine in general, and is called, by the French, *Cachibou*, *Chibou*, and *Résine de Gomart*.

BURSITIS, see Synovitis.

BURST, Hernia; Hernial.

BURSTEN, see Hernial.

BURSULA, Scrotum.

BURTHISTLE, *Xanthium*.

BURTREE, *Sambucus*.

BURUNHEM, *Monesia*.

BURWEED, *Xanthium*.

BURWORT, *Ranunculus acris*.

BUSH, JEW, *Pedilanthus tithymaloides*.

BUSSANG, MINERAL WATERS OF. *Bussang* is a village in the department of Vosges, France. The waters are acidulous chalybeates.

BUSSEROLLE, *Arbutus uva ursi*.

BUSII SPIRITUS BEZOARTICUS, *Bezoardic Spirit of Bussius*. A preparation, regarded as sudorific, diuretic, and antispasmodic; obtained by distilling subcarbonate and muriate of ammonia, amber, oil of cedar or juniper, &c.

BUTE, ISLAND OF, CLIMATE OF. This island is in the Frith of Clyde, about 18 miles below Greenock. The climate is mild and equable, but rather moist; and, as a winter residence, it holds out advantages for those only that appear to demand such a condition of the atmosphere. The climate resembles, in character, that of the S. W. of England and France, and the Channel islands; although its temperature is lower.

BUTEA FRONDO'SA, *Erythrina monosperma*.

ma, *Rudolphia frondosa*, see *Kino*. A tree, common in Bengal, and in the mountainous parts of India; *Nat. Ord.* *Leguminosæ*; from which gum butea flows. Dr. Pereira found this gum to be identical with a specimen marked *gummi rubrum astrigenum*—the *gomme astrigente* de *Gambie* of M. Guibourt. By some, this gum has been confounded with kino.

BUTIGO, Gutta rosea.

BUTOMON, *Iris pseudacorus*.

BUTTER, from *butropov*; itself from *βους*, 'ox,' and *τεπος*, 'any thing coagulated.' *Butyrum*, *Pice'rium*, (F.) *Beurre*. A sort of concrete oil, obtained from the cream that forms on the surface of the milk furnished by the females of the mammalia, especially by the cow and the goat. Fresh butter is very nutritious, whilst the rancid is irritating. The ancient chemists gave the name *Butter* to many of the metallic chlorides. It has also been applied to vegetable substances, which resemble, in some respects, the butter obtained from milk. The essential fatty matter in it is *butyryn* or *butyrate of glycerin*.

BUTTER OF ANTIMONY, *Antimonium muriatum*.

BUTTER OF BAMBOOC or BAMBOC, (F.) *Beurre de Bambouc* ou *Bambuk*. A vegetable oil obtained from a species of almond, and used in Senegal in neuralgic and rheumatismal pains.

BUTTER OF CA'CAO, Oil of *Ca'cao*, *Oleum Cacao spissatum*, O. *Theobroma cacao expressum*, (F.) *Beurre de Cacao*, *Huile de Cacao*. A fat substance, of a sweet and agreeable taste, obtained from the *Theobroma cacao* or chocolate nut. Owing to its firmness and fusibility, it is well adapted to serve as the constituent of suppositories.

BUTTER OF COCOA, (F.) *Buerre de Coco*. A fatty, concrete substance, which separates from the milk of the cocoa nut. It is sweet and agreeable.

BUTTER AND EGGS, *Narcissus pseudonarcissus*—b. *Kokum*, see *Garcinia purpurea*—b. of *Nutmegs*, see *Myristica*.

BUTTERBUR, *Tussilago petasites*.

BUTTERCUPS, *Ranunculus acris*, and *R. bulbosus*.

BUTTERFLY-WEED, *Asclepias tuberosa*.

BUTTERMILK, (Prov.) *Churn-milk*, *Kern-milk*, *Whig*, (Sc.) *Bladoch*, *Bledoch*, *Bladda*, *Kirmilk*, *Sourmilk*, (F.) *Babeurre*, *Lait de Beurre*. The thin, sour milk, separated from the cream by churning. It contains caseum and a little butter. It is a refreshing drink when newly made.

BUTTERTEETH, Incisive teeth.

BUTTER WEED, *Erigeron Canadensis*.

BUTTERWORT, *Pinguicula vulgaris*.

BUTTOCK-HUMP, *Steatopyga*.

BUTTONBUSH, *Cephalanthus occidentalis*. BUTTONWOOD SHRUB, *Cephalanthus occidentalis*.

BUTUA, Pareira brava.

BUTYRATE OF GLYCERIN, see *Butter*.

BUTYRIN, see *Butter*.

BUTYRUM, *Butter*—b. *Amygdalarum dulcium*, Confection (almond)—b. *Saturni*, Unguentum plumbi superacetatis—b. *Zinci*, *Zinci chloridum*.

BUVEUR, *Rectus internus oculi*.

BUXTON, MINERAL WATERS OF, *Buxtonien'ses Aquæ*. Buxton is a village in Derbyshire. The springs are thermal, and about 82° Fahrenheit. They contain sulphate of soda, chloride of calcium, chloride of sodium, chloride of magnesium, carbonate of lime, carbonic acid, and azote. They are used in cases in which thermal springs, in general, are recommended. They contain little or no mineral impregnation.

BUXUS, *Buxus sempervirens*. The Box-tree, (F.) *Buis* ou *Bouis*. The leaves are bitter and

aromatic, and, as such, have been used in medicine, in cases of worms, dyspepsia, &c., in the form of decoction. They are sometimes, also, added to beer. The seed was anciently called *Certhe'gan*.

BYNE, Malt.

BY'RETHURUM. A sort of cap or *Couvrechef*, filled with cephalic substances.—Forestus.

BYRON ACID or SOUR SPRINGS. These Springs are in the town of Byron, Genesee county, New York. The water is a nearly pure dilute sulphuric acid. They are powerfully astringent and tonic.

BYRSA, *Bypsa*. A leather skin to spread plasters upon.

BYRSODEP'SICON. A tan stuff, with which CÆLIUS AURELIANUS sprinkled wool, which he

applied in certain cases to the umbilical region; from *Bypsa*, 'leather,' and *de'vco*, 'I tan.'

BYRSODEP'SICON PRINCIPIMUM, Tannin. BYSAU'CHEN, from *βυσ*, 'I stop up,' and *αυχνη*, 'the neck.' A morbid stiffness of the neck. One with a short neck,—*Simotrache'us*.

BYSSOS, Vulva.

BYSSUS, *Byssum*. The ancients gave this name to several vegetable substances, which were used for the fabrication of stuffs prized for their fineness, colour, and rarity of material. It is now chiefly applied to the filaments, by the aid of which the acephalous mollusca attach their shells to the rocks. Byssus was formerly also applied to the *female pudendum*.

BYTHOS, *βυθος*, 'depth.' An epithet used by Hippocrates for the fundus of the stomach.

C.

C. This letter in the chemical alphabet signifies nitre. It is also sometimes used in prescriptions for calx.

CAA-AP'IA, *Dorstenia Brazilian'sis* seu *cordifolia* seu *placentoi'des* seu *vitell'la*. Order, Urticæ. The root, according to Piso, is employed as emetic and anti-diarrhoeic.

CAA-ATAY'A. A plant of Brazil, supposed to be a species of *gratiola*. It is very bitter, and considered to be one of the best indigenous cathartics.

CAACICA, *Euphorbia capitata*.

CAA-GHIYU'YO, *Frutex bac'cifer Brazilian'sis*. A shrub of Brazil, whose leaves, in powder, are considered detersive.

CAAOPIA, *Hypericum bacciforme*.

CAAPEBA, *Pareira brava*, *Pothomorpha pelata* and *P. umbellata*.

CAAPONGA, *Cithrum maritimum*.

CAAROBA. A Brazilian tree, whose leaves, in decoction, promote perspiration. See *Ceratonia*.

CABAL, *Cab'ala*, *Ombal'la*, *Cal'bala*, *Caba'lia*, *Kab'ala*, *Gaballa*. This word is from the Hebrew, and signifies knowledge transmitted by tradition. Paracelsus and several authors of the 16th and 17th centuries have spoken much of this species of magic, which they distinguished into *Juda'ic* or *theolo'gian*, and *Hermel'ic* or *medic'inal*; the latter being, according to them, the art of knowing the most occult properties of bodies by an immediate communication with spirits,—the knowledge being thus acquired by inspiration, and incapable of inducing error. It was also called *Ars cabalistic'ca* seu *signa'ta*, 'cabalistic art.'

CABAL'HAU. A plant of Mexico, according to Dalechamps, which passes for an antidote to white hellebore, and yet is used for poisoning arrows. It is unknown to botanists.

CABALIST, *Cabalie'ta*. One instructed in the Cabal.

CABALLATION, *Cynoglossum*.

CABARET, *Asarum*.

CABBAGE, *Brassica*.—c. Cow, *Nymphæa odorata*.—c. Irish, *Dracontium foetidum*.—c. Skunk, *Dracontium foetidum*.—c. Swamp, *Dracontium foetidum*.—c. Water, *Nymphæa odorata*.—c. Tree, *Geoffræa inermis*.—c. Bark tree, *Geoffræa inermis*.—c. CABBAGIUM, *Geoffræa inermis*.

CABOTZ, *Hagenia Abyssinica*.

CABUREIBA, see *Myroxylon Peruiferum*.

CABUREICIBA, see *Myroxylon Peruiferum*. CACÆMIA, *Cachæ'mia*, from *kakos*, 'bad,' and 'αμα,' 'blood.' A faulty or morbid condition of the blood.

CACÆSTHE'SIS, *Cacæsthe'sis*, *Cacæsthe'sis*, from *kakos*, 'bad,' and *αισθησις*, 'feeling.' Morbid sensation. Morbid general feeling. Indisposition.

CACAFERRI, *Ferri subcarbonis*.

CAC'AGOGUE, *Cacago'gus*, *Cac'cagogue*, *Cacago'gus*, from *κακη*, 'excrement,' and *αγειν*, 'to expel.' An ointment, composed of alum and honey; which, when applied to the anus, produced an evacuation.—Paulus of Ægina.

CACALEXITERIA, *Alexiteria*.

CACA'LLIA ANTEUPHORB'BIUM, *Anteuphor'bium*. Ord. Compositæ. A plant, which Dodoens and others considered to be capable of tempering the caustic properties of euphorbium. It is also called *Klein'ia*.

Many varieties of the *Cacalia* are used, in different countries, chiefly as condiments.

CA'CAO, *Ca'coa*, *Caca'vi*, *Quahoil*, *Cacava'ta*. The cocoa or chocolate nut; fruit of *Theobroma cacao*, *Co'coa* *Cacavif'era*, *Ca'cao minor* seu *sati'va*, *Cacao theobro'ma*, (F.) *Cacoyer* ou *Cacotier ordinaire*. Family, Malvacæ. Sex. Syst. Polydelpbia Pentandria.

CACAO'TIER ORDINAIRE, see *Cacao*.

CACAOYER ORDINAIRE, see *Cacao*.

CACATION, Defecation.

CACATORIA, *Diarrhoea*.

CAC'ATORY, *Cacato'rius*, from *cacare*, 'to go to stool.' *Febris cacato'ria*; a kind of intermittent fever, accompanied by copious alvine evacuations.—Sylvius.

CACAVATA, *Cacao*.

CACAVI, *Cacao*, *Jatropha manihot*.

CACCAGOGUE, *Cacagogue*.

CACCE, Excrement.

CACCION'DE. A sort of pill, chiefly formed of catechu, recommended by Baglivi in dysentery.

CACEPHEBOTE'SIA, from *kakos*, 'bad,' and *εφεβον*, 'puberty.' Morbid puberty. Disease occurring at the period of puberty.

CACHALOT, see *Cetaceum*.

CACHANG-PARANG. A sort of bean of Sumatra, mentioned by Marsden, whose seeds are given in pleurisy. Jussieu considered it to be the *Mimosa scandens*.

CACHECTIC, *Cachec'tes*, *Cachec'ticus*, (F.) *Cachectique*, same etymon as *Cachexia*. One at

tacked with cachexia. Belonging to cachexia. *Cachectica remedia* are remedies against cachexia.

CACHELCOMA. Hellococae.

CACHEN-LAGUEN, Chironia Chilensis.

CACHET (F.), 'a seal.' A stamp or medicine stamp, used by the ancient Romans for marking their drugs, especially those prescribed in diseases of the eyes; and hence called, by some, *oculist stamps*. Some of those more lately discovered amongst Roman antiquities, have been described by M.M. Siehel and Duchalais, and by Prof. Simpson, of Edinburgh.

CACHEXIA, from *kakos*, 'bad,' and *εξις*, 'habit.' *Status cachecticus*, *Cach'icay*, *Dysthe'sis*, (F.) *Cachexie*. A condition in which the body is evidently depraved. A bad habit of body, chiefly the result of scorbutic, cancerous, or venereal diseases when in their last stage. Hence we hear of a *Scorbutic Cachexia*, *Cancerous Cachexia*, &c. Sauvages and Cullen have included under this head a number of diseases—consumptions, dropsies, &c. Cachexia has been sometimes confounded with diathesis. *Cachexia Icteric* is jaundice or icterus itself, or a disposition thereto. Fluor albus is sometimes called *Cachexia Uteri*na.

CACHEXIA AFRICANA, Chthonophagia—c. Calculosa, Lithia—c. Cancerous, see Cancer—c. Chlorotic, Chlorosis—c. Dysthetica, Dyscrasia.

CACHEXIA, GAOL. The deteriorated organic actions induced by confinement in prisons, which is so often the precursor of scrofula.

CACHEXIA ICTERICA, Icterus.

CACHEXIA LONDINENSIS. The paleness and other evidences of impaired health presented by the inhabitants of London. A similar cachexia is seen in those of other crowded cities.

CACHEXIA, LYMPHATICA FARCINOSA, see Equinia.

CACHEXIA, MARSH, (F.) *Cachexie paludéenne*, on *palustre*. The state of cachexy observed in malarious districts.

CACHEXIA, SATURNINA, Saturnismus—c. Scorbutic, see Purpura—c. Scrophulosa, Scrofula.

CACHEXIA SPLE'NICA. The state of scorbutic cachexia, which often accompanies diseases, especially enlargement of the spleen, *Splenal'gia Bengalen'sis*, in India.

CACHEXIA VENEREA, Syphilis—c. Venous, Venosity—c. Virginum, Chlorosis.

CACHEXIE, Cachexia—c. *Iodée*, see Iodine—c. *Paludéenne*, Cachexia, marsh—c. *Palustre*, Cachexia, marsh.

CACHEXY. Cachexia.

CACHIBOU, see *Bursera gummifera*.

CACHINLAGUA, Chironia chilensis.

CACHINNA'TIO, from *cachinno*, 'I laugh aloud.' A tendency to immoderate laughter, as in some hysterical and maniacal affections.

CACHIRI. A fermented liquor made, in Cayenne, from a decoction of the rasped root of the manioc. It resembles perry.

CACHIX. A small stone or pebble, found on the sea shore. One of these, when heated in the fire, and cooled in whey, communicates an astringency to the liquid, so that it was anciently esteemed to be useful in dysentery.—Galen.

CACHOS. An oriental fruit, apparently of a Solanum, which is esteemed lithontriptic.

CACHOU, Catechu.

CACHRYS LIBANO'TIS. An umbelliferous plant which grows in Africa and the south of Europe. It is aromatic and astringent. Its seeds are extremely acrid.

CACHRYS MARITIMA, Crithmum maritimum.

CACHUN'DE. An Indian troch or pastile composed of amber, mastic, musk, cinnamon, alices, rhubarb, galanga, pearls, rubies, emeralds,

garnets, &c. It is regarded by the people of India as an antidote, stomachic and anti-spasmodic.

CACO, *kako*, properly only an abbreviation of *kakos*. In composition it means something defective; as in the following words:

CACOE'STHESIS, Cacæsthesi.

CACO-ALEXIT'ERIA, Alexipharmic.

CACOCOL'IA, from *kakos*, 'bad,' and *χολη*, 'bile.' Diseases induced by a depraved condition of the bile.

CAC'OCRO'LI, *Cac'ochri*, from *kakos*, 'bad,' and *χρσα*, 'colour.' Diseases in which the complexion is morbidly changed in colour.

CACOCYL'IA, from *kakos*, 'bad,' and *χυλος*, 'chyle.' Depraved chylification.

CACOCY'M'IA, *Kakochym'ia*, *Corruptio Humo'rum*, from *kakos*, 'bad,' and *χυμος*, 'juice,' 'humour.' *Cacoch'ymy*. Depravation of the humours.

CACOCY'MIA PLUMBEA, Lead poisoning—c. Scorbutica, see Purpura—c. Scrophulosa, Scrofula—c. Veneræ, Syphilis.

CACOCY'MUS, *Cacochym'icus*. One attacked with cacochymia. Belonging to cacochymia.

CACOCNE'MUS, *Cacocnem'icus*, *Malis suris præditus*; from *kakos*, 'bad,' and *κνημη*, 'the leg.' One who has bad legs.

CACOCORE'MA, from *kakos*, 'bad,' and *κορεω*, 'I purge, or cleanse.' A medicine which purges off the vitiated humours.

CACODE'MON, from *kakos*, 'bad,' and *δαίμων*, 'a spirit.' An evil spirit, to which were ascribed many disorders. The nightmare.

CACOD'ES, from *kakos*, 'bad,' and *οειν*, 'to smell'—*malè olens*. Having a bad smell; *Cacodia*, *Cacos'mia*.

CACODIA, see Cacodes.

CACOE'TH'ES, *Cacoe'th'icus*, (F.) *Cacoe'the*, from *kakos*, 'bad,' and *εθος*, 'disposition, habit,' &c. Of a bad or vitiated character, as *ulcus cacoe'thes*, an ulcer of a malignant character.

CACOETHICUS, Cacoethes.

CACOGALACT'IA, *Cacogal'lia*, from *kakos*, 'bad,' and *γαλα*, gen. *γαλακτος*, 'milk.' A bad condition of the milk.

CACOGALACTICA. Same etymon as the last. One who suffers from a bad condition of the milk.

CACOGALIA, Cacogalactia.

CACOGENESIS, (F.) *Cacogénese*, from *kakos*, 'bad,' and *γενεσις*, 'generation.' A morbid formation. A monstrosity.

CACOMORPHIA, Deformation.

CACOMORPHOSIS, Deformation.

CACOPATH'IA, *Pas'sio Muli*, from *kakos*, 'bad,' and *παθος*, 'affection.' A distressed state of mind.—Hippocrates.

CACOPHO'NIA, from *kakos*, 'bad,' and *φωνη*, 'voice,' *vitia'ta vox*. A dissonant condition of voice.

CACOPLAS'TIC, *Cacoplas'ticus*, *Dysplasmatic*; from *kakos*, 'bad,' and *πλασσω*, 'I form.' Susceptible of only a low degree of organization, as the indurations resulting from low or chronic inflammation, fibro-cartilage, cirrhosis, &c.

CACOPRA'GIA, *Cacoprax'is*, from *kakos*, 'bad,' and *πραττω*, 'I perform.' Depraved condition of the organic functions.

CACOPRAXIS, Cacopragia.

CACORRHACHI'TIS, from *kakos*, 'bad,' and *ραχις*, 'the spine.' *Cacorrhachis*, *Cacorrhachis*, *Cacorrhach'itis*, *Spondylal'gia*. Deformity of the spine. Disease of the spine. Spontaneous luxation of the vertebrae and ribs dependent upon internal causes.

CACORRHYTH'MUS, *Arrhyth'mus*, from *kakos*, 'bad,' and *ρhythμος*, 'rhythm,' 'order.' Irregular.

CACOS'IS. *Mala disposit'io*, (F.) *Vice*. A

bad condition of body.—Hippocrates. A diseased condition in general.

CACOSIT'IA, from *κακος*, 'bad,' and *σιτιον*, 'aliment.' Disgust or aversion for food.—*Fæstidium ciborum*.

CACOSMIA, see *Cacodes*.

CACOSOM'UM, from *κακος*, 'bad,' and *σωμα*, 'the body.' An hospital for leprosy, and incurable affections in general.

CACOSPERMA'SIA, *Cacospermattia*, *Cacospermia*, from *κακος*, 'bad,' and *σπερμα*, 'sperm.' A bad condition of the sperm.

CACOSPHYX'IA, from *κακος*, 'bad,' and *σφυξις*, 'pulse.'—*Vitiosus pulsus*. Bad state of pulse.—Galen.

CACOSPLANCH'NIA, from *κακος*, 'bad,' and *σπλῆγνον*, 'a viscous.' Indigestion. The emaciation dependent upon imperfect digestion.—Siebenhaar.

CACOSTOM'ACHUS, from *κακος*, 'bad,' and *στομαχος*, 'the stomach.' What disagrees with the stomach. Indigestible.—Gorræus.

CACOSTOMATOPHRESIA, Breath, offensive.

CACOS'TOMUS, from *κακος*, 'bad,' and *στομα*, 'a mouth.' Having a bad mouth, or a bad breath.

CACOTHANASIA, see *Euthanasia*.

CACOTHYM'IA, *Vitium Animi*, from *κακος*, 'bad' and *θυμος*, 'mind,' 'disposition.' A vicious state of mind.—Linden.

CACOTRIBULUS, *Centaurea calcitrapa*.

CACOTRICH'IA, from *κακος*, 'bad,' and *τριχ*, 'hair.' Disease of the hair.

CACOTROPH'IA, from *κακος*, 'bad,' and *τροφη*, 'nutrition.'—*Vitiosa nutritio*;—disordered nutrition.—Galen.

CACOU, *Cagot*, *Catechu*.

CACOU'CIA COCCIN'EA, *Coccin'ea*, *Coccin'ea*, *Schousbæa coccin'ea*, *Tikimma*. A perennial twining shrub of South America, the plant of which, as well as the fruit, is possessed of emeto-cathartic properties.

CACTIER, *Cactus opuntia*.

CACTUS, *Cynara*.

CACTUS OPUN'TIA, *Opuntia*, *O. vulg'ris*. The Indian Fig. (F.) *Cactier*, *Raquette*, *Figuier d'Inde*. Ord. *Cactaceæ*. This plant grows in South America, Spain, Italy, &c. Its fruit, which has the shape of the fig, is of a sweetish taste, and colours the urine red when eaten. Its leaves are considered refrigerant.

The fruits of different species of cactus are called *Tunas*.

CADA'BA, *Stroëmia*. A genus of the family *Capparideæ*, natives of India and Arabia. The young shoots of the *Cada'ba farinosa* are considered to be an antidote against venomous bites.

CADA'VER, *Ptoma*, *Neeron*. A dead body, *corps*, *corpse*, (Sc.) *Corp*; a subject; a *carcass*, (F.) *Cadavre*. The word has been supposed to come from *cado*, 'I fall'; and by some to be a contraction from *caro data vermis*, 'flesh given to the worms.' [?]]

CADAVERIC, *Cadaverous*.

CADAVEROUS, *Cadav'eric*, *Cadavero'sus*, *Necro'idæ*, (F.) *Cadav'reux*, *Cadav'érique*. Belonging to the dead body; as *cadaverous smell*. The *Cadav'ericus* or *Hippocrat'ic face*, (see *Face*), is an unfavourable sign in disease, and generally denotes a fatal termination.

CADAV'EROUS or CADAV'ERIC HYPER'EMIA. The hypostatic hyperæmia observed in dependent parts of the dead body.

CADDY INSECT, see *Ectoza*.

CADE, *Juniperus oxycedrus*.

CADEJ-INDI, *Malabathrum*.

CADEL-AVANACU, *Croton tiglium*.

CADIA. An Egyptian, leguminous plant.

The Arabs attribute to its fresh leaves the power of relieving colic.

CADIVA INSANIA, Epilepsy.

CAD'MIA, *Calamina*, *Tutia*.

CADMI'I SULPHAS, *Cadm'ium sulphuricum*, *Sulphas Cadm'ius*, *Melini Sulphas*, *Klaprothii Sulphas*, *Klaprothium Sulphuricum*, *Melinum Sulphuricum*, *Sulphate of Cad'mium*. Used in spots on the cornea, and in chronic torpid inflammation of the conjunctiva, in the quantity of half a grain to a grain to the ounce of water. It does not differ materially in its properties from sulphate of zinc. The nitrate, in the dose of ʒth of a grain, induces vomiting and purging.

CADM'UM SULPHURICUM, *Cadmii Sulphas*.

CADRE DU TYMPAN, *Tympanal*. The portion of the temporal bone which supports the membrana tympani.

CADTCHU, *Catechu*.

CADUCA HUNTERI, *Decidua*—c. *Passio*, Epilepsy.

CADU'CITY, *Imbecillitas*, *Debil'itas*, *Cadu'citas*, from *cadere*, 'to fall.' The French use the word *Caducité* for the portion of human life which is comprised generally between 70 and 80 years. The age which precedes decrepitude. It is so termed in consequence of the limbs not usually possessing sufficient strength to support the body. The precise age must of course vary in individuals.

CADUQUE, *Decidua membrana*—c. *Réfléchie*, see *Decidua membrana*—c. *Uterine*, *Decidua*—c. *Vraie*, *Decidua membrana*.

CADURCUS, *Vulva*.

CADUS, *καδος*. A Greek measure equal to ten gallons English.—Pliny. *Amphora*.

CÆCA FORAM'INA (ANTERIUS ET POSTERIUS) are situate at the fore and back parts of the tuber annulare of the brain, and at the extremities of the depression made by the vertebral artery. The former is placed between the nerves of the third, and the latter between those of the sixth pair.

CÆCÆ HÆMORRHOÏ'DES, *Blind Piles*, (F.) *Hémorrhoides aveugles*, are those unaccompanied by any discharge.

CÆCÆ, *Cæc'lis*. Belonging to the Cæcum, from *cæcus*, 'blind, hidden.' The *Cæcal arteries* and *veins* are the branches of the *Arteria et vena colica dextra inferiores*, distributed to the cæcum. The *Cæcal* or *blind* extremity of a duct is its closed termination. See *Cæcus*.

CÆCATRIX, *Cicatrix*.

CÆ'CITAS, *Cæ'citas*, *Cæcitu'as*, *Ablep'sia*, *Obeæcat'io*, *Ocæcat'io*, *Anap'sia*, *Typhlo'es*, *Ce'city*, *Blindness*, (F.) *Aveuglement*, *Cécité*, *Perte de la vue*. *Cæcitas* may be dependent upon many different diseases—as upon amaurosis, specks, hypopyon, cataract, glaucoma, ophthalmia, atrophy of the eye, &c.

CÆCITAS CREPUSCULARIS, *Hemeralopia*—c. *Diurna*, *Nyctalopia*—c. *Nocturna*, *Hemeralopia*.

CÆCITE, *Typhlo-enteritis*.

CÆCITUDO, *Cæcitas*.

CÆCUM, *Cæcum*, *Intest'ium cæcum*, *Monomæchum*, *Monom'æcum*, *Monocælon*, *Monocælum*, *Typhlo'teron monocælon*, *Typhlo'terum*, *Typhlo'enterum*, *Int'ium intest'ini crassi*, *Succus Intest'ini crassi seu Coli*, *Cæcum Caput coli*, *Caput coli*, *Prima cella coli*, *Int'ium ext'uberans coli*, from *cæcus*, 'blind.' The *Blind Gut*, so called from its being open at one end only. That portion of the intestinal canal which is seated between the termination of the ileum and commencement of the colon; and which fills, almost wholly, the right iliac fossa; where the peritoneum retains it immovably. Its length is about three or four fingers' breadth. The *Ileo-cæcal*

valve or *Valve* of Bauhin shuts off all communication between it and the ileum; and the *Appendix vermiformis cæci* is attached to it.

CÆCUM FORAMEN of the frontal bone is a small cavity at the inferior extremity of the internal coronal crest or crista. — *Fronto-ethmoidal foramen*, (F.) *Trou aangle ou borgne*. Morgagni has given the same name to the small cavity in the middle of the upper surface of the tongue, near its base; the sides of which are furnished with mucous follicles — *Lacunes de la langue* — (Ch.)

CÆCUM, PHLEGMONOUS TUMOUR OF THE, Typhlo-enteritis.

CÆCUS. 'Blind.' One deprived of sight, *Typhlops*, (F.) *Aveugle, Borgne*. In anatomy, it is used to designate certain holes or cavities, which end in a *cul-de-sac*; or have only one opening. See *Cæcal*.

Blind Ducts of the Ure'thra, (F.) *Conduits aveugles de l'urèthre*, are the *Mucous Lacunæ of the Ure'thra*.

CÆLA-DOLO, *Torenia Asiatica*.

CÆMENTUM, Lute.

CÆRULEUM BEROLINENSE, Prussian blue — *c. Borussiae*, Prussian blue.

CÆRULOSIS NEONATORUM, Cyanopathy.

CÆSALPINIA, *C. sappan*, *Sappan* or *Sampfen wood*, (F.) *Brésillet, Bois de Suppan*. Ord. Leguminosæ. Called after Cæsalpinus. A small Siamese tree, the wood of which is used in decoction, in cases of contusion.

Brazil wood, *Pernambuco* or *Fernambuco wood*, formerly used as an astringent, is the wood of *CÆSALPINIA ECINATA*. This is the proper *Brazil wood*; but another variety in commerce is the *Brasileto* from *Cæsalpinia Brasiliensis* and *C. cristata*, which grow in the West Indies.

The *Nicaragua* or *Peach-wood* is analogous to this, and is said to be derived from a species of *Cæsalpinia*.

The kernel of *CÆSALPINIA BONDUCELLA*, the seed of which is called in India *Kutuleja* and *Kutoo Kurnja*, is given as a febrifuge tonic. Dose, ten grains.

CÆSAREAN SECTION, *Cæsa'rean operation*, *Tomotoc'ia*, *Cæsa'rea sectio*, *Partus cæsa'reus*, *Operatio cæsa'rea*, *Metrotom'ia*, (F.) *Opération Césarienne*, from *cædère*, *cæsum*, 'to cut.' An incision made through the parietes of the abdomen and uterus to extract the foetus. In this manner, Julius Cæsar is said to have been extracted. — Pliny. It is also called *Hysterotom'ia*, *Hysterotomotoc'ia*, *Gastrometrotom'ia*, *Gusterhysterotomy*, *Gastrometrotom'ia*, *Gastrohysterotomy*. An incision has been made into the uterus through the vagina, constituting the *Vaginal Cæsarean Section*, *Gastroelytrotom'ia*, *Gastrellytrotom'ia*, *Gastrocolpotom'ia*, *Laparacolpotom'ia*, *Laparoelytrotom'ia*, (F.) *Opération césarienne vaginale*. The Cæsarean section may be required when the mother dies before delivery; when there is some invincible obstacle to delivery from the faulty conformation of the pelvis; or when the child has passed into the abdominal cavity in consequence of rupture of the uterus.

CÆSARIES, Capillus, Scalp.

CÆSIUS, Glaucoma.

CÆSONES, *Cæ'sares*. Children brought into the world by the Cæsarean operation.

CÆSULIÆ. They who have gray eyes.

CÆSURA, Cut.

CÆTCHU, *Catechu*.

CAF, Camphor.

CAFAL, Agrimony.

CAFAR, Camphor.

CAFÉ, Coffee.

CAFÉ À LA SULTANE. This name has

been given to an infusion or decoction of the ground *coques* or pericarps which surround the coffee.

CAFÉ CITRIN. The aqueous infusion of unroasted coffee, so called on account of its yellowish tint.

CAFEIN, see *Coffea Arabica*.

CAFÉIER and **CAFÉYER**, *Coffea Arabica*.

CAFFA, Camphor.

CAFFEIN, see *Coffea Arabica*.

CAFIER, *Coffea Arabica*.

CAFUR, Camphor.

CAGAS'TRUM. The principal or germ of diseases which are communicable. — *Paracelsus*.

CAGNEUX, *Cagot*. See *Kyllosis*.

CAGOSANGA, *Ipecacuanha*.

CAGOTS (F.). A name given to deformed and miserable beings, met with in the Pyrénées, Bern, and Upper Gascony, in France, where they are also called *Capots*. In other districts they are called *Gézits*, *Gézitains*, *Crétins*, *Gahets*, *Cayons*, *Culiberts*, *Cacous*, *Cagneux*, &c. See *Crétin*. The word *Cagot* is supposed to be an abbreviation of *Canis Gothus*, 'Dog of a Goth.'

CAGUE-SANGUE, *Caguesangué*.

CAHINCÆ RADIX, *Caincæ radix*.

CAJEPUT OIL, *Caj'eput oil*, *Kyaput'ty*, *Cajuput'i O'leum*. The volatile oil of the leaves of *Melaleuca Cajuputi*, Ord. Myrtaceæ, a native of the Moluccas. The oil has a strong, fragrant smell, like camphor; taste pungent and aromatic. It is stimulant, and useful where the essential oils in general are employed. It has also been called *Oil of Witneben*, from the person who first distilled it.

CAIK, *Pleurodynia*.

CAIL-CEDRA, *Swietenia Senegalensis*.

CAILLE, *Tetrao coturnix*.

CAILLÉ, see *Curd*.

CAILLEAU, *Lantana*.

CAILLEBOTTE, see *Curd*.

CAILLE-LAIT, *Rennet* — *c. Blanc*, *Galium mollugo* — *c. Vraie*, *Galium verum*.

CAILLETTE, *Abomasus*.

CAILLOT, *Coagulum*.

CAINANÆ RADIX, *Caincæ radix*.

CAÏN'CÆ RADIX, *Radix Chiococ'ea*, *R. Caina'næ* seu *Cainana'næ* seu *Cahincæ* seu *Kahincæ* seu *Serpenta'riæ Brasilien'sis*, *Cainca Root*. The bark of the roots of *Chiococ'ea anguif'uga* seu *densif'olia*, and, perhaps, *Ch. racem'osa*, a plant of the Order *Rubiaceæ*. *Sex. Syst.* *Pentandria Monogynia*, of Linnaeus. It is bitter, tonic, and diuretic, but has not been long introduced. Dose of the powder, from ʒj to ʒss.

Dr. John H. Griscom, of New York, considers there is a remarkable analogy between the *Cainca* and the *Apocynum cannabinum*.

CAINITO, *Chrysophyllum Cainito*.

CAIPA SCHORA. A cucurbitaceous Malabar plant, the fruit of which has a pyriform shape. The juice is drunk in that country for the purpose of arresting hiccough. The fruit, when unripe, is emetic.

CAISSE, Case — *c. du Tambour*, *Tympanum* — *c. du Tympan*, see *Tympanum*.

CAITCHU, *Catechu*.

CAJAN, *Phaseolus creticus*.

CAJUPUTI, *Caieput*.

CAKES, WORM, STORY'S. These were composed of *calomel* and *jalap*, made into cakes, and coloured with *cinnabar*.

CALABASH-TREE, **NARROW-LEAVED**, *Crescentia Cujete*.

CALADIUM ESCULENTUM, *Arum esculentum*.

CALAD'NIUM SEGU'NUM, *Nat. Ord. Araceæ*. A plant of India, whose acrid juice has been given

there in gout and rheumatism, and as an anaphrodisiac to women. The tincture has been prescribed in pruritus vulvæ.

CALAF, *Salix Ægyptiaca*. A large-leaved Egyptian willow, called, also, *Ban*. The distilled water of the flowers, called *Macahalef*, passes, in that country, for an excellent antaphrodisiac. It is also used as an antiloinic, antiseptic, and cordial.

CALAGUALA, see *Calagnalæ radix*.

CALAGERI, *Vernonia anthelmintica*.

CALAGIRAH, *Vernonia anthelmintica*.

CALAGUALÆ RADIX, *Calagne'le Radix*. The root of *Polypodium Calagua'la seu adiantifor'me seu coria'ceum seu ammi'folium seu argenteum seu pol'itum*, *Aspidium coria'ceum seu ferrugin'eum seu dis'color*, *Tectaria calahuala seu ferrugin'ea*, *Calagnala*, *Calahuala*. Order, Felices. It has been exhibited in Italy in dropsy, pleurisy, contusions, abscesses, &c. Its properties are not, however, clear.

CALAHUALA, see *Calagnalæ radix*.

CALAMANDRINA, *Teucrium chamædrys*.

CALAMBAC, *Agallochum*.

CALAMBOUK, *Agallochum*.

CALAME'DON, from *καλαμος*, 'a reed.' This word has had various significations. Some have used it for an oblique fracture of a bone; the fractured portions having the shape of the nib of a pen. Others have used it for a longitudinal fracture; and others, again, for one that is comminuted.

CALAMENT, *Melissa calamintha*.

CALAMINA, *Calamine*, from *calamus*, 'a reed,' so called from its reed-like appearance. *Cadm'ia*, *C. lapidea aëro'sa seu fos'silis*, *Lapis Aëro'sus*, *Lapis Calaminaris*, *Calaminaris*, *Car'bonas Zinc' impu'rus*, *Cathmir*, (F.) *Pierre calaminaire*. Native impure carbonate of zinc. Calamine is chiefly used for pharmaceutical purposes in the form of the *CALAMINA PRÆPARATA* (Ph. U. S.), *Lapis Calaminaris præparatus*, *Car'bonas zinc' impu'rus præparatus*, *Zinc' car'bonas præparatus*, *Prepared Calamine*:—Calamine reduced to an impalpable powder by roasting and levigation. In this state it is sprinkled or dusted on excoriated parts, or to prevent excoriation, &c.

CALAMINARIS, *Calamina*.

CALAMINT, *Melissa Calamintha*—c. Field, *Melissa nepeta*—c. Mountain, *Melissa grandiflora*—c. Spotted, *Melissa nepeta*.

CALAMINTA HUMILIOR, *Glechoma hederacea*.

CALAMINTHA, *Melissa C.*—c. *Anglica*, *Melissa nepeta*—c. *Erecta Virginiana*, *Cunila Mariana*—c. *Hederacea*, *Glechoma hederacea*—c. *Magno flore*, *Melissa grandiflora*—c. *Montana*, *Melissa grandiflora*—c. *Nepeta*, *Melissa nepeta*—c. *Parviflora*, *Melissa nepeta*—c. *Pulegii odore*, *Melissa nepeta*—c. *Trichotoma*, *Melissa nepeta*.

CALAMUS, *καλαμος*, 'the reed.' In the Pharmacopœia of the U. S., the rhizoma of *acorus calamus*.

CALAMUS ALEXANDRI'NUS. Celsus has thus called a medicine, which was long confounded with *Calamus Aromaticus*. It is not a root, however, but the stalk of a plant of India and Egypt, probably the *Andropogon Nardus*. It entered into the theriaca, and has been regarded as antihysteric and emmenagogue;—*Calamus aromaticus verus*.

CALAMUS AROMATICUS, *Acorus calamus*—c. *Aromaticus verus*, *Calamus Alexandrinus*—c. *Draco*, *C. rotang*—c. *Indicus*, see *Saccharum*—c. *Odoratus*, *Acorus calamus*, *Juncus odoratus*.

CALAMUS ROTANG, *C. Draco*. Order, Palmæ. The systematic name of a plant, whence *Dragon's Blood*, *Sanguis Draco'nis*, *Cinnab'aris Græcorum*,

Draconthæ'ma, (F.) *Sang-Dragon*, is procured. It is the red, resinous juice, obtained, in India, from wounding the bark of the *Calamus Rotang*. It has been used as an astringent in hemorrhages, &c.; but is now rarely employed.

CALAMUS SCRIPTO'RIVUS, *Anag'typhæ*, 'a writing pen,' (F.) *Fossette angulaire du quatrième ventricule*. A small, angular cavity, situate at the superior extremity of the medulla, in the fourth ventricle of the brain, which has been, by some, supposed to resemble a pen.

CALAMUS VULGARIS, *Acorus calamus*.

CALAPPITE. Rumphius has given this name to calculeous concretions, found in the interior of certain cocoa-nuts. The cocoa-tree itself the Malays call *Calappa*. These stones are, likewise, termed *Vegetable Bezoards*. The Malays attribute potent virtues to them, and wear them as amulets.

CALASAYA, *Cinchonæ cordifoliæ cortex*.

CALBALA, *Cabal*.

CALBIA'NUM. The name of a plaster in Myrepsus, the composition of which we know not.

CALCADINUM, *Ferri sulphas*.

CALCAIRE, *Calcareous*.

CALCA'NEAL, *Calca'neus*, from *calx*, 'the heel.' Having relation to the calcaneum, as 'calcaneal arteries.'

CALCANEO-PHALANGIEN DU PETIT ORTEIL, *Abductor minimi digiti pedis*—c. *Phalanginiæ communis*, *Extensor brevis digitorum pedis*—c. *Sous-phalangietien communis*, *Flexor brevis digitorum pedis*—c. *Sous-Phalangien communis*, *Flexor brevis digitorum pedis*—c. *Sous-phalangien du petit orteil*, see *Abductor minimi digiti pedis*—c. *Sus-phalangietien communis*, *Extensor brevis digitorum pedis*.

CALCA'NEUM, from *calx*, 'the heel.' *Calca'neus*, *Calcæar*, *Cal'cia*, *Ichnus*, *Os Calcis*, *Pterna*, *Pter'nium*. The largest of the tarsal bones: that which forms the heel. It is situate at the posterior and inferior part of the foot; is articulated above and a little anteriorly with the astragalus; anteriorly, also, with the os cuboides. Its posterior surface—called *Heel*, *Talus*, *Calx*, (F.) *Talon*—gives attachment to the tendo-achillis: the lower has, posteriorly, two tuberosities, to which the superficial muscles of the sole of the foot are attached. The small *Apoph'ysis* or *lateral Apoph'ysis* of the *Calca'neum*, (F.) *Petit Apophyse* ou *Apophyse latérale du Calcaneum*, is a projection at the upper surface of this bone, on which is formed the posterior portion of the cavity that receives the astragalus. The great *Apoph'ysis*, anterior *Apoph'ysis* of the *Calca'neum*, is the projection which corresponds, on one side, with the cuboides; and on the other forms the anterior part of the facette which receives the astragalus.

CALCAR, *Calcaneum*, *Ergot*—c. *Avis*, *Hippocampus minor*.

CALCA'REOUS, *Calca'reus*, *Calca'rius*, from *calx*, 'lime.' (F.) *Calcaire*. Containing lime:—as *calcareous concretions*, *C. depositions*, &c.

CALCAREUS CARBONAS, *Creta*.

CALCARIA CHLORATA, *Calcis chloridum*—c. *Chlorica*, *Calcis chloridum*—c. *Phosphorica*, see *Cornu cervi*—c. *Pura*, *Calx*—c. *Pura liquida*, *Liquor calcis*.

CALCARIÆ CHLORUM, *Calcis chloridum*.

CALCATAR, *Ferri sulphas*.

CALCATREPPOLA, *Centaurea calcitrapa*.

CALCE'NA, **CALCE'NON**, **CALCENO'NTIA**, **CALCINO'NTIA**. Words employed by Paracelsus to designate the concretions of tartrate of lime which form in the human body.

CALCENOS, *Calcetus*.

CALCEOLARIA, from *calceolus*, 'a small slipper;' *Slipperwort*.

CALCEOLARIA PRIMATA is used in Peru as a laxative.

CALCEOLARIA TRIFIDA is esteemed to be febrifuge.

CALCETUS, *Calcetaninus*, *Calcetinus*. That which abounds in tartrate of lime. An adjective used by Paracelsus in speaking of the blood; *Sanguis calcetinus*. Hence came the expression, *Calcined blood*, *Sang calciné*.

CALCEUM EQUINUM, Tussilago.

CALCHOIDEA (OS), Cuneiform bone.

CALCIA, Calcareum.

CALCIFEROUS CANALS, Canaliculi califeri.

CALCIFICATION, see Cretification.

CALCIGEROUS CELL, see Tooth.

CALCIGRADUS, *Pternobates*, from *calx*, the heel, and *gradus*, 'a step.' One who walks on his heels.

CALCII CHLORIDUM, Calcis murias—c. Chloruretum, Calcis murias—c. Oxychloruretum, Calcis chloridum—c. Oxydum, Calx viva—c. Protochloruretum, Calcis chloridum—c. Sulphuretum, Calcis sulphuretum.

CALCINATION, *Calcina'tio*, *Calc'i'non*, *Cop'rena'tio*, from *calx*, 'lime.' The act of submitting to a strong heat any infusible mineral substance, which we are desirous of depriving either of its water, or of any other volatilizable substance that enters into its composition; or which we wish to combine with oxygen. *Alum* is calcined to get rid of its water of crystallization;—*chalk*, to reduce it to the state of pure lime, by driving off the carbonic acid; and certain metals are subjected to this operation to oxidize them.

CALCINATUM MAJUS POTERII, Hydrargyrum præcipitatum.

CALCINONIA, Calceana.

CALCIS BICHLORURETUM, Calcis chloridum—c. Carbonas, Creta—c. Carbonas durus, Creta, Marmor—c. Carbonas friabilis, Creta.

CALCIS CARBONAS PRÆCIPITATUS, *Præcipitated Carbonate of Lime*, *Precipitated Chalk*. This preparation, introduced into the last edition of the Pharmacopœia of the United States, is prepared as follows: *Liq. Calcii Chlorid.* Ovs. *Soda Carb. sat.* lbvj; *Aqua destillat.* q. s. Dissolve the carbonate of soda in six parts of distilled water; heat this and the solution of chloride of calcium, separately, to the boiling point, and mix. Wash the precipitate repeatedly with distilled water, and dry on bibulous paper. It has the same properties as creta præparata, and is preferred to it in certain cases,—for example, as an ingredient in tooth powders, owing to its freedom from gritty particles.

CALCIS CHLORIDUM, *Chlo'ride* or *Chlo'ruret* or *Hypochlo'rite* or *Chlorite* or *Oxy muriate of Lime*, *Calx chlorinata* (Ph. U. S.) seu *Oxymuriatica*, *Protochloruret of Calcium*, *Calc'ria chlora'ta* seu *Chlorica*, *Chlorum* seu *Chloratum Calca'ria*, *Oxychlorure'tum* seu *Protochlorure'tum Calcii*, *Chlorur'tum Oxidi Calcii*, *Bichlorure'tum* seu *Oxymuriatus Calcis*, *Calcis Hypochloris*, *Bleaching or Tennant's Powder*, (F.) *Protochlorure de Calcium*, *Chlorure d'Oxide de Calcium*, *Chlorure ou Oxichlorure ou Bichlorure ou Oximuriate ou Muriate suraigéné ou origéné de Chaux*, *Poudre de Blanchement*, *P. de Tennant*. A compound resulting from the action of chlorine on hydrate of lime. Chloride of lime is a most valuable disinfecting agent, (see Disinfection,) when dissolved in the proportion of one pound to six gallons of water. It has likewise been employed both internally and externally in various diseases, as in scrofula, fœtor oris, foul ulcers, &c. &c.

CALCIS HEPAR, Calcis sulphuretum—c. Hydras, see Calx—c. Hypochloris, Calcis chloridum.

CALCIS MURIAS, *Muriate of Lime*, *Calx salita*, *Calcii Chlorure'tum* seu *Chlo'ridum*, *Chloride of calcium*, (F.) *Chlorure de calcium*, *Muriate ou Hydrochlorate de Chaux*. This salt has been given, in solution, as a tonic, stimulant, &c., in scrofulous tumours, glandular obstructions, general debility, &c. A *Solutio Muriat'is Calcis*, *Liquor Calcis Muriat'is*, *Solutio of Muriate of Lime*, *Liquid Shell*, may be formed of *Muriate of Lime*, ℥j; dissolved in distilled water, f℥ij. The LIQUOR CALCII CHLORIDUM or *Solutio of Chloride of Calcium*, of the Pharmacopœia of the United States, is prepared as follows:—*Marble*, in fragments, ℥ij, *Muriatic acid*, Oj; *Distilled water*, a sufficient quantity. Mix the acid with a half pint of the water, and gradually add the marble. Towards the close of the effervescence apply a gentle heat, and, when the action has ceased, pour off the clear liquor and evaporate to dryness. Dissolve the residuum in its weight and a half of distilled water, and filter. Dose, from grt. xxx to f℥j, in a cupful of water.

CALCIS OXYMURIAS, Calcis chloridum—c. Phosphas, see Cornu cervi—c. Phosphas præcipitatum, see Cornu ustum.

CALCIS SULPHURETUM, *Calcii sulphure'tum*, *Hepar Calcis*, *Sulphuret of Lime*, *S. of Calcium*, *Hydrosulphate of Lime*, (F.) *Proto-hydrosulphate de Calcium*, *Hydrosulphate de chaux*. Principally used in solution, as a bath, in itch and other cutaneous affections.

CALCITEA, Ferri sulphas,

CALCITEOSA, Plumbi oxydum semivitreum.

CALCITHOS, Cupri subacetas.

CALCITRAPA, Centaurea Calcitrapa, Delphinium consolida—c. Hippophastrum, Centaurea calcitrapa—c. Stellata, Centaurea calcitrapa.

CALCIUM, CHLORIDE OF, Calcis murias—c. Chlorure de, Calcis murias—c. Chlorure d'oxide de, Calcis chloridum—c. Protohydrochlorate de, Calcis sulphuretum—c. Protochlorure de, Calcis chloridum—c. Protochloruret of, Calcis chloridum—c. Protoxide of, Calx—c. Sulphuret of, Calcis sulphuretum.

CALCO-SUBPHALANGÆUS MINIMI DIGITI, Abductor minimi digiti pedis—c. Sulphalangeus pollicis, Abductor pollicis pedis.

CALCOCOS, Bell-metal.

CALCHOIDEA, (ossicula,) Cuneiform bones.

CALCOTAR, Ferri sulphas.

CALCUL, Calculus.

CALCULEUX, Calculous.

CALCULI, see Calculus—c. Articular, see Calculi, arthritic; and Concretions, articular.

CALCULI, ALTERNATING, see Calculi, urinary.

CALCULI, ARTHRITIC, *Tophi*, *Tuber'cula arthritica*, *Calculi arthritici* seu *podagrichi*, *Chalk-stones*, *Gout-stones*, *Nodis*, (F.) *Pierres crayeuses*, *Calculs arthritiques*, *Nœuds*. Concretions, which form in the ligaments, and within the capsules of the joints, in persons affected with gout. They are composed of uric acid, soda, and a little animal matter; very rarely, urate of lime and chloride of sodium are met with. Similar calculi are found in other parts beside the joints.

CALCULI, BILIARY, *Calculi bili'osi* seu *fellei* seu *biliarii*, *Biliary Concretions*, *Gall-stones*, *Chololithus*, *Cholelithus*, (F.) *Calculs biliâires*, *Pierres au fiel*, *Angichololithe*,—*Piorry*. Some of these contain all the materials of the bile, and seem to be nothing more than that secretion thickened. Several contain *Pieromel*; and the greater part are composed of from 88 to 94 parts of *Cholesterin*, and of from 6 to 12 of the yellow matter of the bile. Biliary calculi are most frequently found in the gall-bladder: at other times,

In the substance of the liver, in the branches of the *Ductus hepaticus*, or in the *Ductus Communis Caudoductus*. The first are called *Cystic*; the second *Hepatic*; and the last, sometimes, *Hepatocystic*. The causes which give rise to them are very obscure. Often they occasion no uneasiness, and at other times the symptoms may be confounded with those of hepatitis. At times, they are rejected by the mouth, or by the bowels, along with a considerable quantity of bile, which had accumulated behind them; at other times they occasion violent abdominal inflammation, abscesses, and biliary fistulae, rupture of the gall-bladder, and fatal effusion into the peritoneum. The passage of a gall-stone is extremely painful; yet the pulse is not at first affected. Antiphlogistics, when there is inflammatory action, and strong doses of opium, to allay the pain and spasm, with the warm bath, are the chief remedies. Solvents are not to be depended upon. They cannot reach the calculi.

CALCULI BONE EARTH, see Calculi, urinary—c. Compound, see Calculi, urinary—c. Cystic, see Calculi, urinary.

CALCULI OF THE EARS, (F.) *Calculi de l'Oreille*. Hard, light, and inflammable concretions, which occur in the *meatus auditorius externus*, and are merely indurated cerumen. They are a frequent cause of deafness. They can be easily seen, and may be extracted by appropriate forceps, after having been detached by injections of soap and water.

CALCULI FELLEI, Calculi, biliary—c. Fusible, see Calculi, urinary.

CALCULI, LACHRYMAL, (F.) *Calculi lacrymaux*. Concretions sometimes, but rarely, form in the lachrymal passages, where they occasion abscesses and fistulae, which do not heal until they are extracted. No analysis has been made of them.

CALCULI, LITHIC, see Calculi, urinary.

CALCULI OF THE MAMMÆ, (F.) *Calculi des Mamelles*. Haller gives a case of a concretion, of a yellowish-white colour, which had the shape of one of the excretory ducts of the mammary gland, having been extracted from an abscess seated in that organ.

CALCULI, MULBERRY, see Calculi, urinary.

CALCULI OF THE PANCREAS, (F.) *Calculi du Pancréas*. These are but little known. Analogy has induced a belief that they resemble the salivary. Some have supposed that certain transparent calculi, rejected by vomiting, or passed in the evacuations, have proceeded from the pancreas, but there seems to be no reason for this belief.

CALCULI OF THE PINEAL GLAND, (F.) *Calculi de la Glande Pinéale*. These have been frequently met with. No symptom announces their presence during life. They are composed of phosphate of lime.

CALCULI PODAGRICI, Calculi, arthritic.

CALCULI OF THE PROSTATE, *Prostat'ic calculi*. These are not very rare. They have generally the same composition as the preceding. They usually present the symptoms common to every tumefaction of the prostate, and sometimes those of calculi in the bladder.

CALCULI PULMONARY, (F.) *Calculi pulmonaires*. These concretions are very frequently met with in the dead body, without seeming to have produced unpleasant symptoms during life. At other times, they are accompanied with all the symptoms of phthisis, *Litho-phthisis*, *Phthisis calculeuse*, of Bayle. See *Lithiasis pulmonum*. At times they are expectorated without the supervention of any unpleasant symptom. They

are usually formed of carbonate of lime and animal matter.

CALCULI, SALIVARY, *Calculi salivales*, *Sialolithi*, (F.) *Calculi salivaires*. Concretions, usually formed of phosphate of lime and animal matter, which are developed in the substance of the salivary glands or in their excretory ducts. In the first case, they may be mistaken for a simple swelling of the gland; in the second, they may generally be detected by the touch. They may be extracted by incision in the interior of the mouth. The calculus developed in the sublingual ducts has been called *Calculus sublingualis* and *Ranula lapide'a*.

CALCULI, SPERMATIC, (F.) *Calculi spermaticus*. These have been sometimes found in the vesiculæ seminales after death. They cannot be detected during life. No analysis has been made of them.

CALCULI OF THE STOMACH AND INTESTINES, *Enterolithus*, *E. Calculus*, *Coprolithus*, *Concretiones albiæ*, (F.) *Calculi de l'estomac*, *C. intestinaux* ou *stercoraux*, *Pierres stercorales*, *Concretions intestinales*. Calculi of the stomach are rare, and have almost always been carried thither by the antiperistaltic action of the intestines. The symptoms occasioned by them are those of chronic gastritis. It has been imagined that the continued use of absorbent powders, as magnesia, will give occasion to them.

Intestinal concretions, (F.) *Calculi intestinaux*, are not uncommon in animals (see BEZOARD:) but they are rare in man. The causes which give rise to them are little known: sometimes a biliary calculus affords them a nucleus. Their composition varies. They are light, hard, very fetid, and not inflammable. They are formed, ordinarily, between the valvulæ of the small intestines, or in the cells of the large, and sometimes in old herniæ. Whilst they do not obstruct the passage of the alimentary mass, they produce no unpleasant symptoms. At times, the movable tumour which they form may be felt through the parietes of the abdomen. They are generally evacuated *per anum*.

CALCULI OF THE TONSILS. Calculous concretions, which sometimes form in the tonsils. (F.) *Calculi des Amygdales*. They are easily recognised by the sight and touch: sometimes they are discharged by spitting, either alone or with the pus of an abscess occasioned by their presence. They have not been analyzed.

CALCULI, TRIPLE, see Calculi, urinary—c. Uric, see Calculi, urinary.

CALCULI, URINARY, *Urolithi*, (F.) *Calculi urinares*, *Pierres urinares*. Concretions which form from the crystallizable substances in the urine, and which are met with not only in the whole course of the urinary passages, but in fistulous openings wherever the urine stagnates naturally or accidentally. Their causes are but little known. They are more common at the two extremities of life than at the middle, and more so in some countries and districts than in others. At times, a clot of blood, a portion of mucus, &c., form the nucleus. The symptoms and treatment vary according to the seat of the calculus. There is no such thing probably as a medical solvent. See Urinary Calculi.

Chymists have demonstrated the existence of several components of urinary calculi, viz., *Lithic Acid*, *Phosphate of Lime*, *Ammonio-Magnesian Phosphate*, *Oxalate of Lime*, *Cystic Oxide*, and *Xanthic Oxide*, with an animal cementing ingredient. The varieties of calculi, produced by the combination or intermixture of these ingredients, are thus represented by Dr. Paris.

A TABULAR VIEW OF DIFFERENT SPECIES OF URINARY CALCULI.

SPECIES OF CALCULI.	EXTERNAL CHARACTERS.	CHYMICAL COMPOSITION.	REMARKS.
1. LITHIC OR URIC.	Form, a flattened oval. S. G. generally exceeds 1.500. Colour, brownish or fawn-like. Surface, smooth. Texture, laminated.	It consists principally of <i>Lithic Acid</i> . When treated with nitric acid, a beautiful pink substance results. This calculus is slightly soluble in water, abundantly so in the pure alkalis.	It is the prevailing species; but the surface sometimes occurs finely tuberculated. It frequently constitutes the <i>nucleus</i> of the other species.
2. MULBERRY. (F.) <i>Calculus muricatus</i> .	Colour, dark brown. Texture, harder than that of the other species. S. G. from 1.428 to 1.976. Surface, studded with tubercles.	It is <i>oxalate of lime</i> , and is decomposed in the flame of a spirit lamp swelling out into a white efflorescence, which is <i>quicklime</i> .	This species includes some varieties, which are remarkably smooth and pale-colored, resembling <i>hemispheres</i> .
3. BONE EARTH	Colour, pale brown or gray; surface, smooth and polished; structure, regularly laminated; the laminae easily separating into concrete crusts.	Principally <i>phosphate of lime</i> . It is soluble in muriatic acid.	
4. TRIPLE.	Colour, generally brilliant white. Surface, uneven, studded with shining crystals, less compact than the preceding species. Between its laminae small cells occur, filled with sparkling particles.	It is an <i>ammoniaco-magnesian phosphate</i> , generally mixed with phosphate of lime. Pure alkalis decompose it, extracting its ammonia.	This species attains a larger size than any of the others.
5. FUSIBLE.	Colour, grayish white.	A compound of the two foregoing species.	It is very fusible, melting into a vitreous globule.
6. CYSTIC.	Very like the triple calculus, but it is unstratified and more compact and homogenous.	It consists of <i>cystic oxide</i> . Under the blowpipe it yields a peculiarly fetid odour. It is soluble in acids, and in alkalis, even if they are fully saturated with carbonic acid.	It is a rare species.
7. ALTERNATING.	Its section exhibits different concentric laminae.	Compounded of several species, alternating with each other.	
8. COMPOUND.	No characteristic form.	The ingredients are separable only by chymical analysis.	

1. *Renal Calculi*, (F.) *Calculus rénauz*. These have almost always a very irregular shape: at times, there is no indication of their presence: at others, they occasion attacks of pain in the kidneys, sometimes accompanied with bloody or turbid urine. Often, they cause inflammation of the kidneys, with all its unpleasant results. They are generally formed of uric acid, animal matter, and oxalate of lime, with, sometimes, phosphates. The treatment will have to vary, according to the absence or presence of inflammatory signs,—relieving the irritation by opiates. A surgical operation can rarely be applicable.

2. *Calculi of the Ureters*, (F.) *Calculus des Urètres*. These come from the kidneys, and do not produce unpleasant effects, unless they are so large as to obstruct the course of the urine, and to occasion distension of the whole of the ureters above them; or unless their surface is so rough as to irritate the mucous membrane, and occasion pain, hemorrhage, abscesses, &c. The pain, during the passage, is very violent, extending, at times, to the testicle of the same side in the male; and occasioning a numbness of the thigh in both sexes. The treatment consists in general or local blood-letting, warm bath, and opiates.

3. *Calculi, Vesical; Stone in the Bladder*, *Lith in Vesica'lis*, *Lith'asis cystica seu vesica'lis*, *Cysto-lith'asis*, *Dysuria calculosa seu irritata*, *Calculus vesicæ*, (F.) *Calculus vésicaux*. These are the most common. Sometimes, they proceed from the kidneys: most commonly, they are formed in the bladder itself. Sense of weight in the perinæum, and sometimes of a body rolling when the patient changes his position; pain or itching at the extremity of the glans in men; frequent desire to pass the urine; sudden stoppage to its flow; and bloody urine—are the chief

signs which induce a suspicion of their existence. We cannot, however, be certain of this without sounding the patient. Sometimes, when of a small size, they are expelled: most commonly, they remain in the bladder, the disorganization of which they occasion, unless removed by a surgical operation.

4. *Calculi, Urethral*. These almost always proceed from the bladder. The obstruction, which they cause to the passage of the urine, the hard tumour, and the noise occasioned when struck by a sound, indicate their presence. They are removed by incision.

5. *Calculi of Fistulous passages*. These arise when there is some fistulous opening into the urethra. They can be readily recognised, and may generally be extracted with facility. (F.) *Calculus placés hors des voies urinaires*. See Urinary Calculi.

CALCULI OF THE UTERUS, (F.) *Calculus de l'Utérus*. These are very rare. The signs, which indicate them during life, are those of chronic engorgement of the uterus. Their existence, consequently, cannot be proved till after death.

CALCULIFRAGUS, Lithontriptic.

CAL'CULOUS, (F.) *Calculoux, Graveloux*. That which relates to calculi, especially to those of the bladder.

CALCULOS' OXYD, Oxyd, cystic.

CALCULS (F.), pl. of *Calcul*, a calculus.

CALCULS BILIAIRES, Calculi, biliary—*c. de l'Estomac*, Calculi of the stomach—*c. de la Glande Pinéale*, Calculi of the pineal gland—*c. Intestinaux*, Calculi of the stomach and intestines—*c. Lacrymaux*, Calculi, lachrymal—*c. des Mamelles*, Calculi of the mammae—*c. Mûrains*, see Calculi, urinary—*c. de l'Oreille*, Calculi in the ears—*c. du Pancreas*, Calculi of the Pancreas—*c. Placés hors des voies urinaires*, Calculi of fistulous passages—*c. Pulmonaires*, Calculi, pulmo-

nary — *c. Rénaux*, Calculi, renal — *c. Salivaires*, Calculi, salivary — *c. Spermatiques*, Calculi, spermatic — *c. Stercoraux*, see Calculi of the stomach and intestines — *c. Urinaires*, Calculi, urinary — *c. des Urètres*, Calculi of the ureters — *c. de l'Utérus*, Calculi of the uterus — *c. Vésicaux*, Calculi, vesical.

CALCULUS, *Lapis, Lithos*. A diminutive of *calx*, a lime-stone. (F.) *Calcul, Pierre*. Calculi are concretions, which may form in every part of the animal body, but are most frequently found in the organs that act as reservoirs, and in the excretory canals. They are met with in the tonsils, joints, biliary ducts, digestive passages, lachrymal ducts, mammae, pancreas, pineal gland, prostate, lungs, salivary, spermatic and urinary passages, and in the uterus. The causes which give rise to them are obscure.

Those that occur in reservoirs or ducts are supposed to be owing to the deposition of the substances, which compose them, from the fluid as it passes along the duct; and those which occur in the substance of an organ are regarded as the product of some chronic irritation. Their general effect is to irritate, as extraneous bodies, the parts with which they are in contact; and to produce retention of the fluid, whence they have been formed. The symptoms differ, according to the sensibility of the organ and the importance of the particular secretion whose discharge they impede. Their solution is generally impracticable: spontaneous expulsion or extraction is the only way of getting rid of them.

CALCULUS BEZOAR, Bezoard — *c. Dentalis*, Odontolithus — *c. Encysted, Calcul chatonné* — *c. Sublingualis*, see Calculi, salivary — *c. Vesicae*, Calculus, vesical.

CALDAS, WATERS OF. Caldas is a small town, ten leagues from Lisbon, where are mineral springs, containing carbonic and hydrosulphuric acid gases, carbonates and muriates of lime and magnesia, sulphates of soda and lime, sulphuret of iron, silica, and alumina. They are much used in atonic gout. They are thermal. Temperature 93° Fahrenheit.

CALDERIÆ ITALIÆ. Warm baths in the neighbourhood of Ferrara, in Italy, much employed in dysuria.

CALEBASSES, *Cucurbita lagenaria*.

CALEDONIA SPRINGS. These are in Pennsylvania, fifteen miles from Chambersburg. The waters are pure; temperature 52° Fahr.

CALEDONIA SPRINGS, see Canada, mineral Waters of.

CALEFA'CIENTS, *Calefacien'tia, Therman'tica*, from *calidus*, 'warm,' and *facio*, 'I make.' (F.) *Échauffante*. Substances which excite a degree of warmth in the part to which they are applied, as mustard, pepper, &c. They belong to the class of stimulants.

CALEFACTIO, *Échauffement*.

CALENDULA ALPINA, *Arnica montana*.

CALENDULA ARVEN'SIS, *Caltha Arven'sis* seu *officina'lis*, Wild Mar'igold, (F.) *Souci des Champs*. Order, Compositæ. This is, sometimes, preferred to the next. Its juice has been given, in the dose of from $\mathfrak{f}\mathfrak{ss}$ to $\mathfrak{f}\mathfrak{ss}\mathfrak{iv}$, in jaundice and cachexia.

CALENDULA OFFICINA' LIS, *C. Sati'va, Oxy-santhemum, Sponsa solis, Caltha vulgari's; Ver-rucaria*, Single or Garden Mar'igold, (Prov.) *Golding*, (F.) *Souci, S. ordinaire, Fleur de tous les mois*. Family, Syntantheræ, Syngenesia necessaria, Linn. So called from flowering every calend. The flowers and leaves have been exhibited as aperients, diaphoretics, &c., and have been highly extolled in cancer.

CALENDULÆ MARTIALES, Ferrum ammoniatum.

CALENTURA, *Cal'enture*, from *calere*, 'to be warm.' The word, in Spanish, signifies fever. A species of furious delirium to which sailors are subject in the torrid zone:—a kind of phrenitis, the attack of which comes on suddenly after a broiling day, and seems to be characterized by a desire in the patient to throw himself into the sea. It is only a variety of phrenitis.

CALENTURA CONTINUA, Synocha.

CALENTURAS; *Palo de Calentu'ras*. Pomet and Lémery say, that these words are sometimes applied to cinchona. Camelli says, they mean, also, a tree of the Philippine Isles, the wood of which is bitter and febrifuge.

CALF KNEE, Entogonynecon — *c. of the Leg*, Sura.

CALICE, Calix.

CALICES RÉNALES, see Calix.

CALICO BUSH, *Kalmia latifolia*.

CALIDARIUM, see Stove.

CALIDUM ANIMALE, Animal heat — *c. Innatum*, Animal heat.

CALIFORNIA, MINERAL WATERS OF. Several mineral springs exist in California, near 40° N. Lat. and 40½° longitude west of Washington, directly upon the California trail, leading from the sink of Humboldt river to Pyramid lake. Their temperature varies from cold to almost boiling. They ebb and flow at irregular intervals; have not been analyzed; but some have the taste of a strong solution of sulphate of magnesia. They have had the singular effect of inducing blennorrhæal symptoms, both in man and animals.

CALIGATIO, Dazzling.

CALIGO. 'A mist.' *Achlys*, (F.) *Brouillard*. An obscurity of vision, dependent upon a speck on the cornea: also, the speck itself; *Caligo cor-neæ* seu *a Nephel'io* seu *a Lenc'o'natē*, *Mac'ula cor-neæ* seu *semipell'u'cida*, *Phtharma caligo*, *Hebetu'do visus*, *Neb'ula*, *Opake cornea*, *Web-eye*, *Web*, *Pin-and-Web*, (F.) *Nouage de la Cornée*, *Taie ou Tays*, *Obscurcissement de la vue*, *Akéra-todiaphanie*, — Piorry.

CALIGO LENTIS, Cataract — *c. Pupillæ*, Syne-zisis — *c. Synzesis*, Syne-zisis — *c. Tenebrarum*, Hemeralopia.

CALIHACHA CANELLA, *Laurus cassia*.

CALISAYA BARK, *Cinchonæ cordifoliæ* cortex.

CALISTHENICS, *Callisthen'ics, Calisthen'i'a, Callisthen'i'a*, (F.) *Callisthénie*, from *kalos*, 'beautiful, and *sthenos*, 'strength.' The art of promoting, by appropriate exercises, strength of body, and grace of movements.

CALIX, *Calyx, Infundib'ulum*, from *καλῆς*, 'a cup,' 'a flower cup.' (F.) *Calice, Entonnoir*. Anatomists have given this name to small cup-like membranous canals, which surround the papillæ of the kidney, and open into its pelvis, whither they convey the urine:—*Calices renal'es, Cylind'ri membrana'cei Renum, Fis'tulæ ure'terum renum, Canales membrana'ci Renum, Tri'buli pelvis renum*. Their number varies from 6 to 12 in each kidney.

CALIX VOMITORIA, Goblet, emetic.

CALLA PALUS'TRIS, *Water Arum*; indige-nous. Ord. Araceæ: has similar virtues to those of the Arums.

CALLA VIRGINICA, *Peltandra Virginica*.

CALLEUX, Callous.

CALLIBLEPH'ARUM, from *καλλος*, 'beauty, and *βλεφαρον*, 'eyelid.' A remedy for beautify-ing the eyelids.

CALLICANTHUS, *Calycanthus*.

CALLICAR'PA AMERICA'NA, *French Mulberry*; indigenous. *Order*, Verbenaceæ. Used as a euphotic in cutaneous diseases. Said to be useful in dropsy.

• **CALLICOCCA IPECACUANHA**, *Ipecacuanha*.

CALLICREAS, *Pancreas*.

CALLIOMARCHUS, *Tussilago*.

CALLIPEDIA, from *καλλος*, 'beauty,' and *παις*, *παῖς*, 'a child.' The art of begetting beautiful children. This was the title of a poem by Claude Quillet, in 1655; "*Callipedia sive de pulchra proles habenda ratione*." The author absurdly supposes, that the beauty of children is affected by the sensations which the mother experiences during her pregnancy.

• **CALLIPERS OF BAUDELOCQUE**, see *Pelvimeter*.

CALLIPESTRIA, *Cosmetics*.

CALLIPHYLLUM, *Asplenium trichomanoides*.

CALLIPYGOS, from *καλλος*, 'beauty,' and *πυγή*, 'buttocks.' A cognomen of Venus, owing to her beautiful nates.

CALLISTHENICS, *Calisthenics*.

CALLITRICHE AQUATICA, *C. Verna* — *c. Heterophylla*, *C. Verna* — *c. Intermedia*, *C. Verna*.

CALLITRICHE VERN, (*καλλος*, 'beautiful,' and *δαίς*, *τριχός*, 'hair,' from its capillary and tufted stems.) *C. aquatica seu intermedia seu heterophylla*, *Water chickweed*, *Water starwort*: indigenous; *Order*, *Callitrichaceæ*; is diuretic, and used in dropsy, in the form of tincture or of decoction.

CALLITRIS ECKLO'NI. A South African tree, *Ord. Coniferae*, from the branches and cones of which a gum exudes, that resembles Gum Sandarac. This is successfully used in the form of fumigations in gout, rheumatism, cedematous swellings, &c.

CALLITRIS CUPRESSOIDES, a common shrub in the neighbourhood of Cape Town, exudes a similar substance.

CALLOSITAS, *Induration* — *c. Palpebrarum*, *Scleriosis* — *c. Vesicæ*, *Cystauxe*.

CALLOSITY, *Callositas*, *Scyros*, *Tylē*, *Tylus*, *Tylo'ma*, *Tylo'sis*, *Dermatoclerosis*, *Dermatotylo'ma*, *Dermatotylo'sis*, *Dermatotylo'gylus*, *Porus*, *Ephy'ma Callus*. Hardness, induration, and thickness of the skin, which assumes a horny consistence in places where it is exposed to constant pressure. (F.) *Durillon*. Also the induration, which is observed in old wounds, old ulcers, fistulous passages, &c.

CALLOUS, *Callo'sus*, *Ochtho'des*, from *callus*, 'hardness.' (F.) *Callex*. That which is hard or indurated. A *Callous Ulcer* is one whose edges are thick and indurated.

CALLUNA ERICA, *Erica vulgaris* — *c. Vulgaris*, *Erica vulgaris*.

CALLUS, *Calus*, *Callum*, *Osteo'gylus*, (F.) *Cal*. The bony matter, thrown out between the fractured extremities of a bone, which acts as a cement, and as a new bony formation. The words are, likewise, used occasionally in the same sense as *Callosity*.

CALLUS, PROVISIONAL. When the shaft of a long bone has been broken through, and the extremities have been brought in exact juxtaposition, the new matter, first ossified, is that which occupies the central portion of the deposit, and thus connects the medullary cavities of the broken ends, forming a kind of plug, which enters each. This was termed by M. Dupuytren the *provisional Callus*.

CALMANTS, *Sedatives*.

CALME, (F.) The interval that separates the paroxysms of an acute or chronic disease. When the type is intermittent, the word *intermission* is used.

CALOMBA, *Calumba*.

CAL'OMEL, from *καλος*, 'good,' and *μελος*, 'black.' A term, formerly applied to *Astibeg mineral*. By some, the mild chloride of mercury is said to have been so called by Sir Theodora Turquet de Mayerne, in consequence of its having been prepared by a favourite black servant, or, according to others, because it was a good remedy for black bile. It is now applied to the mild chloride of mercury only. See *Hydrargyri submuriæ*.

CALOMEL STOOLS. A term applied to the green, spinach-like, evacuations occasioned by the internal use of the mild chloride of mercury.

CALOMEL'ANOS TURQUETI, *Hydrargyri submuriæ*. A name, also, given by Riverius to purgative pills, prepared with calomel, sulphur, and resin of jalap. — *Dictionaries*.

CALOMELAS, *Hydrargyri submuriæ*.

CALO'NIA, *καλονία*. An epithet formerly given to myrrh. — *Hippocrates*. See *Myrrha*.

CALOPHYLLUM INOPHYLLUM, see *Fagara octandra*.

CALOR, *Heat* — *c. Animalis*, *Animal heat* — *c. Nativus*, *Animal heat*.

CALORICITÉ, (F.) *Caloric'itas*. The faculty possessed by living bodies of generating a sufficient quantity of caloric to enable them to resist atmospheric cold, and to preserve, at all times and in every part, a temperature nearly equal. See *Animal Heat*.

CALORIFACIENT, *Calorifi'ant*, *Calorificiens*, *Calorifi'ans*: from *calor*, 'heat,' and *facere*, 'to make.' Having the power of producing heat. Relating to the power of producing heat.

CALORIFICA'TION, *Calorifica'tio*, from *calor*, 'heat,' and *fieri*, 'to be made.' The function of producing animal heat.

CALORINÆSES, from *calor*, 'heat.' The name under which M. Baumes proposes to arrange all diseases characterized by a sensible change in the quantity of animal heat. The *Calorinæses* form the first class of his Nosology.

CALOTROPIS GIGANTEA, *Mudar* — *c. Mudarii*, *Mudar*.

CALOTTE, (F.) *Pile'olum*. Anatomists sometimes give the name, *Calotte aponeurotique*, to the aponeurosis of the occipito-frontalis muscle, which covers it externally; and that of *Calotte du crâne* to the *scalp-cap*.

Calotte is also applied to an adhesive plaster, with which the head of a person labouring under *tinea capitis* is sometimes covered, after the hair has been shaved off. This plaster is pulled suddenly and violently off, in order to remove the bulbs of the hair. It means, also, a sort of cap made of boiled leather, worn by those who have undergone the operation of trepanning, &c.

CALOTTE D'ASSURANCE, *Condom*.

CALTHA ALPINA, *Arnica montana* — *c. Arvensis*, *Calendula arvensis* — *c. Marsh*, *C. Palustris* — *c. Officinalis*, *Calendula arvensis*.

CALTHA PALUSTRIS, *Marsh Mar'igold*, *Marsh Caltha*, *Cowslips*. *Order*, *Ranunculaceæ*: indigenous. A syrup prepared from it is a popular cough remedy.

CALTHA VULGARIS, *Calendula officinalis*.

CALTROPS, see *Trapa natans*.

CALUM'BA, *Calom'ba*, *Calom'ba*, *Calom'ba*, (Ph. U. S.): *Columbo*, *Radic Columba*. (F.) *Columbe* ou *Columbe*. The root of *Menispermaceæ*.

palma'tum, Cocculus palma'tus. Order, Menispermaceæ: indigenous in Mozambique. Its odour is slightly aromatic; taste unpleasantly bitter. It is tonic and antiseptic. Dose, gr. 10 to 5j in powder.

CALUMBA, AMERICAN, Frase'ra Walteri seu *Carolinensis* seu *Officinalis* seu *verticillata*, *Sce'r'tia difformis* seu *Frase'ra, American* or *Marietta Columbo, Indian Lettuce, Yellow Gentian, Golden Seal, Meadow pride, Pyr'amid.* The root *Frase'ra* (Ph. U. S.) is used in the same cases as the true Calumba.

CALUS, Callus.

CALVA, Cranium.

CALVA, Calva'ria. The cranium; the upper part especially; the skull-cap;—the *Vault of the Cranium, Cam'era.*

CALVARIA, Cranium.

CALVA'TA FERRAMEN'TA. Surgical instruments, which have a head or button.

CALVIT'IES, Calvit'ium, Phal'acia, Phalarosis, Glabrit'ies, Ophi'asis, Depilatio Capitis, Phalaro'ma, Madaro'sis, Lipsotrich'ia, Baldness, (Old Eng.) *Baldness, Cal'city, &c., from calvus, 'bald,' (F.) Calvitie, Chauvete.* Absence of hair, particularly at the top of, and behind, the head.

CALVIT'IES PALPEBRA'RUM is loss of the eyelashes.

CALVUS, Athrix.

CALX, Lime, Cal'rium Terræ, Protox'ide of Calcium, Calca'ria pura, (F.) Chaux. The lime, employed in pharmacy, should be recently prepared by calcination. When water is sprinkled over caustic lime, we have *slaked lime, hydrate of lime,*—the *Calcis Hydras* of the London pharmacopœia.

CALX, see Calcanæum — c. Bismuthi, Bismuth, subnitrate of — c. Chlorinata, Calcis chloridum — c. Cum kali puro, Potassa cum calce — c. Salita, Calcis murias — c. Oxymuriatica, Calcis chloridum.

CALX E TESTIS; lime prepared from shells. It has probably no medicinal advantages over that prepared from marble.

CALX VIVA, C. recens seu *usta, Calx et Calx viva, Ox'idum Cal'ci, Fumans nix, Lime* or *Quicklime, (F.) Chaux vive.* The external operation of calx viva is escharotic, but it is rarely used. Lime is a good disinfecting agent. It is employed internally in the form of Liqueur Calcis.

CALYCANTH'US, C. Flor'idus, Callican'thus, [?] Carol'na Allspice, Sweet-scented shrub, Sweet shrub, from calyx, 'a flower-cup,' and andos, 'a flower.' An indigenous plant; Order, Calycanthaceæ; with purplish flowers, of strong, agreeable odour, which appear from March to June. The root is possessed of emetic properties.

CALYPTRANTHES CARYOPHYLLATA, Myrtus caryophyllata.

CALYSTEGIA SEPIUM, Convolvulus sepium — c. *Soldanella, Convolvulus soldanella.*

CALYX, Calix.

CAMARA, Calva.

CAMARÉZ, MINERAL WATERS OF. Camaréz is a small canton near Sylvanès, in the department of Aveyron, France, where there are acidulous chalybeates.

CAMARO'SIS, Camaro'ma, from kapapa, 'a vault,' Camera'tio, Testudina'tio Cra'nii. A species of fracture of the skull — *cam'rated fracture*—in which the fragments are placed so as to form a vault, with its base resting on the dura mater. — Galen, Paulus of Ægina.

CAMBE-LEAF, Nymphaea alba et lutea.

CAMBING. A tree of the Molucca Islands, from the bark of which a kind of gum-resin ex-

udes, which has been highly extolled in dysentery. It appears to have some resemblance to the simarouba — Rumphius.

CAMBIUM, 'Exchange.' A name formerly given to a fancied nutritive juice, which was supposed to originate in the blood, to repair the losses of every organ, and produce their increase. — Sennertus.

CAMBO, MINERAL WATERS OF. A village in the department of Basses Pyrénées, France, where there are two mineral springs; the one an acidulous chalybeate, the other sulphureous. Temperature, 62° to 69° Fahrenheit.

CAMBODIA, Cambogia.

CAMBO'GIA, from Cambodia, in the East Indies, where it is obtained. Hence, likewise, its names *Cambol'dia, Cambol'gium, Cambol'gia, Cambol'gium, Gamba'gium, Gamba'dia.* It is called also, *Gutta, Gutta gamba* seu *Gambandra, Gummi Gutta, Cataguan'na, Cataguan'na, Chrys'osopus, Laxat'ivus Ind'icus, Gummi Bo'gia* seu *gambal'dra* seu *de Goa* seu *de Jenu* seu *ad Pod'agram, Chitta jemoco, Camboe* or *Gamboge, (F.) Gomme Gutte.* Ord. Guttiferae. A yellow juice obtained from *Hebradendron Cambogioi'des*, and other plants of the natural family Guttiferae, but it is not known from which of them the official camboge is obtained. It is inodorous, of an orange yellow colour; opaque and brittle; fracture, glassy; is a drastic cathartic, emetic and anthelmintic; and is used in visceral obstructions and dropsy, and wherever powerful hydragogue cathartics are required. Dose from gr. ij to vi, in powder, united with calomel, squill, &c.

CAMBODIA GUTTA, Garcinia cambogia.

CAMBU'CA, Cambuc'ca membra'ta. Buboes and venereal ulcers, seated in the groin or near the genital organs. — Paracelsus. See Bubo.

CAMELÉE, Cneorum tricocum.

CAMERA, Chamber, Fornix, Vault — c. *Cordia, Pericardium* — c. *Oculi, Chamber of the eye.*

CAMERATIO, Camarosis.

CAMFOROSMA, Camphorosma.

CAMINGA, Cavella alba.

CAMISIA FETÛS, Choron.

CAMISOLE, Waistcoat, strait.

CAMMARUS, Crab.

CAMOMILLE FÉTIDE, Anthemis cotula — c. *Puante, Anthemis cotula* — c. *Romane, Anthemis nobilis* — c. *des Teinturiers, Anthemis tinctoria* — c. *Vulgaire, Matricaria chamomilla.*

CAMOSIERS, WATERS OF. Camosiers is a canton, two leagues from Marseilles, where are two springs containing carbonate of lime, sulphur, chloride of sodium, &c. They are purgative, and used in skin complaints.

CAMOTES, Convolvulus batatas.

CAMPAGNE, MINERAL WATERS OF. Campagne is in the department of Aude, France. The waters contain sulphate and chlorhydrate of magnesia. Temperature, 80° Fahrenheit.

CAMPAN'ULA. Diminutive of *campana*, a bell.

CAMPANULA TRACHE'LIUM, Canterbury Bell or *Throatwort*, was formerly used, in decoction, in relaxation of the fauces. It is, also, called *Cervicaria.*

CAMPE, Flexion.

CAMPHENE. The radical of camphor, which, according to Dumas, consists of camphene and one equivalent of oxygen. Camphene or *camphine* is represented by pure oil of turpentine. That which is used for burning in lamps, and is occasionally employed therapeutically, is oil of turpentine purified by distillation from a solution of caustic potassa.

CAMPHIRE, Camphor.

CAMPHOR, from Arab. *Ca'phur* or *Kan'phur*, *Camphora*, *Caphura*, *Cajja*, *Caf*, *Cafur*, *Caphora*, *Altafor*, *Camphire*, (F.) *Camphre*. A concrete substance, prepared, by distillation, from *Laurus Camphora*, *Persea Camfura*, *Campona officinarum*, an indigenous tree of the East Indies. Order, Laurineæ. Its odour is strong and fragrant: it is volatile, not easily pulverizable; texture crystalline. Soluble in alcohol, ether, oils, vinegar, and slightly so in water. Its properties are narcotic, diaphoretic, and sedative. Dose, gr. v to ℥j. Dissolved in oil or alcohol, it is applied externally in rheumatic pains, bruises, sprains, &c.

CAMPHOR WATER, *Mistura Camphoræ*.

CAMPHORA'CEOUS, *Camphora'ceus*. Relating to or containing camphor;—as a 'camphoraceous' smell or remedy.

CAMPHORATA HIRSUTA, and C. MONSPELIENSIS, *Camphorosma Monspelica*.

CAMPHORATED, *Camphoratus*, (F.) *Camphré*. Relating to camphor; containing camphor: as a *camphorated* smell, a *camphorated* draught.

CAMPHOROS'MA MONSPELIACA, *C. Perennis*, from *Camphor*, and *οσμη*, 'odour.' *Sela'go*, *Camphorata hirsuta* seu *Monspelien'sium*, *Hairy Camphoros'ma*, (F.) *Camphrée de Montpellier*. Order, Chenopodiaceæ. See *Syst. Tetrandria Monogynia*. This plant, as its name imports, has an odour of camphor. It is regarded as diuretic, diaphoretic, cephalic, antispasmodic, &c. It is also called *Chamaenecée* and *Stinking Ground Pine*.

CAMPHOROSMA PERENNIS, C. Monspelica.

CAMPHRE, Camphor.

CAMPHRE, Camphorated.

CAMPHRÉE DE MONTPELLIER, *Camphorosma Monspelica*.

CAMPION, ROSE, *Lychnis githago*.

CAMPOMANE'SIA LINEATIFOLIA. A tree, twenty to thirty feet high, which grows in Peru, and whose fruit—*palillo*—of a bright yellow colour, and as large as a moderate-sized apple, has an exceedingly agreeable scent, and is one of the ingredients in making the perfumed water called *mistura*.—Tschudi.

CAMPSIS, *Flex'io*, *Curva'tio*, *Inflex'io*. Bone or cartilage, forcibly bent from its proper shape, without breaking.—Good.

CAMPSIS DEPRESSIO, Depression.

CAMPYLORRHACHIS; from *καμπυλος*, 'crooked,' and *ραχis*, 'spine.' A monster whose spine is crooked.—Gurlt.

CAMPYLORRHINUS; from *καμπυλος*, 'crooked,' and *ριν*, 'nose.' A monster whose nose is crooked.—Gurlt.

CAMPYLOTIS, Cataclasis.

CAMPYLUM, Cataclasis.

CAMUS, (F.) *Simus*, *Res'mus*, *Simo*, *Silo*, *Silva*. One who has a short, stumpy nose. The French speak of *Naz camus*, 'short nose.'

CANADA BURNET, *Sanguisorba canadensis*.

CANADA, MINERAL WATERS OF. *St. Catherine's*, at the head of Lake Ontario, at the entrance of the Welland Canal, 12 miles from Niagara Falls, and on the Great Western Railroad, has a mineral spring, much frequented, which Dr. Chilton, of New York, analyzed and found to contain chlorides of calcium, magnesium, sodium, and iron, sulphate of lime, carbonates of lime and magnesia, bromide and iodide of magnesium, silica and alumina.

Tuscarora Acid Spring is in the township of

Tuscarora, about 20 miles north of Pass Dover. It contains a large amount of free sulphuric acid; beside sulphates of the alkalies, lime, magnesia, alumina and iron.

Charlottesville Sulphur Spring is a few miles from Dover, on Lake Erie. It contains an unusual quantity of sulphuretted hydrogen.

The *Caledonia Springs*—a well known watering place—are a few miles south of the Ottawa river, and 40 miles from Montreal. There are various sources; one—the *Gas Spring*—discharging carburetted hydrogen; a *Saline* and a *Sulphur Spring*; besides an *Intermitting Spring*, which discharges large quantities of carburetted hydrogen; and in which iodine and bromine have been detected.

CANAL, *Canalis*, *Ductus*, *Med'tus*, *Poros*, *Och'etos*, (F.) *Conduit*. A channel for affording passage to liquids, or solids, or to certain organs.

CANAL, ALIMEN'TARY, *C. Digestive*, *Canalis seu Ductus cibarius seu digestivus seu alimentarius*, *Tubus alimentarius seu intestinalis*, *Digestive Tube*, *Alimentary Duct or Tube*. The canal extending from the mouth to the anus.

CANAL, ARACH'NOID, *Canalis Bichat'ii*, *Canal of Bichat*. A canal, formed by the extension of the arachnoid over the transverse and longitudinal fissures of the brain, which surrounds the vena magna Galeni. The orifice of the canal has been termed the Foramen of Bichat.

CANAL ARTERIEL, Arterial duct—*c. de Bartholin*, *Ductus Bartholinus*—*c. of Bichat*, *Canal*, *arachnoid*—*c. Bullular*, of *Petit*, *Godronné canal*—*c. Carotidien*, *Carotid canal*—*c. Choldoquo*, *Choledoch duct*—*c. Ciliary*, *Ciliary canal*—*c. of Cotonnius*, *Aqueductus vestibuli*—*c. Défèrent*, *Deferens vas*—*c. of Fontana*, *Ciliary canal*—*c. Goudronné*, *Godronné canal*—*c. Hépatique*, *Hepatic duct*.

CANAL OF HUGUIER. A small canal at the inner side of the fissure of Glaser, through which the chorda tympani nerve leaves the cavity of the tympanum.

CANAL, HY'ALOID. A cylindrical passage, described by M. J. Cloquet as formed by the reflection of the hyaloid membrane into the interior of the vitreous body around the nutritious artery of the lens. M. Cruveilhier has never been able to see it.

CANAL, INCI'SIVE, see *Palatine canals*—*c. Infra-orbital*, *Suborbital canal*—*c. Inferece de Vos temporal*, *Carotid canal*—*c. Intermédiaire des ventricules*, *Aqueductus Sylvii*.

CANAL, INTES'TINAL, *Canalis seu Ductus intestinalis*. The portion of the digestive canal formed by the intestines.

CANAL OF JACOBSON, *Canal*, *tympanic*.

CANAL, MED'ULLARY. The cylindrical cavity in the body or shaft of a long bone, which contains the marrow.

CANAL, NASAL, *Lachrymal canal*.

CANAL OF NUCK. A cylindrical sheath formed around the round ligaments of the uterus by a prolongation of the peritoneum into the inguinal canal.

CANAL DE PETIT, *Godronné canal*—*c. Pulmo-aortique*, *Arterial duct*—*c. Rachidian*, *Vertebral canal*.

CANAL OF SCHLEMM. A minute circular canal, discovered by Professor Schlemm, of Berlin. It is situate at the point of union of the cornea and scleroticæ.

CANAL, SPINAL, *Vertebral canal*—*c. Spiræde de Vos temporal*, *Aqueductus Fallopii*—*c. de Senon*, *Ductus salivialis superior*—*c. Thoracique*, *Thoracic duct*—*c. Veineux*, *Canal*, *venous*—*c. Vulvo-uterine*, *Vagina*—*c. de Warthon*, *Ductus salivalis inferior*.

Fig. 5



ERYSIPELAS
ULCERATED.

Fig. 6



SIMPLE PURULENT
ULCER.

SCALD HEAD


Fig. 7



Fig. 8



OEDEMATOUS
SWOLLEN SORE LEG



CANAL, TYM'PANIC, *Canalis tympanicus*, *Canal of Jacobson*. A canal which opens on the lower surface of the petrous portion of the temporal bone, between the carotid canal and the groove for the internal jugular vein. It contains Jacobson's nerve.

CANAL, VECTOR, Tube, Fallopiun.

CANAL, VEINOUS, *Canalis seu Ductus venosus seu D. v. Aran'tii*, (F.) *Canal veinous*. A canal which exists only in the fœtus. It extends from the bifurcation of the umbilical vein to the vena cava inferior, into which it opens below the diaphragm. At times, it ends in one of the infra-hepatic veins. It pours into the cava a part of the blood, which passes from the placenta by the umbilical vein. After birth, it becomes a fibro-cellular cord.

CANAL OF WIRSIUNG, see Pancreas.

CANALE AERIFERI, see Bronchia.

CANALE BRESCHET'I, Canals in the diploë for the passage of veins; so called after M. Breschet.

CANALE CIRCULARES, Semicircular canals—c. Cochleæ, Scape of the cochlea—c. Lacrymales, Lacrymal ducts—c. Membrani renium, see Calix—c. Tubæformes, Semicircular canals.

CANALICULATED, Grooved.

CANALICULATUS, *Cannelé*, Grooved.

CANALICULE, Grooved.

CANALICULES CALCIFÈRES, Canaliculi calciferi.

CANALICULI, plural of Canaliculus—c. Bili-feri, Pori biliarii—c. of Bone, see Lacunæ of Bone.

CANALIC'ULI CALCIF'ERI, *Calciferous canals*, (F.) *Canalicules Calcifères*. Minute canaliculi, observed with the microscope, in the cartilages of bones, which are full of calcareous salts.

CANALICULI HAVERSIANI, Canals, nutritive—c. Lacrymales, Lacrymal ducts—c. Iimæum, Lacrymal ducts—c. Semicirculares, Semicircular canals—c. Seminales, see Testicle—c. Seminiferi, see Testicle—c. Vasculosi, Canals, nutritive.

CANALIC'ULUS, diminutive of *canalis*, 'a channel.' A small channel. See Canaliculi.

CANALIS, Meatus—c. Alimentarius, Canal, alimentary—c. Arteriosus, Arterial duct—c. Bichatii, Canal, arachnoid—c. Canaliculatus, Gorge—c. Caroticus, Carotid canal—c. Cervicis, see Uterus—c. Deferens, Deferens vas—c. Eminentie quadrigemine, Aquæductus Sylvii—c. Fallopii, Aquæductus Fallopii—c. Intestinorum, Intestinal tube—c. Lacrymalis, Lacrymal or nasal duct—c. Medius, Aquæductus Sylvii—c. Medullæ Spinalis, see Vertebral column—c. Nervous fistulosus renium, Ureter—c. Orbitæ nasalis, Lacrymal or nasal duct—c. Scalarum communis, Infundibulum of the cochlea—c. Semicircularis horizontalis, see Semicircular Canals—c. Semicircularis verticalis posterior, see Semicircular Canals—c. Semicircularis verticalis superior, see Semicircular canals—c. Tympanicus, Canal, tympanic—c. Uriauius, Urethra—c. Vidianus, Pterygoid canal.

CANALS OF HAVERS, Canals, nutritive—c. Haversian, Canals, nutritive.

CANALS, NUTRITIVE, *Canals for the nutrition of bones*, *Ductus nutritii*, *Canaliculi vasculosi seu Haversiani*, *Haversian Canals*, *Canals of Havers*, (F.) *Canaux nourriciers ou de Nutrition des os*, *Conduits nourriciers ou nutritiers*. The canals through which the vessels pass to the bones. They are lined by a very fine lamina of

compact texture, or are formed in the texture itself. There is, generally, one large nutritious canal in a long bone, situate towards its middle.

CANAPACIA, *Artemisia vulgaris*.

CANARIES, CLIMATE OF. The climate of the Canaries greatly resembles that of Madeira. That of the latter, however, is more equable, and the accommodation for invalids much superior.

CANARIUM COMMUNE, see *Amyris elemifera*.

CANARY-SEED, *Phalaris Canariensis*.

CANAUX AQUEUX, see Aqueous—c. *Demicirculaires*, Semicircular canals—c. *Éjaculateurs*, Ejaculatory ducts—c. *Nourriciers*, Canals, nutritive—c. *de Nutrition des os*, Canals, nutritive.

CANAUX DE TRANSMISSION. According to Bichat, the bony canals intended to give passage to vessels and nerves going to parts more or less distant—as the *Canalis Caroticius*, &c.

CANAUX VEINEUX, *Venous Canals*. The canals situate in the diploë, which convey venous blood.

CAN'CAMUM. A mixture of several gums and resins, exported from Africa, where it is used to deterge wounds. Dioscorides calls, by the name *κακκαρον*, the tears from an Arabian tree, which are similar to myrrh, and of a disagreeable taste. He advises it in numerous diseases. This name is given, also, to the Anime.

CANCAMY, Anime.

CAN'CELLATED, *Cancellatus*, (F.) *Cancellé*; from *Cancelli*, 'lattice-work.' Formed of cancelli, as the 'cancellated structure of bone.'

CANCEL'LI, 'Lattice-work.' The *Cellular* or *Spongy Texture of Bones*, (F.) *Tissu cellulaire*; consisting of numerous cells, communicating with each other. They contain a fatty matter, analogous to marrow. This texture is met with, principally, at the extremities of long bones; and some of the short bones consist almost wholly of it. It allows of the expansion of the extremities of bones, without adding to their weight, and deadens concussions.

CANCEL'LUS, from *cancer*, 'a crab.' A species of crayfish, called the *Wrong Heir*, and *Bernard the Hermit*: which is said to cure rheumatism, if rubbed on the part.

CANCER, 'a crab.' *Carcinosis*, *Lupus cancerosus*. A disease, so called either on account of the hideous appearance which the ulcerated cancer presents, or on account of the great veins which surround it, and which the ancients compared to the claws of the crab: called also *Carcinoma*. It consists of a scirrhus, livid tumour, intersected by firm, whitish, divergent bands; and occurs chiefly in the secretory glands. The pains are acute and lancinating, and often extend to other parts. The tumour, ultimately, terminates in a fetid and ichorous ulcer, — *Ulcus cancerosum*. It is distinguished, according to its stages, into *occult*, *Cancer occultus*, and *open*, *Cancer aper'tus*; the former being the scirrhus, the latter the ulcerated condition. At times, there is a simple destruction or erosion of the organ, at others, an *encephaloid* or *cerebriform*, and, at others, again, a *colloid* degeneration.

For its production, it requires a peculiar diathesis, or cachexia. The following table, from Dr. Walshe, exhibits the characters of the three species of carcinoma:

The use of irritants in cancerous affections is strongly to be deprecated. When the disease is so situate that excision can be practised, and the constitution does not exhibit great contamination, the sooner it is removed the better.

Encephaloid.	Scirrhus.	Colloid.
<p>Resembles lobulated cerebral matter.</p> <p>Is commonly opaque from its earliest formation.</p> <p>Is of a dead white colour.</p> <p>Contains a multitude of minute vessels.</p> <p>Is less hard and dense than scirrhus.</p>	<p>Resembles rind of bacon traversed by cellular-fibrous septa.</p> <p>Has a semi-transparent glossiness.</p> <p>Has a clear whitish or bluish yellow tint.</p> <p>Is comparatively ill-supplied with vessels.</p> <p>Is exceedingly firm and dense.</p>	<p>Has the appearance of particles of jelly imbed in a regien of alveolar matter.</p> <p>The contained matter is strikingly translucent.</p> <p>Greenish yellow is its predominant hue.</p> <p>Its vessels have not been sufficiently examined as yet.</p> <p>The jelly like matter is exceedingly soft; a roll of mass is, however, firm and resisting.</p> <p>The pulsatious variety has been detected in the vessels.</p> <p>Is composed of shills in a state of emboisement.</p>
<p>Is frequently found in the veins issuing from the diseased mass.</p> <p>The predominant microscopical elements are globular not always distinctly cellular, and caudate corpuscula.</p> <p>Occasionally attains an enormous bulk.</p> <p>Has been observed in almost every tissue of the body.</p> <p>Very commonly co-exists in several parts or organs of the same subject.</p>	<p>Has not been distinctly detected in this situation.</p> <p>The main microscopical constituents are juxtaposed nuclear cells; caudate corpuscula do not exist in it.</p> <p>Rarely acquires larger dimensions than an orange.</p> <p>Its seat as ascertained by observation, is somewhat more limited.</p> <p>Is not unusually solitary.</p>	<p>Observes a mean in this respect.</p> <p>Has so far been seen in a limited number of parts only.</p> <p>Has rarely been met with in more than one organ.</p>
<p>Is remarkable for its occasional vast rapidity of growth.</p> <p>Is frequently the seat of interstitial hemorrhage and deposition of black or bistre-coloured matter.</p> <p>When softened into a pulp, appears as a dead white or pink opaque matter of creamy consistence.</p> <p>Subcutaneous tumours are slow to contract adhesion with the skin.</p> <p>Ulcerated encephaloid is frequently the seat of hemorrhage, followed by rapid fungous development.</p>	<p>Ordinarily grows slowly.</p> <p>Is comparatively rarely the seat of these changes.</p> <p>Resembles, when softened, a yellowish brown semitransparent gelatinous matter.</p> <p>Scirrhus thus situate usually becomes adherent.</p> <p>Scirrhus ulcers much less frequently give rise to hemorrhage; and fungous growths (provided they retain the scirrhus character) are now more slowly and less abundantly developed.</p>	<p>Grows with a medium degree of rapidity.</p> <p>Undergoes no visible change of the kind.</p>
<p>The progress of the disease after ulceration is commonly very rapid.</p> <p>It is the most common form under which secondary cancer exhibits itself.</p> <p>Is the species of cancer most frequently observed in young subjects.</p>	<p>There is not such a remarkable change in the rate of progress of the disease after ulceration has set in.</p> <p>Is much less common before puberty.</p>	<p>Has so far been observed in adults only.</p>

CANCER ALVÉOLAIRE, Colloid—c. Alveolaris, Colloid—c. Apertus, see Cancer.

CANCER AQUATICUS, *Gangrenous stomatitis*, *Cancerum Oris*, *Gangrenop'sis*, *Canker of the mouth*, *Gangrenous sore mouth*, *Gangrenous erosion of the cheek*, *Sloughing Phagedæna of the mouth*, *Water Canker*: called, also, *Aphthæ serpentes*, *Gangræna Oris*, *Noma*, *Noma*, *Nomus*, *Pseudocarcinoma labii*, *Stomacacæ gangrenosa*, *Cheilocacæ*, *Uloacæ*, *Ullitis septica*, *Cheilomacæ*, *Scorbutus Oris*, *Stomatomacæna putrida*, *Stomatopsepsis*, *Stomatonecrosis*, *Carbunculus labiorum et genarum*, (F.) *Cancer aquaticus*, *Stomatitis gangrenosa*, *S. Charbonnosa*, *Gangrène ou Syphilis de la Bouche*, *Égérie*, *Aphthæ gangrèneuse*. Certain sloughing or gangrenous ulcers of the mouth—so called, perhaps, because they are often accompanied with an afflux of saliva. The disease is not uncommon in children's asylums, and demands the same treatment as hospital gangrene—the employment of caustics, and internal and external antiseptics.

CANCER AQUATICUS, *Cancer aquaticus*, *Stomacacæ*—c. *Alvéolaire*, Colloid—c. *Astutus*, see *Cancerum chelæ*—c. *Black*, *Cancer melanotic*—c. of the Breast, *Mastocarcinoma*—c. *Caminiariorum*, *Cancer chimney-sweepers'*—c. *Cellular*, *Encephaloid*—c. *Cerebriforme*, see *Encephaloid*.

CANCER, CHIMNEY-SWEEPERS', *Sootwart*, *Cancer munditum seu purgatorius infanticuli seu caminiariorum*, *Cancer seu carcinoma seroti*, *Oschocarcinoma*, *Oschocarcinoma*, (F.) *Cancer des Ramoneurs*. This affection begins with a superficial, painful, irregular ulcer, with hard and elevated edges, occupying the lower part of the

serotum. Extirpation of the diseased part is the only means of effecting a cure.

CANCER COLLOÏDE, Colloid.

CANCER, DAVIDSON'S REMEDY FOR, see *Conium maculatum*.

CANCER DUR, Scirrhus.

CANCER ÉBURNÉ. A kind of waxy degeneration of the breast, so called by M. Alibert, but which appears to be in no way allied to cancer.

CANCER ENCÉPHALOÏDE, Scirrhus—c. Epithelial, see Epithelial—c. *Fibrous*, Scirrhus—c. *Fibrous*, Scirrhus—c. *du Foie*, *Hépatocarcinome*.

CANCER GALIENI, (F.) *Cancer de Galien*. A bandage for the head, to which Galien gave the name *cancer*, from its eight heads resembling, rudely, the claws of the crab. It is now supplied by the bandage with six claws or heads, which is called the *Bandage of Galien* or *B. of the Poor*.

CANCER DE GALIEN, *Cancer Galeni*—c. *Gelatiniform*, Colloid—c. *Gelatinous*, Colloid—c. *Gommeux*, Colloid—c. *Hard*, Scirrhus—c. *Intestinorum*, *Enteropathia cancerosa*—c. *des Intestins*, *Enteropathia cancerosa*—c. of the Lung, *Platibus cancerous*—c. *Lupus*, *Lupus*—c. *Mastoid*, see *Mastoid cancer*—c. *Medullaris*, *Encephaloid*—c. *Melanous*, *Cancer melanotic*—c. *Mélan*, *Cancer melanotic*—c. *Melanodes*, *Cancer melanotic*.

CANCER, MELANOTIC, *Cancer melanoticus*, *Carcinoma melanoticum seu melanoticum seu melanous*, *Fungus melanoides*, *Melanocarcinoma*, *Black* or *Malanoid Cancer*, (F.) *Cancer Mélan*. A combination of cancer and melanosis.

CANCER, MILTLIKE, see *Miltlike*.

CANCER MOLLIS, see *Encephaloid*—*c. Mou*, *Encephaloid*—*c. Munditorum*, Cancer, chimney-sweepers'—*c. Nephroid*, see *Nephroid*—*c. Occultus*, see *Cancer*—*c. Oculi*, *Scirrhopthalmus*—*c. Oris*, Stomachæ—*c. Ossis*, *Spina veniosa*—*c. Pharyngis et œsophagi*, *Lænoscirrhus*—*c. Pul-tacæ*, Colloid—*c. Purgatoris infumiculi*, Cancer, chimney-sweepers'—*c. des Ramoneux*, Cancer, chimney-sweepers'—*c. Scirrhus*, *Scirrhus*—*c. Scroti*, Cancer, chimney-sweepers'—*c. Soft*, *Hæmatodes fungus*—*c. Solanoid*, see *Solanoid*—*c. Squirrhens*, *Scirrhus*—*c. of the Stomach*, *Gastrostentosis cardiaca* et *pylorica*—*c. Uteri*, *Metrocarcinoma*.

CANCÉREUX, Cancerous.

CANCER ROOT, *Orobanche Virginiana*, *Phytolacca decandra*—*c. Weed*, *Goodyera pubescens*, *Salvia lyrata*.

CANCEROMA, Carcinoma.

CAN'CEROUS, *Cancro'sus*, *Carcino'sus*, (F.)

Cancéreur. Relating to cancer; as *Cancerous ulcer*, *Cancerous diathesis*.

CANCHALAGUA, *Chironia Chilensis*.

CANCRENA, Gangrene.

CANCRODES, Cancroid.

CANCROID, *Kankroid*, *Cancro'des*, *Cancroï'des*, *Carcino'des*, *Carcinoides*, *Cancroïdous*, from *cancer* and *aidos*, 'form.' That which assumes a cancerous appearance. Cancroid is a name given to certain cutaneous cancers by Alibert: called also *Cheloid* or *Keloid*, *Chel'ois*, *Can'crois*, *Kel'oides*, (χελύς, 'a tortoise,' and αἶος, 'likeness,' or from κηλῖς, 'a spot or stain,' and αἶος, 'likeness,') *Kelis*, from their presenting a flattish raised patch of integument, resembling the shell of a tortoise.

CANCROID OF THE SKIN, see *Epithelial*.

CANCROIS, see *Cancroid*.

CANCROMA, Carcinoma.

CANCORUM CHELÆ, *Oc'uli seu Lap'idēs seu Lapill'i Cancor'um*, *Concremen'ta As'taci fluvial'itis*, *Crab's stones* or *eyes*, (F.) *Yeux d'écrevisse*, *Pierre d'Écrevisse*. Concretions found, particularly, in the *Cancer As'tacus* or *Cray-fish*. They consist of carbonate and phosphate of lime, and possess antacid virtues, but not more than chalk.

CANCROSUS, Cancerous, *Chancreuse*.

CANCROUM ORIS, *Cancer Aquaticus*, *Stomacæ*.

CANDELA, *Bougie*—*c. Fumalis*, *Pastil*—*c. Medicata*, *Bougie*—*c. Regia*, *Verbascum nigrum*.

CANDELARIA, *Verbascum nigrum*.

CANDI, *Candum*, *Canthum*, *Can'tion*, *Candy*, from *candidus*—'white, bleached, purified.' Purified and crystallized sugar. See *Saccharum*.

CANDIDUM OVI, *Albumen ovi*.

CANDLE, *Pupil*.

CANDLE, MERC'URAL. A candle made of *wax* and *vermilion* for mercurial fumigation.

CANDLE TREE, *Parmentiera cerifera*.

CANDY, *Candi*.

CANDYTUFT, *BITTER*, *Iberis amara*.

CANE, SUGAR, see *Saccharum*—*c. Sweet*, *Acorus calamus*.

CANEFICIER, *Cassia fistula*.

CANELA, *Nectandra cinnamomoides*.

CANELÉ, Grooved.

CANELLA, see *Canella alba*.

CANELLA ALBA, diminutive of *canna*, 'a reed.' A West India tree. *Fam. Magnoliacæ*: *Sez. Syst.* *Dodecandria Monogynia*. It is so called because its bark is rolled up like a reed. *Cortex Winteranus epu'rius*, *Canella Cuba'na seu Winter'ania*, *Cinnamomum album*, *Cortex Antiscorbuticus seu Aromat'icus*, *Costus cortico'sus*, *Cam'inga*, *Canella* (Ph. U. S.), *Canella Bark*, (F.) *Canelle* ou *Canelle blanche*, *Fausse Écorce de*

Winter, *Ecorce Carriocostine*; *Ord.* *Guttifereæ*, is a pungent aromatic. Its virtues are partly extracted by water; entirely by alcohol. It is a stimulant, and is added to bitters and cathartics.

CANELLA CARYOPHYLLATA, *Myrtus caryophyllata*—*c. Cubana*, *C. alba*, *Laurus cassia*—*c. Malabarica* et *Javensis*, *Laurus cassia*.

CANELLIFERA MALABARICA, *Laurus cassia*.

CANEPIN, (F.) A fine lamb's skin or goat's skin, used for trying the quality of lancets.

CANICACEOUS, *Furfuraceus*.

CAN'ICÆ, Meal, in which there is much bran. Also, coarse bread; or bread in which there is much bran—*Panis Canica'ceus*.

CANICIDA, *Aconitum*.

CANIC'ULA; the *Dogstar*, from *canis*, 'a dog,' *Erupio*, *Sirius*, (F.) *Canicule*. This star, which gives its name to the *Dogdays*, *Dies canicula'res*, because they commence when the sun rises with it, was formerly believed to exert a powerful influence on the animal economy. The *Dogdays* occur at a period of the year when there is generally great and oppressive heat, and therefore—it has been conceived—a greater liability to disease.

CANIF, Knife.

CANIN, Canine.

CANINANÆ RADIX, *Caincæ radix*.

CANINE, *Caninus*, *Cyn'ius*, from *canis*, 'a dog,' (F.) *Canin*. That which has a resemblance to the structure, &c., of a dog.

CANINE FOSSA, *Fossa Cani'na*, *Infra-orbital* or *Suborbital fossa*, (F.) *Fosse Canine*. A small depression on the superior maxillary bone, above the *dens caninus*, which gives attachment to the *caninus* or *levator anguli oris* muscle.

CANINE LAUGH, *Sardon'ic laugh*, *Risus Cani'nis seu Sardon'icus seu Sardo'nus seu Sardon'ios seu de Sardo'nia seu involuntari'us seu spas'ticus*, *Tortu'ra seu Distor'sio Oris*, *Gelas'mus*, *Sardi'asis*, *Sardon'asis*, *Trismus Sardon'icus seu cyn'icus*, *Spasmus musculorum faciei seu cyn'icus*, *Prosopospas'mus*, (F.) *Ris canin*, *R. Sardonique*, *R. Sardonien*, *R. moqueur*. A sort of laugh, the facial expression of which is produced particularly by the spasmodic contraction of the *Caninus* muscle. Probably, this expression, as well as *Cynic Spasm*, *Spasmus canin'us seu cyn'icus*, *Convul'sio cani'na*, *Trismus cyn'icus*, may have originated in the resemblance of the affection to certain movements in the upper lip of the dog. The *Risus Sardon'icus* is said to have been so called from similar symptoms having been induced by a kind of *Ranunculus*—*R. sceleratus*—that grows in Sardinia.

CANINE TEETH, *Dentes Cani'ni seu Lania'rii seu angula'res seu cuspidati seu columella'res seu ocula'res seu morden'tes*, *Cynodon'tes*, *Eye Teeth*, *Pug Teeth*, (F.) *Dents canines* ou *laniaires* ou *angulaires* ou *oculaires* ou *aillères* ou *convôides*. The teeth between the lateral incisors and small molars of each jaw;—so named because they resemble the teeth of the dog.

CANINUS, *Levator anguli oris*—*c. Sentis*, *Rosa canina*—*c. Spasmus*, see *Canine Laugh*.

CANIRAM, *Strychnos nux vomica*.

CANIRAMINUM, *Brucine*.

CANIRUBUS, *Rosa canina*.

CANIS INTERFECTOR, *Veratrum sabadilla*—*c. Ponticus*, *Castor fiber*.

CANIT'IES, from *canus*, 'white.' *Blanching*, *Whiteness* or *grayness* of the hair, and especially of that of the head—*Trichon'osis polio'sis*. (F.) *Canitie*. When occurring in consequence of old age, it is not a disease. Sometimes it happens suddenly, and apparently in consequence of severe mental emotion. The causes, however, are not clear. See *Poliosis*.

CANKER, Stomacae—c. of the Mouth, *Cancer aquaticus*.

CANKER-RASH, see *Cynanche maligna*.

CANKER, WATER, *Cancer aquaticus*.

CANNA, see *Tous-les-Mois*, *Cassia fistula*, *Trachea*—c. *Brachii*, *Ulna*—c. *Domestica eruris*, *Tibia*—c. *Edulis*, *Tous-les-Mois*—c. *Fistula*, *Cassia fistula*—c. *Ladica*, *Sagittarium alexipharmacum*—c. *Major*, *Tibia*—c. *Minor*, *Fibula*, *Radius*—c. *Solutiva*, *Cassia fistula*—c. *Starch*, *Tous-les-Mois*.

CANNABIN, *Bangue*.

CANNAB'INA, from *kavvaßis*, 'hemp.' Remedies composed of *Cannabis Indica*.—*Pereira*.

CANNABINA AQUATICA, *Eupatorium cannabinum*.

CANNABIS INDICA, *Bangue*. See, also, *Churrus*.

CAN'NABIS SATI'VA, (F.) *Chanvre*, *Chambrie*. The seed of this—*Hempseed*, *Seed'ina Can'nabis*, (F.) *Chênevis*—is oily and semiluminous. The decoction is sometimes used in gonorrhœa.

CANNACORUS RADICE CROCEA, *Curcuma longa*.

CANNAMELLE, see *Saccharum*.

CANNE AROMATIQUE, *Acorus calamus*—c. *Congo*, *Costus*—c. de *Rivière*, *Costus*—c. à *Sucre*, see *Saccharum*.

CANNEBERGE, *Vaccinium oxycoccos*—c. *Poneticæ*, *Vaccinium vitis idæa*.

CANNEL, *Laurus cinnamomum*.

CANNEL-BONE, *Clavicle*.

CANNEL-WATER, *Aqua cinnamomi*.

CANNELÉ ou CANELÉ (F.), from *canalis*, 'a canal.' *Sulcat'us*, *Striat'us*, *Canaliculat'us*. Having a canal or groove—as *Muscle cannelé* (Lieutenant), the Gemini; *Corps cannelés ou striés*, the *Corpora striata*; *Sonde cannelée*, a grooved sound, &c. See *Grooved*.

CANNEL BAYNE, *Clavicle*.

CANNELLE, *Laurus cinnamomum*—c. *Blanche*, *Canella alba*—c. de la *Chine*, *Laurus cassia*—c. de *Coromandel*, *Laurus cassia*—c. *Fausse*, *Laurus cassia*—c. *Giroflée*, *Myrtus caryophyllata*—c. des *Indes*, *Laurus cassia*—c. de *Java*, *Laurus cassia*—c. de *Malabar*, *Laurus cassia*—c. *Matte*, *Laurus cassia*—c. *Officinale*, *Laurus cinnamomum*—c. *Poirée*, see *Winters aromaticæ*.

CANNULA, *Canula*.

CANNULÉ PULMONUM, *Bronchia*.

CANOPUM, see *Sambucus*.

CANOR STETHOSCOPICUS, *Tintement métallique*.

CANQUOIN'S CAUSTIC PASTE, *Pasta zinci chloridi*.

CANTABRICA, *Convolvulus Cantabrica*.

CANTABRUNO, *Furfur*.

CANTARELLUS, *Meloe proscarabæus*.

CANTATIO, *Charm*.

CANTEL, *Vertex*.

CANTERBURY, WATERS OF, *Aquæ Cantuarien'ses*. The waters of *Canterbury*, in *Kent*, *England*, are impregnated with *iron*, *sulphur*, and *carbonic acid*.

CANTERIUM, *Cantherius*.

CANTHARIDE TACHETÉE, *Lytta vittata*.

CANTHARIDINE, see *Cantharis*.

CANTHARIS, from *kavðapos*, 'a scarabæus.' *O. vesicato'ria*, *Musca Hispanica*, *Mel'oe vesicato'ria*, *Lytta vesicato'ria*, *Blistering Fly*, *Blister-fly*, *Blisterbeetle*, *Spanish Fly*, *Fly*, (F.) *Cantharides*, *Mouches*, *M. d'Espagne*. This fly—Order, *Coleoptera*—originally, perhaps, a native of *Italy* and *Spain*, is now found in *France*, *Germany*, *Hungary*, *Russia*, *Siberia*, and *England*. It is, however, rare in the last-named country. It is found in species of *Oleaceæ*—as the *ash*, *privet*, and *lilac*; and of *Caprifoliaceæ*—as the

elder and *lonicera*. It is much employed in *medicine*, and is the most common *vesicatory*. Given internally, and even when absorbed from the skin, it affects the urinary organs, exciting *strangury*. This may be prevented, in cases of blisters, by interposing between the blistering plaster and skin a piece of tissue-paper. Diluents relieve the *strangury*. Dose, half a grain to one grain. If kept dry, the flies will retain their activity for many years. Their active principle—*Cantharidin*, *Cantharidin'na*—has been separated from them.

Cantharis vesicatoria is the ordinary blistering insect used in *Europe*.

Cantharis seu *Lytta vittata*, (which see,) and *C. atrata*, *C. marginata*, and *C. cinerea*, of *America*; *C. atomaria*, of *Brazil*; *C. rubripes*, of *Sumatra* and *Java*; *C. gigas*, *Lytta carulea*, of *Guinea* and the *East Indies*; *C. violacea*, *Lytta gigas mus*, of the *East Indies*; *C. Syriaca*, *Lytta sog'etum*, of *Arabia*; *Myla'bris*, *M. punctata*, *M. pustulata*, and *M. cichorii*, of *China* and the *East Indies*; *Mel'oe proscarabæus*, and *M. majalis* or *True Mayworm*—possess similar properties.

CANTHE'RIUS, *Cante'rium*. The cross-piece of wood in the apparatus used by *Hippocrates* for reducing luxations of the humerus.

CANTH'ITIS. Inflammation of the canthus of the eye.

CANTHIUM CORONATUM, *Randia dume-torum*.

CANTHOPLAS'TICÉ, *Can'thoplasty*, (F.) *Canthoplastic*, from *kanthos*, 'the angle of the eye,' and *πλαστικός*, 'formative.' The formation, by plastic operation, of the angle of the eye. An operation proposed by *Ammon*, when the eyelids are not sufficiently cleft.

CANTHIUM, *Candi*.

CANTHUS, *Epican'this*, *Angulus ocularis*, *Fons lacrymarum*. The corner or angle of the eye. The greater canthus is the inner angle, *Hircus*, *Hir'gus*, *Rhanter*; the lesser canthus, the outer angle, *Parop'ia*, *Pega*.

CANTIANUS PULVIS, 'Kent Powder.' A cordial powder, known under the name 'Countess of Kent's Powder,' composed of coral, amber, crabs' eyes, prepared pearls, &c. It was given in cancer.

CANTION, *Candi*.

CAN'ULA, *Can'nula*, *Au'tiscus*, *Aulos*. Diminutive of *Canna*, 'a reed.' *Tu'tulus*, (F.) *Canule* ou *Canule*. A small tube of gold, silver, platinum, iron, lead, wood, elastic gum, or gutta percha, used for various purposes in surgery.

CA'OUTCHOUC. The Indian name for *Indian Rubber*, *Elastic Gum*, *Gum Elastic*, *Gomme elastique*, *Cauchou*, *Res'ina elastica* seu *Cayenne'sis*, *Cayenne Resin*, *Cautchuc*, (F.) *Gomme elastique*, *Résine elastique* ou de *Cayenne*. A substance formed from the milky juice of *Hevea seu Hevea Guianen'sis*, *Jatropha elastica*, *Siphonia Cahucho*, *S. elastica*, *Seringue Tree*, and of *Ficus Indica*, and *Artocarpus integrifolia*—South American trees. It is insoluble in water and alcohol; but boiling water softens and swells it. It is soluble in the essential oils and in ether, when it may be blown into bladders. It is used in the fabrication of catheters, bougies, pessaries, &c.

CAP, PITCH, see *Depilatory*.

CAPACITY, LUNG, see *Vital Capacity*—c. *Pulmonic*, see *Vital Capacity*.

CAPA-ISIAKKA, *Bromelia ananas*.

CAPBERN, WATERS OF. *Capbern* is in the department *Hautes-Pyrénées*, *France*. The waters contain sulphates and carbonates of lime and magnesia, and chloride of magnesium. Temperature, 75° Fahrenheit. They are purgative.

CAPELET, *Myrtus carophyllata*.

CAPEL'YNA, *Capelli'na*, (F.) *Capeline*. A Woman's Hat, in French; *Capis'trum*, from *caput*, 'head.' A sort of bandage, which, in shape, resembles a riding-hood. There are several kinds of *Capelines*:—1. That of the head, *C. de la tête*, *Fascia capiti'lis*. See *Bonnet d'Hippocrate*. *C. of the clavicle*, employed in fractures of the acromion, clavicle, and spine of the scapula. *C. of an amputated limb*—the bandage applied round the stump.

CAPELLINA, *Capelina*.

CAPER BUSH, *Capparis spinosa*.

CAPER PLANT, *Euphorbia lathyris*.

CAPERS, see *Capparis spinosa*.

CAPETUS, Imperforation.

CAPHORA, *Camphor*.

CAPHURA, *Camphor*.

CAPILLAIRE, *Capillary*, see *Adiantum capillus veneris* — *c. du Canada*, *Adiantum pedatum* — *c. de Montpellier*, *Adiantum capillus veneris*.

CAPILLAMEN'TUM, from *capillus*, 'a hair,' *Capilli'tium*, *Tricho'ma*, *Trichoma'tion*. Any villous or hairy covering. Also, a small fibre or fibril.

CAPILLARY, *Capilla'ris*, *Capilla'ceus*, from *capillus*, 'a hair,' (F.) *Capillaire*. Hair-like; small.

CAPILLARY VESSELS, *Vasa capilla'ria*, *Micrang'i'a*, *Trichang'i'a*, (F.) *Vaisseaux capillaires*, *Capillicules*, are the extreme radicles of the arteries and veins, which together constitute the capillary, intermediate, or peripheral vascular system — the *methamata* or *methamatus* blood-channels of Dr. Marshall Hall. They possess an action distinct from that of the heart, but not one of rhythmic contraction and dilatation. Formerly, it was conceived that white vessels — *Vasa sero'sa* — or vessels so minute as not to admit red corpuscles, were distributed to the tissues which do not receive red blood.

CAPILLATIO, *Trichismus*.

CAPILLATUS, *Impuber*.

CAPILLICULES, *Capillary vesse's*.

CAPILLITIUM, *Capillamentum*, *Entropion*, *Scalp*.

CAPILLORUM DEFLUVIUM, *Alopecia*.

CAPIL'PILUS, quasi *Capitis Pilus*, *Coma*, *Chaté*, *Crinis*, *Pilus*, *Thrix*, *Cana'ries*, (F.) *Cheveu*. This term is generally applied to the hair of the head, *Pili seu Honor cap'itis*, the characters of which vary according to races, individuals, &c. Hairs arise in the areolar membrane, where the bulb is placed, and are composed of two parts—one, external, transparent, and of an epidermoid character; the other, internal and *sui generis*, which, in the opinion of some, communicates to them their colour. The hair is insensible, and grows from the root.

CAPILLUS VENERIS, *Adiantum capillus veneris* — *c. v. Canadensis*, *Adiantum pedatum*.

CAPILE'NIUM, *Capiti'plenum*, from *caput*, 'the head,' and *plenum*, 'full.' A word employed with different significations. A variety of catarrh. — *Schneider*. A heaviness or disorder in the head common at Rome, like the *καρβαρία*, *Carebaria*, of the Greeks. — *Baglivi*.

CAPISTRATIO, *Phimosis*.

CAPISTRUM, *Capeline*, *Chevestre*, *Trismus* — *c. Auri*, *Borax*.

CAPIS'TRUM, *Phimos*, *Cemos*, *κνυος*, 'a halter.' This name has been given to several bandages for the head. See *Capeline*, *Chevestre*.

CAPITALIA REMEDIA, *Cephalic remedies*.

CAPITALIS, *Cephalic*.

CAPITELLUM, *Alembic*, see *Caput*.

CAPITEUX, *Heady*.

CAPITILU'VIUM, from *caput*, 'the head,' and *lavare*, 'to wash.' A bath for the head.

CAPITIPLENIUM, *Capiplenium*.

CAPITIPURGIA, *Caput purgia*.

CAPITIT'RAHA, from *caput*, 'the head,' and *trahere*, 'to draw.' Instruments which, like the forceps, draw down the head of the foetus when impacted in the pelvis.

CAPITO'NES, from *caput*, 'the head.' *Macroceph'ali*, *Proceph'ali*. Foetuses whose heads are so large as to render labour difficult. The condition is called *Macroceph'alia*.

CAPITULUM, *Alembic*, *Condyle*, see *Caput* — *c. Costæ*, see *Costa* — *c. Laryngis*, *Corniculum laryngis* — *c. Martis*, *Eryngium campestre* — *c. Santorini*, *Corniculum laryngis*.

CAPITULUVIUM, *Bath (head)*.

CAPNISMOS, *Fumigation*.

CAPNITIS, *Tutia*.

CAPNOIDES CAVA, *Fumaria bulbosa*.

CAPNORCHIS, *Fumaria bulbosa*.

CAPNOS, *Fumaria*.

CAPON, *Cagot*.

CAPON SPRINGS. A pleasant summer retreat, situated in a gorge of the North Mountain, in Hampshire co., Va., 23 miles W. of Winchester. The waters in the vicinity are sulphurous and chalybeate; — those at the springs alkaline and diuretic.

CAPOT, *Cagot*.

CAP'PARIS SPINO'SA, *Cap'paris*, *Cappar*, *Ca'pria*, *Prickly Caper Bush*, (F.) *Câprier*. Ord. *Capparideæ*. Ser. *Nyct*. *Polyandria Monogynia*. The bark of the root and the buds have been esteemed astringent and diuretic. The buds are a well-known pickle — *Capers*, (F.) *Câpres*.

CAPPARIS BADUC'CA, *Baduk'ka*. A species of caper, cultivated in India on account of the beauty of its flowers. The Orientals make a liniment with its juice, with which they rub pained parts. The flowers are purgative.

CAPPONE, WATERS OF. At Cappone, in the isle of Ischia, are waters containing carbonate of soda, chloride of sodium, and carbonate of lime. Temp. 100° Fahr.

CAPREOLA'NIS, from *capreolus*, 'a tendril,' *Cissoi'des*, *Eliticoi'des*, (F.) *Caprilaire*. Twisted.

CAPREOLA'RIA VASA. Some have called thus the spermatie arteries and veins, on account of their numerous contortions.

CAPREOLUS, *Helix*.

CÁPRES, see *Capparis spinosa*.

CAPRIA, *Capparis spinosa*.

CAPRICORNUS, *Plumbum*.

CÁPRIER, *Capparis spinosa*.

CAPRIFOLIA, *Lonicera periclymenum*.

CAPRIFOLIUM DISTINCTUM, *Lonicera periclymenum* — *c. Periclymenum*, *Lonicera periclymenum* — *c. Sempervivens*, *Lonicera sempervivens*, — *c. Sylvaticum*, *Lonicera periclymenum*.

CAPRILUQUIUM, *Egophony*.

CAPRIZANS PULSUS, see *Pulse*, *caprizant*.

CAPSA, *Boîte*, *Capsule*, *Case* — *c. Cordis*, *Pericardium*.

CAPSARIUM, *Boítier*.

CAPELLA BURSA PASTORIS, *Thlaspi bursa*.

CAPSICUM, see *Capsicum annum*.

CAP'SICUM ANNUUM, *C. Hispan'icum*, from *καττω*, 'I bite.' The systematic name of the plant whence, in part, *Cayenne Pepper* is obtained, — *Piper In'dicum* seu *Hispan'icum* seu *Brazilia'num* seu *Guineen'sè* seu *Calec'iticum* seu *Tur'cicum* seu *Lusitan'icum*, *Sola'mum urens*, *Siliquis'trum Plin'i'i*, *Cayenne* or *Guinea* or *Chilli Pepper*, *Garden Ginger*, (F.) *Piment*, *Poivre d'Inde*, *Poivre de Guinée*, *Corail des Jardins*. Order, *Solanææ*. The pungent, aromatic properties of *Bacca Cap-*

ici, *Capsicum Berries*, *Capsicum* (Ph. U. S.), are yielded to ether, alcohol, and water. They are highly stimulant and rubefacient, and are used as a condiment. Their active principle is called *Capsicin*.

CAPSICUM HISPANICUM, *Capsicum annum*.

CAPSIQUE, *Capsicum annum*.

CAPSITIS, see *Phacitis*.

CAPSULA, *Boitier*—c. *Articularis*, *Capsular* ligament—c. *Cordis*, *Pericardium*—c. *Dentis*, *Dental follicle*—c. *Humoris aquei*, see *Aqueous humour of the eye*—c. *Lentis*, see *Crystalline*—c. *Nervorum*, *Neurilemma*—c. *Sequestralis*, see *Sequestrum*.

CAPSULE SEMINALES, *Vesiculæ S.*—c. *Synoviales*, *Bursæ mucosæ*.

CAPSULAIRE, *Capsular*.

CAPSULAR, *Capsula'ris*, (F.) *Capsulaire*. Relating to a capsula or capsule.

CAPSULAR ARTERIES, *Suprarenal Arteries and Veins*. Vessels belonging to the suprarenal capsules. They are divided into superior, middle, and inferior. The first proceed from the inferior phrenic, the second from the aorta, and the third from the renal artery. The corresponding veins enter the phrenic, vena cava, and renal.

CAPSULAR CATARACT, see *Cataract*.

CAPSULAR LIGAMENT, *Ligamentum capsula're*, *Capsula articul'ris*, *Articular or Fibrous capsule*, (F.) *Ligament capsulaire*, *Capsule articulaire*, *Capsule fibreux*, &c. Membranous, fibrous, and elastic bags or capsules, of a whitish consistence, thick, and resisting, which surround joints.

CAPSULE, *Capsula*, *Capsa*, a box, or case, (F.) *Capsule*. This name has been given, by anatomists, to parts bearing no analogy to each other.

CAPSULE, *BOWMAN'S*, see *Kidney*—c. *Cellular*, of the Eye, see *Eye*—c. *Fibrous*, *Capsular ligament*.

CAPSULE, *GELATINOUS*, *Capsula gelatinæ*, (F.) *Capsule gélatineuse*, *Capsule of gelatin*. A modern invention by which capsaiba and other disagreeable oils can be enveloped in gelatin so as to conceal their taste.

CAPSULE OF GLISSON, *Capsula GLISSONII* seu *communis GLISSONII*, *Vag'ina Porta*, V. *GLISSONII*. A sort of membrane, described by Glisson, which is nothing more than dense areolar membrane surrounding the vena porta and its ramifications in the liver.

CAPSULE OF THE HEART, *Pericardium*—c. *Ocular*, see *Eye*.

CAPSULE, *RENAL*, *Suprarenal* or *Atrabiliary C.*, *Renal Gland*, *Glandula suprarenalis*, *Capsula rena'lis* seu *suprarenalis* seu *atrabili'ris*, *Ren succenturiatus*, *Nephridium*, (F.) *Capsule surrénale* ou *atrabilaire*. A flat triangular body, which covers the upper part of the kidney, as with a helmet. A hollow cavity in the interior contains a brown, reddish or yellowish fluid. The renal capsules were long supposed to be the secretory organs of the fancied atrabiles. They are much larger in the fetus than in the adult. They are probably concerned in lymphosis.

A singular condition of cachexia, the leading characteristics of which are anemia, general languor and debility; remarkable feebleness of the heart's action; irritability of the stomach, and a peculiar bronzed skin, was first described by Dr. Thos. Addison, of London, as connected with a diseased condition of the supra-renal capsules. It has been proposed to call it the *Disease of Addison*, (F.) *Maladie d'Addison*.

CAPSULE, *SEMINAL*, *Capsula semina'lis*. *BARTOLINE* thus designates the extremity of the vas deferens, which is sensibly dilated in the vicinity of the vesiculæ seminales. Some anatomists apply this name to the vesiculæ themselves.

CAPSULE, *SYNOVIAL*, *Capsula Synovia'lis*, A

membranous bag, surrounding the movable articulations and canals, which gives passage to tendons. Synovial capsules exhale, from their articular surface, a fluid, whose function is to favour the motions of parts upon each other. See *Bursa mucosa*, and *Synovia*.

CAPSULE SURRÉNALE ou *ATRABILIAIRE*, *Capsule*, renal.

CAPSULITIS, see *Phacitis*.

CAPUCHON, *Trapezius*.

CAPUCINE, *Tropæolum majus*.

CAPULIES, *Prunus capulin*.

CAPULUS, *Scrotum*.

CAPUT, 'the head.' Also, the top of a bone or other part, (F.) *Tête*. The head of small bones is sometimes termed *capitulum*, *capitellum*, *cephalidium*, *ceph'alis*, *cephalum*. Also, the glans penis.

CAPUT ASPERÆ ARTERIÆ, *Larynx*—c. *Coli*, *Cæcum*—c. *Gallinaceum*, see *Gallinaginis caput*—c. *Gallinaginis*, see *Gallinaginis caput*—c. *Genitale*, *Glans*—c. *Lubricum*, *Penis*—c. *Major*, see *Epididymis*—c. *Minor*, see *Epididymis*—c. *Machi*, *Leontodon Taraxacum*—c. *Obstipum*, *Torticollis*—c. *Penis*, *Glans*.

CAPUT PUR'GIA, *Capitipur'gia*. Remedies, which the ancients regarded as proper for purging the head:—*errhines*, *sternutatories*, *apophlegmatisantia*, &c. Prosper Alpinus makes the *caput purgia* to be the same as *errhines*; and the *apophlegmatisantia* the same as the masticatories of the moderns.

CAPUT SCAPULÆ, *Acromion*.

CAPUT SUCCEANÆUM. A term sometimes used for the tumefied scalp, which first presents in certain cases of labour.

CAPUT TESTIS, *Epididymis*.

CAQUE-SANGUE, *Cague-sangue*. Old French words which signify *bloody evacuations*, (F.) *Dijections sanguinolentes*. They come from *cacare*, 'to go to stool,' and *sanguis*, 'blood.' Under this term was comprehended every affection in which blood is discharged from the bowels.

CARA SCHU'LLI, *Frutex Ind'icæ spinosus*, *Barleria burxifolia*. A Malabar plant, which, when applied externally, is maturative and resolvent. The decoction of its root is used, in the country, in ischuria.

CARABAC'CIUM. An aromatic wood of India, of a yellowish colour, and a smell like that of the clove. Its decoction and infusion are given as stomachics and antiscorbutics.

CARABUS. A genus of coleopterous insects. Two species, the *chrysocephalus* and *ferrugineus* have been recommended for the toothach. They must be pressed between the fingers, and then rubbed on the gum and tooth affected.

CARACTERE, *Character*, *Symbol*.

CARAGNA, *Caranna*.

CARAMATA, *Amumti*. A tree in the inland parts of Pomeronia. It furnishes a febrifuge bark, which Dr. Hancock says may be used in typhoid and remittent fevers where cinchona is either useless or pernicious.

CARAMBOLO, *Averrhoa carambola*.

CAREMEL, *Saccharum percoctum* seu *tosum*. Sugar subjected to the action of heat, until it is partly decomposed, deliquescent, of a brown colour, and a strong, agreeable and empyreumatic odour.

CARAN'NA, *Caragna*, *Tacamaha'ca*, *Caragna*, *Caran'na Gummi*, *G. Bre'vis*, *Gum Caran'na*, (F.) *Caragne*, *Gomme Caragne* ou *Carane*. A gum-resinous substance, which flows from a large tree in New Spain, and is obtained from South America in impure masses. It preserves its softness for a long time, has an aromatic smell, and a slightly acid and bitter taste. It was formerly used as a vulnerary and in plasters.

CARAWAY, Carum.

CARAWAY SEEDS, see Carum.

CARBASA, Linteum.

CARBASUS, Linteum.

CARBO, Carbon, (F.) *Carbone*. An elementary body, extensively distributed in nature; but of which the two following forms are official in the Pharmacopœia of the United States. Also, a coal, charcoal.

CARBO LIGNI, Charcoal (F.) *Charbon*. Fresh Charcoal is antiseptic. It is used to improve the digestive organs in cases of worms, dyspepsia, &c.; as a cataplasm to gangrenous and fetid ulcers, tinea, &c., and forms a good tooth-powder. Dose, gr. x to ʒi. Also, Anthrax.

CARBO ANIMALIS, *C. carnis*, Animal charcoal, (F.) *Charbon animal*, usually prepared by subjecting bones to a red heat in close vessels. The result is *Bone black*, commonly called *Ivory black*. (F.) *Noir animal ou d'os*. It is given in the same cases as *Carbo Ligni*, and has been extolled in cancer. Dose, gr. ss to gr. iij.

The Pharmacopœia of the United States contains a formula for the preparation of CARBO ANIMALIS PURIFICATUS, *Purified animal charcoal* (Carbon. animal. ʒij; Acid. muriat., Aqua ʒā f ʒxij). Pour the muriatic acid, previously mixed with the water, gradually upon the charcoal, and digest with a gentle heat for two days, occasionally stirring the mixture. Having allowed the undissolved portion to subside, pour off the supernatant liquor, wash the charcoal frequently with water, until it is entirely free from acid, and lastly dry it.

CARBO FOS'ILIS, *Lithanthrax*, Stone coal.

CARBO HUMA'NUM. The human excrement.—*Paracæsus*.

CARBO LIGNI, Carbo—c. Mineralis, Graphites—c. Palpebrarum, Anthracosis—c. Spongiae, Spongia usta.

CARBON, Carbo—c. Sesqui-iodide of, Carbonis sesqui-iodidum—c. Bisulphuret of, Carbonis sulphuretum—c. Sesquichloride of, Carbonis trichloridum—c. Sulphide of, Carbonis sulphuretum—c. Sulphuret of, Carbonis sulphuretum—c. Terechloride of, Carbonis trichloridum, see Chloroform.

CARBONAS or CARBO'NAS. A carbonate. (F.) *Carbonate*. A salt, formed by the combination of carbonic acid with a salifiable base.

CARBONAS NATRICUM, *Sodæ carbonas*—c. Plumbeus, *Plumbi subcarbonas*.

CARBONATE D'AMMONIAQUE, *Ammonie carbonas*.

CARBONATED, *Carbonatus*, *Aëriatus*, (F.) *Carboné*, *Aéré*. That which is impregnated with carbonic acid.

CARBONE, Carbon—c. *Trichlorure de*, Carbonis trichloridum.

CARBONÉ, Carbonated.

CARBONEUM CHLORATUM, Chloroform.

CARBONIC ACID, *Ac'idium Carbonicum*, *Solid Air of Hales*, *Fæctitious or Fixed Air*, *Carbonaceous or Calcareous or Aërial or Mephitic Air*, *Mephitic Air*, *Spir'itus lethalis*, (F.) *Acide Carbonique*. This gas, which neither supports respiration nor combustion, is not often used in medicine. It is the main agent in effervescent draughts, fermenting poultices, &c. It is often found occupying the lower parts of mines—when it is called the *choke damp*—caverns, tombs, wells, brewers' vats, &c., and not unfrequently has been the cause of death. Lime thrown into such places soon absorbs the acid.

CARBONIS SESQUICHLORIDUM, Carbonis trichloridum.

CARBONIS SESQUI-IODIDUM, *C. Sesqui-ioduretum*, *Sesqui-Iodide or Sesqui-Ioduret of Carbon*. This is made by mixing concentrated alcoholic solutions of iodine and potassa, until

the former loses its colour; a solution is obtained from which water throws down a yellow precipitate—the sesqui-iodide of carbon. It has been used in enlarged glands and in some cutaneous affections, applied externally, (ʒss to ʒvj of cerate). See Iodoform.

CARBONIS SULPHURETUM, *Sulph'uris Carbyretum*, *Sulphuretum seu Sul'idum Carbo'ni*, *Carbo'nium Sulphuratum*, *Al'cohol Sul'phuris*, *Bisulphuretum Carbo'ni*, *Sulphuret*, *Sulphide* or *Bisulphuret of Carbon*, *Carburet of Sulphur*, (F.) *Sulfure de Carbone*, *Carbure de Soufre*, *Soufre Carburé*, *Alcool de Soufre*. This transparent, colourless fluid, which has a very penetrating, disagreeable odour, and a taste which is cooling at first, but afterwards acid and somewhat aromatic, is a diffusible excitant. It is diaphoretic, diuretic, and has been said to have proved emmenagogue. It is also used in nervous diseases as an antispasmodic. Dose, one drop to four, repeated frequently.

It is used externally, where a cooling influence has to be rapidly exerted, and has been inhaled as an anæsthetic.

CARBONIS TRICHLORIDUM, *C. Sesqui-chloridum*, *Terchloride or Sesquichloride of Carbon*, (F.) *Trichlorure ou Sesquichlorure de Carbone*, is formed by the action of Chlorine on Chloroform, under the influence of sun-light. It has been given in cholera, and applied to correct the fetor of foul ulcers. Dose, four grains or more.

CARBONIUM SULPHURATUM, Carbonis sulphuretum.

CARBUNCLE, Anthrax.

CARBUNCLE, FUNGUS, Terminusus.

CARBUNCLE OF THE TONGUE, Glossanthrax—c. Berry, Terminusus.

CARBUNCLED FACE, Gutta rosea.

CARBUNCULAR EXANTHEM, Anthracia.

CARBUNCULATIO OCULI, Blepharanthracosis.

CARBUNCULUS, Anthrax—c. Anginosus, Cynanche maligna—c. Contagiosus, see Anthrax—c. Gallicus, see Anthrax—c. Hungaricus, see Anthrax—c. Labiorum et genarum, Cancer aquaticus—c. Polonicus, see Anthrax—c. Pulmonum, Necropneumonia—c. Septentrionalis, see Anthrax.

CARBUNCULUS RUBINUS. A red, shining, and transparent stone, from the Isle of Ceylon; formerly employed in medicine as a preservative against several poisons, the plague, &c.

CARBUNCULUS ULCUSCULOSUS, Cynanche maligna.

CARBURE DE SOUFRE, Carbonis sulphuretum.

CAR'CAROS, from *καρκαρω*, 'I resound,' 'I tremble.' A fever, in which the patient has a general tremor, accompanied with an unceasing noise in the ears.

CARCINODES, Canceroid, *Chancreuse*.

CARCINOIDES, Canceroid.

CARCINO'MA, *Karkino'ma*, *Cancero'ma*, *Cancro'ma*, from *καρκινος*, 'a crab.' Some authors have thus called indolent tumours different from cancer; others, incipient cancer; and others, again, the species of cancer in which the affected structure assumes the appearance of cerebral substance; but the majority of authors use Carcinoma in the same sense as Cancer.

CARCINOMA ALVEOLARE, Colloid—c. Epithelioides, Epithelial Cancer—c. Fibrosus, Scirrhus—c. Hæmatodes, Hæmatodes fungus—c. Intestinorum, Enteropathia cancerosa—c. Lingue, Glosso-carcinoma—c. of the Liver, Hepatoscirrhos—c. Medullare, Encephaloid—c. Melanodes, Cancer, melanotic—c. Melanoticum, Cancer, melanotic—c. Simplex, Scirrhus—c. Spongiosum, Encephaloid, Hæmatodes fungus—c. Scroti, Cancer, chimney-sweepers—c. Uteri, Metrocarcinoma, Metro-

scirrhus—c. *Ventriculi*, *Gastroscirrhus*; see *Gastrostenosis cardiaca et pylorica*.

CARCINOMATOUS, *Carcinomatous*, (F.) *Carcinomateux*. Relating to Carcinoma.

CARCINOME MOU ET SPONGIEUX, *Encephaloid*—c. *Sanguifant*, *Encephaloid*, *Inflammationes fungus*.

CARCINOS, *Cancer*.

CARCINOMES, (G.) *Karcinosen*, from *καρκινος*, 'a crab.' A family of diseases, according to the classification of Fuchs; which embraces the different forms of *Cancer*.

CARCINOSUS, *Cancerous*.

CARCINUS SPONGIOSUS, *Encephaloid*.

CARDAMANTICA, *Cardamine pratensis*, *Lepidium Iberis*.

CARDAMINDUM MAJUS, *Tropaeolum majus*.

CARDAMINE FONTANA, *Sisymbrium nasturtium*—c. *Nasturtium*, *Sisymbrium nasturtium*.

CARDAMINÆ PRATEN'SIS, *Cardaminæ*, *Cardamantica*, *Nasturtium Aquaticum*, *Cardamon*, *Culi flos*, *Iberis sophia*, *Nasturtium pratense*, *Ladies-mauve*, *Cuckoo-flower*, *Common Bitter Cress*, *Meadow Cress*, (F.) *Cresson élégant ou des prés*, *Passerage sauvage*. *Ord.* *Cruciferae*. The flowers have been considered useful as antispasmodics, in the dose of ʒi. to ʒij. They are probably inert.

CARDAMOM, *Cardamomum* (F.) *Cardamome*. The name of the fruit of various species of *Amomum* and *Elettaria*, respecting the botanical history of which there has been much confusion.

CARDAMOM, *BENGAL*, *Amomum maximum*—c. *Cluster*, *Amomum cardamomum*—c. *Greater*, *Amomum Grana Paradisi*—c. *Java*, *Amomum maximum*—c. *Lesser*, *Amomum cardamomum*—c. *Nepal*, *Amomum maximum*—c. *Round*, *Amomum cardamomum*.

CARDAMOME, *Amomum cardamomum*—c. *de la Côte de Malabar*, *Amomum cardamomum*.

CARDAMOMUM MAJUS, *Amomum grana paradisi*—c. *Minus*, *Amomum cardamomum*—c. *Piperatum*, *Amomum grana paradisi*—c. *Rotundum*, *Amomum cardamomum*—c. *Wild*, *Fagarastrum Capense*.

CARDAMON, *Cardamine pratensis*.

CARDAMUM MAJUS, *Tropaeolum majus*.

CARDÈRE, *Dipsacus sylvestris*—c. *Cultivé*, *Dipsacus fullonum*.

CARDIA, *καρδία*, 'the heart.' *Stomachus*, *Orificium sinistrum* seu *Ingressus superior* seu *Ostium oesophageum ventriculi*. The superior or oesophageal orifice of the stomach. Also, the Heart.

CARDIAC, *Cardiacus*, from *καρδία*, 'the heart;' or the upper orifice of the stomach. (F.) *Cardiaque*. Relating to the heart, *Carditic*, *Cardial*, (F.) *Cardiaque*, *Carditique*,—or to the upper orifice of the stomach. A *Cordial*.

CARDIAC ARTERIES, *Coronary arteries*, (F.) *Artères cardiaques ou coronaires*, are two in number. They arise from the aorta, a little above the free edge of the sigmoid valves, and are distributed on both surfaces of the heart.

CARDIAC GANGLION, *Ganglion cardiacum*, situated beneath the arch of the aorta, to the right side of the ligament of the ductus arteriosus. It receives the superior cardiac nerves of opposite sides of the neck, and a branch from the pneumogastric, and gives off numerous branches to the cardiac plexuses.

CARDIAC NERVES, (F.) *Nervi cardiaques*. These are commonly three on each side; a *superior*, *middle*, and *inferior*, which are furnished by corresponding cervical ganglia. Commonly, there are but two on the left side; the upper and middle, which draw their origin from the last two cervical ganglia. Scarpa calls the *superior*—

Cardiacus superficialis; the *middle*—*C. profundus* seu *magnus*; and the *inferior*—*C. parvus* seu *minor*. There are, besides, *Cardiac filaments*, (F.) *Filicæ cardiaques*, furnished by the par vagum or pneumo-gastric nerve, which become confounded with the above.

CARDIAC PLEXUS, *Plexus cardiacus*. There are three cardiac plexuses. 1. The *great cardiac plexus* is situated upon the bifurcation of the trachea. It is formed by the convergence of the middle and inferior cardiac nerves; and by branches from the pneumogastric, descendens noni, and first thoracic ganglion. 2. The *anterior cardiac plexus* is situated in front of the ascending aorta near its origin. It is formed by filaments from the superior cardiac nerves; from the cardiac ganglion; and from the great cardiac plexus. Filaments from this plexus accompany the left coronary artery, and form the *anterior coronary plexus*. 3. The *posterior cardiac plexus* is seated upon the posterior part of the ascending aorta near its origin. It is formed by numerous branches from the great cardiac plexus. It divides into two sets of branches, which together constitute the *posterior coronary plexus*.

CARDIAC VEINS, *Coronary Veins*, (F.) *Veines Cardiaques*, are commonly four in number; two anterior and two posterior. They open into the right auricle by one orifice, which is furnished with a valve, and is called, by Portal, *Sinus coronaire du Cœur*.

CARDIACA CRISPA, *Leonurus cardiaca*—c. *Passio*, *Cardialgia*—c. *Trilobata*, *Leonurus cardiaca*—c. *Vulgaris*, *Leonurus cardiaca*.

CARDIACUS, *Cordial*, *Stomachal*.

CARDIAGMUS, *Cardialgia*.

CARDIAGRA, *Affectio arthritica cordis*; from *καρδία*, 'the heart,' and *αγρᾱ*, 'seizure.' Gout of the heart, *Cardialgia*.

CARDIAGRAPHY, *Cardiagraphia*, from *καρδία*, 'the heart,' and *γραφῆν*, 'a description.' An anatomical description of the heart.

CARDIATRE, see *Cardiac*.

CARDIAL, see *Cardiac*.

CARDIALGIA, *Cardiaca Passio*, *Colica Ventriculi*, *Spasmus Ventriculi*, *Perodyn'ia*, *Cordolium*, *Cardilæ'a*, *Dyspeptodyn'ia*, *Dyspeptodyn'ia*, *Dyspeptodyn'ia*, *Peratodyn'ia*, *Cordiodyn'ia*, *Gastrodyn'ia*, *Gastralgia*, *Gasteralgia*, *Gastrocol'ia*, *Gastrodyn'ia*, *Passio Cardiacæ*, *Stomachalgia*, *Stomacalgia*, *Cardiacus Morbus*, *Cardiagmus*, *Cardialgy*; from *καρδία*, 'the cardiac orifice of the stomach,' and *αλγος*, 'pain.' Pain of the stomach, (F.) *Douleur de l'Estomac*, *D. névralgique de l'Estomac*. Also, *Heartburn*, (F.) *Cardialgie*, *Ardeur d'Estomac*, *A. du Cœur*. Impaired appetite, with gnawing or burning pain in the stomach or epigastrium,—*Morsus seu ardor ventriculi*, *Morsus stomachi*, *Sula*, *Limo'sia cardialgia mordens*, *Roxia Stomachi seu Ventriculi*—a symptom of dyspepsia.

CARDIALGIA INFLAMMATORIA, *Gastritis*—c. *Sputatoria*, *Pyrosis*.

CARDIALOG'IA, from *καρδία*, 'the heart,' and *λογος*, 'a discourse.' A treatise on the heart.

CARDIANASTROPHE, *Ectopia cordis*.

CARDIARCTIE, *Heart*, concentric hypertrophy of the. Contraction of the cavities of the heart.

CARDIARIUS; same etymology. A name given to a worm, said to have been found in the heart or pericardium.

CARDIATOM'IA, from *καρδία*, 'the heart,' and *τομή*, 'to cut.' Dissection of the heart.

CARDIATROPHIA, *Heart*, atrophy of the.

CARDIAUXE, *Heart*, hypertrophy of the.

CARDIECTASIS, *Dilatation* of the heart, see *Aneurism* of the heart—c. *Partialis*, *Aneurism* of the heart.

CARDIELCOSIS: from καρδία, 'the heart,' and 'ελκος,' 'an ulcer.' Ulceration of the heart.
CARDIETHMOLIPOSIS, Steatosis cordis.
CARDIEURYSMA, Aneurism of the heart.
CARDILÆA, Cardialgia.

CARDIM'ELECH, from καρδία, 'the heart,' and מֶלֶךְ, *Melek*, (Hebr.) 'a governor.' A supposititious active principle seated in the heart, and governing the vital functions.—Dolæus.

CARDINAL FLOWER, *Lobelia cardinalis*—c. f. Blue, *Lobelia spiliotica*.

CARDINAL PLANT, *Lobelia cardinalis*.

CARDINAMENTUM, Ginglymus, Gomphosis.

CARDIOBOTANUM, *Centaurea benedicta*.

CARDIOCE'LE, from καρδία, 'the heart,' and κλη, 'rupture.' Hernia of the heart, especially into the abdominal cavity.

CARDIOCLASIE, *Cardiorrhæxis*.

CARDIODYNÊ, *Cardiodyn'ia*; from καρδία, 'the heart, the stomach,' and δυνν, 'pain.' Pain in the heart. Also, Cardialgia.

CARDIDYNE SPASMODICA INTERMITTENS, *Angina pectoris*.

CARDIOG'MUS. Hippocrates employed this word καρδιωγμος, synonymously with cardialgia. In the time of Galen it was used, by some writers, for certain pulsations of the heart, analogous to palpitations. Sauvages understood by *Cardiog-mus* an aneurism of the heart or great vessels, when still obscure. Also, *Angina pectoris*.

CARDIOG'MUS CORDIS SINISTRI, *Angina pectoris*.

CARDIOMALA'CIA, *Malaco'sis seu Mala'cia seu Malar'is seu Molli'ties Cordis*, (F.) *Ramollissement du Cœur*, from καρδία, 'the heart,' and μαλακία, 'softness.' Softening of the heart, caused by inflammation of the organ, or a consequence of some lesion of the function of nutrition.

CARDIOMETRY, *Cardiomet'ria*, from καρδία, heart, and μετρον, 'measure.' Measurement of the heart, as by percussion and auscultation.

CARDIOMYOLIPOSIS, *Steatosis cordis*.

CARDIONCHI, see Aneurism.

CARDIONEURALGIA, *Angina pectoris*.

CARDION'OSUS, *Morbis cordis*, from καρδία; 'heart,' and νοσος, 'disease.' Disease of the heart. Heart disease.

CARDIOPALMUS, *Cardiotromus*.

CARDIOPERICARDITIS, see Pericarditis.

CARDIORRHEU'MA, *Rheumat'is mus cordis*; from καρδία, 'the heart,' and ρευμα, 'defluxion, rheumatism.' Rheumatism of the heart.

CARDIORRHEX'IS, *Cardioclasis*, (Piorry,) *Ruptura cordis*, (F.) *Rupture du Cœur*, from καρδία, 'the heart,' and ρηξις, 'laceration.' Laceration of the heart.

CARDIOSCLÉROSIE, (Piorry), from καρδία, 'the heart,' and σκληρος, 'hard.' (F.) *Endurcissement du Cœur*. Induration of the heart.

CARDIOSTENO'SIS, *Stenocard'ia*, from καρδία, 'the heart,' and στενωσις, 'contraction.' Contraction of the openings of the heart.

CARDIOTRAU'MA, from καρδία, 'the heart,' and τραυμα, 'a wound.' A wound of the heart.

CARDIOTROMUS, *Palpitatio Cordis trepidans*, *Cardiopal'mus*, *Trepidatio Cordis*, from καρδία, 'the heart,' and τρομος, 'tremor.' Rapid and feeble palpitation or fluttering of the heart.

CARDIOTROTUS, from καρδία, 'the heart,' and τροσκω, 'I wound.' One affected with a wound of the heart.—Galen.

CARDIPERICARDITIS, see Pericarditis.

CARDITE, *Carditis*.

CARDITIC, *Cardiac*. Also, relating or belonging to carditis.

CARDITIS, from καρδία, 'the heart,' and the termination *itis*. Inflammation of the fleshy substance of the heart. *Empres'ma Carditis*, *Inflamma'tio Cordis*, *I. Carditis*, *Cuma Carditis*, *Myocarditis*, *Carditis Muscularis*, (F.) *In-*

flammation du Cœur, *Cardite*. The symptoms of this affection are by no means clear. They are often confounded with those of pericarditis, or inflammation of the membrane investing the heart. Carditis, indeed, with many, includes both the inflammation of the external investing membrane and that of the interior of the heart. See Pericarditis, and Endocarditis.

CARDITIS EXTERNA, Pericarditis—c. Interna, Endocarditis—c. Muscularis, Carditis—c. Membrana, Pericarditis—c. Polyposa, Polypi of the heart—c. Serosa, Pericarditis.

CARDO, Ginglymus.

CARDOPATIUM, *Carlina acaulis*.

CARDOPERICARDITIS, see Pericarditis.

CARDUUS ALTILIS, *Cynara scolymus*—c. Benedictus, *Centaurea benedicta*—c. Brazilianus, *Bromelia ananas*—c. Domesticus capite majori, *Cynara scolymus*—c. Hemorrhoidalis, *Cirsium arvense*.

CARDUUS MARIA'NUS, *C. Maria'na seu lactens*, *Styl'bium*, *S. Maria'num seu maculatum*, *Carthamus maculatus*, *Cirsium maculatum*, *Spina alba*, *Common Milk Thistle*, *Ladies' Thistle*, (F.) *Chardon-Marie*. Ord. Compositæ. The herb is a bitter tonic. The seeds are oleaginous. It is not used.

CARDUUS PINEUS, *Atractylis gummifera*—c. Sativus, *Carthamus tinctorius*—c. Sativus non-spinosus, *Cynara scolymus*—c. Solstitialis, *Centaurea calcitrapa*—c. Stellatus, *Centaurea calcitrapa*—c. Tomentosus, *Onopordium acanthium*—c. Veneris, *Dipsacus fullonum*.

CAREBARE'SIS, *Carebaria*.

CAREBARIA or **CAREBARI'A**, *Carebare'sis*, from καρν, 'the head,' and βαρος, 'weight.' *Scordine'ma*, *Cerebaria*, *Scordini'smus*, *Cordine'ma*. Heaviness of the head.—Hippocrates, Galen.

CARE'NA, *Kare'na*. The twenty-fourth part of a drop.—Ruland and Johnson.

CARE-TAKER, Nurse.

CAREUM, Carum.

CAREX ARENARIA, *Sarsaparilla Germanica*.

CARIACOU. A beverage, used in Cayenne, and formed of a mixture of cassava, potato, and sugar, fermented.

CAR'IOA, from *Caria*, where the fig was cultivated. See *Ficus carica*.

CAR'ICA PAPA'YA, *Papaw tree*, *Pawpaw*, (F.) *Papayer*. Ord. Cucurbitaceæ. A native of America, India, and Africa. The fruit has somewhat of the flavour of the pumpkin, and is eaten like it. The milky juice of the plant and the seed and root have been regarded as anthelmintic.

The *papaw*, of North America, belongs to the order Anonaceæ, (*Ano'na seu Asi'mina seu Porcel'ia seu Uva'ria tri'loba*, *Ficus Indica*, *Orchidocar'pum orietinum*), Custard Apple Family.

CAR'IOUM. Said to have been named after its inventor Caricus. *Car'yum*. A detergent application to ulcers; composed of black hellebore, sandarach, copper, lead, sulphur, orpiment, cantharides, and oil of cedar.—Hippocrates.

CARIE, Caries—c. *des Dents*, Dental gangrene.

CARIÉ, Carious.

CARIES, *Nigrit'ies Os'sium*. An ulceration of bone, *Osteoche'losis*.—Necrosis being death of a bone. It resembles the gangrene of soft parts. Hence it has been termed *Caries gangranosa*, *Gangra'na Caries seu Os'sium*, *Tere'do*, *Arro'sio*, *Euros*, (F.) *Carie*. It is recognised by the swelling of the bone which precedes and accompanies it; by the abscesses it occasions; the fistula which form; the sanious character, peculiar odour and quantity of the suppuration, and by the evidence afforded by probing. The most common causes of caries are blows, the action of some virus, and morbid diatheses. When dependent on

any virus in the system, *this* must be combated by appropriate remedies. When entirely local, it must be converted, where practicable, into a state of necrosis or death of the affected part. For this end, stimulants, the actual cautery, &c., are applied.

CARIES, DENTUM, Dental gangrene—c. Pudentum, see CHANCRO—c. of the Vertebrae, Vertebral disease—c. Vertebrarum, Vertebral disease.

CARIUEUX, Carious.

CARIM CURINI, *Justitia ecobolium*.

CAR'NA, 'a ship's keel.' The vertebral column, especially of the fetus. Also, the breast-bone bent inwards. Hence, *Pectus carinatum*:—the chest affected with such deformity.

CAR'IOUS, *Cariousus*, *Euro'des*, (F.) *Carié*, *Carieus*. Affected with caries.

CARIUM TERRE, Calx.

CARIVE, *Myrtus pimenta*.

CARIVILLANDI, *Smilax sarsaparilla*.

CARLINA, 'Carlina Thistle.'

CARLINA ACAULIS, *C. chamaeleon*, *Chamaeleon album*, *Card-patium*, (F.) *Carlina sans tige*. Ord. Compositæ, which grows in the Pyrenees, and on the mountains of Switzerland, Italy, &c., has been recommended as a tonic, emmenagogue, and sudorific.

CARLINA CHAMELEON, *C. acaulis*.

CARLINE SANS TIGE, *Carlina acaulis*.

CARLISLE SPRINGS. These are about five miles from the town of Carlisle, in Pennsylvania. The water is mildly sulphureous.

CARLO SANCTO RADIX. 'St. Charles's Root,' found in Mechoacan in America. The bark is aromatic, bitter and acrid. It is considered to be sudorific, and to strengthen the gums and stomach.

CARLSBAD, MINERAL WATERS OF. Carlsbad is a town in Bohemia, 24 miles from Egra, celebrated for its hot baths. The water contains about 47 parts in the 100 of purging salts. It is a thermal saline; temperature 121° to 147° Fahrenheit. The constituents are—carbonic acid, sulphate of soda, carbonate of soda, and chloride of sodium.

CARMANTINE, *Justitia pectoralis*—c. *Pectorale*, *Justitia pectoralis*.

CARMEN, 'a verse.' An amulet. A charm, which, of old, often consisted of a verse. See Charm.

CARMINANTIA, Carminatives.

CARMINATIVA, Carminatives.

CARMINATIVES, *Carminant'ia seu Carminat'ia*, from *carmen*, 'a verse,' or 'charm,' *Antiphys'ica*, *Physagoga*, *Xan'tica*, *Antiphysel'ica*, (F.) *Carminat'is*. Remedies which allay pain, 'like a charm,' by causing the expulsion of flatus from the alimentary canal. They are generally of the class of aromatics.

THE FOUR GREATER CARMINATIVE HOT SEEDS, *Quatuor sem'ina cal'ida majo'ra carminat'iva*, were, of old, anise, carui, cummin, and fennel.

THE FOUR LESSER CARMINATIVE HOT SEEDS, *Quatuor sem'ina cal'ida mino'ra*, were bishop's weed, stone parsley, smallage, and wild carrot.

CARMOT. A name given, by the alchemists, to the matter which they believed to constitute the Philosopher's stone.

CARNABADIA, Carum, (seed.)

CARNABADIUM, *Cuminum cyminum*.

CARNATIO, Syssarcosis.

CARNATION, *Dianthus caryophyllus*.

CARNELIAN, Cornelian.

CARNEOLUS, Cornelian.

CAR'NEOUS, *Carno'us*, *Sarco'des*, *Incarna'tus*, from *caro*, *carnis*, 'flesh,' (F.) *Charnu*. Consisting of flesh, or resembling flesh.

CARNEOUS COLUMNS, *Fleshy Columns*, *Colum'*

na Carneæ, of the heart, (F.) *Colonnes charnues*, are muscular projections, situate in the cavities of the heart. They are called, also, *Musculi Papillares*.

CARNEOUS FIBRES, *Fleshy Fibres*, *Muscular Fibres*, (F.) *Fibres charnues ou musculaires*, are fibres belonging to a muscle.

CARNEUM MARSUPIUM, Ischio-trochanterianum.

CARNIC'ULA. Diminutive of *caro*, *carnis*, 'flesh.' The gum.—Fallopian.

CARNIFICA'TIO, Carnification—c. Pulmonum, Hepatification of the lungs.

CARNIFICA'TION, *Carnificat'io*, from *caro*, *carnis*, 'flesh,' and *geri*, 'to become.' Transformation into flesh. A morbid state of certain organs, in which the tissue acquires a consistence like that of fleshy or muscular parts. It is sometimes observed in hard parts, the texture becoming softened, as in *Osteo-sarcoma*. When it occurs in the lungs, they present a texture like that of liver. Such is the condition of the foetal lung.

CARNIFORMIS ABSCESS'US. An abscess, which ordinarily occurs in the neighborhood of the articulations, and whose orifice is hard, the sides thick and callous.—M. A. Severinus.

CARNIVOROUS, *Carniv'orus*, *Sarcoph'agus*, *Creatoph'agus*, *Crepoph'agus*, *Zoiph'agus*, *Creatoph'agous*, *Crepoph'agous*, *Creatoph'agous*, (F.) *Carnivore*; from *caro*, *carnis*, 'flesh,' and *voro*, 'I eat.' That which eats flesh. Any substance which destroys excrescences in wounds, ulcers, &c.

CARNOSA CUTIS, *Panniculus carnosus*.

CARNOS'ITAS, (F.) *Carnosité*, from *caro*, *carnis*, 'flesh.' A fleshy excrescence.

CARNOSITIES OF THE URE'THRA, *Car'uncles in the Ure'thra*, (F.) *Carnosités ou Caroncules de l'urètre*. Small fleshy excrescences or fungous growths, which were, at one time, presumed to exist in the male urethra, whenever retention of urine followed gonorrhœa.

M. Cullérier uses the term *Carnosité vénérienne* for a cutaneous, cellular, and membranous tumour, dependent upon the syphilitic virus. See, also, *Polysarcia*.

CARNOSUS, Carneous.

CARO, Flesh—c. Accessoria, see *Flexor longus digitorum pedis profundus perforans*, (*accessorius*)—c. *Excrecens*, *Excrescence*—c. *Fungosa*, *Fungosity*—c. *Glandulosa*, *Epiglottic gland*—c. *Luxurians*, *Fungosity*—c. *Orbicularis*, *Placenta*—c. *Parenchymatica*, *Parenchyma*—c. *Quadrata*, *Palmaris brevis*—c. *Quadrata Sylvii*, see *Flexor longus digitorum pedis profundus perforans*, (*accessorius*)—c. *Viscerum*, *Parenchyma*.

CAROB TREE, *Ceratonia siliqua*.

CAROA ALNABATI, *Ceratonium siliqua*.

CARODES, Carotic.

CAROLI, see *Chancro*.

CAROL'NA, NORTH, MINERAL WATERS OF. In the counties of Warren, Montgomery, Rockingham, Lincoln, Buncomb, and Rowan, there are mineral springs. They belong generally to the sulphureous or acidulous saline.

CAROLINA, SOUTH, MINERAL WATERS OF. They are numerous. Paocelet Springs, on the west bank of Paocelet River, contain sulphur and iron. Many, with similar properties, but not held in estimation, are scattered about the State.

CARONCULE, Caruncle—c. *Lachrymale*, *Caruncle*.

CARONCULES MYRTIFORMES, *Caruncule myrtiformes*—c. *de l'Urètre*, *Carnosities of the urethra*.

CARONCULEUX, Carunculous.

CAROPI, *Amomum cardamomum*.

CAROSIS, Somnolency.

CAROTA, see *Daucus carota*.

CAROT'IC, *Carot'icus*, *Carot'id*, *Carot'idus*, *Carotidæus*, *Caro'des*, *Com'atose*, from *καρος*, 'stupor,' (F.) *Carotique*. Relating to stupor or carus—as a *carotic state*;—or to the carotids, (F.) *Carotidien*.

CAROTIC ARTERIES, Carotids—c. Ganglion, see Carotid Nerve—c. Nerve, Carotid Nerve—c. Plexus, see Carotid Nerve.

CAROTICA, Narcotics.

CAROTICUS, Carotid.

CAROTID, Carotic.

CAROTIDÆUS, Carotic.

CAROTIDIEN, Carotic.

CAROTIDS, *Carot'ides*, *Carot'icæ*, *Carotidæ*, *Capita'les*, *Jugula'res*, *Sopora'les*, *Sopora'riæ*, *Soporif'era*, *Somnif'era*, *Apoplec'ticæ*, *Lethargicæ* (*Arter'icæ*), *Carot'id Ar'teries*, *Cephal'ic Ar'teries*, (F.) *Ar'tères Carotides*; from *καρος*, 'stupor,' [?]. The great arteries of the neck, which carry blood to the head. They are divided into, 1. *Primitive* or *common*; the left of which arises from the aorta, and the right from a trunk, common to it and the subclavian. 2. *External* or *pericephalic*, branch of the primitive, which extends from the last to the neck of the condyle of the lower jaw; and, 3. *Internal*, *Ar'teria cerebra'lis* vel *encephalicæ*, another branch of the primitive, which, arising at the same place as the external, enters the cranium, and terminates on a level with the fissure of Sylvius, dividing into several branches.

CAROTID or **CAROTIC CANAL**, *Canalis Carot'icus*, *Canal inflexe de l'os temporal*—(Ch.). *Canal carotidien*, is a canal in the temporal bone, through which the carotid artery and several nervous filaments pass.

CAROTID or **CAROTIC FORAM'INA**, *Foram'ina Carot'ica*, (F.) *Trois carotidiens*, are distinguished into *internal* and *external*. They are the foramina at each extremity of the *Canalis Carot'icus*.

CAROTID GANGLION, see Carotid nerve.

CAROTID NERVE, *Carot'ic nerve*, *Nervus carot'icus*. A branch from the superior cervical ganglion of the great sympathetic, which ascends by the side of the internal carotid. It divides into two portions, which enter the carotid canal, and, by their communication with each other and the petrosal branch of the vidian, form the *carotid plexus*. They also frequently form a small gangliform swelling on the under part of the artery—the *carotic* or *carotid* or *cavernous ganglion*, *ganglion de Laumonier*.

CAROTID PLEXUS, see Carotid nerve.

CAROTTE, *Daucus carota*.

CAROUA, *Carum* (seed).

CAROUBIER, *Ceratonium siliqua*.

CAROUGE, see *Ceratonium siliqua*.

CARPAL, *Carpa'lis*; from *καρπος*, 'the wrist,' Belonging or relating to the carpus or wrist.

CARPASA, *Carbasa*.

CARPA'SIUM, *Car'pasum*, and *Carpe'sium*. Dioscorides, Pliny, Galen, &c., have given these names, and that of *Carpasos*, to a plant, which cannot now be determined, and whose juice, called *Opunc'arpasos*, *στακαρισσος*, passed for a violent, narcotic poison, and was confounded with myrrh.

CARPATICUM, see *Pinus cembra*.

CARPE, *Carpus*.

CARPENTARIA, *Achillea millefolium*.

CARPENTER'S LEAF, *Galax aphylla*.

CARPESIUM, *Carpasium*.

CARPHODES, *Carpho'des*, from *καρφος*, 'flocculus,' and *ειδος*, 'resemblance.' Flocculent, stringy;—as *mucus carphodes*, flocculent or stringy mucus.

CARPHOLOG'IA, *Tilimus*, *Carpholog'ia*, *Croc'idismus*, *Crocylidismus*, *Floccid'rum venutii*, *Floccid'gium*, *Tricholog'ia*, *Croc'idic'is*, *Floccid'ation*, *Floccid'ation*, from *καρφος*, 'flocculus,' and *λεγω*,

'I collect,' or 'pluck,' (F.) *Carphologie*. Action of gathering flocculi. A delicious picking of the bed-clothes, as if to seek some substance, or to pull the flocculi from them. It denotes great cerebral irritability and debility, and is an unfavourable sign in fevers, &c.

CARPHOS, *Trigonella fœnum*.

CARPIA, *Lintum*.

CARPIÆUS, *Palmaris brevis*.

CARP'IAL, *Car'pian*, *Carpi'anus*, *Carpi'al'is*, (F.) *Carpien*. Belonging to the Carpus.

CAR'IAL LIG'AMENTS, (F.) *Ligaments Carpiens*, are, 1. The fibrous fasciæ, which unite the bones of the carpus; and, 2. The annular ligaments, anterior and posterior.

CARPIAN, *Carpial*.

CARPIEN, *Carpial*.

CARPISMUS, *Carpus*.

CARPOBALSAMUM, see *Amyris opobalsanum*.

CARPOLOGIA, *Carphologia*—c. *Spasmodica*, *Subsultus tendinum*.

CARPO-METACARPEUS MINIMI DIGITI, *Adductor metacarpi minimi digiti*—c. *Métacarp'ien du petit doigt*, *Opponens minimi digiti*—c. *Métacarp'ien du pouce*, *Opponens pollicis*—c. *Phalangeus minimi digiti*, *Abductor minimi digiti*—c. *Phalangien du petit doigt*, *Abductor minimi digiti*, *Flexor parvus minimi digiti*—c. *Phalangien du pouce*, *Flexor brevis pollicis manus*—c. *Sus-phalangien du pouce*, *Abductor pollicis brevis*.

CARPO-PEDAL, from *carpus*, 'the wrist,' and *pes*, *pedis*, 'the foot.' Relating to the wrist and foot.

CARPO-PEDAL SPASM, *Cer'eb'ral spasmod'ic croup*. A spasmodic affection of the chest and larynx in young children, accompanied by general or partial convulsions. The disease commonly occurs between the third and ninth month, and is characterized by excessive dyspnœa, accompanied by a loud croupy noise on inspiration; the thumbs being locked, and the hands and feet rigidly bent for a longer or shorter period. The seat of the disease is evidently in the cerebro-spinal axis, primarily or secondarily: generally, perhaps, it is owing to erethism seated elsewhere, but communicated to the cerebro-spinal centre, and reflected to the respiratory and other muscles concerned. It seems to be connected with dental irritation; and consequently, in the treatment, where such is the case, the gums should be freely divided; after which, cathartics and revulsives, with the use of narcotics and appropriate diet, will generally remove the affection; for although extremely alarming, it is often not attended with great danger. See *Asthma thymicum*.

CARPOS, *Fruit*.

CARPOTICA, from *καρπος*, 'fruit.' Diseases affecting impregnation. Irregularity, difficulty or danger produced by parturition:—the 3d order, class *Genetica*, of Good.

CARPUS, *Carpi'smus*, *Brachia'l'is*, *Rasce'ta*, *Rasce'ta*, *Rascha*, *Rasce'ta*, *Rasce'ta*, *Wrist*, (Prov.) *Hand-wrist*, *Shackle*, (F.) *Carpe*, *Poignet*. The part between the fore-arm and hand. Eight bones compose it (in two rows). In the superior row there are, from without to within—the *Scaphoides* or *naviculat'rè*, *Luna'rè* or *semiluna'rè*, *Cuneifor'mè*, and *Orbicula'rè* or *pisifor'mè*. In the lower row—*Trapezium*, *Trapezoi'des*, *Magnus*, and *Uncifor'mè*.

CARRAGEEN MOSS, *Fucus crispus*.

CARRÉ, *Quadratus*—c. *de la Cuissè*, *Quadratus femoris*—c. *des Lombes*, *Quadratus lumborum*—c. *du Menton*, *Depressor labii inferioris*—c. *du Pied*, *Extensor brevis digitorum pedis*.

CARREAU, *Tabes mesenterica*.

CARRÉE, see *Flexor longus digitorum pedis profundus perforans* (accessorius).

CARRELET, (F.) *Acus triangularis*. A straight needle, two or three inches long, the point of which is triangular; and which the ancients used in different operations. Also, a wooden, triangular frame for fixing a cloth through which different pharmaceutical preparations are passed.

CARRON OIL, *Linimentum aquæ calcis*.

CARROT, CANDY, *Athamanta cretensis*—c. *Deadly*, *Thapsia*.

CARROT PLANT, *Daucus carota*.

CARTHAGENA BARKS, see *Cinchona*.

CARTHAMUS MACULATUS, *Carduus marianus*.

CARTHAMUS TINCTORIUS, *Am'yrion*, *Cnicus*, *Crocus Germanicus* seu *Saracenicus*, *Carthaminum officinarum*, *Car'duus sativus*, *Saffranum*, *Saffron-flower*, *Safflower*, *Saffron*, *Barbadoe Saffron*, *Dyer's Saffron*, (F.) *Carthame*, *Safran bâlard*, *Carthame des Teinturiers*. Ord. *Compositæ*. Sex. Syst. *Syngenesia Polygamia æqualis*. The seeds are aromatic, cathartic, and diuretic; yet to the parrotquet they are an article of food; hence their name, *Graines de Parrotquet*. The flowers, *Car'thamus* (Ph. U. S.), are employed as a cosmetic, and are a reputed diaphoretic. [?]

CARTHEGON, see *Buxus*.

CARTILAGE, *Chondros*, *Cartilago*, *Gristle*, (Sc.) *Girle*, (F.) *Cartilage*. A solid part of the animal body, of a medium consistence between bone and ligament, which in the fœtus is a substitute for bone, but in the adult exists only in the joints, at the extremities of the ribs, &c. Cartilages are of a whitish colour, flexible, compressible, and very elastic, and some of them apparently inorganic. They are composed, according to J. Davy, of .44 albumen, .55 water, and .01 phosphate of lime.

CARTILAGE ANONYME, *Cricoid*, (cartilage)—c. *Epiglotticæ*, *Epiglottidis*—c. *Mucronæ*, *Xiphoid Cartilage*—c. *Supra-arytenoid*, *Corniculum laryngis*—c. *Tarsal*, see *Tarsus*.

CARTILAGES, ARTICULAR, *Obdu'cent Cartilages*, invest bony surfaces, which are in contact; hence they are called *investing* or *incrusting cartilages*, (F.) *Cartilages de revêtement ou d'encroûtement*.

CARTILAGES, CUNEIFORM, OF THE LARYNX, see *Cuneiform*.

CARTILAGES EBURNÉS, see *Eburnification*.

CARTILAGES, INTERARTICULAR, *Menisci*, are such as are situate within the joints, as in the knee joint.

CARTILAGES, INTERVERTEBRAL, see *Intervertebral*.

CARTILAGES OF OSSIFICATION are such as, in the progress of ossification, have to form an integrant part of bones; as those of the long bones in the new-born infant. They are termed *temporary*; the others being *permanent*. All the cartilages, with the exception of the articular, are surrounded by a membrane analogous to the periosteum, called *Perichondrium*.

CARTILAGES OF THE RIBS are, in some respects, only prolongations of the ribs. Those of the nose, of the meatus auditorius, and Eustachian tube, present a similar arrangement. Other cartilages resemble a union of fibrous and cartilaginous textures; hence their name *Fibro-cartilages*.

CARTILAGES OF SANTORINI, *Corniculum Laryngis*—c. *Semilunar*, see *Semilunar*—c. *Sigmoid*, *Semilunar cartilages*—c. of *Wrisberg*, *Cuneiform Cartilages of the Larynx*.

CARTILAGINES GUTTURALES, *Arytenoid cartilages*—c. *Semilunares*, *Semilunar cartilages*—c. *Sigmoideæ*, *Semilunar cartilages*.

CARTILAGINIS ARYTENOIDÆ CAPITULUM, *Corniculum laryngis*.

CARTILAGINOUS, *Cartilagineus*, *Cartilagin'ous*, *Chondro'ides*, *Chondroi'des*, (F.) *Cartilagineus*. Belonging to, or resembling cartilage.

CARTILAGINOUS TISSUE, see *Tissue*.

CARTILAGO, *Cartilage*—c. *Clypealis*, *Thyroid cartilage*—c. *Ensiformis*, *Xiphoid cartilage*—c. *Gutturalis*, *Arytenoid cartilage*—c. *Innominata*, *Cricoid*—c. *Mucronata*, *Xiphoid cartilage*—c. *Peltalis*, *Thyroid cartilage*, *Xiphoid cartilage*—c. *Peltatus*, *Thyroid cartilage*—c. *Scutiformis*, *Thyroid cartilage*—c. *Uvifer*, *Uvula*—c. *Xiphoides*, *Xiphoid cartilage*.

CARUM, from *Caria*, a province of Asia. *A'pium* seu *Bu'nium* seu *Ligusticum* seu *Soseli carvi* seu *carum*, *Sium carvi*, *Car'ueum*, *Carum carvi*, *Carvi*, *Quinnum pratense*, *Cucius*, *Car'oon*, the *Car'away*, (F.) *Carvi*, *Camin des prés*. Order, *Umbelliferae*. Sex. Syst. *Pentandria Digynia*. The seeds, *Caraway seeds*, (Prov.) *Carvy seeds*, *Carnab'dia*, *Car'oua*, *Carum* (Ph. U. S.), are carminative. Dose, gr. x to ʒj. swallowed whole or bruised. The oil, *Oleum Car'ui*, (F.) *Huile de carvi*, has the properties of the seeds. Dose, gtt. ij to vj. *Aqua Car'ui*, *Car'away Water*, is official in the Pharmacopœias of London and Dublin.

CARUM BULBOCASTANUM, *Bunium bulbocastanum*.

CAR'UNCLE, *Carun'cula*, diminutive of *caro*, 'flesh.' A small portion of flesh, *Sar'cium*, *Sarcid'ium*. A fleshy excrescence,—*Ephy'na carun'cula*, (F.) *Caruncule*.

CARUNCLE, *Carnositas*.

CARUNCLES IN THE URETHRA, *Carnosities*.

CARUN'CULA LACRYMALIS, (F.) *Caruncule lacrymale*. A small, reddish, follicular body, situate at the inner angle of the eye. It secretes a gummy substance.

CARUN'CULA SEMINALIS, *Gallinaginis caput*.

CARUNCULE CUTICULARES, *Nymphæ*.

CARUNCULE MAMMILLARES. The extremities of the lactiferous tubes in the nipples. The olfactory nerves have been so called by some.

CARUNCULE MYRTIFORMES, *C. Vaginal'es*, *Glandulæ myrtiformes*, (F.) *Caruncules myrtiformes*. Small, reddish tubercles, more or less firm, of variable form, and uncertain number, situate near the orifice of the vagina, and formed by the mucous membrane. They are regarded as the remains of the hymen.

CARUNCULE PAPILLARES, *Papillæ of the kidney*.

CARUN'CULOUS, *Carun'cular*, (F.) *Carunculeux*. Relating to caruncles or carnosities.

CARUON, *Carum*.

CARUS, *kapos*, *Supor caroticus*, *Profound sleep*. The last degree of coma, with complete insensibility, which no stimulus can remove, even for a few instants. *Sopor*, *Coma*, *Lethargia*, and *Carus*, are four degrees of the same condition.

CARUS APOPLEXIA, *Apoplexy*—c. *Asphyxia*, *Asphyxia*—c. *Catalepsia*, *Catalepsy*—c. *Ectasis*, *Ectasis*—c. *Hydrocephalus*, *Hydrocephalus internus*—c. ab *Insolatione*, *Coup de soleil*—c. *Lethargus*, *Lethargy*—c. *Lethargus cataphora*, *Somnolency*—c. *Lethargus vigil*, *Coma vigil*—c. *Paralysis*, *Paralysis*—c. *Paralysis paraplegia*, *Paraplegia*—c. *Vernus*, *Lethargy*.

CARVI, *Carum*.

CARVY SEEDS, see *Carum*.

CARYA, *Hickory*, *Juglans regia*—c. *Basilica*, *Juglans regia*.

CARYDION, *Corylus avellana*.

CARYEDON CATAGMA, see *Fracture*.

CARYOCOSTINUS, *Caryocostinum*. An electuary prepared of the costus and other aromatic substances, &c. It was cathartic. See *Confectio scammonia*.

CARYON, *Juglans regia*—c. Ponticon, *Corylus avellana* (nut.).

CARYOPHYLLA, *Geum urbanum*.

CARYOPHYLLATA AQUATICA, *Geum rivale*—c. Nutans, *Geum rivale*—c. Urbana, *Geum urbanum*—c. Vulgaris, *Geum urbanum*.

CARYOPHYLLUM RUBRUM, *Dianthus caryophyllus*.

CARYOPHYLLUS AMERICANUS, see *Myrtus pimenta*—c. Aromaticus, *Eugenia caryophyllata*—c. Hortensis, *Dianthus caryophyllus*—c. Pimenta, *Myrtus Pimenta*—c. Vulgaris, *Geum urbanum*.

CARYOTL. The best kind of dates.—Galen.

CAS RARES (F.), *Rare cases*. This term is used, by the French, for pathological facts, which vary from what is usual. See a celebrated article under this head in the *Dictionnaire des Sciences Médicales*, Vol. IV.

CASAMUM, *Cyclamen*.

CASAMUNAR, *Cassumuniar*.

CASCADING, Vomiting.

CASCARA, CASCARILLA. Spanish words, which signify *bark* and *little bark*, under which appellations the bark (*Cinchona*) is known in Peru. They are now applied to the bark of *Croton cascarilla*. The bark-gatherers are called *Cascarilleros*.

CASCARILLA, *Cascara*, *Croton cascarilla*—c. Carabaya, see *Cinchona cordifolia* cortex—c. Colorado, *Cinchona oblongifolia* cortex—c. Roxa, *Cinchona oblongifolia* cortex.

CASCARILLEROS, see *Cascara*.

CASCHEU, *Catechu*.

CASE, *Capsa*, *Theca*, (F.) *Cistise*. This name is given to boxes for the preservation of instruments, or of medicines necessary in hospital or other service. We say, e. g.—*A case of amputating*, or of *trepanning instruments*. A pocket case, *Armamentarium portabile*, (F.) *Trousse*, contains the smaller instruments in constant use with the surgeon.

CASE, *Casus*, from *cadere*, *casum*, 'to fall.' The condition of a patient;—as a *case of fever*, &c. (F.) *Observation*. Also, the history of a disease.

CASEARIUS, Cheesy.

CASEIN, *Caseine*, *Cas'eum*, *Galactine*, *Tyrine*, *Lactalbumen*, *Caseous matter*, (F.) *Caseïne*, *Matière caseuse*, from *caseus*, 'cheese.' The only nitrogenized constituent of milk. It is identical in composition with the chief constituents of blood,—fibrin and albumen, all being compounds of protein. A similar principle exists in the vegetable, *Vegetable Casein* or *Legu'min*, *Veg'etable Gluten*. It is chiefly found in leguminous seeds—peas, beans, lentils. Like vegetable albumen, Casein is soluble in water; and the solution is not coagulable by heat.

CASEIN, BLOOD Globulin—c. of the Saliva, see *Saliva*—c. of the Small intestine, *Albuminose*.

CASEOSUS, Cheesy.

CASEOUS MATTER, Casein.

CASEUM, Casein.

CASEUS, Cheese—c. Equinus, Hippace.

CASEUX, Cheesy.

CASHEW, *Anacardium occidentale*.

CASHOO. An aromatic drug of Hindoostan, said to possess pectoral virtues.

CASHOW, *Catechu*.

CASIA, *Laurus cassia*.

CASMINA, *Cassumuniar*.

CASMONAR, *Cassumuniar*.

CASSA, *Thorax*.

CASSADA ROOT, *Jatropha manihot*.

CASSAVA, BITTER, *Jatropha manihot*—c. Root, *Jatropha manihot*.

CASSE AROMATIQUE, *Laurus cassia*—c. en Bâton, *Cassia fistula*—c. en Bois, *Laurus cas-*

sia—c. des Boutiques, *Cassia fistula*—c. Séné, *Cassia senna*.

CASSE-LUNETTE, *Centaurea cyanus*, *Cyanus segetum*, *Euphrasia officinalis*.

CASSEENA, *Ilex vomitoria*.

CASSENOLES, see *Quercus infectoria*.

CASSIA, *Laurus cassia*—c. Absus, *Absus*—c. Acutifolia, *C. senna*—c. Egyptian, *C. senna*—c. Alexandrina, *C. fistula*—c. Bonplandiana, *C. fistula*.

CASSIA CHAMÆCRISTA, *Prairie senna*, *Partridge Pea*, *Wild Senna*, an indigenous plant, Order, Leguminosæ, which flowers in August. It resembles *Cassia Marilandica* in properties.

CASSIA CINNAMOMEA, *Laurus cassia*—c. Caryophyllata, *Myrtus caryophyllata*—c. Canella, *Laurus cassia*—c. Egyptian, *Cassia senna*—c. Excelsa, *C. fistula*.

CASSIA FISTULA, *C. nigra* seu *fistularis* seu *Alexandrina* seu *coccinea* seu *Bonplandiana*, *Canna*, *C. solutaria* seu *fistula*, *Cathartocarpus*, *Bactrilo'bium fistula*, *Purging Cassia*, (F.) *Casse Canefier*, *Canefier* ou en bâtons ou des Boutiques. The pulp of *Cassia Fistula* or *Cathartocarpus Fistula*, *Pudding pipe tree*; Fam. Leguminosæ; See. Syst. Decandria Monogynia, *Pulpa Cassia*, *Cassia Aramentum*, *Cassia Fistula Pulpa*, (Ph. U. S.), which is obtained in long pods, is black, bright, and shining; sweet, slightly acid, and inodorous. It is laxative in the dose of ʒiv to ʒj.

CASSIA LANCEOLATA, *C. senna*—c. Lignea, *Laurus cassia*—c. Lignea Malabarica, *Laurus cassia*.

CASSIA MARILANDICA, *Senna Americana*, *American* or *Wild Senna*, *Locust plant*, (F.) *Séné d'Amérique*. The leaves of this plant are similar, in virtue, to those of *cassia senna*. They are, however, much inferior in strength.

CASSIA NIGRA, *C. fistula*—c. Officialis, *C. senna*—c. Orientalis, *C. senna*—c. Purging, *Cassia fistula*.

CASSIA SENNA, *C. lanceolata* seu *acutifolia* seu *orientalis* seu *officinalis*. The name of the plant which affords *senna*. It is yielded, however, by several species of the genus *cassia*. The leaves of *senna*, *Senna Folia*, *Senna Alexandrina* seu *Italica*, *Senna*, *Senna* or *Egyptian Cassia*, (F.) *Séné*, *Casse Séné*, have a faint smell, and bitterish taste. The active part, by some called *Cathartin*, is extracted by alcohol and water. Their activity is injured by boiling water. They are a hydragogue cathartic, and apt to gripe. Dose of the powder, ʒj to ʒj. Infusion is the best form.

The varieties of *senna*, in commerce, are *Tinnivelly Senna*, *Bombay* or *Common India Senna*, *Alexandrian Senna*, *Tripoli Senna*, and *Aleppo Senna*.

CASSIÆ ARAMENTUM, see *Cassia fistula*—c. *Fistula pulpa*, see *Cassia fistula*—c. Flores, see *Laurus cinnamomum*.

CASSIALA, *Hyssopus*.

CASSIDA GALERICULATA, *Scutellaria galericulata*.

CASSIDE BLEUE, *Scutellaria galericulata*.

CASSINA, *Ilex vomitoria*.

CASSINE CAROLINIANA, *Ilex paraguensis*—c. Evergreen, *Ilex vomitoria*—c. Peragua, *Ilex paraguensis*.

CASSIS, *Ribes nigrum*.

CASSITEROS, Tin.

CASSUMUNIAR, *Casamun'ar*, *Casmonar*, *Zerumbet*, *Casmina*, *Risajon*, *Ben'galé Indo-rum*, *Bengal Root*, (F.) *Racine de Bengale*. A root, obtained from the East Indies, in irregular slices of various forms; some cut transversely, others longitudinally. It is an aromatic bitter, and is consequently tonic and stimulant. It was once

down,' and γλῶσσα, 'the tongue.' An instrument for pressing down the tongue. See Glossocatochus.

CATAGRAPHOLOGIA, Pharmacocatagraphologia.

CATAIRE, Nepeta, see *Frémissement cataire*.

CATALEN'TIA, Epilepsy, or some disease resembling it — Paracelsus.

CATALEPSIA SPURIA, Ecstasis.

CATALEPSY, *Catalepsi'sia*, *Catalep'sis*, *Cat'ochē*, *Cat'ochus*, *Cat'ocha Gale'ni*, *Morbus atton'itus Celsi*, *Hystē'ria catalept'ica*, *Congela'tio*, *Deten'tio*, *Eucatalepsi'sis*, *Ap'honia* — (Hippocr.,) *Anau'dia* — (Antigenes,) *Apprehen'sia*, *Contempla'tio*, *Stupor vig'ilans*, *Prehen'sio*, *Carus Catalepsi'sia*, *Oppres'sio*, *Comprehensio* — (Cael. Aurelian,) *Comprehensio*, *Apoplec'sia Catalepsi'sia*, from *καταλαμβάνω*, 'I seize hold of.' *Trance* [?] (F.) *Catalepsi'e*. A disease in which there is sudden suspension of the action of the senses and of volition; the limbs and trunk preserving the different positions given to them. It is a rare affection, but is seen, at times, as a form of hysteria. Some of the Greek writers have used the word in its true acceptance of a seizure, surprise, &c.

CATALEPTIC, *Catalept'icus*, (F.) *Cataleptique*. Same etymon. Relating to catalepsy. Affected with catalepsy.

CATALEPTIC METHOD, *Meth'odus Catalept'ica*. The administration of external agents when internal agents are inapplicable.

CATALOT'IC, *Catalot'icus*, from *καταλωω*, 'to break or grind down.' A remedy which removes unseemly cicatrices.

CATAL'PA, *C. Arbo'rea* seu *Cordifo'lia* seu *Arbores'cens* seu *Bignoni'oides* seu *Syringifo'lia*, *Bigno'via Catal'pa*, *Cataw'ba tree*, *Bean tree*, *Indian Bean*. A decoction of the pods of the *Catalpa*, an American tree, of the Order *Bignoniaceæ*, *Didymia Angiospermia*, has been recommended in chronic nervous asthma.

CATAL'PA ARBOREA, *Catalpa* — c. *Bignonioides*, *Catalpa* — c. *Cordifolia*, *Catalpa* — c. *Syringefolia*, *Catalpa*.

CATALY'YSIS, Paralysis, from *kata*, and *λυω*, 'I dissolve or decompose.' The action of presence in producing decomposition; as when a body which possesses what has been termed *catalytic force* — (F.) *Force catalytique* — resolves other bodies into new compounds by mere contact or presence, without itself experiencing any modification.

CATALYT'IC, *Catalyt'icus*. Same etymon. A medicine, which is presumed to act by the destruction or counteraction of morbid agencies in the blood. — Headland. See *Catalysis*.

CATALYTIC FORCE, see *Catalysis*.

CATAMENIA, Menses — c. *Alba*, *Leucorrhœa*.

CATAMENIAL, *Catamenia'lis*, *Men'strual*, *Men'struous*, *Men'struous*, (F.) *Menstruel*, from *kata*, and *μην*, 'a month.' Appertaining or relating to the catamenia.

CATAMENIORUM FLUXUS IMMODICUS, Menorrhagia.

CATANANCE, *Cichorium intybus*.

CATANGELOS, Ruscus.

CANTLE'MA, *Catantle'sis*, from *kata*, 'upon,' and *αντλω*, 'I pour.' *Cateone'sis* and *Catæone'sis*. Ablution with warm water. A fomentation. — Mosehion, Marcellus Empiricus.

CATAPAS'MA, from *καταπασσω*, 'I sprinkle.' *Catapas'tum*, *Consep'er'sio*, *Epipas'ton*, *Pasma*, *Sympas'ma*, *Empas'ma*, *Diapas'ma*, *Xer'ion*, *Asper'sio*, *Epipas'tum*, *Pulvis asperso'rins*. A compound medicine, in the form of powder, employed by the ancients to sprinkle on ulcers, absorb perspiration, &c. — Paulus of Ægina.

CATAPH'ORA, 'a fall,' from *καταφερω*, 'I throw down.' A state resembling sleep, with

privation of feeling and voice. Somnolency. According to others, *Cataphora* is simply a profound sleep, which it is difficult to rouse from — in this sense being synonymous with *Sopor*.

CATAPHORA COMA, s. c. *Apoplexy* — c. *Hydrocephalica*, see *Apoplexy* — c. *Cymini*, *Thiaca Londinensis* — c. *Magnetica*, *Somnambulism*, *Magnetie*.

CATAPHRAC'TA, *Cataphrac'tes*, a *Chirasa*, from *καταφρασσω*, 'I fortify.' A name given by Galen to a bandage applied round the thorax and shoulders. It was also called *Quadr'yga*.

CATAPIESIS, Depression.

CATAPINOSIS, Absorption.

CATAP'LASIS, from *καταπασσω*, 'to besmear.' The act of besmearing or overlaying with plaster.

CAT'AP'PLASM, *Cataplas'ma*, *Epiplas'ma*, *Poultice*, *Pultis*, (Prov.) *Pulsey*, from *καταπασσιν*, (*kata* and *πασσιν*, 'to form or mould,') 'to besmear.' (F.) *Cataplasme*. A medicine applied externally, under the form of a thick pap. Cataplasms are formed of various ingredients, and for different objects. They may be *anodyne*, *emollient*, *tonic*, *antiseptic*, *irritating*, &c. A simple poultice acts only by virtue of its warmth and moisture. Mealy, fatty substances, leaves of plants, certain fruits, crumb of bread, &c., are the most common bases. The chief poultices which have been official are the following: — *Anodyne* — c. *Cicutæ*, c. *Digitalis*. *Antiseptic* — c. *Carbonis*, c. *Dauci*, c. *Fermenti*, c. *Acetosæ*, c. *Cumini*. *Emollient* — c. *Lini*, c. *Panis*, c. *Mali maturi*. *Irritating* — c. *Sinapis*, c. *Sodii chloridi*, c. *Quercus Marinzæ*. *Tonic* and *Astringent* — c. *Alum*, c. *Goulard*, c. of *Roses*.

The Parisian Codex had some other official cataplasms: — 1. *Cataplas'ma anod'gnum*, made of poppy and hyoscyamus. 2. *Cataplas'ma emolliente*, made of meal and pulps. 3. *Cataplas'ma ad suppuratio'nem promovent'um*, of pulps and basilicon. 4. *Cataplas'ma rubefaciens vel antipleurit'icum*, vinegar of pepper and vinegar.

The only cataplasms, the preparation of which it is important to describe, are some of the following: —

CATAPLASM, ALUM, *Coagulum Aluminosum* — c. of Beer grounds, see *Cataplasma Fermenti* — c. Carrot, *Cataplasma Dauci* — c. Charcoal, *Cataplasma carbonis ligni* — c. of Chlorinated Soda, *Cataplasma Sodæ chlorinatæ* — c. Flaxseed, *Cataplasma Lini* — c. Hemlock, *Cataplasma Conii* — c. Ice, see Ice — c. Linseed, *Cataplasma Lini* — c. Slippery Elm, *Cataplasma Ulmi* — c. Yeast, *Cataplasma Fermenti*.

CATAPLASMA, Cataplasma — c. Bynes, see C. Fermenti.

CATAPLAS'MA CARBO'NIS, *Charcoal Cataplasma* or *poultice*, (F.) *Cataplasme au charbon*. Made by adding powdered charcoal to a common cataplasma. Used as an antiseptic to foul ulcers, &c.

CATAPLAS'MA CON'I, *Hemlock Cataplasma*, made by spreading soft Extract of Hemlock on a flaxseed poultice. Used in malignant painful ulcers.

CATAPLAS'MA DAUCI, *Carrot Cataplasma* or *poultice*. Made by boiling the root of the Carrot until it is soft enough to form a poultice. Used in fetid ulcers.

CATAPLASMA FÆCULÆ CEREVISIÆ, see C. Fermenti.

CATAPLAS'MA FERMENT'I, *C. efferves'cens*, *Yeast Cataplasma* or *poultice*, (F.) *Cataplasme de Levure*. (Take of wheaten flour, ℥ij; yeast and water, aa, f℥v. Expose to a gentle heat.) It is antiseptic, and a good application to bruises. A *Cataplasma of Beer Grounds*, *Cataplasma Fæculæ Cerevisiæ*, C. Bynes, is used in the same cases.

CATAPLAS'MA LINI, *Linseed or Flaxseed Cataplasma*, (F.) *Cataplasme de farine de Lin*, *Cata-*

plasma common, made by mixing boiling water with *linseed meal*, is an excellent emollient.

CATAPLASMA SINA'PIS, *C. Sina'peos*, *Sin'apism*, Mustard Cataplasma, poultice or plaster, (F.) *Cataplasme de Moutard ou Sinapisme*. (Mustard and *linseed meal* or meal, $\bar{a}\bar{a}$, equal parts: warm vinegar or water, q. s.) A rubefacient and stimulant applied to the soles of the feet in coma, low typhus, &c., as well as to the pained part in rheumatism, &c.

CATAPLASMA SODE CHLORINAT'E, *Cataplasma of Chlorinated Soda*. Made by adding solution of chlorinated soda to a common *linseed cataplasma*. Applied to foul, offensive ulcers.

CATAPLASMA ULMI, *Slippy Elm poultice*, is made by adding boiling water to the powdered bark of *Ulmus fulca*, stirring constantly. It is a light, agreeable cataplasma.

CATAPLASME, *Cataplasma* — *c. au Charbon*, *Cataplasma carbonis ligni* — *c. de Farine de Lin*, *Cataplasma lini*.

CATAPLEX'IS, *Stupor*, from *κατα*, and *πλησσω*, 'I strike.' The act of striking with amazement. Appearance of astonishment, as exhibited by the eyes in particular. See *Hæmodia*.

CATAPOSIS, Deglutition.

CATAPOTIA, Pilula.

CATAPOTION, Pilula.

CATAPSYN'IS, from *καταψυχω*, 'I refrigerate.' *Peripne'ia*. Considerable coldness of the body, without rigor and horripilation. — Galen. *Perfric'tio*. Coldness in the extreme parts of the limbs. — Hippocrates.

CATAPTO'SIS, *Deciden'tia*, a fall. This word, at times, expresses the fall of a patient, attacked with epilepsy or apoplexy; at others, the sudden resolution of a paralytic limb.

CATAPULTA VIRILIS, Penis.

CATAPUTIA MINOR, *Euphorbia lathyris*, *Ricinus communis*.

CAT'ARACT, *Catarac'ta*, *Catarrhac'ta*, *Sufful'tio* Oculi, *S. Lentis crystallinæ*, *Phtharma catarac'ta*, *Calig'o lentis*, *Gutta opac'ea*, *Hypoc'hyma*, *Hypoc'hymis*, *Hypoph'ymis*, *Phacoscolo'ma*, *Parop'h'is catarac'ta*, *Glauc'o'ma Wauthou'si*, from *καταπασσειν*, (*κατα* and *πασσειν*), 'to tumble down.' A deprivation of sight, which comes on as if a veil fell before the eyes. Cataract consists in opacity of the crystalline lens or its capsule, which prevents the passage of the rays of light, and precludes vision. The causes are obscure. *Diagnosis*. — The patient is blind, the pupil seems closed by an opaque body, of variable colour, but commonly whitish: — the pupil contracting and dilating. Cataracts have been divided, by some, into *spurious* and *genuine*. The former, where the obstacle to vision is between the capsule of the lens and the uvea: the latter, where it is in the lens or capsule. A *lenticular cataract* is where the affection is seated in the lens: — a *capsular* or *membranous*, in the capsule. The *capsular* is divided again, by Beer, into the *anterior*, *posterior*, and *complete capsular* cataract. When the capsule is rendered opaque, in consequence of an injury, which cuts or ruptures any part of it, it thickens, becomes leathery, and has been called *Catarac'ta akida siliq'uosa*. *Catarac'ta Morgagniana lactea seu puriformis* is the milky variety, in which the crystalline is transformed into a liquid similar to milk, (F.) *Cataracte lacteuse*; or, as generally defined, in which there is opacity of the fluid situate between the lens and its capsule. The *cap'sulo-lenticular* affects both lens and capsule, (*Achry'stallodiaphanie*, *Adiaphanie p'rochry'stallique* — Piory), and Beer conceives the liquor Morgagni, in an altered state, may contribute to it. Cataracts are also called *hard*, *soft* (*Phacoma'licia*,), *stony*, (F. *pierruse*.) *milky* or *cheery* (*lactese* or *ca'uese*), *Galactocata-*

rac'ta, *Catarac'ta lactic'olor*, according to their density: — white, pearly, yellow, brown, gray, green, black, (F.) *blanche*, *perlée*, *jaune*, *brune*, *grise*, *verte*, *noire*, according to their colour: — fixed or oscillating — *catarac'ta capsulo-lenticularis fixa vel tremula*, (F.) *fixe* on *branlante*, according as they are fixed or movable behind the pupil. They are likewise called *Catarac'ta marmor'icia*, *fene'strac'ta*, *stellac'ta*, *punctac'ta*, *dimidiac'ta*, &c., according to the appearances they present.

They may be *simple*, or *complicated* with adhesion, amaurosis, specks, &c.; and *primary* or *primitive*, when opaque before the operation; — *secondary*, when the opacity is the result of the operation.

The following classification of cataracts is by M. Desmarres:—

CLASS I.—True Cataracts.

		Green.
		Black.
	Hard.	Osseous.
		Stony or chalky.
		Striated-etiolated,
		barred, debiscent, with three branches, &c.
	Soft.	Disseminated, or dotted.
a. Lenticular Cataracts.		Congenital.
		Traumatic.
		Glaucomatous.
		Morgagnian, or interstitial.
	Liquid.	Cystic, purulent, fetid.
	Other varieties, soft, hard, or liquid.	Shaking, or floating.
		Luxated.
b. Capsular Cataracts.	Anterior.	Pyramidal or vegetant.
	Posterior.	Arid siliquose.
c. Capsulo-lenticular Cataracts.	All the varieties of lenticular and capsular Cataracts.	
d. Secondary Cataracts.	Lenticular.	
	Capsular.	
	Capsulo-lenticular.	

CLASS II.—False Cataracts.

Fibrinous.
Purulent.
Sanguineous.
Pigmentous.

Cataract is commonly a disease of elderly individuals, although, not unfrequently, *congenital*. It forms slowly; objects are at first seen as through a mist; light bodies appear to fly before the eyes, and it is not until after months or years that the sight is wholly lost. No means will obviate the evil except an operation, which consists in removing the obstacle to the passage of the light to the retina. Four chief methods are employed for this purpose. 1. *Couching* or *Depression*, *Hyaloniz'is*, *Hyalonyx'is*, (F.) *Abaissement*, *Déplacement de la Cataracte*. This consists in passing a cataract needle through the sclerotics and subjacent membranes, a little above the transverse diameter of the eye; and at about two lines' distance from the circumference of the transparent cornea, until the point arrives in the posterior chamber of the eye. With this the crystalline is depressed to the outer and lower part of the globe of the eye, where it is left. 2. *By absorption*, — by the French termed *bruitement* or *bruising*. This is performed in the same manner as the former; except that, instead of

turning the crystalline from the axis of the visual rays, it is divided by the cutting edge of the needle, and its fragments are scattered in the humours of the eye, where they are absorbed. 3. *By extraction*, which consists in opening, with a particular kind of knife, the transparent cornea and the anterior portion of the capsule of the crystalline; and causing the lens to issue through the aperture. Each of the processes has its advantages and disadvantages, and all are used by surgeons. 4. Some, again, pass a cataract needle through the transparent cornea and pupil to the crystalline, and depress or cause its absorption. This is called *Ceratonyxis*. See, also, *Reclination*.

CATARACT, BLACK, Amaurosis — c. Capsular, see Cataract — c. Capsulo-lenticular, see Cataract — c. Central, Centradiaphanes — c. Cheesy, see Cataract — c. Congenital, see Cataract — c. Complicated, see Cataract — c. Fixed, see Cataract — c. Genuine, see Cataract — c. Hard, see Cataract — c. Lenticular, see Cataract — c. Membranous, see Cataract — c. Milky, see Cataract — c. Opaque, see Cataract — c. Primary, see Cataract — c. Primitive, see Cataract — c. Secondary, see Cataract — c. Simple, see Cataract — c. Soft, see Cataract — c. Spurious, see Cataract — c. Stony, see Cataract — c. Vacillating, see Cataract.

CATARACTA, Cataract — c. Arida siliquosa, see Cataract — c. Capsulo-lenticularis, see Cataract — c. Centralis, Centradiaphanes — c. Dimidiata, see Cataract — c. Fenestrata, see Cataract — c. Glauca, Glaucoma — c. Lacticolor, see Cataract — c. Liquida, Hygrocataracta — c. Marmoracea, see Cataract — c. Morgagniana, see Cataract — c. Nigra, Amaurosis — c. Punctata, see Cataract — c. Stellata, see Cataract.

CATARACTE, ABASSEMENT DE LA, see Cataract — c. Blanche, see Cataract — c. Brûlante, see Cataract — c. Brune, see Cataract — c. Caséuse, see Cataract — c. Déplacement de la, see Cataract — c. Fixe, see Cataract — c. Grise, see Cataract — c. Jaune, see Cataract — c. Laiteuse, see Cataract — c. Noire, Amaurosis, see Cataract — c. Perlée, see Cataract — c. Pierreuse, see Cataract — c. Verte, see Cataract.

CATARACTÉ (F.), *Cataractus*, *Cataractâritus*. One affected with cataract. The French use this term, both for the eye affected with cataract and the patient himself.

CATARIA, see Nepeta — c. Vulgaris, Nepeta. CATARRH', *Catar'rh*, *Catar'rhopus*, *Catar'rhœma*, *Rheum*, *Deflux'io*, *Catarr'hus*, *Phlegmat'rhay'ia*, *Phlegmat'orrhœa*, from *kata*, 'downwards,' and *rho*, 'I flow.' A discharge of fluid from a mucous membrane. The ancients considered catarrh as a simple flux, and not as an inflammation. Generally it partakes of this character, however. *Catarrh* is, with us, usually restricted to inflammation of the mucous membrane of the air-passages: the French extend it to that of all mucous membranes; (F.) *Flux muqueus*, *Fluxion catarrhale*.

Catarrh, in the English sense, *Broncho catarrhus*, *Pulmonary Catarrh*, *Ling fever* (vulgarly), *Rheuma Pectoris*, *Destillatio Pectoris*, *Catar'rh*, *Pec'toris*, *C. Pulmonum*, *C. Pulmonalis*, *C. Bronchialis*, *Blennop'tysis*, *Tussis catarrhalis simplex*, *Gravello* (of many), *Febris Catarrhalis*, *Blennorrhœa*, *Bronchitis*, *Catar'rh*, *à Frigoré*, a Cold, (Prov.) *Cooth*, (F.) *Catarrhe pulmonaire*, *Fièvre Catarrhale*, *Rhume de Poitrine*, is a superficial inflammation of the mucous follicles of the trachea and bronchi. It is commonly an affection of but little consequence, but apt to relapse and become chronic. It is characterized by cough, thirst, lassitude, fever, watery eyes, with increased secretion of mucus from the air-passages. The antiphlogistic regimen and time usually remove it. Sometimes, the inflam-

mation of the bronchial tubes is so great as to prove fatal.

CATARRH, ACUTE, OF THE UTERUS, see *Metritis* — c. Chronic, *Bronchitis*, (chronic) — c. Dry, see *Bronchitis* — c. Pulmonary, *Bronchitis*, *Catarrh* — c. Rose, Fever, hay — c. Suffocating nervous, *Asthma Thymicum* — c. Summer, Fever, hay.

CATARRH', EPIDEM'IC, *Catar'rh*, *epidem'icus*, *C. à conta'gio*, *Rheuma epidem'icum*. Catarrh prevailing owing to some particular *Constitutio aëris*, and affecting a whole country, — *Influenza*.

CATARRHACTA, Cataract.

CATARRHAL, *Catar'rh*, *alis*, *Catarrho'icus*, *Catarrho'itis*, *Catarrho'iticus*. Relating to catarrh, — as *Catarrhal* Fever.

CATARRHE AIGUE DE L'UTÉRUS, see *Metritis* — c. Buccal, *Aphthæ* — c. Conclusive, *Bronchitis* — c. Gastric, *Gastritis* — c. Gutta'al, *Cynanche tonsillaris* — c. Intestinal, *Diarrhœa* — c. Laryngien, *Laryngitis* — c. Nasal, *Coryza* — c. Oculaire, *Ophthalmia*, *c. de l'Oreille*, *Otitis* — c. Pharyngien, *Cynanche parotidea* — c. Pituiteux, *Bronchorrhœa* — c. Pulmonaire, *Catarrh* — c. Sec; see *Bronchitis* — c. Stomacal, *Gastro'rhœa* — c. Uterin, *Leucorrhœa* — c. Vesical, *Cystorrhœa*.

CATARRHEC'TICA, from *katarrhynvvti*, 'I break down.' Remedies considered proper for evacuating; — as diuretics, cathartics, &c. — *Hippocrates*.

CATARRHEUMA, Catarrh.

CATARRHEUX (F.) *Catarrho'sus*. One subject to catarrh; affected with catarrh.

CATARRHÆA IA, *Catarrhœ'is*; same etymon as *Catarrh etica*. The action of *Catarrhætica*. Also, effusion; evacuation of the bowels.

CATARRHEXIS, *Catarrhexia*, *Excrement* — c. Vera, *Hamatochezia*.

CATARRHŒA, Rheumatism.

CATARRHOËT'ICUS, from *katarrho*, 'I flow from.' An epithet for disease produced by a discharge of phlegm; catarrhal.

CATARRHOPA PHY'MATA, from *katarrhōpos*, *katarrhōpos*, 'sloping downwards.' Tubercles tending downwards, or with their apices downwards.

CATARRHOPHE, Absorption.

CATARRHOPHESIS, Absorption.

CATARRHOP'IA, *Catar'rhysis*, from *kata*, 'downwards,' and *pos*, 'inclination.' An afflux of fluids towards the inferior parts, and especially towards the viscera of the abdomen. The Greek word *avaporia* expresses an opposite phenomenon, or a tendency towards the upper parts.

CATARRHOPUS, Catarrh.

CATARRHOS'CHESIS, from *katarrhos*, 'catarrh,' and *oxycsis*, 'suppression.' The suppression of a mucous discharge.

CATARRHIUS, *Defluxion*, *Tussis* — c. *Æstivus*, fever, hay — c. *Bellinulanus*, *Cynanche parotidea* — c. *Bronchialis*, *Catarrh* — c. *Bronchiorum*, *Bronchitis* — c. *à Contagio*, *Influenza* — c. *Epidemicus*, *Influenza*, *Catarrh*, *epidemic* — c. *Genitalium*, *Leucorrhœa* — c. *Gonorrhœa*, *Gonorrhœa* — c. *Intestinalis*, *Diarrhœa* — c. *Laryngeus*, *Laryngo-catarrhus* — c. *ad Nares*, *Coryza* — c. *Nasalis*, *Coryza* — c. *Pulmonalis*, *Catarrh* — c. *Pulmonum*, *Bronchitis*, *Catarrh* — c. *Senilis*, *Bronchitis*, (chronic) — c. *Suffocativus* *Barbadensis*, *C. trachealis* — c. *Trachealis*, *Laryngo-catarrhus* — c. *Urethrae*, *Gonorrhœa pura* — c. *Urethralis*, *Gonorrhœa* — c. *Vesicæ*, *Cystorrhœa*.

CATARRHYSIS, *Catarrhopia*, *Defluxion*.

CATARTISIS, *Catartismus*.

CATARTIS'MUS, *Catar'tis*, from *katartisein*, 'to repair, replace.' The coaptation of a luxated or fractured bone, or hernia.

CATASARCA, Anasarca.
 CATASCEUE, Structure.
 CATASCHASMUS, Bloodletting, Scarification.
 CATASTAGMUS, Catarrh, Coryza.
 CATASTALAGMUS, Coryza, Distillation.
 CATASTALTICA, Hæmatostatica, Sedatives, Styptics.

CATAS'TASIS, from *καθίστημι*, 'I establish.' The constitution, state, condition, &c., of any thing. — Hippocrates. Also, the reduction of a bone. See Constitution, and Habit of Body.

CATAT'ASIS, from *κατατείνω*, 'I extend.' Extension. The extension and reduction of a fractured limb. — Hippocrates.

CATATHLIPSIS, Oppression.

CATAWBA TREE, Catalpa.

CATAxis, Fracture.

CATCH FLY, Apocynum androsæmifolium, Silene Virginica.

CATCHING, see Contagious.

CATCHUP, Ketchup.

CAT'CHU. The extract of the wood of *Aca'cia* seu *Mimo'sa Cat'chu*, *Cuat'chu*, and of the leaves of *Nau'clea* seu *Uncaria gambir*, oriental trees. The drug is also called *Extrac'tum Cat'chu*, *Terra Japonica*, *Japan Earth*, *Caschen*, *Cat'chu*, *Cashou*, *Cat'chu*, *Castjoe*, *Cacan*, *Cate*, *Kaath*, *Cuti*, *Cuteh*, *Caira*, *Succus Japonicus*, *Gambir*, (F.) *Cachou*. It is a powerful astringent, and is used in diarrhoea, intestinal hemorrhage, &c. Dose, gr. xv to ʒss, in powder.

CATECHU, SQUARE, see Nauclea gambir.

CATEIAD'ION, from *καρα*, and *εἰα*, 'a blade of grass.' A long instrument thrust into the nostrils to excite hemorrhage in headach. — Aretæus.

CATENÆ MUSCULUS, Tibialis anticus.

CATEONESIS, Catantlema.

CATGUT, Galega Virginiana.

CATH'ERESIS, *καθάρσις*, 'subtraction, diminution.' Extenuation or exhaustion, owing to forced exercise. — Hippocrates. The action of cathartics.

CATH'ERETICUS, Cathartic.

CATHARETICUS, Cathartic.

CATHARISMOS, Depuration.

CATHARMA, *Purgamentum*. The matter evacuated by a purgative, or by spontaneous purging; also, a cathartic.

CATHAR'MUS. Same etymon; a purgation. — Hippocrates. Also, the cure of a disease by magic, &c.

CATHAR'SIS, from *καθαίρειν*, (*καθ'* and *αιρειν*, 'to take away,') 'to purge.' *Purga'tio*, *Apocathar'sis*, *Copropho'ria*, *Coprophore'sis*. A natural or artificial purgation of any passage; — mouth, anus, vagina, &c.

CATHAR'TIC, *Cathar'ticus*, *Cathare'ticus*, *Cathar'ma*, *Coprocril'ticum*, *Coprago'gum*, *Lustramentum*, *Purgans medicamentum*, *Trichilium*, *Dejecto'rium Remed'ium*, *Eccathar'ticus*, *Hypocriticus*, *Hypochoret'icus*, *Alvum evac'ans*, *Hypel'atos*, *Lapac'ticus*, *Apocathar'ticus*. Same etymon. (F.) *Cathartique*. A medicine which, when taken internally, increases the number of alvine evacuations. Some substances act upon the upper part of the intestinal canal, *calomel* and *colocynth*; others, on the lower part, as *aloes*; and some on the whole extent, as *saline* purgatives. Hence a choice may be necessary. Cathartics are divided into purgatives and laxatives. The following is a list of the chief cathartics:

Aloe, Cassia Marilandica, Colocynthis, Elaterium, Gambogia, Hydrargyri Chloridum mite, Hydrargyri Oxidum nigrum, Hydrarg. cum Magnesîa, Jalapa, Juglans, Magnesia, Magnesiæ Carbonas, Magnesiæ Sulphas, Manna, Mannita, Oleum Euphorbiæ, Lathyridis, Oleum Ricini, Oleum Tiglli, Podophyllum, Potassæ Acetas, Potassæ Bisulphas, Potassæ Sulphas, Potassæ Bi-

tartras, Potassæ Tartras, Rheum, Scammonium, Senna, Sinapis, Soda et Potassæ Tartras, Soda Phosphas, Soda Sulphas, Sodii Chloridum, Sulphur, Veratrica, Aquæ Minerales Sulphureæ et Salinae, Emomata, Suppositoria.

CATHARTIN, see Cassia Senna, and Convolvulus jalapa.

CATHARTIQUE, Cathartic.

CATHARTOCARPUS, Cassia fistula.

CATHEDRA, Anus.

CATHEMERINUS, Quotidian.

CATHEMERUS, Quotidian.

CATHERET'IC, *Catharet'icus*, *Ectylo'ticus*, *Sarcoph'agus*, from *καταίρειν*, 'to eat,' 'destroy.' Substances applied to warts, exuberant granulations, &c., to eat them down. *Mild caustics*.

CATH'ETER, from *καθέναι* (*καθ'* and *αινειν*, 'to send,') 'I explore.' *Æne'a*, *Alg'alie*, *Alyalie*, *Cathete'ria*, *Demissor*, *Inmiss'or*. A hollow tube, introduced by surgeons into the urinary bladder, for the purpose of drawing off the urine. Catheters are made of silver or elastic gum. See Bougie. The French generally use the word *catheter* for the solid *sonde* or *staff*; and *alg'alie* and *sonde* for the hollow instrument.

CATHETER, NASAL. An instrument, invented by M. Gensoul, of Lyons, for catheterizing the ductus ad nasum. It is hook-shaped; the extremity, bent at a right angle, is about an inch in length, suited to the distance of the lower orifice of the duct from the nostril, and likewise to the length and form of the duct, with a slight spiral turn.

CATHETERIS, Catheter.

CATHETERISIS, Catheterismus.

CATHETERIS'MUS, *Cathete'ria*, *Catheteri'satio*, *Catheterism*, *Catheterizatio*, *Inmissio Cathete'ria*, same etymon. The introduction of a catheter or sound into a natural passage, as through the urethra into the bladder or into the Eustachian tube, larynx, &c. Also, probing a wound. Melosis.

CATHETERIZATION, Catheterismus.

CATHETERIZE. To perform the operation of catheterism, — in other words, to introduce the catheter: to probe or sound a cavity.

CATHID'YSIS, from *καθίστημι*, 'I place together.' Reduction of a part to its natural situation.

CATHMIA, Plumbi oxydum semi-vitreum.

CATHMIR, Calamina.

CATH'ODIC, *Cathod'icus*; from *καθ'*, 'downwards,' and *ὁδός*, 'a way.' An epithet applied by Dr. Marshall Hall to a downward course of nervous action.

CATH'OLIC HUMOURS. (F.) *Humours Catholiques*, are the fluids spread over the whole body.

CATHOLICON, Panacea.

CATHOLICON DUPLEX. An ancient purging electuary, chiefly composed of cassia, tamarinds, rhubarb, senna, &c.

CATIL'LIA. A weight of nine ounces.

CATILLUS, Cup.

CATINUS FUSORIUS, Crucible.

CATLING, Knife, double-edged.

CATNEP, Nepeta Cataria.

CATO, *κατω*, 'below,' 'beneath.' This word, in the writings of Hippocrates, is often used for the abdomen, especially the intestines. When he advises a remedy *κατω*, he means a purgative; when *ανω*, 'above or upwards,' an emetic. As a prefix, *Cato* means 'beneath,' as in

CATOCATHARTIC, *Catocathar'ticus*, from *κατω*, 'downwards,' and *καθαίρειν*, 'I purge.' A medicine which purges downwards. One that produces alvine evacuations. The antithesis to *Anacathartic*.

CATOCCHA GALENI, Catalepsy.

CAT'OCHEË, *Cat'ochæis*, *Cat'ochus*, from *κατ'οχεύω*,

'I retain,' 'I hold fast.' This word has, by some, been used synonymously with Catalepsy; by others, with Coma vigil; by others, with Tetanus.

CATOCHUS. Catache, Ecstasis—c. Cervinus, Tetanus—c. Holotonicus, Tetanus—c. Infantum, Induration of the cellular tissue.

CATOMIS'MOS, from *κατω*, 'beneath,' and *σπας*, 'shoulder,' *Subhumeral'io*. A mode with the ancients of reducing luxation of the humerus by raising the body by the arm.—Paulus of Ægina.

CATOPTER, Speculum.

CATOP'TRIC, *Catop'tricus*, (F.) *Catoptrique*, from *κατοπτρις* or *κατοπτρον*, 'a mirror.' Relating or appertaining to Catoptrics or the reflection of light.

CATOP'TRIC EXAMINATION OF THE EYE. When a lighted candle is held before the eye, the pupil of which has been dilated by belladonna, three images of it are seen—two erect, and one inverted:—the former owing to reflection from the cornea and anterior surface of the crystalline; the latter owing to reflection from the posterior layer of the crystalline inverted, and between the other two. This mode of examining the eye has been proposed as a means of diagnosis between cataract and amaurosis. In the latter, all the images are seen.

CATOPTROMANCY, from *κατοπτρον*, (*κατα*, and *οπτοιαι*), 'a mirror,' and *μαντιαι*, 'divination.' A kind of divination by means of a mirror.

CATOPTRON, Speculum.

CATORCHITES. A kind of sour wine, prepared with the orchis and black grape, or dried figs. It was formerly employed as a diuretic and emmenagogue.—Dioscorides. Called, also, *Sycites*.—Galen.

CATORETICUS, Purgative.

CATOTERICUS, Purgative.

CATOTICA, from *κατω*, 'beneath.' Diseases infecting internal surfaces. Pravity of the fluids or emunctories, that open on the internal surfaces of organs. The second order in the class *Ecetica* of Good.

CATOX'YS, *Peracutus*, from *kata*, 'an intensive,' and *αξυς*, 'acute.' Highly acute: as *Morbus Catoxys*, *M. Peracutus*, a very acute disease.

CATS' EYE. A name, under which several morbid conditions of the eye would appear to have been confounded—all agreeing, however, in presenting an opalescent appearance of the pupil or of the bottom of the eye; those parts reflecting the light in various colours, or, at least, with various degrees of intensity, according to the direction in which the eye is turned. This appearance was compared by Beer to the reflection from the tapetum of the eye of the cat.—MacKenzie.

CAT'S EYE, AMAUROTIC, see Amaurotic.

CAT'S FOOT, *Antennaria dioica*.

CAT'S PURR, *Frémissement Cataire*.

CATTAGAUMA, Cambogia.

CATTITEROS, Tin.

CATULOTICA, *Cicatrisantia*.

CATURUS SPICIFLORUS, (*katra*, *Catus*, 'a cat,' and *οψα*, 'tail,') *Cauda felis*, from the shape of its flowers. *Acalypha hispida*. An East Indian shrub, Order, Euphorbiaceæ: used in its native country, in decoction, in diarrhoea and dysentery.

CATU-TRIPALI, *Piper longum*.

CAUCALIS CAROTA, *Daucus carota*—c. *Sanicula*, *Sanicula*.

CAUCALOIDES, *Patella*.

CAUCASIAN, see Homo.

CAUCHEMAR, Incubus.

CAUCHEVIELLE, Incubus.

CAUCHUC, Caoutchouc.

CAUDA, see Clitorism, Coccyx, Penis.

CAUDA EQU'NA. The spinal marrow, at its termination, about the second lumbar vertebra, gives off a considerable number of nerves, which, when unravelled, resemble a horse's tail; hence the name; (F.) *Queue de Cheval*, *Q. de la Moëlle Épinère*. See *Medulla Spinalis*.

CAUDA FELIS, *Caturus*—c. f. *Agrestis*, *Acalypha betulina*.

CAUDA SALAX, Penis.

CAUDAL, *Caudate*, *Caudalis*, *Caudatus*; from *cauda*, 'a tail.' Relating or appertaining to a tail. Having a tail or tail-like appendage:—as '*caudal* or *caudate* cells or corpuscles'—cells or corpuscles having a tail-like appendage, as in cancerous growths.

CAUDATE, Caudal.

CAUDATIO, Clitorism.

CAUDATUS, Bicusidatus.

CAUDEX CEREBRI, Peduncles of the brain.

CAUDIEZ, MINERAL WATERS OF. Caudiez is a small town, nine leagues from Perpignan, in France, where there is a thermal spring, containing a little sulphate of soda and iron.

CAUDLE, *Cowdel*, (F.) *Chaudéau*, from *choud*, 'warm or hot.' A nourishing gruel given to women during the childbed state. The following is a form for it: Into a pint of fine gruel, not thick, put, whilst it is boiling hot, the yolk of an egg beaten with sugar, and mixed with a large spoonful of cold water, a glass of wine, and nutmeg. Mix the whole well together. Brandy is sometimes substituted for the wine, and lemon peel or capillaire added. It is also sometimes made of gruel and beer, with sugar and nutmeg.

CAUL, from (L.) *caula*, 'a fold,' *Pilus*, *Pileolus*, *Gal'lea*, *Vitta*, (F.) *Coeffe*, *Coiffe*—(*Être né coiffé*—'to be born with a caul.') The English name for the omentum. When a child is born with the membranes over the face, it is said to have been 'born with a caul.' In the catalogue of superstitions, this is one of the favourable omens. The caul itself is supposed to confer privileges upon the possessor; hence the membranes are dried, and sometimes sold for a high price. See Epilploon.

CAULE'DON, *Ciccy'don*, from *καυλος*, 'a stalk.' A transverse fracture.

CAU'LIFLOWER, (G.) Kohl, 'cabbage,' and *flower* (?), *Brassica Florida*.

CAULIFLOWER EXCRESCENCE, (F.) *Chonfleur*. An excrescence, which appears about the origin of the mucous membranes, chiefly about the anus and vulva, and which resembles, in appearance, the head of the cauliflower. It is often syphilitic in its character.

CAULIFLOWER EXCRESCENCE OF THE UTERUS, *Metrobot'rytes*. A morbid growth from a part, or the whole, of the circumference of the os uteri; and, at times, from the surface of the cavity of the uterus. It is doubtful whether its progress can be arrested except by excision.

CAULIS, Penis—c. Florida, *Brassica Florida*.

CAULOPHYLLUM THALICTROIDES, (*καυλος*, 'a stalk or stem,' and *φυλλον*, 'a leaf,' so called because the leaf-stalks appear like a continuation of the stem.) *Leon'tice thalictroides*, *Blueberry Cohosh*, *Cohosh*, *Cohush*, *Blue Cohosh*, *Blueberry*, *Papoose Root*, *Squaw Root*, *Blue Ginseng*, *Yellow Ginseng*, a plant of the Family *Berberidæ*; *Sex. Syst.* Hexandria Monogynia, which grows all over the United States, flowering in May and June. The infusion of the root is much used by the Indians in various diseases. To it are ascribed emmenagogue and diaphoretic virtues.

CAULOPIE'GIA, from *καυλος*, 'the male organ,' and *εληνη*, 'a wound,' or 'stroke.' An injury or paralysis of the male organ.

CAULORRHAGIA, Stomatosis—c. Ejaculato-

ria, Spermato-cystidorrhagia.—c. Stillatitia, Urethrorrhagia.

CAULORRHŒA BENIGNA, Gonorrhœa pura.

CAULUS, Penis.

CAUMA, *καυμα*, 'a burnt part,' from *καω*, 'I burn.' Great heat of the body or atmosphere. Synocha, Empresma.

CAUMA BRONCHITIS, Cynanche trachealis.—c. Carditis, Carditis.—c. Enteritis, Enteritis.—c. Gastritis, Gastritis.—c. Hemorrhagicum, Hemorrhagia activa.—c. Hepatitis, Hepatitis.—c. Ophthalmitis, Ophthalmia.—c. Peritonitis, Peritonitis.—c. Phrenitis, Phrenitis.—c. Pleuritis, Pleuritis.—c. Podagricum, Gout.—c. Rheumatismus, Rheumatism, acute.

CAUMATO'DES, *Caumato'rus*, from *καυμα*, 'fire, heat.' Burning hot. *Febris caumato'des*, *F. causo'des*. Inflammatory fever. Synocha.

CAUNGA, Areca.

CAUSA CONJUNCTA, Cause, proximate.—c. Continens, Cause, proximate.

CAUSÆ ABDITÆ, Causes, predisponent or remote.—c. Actuales, Causes, occasional.—c. Præincipientes, Causes, procatactic.—c. Prœgumene, Causes, predisponent.

CAUSE, *Causa*, *Aitia*, *Aition*. An act which precedes another, and seems to be a necessary condition for the concurrence of the latter. The causes of disease are generally extremely obscure; although they, sometimes, are evident enough. The *predisponent* and *occasional* causes are the only two on which any stress can be laid; but as authors have divided them differently, a short explanation is necessary.

CAUSE, AC'CESSORY, (F.) *Cause Accessoire*. One which has only a secondary influence in the production of disease.

CAUSES, ACCIDENT'AL, *Common Causes*, (F.) *Causes Accidentelles*, are those which act only in certain given conditions; and which do not always produce the same disease. Cold, e. g., may be the accidental cause of pneumonia, rheumatism, &c.

CAUSES CACHÉES, C. occult.—c. Common, C. accidental.—c. Exciting, C. Occasional.—c. Essential, C. Specific.—c. *Déterminantes*, C. Specific.—c. *Éloignées*, C. Predisponent.

CAUSES, EXTERNAL, (F.) *Causes externes*, are such as act externally to the individual; as air, cold, &c.

CAUSES FORMELLES (F.), are such as determine the form or kind of disease. They differ from the *Causes matérielles*, which are common to a set of diseases; as, to the neuroses, phlegmasiæ, &c.

CAUSES, HIDDEN, C. Occult.—c. Immediate, C. essential.

CAUSES, INTERN'AL, (F.) *Causes Internes*, are those which arise within the body; as mental emotions, &c.

CAUSES, MECHAN'ICAL, (F.) *Causes mécaniques*, are those which act mechanically, as pressure upon the windpipe in inducing suffocation.

CAUSES, NEGATIVE, (F.) *Causes négatives*, comprise all those things, the privation of which may derange the functions;—as abstinence too long continued. They are opposed to *positive causes*, which, of themselves, directly induce disease;—as the use of indigestible food, spirituous drinks, &c.

CAUSES, OBSCURE, C. Occult.

CAUSES, OCCASIONAL, *Exciting Causes*, *Causes actua'les*, (F.) *Causes occasionelles*, are those which immediately produce disease. The occasional causes have been divided into the *cognizable* and *non-cognizable*.—C. J. B. Williams.

I. Cognizable Agents.

1. Mechanical.
2. Chemical.
3. Ingesta.
4. Bodily exertion.
5. Mental emotion.
6. Excessive evacuation.
7. Suppressed or defective evacuation.
8. Defective cleanliness, ventilation, and draining.
9. Temperature and changes.

EXCITING
CAUSES
OF
DISEASE.

II. Non-Cognizable Agents.

1. Endemic.
2. Epidemic.
3. Infectious.

} Poisons.

CAUSES, OCCULT, *Hidden causes*, *Obscure causes*, (F.) *Causes occultes ou cachées ou obscures*. Any causes with which we are unacquainted; also, certain inappreciable characters of the atmosphere, which give rise to epidemics.

CAUSES, PHYSICAL, (F.) *Causes Physiques*,—those which act by virtue of their physical properties; as form, hardness, &c. All vulnerating bodies belong to this class.

CAUSES, PHYSIOLOGICAL, (F.) *Causes Physiologiques*, those which act only on living matter;—narcotics, for example.

CAUSES, PREDISPO'NENT, C. *predispos'ing*, *Remote causes*, *Causa prægumena*, *Causa abditæ*, *Causa remotæ*; (F.) *Causes prédisposantes*, *Causes éloignées*,—those which render the body liable to disease. They may be *general*, affecting a number of people, or *particular*, affecting only one person.

CAUSES, PRIN'CIPAL, (F.) *Causes principales*,—those which exert the chief influence on the production of disease, as distinguished from the *accessory causes*.

CAUSES, PROCATARC'TIC, *Causa procatartica*, *Causa præincipien'tes*, from *προκαταρκτος*, 'the origin or beginning of a thing,' (*προ*, 'before,' and *καταρχω*, 'I begin.') These words have been used with different significations. Some have employed them synonymously with *predisponent* or *remote causes*; others with *occasional* or *exciting causes*.

CAUSE PROCHAINE, C. proximate.

CAUSE, PROXIMATE, *Causa proxima* seu *continens seu conjuncta*, (F.) *Cause continente ou prochaine*, may be the disease itself. Superabundance of blood, e. g., is the proximate cause of plethora.

CAUSES, Remote, C. predisponent.

CAUSES, SPECIFIC, *Essential* or *Immediata causes*, &c., (F.) *Causes spécifiques*, *C. essentielles*, *C. déterminantes*;—those which produce a determinate disease.

CAUSIS, Burn, Ebullition, Fermentation, Incendium, Ustion.

CAUSOMA, Inflammation.

CAUSTIC, *Causticus*, *Canther'icus*, *Dieret'icus*, *Ero'dens*, *Adu'rens*, *Urens*, *Pyrot'icus*, from *καω*, 'I burn,' *καυσος*, 'heat.' (F.) *Caustique*. Bodies which have the property of causticity; and which, consequently, burn or disorganize animal substances. The word is also used substantively. The most active are called *Eucharot'ics*. Caustics are also termed 'corrosives.'

CAUSTIC BEARER, *Porte-pierre*—c. *Paste*, *Canquoin's*, *Pasta Zincæ chloridi*.

CAUSTICA ADUSTO, Cauterization.

CAUSTICITY, *Caustic'itas*, from *καυσω*, 'that which burns,' (*καω*, 'I burn.') The impression which caustic bodies make on the organ of taste; or, more commonly, the property which distinguishes those bodies.

CAUSTICOPHORUM, *Porte-pierre.*

CAUSTICUM ÆTHIOPICUM, Unguentum Melancholicum causticum. A sort of paste, made by rubbing powdered saffron with concentrated sulphuric acid, recommended by Velpéau as a caustic in cases of gangrenous and carcinomatous ulcers. The acid is the caustic: the saffron, the constituent merely.

CAUSTICUM ALKALINUM, Potassa fusa—c. Americainum, Veratrum sabadilla—c. Antimoniale, Antimonium muriatum.

CAUSTICUM COMMUNĒ, Potent'ial Caustery, Common Caustic, Cauterium potentia'le, Lapis septicus, Crusticium communĒ mil'ius. This consists of quicklime and black soap, of each equal parts.

CAUSTICUM COMMUNE, Potassa fusa—c. Commune acerrimum, Potassa fusa—c. Commune fortius, Potassa cum calce—c. Lunare, Argenti nitras—c. Potentiale, Potassa fusa—c. Salinum, Potassa fusa—c. Viennense fusum Filhos, see Powder, Vienna.

CAUSTIQUE, Caustic—c. Filhos, see Powder, Vienna—c. de Viennæ, Powder, Vienna.

CAUSUS, from καω, 'I burn,' καυσος, 'heat.' A highly ardent fever: *Deut'reus*. Pincel regards it as a complication of bilious and inflammatory fever; Broussais, as an intense gastritis, accompanied with bilious symptoms. See Synocha.

CAUSUS, ENDEMIAL, OF THE WEST INDIES, Fever, Yellow—c. *Tropicus endemicus*, Fever, Yellow.

CAUTER, Cauterium.

CAUTÈRE, Cauterium, Fonticulus—c. *Inhérent*, Inherent cautery.

CAUTERETICUS, Caustic.

CAUTERETS, MINERAL WATERS OF. Cauterets is a *bourg* seven leagues from Barèges (*Hautes-Pyrénées*), France. The waters are hydrosulphurous and thermal—temperature 123° F. They are used in the same cases as the Barèges water.

CAUTERIASMUS, Cauterization.

CAUTERIUM, C. actual'le, Cauter, Caut'ery, Inust'rium, Rupt'rium, Ignis actual'is, from καω, 'I burn,' καυρ'ν, 'a burner.' (F.) *Cautère, Feu actuel.* A substance, used for 'firing,' burning or disorganizing the parts to which it is applied. Cauteries were divided by the ancients into *actual* and *potential*. The word is now restricted to the red-hot iron; or to positive burning. It was, formerly, much used for preventing hemorrhage from divided arteries; and also with the same views as a blister. The term *Potent'ial Caustery, Caut'rium potentia'le, Ignis potentia'lis*, (F.) *Feu potentiel*, was generally applied to the causticum commune, but it is now used synonymously with caustic in general. *Cautère* also means an issue.

CAUTERIUM ACTUALE, Cauterium.

CAUTERIZATION, Cauteria'tio, Cauterias'mus, Exus'tio, Inus'tio, Caustica Adus'tio. Firing. The effect of a cautery or caustic. The French, amongst whom cauterization is much used, distinguished five kinds: 1. *Cautérisation Inhérente*, which consists in applying the actual cautery freely, and with a certain degree of force, so as to disorganize deeply. 2. *Cautérisation transcurrente*, which consists in passing the edge of the *Cautère cutellaire*, or the point of the *Cautère conique* lightly, so as not to disorganize deeply. 3. *Cautérisation par pointes*, which consists in applying on the skin, here and there, the hot point of the conical cautery, with sufficient force to cauterize the whole thickness of the skin. 4. *Cautérisation lente*, slow cauterization, by means of the moxa. 5. *Cautérisation objective*, which

consists in holding the cautery at some distance from the part to be acted upon by it.

CAUTÉRIZE; Caustico adurere; (F.) *Cautériser.* To apply a cautery or caustic. To burn with a cautery or caustic.

CAUTERY, Cauterium—c. Galvanic, see Moxa—c. Potential, Causticum commune, Cauterium.

CAVA, Vulva.

CAVA VENA, Vena hepatis. The hollow or deep-seated vein. (F.) *Veine cave.* A name given to the two great veins of the body, which meet at the right auricle of the heart. The *vena cava superior, thoracica seu descendens*, is formed by the union of the subclavians; and receives successively, before its termination at the upper part of the right auricle, the *inferior thyroid, right internal mammary, superior diaphragmatic, azygos, &c.* The *vena cava inferior, abdominalis seu ascendens*, arises from the union of the two *primary iliacs*, opposite the fourth or fifth lumbar vertebra, receives the *middle sacral, lumbar, right spermatic, hepatic, and inferior diaphragmatic*, and opens at the posterior and inferior part of the right auricle.

CAVATIO, Cavity.

CAVEA, Cavity—c. Narium, Nares.

CAVERNA, Antrum. 'A cavern.' This term has been used for the female organs of generation. See Cavity, and Vulva.

CAVERNA NARIUM, Nares.

CAVERNÆ DENTIUM, Alveoli dentium—c. Frontis, Frontal Sinuses.

CAVERNEUX, Cavernous.

CAVERNOUS, Cavernous, (F.) *Caverneux*. Filled with small cavities or caverns,—as a sponge.

CAVERNOUS BODIES; Cor'pora Cavernosa of the penis, *Cor'pora nervosa seu Ner'vco-spongio'sa Penis*, (F.) *Corps Caverneux*. The corpus cavernosum is a kind of cylindrical sac, composed of cells; separated, through its whole extent, by a vertical, incomplete septum, *Septum pectiniforme, Trabecula cor'porum cavernosorum*, and forming nearly two-thirds of the penis. The corpus cavernosum, on each side, arises from the ascending portion of the ischium, and terminates obtusely behind the glans. The arteries of the corpora cavernosa come from the internal pudic. See Helicine Arteries. Nerves are found on the surface of the outer membrane, but they do not appear to penetrate the substance, and the smooth muscular fibre has been traced into the fibrous parietes of the cells, as in the case of all erectile tissues.

J. Müller's researches have led him to infer, that both in man and the horse, the nerves of the corpora cavernosa are made up of branches proceeding from the organic as well as the animal system, whilst the nerves of animal life alone provide the nerves of sensation of the penis.

CAVERNOUS BODIES, Corpora Cavernosa of the Clitoris, are two hollow crura, forming the clitoris.

CAVERNOUS BODY OF THE VAGINA, Corpus Cavernosum Vaginae, Plexus retiformis, is a substance composed of blood-vessels and cells, similar to those of the penis and clitoris, which covers the outer extremity of the vagina, on each side. It serves to contract the entrance to the vagina during coition.

CAVERNOUS GANGLION, see Carotid or Carotid Nerve.

CAVERNOUS RESPIRATION. When a cavity exists in the lungs, and one or more ramifications of the bronchia terminate in it, a loud tubal noise is emitted, provided the cavity be not filled with fluid, which is called *cavernous respiration*. In this condition, the cough is *cavernous* likewise, (F.) *Toux Caverneuse*. When the capacity of the

ravern is very great, the sound of the respiration is like that produced by blowing into a decanter, with the mouth at a little distance from the neck. This kind of cavernous respiration has been called *amphoric*, from *amphora*, 'a flask;' (F.) *Respiration amphorique*, *Souffle amphorique*, *S. métallique*.

The *Veiled Puff*, (F.) *Souffle voilé*, is a modification of the cavernous respiration, in which, according to Laënnec, "a sort of movable veil interposed between the excavation and the ear" seems to be agitated to and fro. It is a sign which is not attended to.

CAVERNUS RHONCHUS, see Cavernous Respiration, and Gurgling.

CAVERNUS SINUS, *Sinus Cavernosus* seu *polymorphus* seu *sphenoidalis*, *Receptaculum sellæ equinæ lateribus oppositum*, (F.) *Sinus cavernosus*. The *Cavernous Sinuses* are venous cavities of the dura mater, filled with a multitude of reddish, soft filaments, intersecting each other; and, as it were, reticulated. They commence behind the inner part of the sphenoid fissure, pass backwards on the sides of the fossa pituitaria, and terminate by opening into a cavity, common to the superior and inferior petrosal sinuses. They receive some meningeal veins, the ophthalmic veins, &c. The anterior extremity of each cavernous sinus has been named the *ophthalmic sinus*.

CAVERNUS TEXTURE or TISSUE, *Tela cavernosa*, (F.) *Tissu caverneux*. The spongy substance which forms the greater part of the penis and clitoris. It seems to consist of a very complicated lace-work of arteries and veins; and, probably, of nervous filaments, with small fibrous plates, which form by their decussation numerous cells communicating with each other. This spongy texture produces erection, by dilating and swelling on the influx of blood; and probably, also, by virtue of some property inherent in it.

CAVERNUS WHISPER. A blowing sound, heard over cavities when words are whispered.—A. Flint.

CAVERNULOUS RHONCHUS, see Gurgling.

CAVIALE, Caviare.

CAVIARE', *Caviar*, *Caviale*, *Kaviar*. A culinary preparation, much used by certain people, and made, on the shores of the Black and Caspian Seas, from the roe of the sturgeon, mixed with salt and other condiments.

CAVIC'ULA, *Cavilla*, from *cavus*, 'hollow.' The ankle or space between the malleoli. Some have given this name to the os cuneiforme. See *Astragalus*.

CAVICULÆ PEDIS NODUS, Tarsus.

CAVIL'LA, *Astragalus*, *Cavila*.

CAVITAS ANTROSA AURIS, Tympanum—c. Buccinata. Cochlea—c. Cochleata, Cochlea—c. Digitata ventriculi lateralis, Cornu posterius ventriculi lateralis.

CAVITAS ELLIPTICA, *Ampulla*, *Sinus ampullæformis*. A dilatation at one end of the semicircular canals of the ear.

CAVITAS HUMERI GLENOIDES, see Glenoid—c. Narium, Nares—c. Oculi, Orbit—c. Oris, Mouth—c. Pulpæ, see Tooth.

CAVITATES CEREBRI, Ventricles of the brain—c. Dura matris, Sinuses of the dura mater—c. Innominata, Arteries of the heart—c. Interscapulares, see Interscapularis.

CAVITÉ, Cavity—c. Dentaire, Dental cavity—c. des Épiploons, see Peritonæum—c. du Tympan, Tympanum.

CAVITY, *Cæcitas*, *Cavum*, *Cælotas*, *Colom*, *Calvea*, *Caver'na*, *Cavatio*, (F.) *Cavité*. The hollow of anything, as of the cranium, mouth, nasal fossæ, &c.

CAVITIES, SPANCHNIC, (F.) *Cavités splanchniques*, are those which contain the viscera. They are three in number:—the cranium, chest, and abdomen. The cavities of bones, connected with joints or otherwise, are described under their particular denominations.

CAVUM, Cavity—c. Abdominis, see Abdomen. CAVUM CRA'NI, *Venter Supre'mus*. The cavity formed by the proper bones of the cranium.

CAVUM DENTIS, see Tooth—c. Mediastini, see Mediastinum—c. Narium, Nasal fossæ, Nares—c. Oris, Mouth—c. Pericardii, see Pericardium—c. Thoracis, see Thorax—c. Tympani, Tympanum.

CAYAN, *Phaseolus Creticus*.

CAZABI, *Jatropha manihot*.

CEANOTHOS, *Cirsium arvense*.

CEANOTHUS AMERICANUS, *Celastrus*—a. Trinervis, *Celastrus*.

CEAR, Heart.

CEASMA, Fissure.

CEBI GALLINÆ. The liver of the fowl, bruised.—Castelli.

CEBIP'ARA. A large Brazilian tree, whose bitter and astringent bark is used in making anti-rheumatic baths and fomentations.

CÉBOCÉPHALE, *Cebocéphalus*, from *κεβος*, 'an ape,' and *κεφαλή*, 'a head.' A genus of monsters, in which the nose does not project, and the interocular region is narrow and plane, so as to give the physiognomy a striking resemblance to that of the American ape.—I. G. St. Hilaire.

CECES, see *Quercus alba*.

CÉCITÉ, Cæcitas.

CECITY, Cæcitas.

CEDAR, RED, *Juniperus Virginiana*—a. White, *Cupressus thyoides*.

CEDEIA, Embalming.

CEDMA, Aneurism, Varix.

CED'MATA, *κιδάρτα*. Rheumatic pains of the joints, especially of the hips, groin, or genital organs. A form of gout or rheumatism.

CÉDRAT, *Citrus medica*.

CEDRELA FEBRIFUGA, *Swietenia febrifuga*.

CEDRELE'UM, from *κεδρος*, 'the cedar,' and *ελαϊον*, 'oil.' The oil of cedar.—Pliny.

CE'DRIA, *Ced'rium*, *Ced'rinum*, *Cedri lacryma*, *Alkitran*. The oil or resin which flows from the cedar of Lebanon. It was supposed to possess great virtues.—Hippocrates, *Foësius*, *Scribonius Largus*, *Dioscorides*. It has been supposed to be the same as the pyroligneous acid. See *Pinus Sylvestris*.

CE'DRINUM VINUM, *Cedar Wine*. A wine prepared by steeping half a pound of bruised cedar berries in six French pints of sweet wine. It is diuretic and subastringent.

CEDRI'TES, from *κεδρος*, 'the cedar.' A wine prepared from the resin of cedar and sweet wine. It was formerly employed as a vermifuge, &c.

CEDRIUM, *Cedra*.

CEDROMELA, see *Citrus medica*.

CEDRON, see *Simaba cedron*.

CEDRONELLA, *Melissa*—c. *Triphylla*, *Dra-cocephalum canariense*.

CEDROS, *Juniperus lycia*.

CE'DROSTIS, *Bryonia alba*.

CEDRUS BACCIFERA, *Juniperus sabina*—c. *Mahogani*, *Swietenia mahogani*.

CEINTURE, Cingulum, *Herpes zoster*—c. *Blanche de la choroïde*, *Ciliary ligament*—c. *Dactyosa*, *Herpes zoster*—c. *de Hildane*, Cingulum Hildani—c. *de Saint-Jean*, *Artemisia vulgaris*—c. *de Vif Argent*, Cingulum mercuriale.

CELANDINE, *Impatiens*—c. Common, *Chelidonium majus*—c. Lesser, *Ranunculus ficaria*—c. Poppy, *Stylophorum diphyllum*.

CELAS'TRUS, *Celas'tus*, *Ceanothus America'nus* seu *triner'cis*, *New Jersey Tea*, *Red Root*, (F.) *Thé de Jersey*. Used by the American Indians, in the same manner as lobelia, for the cure of syphilis. It is slightly bitter and somewhat astringent. A strong infusion of the dried leaves and seeds has been recommended in aphthæ, and as a gargle in scarlatina.

CELASTRUS SCANDENS, *Climbing Stafftree*. A climbing American shrub, the bark of which is said to possess emetic, diaphoretic, and narcotic properties.

CÉLATI'ON, (F.) *Concealment*, from *celare*, 'to conceal.' A word used by French medical writers for cases where there has been concealment of pregnancy or delivery.

CELĒ, κηλη, 'a tumour, protrusion, or rupture,' a very common suffix, as in hydrocele, bubonocèle, &c. See Hernia.

CELERY, (F.) *Céleri*. The English name for a variety of *Apium graveolens*.

CELERY, WILD, *Bubon galbanum*.

CELETA, see Hernial.

CELIA, *Cerevisia*.

CÉLIAQUE, *Cœliac*.

CÉLINE, *Melissa*.

CELLIS, κηλῖς, 'a spot, a stain.' A *macula*, or spot on the skin.

CELL, *Cella*. A small cavity. The same signification as *cellule*. Also, a vesicle composed of a membranous *cell-wall*, with, usually, liquid contents. The whole organized body may be regarded as a congeries of cells having different endowments, each set being concerned in special acts, connected with absorption, nutrition, and secretion, wherever an action of selection or elaboration has to be effected. These cells are generally termed *primary*, *elementary*, or *primordial*. When they give rise to other cells, they are, at times, termed *parent* or *mother cells*; the resulting cells being termed *daughter cells*.

CELL, **APOPLECTIC**, see Apoplectic cell—c. Bone, *Lacuna* of bone—c. Bronchic, *Cellule*, bronchic—c. Calcigerous, see Tooth—c. Daughter, see Cell—c. Elementary, see Cell.

CELL, **EPIDERMIC** or **EPITHELIAL**. The cells or corpuscles that cover the free membranous surfaces of the body, and which form the epidermis and epithelium, are termed '*epidermic* or '*epithelial cells*.' They are developed from germs furnished by the subjacent membrane.

CELL, **EPITHELIAL**, Cell, epidermic—c. Fat, see Fatty vesicles.

CELL FORCE. The plastic or formative force seated in a cell, by the agency of which the different tissues are developed. See Cell Life.

CELL FORMATION, Cytogeny—c. Genesis, Cytogeny—c. Germ, Cytoblast, see Molecule—c. Germinal, see Cytoblast—c. Nucleated, see Cytoblast.

CELL LIFE. The life which is possessed by the separate cells that form the tissues, and by which the nutrition of the tissues is presumed to be effected.

CELL, **MOTHER**, see Cell—c. Osseous, *Lacuna* of bone—c. Parent, see Cell.

CELL, **PIGMENT**. Pigment cells are mingled with the epidermic cells, and are most manifest in the coloured races. They are best seen on the inner surface of the choroid of the eye, where they form the *pigmentum nigrum*.

CELL, **PRIMARY**, see Cell—c. Primordial, see Cell—c. Sperm, see Sperm—c. Wall, see Cell.

CELLA, Cell—c. Media, see Ventricles of the Brain—c. Turcica, *Sella Turcica*.

CELLOID, *Celloi'des*, badly compounded from

cella, and αἶος, 'resemblance.' Resembling a cell or cellule.

CELLULA, *Cellule*.

CELLULÆ COLL, see Colon—c. *Aerem*, *Cellules*, bronchic—c. *Bronchicæ*, see Cellule—c. *Mastoideæ*, *Mastoid cells*—c. *Medullares*, see *Medullary membrane*—c. *Pulmonales*, *Cellules bronchicæ*, see Pulmo—c. *Sanguinis*, *Globules* of the blood.

CELLULAR, *Cellularis*, *Cellulo'sus*, (F.) *Cellulaire*. Composed of cells or cellules, from *cella* or *cellula*, 'a cell.'

CELLULAR MEMBRANE, *Membrana cellulo'sa* seu *Cellularis*—*M. adiposa* seu *pinguedino'sa*, of some, *Panniculus adiposus*,—Membrane formed of cellular tissue, (F.) *Membrane cellulaire*. Generally used for the tissue itself.

CELLULAR SYSTEM. The whole of the cellular tissue of the human body.

CELLULAR TISSUE, *Tela cellularis* seu *cellulo'sa* seu *Hippoc'ratis cribro'sa*, *Eth'myphæ*, *Contextus cellulo'sus*, *Reticulated*, *filamentous*, *laminated*, *crib'riform*, *porous*, *areolar*, *connective*, and *mucous Tissue*, *Reticular* or *cellular substance*, (F.) *Tissu cellulaire*, *réticulé*, *lamineux*, *cribleux*, *poreux*, *aréolaire*, *muqueux*, *conjonctif*, *Ethmose* of Piorry, &c., is the most common of all the organic tissues. It contains irregular *areolæ* between the fibres, as well as serum, fat, and the adipous tissue. Of the fibres, some are of the yellow elastic kind; but the greater part are of the white fibrous tissue, and they frequently present the form of broad flat bands, in which no distinct fibrous arrangement is perceptible. See Fibrous.

The cellular tissue or texture unites every part of the body, determines its shape, and by its elasticity and contractility, and by the fluid which it contains in its cells, facilitates the motion of parts on each other.

Cellular tissue has been divided by anatomists into the external, general or common cellular tissue—*textus cellularis intermedius* seu *laxus*, which does not penetrate the organs,—the cellular texture which forms the envelopes of organs—*textus cellularis strictus*, and that which penetrates into the organs, accompanying and enveloping all their parts,—the *textus cellularis stipatus*, constituting the basis of all the organs. It has likewise been termed *Textus organicus* seu *parenchymatis*.

CELLULAR TISSUE OF BONES, see Canecelli.

CELL'ULE, *Cell'ula*, diminutive of *cella*, 'a cavity.' A small cavity. (F.) *Cellule*. *Cellules* are the small cavities between the laminae of the cellular tissue, corpora cavernosa, &c.

CELLULES OF CELLS, **BRONCHIC**, *Cell'ulæ Bronchicæ* seu *Pulmonales* seu *Ac'rcæ*, *Porî pulmonum*, *Vesic'ulæ pulmonales*. The air-cells of the lungs. See Pulmo.

CELLULES BRONCHIQVES, Bronchial cells—c. *des Os*, *Lacuna* of Bone—c. *Ossenses*, *Lacuna* of Bone.

CELLULITIS VENENATA, see Wound.

CELLULO-FIBROUS TUMOUR, Tumour, fibro-cellular.

CELLULOSE, same etymon as *Cellule*. The substance which is left after the action upon any kind of vegetable tissue of such solvents as are fitted to dissolve out the matter deposited in its cavities and interstices. The tunicated or ascidian mollusca have, in their integuments, a considerable quantity of it, and it exists in the Corpora amylacea of the brain and other parts. It is identical with starch, in the proportion of its constituents, and forms the principal part of the thickness of the walls of the cells, vessels, &c., of which the vegetable organism is composed.

CELLULOSUS, *Cellular*.

CELOLOG'IA, from κηλη, 'rupture,' and λογος,

a discourse.' The doctrine of hernia. A treatise on hernia.

CELO'SOMUS, from *κηλη*, 'a rupture,' and *σωμα*, 'body.' A monster in which the trunk is malformed, and evagination or displacement of the viscera exists.

CELOTES, see Hernia.

CELOTOM'IA, *Kelotom'ia*, *Celot'omy*, from *κηλη*, 'a rupture,' and *τεμνειν*, 'to cut.' An operation, formerly employed for the radical cure of inguinal hernia; which consisted, principally, in passing a ligature round the hernial sac and spermatic vessels. It necessarily occasioned atrophy and loss of the testicle; and did not secure the patient against the return of the disease. The intestines were, of course, not included in the ligature. Also, the operation for hernia in general.—*Herniot'omy*.

CELOT'OMUS, same etymon. *Herniot'omus*. A knife used in the operation for hernia. Adjectively, it means relating to celotomy, like *Celotom'icus*.

CELSA. A term, used by Paracelsus for a cutaneous disease, dependent, according to him, on a false or heterogeneous spirit or vapour, concealed under the integuments, and endeavouring to escape. Perhaps the disease was *Urticaria*.

CELSUS, METHOD OF, see Lithotomy.

CELTIS OCCIDENTALIS, *Sugarberry*, *Hackberry*, *Nettle-tree*, *Beverwood*. Order, Ulmaceæ: indigenous, flowering in May. The bark is said to be anodyne and cooling; the berries are sweet and astringent. It has been used in dysentery.

CEMBRO NUTS, see *Pinus cembra*.

CEMENT. A glutinous substance introduced into a carious tooth to prevent the access of air or other extraneous matters. The following is an example: *R. Sandarac. ʒij. Mastich. ʒi. Succin. gr. x. Ether. ʒi.* Dissolve with the aid of heat. *Osteo-medic's Cement for the teeth* is prepared of finely powdered *caustic lime*, thirteen parts; *anhydrous phosphoric acid*, twelve parts. When introduced into a carious tooth, it becomes solid in about two minutes.

CEMENTERIUM, Crucible.

CEMENTUM, see Tooth.

CENANGIA, *Ceneangia*.

CENCHRON, *Panicum miliaceum*.

CENDRE DU LEVANT, Soda.

CENDRÉ, Cineritious.

CENDRÉS GRAVELÉES, see Potash—*c. de Sarcenet*, see Potash.

CENEANG'IA, *Cenangi'a*, from *κενος*, 'empty,' and *αγγειον*, 'a vessel.' Inanition. Empty state of vessels.—Galen.

CENEMBATE'SIS, from *κενος*, 'empty,' and *εμβαλλω*, 'I enter.' Paracentesis. Also, the act of probing a wound or cavity; *Melo'sia*.

CENEONES, Flanks.

CENESTHÉSIE, *Cenæsthesia*.

CENIGDAM, *Ceniplam*.

CENIGOTAM, *Ceniplam*.

CENIPLAM, *Ceniglam*, *Cenigotam*, *Cenipolam*. The name of an instrument anciently used for opening the head in epilepsy.—Paracelsus.

CENIPOLAM, *Ceniplam*.

CENO'SIS, from *κενος*, 'empty.' *Ine'sis*, *Inethmos*. Evacuation. It is sometimes employed synonymously with inanition, and opposed to repletion.—*Emeric'io*.

CENOSPUDIA, *Aphelxia otiosa*.

CENOTIC, Drastic.

CENOTICA, from *κενωσις*, 'evacuation.' Diseases affecting the fluids. Morbid discharges or excess, deficiency or irregularity of such as are natural. The first order, class *Genetica*, of Good. Also, Drastics.

CENTAUREA BEHEN, *Serrat'ula behen*, *Be-*

hen abiad, *Behen album*, *Been*, *White Behen*. Ord. Gentianeæ. Astringent.

CENTAUREA BENEDICTA, *Carduus benedictus*, *Chnicus sylvestris* seu *benedictus*, *Cardiobolbanum*, *Blessed or Holy Thistle*, *Virgin-Mary Thistle*, (F.) *Chardon béat*. Fam. Cynarocephaleæ. S.S. Syst. Syngenesia Polygamia fruticosa. A strong decoction of the herb is emetic;—a strong infusion, diaphoretic [?]; a light infusion, tonic and stomachic. Dose, gr. xv to ʒj of the powder.

CENTAUREA CALCITRATA, *Calcitrap'ea*, *Calcitrep'pola*, *Carduus solstitialis* seu *stellatus*, *Jalisco ramosissima*, *Caecotribulus*, *Calcitrap'ea stellata* seu *hippophæstum*, *Stellata rapi*, *Centaurea stellata*, *Common Star-Thistle*, *Star-Knapweed*, (F.) *Centauree étoilée*, *Chardon étoilé*, *Chaussetrappe*, *Pigneroles*. It is possessed of tonic properties, and has been given in intermittents, dyspepsia, &c. It is not much used.

CENTAUREA CENTAURIUM, *Rhepon'ticum vulgare*, *Centauri'um magnum* seu *majus* seu *officinale*, *Greater Centaury*, (F.) *Centauree grande*. It is a bitter; and was formerly used as a tonic, especially the root.

CENTAUREA CYANUS, *Cy'annus*, *Blue bottle*, *Corn-flower*, *Blue bonnets*, (Sc.) *Blawort*, (F.) *Blavelle*, *Blavéole*, *Blavérolle*, *Bluet*, *Barbeau*, *Aubifoin*, *Casse-lunette*. The flowers were once much used as a cordial, tonic, &c. They are now forgotten.

CENTAUREA STELLATA, *Centaurea calcitrapa*.

CENTAURÉE ÉTOILÉE, *Centaurea calcitrapa*—*c. Grande*, *Centaurea centaurium*—*c. Petite*, *Chironia centaurium*.

CENTAUREUM, *Chironia centaurium*.

CENTAURIS, *Chironia centaurium*.

CENTAURIUM MAGNUM, *Centaurea centaurium*—*c. Minus vulgare*, *Chironia centaurium*—*c. Officinale*, *Centaurea centaurium*—*c. Parvum*, *Chironia centaurium*.

CENTAURY, AMERICAN, *Chironia angularis*—*c. European*, *Chironia centaurium*—*c. Greater*, *Centaurea centaurium*—*c. Lesser*, *Chironia centaurium*.

CENTESIS, Paracentesis, Puncture.

CENTIGRAMME, (F.) from *centum*, 'a hundred,' and *γρᾶμμα*, 'gramme,' *Centigramma*. The hundredth part of a gramme. A centigramme is equal to about the fifth part of a French grain, gr. .1543 Troy.

CENTILITRE, *Centili'tra*, from *centum*, 'a hundred,' and *λίτρον*, 'litre.' An ancient Greek measure for liquids:—the hundredth part of a litre—equal to nearly 2.7053 fluidrachms.

CENTIMÈTRE, *Centim'eter*; the hundredth part of a metre—equal to about four lines, .3937 English inch.

CENTIMORBIA, *Lysimachia nummularia*.

CENTINERVA, *Plantago*.

CENTINODE, *Polygonum aviculare*.

CENTINODIA, *Polygonum aviculare*.

CENTO VIRGINALIS, Hymen.

CENTRAD, see Central aspect.

CENTRADIAPHANES, *Cataracta centralis*, from *κεντρον*, 'centre,' *a*, privative, and *εἰσφανς*, 'transparent.' Cataract owing to obscurity of the central portion of the crystalline.

CENTRAL, *Centra'tis*, from *centrum*, 'the centre.' Relating or appertaining to the centre.

CENTRAL ARTERY OF THE RETINA, *Arteria Centra'tis Retinae*, *Central Artery of Zinn*. This artery is given off from the *arteria ophthalmica*, and penetrates the optic nerve a little behind the ball of the eye; running in the axis of the nerve, and spreading out into many small branches upon the inside of the retina. When the nerve is cut across near the eye, the

orifice of the divided artery is observable. This was formerly called *Porus Opticus*.

CENTRAL ASPECT. An aspect towards the centre of an organ.—Barelay. Centrad is used by the same writer adverbially, to signify 'towards the central aspect.'

CENTRE, *Κέντρον*, (F.) Centre. The middle point of a figure or body.

CENTRE OF ACTION. The visus in which the whole or a great part of any function is executed, and to which several other organs contribute. Thus, the vital activity seems to be wholly centred in the stomach, during chymification; in the duodenum, during chylication. In like manner, the uterus becomes a centre of action during gestation.

CENTRE, EPIGASTRIC. The ganglions and nervous plexuses, formed by the great sympathetic and pneumogastric nerves, in the epigastrium, around the coeliac artery; where the impressions received from various parts of the body seem to be centred.

CENTRE OF FLUXION. The part towards which fluids are particularly attracted. An irritated organ is said to be a centre of fluxion.

CENTRES, NERVOUS, (F.) Centres nerveux. The organs whence the nerves originate; as the brain and spinal marrow.

CENTRE, OPTIC, see Optic centre.

CENTRE, OVAL, *Centrum Ovale*, C. O. Vienssenii, *Tegmen tum ventriculo rum cerebri*. When the two hemispheres of the brain are sliced away, till on a level with the corpus callosum, the medullary part in each is of an oval shape: hence called *centrum ovale minus*, (F.) *centre médullaire hémiphéral*. The two centres of the opposite sides, together with the corpus callosum, form the *centrum ovale of Vienssens*. Vienssens supposed all the medullary fibres to issue from that point, and that it was the great dispensatory of the animal spirits.

CENTRE, PHRENIC, *Tendineum Centre of the Diaphragm, Centrum Phrenicum seu Nervinum seu Tendineum seu Tendineum*, (F.) *Centre phrénique ou tendineux du Diaphragme*. The central aponeurosis or cordiform tendon of the diaphragm.

CENTRE OF SYMPATHETIC IRRADIATIONS, (F.) *Centre d'irradiations sympathiques*. Any organ which excites, sympathetically, the action of other organs, more or less distant from it; and with which it seems to have no immediate communication.—Marjolin.

CENTRE, TENDINOUS, OF THE DIAPHRAGM, Centre, phrenic.

CENTROMYRINE, Rusens.

CENTROSTAL'TIC, *Centrostalticus*, badly formed from *centrum*, 'the centre,' and *στέλλω*, 'I send.' An epithet applied by Dr. Marshall Hall to the action of the *vis nervosa* in the spinal centre.

CENTRUM, see Vertebrae—c. Commune, Solar plexus—c. Nerveum, Centre, phrenic—c. Opticum, Optic centre—c. Ovale, Centre, oval—c. Ovale minus, see Centre, oval—c. Ovale of Vienssens, Centre, oval—c. Semicircularis geminum, Tænia semicircularis—c. Tendinosum, Centre, phrenic.

CENTRUM VITA'LE, *Nodus seu Fons vitalis*, (F.) *Nœud vital*. A term applied, at times, to the medulla oblongata; at others, to the medulla oblongata, and the medulla spinalis as far as the second cervical nerve of the spinal marrow, in any part of which a wound would seem to be instantly fatal. It is the nervous centre of respiration and deglutition.

CENTRY, Chironia angularis.

CENTUM CAPITA, *Eryngium campestre*.

CENTUMNODIA, *Polygonum aviculare*.

CEPA ASCALONICA. Bulbus esculentus, Echalotte—c. Marina, Scilla—c. Victorialis, Allium victorale—c. Vulgaris, Allium cepa.

CEPÆA, Veronica beccabunga.

CEPHAELINUM, Emetine.

CEPHAELIS IPECACUANHA, Ipecacuanha.

CEPHALÆA, *Headach*, (F.) *Céphalée*, from *κεφαλή*, 'head.' Some use the term synonymously with cephalalgia; others, for a periodical head-ach; others, again, for a more violent headach than cephalalgia implies; and others for a chronic headach. The last was its ancient signification.

Cephalæa spasmodica, Cephalalgia spasmodica, C. Nauseosa, Sick-headach, is characterized by partial, spasmodic pain: often shifting from one part of the head to another: chiefly commencing in the morning, with sickness and faintness. It is extremely apt to recur, notwithstanding every care.

CEPHALÆA ARTHRITICA, Cephalagra—c. Hemierania, Hemierania—c. Nauseosa, C. Spasmodica—c. Pulsatilis, Crotaphæ.

CEPHALÆMATOMA, (F.) *Céphalœmatome*, from *κεφαλή*, 'head,' and *αἷμα*, 'blood.' *Cephalæmatoma neonatorum, Ecchymoma capitis, E. capitis recens natorum, Thrombus neonatorum, Abscessus capitis sanguineus neonatorum, Tumor capitis sanguineus neonatorum, Cephalophyma, Craniohæmatomus.* A sanguineous tumour, sometimes developed between the pericranium and the bones of the head of new-born children. Similar tumours are met with occasionally above other bones, and at all periods of existence.

CEPHALÆMATOMA NEONATORUM, Cephalæmatoma.

CEPHALÆMIA, *Hyperæmia cerebri seu Capitis, Encephalohæmia*, (F.) *Hyperémie ou Congestion du cerveau, Encephalohémie, H. cérébrale, Congestion cérébrale*. Accumulation of blood in the vessels of the brain.

CEPHALOGÆGUS, *Cephaloductor, Capiti-ductor*, from *κεφαλή*, 'head,' and *αγωγός*, 'a leader, a driver.' An instrument used for drawing down the fetal head.

CEPH'ALAGRA, from *κεφαλή*, 'the head,' and *αγρα*, 'seizure.' *Cephalæa arthritica, Menin-gitis arthritica.* Gout in the head.

CEPHALAGRA'PHIA, from *κεφαλή*, 'the head,' and *γραφη*, 'a description.' An anatomical description of the head.

CEPHALAL'GIA, *Cephaloponia, Cephalodyn'ia, Encephalodyn'ia, Homonop'agia*, from *κεφαλή*, 'the head,' and *αλγος*, 'pain'; *Encephal'gia, Dolor Capitis seu cephal'icus, Suda, Pain in the head; Headach, (Sc.) Hedeveerk, (Prov.) Hedeveark, (F.) Céphalalgie, Mal à tête*. Every kind of headach, whether symptomatic or idiopathic, is a cephalalgia. It is ordinarily symptomatic, and has to be treated accordingly.

CEPHALALGIA CONTAGIOSA, Influenza—c. Inflammatoria, Phrenitis.

CEPHALALGIA PERIODICA, *Febris intermittens cephalica larvata, Intermittent headach*. Headach which returns periodically; properly, perhaps, a form of neuralgia.

CEPHALALGIA PULSATILIS, Crotaphæ—c. Spasmodica, see Cephalæa.

CEPHALALOG'IA, from *κεφαλή*, 'the head,' and *λογος*, 'a discourse.' An anatomical dissection on the head.

CÉPHALANTHE D'AMÉRIQUE, Cephalanthus occidentalis.

CEPHALANTHUS OCCIDENTALIS, Buttonwood shrub, Buttonbush, White Ball, Little Snowball, Scaumpwood, Pond Dogwood, Globe-flower, (F.) *Céphanthe d'Amérique, Bois de Marais*. An ornamental shrub, *Ord. Rubiaceæ; Ser. Syst. Tetrandria Monogynia*, which grows all

over the United States, near streams and ponds, and flowers in July and August. The bark of the root has been used as an antiperiodic tonic.

CEPHALARTICA, Cephalic remedies.

CEPHALATOMIA, *Cephalotomia*, from κεφαλή, 'the head,' and τέμνω, 'to cut.' Anatomy, or dissection, or opening of the head.

CEPHALE, Head.

CEPHALÉMATOME, Cephalæmatoma.

CEPHAL'IC, *Cephal'icus*, *Capit'alis*, from κεφαλή, 'the head,' (F.) *Céphalique*. Relating to the head.

CEPHAL'IC REM'EDIES, *Cephal'ica* seu *Capit'alia remedia*, are remedies capable of relieving affections of the head, especially headach:—*Cephalartica*.

CEPHAL'IC VEIN, *Vena Cephal'ica* seu *Capit'itis*, (F.) *Veine céphalique*, *Veine radiale cutanée* of Chaussier. The great superficial vein at the outer part of the arm and fore-arm. It begins on the back of the hand, by a number of radicles, which unite into a single trunk, called the *Cephal'ic of the Thumb*, *Cephal'ica Pollicis*, (F.) *Veine céphalique du pouce*. It ascends along the anterior and outer part of the fore-arm, where it forms the *superficial radial*. At the fold of the elbow it receives the *median cephalic*, ascends along the outer edge of the biceps, and opens into the axillary vein. The name *Cephalic* was given to it by the ancients, because they thought it had some connection with the head, and that blood-letting ought to be performed on it, in head affections.

Chaussier calls the internal jugular, *Veine céphalique*, and the primary or common carotid, *Artère céphalique*.

CEPHALIDIUM, see Caput.

CEPHALINÆ. The base or root of the tongue. —Glossans.

CEPHALIS, see Caput.

CEPHALITIS, Phrenitis.

CEPHALUM, see Caput.

CEPHALODUCTOR, Cephalagogus.

CEPHALODYMIA, *Encephalodymia*; from κεφαλή, 'head,' and δύνω, 'I enter into.' A class of double monstrosities, in which the heads are united. It is divided into two genera, *Frontodymia* and *Bregmatodymia*: in the former the union being between the ossa frontis; in the latter between the bregmata.—Cruveilhier.

CEPHALODYNIA, Cephalalgia.

CEPHALŒDEMA, Hydrocephalus.

CEPHALOGRAPHY, *Cephalographia*; from κεφαλή, 'head,' and γραφή, 'a description.' An anatomical description of the head.

CEPHALOID, Encephaloid.

CEPHALOLOG'Y, *Cephalolog'ia*; from κεφαλή, and λόγος, 'a discourse.' A treatise on the head.

CEPHALOMA, Encephaloid.

CEPHALOM'ELUS, (F.) *Céphalomèle*; from κεφαλή, 'head,' and μέλος, 'a limb.' A monster having one or two accessory limbs on the head. —I. G. St. Hilaire.

CEPHALO-MENINGITIS, Meningo-cephalitis.

CEPHALOM'ETER, (F.) *Céphalomètre*, from κεφαλή, 'the head,' and μέτρον, 'measure.' An instrument for measuring the different dimensions of the foetal head, during the process of accouchement. A kind of forceps.

CEPHALON'OSUS, from κεφαλή, 'the head,' and νόσος, 'disease.' This term has been applied to the *Febria Hungar'ica*, in which the head was much affected. See Fever, Hungaria. Others have so called any cerebral disease or fever.

CEPHALOPAGES, Symphyoccephalus.

CEPHALO-PHARYNGE'US, from κεφαλή, 'the head,' and φάρυγξ, 'the pharynx,' belonging to the head and pharynx. Winslow has given

this name to the portion of the *constrictor pharyngis superior*, which is attached, above, to the inferior surface of the basiliary process of the os occipitis. The *Cephalo-pharynge'al Aponeurosis* is a thin, fibrous membrane, which is attached to the basiliary process, and gives insertion to the fibres of the *constrictor superior pharyngis*.

CEPHALOPHYMA, Cephalæmatoma.

CEPHALOPONTA, Cephalalgia.

CEPHALO-RACHIDIAN, Cephalo-spinal.

CEPHALOSOMATODYM'IA, *Encephalosomatodymia*; from κεφαλή, 'head,' σῶμα, 'body,' and δύνω, 'I enter into.' A double monstrosity, in which the union is between the heads and the trunks. Of this there are varieties:—for example, *Infra-maxillosternodymia*, where the union is with the inferior maxillary bones and sternum; and *Prosoposternodymia*, between the faces and sternum.—Cruveilhier.

CEPHALO-SPINAL, *Cephalo-spinal'is*, *Cephalo-rachid'ian*, *Enceph'alo-rachid'ian*, *Cer'ebro-rachid'ian*, *Cranio-spinal*. A hybrid term, from κεφαλή, 'head,' and σπία, 'spine.' Belonging to the head and spine.

CEPH'ALO-SPINAL FLUID, *Cephalo-rachid'ian fluid*, *Cer'ebro-spinal fluid*, *Flu'idum cer'ebro-spinal'e*, *Subarachnoid'ean fluid*, (F.) *Liquide céphalo-rachidien*, is an exhaled fluid, which is found beneath the arachnoid, wherever pia mater exists in connexion with the brain and spinal cord. It seems to have a protecting office, and to keep up a certain degree of pressure on the organ—at least in the spinal canal.

CEPHALOTHORACOSTERUMENUS, from κεφαλή, 'head,' θώραξ, 'the chest,' and στήναι, 'to rob.' A monster without head or chest.

CEPHALOTOMIA, Eccephalosis.

CÉPHALOTRIBE, (F.) An instrument invented by Baudelocque, the nephew, for crushing the head of the foetus in utero; from κεφαλή, 'the head,' and τριβω, 'I bruise.' It consists of a strong forceps, the blades of which are solid: 16 lines broad, and 3 thick. The handles are perforated at their extremity to receive a screw with three threads, the direction of which is very oblique, so as to allow great rapidity of rotation, and the screw is moved by a winch 6 inches long, to increase the force of the pressure. The bones of the head are easily crushed by it.

CEPHALOTRIP'SY, *Cephalotrip'sis*; same etymon as *Cephalotribe*. The operation of crushing the head of the foetus in utero.

CEPHALOTRYPESIS, Trepanning.

CEPHALOXIA, Torticollis.

CEPULLA, Allium cepa.

CER, Heart.

CERA FLAVA et CERA ALBA, *Ceros*, *Yellow and White Wax*, (F.) *Cire Jaune et Blanche*. An animal substance prepared by the bee, and by some plants, as the *Cerox'ylon* and *Myrica cerif'era*. Its colour is yellow, and smell like that of honey, but both are lost by bleaching. It is demulcent and emollient; is sometimes given in the form of emulsion, in diarrhoea and dysentery, but is chiefly used in cerates and ointments.

CERÆ'Æ, from κερας, 'a horn,' κεραία. The Cornua of the uterus.—Rufus of Ephesus.

CERAM'ICÆ, *Cerami'tis*, from κεραμος, 'potter's earth.' A sort of earth used as a cataplasm in peripneumony.—Hippocrates.

CERAMIUM, Amphora—c. Helminthochortus, Corallina Corsianna.

CEKAMNIUM, Amphora.

CERAMURIA, see Urine.

CERANTHEMUS, Propolis.

CERAS, κερας, 'genitive,' κερατος, 'horn,' Cornu; also, the Cornua. Hence, *Ceratotomy*, *Ceratocle*, &c.

CERASION, see *Prunus cerasus*.

CERAS'MA, from *κεραυνωμι*, 'to mix.' something mixed. A mixture of hot and cold water. *Meleoceras'ma*.—Gorrieus.

CERAS'UM, see *Prunus cerasus*.

CERAS'US ACIDA, *Prunus cerasus*—c. *Avium*, *Prunus avium*, *P. nigra*—c. *Dulcis*, *Prunus nigra*—c. *Hortensis*, *Prunus cerasus*—c. *Lauro-cerasus*, *Prunus lauro-cerasus*—c. *Padus*, *Prunus padus*—c. *Racemosus sylvestris*, *Prunus padus*—c. *Rubra*, *Prunus cerasus*—c. *Serotina*, *Prunus Virginiana*—c. *Virginiana*, *Prunus Virginiana*—c. *Vulgaris*, *Prunus cerasus*.

CÉRAT BLANC' ou DE GALIEN, *Ceratum Galeni*—c. *de Blanc de Baleine*, *Ceratum cetacei*—c. *de Goudard*, *Ceratum plumbi*—c. *pour les Lèvres*, *Cerate for the lips*—c. *de Plomb composé*, *Ceratum plumbi compositum*—c. *de Savon*, *Ceratum Saponis*—c. *de Suracétate de plomb*, *Ceratum plumbi superacetatis*.

CERATE, *Ceratum*, from *κησας*, Lat. *cera*, 'wax,' *Cerelæum*, *Cero'ma*, *Cero'nium*, *Cero'tum*, *Ceroleum*, *Oleo-ceratum*, *Ceratomalag'ma*, (F.) *Cérat*, *Oléo-cérat*, *Eléocérat*, *Cérat*, *Lipuroid*. A composition of wax, oil, or lard, without other ingredients.

CERATE, *Simple Cerate*, *Ceratum*, *Ceratum simplex* (Ph. U. S.), (F.) *Cérat Simple*. (White wax, \mathfrak{z} iv, Lard, \mathfrak{z} vii.) It is applied as an emollient to excoriations, &c. See *Ceratum Cetacei*.

CERATE, BELLEVILLE'S, see *Unguentum Hydrargyri nitrico-oxidi*.

CERATE OF CALAMINE, *Ceratum Calami'næ seu Calami'næ preparat'æ seu Carbonat'is zinci impur'is seu Zinci Carbonat'is seu lap'idis Calami'næ seu epulot'icum*, *Cerate of Carbonate of Zinc*, *Turner's Cerate*, *Healing Salve*, (F.) *Cérat de Pierre Calami'næ*, *C. de Calamine*. (Calamine, *Cera flava*, \mathfrak{aa} \mathfrak{z} ij, *alapis*, lbj. Melt the wax and lard together, and, on cooling, add the carbonate of zinc and stir till cool.—Ph. U. S.)

CERATE OF CANTHARIDES, *Ceratum Canthar'idis seu Lytta*, *Blister Ointment*, *Ointment of Spanish Flies*, *Unguentum ad vesicato'ria seu Pulveris Mel'is vesicato'rii seu epispasticum fortius*, (F.) *Cérat de Cantharides*. (*Spermaceti cerate* \mathfrak{z} ij, *Cantharides in powder*, \mathfrak{z} j. The cerate being softened by heat, stir in the flies.) This cerate of the European Pharmacopæias is used to keep blisters, issues, &c., open. See *Unguentum Lyttæ*. For the Cerate of Spanish flies of the U. S. Pharmacopæia, see *Emplastrum Lyttæ*.

CERATE, GOULARD'S, *Ceratum plumbi compositum*.

CERATE, KIRKLAND'S NEUTRAL. (*Diachyl.* \mathfrak{z} vii, *aloe oil* \mathfrak{z} iv, *prepared chalk* \mathfrak{z} iv: when nearly cool, add *Acet. dest.* \mathfrak{z} iv, *plumb superacet.* \mathfrak{z} ij.) A cooling emollient.

CERATE OR POMATUM FOR THE LIPS, *Ceratum labiat'æ cœbrum*, *Pommatum ad labia demulcent'ia*.—Ph. P. (F.) *Cérat ou Pommade pour les Lèvres*. (Wax 9 parts; oil 16 parts;—coloured with alkane.)

CERATE, LEAD, COMPOUND, *Ceratum plumbi compositum*.

CERATE, MARSHALL'S. (*Palm oil* \mathfrak{z} vi, *calomel* \mathfrak{z} j, *sugar of lead* \mathfrak{z} ss, *ointment of nitrate of mercury* \mathfrak{z} ij.)

CERATE, RESIN, COMPOUND, *Ceratum Resinæ compositum*—c. *Savine*, *Ceratum sabine*—c. *Soap*, *Ceratum saponis*—c. *Spermaceti*, *Ceratum cetacei*—c. of *Superacetate* or *sugar of lead*, *Ceratum plumbi superacetatis*—c. *Turner's Cerate of Calamine*—c. of *Carbonate of zinc*, *Cerate of calamine*.

CERATECTOM'IA, from *κερας*, *keratos*, 'the

cornea,' and *εκτομος*, 'cut out.' An incision through the cornea. See *Ceratotomy*.

CERATIA, *Ceratonium siliqua*.

CERATIASIS, *Kerati'asis*, from *κερας*, *keratos*, 'horn.' A morbid condition characterized by corneous growths.

CERATION, *Siliqua*.

CERATITIS, *Kerati'tis*, from *κερας*, *keratos*, 'the cornea,' and *itis*, 'inflammation.' Inflammation of the cornea, *Kerati'tis*, *Ceratodē'tis*, *Keratodē'tis*, *Ceratomeningi'tis*, *Cornei'tis*, *Injunctio cor'neæ*.

CERATIUM, *Ceratonium siliqua*.

CER'ATO. In composition, relating or appertaining to the cornua of the os hyoides, or to the cornea.

CERATOCE'LE, *Keratoce'lē*, *Aquila*, *Urat'io*, *Promine'tia Cor'neæ*, *Hernia Cor'neæ*, *Ceratodē'ce'lē*, from *κερας*, *keratos*, 'horn,' and *κηλη*, 'tumour.' A protrusion of the transparent cornea, or rather of the membrane of the aqueous humour, through an opening in the cornea.

CERATODEITIS, *Ceratitis*.

CERATODEOCELE, *Ceratocele*.

CERATODEONYXIS, *Ceratonyxis*.

CERATODES MEMBRANA, *Cornea*.

CERATOGLOS'SUS, *Keratoglossus*, from *cerato*, and *γλωσσα*, 'the tongue.' A muscle, extending from the great cornu of the os hyoides to the base of the tongue. It is a part of the hyoglossus.

CERATOIDES, *Cornea*.

CERATOLEUCOMA, *Leucoma*.

CERATO'MA, *Cerato'sis*, from *κερας*, *keratos*, 'horn.' A horny growth, or horny formation.

CER'ATO-MALA'CIA, *Ker'ato-mala'cia*, from *κερας*, *keratos*, 'cornea,' and *μαλακία*, 'softness.' Softening of the cornea, the result of inflammation or of starvation.

CERATO-MALAGMA, *Cerate*.

CERATO-MENINGITIS, *Ceratitis*.

CERATO-MENINX, *Cornea*.

CERATO'NIA SIL'IQUA, from *κερας*, *keratos*, 'a horn'—from its hornlike pod. The *Carob Tree*, *Ceratum*, *Ceratia*, *Sil'iqua dulcis*, *Ceratoba Alnabati*, *Sweetpod*, (F.) *Caroubier*, (*Fruit*, *Carouge*.) This—the fruit of the *Ceratonia siliqua*—is mucilaginous, and employed in decoction where mucilages are indicated.

CERATONYX'IS, *Keratonyx'is*, *Ceratodeonyx'is*, from *κερας*, *keratos*, 'the cornea,' and *ωνωω*, 'I puncture.' An operation by which the crystalline is depressed by means of a needle introduced into the eye through the cornea. Some divide the crystalline into fragments with the needle, and leave them to the action of the absorbents. The operation is as old as the 17th century.

CER'ATO-PHARYNGE'US, *Ker'ato-Pharynge'us*, from *cerato*, and *φαρυγξ*, 'the pharynx.' The great and small *Cer'ato-pharyngei* are small fleshy bundles, forming part of the *Hypopharyngens* of Winslow.

CERATOPLAS'TICĒ, *Cer'atoplasty*, *Ker'atoplasty*, from *κερας*, *keratos*, 'the cornea,' and *πλαστικός*, 'forming, formative.' The operation for the formation of an artificial cornea. It has not been practised on man.

CERATORRHEX'IS, *Rupt'ra cor'neæ*, from *κερας*, *keratos*, 'the cornea,' and *ρηξις*, 'rupture.' Rupture of the cornea.

CERATOSIS, *Ceratoma*.

CER'ATO-STAPHYLINUS, *Ker'ato-staphylinus*, from *cerato*, and *σταφυλη*, 'the uvula.' Some fleshy fibres of the *Thyro-staphylinus* of Winslow.

CERATOTO'MIA, *Ceratotomy*, from *κερας*, *keratos*, 'cornea,' and *τεμνειν*, 'to cut.' Section of the transparent cornea. This incision is used

in the operation for cataract, to give exit to pus effused in the eye, in case of hypopyon, &c.

CERATOTOMUS, *Keratotomus*, *Kerat'omus*, from *keras*, κερατος, 'cornea,' and *tomos*, 'to cut.' A name given by Wenzel to his knife for dividing the transparent cornea, in the operation for cataract. Many modifications of the instrument have been made since Wenzel's time. See Knife, cataract.

CERATUM, Cerate—*c. Album*, *Ceratum cetaeci*, *Ceratum Galeni*—*c. de Althæa*, *Unguentum de Althæa*—*c. Calamina*, *Cerate of Calamine*—*c. Cantharidis*, *Cerate of Cantharides*, *Emplastrum Lyttæ*—*c. de Cerussa*, *Unguentum plumbi subcarbonatis*.

CERATUM CETAECI, *C. Spermaceti seu album seu ceti seu simplex* (Ph. E.), *Unguentum Cetaeci* (Ph. D.) seu *adipocetæ cetaeum*, *Linimentum album*, *Emplastrum Spermatis Ceti*, *Spermaceti Cerate*, (F.) *Cérat de blanc de baleine*, *Onguent blanc*. (*Spermaceti*, ʒij; *white wax*, ʒiij; *olive oil*, fʒvi. Ph. U. S.) A good emollient to ulcers, &c.

CERATUM CETI, *Ceratum cetaeci*—*c. Cicutæ*, *Ceratum conii*—*c. Citrinum*, *Ceratum resinae*.

CERATUM CONII, *Ceratum Cicutæ*. (*Ung. conii*, ʒij; *cetucci*, ʒij; *cera alba*, ʒiij.) A formula in Bartholomew's Hospital; occasionally applied to cancerous, scrofulous sores, &c.

CERATUM EPULOTICUM, *Cerate of calamine*.

CERATUM GALENI, *C. album seu refrigerans Galeni*, *Unguentum ceratum seu amygdalinum seu simplex*, *Emplastrum ad fonticulos*, *Oleocera-tum aquâ subactum*, *Cold Cream*, (F.) *Cérat blanc ou de GALEN*. (*White wax*, 4 parts; *oil of sweet almonds*, 16 parts; *add, when melted, water or rose-water*, 12 parts. Ph. P.) A mild application to chaps, &c.

CERATUM HYDRARGYRI COMPOSITUM, *Compound Cerate of Mercury*. (*Ung. Hydrarg.*, *Cerat. Sapon. Comp.*, ʒā, ʒvj; *Camphor*, ʒiiss. Rub together. Ph. L.) A discutient to indolent tumours.

CERATUM LABIALE RUBRUM, *Cerate for the lips*—*c. Lapidis calaminaris*, *Cerate of calamine*—*c. Lithargyri acetati compositum*, *Ceratum plumbi compositum*—*c. Lyttæ*, *Cerate of cantharides*—*c. Mercuriale*, *Unguentum hydrargyri*—*c. Picatum*, *Pisseleum*.

CERATUM PLUMBI COMPOSITUM, *C. Lithargyri Acetati Compositum seu Subacetati plumbi medicatum seu Plumbi Subacetatis* (Ph. U. S.) seu *Satur'ni*, *Compound Lead Cerate*, *Goulard's Cerate*, *Goulard's Ointment*, (F.) *Cérat de Goulard*, *C. de Plomb composé*. (*Liq. plumb. subacet.*, ʒiiss; *cera flavæ*, ʒiv; *ol. oliv.*, ʒix; *camphoræ*, ʒss. Ph. U. S.) Its virtues are the same as the next.

CERATUM PLUMBI SUPRACETATIS, *C. Plumbi Acetatis*, *Unguentum Cerusæ Acetatis*, *Cerate of Supracetate or Sugar of Lead*, *Unguentum Acetatis Plumbi*, (F.) *Cérat de sucrétate de Plomb*. (*Acetate of lead*, ʒij; *white wax*, ʒij; *olive oil*, ʒss.) Cooling and astringent.

CERATUM REFRIGERANS GALENI, *Ceratum Galeni*.

CERATUM RESINÆ, *C. Resinæ flavæ seu citrini*, *Unguentum basilicon flavum seu Resinæ flavæ seu Resinosum*, *Resin Cerate or Ointment*, *Yellow Basilicon*, *Basilicon Ointment*. (*Resin. flav.*, ʒv; *Cera flavæ*, ʒij; *Adipis*, ʒviij. Ph. U. S.) A stimulating application to old ulcers, &c. Digestive.

DR. SNELLOME'S Ointment for the Eyes consists of finely powdered *verdigris*, ʒss, rubbed with oil, and then mixed with an ounce of *ceratum resinae*.

CERATUM RESINÆ COMPOSITUM, *Compound Resin Cerate*, (*Resin.*, *Sevi*, *Cera flavæ*, ʒā, ʒij; *Terebinth.*, ʒss; *Ol. Lini*, ʒss. *Deasler's Salve*.

Melt together, strain through linen, and stir till cool. Ph. U. S.)

CERATUM SABINÆ, *Unguentum Sabinæ*, *Savinæ Cerate*, (F.) *Cérat de Sabine*. (*Savinæ*, in powder, ʒij; *Resin Cerate*, ʒij. Ph. U. S.) Irritative, 'drawing.' Used in the same cases as the cerate of cantharides.

CERATUM SAPONIS, *C. Saponis Compositum*, *Soap Cerate*, (F.) *Cérat de Savon*. (*Liq. Plumb. subacetat.*, Oj; *Sapon*, ʒvj; *Cera alba*, ʒx; *Ol. olivæ*, Oj. Boil the solution of subacetate of lead with the soap over a slow fire, to the consistence of honey; then transfer to a water-bath, and evaporate until all the moisture is dissipated: lastly, add the wax, previously melted with the oil, and mix. Ph. U. S.) It is applied in cases of sprains or fractures.

CERATUM SATURNI, *Ceratum plumbi compositum*—*c. Simplex*, *Cerate*, *simple*, *Ceratum cetaeci*—*c. Spermaceti*, *Ceratum cetaeci*—*c. Subacetati plumbi medicatum*, *Ceratum plumbi compositum*—*c. Tetrapharmacum*, *Pisseleum*.

CERATUM ZINCI CARBONATIS, *Cerate of Carbonate of Zinc*. (*Zinci carbonat. preparat.*, ʒij; *Ung. simpl.*, ʒx. Ph. U. S.) Used in the same cases as the *Ceratum calamine*.

CERAUNION, from κεραυνος, 'thunder,' 'a thunderbolt.' *Lapis fulminans*. A kind of stone, which was believed to be formed during thunder; and to be possessed of the power of inducing sleep, and numerous other prophylactic virtues. It was rubbed on the knee, breast, &c., in swellings of those parts.

CERBERA AHOVAL, see *Strychnine*.

CERBERA TANGHIN, *Tanghinia*, *T. venenifera*. A tree of Madagascar, *Order*, *Apocynaceæ*; the kernel of whose fruit is a deadly poison; and although not larger than an almond is sufficient to destroy twenty people.

CERBERA THEVETIA, see *Strychnine*.

CERBERUS TRICEPS, *Pulvis cornachini*.

CERCA'RIA, from κερκος, 'a tail.' A genus of agastrie, caudate, infusory animalcules, one of the most curious of which inhabits the tartar of the teeth. The, so called, spermatozoa have been presumed by some to belong to this genus.

CERCHNASMUS, *Corchnus*.

CERCHNOMA, *Cerchnus*.

CERCHNUS, *Cerchnas'mus*, *Cerchnum*, *Cerchno'ma*, from κερχνω, 'I render hoarse.' A rough voice produced by hoarseness. See *Rattle*.

CERCIFIS, *Tragopogon porrifolium*.

CER'CIS, *kepris*. A sort of pestle for reducing substances to powder. Also, the radius or small bone of the arm. See *Pilum*, and *Radius*.

CEROLE, *Circulus*—*c. de la Choroïde*, *Ciliary ligament*—*c. Cilare*, *Ciliary ligament*.

CERCO'SIS, from κερκος, 'a tail.' *Mentula mulie'bris*, the *Clitoria*. Some authors have employed the word synonymously with *nymphomania* and elongation of the clitoris; and with *Poly-pus Uteri*, the *Sarco'ma Cerco'sis* of Sauvages.

CERCOSIS CLITORIDIS, *Clitorism*—*c. Externa*, *Clitorism*.

CEREA, *Cerumen*.

CEREA'LIA, from CERES, 'goddess of corn' (F.) *Céréales* (*Plantæ*). The cerealia are gramineous plants, the seed of which serve for the nourishment of man:—as wheat, barley, rye, &c. At times, the same term is applied to some of the leguminous plants.

CEREBARIA, *Carebaria*.

CEREBEL'LA URINA. Urine of a whitish appearance, of the colour of the brain or cerebellum, from which Paracelsus thought he could distinguish diseases of that organ.

CEREBELLI'TIS, badly formed from *cerebellum* and *itis*, denoting inflammation. *Parentcephali'tis*, *Inflammati'o cerebel'li*. Inflammation

of the cerebellum: a variety of phrenitis or encephalitis.

CEREBEL'LOUS, *Cerebello'us*, from *cerebellum*, 'the little brain.' (F.) *Cérébelleux*. Chaussier has given this epithet to the vessels of the cerebellum. These are three in number; two of which are *inferior*: the larger, *inferior cerebelli*, which arises from the posterior cerebral or vertebral; and the smaller, whose existence is not constant, from the meso-cephalic or basiliary: —the third, called *A. cerebelleuse supérieure* (*superior cerebelli*), is also a branch of the basiliary.

CEREBEL'LOUS AP'OPLEXY, *Apoplex'ia cerebelli*: apoplexy of the cerebellum.

CEREBEL'LUM, diminutive of *Cerebrum*; *Cerebrum parvum* seu *posterius*, *Appendix ad cerebrum*, *Encaenion*, *Encaenis*, *Epynerod'nis*, *Parvencyph'alis*, *Parvencyph'alus*, *Parvencyph'alum*, *Enceph'illum*, *Enceph'alus opist'hius*, *Micrenceph'alius*, *Micrenceph'alum*, *Little brain* (F.) *Cervelet*. A portion of the medullary mass, contained in the cavity of the cranium. It fills the lower occipital fossæ below the tentorium, and embraces the tuber annulare and medulla. It is composed, like the brain, of vesicular and tubular substance, arranged in laminae, as it were; so that, when a section is made of it, it has an arborescent appearance, called *Arbor vitæ*. The cerebellum is divided into two lobes or hemispheres or lateral masses, *Lobus superior anterior* seu *quadrangul'aris*, and *Lobus superior posterior* seu *semilun'aris*, and each lobe is again subdivided into *Monticuli* or *Lobules*. In the cerebellum are to be observed the *crura cerebelli*, the *fourth ventricle*, the *valcula magna cerebri*, the *processus vermiculares*, superior and inferior, &c.

CEREBRAL, *Cerebra'lis*, (F.) *Cérébral*, from *cerebrum*, 'the brain.' Belonging to the brain; similar to brain.

CEREBRAL APOPHYSIS, Pineal gland.

CEREBRAL ARTERIES are three on each side: —the *anterior* or *artery of the corpus callosum*, and the *middle*, *arteria Sylviana*, are furnished by the internal carotid: —the *posterior* or *posterior and inferior artery of the brain*, *A. profunda cerebri*, arises from the vertebral. Chaussier calls these arteries *lobaires*, because they correspond with the anterior, middle, and posterior lobes, whilst he calls the trunks, whence they originate, *cerebral*.

CEREBRAL NERVES are those which arise within the cranium, (F.) *Nervi craniens*, all of which, perhaps, with the exception of the olfactory, originate from the medulla oblongata. See *Nerves*.

In *Pathology*, an affection is called *cerebral*, which specially occupies the brain. *Fièvre cérébrale*, *Cerebral fever*, is a variety in which the head is much affected.

CEREBRA'TION, *Cerebratio*, from *cerebrum*, 'the brain.' The action of the brain during any mental or moral manifestation.

CEREBRIFORM, *Encephaloid*.

CEREBRITIS, *Phrenitis*.

CEREBROPATHY, see *Nervous diathesis*.

CEREBRO-MALACIA, *Mollities cerebri*.

CEREBRO-RACHIDIAN, *Cephalo-spinal*.

CEREBRO-SPINAL, *Cephalo-spinal*. A *cerebro-spinal* or *cerebro-spinant* is a neurotic, which exercises a special influence over one or more functions of the brain and spinal cord, and their respective nerves. — *Pereira*.

CEREBRO-SPINAL AXIS, see *Encephalon*.

CEREBRO-SPINANT, *Cerebro-spinal*.

CEREBRUM or **CEREBRUM**, from *capa*, 'the head.' The brain. (F.) *Cerveau*, *Cervelle*. This term is sometimes applied to the whole of the contents of the cranium: at others, to the upper portion: —the posterior and inferior being called cerebellum. The brain, properly so called,

extends from the os frontis to the superior occipital fossæ. Anteriorly, it rests on the orbital vault: behind this, on the middle fossæ of the base of the cranium; and, posteriorly, on the *tentorium cerebelli superextensum*. The upper surface is divided by a deep median cleft (*Scissura interlobaire*, — Ch.) into two halves, called *hemispheres*, which are united at the base by the *corpus callosum*. At its surface are numerous *convolutions*. The inferior surface exhibits, from before to behind, three lobes, distinguished into *anterior*, *middle*, and *posterior*. The middle is separated from the anterior by the *fissure of Sylvius*; and from the posterior, by a shallow furrow which corresponds to the upper portion of the *pars petrosa*. Internally, the brain has, on the median line, the *corpus callosum*, *septum lucidum*, *fornix*, *pineal gland*, and *third ventricle*: —and laterally, the *lateral ventricles*, in which are the *corpora striata*, *optic thalami*, &c. It is contained in a triple envelope, (see *Meninges*.) Its texture is pulpy, and varies according to age. Two substances may be distinguished in it — the *white*, *medullary*, *tubular* or *fibrous* — *medulla cerebri*, (F.) *Pulpe cérébral*, and the *cortical*, *cineritious*, *vesicular*, or *gray*. The former is white; and occupies all the interior and base of the brain. The latter is grayish and softer. It is situate particularly at the surface of the organ.

The brain receives several arterial vessels, furnished by the internal carotid and vertebral. Its veins end in the sinuses. It is the material organ of the mental and moral manifestations. According to Gall, each part is 'the special seat of one of those faculties, and the brain and cerebellum, inclusive, are called by him 'the nervous system of the mental faculties.' See *Craniology*.

The substance of the nervous system — *Neurine* — has been analyzed by Vauquelin, and found to contain water, 80·00; white fatty matter, 4·53; red fatty matter, called *ceresine*, 0·70; osmazome, 1·12; albumen, 7·00; phosphorus, 1·50; sulphur, acid phosphates of potass, lime, and magnesia, 5·15.

CEREBRUM ABDOMINALE, *Solar plexus* — c. *Elongatum*, *Medulla oblongata* — c. *Parvum*, *Cerebellum* — c. *Posterius*, *Cerebellum*.

CEREC'TOMY, *Cerecot'mia*, *Kerec'tomy*, from *kepas*, 'cornea,' and *ekronai*, 'excision.' Excision of a layer or portion of the cornea.

CEREFOLIUM, *Scandix cerefolium* — c. *Hispanicum*, *Chærophyllum odoratum* — c. *Sylvestre*, *Chærophyllum sylvestre*.

CERELÆUM, *Cerate*.

CÉRÉOLÉ, *Cerate*.

CEREOLUS, *Bougie*.

CERERISIA, *Cerevisia*.

CERERIUM, *Cerium*.

CEREUM MEDICATUM, *Bougie*.

CEREUS, *Bougie*.

CEREVIS'IA, quasi *Cereris'ia*, *Cervis'ia*, *Ce'lia*, *Zythus*, *Zythum*, *Liquor Cer'eris*, *Oinos xptōns*, *Vinum hordeic'um*, *Bira*, *Bryton*, *Spvtop*, *Barley wine*, from *Ceres*, 'corn;' whence it is made. *Ale* (*Alla*). (Sc.) *Barley-bree*, *Barley-corn*, *Beer*, *Porter*, (F.) *Bière*, *Cervoise*. These fluids are drunk by the inhabitants of many countries habitually, and in Great Britain and Germany more than in others. They are nourishing, but not very easy of digestion. The old dispensaries contain numerous medicated ales, which are no longer in use. Some, however, are still admitted. They are the *Brutolés* or *Brytolés* of Chéreau, *Brytolatures* of Bérail; — *Bières Médicinales*.

CEREVISIA ABIETIS, *Beer*, spruce — c. *Nigra*, see *Falltranck*.

CERFEUIL, *Scandix cerefolium* — c. *Musqué*,

Chærophyllum odoratum—*c. Sauvage*, *Chærophyllum sylvestre*.

CERION, see *Favosus*, *Porrigio favosa*.

CERISIER, *Prunus cerasus*—*c. à Grappes*, *Prunus nalis*—*c. de Virgiate*, *Prunus Virginiana*.

CERUM, *Cerôrium*. A metal which is never found isolated, being usually combined with lanthanum or lantanum and *didymium*. The salts of cerium are supposed to be sedative and tonic. They have been used in dyspepsia in its various forms: the vomiting of pregnancy, &c. Dose of the oxalate or nitrate, one or two grains.

CERNIN, SAINT, WATERS OF. St. C. is a parish in the diocese of St. Flour, Upper Auvergne, France. The water is a chalybeate. It is called *Eau du Combou*.

CERNOS, *Capistrum*.

CEROËNË, *Cerou'nê*, or *Ciron'ê'nê*, *Cerô'num*, from *κeros*, 'wax,' and *ovos*, 'wine.' A plaster composed of yellow wax, mutton suet, pitch, Burgundy pitch, bals *Armeniac*, thus and wine. It was used as a strengthening plaster. Sometimes it contained neither wax nor wine.

CEROLEUM, *Cerate*.

CEROMA, *Cerate*. See *Lardaceous*.

CEROMANTIA, from *κeros*, 'wax,' and *μαντεια*, 'divination.' The art of foretelling the future, from the figures which melted wax assumes, when suffered to drop on the surface of water.

CERONTUM, *Cerate*.

CEROPIS'SUS, from *κeros*, 'wax,' and *πισσα*, 'pitch.' A depilatory plaster, composed of pitch and wax.

CEROS, *Cera*.

CEROSTROSIS, *Hystriasis*.

CEROTUM, *Cerate*.

CEROXYLON, see *Cera flava et alba*.

CERUA, *Ricinus communis*.

CERUMEN, from *cera*, 'wax,' *ἄ. Aurium*, *Ce'ra*, *Aurium Sordes*, *Sordic'ula aurium*, *Marmora'ta Aurium*, *Cypselê*, *Cerum'inos* *Humour*, *Ear-wax*, *Cyp'selis*, *Pu'gile*, (F.) *Cire des Oreilles*. A name given to the unctuous humour, similar to wax in its physical properties, which is met with in the *meatus auditorius externus*. It is secreted by glands, situate beneath the skin lining the meatus. It lubricates the meatus, preserves the suppleness of the lining membrane, prevents the introduction of bodies floating in the atmosphere, and by its bitterness and unctuousness prevents insects from penetrating.

CERUMINOUS, *Cerum'inosus*, (F.) *Cérumineux*. Relating to cerumen.

CERUMINOUS GLANDS, *Cerum'inosus Follicles*, (F.) *Glandes ou Follicules cérumineuxes*. Glands or follicles which secrete the cerumen.

CERUSE, *Plumbi subcarbonas*.

CERUSSA ACETATA, *Plumbi superacetas*—*c. Alba Hispanica*, *Plumbi subcarbonas*—*c. Alba Norica*. *Plumbi subcarbonas*—*c. Nigra*, *Graphites*—*c. Psymmithron*. *Plumbi subcarbonas*—*c. Serpentina*, see *Arum maculatum*.

CERUSSE, *Plumbi subcarbonas*.

CERUSSEA URINA. A term used by Paracelsus for the urine, when of a colour like cerusse.

CERVARIA ALBA, *Laserpitium latifolium*.

CERVEAU, *Cerebrum*.

CERVELET, *Cerebellum*.

CERVELLE, *Cerebrum*.

CERVI BOLETUS, *Elaphomyces Granulatus*.

CERVI ELAPHI CORNU, *Cornu cervi*, see *Cervus*.

CERVICAL, *Cervica'lis*, *Trache'lian*, (F.) *Trachélien*, from *cervix*, *cervicis*, the neck, 'the back of the neck.' Every thing which concerns the neck. *Cervi'cal*, in antiquity, meant a pillow or bolster.

CERVICAL ARTERIES are three in number: 1.

The ascending, anterior, or superficial, a branch of the inferior thyroïd, distributed to the scutular muscles and integuments. 2. The transverse (*Cervico-scapularis*—(Ch.) a branch of the axillary artery, or of the subclavian; distributed to the levator scapulae, trapezius, &c. 3. The posterior or profound. A. *transversa'lis colli*, *Trachilo-cervicale*—(Ch.) a branch of the subclavian, distributed to the deep-seated muscles on the anterior and posterior parts of the neck. See, also, *Princeps Cervicis (arteria)*.

CERVICAL GANGLIONS. The three ganglia of the great sympathetic. The cervical glands or lymphatic glands of the neck are, also, so called. See *Triplanchnic nerve*.

CERVICAL LIG'AMENTS. These are two in number. 1. The anterior, extending from the basilar process of the occipital bone to the anterior part of the first cervical vertebra. 2. The posterior or *supraspin'ous*, *Ligamentum Nuchæ*, which extends from the outer occipital protuberance to the spinous process of the seventh cervical vertebra. In animals with large heads it is very strong.

CERVICAL NERVES are eight in number on each side, and form the eight cervical pairs, which are the first given off from the spinal marrow.

CERVICAL PLEXUS, *Plexus Trachelo-scapularis* (Ch.) The nervous network formed by the anterior branches of the first three cervical nerves, above the posterior scutular muscle, and at the outer side of the pneumogastric nerve, carotid artery, and jugular vein.

CERVICAL REGION, ANTERIOR DEEP, *Prevertebral region*. The region of the neck, occupied by three pairs of muscles placed immediately in front of the cervical and three superior dorsal vertebrae—viz. the rectus capitis anticus major the rectus capitis anticus minor, and longus colli;—hence termed *prevertebral muscles*.

CERVICAL VEINS have nearly the same distribution as the arteries.

CERVICAL VERTEBRÆ. The first seven vertebrae of the spine.

CERVICALIS DESCENDENS, see *Hypoglossus* and *Sacro-lumbalis*.

CERVICARIA, *Campanula trachelium*.

CERVICO-DORSO-SCAPULAIRE. Rhomboideus—*c. Dorsocostal*, *Serratus posticus superior*—*c. Dorsocostoidien et dorso-trachélien*, *Splenius*—*c. Mastoïdien*, *Splenius*.

CERVICO-FACIAL, *Cervi'co-facialis*. Belonging to the neck and face.

CERVICO-FACIAL NERVE, *Nervus cervico-facialis*. A branch of the facial nerve, distributed to the neck and face.

CERVISIA, *Cerevisia*.

CERVISPINA, *Rhamnus*.

CERVIX, *Collum*. The neck, especially the back part. A neck.

CERVIX OBSTIPA, *Torticollis*—*c. Uteri*, *Collum uteri*.

CERVOISE, *Cerevisia*.

CERVUS, *Cerva*. The horn of the *Cervus Elaphus*, called *Cornu*, *Cervi Elaphi Cornu*, *Cornu Cervi'num*, *Hartshorn*. (F.) *Corne de cerf*, contains 27 parts of gelatin in the 100. A jelly made from the shavings is emollient and nutritive.

The Stag's Pizzle, *Præputius Cervi*, was once considered to be aphrodisiac. Dose, ʒj to ʒi, in powder.

CERVUS ALCES. The Elk, (F.) *Élan*. The hoof of this animal was anciently used as an anti-epileptic. The animal, it was asserted, was subject to attacks of epilepsy, and always cured them by putting its hoof into the ear. The hoof was also worn as an amulet.

Cornu Ustum, *Burnt Hartshorn*, has been used as an antacid, but it consists of 57 parts of phos-

phate, and only one of carbonate of lime. It is, therefore, not of much use.

CÉSARIENNE OPÉRATION, Cæsarean section.

CESSATIO MENSIIUM, Amenorrhœa.

CESTRITES. A wine prepared from betony, (*κεστρύου*, 'betony'.)

CESTRON, *Betonica officinalis*.

CETA'CEUM, from *κῆτος*, *cetus*, 'a whale.' *Album Ceti*, *Adipocera cetacea*, *Stercorium cetaceum*, *Spermaceti*, *Parmaceti*, *Sperm.* (F.) *Blanc de Baleine*, *Cétine*, *Adipocire de Baleine*. An inodorous, insipid, white, crystallized, friable, unctuous substance, obtained from the brain of the *Physalus Macrorhynchus* or *Spermaceti Whale*, (F.) *Cachalot*, and other varieties of whale. S. g. 9433; melts at 112°. It is demulcent and emollient, and has been given in coughs and dysentery, but is mostly used in ointments. Dose, ʒss to ʒiiss, rubbed up with sugar or egg.

CETERACH OFFICINARUM, *Asplenium ceterach*.

CÉTINE, *Cetaceum*.

CETRARIA ISLANDICA, *Lichen islandicus*.

CETRARIIN, see *Lichen islandicus*.

CETRARIUM, see *Lichen islandicus*.

CEVADILLA, see *Veratrum sabadilla*.—c. Hispanorum, *Veratrum sabadilla*.

CÉVADILLE, *Veratrum sabadilla*.

CHAA, *Thea*.

CHACRILLE, *Croton cascarilla*.

CHADLOCK, *Sinapis arvensis*.

CHEREFOLIUM, *Scandix cerefolium*.

CHEROPHYLLUM, from *χαίρω*, 'I rejoice,' and *φύλλον*, 'a leaf,' owing to its luxuriant foliage. *Scandix cerefolium*.

CHEROPHYLLUM CEREFOLIUM, *Scandix cerefolium*.—c. *Angulatum*, *Ch. sylvestre*.

CHEROPHYLLUM ODORATUM, *Scandix seu Cicutaria Odorata*, *Myrrhis Odorata* seu *Major*, *Cerefolium Hispanicum*, *Sweet Cic'ely*, (F.) *Cerfeuil musqué ou d'Espagne*; *Order*, *Umbelliferae*; has the smell of aniseed, and is cultivated on account of its aromatic properties.

CHEROPHYLLUM MONOGYNUM, *Ch. sylvestre*.—c. *Sativum*, *Scandix*.

CHEROPHYLLUM SYLVESTRÈ, *Cicutaria*, *Cherophyllum temulum* seu *monogynum* seu *angulatum* seu *verticillatum*, *Anthriscus humilis* seu *procerus*, *Cerefolium Sylvestre*, *Bastard Hemlock*, *Wild Chervil* or *Cow-weed*, (F.) *Cerfeuil sauvage*, *Perail d'Ane*, is a slightly fetid aromatic, but is not used.

CHEROPHYLLUM TEMULUM, *Cherophyllum sylvestre*.—c. *Verticillatum*, *Ch. sylvestre*.

CHETE, *Capillus*.

CHAFING, *Erythema Intertrigo*, *Intertrigo*, *Paratrimma*, *Paratrimma*, *Diatripsia*, *Diatripsia*, *Attritia*, *Attritio*, from (F.) *exchauffer*, *échauffer*, 'to heat'; *Fret*, *Gall of the skin*; *Attritua*, (F.) *Échauffement*, *Écorchures*. The red excoriations which occur in consequence of the friction of parts, or between the folds of the skin, especially in fat or neglected children. Washing with cold water and dusting with hair-powder is the best preventive. When occurring between the nates and in the region of the perineum, from long walking,—*Intertrigo podicis*, *Proctalgia intertriginosa*,—it is vulgarly designated by the French *Entrefesson*.

CHAFFBONE, *Jawbone*.

CHAIR, *Flesh*.

CHAIR, OBSTETRIC, *Labour chair*.

CHAISE PERCÉE, *Lissanum*.

CHALASIS, *Relaxation*.

CHALASMUS, *Relaxation*.

CHALASTICUS, from *χαλασ*, 'I relax.' A

medicine proper for removing rigidity of the fibres.—Galen. An emollient or relaxant.

CHALAZA, *Chalazion*, *Chalazion*, *Chalaziosis*, *Porosis*, *Grando*, *Tophus*, *Hail*, (F.) *Grêle*, *Gravelle*, from *χάλαζα*, 'hail.' A hard, round, transparent tumour, developed in different parts of the body, more especially in the substance of the eyelids.—Also, the *Cicatricula* of the egg. Generally, however, in the language of oöologists, the *chalaza* or *pole* are, in the egg of the bird, the more dense internal layer of the albumen, which adheres to the yolk, and is continued, in the form of two spirally twisted bands, towards the extremities of the egg. The twisting is considered to be produced by the revolving motion of the egg in its descent through the oviduct.

CHALAZÆ, see *Chalaza*.

CHALAZONEPHRITIS, *Kidney*, *Bright's disease* of the.

CHALCANTHUM, *Atramentum*, *Ferri sulphas*.—c. *Album*, *Zinci sulphas*.

CHALCEDONIUS, *Cornelian*.

CHALCITAS, *Bismuth*.

CHALCITES, *Coleothar*.

CHALCOIDEUM (os), *Cuneiform bone*.

CHALCOS, *Brass*, *Cuprum*, *Æreolum*.

CHALCUS, *Æreolum*.

CHALÈRE, *Heat*.—c. *Acere*, see *Acrid*.—c. *Animale*, *Animal heat*, see *Heat*.

CHALEURS DES ANIMAUX, see *Heat*.—c. *du Foie*, *Heat*.

CHALINI, see *Lip*.

CHALINOPLASTY, *Chalinoplastice*; from *χαλινος*, 'frenum,' 'a bridle,' and *πλασσω*, 'I form.' The operation for forming a new frenum.

CHALK, *Creta*.—c. *Red*, *Rubrica febrilis*.—c. *Stones*, *Calculi*, *arthritis*.

CHALL, *Maxillary bone*.

CHALYBÆ, *Chalybeate*.

CHALYBÆATE, *Chalybeatus*, *Ferruginus*, *Ferruginosus*, *Ferratus*, *Martialis*, *Ferruginous*, *Martialis*, (F.) *Ferrugineux*, *Chalybæ*, from *chalybs*, 'iron or steel.' Of, or belonging to iron; containing iron. Any medicine into which iron enters, as *chalybeate mixture*, *pills*, *waters*, &c. See *Waters*, *Mineral*.

CHALYBIS RUBIGO, *Ferri subcarbonas*.

CHALYBOKRENÆ, *Water*, *mineral*, (*chalybeate*.)

CHALYBOPEGÆ, *Water*, *mineral*, (*chalybeate*.)

CHALYBS, from *Chalybes*; a people of Pontus, who dug iron out of the earth; *Acies*, *Sæcl*. The *Proto-carburet of iron*, (F.) *Acier*. As a medicine, steel does not differ from iron.

CHALYBS TARTARIZATUS, *Ferrum tartarizatum*.

CHAMA, *Cheme*.

CHAMÆACTE, *Sambucus ebulus*.

CHAMÆBATUS, *Fragaria*.

CHAMÆCEDRIS, *Artemisia santonica*.

CHAMÆCISSUS, *Glechoma hederacea*.

CHAMÆCLEMA, *Glechoma hederacea*.—c. *Hederaceæ*, *Glechoma hederacea*.

CHAMÆCYPARISSUS, *Artemisia santonica*.

CHAMÆDROPS, *Teucrium chamædrys*.

CHAMÆDRYTES. A wine, in which the *Teucrium Chamædrys* has been infused.

CHAMÆDRYS, *Rubus chamaemorus*, *Teucrium Ch.*, *Veronica*.—c. *Incana maritima*, *Teucrium marum*.—c. *Marum*, *Teucrium marum*.—c. *Minor repens*, *Teucrium Ch.*.—c. *Palustris*, *Teucrium scordium*.—c. *Scordium*, *Teucrium scordium*.—c. *Vulgaris*, *Teucrium Ch.*

CHAMÆGEIRON, *Tussilago*.

CHAMÆLEAGNUS, *Myrica gale*.

CHAMÆLAÏTES. A wine impregnated with *Chamelea*, *Daphnè Alpinæ*.

CHAMÆLEA, *Cneorum tricoceum*.

CHAMÆLEON ALBUM. *Carlina aculis*.

CHAMÆLEUCE, Tussilago.

CHAMÆLINUM, *Linum catharticum*.

CHAMÆLIRIUM LUTEUM, *χάμα*, 'on the ground,' and *lagon*, 'a lily,' [?] *l'cal'trum luteum*, *Helo'nias lutea* seu *Dio'ca*, *Devil's Bit*, *Blazing star*, *Unicorn root*; indigenous: Order, *Melanthaceae*; flowering in June; is acrid. An infusion of the root has been given as an anthelmintic: a tincture, as a tonic.

CHAMÆMELUM, *Anthemis nobilis*—c. *Fœtidum*, *Anthemis cotula*—c. *Nobile*, *Anthemis nobilis*—c. *Oloratum*, *Anthemis nobilis*—c. *Vulgare*, *Matricaria chamomilla*.

CHAMÆMORUS, *Teucrium chamæpitys*, *Rubus chamæmorus*—c. *Norwegica*, *Rubus chamæmorus*.

CHAMÆPEUCE, *Camphorosma Monepeliaca*.

CHAMÆPITUINUM VINUM. A wine, in which the leaves of the *Chamæpitys*, *Teucrium Chamæpitys*, have been infused.

CHAMÆPITYS, *Teucrium chamæpitys*—c. *Anthyllus*, *Teucrium iva*—c. *Moschata*, *Teucrium iva*.

CHAMÆPLTON, *Erysimum*.

CHAMÆRAPHANUM, from *χαμαί*, 'on the ground,' and *ραφανός*, 'the radish.' So Paulus of Ægina calls the upper part of the root of the *Apium*.

CHAMÆTROPIS SERRATULA, *Sarc Palmetto*, from *χαμαί*, 'on the ground,' and *πύλη*, 'a shrub.' A farina is prepared from the roots of this plant, which is used by the Indians, in Florida, as diet.

CHAMBAR, *Magnesia*.

CHAMBER, *Cam'era*, (F.) *Chambre*. A term used in speaking of the eye, in which there are two chambers, *Cam'era oculi*:—an anterior and a posterior; (F.) *Chambre antérieure et postérieure*. The anterior is the space between the cornea and the anterior part of the iris:—the posterior, the space between the iris and anterior surface of the crystalline. They are filled with the aqueous humour, and communicate by the opening in the pupil.

CHAMBERLAIN'S RESTORATIVE PILLS. This nostrum, recommended in scrofula, and all impurities of the blood, has been analyzed by Dr. Paris, and found to consist of cinnabar, sulphur, sulphate of lime, and a little vegetable matter. Each pill weighs 3 grains.

CHAMBRE, Chamber.

CHAMBRIE, *Cannabis sativa*.

CHAMELEA, *Daphne Alpina*.

CHAMOMILE DOGS', *Matricaria chamomilla*—c. *Dyers'*, *Anthemis tinctoria*—c. *German*, *Matricaria chamomilla*—c. *Spanish*, *Anthemis pyrethrum*—c. *Stinking*, *Anthemis cotula*—c. *Wild*, *Anthemis cotula*, *Matricaria glabrata*.

CHAMOMILLA FETIDA, *Anthemis cotula*—c. *Nostras*, *Matricaria chamomilla*—c. *Romana*, *Anthemis nobilis*—c. *Spuria*, *Anthemis cotula*.

CHAMPACA, *Michelia champaca*.

CHAMPIGON, Fungus—c. *de l'Appareil des Fractures*, *Clavaria*—c. *de Couché*, see *Agaric*—c. *du Ferment*, *Torula cerevisia*—c. *de Malté*, *Cynomorion coccineum*—c. *du Muguet*, *Aplthaphyte*.

CHAMPOOING, Shampooing.

CHANCEBONE, Ischion.

CHANCELAGUA, *Canchalagua*.

CHANCRE (F.), *Shanker*, *Ulcus seu Ulcus enlum cancerosum*. A sore, which arises from the direct application of the venereal virus; hence it is almost always seated, in men, on the penis. The French use the word *Chancre*, in popular language, for cancerous ulcers, the malignant aphthæ of children, &c. Formerly, the terms

Car'oli and *Ca'ries pudendo'rum* were used for venereal pustules or sores on the parts of generation.

CHANCRE LARVÉ. A concealed chancre, such as has been supposed by M. Ricord to give occasion to gonorrhoea virulenta.

CHANCREUX, *Chancreuse* (F.), *Cancro'sus*, *Carcino'des*. Having the nature of chancre, or of cancer.

Bouton Chancreux. A small tumour of a cancerous nature, which makes its appearance on the face—most frequently on the upper lip—*Noli me tangere* [?].

CHANDOO. An aqueous extract of opium used by the Chinese for smoking.

CHANGE OF LIFE, see *Menses*.

CHANNELBONE, Clavicle.

CHANT DES ARTERES, *Sifflement modulé*.

CHANVRE, *Cannabis sativa*—c. *Bâtard*, *Galeopsis grandiflora*—c. *Indien*, *Bangue*.

CHAOMANTIA. The alchemists meant, by this word, the art of predicting the future from observation of the air. The word *Chaos* was used by Paracelsus for the air; (*μαντεια*, 'divination'.)

CHAOSDA, Plague.

CHAPERON DE MOINE, *Aconitum napellus*.

CHAPETONNADE, *Chappetonnade*.

CHAPITEAU, Alembic.

CHAPPEDONADE, *Chappetonnade*.

CHAPPETONADE, *Chappetonnade*, *Chapetonnade* (F.), *Vom'itus rabio'sus*. Vomiting, accompanied by furious delirium, attacking strangers in hot countries.

CHAPS, Rhagades.

CHARACTER, *χαρακτρη*, 'a mark or impression.' (F.) *Caractère*. In Pathology it is used synonymously with stamp or appearance. We say, "A disease is of an unfavourable character," &c. The prevailing epidemic has a bilious character," &c. In *Mental Philosophy* it means—that which distinguishes one individual from another, as regards his understanding and passions. See *Symbol*.

CHARADRIUS. Ælian thus calls a bird, which was reputed to cure jaundice. The word now means the plover.

CHARANTIA, *Momordica elaterium*.

CHARBON, *Anthrax carbo*, *Carbo ligni*—c. *Animal*, *Carbo animalis*—c. *Minéral*, *Graphites*—c. *Pulmonaire*, *Anthracoësis*.

CHARBONNEUX, *Anthraxoid*.

CHARCOAL, *Carbo*—c. *Animal*, *Carbo animalis*.

CHARDON AUX ANES, *Onopordium acanthium*—c. *Béni*, *Centaurea benedicta*—c. *à Bonnetier*, *Dipsacus fullonum*—c. *Étoilé*, *Centaurea calcitrapa*—c. *à Foulon*, *Dipsacus fullonum*—c. *Hémorrhoidal*, *Cirsium arvense*—c. *Marie*, *Carduus Marianus*—c. *Roland*, *Eryngium campestre*.

CHARLATAN, from Ital. *ciarlatano*, 'to talk much,' *Circulato'ry*, *Circumfora'te'us*, *Periodontes*, *Pseudomed'icus*, *Aggr'ta*, *Anag'ellon*, a Quack, Quacksalber, Empirical Pretender, Empiric. Originally, one who went from place to place to sell a medicine, to which he attributed marvellous properties. By extension—any individual who endeavours to deceive the public by passing himself off as more skillful than he really is. According to Ménage, the word comes from *circulatus*, a corruption of *charlatan*.

CHARLATANRY, *Aggr'tia*, *Banaus'tia*, the conduct or action of a charlatan. (F.) *Charlatanerie*, *Charlatanisme*, *Quackery*, *Empiricism*.

CHARLOCK, *Sinapis arvensis*.

CHARLOTTEVILLE SULPHUR SPRING, see *Canada*, *Mineral Waters* of.

CHARM, Lat. *Carmen*, 'a verse,' because charms often consisted of verses, whence comes the Italian, *Giarma*, (F.) *Charme*, with the same signification; *Canta'tio*, *Incantamen'tum*. A trick, a spell, an enchantment. A sort of magic, or superstitious practice, consisting of words, characters, &c., by which, it was believed, individuals might be struck with sickness or death, or be restored to health.

The following are specimens of old charms—*verse charms*—

For stanching Blood, (Pepys.)

Sanguis mane in te
Sicut Christus fuit in se;
Sanguis mane in tuâ venâ;
Sicut Christus in suâ penâ;
Sanguis mane fixus,
Sicut Christus quando fuit crucifixus.

For Cramp, (Pepys.)

Cramp be thou faintless,
As our lady was sinless,
When she bare Jesus.

For the Foot when asleep, (Coleridge.)

Foot! foot! foot! is fast asleep!
Thumb! thumb! thumb! in spittle we steep;
Crosses three we make to ease us,
Two for the thieves, and one for Christ Jesus.

The same charm served for cramp in the leg, with this substitution:

The devil is tying a knot in my leg!
Mark, Luke, and John, unloose it, I beg!—
Crosses three, &c.

For a Burn, (Pepys.)

There came three angels out of the East;
The one brought fire, the other brought frost.
Out fire! in frost.
In the name of the Father and Son and Holy Ghost.
Amen.

CHARNIÈRE, Ginglymus.

CHARNU, Carneous.

CHARPENTE OSSEUSE, Skeleton.

CHARPIE, Lintenum—c. *Brite*, see Lintenum—c. *Rupie*, see Lintenum.

CHARTA ANTARTHRITICA, Gout paper—c. *Antirheumatica*, Gout paper—c. *Vesicatoria*, see *Sparadrapum Vesicatorium*—c. *Virginica*, Amnios.

CHARTRE, Tabes mesenterica.

CHAN (F.), *Acus fora'tum*. The eye of a needle. Sometimes, this opening is near the point of the instrument, as in the ligature needle.

CHASCHISCH, see Bangle.

CHASME, Yawning.

CHASPE, Variola.

CHASSE (F.), *Mam'mbrum*. A kind of handle composed of two movable laminæ of horn, shell, or ivory, united only at the extremity, which holds the blade of the instrument,—as in the common bleeding lancet. Also, a *liqueur* or alcoholic liquor, taken after a meal; hence *chasse-café*, *chasse-tout*, &c., from (F.) *chasser*, 'to drive,' because presumed to aid digestion.

CHASSIE (F.), *Lema*, *Lippa*, *Glama*, *Glemë*, *Gra'mia*, *Lenos'tus*, *Sebum palpebra'le*; the gum of the eye. (Prov.) *Gound* or *Gownde*, from *chasser*, 'to drive out.' A sebaceous humour, secreted mainly by the follicles of Meibomius, which sometimes glues the eyelids together.

CHASSIEUX (F.), *Lippus*; covered with *Chassie*—as *Paupièrès chassieuxes*.

CHASTE TREE, Vitex.

CHÂTAIGNE, see *Fagus castanea*—c. *d'Eau*, *Truva natans*.

CHÂTAIGNIER COMMUN, *Fagus castanea*—c. *Nain*, *Fagus castanea pumila*.

CHATEAU-LANDON, WATERS OF. A town three leagues from Nemours, in France. The waters contain alum and iron.

CHATEAU-SALINS, WATERS OF; a town in the department of La-Meurthe, France. The waters contain carbonate of lime, sulphates of lime and magnesia, and chlorides of magnesium and sodium.

CHATELDON, MINERAL WATERS OF. Chateldon is in the department of Puy-de-Dôme, France. The waters contain carbonic acid and iron.

CHATEL-GUYON, MINERAL WATERS OF. A village in France, in the department Puy-de-Dôme, near which there are five thermal acidulous springs. Temperature, 86° Fahrenheit.

CHATON (F.), 'a husk.' In pathology, it means a *funda* or cavity formed by the irregular or *hour-glass* contraction of the uterus, in which the placenta is often retained or *enchatonné* after the birth of the child. It is detected by passing the fingers along the cord as far as the part which is contracted, when the placenta will not be discoverable.

The treatment consists in relaxing by a large dose of an opiate, then passing the fingers along the cord, and gradually dilating the opening through which it passes, until it permits the hand to go through. The placenta must then be grasped and gently withdrawn.

CHATON, Vaginal process.

CHATONNÉ, *Incarcera'tus*, from *chaton*. An epithet in French for anything shut up in a sac, or imprisoned in any manner—as

CHATONNÉ, CALCUL (F.), *Cal'culus incar-cera'tus*, *Encyst'ed Cal'culus*, *Calcul enkysté*. A urinary calculus, adherent to the inner surface of the bladder, so that it is immovable, and cannot pass to the different parts of that organ. This happens when calculi form in some natural or accidental cavity of the bladder; or when the organ, by ulceration, gives rise to fungi, which surround the calculus; or when it is lodged in the orifice of the ureter or urethra.

CHATONNÉ, PLACENTA (F.). The placenta when retained as above described. See *Chaton*.

CHATONNEMENT (F.), *Enchatonnement*, *Incarcera'tio*, *Chatonnement du placenta*, *Enkys-tement*.—Hour-glass contraction of the uterus. See *Chaton*.

CHATOUILLEMENT (F.). This word sometimes means the action of tickling or titillation (*Titilla'tio*), and, at others, the sensation which gives rise to the action (*Pruri'tus*), Itching.

CHÂTRÉ, Castratus.

CHÂTRURE, Castration.

CHAUDEAU, Caudle.

CHAUDEBOURG, MINERAL WATERS OF. C. is three quarters of a league from Thionville, in France. The waters contain iron, sulphate of lime, sulphate of magnesia, and carbonate of lime.

CHAUDEPISSE, Gonorrhœa impura—c. *Cor-dée*, Gonorrhœa cordata—c. *Tombée dans les Bourses*, Hernia humoralis.

CHAUDS-AIGUES, MINERAL WATERS OF. A small town in the department of Cantal, France, where there is a number of saline springs containing carbonic acid, carbonate of soda, and chloride of sodium. Temp. 190° Fahrenheit.

CHAUDIÈRE, see Alembic.

CHAUFFOIR (F.), *Lintenum Calefacto'rium*. According to the *Académie*, a warmed cloth, used either for the purpose of warming a patient, or to apply to a female recently delivered.

CHAULE, Maxillary bone.

CHAULMOOGRA. The fruit of the *Chaulmoogra*, *Gynocarpia odorata*. Order, Capparidæ, of India. A bland fixed oil is obtained by expressing the dried seed, which is used by the natives in leprosy. The seed are given internally, in the dose of about 3ss in the day, and the eruption is rubbed with the oil.

CHAUSSE (F.), *Chausse d'Hippocrate*, *Manche d'Hippocrate*, *Mou'ica Hippocratias*, *Mou'ica Hippocrates* Steece. A conical bag, made of flannel, for straining liquids.

CHAUSSE-TRAPPE, *Centaurea calcitrapa*.

CHAUVETÉ, Calvities.

CHAUX, Calx—*c. Bichlorure de*, *Calcis chloridum*—*c. Chlorure de*, *Calcis chloridum*—*c. Hydrochlorate de*, *Calcis murias*—*c. Hydrosulfate de*, *Calcis sulphuretum*—*c. Muriate de*, *Calcis murias*—*c. Muriate origéné de*, *Calcis chloridum*—*c. Muriate suavisigéné de*, *Calcis chloridum*—*c. Orichlorure de*, *Calcis chloridum*—*c. Oximiniate de*, *Calcis chloridum*—*c. Phosphate de*, see *Cornu cervi*—*c. Vine*, Calx viva.

CHAVEL, Maxillary bone.

CHAVICA BETLE, Betel—*c. Roxburghii*, *Piper longum*.

CHAWBONE, Maxillary bone.

CHAWD PYS, Gonorrhœa impura.

CHAWING, Mastication.

CHECKER-BERRY, *Ar utus uva ursi*.

CHEEK, Gena—*c. Teeth*, Molar teeth.

CHEESE, Sax. *cere*, (L.) *Caseus*, *Tyros*, *Pectē*, (Sc.) *Furmage*, (F.) *Fromage*. An aliment, prepared from the caseous and oleaginous parts of milk. Fresh cheeses owe their chief medical properties to the immediate principle, essentially cheesy, to which the name *caseini* or *casein* has been applied. Those, which have been recently salted, are digested with comparative facility. The flavour of cheese is owing to an ammoniacal caseate. On the whole, cheese itself is not easy of digestion, although it may stimulate the stomach to greater exertion, and thus aid in the digestion of other substances.

CHEESE RENNET, *Galium verum*.

CHEE'SY, *Casearius*, *Caseosus*, *Tyro'des*, (F.) *Caseux* ou *Caseux*. Having the nature of cheese.

CHEF, Bandage (tail).

CHEGÔE, *Chique*.

CHEGRE, *Chique*.

CHEILY'TIS, *Chil'y'tis*, from *χειλος*, 'a lip.' Inflammation of the lip. See *Chilon*.

CHEILO'CACE, from *χειλος*, 'a lip,' and *κακος*, 'evil,' *Labrisul'eium*. A disease, characterized, it is said, by swelling, induration, and slight redness of the lips without inflammation; reputed, but without any authority, to be common in England and Scotland, amongst children. Also, the thickness of the upper lip of scrofulous children. See *Stomacæ*, and *Cancer aquaticus*.

CHEILOCARCINO'MA, from *χειλος*, 'a lip,' and *καρκινωμα*, 'a cancer.' Cancer of the lip.

CHEILOMALACIA, *Cancer aquaticus*, *Stomacæ*.

CHEILON, *Chilon*.

CHEILON'CUS, *Cheilophy'ma*, from *χειλος*, 'lip,' and *ογκος*, 'swelling.' A swelling of the lip.

CHEILOPHYMA, *Cheiloncus*.

CHEILOPLAS'TICÊ, *Chiloplas'ticê*, *Cheiloplasty*, from *χειλος*, 'lip,' and *πλαστικός*, 'forming.' The operation for an artificial lip.

CHEILOS, Lip.

CHEIMA, Cold.

CHEIMETLON, Chilblain.

CHEIMIA, Rigor.

CHEIR, Manus, see *Chir*.

CHEIRANTHUS CHEIRI, from *χειρ*, 'the hand,' and *ανθος*, 'flower,' a flower for the hand.

The systematic name of the *Common Yellow Wall Flower*, *Viola lutea*, *Lonicum luteum*, *Koyel*, *Cheiri*, (F.) *Gérolée*, *Gérolée* ou *Violet jaune*, *Ord. Cruciferae*. The flowers have been esteemed nervine, narcotic, and deobstruent.

CHEIRAP'SIA. The action of rubbing or scratching, from *χειρ*, 'the hand,' and *πρω*, 'I touch.' A troublesome symptom in the itch.

CHEIRARTHRO'CACÊ, from *χειρ*, 'the hand,' *αρθρον*, 'a joint,' and *κακία*, a 'disease.' Inflammation of the articular surfaces of the wrist.

CHEIRIATER, Surgeon.

CHEIRIS'MA, *Cheiris'mus*. The act of touching—handling. Any manual operation.

CHEIRIXIS, Surgery.

CHEIRONOM'IA, *Chironom'ia*, from *χειρ*, 'I exercise with the hands.' An exercise, referred to by Hippocrates, which consisted in using the hands, as in our exercise of the dumb-bells.

CHEIROPLETHES, Fasciculus.

CHEIROSIS, Subactio.

CHELA, *Chela*. This word has several significations. *Chela*, a forked probe used for extracting polypi from the nose. *Chela*—chaps, or cracks on the feet, organs of generation, &c. *Chela* likewise means claws, especially those of the crab. See *Cancerum Chela*.

CHELE PALPERARUM, see *Tarsus*.

CHELAPA, *Convolvulus jalapa*.

CHELE, *Chela*.

CHELIDOINE GRANDE, *Chelidonium majus*—*c. Petite*, *Ranunculus ficaria*.

CHELIDON, *χελιδων*, the hollow at the bend of the arm; *Hirundo*.

CHELIDONIA ROTUNDIFOLIA MINOR, *Ranunculus ficaria*.

CHELIDONTIUM, *Bryonia alba*—*c. Diphylum*, *Stylophorum diphylum*—*c. Hamatodes*, *Ch. majus*.

CHELIDONIUM MAJUS, *Ch. hamatodes*, from *χελιδων*, 'a swallow,' because its flowering coincides with the appearance of the swallow. *Papaver Corniculatum* seu *luteum*, *Common Chelidone*, *Tetterwort*, (F.) *Chélidoine grande*, *L'Éclaire*, *Herbe à l'hirondelle*. *Ord. Papaveraceæ*. *Sex. Syst.* Polyandria Monogynia. The root and recent plant have been considered aperient and diuretic. Externally, the juice has been employed in some cutaneous diseases.

CHELIDONIUM MINUS, *Ranunculus ficaria*.

CHELIDONIUS LAPIS. A name given to stones, which, it was pretended, existed in the stomach of young swallows. They were formerly believed capable of curing epilepsy.

CHELOID, *Canceroid*.

CHELOIS, see *Canceroid*.

CHELO'NE, *χελων*, 'a tortoise.' An instrument for extending a limb; so called, because in its slow motions it resembled a tortoise.—*Oribasius*. See *Testudo*.

CHELO'NE GLABRA, *Common Snake head*, *Turtle head*, *Turtle bloom*, *Shellflower*, *Balmoney*. An indigenous plant. *Sex. Syst.* Didynamia Angiospermia; blossoming from July to November. The leaves are bitter and tonic; without any aromatic smell, and with very little astringency.

CHELO'NIA MYDAS, *The Green Turtle*. This species of turtle abounds on the coast of Florida. It is the one so prized by the epicure.

CHELO'NION, *Chelo'nium*, from *χελων*, 'a tortoise,' from its resembling in shape the shell of that animal. The upper, gibbous part of the back.—*Gorræus*. The scapula.

CHELONOPH'AGI, from *χελων*, 'the tortoise,' and *φαγω*, 'I eat.' An ancient name for certain tribes, who dwelt on the coast of the Red

Sea, and who lived only on tortoises.—Pliny, Diodorus of Sicily.

CHELSEA PENSIONER. An empirical remedy for rheumatism and gout, sold under this name. (*Gum guaiac*, 3j; powdered rhubarb, 3ij; cream of tartar, 3j; flowers of sulphur, 3j; one nutmeg, finely powdered; made into an electuary with a pound of clarified honey). Dose, two spoonfuls.

CHELTENHAM, MINERAL WATERS OF. C. is a town in Gloucestershire, England, nine miles from Gloucester, and 91 W. of London. Its water is one of the most celebrated natural purgatives in England. It is a saline, acidulous chalybeate, and is much frequented. Its main constituents are chloride of sodium, sulphate of soda, sulphate of magnesia, carbonic acid and carbonate of iron.

CHELTENHAM SALTS. These are sometimes made from the waters; at others, factitiously. The following is a formula: *Sodii chlorid*, *magnæ sulphat*, *sodæ sulph*, $\text{aa } 1\text{ij}$; dissolve, filter, evaporate to dryness; then add *Ferri sulph*, 3ss.

CHELTENHAM WATER, ARTIFICIAL, may be made of *Epsom salts*, gr. xij; *iron filings*, gr. j; *Glenber's salt*, 3iv; water, 4 gallons; impregnated with the gas from marble powder and sulphuric acid, $\text{aa } 3\text{ij}$.

CHELYS, Thorax.

CHELYS'CION, from *χελύς*, 'the chest.' A short, dry cough.—Galen, Hippocrates, Fœsius.

CHEMÉ, Chama, Cher'amia. An ancient measure equivalent to about two teaspoonfuls.

CHEMEUTICE, Chymistry.

CHEMIA, Chymistry.

CHEMIATER, Chymiatier.

CHEMIATRIA, Chymiatria.

CHEMICO-HISTOLOGY, see Chymico-history.

CHEMIOUS, Chymical, Chymist.

CHEMISMUS, Chymism.

CHEMIST, Chymist.

CHEMISTRY, Chymistry.

CHEMO'SIS, Chymo'sis, from *χρημ*, 'an aperture,' or rather, perhaps from *χυμος*, 'an humour.' A name given to ophthalmia, when the conjunctiva, surrounding the cornea, forms a high ring, making the cornea seem, as it were, at the bottom of a well. By some, it is used synonymously with *ophthalmia membranarum*. See Ophthalmia.

CHEMOTICE, Chymistry.

CHEMOTICUS, Chymical.

CHENAY, MINERAL WATERS OF. Chenay is a town in France, two leagues from Rheims. The waters are chalybeate.

CHÊNE, Quercus alba—c. *Marin*, *Fucus vesiculosus*—c. *Petit*, *Teucrium chamædrys*—c. *Vert*, *Ilex aquifolium*.

CHÊNETTE, Teucrium chamædrys.

CHÊNEVIS, see Cannabis sativa.

CHENOBOSCON, Potentilla anserina

CHENOCOPRUS, from *χην*, 'a goose,' and *κοπος*, 'dung.' The dung of the goose is so designated in some old Pharmacopœias. It was formerly employed as a febrifuge and diuretic.

CHENOPODE, Chenopodium.

CHENOPODIUM AMBROSIACUM, Ch. ambrosioides.

CHENOPODIUM AMBROSIODES, *Oh. suffructicosum* seu *ambrosiacum* seu *Mexicanum*, from *χην*, 'a goose,' and *ποδος*, *podos*, 'a foot'; *At'riplex Mexicana*, *Ambri'na ambrosioides*, *Botrys*, *B. Mexicana* seu *Americana*, *Ambrosioides Mexicana*, *Artemis'ia Botrys*, *Mexico* or *Spanish Tea*, (F.) *Anserine*, *Thé du Mexique*, *Ambrosie* ou *Ambrosie du Mexique*. *Ord. Chenopodiaceæ*. The infusion was once drunk as tea. It has been

given in paralytic cases; and in the United States is said to be used as an anthelmintic indiscriminately with Ch. anthelminticum.

CHENOPODIUM ANTHELMINTICUM, *Chenopodium*, *Botrys* seu *Ambri'na anthelmintica*, *Wormseed*, *Wormgoosefoot*, *Wormseed goosefoot*, *Jerusalem Oak* of America, *Goosefoot*, *Stinkweed*, (F.) *Anserine anthelmintique*, *A. vermifuge*. This plant grows plentifully in the United States. The fruit—*Chenopodium*, (Ph. U. S.)—is much used in cases of worms. Dose of the powder, from a teaspoonful to a tablespoonful or more. The oil, *O'leum Chenopodii*, (Ph. U. S.), from 8 to 10 drops, is more frequently exhibited. It is as much used in America as the *Semen Santonici* is in England.

CHENOPODIUM BONUS HENRICUS, *Chrysolachnanum*, *Mercurialis*, *Bonus Henricus*, *Tota bona*, *Lap'athum unctuosum*, *Chenopodium*, *Ch. sagittatum*, *Pes anserinus*, *English Mercury*, *Allgood*, *Angular-leaved goosefoot*, *Mercury goosefoot*, *Good King Henry*, (F.) *Anserine Bon Henri*, *Épinard sauvage*. The leaves are emollient, and have been applied to ulcers, &c. It has also been considered refrigerant and ecceprotic.

CHENOPODIUM BOTRYS, *Ambrosia*, *Artemisia Chenopodium*, *At'riplex odorata* seu *sua'volens*; *Jerusalem Oak*, (Eng.), (F.) *Anserine Botrys*, possesses anthelmintic properties, and was once given in diseases of the chest, palsy, &c. It is useless.

CHENOPODIUM FETIDUM, *Chenopodium vulvaria*—c. *Olidum*, *Ch. vulvaria*.

CHENOPODIUM QUINOA, Quinoa. A nutritious, wholesome, and agreeable article of food with the Peruvians. The leaves, before the plant attains maturity, are eaten as spinach; but the seeds are most generally used as food, boiled in milk or broth, and sometimes cooked with cheese and Spanish pepper.

CHENOPODIUM SAGITTATUM, Ch. *Bonus Henricus*—c. *Suffructicosum*, Ch. *ambrosioides*.

CHENOPODIUM VULVARIA seu *fætidum* seu *ol'idum*, *At'riplex fætida* seu *ol'ida*, *Vulvaria*, *Garosum*, *Raphæ*, *Stinking Orache* or *Goosefoot*, (F.) *Vulvaire*, *Arroche puant*, *Anserine fætide*. The fetid smell has occasioned it to be used as an antispasmodic and nerveine.

CHEOPINA, Cheopine.

CHEQUERBERRY, Gaultheria.

CHERAMIS, Cheme.

CHERBACHEM, Veratrum album.

CHERBAS, Lettuce.

CHERMES, Kermes—c. *Mineral*, *Antimonii sulphuretum præcipitatum*, and *A. sulphuretum rubrum*.

CHERNIBIUM, Urinal.

CHERRY, BIRD, Prunus padus—c. *Choke*, *Prunus Virginiana*—c. *Ground*, *Physalis viscosa*.

CHERRY TREE, RED, Prunus cerasus—c. *Tree*, black, *Prunus avium*—c. *Tree*, wild, *Prunus Virginiana*—c. *Water*, *Kirschwasser*.

CHERRY, WILD CLUSTER, Prunus padus—c. *Winter*, *Physalis*.

CHERSÆ, Fæces.

CHERVI, Sium sisarum.

CHERVIL, Scandix cerefolium—c. *Wild*, *Chærophylum sylvestre*.

CHESIS, from *χέειν*, 'to go to stool.' A more frequent desire to evacuate the bowels.

CHEST, Thorax—c. *Explorer*, see *Explorer*, chest.

CHEST-MEASURER, SIBSON'S. An instrument, described by Dr. Sibson, which is used for determining the mobility of the chest. It is a species of spring, which, when applied to the parietes of the chest, measures the modifications of its diameters, and indicates, by the motion of

the index hand on a dial, any movement of respiration, to the hundredth of an inch.

CHEST, NARROW, see Lordosis.

CHESTNUT TREE, *Fagus castanea*.

CHEVAUCHEMENT (F.), *Os'sium superpositio seu equitatio*, *Parallaxis*, *Parallaxis*. From *chevaucher*, (*cheval*, 'a horse'), 'to ride on horseback.' The riding of one bone over another after fracture, giving rise to shortening of the limb. See Riding of Bones.

CHEVELURE, Scalp.

CHEVSTRE, *Chestre*, *Capistrum*, (from *caput*, 'the head.') A bandage, applied round the head in cases of fracture or luxation of the lower jaw. According to the mode in which it is made, it is called *simple*, *double*, *oblique*, &c.

CHEVEU, Capillus.

CHEVILLE DU PIED, Malleolus.

CHÈVREFEUILLE, *Lonicera periclymenum*.

CHEVROTAIN PORTE-MUSC, see Musk.

CHEVROTANTE (VOIX), Egophony.

CHEWING, Mastication.

CHEYLETUS SCABIEL, see Psora.

CHEZANAN'CÈ, from *χέω*, 'I go to stool,' and *awayen*, 'necessity.' An ointment composed of honey and alum, and rubbed on the anus to occasion evacuations.—Paulus of Ægina.

CHIA, *Chia terra*, from *Chios*, an island where it was found. A kind of white earth, formerly used for burns.—Galen.

CHI'ACUM COLLYRIUM. A collyrium consisting of several drugs and Chian wine.—Paulus of Ægina.

CHIADUS, Furunculus.

CHIASMUS, *Chias'ma*, *Chiasm*, from *χίαω*, to form like the letter *χ*. The crucial union of parts,—as the *optic commissure* or *chiasm* of the optic nerve,—*Chias'mus seu Chias'ma nervo'rum optico'rum*.

CHIASTER, Kiaster.

CHIASTOS. Same etymon. A bandage, so called because it resembles the letter *χ*—*Oribasius*.

CHIBOU, see *Bursera gummifera*.

CHICHA. A drink made in Peru with Indian meal dried in the sun, and fermented with water. Its taste is that of bad cider. It is also made from rice, peas, barley, &c.

CHICK'S SPRINGS, see South Carolina, mineral waters of.

CHICKEN-BREASTED, see Lordosis.

CHICKEN-BROTH. When chicken-tea is boiled down one-half, with the addition of a little *parsley* or *celery*, and the yolk of an egg previously beaten up in two ounces of soft water, it forms a soup much relished by the convalescent.

CHICKEN-PEPPER, *Ranunculus abortivus*.

CHICKENPOX, *Varicella*.

CHICKEN-TEA, *Chicken-water*. This may be prepared as follows: Take a small chicken, freed from the skin and fat between the muscles; and having divided it longitudinally, remove the lungs, liver, and every thing adhering to the back and side-bones: cut the whole—bones and muscles—into very thin slices; put into a pan with a sufficient quantity of boiling water; cover the pan; and simmer with a slow fire for two hours. Put the pan upon the stove for half an hour, and strain through a sieve.

Used where the lightest animal diet is indicated.

CHICKEN-WATER, *Chicken-tea*.

CHICKENWORT, *Alsine media*.

CHICKWEED, *Alsine media*—c. *Water*, *Calitriche verna*.

CHICORÉE DES JARDINS, *Cichorium endivia*—c. *Sauvage*, *Cichorium intybus*.

CHICORY, *Cichorium intybus*.

CHICOT, Stump (of a tooth).

CHICOTIN (F.). The bitter powder or juice of the colocynth, with which the nurse, at times, rubs her nipple when desirous of weaning an infant.

CHIENRENT, *Triticum repens*—c. *Rouge*, *Sarsaparilla Germanica*.

CHIGGO, *Chique*.

CHIGGRE, *Chique*.

CHIGOE, *Chique*.

CHILBLADDER, *Chilblain*.

CHILBLAIN, (Prov.) *Chilbladder*, *Per'nio*, *Bagan'tia*, *Erythema Per'nio seu ? Eri'gae*, *Chimet'lon*, *Chimet'um*, *Chimon*, *Mules*, from *chill*, 'cold,' and *blain*, 'a pustule.' (F.) *Engelure*. An erythematous inflammation of the feet, hands, &c., occasioned by cold. It is very common in youth—not so in the adult or in advanced age. It is apt to degenerate into painful, indolent ulcerations, called *Kibes*, (Sc.) *Mules*. Chilblains are prevented by accustoming the parts to exposure; and are treated by stimulant, terebinthinate, and balsamic washes, ointments, and liniments.

CHILD, Infans.

CHILD-BEARING, Parturition.

CHILD-BED, Parturient, Parturition.

CHILD-BED FEVER, Puerperal fever.

CHILD-BIRTH, Parturition.

CHILD-CROWING, *Asthma thymicum*.

CHILDHOOD, Infancy.

CHILD-ILL, Parturition.

CHILD-MURDER, Infanticide.

CHILDREN'S BANE, *Cicuta maculata*.

CHILL, CLIMATE OF, see Lima.

CHILI, MINERAL WATERS OF. The most celebrated mineral springs of Chili, in South America, are those of Peldehués and Cauquenes. The former are not far from St. Jago. They consist of two springs, one thermal, the other cold. The hot spring is clear, inodorous, and contains soda and carbonic acid. The cold spring contains iron and sulphate of soda. Cauquenes is much resorted to by invalids during the summer. Mineral waters are very common in Chili. Those of Apoquindo, Colina, and Cauquenes, have been analyzed by Prof. J. Lawrence Smith, M.D.

CHILIOGRAMMA, Kilogramme.

CHILLOPHYLLON, *Achillea millefolium*.

CHILITES, *Chelitis*.

CHILL, Rigor.

CHILLI, Capsicum.

CHI'LOIN, *Choi'lon*, *Cheli'tis*, from *χελος*, 'a lip.' Inflammation of the lips.—Vogel. One who has a thick lip; *Labeo*, *Labeo*.

CHILOPLASTICE, *Cheloplastice*.

CHIMAPHILA, *Pyrola umbellata*—c. *Umbellata*, *Pyrola umbellata*.

CHIMETLUM, *Chilblain*.

CHIMIA, Chymistry.

CHIMIATER, Chymiatier.

CHIMIATRIA, Chymiatria.

CHIMIE, Chymistry.

CHIMISME, Chymism.

CHIMISTE, Chymist.

CHIMON, *Chilblain*, Cold.

CHINA, *Cinchona*, *Smilax china*—c. *Americana* or *West India*, *Smilax pseudo-china*—c. *Calisaya*, *Cinchonæ cordifolia cortex*—c. *Ocidental*, *Smilax pseudo-china*—c. *Orientalis*, *Smilax china*—c. *Ponderosa*, *Smilax china*—c. *Regia*, *Cinchonæ cordifolia cortex*—c. *Root*, *Smilax china*—c. *Spuria nodosa*, *Smilax pseudo-china*—c. *Vera*, *Smilax china*.

CHINCAPIN, *Fagus castanea pumila*—c. *Water*, *Nelumbium luteum*.

CHINCHE, Cimex.

CHINCHINA, *Cinchona*.

CHINCHUNCHULLI, *Tonidium marcucel*.

CHINCOUGH, Pertussis.

CHINESE, MEDICINE OF THE, *Medici'na Sin'ica*. Medicine has been long, but most imperfectly, practised by the Chinese. From their therapeutics we have obtained the old operations of acupuncture and moxibustion.

CHING'S WORM LOZENGES, see Worm Lozenges, Ching's.

CHINIDIN, Quinidia.

CHININUM, Quinine. See Chinium.

CHINIO'DINE, *Chino'idine*, *Chinoidi'na*, *Chinoidin*, *Quinoid'ine*, *Quinoid'in*, *Qui'nodin*; from *China*, 'Cinchona.' A substance presumed to be an alkaloid by Sertürner, who separated it from cinchona. It has been supposed to be a mixture of quinia, cinchonia, and a peculiar resinous matter, but according to Liebig it is simply the alkaloid quinia in an amorphous state—*amorphous quinia*.

CHINIUM ACETICUM, Quiniæ acetas—c. Arsenicosum, Quiniæ arsenias—c. Citricum, Quiniæ citras—c. Ferrocyanogenatum, Quiniæ Ferrocyanas—c. Hydrochloricum, Quiniæ murias—c. Hydroiodicum, Quiniæ hydriodas—c. Lacticum, Quiniæ lactas—c. Muraticum, Quiniæ murias—c. Nitricum, Quiniæ nitras—c. Phosphoricum, Quiniæ phosphas—c. Salitum, Quiniæ murias—c. Sulphuricum, Quiniæ sulphas—c. Tannicum, Quiniæ et Cinchoninæ tannas—c. Valerianicum, Quiniæ valerianas.

CHINNEYWEED, Lichen roccella.

CHINOLEINUM, Leukoleinum.

CHINQUAPIN, Fagus castanea pumila.

CHINWHELK, Syosis.

CHIOCOCCÆ RADIX, Caincæ radix.

CHIOLI, Furunculus.

CHION, Snow.

CHIONAN'THUS VIRGIN'ICA, *Fringe Tree*, *Snowdrop Tree*; indigenous; *Nat. Ord.* Oleaceæ; from *χίων*, 'snow,' and *ανθος*, 'a flower.' An infusion of the root has been used as an antiperiodic and tonic.

CHIQUE (F.), *Puce pénétrante*, *Pulex Penetrans*, *Dermatophilus*, *Sarcophyll'a penetrans*, *Tick*, *Chigge*, *Chig'oe*, *Chiggo*, *Chegre*, *Cheg'oe*, *Jigger*. A small insect in America and the Antilles, which gets under the epidermis, and excites great irritation.

CHIR, Manus, see Chir.

CHIRAETA, Gentiana chirayta.

CHIRAGRA, from *χειρ*, 'hand,' and *αγχα*, 'a seizure.' Gout in the hand.

CHIRAPOTHECA, Arsenal.

CHIRAPSLIA, Friction.

CHIRARTHRI'TIS, from *χειρ*, 'hand,' *αρθρον*, 'joint,' and *itis*, denoting inflammation. Inflammation of the joints of the hand.

CHIRAYITA, Gentiana chirayta.

CHIRAYTA, Gentiana chirayta.

CHIRETTA, Gentiana chirayta.

CHIRHEUMA, Chirrhœuma.

CHIRIATER, Surgeon.

CHIRIATRIA, Surgery.

CHIRIMOYA, Anona tripetala.

CHIRISIS, Surgery.

CHIRISMUS, Surgery.

CHIRIXIS, Surgery.

CHIROCYRTO'SIS, from *χειρ*, 'the hand,' and *κυρτωσις*, 'crookedness.' Crookedness of the hand.

CHIROMANCY, *Chironanti'a*, *Vaticini'um chironanti'cum*, *Pal'mistry*, from *χειρ*, 'the hand,' and *μαντεία*, 'divination.' (F.) *Chironancie*. Art of divining by inspection of the hand.

CHIRONAX, Surgeon.

CHIRO'NIA. (From Chiron, *χαιρον*, the Centaur, who is said to have discovered its use.) A genus of plants. *Ord.* Gentianaceæ. Also, *Tarax communis*.

CHIRO'NIA ANGULA'RIS, *Amer'ican Cent'ary*,

Rosepink, *Wild Sue'ory*, *Bitterbloom*, *Centry Sabba'tia*, *S. Angula'ris*. Every part of this plant is a pure and strong bitter, which property is communicated alike to alcohol and water. It is used as a tonic and stomachic.

CHIRO'NIA CENTAURI'UM, *Centauri'um minus seu vulga'rè seu parvum*, *Gentia'na centauri'um seu Gerar'di*, *Erythra'a Centauri'um*, *Hippocentaure'a centauri'um*, *Centaur'um*, *Centaur'ris*, *Smaller or Lesser Cent'ary*, *Europe'an Cent'ary*, (F.) *Centaurie petite*, *Herbe au centaure ou à la fievre*. The tops of the lesser Centaury, *Centauri'rii Cacum'ina*, are aromatic and tonic, and are sometimes employed as such.

CHIRO'NIA CHILEN'SIS, *Gentia'na Cuchenhauen*, *Cachen-laguen*, *Chachinlagua*, *Chancelagua*, *Erythra'a Chilensis*, *Gentia'na Peruvia'na*. A very bitter plant, indigenous in Chili. It possesses the virtues of the *Chironææ*. Given in infusion—(3j, to water Oj.)

CHIRONI'UM, from *χαιρον*, 'bad, malignant.' An ulcer difficult of cure;—of a swollen, hard, and callous nature.—Galen. Some have supposed the word to come from Chiron, the Centaur, who was unable to cure such ulcers.

CHIRONOMIA, Cheironomia.

CHIROPODALGIA, Acrodynia.

CHIROP'ODIST, (F.) *Pédi'cure*. One who treats diseases of the hands and feet, or rather whose profession it is to remove corns and bunions; from *χειρ*, 'the hand,' and *πους*, *podos*, 'the foot.'

CHIROBRHEUMA, Chirrhœuma.

CHIROSI'S, Subactio.

CHIROSTROPHO'SIS, from *χειρ*, 'the hand,' and *στροφειν*, 'to turn.' Distortion of the hand.

CHIROTHE'CA, from *χειρ*, 'the hand,' and *θηκη*, 'a sheath.' A bandage for the hand. A kind of bandage in which the fingers and hand are enveloped in spiral turns. When the whole hand and fingers are covered, it is called the double or complete *Chirotheca*, *Ch. comple'ta*, *Vinctu'ra omnibus dig'itis*; and when only a finger is covered, the half or incomplete, *Ch. incomple'ta*, *Vinctu'ra pro uno dig'ito*. See *Gantelet*.

CHIROTRI'BA, from *χειρ*, 'the hand,' and *τριβω*, 'I rub.' Friction with the hand. According to others, dexterity in an art.—Hippocrates, Galen.

CHIRRHEUMA, *Chirorrhœuma*, *Rheumatis'mus manûs*, from *χειρ*, 'the hand,' and *ρευμα*, 'flux.' Rheumatism of the hand.

CHIRURGEON, Surgeon.

CHIRURGI PHYSICI, see Surgeon.

CHIRURGIA, Surgery—c. Anaplastica, Morioplastice—c. Curtorum, Morioplastice—c. Infusoria, Infusion of Medicines—c. Militaris, Surgery, military—c. Transfusoria, Transfusion.

CHIRURGICAL, Surgical.

CHIRURGICUS, Surgical.

CHIRURGIE, Surgery—c. Militaire, Surgery, military.

CHIRURGIEN, Surgeon—c. Consultant, Consulting Surgeon—c. Hernieur, see Hernial.

CHIRURGIQUE, Surgical.

CHIRURGUS, Surgeon.

CHIST. An Arabic word which signifies the sixth part. The sixth part of the *Congius* or gallon.

CHITON, Tunic.

CHITONISCUS, Indusium.

CHITTICK'S NOSTRUM, see Nostrum.

CHIUUM VINUM. From *Chios*, the island where it was produced; *Chian wine*; used by the physicians of antiquity in cases of defluxions and ophthalmia.—Scribonius Largus.

CHLIABOS, *χλιαρος*, 'tepid.' A name given to slight fevers, in which the heat is not great.—Galen.

CHLIAS'MA, *χλίσμα*, same etymon. A tepid and moist fomentation.—Hippocrates.

CHLOAS'MA, *Pityriasis versicolor*, *Macula hepatica*, *Pannus hepaticus*, *Hepaticum*, *Phaze*, *Phalcea*, *Phaeus*, (F.) *Taches hépatiques*, *Chaleurs du foie*, *Éphélide scorbutique*, *Liverapot*, from *χλωος*, 'a greenish-yellow colour, verdure.' A cutaneous affection, characterized by one or more broad, irregular-shaped patches, of a yellow or yellowish-brown colour, occurring most frequently on the front of the neck, breast, abdomen, and groins. The patches do not generally rise above the surface. There is usually some degree of itching. A cryptogamous growth has been found in it, to which M. Robin gives the name *Microsporon furfur*.

The causes are not very evident. Sulphur externally—in any and every form—generally removes it speedily. Should there be difficulty, the external use of the remedy in baths or fumigations may succeed.

CHLOASMA ALBUM, Achroma.

CHLORA, Chlorine.

CHLORAS KALICUS DEPURATUS, Potasse murias hyperoxygenatus.

CHLORASMA, Chlorosis.

CHLORE, Chlorine—c. *Liquide*, see Chlorine.

CHLORÉTHÉRIDE, Chloroform.

CHLORETUM CALCARIÆ, Calcei chloridum.

CHLORIASIS, Chlorosis.

CHLORIC ETHER, CONCENTRATED, see Ether, chloric—c. Ether, strong, see Ether, chloric.

CHLORIDE OF BROMINE, see Bromine, chloride of.

CHLORINE, from *χλωρος*, 'green.' *Chlorinum*, *Chlorinum*, *Chlora*, *Oxy muriatic Acid Gas*, *Oxygenated Muriatic Acid Gas*, *Dephlogisticated Marine Acid*, *Hal'ogene*, *Mu'rigene*, *Chlorum*, (F.) *Chlore*. So far as we know, this is an elementary substance. It is a greenish, yellow gas, of a strong suffocating smell, and disagreeable taste: incapable of maintaining combustion and respiration, and very soluble in water. One of its characteristics is, that of destroying, almost immediately, all vegetable and animal colours. It is employed in fumigations as a powerful disinfecting agent. A very dilute solution, *Aqua seu Liquor Chlorini*, *Chlorin'ii Liquor* (Ph. D.), *Chlorin'ei aqua* (Ph. E.), *Chlorine water*, (F.) *Chlore liquide*, has been administered internally, in certain cases of diarrhoea and chronic dysentery. Immersion of the hands and arms in it has often removed itch and other cutaneous affections. It has also been inhaled in a dilute state in the early stage of phthisis, but it is of doubtful efficacy, and is better adapted for chronic bronchitis.

CHLORINE, BISULPHURET OF, Sulphur, chloride of.

CHLORO-ANÉMIE, Chlorosis.

CHLOROFORM, *Chloroformum*, *Chloroformum*, *Carb'oneum chloratum*, *Superchloridum formyl'icum*, *Perchloride* and *Tetrachloride of Formyl*, *Chloroform'yl*, called also, but not correctly, *Tetrachloride of Carbon*, and *Chloric ether*, *Æther chloricus*;—(F.) *Chloroforme*, *Chloréthér'ide*, so called on account of the connexion of chlorine with formic acid, is a colourless, oleaginous liquid, of a sweetish ethereal odour, hot, aromatic, and peculiar taste. The specific gravity of that of the Ph. U. S. is 1.49. It may be obtained by distilling from a mixture of chlorinated lime and alcohol,—rectifying the product by redistillation, first from a great excess of chlorinated lime, and afterwards from strong sulphuric acid. It has been used with advantage in asthma, and in diseases in which a grateful soothing agent is required. Dose fʒss to fʒj, diluted with water.

It has likewise been prescribed with great success as an anæsthetic agent in spasmodic diseases; and to obtund sensibility in surgical operations and in parturition,—especially in the way of inhalation; but its use requires caution. See Anæsthetic.

CHLOROFORM, TINCTURE OF, Ether, chloric.

CHLOROFORMIZA'TION, *Chloroformisatio* The aggregate of anæsthetic phenomena occasioned by the inhalation of chloroform.

CHLOROFORMYL, Chloroform.

CHLORO'MA, from *χλωρος*, 'green.' A peculiar form of tumour containing a yellow green substance, which takes the place, at times, of muscles and periosteum.

CHLOROPHYLL, Fecula, green.

CHLORO'SIS, from *χλωρος*, 'green,' *Pallidus Morbus*, *Fædus Virginum color*, *Pallidus color virgin'us*, *Pallor virginum*, *Jeterit'ia alba*, *Jeterus albus*, *Leucopathi'a*, *Morbus virgin'us seu Parthen'us*, *Fædi col'ores*, *Dyspep'ia chlor'is*, *Feb'ris amato'ria*, *Cachex'ia seu Feb'ris virginum*, *Feb'ris alba*, *Anepithym'ia chlor'is*, *Chloros'mum*, *Chloros'ma*, *Chlori'asis*, *Citto'is*, *Green-sickness*, (F.) *Chlorose*, *Pâles-couleurs*, *Chloro-anémie*. A disease which affects young females, more particularly those who have not menstruated. It is characterized by a pale, lurid complexion, languor, listlessness, depraved appetite and digestion, palpitation, &c. The disease generally goes off on the occurrence of the menstrual flux; but sometimes it is long before this is established, and, at times, the catamenia are in much larger quantity than usual. To this last form M. Trousseau has given the name *chlorose hémorrhagique*.

The blood of chlorosis is generally thin, light-coloured and deficient in red corpuscles; and, according to Piorry, in iron,—*achalybémie*; and the clot is in less proportion to the serum than in health. On auscultation, a bellows' sound has been almost invariably detected over the heart, and a continuous blowing sound in the larger arteries, (especially the carotids and subclavians,) re-enforced by each systole of the ventricle, and resembling the buzzing of a humming-top, the cooing of doves, the whistling of air through a key-hole, &c., (see *Bruit*.) Very similar sounds are heard in the arteries after copious hemorrhage: they seem, therefore, to coincide with enfeebled circulation.

Tonics—as iron—are usually required in the treatment,—the disease most commonly occurring in those in whom there is considerable torpor of the system.

CHLOROSIS ÆTHIOPUM, Chthonophagia—c. *Amatoria*, *Hætic fever*—c. *Gigantea*, see *Poly-sarcia*.

CHLOROSMA, Chlorosis.

CHLOROT'IC, *Chlorot'icus*, (F.) *Chlorotique*. Affected with chlorosis, or pertaining to chlorosis;—as *chlorotic female*, *chlorotic symptoms*, &c.

CHLORUM, Chlorine.

CHLORURE DE CARBON, Chloroform—c. *d'Or*, *Gold*, *Muriate of*.

CHLORURE AMMONICUM, *Ammonia murias*—c. *Oxidi Calcii*, *Calcei chloridum*.

CHOA, Chu.

CHOA'CUM EMPLAS'TRUM NIGRUM. A black plaster, mentioned by Celsus, and composed of equal parts of litharge and resin. The litharge was first boiled in oil.

CHOAK, *Cynanche trachealis*—c. *Wolf*, *Ly-canthe*.

CHOANA, *Pelvis*—c. *Cerebri*, *Infundibulum of the brain*.

CHOANE, *Infundibulum*.

CHOANORRHAGIA, *Epistaxis*.

CHOAVA, *Coffea Arabica*.

CHOCOLAT, see Chocolate.

CHOCOLATA, Chocolate—c. cum Osmazomâ, see Osmazome—c. Salutis, see Chocolate—c. Simplex, see Chocolate.

CHOCOLATE, *Chocolatatum*, *Chocola'ta*, *Succola'ta*, *Succocolla'ta*. Dr. Alston says, that this word is compounded from two East Indian words:—*choco*, 'sound,' and *alte*, 'water,' because of the noise made in its preparation [?]. An alimentary paste prepared from the kernels of *Theobroma cacao* or *Cacao*, with sugar, and often aromatics. (See *Cacao*.) The chocolate thus simply prepared—as it is met with, indeed, in commerce—is called in France, *Chocolat de santé*, *Chocola'ta simplex* seu *Salu'tis*. It is not very easy of digestion.

The *chocolat à la vanille* contains three ounces of *vanilla* and two of *cinnamon* to twenty pounds of common chocolate. The addition of the aromatic renders it somewhat more digestible. Chocolates may likewise be medicated.

CHOCOLATE, OSMAZOME, see Osmazome.

CHOCOLATE ROOT, *Geum rivale*, G. Virginianum.

CHOCUS, Chu.

CHENICIS, Trepan.

CHENION, Cord.

CHENOS, Cord.

CHERADO'DES, from *χοιρας*, *χοιραδος*, 'scrofula,' and *ειδος*, 'resemblance.' Of the nature of scrofula. Affected with scrofula.

CHERAS, Scrofula.

CHOIROS, Vulva.

CHOKE CHERRY, *Prunus Virginiana*—c. Dump, Carbonic acid.

CHOKES, Throat.

CHOLA, Chole.

CHOLÉ'MIA, *Cholhæ'mia*, *Choli hæ'mia*, from *χολη*, 'bile,' and *αἷμα*, 'blood.' A morbid state, in which bile exists in the blood. Jaundice.

CHOLAGO, Cholas.

CHOL'AGOGUE, *Cholago'gus*, *Chole'gos*, *Fel'iducus*, *Bili'ticus*, from *χολη*, 'bile,' and *αγω*, 'I expel.' The ancients give this name to cathartics, which were reputed to cause the flow of bile.

CHOLANSIS, Cholosis.

CHOLAS, *χολας*, plur. *χολαδες*. *Chola'go*. The epigastric region. The intestines.—Homer.

CHOLASMA, Cholosis.

CHOLÉ, *Cholus*, *Chola*, 'bile,' in composition. Hence:

CHOLEC'CHYSIS, *Cholen'chysis*; from *χολη*, 'bile,' and *εχχσις*, 'effusion.' Effusion of bile.

CHOLECYST, Gall-bladder.

CHOLECYSTEURYS'MA, from *χολη*, 'bile,' *κυστις*, 'bladder,' and *ευρυσμα*, 'dilatation.' Dilatation of the gall-bladder.

CHOLECYSTITIS, *Inflammat'io Vesicæ fell'æ*, *I. cyst'idis fell'æ*, *Cysti'tis fell'æ*, *Hepati'tis cyst'idis*, from *χολη*, 'bile,' and *κυστις*, 'bladder.' (F.) *Inflammation de la Vésicule du Fieci Cholécystite*. Inflammation of the gall-bladder.

CHOLEDOCH, *Choledochus*, (F.) *Cholé-douque*, from *χολη*, 'bile,' and *δοχος*, 'containing or receiving.' The *Ductus choledochus* seu *hepato-cysticus*, *Ductus communis choledochus*, (F.) *Conduit ou Canal Cholédouque*, is the duct formed by the union of the hepatic and cystic ducts, which pour the hepatic and cystic bile into the duodenum.

CHOLEDOCITIS, from *choledochus*, and *itis*; a suffix denoting inflammation. Inflammation of the choledoch duct.

CHOLEDOG'RAPHY, *Choledograp'hia*, *Cholepgraph'ia*, *Cholograph'ia*, from *χολη*, 'bile,' and *γραφειν*, 'to describe.' A description of what relates to the bile and biliary organs.

CHOLEDOL'O'GY, *Choledolog'ia*, *Cholelog'ia*, *Cholog'ia*, from *χολη*, 'bile,' and *λογος*,

'a discourse.' A treatise on the bile and biliary organs.

CHOLÉDOQUE, *Choledoch*.

CHOLEGOS, *Chologogue*.

CHOLEGRAPHIA, *Choledography*.

CHOLEHÆMIA, *Icterus*.

CHOLEIA, *Claudication*.

CHOLELITHIA, *Cysthepatolithiasis*—c. *Icterus*, *Icterus*.

CHOLELITHIASIS, *Cysthepatolithiasis*.

CHOLELITHUS, *Calculi*, *biliary*.

CHOLELOGIA, *Choledology*.

CHOLEMES'IA, *Cholem'esia*, from *χολη*, 'bile,' and *εμεσις*, 'vomiting.' Vomiting of bile.

CHOLENCHYSIS, *Cholecchysis*.

CHOLEPYRA, *Fever*, *bilious*.

CHOLEPYRETUS, *Fever*, *bilious*.

CHOLEPYRRIN, see *Bile*.

CHOLER, *Bile*. Anger was supposed to be produced by a superabundance of bile; hence the term *Choler* for anger.

CHOL'ERA, *Chol'era-morbus*, *Cholera nostras* seu *vulgaris*, *Sporad'ic Chol'era*, *Cholerrhæ'gia*, *Passio choleric'a*, *Felliflua passio*, *Morbus fellifluus*, *Hol'era*, *Bilis flux'io*, (F.) *Choladræ lymphatique*, *Hydrocholadræ*, *Choléra-morbus sporadique*, *Ch. Européen*, *Trousse-galant*, from *χολη*, 'bile,' and *ερω*, 'I flow.' According to others, from *χολαδες*, 'intestines,' or from *χολερα*, 'the gutter of a house to carry off the rain.' The higher degrees have been called *Centrogangli'itis*, *Myelogaugli'itis*, and *Hologangli'itis*. A disease characterized by anxiety, gripings, spasms in the legs and arms, and by vomiting and purging (generally bilious:) vomiting and purging are, indeed, the essential symptoms. The disease is most common in hot climates,—and in temperate climates, during summer. *Spasmod'ic*, *Asiat'ic*, *Malig'nant*, *Ind'ian*, *Epidem'ic*, *Pestilential*, *Convulsive Nervous*, *Eastern*, or *Oriental Cholera*, *Asphyx'ia pestilen'ta*, *Pestilential asphyx'ia*, *Choleric Pest'ience*, *Cholera orientalis* seu *Indica* seu *Epidem'ica* seu *Asphyx'ia*, *Typhus Bengalensis*, *Chol'ero-typhus*, *Ganglion'is peripher'ica et medullaris*, *Hymenogaugli'itis*, *Pantogaugli'itis*, *Cholerrhæ'a lymphatica*, *Psorent'ia*, *Achol'ia*, *Typhoid Fever of India*, *Hyperanthrax'is*, *Enterop'ya Asiat'ica*, *Triplanchn'ia*, *Triplanchni'tis*, *Hæmatoporrh'osis*, *Hæmatorrh'osis*, *Morbus ory'zeus*, *Rice disease*, (because supposed by Tytler to be caused by damaged rice,) (F.) *Mort de Chien*, is frightful in the rapidity of its progress, the patient sometimes dying in a few hours from the first onset. In temperate climates, common cholera is not usually a disease of much consequence. It requires that the stomach and bowels should be cleared, and afterwards the irritation be allayed by full doses of opium. In malignant cholera, large doses of calomel, with opiates, form the great reliance of many practitioners. In the worst forms, however, it runs its course, unmodified by treatment.

CHOLÉRA EUROPÉEN, see *Cholera*—c. Indian, see *Cholera*—c. Indica, see *Cholera*.

CHOL'ERA INFANTUM, *Choleric Fever of Infants*; *Cholera of Infants*. A disease so termed by American physicians. It occurs, generally, in the middle states of the Union, in June or July, and continues during the hot weather, hence called the 'summer complaint.' The chief symptoms are vomiting, purging of green or yellow matter, slime or blood, attended with pain or uneasiness; and swelling of the abdomen, with some prexia, generally. The heat of the weather seems to be the predisposing, if not the exciting, cause. It is a fatal disease in towns; differing little, if at all, from what is vulgarly called the *Watery Gripes* in England. Clearing, gently,

the alimentary canal, so as to remove the offending matter, and then exhibiting chalk mixture and laudanum, with counter-irritants, as sinapisms to the abdomen, is the most satisfactory plan of treatment.

CHOLERA, MALIGNANT, see Cholera—c. Morbus. Cholera—c. *Morbus Sporadicus*, see Cholera—c. Nostras, Cholera morbus—c. Oriental, see Cholera—c. Pestilential, see Cholera—c. Poison, see Poison—c. Spasmodic, see Cholera—c. Sporadic, see Cholera—c. *Sec*, Cholera sicca—c. *Vulgaris*, Cholera morbus.

CHOLERA SICCA, (F.) *Choléra sec.* Pnéumatosis of the digestive passages in which there is a copious discharge of gas upwards and downwards. CHOLERAIC, Choleric.

CHOLERAPHOBIA, Choleromania.

CHOLERIC, *Choler'icus*, (F.) *Cholérique*. Belonging to bile; also to cholera; *Cholera'ic*,—as the 'choleric or choleraic virus.' The French use the term *Fèvre cholérique* for the fever accompanying cholera.

CHOLERIC FEVER OF INFANTS, Cholera infantum.

CHOLERIC TEMPERAMENT. The bilious temperament.

CHOLERINE, (F.) *Cholérine*. A diminutive of cholera. The first stage of epidemic cholera; also, the precursory symptoms of cholera.

CHOLÉRIQUE, Choleric.

CHOLEROID, *Choleroïdes*, from *cholera*, and *εἶδος*, 'resemblance.' Resembling cholera; as a 'choleroïd affection.'

CHOLEROMANIA, *Cholerapho'bia*, *Cholero-pho'bia*, from *cholera* and *mania*. A dread of cholera to such an extent that the individual fancies himself affected or threatened by it.

CHOLEROPHOBIA, Choleromania.

CHOLEROPHONÉ, from *cholera*, and *φωνή*, 'voice.' The peculiar voice of one in cholera.

CHOLEROPROSPON, *Cholero'psis*, from *cholera*, and *προσῶπον*, 'countenance.' The facial expression of cholera.

CHOLEROPSIS, Choleroprosopon.

CHOLERO-TYPHUS, Cholera, (spasmodic.)

CHOLERRHAGIA, Cholera.

CHOLERRHŒA LYMPHATICA, Cholera.

CHOLESTEATINOMA, Cholesteatoma.

CHOLESTEATINE, Cholesterin.

CHOLESTEATOMA, from *χολή*, 'bile,' and *στέατομα*. An encysted tumour, which presents upon the interior of the cyst several superimposed layers of cholesterin, which glisten like pearl. Hence it was named by Cruveilhier the *laminated nacreous fatty tumour*.

CHOLESTERIN, *Chol'esterine*, *Cholesteri'na*, *Cholosteri'na*, *Cholestearine*, *Cholesteatine*, from *χολή*, 'bile,' and *στερεός*, 'solid,' or *στέαρ*, 'suet,' *Ad'ipocire*, *Choline*. An inodorous, insipid substance, in white, shining scales; fusible and crystallizing, on cooling, in radiated fibres. Soluble in alcohol. It forms the crystalline part of certain biliary calculi. See *Adipocire*.

CHOLESTERIN DISEASE, see *Lardaceous*.

CHOLEUMA, Choleosis.

CHOLŒMIA, Cholæmia.

CHOLIA, Choleosis.

CHOLICE'LE, from *χολή*, 'bile,' and *κλήη*, 'tumour.' A swelling, formed by the bile morbidly accumulated in the gall-bladder.

CHOLICUS, Bilious.

CHOLIHÆMIA, Cholæmia.

CHOLINE, Cholesterin.

CHOLICYSTIECTASIE, *Turgescentia vesicæ felleæ*.

CHOLICYSTITE, Cholecystitis.

CHOLIUS, Bilious.

CHOLOCATARRHOPHŒ, *Cholocatarrho'phæ*, from *χολή*, 'bile,' and *καταρροῦν*, 'to sup up.' The absorption of bile.

CHOLQGRAPHIA, Choleodography.

CHOLOLITHUS, see *Calculi*, biliary.

CHOLO'MA, from *χωλός*, 'lame, maimed.' Distortion of a limb; incapacity of moving it. Also, the act of limping, claudication.

CHOLOPLANIA, Icterus.

CHOLORRHŒA, from *χολή*, 'bile,' and *ρῆσις*, 'I flow.' An abnormal discharge of bile.

CHOLOS, Bile, Chole.

CHOLO'SES, from *χολή*, 'bile.' Alibert includes, under this head, every morbid affection of the liver and spleen.

CHOLOSIS, from *χωλός*, 'lame.' *Cholotes*, *Cholas'ma*, *Cholan'sis*, *Cholan'ma*, *Chol'ia*, *Claudication*. Hippocrates employs this word particularly for a lameness of the hand, which renders a person one-handed, (F.) *manchet*. Also, icterus; bilious dyscrasy.

CHOLOSIS AMERICANA, Fever, yellow.

CHOLOSTERINA, Cholesterin.

CHOLOTES, Choleosis.

CHOLEZEMIA FEBRILIS, Fever, bilious.

CHONDRIIN, from *χονδρός*, 'cartilage,' (F.) *Gélatine des cartilages*. A variety of gelatin, obtained by boiling the cornea, the permanent cartilages, and the bones before they are ossified.

CHONDRI'TIS, *Inflammatio Cartilag'ina*, (F.) *Chondrite*, from *χονδρός*, 'cartilage,' and *itis*, a termination denoting inflammation. Inflammation of cartilage.

CHONDROCLASIS, from *χονδρός*, 'cartilage,' and *κλάσις*, 'fracture.' Fracture of a cartilage.

CHONDRODES, Cartilaginous, Chondroid.

CHONDROGENESIS, *Chondrogen'esis*, from *χονδρός*, 'a cartilage,' and *γενεσις*, 'formation.' Formation of cartilage. A morbid conversion of parts into cartilage.

CHONDROGLOS'SUS, from *χονδρός*, 'a cartilage,' and *γλῶσσα*, 'the tongue.' A fasciculus of fleshy fibres, passing from the lesser cornua of the os hyoides to the tongue. It forms part of the hyoglossus.

CHONDROGRAPHY, *Chondrogra'phia*, from *χονδρός*, 'cartilage,' and *γραφη*, 'a description.' A description of cartilages.

CHONDROID, *Chondroï'des*, *Chondroï'des*, from *χονδρός*, 'cartilage,' and *εἶδος*, 'resemblance.' Cartilaginous. Resembling cartilage.

CHONDROLOGY, *Chondrolog'ia*, from *χονδρός*, 'a cartilage,' and *λογία*, 'a discourse.' A treatise on cartilages.

CHONDROMA, *Enchondro'ma*, from *χονδρός*, 'cartilage,' A cartilaginous growth proceeding from bones, including spina ventosa, osteosarcoma, &c., of many.

CHONDRO-PHARYNGE'US, from *χονδρός*, 'a cartilage,' and *φάρυγξ*, 'the pharynx.' Fibres of the muscular coat of the pharynx, which arise from the lesser cornua of the os hyoides, and form part of the *Constrictor medius*. See *Constrictor pharyngis*.

CHONDROSIS, from *χονδρός*, 'cartilage.' A morbid formation or condition of a cartilage.

CHONDROSTERNAL, *Chondrosterna'lis*, from *χονδρός*, 'a cartilage,' and *στερνον*, 'the breast-bone.' Having relation to the cartilages of the ribs and breast-bone,—as the *chondrosternal ligaments*, articulations, &c.

CHONDROSYNDESMUS, from *χονδρός*, 'a cartilage,' and *σύνδεσμος*, 'a ligament.' Union of bones, by means of fibro-cartilage. Synchondrosis.

CHONDROTOMY, *Chondrotom'ia*, from *χονδρός*, 'a cartilage,' and *τεμνειν*, 'to cut.' Dissection of cartilages.

CHONDRO-XIPHOID, *Chondro-xiphoid'ia*, from *χονδρός*, 'a cartilage,' and *ξίφος*, 'a sword.' Resembling a sword.

from *χόνδρος*, 'a cartilage,' and *xiphoid*, the cartilage so called. Having relation to the xiphoid cartilage, as '*Chondro-xiphoid ligament*.'

CHONDRUS, Cartilage. *The Xiphoid cartilage*, in particular.

CHONDRUS, *Alica*, *Fucus crispus*—c. *Crispus*, *Fucus crispus*—c. *Polymorphus*, *Fucus crispus*.

CHONOS, Infundibulum.

CHOPINE, *Chopi'no*, *Choepli'na*. A measure, which contains 16 ounces. A pint measure.

CHORA, *χωρα*, 'region.' Any void space. Also, the orbit of the eye.

CHORD, **TESTICULAR**, Spermatic chord.

CHORDA, *Corda*, from *χορδή*, 'a gut,' and hence, a string made of gut. A string of an instrument. It has several meanings:—as a *tendon*,—the *intestines* (*Chordæ*.) Paracelsus calls the genital organs *Chordæ*. *Chorda seu Funiculus Tympani*, a branch of the 7th pair of nerves, according to some, of the 5th pair, according to others, and to probability, which passes through the tympanum, (F.) *Nervus Tympanicus*—(Ch.) *Corde du Tambour ou du Tympan*. *Chordæ Willisii*, *Trabeculæ Willisii*, the small fibres which cross the sinuses of the dura mater: so called from their first describer, Willis. *Chordæ Tendinæ*; the tendinous and cord-like substances which connect the *columnæ carneæ* of the ventricles of the heart to the valves, &c.

CHORDA, Intestine.

CHORDA DORSALIS, *Dorsal cord*. The axis, around which, according to Von Baer, the first parts of the fœtus are formed,—the rudiment of the future vertebral column.

CHORDA HIPPOCRATIS, *Achillis Tendo*—c. *Longitudinalis Corporis Callosi*, see *Raphe*—c. *Penis*, *Chordee*—c. *Tympani*, see *Chorda*—c. *Veneris*, *Chordee*.

CHORDÆ, *Chordee*—c. *Ferrenii*, *Cords*, vocal—c. *Longitudinales lanciai*, see *Raphe*—c. *Tendinæ*, see *Chorda*—c. *Transversales Willisii*, see *Corpus callosum*—c. *Vocales*, *Cords*, vocal—c. *Willisii*, see *Chorda*.

CHORDAP'SUS, from *χορδή*, 'intestine,' and *αἴσιν*, 'to tie,' *Intususceptio*. Constriction or twisting of the intestines.—Celsus. The *Colica Ileus*. See *Enteritis* and *Ileus*.

CHORDEE', *Cordee'*, *Chorda Ven'eri* seu *Penis*, *Phallanculo'sis*, *Phallogamp'sis*, *Chordæ*, *Chorda'ta Gonorrhoe'a*. A painful affection of the penis, attending gonorrhoea, and happening chiefly at night. It occurs when the violence of the inflammation is such, that the urethra cannot expand as much as the corpora cavernosa during erection, and consequently the glands seem painfully drawn downwards. The application of cold water topically, and the administration of opium internally, palliate it.

CHORE'A, *Ch. Sancti Viti* seu *Sancti Modesti* seu *gesticulato'ria*, *Choromania*, *Syn'clonus saltans*, *Tripudia'tio spastica*, from *χορεία*, 'a dance,' sometimes called *Scelotyr'bæ*, *S. Tarantismus*, *Viti Saltus*, *Saltatio SANCTI VITI*, *Ballis'mus*, *Convulsio habitua'lis*, *Siph'ia prava*, *Syn'clonus Choreæ*, *Myotyr'bæ*, *Choreoman'ia*, *Orchestromania*, *Epileps'ia Saltatoria*, *Morbus Saltatorio'rius* seu *gesticulato'rius* seu *Lasciv'us*, *Illecan'osus*, *St. Vitus's Dance*, *Viper's Dance*, because the movements resemble dancing; (F.) *Chorée*, *Danse de SAINT WIT*, *Danse de ST. GUY*. The characteristics are:—irregular and involuntary motions of one or more limbs, and of the face and trunk. It is a disease which usually occurs before puberty; and is generally connected with torpor of the system, and of the digestive organs in particular. Its duration is long, but it is usually devoid of danger; although frequently but little under the control of medicine. The spasms do not continue

during sleep. The indications of treatment are: to strengthen the general system, and stimulate the intestinal canal. Purgatives, once or twice a week, and chalybeates, with appropriate regimen, will fulfil these.

CHOREA, **CHRONIC**, see *Tic*—c. *Faciei*, *Tic*—c. *Gesticulatoria*, *Chorea*—c. *Partial*, see *Tic*—c. *Sancti Modesti*, *Chorea*—c. *Sancti Valentini*, *Tarantismus*—c. *Sancti Viti*, *Chorea*.

CHORÉE, *Chorea*.

CHORE'GIA, from *χορος*, 'a troop of dancers and singers,' and *αγω*, 'I lead.' Hippocrates uses this term for the whole of the apparatus necessary for a physician or surgeon.

CHOREOMANIA, *Chorea*.

CHORIODEITIS, *Chorioide'itis*, *Chorioidi'tis*, *Chorioidi'tis*, (F.) *Chorioidite*, from *χοριοειδής*, 'the choroid,' and *itis*. Inflammation of the choroid coat of the eye.

CHORIODES, *Choroid*.

CHORIOIDEA, *Choroid*.

CHORIOIDES, *Choroid*.

CHO'RION, *χοριον*, 'skin,' from *χωρειν*, 'to contain,' include; (?) *Camis'ia Fœtalis*: the chemise or shirt of the fœtus in utero. A thin, transparent membrane, formerly confounded with the decidua, which surrounds the fœtus in utero on every side, and appears to be developed from nucleated cells formed in the Fallopian tube. Some histologists, however, consider, that it exists in the ovary. The general opinion is, that it is formed as above described; and perhaps, also, from the zona pellucida, which disappears in the tube. In the uterus, villous prolongations are formed on its surface, which have given it, with more recent writers, the name 'shaggy chorion.' These villi are probably the agents of the absorption of nutritive matter furnished from the lining membrane of the uterus.

By many anatomists, the chorion is considered to be formed of two layers; the outer, called by Burdach *Exocho'rion*: the inner, *Endocho'rion*. By others, the distinction of laminæ is denied. It is exterior to the amnion.

CHORION also means the true skin. See *CUTIS*. By some, the *Zona pellucida* is so called.

CHORION FUNGOSUM, *Decidua*—c. *Reticulatum*, *Decidua*—c. *Shaggy*, see *Chorion* and *Decidua*—c. *Spongiosum*, see *Decidua*—c. *Spongy*, see *Decidua*—c. *Tomentosum*, see *Decidua*.

CHORIONITIS. Induration of the Cellular tissue.

CHOROID, *Chorio'deus*, *Chorio'des*, *Chorioi'des*, *Chorio'des* from *χοριον*, 'the chorion,' and *ειδος*, 'shape,' 'resemblance.' Several parts are so called, which resemble the chorion in the multitude of their vessels.

CHOROID MUSCLE, *Ciliary muscle*.

CHOROID PLEXUS, *Plexus chorio'des* seu *chorio'deus* seu *reticul'a'ris*, *Vermes cer'ebri*. Two membranous and vascular duplicatures of the pia mater, situate in the lateral ventricles. They are fixed to the *Tela chorioidea* by one edge, and are loose and floating at the other.

CHORIOIDEA TELA, *Plexus chorio'deus medius*, (F.) *Toile chorio'dienne*. A kind of vasculomembranous prolongation of the pia mater, which lines the lower surface of the fornix united with the corpus callosum. It is stretched above the third ventricle and covers the posterior commissure and corpora quadrigemina. Anteriorly, the *tela chorioidea* is continuous with the plexus chorioides.

CHORIO'DEA seu **CHORIO'DEA TU'NICA**, *Ch. Membra'na*, or simply the *Choroid*, *Tunica vasculo'sa* seu *acini'form'is* seu *rhagio'des Oculi*. (F.) *Membrane chorioide*, *Chorioide*. A thin membrane, of a very dark colour, which lines the sclerotica

internally. The part behind the iris is called *Uvea*. It is situate between the sclerotica and retina, has an opening, posteriorly, for the passage of the optic nerve; and terminates, anteriorly, at the great circumference of the iris, where it is continuous with the ciliary processes. According to Ruysch, the choroid consists of two layers, to the innermost of which his son gave the name *Tunica seu Membrana Ruyschiana*, *M. chorio-capillaris*, (F.) *Membrane Ruyschienne*. The internal surface of the membrane is covered with a dark pigment, consisting of several layers of pigment cells, called *Pigmentum nigrum*, *Stratum pigmenti*, *Ophthalmochroites*, *Æthiops animal*, *Mélaine*, *Mélanine*, *Matière ou Principe de la Mélanose*, (F.) *Enduit choroidien*. Its use seems to be, to absorb the rays of light after they have traversed the retina.

CHOROÏDEÆ VENÆ, *Venæ Gale'ni*, (F.) *Veines choroïdiennes*. Two veins, that creep along the tela choroidea; into which almost all those of the lateral ventricles, of the upper part of the cerebellum, of the pineal gland, and the corpora quadrigemina open. The Venæ Galeni open into the *sinus quartus* or *fourth sinus*.

CHOROÏDE, CEINTURE BLANCHE DE LA, Ciliary ligament—*c. Commissure de la*, Ciliary ligament.

CHORODITIS, Choriodeitis.

CHOROÏDO-RETINITIS. Inflammation of the choroid and retina.

CHOROMANIA, Chorea.

CHOSÉS CONTRE NATURE, *Res contra naturam*—*c. Naturæ*, *Res naturales*—*c. non Naturæ*, *Res non naturales*.

CHOSIS, Arenatio.

CHOU, Brassica—*c. Cabus*, Brassica capitata—*c. Crœtæ*, Sauer Kraut—*c. Fleur*, Brassica Florida, Cauliflower excrecence—*c. Marin*, Convolvulus soldanella—*c. Navet*, Brassica rapa—*c. Pommé*, Brassica capitata—*c. Potager*, Brassica—*c. Roquette*, Brassica crua.

CHOWDER, a favorite New England dish, made of fish, pork, onions, and biscuit, stewed together.

In Devonshire, England, *chowder* is a fish-seller.—Halliwell. In the west of England *chowder beer* is a liquor made by boiling *black spruce* in water, and mixing it with molasses.—Webster.

CHREMMMA, Sputum.

CHREMPISIS, Exspuition.

CHRISIS, from *χρῶ*, 'I anoint.' The action of anointing. Inunction.

CHRISMA, same etymon. The act of anointing. The salve or liniment used, *Chrism*, *Creme*, *Prurigo*.

CHRISTI MANUS. Troches prepared from refined sugar boiled in rose-water with or without prepared pearls.

CHRISTOPHER, HERB, *Actæa spicata*.

CHRISTOPHORIANA SPICATA, *Actæa spicata*.

CHRISTOS, *χριστός*, from *χρῶ*, 'I anoint.' Any medicine applied under the form of liniment or ointment.

CHROA, *Chroæ*, *Chroma*. Colour in general. The surface of the body. The skin.

CHROMA, Chroa.

CHROMATISME, Aberration of refrangibility.

CHROMATOG'ENOUS, from *χρῶμα*, *χρῶμα-ros*, 'colour,' and *γενῶ*, 'I make.'

CHROMATOG'ENOUS APPARATUS. A supposititious glandular apparatus for producing the colouring matter of the skin, composed of a glandular or secreting parenchyma, situate a little below the papillæ, and presenting special excre-

tory ducts, which pour out the colouring matter on the surface of the true skin.—Breschet.

CHROMATOMETABLEPSIA, *Achromatopsia*.

CHROMATOPHO'BIA, from *χρῶμα*, *χρῶμα-ros*, 'colour,' and *φῶβος*, 'dread.' Morbid sensibility to certain colours.

CHROMATOPSEUDOPSIA, *Achromatopsia*.

CHROMATOPSIA, *Chromopsia*.

CHROMIC ACID, *Ac'idum Chro'micum*, (F.) *Acide chromique*. Obtained by crystallization from a mixture of *bichromate of potassa*, and *oil of vitriol*. It has been used as an escharotic in external hemorrhoids.

CHROMIDROSIS, from *χρῶμα*, 'colour,' and *ῥῖσος*, 'sweet.' Abnormal coloration of the perspiratory secretion.

CHROMOP'SIA, *Chromop'ia*, *Chromatop'ia*, *Chrotop'ia*, *Chrup'ia*, *Crop'ia*, *Virus coloratus*, *Suffusio colorans*, *Chro'p'sy*, *Chromop'sy*, *Iridescent vision*, from *χρῶμα*, 'and *ὄψις*, 'vision.' A state of vision in which a colored impression, as of the rainbow, is made on the retina. Said to be occasionally observed in jaundice.

CHRONIC, *Chron'icus*, *Chro'nicus*, *Polychro'nus*, *Inceteratus*, *Anti'quus*, (F.) *Chronique*, from *χρῶς*, 'time.' Of long duration.

CHRONIC DISEASES, *Morbi chronici*, *Macronos'ia*, *Macro'sia*, (F.) *Maladies Chroniques*, are those whose duration is long, or whose symptoms proceed slowly. The antithesis to *chronic* is *acute*.

CHRONIC'ITY, *Chronic'itas*, (F.) *Chronicité*. The state of a chronic malady.

CHRONIQUE, Chronic.

CHRONO, from *χρῶς*, 'time.' A prefix to terms denoting inflammation of a part, to show that such inflammation is chronic.—Piorry.

CHRONOGYNIA, Menstruation.

CHRONO-HÉPATITE, Hepatitis, chronic.

CHRONO-NÉPHRITE, Nephritis (chronic).

CHRONO-THERMAL, from *χρῶς*, 'time,' and *θερμῶν* 'heat.' Relating to time and temperature. An epithet given to a fanciful 'system' by Dr. Samuel Dickson, which maintains, that there can be no increase or diminution of temperature without motion; no motion without time; that motion consists in attraction and repulsion; that attraction and repulsion are peculiar to electric action; and hence, that medicines must change the motions of the system, and be electrical in their operation.

CHROOPSY, Chromopsia.

CHROTOPSIA, Chromopsia.

CHRUPSIA, Chromopsia.

CHRYSALEA, Nitro-muriatic acid.

CHRYSAN'THEMUM, *Calendula officinalis*.

CHRYSAN'THEMUM LEUCANTHEMUM, from *χρῶς*, 'gold,' and *ανθεῶν*, 'a flower.' The *Or-eye daisy*, *Daisy*, *Whiteweed*, *Goldens*, *Maudlinwort*, *Bellis major seu pratensis*, *Baythall'mon majus*, *Leucanthemum vulgare seu Bellidiod'es*, *Matricaria*, *Consolida media*, *Oculus Bovis*, (F.) *Chrysanthème*, *Chrysène*, *Grand Marguerite des prés*. *Ord. Compositæ*. The flowers and herbage slightly acid; and were once used in pulmonary diseases.

CHRYSAN'THEMUM PARTHENIUM, *Matricaria parthenium*.

CHRYSE, from *χρῶς*, 'gold.' The name of a yellow plaster, described by Paulus of Ægina, and composed of thus, alum, lead, colophony, resin, oil, and orpiment, boiled in vinegar.

CHRYSENE, *Chrysanthemum leucanthemum*.

CHRYSITIS, see *Plumbi oxidum semivitreum*.

CHRYSOBALANUS GALENI, see *Myristica mosehata*.

CHRYSOCALIS, *Matricaria*.

CHRYSOCHALCOS, Brass.

CHRYSOCOLLA, Borax.

CHRYSOCOMA, Millefolium.

CHRYSolACHANUM, *Chenopodium bonus* Heriencus.

CHRYSolITHUS, *Chrysolite*, from χρυσος, 'gold,' and λίθος, 'stone.' A precious stone, of a golden colour, regarded by the ancients as cardiac, cephalic, &c.

CHRYsOMELIA, see *Citrus aurantium*.

CHRYsOPHYLLUM BURANHEM, Monesia.

CHRYsOPHYLLUM CAINITO, from χρυσος, 'gold,' and φύλλον, 'a leaf.' *Cainito*, *Sideroxylon*, *Broad-leaved Star-apple*. A tree of the Antilles, Ord. Sapotaceæ, which produces one of the best fruits of the country. There are several varieties of it.

CHRYsOPHYLLUM GLYCIPLHEUM, Monesia.

CHRYsOPUS, Cambogia.

CHRYsOS, Gold.

CHRYsOSPERMUM, *Sempervivum tectorum*.

CHRYZA FIBRAUREA, Coptis.

CHTHONOPHAGIA, *Cucheeia Africana*, *Maldacia seu Pica Africana*, *Leucophlegma* seu *Chlorosis Ethiopia*, *Dirt-eating*, (F.) *Mal d'Estomac* from χθον, 'earth,' and φάγω, 'I eat.' A disorder of the nutritive functions observed amongst the negroes of the South and of the West Indies, in which there is an irresistible desire to eat dirt. It is accompanied by most of the signs of chlorosis.

The Ottomacs on the Orinoco, and the natives of portions of the Hudson's Bay territory, are *Chthonophagi* or 'earth eaters.' The earth is a form of stearite.

CHU, Choa or Chus, χους, *Chocus*. A liquid measure amongst the Greeks, answering to the *Congius* of the Romans, and containing six sextarii, or twelve Attic cotylæ, or nine pints.—Galen.

CHUMOSIS, Chemosis.

CHURNMILK, Buttermilk.

CHURRUS, see *Bangue*.

CHUTE, Protopus—c. du Fondement, Proctocœle—c. de la Matrice, Procidencia uteri—c. des Oeufs, see Parturition—c. du Rectum, Proctocœle.

CHYLAIKE, Chylous.

CHYLAR, Chylous.

CHYLARION, Chyle.

CHYLE, *Chylus*, *Succus nutritivus*, from χυω, 'I flow.' The word, in Hippocrates, means *Tisane* or *Decoction* of barley, *Chylarion*, χυλαριον. Galen first used it in its present sense—i. e. for a nutritive fluid, extracted by intestinal absorption, from food which has been subjected to the action of the digestive organs. It is of a whitish appearance; and is formed from the chyme in the duodenum, and the rest of the small intestines, by the chyloferous vessels, which arise at the mucous surface of the intestine. Along these it passes through the mesenteric glands to the thoracic duct, and is finally poured into the left subclavian. It is composed, like the blood, of a fluid—*liquor chyli*—and of *Chyle corpuscles* or *globules*, the average size of which is about 1-1600th of an inch. See *Chyme*.

CHYLE CORPUSCLES, see *Chyle*.

CHYLETIX, Chylous.

CHYLIFEROUS, *Chylifer*, *Chyliferus*, *Chyliphorus*, (F.) *Chylifère*, from *chylus*, 'chyle,' and *ferre*, 'to carry.' Chyle-bearing.

CHYLIFEROUS VESSELS, *Vasa Chylifera* seu *Chylifera*, *Via chylifera*, *Vena lactea*, *Vasa lactea*, *Lacteals*, (F.) *Vaisseaux chylifères*, *V. Lactés*. Vessels which convey the chyle from the intestines to the thoracic duct. They have also been called *galactophorous vessels*.

CHYLIFICATION, *Chylificatio*, *Chylosis*, *Chylopoesis*, *Præparatio chyli*, from *chylus*,

'chyle,' and *facere*, 'to make.' Formation of chyle during the digestive processes. It has, at times, been applied to the changes produced on the aliment in the small intestine; but is now restricted to the act of forming chyle at the radicles of the chyloferous vessels.

CHYLINE, Cyclamen.

CHYLISMA, *Succus expressus*.

CHYLIS'MUS, from χυλος, 'juice.' The act of expressing the juice of vegetables, &c.

CHYLOCYSTIS, *Receptaculum chyli*.

CHYLODES, Chylous.

CHYLODIABETES, Chyluria.

CHYLODIARRHŒA, *Cœliac flux*.

CHYLODOCHUM, *Receptaculum chyli*.

CHYLOGRAPHY, from χυλος, 'chyle,' and γραφή, 'a description.' A description of the anatomy, &c., of the chyloferous vessels.

CHYLOPOESIS, Chylification.

CHYLOPOËTIC, *Chylopoëticus*, *Chylopoëus*, from χυλος, 'chyle,' and ποιω, 'I make.' Relating to or connected with the formation of chyle. Chiefly applied to the organs immediately concerned in it; as the stomach, intestines, omenta, and mesentery. *Assistant Chylopoetic*—applied to viscera which aid in the formation of chyle, as the liver and pancreas.

CHYLORRHŒA, *Cœliac flux*—c. Pectoris, *Chylorhorax*—c. Renalis, *Chyluria*—c. Urinalis, *Chyluria*.

CHYLOSIS, Chylification.

CHYLOSTAGMA DIAPHORETICUM MINDERERII (χυλος, 'juice,' and σταζω, 'I distil.') A compound prepared by distilling the *theriac* of Andromachus, the *mithridate* of Damocrates, and other alexipharmics, &c. It is nearly the same preparation as the *Aqua Theriacalis Bezoardica*.

CHYLOTHORAX, *Pleurorrhœa chylousa*, *Chylorrhœa Pectoris*, *Hydrothorax chylous*; from χυλος, 'chyle,' and θώραξ, 'the chest.'—Effusion of chyle into the chest, owing to the rupture of a chyloferous vessel.

CHYLOUS, *Chylar*, *Chylous* seu *Chylaris*, *Chylodes*, (F) *Chyleux*, *Chylaire*. Relating to the chyle; or having some analogy to that fluid.

CHYLURIA, *Diabète lacteus* seu *chylous*, *Chylodiabète*, *Galaeturia*, *Fluxus cœliacus per Renes*, *Pyeluria lactea* seu *Chylous*, *Cœliaca urinaria* seu *renalis*, *Chylorrhœa urinaria* seu *renalis*, from χυλος, 'chyle,' and ούρον, 'urine.' (F.) *Diabète chyleux*. A discharge of milky urine, without any apparent lesion of the kidneys or bladder.

CHYLUS, Chyle, *Decoction*, *Succus*.

CHYME, *Chymus*, χυμος, 'juice,' from χυω, 'I flow.' The pulp, formed by the food, mixed with the supra-diaphragmatic and gastric secretions, after it has been for some time in the stomach. In this it continues until it reaches the biliary and pancreatic ducts, which open into the duodenum, where the conversion into chyle occurs, which is absorbed by the chyloferous vessels,—the excrementitious portion of the food traversing the large intestine to be evacuated *per anum*. Castelli asserts, that *Chyme* and *Chyle* were used in an opposite sense by the ancients, to that accepted at present.

CHYMI, Humours.

CHYMIA, Chymistry—c. Organica, Chymistry, organic—c. Pharmaceutica, see Chymistry.

CHYMIATER, *Chimioter*, *Chemia'ter*, from χυμια or χημια, 'chymistry,' and ιατρος, 'a physician.' *Intro-chymicus*. A chemical physician.

CHYMIATRIA, *Chimiatria*, *Chemiatria*, *Intro-chemia*, *Medicina spagyrica*, *Arts Chymiatrica*, (F.) *Chimistrie*, *Chimisme* (of some), from χυμια or χημια, 'chymistry,' and ιατρεια, 'cure.' The art of curing by chemical means.

CHYMICAL, *Chemical*, *Chemicus*, *Chemo'-licus*. A medicine formed by the aid of chymistry, in contradistinction to *Galenic*.

CHYMICO-HISTOLOG'Y, *Chymico-histolog'ia*, *Chem'ico-histology*. The doctrine of the organic chemistry and morphology of tissues.

CHYMICOPHANTA, *Chymist*.

CHYMICUS, *Chymical*, *Chymist*.

CHYMIE, *Chymistry*.

CHYMIFICAT'ION, *Chymifica'tio*, *Chymo'-ria*, from *χυμος*, 'juice,' and *facere*, 'to make.' Formation of chyme.

CHYMISM, *Chymism'us*, *Chemism'us*, (F.) *Chimisme*. The abuse of chymistry in its application to the science of organized bodies. By some used synonymously with *Chymiat'ria*. Also, an account of the chemical phenomena appertaining to a function—as the *chemismus* of respiration.

CHYMIST, *Chem'ist*, *Chemicus*, *Chymico-phan'ta*, *Chym'icus*, (F.) *Chimiste* ou *Chymiste*. One acquainted with chymistry. In Great Britain it has, also, the signification of "one who sells chemicals."

CHYMISTE, *Chymist*.

CHYMISTRY, *Chem'istry*, *Chemi'a*, *Chymi'a*, *Chimi'a*, *Chementic'e*, *Chemot'ic'e*, *Philosoph'ia per ignem*, *Spagy'ria*, *Pyrotechn'ia*, *Pyrosoph'ia*, *Ars hermet'ica* seu *mago'rum* seu *separato'ria* seu *spagy'ria*, *Archim'd'gia*, from *χυμος*, 'juice,' or from Arab, *chem'a*, 'a secret.' (F.) *Chimie* ou *Chymie*. A branch of the natural sciences, whose object is to investigate the nature and properties of bodies, simple and compound, inorganic and organized; and to study the force or power, by virtue of which every combination is effected. It investigates the action between the integrant molecules or atoms of bodies.

Organ'ic Chemistry, *Chymi'a organ'ica*, *Organo-chemi'a*, is the chymistry of organized substances, --animal and vegetable. *Animal Chym'istry*, *Zoöch'em'y* or *Zoöch'ymy*, *Zoöchem'i'a*, is the chymistry of substances afforded by the dead or living animal body. This branch of chymistry has been farther subdivided into *physiological*, when it considers the changes produced in organized bodies in health, --*pathological*, when it regards those produced by organic or other diseases. *Anthropoch'ymy*, *Anthropochem'i'a*, is the chymistry of the human body. Chymistry is called *Therapeutical* or *Pharmaceut'ical*, *Pharmaco-chymi'a*, *Chymi'a pharmaceut'ica*, when it is engaged in the analysis of simple medicines; in improving the prescribing and preparing of chemical and Galenic medicines; in the means of preparing them, and detecting adulterations, &c. *Hygien'ic Chym'istry* is that which is applied to the means of rendering habitations healthy, of analyzing the air we breathe, preventing the occurrence of disease, pointing out healthy aliments, and appreciating the influence of professions, &c., on the health of man. All these different subdivisions, with vegetable chymistry, are, at times, included under the head of *Medical Chym'istry*, *Phytochym'istry*; at others, the term comprehends only the *Animal*, *Vegetable* and *Pharmaceutical* subdivisions. *Vital Chemistry*, *Biochymi'a*, is that which is exerted under the influence of vitality. *Histocho'ymy*, *Histochem'i'a*, *Histo-chem'istry*, is the chemistry of the tissues, and *Phlegmatoch'em'y*, that of the animal humours, (Lehmann).

A knowledge of chemistry is of great importance to the physician. Many of the functions are of a chemical nature: many diseases require a chemical mode of treatment; and, without an acquaintance with it, two or more substances might be given in combination, which, by forming a chemical union, might give rise to other compounds, possessing very different virtues from the

components taken singly, and thus the prescriber be disappointed in the results.

CHYMISTRY, *ANIMAL*, see *Chymistry* — c. *Hygienic*, see *Chymistry* — c. *Medical*, see *Chymistry* — c. *Organic*, see *Chymistry* — c. *Pharmaceutic*, see *Chymistry* — c. *Therapeutical*, see *Chymistry* — c. *Vegetable*, see *Chymistry* — c. *Vital*, see *Chymistry*.

CHYMOCHEZIA, *Cœliac flux*.

CHYMOPLANIA, (G.) *Chymoplanien*, *Dyschymosen*, from *χυμος*, 'juice,' and *πλανω*, 'wandering.' A transposition of secretions; — a family of diseases in the classification of Fuchs, which includes icterus, uroplania, menoplania and galactoplania.

CHYMORRHEA, *Cœliac flux*, *Lientery*.

CHYMOSIN, *Pepsin*.

CHYMOSIS, *Chemosis*, *Chymification*.

CHYMOUS, *Chymo'sus*, (F.) *Chymeux*. Relating to, or resembling chyme.

CHYMOZEMIA, *Hypercæmia*.

CHYTLEN, RADIX. A cylindrical root, bitter and inodorous, brought from China. It is held by the Chinese to be stomachic. —Murray.

CHYT'LOX, *χυτλον*, from *χυω*, 'I pour out.' A liquid formerly used for rubbing the body after bathing.

CIBARIUM, *Aliment*.

CIBARIOUS, *Esulent*.

CIBARIUS PANIS, 'Coarse bread.' Bread made of second flour. —Celsus.

CIBAT'IO, *Trophê*. The taking of food. In Pharmacy, it is the same as *Incorporation*.

CIBUS, *Aliment* — c. *Albus*, *Blancmanger* — c. *Deorum*, *Asafetida*.

CICATRICE, *Cicatrix*.

CICATRICES OVARIORUM, *Stigmata ovariorum*.

CICATRIZ'ULA. Diminutive of *Cicatrix*. A small *cicatrix*, *Stigma*, (F.) *Cicatriceule*. The term is, also, applied to a small white spot, called the *tread*, *chal'asa*, *chala'zium*, observable at the surface of a fecundated egg. See *Molecule*.

CICATRISAN'TIA, *Epsulo'tica*, *Synulo'tica*, *Apulo'tica*, *Catulo'tica*, *Ulo'tica*, (F.) *Cicatrisants*. Remedies formerly considered to be capable of producing cicatrization.

CICA'TRIX, *Cæca'trix*, *Ule*, *Oulê*, *Cicatrice*, from *cæcare*, 'to conceal,' because it conceals the wound. (F.) *Cicatrice*. A scar. A seam. (Sc.) *Arr*. The union of parts, which have been divided. A scar or formation, of a reddish colour, afterwards whitish, and of variable thickness, which takes place at the surface of wounds or ulcers after their cure. A cicatrix may vary much in shape, consistence, and thickness. The cicatrix of a bone is called *Callus*. A *vic'iosa cicatrix*, (F.) *Cicatrice vicieuse*, is one which interferes with the action of the parts on which it occurs. The *scars*, (F.) *Coutures*, after small-pox, are called *Pits* or *Pock-marks*, (Sc.) *Pock-arrs*, (F.) *Coutures par la petite vérole*. See *Pock-mark*.

CICATRIX VARIOLÆ, *Pock-mark*.

CICATRIZA'TION, *Cicatrisa'tio*, *Epsulo'sis*, *Synulo'sis*. The process by which a cicatrix is formed. Every tissue, except the nails, epidermis, hair, and enamel is, probably, capable of cicatrization.

CICELY, SWEET, *Chærophylllum odoratum*, *Osmorrhiza longistylis*, *Scandix odorata*.

CICERARIETINUM. The *Cicer* plant, *Erb'bin'thus*, (F.) *Cicérole*, *Pois Chiche*, *Garvance*, *Pesette*. The seeds are ground into flour, and used as bread in some countries.

CICER LENS, *Ervum lens*.

CICERA TAR'TARI. Small pills of turpentine and cream of tartar — of the size of a vetch or *cicer*.

CICERBITA, *Sonchus oleraceus*.

CICÉROLE, *Cicer arietinum*.

CICHO'RIMUM ENDIV'IA. The systematic name of the *Endive*, *Endiv'ia*, *Endi'va*, *Intubum*, *Intybum* (*Antiq.*), *Scariola*, *Intybus hortensis*, (*F.*) *Chicorée des Jardins*, *Scariola*. *Ord.* *Chicoraceæ*. *Sex. Syst.* Syngenesia Polygamia æqualis. It is a common pot herb, and is eaten as salad.

CICHO'RIMUM INTYBUS, *Serie*, *Seriola*, *Intubum erradicum*. The systematic name of the *Wild Sarcory*, *Wild Chic'ory*, *Chic'ory*, *Chic'ory*, *Wild Endive*, *Ambuleia*, *Heliotropion*, *Catanan'cè*, *Chic'o'reum*, (*F.*) *Chicorée sauvage*. It is bitter, and was once given as a tonic. The root, roasted and ground, is often used instead of, or mixed with, coffee.

CICHORY, *Cichorium intybus*—*c.* *Wild*, *Cichorium intybus*.

CICI, *Ricinus communis*.

CICINDE'LA, *Lam'pyris Noctilu'ca*, *Nited'ula*. The *Glow-worm*. (*F.*) *Ver luisant*. This insect was once thought to be anodyne and lithon- triptic.

CICIS, see *Quercus infectoria*.

CICON'GIUS; an ancient measure, containing 12 pints.

CICUTA, *Conium maculatum*.

CICU'TA AQUAT'ICA, *C. viro'sa*, *Cicuta'ria aquat'ica*, *Corian'drum cicuta*, *Water Hemlock*, *Cowbane*, (*F.*) *Cigué aquatique ou vireuse*, *Cicutaire aquatique*. *Family*, *Umbelliferae*. *Sex. Syst.* *Pentandria Digynia*. A violent poison, often eaten by mistake for *Wild Smallage*, *Apium Graveolens*. It produces tremors, vertigo, burning at the stomach, and all the symptoms occasioned by the *Narcotico-acrid* class of poisons.

CICU'TA MACULA'TA, (*F.*) *Cigué d'Amérique*, *American water hemlock*, *American Hemlock*, *Snakeweed*, *Death of man*, *Water parsley*, *Poison root*, *Wild Hemlock*, *Children's bane*, *Spotted Cowbane*, is analogous in botanical character and medical properties to the European species. See *Conium maculatum*.

CICUTA MAJOR, *Conium maculatum*—*c.* *Major foetida*, *Conium maculatum*—*c.* *Stoerkii*, *Conium maculatum*—*c.* *Terrestris*, *Conium maculatum*—*c.* *Virosa*, *Cicuta aquatica*—*c.* *Vulgaris*, *Conium maculatum*.

CICUTAIRE AQUATIQUE, *Cicuta aquatica*.

CICUTARIA, *Chærophyllym sylvestre*—*c.* *Aquatica*, *Cicuta aquatica*, *Phellandrium aquaticum*—*c.* *Odorata*, *Chærophyllym odoratum*.

CICUTINE, see *Conium*.

CIDER, *Poma'ceum*, (*F.*) *Cidre*. This word is said to have been formerly written *sidre*; and to have come from *Sic'era*, *ikepa*, which signifies any kind of fermented liquor other than wine. It is made from the juice of apples, and, when good, is a wholesome drink.

CIDRE, *Cider*.

CIERGE DE NOTRE DAME, *Verbascum thapsus*.

CIGAR', *Cigar'ra*, (*S.*) *Cigarro*, (*F.*) *Cigare*. A small tubular roll of tobacco, used for smoking. **MEDIC'INAI** or **MED'ICATED CIGARS**, (*F.*) *Cigares Médicinaux*, are made of dried plants, with, or without, the addition of medicinal articles. When rolled in paper, they form *cigarettes*, which may be used with or without combustion, as in the case of the *Cigarettes de Camphre*. The *Cigarettes arsénicales* de Boudin are made of paper impregnated with an aqueous solution of arsenious acid, or of arseniate of soda, and dried.

CIGARETTE, see *Cigar*—*c.* *Arsénicale* de Boudin, see *Cigar*—*c.* de *Camphre*, see *Cigar*.

CIGARRA, *Cigar*.

CIGNUS; an ancient measure, which contained about two drachms.

CIGUÉ AQUATIQUE, *Cicuta aquatica*—*c.* *d'Amérique*, *Cicuta maculata*—*c.* *d'Eau*, *Phellandrium aquaticum*—*c.* *Grande*, *Conium maculatum*—*c.* *Ordinaire*, *Conium maculatum*—*c.* *Officinale*, *Conium*—*c.* *Petite*, *Æthusa cynapium*—*c.* *Vireuse*, *Cicuta aquatica*.

CIL'IA (singular *Cilium*), *Blephar'ides*, *Epice'rides*, *Pili palpebra'rum*. The *eyelashes*, (*Sc.*) *Winkers*. The hairs on the eyelids. (*F.*) *Cils*. Their use seems to be, to prevent the entrance into the eye of light bodies flying in the atmosphere; and to diminish, in certain cases, the intensity of light. Also, the tarsi. Also, a peculiar sort of moving organs, resembling small hairs, *vibratory* or *vibratile cil'ia*, *Cil'ia vibrato'ria*, (*F.*) *Cils vibratiles*, which are visible with the microscope in many animals. These organs are found on parts of the body which are habitually in contact with water, or other more or less fluid matters, and produce motion in these fluids, impelling them along the surface of the parts. Cilia have been found to exist in all vertebrated animals except fishes, having been discovered on the respiratory, uterine, and other membranes of mammalia, birds, and reptiles.

The terms "*vibratory motion*" and "*ciliary motion*" have been used to express the phenomena exhibited by the moving cilia; and it is probable, that this motion is concerned in the progression of fluids along the membranes. As yet, the motion has only been observed in the direction of the outlets of canals.

CILIAIRE, *Ciliary*.

CILIARY, *Cilia'ria*, (*F.*) *Ciliaire*. Relating to the eyelashes, or to *cilia*. This epithet has, also, been applied to different parts, which enter into the structure of the eye; from the resemblance between some of them (*the ciliary processes*) and the eyelashes.

CILIARY ARTERIES, *Arte'ria cilia'res*, (*F.*) *Artères ciliaires*. These are furnished by the ophthalmic artery. They are distinguished into, 1. *Short* or *posterior* (*Art. uéales*—*Chauss.*) 30 or 40 in number, which are distributed to the ciliary processes. 2. *Long*, (*Art. Iriennes* of *Chauss.*) two in number, which, by the anastomoses of their branches, form two arterial circles at the anterior surface of the iris; and 3. The *anterior*, *Arte'ria cilia'res antero'ries* of Haller, the number of which is variable. These pierce the sclerotic a few lines from its union with the cornea; and are principally distributed to the iris.

CILIARY BODY, *Corpus Cilia're*, *Nexus Stamin'eus Oculi*, *Coro'na Cilia'ris*, *Ciliary Disc*, (*F.*) *Corps ciliaire*. A ring of the choroid surrounding the crystalline in the manner of a crown; placed behind the iris and the ciliary circle. It resembles the disk of a radiated flower, and is formed by the union of the ciliary processes. See *Ciliary Muscle*.

CILIARY CANAL, *Canal of Fonta'na*. A small, extremely narrow circular space, formed between the ciliary circle, the cornea, and the sclerotic. It can be filled with injection, and it is not certain that it is not the cavity of a blood-vessel.

CILIARY CIRCLE, *Ciliary ligament*—*c.* *Disc*, *Ciliary body*—*c.* *Ganglion*, *Ophthalmic ganglion*.

CILIARY LIG'AMENT, *C. Circle* or *Ring*, *Ligamentum seu Inserti'tium cilia'rè seu Pridis*, *Plexus cilia'ris*, *An'nulus seu Cir'culus seu Orbic'ulus cilia'ris*, *A. cellulo'sus*, *Com'missure of the Uvea*, (*F.*) *Commissure de la Choroïde*—(*Ch.*) *Ligament on Cercle ciliaire*, *Cercle de la Choroïde*, *Ceinture blanche de la Choroïde*. A species of greyish ring, of a pulpy consistence, situate between the choroid, iris, and sclerotic. The internal surface of the choroid is uniform, until it approaches within ten lines and a half of the edge of the cornea; here a dentated line is ob-

served, termed *ora serrata*. The outer surface presents the *an'nulus al'bidus seu gangliiformis*, the anterior edge of which unites to the inner surface of the sclerotica and constitutes the *ciliary ligament*.

CILIARY or TARSAL MARGIN of the eyelids; (F.) *Bord ciliaire*. The edge in which the cilia or eyelashes are situate.

CILIARY MOTION, see *Cilia*.

CILIARY MUSCLE, *Musculus ciliaris*. The part of the orbicularis palpebrarum in the vicinity of the ciliary margin. Also, the greyish, semi-transparent structure behind the ciliary ligament and covering the outside of the ciliary body. By its contraction the ciliary processes, and with them the lens, must be drawn towards the cornea. It appears to be the same muscle as the *Tensor choroideæ*, *Protrahens lentis*, or *choroid muscle* of some anatomists.

CILIARY NERVES (*Nerfs triens*,—Chauss.), (F.) *Nerfs ciliaires*. These are 12 to 16 in number. They arise from the nasal nerve, and particularly from the anterior part of the ophthalmic ganglion; and unite in two fasciculi, which pass around the optic nerve, and pierce the sclerotica near the entrance of that nerve into the eye. They are lost in the ciliary ligament.

CILIARY PLEXUS, *C. Ligament*.

CILIARY PROCESSES, *Processus seu Rad'ii seu Striæ ciliares*, (F.) *Process ciliaires*, *Rayons sous-triens*—(Ch.). Triangular folds, sixty or eighty in number, placed at the side of each other, and radiating, so as to resemble the disk of a radiated flower. They are lodged in depressions at the anterior part of the vitreous humour. The uses of these processes are not known.

CILIARY RING, *Ciliary ligament*.

CILIARY STRIÆ are numerous pale, radiated striæ in the posterior portion of the *Corpus ciliare*, but so covered by the *Pigmentum nigrum* as not to be distinctly seen till the paint is removed. The ciliary processes are formed by these striæ.

CILIARY VEINS, (F.) *Veines ciliaires*, follow nearly the same course as the arteries. In the choroid they are so tortuous, that they have received the name *Vasa vorticosa*. They open into the ophthalmic vein.

CILIARY ZONE, *Zona seu Zonula seu Lam'ina Ciliaris*, *Membran'ula Cor'ona Ciliaris*. Under the corpus ciliare, the capsule of the vitreous humour sends off an external lamina, which accompanies the retina, and is inserted, with it, into the forepart of the capsule of the lens, a little before its anterior edge. This is the *Zonula ciliaris*, *Zonula Zinnii* or *Zonula of Zinn*, *Cor'ona Ciliaris*, *Orb'iculus Ciliaris*. It is of a striated appearance and circular form, and assists in fixing the lens to the vitreous humour.

CILIATED, *Ciliatus*, (F.) *Cilié*, from *cilia*. Provided with cilia—as "*ciliated epithelium*," the epithelium to which vibratory cilia are attached.

CILIUM, see *Cilia*.

CILLEMENT, *Nictation*.

CILLO. A name given by some authors to those whose upper eyelid is perpetually tremulous;—a trembling, which in some cases is called *Life's blood*. "To have life's blood in the eye," in other words, is to have this affection. Vogel calls it *Cillo'sis*.

CILLOSIS, *Cillo*.

CILS, *Cilia*—*c. Vibratilis*, see *Cilia*.

CIMEX, *C. lectularius*, *Acan'thia lectularia*, *Koris*, *kops*. The Bug; *Wall*, *House* or *Bed Bug* or *Chinche*. (F.) *Punaise*. Six or seven of these, given internally, are said to have prevented ague! There is scarcely anything which is sufficiently disgusting, that has not been exhibited for this

purpose, and with more or less success. The bug has also been esteemed emmenagogue.

CIMICIFUGA, *Actæa racemosa*.

CIMO'LIA PURPURES'GENS, *Terra Saponaria seu Fullonica*, *Fuller's Earth*. A compact bolar earth, employed in the arts. Used at times as a cooling application to inflamed nipples, &c.

CIMO'LIA TERRA, *Cimo'lia alba*, *Cimo'lius*, *Smectis*, *Smectris*; from *Κίμωλος*, an island in the Cretan Sea, where it is procured. It was formerly used as an astringent, &c.—Scribonius Largus, Pliny. Probably, the same as the last.

CINA CINA, *Cinchona*—*c. Levantica*, *Artemisia Santonica*.

CINABARIS, *Hydrargyri sulphuretum rubrum*.

CINABARIUM, *Hydrargyri sulphuretum rubrum*.

CIN'ABRA, *Grana*. The smell of a hog-scent. A rank smell, like that of the armpit, *Hicous alacrum*.

CINABRE, *Hydrargyri sulphuretum rubrum*.

CINÆDIA, *Masturbation*.

CINARA HORTENSIS, *Cynara scolymus*—*a. Scolymus*, *Cynara scolymus*.

CINCHONA. So called from the Spanish Viceroy's lady, the Countess de Cinchon, who was cured of fever by it at Lima, about 1638. Called also *Cortex seu Pulvis Jesuiticus*, *Jesuit's Bark* or *Powder*, *Cortex Patrum*, because it was introduced into Europe by the Jesuits; also *Pulvis Comitisæ* or the *Countess's Powder*, and *Cardinal del Lugo's Powder*, *Cortex Cardinalis de Lugo*, because he introduced it at Rome; and in France, *Talbor's Powder* and *English remedy*, because successfully used there by Sir Robert Talbor, who kept it a secret. It is the pharmacopœial name of several kinds of barks from various species of *Cinchona*, from the western coast of South America. *Order*, *Cinchonaceæ*, *Sex. Syst.* *Pentandria Monogynia*. Called, also, *Cortex*, *Bark*, *Peruvian Bark*, *Cortex China seu China*, *Chinchina*, *Polos de Calcutra*, *Kina Kina*, (*Bark of Barks*), *Kinkina*, *Cina Cina*, *Quina Quina*, *Quinquina*, *Magnum Dei domini*, (F.) *Quinquina*; and in Peru *Cascara* and *Cascarilla*. See *Cascara*.

CINCHONA AMYGDALIFOLIA, see *Cinchonæ cordifoliæ cortex*—*c. Boliviana*, see *Cinchonæ cordifoliæ cortex*—*c. Calisaya*, see *Cinchonæ cordifoliæ cortex*—*c. Micrantha*, see *Cinchonæ cordifoliæ cortex*—*c. Officinalis* (*Cortex Flavus*), *Cinchonæ cordifoliæ cortex*—*c. Pallida*, *Cinchonæ lancifoliæ cortex*—*c. Ovata*, see *Cinchonæ cordifoliæ cortex*—*c. Pitaya*, *Pitaya bark*—*c. Pubescens*, see *Cinchonæ cordifoliæ cortex*—*c. Scrobicula*, see *Cinchonæ cordifoliæ cortex*—*c. of Virginia*, *Magnolia glauca*.

CINCHONA CARIBÆÆ CORTEX, from *Exotema Caribæum*, *Caribæan* or *Saint Lucia Bark*, (F.) *Écorce de Saint Lucie*, *Quinquina Piton*, from *Exotema floribundum*; and the *Pitaya Bark*, *Quinquina bicolor*, from an exotema [?] or from *strychnos pseudoquina* [?], are useful substitutes for the cinchona of Peru. These are the most important spurious barks. They contain neither quina nor cinchonia.

CINCHONÆ CORDIFOLIÆ CORTEX, *Cortex flavus*, *Cinchonæ officinalis cortex flavus*, *China regia seu Calisaya*, *Cortex Chinæ regius seu sumu seu luteus*, *Yellow Royal* or *Calisaya Bark*, (F.) *Quinquina jaune* ou *jaune royal*, *Calisaya*. Odour aromatic; taste strong, bitter, astringent. Not rolled; often without the epidermis, which is very thick and inert; light, friable; fracture fibrous. Active principle *Quina*. The false or spurious *Calisaya* barks are those of *Cinchona*

Calisaya, variety *Josephiana*; *C. Boliviana*. *C. ovata*, variety *rufinervis*, called, in Peru, *Cascarilla Carabaya*—the Carabaya bark of commerce;—of *C. scrobiculata*, two varieties, *Cusco Bark* and *Bark of St. Ann*; of *C. pubescens*, which furnishes the *Cusco* or *Arica* bark; of *C. micrantha* and of *C. amygdalifolia*.

CINCHONÆ LANCEOLATÆ CORTEX, *Cortex Peruviana* seu *pallidus*, *Cinchonæ officinalis cortex canariensis*, *Cinchona pallida*, *Pale*, *Loxa* or *Crown Bark*, (F.) *Quinquina gris de Loxa*, *Quinquina Orange*. Its odour is aromatic; taste pleasant, bitter, and astringent. The pieces are rolled in double or single quills. Epidermis brown, cracked; fracture resinous. Internally of a cinnamon colour. Its active principle is *Cinchonia*.

CINCHONÆ OBLONGIFOLIÆ CORTEX, *Cortex ruber*, *Cinchonæ officinalis cortex ruber*, *Red Bark*, (F.) *Quinquina rouge*, (Sp.) *Cascarilla roxa* and *C. colorada*, of S. America. Odour and taste the same as the pale, but more intense: in large flat pieces, solid, heavy, dry; fracture short and smooth; of a deep brownish-red colour. Although this variety of bark is assigned to the *Cinchona oblongifolia* by some, it would seem, that nothing is certainly known as to its source. Active principles, *Cinchonia* and *Quinia*.

The last three are the only official varieties in the Pharmacopœia of the United States. There are many other varieties, however, which are genuine cinchona barks, and yet have not been considered worthy of an official position. The Edinburgh Pharmacopœia admits, indeed, *Cinchona cinerea*, *Grey bark*, *Silteer bark* or *Huanuco* or *Lima bark*, which is obtained around Huanuco in Peru, and, with the *Jaen* or *Ash bark* and the *Huamiles bark*, belongs to the class of pale or Loxa barks. Amongst the genuine but inferior barks are those brought from the northern Atlantic ports of South America, which, in commerce, are variously called *Pitaya*, *Bogota*, *Carthagena*, *Maracaybo*, and *Santa Martha barks*. The *Bogota* is also called *Fusa gasuga* and *Coquetta bark*.

All these barks are bitter, astringent, tonic, and eminently febrifuge. The yellow bark has been thought equal to any of the others, but the red contains more active principle. The discovery of their active principles is one of the most important gifts of modern chemistry. Still, in pernicious intermittents, the bark, in substance, is often demanded. It is employed in every disease in which there is deficient tone; but in cases where the stomach is much debilitated, the powder had better be avoided, in consequence of the woody fibre, which might disagree. Externally, it is used in enemata, gargles, &c., and in gangrenous ulcerations. When it excites nausea, an aromatic may be added to it; if purging, opium; if costiveness, rhubarb, &c. Dose, ℥ss to ʒj or more.

ESSENTIAL SALT OF BARK, as it is called, is an extract, prepared by macerating the bruised substance of bark in cold water, and submitting the infusion to a very slow evaporation.

CINCHONIA, Cinchonine—c. Tannate of, Quinine et cinchoninæ tannas.

CINCHONICINE, *Cinchonici'na*, *Cinchonic'ia*. When a salt of cinchonia is exposed to heat, with certain precautions, the alkaloid is changed into another, isomeric with itself, to which this name has been given. Cinchonidine, and its sulphate, possess the same medical properties as cinchonia and its salts. Dose, gr. v. to gr. xv.

CINCHONINE, *Cinchoni'na*, *Cinch'otin*, *Cincho'nia*. The active principle of *Cincho'na lanceolata*. An organic, crystalline alkali; of a white colour, and bitter, slightly astringent taste;

very soluble in alcohol and ether, but almost insoluble in water.

Sulphate of *Cinchonia*, which is formed directly from cinchonia, is soluble in water and alcohol. The action of the sulphate of cinchonia is similar to that of the sulphate of quinia: but it is less energetic, and consequently requires to be given in a larger dose.

CINCHONINE, TARTRATE OF, see Quinine, tartrate of.

CINCHONISM, Quinism.

CINCHOTIN, Quinidia.

CINCHINULUS. A little lock or curl of hair.

CINCHINUS. A curled or frizzled lock. The hair on the temples.

CINC'LICIS, *Cincli's'mus*, 'agitation; rapid and frequent motion.' The movement of the thorax in dyspnoea.—Hippocrates. It has been used, also, synonymously with nictation.

CINCLISMUS, Cinclisis.

CINEFACTIO, Incineration.

CINE'MA, *Cine'sia*, from κινεω, 'I move.' Motion.

CINERARIA MARITIMA, Achevon, Abiat.

CINERATION, Incineration.

CINERES CLAVELLATI, see Potash—c. Gravelati, see Potash—c. Russici, Potash of commerce.

CINEREUS, Cineritious.

CINERITIOUS, *Ciner'eus*, from cinerea, 'ashes;' (F.) *Cendré*. Of the colour of ashes. The cortical substance of the brain, and the vesicular neurine in general, have been so called. See Cortex Cerebri, and Neurine.

CINESIPATHY, Kinesipathy.

CINESIS, Cinema, Motion.

CINETH'ICS, from κινεω, 'I move.' The science of movements in general.

CINETIC, Motory.

CINETICA. Same etymon. Diseases affecting the muscles, and characterized by irregular action of the muscles or muscular fibres, commonly denominated *Spasm*. The 3d order in the class *Neurotica* of Good. Also, agents that affect the voluntary or involuntary motions.—Pereira.

CINETUS, Diaphragm.

CINGULARIA, Lycopodium.

CIN'GULUM, *Zone*, from cingo, 'I bind.' (F.) *Ceinture*. A cincture. A girdle. The part of the body, situate below the ribs, to which the girdle is applied. The waist. Herpes zoster.

CIN'GULUM HILDA'NI, *Zo'nula Hilda'ni*, (F.) *Ceinture de Hildane*. A leathern girdle formerly used for the reduction of luxations and fractures of the extremities.

CIN'GULUM MERCURIALE, *C. Sapientia* seu *Stultit'ia*. A woollen girdle, containing mercurial ointment. It was used as an antisyphilitic, and in diseases of the skin. (F.) *Ceinture de vif argent*.

CINGULUM SANCTI JOANNIS, *Artemisia vulgaris*.

CINIS, Santonine.

CINIS FÆCUM, see Potash—c. Infectorius, see Potash.

CINNABAR, Hydrargyri sulphuretum rubrum—c. Græcorum, see Calamus rotang.

CINNABRE, Hydrargyri sulphuretum rubrum.

CINNAMOMUM, *Laurus cinnamomum*—c. Album, *Canella Alba*—c. Aromaticum, see *Laurus cinnamomum*—c. Culilawan, *Laurus Culilawan*—c. Indicum, *Laurus cassia*—c. Magellanicum, *Winters aromaticæ*—c. Malabaricum, *Laurus cassia*—c. Sylvestre Americanum, *Nectandra cinnamomoides*—c. Zeylanicum, *Laurus cinnamomum*.

CINNAMON, see *Laurus cinnamomum*—c. Malabar, *Laurus cassia*—c. Wild, *Laurus cassia*.

CINON'OSI, from *κινω*, 'I move,' and *νοσος*, 'a disease.' Diseases of motion.

CINOPER, Hydargyri sulphuretum rubrum.

CINOPLANE'SIS, from *κινω*, 'I move,' and *πλάνησις*, 'a wandering about.' Irregularity of motion.

CINQUEFOIL, *Potentilla reptans* — c. Marsh, *Comarum palustre* — c. Norway, *Potentilla Norvegica*.

CINZILLA, *Herpes zoster*.

CION, *Uvula*.

CIONIS. The *Uvula*. Also, tumefaction, or elongation of the *uvula*; *Staphylodial'ysis*.

CIONITIS, from *κίονις*, 'the uvula,' and *itis*, 'inflammation.' Inflammation of the *uvula*, *Uvulitis*.

CIONORRHAPHIA, *Staphyloraphy*.

CIONOTOME, from *κίον*, 'the uvula,' and *τομή*, 'incision.' Curved scissors for cutting off the *uvula*.

CIONOT'OMY, *Cionotomia*, from *κίον*, 'the uvula,' and *τομή*, 'incision.' Excision of the *uvula* when too long.

CIPIPA, see *Jatropha manibot*.

CIRCÆA, *Atropa mandragora*, *Circæa Luteiana*.

CIRCÆA LUTETI'ANA, *Circæa*, *Paris'ian*. *Circæa*, from *Circe*, the Enchantress; *Enchanters' Nightshade*, (F.) *Herbe de Saint Étienne* ou *aux Sorciers*. This plant, common in the vicinity of Paris, was formerly considered to be resolvent and vulnerary. It was also supposed to possess wonderful magical and enchanting properties.

CIRCINUS, *Herpes zoster*.

CIRCLE, *Circulus* — c. Ciliary, Ciliary ligament — c. of Willis, see *Circulus*.

CIRCOCELE, *Cirsocele*.

CIRC'ONCISION, *Circumcision*.

CIRC'ONFLEXE, *Circumflexus*.

CIRC'ONSCRIT, *Circumscribed*.

CIRC'ONVOLUTION, *Convolution*.

CIRCUIT, *Circuitus*; in pathological language, generally means 'period,' 'course.'

CIRCUITUS, *Period*, *Circuit*.

CIR'CLAR, *Circularis*, from *circulus*, 'a circle,' (F.) *Circulaire*. Having the form of a circular; as *Circular Amputation*, &c.

The French use the expression "*Une circulaire*," for a turn of a bandage around any part.

CIRCULAR SINUS of Ridley, *Sinus coronarius*.

CIRCULA'TION, *Circula'tio*, *Cyclophor'ia*, *Cyclo'sis*, *Periodus sanguinis*, from *circulus*, 'a circle;' or rather, from *circum*, 'around,' and *ferre, latum*, 'to carry.' (F.) *Circulation*. Physiologists give this name to the motion of the blood through the different vessels of the body — *sanguimotion*; — to that function, by which the blood, setting out from the left ventricle of the heart, is distributed to every part of the body by the arteries; — proceeds into the veins, returns to the heart, enters the right auricle, and passes into the corresponding ventricle, which sends it into the pulmonary artery to be distributed to the lungs, whence it issues by the pulmonary veins, and passes into the left auricle. From this it is sent into the left ventricle, and is again distributed by means of the arteries.

CIRCULA'TION, *CAP'ILLARY*, *C. des Parenchymæ*, is that which takes place in the capillary vessels; and is, in some measure, independent of the action of the heart. See *Capillary Vessels*.

CIRCULATION, *PULMON'IC* or *LESSER*, is the circle from the right to the left side of the heart by the lungs. — The *GREATER* or *SYSTEMAT'IC* or *SYSTEM'IC*, is that through the rest of the system.

CIRCULATOIRE, *Circulatory*.

CIRCULATOR, *Charlatan*.

CIR'CLATORY, *Circulato'rius*, (F.) *Circula-*

toire; same etymon as *circulation*. Relating to the circulation of the blood; — *sanguimot'ory*.

CIR'CVLUUS. A circle or ring; *Cyclus*, *Cy'rios*, (F.) *Cercle*. Any part of the body which is round or annular, as *Circulus Oculi* — the *globe, bulb, or orb of the eye*. — Hippocr., Galen. It is, also, applied to objects, which by no means form a circle, — as to the *Circle of Willis*, *Circulus arterio'sus Willis'ii*, (F.) *Hexagone artérielle*, which is an *anastomotic circle* at the base of the brain, formed by the anterior and the posterior cerebral arteries and the communicating arteries of Willis.

CIRCULUS ARTERIO'SUS I'RIDIS. The artery which runs round the iris, and forms a circle.

CIRCULUS ARTERIOSUS WILLISII, *Circle of Willis*, see *Circulus* — c. *Ciliaris*, *Ciliary ligament* — c. *Membranous*, *Hymen*.

CIR'CVLUS CALLO'SUS HALLERI. *Tendo cordis veno'sus*. The fibro-cartilaginous rings, around the auriculo-ventricular openings of the heart, to which the tricuspid and mitral valves are attached.

CIRCULUS QUAD'RUPLEX; a kind of bandage used by the ancients.

CIRCULUS TONSILLA'RIS. A plexus formed by the tonsillitic branches of the glosso-pharyngeal nerve around the base of the tonsil.

CIRCULUS VENO'SUS, *Figura veno'sa*, *Vena seu Sinus termina'tia*. The venous circle in the embryo, which bounds the *Area Vasculosa* or *Vascular Area*.

CIR'CVLUS VENO'SUS ARE'OLÆ. The venous circle, formed by the union of the veins around the nipple. It embraces, however, only two-thirds of the circuit.

CIRCUMAGENTES, *Oblique muscles* of the eye.

CIRCUMCAULALIS MEMBRANA, *Conjunctiva*.

CIRCUMCISIO, *Circumcision* — c. *Fœminarum*, see *Circumcision*.

CIRCUMCIS'ION, *Circumcis'io*, *Posthetomy*, *Præcis'io seu Abscis'io Præputii*, *Circumcis'ionis*, *Circumsec'tio*, *Perit'omè*, from *circum*, 'around,' and *cædere*, 'to cut.' (F.) *Circumcision*. An ancient operation, performed by some nations as a religious ceremony. It consists in removing circularly a portion of the prepuce of infants; — a custom, which was probably suggested with a view to cleanliness. In cases of extraordinary length of prepuce, or when affected with disease, the operation is sometimes undertaken by surgeons. A similar operation is performed, amongst the Egyptians, Arabians, and Persians, on the female, *Circumcis'io femina'rum*, by removing a portion of the nymphæ, and at times the clitoris.

CIRCUMCISURA, *Circumcision*.

CIRCUMDUCTIO, *Perisphalix*.

CIRCUMDUCTIONIS OPIFEX, *Obliquus superior oculi*.

CIRCUMFLEX, *Circumflex'us*, from *circum*, 'around,' and *flexus*, 'bent.' (F.) *Circumflexe*. Curved circularly. A name given to several organs.

CIRCUMPLEX or *ARTIC'ULAR ARTERIES* of the arm are distinguished into *anterior* and *posterior*. They arise from the axillary, and are distributed around the shoulder.

CIRCUMPLEX ARTERIES of the THIGH are distinguished into *external* and *internal*. — *A. Sastrochanterienne* — Ch. They are given off from the *Profunda*, and surround the head of the thigh bone.

CIRCUMPLEX MUSCLE, *Circumflexus Musculus*, *C. Palati Mollis*, *Tensor Palati*, *Peristaphyl'ius externus seu inferior*, *Spheno-salpingo-staphyl'ius seu Staphyl'ius externus*, *Musculus tubæ novæ*, *Palato-salpingeus*, *Pter'gyo-staphyl'ius*,

Petro-salpin-go-staphyli-nus, *Spheno-pter-ygo-palati-nus*, *Sulpingo-staphyli-nus*, (F.) *Palato-salpingien*. A muscle, which arises from the spinous process of the sphenoid bone, and is inserted into the *velum pendulum palati*. Its use is to stretch the velum.

CIRCUMFLEX NERVE, Axillary Nerve.

CIRCUMFLEX VEINS follow the arteries.

CIRCUMFORANEUS, Charlatan.

CIRCUMFU'SA. Hallé has thus designated the first class of subjects that belong to Hygiene—as atmosphere, climate, residence, &c.; in short, every thing which acts constantly on man externally and internally.

CIRCUMGYRATIO, Vertigo.

CIRCUMLIGATURA, Paraphimosis.

CIRCUMLIT'IO, from *circumlitino*, 'I anoint all over.' *Perich'risis*, *Perich'riston*. A term formerly used for liniments, but especially for those applied to the eyelids.

CIRCUMOSSALE, Periosteum.

CIRCUMSCRIBED, *Circumscriptus*, (F.) *Circomscrit*. A term applied, in pathology, to tumours, which are distinct at their base from the surrounding parts.

CIRCUMSECTIO, Circumcision.

CIRCUMVALLATÆ PAPILLÆ, see *Papillæ of the Tongue*.

CIRE, Cera.

CIRE JAUNE ET BLANCHE, *Cera flava et alba*—*c. des Oreilles*, Cerumen.

CIRIOS, Circulus.

CIRON, *Acarus*, Psora.

CIRRHAGRA, Plica—*c. Polonorum*, Plica.

CIRRHON'OSUS; from *kirpos*, 'yellow,' and *voros*, 'disease.' A disease of the fetus, in which there is a yellow coloration of the serous membranes.—*Siebenhaar*.

CIRRHOSE, *Cirrhosis*—*c. du Foie*, *Cirrhosis*.

CIRRHOS'IS, *Kirrho'sis*, *Cirrhono'sis*, *Kirrhono'sis*, (F.) *Cirrhose*, from *kirpos*, 'yellow.' A yellow colouring matter, sometimes secreted in the tissues, owing to a morbid process. Also, called *Kirrho'sis* or *Kirrho'sis*.

CIRRHOS'IS HEPATIS, see *Hepatitrophia*. *Gran'ulated*, *gran'ular*, *mam'millated*, *tuber'culated*, and *hob-nailed liver*, (F.) *Cirrhose du Foie*. It appears to be dependent upon repletion of the terminal extremities of the biliary ducts with bile, along with atrophy of the intervening parenchyma. Hence the liver is smaller in size, or atrophied.

CIRRHOSIS HEPATIS, see *Cirrhosis*.

CIRRHOSIS OF THE LUNG, *Cirrho'sis pulmo'num*. Dr. Corrigan has described a condition of the lung under this name, the general character of which he considers to be a tendency to consolidation or contraction of the pulmonary tissue, with dilatation of the bronchial tubes.

CIRRHOSIS PULMONUM, C. of the Lung.

CIRRHOTIC, *Cirrho'ticus*. Same etymon as *cirrhosis*. Affected with, or having the character of *cirrhosis*.

CIRSIUM ARVEN'SE, *Car'duus hemorrho'ida'lis*, *Ceano'thos*, (F.) *Chardon hémorrhoidal*. A common plant, used in France, in the form of emulsion in hemorrhoids; and worn as an amulet.

CIRSUM MACULATUM, *Carduus marianus*.

CIRSOCE'LE, *Cirsoce'le*, *Cirso'cheum*, from *kirpos*, 'varix,' and *κηλη*, 'hernia;' *Varicose Hernia*. The greater part of authors have employed the term synonymously with *Varicocele*. Pott gives it a different signification. *Varicocele*, he calls the tumour formed by the veins of the scrotum; *Cirsocele*, *Funiculus varico'sus*, the varicose dilatation of the spermatic veins. The scrotum feels as if it contained earthworms. It is

commonly an affection of no consequence, demanding merely the use of a suspensory bandage.

CIRSOI'DES, *Cirso'ides*, from *kirpos*, 'varix,' and *ειδος*, 'resemblance.' Varicose, or resembling a varix. Rufus of Ephesus, according to James, applies this term to the upper part of the brain, as well as to the spermatic vessels.

CIRSOM'PHALUS, from *kirpos*, 'varix,' and *ομφαλος*, 'navel.' Varicose dilatation of the veins surrounding the navel. The term has, likewise, been applied to the aneurismal dilatation of the arteries of that region; called also, *Varicomp'h-alus*, (F.) *Hernie anévrymale*, *Aneurismal Hernia*.

CIRSOPHTHAL'MIA, *Cirsophthal'mus*, *Telangiecta'sia oculi*, from *kirpos*, 'varix,' and *οφθαλμος*, 'the eye;' *Varicose ophthalmia*, *Ophthalmia varico'sa*, *Varicos'itis conjunctivæ*. A high degree of ophthalmia, in which the vessels of the conjunctiva are considerably injected.

CIRSOCEUM, *Cirsocele*.

CIRSOT'OMY, *Cirsotom'ia*, from *kirpos*, 'varix,' and *τομή*, 'an incision.' Any operation for the removal of varices by incision.

CIRSUS, Varix.

CIRSYDROSCHEOCE'LE, from *kirpos*, 'varix,' *ὕδωρ*, 'water,' *σχέον*, the 'scrotum.' *Varicocele* with water in the scrotum.

CISEAU (F.), *Forfex*. An instrument, composed of a flattened rod of metal, sharpened at one extremity, and used in anatomical preparations, and in certain surgical operations, for dividing bones.

CISEAUX, Scissors.

CISSA, Malacia.

CISSAM'PELOS CAPEN'SIS, *Nat. Ord. Menispermaceæ*, grows in almost every mountainous part of the Cape of Good Hope. The root is used as an emetic and cathartic by the Boers.

CISSAMPELOS GLABERRIMA, see *Pareira brava*—*c. Pareira*, *Pareira brava*.

CISSARUS, *Cistus Creticus*.

CISSINUM, from *κισσος*, 'ivy.' Name of a plaster of ivy, used in wounds of the nerves or tendons.—*Paulus of Ægina*.

CISSOIDES, *Capreolaris*.

CISSOS, *Hedera helix*.

CISTERN, LUMBAR, *Receptaculum chyli*.

CISTER'NA, from *κίστη*, (L.) *Cista*, 'a chest.' (F.) *Citerne*. This term has been applied to various parts of the body, which serve as reservoirs for different fluids. The fourth ventricle of the brain has been so called.—*Arantius*.

CISTERNA CHYLI, *Receptaculum chyli*.

CISTHORUS, *Cistus Creticus*.

CISTOCLE, *Cystocoele*.

CISTUS CANADENSIS, *Helianthemum Canadense*.

CISTUS CRE'TICUS, *C. salvifolius* seu *taur'icus*, *Cisthorus*, *Cis'arvus*, *Dorycin'um*, *Gum Cistus*, *Ord. Cystineæ*; *Sec. Syst. Polyandria Monogynia*. The systematic name of the plant whence the *Lad'danum*, *Lad'damen* or *Lad'danum*, *Gum'mi Lad'danum*, is obtained. *Lad'danum* is a gummy-resinous substance, of a very agreeable smell, found in the shops in great masses. Its colour is blackish-green; taste, warm and bitter. It is but little used now. Formerly, it was a component of warm plasters, and was prescribed internally as a stomachic. *Ladanum* is also obtained from *Cistus ladanif'erus*, and *C. laurifolius*.

CISTUS, Gum, *Cistus Creticus*—*c. Salvifolius*, *C. Creticus*—*c. Tauricus*, *C. Creticus*.

CITERNE LOMBAIRE, *Receptaculum chyli*.

CITHARUS, Thorax.

CITRAGO, *Melissa*.

CITRARIA, *Melissa*.

CITRAS CHINICUS, *Quinix citras*.

CITREA MALUS, see *Citrus medica*.

CITREOLUS, *Cucumis sativus*.

CITRIC ACID, *Ac'idum cit'ricum*, *Ac'idum Limo'nis*, *Acid of Lemons*, *Ac'idum Limo'num*, (F.) *Acide citrique*. This acid is found in the lemon, orange, &c. It is in rhomboidal prisms, which slightly effloresce on exposure to the air. It dissolves in a twelfth part of its weight in boiling water, and has an extremely acid but agreeable taste. It is employed in medicine as antiseptic, refrigerant and diuretic. Rubbed up with sugar and with a little of the essence of lemon, it forms the dry *Lemonade*, (F.) *Limonade sèche*.

CITRINE OINTMENT, *Unguentum hydrargyri nitratis*.

CITRON, see *Citrus medica*—c. Tree, see *Citrus medica*.

CITRONELLE, *Artemisia abrotanum*, *Melissa*.

CITRONNADE, *Lemonade*.

CITROUILLE, *Cucurbita*.

CITRUL, *SICILIAN*, *Cucurbita citrullus*.

CITRULLUS, *Cucurbita citrullus*.

CITRULLUS AMARUS. An African plant, *Ord. Cucurbitaceæ*, called by the Boers *Bitterappel* or *Wild Watermelon*, the pulp of which, like that of *colocynthis*, is a drastic cathartic.

CITRULLUS COLOCYNTHIS, *Cucumis colocynthis*.

CITRUS, see *Citrus medica*—c. *Acida*, see *Lime*.

CITRUS AURANTIIUM. The systematic name of the *Orange Tree*, *Aurantium*, *A. Hispani'se*, *Malus Aurantia*, *Malus Aurantia major seu vulgaris*, *Aurantium vulgare*, *Citrus vulgaris*. *Ord. Aurantiaceæ*. *Sex. Syst. Polyadelphia Icosandria*. The fruit are called *Mala Aurantia*, *Chrysomelia*, *Nerantia*, *Martiana Poma*, *Poma Aurantia* seu *Chinesia*, *Aurantia Curassavica*, *Oranges*. The *Flowers of the Orange*, *Flores Naphæ*, are highly odoriferous, and used as a perfume. On distillation, they yield a small quantity of essential oil—*Oleum Aurantii*, *Oleum seu Essentia Nero'li*,—with spirit and water, the *Aqua Florum Aurantii*, *Aurantii floris aqua*, *Aqua aurantii*, *Orange-flower water*. They were once used in convulsive and epileptic cases. The leaves, *Folia Aurantii*, have a bitterish taste, and furnish an essential oil. They have been used for the same purposes as the flowers. The yellow rind of the fruit, *Aurantii Cortex* (Ph. U. S.), *Citrus Bigaradia* (Ph. L.), *Bigarade* or *Bitter Orange*, *Orange Peel*, is an aromatic bitter, and is used in dyspepsia, and where that class of remedies is required. The *Juice*, *Succus Aurantii*, *Orange juice*, is a grateful acid, and used as a beverage in febrile and scorbutic affections.

CITRUS BERGAMIA, *Citrus mella rosa*—c. *Bigaradia*, see *Citrus aurantium*—c. *Decumana*, *Shaddock*—c. *Limetta*, see *Citrus mella rosa*—c. *Limonum*, *C. medica*.

CITRUS MEDICA, *C. Limo'num*. The systematic name of the *Lemon Tree*. The *Lemon*, *Limon* (Ph. U. S.), *Limo'num Malum* seu *Bacca*, *Malus Medica*, *M. Limo'nia* *Ac'ida*, *Limon*, *Cit'rea*, *Malus*, *Citrus*, (F.) *Citron*, *Cédrat*, has a fragrant odour, depending upon the essential oil, *Oleum Limo'nis*, of the rind. The outer rind, *Cortex Limo'num*, *Limo'nis Cortex* (Ph. U. S.), *Lemon Peel*, *Zest*, *Flave'do Corticum Citri*, is used in the same cases as the *Cortex Aurantii*.

The *juice*, *Succus Limo'nis*, *Limo'num Succus* (Ph. L.), (F.) *Suc du Limon*, *Suc de Citron*, is sharp, but gratefully acid, the acidity depending upon the citric acid it contains, and is given as a refrigerant beverage in febrile affections. In doses of half an ounce to an ounce, three times a day, it has appeared to exert a markedly sedative

influence on the circulation, and has been given, apparently with benefit, in acute rheumatism and rheumatic gout. Alone, or combined with wine, it is prescribed in scurvy, putrid sore throat, &c. Its general properties are refrigerant and antiseptic. Sweetened and diluted, it forms *Lemonade*. *Artificial lemon-juice* is made by dissolving an ounce of citric acid in fourteen fluidounces of water; adding a few drops of essence of lemon.

Lemonpulp tea, or *water*, is made by paring the rind of one lemon, previously rubbed with half an ounce of sugar: the peelings and sugar are then put into a jar, and a quart of boiling water is poured over them. When cold, the fluid must be poured off, and a tablespoonful of lemon juice be added.

It is an agreeable drink in fevers.

Citron Tree is likewise considered to belong to the same species—*Citrus Medica*. Its fruit is called *cedrome'la*. It is larger and less succulent than the lemon. *Citron juice*, when sweetened with sugar, is called by the Italians *Agri di Cedro*.

CITRUS MELLA ROSA of De Lamarck, another variety of *Citrus Medica*, affords the *Bergamote*, as also do *Citrus Limetta* and *C. Bergamia*.

CITRUS VULGARIS, *Citrus aurantium*.

CITTA, *Malacia*.

CITTARA, **MINERAL WATERS OF**. These springs are in the Isle of Ischia, near the sea. They contain carbonate and sulphate of lime, and chloride of sodium. Their temperature is 100° Fahrenheit.

CITTOS, *Hedera helix*.

CITTOSIS, *Chlorosis*, *Malacia*.

CIVETTA, *Zib'ethum*, *Civet*, (F.) *Civette*. An unctuous perfume, of a very penetrating odour, obtained from different mammalia of the *Viverra* kind, particularly from *Viverra civetta*. It is contained in a fold of the skin, situate between the anus and the organs of generation.

CLABBER, *Bonnyclabber*.

CLABBERGRASS, *Galium verum*.

CLADES GLANDULARIA, *Plague*.

CLADISCOS, *Ramusculus*.

CLADONIA ISLANDICA, *Lichen islandicus*.

CLADONIA RANGIFERINA. The ancients regarded this European plant as pectoral and stomachic. It enters into the composition of the *Poudre de Chypre*.

CLADOS, *Ramus*.

CLADRAS'TIS TINCTO'RIA, *Virgil'ia Yellow Ash*, *Fustic Tree*, *Yellow Locust*. An indigenous tree, which flourishes from Kentucky to Alabama. The bark of the tree and the roots are cathartic.

CLAIRET, *Claret*.

CLAIRVOYANCE (F.), 'Clear-seeing.' A clearness of sight, said to be communicated by animal magnetism, which not only enables the magnetized person to see in the dark, through stone walls, &c., but even to observe prospects, which he has never seen previously, whilst he may fancy he is flying in the air. It need hardly be said, that the possession of such powers is fabulous.

CLAMMY, *Viscidus*, *Glutinosus*, (F.) *Pâteux*, (D.) *Klam*, 'moist.' Viscous; glutinous. A clammy mouth, (F.) *Bouche pâteuse*, is one that is covered with a mucous coat.

CLAMMY WEED, *Polanisea graveolens*.

CLAMOR, *Cry*.

CLANGOR, *Oxyphonia*.

CLAP, *Gonorrhœa impura*.

CLAPIER (F.). A clapper, *Latibulum*, from *clapper*, 'to conceal.' A purulent foyer of disease; concealed in the flesh or under the skin. See Sinus.

CLAPWORT, *Orobancha Americana*.

CLAQUEMENT, *Odontosynerismus*.

CLARENDON, MINERAL WATERS OF. Mild acidulous springs in Vermont; temperature from 48° to 54° Fahr. The gas, given off from the water, consists of carbonic acid, oxygen, and nitrogen. The springs have a reputation in cutaneous diseases, chronic bronchitis, &c.

CLAR'ET, (*Vin clair* [?]), *Clare'tum*, *Vin de Bordeaux*, (F.) *Clairét*. A pleasant French wine, which may be used whenever wine is required. Also, a wine impregnated with spice and sugar, called likewise *Vinum Hippocrat'icum* seu *Medicatum*, *Potus Hippocrat'icus*, *Hippocras*, *Hyp'ocras*. Schröder speaks of a *Clare'tum alt'erans*, and a *C. purgans*.

CLARETA, *Albumen ovi*.

CLARETUM, *Claret*.

CLARIFICA'TION, *Clarifica'tio*, *Depura'tion*, from *clarus*, 'clear,' and *facio*, 'I make.' A pharmaceutical operation, which consists in separating from a liquid every insoluble substance, held in suspension by it, that affects its transparency. *Decanting* and *filtering* are the operations necessary for this purpose.

CLARY, COMMON, *Salvia sclarea*.

CLASIS, *Fracture*.

CLASMA, *Fracture*.

CLASS, *Clas'is*, (F.) *Classe*. An assemblage of a certain number of objects. In *Natural History* and in *Medicine*, a group of objects or individuals having one or more common characters. The *classes* are divided into *orders*, the *orders* into *genera*, the *genera* into *species*, and these last into *varieties*.

CLASSIFICA'TION, *Classifica'tio*, from *classis*, 'a class,' and *facio*, 'I make.' The formation of classes. A methodical distribution of any objects whatever into classes, orders, genera, species, and varieties. See Nosography, and Nosology.

CLASSY, MINERAL WATERS OF. Classy is near Laon in Picardy, France. The waters are chalybeate.

CLAUDICATIO, *Claudication*—c. *Anatica*, *Vællatio*.

CLAUDICA'TION, *Claudica'tio*, from *claudicare*, 'to be lame.' The act of halting or limping. *Lameness*, (Prov.) *Hirpling*, *Clau'ditas*, *Cholo'sis*, *Cholei'a*, *Cholo'ma*, (F.) *Claudication*, *Boitement*, *Boiterie*. This condition does not constitute any special disease, but is produced by different causes or affections. It may be the result of the shortening or elongation of one of the lower limbs, of ankylosis of the joints, palsy of the muscles, pain, &c.

CLAUDITAS, *Claudication*.

CLAUSTRUM GUTTURIS, Isthmus of the fauces—c. *Palati*, *Velum pendulum palati*—c. *Virginittis*, *Hymen*.

CLAUSU'RA, from *claudere*, 'to shut.' An imperforation of any canal or cavity.

CLAUSU'RA UTERI. Preternatural imperforation of the uterus.

CLAVA RUGOSA, *Acorus calamus*.

CLAVALIER À FEUILLES DE FRÊNE, *Xanthoxylum clava Herculis*.

CLAVARIA CORALLOIDES, *Coralloides Fungus*, *Coralwort*. *Ord. Fungi*, (*clava*, 'a knotty branch, a club.') Said to be corroborant and astringent. A kind of *clavaria*, called (F.)

Digital blanc, *Digital humain*, *Champignon de l'appareil des fractures*, formed of digitations, grouped together, and two or three inches in length, is said to have been often found, formerly at the *Hôtel Dieu* of Paris, on the splints of white wood used in the treatment of fractures, in autumn.—H. Cloquet.

CLAVATIO, *Gomphosis*.

CLAVEAU, *Murr*.

CLAVELEE, *Murr*.

CLAVES CALVARIÆ, *Wormiana Ossa*.

CLAVICLE, *Clavis*, *Clavic'ula*, *Clavic'ulus*, *Lig'ula*, *Fur'cula*, *Os Jug'uli*, *Jug'ulum*, *Cleis*, *Clei'dion*, from *clavis*, 'a key,' (F.) *Clavicule*. The collar-bone, (Old Eng.) *Cannelbone*, *Channell-bone*, (Sc.) *Cannelbayne*, *Cragbane*, *Halsbane*. The clavicle is shaped like the letter *S*, and is placed transversely at the upper part of the thorax. It is articulated, at one extremity, with the sternum; at the other with the acromion process of the scapula. It gives attachment, above, to the *Sterno-cleido mastoideus*; below, to the *Subclavius*; before, to the *Pectoralis major* and *Deltoides*; and behind, to the *Trapezius*. It serves as a point of support for the muscles of the arm, and protects the vessels and nerves passing to that extremity.

The fibres, connecting the lamellæ or plates of bones, have also been called *Claviculi* or *Nails*.

CLAVICULAR, *Clavicula'ris*, (F.) *Claviculaire*; same etymon. Relating to the clavicle or collar-bone.

CLAVICULAR NERVES, *Nervi clavicula'res*. Branches of the fourth cervical nerve, which are distributed to the clavicular region.

CLAVICULE, *Clavicle*.

CLAVICULI, see *Clavicle*.

CLAVIS, *Clavicle*, *Key*—c. *Anglica*, *Key*.

CLAVUS. A nail. *Helos*, *Gomphos*, (F.) *Clou*. This word is employed in medicine in various senses. It means, 1. A *Corn*, from its resemblance to the head of a nail. 2. Certain condylomatous excrescences of the uterus. 3. A callous tumour, which forms on the white of the eye, and resembles a nail, the *Clavus Oculi*, (F.) *Clou de l'œil*. This last, by some, is considered to be synonymous with *staphyloma*; by others, with *staphyloma of the cornea*. Also, the penis.

CLAVUS HYSTERICUS, *Monopa'gia*, *Monope'gia*, (F.) *Clou Hystérique*. An acute pain, confined to a small point of the head, described by the sick as resembling that which would be produced by a nail driven into the head. It has been particularly noticed in hysterical females;—hence its name. It is called *Ovum hystericum*, when the pain occupies a greater extent.

CLAVUS SECALINUS, *Ergot*—c. *Secalis*, *Ergot*—c. *Siliginis*, *Ergot*.

CLEANSINGS, *Lochia*.

CLEAR-SEEING, *Clairvoyance*.

CLEARWEED, *Pilea pumila*.

CLEAVAGE, from Anglo-Saxon *cleofan*, 'to split.' The natural line of separation exhibited by certain substances, as minerals, when subjected to mechanical force. The term has been applied to the separation of muscles into longitudinal and circular striæ, when mechanical violence is used.

CLEAVERS, *Galium aparine*.

CLEAVERS' BEES, *Galium aparine*.

CLEAVEWORT, *Galium verum*.

CLEF, *Key*—c. *du Crane*, *Wormiana ossa*—c. *de Gargenot*, *Key*—c. *à Noix*, see *Key*—c. *à Pivot*, see *Key*—c. *à Pompe*, see *Key*.

CLEFT, Rima, see Monster — c. Palato, see Harelip.

CLEIDAGRA, Cleisagra.

CLEIDION, Clavicle.

CLEIDO-COSTAL, Costo-clavicular.

CLEIDO-MASTOÏDEUS. Albinus thus designates the posterior portion of the sterno-cleido-mastoideus, which he considers a separate muscle. It has been corrupted into *clino-mastoideus*.

CLEIS, Clavicle, Key.

CLEIS'AGRA, *Cleid'agra*, from *κλεις*, 'the clavicle,' and *αγρα*, 'a seizure.' Gout in the clavicle. — A. Paré.

CLEITHORION, Clitoris.

CLEITHORIS, Clitoris.

CLEITORION, Clitoris.

CLEITORIS, Clitoris.

CLE'MATIS, from *κλημα*, *κλήματος*, 'a branch of a vine, a tendril.' A genus of plants in the Linnean system.

CLEMATIS DAPHNOIDES MAJOR, Vinca minor — c. *Corymbosa*, C. *erecta*.

CLEMATIS ERECT'A, *C. erecta seu corymbo'sa*, *Clematis erecta*, *Flammula Jovis*, *Upright Virgin's Bower*, (F.) *Olématis droite*. Order, Ranunculaceæ. *Sex. Syst.* Polyandria Polygynia. The leaves contain an acrid principle. They have been esteemed anti-venereal; and, in the form of powder, have been used as an escharotic.

CLEMATIS FLAMMULA, see C. *vitalba* — c. *Recta*, C. *erecta* — c. *Sepium*, C. *Vitalba* — c. *Viorna*, see C. *Vitalba* — c. *Virginica*, see C. *Vitalba*.

CLEMATIS VITAL'BA, *C. sepium seu sylvestris*, *Vital'ba*, *Vior'na*, *Atra'gené*, *Traveller's Joy*, *Common Virgin's Bower*, (F.) *Olématis*, *Herbe aux gueux*, *Aubevine*. It has been used in the same cases as the former. In infusion it has been applied in cases of itch.

The leaves of CLEMATIS CRISPA — C. FLAMMULA, sweet-scented Virgin's bower — C. VIRGINICA, common Virgin's bower — and C. VIOR'NA, Leather-flower, have similar properties.

CLÉMATITE, Clematis vitalba — c. *Droite*, Clematis *recta*.

CLEMATITIS ERECTA, Clematis *erecta*.

CLEOME PENTAPHYLLA, Gynandropsis pentaphylla.

CLEONIS COLLYRIUM. A collyrium described by Celsus, composed of equal parts of Samian earth, myrrh, and thus mixed with white of egg: called doubtless after its inventor.

CLEONIS GLUTEN. An astringent formula of myrrh, frankincense, and white of egg.

CLEP'SYDRA, from *κλεπτω*, 'I conceal,' and *υδωρ*, 'water.' An instrument contrived by Parnacelus to convey fumigation to the uterus.

CLEPTOMANIA, Kleptomania.

CLETHRA ALNIFO'LIA, *Sweet pepper-bush*, *White alder*; indigenous. Order, Ericaceæ. The leaves and flowers are diaphoretic and excitant.

CLETORIS, Clitoris.

CLEVES, MINERAL WATERS OF. This spring is a quarter of a league from Cleves, in Westphalia. It contains carbonate and sulphate of iron.

CLICKING, see *Râle sibilant*.

CLIDER, Galium aparine.

CLIFFORT'IA ILICIFO'LIA. Ord. Rosaceæ. A common South African plant, used by the Boers as an emollient expectorant in catarrh.

CLIFTON, CLIMATE OF. The vicinity of Clifton and of Bristol, England, appears to be

the mildest and driest climate in the west of England; and, consequently, the best winter residence, in that part of the country, for invalids. It is, also, a favorable summer climate, and is surrounded by numerous places of agreeable resort, suited for those who may pass the season there.

For the mineral waters of Clifton, see Bristol Hot Well.

CLIGNEMENT, Scardamygmus.

CLIGNOTEMENT, Nictation.

CLIMA, Climate.

CLIMAC'TERIC, *Climacter'icus*, *Climacter'icus*, from *κλιμακτηρ*, 'a step.' (F.) *Climacter'ique* ou *Climatérique*. A word, which properly signifies 'by degrees.' It has been applied to certain times of life, regarded to be critical.

At present, the word *Climacter'ic* is chiefly applied to certain periods of life, at which great changes occur, independently of any numerical estimate of years. Such are the period of puberty in both sexes; that of the cessation of the menses in women, &c.

CLIMACTERIO YEARS, *Anni Climacter'ici*, are, according to some, all those in the life of man, which are multiples of the number 7, *Septenn'ia*s. Others have applied the term to years, resulting from the multiplication of 7 by an odd number. Some have admitted only three *climacterics*; others, again, have extended them to multiples of 9. Most, however, have considered the 63d year as the *Grand Climacteric*; — 63 being the product of the multiplication of 7 by 9, and all have thought that the period of three, seven, or nine, which they respectively adopted, was necessary to the entire renewal of the body; so that there was, at these times, in the economy, none of the parts of which it had previously consisted. The climacteric years have also been called, (*Anni*) *hebdomad'ici*, *scala'res*, *grad'arii*, *scan'siles*, *genethliaci*, *natali'i*, *fata'les*, *crit'ici*, *decreto'rii*, *hero'ici*, &c. All the notions on the subject are essentially allied to the doctrine of numbers of Pythagoras.

CLIMATAL, Climatic.

CLIMATE, *Clima*, *Inclina'tio cæli*, (F.) *Climat*, Gr. *κlima*, 'a region.' In geography, the word *climate* is applied to a space on the terrestrial globe, comprised between two circles parallel to the equator, and arbitrarily measured according to the length of the days. In a hygienic point of view, we understand by *climate*, since Hippocrates, a country or region, which may differ from another in respect to season, qualities of the soil, heat of atmosphere, &c. Climate, indeed, embraces, in a general manner, all the physical circumstances belonging to each region, — circumstances which exert considerable influence on living beings. The dark complexion of the inhabitants of the torrid zone is easily distinguishable from the paleness of those of the frigid, — so are the diseases. They are all modified, more or less, by climate or locality. Hot climates predispose to abdominal complications in febrile affections; cold climates to the ratic, &c.

One of the most important considerations with regard to climates is their comparative fitness for the residence of invalids, and especially of those who are liable to, or suffering under, catarrhal or consumptive affections. The great object, in such cases, is to select a climate which will admit of regular and daily exercise in the open air, so that the invalid may derive every advantage which this form of revulsion is capable of effecting. To an inhabitant of the northern and middle portions of the United States — and the same

applies to Great Britain, France, and the northern parts of the old world—a more southern climate alone affords these advantages in an eminent degree. During the summer months there are few, if any, diseases, which require a milder climate than that of the United States, or of the milder districts of Europe. The temperature of the winter months is, consequently, the most important object of attention. Equability of temperature is essential, inasmuch as all sudden changes interfere with the great desideratum—exercise in the open air. In the whole continent of North America the changes are very sudden and extensive. It is not uncommon for the range to be 40°, between two successive days. So far, therefore, as this applies, the American climate is not well adapted to the invalid. In the southern portions, however, of the Union, this objection is counterbalanced by many advantages.

The following tables exhibit the mean temperature of the year, and of the different seasons—with the mean temperature of the warmest and

coldest months at different places in America, Europe, Africa, &c., as deduced from the excellent paper of Von Humboldt on Isothermal Lines, the Meteorological Registers kept by the surgeons of the United States army, under the direction of Surgeon-generals Lovell and Lawson, 1822 to 1854, inclusive, the work of Sir James Clark on Climate, &c.

Certain of the tables show the mean monthly temperature, maximum, minimum and range, as well as the greatest daily, and mean daily range during the corresponding months—but of different years—at some of the prominent retreats for the valetudinarian in Great Britain, on the continent of Europe, and in the African islands. It is proper, however, to remark, that in no situations, except in those to which an asterisk is affixed, was the register thermometer used. In the others, the observations were made during the *day* only, and consequently the numbers given are far below the real range throughout the twenty-four hours. The places are ranged in the order of their mean temperature.

TABLE OF MAXIMUM, MINIMUM, AND RANGE OF TEMPERATURE.

PLACES.	DECEMBER.			JANUARY.			FEBRUARY.			MARCH.			APRIL.		
	Max.	Min.	Range	Max.	Min.	Range	Max.	Min.	Range	Max.	Min.	Range	Max.	Min.	Range
Sidmouth	54	25	29	47	21	26	52	27	25	56	26	30	60	31	29
Penzance	56	34	22	54	28	26	55	33	22	59	34	25	62	36	26
Pau.....	56	25	31	56	21	35	60	35	25	65	35	30	71	43	28
Montpellier.....	57	32	25	53	27	26	55	30	25	58	35	23	64	41	23
Nice.....	59	40	19	58	27	31	58	37	21	65	41	24	69	46	23
Rome.....	60	31	29	58	29	29	60	33	27	65	37	28	74	44	30
Naples.....	61	34	27	58	29	29	60	31	29	69	38	31	78	43	35
Madeira.....	68	52	16	69	50	19	68	51	17	69	51	18	72	55	17

TABLE OF MEAN TEMPERATURE.

PLACES.	DECEMBER.	JANUARY.	FEBRUARY.	MARCH.	APRIL.
Sidmouth.....	43.00	36.30	42.00	45.00	51.00
Penzance.....	46.50	43.00	44.50	46.50	48.50
Pau.....	41.53	38.89	44.96	46.80	55.79
Montpellier.....	46.00	42.00	45.00	47.00	53.00
Nice.....	48.60	45.85	49.00	51.45	57.00
Rome.....	49.62	47.65	49.45	52.05	56.40
Naples.....	50.50	46.50	48.50	52.00	57.00
Madeira.....	60.50	59.50	58.50	61.06	62.50

TABLE OF DAILY RANGE OF TEMPERATURE.

PLACES.	DECEMBER.		JANUARY.		FEBRUARY.		MARCH.		APRIL.	
	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.	Mean daily range.	Greatest daily range.
Sidmouth.....		13		13		12		12		13
Penzance.....	3		4		8		8		9	
Pau.....	7	13	7	16	16		17		8	18
Montpellier.....	9		8		14		14		14	
Nice.....	10	14	8	16	9	18	9	17	11	18
Rome.....	10	15	11	16	10	18	12	19	13	20
Naples.....	9	13	9	14	11	19	11	18	14	20
Madeira.....	11	14	11	17	13		10	14	11	13

1. AMERICA, &c.

PLACES.	Latitude.	Mean tem- perature of several years.	Mean temperature of different seasons.				Mean temperature of	
			Winter.	Spring.	Summer.	Autumn.	Warmest month.	Cool- est month.
Nain.....	57° 08'	26° 42'	0° 60	23° 60	48° 38	33° 44	51° 80	11° 20
Fort Brady, Mich....	46.39	41.37	14.09	37.89	61.83	43.94	62.87	12.63
Quebec, L. C.....	46.47	41.74	14.18	38.04	68.00	46.04	73.40	13.81
Eastport, Me.....	44.54	42.44	23.44	38.58	60.54	45.43	63.52	20.31
Fort Howard, Mich....	44.40	44.50	20.82	41.40	68.70	45.18	73.67	17.95
Fort Crawford, Miss..	43.03	45.52	23.76	43.09	69.78	46.74	71.34	20.14
Cambridge, Mass....	42.21	50.36	33.98	47.66	70.70	49.82	72.86	29.84
Council Bluffs, Miss..	41.25	50.82	27.38	46.38	72.84	48.60	75.92	27.19
Newport, R. I.....	41.30	51.02	33.82	46.87	68.70	53.83	71.46	32.14
Philadelphia.....	39.56	53.42	32.18	51.44	73.94	56.48	77.00	32.72
New York.....	40.40	53.78	29.84	51.26	79.16	54.50	80.78	25.34
Cincinnati.....	39.06	53.78	32.90	54.14	72.86	54.86	74.30	30.20
Monticello, Va.....	37.58	55.40	37.67	54.67	73.33	56.50	75.00	36.00
Washington, D. C....	38.53	55.56*	36.80	53.83	75.90	56.59	79.13	34.66
Smithville, N. C....	34.00	58.88	53.44	64.76	80.46	68.15	82.93	50.69
Charleston, S. C....	32.47	60.18	51.09	66.73	80.89	67.55	82.81	49.43
Natchez, Miss.....	31.34	64.76	48.56	65.48	79.16	66.02	79.70	46.94
Pensacola, Flor.....	30.28	68.77†	55.13	69.67	82.57	69.05	83.55	53.80
St. Augustine, do....	29.48	72.23	59.29	71.47	82.73	75.15	83.94	56.60
Tampa Bay, do....	27.57	72.37	61.24	72.93	80.14	75.28	80.72	58.70
Vera Cruz.....	19.11	77.72	71.96	77.90	81.50	78.62	81.86	71.06
Havana.....	23.10	78.08	71.24	78.98	83.30	78.98	83.84	69.98
Bahamas.....	26.40	78.3*	71.	77.	83.	80.	90.	64.
Barbadoes.....	13.10	79.3	76.7	19.	81.	80.		
Cumana.....	10.27	81.86	80.24	83.66	82.04	80.24	84.38	79.16
St. Louis, Missouri, Lat. 38° 46'. Mean temp. 55° 86. New Harmony, Lat. 38° 11'. Mean temp. 56° 74. New Orleans, Lat. 30°. Mean temp. 69° 01. Baton Rouge, Lat. 30° 26'. Mean temp. 68° 07. Jamaica, coast, Mean temp. 80° 6.								

RANGE OF MEAN TEMPERATURE IN THE U. STATES FOR THE EXTREME MONTHS, SEASONS, AND YEAR
(Army Meteorological Register, 1855.)

STATIONS.	YEARS.	Latitude.	January.	July.	Spring.	Summer.	Autumn.	Winter.	Year.
Hancock Barracks, Maine.....	17	46° 07'	16.0	8.7	6.4	6.8	7.2	7.1	5.0
Fort Sullivan, Maine.....	25	44.54	14.0	9.9	7.7	6.8	4.7	7.8	3.7
Fort Constitution, New Hampshire, 25	43.04	14.0	9.0	7.0	7.4	5.7	11.9	6.7	
Fort Independence, Massachusetts, 17	42.20	9.9	8.6	7.8	4.9	6.9	11.0	4.7	
Fort Columbus, New York.....	33	40.42	12.6	11.8	7.8	7.1	9.9	10.3	7.3
Watervliet, New York.....	31	42.43	15.3	10.5	8.8	6.7	9.2	12.8	7.4
Alleghany Arsenal, Pennsylvania... 22	40.32	18.9	14.5	12.9	7.8	12.0	11.2	7.8	
Fort McHenry, Baltimore, Md..... 24	39.17	14.2	6.8	8.9	6.0	7.2	11.2	4.9	
Fort Monroe, Virginia, (Norfolk)... 30	37.00	16.9	7.4	10.1	5.9	7.6	16.0	9.0	
Fort Moultrie, S. C., (Charleston)... 28	32.45	18.3	6.1	7.1	7.3	9.5	16.7	6.9	
Fort Marion, St. Augustine, Florida, 20	29.43	16.1	8.6	11.3	6.7	7.1	15.4	6.7	
Key West, (Florida),.....	14	24.32	12.6	5.9	8.5	3.2	4.2	8.2	3.5
Fort Brooke, Florida.....	25	28.00	17.7	6.8	6.8	5.4	4.8	11.9	4.0
Fort Barrancas, Pensacola, Florida, 17	30.18	18.2	5.3	7.1	3.2	4.5	9.7	3.0	
Mount Vernon Arsenal, Mobile, Ala., 14	31.12	14.2	6.1	4.7	4.3	5.6	6.8	3.5	
Fort Pike, Louisiana.....	14	30.10	10.7	6.2	6.2	3.0	7.0	13.3	4.8
New Orleans, Louisiana.....	20	29.57	18.0	6.8	7.4	5.2	5.5	8.9	4.7
Baton Rouge, Louisiana.....	24	30.26	15.4	7.1	9.3	4.8	6.9	11.1	4.2
Fort Jessup, Louisiana.....	23	31.33.	17.5	7.8	9.0	4.6	7.6	10.6	6.6
Fort Gibson, Indian Territory..... 27	34.47	19.5	7.8	9.2	8.6	11.3	14.4	6.9	
Jefferson Barracks, Missouri..... 26	38.28	17.6	12.0	15.0	9.1	10.2	19.2	7.0	
St. Louis, Missouri.....	12	38.40	14.0	7.2	7.2	4.7	7.4	7.3	3.7
Fort Gratiot, Michigan.....	17	42.55	13.6	11.8	11.8	8.2	8.9	10.7	6.1
Fort Mackinac, Michigan.....	24	45.51	12.3	10.3	8.1	5.8	7.9	9.9	6.1
Fort Brady, Michigan.....	31	46.30	13.0	13.3	13.0	8.5	9.8	9.0	7.1
Fort Howard, Wisconsin.....	21	44.30	14.5	11.8	9.2	7.6	9.9	16.6	5.1
Fort Crawford, Wisconsin.....	19	43.05	19.6	10.4	17.7	8.1	9.2	16.4	8.9
Fort Armstrong, Illinois.....	11	41.30	18.5	10.3	6.8	8.3	6.2	17.4	7.3
Fort Snelling, Minnesota.....	35	44.53	27.9	14.6	17.3	11.0	10.4	16.0	8.6
Fort Leavenworth, Kansas.....	24	39.21	24.7	10.0	17.8	8.0	9.4	13.3	8.0
Fort Kearney, Nebraska.....	6	40.38	19.9	5.2	5.8	4.1	7.0	13.0	5.3
Fort Laramie, Nebraska.....	6	42.12	13.2	4.5	6.4	2.6	10.8	4.8	5.8
Fort Brown, Texas.....	7	25.54	10.0	2.9	2.5	2.2	3.1	4.2	1.6
Fort McIntosh, Texas.....	6	27.31	10.2	3.7	4.8	5.5	3.1	3.6	1.9
Santa Fé, New Mexico.....	5	35.41	4.3	6.9	2.3	1.8	5.0	3.7	0.8
San Diego, California.....	5	32.42	3.7	6.9	40.7	2.7	3.6	2.6	2.3
Benicia, California.....	6	38.03	6.9	4.9	4.6	2.9	3.4	3.2	8.2
Fort Vancouver, Oregon.....	6	45.40	10.1	3.4	4.0	2.5	2.2	2.3	1.4
Fort Steilacoom, Washington Ter., 6	47.10	12.6	3.8	2.3	1.8	2.8	4.0	2.0	

2. EUROPE, AFRICA, &c.

PLACES.	Latitude.	Mean temperature of several years.	Mean temperature of different seasons.				Mean temperature of	
			Winter.	Spring.	Summer.	Autumn.	Warmest month.	Cooldest month.
Geneva.....	48° 12'	49° 28'	34° 70	47° 66	64° 94	50° 00	66° 56	34° 16
Gosport.....	48 .1	50 .24	40 .44	47 .63	62 .00	50 .88		
Newport, Isle of Wight	50 .40	51 .00	40 .31	49 .00	63 .09	51 .63		
Sidmouth.....		52 .10	40 .43	50 .66	63 .83	53 .50		
Penzance.....	52 .11	51 .80	44 .03	49 .63	60 .70	53 .36		
Undercliff.....		51 .11	42 .14	29 .26	60 .28	52 .76		
Hastings.....	50 .52	57 .00	40 .11	45 .77	60 .45	51 .00		
Bute.....	55 .42	48 .25	39 .62	46 .66	58 .02	48 .59		
Cove of Cork.....	51 .54	51 .58	43 .90	49 .43	61 .26	51 .73		
Jersey.....	49 .13	53 .06	43 .82	50 .97	62 .84	54 .63		
Paris.....	48 .50	51 .08	38 .66	49 .28	64 .58	51 .44	65 .30	36 .14
Pau.....	43 .7	54 .95	41 .79	54 .96	67 .41	55 .64		
Sienna.....	43 .24	55 .60	40 .50	54 .10	70 .80	57 .10		
Nautes.....	47 .13	55 .62	42 .23	53 .10	70 .73	56 .41	70 .52	39 .02
Bordeaux.....	44 .50	56 .48	42 .08	56 .46	70 .88	56 .30	73 .04	41 .00
Montpellier.....	43 .36	57 .60	44 .20	53 .33	71 .30	61 .30		
Avignon.....	43 .7	58 .20	42 .60	57 .13	74 .66	59 .00		
Florence.....	43 .46	59 .00	44 .30	56 .00	74 .00	60 .70		
Nice.....	43 .42	59 .48	47 .82	56 .23	72 .26	61 .63		
Marseilles.....	43 .17	59 .50	45 .50	57 .56	72 .50	60 .08		
Toulon.....	43 .07	59 .90	43 .30	53 .70	74 .30	59 .00		
Leghorn.....	43 .33	60 .00	46 .30	57 .60	74 .10	62 .00		
Genoa.....	44 .25	60 .37	44 .57	53 .60	74 .03	62 .94		
Pisa.....	43 .43	60 .60	46 .03	57 .20	75 .15	62 .80		
Rome.....	41 .53	60 .40	45 .86	57 .74	75 .20	62 .78	77 .00	42 .26
Naples.....	40 .54	61 .40	48 .50	53 .50	70 .83	64 .50		
St. Michaels, Azores..	37 .47	62 .40	57 .83	61 .17	68 .33	62 .33		
Cadiz.....	36 .32	62 .88	52 .90	59 .53	70 .43	65 .35		
Madeira, Funchal....	32 .37	64 .56	59 .50	62 .20	69 .33	67 .23		
Algiers.....	36 .48	69 .98	61 .52	65 .66	80 .24	72 .50	82 .76	60 .08
Canaries, Santa Cruz.	28 .28	70 .94	64 .65	68 .87	76 .68	74 .17		
Cairo.....	30 .02	72 .32	58 .46	73 .58	85 .10	71 .42	85 .82	56 .12
London, Lat. 51° .30'. Mean temp. 50° .36.			Enviions of London, Mean temp. 48° .81.					
Perpignan, Mean temp. 59° .54.			Lyons, Mean temp. 55° .76.					
Nismes, Mean temp. 60° .26.								

In the United States, the most favourable region for the phthisical invalid is that of Florida,—especially of Pensacola. St. Augustine is frequently chosen, but it is liable to north-east storms, which interfere with the out-door movements of the valetudinarian, and are the source of much discomfort. Still, great benefit has often been derived from it as a winter retreat. Of the Atlantic Isles, Madeira appears to be best adapted for the consumptive, and those affected with chronic bronchitis. In Italy, Rome, and Pisa,—and in England, Torquay and Undercliff, are to be preferred. Chronic rheumatism and gout are benefited by a warm climate, which, again, is unfavourable to those who are predisposed to cerebral diseases, especially to such as are characterized by debility and mobility of the nervous system—as paralysis, epilepsy, mania, &c. Hypochondriasis and dyspepsia require rather change of climate and travelling exercise than a sojourn in any one. (See the Author's Human Health, Philad., 1844).

For the mortality of different countries and cities, see Mortality.

CLIMATERIC, Climacteric.

CLIMATIC, *Climaticus*, *Climal*. Belonging to, or dependent upon climate.

CLIMATIC DISEASES, *Morbi climatici*, are such as are caused by climate.

CLIMATOLOGY, *Climatologia*, (F.) *Climatologie*, from *κλιμα*, 'a region,' and *λογος*, 'a discourse.' A treatise on climates, their effects on organized bodies, &c., &c.

CLIMBER, WOODY, *Ampelopsis quinquefolia*.

CLINE, *κλινη*, 'a bed.' Hence:

CLINIATER, Clinical physician.

CLINIATRIA, Clinical medicine.

CLINIATRUS, Clinical physician.

CLINIC, see Clinique.

CLINICA, Clinical medicine.

CLINICAL, *Clinicus*, (F.) *Clinique*, from *κλινη*, 'a bed.' Relating to a bed.

CLINICAL LECTURE, (F.) *Leçon Clinique*. One given at the bed-side, or on a particular case or cases.

CLINICAL MEDICINE, *Cliniatri'a*, *Clinica*, *Clinice*, (F.) *Médecine clinique*. That which is occupied with the investigation of diseases at the bed-side, or individually.

CLINICAL PHYSICIAN, *Clinia'ter*, *Clinia'trus*, (F.) *Clinicien*. One who practices or teaches clinical medicine.

CLINICE, Clinical medicine.

CLINICIEN, Clinical physician.

CLINICUM, *Clinique*.

CLINIQUE (F.), Clinical. *Clinique*, *Clinicum*, is also used substantively for *École Clinique* or *Clinical School*: a school in which medicine is taught by examining diseases on the patients themselves, at the bed-side, as it were. Thus, the French say,—"La Clinique de la Charité," "The Clinical School of the Hospital La Charité." The term has been introduced into this country, and anglicised *Clinic*.

CLINO-CEPHA'LIA, *Klino-cepha'lia*, from *κλινη*, 'a bed,' and *κεφαλη*, 'head.' Monstrosity in which the head is unusually flattened, owing to synostosis of the parietal and sphenoid bones.

CLINODES, Clinoid.

CLINOID, *Clinoi'des*, *Clino'des*, *Clinoi'deus*, from *κλινη*, 'a bed,' and *ειδος*, 'form.' Resembling a bed.

CLINOID PROC'ESSES, (F.) *Apophyses clinoi'des*, are four processes at the upper surface of the

sphenoid bone, which have been compared to the posts of a bed. On them the pituitary gland rests. They are divided into *anterior* and *posterior*. Each of the anterior terminates in a point called *Transverse Spinous Process*.

CLINOIDEUS, Clinoid.

CLINOPODIA, Thymus mastichina.

CLINOPODIUM ARVENSE, C. vulgare—c. Incanum, Pycnanthemum incanum—c. Majus, C. vulgare.

CLINOPODIUM VULGARĒ, from κλιν, 'a bed,' and πους, 'foot,' so called from the shape of its flowers, *Clinopodium arven'sē seu majus*, *Ocimum sylvestre*, *Melissa clinopodium*, *Thymus sylvestris*, *Wild Basil*, (F.) *Grand Basilic sauvage*. A European plant, Ord. Labiatae, which was formerly considered to be an antidote to the bites of venomous animals, to facilitate labour, relieve strangury, stop diarrhoea, &c.

CLIQUEETIS, Crepitation—c. *Métallique*, *Tintement métallique*.

CLISEOMETER, *Clisiometer*, (F.) *Klisiomètre*, from κλίσις, 'inclination,' and μέτρον, 'a measure.' An instrument, intended to measure the degree of inclination of the pelvis and to determine the relation between the axis of the pelvis and that of the body.—Osiander, Stein.

CLITBUR, *Arctium lappa*.

CLITHE, *Arctium lappa*.

CLITHEREN, *Galium aparine*.

CLITORICARCINOMA, from κλειτορίς, 'clitoris,' and καρκίνωμα, 'cancer.' Cancer of the clitoris.

CLITORIDES INFERIORES LATI ET PLANI MUSCULI, Constrictores cunni.

CLITORIS, from κλητρον, 'a servant who invites guests' [?]. *Clei'toris*, *Clei'toris*, *Clei'thoris*, *Clei'torion*, *Clei'thorion*, *Dulce'do Amor'is*, *Venus*, *Myrtos*, *Murtos*, *Murtum*, *Estrium*, *Extrus*, *Coles femina'rum*, *Penis mulie'bris seu femineus*, *Men'tula mulie'bris*, *Membrum mulie'bre*, *Supercl'bia*, *Cercoc'his*, *Hypoder'mis*, *Nympha*, *Nymphē*, *Epider'his*, *Libi'dinis Sedes*, *Tunic'ula*, *Crista*. A small, round organ, situate at the upper part of the vulva, and separated by a small space from the anterior commissure of the labia. Its base is surrounded by a slight fold, similar to the prepuce; and it is, internally, of the same structure as the *corpora cavernosa penis*. The artery of the clitoris, (F.) *A. Clitorienne* ou *Clitoridienne*, is a branch of the internal pudic. The vein communicates with the internal pudic, and the nerve with the pudic.

CLITORISM, *Clitoris-mus*. The abuse made of the clitoris. Also, an unusually large clitoris, *Cercoc'his extern'a seu Clitor'idis*, *Cauda'tio*, *Cauda*.

CLITORITIS, *Clitorit'itis*, from κλειτορίς, 'clitoris,' and *itis*, 'denoting inflammation.' Inflammation of the clitoris.

CLITORITITIS, Clitoritis.

CLITORIUM, Vulva.

CLIVER, *Galium aparine*.

CLIVUS BLUMENBACH'IL. An inclining surface behind the dorsum ephippii of the sphenoid bone, which is continuous with the basilar process of the occipital bone.

CLOACA, (F.) *Cloaque*, akin to κλυζω, 'I wash' [?]. The pouch at the extremity of the intestinal canal, in which the solid and liquid excretions are commingled in birds, fish and reptiles. In the male, it gives exit to the excrements, sperm and urine: in the female, to the eggs, fecal matters, and urine.

CLOANX, Orpiment.

CLOAQUE, Cloaca.

CLOCHE (F.). A popular expression in France for a blister or other vesicle.

CLOISON, Septum—c. *des Fosses nasales*, *Septum narium*—c. *Transparente*, *Septum lucidum*.

CLOISONNEMENT, see Generation. Also, the division of a hollow organ—as the bladder, vagina, and uterus, into two portions by a septum,—a malformation—*anomalie par cloisonnement*.

CLONIC, *Clon'icus*, from κλονος, 'agitation,' 'motion.' (F.) *Clonique*. Irregular convulsive motions.' Convulsion with alternate relaxation; in contradistinction to *tonic*, which signifies a constant rigidity;—*Clonus*, *Clon'ici partia'les*.

CLONIC SPASM, see Spasim.

CLONICI UNIVERSALES, Synclonus.

CLONIQUE, Clonic.

CLONIS, Sacrum.

CLONISM, *Clon'ismus*; same etymon. Clonic spasm.—Baumes.

CLONODES, from κλονος, 'agitation,' and εἶδος, 'resemblance.' Convulsive. Galen applies this term to the pulse, when convulsive, as it were, and unequal. He compares the sensation it communicates to the finger to that produced by a bundle of sticks or of rods in vibration.

CLONOS EPILEPSIA, Epilepsy—c. *Hydrophobia*, *Hydrophobia*.

CLONUS, Synclonus—c. *Nictitatio*, *Nictation*—c. *Palpitatio*, *Palpitation*—c. *Pandiculatio*, *Pandiculation*—c. *Pandiculatio maxillarum*, *Yawning*—c. *Singultus*, *Singultus*—c. *Sternutatio*, *Snecing*—c. *Subsultus*, *Subsultus tendinum*.

CLOPEMANIA, Kleptomania.

CLOPORTES ORDINAIRES, Onisci aselli.

CLOSE-STOOL, Lasanum.

CLOT, Coagulum—c. of Blood, see Blood.

CLOTBUR, Xanthium.

CLOTHING, Vestitus.

CLOTTY, *Grumo'sus*, (F.) *Grumelleux*. Composed of clots.

CLOU, Clavus, Furunculus—c. *de Girofle*, see *Eugenia caryophyllata*—c. *Hystérique*, *Clavus hystericus*—c. *de l'Œil*, *Clavus oculi*—c. *de Seigle*, *Ergot*.

CLOUDBERRIES, *Rubus chamæmorus*.

CLOUDBERRY TREE, *Rubus chamæmorus*.

CLOVE, see *Eugenia caryophyllata*—c. *Bark*, see *Myrtus caryophyllata*—c. *Pink*, *Dianthus caryophyllus*—c. *July flower*, *Dianthus caryophyllus*.

CLOVE-TONGUE, *Helleborus niger*.

CLUBFEET, Kyllosis.

CLUBMOSS, *Lycopodium*—c. *Common*, *Lycopodium complanatum*—c. *Fir*, *Lycopodium selago*—c. *Upright*, *Lycopodium selago*.

CLUNES, Nates.

CLUNE'SIA, from *Clunes*, 'the nates.' *Proctal'gia*, *Proctit'is*. An inflammation of the buttocks.—Vogel.

CLUNIS, Sacrum.

CLUPEA THYRSSA. The *Yellow-billed Sprat*; a very poisonous fish of the West Indies.

CLUTIA ELUTERIA, *Croton cascarrilla*.

CLYDON, Flatulence.

CLYPEALIS (Cartilago.) Thyroid cartilage.

CLYSANTLIUM, Syringe, Clyster.

CLYSIS, *Clymus*. Same etymon as clyster. The application of a clyster. Washing out by means of a clyster.

CLYSMA, Clysis, Clyster, Enema—c. *Tonicum*, *Enema fetidum*.

CLYSMATIÖN, same etymon as clyster. A diminutive of clysmas. A small clyster.

CLYSMUS, Clysis, Enema.

CLYSOIR ou CLYSOIRE (F.). An apparatus for administering enemata, consisting of a long funnel-shaped tube of elastic gum, furnished with a pipe at the apex. An instrument, similar to this, with a syringe attached, is called a *Clyso-pompe*.

CLYSOPOMPE, *Clysoir*, Syringe.

CLYSTER, *Clysterium*, *Clysmas*, *Enclys'ma*,

Clymus, Enema, Lavamentum, κλυστήρ, from *κλύω*, 'to wash.' (F.) *Clystère, Lavement, Remède*. A liquid, thrown into the large intestines by means of a syringe, or bladder and pipe properly prepared, &c.; the nozzle of the syringe or pipe being introduced into the anus. See *Enema*.

CLYSTÈRE OF ALOES, *Enema aloes*—c. *Anodyne, Enema anodynum*—c. of *Assafœtida, Enema assafœtida*—c. *Cathartic, Enema catharticum*—c. of *Colocynthis, Enema colocynthis*—c. *Common, Enema commune*—c. *Domestic, Enema commune*—c. *Fetid, Enema fetidum*—c. *Pipe, Elasma*—c. *Purgine, Enema cataarticum*—c. *Starch and opium, Enema anodynum*—c. *Tobacco, Enema tabaci*—c. *Turpentine, Enema terebinthinae*—c. *Uterinus, Sparallium*.

CLYSTÈRE, *Enema*.

CNEME, *Leg, Tibia*.

CNEMODACTYLÆUS, *Extensor communis digitorum pedis*.

CNEMOLORDOSIS, from *κνήμη*, 'the leg,' and *λόδος*, 'the state of being bent forward.' Bending of the leg forward.

CNEMOSCOLIOSIS, from *κνήμη*, 'leg,' and *σκολιωσις*, 'bending,' especially sideways. Bending of the leg sideways. The state of being bow-legged, or bandy-legged, (Prov.) *Serog-legged, shackle-hamned, (L.) Valgus*.

CNEORON, *Daphne gnidium*.

CNEORUM TRICOC'CUM, *Almeze'rión, Aconit'us, Chamael'a, Widow-wail, Spurge-Olive, (F.) Caméléé*. This plant, *Ord. Terebinthaceæ*, a native of Southern Europe, contains a very irritating, acrid principle. The ancients employed its leaves as a powerful purgative. It is now sometimes used for deterring ulcers.

CNESIS, *Cnesmos, Cnismos*. A painful itching. —Galen.

CNESMA, *Itching*.

CNEMOS, *Cnesis, Itching, Prurigo*.

CNICELE'UM, from *κνικος*, 'carthamus,' and *λαιον*, 'oil.' Oil of carthamus. —Dioscorides.

CNICUS, *Carthamus tinctorius*—c. *Sylvestris, Centaurea benedicta*.

CNIDELÆ'ON, *Cnidelæ'um*, from *κνίδεις*, 'cnidian,' and *λαιον*, 'oil.' Oil made from the grana cnidia or mezereon berries. —Dioscorides.

CNID'IA GRANA, *Cnidii Cocci, Coccognid'ia, Eto'lión, Coccum*. The berries of the *Daphne gnidium*. —Fœsius.

CNIDIUM SILAUS, *Peucedanum silaus*—c. *Tenuifolium, Sison ammi*.

CNIDOSIS. A pungent itching, compared to that produced by the *Urtica urens* or *Nettle*. (*κνίδω*). —Hippoc. *Urticaria; urticatio*.

CNIPOTES, *Pruritus*. Itching. The dry ophthalmia, *Xerophthalmia*.

CNISMOREGMIA, *Pyrosis*.

CNISMOS, *Cnesmos*.

CNISSA, see *Nidorous*.

CNISSOREGMIA, from *κνισσα*, 'the smell of burnt fat,' and *αγρυς*, 'I put forth,' *Ructus nidorous*. A nidorous eructation, as of rotten eggs.

CNYMA, *κνύμα*. A slight itching. Also, a puncture or vellication. —Galen.

COACTUS, see *Curds*.

COACUM, *Phytolacca decandra*.

COAGMENTATIO, *Gomphosis*.

COAGULABLE LYMPH, *Fibrin, Liquor sanguinis*.

COAGULANTS, *Coagulan'tia*, from *coagulare*,—itself from *co* and *agere*, 'to act together,' or from *cogo*, 'to drive or bring together.' Remedies, or poisons, supposed to possess the power of coagulating the blood, or to give consistency to animal fluids. The word and the absurd notions connected with it are equally abandoned.

COAGULA'TION, *Coagula'tio, Thrombo'sis*. The conversion of a liquid into a more or less soft and tremulous mass. Many animal and vegetable fluids are capable of coagulation.

COAGULATUS, see *Curds*.

COAG'ULUM, *Grumus, (F.) Caillot, Grumeau*, ('a small clot.') A soft mass formed in a coagulable liquid. The *Clot of the Blood* is particularly so 'called—the *Cruor, Insula, Placenta, Hepar San'guinis, Crassamentum, Sanguis concretus*—the red mass, composed of fibrin, serum, and colouring matter, which separates when the blood is left to itself. See *Blood*. The word is also applied, in pathology, to the sanguineous concretions, which form in different natural and accidental cavities; and which, when they occur at the mouth of a divided artery, sometimes suspend the flow of blood. This is, indeed, one of the means by which hemorrhage is arrested.

COAG'ULUM ALU'MINIS, *Coagulum Aluminosum, Cataplasma aluminis, Alum curd or cataplasm*. This is made by beating the white of egg with a little alum, until a coagulum is formed. It is applied in cases of ophthalmia, where an astringent is necessary.

COALES'CENCE, *Coalescent'ia, Coalit'io partium, from coalescere, 'to run together,' (co, and alere, 'to nourish,') Sym'physes, Prost'ophysis*. The adhesion or union of parts previously separated, as in case of wounds and preternatural adhesions or malformations. See *Monster*.

COALFISH, see *Oleum jecoris aselli*.

COALITIO, *Coalition, Symphysis*.

COALITIO PARTIUM, *Coalescence*.

COALIT'ION, *Coalit'io*. The same etymon as *coalescence*. It has been used in the same cases; as well as to express the action of several parts of the frame, which have the same nutrition.

COALITUS, *Symphysis*.

COALTER'NÆ FEBRES. A name given to two intermittents, which attack a person at the same time, but whose paroxysms are distinct: so that the attack of one supervenes when the other has ceased. The term *Double Intermittent* expresses the same meaning.

COAPTA'TION, *Coapta'tio*, from *co*, and *aptare*, 'to adjust,' 'adapt,' *Paragoge*. The act of adapting the two extremities of a fractured bone to each other; or of restoring a luxated bone to its place. *Coaptation* must be effected gently. Usually, extension and counter-extension are, in the first place, necessary.

COARTICULATIO, *Diarthrosis, Synarthrosis, Coarctatio, Arctatio, Coarctation, Stricture*—c. *Ventriculi, Stricture of the Stomach*.

COARCTA'TION, *Coarctatio*, from *coarctare*, 'to straiten.' Stricture. Avicenna speaks of *Coarctation of the Pulse*.

COARCTOTOMIE, *Coarctotom'ia*, a hybrid term, from *coarctare*, 'to narrow,' and *τομή*, 'incision.' Urethrotomy, when limited to the simple division of a stricture.

COAT, see *Enduit*—c. *Buffy, Corium phlogisticum*.

COATED, see *Enduit*.

COBALT, PROTOXIDE OF, *Smalt*.

COBHAM, MINERAL WATERS OF. Cobham is seven miles from Kingston, in Surrey, England. The waters are weak saline purgatives.

COBRA DI CAPELLO, *Crotalus horridus*.

COBWEB, *Araneæ tela*.

COCASH, *Erigeron Philadelphicum*.

COCCA'RUM. A very small pill.

COCCHIA, *Cochia*.

COCCI GRANUM, *Kermes*—c. *Orientales*, see *Menispermum cocculeus*.

COCCIGIO-CUTANÉ SPHINCTER, Sphincter ani externus.

COCCINELLA, diminutive of *kokkos*, *coccus*, 'a berry' which it resembles. *Coccus caeti*.

COCCINELLA SEPTEMPUNCTATA, *Lady-bird*, *Lady-cow*, *Lady-bug*, *Cow-lady*, *Cushy-cow-lady*, (Prov.) *Fly-golding*, *Golden-bug*, *Golden-knop*. This insect, bruised upon an aching tooth, has long been regarded as anti-odontalgic.

COCCION, *κοκκίον*. A weight, mentioned by Myrseus: the same as the *silqua*.

COCCIONELLA, *Coccus caeti*.

COCCOBALSAMUM, see *Amyris opobalsamum*.

COCCOGNIDIA, *Cnidia grana*.

COCCOLOBA UVIFERA, see *Kino*.

COCCONES, see *Punica granatum*.

COCCULE OFFICINARUM, see *Menispermum cocculus*.

COCCULUS CRISPUS, *Menispermum tuberculatum*—*c. Indi aromaticus*, see *Myrtus pimenta*—*c. Indicus*, see *Menispermum cocculus*—*c. Palmatus*, *Columba*—*c. Suberosus*, *Menispermum cocculus*.

COCCUM, *Cnidia grana*, *Kermes*—*c. Baphicum*, *Kermes*—*c. Infectarium*, *Kermes*—*c. Scarlaticum*, *Kermes*—*c. Tinctorum*, *Kermes*.

COCCUS, (*κοκκος*, 'a berry,') *Coccus caeti*—*c. Americanus*, *Coccus caeti*.

COCCUS CACTI. The systematic name of the *Cochineal Insect*, *Coccinella*, *Cochinilla*, *Coccinella*, *Coccinella*, *Ficus Indica grana*, *Coccus Polonicus seu Americus seu Indicus Tinctorius*, *Scarabaeolus hemisphaericus*, *Cochinelifera cochinilla*, *Coccus Americus*, *Cochinella*, *Coccus*; the *Cochineal Animal*, (F.) *Cochenille*, *Graine d'Écarlate*. *Class*, *Insecta*. *Order*, *Hemiptera*. The cochineal insects have a faint, heavy odour; their taste is acrid, bitterish, and astringent; colour blackish-red externally, purple-red within. They are used chiefly for giving a red colour to tinctures, &c. They were, at one time, esteemed astringent, stimulant, diuretic, and anodyne.

COCCUS ILICIS, *Kermes*—*c. Indicus tinctorius*, *Coccus caeti*—*c. Indicus*, see *Menispermum cocculus*—*c. Polonicus*, *Coccus caeti*.

COCCYCEPHALUS, *Coccygo-cephalus*, (F.) *Coccycephale*, from *coccyx* and *κεφαλή*, 'the head.' A monster whose head has the shape of the os coccygis.

COCCYGEAL, *Coccygeus*—*c. Nerve*, see *Sacral Nerves*.

COCCYGEUS, *Coccygeal*, (F.) *Coccygien*, from *κοκκυς*, because it is inserted into the coccyx; *Ischio-Coccygeus*, (F.) *Ischio-coccygien*. Belonging both to the ischium and coccyx. The muscle *Ischio-coccygeus*, *Levator Coccygis*, *Triangularis Coccygis*. It arises from the spinous process of the ischium, and is inserted into the extremity of the sacrum; and into nearly the whole length of the os coccygis laterally. It retains the coccyx in place, and prevents it from being forced backwards during the evacuation of the faeces.

COCCYGIO-ANAL, Sphincter ani externus.

COCCYGIO-CUTANÉ SPHINCTER, Sphincter ani externus.

COCCYGOCEPHALUS, *Coccycephalus*.

COCCYMELEA, *Prunus domestica*.

COCCYX, 'a cuckoo,' whose bill it is said to resemble; *Os Coccygis seu Alagus*, *Cauda*, *Ossis sacri acutamen*, *Rump* or *Crupper Bone*, *Cuculus*, *Uropygion*, *Spondylis*, *Spondylium*. An assemblage of small bones, attached to the lower part of the sacrum; the curvature of which it prolongs, and of which it seems to be an appendage. Generally, it consists of four bones. Behind the

base of the coccyx are two small tubercular eminences. These are called *Cornua of the Coccyx*.

COCHEMAR, *Incubus*.

COCHENILLE, *Coccus caeti*.

COCHERIA, *Cochia*.

COCHIA, *Cochia*, *Cocheria*, from *κοκκος*, 'berry,' 'seed,' or from *κοχμα*, 'to flow profusely.'

An ancient name for several official purgative pills; thus called, either because they produced copious evacuations, or were shaped like a seed.

COCHIN LEG, see *Elephantiasis*.

COCHINEAL, *Coccus caeti*.

COCHINELIFERA COCHINILLA, *Coccus caeti*.

COCHINILLA, *Coccus caeti*.

COCHLEA. A *Snail's shell*, (F.) *Limacon*, *Coquille*, *Cochlée*. Anatomists have given this name to the most anterior of the three cavities, which constitute the labyrinth of the ear, the *Pelvis Auditoria*, *Concha auris interna seu Labyrinthi*; *Cavitas cochleata seu buccinata*, *Antrum buccinae seu Trochlea labyrinthi*:—and that of *Scala of the Cochlea*, (F.) *Rampes du limacon*, to two spiral cavities in the interior of the cochlea. One of these scales terminates at the *Fenestra rotunda*, and is called *Scala tympani*: the other opens at the anterior and inferior part of the vestibule: it is called *Scala vestibuli*.

COCHLEA, *Scala of the*, see *Cochlea*.

COCHLEAR, *Cochlearis*, (F.) *Cochlaire*, *Cochléarien*. Relating or appertaining to the cochlea.

COCHLEAR, *Cochleare*—*c. Auriculare*, *Ear-pick*—*c. Nerve*, see *Auditory Nerve*.

COCHLEARE, *Cochlear*, *Cochlearium*, from *cochlea*; its bowl resembling a shell. A *spoonful*, (F.) *Cuillerée*; abbreviated in prescriptions usually to *coch*. See *Abbreviation*. Also, a *scop*.

COCHLEARE MAGNUM, a *tablespoonful*; *C. Medium*, a *dessert or pap-spoonful*; and *C. Minimum*, a *teaspoonful*.

COCHLEARIA, from *cochleare*, 'a spoon,' so called from its resemblance. *C. officinalis*.

COCHLEARIA ARMORACIA, *Raphanus rusticus seu marianus seu sylvestris seu magna*, *Armoracia*, *A. sativa seu rustica*, *Horseradish*; (Sc.) *Rotcoll*. *Fam.*, *Cruciferae*. *Ser. Syst.* *Tetradynamia Siliculosa*. (F.) *Raifort sauvage*, *Cran*, *Cran de Bretagne*, *Cranson*, *Montagne des Moines* ou *des Allemands*, *Radis de cheval*. The root of horseradish is frequently used at table; and has long been in the *Materia Medica*. It is stimulant and diuretic. Externally, it is rubefacient.

COCHLEARIA CORONOPUS, *Coronopus*, *Coronopus Ruelii* seu *depressus* seu *vulgaris*, *Lepidium squamatum*, *Senebiera coronopus*, *Wild Scoury Grass*, *Swine's Cress*, (F.) *Corne de Cerf*. This European plant is considered to be diuretic and antiscorbutic. The term *Coronopus* was given, by the ancients, to various plants.

COCHLEARIA HORTENSIS, *Cochlearia officinalis*.

COCHLEARIA OFFICINALIS, *Cochlearia*, *C. hortenensis* seu *pyrenaica* seu *vulgaris*, *Lemon Scoury Grass*, *Common Scoury Grass*, (Sc.) *Scrub-grass*, (Prov.) *Seroby-grass*, (F.) *Cranson*, *Herbe aux cuillers*. It has been considered a powerful antiscorbutic. It is sometimes eaten in salad.

COCHLEARIA PYRENAICA, *C. officinalis*—*c. Vulgaris*, *C. officinalis*.

COCHLEARIFORMIS, from *cochleare*, 'a spoon,' and *forma*, 'shape,' (F.) *Bec de Cuiller*.

COCHLEARIFORMIS PROCES'SUS, *Cochleariform process*. A small, very thin plate, which separates the bony portion of the Eustachian tube from the canal for the passage of the tensor tympani.

COCHLEARIS. A gelatinous looking tissue, seen on opening the cochlea, by which the mem-

braneous zone is connected, at its outer or convex margin, with the outer wall. It is supposed by Todd and Bowman to be muscular; and to have a preservative office, being placed to defend the cochlear nerves from undue vibrations of sound, in a way analogous to that in which the iris protects the retina from excessive light.

COCHLEARIUM, Cochleare.

COCHO'NĒ, κοχυνή. The junction of the ischium, near the seat or breech.—Fœsius. The breech proper, from the hip-bones to the anus. The perinaeum. The coccyx.

COCL'IO. A weight of eleven ounces.

COCK-EYE, see Luscitas.

COCK-GRASS, Lolium temulentum.

COCKLE, CORN, Lychnis githago.

COCKLE-BUR, Agrimony, Xanthium.

COCKLES, INDIAN, see Misperspermum cocculus.

COCKMINT, Tanacetum balsamita.

COCKSPUR, Ergot—c. Rye, Ergot.

COCKUP HAT, Stillingia.

COCKYBABY, Arum maculatum.

COCLES, *Borgue*.

COCO, Cocos nucifera.

COCO OF THE MALDIVES', *Cocos de Maldive*.

The fruit of a palm, called *Lodoicea* by Comerson. It was formerly termed, in the shops, *Nux Medica*, and enjoyed great reputation.

COCOA, Cacao, Cocos nucifera.

COCOA CACAVIFERA, Cacao.

COCOBAY, Mal de San Lazaro.

COCO-OLEIN, see Cocos nucifera.

COCOS BUTYRACEA. The name of the plant which affords the palm oil, *Oleum palmæ*, obtained chiefly from the fruit, by bruising and mixing the kernels with water, without the aid of heat, by which the oil is separated and rises to the surface. It is of the consistence of an ointment, with little taste, and a strong, though not disagreeable, smell. It is applied to sprains, &c.; but has no advantages over other oils. It has been used as the excipient for suppositories, and has been called, also, *Oleum Palmae sebaeum*, *O. fixum nucum cocos butyrateæ* and *Mackeani fat.* It is procured likewise from the *Ela's Guineensis*, and *Ela's Occidentalis*, two species of palms. Cocos is said to be contracted from (P.) *maccoco* or *maccaco*, 'a monkey,' the three holes at the end of the cocoa-nut shell causing it to resemble somewhat a monkey's head.

COCOS NUCIFERA, *Palma cocos*, (F.) *Cocotier*. Order, Palmæ. The systematic name of the plant whose fruit is the cocoa-nut. (F.) *Coco*. It is an agreeable kernel, but not easy of digestion. *Emulsions*, *orgeat*, &c., are made from it. The juice of the cocoa, *Toddy*, when fermented, forms wine, and arrack is distilled from it. The expressed oil—*Coco-olein*—has been found almost as efficacious as cod-liver oil in tuberculosis.

COCOTE, Influenza.

COCOTIER, Cocos nucifera.

COCOWORT, *Thlaspi bursa*.

COC'IO, Coction, Digestion—c. *Ciborum*, Digestion—c. *Morbi*, Coction.

COC'TION, *Coc'tio*, *Pepsis*, *Pepan'sis*, *Pepas'mos*, *Symp'es'sis*, *Concoct'ion*, from *coquere*, *coc'tum*, 'to boil.' This word has been employed in various senses. 1. With the ancients, *coction* meant the particular kind of alteration which the food experiences in the digestive organs, particularly in the stomach. It meant a preparation from its crude state. 2. It expressed the maturation or change, which the humoral pathologists believed morbid matter experiences before elimination. It was considered, that coction, *Coc'tio morbi*, was produced during the violence of the disease; and hence this was called the *Period of Coction*. See Humorism.

COC'YTA, (F.) *Cocyle*, from *κωκυτος*, 'lamontation.' The pain caused by the introduction of poison under the skin.—Linnæus.

COD, or CODDS, Scrotum.

COD-OIL, Oleum jecoris aselli.

COD-LIVER OIL, Oleum jecoris aselli—c. Olein, see Oleum jecoris aselli.

CODAGAPALA BARK, Nerium antidysentericum.

CODE, Codex.

CODĒIA, Codeine.

COD'EINE, *Codei'a*, *Codei'num*, *Papaveri'num*, from *κωδία*, 'a poppy head.' An alkaloid discovered by Robiquet in opium, in 1832. It is soluble in water, alcohol and ether, and unites readily with acids. As a hypnotic, Magendie thinks one grain of codeia equal to half a grain of morphia. The muriate of codeia appears to be stronger than the pure codeia.

CODESELLA, Anthrax.

CODEX. A collection of laws. (F.) *Code*. By extension, a collection of approved medical formulæ, with the processes necessary for forming the compounds referred to in it. The Parisian Pharmacopœia is styled *Codex medicamentarius*.

CODEx MEDICAMENTARIUS, Pharmacopœia, see Codex.

CODIA, Papaver (capsule.)

CODOCELE, Bubo.

CODOSCELLA, Bubo.

CŒCAL, Cæcal.

CŒCITAS, Cæcitas.

CŒCUM, Cæcum.

COEFFE, Caul.

CŒ'LIA, κοιλία, κοιλῆ, 'a hollow place.' This word generally meant a cavity in the body:—the abdomen, in particular. It has also been used for the alimentary canal, &c.:—*ανὰ κοιλία*, 'the stomach,' *κατὰ κοιλία*, 'the abdomen.' Also, an alvine evacuation; excrement.

CŒ'LIAC, *Cœ'liacus*, *Gastrocœ'liacus*, *Gastrocœ'licus*, (F.) *Cœlique* ou *Cœlique*, from *κοιλία*, 'the cavity of the abdomen.' Relating to the cavity of the abdomen.

CŒLIAC ARTERY, *Arte'ria Cœ'liaca*, *Cœliac axis*, *Artère opistogastrique*, (Ch.) *A. Cœliaque*, *Tronc cœliaque*, *Trépid de la cœliaque*, is situated above the pancreas, and behind the upper part of the stomach. It arises from the anterior part of the abdominal aorta, where it passes between the pillars of the diaphragm, and soon divides into the *coronaria ventriculi*, *hepaticæ*, and *splenicæ* arteries.

CŒLIAC FLUX, *Cœliac Passion*, *Cœliaca chylo'sa*, *Diarrhœ'a chylo'sa* seu *chylo'sa*, *Fluxus chylo'sus* seu *cœliacus*, *Passio calinea* seu *Ventriculo'sa*, *Chymoche'zia*, *Fluor caline intestin'um*, *Chylorrhœ'a*, *Chymorrhœ'a*, *Cœliaca lactea*, *Morbus cœliacus*, *Chylodiarrhœ'a*, *Galactodiarrhœ'a*, *Sedes lactescen'tes*, *Gastrorrhœ'a*, (F.) *Flux cœliaque*. A species of diarrhœa, in which the food is discharged by the bowels in an undigested condition. By some, defined to be diarrhœa attended with discharges of chyle or chyme. It is, in general, symptomatic of tubercular disease of the mesenteric glands. See *Lientery*.

CŒLIAC PASSION, Cœliac flux.

CŒLIAC PLEXUS, *Solar Plexus*, *Plexus mesenter'ii pri'us et max'imus*, *P. gangliofor'mis semilunaris*, (F.) *Plexus mœdian* ou *opistogastrique*, (Ch.), *Pl. Cœliac* ou *solaire*, *Ganglion de Vieussens*, is formed of numerous nervous filaments, which proceed from the semilunar ganglia of the great sympathetic. It is strengthened by several branches of the right and left pneumogastric nerves; is seated around the trunk of the cœliac artery, behind the stomach, and furnishes the secondary plexuses—the *diaphrag-*

matic, coronary of the stomach, epenic and hepatic, which accompany the arteries of the same name.

CÆLIACA, from *κοιλιακος*, 'celiac.' Diseases of the digestive organs; the 1st class in Good's *Nosology*. It comprises two orders, *Enterica* and *Splanchnica*. Also, medicines that act on the digestive organs.—Pereira.

CÆLIACA CHYLOSA, Celiac flux—c. Lactea, Celiac flux—c. Renalis, Chyluria—c. Urinalis, Chyluria.

CÆLIADELPH'US, from *καλῖα*, and *ἀδελφος*, 'brother.' *Anocæliadelph'us*. A double monster, in which the two bodies are united at the abdomen.

CÆLIÆMIA, *Hyperæmia abdom'inis*, *Congestio abdomin'alis*, from *κοιλια*, 'abdomen,' and *αἷμα*, 'blood.' Hyperæmia or congestion of the blood-vessels of the abdomen.

CÆLIAGRA, *Gastrit'is seu Enterit'is seu Colic'ia seu Diarrhœa arthrit'ica*. Gout of the abdomen.

CÆLIALGIA, Tormina.

CÆLIAQUE, *TRÉPIED DE LA*, Celiac artery.

CÆLIOCELE, see *Hernia*, hypogastric.

CÆLIOCHYSIS, Ascites.

CÆLIOCYESIS, Pregnancy, abdominal.

CÆLIODYNIA, Colic.

CÆLIOLYSIA, Diarrhœa.

CÆLION'CUS, *Cæliophy'ma*, from *κοιλια*, 'the abdomen,' and *ογκος*, 'a tumour.' A tumour of the abdomen.

CÆLIOPHYMA, Cælioncus.

CÆLIOPHY'MATA, *Tuber'cula peritonæ'i*, from *κοιλια*, 'the cavity of the abdomen,' and *φύμα*, 'a hard tumour.' Tubercles of the peritoneum.

CÆLIOPSOPHIA, Borborygmus.

CÆLIOPYO'SIS, from *κοιλια*, 'the abdomen,' and *πυωσις*, 'suppuration.' Suppuration of the abdomen or its parietes.

CÆLIORRHŒA, Diarrhœa.

CÆLIORRHŒUMA, *Rheumatismus abdom'inis*, from *κοιλια*, 'the abdomen,' and *ρευμα*, 'de-fluxion, rheumatism.' Rheumatism of the muscles of the abdomen.

CÆLIOSPASM'US, from *κοιλια*, 'abdomen,' and *σπασμος*, 'spasm.' Spasm or cramp of the abdomen.

CÆLO'MA, from *κοιλος*, 'hollow.' A round ulcer of the cornea, broader and deeper than that described under the name *Bothrion*.

CÆLON, Cavity.

CÆLOPHLEB'TIS, from *κοιλος*, 'cavus,' *φλεψ*, *φλεβος*, 'a vein,' and *itis*, denoting inflammation. Inflammation of the vena cava.

CÆLOPHTHAL'MUS, from *κοιλος*, 'hollow,' and *οφθαλμος*, 'eye.' One who is hollow-eyed.

CÆLOSTOM'IA, from *κοιλος*, 'hollow,' and *στομα*, 'mouth.' A defect in speaking, when the voice seems as if it came from a cavern;—that is, obscure, and as if at a distance.

CÆLOSTOMUS, same etymon. One who has a hollow voice.

CÆLOTES, Cavity.

CÆNADELPH'US, from *κοινος*, 'common,' and *ἀδελφος*, 'brother.' A double monster, the two portions having, in common, one or more organs indispensable to life.

CÆNÆSTHES'IS, from *κοινος*, 'common,' and *αισθησις*, 'feeling.' *Cænæsthêsia*, (F.) *Cænæsthêsie*. Common feeling. Some German writers mean, by this, a sixth sense. It is the feeling of self-existence or individuality, and is manifested by the sense of buoyancy or depression, which we experience without any known cause;—by involuntary shuddering, feeling of chill or glow, &c.

CÆNOLOGIA, Consultation.

CÆNOTES, *κοινωτης*, 'commonness,' from *κοινος*, 'common.' The physicians of the methodic sect asserted, that all diseases arise from *relaxation, stricture*, or a mixture of both. These were called *Cænotes*: or what diseases have in common.

COËTUS, Coition.

COËUR, Heart.

COFFŒA ARAB'ICA, *C. vulg'aris*, *Jas'minum Arab'icum*, (F.) *Caffier*, *Caffier*, and *Caffier*. Order, Rubiaceæ. Sex. Syst. Pentandria Monogynia. The plant which affords coffee, *Choua*, *Bon*, *Buna*, (F.) *Café*. Originally from Yemen.

The infusion of coffee is an agreeable and wholesome article of diet. It is, to a certain extent, a tonic, and is employed as such in convalescence, especially from fevers, &c. In cases of poisoning by opium, and in old asthmas, its use has been extolled. For this purpose the *Moka* is the best. It ought to be newly torrefied, but not too much burnt; should be strong, and drunk soon after grinding.

A highly azoted alkaloid was first found in coffee, and called *Caffein* or *Caffein*. It was afterwards discovered in the leaves of the tea plant, and called *Thein*, and in the Guarana of Brazil, and called *Guaranin*. In doses of from 2 to 10 grains, it induces violent nervous and vascular excitement. Caffein—as well as its various salts—the citrate, lactate, malate, and murate, have been given as an antiperiodic in grain doses.

Facitious Coffees have been, from time to time, recommended, but they are infinitely inferior to the genuine. Various substitutes have been proposed; wheat, barley, hollyberries, acorns, sunflower seeds, beech-mast, peas, beans, succory-root, seeds of gooseberries and currants left in making wine, and washed,—sliced turnips, &c. These have been roasted, with the addition of a little butter or oil: but they have not the aroma of coffee. The best substitute is said to be the seeds of the Yellow water flag, *Gladiolus luteus* or *dris pseudacorus*.

Hunt's Economical Breakfast Powder consisted of rye, roasted with a little butter.

COFFEE-BEAN TREE, KENTUCKY, *Gymnocladus Canadensis*.

COFFEE-TREE, *Gymnocladus Canadensis*.

COFFEE, WILD, *Triosteum*.

COGNASSIER, *Pyrus cydonia*.

COGNITIO PHYSIOLOGICA, Physiology

COHABITATIO, Coition.

COHABIT'ATION, *Cohabita'tio*, from *cum*, 'with,' and *habitare*, 'to dwell.' The act of dwelling together. In legal medicine, it means the consummation of marriage.—*Copulation*.

COHES'ION, *Cohæsi'o*, from *co*, and *herere*, 'to stick.' *Vis cohæsi'o'nis seu adhæsi'o'nis seu attracti'o'nis*, Force of cohe'sion, Attraction of cohe'sion or of aggrega'tion, is that force in the particles of matter, whereby they are connected in such a way as to resist any attempt towards their removal or separation. This force has to be attended to, in the management of disease. Emollients, rubbed into a part, act by diminishing the cohesion.

COHIBENS, Epischeticus.

COHOBAT'ION, *Cohoba'tio*, *Cohob'ium*, *Co'hob*, *Co'hoph*. Distillation of a liquid—already distilled—on the same substances. When this is repeated three or four times, it is called *Recohob'ation*.

COHOL. Synonym of Alcohol. Also, a dry collyrium.—*Avicenna*.

COHOSH, *Actæa racemosa*, *Caulophyllum thalictroides*—c. Black, *Actæa racemosa*—c. Blue, *Caulophyllum thalictroides*—c. Blueberry,

Caulophyllum thalictroides—c. Red, *Actæa rubra*—c. White, *Actæa alba*.

COHUSH, *Caulophyllum thalictroides*.

COIFFE, Caut.

COIGN, see *Pyrus cydonia*.

COIGNASSIER, *Pyrus cydonia*.

COINCIDENTIA, from *co*, and *incido* (in, and cado), 'to fall or happen.' Some authors have translated, by this term, the word *parempto'sis*, used by Galen to designate the occlusion of the foramen opticum by a humour, proceeding from the base of the brain, and occasioning blindness!—Castelli.

COIN'DICANT, *Coin'dicans*, from *co*, and *indico*, 'I indicate.'

COIN'DICANT SIGNS are those which furnish the same indications; or which confirm the indication afforded by another sign—*συμπεδικνυσμα*.

COIRA, Catechu.

COIRAS, *Scrofula*.

COIT, Coition.

COIT'ION, *Co'itus*, *Co'etus*, from *co'zo*, (co, and eo, 'to go,') 'to go together.' *Co'iture*, *Copulation*, *Copulatio*, *Cop'ula* *Carna'lis*, *Aphrodis'ia*, *Aphrodisiasm'us*, *Epip'locæ*, *Acces'us*, *Ba'ia'tio*, *Amplexa'tio*, *Amplex'us*, *Conven'tus*, *Comple'xio*, *Conj'ugium*, *Agglutina'tio*, *Lagne'a*, *Lagne'ma*, *Lagne'a*, *Micis*, *Perm'itio*, *Perm'itio*, *Syndys'mus*, *Syn'u'sia*, *Concu'bitus*, *Congres'sus*, *Concubita'tio*, *Venus*, *Res Ven'e'nea*, *Concu'bitus ven'e'reus*, *Præ'tium*, *Duell'um ven'e'reum*, *Noc'turna bella*, *Concu'via corpora'lia*, *Homil'ia*, *Ven'ery*, *Sexual intercourse*, (F.) *Coit*, *Approche*, *Acco'plement*. The carnal union of the sexes.

COITURE, Coitus.

COITUS, Coition—c. *Difficilis*, *Dyssynodus*—c. *Sodomiticus*, Buggery.

COL, Collum—c. *de la Matrice*, Collum uteri—c. *Uteri*, Collum uteri.

COLA, Articulation.

COLATIO, see *Colatura*.

COLATO'RIMUM, *Hylister*, from *colare*, 'to strain.' A strainer of any kind. (F.) *Couloir*. A term by which the ancient physicians described every canal or conduit through which the excrementitious humours of the body are evacuated. Ulcers, fistulae, setons, caustics, &c., have been called artificial or accidental *Colatoria*, because they were considered to be passages by which the animal economy is freed from some morbid matter.

COLATUM, see *Colatura*.

COLATU'RA, from *colare*, 'to strain.' *Colatura*, (F.) *Colature*. A filtered or strained liquor. It likewise means straining a liquid, — *Colatio*, *Dy'lysis*, *Dy'lysmus*, *Hyl'isis*, *Hyl'ismus*.

COLCHESTER, MINERAL WATERS OF. *Aqua Colcestren'sis*. The waters of Colchester are of the bitter purging kind, similar to those of Epsom.

COLCHICIA, see *Colchicum autumnale*.

COLCHICUM, from *Colchis*, in Asia, where it is said to have been common. *Colchicum autumnale*.

COLCHICUM AUTUMNA'LE, from the country *Colchis*, *Col'chicum*, *Coun*, *Meadow Saffron*, (F.) *Colchique*, *Tue-chien*, *Mort aux chiens*, *Sa'fran des prés ou bâtarde*, *Narcisse d'automne*, *Vielotte*. *Nat. Ord.* Colchicaceæ. *Class.* Hexandria. *Order*, Monogynia. The bulb or root (*Bulbus seu Radix seu Cormus*), and the seeds, *Col'chici semen*, are the parts used in medicine. The taste is acrid, excoiating the mouth; but the acrimony is lost by drying. It is narcotic, diuretic, and cathartic; and has been given in dropsy, gout, and rheumatism. Dose, gr. j to vj of the fresh bulb. It is supposed to be the active ingredient of the *Eau médicinale d'Hiisson*. The active

principle is called *Colchic'ia*. *Colchicum*, in an over-dose, is an acro-narcotic poison.

Dr. Wilson's Tincture for the Gout is said to be merely an infusion of *Colchicum*, or *Col'chicin*.

COLCHICUM ZEYLANICUM, *Kæmpferia rotunda*.

COLCHIQUE, *Colchicum autumnale*.

COL'COTHAIR, *C. Vitrioli*, *Henri'cus rubens*, *Chalc'i'tis*, *Brown red*, *Rouge*, *Crocus*, *Oxidum Ferri rubrum*, *Tritox'ydum Ferri*, *Sulphas Ferri calcina'tus*, *Ferrum vitriola'tum ustum*, *Terra vitrioli dulcis*, *Crocus martis vitriola'tus seu astrin'gens*, (F.) *Rouge d'Angleterre ou de Prusse*, *Safran de Mars astringent*. The red oxide of iron, obtained by calcining sulphate of iron to redness, with or without the contact of air. It possesses the general properties of the preparations of iron, and has been applied to staunch blood, &c.

COLD, (Sax.) *colb*, *Frigus*, *Psychos*, (F.) *Froid*. The sensation produced by the abstraction of caloric from our organs,—*Cheima*, *Chimon*. See *Heat*.

Three degrees of cold are generally distinguished in disease. 1. The simple feeling of cold (*Algor*), 2. *Chilliness* (*Horror*), and 3. *Shivering* (*Rigor*). Cold is employed in medicine, chiefly, as a refrigerant and sedative.

Cold is, also, used for a morbid affection induced by cold. 'To take cold,' (F.) *s'enrhumer*, is to be affected by a disease presumed to be caused by cold. See *Catarrh* and *Coryza*.

COLD-CHILL, Intermittent fever.

COLD IN THE EYE, Ophthalmia, catarrhal—c. in the Head, *Coryza*.

COLEITIS, Colpositis.

COLEOCLE, see *Hernia*.

COLECEDEMA, *Elytrocedema*.

COLEOPTOSIS, *Prolapsus vaginæ*.

COLEORRHÆXIS, from *κολεος*, 'a vagina or sheath,' and *ρῆξις*, 'rupture,' Laceration or rupture of the vagina.

COLEOSITIS, *Leucorrhœa*.

COLEOSTEGNO'SIS, *Colpostegno'sis*, *Colpostenochor'ia*, *Colpospynia'sis*, from *κολεος*, 'a vagina or sheath,' and *στεγνσις*, 'constriction.' Narrowness of the vagina, natural or acquired.

COLÈRE, Rage.

COLES, Penis—c. *Feminarum*, *Clitoris*.

COLEUS, Vagina.

COLIBERT, *Cagot*.

COLIC, *Col'icus*, from *κωλον*, 'the colon.' (F.) *Colique*. Relating to the colon.

COLIC ARTERIES, *Arte'ria colica*, (F.) *Artères Coliques*, are six in number, three of which, given off by the superior mesenteric, are called *Colica dextra*; and three, proceeding from the inferior mesenteric, *Colica sinist'ra*. All proceed towards the colon, passing between the two laminae of the peritoneum, which form the mesocolon, where they ramify and anastomose with each other. The first, *Colica dextra*, *Ramus colica dexter*, is called *C. dextra superior*, (*Meso-colique*,—Ch.) The second, *C. dextra media*, *Colica media*, *Ramus colicus medius*, *Arteria media anastomot'ica*, (*C. droite*, Ch.), and the third, *C. dextra infer'ior* or *Neo-colica*, (*A. cæcale*,—Ch.) Of the three *Colica sinist'ra*, the first or superior is called, by *Chaussier*, *Colica magna sinist'ra*; the second or media is not distinguished from the preceding, as they often arise from a common trunk; and the third is called by *Chaussier* *Colica parva sinist'ra*. To these arteries as many veins correspond, which open into the great and little mesenteric.

COLIC LOBE OF THE LIVER is the great lobe of that organ.

COLIC, *Col'lica*, *C. Passio*, *Colica*, *Colicodyn'ia*, *Cœliodyn'ia*, *Dolor'es intestino'rum*, *Dolor col'icus*, *Dysenteronew'ia*, *Enteral'gia*, *Gripes*, *Mulligrubs*, *Bellyache*, (Sc.) *Bellythra*, *Weam-ith*.

(Prov.) *Bellypharm*, *Bellypark*, *Gullion*. In its etymological acceptation, Colic signifies an affection or pain in the colon. But it is employed in a more extensive signification. It includes every acute pain of the abdomen, aggravated at intervals. The word has often, however, epithets affixed to it, which render it more or less precise. See *Tormina*.

COLIC, BILIOUS, Colica biliosa—c. Crapulent, Colica crapulosa—c. Devonshire, Colica metallica—c. Horn, Priapismus—c. Lead, Colica metallica—c. Madrid, Colica Madridensis—c. Menstrual, Colica menstrualis—c. Metallic, Colica metallica—c. Nephretic, Colica nephretica—c. Normandy, Colica metallica—c. Painters', Colica metallica—c. Plumbers', Colica metallica—c. of Poitou, Colica metallica—c. of Prostitutes, Colica scortorum—c. Renal, Colica nephretica—c. Saturnine, Colica metallica—c. Root, Aletris—c. of Surinam, Colica metallica—c. Uterine, Colica uterina—c. Vegetable, Colica metallica—c. Weed, Dientra Canadensis—c. Worm, Colica verminosa.

COLICA ACCIDENTALIS, Colica crapulosa—c. Acuta, Enteritis—c. Arthritica, Cœliagra.

COLICA BILIOUSA, *Bilious Colic*, (F.) *Colique Biliense*. Colic, occasioned by an accumulation of bile in the intestines or in its own passages. The treatment required resembles that proper for Colica convulsiva.

COLICA CALLOSÆ. Colic attended with sense of stricture in some parts of the intestinal canal; often of flatulency and pain: the flatulency gradually passing off by the stricture; the bowels tardy; and at length discharging small liquid stools.

COLICA CONVULSIVA, *C. Spasmodica* seu *pituitosa* seu *nervosa* seu *idiopathica*, *Enteroprasmus*. Colic, not the symptom of any other affection. It is characterized by griping pain in the bowels, chiefly about the navel, with vomiting and costiveness—the pain increasing at intervals. The indications are to clear the intestines, and allay spasm. Calomel and opium—castor oil and opium—emollient and cathartic enemata, with fomentations, wet or dry, to the abdomen, usually succeed.

COLICA CRAPULOSA, *C. accidentalis* seu *heluolum*, *Crapulent colic*, *Surfeit*, *Colic from overeating*, (F.) *Colique d'Indigestion*. A colic, arising from eating indigestible aliments, or digestible aliments in too great abundance. The remedy is obvious.

COLICA DAMNORUM, Colic, metallic—c. Ferriosa, Colica inflammatoria—c. Figulorum, Colica metallica.

COLICA FLATULENTA, *Inflata*, *Gastrodynia flatulenta*, *Physocapnus*, *Pneumatosis enterica*, (F.) *Colique flatulente* ou *flatuense* ou *ventreuse*. Colic, arising from an accumulation of air in the intestines. It is very common in infants, and may be relieved by aromatics, especially when combined with antacids—for example, oil of aniseed with magnesia.

COLICA HELLUOLUM, Colica crapulosa.

COLICA HEMORRHOIDALIS, *Hemorrhoidal Colic*, (F.) *Colique hémorrhoidale*. A kind of colic, supposed to precede hemorrhoids or to supervene on their suppression.

COLICA HEPATICA, *Hepatalgia*, *Hepatalgia Calculosa*, *Hepatic colic*, (F.) *Colique hépatique*. Pain in the region of the liver, chiefly about the gall-bladder, occasioned by the passing of a biliary calculus through the cystic and choledoch ducts.

COLICA IDIOPATHICA, Colica convulsiva—c. Ileus, Ileus.

COLICA INFLAMMATORIA, *C. Phlogistica* seu *plethorica* seu *febricosa* seu *pulsatilis*, *Inflammatory colic*. The same as enteritis.

COLICA LAPPONICA, see *Seta equina*—c. *Loethalis*, *Dyslochia*—c. *Madridensis*, *Colic of Madrid*—c. *Nervosa*, *Colica convulsiva*, *Colic, metallic*—c. *Phlogistica*, *Colica inflammatoria*.

COLICA MADRIDENSIS, *Colic of Madrid*, *Madrid Colic*. A species of colic, endemic in several provinces of Spain, whose nature is not clear. Its symptoms resemble those occasioned by lead. See *Colica metallica*.

COLICA MENSTRUALLIS, *Menstrual Colic*, (F.) *Colique menstruelle*. Colic, which precedes or accompanies the menstrual evacuation, or which is owing to the suppression of that evacuation.

COLICA METALLICA, *Metallic Colic*, *Painters' Colic*, *Colic of Poitou*, *Colic of Surinam*, *Billon*, *Devonshire Colic*, *Dry Bellyache*, *Dry Gripes*, *Saturnine Colic*, *Lead Colic*, *Plumbers' Colic*, *Rachialgia*, *R. Pictonum* seu *Pictariensium*, *Morbus Metallicus*, *Colicoplegia*, *Colica Rachialgia* seu *Damnorum* seu *Plumbariorum* seu *Pictonum* seu *Saturnina* seu *Figulorum* seu *Nervosa* seu *Pictorum* seu *sicca*, *Paralysis rachialgia*, *Palms Plumbarius*, *Molybdocten*, (F.) *Dysentéronervie Saturnine*, *Colique de Poitou* ou *végétale* ou *des peintres* ou *de plomb* ou *métallique* ou *Saturnine* ou *des barbouilleurs* ou *sèche*. Under this head is generally described the colic produced by lead, as well as the other colics mentioned in the synonymy; and they certainly resemble each other greatly, although some of them are more like bilious colic. There is not much to distinguish this variety of colic from others. The same violent pain about the navel is present, with retraction of the abdomen towards the spine. It is apt also to occasion palsy. The only difference of treatment is in the necessity for employing more opium along with the purgative. The paralytic sequelæ must be treated by change of air, rubbing the spine, exercise, &c. Treating the disease upon general principles is infinitely more philosophical, and more successful than the former empirical management at *La Charité*, Paris, which it is unnecessary to detail.

The *Colic of Madrid*, *Poitou*, *Devonshire*, *Normandy*, &c., has been called *vegetable colic*, (F.) *Colique végétale*, because ascribed to the use of new wine or cider, unripe fruit, &c., although really, perhaps, owing to the adulteration of wines and cider by litharge. All the symptoms, indeed, are those of Colica metallica.

COLICA NEPHRETICA, *Nephrocolica*, *Colica renalis*, *Nephretic* or *Renal Colic*, (F.) *Colique Néphrétique*. Acute pains, which accompany nephritis, and especially calculous nephritis, or the passage of a calculus into the ureter.

COLICA PICTONUM, Colica metallica—c. *Pictonum*, Colica metallica—c. *Pituitosa*, Colica convulsiva—c. *Plethorica*, Colica inflammatoria—c. *Plumbariorum*, Colica metallica—c. *Pulsatilis*, Colic, inflammatory—c. *Rachialgia*, Colica metallica—c. *Renalis*, Colica nephretica—c. *Saturnina*, Colica metallica.

COLICA SCORTORUM, *Colic of Prostitutes*. A form of colic said, by Dr. Martin Hassing, of Copenhagen, to have been frequently observed by him amongst that unfortunate class of beings. It may well be doubted whether any special affection of the kind appertains to them.

COLICA SICCA, C. metallica—c. *Spasmodica*, Colica convulsiva, Ileus.

COLICA STERCORÆA, *Colica Stipata*, *Stercoraceous Colic*, (F.) *Colique stercorale*. A species of colic, ascribed to the retention of fecal matters in the intestines. The retention is itself, however, an effect, that may be caused in the same manner as the colic pains themselves.

COLICA STIPATA, Colica stercoræa.

COLICA UTERINA, *Hystero-colica*, *Uterine*

Colic, (F.) *Colique utérine*. Pain seated in the uterus, sometimes called *Hystericalgia*.

COLICA VENTRICULI, Cardialgia.

COLICA VERMINOSA, *Helminthocolica*, *Worm Colic*, (F.) *Colique vermineuse*. Abdominal pain, occasioned by the presence of worms in the intestines.

COLICODYNIA, *Colica*.

COLICOPLEGIA, *Colic*, metallic.

COLIMAC'ON, *Helix pomatia*, *Limax*.

COLIQUE, *Colic*—*c. des Barbouilleurs*, *Colica metallica*—*c. Hépatique*, *Colica hepatica*—*c. d'Indigestion*, *Colica crapulosa*—*c. Métallique*, *Colica metallica*—*c. de Miserere*, *Ileus*—*c. de Miséricorde*, *Ileus*—*c. des Peintres*, *Colica metallica*—*c. de Plomb*, *Colica metallica*—*c. de Poitou*, *Colica metallica*—*c. Saturnine*, *Colica metallica*—*c. Sèche*, *Colica metallica*—*c. Végétale*, *Colica metallica*—*c. Ventreuse*, *Colica flatulenta*.

COLIQUES, Pains, (after.)

COLITIS, from *κωλον*, 'the colon,' and *itis*, denoting inflammation; *Coloni'tis*, *Enteri'tis colica*, (F.) *Inflammation du colon*. Inflammation of the peritoneal or mucous membrane of the colon. The former is termed *Serocoli'tis*, and *Encocoli'tis*; the latter, *Endocolitis* and *Dysentery*.

COLIX, *Trochiscus*.

COLLA PISCII, *Ichthyocolla*.

COLLAPSE, *Collapsus*—*c. of the Lung*, see *Atelectasis*—*c. Pulmonary*, see *Atelectasis*.

COLLAP'SUS, *Collapse*, *Conciden'tia*; from *col* or *enm*, 'with,' and *labor*, *lupus*, 'to fall.' (F.) *Affaïsement*. A complete prostration of strength, either at the commencement or in the progress of a disease.

COLLAR-BONE, *Clavicla*.

COLLARD, *Dracontium foetidum*—*c. Cow*, *Dracontium foetidum*—*c. Polecat*, *Dracontium foetidum*.

COLLARIUM SALINUM, *Haloderæum*.

COLLATERAL, *Collateralis*, (F.) *Collate'ral*, from *col*, 'with,' and *latus*, 'side.' That which accompanies or proceeds by the side of another.

COLLATERAL ARTERIES OF THE ARM, *Arte'riæ Collaterales Bra'chii*, (F.) *Artères collatérales du bras*. They are given off by the brachial, and are distinguished into, 1. The collateral—superior or external (*Grandes musculaires du bras*—Ch.) which arise from the inner part of the brachial, and extend as far as the inferior and external part of the arm. 2. The collateral—inferior or internal (*Collatérales du Coudé*—Ch.) which arise from the brachial, near the elbow-joint, and descend towards the upper part of the fore-arm.

The vessels which pass to the fingers and toes are also called *collateral*.

Speaking generally, *collateral branches* are those which follow nearly the same course as the vessel whence they emanate.

COLLATERALIS, *Ischio-cavernosus*.

COLLE-CHAIR, *Sarcocolla*.

COLLE DE POISSON, *Ichthyocolla*.

COLLECTION, *Collectio*, from *colligere*, (*col* and *lego*) 'to gather.' This word is often united to the epithet *purulent*, *serous*, &c., to express a gathering of pus, serum, &c.

COLLESIS, *Agglutination*.

COLLET DES DENTS, see *Tooth*.

COLLETICUS, *Agglutinant*.

COLLEY'S DEPIATORY, see *Depilatory*.

COLLIC'IE. Same etymon as *Collection*. 'Drains to collect and convey away water.' See *GOUTTIÈRE*. Union of the ducts passing from the puncta lacrymalia, *Collic'ie puncto'rum lacryma'tium*.

COLLICULI NERVI ETHMOIDALIS, *Cor-*

pora striata—*c. Nervorum opticorum*. *Thalami nervorum opticorum*—*c. Vagine*, *Nymphæ*.

COLLICULUS, see *Optic nerve*.

COLLICULUS CAVEÆ POSTERIORIS VENTRICULORUM LATERALIS, *Hippocampus minor*—*c. Seminalis*, *Gallinaginis caput*.

COLLIER (F.). A collar, from *collum*, 'the neck.' A name given to certain eruptions which surround the neck like a collar.

COLLIER ANODIN, *Necklace*, *anodyne*.

COLLIGAMEN, *Ligament*.

COLLIGATIO, *Syndesmosis*.

COLLINSONIA, *C. Canaden'sis* seu *decussa'ta*, *Horseweed*, *Horsebalm*, *Richweed*, *Richleaf*, *Heal-all*, *Stoneroot*, *Knotroot*, *Knotweed*, *Knobweed*. Order, *Labiatae*. Called after Peter Collinson. This indigenous plant is possessed of diuretic properties, which seem to reside in a volatile oil. Infusion is the best form of administration. The leaves in domestic practice are applied to wounds and bruises.

COLLINSONIA DECUSSATA, *C. Canadensis*.

COLLIQUAMENTUM, from *colliqueo*, (*col* and *liqueo*), 'I melt.' The first rudiments of an embryo.—Harvey.

COLLIQUA'TION, *Colliquatio*, *Eliquatio*, *Syntex'is*, *Etetex'is*, *Solutio*, *Dissolutio*. 'The act of melting.' (F.) *Fonte*. The ancients meant, by this term, the progressive diminution of the solid parts, with copious excretion of liquids by one or more passages. They thought, that all the solids melted; and that the liquids, and particularly the blood, lost a portion of their density.

COLLIQUATIVE, *Colliquativus*, *Colliquescens*, from *colliquescere*, (*col* and *liqueo*), 'to grow liquid.' (F.) *Colliquatif*. An epithet given to various discharges, which produce rapid exhaustion. Hence we say, *Colliquative sweats*, *Colliquative diarrhœa*, &c.

COLLIQUESCENS, *Colliquative*.

COLLISIO, *Contusion*.

COLLIX, *Trochiscus*.

COLLODES, *Glutinosus*.

COLLODION, *Collo'dium*, *Ethe'real Solution of Gun-cotton*, *Maynard's Adhesive Liquid*; from *κόλλα*, 'glue.' A solution obtained by dissolving *gun-cotton*, *Pyrox'ylin*, (F.) *Fulmicoton*, *Coton-poudre*, *Poudre-coton*, *Coton fulminant*, *Coton azotique*, in a mixture of rectified ether and alcohol, in the proportion of about 16 parts of the former to 1 of the latter. When applied to a part, the ether evaporates, and the solid adhesive material is left, which contracts. Hence it is used in cases of wounds, to keep their edges together. It forms, also, a coating, and has been applied in abrasions, and in cases of burns. In various chronic cutaneous diseases, it has been applied with advantage; and has been employed to give a coating to pills, which it deprives of their taste, without interfering with their action.

Collodion is in the last edition of the Ph. U. S. (1851.)

COLLODION, CANTHARIDAL, *Collo'dium vesicans* seu *cantharida'le*. Prepared by exhausting, by displacement, a pound of coarsely-powdered *cantharides* with a pound of sulphuric ether, and three ounces of acetic ether. In two ounces of this saturated solution of cantharides, twenty-five grains of *gun-cotton* are dissolved. By painting the surface with a little of this solution, vesication is produced, as with the plaster of cantharides.

COLLOIDIUM, *Collodion*—*c. Cantharidale*, *Collodion*, *cantharidal*—*c. Vesicans*, *Collodion*, *cantharidal*.

COLLOID, *Colloï'des*, from *κόλλα*, 'glue,' and *ειδος*, 'resemblance;' *Cancer alveola'ris*, *Ca'ci-no'ma alveola'rè*, (F.) *Cancer alvéolaire* ou *Col-*

loide ou Gélatiniforme, Cancer alvéolaire ou aréolaire ou pulvace ou gommeux, gelatinous cancer. An epithet applied to a product of morbid secretion, resembling glue, or particles of jelly inlaid in a regular alveolar bed. The three species of cancer or carcinoma are,—*Encephaloid, Scirrhus, and Colloid.* See *Cancer*.

COLLO'MA, from *κόλλα*, 'glue.' A name proposed by Dr. Walshe for the gelatiniform matter, which is of common occurrence in cysts. It is transparent, amorphous, and devoid of vessels and nerves.

COLLONE'MA, from *κόλλα*, 'glue.' A peculiar gelatinous tumour, consisting of a remarkably soft gelatiniform tissue, which trembles on being touched.—J. Müller.

COLLOSTRUM, Colostrum.

COLLOURION, Collyrium.

COLLUM, *Cervix, Trache'os, Auchen, Neck*, (Prov.) *Hause, Hose*, (F.) *Col, Cou*. The part of the body situate between the head and chest. Also, a part of an organ resembling the neck, as in the following cases.

COLLUM ASTRAG'ALI, *Cervix Astrag'ali, Neck of the Astragalus*, (F.) *Col de l'astragale*. A depression, which separates the anterior extremity of the astragalus from its body.

COLLUM COSTA'RUM, *Cervix Costa'rum, Neck of the Ribs*, (F.) *Col des Côtes*. The narrow part of the ribs, between the head and tubercle.

COLLUM DENT'IVM, *Cervix Dentium, Neck of the Teeth*, (F.) *Col ou Collet des Dents*. The part of the teeth between the corona and fang, which is embraced by the gum.

COLLUM FEM'ORIS, *Cervix Fem'oris, Neck of the Thigh-bone*, (F.) *Col du Fémur*. The long, narrow, and oblique portion of the os femoris, which separates the head from the two trochanters.

COLLUM FIB'ULÆ, *Cervix Fib'ulæ, Neck of the Fibula*, (F.) *Col du Péron*. A slight narrowness seated below the head or upper extremity of the fibula.

COLLUM GLANDIS, see *Glans*.

COLLUM HU'MERI, *Cervix Hu'meri, Neck of the Humerus*. A circular, depressed portion, which separates the head of the os humeri from its two tuberosities. Some surgeons place the neck below the tuberosities, no precise line of demarcation indicating its extent.

COLLUM MANDIB'ULÆ seu MAXILLÆ INFERIO'RI, *Cervix Mandib'ulæ seu Maxillæ Inferio'ris, Neck of the lower jaw*, (F.) *Col de l'os maxillaire inférieure*. A depression observable on each side of the lower jaw, immediately below the condyles.

COLLUM OBSTIPUM, *Torticollis*.

COLLUM OSSIS MAGNI seu CAPITATI, *Cervix ossis magni seu capitati, Neck of the Os Magnum*, (F.) *Col du grand Os*. A circular depression beneath the head of this bone of the carpus.

COLLUM PEDIS, *Instep*.

COLLUM RADII, *Cervix Radii, Neck of the Radius*, (F.) *Col du Radius*. A narrow portion of the radius, which supports the head of the bone.

COLLUM SCAP'ULÆ, *Cervix Scap'ulæ, Neck of the Scap'ula*, (F.) *Col de l'Omoplate*. A narrow portion, observable below the glenoid cavity of the scapula, which seems to separate it, as it were, from the rest of the bone.

COLLUM UTERI, *Cervix Uteri, Jugulum Uteri, Neck of the Uterus*, (F.) *Col de la Matrice, Col utérin*. A narrow, cylindrical, and flattened portion of the uterus, which terminates inferiorly, and opens into the vagina by the *Os Uteri* or *Os Tincæ*. This neck is perceptible on examination per vaginam, until near the end of utero-gesta-

tion. As the uterus enlarges, however, it becomes shorter and wider, so that a manifest difference exists between its condition at seven and nine months.

COLLUM VESICÆ, *Cervix Vesicæ, Neck of the Bladder*, (F.) *Col de la Vessie*. The anterior part of the base of the bladder, which resembles the neck of a bottle, and is continuous with the urethra.

COLLURIUM, Collyrium.

COLLUTIO, Gargarism.

COLLUTO'RIVM, (F.) *Collutoire*, from *colluere*, 'to wash out.' A mouth wash.

COLLUTORIVM ADSTRINGENS, *Mel boracis*.

COLLUVIES, from *colluo*, 'I cleanse.' Filth, excrement. Discharge from an old ulcer.

COLLUVIES GASTRICA, *Embarras gastrique*.

COLLYR'IVM, *Collu'rium, Collu'rium*, (F.) *Collyre*, from *κωλυω*, 'I check,' and *ρεω*, 'I flow,' or from *κόλλα*, 'glue,' and *οὐρα*, 'tail' [?], (F.) *Collyre*. The ancients designated, by this term, a solid medicine, of a long, cylindrical shape; proper to be introduced into the vagina, anus, or nostril. They are said to have given it this name because it was shaped like a *Rat's Tail*, and because there entered into its composition powders and glutinous matters.—Celsus, Oribasius, Scribonius Largus. At the present day, Collyrium means an application to the eye. Some are *pulverulent* and *dry*, *Ophthalmempas'mata*, *Collyria sicca*, but the greatest part are *liquid*, *Hygrocollyria*; and receive different epithets, as *stringent*, *emollient*, &c. The term is now little more than synonymous with *Eye-water*. Collyria are generally extemporaneous formulae.

COLLYRIUM BATEANVM, see *Lotion*, camphorated, of Bates—c. *Sicum*, see *Collyrium*—c. *Sicum Alexandrinum*, see *Alexandrine*.

COLOBO'MA, *κολοβωμα*, 'any thing truncated or shortened.' A mutilated or maimed organ.

COLOBO'MA IRIDIS, *Irido-coloboma*. A congenital peculiarity of the iris, consisting in a fissure of its lower portion, and a consequent prolongation of the iris to the margin of the cornea. See *Iridorrhagias*.

COLOBO'SIS, *κολοβωσις*, The act of curtailing or mutilating; mutilation.

COLOCHOL'O'SIS, from *κωλον*, 'the colon,' and *χολη*, 'bile.' Bilious dysentery, *Dysenteria biliosa*.

COLOCYNTH, *Cucumis colocynth*.

COLOMBINE, *COMMON*, *Aquilegia vulgaris*.

COLOMBA, *Calumba*.

COLON, *C. Cæcum, Monen'terum, Colum, Intestinum majus, I. cellulatum, I. crassum et plenum, I. grande, I. laxum, Physce, φυσικη*. That portion of the large intestines which extends from the cæcum to the rectum. It is said to have been so called from *κωλος*, 'hollow,' or from *κωλυω*, 'I arrest,' because the excrements are arrested, for a considerable time, in its sacs,—*cel'ulæ seu haustra seu loculamenta coli*. The colon is usually divided into four portions. 1. The *right lumbar* or *ascending*, *Colon dextrum*, situate in the right lumbar region, and commencing at the cæcum. 2. *Transverse colon*, *Colon transversum*, *transverse arch of the colon*, the portion of the colon which crosses from the right to the left side, at the upper part of the abdomen. 3. The *left lumbar* or *descending colon*, *Colon sinistrum*, extending from the left part of the transverse arch, opposite the outer portion of the left kidney, to the corresponding iliac fossa. 4. The *Iliac colon* or *Sigmoid flexure of the colon*, (F.) *Colon iliaque* ou *S. du colon*, the portion of the intestine which makes a double curvature in the left iliac fossa, and ends in the rectum.

The muscular fibres, as in the cæcum, are in three flat stripes, *Tænia seu Fascia Ligamentosa Coli*, *Tænia Valsalva seu Ligamentum coli*.

COLON, Mombrium—c. Inflammation of the, Colitis.

COLON, TORPOR OF THE. A disease in which the muscular coat of the colon acts with deficient energy; giving occasion to distension of the intestine, which, by pressing upon the other organs, may interfere so much with their functions, as to lead to distressing gastric, cardiac and other disorders.

COLONITIS, Colitis, Dysentery.

COLONNE, Columna—c. *Vertébrale*, Vertebral column.

COLONNES CHARNUES, Carneæ Columnæ—c. *Charnues du Cœur*, Columnæ carneæ.

COLOPHANE, Colophonia.

COLOPHANY, Colophonia.

COLOPHONIA, so called from Colophon, a city of Ionia; *Phrygæ, Fricæ, Pix Græca, Resina nigra, Collophany, Colophony, Black Resin, Pitch, Becca Rosin*, (F.) *Colophane, Colophane, Arcanson, Brai sec*. The black resin which remains in the retort, after the distillation, by means of a strong fire, of common turpentine. It is used like the turpentine in general, and in some pharmaceutical preparations.

COLOPHONY, Colophonia.

COLOQUINTE, Cucumis colocynthis.

COLOQUINTIDA, Cucumis colocynthis.

COLOR VIRGINEUS PALLIDUS, Chlorosis—c. *Virginum foedus, Chlorosis*.

COLO-RECTITIS, Dysentery.

COLOSTRA, Colostrum.

COLOSTRATIO, Colostration.

COLOSTRATION, *Colostratio*. Disease in new-born children, attributable to the colostrum.

COLOSTRUM, *Colostrum, Collustrum, Colustrum, Trophæalis, Protogala, Neogala, Primum Puerperæ lac, Pytia, Pyetia, Green Milk, (Beastings, Beestings, Biestings or Brustings, Brust, (Sc.) Beist, Beistyn, (Prov.) Fore-milk, Poad-milk, Pored milk, Pound-milk, in the cow, &c.)*, from *kolos*, 'food' [?], or from *kollos*, 'glue', on account of the viscosity of the first milk. (F.) *Bézon*. The first milk after accouchement. It contains more serum and butter, and less casein than common milk, and seems to possess an evacuant property, which renders it fit to aid in the expulsion of the meconium. *Colostrum* formerly meant an emulsion prepared of turpentine and yolk of egg.

COLOUR-BLINDNESS, Achromatopsia.

COLOURS, ACCIDENTAL, *Op'posite colours, Complementary or Harmonic colours*, (F.) *Couleurs Complémentaires*. If the eye has been for some time regarding a particular colour, the retina becomes insensible to this colour; and if, afterwards, it be turned to a sheet of white paper, the paper will not seem to be white, but will be of the colour that arises from the union of all the rays of the solar spectrum, except the one to which the retina has become insensible. Thus, if the eye be directed for some time to a red wafer, the sheet of paper will seem to be of a bluish-green, in a circular spot of the same dimensions as the wafer. This bluish-green image is called an *ocular spectrum*, because it is impressed upon the eye and may be retained for a short time; and the colour bluish-green, is said to be the *accidental colour* of the red. If this experiment be made with wafers of different colours, other accidental colours will be observed, varying with the colour of the wafer employed, as in the following table:—

Colour of the
Wafer.

Accidental colour, or colour of the
ocular spectrum.

Red	Bluish green.
Orange	Blue.
Yellow	Indigo.
Green	Violet with a little red.
Blue	Orange red.
Indigo	Orange yellow.
Violet	Yellow green.
Black	White.
White	Black.

If all the colours of the spectrum be ranged in a circle, in the proportions they hold in the spectrum itself, the accidental colour of any particular colour will be found directly opposite. Hence, the two colours have been termed *opposite colours*. It will follow from what has been said, that if the primary colour, or that to which the eye has been first directed, be added to the accidental colour, the result must be the same impression as that produced by the union of all the rays of the spectrum—white light. The accidental colour, in other words, is what the primitive colour requires to make it white light. The primitive and accidental colours are, therefore, *complements* of each other; and hence accidental colours have also been called *complementary colours*. They have likewise been termed *harmonie*, because the primitive and its accidental colour *harmonize* with each other in painting.

COLPAL'GIA, from *κολπος*, 'vagina,' and *algos*, 'pain.' Pain in the vagina.

COLPATRE'SIA, *Elytratre'sia, Aneylocol'pus, Atretely'tria*, from *κολπος*, 'vagina,' and *αρετος*, 'without opening.' Imperforation of the vagina.

COLPEMPHRAX'IS, from *κολπος*, 'vagina,' and *εμφραξις*, 'obstruction.' Obstruction of the vagina by foreign bodies.

COLPEURYN'TER, Speculum vaginae. The same term is given by Dr. Braun to an instrument for dilating the os uteri, and upper part of the vagina. The operation is called *Colpeury'sis*.

COLPITIS, Colposis.

COLPOC'ACE, *Edæotitis gangræno'sa, Gangræna genitalium et vaginæ*, from *κολπος*, 'vagina,' and *κακος*, 'evil.' Putrescence or gangrene of the vagina and labia.

COLPOCACE INFANT'ILIS, *Edæotitis gangræno'sa puella'rum, Noma pudenda'rum*. Gangrene or putrescence of the vagina and genitals in young children.

COLPOCACE PUERPERA'RUM, *Edæotitis Gangræno'sa puerperarum, Tocolomycol'itis maligna vagina'lis* (Ritgen). Sloughing of the vagina and genitals in puerperal women.

COLPOCELE, Elytrocele.

COLPOCYSTOTOM'IA, from *κολπος*, 'vagina,' *κυστις*, 'bladder,' and *τομή*, 'incision.' *Scelus vagi'no-vesical'is*. Lithotomy through the vagina.

COLPODESMORRAPH'IA, from *κολπος*, 'vagina,' *δεσμος*, 'ligament,' and *ραφή*, 'suture.' The removal of a part of the mucous membrane of the vagina for the radical cure of prolapsus vaginae et uteri.

COLPÆDEMA, Elytroedema.

COLPOPHLEGHMENTITIS, Encolpitis.

COLPOPTOSIS, Prolapsus vaginae.

COLPORRHAG'IA, *Elytrorrhag'ia*, from *κολπος*, 'vagina,' and *ρρηνυμι*, 'I break forth.' Discharge of blood from the vagina.

COLPORRHAPHY, Elytrorrhaphy.

COLPORRHEX'IS, *Ruptura vaginæ*, from *κολπος*, 'vagina,' and *ρηξις*, 'rupture.' Rupture of the vagina. Also, colporrhagia.

COLPORRHEA, Leucorrhœa.

COLPOS, see Mamma: Sinus, Vagina.

COLPOS'IS. Inflammation of the vagina.

Synonymous with *Elytritis*, *Elytritis*, *Coleitis*, *Cupitis*. See *Leucorrhœa*.

COLPOSTEGNOSIS, *Coleostegno'sis*, from *κολπος*, 'vagina,' and *στεγνω*, 'I close.' Atresia, or obliteration of the vagina.

COLPOSTENOCHORIA, *Coleostegnosis*.

COLPOSYNIZESIS, *Coleostegnosis*.

COLPOTOMY, *Colpotomia*, from *κολπος*, 'vagina,' and *τομή*, 'incision.' An incision of the vagina in parturition.

COLT'S FOOT, *Asarum Canadense*, Tussilago.

COLUBER BERUS. The systematic name of the viper, *Vipera*, (F.) *Ouleuvre*, *Vipère*. Class, Reptilia, Ord. Ophidia. A poisonous reptile—the poison lying in small sacs near its teeth. The flesh is innocent, and has been often taken in scrofula, and in cutaneous disorders in general, but it is inefficacious.

COLUBRINA, *Polygonum bistorta*—c. *Dracontia*, *Arum dracuncul*—c. *Lusitanica*, *Euphorbia capitata*—c. *Virginiana*, *Aristolochia serpentaria*.

COLUM, *Colon*.

COLUMBINE, *Aquilegia vulgaris*—c. *Wild*, *Aquilegia Canadensis*.

COLUMBO, *Calumba*—c. *American*, see *Calumba*—c. *Marietta*, see *Calumba*.

CUMELLA, *Pillar*, *Uvula*—c. *Cochleæ*, *Modiolus*.

COLUMNA, *Column*, (F.) *Colonne*. Anatomists use this word for parts which resemble a column or pillar; hence for the penis.

COLUMNA ADSTANS INGUINIBUS, *Penis*—c. *Dorsi*, *Vertebral column*—c. *Fornicis*, see *Fornix*.

COLUMNA NASI. The cartilaginous part of the septum of the nostrils. See *Nares*.

COLUMNA ORIS, *Uvula*—c. *Spinalis*, *Vertebral column*—c. *Virginitalis*, *Hymen*—c. *Foraminis ovalis*, see *Ovalis fossa*—c. *Valvulæ Viuessenii*, see *Valvula Viuessenii*—c. *Vertebris*, *Vertebral column*.

COLUMNÆ BERTINI, *Columns of Bertin*. Processes of the cortical substance of the kidney, which extend between the pyramids of Malpighi, as far as the hilus.

COLUMNÆ CARNÆÆ, *C. Cordis*, *Lacer'ti* seu *Lacer'tuli* seu *Funes* seu *Fasciculi teretes Cordis*, *Trabes* seu *Trabeculæ Cordis*, (F.) *Columnæ charnues du cœur*. Small, fleshy columns, which project, more or less, into the auricles and ventricles of the heart, whose use appears to be to prevent too great dilatation of those cavities. A few of these *columnæ*—see *Musculi papillares*--are attached by one extremity to the walls of the heart, and, by the other, give insertion to chordæ tendinæ. See *Carneous columns*.

COLUMNÆ CARNÆÆ of the Rectum, see *Rectum*—c. *Papillares*, see *Columnæ Carnææ*.

COLUMNÆ LONGIFOLIA, *Bahel*.

COLUMNS OF BERTIN, *Columnæ Bertini*.

COLUMNS, POSTERIOR MEDIAN, OF THE MEDULLA OBLONGATA, *Funiculi graciles*—c. of *Morgagni* or of the *Rectum*, see *Rectum*—c. of the *Spinal Marrow*, see *Vertebral Nerves*.

COLUS JOVIS, *Salvia sclarea*.

COLUSTRUM, *Colostrum*.

COLUTEA, *C. Arborescens* seu *hirsuta*, *Senna Germanica*, *Bladder Senna*, (F.) *Baughendier*, *Faux Séné*, *Séné indigène*. Ord. Leguminosæ. Sex. Syst. Diadelphia Decandria. The leaves are slightly purgative, and are often mixed with those of the cassia senna.

COLUVRINE DE VIRGINIE, *Aristolochia serpentaria*.

COLYMBADES, *Pickled Olives*. These, when bruised and applied to a burnt part, were supposed to be able to prevent vesication.—*Dioscorides*.

COLYMBIFERA MINOR, *Mentha crispa*.

COMA, *Semisomnis*, *Semisopitus*, *Semisoporus*, *Subeth*, (Arab). A profound state of sleep, from which it is extremely difficult to rouse the individual. It is a symptom which occurs in many diseases. Two varieties are distinguished, 1. *Coma vigil* seu *agrypnodes*, *Pervigilium*, *Vigilia nimia*, *Typhoidia*, *Veterinus*, *Veterinitas*, *Agrypnocoma*, *Carus lethargus vigil*, *Typhomania*, which is accompanied with delirium. The patient has his eyes closed, but opens them when called; and closes them again immediately. This state is accompanied with considerable restlessness. 2. *Coma Somnolentum*, *C. Comatodes*;—in which the patient speaks when roused, but remains silent and immovable in the intervals. Coma is a deeper sleep than *sopor*, but less so than *lethargy* and *carus*.

COMA, *Capillus*—c. *Agrypnodes*, see *Coma*—c. *Apoplexia*, *Apoplexy*—c. *Cæsarea*, *Plica*—c. *Comatodes*, see *Coma*—c. *Somnolentum*, *Somnolency*, see *Coma*—c. *Vigil*, see *Coma*.

COMACON, *Myristica moschata*.

COMAN'DRA UMBELLATA, *Bastard Tondax*; indigenous; Order, *Santalacæ*; flowering in May and June; has been used in fevers by some of the Indian tribes.

COMAROS, *Arbutus unedo*.

COMARUM PALUSTRE, *Potentilla palustris*, *Marsh Cinquefoil*. An indigenous plant, Family, *Rosacæ*, which flowers in June. It is possessed of astringent virtues.

COMATA, from *κωπα*. Diseases characterized by diminution of the powers of voluntary motion, with sleep or impaired state of the senses.—*Cullen*.

COMATEUX, *Comatose*.

COMATODES, *Comatose*.

COMATOSE, *Comatodes*, *Caroticus*, *Caroleus*, (F.) *Comateux*. Relating to or resembling coma:—as *comatose sleep*, *comatose fever*, &c.

COMBUSTIBILITY, *PRETERNATURAL*, *Combustion*, human.

COMBUSTIO, *Burn*—c. *Spontanea*, *Combustion*, human.

COMBUS'TION, HUMAN, *Spontaneous Combustion* or *Preternatural Combustibility*, *Catacan'sis*, *Tachencan'sis*, *Incedidum spontaneum*, *Combustio spontanea*, *Autempressus*, *Idiopcanc'sis*, *Idiempres'sis*, *Catacan'sis ebriosa*, (F.) *Combustion humaine*, *C. humaine spontanée*; from *comburare*, *combustum*, (*com*, and *urere*, *ustum*) 'to burn.' These terms have been applied to the extraordinary phenomenon of a rapid destruction of the human body, by being reduced to ashes either spontaneously or by the contact of an ignited substance. It is said to have occurred in the aged, and in those that were fat and hard drinkers. In such, Dr. Traill has found a considerable quantity of oil in the serum of the blood. Vicq d'Azyr, Lair, and Dupuytren think it necessary, that the body should be placed in contact with an ignited substance. Le Cat, Kopp, and Marc are of opinion that this is not necessary. The former appears to be the more probable view.

COMBUSTURA, *Burn*.

COMEDONES, see *Acene*.

COMES ARCHIATRORUM, see *Exarchiater*.

COMESTIBLE (F.), *Edulis*, from *comedere*, 'to eat.' *Eatable*, (F.) *Édible*. Esculent. When this word is used substantively, in French as in English, it means solid food.

COMEZ. Half a drop.—*Ruland*.

COMFREY, *Symphytum*—c. *Spotted*, *Pulmonaria*—c. *Wild*, *Cynoglossum Virginicum*.

COMISTE, *Aliment*.

COMMANDUCATIO, *Mastication*.

COMMEMORATIVE, *Commemorativus*, (F.) *Commemoratif*, from *commemorare*, (*com* and

memor.) 'to cause to remember.' That which recalls, or rather which is recalled.

COMMEMORATIVE SIGNS, (F.) *Signes commémoratifs*, are those deduced from circumstances which have gone before; or rather, according to others, those which point out the preceding condition of the patient. They are also called *Anamnesis signs*.

COMMENSUM, Symmetry.

COMMERCIUM, Sympathy.

COMM. Gummi.

COM'MINUTED, *Comminutus*, from *comminere*, (com and *minuo*), 'to break to pieces.' (F.) *Comminutif*. A comminuted fracture is one in which the bone is broken into a number of pieces.

COMMINUTION, *Comminutio*, *Thrypsis*, *Leio'sis*. Same etymon. Fracture of a bone into a number of pieces.

COMMISSURA, Articulation.

COMMISSURA ANTERIOR CEREBRI, Commissure, anterior, of the Brain.

COMMISSURA BREVIS. A lobule or prominence of the inferior vermiform process of the cerebellum, situate in the incisura posterior, below the horizontal fissure.

COMMISSURA LABIORUM, Prostomia—c. Magna cerebri, Corpus callosum—c. Maxima cerebri, Corpus callosum—c. Nervea, Syndesmosis—c. Ossium carnea, Syssarcosis.

COMMISSURA POSTERIOR CEREBRI, Commissure, posterior, of the Brain.

COMMISSURA SIMPLEX. A small lobule or prominence of the superior vermiform process, near the incisura posterior of the cerebellum.

COMMISSURAL; same etymon as commissure. Of or belonging to a commissure.

COM'MISSURE, *Commissura*, *Compages*, *Compaginatio*, *Syn'physis*, *Syn'bolē*; from *com-mitto*, (com and *mittere*, *missum*), 'to join together.' A point of union between two parts: thus, the commissures of the eyelids, lips, &c., are the angles, which they form at the place of union. See *Fibres*, converging.

COM'MISSURE, ANTE'RIOR, OF THE BRAIN, *Commissura ante'rior cer'ebri*, (F.) *Commissure antérieure du cerveau*. A small medullary fasciculus, situate transversely at the anterior part of the third ventricle, and uniting the two hemispheres.

COMMISSURE, GREAT, OF THE BRAIN, *Commissura Magna Cer'ebri*, (F.) *Grande commissure du cerveau*, which unites the two hemispheres for some extent, is the *Corpus callosum*.

COMMISSURE, MIDDLE, OF THE BRAIN. A layer of gray substance uniting the thalami optici.

COMMISSURE, OBLIQUE OR INTERCEREBRAL, see *Valvula Vieussensii*—c. Optic, see *Chiasmus*.

COMMISSURE, POSTERIOR OF THE BRAIN. *Commissura poste'rior cer'ebri*. A medullary band, situate at the posterior part of the third or middle ventricle.

COMMISSURE OF THE UVEA, Ciliary ligament.

COMMISSURE DE LA CHOROÏDE, Ciliary ligament.

COMMO'SIS, from *κομω*, 'I adorn.' The art of concealing natural deformities, as by painting the face. See, also, *Protopis*.

COMMO'TICĒ, in the older writers, meant the art of communicating factitious beauty to a person. Painting the face; *Comoporia*.

COMMOTIO, Motion, Concussion—c. Cerebri, Concussion of the brain.

COMMOTION, Concussion—c. du Cerveau, Concussion of the brain.

COMMUNICABLE, see *Contagious*.

COMMUNICANS, from *communis*, (com and *munus*, 'an office'), 'common.' *Conjun'gens*. That which communicates or establishes a communica-

tion. Communicant. There are two *Arteries Communicantes*, both within the cranium;—the one anterior, very short, and extending transversely from one anterior p cerebral artery to the other,—the other posterior, called also *Communicans Willis'ii*, which passes from the internal carotid to the posterior cerebral artery. It is a branch of the basiliary.

COMMUNICANS NONI. A long slender nervous branch, formed by filaments from the first, second and third cervical nerves, which descends upon the outer side of the internal jugular vein, and forms a loop with the descendens noni over the sheath of the carotids.

COMMUNICANS PERONEI, see *Communicans poplitei*.

COMMUNICANS POPLITEI, *C. tibialis (nervus)*. A large nerve, which arises from the popliteal, and, at a variable distance below the articulation of the knee, receives the *communicans peronei* from the peroneal nerve,—the two forming the *external saphenous nerve*.

COMMUNICANS TIBIALIS, *Communicans poplitei*.

COMPACT, *Compactus*, from *com* and *pangere*, *pacum*, 'to strike, to fix.' Solid, close. (F.) *Compacte*. The term *Compact Tissue* is given to the hardest and closest parts of bone.

COMPAGES, Articulation, Commissure—c. Ossium per Lineam Simplicem, Harmony—c. Vertebrarum, Vertebral column.

COMPAGINATIO, Commissure.

COMPAS DÉPAISSEUR, see *Pelvimeter*.

COMPASS PLANT, *Silphium laciniatum*.

COMPASSIO, Sympathy.

COMPEBA, Piper cubeba.

COMPENSATION, *Balancement*.

COMPEPER, Piper cubeba.

COMPER'NIS; from *com* and *perna*, 'a gammon of bacon with the leg on.' One who has his knees turned inwards. A case of distortion of the legs.

COMPETENTIA MEMBRORUM OMNIUM, Symmetry.

COMPLAINT, Disease.

COMPLAINT, FAMILY, see *Hereditary*.

COMPLEMENTAL AIR, see *Respiration*.

COMPLEMENTARY AIR, see *Respiration*.

COMPLETIO, Plethora.

COMPLEX, *Complexus*, from *com* 'with,' and *plectere*, *plexus*, 'to twist.' Embracing several distinct things. Chaussier uses this term, in his anatomical descriptions, for *complicated*.

COMPLEXIO, Coition, Complexio, Confusio, Temperament.

COMPLEXION, *Complexio*. Often employed, in English, for the colour of the face, as "He has a good complexion,"—a "sallow complexion," &c. It formerly had a more extensive signification, and still has in France. It signifies the aggregate of physical characters presented by any individual, considered with respect to his external arrangement or condition. It means more than *constitution*, for which it is used synonymously in many cases; and differs from *temperament*, which is less the external condition of the body than the state or disposition of the organs in health.—H. Cloquet.

COMPLEXUS, Complex.

COMPLEXUS MINOR, *Mastoidæus lateralis*, *Trachelo-mastoidæus*, (F.) *Trachélo-mastoidien*, *Muscle petit Complexus*. It arises from the transverse processes of the last four cervical vertebrae, and is inserted into the mastoid process.

COMPLEXUS MUSCULUS, *Biven'ter Cervicis*, *Complexus Major*, *Dorso-trachélon-occipital*, *Trachélo-occipital* (Ch.), (F.) *Muscle grand complexus*. A muscle, situate at the hind part of the neck, where it extends from the interval that separates the two prominent ridges on the posterior surface

of the os occipitis to the transverse and articular processes of the last six cervical vertebra, as well as to those of the first five dorsal. It serves to straighten, incline, and turn the head.

COMPLICATION, *Complicatio*, from *com* 'with,' and *placere*, 'to fold.' In medicine, it means the presence of several diseases, *morbis complicati seu perplexi*, or of several adventitious circumstances foreign to the primary disease.

COMPOSIT'ION, *Compositio*, from *componere*, (*com* and *ponere*, *positum*, 'to place') 'to place together.' *Agglutis*. The act of composing or compounding, — of medicines for example; *Iamatozyntar'is*. Also, the *Compound*, *Compositum*, or thing compounded. Likewise, a combination.

COMPOSITUM, *Compound*.

COMPOUND. Same etymology; to mix medicines. To mingle different ingredients into one whole. Used adjectively, *compound* signifies the result of the union of several medicinal agents, as "a compound medicine."

COMPREHENSIO, *Catalepsy*.

COMPRESSIO, *Catalepsy*.

COMPRESS, *Compress'a*, *Splenium*, *Spleniola*, *Splenis'eus*, *Plat'uga*, *Penicill'um*, *Penic'ulum*, from *comprimere*, (*com* and *primere*, *pressum*, 'to press') 'to press together.' (F.) *Compresse*. Folded pieces of lint or rag, so contrived as, by the aid of a bandage, to make due pressure upon any part. According to their shape, direction, and use, compresses have been called *long* (F.) *longuettes*, *square* (*carrees*), *triangular*, *prismatic*, *graduated* (*graduées*), *split* (*fendues*), *fenêtrées*, *criblées*, *crois de Malte*, *oblique*, *circular*, *dividing* (*divisives*), *uniting* (*unissantes*), *cribelliform*, &c.

The *Umschlag* or compress of the hydropathists is a cloth, well wetted with cold water, applied to the surface near the supposed seat of disease, securely covered with a dry cloth, and changed as often as it becomes dry. It is sometimes covered with a layer of oiled silk, to prevent evaporation.

COMPRESSEUR DE DUPUYTREN, *Compressor* of Dupuytren — *c. du Nez*, *Compressor nasi* — *c. de la Prostata*, *Compressor prostate*.

COMPRESSIF, *Compressive*.

COMPRESSIO, *Compression*, *Thlipsis*.

COMPRES'SION, *Compression*, *Ener'e'sis*. Same etymology. Pressure; methodical compression. An agent frequently had recourse to in surgery. We compress a limb, affected with *anæmia*, *varices*, *hydrops articuli*, *callous ulcer*, &c. The abdomen is compressed after delivery, after *paracentesis abdominis*, &c. The compression is produced by means of the roller, laced stocking, &c., according to the part, and to the particular case. Moderate pressure aids the contractility of parts, and excites the action of the absorbents; so that large tumours at times disappear after it has been used for some time. A greater degree of pressure occasions, still more, the emaciation of the part, but it is apt to impede the circulation. Pressure is often used to stop or moderate the flow of blood in cases of aneurism or wounds of arteries and veins. In such cases, the compression may be immediate, when applied upon the artery itself, or it may be mediate, when applied through the integuments and soft parts. The French use the term *Compression immédiate*. *1 mode* for that which is exerted perpendicularly to the axis of a vessel, so as to flatten its sides. It is practised with the finger, forceps, tourniquet, compresses, &c.

COMPRES'SION OF THE BRAIN. This may arise either from coagula of blood, a soft tumour, a bony excrescence, a depressed portion of the skull, or the presence of some foreign body. The

effects vary, according as the compression takes place suddenly or gradually. When sudden, the symptoms are of the comatose or apoplectic character. When gradually, mania, convulsions, &c., are more likely to occur. Compression, arising from a depressed portion of skull, requires the use of the trephine.

COMPRES'SIVE, *Compressiv'us*, (F.) *Compressif*. That which compresses. A *compressive bandage* is one that compresses the parts to which it is applied.

COMPRESSOR OR COMPRESSORIUM OF DUPUYTREN, (F.) *Compresseur de Dupuytren*. An instrument for compressing the femoral artery, invented by Dupuytren. It is constructed on the same principles as the tourniquet of J. L. Petit, from which it only differs in this respect: — that, instead of being maintained in its place by a strap, which always compresses more or less the circumference of the limb, the two pads are placed at the extremities of a semicircle of steel, which, by passing from one to the other without touching the parts, limits the pressure to two opposite points of the thigh, and permits the collateral circulation to go on.

COMPRESSOR NARIS, *Rina'us*, *Nasa'lis*, *Transversal'is Nasi*, *Myrtiform'is*, *Dilatator* *nasal'is*, *Constrictor Nasi seu Natrium seu Naris*, *Triangula'ris Nasi*, (F.) *Incillo-nasal*, *Somacillo-nasal*, — (Ch.) *Compresseur du nez*, *Transversal du nez*. A muscle, situate at the sides of the nose; flat and triangular. It arises from the inner part of the fossa canina, and passes to the dorsum of the nose; where it is confounded with that of the opposite side.

COMPRESSOR OR CONSTRICTOR OF NECK. An instrument for compressing the urethra, to obviate incontinence of urine. It consists of a girdle of iron, which surrounds the pelvis, to which is fixed a plate of the same metal, that compresses the urethra in *perinaeo*.

COMPRESSOR PROSTATÆ, *Prostat'icus superior*, *Pubio-prostat'icus*, *Sub-pubio-prostat'icus*, (F.) *Compresseur de la prostate*. A muscle, admitted by ALBINUS, which is formed of the anterior fibres of the *Levator ani*, that embrace the prostate. It is the *Prostat'icus superior* of WINSTON.

COMPRESSOR URETHRÆ. A muscle consisting of two portions — one of which is *transverse* in its direction, and, in consequence of its having been particularly described by Mr. Guthrie, has been called *Guthrie's muscle*. It arises from the ramus of the ischium, and passes inwards to embrace the membranous urethra. The other portion is *perpendicular*, descending from the pubis and passing down to be inserted into the transverse portion of the muscle: — *Pubio-urethra'lis*. This portion has been considered by many to be only the anterior fibres of the levator ani; and having been described by Mr. Wilson, it has been called *Wilson's muscle*.

COMPRESSOR VENE DORSALIS PENIS. A small muscle, distinctly seen in animals, less distinctly in man, which arises from the ramus of the pubis, and ascending in a direction forwards is inserted above the vena dorsalis, joining with its fellow of the opposite side on the mesial line. Its use is supposed to compress the vein in erection. It is sometimes called the *muscle of Houston*, after Dr. Houston of Dublin.

COMPRESSORIUM, *Compressor*.

COMPTONIA ASPLENIFO'LIA, *Liquidambar peruviana* seu *asplenifolia*, *Mycra asplenifolia*, *Sweet Fern*, *Shubby Sweet Fern*, *Sweet bush*, *Fern bush*, *Fern gale*, *Splenaceous bush*, *Meadow fern*, *Astringent root*. An indigenous shrubby plant, which grows in sandy or stony woods from New England to Virginia. It pos-

esses tonic and astringent properties, and is used as a domestic remedy in diarrhoea, &c., in the form of decoction.

COMPUNCTIO, Paracentesis, Puncturing.

CON-ESTHESIS, Cænesthesis.

CONARIUM, Pineal gland.

CONATUS, Effort.

CONCARNATIO, Syssarcosis.

CONCASSER, (F.), from *conquassare*, (con and *quassare*, 'to shake much,') 'to break to pieces,' 'to comminute.' To reduce roots, woods, &c. into small fragments, in order that their active principles may be more readily separated from them.

CONCAUSA, *Concausa*, (con and *causa*,) Synaption. A cause which co-operates with another, in the production of disease.

CONCAVITAS CONCHULARIS CEREBRI, Infundibulum of the brain.

CONCAVUM PEDIS, Sole.

CONCENTRANTIA, from *con* and *centrum*, 'a centre.' A name once given to absorbents of acids.

CONCENTRATIO, *Concentratio*. A word sometimes used, in medical theories, to express an afflux of fluids, or a convergence of sensibility or of vital force, towards an organ. It is applied, also, to the pulsation of arteries, when not easily felt under the finger.

POULS CONCENTRÉ, (F.) A term applied by the French to a pulse of the above character.

CONCEPTACULA SEMINARIA, Vesiculæ seminarie.

CONCEPTACULUM, Uterus, Vessel.

CONCEPTIO, Conception — c. Vitiosa, Pregnancy, extra-uterine.

CONCEPTION, *Conceptio*, *Conceptus*, *Cyesis*, *Syllepsis*, *Androlepsia*, from *concepio*, (con and *capio*,) 'I conceive.' The impregnation of the ovum by the positive contact of the male sperm, whence results a new being. The whole subject of conception is most mysterious. It seems to occur as follows. During the sexual union, the male sperm passes along the uterus and Fallopian tubes: the fimbriated extremities of the latter seize hold of the ovarium; and the sperm in this manner comes in contact with a matured ovum, and impregnates it. The fecundated ovum remains some time in the ovarium, but at length bursts its covering, is laid hold of by the fimbriated extremity of the Fallopian tubes, and passes along the tube into the cavity of the uterus, where it remains for the full period of utero-gestation. Some are of opinion, that the ovum is not impregnated until it has entered the Fallopian tube, or uterus.

CONCEPTION, FALSE, *Falsus Conceptus*, *Spirium geminum*, (F.) *Fausse Conception*, *Faux germe*. An irregular, preternatural conception, the result of which is a mole or some similar production, instead of a properly organized fetus. See Mole.

CONCEPTUS. The first rudiments of the fetus, after conception. Also, conception.

CONCEPTUS FALSUS, Conception, false.

CONCHA, *Conchus*, *κογχη*, 'a concave shell.' A liquid measure, amongst the Athenians, equal to half an ounce. Anatomists apply this term to several hollow parts of the body:—as the *Concha* of the Ear, — *Concha Auris*, *Concha Auricularis*; (F.) *Conque*—the hollow part of the cartilage of the external ear. It has, also, been applied to the genital organs of the female; to the patella, &c.

CONCHA, Patella, Turbinate bone (middle,) Vulva — c. *Auris interna*, Cochlea — c. *Cerebri*, Infundibulum of the brain — c. *Genu*, Patella — c. *Inferior*, Turbinate bone, (inferior) — c. *Labyrinthi*, Cochlea — c. *Morgagniana*, Turbinate

bone, (superior) — c. *Narium superior*, Turbinate bone, (superior.).

CONCHLE NARIUM, Turbinate bones — c. *Veneris*, Turbinate bones.

CONCHARUM ANTIFEBRILE, A febrifuge and sudorific preparation in Bates's *Pharmacopæia*. It was composed of vinegar, mussel-shells, and water of *Carduus benedictus*.

CONCHO-HELIX. A small, fleshy fasciculus, attached to the concha of the ear and helix. It is also called the *small muscle of the helix*.

CONCHUS, *Concha*. The cranium. In the plural, it means the orbital cavities.—Castell.

CONCHUS ORBIT, Orbit.

CONCHYLIA, Turbinate bones.

CONCIDENTIA, Collapse.

CONCILIA CORPORALIA, Coition.

CONCILIUM. A milky plant, referred to by Pliny as aphrodisiac and antiphthical. Adanson considers it to be a *Complanula*.

CONCOTED, *Concoccus*, *Natura'tus*, *Papa'tus*; from *con* and *coquere*, 'to boil.' Brought to maturity; ripe; concocted; digested.

CONCOCTIO, Coction — c. *Tarda*, Dyspepsia.

CONCOCTION, Coction.

CONCOMBRE ORDINAIRE, *Cucumis sativus* — c. *Sauvage*, *Momordica elaterium*.

CONCOMITANS, Concomitant.

CONCOMITANT, *Concomitans*, from *con* and *comitare*, (itself from *comire* — cum and *ire*,) 'to go with.' That which accompanies. A symptom which accompanies others.

CONCREMATIO, Calcination.

CONCREMENTA ZOOHYLICA, see Zoohylica.

CONCREMENTUM, Concretion.

CONCRETIO, Adherence, Concretion — c. *Palpebrarum cum bulbo oculi*, Symbblepharisis.

CONCRETION, *Concretio*, *Concrementum*, from *concrecere*, *concretum*, (con and *crecere*,) 'to condense, thicken, become solid.' *Pecis*, *Synpæcis*, *πέσις, συνπέσις*. The act of becoming thick or solid. It was once used synonymously with adhesion or growing together—as, "*concretion of the toes*." Most commonly, it is applied to extraneous and inorganic bodies, of a solid character, met with in different textures, after certain chronic inflammations; or which make their appearance in the joints or in the reservoirs for excremental fluids. Concretion is, therefore, frequently synonymous with *Calculus*, and is then rendered, in Latin, by the word *Concrementum*. But *Concretion* has a more extensive signification than *Calculus*; thus, accidental ossifications or deposits of phosphate of lime in certain organs, and especially in the liver and lungs, are properly called *osseous concretions*. They could not well be called *osseous calculi*.

CONCRETION, FIBRINOUS, SANGUINEOUS, POLYPIFORM, or POLYPOUS, OF THE HEART, see Polypus — c. *Intestinalis*, *Calculus of the Stomach and Intestines*.

CONCRETIONES ALVINÆ, Calculi of the stomach and intestines.

CONCUBITUS, Coition — c. *Veneris*, Coition.

CONCURSUS, from *concurrere*, (con and *currere*, *cursum*, 'to run,') 'to meet together.' *Syndromè*. The congeries or collection of symptoms, which constitute and distinguish a particular disease.

CONCUS'SION, *Commo'tio*, from *concutio*, *concussus*, (con and *quater*, 'to shake,') 'I shake together.' *Concus'sio*, *Thlasma Concus'sio*, *Commo'tio*, *Anasim'sus*, *Tinagnus*, (F.) *Commo'tio*. In Surgery, it is used for the agitation often communicated to one organ by a fall upon another; as to the brain from a fall on the breech, &c.

In all severe injuries, in sudden encephalic hemorrhage, and in overwhelming emotions, a

concussion or shock is felt to a greater or less extent in the nervous system, which requires the careful attention of the physician.

CONCUSSION OF THE BRAIN, *Commo'tio Cerebri*, *Apoplex'ia nervosa traumatica*, *Encephalosis'mus*, (F.) *Commo'tion du Cerveau*, sometimes gives rise to alarming symptoms, even to abolition of the functions of the brain, yet without any sensible organic disease. Slight concussion of the brain, called *stunning*, consists in vertigo, tinnitus aurium, loss of memory, and stupefaction; all these being temporary. When more severe, there is instant loss of sensation and volition, vomiting, the patient being as if in a sound sleep, but there is no stertorous breathing. Pulse variable, generally more rapid and feeble than in compression; extremities cold. Little can be done here, till reaction has occurred: after this, the case must be treated according to general principles,—by bleeding, blisters, cold applied to the head, &c. After severe concussion, a patient, although apparently well, is not safe till some time after the accident.

CONDENSAN'TIA, from *con* and *densus*, 'dense;' *Incrassantia*. Medicines esteemed proper for inspissating the humours.

CONDENSATIO, Condensation, Stegnosis.

CONDENSATION, *Condensatio*, *Inspissatio*. Increase in density of the blood and other liquids, or of the solids, which enter into the composition of the human body.

CONDENSER, see Alembic.

CONDENSER, LIEBIG'S. A distillatory arrangement, invented by Liebig, in which the tube conveying the vapour is made to pass through another tube, the calibre of which is such as to leave a space between the two, through which a stream of water may be made to run.

CONDIMENT, *Condimentum*, *Artyma*, *Hedys'ma*, *Conditu'ra*, from *condire*, 'to season.' (F.) *Assaisonnement*. Some substances are called, at times, *aliments*, and at others, *condiments*, according as they constitute the basis or the accessory to any dish: such are cream, butter, mushrooms, olives, &c. Others are always *condiments*, as they are only used to improve the savour of food, and contain but little nutritive matter. Such are pepper, salt, cinnamon, &c. Almost all condiments are possessed of stimulant properties.

CONDIT, Confectio.

CONDITUM, same etymon. A pharmaceutical compound of wine, honey, and some aromatics, especially pepper. Also, a confection.

CONDITURA CADAVERUM, Embalming.

CONDOM, *Armour*, (F.) *Baudruche*, *Redingote Anglaise*, *Gant des Dames*, *Calotte d'assurance*, *Peau divine*. The intestinum cæcum of a sheep, soaked for some hours in water, turned inside out, macerated again in weak, alkaline ley, changed every twelve hours, and scraped carefully to abstract the mucous membrane, leaving the peritoneal and muscular coats exposed to the vapour of burning brimstone, and afterwards washed with soap and water. It is then blown up, dried, cut to the length of seven or eight inches, and bordered at the open end with a ribbon. It is drawn over the penis prior to coition, to prevent venereal infection and pregnancy. It received its name from its proposer, Dr. Condom.

CONDUCTIO, Tonic spasm, Convulsion.

CONDUCTION, VIBRATIONS OF, see Sound.

CONDUCTOR, *Director*, *Itinera'rium*, (F.) *Conducteur*; from *con*, 'with,' and *ducere*, *ducere*, 'to lead.' That which conducts. The *Conductor* was an instrument, formerly used in the high operation for the stone, for directing the forceps into the bladder.

CONDUIT, Canal—*c. Auditif externe*, Auditory canal, external—*c. Auditif interne*, Auditory canal, internal—*c. Auriculaire*, Auditory canal, external—*c. Cholédoque*, Choledoch duct—*c. Défèrent*, Deferens—*c. d'Eustache*, Eustachian tube—*c. Gutturale de l'oreille*, Eustachian tube—*c. Labyrinthique*, Auditory canal, internal—*c. Pterygoïdien*, Pterygoid canal—*c. de la Pudeur*, Vagina—*c. Sossorbitaire*, Suborbital canal—*c. Spermatique*, Deferens (vas)—*c. Thoracique*, Thoracic duct—*c. Vidien*, Pterygoid canal.

CONDUITS ADIPEUX, Fatty canals—*c. Aqueux*, see Aqueous—*c. Aëngles de l'urètre*, see Cæcus—*c. Dentaires*, Dental canals—*c. Ejaculateurs*, Ejaculatory ducts—*c. Lacteymaux*, Lachrymal ducts—*c. Lactifères*, Lactiferous vessels—*c. Nourriciers*, Canals, nutritive—*c. Nutriciers*, Canals, nutritive.

CONDYLARTHRO'SIS, from *κονδύλος*, 'a condyle,' and *arthron*, 'a joint.' Articulation by condyles. An elongated head or condyle, received into an elliptical cavity.

CONDYLE, *Con'dylè*, *Con'dylus*, *Capitulum*, from *κονδύλος*, 'a knot, eminence.' An articular eminence, round in one direction, flat in the other. A kind of process, met with more particularly in the ginglymoid joints;—such as the condyles of the occipital, inferior maxillary bone, &c. Some anatomists have applied the term, however, to eminences that are not articular,—as to the lateral tuberosities at the inferior extremity of the *os humeri*, and even to certain depressions,—as to the concave articular surfaces at the upper extremity of the tibia. Chaussier calls the transverse root of the zygomatic process *Condyle of the temporal bone*.

CONDYLI DIGITORUM MANUS, Phalanges of the fingers.

CONDYLIEN, see Condyloid Foramina.

CONDYLIUS, Condyloid.

CONDYLOID, *Condyloïdeus*, *Condyloïdes*, *Condyli'us*, from *κονδύλος*, 'a condyle,' and *αἶος*, 'shape.' Having the shape of a condyle.

CONDYLOID FORAMINA, *Foram'ina Condyloïden*, (F.) *Trous condyloïdiens*, *Fosses condyloïdiennes*. They are distinguished into *anterior* and *posterior*. They are four in number, seated in the occipital bone;—two anterior, and two posterior to the *condyles* or *condyloid processes* of the same bone, and to depressions, which are called *Fossæ Condyloideæ*.

As the word *Condyloïdien* means, 'that which has the shape of a condyle,' it has been judiciously proposed by some French anatomists that *condylien* should be used in preference, in the cases of the foramina and fossæ.

CONDYLOÏDIEN, see Condyloid Foramina.

CONDYLO'MA, *Condyloïsis*, *Con'dylus*, *Veruca carnos'a*, from *κονδύλος*, 'a knot,' 'an eminence.' A soft, fleshy excrescence, of an indolent character, which appears near the orifice of the genital organs and rectum, and occasionally on the fingers and toes. It is a consequence of the syphilitic virus. Such tumours are also called *Dermophymata venerea*.

CONDYLOSIS, Condylosoma.

CONDYLUS, Condylosoma.

CONFLOWER, *Rudbeckia laciniata*—*c. Purple*, *Echinacea purpurea*.

CONFÈINE, see Conium.

CONESSI CORTEX, *Nerium antidysentericum*.

CONFECTIO, *Confection*, from *conficio*, *confectum*, (*con* and *facere*, 'to make,') 'I make up.' *Aliquidus*, (F.) *Confit*, *Confiture*, *Condit*. In general, it means anything made up or preserved with sugar. In the Pharmacopœias of the United States, London, and Dublin, it is used to denote arti-

cles before called electuaries and conserves. *Confectio* or *Confectum* also means Confectionery.

CONFECTIO ALKERMES, Alkermes.

CONFECTIO AMYGDALA'RUM, *C. Amygdalæ*, *Conserua amygdalarum*, Almond Confection, Almond Paste, *Pasta regia* seu *Amygdali'na* seu *Emulsi'va*, Butyrum Amygdalarum Dulcium, (F.) Confection d'Amandes. (Sweet almonds, 3viij; gum acacia, 3j; white sugar, 3iv. Blanch the almonds, and beat into a paste.) A good mode of keeping almonds in a state fit for making emulsions.

CONFECTIO ARCHIG'ENI, *C. Pauli'na*. (Castor, long pepper, black pepper, storax, galbanum, costus, opium, of each 3ss; saffron, 3ij; syrup of wormwood, 3ij.) It was much recommended as a stimulant in nervous affections.

CONFECTIO AROMATICA, *C. Cardi'aca* seu *Raleigh'na*, *Electuarium Aromaticum*, *Aromatic Confection*, *Sir Walter Raleigh's Cordial*. The following is the formula, (Ph. U. S.): Pulv. aromat. 3vss; croci, in pulv. 3ss; syrup. aurant. 3vj; Mel despumat. 3ij. Rub the aromatic powder with the saffron; then add the syrup and honey, and beat together until thoroughly mixed. Dose, ʒj to 3j.

CONFECTIO AURANTIO'RUM, *C. Aurantii Corticis*, (Ph. U. S.), *Conserua Corticium Aurantiorum* seu *Corticis Exterioris Aurantii Hispanensis* seu *Flavedinis Corticium Aurantium Hispanicum* seu *Aurantii* seu *Citri Aurantii*, Confection of the Orange, Conserve of Orange Peel. (Yellow part of the peel of the orange, ʒij; rectified sugar, ʒiij; beat into a conserve.) It is stomachic, and an agreeable vehicle, corrigent, and adjuvant for other remedies. Dose, 3ss.

CONFECTIO CARDIACA, Confectio aromatica.

CONFECTIO CAS'SIA, *Conserua cassia*, *Electuarium Cassia Fistula* seu *Cassia* seu *e Cassia* seu *laxativum* seu *Cassia tamarindatum* seu *lenitivum* seu *e Cassia*, *Diacas'sia cum Manna*, *Cassia Confection*. (*Cassia* pulp, ʒbss; manna, 3ij; tamarind pulp, 3j; syrup of roses, f3viij. Bruise the manna, dissolve by heat, mix the pulp, and evaporate.—(Ph. L.) It is gently laxative. Dose, 3ss.

CONFECTIO CAT'ECU COMPOS'ITA, *Electuarium Catechu*, Compound Confection of Catechu. (Pulv. catechu comp. 3v (avoirdupois); Syrup. f3v, mix. (Ph. D.) The Edinburgh confection is formed of catechu and kino, each 3iv; cinnamon and nutmeg, each 3j; opium, diffused in a little sherry, 3iss; Syrup of Red Roses, of the consistence of honey, Oiss (Imp. meas.): mixed and thoroughly beaten together. A grain of opium is contained in about 200 grains of the mass. Dose, as an astringent, 3ss to 3j.

CONFECTIO DAMOCRATIS, Mithridate.

CONFECTIO HAMEC—so called from an Arabian physician—was composed of the bark of the yellow myrobalans, black myrobalans, violets, pulp of colocynth, polypodium of the oak, leaves of wormwood, thyme, aniseed, fennel, red roses, pulps of prunes, raisins, sugar, honey, senna, rhubarb, &c. It was used as a purgative in glysters.

CONFECTIO HYACIN'THI, *Hyacinth Confection*, *Electuarium* seu *Confectio de Croco emendata olim dicta de hyacinthis*, (Ph. P.) A tonic and slightly astringent confection, composed, according to Baumé, of terra sigillata, crabs' stones, cinnamon, leaves of the dittany of Crete, myrrh, saffron, syrup of lemon, camphor, Narbonne honey, oil of lemon, &c.: and formerly the hyacinth was added, whence its name.

CONFECTIO O'PII, *C. opia'ta*, *Electuarium Opia'tum* seu *O'pii*, *Philo'nium Londinen'se* seu *Roma'nium*, *Theriaca Edinen'sis*, *Electuarium Thebaicum*, *Opium Confection*, (Opium, in powder, 3vss; Pulv. aromat. 3vj; Mel despumat, 3xiv:

mix thoroughly. One grain of opium in 36:—Ph. U. S.) It is narcotic and stimulant. Dose, gr. x to ʒj.

CONFECTIO PAULINA, Confectio Archigeni—c. ex Capitis papaveris, Diacodium.

CONFECTIO PIP'ERIS, *C. Piperis nigri*, *Electuarium Piperis*, *Ward's Paste for Fistula*. (Black pepper, elecampane root, each ʒij; fennel seed, ʒiij; honey and sugar, each ʒiij, (Ph. U. S.) Dose, size of a nutmeg, three or four times a day.

CONFECTIO RALEIGHANA, Confectio aromatica.

CONFECTIO ROSÆ, *C. Rosæ Gallicæ* seu *Rosæ rubræ*, *Conserua Rosæ* seu *Rosæ Gallicæ* seu *Florum Rosarum Rubrarum* seu *Florum Rosæ Rubræ*, *Rhodosa'charum*, *Sac'charum Rosaceum*, Confection of the Red Rose. (Red roses, in powder, 3iv; Sugar, in powder, 3xxx; Clarified honey, 3vj; Rose water, f3viij. Rub the roses with the rose water at a boiling heat; then add gradually the sugar and honey, and beat until thoroughly mixed.—Ph. U. S.) It is astringent, and chiefly used as a vehicle for other remedies.

CONFECTIO ROSÆ CANI'NÆ, *Conserua Rosæ Cani'næ* seu *Rosæ Fructus* seu *Cynos'bati* seu *Cynor'rhodi*, *Rob Cynos'bato*, Confection or Conserve of Dog Rose. (Pulp of dog rose, ʒij; sugar, 3xx. Incorporate. Ph. L.) It is chiefly used as a vehicle for other remedies.

CONFECTIO RUTÆ, Confection of Rue. (Rue leaves, dried, caraway seeds, bay berries, of each 3iss; sagepenny, 3iv; black pepper, 3ij; honey, 3xvi. Mix.—Ph. L.) It is given in clysters, as an antispasmodic and carminative.

CONFECTIO DE SAN'TALIS, Confection of the Sanders, (F.) Confection de Sandaure. (Sandal wood, red coral, bole armeniac, terra sigillata, kermes berries, tormentil root, dittany, saffron, myrtle, red roses, calcined hartshorn, and cloves.) It was formerly used as an astringent.

CONFECTIO SCAMMO'NI, *Electuarium Scammoni* seu *e Scammoni*, *Caryocostivum*, Confection of Scammony. (Powdered scammony, 3iss; bruised cloves, powdered ginger, āā 3vj; oil of caraway, 3ss; syrup of roses, q. s.—(Ph. L.) A stimulating cathartic. Dose, 3ss to 3j.

CONFECTIO SENNÆ, *Electuarium Cassia Sennæ*, *E. Sennæ cum Pulpis* seu *Sennæ compos'itum* seu *ape'riens* seu *cathol'icum commu'nē* seu *diap'ri'um* seu *ecoprot'icum* seu *Sennæ* seu *e Sennæ* seu *lenitivum*, *Benedictivum Laxativum*, Confection of Senna, *Lenitive Electuary*, &c. (Senna leaves, 3viij; figs, ʒij; tamarind pulp, cassia pulp, pulp of French prunes, each ʒbss; coriander seed, 3iv; liquorice root, 3ij; sugar, ʒbss; water, Oiv. Rub the senna and coriander together: separate 10 ounces of the powder with a sieve; boil the residue with the figs and liquorice root in the water to one-half; press out the liquor and strain. Evaporate the liquor by means of a water bath to a pint and a half; add the sugar, and form a syrup. Rub the pulps with the syrup, gradually added; throw in the sifted powder, and beat till thoroughly mixed.—Ph. U. S.) It is a laxative, and is used in habitual constipation, and in constipation during pregnancy. Dose, 3ss to 3ss.

CONFECTIO SULPH'URIS, Confection of Sulphur (Sulphur. sublim. 3ij; Potass. Bitart. 3j; Mel. despumat. 3j; Syrup. Zingib. Syrup. Croci, āā f3ss. M.—Ph. D.) As a laxative and alterative. Dose, 3j to 3j.

CONFECTIO TEREBINTHINÆ, Confection of Turpentine. (Ol. Tereb. f3j; Pulv. Glycyrrhiz. 3j; Mel. despumat. 3ij.—Ph. D.) An agreeable mode of administering turpentine. Dose, a scruple to a drachm.

CONFECTIO DE THURĒ, *Frank'incense Confection*, (F.) Confection d'Encens. A compound of coriander seeds, nutmeg, thus, liquorice, mastich

cubus, prepared hartshorn, conserve of red roses, sugar, &c.

CONFECTION, Confectio—*c. d'Amandes*, Confection, almond—*c. of Catechu*, compound, Confectio catechu compositum—*c. d'Encens*, Confectio de Thure—*c. Frankincense*, Confectio de Thure *c. of the Orange*, Confectio aurantiorum—*c. of the Red rose*, Confectio rosæ Gallicæ—*c. of Rue*, Confectio rutæ—*c. de Sandaur*, Confectio de santalis—*c. of the Sanders*, Confectio de santalis—*c. of Sulphur*, Confectio sulphuris—*c. of Turpentine*, Confectio terebinthinae.

CONFECTUM, Confection.

CONFERVA HELMINTHOCORTOS, Coralina Corsicana.

CONFERVA RIVA'LIS. This species of *River Weed* has been recommended in cases of spasmodic asthma, phthisis, &c.

CONFINEMENT, Parturient state.

CONFIRMANTIA, Tonics.

CONFIRMED, Consummatus.

CONFIT, Confectio.

CONFITURE, Confectio.

CONFLUENT, *Confluens*, from *con*, and *fluere*, 'to flow' 'Running together.' An epithet for certain exanthematous affections, in which the pimples, pustules, &c., run together. It is particularly applied to small-pox, so circumstanced. Some authors have called scarlatina or scarlet fever *Confluent Measles*, *Morbili Confluentes*.

CONFLUENT DES SINUS, Torcular Herophili.

CONFLUENTIA, *Confœderatio*; same etymon as *Confluent*. A term, employed by Paracelsus to express the concordance between a disease and its remedies.

CONFLUXIO, Sympathy.

CONFEDERATIO, Confluentia.

CONFORMATIO, Conformation, Structure.

CONFORMATIO, *Conformatio*, *Diapylæsis*, *Diapylæmus*, Structure, from *conformare*, (*con*, and *formare*, *forhatum*, 'to form,' 'to arrange,' 'dispose.' The natural disposition or arrangement of the body.

Faulty conformation, (*F.*) *Vice de conformation*, is vice of original formation; existing, of course, from birth. In French surgery, *Conformation* is used synonymously with *Coaptation*, and both it, *Diapylæsis* and *Anapylæsis* mean, also, restoration to the original form—as in fractures, &c.

CONFORTANTIA, Tonics.

CONFORTATIVA, Tonics.

CONFORTER (*F.*), *Confirma'rê*, *Conforta'rê*, *Corroborat'rê*, (*con*, and *fortis*, 'strong.') To make stronger—to give energy. *Conforter l'estomac*, 'to strengthen the stomach.'

CONFRICATIO, *Confricatio*, from *confricare*, (*con*, and *fricare*, *fricatum*,) 'to rub.' The action of reducing a friable substance to powder, by rubbing it between the fingers; and of expressing the juice of a plant with the hand.

CONFRICA'TRIX, from *con*, 'with,' and *fricare*, *fricatum*, 'to rub.' A female who practises masturbation.

CONFUSÆ FEBRES. Intermittents, whose paroxysms are irregular and confused.—Bellini.

CONFUSIO, from *confundo*, (*con* and *fundere*, *fundere*, 'to pour,') 'I mix together.' *Syn'chysis*. A disease of the eye, which consists in the mixture of the humours.—Galen. In modern times, *Synchysis* has been applied to a morbid state of the vitreous body, in which it is reduced to a diffident condition. Occasionally, sparkling bodies are perceptible, which seem to be seated behind the crystalline, in the vitreous humour: hence, the disease has been called *sparkling synchysis* or *spintheropia*, from *σπινθηρ*, 'a spark,' and *ωψ*, 'the eye;' (*F.*) *Spintheropie*, *Scintille-*

ment de l'œil. *Confusio* has, also, been used synonymously with *Complexio*.

CONFUSIONES ANIMI, Affections of the mind.

CONGEE DISCHARGES, Rice-water discharges.

CONGELANTIA, Congelativa.

CONGELATIO, Catalepsy.

CONGELATIO, *Congelatio*, *Conglaciatio*, *Gelatio*, from *congelare*, (*con*, and *gelare*, *gelatum*,) 'I congeal,' 'I freeze.' The action of congealing, of passing to the solid state by the abstraction of heat; as *congelation of water*, *mercury*, &c. The term had once other acceptations. 1. It was synonymous with *concretion*. 2. With *coagulation*, in which sense it is still occasionally employed. 3. 'The ancients called all diseases, produced by cold, *congelations*, as well as those in which there was a kind of stupor or torpor—particularly catalepsy. Also, *Frostbite*, (*Prov.*) *Krebnitz*.

Congelation of a part, by the application of powdered ice or of a freezing mixture, in a bladder or gauze bag, has been employed to induce anesthesia in the lesser surgical operations. It has also been used in external inflammation.

CONGELATIVA MEDICAMENTA, *Conglutinantia*, *Congelantia*. Medicines, considered capable of uniting or consolidating wounds, &c.

CON'GENER, *Congenerous*, from *con*, and *genus*, *generis*, 'kind,' (*F.*) *Congénère*. Of the same kind or species. Resembling each other in some manner. When applied to muscles, it means, that they concur in the same action; in opposition to the word *antagonist*, or that which acts in an opposite direction.

In France *Congénères* is applied to those who join in the dissection of the same subject.

CONGÉNIAL, Congenital.

CONGENIALIS, Congenital.

CONGENITE NOTÆ, Nævus.

CONGENITAL, *Con'genite*, *Congenialis*, *Con'genitus*, *Syngenicus*, from *con*, and *genitus*, 'begotten,' (*F.*) *Congénial* or *Congénital*. Diseases which infants have at birth: hence, *Congenital affections* are those that depend on faulty conformation: as *congenital hernia*, *congenital cataract*, &c. See *Connate*.

CONGESTED, Hyperæmic.

CONGESTIO, Congestion—*c. Abdominalis*, *Cœliæmia*—*c. Pectoris*, *Stethæmia*—*c. Pulmonum*, *Stethæmia*—*c. Sanguinis*, *Congestion*.

CONGESTION, *Congestio*, *Rhopê*, from *congerere*, (*con*, and *gerere*, *gentum*, 'to carry,') 'to amass,' 'accumulate,' &c. *Symph'ora*, *Hæmatopogo'gê*, *Hæmatosymphore'sis*, *Hæmatosymphogê*, *Hæmorme'sis*, *Symphore'ma*, *Symphore'sis*, *Synthrois'mus*, *Synthrois'sis*, *Syn'guinis Congestio*, *Engorgement*. Accumulation of blood—*hyperæmia*—in an organ. It is an important symptom in febrile and other disorders. It may arise either from an extraordinary flow of blood by the arteries, or from a difficulty in the return of blood to the heart by the veins. More often, perhaps, it is owing to the latter cause, and is termed *venous congestion*, *stasis* or *stagnation*—being not usually attended with symptoms of oppression and collapse. See *Hyperæmia*.

CONGESTION OF THE ABDOMEN. *Cœliæmia*—*c. of the Brain*, *Stethæmia*—*c. Cérébrale*, *Cephalæmia*—*c. du Cerveau*, *Cephalæmia*—*c. of the Lungs*, *Stethæmia*—*c. des Poumons*, *Stethæmia*—*c. Sanguine rachidienne*, *Hypermyelohæmia*.

CONGESTIVE, *Congestivus*, (*F.*) *Congestif*. Belonging or relating to, or affected with, *congestion*—as '*congestive fever*.'

CONGESTIVE FEVER, see *Fever*, *congestive*.

CONGLACIATIO, Congelation.

CONGLOBATE, *Conglobatus*, from *conglobare*,

(*oon*, and *globus*, 'a ball,') 'to collect,' 'to gather into a ball.' (F.) *Conglobé*.

CONGLOBATE GLAND, *Glandula conglobata*, *Hydraden*, *Globate gland*, *Lymphatic gland*, (F.) *Glande Conglobée*, *Ganglion lymphatique*. A round body, formed of lymphatic vessels, connected together by cellular structure, but having neither a cavity nor excretory duct. The mesenteric, inguinal and axillary glands are of this class.

CONGLOBÉ, Conglobate.

CONGLOMERATE. *Conglomeratus*, from *con*, and *glomerare*, *glomeratum*, 'to gather in a heap,' *Glomerate*, *Glomera'tus*, (F.) *Congloméré*. Heaped together.

CONGLOMERATE GLANDS, *Glandula conglomerata* seu *vasculosa*, are those whose lobules are united under the same membrane; as the liver, kidney, testicle, &c.

CONGLOMERATIO INTESTINORUM, Epiploce intestinales.

CONGLUTINANTIA, Congelativa.

CONGRÈS (F.), *Congress*, *Congressus*, from *congregari*, *congressus*, (*con*, and *gradi*, 'to go,') 'to go together.' This term, which has often been used synonymously with *Coition*, means, also, the ocular proof, formerly ordered by judicial authority, in the presence of surgeons and matrons, to test the impotence or capabilities of parties:—a most unsatisfactory and indecent exhibition. It was forbidden by the Parliament of Paris in the year 1667.

CONGRESSUS, Coition.

CONI VASCULO'SI, *Cor'pora pyramidalia testis*. Conical bundles, formed by the vasa efferentia of the testis; having their base towards the epididymis, into the tube of which they enter.

CONIA, *kovia*. A wine, prepared by fermenting the must of the grape on tar previously washed in sea-water.—*Orfila*. See, also, *Conium*, and *Lixivium*.

CONIASIS, Incrustation.

CONICINE, see *Conium*.

CONII FOLIA, see *Conium*—c. *Semen*, see *Conium*.

CONIINE, see *Conium*.

CONINE, see *Conium*.

CONIOSTOSIS, Pulverization.

CONIS, Pulvis.

CONISTERIUM, Apodyterium.

CONIUM, *C. maculatum*, *Abi'otes*, *Corian'drum maculatum*, *Cicuta major* seu *maculata* seu *Storckii* seu *terrestris* seu *major foetida* seu *vulgaris*, *Common* or *Poison Hemlock*, *Hemlock*, *Poison parsley*, *Spotted parsley*, (Sc.) *Humlock*, (F.) *Ciguë ordinaire*, *C. grande* ou *officinale*. *Ord. Umbelliferae*. *Sex. Syst.* Pentandria Digynia. The leaves, *Coni'i Folia*, and seeds, *Coni'i Semen*, are narcotic and poisonous in an overdose. Light destroys the virtues of the leaves; and, therefore, the powder should be kept in opaque bottles, well corked. It has been used as a palliative in cancer and other painful affections; but is not equal to opium. Externally, it has been applied in fomentation to cancerous and scrofulous ulcers. Dose, gr. ij to x.

Its active principle is *Conia*, *Conine*, *Coniine*, *Confine*, *Conicine*, *Cicutine*. It, or its salts, have been given as sedatives to the nervous centres in various neuralgic and spasmodic diseases. It is a most energetic poison.

Da'vidson's Remedy for Cancer is said to consist of powdered hemlock and arsenious acid.

CONIUM MOSCHA'TUM, *Aracacha*: A very agreeable and nutritive kind of tuberous vegetable, in flavour not unlike celery, which grows on the coast of Peru, but is more abundant on the projecting ridges of the Cordilleras, and on the east-

ern declivity of the Andes. It is cooked by being either simply boiled in water, or made into a kind of soup.—*Tschudi*.

CONJUNCTIVE, Conjunctiva.

CONJUNCTIVITE, see *Ophthalmia*—c. *Blennorrhagica*, see *Ophthalmia*.

CONJUGAISON, Conjugation.

CONJUGATION, *Conjugatio*, from *conjungere*, *conjunctio*, (*con*, and *jugum*, 'a yoke,') 'to yoke together.' (F.) *Conjugaison*. Assemblage, union, —*CONJUGIUM*.

CONJUGATIONIS FORAMINA, (F.) *Trous de conjugaison*. The apertures at the sides of the spine, formed by the union of the notches of the vertebrae. They give passage to the nerves of the spinal marrow, and to the vessels which enter or issue from the vertebral canal.

CONJUGIUM, Coition, Conjugation.

CONJUNCTI (MORBI), from *conjungere*, (*con*, and *jungere*, *junctum*,) 'to join together.' Diseases joined together. Authors have distinguished two kinds of these: one, in which the diseases go on simultaneously—*morbi connexi*; the other, in which they succeed each other—*morbi consequentes*.

CONJUNCTIO, Articulation.

CONJUNCTIVA, CUTICULAR, Xerophthalmia—c. *Granular*, *Trachoma*.

CONJUNCTIVA MEMBRANA, *Circumcunctilis*, *Epipheph'geos*, *Tunica aquata* seu *adnata* seu *conjunctiva*, *Tunda oculi*, (F.) *Conjunctive*, *Membrane adnée*. A mucous membrane, so called because it unites the globe of the eye with the eyelids. It covers the anterior surface of the eye, the inner surface of the eyelids, and the *caruncula lacrymalis*. It possesses great general sensibility, communicated to it by the fifth pair.

CONJUNCTIVITIS, Ophthalmia—c. *Ægyptiaca*, *Ophthalmia*, *purulent*—c. *Blennorrhagica*, see *Ophthalmia*—c. *Catarrhalis*, *Ophthalmia*, *catarrhal*—c. *Gonorrhoeica*, see *Ophthalmia*—c. *Puro-mucosa catarrhalis*, *Ophthalmia*, *catarrhal*—c. *Puro-mucosa contagiosa* seu *Ægyptiaca*, *Ophthalmia*, *purulent*.

CONJUGENS, Communicans.

CONNATE, *Conna'tus*, (F.) *Conné*, from *con*, and *natus*, 'born with.'

CONNATE DISEASES, (F.) *Maladies connées*, *Morbi connati*, are such as an individual is born with:—*connate* having the same signification as *congenital*. A difference has been made by some, however; those diseases or conditions which are dependent upon original conformation, being called *congenital*;—whilst the diseases or affections that may have supervened during gestation or delivery, are termed *connate*.

CONNÉ, Connate.

CONNECTICUT, MINERAL WATERS OF. There is a mineral spring at Stafford, in this State, twenty-four miles from Hartford, which has obtained more celebrity than any one in New England. Its principal ingredients are iron and carbonic acid. It, consequently, belongs to the class of acidulous chalybeates. There are other springs in the State, of which, however, little that is accurate is known.

CONNECTIVE TISSUE, Cellular tissue.

CONNERVATIO, Syndesmosis.

CONNEXIO OSSIIUM CARNOSA, *Syssa'reosis*—c. *Cartilaginea*, *Synchondrosis*—c. *Ligamentosa*, *Syndesmosis*.

CONNUTRI'TUS, *Syn'trophos*, from *con*, 'with,' and *nutrio*, *nutritus*, 'to be nourished.' A disease is so called which has grown up, as it were, with an individual, or has been connate with him.—*Hippocrates*, *Galen*.

CONOID, *Conoi'deus*, *Conoi'des*, from *κωνος*, 'a cone,' and *eidos*, 'shape.' (F.) *Conoïde*. Resembling a cone.

CONOID LIG'AMENT, *Ligamentum conoïdes*. A ligament, passing from the coracoid process to the scapula, and forming part of the coraco-clavicular ligament of some anatomists.

CONOIDAL SUBSTANCE OF THE KIDNEY, see Kidney.

CONOIDES CORPUS, Pineal gland.

CONOPHTHALMIA, Staphylocoma cornæ.

CONQUASSANT, (F.) *Conquassans*, from *con*, 'with,' 'together,' and *quassare*, 'to shake.' *Douleurs conquassantes* are the pains of parturition, at the time of their greatest intensity, when the head is engaged in the pelvis.

CONQUASSATION, *Conquassatio*, *Quassatio*, *Quassatura*. Same etymon. A pharmaceutical operation, which consists in dividing, with a pestle, fresh vegetables, fruits, &c. See Confection.

CONQUASSATIONES ANIMI, Affections of the mind.

CONQUE, Concha.

CON'SCIOUSNESS, DOUBLE. A somnambulist condition, in which the individual leads, as it were, two lives, recollecting in each condition what occurred in previous conditions of the same character, but knowing nothing of the occurrences of the other. See Duality of the Mind.

CONSEC'UTIVE, *Consecutivus*, from *con*, 'with,' and *sequor*, *secutus*, 'to follow.'

CONSECUTIVE PHENOM'ENA OR SYMPTOMS, (F.) *Phénomènes ou accidens consécutifs*, are such as appear after the cessation of a disease, or, according to others, during its decline; but without having any direct connexion with it.

CONSENSUAL, see Instinctive.

CONSENSUS, Consent of parts, Sympathy.

CONSEN'SUS OCULO'RUM. The intimate association between the two eyes, as exemplified in their simultaneous action in health, and often in disease.

CONSENT OF PARTS, *Consensus*, *Consensus partium*, *Sympathia*, from *con*, and *sentire*, *sensum*, 'to feel.' (F.) *Consentement des Parties*. That relation of different parts of the body with each other which is more commonly denominated sympathy.

CONSEITEMENT DES PARTIES, Consent of parts.

CONSERVATION, *Conservatio*, *Phylaxis*, from *conservare*, (*con*, and *servare*, *servatum*), 'to preserve.' (F.) *Conservation*, *Asservation*. The art of preserving any object of pharmacy, any remedial agent, &c., from decay.

CONSERVA, *Conserve*, (F.) *Conserve*, *Electuaire simple*, *Saccharolé mou*. Same etymon. A pharmaceutical preparation, composed of a vegetable substance and a sufficient quantity of sugar beaten into a uniform mass. See Confection.

CONSERVA ABSIN'THII, *C. absin'thii marit'imi*, *Conserve of Wormwood*. (Leaves lbj, sugar lbij.) It has been employed as a tonic, stomachic, and vermifuge.

CONSERVA AMYGDALARUM, *Confectio amygdalarum*.

CONSERVA ANGEL'ICÆ, (Ph. P.) *Conserve d'Angélique*, *C. of Angel'ica*. (Pulp of root 250 parts; white sugar, boiled in a decoction of the root, and reduced to the consistence of a solid electuary, 1000 parts.) It is tonic, aromatic, and stomachic.

CONSERVA DE A'PIO GRAVEOLEN'TE (Ph. P.), *Conserve d'Ache*, *Conserve of Smallage*. Prepared like the preceding. Reputed to have the same properties.

CONSERVA ARI, *Conserve of Arum*. (Fresh root lbss, sugar lbss.) Esteemed to be diuretic and stimulant.

CONSERVA AURANTII, *Confectio aurantiorum*—c. Cassiæ, *Confectio cassiæ*—c. Citri aurantii, *Confectio aurantiorum*.

CONSERVA COCHLEA'RIS HORTEN'SIS, *Conserve of Lemon Scurvy Grass*. (Leaves lbj, sugar lbij.) Reputed to be stimulant and antiscorbutic.

CONSERVA CORTICIS EXTERIORIS AURANTII HISPALENSIS, *Confectio aurantiorum*—c. *Corticum aurantiorum*, *Confectio aurantiorum*—c. *Florum rosarum rubrarum*, *Confectio rosæ Gallicæ*—c. *Cynosbati*, *Confectio rosæ caninæ*—c. *Cynorrhodi*, *Confectio rosæ caninæ*—c. *Flavodinis corticis aurantiorum Hispalensium*, *Confectio aurantiorum*.

CONSERVA LU'JULÆ, *C. Folio'rum lu'julæ*, *Conserve of Woodsorrel*. (Leaves lbj, sugar lbij.) Gratefully acid and refrigerant.

CONSERVA MENTHÆ, *C. Menthæ folio'rum*, *C. Menthæ sativæ*, *Conserve of Mint*. (Leaves lbj, sugar lbij.) Stomachic in nausea and vomiting.

CONSERVA PRUNI SYLVEST'IS, *C. Prunæ sylvestris*, *Pulpa pruno'rum sylvestrium cond'ita*, *Conserve of Sloes*, (Pulp 1 part, sugar 3 parts,) Possessed of astringent properties.

CONSERVA ROSÆ, *Confectio rosæ Gallicæ*—c. *Rosæ caninæ*, *Confectio rosæ caninæ*—c. *Rosæ fructūs*, *Confectio rosæ caninæ*—c. *Rosæ Gallicæ*, *Confectio rosæ Gallicæ*.

CONSERVA SCILLÆ, *Conserve of Squill*. (Fresh squills 3j, sugar 3x.) Possesses the diuretic and other virtues of the squill.

CONSERVE, *Conserve*, see *Confectio*—c. *d'Ache*, *Conserve de apio graveolente*—c. of Aloes, *Conserve pruni sylvestris*—c. of Lemon scurvy grass, *Conserve cochleariæ hortensis*—c. of Mint, *Conserve menthæ*—c. of Orange, *Confectio aurantiorum*—c. of Roses (red), *Confectio rosæ gallicæ*—c. of Smallage, *Conserve de apio graveolente*—c. of Woodsorrel, *Conserve lujulæ*—c. of Wormwood, *Conserve absinthii*.

CONSERVES, Spectacles.

CONSIDEN'TIA, from *considera*, (*con* and *sedo*) 'to settle.' This word has two acceptations. 1. It is synonymous with *Apocatastasis*, and means restoration or cessation, and the subsiding of a humour, as thearic; and 2. It signifies contraction of any cavity or canal.—See Synæsis.

CONSISTENTIA, *Consistere*, (*con* and *sistere*) 'to stand still.' A term employed in two senses. 1. When joined to the word *Morbi* or *Ætiæ*, it expresses the acme of a disease, or the age at which the constitution has acquired its full strength. 2. By *Consistentia humoris* is meant the density of a humour.

CONSOLIDA MAJOR, *Symphytum*—c. *Media*, *Ajuga*, *Chrysanthemum leucanthemum*—c. *Minor*, *Prunella*—c. *Regalis*, *Delphinium consolida*—c. *Rubra*, *Tormentil*—c. *Saracenica*, *Solidago virgaurea*.

CONSOLIDAN'TIA, *Consolidativa Medica*, from *con*, and *solidus*, solid. Substances, formerly given for the purpose of consolidating wounds, or strengthening cicatrices.

CONSOLIDATIVA, *Consolidantia*.

CONSUMME (F.), *Consumma'tum*, from *consummo*, (*con* and *summa*) 'I add together,' *Zonot*. Soup strongly charged with gelatin, and consequently very nutritious, although not proportionably easy of digestion.

CONSUMPTION, Consumption

CONSONANCE, see Sound.

CON'SONANT, *Con'sonans*, from *consono*, 'to sound together,' (*con*, 'with,' and *sono*, 'to sound,') because it is generally believed that a consonant cannot be properly expressed, except when conjoined with a vowel. Physiologically, a breath, or sound produced in the larynx, which suffers more or less interruption in its passage through the vocal tube.

CONSORTIUM. Sympathy.

CONSOUDE, *GRANDE*, Symplytum.

CONSOUND, MIDDLE, Ajuga.

CONSPERSIO, Catapasma.

CONSPICILLA, Spectacles.

CONSPIRATIO, Sympathy.

CONSTELLATUM UNGUENTUM. An ointment composed of cleansed earthworms! dried and pulverized; and of the fat of the bear or wild boar. It was formerly employed in tooth-ach, and to hasten the cicatrization of wounds.

CONSTERNATIO, Stupor.

CONSTIPATED, *Constipatus*, (Prov.) *Maw-bound*, (F.) *Constipé*; same etymon as the next. Affected with constipation; Costive; *Bound*, *Belly-bound*.

CONSTIPATIO, Constipation, Stegosis.

CONSTIPATION, *Constipatio*, from *constipare*, (con and *stipare*, *stipatum*), 'to cram close.' *Obstipatio*, *Adstrictio*, *Arctatio*, *Obstipatio* seu *Retentio alvi*na, *Ischocolia*, *Alvus astricta* seu *adstricta* seu *tarda* seu *dura* seu *seguis*, *Obstructio* seu *Suppressio alvi*, *O. Ductus alimentarii*, *O. intestinalis*, *Torpor intestinum*, *Stypsis*, *Constipatio alvi*, *Coprosistis*, *Acopria*, *Acoprosis*, *Coproëpis*, *Ischocolia*, *Ischocolia*, (F.) *Échauffement*, *Ventre reserré*, *Costiveness*, *Fæcal Retention*, *Alvine obstruction*. A state of the bowels, in which the evacuations do not take place as frequently as usual; or are inordinately hard, and expelled with difficulty. It may be owing either to diminished action of the muscular coat of the intestines, or to diminished secretion from the mucous membrane, or to both. Cathartics will usually remove it; after which its exciting and predisponent causes must be inquired into and obviated, to render the cure permanent.

CONSTIPATUS, Constipated.

CONSTIPÉ, Constipated.

CONSTITUENS, Vehicle, see Prescription.

CONSTITUTIO, Constitution, Habit of body—c. *Aeris*, Constitution of the atmosphere—c. *Epidemica*, Constitution, epidemic—c. *Nervosa*, Nervous diathesis.

CONSTITUTION, *Constitutio*, *Catasistis*, *Status*, from *con*, and *statuere*, from *stare*, 'to stand.' A collection of several parts, forming a whole. In medicine, *Constitution* means the state of all the organs of the human body considered in regard to their special and relative arrangement, order, or activity. A good constitution is one in which every organ is well developed, and endowed with due energy, so that all perform their functions with equal facility. Any want of equilibrium in their development and energy forms a difference in the constitution. We say that a man is of a good or robust, a delicate or weak constitution, when he is commonly healthy, or commonly labouring under, or unusually susceptible of, disease.

CONSTITUTION, ARTERIAL, Plethora arteriosa.

CONSTITUTION OF THE ATMOSPHERE, *Constitutio Aëris*, (F.) *Constitution Atmosphérique*. The condition of the atmosphere, as regards dryness and humidity, temperature, heaviness, direction of the winds, &c., considered as respects its influence on the animal economy.

CONSTITUTION, EPIDEMIC, *Constitutio epidemica*, *Medical Constitution*, (F.) *Constitution médicale*, *C. Epidémique*. The aggregate of meteorological conditions, so far as they are appreciable, during which diseases prevail epidemically.

CONSTITUTIONAL, (F.) *Constitutionnel*. Belonging to the constitution of an individual; to his manner of being; as *constitutional phthisis*, *c. gout*, &c. By some, this epithet has been given to diseases, produced by the constitution

of the atmosphere; but this acceptance is not common.

CONSTRICTEUR D'HERBINEAUX, see *Serre-neud*.

CONSTRICTEUR DE LA VUIVE, *Constrictores cunni*—c. *du Vagin*, *Constrictores cunni*. *CONSTRICTO*, *Astrictio*, *Systole*.

CONSTRICTIVA, Styptics.

CONSTRICTOR, from *constringere*, (con, and *stringere*, *strictum*), 'to bind.' (F.) *Constrictor*. That which binds in a circular direction. A sphincter. Different muscles are so called.

CONSTRUCTOR ANI, Sphincter ani externus—c. of Nuck, Compressor of Nuck.

CONSTRICTORES ALARUM NASI, Depressor alæ nasi.

CONSTRICTORES CUNNI, *C. Vaginé* seu *Vulvæ*, *Clitoridis inferiores latè et plani musculi*, *Sphincter Vaginé*, (F.) *Constrictores du vagin*, *C. de la Vulve*. Small muscles which originate beneath the clitoris, descend along the sides of the vagina, and terminate by becoming confounded with the *transversus perinæi* and external sphincter ani muscles. Their use is to contract the entrance of the vagina.

CONSTRICTORES ISTHMI FAUCIUM, *Glossostaphylinus*.

CONSTRICTORES LARYNGIS, Lieutaud describes, under the name *Grand constrictor du Larynx*, the muscle *Cricico-arytenoideus lateralis* with the *Thyro-arytenoideus*.

CONSTRICTORES NASI, Compressor nasii.

CONSTRUCTOR ŒSOPHAGI, *Constrictor of the Œsophagus*, (F.) *Constricteur de l'Œsophage*, *Muscle œsophagien*. A fasciculus of fleshy, circular fibres, at the upper part of the œsophagus.

CONSTRICTORES ORIS, Orbicularis oris—c. *Palpebrarum*, *Orbicularis palpebrarum*.

CONSTRICTORES PHARYNGIS, *Constrictors of the Pharynx*, *Sphincter Gulae*. Muscular expansions which assist in forming the parietes of the pharynx. Three of these are generally admitted.

1. The *Constrictor Pharyngis inferior*, *Cricopharyngeus* and *Thyro-pharyngeus*, (F.) *Cricothyro-pharyngien*. It is broad, very thin, quadrilateral, seated superficially, extending from the thyroid and cricoid cartilages as far as the middle of the pharynx, and uniting on the median line, with that of the opposite side. 2. The *Constrictor Pharyngis Medius*, *Hyo-pharyngeus* and *Chondro-pharyngeus*, *Syndesmo-pharyngeus*, (F.) *Hyo-glossobasi-pharyngien*, occupies the middle part of the pharynx. It is triangular, and attached anteriorly to the great and little cornu of the os hyoides; to the stylohyoid ligament; and terminates, posteriorly, by joining its fellow of the opposite side. 3. The *Constrictor Pharyngis superior*, *Cephalo-pharyngeus*, *Glossopharyngeus*, *Mylo-pharyngeus*, *Pterygo-pharyngeus*, (F.) *Pterygo-syndesmo-staphili-pharyngien*, is quadrilateral, and extends from the internal ala of the pterygoid process, from the inter-maxillary ligament, from the internal oblique line of the jaw, and from the base of the tongue to the posterior part of the pharynx.

The constrictors of the pharynx contract it. They can likewise approximate the ends to each other. Chaussier considers those and the *stylopharyngeus* as but one muscle, and includes all under this last name.

CONSTRICTORES VAGINÆ, *C. cunni*—c. *Vulvæ*, *C. cunni*.

CONSTRINGENTIA, Astringents, Styptics.

CONSUETUDO, Habit—c. *Menstruæ*, *Menses*.

CONSULTANT, Consulting physician or surgeon.

CONSULTATION, *Consulta*, *Consultatio*, *Deliberatio*, *Cœnologia*, *Contentio*, *Symbolus*, from *consuere*, *consultum*, 'to hold council.' This word has

several acceptations. In English, it means, almost always, the meeting of two or more practitioners, to deliberate on any particular case of disease. In France, it signifies the written result of such deliberations, as well as the opinion of a physician, given to a patient, who consults him, either personally or by writing.

CONSULTING PHYSICIAN or SURGEON, (F.) *Médecin ou Chirurgien consultant, Consultant*. One who consults with the attending practitioner, regarding any case of disease. Some physicians, surgeons, and accoucheurs confine themselves to consulting practice.

CONSUMMATUM, *Consummé*.

CONSUMMATUS, from *con, cum*, 'with,' and *summus*, 'the whole.' Confirmed; established; developed,—as *Phthisis consummata*, 'confirmed consumption.'

CONSUMPTION, from *consumere*, (*con* and *sumere, sumptum*), 'to waste away:' *Consumptio, Consumptio, Syntexis*, Wasting, Wearing, (F.) *Consumption*. Progressive emaciation or wasting away. This condition precedes death in the greater part of chronic diseases, and particularly in *phthisis pulmonalis*: on this account it is, that phthisis has received the name *consumption*—See *Phthisis*. *Fèvre de Consumption, Consumptive fever*, is the same as *Hectic fever*.

CONSUMPTION OF THE BOWELS, Enterophthisis.

CONSUMPTION, GALLOPING, *Phthisis acuta*, (F.) *Phthisis galopante ou aiguë*. Phthisis pulmonalis, which rapidly runs through its course to a fatal termination.

CONSUMPTION, PULMONARY, Phthisis pulmonalis.

CONSUMPTIVA. Same etymology. Cauties, used for the destruction of fungi. *Burnt alum, lunar caustic*, &c., were formerly so called.

CONSUMPTIVE, Phthisicus.

CONSUMPTIO, Consumption.

CONTABESCENCIA, from *contabescere*, (*con* and *tabescere*), 'to grow lean.' Consumption, marasmus, atrophy, &c.

CONTACT, *Contactus*, from *con* and *tangere, tactum*, 'to touch.' The state of two bodies that touch each other. In the theory of contagious diseases, we distinguish *immediate or direct contact*, as when we touch a patient labouring under one of those diseases; and *mediate or indirect contact*, when we touch, not the patient himself, but objects that have touched or emanated from him. The air is, most commonly, the medium by which this last kind of contact is effected.

CONTAGION, *Contagio, Contages, Contagium, Aparcheia, Apocrysis*. Same etymon. The transmission of a disease from one person to another by direct or indirect contact. The term has, also, been applied, by some, to the action of miasmata arising from dead animal or vegetable matter, hogs, fens, &c., but in this sense it is now abandoned. Contagious diseases are produced either by a virus, capable of causing them by inoculation, as in small-pox, cow-pox, hydrophobia, syphilis, &c., or by miasmata, proceeding from a sick individual, as in plague, typhus gravior, and in measles and scarlatina. [?] Scrofula, phthisis pulmonalis, and cancer, have, by some, been esteemed contagious, but apparently without foundation. Physicians are, indeed, by no means unanimous in deciding what diseases are contagious, and what not. The contagion of plague and typhus, especially of the latter, is denied by many. It seems probable, that a disease may be contagious under certain circumstances and not under others. A case of common fever, arising from *common causes*, as from cold, if the patient be kept in a close, foul situation, may be converted into a disease, capable of producing emanations, which may excite a similar

disease in those exposed to them. *Contagion and infection* are generally esteemed synonymous. Frequently, however, the former is applied to diseases not produced by contact; as measles, scarlet fever, [?] &c., whilst *infection*, (Prov. *Sæti*), is used for those that require positive contact; as itch, syphilis, &c., and conversely. Diseases, which cannot be produced in any other way than by contagion, are said to have their origin in *specific contagion*; as small-pox, cow-pox, measles, hydrophobia, syphilis, &c. Those which are produced by contagion, and yet are supposed to be sometimes owing to other causes, are said to arise from *common contagion*; as typhus, cynanche parotidea, &c.

CONTAGIONIST, (F.) *Contagioniste*. One who believes in the contagious character of a particular disease,—as of yellow fever.

CONTAGIOUS, *Contagiosus*, (F.) *Contagieux*. Capable of being transmitted by mediate or immediate contact. *Communicable*,—as a *contagious disease, contagious fever, contagious effluvia*, &c. Commonly, the epithet *infectious*, (Prov.) *Taking, Smittle, Catching*, is applied to those that are communicated by immediate contact, as itch, syphilis, &c.

CONTAGIUM, Contagion. Miasm.

CONTEMPLABILES DIES, Critical days.

CONTEMPLATIF, (F.) *Contemplatifus*, (*con* and *templum*). Appertaining to contemplation. The predominant idea of the melancholic — of the monomaniac — is sometimes called *contemplative*.

CONTEMPLATIO, Catalepsy.

CONTEMPLATIVUS, *Contemplatif*.

CONTENSIO, Tension.

CONTENTIO, Consultation.

TEXTURE, *Contextura, Contextus*, from *con* and *texere*, (*quasi tegere*, from *tegere*, 'to cover.') *textum*, 'to weave,' 'to make a web.' Arrangement of parts;—texture. A name given, metaphorically, to the structure of organized bodies; as the *texture of muscles, fibres*, &c. See *Tissue*, and *Texture*.

CONTIGUITY, DIARTHROSIS OF, see *Continuity*.

CONTINENCE, *Continentia*, from *continere*, (*con* and *tenere*, 'to hold or keep,') 'to contain oneself,' 'to restrain.' Restraint. Abstinence from, or moderation in, the pleasures of physical love.

CONTINENS, Continent.

CONTINENT, *Continens*. Restrained. This word is synonymous, also, with *Continued*; (F.) *Continu*.

CONTINENT CAUSE, *Causa conjuncta*, is a cause, real or presumed, which having given rise to a disease, continues to act during the whole of its duration. It may be considered synonymous with proximate cause. A *continent fever*, *Febris continens*, is one which preserves during its whole course, the same degree of intensity, without any remission or sensible exacerbation. A disease which continues uninterrupted, has been also called *Febrilis continua, Aciopathia* or *Aipathia*.

CONTINENTIA, Continence.

CONTINUED FEVER, *Febris continua* seu *continens* seu *anabolia* seu *assidua*. A fever which presents no interruption in its course. Continued fevers form a division in the class *Pyrexia* of Cullen, and include three genera—*Synocha, Synochus*, and *Typhus*. It is proper to remark, that some of the older writers make a distinction between the continual fever, *Febris continua*, and the *synochus* or *febris continens*. Thus, Rhases states that the *synochus* or *continens* is a fever, which consists of one paroxysm from beginning to end; whilst the *continua* is allied to intermittents.

CONTINUITY, *Continuitas*. An adhesion of two things between each other, so that they cannot be separated without fracture or laceration.

CONTINUITY, DIARTHROSES OF, (F.) *Diarthroses de Continuité*, are movable joints, in which the bones are continuous, but do not touch immediately, there being between them a ligamentous substance, whose flexibility permits motion. The vertebral articulations are examples of this. **DIARTHROSES OF CONTINUITY**, *Diarthroses de Continuité*, on the other hand, are movable articulations, in which the bones are not continuous, but touch by surfaces covered with a cartilaginous layer, which is always moistened by synovia.

CONTINUITY, LESION OF, see Continuity, Solution of.

CONTINUITY, SOLUTION OF, *Solutio contin'is*, is any division of parts, previously continuous. Wounds and fractures are *solutions of continuity*. The word *Continuity* is opposed to *Contiguity*: the latter meaning the condition of two things which are near each other, or touch without uniting. There is *contiguity* between the head of the humerus and the glenoid cavity of the scapula, but not *continuity*.

A *lesion of continuity*, *Læsio contin'is*, is a destruction of continuity caused by disease.

CONTENDING, *Contundens*.

CONTORSIO, *Contorsio*—c. Columnæ vertebralis, Rheuchistrophosis.

CONTORSION, *Contortio*, from *contorqueo*, 'con and torquere, tortum,' 'to wring,' 'I twist about.' Violent movement of a part, accompanied with a kind of torsion, twist or cast; as *contortion of the face*.

CONTRAPERTURA, from *contra*, 'against,' and *aperio*, 'I open.' A counter-opening: *Contraincisio*, *Incisio prioris opposita*, (F.) *Contreouverture*. An incision, made in the most depending part of a wound or abscess, when the first opening is not situate in a manner favourable for the discharge of the pus.

CONTRAEXTENSIO, *Counter-extension*.

CONTRACTILE, *Contractilis*, from *contra-* here, (con and trahere, tractum,) 'to draw together.' Capable of contracting. The fibre of muscles is *contractile*.

CONTRACTILITÉ, *Contractility*—c. *par Défaut d'Extension*, *Elasticity*—c. *de Tissu*, *Elasticity*.

CONTRACTILITY, *Contractilitas*: same etymon; (F.) *Contractilité*. That vital property, which gives, to certain parts, the power of contracting. The muscles of locomotion are endowed with a power of *voluntary contractility*, or one dependent immediately on the action of the brain:—the muscles of the viscera of digestion, and other internal organs, enjoy an *involuntary contractility*. *Contractility* and *irritability* are frequently used synonymously to signify the property possessed by any tissue of *contracting* on the application of an appropriate stimulus.

CONTRACTILITY, *Irritability*.

CONTRACTIO, *Contraction*—c. *Cordis*, *Systole*.

CONTRACTION, *Contractio*. Same etymon; *Systole*. Action of contracting. When we speak of the *contraction of a muscle*, we mean the phenomenon it exhibits during the time it is in action.

CONTRACTOR UTERI, *Abortive*.

CONTRACTURA. Same etymon. *Acamp'sia*, *Entesis articuli*, *Rigor artuum*, *Muscular Stiff-joint*, (F.) *Contracture*. A state of permanent rigidity and progressive atrophy of the flexor muscles, which prevents the motions of extension beyond a certain limit. The affected muscles form hard cords beneath the skin. On dissection, they are found converted into tendinous fibres, the fleshy fibres having almost disappeared, when the disease has been of any duration. It succeeds, frequently, other diseases, particularly

rheumatism, neuralgia, convulsions, syphilis, colica pictionum, &c. The warm bath, vapour bath, or thermal waters, oleaginous embrocations, mechanical extension of the limbs, &c., are the chief means of treatment.

CONTRAFISSURA, from *contra*, 'against,' and *fendere, fissum*, 'to cleave'; *Repercussio*, *Resonitus*, *Catagma Fissura contrajacens*, *Apeche-ma*, *Anticom'ma*, *Antic'ope*, *Antic'ris*, *Anticus'ma*, *Infart'ium*, *Counterstroke*, (F.) *Contre-coup*, *Contre-fente*, *Contre-fracture*. A fracture, contusion, or injury, produced, by a blow, in a part distant from that which is struck. Five species of *contrafissura* or *contre-coups* may occur in the skull. 1. When the internal table yields and fractures. 2. When the bone breaks in any other part than the one struck. 3. When a bone, which has been struck, remains uninjured, and its neighbour is fractured. 4. When the bone is fractured in a place diametrically opposite to that struck, as in fractures at the base of the cranium, from a fall on the vertex; and lastly, when the violence of the blow produces a separation of the neighbouring or distant sutures. These fractures of the skull are also called *Fractures par résonance*.

CONTRAHENTIA, *Astringents*, *Styptics*.

CONTRAINCISIO, *Contra-apertura*.

CONTRAINDICATIO, *Counter-indication*.

CONTRAIRRITATIO, *Counter-irritation*.

CONTRALUNARIS, from *contra*, 'against,' and *luna*, 'the moon.' An epithet for a woman who conceives during the menstrual discharge.—*Dietrich*.

CONTRASTIMULANS, *Contro-stimulants*.

CONTRAYERVA, *Dorstenia contrayerva*—c. *Balls*, *Pulvis contrayerve compositus*.

CONTRAYERVA, *Lissox*, *Dorstenia contrayerva*—c. *Mexican*, *Psoralea pentaphylla*—c. *Nova*, *Psoralea pentaphylla*—c. *Virginiana*, *Aristolochia serpentaria*.

CONTRECoup, *Contra-fissura*.

CONTREEXTENSION, *Counter-extension*.

CONTREFENTE, *Contra-fissura*.

CONTREFRACTURE, *Contra-fissura*.

CONTREINDICATION, *Counter-indication*.

CONTREOUVERTURE, *Contra-apertura*.

CONTREPOISON, *Antidote*.

CONTRESTIMULISME, *Controstimulus*.

CONTRESTIMULUS, *Controstimulus*.

CONTREXEVILLE, *MINERAL WATERS*

OF. *Contrexeville* is a town in France in the department of Vosges, four leagues from Neufchâteau. The waters contain carbonate of iron, chloride of calcium, and carbonate of lime, chloride of sodium, a bituminous substance, and free carbonic acid. They are frequented by those labouring under cutaneous, scrofulous, and calculous affections.

CONTRITIO, *Syntrim'ma*, *Syntrip'sis*, *Tritura*, *Trius*, *Tri'tio*, *Tripsis*, from *contero*, 'I bruise or make small,' *Comminution*, *Trituration*.

CONTROSTIMULANT, *Controstimulans*, *Hypothen'ic*, from *contra*, 'against,' and *stimulus*, 'that which excites.' A substance that possesses a particular, debilitating property, acting upon the excitability in a manner opposite to stimulus. A name given to therapeutical agents, which, according to the Italian theory of *contro-stimulus*, are endowed with the property of diminishing excitement by a specific action. These agents are by no means clearly defined.

CONTROSTIMULUS or **CONTRESTIMULUS**, *DOCTRINE OF*, (F.) *Contrestimulisme*. Same etymon. The name given by Rasori, about thirty years ago, to a new medical doctrine, of which he was the originator—*La nuova Dottrina Medica Italiana*. It is founded on the *contro-stimulant* property attributed to a certain number of

medicines. In this doctrine, as in that of Brown, under the name *excitability*, a fundamental principle of physiology is admitted, by virtue of which living beings are endowed with an aptitude for feeling the action of external agents or exciting influences, and of reacting on these influences. When this excitability is too great, there is excess of stimulus or *Hypersthenia*: when too little, there is deficiency or *Hyposthenia*. Diseases, general and local, are divided into three great classes, or into, 1. *Hypersthenic*; 2. *Hyposthenic*; 3. *Irritative*. The contro-stimulant physicians admit only two classes of medicines—*stimulants* and *contro-stimulants*.

CONTUND'ING, *Contu'sing*, *Contun'dens*, from *contundere*, (con and *tundere*, *tuum*), 'to bruise.' (F.) *Contondant*. That which causes contusions. An epithet given to round, blunt, vulnerating projectiles, which bruise or lacerate parts without cutting them.

CONTUS, *Contused*.

CONTUS, *Penis*.

CONTU'SED. Same etymon as *Contunding*. (F.) *Contus*. Affected with contusion. Thus we say—a *contused wound*.

CONTU'SION, *Contu'sio*, *Bruise*, (Prov.) *Doke*, *Collis'io*, *Phlasma*, *Thlasia*, *Thlas'ma*, *Th. contu'sio*. *Rhegē*, *Rhegma*, *Rhegmus*, *Famex*, *Famis*, *Famix*, (F.) *Meurtrissure*. Same etymon. An injury or lesion—arising from the impulse of a body with a blunt surface—which presents no loss of substance, and no apparent wound. If the skin be divided, the injury takes the name of *contused wound*. The differences of contusions, as to extent, are of course infinite. When slight, the blood stagnates in the capillaries of the skin, or is effused into the subcutaneous areolar tissue. Time and cold applications remove it. When the texture has been lacerated, there is effusion of blood, with more or less torpor in the part affected. Cooling applications, general or topical bleeding, emollients, poultices, &c., are here necessary, according to circumstances. In the severest kinds of contusion, all the soft and solid parts, except the skin, are bruised, and, at times, reduced to a kind of pap. When the disorganization is to this extent, there is no hope except in amputation. A deep contusion of the soft parts has been called *Sarcoth'lasis*, and *Sarcoth-las'ma*.

CONTUSION, *WIND*, see *Wind of a Ball*.

CONUS ARTERIO'SUS, *Infundibulum* of the heart. The portion of the right ventricle from which the pulmonary artery proceeds, forms a prominence on the right side of the anterior furrow of the heart, and is prolonged towards the left, becoming narrower at the same time, so as to form a funnel-shaped projection, which extends a little beyond the base of the ventricles. This is the *Conus arteriosus*.

CONUS VASCULOSUS, see *Coni Vasculosi*.

CONVALES'GENCE, *Convalescentia*, *Analep'sis*, *Anas'tasis*, *Reconvalescentia*, from *convalescere*, (con and *valescere*) 'to grow well.' *Excantrophē*. Recovery of health after disease. The time which elapses between the termination of a disease and complete restoration of strength.

CONVALLARIA ANGULOSA, *C. polygonatum*.

CONVALLARIA CANALICULA'TA, *Polygo'natum*, *canaliculatum* seu *latifolium*, *Great Solomon's Seal*; and **CONVALLARIA PUBES'CENTS**, *Polygo'natum pubescens*, *Smaller Solomon's Seal*: indigenous. A decoction of the roots is said to be cathartic, diuretic, and diaphoretic.

CONVALLA'RIA MAIA'LIS, from *convallis*, 'a valley,'—from its abounding in valleys. *Lil'ium Convall'ium*, *Convalla'ria*, *C. mappi*, *Maion'themum*, *Lily of the Valley*, *May Lily*, (F.) *Muguet*,

Muguet de Mai. The recent flowers are reputed to be aromatic, cephalic, errhine, and cathartic. They are not used.

CONVALLARIA MAPPI, *C. Maialis*.

CONVALLA'RIA MULTIFLO'RA, *Polygo'natum multiflorum*, which grows in this country and in Europe, has analogous properties.

CONVALLA'RIA POLYGO'NATUM. The systematic name of *Solomon's Seal*, *Convalla'ria angulosa*, *Polygo'natum uniflorum* seu *anceps* seu *culga'rē*, *Sigil'ium Salomo'nis*, *Polygo'natum*, (F.) *Sceau de Salomon*, *Genouillet*. The root was once used as an astringent and tonic. It is, also, a reputed cosmetic.

CONVENTUS, *Coition*.

CONVER'SION, *Conver'sio*, from *con* and *vertere*, *verum*, 'to turn.' Change from one state into another.

CONVERSION OF DISEASES, (F.) *Conversion des maladies*, is the change or transformation of one disease into another.

CONVOLUTION, *Convolutio*, from *convolvere*, (con and *volvere*, *volutum*) 'to entwine;' *Epiphallion*, *Gyrus*, *Helig'mus*, (F.) *Circonvolution*. The rolling of any thing upon itself.

CONVOLUTIONS, *Cer'eb'ral*, *Gyri* seu *Plicat'ura* seu *Spiræ* seu *Processus entero'dei* seu *Intest'icula Cer'ebri*, are the round, undulating, tortuous projections observed at the base of the brain. In them Gall's organs, of course, terminate.

CONVOLUTION, *INTER'NAL*, *C. of the Corpus Callo'sum*, *Convolution d'Ourlet* (Foville). A cerebral convolution of great extent, the principal portion of which is found on the inner surface of each hemisphere above the corpus callosum. In front it bends downwards and backwards to the fissure of Sylvius, and behind, it extends to the middle lobe and forms the hippocampus major.

CONVOLUTIONS, *INTES'tINAL*, are the turns made by the intestines in the abdomen.

CONVOLUTION, *SUPRA-OR'BITAR*. A convolution of the brain, which exists on the inferior surface of the anterior lobe, and rests upon the roof of the orbit.

CONVOLVULUS, from *convolvere*, (con and *volvere*) 'to entwine.' *Ileus*, *Intussusceptio*.

CONVOLVULUS BATA'TAS. Same etymon. *C. Indicus*, *Camotes*; the *Sweet Potato*, *Spanish Potato*. (F.) *Potato*. *Ord. Convolvulaceæ*. This is the only esculent root of the genus *Convolvulus*. It is much eaten in the United States.

CONVOLVULUS CANTAB'RICA, *Cantab'rica*, *Lavender-leaved Bindweed*, has been considered antelmintic and actively cathartic.

CONVOLVULUS INDICUS, *C. Batatas*.

CONVOLVULUS JALA'PA. A systematic name, generally thought to be the *Jalap* plant, which is, properly, *Erogo'nium purga*, *Ipome'a Jalapa* seu *purga* seu *Schiedeana* seu *macrorrhiza*, *Jala'pium*, *Jala'pa*, *Mechoacan'a nigra*, *Jalapapa*, *Jalapa*, *Jalap Root*, *Gialappa*, *Xolappi*, *Bryo'nia Mechoacan'a nigricans*, *Bryo'nia Peruviana*, *Chela'pa*, *Rhabar'barum Nigrum*, *Gelap'pium*, (F.) *Jalap*, is produced from South America. Its odour is nauseous; taste sweetish and slightly pungent. It is solid, hard, heavy, brittle; fracture, resinous; internally, light gray; externally, covered with a deep brown, wrinkled bark. Its operation is cathartic, the resinous part griping violently. Dose, 10 gr. to ʒss. A drop or two of any essential oil may prevent it from griping. An active principle has been separated from *Jalap*, to which the names *Jalapin* and *Cathartin* have been given.

An inferior kind, called *light*, *fusiform* or *male Jalap*, is yielded by *Ipome'a* seu *Convolvulus Orizabensis*, *I. batato'des* seu *Mestitlan'ica*, which grows near Orizaba and Mestitlan, in Mexico.

CONVOLVULUS MAJOR ALBUS, *Convolvulus sepium*—c. *Maritimus*, *Convolvulus soldanella*.

CONVOLVULUS MECOCHACAN, *Mechoaca'na Radix*, *Jalap'pa alba*, *Rhabar'barum album*, *Piptostegia Piao'nis*, *Mechoacan*, *Jeticucu*, (F.) *Rhabar'ba* ou *Rhubarbe blanche* ou *des Indes*, *Jalap blanc*, *Bryone d'Amérique*. A Mexican convolvulus, the root of which possesses aperient properties, and was once extensively used instead of jalap. See *Batata de purga*.

CONVOLVULUS MEGALORHIZUS, c. *Panduratus*.

CONVOLVULUS NIL, *Ipomœa nil*, *Blue Morning-glory*. An East Indian plant, naturalized in some of the Western States. The seeds are sold in Calcutta as a cathartic. The roots have the same properties.

CONVOLVULUS OPERCULATUS, see *Batata de purga*—c. *Orizabensis*, see *Convolvulus Jalapa*.

CONVOLVULUS PANDURATUS, c. *Megalorhizus*, *Pseudo-mechoaca'na*, *Fiddle-leaved Bindweed*, *Hog Potato*, *Virginian Bindweed*, *Wild Potato*, *Mech'ameck*, *M. Bindweed*, *Wild Jalap*, *Man in the ground*, *Man of the Earth*, *Wild Rhubarb*, *Wild Potato-Vine*, *Kassau'nder*, *Kassau'nder*, *Kussau'nder*, (F.) *Liseron Mechacane*. In Virginia, and some other parts of the United States, the root of this plant has been much recommended in cases of gravel. It is used either in powder or decoction.

CONVOLVULUS PERENNIS, *Humulus lupulus*.

CONVOLVULUS PES CA'PRE, *Bargada*. A plant used in India as a cataplasm in arthritic cases.

CONVOLVULUS REPENS, c. *sepium*.

CONVOLVULUS SCAMMONEA seu SCAMMONEIA, c. *Syriacus*. The systematic name of the *Scammony Plant*. A Syrian and Mexican plant; the concrete gummi-resinous juice of which, *Scammonia*, *Scammonia Gummi Lesi'na*, *Scammonium*, *S. Syriacum*, *Diagryd'ium*, *Dacrydion*, *Scammony*, *Mahmoudy*, (F.) *Scammonée d'Alep*, comes to us in blackish-gray fragments, becoming whitish-yellow when touched with wet fingers. It is a drastic, hydragogue cathartic. Dose, gr. iij to gr. xv, triturated with sugar.

CONVOLVULUS SEA, *Convolvulus Soldanella*.

CONVOLVULUS SEPIUM, *Convolvulus major albus* seu *repens* seu *Tuguriurum*, *Calyste'gia sepium*, *Great* or *Hedge Bindweed*, (F.) *Liseron des Haies*, *Grand Liseron*. The juice of this plant is violently purgative, and is given in drop-sical affections.

CONVOLVULUS SOLDANEL'LA. The systematic name of the *Sea Convolvulus*, *Convolvulus Maritimus*, *Bras'tica Mari'na*, *Calyste'gia Soldanella*, *Sea Bindweed*, *Soldanel'la*, (F.) *Chou Marin*. The leaves of this plant are said to be drastic cathartic; but they are not much, if at all, used.

CONVOLVULUS SYRIACUS, *Convolvulus scammonia*—c. *Tuguriurum*, c. *sepium*.

CONVOLVULUS TURPETHUM, *Turpethum*. The systematic name of the *Turbith plant*. (F.) *Racine de Turbith*. The cortical part of the root of this species of convolvulus is brought from the East Indies. It is a cathartic, but not used.

CONVULSÉ, *Convulsed*.

CONVULSED, *Convulsus*, (F.) *Convulsé*. Same etymon as *convulsion*. Affected with convulsions.

CONVULSIBILITAS, see *Subsultus tendinum*.

CONVULSIF, *Convulsive*.

CONVULSIO, *Convulsion*—c. *Canina*, see *Canine laugh*.

CONVULSIO CEREALIS, *Convulsio* ab *Ustilaginæ* seu *Solentis'is*, *Ergotism* seu *spasmodicus*, *Myrmec'asia*, *Myrmecismus*, *Myrmecismus*, *Myrmecosis*, *Cereal Convulsion*, (F.) *Convulsion céréale*, *Ergotisme convulsif*, *Convulsion de Sologne*. A singular disorder of the convulsive

kind, attended with a peculiar tingling and formation in the arms and legs; hence called by the Germans *Kriebelkrankheit*. It is said to be endemic in some parts of Germany, and to arise often from the use of spoiled corn. See *Raphania*.

CONVULSIO HABITUALIS, *Chorea*—c. *Indica*, *Tetanus*—c. *Raphania*, *Raphania*—c. *Soloniensis*, c. *cerealis*, *Ergotism*—c. *Tonica*, *Tonic Spasm*—c. *Uteri*, *Abortion*.

CONVULSION, *Spasmus*, *Convulsio*, from *convellere*, (con and *vellere*, *vulsum*), 'to tear, 'to pull together;' *Conduc'tio*, *Hieran'osus*, *Disten'tio nervo'rum*, *Spasmus clon'icus*, *Convulsio clon'ica*, *Eclamps'ia*, *Syspa'ia Convulsio*, *Hyperspasm'ia*, *Clonic Spasm*. This word has several acceptations. It means any violent perversion of the animal movements. The word *Convulsions* generally, however, signifies alternate contractions, violent and involuntary, of muscles, which habitually contract only under the influence of the will. This alternate contraction, when slight, is called *tremor*; when strong and permanent, *tetanus*, *trismus*, &c. *Spasms*, *Cramp*, *Risus Sardonius*, and *St. Vitus's Dance* are *convulsions*.

CONVULSION, SALAAM', *Eclamps'ia* seu *Spasmus nutans*. A name given to a singular kind of convulsion in children, in which there is a peculiar bobbing of the head forward.—Sir Charles Clarke.

CONVULSION DE SOLOGNE, *Convulsio cerealis*.

CONVULSIONNAIRE (F.). A name given, during the last century, to individuals who had, or affected to have, convulsions, produced by religious impulses. The epithet was first given to fanatics, who exhibited the most wonderful and varied seizures, at the tomb of Paris, a Jansenist, who died in the year 1727, and was buried in the cemetery of St. Médard. Some of these were called *Sautseuses* or *Jumpers*, of whom there have been examples in more recent times: others, from barking like a dog, were called *aboyseuses*; and others, from mewing like a cat, *miaulantes*.

A form of convulsion, induced by religious frenzy, has been vulgarly called the "jerke," and the gymnastic movements of the *Shakers* may be placed in the same category.

CONVULSIONS OF CHILDREN, *Eclampsia*—c. *des Enfants*, *Eclampsia*—c. *des Femmes enceintes* et *en couches*, *Eclampsia gravidarum* et *parturientium*—c. *Puerperal*, *Eclampsia gravidarum* et *parturientium*. See *Mania*, *dancing*.

CONVULSIVE, *Convulsivus*, *Spasmodes*, *Agitato'rius*, (F.) *Convulsif*. That which is accompanied by, or is analogous to, convulsions, as *convulsive cough*, *convulsive disease*.

CONYZA, *Inula dysenterica*—c. *Bifoliata*, *Serico'carpus tortifolius*—c. *Coma aurea*, *Solidago Virgaurea*—c. *Major*, c. *squarrosa*—c. *Media*, *Inula dysenterica*—c. *Pycnostachya*, *Pterocaulon pycnostachyum*.

CONYZA SQUARRO'SA, c. *Major*, *Brepho'tonon*, *In'ula squarro'sa*, *Great Fleabane* or *Spikenard*, (F.) *Herbe aux mouches*. Ord. *Compositæ*. A European plant, whose strong and disagreeable odour was formerly considered to be destructive to flies, fleas, &c. Its infusion in wine was once used as an emmenagogue and anti-icteric; and in vinegar as an anti-epileptic.

COOKERY, *Culinary art*.

COOLWEED, *Pilea pumila*.

COONTIE or COONTI, see *Arrowroot*.

COOPER'S WELL, see *Mississippi*, *Mineral Waters* of.

COOPERCULUM OCULI, *Palpebra*.

COOPERTORIUM, *Thyroid cartilage*.

COOSTRUM. The middle part of the diaphragm.—Ruland.

COOT, Malleolus.

COOTH, Catarrh.

COPAITER, Copaiba.

COPAHU, Copaiba.

COPAIBA. The resinous juice of *Copaifera officinalis* seu *Jacquinii*, *Copaiba officinalis*, and other species of *Copaifera*; (F.) *Copehiet*. Ord. Leguminosæ. *Secr. Syst.* Decandria Monogynia. It is the *Copaifera officinalis* *Rosina*, *Ralsmanni* *Copaiba* seu *Brazilien* seu *de Copahú* seu *Copivai*, *Balsam of Copaiba* or *Copaiba*, (vulgarily pronounced *capayba*.) (F.) *Copahu*, *Bois de Copahu*, *B. du Brésil*, *Térébinthe de Copahu*. Its odour is peculiar, but not unpleasant; taste pungent, bitter; consistence, syrupy; colour yellowish, and transparent. It is soluble in alcohol, ether, and the expressed oils. S. g. 0.950. Its properties are stimulant and diuretic; in large doses it is purgative. It acts on the lining membrane of the urethra, and on mucous membranes in general. It is given in gonorrhœa, gleet, leucorrhœa, &c., in the dose of gr. x to ʒi, twice or thrice a day. Should symptoms of urticaria or diarrhœa arise, the dose is too large, and must be diminished. It can be inspissated by means of magnesia so as to be made into pills, and a plan has been devised for enveloping it in gelatin, so that its taste is entirely concealed. See Capsules, gelatinous.

COPAIFERA JACQUINI, see *Copaiba*—c. *Officinalis*, see *Copaiba*.

COPAIVA OFFICINALIS, see *Copaiba*.

COPAL, *Copale*, *Rosina Copal*, *Gummi copal*, *Uatum*. A resinous substance brought from the East Indies, South America, and the western coast of Africa, which flows spontaneously from *Elaeocarpus Copalliferus*, *Vateria Indica*, and probably from different species of *Hymenæa*. Ord. Leguminosæ. It is a stimulant like all the resins, and, dissolved in rectified spirits of wine, has been used in cases of spongy gums and looseness of the teeth; but it is now only employed in varnishes.

COPALCHE BARK, see *Croton pseudo-china*.

COP-BONE, Patella.

COPE, Cut.

COPHOMA, Cophosis.

COPHONIA, Acouphonia.

COPHOSIS, *Cophoma*, *Cophotes*, *Surditas*, *Paracosis*, *Dysacosis* *auditoria*; from *κοπος*, 'deaf.' (F.) *Surdité*. Diminution or loss of hearing. Cullen uses the word synonymously with *Dyscœcia*, and Pinel with *Paracosis* or *Paracœd*. According to Sauvages, *Cophosis* differs from both,—from *Dyscœcia*, because in it the sonorous rays cannot reach the labyrinth; and from *Paracosis*, which consists in a confused state of hearing.

Cophosis seems, usually, to be synonymous with deafness,—*Paracosis*, *Surditas*.

COPHOTES, Cophosis.

COPIOPIA, Copyopia.

COPOS, *Lassitud*, *Fatigatio*, *Lassitudo*, *Fatigue*. A state of body in which most of the animal functions are exerted with less promptitude and vigour than common. The ancients admitted three species: 1. That arising from plethora, *Lassitud tensiva*, *tonodes*; 2. From plethora and increased heat combined, *Lassitud phlegmonosa*, *astuosa*, *phlegmonodes*; and 3. Owing to a morbid condition of the humours, *Lassitud nigra*, *heleodes*.

COPPER, Cuprum—c. Ammoniated, Cuprum ammoniatum—c. Ammonio-sulphate of, Cuprum ammoniatum—c. Rose, Papaver rheas—c. Subacetate of, Cupri subacetatas—c. Sulphate of, Cupri sulphas—c. Ammoniacal sulphate of, Cuprum ammoniatum.

COPPERAS, Ferri sulphas—c. White, Zinci sulphas.

COPPER-NOSE, Gutta rosea.

COPRACRASIA, Scoracrasia.

COPRACRATIA, Scoracrasia.

COPRAGOGUM, from *κοπος*, 'fæces,' and *αγω*, 'I bring away.' *Stercus æ primis rictu obdæna*. A cathartic. The name of a laxative electuary, mentioned by Ruland.

COPRECRITICUS, Coprocriticus.

COPREMESIS, *Copriemesis*, *Vomitus feculentus* seu *stercoris*: same etymon as the next. Vomiting of fæces.

COPREMETUS, *Coproemetus*, *Copriemetus*, *Mordicimus*, from *κοπος*, 'excrement,' and *αγω*, 'I vomit.' One who vomits fæces.—Hippocrates.

COPRIEMESIS, Copremesis.

COPRIEMETUS, Copremetus.

COPROCRITICUS, *Coproecriticus*, from *κοπος*, 'excrement,' and *χωμα*, 'I separate.' A mild cathartic; an ecoprotic.

COPROEMETUS, Copremetus.

COPROEPTICUS, Constipation.

COPROLITHUS, see *Calculi* of the stomach and intestines.

COPROPHORESIS, Catharsis.

COPROPHORIA, Catharsis.

COPORRHŒA, Diarrhœa.

COPROS, Excrement.

COPROSCLEROMA, Coprosclerosis.

COPROSCLEROSIS, from *κοπος*, 'excrement,' and *σκληρω*, 'I harden.' Induration of fecal matters: *Coproscleroma*.

COPROSTASIS, Constipation.

COPTÈ, *Copton*, from *κοπω*, 'I beat or pound.' A sort of cake, composed of vegetable substances, which the ancients administered internally, and applied to the epigastric region in the form of cataplasm.

COPTIS, *C. trifolia*, *Nigella*, *Helleborus trifolius*, *Fibra aërea*, *Chrysa fibrata*, *Anemone Grœlandica*, *Gold thread*, *Manch root*: from *κοπω*, 'to cut,' in reference to the divided leaves. The root of this evergreen—*Coptis*, (Ph. U. S.)—is much used in Massachusetts, in aphthous and other ulcerations of the mouth, as a local application. It is a pure bitter, and can be used wherever such is indicated.

COPTIS TEETA, *Mishme Teeta* (Upper Assam), *Honglane* (Chinese). The root of this plant is considered to be a powerful tonic and stomachic.

COPTON, Copte.

COPULA, Ligament—c. Carnalis, Coition—c. Cartilaginea, see *Synchondrosis*—c. Magna cerebri, Corpus callosum.

COPULATION, Coition.

COPYOPIA, *Copio'pia*, *Kopyo'pia*, *Kopio'pia*, *Ophthalmocopia*, (F.) *Lassitudo oculorum*, from *κοπος*, 'fatigue,' and *ωπ*, 'the eye,' *Fatigue of vision*. Weakness of sight. Inability of the eye to sustain continued exertion.

COQ, Phasianus Gallus—c. *des Jardins*, *Tanacetum*.

COQUE DU LEVANT. see *Menispermum coeculus*—c. *d'Enf*, see *Ovum*—c. *Levand*, see *Menispermum coeculus*.

COQUELICOT, Papaver rheas.

COQUELOURDE, Anemone pulsatilla.

COQUELUCHE, Influenza, Pertussis.

COQUENTIA MEDICAMENTA, from *coquere*, 'to digest.' Medicines which were formerly believed to be proper for favouring the coction or digestion of food.

COQUERET, Physalis.

COQUETTE, Influenza.

COQUILLE, Cochlea—c. *d'Enf*, see *Ovum*.

COR, a corn. Also, Heart—c. Aorticum, see *Heart*—c. Arteriosum, see *Heart*—c. Bovinum,

Heart, hypertrophy of the—c. *Hirsutum*, *Cor villosum*—c. *Hispidum*, *Cor villosum*—c. *Pulmonale*, see Heart—c. *Taurinum*, Heart, hypertrophy of the—c. *Tomentosum*, *Cor villosum*—c. *Venosum*, see Heart.

COR VILLOSUM, *C. tomentosum* seu *hirsutum* seu *hispidum*. The condition of the heart, at times, after pericarditis, when it is studded with projections of fibrinous matter from its surface.

COR'ACO. In composition, appertaining or relating to the coracoid process.

CORACOBRACHIÆUS, *Coracobrachialis*.

COR'ACO-BRACHIAL, *Cor'aco-Brachialis*. Belonging both to the coracoid process and arm.

CORACO-BRACHIALIS (Muscle), *Coracobrachiæus*, *Coraco-humeral*—(Ch.), *Perforatus Casse-rii*, *Perforatus*, is situated at the inner and upper part of the arm. It arises from the coracoid process of the scapula, and is inserted at the middle part of the inner side of the humerus. It carries the arm forwards and inwards, raising the humerus a little. It can, also, by acting inversely, depress the shoulder.

CORACO-CLAVICULAR, *Coraco-clavicu- laris*. Belonging to the coracoid process and clavicle.

CORACO CLAVICULAR LIGAMENT—called, also, *Omo-clavicular*—serves to unite the clavicle to the coracoid process. It is very irregular, and formed of two fasciculi, which the greater part of anatomists have described as particular ligaments, under the names *Conoid* and *Trapezoid*.

CORACODES, *Coracoid*.

CORACO-HUMERALIS, *Coraco-brachialis*—c. *Hyoideus*, *Omo-hyoideus*—c. *Radialis*, *Biceps flexor cubiti*.

COR'ACOID, *Coracoïdeus*, *Coracoïdes*, *Coracoïdes*, from *καρὰ*, *καρπος*, 'a crow,' and *ειδός*, 'resemblance.' (F.) *Coracoïde*. Resembling the beak of a crow. A name given by Galen—and still retained—to the short, thick process, situated at the anterior part of the upper margin of the scapula, which has some resemblance to the beak of a crow. *Processus Coracoïdeus* seu *uncinatus* seu *corniculatus* seu *anchoralis* seu *rostriformis* seu *angryoides*, *Crow's beak-like process*. This process gives attachment to the *Coraco-clavicular* and *Coraco-acromial* ligaments, and to the *Coraco-brachialis*, *Pectoralis minor*, and *Biceps* muscles.

CORACOID LIGAMENT, *Ligamentum coracoïdeum*, (F.) *L. Coracoïdien*. This name is given to the small fibrous fasciculus, which converts the notch, at the superior margin of the scapula, into a foramen.

CORAIL, *Coral*—c. *des Jardins*, *Capsicum annum*.

CORAL, *Corallium*, *Corallus*, *Courea'llium*, *Arbor Maris*, *Azur*, *Bole'sis*, from *κορσ*, 'I adorn,' and 'als,' 'the sea;' and, according to some, from *κορη*, 'a daughter,' and 'als,' 'the sea.' (F.) *Corail*. One of the most beautiful productions of the deep. It is fixed to submarine rocks, in the form of a shrub, and is of a bright red colour. It is the habitation of a multitude of animals, of the Zoophyta order, and is formed of a calcareous substance, secreted by the animals themselves. It is in very hard, concentric layers; covered, externally, by a species of porous bark, full of cells, each of which contains one of these animals. Linnæus calls the red coral, *Isis nobilis*, and M. de Lamarck, *Corallium rubrum*. It is much fished for on the coasts of Barbary and Sicily. Coral was formerly esteemed tonic, absorbent, astringent, &c.; but analysis has shown, that it contains only carbonate of lime and a little gelatin.—Dioscorides, Pliny, Oribasius, the *Geoponica*, &c. *Corallium album* is a hard, white, calcareous, brittle substance, the nidus of

the *Madrep'ora oculata*. It has been given as an absorbent.

CORALLINA. Diminutive of *Corallium*: *Muscus muritinus*, *Corallina officinalis*, *Lichen*, *Corallina alba*, *Seu Cor'alline*, *White Worm-weed*, (F.) *Coralline blanche*. The production of an animal, which belongs to the class *Pol'yp't*, and which is found in all the seas of Europe;—particularly in the Mediterranean. It has the appearance of a plant, is homogeneous, an inch or two in height, of a white, reddish, or greenish colour, salt taste, and marine smell. It contains gelatin, albumen, chloride of sodium, phosphate, carbonate, and sulphate of lime, carbonate of magnesia, silica, oxide of iron, and a colouring principle. It was once much used as a vermifuge; but is not now employed. Dose, ʒss to ʒj, in powder.

CORALLINA CORSICANA, *C. rubra*, *Helminthochorton*, *Helminthochortum*, *Elminthochorton*, *Muscus helminthochortos*, *Melithochorton*, *Muscus corallinus* seu *Marinus* seu *Cor'sicus*, *Conferva Helminthochortos*, *Spharococcus helminthochortos*, *Gigartina* seu *Plocaria* seu *Fucus helminthochorton*, *Coralli'na melitochorton*, *Lemithochorton*, *Ceramium helminthochortus*, *Corsican Worm-weed*, (F.) *Coralline de Corse ou noire*, *Mousse de Corse*. It is a mixture of several marine plants and zoophytes, as the *fucus*, *ceramium*, *ulva*, *coralline*, *conferva*, &c., and has gained great reputation for destroying all kinds of intestinal worms, when given in strong decoction. The Geneva Pharmacopœia directs an official syrup—the *Sirap de Coralline*.

CORALLINE BLANCHE, *Corallina*—c. *de Corse*, *Corallina Corsicana*—c. *Noire*, *Corallina Corsicana*.

CORALLOIDES FUNGUS, *Clavaria coral- loides*.

CORALWORT, *Clavaria coralloides*.

CORCULUM GERMINIS, see *Moleculæ*.

CORD, *Funis*, *Funiculus*, *Chanos*, *Cha'nion*, *χοινος*, *χοινιον*, from the Latin *Chorda*, which is itself derived from *χορδή*, 'intestine;' and, afterwards, was applied to musical cords or strings, made of the intestines of animals. See *Chorda*.

CORD, **SPERMATIC**, see *Spermatic cord*—c. *Umbilical*, *Funiculus umbilicalis*.

CORDS, **VOCAL**, *Cords of Ferrein*, *Chordæ vocales*, *Ch. Ferre'nii*. A name given to the ligaments of the glottis, which Ferrein compared to stretched cords, and to which he attributed the production of voice. See *Thyreo-arytenoid Ligament*.

CORDA, *Chorda*—c. *Hippocraticis*, *Achillis tendo*—c. *Magna*, *Achillis tendo*—c. *Spinalis*, *Medulla spinalis*.

CORDE DU TAMBOUR, *Chorda tympani*—c. *du Tympan*, *Chorda tympani*.

CORDEE, *Chordee*.

CORDIA AFRICANA, *Sebestina*—c. *Domes- tica*, *Sebestina*—c. *Myxa*, *Sebestina*—c. *Obliqua*, *Sebestina*—c. *Sebestina*, *Sebestina*—c. *Smooth- leaved*, *Sebestina*.

CORDIAL, *Cordia'lis*, *Cardi'acus*, from *cor*, *cordis*, 'the heart.' A tonic or excitant medicine, judged to be proper for exciting the heart. A warm stomachic.

CORDIAL, **GODFREY'S**, see *Godfrey's Cordial*—c. *Nervous*, *Brodum's*, see *Tinctura gentianæ composita*—c. *Sir Walter Raleigh's*, *Confectio aromatica*—c. *Warner's*, see *Tinctura rhei et sennæ*.

CORDIFORM TENDON OF THE DIA- PHRAGM, *Centre*, *phrenic*.

CORDINE'MA, from *kapa*, 'the head,' [?] and *divew*, 'I move about.' Headache, accompanied with vertigo. See *Carebaria*.

CORDIS EMUNCTORIUM, *Axilla*.

CORDOLIUM, *Cardialgia*.

CORDON (F.), *Funiculus*, diminutive of *funis*, 'a cord.' A term applied to many parts, which resemble a small cord.

CORDON NERVEUX, *Ramus Nervo-us*. A principal division of a nerve, or the nervous trunk itself.

CORDON OMBILICALE, *Funiculus umbilicalis*—*c. Spermaticus*, Spermatic chord—*c. Testiculaire*, Spermatic chord.

CORDONS SUS-PUBIENS, Round ligaments of the uterus—*c. Vasculares*, Round ligaments of the uterus.

CORDS, see **Cord**.

CORE, Pupil, see **Furunculus**.

COREC'LISIS, *Ankylos'orē*, *Atretop'ia*, from *κορη*, 'pupil,' and *κλειω*, 'I close.' Closure or obliteration of the pupil.

CORECTOMIA, see **Coretomia**.

CORECTOP'IA, from *κορη*, 'the pupil,' *εκ*, 'out of,' and *τοπος*, 'place.' Displacement of the pupil. A condition of the iris in which one segment is larger than the other; so that the pupil is not in the centre.—Von Ammon.

COREDIALYSIS, see **Coretomia**.

CORE'MATA, from *κορεω*, 'I cleanse.' Remedies proper for cleansing the skin.—Paulus of Ægina.

COREMETAMORPHOSIS, *Dyscoria*.

COREMORPHO'MA, same etymon as the next. A morbid change in the shape of the pupil.

COREMORPHO'SIS, *Conforma'tio pupill'æ artificia'lis*, *Cor'eplasty*, from *κορη*, 'the pupil,' and *μορφωσις*, 'formation.' The operation for artificial pupil. See **Coretomia**.

CORENCLEP'SIS, from *κορη*, 'the pupil,' and *εγκλειω*, 'to include.' Operation for artificial pupil, which consists in drawing out, through an incision in the cornea, a portion of the iris, and cutting it off.

COREON'CION, *Coron'cion*, *Coreon'cium*, from *κορη*, 'the pupil,' and *ογκις*, 'a small hook.' An instrument, used for the formation of artificial pupil by Langenbeck. It is hooked at its extremity. A double-hooked forceps, used by Von Gräfe, is similarly named.

COREOP'SIS TRICHOSPER'MA, *Tickweed sunflower*, *Tickweed sunflower*; from *κορις*, 'the bug,' and *οψις*, 'resemblance.' An indigenous plant, of the Composite Family, with large golden yellow rays, which flowers in September. It is said to have been used as an alternative.

COREPLASTY, *Coremorphosis*.

CORETODIALYSIS, see **Coretomia**.

CORETOMEDIALYSIS, see **Coretodialysis**.

CORETOM'IA, from *κορη*, 'the pupil,' and *τεμνω*, 'to cut.' *Iridotom'ia*, *Coretotom'ia*, *Corotom'ia*, *Iridectom'ia*, *Coretonectom'ia*, *Corectom'ia*, *Iridodial'yeis*, *Corodial'yeis*, *Corodial'yeis*, *Coretodial'yeis*, *Coretomedial'yeis*, *Iridotomedial'yeis*, *Iridectomedial'yeis*. Various operations for the formation of artificial pupil are so termed. The first four signify the simple incision of the iris for that purpose; the next three, excision of the iris, or incision with loss of substance; the next four the simple separation or tearing asunder of the iris from the ciliary ligament; the next two such separation with incision of the iris; and the last such separation with excision of the part of the iris detached. When a portion of the iris is left strangulated in the wound, it is termed *Iridencle'isia*, *Iridencle'ismus*, and *Iridotencle'isia*.

CORETONECTOMIA, see **Coretomia**.

CORETOTOMIA, *Coretomia*.

CORIAN'DER, from *κορις*, 'the bug:' the smell of the fresh herb resembling that of the bug. [?] *Coriandrum sativum*.

CORIANDRUM CICUTA, *Cicuta virosa*—*c. Maculatum*, *Conium maculatum*.

CORIAN'DRUM SATIVUM, *Corian'der*, *Corian'*

non, (F.) *Coriandre*. Family, *Umbelliferae*. *Ses. Syst.* *Pentandria Digynia*. The seeds of the coriander—*Corian'drum* (Ph. U.S.), *Coriannon*—have an aromatic odour, and grateful, pungent taste. They are carminative, but are chiefly used to cover the taste of other medicines.

CORIANNON, *Coriandrum sativum*.

CORIGEEEN, *Fucus crispus*.

CORIITIS, *Cytitis*.

CORINTHIACÆ, see *Vitis Corinthiaca*.

CORION, *Corium*, *Hypericum perforatum*.

CORIS, *Cimex*—*c. Mospeliensis*, *Symphytum Petraeum*.

CO'RIMUM, *Corion*, (F.) *Cuir*. The skin of animals is so called, especially when tanned. The *cutis vera*, or the thickest part of the human skin.

CORIUM PHLOGIS'TICUM, *Crusta pleuret'ica seu inflammato'ria seu phlogis'tica*, *Pegmin*, *Inflam'matory Crust* or *Buff*, *Buffy Coat*, (F.) *Couenne*, *C. Pleurétique*, *C. Inflammatoire*. The grayish crust or buff, varying in thickness, observed on blood drawn from a vein during the existence of violent inflammation, pregnancy, &c. It is particularly manifest in pleurisy, and hence one of its names. For its production, it appears to be requisite, that there should be an increase in the proportion of the fibrinous element of the blood over that of the red corpuscles, with increased aggregation of those corpuscles. Under such circumstances, the buffy coat assumes a concave appearance on its upper surface, and the blood is, therefore, said to be *cupped*.

The buff is generally believed to consist of fibrin; but, according to the researches of Mulder, it is composed of a binoxide of protein, which is insoluble in boiling water, and a tritoxide which is soluble. These oxides are comprehended by him under the name *oxyprotein*.

When the blood presents the above appearance, it is said to be *buffy*.

CORK, *Suber*.

CORMIER, *Sorbus domestica*.

CORMUS. In botany, when the stem of a plant, without creeping or rooting, is distended under ground, retaining a round or oval form, it is so called. The *Cormus* is vulgarly termed a root,—*radix*.

CORN, (Saxon *corn*), (G.) *Kern*. In England, this word means the *Cerealia*, or those seeds, which grow in ears, not in pods. In the United States, *Corn* always means *Indian Corn*. Its English sense corresponds to the French *Blé* or *Bled*.

CORN, *Clavus*, *Clavus Pedis*, *Ecphy'ma Clavus*, *Gemur'sa*, (F.) *Cor*, *Ognon*, from *cornu*, 'a horn.' A small, hard, corneous tumour, which forms upon the foot, generally on the toes; and is commonly produced on the most projecting parts, by the pressure of too tight shoes. A part of the corn is raised above the skin, and forms a round tumour, like the head of a nail: the other portion, which serves as its base, is buried more or less deeply in the integuments, and occasionally extends as far as the tendons and periosteum. Corns may, sometimes, be removed, by immersing the feet in warm water, but commonly they return. They can, likewise, be destroyed by the knife or caustic, or by paring them down and pulling them out by the roots; but these operations are not always as simple as they seem. In the way of palliation, they must be constantly pared; and, for the purpose of preventing pressure, any soft plaster, spread upon linen or leather, may be applied, with a hole in the centre to receive the corn; and layer after layer of plaster be added, until they attain the level of the corn. When very irritable, the lunar caustic, rubbed over the surface, will generally diminish irritability.

lity surprisingly, and in a mode not easy of explanation.

CORN, Zea mays.

CORN COCKLE, *Lychnis githago*.

CORN, GUINEA, *Panicum Italicum*—c. Indian, Zea mays—o. Wild, *Matricaria chamomilla*.

CORN FLAG, *Gladiolus vulgaris*—c. Flower, *Centaurea cyaneus*—c. Salad, *Valeriana dentata*.

CORN, SQUIRREL, *Dicentra Canadensis*.

CORNALINE, Cornelian.

CORNE, Cornu—c. d'Ammon, Cornu ammonis—c. de Béliet, Cornu ammonis—c. de Cerf, Cervus, Cornu cervi, Cochlearia coronopus—c. de Chamois, Cornu rupicaprae—c. Descendante ou inférieure du ventricule latéral, Cornu descendens ventriculi lateralis—c. de la Peau, Cornu—c. Postérieure du ventricule latéral, Cornu posterius seu posticum ventriculi lateralis.

CORNÉ, Corneous.

CORNEA, *C. transparentis* seu *pellucida* seu *lucida*, Ceras, *Scleroflaca ceratoides*, *Ceratoides* seu *Ceratoides membrana*, *Ceratome'ninx*, *Membrana cornea*; from cornu, 'horn.' The transparent cornea. (F.) Cornée. One of the coats of the eye, so called because it has some resemblance to horn. It is termed transparent to distinguish it from the opaque—*Cornea opaca* or *Sclerotic*. It is convex, anteriorly; concave, posteriorly; forming nearly one-fifth of the anterior part of the eye, and representing a segment of a sphere about seven lines and a half, or in. 0.625 in diameter. It seems to be constituted of laminae in superposition, but of the precise number anatomists are not agreed. Henle assigns it four; the third, a very solid cartilaginous lamella, being called *Membrane de Demours* or *M. de Descemet*: see *Aqueous humour*. Messrs. Todd and Bowman assign it five layers.

CORNEA, CONICAL, Staphyloma of the cornea—c. Opake, Caligo—c. Opaca, Sclerotic—c. Sugar-loaf, Staphyloma of the cornea.

CORNÉE, Cornea.

CORNEITIS, Ceratitidis.

CORNEL, ALTERNATE-LEAVED, *Cornus alternifolia*—c. American Red-rod, *Cornus sericea*—c. Large-flowered, *Cornus Florida*—c. Panicle, *Cornus paniculata*—c. Round-leaved, *Cornus circinata*—c. White, *Cornus paniculata*.

CORNE'LIAN, *Corne'lian*, *Chalcedonius*, *Corne'olus*, *Lapis Corne'olus*, *Corne'olus*, *Corne'olus*, *Lapis Sard'ius*, (F.) Cornaline. A precious, semi-transparent stone, found in Sardinia. The ancients ascribed to it a number of absurd properties.

CORNELUS, Cornelian.

CORNEOLUS, Cornelian.

CORNEOUS, *Corn'neus*, Horny, (F.) Corné, from cornu, 'a horn.' Having the nature or appearance of horn.

CORNEOUS TISSUE, *Tela cor'nea*, is that which forms the nails. The corneous membrane is the cornea.

CORNES DU CARTILAGE THYROÏDE, *Cornua cartilaginis thyroideae*—c. du Cœcyx, *Cornua cœcygis*—c. de l'Hyoïde, *Cornua hyoidei ossis*. CORNES DE LIMAÇON (F.), *Snail's Horns*. A name given by Anel to the lachrymal puncta and ducts.

CORNES DE LA MATRICE, *Cornua uteri*—c. de la Peau, Horny excrescences.

CORNESTA, Retort.

CORNET ACOUSTIQUE, Ear-trumpet—c. Moyen, Turbinated bone, middle—c. de Morgagni, Turbinated bone, superior.

CORNETS, Turbinated bones—c. de Bertin, Sphenoidal cornua—c. Sphénoidal (F.), Sphenoidal cornua.

CORNICHON, see *Cucumis sativus*—c. de Cerf, Cornu cervi.

CORNICULARIS PROCESSUS, Coracoid process.

CORNIC'ULUM, diminutive of cornu, 'a horn,' 'a little horn.' A species of cupping instrument, shaped like a trumpet, having a hole at the top for sucking the air out, to diminish the pressure in its interior.—Scultetus, Hildanus.

CORNICULUM LARYNGIS, *Capitulum Santorini* seu *Laryngis* seu *Cartilago'nis arytenoideæ*, *Supra-arytenoid Cartilage*, *Cartilage of Santorini*. A small, very movable, cartilaginous tubercle, found on the arytenoid cartilages.

CORNIER, Cornus Florida.

CORNIFICATION, *Racornissement*.

CORNINE, see Cornus Florida.

CORNOUILLER, Cornus Florida—c. à Feuilles arrondies, *Cornus circinata*—c. à Grandes fleurs, *Cornus Florida*—c. Soyeux, *Cornus sericea*.

CORNU, Ceras, Corn, Horn. (F.) Corne. A conical, hard, epidermous projection, which grows on the heads of certain animals, serving them as a weapon of offence and defence. Anatomists have given this name to parts of the human body, which have nearly the same shape as the horns of animals.

CORNU. A horny excrescence; a corneous wart, which occasionally forms on the skin, and requires the use of the knife; (F.) *Corne de la Peau*.—See Corn. Also, Cornu Cervi.—See Cervus. Also, a Retort.

CORNU ACUSTICUM, Ear-trumpet.

CORNU AMMO'NIS, *C. Arietis*, *Hippocampus major*, *Pes hippocampi major*, *Pes Hippopotami major*, *Protuberantia cylindrica*, *Vermis Bombycinus*, *Processus cerebri lateralis*, (F.) *Corne d'Ammon* ou de Béliet, *Grande Hippocampe*, *Pied de cheval marin*, *Protubérance cylindroïde* (Ch.), *Bourrelet roulé*. A broad, considerable eminence, curved on itself, and situate at the posterior part of the lateral ventricle. Its surface presents two or three tubercles separated from each other by shallow grooves.

The *Accessorius Pedis Hippocampi*, (F.) *Accessoire du pied d'hippocampe*, is a prominence, usually formed by the base of the inferior cornu of the lateral ventricle of the brain. It is merely a fold of the hemisphere, and was by Malacarne called *Cuisart* ou *Armure des Jambes*.

CORNU ANTERIUS seu ANTI'CUM VENTRICULI LATERALIS; Anterior Cornu of the Lateral Ventricle. The portion of the lateral ventricle of the brain, which is lodged in the middle of the lobe, and forms the commencement of that cavity.

CORNU CERVI, Cornu, C. Cervinum, Cervi Elaphi Cornu, Hartshorn, (F.) *Corne de cerf*, *Bois de cerf*, *Cornichon de cerf*. The horns of various species of the stag. They contain about 27 per cent. of gelatin. The Shavings, *Raspatura* seu *Ras'ura Cornu Cervi*, C. C. *raspatum*, boiled in water, have, consequently, been esteemed emollient and nutritive.

Hartshorn Jelly may be made as follows:—Hartshorn shavings, 3vj; boil in water Oiv to Oij; strain, and add, whilst hot, of lemon-juice, two tablespoonsful; white sugar, 3vj; and Sherry wine, two glasses. A good nutriment for the sick, where wine is not improper.

When burnt, the shavings constitute the *Cornu cervi calcinatum*, *Cornu ustum*, *Phosphas Calcis*, *Calcaria phosphorica*, (F.) *Corne de cerf calcinée*, *Phosphate de chaux*, which consists of 57.5 per cent. of phosphate of lime. It has been used as an antacid, but is wholly inert, as its composition would indicate. It contains only 1 per cent. of carbonate of lime.

Phosphate of lime has also been given in osteomalacia, from a presumption that the disease is owing to a deficiency of earthy matter in the

blood, which is doubtless an error—the fault being in the agents of assimilation or the cells of nutrition. With as little propriety, it has been proposed as an excitant to the healthy action of the cells in tuberculosis.

Bone earth is usually sold in the shops for burnt hartshorn.

The Dublin Pharmacopoeia has a *Calcis phosphas precipitatum*, prepared by dissolving burnt ox bones in muriatic acid, and precipitating by solution of ammonia. The dried precipitate, called *bone phosphate of lime*, has no virtues not possessed by the cornu ustum.

Hartshorn was once supposed to possess a bezoardic power.

CORNU CERVINUM, Cervus, Plantago coronopus.

CORNU DESCENDENS VENTRICULI LATERALIS, (F.) *Corne descendante ou inférieure du ventricule latéral, Dig'ital Cav'ity, Descending or inferior cornu of the lateral ventricle.* The termination of the lateral ventricle in the middle lobe of the brain, behind the fissure of Sylvius.

CORNU ETHMOIDAL, Turbinated bone, middle—c. Middle, Turbinated bone, middle.

CORNU POSTERIUS seu POSTICUM VENTRICULI LATERALIS, *Cavitas digitalis, Posterior Cornu of the Lateral Ventricle.* (F.) *Corne postérieure du ventricule latéral.* The triangular prolongation of the lateral ventricle of the brain into the substance of the occipital lobe.

CORNU RUPICAPRÆ, (F.) *Corne de Chamois.* The horn of the chamois. It has the same properties as the Cornu Cervi.

CORNUA, Turbinated bones.

CORNUA CARTILAGINIS THYROIDÆ, (F.) *Cornes du Cartilage thyroïde.* Eminences on the thyroid cartilage, distinguished into *great or superior*, which are articulated with the great cornu of the os hyoides;—and into *small or inferior*, united with the cricoid cartilage.

CORNUA COCCYGIS, *Cornua of the Coccyx*, (F.) *Cornes du Coccyx.* Two small, tubercular eminences at the base of the coccyx, which are articulated with those of the sacrum.

CORNUA CUTANEA, Horny excrescences.

CORNUA HYOIDEI OSSIS, *Radi'ces ossis hyoïdei, Cornua of the Hyoid Bone*, (F.) *Cornes de l'hyoïde.* Four fragments of the os hyoides, situate above the body of the bone, and distinguished into the *small or superior*, and the *great or lateral*.

CORNUA LACRYMALIA, Lachrymal ducts—c. Limacum, Lachrymal ducts, see Lachrymal puncta.

CORNUA SACRALIA, *Cornua of the Sacrum.* Two tubercles, situate at the posterior and inferior surface of the sacrum, which are sometimes united.

CORNUA SPHENOIDAL, see Sphenoidalia cornua—c. Styloid, see Hyoides, os.

CORNUA UTERI, *Cornua of the Uterus, Værææ, κεραι, Plectanæ*, (F.) *Cornes de la Matrice.* The angles of the uterus, where the Fallopian tubes arise. Sometimes applied to the Fallopian tubes themselves.

CORNUE, Retort—c. *Tubulée*, see Retort.

CORNUMUSA, Retort.

CORNUS ALTERNIFOLIA, *Alternate-leaved Cornel*; indigenous. *Nat. Ord. Cornaceæ.* The bark is one of the "Shaker medicines," being considered diaphoretic and astringent.

CORNUS AMOMUS, C. sericea—c. Blueberried, C. sericea.

CORNUS CIRCINATA, C. tomentosa, *Round-leaved Cornel or Dogwood*, (F.) *Cornouiller à feuilles arrondies.* Order, Cornaceæ. The bark of this variety has been used for similar purposes with the next.

CORNUS FLORIDA, *Dogwood, Dogtree, Boxtree, Bitter Redberry, Large-flowered Cornel, Malebrant, flowered Dogwood, Florida Dogwood, Virginia Dogwood, Boxwood (New England)*, (F.) *Cornouiller, Cornier, C. à grandes fleurs.* The bark of this beautiful tree, which grows everywhere in the United States, has been long employed as a substitute for cinchona. Dose, from ʒss to ʒi. Its active principle has been separated from it, and received the name of *Cornine*.

CORNUS FEMINA, C. sericea—c. Mas odorata, *Laurus sassafras*.

CORNUS PANICULATA, *Pan'icled or White Cornel or Dogwood*, indigenous, has been used as a substitute for Cornus Florida.

CORNUS RUBIGINOSA, C. sericea—c. Sanguinea, *Sebestina*, C. sericea.

CORNUS SERICEA, C. amo'mus seu f'e'mina seu rubigino'sa seu Sanguinea, *Swamp Dogwood, Red Willow, Rose Willow, New England Dogwood, Female Dogwood, Silky-leaved Dogwood, American Red-rod Cornel, Red-rod, Red O'sier, Blueberried Dogwood, Blueberried Cornus*, (F.) *Cornouiller soyeux.* The bark, it is said, has been found little inferior to *Cinchona Pallida* in intermittents.—Barton.

CORNUS TOMENTOSA, C. circinata.

CORNUTA, Retort.

CORODIALYSIS, see Coretomia.

COROLLA INFUNDIBULIFORMIS, see Tube, Fallopian.

CORONA, Crown—c. Ciliaris, Ciliary body—c. Dentis, Crown of a tooth—c. Glandis, Crown of the glans—c. Posterior unguæ, Olecranon.

CORONA RADIANIS, *Ra'diating Crown of Reil.* Fasciculi of white fibres radiate in all directions from every part of the surface of the optic thalamus, excepting its inner side, which is free and corresponds to the third ventricle; the anterior of these fibres pass directly forwards, the middle fibres outwards, and the posterior backwards, forming the *coro'na ra'diana*.

CORONA REGIA, Trifolium melilotus—c. Unguæ, Olecranon—c. Veneris, Crown of Venus.

CORONÆ CILIARIS MEMBRANULA, Ciliary zone—c. Palpebrarum, see Tarsus.

CORONAD, see Coronal Aspect.

CORONAL, *Corona'lis, Corona'rius.* Relating to the crown; from *corona*, 'a crown.' A name formerly given to the frontal bone, because on it partly reposes the crown of kings.

CORONAL ASPECT. An aspect towards the plane of the corona or crown of the head. *Coronad* is used adverbially to signify 'towards the coronal aspect.'—Barclay.

CORONAL SUTURE, *Sutu'ra corona'lis, S. Frontoparietalis, Puppis Sutu'ra, S. Arcua'lis.* The suture of the head, which extends from one temporal bone to the other, over the crown of the head, and unites the parietal bones with the frontal. The *Suture Coronale* of the French anatomists is the suture which unites the two halves of the os frontis at the early period of life. It is a prolongation of the sagittal.

CORONALE, MINERAL WATERS OF. These waters are found near Lucca, in Italy. They received their name from an erroneous notion, that they are particularly adapted for curing diseases of the os frontis. Their temperature is 95° Fahrenheit. They contain free carbonic acid, sulphates of lime and magnesia, chlorides of sodium and magnesium, &c., and some iron.

CORONARIUS, Coronal.

CORONARY, *Corona'rius*, (F.) *Coronaire*, from *corona*, 'a crown.' Resembling a crown.

CORONARY ARTERY OF THE STOMACH, *Arteria Corona'ria Ventriculi seu Gas'trica superior*, (F.) *Artère coronaire stomacique. A. Stomogast-*

trique (Ch.), is one of the branches of the cœliac, which passes towards the superior orifice of the stomach, extends along its lesser curvature, and ends by anastomosing with the pyloric branch of the hepatic. This artery furnishes branches to the inferior part of the œsophagus, the cardiac orifice, the great *cul-de-sac*, the middle part of the stomach, the lesser omentum, and anastomoses with the other arteries of those organs.

CORONARY LIGAMENT of the liver is a reflection of the peritoneum, which surrounds the posterior margin of the liver. The same term is likewise applied to ligaments which unite the radius and ulna.

CORONARY PLEXUS OF THE HEART. The anterior and posterior coronary plexuses of the heart are derived from the anterior and posterior cardiac plexuses. See Cardiac Plexus.

CORONARY PLEXUS OF THE STOMACH. This plexus of nerves is given off from the upper part of the solar plexus.

CORONARY SINUS OF THE HEART, Sinus, coronary, venous—c. Sinus of Ridley, Sinus coronarius.

CORONARY TENDONS, *Tendines corona'riæ*. The fibro-cartilaginous rings at the orifices of the coronary arteries and aorta.

CORONARY VEIN OF THE STOMACH, *Vena Coronaria Ventriculi*, accompanies the artery, and terminates in the vena porta. Sömmering, and some other anatomists, call all the four arteries of the stomach *Coronariæ Stomachicæ*.

CORONCION, Coreoncion.

CORO'NĒ, from *κρῶν*, 'a crow.' The coronoid process of the lower jaw.

COR'ONOID, *Coronoï'des*, from *κρῶν*, 'a crow,' and *αἶδος*, 'shape,' 'resemblance.' Resembling the beak of a crow. The name has been given to two processes. One, situate at the anterior and superior part of the ramus of the os maxillare inferius, and affording attachment to the temporal muscles: the other, called, also, *sharp process*, situate at the superior part of the ulna, anterior to the great sigmoid fossa, and forming a part of the hinge of the elbow-joint.

CORONOPodium, *Plantago coronopus*.

CORONOPUS, *Cochlearia coronopus*—c. Depressus, *Cochlearia coronopus*—c. Ruellii, *Cochlearia coronopus*—c. Vulgaris, *Cochlearia coronopus*.

COROTOMIA, Coretomia.

CORP, Cadaver.

CORPORA, see Corpus—c. Albicantia, Mammary tubercles.

CORPORA AMYLA'CEA, *Corpuscula amyla'cea*, *Amyla'ceous* or *Am'ylloid bodies*, *Starch corpuscles*, *Animal starch*. A name given by Virchow to certain bodies of microscopic size, found chiefly in the human brain and spinal marrow—in the *ependyma ventriculorum* and its prolongations—mingled with the proper nerve elements; and having most of the chemical characters of cellulose.

CORPORA ARANTII, *Noduli Arantii*, *Tubercula A.*—c. Bigemina, *Quadrigenina corpora*—c. Candicantia, *Albicantia corpora*, *Mamillary tubercles*—c. Cavernosa, *Cavernous bodies*—c. Fibrosa, *Corps Fibreux*.

CORPORA FIMBRIA'TA, *Tænia Hippocampi*, *Fimbriated* or *fringed bodies*, (F.) *Corps Frangés* ou *Bordés*, *Banquettes des Cornes d'Ammon*, *Banquette de l'Hippocampe*. The thin, flattened, and very delicate band, situate along the concave edge of the cornu ammonis, which is a continuation of the posterior crura of the fornix.

CORPORA GENICULA'TA, (F.) *Corps Géniculés*, *C. génouillés*. Eminences situate at the lower and outer part of the optic thalami. Each antie-

tract commences at the *corpus geniculatum externum*. The *corpus geniculatum internum* is merely a tubercle inserted into the bend or knee, formed by the corpus geniculatum externum.

CORPORA GLOBOSA CERVICIS UTERI, *Nathothi glandule*—c. Lutea, see *Corpus luteum*.

CORPORA MALPIGII'NA, *Malpighian Bodies*, *Ac'ini*, *Corpuscles* or *Glomerules* of *Malpighi*, *Glomeruli renales*, *Corpuscula Malpighii*. Scattered through the plexus formed by the blood-vessels and uriniferous tubes in the kidney, a number of small dark points may be seen with the naked eye, which received their name from Malpighi, their describer. Each of these, under the microscope, is found to consist of a convoluted mass of blood-vessels, which constitutes the true *glandule, corpuscle*, or *glomerule* of *Malpighi*. It was at one time supposed that the tubuli uriniferi originate in them; but this does not appear to be the case. Their use is not positively known, but as they have been traced by Mr. Bowman into the commencement of the urinary tubes, in which they lie uncovered, it has been supposed that their office may be to separate the watery portions of the blood to be mixed with the proper urinous matter.

CORPORA MAMMILLARIA, *Mamillary tubercles*—c. Nervio-spongiosa *Penis*, *Corpora cavernosa*—c. Nervosa, *Corpora cavernosa*.

CORPORA OLIVA'RIA, *C. ovata*, *Eminentia Olivares* seu *ovales laterales*, *Olivæ*, *Prominentia Semiorales Medullæ Oblongatæ*, (*Eminences latérales*, Ch.) Oblong, whitish eminences, situate at the occipital surface of the medulla oblongata, exterior to the corpora pyramidalia.

CORPORA OVATA, *Corpora olivaria*—c. Pisiformia, *Mamillary tubercles*.

CORPORA PYRAMIDA'LIA, *Eminentia pyramidalis* seu *medullæ inferiæ*, *Pyramides*, (F.) *Corps pyramidaux*, *Pyramides antérieures* (Gall), *Eminences pyramidales* (Ch.) Two medullary bundles, placed on either side of the anterior fissure. These bodies have also been called *Corpora Pyramidalia antica*, to distinguish them from the *C. olivaria*, which have been called *Corpora Pyramidalia lateralia*.

Over a space, commencing 8 or 10 lines below the pons and extending to the lower end of the medulla oblongata, a portion of each pyramid crosses to the opposite side—*decussation of the pyramids*, *Decussatio pyramidum*.

CORPORA PYRAMIDALIA POSTERIORA, see *Corpora restiformia*—c. *Pyramidalia testis*, *Coni vasculosi*—c. *Quadrigenina*, *Q. terbecula*.

CORPORA RESTIFORMIA, *Crura* seu *Pedunculi medullæ oblongatæ*, *Corpora pyramidalia posteriora*, *Processus a cerebell'o ad medullam oblongatam*, *Posterior pyramids*, *Peduncles of the medulla oblongata*, *Inferior peduncles of the cerebellum*, (F.) *Corps restiformes*, *Cuisses postérieures*, *Pyramides postérieures*, *Racines*, *Bras ou Jambes du cercelet*, *Petites branches de la moëlle allongée*, *Pédoncules du cercelet*. Two medullary projections, oblong, and of a whitish appearance, which proceed from each side of the upper extremity of the medulla oblongata, and contribute to the formation of the cerebellum. The *posterior pyramids*—*fasciculi graciles*—are described as distinct from the restiform bodies. They are situate in contact with each other. one on each side of the posterior median fissure. They become closely applied to the restiform bodies, and have been considered to be blended with them, and to proceed to form the cerebellum. According to some, they ascend to the cerebrum.

CORPORA STRIATA, *Grand ganglion supérieur du cerveau* (Gall), *Couches des nerfs éthmoïdaux*, *Corps cannelés*, *Eminentia Lenticulares*, *Colliculi Nervi Ethmoidalis*, *Apices Crurum medullæ*

oblongata, *Ganglion cerebri anterius*, *Anterior cerebral ganglion*, (F.) *Corps striés*. Pyriform eminences of a slightly brownish-gray colour, which form part of the floor of the lateral ventricles of the brain. When cut, a mixture of gray and white substance is seen, arranged alternately, to which they owe their name. The tract of fibres that ascends from the anterior pyramids passes chiefly into them. Willis considered that the soul resided there.

CORPORA STRIATA SUPERNA POSTERIORA, *Thalami nervorum optico-rum*—c. *Triticea*, see *Hyoides os*—c. *Wolfiana*, see *Corpus Wolfianum*.

CORPUS, Body—c. *Bordés*, *Corpora fimbriata*—c. *Callez*, *Corpus callosum*—c. *Cannelés*, *Corpora striata*—c. *Caverneux*, *Corpora cavernosa*—c. *Cendré*, *Corpus dentatum*—c. *Ciliaire*, *Corpus dentatum*—c. *Dentelé*, *Corpus dentatum*—c. *Étranger*, *Extraneous body*—c. *Festonné*, *Corpus dentatum*.

CORPUS FIBREUX, *Corpora Fibrosa*. Bayle has given this name to adventitious fibrous productions of a round shape, more or less adherent, and sometimes having a pedicle, which form in certain parts of the body, particularly in the substance of the uterus.

CORPUS FOLLICULAIRE VAGINALE, *Cowper's glands in the female*—c. *Frangés*, *Corpora fimbriata*—c. *Géniculés*, *Corpora geniculata*—c. *Genouillés*, *Corpora geniculata*—c. *Godronné*, *Fascia dentata*—c. *Humain*, see *Body*—c. *Hyalode*, *Corpus vitreum*—c. *d'Hygmore* ou *d'Highmore*, *Corpus Highmori*—c. *Jaune*, *Corpus luteum*—c. *Muqueux*, *Corpus mucosum*—c. *d'Oken*, *Corpus Wolfianum*—c. *Organisés*, *Organized bodies*—c. *Pampiniforme*, *Corpus pampiniforme*—c. *Phacode*, *Crystalline*—c. *Pyramidaux*, *Corpora pyramidalia*—c. *Restiformes*, *Corpora restiformia*—c. *Rhomboïde*, *Corpus dentatum*—c. *Striés*, *Corpora striata*—c. *Thyroïde*, *Thyroid gland*—c. *Variéforme*, *Epididymis*—c. *Variqueux*, *Corpus pampiniforme*, *Epididymis*—c. *Vitré*, *Corpus vitreum*—c. *de Wolff*, *Corpus Wolfianum*.

CORPSE, *Cadaver*.

CORPULENT, *Corpulentus*, *Obe'sus*, *Crassus*, *Fat*, *Fleshy*, (Sc.) *Corax*, (Prov.) *Sobby*; from *corpus*, 'the body,' and *lentus*, 'thick.' Having an unusual development of fat or flesh in proportion to the frame of the body.

CORPULENTIA, *Corpulence*, is synonymous with *Obesity* and *Polysarcia*.

CORPULENTIA CARNOSA, *Torositas*.

CORPUS, *Soma*, a *Body*. Any object which strikes one or more of our senses. Gases, liquids, metals, vegetables, animals, are so many bodies. Natural bodies have been divided into *animal*, *vegetable*, and *mineral*; or into *inorganic*, including the mineral kingdom; and *organised*, including the animal and vegetable. The chief differences between organized and inorganic bodies consists in the former having an origin by generation, growth by nutrition, and termination by death; the latter a fortuitous origin, external growth, and a termination by chemical or mechanical force.

Many parts of the frame have, also, been distinguished by this name, as *Corpus Callosum*, *C. Mucosum*, &c. See *Body*.

CORPUS ADENIFORME, *Prostate*—c. *Adenoides*, *Prostate*—c. *Adiposum*, *Pinguedo*—c. *Alienum*, *Extraneous*—c. *Annulare*, *Pons varolii*.

CORPUS CALLOSUM, *Commissura Magna* seu *maxima cerebri*, *Trabs Medullaris* seu *Cerebri*, *Trabecula* seu *Copula magna cerebri*, (F.) *Corps Callez*, *Voûte médullaire*, *Plafond des ventricules du cerveau*, *Méolobe* (Ch.) A white, medullary band, perceived on separating the two hemispheres of the brain, which it connects with

each other. La Peyronie regarded it as the seat of the soul. On it are seen longitudinal and transverse fibres—*Stria longitudinalis Lancisii*, (see *Raphe*), and *Stria seu Chorda transversales Willisii*. The anterior portion, which bends downwards, is termed *genu*; the posterior flexure, *epilegium*, *tuber*, (F.), *bouretlet*. The fibres which curve backwards into the posterior lobes from the posterior border of the corpus callosum, have been termed *Forceps*; those which pass directly outwards into the middle lobes from the same point, *Tape'tum*; and those which curve forwards and inwards from the anterior border to the anterior lobe, *forceps anterior*.

CORPUS CILIARE, *Ciliary Body*, *Corpus dentatum*, see *Ciliary*—c. *Cinereum*, *Corpus dentatum*—c. *Conoides*, *Pineal gland*—c. *Crystalloides*, *Crystalline*.

CORPUS DENTATUM, *C. Denticulatum* seu *Ciliaire*, (F.) *Corps dentelé* ou *festonné*. A central, oval nucleus, of cineritious substance, met with in the cerebellum; the circumference of which exhibits a number of indentations, surrounded by medullary substance.—*Vicq d'Azyr*. It is seen by dividing the cerebellum vertically into two equal parts.—The same body has been called *Corps cendré* ou *ciliaire* ou *rhomboïde*, *Corpus Cine'reum* seu *Rhomboïdeum* seu *Rhomboïdale*, *Ganglion du cervelet*, *Noyau central des Peduncules du cervelet*, *Nucleus dentatus* seu *fimbriatus* seu *centralis* seu *rhomboidalis*, *Substantia rhomboïde*, *Ganglion ciliaire*, (F.) *Cerebelli*. The term *Corpus dentatum*, *Nucleus olivæ*, is also given to the ganglion of the corpus olivæ, which, like that of the cerebellum, is a yellowish-gray dentated capsule, open behind, and containing medullary matter, from which a fasciculus of fibres proceeds upwards to the corpora quadrigemina and thalami optici.

CORPUS DENTICULATUM, c. *Dentatum*—c. *Discoïdes*, *Crystalline*—c. *Externum*, *Extraneous body*—c. *Extraneous*, *Extraneous body*—c. *Geniculatum*, see *Corpora Geniculata*—c. *Glandiforme*, *Prostate*—c. *Glandosum*, *Prostate*—c. *Glandulosum*, *Prostate*.

CORPUS GLANDULOSUM MULIERUM, *Glandula Prostata mulierum*. A vascular spongy eminence, which surrounds the orifice of the urethra, and projects at its under part.

CORPUS HIGHMORI, *C. Highmoria'nium*, *Mediastinum testis*, *Medusta semina'ria*, (F.) *Corps d'Hygmore* ou *d'Highmore*, *Sinus des Vaisseaux séminifères*, (Ch.) An oblong eminence, along the superior edge of the testicle, which seems formed of a reflection of the tunica albuginea, through which the principal trunks of the seminiferous vessels pass before they reach the epididymis.

CORPUS INCOMPREHENSIBILE, *Thymus*.

CORPUS LUTEUM, (F.) *Corps jaune*, 'yellow body.' A small yellowish body, perceived in the ovary, and left after the rupture of one of the vesicles. It was, for a long time, considered an evidence of previous impregnation; but *Corpora lutea* are met with in unquestionable virgins; although the corpora lutea of virgins have been generally regarded to differ materially, in size and character, from those of impregnation, which have been called *true corpora lutea* in contradistinction to the other, called *false corpora lutea*.

CORPUS MUCCOSUM, *C. reticulare*, *Reté mucosum*, *Mucus* seu *Reté seu Stratum Malpighii*, *Reticulum cutaneum* seu *mucosum*, *Meodermum*, *Mucous web*, (F.) *Corps muqueux*, *Couche de Malpighi*. The second layer of the skin has been so called. It is situate between the *cutis vera* and *cuticle*, and gives colour to the body. In the white varieties of our species it is colourless; in the negro, black. By most anatomists the exist-

ence of such a layer, distinct from the epidermis, is denied.

CORPUS OKENSE, Corpus Wolffianum.

CORPUS PAMPINIFORME, from *pampinus*, 'a tendril.' *C. Varicosum* seu *pyramidalis*, *Hederaceus plexus*, *Plexus pampiniformis* seu *vasculosus funiculi spermatici pampiniformis*, (F.) *Corpus pampiniforme*, *C. Varicose*. The plexus or retiform arrangement of the spermatic arteries and veins in the cavity of the abdomen, anterior to the psoas muscle.

CORPUS PAPILLARE, *Textus Papillaris*, *Corpus reticularis*. The nervous and vascular papillae situate beneath the epidermis, called by Breschet *Neurothel'ic apparatus*.

CORPUS PHACOIDES, Crystalline—c. Pituitare, Pituitary gland—c. Psalloides, Lyra—c. Pyramidale, Corpus pampiniforme—c. Reticulare, Corpus mucosum, Corpus Papillare—c. Rhomboidale, Corpus dentatum—c. Rhomboideum, Corpus dentatum—c. Thynianum, Thymus—c. Thymicum, Thymus—c. Thyroideum, Thyroid gland—c. Turbinatum, Pineal gland—c. Varicosum, Corpus pampiniforme, Spermatic cord—c. Varicosum testis, Epididymis.

CORPUS SPONGIOSUM URETHRÆ, *Substantia spongiosa urethræ*. This substance arises before the prostate gland, surrounds the urethra, and forms the bulb. It then proceeds to the end of the corpora cavernosa, and terminates in the glans penis, which it forms. Kobelt describes, in the female, as the analogue to the corpus spongiosum of the male, a venous plexus, which, as it lies between the glans clitoridis, and the part that corresponds, in the male, to the bulb of the urethra, he terms *pars intermedia*, (F.) *Réseau intermédiaire*.

CORPUS STRIATUM, see *Corpora striata*—c. Trigonom, see Urinary Bladder.

CORPUS VITREUM, *Humor Vitreus* seu *Hyaloides* seu *hyalinus*, *Glacia'lis humor*, *Vitreous humor*, *Vit'rina ocularis*, (F.) *Corps vitré ou hyaloïde*, *Humeur hyaloïde*. The transparent mass, of a gelatinous consistence, which fills the eye, behind the crystalline. It is contained in cells, formed by the *tunica hyaloïde*.

CORPUS WOLFFIANUM, *C. Oken'sē*, *Wolffian body*, (F.) *Corps de Wolff* ou *d'Oken*. At a very early period of foetal formation, bodies are perceptible, which were first described by Wolff, as existing in the fowl, and in the mammalia by Oken. According to Müller, they disappear in man very early, so that but slight remains of them are perceptible after the 9th or 10th week of pregnancy. They cover the region of the kidneys and renal capsules, which are formed afterwards; and they are presumed to be the organs of urinary secretion during the first periods of foetal existence.

CORPUSCLES, BLOOD. Globules of the blood—c. Bone, see *Lacunæ of Bone*—c. Caudate, see *Caudate*—c. Chyle, see *Chyle*.

CORPUSCLES, EXUDATION. The organizable nuclei contained in fibrinous fluids, which are the origin of the new tissues formed from such fluids.

CORPUSCLES, GANGLION, see *Neurine*—c. Glandiform, *Acinus*—c. Lymph, see *Lymph*—c. of Malpighi, *Corpora Malpighiana*—c. Mucous, see *Mucus*—c. Nerve, see *Neurine*—c. Osseous, see *Lacunæ of bone*.

CORPUSCLES, PACINIAN, (F.) *Corpuscules de Pacini*, so called from Filippo Pacini, an Italian physician, who, it is generally conceived, first noticed them in 1830. They appear, however, to have been depicted in 1741 by Lehmann, from a preparation by A. Vater, who called them *Papillæ* and *P. nervæ*. Hence, it has been proposed by J. C. Strahl (1848) to call them *Vaterian corpuscles* or *Corpuscles of Vater* (*Vater'sche*

Körperchen.) Small bodies connected with the cutaneous nerves of the palm and sole. They have also been found sparingly and inconstantly in nerves at the wrist and elbow; in the upper arm, fore-arm, and thigh, and inter-costal nerve, the sacral plexus, solar plexus and the plexuses adjacent to it. In each corpuscle there is the termination of a nervous filament. Their uses are not known.

CORPUSCLES OF PURKINJE, see *Canaliculus*—c. Pus, see *Pus*—c. Pyoid, see *Pus*—c. Splenic, see *Spleen*—c. Starch, *Corpora amylacea*—c. Tactile, *Corpuscula tactus*—c. Touch, *Corpuscula tactus*—c. of Vater, c. Pacinian—c. Weberian, see *Ejaculatory Ducts*—c. White, see *Globules of the blood*—c. White granulated, see *Globulin*.

CORPUSCULA ARANTII, *Tubercula A.*, see *Sigmoid valves*—c. *Glandularum similia intestinalium*, *Peyeri glandulæ*—c. *Globosa cervicis uteri*, *Nabothi glandulæ*.

CORPUSCULA LÆNNÆCII. Fibroid granulations, occasionally observed on thickened portions of the pericardium.

CORPUSCULA MALPIGHII, *Corpora Malpighiana*—c. Ossium, see *Lacunæ of bone*—c. Sanguinis, *Globules of the blood*—c. Seminis, *Spermatozoa*—c. Sesamoidea, see *Sigmoid valves*.

CORPUSCULA TACTUS, *Tactile* or *Touch Corpuscles*, (F.) *Corpuscules du tact*, *Azile bodies*, (G.) *Axen Körper*. Bodies, composed of a horizontally laminated mass of areolar tissue, which are found in the papillæ of parts endowed with great tactile sensibility. The nerves of touch communicate with them, and their function is probably to render the sense more acute.

CORPUSCULA TRITICEA, see *Hyoides*, os—c. Weberiana, see *Ejaculatory Ducts*.

CORPUSCULAR, *Corpuscularis*. Relating to, or containing, corpuscles, as *corpuscular fibrin*. Fibrin which tends to disintegration, and consists of granules in considerable proportion.

CORPUSCULES CALCAIRES, *Lacunæ of bone*—c. *Noirs*, *Lacunæ of bone*—c. *Des Os*, *Lacunæ of bone*—c. *Ramiflès*, *Lacunæ of bone*.

CORPUSCULUM ARANTII, see *Sigmoid valves*—c. *Conicum Rosenmülleri*, *Parovarium*—c. *Sesamoideum*, see *Sigmoid valves*.

CORRAGO, Borago officinalis.

CORRÊ, *Corrê*, from *καρπ*, 'I shave.' [?] The temples or the part of the jaws, which it is usual to shave.—Gorræus.

CORRECTIF, *Correctif*.

CORRECTION, *Correctio*, from *corrige*, (con and *regere*, *rectum*, 'to rule or order') to correct.' The act of correcting medicines; that is, of diminishing their energy or obviating unpleasant effects, by mixing them with substances which mitigate their operation.

CORRECTORIUS, *Corrigens*.

CORRELATION, see *Synergy*.

CORRIGEEN MOSS, *Fucus crispus*.

CORRIGENT, *Corrigens*, *Correctorius*, *Castigans*, *Infringens*, *Emendans*: same etymon. (F.) *Correctif*. That which corrects. A corrigent, in a pharmaceutical formula, is a substance added to a medicine to mollify or modify its action. In the following formula, the aloes, if not corrected, might induce tormina. The *Oleum Menthae* is added as a corrigent.

R Aloes

Olei Menthae

Syrup q. s. ut fiant Pilulæ

xvj

CORRIG'IA. Akin to *corium*, 'leather. Leather straps. By extension, the term has been applied to the tendons and ligaments.

CORROBORANT, *Corrob'orative*, *Corrob'orans*, *Corroborati'vus*, *Rob'orans*, *Muscularis*, *Restan'rans*, *Restor'ative*, *Bracing*, from *corroborare* (con, and *robur*, 'strength,') 'to strengthen.

(F.) *Corroborant, Corroboratif, Fortifiant*. Any substance which strengthens and gives tone. Wine, for example, is a corroborant. See Tonic.

CORROBORANTIA, Tonics.

CORROBORATIF, Corroborant.

CORROBORATIVE, Corroborant.

CORRODANT, Corrosive.

CORRODENS, Corrosive.

CORROSIF, Corrosive.

CORROSION, *Corro'sio, Diabro'sis, Anabro'sis, Ero'sio*, from *con*, and *rodere, rosus*, 'to gnaw.' The action or effect of corrosive substances.

CORRO'SIVE, *Corro'dens, Diabro'icus, Corros'ous*, same etymon. (F.) *Corrosif, Corrodant*. Corrosives are substances, which, when placed in contact with living parts, gradually disorganize them. *Caustic alkalies, Mineral acids, Corrosive sublimate*, are *corrosives*. They act either directly, by chemically destroying the part, — or indirectly, by causing inflammation and gangrene.

CORROSIVE POISON, see Poison.

CORRUGATIO, Corrugation — *c. Cutis*, Wrinkle.

CORRUGA'TION, *Corrugat'io, Synæ'rema*, from *con*, and *rugā*, 'a wrinkle.' Wrinkling, Frowning, (F.) *Frontement*; the contraction of the *Corrugato'res Supercil'ii* muscles. Corrugation of the skin is often owing to the application of styptic medicines: it is rendered by them unequal and rugous.

CORRUGATOR COITERII, Corrugator supercilii.

CORRUGA'TOR SUPERCIL'II, from *corrugare*, 'to wrinkle,' same etymon. *Musculus supercil'ii seu frontalis versus seu Corrugator Coite'rii, Supercil'aris*, (F.) *Cutanéo-sourcilier, Muscle Sourcilier ou Surcilier, M. Fronto-Sourcilier*. A muscle situate in the eyebrows. It is attached, by its inner extremity, to the superciliary ridge, and is confounded, externally, with the occipitofrontalis and orbicularis palpebrarum. It carries the eyebrow inwards, and wrinkles the skin of the forehead.

CORRUPT'ION, *Corrupt'io, Phthora, Diaphthora*, from *corrumpere, corruptum*, (*con*, and *rumpere*, 'to break,') 'to destroy.' Act of corrupting. State of being corrupted. Reaction of the particles of a body upon each other. It is probable that something like corruption may take place even in the living body. The word is used, at times, like *Boue* with the French, in a vague manner for a state of puriform, putrid, or other breaking down of parts or humours. 'To spit corruption,' is a vulgar expression.

CORSE, *Cadaver, Corce*.

CORSET, from *Corse*, (F.) *corps*, 'the body.' *Stethodesm'ium, Stethodesm'is, Stethodesm'us, Tunica Thorac'is, Thorax, Pectoral'is*. An article of dress, which closely embraces the trunk, and is much used by females in civilized countries. When corsets or stays are worn very tight, many serious evils result from the unnatural compression.

Different bandages, more or less complicated, which embrace the greater part of the trunk, are likewise so called.

An *orthopedic corset* is one intended to obviate deviation of the stature.

CORSET DE BRASDOR. The name of a bandage invented by one Brasdor, to keep *in situ* the fragments of a fractured clavicle.

CORSSY, Corpulent.

CORTALON, Senecio.

CORTEX, *Phloios, Phloos, Phloos, Lemma, Bark*, (F.) *Écorce*. This word has often been applied exclusively to *Cinchona*: thus, we say *Bark* — the cortex or bark, *καὶ ἑξοχῇ*. It means, also, any bark.

CORTEX ADSTRING'ENS BRASILIIENSIS. An as-

tringent bark introduced from Brazil into Germany in the year 1828. It is said to be obtained from *Mimosa cochlearifolia seu virginiana*. It has been used with advantage in all cases in which astringent barks in general are indicated. Dose of the powder ʒj to ʒss.

CORTEX ALCOMOCO, Alcomogue — *c. Anisi stellati*, see *Illicium anisatum* — *c. Antiscorbuticus*, *Canella alba* — *c. Aromaticus*, *Canella alba* — *c. Aurantii*, see *Citrus aurantium* — *c. Canellæ Malabaricæ*, *Laurus cassia* — *c. Cardinalis del Lugo*, *Cinchona* — *c. Caryophyllatus*, see *Myrus caryophyllata* — *c. Caryophylloides*, *Laurus cullawan*.

CORTEX CEREBRI. The *Cortical, Cinerit'ious, Vesicular or Gray substance of the Brain*. The gray portion observed at the exterior of the cerebrum and cerebellum; so called because it forms a kind of bark to the medullary substance. Gall considers, that this substance forms the nerves; and therefore calls it *Substantia matrice des Nerfs*.

The name *cortical* is likewise given to the external substance of the kidneys, because it is of a deeper colour than the inner part of the organ, and forms a kind of envelope to it.

CORTEX CHACARILLÆ, *Croton Cascarilla* — *c. Chinæ*, *Cinchona* — *c. Chinæ flavus*, *Cinchonæ cordifoliæ cortex* — *c. Chinæ luteus*, *Cinchonæ cordifoliæ cortex* — *c. Chinæ regius*, *Cinchonæ cordifoliæ cortex* — *c. Crassior*, *Laurus cassia* — *c. Cullaban*, see *Laurus Cullawan* — *c. Cullawan*, *Laurus Cullawan* — *c. Eleutheriæ*, *Croton cascarilla* — *c. Flavius*, *Cinchonæ cordifoliæ cortex* — *c. Lavola*, see *Illicium anisatum* — *c. Magellanicus*, see *Wintera aromatica*.

CORTEX OVI, Cortical membrane. This membrane, so called by Boer and Granville, is usually regarded as a uterine production, and designated *Decidua reflexa*. They consider it to surround the ovule, when it descends into the uterus, and to enclose the shaggy chorion. It is absorbed during the first months of utero-gestation, so as to expose the next membrane to the contact of the decidua, with which a connexion takes place at the part where the placenta is to be formed. In that part, Boer and Granville consider, that the Cortex Ovi is never altogether obliterated, but only made thinner, and in process of time is converted into a mere pellicle or envelope, which not only serves to divide the filiform vessels of the chorion into groups or cotyledons, in order to form the placenta, but also covers those cotyledons. This, Dr. Granville calls *membrana propria*. See *Decidua membrana*.

CORTEX PALLIDUS, *Cinchonæ lancifoliæ cortex* — *c. Patrum*, *Cinchona* — *c. Peruvianus*, *Cinchona* — *c. Profluvii*, *Nerium antisyderiticum* — *c. Ruber*, *Cinchonæ oblongifoliæ cortex* — *c. Striata dentium*, Enamel of the teeth — *c. Thuris*, *Croton cascarilla* — *c. Winteranus*, see *Wintera aromatica* — *c. Winteranus spurius*, *Canella alba*.

CORTICAL, Cortical'is; from *cortex*, 'bark.' Belonging to bark.

CORTICAL MATTER OF THE BRAIN, Cortex Cerebri — *c. Membrane*, *Cortex Ovi* — *c. Substance of the Kidney*, see *Kidney* — *c. Substance of the Teeth*, see *Tooth*.

CORTUSA AMERICANA, *Heuchera cortusa*.

CORU. An Indian tree, the bark of whose root furnishes a milky juice, which is employed in diarrhoea and dysentery. It is also called *Coru Canar'ica*.

CORUSCATIO, Astrape.

CORVISARTIA HELENIUM, *Inula helenium*.

CORYBANTIASM. *Corybantias'mus, Corybantis'mus*, from *Κορυβας*, one of the Corybantes. A name formerly given to a kind of frenzy, in which the sick were tormented by fantastic visions, and perpetual want of sleep.

CORYDALIS BULBOSA, *Fumaria bulbosa*—c. *Canadensis*, *Dicentra Canadensis*—c. *Cava*, *Fumaria bulbosa*—c. *Tuberosa*, *Fumaria bulbosa*.
 CORYLUS, (from *κορυς*, 'a helmet,' from the involucre,) *Corylus avellana*.

CORYLUS AVELLANA, *Corylus*, *Bundurh*, *Avella'na*, *Caryd'ion*, *The Hazel-nut Tree*, (F.) *Coudrier*, *Noisetier*; *Ord. Amantaceæ*; *Sex. Syst. Monœcia Polyandria*. The nut,—*Filbert*, (F.) *Aveline*; *Hazel-nut*, (F.) *Noisetie*,—of this tree, is much eaten in many countries. Like all nuts, it is by no means easy of digestion. It is the *Nux avellana*, *καρυον Πωρικον*, of the ancients.

CORYLUS ROSTRATA, *Beaked Hazel*. An indigenous shrub, *Nat. Ord. Amantaceæ*; *Sub-order*, *Cupuliferæ*; *Sex. Syst. Monœcia Polyandria*; the nut of which is surrounded by a coriaceous and scaly involucre, terminating in a tube covered with short and thick bristles. These bristles have been given as an anthelmintic in the same cases and doses as *mucuna*.

CORYMBETRA, *Hedera helix*.

CORYNE, *Penis*.

CORYPHE, *Acme*, *Vertex*—c. *Cordis*, see *Mucro*.

CORYZA, *Grave'do*, *Rhinitis*, *Catarrh'os*, *Catarrhal'g'mos*, *Stillicid'ium Nas'rium*, *Phlegmatorrhag'ia*, *Blennorrhœa nasalis*, *Blennorrhin'ia*, *Des'tillatio*, *Catarrhus ad Nares seu nasalis*, *Rhinocatar'rhus*, *Angina nasalis*; vulgarly, 'running at the nose,' 'cold in the head;' (Prov.) *Snuurle*, in French, *Rhume de cerveau*, *Catarrhe nasal*, *Enchiffrement*; in old English, *Pose* or *Mur*; whence *Murren* and *Murrain*; from *κορυς*, or *καρυς*, 'the head,' and *ζωω*, 'I boil.' Inflammation, attended with increased discharge, of the Schneiderian membrane lining the nose, and the sinuses communicating with it. The affection generally subsides without any medical treatment.
Chronic coryza is termed, also, *Ozæna benigna*.

CORYZA ENTONICA, *Ozæna*—c. *Maligna*, *Ozæna*—c. *Ozænosa*, *Ozæna*—c. *Purulenta*, *Ozæna*—c. *Scarlatiosa*, *Rhinocaceæ*—c. *Ulcerosa*, *Ozæna*—c. *Virulenta*, *Ozæna*.

COSCINISMUS, *Cribration*.

COSCINOI, see *Cribration*.

COSMETIS, *Cosmetics*.

COSMETICS, *Arts cosmetica*, *Callipist'ria*, *Cosm'ia*, (F.) *Cosmétique*, from *κοσμεω*, 'I adorn,' 'I embellish.' The art of improving the beauty. *Cosmetic*, *Stilbo'ma*, is, also, used for the different means employed for that purpose; as the compounds, into which enter the oxides of lead, bismuth, mercury, arsenic, &c. All these, however, injure the skin, and often give rise to unpleasant cutaneous affections. Frequent ablution with cold water, and bathing are the best cosmetics. Essences, soaps, and all the preparations intended for the toilet, fall, also, under this head.

COSMÉTIQUE, *Cosmetics*.

COSMETOLOG'Y, *Cosmetolog'ia*, from *κοσμεω*, 'to adorn,' and *λογος*, 'a discourse.' A treatise on the dress, and cleanliness of the body.

COSMOS, from *κοσμος*, 'the world,' 'order,' &c. The order which was supposed to preside over critical days. Hippocrates and others have termed *κοσμοι*, 'bracelets,' employed, not only as ornaments (*κοσμευ*, 'to adorn,') but as therapeutical agents.

COSSA, *Haunch*.

COSSUM. A malignant ulcer of the nose, often of a syphilitic character.—*Paracelsus*.

COSSUS, *Cossia*. A sort of white, short, thick worm or larve, found in trees, logs of wood, &c., and used by the Romans as a great article of *gourmandise*. They were, also, applied to ulcers.—*Pliny*. The term has, likewise, been given to

small vermiform pimples on the face—*acne*—which arise from inflammation of the sebaceous follicles.

COSTA, *Pæura*, *Pleurum*, *Pleuro'ma*. A *Rib*, from *custodire*, 'to guard,' 'defend.' (F.) *Côte* old French, *Coste*. The ribs are 24 in number;—12 on each side. They are irregular, long, bony curves; slightly flattened, and situate obliquely at the sides of the chest. The intervals between them are called *Intercostal spaces*, (F.) *Espaces intercostaux*, and they are numbered first, second, third, &c., reckoning from above to below. They have been distinguished into *Costæ veræ*, *Pleuro-poph'yses* of *Owen*, *True ribs*, (F.) *Vraies Côtes*, *Côtes sternales*, *Côtes vertébro-sternales* (Ch.), and into *Costæ spur'ia*, *Mendo'sæ* seu *Nothæ Costæ*, *False ribs*, *Côtes asternales* (Ch.), *Faussees Côtes*. The true or sternal ribs, as they have also been called, are the first 7; which are articulated at one extremity to the spine, and at the other, by means of their cartilages, *hamapoph'yses* of *Owen*, to the sternum. The false or short ribs are the remaining 5: the uppermost three being united, by means of their cartilages, to the cartilage of the last true rib. The others are free at their external extremity, and, hence, have been called *Floating ribs*, *Costæ fluctuantes*, (F.) *Côtes flottantes*. The vertebral extremity of each rib is slightly expanded. It is called the head of the rib—*Capitulum Costæ*: the space between this and the tubercle is the *collum* or *neck*. Anterior to the tubercle is the *angle*—*Angulus* seu *Cubitus Costæ*. The angle is the part where the bone bends to form the lateral part of the thorax.

COSTÆ CAPITULUM, see *Costa*—c. *Fluctuantes*, see *Costa*—c. *Mendosa*, see *Costa*—c. *Spuria*, see *Costa*—c. *Veræ*, see *Costa*.

COSTAL, *Costalis*, from *costa*, 'a rib.' Appertaining or relating to a rib—as 'costal cartilage.'

COSTE, *Costa*.

COSTIVE, *Constipated*.

COSTIVENESS, *Constipation*.

COSTMARY, *Tanacetum balsamita*.

COSTO, from *costa*, 'a rib.' In composition, relating or appertaining to a rib.

COSTO-ABDOMINAL, *Obliquus externus abdominis*—c. *Basi-scapulaire*, *Serratus magnus*—c. *Claviculaire*, *Subclavian muscle*.

COSTO-CLAVICULAR, *Costo-Clavicula'ris*. Belonging to the ribs and clavicle.

COSTO-CLAVICULAR LIGAMENT, *Gleidocostal Ligament*, is a fibrous, flattened fascia, which extends, obliquely, from the cartilage of the first rib to the inferior surface of the clavicle.

COSTO-CORACOIDIEN, *Pectoralis minor*—c. *Scapulaire*, *Serratus magnus*.

COSTO-STERNAL, *Costo-Sterna'lis*. Relating to the ribs and sternum. The articulation of the sternum with the anterior extremity of the first seven ribs.

COSTO-TRACHELIANUS. Relating to the ribs and to the tracheal or transverse processes of the neck. Under the name *Costo-trache'lian*, *Chaussier* designates the anterior and posterior scaleni.

COSTO-TRANSVERSA'RIUS. Relating to the ribs, and to the transverse processes.—*Bichat* gave this name to the articulation of the tuberosities of the ribs with the transverse processes of the spine.

COSTO-VERTEBRAL, *Costo-vertebra'lis*. Belonging to the ribs and vertebra.—*Bichat* gave this name to the articulation of the head of the ribs with the vertebra, and to the ligaments connected with it.

COSTO-XIPHOID, *Costo-xiphoi'deus*. The name of a ligament, which unites the cartilage of the seventh rib to the xiphoid or ensiform cartilage.

COSTUS. From the Arabic. The ancients seem to have given this name to several plants. One has borne the appellation, since the time of Dioscorides,—the *Costus Arabicus* seu *Indicus* seu *speciosus* seu *amarus*, *dulcis*, *orientalis*, *Amomum hirsutum*, *Helleb'ia grandiflora*, *Bank'sia speciosa*, *Teiana*, (F.) *Canne Congo*, *Canne de Rivière*: *Family*, *Amomeæ*; *Sex. Syst.* *Monandria Monogynia*; the root of which is aromatic, and has been considered tonic, carminative, diuretic, emmenagogue, &c. The virtues of the ancient *costus* are highly extolled.—Theophrastus, Dioscorides, Pliny, Galen.

COSTUS AMARUS, &c., *Costus*—c. *Arabicus*, *Costus*—c. *Corticosus*, *Canella alba*—c. *Hortorum*, *Tanacetum balsamita*—c. *Hortorum minor*, *Achillea ageratum*—c. *Indicus*, *Costus*—c. *des Jardins*, *Balsamita suaveolens*—c. *Nigra*, *Cynara scolymus*—c. *Speciosus*, *Costus*.

COSTYLE, *Cotyle*.

COTA, *Anthemis cotula*.

COTARONIUM; an obscure term, used by Paracelsus for a universal solvent: such a thing as does not exist.

CÔTE, *Costa*.

CÔTES, *COL DES*, *Collum costarum*—c. *Asternales*, see *Costa*—c. *Faussee*, see *Costa*—c. *Flotantes*, see *Costa*—c. *Sternales*, see *Costa*—c. *Vertébrosternales*, see *Costa*—c. *Vraies*, see *Costa*.

COTIGNAC, (pron. *cotinniac*.) A kind of conserve or preserve, prepared from quinces not entirely ripe, and sugar. It is esteemed stomachic and astringent.

COTON, *Gossypium*—c. *Azotique*, see *Collodion*—c. *Fulminant*, see *Collodion*—c. *Poudre*, see *Collodion*.

COTONIA, *Pyrus cydonia*.

COTTON, *Gossypium*.

COTULA, *Anthemis cotula*—c. *Fetida*, *Anthemis cotula*.

COTULA MULTIFIDA. A South African plant, which is used by the Hottentots in rheumatism, scalds and cutaneous diseases.

COTURNIUS, LIQUOR OF, *Liquor Coturni*, *L. of Cotugno*, *Aquila acustica*, *Aqua auditoria*, *Perilymph*, (F.) *Lympha de Cotugno*. A transparent, slightly viscid fluid, which fills all the cavities of the internal ear, and of which Cotugno made mention. It is also called *Aqua Labyrinthi*.

COTURNIX, *Tetrao coturnix*.

COTYLĒ, *Cotyla*, *Cotyle don*, *Oostylē*; the same as *Acetabulum*. A hollow cavity in a bone, which receives the head of another bone: particularly 'the cotyloid cavity.' *Koruly* signified a drinking cup, and, indeed, any thing hollow, as the hollow of the hand.—Athenæus.

COTYLEDON, *Cotyle*—c. *Marina*, *Umbilicus marinus*.

COTYLĒDON ORBICULATA. A plant of South Africa, *Nat. Ord.* *Crassulaceæ*. The fresh juice is used in epilepsy; and Dr. Pappe speaks well of it. The leaves form a good application to hard corns. *Crassula arborea* *cens* has the same properties.

COTYLĒDON UMBILICUS, *C. U. Ven'eria*, *Navelwort*, *Venus's Navelwort*, (F.) *Nomb'ril de Venus*. A plant which grows in Europe on old walls and rocks. The leaves are emollient, and applied externally to piles, inflamed parts, &c. Internally, the juice has been given in epilepsy. The flowers have been used in calculous cases, and in dropsy.

COTYLĒDONS, (*κοτυληδών*, 'the hollow of a cup,') *Acetab'ula uteri'na*. The lobes which, by their union, form the placenta.

COTYLOID, *Cotyloides*, *Cotyloides*, *Cotyloides*, from *κοτυλη*, 'a drinking cup,' and *ειδος*,

'form.' Resembling the ancient *κοτυλη*. The name of a hemispherical cavity, situate in the os innominatum, which receives the head of the os femoris.—*Fossa cotyloidea*, *Sinus Coxæ*, *Acetabulum Pyxis*.

COTYLOID LIGAMENT, *Ligamentum Cotyloideum*, (F.) *Ligament cotyloïdien*, is a very thick, fibro-cartilaginous ring, surrounding the cotyloid cavity, the depth of which it increases.

COTZ, *Hagenia Abyssinica*.

COU, *Collum*—c. *du Pied*, *Instep*—c. *Gros*, *Bronchocele*.

COUCH GRASS, *Triticum repens*.

COUCHE, *Couches*, (F.), from *coucher*, 'to lie down.' This word is used, 1. For *parturition*, *accouchement* or *delivery* (*puerperium*, *parturition*;) hence, *une couche heureuse*, 'a happy delivery'; *une fausse couche*, 'a premature delivery'; and, 2. For the time during which a female remains in bed on account of delivery.—*Tempus puerperii*, the *child-bed state*. The *Lochia* have been termed *Suites de couches*.

Couche also means a layer, bed or thalamus, as, **COUCHE CELLULEUSE**, *Membrana granulosa*—c. *de Malpighi*, *Corpus mucosum*.

COUCHER, *Decubitus*.

COUCHES DES NERFS ETHMOÏDAUX, *Corpora striata*—c. *des Nerfs oculaires*, *Thalami nervorum optico-rum*—c. *des Nerfs optiques*, *Thalami nervorum optico-rum*—c. *Optiques*, *Thalami nervorum optico-rum*.

COUCHING, see *Cataract*.

COUCINEA COCCINEA, *Cacoucia coccinea*.

COUDE, *Elbow*.

COUDE-PIED, *Instep*.

COUDÉE, *Cubitus*.

COUDRIER, *Corylus avellana*.

COUENNE (F.), *Cutis suilla*. A term given to various parts of the human skin, which are prominent, hard, brownish, and often covered with rough hairs, so as to form patches not very unlike the skin of the hog. It is a malformation, occurring during intra-uterine existence, and remaining for life. See, also, *Nævus*, and *Corium phlogisticum*.

COUENNE INFLAMMATOIRE, *Corium phlogisticum*—c. *Pleurétique*, *Corium phlogisticum*.

COUF, *Tussis*.

COUGH, *Tussis*—c. *Bronchial*, *C. tubal*—c. *Root*, *Trillium latifolium*.

COUGH, TUBAL, *Bronchial cough*, *Tussis bronchica*, (F.) *Toux tubaire ou bronchique*. Cough is so termed, when the succussion communicated by it to the parietes of the chest is very energetic, and a sensation is experienced by the ear as if a column of air was traversing tubes with solid walls, with much noise, strength, and rapidity.

COUGH, WINTER, *Bronchitis*, (chronic.)

COUGHING, *Tussis*.

COUL, *Abscess*, *Swelling*.

COULER, *To strain*.

COULEURS COMPLÉMENTAIRES, *Colours*, *accidental*.

COULEUVRE, *Coluber berus*.

COULEUVRÉE, *Bryonia alba*—c. *de Virginie*, *Aristolochia serpentaria*.

COULEUVRINE, *Polygonum bistorta*.

COULISSE (F.), 'A groove, a gutter.' Anatomists designate, by this name, every deep groove or channel in a bone, in which a tendon plays;—such as the *Couliſſe Bicpitale* or *Bicipital Groove* of the *Humerus*.

COULISSE BICPITALE, *Bicipital groove*—c. *Humérale*, *Bicipital groove*.

COULOIR, *Colatorium*.

COUM, *Colchicum autumnale*.

COUMAROUNA ODORATA, Dipterix odorata.

COUN'TENANCE, *Vultus, Voltus, Visage*, (F.) *Figure, Visage*: from *cum*, 'with,' and *teneo*, 'I hold.' The form of the face. The system of the features.

COUNTER-EXTEN'SION, *Contra-exten'sio, Anti'tas*, (F.) *Contre-extension*, from *contra-extendere*, 'to extend in a contrary direction.' It consists in retaining firmly and immovably the upper part of a limb, whilst extension is practised on the lower, in cases of fracture or luxation.

COUNTER-INDICA'TION, *Contra-indica'tio, Antideix'is, Antendix'is, Antideix'is*. An indication contrary to another. (F.) *Contre-indication*. Any circumstance, which acts as an obstacle to the employment of such therapeutical means as seem to be indicated by other circumstances.

COUNTER-IRRITANT, see Counter-irritation—*c. Granville's*, (Lotion), Granville's Counter-irritant.

COUNTER-IRRITA'TION, *Contra-irrita'tio*. An irritation, excited in a part of the body, with the view of relieving one existing in another part. The remedies used for this purpose are called *Counter-irritants*, and form a most valuable class of remedial agents. See Derivation.

COUNTER-POISON, Antidote.

COUNTERSTROKE, *Contra'ssura*.

COUNTING, METHOD OF, Method, numerical.

COUP, Blow—*c. de Feu*, Wound, gunshot.

COUP DE MAÎTRE, *Tour de Maître* (F.). A masterly stroke or performance. Applied to a mode of introducing the sound or catheter into the bladder, which consists in first passing it with its convexity towards the abdomen of the patient, and giving it a half turn towards the right groin, when its extremity has reached the root of the penis under the symphysis pubis. There is no advantage in this mode of introduction.

COUP DE SANG (F.), *Blood-stroke*. A common term, used by some physicians in France, to designate the loss of sensation and motion, which results from hemorrhage in the brain, or from simple congestion in the vessels of that organ. See Apoplexy. Some authors have comprehended, under the same denomination, different hemorrhages, which occur in the areolar texture of the face, lungs; skin, &c.

COUP DE SOLEIL (F.), *Siri'asis, Seiri'asis, Sîriasis Egyptiaca, Ictus solis seu solar'is, In-sola'tio, Encephali'tis insolatio'nis, Phreni'tis calentu'ra, Stroke of the sun, Egyptian Starstroke or Sunstroke*. An affection produced by the action of the sun on some region of the body;—head, hands, arms, &c. A very common effect of exposing the naked head to the sun is inflammation (?) of the brain or its meninges, *Helien-cephali'tis*, which Sauvages calls *Carus ab Insolatio'nē, Morbus solstitial'is*. It has, also, been called *heat apoplexy*; although the morbid phenomena observable after death are generally not confirmatory of either inflammation or apoplexy.

COUP DE VENT. A windstroke. The aggregate of morbid phenomena ascribed to exposure to a blast of air.

COUPE-BRIDE, Kiotome.

COUPEROSE, *Gutta rosea*.

COUPURE, Cut, Wound (incised.)

COURALIUUM, Coral.

COURANTE, Diarrhoea.

COURAP. A distemper, very common in India, in which there is a perpetual itching of the surface, and eruption. It is of an herpetic character, and appears chiefly on the axilla, groins, breast, and face.

COURBARIL, see Anime.

COURBATURE (F.), *Acer'ba lassitu' do, Violent lassitude, Fatigue*. An indisposition, characterized by a sensation, as if the limbs were bruised; general feeling of debility, extreme lassitude; and, sometimes, slight fever. It appears immediately after severe exercise, but sometimes not till the next day. Rest removes it.

COUREBURE, Curvature.

COUREGE, *Cucurbita pepo*.

COURMI or CURMI, *kouput, kuput*. A fermented liquor, made from barley. A kind of ale or beer.—*Dioscorides*.

COURO-MOELLI. An Indian tree, the bark of which is said to be anti-venomous.

COURON'DI, *Couron'do*. An evergreen tree of India, the juice of which, mixed with warm whey, is said to cure dysentery.—*Ray*.

COURONNE, Crown—*c. Ciliare, Godronné canal—c. du Dent*, Crown of a tooth—*c. du Gland*, Crown of the glans—*c. de Moine*, Leontodon taraxacum—*c. de Saint-Jean*, *Artemisia vulgaris*—*c. du Trépan*, Crown of the trepan—*c. de Vénus*, Crown of Venus.

COURONNE RAYONNANTE (F.) A term given by Reil to the fan-shaped terminations of the cruri cerebri in the brain.

COURONNEMENT (F.), *Coro'na*. A vulgar expression, sometimes used to designate the circular ring, which the os uteri forms around the head of the child at a certain period of accouchement—the head seeming to be surrounded, as it were, by a crown. The head is then said to be au couronnement.

COURS DE VENTRE, Diarrhoea.

COURSE, Running.

COURSES, Menses.

COURT, Short.

COURTE HALEINE, Dyspnoea.

COUSSIN, Culex.

COUSSINET, Pad.

COUSSO, *Hagenia Abyssinica*.

COUTEAU, Knife—*c. à Amputation*, Knife, amputation—*c. à Cataracte*, Knife, cataract—*c. Courbe*, Knife, crooked—*c. à Crochet*, Knife, crochet—*c. Désarticulateur*, Knife, double-edged—*c. Droit*, Knife, amputation—*c. de Feu, Couteil-laïre—c. Interosseux*, Knife, double-edged—*c. Lenticulaire*, Knife, lenticular—*c. Lithotome*, Knife, lithotomy—*c. en Serpette*, Knife en serpette—*c. Symphysien*, see Symphysectomy—*c. à deux Tranchans*, Knife, double-edged.

COUTOU'BEA ALBA, *Ex'acum spica'tum, P'crium spica'tum*. A plant of Guiana, which is very bitter, and is considered, in the country, to be emmenagogue, anthelmintic, and antidyspeptic.

COUTURES PAR LA PETITE VÉROLE, see Cicatrix.

COUTURIER, Sartorius.

COUVERCLE, see Crucible.

COUVERCHEF (F.), *Cucul'lus, Fascia'tio cuculla'ta, Seepaster'ium, Scepas'trum*. A bandage, applied to the head for retaining certain dressings, &c., *in situ*. The French surgeons distinguish two kinds: 1. The *Grand Couverchef* or *Servette en carré* (*Cucul'lus major*), which is formed of a napkin or large square compress; and, 2. The *Petit Couverchef* or *Mouchoir en triangle* (*Cucul'lus minor*), formed of a napkin or other square compress, folded from one angle to the other diagonally opposite to it.

COUVERCHEF, see Bandage, Galen's.

COVE, CLIMATE OF. On the northern side of Cork Harbour, in Ireland, is the Island of Cove. The town of Cove is on the southern acclivity of a hill running from east to west. It is, consequently, sheltered from the north winds.

and receives the full force of the sun. It is one of the mildest climates in Great Britain, and corresponds in its influence on disease with the south-west of England. It is well adapted as a winter retreat for the phthisical.

COVOLAM, *Crataeva Marmelos*. The bark of this tree is tonic; the unripe fruit is astringent; but, when ripe, is delicious.

COWBANE, *Cicuta aquatica*—c. Spotted, *Cicuta maculata*.

COWBERRY, *Vaccinium vitis idæa*.

COWCUMBER, *Cucumis sativus*.

COWDEL, Caudle.

COWHAGE, *Dolichos pruriens*.

COWITCH, *Dolichos pruriens*.

COWLADY, *Coccinella septempunctata*.

COWMUMBLE, *Heracleum spondylium*.

COWPARSNIP, *Heracleum lanatum*, H. spondylium.

COWPER'S GLANDS, *Accessory glands*, (F.) *Glandes accessoires ou de Cowper, Prostates inférieures ou Petites prostates*. Called after their discoverer. Two small ovoid groups of mucous, reddish follicles, situate behind the bulb of the urethra, before the prostate, the excretory ducts of which open into the bulbous portion of the urethra. They are also called *Glandulæ anteprosatæ*, and *G. prostatæ inferiores*. The terms *anterior prostate*, and *anteprostatic gland*, are generally, however, given to a third glandular body, occasionally found in front of, and between, Cowper's glands.

COWPER'S GLANDS IN THE FEMALE, *Glandulæ Couperi, Glands of Duverney or of Bartholin, Prostatæ muliebres seu Bartholini, Glandes de Couper ou de Méry, G. vulvaire conglomérée ou vulvo-vaginale, Corps folliculaire vaginal*, are situate at each side of the entrance of the vagina, beneath the skin covering the posterior or inferior part of the labia. They are rounded, but elongated, flat, and bean-shaped; their long diameter varying from five to ten lines; their transverse, from two and a half to four and a quarter; and their thickness from two and a quarter to three lines. Like Cowper's glands in the male, they are not invariably present. The secretion from them is a thick, tenacious, grayish-white fluid, which is emitted in great quantity during sexual intercourse; and is probably the fluid supposed, of old, to be the female sperm.

COWPOX, *Vaccina*—c. Inoculation, Vaccination—c. Itch, see Itch, cowpox—c. Vesicle, see Vaccination.

COWRAP, Impetigo.

COWSLIP, *Caltha palustris*, *Primula veris*—c. Jerusalem, *Pulmonaria*.

COWSTRIPLINGS, *Primula veris*.

COWSTROPPE, *Primula veris*.

COWTHWORT, *Matricaria*.

COWTREE, *CEYLON, Gymneura lactiferum*.

COWTREE, OF SOUTH AMERICA, *Galactodendron utile*.

COWWEED, *Chærophyllum sylvestre*.

COXA, Haunch.

COXÆLUVIUM, Bath, hip.

COXAGRA, *Neuralgia femoro-poplitæa*.

COXAL'GIA, *Merocoxal'gia, Osphyal'gia, Osphalgema*. A word of hybrid origin, from *coxa*, 'hip,' and *algos*, 'pain.' Pain in the hip. A sign of rheumatic or other inflammation—*Coxitis*, or of some disease about the hip-joint. See *Neuralgia femoro-poplitæa*, and *Coxarum morbus*.

COXARTHROSIS, *Coxitis*.

COXARTHROACE, *Coxarum morbus*.

COXARUM MORBUS, *Coxen'dicus seu Coxarius morbus, Arthroacæ coxarum, Coxarthroacæ, Ospharthroacæ, Coxal'gia* (of some), *Hip Disease*. A scrofulous caries, and often spontaneous luxation of the head of the os femoris, occasioning permanent shortening of the limb, and not unfrequently hectic and death—*Ischiophthisis*.

COXEN'DIX, Haunch. This word has been used synonymously with ischium; and anatomists have also applied it to the ilia, *Ossa Coxen'dicia*.

COXITIS, *Coxarthritie*, a hybrid term, from *Coxa*, 'the hip,' and *itis*, denoting inflammation. Inflammation of the hip-joint.

COXO, in composition. Relating or appertaining to the coxal bone.

COXO-FEMORAL, *Coxo-femoralis*. Belonging to the coxal bone or ileum, and to the os femoris. *Ilio-femoral* has the same signification.

COXO-FEMORAL ARTICULATION, *Ilio-Femoral A.*, (F.) *Articulation de la Hanche, Hip-joint*, (Prov.) *Hanch, Locking, Lonk*. The head of the femur and the articular cavity are covered by a *diarthrodial cartilage and synovial capsule*; and by a very strong capsular or articular ligament attached to the circumference of the cotyloid cavity, and to the neck of the femur. There is, also, a *round or interarticular ligament*, passing from the inferior notch of the cotyloid cavity to the rough depression at the top of the caput femoris; and a *cotyloid ligament*—a sort of thick, cartilaginous ring, situate on the edge of the cotyloid cavity, and converting the inferior notch into a foramen. These are the great means of union in this extensive articulation.

CRAB, *Asiacus fluviatilis, Cancer, Gammarus, Gambarus, Cammarus*, (F.) *Crabbe, Écrevisse, Escrèce*. Class, Crustacea. A shell-fish much used as an article of diet. Like other shell-fish, it is apt to disagree, and excite urticaria, &c.

CRABE, Crabyaws.

CRABLOUSE, *Pediculus pubis*.

CRABS' EYES, *Cancrorum chelæ*—c. Stones, *Cancrorum chelæ*.

CRABYAWS, (F.) *Crabe*. A name in the West Indies for a kind of ulcer on the soles of the feet, with edges so hard, that they are difficult to cut.

CRACHAT, Sputum.

CRACHEMENT, Excreation—c. *de Pus, Vomica*, Phthisis pulmonalis—c. *de Sang, Hæmoptysis*.

CRACHOTEMENT (F.), from *cracher*, 'to spit.' The frequent spitting of a small quantity of saliva.

CRACKED POT SOUND, *Bruit de pot fêlé*.

CRACKLING OF LEATHER, see *Craquement de cuir*.

CRACOW GROATS, *Semolina*.

CRADLE, *Sax. opavel, Arculus, Solen*, (F.) *Arceau on Archet*. A semicircle of thin wood, or strips of wood, used for preventing the contact of the bed-clothes in wounds, fractures, &c. An ordinance of the Grand Duke of Tuscany forbade mothers and nurses to sleep with a child near them, unless it was placed under a solid cradle of this kind, in order that no accident might arise from overlaying.

CRÆPALE, from *κραπαλη*, *Cræ'ula*, 'drunkenness, surfeit.' A derangement of the functions of the brain, &c., produced by wine or any other fermented liquor.—Galen.

CRAGBANE, *Clavicle*.

CRAIE, *Creta*—c. *Barotique, Baryta, carbonate of—c. Magnésienne, Magnesias carbonas—c. de Plomb, Plumbi subcarbonas*.

CRAISEY, *Ranunculus acris*.

CRAMA, *κραμα, from κραω, 'I mix.'* A mixture of any kind. Dilute wine.

CRAMBE, Cabbage.

CRAMBION, from *κραμβη*, 'cabbage.' A decoction of cabbage.—Hippocrates.

CRAMERIA, Krameria.

CRAMP, from (G.) *krampfen*, 'to contract.' *Crampus*, *Enta'sia systrem'ma*, *Tetanus dolorif'icus*, *Myospasmus*, *Spasmus muscula'ris*, *Myul'gia*, *Rheg*, *Rhegma*, *Rhegmus*. (F.) *Crampe*. A sudden, involuntary, and highly painful contraction of a muscle or muscles. It is most frequently experienced in the lower extremities, and is a common symptom of certain affections—as of *Colica Pictorum* and *Cholera Morbus*. Friction and compression of the limb, by means of a ligature applied around it above the muscles affected, will usually remove the spasm.

CRAMP, SCRIVENERS', Cramp, writers'.

CRAMP OF THE STOMACH, (F.) *Crampe de l'Estomac*. A sudden, violent, and most painful affection of the stomach, with sense of constriction in the epigastrium. It seems to be the effect of the spasmodic contraction of the muscular coat of the stomach, and requires the use of the most powerful stimulants and antispasmodics, of which opium is the best.

CRAMP, WRITERS', *Graphospasm'us*, *Mogigraph'ia*, (F.) *Crampe des Écrivains*, *Scriv'eners' Cramp*, *Stam'mering of the Fingers*. A condition of the fingers, in which they are unable to hold the pen, or in which one or more of the muscles of the fingers are irregularly and irresistibly contracted.

CRAMPE DES ÉCRIVAINS, Cramp, writers'.

CRAMPE DE L'ESTOMAC, Cramp of the stomach.

CRAN DE BRETAGNE, Cochlearia armoracia.

CRANBERRY, *Vaccinium oxycoccus*—c. Alpine, *Vaccinium vitis idæa*—c. American, common, see *Vaccinium oxycoccus*—c. Upland, *Arbutus uva ursi*.

CRÂNE, Cranium—c. *Humain*, Cranium humanum.

CRANE'S BILL, *Geranium*—c. Spotted, *Geranium maculatum*—c. Stinking, *Geranium Robertianum*.

CRANIAL, *Crania'lis*, *Crania'nus*, (F.) *Crânien*; same etymon as cranium. Relating or appertaining to the cranium—as 'cranial nerves.'

CRÂNIEN, Cranial.

CRANIOCELE, Encephalocele.

CRANIOHÆMATONCUS, Cephalæmatoma.

CRANIOLOGY, *Cranios'copy*, *Craniology'ia*, *Cranionan'cy*, *Phrenol'ogy*, *Cranioscop'ia*, *C. Gallia'na*, *Encephaloscop'ia*, *Doctri'na Gallia'na*. (F.) *Cranilogie*, *Cranioscopie*. Words introduced, since Gall, into medical language. The first two terms are respectively derived from *κρανιον*, 'cranium,' *λογος*, 'a discourse,' and *σκοπειν*, 'to examine.' They signify a description, or simply an examination, of the different parts of the external surface of the cranium, in order to deduce from thence a knowledge of the different intellectual and moral dispositions. Strictly speaking, it is by *Cranioscopy* that we acquire a knowledge of *Craniology*, *Organology*, or *Cranology*, as it has been variously termed. These words are generally, however, used in the same sense. The cranium being moulded to the brain, there are as many prominences on the bone, as there are projections at the surface of the brain. According to Gall, each projection, which he calls an *organ*, is the seat of a particular intellectual or moral faculty, and all persons endowed with the same faculty, have, at the same part of the brain, a prominence, which is indicated, externally, by a bump or projection in the bony case. The *System of Gall* is made to comprise 27 prominences, which answer to 27 faculties. The following Table exhibits these supposed organs, and their seat:

CEREBRAL ORGANS AND THEIR SEAT, ACCORDING TO GALL.

1. *Instinct of generation, of reproduction; amateness; instinct of propagation; venereal instinct.*
(German.) *Zeugungstrieb*, *Fortpflanzungstrieb*, *Geschlechtstrieb*.
2. *Love of progeny; philoprogenitiveness.*
(G.) *Jungenliebe*, *Kinderliebe*.
3. *Attachment, friendship.*
(G.) *Freundschaftsinn*.
4. *Instinct of defending self and property; love of strife and combat; combativeness; courage.*
(G.) *Muth*, *Raufsinn*, *Zanksinn*.
5. *Carnivorous instinct; inclination to murder; destructiveness; cruelty.*
(G.) *Wurgsinn*, *Mordsinn*.
6. *Cunning; finesse; address; secretiveness.*
(G.) *List*, *Schlaueit*, *Klugheit*.
7. *Desire of property; provident instinct; cupidity; inclination to robbery; acquisitiveness.*
(G.) *Eigenthumssinn*, *Hang zu stehlen*, *Einsammlungssinn*, *Diebsinn*.
8. *Pride; haughtiness; love of authority; elevation.*
(G.) *Stolz*, *Hochmuth*, *Hohensinn*, *Herrschaftsinn*.
9. *Vanity; ambition; love of glory.*
(G.) *Eitelkeit*, *Ruhmsucht*, *Ehrgeiz*.
10. *Circumspection; foresight.*
(G.) *Behutsamkeit*, *Vorsicht*, *Vorsichtigkeit*.
11. *Memory of things; memory of facts; sense of things; educability; perfectibility; docility.*
(G.) *Sachedächtniss*, *Erziehungsfähigkeit*, *Sachsinn*.

Seated in the cerebellum. It is manifested at the surface of the cranium by two round protuberances, one on each side of the nape of the neck.

Indicated at the external occipital protuberance.

About the middle of the posterior margin of the parietal bone, anterior to the last.

Seated a little above the ears, in front of the last, and towards the mastoid angle of the parietal bone.

Greatly developed in all the carnivorous animals; forms a prominence at the posterior and superior part of the squamous surface of the temporal bone, above the mastoid process.

Above the meatus auditorius externus, upon the sphenoidal angle of the parietal bone.

Anterior to that of cunning, of which it seems to be a prolongation, and above that of mechanics, with which it contributes to widen the cranium, by the projection which they form at the side of the frontal bone.

Behind the top of the head, at the extremity of the sagittal suture, and on the parietal bone.

Situate at the side of the last, near the posterior internal angle of the parietal bone.

Corresponds to the parietal protuberance.

Situate at the root of the nose, between the two eyebrows, and a little above them.

12. *Sense of locality; sense of the relation of space; memory of places.*
(G.) Ortsinn, Raumsinn.
13. *Memory of persons; sense of persons.*
(G.) Personensinn.
14. *Sense of words; sense of names; verbal memory.*
(G.) Wordgedächtniss, Namensinn.
15. *Sense of spoken language; talent of philology; study of languages.*
(G.) Sprachforschungssinn, Wort-sinn, Sprachsinn.
16. *Sense of the relations of colour; talent of painting.*
(G.) Farbensinn.
17. *Sense of the relations of tones; musical talent.*
(G.) Tonsinn.
18. *Sense of the relations of numbers; mathematics.*
(G.) Zahlensinn.
19. *Sense of mechanics; sense of construction; talent of architecture; industry.*
(G.) Kunstsinn, Bausinn.
20. *Comparative sagacity.*
(G.) Vergleichender Scharfsinn.
21. *Metaphysical penetration; depth of mind.*
(G.) Metaphysischer Tiefsinn.
22. *Wit.*
(G.) Witz.
23. *Poetical talent.*
(G.) Dichtergeist.
24. *Goodness; benevolence; mildness; compassion; sensibility; moral sense; conscience; bonhommie.*
(G.) Gutmüthigkeit, Mitleiden, moralischer Sinn, Gewissen.
25. *Imitation; mimicry.*
(G.) Nachahmungssinn.
26. *God and religion; theosophy.*
(G.) Theosophisches Sinn.
27. *Firmness; constancy; perseverance; obstinacy.*
(G.) Stetigkeit, Festersinn.

The first nineteen of these, according to Gall, are common to man and animals; the remaining eight, man possesses exclusively. They are, consequently, the attributes of humanity.

ORGANS ACCORDING TO SPURZHEIM.

1. Amativeness.—2. Philoprogenitiveness.—3. Inhabitiveness.—4. Adhesiveness or Attachment.—5. Combativeness.—6. Destructiveness.—7. Constructiveness.—8. Acquisitiveness.—9. Secretiveness.—10. Self-esteem.—11. Love of Approbation.—12. Cautiousness.—13. Benevolence.—14. Veneration.—15. Firmness.—16. Conscientiousness or Justice.—17. Hope.—18. Marvelousness.—19. Wit.—20. Ideality.—21. Imitation.—22. Individuality.—23. Form.—24. Size.—25. Weight and resistance.—26. Colour.—27. Locality.—28. Numeration.—29. Order.—30. Eventuality.—31. Time.—32. Melody or Tune.—33. Language.—34. Comparison.—35. Causality.

A fundamental principle with the *Craniologists* or *Phrenologists* is,—that the brain does not act as a single organ, but that it consists of a plurality of organs; but, were we able to admit this, the assignment of the seat of different faculties could not but be considered premature.

CRANIOMANCY, Craniology.

CRANIOMETRY, *Craniometria*; from *κρανιον*, 'the cranium,' and *μετρον*, 'measure.' Measurement of the skull. A form of compasses

Answers to the frontal sinus, and is indicated externally by two prominences at the inner edge of the eyebrows, near the root of the nose, and outside of the organ of memory of things.

At the inner angle of the orbit.

Situate at the posterior part of the base of the two anterior lobes of the brain, on the frontal part of the bottom of the orbit, so as to make the eye prominent.

Also at the top of the orbit, between the preceding and that of the knowledge of colour.

The middle part of the eyebrows, encroaching a little on the forehead.

A little above and to one side of the last; above the outer third of the orbitar arch.

On the outside of the organ of the sense of the relation of colour, and below the last.

A round protuberance at the lateral base of the frontal bone, towards the temple, and behind the organs of music and numbers.

At the middle and anterior part of the frontal bone, above that of the memory of things.

In part confounded with the preceding. Indicated at the outer side of the last by a protuberance, which gives to the forehead a peculiar hemispherical shape.

At the lateral and outer part of the last; and giving greater width to the frontal prominence.

On the outer side of the last; divided into two halves by the coronal suture.

Indicated by an oblong prominence above the organ of comparative sagacity; almost at the frontal suture.

At the outer side of the last.

At the top of the frontal bone and at the superior angles of the parietal bone.

The top of the head; at the anterior and most elevated part of the parietal bone.

used for this purpose is called a *craniometer*, (F.) *Craniomètre*.

CRANIOPATHIA; from *κρανιον*, 'the cranium,' and *παθος*, 'disease.' Disease of the cranium.

CRANIOPLASTY, *Cranioplasticæ*, from *κρανιον*, 'the skull,' and *πλασσω*, 'I form.' An operation for the restoration of the soft parts and bones of the cranium.

CRANIOSCOPY, Craniology.

CRANIO-SPINAL, Cephalo-spinal.

CRANIO-SPINAL Axis, see Encephalon.

CRANIOSTENOSIS, see Microcephalus.

CRANIOTABES, from *κρανιον*, 'the skull,' and *tabes*, 'wasting.' A softening of the bones of the cranium, and a consequent thinness of those bones,—as the occiput, (*soft occiput*),—which are much exposed to pressure. It is a variety of rickets.—Elsässer.

CRANIOTOMY, *Craniotomia*, (F.) *Craniotomie*; from *κρανιον*, 'the cranium,' and *τομή*, 'incision.' The operation of opening the head in parturition.

CRANIOTOMY FORCEPS. An instrument, resembling the lithotomy forceps, for laying hold of and breaking down the bones of the head in parturition.

CRANIUM, *Cra'nion*, *Cranum*, *Calog'ria*, *Sca'phion*, *Conchus*, *Calva*, *Olla cap'itis*, *Theca cer'e'bra*, *Brain-pan*, (Old Eng.) *Pannikell*, (Prov.)

Harnpan, the skull; from *κρανος*, 'a helmet,' or from *κρανον*, 'head,' (F.) *Crâne*. The collection of bones which form the case for lodging the brain and its membranes, as well as their vessels, and some of the nerves. These bones are eight in number—the *frontal*, *occipital*, two *parietal*, two *temporal*, the *sphenoid*, and *ethmoid*. Besides these, there might be considered, as belonging to the cranium, the *cornua sphenoidal*, the bones of the ear, and the *ossa Wormiana*. Dr. Prichard has characterized the primitive forms of the skull according to the width of the *Bregma*, or space between the parietal bones: 1. The *Stenobregmate*, (*στενος*, 'narrow,') or Ethiopian variety. 2. The *Mesobregmate*, (*μεσος*, 'middle,') or Caucasian variety; and 3. The *Platybregmate*, (*πλατος*, 'broad,') or Mongolian variety.

CRANIUM HUMANUM, *Human Cranium* or *Skull*. (F.) *Crâne humain*. This was anciently much used in prescriptions, and was considered anti-epileptic, alexipharmic, antiloimic, &c.

CRANOMANCY, *Cranomanti'a*, from *κρανον*, 'the head,' and *μαντεια*, 'divination.' (F.) *Cranomancie*. The art of divining—from the inspection of the head or cranium—the moral dispositions and inclinations of individuals.

CRANSON, *Cochlearia armoracia*, *C. officinalis*. **CRANSSAC**, **MINERAL WATERS OF**. *Cranssac* is a village in the department of Aveyron, six leagues from Rhodes, which possesses acidulous chalybeate springs, that have been known for a long time.

CRANTER, from *κραναιν*, 'to finish,' 'render perfect.' The *Dens Sapien'tiæ* has been so called.

CRANUM, *Cranium*.

CRAPULA, *Crepale*.

CRAQUEMENT DE CUIR NEUF (F.), *Necrocyte'chus*, 'crackling of new leather.' A sound like the crackling of new leather, sometimes heard on examining the heart with the stethoscope, which is symptomatic of pericarditis.

CRAQUEMENT PULMONAIRE, (F.) 'Pulmonary crackling.' This *bruit* or sound consists in a succession of small cracklings, heard during inspiration, and almost always at the top of the lung. It is heard at the commencement of phthisis.

CRASIS, from *κραννυμι*, 'I mix.' A mixture of the constituents of a fluid; as the *crasis of the blood*, *humours*, &c. The word has also been employed in a more extensive signification, as synonymous with *Constitution*, *Temperament*, &c.

CRASPEDON, *Staphyloedema*.

CRASSAMEN'TUM, from *crassus*, 'thick.' The thick part or deposit of any fluid. It is particularly applied to the clot of the blood.

CRASSAMENTUM SANGUINIS, see *Blood*.

CRASSE, (F.) *Dirt* or *impurity*. A sort of *layer* or *enduit*, which covers the skin, where cleanliness is not attended to; and which is sometimes the result of the cutaneous exhalation; at others, of extraneous matter adhering to the surface of the body.

CRASSE'NA. A term by which Paracelsus designated the saline, putrescent, and corrosive principles, which, he thought, gave rise to ulcers and tumours of different kinds.

CRASS'ULA, from *crassus*, 'thick,' owing to the thickness of its leaves. *Sedum telephium*.

CRASSULA ARBORESCENS, see *Cotyledon orbiculata*—c. *Portulacaceæ*, c. *Tetragona*.

CRAS'SULA TETRAGO'NA. A South African succulent plant, *Ord. Crassulaceæ*, which is somewhat astringent. Boiled in milk it is used in diarrhoea. *Crassula portulaca'cea* is said to be used in similar cases.

CRASSUS, *Corpulent*.

CRATÆGUS A'RIA, from *κρατος*, 'strength,' owing to the hardness of the wood. *Mes'pilus*,

M. A'ria, *Aria seu Sorbus*, *S. Aria seu Alpi'na*, *White Bean Tree*, (F.) *Alisier blanc*, *Alouche*, *Alouchier*. The fruit, which is of the size of a small pear, is slightly astringent, and somewhat agreeable to the taste. It has been employed in diarrhoea.

CRATÆVA MARMELOS, *Covolan*.

CRATERAU'CHEN, from *κρατερος*, 'strong,' and *αυχνη*, 'neck.' One with a strong neck: as well as a strong neck itself.

CRATEVÆSIUM, *Sisymbrium nasturtium*.

CRATOS, *Force*.

CRAUTE, **MINERAL WATERS OF**. *Craute* is a village five leagues from Autun, in France, where are mineral springs, containing sulphohydrate of magnesia, chloride of sodium, chloride of lime, sulphate of lime, and carbonate of magnesia.

CRAVATE SUISSE (F.), 'Swiss cravat.' A band of the longitudinal or superficial layer of the muscular fibres of the stomach, along the lesser curvature of the organ, the shape of which it assists in preserving;—so called, on account of a fancied resemblance to a Swiss cravat.

CRAW-CRAW. A species of itch, common on the African coast.—Boyle.

CRAYEUX, *Cretaceous*.

CRAYON NOIR, *Graphites*.

CRAZINESS, *Insanity*.

CRAZY, *Insane*, *Pain*.

CREA, *Shin*.

CREAM, *Cremor seu Flos lactis*, (Prov.) *Ream*, (F.) *Crème*. A thick, unctuous matter, of a yellowish-white colour, and sweet, agreeable taste, which rises to the surface of milk, if kept at rest. It is composed of butter, serum, and casein.

Crème is a name given in France to *bouillies* prepared with farina of different kinds; as the *Cream of Rice*, (F.) *Crème de Riz*, *C. of Barley*, (F.) *C. d'Orge*, &c. The same term, and 'medicinal cream,' (F.) *Crème médicinale*, are used by M. Bérar for a preparation, resulting from the union of yolk of egg and sugar with milk, with or without the addition of medicinal agents.

CREAM, **Rigor**—c. *Cold*, *Ceratum Galeni*, *Unguentum Aquæ rosæ*—c. of *Tartar*, *Potassæ supertartaras*.

CREAS, *Rubeola*.

CREASOTE, *Cre'asote*, *Creaso'tum*, *Kre'asote*, *Creaso'ton*, *Creaso'ton*; *Oxy-hydro-carbure'tum ex o'leo pyroxyli'co para'tum*, (Ph. L.) from *κreas*, 'flesh,' and *ωσνω*, 'a preserver.' A substance discovered not long ago by Reichenbach. It is obtained from tar, by distillation, and appears to be the active antiseptic and medicinal agent in tar-water and crude pyroligneous acid. It is a colourless transparent fluid, of a penetrating and disagreeable odour, and is freely soluble in alcohol and acetic acid. Its taste is bitter. S. g. 1.037. It coagulates albumen, whence its hæmstatic power. It is a most powerful antiseptic; and has been largely administered in hemorrhages both internally and externally. As an external application, it has been used in burns, ulcers, especially those of a sloughing character, chronic cutaneous affections, and has been applied to tapeworm when protruded, &c. Internally, it has been administered in phthisis, vomiting, diabetes mellitus, nervous diseases, chronic glanders, &c. The dose internally is one or two drops. Externally, it is sometimes applied pure; at others, diluted, and commonly with water, (f3ss to f3vj) or in the form of ointment, (f3ss to ʒj of cerate.)

CREASOTE PLANT, *Larrea Mexicana*.

CRE'ATINE, *Cre'atin*, *Creati'na*, *Kre'atine*, from *κreas*, 'flesh.' A nitrogenized crystallizable neutral substance, obtained by the agency of

water and heat in making broths and soups. It does not combine either with acids or alkalies.

By the action of strong acids, creatine is converted into *Creatinine* or *Kreatinine*, a substance which has a strong alkaline reaction, and forms crystallizable salts with acids. It preëxists to a small extent in the juice of flesh; and is found in conjunction with creatine in urine.

CREATININE, see Creatine.

CREATOPHAGUS, Carnivorous.

CREAZOTON, Creasote.

CREEPER, TRUMPET, *Tecoma radicans*—c. Virginia, *Ampelopsis quinquefolia*, *Tecoma radicans*.

CREEPERS, Fidgets.

CREMASON, Pyrosis.

CREMASTER, *Suspende'ulum*, *Eleva'tor Testiculi*, *Musculus Testis* seu *Testicon'dus*. That which suspends; from *κρεμαω*, 'I suspend.' The Greeks designated, by this term, the spermatic cord, or all the parts by which the testicle is, as it were, suspended; but, since Vesalius, it has received its present limited meaning. The cremaster is a very thin, muscular fascia—sometimes hardly perceptible—which detaches itself from the internal oblique muscle; passes through the abdominal ring, and vanishes around the tunica vaginalis; serving to draw up the testicle, and to move it slightly. It has been, also, called *Tunica Brythroi'des* and *Suspensorium Testis*.

CREMASTERIC, *Cremaster'icus*, same etymon as *Cremaster*. Appertaining or relating to the cremaster,—as '*cremasteric artery*,'—a branch of the epigastric.

CRÈME, Chrisma.

CRÈME, Cream—c. *Médecinal*, see Cream—c. *de Soufre*, Sulphur præcipitatum—c. *de Tartre*, Potassæ supertartaras.

CREMER. The common name for a disease frequent in Hungary, which is produced by excess in eating and drinking.—Chomel.

CREMNOI, plural of *κρηνος*, 'a steep bank.' The lips of an ulcer. The *Labia pudendi*.—Hippocrates, Foësius.

CREMNON'CUS, from *κρηνοει*, 'the labia pudendi,' and *ογκος*, 'a tumour.' A swelling of the labia pudendi.

CREMOR LACTIS, Cream—c. *Tartari*, Potassæ supertartaras—c. *Urinæ*, see Urine.

CRENA, *Crenatu'ra*, *Dentic'ulus*, (F.) *Crênelure*. In the plural, the small teeth or projections in the bones of the cranium, by means of which an accurate junction is formed at the sutures. *Crênelure* has also been used in surgery, for the gutter or groove in certain instruments, which is generally intended to secure the passage of cutting instruments; the groove, for example, of a director.

CRENATURA, Crena.

CRÊNELURE, Crena.

CRE'OLE, (S.) *Crillo*; from (S.) *criar*, 'to create or foster.' A native of America, or of the West Indies, whose ancestors emigrated from the Old World, or from Africa. Hence there may be black as well as white creoles.—Tschudi.

CREOPHAGUS, Carnivorous.

CREPALIA TEMULENTA, *Lolium temulentum*.

CREPANELLA, *Plumbago Europæa*.

CREPA'TIO, *Crepatu'ra*, from *crepare*, *crepatum*, 'to make a noise.' The action of bursting any seed by ebullition. *Coque ad crepatu'ram*, 'Boil till it bursts.'

CREPATURA, Hernia.

CREPIDINES, PALPEBRARUM, see Tarsus.

CREP'ITANT, *Crep'itating*, *Crepitans*; same etymon as *Crepatio*. Crackling.

Rule Crépitant Sec à Grosses Bulles, 'dry crackling noise with formation of large bubbles,' is heard in pulmonary emphysema, when the distension becomes greater and greater, and is followed by rupture of the vesicles. The air forcing itself a passage in the interlobular areolar tissue, gives rise to this *Rôle* during inspiration.

Rôle Crépitant Sec de Laënnec, R. vésiculaire. The *crepitant rattle*, *Crepita'tio vesicula'ris*, heard, during respiration, in severe pneumonia and in œdema of the lung; so termed on account of the analogy between the sound and that occasioned by pressing a healthy lung between the fingers. It resembles the sound produced by rubbing slowly and firmly between the finger and thumb a lock of hair near the ear. The *Rhonchus crepitans redux*, (F.) *Rôle crépitant redux*, is the sound heard in respiration coexistent with the resolution of pneumonia. It indicates the return of the cells to the puerious condition.

CREPITATIO, Crepitation—c. *Vesicularis*, Crepitation.

CREPITA'TION, *Crepita'tio*, *Crep'itus*, from *crepitare*, 'to make a noise.' Crackling. Crepitation or *crepitus*, (F.) *Cluquitis*, has been used, in Surgery, to designate the noise occasioned by the friction of fractured bones, when the surgeon moves them in certain directions. When it cannot be heard at a distance, it may be detected by the immediate application of the ear, or by the use of the stethoscope. *Crepitus* or *crackling* is, likewise, met with in cases of gangrene, when air is effused into the areolar membrane—provided the part be carefully examined with the fingers. The same term is used for the cracking of the joints in health or disease.

CREPITUS, Crepitation, Fart—c. *Lupi*, Lycoperdon.

CRESCENTIA, Growth.

CRESCENT'IA CUJETE; called after *Crescentio*, an Italian writer on agriculture; *Cujete*, *Narrow-leaved Calabash Tree*. Ord. Solanææ. The pulp of the fruit of this West India plant is acidulous. It is used in diarrhoea and headach; and, in syrup, for diseases of the chest.

CRESCENTILÆ, Waxing kernels.

CRESERA, see Cribration.

CRESPINUS, *Oxyacantha Galeni*.

CRESS, BITTER, COMMON, Cardamine pratensis—c. Garden, Cardamine sativum—c. Indian, *Tropæolum majus*—c. Meadow, Cardamine pratensis—c. Penny, Thlaspi—c. Swines', Cochlearia coronopus—c. Water, *Sisymbrium nasturtium*—c. Water, marsh, *Nasturtium palustre*.

CRESSES, DOCK, Lapsana—c. *Sciatica*, Lepidium Iberis.

CRESSI, *Sisymbrium nasturtium*.

CRESSON ALÉNOIS, *Lepidium sativum*—c. *Élégant*, Cardamine pratensis—c. *de Fontaine*, *Sisymbrium nasturtium*—c. *des Indes*, *Tropæolum majus*—c. *des Jardins*, *Lepidium sativum*—c. *du Mexique*, *Tropæolum majus*—c. *de Para*, *Spilanthus oleraceus*—c. *des Prés*, Cardamine pratensis.

CREST, Crista—c. of the Ilium, see Crista of the Ilium—c. of the Pubis, see Crista of the Pubis—c. of the Tibia, see Crista of the Tibia—c. of the Urethra, see Crista urethralis.

CRETA, *Car'bonas calcis friab'ilis*, *Car'bonas calca'reus*, *Glimmar'go*, *Melia Terra*, Chalk, Carbonate of Lime, (F.) *Craie*. Native friable carbonate of lime.

CRETA PREPARA'TA, *Car'bonas Calcis præpara'tus*, Prepared Chalk. (Prepared by levigation.) Used externally, as an absorbent; internally, as an antacid. Dose, gr. x to ʒj or more.

CRETA'CEOUS, *Creta'ceus*, (F.) *Crayeux*, from *creta*, 'chalk.' Containing, or relating to, or

having the characters of, chalk; 'as 'cretaceous mixture, cretaceous tubercles.'

CRÊTE, Crista—*c. de Coq*, Crista galli—*c. de l'Ethmoïde* Crista galli—*c. Uréthrale*, Gallinaginis caput.

CRETAFAC'TION, *Cretefac'tio*, from *creta*, 'chalk,' and *facere, factum*, 'to make.' Formation of, or conversion into chalk, as the conversion of tubercles into calcareous or cretaceous concretions; *Calcification*.

CRÊTES DE COQ, see Crista.

CRÊTIN. One affected with cretinism. The word is said to come from *Chrétien*, 'Christian,' because the Crétin, being in a state of idiocy, is incapable of committing any sin [?] See Cagot.

CRÊTINISM, *Cretinis'mus*, *Crêt'inage*, *Cyrt'o'is Cretinis'mus*, *Kretinis'mus*, *Idiot'i'a endem'ica*, *Idiotism'us endem'icus*, *Fatu'itus concall'inus* seu *Alpicola'rum*, *Micrenceph'alon*, (F.) *Crêtinisme*. An epidemic affection in the low, deep, narrow situations of the Valais; in the valley of Aost, Maurienne, a part of Switzerland, the Pyrenees, Tyrol, &c. It is a state of idiocy, commonly accompanied by an enormous goitre, and is often hereditary. The unfortunate crétin is little better than the animals around him. He rarely attains an advanced age. Like idiocy, cretinism has been divided into *complete* and *incomplete*.

CREUSET, Crucible.

CREUX DE L'AISSÉE, see Axilla—*c. de l'Estomac*, *Fosselle du cœur*—*c. de la Main*.

CREUZNACH, MINERAL WATERS OF. The springs of Creuznach are in Germany. They contain iodine, bromine, and the chlorides of sodium and calcium.

CREUZOT, MINERAL WATERS OF. C, is a mountain to the north-east of Mount Cenis. Near it is a saline chalybeate spring.

CREVASSE, (F.) *Rima*, from *crever*, 'to break or crack.' *A crack, a cleft*. The words *crevasse*, *gerçure*, *fissure*, and *rhagade* are often used synonymously for small longitudinal cracks or chaps of a more or less painful character. Sometimes *crevasse* is employed to designate the solutions of continuity or ruptures, which supervene in distended parts, in the urinary passages, uterus, &c.: it is then synonymous with rupture.

CREWDS, Rubeola.

CREWEL, Primula veris.

CRI, Cry—*c. de Cuir*, *Bruit de Craquement*, *Craquement de cuir neuf*.

CRIBLÉ, Cribratus.

CRIBRA, see Cribration.

CRIBRA'TION, *Cribra'tio*, *Coscinis'mus*, from *cribrare*, *cribratum*, 'to sift.' *Sifting*. A pharmaceutical operation, which consists in separating the finer parts of drugs from the coarser. *Sieves*, *Drumsieves*, are used for this purpose. These were formerly called *Cribra*: *Cos'cinoi*. A sieve for separating the bran from meal was termed *Cre'sera*, *Aleuote'is*, *Cribrum Pollina'rium*.

CRIBRA'TUS, *Cribr'o'us*, from *cribrum*, 'a sieve.' (F.) *Criblé*. Having holes like a sieve. *Lame criblée*, *Crib'riform plate*, is the horizontal lamina of the ethmoid bone—so called because it is perforated like a sieve. Through the perforations, *Param'ina cribr'o'sa*, the olfactory nerves pass.

La'mina Cribr'o'sa of Albinus; *Crib'riform la-mel'la*—a circular spot, perforated with small holes, seen when the optic nerve is regarded from the inside, after removing the retina and choroid. From these holes the medullary matter may be expressed.

CRIB'RIFORM, *Cribriform'is*, from *cribrum*,

'a sieve,' and *forma*, 'form.' The ethmoid bone was formerly so called.—*Os Cribrifor'me*.

CRIBRIFORM COMPRESS. A square piece of linen pierced with a number of holes. This is spread with cerate and applied to a suppurating surface, the holes being intended for the escape of the pus.

CRIBRIFORM FASCIA, see Fascia, cribriform—*c. Lamella*, see Cribratus—*c. Plate* of the Ethmoid, see Cribratus.

CRIBROSUS, Cribratus.

CRIBRUM POLLINARIUM, see Cribration.

CRIBRARYTENODES, *Crico-arytenoid*.

CRICELLA'SIA, *Cricela'sia*, *κρικηλασια*, from *κρικος*, 'a circle,' and *ελασσα*, 'I drive.' An exercise with the ancients, which appears to have been the same as the childish play of rolling the hoop.

CRICK IN THE NECK, *Shoot*, (Prov.) *Hod*. A painful rheumatic affection of the muscles of the neck, which causes the person to hold his head to one side in a characteristic manner.

CRICO. In composition, appertaining or relating to the cricoid cartilage.

CRICO-ARYTENOID, *Crico-arytenoïdeus*, *Cricaryteno'des*. Relating to the cricoid and arytenoid cartilages.

CRICO-ARYT'ENOID MUSCLE, *Crico-arytenoïdeus latera'lis*, *Crico-latéro-aryténoidien*. A muscle which proceeds from the lateral part of the superior edge of the cricoid cartilage to the outer and anterior part of the base of the arytenoid cartilage. It carries the arytenoid cartilage outward and forward.

CRICO-ARYT'ENOID, *Poste'rior*; *Crico-aryténoides posti'cus*, *Crico-crêti-aryténoidien*, *Dilatateur postérieur du Larynx*. A small, triangular muscle, seated in the back part of the larynx. It extends from the prominent line, at the middle of the posterior surface of the cricoid cartilage, to the outer and posterior part of the base of the arytenoid cartilage.

CRICO-ARTENOIDEUS SUPERIOR, of Winslow, forms part of the *Arytenoïdeus* of modern anatomists.

CRICO-CRÊTI-ARYTÉNOÏDIEN, *Crico-arytenoid*, posterior—*c. Latéro-aryténoidien*, *Crico-arytenoid*.

CRICO-PHARYNGE'AL, *Crico-Pharynge'us*. Belonging to the cricoid cartilage and pharynx.

CRICO-PHARYNGEUS muscles of Winslow, is a fleshy bundle, which forms part of the *Constrictor Pharyngis inferior*. He calls the other part *Thyro-pharynge'us*, and proposes to call the whole *Thyro-crico-pharyngeus*.

CRICO-THYREOIDES, *Crico-Thyroidean*.

CRICO-THYROIDEAN, *Crico-thyroid*, *Crico-thyro'des*, *Crico-thyroo'des*, *Crico-thyroïdeus*. Belonging to the cricoid and thyroid cartilages.

CRICO-THYROID MEMBRANE is of a fibrous nature, and extends from the upper edge of the cricoid cartilage to the inferior edge of the thyroid.

CRICO-THYROID MUSCLE, *Crico-thyroïdeus*, (F.) *Dilatateur antérieur du larynx*, is a small fleshy bundle, of a triangular shape, at the anterior and inferior part of the larynx. It extends from the outer surface of the cricoid cartilage to the lateral parts of the inferior margin of the thyroid cartilage; and its use is to approximate, anteriorly, the corresponding margins of the cricoid and thyroid cartilages; and thus to stretch the ligaments of the glottis, which it contracts by separating the thyroid cartilage from the arytenoid.

CRICO-THYRO-PHARYNGE'US, *Constrictor pharyngis*.

CRICOID, *Cricoi'deus*, *Cricoi'des*, *Crico des*, *Annula'ris*, *Cymbala'ris*, from *ρικος*, 'a ring,' and *ειδος*, 'form.' Having the form of a ring.

A name given to one of the cartilages of the larynx, *Cartilago innominata*, (F.) *Cartilage anonyme*, which is situate at its lower part, between the thyroid and first ring of the trachea. It is much higher behind than before.

CRICOIDES, Annular, Cricoid.

CRICOS, Ring.

CRIMNO'DES, *Crimnoï'des*, from κρινον, 'coarse meal, bran,' and εἶδος, 'resemblance.' Resembling meal or bran. Applied to urine, which deposits a bran-like sediment, *Crimno'des Uri'na*.

CRIMNON, Façina.

CRINA'LE, from *crinis*, 'hair.' An instrument formerly used to compress in cases of fistula lacrymalis. It has its name from the circumstance of its having at one end a small cushion stuffed with hair.—Scultetus.

CRINA'TUM. A species of fumigation, used by Paulus of Ægina. The roots of lilies entered into the process; hence its name, — from κρινον, 'the lily.'

CRINES, Hair.

CRINIS, Capillus.

CRINKLE-CRANKLE, Wrinkle.

CRINOM'YRON, from κρινον, 'the lily,' and ὑγρον, 'ointment.' *Unguentum liliaceum*, U. *Ægyptium album*, U. *Sus'num*. An ointment, composed of lilies and some aromatic plants.—Gorræus.

CRINON, *Lilium candidum*.

CRINONES, see *Aene punctata*.

CRIOLO, Creole.

CRIOMYX'OS. In antiquity, one who had much mucus flowing from his nasal fossæ, like the ram: from κριος, 'a ram,' and μυξα, 'mucus.'

CRISIS, *Diac'risis*, *Dijudica'tio*, 'decision;' from κρινω, 'I decide.' This word has been used in various acceptations. Some mean by *crisis* of a disease, when it augments or diminishes considerably, becomes transformed into another, or ceases entirely. Some have used the word to signify only the favourable changes which supervene in disease; others, for the change going on in the acme or violence of the disease. Others, again, have given this name only to a rapid and favourable change, joined to some copious evacuation or eruption; whilst others have applied the term to the symptoms that accompany such change, and not to the change itself;—thus including, under the same denomination, the *critical* phenomena and the *crisis*.

CRISPA'TION, *Crispat'ura*, from *crispare*, *crispatum*, 'to wrinkle.' A contraction or spasmodic constriction, which supervenes in certain parts, either spontaneously or by the influence of some morbid cause or therapeutical agent. The capillary vessels of a wound are, by the French, termed *crispées*, when, immediately, after an operation, the blood does not flow from them. The skin is said to be *crispée*, when contracted, and the bulbs of the hair become more prominent. *Crispation* of the nerves is a slight convulsive motion of external or internal parts, much less than that which occurs in convulsion.

CRISTA, *Ambe*, *Ambon*. A crest. (F.) *Crête*. A name given to several bony projections; also, to the clitoris.

The word *Crêtes* (F.), *Cristæ*, is also used, in France, for fimbriated excrescences, which form at the anus, and near the genital organs; and are commonly owing to the syphilitic virus. *Crêtes de Coq* are syphilitic excrescences, resembling, in form, the crest of the cock.

CRISTA GALLI; two Latin words, signifying the comb of a cock, (F.) *Crête de l'ethmoïde*, *C. de Coq*. A flat, triangular process, rising above the cribriform plate of the ethmoid bone, and re-

sembling a cock's comb. It gives attachment to the anterior part of the falx cerebri.

CRISTA OR CREST OF THE ILIUM, is the superior margin of the ilium:—the *Crista* or *Crest* of the *Tibia*, the anterior edge, the shin; and the *Crista* or *Crest* of the *Pubis*, the posterior sharp edge on the upper surface of the bone. Winslow calls the nymphæ, *Crista* of the *clitoris*; and Chaussier, the verumontanum, *Crista Urethralis*, *Crest* of the *Urethra*.

CRISTA INTERNA, Frontal Spine—c. *Sphenoidalis*, Sphenoidal spine—c. *Tibiæ*, Shin—c. *Urethralis*, Gallinaginis caput.

CRISTA VESTIB'ULI, *Eminentia pyramidalis*. A crest which divides the vestibule of the ear into two fossæ, — one inferior and hemispherical, called *Fovea hemispha'rica*, *Sinus rotundus*; the other superior and semi-elliptical, *Fovea elliptica* seu *semi-elliptica* seu *hemis-elliptica*, *Sinus oratus*. Morgagni has described a third groove-like fossa, *Recessus* seu *Fovea sulciformis*, situate at the mouth of the common orifice of the two superior semicircular canals.

CRISTÆ CLITORIDIS, Nymphæ.

CRISTALLIN, Crystalline.

CRISTALLINE, Mesembryanthemum crystallinum.

CRISTALLOIDITIS, Phacitis.

CRITHE, Hordeolum, Hordeum, Penis.

CRITHIDION, Hordeolum.

CRITHMUM MARITIMUM, *Crithmum*, *Cachrys maritima*, *Feniculum maritimum*, *Sonchifera*, (F.) *Passie-pierre*, *Perce-pierre*, *Fenouil marin*, *Bacile*. Family, Umbellifera. Sec. Syst. Pontandria Dignia. A plant which grows on the sea-coasts, has a spicy, aromatic flavour, and is used, pickled, as a condiment.

The *Cauponga* of Brazil is a kind of erithmum.

CRITICAL, *Criticus*, (F.) *Critique*. Belonging to a crisis.

CRITICAL DAYS, *Dies critici* seu *judicatorii* seu *Decretorii* seu *Principes* seu *Radicales* seu *Contemplabiles* seu *Intermediarii*, (F.) *Jours critiques*, are those on which a crisis, it is imagined, is most likely to happen. According to Hippocrates and Galen, the greatest number of fevers terminate favourably on the 7th day, and many on the 11th;—these two days being the most propitious. Next to these come, in order of efficiency, the 9th, 11th, 20th or 21st, 17th, 5th, 4th, 3d, 18th, 27th, and 28th. The sixth day was called by Galen, the *Tyrant*, *τυραννος*, because the crises that happened then were generally unfavourable. After this, the most unfavourable were the 8th, 10th, 12th, 16th, and 19th. The 13th was a sort of neutral day; the crises which happened on it being neither favourable nor unfavourable. Days were, also, divided into *Inter-calary*, on which the crises happened less frequently, and were less complete than on the *critical* or *indicatory*;—and into *vacant* and *non-dec'retory*, on which a crisis hardly ever occurred. According to this division, they were enumerated as follows:

Critical days.....7th, 14th, 20th, 27th, 34th, 40th, 60th, &c.

Indicatory days.....4th, 11th, 17th, 24th, &c.

Intercalary days.....3d, 5th, 6th, 9th, &c.

Non-decretory days..2d, 8th, 10th, 12th, 13th, &c.

Fortunate crises were considered to be indicated by favourable signs appearing three days before.

CRITICAL PERIOD, see *Menses*.

CRITIQUE, Critical.

CROCHET, Crochet.

CROCIDISMUS, Carphologia.

CROCIDIXIS, Carphologia.

CRO'CINUM, from *κροκος*, 'saffron.' Made with saffron; coloured with saffron.

CROCO'DES, from *κροκος*, 'saffron,' and *ειδος*, 'resemblance.' Certain troches into which saffron entered as an ingredient.—Paulus of Ægina.

CROCODI'LEA, *Stercus Lacer'tæ*. The excrements of the crocodile, which the Arabists extolled as a remedy for cutaneous diseases, and which was long used as a cosmetic.

CROCODILIUM, Echinops.

CROCOMAG'MA, from *κροκος*, 'saffron,' and *μαγμα*, 'a kneaded or squeezed mass.' A kind of troch, composed of saffron and spices.—Galen, Dioscorides, Paulus of Ægina, Scribonius Largus.

CROCUS, *C. sativus* seu *officinæ* seu *Austriacus* seu *Orientalis*, *Medic'na Tristit'ia*, *Panicula vegetabilis*, *Zafferan*, *Antina Pulmo'num*, *Javis flus*, *Saffron*, (F.) *Safran*. Order, Iridaceæ. The stigmata, *Crocus* (Ph. U. S.), are the parts used in medicine. They are brought from the East. The odour is aromatic, and the taste aromatic, pungent, and bitter:—the colour deep orange red. Its virtues are yielded to alcohol, wine, vinegar and water. Its operation has been considered stimulant, exhilarating and diaphoretic. It is not much used.

Cake Saffron is sometimes met with. It consists of one part of saffron and nine of marigold, made into a cake with oil, and pressed.

Crocus, Colcothar—c. Antimonii, Oxydum stibii sulphuratum—c. Austriacus, Crocus—c. Germanicus, Carthamus tinctorius—c. Indicus, Carema longa—c. Martis adstringens, Colcothar—c. Martis aperiens, Ferri subcarbonas—c. Martis vitriolatus, Colcothar—c. Metallorum, Oxydum stibii sulphuratum—c. Officinalis, Crocus—c. Orientalis, Crocus—c. Saracenicus, Carthamus tinctorius—c. Sativus, Crocus.

CROC'DISMUS, Carphologia.

CROIL, Nanus.

CROISÉ (F.), *Crucia'tus*, *Crossed*. An epithet given to paralysis, when it attacks the arm of one side and the leg of another.

CROISEMENT, Crossing.

CROISSANCE, Growth.

CROISSANCES, Waxing kernels.

CROISSANTS, Waxing kernels.

CROIX DE MALTE, *Sple'nium Crucia'tum*, *Maltese Cross*, *Cross of Malta*. A compress, having the form of the Maltese cross. It is made of a piece of square linen, folded in four, and divided with scissors from each angle to a small distance from the centre. It is used especially after amputation of the fingers, penis, and limbs.

When the compress is cut in two of its angles only, it is called the *Half Maltese Cross*,—*Demi-croix de Malte*.

CROK, Nanus.

CROMMYON, Allium cepa.

CROMMYOXYREG'MIA, from *κρομμυον*, 'an onion,' *οξύς*, 'acid,' and *πρυγγυι*, 'to break out.' Sour, fetid eructations, exhaling a smell similar to that of onions.

CROPALE, Nerium antidysentericum.

CROOPBACKED, see Hump.

CROOPY, Croupy.

CROPSIA, Chromopsia.

CROSS, MALTESE, *Croix de Malte*.

CROSS-EYE, Strabismus.

CROSSE DE L'ARTÈRE, Arch of the aorta.

CROSSE MITE, *Acarus Crossii*.

CROSSES PALMAIRES, Palmar arches.

CROSSFOOT, see Kylosis.

CROSSING, (F.) *Croisement*, *Métissage*. The union of an animal of one race with one of another, in order to improve the breed. In man, as well as in animals, the union of near relatives

is apt to deteriorate the offspring, and hence crossing is important in his case.

CROSSWORT, *Eupatorium perfoliatum*, *Lysimachia quadrifolia*.

CROTALOPHORUS, *Crotalus horridus*.

CROT'ALUS HOR'RIDUS, *Crotaloph'orus*. The Rattlesnake, *Cobra di Capello*, (F.) *Crotale*. Class, Reptilia, Ord. Ophidia. A venomous reptile of North America. Its poison is virulent. It is so called from the rattle in its tail, (*κραταλον*, 'a rattle,' 'a small bell.')

CROT'APHĒ, *Crotaphium*, from *κροταφος*, 'temple.' *Cephalalg'ia Pulsat'ilis*, *Cephalæ'a Pulsat'ilis*, *Sphygmoceph'alus*. A pulsatory pain, chiefly in the temples, with drumming in the ears.

CROTAPHITE ARTERIES, Temporal arteries.

CROTAPHITES, Temporal muscle.

CROTAPHIUM, Crotaphe.

CROTAPHUS, Temple, Temporal bone.

CROTCHET, *Hamus*. A small hook or crook. (F.) *Crochet*. An obstetrical instrument, whose name indicates its shape, and which is used in the extraction of the fetus, when it becomes necessary to destroy it to expedite delivery. Crotchets are differently formed; some are sharp, others blunt; some contained in a sheath, others naked.

CROTON BENZOE, Benjamin.

CROTON CASCARILLA, *Cortex Eleuther'ia*, *Croton Eleuther'ia* seu *Elute'ria*, *Clut'ia Elute'ria*, *Thus Judæ'rum*, *Chacari'lla cortex*, *Cuscari'lla*, *Elute'ria*, *Gascari'lla*. Order, Euphorbiaceæ. Sex. Syst. Monœcia Monadelphia. (F.) *Quinquina aromatique*, *Cascarille*, *Chacirille*. The bark of *Croton Cascarilla* of the Bahamas, *Cortex Thuris*, *Cascarilla*, (Ph. U. S.), (F.) *Écorce éluthérienne*, has a slightly aromatic odour, and bitterish aromatic taste. The smoke has the odour of musk. The active parts are an essential oil and bitter extractive. They are completely extracted by proof spirit. It is tonic and stomachic. Dose, gr. xv to ʒss or more, in powder.

CROTON ELEUTHERIA, *Croton cascarilla*—c. *Jamaigota*, *C. tiglium*—c. *Lacciferum*, see *Lacca*—c. *Oil*, *Croton tiglium*.

CROTON PSEUDO-CHINA, *Copalche bush*. A South American plant, which yields a bark—*Cortex Copalche* seu *Copalchi*, very like that of *Cascarilla*.

CROTON RACEMOSUM, *Beenel*. A small Malabar tree, whose aromatic root, boiled in oil of sesame, is employed, by the orientals, as a liniment in headach and rheumatism.

CROTON SEEDS, see *Croton tiglium*.

CROTON TIG'LIUM, *C. Jambaly'ia*, *Cudel-Avanacu*. (F.) *Petit Pignon d'Inde*. A Ceylonese plant, every part of which is endowed with medicinal power. The root is a drastic cathartic: the wood, *Lignum Pava'na*, *Pavana wood*, and the seed, have like virtues. The seeds have been long known under the names, *Grana Moluc'cæ*, *Til'li Grana*, and *Grana Tig'lii* seu *Tig'lia*, *Croton seeds*, *Purgum nuts*. From these seeds the *Croton Oil*, *O'leum Tig'lii*, is expressed. It is of a pale, brownish-yellow colour, and hot, biting taste; and is a most powerful drastic cathartic. Dose, from half a drop to three drops, made into pills with crumb of bread. It is also applied externally as a rubefacient, 3 to 5 drops being rubbed on the part; or one part of the oil and three parts of olive oil may be added together, and a little of this be rubbed on.

CROTONĒ, *κροτωνη*. A fungus, which grows on trees, and is produced by an insect, from *κροτων*, 'a tick.' By extension, applied to a fungous tumour developed on the periosteum.

CROUP, *Cynanche trachealis*—c. *Bronchial*, *Polypus bronchialis*—c. *Cerebral*, *Asthma thymi-*

cum.—c. Cerebral spasmodic, Carpo-pedal spasm.—c. Chronic, Polypus bronchialis.—c. *Faux*, Asthma thymicum, see Cyanche trachealis.

CROUP, Hysterical. A spasmodic affection of the laryngeal muscles by no means unfrequent in hysterical females,—the paroxysm consisting in a long protracted, loud and convulsive cough, followed at times by crowing respiration, and by dyspnoea so great as to threaten suffocation. The treatment is that advised for hysteria.

CROPE, Pseudo, Asthma thymicum—c. *Pseudo-nerveux*, Asthma thymicum—c. Spasmodic, see Asthma thymicum, and Cyanche trachealis—c. Spurious, Asthma thymicum.

CROPE, see Croupion.

CROUPION (F.), *Uropygium*, *Orus*, *Orrhos*, the *Rump*. The region of the coccyx. The *Croupier*, (F.) *Croupe*.

CROUP-LIKE INSPIRATION OF INFANTS, Asthma thymicum.

CROUPAL, Croupy.

CROUPOUS, Croupy.

CROUPY, Croupous, Crupo'sus, (F.) Croupal. Relating, or appertaining to, or resembling croup; as a 'croupy cough.' Hoarse. (Prov.) *Croupy*.

CROUPY INFLAMMATION, Inflammatio crupo'sa, is the inflammation of a mucous membrane, attended with a pseudo-membranous exudation.

CROUTE, Crusta—c. *de Lait*, Porrigo larvalis—c. *Laitesse*, Porrigo larvalis.

CROUTEUX, Crustacé.

CROWFOOT, Geranium maculatum—c. Bristly, Ranunculus Pennsylvanicus—c. Bulbous, Ranunculus bulbosus—c. Bulbous-rooted, Ranunculus bulbosus—c. Celery-leaved, Ranunculus sceleratus—c. Crushed, Ranunculus sceleratus—c. Marsh, Ranunculus sceleratus—c. Meadow, Ranunculus acris—c. Small-flowered, Ranunculus abortivus—c. Tall, Ranunculus acris—c. Water, smaller, Ranunculus flammula.

CROWLING, Borborygmus.

CROWN, Coro'na, Steph'ané, (G.) Krone, (F.) Couronne. In anatomy, this name is given to parts of a circular form, which surmount other portions of the same body. Thus, the *Crown of a Tooth*, *Coro'na Dentis*, (F.) *Couronne du Dent*, is the portion of the tooth which projects above the gum.

CROWN OF THE HEAD, Vertex.

CROWN OF THE GLANDS, Coro'na seu Tor'ulus glandis, (F.) *Couronne du gland*, is the round, almost circular, ring, which circumscribes the base of the glands.

CROWN OF THE TREPAN, Modi'olus, (F.) Couronne du Trépan, is a species of saw, in form of a crown, or rather of a portion of a cylinder, having grooves on its external surface, and teeth at its lower extremity; the other being fitted to the handle of the trepan.

CROWN OF VENUS, Coro'na Ven'erie, Gutta Rosacea Syphilitica, (F.) *Couronne de Venus*. Red, rosy pustules, dry or moist, on the face; but particularly on the forehead and temples, owing to constitutional syphilis.

CROWS' BEAK-LIKE PROCESS, Coracoid process.

CRU, Crude.

CRUCIAL, from cruz, crucis, 'a cross.' Crucial'is, Crucia'tus. Having the shape of a cross. Appertaining to a cross.

CRUCIAL BANDAGE, T Bandage.

CRUCIAL INCISION, Incisura crucia'lis, (F.) Incision cruciale. An incision made in the form of a cross. Often employed for exposing the cranium, for the purpose of applying the trepan.

CRUCIAL LIGAMENT OF THE ATLAS, Annular ligament of the Atlas.

CRUCIAL LIG'AMENTS, (F.) Ligaments croisés

on cruciformes. Two strong ligaments within the knee-joint. The anterior passes obliquely from a depression anterior to the spine of the tibia to the posterior and inner part of the external condyle of the femur:—the other, the posterior, extends from the posterior part of the spine of the tibia to the anterior and outer part of the internal condyle.

CRUCIATI DOLO'RES, Excruciating pains.

CRUCIATUS, Croisé.

CRUCIBLE, from cruz, crucis, 'a cross,' so called—it has been supposed—from being made in the shape of a cross, or from having a cross impressed upon it. [?] *Crucibulum, Cati'nus fuso'rius, Tigill'um, Albot, Cementerium, (F.) Creuset.* A vessel of earth, silver, platinum, gold, blacklead, &c., for receiving substances, which have to be exposed to a strong heat. It is sometimes covered with a *top* or *lid*. (F.) *Couvercle*.

CRUCIFORM, from cruz, crucis, 'a cross,' and forma, 'shape,' Cruciform'is, Cross-shaped. A name given to the ligaments which close the articulations of the phalanges; and likewise to the crucial ligaments.

CRUDE, Crudus, Omus, (F.) Cru. Raw, unripe, not concocted.

CRUDITAS MORBI, see Crudity—c. Ventriculi, see Crudity.

CRUDITY, Cru'ditas, Om'otes, Status Crudi'tatis, from crudus, 'crude,' 'unprepared,' Incoctus. Rawness, Crudeness. This has received several acceptations. 1. It expresses the quality of certain aliments, which have not experienced the action of fire. 2. The condition of matters in the digestive tube, which have not undergone the digestive changes—*cru'ditas ventriculi*; and, 3. In the language of the Humorists, it means the condition of the morbid matter in a sick individual, *cru'ditas morbi*, when it has not yet been prepared or concocted by the action of the organs—*Azym'ia humo'rum*.

The word is used in the plural, *Cru'dities*, synonymously with *crude matters*. (F.) *Matidres crues*. It is applied to those, when contained in the stomach and intestines.

CRUE, Growth.

CRUELS, Herpes Zoster, Scrofula.

CRUENTA EXSPUITIO, Hæmoptysis.

CRUES (Matidres), see Crudity.

CRUME'NA VESI'CÆ. The cavity of the urinary bladder.

CRUOR. Coagulated blood, gore. The signification of this word is very vague. It has been used to designate blood in general, venous blood, extravasated or coagulated blood, and the colouring matter.

CRUOR SANGUINIS, see Blood.

CRUORIN, Hæmatin.

CRUPOSUS, Croupy.

CRUPPER, Croupion—c. Bone, Coccyx.

CRUPSIA, Metamorphopsia.

CRURA. The plural of *crus*, 'a leg.' (F.) *Crusæ*. Applied to some parts of the body, from their resemblance to legs or roots; as the *Crura cerebri, Crura cerebelli, &c.*

CRURA ANTERIORA MEDULLÆ OBLONGATÆ, Peduncles of the brain—c. Cerebelli, Peduncles of the cerebellum—c. Cerebelli ad Corpora Quadrigemina, see Peduncles of the cerebellum—c. Cerebelli ad Pontem, see Peduncles of the cerebellum—c. Cerebri, Peduncles of the brain—c. Clitoridis interna, Bulbus vestibuli—c. Medullæ oblongatæ, Corpora restiformia, Thalami nervo-urum opticorum, see Peduncles of the cerebellum—c. Posteriora medullæ oblongatæ, Peduncles of the cerebellum.

CRURÆUS, Cruralis.

CRURAL, Crura'lis, from crus, cruris, 'the

thigh, and lower limb.' What belongs to the thigh or lower limb.

CRURAL ARCH, Inguinal or Femoral Arch, (F.) *Arcade crurale* or *inguinale* (Ch.), *Poupart's Ligament, L. of Fallopius*. This arch is formed by the internal portion of the inferior edge of the aponeurosis of the obliquus externus muscle, which is attached, at one end, to the pubis; at the other, to the anterior and superior spinous process of the ilium. At its posterior and inner part, the aponeurosis, forming the arch, sends off a falciform reflection, which is attached along the crest of the pubis, and is known under the name of *Gimbernats's Ligament*. Beneath this arch, the vessels, nerves, and muscles make their exit from the pelvis to the thigh.

CRURAL ARTERY, Artère Crurale, Femoral Artery, (F.) *Artère crurale, A. Pelvi-crurale*,—(Ch.,) is the continuation of the external iliac. It extends from the crural arch to the aperture in the triceps, or to the ham. Chaussier applies the name *Artère crurale* to the trunk, which extends from the primitive or common iliac to the tibial arteries—embracing, of course, the external iliac, femoral, and popliteal.

CRURAL CANAL, Crural Ring, Femoral canal or ring. M. Jules Cloquet has described this canal with minuteness, and given it the name *Anneau crural, Anneau femoral*. It is nearly an inch long, triangular, more spacious above than below, and shorter and broader in the female than in the male. Its upper orifice is bounded, anteriorly, by the crural arch; posteriorly, by the crista of the pubis; on the outer side by the psoas and iliacus muscles, covered by the iliac aponeurosis, and, at the inner, by Gimbernats's ligament. This orifice is covered by the peritoneum, and, according to M. Cloquet, is closed by a more or less resisting septum, which he has named *Septum crurale, Crural septum*. In its course, the crural canal has its anterior parietes formed by the superficial expansion of the fascia lata: the posterior by the pectineus, covered by the deep-seated expansion of the fascia; and more externally by the psoas and iliacus muscles, covered by an expansion of the fascia iliaca. Its inferior orifice is formed by the foramen of the fascia lata, which gives passage to the vena saphæna. It is at the upper orifice of this canal, that *Femoral or Crural Hernia, Hernie inguinale* of Chaussier, occurs: which would be more common, were it not for the fibrous cellular septum there situate.

CRURAL NERVE proceeds from the lumbar plexus, and is situate at the outer side of the psoas muscle and crural artery. After it has passed under the crural arch, it divides into cutaneous and muscular branches. One of the branches, which is larger than the rest, is called the *Saphæna nerve*. It gives off filaments to the integuments of the knee, to the inner part of the leg, and to the dorsal surface of the foot. The remainder of the branches of the crural are distributed to the anterior and inner part of the thigh. The *Accessory of the Crural Nerve* is a term given to the 4th and 5th pairs of lumbar nerves.

CRURAL PLEXUS of Chaussier is the union of the anterior branches of the last four pairs of lumbar nerves, and the first four sacral; forming the *lumbar and sacral plexuses* of most anatomists.

CRURAL RING, Crural canal—c. Septum, see Crural canal.

CRURAL VEIN, Femoral Vein, has the same arrangement as the artery. It receives only one great branch, the saphæna.

CRURAL, Triceps cruris.

CRURALS, Cruræus. A part of the *Triceps crural* of the French, or of the *Trifémorotulien*

of Chaussier. The cruralis is situate at the anterior, outer, and inner part of the thigh. It arises fleshy, from between the two trochanters, adheres firmly to most of the fore part of the os femoris, and is inserted, tendinous, into the upper part of the patella, behind the rectus. Its use is to assist the vasti and rectus in the extension of the leg. Under *Muscle Triceps Crural*, the French describe the cruralis and the two vasti. Some small muscular slips, sometimes found under the cruræus muscle, and inserted into the capsular ligament of the knee-joint, have been called *Sub-cruræi*.

CRURIS RADIUS, Fibula.

CRUS, Leg, Thigh—c. Fornicis, see Fornix.

CRUST, Eschar—c. Milk, Porrigo larvalis.

CRUSTA. A crust or scab: (F.) *Croûte*. An assemblage of small flakes, formed by the drying up of a fluid secreted by the skin.

The lining membrane of the stomach and intestines has been called *Crusta villosa*.

CRUSTA ADAMANTINA DENTUM, Enamel of the teeth.

CRUSTA GENU EQUINÆ, Hippogonyol'epus, Sweat or Knee Scab, Mock or Encircled Hoof Knees, Hangers, Dew Claws, Night Eyes, Horse Crust. This morbid secretion from the horse has been advised in cases of epilepsy. It is used in the form of powder (gr. ij to gr. xx), and of tincture.

CRUSTA INFLAMMATORIA, Corium phlogisticum—c. Lactea, Porrigo larvalis, P. lupinosa—c. Membranacea, Peristroma—c. Petrosa, see Tooth—c. Phlogistica, Corium phlogisticum—c. Pleuretica, Corium phlogisticum—c. Pruriginosa, Gutta rosea—c. Serpiginosa, Gutta rosea—c. Vermicularis, Peristroma—c. Villosa linguæ, see Tongue—c. Villosa ventriculi, see Stomach.

CRUSTACÆ, (F.) Alibert has substituted this word; in his Nosology, for *croûteux*, 'crusty.' Having crusts or scabs; as *Dartre crustacée*.

CRUSTULA. Diminutive of Crusta. A small shell or scab. An ecchymosis of the conjunctiva.

CRUSTUMINATUM. A rob, prepared from the *Pyra Crustumina* or *Crustumina*, (so called from Crustumium, a town in Italy, where they grew,) boiled with honey or in rain-water.—Aëtius.

CRUT, Nanus.

CRUX CERVI. An ancient appellation for the bone in the heart of the stag. It was once considered useful in diseases of the heart.

CRY, Clamor, Boë, (F.) Cri. The sound of the unarticulated voice. The native voice, which the idiot and deaf possess equally with the man of genius and bearing. The cry of the new-born child has been called *Vagi'us*, (F.) *Vagissement*. We say, "A cry of joy, of pleasure, of pain," &c., according to the expression which it may convey to the hearer.

CRYING, Parturition.

CRYING-OUT, Parturition.

CRYMO'DES, from κρυος, 'cold.' A continued fever, *Febris crymo'des*, in which the internal parts feel hot, and the external cold; and which was attributed to an erysipelatous inflammation of the lungs.—Aëtius, Goræus.

CRYMODYN'IA, from κρυος, 'cold,' and δυνα, 'pain.' Chronic rheumatism, and all its modifications.—Baumes.

CRYMO'SIS, from κρυος, 'cold.' Diseases caused by the action of cold.—Baumes.

CRYPsor'CHIS, from κρυος, 'cold,' and χις, 'a testicle.' *Cryptor'chis, Testicon'dus*. One in whom the testes have not descended into the scrotum. The state is called, *Cryptorchidism, Parorchid'ium*.

CRYPTA, from κρυος, 'concealed.' *Folliculo-se gland, Lacu'na, Follic'ulus*, (F.) *Crypte*,

Follicule. A *crypt* or *follicle* is a small, roundish, hollow body, situate in the substance of the skin or mucous membranes—*Crypta mucosa*, *Folliculus mucosus*—and constantly pouring the fluid, which it secretes, on their surfaces.

The use of the cryptal or follicular secretion, is to keep the parts on which it is poured supple and moist, and to preserve them from the action of irritating bodies with which they have to come in contact.

The little rounded appearances at the ends of the small arteries, in the cortical substance of the kidney, are also called *Crypta*.

CRYPTA MUCOSA, see *Crypta*.

CRYPTÆ LIEBERKÜHNIANÆ, Lieberkühn's glands—*c. Sebaceæ*, Sebaceous glands.

CRYPTE, *Crypta*.

CRYPTOCEPHALUS, from *κρυπτος*, 'concealed,' and *κεφαλή*, 'head.' A monster whose head is excessively small, and does not appear externally.—G. St. Hilaire.

CRYPTOCOCCUS, Fermentum, *Torula cerevisiæ*—*c. Cerevisiæ*, *Torula cerevisiæ*.

CRYPTODIDYMUS, *Fœtus in fœtu*. A monstrosity, in which one fœtus is found contained in another.

CRYPTOGAMOUS, *Cryptogam'ic*, *Cryptogam'icus*; from *κρυπτος*, 'concealed,' and *γamos*, 'marriage.' An epithet applied by botanists to plants whose organs of fructification are concealed or not manifest. *Agamous* plants are those whose sexual organs are not known.

CRYPTOPYIC, *Cryptopy'icus*, from *κρυπτος*, 'I conceal,' and *πυον*, 'pus.' A state of disease, kept up by an occult abscess.

CRYPTOPYICUS, *Cryptopyic*.

CRYPTORCHIDISMUS, see *Crypsorchis*.

CRYPTORCHIS, *Crypsorchis*.

CRYPTS OF LIEBERKÜHN, see *Intestine*—*c. Sænival*, *Bursæ mucosæ*.

CRYSTAL MINERAL, *Potassæ nitras fusus sulphatis paucillo mixtus*.

CRYSTAL'LI, Vesicles filled with a watery fluid. They are also called *crystal'linæ*. Probably the pemphigus of modern writers.

CRYSTALLI LENSÆ, *Argentii nitras*.

CRYSTALLINI, *Crystalline*.

CRYSTALLINA, from *κρυσταλλος*, 'crystal.' A vesicle or phlyctæna, filled with serum, and appearing on the prepuce or in the vicinity of the anus, surrounded by a reddish extravasated aureola. It may be syphilitic or not. See *Crystalli*.

CRYSTALLINA TUNICA, Arachnoid membrane.

CRYSTALLINE, *Crystal'linus*. Having the appearance of crystal.

CRYSTALLINE, *Crys'talline humour* or *Lens* or *Body*, *Crystall'linus*, *Lens cristall'lini* seu *crystall'loides*, *Corpus cristall'linum* seu *Discoides* seu *Crystalloides* seu *Phacoides*, *Humor cristall'linus* seu *glaci'lis*, *Phacæ*, *Phuceæ*, *Phucus*, *Gemma Oculi*, (F.) *Cristallin* ou *Crystallin*, *Humeur cristalline*, *Corps cristallin* seu *phacôide*, *Lentille cristalline*. A lenticular, transparent body, situate between the vitreous and aqueous humours of the eye, at the union of the anterior third with the two posterior thirds of the organ. It is composed of a soft exterior substance; and an interior, composed of an albuminoid substance, called *Crys'talline* or *Krys'talline* by Hünefeld, (F.) *Crystalline*, forming a solid nucleus, in which a number of elliptical layers is perceptible. It is contained in a capsule, called *Tunica ar'nea* seu *crystall'ina*, *Capsula lentis*, and receives, at its posterior surface, a small branch of the central artery of the retina, which is always readily distinguishable in the fœtus, prior to the seventh month of utero-gestation.

The use of the crystalline is to refract the rays of light, and to serve as an achromatic glass: for

which its laminæ or layers, increasing in refractive power from the circumference to the centre, admirably adapt it.

CRYSTALLINO-CAPSULITIS, see *Phacitis*.

CRYSTALLION, *Plantago psyllium*.

CRYSTALLITIS, *Phacitis*.

CRYSTALLOID, *Crystall'loides*, from *κρυσταλλος*, 'crystal,' and *αἶδος*, 'form,' 'resemblance.' Resembling crystal or the crystalline. The capsule or membrane of the crystalline. Also, the crystalline itself.

CRYSTALLUS MINERALIS, *Potassæ nitras fusus sulphatis paucillo mixtus*.

CRYSTALS, BLOOD, *Hæmatoidin*.

CRYSTAUX D'HÉMATINE, *Hæmatoidin*.

CRYPHE, *Hordeolum*.

CTEDON, *Fibre*.

CTEIS, *Pubes*.

CTESTIPHON'TIS MALAG'MA. A plaster employed and described by Celsus.

CUBA, see *Havana*.

CUBAL SINI, *Piper cubeba*.

CUBATIO, *Decubitus*.

CUBEBA, *Piper Cubeba*.

CUBIFORMIS, *Cuboid*.

CUBIT, *Ulna*—*c. Top* of the, *Olecranon*.

CUBITAL, *Cubita'lis*, *Ulnar*, *Ulna'ris*. Connected with or relating to the cubitus, or to the inner and posterior part of the forearm.

CUBITAL ARTERY, *Arte'ria cubita'lis*, *A. ulna'ris*, arises from the humeral a little below the bend of the elbow; proceeds along the anterior and ligament part of the forearm; passes anterior to the ligamentum annulare of the carpus, and goes to form, in the palm of the hand, the superficial palmar arch. Besides the numerous muscular branches, which it gives off in its course, it sends posteriorly the common trunk of the *interosseous arteries*, and internally, the two *cubital recurrents*, anterior and posterior,—articular branches, which anastomose on the inside of the elbow with the divisions of the humeral artery.

CUBITAL MUSCLES are two in number. 1. The *Cubita'lis anterior* seu *inter'nus*, *Flexor carpi ulna'ris*, *Ulna'ris inter'nus*, *Epitrochleo-cubito-carpien*, *Cubito-carpien*—(Ch.), is a long muscle, situate at the anterior and inner part of the forearm. It arises from the inner condyle of the os humeri, at the inner side of the olecranon, and from the posterior edge of the ulna, and is inserted by a tendon into the os pisiforme. Its use is to bend the hand on the forearm, by directing it slightly inwards. 2. The *Cubitalis posterior* seu *exter'nus*, *Extensor Carpi Ulna'ris*, *Ulna'ris exter'nus*, *Cubite'us exter'nus*, *Cubito-supræmetacarpien*—(Ch.), is situate at the posterior part of the forearm. It arises from the external condyle of the os humeri, and is inserted into the superior extremity of the fifth bone of the metacarpus. Its use is to extend the hand, inclining it a little inwards.

CUBITAL NERVE, *Ulnar nerve*, *Cubito-digital*—(Ch.), is furnished by the last two or three nerves of the brachial plexus, and is distributed to the inner and anterior side of the forearm; to the inner part of the palm and of the back of the hand, and to the last two or three fingers.

CUBITAL VEINS, *DEEP-SEATED*, and the *Recurrent cubital veins*, accompany the corresponding arteries. The superficial cubital veins belong to the basilic. Chaussier calls them *cutanéous cubital*.

CUBITALE (OS), *Cuneiform bone*.

CUBITALIS RIOLANI, see *Anconeus*.

CUBITEUS EXTERNUS, see *Cubital muscles*.

CUBITO-CARPIEN, see *Cubital muscles*—*c. Cutané (nerf)*, Cutaneous nerve—*c. Phalangien commun*, *Flexor profundus perforans*—*c. Radi-supræmetacarpien du ponce*, *Abductor longus pollicis*—*c. Radial*, *Pronator radii quadratus*.

CUBITO-SUPRAPALMARIS. Belonging to the cubitus and to the supra-palmar or dorsal surface of the hand. Chaussier gives this name: 1. To a small artery, which is given off by the cubital or ulnar, a little above the wrist. 2. To a vein, which accompanies this artery.

CUBITO-SUS-MÉTACARPIEN, see Cubital muscles—*c. Sus-métacarpien du ponce*, Abductor ongus pollicis—*c. Sus-Phalangien de l'index*, Extensor proprius indicis—*c. Sus-phalangien du ponce*, E. longus pollicis—*c. Sus-phalangien du ponce*, E. pollicis brevis.

CUBITUS, *кубитовъ*, *Cybiton*. The *Elbow*. Also, one of the bones of the forearm. See *Ulna*, and *Forearm*.

CUBITUS, (F.) *Coudée*, the ancient name of a measure 18 inches long.

CUBITUS COSTÆ, see *Costa*—*c. Supinus*, see *Decubitus*.

CUBOID, *Кубои́десъ*, *Cuboi'deus*, *Cubo'des*, *Cubifor'mis*, *Cyboi'des*, *Cyrtoi'des*, *Grandino'sum os*, *Tes'acro*, *Tessera os*, *Os cu'rium*, from *кубосъ*, 'a cube,' and *αἰός*, 'form.' Having the form of a cube. This name was given by Galen to one of the bones of the tarsus, and is still retained. It is situate at the anterior and outer part of the tarsus; and is articulated, behind, with the calcaneum; before, with the last two metatarsal bones, and within, with the third os cuneiforme, and sometimes with the scaphoides. Its inferior surface has an oblique groove for the tendon of the *peroneus longus*.

CUCKOLD, *Arctium lappa*.

CUCKOO FLOWER, *Cardamine pratensis*.

CUCKOW BREAD, *Oxalis acetosella*.

CUCKOW PINT, *Arum maculatum*.

CUCC'BALUS BEHEN, *Behen officinarum seu vulg'ris*, *Silēnē inflat'a seu crassifolia seu Tho'rei*, *Visca'go behen*. This plant was once considered alexipharmic and cordial. It is the *Spatling Poppy*.

CUCULLARIS, *Trapezius*.

CUCULA'TUM MAJUS. A barbarous term, used by Ruland, for brandy and spirit of wine.

CUGULLATA, *Sanicula*.

CUGULLUS, *Couvrechef*, *Cucupha*, *Infundibulum* of the cochlea.

CUCULUS, *Coccyx*, *Pertussis*.

CUCUMBER, *Cucumis sativus*—*c. Asses*, *Momordica elaterium*—*c. Indian*, *Medeola Virginica*—*c. Squirting* or wild, *Momordica elaterium*—*c. Star*, one-seeded, *Sycios angulatus*—*c. Tree*, *Magnolia acuminata*.

CUCUMER, *Cucumis sativus*.

CUCUMIS, *Penis*—*c. Agrestis*, *Momordica elaterium*.

CUCUMIS COLOCYN'THIS, *Citrullus 'Colocynthis*. Official names of the *Colocynth* or *Bitter Apple*, *Colocyn'this*, *Coloquin'tida*, *Athan'dal* (Arab.), *Bitter Gourd*, *Bitter Cucumber*, (F.) *Coloquinte*. Ord. *Cucurbitaceæ*. *Ser. Syst. Monœcia Monadelphica*. A Turkey and Cape of Good Hope plant. The spongy part or medulla of the fruit, *Colocyn'thidis seu Cucumeris Colocyn'thidis Pulpa*, *Colocynthis* (Ph. U. S.), has a bitter, nauseous, and acrimonious taste, and is a strong cathartic, acting chiefly on the upper part of the intestines. It is scarcely ever used, except in combination.

CUCUMIS MELO. The systematic name of the *Melon Plant*, *Melo*, *Common Melon*, (F.) *Melon*. The fruit is an agreeable article of diet, but not very digestible, unless when ripe. The seeds possess mucilaginous properties.

CUCUMIS SATIVUS. The systematic name of the *Cucumber plant*, *Cucumis*, *Angourion*, *Citrullus*, *Cucumer* or *Cu'vimer*, from its curved shape. (F.) *Concombre ordinaire*. The *cucumber*, *cucumber*, is used, when young, as a pickle,

when it is called a *Gherkin*, (F.) *Cornichon*. It is not a fruit easy of digestion. The seeds are mucilaginous.

An ointment of green cucumbers—*cucumber ointment*—is used as a cooling salve. It is made by combining, by means of heat, the juice of green cucumbers with lard and cast snet.

CUCUMIS SYLVESTRIS, *Momordica elaterium*.

CUCUPHA, *Cu'culus*, *Pileus*, *Sacculus cephalic'us*. A sort of coif or cap, with a double bottom, between which is enclosed a mixture of aromatic powders, having cotton for an excipient. It was formerly used as a powerful cephalic.

CUCURBITA, à *curvata*, owing to its shape. A gourd. (F.) *Citrouille*. See *Cupping-glass*.

CUCURBITA ANGURIA, *C. citrullus*.

CUCURBITA CITRULLUS, *C. Anguria seu pratinat'ida*. The systematic name of the *Watermelon plant*; *Citrullus*, *Anguria*, *Tetrangu'ria*, *Sicilian Citrul*, *Water-melon*. Ord. *Cucurbitaceæ*; *Ser. Syst. Monœcia Monadelphia*. (F.) *Melon d'eau*, *Pastèque*. The juice of the fruit is very abundant, whence its name. The *Watermelon* is extremely refreshing and agreeable, when made cool, and is eaten like the common melon. It is very much used in the United States.

CUCURBITA LEUCANTHA, *C. lagenaria*.

CUCURBITA PEPO, *Pepo*. The systematic name of the *Common Pompon* or *Pumpkin*, *Cucur'bita*. The seeds of this plant, as well as those of *Cucur'bita lagenaria*, *Bottle-gourd*, contain a large proportion of oil, capable of forming emulsions. They have been recommended in cases of tapeworm, in the dose of fʒss. The fixed oil of the seeds has been given, with success, in the same cases, in the dose of half a fluid ounce.

Both the fruit of *CUCURBITA LAGENARIA*, *C. leucantha*, *Pepo lagenaria*, (F.) *Calebasses*, and that of *CUCURBITA PEPO*, *Pepo vulg'ris*, (F.) *Potiron*, *Courge*, are eaten.

CUCURBITA PINNATIFIDA, *C. citrullus*.

CUCURBITAIN, *Cucurbitinus*.

CUCURBITATIO, *Cupping*.

CUCURBITE, see *Alembic*.

CUCURBITINUS, (F.) *Cucurbitin*. This name was formerly given to the *Tænia solium*, because composed of rings which resemble the seeds of the gourd,—*cucurbita*. The ancients believed, that the rings, which are sometimes discharged, were so many separate worms. See *Tænia*.

CUCURBITULA, *Cupping-glass*.

CUCURBITULÆ CRUENTÆ, *Cupping* with the scarificator—*c. cum Ferro*, *Cupping* with the scarificator—*c. Siccæ*, *Cupping*, dry.

CUDWEED, *Antennaria plantaginifolia*, *Gnaphalium margaritaceum*.

CUILLERÉE, *Cochlearo*.

CUIR, *Corium*—*c. Chev'leu*, *Scalp*.

CUISSART. A wooden leg. See *Cornu ammonis*.

CUISSE, *Thigh*, *Crus*—*c. Postérieure du cervelet*, *Corpus testiforme*.

CUISSON (F.). A smarting, burning pain.

CUIVRE, *Cuprum*—*c. Ammoniacal*, *Cuprum ammoniatum*—*c. et Ammoniaque*, *sulfate de*, *Cuprum ammoniatum*—*c. Limaille de*, see *Cuprum*—*c. Sulfate de*, *Cupri subacetat*—*c. Sulfate de*, *Cupri sulphas*.

CUIVREUX (F.). *Copper-coloured*. A tint of complexion, (F.) *Teint cuivreux*, such as is observed in cancerous affections. Syphilitic ulcers of the throat, &c., are often copper-coloured.

CUJETE, *Cressentia cujete*.

CULBIC'IO. A Latin word, employed by old writers as synonymous with *ardor urinæ* and *gonorrhœa*.

CULBUTE (F.). 'A tumble head-over-heels.' A movement which the fœtus has been supposed to execute at the end of the 7th month of utero-gestation; and by means of which, it was presumed, the head presented towards the orifice of the uterus; a change of situation which is impracticable in ordinary circumstances.

CULCITA SALINA, Halotyle.

CULEON, Anus.

CULEUS. A measure containing 20 barrels, or 40 urns, equal to 180 gallons.—Pliny, Gormæus.

CULEX, (F.) *Cousin*. A genus of insects, unhappily too well known in almost every part of the world, on account of their bites, which give rise to painful, local inflammation. The gnats and mosquitoes belong to this genus.

CULI FLOS, Cardamine pratensis.

CULILAWAN, see *Laurus culilawan*.

CULINARY ART, from *culina*, 'a kitchen.' *Cookery*, *Res culinaria seu coquinaria*, *Ars culinaria seu coquinaria*, *Magisteriæ*. The art of preparing meats for the table. In judging of the dietetic properties of various kinds of aliment, the culinary process to which they have been subjected will always have to be considered. Many of the writers on the culinary art have been physicians.

CULLITLAWAN, see *Laurus culilawan*.

CULMINATIO, Acme.

CUTELLAIRE (F.), from *cutellus*, a 'little knife.' *Le couteur cutellaire*, known also under the name *Couteau de feu*, *Fire-knife*, is used for what the French term the *Couteur transcurrente*. (See *Cauterization*.) It is shaped like a small hatchet.

CUTELLUS, Culter, Knife.—c. *Anceps*, Knife, double-edged.—c. *Uncus*, Knife, crotchets.

CUTLER, *Cutellus*, from *colo*, *cultum*, 'I cultivate.' A *coulter*, a *knife*, *sculptel*, *machœra*, *machœrion*, *machœria*. Also, the third lobe of the liver, so called from some fancied resemblance.—Theophilus Protospatarius.

CUTLER CURVUS, Knife, crooked.—c. *Falcatus*, Knife, crooked.—c. *Lenticularis*, Knife, lenticular.—c. *Rectus*, Knife, amputation.—c. *Tonsorius*, Razor.

CULTRIVOROUS, *Cultricorvus*, from *culter*, 'a knife,' and *vorare*, 'to devour.' Individuals, who have seemed to swallow knives with impunity, have been so called.—*Knife-eaters*.

CULUS, Anus.

CULVER'S PHYSIC, *Leptandra Virginica*.

CULVER'S ROOT, *Leptandra purpurea*.

CUMAMUS, Piper cubeba.

CUMIN, *Cuminum cyminum*—c. *Fauz*, *Nigella*—c. *des Près*, *Carum*.

CUMINUM CYMINUM. The systematic name of the *Cumin* or *Cumin plant*, *Cuminum minimum* seu *Romanium*, *Cymimum*, *Cumimum*, *Carabadium*. Ord. Umbelliferae. Sex. Syst. Pentandria Digynia. (F.) *Cumin*, *Anis aigre*. The seeds of cummin, which is a native of Egypt, have a warm, bitterish, disagreeable taste. Water extracts their odour, and spirit takes up both odour and taste. They are not used, except in a plaster, which bears their name. When drunk in wine, the ancients believed they produced paleness; hence, Horace called cummin *exanguis*; and Juvenal, *pallens*.—Dioscorides, Pliny.

CUMINUM MINTUM, *C. cyminum*—c. *Nigrum*, *Nigella*—c. *Pratense*, *Carum*—c. *Romanum*, *C. cyminum*.

CUMMIN, *Cuminum cyminum*.

CUMULUS, 'a heap or pile.' A rounded prominence, in the centre of the proligerous disk, in which there is a small opaque cavity that contains the ovum. See *Tache embryonnaire*.

CUMULUS, *GERMINAL*, *Tache embryonnaire*—c.

Germinativus, *Tache embryonnaire*—c. *Proligerus*, *Tache embryonnaire*.

CUNEA/LISSUTURA, from *cuneus*, 'a wedge.' The suture formed between the great and little alæ of the sphenoid bone and the os frontis.—Blasius.

CUNÆN (F.), *Cuneanus*. Relating to the cuneiform bones.

Articulations Cunæennes;—the joints between the cuneiform bones, as well as between them and other parts.

Ligaments Cunæennes;—the ligaments which hold the cuneiform bones together.

CUNEIFORM, *Cuneiformis*, *Sphenoides*, from *cuneus*, 'a wedge,' and *forma*, 'shape.' *Wedge-shaped*. This name has been given to several bones. 1. To the sphenoid. 2. To a bone of the carpus, situate between the os lunare and os orbiculare. It is, also, called *Os Pyramidale*, *Os Triquetrum*, and *Os Cubitale*. 3. To the basilar process of the occipital bone; and, 4. To three of the bones of the tarsus, which are distinguished, according to situation, reckoning from within outwards, into *first*, *second*, and *third*,—or *internal*, *middle*, and *external*; and according to size, reckoning in the same order, into *great*, *small*, and *middle-sized*. The posterior surface of these bones is united to the anterior face of the scaphoides; the anterior surface with the corresponding metatarsal bones; and, in addition, the external surface of the third is articulated, behind, with the cuboides. They are also called *Chalcoides ossa seu ossicula*.

CUNEIFORM CARTILAGES OF THE LARYNX, *Cartilages of Wrisberg*, are two small cylinders of fibro-cartilage, about seven lines in length, and enlarged at each extremity. By the base, the cartilage is attached to the middle of the external surface of the arytenoid; and its upper extremity forms a prominence on the border of the aryteno-epiglottidean fold of membrane. They are sometimes wanting.

CUNEO. In composition, appertaining, or relating, to the cuneiform bones.

CUNEO-CUBOID, *Cuneo-cuboides*. Belonging to the cuneiform bones and cuboides.

CUNEO-CUBOID ARTICULATION is formed by the third cuneiform bone and cuboides. It is furnished with a synovial capsule, and two cuneo-cuboid ligaments:—a *dorsal* and a *plantar*.

CUNEO-SCAPHOID, *Cuneo-scaphoides*. Belonging to the cuneiform bones and scaphoid.

CUNEO-SCAPHOID ARTICULATION is formed by the posterior surfaces of the three ossa cuneiformia, and the anterior surface of the scaphoid. It is furnished with a synovial capsule and ligaments, some of which are *dorsal*, others *plantar*.

CUNILA, *Satureia capitata*—c. *Bubula*, *Origanum*.

CUNILA MARIA'NA, *Satureia origanoides*, *Calamintha erecta Virginiana*, *Dittany*, *American* or *Mountain Dittany*, *Mint-leaved* or *Maryland Cunila*, *Stemonit*, *Wild Basil*, *Sweet Horsemint*, (F.) *Cunille d'Amérique*. A small indigenous herb, *Order*, *Labiata*, growing on dry, shady hills, from New England to Georgia, and flowering in June and July. Its medical properties are dependent upon essential oils, like the mints.

CUNILA, *MARYLAND*, *Cunila mariana*—c. *Mint-leaved*, *C. mariana*—c. *Muscula*, *Inula dysenterica*—c. *Pulegioides*, *Hedeoma pulegioides*—c. *Sativa*, *Satureia hortensis*.

CUNILAGO, *Inula dysenterica*.

CUNILE D'AMÉRIQUE, *Cunila mariana*.

CUNNUS, *Vulva*.

CUP, (Sax.) *eopp*, *Scutella*, *Catillus*, *Patella*, *Excipula*, (F.) *Palette*, *Poëlette*, *Poëlette*, *Vase à saigner*. A small vessel of a determinate size,

for receiving the blood during venesection. It generally contains about four ounces. A bleeding of two cups is, consequently, one of eight ounces.

CUPAMENI, *Acalypha Indica*.

CUPIDITAS, Voluntas—c. Desedendi, Voluntas desedendi.

CUPIDO, Appetite. Also, Cupid, the god of love, in ancient mythology.—*Deus copulatio'nis*.

CUPOLA, see Infundibulum of the cochlea.

CUPPED, see Corium phlogisticum.

CUPPING, *Catacam'us*, from (F.) *couper*, 'to cut,' or to draw blood in vessels resembling cups; *Applicatio cucurbita'rum* seu *cucurbitula'rum*, *Cucurbitatio*. A species of blood-letting, performed by a scarificator, and glass, called a *cupping-glass*, *Cucurbit'ula*, *Cucurbita*, *Sic'na*, (Old Engl.) *Bleeding-boist*, (F.) *Ventouse*. The lancets are placed in such a manner in the scarificator, that, when it is applied upon the affected part, the whole are, by means of a spring, pushed suddenly into it. After scarification, the cupping-glass, which has been previously exhausted by heat, or by an exhausting syringe, is applied. The pressure of the air within the glass being thus diminished, the necessary quantity of blood may be drawn. See Bellometer. *Dry cupping*, *Cucurbit'ula sicca*, is the application of the glasses, without previous scarification; (F.) *Ventouses sèches*. It is used to prevent the activity of absorption from any wounded part; occasionally, to excite suppuration in indolent abscesses; and to remove the pus when an abscess is opened. *Cupping*, without any epithet, means the abstraction of blood by means of the scarificator and cups:—(F.) *Ventouses scarifiées*, (L.) *Cucurbitula cruenta*, *C. cum Ferro*, *vet cupping*. The verb 'to cup,' signifies to draw blood by cupping.

CUP-PLANT, *Silphium perfoliatum*.

CUPRESSUS, *κυνάριστος*, *C. sempervirens*—c. Arbor vitæ, *Thuja occidentalis*—c. Disticha, *Taxodium distichum*.

CUPRESSUS SEMPERVIRENS, *C. pyramidalis*. The systematic name of the *Cupressus*, *Cypariss'ens*, *Cypress*, (F.) *Cyprès*. Nat. Ord. Coniferae. The berries, leaves, and wood, have been considered astringent and useful in intermittents. The whole plant abounds with a bitter, aromatic, and terebinthinate fluid.

CUPRESSUS THYROIDES, *White cedar*, indigenous. An infusion of the tops is said to be stomachic, and, when taken warm, diaphoretic.

CUPRI AMMONIO-SULPHAS, *Cuprum ammoniatum*—c. et *Ammoniae sulphas*, *Cuprum ammoniatum*—c. *Diacetas*, *C. subacetas*.

CUPRI LIMATURA, *Filings of Copper*, (F.) *Limailles de Cuivre*, have been used in hydrophobia. It has been remarked under *Cuprum*, that they are inert.

CUPRI RUBIGO, *Cupri subacetas*.

CUPRI SUBACETAS, *C. Diacetas* seu *Rubi'go*, *Æru'go*, *Ver'digris*, *Hæpanti'um vir'idè*, *Prasi'um viridè*, *Crystals of Venus*, *Vir'idè Æ'ris*, *Æru'ca*, *Cal'cithos*, *Subac'etate of Copper*, (F.) *Sous-acétate de Cuivre*, *Vert-de-gris*, *Crystaux de Vénus*, *Verdet*. Impure subacetate of copper. This, as usually met with, is in masses, difficult to break; not deliquescent; foliaceous; of a fine bluish-green colour, and salt taste. It is tonic, emetic, escharotic, and detergent; but scarcely ever employed internally. Chiefly used in detergent ointments. Dose, as a tonic, under $\frac{1}{2}$ gr.: emetic, from gr. j. to gr. ij. Powdered verdigris appears to be the active ingredient in *Smellone's Eye-salve*, which may be imitated by rubbing half a drachm of finely-powdered verdigris with a little oil, and then mixing it with an ounce of yellow basilicon.

An ointment composed of one drachm of finely-

powdered verdigris, with an ounce of lard or spermaceti ointment, is used in psoriasis, tetter, &c.

CUPRI SULPHAS, *Vitriolum Cupri* seu *Veneris* seu *Cyprium* seu *Cyprium* seu *Carad'ennu* seu *Roma'num*, *Cuprum Vitriola'tum*, *Lapis Carad'enus*, *Sulphate of Copper*, *Blue Stone*, *Blue Vitriol*, *Roman Vitriol*, *Mortooth*, (F.) *Sulfate de Cuivre*, is in rhomboidal, rich, blue, semi-transparent, efflorescing crystals. The taste is harsh, styptic, and corrosive. It is soluble in four parts of water at 60°, and is tonic, emetic, astringent, and escharotic. As a tonic, it has been used in epilepsy, intermittents, &c. Dose, as a tonic, gr. $\frac{1}{2}$ to gr. ij, in pill; as an emetic, gr. ij to x, in water. A very weak solution is sometimes used in ophthalmia and in gleet; and it forms the basis of BATES'S *Aqua camphorata*, which has been recommended, diluted with sixteen parts of water, in the purulent ophthalmia of children.

CUPRI SULPHAS AMMONIACALIS, *Cuprum ammoniatum*—c. *Vitriolum*, *Cupri sulphas*.

CUPRUM, *Chalcos*, *Æs*, *Venus* of the Alchemists: *Copper*, *Cyprium*, *Vir'idè montanum*, (F.) *Cuivre*. Its odour is peculiar and sensible when rubbed; taste disagreeable and metallic; colour red yellow. S. g. 7.87; ductile; very malleable; less hard than iron; easily oxidized. In its metallic state, it exerts no action on the system. When swallowed, it undergoes no deleterious change. Copper cannot be dissolved whilst tin is in the mixture, and hence the utility of tinning copper vessels. Copper culinary vessels are harmless under ordinary cleanliness, provided the substances be not suffered to remain in them till cold. The salts of copper are very deleterious.

CUPRUM ALUMINATUM, *Lapis divinus*.

CUPRUM AMMONIATUM, *C. Ammoniacale* seu *Ammoni'acum* seu *ammoni'aco-sulphur'icum*, *Ammonia'ted Copper*, *Ammoni'acal Sulphate of Copper*, *Ammonia'tum Cupri*, *Sal anti-epilepticus* of WEISSMAN, *Sulfas Cupri ammoniacalis* seu *ammoni'aca cupra'tus* seu *Sulfus Cupri et Ammonia'*, *Subsulfus ammonio-cupricus* *Dento-Sulfas Cupri et Ammonia'*, *Cupri Ammonio-sulfas*, *Ammonia' Cupro-sulphas*, *Ammonio-sulphate of Copper*, (F.) *Sulfate de cuivre et d'ammoniaque*, *Cuivre ammoniacal*. (*Cupri Sulph. 3ss; Ammon. Carb. 3vj*. Rub in a glass mortar till effervescence ceases. Wrap the ammoniated copper in bibulous paper, and dry with a gentle heat. Keep in a well-stopped glass bottle.—Ph. U. S.) A crystalline powder of a rich violet colour, and hot, styptic taste. By exposure to air, it becomes partly converted into carbonate of copper. It is tonic and astringent, and has been chiefly employed in epilepsy and other obstinate spasmodic diseases.—Dose, gr. $\frac{1}{2}$ gradually increased to gr. iv, in pill.

CUPRUM VITRIOLATUM, *Cupri sulphas*.

CUPULAR, *Cupularis*, (F.) *Cupulaire*; from *cupula*, 'a small cup.' Of or belonging to a cupula.

CUPULAR CAUTERY, (F.) *Cautère cupulaire*, is an iron in the shape of a cupula, formerly used to cauterize the skin of the cranium in certain diseases, as epilepsy, chronic headach, &c.

CURA, *Cura'tio*, *Merim'na*, *Merim'nè*. Attention to, or treatment or cure of, a disease. Keuchen defines *Cura*, 'medicine,' and *Curator*, 'the physician.' *Curatio*, also, sometimes means purification; as, *Adeps suillus curatus*.—Scribonius Largus.

CURA AYENACEA. A decoction of oats and succory roots, in which a little nitre and sugar are dissolved. Used as a refrigerant.

CURA DERIVATIVA, *Derivation*—c. *Fumigatoria*, *Thymiatechny*.

CURA MAGNA. 'Great cure.' A term employed, at times, for a method of treatment preferable to all others. Thus pyralism has, by

many, been considered the "*cura magna*" for syphilis.

CURA MEDEANA, Transfusion—c. Palliativa, see Palliative—c. Radialis, see Palliative—c. Revulsoria, see Derivation.

CURABLE, *Sanab'ilis*, *Aces'tos*, *Aces'mius*, *Pir'inos*, *Medicab'ilis*, *Med'icable*, *San'able*, (F.) *Guérissable*. That which is susceptible of cure. An epithet applied to both patients and diseases.

CURAGE, *Polygonum hydropiper*.

CURA'RE, *Wourali*, *Woorali*, *Woorara*, *Waurari*, *Woorara*, *Wuadi*, *Uvari*, *Uvadi*, *Ouvari*, *Yoorara*. A very energetic vegetable poison, employed by the South American Indians to poison their arrows. It is said to be obtained from the bark of a species of convolvulus, called *Vajeco de Maracare*, but is referred by Martius to *Stychnos Guianensis*, and by Dr. Schomburg to *S. torricellia* seu *toxicifera*. The juice of *Echites euberec'ta*, another apocyanaceous plant, is said to enter into its composition. Dr. Brainard thinks it contains the poison of serpents as its main ingredient.

CURATIO, *Mede'la*, *Sana'tio*, *A'cesis*, *Al-thae'is*, *Althae'is*, *Therape'ia*, *I'isus*, *Curation*, *Treatment* (F.) *Traitement*. The aggregate of means employed for the cure of disease. See Therapeutics.

CURATIO, Cura, Cure—c. Contrariorum per Contrarium, Allopathy—c. Morbi per Inediam, Linothepapia.

CURATIVE, *San'ative*, *San'atory*, *Healing*, *Acesp'orus*, *Acesino'sus*. Relating to the cure of a disease. Curative Indications are those which point out the treatment to be employed. Curative Treatment, that employed in the cure of disease, in opposition to preservative or prophylactic treatment. We say, also, Curative Process, Curative Means, &c.

CURATOR INFIRMORUM, *Infirmer*.

CURCAS PURGANS, *Jatropha curcas*.

CURCULIO, Penis.

CURCUMA ANGUSTIFOLIA, see Arrow-root—c. Aromaticæ, *Kæmpferia rotunda*.

CURCUMA LONGA, *Curcuma*, *Amomum Curcuma*, *Borri*, *Borriberri*, *Cober'ri*, (Hindoostan,) *Crocus Indicus*, *Terra Mer'ita*, *Cinnac'orus rad'ice crocea*, *Mayp'ila*, *Kua Kaha*, *Cype'rus Indicus*, *κρυπτος ἴνδικος* of Dioscorides [?], *Turmeric*, (F.) *Racine de Safran*, *Safran des Indes*, *Souchet des Indes*. Ord. Zingiberaceæ. Turmeric root—the rhizoma of *curcuma longa*—*Curcuma* (Ph. U. S.), is brought from the East Indies; but is possessed of very little, if any, medicinal efficacy. It is one of the ingredients in *Curry Powder*.

CURCUMA ZEDOARIA, see *Kæmpferia rotunda*—c. Zerumbet, see *Kæmpferia rotunda*.

CURD, *Curd of Milk*, *Lac pres'mum*, (F.) *Caillé*, *Lait caillé*. The coagulated or curdled (*Coac'tus*, *Coagulat'us*, (F.) *Caillé* ou *Caillebotte*), part of milk.

CURD, ALUM, *Coagulum aluminosum*.

CURE; from *cura*, 'care.' *Aces'ia*, *A'cesis*, *Aces'mus*, *Curat'io*, *C. felix*, *Sanat'io*, *San'itas*, (F.) *Guérison*. A restoration to health; also, a remedy; a restorative.

CURE-ALL, *Geum Virginianum*, *Oenothera biennis*.

CUREDENT, *Dentiscapulum*.

CURE DU RAISIN, Grape-cure.

CURE-LANGUAGE, (F.) *Lingua scalpium*, *Tongue-scraper*. An instrument of ivory, tortoise-shell, &c., shaped like a knife or rasp, for removing the mucous coating which covers the tongue after sleep, &c.

CURE-OREILLE, Ear-pick.

CURETTE, Scoop.

CURMI, Courmi.

CURRANT, BLACK, *Ribes nigrum*—c. Red, *Ribes rubrum*.

CURRENTS, see *Vitis Corinthis*.

CURRY or CURRIE POWDER. A condiment, formed of various spices, and eaten with rice, particularly in India. The following is one of the forms of its preparations: *Sem. coriand.* 3xviii. *pip. nigr.* 5ij, *cayen.* 5j, *rad. curcumæ*, *sem. cumini*, aa 5ij, *sem. fienugr.* 5iv: mix.

CURSUM HÆMORRHOIDALIS HERBA, *Ranunculus ficaria*.

CURSUS, Running—c. Matricis, *Leucorrhœa*—c. Menstruus, Menses.

CURVAMEN, Curvature.

CURVATEUR DU COCCYX, Curvator Coccygis.

CURVATIO, Campsis.

CURVATOR COCCYGIS, (F.) *Curvateur du Coccyx*. Sömmering gives this name to a small bundle of fleshy fibres, which descends on the middle of the coccyx, uniting on each side with the ischio-coccygei muscles. It is a part of those muscles.

CURVATURE, *Curvatur'a*, *Curva'men*, *Flexura*, *Gnac'm'is*, from *curvare*, *curvatum*, 'to bend.' *Cyrto'ma*, (F.) *Courbure*. The condition of a line or surface, which approximates more or less to the form of an arc; as the curvatures of the spine, duodenum, stomach, &c. Accidental curvatures of bones are the effect of rickets, or *Mollities ossium*. The Greeks called the curvature of the spine, *ὑβος*, *σκολιός*, and *λόφος*, according as the deviation was backwards, laterally, or forwards.

CURVATURE, POTT'S, Vertebral disease.

CURVE OF CARUS, see Pelvis.

CURVED LINES, *Lin'æ semicircula'res exte'ri'na* seu *arcu'atæ*, (F.) *Courbes Lignes*. Two crooked lines or projections on the posterior surface of the occipital bone. They are distinguished into *superior* and *inferior*. Some lines on the os innominatum are also so called.

CURVIMER, *Cucumis sativus*.

CUSCO-CINCHONIA, *Aricina*.

CUSCUS, *Andropogon muricatus*.

CUSCUTA, C. *Europæa*.

CUSCUTA CHLOROCARPA, see C. *Glomerata*.

CUSCUTA EPITHYMIUM, C. *minor*. The systematic name of the Dodder of Thyme, *Epithymum*, *Epithymum Cuscut'a* seu *Creticum*. Ord. Convolvulaceæ. A parasitical plant, possessed of a strong, disagreeable smell, and a pungent taste very durable in the mouth. It was once used as cathartic in melancholia.

CUSCUTA EUROPA'IA seu *major* seu *vulga'ris* seu *tetrandra* seu *filiform'is*, *Cuscut'a*, *Epithymum officinarum*, was conceived to possess similar properties.

CUSCUTA FILIFORMIS, C. *Europæa*.

CUSCUTA GLOMERATA, and CUSCUTA CHLOROCARPA, Dodder, *American Dodder*, indigenous plants, are bitterish, subastringent, tonic, and antiperiodic.

CUSCUTA MAJOR, C. *Europæa*—c. Minor, C. *Epithymum*—c. *Tetrandra*, C. *Europæa*—c. *Vulgaris*, C. *Europæa*.

CUSHIA, *Heraclæum spondylium*.

CUSHY-COW-LADY, *Coccinella septempunctata*.

CUSPARIA BARK, see C. *Febrifuga*—c. Bark, False, *Brucea antidysenterica*, and *Stychnos nux vomica*.

CUSPARIA FEBRIFUGA, *Bonplan'dia trifoliata* seu *angustura*, *Angustura* (Ph. U. S.), *Galipæa febrifuga*. Order, Rutaceæ. The South American tree which furnishes the *Cusparia* or *Angustura Bark*, (F.) *Angusture* ou *Angosture vraie*, *Quinquina faux de Virginie*. According to Dr. Hancock, however, the *Angustura bark*

is derived from *Galipã'a officinalis*, *Sex. Syst.* Decandria Monogynia, *Nat. Ord.* Diosmeæ; and this view has been adopted in the Pharmacopœia of the United States. Its odour is peculiar; taste intensely bitter and slightly aromatic; pieces thin; externally, gray and wrinkled; internally, yellowish-fawn; fracture, short and resinous. It yields its virtues to water and to proof spirit. It is tonic, stimulant, and aromatic. Dose, gr. v to xx or more, in powder.

CUSPIS. A spear. This name has been given to the glans penis.—Röfink. It meant, also, a kind of bandage.

CUSTODIA VIRGINITATIS, Hymen.

CUSTOS, Vulva.

CUT. from (F.) *couteau*, 'a knife,' or from West Gothic, *kota*, 'to cut;' or from *κόττω*, 'I cut.' *Casura*, *Tome*, *Incisio*, *Vulnus simplex*, *Tres vulnus simplex*, *Cope*, (F.) *Coupure*. A common expression for the division or solution of continuity made by a sharp instrument.

CUTAMBULUS, from *cutis*, 'skin,' and *ambulo*, 'I walk.' 'Walking in the skin.' An epithet given to certain parasitical animals, which creep under the skin—such as the Guinea-worm; and to certain pains felt between the skin and flesh, as it were.

CUTANEAL, Cutaneous.

CUTANEO-SOURCILIER, Corrugator supercilii.

CUTANEOUS, *Cuta'neal*, *Cuta'neus*, (F.) *Cutané*, from *cutis*, 'the skin.' Belonging to the skin.

CUTANEOUS DISEASES, *Eruptions*, *Epiphy'mata*, are the numerous affections of a morbid character to which the skin is liable—*Dermatopathia*, *Dermato'ses*.

Chronic cutaneous diseases may be thus classified:

- | | | |
|----------------------|---|--------------|
| 1. Exanthematous | { | Urticaria. |
| | | Roseola. |
| | | Erythema. |
| | | Pemphigus. |
| | | Rapia. |
| 2. Vesicular | { | Herpes. |
| | | Scabies. [?] |
| | | Eczema. |
| 3. Pustular | { | Impetigo. |
| | | Ecthyma. |
| | | Lichen. |
| 4. Papular | { | Straphulus. |
| | | Prurigo. |
| | | Lepa. |
| 5. Squamous | { | Psoriasis. |
| | | Pityriasis. |
| | | Aene. |
| | | Sycosis. |
| 6. Folliculous | { | Ichthyosis. |
| | | Trichosis. |
| | | Favus. |

CUTANEOUS EXHALATION AND ABSORPTION are those which take place from the skin.

CUTANEOUS, MIDDLE POSTERIOR, see Sciatic nerve, lesser.

CUTANEOUS NERVES, *Cutaneal Nerves*, so called, of the upper extremity, are two in number. They are given off from the brachial plexus. The *internal cutaneous*, *Cubito-cutané*, (Ch.) descends along the inner part of the arm, and divides above the elbow into two branches; the outermost of which follows the outer edge of the biceps, and spreads over the anterior and inner part of the forearm; and the innermost accompanies the basilic vein, passing to the corresponding side of the hand and little finger. The *external cutaneous*, *Radio-cutané* (Ch.) *Musculo-Cutaneus*, *Perforans CASSERII*, passes outwards; perforates the coraco-brachialis; descends along the anterior and outer part of the arm; passes

as far as the middle fold of the elbow under the median cephalic; and descends along the anterior and outer edge of the forearm. At some distance above the wrist, it divides into two branches; an *outer*, which is distributed on the back of the hand, the thumb and index finger; and an *inner*, which descends on the muscles of the thumb into the palm of the hand, and divides into fine filaments, which may be traced to the fingers.

Cutaneous nerves, *Middle cutaneous*, two in number, are branches of the crural or femoral nerve. They are distributed to the integument of the middle and lower part of the thigh and of the knee.

Nervus Cutaneus Minor, *Lesser internal cutaneous nerve*, *Nerve of Wrisberg*, takes its origin from the axillary plexus, but is more particularly connected with the ulnar nerve. It soon separates from the ulnar, running afterwards between it and the inner side of the arm. A little below the axilla it divides into two branches, which are distributed to the arm.

Chaussier calls the *cephalic* and *basilic veins* the *Radial* and *Cubital Cutaneous*.

CUTCH, Catechu.

CUTE, Malleolus.

CUTI, Catechu.

CUTICLE, Epidermis—c. of the Enamel, see Enamel.

CUTICULA, Epidermis.

CUTIO, Oniscus.

CUTIS, *Pellis*, *Pella*, *Co'rium*, *Derma*, *Deris*, *Anthro'pé*, *Anthro'pen*, *Skin*, (F.) *Peau*, *Dermé*. A dense, resisting membrane, of a flexible and extensible nature, which forms the general envelope of the body; and is continuous with the mucous membranes, through the different natural apertures. It is generally considered to be formed of three distinct layers—the *epidermis*, *rete* or more properly *corpus mucosum*, and *corium* (*cutis vera*, *derma*). Some anatomists, however, separate it into several others. Its outer surface is covered by a number of small eminences, called papillæ, which are generally regarded as essentially nervous and vascular. The skin is our medium of communication with external bodies. It protects the subjacent parts; is the seat of touch; and through it are exhaled the watery parts of the blood, which are not needed in the nutrition of the body. The state of the skin, as regards heat and dryness, affords useful information in pathological investigations. Its colour, too, requires attention: the paleness of disease is as characteristic as the rosy complexion of health. The colour of the skin varies according to the age, sex, &c. As a general rule, it is finer in the female and child than in the male and adult. In old age it becomes light-coloured, thin, and dry. It likewise varies according to the races, &c.

CUTIS ANSERINA, *Horrida cutis*—c. *Carnosa*, *Panniculus carnosus*—c. *Extima*, *Epidermis*—c. *Lingux*, see Tongue—c. *Sailla*, *Conenne*—c. *Summa*, *Epidermis*—c. *Tensa Chronica*, *Induration* of the cellular tissue—c. *Ultima*, *Epidermis*—c. *Vera*, *Corium*, see Cutis.

CUTITIS, *Cytitis*, Erysipelatous inflammation.

CUTLINS, Gfroats.

CUTTING ON THE GRIPE, see Lithotomy.

CUTTLE FISH, Sepia.

CUTTUBUTH, *Cutubuth*, *Kutubuth*, *Leucomo'ria*, *Melancholia errabunda*. The Arabian physicians gave this name to a species of melancholy, accompanied with so much agitation, that the patients cannot remain tranquil for the space of an hour.

CUURDO CANELLA, *Laurus cinnamomum*.

CYANIC, *Cyan'icus*, (F.) *Cyanique*; from

κυανος, 'blue.' An epithet for the blue stage of a malady—as cholera.

CYANEPHIDROSIS, from *κυανος*, 'blue,' *επι*, an augmentative preposition, and *ιδρωσ*, 'I sweat.' Copious perspiration, which colours the linen blue.

CYANODERMIA, Cyanopathy.

CYAN'OGEN, (F.) *Cyanogène*; from *κυανος*, 'blue,' and *γενναω*, 'I generate.' So called from its being an ingredient in Prussian blue. It forms, with oxygen, the cyanic and other acids; with hydrogen, the hydrocyanic acid.

CYANOP'ATHY, *Cyanopathia*, *Cyano'sia*, *C. cardiaca*, *Cyanoder'mia*, *Hæmato-cyano'sis*, *Παματο-cyanopathia*, *Kyano'sis*, *Morbus cæruleus*, *Cærule'sis neonatorum*, from *κυανος*, 'blue,' and *αθος*, 'affection.' (F.) *Cyanose*, *Maladie bleu*, *leître bleu*. A disease in which the surface of the body is coloured blue. It is often symptomatic, and commonly depends on a direct communication remaining between the cavities of the right and left side of the heart; *Patency of the foramen ovale*, *Aclitrocardie*.—Piorry. Such communication does not, however, always occasion the blue disease, but it is generally thought that the disease never exists without this state of parts; or without some obstacle to the circulation in the right side of the heart. The blueness does not seem to be owing to the admixture of black and red blood. A child affected with blueness is said to be *cyanosed*, (F.) *Cyanosé*.

CYANOSE, see Cyanosis.

CYANOSIS, Cyanopathy—c. Pulmonalis, Atelectasis pulmonum.

CYANURETUM FERRO-ZINCICUM, *Zinci ferro-hydrocyanas*.

CYANUS, *Centaurea cyanus*—c. *Ægyptiacus*, *Nymphaea nelumbo*.

CY'ANUS SEG'ETEM, *Bluebottle*, (F.) *Bluet* ou *Bluet des Moissons*, *Barbeau*, *Aubifoin*, *Casse-Lunettes*. Ord. Compositæ. The blue flowers of this European plant, (*κυανος*, 'blue,') when distilled with water, have been used in ophthalmia.

CYAR, *κυαρ*. 'The eye of a needle,' 'a small hole.' The *Meatus auditivus internus*. See Auditory canal (internal).

CYATHIS'CUS, diminutive of *κυαθος*, 'a bowl.' The concave part of a sound, made like a small spoon, as in the case of the ear-pick.

CY'ATHUS, 'a bowl.' A measure, both of the liquid and dry kind, equal to about an ounce and a half, or to the tenth part of a pint. According to Pliny and Galen, about 10 drachms.

CYATHUS CEREBRI, Infundibulum of the brain.

CYBE, Head.

CYBITON, Cubitus.

CYBOIDES, Cuboid.

CYCAS CIRCINALIS, see Sago—c. *Revoluta*, see Sago.

CY'C'EON, from *κυκλωω*, 'I mix together.' The ancient name of a medicine of the consistence of pap, composed of wine, water, honey, flour, barley meal, and cheese.—Hippocrates.

CYCLAMEN EUROPÆ'UM, *Arthamita*, *A. cyclamen*, *Cyclam'inus*, *Cyssaanthemon*, *Cyssaophyllon*, *Panis porc'inus*, *Cas'amum*, *Chyl'inē*, *Sov-bread*. Ord. Primulacæ. Sez. Syst. Pentandria Monogynia. (F.) *Pain du Porceau*. The fresh root is said to be acrid, bitter, drastic, and anthelmintic. Dose, ʒi. For external use, see Arthanita.

CYCLE, *Cyclus*, from *κυκλος*, 'a circle.' A period or revolution of a certain number of years or days. The Methodists gave this name to an aggregate of curative means, continued during

a certain number of days. Nine was the usual number.

CYCLE, *Hebdom'adal* or *Heptal*. A period of seven days or years, which, according to some, either in its multiple or submultiple, governs an immense number of phenomena of animal life.

CYCLISCUS, *Cyclismus*.

CYCLIS'MUS. A *Trach*, *Trachis'eus*. This name, as well as *Cyclis'eus*, was also given to a circular kind of rasp. They have the same etymon as *cycle*.

CYCLOCEPH'ALUS, (F.) *Cyclocephale*, from *κυκλος*, 'a circle,' and *κεφαλη*, 'head.' A monster whose eyes are in contact, or united in one.

CYCLOPHOR'IA, from *κυκλος*, 'a circle,' and *φωω*, 'I bear.' The circulation of the blood or other fluids.

CYCLOPIA. Same etymon as *Cyclops*. State of a monster that has both eyes united into one. Called, also, *Monops'ia* and *Rhinencephal'ia*.

CYCLOPIA GENISTO'DES; *Not. Ord. Leguminosæ*. A South African plant, the decoction and infusion of which are used as expectorants in chronic bronchitis and in phthisis.

CYCLOPS, *κυκλωψ*, from *κυκλος*, 'an orb or circle,' and *ωψ*, 'an eye.' *Monoc'ulus*, *Monops*, *Monophthal'mus*, *Monom'matus*, *Unic'ulus*, *Unoc'ulus*. A monster having but one eye, and that placed in the middle of the forehead, like the fabulous Cyclops.

CYCLOSIS, Circulation.

CYCLOTOME, *Cyclot'omus*, from *κυκλος*, 'a circle,' and *τεμνειν*, 'to cut.' An instrument, composed of a ring of gold and a cutting blade, by means of which the ball of the eye can be fixed, whilst the cornea is cut. It was invented by a surgeon of Bordeaux, named Guérin, for extracting the cataract. It is not used.

CYCLUS, *Circulus*.

CYDONIA, *Pyrus cydonia*—c. *Maliformis*, *Pyrus cydonia*—c. *Vulgaris*, *Pyrus cydonia*.

CYDONIA'TUM, *Cydonat'um*, *Dyacydoni'tes*. A composition of the *Cydonia mala* or quinces, with the addition of spices.—Paulus of Ægina.

CYDONIUM, see *Pyrus Cydonia*—c. *Malum*, see *Pyrus cydonia*.

CY'E'MA, *κυημα*, from *κυω*, 'I conceive.' Conception. Likewise the product of conception. See Embryo, and Fœtus.

CYESIOGNO'MON, from *κυωσις*, 'pregnancy,' and *γνωμων*, 'a sign, a token.' A sign of pregnancy.

CYESIOGNO'SIS, from *κυωσις*, 'pregnancy,' and *γνωσις*, 'knowledge.' Diagnosis of pregnancy.

CYESIOL'OGY, *Cyesiology'ia*, from *κυωσις*, 'pregnancy,' and *λογος*, 'a description.' The doctrine of gestation.

CYESIS, Conception, Feundation, Pregnancy.

CYESTEINE, *Kiesteine*.

CYLICH'NĒ, *Cyllich'nia*. A pill-box or earthenware pot, or small cup; from *κυλιξ*, 'a cup.'—Galen, *Œsius*.

CYLINDRI see Villous membranes—c. *Membranæci renum*, see *Calix*.

CYLINDROID, *Cylindroi'des*, from *κυλινδρος*, 'cylinder,' and *ειδος*, 'form.' Having the form of a cylinder. Chaussier calls the *Cornu Ammonis*, *Protuberance cylindroïde*.

CYLINDROIDES, *Teres*.

CYLOEPUS, see *Kyllosis*.

CYLOPODA, see *Kyllosis*.

CYLOPODION, see *Kyllosis*.

CYLO'SIS, *κυλλωσις*. Lameness, mutilation, or vicious conformation.—Hippocrates, Galen. See *Kyllosis*.

CYLOSOM'US, from *κυλλος*, 'crooked,' and *σωμα*, 'body.' A malformation by defect, in which the fissure and eventration are lateral, chiefly in the lower part of the abdomen, the in-

ferior extremity of the side affected with the fissure absent, or very little developed.—Vogel.

CYMATODES. *Undosus* from *κύμα*, 'a wave,' and *εδος*, 'resemblance.' The vacillating, undulatory character of the pulse in weak individuals.

CYMBÆ, Vulva.

CYMBALARIA ELATINE, Antirrhinum elatine.—c. Muralis, Antirrhinum linaria.

CYMBALARIS, Cricoid.

CYMBIFORME OS, Scaphoid bone.

CYMOPOGON SCHÖENANTHUS, Juncus odoratus.

CYMINUM, Cuminum cyminum.

CYMOGRAPHION, Kymographion.

CYN'ANCHĒ, *Angina*, from *κύνω*, *κύνος*, 'a dog,' and *αἶμα*, 'I suffocate' [?], 'Dog choak,' *Sore Throat*. (Old E.) *Stenosis*, *Stenocæcie*; *Paracynan'chē*, *Synan'chē*, *Prunella*, *Squinan'thia*,

Empres'ma Paristhmī'tis, *Cuma Paristhmī'tis*, *Isthmī'tis*, *Paristhmī'tis*, *Inflammatio Fau'cium*. Inflammation of the supra-diaphragmatic portion of the alimentary canal, and of the lining membrane of the upper part of the air-passages:—(F.) *Angine*, *Esquinancie*, *Mal de Gorge*, *Synancie*, *Squinancie*, *Kinancie*. It comprises the following chief varieties.

CYNANCHE EPIDEMICA, Cynanche maligna.—c. Externa, C. Parotidæa.—c. Fau'cium, C. tonsillarīis.—c. Gangrenosa, Cynanche maligna.—c. Laryngea, Laryngitis.

CYNANCHE MALIG'NA, C. gangrenosa seu epidemica seu pur'puro-parotidæa seu ulcero'sa, *Febris epidemica cum Angina*, *Empres'ma Paristhmī'tis Malig'na*, *Tonsillarum gangrenosa*, *Tonsillæ pestilentes*, *Epidemica gutturis lues*, *Pestilens fau'cium affectus*, *Pædan'chonē* [?], *Gangræna Tonsillarum*, *Ulcus Syriacum*, *Gorotill'o*, *Carbunculus angino'sus*, *Angina ulcero'sa seu ulcuscuto'sa seu epidemica seu gangrenosa seu malig'na seu fau'cium malig'na*, *Isthmoty'phus*, *Putrid or Ulcerous Sore Throat*, *Gangrenous Inflammation of the Pharynx*, (F.) *Angine gangræneuse ou Maligne*. It is characterized by crimson redness of the mucous membrane of the fauces and tonsils; ulcerations, covered with mucus; and spreading sloughs, of an ash or whitish hue; the accompanying fever typhus. It is often epidemic, and generally contagious [?]; and is frequently found accompanying scarlet fever,—giving rise to the variety, *Scarlatina maligna*, *Canker rash*. Cynanche maligna has been made to include both diphtheritic and gangrenous pharyngitis. See Pharyngitis, diphtheritic.

The general treatment is the same as in typhus; and stimulant antiseptic gargles must be used, consisting, for example, of the decoction of bark and muriatic acid.

CYNANCHE MAXILLARIS, c. Parotidæa.—c. *Esophæga*, *Esophagitis*.

CYNANCHE PAROTIDÆA, C. Parotides seu externa seu maxillaris, *Empres'ma Parotit'is*, *Parotitis*, *P. epidemica seu erysipelato'sa seu contagio'sa seu spu'ria seu sero'so glu'tinē tu'ens*, *Parot'ia*, *Parotidit'is*, *Angina maxillaris seu externa seu parotidæa exter'na*, *Gissa*, *Erythrochæ'ras*, *Genyocynan'chē*, *Gnathocynan'chē*, *Inflammat'io parotidum*, *Infla'tio parotidum*, *Catar'rhus Bellinulæ'nus*, *Branks* (Scotch), *Mumps*, (F.) *Inflammation de la Parotide*, *Oreillons*, *Ourles*. The characteristic symptoms are:—a painful tumour of the parotid gland, or of the cellular tissue surrounding it, or of both, not of the suppurative kind; frequently extending to the maxillary gland. It is very conspicuous externally, and is often accompanied with swelling of the testes in the male, and of the mammae in the female; the testes being sometimes absorbed afterwards. It is generally epidemic, and apparently contagious. [?] The treatment is very

simple; the adoption merely of the antiphlogistic plan, under which it usually soon disappears. When inflammation of the mammae or testes supervenes, it must be treated as if idiopathic.

Epidem'ic Parotit'is or *Mumps* is also termed *Parotit'is polymor'pha seu epidem'ica seu specifi'ca*.

CYNANCHE PHARYNGE'A, *Empres'ma Paristhmī'tis Pharyngea*, *Isthmī'tis*, *Pharyngit'is*, *Inflammat'io Pharyn'gia*, *Parasynan'chē*, *Angine inflammatoriæ*, *Sp. iv.*—(Boerhave.) (F.) *Angine Pharyngée*, *Catarre pharyngien*. Inflammation of the pharynx.

This disease can hardly be said to differ, in pathology or treatment, from Cynanche tonsillarīis. The same may be remarked of the *Cynan'chē Esophagæ'a*, *Esophagitis*.

CYNANCHE PRUNELLA, C. Tonsillarīis.—c. *Purpuro-parotidæa*, Cynanche maligna.—c. *Simplex*, *Isthmitis*—c. *Stridula*, C. trachealis.

CYNANCHE TONSILLARIS, C. fau'cium seu *Prunella*, *Empres'ma Paristhmī'tis Tonsillar'is*, *Synan'chē*, *Amygdalit'is*, *Branci*, *Branchi*, *Hyan'chē*, *Tonsillit'is*, *Paristhmī'tis*, *Paristhmī'tis*, *Dysphag'ia inflammatoriæ*, *Antiadit'is*, *Angine inflammatoriæ seu cum tumore seu tonsillar'is seu synochal'is seu sanguinæ seu vera et legit'ima*, *Inflammatio Tonsillarum*, *Antidoneus inflammatori'is*, *Inflammation Sore Throat*, *Common Squinancy*, *Squinzey*, *Squinsy* or *Quinsy*, (F.) *Amygdalitis*, *Inflammation des Amygdales*, *Angine tonsillaire*, *Pharyngite tonsillaire*, *Esquinancie*, *Squinancie*, *Catar'rhé gutturale*, *Angine gutturale inflammatoire*. The characteristic symptoms of this affection are, swelling and florid redness of the mucous membrane of the fauces, and especially of the tonsils; painful and impeded deglutition, accompanied with inflammatory fever. It is generally ascribed to cold, and is one of the most common affections of cold and temperate climates. It usually goes off by resolution, but frequently ends in suppuration.

Common sore throat is an affection of no consequence. It requires merely rest, and the observance of the antiphlogistic regimen. When more violent,—in addition to this,—bleeding, local or general, or both,—purgatives, inhalation of the steam of warm water; acid, or emollient gargles; rubefacients externally, or sinapisms or blisters. When suppuration must inevitably occur, the continued use of the inhaler must be advised, and an opening be made into the abscess as soon as pus shall have formed. If the patient be likely to be suffocated by the tumefaction, bronchotomy may be necessary.

CYNANCHE TRACHEALIS, Cynanche larynge'a of some, C. strid'ula, *Suffocatio strid'ula*, *Asthma infantum*, *Catar'rhus suffocati'vus* *Barbarden'is*, *Angina polypo'sa seu membrana'cea seu pulpo'sa seu pernici'o'sa seu inflammatoriæ seu suffocatoriæ seu stridit'is seu can'ina seu exsudatoriæ seu tracheal'is*, *Empres'ma bronchit'is seu bronchlemmit'is*, *Laryngit'is* et *Tracheit'is infantilis*, *Laryngit'is exsudati'ca seu membrana'cea seu polypo'sa*, *Laryngos'tasis seu Laryngo-tracheit'is*, *Laryngo-tracheitis with diphtherit'ic exudation*, *Laryngocac'hē*, *Orthopnœ'a cynan'chica*, *Cynanchorthopnœ'a*, *Pædan'chonē* [?], *Morbus Strangulato'rius*, *Trachit'is*, *Tracheit'is*, *Trachelit'is*, *Diphtherit'is tracheal'is*, *Expectoratio Sol'ida*, *Cuma Bronchit'is*, *Croup*, *Roup*, *Hives*, *Bold Hives*, *Choak*, *Stiffing*, *Rising of the lights*, (F.) *Angine laryngée et trachéale*, *Laryngite avec production de fausses membranes*, *Laryngite pseudo-membraneuse*. A disease characterized by sonorous and suffocative breathing; harsh voice; cough, ringing, or like the barking of a dog; fever, highly inflammatory. It is apt to be speedily attended with the formation of a

false membrane, which lines the trachea beneath the glottis, and occasions violent dyspnoea and suffocation, but is sometimes expectorated. The dyspnoea, as in all other affections of the air-passages, has evident exacerbations. It differs in its character in different situations: being infinitely more inflammatory in some places than in others, and hence the success obtained by different modes of treatment. It chiefly affects children, and is apt to recur, but the subsequent attacks are usually less and less severe.

As a general rule, it requires the most active treatment; bleeding from the arm or neck, so as to induce paleness; leeches applied to the neck, &c., according to the age,—the warm bath, blisters to the neck or chest, and purgatives. Formidable as the disease may be, if this plan be followed *early*, it will generally be successful. Many specifics have been recommended, but the search has been idle and fruitless. The majority of cases of what are called croup are not of this inflammatory cast; but are more of a spasmodic character, and have been termed by the French *faux croup*, and with us are occasionally termed *spasmodic croup*. They generally yield to an emetic and the warm bath.

CYNANCHE TRACHEALIS SPASMODICA. Asthina acutum. A. Thymicum—c. Ulcerosa, Cynanche maligna.

CYNANCHICA. Medicines used in cases of quinsy were formerly so called.

CYANCHORTHOPNŒA, Cynanche trachealis.

CYNANCHUM. Same etymon. A genus of plants, *Ord.* Asclepiadaceæ, of which the following are used in medicine.

CYNANCHUM ANGEL. *C. oleæfolium*—c. *Ipecacuanha*, *Tylophora asthmatica*.

CYNANCHUM MONSPELIACUM, *Scammony* of *Montpellier*. The plant furnishes a blackish kind of gum-resin, which is purgative, and but little used.

CYNANCHUM OLEÆFOLIUM, *C. argel*, *Solenostemma argel*, *Argel*. An Egyptian, Nubian and Arabian shrub, the leaves of which form a portion of most samples of Alexandrian senna. They resemble senna in their action.

CYNANCHUM TOMENTOSUM, *Tylophora asthmatica*—c. *Vincetoxicum*, *Asclepias vincetoxicum*—c. *Viriditorum*, *Tylophora asthmatica*—c. *Vomitiorum*, *Tylophora asthmatica*.

CYNANTHEMIS, *Anthemis cotula*.

CYNANTHROPŒIA, from *κυνος*, *kyvos*, 'dog,' and *ανθρωπος*, 'man.' A variety of melancholia, in which the patient believes himself changed into a dog; and imitates the voice and habits of that animal.

CYNARA. *C. Scolymus*, *Cinara scolymus*, *κίναρα*. The systematic name of the *Artichoke*, *Artichoculum*, *Articoaculus*, *Articoaculus lavis*, *Cotus nigra*, *Carduus sativus non spinosus*, *Cinara hortensis*, *Scolymus sativus*, *Carduus sativus* seu *domesticus* *cypriote* *major* seu *altilis*, anciently *κακτος*, *caetus*, (F.) *Artichaut*. *Ord.* Compositæ. *Sex. Syst.* Syngenesia Polygamia æqualis. Indigenous in the southern parts of Europe. Much used as an agreeable article of diet. The juice of the leaves, mixed with white wine, has been given in dropsies.

CYNICUS. *Cynoidea*, from *κυνος*, *kyvos*, 'a dog.' *Cynic*, (F.) *Cynique*. Relating to, or resembling a dog. *Cynic spasmus* is a convulsive contraction of the muscles of one side of the face,—dragging the eye, cheek, mouth, nose, &c., to one side. See *Canine*.

CYNIPS QUERCUS FOLII, see *Quercus infectoria*.

CYNOCOPRUS, *Album græcum*.

CYNOCRAMBE, *Mercurialis perennis*.

CYNOCTONON, *Aconitum*.

CYNOCTYTISUS, *Rosa canina*.

CYNODECTOS, from *κυνος*, 'a dog,' and *εχγω*, 'I bite.' One who has been bitten by a dog.—*Dioscorides*.

CYNODES, *Cynicus*.

CYNODESMION, *Frænum penis*.

CYNODESMUS, *Frænum penis*.

CYNODON DACTYLON, (*κυνος*, 'a dog,' and *οδον*, *οδοντος*, 'a tooth,') *Panicum* seu *Digitaria dactylon*, *Bermuda grass*; indigenous. *Ord.* Gramineæ. The roots have been used as a substitute for *sarsaparilla*.

CYNODONTES, *Canine teeth*.

CYNOGLOSSUM, from *κυνος*, *kyvos*, 'a dog,' and *γλωσσα*, 'a tongue.' *Cynoglossum officinale* seu *bicolor*, *Lingua Canina*, *Hound's tongue*, *Caballatton*, (F.) *Langue de Chien*. *Ord.* Boraginæ. See *Syst. Pentandria Monogynia*. It is aromatic and mucilaginous, and has been supposed to be possessed of narcotic properties.

CYNOGLOSSUM AMPLEXICAULE, *C. Virginicum*.

CYNOGLOSSUM VIRGINICUM, *C. amplexicaule*, *Wild Camfrey*; indigenous; has mucilaginous properties. The leaves are smoked, and are said to produce effects like those of tobacco. [?]

CYNOLOPHOL, Spinous processes of the vertebrae.

CYNOLYSSA, *Hydrophobia*.

CYOMETRA AGALLOCHUM, *Agallochum*.

CYNOMOIA, *Plantago psyllium*.

CYNOMORIUM COCCINEUM, from *κυνος*, 'a dog,' and *μορον*, 'the penis.' (F.) *Cham-pignon de Malte*. *Ord.* *Balanophoreæ*. Improperly called *Fungus Meliten'sis*, or *Fungus of Malta*, as it is not a fungus. The powder has been given as an astringent in hemorrhage, dysentery, &c.

CYNOREXIA, *Boulimia*.

CYNORRHODON, *Rosa canina*.

CYNOSEATOS, *Rosa canina*.

CYNOSORCHIS, *Orchis mascula*.

CYNOSPASTUM, *Rosa canina*.

CYON, *κυνος*. The word sometimes signifies the frænum of the prepuce; at others, the penis.

CYOPHORIA, *Pregnancy*.

CYTOTROPHY, *Cytotrophia*, *Embryotrophy*, *Embryotrophia*, from *κυος*, 'embryo,' and *τροφειν*, 'to nourish.' Nutrition of the embryo. Foetal nutrition.

CYPARISSUS, *Cupressus sempervirens*.

CYPERUS ANTIQUORUM, *κυνειπος*, *Lawsonia inermis*—c. *Indicus*, *Curcuma longa*.

CYPERUS LONGUS, *C. Romanus*, *Galungale*, (F.) *Souchet odorant*. *Ord.* *Cyperaceæ*. It possesses aromatic and bitter properties, but is not used. See, also, *Dorstenia contrayerva*.

CYPERUS OBOBUS, *Dorstenia contrayerva*—c. *Peruanus*, *Dorstenia contrayerva*—c. *Romanus*, *C. longus*.

CYPERUS ROTUNDUS, *C. tetraschachys*, the *Round Cyperus*, (F.) *Souchet rotund*. It is a more grateful aromatic bitter than the *C. longus*.

CYPERUS TETRASTACHYS, *C. rotundus*.

CYPHEL, *Sempervivum tectorum*.

CYPHOMA, *Cyphos*, *Cyphosis*, *Cyrto'ma*, *Cyrto'sia*, *Opiethocyphosis*, from *κυφος*, 'gibbous,' Gibbosity of the spine. See *Vertebral Column*, &c. *Cyrto'sia* forms a genus in the order *Dys-thet'ica*, and class *Hæmat'ica* of Good, and is defined: 'head bulky, especially anteriorly; stature short and incurvated; flesh flabby, rhabid, and wrinkled.' It includes *Cretinism*, and *Rickets*.

CYPHOSIS, *Gibbositatis*, see *Hump*, and *Kyphosis*.

CYPRÈS, *Cupressus sempervirens*.

CYPRESS, *Cupressus sempervirens*, *Taxodium distichum*—c. Bald, *Taxodium distichum*—c. Tree, *Liriodendron*.

CYP'RINUM (O'LEUM), *κυπρινον* (ελαιον), *Oil of Cypress*; prepared of olive oil, cypress flowers, calamus, myrrh, cardamom, inula, bitumen of Judea, &c. It was formerly employed in certain diseases of the uterus, pleurisies, &c., and was regarded to be both stimulant and emollient.

CYPRIPEDIVM ACAULĒ, (*Κυπρις*, 'Venus,' and *πόδιον*, 'a sock or buskin.') *C. hultmii*, *Stemless Ladies' Slipper*; *Moccasin Flower*; *Noah's Ark*. Ord. Orchidaceæ. Indigenous; flowers in May and June. The roots are used by steam-doctors in nervous diseases, like valerian.

CYPRIPEDIUM CALCEOLUS, *C. Luteum*, *C. spectabile*—c. *Canadense*, *C. spectabile*—c. *Flavescens*, *C. Luteum*—c. *Humile*, *C. acule*.

CYPRIPEDIUM LUTĒUM, *C. Calceolus* seu *Flavescens* seu *Pubescens* seu *Parviflorum*, *Yellow ladies' slipper*, *Moccasin flower*, *Moccasin plant*, *Yellows*, *Bleeding heart*, *American valerian*, *Yellow umbil*, *male Mervine*, *Noah's ark*, (F.) *Sabat de Venus jeune*. An indigenous plant, which is found all over the United States; blossoming in May and June. The root is considered to be antispasmodic, and is used in the same cases as valerian.

CYPRIPEDIUM PARVIFLORUM, *C. Luteum*—c. *Pubescens*, *C. Luteum*.

CYPRIPEDIUM SPECTABILĒ, *C. Calceolus* seu *Canadense*, *Showy ladies' slipper*, is the most beautiful of the genus. It is said to possess properties identical with those of *C. Luteum*.

CYPRIVM, *Cuprum*.

CYPSELE, *Cerumen*.

CYRCEON, *Anus*.

CYRTOIDES, *Cuboid*.

CYRTOMA, *Curvature*, *Cyphoma*.

CYRTONOSOS, *Rachitis*.

CYSSANTHEMON, *Cyclamen*.

CYSSARUS, *Anus*, *Rectum*.

CYSSOPHYLLON, *Cyclamen*.

CYS'SOTIS, from *κυσος*, 'the anus.' The lower part of the rectum: tenesmus.

CYST, *Kyst*.

CRYSTAL'GIA, *Cystidall'gia*, from *κυστις*, 'the bladder,' and *αλγος*, 'pain.' Pain in the bladder.

CYSTANENCEPHALIA, from *κυστις*, 'a bladder,' and *anencephalia*, 'absence of brain.' A monstrosity, in which, in place of a brain, a bladder is found filled with fluid.—G. St. Hilaire.

CYSTATROPH'IA, from *κυστις*, 'bladder,' and *ατροφία*, 'atrophy.' *Acysturo trophic*,—*Piorry*. Atrophy or wasting of the bladder.

CYSTAUCHENOTOM'IA, *Cystotrachelotomia*, *Cystidotrachelotomia*, from *κυστις*, 'a bladder,' *αυχνη*, 'the neck,' and *τομή*, 'incision.' An incision into the neck of the bladder. See *Lithotomy*.

CYSTAUX'Ē, *Hypertroph'ia vesicæ urinariæ*, *Callositas vesicæ*; from *κυστις*, 'bladder,' and *αὐξή*, 'increase.' Hypertrophy of the coats of the urinary bladder.—Fuchs. See *Cysthypersarcosis*.

CYSTECTASY, *Lithectasy*.

CYSTENCEPH'ALUS, from *κυστις*, 'bladder,' and *κεφαλή*, 'head.' A monster having a head with a vesicular brain.—G. St. Hilaire.

CYSTEOLITHOS, from *κυστις*, 'the bladder,' and *λίθος*, 'a stone.' Stone in the bladder. Also, a medicine, employed to dissolve or break stone.

CYSTERETHISMUS, *Irritabil'itas seu Impatientia Vesicæ*, from *κυστις*, 'the bladder,' and *ερεθισμός*, 'I irritate.' Irritability of the bladder.

CYSTEPAT'ICUS, from *κυστις*, 'the bladder,' and *πάρα*, 'the liver.' Belonging to the gall-bladder and liver. This name was given, formerly, to imaginary excretory ducts for the bile,

which were supposed to pass directly from the liver to the gall-bladder.

CYSTHEPATOLITH'ASIS, *Cystidopatlith'asis*, *Cholelith'ia*, *Cholelith'asis*, from *κυστις*, 'the bladder,' *πάρα*, 'the liver,' and *λίθωσις*, 'pain caused by a calculus.' The aggregate of phenomena caused by the presence of biliary calculi. See *Calculi*, *biliary*.

CYTHITIS, *Kysthitis*.

CYTHIUS, *Anus*, *Vulva*.

CYTHIPERSARCO'SIS, from *κυστις*, 'bladder,' *ὑπερ*, 'over,' and *σαρκώσις*, 'a fleshy growth,' *Excrescentia Vesicæ urinariæ*. A fleshy thickening of the coats of the bladder.

CYSTIC, *Cysticus*, (F.) *Cistique*, from *κυστις*, 'a bladder.' Belonging to the gall-bladder. Also, containing *Cystæ*, *Cystous*.

CYSTIC ARTERY, *Arteria Cystica*, is given off from the right branch of the hepatic, and divides into two branches, which proceed to the gall-bladder. It is accompanied by two cystic veins, which open into the vena porta abdominalis.

CYSTIC BILE. Bile contained in the gall-bladder.

CYSTIC CAL'CULI. Calculi formed in the gall-bladder. See *Calculi*.

CYSTIC DUCT, *Ductus cysticus*, *Meatus cysticus*. The duct proceeding from the gall-bladder, which, by its union with the hepatic, forms the *ductus communis choledochus*.

CYSTIC OXYD, *Oxyd*, *cystic*—**CYSTIC OXYD CAL'CULI**, see *Calculi*, *Urinary*.

CYSTIC SARCO'MA, of *AB'ERNETHY*, *Emphy'ma Sarcoma cellulolum*. Tumour, cellular or cystous; cells oval, currant-sized or grape-sized, containing serous fluid; sometimes caseous. Found in the thyroid gland (forming bronchocele,) testis, ovarium, &c.

CYSTICA, *Cystic Remedies*. Such medicines as were formerly believed proper for combating diseases of the bladder.

CYSTICER'CUS, from *κυστις*, 'a bladder,' and *κερκος*, 'a tail.' A genus of entozoa of the family of the hydatids, distinguished by the caudal vesicle in which the cylindrical or slightly depressed body of the animal terminates. The *Cysticer'cus cellulosa*, (F.) *Cysticerque du Tissu cellulaire*, *Hydatis finna*, has been often found in the cellular membrane.

CYSTICERQUE DU TISSU CELLULAIRE, *Cysticerus cellulosa*.

CYSTIDALGIA, *Cystalgia*.

CYSTIDELCO'SIS, from *κυστις*, 'bladder,' and *ἐλκωσις*, 'ulceration.' Suppuration or ulceration of the urinary bladder.

CYSTIDEPAT'ICUS, *Hepatocystic*.

CYSTIDEPATOLITHIASIS, *Cysthepatolithiasis*.

CYSTIDOBLENNORRHŒA, *Cystirrhœa*.

CYSTIDOCATARRHUS, *Cystirrhœa*.

CYSTIDOCELE, *Cystocele*.

CYSTIDOPLEGIA, *Cystoparalysis*.

CYSTIDORRHAGIA, *Cystorrhagia*.

CYSTIDORRHEXIS, *Cystorrhæxis*.

CYSTIDORRHŒA, *Cystirrhœa*.

CYSTIDOSMATOTOMIA, *Cystosomatotomy*.

CYSTIDOSPASMUS, see *Cystospastic*.

CYSTIDOSTENOCHORIA, *Stricture of the urinary bladder*.

CYSTIDOTOMIA, *Cystotomy*.

CYSTIDOTRACHELOTOMIA, *Cystauchentomy*.

CYSTINE, *Oxyd*, *cystic*.

CYSTINURIA, *Urine*, *cystinic*.

CYSTINX, *Vesicula*.

CYSTIPHLOGIA, *Cystitis*.

CYSTRIRHAG'IA, from *κυστις*, 'the bladder,' and *ρρηγνυσι*, 'I break forth.' Hemorrhage from the

bladder, (F.) *Hémorrhagie de la Vessie*. By some used synonymously with *cystirrheuma*.

CYSTIRRHEUMA, *Cystirrheuma*.

CYSTIRRHEA, from *κυστις*, 'the bladder,' and *ρῆω*, 'I flow.' *Parduria Stillatitia Mucosa*, *Blennorrhia*, *Blennorrhoea urinalis seu vesicæ*, *Cysto-blennorrhoea*, *Cystorrhoea*, *Cystido-blennorrhoea*, *Ischuria*, *Cysto-phlegmaticea*, *Tenesmus Vesicæ Mucosus*, *Urina mucosa*, *Cysticater-rhus*, *Cystido-catar-rhus*, *Cystidorrhoea*, *Urocysto-catar-rhus*, *Dysuria Mucosa*, *Pyuria Mucosa seu viscidula seu serosa*, *Morbus cystophlegmaticeus*, *Catar-rhus vesicæ*, (F.) *Cystite muqueuse*, *Blennocystite*, *Flux muqueux de la vessie*, *Catar-rhe vésical*, *Blennocystorrhée*,—Piorry. A copious discharge of mucus from the bladder, passing out with the urine, and generally attended with dysuria. It is commonly dependent upon an inflammatory or subinflammatory condition of the lining membrane. The treatment must be regulated by the cause. If it be not produced by an extraneous body; the antiphlogistic plan,—the exhibition of warm diluents, and keeping the surface in a perspirable state, by wearing flannel, are indicated. Some of the turpentine may, at times, be given with advantage; and astringent or other appropriate injections be thrown into the bladder.

CYSTIS, Follicle, Urinary bladder—c. *Chole-dochus*, Gall-bladder—c. *Fellea*, Gall-bladder—c. *Serosa*, Hygroma.

CYSTITE, *Cystitis*—c. *Muqueuse*, *Cystitis*.

CYSTITIS, from *κυστις*, 'the bladder.' *In-flammatio Vesicæ*, *Empresma Cystitis*, *Cystitis urica*, *Uro-cystitis*, *Cystiphlogia*, *Cystophlogia*, *Inflammation of the bladder*, (F.) *Cystite*, *Inflammation de la vessie*, *Cysturite*,—Piorry; characterized by pain and swelling in the hypogastric region; discharge of urine painful or obstructed, and tenesmus. It may affect one or all of the membranes; but commonly it is confined to the mucous coat. In the chronic condition, it appears in the form of *cystirrheuma*. It must be treated upon the same energetic principles as are required in other cases of internal inflammation; venesection, general and local, the warm bath, warm fomentations, warm, soothing enemata, diluents, &c. *Cantharides* must be avoided, even in the way of blisters, unless with precautions, as the disease is often occasioned by them.

CYSTITIS FELLEA, *Cholecystitis*—c. *Urica*, *Cystitis*.

CYSTITOME, *Cystit'omus*, *Cyst'otome*, *Kibis'itome*, *Kystitome*, *Kystotome*, from *κυστις*, 'the bladder,' and *τεμνειν*, 'to cut.' An instrument, invented by Lafaye, for dividing the anterior part of the capsule of the crystalline in the operation for extracting cataract. It was formed like the *Pharyngotome*.

CYSTO. In composition, relating or appertaining to a bladder, and generally to the urinary bladder.

CYSTOBLAST, *Cytoblast*.

CYSTO-BLENNORRHOEA, *Cystirrheuma*.

CYSTO-BUBONCELE, from *κυστις*, 'the bladder,' *βουβων*, 'the groin,' and *κηλη*, 'a tumour.' Hernia of the bladder through the abdominal ring.

CYSTO-CATARRHUS, *Cystirrheuma*.

CYSTOCELE, *Cystocelia*, *Cistocelē*, *Cystido-celē*, from *κυστις*, 'the bladder,' and *κηλη*, 'a tumour.' *Hernia vesicæ urinaria*, *Hernia of the bladder*, (F.) *Hernie de la Vessie*. It is not common. It occurs, most frequently, at the abdominal ring; less so at the crural arch, perinæum, vagina, and foramen thyroideum. It may exist alone, or be accompanied by a sac, containing some abdominal viscus. The tumour is soft and

fluctuating; disappears on pressure, and increases in size when the urine is retained. It must be reduced and kept in position by a truss. *Vaginal cystocele* is kept in place by a pessary.

CYSTOCLE BILIOSA, *Turgescentia vesiculæ fellæ*.

CYSTODYN'IA, from *κυστις*, 'the bladder,' and *ὄδυν*, 'pain.' Pain of the bladder; particularly rheumatic pain.

CYSTOHÆM'IA, (F.) *Cystohémie*, from *κυστις*, 'bladder,' and *αἷμα*, 'blood.' Congestion of blood in the bladder.

CYSTOID, *Cystoides*, *Cyst-like*, from *κυστις*, 'a bladder or cyst,' and *εἶδος*, 'resemblance.' That which resembles a cyst, as 'cystoid tumour.'

CYSTO-LITHIASIS, see *Calculi*, vesical.

CYSTO-LITH'IC, *Cystolithicus*, (F.) *Cystolithique*, from *κυστις*, 'the bladder,' and *λίθος*, 'a stone.' Relating to stone in the bladder.

CYSTOMA, see *Kyst*.

CYSTO-MEROCE'LE, from *κυστις*, 'the bladder,' *μερος*, 'the thigh,' and *κηλη*, 'hernia.' Femoral hernia, formed by the bladder protruding beneath the crural arch.

CYSTON'CUS, from *κυστις*, 'the bladder,' and *ογκος*, 'tumour.' Swelling of the bladder.

CYSTOPARALYSIS, *Cystidoparalysis*, *Cystidoplegia*, *Cystoplegia*, *Cystoplex'ia*, *Acystinervia*, *Acystinervia*, *Amysthénie cystique*,—Piorry, from *κυστις*, 'bladder,' and *παράλυσις*, 'palsy.' Paralysis of the urinary bladder. See *Enuresis*.

CYSTO-PHLEGMAT'IC, *Cystophlegmaticus*, from *κυστις*, 'the bladder,' and *φlegμα*, 'mucus, phlegm.' Belonging to the vesical mucus. *Morbus cystophlegmaticus*. *Cystirrheuma*.

CYSTOPHLOGIA, *Cystitis*.

CYSTOPHTH'ISIS, *Phthisis vesicalis*, from *κυστις*, 'the bladder,' and *φθίω*, 'I consume.' Consumption from ulceration of the bladder,—*Ul'cera seu Helco'sis vesicæ*.

CYSTOPLAS'TIC, *Cystoplasticus*, from *κυστις*, 'the bladder,' and *πλασσω*, 'I form.' An epithet for an operation for the cure of fistulous openings into the bladder, *Cystoplasty*, (F.) *Cystoplastie*; sometimes restricted to the cure by translation of skin from a neighbouring part.

CYSTOPLEGIA, *Cystoparalysis*.

CYSTOPLEG'IC, *Cystopleg'icus*. Belonging to paralysis of the bladder; from *κυστις*, 'the bladder,' and *πλησσω*, 'I strike.'

CYSTOPLEX'IA, *Cystoparalysis*.

CYSTOPTO'SIS, from *κυστις*, 'the bladder,' and *πτειν*, 'to fall.' Relaxation of the inner membrane of the bladder, which projects into the canal of the urethra.

CYSTOPY'IC, *Cystopy'icus*, (F.) *Cystopyique*, from *κυστις*, 'the bladder,' and *πυω*, 'pus.' Relating to suppuration of the bladder.

CYSTORRHAG'IA, *Hæmaturia cystica*, *Hæmorrhag'ia vesicæ*, *Strangu'ria cruent'a*, *San-guinis fluor vesicæ*, from *κυστις*, 'the bladder,' and *ρῆγν*, 'rupture.' A discharge of blood from the vessels of the urinary bladder.

CYSTORRHEU'MA, *Cystirrheu'ma*, *Rheuma-tismus vesicæ urinaria*; from *κυστις*, 'the bladder,' and *ρῆγμα*, 'defluxion,' 'rheumatism.' Rheumatism of the bladder.

CYSTORRHEX'IS, *Cystidorrhex'is*; from *κυστις*, 'bladder,' and *ρῆξις*, 'rupture.' Rupture of the urinary bladder.

CYSTORRHEA, *Cystirrheuma*.

CYSTOSARCO'MA, from *κυστις*, 'a bladder or cyst,' and *sarcoma*. A tumour consisting of a combination of cysts and cystoids, so called by J. Müller.

CYSTOSOMATOM'IA, *Cystosomatotom'ia*, *Cystidosomatotom'ia*, from *κυστις*, 'the bladder,' *σωμα*,

'body,' and *τομή*, 'incision.' An incision into the body of the bladder.

CYSTOSPASMUS, see Cystospastic.

CYSTOSPAS'TIC, *Cystospas'ticus*, from *κυστίς*, 'the bladder,' and *σπασσω*, 'I contract.' Relating to spasm of the bladder, and particularly of its sphincter;—*Cystidospas'mus*, *Spasmus Vesicæ*, *Ischuria spasmodica*, (F.) *Spasme de la Vessie*.

CYSTOSPERMI'TIS, (F.) *Cytospermite*, from *κυστίς*, 'bladder,' *σπέρμα*, 'sperm,' and *itis*, 'denoting inflammation.' Inflammation of the vesiculae seminales.

CYSTOSTENOCHOR'IA, *Cystidostenochoria*, *Strictura vesicæ*, *Vesicæ succata*. A stricture, narrowness, inequality or saccated condition of the urinary bladder.

CYSTOTHROMB'OID, *Cystothromboïdes*, from *κυστίς*, 'the bladder,' and *σπῆγμα*, 'a clot.' Relating to the presence of clots in the bladder.

CYSTOTOME, *Cystotomus*, from *κυστίς*, 'the bladder,' and *τομή*, 'to cut.' An instrument intended for cutting the bladder. Instruments of this kind have been more frequently, although very improperly, called *Lithotomes*.

CYSTOTOM'IA, *Cystidotomia*, same etymon. Incision of the bladder; *Sectio vesicæ*. *Cystotomia* means cutting into the bladder for any purpose; (F.) *Incision de la vessie*. Commonly, it is applied to the puncturing of the bladder for the purpose of removing the urine; whilst *Lithotomy* has been employed for the incisions made with the view of extracting calculi from the bladder. See *Lithotomy*.

CYSTOTRACHELOTOMIA, *Cystauchenotomia*. See *Lithotomy*.

CYSTOUS, *Cystis*.

CYSTOVAR'IIUM, from *κυστίς*, 'a cyst or bladder,' and *οἰστίον*, 'ovarium.' A morbid condition, which consists in the formation of cysts in the ovary.

CYSTURITE, *Cystitis*.

CYT'INUS, *C. Hypocist'is*, *Hypocist'is*, *As'arum Hypocist'is*. Ord. Cistineæ. A small parasitical plant, which grows in the south of France and in Greece, on the roots of the woody cistus. The juice of its fruit is acid, and very astringent. It is obtained by expression, and converted into an extract, which was called *Succus Hypocist'idis*,

and was formerly much used in hemorrhages, diarrhoea, &c.

CYTIS'INA, *Cytisine*. An immediate vegetable principle, discovered by Chevalier and Lassaigne, in the seeds of *Cytisus Laburnum* or *Bean-Trefoil Tree*. Ord. Leguminosæ. *Sex. Syst.* Diadelphia Decandria. Cytising has analogous properties to emetine. In the dose of one or two grains, it produces vomiting and purging; and, in a stronger dose, acts as an acrid poison. The seeds of the *Cytisus Laburnum*, (F.) *Aubours*, have been long known to produce vomiting and purging.

CYTISMA ECZEMA, Eczema—c. Herpes, Herpes.

CYTISO-GENISTA, *Spartium scoparium*.

CYTISUS LABURNUM, see *Cytisina*—c. *Scoparium*, *Spartium scoparium*.

CYTIT'IS, *Cyti'tis*, *Dermati'tis*, *Dermi'tis*, *Cuti'tis*, *Cori'i'tis*, from *κυστίς*, 'the skin,' and *itis*, 'denoting inflammation.' Inflammation of the skin.

CYTOBLAST, *Cytoblast*, from *κυτος*, 'cell,' and *βλαστος*, 'germ.' *Cell-germ*, *Nu'cleus*, (F.) *Noyau*. A granule, from which all animal and vegetable bodies are presumed to be formed. When the nucleus or cytoblast forms a cell, and is attached to its walls, the *germinal cell*, thus formed, is called a *nucleated cell*. When the nucleus contains a simple granule, the latter is termed a *nucleolus*, (F.) *Nucléole*, *Nucléole*.

CYTOBLASTE'MA; (F.) *Cytoblastème*, same etymon. *Intercell'ular substance*, *Hy'aline substance*, *Substantia vit'rea seu hyal'ina*, *Matrix*. The gum or mucus in the vegetable, and probably the liquor sanguinis after transudation from the vessels in the animal, in a state fully prepared for the formation of the tissues.—Schwann and Schleiden. By many, *Blastema* is preferred, inasmuch as it does not convey the idea of cellular development.

CYTOGENET'IC, *Cytogenet'icus*, (F.) *Cytogénétique*, same etymon as the next. Relating or belonging to cell formation.

CYTOG'ENY, *Cytogen'esis*, from *κυτος*, 'cell,' and *γενεσις*, 'generation.' Cell formation, cell genesis.

CYT'TAROS, Glans.

D.

The figure of the Greek Δ, according to Galen, was the sign for quartan fever.

DABACH, Viscum album.

DACNE'RON, *δακνηρον*, from *δακνεν*, 'to bite.'

An ancient name for a collyrium, composed of oxide of copper, pepper, cadmia, myrrh, saffron, gum Arabic, and opium.

DACRY, *Dac'ryma*, *Dac'ryon*, *δακρυ*, *δακρυμα*, *δακρυον*, 'a tear.' Hence:

DACRYADENAL'GIA, *Dacryoadenal'gia*, from *δακρυω*, 'I weep,' *αδην*, 'a gland,' and *αλγος*, 'pain.' Disease or pain in the lachrymal gland.

DACRYADENIT'IS, *Dacryoadenit'is*; from *δακρυω*, 'I weep,' *αδην*, 'a gland,' and *itis*. Inflammation of the lachrymal gland.

DACRYALLGEO'SIS, from *δακρυ*, 'a tear,' and *αλλοιωσις*, 'change.' A morbid condition of the tears.

DACRYDION, *Convolvulus scammonia*.

DACRYGEO'SIS, from *δακρυω*, 'I weep,' and *γελω*, 'I laugh.' A kind of insanity, in which the patient weeps and laughs at the same time.

DACRYHEMOR'RHYSIS; *Sanguineous Lachrymation*, from *δακρυ*, 'a tear,' and *αιμorrhesis*, 'hemorrhage.' A flow of bloody tears.

DACRYNOMA, Epiphora.

DACRYOADENALGIA, *Dacryadenalgia*.

DACRYOADENITIS, *Dacryadenitis*.

DACRYOBLENNORRHŒ'A, from *δακρυω*, 'I weep,' *βλεννα*, 'mucus,' and *ρῶ*, 'I flow.' Discharge of tears mixed with mucus.

DACRYOCYSTAL'GIA, from *δακρυω*, 'I weep,' *κυστίς*, 'a sac,' and *αλγος*, 'pain.' Disease or pain in the lachrymal sac.

DACRYOCYSTIS, Lachrymal sac.

DACRYOCYSTITIS, from *δακρυν*, 'a tear,' and *κυστις*, 'bladder.' Inflammation of the lachrymal sac.

DACRYOCYSTOSYRINGOKATAKLEISIS; from *dacryocystis*, the 'lachrymal sac,' *εuryzē*, 'a pipe, a fistula,' and *κατακλεισις*, 'a locking up.' A term, proposed by Dieffenbach for the healing of lachrymal fistulæ by transplantation. The operation consists in paring the edges of the fistula, loosening the borders, and assisting the requisite tegumental displacement by lateral incisions.

DACRYO'DES, *Lacrymō'us*, from *δακρυω*, 'I weep.' Resembling tears:—hence, *Ulcus dacryodes*. A sanious ulcer, a weeping sore.

DAC'RYOLITE, *Dacryol'ithus*, *Dac'ryolith*, (F.) *Dacryolithe*, from *δακρυω*, 'I weep,' and *λιθος*, 'a stone.' A concretion found in the lachrymal passages.

DACRYOLITHI'ASIS; same etymon as the last. The formation of concretions in the tears.

DACRYO'MA. Same etymon. The effusion of tears, occasioned by an occlusion of the puncta lachrymalia.—Vogel.

DACRYON, Tear.

DACRYOPCE'US, from *δακρυω*, 'I weep,' and *ποιω*, 'I make.' A substance which excites the secretion of tears,—as the onion, horse-radish, garlic, &c.

DACRYOPUS, from *δακρυω*, 'I weep,' and *ωψ*, 'the eye,' a weeping eye. A tumefaction of the lachrymal passages.

DACRYOPYORRHŒ'A, *Pyorrhœ'a via'rum lacrymalium*, from *δακρυ*, 'a tear,' *πυον*, 'pus,' and *ρεω*, 'to flow.' A discharge of tears mixed with purulent matter.

DACRYORRHŒ'A, *Dacryorrhœ'a*, *Dacryor'rhysis*, *Dacryor'rhysis*, from *δακρυ*, 'a tear,' and *ρεω*, 'to flow.' A morbid flux of tears.

DACRYORRHYSIS, *Dacryorrhœa*.

DACRYSOLENT'ITIS, from *δακρυ*, 'a tear,' *σωλην*, 'a canal,' and *itis*, denoting inflammation. Inflammation of the lachrymal ducts.

DACRYOSYRINX, *Fistula lacrymalis*.

DACRYRRHŒA, *Dacryorrhœa*.

DACRYRRHYSIS, *Dacryorrhœa*.

DACTYLE'THRA, *Dactyl'i'thra*, from *δακτυλος*, 'a finger.' A name given by the ancients to different topical applications, having the form of a finger, and proper for being introduced into the throat to excite vomiting.

DACTYLETUS, *Hermodactylus*.

DACTYL'ION, *Dactyl'ium*, (F.) *Doigts palmés*, *Webbed fingers*, from *δακτυλος*, 'a finger.' The union of the fingers with each other. This affection is generally congenital; but it may be owing to burns, ulcerations, inflammation of the fingers, &c.

DACTYL'IOS, from *δακτυλος*, 'a finger.' A troch or lozenge, when shaped like a finger. The anus.

DACTYLITIS, *Paronychia*.

DACTYL'IOS, from *δακτυλος*, *annulus*, 'a ring.'

DACTYL'IOS ACULEA'TUS. A worm of a light colour, annulated, cylindrical, but tapering slightly towards both extremities, from two-fifths to four-fifths of an inch long, which has been found in the urine.

DACTYLODOCHME, *Dochme*.

DACTYLOSYM'PHYSIS, from *δακτυλος*, 'a finger,' and *συνφυσις*, 'union.' Adhesion of the fingers to each other.

DACTYLOTHE'KĒ, from *δακτυλος*, 'a finger,' and *θηκη*, 'a case or sheath.' An instrument for keeping the fingers extended when wounded.—Ambrose Paré.

DAC'TYLUS, *Dig'itus*, 'a finger.' The smallest measure of the Greeks, the sixth part of a foot. Also, the Date.

DÆDA'LEA SUAV'EOLENS. *Boletus seu Fungus Sa'ticis*, *Boletus discoidens seu suaveolens*, *Fungus albus salig'neus*, (F.) *Agaric odorant*, *Bolet odorant*. Ord. Fungi. A champignon, which grows on the trunks of old willows. It has a smell of anise, which is penetrating and agreeable; and has been recommended in phthisis pulmonalis in the dose of a scruple four times a day.

DÆDALUS, *Hydrargyrum*.

DÆDION, *Baugie*.

DÆMONIACUS, Possessed.

DÆMONOMANIA, *Demonomania*.

DÆS, *Tæda*.

DAFFING, *Insanity*.

DAFFODIL, *Narcissus pseudonarcissus*.

DAFFY'S ELIXIR, *Tinctura senæ compo-sita*.

DAFT, *Insane*.

DAG, *Stitch*.

DAISY, *Chrysanthemum leucanthemum*, *Eri-geron Philadelphicum*—d. Common, *Bellis*—d. Ox-eye, *Chrysanthemum leucanthemum*.

DALBY'S CARMIN'ATIVE. A celebrated empirical remedy, much used as a carminative for children. The following is a form for its preparation:—(*Magnesie alb. ℥ij*; *ol. mentha piper. gtt. j*; *ol. nuc. moschat. gtt. iij*; *ol. anisi, gtt. iij*; *tinct. castor. gtt. xxx*; *tinct. asafoetid. gtt. xv*; *tinct. opii, gtt. v*; *ep. pulegii, gtt. xv*; *tinct. cardam. c. gtt. xxx*; *aquæ mentha pip. ℥ij. M.*) A Committee of the Philadelphia College of Pharmacy recommend the following form:—(*Aquæ, Ox*; *Succchar. alb. ℥xxxij*; *Carbon. Potass. ℥ss*; *Carb. Mag. ℥xij*; *Tinct. Opii. f℥vj*; *Ol. ment. pip., Ol. Anethi Fœnicul. āā f℥ij. M.*)

DALCOP, *Idiot*.

DALTO'NIAN. An absurd name given to one who cannot distinguish colours; because the celebrated chemist Dalton had the defect. See *Achromatopsia*.

DALTONISM, *Achromatopsia*.

DAMSON, *Prunum Damascenum*—d. Mountain, *Quassia simarouba*—d. Tree, *Prunus domestica*.

DANCE, see *Mania*, dancing—d. St. John's, see *Mania*, dancing—d. St. Vitus's, *Chorea*, see *Mania*, dancing.

DANCING, *Salta'tio*, (F.) *Danse*. A kind of exercise and amusement, composed of a succession of motions, gestures, and attitudes, executed by measured steps to the sound of the voice or musical instrument. It is a healthy exercise.

DANCING MANIA, see *Mania*, dancing—d.

Plague, see *Mania*, dancing.

DANDELION, *Leontodon taraxacum*.

DANDRIFF, *Pityriasis*.

DANDRUFF, *Pityriasis*.

DANDY, *Dengue*.

DANEVERT, **MINERAL WATERS OF**. A spring, a league and a half from Upsal, in Sweden. The waters contain carbonic acid, holding in solution carbonate of iron, sulphate of iron, sulphates of soda and lime, chloride of sodium, and silica. It is frequently employed in medicine.

DANEWORT, *Sambucus ebulus*.

DANICH; an Arabic word, signifying the weight of 8 grains.

DANIEL'LIA THURIF'ERA. The *Frankincense Tree* of Western Africa, which furnishes the product termed *African Frankincense*.

DANSE, *Dancing*—d. de *St. Guy*, *Chorea*—d. de *St. Witt*, *Chorea*.

DAOUN SETAN, see *Urtica*.

DAPHNĒ, *Laurus*.

DAPHNE ALPI'NA, *Chama'ie'a*, *Chama'la'a*, *Widow wail*. Ord. *Thymelææ*. Ser. *Syot*. *Octandria Monogynia*. A sort of dwarf-olive. An acrid, volatile, alkaline principle has been

separated from the bark of this plant by M. Vauquelin, to which he has given the name *Daphnina*. The plants of the genus owe their vesicating property to this principle.

DAPHNÉ BOISGENTIL, *Daphne mezereum* — d. Flax-leaved, *Daphne gnidium* — d. Garou, *Daphne gnidium*.

DAPHNE GNIDIUM, *D. paniculata*, *Thymelæa*, *Th. Mouspeliaca*, *Thymelæa*, *Cneoiron*, *Spurge Flax*, *Flax-leaved Daphne*. The plant which furnishes the *Garou Bark*, (F.) *Daphné Garou*, *Sain-bois*. It is chiefly used, when used at all, for exciting irritation of the skin. The *Grana Gnidiæ*, (see *Chid'ia grana*), are acrid poisons, like all the plants of this genus, when taken in quantity. According to others, the garou bark and grana gnidia are obtained from the *daphne laureola*.

DAPHNE LAUREOLA, *D. major*, *Thymelæa laureola*. The systematic name of the *Spurge Laurel*, *Laureola*. The bark of this plant has similar properties to the last.

DAPHNE LIOTTARDI, *D. laureola* — d. Major, *D. laureola*.

DAPHNE MEZE'REUM, *D. Liottardi*, *Thymelæa mezereum*. The systematic name of the *Meze'reon*, *Mezereum*, *Spurge olive*, (F.) *Daphné Boisgentil*. The bark of the mezereon, *Meze'reum*, possesses analogous properties to the other varieties of *Daphne*. It is considered stimulant and diaphoretic; and, in large doses, is emetic. It has been employed in syphilitic cases, but its efficacy is doubtful. Soaked in vinegar, — *meze'reum acetatum*, like the other varieties of *daphne*, it has been employed to irritate the skin, especially to keep issues open.

DAPHNE PANICULATA, *D. Gnidium*.

DAPHNE LAURON, *Oleum Laurinum*, from *dagyn*, 'the laurel or bay tree,' and *elaion*, 'oil.' *Oil of Bay*.

DAPHNINE, see *Daphné Alpina*.

DARNEL, *Lolium temulentum*.

DARSEN, *Laurus cinnamomum*.

DARSINI, *Laurus cinnamomum*.

DARSIS, from *depa*, 'I excoriate,' 'I skin.'

The Greek physicians seemed to have used this word to designate the anatomical preparation, which consists in removing the skin for exposing the organs covered by it.

DARTA, *Impetigo* — d. *Excoriativa*, *Herpes exedens* — d. *Maligna*, *Herpes exedens*.

DARTOS. Same derivation: *Membræa carnosa*, *Tunica muscularis seu rubicunda scroti*, *T. duras*, *Marsupium musculo-sum*; from *dapros*, 'skinned.' A name given to the second covering of the testicle, which the ancient anatomists conceived to be muscular, but which is merely areolar. Its external surface is towards the scrotum; the internal towards the tunica vaginalis. Frederick Lobstein and Breschet consider, that it proceeds from an expansion of the fibrous cord, known by the name *Gubernaculum Testis*.

DARTRE, *Herpes*, *Impetigo*, *Pityriasis* — d. *Crustacea*, *Ecthyma impetigo* — d. *Crustacea flavescente*, *Porrigio lupinosa* — d. *Croûteuse*, *Impetigo* — d. *Écailleuse*, *Psoriasis* — d. *Fongueuse*, *Ecthyma* — d. *Furfuracée arrondie*, *Lepra*, *Lepra vulgaris* — d. *Furfuracée volante*, *Lichen*, *Pityriasis* — d. *Phlycténoidé*, *Herpes phlycténoides* — d. *Pustulense couperose*, *Gutta rosea* — d. *Pustulense dissimulé*, *Ace* — d. *Pustulense mentagra*, *Sycosis* — d. *Rongeante*, *Herpes exedens*; see *Esthimenus* — d. *Squammeuse humide*, *Eczema* — d. *Squammeuse lichénoidé*, *Psoriasis* — d. *Vive*, *Eczema*.

DARTREUX (F.), *Herpetic*. Participating in the characters of *Dartre* or *Herpes*. Also, one affected with *dartre*. *Dartre* has been used, at

one time or other, for almost every disease of the skin. See *Herpes*.

DAS'YMA, from *daus*, 'rough,' 'hairy.' A disease of the eye—the same as *trachoma*, but less in degree.—*Aëtius*, *Gorreaus*.

DAS'YTES. Same etymon. Roughness, particularly of the tongue and voice. Hairiness, *Hirsuties*.

DATE, *Pal'mula*, *Dactylus*, *Bal'anos*, *Phœ'nico*, the fruit of the *Phoenix dactylifera* seu *exzel'sa*, *Palma dactylifera*, (F.) *Datte*. The unripe date is astringent. When ripe, it resembles the fig. The juice of the tree is refrigerant.

DATE PLUM, *INDIAN*, *Diospyrus lotus*.

DATTE, *Date*.

DATURA, *D. Stramonium*.

DATU'RA SANGUIN'EA, *Red Thorn Apple*; called by the Indians of Peru *Huacacachu*, *Yerba de Huaca* (*huaca*, a grave) or *Grave-plant*, and *Bo-vachevo*. *Ord.* *Solanaceæ*. A plant from which the Peruvian Indians prepare a narcotic drink called *Tonga*.

DATU'RA STRAMO'NIUM, *Stramo'nia*, *Barynoc-calon*, *Sola'nium foetidum*, *Stramo'nium*, *S. majus* album seu *spino'sum* seu *vulgarium* seu *foetidum*, *Pomum* seu *Malum spino'sum*, *Nux methel*, *Datu'ra*, *Dutro'a*, *Daty'ra*, *Thorn Apple*, *Apple of Peru*, *Jamestown Weed*, *Jimson* or *Jimson Weed*, *Stinkweed*, (F.) *Stramaine*, *Pomme épineuse*, *Endormie*, *Herbe aux sorciers* ou *de diable*. The leaves, *Stramo'ni fo'lia*, the seeds, *Stramo'ni Semen*, and the root, *Stramo'ni Radix*, are the parts used in medicine. They are narcotic and poisonous; are given internally as narcotics and antispasmodics; and applied externally as sedatives, in the form of fomentation. The seeds are smoked like tobacco, in asthma. The dose of the powder is, gr. j to gr. viii.

DA'TURINE, *Datur'ina*, *Datu'ria*, *Datu'rium*. The active principle of the *Datura Stramonium*, separated by Brandes, a German chemist. It has not been rendered available in medicine.

DATURIUM, *Daturine*.

DATYRA, *Datura stramonium*.

DAUCI RADIX, see *Daucus Carota*.

DAUCITES VINUM. Wine, of which the *Daucus* or *Wild Carrot* was an ingredient. The seeds were steeped in must. It was formerly used in coughs, convulsions, hypochondriasis, diseases of the uterus, &c.

DAUCUS CANDIANUS, *Athamanta cretensis*.

DAUCUS CARO'TA. The systematic name of the *Carrot Plant*; *Daucus*, *D. sylvestris* seu *vulgaris* seu *sativus*, *Caucalis carota*, *Postina'ca sylvestris tenuifolia officinarum*, *Ado'rian*; *Order*, *Umbelliferæ*. (Sc.) *Meeran*, *Mirrot*, (F.) *Carotte*; *Ord.* *Umbelliferæ*. The fruit, *Carota*, — (Ph. U.S.) — and the root, *Dauci radix*, have been used in medicine. The root is sweet and mucilaginous; and the seeds have an aromatic odour, and moderately warm, pungent taste. The root has been used, as an emollient, to fetid and ill-conditioned sores. The seeds have been regarded as stomachic, carminative, and diuretic; but they have little efficacy. The seeds of the wild plant are, by some, preferred to those of the garden.

DAUCUS CRETICUS, *Athamanta cretensis* — d. *Cyanopus*, *Pimpinella magna* — d. *Sativus*, *D. carota* — d. *Sepirinus*, *Scandix cerefolium* — d. *Sylvestris*, *D. carota* — d. *Vulgaris*, *D. carota*.

DAUPHINELLE, *Delphinium staphisagria*.

DAVIDSON'S REMEDY FOR CANCER, see *Conium maculatum*.

DAVIER, *Dentagra*.

DAWLISH, CLIMATE OF. A town in Devonshire, frequented by phthical invalids during the winter. It is well protected from northerly winds, and also from the violence of the south

westerly gales. It offers, however, but a confined space.

DAWPIT, see Fatuitas.

DAX, MINERAL WATERS OF. Dax is a city two leagues from Bordeaux, where there are four springs, that are almost purely thermal; containing only a very small quantity of chloride of magnesium, and sulphate of soda and lime. Temperature 76° to 133° Fahrenheit.

DAYBLINDNESS, Nyctalopia.

DAYMARE, Incubus vigilantium.

DAYNETTLE, Lamium album.

DAYSIGHT, Hemeralopia—d. Vision, Hemeralopia.

DAZZLING, *Caligatio*, (F.) *Éblouissement*. A momentary disturbance of sight, occasioned either by the sudden impression of too powerful a light, or by some internal cause; as plethora.

DE VENTRE INSPICIENDO, 'of inspecting the belly.' Where there is reason to suppose that a woman feigns herself pregnant, a writ *de ventre inspiciendo* may be issued to determine whether she be so or not. Until recently, in England, the decision was left to twelve matrons and twelve respectable men, according to the strict terms of the ancient writ.

DEAD, (Sax.) *deaf*, *Mortuus*, (F.) *Mort*. Deprived of life, exanimate.

DEADHOUSE, *Morgue*.

DEADLY, Lethiferous, Mortal.

DEAD MEN'S BELLS, Digitalis.

DEADNETTLE, Lamium album.

DEAF: same etymon as Deafness. *Surdus*, (F.) *Sourd*, (Sc.) *Dull*. Devoid of hearing. Hard of hearing.

DEAF-DUMBNESS, Mutitas surdorum.

DEAFNESS, (Sax. *deaf*), *Surditas*, *Cophosis*, *Dyscoi'a*, *Dysecoi'a organica*, *Barycoi'a*, *Hardness of hearing*, *Hypocophosis*, *Subsurditas*, *Bradycoi'a*, *Bradydycoi'a*, (F.) *Surdité*, *Durété de l'ouïe*, *Dysécécé*. Considerable diminution or total loss of hearing. It may be the effect of acute or chronic inflammation of the internal ear, paralysis of the auditory nerve or its pulpy extremity, or of some mechanical obstruction to the sonorous rays. In most cases, however, the cause of the deafness is not appreciable, and the treatment has to be purely empirical. Syringing the ears, dropping in slightly stimulating oils, fumigations, &c., are the most likely means to afford relief.

DEAFNESS, TAYLOR'S REMEDY FOR, see Allium.

DEALBATIO, Dealbation, Paleness.

DEALBATION (F.), *Dealbatio*, *Blanching*. Applied, particularly, to the whitening of bones for anatomical purposes.

DEAMBULATIO, Walking.

DEARCH, Nanus.

DEARTICULATIO, Diarthrosis.

DEASCIATIO, Apocrepnismum.

DEATH, (Sax. *deaf*), (Sc.) *Dede*, *Deid*; *Apotheosis*, *Abiosis*, *Extinctio*, *Obitus*, *Psychoraghi'a*, *Psychorrhagi'a*, *Letum*, *Letum*, *Mors*, *Interitus*, *Thanatos*, *Decease*, (F.) *Mort*. Definitive cessation of all the functions, the aggregate of which constitute life. *Real Death*, *Apothanasia*, is distinguished from asphyxia or apparent death—the latter being merely a suspension of those same functions. But it is often difficult to judge of such suspension, and the only certain sign of real death is the commencement of putrefaction. At times, therefore, great caution is requisite to avoid mistakes. Death is commonly preceded by some distressing symptoms, which depend on lesion of respiration, circulation, or of the cerebral functions, and which constitute the *agony*. That which occurs suddenly, and without any, or with few, precursory signs, is called *sudden death*. It is ordinarily caused by disease of the heart, apoplexy, the rupture of an aneurism, or by some

other organic affection. Death is *natural*, (Sc.) *Stræ-death*, when it occurs as the result of disease: *violent*, when produced by some forcible agency. It may likewise affect the whole body, or a part only; hence the difference between *somatic* and *molecular* death.

The chief varieties of the modes of death may be thus given:

Death beginning at the heart,	{	Suddenly—Syncope	{	Suddenly—Syncope
		Gradual—Asthénia.		
"	"	in the lungs—Asphyxia.		
"	"	in the brain—Apoplexy.		
"	"	in the gray matter of the medulla,	{	Paralysis of pneumogastriacs, &c.
"	"	in the blood [?]		necræmia.

As it was at one time conceived, with Bichat, that death must commence in the heart, lungs, or brain, these organs were called *a'tria mortis*, 'the halls of death.'

DEATH, APPARENT, Asphyxia.

DEATH, BLACK. *Pestis nigra*, *Black Plague*, (F.) *Peste noire*. The plague of the 14th century was so called, which is supposed to have proved fatal in Europe to 25,000,000 of people.—Hecker.

DEATH OF MAN, Cicuta maculata—d. Stiffening, Rigor mortis.

DEBILIS, Infirmitas.

DEBILITANTS, *Debilitantia*, *Antidynamica*, *Philadynamica*, from *debilitare*, itself from *debilis*, quasi *dehibilis*, 'weak,' 'to weaken.' Remedies exhibited for the purpose of reducing excitement. Antiphlogistics are, hence, debilitants.

DEBILITAS, Debility—d. Erethisica, see Irritable—d. Nervosa, Neurasthenia, see Irritable—d. Visus, Asthenopia.

DEBILITY, *Debil'itas*, *Astheni'a*, *Hypostheni'a*, *Blaci'a*, *Anenerge'sia*, *Anergic'a*, *Weakness*, (F.) *Faiblesse*. A condition, which may be induced by a number of causes. It must not be confounded with *fatigue*, which is temporary, whilst debility is generally more permanent.

Debility may be *real*, or it may be *apparent*; and, in the management of disease, it is important to attend to this. At the commencement of fever, for example, there is often a degree of apparent debility, which prevents the use of appropriate means, and is the cause of much evil. Excitement is more dangerous than debility.

DÉBOITEMENT, Luxation.

DÉBORDEMENT (F.), from *déborder* (*de* and *border*), 'to overflow.' A popular term for one or more sudden and copious evacuations from the bowels. It is chiefly applied to bilious evacuations of this kind—*Débordement de Bile*.

DÉBRIDER (F.), *Frangere solutio*, from (F.) *débrider* (*de* and *brider*), 'to unbridle.' The removal of filaments, &c., in a wound or abscess, which prevent the discharge of pus. In a more general acceptation, it means the cutting of a soft, membranous, or aponeurotic part, which interferes with the exercise of any organ whatever; thus, in paraphimosis, *débridement* of the prepuce is practised to put an end to the inflammation of the glans: in strangulated hernia, *débridement* of the abdominal ring is had recourse to, to remove the stricture of the intestine, &c.

DEC'AGRAMME, from *deka*, 'ten,' and *γραμμα*, 'a gramme.' The weight of ten grammes, 154.34 grains Troy.

DECAM'YRON, from *deka*, 'ten,' and *μυρον*, 'ointment.' An ancient cataplasm, composed of *malabathrum*, *mastic*, *euphorbium*, *spikenard*, *styrax calgmita*, carbonate of lime, *common pepper*, *unguentum nardi*, *opobalsamum*, and *uax*.

DECANTATION, *Decanta'tio*, *Defusio*, *Mo-*

tangien's, *Catach'ysis*. A pharmaceutical operation, which consists in pouring off, gently, by inclining the vessel, any fluid which has left a deposit.

DECAPITATIO ARTICULORUM, see Resection.

DECAPITATION, Decollation.

DECARBONIZATION, Hæmatosis.

DECEASE, Death.

DECESSIO, Epiesima.

DÉCHARNÉ, Demusculatus.

DÉCHAPELLEMENT. An old French term, employed by Paré and others for the operation of cutting and removing, with a strong forceps, the corona of a carious tooth, where it is desired to preserve the fang.

DÉCHAUSSEMENT (F.), (*de* and *chausser*.) The state in which the gums have fallen away from the teeth, as in those affected by mercury, in old persons, &c. Also, the operation of lancing the gums. See Gum lancet.

DÉCHAUSOIR, Gum lancet.

DÉCHIREMENT, Laceration.

DÉCHIRURE, Wound, lacerated—*d. de l'Uterus*, Uterus, rupture of the.

DECIDENTIA, Cataptonsis, Epilepsy.

DECIDUA, Decidua membrana.

DECID'UA MEMBRA'NA, *Decid'ua*, *Decid'uons Membrana*, from *decidere*, 'to fall off,' (*de* and *cadere*.) So called on account of its being considered to be discharged from the uterus at parturition. A membrane, formerly defined to be the outermost membrane of the fœtus in utero; and still so defined by some—as by Dr. Lee. Chaussier calls it *Epicho'rion*; by others, it has been called *Membra'na cadu'ca Hunteri*, *M. flocculenta seu cellulosa seu sinuosa seu communis seu præcis' tens*, *Decidua externa seu spongio'sa*, *Tu'nica exterior ovi seu cadu'ca seu crassa*, *Membra'na cribrosa seu ovi mater'na seu mucosa seu uteri interna evoluta*, *Epio'nê*, *Placenta uteri succenturiatus*, *Subplacenta*, *Nidamentum*, *Anhistous membrane*, (F.) *Caducue*, *C. vraie*, *Membrane caducue*, *Épione*, *Périone*;—prior to the time of the Hunters, called *Chorion spongio'sum seu tomento'sum seu fungo'sum seu reticulatum*, &c., *Tu'nica filamento'sa*, *Shaggy Chorion*, &c. Great diversity has prevailed regarding this membrane and its reflected portion. It exists before the germ arrives in the uterus—as it has been met with in tubal and ovarian pregnancies; and is occasioned by a new action, assumed by the uterine vessels at the moment of conception. Chaussier, Lobstein, Gardien, Velpéau, and others, consider it to be a sac, without apertures, completely lining the uterus, and that when the ovum descends through the tube, it pushes the decidua before it, and becomes enveloped in it, except at the part destined to form the placenta. That portion of the membrane which covers the ovum, forms the *Membra'na decid'ua reflex'a*, (F.) *Caducue réfléchie*, *Membrane caducue réfléchie*, according to them—the part lining the uterus being the *Decidua Uteri*, (F.) *Caducue utérine*. Towards the end of the fourth month, the decidua reflexa disappears. The very existence of a *Tunica decidua reflexa* has, however, been denied. This last membrane has received various names. Dr. Granville, regarding it as the external membrane of the ovum, has termed it *Cortex ovi*. It has also been termed *Involuerum membrana'ceum*, *Membra'na retiformis cho'rii seu filamento'sa seu adventitia seu crassa*, *Ovuline* and *Decid'ua protrusa*. To the membrane which, according to Bojanus and others, is situate between the placenta and the uterus, and which he considers to be produced

at a later period than the decidua vera, he gave the name *membra'na decid'ua serotina*.

Histological researches seem to show, that the decidua is an altered condition of the lining membrane of the uterus, with a whitish secretion filling the uterine tubular glands. The decidua reflexa is probably formed by the agency of nucleated cells from the plastic materials thrown out from the decidua uteri; in the same manner as the chorion is formed in the Fallopian Tube from plastic materials thrown out from its lining membrane. That the decidua reflexa is not a mere inverted portion of the decidua uteri is shown by the fact, that the texture of the two is by no means identical.

DECIDUA PROTRUSA, Decidua reflexa.

DECIDUOUS MEMBRANE, Decidua (membrana).

DECIGRAMME, *Decigram'ma*, from *decimus*, 'the tenth part,' and *γραμμα*, 'gramme.' The tenth part of the gramme in weight; equal to a little less than two grains, French; 1.543 Troy.

DECIMA'NA FEBRIS, from *decem*, 'ten.' An intermittent, whose paroxysms return every 10th day or every 9 days. It is supposititious.

DECLAMATION, *Declama'tio*, from *de*, and *clamare*, *clamatum*, 'to cry out.' The art of depicting the sentiments by inflections of the voice, accompanied with gestures, which render the meaning of the speaker more evident, and infuse into the minds of the auditors the emotions with which he is impressed. Declamation may become the cause of disease: the modification, produced in the pulmonary circulation,—accompanied by the great excitement, sometimes experienced,—is the cause of many morbid affections; particularly of pneumonia, hæmoptysis, and apoplexy. In moderation, it gives a healthy excitement to the frame.

DÉCLIN, Decline.

DECLINATIO, Decline.

DECLINE, *Declina'tio*, *Inclina'tio*, *Decrementum*, *Remis'sio*, *Parac'mê*, *Parac'masis*, (F.) *Déclin*, from *de*, and *clinare*, 'to bend.' That period of a disorder or paroxysm, at which the symptoms begin to abate in violence. We speak, also, of the decline of life, or of the powers, (F.) *Déclin de l'âge*, *L'âge de déclin*, when the physical and moral faculties lose a little of their activity and energy. See Phthisis, and Tabes.

DÉCLIVE, (F.) *Declivis*, from *de*, 'from,' and *clinus*, 'acclivity.' Inclining downwards. This epithet is applied to the most depending part of a tumour or abscess.

DÉCOCTÉ, Decoction.

DECOCTION, *Decoctio*, from *decoquere*, 'to boil,' (*de* and *coquere*, *coctum*.) *Eps'e'sis*, *Apoz'esia*, *Zesis*, *Hepse'sis*, *Aphepsis*, (F.) *Décoction*. The operation of boiling certain ingredients in a fluid, for the purpose of extracting the parts soluble at that temperature. Decoction, likewise, means the product of this operation, to which the terms *Decoction*, *Zema*, *Aphepse'ma*, *Ap'ozem*, *Apoz'ema*, *Hepse'ma*, *Chylus* and *Eps'e'ma*, (F.) *Décocté*, *Hydrolé*, *Hydrolite*, have been applied according to ancient custom, in order to avoid any confusion between the operation and its product;—as *præparatio* is used for the act of preparing; *præparatum*, for the thing prepared.

DECOCTION OF ALOES, COMPOUND, *Decoctionum aloes compositum*—d. of Bark, *Decoctionum Cinchonæ*—d. of Bark, pale, see *Decoctionum Cinchonæ*—d. of Bark, red, see *Decoctionum Cinchonæ*—d. of Bark, yellow, see *Decoctionum Cinchonæ*—d. of Barley, *Decoctionum hordei*—d. of Barley, compound, *Decoctionum hordei compositum*—d. Bitter, *Decoctionum amarum*—d. of Broom, *Decoctionum scoparii*—d. of Broom, compound, *Decoctionum sco-*

parii compositum—d. of Cabbage tree bark, Decoction geoffrae inermis—d. of Cassia, Decoction cassiae—d. of Chamomile, Decoction anthemidis nobilis—d. of Cinchona, Decoction Cinchonæ—d. of Cinchona, compound laxative, Decoction kinæ kinæ compositum et laxans—d. of Colomba, compound, Decoction Colombar compositum—d. of Dandelion, Decoction Taraxaci—d. of Dogwood, Decoction Cornûs Floridæ—d. of Elm bark, Decoction ulmi—d. of Flaxseed, compound, Decoction Lini compositum—d. of Foxglove, Decoction digitalis—d. de Gayac composée, &c., Decoction de Gayaco compositum—d. of Guaiacum, compound, Decoction Guaiaci compositum—d. of Guaiacum, compound purgative, Decoction de Guayaco compositum—d. of Hartshorn, burnt, Mistura cornu uesti—d. of Hellebore, white, Decoction veratri—d. of Iceland moss, Decoction cetrariæ—d. of Logwood, Decoction hamatoxyli—d. of Marshmallows, Decoction althææ—d. de Mauve composée, Decoction malvæ compositum—d. of Oak bark, Decoction quercûs—d. d'Orge, Decoction hordei—d. d'Orge composée, Decoction hordei compositum—d. of Pereira Brava, Decoction Pareiræ—d. of Pipsissewa, Decoction chinaphilæ—d. of Poppy, Decoction papaveris—d. of Quince seeds, Decoction cydoniæ—d. de Quinquina composée et laxative, Decoction kinæ kinæ compositum et laxans—d. de Salsepareille composée, Decoction sarsaparillæ compositum—d. of Sarsaparilla, Decoction sarsaparillæ—d. of Sarsaparilla, compound, Decoction sarsaparillæ compositum—d. of Sarsaparilla, false, Decoction araliæ nudicaulis—d. of Squill, Decoction sollæ—d. of Tormentil, Decoction Tormentillæ—d. of Uva ursi, Decoction uvæ ursi—d. of the Woods, Decoction Guaiaci compositum—d. of Woody nightshade, Decoction dulcamaræ—d. of Zittmann, Decoction Zittmanni.

DECOCTUM, Decoction—d. Album, Mistura cornu uesti.

DECOCTUM AL/OËS COMPOSITUM, Decoction Aloës, Balsam of Life, Compound Decoction of Aloës. (Ext. glyc. ʒvij, potass. carb. ʒi, aloës, myrrhæ contrit.: croci aa ʒiss, aquæ Oiss (Imp. meas.). Boil to Oj: strain, and add tinct. card. c. ʒvij. Pharm. L.) The gum and extractive are dissolved in this preparation. The alkali is added to take up a little of the resin. The tincture prevents it from spoiling. It is gently cathartic. Dose, fʒss to ʒij.

DECOCTUM ALTHÆÆ, Mistura Althææ, D. althææ officinalis: Decoction or Mixture of Marsh-mallows. (F.) Décoction de Guimauve. (Rad. althææ sic. ʒiv. uvar. passur. ʒij, aquæ. Ovj. Boil to Ov. Pour off the clear liquor. Pharm. E.) It is used as a demulcent.

DECOCTUM AMARUM; Bitter Decoction. (Rad. gent. ʒi, aquæ Ojss. Boil for a quarter of an hour; add species amara ʒij. Infuse for two hours and filter, without expressing. Pharm. P.) It is tonic. Dose, fʒss to ʒij.

DECOCTUM AMYLI, Mucilage amyli.

DECOCTUM ANTHEMIDIS NOB'ILIS, D. Chamæmelis: Decoction of Chamomile. (Flor. anthemid. nobil. ʒij, sem. carui ʒiv, aquæ Ov. Boil for fifteen minutes and strain. Pharm. E.) It contains bitter extractive and essential oil, dissolved in water. It is used, occasionally, as a vehicle for tonic powders, pills, &c., and in fomentations and gylsters; but for the last purpose, warm water is equally efficacious. The Dublin college has a compound decoction.

DECOCTUM ARA'LIE NUDICAULIS, Decoction of False Sarsaparilla. (Araliæ nudicaul. ʒvij, aquæ Oviij. Digest for four hours, and then boil to four pints: press out and strain the decoction. Former Ph. U. S.) It is used as a stomachic,

but it is an unnecessary and laborious preparation.

DECOCTUM CAS'SIÆ; Decoction of Cassia. (Cassia pulp. ʒij, aquæ Oj. Boil for a few minutes, filter without expression, and add syrup. violar. ʒij, or manna pur. ʒij. Pharm. P.) It is laxative, in the dose of ʒvj.

DECOCTUM CETRA'RIÆ, Decoction Liche'nis Island'ici, Decoction of Iceland Moss. (Cetrar. ʒss, aquæ Oiss. Boil to a pint, and strain with compression. Ph. U. S.) It consists of bitter extractive and fœcula, dissolved in water, and its operation is tonic and demulcent. Dose, fʒj to ʒiv. It is also nutrient, but hardly worthy of the rank of an official preparation.

DECOCTUM CHAMÆMELI, Decoction anthemidis nobilis.

DECOCTUM CHIMAPH'ILE, Decoction of Pipsissewa. (Chimaph. contus. ʒij, aquæ Oiss. Boil to a pint, and strain. Pharm. U. S.) A pint may be given in the course of the twenty-four hours.

DECOCTUM CINCHO'NÆ, D. Corticis Cinchonæ seu Kinæ Kinæ, Decoction of Cinchona or of Bark. (Cinchon. cort. contus. ʒj, aquæ Oj. Boil for ten minutes, in a slightly-covered vessel, and strain while hot. It contains quinia and resinous extractive, dissolved in water. Long coction oxygenates and precipitates the extractive. It can be given, where the powder does not sit easy, &c. Dose, fʒj to ʒiv.

The Pharmacopœia of the United States has a Decoction Cinchonæ flavæ, Decoction of yellow bark, and a Decoction Cinchonæ rubræ, Decoction of red bark, both of which are prepared as above. The London, Edinburgh, and Dublin Colleges have a Decoction Cinchonæ pallidæ, Decoction of Pale Bark.

DECOCTUM COLOM'BÆ COMPOSITUM; Compound Decoction of Colomba. (Colomb. contus., quassia, aa ʒij, cort. avant. ʒi, rhej pulv. ʒi, potassæ carbonat. ʒss, aquæ ʒxx. Boil to a pint, and add tinct. lavand. fʒss. Former Ph. U. S.) Given as a tonic, but not worthy an official station.

DECOCTUM COMMUNE PRO CLYSTERE, D. malvæ compositum—d. Cornu cervini, Mistura cornu uesti.

DECOCTUM CORNÛS FLORIDÆ, Decoction of Dogwood. (Cornûs Florid. cont. ʒj, aquæ Oj. Boil for ten minutes in a covered vessel, and strain while hot. Pharm. U. S.) Dose, fʒss.

DECOCTUM CYDO'NLE, Mucilago Sem'inti Cydonii Mali seu Sem'inum Cydoniorum; Decoction or Mucilage of Quince seeds. (Cydoniæ sem. ʒij, aquæ Oj. Boil for ten minutes over a gentle fire, and strain. Pharm. L.) It is merely a solution of mucilage in water, and is used as a demulcent.

DECOCTUM DAPHNES MEZE'REI, Decoction Mezere'i, Decoction of Meze'reon. (Meze'rei, ʒij, glycyrrh. cont. ʒi, aquæ Oij. Boil over a gentle fire to Oiss, and strain. Ph. E.) The acrimony of the meze'reon and the saccharine mucilage of the liquorice root are imparted to the water. It is somewhat stimulant, and has been used in secondary syphilis; but is devoid of power. Dose, fʒij to ʒvj.

DECOCTUM DIAPHORETICUM, D. Guaiaci compositum—d. pro Enemate, D. malvæ compositum.

DECOCTUM DIGITA'LIS, Decoction of Fox-glove. (Fol. digit. sic. ʒi, aquæ q. s. ut colentur fʒviij. Let the liquor begin to boil over a slow fire, and then remove it. Digest for fifteen minutes, and strain. Pharm. D.) It possesses the properties of the plant. Dose, fʒij to ʒij.

DECOCTUM DULCAMARÆ, Decoction of Woody Nightshade. (Dulcamaræ cont. ʒi, aquæ Oiss. Boil to Oj, and strain. Pharm. U. S.) This decoction has been considered diuretic and diapho-

retic, and has been administered extensively in skin diseases. It is probably devoid of efficacy.

DECOCTUM PRO FOMENTO, D. papaveris.

DECOCTUM GEOFFRÆ INERMIS; *Decoction of Calbage-Tree Bark.* (Cort. geoffr. inermis, in pulv. ʒi, Aquæ Oij. Boil over a slow fire to a pint, and strain. Pharm. E.) It is possessed of anthelmintic, purgative, and narcotic properties, and has been chiefly used for the first of these purposes. Dose, to children, fʒij — to adults, fʒss to ʒij.

DECOCTUM GUAIACI COMPOSITUM, D. Guaiaci officinalis compositum seu Ligno'rum seu de Guyaco compositum seu sudorificum seu diaphoreticum; Compound decoction of Guaiacum, *Decoction of the Woods.* (Lign. guaiac. rasur. ʒiij, fruct. sicc. vitis vinifer. ʒij, rad. lauri. sassafr. concis., rad. glycyrrh. aa ʒj, aquæ Ox. Boil the Guaiacum and raisins over a slow fire to Oj: adding the roots towards the end, then strain. Pharm. E.) It is possessed of stimulant properties, and has been given in syphilitic, cutaneous, and rheumatic affections. The resin of the guaiacum is, however, insoluble in water, so that the guaiac wood in it cannot be supposed to possess much, if any, effect. The Parisian Codex has a

DECOCTUM DE GUYACO COMPOSITUM ET PURGANS, (F.) *Décotion de Gayac Composée et Purgative*; Compound purgative Decoction of Guaiacum. (Lign. guaiac. rasp., rad. sarsap. aa ʒj, potass. carbonat. gr. xxv. Macerate for twelve hours, agitating occasionally, in water Oiv, until there remain Oij. Then infuse in it fol. sennæ ʒij, rheæ ʒj, lign. sassafr., glycyrrh. rad. aa ʒij, sem. coriand. ʒj. Strain gently, suffer it to settle, and pour off the clear supernatant liquor. The title sufficiently indicates the properties of the composition.

DECOCTUM HÆMATOXYLLI, *Decoction of Log-wood.* (Hæmatoxyl. rasur. ʒj, aquæ Oij. Boil to a pint, and strain. Pharm. U. S.)

DECOCTUM HELEBORI ALBI, D. veratri.

DECOCTUM HORDEI, *Decoction Hordei Distichi, Ptis'ana Hippocrati'ca, Tipts'aria, Tapas'aria, Aqua Horde'ata, Hydrocrith's, Barley Water, Decoction of Barley.* (F.) *Décotion d'Orge, Tisanæ Commune.* (Hord. ʒij, Aquæ Oivss. Wash the barley well, boil for a few minutes in water Oss: strain this, and throw it away, and add the remainder, boiling. Boil to Oij, and strain. Pharm. L.) It is nutritive and demulcent, and is chiefly used as a common drink, and in glysters.

DECOCTUM HORDEI COMPOSITUM, *Decoction pectora'le, Mistu'ra Hor'dei, Ptis'ana communis*; Compound Decoction of Barley, (F.) *Décotion d'Orge composée.* (Decoct. hord. Oij, carica fruct. concis. ʒiiss, glycyrrh. rad. concis. et contus. ʒvj, urarum pass. deuptis acinis. ʒijss, aquæ Oij. Boil to Oij, and strain. Pharm. L.) It has similar properties to the last.

DECOCTUM KINÆ KINÆ, D. Cinchonæ.

DECOCTUM KINÆ KINÆ COMPOSITUM ET LAXANS, *Décotion de quinquina composée et laxative*; Compound laxative decoction of Cinchona. (Cort. cinchon. ʒj, aquæ Oij. Boil for a quarter of an hour, remove it from the fire; then infuse in it, for half an hour, fol. sennæ, sodæ sulph. aa ʒij: add syrup. de senad fʒj.) Its title indicates its properties.

DECOCTUM LICHENIS ISLANDICI, *Decoction Cetrariæ*—d. Lignorum, D. Guaiaci compositum.

DECOCTUM LINI COMPOSITUM, *Compound Decoction of Flaxseed.* (Lini ʒj, Glycyrrhiz. contus. ʒss, aquæ Oiss. Boil for ten minutes, and strain. Ph. D.)

DECOCTUM MALVÆ COMPOSITUM, *Decoction pro enem'atē seu commu'nē pro clyste'rē*; Com-

pound Decoction of Mallow, (F.) *Décotion de Mauve composée.* (Malvæ exsicc. ʒj, anthemid. flor. exsicc. ʒss, aquæ Oj. Boil for fifteen minutes and strain.) It consists of bitter extractive, and mucilage in water, and is chiefly used for clysters and fomentations. It is unworthy a place in the pharmacopœias.

DECOCTUM MEZEREI, D. daphnes mezerei.

DECOCTUM PAPAV'ERIS, *Decoction pro Fomen'to, Fotus communis*; *Decoction of Poppy.* (Papav. sonnif. capsul. concis. ʒiv, aquæ Oiv. Boil for fifteen minutes, and strain. Pharm. L.) It contains the narcotic principle of the poppy, and mucilage in water: is anodyne and emollient, and employed as such in fomentation, in painful swellings, ulcers, &c.

DECOCTUM PAREIRÆ, *Decoction of Pareira Brava.* (Pareir. brav. ʒx, Aq. destillat Oiss, (Imp. meas.) Boil to a pint and strain. Dose, fʒj to ʒij.

DECOCTUM PECTORALE, D. hordei compositum —d. Polygalæ Senegæ, D. Senegæ.

DECOCTUM QUERCUS ALBÆ, D. Quercus Roboris; *Decoction of White Oak Bark.* (Quercus cort. ʒj, aquæ Oiss. Boil to a pint, and strain. Pharm. U. S.) It is astringent, and used as such, in injections, in leucorrhœa, uterine hemorrhage, &c., as well as in the form of fomentation to unhealthy ulcers.

DECOCTUM SARSÆ, *Decoction Sarsaparillæ.*

DECOCTUM SARSAPARILLÆ, D. Smilacis Sarsaparilla, D. Sarsæ, D. Sarsæ; *Decoction of Sarsaparilla.* (Sarsaparill. rad. concis. ʒv, aq. fervent. Oiv. (Imp. meas.) Macerate for four hours near the fire, in a lightly-covered vessel; then bruise the root; macerate again for two hours; then boil to Oij, and strain. Pharm. L.) It contains bitter extractive, and mucilage in water; is demulcent, and has been used, although it is doubtful with what efficacy, in the sequelæ of syphilis. Dose, fʒiv to Oss.

DECOCTUM SARSAPARILLÆ COMPOSITUM; Compound Decoction of Sarsaparilla, (F.) *Décotion de Salsepareille Composée.* (Sarsaparill. concis. et contus. ʒvj, Sassafr. Cort. concis., Lign. Guaiac. rasur., Rad. Glycyrrhiz. contus., aa ʒj, Mezerei, concis. ʒij, Aquæ Oiv. Boil fifteen minutes, and strain. Pharm. U. S.) This is considered to be possessed of analogous properties to the celebrated Lisbon Diet-Drink, *Decoctum Lusitanicum*, which it resembles in composition.

DECOCTUM SARZÆ, *Decoction Sarsaparillæ.*

DECOCTUM SCILLÆ; *Decoction of Squill.* (Scillæ, ʒij, juniper. ʒiv, Senegæ, ʒij, Aquæ Oiv. Boil till one half the liquor is consumed; strain, and add spirit of nitrous ether, fʒiv. Former Pharm. U. S.) Dose, fʒss.

DECOCTUM SCOPARIÏ, *Decoction of Broom* (Scopar. exsiccate ʒss, (avoir'dupois); Aquæ Oss. (Imp. meas.) Boil for ten minutes and strain. —Ph. D.)

DECOCTUM SCOPARIÏ COMPOSITUM, D. Scopariï (Ph. E.), Compound Decoction of Broom (Scopariï, Junip. cont., Tarazac. cont., aa ʒss; Aq. destillat. Oiss, (Imp. meas.) Boil to a pint, and strain. Ph. L.) Dose, half a pint to a pint, in the day as a diuretic, in dropsy.

DECOCTUM SENEGÆ, D. Polygalæ Senegæ; *Decoction of Senega.* (Seneg. cont. ʒj, aquæ Oiss. Boil to Oj, and strain. Ph. U. S.) It is reputed to be diuretic, purgative, and stimulant, and has been given in dropsy, rheumatism, &c.

DECOCTUM SMILACIS SARSAPARILLÆ, D. sarsaparillæ—d. Sudorificum, D. Guaiaci compositum.

DECOCTUM TARAX'ACI, *Decoction of Dandelion.* (Tarazac. contus. ʒiv, aquæ Oiss. (Imp. meas.) Boil to a pint, and strain. Ph. L.) Dose fʒiiss.

DECOCTUM TORMENTILLÆ, *Decoction of Tormentil.* (Tormentil. cont. ʒij, Aquæ destillat. Oiss.

(Imp. meas.) Boil to a pint, and strain. Dose, as an astringent, one or two fluid ounces.

DECOCTUM ULMI, *D. Ulmi Campestris*, *Decoction of Elm Bark.* (*Ulm* cort. recent. cont. $\frac{3}{4}$ iss, *aqvæ Oij.* (Imp. meas.) Boil to Oj, and strain. *Ph. L.*) It is a reputed diuretic, and has been used in lepra and herpes; but, probably, has no efficacy. Dose, f $\frac{3}{4}$ iv to Oss.

DECOCTUM UVE URSI, *Decoction of uva ursi.* (*Uva ursi*, $\frac{3}{4}$ j, *aqvæ*, f $\frac{3}{4}$ xx. Boil to a pint and strain. *Ph. U. S.*) Dose, f $\frac{3}{4}$ iss.

DECOCTUM VERA'TRI, *Decoction Helleb'ori albi*; *Decoction of White Hellebore.* (*Veratri rad.* cont. $\frac{3}{4}$ j, *aqvæ Oij.* *spir. rec.* f $\frac{3}{4}$ ij. Boil the water decoction to Oj, and, when it is cold, add the spirit. *Pharm. L.*) It is stimulant, acrid, and cathartic; but is used only externally. It is a useful wash in *tinea capitis*, *psora*, &c. Should it excite intense pain on being applied, it must be diluted.

DECOCTUM ZITTMANNI, *Zittmann's Decoction.* A most absurd farrago, extolled by Theden in venereal diseases, the formula for which, according to Jourdan, "some blockheads have lately reproduced among us as a novelty." It is composed of *sarsaparilla*, *pulvis stypticus*, *calomel* and *cinnabar*, boiled in water with aniseed, fennel seed, and liquorice root. A stronger and a weaker decoction were directed by Zittmann. Formulae for its preparation are contained in Jourdan's *Pharmacopœia Universalis*; and in Linck's *Vollständiges Recept-Taschenbuch.* Leipzig, 1841.

DECOLLATION, *Decapitation*, *Decapitatio*, *Obtruncatio*, from *de* and *collum*, 'the neck.' The act of separating the head from the trunk, by the section of the neck, as in cases of dystocia.

DÉCOLLEMENT, (F.) *Deglutina'tio*, (from *de* and *coller*, 'to glue.') The state of an organ that is separated from the surrounding parts, owing to destruction of the areolar membrane which united them. The skin is *décollée*, i. e. separated from the subjacent parts, by a burn, subcutaneous abscess, &c.

DECOLEMENT DU PLACENTA is the separation or detachment of the whole or a part of the placenta from the inner surface of the uterus. *Décollement*, from *de*, 'from,' and *collum*, 'the neck,' *obtruncatio*, also means the separation of the head of the fetus from the trunk, the latter remaining in the uterus.

DECOLORATION, *Decolora'tio*, *Discolora'tio*, from *de*, 'from,' and *colorare*, *coloratum*, 'to colour.' Loss of the natural colour; *Parachro'sis*. Devoid of colour—*Achroma'ia*.

In Pharmacy, any process by which liquids, or solids in solution, are deprived wholly or in part of their colour,—as by bringing them in contact with animal charcoal.

DÉCOMPOSÉE (F.), from *de*, 'from,' and *componere*, *compositum*, 'to compose.' *Decomposed*, *Dissolutus*. An epithet, applied to the face when extensively changed in its expression and colour; as in the choleric or moribund.

DECORTICA'TION, from *de*, 'from,' and *cortex*, *corticis*, 'bark.' An operation, which consists in separating the bark from roots, stalks, &c.

DECOS'TIS, *Apleu'ros*, from *de*, *priv.*, and *costa*, 'a rib.' One who has no ribs.

DECREMENTUM, Decline.

DECREPIT, see *Decrepitude*.

DECREPITUDE, *Decrepitu'do*, *Ætas decrep'ita*, *Ul'tima senectus*, *Senect'a decrep'ita* seu *extre'ma seu summa seu ul'tima*, from *decrepitis*, (*de*, and *crepere*, *crepitum*, 'to creak,' 'very old.' The last period of old age, and of human life; which ordinarily occurs about the eightieth year. It may, however, be accelerated or protracted.

Its character consists in the progressive series of phenomena which announce the approaching extinction of life. One in this stage of existence is said to be *decrep'it*, *decrep'itus*, (F.) *décrépit*.

DECRETORII DIES, Critical days.

DECURBITUS, from *decumbere*, (*de*, and *cubere*, *cubitum*, 'to lie,') 'to lie down.' *Cuba'tio*, Lying down. Assuming the horizontal posture, *cu'bitus sup'inus*; (F.) *Coucher*. *Horâ decubitis*, 'at bed time.' The French say—*Decubitus horizontal*, *sur le dos*, *sur le côté*, for, lying in the horizontal posture, on the back or side.

DECURTA'TUS, *μειωπος*, or *μειωπισ'ων*, from *de*, and *curtus*, 'short,' 'shortened, curtailed,' 'running to a point.' When applied to the pulse, it signifies a progressive diminution in the strength of the arterial pulsations, which, at last, cease. If the pulsations return and gradually acquire all their strength, it is called *Pulsus decurta'tus recip'ocus*.—Galen.

DECUSSATIO, *Decussation*—d. Pyramidum, see *Corpora pyramidalia*.

DECUSSA'TION, *Decussa'tio*, *Chias'mos*, *Incrucia'tio*, *Intersec'tio*, *Intracuta'ra*, from *decussis*, that is, *decem asses*; also, the figure of the letter X. Union in the shape of an X or cross. Anatomists use this term chiefly in the case of the nerves—as the *decussation of the optic nerves*, which cross each other within the cranium.

DECUSSATION OF THE PYRAMIDS, see *Corpora pyramidalia*.

DECUSSO'RIMUM, from *decutio*, *decussum*, (*de*, and *quatio*,) 'I shake down,' (F.) *Décuatoire*. An instrument used by the ancients, for depressing the dura mater, and facilitating the exit of substances effused on or under that membrane. It is described by Scultetus, Paré, &c. See *Meningophylax*.

DÉDAIGNEUX, *Rectus superior oculi*.

DEDE, Death.

DEDENTITION, see *Dentition*.

DEDOLA'TION, *Dedola'tio*, from *dedolare*, (*de*, and *dolare*, *dolatum*,) 'to cut and hew with an axe.' This word has been used by surgeons to express the action by which a cutting instrument divides obliquely any part of the body, and produces a wound with loss of substance. It is commonly on the head, that wounds by dedolation are observed. When there was a complete separation of a portion of the bone of the cranium, the ancients called it *Apocepharnis'mus*.

DEERBALLS, *Elaphomyces granulatus*.

DEERBERRY, *Gaultheria*, *Vaccinium stamineum*.

DEERFOOD, *Brasenia hydropeltis*.

DEERS' TEARS, Bezoar of the Deer.

DÉFAILLANCE, Syncope.

DEFECATION, *Defæca'tio*, from *de*, and *fæx*, *Pl. fæces*, 'excrement.' The act by which the excrement is extruded from the body. *Caca'tio*, *Ecre'tio alvi* seu *alv'ina* seu *fæcum alv'inarum*, *Dejectio alvi*, *Sece'sio*, *Expul'sio* seu *Ejectio fæcum*, *Apago'gê*, *Hypochore'sis*, *Echore'sis*, *Ecopro'sis*, *Eges'tio*. The fæces generally accumulate in the colon, being prevented by the annulus at the top of the rectum from descending freely into that intestine. In producing evacuations, therefore, in obstinate constipation, it is well, by means of a long tube, to throw the injection into the colon.

In Pharmacy, defecation means the separation of any substance from a liquid in which it may be suspended. See *Clarification*.

DEFECTIO ANIMI, Syncope.

DEFECTUS LOQUELÆ, *Aphonia*—d. *Veneris*. *Anaphrodisia*.

DEFENSIVUM, from *defendere*, *defensum*, 'to defend,' (F.) *Défensif*. A preservative or defence. The old surgeons gave this name to dif-

ferent local applications, made to diseased parts, for the purpose of guarding them from the impression of extraneous bodies, and particularly from the contact of air.

DEFERENS, from *defero*, (*de*, and *ferre*), 'I bear away,' 'I transport.'

DEFERENS, VAS, *Ductus seu Canalis deferens*, *Fibrator*, (F.) *Conduit spermaticus*, *Canal seu Conduit deferens*, is the excretory canal of the sperm, which arises from the epididymis, describes numerous convolutions, and with the vessels and nerves of the testicle concurs in the formation of the spermatic chord, enters the abdominal ring, and terminates in the ejaculatory duct.

DEFIBRINATED, Desfibrinized.

DEFIBRINATION, see Desfibrinized.

DEFIBRINIZED, (F.) *Désfibriné*, *Desfibrinated*. From *de*, and *fibrin*. Deprived of fibrin. A term applied to blood from which the fibrin has been removed, as by whipping. The act of removing fibrin from the blood has been termed *defibrination*.

DEFIGURATIO, Deformation.

DEFLAGRATIO, *Deflagratio*, from *deflagrare*, (*de*, and *flagrare*, *flagratum*, 'to burn,') 'to set on fire.' In pharmacy, the rapid combustion of a substance with flame; great elevation of temperature; violent motion, and more or less noise. Thus, we speak of the deflagration of nitrate, and of chlorate, of potassa, of gunpowder, &c.

DEFLECTENS, Derivative.

DEFLECTIO, Derivation.

DEFLOURATION, Stuprum.

DEFLOURATION, *Depucella*, *Depudication*, *Devirginatio*, *Defloratio*, *Devirginatio*, *Virginis vitatio*, *Virginitas deflorata*, from *deflorescere*, (*de*, and *florescere*), 'to shed flowers.' The act of depriving a female of her virginity. Inspection of the parts is the chief criterion of defloration having been forcibly accomplished; yet inquiry must be made, whether the injury may not have been caused by another body than the male organ. Recent defloration is infinitely more easy of detection than where some time has elapsed.

DEFLUVIUM, Aporrhœa—d. Capillorum, Alopecia—d. Pilorum, Alopecia.

DEFLUXIO, Catarrh, Defluxion, Diarrhœa—d. Catarrhalis, Influentia.

DEFLUXION, *Defluxio*, (*de*, and *fluere*, *fluxum*, 'to flow,') *Defluxus*, *Catarrhysis*, *Hyporrhysis*, *Catarrhus*. A falling down of humours from a superior to an inferior part. It is sometimes used synonymously with inflammation.

DEFLUXUS DYSENTERICUS, Dysentery.

DEFORMATIO, *Deformatio*, *Cacomorpha*, *Cacomorpha*, *Dysmorphia*, *Dysmorphia*, *Dysmorphia*, *Deformitas*, *Informitas*, *Defiguratio*, *Disfiguratio*, *Deformitas*, (F.) *Difformité*, from *de*, and *forma*. Morbid alteration in the form of some part of the body, as of the head, pelvis, spine, &c. A deformity may be natural or accidental.

DEFRUTUM, for *Deservitum*, (*de*, and *ferveo*, 'I boil,') (F.) *Vin cuit*. Grape must, boiled down to one half and used as a sweetmeat.—Plin. Columell. Isidor. See Rob.

DEFURFURATIO, see Desquamation.

DEFUSIO, Decantation.

DEGENERATIO, Degeneration—d. Adiposa Cordis, Steatosis cordis.

DEGENERATION, *Degeneratio*, *Nothenis*, *Nothia*, from *degener*, 'unlike one's ancestors,' (*de*, and *genus*, *generis*, 'family,') *Degen'eracy*, (F.) *Dégénération*, *Abatardissement*. A change for the worse—*degradation*—in the intimate composition of the solids or fluids of the body. In

pathological anatomy, *degeneration* means the change which occurs in the structure of an organ, when transformed into a matter essentially morbid; as a *cancerous*, or *tubercular*, degeneration.

Dégénérescence is, by the French pathologists, employed synonymously with Degeneration.

DÉGÉNÉRESCENCE, Degeneration—d. *Graisseuse du Foie*, Adiposis hepatica—d. *Graulée du Rein*, Kidney, Bright's disease of the—d. *Noire*, Melanosis.

DEGLUTINATIO, *Décollement*.

DEGLUTITIO, Deglutition—d. *Difficilis*, *Dysphagia*—d. *Impedita*, *Dysphagia*—d. *Læsa*, *Dysphagia*.

DEGLUTITION, *Deglutitio*, *Catap'osis*, from *de*, and *glutire*, *glutitum*, 'to swallow.' The act by which substances are passed from the mouth into the stomach, through the pharynx and œsophagus. It is one of a complicated character, and requires the aid of a considerable number of muscles; the first step being voluntary, the remainder executed under spinal and involuntary nervous influence.

DEGMUS, *Dexis*, *Morus*, (F.) *Morsure*,—a bite in general, *Anygos*, from *δακνυς*, 'I bite.' A gnawing sensation about the upper orifice of the stomach, which was once attributed to acrimony of the liquids contained in that viscous.

DÉGORGEMENT, Disorgement.

DÉGOUT, Disgust.

DEGRADATION, Degeneration.

Degree, from *dégré*, originally from *gradus*, 'a step.' A title conferred by a college,—as the 'degree of Doctor of Medicine.' Galen used this expression to indicate the qualities of certain drugs. Both he and his school admitted cold, warm, moist, and dry medicines, and four different 'degrees' of each of those qualities. Thus, *Opium* was warm in the first degree, *Agrimony* in the second, *Roche Alum* in the third, and *Garlic* in the fourth. *Bedegar* was cold in the first, the flower of the Pomegranate in the second, the *Sempervivum* in the third, *Opium* in the fourth, &c. The French use the term *dégré* to indicate, 1. The intensity of an affection: as a burn of the first, second, third degree, &c. 2. The particular stage of an incurable disease, as the third degree of phthisis, cancer of the stomach, &c.

DEGUSTATION, *Degustatio*, from *de*, and *gustare*, *gustatum*, 'to taste.' *Gustation*. The appreciation of sapid qualities by the gustatory organs.

DEHISCENCE, *Dehiscencia*, from *dehiscere* (*de*, and *hiscere* 'to gape'), 'to break open in chinks.' Separation of a shut organ along the line of union. A regular and determinate separation, which takes place in a shut organ.

DEICTICOS, Index.

DEID, Death.

DEIL'S SNUFFBOX, *Lycopodon*.

DEIRONCUS, *Bronchocele*.

DEJECTEDNESS, Depression.

DEJECTIO ALVI, Defecation—d. *Alvina*, Excrement.

DEJECTION, from *dejacere*, *dejectum* (*de*, and *jacere*), 'to cast down.' *Dejectio*, *Subductio*, *Hypogag'ge*, *Hypochore'sis*, *Hypochore'sis*, *Apopl'atus*, *Hypoph'atus*, *Hypoph'ora*. The expulsion of the fæces;—*Ejectio*, *Eges'tio*. Also, a faecal discharge or stool,—generally, however, with *alvina* prefixed,—as an *alvina Dejection*, *Alvus vir'idis*, *Dejectio alvi'na*. Also depression of spirits.

DEJECTIONES NIGRÆ, Melæna.

DEJECTORIUM, Cathartic.

DELACRYMATO, Epiphora.

DELACRYMATIVUS, Apodacryticus.

DELAPSIQ, Prolapsus.

DELAPSUS, Prolapsus—d. Palpebræ, Blepharoptosis.

DELATIO, Indication.

DÉLAYANTS, Diluentia.

DELCROIX'S DEPILATORY, see Depilatory, Colley's.

DELETERIOUS, *Deleterius*, *Perniciosus*, (F.) *Délétère*, *Pernicieux*, from *ἐλέω*, 'I injure.' That which produces destructive disorder in the exercise and harmony of the functions.

DELETERIUM, Poison.

DELIERET, Delirious.

DELIGATIO, Deligation, Ligature, see Bandage.

DELIGATION, *Deligatio*, *Epidei'sis*, *Deligatura*, *Vulnerum deligatio* seu *vinctura*, *Fascia*, *rum Applicatio*, *Plagæ*, *rum Vinctura*, *Fascia*, *rum*, from *deligare*, *deligatum*, (*de*, and *ligo*), 'to bind.' The deligation of wounds formerly embraced the application of apparatus, dressings, &c.,—the denomination *Deligator Plagæ*, *rum* being synonymous with *Medicus Vulnerarius*, and in derivation, with the Wundarzt, 'wound physician' or surgeon, of the Germans. Deligation is hardly ever used now as an English word. In France, it is applied to the regular and methodical application of bandages, and to the ligation of arteries.

DELIGATURA, Deligation.

DELIQUESCENT, *Deliquescent*, from *deliquescere*, (*de* and *liquescere*), 'to melt,' 'to dissolve.' Any salt which becomes liquid by attracting moisture from the air. The deliquescent salts require to be kept in bottles, well stopped. Chloride of lime, acetate of potassa, and carbonate of potassa, are examples of such salts. The ancient chemists expressed the condition of a body, which had become liquid in this manner, by the word *Deliquium*.

DELIQUIUM ANIMI, Syncope.

DELIRAMENTUM, Delirium.

DELIRANS, Delirious.

DÉLIRANT, Delirious.

DELIRATIO, Delirium—d. *Senum*, see Dementia.

DÉLIRE, Delirium—d. *Crapuleux*, Delirium tremens—d. *des Ivrognes*, Delirium tremens—d. *Nerveux*, Delirium nervosum.

DÉLIRE DES PERSÉCUTIONS, Mania, in which the morbid hallucination is, that the individual is persecuted.

DÉLIRE TRAUMATIQUE, Delirium nervosum—d. *Tremblant*, Delirium tremens.

DELIRIA, Insanity.

DELIRIE, Delirious.

DELIRIOUS. Same etymon as **DELIRIUM**. *Delirans*, *Delirius*, *Excerebra'tus*, *Alloch'oös*, *Ali'enus*, *Parale'rus*, *Raving*, (Sc.) *Delieret*, *Delirie*, (F.) *Délirant*. One who is in a state of delirium. That which is attended by delirium. The French use the term *Fièvre pernicieuse délirante* for a febrile intermittent, in which delirium is the predominant symptom.

DELIRIUM, from *de*, 'from,' and *lira*, 'a ridge between two furrows' [?]: more probably from *de*, and *ληρημα*, 'a silly saying or action.' *Parac'opë*, *Phreni'tis*, *Phledoni'a*, *Desipien'tia*, *Aphros'ia*, *Paralere'ma*, *Paralere'sis*, *Paralog'i'a*, *Phanta'sia*, *Paraphros'yne*, *Emo'tio*, *Leros*, *Parane'a*, *Alloph'asia*, *Delira'tio*, *Delirament'um*, (Sc.) *Ravery*, *Roving*, (F.) *Délire*, *Égarement d'esprit*, &c., *Transport*, *Ideosynchysie*. Straying from the rules of reason; wandering of the mind. Hippocrates used the word *mania*, for delirium sine febre, and the Greek words given above for delirium cum febre. In different authors, also, we find the words, *Paraph'ora*, *Paraph'otes*, *Paraphrene'sis*, *Paraphren'ia*, *Phrene'sis*, *Phrenet'asia*, &c., for different kinds of delirium. Delirium is usually symptomatic.

DELIRIUM, Delirium is usually symptomatic.

DELIRIUM EBRIOSITATIS, Drunken madness; also *D. tremens*—d. Epileptic, see Epilepsy—d. *Furiosum*, *Mania*—d. *Maniacum*, *Mania*.

DELIRIUM NERVOSUM, (F.) *Délire nerveux* ou *traumatique*. A form of delirium which supervenes after surgical operations, or severe injuries.

DELIRIUM POTATORUM, *D. tremens*—d. *Tremefaciens*, *D. tremens*—d. *Senile*, see Dementia.

DELIRIUM TREMENS, *D. potato'rum* seu *obri'osita'tis* seu *tremifa'ciens* seu *vig'ilians*, *Ma'nia à potu* seu *e temulentia*, *Enoma'nia*, *Erethis'mus obrioso'rum*, *Dipsoma'nia*, *Meningi'tissen* *Phreni'tis* *potato'rum*, *Tromoma'nia*, *Tromoparana'a*, *Potoparana'a*, *Pototromoparana'a*, (F.) *Encephalopathie crapuleuse*, *Délire tremblant* ou *crapuleux* ou *des Ivrognes*, *Folie des Ivrognes* ou *Alcoolique*. A state of delirium and agitation, peculiar to those addicted to spirituous liquors, with great sleeplessness. It is preceded by indisposition, lassitude, watchfulness, headache, and anorexia; the delirium and tremors, *Tremor potato'rum*, most commonly recurring in paroxysms. It is caused by the habitual and intemperate use of ardent spirits or of opium or tobacco; or rather by abandoning them after prolonged use. The treatment is various. Many have regarded stimulants, with large doses of opium to induce rest, to be indispensable. It is certain, however, that the expectant system will often, if not generally, be successful; and, it is probable, a cure effected in this manner will be more permanent than when produced by excitants.

DELIRUS, Delirious.

DÉLITESCENCE, see Repercussion.

DELITESCENTIA, see Repercussion.

DELIVER, (Prov.) *lay*, same etymon as the next. 'To put to bed.' To disburden of a child.

DÉLIVRANCE (F.), (*de*, and *liver*, 'to set free.') An expression, which, in common language, signifies the action of delivering, *libera'tio*, but in the practice of obstetrics, means the extrusion of the secundines, either spontaneously or by the efforts of art. This completion of delivery—*partus secunda'rius*, *secundina'rum expul'sio* seu *extra'ctio*, is produced by the same laws as the expulsion of the fœtus. Sometimes, the after-birth follows the child immediately; at others, it is retained; and requires manual interference to remove it. The following are the chief cases in which this becomes necessary. 1. Intimate adhesion between the placenta and paries of the uterus. 2. Spasmodic contraction of the orifice of the uterus. 3. Hour-glass contraction. 4. Torpor or hemorrhage after the expulsion of the child, and,—5. Insertion of the placenta at the orifice of the uterus.

DELIVERY, Parturition—d. *False*, *False water*.

DÉLIVRE, Secundines.

DELOCATIO, Luxatio.

DELPHINE, see Delphinium staphisagria.

DELPHININE, see Delphinium staphisagria.

DELPHINIUM, from *δελφίς*, or *δελφίν*, 'a dolphin,' which the flowers resemble. *D. consolida* seu *Seg'etum* seu *Versic'olor*, *Calcitra'pa*, *Consolida Regu'tis*, *Branching Larkspur*, *Stag'weeed*. Ord. Ranunculaceæ. Sez. Syst., *Polyandria Trigynia*. (F.) *Pied d'alo'nette des champs*. It has been employed as a vermifuge. The flowers have been used in ophthalmia, and the seeds have the same property as those of *Stavesacre*.

DELPHINIUM CONSOLIDA, Delphinium—d. *Segetum*, Delphinium.

DELPHINIUM STAPHISAGRIA. The systematic

name of the *Staves'acre*, *Staphisagria*, *Phtheiroc-*
torum, *Phtheirium*, *Staphis*, *Pedicularia*, (F.)
Staphisagria, *Herbe aux Poux*, *Dunphinelle*.
The seeds, *Staphisagria*, (Ph. L. and E.) have a
disagreeable smell; a nauseous, bitterish, hot
taste; and are cathartic, emetic, and vermifuge:
owing, however, to the violence of their opera-
tion, they are seldom given internally, and are
chiefly used in powder, mixed with hair powder,
to destroy lice. The active principle of this
plant has been separated, and received the name
Delphinia, (F.) *Delphine*, *Delphinine*. It is ex-
tremely acrid, and has been recently used, like
veratrum, in tic douloureux, paralysis and rheu-
matism. It is used in the form of ointment, or
in solution in alcohol, (gr. x to xxx, or more, to
℥i.) applied externally.

DELPHINIUM VESICOLOR, Delphinium.

DELPHYS, Uterus, Vulva.

DELTA, Vulva.

DELTIFORMIS, Deltoid.

DELTOID, *Deltoides*, *Delto'des*, *Deltoïdeus*,
Deltiformis, (F.) *Deltoides*, from the Greek cap-
ital letter Δ, *deltā*, and *adōs*, 'resemblance.' *Sous-*
acromio-clavi-huméral of Dumas; *Sus-acromio-*
huméral of Chaussier. A triangular muscle
forming the fleshy part of the shoulder, and
covering the shoulder-joint. It extends from the
outer third of the clavicle, from the acromion
and spine of the scapula, to near the middle and
outer part of the os humeri, where it is inserted
by means of a strong tendon. This muscle
raises the arm directly upwards, when the shoulder
is fixed, and carries it anteriorly or posteri-
orly, according to the direction of the fibres,
which are thrown into action. If the arm be
rendered immovable, the deltoid acts inversely
and depresses the shoulder.

DELTOÏDE, Deltoid.

DELUSION, Hallucination.

DEM EL MUCA. A name given by Prosper
Alpinus to a disease, which, he says, is proper to
Egypt. He considers it to be inflammation of the
brain; but others describe it as a pernicious in-
termittent.

DEMAGNETIZATION, *Demagnetisation*.
The act of removing the condition of magneti-
zation.

DÉMANGEAISON, Itching.

DEMANUS, Acheir.

DÉMENCE, Dementia—*d. Innée*, Idiostism.

DEMENTED, Insane.

DEMENTIA, *Amentia*, *Fatuitas*, *Ano'a*,
Ano'sia, *Dementia*, *Ecnia*, *Parano'a*, *Mo-*
ria demens, *Nuisthenia*, *Incoherence*, *Imbecil-*
ity, *Incoherent Insanity*; from *de*, 'from,' or
'out of,' and *mens*, *mentis*, 'mind,' or 'reason,'
(F.) *Démence*, *Bêtise*. In common parlance, and
even in legal language, this word is synonymous
with insanity. Physicians, however, have applied
it to those cases of unsound mind which are char-
acterized by a total loss of the faculty of thought,
or by such an imbecility of intellect that the
ideas are extremely incoherent, there being at
the same time a total loss of the power of rea-
soning. Mania and melancholy are apt to end
in this, if possible, more deplorable state. *Dotage*
is the *Dementia* of the aged—*Mo'ria demens tene-*
ma, *Lere'ma*, *Lere'sia*, *Lerus*, *Deliratio Senum*,
Amentia Senilis, *Puerilitas*, *Delirium Senile*,
Anil'itas, *Senile Insan'ity*, *Senile Dementia*, *In-*
sanity of the aged, (Sc.) *Dottrice*,—a form of moral
insanity, in which the whole moral character of
the individual is changed.

DEMENTIA ACCIDENTALIS, see Lunatic—*d. Ad-*
ventitia, see Lunatic—*d. Naturalis*, Idiostism—*d.*
Paralytica, Paralysis of the Insane.

DEMI, Semi—*d. Bain*, Semicupium—*d. Circu-*

laire, Semi-circular—*d. Épineux*, Semi-spinalis
colli—*d. Épineux du dos*, Semi-spinalis dorsi—
d. Gantelet, *Gantelet*—*d. Interosseus du poince*,
Flexor brevis pollicis manus—*d. Lunaire*, Semi-
lunar—*d. Membraneux*, Semi-membranosus—*d.*
Nerveux, Semitendinosus—*d. Orbicularis*, *Orbicu-*
laris oris—*d. Tendineux*, Semi-tendinosus—*d.*
Tierce, Hemitritæa (febris).

DEMISSIO ANIMI, Depression.

DEMISSOR, Catheter.

DEMODEX FOLLICULORUM, (ὄνκος, 'lard,'
and ὄζω, 'a boring worm,') *Acarus folliculorum*.

DEMOMANIA, *Dæmonomania*, *Thoma-*
nia, *Ephemonia*, from *dæmon*, 'demon,' a spirit
of good or evil, and *mania*, 'madness.' A variety
of madness, in which the person conceives him-
self possessed of devils, and is in continual dread
of malignant spirits, the pains of hell, &c. Also,
religious insanity, *Sebastomania*.

DEMONSTRATOR, from *demonstrare*, (*de*
and *monstrare*,) 'to show,' 'exhibit.' One who
exhibits. The index finger. See *Digitus*.

DENOMSTRATOR OF ANATOMY. One who ex-
hibits the parts of the human body. A teacher
of practical anatomy.

DEMOTIVUS LAPSUS, *Repentina mora*,
from *demovere*, (*de* and *moveo*,) 'to move off,'
'remove,' and *lapsus*, 'a fall.' Sudden death.

DEMULCENTS, *Demulcentia*, *Involventia*,
Obvolventia, *Lubricantia*, from *demulcere* (*de*,
and *mulcere*,) 'to soothe,' 'to assuage.' (F.)
Adoucissants. Medicines supposed to be capable
of correcting certain acid conditions imagined
to exist in the humours. Substances of a mucil-
aginous or saccharine nature belong to this class.
Demulcents may act directly on the parts with
which they come in contact; but in other cases,
as in catarrh, their effect is produced by con-
tiguous sympathy; the top of the larynx being
soothed by them first, and indirectly the inflamed
portion of the air-passages. In diseases of the
urinary organs, they have no advantage over
simple diluents.—See *Diluents* and *Emollients*.
The following are demulcents;—*Acaciæ Gummi*;
Althææ Folia et Radix; *Amygdalæ*; *Amylum*;
Avenæ Farina; *Cera*; *Cetaceum*; *Cydoniæ Se-*
mina; *Fucus Crispus*; *Glycerhiza*; *Hordeum*;
Lichen; *Linum*; *Olivæ Oleum*; *Sassafras Medu-*
la; *Sesamum*; *Sevum*; *Tragacantha*, and *Ulmus*.

DEMUSCULATUS, from *de*, and *musculus*,
'a muscle.' (F.) *Décharné*. Emaciated, devoid
of flesh.

DENARIUS; a Roman coin, equal in value
to about 10 cents, or 8 pence English. It was
marked with the letter X, to signify 10 asses.

Also, the 7th part of the Roman ounce.

DENDROLIBANUS, Rosmarinus.

DENERVATIO, Aponeurosis.

DENGUE, *Dunge*, *Dunga*, *Dandy*, *Bouquet*,
Bucket Fever, *Eruptive articular fever*, *E. rheu-*
matic fever, *Breakbone*, *Neuralgia*, *Solar* or *Sun*
Fever, *Rheumatismus febrilis*, *Scarlatina rheu-*
matica, *Exanthematis arthrosia*, *Plantaria*, *Febris*
exanthemat'ica articula'ris, (F.) *Giraffe*. A dis-
ease, which first appeared in the years 1827 and
1828, in the West Indies, and in the southern
states of North America. It was extremely
violent in its symptoms, but not often fatal. It
usually commenced with great languor, chilliness,
and pain in the tendons about the smaller joints.
To these symptoms succeeded burning heat and
redness of the skin, pains in the muscles of the
limbs or in the forehead, with vomiting or nausea.
The fever continued for one, two or three days,
and usually terminated by copious perspiration.
In different places, it put on different appear-
ances; but seems in all to have been a singular
variety of rheumatic fever. The usual antiphlo-
gistic treatment was adopted, and successfully.

DENIACH, MINERAL WATERS OF. Deniach is a village in Swabia, at the entrance of the Black Forest. The waters contain carbonic acid, carbonate of iron, carbonate of lime, sulphate of magnesia, and carbonate of soda.

DENIGRATIO, from *denigrare*, (de, and niger, 'black,') 'to become black.' The act of becoming black, as in cases of sphaelus, sugillation, &c.

DENS, Tooth—d. Leonis, *Leontodon taraxacum*.

DENS PROLIFER. A term used by Bartholin for a supernumerary tooth, which appears to grow upon a primitive or parent tooth.

DENT, Tooth—d. de Lion, *Leontodon taraxacum*—d. de Sagesse, see *Dentition*.

DENTAGOGUM, *Dentagum*.

DENTAGRA, *Den'ticeps*, *Dentar'paga*, *Denti'ducum*, *Dentic'ulum*, *Dentago'gum*, *Denta'lis forfex*, *Forfex denta'ria*, *Odonthar'paga*, *Odon'tagra*, *Odontago'gon*, from *dens*, *dentis*, 'a tooth,' and *ayaa*, 'a seizure.' (F.) *Davies*. A tooth-forceps.

DENTAIRES, *Dental*.

DENTAL, *Denta'lis*, *Denta'rius*, *Denticula'tus*, (F.) *Dentaire*, from *dens*, *dentis*, 'a tooth.' That which concerns the teeth. See *Dentiformis*.

DENTAL ARCHES, (F.) *Arcades dentaires*, are the arches formed by the range of alveoli in each jaw.

DENTAL ARTERIES are those arteries which nourish the teeth. They proceed from several sources. The teeth of the upper jaw, e. g., receive their arteries from the *infraorbital* and *superior alveolar*,—themselves branches of the *internal maxillary*. The teeth of the lower jaw receive their branches from the *inferior dental* or *inferior maxillary*, which is given off by the *internal maxillary*, and runs through the *dental canal*, issuing at the *mental foramen*, after having given numerous ramifications to the teeth and jaw.

DENTAL CANALS, (F.) *Conduits dentaires*, *Maxillary canals*. The bony canals, through which the vessels and nerves pass to the interior of the teeth.

DENTAL CARTILAGE, *Cartilago denta'lis*. The cartilaginous elevation, divided by slight fissures, on the biting margins of the gums in infants, prior to dentition. It is a substitute for the teeth.

DENTAL CAVITY, (F.) *Cavité dentaire*. A cavity in the interior of the teeth in which is situated the *dental pulp*.

DENTAL EXCAVATOR. An instrument for the removal of the decayed part of a tooth, preparatory to the operation of filling.

DENTAL FOLLICLE, *Folliculus seu Capsula dentis*, (F.) *Follicule dentaire*. A membranous follicle, formed of a double lamina, in which the teeth are contained before they issue from the alveoli, and which, consequently, aids in the formation of the alveolo-dental periosteum, and of the membrane that envelops the pulp of the teeth. The dental follicles are lodged in the substance of the jaws.

DENTAL GANGRENE, *Caries Dentium*, *Odontalg'ia cariosa*, *Odontonecro'sis*, *Odontosphacel'is*, *Odontosphacelis'mus*, *Necro'sis Dentium*, (F.) *Carie des Dents*. Gangrene or caries of the teeth. See *Caries*.

DENTAL NERVES, (F.) *Nerfs dentaires*. Nerves which pass to the teeth. Those of the upper incisors and canines are furnished by the *infra-orbital nerve*, a branch of the *superior maxillary*, and those of the molars by the trunk of the same nerve. The teeth of the lower jaw receive the nerves from the *inferior maxillary*, which, as well as the *superior maxillary*, arises from the 5th pair.

DENTAL PULP, *Pulpa seu Nucleus seu Substan'tia pulposa dentis*, *Germen denta'le*, (F.) *Pulpe dentaire*. A pulaceous substance, of a reddish-gray colour, very soft and sensible, which fills the cavity of the teeth. It is well supplied with capillary vessels.

DENTAL SURGEON, *Dentist*.

DENTAL VEINS have a similar distribution with the arteries.

DENTALIS, *Odontoid*—d. *Forfex*, *Dentagra*.

DENTALIUM, from *dens*, *dentis*, 'a tooth,' the *dog-like tooth shell*. A genus of shells in the shape of a tooth. They formerly entered into several pharmaceutical preparations, but were useless ingredients; consisting—like shells in general—of carbonate of lime and gelatin.

DENTARIA, *Anthemis Pyrethrum*, *Plumbago Europæa*.

DENTARIUS, *Dentist*.

DENTARPAGA, *Dentagra*.

DENTATA VERTEBRA, *Axis*.

DENTATE, *Denta'ted*, *Denta'tus*, (F.) *Denté*. Cut in the shape of teeth.

DENTATURE, *Digitation*.

DENTATUS, *Dentate*.

DENTELAIRE, *Plumbago Europæa*.

DENTELARIA, *Plumbago Europæa*.

DENTELÉ, *Dentate*—d. *Antérieur petit*, *Pectoralis minor*—d. *Grand*, *Serratus magnus*—d. *Postérieur et inférieur*, *petit*, *Serratus posticus inferior*—d. *Supérieur*, *petit*, *Serratus posticus superior*.

DENTELURE, *Digitation*.

DENTES, *Teeth*, see *Tooth*—d. *Angulares*, *Canine teeth*—d. *Bicuspidati*, *Molar teeth*—d. *Buccales*, *Molar teeth*—d. *Canini*, *Canine teeth*—d. *Clavales*, *Molar teeth*—d. *Columellares*, *Canine teeth*—d. *Ctenes*, *Incisive teeth*—d. *Cuspidati*, *Canine teeth*—d. *Dichasteres*, *Incisive teeth*—d. *Gelasini*, *Incisive teeth*—d. *Gomphii*, *Molar teeth*—d. *Incisores*, *Incisive teeth*—d. *Lactei*, see *Dentition*—d. *Laniarii*, *Canine teeth*—d. *Maxillares*, *Molar teeth*—d. *Molares*, *Molar teeth*—d. *Mordentes*, *Canine teeth*—d. *Oculares*, *Canine teeth*—d. *Primores*, *Incisive teeth*—d. *Rasorii*, *Incisive teeth*—d. *Sapientiae*, see *Dentition*—d. *Serotini*, see *Dentition*—d. *Sophroretici*, see *Dentition*—d. *Sophronistæ*, see *Dentition*—d. *Sophronisteres*, see *Dentition*—d. *Tomici*, *Incisive teeth*.

DENTICEPS, *Dentagra*.

DENTICOLA HOMINIS, see *Odontolithos*.

DENTICULATUM, (*Ligamentum*), diminutive of *dens*, *dentis*, 'a tooth.' *Ligamentum denta'tum*, *Membra'na denta'ta*, (F.) *Ligament denticulé* or *denticulé*. A slender cord, situated between the anterior and posterior fasciculi of the spinal nerves, and between the tunica arachnoidea and pia mater. It is attached to the dura mater, where that membrane issues from the cranium, and accompanies the spinal marrow to its inferior extremity. It sends off, from its outer edge, about twenty slender processes, in the form of *denticuli*, each of which passes outwards and connects itself with the dura mater in the intervals between the anterior and posterior roots of the nerves.

DENTICULATUS, *Dental*.

DENTICULUM, *Dentagra*.

DENTICULUS, *Crena*.

DENTIDUCUM, *Dentagra*.

DENTIER, (F.) from *dens*, *dentis*, 'a tooth.' *Denture*. A set or row of teeth, mounted on metal or ivory, to be adjusted to the alveolar margin. *Dentiers* are simple or double. To the latter, that is, to the full set, the name *Râtelier* is given.

DENTIFORM, *Dentiformis*, from *dens*, *dentis*, 'a tooth,' and *forma*, 'form.' Resembling a tooth in shape.

DENTIFORMIS, Odontoid.

DENTIFRICE, *Dentifricum*, *Remedium dentifricum*, *Odontotrimma*, *Odontomesma*, *Tooth powder*, *Tooth paste*, from *dens*, *dentis*, 'a tooth,' and *fricare*, 'to rub,' a name given to different powders and pastes proper for cleansing the enamel of the teeth, and removing the tartar which covers them. Powdered bark and charcoal, united to any acidulous salt—as cream of tartar—form one of the most common dentifrices.

Electuaire ou Opiate Dentifrice, of the Parisian Codex, consists of coral $\mathfrak{z}\text{iv}$, cuttlefish bones and cinnamon, \mathfrak{aa} $\mathfrak{z}\text{ij}$, cochineal \mathfrak{ss} , honey $\mathfrak{z}\text{x}$, alum gr. iv or v .

Poudre Dentifrice of the Codex is composed of Bole Armeniac, red coral, [prepared,] and cuttlefish bones, \mathfrak{aa} $\mathfrak{z}\text{vj}$, dragon's blood $\mathfrak{z}\text{ij}$, cochineal $\mathfrak{z}\text{j}$, cream of tartar $\mathfrak{z}\text{ix}$, cinnamon $\mathfrak{z}\text{ij}$, cloves, gr. xij , well mixed. The one used by the author, is composed of charcoal and orris root, each one part; prepared chalk, two parts.

Charcoal, finely powdered and mixed with chalk, forms as good a dentifrice as any.

DENTIGEROUS, *Dentiger*, from *dens*, *dentis*, and *gero*, 'I carry.' Tooth carrying, as dentigerous cyst: one containing teeth.

DENTILION, *Leontodon taraxacum*.

DENTINAL, *Dentinalis*, from *dens*, *dentis*, 'a tooth.' Relating to the dentine of the teeth.

DENTINE, see Tooth—d. Secondary, see Tooth.

DENTISCALPIUM, from *dens*, *dentis*, 'a tooth,' and *scalpere*, 'to scrape.' *Odontoglyphum*. This word has been applied to the instrument used for scaling the teeth; to the tooth-pick, (F.) *Cure-dent*; and to the gum lancet,—the *déchaussoir* of the French.

DENTIST, *Dentista*, *Odontoter*, *Dentarius*, *Dental Surgeon*, *Surgeon-Dentist*, (F.) *Dentiste*, from *dens*, *dentis*, 'a tooth.' One who devotes himself to the study of the diseases of the teeth, and their treatment.

DENTISTRY, *Odontotechny*, *Odontiatry*, *Odontotherapy*, *Dental Surgery*, Same etymon. The art of the dentist.

DENTITIO, *Dentition*—d. *Difficilis*, *Dysodontiasis*.

DENTITION, *Teething*, *Dentitio*, *Dentitia*, *Odontophyia*, *Odontiasis*, *Odontosis*, from *dentire*, (*dens*, *dentis*, 'a tooth,') 'to breed teeth.' The exit of the teeth from the alveoli and gums; or rather the phenomena which characterize the different periods of their existence. The germs of the first teeth, *dentes lactei* or *milk teeth*, (F.) *dens de lait*, are visible in the fetus about the end of the second month; and they begin to be ossified from the end of the third to that of the sixth month. At birth, the corona of the incisors is formed, but that of the canine is not completed; and the tubercles of the molares are not yet all united. Gradually the fang becomes developed; and at about six or eight months begins what is commonly called, the first dentition, *Odontia dentitio*, *Odontia lactantium*. The two middle incisors of the lower jaw commonly appear first; and, some time afterwards, those of the upper jaw; afterwards, the two lateral incisors of the lower jaw; and then those of the upper, followed by the four anterior molares; the canine, or eye-teeth, at first, those of the lower, and, afterwards, those of the upper jaw, next appear; and, subsequently and successively, the first 4 molares—2 above and 2 below, 1 on each side. The whole number of the primary, temporary, deciduous, shedding or milk-teeth, (*dentes temporarii*,) (F.) *Dents de lait*, is now 20.

The eruption of the milk teeth takes place, approximately, in the following order:

Central incisors.....	6th to 8th month.
Lateral incisors.....	7th to 10th month.
First molar.....	12th to 14th month.
Canines.....	15th to 20th month.
Second molar.....	20th to 30th month.

The second dentition or shedding of the teeth *odontia dentitio*, *odontia puerilis*, *Dedentition*, begins about the age of 6 or 7. The germs or membranous follicles of these second teeth—to the number of 32—as well as the rudiments of the teeth themselves, are visible, even in the fetus, with the exception of those of the small molares, which do not appear till after birth. They are contained in alveoli of the same shape as those of the first dentition. Their ossification commences at from 3 to 6 months after birth, in the incisors and first molares; at eight or nine months, in the canine; about three years, in the molares; $3\frac{1}{2}$ in the second great molares, and about 10 years in the last. As the alveolus of a new tooth becomes gradually augmented, the septum between it and that of the corresponding milk tooth is absorbed, and disappears. The root of the milk teeth is likewise absorbed; its corona becomes loose and falls out, and all the first teeth are gradually replaced by the permanent teeth, *Dentes serotini*. This second dentition becomes necessary in consequence of the increased size of the jaws. The new teeth have neither the same direction nor the same shape as the old; and they are more numerous, amounting till the age of 25, (sooner or later,) to 28. About this period, a small molaris appears at the extremity of each jaw, which is called *Dens sapientie* or *wisdom tooth*, *wit tooth*, *Dens serotinus*, *Dens saphoroticus*, *D. saphronista*, *D. saphronister*, (F.) *Arrière dent*, *Dent de sagesse*, making the whole number of permanent teeth 32.

The eruption of the permanent teeth is remarkable for its general regularity; so that it constitutes an important means for ascertaining the age of the individual during the early period of life.

First molars.....	7th year.
Central incisors.....	8th year.
Lateral incisors.....	9th year.
First bicuspids.....	10th year.
Second bicuspids.....	11th year.
Canines.....	12th year.
Second molars.....	13th year.

The teeth of the lower jaw precede by a few weeks those of the upper.

During the period of dentition, that is, of the first dentition, the infant is especially liable to disease;—the irritation, produced by the pressure of the tooth on the superincumbent gum, sometimes occasioning pyrexia, convulsions, diarrhoea, &c., which symptoms are often strikingly relieved by a free division of the distended gum. This disordered condition is called *Teething*, *Odontia dentitio*, *Odontiasis*, *Odontalgia dentitio*, *Odontiasis*, *Odontalgia dentitio*, *Odontiasis*.

DENTITIS, *Dentition*.

DENTIUM CORTEX, Enamel of the teeth—d. *Dolor*, *Odontalgia*—d. *Nitor*, Enamel of the teeth—d. *Scalptura*, (Lancing the gums,) see Gum lancet—d. *Vacillantis*, *Odontocisis*.

DENTO, from *dens*, *dentis*, 'a tooth.' One whose teeth are prominent.

DENTOIDEUS, Odontoid.

DENTS BICUSPIDÉES, Bicuspid teeth—d. *Col des*, *Collum dentium*—d. *Conoides*, Canine teeth—d. *de Lait*, see Dentition—d. *Mâchelières*, Molar teeth—d. *Molaires*, Molar teeth—d. *Multi-*

cuspidées, Molar teeth, great—*d. Œillères*, Canine teeth.

DENTURE, Dentier.

DENU'DATION, *Denu'datio*, *Gymno'sis*, from *dennare*, (*de*, and *nudare*, *nudatum*), 'to lay bare.' Condition of a part, deprived of its natural coverings, whether by wounds, gangrene, or abscess. It is particularly applied to the bones, when deprived of their periosteum, and to the teeth when they lose their enamel or dental substance, or when the gums recede from them, and their sockets are destroyed.

DEOB'STRUENT, *Deob'struens*, *Dephrac'ticum*, *Deopp'ilans*, *Deoppilat'ivum*, *Ephrac'tic*, from *de*, and *obstruere*, (*ob*, and *struere*,) 'to obstruct.' (F.) *Désobstruant*, *Désobstructif*, *Désopilatif*, *Désopilant*. Medicines given with the view of removing any obstruction. The word corresponds to *aperient*, in its general, not in its particular sense. It is now almost abandoned, and when used, conveys by no means definite ideas.

DEODORIZER, Antibiotic.

DEONTOL'OGY, *Deontology*, from *τα δέοντα*, 'what are fitting or necessary,' and *λογος*, 'a description.' A word introduced by Bentham to signify morals, or the science of duties.

MEDICAL DEONTOL'OGY, *Deontology*, *in med'ica*, (F.) *Déontologie médicale*, *Medical ethics*, *Medical etiquette*. The duties and rights of medical practitioners.

DEOPILATIVUM, Deobstruent.

DEPASCENS, Phagedenic.

DAPACPERATUS, Impoverished.

DEPERDITIO, Abortion.

DÉPÉRISSEMENT, Emaciation.

DEPHRACTICUM, Deobstruent.

DÉPILATIF, Depilatory.

DEPILATIO, Alopecia—*d. Capitis*, Calvities.

DEPILATION, *Depilatio*, *Dropac'smus*, *Mad'e'sis*, *Mad'isia*, *Pilo'sis*, from *de*, and *pilus*, 'hair.' Loss of hair, either spontaneously or by art.

DEPILATORY, *Depilato'rium*, *Dropac*, *Pilo'thron*, *Ectillot'icus*, *Epilato'rium*, (F.) *Dépilatoire*, *Dépilatif*, *Épilatoire*. Same etymon. Any thing which causes the loss of the hair. Depilatories are usually caustic applications, in which quicklime or some other alkaline substance, sulphuret of iron, &c., enter.

DEPILATORY, COLLEY'S, seems to consist of quicklime and a portion of sulphuret of potassa.

Delcroix's depilatory, and *Plenck's depilatory*, have a similar composition. They all, probably, contain arsenic, in the form of orpiment. The following formula has been given for that of Colley: quicklime, 30 parts; soap lees, 125 parts; sulphur, 4 parts; nitre, 4 parts; orpiment, 12 parts. Another depilatory by F. Boudet is formed of sulphuret of sodium, 3 parts; quicklime, in powder, 10; starch, 10; mix. Rub a little of this powder with water; apply it to the part, and remove the hair in a minute or two with a wooden knife. Dervault recommends the *Depilatory of Martine*, which is a sulphuretted sulphuret of calcium; made by passing sulphohydric acid into quicklime to saturation. A little of this is applied to the part, and in eight or ten minutes it becomes solid. It must then be washed off with cold or hot water, when the skin will be found denuded. Another, given by Redwood, is a strong solution of the sulphuret of barium, made into a paste with starch powder. It is used like the one advised by Boudet. Those that do not contain arsenic are, of course, safer.

A *pitch plaster*, *Pitch-cap*, is sometimes used as a *depilatory*. It of course pulls the hair out by the roots.

DEPILIS, same etymon. Devoid of hair. Hairless.

DÉPLACEMENT DE LA CATARACTE, see Cataract.

DÉPLÉTIF, Depletory.

DEPLETION, *Deple'tio*, from *depleo*, *depletum*, 'I unload.' The act of emptying or unloading, for example, the vessels, by blood-letting and the different evacuations. Also, inordinate evacuation.

DEPLETORY, *Deple'ting*, (F.) *Déplétif*. Having relation to depletion—as 'a depletory or depleting agent.'

DEPLUMA'TIO, *Ptilo'sis*, from *deplumio* (*de* and *pluma*), 'without feathers.' A disease of the eyelids, in which they are swollen, and the eyelashes fall out. See Madarosis.

DEPOSIT, *Depos'itum*, from *depono*, (*de* and *pono*, *positum*, 'to lay or put,') 'to lay or put down.' (F.) *Déposé*. Any thing laid or thrown down. In physiology and pathology, a structureless substance, separated from the blood or other fluid, as the typhous, tuberculous, purulent, melanic, and diphtheritic deposits.

DEPOSIT, Fecculence.

DEPOSITIVE, *Depositi'vus*, same etymon. An epithet used by Mr. Erasmus Wilson to express that condition of the membrane in which plastic lymph is exuded into the tissue of the derma, so as to give rise to the production of small hard elevations of the skin, or pimples. Under 'depositive inflammation of the derma,' he comprises strophulus, lichen, and prurigo.

DÉPÔT, Abscess, Sediment—*d. Laitouse sur la Cuisse*, Phlegmatia dolens—*d. de l'Urine*, Sediment of the urine.

DEPRAVATION, *Deprava'tio*, from *de* and *pravus*, 'bad.' Perversion, corruption—as depravation of the taste, &c.

DEPREHENSIO, Diagnosis, Epilepsy.

DEPRESSIO, *Depres'sio*, *Impres'sio*, from *deprimere*, *depressum* (*de* and *primere*), 'to depress;' *Esphla'sis*. In *Anatomy*, it means an excavation, hollow, or fossa. In *Surgery*, it is applied to a fracture of the cranium, in which the portions of fractured bone are forced inwards; (F.) *Subgrondation*, *Entablement*; called also, *Catap'lexis*, *Campsis*, *Depres'sio*, *Thlasis*, *Depres'sio*. Depression, (F.) *Abaissement*, means Couching.—See Cataract.

Depression also means deflection or dejectedness—*Ademon'ia*, *Ademo'syné*, *Demis'sio animi*.

DEPRESSION, Nervous, Shock, nervous—*d. Vital*, Shock, nervous.

DÉPRESSOIRE, Meningophylax.

DEPRESSOR, (F.) *Abaisseur*. Same etymon. Several muscles have been so termed, because they depress the parts on which they act.

DEPRESSOR ALÆ NASI, *D. labii superioris alæque nasi* seu *Labii superioris proprius*, *Incis'vus me'dius*, *Myrtiformis*, *Contracti'vus alarum nasi* ac *depresso'vus labii superioris*, *Maxillo-alveoli-nasal*—part of the *labialis*, (Ch.) (F.) *Abaisseur de l'aile du nez*. It arises from the superior maxillary bone immediately above the junction of the gums with the two incisor and canine teeth; and passes upwards to be inserted into the upper lip and root of the ala nasi, which it pulls downwards.

DEPRESSOR AN'GULI ORIS, *Triangula'ris*, *Depressor labio'rum communi's*, *D. labiorum*, (F.) *Sousmaxillo-labial*, *Maxillo-labial* (Ch.), *Abaisseur de l'angle des lèvres* ou *Muscle Triangulaire*. A muscle, situate at the lower part of the face. Its form is triangular. It arises from the outer oblique line on the lower jawbone, and terminates in a point at the commissure of the lips, which it pulls downwards.

DEPRESSOR LABII INFERIORIS; *Quadratus*, *Q. Menti*, *Depressor labii inferioris proprius*—

(F.) *Mentonnier-labial*, *Mento-labial* (Ch.), *Carré du Menton*, *Houppé du Menton*, *Abaisseur de la lèvre inférieure*. A small, thin, and quadrilateral muscle, which arises from the external oblique line of the lower jaw, and ascends to the lower lip, where it becomes confounded with the orbicularis oris. It pulls the lower lip downwards and outwards.

DEPRESSOR LABII SUPERIORIS PROPRIUS. D. Alæ nasi—d. Labiorum communis, D. Anguli oris—d. Oculi, Rectus inferior oculi.

DEPRESSOR PALPEBRÆ INFERIORIS. A fleshy bundle, which forms part of the palpebralis muscle. Heister describes it separately, but it is not admitted now.

DEPRESSORIUM, Meningophylax.

DEPRIMENS AURICULÆ, Retrahens auris—d. Maxillæ biventer, Digastricus—d. Oculi, Rectus inferior oculi.

DEPRIMENTIA, Sedatives.

DEPUCILATION, Defecation.

DEPUDICATION, Defecation.

DEPURANTIA, from *depurare* (de and *purus*), 'to purify.' (F.) *Dépurans*, *Dépuratifs*. Medicines were formerly so called, which were supposed to possess the property of removing, from the mass of blood or humours, those principles which disturbed their purity; and of directing them towards some one of the natural excretories. The juices of what were called anti-scorbutic herbs, sulphur, and many other medicines, were ranked under this class.

DEPURATIFS, Depurantia.

DEPURATION, *Depuratio*. Same etymon. *Catharsis*, *Mundatio*. In Pathology, depuration has been used for the process by which nature purifies the animal economy, either by the agency of some eruptive disease, or some spontaneous evacuation, or by the assistance of medicine. See Clarification, and Depuratory.

DEPURATIVE, Depuratory.

DEPURATORY. Same etymon. *Depuratorium*, *Depurative*, (F.) *Dépuratoire*. That which causes depuration, as the urinary and cutaneous depurations. Applied, also, to diseases which have been considered capable of modifying the constitution advantageously, by acting on the composition of the fluids—such as certain eruptions, intermittents, &c. The word is, also, appropriated to medicines and diet, by which the same effect is sought to be induced.

DERADELPH'US, (F.) *Déradelphe*, from *δερν*, 'the neck,' and *ἀδελφός*, 'a brother.' A double monocephalous monster, whose two trunks are united above the umbilicus, and separated below.—G. St. Hilaire.

DER'ADEN, from *δερν*, 'the neck,' and *ἀδην*, 'a gland.' A gland in the neck.

DERADENT'IS, from *δερν*, 'neck,' *ἀδην*, 'a gland,' and *itis*, denoting inflammation. Inflammation of the glands of the neck.

DERADENON'CUS, from *δερν*, 'the neck,' *ἀδην*, 'a gland,' and *οὔκος*, 'a swelling.' Tumor of the glands of the neck.

DERANENCEPHALIA, *Thipsencephalia*, from *δερν*, 'neck,' and *anencephalia*, 'absence of brain.' A monstrosity in which only a small portion of the brain exists, resting on the cervical vertebrae—more properly *derencephalia*.—G. St. Hilaire. According to Is. G. St. Hilaire, a monstrosity characterized by the total absence of encephalon, and of the medulla spinalis into the cervical region.

DERANGED, Insane.

DERANGEMENT, Insanity.

DERATÉ (F.), *Lie'né privatus*, from *de* and *rate*, 'spleen.' Deprived of spleen; as by an operation.

DERCH, Nanus.

DERBIA, Impetigo.

DERENCEPHALIA, *Deranencephalia*.

DERENCEPHALUS, from *δερν*, 'the neck,' and *κεφαλή*, 'head.' A monster whose brain is in the neck.—G. St. Hilaire.

DERIS, Cutis.

DERIVANS, Derivative.

DÉRIVATIF, Derivative.

DERIVATION, *Derivatio*, *Deflectio*, *Parochetensis*, *Antilepsis*, *Antispasmodia*, *Revolusio*, *Cura derivativa seu revulsoria*, from *derivare*, *derivatum*, (de, and *rivus*, 'a river,') 'to turn water from its regular course.' When a 'centre of fluxion' is established in a part, for the purpose of abstracting the excited vital manifestations from some other, a derivation is operated.

The term DERIVATION has likewise been applied to the suction power of the heart—a presumed agency in the circulation of the blood.

DERIVATIVE, *Revellent*, *Revulsivæ*, *Deflectens*, *Derivans*, *Derivativus*, *Antispasmodicus*, *Revulsivus*, *Revulsorius*, (F.) *Dérivatif*, *Réculsif*. Same etymon. A remedy, which by producing a modified action in some organ or texture derives from the morbid condition of some other organ or texture. Revellents are amongst the most important remedies: they include, indeed, every physical and moral agent which is capable of modifying the function of innervation, and therefore almost every article of the materia medica. The following is a list of the chief local Derivatives:

1. EPISPASTICS.—Acidum Aceticum, Acidum Nitricum, Acidum Sulphuricum, Allium, Ammonia, Ammoniacum, Asafoetida, Cantharis, Capsicum, Galbanum, Olea Essentialia, Pix Abietis, Sinapis, Caloric, Friction.

2. VESICANTS.—Ammonia, Argenti Nitras, Cantharis, Cantharis Vittata, Hydrargyri Iodidum rubrum, Ranunculus, Sinapis, Caloric.

3. SUPPURANTS.—Acida Mineralia, Antimonii et Potassæ Tartras, Cantharis, Mezereum, Pix Abietis, Sabina, Tigllii Oleum, Fonticulus, Setaeum.

4. —ESCHAROTICS.—A. *Erc lents*.—Acidum Aceticum, Acida Mineralia, Alumen Exsiccatum, Argenti Nitras, Cupri Sub-Acetas, Cupri Sulphas, Sabina, Saccharum Purissimum.

B. *Actual Canterants*.—Caloric, White Hot Iron, Moxa.

C. *Potential Canterants*.—Acidum Arseniosum, Acidum Nitricum, Acidum Sulphuricum, Antimonii Murias, Argenti Nitras, Calx, Potassa, Potassa cum Calce, Zinci Chloridum.

DERIVATORIUS, Derivative.

DERMA, Cutis.

DERMAD, see Dermal Aspect.

DERMAL, *Dermic*, *Dermatis*, *Dermicus*, from *δερμα*, 'the skin.' Relating or belonging to the skin.

DERMAL ASPECT. An aspect towards the skin or external surface.—Barclay. *Dermad* is used adverbially by the same writer to signify 'towards the dermal aspect.'

DERMAL'GIA, properly *Dermatalgia*, *Dermatodyn'ia*, from *δερμα*, 'the skin,' and *αλγος*, 'pain.' Pain in the skin. Neuralgia of the skin. Rheumatic Dermalgia or Rheumatism of the skin is a form of neuralgia, which is referred, at times, to the nervous trunks, muscles, &c., but appears to be seated in the cutaneous nerves.

DERMATAGRA, Pellagra.

DERMATALGIA, Dermalgia.

DERMATANEU'RIA, *Adermoner'via*, from *δερμα*, *depparos*, 'skin,' a, priv., and *νευρον*, 'nerve.' Paralysis or loss of sensibility of the skin.

DERMATAUZE, Dermatophyma.

DERMATIATRIA, from *δερμα*, *depparos*, 'skin,'

and *carpea*, 'healing.' Healing of cutaneous diseases. The treatment of diseases of the skin—*Diadermatr'ia*, *Dermatocrati'a*.

DERMATITIS, Cytitis, Erysipelatous inflammation.

DERMATOCHOLOSIS, Icterus.

DERMATOCHYSIS, Anasarca.

DERMATOCRATIA, see *Dermatitria*.

DERMATODES, Dermatoid.

DERMATODYNIA, Dermalgia.

DERMATOID, *Dermatoï'des*, *Dermato'des*, *Dermoi'des*, *Dermo'des*, *Dermoid*, from *derma*, 'the skin,' and *eidos*, 'form.' That which is similar to the skin. This name is given to different tissues, which resemble the skin. The dura mater has been so called by some.

Morbi dermatodes, chronic cutaneous diseases

DERMATOL'OGY, *Dermatolog'ia*, *Dermol'ogy*, from *derma*, *dermatos*, 'the skin,' and *logos*, 'a discourse.' A discourse or treatise of the skin.

DERMATOPATHIA, Cutaneous disease.

DERMATOPERISCLERISMUS, Induration of the cellular tissue.

DERMATOPERISCLEROSIS, Induration of the cellular tissue.

DERMATOPHILUS, *Chique*.

DERMATOPHY'MA, *Dermatoph'ia*, from *derma*, *dermatos*, 'skin,' and *phma*, 'tumour.' A tumefaction of the skin.

DERMATOPHY'MATA VENEREA, Condylomata.

DERMATOPHY'TÆ, properly *Dermatoph'ytæ* (*Morbi*), from *derma*, *dermatos*, 'skin,' and *phyton*, 'a vegetable.' An order of cutaneous diseases, including those that depend on, or are characterized by, the presence of parasitic plants. It is made to include Porrhigo and Sycois.

DERMATORRHAG'IA, from *derma*, *dermatos*, 'skin,' and *ragh*, 'rupture.' A discharge of blood from the skin.

DERMATORRHE'A, from *derma*, *dermatos*, 'skin,' and *rho*, 'I flow.' A morbidly increased secretion from the skin.

DERMATOSCLEROSIS, Callosity.

DERMATOSES, Cutaneous diseases.

DERMATOSIES VÉROLEUSES, *Syphilides*.

DERMATOSPASMUS, Horrida cutis.

DERMATOTYLOMA, Callosity.

DERMATOTYLOSIS, Callosity.

DERMATOTYLUS, Callosity.

DERME, Cutis.

DERMIC, *Der'micus*, *Dermat'icus*, *Dermat'icus*, *Dermat'icus*, *Dermat'icus*, from *derma*, 'skin.' Relating to the skin.

DERMITIS, Cytitis.

DERMODES, Dermatoid.

DERMOG'RAPHY, *Dermograph'ia*, from *derma*, 'the skin,' and *grapho*, 'I describe.' An anatomical description of the skin.

DERMOHE'MIA, from *derma*, 'skin,' and *haima*, 'blood.' Hyperemia or congestion of the skin.

DERMOID, Dermatoid.

DERMOLOGY, Dermatology.

DERMOPHYTE, *Dermatophyte*, same etymon. A parasitic vegetable growth on the skin.

DERMO-SKELETON, see Skeleton.

DERMO-SYPHILIDES, *Syphilides*.

DERMOT'OMY, *Dermotom'ia*, from *derma*, 'the skin,' and *temno*, 'to cut.' The part of anatomy which treats of the structure of the skin.

DERODYMUS, Diaphegma.

DERONCUS, Bronchocele.

DERTRON. This word, which is used by Hippocrates, signified, according to some, the omentum or peritoneum, but according to others, the small intestine. See Epiploon.

DÉSALTÉRANTS, Refrigerants.

DÉSARTICULATION (F.), from *de*, and *articulus*, 'a joint.' *Disjointing*. A word used to express the part of the operation, in amputation at an articulation, which consists in dividing the ligaments, and separating the articular surfaces. The word has, also, been used for that kind of anatomical preparation, the object of which is to separate the different bones of the skeleton, and especially those of the head.

DÉSASSIMILATION, see Assimilation.

DESCALORINÈSES, from *de*, 'from,' and *calor*, 'heat.' A name given by Baumes to diseases which are characterized by diminished heat.

DESCEMET, MEMBRANE OF, see Aqueous Humour, and Cornea.

DESCEMETITIS. A term improperly formed, and really signifying 'inflammation of Descemet.' Inflammation of the membrane of Descemet.

DESCENDENS NONI, see Hypoglossus.

DESCENSIO, Catapasis.

DESCENSUS, Catapasis—d. Testiculorum Orchido-catapasis.

DESCENTE, Hernia—d. *de la Matrice*, *Procidencia uteri*.

DESECTUS, Castratus.

DÉSENFLEURE, *Détumescence*.

DESHLER'S SALVE, *Ceratum resinæ compositum*.

DESICCANTIA, Desiccativa.

DESICCATIO, Draining—d. *Herbarum*, *Drying of plants*.

DESICICATION, Drying.

DESICCATIVA, *Desiccat'ia*, *Siccat'ia*, *Exsiccat'iva*, from *desicare*, (*de*, and *sicare*, *siccum*, 'to dry,') 'to dry up.' (F.) *Désiccatifs* Remedies, which, when applied externally, dry up the humours or moisture from a wound.

DÉSINFECTANT, Disinfectant.

DÉSINFECTIO, Disinfection.

DESIPIENTIA, Delirium.

DÉSIRE, Libido—d. *Veneréal*, *Appetite*, *veneréal*, see Libido.

DES'MA, *Des'mē*, *Des'mus*. A bandage, a ligament. Hence:

DESMATUR'GIA, from *desma*, 'bandage,' and *ergon*, 'work.' The doctrine of the application of bandages. Bandaging.

DESMEDION, Fasciculus.

DESNECX, Ligamentous.

DESMIOGNA'THUS, (F.) *Desmiognathe*, from *desmos*, 'ligament,' and *gnathos*, 'jaw.' A double parasitic monster, *Order*, *Polygnathiens*, characterized by a supernumerary imperfect head, which is united, under the neck, by muscular and cutaneous, but no bony attachments, to the principal subject.

DESM'TIS, *Desmophlog'osis*, *Desmophlog'ia*, from *desm*, 'a ligament,' and *itis*, denoting inflammation. Inflammation of ligaments.

DESMOCHAUNO'SIS, from *desmos*, 'ligament,' and *chaunos*, 'relaxation.' Relaxation of an articular ligament.

DESMODYN'IA, from *desm*, 'ligament,' and *odyn*, 'pain.' Pain in the ligaments.

DESMOG'RAPHY, *Desmograph'ia*, from *desmos*, 'a ligament,' and *grapho*, 'a description.' An anatomical description of the ligaments.

DESMOID, *Desmoïdes*, *Desmo'des*, from *desmos*, 'a ligament,' and *eidos*, 'resemblance.' Having the character of, or resembling, ligament.

DESMOD TISSUE, from *desmos*, 'a ligament,' and *eidos*, 'shape.' *Ligamentous Tissue*, *Textus desmoeus*. This tissue is very generally diffused over the human body; has a very close connexion with the articular tissue, and is continuous with it.

in divers places. It constitutes the ligaments, aponeuroses, &c.

DESMOLOGY, *Desmolog'ia*, from *δεσμος*, 'a ligament,' and *λογος*, 'a discourse,' 'a treatise.' That part of anatomy which describes the ligaments.

DESMOPATHY, *Desmopath'ia*, from *δεσμος*, 'ligament,' and *παθος*, 'disease.' Disease of the ligaments.

DESMOPHLOGIA, *Desmitis*.

DESMORRHÆSIS, from *δεσμος*, 'a ligament,' and *ρῆσις*, 'rupture.' Rupture of an articular ligament.

DESMOS, Ligament.

DESMOSUS, Ligamentous.

DESMOTOMY, *Desmotom'ia*, from *δεσμος*, 'a ligament,' and *τεμνειν*, 'to cut.' The part of anatomy which teaches the mode of dissecting the ligaments.

DÉSOSTRUAANT, Deobstruent.

DÉSOSTRUCTIF, Deobstruent.

DÉSOPILATIF, Deobstruent.

DÉSORGANISATION, Disorganization.

DÉSOTOXYGENÈSES, from *de*, and *oxygen*. M. Baumes includes under this title an order of diseases, which he considers dependent upon a diminution in the quantity of the oxygen necessary for the animal economy.

DESPOTATS (F.). Inferm soldiers, formerly charged with the office of removing the wounded from the field of battle: perhaps from *desportator*, 'one who bears away.' The class of *Despotats* was introduced by Leo VI., at the commencement of the 9th century.

DESPUMATION, *Despumatio*, *Apaphris'mos*, *Epaphris'mos*, from *despumare*, (*de*, and *spuma*), 'to skim,' 'to remove the froth.' The separation of the froth and other impurities, which rise, by the action of the fire, to the surface of any fluid.

DESQUMATION, *Desquamatio Cutis*, *Eccl'ep'is*, *Aposyr'ma*, *Apolep'is*, *Apolep'simus*, *Aluuting*, from *desquamare*, (*de*, and *squama*, 'a scale,'), 'to scale off.' Exfoliation, or separation of the epidermis, in the form of scales, of a greater or less size, *Furfuratio*, *Defurfuratio*. This affection is a common consequence of exanthematous diseases.

DESQUMATORIUS, Exfoliative.

DESSÈCHEMENT, Atrophy, Draining.

DESSICATIF, Desiccative.

DESSICATION, Desiccation.

DESTILLATIO, *Coryza*, Distillation—*d. Pectoris*, Catarrh—*d. Uteri*, Leucorrhœa.

DESTRUCTIO, Diaphthora.

DESUDATIO, *Ephidro'sis*, *Sudam'ina*, *Hidro'a*, from *desudare*, (*de*, and *sudare*), 'to sweat.' *Desudation* means a profuse and inordinate sweating, a mucksweat; but, most commonly, the term is applied to an eruption of small pimples, similar to millet seed, which appears chiefly on children, and is owing to want of cleanliness.

DÉSYMPHYSER, see Symphyseotomy.

DETFENTIO, Catalepsy.

DETERGENS, Abstergent.

DETERGENTS, from *detergere*, (*de*, and *tergere*, 'to clean,' 'to cleanse.' *Detergentia*, *Detersoria*, *Etergentia*, *Abtergentia*, *Abstersiva*, *Abtersoria*, *Abluentia*, *Traumat'ica*, *Smect'ica*, *Emundant'ia*, *Mundificat'iva*, *Mundifican'ia*, *Rhypt'ica*, *Abster'sives*, (F.) *Détersifs*, *Mundificatifs*, *Abluenta*. Medicines, which possess the power to deterge or cleanse parts, as wounds, ulcers, &c. They belong to the class of stimulants, or to that of emollients.

DETERMINATION, *Determinatio*, from *de*, and *terminus*, 'a boundary.' Strong direction to a given point;—as 'a determination of blood to the head.'

DÉTTERSIFS, Detergents.

DETERSORIA, Detergents.

DETERSORIUM, from *detergere*, *deteraum*, 'to cleanse.' The place, in ancient bathing establishments, where the bather was cleansed and dried.

DETESTATIO, Castratio.

DÉTORSE, Sprain.

DETRACTIO SANGUINIS, Bloodletting.

DETRITUS, from *deterere*, (*de*, and *terere*, *trium*, 'to rub,'), 'to bruise or wear out.' The residuum, occupying the place of the organic texture of parts which have undergone disorganization.

DÉTROIT ABDOMINALE, Pelvis (brim)—*d. Inférieur*, Pelvis (outlet)—*d. Périnéal*, Pelvis (outlet)—*d. Supérieur*, Pelvis (brim.)

DÉTRONCATION, Detruncatio.

DETRUNCATION, *Detrunca'tio*, (F.) *Détroncation*, from *de*, and *truncus*, 'a trunk.' Separation of the trunk from the head of the foetus, the latter remaining in the uterus.

DETRUSION, *Detru'sio*; same etymon as the next. The act of thrusting or forcing down or away. Applied by Dr. Walshe to lateral displacement of the heart by extraneous pressure.

DETRUSOR URINÆ, *Protru'sor*, from *detrudere*, (*de*, and *trudere*, *trusum*, 'to thrust,'), 'to thrust down or from.' The muscular coat of the urinary bladder was formerly so called. It was, also, named *Constrictor Vesicæ Urinæ*.

DÉTUMESCENCE (F.), *Detumescent'ia*, from *detumere*, (*de*, and *tumere*, 'to swell,'), 'to cease to swell.' A diminution of swelling. This word has nearly the same signification as the French word *Désenflure*. The latter is, however, more particularly applied to the diminution of œdema or anasarca; the former, to the resolution of a tumour properly so called.

DEUNX. The ancient name of a weight of 11 ounces, supposing the pound to consist of 12.

DEURENS (FEBRIS,) Causus.

DEUS COPULATIONIS, Cupido.

DEUTERANGÆA, Dementia.

DEUTERIA, from *deuteros*, 'the second.' Vogel has used this term for the symptoms produced by retention of the secundines. The word was also applied, by the Greeks, to a second or inferior wine.

DEUTERION, Secundines.

DEUTEROPATHIA, *Hysteroopath'ia*, *Morbus secundarius*, from *deuteros*, 'the second,' and *παθος*, 'disease.' A secondary disease. One produced by another, and of which it is only, in some measure, symptomatic, or the sympathetic effect.

DEUTEROSCOPY, *Deuteroscop'ia*, from *deuteros*, 'the second,' and *σκοπεω*, 'I view.' Second sight. A fancied power of seeing future things or events.

DEUTO, *deuteros*, 'second.' A prefix denoting two, or double,—as *deutoxide*, having two degrees of oxidation.

DEUTOAZOTATE ACIDE LIQUIDE DE MERCURE, see *Hydrargyri nitras*.

DEVELOPMENT, *Evolutio*, from (F.) *développer*, 'to unfold,' (F.) *Développement*. In Physiology, it means growth or increase; and in Pathology, its signification is similar. It—as well as *Differentiation*—is at times used, in Physiology, for the kind of increase which takes place by the formation of new tissues from those already existing, as in the embryo, whilst *growth* is restricted to the increase of the same tissue. By development of the pulse is understood an increase in its strength and fullness. Diseases of development, *Machi evolutio'nis*, are such as are peculiar to the period of growth.

'TAKING DEVELOPMENTS.' A term used by practical craniologists to signify the act of measuring prominences of the skull, which are regarded by them as indicating the size of corresponding cerebral organs.

DEVELOPMENT, VESICLE OF, see Vesicle of Development.

DEVERTICULUM, Diverticulum.

DEVIA'TION, *Devia'tio*, from *de*, 'from,' and *via*, 'the way.' Out of the way. By this word is meant—a vicious curvature of the spine or other bones;—faulty direction of the teeth or other part;—the passage of blood, bile, urine, milk, &c., into vessels not natural to them.

DÉVIATION DES RÉGLES, Menstruation (vicarious.)

DEVIL IN A BUSH, *Nigella*.—*d.* Bit, *Veratrum viride*.

DEVIL'S BIT, *Aletris farinosa*, *Scabiosa succisa*, *Chamelirium luteum*.—*d.* Bite, *Liatriis*.—*d.* Dung, *Asafœtida*.—*d.* Shoestrings, *Galega Virginiana*.

DEVIRGINATION, Deforation, Stuprum.

DEVISERATION, Exenterismus.

DÉVOIEMENT, Diarrhœa.

DEWBERRY, AMERICAN, see *Rubus cæsius*.—*d.* Plant, *Rubus cæsius*.

DEWCLAWS, *Crusta genu equinæ*.

DEXIOCAR'DIA, from *δεξις*, 'right,' and *καρδία*, 'the heart.' A case in which the heart is found to beat on the right side. It is met with occasionally in pleurisy and pneumothorax.

DEXIS, Degmus.

DEX'TANS. A weight of 10 ounces, supposing the pound to consist of 12.

DEXTERINA, Dextrine.

DEXTRAD, from *dexter*, 'right-handed.' A term used adverbially by Dr. Barclay to signify 'towards the dextral aspect.' See Mesial.

DEXTRAL ASPECT, see Mesial.

DEXTRALITY, from *dexter*, 'right.' The state of being on the right side. Right-handedness. The state of being right-handed.

DEXTRIN, *Dextrine*, *Dextrinum*, *Dexteri'na*, *British gum*, *Artific'ial gum*, from *dexter*, 'right-handed.' So called, from its refracting the rays, in the polarization of light, more to the right hand than any substance known. A substance obtained by the continued action of diluted sulphuric acid upon starch at the boiling point. It is used in the treatment of fractures, by the 'immovable apparatus.' The bandages are soaked in a solution, in water, of the dextrine—previously moistened thoroughly with tincture of camphor, to prevent it from leaking when the water is added. The solution should be of the consistence of molasses.

DEXTRINUM, Dextrine.

DI, *di*, *dis*, 'bis, twice, double.' Hence, *Dicrotius*, *Digastrius*, &c.

DIA, *dia*, in composition, 'through, asunder, out of, separated.' When prefixed to any therapeutical agent, it meant, in ancient pharmacy, a preparation into which that agent entered.

DIABEBOS, *Astragalus Malleolus*.

DIABÊTE, Diabetes.—*d.* *Chylæna*, Chyluria.—*d.* *Faucæ*, see Diabetes.—*d.* *Insipida*, see Diabetes.—*d.* *Sucræ*, Diabetes (mellitus.)

DIABETES, from *dia*, 'through,' and *βαίω*, 'I pass.' *Urinae profusivum*, *Hyperdiuresis*, *Sipho urinæ*, *Urorrhœgia*, *Polymuria*, *Hydrops ad Mat'ricem seu Mat'ricæ*, *Polyuresis*, *Urocinia*, *Uroorrhœa*, *Dipsacosis*, *Diarrhœa in Urinâ seu urinosa*, (F.) *Diabète*, *Flux d'Urine*. A disease, characterized by great augmentation and often manifest alteration in the secretion of urine; with excessive thirst, and progressive emaciation. Cullen has described two species:—*Diabetes insip-*

idus and *D. Mellitus*; the former, (F.) *Diabète faucæ ou insipide*, *Diabète*, being, simply, a superabundant discharge of limpid urine, of its usual, urinary taste: the latter, *D. Mellitus*, called, also, *Pac'cia Mellita*, *Diabetes Anglicus seu verus*, *Melituria*, *Melithyperuria*, *Glucosuria*, *Glycyrrhœa urinosa*, *Urocinia mellita*, *Saccharorrhœa urinosa*, *Phthisuria*, *Uro-phthisis*, *Tabea diuretica seu diabetica*, *Dyspepsia saccharigena*, *Apocenosia Diabetes Mellitus*, *Saccharine diabetes*, (F.) *Diabète sucré*, *Hyperurorrhœa saccharine*, *Phthisurie sucrée*,—falls under the definition given above. The quantity of urine, discharged in the 24 hours, is sometimes excessive, amounting to 30 pints and upwards; each pint containing sometimes 24 oz. saccharine matter. This replaces the urea, which is not found in quantity in the urine of those labouring under diabetes. Where the disease is situate is not clear. The whole system of nutrition, however, seems to be morbidly implicated. A part of the urine must be formed at the expense of the system, as the egesta frequently far exceed the solid and liquid ingesta. On dissection, no morbid appearance is met with, sufficient to enable us to fix on the seat of this distressing affection.

All the remedies that have been tried have usually been found insufficient in *D. Mellitus*.

D. insip'idus, *Hyperuresis aquosa*, *Hydruria*, *Hydruresis*, *Paruria inconfinens aquosa*, *Diabetes spurius*, *Urorrhœa*, *Ureal Diabetes*, (F.) *Polyurie*, *Hyperurorrhœa*, *Diabète insipide*, *Faucæ diabète*, which occurs in hysterical habits, and has, hence, been called *D. hyster'icus*, is of comparatively trifling moment. Exclusive diet, and attention to the state of the cutaneous transpiration, which have sometimes produced good effects in *D. Mellitus*, have most commonly failed.

DIABETES ANGLICUS, see Diabetes.—*d.* Chylous, Chyluria.—*d.* Insipidus, see Diabetes.—*d.* Lacteus, Chyluria.—*d.* Mellitus, see Diabetes.—*d.* Spurius, see Diabetes.—*d.* Ureal, see Diabetes.—*d.* Verus, Diabetes (mellitus.)

DIABET'IC, *Diabêticus*, (F.) *Diabétique*. Same etymon. Relating to diabetes.

DIABETIC SUGAR, see Glucose.

DIABOT'ANUM, from *δια*, and *βοτάνη*, 'an herb.' A medicine, prepared with herbs.—Galen.

DIABROSIS, Erosion, Corrosion.

DIABROTICUS, from *δια*, and *βρωσκω*, 'I eat or corrode.' A substance, capable of causing erosion of the part to which it is applied. It ordinarily means a medicine, whose activity places it between escharotics and caustics. See Corrosive.

DIACAR'YON, from *δια*, and *καρυον*, 'a nut.' *Rob nucum*. The rob of nuts or of walnuts.

DIACASSIA CUM MANNÂ, Confectio cassiæ.

DIACATHOLICON, *Diacathol'icum*, from *δια*, and *καθολικός*, 'universal.' The name of a purge, so called from its general usefulness. It was an electuary, and composed of the pulp of cassia, tamarinds, leaves of senna, root of polyrrhiza, flowers of the violet, rhubarb root, aniseed, sugar, liquorice, and fennel.

DIACAUSIS, from *διακαίω*, (*δια*, and *καίω*), 'I burn through.' Excessive heat. Over-heating.

DIACAUSTIC, *Diacaust'icus*. Same etymon. That which is caustic by refraction; as a double convex lens, which has been sometimes used for cauterizing an ulcer by directing the sun's rays upon it.

DIACELTATESSON PARACELSI, Pulvis Cornachini.

DIACHALASIS, from *διαχαλαιν*, (*δια*, and *χαλαιν*), 'to be open or relaxed.' Hippocrates uses this word for fracture of the bones of the skull; or for relaxation and separation of the sutures, in consequence of a wound of the head.—Hippocrates.

DIACHALCITEOS, from *δια*, and *χαλκίτις*, 'chalcitis or colcothar.' A plaster, whose composition is the same as that of the diapalma, except that, in place of the sulphate of zinc, a mixture of oil and colcothar is substituted.

DIACHEIRIS/MOS, *Diacheir'isios*, *Tractatio manu'ria*, from *δια*, and *χεῖρ*, 'the hand.' The preparation, administration, and dispensing of medicines.—Hippocrates.

DIACHORE'MA, *Diachore'sis*, from *διαχωρῶ*, (*δια*, and *χωρῶ*, 'I retire,') 'I separate from.' Every kind of excreted matter and excretion; but more particularly the faeces and alvine excretion.—Fœsius, Gorreus. See Excrement.

DIACHOREMA XYSMATODES, see *Ramenta intestinorum*.

DIACHORESIS, Excretion.

DIACHRISIS, Inunction.

DIACHRIST'A, from *δια*, and *χρίω*, 'I anoint.' Medicines, applied as astringents to the velum palati, the palate itself, the tongue, &c. Probably gargles.—Paulus of Ægina.

DIACH'Y/LON, *Diach'yllum*, *Emplastrum diach'yllum*, from *δια*, and *χυλος*, 'juice,' i. e. composed of juices. The plaster of this name was formerly made of certain juices. The term is now confined to the *EMPLASTRUM PLUMBI* or *Lend Plaster*.

DIACHY/LON CUM GUMMI, *Emplastrum gummosum*—d. *Gomme*, *Emplastrum cum gummi-resinis*—d. Gum, *Emplastrum gummosum*—d. *Magnum cum gummi*, *Emplastrum galbani comp.*—d. Simplex, *Emplastrum plumbi*—d. White, *Emplastrum plumbi*—d. Yellow, *Emplastrum gummosum*.

DIACHYTICA, from *δια*, and *χυω*, 'I pour out,' 'I melt.' Medicines which discuss tumours.

DIACINE'MA, from *δια*, and *κινεω*, 'I move.' A slight dislocation.—Celsus, Galen. A subluxation.

DIACLASIS, Refraction.

DIAC'LYSIS, *Diacly'smus*, from *δια*, and *κλυζω*, 'to wash out.' Rinsing or cleansing—especially of the mouth.

DIACLYSMA, Gargarism.

DIACOC'CYMELON, *Diaprunum*.

DIACODION, *Syrupus papaveris*.

DIACODIUM, *Confectio ex Capit'ibus Papaveris*, from *δια*, and *κωδία*, 'a poppyhead.' (F.) *Diacode*. The ancients had various forms for preparing it. The *Syrup of Poppies*—*Syrupus Papaveris* seu *Diaco'dion*—is now substituted for it.

DIACOPE, Abscission, Dissection, Intersection.

DIAC'OPĒ, *En'copē*, from *δια*, and *κοπτεν*, 'to cut.' A cut, incision, fissure, or longitudinal fracture. When used, since Galen, it generally signifies an oblique incision, made in the cranium by a sharp instrument, without the piece being removed. It is not now employed.

DIAC'OPĒ CRA'NI, *Præcis'io* seu *Dissectio Cra'ni*. Opening the head; and separation of the bones of the cranium.

DIACOPRÆ'GIA, from *δια*, *κοπρος*, 'excrement,' and *αἶς*, *αἶγος*, 'a goat.' A name given, in Blancard's Lexicon, to a medicine, composed of goat's dung, which the ancients praised in diseases of the spleen, parotids, &c.

DIAC'RISES, from *δια*, and *κρίνω*, 'I separate.' A class of diseases characterized by alterations of secretion.—Gendrin.

DIACRISIOG'RAPHY, *Diacrisiograph'ia*, from *διακρίσις*, 'separation,' and *γραφῆν*, 'a description.' A description of the organs of secretion.

DIACRIT'ICA SIG'NA. Same etymon. Signs by which one disease can be accurately discriminated from another.—*different'ial diagnō'sis*.

DIACYDONITES, *Cydoniatum*.

DIAD'EMA, *Fascia cap'itis*, *Redimic'ulum*, from *διαδω*, (*δια*, and *δω*, 'to bind,') 'I bind round.' A sort of bandage; advised in headach, in which relaxation of the sutures was apprehended.—Forestus.

DIADERMATRIA, Endermic method.

DIAD'EX'IS, *Diad'ochē*, *Metatopto'sis*, from *διαδεχομαι*, (*δια*, and *δεχομαι*, 'to take or receive,') 'I transfer,' 'I succeed to.' A transformation of a disease into another, differing from the former both in its nature and seat.

DIADOCHE, *Diadexis*.

DIAD'OSIS, from *διατίθεω*, 'to distribute.' In some authors, it means the distribution of nutritive matter over the whole body,—in other words, *nutrition*; whilst, in others, it is synonymous with the remission or cessation of a disease.

DIAD'Œ'US, from *δι*, 'double,' and *αἰδω*, 'the parts of generation.' A monster whose organs of generation and urinary bladder are double. It has only been observed in animals.

DIÆ'RESIS, from *διαίρω*. (*δια*, and *αίρω*, 'I take away,') 'I divide,' 'I separate.' A division or solution of continuity. A surgical operation, which consists in dividing any part of the body. *Hæmorrhag'ia per diæ'resin* is hæmorrhage owing to separation or division of vessels.

DIÆRESIS UNGULÆ, *Onychoptosis*.

DIÆRETICUS, Caustic.

DIÆTA, Diet—d. *Lactea*, *Galactodiæta*.

DIÆTE'MA, has the same signification as diet, with most authors. Galen gives it a more extensive meaning, comprising, under it, what constitutes Hygiene.

DIÆTETICA, Dietetics.

DIAGNOSE, *Diagnosticate*.

DIAGNOS'IS, *Dignō'tio*, *Diagnō'sticē*, *Deprehens'io*, from *δια*, and *γινωσκω*, 'I know.' *Discrimination*, (F.) *Diagnose*, *Diagnosticque*. That part of medicine whose object is the discrimination of diseases, and the knowledge of the pathognomonic signs of each. It is one of the most important branches of general pathology. See Symptoms.

DIAGNOSIS, DIFFERENTIAL, see *Diacritica signa*.

DIAGNOS'TIC, *Discreti'vus*, *Diagnō'sticus*; (F.) *Diagnosticque*. Same etymon. A symptom which is characteristic of a disease.

DIAGNOS'TICATE, — sometimes *diagnose*. To discriminate one disease or phenomenon from another.

DIAGRYDIUM, *Convolvulus scammonia*.

DIAGRY'TUM CYDONIA'TUM, from *diagrydium*, 'scammony,' and *cydonium*, 'quince.' A pharmaceutical preparation, obtained by inspissating and drying, by means of heat, two parts of *scammony*, and one of *quince juice*. It was formerly used as an energetic purgative.

DIAGRY'TUM GLYC'YRRHIZA'TUM. An analogous preparation, containing extract of liquorice in place of quince juice.

DIAGRY'DIUM SULPHURA'TUM. Scammony, which has been exposed to the vapour of burning sulphur. These *diagrydia* are not now used.

DIALEIPSIS, *Apyrexia*, *Intermission*.

DIALEIPYRA, *Intermittent fever*.

DIALEMMMA, *Apyrexia*.

DIALEP'SIS, *Intercept'io*, from *διαλαμβάνω*, 'I intercept.'—Hippocrates employs this word for

the interstices, or intervals, left between the turns of a bandage.

DIALIPSIS, Apyrexia, Intermissio.

DIALYSIS, *Dissolutio*, from *δια*, and *λυσις*, 'solution.' A dissolution or loss of strength; *Resolutio virium*. Weakness of the limbs. Also, a solution of continuity.

DIAMANT, Diamond.

DIAMASTEMA, Masticatory.

DIAMBRÆ SPECIES. A name given by the ancients to powders, one of which bore the name — *Species diambræ sine odore*, — the other, that of *Species diambræ cum odore*. The former was composed of cinnamon, angelica root, cloves, mace, nutmeg, galanga, cardamom, and numerous other substances; the latter, besides, had ambergris and musk. These powders were used as tonics, in cases of debility of the stomach, and in certain nervous affections.

DIAMETERS OF THE PELVIS, see Polvis.

DIAMNES, Enuresis.

DIAMOND, *Adamas*, (from *α*, privative, and *επαω*, 'I conquer.' 'Invincible'; (F.) *Diamant*. So called from its hardness. It is the most precious of all stones, and was formerly conceived to possess extraordinary cordial virtues.

DIAMORUM, *Rob ex moris*, from *δια*, and *μορον*, 'a mulberry.' An ancient syrup prepared with honey and mulberry juice. It was employed as a gargle in sore throat. — Galen.

DIAMOTOSIS, from *μοτος*, 'charpie,' 'lint.' The introduction of lint into an ulcer or wound.

DIANA, Argentum.

DIANANCAMUS, from *δια*, and *αναγκαω*, 'I force.' Coaptation, reduction of a fractured or luxated limb.

DIANOEMA, Imagination.

DIANTHUS CARYOPHYLLUS, from *Δις*, *Δις*, 'Jove.' [?] *ανθος*, 'flower,' and *caryophyllum*, 'the clove.' *Clove Pink*. Also called *Caryophyllum rubrum*, *Tulnica*, *T. horten'sis* seu *rubra*, *Caryophyllum horten'sis*, *Clove July flower*, *Gillflower*, *Carnation*, (Sc.) *Jeraplöris*, *Geraflouris*. Order, Caryophyllæ. (F.) *Œillet giroflée*. The flowers were once much used; but are now only employed in syrup, as a useful and pleasant vehicle for other medicines.

DIAPALMA, *Phœnicium Emplastum*. A plaster composed of equal parts of litharge, olive oil, orange, water, a certain quantity of sulphate of zinc dissolved in water, and white wax. It is classed amongst the topical, desiccative, emollient, resolvent, detersive, and cicatrizing medicines. Mixed with a quarter of its weight of olive oil, it acquires the consistence of an ointment, and forms the *Cerate of Diapalma*.

DIAPASMA, Catapasma.

DIAPEDESIS, *Transudatio*, *Persudatio*, *Persultatio*, from *διαπιδω*, (*δια*, and *πιδω*), 'I leap through.' Exhalation, as of blood, in the form of dew, at the surface of the skin, or of any membrane; *Sweating of blood*, (F.) *Sueur de Sang*, *Hæmorrhag'ia per diapedesin*, *Hæmatopedesin*, *Hæmidrosin*.

DIAPENSIA CORTUSA, Sanicula.

DIAPENTES or DIAPENTE, from *δια*, and *πεντη*, 'five.' A medicine composed of five ingredients. See *Diatesaron*.

DIAPHONICON, *Diaphænix*, *Medicamentum ex Pul'nullis*, from *δια*, and *φωνιξ*, *φωνικος*, 'a date.' A drastic electuary, of which the date was the chief excipient.

DIAPHORESIS, from *διαφωρεω* (*δια*, and *φωρεω*, 'I convey,' 'I dissipate.' (F.) *Diaphorèse*. A greater degree of perspiration than natu-

ral, but less than in sweating. Every kind of cutaneous evacuation.

DIAPHORETIC. Same etymon. *Diapnoëticus*, *Diaphoreticus*, (F.) *Diaphorétique*. A medicine which excites diaphoresis. Diaphoretics are very uncertain as a class. The following is a list of the most reputed:—*Ammoniac Acetatis Liqueur*; *Ammoniac Carbonas*; *Antimonialis pulvis*; *Antimonii et Potassæ Tartaras*; *Antimonii Sulphuretum præcipitatum*; *Asclepias tuberosa*; *Camphora*; *Contraÿerva*; *Dulcanara*; *Eupatorium perfoliatum*; *Guaiaci Lignum*; *Guaiacum*; *Ipecacuanha*; *Mezereum*; *Opium*, and its active principle *Morphia*; *Sarsaparilla*; *Sassafras*; *Serpentaria*; *Spiritus Aetheris Nitrici*; *Sulphur*; *Xanthoxylum*; *Caloric*; *Exercise*, (active); and *Friction*.

The epithet *Diaphoretic* has also been given, by some, to continued fever, accompanied with constant perspiration.

DIAPHORETIC, MINERAL, *Antimonium diaphoreticum*.

DIAPHORETICUM JOVIALE, see Antihæctic.

DIAPHRAGM, *Diaphragma*, *Diaphraç'is*, *Respiratorium Ventris*, *Discretorium*, *Phrenes*, *Septum transversum*, *Discrimen Thoracis et Ventris*, *Disceptum*, *Cinctus*, *Diaçma*, *Præcinctus*, *Diazoma*, *Diazost'ra*, *Hyppocinctus*, *Hyppozoma*, *Perizonia*, *Dissep'tum*, *Dissip'tum*, *Præcordia*, *Succin'gens membra'na seu musculus*, *Succinctus*, *Succinctura*, the *Midriff*, (Prov.) *Heart-seirts*, from *δια*, 'between,' and *φρασσα*, 'I close.' A large, azygous muscle; stretched transversely between the thoracic and abdominal cavities, which it separates from each other; tendinous in the centre; thin, almost circular, and unequally convex, upwards. It is fleshy at its circumference, which is attached to the cartilago ensiformis, to the last six ribs, to the aponeurosis stretched from the last rib to the transverse process of the first lumbar vertebra; and, lastly, to the bodies of the first three or four lumbar vertebrae. When it contracts, its fibres become straight, the chest is enlarged, and the abdomen diminished. It is then an inspiratory muscle. It may, also, diminish the capacity of the chest, and be an expiratory muscle. The muscle plays an important part in sighing, yawning, coughing, sneezing, laughing, sobbing, crying, hicoughing, singing, vomiting, and excretion of the fæces and urine, the expulsion of the foetus, &c.

DIAPHRAGMA AURIS, see Tympanum—d. Cerebri, Tentorium—d. Narium, Septum narium—d. Ventriculorum lateraliū cerebri, Septum lucidum.

DIAPHRAGMAL'GIA, *Diaphragmat'gia*, from *διαφραγμα*, *διαφραγματος*, 'the diaphragm,' and *αλγος*, 'pain.' Pain in the diaphragm.

DIAPHRAGMAT'GIA, *Diaphragmal'gia*.

DIAPHRAGMATIC, *Diaphragmaticus*, (F.) *Diaphragmatique*. Belonging to the diaphragm. A name given to several vessels and nerves.

DIAPHRAGMATIC or PHRENIC ARTERIES. These are distinguished into *superior* and *inferior*. The former, called, also, *supradiaphragmatic*, are two in number, one on each side. They arise from the internal mammary, and descend along the phrenic nerve, to be distributed on the upper surface of the diaphragm. The latter or *infradiaphragmatic* are also two in number. They arise from the upper part of the abdominal aorta, or from the celiac artery, and divide into two principal branches, which are distributed on the lower surface of the diaphragm and in its substance.

The *superior diaphragmatic veins* follow the same course as the arteries, and empty themselves — the *right*, into the *vena cava superior*;

the left, into the corresponding subclavian vein. The two inferior diaphragmatic veins open into the vena cava inferior.

DIAPHRAGMATIC HERNIA, *Phrenic Hernia*, *Diaphragmatocele*. The abdominal viscera are occasionally protruded through the diaphragm, either through some of the natural apertures in the muscle, or through deficiencies, or wounds, or lacerations in it.

DIAPHRAGMATIC OR PHRENIC NERVES, *Internal respiratory* of Sir Charles Bell, are two in number; one on the left side, the other on the right. They arise from the second and third nerves of the cervical plexus, about the middle of the neck, and receive two or three filaments from the brachial plexus, after which they descend into the chest at the sides of the pericardium, and are distributed on the diaphragm.

DIAPHRAGMATIC PLEXUSES are two in number; one right, and the other left. They arise from the upper part of the solar plexus, by a small number of branches, which are distributed to the diaphragm, following exactly the branches of the inferior diaphragmatic arteries.

DIAPHRAGMATIC RING, (F.) *Anneau diaphragmatique* of Chaussier, is a name given to the irregularly quadrilateral aperture by which the vena cava inferior passes through the diaphragm.

DIAPHRAGMATITIS, Diaphragmitis.

DIAPHRAGMATOCELE, see Hernia.

DIAPHRAGMATITIS, from *διαφραγμα*, 'the diaphragm,' and *itis*, a suffix denoting inflammation. *Diaphragmatitis*, *Inflammatio septi transversi*, *Paraphrenitis*, *Empyema Pleuritidis Diaphragmatica*, *Paraphrenesis Diaphragmatica*, *Inflammation of the Diaphragm*. The terms, *Paraphrenitis* and *Paraphrenesis* have been obtained from the Peripatetic philosophy, which supposed the seat of the *πνευ* or soul to be the præcordia. The essential symptoms of diaphragmitis are:—painful constriction around the præcordia, with small, quick, laborious breathing. It is a rare disease.

DIAPHRATON HYMEN, Mediastinum—d. Membrana, Mediastinum.

DIAPHRAXIS, Diaphragm.

DIAPHYTHORA, *Destructio*, from *δια*, and *φθειρω*, 'to corrupt.' Corruption in general; more especially corruption of the foetus in utero.—Hippocrates. Also, corruption of the blood in the stomach.—Galen. See Abortion.

DIAPHYLACTIC, Prophylactic.

DIAPHYSIS, from *διαφω*, 'I rise between.' *Interstitium*, *Discriminatio*: 'an interstice, interval, division.' Any thing that separates two bodies. Also, the middle part or body of a long bone, *Corpus Ossis*. One of the ligaments of the knee.—Hippocrates, Paré.

DIAPYIA. Some lexicographers use this word synonymously with DIAPYESIS or SUPPURATION, others have employed it in opposition to MYOPIA.

DIAPLASIS, Conformation, Reduction.

DIAPLASMUS, Conformation, Reduction.

DIAPNEUSIS, Perspiration.

DIAPNOE, Perspiration.

DIAPNŒA, Perspiration.

DIAPNOGENOUS APPARATUS, see Perspiration.

DIAPNOICUS, Diaphoretic.

DIAPOPHYSES, Transverse processes of the vertebrae.

DIAPOREMA. Anxiety, jactitation; from *διαπορεω*, 'I doubt.'

DIAPRUNUM, *Diacoccyne* lon. A purgative

electuary, of which the pulps of prunes and rhubarb formed the basis. By adding to the *diaprunum simplex* a 24th part of powdered scammony, the *Diaprunum resolutivum seu compositorium* was formed. It was more active than the former.

DIAPYEMA, Empyema, Suppuration.

DIAPYESIS, Suppuration—d. Oculi, Hypopyon.

DIAPYETICA, *Dyapye'mata*, from *διαπυνω*, *διαπυνωσις*, (*δια*, and *πυνω*, 'pus,') 'suppuration.' Medicines which promote suppuration.

DIAPYETICUM, Suppurative.

DIARŒMIA, (F.) *Diarrémie*, from *δια*, 'through,' *ρῶω*, 'I flow,' and *αἷμα*, 'blood.' A pathological condition, said to be common in sheep, in which the globules of the blood are diminished in quantity; the blood itself thinner, and transuding through the coats of the vessels into the cavities.—Delafore.

DIARŒMIE, Diarræmia.

DIARIA, Ephemera.

DIARRHAGE, Fracture.

DIARRHŒA, from *δια*, 'through,' and *ρῶω*, 'I flow.' *Enterorrhœa*, *Incontinentia alvi*, *Alvi profluivum seu fluxus aquosus*, *Ventris profluivum*, *Catiorrhœa*, *Catololysis*, *Alvus cita*, *Cacatoria*, *Coprorrhœa*, *Catarrhus intestinalis*, *Alvi fluxus*, *Rheuma*, *Epiphora Alvi*, *Fluvius alcinus*, *Lacitas alvi*, *Defluxio*, *Lax*, *Looseness*, *Purging*, *Scouring*, (Sc.) *Scour*, (Prov.) *Ray*, *Scutter*, *Skitter*, (F.) *Diarrhée*, *Dévolement*, *Catarrhe intestinal*, *Flux de Ventre*, *Cours de Ventre*, *Courante*. A disease characterized by frequent liquid alvine evacuations, and generally owing to inflammation or irritation of the mucous membrane of the intestines. It is commonly caused by errors in regimen, the use of food noxious by its quality or quantity, &c., constituting the *Diarrhœa stercoraria*. D. *Crapulosa* of writers. It may be acute or chronic. Many varieties have been made by some nosologists—e. g. mucous, —*Diarrhœa mucosa*, *Blennorrhœa*, *Blennorrhœa*, *Mycodiarrhœa*; bilious, —*Ileo-cholosa*, *Diarrhœa biliosa*; serous, —*Hydrochœzia*, *Hydrodiarrhœa*, *Orrhœchia*; dependent upon the matters evacuated. Diarrhœa requires different treatment, according to its nature. If caused, as it often is, by improper matters in the intestinal canal, these must be evacuated; and the astringent plan of treatment must not be adopted, unless the discharges seem kept up by irritability of the intestines, or unless they are colligative. The indiscriminate use of astringents is to be deprecated.

A very fatal diarrhœa prevails amongst the native inhabitants of India, to which Mr. Tytler has given the name *Diarrhœa hecticæ*, because, like hectic fever, it seems to obtain habitual possession of the constitution, to operate upon it with scarcely any perceptible intermission, and, in general, to defy the most powerful remedies.

DIARRHŒA ADIPOSA, *Gras-fondure*—d. cum Apepsiâ, Lientery—d. Arthritica, Coeliagra—d. Biliosa, see Diarrhœa—d. Carnosa, Dysentery—d. Chylosa, Coeliac flux—d. Chymosa, Coeliac flux—d. Crapulosa, see Diarrhœa—d. Cruenta, Hæmatochœzia—d. Dyspeptica, Lientery—d. Hepatica, Hepatirrhœa—d. Ingestorum, Lientery—d. Lienteria, Lientery—d. Mucosa, see Diarrhœa—d. Purulenta, Pyochœzia—d. Sanguinolenta, Hæmatochœzia—d. Stercoraria, see Diarrhœa—d. in Urinâ, Diabetes—d. Urinosa, Diabetes.

DIARRHŒIS/CHESIS, from *Diarrhœa*, and *αἵσσις*, 'arrest.' Arrest of a diarrhœa.

DIARTHRODIAL, *Diarthrodialis*. Relating

to diarthrosis or movable articulations; as *diarthro'dial articulation*. *Diarthro'dial cartilages* or *incrusting cartilages* are the cartilages which invest the articular extremities of bones.

DIARTHROSIS, from *διαρρῶς*, (*dia*, and *ῥῶς*), 'I articulate.' *Dearticulatio*, *Proarthro'sis*, *Aparthro'sis*, *Abarticulatio*, *Coarticulatio*, *Perarticulatio*, *Rota'tio*, (F.) *Emboiture*. A movable articulation. One which permits the bones to move freely on each other in every direction, as in the case of the shoulder joint.

DIASATYR'ION, from *δια*, and *σατυρον*, 'the orchis mascula.' An electuary, of which this plant formed the basis. [?].—Myrepsus. The ancients attributed to it the faculty of exciting the organs of generation.

DIASINCI ANTIDOTUS, Mithridate.

DIASCOR'DIUM, from *δια*, and *σκορδιον*, 'the water germander.' An electuary, so called because this plant entered into its composition. The Parisian codex had a formula for its preparation, under the title, *Electuarium opiatum astrin'gens seu diascor'dium*. (R. fol. scord. \mathfrak{z} ss, rosar. rubr., distort. rad., gentianæ, tormentillæ, sem. berber. aa \mathfrak{z} ss, zingib., piper. long. aa \mathfrak{z} ij, cassia lignea, cinnamon., dictamn. Cretus., styrac. calamit., galban., gum. acacia aa \mathfrak{z} ss, bol. oriental. præpar. \mathfrak{z} ij, extract. vinos. opii. \mathfrak{z} ij, mel. rosat. præp. lbij, vin. hispan. lbss: fiat electuarium.) In place of the *styrax calamita*, the balsam of tolu or benjamin may be used. The opium is, in this preparation, in the proportion of 1 to 384. The diascordium is employed in diarrhoea and dysentery, as a tonic, stomachic, and astringent. The common dose is from a scruple to a drachm and a half. See *Pulvis cretæ compositus*.

DIASOSTIC, Prophylactic.

DIASPASIS, Divulsiō.

DIASPER'MATON, from *δια*, and *σπερμα*, *σπερματος*. The ancient name of two cataplasms, composed of seeds.—Galen, Paulus.

DIASPHYX'IS, from *δια*, and *σφυζω*, 'I strike, I beat.' Pulse. Also, a violent beat of the pulse.

DIASTAL'TIC, *Diastalticus*; from *δια*, 'through,' and *στέλλω*, 'I send.' An epithet applied by Dr. Marshall Hall to the reflex or excitomotor system of nerves; because the actions they induce are performed 'through' the spinal marrow as their essential centre.

DIASTALTICUS, Diastolic.

DIASTASÉ'MIA, (F.) *Diastasémie*, from *διασάσις*, 'separation,' and *αἷμα*, 'blood.' A pathological condition, characterized by a separation of the elements of the blood globules;—the fibrin and albumen separating also from the colouring matter, whilst the fibrin attaches itself to the valves of the heart.—Delafore.

DIASTASE. Same etymon as Diastasis. A vegetable principle, allied in its general properties to gluten, which appears in the germination of barley and other seeds, and, by its presence, converts the starch into sugar and gum.

DIASTASÉMIE, Diastasæmia.

DIAS'TASIS, from *δια*, and *τάσσειν*, 'to place,' 'separation,' 'distance.' *Diastéma*, *Disiden'tia*. A separation of bones, and particularly of the bones of the cranium, from each other; of the radius from the ulna, and the fibula from the tibia. The ancients used this word to designate the three dimensions of the body,—length, breadth, and thickness; for the interval separating the patient from the physician; the swelling

of varicose veins; the time at which some change occurred in disease, &c.

DIASTEMA, Diastasis, Interstice.

DIASTEMATELYT'RIA, from *διαστημα*, 'interstice,' and *ελυτρον*, 'vagina.' An organic deviation, characterized by a longitudinal division or fissure of the vagina.—Breschet.

DIASTEMATENCEPHALIA, from *διαστημα*, and *εγκεφαλος*, 'the brain.' An organic deviation, consisting in a longitudinal division of the brain.—Breschet.

DIASTEMA'TIA, from *διαστημα*. A term employed by Breschet for an organic deviation, characterized by the presence of a fissure in the mesial line of the body.

DIASTEMATOCAU'LIA, from *διαστημα*, and *καυλος*, 'trunk.' An organic deviation, characterized by a longitudinal division of the trunk.

DIASTEMATOCHET'LIA, from *διαστημα*, and *χελος*, 'the lip.' An organic deviation, consisting in a longitudinal division or fissure of the lip.

DIASTEMATOCRA'NIA, from *διαστημα*, and *κρανιον*, 'the cranium.' An organic deviation, consisting in a longitudinal division of the cranium.

DIASTEMATOCYS'TIA, from *διαστημα*, and *κυστις*, 'bladder.' An organic deviation, characterized by a longitudinal division of the urinary bladder.

DIASTEMATOGAS'TRIA, from *διαστημα*, and *γαστηρ*, 'the stomach.' An organic deviation, characterized by a longitudinal division of the stomach.

DIASTEMATOGLOS'SIA, from *διαστημα*, and *γλωσσα*, 'tongue.' An organic deviation, characterized by a longitudinal division or fissure of the tongue.

DIASTEMATOGNA'THIA, from *διαστημα*, and *γναθος*, 'jaw.' An organic deviation characterized by a longitudinal division of the jaw.

DIASTEMATOME'TRIA, from *διαστημα*, and *μητρα*, 'womb.' An organic deviation, characterized by a longitudinal division or fissure of the womb.

DIASTEMATOPYEL'IA, from *διαστημα*, and *πελως*, 'pelvis.' An organic deviation, characterized by a longitudinal division or fissure of the pelvis.

DIASTEMATORA'CHIA, from *διαστημα*, and *ραχίς*, 'spine.' An organic deviation, characterized by a longitudinal division or fissure of the spine.

DIASTEMATORHI'NIA, from *διαστημα*, and *ριν*, 'the nose.' An organic deviation, characterized by a longitudinal division of the nose.

DIASTEMATOSTAPHYL'IA, from *διαστημα*, and *σταφυλη*, 'uvula.' An organic deviation, characterized by a longitudinal division of the uvula.

DIASTEMATOSTERN'IA, from *διαστημα*, and *στερνον*, 'the sternum.' An organic deviation, characterized by a longitudinal division of the sternum.

DIASTEMENTER'IA, from *διαστημα*, and *εντερον*, 'intestine.' An organic deviation, characterized by a longitudinal division of the intestine.

DIAS'TOLÉ, from *διαστέλλω*, (*δια* and *στέλλω*, 'I send,' 'I dilate,' 'I open.' *Relaxa'tio* seu *Remis'sio cordis et arteriarum*. Dilatation of the heart and arteries, when the blood enters their cavities. It is the opposite movement to *systole*, in which the heart and arteries contract

to send forth the blood. *Diastole* and *systole* are, consequently, successive movements. *Diastole*, *Motus cordis diastalticus*, occurs simultaneously in the two ventricles. The almost inappreciable time, which elapses between the diastole and systole has been called *perisystole*, and that which succeeds to the diastole, *peridiastole*. When we speak of the contraction or *systole* of the heart, as well as of its *diastole* or *dilatation*, we mean that of the ventricles. This dilatation is active.

DIASTOLEUS, Dilator.

DIAS'TOLIC, *Diastol'icus*, *Diastalt'icus*; same etymon. Belonging to the diastole of the heart—as 'diastolic impulse of the heart.'

DIASTOLIC IMPULSE OF THE HEART, see Impulse, diastolic.

DIATOMOTRIS, from *διατομω*, (*δια*, and *τομα*, 'mouth,') 'I dilate an aperture.' Dilating instruments, such as the different kinds of specula for the mouth, anus, vagina, &c.

DIASTREMA, Distortion, Perversion, Sprain.

DIASTROPHE, Distortion, Perversion, Sprain.

DIAT'ASIS, *Disten'sio*, from *διατενω*, (*δια*, and *τενω*, 'I stretch,') 'I distend.' *Tension*. The reduction of a fracture by extension and counter-extension.

DIATES'SARON, from *δια*, and *τεσσαρες*, 'four.' An electuary, into the composition of which entered four medicines; viz. the roots of *gentian*, *aristolochia rotunda* and *bay-berries*, each ℥ij. honey ℔ij. The whole was incorporated with extract of *juniper*. The *diapentes* is nothing more than this electuary, mixed with two ounces of *ivory shavings*. The *diatesसारon* is tonic, and it was formerly employed in cases of stings and bites of venomous animals. It was regarded as emmenagogue, alexiterial, and alexipharmic.

DIATHE'SIS, from *διατίθημι*, (*δια*, and *τίθημι*, 'to place,') 'I dispose.' Disposition, constitution, affection of the body: predisposition to certain diseases rather than to others;—(F.) *Imminence morbose*. The principal diatheses, mentioned by authors, are the cancerous, scrofulous, scorbutic, rheumatic, gouty and calculous.

DIATHESIS HÆMORRHOIDALIS, see Hæmorrhoids—d. Hemorrhagic, Hematophilia—d. Nervosa, Nervous diathesis—d. Purulenta, see Pyohæmia—d. Rheumatic, see Rheumatic—d. Sthenica, Sthenia.

DIATRAGACANTHUS. A powder, composed of gum tragacanth, ℥ij. gum Arabic, ℥j and ℥ij. starch, ℥ss, liquorice, ℥ij, as much of the seeds of the melon, and white poppy; seeds of the water-melon, cucumber, and gourd, and sugar candy ℥ij. It was used as a demulcent.

DIATRESIS, Perforation.

DIATRIMMA, Chafing.

DIATRINSANTALON, see Diatrion.

DIATR'ON, *Diatri'um*, from *δια*, and *τρεῖς*, 'three.' A medicine composed of three ingredients. There were formerly two kinds. The first was called *Diatri'um Piperis* or *spec'ies*, and was formed of black and long pepper, aniseed, thyme, and ginger. It was highly stimulating. The second species was known under the name, *Diatrinsantal'on*, *Diatri'um Santal'orum pulvis*, or *Powder of the three Sanders*. They were considered diaphoretic.

DIATRITA'RIL. The Methodists were so called, who pretended to cure all diseases by subjecting the patients to treatment ever third day.

DIAT'RITOS. Relating to every third day, and the means then used by the Methodists to cure disease.

DIAZOMA, Diaphragm.

DIAZOSMA, Diaphragm.

DIAZOS'TER, from *διαζωωωμι*, (*δια*, and *ζωωωμι*, 'to gird,') 'I surround.' The twelfth vertebra of the back, because it corresponds to the girdle, *ζωστήρ*.

DIAZOSTRA, Diaphragm.

DIBOTHRUM LATUM, Bothriocephalus latus.

DICEN'TRA CANADEN'SIS, *Cory'dalis Canadensis*, (*dis*, 'twice,' and *κεντρον*, 'a spur,') *Squirrel corn*, *Colic weed*. Family, *Fumariaceæ*. An indigenous plant, growing from Maine to Wisconsin, which flowers in May, the flowers having the odour of hyacinths. It has been given internally in syphilis, and applied externally in syphilis and gonorrhœa.

DICEPHALIUM, Bicephalum.

DICEPH'ALUS, *Biceph'alus*, *Derod'ymus et Id'ymus*, *Janus*, from *δι*, 'double,' and *κεφαλη*, 'head.' A monster with two heads.

DICERAS RUDIS, *Ditrachyceras rudis*, see Worms.

DICHALCON. A weight, equal to a third part of the obolus.

DICHOPHY'IA, from *διχα*, 'double,' and *φυω*, 'I grow.' A disease of the hairs, in which they split and grow forked.

DICHROMOS, *Verbena officinalis*.

DICIATRIA, Medicine, legal.

DICLIDOSTO'SIS, from *δελος*, 'a double door,' and *οστωσις*, 'ossification.' Ossification of valves—as of the heart.

DICLIS, Valve.

DICOR'YPHUS, *Dicra'nus*, from *δι*, 'double,' *κορυφη*, 'the crown of the head.' A monster with a double vertex or cranium.

DICORYPHUS DIHYPOGASTRIUS, Hemipages.

DICRANUS, Dicoryphus.

DI'CROTUS, *Bisfer'iens*, from *dis*, 'twice,' and *κροτω*, 'I strike.' An epithet given to the pulse, when it seems to beat twice as fast as usual. It is synonymous with the term *rebounding*, the artery rebounding after striking, so as to convey the sensation of a double pulsation. It has been considered, and with truth, to frequently foretell hemorrhage. In bad cases of typhus, it certainly announces such a tendency.

DICTAMNE, *Dictamnus albus*—d. de Crète, *Origanum dictamnus*.

DICTAM'NUS ALBUS, from *Dictamnus*, a town in Crete; *D. Fraxinella*, *Fraxinella*, *F. Dictamnus*, *White Fraxinella*, *Bastard Dictamnus*, (*F.*) *Dictamnus*, *Fraxinella*. The fresh root has been considered nervine, anthelmintic, and emmenagogue. It is not used.

DICTAM'NUS CRETICUS, *Origanum dictamnus*—d. *Fraxinella*, *Dictamnus albus*.

DICTYITIS, Retinitis.

DICTYON, Rete.

DIDELPHYS, Dihysteria.

DIDYMAL'GIA, from *διδυμοι*, 'the testicles,' and *αλγος* 'pain.' Pain in the testicles.

DIDYMIS, Epididymis.

DIDYMITIS, *Hernia humoralis*.

DIDYMOS, Gemellus.

DIDY'MUS, Testicle—d. *Symphogastrius*, *Gastrodidymus*—d. *Symphohypogastrius*, *Hy-*

pogastrodidymus—d. Symphyoperinæus, Pygoidymus—d. Symphyothoracogastrius, Thoracogastrodidymus.

DIECBOLION, from *dia*, and *εκβαλλω*, 'I cast out.' A name given, by the ancients, to a remedy which they believed capable of producing abortion.

DIERENBACH, MINERAL WATERS OF. Dierenbach is a city in Bavaria, two leagues from which is a sulphurous spring.

DIERVIL'LA TRIFIDA, *D. Canadensis* seu *lutea*, Bush honeysuckle. An indigenous plant of the Honeysuckle tribe—Lonicereæ. Ord. Caprifoliaceæ—whose flowers appear from June to August. It has been used as a diuretic; and in gonorrhœa and syphilis.

DIES, *He'mera*, A day, (F.) *Jour*. The day is, properly, the period during which the solar light illumines our horizon: but commonly, also, we designate by the word *day* the period of 24 hours or *Nycthe'meron*, which is frequently divided into four parts—morning, midday, evening, and midnight. In antiquity, great importance was attached to the observation of days in disease. The medical day is usually reckoned at 24 hours,—universally, in estimating the duration of a disease. In parts of the United States, it comprises only the time when the sun is above

the horizon, as regards the administration of medicine, so that if a medicine be ordered to be taken four times a day, it is understood to mean during the 12 hours of day.

DIES CANICULARES, see *Canicula*—d. *Contemplabiles*, Critical days—d. *Contemplantes*, Indicating days—d. *Critici*, Critical days—d. *Decretorii*, Critical days—d. *Indicantes*, Indicating days—d. *Indicatorii*, Indicating days—d. *Indices*, Indicating days—d. *Internuntii*, Critical days—d. *Judicatorii*, Critical days—d. *Radicales*, Critical days.

DIET, *Diæ'ta*, *Diæ'tema*, *Victûs Ra'tio*, (F.) *Diète*. Originally, this word signified nearly the same thing as *Hygiene* and *Regimen*,—that is, Diet was the employment of every thing necessary for the preservation of health and life. At the present day, it signifies a particular kind of food, and, at times, a privation of food and drink;—abstinence. To put any one upon diet, (F.) *mettre quelqu'un à la diète*, means to deprive him of his usual nourishment:—*milk diet* means a diet of milk, &c. See *Aliment*, *Dietetics*, *Hygiene*, and *Regimen*.

Every well regulated hospital has certain dietetic regulations. The following *Table of Dietary* shows the particular regimen that has been selected for the sick, in certain hospitals.

DIET-TABLE OF DIFFERENT HOSPITALS OF GREAT BRITAIN, IRELAND, AND THE UNITED STATES.

I. ENGLAND.

LONDON HOSPITALS.

1. London Hospital.

	COMMON DIET.	MIDDLE DIET.	LOW DIET.	MILK DIET.
<i>Per Day</i>	12 oz. Bread. 1 pint Porter, <i>Men</i> . ½ pint do., <i>Women</i> .		8 oz. Bread.	12 oz. Bread.
<i>Breakfast</i>	Gruel.		Gruel.	Gruel.
<i>Dinner</i>	8 oz. Beef, with Potatoes, thrice a week. 8 oz. Mutton, with Potatoes, twice a week. 8 oz. Potatoes and Soup, with vegetables, twice a week.	The same, except that 4 oz. of Meat shall be given instead of 8 oz.	Broth.	1 pint Milk.
<i>Supper</i>	1 pint of Broth.		Gruel or Broth.	1 pint Milk.

2. St. Bartholomew's Hospital.

	COMMON DIET.	BROTH DIET.	THIN OR FEVER DIET.	MILK DIET.
<i>Daily</i>	Milk Porridge. 12 oz. Bread. 6 oz. Mutton or Beef. 1 pint Broth [with Milk Porridge. Peas or Potatoes, 12 oz. Bread. 4 times a week.] 2 pints Broth. 2 pints Beer, <i>Men</i> . 1 pint, <i>Women</i> . 1 oz. Butter, twice a week.	1 pint Beer. 1 oz. Butter.	Milk Porridge. 12 oz. Bread. 1 pint of Milk, with Tapioca, Arrow-root, Sago, or Rice, as may be prescribed. Barley water.	Milk Porridge. 12 oz. Bread. 2 pts. Milk, with Tapioca, Arrow-root, Sago, or Rice, as may be prescribed. Barley water. 1 oz. Butter. Bread Pudding, three times a week, when ordered.

3. St. Thomas's Hospital.

	FULL DIET.	MILK DIET.	DRY DIET.	FEVER DIET.
Daily.....	2 pints Beer; 14 oz. Bread.	12 oz. Bread.	14 oz. Bread, 2 pints Beer.	12 oz. Bread; two pints Beer.
Breakfast ...	Water Gruel. ½ lb. of Beef when dressed, twice a week; 4 oz. Butter, or 6 oz. of Cheese, thrice a week; ½ lb. Mutton when boiled, thrice a week.	1 pint Milk. 1 pint Milk, 4 times a week. Rice Pudding, thrice a week.	Water Gruel. 4 oz. Butter, 4 times a week; Rice Pudding and 4 oz. of Butter, three times a week.	Water Gruel. ½ lb. Beef, for tea.
Dinner.....				
Supper.....	1 pt. Broth, 4 times a week.	1 pint Milk.		

4. St. George's Hospital.

	EXTRA DIET.	ORDINARY DIET.	FISH DIET.	FEVER DIET.	BROTH DIET.	MILK DIET.
Daily.....	12 oz. Bread. <i>Men.</i> 2 pints Beer. <i>Women.</i> 1½ pint Beer.	12 oz. Bread. 1 pint Beer.	12 oz. Bread.	12 oz. Bread. Barley Water <i>ad libitum.</i>	12 oz. Bread.	12 oz. Bread.
Breakfast .	1 pint Tea. ½ pint Milk. 12 oz. Meat, roasted (weighed with the bone before it is dressed) four days,--boiled; three days, ½ lb. Potatoes.	1 pint Tea. ½ pint Milk. One-half of the meat allowed for extra diet. ½ lb. Potatoes.	1 pint Tea. ½ pint Milk. 4 oz. of plain boiled white fish (as Whiting, Plaice, Flounders, or Haddock.)	1 pint Tea. ½ pint Milk. Arrow-root &c., must be specially directed.	1 pint Tea. ½ pint Milk. 1 pint Broth 6 oz. light Pudding.	1 pint Tea. ½ pint Milk. 1½ pint Rice. Milk four days. ½ lb. Bread or Rice Pudding three days.
Dinner....						
Supper....	1 pint Gruel. ½ pint Milk.	1 pint Gruel. ½ pint Milk.	1 pint Gruel. ½ pint Milk.	1 pint Tea. ½ pint Milk.	1 pint Gruel. ½ pint Milk.	½ pint Milk.

5. Guy's Hospital.

	FULL DIET.	MIDDLE DIET.	LOW DIET.	MILK DIET.	FEVER DIET.
Daily .	14 oz. Bread. 1½ oz. Butter. 1 qt. Table Beer. 8 oz. Meat, when dressed.	12 oz. Bread. 1½ oz. Butter. 1 pt. Table Beer. 4 oz. Meat, when dressed, and ½ pint Broth.	12 oz. Bread. 1 oz. Butter. Tea and Sugar. Half a pound of Beef, (for Beef-tea,) or Arrow-root or Sago, when ordered.	12 oz. Bread. 1 oz. Butter. 2 pints Milk.	6 oz. Bread. 1 oz. Butter. Tea and Sugar.
For each Diet, Gruel or Barley-water, as required.					

6. Westminster Hospital.

	FULL DIET.	MIDDLE DIET.	LOW DIET.		SPOON, OR FEVER DIET.	INCURABLES' DIET.
Daily.....	14 oz. Bread.	10 oz. Bread.	<i>Fixed.</i> ½ lb. Bread.	<i>Casual.</i> —	½ lb. Bread.	½ lb. Bread. ½ lb. Meat. ½ lb. Potatoes. ½ pint Milk. 1 pint Porter.
Breakfast	1 pint Milk Porridge, or Rice Gruel. ½ lb. Meat, roasted, boiled, or chops.	1 pint Milk Porridge, or thin Gruel. ½ lb. Meat, roasted, boiled, or chops.	1 pint Tea, with Sugar and Milk. No fixed Diet for Dinner.	—	1 pint Tea, with Sugar and Milk. Barley Water.	
Dinner..	½ lb. of Potatoes.	½ lb. of Potatoes.		1 pint Broth, or ½ lb. of Bread, or Rice Pudding, or 1 pt. Beef Tea, or a Chop, or Fish.		
Supper..	1 pint Milk Porridge, or Rice Gruel.	1 pint Milk Porridge, or thin Gruel.	1 pint Tea, with Sugar and Milk.	—	1 pint Tea, with Sugar and Milk.	

7. *Middlesex Hospital.*

	DIETA CARNIS, OR MEAT DIET.	DIETA JUSCULI, OR SOUP DIET.	DIETA LACTIS, OR MILK DIET.	DIETA SIMPLEX, OR SIMPLE DIET.	CANCER DIET.
<i>Daily . . .</i>	12 oz. Bread.	12 oz. Bread.	12 oz. Bread.	6 oz. Bread.	12 oz. Bread. $\frac{1}{2}$ lb. Meat. $\frac{1}{2}$ lb. Potatoes. 1 pint Milk.
<i>Breakfast.</i>	1 pint Milk.	1 pint Milk.	1 pint Milk.	1 pint Barley-water.	
<i>Dinner .</i>	<i>Physician's Patients.</i> $\frac{1}{2}$ lb. of Potatoes, 4 oz. dressed meat, (beef or mutton,) roast and boiled alternately, 4 days. $\frac{1}{4}$ oz. Meat in Soup, 3 days. <i>Surgeon's Patients.</i> $\frac{3}{4}$ lb. of Potatoes, $\frac{1}{4}$ oz. dressed meat, (beef or mutton,) roast and boiled alternately. 1 pint Gruel alternately with 1 pint of Barley-water.	1 pint Soup, made with $\frac{1}{4}$ oz. Beef, alternately with 1 pint of Broth with Barley.	$\frac{1}{2}$ pint of Milk with Rice-pudding, 4 days, and with Batter-pudding, 3 days.	1 pint Gruel.	
<i>Supper .</i>		1 pint Gruel.	$\frac{1}{2}$ pint Milk or 1 pint Gruel.	1 pint Gruel or Barley-water.	

8. *North London Hospital.*

	FULL DIET.	MIDDLE DIET.	LOW DIET.	MILK DIET.
<i>Daily</i>	16 oz. Bread. $\frac{1}{2}$ pint Milk. $\frac{1}{2}$ lb. Meat and $\frac{1}{2}$ lb. Potatoes 4 days. 1 pt. Soup or Rice 3 days.	16 oz. Bread. $\frac{1}{2}$ pint Milk. 1 pint Soup or Rice.	8 oz. Bread. $\frac{1}{2}$ pint Milk. Oatmeal for Gruel.	17 oz. Bread. 2 pints Milk.

9. *King's College Hospital.*

	FULL DIET.	MIDDLE DIET.	MILK DIET.	LOW DIET.	FEVER DIET.
<i>Daily</i>	1 pint Beer, or $\frac{1}{2}$ pint Porter. 14 oz. Bread.	14 oz. Bread.	1 lb. Bread.	8 oz. Bread.	—
<i>Breakfast . . .</i>	1 pt. Milk Porridge.	1 pt. Milk Porridge.	1 pint Milk.	1 pint Gruel.	1 pint Gruel.
<i>Dinner</i>	$\frac{1}{2}$ lb. Meat. $\frac{1}{2}$ lb. Potatoes.	$\frac{1}{2}$ lb. Meat. $\frac{1}{2}$ lb. Potatoes.	1 pint Milk.	1 pint Broth.	2 pints Barley-water.
<i>Supper</i>	1 pt. Milk Porridge.	1 pt. Milk Porridge.	1 pint Gruel.	1 pt. Milk Porridge.	1 pt. Milk Porridge.

10. *Dreadnought Hospital Ship.*

	FULL DIET.	ORDINARY DIET.	LOW DIET.	MILK DIET.	FEVER DIET.
<i>Breakfast . . .</i>	1 pint Tea. 1 lb. Bread. $\frac{3}{4}$ lb. Meat.	Ditto. Ditto. $\frac{1}{2}$ lb. Meat.	Ditto. $\frac{1}{2}$ lb. Bread.	Ditto. 1 lb. Bread.	Ditto.
<i>Dinner</i>	$\frac{3}{4}$ lb. Potatoes. 2 pints Beer, (if ordered.)	$\frac{1}{2}$ lb. Potatoes. 1 pint Beer, (if ordered.)	1 pint of Beef Tea.	1 pint Milk.	Gruel.
<i>Supper</i>	1 pint Broth.	1 pint Broth or Gruel.	1 pint Gruel or Milk, (if ordered.)	1 pint Milk.	Gruel or Barley-water.

HOSPITALS.	ORDINARY DIET.	LOW DIET.
LIVERPOOL.	<p><i>Breakfast.</i>—A pint milk porridge, breaded every morn'g.</p> <p><i>Dinner.</i>—(1, 5, 7*)—Boiled beef and vegetables.—(2)—Rice, milk, and bread.—(3)—Stewed beef and potatoes.—(4)—Pea soup and bread.—(6)—Ale, gruel, and bread.</p> <p><i>Supper.</i>—A pint of broth and bread on Sunday and Thursday. A pint of milk and bread on the other days.</p> <p>FULL DIET.—The same as the ordinary diet.</p>	<p>Consists throughout the day of milk porridge, common batter, or rice, pudding.</p> <p>The ale and beer are bought.</p>
BRISTOL.	<p><i>Breakfast.</i>—On Sunday, Tuesday, Thursday, and Saturday, milk porridge; Monday, Wednesday, and Friday, meat broth; 12 oz. of bread on meat days; 14 oz. on the other days.</p> <p><i>Dinner.</i>—(1, 3, 5)—Three-fourths of a pound of meat, with vegetables; two pints of beer daily.—(2, 4, 6, 7)—A pint of gruel or pap.</p> <p><i>Supper.</i>—A wine pint of gruel of meal broth on Sunday; 2 oz. of cheese for the men, one-fourth of an oz. of butter for the women, on Monday, Wednesday, Friday, and Saturday.</p> <p>FULL DIET.—The patients have meat every day.</p>	<p><i>Breakfast.</i>—A wine quart of milk porridge or milk.</p> <p><i>Dinner.</i>—A wine quart of weak broth.</p> <p><i>Supper.</i>—The same as breakfast, 14 oz. of bread, and barley-water for common drink. 20 bushels of malt, 15 lbs. of hops to 14 gals. of strong ale; 21 bush. of malt, and 12 pounds of hops, to 360 gals. of ale; 11 bush. of malt, 7 pounds of hops, to 360 gals. of small beer.</p>
BIRMINGHAM. Revised 1819.	<p><i>Breakfast.</i>—To each man a pint and a half of milk porridge, with 4 oz. of bread added to it. To each woman or child, one pint of milk porridge without bread. To each patient 2 oz. of bread daily.</p> <p><i>Dinner.</i>—(1, 3, 5)—To each man 8 oz. of baked or boiled meat; to each woman or child, 6 oz. To each patient, 6 oz. of vegetables, and one pint of beer daily.—(2)—To each man, a quart of rice or barley broth, made with a variety of vegetables; to each woman or child, one pint and a half.—(4)—Twelve ounces of boiled rice or bread pudding.—(6)—Four oz. of boiled or baked meat, a pint of rice or barley broth, made with a variety of vegetables.—(7)—Six oz. of baked rice or bread pudding, or a pint of rice or barley broth, made with a variety of vegetables.</p> <p><i>Supper.</i>—A pint of broth, milk porridge, or gruel.</p> <p>FULL DIET.—Breakfast the same as in ordinary diet. Dinner, 6 oz. of boiled or baked meat, 6 oz. of vegetables, 6 oz. of baked rice or bread pudding. Supper, the same as ordinary diet.</p>	<p><i>Breakfast.</i>—The same as ordinary diet.</p> <p><i>Dinner.</i>—A pint of broth or rice milk, 6 oz. of baked rice or bread pudding to each patient every day.</p> <p><i>Supper.</i>—The same as ordinary diet.</p>

II. SCOTLAND.

EDINBURGH.	<p><i>Breakfast.</i>—One mutchkin of porridge, three gills of milk or beer; or five and one-fourth ounces of fine bread, milk or beer.</p> <p><i>Dinner.</i>—(1, 4)—One chopin of broth, 8 ounces of butcher's meat boiled in the broth, or beef-steak; five and one-fourth ounces of bread.—(2, 5, 7)—A chopin of broth made of beef and bones, barley, groats, potatoes, and vegetables; five and a-half ounces of bread.—(3, 6)—Potato soup, with beef and veal, or bones; bread as above.</p> <p><i>Supper.</i>—As the breakfast each day.</p> <p>FULL DIET.—At discretion.</p>	At discretion.
GLASGOW.	<p><i>Breakfast.</i>—Milk porridge, quantity not limited, with half a mutchkin of sweet milk, or one mutchkin of buttermilk or beer.</p> <p><i>Dinner.</i>—(1)—Broth made of barley, vegetables, and the dripping of the meat roasted during the week, with a quartern loaf to a man, and half to a woman.—(2, 4)—Beef boiled: 8 oz. to the men, and 6 oz. to the women; a quartern loaf to a man, and half to a woman,—or vegetables.—(3, 6)—Broth, made with beef, barley, and vegetables; a quartern loaf to men, and half to women.—(5)—Potato soup, with cow heels, bones, &c.—(7)—Six oz. of cheese to men, 4 oz. to women; bread as above.</p> <p><i>Supper.</i>—As the breakfast each day.</p> <p>FULL DIET.—At discretion.</p>	At discretion.

* The figures in parentheses denote the days of the week.

III. IRELAND.

5. HOSPITALS of House of Industry.	<i>Per diem</i> .—Sixteen oz. of white bread, one quart of new milk, and one quart of buttermilk for whey. FULL DIET .—Two ounces of bread <i>per diem</i> , one quart of broth, one quart of new milk.	One pint of flummery <i>per diem</i> , one quart of new milk, and one quart of buttermilk for whey.
STEPHEN'S HOSPITAL.	<i>Breakfast</i> .—Half a pound of bread, one pint of milk. <i>Dinner</i> .—(1, 2, 3, 5, 7)—One quart of soup, half a pound of bread, or two pounds of potatoes; one pint of milk or beer.—(4, 6)—Twelve oz. of bread, one quart of sweetened gruel.	Daily, half a pound of bread, two quarts of new milk, and one quart of buttermilk.
ROYAL HOS- PITAL, Phœ- nix Park.	<i>Breakfast</i> .—One pint of oatmeal or rice gruel. <i>Dinner</i> .—Half a pound of meat, three-fourths of a pound of bread; one pound of potatoes. <i>Supper</i> .—One pint of oatmeal or rice gruel. FULL DIET .—Three-fourths of a pound of meat, one pound of bread, half a pound of potatoes, one quart of beer.	<i>Breakfast</i> .—Tea. <i>Dinner</i> .—Half a pound of bread made into panada or pudding.
RICHMOND HOSPITAL.	<i>Breakfast</i> .—One quart of stirabout, one pint of new milk. <i>Dinner</i> .—Bread, 8 oz.; soup, 1 quart. <i>Supper</i> .—Bread, 4 oz.; new milk, one pint. FULL DIET .—Breakfast, bread, 8 oz.; new milk, one pint.—Dinner, bread, 8 oz.; mutton or beef, 3 oz.—Supper, bread, 4 oz.; new milk, one pint.	<i>Breakfast</i> .—Flummery, one pint; new milk, one pint. <i>Supper</i> .—Half a pound of bread, one pint of milk.
BELFAST HOSPITAL.	<i>Breakfast</i> .—One pint of stirabout, one pint of new milk. <i>Dinner</i> .—Half a pound of bread, one pint of new milk. <i>Supper</i> .—One pint of flummery, one pint of new milk. FULL DIET .—Breakfast, one quart of stirabout, one pint of new milk.—Dinner, 2 lbs. of potatoes, one pint of milk.—Supper, one pint of flummery, one pint of new milk.	<i>Per diem</i> .—Quarter of a pound of bread, one quart of gruel, three pints of new milk, half a pint of flummery. Barley water at occasions.
CORK FEVER HOSPITAL.	<i>Breakfast</i> .—One half quartern loaf for every four, and one pint of new milk each. Under 12 years, half a quartern loaf for every eight. <i>Dinner</i> .—(1, 3, 5)—One pound of beef, and two pounds of potatoes. Under 12 years, half a pound of beef, and one pound of potatoes.—(2, 4, 6, 7)—Potatoes and milk. <i>Supper</i> .—One pint of milk and one of stirabout, for adults; half do. for children.	<i>Breakfast</i> .—One-fourth of a lb. of bread, with milk and water sweetened. <i>Dinner</i> .—Gruel, broth, wine, and porter, as ordered by the physician.

IV. UNITED STATES.

PENNSYLVANIA HOS- PITAL, Philada.	<i>Breakfast</i> .—Tea, coffee, or chocolate, with sugar or molasses and milk, and common baker's bread at discretion. <i>Dinner</i> .—Soup always; meat of two kinds—mutton and beef, generally—pork frequently; vegetables, according to the season; potatoes and rice, always. <i>Supper</i> .—Tea and bread: no butter allowed either to breakfast or supper, unless prescribed.	Gruel, gum water, barley water, and other articles prescribed by the physician.
PHILADEL- PHIA HOSPITAL, (Blockley.)	HOUSE DIET .—Arrow-root, gruel, sago, tapioca, rice, beef tea, beef essence, chicken water, rice water, barley water, gum water, flaxseed tea, lemonade. (The diet on which the sick are placed on entering the house until otherwise directed.) MODERATE DIET .—Tea, crackers, broth, rice, mush, milk, potatoes, &c. FULL DIET .—Bread, coffee, tea, white meat, mutton, beef, ham, eggs, butter, soup, potatoes, &c.	The kind and quantity left to the physician. The House Diet may be regarded as low diet.
NEW YORK HOSPITAL, N. York City.	<i>Breakfast</i> .—Bread and black tea, one ounce of tea to every six, and a pint of milk to every eight patients. <i>Dinner</i> .—Tuesdays, Wednesdays, Thursdays, and Saturdays, beef soup, with beef and potatoes, and bread. On Mondays, boiled rice, with one gill of molasses. <i>Supper</i> .—The same as breakfast.	Special diet is directed by the attending physician, and adapted to each case.

The *Diet Scale of the British Navy* allows from 31 to 35½ ounces of dry nutritious matter daily; of which 26 ounces are vegetable, and the rest animal—9 ounces of salt meat, or 4½ ounces of fresh.

That of the Navy of the United States is as follows:—*Three days in the week*—Pork, 16 oz.; beans or peas, 7 oz.; biscuit, 14 oz.; pickles or cranberries, 1 oz.; sugar, 2 oz.; tea, ½ oz.;—40½ oz. *Two days in the week*—Beef, 16 oz.; flour, 8 oz.; fruit, dried, 4 oz.; biscuit, 14 oz.; tea and sugar, 2½ oz.; pickles or cranberries, 1 oz.;—45½ oz. *Two days in the week*—Beef, 16 oz.; rice, 8 oz.; butter, 2 oz.; cheese, 2 oz.; biscuit, 14 oz.; tea and sugar, 2½ oz.; pickles or cranberries, 1 oz.;—45½ oz.

In the Edinburgh workhouse the total allowance of dry food is about 17 ounces—13 ounces vegetable, and 4 ounces animal. In the Edinburgh children's poor-house, the diet consists of milk and porridge, barley broth and bread, amounting to 13 ounces of vegetable food to 4 ounces of animal. These allowances have been found ample for the maintenance of health. Perhaps the case of the smallest quantity of food on which life was vigorously supported was that of Cornaro,—not more than 12 ounces a day, chiefly of vegetable matter, for a period of 58 years.

DIET DRINK. A decoction or potion, variously composed, and used in considerable quantity, for the purpose of purifying the blood. The *Decoctum Lusitanicum* or *Lisbon Diet-drink* is one of the most celebrated. See *Decoctum Sarsaparillæ Compositum*.

DIET, DRY. A diet in which there is but a small or no allowance of fluid.

DIETARY, TABLE OF, see *Diet Scale*.

DIËTE, Diet.

DIETETICS, *Diätetikē, Diätet'ica, Medicīna Diätet'ica*; same etymon. (F.) *Diététique*. A branch of medicine, comprising the rules to be followed for preventing, relieving, or curing diseases by diet. Dietetics is diet administered according to principle. It is an important part of Hygiene. A well regulated system of diet has great power in checking disease, and likewise in preventing it. A proper knowledge of dietetics is, indeed, as important as that of the *Materia Medica*, strictly so called.

Dietetics has been used, also, synonymously with Hygiene.

DIËTÉTIQUE, Dietetics.

DIETETISTS, *Diätetistæ, (F.) Diététistes*. Physicians who apply only the rules of dietetics to the treatment of disease.

DIEU-LE-FILT, MINERAL WATERS OF. The waters of Dieu-le-filt, in France, are chalybeate, and much sought after.

DIEURYSMUS, Dilatation.

DIEX'ODOS, from *δια*, and *εξόδος*, 'an exit or way out.' *Di'odos*. Any opening by which an excretion takes place.

DIFFERENTIAL DIAGNOSIS, see *Diacritica signa*.

DIFFERENTIATION, Development.

DIFFICULTAS·INTESTINORUM, Dysentery.

DIFFLATIO, Perspiration.

DIFFORMITÉ, Deformation.

DIFFUSIBLE, (stimulants) from *diffundere*, (*dis*, and *fundere*, *fundum*, 'to pour,') 'to pour apart or abroad.' Those stimulating medicines are so called, which augment the action of the vascular and nervous systems in an acute but transitory manner.

DIGASTRICUS, from *dis*, 'twice,' and *gαστήρ*, 'a belly:' *Biven'ter, B. Maxill'æ, Dep'rimentæ Maxill'æ Biven'ter, Bigas'ter, (F.) Mastoïdohyogénien, Mastoïdo-génien*—(Ch.), *Digastrique, Abaisseur de la mâchoire inférieure*. The name *Digastricus* was formerly given to several muscles. It is now restricted to one of the muscles of the superior hyoid region. The digastricus is thick and fleshy at its extremities, thin and tendinous at its middle. It is attached to the mastoid groove of the temporal bone, and to a fossa at the side of the symphysis menti. Its tendon passes through an aponeurotic ring, which is attached to the os hyoides.

The use of the digastricus is to depress the lower jaw, or to raise the os hyoides, and to carry it forwards or backwards, as in deglutition.

The strong double-bellied muscle, which forms the gizzard of birds, is also called *Digastricus*.

DIGASTRICUS CRANII, Occipito-frontalis.

DIGASTRIQUE, Digastricus.

DIGENY, see *Generation*.

DIGERENTIA, Digestives.

DIGESTIBLE, *Concoctū hab'ilis*. Capable of being digested. All food is not equally digestible, and some of the most important is the least so:—the fat of meat, for example. Certain substances, again, are entirely rebellious. The following table exhibits the time required for the stomachal digestion of different alimentary substances, in a well-known case, which fell under the care of Dr. Beaumont. The table is extracted from the Author's *Human Health*, Philadelphia, 1844. The most digestible substances are taken as the standard, which has been arbitrarily fixed at 1000; and accordingly, *aponeurosis*, the first article in the table, requiring 3 hours, whilst *pigs' feet soured, rice, &c.*, require but one, its digestibility, compared with that of these aliments, is placed as 333 to 1000; and so of the others. It need scarcely be said, that all these tabular results apply, in strictness, to the individual concerned only; yet they afford useful comparative views, which with exceptions depending upon individual peculiarities, may be regarded as approximations applicable to mankind in general.

ALIMENTS.	Form of preparation.	Time required for stomachal digestion.	Ratio of digestibility compared with the most digestible articles in the table.*	ALIMENTS.	Form of preparation.	Time required for stomachal digestion.	Ratio of digestibility compared with the most digestible articles in the table.
		<i>h. m.</i>				<i>h. m.</i>	
Apocentrosis	boiled	3	333	Marrow, animal, spinal ..	boiled	2	40
Apples, mellow	raw	5	500	Meat and vegetables	hashed	2	30
Do. sour, hard	do.	2	50	Milk	boiled	2	500
Do. sweet, mellow	do.	1	50	Do. do.	raw	2	15
Barley	boiled	2	500	Mutton, fresh	roasted	3	15
Bass, striped, fresh	broiled	3	333	Do. do.	broiled	3	333
Beans, pod	boiled	2	30	Do. do.	boiled	3	333
Do. and green corn	do.	3	45	Oysters, fresh	raw	2	55
Beef, fresh, lean, rare	roasted	3	333	Do. do.	roasted	3	15
Do. do. dry	do.	3	30	Do. do.	stewed	3	30
Do. do. steak	broiled	3	333	Parsnips	boiled	2	30
Do. with salt only	boiled	2	45	Pig, sucking	roasted	2	30
Do. with mustard, &c.	do.	3	30	Pigs' feet, soured	boiled	1	1000
Do.	fried	4	250	Pork, fat and lean	roasted	5	15
Do. old, hard salted	boiled	4	15	Do. recently salted	boiled	4	30
Beets	boiled	3	45	Do. do.	fried	4	15
Brains, animal	boiled	1	45	Do. do.	broiled	3	15
Bread, corn	baked	3	15	Do. do.	raw	3	333
Do. wheat, fresh	baked	3	30	Do. do.	stewed	3	333
Butterf	melted	3	30	Potatoes, Irish	boiled	3	30
Cabbage, head	raw	2	30	Do. do.	roasted	2	30
Do. with vinegar	do.	2	500	Do. do.	baked	3	20
Do.	boiled	4	30	Rice	boiled	1	1000
Cake, corn	baked	3	333	Do. do.	do.	1	45
Do. sponge	do.	2	30	Salmon, salted	do.	4	250
Carrot, orange	boiled	3	15	Sausage, fresh	broiled	3	20
Cartilage	do.	4	15	Soup, barley	boiled	1	30
Catfish, fresh	fried	3	30	Do. bean	do.	3	333
Chesee, old, strong	raw	3	30	Do. beef, vegetables, and			
Chicken, full grown	fricasseed	2	45	bread	do.	4	250
Codfish, cured dry	boiled	2	500	Do. chicken	do.	3	333
Corn (green) and beans	boiled	3	45	Soup marrow bones	do.	4	15
Custard	baked	2	45	Do. mutton	do.	3	30
Duck, domesticated	roasted	4	250	Do. oyster	do.	3	30
Do. wild	do.	4	30	Suet, beef, fresh	do.	5	30
Dumpling, apple	boiled	3	333	Do. mutton	do.	4	30
Eggs, fresh	hard boiled	3	30	Tapioca	do.	2	500
Do. do.	soft boiled	3	333	Tendon, boiled	do.	5	30
Do. do.	fried	3	30	Tripe, soured	do.	1	1000
Do. do.	roasted	2	15	Trout, salmon, fresh	do.	1	30
Do. do.	raw	2	500	Do. do.	fried	1	30
Do. do.	whipped	1	30	Turkey, domestic	roasted	2	30
Flounder, fresh	fried	3	30	Do. do.	boiled	2	25
Fowls, domestic	boiled	4	250	Do. wild	roasted	2	18
Do. do.	roasted	4	250	Turnips, flat	boiled	3	30
Gelatin	boiled	2	30	Veal, fresh	broiled	4	250
Goose, wild	roasted	2	30	Do. do.	fried	4	30
Heart, animal	fried	4	250	Vegetables and meat			
Lamb, fresh	boiled	2	30	hashed	warmed	2	30
Liver, beef's, fresh	do.	2	500	Venison, steak	broiled	1	35

* Pigs' feet soured, rice, and tripe soured, being the most digestible articles in the table, are estimated at 1000.

† In the case of oils, and other substances of similar nature, which undergo little digestion in the stomach, the time merely indicates the period that elapses before they are sent into the duodenum.

DIGESTIFS, Digestives.

DIGESTIO, Digestion—d. Depravata, Dyspepsia—d. Difficilis, Dyspepsia—d. Laesa, Dyspepsia.

DIGESTION, Digest'io, from *digere*, 'to dissolve'; *Cœtio*, *C. Ciborum*, *Pepsis*, *Digestive Process*. Digestion is a function, by means of which alimentary substances, when introduced into the digestive canal, undergo different alterations. The object of this is to convert them into two parts; the one, a reparatory juice, destined to renew the perpetual waste occurring in the economy; the other, deprived of its nutritious properties, to be rejected from the body. This function is composed of a series of organic actions, differing according to the particular organization of the animal. In man they are eight in number, viz: 1. Prehension of food. 2. Mastication. 3. Insalivation. 4. Deglutition. 5. Action of the stomach. 6. Action of the small intestine. 7. Action of the large intestine. 8. Expulsion of the fæces.

DIGESTION is also a pharmaceutical operation,

which consists in treating certain solid substances with water, alcohol, or other menstruum, at a slightly elevated temperature,—in a sand-bath, for example, or by leaving them exposed for some time to the sun.

DIGESTIVE, see Digestives—d. Principle, Pepsin—d. Process, Digestion.

DIGESTIVE TEXTURE. The particular organic condition of substances which affects their digestibility.

DIGESTIVE TUBE, Canal, alimentary.

DIGESTIVES, Digest'iva, *Digerentia*; same etymon as Digestion. (F.) *Digestifs*. A term given, by surgeons, to substances, which, when applied to a wound or ulcer, promote suppuration; as the *ceratum resinæ*, warm cataplasms, fomentations, &c. Such substances are vulgarly termed drawing.

DIGITAL, *Digitalis*; from *digitus*, 'a finger;' having the shape of a finger; *digitated*. Belonging to the fingers.

The *Appendix vermiformis cæci* is sometimes called DIGITAL APPENDIX.

DIGITAL ARTERIES, VEINS, and NERVES, are those distributed to the fingers.

DIGITAL BLANC, Clavaria.

DIGITAL CAVITY, *An'cyroid cavity*, *Cornu decedens ventriculi lateralis*. The occipital portion of the lateral ventricle of the brain.

DIGITAL HUMAIN, Clavaria.

DIGITAL IMPRESSIONS, see Impressions Digital, are the slight depressions observable on the inner surface of the bones of the cranium, which correspond to the cerebral convolutions.

DIGITALE PETITE, *Gratiola officinalis*.

DIGITAL'LE. Same etymon as Digital. (F.) *Doigtier*. A finger stall. The term *Doigtier d'Asdrubali* has been given to a small iron instrument used for measuring the dimensions of the pelvis. Placed at the end of the index finger, it adds to its length and enables it to reach the promontory of the sacrum.

DIGITALINE, see Digitalis.

DIGITAL'IS, from *digitus*, 'a finger,' because its flower represents a finger; *Digital'is purpurea*, *Baccharis*, *Bacchar*, *Baccar*, [?] *Fox-glove*, (Sc.) *Deadmen's bells*, (Prov.) *Flapdock*. Ord. *Scrophularinææ*. *Sex. Syst.* *Didynamia Angiospermia*. (F.) *Digitale*, *Gants de notre dame*, *Doigtier*. The leaves of this plant, which are indigenous in Great Britain, are powerfully sedative, diminishing the velocity of the pulse, diuretic, and sorbafacient. In over-doses, Digitalis causes vomiting, purging, dimness of sight, vertigo, delirium, hiccough, convulsions, and death:—all the symptoms, in short, which characterize the acro-narcotic class of poisons. Its active principle has been called *Dig'italine*. It is a hundred-fold stronger than the most active preparation of digitalis.

Digitalis has been administered in inflammatory diseases, phthisis, active hemorrhage, dropsy, &c.; but although it is a powerful remedy, it has not been as much employed as it probably would have been in particular cases, owing to the over-strained eulogiums, which many have passed upon it in almost all diseases. The average dose is one grain, in the form of pill, which may be repeated every six or eight hours.

DIGITALIS MINIMA, *Gratiola officinalis*.

DIGITARIA DACTYLON, *Cynodon dactylon*.

DIGITATION, *Digital'tio*, *Productio dentata*, *Inner'tio denticulata*, *I. digitata*, (F.) *Dentature*, *Dentelure*. A division into processes having the form of fingers. Several muscles, as the *serati*, exhibit digitations, similar to those which the fingers form, when held separate.

DIGITATIONES TUBARUM FALLOPII, see Tube, Fallopian.

DIGIT'IUM. Desiccation or atrophy of the fingers.—Linnæus. Sauvages calls the same affection *Paronychia Digit'ium*, see *Paronychia*.

DIGITORUM TENSOR, *Extensor brevis digitorum pedis*.

DIG'ITUS, *Dactylos*, *Finger*, (F.) *Doigt*. A name given to the prolongations which form the extremity of the hand. There are five on each hand: the first, the thumb, *Anticheir*, *Pollex*, *Allus Pollex*, *Manus parva majori adjutrix*, *Dig'itus primus seu magnus*, *Pro'manus*, (F.) *Pouce*; the second, the index, *Digitus index seu secundus seu salutaris seu salutarius seu demonstratorius seu demonstrativus*, *Indic'ator*, *Demonstrator*, *Li'chanos*; the third, *D. medius seu tertius seu famulus seu impudicus seu obscenus seu infamis seu verpus*, (F.) *Doigt du milieu*, *middle finger* or *long finger*; the fourth, the ring finger, *Digitus annularis*, *Param'egos*, *D. quartus seu medicinalis seu medicus seu cordis*, *lat'ricus*, (F.) *Annulaire*; and the little finger, *Oit'es*, *Dig'itus annularis seu minimus seu parvus seu quintus seu myops seu ot'ites*, (F.)

Auriculaire, *Petit doigt*. All of these have three phalanges, except the first, which has only two.

DIGITUS ANNULARIS, *Annular finger*—d. *Auricularis*, see *Digitus*—d. *Cordis*, see *Digitus*—d. *Demonstrativus*, see *Digitus*—d. *Demonstratorius*, see *Digitus*—d. *Famosus*, see *Digitus*—d. *Impudicus*, see *Digitus*—d. *Index*, *Index*, see *Digitus*—d. *Indic'atorius*, *Index*, see *Digitus*—d. *Infamis*, see *Digitus*—d. *Magnus*, *Pollex*, see *Digitus*—d. *Medicinalis*, see *Digitus*—d. *Medicus*, see *Digitus*—d. *Medius*, see *Digitus*—d. *Minimus*, see *Digitus*—d. *Myops*, see *Digitus*—d. *Obscenus*, see *Digitus*—d. *Oit'es*, see *Digitus*—d. *Parvus*, see *Digitus*.

DIG'ITUS PEDIS, *Toe*, (Sc.) *Tae*, (F.) *Orteil*. The toes are five in number, and distinguished numerically, reckoning from within to without. The first is, also, called *great toe*, *tom-toe*, (F.) *gros orteil*; the fifth, the *little toe*, *petit orteil*. They have nearly the same organization as the fingers.

DIGITUS PRIMUS, *Pollex*, see *Digitus*—d. *Quartus*, see *Digitus*—d. *Quintus*, see *Digitus*—d. *Salutaris*, see *Digitus*—d. *Salutarius*, see *Digitus*—d. *Salutatorius*, *Index*—d. *Secundus*, *Index*, see *Digitus*.

DIGITUS SEMINOR'TUUS, 'Half dead finger.' A torpid condition of the fingers, met with in connection with chlorosis, and cerebro-spinal affections.

DIGITUS SUPERNUMERARIUS, *Supernumerary finger*, (F.) *Doigt surnuméraire*. An occasional monstrosity occurs, in which there are supernumerary fingers; and at times owing to hereditary transmission.

DIGITUS TERTIUS, see *Digitus*—d. *Verpus*, see *Digitus*.

DIGNA'THUS, from *di*, 'double,' and *γῶβος*, 'lower jaw.' A monster having two lower jaws. —Gurlt.

DIGNOTIO, *Diagnosis*.

DIHYPOGASTRIUS, from *di*, 'double,' and *υπογαστριον*, 'the hypogastrium.' A monster whose pelvis, together with the lower portion of the abdomen, is double.

DIHYSTERIA, *Dime'tra*, *Didel'phys*, *U'terus duplex*, from *di*, 'double,' and *υἱστρον*, 'uterus.' The state in which there is a double uterus.

DIJUDICATIO, *Crisis*.

DILACERATIO, *Laceration*.

DILATANTS, *Dilating agents*.

DILATEUR, *Dilator*—d. *Antérieur du larynx*, *Crico-thyroid muscle*—d. *Postérieur du larynx*, *Crico-arytenoid*, *posterior*.

DILATATIO BRONCHIORUM, *Bronchia*, *dilatation of the*—d. *Ventriculi*, *Gastroectasis*—d. *Intestinorum*, *Enterectasis*.

DILATATION, *dilatatio*, from *dilatare*, *dilatatum*, (*latum* *fivere*) 'to enlarge;' *Eury'smus*, *Aneurys'mus*, *Dieury'smus*. Augmentation of the bulk of a body, occasioned by a separation of some of its molecules. Caloric has the property of dilating all bodies. In Surgery, it means the accidental or preternatural augmentation of a canal or opening: as in aneurisms, varices, &c., or the process of enlarging any aperture or canal. When used so as to obtain a view of parts, as by the speculum, it is termed *Dioptris'mus*.

DILATATOIRE, *Dilator*.

DILATATORIUM, *Dilator*.

DILATORIUS, *Dilator*.

DILATING AGENTS, *Dilatan'tia*, (F.) *Dilatans*. Certain substances used in surgery, either to keep parts separate which have a tendency to unite—as after opening an abscess, to prevent the edges of the incision from uniting; or to increase and dilate openings of canals, either when natural, or formed accidentally or artificially. These agents differ from each other: the

chief are—*prepared sponge tents, gentian root, bougies, summts, dried peas for issues, &c.* •

DILATOR, *Dilatatorius*, *Diastoleus*, (F.) *Dilatateur* ou *Dilatatoire*. A muscle, whose office it is to dilate certain parts; such as the inspiratory muscles, which dilate the chest.

DILATOR, *Dilatatorium*, (F.) *Dilatateur*. An instrument, used for dilating a wound, excretory canal, or other natural or artificial opening. When employed to obtain an inspection of internal parts, it is termed *Speculum*, *Dioptra*, or *Dioptron*. There are several instruments of this kind, each taking its name from the part to which it is applied; as *Speculum Oris*, *S. Nasi*, *S. Uteri*, &c.

DILATOR, ARNOTT'S. A modification of the old dilators for strictures of the urethra. It consists of a tube of oiled silk, lined with the thin gut of some small animal to make it air-tight, and fixed on the extremity of a small canula, by which it is distended with air or water, from a bag or syringe at the outer end, whilst a stop-cock or valve serves to keep the air or water in, when received. As soon as the bag is passed within the stricture or strictures, as much air is to be injected into it as the patient can easily bear. The instrument is not much used.

DILATORES ALARUM NASI, Compressor Naris.

DILATRIS TINCTORIA, *Lachnanthes tinctoria*.

DILL, *Anethum graveolens*.

DILLY, *Anthemis cotula*.

DILUENTIA, from *diluo*, (*dis*, and *here*), 'I wash away.' (F.) *Dilutants*. Medicines which have been conceived proper for augmenting the fluidity of the blood and other animal liquids. All aqueous drinks are diluents. They are administered, with great advantage, in various diseases. In fever, water, which is the most familiar diluent, may be freely allowed; the only precaution being to give it *hot* in the cold stage, *cold* in the hot, and *tepid* in the sweating. In diseases, where it is considered necessary to abstract blood largely, diluents should not be given too freely. The abstraction of blood occasions activity of absorption, and the mass is speedily restored. It is also obvious, that in cases of inflammation of the mammae, in nurses, diluents should not be freely allowed, as they increase the secretion of milk, and add to the irritation. When *demulcents* are exhibited in cases of urinary disease, they act simply as diluents: their mucilaginous portion is digested in the stomach and small intestine,—the watery portion alone being separated by the kidney.

DILWEED, *Anthemis cotula*.

DIMETRA, *Dihysteria*.

DINANT, MINERAL WATERS OF. Dinant is a small town, six leagues from St. Malo, in France, where are mineral waters, containing carbonate of iron, chloride of sodium, &c. They are much esteemed.

DINGEE, *Dengue*.

DINIC, *Antidinic*.

DINKHOLD, MINERAL WATERS OF. A rich carbonated water, situate near the junction of the Lahn with the Rhine, in the duchy of Nassau. It contains sulphate of soda, chloride of sodium, carbonate of soda, sulphate of lime, carbonate of lime, and sulphate of magnesia.

DINOMANIA, *Tarantismus*.

DINUS, *Vertigo*—*d. Scotoma*, *Scotodynia*—*d. Vertigo*, *Vertigo*.

DIOBOLON, *Scrupule*.

DIOCRES, *Psatill*.

DIODONCEPHALUS, (F.) *Diodoncéphale*, from *dis*, 'twice,' *odous*, *odontos*, 'a tooth,' and *κεφαλη*, 'head.' A monster with a double range of teeth, or double jaw.

DIODOS, *Dioxodos*.

DIÖCESIS, *Dispensation*.

DIONCOSIS, from *dia*, and *oikos*, 'a tumour.' The Methodists applied this name to a sort of tumefaction or plethora, occurring either directly from too great a quantity of fluid circulating in the system, or owing to the retention of substances which ought to be excreted. It is the antithesis to *symplocosis*. See *Intumescence*.

DIONYSIANUS, (F.) *Dionysien*, from *Διονυσος*, 'Bacchus,' who is represented by the poets as wearing horns. One who has long or horn-like excrescences.

DIONYSIS/CUS, same etymon. One who has a long horn-like excrescence on the frontal region. —Vogel.

DIOPHTHALMUS, *Binoculus*.

DIOPTRA, *Speculum*, see *Dilator*.

DIOPTRISMUS, *Dilatation*.

DIOPTRON, *Speculum*, see *Dilator*.

DIORTHOSIS, from *διορθω*, 'I make straight.' The reduction of a fractured or luxated limb.

DIOSCOREA, (after *Dioscorides*.) see *Yam*.

DIOSCOREA VILLOSA, *Wild Yamroot*; indigenous: *Order*, *Dioscoriaceæ*; flowering in July. A decoction of the root has been prescribed in bilious colic. It is said to be expectorant, diaphoretic, and, in large doses, emetic.

DIOSCURI, *Parotis*.

DIOSMA, *D. crenata*.

DIOSMA CRENATA, from *dios*, 'divine,' and *oσm*, 'odour.' *Bavosma crenata*, *Agathosma crenatum*, *Buchu Leaves*, *Diosma* (Ph. U. S., 1842, *Buchu*, 1851), *Bucku*, *Bocchoe*, *Bocho*, *Boccho*, *Buckho*, (F.) *Diosmæ crénelée*. *Ord. Diosmææ*. A South African plant, the powder of whose leaves is used by the Hottentots to perfume their bodies. It has been employed in chronic affections of the bladder and urinary organs in general. It has also been given in cholera. It is often adulterated in commerce, by the substitution of less potent plants of the same family, as *Diosma serratifolia*, and *Euplexurum serrulatum*.

DIOSMA SERRATIFOLIA, *D. crenata*.

DIOSMÉE CRÉNELÉE, *Diosma crenata*.

DIOSPYROS LOTUS, *Faba Græca*, *Indian Date Plum*, (F.) *Plaqueminier d'Europe*. *Ord. Ebenaceæ*. This tree grows in some of the southern parts of Europe. Its fruit is very astringent, and has been recommended in dysentery and hemorrhage.

DIOSPYROS VIRGINIANA, *Lotus Virginia'na*, *Persimmon*. A common tree in the middle parts of the United States. The fruit, *Persimmons*, *Yellow Plums*, *Winter Plums*, *Seeded Plums*, which is only eatable after frost, (when it is tolerable,) is sometimes made into cakes with bran. These, being dried in an oven, are kept to make beer. When bruised in water, fermentation takes place. The unripe fruit, *Diospyros* (Ph. U. S.), is distressingly acerb and astringent. It has been used as an astringent. The bark of the tree is extremely bitter, and may be used where bitters are indicated.

DIO'TA, *Dyo'ta*, from *dis*, and *ovs*, *wros*, 'ear.' Two-eared, two-handled. Applied to a wooden cup, lined with a composition of *resin*, *cinnamon*, *cloves*, and *ginger*, to give more flavour to beer. It was formerly much used in the north of Europe.

DIPHORUS, *Diphros*.

DIPHRUS, *Diphorus*, from *dis*, 'two,' and *φέρω*, 'I carry.' Properly a seat for two. A close stool.

DIPHRUS MAIEUTICUS, *Sella obstetric'ia*. An obstetric chair.

DIPHYRGES, from *dis*, and *φρυγω*, 'I torrefy.' The oxide of copper, more or less pure. The ancients reckoned three kinds of diphryges, which they used as astringents.

DIPHTHERIA, *Diphtheritis*.

DIPHThÉRITE, Diphtheritis—*d. Buccale*, Stomatitis, pseudomembranous.

DIPHThÉRITIS, *Diphther'ia*, *Diphtherit'ic Inflammation*, *Pellic'ular Inflammation*, (F.) *Diphthérie* from *διφθέρα*, 'a membrane.' A name given by M. Bretonneau to a class of diseases, which are characterized by a tendency to the formation of false membranes; and affect the dermoid tissue,—as the mucous membranes, and even the skin.

DIPHThERITIS OF THE THROAT, Angina pellicularis—*d. Trachealis*, Cynanche trachealis.

DIPLASIASMUS, Duplication.

'DIPLOCEPHALIA, from *διπλος*, 'double,' and *κεφαλή*, 'head.' A monstrosity characterized by the presence of two heads on the same body.

DIPLECOIA, Double hearing.

DIPLOË, from *διπλω*, 'I double.' *Diplo'sis*, *Meditull'ium*, *Medium Calva'ria Discrimen*. The cellular structure, which separates the two tables of the skull from each other. The ancients applied the term, also, to the proper coat of the uterus [?].—Rolfink, in Castelli.

The Diploë has the same use as the cellular structure of bones in general. See Cancelli.

DIPLOGENESIS, *διπλος*, 'double,' and *γενεσις*, 'generation,' (F.) *Diplogénese*. An organic deviation, which consists of the duplication of parts. Monstrosity by duplication. *Duplic'itas monstro'sa*, constituting *double monsters*.

DIPLOLEPIS GALLÆ TINCTORIÆ, see Quercus infectoria.

DIPLO'MA. Same etymon as Diploë. A letter or writing conferring some privilege: usually applied to the document, certifying that a person has obtained the title of Doctor. It was so called because formerly written on waxed tables, folded together. Diploma is also used in pharmacy for a vessel with double walls,—as a water-bath.

DIPLO'PIA, from *διπλος*, 'double,' and *ὀπτομα*, 'I see.' *Visus Duplicat'is*, *Ditto'pia*, *Dit-top'sia*, *Amphamphoterodiplo'sia*, *Amphodiplo'pia*, *Ambio'pia*, *Double Vision*, (F.) *Bévue*. An affection of the sight, in which two distinct sensations are produced by the same object, which consequently seems double. Sometimes more than two are seen; but still the disease is termed *Diplo'pia* and *Suffusio Multi'plicans*. This affection arises from some derangement in the visual axes, in consequence of which, the images are not impressed on corresponding parts of the retina of each eye. At times, the diplopia affects one eye, or each eye when used alone, owing to some irregularity in the curvature, density, or position of the dioptric media. It is then called *Unioct'ular Diplopia*.

The diplopia of both eyes has been termed *Amphodiplo'pia* and *Amphoterodiplo'pia*.

DIPLO'SIS, Diploë, Duplication.

DIPLOSO'MA, from *διπλος*, 'double,' and *σῶμα*, 'body.' The *Diplosoma crenata* is an imperfectly described entozoon, which has been passed from the urinary bladder. It varies in length from four to six or eight inches, and is thinnest in the middle, where it is bent at an acute angle upon itself, so that the two halves hang nearly parallel, and give to it an appearance as if two worms had been tied together by their heads. It has been confounded with the *Spiroptera hominis*.

DIPLOSO'MATIA, *Diplosom'ia*, from *διπλος*, 'double,' and *σῶμα*, *σώματος*, 'body.' An organic deviation, characterized by the presence of two complete bodies, united by one or more of their parts.

DIPNOÏS, from *dis*, and *πνεν*, 'breath.' Having two vent holes; *Bispi'rus*. An epithet ap-

plied to wounds which pass through a part, and admit the air at both ends.—Galen.

DIPROSO'PUS, *Irid'ymus* et *Opod'ymus*, from *δι*, 'double,' and *προσωπον*, 'countenance.' A monster having a double face.

DIPROSO'PUS DIÆDÆ'us, from *δι*, 'double,' and *προσωπον*, 'countenance,' and *δι*, 'double,' and *αἰδία*, 'parts of generation.' A double monster, in whom the duplication affects superiorly the face, and inferiorly the anterior pelvic region.—Barkow.

DIPROSO'PUS DIHYPOGAS'TRIS, from *δι*, 'double,' and *προσωπον*, 'countenance,' and from *δι*, 'double,' and *υπογαστριον*, 'the hypogastrium.' A double monster in whom the duplication affects superiorly the face, and inferiorly the lower part of the body, four lower extremities being always present.—*Tetrast'celus*.

DIPSA, Thirst.

DIPSACOS, Diabetes.

DIPSACUS FULLO'NUM, *Herba Card'ui Ven'eri*, *Card'ius Ven'eri*, (F.) *Cardère cultivé*, *Chardon à foulon*, *Chardon à bonnetier*, has had similar properties ascribed to it. Both have been also regarded as stomachic.

DIP'SACUS SYLVES'TRIS, from *διψα*, 'thirst,' said to be so called, owing to the leaves being so placed as to hold water. *Cultivated Teasel*, (F.) *Cardère*. *Ord.* Dipsacæ. The roots of this European plant are diuretic and sudorific. The water, which collects at the base of the leaves, has been recommended as an eye-water.

DIPSET'ICOS, from *διψα*, 'thirst,' (F.) *Dip-sétique*. A remedy believed to be capable of exciting thirst. See Alternative.

DIPSO'DES, *Sit'iens*, *Siticulo'sus*, (Old Eng.) *Affort*, (Prov.) *Thrusty*; from *διψα*, 'thirst,' and *ειδος*, 'resemblance.' Thirsty. Causing thirst.

DIPSOMAN'IA, from *διψα*, 'thirst,' and *mania*. Really, 'thirst-mania.' Often, however, applied to habitual drunkenness and to delirium tremens. An insatiable desire for intoxicating liquors.

DIPSOPATHY, *Dipsopath'ia*, from *διψα*, 'thirst,' and *παθος*, 'disease.' A mode of treatment, which consists in abstaining from drinks.

DIPSO'SIS. Same etymon. *Morbid thirst*. The desire for drinking, excessive or impaired. A genus in the class *Caliacea*, order *Entericea*, of Good.

DIPSO'SIS AVENS, Polydipsia—*d. Expers*, *Adipsia*.

DIP'TERIX ODORA'TA, *Coumarou'na odorata*. The fruit of this large tree of Guiana is a pod, inclosing a single seed. The bean, *Tonka bean*, has a strong, agreeable, aromatic odour, and is used to flavour snuff.

DIPY'GUS, from *δι*, 'double,' and *πυγη*, 'nates.' A double monster, the duplication being confined to the posterior portion of the lower end of the trunk—the coccygeal region.

DIPYRE'NON, from *δισ*, 'twice,' and *πυρην*, 'a kernel.' A *specil'um* or probe with two buttons or kernels—one at each end.

DIPYRITES, Biscuit.

DIPYROS, Biscuit.

DIRCA PALUS'TRIS, *Leatherwood*, *Swamp Leatherwood*, *Moosewood*, *Sicamprood*, *Rop-bark*, *Bois de plomb*, (Canada.) *Ord.* Thymelæacæ. An indigenous shrub, which grows in boggy woods and low wet places throughout the United States. It is analogous to mezereon in its action—six or eight grains of the fresh bark producing violent vomiting, preceded by a sense of heat in the stomach, often followed by purging. Applied to the skin, the bark vesicates.

DIRECTOR, *Itinera'trium*, from *dirigere*, *directum*, (*dis*, and *regere*, 'to rule,') 'to direct.' A Conductor. A grooved sound for guiding a knife, in dividing any part; (F.) *Londe cannelée*.

DIRECTOR PENIS, Ischio-cavernosus

DIRIBITORIUM, Mediastinum.

DIRLING, Tingling.

DIRT-EATING, Chthonophagia.

DIRUPTIO, Rhexis.

DIS, Di.

DISC, see Disk.

DISCEPTUM, Diaphragm.

DISCHARGE', (*dis*, and *charge*), *Ec'roë*, *Flucus*, *Proflu'vium*, (F.) *Écoulement*. In pathology, an increased discharge from any part, that naturally secretes a fluid.

DISCHROA, Dyschroëa.

DISCREET', from *discernere*, *discretum*, (*dis*, and *cernere*), 'to separate.' *Discre'tus*, *Intertine'tus*, *Sep'arated*, (F.) *Discret*. This epithet is given to certain exanthemata, in which the spots or pustules are separated from each other. It is opposed to confluent.

DISCRE'T, Discreet.

DISCRE'TA PURGA'TIO. The purgation or expulsion of some particular matter.

DISCRETIVUS, Diagnostic.

DISCRETORIUM, Diaphragm.

DISCRI'MEN, 'Separation, Division.' Same etymon as Discreet. A bandage, used in bleeding from the frontal vein; so called, because, in passing along the sagittal suture, it divides the head into two equal parts.

DISCRIMEN CALVARIE MEDIUM, Diploë—d. Narium, Septum N.

DISCRIMEN NASI; a bandage, in the form of the letter X, intended to support the nose, in transverse wounds of the organ.

DISCRIMEN THORACIS ET VENTRIS, Diaphragm.

DISCUS PROLIGERUS, Proligerous disc—d.

Vitellinus, Proligerous disc.

DISCUNNIES, Discutients.

DISCUSSIO, Resolution.

DISCUSSIVA, Discutients.

DISCUSSORIA, Discutients.

DISCUTIENTS, *Discutien'tia*, *Resolven'tia*, *Discussi'va*, *Discussio'ria*, from *discutere*, 'to shake apart, (*dis*, and *cutere*, 'to shake.')

(F.) *Discussi's*, *Résolutifs*. Substances which possess the power of repelling or resolving tumours.

DISEASE', old French *désaise*; from *dis*, and *ease*: *Morbus*, *Noson*, *Nos'e'ma*, *Noscu'ma*, *Nusus*, *Pathos*, *Path'e'ma*, *Lucs*, *Malum*, *Passio*, *Ægritu'do*, *Ægrotat'io*, *Vit'ium*, *Arrhos'tia*, *Arrhost'e'ma*, *Arrhosten'tia*, *Vale't'do ad'ver'sa*, *Mal'ady*, *Complaint*, *Sickness*, *Distem'per*, *Ailment*, *Illness*, (F.) *Maladie*. An opposite state to that of health, consisting in a change either in the position and structure of parts, or in the exercise of one or more of their functions, or in both.

By some, *Disease* is applied to structural change, whilst *Disorder* is restricted to functional derangement.

The following table, essentially that of Dr. C. J. B. Williams, comprises the chief elements of structural disease:

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

which is harmless. Disinfection also includes the action of antiseptics.

Chlorine or Oxymuriatic Fumigation, Solutio alexiteria Gaubiana, Fumigatio antiloinfugantionensis, is made by adding common salt $\mathfrak{z}\text{ij}$, to black oxide of manganese $\mathfrak{z}\text{j}$, sulphuric acid $\mathfrak{z}\text{j}$, and water $\mathfrak{f}\mathfrak{z}\text{ij}$. This may be carried through an apartment, or be shut up in it.

Nitrous Fumigation, Alexiterium Nitricum, Fumigatio Smythiana, may be formed by placing nitrate of potass $\mathfrak{z}\text{iv}$, and sulphuric acid $\mathfrak{z}\text{ij}$, in a saucer on hot sand.

DISJUNCTI MORBI. (From *dis*, and *jungere, junctum*, 'to join.') *Disjoined diseases.* *Fennelus* has thus denominated diseases which occupy different organs, and are not produced by each other.

DISK, BLOOD, Globule of the blood—*d.* Intervertebral, Intervertebral Cartilage—*d.* Proligerous, see *Proligerous Disk*.

DISLOCATIO LIENIS SEU SPLENIS, Splenoptopia.

DISLOCATION, Luxation.

DISMAL, Melancholy.

DISODICUS, from *dis*, 'twice,' and *odos*, 'threshold.' Having a double opening.

DISOMATOUS, see *Disomus*.

DISOMUS, (F.) *Disome*, from *dis*, and *σωμα*, 'body.' A monster with two bodies. Such a monster is said to be *disom'atous*.

DISORDER, see *Disease*.

DISORDERED, Sick.

DISORGANIZATION, Organo'rum destructio, (F.) *Désorganisation.* A complete morbid change in the structure of an organ, or even total destruction of its texture. In sphacelus and sloughy ulcers, we have examples of this change.

DISPENSARE, Dispensary, Dispensary.

DISPEN'SARY, Dispensa'rium, from *dispendere*, (*dis*, and *pendere, pensus*, 'to weigh,') 'to take from a weight or mass,' 'to distribute.' The shop or place in which medicines are prepared. Also, an institution in which the poor are furnished with advice and necessary medicines. (F.) *Dispensaire.* This word is also used by the French synonymously with *Dispensary*.

DISPENSATION, Dispensa'tio, Diaec'sis, Epineme'sis. Same etymon. The composition and distribution of medicines according to prescription. In France, it is more commonly applied to the weighing, measuring, and arranging of the articles which have to enter into a formula, prior to combining them.

DISPENSATOR, Apothecary.

DISPEN'SATORY, Dispensato'rium; same etymon. *Antidota'rium, Liber Pharmaceuticus*, (F.) *Dispensaire.* A book which treats of the composition of medicines. A *Dispensatory* differs from a *Pharmacopœia*, in containing the physical and medical history of the various substances; whilst the *Pharmacopœia* is mainly restricted to the mode of preparing them. The *Pharmacopœia* too, is published under the authority of, and by the members of a college or association; whilst a *Dispensatory* contains the whole of the *Pharmacopœia* or *Pharmacopœias*, with additions by the author, under whose authority alone it usually appears. The *Pharmacopœia*, in other words, is *official*; the *Dispensatory*, generally, *private*. Formerly, the words were used indiscriminately. See *Formulary*, and *Pharmacopœia*.

DISPLACED, see *Percolation*.

DISPLACEMENT, Percolation.

DISPOSIT'ION, Disposit'io, from *dis*, and *ponere, positum*, 'to put or set.' A particular condition of the body, which renders it susceptible of altering, suddenly, to a state of health or disease;—of improving, or becoming changed for the worse.

The *disposition* to phthisis pulmonalis is sometimes so strong, owing to original conformation, that the disease will make its appearance, notwithstanding every care. See *Diathesis*.

DISPUTATIO, Thesis.

DISQUE PROLIGÈRE, Proligerous Disk.

DISQUES INTERVERTEBRAUX, Intervertebral cartilages—*d. Sanguins, Globules of the blood.*

DISRUPT'IO, from *disrumpere*, (*dis*, and *rumpere*,) 'to break off.' A puncture, which interests deep-seated parts.

DISSECTING ANEURISM, see *Aneurism*.

DISSECTIO, Dissection—*d. Tendinum, Tenotomy.*

DISSECT'ION, Dissec'tio, from *dissecare*, (*dis*, and *secare, sectum*, 'to cut,') 'to cut open;' *Diagnosis, Sectio Anatomica, Practical Anatomy, Necro'tomy.* An operation, by which the different parts of the dead body are exposed, for the purpose of studying their arrangement and structure. Dissection has received various names, according to the organ concerned;—as *Osteotomy, Syndesmotomy, Myotomy, Angiotomy, Neurotomy, Desmotomy, &c.*

DISSECTOR, Prosector, Pro'tomus, same etymon, (F.) *Prosecteur.* A practical anatomist. One who prepares the parts for the anatomical lecture.

DISSEPIPMENTUM NARIUM, Septum narium—*d. Thoracis, Mediastinum.*

DISSEPTUM, Diaphragm.

DISSERTATION, INAUGURAL, Thesis.

DISSIDENTIA, Diastasis.

DISSIPIUM, Diaphragm.

DISSOLUTIO SANGUINIS, Hæmateclysis—*d. Ventriculi, Gastromalacia.*

DISSOLUTION, Dissol'd'tio, Dial'y'sis, from *dissolvere*, (*dis*, and *solvere, solutum*,) 'to loosen,' 'to melt.' This word is in frequent use, in the writings of the humorists. It is employed, particularly, in speaking of the blood,—to designate, not its entire decomposition or putrefaction, but a diminution in its consistence.

DISSOLUTION, Solution.

DISSOLVANTS, Dissolventia.

DISSOLVENTIA, Solvents, (F.) *Dissolvants.* Same etymon. Medicines believed to be capable of dissolving swellings, concretions, &c. *Discutients, Resolvents.*

DISTAD; from *disto*, (*di*, and *sto*,) 'to stand apart.' Away from a centre. Towards the far extremity. In anatomy, used by Dr. Barclay adverbially, to signify 'towards the distal aspect.'

DISTAL ASPECT. An aspect of a bone from the trunk, or towards the extremity.—Barclay.

DISTEMPER, Disease.

DISTEMPERANTIA, Dyscras

DISTEMPERED, Sick.

DISTENSIO, Diastasis, Tension, Tetanus—*d. Nervorum, Convulsion.*

DISTENTION DE LA VÉSICULE DU FIEL, *Turgescencia vesicæ fellæ.*

DISTICH'ASIS, Distich'ia, Districh'ia, Districh'asis, Distac'hia, Distac'h'asis, from *dis*, 'double,' and *στίχος*, 'a row.' An increased number of eyelashes; some turned towards the eye, and irritating the organ; whilst others preserve their natural direction.—Galen, Paulus of Ægina. See *Entropion*.

DISTILLATION, Destilla'tio, Stalag'mos, from *destillare*, (*de*, and *stillare*,) 'to drop, little by little.' *Catantag'mos.* An operation, by which—by the aid of heat and in close vessels—the volatile are separated from the fixed parts of bodies, with the view of collecting the former, when condensed in appropriate receivers. The ancients distinguished distillation *per latius, per ascen'sum*, and *per descen'sum*, according to the direction

which the volatized matters were made to take. In *distillation per latus*, the apparatus is so arranged, that the vapour passes horizontally through a succession of spaces before reaching the receiver. *Distillation per ascensum* is the ordinary method by the still. In *distillation per descensum*, the fire is applied above and around the top of the apparatus; and it is so arranged that the vapour must pass downwards. When dry organic matter is placed in an apparatus for distillation, and heat is applied until all volatile matter is driven out, the process is called *dry or destructive distillation*.

DISTILLATION, DESTRUCTIVE, see Distillation—*d. Dry*, see Distillation—*d. Per Ascensum*, see Distillation—*d. Per Descensum*, see Distillation—*d. Per Latus*, see Distillation.

DISTOCIA, *Distoc'ia*, from *dis*, 'twice,' and *σκος*, 'birth.' Delivery of twins.

DISTECHIA, Distichiasis.

DISTECHIASIS, Distichiasis.

DISTOMA HEPATICOUM, *Dis'tomum hepaticum*, from *dis*, and *στομα*, 'mouth.' *Fasciola hepatica* seu *huma'na* seu *lanceola'ta*, *Planaria latius'cula*, *Gowd-worm*, *Fluke*, *Liver Fluke*, (F.) *Douve*, *D. du Foie*. An obovate, flat worm, an inch in length, and nearly an inch broad; sometimes found in the gall-bladder of man, whence it occasionally passes into the intestinal canal. It is one of the most common varieties of worms, infesting the livers of the sheep, goat, ox, stag, fallow-deer, horse, ass, hog, hare, &c. The treatment is like that required for other worms. **DISTOMA** seu **DISTOMUM LANCEOLA'TUM**, a different species, has been found in the biliary ducts of man,—and of different phytivorous mammalia.—Méhli.

DISTOMA OCULI HUMANI, *Dis'tomum oculi huma'ni*. A parasite once observed in the eye of a child who had suffered from lenticular cataract.—Gescheidt.

DISTOMUM HÆMATOBIUM, see Worms—*d. Hepaticum*, *Distoma hepaticum*.

DISTORSIO, Distortion, Sprain, Strabismus—*d. Oris*, Canine laugh.

DISTORTION, *Distort'sio*, *Intor'sio*, from *distorquere*, (*dis*, and *torquere*, *tortum*), 'to wrest aside.' *Diastrem'ma*, *Dia'strophē*, *Liggy'sia*, (F.) *Distorsion*. Usually applied to the preternatural curvature of a bone; as *distortion of the spine*, *limbs*, &c. It signifies, also, a morbid state of the muscles of the eye, constituting squinting or strabismus.

DISTORTOR ORIS, *Zygomatikus major*.

DISTRIBUTORIA LACTEA THORACICA, Thoracic duct.

DISTRICHIA, Distichiasis.

DISTRICHIASIS, Distichiasis.

DISTRIX, from *dis*, 'double,' and *σπῆ*, 'the hair.' *Trichio'sis distrix*, *Fissu'ra capillo'rum*. Hairs of the scalp, weak, slender, and splitting at their extremities.

DITOCIA, Distocia.

DITRACHYCERAS, from *dis*, 'two,' *τραχυς*, 'rough,' and *κερας*, 'horn.' A genus of intestinal worms. The *Ditrachy'ceras rudis*, *Dic'eras rudis*, (F.) *Bicorne rude*. This fancied entozoon was first observed in the evacuations of a female, by M. Sultz, of Strasburg. It is of a fawn colour, from 3 to 5 lines in length, with an oval head, flattened and terminating in a point posteriorly; contained in a membranous sac, and furnished anteriorly, with a bifurcated, rugous horn. Laënnec considers it a vesicular worm or hydatid. According to Von Siebold, there is no such entozoon. The undigested seed of the mulberry, which had been eaten, was mistaken for it.

DITTANDER, *Lepidium sativum*.

DITTANY, *Conula Mariana*—*d. American*,

Conula Mariana—*d. Bastard*, *Dictamnus albus*—*d. of Crete*, *Origanum dictamnus*—*d. Mountain*, *Conula Mariana*.

DITTECOIA, Double hearing.

DITTOPIA, Diplopia.

DITTOPSIA, Diplopia.

DIURESIS, *Diuresis*, *Diureticæsthesis*, *Ureicæsthesis*; from *διουρειν*, 'to pass the urine,' and *αἰσθησις*, 'feeling.' The desire or want to pass the urine.

DIURESIS, *Diurism*, from *δια*, 'through or by,' and *ουρειν*, 'I pass the urine.' (F.) *Diurèse*. An abundant excretion of urine.

DIURETIC, *Diureticus*, *Ischurætic*, same etymon; *Ureticus*, *Urinatis*. (F.) *Diurétique*. A medicine which has the property of increasing the secretion of urine. Diuretics act by producing a discharge of the watery and other parts of the blood; and, by such discharge, indirectly promote absorption over the whole system. Hence, they are employed in dropsy. The following are the chief Diuretics:—*Cantharis*; *Cantharis Vitata*; *Potassæ Acetas*; *Potassæ Liquor*; *Cahinea*; *Colchici Radix*; *Colchici Semen*; *Digitalis*; *Diosma Crenata*; *Juniperus*; *Oleum Terebinthina*; *Potassæ Nitras*; *Potassæ Bitartas*; *Scilla*; *Sodæ Sales*, and *Spiritus Ætheris Nitrici*.

DIURETICA, *Arnica Montana*.

DIURETICO-ÆSTHESIS, *Diuresæsthesis*.

DIURISM, *Diuresis*.

DIVARIATIO, *Ectropion*.

DIVERSORIUM CHYLI, *Receptaculum chyli*.

DIVERTICULA SPIRITUM ANIMALIUM, *Ganglions*, *nervous*.

DIVERTICULUM, *Deverticulum*. (F.) *Diverticule*. 'A turning; from *divertere*, (*di*, and *vertere*), 'to turn aside.' A blind tube branching out of the course of a longer one. An organ which is capable of receiving an unusual quantity of blood, when the circulation is obstructed or modified elsewhere, is said to act as a diverticulum. Also, a malformation or diseased appearance of a part, in which it passes out of its regular course. It is sometimes applied to such a condition of the alimentary canal. Also, a hole to get out at. A by-passage. See *Ectrope*.

DIVERTICULUM NUCULI. The opening through which the round ligaments of the uterus pass.—Parr.

DIVERTICULUM PHARYNGIS, *Pharyngocele*.

DIVIDING, *Div'idens*, from *dividere*, *divium*, (*di* or *dis*, and the *Hetruscan* verb *iduo*, 'to part or portion.') (F.) *Divisif*. That which divides or separates.

DIVIDING BANDAGE, *Fas'cia div'idens*, (F.) *Bandage divisif*, is a bandage employed for the purpose of keeping parts separated from each other. It is used particularly to prevent improper union; as in cases of burns of the neck or other parts.

DIVINATIO, *Mantia*.

DIVISIF, *Dividing*.

DIVISION, *Divis'io*, *Dia'resis*. Same etymon. The accidental separation of parts naturally united; in other words, a wound or solution of continuity. Most frequently, however, it means an operation, which consists in cutting certain parts, with the view of fulfilling some therapeutic indication.

DIVULSIO, *Dia'spasis*, from *divellere*, (*dis*, and *vellere*, *vulsum*), 'to pull asunder.' A term used in Surgery, to express the rupture or laceration of organs by external violence.

DIVULSIO URINÆ, Cloudiness of urine.—See *Enæorema*.

DIXON'S ANTIBILIOUS PILLS, *Pilule antibiliosæ*.

DIYLISIS, *Colatio*.

DIYLISMUS, *Colatio*.

DIZZINESS, Vertigo.

DIZZY, Vertiginous.

DOCCIONE, MINERAL WATERS OF. This spring is at Lucca, in Italy. It is thermal saline.

DOCH'ME, δοχμή. A measure of the Greeks, equal to about four fingers' breadth: *Dactylo-doch'me*.

DOCIMA'SIA, *Docimas'ticē*, *Docimasilog'ia*, from δοκιμαζω, 'I try or prove the quality of any thing.' The act of assaying.

DOCIMA'SIA MEDICAMENTO'RUM ET VENENO'RUM. The testing of medicines and poisons.

DOCIMA'SIA PULMO'NUM, *d. Pulmona'lis*, *Pneobiomanti'a*, *Pneobiomanti'a*, *Pneobiomanti'a*. Lung proof, Respiration proof. Different proofs to which the organs of respiration of a new-born child are subjected, for the purpose of detecting whether it has or has not respired after birth; in other words, whether it was born alive or dead; — *Pneusiobiogno'sis*, *Pneusiobios'copē*. These consist, chiefly, 1. In testing them with water, for the purpose of seeing whether the lungs are specifically heavier or lighter than that fluid. This is called *Docima'sia Pulmo'num hydrostat'ica* or the *Hydrostatic Test*. If lighter, it would be some evidence that the fœtus had respired: 2. In comparing the weight of the lungs with that of the whole body; the weight of lungs in which respiration has taken place being nearly twice as great. This is *Docima'sia Pulmo'num Stat'ica*, or *Ploucquet's Test*; and, 3. By measuring the circumference of the thorax and lungs; and comparing their dimensions with those of an infant which has not respired. That is *Daniel's Test*. These tests, singly, afford only probable evidence; but when united, the deductions may be more conclusive.

DOCIMASIOLOGIA, *Docimasia*.

DOCK, *Rumex*—*d. Bitter*, *Rumex obtusifolius*—*d. Bloody*, *Rumex sanguineus*—*d. Bloody-reined*, *Rumex sanguineus*—*d. Blunt-leaved*, *Rumex obtusifolius*.

DOCK CRESSSES, *Lapsana*.

DOCK, CURLED, *Rumex crispus*—*d. Sour*, *Rumex acetosus*, *R. crispus*—*d. Sour, boreal*, *Oxyria reniformis*—*d. Spatter*, *Nuphar advena*—*d. Water*, *Rumex hydrolapathum*—*d. Wild, sharp-pointed*, *Rumex acutus*—*d. Yellow*, *Rumex crispus*.

DOCKEN, *Rumex*.

DOCNA SURA, MINERAL WATERS OF. This spring is in the Krapach mountains. The water contains sulphate of soda, carbonate of soda, chloride of sodium, carbonate of lime, silica, and iron.

DOCTEUR, Doctor, Physician.

DOCTOR, *Med'icus*, from *doctus*, 'learned'; 'un homme qui devrait être docte.' *A Physician*. (*Sc.*) *Medicinar*, *Medicinar*. (*F.*) *Docteur*, *Médecin*. Frequently applied to any one who practises medicine; although properly confined to him who has received his degree of Doctor of Medicine. See Physician.

To **DOCTOR** is sometimes used, vulgarly, for to 'treat with drugs.'

DOCTORS' STUFF, see Medicament.

DOCTRINA GALLIANA, Craniology—*d. So-teria*, *Medicina*.

DOCTRINE HOMŒO-ORGANIQUE, Homœopathy.

DOCTRINE, MEDICAL, *Doctri'na Med'ica*. The principles or positions of any medical sect or master. Medicine has been too full of doctrines. One of the first was that of Herodicus of Selivriæ, who recommended gymnastic exercises in disease. The chief founders of doctrines have been Hippocrates, Serapion of Alexandria, Philinus of Cos, Herophilus, Asclepiades, The-mison of Laodicea, Thessalus of Tralles, Soranus of Ephesus, Leonides of Alexandria, Athenæus

of Attalia, Archigenes of Apamæa, Agathinus of Sparta, Galenus, Paracelsus, John Baptist van Helmont, Sylvius de le Boe, Keill, Hamberger, Pitcairne, H. Boerhaave, J. D. Stahl, Frederick Hoffmann, George Baglivi, Cullen, Darwin, John Brown, Beddoes, Girtanner, Reil, Rush, Rasori, and Broussais.

DOCTRINE, PHYSIOLOGICAL, Broussaism.

DODDER, *Cuscuta glomerata*—*d. American*, *Cuscuta glomerata*—*d. of Thyme*, *Cuscuta epithymum*.

DODECADACTYLITIS, Duodenitis.

DODECADACTYLON, Duodenum.

DODECAPHAR'MACUM, from δώδεκα, 'twelve,' and φάρμακον, 'a medicine.' An ancient name given to all medicines which consisted of twelve ingredients. See *Apostolorum Unguentum*.

DODECATH'ËON, from δώδεκα, 'twelve,' and ἔθνη, 'I put.' An antidote, consisting of twelve simple substances.—Paulus of Ægina.

DODECATHËON, *Sanicula*—*d. Plinii*, *Pinguicula vulgaris*.

DODONÆA, *Myrica gale*.

DODONÆA THUNBERGIANA. A shrub of the *Nat. Ord.* Sapindaceæ, which grows at the Cape of Good Hope. A decoction of the root is used as a gentle cathartic in fever.

DODRANS, *Spithama*.

DOGCHOAK, *Cynanche*.

DOGDAYS, (*F.*) *Jours Caniculaires*. During these days, comprised between the 24th of July and the 23d of August, the temperature of the air in Europe is generally high and oppressive. They have received this name from the dogstar, *Canic'ula*, *Σείριος*, *Sir'ius*—a very brilliant star in the constellation of the great dog, which rises and sets, at this time, with the sun. It was formerly believed to be a period particularly unpropitious to health; that it was dangerous to purge during it; and other phantasies were indulged in regard to it.

DOGGRASS, *Triticum repens*.

DOGHIP, see *Rosa canina*.

DOGMATIC, from the Greek δόγμα, *dogmatos*, from δόκεω, 'I think.' *Dogmat'icus*, (*F.*) *Dogmat'ique*. The name of an ancient medical sect; so called, because its members endeavoured, by reasoning, to discover the essence of diseases and their occult causes; whilst the *Empirics*, their rivals, confined themselves strictly to experience; i. e., to the observation of facts. The union of the two modes of investigation makes the rational physician. These sectarians are likewise called *Dog'matists*, and their doctrine *Dog'matism*. The founders of the sect were Hippocrates, Thessalus, Draco, and Polybius; and the most celebrated of its supporters were Diocles of Carysta, Praxagoras of Cos, Chrysippus of Soli, Herophilus, Erasistratus, &c.

DOGS' BANE, *Apocynum androsæmifolium*—*d. Bane*, bitter, *Apocynum androsæmifolium*.

DOGS' GRASS, *Triticum repens*.

DOGS' STONES, *Orchis mascula*.

DOGSTAR, *Canicula*.

DOGTREE, *Cornus Florida*.

DOGWOOD, *Cornus Florida*—*d. Blueberried*, *Cornus sericea*—*d. Female*, *Cornus sericea*—*d. Florida*, *Cornus Florida*—*d. Great flowered*, *Cornus Florida*—*d. Jamaica*, *Piscidia erythrina*—*d. New England*, *Cornus sericea*—*d. Panicle*, *Cornus paniculata*—*d. Pond*, *Cephalanthus occidentalis*—*d. Silky-leaved*, *Cornus sericea*—*d. Virginian*, male, *Cornus Florida*—*d. Round-leaved*, *Cornus circinata*—*d. Striped*, *Acer Pennsylvanicum*—*d. Swamp*, *Cornus sericea*, *Ptelea trifoliata*—*d. White*, *Cornus paniculata*.

DOIGT, *Digitus*—*d. Auricular*, *Auricular finger*—*d. Milken*, see *Digitus*—*d. Petit*, see *Digitus*—*d. Surnuméraire*, *Digitus supernumerarius*.

DOIGTIER, Digitale, Digitalis — *d. d'Asdrualti*, see Digitale.

DOIGTS PALMÉS, Dactylon.

DOITTRIE, see Dementia.

DOKE, Contusion.

DOLABRA, Ascia, *Doloire*.

DOLIC, Dolichos.

DOLICHOCEPHALÆ (GENTES); from *δολχος*, 'long,' and *κεφαλη*, 'head.' 'Longheads.' Nations of men whose cerebral lobes completely cover the cerebellum — as the Kelts, Germans, negroes, &c. Retzius.

DOLICHOCEPHALIA; same etymon. Monstrosity, in which the head is abnormally long.

DOL'ICHOS, from *δολχος*, 'long.' A genus of plants of the leguminous family, (F.) *Dolie*, *Dolique*. It includes a number of species, generally indigenous in India or America. The *Dol'ichos Lablab*, or *Lablab*, is found in Egypt. (Prospero Alpini.) Its fruit is eaten there, like the haricot with us. The *Dol'ichos Stenensis*, is eaten in China, and is stored up as a provision for long voyages. The *Dol'ichos Tuberosus*, (F.) *Pois Putate* of Martinique, has tuberous roots of the size of both fists, and has the consistence and taste of the turnip. The *Dol'ichos Bulbosus* of the West Indies resembles the turnip; — and from the seeds of the *Dol'ichos Soja*, the Japanese prepare the well-known sauce *Soy*, which they term *Sooja*. The most important in medicine is the

DOL'ICHOS PRURIENS, *Dol'ichos*, *Stizolobium*, *Mucuna pruriens*, *Nepretia pruriens*, *Cochage*, *Cowitch*, *Adsarria Pala*, Order, Leguminosæ. (F.) *Pois à gratter*. The stiff hairs of the *Dol'ichos Pods*, called *Dolichos Pubes*, *D. Pruriens* *tis pubes*, *Dolichi Setæ Leguminum*, *Lanu'go Sil'iquæ hirsuta*, *Mucuna* (Ph. U. S.), are the parts used in medicine. They excite an intolerable, prurient sensation, when applied to the skin; but do not irritate the mucous membrane over which they pass, when administered internally. The *Dolichos* is a mechanical anthelmintic, and is useful in cases of ascarides lumbricoides, and oxyures vermiculares. Dose, gr. v to x, of the pubes, in molasses.

Mucuna pruri'ta, a distinct species, but possessing similar properties, grows in the East Indies.

DOLIQUE, Dolichos.

DOLOIRE (F.), *Ascia*, *Dola'bra*, 'a carpenter's axe.' A *Bandage en doloire* is one in which the turns are so placed, that each one covers two-thirds of that which is immediately beneath it. It has received the name *Doloire* from its representing the obliquity of the edge of the instrument whose name it bears. See *Bandage*.

DOLOR, Pain — *d. Ani*, Proctalgia — *d. Capitis*, Cephalalgia — *d. Cephalicus*, Cephalalgia — *d. Colicus*, Colic — *d. Crucians faciei*, Neuralgia, facial — *d. Dentium*, Odontalgia — *d. Dentium à stridore*, Hæmodia — *d. Faciei*, Neuralgia, facial — *d. Ischiadicus nervosus*, Neuralgia femoro-poplitea — *d. Lenis*, Hypodynia — *d. Mitis*, Hypodynia — *d. Nephreticus*, Nephralgia — *d. Pectoris externus*, Pleurodynia — *d. Pudendorum*, Pudendagra — *d. Rheumaticus* et *arthriticus*, Rheumatism.

DOLORES AD PARTUM, Pains, labour — *d. Intestinalium*, Colic — *d. Parturientis*, Pains, labour — *d. Post partum*, Pains, after — *d. Puerperarum*, Pains, after — *d. Rodentes*, Pains, gnawing.

DOLP, Nates.

DOMBEY'A EXCEL'SA, *Araucaria Dombey'i*. A tree of Chili, which furnishes *Dombeya turpentine*; a glutinous, milky-looking fluid of a strong odour and taste.

DOMES'TIC, *Domes'ticus*, from *domus*, 'a house.' The term *Domestic* or *Popular Medicine*, has been given to treatises written for the purpose of enabling those who are not of the pro-

fession to treat diseases, which may occur in their families, without the necessity of calling in a physician. The term, likewise, signifies — Medicine, when thus practised. It is probable, that such works have been attended with mischievous as well as advantageous results.

DOMINARUM AQUA. A medicine described by Myrepsus, which he considered emmenagogue.

DOMPTE-YE'NIN, *Asclepias vincetoxicum*.

DOMUS LEPROSARIA, *Ladrerie*.

DONDO, Albino.

DONESIS, Agitation.

DOORWEED, Polygonum.

DORADILLA, *Asplenium ceterach*.

DORCADIZON, Caprizans.

DOREA, Hæmeralops.

DOREMA AMMONIACUM, see *Ammoniac*, Gum.

DORMITATIO, Somnolency.

DORMITTO, Sleep — *d. Lucumoriana*, see *Lucumorianus*.

DORONIC, *Doronicum pardalianches* — *d. d'Allemagne*, *Arnica montana*.

DORONICUM ARNICA, *Arnica montana* — *d. Cordatum*, *D. Pardalianches* — *d. Germanicum*, *Arnica montana* — *d. Officinale*, *D. Pardalianches* — *d. Oppositifolium*, *Arnica montana*.

DORONICUM PARDALIANCHES, *D. Romanum* seu *Cordatum* seu *Officinale*, *Roman Leopard's Bane*, (F.) *Doronic*, *Doronique*, *Mort aux Panthères*. The root of this plant resembles *Arnica Montana* in its properties.

DORONICUM PLANTAGINIS FOLIO, *Arnica montana* — *d. Romanum*, *D. Pardalianches*.

DORSAD, see *Dorsal Aspect*.

DORSAL, *Dorsalis*, from *dorsum*, 'the back.' *Notiæ's*, *Notal*, *Tergal*. Relating to the back of the body, or of one of its parts; as the *Dorsal vertebrae*, nerves, &c.; *Dorsal artery of the tongue*, penis, &c.; *Dorsal region of the foot*, hand, &c.; *Dorsal Consumption*, &c.

DORSAL ASPECT. An aspect towards the *dorsum* or backbone. — Barclay. *Dorsad* is used by the same writer adverbially, to signify 'towards the dorsal aspect.'

DORSAL, LONG, *Longissimus dorsi*.

DORSALIS, *Dorsal*.

DORSALIS PENIS, (Nervus.) The branch of the internal pudic nerve, which is distributed to the upper part of the male organ; and to the clitoris of the female.

DORSCH, see *Oleum jecoris aselli*.

DORSE, see *Oleum jecoris aselli*.

DORSO-COSTAL, *Serratus posticus superior* — *d. Lombo-costal*, *Serratus posticus inferior* — *d. Lombo-sacro-humeral*, *Latissimus dorsi* — *d. Susacromien*, *Trapezius* — *d. Trachelon-occipital*, *Complexus*.

DORSTENIA BRASILIENSIS, *Caa-apia*.

DORSTENIA CONTRAYER'VA, called after Dr. Dorsten; *d. Housto'nii*, *Contrayer'va*, *Drake'na*, *Cyp'rus longus odor'us* seu *Peru'rus*, *Bezoar'dica Radix*, *Lisbon Contrayer'va*, (F.) *Racine de Charcis* ou de *Dracke* ou des *Philippines*. Order, *Urticæ*. *Sex. Syst.* Tetrandria Monogynia. A plant of South America, whose root, *Contrayer'va* (Ph. U. S.), is aromatic, bitter, and astringent. It has been given as a tonic, stimulant and sudorific. Dose, gr. xij to ʒss.

DORSTENIA CORDIFOLIA, *D. Brasiliensis* — *d. Placentoides*, *D. Brasiliensis* — *d. Vitella*, *D. Brasiliensis*.

DORSUM, *Notos*, *Noton*. The back. *Metaph'renon*, *Tergum*, (Prov.) *Rig*, (F.) *Dos*. The posterior part of the trunk, extending from the inferior and posterior region of the neck as far as the loins. The back of the foot, *Dorsum pedis*, is the upper part of that organ, opposite the sole: the

back of the hand, *Dorsum manûs*, the part opposed to the palm. In the same sense, we say *Dorsum penis*, *Dorsum nasi*, for the upper part of those organs.

DORSUM, Vertebral column—*d. Ephippii*, see *Sella turcica*—*d. Manûs*, *Opisthenar*.

DORYCIMUM, *Costus creticus*.

DOS, Dorsum.

DOSAGE, Posology.

DOSE, *Dosis*, *Præbium*. The quantity of any substance, which ought to form part of a compound medicine, or ought to be exhibited singly, to produce a desired effect. Many circumstances influence the doses of medicine. Women require smaller doses, as a general principle, than men. *Habit* has a great effect, as well as *climate*, *age*, and *idiosyncrasy*: all these, and other circumstances, must be taken into account; and every general rule on the subject will be found to have numerous exceptions. Some of the mechanical physicians laid it down as a rule, that the doses of medicines must always be as the square of the constitution!—A matter not easy of calculation.

The following Tables will exhibit an approximation to the proper doses (according to age) of most substances.

TABLE OF DOSES ACCORDING TO AGE.

Age	Let the full dose be	1	1 drachm.
24	will require	2-3ds.	2 scruples.
18	" " " "	half.	$\frac{1}{2}$ drachm.
14	" " " "	1-3d.	$\frac{1}{3}$ scruple.
7	" " " "	1-4th.	15 grains.
4	" " " "	1-6th.	10 grains.
3	" " " "	1-8th.	8 grains.
2	" " " "	1-12th.	5 grains.
1	" " " "		

The table of doses, according to age, recommended by Dr. Thomas Young, differs in some respects from the above. Either affords a sufficient general approximation. His rule is, that

For children, under twelve years of age, the doses of most medicines must be diminished in the proportion of the age to the age increased by twelve:

$$\text{Thus, at two years, to 1-7th; i. e., } 1-7\text{th} = \frac{2}{2 + 12}$$

At twenty-one the full dose is given.

DOSES, BROKEN. When an agent is given in small portions it is said to be in *broken doses*,—*refractis dosibus*.

DOSIOLOGIA, Posology.

DOSIS, Dose.

DOSSIL, *Bourdonnet*.

DOTAGE, Dementia.

DOTHEN, *Furunculus*.

DOTHIENTENTERIA, *Dothinenteritis*.

DOTHIENTÉRIE, *Dothinenteritis*.

DOTHIENTERIA, *Dothinenteritis*.

DOTHIENTERITIS, properly *Dothiententeritis*, *Dothiententeria*, *Dothinenteria*, *Enteritis pustulosa*, *Enterodothientia*, *Helcenteritis*, *Follicular Gastroenteritis*, from *dothiv*, 'a pustule,' and *enteron*, 'intestine.' (F.) *Dothientérie*, *Dothientérie*, *Dothientérite*. An inflammation and ulceration of the glands or follicles of Peyer and Brunner, which Bretonneau considers to be the essence of a large class of fevers, particularly of those of the typhoid character. See Typhus.

DOTHEN, *Furunculus*.

DOTTRINA MEDICA ITALIANA, *Contro-stimulus*, (doctrine of.)

DOUBLE, *Ingluvies*.

DOUBLE-CONSCIOUSNESS, see *Consciousness*, double.

DOUBLE-HEARING, *Dipleco'ia*, *Dittec'o'ia*, *Parac'o'ia duplica'ta seu imperfec'ta*. The action of the one ear unaccordant with that of the other: sounds heard doubly, and in different tones or keys.

DOUBLE-MONSTERS, see *Duplication*.

DOUBLE-QUARTE, see *Quartan*.

DOUBLE-QUOTIDIENNE, see *Quotidian*.

DOUBLE-TIERCE, see *Tertian*.

DOUCE-AMÈRE, *Solanum dulcamara*.

DOUCHE (F.). In Italian, *doccia*: modern Latin, *ducia*; *Cataclysmus*, *Douche*. This term is applied to a column of fluid, of a determinate nature and temperature, let fall upon the body. *Pumping* is a variety of the *Douche*. In using this kind of bath, the fluid is directed upon the part on which we are desirous of acting. The *douches descendantes* are those in which the fluid falls from a height,—the *douches ascendantes*, those administered in diseases of the uterus,—the *douches horizontales*, where the fluid is impelled horizontally, &c. They may be cold or warm, according to circumstances. The apparatus consists of a reservoir of water having a pipe or plug, by means of which the water can be directed as the practitioner may desire. The *Douche* communicates a considerable and peculiar shock to the nervous system; and is one of the most successful means for taming the furious maniac. It is, also, useful in chronic rheumatism, stiff joints, &c.

Douches of air are, also, occasionally used, as in cases of obstruction of the Eustachian tube by mucus. They are sent from an air-press—of which Deleau and Kramer have invented one each—through a catheter introduced through the nose into the tube.

DOULEUR, Pain—*d. de Côté*, *Pleurodynia*—*d. des Dents*, *Odontalgia*—*d. de l'Estomac*, *Cardialgia*—*d. Névralgique de l'Estomac*, *Cardialgia*—*d. Pulsative*, see *Throbbing*.

DOULEURS, Pains, labour—*d. Conquassantes*, see *Conquassant*.

DOUNT, *Palpitation*.

DOUP, Nates.

DOUSE, *Douche*.

DOUVE, *Distoma hepaticum*—*d. du Foie*, *Distoma hepaticum*—*d. Petite*, *Ranunculus flammula*.

DOWN, see *Attack*.

DOWP, Nates.

DOYLE, *Strabismus*.

DRACHION, *Pugillus*.

DRACHM, *Drachma*, *Dram*, (F.) *Gros*. The ancient name of a piece of money, weighing the eighth part of an ounce. At the present day it is used for this weight.

DRACO MITIGATUS, *Hydrargyri submuriæ*—*d. Sylvestris*, *Achillea ptarmica*.

DRACOCEPH'ALUM CANARIEN'SÉ, *D. Moldavicum*, from *δρακων*, 'a dragon,' and *κεφαλή*, 'head,' from the shape of the corolla, *Melissa Turcica seu Canariensis*, *Cedronella triphylla*, *Alpinia Balsamum*, Turkey or *Cana'ry Balsam*, *Balm of Gil'ead Tree*, (F.) *Melisse de Moldavie*. Ord. Labiate. A Turkish and Siberian plant, which has an aromatic taste, joined with an agreeable flavour. It has been used as a tonic.

DRACONTHÈMA, see *Calamus rotang*.

DRACONTIUM, *Dracunculus*. See, also, *Dracuncium foetidum*.

DRACONTIUM FÆTIDUM, *Dracontium*, from *δρακων*, 'a dragon,' whose tail the roots resemble, *Ictodes fætidus*, *Symplocarpus fætidus*, *Pothos fætida seu Puto'ia*, *Arum Americanum*

beta folio, Spathyema foetida, Skunk-cabbage, Skunk-weed, Polecat-collard, Cow-collard, Collard, Ich-weed, Stink-poke, Nicotiana-cabbage, Polecat-weed, Hel'ebore, El'ebore, Irish cabbage. Ord. Aroidæ or Araceæ. *See. Syst. Tetrandria Monogynia.* This indigenous plant, as some of its names import, is extremely fetid. The property on which its medical virtues are dependent, resides in a volatile principle, which is impaired by long keeping, especially in powder. Dose, of the dried root, *Dracontium*, (Ph. U. S.), ten to twenty grains. It resembles asafetida and other fetid gums in its properties; in other words, belongs to the class of reputed antispasmodics.

DRACONTIUM ANGUSTIS-PATHA, Symplocar'pus Angustis'patha, Narrow-spathed Skunk-cabbage, is possessed of similar properties.

DRACUNCULUS, diminutive of δράκων, 'a dragon or serpent,' *D. Gor'dius, Dracontium, Vena Medinæ Ar'abum, Vena seu Gor'dius Medinensis, Medinogor'dius, Vermiculus Capilla'ris, Filic'ria Medinensis seu Guineensis, Muls Dracontulus seu Gor'dii, Helminthioncus Medinensis, Muscular Hairworm, Bichios, Bicho, Guinea Worm or Threadworm, (F.) Dragonneau, Ver de Guinée, Ver Filaire, V. de Méline, V. cutané, Veine ou Filaire de Médine.* A genus of worms, frequently met with in Indian and African climes. They are characterized by a filiform body, and are smooth and nearly of equal thickness throughout. The Guinea worm, when small, insinuates itself through the cutaneous pores, and penetrates into the areolar membrane and between the muscles; especially between those of the lower limbs, where it occasions a tumour like a boil, which subsequently suppurates, and the head of the worm appears and emerges gradually. The head must then be seized, and the worm be cautiously rolled round a small cylinder of linen or other substance. Care must be taken not to break it, as great pain and obstinate suppuration might be the consequence.

Considerable obscurity rests on this subject. Some even deny that the cases of *Dracunculus*, on record, are really those of worms.

DRACUNCULUS POLYPHYLLUS, *Arum dracuncul—d. Pratensis, Achillea ptarmica.*

DRAGÉES, (F.) Almonds or dried preserves, covered with white sugar; *Sugar-plums.*

DRAGÉES DE KEYSER. A pharmaceutical preparation, formerly much celebrated in syphilis. It was composed of acetate of mercury, menna, starch, mucilage and gum tragacanth.

DRAGMA, Pugillus.

DRAGMIS, Pugillus.

DRAGON (F.), *Dragon.* Some authors have given this name to opaque spots on the cornea, and to cataract.

DRAGON CLAW, SCALY, Pterospora andromedea—d. Green, Arum dracontium—d. Root, Arum triphyllum, Pterospora andromedea—d. Turnip, Arum triphyllum.

DRAGON'S BLOOD, see *Calamus rotang.*

DRAGONNEAU, *Dracontulus.*

DRAINING, *Desiccatio, (F.) Dessèchement.* The act of drawing off the water from marshes, for the purpose of destroying the unhealthy emanations which proceed from them. It is a well known fact, that from marshes arise certain emanations or miasmata, with the nature of which we are, however, totally unacquainted, but which are the fertile source of intermittents and other diseases. Draining the soil and converting it into arable land changes its character, and the malaria ceases to be given off. It has happened, however, that although in some such situations intermittents have been got rid of, consumptions have taken their place.

DRAITSCH WATER, Godesberg, mineral waters of.

DRAKENA, *Dorstenia contrayerva.*

DRAM, Melancholic.

DRAPEAU (F.), *Vexil'um*, from *drap*, 'cloth.' A bandage, used in wounds, &c., of the nose, to keep the dressings *in situ*. It is composed of a small triangular compress, having two openings at its inferior part, corresponding to those of the nostrils. It is fixed by means of a bandage, passing from its superior angle over the head, and another passing round the head, under the orbits, so as to join the first at the nape of the neck. Also, *Pterygion.*

DRAP FANON, see *Fanon.*

DRASTIC, *Dra'sticus, Cenot'ic, (F.) Drastique*, from *δραω*, 'I operate strongly.' Active. An epithet given to purgatives, which operate powerfully; as *elaterium*, oil of *croton*, &c.

DRAUGHT, *Haustus.* A term also applied, by nurses, to the sudden rush of blood to the mammae, occasioned in the mother by the sight, or even thought, of her infant, and which occasions a greatly-increased secretion of milk.

DRAUGHT, BLACK, see *Infusum sennæ composition.*

DRAWING, see *Digestives.*

DRAX, Pugillus.

DREAM, *Somnium—d. Waking, Hallucination.*

DRÊCHE, Malt.

DREGISTER, Druggist.

DREGS, *Feculence.*

DRENCH, (Sax. *spencean*, 'to soak.') To give liquid physic by force. A liquid medicine, given by violence, is called "a *drench*."

DREPANOIDES, *Falciform.*

DRESS, *Vestitus.*

DRESSER, from (F.) *dresser*, 'to put right.' An hospital assistant, whose office it is to dress wounds, ulcers, &c. He corresponds in function to the *Élève externe* of the French hospitals.

DRESSING, *Cura, Cura'tio, (F.) Pansement.* Same etymon. The methodical application of any remedy or apparatus to a diseased part. Also, the remedy or apparatus itself.

DRIBURG, MINERAL WATERS OF. At the small town of Driburg, near Pymont, in Germany, there are nine springs, which are rich in saline ingredients, iron, and carbonic acid gas. Their action on the economy is like that of Pymont water. They contain chloride of sodium, sulphate of magnesia, sulphate of lime, carbonate of lime, carbonate of magnesia, carbonate of iron, chloride of calcium, and chloride of magnesium.

DRIMYPHA'GIA, from *δριμυς*, 'acid,' 'aromatic,' and *φαγω*, 'I eat.' An aromatic and exciting diet.

DRIMYS WINTERI, *Winters aromatica.*

DRINK, from Sax. *spencean, Poma, Potio, Potus, Ber'eage, (F.) Boisson.* Every liquid introduced into the alimentary canal for the purpose of repairing the loss of the fluid parts of the body. The necessity for its use is indicated by the sensation of thirst. Fluid, taken during a meal, aids in the digestion of the food. Some drinks are exciting and tonic, as the different varieties of beer, wine, and spirits, which we use at table. In a therapeutical point of view, drinks are used to appease the thirst which prevails in febrile affections, or to act as diluents in those and other cases.

The ordinary drinks, according to their chemical composition, are—1. *Water*, spring water, river water, well water, &c.—2. *Juices and infusions of Animal and Vegetable substances*, lemon juice, currant juice, whey, tea, coffee, mattee, &c.—3. *Fermented Liquors*, wines, ale, beer, cider, perry.—4. *Spirituous Liquors*, brandy, alcohol,

ether, kirschwasser, rum, arack, gin, whisky, ratafas, cordials, &c.

DRIVELLING, Slaverling.

DROAT, Throat.

DRODDUM, Nates.

DROGUE, Drug.

DROGUIER (F.). Same etymon as drug. A collection of different simple medicinal substances. A methodical collection of specimens of drugs.

DROQUISTE, Druggist.

DROICH, Nanus.

DROIT, Rectus—*d. de l'Abdomen*, Rectus abdominis—*d. Antérieur de la cuisse*, Rectus femoris—*d. Antérieur de la tête*, Rectus capitis internus minor—*d. Antérieur de la tête, grand*, Rectus capitis internus major—*d. Externe de l'œil*, Rectus externus oculi—*d. Inférieur de l'œil*, Rectus inferior oculi—*d. Interne de la cuisse*, Gracilis—*d. Interne de l'œil*, Rectus internus oculi—*d. Latéral de la tête*, Rectus capitis lateralis—*d. Postérieur de la tête, grand*, Rectus capitis posterior major—*d. Postérieur de la tête, petit*, Rectus capitis posterior minor—*d. Supérieur*, Rectus superior oculi.

DROP, Gutta—*d. Ague*, tasteless, Liqueur arsenicalis—*d. Black*, Guttae nigrae—*d. Red*, Ward's, see Vinum antimonii tartarizati—*d. Serene*, Amaurosis.

DROPACISMUS, Depilation.

DROPAX, Depilatory.

DROPPED HANDS, see Hands, dropped.

DROPS, ABBÉ ROUSSEAU'S, Landanum Abbatis Rousseau—*d. Anodyne*, Liqueur morphinæ acetatis—*d. Antiscorbutic*, Marsden's, see Infusum gentianæ compositum—*d. Anti-venereal*, see Tinctura ferri muriatis—*d. Chamomile*, see Anthemis nobilis—*d. Dutch*, Balsam of sulphur, Balsam of Turpentine—*d. Fit*, Spiritus ammoniæ fetidus, see Fuligo—*d. Golden*, de la Motte's, Tinctura seu alcohol sulfurico-æthereus ferri—*d. Green's*, see Liqueur Hydrargyri oxymuriatis—*d. Jesuit's*, Tinctura benzoini composita—*d. Lavender*, Spiritus Lavandulæ compositus—*d. Nitre*, Spiritus ætheris nitrici—*d. Norris's*, see Antimonium tartarizatum—*d. Norton's*, Liqueur Hydrargyri Oxymuriatis—*d. Pectoral*, Pectoral drops, Bateman's—*d. Peppermint*, Pastilli de menthâ piperitâ—*d. Soot*, see Fuligo—*d. Wade's*, Tinctura benzoini composita—*d. White*, Ward's, see Hydrargyri nitratis.

DROPS, WARBURG'S. A secret preparation used in Demarara under the name of Warburg's Fever Drops. See Bebeeru.

DROPSICAL, Hydropic.

DROPSIED, Hydropic.

DROPSY, Hydrops—*d. of the Belly*, Ascites—*d. of the Brain*, Hydrocephalus.

DROPSY, CARDIAC. Hydrops cardiacus, (F.) *Hydropisie cardiaque*. Dropsy, dependent on disease of the heart.

DROPSY OF THE CELLULAR MEMBRANE, Anasarca—*d. of the Chest*, Hydrothorax—*d. of the Eye*, Hydrophthalmia.

DROPSY, FIBRINOUS. Dropsy in which the effused fluid contains fibrin.

DROPSY, GENERAL, Anasarca—*d. of the Head*, Hydrocephalus.

DROPSY, HEPATIC, Hydrops hepaticus, (F.) *Hydropisie hépatique*. Dropsy, dependent on disease of the liver.

DROPSY OF THE LACHRYMAL SAC, Fistula lacrymalis, Lachrymal hernia—*d. of the Pericardium*, Hydropericardium—*d. of the Peritoneum*, Ascites—*d. of the Pleura*, Hydrothorax.

DROPSY, RENAL, Hydrops renal'is, (F.) *Hydropisie rénale*. Dropsy, dependent on disease of the kidney.

DROPSY, WATER OF, Serum of Serous membranes—*d. Wind*, Emphysema.

DROPSY OF THE WOMB, Hydrometra.

DROWORT, Spirea filipendula—*d. Hemlock*, Ceanothe—*d. Water*, Hemlock, Ceanothe—*d. Western*, Gillenia trifoliata.

DROPYK, Hydrops.

DROS'ERA ROTUNDIFO'LIA, from *ῥοσος*, 'dewy,' *ῥοσος*, 'dew,' the leaf glands exuding a clear fluid like dew drops. The systematic name of the *Sundew*, *Ros solis*, *Drosiobotanon*, *Drosium*, *Rorella*, (F.) *Rossolis*, *Rosée du soleil*, *Herbe à la rosée*. Ord. *Droseraceæ*. This plant has a bitter, acrid, and caustic taste. It has been used as a rubefacient, and to destroy warts and corns. It has, also, been regarded as a pectoral.

DROSIOBOTANON, *Drosera rotundifolia*.

DROSISTICON, see Bosa.

DROSIMUM, *Drosera rotundifolia*.

DROSOMELI, *Fraxinus ornus*.

DROW, Syncope.

DRUG, *Medicamentum*, (F.) *Droque*. A name ordinarily applied to simple medicines, but, by extension, to every substance employed in the cure of disease. Ménage derives it from *droga*, and this from the Persian *droa*, 'odour,' because many drugs have a strong odour. It is, doubtless, from the Teutonic *trocken*, Sax. *trigan*, 'to dry.'

DRUG. To prescribe or administer drugs. Most commonly, perhaps, to dose to excess with drugs. One who so doses is sometimes called "a druggier." "To drug," also means to tincture with some medicinal article.

DRUG DISEASE. A morbid condition, which is—or is presumed to be—caused or kept up by the administration of drugs.

DRUG-GRINDER, see Pulverization—*d. Mill*, see Pulverization.

DRUGGER, Druggist, see Drug.

DRUGGIST, *Drugger*, *Drugster*, (Prov.) *Dregister*. Same etymon. *Pharmacopola*, *Materialista*, *Seplasia'rius*, *Pharmacopæus*, (F.) *Drugiste*. One who sells drugs.

DRUGSTER, Druggist.

DRUM, Melancholic.

DRUM OF THE EAR, Tympanum.

DRUM-SIEVE, see Cribration.

DRUNKENNESS, Temulentia.

DRUNKWORT, *Nicotiana tabacum*.

DRYING, *Desiccation*, *Sicca'tio*, *Desicca'tio*, *Exsicca'tio*, *Xeran'sis*, *Auan'sis*, same etymon as Drug, (F.) *Dessiccation*. Evaporation or removal of the superfluous humidity in a body.

DRYING OF PLANTS, *Desicca'tio herbarum*, *Desicca'tion of Plants*. Those which are very succulent should be dried quickly. They must be spread upon lattice work, covered from the light, and exposed to the heat of the sun or to that of a stove, not exceeding 110° Fahrenheit. The drying of less succulent plants can be effected at a lower temperature. *Flowers* must be dried very carefully, excluding light. Seeds are dried in a place where there is a free current of air. *Pulpy fruits* may be exposed to a gentle heat in a stove, which may be gradually elevated. *Roots* may also be dried in the stove: the tuberous require to be cut in slices.

DRY NURSE, see Nurse

DTHOKE, *Frambesia*.

DUALITY, *Dual'itas*, from *duo*, 'two.' The state or quality of being two.

DUALITY OF THE MIND, OR BRAIN. As the organ consists of two hemispheres, they have been regarded by some as separately and distinctly concerned in the mental and moral manifestations.

DUALM, Syncope.

DUCKFOOT, see Kylosis.

DUCKLEGGED. Having short legs like a duck.

DUCKSFOOT, *Podophyllum montanum*.

DUCT, ALIMENTARY, Canal, alimentary, Thoracic duct.

DUCT OF BELLINI, Uriniferous tube.

DUCT, NASAL, Lachrymal duct—d. Pancreatic, see Pancreas.

DUCT OF PECQUET, Thoracic duct.

DUCT, VITELLINE, see *Vesicula umbilicalis*.

DUCT OF WIRSUNG, see Pancreas.

DUCTIO PRÆPUTII, Masturbation.

DUCTLESS GLANDS, see Ganglion.

DUCTOR CANALICULATUS, Gorget—d. Urinae, Ureter.

DUCTULI RECTI, see Rete testis.

DUCTUS, Canal, Meatus—d. Arteriosus, Arterial duct—d. Auris palatinus, Eustachian tube.

DUCTUS BARTHOLINIANUS, BARTHOLIN'S Duct, (F.) *Canal de BARTHOLIN*. The excretory duct of the sublingual gland.

DUCTUS BELLINIANI, Uriniferous tubes—d. Biliarii, Pori biliarii—d. Bifleri, Pori biliarii—d. Botalli, Arterial duct—d. Choledochus, Choledoch duct—d. Chyliferus, Thoracic duct—d. Cibarius, Canal, alimentary—d. Excretorius, Excretory duct—d. Ferreini, Ferrein, canal of—d. Hepaticus, Hepatic duct—d. Incisivus, Palatine duct (anterior)—d. Intestinalis, Canal, alimentary—d. Lacteus, Thoracic duct—d. Lacrymalis, Lachrymal duct—d. Lacrimalis, Lachrymal duct—d. Nasalis orbitæ, Lachrymal or nasal duct—d. Nasalo-palatinus, Palatine duct (anterior)—d. ad Nasum, Lachrymal or nasal duct—d. Nutritii, Canals, nutritive—d. Omphalo-mesentericus, see *Vesicula umbilicalis*—d. Pancreaticus, see Pancreas—d. Pecqueti, Thoracic duct—d. Pneumaticus, see Air-bladder—d. Punctorum lacrymalium, Lachrymal ducts—d. Riviniani, see Sublingual gland—d. Rorifer, Thoracic duct.

DUCTUS SALIVARIS INFERIOR, *D. Whartoniana*, (F.) *Canal de WARTHON*. The excretory duct of the submaxillary gland.

DUCTUS SALIVARIS SUPERIOR, *D. Stenoniana*, STENO'S Duct, (F.) *Canal de STENON*. The excretory duct of the parotid gland which opens into the mouth opposite the second upper molar tooth.

DUCTUS SEMI-CIRCULARES LABYRINTHI, Semi-circular canals—d. Serosi, Lymphatic vessels—d. Spirales cochleæ, Scalæ of the cochlea—d. Stenonianus, Ductus salivaris superior—d. Thoracicus, Thoracic duct—d. Urinae, Ureter—d. Urinarius, Urethra—d. Uriniferi Bellini, Uriniferous tubes—d. Varicosi uteri, Tubæ Fallopiannæ—d. Venosus, Canal, venous—d. Vitellarius, see *Vesicula umbilicalis*—d. Vitello-intestinalis, see *Vesicula umbilicalis*—d. Waltheriani, see Sublingual gland—d. Whartonianus, Ductus salivaris inferior—d. Wirsungianus, see Pancreas.

DUELECH, Dulech.

DUEL/LA. The ancient name of a weight, which was equivalent to eight scruples.

DUELLUM VENEREUM, Coition.

DUG, Nipple.

DULCAMARA, *Solanum dulcamara*—d. Flexuosa, *Solanum dulcamara*.

DULCEDO AMORIS, Clitoris—d. Sputatorium, see Salivation.

DULECH, *Duelech*. A term employed by Paracelsus and Van Helmont to designate a pretended tartarized substance, which forms in the human body, and produces acute pain, accompanied with great danger.

DULL, Deaf.

DULSE, *Rhodomela palmata*—d. Pepper, *Laurentia pinnatifida*.

DCMB, see Mutitas.

DUMBNESS, Mutitas.

DUNBLANE, MINERAL WATERS OF. The springs of Dunblane, in Scotland, contain sulphate of soda, chlorides of sodium and calcium, and carbonate of iron.

DUNG, Fimus—d. Cow's, Fimus Vaccæ—d. Goose, Fimus anseris—d. Stone-horse, Fimus equinus.

DUNGA, Dengue.

DUNT, Palpitation.

DUODENAL, *Duodena'lis*, (F.) *Duodénal*. Relating or appertaining to the duodenum.

DUODENITIS, *Dodecadactylitis*, (F.) *Duodénite*. A hybrid term, from *duodenum*, and *itis*, denoting inflammation. Inflammation of the duodenum, characterized by white tongue, bitter taste, anorexia, fulness and tenderness in the region of the duodenum, and often yellowness of skin, along with the ordinary signs of febrile irritation.

DUODENUM, from *duodeni*, 'twelve.' *Ventriculus Succenturiatus*, *Ec'physis* seu *Appendix* seu *Processus Ventriculi*, *Portona'rium*, *Dodecadactylon*, (from *dōdeka*, 'twelve,' and *δάκρυλος*, 'a finger.') The duodenum is the first part of the intestinal canal, commencing at the pyloric orifice of the stomach, and terminating in the jejunum. Its length is about twelve fingers' breadth, and as it is only partially covered by the peritoneum, it is susceptible of considerable dilatation; hence its name *Ventriculus succenturiatus*. In the duodenum, *chylification* takes place after the admixture of the biliary and pancreatic fluids with the chyme.

DUO-STERNAL. Bécларd thus names the second osseous portion of the sternum, which corresponds to the second intercostal space.

DUPLICATION, (*duo*, 'two,' and *plicare*, *plicatus*, 'to fold.') *Duplicatio*, *Duplic'itas*, *Diplasias'mus*, *Diplo'sis*, (F.) *Duplicité*. A species of malformation or monstrosity, characterized by the parts concerned being doubled. See *Diplogenesis*.

DUPLICATION OF THE FÆTUS, Evolution, spontaneous.

DUPPLICATION, *Duplicatu'ra*, *Reflec'tion*. Same etymon. The folding or reflection of a membrane upon itself; as *Duplication* or *reflec'tion of the pleura, peritoneum, &c.*

DUPPLICITAS MONSTROSA, see *Diplogenesis*.

DUPPLICITÉ, Duplication.

DUPONDIUM. A weight of four drachms.—Galen.

DUR, Hard.

DURA MATER, *Crassa* seu *Dura meninx*, *Sclerome'ninx*, *Meninx exte'rior* seu *sclera* seu *dermato'des* seu *pache'ia*, of Galen, *Dura membr'na cere'brum am'biens*, *Cuticu'laris membr'na*, *Méninge*, (Ch.) It has been called *dura*, because of its great resistance; and *mater*, because it was believed to give rise to every membrane of the body. (F.) *Dure mère*, *Dure-tête* (Paré) It is a fibrous, semi-transparent membrane, of a pearly-white colour, thick, and very resisting; lines the cavity of the cranium, and contains the spinal marrow; protects the brain and marrow; and, by its various expansions—the *falz cerebri*, *tentorium*, &c.,—supports the different parts of the cerebral mass. The largest artery of the dura mater is the *A. meningea me'dia*.

DURA MATER, LATERAL PROCESSES OF THE, Tentorium—d. m. Testis, Albuginea.

DURATION OF LIFE, see Life—d. of Pregnancy, see Pregnancy.

DURE-MÈRE, Dura mater.

DURE-TAYE, Dura mater.

DURÉE DE LA VIE, see Longevity.

DURETÉ D'OREILLE, Baryecolia—*d. de l'Oie*, Deafness.

DURILLON, Callosity.

DURUS, Hard.

DUSTING BAG. In pharmacy, a kind of sieve, which consists of a bag made of lawn or other like material, hung inside of a wide-mouthed bottle or tin canister, to the mouth of which it is secured. The powdered substance is put into the bag, and the mouth being closed with a cover, the apparatus is shaken, and the finer particles pass into the bottle or canister.

DUTCHMAN'S PIPE, *Aristolochia hirsuta*.

DUTROA, *Datura stramonium*.

DWALE, COMMON, *Atropa belladonna*.

DWARF, Nanus—*d. Flag*, *Iris lacustris*.

DWARFISH, see Nanus.

DWARFISHNESS, Nanosomus.

DYCTOIDES, Reticular.

DYERS' BROOM, *Genista tinctoria*—*d. Weed*, *Genista tinctoria*, *Reseda luteola*.

DYNAMETER, Dynamometer.

DYNAMIC, *Dynamical*, *Dynam'ic*, (F.) *Dynámique*. Same etymon. In Mechanics, *Dynamics* investigates the powers whereby bodies are put in motion, and the laws connected therewith. In Biology, that which relates to the vital forces, to the organism in action, *Vital Dynamics*. The influences of agents on the organism, which are ascribable to neither mechanical nor chemical causes, are sometimes termed *dynamic*. In this sense, it is a favorite term with the homœopathists, who maintain that, by certain processes, called, in the aggregate, *dynamization* and *potentizing*, the dynamic powers of a medicine may be set free and developed—as by shaking the bottle in which the article is contained, or by mixing, an unlimited number of unmedicated globules with one that is medicated, and shaking them together. In this way, the former become *potentized*!

DYNAMIC DISEASES, see Organic.

DYNAMICS, VITAL, see Dynamic.

DYNAMIUS, Dynamic.

DYNAMIS, Faculty, Force.

DYNAMIZATION, see Dynamic.

DYNAMOMETER, *Myodynamom'eter*, *Myodynamom'eter*, (F.) *Dynamomètre*, from *δυναμις*, 'force,' 'vital power,' and *μετρον*, 'measure.' An instrument, contrived by M. Regnier, for measuring the comparative muscular strength of man and animals. It consists of a spring, which, when pressed upon, causes a needle to move upon a portion of a circle, furnished with a scale of kilogrammes and one of myriagrammes. To measure the strength of the hands, the two branches of the spring are firmly grasped, and brought as near together as the force can carry them. This effort makes the needle traverse, and indicates, on the scale of kilogrammes, the strength of the experimenter's hands. A man, 25 or 30 years of age, exerts a force commonly equal to 50 kilogrammes or 100 pounds.

The strength of the loins of a man, about 30 years of age, as indicated by this instrument, is usually about 30 myriagrammes, or 265 pounds, which shows the weight he is capable of raising.

From experiments made by Peron, in his voyage, on 12 individuals of Van Diemen's Land, 17 of New Holland, 56 of the Island of Timor, 17 Frenchmen belonging to the expedition, and 14 Englishmen in the colony of New South Wales, he found their comparative strength, indicated by the dynamometer, to be as follows:

NATIVES OF	STRENGTH			
	of the arm.	of the loins.		
	Kilog.	lbs.	Myri.	lbs.
1. Van Diemen's Land.	50.6	101.2		
2. New Holland.	50.8	101.6	10.2	208.08
3. Timor.	58.7	117.4	11.6	238.64
4. France.	69.2	138.4	15.2	310.08
5. England.	71.4	142.4	15.2	332.52

DYNAMOMETER OF DYNAM'ETER, MEDIC'INAL. An instrument, invented by Dr. Paris, for the purpose of showing the quantity of active matter contained in a given weight or measure of any official compound, with the dose of any preparation, which will be equivalent in strength to a given quantity of any other of the same class. The instrument is more ingenious than useful.

DYNAMOPATHY, Homœopathy.

DYOTA, Diota.

DYS, *δυσ*, in composition, 'difficult, faulty,' sometimes privative; mostly answering to the particles *dis*, *in*, *mis*, or *un*, in English. Hence:

DYSÆMIA, from *δυσ*, 'with difficulty,' and *αἷμα*, 'blood.' A morbid condition of the blood.

DYSÆSTHESIA, *Dysæsthe'sis*, from *δυσ*, 'with difficulty,' and *αἰσθάνομαι*, 'I feel.' Obscure, diminished, or even abolished sensation.

DYSÆSTHESIA AUDITORIA, Cophosis—*d. Gustatoria*, *Ageusia*—*d. Interna*, *Amnesia*—*d. Olfactoria*, *Anosmia*.

DYSÆSTHESIA VISUALIS, *Parop'sis*. Sense of sight vitiated or lost.

DYSÆSTHESIS, Dysæsthesia.

DYSANAGOGOS, from *δυσ*, 'with difficulty,' and *αἰνῶ*, 'I bring up.' That which is expectorated with difficulty. An epithet given, by the Greek writers, to the sputa, when expectorated with difficulty, on account of their visciditv.

DYSANARRHOPHESIS, *Dysanarrhoph'ia*; from *δυσ*, 'with difficulty,' and *αναρροήσις*, 'absorption.' Diminished absorption from morbid causes.

DYSAPHĒ, *Dysaph'ia*, from *δυσ*, and *ἀφῆ*, 'feeling.' Morbid feeling.

DYSAPULOTUS, *Dysapu'tus*, *Dyscuplotos*, *Dyscuplō'ticus*, *Dyspu'tus*, from *δυσ*, and *απουλω*, to heal.' Healing with difficulty.

DYSARTHRTIS, Gout, irregular.

DYSCATABROSIS, Dysphagia.

DYSCATAPOISIS, Dysphagia.

DYSCATAPO'TIA, from *δυσ*, and *καταπινω*, 'I drink.' Difficulty of swallowing liquids. This term is recommended by Dr. Mead as a substitute for hydrophobia, which means dread of liquids. The dread seems to be partly caused by the difficulty of deglutition.

DYSCHEZIA, from *δυσ*, and *χεῖν*, 'to go to stool.' Difficult and painful defecation.

DYSCHRŒA, from *δυσ*, 'with difficulty,' and *χρῶα* or *χρῶα*, 'colour.' *Dyschrō'a*. Sickly and unhealthy colour of the skin. Used synonymously with the *maculæ* of Willan.

DYSCHROMATOPSIA, Achromatopsia.

DYSCHYMOSEN, Chymoplasia.

DYSCINE'SIA, (F.) *Dyscinē'sie*, from *δυσ*, 'with difficulty,' and *κινεω*, 'I move.' Difficulty or utter incapability of moving.—Galen.

DYSCOILIA, Constipation.

DYSCOPHOSIS, from *δυσ*, 'with difficulty,' and *κοφω*, 'I am deaf.' A defect in the sense of hearing.—Hippocrates.

DYSCORIA, *Coremetamorpho'sis*, from *δυσ*, and *κορη*, 'the pupil.' Irregularity of shape of the pupil.

DYSCRASIA, from *δυσ*, and *κρῶσις*, 'tempera

ment. *Intempe'ries, Dintemperan'tia, Dys'crasy.* A bad habit of body. See *Dyssthetica*.

DYSCRASIA POTATO'RUM. Dyscrasia of drinkers. The aggregate of cachectic phenomena exhibited by those who indulge in alcoholic liquors.

DYSCRASIA SATURNINA. Saturnismus—d. Scrofulosa, Scrofula—d. Tuberculosa, see Tubercle.

DYSCRASIACUM, Spanamic.

DYSCRASY, Dyscrasia—d. Bilious, Cholosis, Icterus.

DYSCRITOS, from *δυσ*, 'with difficulty,' and *κρισις*, 'judgment.' That which it is difficult to judge of.—*Hippocrates*.

DYSDACRIA, Dysdacryo'sis, from *δυσ*, and *δακρουν*, 'a tear.' A morbid condition of the tears.

DYSDACRYOSIS, Dysdacria.

DYSECCRIS'IA, from *δυσ*, and *εκκρισις*, 'excretion.' Difficult or defective excretion.

DYSECOIA, Baryecolia, Deafness.

DYSEL'CIA, Dysepulo'tus, from *δυσ*, 'with difficulty,' and *ελκος*, 'an ulcer.' An ulcer difficult to heal.—*Hippocrates, Fœsius*.

DYSEMESIA, Vomiturition.

DYSENTERIA, Dysentery—d. Biliosa, Colocolosis—d. Hæmatera, Dysentery—d. Hepatica, Hepatirrhœa—d. Maligna, Enterocæce—d. Putrida, Enterocæce—d. Scorbutica, Enterocæce—d. Splenica, Melæna—d. Typhodes, Enterocæce.

DYSETER'IC, Dysenter'icus, Dysenter'us, (F.) *Dysentérique, Dysentérique.* Same etymon as *Dysentery*. Relating to dysentery.

DYSENTÉRIE, Dysentery.

DYSETERIUM, Dysentery.

DYSETERONERVIA, Colic—d. Saturnina, Colica metallica.

DYSE'NTERY, Dysenter'ia, D. hæmate'ra, Dysenter'ium, from *δυσ*, and *εντερων*, 'an intestine.' *Difficul'tas intestino'rum. Dissolutus morbus, Diarrhœa carnosæ, Coli'tis, Coloni'tis, Colo-recti'tis, Endocoli'tis, Encoli'tis, Deslux'us Dysenter'icus, Febris Dysenter'ica, Fluxus dysentericum, Fluxus dysenter'icus seu cruent'us cum Tenes'mo, Rheumatismus intestino'rum cum ul'cerè, Tor'mina, T. Celèr', Blennenter'ia, Morbus dissolutus, Sedes cruentæ. Luca dysenter'ica, Bloody Flux; Flux, (F.) Dysentérie, Dysentérie, Flux dysentérique, Flux de Sang.* Inflammation of the mucous membrane of the large intestine; the chief symptoms of which are:—fever, more or less inflammatory, with frequent mucous or bloody evacuations; violent tormina and tenesmus. When the evacuations do not contain blood, it has been called *Dysenter'ia alba* or *simple Dysentery, Myco-dysenter'ia*. The seat of the disease is, generally, in the colon and rectum. It occurs, particularly, during the summer and autumnal months, and in hot climates more than in cold: frequently, also, in camps and prisons, in consequence of impure air, and imperfect nourishment: and is often epidemic. Sporadic cases of dysentery are, generally, easily managed; but when the disease occurs epidemically, it often exhibits great malignancy. Generally, it yields to mild laxatives, as castor oil, combined with diaphoretic narcotics, such as the *pulvis ipecacuanhæ compositis*, and counter-irritants to the abdomen; but, at times, the inflammation runs on so speedily to ulceration, that, unless a new action be rapidly excited, death will be the consequence. In such cases, mercury must be rapidly introduced into the system, and narcotics may be combined with it.

The whole management in acute dysentery must, of course, be strictly antiphlogistic.

DYSE'NTERY, Bilious, Colocolosis.

DYSEPULOTICUS, Dysapulotus.

DYSEPULOTOS, Dysapulotos, Dyselcia.

DYSGALACTIA, Dysgalia.

DYSGA'LIA, Dysgalac'tia, from *δυσ*, and *γαλα, γαλακτος*, 'milk.' An unhealthy condition or depravation of the milk.

DYSGENNE'SIA, from *δυσ*, and *γεννησις*, 'generation.' Lesion of the generative organs or functions.

DYSGEU'SIA, Disgeus'tia, from *δυσ*, and *γευσις*, 'taste.' A morbid condition of the sense of taste.

DYSHÆMORRHŒ'A, from *δυσ*, *αἷμα*, 'blood,' and *ρεω*, 'to flow.' Difficulty in the flow of blood,—according to some, of the hemorrhoidal flux. Also, symptoms occasioned by its diminution or suppression.—*Sagar*.

DYSHÆ'MIA, from *δυσ*, and *αἷμα*, 'blood.' A morbid condition of the blood.

DYSHAPH'TA, Dysaph'hia, from *δυσ*, and *αφή*, 'touch.' A morbid condition of the sense of touch.

DYSHI'DRIA, Dysi'dria, from *δυσ*, and *ἰδρω*, 'sweat.' A morbid state of the perspiration.

DYSI'A TOS, δυσιατος, from *δυσ*, 'with difficulty,' and *ιαωαι*, 'to heal.' *Cura'tu diffic'ilis.* Difficult of cure.—*Hippocrates*.

DYSLALIA, Balbuties, Bradylogia.

DYSLOCH'IA, Col'ica lochia'tis, Hyster'al'gia lochia'tis, from *δυσ*, and *λοχος*; 'relating to parturition.' Diminution or suppression of the lochial discharge.

DYSMASE'SIS, Dysmasse'sis, Bradymasse'sis, from *δυσ*, 'with difficulty,' and *μασσις*, 'mastication.' Difficult or impeded mastication.

DYSMENIA, Dysmenorrhœa.

DYSMENORRHŒE, Dysmenorrhœa.

DYSMENORRHŒ'A, Dym'e'nia, Parame'nia diffic'ilis, Menorrhag'ia stillati'lia, Menstrua diffic'il'ia seu Dolorosa, Amenorrhœ'a diffic'ilis seu partial'is, Menses dolorifici, Menorrh'gia diffic'ilis, Menstruatio diffic'ilis seu dolorific'a, Laborious or Difficult Menstruation, (F.) Dymenorrhœe, Menstruation difficile, Règles difficiles, Strangurie menstruelle. Catamenia passed with great local pain, especially in the loins:—with sometimes a membranous discharge. *Dysmenorrhœa* is very difficult of removal, and prevents conception. In the married female, if she should be able to pass one period without pain, and subsequently become pregnant, the morbid action may be broken in upon by gestation, and a perfect cure be obtained. Change of air, soon after marriage, will sometimes give occasion to this desirable result. The affection generally depends upon erethism of the interior of the uterus, called into action at each catamenial period. The violence of the pain requires the liberal use of narcotics.

DYSMNE'SIA, from *δυσ*, 'badly,' and *μνησις*, 'memory.' Defective memory.

DYSMORPHE, Deformation.

DYSMORPHIA, Deformation.

DYSMORPHOSIS, Deformation.

DYSNÉPHRONERVIE, Nephralgia.

DYODES, Fetid.

DYSO'DIA, Dysod'mia, Dysos'mia, Fortor, 'fœtor,' from *δυσ*, 'badly,' and *οἶω*, 'I smell.' (F.) *Puanteur.* Sauvages has given this generic name to all diseases, characterized by fetid emanations, from whatever part proceeding:—from the mouth, nasal fossæ, bronchia, stomach, axillæ, groins, &c. Also, a stench or stink. *Pœna.*

DYSDOMIA, Dysodia.

DYSDODONTIASIS, Denti'io diffic'ilis, from *δυσ*, 'with difficulty,' and *οδοντιασις*, 'dentition.' Difficult dentition.

DYSONE'ROS, from *δυσ*, 'with difficulty,' and *νειρος*, 'a dream.' Insomnia, with restlessness.

DYSO'PIA, properly Dysop'sia, Dysora'sia, signifying 'shame,' from *δυσ*, 'with difficulty,'

and *οστροπα*, 'I see.' Difficulty of seeing: obscurity of vision.

DYSOPSIA DISSITORUM, Myopia.

DYSOPSIA LATERALIS, *Parop'sis latera'lis*, *Skue-sight*, *Skew-sight*, *Sight askew*. Vision only accurate when the object is placed obliquely. This state is generally caused by some opacity of the cornea.

DYSOPSIA LUMINIS, Nyctalopia—d. Proximum, Presbytia—d. Tenebrarum, Hemeralopia.

DYSOPSIA, Dysopia.

DYSORASIS, Dysopia.

DYSOREX'IA, *Inappeten'tia*, *Appeti'tus defici'ens*, from *δυσ*, 'with difficulty,' and *ορε'is*, 'appetite.' Diminution of appetite.

DYOS'MIA, from *δυσ*, 'with difficulty,' and *οσμη*, 'smell.' Diminution of smell.

DYOSOPHRE'SIA, *Dysophre'sis*, *Dysophra'sia*, *Dysphre'sis*, from *δυσ*, and *οσφρη'is*, 'the sense of smell.' A morbid state of the sense of smell.

DYSOSTO'SIS, from *δυσ*, and *οστρον*, 'a bove.' A faulty conformation or morbid condition of bone.

DYSPATHIA, see Serious.

DYSPEPSIA, from *δυσ*, 'with difficulty,' and *πεψω*, 'I concoct.' *Limo'sis Dyspep'sia*, *Anorez'ia*, *Apep'sia*, *Bradypep'sia*, *Gastroutax'ia*, *Digestio deprava'ta seu diffi'cilis seu laesa*, *Gastro-ato'nia*, *Concoctio tarda*, *Stom'achi resolutio*, *Cruditas*, *Passio stomach'ica*, *Dyspep'sy*, *Indigestion*, *Difficulty of Digestion*. A state of the stomach, in which its functions are disturbed, without the presence of other diseases, or when, if other diseases be present, they are of but minor importance. The symptoms of dyspepsia are very various. Those affecting the stomach itself are:—loss of appetite; nausea; pain in the epigastrium or hypochondrium; heart-burn; sense of fulness, or weight in the stomach; acrid or fetid eructations; pyrosis, and sense of fluttering or sinking at the pit of the stomach. The sympathetic affections are of the most diversified character. Dyspepsia, being generally of a functional nature, is devoid of danger. When arising from disease of the stomach itself, it is, of course, more serious.

It is usually dependent on irregularity of living; either in the quantity or quality of the food taken: and the most successful treatment is, to put the patient on a diet easy of digestion; to combat the causes, where such are apparent; and, by proper remedies and regimen, to strengthen the system in every practicable manner. A great error exists in regarding it as always a disease of debility. It is often connected with an inflammatory or subinflammatory condition of the mucous lining of the stomach, and of course a very different plan of treatment is required in the two cases. Dyspepsia is often attended with too great a secretion of the gastric acids; but, on other occasions, they would appear to be too small in quantity, so as to constitute alkaline indigestion or neutral indigestion.

DYSPEPSIA CHLOROSIS, Chlorosis—d. Hypochondriasis, Hypochondriasis—d. Pyrosis; Pyrosis.

DYSPEPSIODYNIA, Cardialgia.

DYSPEPSODYNIA, Cardialgia.

DYSPEPSY, Dyspepsia.

DYSEP'TIC, *Dyspep'ticus*, *Dyspep'tus*, *Apep'tic*, *Apep'ticus*; same etymon. Having relation to dyspepsia, as 'dyspeptic bread.' One who suffers from dyspepsia.

DYSPEPTICUS, Dyspeptic.

DYSPEPTODYNIA, Cardialgia.

DYSPEPTUS, Dyspeptic.

DYSPERMA'SIA, *Dyspermatism'us*, from *δυσ*, 'with difficulty,' and *σπερμα*, 'sperm.' Difficulty—sometimes incapacity—of voiding the sperm.

DYSPERMATISMUS, Bradyspermatismus.

DYSPHA'GIA, *Dyscatrobo'sis*, *Dyscatap'osis*, *Deglutit'io diffi'cilis*, *D. laesa*, *D. impedita*, from *δυσ*, 'with difficulty,' and *φαγω*, 'I eat.' Difficulty of deglutition. Dysphagia is almost always symptomatic, either of inflammation or of other disease of the organs of deglutition, or of incomplete obstruction of the œsophagus, by some obstacle within it, or by a neighbouring tumour. At times, it is produced by spasms or paralysis of the œsophagus. The prognosis and treatment vary according to the cause.

DYSPHAGIA ATONICA, Pharyngoplegia.

DYSPHA'GIA CONSTRICTA, *D. Pharynge'a seu Œsophage'a seu Callosa*, *Strictu'ra Pharyng'is seu Œsoph'agi vera seu Callosa*, *Stenochori'a Œsoph'agi*, *Œsophagiare'tia*, *Læmosteno'sis*. Stricture of the pharynx and œsophagus is an affection which may be the result of pharyngitis or œsophagitis; but more frequently of malignant disease in the parietes of the tube. The only remedy is the bougie.

DYSPHAGIA INFLAMMATORIA, Œsophagitis—d. Pharyngea, D. Constricta—d. Œsophagea, D. Constricta—d. Spasmodica, Œsophagismus—d. Callosa, D. Constricta—d. Paralytica, Œsophagoplegia, Pharyngoplegia—d. Nervosa, Œsophagismus—d. Thorpida, Pharyngoplegia—d. Globosa, Angone—d. Hysterica, Angone—d. Inflammatoria, Cynanche tonsillaris—d. Lingua, Paraglossa—d. Ranula, Ranula—d. Uvula, Staphyloedema—d. Scirrhus, Læmoscirrus—d. Spastica, Œsophagismus.

DYSPHO'NIA, *Dys'phony*, from *δυσ*, and *φωνη*, 'the voice.' Difficulty of producing and articulating sounds: voice imperfect or depraved. *Apho'nia* (of some).

DYSPHONIA IMMODULATA NASALIS, Rhinophonia—d. Immodulata palatina, Asaphia.

DYSPHOR'IA, *Inquietudo*, *Asta'dia*. Dissatisfaction; restlessness; suffering; indisposition; from *δυσ*, and *φωρ*, 'I bear.'

DYSPHORIA ANXIETAS, Anxiety—d. Nervosa, Fidgets—d. Simplex, Fidgets.

DYSPHOTIA, Myopia.

DYSPHRESIS, Dysosphresia.

DYSPHO'NIA, from *δυσ*, and *φων*, 'fat.' A morbid condition of the adipous substance.

DYSPASMATIC, Cæoplastic.

DYSPLASTICUM, Spanæmic.

DYSPNÉE, Dyspnœa—d. Cardiacque, Asthma, cardiac.

DYSPNŒ'A, from *δυσ*, and *πνω*, 'I breathe.' *Pseudo-asthma*, *Amphipne'ma*, *Respirat'io diffi'cilis seu brevis et rara*, *Brachypnœ'a*, *Retentio aë'rea*, *Anhelat'ion*, *Short breath*, *Difficulty of breathing*, (F.) *Dyspnœe*, *Courte Haleine*. Dyspnœa may be idiopathic or symptomatic. The latter accompanies almost all thoracic diseases. Urgent dyspnœa has been called *amphipne'ma*, *αμφιπνευμα*.—Hippocrates.

DYSPNŒA CONVULSIVA, Asthma—d. Hydrothoracica, Hydrothorax—d. Physiothoracica, Pneumothorax—d. Pinguedinoso, Pursiness—d. Pneumatica, Pneumothorax—d. Pyothoracica, Empyema.

DYSENTÉRIE, Dysentery.

DYSENTÉRIQUE, Dysenteric.

DYSSIA'LIA, from *δυσ*, and *σάλος*, 'saliva.' A morbid condition of the saliva.

DYSSYN'ODUS, *Dyssyn'isia*, from *δυσ*, and *συνωδος*, 'coition.' *Coitus diffi'cilis*. Difficulty in coition.

DYSSYNUSIA, Dyssynodus.

DYSTHAN'ATOS, from *δυσ*, and *θανατος*, 'death.' That which causes a slow and painful death. One who experiences this kind of death.—Hippocrates, Galen.

DYSTHELA'SIA, from *δύς*, and *θηλαζω*, 'I give suck.' Inaptitude for suckling.

DYSTHERAPEUTOS, *Difficiliter cura'bilis*; from *δύς*, and *θεραπεία*, 'medical treatment.' That which is difficult of cure.

DYSTHESIA, from *δύς*, 'badly,' and *τῆσι*, 'I am situated.' *Dys'thesis*, *Cachec'ia*. Morbid habit. Bad humour. Impatience in disease.—Erotian.

DYSTHETICA, *Cachex'ia*, *Dyscrasia*, *Cachex'ies*; same etymon. A morbid condition of the blood or blood-vessels; alone or connected with a morbid state of other fluids, producing a diseased habit. The fourth order in the class *Hæmatica* of Good, including *Plethora*, *Hæmorrhagia*, &c.

DYSTHYMIA, from *δύς*, and *θυμός*, 'mind.' *Depression*, *Despondency*. A bad sign in acute diseases. Also, Melancholy.

DYSTOC'IA, *Mogostoc'ia*, *Bradytoc'ia*, *Reten'tio fœtus*, from *δύς*, and *τοκος*, 'accouchement.' (F.) *Dystocie*. A laborious accouchement, *Labo'rius* or morbid or difficult labour, *Parodynia*, *Partus diffic'ilis*. See Laborious.

DYSTOCIA ABORTIVA, Abortion—d. *Dyscyesis*, Pregnancy, morbid—d. *Dyscyesis extra-uterina*, see Pregnancy, preternatural.

DYSTOCHI'ASIS, *Hispid'itis*, from *δύς*, 'bad,' and *στοχος*, 'order.' Irregular position of the eye-lashes.—Forestus.

DYSTON'IA, from *δύς*, and *τονος*, 'tone.' Morbid condition of the tone of a tissue or organ.

DYSTROPH'IA, from *δύς*, 'with difficulty,' and *τροφή*, 'nourishment.' Imperfect or defective nutrition.

DYSULOTUS, *Dysapulotus*.

DYSURE'SIA, *Dysure'sis*; from *δύς*, and *ουρησις*, 'passing the urine.' Defective secretion and evacuation of the urine.

DYSURIA, *Dys'ury*, *Uri'næ diffic'ilis excre'tio*, from *δύς*, 'with difficulty,' and *ουρον*, 'urine.' *Stran'gury*, (of some.) (F.) *Dysurie*. Difficulty of passing the urine. In this affection the urine is voided with pain, and a sensation of heat in some part of the urethra. Dysuria is the first degree of retention of urine. It differs from strangury, in which the urine can only be passed in drops and with great straining.

DYSURIA CALCULOSA, Calculi, vesical—d. Irritativa, Calculi vesical—d. Mucosa, Cystirrhœa.

DYSURY, Dysuria.

E.

EAGLE-STONE, *Ætites*.

EAR, *Auris*, *Ous*, *ous*, *Ac'oŕ*, Saxon, *eare*, (Prov.) *Lug*. (F.) *Oreille*. The organ of audition. It is composed of a series of more or less irregular cavities, in which the sonorous rays are successively received and reflected, until they agitate the nerves which are destined to convey the impression to the brain. The ear is contained partly in the substance of the temporal bone; and a part projects externally, behind the joint of the lower jaw. It may be divided into three portions;—the *outer or external ear*, formed by the auricle and meatus auditorius; the *middle ear*, comprising the cavity of the tympanum and its dependencies; and the *internal ear*, comprehending the three semicircular canals, the cochlea and the vestibule; which, together, constitute the *osseous labyrinth*. Within the cavity of this labyrinth are contained membranes having nearly the shape of the vestibule and semicircular canals, but not extending into the cochlea. These membranes form the *membranous labyrinth*. Between the osseous and the membranous labyrinth is situated the liquor of Cotunnus, and within the membranous labyrinth is a fluid, termed, by De Blainville, *vitrine auditive*, from its supposed analogy to the vitreous humour of the eye. The form of the membranous vestibule is not an exact imitation of the osseous cavity, being composed of two distinct sacs, which open into each other,—the one termed the *Sac'culus vestib'uli*; the other *Sac'culus*. Each sac contains in its interior a small mass of white calcareous matter resembling powdered chalk, which seems to be suspended in the fluid of the sacs by means of a number of nervous filaments proceeding from the auditory nerve. These are the *otoconies* and *otolithes* of Breschet.

The auditory nerve is distributed to the cavities of the internal ear.

EAR-DOCTOR, Aurist—e. Flap, Proptoma auricularum—e. Kecker, Tonsil.

EARNING, Rennet.

EAR-PICK, *Otolyph'is*, *Otolyphum*, *Coch-*

lear auricula'rē, *Auriscal'pium*, (F.) *Cure-oreille*. A species of small scoop, used for extracting hardened cerumen from the meatus auditorius externus; or to remove foreign bodies from the ear. If carelessly used, it is apt to excite inflammation of the tube.

EAR-SURGEON, Aurist—e. Surgery. Otiatria.

EAR-TRUMPET, *Tubus acus'ticus*, *Acus'ticum Cornu*, (F.) *Cornet acoustique*. An instrument for collecting sound and increasing its intensity, used by those who are hard of hearing. It is, commonly, a kind of cone, formed of silver, tin, or elastic gum, the base of which is turned towards the person who is speaking, and the apex placed in the entrance of the meatus auditorius externus.

EARWAX, Cerumen.

EARWIG, *Forficula auricularia*.

EARTH CLUB, *Orobanchæ Americana*.

EARTH, FULLER'S, *Cimolia purpurescens*.

EARTH GALL, *Veratrum viride*.

EARTH, HEAVY, Baryta—e. Japan, Catechu—e. Lemnian, Terra Lemnia.

EARTH NUT, Pignut, Bunium bulbocastanum.

EARTH, PONDEROUS, Baryta—e. Samian, Sami terra—e. Sealed, Terra sigillata—e. Tale, Magnesia.

EASTINGWORT, Scabious.

EATABLE, Esculent.

EATIN-BERRIES, see Juniperus communis.

EAU, Water—e. d'Aix-la-Chapelle, see Aix-la-Chapelle.

EAU D'ALIBOUR. This compound is made of sulphate of zinc, and sulphate of copper, each 3j; camphor, ten grains; saffron, four grains; water, four fluidounces. Employed in chronic inflammation of the eyelids, and as a vulnerary.

EAU DE FARNIOS, Liqueur amnii—e. d'Amandes amères, Aqua amygdalarum concentrata.

EAU ANTIPUTRIDE DE BEAUFORT. Mineral lemonade prepared with sulphuric acid.

EAU D'ARMAGNAC, Tinctura cinnamomi composita—e. de Balaruc, Balaruc waters—e. de Barèges, Barèges water—e. de Binelli, Aqua Binellii—e. Blanche, Liqueur plumbi subacetatis

dilutus—*e. de Bonferme*, Tinctura cinnamomi composita—*e. de Bonnes*, Bonnes, mineral waters of—*e. de Boule*, see Ferrum tartarizatum—*e. de Bourbonne-les-Bains*, Bourbonne-les-Bains, mineral waters of—*e. de Brocchieri*, Aqua Brocchierii—*e. des Carmes*, see Melissa—*e. de Chaux*, Liqueur calcis—*e. de Chaux composée*, Liqueur calcis compositus.

EAU DE COLOGNE, *Cologne water*. A celebrated perfume, so called from the place where it is made. The following is one formula: *Oil of bergamot*, ℥iij; *Oil of lemon*, ℥ij; *Oil of lavender*, ℥iiss; *Oil of neroli*, ℥iiss; *Oil of origanum*, ℥ij; *Oil of rosemary*, ℥j; *Essence of vanilla*, ℥ij; *Musk*, ten grains; *Rectified spirit*, Oij; *Rose-water*, Oij; *Orange-flower water*, Oj. Macerate for fourteen days, and filter.

EAU DE CUIVRE AMMONIACALE, Liqueur cupri ammoniati—*e. Distillée*, Water, distilled—*e. de Fontaine*, Water, spring—*e. des Fontaines de la Maréguerie*, Rouen, mineral waters of—*e. contre la Gangrène*, Liqueur hydrargyri nitrici—*e. de Goudron*, see Pinus sylvestris—*e. de Goulard*, Liqueur plumbi subacetatis dilutus.

EAU HEMASTATIQUE DE TISSERAND, *Eau de Tisserand*. A hemastatic water, reputed to possess the same properties as the Aqua Brocchierii. It may be prepared by digesting dragon's blood, and turpentine of the Vosges in water.

EAU D'HUSSON, Vinum colchici—*e. des Hydrophiles*, Serum of serous membranes—*e. Hydro sulfurée simple*, Hydrosulphuretted water.

EAU DE JAVELLE, Bleaching liquid, Aqua alkali'na oxymuriat'ica, Labarraque's Solution, (Common salt, ℥ij; black oxide of manganese, ℥ij; water, ℥ij. Put into a retort, and add, gradually, oil of citriol, ℥ij. Pass the vapour through a solution of subcarbonate of potassa ℥ij in water ℥xxix, applying heat towards the last. S. g. 1.087.) It is stimulant, detergent, and antiseptic, —applied externally.

EAU DE LAC, Water, lake.

EAU DE LÉCHELLE. A hæmostatic water, prepared by macerating, in water, various aromatic and astringent herbs, flowers and roots, with tar, and distilling the same. It has been given internally, in hemorrhage, and used, also, as an external styptic.

EAU DE LUCE, Spiritus ammoniæ succinatus—*e. Magnésienne*, Magnesia, fluid—*e. de Maraïs*, Water, marsh—*e. Médicinale d'Husson*, see Colchicum autumnale, Vinum colchici, and Veratrine, sulphate of—*e. Médicinale naturelle*, Water, mineral—*e. de Mer*, Water, sea—*e. Mercurielle*, Liqueur hydrargyri nitrici—*e. Minérale*, Water, mineral—*e. de Monterossi*, Aqua Binellii—*e. de Naples*, Naples water, factitious—*e. de Neige*, Water, snow—*e. de Pagliari*, Hæmostatic, Pagliari's—*e. de Pluie*, Water, rain—*e. de Potasse*, Liqueur potassæ—*e. de Puit*, Water, well—*e. de Rabel*, Elixir acidum Halleri—*e. Régale*, Nitromuriatic acid—*e. Salée*, Water, sea—*e. de Source*, Water, well—*e. Styttique de Brocchieri*, Aqua Brocchierii—*e. Sucrée*, Hydro-saccharum—*e. de Tisserand*, Eau hæmostatique de Tisserand—*e. Végéto-minérale*, Liqueur Plumbi subacetatis dilutus—*e. de Vichy*, Vichy water—*e. de Vie*, Brandy—*e. de Vie Allemande*, Tinctura jalapii composita—*e. de Vie Comphrée*, Spiritus camphoræ.

EAUX, LES, Liqueur amni—*e. Chandes*, Aigues-caudes—*e. Hépatiques*, Waters, mineral, sulfurous—*e. Minérales artificielles*, Waters, mineral, artificial—*e. Minérales factices*, Waters, mineral, artificial—*e. Minérales ferrugineuses*, Waters, mineral, gaseous, &c.—*e. Minérales gazeuses ou acidules*, Waters, mineral, gaseous, &c.—*e. Minérales salines*, Waters, mineral, saline—

e. Minérales sulfureuses, Waters, mineral, sulfurous—*e. Sulfurées*, Waters, mineral, sulfurous.

ÉBEAUPIN, MINERAL WATERS OF. An acidulous chalybeate, in the department of Loire Inférieure, near Nantes.

ÉBLOUISSEMENT, Dazzling.

EBRIECA'SUM, from *ebrius*, 'drunk.' A term employed by Paracelsus to denote a disturbance of the reason, similar to what occurs in drunkenness.

EBRIETAS, Temulencia.

EBRIOSITAS, Temulencia.

EBULLITIO, Ebullition, Strophulus—*e. Stomachi*, Ardor ventriculi, Pyrosis.

EBULLIT'ION, *Ebullit'io*, *Éstual'io*, *Anaz'ois*, *Causis*, from *ebullire*, (*e* and *bullire*), 'to bubble up.' *Boiling*; (*F*) *Bouillonnement*. The motion of a liquid, by which it gives off bubbles of vapour, by heat or fermentation. The boiling point of liquids varies according to the pressure to which they are subjected. For the point of ebullition of different substances, see Heat.

Ebullition is used in France, in a vague manner, for every kind of transient eruption of the skin, occurring without fever or with a slight febrile attack.

EBULUS, Sambucus ebulus.

EBUR, Ivory, see Tooth.

EBURNEATIO, Eburnification.

EBURNIFICATION, *Eburnificatio*, *Eburnatio*, *Eburna'tio*, *Eburna'tion*, from *ebur*, 'ivory,' and *fic*, 'to be made.' An incrustation of the cartilages of articular surfaces of bones with phosphate of lime, which gives them the whiteness and hardness of ivory;—*Cartilages eburnées*.

EC, (before a vowel, *Ex*), *ex*, *ē*, 'out of, from, of.' Hence, *Echynoma*, Eclectic, &c.

ÉCAILLE, Scale.

ÉCAILLES D'HUITRES, *Ostræa testæ*.

ÉCAILLEUX, Squamous.

ECBALIA ELATERIUM, *Momordica elaterium*.

ECBALIUM AGRESTE, *Momordica elaterium*—*e. Officinarium*, *Momordica elaterium*.

ECBESOMENON, Eventus.

ECBLOMA, see Abortion.

ECBOLE, Abortion.

ECBOLIC, Abortive.

ECBOLICUS, Abortive.

ECBOLIUM ELATERIUM, *Momordica elaterium*.

ECBOLIUS, Abortive.

ECBRAS'MATA, from *εβραζω*, 'I boil up.' *Echym'ata*. Hippocrates uses the word for certain burning eruptions.

ECCATHARTICUS, Cathartic.

ECCEPHALO'SIS, *Excerebra'tio*, *Cephalotom'ia*, from *ex*, and *κεφαλη*, 'head.' The removal of the brain of the child to facilitate delivery.

ECCHELYSIS, Expectoration.

ECCHORESIS, Defecation.

ECCHYLOMA, *Succus expressus*.

ECCHYMATA, *Echrasmata*.

ECCHYMOMA, *Echymo'sis*, *Pelidno'ma*, *Pelid'ma*, *Ηυρμα*, *Ηυρμα'tia*, *Ηυρμα'tia*, from *ex*, and *χυρος*, 'juice,' 'humour.' *Ejju'sio*, *Exsuectio*, *Suffu'sio sanguinis*. A livid, black, or yellow spot, *Livor sanguineus*, produced by blood effused into the areolar tissue from a contusion. Spontaneous effusions, occurring as the result of disease or after death, are called *sugillationes*.

ECCHYMOMA ARTERIOSUM, see Aneurism—*e. Capitis* recens natorum, *Cephalomatoma*—*e. Hyponychon*, *Hyponychon*—*e. Lymphaticum*, *Phlegmatia dolens*—*e. Melasma*, *Melasma*.

ECCHYMOSIS, *Echymoma*.

ECCYSIS, Effusion.

ECCLISIS, Luxation.

EC'COPE, *Ec'tomē*, *Ectom'ia*, from *ex*, and

κοπεῖν, 'to cut.' The act of cutting out: also, a perpendicular division of the cranium by a cutting instrument. See *Entaille*.

ECCOPEUS. Same etymon. A knife or instrument for cutting. An ancient instrument—the *pericrator*—used in trepanning.

ECCOPROSISTHESIS, from *εκ*, *κοπος*, 'excrement,' and *αισθησις*, 'sensation.' The sensation or desire to evacuate the bowels.

ECCOPROSIS, Defecation.

ECCOPROTIC, *Eccoproticus*, *Ectoprotic*, from *εξ*, and *κοπος*, 'excrement.' Mild purgatives or laxatives, whose operation is confined to simply clearing out the intestinal canal.

ECCORTHATICUS, from *εκ*, and *κορθω*, 'I collect.' An ancient epithet for remedies to which was attributed the property of evacuating collections of humours.

ECCRINOLOGY, *Eccrinolog'ia*, *Eccrisiolog'ia*, from *εκκρινω*, (*εκ*, and *κρῖνω*), 'I separate,' and *λογος*, 'a discourse.' A treatise on the secretions.

ECCRISIOLOGIA, *Eccrinology*.

ECCRISIOSIS, *Eccrision'is*, from *εκκρίσις*, 'excretion,' and *νοσος*, 'disease.' Diseases of excretion.

ECCRISIOSCHESIS, from *εκκρίσις*, 'excretion,' and *σχῆσις*, 'retention.' Arrest or retention of excretions; or of a critical evacuation.

ECCRISIS, Excretion.

ECCRITICA. Diseases of the excrement functions. The 6th class in Good's Nosology. Also, medicines that act on the excrement system. *Eliminantia*, *Eliminatives*.

ECCYESIS, Pregnancy, extra-uterine—e. Abdominalis, Pregnancy, abdominal—e. Ovaria, Pregnancy, ovarian—e. Tubalis, Pregnancy, tubal.

ECCYLLOSIS, from *εκ*, and *κυλλω*, 'to turn round.' *Morbis evolutio'nis*. A disease of evolution or development.

ECEDEMIOMANIA, *Ecedemion'osus*, from *εκδημω*, 'I travel about,' and *μανία*, 'mania.' A morbid desire to be travelling about.

ECEDEMONOSUS, *Ecedemionian*.

EC'DORA, from *εκ*, and *δερμα*, 'I flay.' *Anad'ora*, *Eccoriat'io*. Excoriation in general, but more especially of the urethra.

EC'DYSIS, from *εκδωω*, (*εκ*, and *δωω*), 'I put off.' Moulting of the skin of animals. Desquamation.

ÉCHALOTTE, *Eschalotte*, *Allium Ascalon'icum*, *Cepa Ascalon'ica*. The *schallot*. A species of allium, employed in culinary preparations.

ÉCHANCRURE (F.) *Emargina'tio*, *Emarginatu'ra*, *Incisu'ra*. A French word employed by anatomists to designate depressions and notches of various shapes, observed on the surface or edges of bones.

ÉCHANCRURE ETHMOÏDALE is on the nasal bone, which unites with the ethmoid. See *Ethmoid*.

ÉCHANCRURE NASALE, *Nasal Notch*, belongs to the os frontis, and is articulated with the bones of the nose.

ÉCHANCRURE PAROTIDIENNE is a triangular space, comprised between the parotidian edge of the inferior maxillary bone and the mastoid process, so called because it lodges the parotid gland.

ÉCHANCRURE SCAPULAIRE, *Notch*, scapular—*é. Sciatique grande*, see *Sciatic Notch*—*é. Sciatique petite*, see *Sciatic Notch*.

ÉCHARDE, Splinter.

ÉCHARPE, Sling.

ÉCHARPE GRANDE et *É. MOYEN*, see *Sling*—*é. de J. L. Petit*, see *Sling*—*é. Petite*, see *Sling*.

ÉCHAUBOULURES (F.) *Échaubouillure*.

Sudam'ina, *Hidro'a*. A word whose meaning is not fixed. It is applied to any eruption on the surface of the body, accompanied with pricking and other uneasy sensations.

ÉCHAUFFANTS, *Calefacients*.

ÉCHAUFFEMENTS, *Chafing*.

ÉCHAUFFEMENT (F.), *Calefac'tio*, *Escale-fu'tio*, from (F.) *échauffer*, (*calefacere*), 'to make warm.' Augmentation of heat in the animal economy; the symptoms of which are a more than ordinary sensation of heat, disposition to perspiration, great thirst, general indisposition, flushed countenance, &c. It goes off by the use of antiphlogistics and abstinence. In the vulgar language it is often used synonymously with *constipation*, and sometimes for simple gonorrhœa, and for chafing.

ÉCHECOL'ON, from *εχω*, 'I have,' and *κολλα*, 'glue.' *Echecollum*. Any topical glutinous remedy.—Goræus, Galen.

ÉCHELLES DU LIMAÇON, *Scalae of the cochlea*.

ÉCHENEIS, *Remora Hildani*.

ÉCHETROSIS, *Bryonia alba*.

ECHID'NA OCELLATA, *Brown ten-inch-long viper*. A most formidable viper in the forests of Peru, the bite of which is so rapidly fatal, that it kills a strong man in two or three minutes. *Tschudi*.

ECHINACEA PURPUREA, from *εχινος*, the hedge hog, owing to the spiny chaff of the roundish disk. *Purple Coneflower*, *Black Sampson*; of the Composite Family; indigenous in Ohio and westward; its dull purple flowers appearing in July. The root is aromatic, and used popularly as a carminative.

ÉCHINE, *Vertebral column*.

ECHINOCOCCUS, *HOMINIS*, see *Worms*—*e. Humanus*, *Hydatid*.

ECHINODERMI, *Porcupine men*.

ECHINOGLOSSUM, *Ophioglossum vulgatum*.

ECHINOPHTHALMIA, from *εχινος*, 'a hedgehog,' and *οφθαλμα*, 'inflammation of the eye.' Ophthalmia of the eyelids, in which the cilia project like the quills of the hedgehog.

ECHINOPS, from *εχινος*, 'hedgehog,' and *ωψ*, 'appearance.' *Crocodil'ion*, *Acanthal'zucca*, *Scabiosa carduifolia*, *Sphæroceph'ala elatior*, *Echinopsis*, *Echinops Sphæroceph'alus*, *Globe thistle*. The root and seeds are reputed to be moderately diuretic.

ECHINOPUS, *Echinops*.

ECHINUS SCANDENS, *Allamanda*.

ECHITES DIFFORMIS, *Forsteriana difformis*—*e. Suberecta*, see *Curare*.

ECHOS, *Sound*, *Tinnitus aurium*.

ECHOSCOPE, *Auscultation*.

ECHOSCOPIUM, *Stethoscope*.

ECHTHYSTEROCYESIS, *Pregnancy, extra-uterine*.

ECLACTIS'MA, *Eclamps'is*, *Eclamp'sia*, *Epileps'is*, *Effulgescentia*, from *εκλακτίζω*, 'I kick.' Epilepsy is often accompanied with flashings of light; and hence Hippocrates has used the last two words for epilepsy. They have all been applied to convulsions.

ÉCLAIRE, *Chelidonium majus*—*é. Petite*, *Ranunculus ficaria*.

ECLAMP'SIA, from *εκλαμ'ναι*, (*εκ*, and *λαμβάνω*, 'I shine,') 'brilliancy.' Flashes of light before the eyes. See *Eclactisma*. Convulsion, as the convulsions of children, *Eclamps'is infantum*, *Epileps'is acut'a infantum seu febrilis infantum seu puerilis*, (F.) *Convulsiones des Enfants*, *Eclampsie*, but probably from *εκ*, and *λαμβάνω*, *λαμβάνω*, 'to seize hold of.'

ECLAMP'SIA GRAVIDA'RUM ET PARTURIEN'TIUM; *Puerperal Convulsions*, (F.) *Convulsions*

des femmes enceintes et en couche. Convulsions of pregnant and parturient women.

ECLAMPsia INFANTUM, see Eclampsia—e. Nutans, Convulsion, Salaam—e. Typhodes, Raphania.

ECLAMPsIE DES ENFANS, Eclampsia infantum.

ELECTIC, (PHYSICIANS), *Eclectici Medici*, from *ἐκλεγω*, (*εκ*, and *λεγω*.) 'I choose.' A sect of physicians, who professed to choose, from other sects, all the opinions which appeared to them best founded. Agathinus of Sparta, master of Archigenes of Apamea, in Syria, was its reputed founder; and Archigenes and Aretæus were its greatest ornaments. The doctrine was called *Eclectism*, *Eclectismus*, *Medicina eclectica*, *Eclectic medicine*. Every judicious physician must be an eclectic.

ELECTISM, see Eclectic.

ELECTOS, *Eclog'na*, *Elig'ma*, *Elis'is*, *Lambit'um*, *Linctus*, *Linctu'dium*, from *ἐκλεχω*, (*εκ*, and *λεχω*.) 'I lick.' (F.) *Looch*. A medicine, of a thick, syrupy consistence, chiefly used to allay cough, and consisting of pectoral remedies. It was formerly sucked from the end of a liquorice stick, made into a kind of pencil; hence its name *Linctus*, from *lingere*, 'to lick.' Although the linctus is usually exhibited in thoracic affections, it may have tonic virtues combined with it.

ELEGMA ALBUM, *Looch album*—e. Gummoso-oleosum, *Looch album*.

ECLIPsIS, Desquamation, Exfoliation.

ECLIPsIS, Desquamation, Exfoliation.

ECLIPsITREPANON, Exfoliative trepan.

ECLIGMA, Eclectos.

ECLIMIA, Boulimia.

ECLIPsIS, Syncope.

ÉCLISSE, Splint.

ECLYSES, Adynamie.

ECLYSIS, *Ecclut'io*; from *ἐκλυω*, 'I loosen.' Resolutio, prostration of strength; faintness.

ECLYSIS PNEUMO-CARDIACA, Asphyxia.

ECMYZESIS, Exsuctio.

ECNEA, Dementia.

ÉCOLE, School.

ECONOMY, *Econom'ia*, from *οικια*, 'a house,' 'a family,' and *νομω*, 'I rule.' By the term *animal economy* is understood,—the aggregate of the laws which govern the organism. The word *economy* is, also, used for the aggregate of parts which constitute man or animals.

ÉCORCE, Cortex—*é. Cariocostine*, *Canella alba*—*é. Élauthérienne*, see *Croton cascarilla*—*é. de Saint Lucie*, *Cinchonæ Caribææ cortex*—*é. de Winter*, see *Winters aromaticæ*—*é. Fausse de Winter*, *Canella alba*.

ÉCORCHURES, Chafing, Excoriation.

ÉCOULEMENT, Discharge, Gonorrhœa—*é. Blanc*, *Leucorrhœa*—*é. de Sang par l'Intestin*, *Hematochezia*.

ÉCOUVILLON, see *Écouvillonnement*.

ÉCOUVILLONNEMENT (F.), from *écouiller*, 'a kind of mop, the sponge of a gun.' A term used by the French therapists for the act of cleansing or applying remedies to a part by means of a mop or brush fixed to the end of a piece of whalebone. Such mop or brush is termed *Écouvillon*.

ECPHLOGOSIS, Inflammation.

ECPHLYSIS, Vesicula—e. Herpes, Herpes—e. Herpes circinatus, Herpes circinatus—e. Herpes exedens, Herpes exedens—e. Herpes miliaris, Herpes phlyctenodes—e. Herpes zoster, Herpes zoster—e. Pompholyx, Pompholyx—e. Rhyphia, Rupia.

ECPHRACTIC, Deobstruent.

ECPIRAXIS, from *ἐκφρασσω*, 'I remove ob-

struction.' The action of ephractic or deobstruent remedies.

ECPHRONIA, Insanity—e. Melancholia, Melancholy.

ECPHYAS, Appendix vermiciformis cæci.

ECPHYMA, Excrescence, Tumour—e. Callus, Callosity—e. Caruncula, Caruncle—e. Clavus, Corn—e. Cœdematium, Cœdema, Phlegmatia dolens—e. Physconia, Physconia—e. Trichoma, Plica—e. Verruca, Verruca.

ECPHYMATA, Rubela.

ECPHYSEsIS, *Ephlat'io*, *Ephl'itus*, from *ἐκφύσσω*, 'I breathe through.' *Exsufflat'io*. A quick and forced expulsion of air from the lungs.

ECPHYsIS, Apophysis—e. Ventriculi, Duodenum.

ECPIES'MA, from *ἐκπιεζω*, (*εκ*, and *πιεζω*.) 'I compress.' *Ejfractura*, *Impaction*, *Decessio*. A fracture of the cranium, with depression of the fragments and compression of the brain.

ECPIES'MOS, *Ecpress'io*, *Ec't'us*, *Ecpiēs'mon*, same etymon. Celsus uses these words to signify the forcing of the eye from the orbital cavity, with apparent but not real augmentation of the organ. See *Exophthalmia*.

ECPLERO'MA, from *εκ*, and *πληρωω*, 'I fill.' *A cushion*, *a pad*. Hippocrates means, by this term, a small pad or ball of leather, or other substance intended to fill the hollow of the armpit; used probably in reducing luxations of the shoulder.

ECPLEXIA, Stupor.

ECPLEXIS, Stupor.

ECPNEUMATOSIS, Expiration.

ECPNEUSIS, Expiration.

ECPNOE, Expiration.

ECPTO'MA, *Ecpto'sis*, *Exciden'tia*; from *εκπιπτο*, (*εκ*, and *πιπτο*.) 'I fall out.' This word has been used in various senses, 1. Synonymously with luxation. 2. For the separation of granenous parts. 3. For the expulsion of the secundines. 4. For the prolapsus of the womb: and 5. For intestinal or omental hernia, &c.

ECPTOSIS, Luxation.

ECPYCTICA, Incrassantia.

ECPYEMA, Abscess, Suppuration, Empyema.

ECPYESIS, Abscess, Empyema, Pustule—e. Impetigo, Impetigo—e. Porrigo, Porrigo—e. Porrigo crustacea, Porrigo larvalis—e. Porrigo favosa, Porrigo favosa—e. Porrigo furfuracea, Porrigo furfurans—e. Porrigo galeata, Porrigo scutulata—e. Porrigo lupinosa, Porrigo lupinosa—e. Scabies, Psora.

ECPYETICUS, Suppurative.

ECPYSCONTUS, Suppurative.

ÉCRASEUR, 'a crusher,' from *écraser*, 'to crush.' A surgical instrument invented by M. Chassaignac, of Paris, which consists of a steel chain, like that of a chain-saw devoid of teeth, which is attached at both ends to a steel mandril, which passes through a hollow steel cylinder. The tightening of the chain is made slowly and with great force, and in the last modification of the instrument is effected by an endless screw with a lever handle, working on a nut clogged on its outer side, which plays on a thread cut on the mandril. The slow bruising it makes is said to be rarely followed by hemorrhage, even in the case of hemorrhoids, and other vascular tumours.

ÉCREVISSE, Crab.

ECREX'IS, *Ruptura*, from *εκρηγνυμι*, (*εκ*, and *ρηγνυμι*.) 'to break.' Rupture, laceration.

ECRHYTH'MUS, *εκ*, and *ρhythmos*, 'rhythm.' A term applied to the pulse, particularly when irregular.—Galen.

ECROE, Discharge.

ÉCROUELLES, Scrofula—*é. Mésentériques*, *Tabes mesenterica*.

EC'RYSIS, from *εκρεω*, 'I run from.' A discharge.

EC'SARCO'MA, from *εκ*, and *σαρξ*, 'flesh.' A fleshy excrecence of various kinds. See *Fungosity*, and *Sarcoma*.

EC'SESMA, *Eczema*.

EC'STASIS, from *εξίστημι*, (*εξ*, and *ίστημι*.) 'I am beside myself.' An *ec'stasy* or *trance*, (*Carus ec'stasis*, *Cat'ochus*, *Ec'stasis*, *Catalep'sia spu'ria*, *Hyperplecia*, (F.) *Ectase*. A state in which certain ideas so completely absorb the mind, that the external sensations are suspended, the voluntary movements arrested, and even the vital action retarded. In *catalepsy*, there is, in addition, complete suspension of the intellectual faculties. This last condition is in general described as *trance*. See, also, *lutation*.

ECSTROPHE, *Exstrophia*.

ECTASIA, *Aneurism*—e. *Venarum*, *Varix*.

ECTEXIS, *Extension*, *Expansion*.

EC'TASIS I'RIDIS, is the extension or expansion of the iris, which occasions diminution of the pupil.

ECTEXIS, *Colligation*.

EC'THETOBREPHOTROPHEUM, *Brephotropheum*.

EC'THIM'MA, *Exulcera'tio*, from *εκθλιβω*, 'I express.' 'I bruise.' Attrition. Chafing, or excoriation, produced by external violence.—*Hippocrates*.

ECTHILIPSIS, *Expression*.

EC'THYMA or **EC'THY'MA**, from *εκθω*, 'I break out.' *Eccyp'sis*, *Phlysis ecthyma*, *Phlyza'cia d'gria*, *Scab'ies Vera*, *Furru'culi aton'ici*, *Pap'u'losus Scall*, (F.) *Dartre crustacee*, *D. fongueuse*. A cutaneous eruption, characterized by large round pustules, always distinct and seated upon an indurated and highly inflamed base. In the course of a day or two the pustules generally break, and olive-brown incrustations are formed which adhere firmly to the skin. These separate in about a fortnight. The disease requires the antiphlogistic treatment. Under the *Ec'thymata*, *Vogel* has designated certain hard, unequal tumours which appear transitorily on the skin. See *Efflorescence*, *Exanthem*, and *Pustule*.

ECTHYSTEROCYESIS, *Pregnancy*, *extra-uterine*.

ECTILLOTICUS, *Depilatory*.

ECTILMOS, *Evisulion*.

ECTOME, *Castration*, *Ecope*, *Entaille*, *Excision*.

ECTOMIA, *Castration*, *Ecope*.

ECTOMIUS, *Castratus*.

ECTONION, *Helleborus niger*.

ECTOPAG'IA, (F.) *Ectopagie*, from *εκτος*, 'outwards,' and *πηγνυμι*, 'to put together.' A genus of double monsters, having a common umbilicus, comprising those which are united naturally throughout the whole extent of the thorax.

ECTOPARASITES, *Epizoa*.

EC'TOPHYTE, *Ectoph'yton*, from *εκτος*, 'outside,' and *φυτον*, 'a vegetable.' A vegetable parasite, which grows on the surface of the body.

ECTOPIA, *Ectop'ia*, *Ectopism'us*, *Entoce'le*, *Situs alie'nus seu perver'nus seu muta'tus*, *Anom'otopie*,—*Piorry*, from *εκ*, and *τοπος*, 'out of place.' Morbid displacement of parts. See *Luxation*.

ECTOPIA ANI, *Proctocele*.

ECTOPIA CORDIS, *Cardianas'trophē*. Displacement, dislocation, or unnatural position of the heart.

ECTOPIA HERNIOSA, *Hernia*—e. *Splenis*, *Splenectopia*.

ECTOPOCYS'TICUS, from *εκτοπος*, 'out of place,' and *κυστις*, 'bladder.' A disease dependent upon displacement of the bladder.

ECTOPROTIC, *Eccoprotic*.

ECTOZO'A, (sing. *Ectozo'on*), *Extozo'a*, *Extoc'o'ria*, (F.) *Extozoaires*; from *εκτος*, 'outside,'

and *ζωον*, 'an animal.' Parasitic animals that infest the exterior of the body,—as lice. A term which, like *Helmin'thia errat'ica*, *Pseudohelmin'thes*, and *Pseudopar'asites*, is applied, also, to worms or larvae of insects that have been introduced into the intestinal canal by accident. Animals, most frequently swallowed, are the hairworm, leech, grub of the fly, caddy insect—*Phala'na penguina'lis*; the larve of the bee, the spider, the *triton pulu'stris*, *lacc'e'ta aquat'ica*, &c. In animals, bots are produced by swallowing the ova of the *æstrus* or gadfly. See *Helminthia erratica*.

ECTRIM'MA, from *εκτριβω*, (*εκ*, and *τριβω*), 'I rub off.' Ulceration of the skin; and particularly that which arises from the pressure of the bed on different parts of the body, after a protracted confinement.—*Hippocrates*.

ECTRODACTYL'IA, from *εκτρωσις*, 'abortion,' and *δακτυλος*, 'a finger.' A malformation, in which one or more fingers or toes are wanting.

ECTROG'ENY, *Ectrogen'ia*, *Ectrogen'esis*, from *εκτρωσις*, 'abortion,' and *γενσις*, 'generation.' Monstrosity by defect.—*Serres*.

ECTRO'MA, *Abortion*.

ECTRO'MELES, from *εκτρωσις*, 'abortion,' and *μελος*, 'a limb.' A genus of monsters, in which the limbs are nearly or altogether deficient, as in the ordinary cetacea.—*I. G. St. Hilaire*.

ECTROPE, *Divertic'ulum*, from *εκτρέπω*, (*εκ*, and *τρέπω*), 'I turn off,' 'divert.' Any duct by which peccant or morbid matter was supposed to be drawn off.—*Hippocrates*.

ECTROP'ION, same etymon. *Ectrop'ium*, *Eber'sio pal'pebræ*, *Blepharopto'sis Ectrop'ium*, *Blepharoto'sis*, *Pal'pebræ inferior extror'tum flexa*, *Divarica'tio seu Reflex'io seu Reclina'tio palpebrarum*, (F.) *Écartement des Paupières*, *Renversement des Paupières*. Eversion of the eyelids, so that they do not completely cover the globe of the eye. It happens more commonly to the lower than to the upper eyelid. It may be owing to the retraction of the skin, after the cure of an ulcer, wound, or burn of the eyelid; or it may depend on tumefaction or relaxation of the conjunctiva. In the majority of cases, removal of a portion of the conjunctiva will effect a cure; but there are many which defy the efforts of art. The ancients called Ectropion of the upper eyelid *lagophthalmia*.

ECTROSIS, *Abortion*.

ECTROSMOS, *Abortion*.

ECTROT'IC, from *εκτρωμα*, (*εκ*, and *τρωσσω*, 'I wound,') 'abortion.' *Ectrot'ica*, *Abortive*. An epithet applied to methods for preventing the development or causing the abortion of any disease—as of chancres by the use of caustic; small-pox pustules by the use of mercurial ointment, &c.

ECTYLOTICUS, *Catheretic*.

ECTYMPANOSIS, *Tympanites*.

ÉCUMEUX, *Frothy*.

ÉCUSSEONS. 'Escutcheons or shields.' Plasters spread upon the skin: or small bags—*sachets*—of the shape of escutcheons, filled with odorous powders, which are applied on the skin. See *Sachet*.

EC'ZEMA, from *εκζω*, (*εκ*, and *ζωω*), 'I loil out,' 'I effervesce.' *Ecze'sma*, *Eczen'ma*, *Pustula ardens*, *Cyrtis'ma Eczema*, *Humid Scall* or *Tetter*, *Running Scall*, (F.) *Dartre squameuse humide*, *D. vive*, *Gale épidémique*. Heat eruption. An eruption of small vesicles on various parts of the skin, usually set close or crowded together; with little or no inflammation around their bases, and unattended by fever.

ECZEMA CAPITIS, E. of the hairy scalp.

ECZEMA OF THE FACE, at an advanced stage,

and occurring in young children, has been described under the names *Crusta lactea* and *Porri-go larvalis*.

ECZEMA OF THE HAIRY SCALP, *Eczema capitis*, *Vesicular scall*, is often confounded with other affections, under the names *Porri-go* and *Tinea*, which are pustular, not vesicular, in their form. It occurs during dentition, and even afterwards, and the discharge is so profuse, that the head appears as if dipped in some glutinous liquid. By and by, the secretion dries into crusts and mats the hair into little separate tufts. The scalp gives evidence of inflammatory excitement, and the lymphatic ganglions of the neck are apt to become inflamed and suppurate.

A variety of humid scalled head, in which the humour from the excoriated surface runs down upon the hairs, and encloses them in little silvery pellicles or sheaths, has received the name *Asbestos Scall*.

ECZEMA IMPETIGINOSUS, *Gall*, *Grocers'* or *Bricklayers' Itch*. Produced by the irritation of sugar or lime.

ECZEMA MERCURIALE, *E. rubrum*, *Erythema mercuriale* seu *ichthiosis*, *Hydrargyria*, *Hydrargyrosis*, *Hydrargyriasis*, *Morbus mercurialis*, *Mercurial lepra* or *rash*. A variety of eczema, arising from the irritation of mercury. The treatment is chiefly palliative, consisting in ablation with mucilaginous infusions or decoctions; mild dressings, where the cuticle has exfoliated; avoiding all irritation; keeping the bowels open; with the use of sulphuric acid and cinchona.

ECZEMA RUBRUM, *Eczema mercuriale*.

ECZEMATOSSES, (*G.*) *Eczematosen*, same etymon. A family of diseases, in the classification of Fuchs, including morbid conditions of the cutaneous secretions—as of the perspiration, sebaceous and colouring matters, &c., and hence many chronic cutaneous affections. His subdivisions are, *epithroses*, *smegmorrhœa*, *acarpæ*, *polycarpæ*, and *monocarpæ*.

ECZESIS, *Effervescens*.

ECZESMA, *Eczema*, *Lichen tropicus*.

EDEMATOUS, *Edeumatous*.

EDENTATUS, *Edentulus*.

EDENTULI, *Nefrendes*.

EDENTULOUS, *Edentatèd*, *Edentulus*, *Edentatus*, from *e*, and *dens*, *dentis*, 'a tooth.' *Anodus*, *Carens dentibus*, *Nodes*, *Nodus*, (*F.*) *Édenté*. One without teeth. This defect can only be remedied by artificial means. See *Nefrendes*.

EDERA, *Hedera helix*.

EDIBLE, *Esculent*.

ÉDOCÉPHALE, from *aîda*, the 'sexual organs,' and *κεφαλή*, 'head.' A monster, which has the two ears near each other, or united under the head, the jaws atrophied; no mouth, and above the eye a tube (*trompe*) resembling a penis.—*I. G. St. Hilaire*.

EDROPPIT, *Hydropic*.

EDUCATIO INFANTUM, *Pædia*.

EDULCORATION, *Glycan'sis*, *Edulcoratio*, from *e*, and *dulcis*, 'sweet.' An operation, the object of which is to deprive a substance of its acid and disagreeable taste, or at least to disguise it. Also, the addition of a saccharine substance to a medicine, whose taste it is desirable to modify agreeably.

ÉDULE, *Comestible*.

EDULIS, *Comestible*.

EE, *Eye*.

EESKIN, *Singultus*.

EEWINKER, *Palpebra*.

EFFERENT, *Eff'rerens*, *Centrifugal*, *Exod'ic*, from *effero*, (*e*, and *fero*), 'I carry,' 'transport.' Conveying outwards, as from the centre to the periphery.

Vasa efferentia are those lymphatic or chyloferous vessels which issue from the glands to convey their lymph to the thoracic duct; so called to distinguish them from those which pass to those glands, and which have been termed *vasa afferentia* seu *inferentia*. Also, nerves that convey the nervous influence from the nervous centres to the circumference. See *Afferent*.

At the upper extremity of the mediastinum testis, the ducts of the rete testis terminate in from 9 to 30 small ducts, called *vasa efferentia*, which forms the *Coni vasculosi*.

EFFERVES'CENTE, *Effervescen'tia*, *Zesis*, *Ec'zesis*, from *effervescere*, (*e*, and *fervescere*,) 'to grow hot.' That agitation, which is produced by the escape of gas through a liquid, independently of the heat of the mixture; such, for instance, as results from the mixture of acetic acid and carbonate of potassa.

In *Pathology* it has a similar signification. It expresses, in the language of the humorists, a sort of ebullition in the fluids of the living body, produced either by elevation of temperature or by the reaction on each other of the principles contained in the fluids in circulation.

EFFETUS, *Impoverished*.

EFFICA'CIOUS, *Eff'icax*, from *efficere*, (*e*, and *facere*), 'to accomplish.' That which produces a great effect,—as 'an efficacious remedy.'

Medici'na eff'icax, *La Médecine efficace*, is a term sometimes applied to surgery.

EFFILA, *Ephelides*.

EFFLATIO, *Echphysis*.

EFFLATUS, *Echphysis*.

EFFLORATIO, *Exanthem*.

EFFLORES'CENTE, *Effloratio*, *Efflorescen'tia*, from *efflorescere*, (*e*, and *florescere*), 'to blow as a flower.' *Stribili'go*, *Eethyma*.

In *Pathology*, efflorescence has the same meaning as *exanthema*; and, in the nosology of Sauvages, the name is given to that order of diseases. Sometimes, it is confined to the cutaneous blush, the *exanth'esis* of Good.

Efflorescence is, also, the conversion of a solid substance into a pulverulent state by exposure to the air. In salts this is generally owing to the loss of a part of their water of crystallization.

EFFLORESCENCE ÉRYSIPELATEUSE, *Roseolæ*.

EFFLORESCENTIA, *Exanthem*.

EFFLORESCENTIE, *Élévures*.

EFFLUVIUM, *Emanation*—*e. Latrinarium*, *Mitte*—*e. Palustre*, *Miasm*, *Marsh*.

EFFLUXION, *Abortion*.

EFFORT, (*e*, and *fortis*, 'strong,') *Nismus*, *Conatus*, *Peira*. A muscular contraction of greater or less strength, the object of which is, either to resist an external force, or to accomplish a function, which has become naturally laborious;—such are, the act of pushing away, or of drawing a body towards us, and the more or less painful efforts used by the mother to cause the expulsion of the fetus. In France, the word *effort* is often used synonymously with *hernia*; and signifies, likewise, the painful twitches of muscles, occasioned by over-exertion, or by the rupture of some of their fleshy fibres. Sauvages calls *Efforts des reins*, *Lumba'go à nisu*, the pain in the loins occasioned by bearing too heavy a burden.

EFFORT, *Hernia*—*e. des Reins*, see *Effort*.

EFFOSSIO, *Exhumation*.

EFFRACTURA, *Ecpiesma*.

EFFRACTURA CRANII, *Enthlasis Cra'nii*, *Fracture* of the Cranium, with depression.—*Paré*.

EFFRENITATIO, *Hypercatharsis*.

EFFUSIO, *Effusion*—*e. Seminis*, *Ejaculation*.

EFFUSION, *Effusio*, *Ec'chysis*, from *effundere*, (*e*, and *fundere*, *fusum*), 'to pour out.' (*F.*) *Épan*.

chement, (Infiltration is the term generally employed for effusion into the areolar membrane. The pouring out of blood or of any other fluid into the areolar membrane, or into the cavities of the body. The effusion of serum or of coagulable lymph, for instance, is a common result of inflammation of serous membranes.)

• *EGARÉ*, Wild.

EGAREMENT D'ESPRIT, Delirium, Insanity.

EGER, MINERAL WATERS OF THE. In the valley of the Eger, at the western extremity of Bohemia, there are several acidulous springs. One of the most frequented of these is Franzensbad.

EGERIA. In ancient mythology, a nymph to whom pregnant females offered sacrifices *ut conceptus alvus facilius egeretur*. By some, supposed to have been identical with Lucina.

EGESTA, see Excretion.

EGESTAS, Want.

EGESTIO, Defecation, Dejection, Excretion.

EGGBED, Ovarium.

EGG-BRANDY, see Ovum.

EGG-PLANT, *Solanum melongena*—*e*. White of, Albumen ovi.

EGLANTIER DE CHIEN, *Rosa canina*—*e*. *Sauvage*. *Rosa canina*.

EGOBRONCHOPHONY, see Egophony.

EGOPHONE, Egophony.

EGOPHONIC, *Égophônicus*, (F.) *Égophonique*. Same etymon as the next. Having the character of, or relating to, egophony.

ÉGOPHONIQUE, Egophonic.

EGOPHONY, *Égophônia*, from αἶς, αἶγος, 'a goat,' and φωνή, 'voice.' *Capriloquium*, *Tragophônia*, Goat's Voice, Bleating Voice, (F.) *Égophonie*, *Voice chevrotante* seu *égophonique* seu *de Polichinelle* seu *sauille*, *Pectoriloquie chevrotante*. Laënnec has designated by this name, the kind of resonance of the voice heard through the stethoscope, when we examine the chest of one labouring under moderate effusion into one of the pleuræ. The voice, which strikes the ear through the cylinder, is more sharp and harsh than usual, and tremulous and broken, like that of the goat. The patient himself is called *Égophoné*. Bouillaud affirms, that the 'bronchial and bleating voice,' (*égobronchophoné*) is the principal symptom of pleuropneumonia. Egophony exists, however, in cases of hepatization where there is no pleural disease.

EGREGOR'SIS, *Vigil'ia*, *Vigili'æ*, *Vigilan'tia*, *Vigila'tio*, *Vigil'ium*, from *εγρεγορεω*, 'I watch.' Watchfulness. A morbid want of sleep.—Galen.

EGRESSUS VENTRICULI, Pylorus.

EIDOS, εἶδος, 'form, resemblance.' The *ε* is often changed into *ω*, at the termination of a word. Thus, *Hæmatoides* or *Hæmato'des*.

ELAMIDES, Meninges.

ELLE'MA, from εἰλεος, Ἰλεος, (εἰλω, 'I roll,') 'a convulsion.' Vogel has given this name to a fixed pain, occupying some portion of the intestinal canal, which the patient compares to the sensation that would be produced by a nail driven into the part.

EILEON, Ileon.

EILEUS, Ileus.

EILOID, (Tumour,) *Eiloid'es*, from εἰλω, 'I roll,' and εἶδος, 'resemblance.' A morbid growth of the cutis, coiled or folded.

EILSEN, MINERAL WATERS OF. Eilsen is about six German miles from Hanover, at the foot of the Harrelberg. It has eleven springs, of which seven are sulphureous and four chalybeate.

EISANTHE'MA, *Eranthe'ma inter'num*, *Entanthe'ma*, from εἰς, 'within,' and ἀνθημα, 'efflo-

rescence. An eruption on a mucous membrane;—apthæ, for example.

EISBOLE, Attack, Injection.

EISPNØE, Inspiration.

EJACULATIO, Ejaculation—*e*. *Seminis Impedita*, *Bradyspermatismus*.

EJACULATIO, *Ejaculat'io*, from *e*, and *jaculor*, (itself from *jacere*,) 'I throw.' *Gonobol'ia*, *Gonobol'ismus*, *Ejaculatio* seu *Profus'io* seu *Effus'io Sem'inis*, *Espatrat'io*, *Patrat'io*, *Spermobol'æ*. The emission of sperm. That which occurs during coition has been termed *Insemination*. The act, by which that fluid is darted out through the urethra.

EJACULATOR. Same etymon. That which effects the emission of sperm. See *Transversus perinæi*.

EJACULATOR SEMINIS, Accelerator urinæ.

EJAC'ULATORY, *Ejaculat'orius*, *Ejac'ulans*; same etymon. (F.) *Éjaculateur*, *Éjaculatoire*. Concerned in the ejaculation of sperm.

EJACULATORY DUCTS OR CANALS, (F.) *Conduits* ou *Canaux éjaculateurs*, are formed by the union of the vasa deferentia with the ducts of the vesiculæ seminales. They open at the lateral and anterior parts of the verumontanum, and convey into the urethra the sperm which is discharged from the vesiculæ, as well as that which comes directly from the testicle by the vas deferens. Between them there is often a depression, sometimes of a large size, which is termed *Utriculus*, *U. prostat'icus*, *Ves'ica* seu *vesic'ula prostat'ica*, *Sinus pocula'ris* seu *prostat'ata*, *Corpusculum Weberianum*, *Weber'ian organ* or *corpuscle*, which has been regarded as the analogue to the uterus in the female, and thence called *Uterus masculinus*.

EJECTIO, Excretion—*c*. *Fæcum*, Defecation.

EJECTION, *Ejec'tio*, from *ejicere*, *ejectum*, (*e*, and *jacere*,) 'to throw out or eject.' The excretion of the feces, urine, sputa, &c.

EL NISPERO, Sapota.

ELABORATIO, *Elaborat'io*, from *e*, and *laborare*, *laboratum*, 'to work.' This word is used, by physiologists, to signify the various changes which substances susceptible of assimilation undergo, through the action of living organs, before they are capable of serving for nutrition. The food is said to be *elaborated* in the stomach during the formation of chyme; the chyme is *elaborated* in the small intestine before it is chyle, &c.

ELEA, Olea Europæa.

ELEAGNUS, *Myrica gale*—*e*. *Cordo*, *Myrica gale*.

ELÆOCARPUS COPALLIFERUS, Copal.

ELÆOCÉROLÉ, Cerate.

ELÆOM'ELI, from ελαιον, 'oil,' and ἐλεῖν, 'honey.' Dioscorides means, by this, an oil thicker than honey, and of a sweet taste, which flows from the trunk of a tree in Syria. It is acrid and purgative; and sometimes occasions serious symptoms, according to that author.—Gorræus.

ELÆOM'ETER, from ελαιον, 'oil,' and μέτρον, 'measure.' A very delicate glass hydrometer, for testing the purity of olive or almond oil, by determining their densities. The 0 or zero of the scale is the point at which the instrument floats in the oil of poppy seeds. In pure olive oil, it floats at 50°, and the space between these points is divided into 50 equal parts, and numbered accordingly. It floats at 35° or 38½° in pure oil of almonds.

ELÆON, Oil.

ELÆOPH'ANES, from ελαιον, 'oil,' and φαίνωμαι, 'I appear.' That which has the appearance of oil; as *Uri'na elæoph'anes*.

ELÆO-SACCHARUM, from *ελαιον*, 'oil,' and *σακχαρ*, 'sugar.' *O'leo-Saccharum*, *Bal'samo-Saccharum*, (F.) *Oléosaccharolé*. A medicine, composed of essential oil and sugar. It was made by pouring upon an ounce of sugar, reduced to an impalpable powder, $\frac{3}{ss}$ or $\frac{3j$ of an essential oil; the mixture being triturated until the two substances were perfectly united.—Rollink.

ELAÏN, see Fat.

ELAIS GUINEENSIS, see *Cocos butyracea*—e. Occidentalis, see *Cocos butyracea*.

ÉLAN, *Cervus alces*.

ÉLANCÉMENT, see Lancinating.

ELAPHOBOSCUM, *Pastinaca sativa*.

ELAPHOMYCES GRANULATUS, *E. officinalis*, (*ελεφας*, 'a stag,' and *μυκης*, 'a fungus.') *Lycoperdon cervinum*, *Cervi Bole'tus*, *Bole'tus cervinus*, *Tuber cervinum*, *Hart's Truffles*, *Deer Balls*, *Lycoperdon nuts*. Nat. Order, Fungi. Formerly used in the preparation of a *Balsamus apoplecticus*, and to promote labour and the secretion of milk. Given, also, as an aphrodisiac. Dose, \mathfrak{z} ss, in powder.

ELAPHRIUM ELEMIFERUM, see *Amyris elemifera*—e. *Tomentosum*, *Fagara octandra*.

ELAPSUS, Luxation.

ELASMA, from *ελαυνω*, 'I drive up.' A clyster pipe.—Linden.

ELASTES, Elasticity.

ELASTICINE, see Fibrous.

ELASTICITY, *Elastic'ity*, *El'ater*, *Elas'tes*, *El'ates*, *Vis elastica*, *Tone*, *Tonic'ity*, (F.) *Contractilité de Tissu*, *Contractilité par défaut d'extension*, from *ελαυνω*, 'I impel,' *ελαστης*, 'a driver.' The property by which certain bodies return to their proper size and shape, where these have been modified by pressure or otherwise. It is possessed by the dead as well as by the living solid.

ELATE, *Pinus picea*—e. *Theleia*, *Pinus rubra*.

ELATER, Elasticity.

ELATERINE, see *Momordica elaterium*.

ELATERIUM, *Extractum elaterii*, *Momordica elaterium*—e. *Cordifolium*, *Momordica elaterium*.

ELATES, Elasticity.

ELATIN, see *Momordica elaterium*.

ELATINE, *Antirrhinum elatine*—e. *Cymbalaria*, *Antirrhinum linaria*—e. *Hastata*, *Antirrhinum elatine*.

ELBOCK, Elbow.

ELBOW, Sax. *elboza*, from *ell*, and *bow*; the bend of the ulna. *Cubitus*, *Ancon*, *Pechys*, *Ulna*, *Umbo*, (Sc.) *Elbock*, *Elbuck*, (F.) *Coude*. This word, abstractedly, means the angle formed by the union of two straight bodies. It is particularly applied to the articulation of the arm with the forearm, and especially to the projection formed by the olecranon process at the posterior part of the joint.

ELBOW, MINERS', see Miners' Elbow.

ELBUCK, Elbow.

ELCAJA, *Trichilia emetica*.

ELCOMA, Ulcer.

ELCOPLASTY, Helicoplasty.

ELCO'SIS, *Helco'sis*, *Elco'ma*, *Helco'ma*, from *ελκος*, 'an ulcer.' Ulceration in general. An ulcer. A deep ulceration of the cornea, in consequence of a blow or of violent inflammation.—Galen, Paré.

Sauvages applies the term to a state of the body, in which there are numerous ulcerations of an obstinate character, complicated with caries, putrescency, low fever, &c.

ELCUSTER, Embryuleus.

ELDER, *Sambucus*—e. Common, *Sambucus Canadensis*—e. Dwarf, *Aralia hispida*, *Sambucus ebulus*—e. Prickly, *Aralia spinosa*—e. Redberried, *Sambucus pubens*.

ELECAMPANE, *Ipula helenium*

ELECTRICITAS, Electricity—e. Animalis, Galvanism—e. Galvanica, Galvanism—e. Metallica, Galvanism.

ELECTRICITY, *Electric'ity*, from *ηλεκτρον*, 'amber,'—the substance in which it was first noticed. Electricity is used medicinally as an excitant. It has been occasionally employed with success in paralysis, rheumatism, accidental deafness, amaurosis, amenorrhœa, &c., but it is uncertain, and not much used; and the cases are not always clear in which it could be of service. It may be communicated by means of the *electric bath*—*Bain électrique*, as it has been called; which consists in placing the patient upon an insulated stool, and connecting him with the prime conductor, of which he thus becomes a part. The fluid may be communicated by points, sparks, or by shocks, according to the required intensity.

ELECTRICITY, CHEMICAL, Galvanism—e. Contact, Galvanism—e. Magnetic, Electro-Magnetism—e. Voltaic, Galvanism.

ELEC'TRIFY, from *electric*, and *fi*, 'I make.' (F.) *Électriser*. To produce the electrical condition in a body, or to render it susceptible of producing electrical phenomena. It is often used to signify the act of communicating the electric fluid to man.

ÉLECTRISER, to electrify.

ELECTRODES, from *ηλεκτρον*, 'amber,' and *ειδος*, 'resemblance.' An epithet for evacuations, which shine like amber.—Hippocrates, Fœsius.

ELECTRO-LITHOTRITY. The disintegration of calculi in the bladder by the mechanical force of the electrical discharge, applied by an appropriate apparatus.

ELECTRO-MAGNETISM, *Magnet'ic electric'ity*. An electro-magnetic apparatus is occasionally used in cases of paralysis. A convenient form consists of a battery of six curved permanent magnets, and an intensity armature, around whose cylinders 1500 yards of fine insulated copper wire are coiled. The ends of the wire communicate respectively, with a pair of directors, each of which holds a piece of sponge, dipped in vinegar or a solution of common salt. When the armature is rotated, and a portion of the body is interposed between the directors, a succession of shocks is received.

ELECTROPUNCTURE, *Electropunctura'tion*, *Electrostix'is*, *Galvanopuncture*. The operation of inserting two or more wires, and then connecting them to the poles of the galvanic apparatus.

ELECTROSTIXIS, Electropuncture.

ELECTRUM, Succinum.

ELECTUAIRE, Electuary—e. *Dentifrice*, *Dentifrice*—e. *Simple*, *Conserva*.

ELECTUARIUM, *Electa'rium*, *Opia'tum*, (Old Eng.) *Allectuary*, an *Electuary*, (F.) *Electuaire* *Saccharolé mou*, from *eligere*, *electum*, 'to make choice.' A pharmaceutical composition of a soft consistence, somewhat thicker than honey, and formed of powders, pulps, extracts, syrup, honey, &c. In the London and American Pharmacopœias, electuaries are classed under Confections.

ELECTUARIUM DE ALÔE, *Opiatum mesentericum*—e. *Aperiens*, *Confectio sennæ*—e. *Aromaticum*, *Confectio aromatica*—e. *Cassia*, *Confectio cassiæ*—e. *Cassia fistulæ*, *Confectio cassiæ*—e. *Cassia sennæ*, *Confectio sennæ*—e. *Cassia tamarindatum seu lenitum*, *Confectio cassiæ*—e. *Catechu*, *Confectio catechu compositum*—e. *Catholicum commune*, *Confectio sennæ*—e. *Cinchonæ*, *Opiatum febrifugum*—e. *de Croco emendatum*, *Confectio hyacinthi*—e. *Diaprunum*, *Confectio sennæ*—e. *Eccoproticum*, *Confectio sennæ*—e. *de Kinâ kinâ*,

Opiatum febrifugum—e. Laxativum, Confectio cassiæ—e. Lenitivum, Confectio sennæ—e. Opiatum, Confectio opii—e. Piperis, Confectio piperis—e. Scammonii, Confectio scammonii—e. Sennæ, Confectio sennæ—e. Sennæ compositum, Confectio sennæ—e. Sennæ cum pulpis, Confectio sennæ—e. Solidum, Tabella—e. Thebaicum, Confectio opii.

ELECTUARY, Electuarium—e. Lenitive, Confectio sennæ.

ELÉNCÉPHALE, from ελαιον, 'oil,' and εγκέφαλον, 'encephalon.' A fatty matter found by Couverbe in the encephalic neurine.

ELELISPHACOS, Salvia.

ELEMENT, *Elementum*, from ancient *eleo* for *oleo*, 'to grow' [?], *Principium*, *P. Primitivum*, *Stoicheion*. A simple, ultimate constituent or principle in the human body, which forms the basis of a fibre or tissue. Also, a constituent of a compound organ. The *inorganic* elements are simple principles. An *organic element*, *proximate principle* or *compound of organization*, results from the union of certain inorganic elements. Oxygen, hydrogen, carbon, and azote, are inorganic elements; fibrin, albumen, osmazome, &c., organic elements.

ELEMENTS, ORGANIC, Principles, immediate—e. Sarcous, see Sarcous.

ELEMENTARY CELL, see Cell.

ELEMENTUM ACIDIFICUM, Oxygen.

ELEMI, Amyris elemifera.

ELENIUM, Inula helenium.

ELEOCHARIS PALUS'TRIS, *Scirpus palustris*; indigenous. *Ord.* Cyperaceæ. The flowering tops, seeds and roots are astringent, and have been used in diarrhœa and hemorrhage.

ÉLÉOLÉS, Olea medicinalis.

ELEOSELINUM, Apium petroselinum.

ELEPHANTIASIS, *Elephantia*, *Elephantiasmus*, *Elephas*, *Lazarus morbus* seu *malum Phœnicæus morbus*, *Phymatosis Elephantiasis*, from ελεphas, ελεφαντος, 'an elephant.' Various affections have been described under this name, by adding an epithet. It is ordinarily and generically applied to a condition, in which the skin is thick, livid, rugous, tuberculate, and insensible to feeling.

ELEPHANTIASIS OF THE ANTILLES, *Barbadoes Leg*, *Glandular disease of Barbadoes*, (F.) *Jambes de Barbade*, is the Elephantiasis of many writers, *Bucne'mia*. It is characterized by the leg being enormously tumid and misshapen; skin thickened, livid, and rugous, often scaly; serotum, arms, or other parts sometimes participating in the affection. The *Bucne'mia Tropica*, *Cochin leg*, is an affection of this kind.

ELEPHANTIASIS ARABICA, *Tyriasis*, *Elephas*, *Elephantia*, *E. Indica*, *Elephantia Arabum*, *Lepra Arabum* (of some), *Maladie glandulaire*, *Ladrerie*, *Yava Skin* of the Polynesians [?]. In this the tubercles are chiefly on the face and joints. There is loss of hair, except on the scalp; voice, hoarse and nasal; and the disease is said to be contagious and hereditary. It most frequently attacks the feet; and gives the lower extremity a fancied resemblance to the leg of an elephant, whence its name. The seat of the disease seems to be in the lymphatic vessels and glands, and in the subcutaneous areolar tissue: the inflammatory condition of which is accompanied with general morbid symptoms. Medicine has little or no effect on this loathsome complaint. See Lepra.

ELEPHANTIASIS OF CAYENNE, *Mal rouge de Cayenne*, seems to be a variety of Lepra, characterized by red and yellow spots, occupying the forehead, ears, hands, loins, &c., afterwards ex-

tending and becoming scaly, with deformity of the parts where they are seated, particularly of the face; and ultimately producing cracks, ulcers, caries, and sometimes death.

ELEPHANTIASIS OF THE GREEKS, *E. Græcorum*, is probably the same disease as Lepra.

ELEPHANTIASIS OF INDIA is characterized by red, livid, or yellowish spots, slightly prominent, to which succeed indolent tumours, formed in the areolar texture. At a more advanced period the phalanges swell, and become ulcerated; the bones of the nose carious, the lips thickened, and emaciation gradually carries off the patient. It likewise belongs to lepra.

ELEPHANTIASIS ITALICA, Pellagra.

ELEPHANTIASIS OF JAVA is likewise a variety of lepra, characterized by large white tumours on the toes and fingers, resembling scrofulous tumefactions. These tumours ulcerate, and the ulcerations spread from the extremities towards the trunk, destroying even the bones. Amputation can alone arrest its progress. The disease is often accompanied by alopecia, and by an eruption of red spots.

ELEPHANTINUM EMPLASTRUM, *Ελεφαντινον*. An ancient plaster, composed of 30 parts of ceruss, 45 of wax, oil *℞ss*, water *℞j*.—Castelli. Oribasius and Celsus both mention a plaster of this name, but they are by no means similar in composition.

ELEPHANTOMMA, Buphthalmia.

ELEPHANTOPUS, (F.) *Éléphantope*; from ελεphas, 'elephant,' and πους, 'foot.' One affected with elephantiasis.

ELEPHANTUSIA, Phylephas macrocarpa.

ELEPHAS, Elephantiasis, Ivory.

ELETTARIA CARDAMOMUM, Amomum cardamomum.

ELEUTERIA, Croton cascarrilla.

ÉLÉVATEUR, Elevator—*É. Commun de l'aile du nez et de la lèvre supérieure*, Levator labii superioris alæque nasi—*É. de l'Œil*, Rectus superior oculi—*É. de la Paupière supérieure*, Levator palpebræ superioris.

ÉLÉVATOIRE, Elevator.

ELEVATIO UTERI. Ascent of the uterus. This displacement occasionally occurs, in the unimpregnated state, in connection with fibrous uterine, and enlarged ovarian tumours.

ELEVATOR, Levator, from *elevare*, (e, and *levare*), 'to lift up' (F.) *Élévateur*. A muscle, whose function it is to raise the part into which it is inserted. See Levator.

ELEVATOR, *Elevatorium*, *Vectis elevatorium*, *Élévatoire*. A name given to different surgical instruments employed for raising portions of bone which have been depressed, for raising and detaching the portion of bone separated by the crown of the trepan, and for removing stumps of teeth.

ELEVATOR ANI, Levator ani—e. Labii inferioris, Levator labii inferioris—e. Labii superioris proprius, Levator labii superioris proprius—e. Labiorum communis, Levator anguli oris—e. Oculi, Rectus superior oculi—e. Patientiæ, Levator scapulæ—e. Scapulæ, Levator scapulæ—e. Testiculii, Cremaster—e. Urethræ, see Transversus perinei.

ELEVATOR, COMMON. This is a mere lever, the end of which is somewhat bent and made rough, in order that it may less readily slip away from the portion of bone which is to be raised. The instrument is used, by forming a fulcrum for it, either on the hand which holds it, or upon the fingers of the other hand; or by making a fixed point for it on the edge of the opening made with the trephine.

ELEVATOR OF LOUIS differed from the last only

in the circumstance of the screw-peg being united to the bridge by a kind of pivot instead of hinge, so that greater extent of motion was permitted.

ELEVATOR OF PETIT consists of a lever mounted on a handle, and straight throughout its whole length, except at its very end, which is slightly curved, in order that it may be more conveniently put under the portion of bone intended to be elevated. The lever is pierced at various distances from its butt-end with several holes, intended for the reception of a movable screw-peg, fixed on the top of a kind of bridge. This part of the instrument consists of an arch, the ends of which are long, and covered with small pads, and on its centre is the screw-peg already mentioned. By means of these holes the arm of the lever can be lengthened at pleasure.

ELEVATOR, TRIPLOID, *Vectis triplœides*. This was so called from its consisting of three branches, uniting above in one common trunk. The latter part was traversed by a long screw, having below a kind of hook, and above a handle for turning it. By turning the screw, the hook was drawn up, and the bone thus elevated.

The simple lever is the only one now used, owing to the want of facility and advantages in the use of the others.

ÉLÈVE EXTERNE, see House-surgeon — *É. Interne*, House-surgeon.

ÉLEVURES (F.), *Efflorescentiæ*. A generic name, including all the exanthemata, in which there is tumefaction of the tissue of the skin. See Exanthem.

ELF-LOCK. A condition of the hair in which it is matted and twisted into locks, as in plica. Supposed, in olden times, to have been the work of elves or fairies.

ELIASTER, *Ilech*.

ELICHRYSUM, *Solidago virgaurea* — *e. Montanum*, *Antennaria dioica*.

ELICOIDES, *Capeolaris*.

ELIGMA, *Electos*.

ELIMINANTIA, *Ecetricia*.

ELIMINATIVES, *Ecetricia*.

ELIQUATIO, *Colliquation*.

ELIXA'TIO, *Epsœsis*, from *elixus*, (*e*, and *lix*, 'water, liquor, lye,') 'boiled,' 'sodden.' This word has been used synonymously with Decoction. The act of boiling.

ELIX'IR, *Istr*, *Izir*, *Quelles*, *Alex'ir*. The etymology of this word is not clear. Lemery derives it from *ἐλκω*, 'I extract;' and also from *αἰκω*, 'I aid.' Others believe it to be from Arabic, *al-ekvir*, or *al-ekvir*, 'chymistry.' An elixir is a medicine composed of various substances held in solution in alcohol. The name has been used, however, for preparations which contain no spirit of wine.

ELIXIR AC'IDUM HALLERI seu DIPP'LI, *E. Antipodag'ricum seu Antinephretic'um seu Sulph'urico-ac'idum, Gutta ac'idula ton'ica, Aqua Rabel'lii, Liquor ac'idus Halleri, Mistu'ra sulph'urico-ac'idula, Æther sulph'uricus ac'idus, Ac'idum sulph'uricum alcoholisatum, A. vitriol'icum vino'sum, Alcohol Sulphuricatum, A. sulph'uricum*, is a mixture of concentrated sulphuric acid and alcohol — in the *Eau de Rabel*, of one part of the former to three of the latter. It is employed as an astringent in hemorrhages, &c.

ELIXIR ALOES, *Tinctura aloes composita* — *e. Anthelminticum Succorum, Tinctura jalapii composita*.

ELIXIR, ANTI-ASTHMAT'IC, OF BOERHAAVE. This elixir is composed of alcohol, aniseed, camphor, orris, asarabacca root, calamus aromaticus, liquorice, and elecampane. It is given in asthma, in the dose of 20 or 30 drops.

ELIXIR ANTHYPOCHONDRIACUM, *Tinctura cin-*

chonæ amara — *e. Antinephreticum, E. acidum Halleri* — *e. Aperitivum, Tinctura aloes composita*.

ELIXIR, ANTISCROFULOUS OF PEYRIÈRE; composed of weak alcohol, subcarbonate of potassa, and gentian root. It is administered in scrofula.

ELIXIR, BOERHAAVE's, see *Tinctura aloes composita*.

ELIXIR, CARMINATIVE, OF SYLVIUS, *Tinctura carminativa Sylvi*.

ELIXIR CLAUDE'RI, CLAUDE'RS *Elixir*. Made by macerating for a few days carbonate of potassa 5j; aloes, guaiacum, myrrh, saffron, rhubarb bruised, of each 5j; water f3viij; and decanting. Dose, a tablespoonful as a laxative and emmenagogue.

ELIXIR, DAFFY'S, *Tinctura sennæ comp.* — *e. Danorum, E. pectorale regis Daniæ* — *e. Fœtidum, Tinctura castorei composita* — *e. of Garus, Tinctura de croco composita* — *e. Guaiaci volatilis, Tinctura Guaiaci ammoniata* — *e. Jalapæ compositum, Tinctura jalapii composita* — *e. of Long Life, Tinctura aloes composita* — *e. de Longue vie, Tinctura aloes composita* — *e. d'Or de M. le Général de la Motte, Tinctura seu Alcohol sulfurico-æthereus* — *e. Paregoric, Edinburgh, Tinctura opii ammoniata* — *e. Paregoricum, Tinctura camphoræ composita* — *e. Pectorale dulce, E. pectorale regis Daniæ*.

ELIXIR PECTORAL'E REGIS DA'NIÆ, *E. Danorum seu Ringelmanni seu ex succo glycyrrhizæ seu pectoral'e dul'cè seu è succo liquorit'icæ* : (*Succ. glycyrrhiz. p. 1; Aq. Fœnicul. p. 2; Alcohol ammoniat. p. 6.* A formula in many of the Pharmacopœias of continental Europe.) With the addition of opium it constitutes the *Elixir ammoniato-opiatum, Extractum thebaïcum ammoniacal'e* of some Pharmacopœias.

ELIXIR PROPRIETATIS, *Tinctura aloes composita* — *e. Purgans, Tinctura jalapii composita* — *e. Radcliffe's, see Tinctura aloes composita* — *e. Rhej dulce, Vinum rhej palmati* — *e. Ringelmanni, E. pectorale regis Daniæ* — *e. Roborans Whyttii, Tinctura cinchonæ amara* — *e. Sacrum, Tinctura rhej et aloes* — *e. Salutis, Tinctura sennæ comp.* — *e. Squire's, see Tinctura camphoræ composita* — *e. Stomachicum, Tinctura gentianæ composita* — *e. Stomachicum spirituosum, Tinctura cinchonæ amara* — *e. Stoughton's, see Tinctura gentianæ composita* — *e. ex Succo glycyrrhizæ, E. pectorale regis Daniæ* — *e. ex Succo liquorit'icæ, E. pectorale regis Danorum* — *e. Sulphurico-acidum, E. acidum Halleri* — *e. Traumaticum, Tinctura Benzoini composita*.

ELIXIR VITÆ OF MATH'OLUS; composed of alcohol and 22 aromatic and stimulating substances. It was formerly employed in epilepsy.

ELIXIR OF VITRIOL, Sulphuric acid, dilute — *e. Vitrioli, Sulphuricum acidum aromaticum* — *e. Vitrioli acido-aromaticum, Sulphuricum acidum aromaticum* — *e. Vitrioli dulce, Spiritus ætheris aromaticus, Sulphuricum acidum aromaticum* — *e. Vitrioli Edinburgensium, Sulphuricum acidum aromaticum* — *e. Vitrioli Mynsichti, Sulphuricum acidum aromaticum* — *e. of Vitriol, sweet, Spiritus ætheris aromaticus* — *e. Vitrioli cum tincturâ aromaticâ, Sulphuricum acidum aromaticum* — *e. of Vitriol, Vignani's, Spiritus ætheris aromaticus* — *e. Whyttii, Tinctura cinchonæ amara*.

ELIXIRIUM ANTISEPTICUM DOCTORIS CHAUSSIER, *Tinctura cinchonæ æthera composita*.

ELIXIS, *Electos*.

ELIXIVIATION, *Lixiviation*.

ELK, *Cervus alces* — *e. Bark, Magnolia macrophylla* — *e. Tree, Andromeda arborea* — *e. Wood, Andromeda arborea, Magnolia macrophylla*.

ELLOPLASTY, *Helcoplasty*.

ELLARNE, *Sambucus*.

ELLEBORASTER, *Helleborus fetidus*.

ELLEBORE, *Dracontium foetidum*.

ELLEBORISMUS, *Helleborismus*.

ELLEBORUM ALBUM, *Veratrum album*.

ELLEBORUS ALBUS, *Veratrum*—*c. Niger*, *Helleborus niger*.

ELLEN-TREE, *Sambucus*.

ELLER, *Alnus glutinosa*.

ELLERMAN'S DEODORIZING FLUID. An antiseptic or remover of odours—said to be a solution of a persalt of iron.

ELLET, *Sambucus*.

ELLIPTIC, Oval.

ELLYCHNIOTOS, from *ελλυχνιον*, 'the wick of a lamp.' A sort of tent, used by the ancient surgeons, so called because it was shaped like a wick, or because it was made of a similar material.

ELM, COMMON, *Ulmus*—*c. Red*, *Ulmus Americana*—*c. Rough-leaved*, *Ulmus Americana*—*c. Slippery*, *Ulmus Americana*.

ELMINTHOCORTON, *Corallina Corsicana*.

ELNORNE, *Sambucus*.

ELODES, *Helo'des*, *Paludal*, *Marshy*, (F.) *Mariéageux*, *Paludéen*, *Palustre*. *Febris elo'des seu helo'des seu paludal/su*, *Helop'yra*, *Helop'yretos*, from *ελος*, 'a marsh,' and *ειδος*, 'resemblance,' Marsh fever. (F.) *Fèvre intermittente paludéenne*, *F. paludéenne* ou *Maremmatique*, because it prevails in the Maremma district of Italy. Also, a kind of fever, characterized by great moisture or sweating.

ELOME, *Orpiment*.

ELONGATION, *Parathro'ma*, *Elongatio*, from *elongare*, *elongatum*, (*e*, and *longus*), 'to lengthen,' 'extend.' An incomplete luxation, in which the ligaments of an articulation are stretched and the limb lengthened, without total luxation. The word has also been used for the extension required in the reduction of fractures and luxations, and for the increased length of limb, (F.) *Allongement*, in diseases and dislocations of the hip-joint.

ELREN, *Sambucus*.

ELTH, see Age.

ELUTRIATIO, Decantation, Elutriation.

ELUTRIATION, *Elutritio*, originally from *eluo*, (*e*, and *luo*, 'I wash,') 'I wash away, I rinse.' In pharmacy a process by which the finer particles of a powder are separated from the coarser. It consists in diffusing the powder in water, allowing the larger and heavier particles to subside, and then decanting the liquor, that the finer particles may subside.

ELUVIES, same etymon. An inordinate discharge of any fluid, and also the fluid itself. In the works of some authors it is particularly applied to the mucus which flows from the vagina in cases of leucorrhœa.

ELUXATION, Luxation.

ELYTRATRESIA, Colpatresia.

ELYTREURYNTER, Speculum vaginæ.

ELYTRITIS, Colpositis, Leucorrhœa.

ELYTROBLENNORRHŒA, Leucorrhœa (vaginal).

ELYTROCELE, from *ελτρον*, 'a vagina or sheath,' and *κηλη*, 'a tumour.' Vogel has given this name to vaginal hernia, *Colpocèle*.

ELYTRODES (tunica), Vaginal coat of the testicle.

ELYTRODEMA, *Colecæde'ma*, *Colpæde'ma*, from *ελτρον*, 'a vagina or sheath,' and *οιδημα*, 'œdema,' 'Edema of the vagina.'

ELYTROITIS, Colpositis.

ELYTRON, from *ελωει*, 'I involve.' A sheath. The vagina. The membranes which envelope the spinal marrow are called *elytra*.

ELYTRONCUS, *Elytroph'y'ma*, from *ελτρον*, 'a vagina or sheath,' and *ογκος*, 'a tumour.' A swelling of the vagina.

ELYTROPAPPUS RHINOCERO'NIS. A South African bush, *Nat. Ord. Compositæ*, the

whole of which is bitter and resinous. The tops of the branches, infused in wine or brandy, make excellent bitters. The tops are also given in powder to children affected with diarrhœa.

ELYTROPHYMA, *Elytroncus*.

ELYTROPLAS'TY, *Elytrophlas'tic*; from *ελτρον*, 'the vagina,' and *πλασσω*, 'I form.' The operation for restoration of the vagina.

ELYTROPTOSIS, Prolapsus vaginæ.

ELYTRORRHAGIA, Colporrhagia.

ELYTRORRHAPHY, *Elyttrorrhaph'ia*, *Colporrhaphy*, *Colporrhaphy*; from *ελτρον*, 'the vagina,' and *ραφη*, 'a suture.' The operation of closing the vagina by suture in cases of prociencia uteri.

EM and EN, *εν*, 'in, into, within,' also 'excess:' frequently used in this last sense by Dr. Good. A common prefix, generally answering to the prefixes *im* and *in*, in English. In composition, before *β*, *π*, *φ*, *χ*, and *μ*, the *ν* is changed into *μ*; before *γ*, *κ*, *ξ*, and *χ*, into *γ*; before *λ*, into *λ*; and before *ρ* generally into *ρ*.

EMACIATE, *Maces'cere*, *Tabes'cere*, (F.) *Anaigrir*; from *emaciare*, *emaciatus*, (*e*, and *maceo*, 'to be lean,') 'to grow lean.' To lose flesh, to become lean, to waste away.

EMACIATION, *Emacia'tio*, *Extenua'tio*, *Ma'cies*, *Maceritudo*, *Maceror*, *Marces'ces*, *Skelet'ia*, *Leptym'us*, *Leptyn'sis*, *Pinguet'inis dimini'tio*, (F.) *Anaigrissement*, *Dépréissement*. That condition of the body, or of a part of the body, in which it grows lean. The state of one who is lean—*Leanness*, *Isch'notes*, (F.) *Maigréur*.

ÉMAIL DES DENTS, Enamel of the teeth.

ÉMAILLOTAGE (F.), from *maillo*, 'swathing or swaddling clothes.' The 'wrapping up' or 'packing up' in dry or wet sheets, which is practised in hydropathic establishments to induce sweating.

EMANA'TION, *Emana'tio*, *Efflu'vium*, from *emanare*, *emanatum*, (*e*, and *manare*,) 'to issue or flow from.' The term is applied to a body which proceeds or draws its origin from other bodies—such as the light which emanates from the sun; the miasm which arises from the putrid decomposition of animal or vegetable substances, &c. See Miasm.

EMAN'SIO MEN'SIUM, from *e*, and *manere*, *mansum*, 'to stay.' This term has been applied to amenorrhœa or obstruction of the menses before they have been established. Some have used it for the retention which occurs even after they have been established. The former is the general acceptance.

EMARGINATIO, *Échanerure*, Notch.

EMARGINATURA, *Échanerure*, Notch.

EMASCULATION, *Emascula'tio*, from *emasculare*, (*e*, and *masculus*, 'a male,') 'to render impotent.' The act of removing or destroying the generative organs of a male animal.

EMASCULATUS, Castratus, Eunuch.

EMBALMING, from *em*, *en*, 'in,' and *balsamum*, 'balsam.' *Balsama'tio*, *Smyrnis'mos*, *Cedai'a*, *Polliet'ia*, *Necroceti'a*, *Comdit'ura Cadaverum*, (F.) *Embaumement*, *Embalamation*. An operation which consists in impregnating the dead body with substances capable of preventing it from becoming putrid, and thus putting it in a condition to be preserved.

EMBA'MMA, *Apoban'ma*, *Bamma*, from *εμβαπτω*, (*εμ*, and *βαπτω*, 'I dip,') 'I immerse.' A medicated pickle or sauce.—Gorræus.

EMBARRAS, Emphraxis.

EMBARRAS GASTRIQUE (F.), Gastric disorder or impediment, *Collic'ies gastrica*, *Sordes primar'um viarum*, *Status gastricus*, *Gaster atax'ia*, (F.) *Gastricité*. Disorder of the stomach, in which there is loss of appetite, with bitterness

and clamminess of mouth, white and yellow tongue, feeling of oppression in the epigastrium, and sometimes pain in that region, nausea and bilious or bitter vomiting; this state being accompanied with headache, lassitude, and pain in the back and limbs.

EMBARRAS GASTRO-INTESTINAL, *Gastro-intestinal disorder*. Slight gastro-enteritis, according to the Broussaists, in which the symptoms of the *Embarras gastrique* and *E. intestinal* are united.

EMBARRAS INTESTINAL, *Intestinal disorder*. The principal characters assigned to this are:—tension of the abdomen colic, borborygmi, discharge of flatus *per anum*, constipation or diarrhoea, &c.

EMBARRURE, Engisoma.

EMBAULEMENT, Embalming.

EMBOÏTEMENT (F.). 'Encasing,' from *em*, 'in,' and *boîte*, 'a box.' *Enadelphi'a*, *Endadelphi'a*, *Enteradelphi'a*. This term has been applied to the theory of generation which considers that the germs are encased in each other in the ovary of the female, in such sort that they are developed in succession after impregnation. It is the theory of encasing of germs, or of monotrosy by inclusion.

EMBOÏTURE, Diarthrosis.

EMBOLE, from *εμβάλλω*, (*εμ* and *βάλλω*), 'I put in place.' Reduction of a luxated bone. *Repositio*. Also, a plug, as of a blood-vessel.

EMBOLIA, (F.) *Embolies*. Diseases produced by plugging or obturation of blood-vessels.—Virchow.

EMBOLIMOS, Intercalary.

EMBOLUM CEREBRI, Infundibulum of the brain.

EMBONPOINT, (F.) 'In good point or plight.' *Bona corporis habitudo*. The state of the animal body when in full health. Excessive *embonpoint* constitutes corpulence and obesity, and may become a morbid condition.

EMBORISMA, Aneurism.

EMBOWELLING, Exenterismus.

EMBREGMA, Embrocation.

EMBROCATIO, Embrocation, Irrigation.

EMBROCA'TION, *Embroca'tio*, *Embreg'ma*, *Embrochê*, *Implu'vium*, from *εμβρέχω*, (*εμ*, and *βρέχω*, 'I wet,' 'I dip in.' A fluid application to be rubbed on any part of the body. It is often used synonymously with liniment. Originally employed in the sense of *Fomentation*.

EMBROCATION, GUESTONIAN, see *Oleum Terebinthinæ rectification*.

EMBROCHE, Embrocation.

EMBRYO, *Em'bryon*, from *ev*, 'in,' and *βρῦν*, 'I grow.' *Cye'ma*. The fecundated germ, in the early stages of its development in utero. At a certain period of its increase, the name *fœtus* is given to it, but at what period is not determined. Generally, the *embryo state* is considered to extend to the period of quickening.

EMBRYOC'TONY, *Fœtus trucidatio*, from *εμβρῦον*, 'the embryo,' and *κρονος*, 'destruction.' The act of destroying a fœtus in utero, when insurmountable obstacles—as certain deformities of the pelvis—oppose delivery.

EMBRYOGRAPHY, *Embryogra'phia*, from *εμβρῦον*, 'the embryo,' and *γραφία*, 'a description.' The part of anatomy which describes the embryo.

EMBRYOLOG'Y, *Embryolog'ia*, from *εμβρῦον*, 'the embryo,' and *λογος*, 'a description.' The doctrine of embryonic description.

EMBRYON'IC, *Embryon'icus*, (F.) *Embryonique*, *Embryonnaire*; same etymon as *embryo*. Relating or appertaining to an embryo:—as 'embryonic life.'

EMBRYONIQUE, Embryonic.

EMBRYONNAIRE, Embryonic.

EMBRYOTHLAS'TA, *Embryothlas'tes*, *Embryothlas'tum*, (F.) *Embryotome*, from *εμβρῦον*, 'the embryo,' and *θλαω*, 'I break.' An instrument for dividing the fœtus piecemeal, in order to effect delivery. A crotchet or other instrument, used, in certain cases of laborious parturition, to break the bones of the fœtus, for the purpose of extracting them with greater facility.—Hippocrates, Galen.

EMBRYOTOCIA, Abortion.

EMBRYOTOME, *Embryothlasta*.

EMBRYOT'OMY, *Embryotom'ia*, from *εμβρῦον*, 'the embryo,' and *τεμνω*, 'to cut.' A division of the fœtus into fragments, to extract it by piecemeal, when the narrowness of the pelvis or other faulty conformation opposes delivery.

EMBRYOT'ROPHY, *Embryotroph'ia*, from *εμβρῦον*, 'the embryo,' and *τροφία*, 'nourishment.' Fœtal nutrition; *Cyot'rophy*.

EMBRYUL'CIA, *Embryusterul'cia*, from *εμβρῦον*, 'embryo, fœtus,' and *ελκω*, 'I extract,' 'I draw.' A surgical operation, by which the fœtus is extracted by means of appropriate instruments, when faulty conformation or other circumstances prevent delivery by the natural efforts.

EMBRYUL'CUS, *Eleus'ter*, same etymon. An iron hook or crotchet, described by Fabricius ab Aquapendente, which was used to extract the fœtus in some cases of laborious labour.

EMBRYUSTERULCIA, *Embryulcia*.

EMENDANS, Corrigent.

ÉMERAUDE, Smaragdus.

EMERODS, Hæmorrhoids.

EMESIA, Vomiturition.

EMESIS, Vomiting.

EMETATROPH'IA, from *εμεω*, 'I vomit,' and *τροφία*, 'want of nourishment.' Atrophy induced by vomiting.

EMETIA, Emetine.

EMET'IC, *Emet'icum*, *Vomit'rium*, *Vom'itory*, *Vomit*, *Puke*, (Prov.) *Pick*, from *εμεω*, 'I vomit.' A substance capable of producing vomiting. (F.) *Émétique*. [This term is also restricted by the French to tartarized antimony—the emetic, as it were, *par excellence*.] *Vomitif*.

Tartarized antimony, emetine, ipecacuanha, and sulphate of zinc, are the chief emetics. They are valuable agents in disease, and may either act *primarily* on the stomach, or *secondarily* on other parts of the system,—the sympathy between the stomach and other parts of the body being very extensive, and an important object of study.

The following are the usual emetics:—Antimonii et Potasse Tartaras; Cupri Acetas; Cupri Sulphas; Emetina; Gillenia; Hydrargyri Sulphas Flavus; Ipecacuanha; Lobelia; Sanguinaria; Scilla; Sinapis, and Zinci Sulphas.

EMETIC ROOT, *Euphorbia corollata*—*e. Tartar*, *Antimonium tartarizatum*—*e. Weed*, *Lobelia inflata*.

EMETICOLOGIA, Emetology.

EM'ETINE, *Emet'ina*, *Emet'ia*, *Vom'itine*, *Cephælinum*. A vegetable alkali, discovered by M. Pelletier in ipecacuanha, and to which it owes its emetic power. It is obtained from different ipecacuanhas, but chiefly from *psychot'ria emet'ica*, *callicoc'ca ipecacuan'ha*, and *vi'ola emet'ica*. It is in transparent scales, of a reddish-brown colour, almost odorless, of a bitter, slightly acid taste; is unchangeable in the air, soluble in water and alcohol, and insoluble in ether. Three grains of impure emetia or one grain of pure are equal to 18 of ipecacuanha. See *Ipecacuanha*.

ÉMÉTIQUE, *Antimonium tartarizatum*, Emetic.

ÉMÉTISER (F.). To cause vomiting by

emetics. In French pharmacy, it is, likewise, used to signify the addition of *émétique*—tartrate of antimony and potassa—to a medicinal preparation.

EMETO-CATHARSIS, see Emeto-cathartic.

EMETO-CATHARTIC, *Em'eto-cathar'ticus*, from *emetos*, 'vomiting,' and *καθαρτικός*, 'a purgative.' A remedy, which at the same time excites vomiting and purging—*Em'eto-cathar'tis*.

EMETOL'OGY, *Emetolog'ia*, *Emeticolog'ia*, from *emetos*, 'vomiting,' and *λογος*, 'a discourse.' A treatise on vomiting, and on emetics.

EM'INENCE, *Em'inen'tia*, *Protuberant'ia*, *Ex'archē*, *Evanthē'ma*. A projection at the surface of a healthy or diseased organ.

ÉMINENCE PORTE ANTÉRIEURE, Lobulus anonymus—*é. Porte postérieure*, Lobulus Spigelli—*é. Sui-pubienne*, Mons veneris.

ÉMINENCES BIGÉMINÉES, Quadrigemina corpora—*é. Latérales*, Corpora olivaria—*é. Pyramidales*, Corpora pyramidalia—*é. Vermiformes du cerellet*, Vermiformis processus.

EMINENTIA, Protuberantia—*e. Annularis*, Pons varolii—*e. Pyramidalis*, Crista vestibuli—*e. Thyreoidica*, Pons Adamii—*e. Tympani*, Pyramid.

EMINENTIE BIGEMINE, Quadrigemina tubercula—*e. Candicantes*, Mammillary Tubercles—*e. Lenticulares*, Corpora striata—*e. Magnæ cerebri*, Thalami nervorum opticoium—*e. Medianæ Internæ*, Corpora pyramidalia—*e. Olivares*, Corpora olivaria—*e. Ouales Laterales*, Corpora olivaria—*e. Pyramidales*, Corpora pyramidalia—*e. Quadrigeminæ*, Quadrigemina tubercula—*e. Teretes*, Processus teretes.

ÉMISSAIRE, Emuncitory,

ÉMISSAIRE SANTORINI, (F.) *Émissaires de Santorini*. A name given to some small veins, which communicate with the sinuses of the dura mater, by apertures in the cranium. Such are the parietal, posterior condyloid, mastoid veins, &c.

EMISSION, Emission—*e. Seminis*, Spermatismus.

EMIS'SION, *Emis'sio*, from *emittere*, (*e*, and *mittere*, *missum*), 'to send out,' 'drive out.' The act by which any matter whatever is thrown from the body. Thus, we say *Emission of urine*, *Emission of semen*, &c.

ÉMISSION SANGUINE, Bloodletting.

EMMENAGOGUES, *Emmenago'ga*, *Emmeni-agō'ga*, *Ame'nia*, from *εμμηνα*, 'the menses,' and *αγω*, 'I drive,' or 'expel.' *Men'agogues*. A name given to medicines believed to have the power of favouring the discharge of the menses. There is probably no substance which possesses this power directly. According to different conditions of the system, the most opposite remedies may act as emmenagogues. *Black hellebore*, *savin*, *madder*, *polygala senega*, and *ergot*, are reputed to be specific emmenagogues. The following list comprises the chief of them:—*Canila Pulegioides*; *Helleborus Niger*; *Montha Pulegium*; *Rosmarinus*; *Rubia*; *Sabina*; *Secale Coranum*; *Senega*, and *Tanaacetum*.

EMMENAGOLOG'IA, from *εμμηναγωγία*, and *λογος*, 'discourse.' A treatise of emmenagogues.

EMMENIA, Menses.

EMMENIAGOGA, Emmenagogues.

EMMENOLOG'IA, from *εμμηνα*, 'menses,' and *λογος*, 'a discourse.' A treatise on menstruation.

EMMENOLOG'ICAL, *Emmenolog'icus*: same etymon as the last. Relating or appertaining to menstruation.

EMMYXIUM ARTICULARE, Hydrarthrus.

EMOLLIENTIA, Emollients.

EMOLLIENTS, *Emollien'tia*, *Malthac'tica*, *Relaxant'ia*, *Epicacanth'ia*, *Malac'tica*, *Lubricant'ia*, *Malacoep'ia*, *Mollien'tia*, from *emollire*, 'to soften,' 'to mollify,' 'to soften.' Substances which

relax and soften parts that are inflamed, or too tense. They are used both internally and externally: as the former, however, consist of mucilaginous substances, they are generally reckoned as demulcents: the latter, or proper emollients, consist of oils, cataplasms, fomentations, &c. Oleaginous bodies, rubbed on a part, act by diminishing its cohesion. Fomentations, in cases of internal inflammation, act probably through contiguous sympathy. The following are the chief emollients:—*Adeps*; *Amygdala Oleum*; *Avena Farina*; *Cera*; *Cetaceum*; *Linum*; *Olivæ Oleum*; *Sesamum*; *Triticum Farina*, and *Ulmus*.

EMOLLITIES, Mollities—*e. Morbosa*, Mollities—*e. Ossium*, Mollities ossium—*e. Uteri Morbosa*, Hysteromalacia.

EMOLLITIO, Mollities—*e. Ventriculi*, Gastro-malacia.

ÉMONCTOIRE, Emuncitory.

EMOPTOE, Hæmoptysis.

EMOTIO, Delirium, Luxation, Passion.

EMOTIONAL, from *emotio*, (*e*, 'from,' and *moveo*, *motum*, 'to move.') Relating to emotion or passion independently of the will:—hence an 'emotional' or instinctive impulse.

EMPASMA, Cataplasma.

EMPÂTEMENT, from *empâter*, 'to render pasty or doughy.' A non-inflammatory engorgement, which retains, more or less, the impression of the finger.

EMPATHE'MA, *Ma'nia a pathe'matē*, (F.) *Manie sans délire*; ungovernable passion; from *εμ*, 'in,' and *παθος*, 'suffering.' Fixed delusion.

EMPEIRIA, Experience.

EMPETRUM, Herniaria glabra.

EMPHLYSIS, from *εμ*, 'in,' and *φλυσις*, 'a vesicular tumour or eruption;' *Ich'orous Exan'them*. An eruption of vesicular pimples, filled progressively with an acid and colourless, or nearly colourless, fluid; terminating in scurf or laminated scabs. A genus in the order *Exanthematica*, class *Hæmatica* of Good.

EMPHLYSIS APHTHA, Aphthæ, Stomatitis, aphthous—*e. Erysipelas*, *Erysipelas*—*e. Miliaria*, *Miliary fever*—*e. Pemphigus*, *Pemphigus*—*e. Vaccina inserta*, *Vaccina*—*e. Varicella*, *Varicella*.

EMPHRACTIC, Emphracticus.

EMPHRACTICA, Phyesonia.

EMPHRACTICUS, *Emphrac'tic*, from *εμφραττω*, (*εμ*, and *φραττω*, 'I shut up,' 'I close,' 'I obstruct.' *Emphas'ticus*, *Emplatton'enos*. Any substance which, when applied to the skin, was presumed to close the pores.

EMPHRAG'MA, same etymon. *Obturatorium*, *Impedimentum*. Anything that obstructs. Hippocrates uses this term to designate the obstacle to delivery on the part of the fœtus, when the presentation is preternatural.

EMPHRAGMA LACRYMALE, Fistula lacrymalis—*e. Salivare*, *Ranula*.

EMPHRAX'IS, *Obstru'ctio*, *Obtura'tio*, *Oppila'tio*, *Infarc'tus*, *Infarc'tio*, *Farc'tus*, *Fartus*, *Infarc'tio*. Same etymon. 'Obstruction.' An Embarras or repletion of canals or cavities by any substance, which is either morbid from quantity or quality.

EMPHRAXIS HEPATIS, Hepatempyphaxis.

EMPHYMA, Tumour—*e. Encystis*, *Encystis*—*e. Encystis atheroma*, *Atheroma*—*e. Encystis ganglion*, *Ganglion*, *Testudo*—*e. Encystis meliceris*, *Meliceris*—*e. Encystis steatoma*, *Steatoma*—*e. Exostosis ossea*, *Exostosis*—*e. Exostosis periosteæ*, *Node*—*e. Sarcoma*, *Sarcoma*—*e. Sarcoma adiposum*, *Adipose sarcoma*—*e. Sarcoma cellulolum*, *Cystic sarcoma*—*e. Sarcoma mammarum*, *Mammary sarcoma*—*e. Sarcoma pancreaticum*, *Pancræatic sarcoma*—*e. Sarcoma scirrhusum*, *Scirrhus sarcoma*—*e. Sarcoma tuberculosum*,

Tuberculate sarcoma—*e. Sarcoma vasculosum*, Sarcoma, vascular.

EMPHYSEMA, from *εμφύσω*, (*εν*, and *φύσω*, 'I blow,' 'I inflate,' *Infusio*, *Empneumatosis*, *Sarcitis flatuosa*, *Emphysema cellulare seu pneumatica*, *Physoncus*, Tumor flatulentus, *Pneumatica*, *Inflatio*, Wind-dropsy, (F.) *Emphysème*, *Aërethymie*.—Piorry. This term is commonly applied to any white, crepitant, shining, elastic, indolent tumour of the integuments, caused by the introduction of air into the areolar texture, *Aërodermea'sia*. Injuries of the larynx, trachea, or lungs; fractures of the ribs, or wounds penetrating the chest, are the most frequent causes of this affection, which is owing to the air escaping from the air-passages, and insinuating itself into the areolar texture surrounding the wound. There are some cases of emphysema, which are owing to internal causes; and hence a division has been made into the *accidental* and *symptomatic*, and the *spontaneous* and *idiopathic*.

EMPHYSEMA ABDOMINIS, Tympanites.

EMPHYSEMA OF THE LUNGS, *E. Pulmonum*, *Asthma aëreum ab Emphysematè Pulmonum*, *Pneumatosis Pulmonum*, *Pneumocœstis*, *Pneumocœstis*, (F.) *Pneumocœstic*, *Emphysème du Poupon*. A considerable pressure or contusion of the chest, or any violent concussion of the lung, may produce a laceration in that viscus, without injury being done to the parietes of the thorax, and may give place to the infiltration of air into the areolar texture, *interlobular emphysema*. Laënnec has described another species of emphysema of the lungs, *Vesicular emphysema*, *Aëropneumonie*, *Aëropneumocœstis* of Piorry, consisting in excessive dilatation of the air-cells, some of which become as large as hemp-seed, ultimately break, and give rise to irregular vesicles at the surface of the lung, some of which are as large as a hazel-nut. *Physical signs*. The thorax externally, generally or locally, appears unnaturally convex and prominent. The intercostal spaces are widened, but depressed. The inspiratory efforts are increased. The sound on percussion is morbidly clear, but not tympanitic. On auscultation, the inspiratory murmur is feeble or suppressed. The expiration, which is more frequently audible, is prolonged, laborious and wheezing. There are no certain physical signs which can distinguish *interlobular emphysema* from the *vesicular*.

Emphysema under the pleuro-pulmonalis is termed, by Piorry, *Aërethymopneumonie*.

EMPHYSEMA PECTORIS, Pneumothorax—*e. Scroti*, Physocœle—*e. Tympanites*, Tympanites—*e. Uteri*, Hysterophysis, Physometra.

EMPHYSEMATOUS, *Emphysematous*; same etymon. Relating or appertaining to emphysema.

EMPHYSEME, *Emphysema*—*e. du Poupon*, *Emphysema* of the lungs.

EMPHYTUM THERMUM, *Emphytnium*.

EMPIRIC, *Empiricus*, from *ἐμπειρία*, (*εν*, and *πειρα*, 'a trial,' 'experience.' One who follows only experience. A sect of physicians, who rejected all theory, and took for their guide experience alone. It was opposed to the dogmatic sect. The Empiric sect prevailed till near the time of Galen. Among its most eminent members, after Philinus and Serapion, were Apollonius, Glaucias, Bacchius of Tanagra, and Zeuxis, both disciples of Herophilus,—Heraclides of Tarentum, Cleophantus, master of Asclepiades, Menodotus of Nicomedia, and Theudas of Laodicea. They occupied themselves, chiefly, with discovering the properties of drugs, and did important service, in this manner, to medicine.

At the present day, the word *Empiric* is only taken in a bad sense, being employed in nearly the same signification as *charlatan* or *quack*.

EMPIRICE, Empiricism.

EMPIRICISM; same etymon. *Empirica Ars*, *Empiricè*, (F.) *Empirisme*. Medicine founded on experience. It generally, at the present day, signifies *quackery*.

EMPIRISME, Empiricism.

EMPLASTICUS. Same etymon as the next. Also, a remedy which adheres, as a plaster, to the surface, and in this manner may obstruct the pores: an emphractic.

EMPLASTRO-ENDERMIC, Endermic.

EMPLASTRUM, from the Greek *επιλασσω*, (*εν*, and *πλάσσω*, 'to form,' 'I spread upon,' (F.) *Emplâtre*, *Stéaraté*. A solid and tenacious compound, adhesive at the ordinary heat of the human body. Some of the plasters owe their consistence to wax and resin; and others to the chemical union which takes place between the semivitreous oxide of lead and oil. Most of them become too consistent by age. When such is the case, they may be remelted by a gentle heat, and oil be added to them.

EMPLASTRUM ADHÆSIVUM, *Emplastrum resinæ*—*e. Adhæsivum* Woodstockii, see *Sparadrapum Adhæsivum*—*e. Alexandri*, Alexandrine—*e. de Althæâ*, Unguentum de Althæâ.

EMPLASTRUM AMMONIACI, *Ammoniæ Plaster*, (*Ammoniæ pur. 3v, acidi aceticæ dil. Oss.* Dissolve the ammoniac in the vinegar and strain: then evaporate the solution by means of a water-bath, stirring constantly until it acquires a proper consistence.—*Ph. U. S.*) It is used as a discutient plaster.

EMPLASTRUM AMMONIACI CUM HYDRARGYRO, *Ammoniæ Plaster with mercury* (*Ammoniæ 1bj, hydrarg. 3ij, olei oliv. f 3j, sulphur. gr. viij.* Rub the mercury with the oil until the globules disappear, then add the ammoniac, previously melted, and mix. *Ph. U. S.*)

EMPLASTRUM ANDRÉE DE CRUZE, *E. de pice et resinis glutinans*—*e. Anglicum*, see *Sparadrapum adhæsivum*.

EMPLASTRUM AROMATICUM, *Aromatic Plaster*. (*Thuris 3ij, cera flava 3ss, pulv. cort. cinnam. 3vj, ol. ess. Piment, ol. ess. Limon. aa 3ij.* Melt the frankincense and wax together, and strain; then add, as it cools, the cinnamon, previously rubbed with the oils, and form a plaster. *Ph. Dubl.*) Used as a stimulating plaster.

A spice plaster, made by incorporating powdered cinnamon and any other spices with melted suet, has been used in cholera infantum. It is spread on leather or linen, and is renewed twice in the twenty-four hours.

EMPLASTRUM ASSAFÆTIDÆ, *E. antihystericum*, *Assafetida plaster*, (*Emp. plumb, Assafet, sing. 1bj, Galban, Cera flava, aa 1bss, Alcohol. dilut. Oij.* Dissolve the assafetida and galbanum in the alcohol, in a water-bath, strain while hot, and evaporate to the consistence of honey; add the lead plaster and wax previously melted together, stir the mixture well, and evaporate to the proper consistence.—*Ph. U. S.*) Used as an antispasmodic and anodyne plaster.

EMPLASTRUM ATTRAHENS, *E. ceræ*—*e. Auriculare*, Hypotium.

EMPLASTRUM BELLADONNÆ, *Plaster of Belladonna*. (*Empl. resin. 3ij, Ext. Belladon. 3iss.* Add the extract to the plaster, previously melted in a water-bath, and mix.—*Ph. U. S.*) An anodyne application in neuralgia and rheumatism.

EMPLASTRUM CALEFACIENS, *E. Piciæ cum cantharidæ*, (*Ph. U. S.*) *Calefacient plaster*, Warm plaster, (F.) *Emplâtre échauffant*. (*Cerat. cantharid (Ph. U. S.), 1bss, Piciæ abiet. 1ijss.* Melt together, and form into a plaster. *Ph. U. S.*) It is rubefacient and stimulant.

EMPLASTRUM CANTHARIDIS, *E. lyttæ*—*e. Can.*

tharidis vesicatorii compositum, E. lyttæ comp.—e. Cephalicum, E. picis comp.

EMPLASTRUM CERÆ, *E. simplex, E. At'trahens. Wax plaster. (Cera flava, See Præp. aa lbij. resina flava lbj. Melt them together and strain. Ph. L.)* It has been considered drawing. It is stimulant.

EMPLASTRUM CICIUTÆ, *E. de Cicutâ seu conii maculati, Unguentum salinum de cicuta, Hemlock plaster, (F.) Emplâtre de Ciguë. (Resina abiet. 960 p. Cera flav. 640 p. Picis albæ, 448 p. Ol. cicuta per decoct. præparat. 128 p. Fol. cicut. recent. 2000 p. Melt the resins, wax and oil: add the cicuta leaves, and boil: strain and add, after having dissolved it in vinegar of squills and cicuta juice, gum ammoniac. 500 p. Ph. Par.)* It is used as a discutient, especially to scirrhus tumours.

EMPLASTRUM CITRINUM, Unguentum de althæâ —e. Commae, E. Plumbi—e. Commune cum resinâ, E. resinæ—e. Conii, E. cicutæ.

EMPLASTRUM CUMINI, *Cumin plaster. (Cumin. semin., cuius sem., lauri bacc. sing. 3ij. picis arida lbij, cera flava 3ij. Melt the pitch and wax together, then add the other ingredients. Ph. L.)* It is used as a warm discutient plaster.

EMPLASTRUM DIACHYLON, Diachylon, E. plumbi —e. Divinum Nicolai, see Magnet—e. Emolliens, Unguentum de althæâ—e. Epispasticum, E. lyttæ.

EMPLASTRUM FERRI, *E. rob'orans, Iron Plaster, Strengthening Plaster. (Ferri subcarb. 3ij, Emp. plumbi, lbj, Picis abietis, lbss. Add the subcarbonate of iron to the lead plaster and Burgundy pitch previously melted together, and stir constantly until they thicken upon cooling.—Ph. U. S.)*

EMPLASTRUM FERRI RUBRI, *E. oxidi ferri rubri —e. Flavum, Unguentum de althæâ—e. ad Fonticulos, Ceratum Galeni.*

EMPLASTRUM GALBANI, *Galbanum plaster, (Empl. litharg. lbij, gum. galban. lbss, cera flava 3iv. Melt the galbanum before adding the plaster and wax, then melt all together. Ph. D.)* A stimulant and discutient.

EMPLASTRUM GALBANI COMPOSITUM, *E. lithargyri compositum, Diachylon magnum cum gummi, Compound galbanum plaster. (Galb. 3vij, Emp. plumbi, lbij, tereb. vulg. 3x, abietis resin. contus. 3ij. Melt the galbanum and turpentine together, then mix in the resin, and afterwards the plaster, previously melted. Ph. U. S.)* It is stimulant and discutient.

EMPLASTRUM GLUTINOSUM, see Sparadrapum adhasivum—e. cum Gummatibus, E. gummosum.

EMPLASTRUM GUMMOSUM, *Emplastrum lithargyri cum gummi seu cum gummatibus seu e gummatibus resinosus seu Oxidi Plumbi semivivæ rei gummosum, Gum plaster, Diachylon cum gummi, Yellow or gum diachylon. (Emp. plumbi p. viii, ammoniac, galbani, cera flava, aa, p. i. Ph. E.)* Properties and uses like the last.

EMPLASTRUM E GUMMATIBUS RESINOSIS, E. gummosum.

EMPLASTRUM CUM GUMMI-RESINIS, (F.) *Emplâtre de gomme résine, Diachylon gommé, Plaster of gum resins. (Emplastr. simpl. p. 1600, cera flava, p. 96, picis albæ, p. 96, terebinth. p. 96. Melt by a gentle heat, and add gum ammoniac, p. 32, bdellium, 32, galban. 32, and sagapenum, p. 32. Dissolve in a sufficient quantity of alcohol, evaporate to the consistence of honey, and mix carefully all together. Ph. P.)* A discutient.

EMPLASTRUM HYDRARGYRI, *Emplastr. lithargyri cum hydrargyro, Mercurial plaster. (Hydrarg. 3vj, ol. olive, resin. aa 3ij, emplastr. plumbi, lbj. Melt the oil and resin together, and when cool rub the mercury with them till the globules disappear; then gradually add the lead plaster,*

previously melted, and mix all. *Ph. U. S.)* It is stimulant, resolvent, and discutient. Applied as a discutient to venereal and other tumours.

EMPLASTRUM HYDRARGYRI COMPOSITUM, *E. de hydrargyro compositum seu de Vignæ cum mercurio emendatum, Compound plaster of mercury. (Empl. simpl. p. 1250, cera flava, p. 64, resina, p. 64. Melt, and before it congeals add pulv. g. ammoniac, p. 20, bdellium, p. 20, alban. p. 20, myrrh, p. 20, saffron, p. 12. Mix carefully, Take of mercury, p. 380, pure turpentine, p. 64, liquid and pure storax, p. 192. Triturate in an iron mortar till the globules disappear: melt all together, and before congealing add essential oil of Lavender, p. 8. Ph. P.)* The same properties as the last.

EMPLASTRUM ICHTHYOCOLLE TELÆ INDUCTUM, see Sparadrapum adhasivum—e. Irritans, E. lyttæ —e. Lithargyri, E. Plumbi—e. Lithargyri compositum, E. galbani compositum—e. Lithargyri cum gummi, E. gummosum—e. Lithargyri cum hydrargyro, E. hydrargyri—e. Lithargyri cum resinâ, E. resinæ—e. Lithargyricum cum resinâ pini, E. resinæ.

EMPLASTRUM LYTTÆ, *E. cantharidis seu cantharidis vesicatorii seu meloës vesicatorii, E. vesicatorium, E. epispasticum seu e cantharidis ubi epispasticum solidius et tenacius hærens seu irritans seu rubeficiens, Plaster of the Spanish or blistering fly, Blistering Plaster, Fly Plaster. (Cantharides, in powder, lbj, Emp. cera lbss, adipis. præp. lbj. Melt the plaster and lard together, and as the mixture becomes thick, on cooling, sprinkle in the flies, and mix. Pharm. L.)* This is the common blistering plaster. Too much heat must not be used in its preparation. It requires to remain on six or eight hours before its full effect is induced, and it acts sufficiently well, provided even a piece of thin gauze or tissue paper be placed between it and the skin, whilst absorption of the flies is thus prevented. See Blister. The Blistering Plaster of the Pharmacopœia of the United States, *Ceratum Cantharidis*, is made as follows:—*Cantharid. in pulv. subtiliss. lbj, cera flava. resinæ, aa 3vij, adipis, 3x.* To the wax, resin, and lard, previously melted, add the Spanish flies, and stir the mixture constantly until cool.

EMPLASTRUM LYTTÆ COMPOSITUM, *E. cantharidis vesicatorii compositum seu meloës vesicatorii compositum, Compound plaster of cantharides or Spanish flies. (Resin. liq. pini laricis. p. 18, resinæ concret. pini abietis, meloës vesicat. aa p. 12, cera flava, p. 4, subacet. cupri, p. ij, semin. sinapis alb., fruct. pip. nigr. aa p. j. Melt the pitch and wax, then add the turpentine, and as these cool sprinkle in the other substances, in the form of powder, so as to make a plaster. Ph. E.)* The same properties as the last, but more energetic and speedy in its action.

EMPLASTRUM MELOES VESICATORII, E. lyttæ —e. Meloës vesicatorii comp., E. lyttæ comp.—e. Mucilaginosum, Unguentum de althæâ—e. Nigrum of Augsburg, see Magnet.

EMPLASTRUM NUREMBERGENSÆ, *E. ex oxido plumbi rubro camphoratum, Emplâtre de Nuremberg, Nuremberg plaster. (Oxid. plumb. rubr. p. 300, ol. olive. p. 600, aquæ, q. s. Boil until the oxide is dissolved, and almost to dryness. Remove the vessel from the fire and add yellow wax, p. 500. Put the vessel again on the fire, and after the wax is melted, add, before it congeals, camphor, p. 24: mix. Ph. P.)* It is considered a desiccative, and has been employed in gangrene.

EMPLASTRUM NYGMATICUM, E. resinæ.

EMPLASTRUM OPII, *Opium Plaster. (Opii in pulv. 3ij, Picis abiet. 3ij; emplastr. plumbi lbj, aq. bullient. f 3iv. Melt together the lead plaster and Burgundy pitch; then add the opium pro-*

viously mixed with the water, and boil over a gentle fire to the proper consistence.—*Ph. U. S.*) It is employed as an anodyne, and to give support.

EMPLASTRUM OX'IDI FERRI RUBRI, E. Ferri rubri seu Rob'orans, Plaster of red oxide of iron, Strengthening Plaster. (*Empl. oxid. plumb. semivitr. p. xxiv, resina pin. p. vi, cera flavæ, olei oleæ Europ. sing. p. iij, oxid. ferri rubr. p. viij.* Rub the red oxide of iron with the oil, and add the other ingredients melted. *Ph. E.*) It is employed as a strengthening plaster.

EMPLASTRUM EX OXIDO PLUMBI SEMIVITREO, E. plumbi—e. Oxidi plumbi semivitrei, E. plumbi—E. Oxidi plumbi semivitrei gummosum, E. gummosum—e. Phoeniceum, Diapalmæ—e. ex Oxido plumbi rubro compositum, E. Norimbergense—e. Picis, E. Picis Burgundicæ, E. Picis compositum.

EMPLASTRUM PICIS BURGUNDICÆ, E. Picis, Burgundy Pitch or Pitch Plaster, (Picis Burgund. lbvj, cera flavæ, lbss.—*Ph. U. S.*)

EMPLASTRUM PICIS COMPOSITUM, E. picis cephalicum, Cephalic plaster, Compound pitch plaster. (*Picis arid. lbij, abietis resinæ lbj, resinæ flavæ, cera flavæ, aa ʒiv, myrist. ol. ʒi, ol. oliv., aquæ, aa f ʒij.* To the pitch, resin, and wax, melted together, add the other matters and mix. *Ph. L.*) It is stimulant and rubefacient. Sometimes used in headach,—applied to the temples. See Depilatory.

EMPLASTRUM PICIS CUM CANTHARIDE, E. Cantharidæ.

EMPLASTRUM DE PICE ET RESINIS GLUTINANS, E. Andræ de Cruce, Emplâtre d'André de la Croix, ou collant de poix et de résines, Adhesive plaster of pitch and resins. (*Picis albæ, p. 128, resin elemi, p. 32, terebinth. pur. p. 16, ol. laur. p. 16.* Melt with a gentle heat, and pass through linen. *Ph. P.*) Used in contusions and fractures as a support.

EMPLASTRUM PLUMBI, E. lithargyri seu communæ seu diachylum seu ox'idi plumbi semivitrei seu ex oxido plumbi semivitreæ, Diachylum simplex, White Diachylum, Lead or Diachylum plaster. (*Plumbi oxid. semivitr. in pulv. lbv., olei oliv., cong., aquæ Oij.* Boil together over a slow fire, stirring constantly until the oil and oxide of lead cohere. *Ph. U. S.*)

EMPLASTRUM POLYCHRESTUM, E. resinæ.

EMPLASTRUM POTASSII IODIDI, Plaster of Iodide of Potassium (Potass. iodid. ʒj, Thuris. ʒvj; Cera, ʒvj, ol. oliv. f ʒij. *Ph. L.*) This plaster, spread on leather, is applied to indolent tumours.

EMPLASTRUM RESINÆ, E. adhasivum seu lithargyri cum resinâ seu resinosum seu nymnativum seu communæ cum resinâ seu Polychrestum seu lithargyricum cum resinâ pini; Resin, Adhesive or Sticking plaster. (*Resin, lbss. emp. plumb. lbij.* Melt the plaster, with a gentle heat, add the resin, and mix. *Ph. U. S.*) Employed in wounds and ulcers.

Bayton's adhesive plaster is made by melting one pound of lead plaster and six drachms of resin together.

EMPLASTRUM RESINOSUM, E. Resinæ—e. Roborans, E. Ferri (Ph. U. S.), E. oxidi ferri rubri—e. Rubefians, E. lyttæ.

EMPLASTRUM SAPONIS, E. saponaceum, Soap plaster. (*Saponis concis. ʒiv, emplastr. plumb. lbij.* Mix the soap with the melted plaster, and boil to a proper consistence. *Ph. U. S.*) It is a mild discutient. Applied to tumours, corns, &c.

EMPLASTRUM SIMPLEX, E. ceræ—e. Spermatitis ceti, Ceratum cetacei.

EMPLASTRUM THURIS, Frankincense plaster. (*Emp. lithargyri, lbij, thuris, lbss. oxid. ferri.*

rubr. ʒij, Ph. D.), *Usc* :—the same as the plaster of red oxide of iron.

EMPLASTRUM VESICATORIUM, Blister, E. lyttæ—e. de Vigo cum mercurio emendatum, E. hydragryri compositum.

EMPLÂTRE, Emplastrum—e. d'André de la Croix, Emplastrum de pice et resinis glutinans—e. de Ciguë, Emplastrum cicuta—e. Collant de poix et de résines, Emplastrum de pice et resinis glutinans—e. de Gomme résine, Emplastrum cum gummi-resinis—e. de Nuremberg, Emplastrum Norimbergense.

EMPLATOMENOS, Emphracticus.

EMPNEUMATOSIS, Emphysema, Inspiration.

EMPOISONNEMENT, Poisoning.

EMPORIUM, εμποριον, (em, 'in,' and πορεω, 'I transport or send.' The brain was so called, of old, because there all the mental affairs are transacted.

EMPORIUM SPIRITUM, Sensorium.

EMPOSIS, Imbibition.

EMPREINTE, Impression.

EMPRESIS, Empresma.

EMPRESMA, Empresmus, Empre'sis, from εμπρησω, 'future,' εμπρησω, 'I burn internally,' Inflammation. Phlegmat'ia membranosæ et paraneumatosa, Phlogist'ici, Febres continuæ inflammatoriæ, Inflammatio interna, Cau'ma, Internal inflammation. A genus in the class hæmatica, order phlogotica, of Good.

EMPRESMA BRONCHITIS, Cynanche trachealis—e. Bronchlemmitis, Cynanche trachealis—e. Carditis, Carditis—e. Cephalitis, Phrenitis—e. Cystitis, Cystitis—e. Enteritis, Enteritis—e. Gastritis, Gastritis—e. Hepatitis, Hepatitis—e. Hysteritis, Metritis—e. Nephritis, Nephritis—e. Orchitis, Hernia humoralis—e. Otitis, Otitis—e. Paristhmitis, Cynanche—e. Paristhmitis tonsillaris maligna, Cynanche maligna—e. Paristhmitis pharyngea, Cynanche pharyngea—e. Paristhmitis tonsillaris, Cynanche tonsillaris—e. Parotitis, Cynanche parotidæa—e. Peritonitis, Peritonitis—e. Peritonitis mesenterica, Mesenteritis—e. Peritonitis omentalis, Epiploitis—e. Pleuritis, Pleuritis—e. Pleuritis diaphragmatica, Diaphragmitis—e. Pneumonitis, Pneumonia—e. Splenitis, Splenitis.

EMPRESMUS, Empresma.

EMPRION, from εν, and πριον, 'a saw.' Serated. Galen has given this name to the pulse, when the sensation produced by the artery under the fingers is analogous to that which would be caused by the unequal teeth of a saw.

EMPROSTHOCYRTOMA, Lordosis.

EMPROSTHOTHONOS, Emprosthoton'ia, Entasia tetanus anti'cus, Tetanus anticus, from εμπροσθεν, 'forwards,' and τενω, 'I stretch,' 'I extend.' A variety of tetanus, in which the body is drawn forwards by the permanent contraction of the muscles,

EMPSYCHO'SIS, from εμψυχω, (em, and ψυχη, 'life,') 'I animate,' 'I vivify.' A word formerly used for the act of animating. The union of the soul with the body.

EMPTOE, Hæmoptysis.

EMPTOICA PASSIO, Hæmoptysis.

EMPTYSIS, Hæmoptysis.

EMPTYSMA, Sputum.

EMPYE, Empyema.

EMPYEMA, Emphy'sis, Em'pye, from em, 'in,' and πυον, 'pus.' Aposte'ma empye'ma. A collection of blood or pus, and, conventionally, of other fluid, in some cavity of the body, and particularly in that of the pleura. Empyema is one of the terminations of inflammation of the pleura, and is called also, Pyothorax verus, Pleurothorax purulentus, Diapye'ma, Ecpye'ma, Ecpye'sis, Emphy'sis seu Pyo'sis pectoris, Hydrothorax purulentus, Dyspnoea pyothorac'ica, Pneumo'ma

suppurato'ria, Absces'sus pec'toris seu Thora'cis, Pleuraposte'ma, Pleuropnye'sia, (F.) Empyëme, Pnyphurite.

The operation for *empyema* properly means the making of an opening into the thorax for the purpose of giving issue to the matter collected in the cavity of the pleura, although it has been used for the operation required for the evacuation of any fluid from the chest, or synonymously with *Parencentesis thoracis*.

EMPYESIS, same etymon. Suppuration. An eruption of phlegmonous pimples, gradually filling with a purulent fluid, and terminating in thick scabs, frequently leaving pits or scabs. *Pust'ulosus Erem'them.* A genus in the order *Eucanthemata*, class *Hæmatica* of Good. See *Empyema*.

EMPIESIS OCULI, *Hypopyon*—e. Pectoris, *Empyema*—e. Variola, Variola.

EMPYMELIUM POLYSARCIA, Polysarcia adiposa.

EMPYOCE'LE, from *em*, 'in,' *πυον*, 'pus,' and *κηλη*, 'tumour,' 'hernia.' A tumour, formed by an accumulation of pus in the scrotum. Different diseases have been described under this name, such as suppuration of the testicle, *empyema* of the tunica vaginalis, accumulation of pus in the cavity of a hernial sac, abscesses of different kinds formed in the cellular texture of the scrotum, &c.

EMPYOMPHALUS, from *em*, 'in,' *πυον*, 'pus,' and *ομφαλος*, 'the navel.' This word has been used to designate a suppurating tumour at the umbilicus; or, at times, umbilical hernia, the sac of which is filled with blood.

EMPYOS, Purulent.

EMPYREU'MA, from *εμπυρεω*, (*em*, and *πυρ*, 'fire,') 'I kindle.' The burnt smell and acrid taste, which volatile products—gaseous and liquid—contract, when animal or vegetable substances are decomposed by a strong heat. The cause of this smell is seated in an oil, called *empyreumatic*, which does not exist in the substance subjected to the operation, but is the result of its decomposition. If the *empyreuma* occurs when the organic substance is placed in a still with a liquid, it is owing to the solid matter touching the bottom of the vessel to which the fire is applied.

EMPYREUMATIC, *Empyreumaticus*; same etymon. Belonging to *empyreuma*,—as an *empyreumatic* odour.

EMPYROS, *Febri'litana*. Same etymon. One who has fever.—Hippocrates.

EMS, MINERAL WATERS OF. Celebrated springs on the river Lahn, duchy of Nassau. They are thermal (from 83 to 115° Fahrenheit,) and carbonated salines, containing carbonic acid, bicarbonate of soda, and chloride of sodium; and are much used in gastric and intestinal affections, &c.

EMUL'GENT, *Emul'gens*, from *emulgere*, (*e*, and *mulgere*, *mulsim*,) 'to milk out,' 'to draw out.' A name given to the renal artery and vein, because the ancients imagined they milked, as it were, the urine through the kidneys. See Renal.

EMULSIFY, (F.) *Émulsionner*, from *emulsion*, and *faire*, 'to become.' To make or form an emulsion.

EMULSIN, see Amygdalin.

EMULSIO, *Emul'sion*, *Emul'sion*; same etymon. A pharmaceutical preparation, of a milky-white opaque appearance, which is composed of oil, divided and held in suspension in water by means of mucilage. Emulsions have been divided into the true and *oily*, and into the false or not *oily*; the latter being composed of resinous

substances, balsams, or camphor, rubbed up with dilute alcohol, mucilage, or yolk of egg.

EMULSIO ACA'CIÆ, *E. Arab'ica*, *Gum Arab'ic Emul'sion*. (*Nuclear*, *amygd. comm.* ʒij, *aqua* lbjss., *mucilag. acac.* ʒij, *sacch.* ʒiv. While beating the decorticated almonds with the sugar and water, add the mucilage.—Ph. E.) Used in the same cases as the next. Under the name *Mistu'ra Acac'ie*, *Gum Arab'ic Misture*, the Edinburgh Pharmacopœia has the following. *Mucilag. Acacia* fʒij; *Amygdal. dulc.* ʒj. and ʒij; *Sacchar.* ʒv; *Aqua* Oij. (Imper. *ingr.*) Blanch the almonds; peel them; beat to a smooth pulp first with the sugar, and then with the mucilage; add the water gradually, stirring constantly, and strain through linen or calico.

EMULSIO AMYGDALÆ, *E. Simplex*, *Lac amygd'alac*, *Amygdal'tum*, *Mistu'ra amygd'al.*, *Emul'sio sive Lac Amygd'alum* or *Almond Emulsion*; *Mixture or Milk*, (F.) *Lait d'amandes*. (*Amygd'al. dulc.* ʒss; *Acacia* in pulv. ʒss; *Sacchar.* ʒij; *Aqua destillat.* fʒvij.) Macerate the almonds in water, and having removed their external coat, beat them with the gum Arabic and sugar, in a marble mortar, till they are thoroughly mixed; then rub the mixture with the distilled water gradually added, and strain.—Ph. U. S.) It is used as a diluent and demulcent.

EMULSIO ANTHYSTERICA, *Mistura asafœtida*—e. Arabica, *Emulsio acacia Arabice*.

EMULSIO CAMPHORÆ, *E. Camphora'ta*, *Mistu'ra Camphora*; *Camphor Emulsion*. (*Camphora* ʒj, *amygd. com.* decortice, *sacch. dur.* aa ʒiv, *aqua* ʒvj.—Ph. E.) A convenient form for giving camphor.

EMULSIO GUAIACINA, *Mistura guaiaci*—e. Leniens, Looch ex ovo.

EMULSIO OL'EI AMYGDAL'ARUM; *Emulsion of Oil of Almonds*, (*Ol. amygd.* ʒj, *gum acac. pulv.* ʒij, *syrup.* ʒj, *aqua destill.* ʒiv. Mix. A good pectoral or cough mixture.

EMULSIO OL'EI RIC'INI; *Castor Oil Emulsion* (*Ol. ricini* ʒss, *vitelli ovi* q. s., *aqua destillat.* ʒj.) An aperient draught.

EMULSIO OL'EI TEREBIN'THINÆ; *Emulsion of Oil of Turpentine*, (*Ol. tereb. rect.* ʒij, *sacch. alb.* ʒss, *vitell. ovi* j, *emuls. amygd.* vel *aqua destillat.* ʒvj. Mix. In rheumatic and nephritic affections. Dose fʒiss.

EMULSIO OLEOSA, Looch ex ovo.

EMULSIO PURGANS CUM JALAP'Æ RESI'NA, *Purging Emulsion with Resin of Jalap.* (*Jalapæ resin.* gr. xij, *sacch. alb.* ʒij. Triturate for some time, and add gradually half the yolk of an egg; continue to triturate, adding by degrees *emuls. simpl.* ʒv, *aq. floz. carant.* ʒij.—Ph. P.)

EMULSIO PURGANS CUM SCAMMONIO; *Purging Emulsion with Scammony*. It is prepared like the preceding, substituting only Scammony for the jalap.

EMULSIO SEMPL'EX, *E. Amygdalæ*.

EMULSION, *Emulsio*—e. Almond, *Emulsio amygdalæ*—e. Camphor, *Emulsio Camphoræ*—e. Castor oil, *Emulsio olei ricini*—e. Gum Arabic, *Emulsio acacia Arabice*—e. *Huileuse*, Looch ex ovo—e. Oil of Almonds, *Emulsio olei amygdalarum*—e. Oil of Turpentine, *Emulsio olei terebinthinæ*—e. Purging, with resin of jalap, *Emulsio purgans cum jalapæ resinâ*—e. Purging, with scammony, *Emulsio purgans cum scammonio*.

ÉMULSIONNER, *Emulsify*.

EMULSIVE, *Emulsivus*; same etymon. An epithet given to seeds whence oil may be extracted by expression; such as almonds, apricots, peaches, hemp, rape, melons, gourds; those of the nut kind, and cucurbitaceous and cruciferous plants in general.

EMULSUM, *Emulsion*.

EMUNCTORIUM, *Emunctory*—e. *Cerebri, Nasus*.

EMUNC'TORY, *Emunctor'rium*, from *emungere*, (e, and *ungere*, *munctum*,) 'to drain off,' 'to cleanse.' *Emissa'rium*, (F.) *Emonctoire*, *Émissionnaire*. Any organ whose office it is to give issue to matters which ought to be excreted. The ancients believed that some organs were more particularly destined to serve as emunctories of others; the nasal fossæ, for example, they believed to be the emunctories of the brain.

EMUNDANTIA, Detergents.

EMYS PALUSTRIS, *Salt Water Ter'rapin*. This species of turtle is found exclusively in salt or brackish waters, near the sea shore, along the whole Atlantic coast. It is much prized by the epicure, and is nutritious and tolerably easy of digestion when dressed plain.

EN, see Em.

ENADELPHIA, *Emboitement*.

ENEMA, Hemostatica.

ENEMON, Styptic.

ENEOREMA, from *ev*, 'in,' and *αἶμα*, 'I lift up,' 'that which hangs or floats in.' (F.) *Enéorème*. The *Neph'elē*, *Nubes*, *Sublimamen'tum*, *Sublimat'io*, *Subl'imē*, *Suspen'sum*, *S. Uri'næ*, *Suspen'sa*, *Nube'cula* or cloud, which is suspended in the urine as it cools.

ENAMEL, *E. of the teeth*, (en, and (F.) *émail*.) *Dent'ium Nator*, *Corte'r*, *C. striat'a*, *Substan'tia filament'osa* of Malpighi, *S. Adamant'ina* seu *vit'rea*, *Crusta Dent'ium adamant'ina*, *Adamant'ine* or *vit'reous substance*. (F.) *Substance vitrée* ou *émaillée*, *Émail des Dents*. The substance which covers the coronæ of the teeth. The enamel is of a milky white colour, and sufficiently hard to strike fire with steel. Its surface is very smooth and polished, and it forms a thicker layer towards the part where the teeth come in contact, and becomes thinner towards the cervix. The fibres of the enamel are perpendicular to the surface of the teeth, on the ivory of which they seem, as it were, planted. This gives them a velvety appearance, when examined by the microscope. The enamel has no blood vessels, and is not renewed when removed. It is formed of phosphate of lime, and a very small portion of animal matter. A delicate membrane covers the enamel—*cuticle of the enamel*, *Nasmyth's membrane*; but so closely united with it, that it can be demonstrated only by the use of chlorohydric acid.

ENAMEL, CUTICLE OF THE, see Enamel.

ENANTHEMA, same etymon as the next. A name recently given to certain eruptions of the mucous membrane, as *exanthema* is to certain eruptions of the skin.

ENANTHE'SIS, from *ev*, 'in,' and *ανθεω*, 'I flourish.' *Rash exan'them*. Eruption of red, level or nearly level, patches, variously figured, irregularly diffused, often confluent, and terminating in cuticular exfoliations. A genus in the order *Exanthematica*, class *Hæmatica*, of Cullen, including scarlet fever, measles, &c.

ENANTHESIS ROSALIA, *Scarlatina*—e. *Urticaria*.

ENANTIOPATHIC, *Palliative*.

ENARTHRO'SIS, *Inarticulat'io*, *Ball and Socket-joint*, from *ev*, 'in,' and *αρθρωσις*, 'an articulation.' A kind of diarthrodial articulation, in which the head of a bone is received into the cavity of another, and can be moved in all directions. The joint of the os femoris with the os innominatum is one of this character.

ENARTH'RUM, from *ev*, 'in,' and *αρθρον*, 'a joint.' A foreign body in a joint.

ENAU'SMA, *Fomites*.

ENCAN'THIS, from *ev*, 'in,' and *κανθος*, 'the

angle of the eye.' A tumour, formed by an increase in size, or a degeneration, of the caruncula lacrymalis. Any morbid growth in the inner angle of the eye.

ENCANTHIS BENIG'NA. Simple excrescence of the caruncula. It commonly yields to astringent collyria.

ENCANTHIS FUNGO'SA. A condition of the semilunar fold and lachrymal caruncle, in which they are the seat of morbid growths.

ENCANTHIS INFLAMMATO'RIA. Inflammation with enlargement—swelling—of the semilunar fold and lachrymal caruncle.

ENCANTHIS MALIG'NA has often a cancerous character, and requires extirpation before it has attained any considerable size.

ENCARDITIS, Endocarditis.

ENCARPOS, Pregnant.

ENCASING, *Emboitement*.

ENCATALEPSIS, *Catalepsy*.

ENCATHIS'MA, *Senec'ipium*. With the ancients *Encathis'ma*, *εγκαθισμα*, (*ev*, and *καθίζω*, 'I seat upon,') meant a vapour-bath taken sitting.

ENCAUMA, from *ev*, 'in,' and *καυω*, 'I burn.' *Epicaum'is*, *Epicaum'is*, *Encaum'is*. A tumour produced by a burn. A burn. Also, an ulcer of the transparent cornea, occasioning loss of the humours.—*Aëtius*, *Gorræus*.

ENCAUSIS, Burn, Encauma, Moxibustion.

ENCAUSSE, MINERAL WATERS OF. Encausse is a village in the department of the Haute-Garonne, which possesses several saline, acidulous springs.

ENCAVURE, Argemæ.

ENCENTE, Pregnant.

ENCENS, Juniperus lycia.

ENCENSIER, Rosmarinus.

ENCEPHALALGIA, Cephalalgia—e. Hydro-pica, Hydrocephalus internus.

ENCEPHAL'IC, *Encephal'icus*, from *ev*, 'in,' and *κεφαλή*, 'the head.' That which is situate in the head. A name given to several parts which relate to the encephalon, as the *encephalic membrane*, *muscles*, &c.

ENCEPHALITIS, from *Encephalon*, and *itis*, denoting inflammation. This term has been used, by some nosologists, synonymously with *Cephal'itis* and *Phren'itis*. By others, it has been appropriated to inflammation of the brain, in contradistinction to that of the membranes.

ENCEPHALITIS EXSUDATORIA, Hydrocephalus internus—e. Infantum, Hydrocephalus internus—e. Insolationis, *Coup-de-soleil*—e. Membranosa, Meningitis—e. Peripherica, Meningitis—e. Potatorum, Delirium tremens.

ENCEPHALIUM, Cerebellum.

ENCEPHALOC'E'LE, from *encephalon*, and *κηλη*, 'a tumour.' *Hernia* seu *Fungus Cerebri*, *Cranioc'e'le*, *Hernia of the Brain*. This may be a congenital or accidental affection. In the former case, it is dependent upon tardy ossification of the fontanelles or some faulty conformation. In the latter, it is owing to some considerable loss of substance in the parietes of the cranium, produced by fractures, wounds with a cutting instrument, caries, the application of the trephine, &c. In slight congenital encephalocoele, gentle pressure may be exerted upon the protruded portion. When the disense is of considerable extent, it is fatal. In accidental encephalocoele, similar means must be used for confining the tumour, and preserving it from the action of external bodies.

ENCEPHALOCHYSIS, Hydrocephalus internus—e. Senilis, Apoplexy, serous.

ENCEPHALODYNIA, Cephalalgia.

ENCEPHALOHÆMIA, Cephalæmia.

ENCÉPHALOHÉMIE, Cephalæmia.

ENCEPH'ALOID, *Encephaloides*, *Ceph'aloid*, *Ceph'aloides*, *Ceph'alo'ma*, from *encephalon*, and *eidos*, 'resemblance.' Lænnec has given the term *Enceph'aloid* or *Cerebriform* matter to one of the morbid substances commonly formed by scirrhous or cancerous tumours. It is either encysted, in irregular masses without cysts, or infiltrated into the texture of the diseased organ. This name was given to it in consequence of its resemblance, when fully developed, to the medullary substance of the brain. It is also called *Fungus medull'ris* seu *Cancro'sus medull'ris* seu *Cerebr'lis*, *Sarco'ma medull'arē*, *Spongoid inflammation*, *Cancer cerebrif'ormē* seu *mollis* seu *medull'ris*, *Carcin'oma medull'arē* seu *spongio'sum*, *Carcin'us spongio'sus*, *Struma fungo'sa*, *Myelo'ma*, *Myelom'yces*, *Myelospon'gus*, *Tela accident'alis medull'ris*, (F.) *Matière cérébriforme*, *Carcinome mou et spongieux*, *Tumeur encéphaloïde ou spongieuse ou osseuse ou pulpeuse ou fonguïde ou fongueuse ou splénoïforme*, *Fungus médullaire*, *Carcinome saignant*, *Cancer mou ou encéphaloïde ou médullaire*; *Milt-like tumour*, *Medullary sarcoma*, *Cellular cancer*, *Fungoid* or *Hæmatocerc'eb'rif'orm disease*. See *Cancer*.

ENCEPHALOID TUMOUR OF THE LUNG, *Phthisis*, *cancerous*.

ENCEPHALOLOG'IA, from *encephalon*, and *λογος*, 'a description.' A description of the *encephalon*.

ENCEPHALOMALACIA, *Mollities cerebri*.

ENCEPHALOMALAXIS, *Mollities cerebri*.

ENCEPHALO-MENINGITIS, *Meningo-cephalitis*.

ENCEPH'ALON, *Enceph'alum*, *Enceph'alus*, *Enkeph'alon*; from *en*, 'in,' and *κεφαλη*, 'the head.' That which is situate in the head. This name has generally been given to the brain, *cerebellum*, and *mesencephalon*. At times, it includes likewise the *medulla spinalis*, when it is also called the *Cerebro-spinal*, and *Neural axis*. Under the term *Cranio-spinal axis* or *centre of automatic actions*, Dr. Carpenter includes the sensory ganglia—the respiratory and stomato-gastric ganglia forming the centre of the medulla oblongata and the spinal ganglia or true spinal cord.

ENCEPHALOPATHIA, (F.) *Encéphalopathie*, from *encephalon*, and *παθος*, 'disease.' A disease of the *encephalon*.

ENCEPHALOPATHIA LITERATO'RUM. The condition of brain, supposed to be produced by over-tasking it.

ENCEPHALOPATHIA PUERPERALIS, *Mania*, *puerperal*—e. *Saturnina*, see *Encéphalopathie saturnine*.

ENCÉPHALOPATHIE, *Encephalopathia*—e. *Crapuleuse*, *Delirium tremens*.

ENCÉPHALOPATHIE SATURNINE. *Encephalopathia Saturni'na*. *Encephalic disorder* occasioned by the poison of lead.

ENCEPHALOPHY'MATA, *Phymato'ses* seu *Struma'ses* seu *Tuber'cula cer'ebri*; from *encephalon*, and *φυμα*, *φυματος*, 'a tumour.' *Tubercles* of the brain.

ENCEPHALOPHTHISIS, see *Encephalopyosis*.

ENCEPHALOPYO'SIS, from *encephalon*, and *πυον*, 'pus.' *Apost'e'ma* seu *Abces'sus* seu *Hæco'sis cer'ebri*. Suppuration of the brain. When accompanied with emaciation and hectic, it is called *Encephalophth'isis*.

ENCEPHALO-RACHIDIAN, *Cephalo-spinal*.

ENCEPHALORRHAGIA, see *Apoplexy*.

ENCEPHALOSCOPIA, *Craniology*.

ENCEPHALOSEPSIS, *Mollities cerebri*.

ENCEPHALOSIS OF THE LIVER, *Hepato-scirrhus*.

ENCEPHALOSISMUS, *Concussion of the brain*.

ENCEPHALOSTRUMOSIS, see *Hydrocephalus internus*.

ENCEPHALUM, *Encephalon*.

ENCEPHALUS OPIS'THIUS, *Cerebellum*.

ENCEROSIS, *Inceration*.

ENCHARAXIS, *Scarification*.

ENCHATONNEMENT, *Chatonnement*.

ENCHEIRE'SIS, *Enchire'sis*, *Enche'ria*, from *en*, and *χαρ*, 'the hand.' Galen uses this term as a part of the title to one of his works, which treats of dissection. It means the manual treatment of any subject.

ENCHEIRIA, *Encheiresis*.

ENCHIFRÈNEMENT, *Coryza*.

ENCHIRESIS, *Encheiresis*.

ENCHONDROMA, *Chondroma*.

ENCHORIONOSUS, *Endemic*.

ENCHORIOS, *Endemic*.

ENCHRISTON, *Liniment*.

ENCHYLOSIS, *Extraction*.

ENCHYMA, *Infusum*, *Plethora*.

ENCHYMO'MA, *Enchymo'sis*, from *en*, and *χυω*, 'I pour.' By the ancients, this word was used to designate the sudden effusion of blood into the cutaneous vessels, which arises from joy, anger, or shame; in the last instance constituting blushing. It differs from *oecymosis* in there being, in the latter, extravasation of blood into the areolar texture, and its being produced by an external cause; a contusion, for example.—*Hippocrates*.

ENCHYSIS, *Infusion*.

ENCLAVÉ, *Wedge*.

ENCLAVEMENT, see *Wedge*.

ENCLUME, *Incus*.

ENCLYDAS'TICOS, *intus fluctuans*, from *en* *κλυδασται*, (*en* and *κλυω*, 'I wash.') 'to float in.' Applied to liquids, e. g., to serum, pus, &c., contained in any cavity.

ENCLYSMA, *Clyster*, *Enema*.

ENCŒ'LIA, from *en*, 'in,' and *κοιλια*, 'the belly,' The abdominal viscera. The entrails.

ENCŒLIALGIA, *Tormina*—e. *Inflammatoria*, *Encœlitis*.

ENCŒLIT'IS, *Encœli'tis*, *Encœlialg'ia inflammato'ria*, *Inflammat'io abdominal'is*; from *en* *κοιλια*, 'the abdominal contents,' and *itis*, inflammation. Inflammation of any of the abdominal viscera.

ENCOLPIS'MUS, from *en*, 'in,' and *κολπος*, 'the vagina.' Injection or introduction of any thing into the vagina.

ENCOLPIT'IS, *Oolpophlegmymenit'is*; from *en*, 'in,' *κολπος*, 'the vagina,' and *itis*, denoting inflammation. Inflammation of the lining membrane of the vagina.

ENCOPE, *Diapope*, *Incision*.

ENCRANION, *Cerebellum*.

ENCRANIS, *Cerebellum*.

ENCRE, *Atramentum*.

ENCYESIS, *Fecundation*, *Pregnancy*.

ENCYMON, *Pregnancy*.

ENCYMOSIA, *Fecundation*, *Pregnancy*.

ENCYSIS, *Pregnancy*.

ENCYST'ED, *Cystidē obductus*, *Sacca'tus*, *Saccula'tus*, *Sac'cated*, *Sac'culated*, *Pouch'd*. Enclosed in a kyst or cyst, or pouch; from *en*, 'in,' and *κυστις*, 'bladder.' (F.) *Enkysté*. An epithet given to certain tumours, or solid or fluid collections enclosed in a particular envelope or cyst. They are movable and often elastic to the touch.

ENCYS'TIS, *Tumor tunica'tus seu cysticus*, *Emphy'ma encys'tia*. Same etymon. *Lu'pia*, *Glan'dula Avicen'næ*, *Nodus*. An encysted tumour.

ENDADELPHIA, see *Emboîtement*.
ENDANGIUM; properly *Endangi'on* or *Endangi'um*, *Membra'næ vascul'arum commun'is*. (F.) *Tunique commune de Bichat*: from *ενδον*, 'within,' and *αγγειον*, 'a vessel.' The serous or lining membrane of vessels.

ENDEICTICOS, Indicant.

ENDEIXIOL'OGY, *Endeixiolog'ia*, from *ενδεικτες*, (*ev*), and *δεικνω*, 'I show,' 'indication,' and *λογος*, 'a discourse.' The doctrine of indications.

ENDEIXIS, Indication.

ENDEMIA, Endemic.

ENDEMIC, *Endem'ical*, *Endem'ia*, *Regional'is morbus*, *Endem'icus*; from *εν*, 'in,' and *εμμος*, 'the people,' *Euchor'ia*, *Enchorion'osus*, *Vernaculus* seu *Endemius morbus*, *Endemy*, (F.) *Endémie*, *Maladie régionale*. A disease said to be endemic, (F.) *Endémique*, or to arise from endemicity, (F.) *endémicité*, when it is owing to some peculiarity in a situation or locality. Thus, ague is endemic in marshy countries; goutre at the base of lofty mountains, &c. Some authors use the term in the same sense as epidemic. We have no accurate knowledge of the emanations or other circumstances which give occasion to endemic affections. We seem to know that some emanation from marshy lands does produce intermittents: but we are ignorant of the nature of such emanation.

ENDÉMICITÉ, see Endemic.

ENDEMICITY, see Endemic.

ENDEMIOL'OGY, *Endemiolog'ia*, from 'endemic,' and *λογος*, 'a discourse.' The doctrine of endemic diseases.

ENDÉMIQUE, Endemic.

ENDEMY, Endemic.

ENDEPIDERMIS, Epithelium.

ENDERMIC, *Enderm'icus*, *Endermat'icus*, *Emplas'tro-endermic*, (F.) *Endermique*, from *εν*, and *δερμα*, 'the skin.' An epithet given to the method of treating diseases by placing the therapeutical agent in contact with the skin, especially after the cuticle has been removed: *Enderm'ism*, *Enderm'ismus*, *Endermo'is*, *Méth'odus endermica* seu *endermat'ica*, *Diadermat'ia*. Morphia, strychnia, &c., are often administered in this way.

ENDERMIS, see Endermic.

ENDERISMUS, see Endermic.

ENDERMOSIS, see Endermic.

ENDESIS, from *εν*, 'in,' and *δεω*, 'I bind.' A ligature, bandage, connexion. Hippocrates has so termed the ankle-joint.

ENDIVE, *Cichorium endivia*—e. Wild, *Cichorium intybus*.

ENDIVIA, *Cichorium endivia*.

ENDIXIS, Indication.

ENDO, from *ενδον*, 'within.' A common prefix, as in the following terms:

ENDO-AORTITIS, from *ενδον*, 'within,' and *aortitis*, 'inflammation of the aorta.' Inflammation of the lining membrane of the aorta.

ENDO-ARTERITIS, see Arteritis.

ENDOBLAST, see Periblast.

ENDOCARDE, Endocardium.

ENDOCARDIAC, *Endocard'ial*, *Endocardi'acus*: same etymon as the next. Relating to the endocardium, or to the interior of the heart; as 'endocardiac sound or murmur,'—a sound produced within the cavities of the heart, in contradistinction to *exocardiac* or *exocardial* sounds or murmurs, which are induced by conditions of the external surface of the organ.

ENDOCARDITIS, *Encarditis*, *Card'itis interna*, *Inflammatio superfic'iei inter'næ cordis*, *Inter'nal Carditis*, *Inflammation of the internal membrane of the heart*, (F.) *Endocardite*, from *endocard'ium*, the lining membrane of the heart,

and *itis*, inflammation. In this disease, the heart's action is visibly increased, and very manifest to the touch; the hand is strongly repelled, and, at moments, is sensible of a trembling vibratory motion. Percussion gives a dull sound over a surface of several inches, owing, according to Bouillaud, to the inflammatory turgescence of the heart, and the engorged state of its cavities. On auscultation, a *bruit de soufflet* is generally heard, masking one or both sounds of the heart; and the ear is sensible of a metallic ringing with each systole of the ventricle. The pulsations are rapid as well as strong, and, with few exceptions, irregular, unequal and intermittent. The pulse, by the way, does not always indicate the force or number of the heart's contractions.

ENDOCARDIUM, from *ενδον*, 'within,' and *καρδια*, 'the heart.' (F.) *Endocarde*. The membrane that lines the interior of the heart.

ENDOCHORION, see Chorion.

ENDOCOLITIS, Dysentery.

ENDOCYMIEN, from *ενδον*, 'within,' and *χυμα*, 'the product of conception.' A double monster by inclusion—Isidore G. St. Hilaire. See *Emboîtement*.

ENDODONTITIS, (F.) *Inflammation de la pulpe dentaire*, *Inflammation of the Dental membrane*, from *ενδον*, 'within,' *odontos*, *odontos*, 'a tooth,' and *itis*, denoting inflammation. Inflammation of the lining membrane of a tooth.

ENDO-ENTERITIS, see Enteritis.

ENDOASTRITIS, *Esogastr'itis*; from *ενδον*, 'within,' and *gastritis*, 'inflammation of the stomach.' Inflammation of the lining membrane of the stomach.

ENDOGEN, see Endogenous.

ENODOGENOUS; from *ενδον*, 'within,' and *γεννω*, 'I engender.' A term first applied to plants—hence called *End'ogens*—in which the new woody matter is deposited within the old, and towards the centre. In the animal, cells are often formed *endogenously*, or within the cells, as in the case of the sperm vesicles.

An *endog'enous aneurism* is one that originates spontaneously from lesions of the inner coat of the arteries.

ENDOLYMPH, *Vitrine auditive*.

ENDOMETRITIS, from *ενδον*, 'within,' and *metritis*, 'inflammation of the uterus.' Inflammation of the lining membrane of the uterus. When accompanied with a diphtheritic exudation, it is sometimes called *Endometrit'is crôupo'sa*.

ENDONARTERITIS, see Arteritis.

ENDOPHLEBITIS, Phlebitis.

ENDOPLAST, see Periplast.

ENDORMIE, *Datura stramonium*.

ENDOSIS, Remission.

ENDO-SKELETON, see Skeleton.

ENDOSMOSE, *Endosmo'sis*, *Double Imbibit'ion*, from *ενδον*, 'within,' and *ωσμος*, 'Impulse.' A term first used by Dutrochet, to express the action by which fluids pass from without to within organic membranes. The action of two fluids on each other, when separated by a membrane. The general conditions of the phenomena are—*first*, that they should have an affinity for the interposed membrane; and *secondly*, that they should have an affinity for each other, and be miscible.

At the present day, *endosmose* is generally used to signify the passage of the more transmissible fluid, whilst *exosmose* signifies that of the least transmissible. The rapidity with which endosmose is accomplished varies according to the nature of the septum or tissue and of the penetrating body, and to the penetrability of the tissue.

ENDOSMIC, Endosmotic.

ENDOSMOTIC, *Endosmot'icus*, *Endos'mic*,

Endos'micus; same etymon. Belonging to endosmose:—as an 'endosmotic current.'

ENDOSTEÏTIS, from *εσθον*, 'within,' *οσσειν*, 'a hope,' and *itis*, denoting inflammation. Inflammation of the internal periosteum.

ENDOSTEUM, Medullary membrane.

ENDÔUTÉRITIS, see Metritis.

ENDUIT (F.), from *induire*, 'to put upon,' 'to put on.' A coat; a fur. This term is often applied to a layer of greater or less thickness which covers the surface of certain organs, and particularly of the tongue and the interior of the mouth. The *enduit* is designated variously, according to its appearance,—*enduit bilieux, jaune, blanc*, &c.—a bilious, yellow, white coat or fur, &c. It is at times owing to the evaporation of the watery portions of the secretions; at others, to a morbid condition of the secretions:—generally, to both causes combined. The tongue, in such a condition, is said to be coated or furred; sometimes, loaded.

ENDUIT CHOROÏDIEN, see Choroid.

ENDURCISSEMENT, Induration—*c. du Cerveau*, Sclerencephalia—*c. Rouge*, see Hepatization—*c. du Cœur*, Cardiosclérosis—*c. du Tissu cellulaire*, Induration of the cellular tissue.

ENDYMA VENTRICULORUM, Ependyma ventriculorum.

ENECHEMA, Tinnitus aurium.

ENECIA, Synocha—*c. Cauma*, Synocha—*c. Synochus Puerperarum*, see Peritonitis—*c. Synochus*, Synochus.

ENEMA, *εμψα*, from *εμψα*, (*ev*, and *εμψα*, 'to inject,' *Clyema*, *Clysmus*, *Enclys'ma*, *Lavamentum*, *Lo'tio*. An Injection, Clyster, (F.) *Clystère*, *Lavement*. A well-known form of conveying medicine into the intestinal canal. See Clyster.

ENEMA ALOËS, Clyster of aloes. (*Aloës*, *Ῥίη*; *Potass. carbon*, gr. xv; *Decoct. Hordei*, Oss, (Imp. meas.), *M. Ph. L.*) Used in cases of ascarides and amenorrhœa.

ENEMA ANODYNUM, *Enema O'pii*; *Anodyne* or *Starch* and *Opium* Clyster. (*Decoct. amyli f3ij*; *tinct. opii M. xxx. M. Ph. L.*) Exhibited in cases of severe diarrhœa or dysentery.

ENEMA CATHARTICUM, Cathartic or Purging Clyster. (*Olei. oliv. f3ij*; *sulph. magnes. 3j*; *muçilag. acacia f3xvi*; *M. Ph. D.*)

ENEMA COLOCYNTHIDIS, Clyster of *Colocyath*. (*Ext. Colocyath 3ss*; *Sapon. mollis 3j*; *aqua, Oj*. (Imp. meas.), *M. Ph. L.*) An efficient cathartic clyster.

ENEMA COMMUNE; Common or Domestic Clyster. (Water gruel or molasses and water Oss or Oj; add a little oil or lard, and a spoonful of common salt.) Given as a cathartic enema; and, without the common salt, as an emollient.

ENEMA FÆTIDUM, *E. assafœtida*, Fetid Clyster, Clyster of *assafœtida*; *Mistura assafœtida pro clysmate*, *Clyema tonicum* et *antispasmodicum* seu *incitans* et *sedans*, (F.) *Lavement antispasmodique*. (The last, with the addition of 3ij of the tincture of *assafœtida*. *Ph. D.*) Given as an antispasmodic and anodyne.

ENEMA NICOTIANÆ; *Tobacco* Clyster. This generally consists of from half a pint to a pint of the *Infusum Tabaci*. It is employed in cases of strangulated hernia; but occasionally acts as a fatal poison when given in this way. The smoke of tobacco is sometimes thrown up the rectum to produce the same medicinal effects as the infusion.

ENEMA OPII, *E. anodynum*.

ENEMA TEREBINTHINÆ; *Turpentine* Clyster. (*Ol. tereb. f3j*. *Decoct. hordei f3xix*, one yolk of egg. Incorporate the turpentine with the yolk, then add the gruel. *Ph. L.*) To be administered in cases of ascarides, (*oxyurea*.)

ENEPIDERMIC, *Enepidermicus*, from *εν*, 'in,' *ενι*, 'upon,' and *δερμα*, 'the skin.' An epi-

thet given to the method of treating diseases, which consists in the application of medicines; as plasters, blisters, &c., to the skin.

ENEREISIS, Compression.

ENERGIA, Action, Force.

ENERGY, *Energi'a*, from *ενεργω*, (*ev*, and *εργον*, 'action') 'I act.' Action. Acting power. Also, vigour; as the 'muscular energy;' the 'brain acts with energy.'

ENERVATIO, Aponeurosis, Enervation.

ENERVATION, *Enerva'tio*, from *en*, 'out of,' and *nervus*, 'strength.' The act of weakening—the state of being weakened. See Debility.

ENERVATIONES TENDINEÆ, Inscripti. ones tendineæ musculorum.

ENERVITAS, Inertia.

ENFANCE, Infancy.

ENFANT, Infans—*c. à Terme*, see Fœtus—*c. Légitime*, see Legitimate.

ENFANTEMENT, Parturition.

ENFLURE, Swelling, Œdema—*c. des Jambes* et *des cuisses* de la femme accouchée, Phlegmatia dolens.

ENFONCEMENT SCAPHOÏDE, Scaphoides fossa.

ENFONCEURE, Arthrodia.

ENFONCURE, Arthrodia.

ENGASTERION, Laboratory.

ENGASTRIMISME, see Engastrimyth.

ENGASTRIMYSME, see Engastrimyth.

ENGASTRIMYTH; *Engastrim'ythus*, *Englottogastor*, *Gastril'oquus*, *Ventril'oquus*, *Gastril'oquist*, *En'rycles*, *Eurycl'i'tus*, *Enteroman'tis*, *Gastroman'tis*, from *εν*, 'in,' *γαστρ*, 'the belly,' and *μυθολογία*, 'I discourse.' A Ventril'oquist. (F.) Ventriloque, Gastriloque. One who possesses the art of modifying his natural voice, so that it seems to come from a greater or less distance, and from different directions. It was formerly believed that such persons spoke from the belly; hence their name. It is an imitative art, and is called VENTRILOQUISM. (F.) *Engastrimisme*, *Engastrimysme*.

ENGESOMA, Engisoma.

ENGELURE, Chilblain.

ENGHIEN MONTMORENCY, MINERAL WATERS OF. *Engbien-les-Bains*. A hydro-sulphurous water, four leagues from Paris, near Montmorency, which is possessed of some celebrity. It contains chloride of sodium, chloride of magnesium, sulphate of magnesia, sulphate of lime, carbonate of magnesia, carbonate of lime, silica, sulphohydrate of lime and magnesia, sulphohydric acid, carbonic acid and nitrogen.

ENGISO'MA, *Engiso'ma*, *Engizo'ma*, from *εγγίζω*, 'I approximate.' (F.) *Embarrure*. A species of fracture of the skull, in which a splinter passes beneath the sound portion of the bone, and compresses the brain.—Galen. Also, a kind of instrument used in fractures of the clavicle.

ENGLISH DISEASE, Rachitis.

ENGLOTTOGASTOR, Engastrimyth.

ENGOMPHOSIS, Gomphosis.

ENGORGEMENT, from *en*, 'in,' and *gorge*, 'the throat.' An obstruction occurring in the vessels of a part, giving rise to augmentation of volume. Congestion.

ENGORGEMENT DES MEMBRES ABDOMINAUX À LA SUITE DES COUCHES, Phlegmatia dolens—*c. Hepatic*, *Hepatœmia*—*c. Laitéux* des membres abdominaux, Phlegmatia dolens.

ENGOUEMENT (F.), *Obstructio*, *Inert'ia*, from *angere*, 'to choke.' Accumulation in a hollow organ, of the matters secreted by it or carried into it. There is said to be *Engouement* des bronches, when the mucus accumulates in the bronchia; and *Engouement* des intestins, when the matters which ought to pass through the in-

testines are detained; as in a case of strangulated hernia.

ENGOUEMENT DES POUMONS, *E. of the lungs*, in Pathological Anatomy, signifies that state of the lungs, in which a mixture of air and thin fluid escapes from them when cut into.

ENGOURDISSEMENT, Torpor.

ENGRENURE, Suture.

ENH-EMATOSIS, Hæmatisis.

ENIXA, Puerpera.

ENIXIO FETÛS, Parturition.

ENIXUS FETÛS, Parturition.

ENKEPHALON, Encephalon.

ENKYSTÉ, Encysted.

ENKYSTEMENT, Chatonnement.

ENMETRITIS, see Metritis.

ENNEAPHARMACOS, from *εννεα*, 'nine,' and *φάρμακον*, 'a medicine.' A medicine, composed of nine simple ingredients. A pessary, so formed.—Galen, Paulus. The *Antidotus Heraclidis*, described by Galen, and some plasters by Aëtius and Celsus, are, likewise, termed *Enneapharmaca*.—Goræus.

ENORMIS, Abnormous.

ENORMITAS CORDIS, Heart, hypertrophy.

ENORMON, *ενόρμων*, *Hormon*, *Impetus faciens*, from *εν*, 'in,' and *ορμω*, 'I rouse,' 'excite.' A word used by Hippocrates in the same sense as *vital principle* is by modern physiologists.

ENOSIS, Insertion.

ENOSTOSIS, *Entostosis*, from *εν*, 'in,' and *οστος*, 'a bone.' A morbid growth of bone inwards—the opposite to *exostosis*.

ENRAGÉ, Rabid.

ENRHUMER, See Cold.

ENROUEMENT, Raucedo.

ENRYTHMOS, from *εν*, and *ρυθμος*, 'number.' Having rhythm. An epithet applied to the pulse when its pulsations occur with some degree of order. It is opposed to *Arythmos*, and differs from *Eurythmos*, which signifies 'regular.'

ENS, A being. Paracelsus meant, by this term, the power exerted by certain beings on our body. He speaks of the *Ens Dei*, *E. Astro-rum*, *E. naturalè*, *E. virtutis*, *E. morborum*, *E. de potentibus spiritibus*, &c. These absurd denominations suggested to some of the ancient chymists a name for certain chymical preparations. The muriate of ammonia and iron was called *Ens martis*; the muriate of ammonia and copper, *Ens veneris*; and *Ens primum* was, with the alchemists, the name of a tincture which they considered possessed of the power of transmuting metals.

ENS MARTIS, Ferrum ammoniatum—*e. Veneris*, Boylei, Ferrum ammoniatum.

ENSIFORM, *Ensiformis*, from *ensis*, 'a sword,' and *forma*, 'form.' Sword-like.

ENSIFORM APOPHYSES or **PROCESSES** are the lesser *alæ* of the sphenoid bone.

ENSIFORM APPENDIX or **CARTILAGE** is the xiphoid appendix of the sternum, &c. See *Xiphoid*.

ENSIFORMIS, Xiphoid.

ENSISTERNAL. Relating to the ensiform process of the sternum. Bécclard gave this name to the last osseous portion of the sternum. He also called it *l'os ultimi-sternal*. See *Ultimo-sternal*.

ENSOMATOSIS, Incorporation.

ENSTALAXIS, Instillation.

ENSTROPHÉ, from *εν*, 'in,' and *στρεφω*, 'I turn.' Inversion of a part, as of the eyelids.

ENTABLEMENT, Depression.

ENTAILLE (*F.*), from *en*, 'in,' and *tailler*, 'to cut.' *Excisio*, *Ec'corpè*, *Ec'tomé*. A deep wound made by a sharp instrument obliquely. *Entailles* and *Taillades* are, also, used to desig-

nate deep scarifications, made for the purpose of producing a speedy *dérgorgement* of any tumefied part; such, for example, as are made in the tongue in certain cases of glossitis.

ENTAMURE (*F.*). A slight incision or laceration.

ENTANTHEMA, Eisanthema.

ENTASIA, Tonic spasm—*e. Articularis*, Contractura—*e. Loxia*, Torticollis—*e. Priapismus*, Priapismus—*e. Systremma*, Cramp—*e. Tetanus*, Tetanus—*e. Tetanus anticus*, Emprosthotonos—*e. Trismus*, Trismus.

ENTASIS, Tonic spasm.

ENTAT'ICUS, *Intensivus*, from *εντελως*, 'I make tense.' A medicine which excites the venereal appetite.

ENTELIPATHIA, Nymphomania.

ENTELMINTHA, Worms.

ENTENDEMENT, Intellect.

ENTERADELPHIA, see *Emboîtement*.

ENTERAD'ENES, from *εντερον*, 'an intestine,' and *αδην*, 'a gland.' The mucous glands of the intestines. See *Peyer's glandulæ*.

ENTERADENOGRAPHY, *Enteradenogra'phia*, from *εντερον*, 'intestine,' *αδην*, 'gland,' and *γραφη*, 'a description.' A description of the intestinal glands.

ENTERADENOLOGY, *Enteradenolog'ia*, from *εντερον*, 'intestine,' *αδην*, 'gland,' and *λογος*, 'a treatise.' That part of anatomy which treats of the intestinal glands.

ENTERAËRETASIA, Tympanites.

ENTERAL'GIA, *Enterodyn'ia*, from *εντερον*, 'intestine,' and *αλγος*, 'pain.' Pain in the intestines. Colic.

ENTERALGIA ACUTA, Enteritis—*e. Inflammatoria*, Enteritis—*e. Saturnina*, Colica metallica.

ENTERANGEMPHRAXIS, Enterangi-emphraxis.

ENTERANGIEMPHRAXIS, *Enterangemphrax'is*, from *εντερον*, 'intestine,' *αγγω*, 'I strangle,' and *εμφρασσα*, 'I obstruct.' Obstruction of the vessels of the intestines.

ENTERATROPH'IA, *Atroph'ia Intestino'rum*, from *εντερον*, 'intestine,' and *atrophia*, 'want of nutrition.' Atrophy of the coats of the intestines.

ENTERAUX'E, from *εντερον*, 'intestine,' and *αυξη*, 'increase.' Hypertrophy of the muscular coat of the intestines.—Fuchs.

ENTERECHE'MA, from *εντερον*, 'intestine,' and *ηχη*, *ηχω*, *ηχημα*, 'sound.' *Son'itus intestinalis*. The sound of the movement of the intestines, heard by the stethoscope.

ENTERECTASIS, *Dilata'tio Intestino'rum*, from *εντερον*, 'intestine,' and *εκτασις*, 'dilatation.' Dilatation of the intestines, as in tympanites.

ENTERELCO'SIS, *Enterhelco'sis*, *Enterohelco'sis*, from *εντερον*, 'intestine,' and *ελκος*, 'ulceration.' Ulceration of intestines—*Ulcerat'io intestinalis*, *Ulcera intestinalia*.

ENTERELESLIA, Ileus.

ENTERELOSIS, Ileus.

ENTEREMPHRAXIS, *Infarctus Intestino'rum*, *Incarcera'tio intestino'rum inter'na*, (*F.*)

Étranglement des Intestins, *E. Intestinale*, from *εντερον*, 'intestine,' and *εμφραξις*, 'obstruction.' Obstruction of the Intestines from accumulation of feces or otherwise. Also, Enterangiempfraxis.

ENTEREN'CHYTA, *Enteren'chytes*, from *εντερον*, 'an intestine,' *εν*, and *χω*, 'I pour.' Any surgical instrument for administering injections.—Scultetus.

ENTEREPIPLOCELE, Enterœpiplocele.

ENTEREPILOMPHALOCELE, from *εντερον*, 'intestine,' *επιπloon*, 'omentum,' *ομφαλος*, 'navel,' and *κηλη*, 'rupture.' Hernia of the umbilicus, with protrusion of the omentum and intestine.

ENTERHELICOSIS, Enterolecosis—e. Nervosa,
Typhus (abdominal.)

ENTERHYPERTROPHIA. *Hypertroph'ia intestinum*, from *enteron*, 'intestine,' and *hypertroph'ia*, 'excessive nutrition.' Hypertrophy of the coats of the intestines.

ENTERIC, Intestinal.

ENTERICA, from *εντερον*, 'an intestine.' Diseases affecting the alimentary canal. Order I., Class I. (*Coliaca*), of Good. Also, medicines affecting the alimentary canal.

ENTERICUS, Intestinal.

ENTÉRIQUE, Intestinal.

ENTÉRITE FOLLICULEUSE, see Typhus
—c. *Typhohémique*, Typhoid fever.

ENTERITIS, from *ερεπον*, 'an intestine,' and *itis*, denoting inflammation. Inflammation of the intestines. *Empresma Enteritis*, *Intestinorum inflammatio*, *Ileo-culitis*, *Chordapneus*, *Cama Enteritis*, *Enterophlogitis*, *Enterophlogosis*, *Colica acuta seu inflammatoria*, *Il'eus inflammatorius*, *Enteralgia acuta seu Inflammatoria*, *Febrius intestinorum seu ill'aca inflammatoria*, *Enteropyra*, (F.) *Entérite*, *Inflammation des Intestins*. The essential symptoms of this disease are:—violent abdominal pain, increased on pressure; with vomiting and inflammatory fever. Enteritis may affect both the peritoneal and the mucous coat of the intestines; and, in violent cases, as the coats may be implicated. The structure of the mucous and peritoneal coats is different; so are their functions in health and disease. The inflammation of the serous coat, *Sero-enteritis*, resembles that of the cellular membrane; the inflammation of the mucous coat, that of the skin. The former is usually, therefore, of a more active character. Inflammation of the mucous coat, *Eso-enteritis*, *Endo-enteritis*, *Muco-enteritis*, *Mucous Enteritis*, *Phlegmymenitis enterica*, is generally attended with diarrhoea, and its pathology is identical with that of dysentery. Inflammation of the peritoneal coat is, on the other hand, generally attended with constipation.

Enteritis of the peritoneal coat, for such usually is the meaning of the word in the abstract, requires the most active treatment. Copious bleeding, followed up by a large dose of opium—and, if the symptoms be not decidedly ameliorated, repeating the bleeding and opium—warm fomentations, and blisters, are the chief agents to be relied upon. Purgatives ought not to be exhibited until the inflammation and spasm are abated by the use of the bleeding and opium. When the physician is called early, this plan will usually succeed. Sometimes, Enteritis passes into a chronic state, requiring much careful management. Broussais considered inflammation of the mucous coat of the stomach and intestines as the proximate cause of the phenomena of fever.

ENTERITIS ARTHRITICA, Coeliagra—e. Colica, Colitis—e. Epiploitis, Epiploitis—e. Follicular, Typhoid fever—e. Folliculosa, see Typhus—e. Mesenterica, Mesenteritis—e. Mucosa, see Enteritis—e. Pustulosa, Doinhenteritis.

ENTERO, from *εντερον*, 'an intestine,' in compound words signifies intestine, as in the following:—

ENTEROBRO'SIS, *Enterorrhœ'is, Perfora'tio*
Intestino'rum, (F.) *Perforation des Intestins*, from
εντερον, 'an intestine,' and *βρωσις*, 'the act of
gnawing.' Perforation of the intestines.

ENTEROCÆCÆ, *Dysenteria putrida* seu typhoides seu scorbutica seu maligna, from εντερων, 'an intestine,' and κακος, 'evil.' Adynamic dysentery, accompanied by phenomena indicating a pseudo-membranous and gangrenous state of the lining membrane of the large intestine.

ENTEROCE'LE, *Hernia intestina'lis*, from

εντερον, 'an intestine,' and κηλη. 'a hernia,' 'tumour.' Abdominal hernia, which contains only a portion of intestine, is so called.

ENTERO-CEPHALOPYRA INFANTUM,
Hydrocephalus Internus.

ENTEROCŒ'LICUS, from *εντερον*, 'intestine,' and *κοιλια*, 'the cavity of the abdomen.' Relating to the cavity of the abdomen.

EN'TERO-COLI'TIS, (F.) *Entéro-colite*, from *εντερον*, 'intestine,' and *colitis*, 'inflammation of the colon.' Inflammation of the small intestine and colon.

ENTEROCYSTOC'E'LĒ, from *εντερον*, 'an intestine,' *κυστις*, 'a bladder,' and *κηλη*, 'a tumour.' Hernia formed by the bladder and a portion of intestine.

ENTERODAR'SIS, from *εντερον*, 'intestine,' and *δαρσις*, 'skinning.' Excoriation of the mucous membrane of the intestines.

ENTERODOTHIA. Dothinerteritis.

ENTERODYNIA, Enteralgia.

ENTERO-EPIPOCE'LE, *Enterepipocele*, *Her'nia intestinalis omenta'lis*, from *εντερον*, 'intestine,' *επιπλουν*, 'the omentum,' and *κληη*, 'tumour.' Hernia, formed by intestine and omentum.

ENTERO-EPIPLOM'PHALUS, from *εντερον*, 'intestine,' *επιπλοον*, 'the omentum,' and *ομφαλος*, 'the umbilicus.' Umbilical hernia, containing intestine and omentum. Almost all umbilical herniæ are of this kind.

ENTEROGRAPHY, *Enterogra'phia*, from *εντερον*, 'intestine,' and *γραφη*, 'description.' The part of anatomy which describes the intestines.

ENTERO-HÆMORRHAGIA. Hæmatochezia.

ENTEROHELICOSIS, Enterelcosis.

ENTERO-HYDROCE'LE, from *εντερον*, 'intestine,' *υδωρ*, 'water,' and *κηλη*, 'tumour.' Intestinal hernia complicated with hydrocele.

ENTERO-HYDROM'PHALUS, from *εντερον*, 'intestine,' *ὕδωρ*, 'water,' and *ομφαλος*, 'the navel.' Umbilical hernia, in which the sac contains, along with a portion of intestine, a quantity of serum.

ENTERO-ISCHIOCE'LE, from *εντερον*, 'intestine,' *ισχιον*, 'ischium,' and *κηλη*, 'tumour.' *Ischiatic hernia*, formed of intestine.

ENTEROLITHIASIS, from *εντερον*, 'intestine,' and *λιθιασις*, 'the formation of stone.' The formation of concretions in the intestines.

ENTEROLITHUS, Calculi of the stomach and intestines—e. Bezoardus, Bezoar—c. Scybalum, Scybala.

ENTEROL'OGY, *Enterolog'ia*, from *εντερον*, 'intestine,' and *λογος*, 'a discourse.' The part of anatomy which treats of the intestines.

ENTEROMALACIA, *Enteromalax'is*, (F.) *Ramollissement de l'Intestin*, from *εντερον*, 'an intestine,' and *μαλασσω*, 'I soften.' Softening of the mucous or other coats of the intestine.

ENTEROMALAXIS. Enteromalacia.

ENTEROMANTIS, Engastrimyth.

ENTERO-MEROCE'LE, from *εντερον*, 'intestine,' *μηρος*, 'the thigh,' and *κηλη*, 'tumour.' Cru-
ral hernia, formed of intestine.

ENTERO-MESENTER'IC, *Entero-mesenter'-icus*, (F.) *Entéro-mésentérique*. Relating to the intestine and mesentery.

ENTERO-MESENTERICA FEBRIS. MM. Petit and Serres have given this name to the typhoid form of adynamic fever, in which the intestines are ulcerated, with enlargement of the corresponding mesenteric glands. See Typhus.

ENTÉRO-MÉSENTÉRITE, Tabes mesenterica—*e. Méésentérite typhoïde*, see Typhus.

ENTEROMORPHIA COMPRESA. One of the algæ, used by the Sandwich Islanders as an esculent.

ENTEROMPHALOCELE, *Enteromphalus*.

ENTEROMPHALUS, *Enteromphalocēlē*, from *εἰς* *teron*, 'intestine,' and *ομφαλός*, 'umbilicus.' Umbilical hernia, formed of intestine.

ENTEROMYCODORITIS, from *εντερον*, 'intestine,' *μυκός*, 'slime,' *δόξα*, 'skin,' and *itis*, denoting inflammation. Inflammation of the mucous membrane of the intestines.

ENTERON, Intestine.

ENTEROPARALYSIS, *Enteroplegia*, from *εντερον*, 'intestine,' and *παράλυσις*, 'paralysis.' Paralysis of the intestines.

ENTEROPARISAGOGE, Intussusception.

ENTEROPATHIA, *Enteropathia*, from *εντερον*, 'intestine,' and *πάθος*, 'disease.' Disease of the intestines in general.

ENTEROPATHIA CANCEROSA, *Enteroscirrhus*, *Enterostenoscirrhosa* seu *organica*, *Enterosarcoma*, *Induratio* seu *Scirrhus* seu *Cancer* seu *Carcinoma* *Intestino-rum*, (F.) *Cancer des Intestins*. Cancer of the intestines.

ENTEROPERISTOLE, *Enterospasmus*, *Hernia incarcerata*, *Incarceratio intestino-rum*, from *εντερον*, 'intestine,' and *περιστέλλω*, 'I contract.' Constriction or obstruction of the intestines, from a cause which acts either within the abdomen, or without it, as in strangulated hernia.

ENTEROPHLOGIA, Enteritis.

ENTEROPHLOGOSIS, Enteritis.

ENTEROPHTHISIS, from *εντερον*, 'intestine,' and *φθίσις*, 'consumption.' Consumption of the bowels. Consumption owing to suppuration in the intestines.

ENTEROPHYMATA, Tubercula intestino-rum.

ENTEROPLASTY, *Enteroplastics*, from *εντερον*, 'intestine,' and *πλάσσω*, 'I form.' A plastic operation for the restoration of an intestine.

ENTEROPLEGIA, Enteroparalysis.

ENTEROPYRA, Enteritis. Also, enteromesenteric fever, *Entéropyrrie*.—Alibert.

ENTEROPYRA ASIATICA, Cholera—e. Biliosa, Fever, bilious.

ENTERORRHAGIA, from *εντερον*, 'intestine,' and *ρῆγμα*, 'violent rupture.' Hemorrhage, or excessive discharge, from the bowels.

ENTERORRHAGIA SIMPLEX, Hæmatochezia.

ENTERORRHAPHIA, *Enterorrhaphia*, from *εντερον*, 'intestine,' and *ρᾶψις*, 'a suture.' Suture of the intestines, for the relief of injuries done to them.

ENTERORRHEUMA, *Rheumatismus Intestinalis*, from *εντερον*, 'intestine,' and *ρευμα*, 'de-fluxion, rheumatism.' Rheumatism of the intestines.

ENTERORRHEXIS, Enterobrosis.

ENTERORRHEA, Diarrhoea, Melæna.

ENTEROSARCOCELE, from *εντερον*, 'intestine,' *σάρξ*, 'flesh,' and *κύλη*, 'a tumour.' Intestinal hernia, complicated with fleshy excrescence, or rather sarcocele.

ENTEROSARCOMIA, Enteropathia cancerosa.

ENTEROSCHEOCELE, *Enteroschocele*, from *εντερον*, 'an intestine,' *σχῆλον*, 'the scrotum,' and *κύλη*, 'tumour.' Scrotal hernia consisting of intestine.

ENTEROSCIRRHUS, Enteropathia cancerosa.

ENTEROSES, from *εντερον*, 'an intestine.' A class of diseases, comprehending all those that are seated in the intestines.—Alibert.

ENTEROSPIGMA, Enteropistole.

ENTEROSTENOSIS, from *εντερον*, 'intestine,' and *στενωσις*, 'stricture.' Stricture or constriction of the intestines. See Ileus.

ENTEROSTENOSIS VOLVULUS, Ileus—e. Organica, Enteropathia cancerosa—e. Scirrhus, Enteropathia cancerosa.

ENTÉROTOME, *Enterotomus*, from *εἰς* *teron*, 'intestine,' and *τεμνω*, 'I cut.' A name given by J. Cloquet to an instrument for promptly opening the intestinal canal through its whole extent. It consists of a pair of scissors, one blade of which is much longer than the other, and rounded at its extremity. This is passed into the intestine.

ENTEROTOMIA, *Enterotomy*, same etymon. In anatomy, it signifies dissection of the intestines. In surgery, it means an operation, little used, which consists in opening the intestines, in order to evacuate the fecal matters accumulated in it; for example, in certain cases of hernia, accompanied with contraction of the intestinal canal; in operations for an artificial anus, performed on the new-born, where the rectum is imperforate or not existing, &c.

ENTEROTOMY, Enterotomia.

ENTEROTYPHUS, see Typhus.

ENTEROZOA, Worms.

ENTERYDROCELE, *Enterohydrocele*, from *εντερον*, 'intestine,' *ὕδωρ*, 'water,' and *κύλη*, 'rupture.' Intestinal hernia with hydrocele.

ENTHELMINTHES, Worms.

ENTHEOMANIA, Demonomania.

ENTHLASIS, *Esphlasia*, *Illisio*, from *εν*, 'in,' and *θλάω*, 'I break.' A fracture of the cranium, with comminution, in which the bone is depressed, or has lost its level.

ENTHLASIS CRANII, Effractura Cranii.

ENTITÉ, Entity.

ENTITY, *Entitas*, (F.) *Entité*; from *ens*, *entis*, 'a being.' The being or essence of anything. It has been used somewhat vaguely, in modern French medicine more especially, to signify apparently a general or essential disease, the nature and seat of which cannot be determined.—Nysten.

ENTOCELE, Ectopia.

ENTOCELE LENTIS. Dislocation of the crystalline into the anterior chamber of the eye.

ENTOGONYANCON, *Genu varum*, *Calf knee*, *X leg*, from *εντος*, 'within,' *γονυ*, 'the knee,' and *αγκων*, 'a bend.' Bending of the knees inwards. The state of being knock-kneed, or in-kneed. (Prov.) *Knar-kneed*, (Sc.) *Nule-kneed*.

ENTOHYALOID MUSCÆ, see Metamorphopsia.

EPIONIA, Tension, Tonic spasm.

ENTONIC, *Entonicus*, from *εν*, denoting excess, and *τονος*, 'tone.' Having great tension or exaggerated action.

ENTONNOIR, Calix, Infundibulum—e. du *Ventricule Moyen* du *Cerveau*, Infundibulum of the brain.

ENTEROPARASITES, Worms.

ENTOPHTHALMIA, from *εντος*, 'within,' and *ὀφθαλμία*, 'inflammation of the eye.' Ophthalmia affecting the interior of the eye.

ENTOPHYTE, *Entophyton*, (pl. *Entophyta*.) from *εντος*, 'within,' and *φυτον*, 'a vegetable.' A vegetable parasite, and especially one growing in the interior of the body. The following are the chief entophytes:—

<i>Alga</i>	of the mouth.
<i>Achorion Schönleini</i>	in Porrigio favosa.
<i>Achorion Lebertii</i>	in Porrigio scutellata.
<i>Microsporon Audouini</i> ..	in Porrigio decalvans.
<i>Mycoderm</i>	of Plica Polonica.
"	of Mentagra.
"	of Muguet.
"	of ulcerated and mucous surfaces.
<i>Sarcina ventriculi</i>	of the stomach.
<i>Torula</i>	" "

ENTOPHYTIC, *Entophyticus*, same etymon. Of or belonging to an entophyte, —as 'an entophytic growth.'

ENTOPTERYGOID, see Pterygoid.

ENTOPTIC, *Entopticus*, from *εντος*, 'within,' and *οφθς*, 'vision.' Relating or appertaining to the vision of objects within the organ of sight,—as of the bloodvessels of the retina.

ENTORSE, Sprain.

ENTOSTHETHIDIA, Entrails.

ENTOSTHIA, Entrails.

ENTOSTOSIS, Entostosis.

ENTOTORRHŒA, Otirrhœa.

ENTOTOZENA, Ozæna.

ENTOXICISMUS, Poisoning.

ENTOXISMUS, Poisoning.

ENTOZOA, (sing. *Entozoon*,) Worms.

ENTOZOAIRES, Worms.

ENTOZOARIA, Worms.

ENTOZOOGENESIS, Helminthiasis.

ENTOZOON FOLLICULORUM, Acarus Folliculorum.

ENTRAILLES, Entrails.

ENTRAILS, *Splanchna*, *Encœlia*, *Viscera*, *Entosthidi'ia*, *Entos'thia*, *Entosthethidi'ia*, *Interal'nea*, from *enteralia*, a word of bad Latin, coming from *εντερον*, 'intestine.' It is used for the viscera enclosed in the splanchnic cavities, and especially for those in the abdomen; *Bowels*, *Guts*, (F.) *Entrailles*, *Viscères*. *Ecta* has had this latter signification, but it was most frequently used by the ancients, as by Pliny, for the thoracic viscera.

ENTREFESSON, Chafing.

ENTRICHOMA, from *εν*, 'in,' and *τριχωμα*, (*τρίς*, *τριχος*,) 'hair.' The tarsal cartilage, and the edge of the eyelids, in which the cilia are implanted.

ENTROPION, *Entrop'ium*, from *εν*, 'in,' and *τροπω*, 'I turn.' *Inver'sio palpebra'rum*, *Capillit'ium*, *Trichi'asis*, *Introrsep'tio* seu *Blepharopto'is entropium*, *Blepharel'o'is*, *Tri'chia*, *Tricho'is*. A name given to the inversion or turning inwards of the eyelids, so that the eye-lashes are directed towards the globe of the eye; irritate and inflame it, and give rise to the affection called *Trichi'asis*.

The contact of the hair with the surface of the eye occasions considerable irritation of the conjunctiva, which is soon followed by chemosis, ulceration of the eye, and other symptoms, such as fever, watchfulness, &c.

If the disease be *entropion*, as above defined, that is, dependent upon the inversion of the eyelids, it must be remedied,—either by dispersing the œdema or swelling of the eyelids, or by cutting out a portion of the skin. When the disease is dependent on a vicious direction of the cilia, they must be plucked out and the bulbs be cauterized.

ENTROPIUM, Entropion.

ENTYPOSIS, see Glene.

ENUCLEATION, *Enucleat'io*, from *enucleare*, *enucleatum*, (e, and *nucleus*,) 'to take out a kernel.' This term has been proposed for the operation of removing tumours, &c., without cutting into them.—Percy.

The word is used in *Pharmacy* for the operation of shelling or removing the kernels of any nut.

ENULA CAMPANA, Inula Helenium.

ENULON, from *εν*, 'in,' and *ουλον*, 'the gum.' The inner part of the gums. The external part was called *ουλον*, and that between the teeth *αρμος*.—Pollux.

ENURESIS, from *ενουρεω*, (*εν*, and *ουρεω*,) 'I void the urine in bed.' *Paru'ria incont'iens*, *Incont'ientia ur'ina*, *Excret'io Ur'ina involunta'ria* seu *inopportu'na*, *Mict'io involunta'ria*, *Uracrat'ia*, *Urorrhœ'a*, *Perirrhœ'a*, *Strangu'ria*, (Galen,) *Anischu'ria*, *Hyperure'is*, *Diam'nes*, *Involuntary discharge of urine*, *Incontinence of*

urine. This affection is most common in advanced life. It may depend on too great irritability of the bladder, or on distension, or injury of the fibres about its neck, paralysis of the organ, *Cysto-paraly'sis*, *Cysto-ple'gia*, the presence of an irregularly shaped calculus impacted in the urethra near its commencement; rupture of the bladder and urethra; renal disease; or on pressure exerted on the bladder by the distended womb or by a tumour. It often occurs after difficult labour, but generally yields, in the course of a week or ten days; the catheter being introduced twice a day in the mean time.

The treatment must of course vary according to the cause; and when the affection on which it is dependent cannot be removed, the discharge of urine may be prevented by exerting a degree of compression on the urethra, by means of appropriate instruments; or a urinary may be employed to receive the urine as it passes off.

ENVIE, Hangnail, Malacia, Nævus—e. de Vomir, Nausea.

ENYPNION, Somnium.

ENYSTRON, Abomasus.

ENZOÏTIA, Enzooty.

ENZOÏTY, *Enzoût'ia*, (F.) *Enzoûtie*, from *εν*, 'in,' and *ζωον*, 'animal.' An endemic disease attacking animals.

EP, EPH, EPI, *επ*, *εφ*, *επι*, 'upon, above;' in composition, generally means, 'augmentation, addition, increase, reciprocal action, repetition.' Hence:

EPACMAS'TICOS, from *επι*, and *ακραω*, 'I increase.' An epithet applied to fevers, *Febres epacmas'tice*, whose symptoms go on augmenting in violence, from the commencement to the termination. Such fevers are, also, called *Epanadidon'tes*.

EPAGOGIUM, Phimosi, Prepuce.

EPANADIPLOSIS, Anadiplosis.

EPANALEPSIS, Anadiplosis.

EPANAS'TASIS, from *επαστασθαι*, 'to cause to rise up.' Pustule. The formation of a tumour. The breaking out of an eruption—*Erupt'io exanthem'atis*. A pustule.

EPANASTEMA, Exanthem, swelling.

EPANCHEMENT, Effusion

EPANESIS, Remission.

EPANETUS, Remittent—e. Hectica, Hectic fever—e. Malignus flavus, Fever, yellow—e. Mitis, Remittent fever.

EPANORTHOSIS, Restauratio.

EPANTHEMA, Exanthem.

EPANTHISMA, Exanthem.

EPAPHŒRESIS, from *επαφαιρω*, (*επι*, *αφω*, and *αιρω*,) 'I take,' 'I take away.' *Repet'ita sublat'io* seu *evacuatio*. A repeated abstraction or evacuation. It was formerly used synonymously with repeated bloodletting; *Phlebotom'ia iterata*.—Galen.

EPAPHRISMOS, Despumation.

EPARMA, Tumour.

EPARSIS, Tumour.

EPAULE, Humerus.

EPENCranIS, Cerebellum.

EPENDYMA, Membrane.

EPENDYMA VENTRICULORUM, *Endy'ma* seu *Indumen'tum ventriculo'rum*, from *επι*, and *ενδυω*, (*εν*, and *δυω*,) 'I enter;' hence, *επενδυμα*, 'an upper garment or cloak.' The lining membrane of the ventricles of the brain, formed by a prolongation of the pia mater, and probably also of the arachnoid.

ÉPERON, Hippocampus minor—e. des Ar-tères, Spur.

ÉPERVIER, Accipiter.

ÉPÉRIÈRE DES MURAILLES, Hieracium murorum.

EPHEBÆON, *Ephebæum*, from *επι*, and *ῥήν*, 'pubes.' The parts of generation: also, the region of the pubes, as well as the hair upon the pubes. It meant, likewise, a part of the ancient gymnasium in which the rhetoricians, philosophers, &c., disputed.

EPHEBEUM, Mons veneris.

EPHEBIA, *Ephebi'a*, *Pubertas*. Same etymology. The age of puberty.

EPHEBOSYNE, Puberty.

EPHEBOTES, Puberty.

EPHEBUS, *Pubens*, *Puber*, (F.) *Pubère*, *Éphèbe*. One who has attained the age of puberty.

EPHEDRANA, Nates.

EPHEL/CIS, from *επι*, and *ἔλκος*, 'an ulcer.' The crust or scab of an ulcer.

ÉPHELIDE SCORBUTIQUE, Chloasma.

EPHELIDES, from *επι*, and *ῥήλος*, 'the sun.' *Ephele'sis*, *Macula solaris*, *Macula fusca seu lenticular'es*, *Vitilig'ines*, *Phaci*, *Pannus lenticular'es*, *Lentic'ula*, *Lenti'go*, *Ephelis Lenti'go seu Lentic'ula seu à solè*, *Nigre'do à solè*, *Spilo'sis ephelis*, *Æst'etes*, *Ephila*, *Freckles*, *Sunburn*, (Prov.) *Ferniti'ckles*, *Fantickles*, *Farn'tickles*, *Fern'freckles*, *Fanfreckles*, *Forn'peckles*, *Freckens*, *Mur'les*, (F.) *Taches de rousseur*, *Hâle*. This term includes not only the yellow lentigines, which appear on persons of a fair skin, and the larger brown patches, which also arise from exposure to the direct rays of the sun, but also those large dusky patches which are very similar in appearance, but occur on other parts of the surface, that are constantly covered. See Chloasma. They do not extend farther than the skin. Many cosmetics have been recommended for their removal. Simple spirituous lotions or weak mineral acids, applied a few times in the day, are as effectual as any other means.

One, affected with freckles, is said to be *freckled*, (Sc.) *Faintickled*, *Fern'tickled*, (Prov.) *Farn'tickled*, *Fan'tickled*, *Fern'freckled*, *Forn'peckled*, *Freckened*, *Mur'led*, (F.) *Tacheté de Rousseurs*.

EPHELIS, see *Ephelides*—e. *Lentigo*, *Ephelides*—e. *à Sole*, *Ephelides*.

EPHELOTES, Leucoma.

EPHEMERA, *Di'aria*, from *επι*, and *ἡμερα*, 'a day.' That which continues a day. This epithet is given to diseases, and particularly to fevers, which last only a day.—*Febris di'aria seu simplex seu Ephe'mera*, *Ephemero'pyra*, *Febri'cula*, *Ephen'eral*, *Di'ary* or *Simple Fever*, (F.) *Fièvre éphémère*. The term prolonged *ephe'mera* is sometimes used for fevers, which cease after two or three days' duration.

EPHEMERA ANGILICA PESTILENS. *Sudor anglicus*—e. *Britannica*, *Sudor Anglicus*—e. *Maligna*, *Sudor Anglicus*—e. *Mortifera*, *Plague*—e. *Pestilentialis*, *Plague*—e. *Sudatoria*, *Sudor Anglicus*.

EPHEMEROPYRA, *Ephemera*.

EPHEMERUS, *Quotidian*.

EPHIALTES HYPOCHONDRIACA, *Incubus vigilans*—e. *Nocturnus*, *Incubus*—e. *Vigilantium*, *Incubus vigilans*.

EPHIDRO'SES, (G.) *Ephidrosen*. A division of the family *Ekzematosen*, of *Fuchs*, comprising morbid varieties of sweating.

EPHIDRO'SIS, from *επι*, and *ἵδρω*, 'I sweat.' *Hidro'sis*, *Idro'sis*, *Sud'dtio*, *Sweating*. Of old, it meant, at times, a slight sweating, *sudat'ion'cula*, of no value, and of bad augury; at others, a good and critical sweat. It has likewise been used for a copious sweat.

EPHIDROSIS PROF'USA, *E. spontanea seu idio-path'ica*, *Hydropede'sis*, *Exuda'tio*, *Exuda'tio*, *Hyper'ephidro'sis*, *Hyper'hydro'sis*, *Hydorr'rhæ'a*, *Hidropede'sis*, (F.) *Flux de Sueur*. An excessive sweating. A colliquative sweat. Such copious

perspirations are generally owing to debility of the exhalants, and require the use of tonics, especially of the mineral acids, opium, &c.

EPHIDROSIS, *Desudatio*—e. *Cruentus*, *Sudor cruentus*—e. *Saccharatus*, *Sudor dulcis*.

EPHIPPIUM, *Sella Turcica*.

EPH'ODOS, from *επι*, and *ὁδός*, 'a way,' 'a road or avenue to a place.' Hippocrates uses this word in three senses: 1. For the vessels or canals, which give passage to the excrements of the body. 2. For a periodical attack or accession of fever; and 3. For the approach of similar or dissimilar things which may be useful or hurtful to the body.—Castelli.

EPH'ATA SPRINGS. These are in Lancaster County, Pennsylvania. The water is pure; and facilities are afforded for bathing—warm and cold.

ÉPI, *Spica*.

EPI'ALOS, *Hepi'alos*, *Quer'cera*, *Quer'quera*. A name given by the ancients to fever, when the hot stage was mingled with irregular chills. Also, the cold stage of fever.

EPIALTES, *Incubus*.

EPIAMA, *Lentive*.

EPIAN, *Frambœsia*.

EPIBLEMA, *Catablema*.

EPIBOLE, *Incubus*.

EPICANTHIS, *Canthus*.

EPICAN'THIS, from *επι*, and *κάνθος*, 'the angle of the eye.' A defective formation, in which a fold of skin passes from the root of the nose over the inner canthus of the eye.

EPICAR'PIUM, *Pericar'pium*, from *επι*, and *καρπος*, 'the wrist.' An application made to the wrists, or to the region of the pulse, was so called.

EPICAUMA, *Encauma*.

EPICAUSIS, *Encauma*.

EPICERASTICA, *Emollients*.

EPICHOLOS, *Bilious*.

EPICHORDIS, *Mesentery*.

EPICHORION, *Decidia*.

EPICHORIUM, *Epidermis*.

EPICHRISIS, *Inunctio*.

EPICHRO'SIS, *Spilo'sis*, from *επι*, and *χρῶμα*, 'colour.' Simple discoloration of the surface.—Good. *Ephelides*.

EPICHRISIS SPILI, see *Nævus*.

EPICHYSIS, *Affusio*.

EPICELIDES, *Cilia*, *Palpebra* (superior.)

EPICE'MASIS, from *επι κοιμασαι*, (*επι*, and *κοιμασαι*, 'to lie down to rest,' 'to sleep.') *De'cubitus ad dormien'dum et Dormit'io*. The position of sleeping, as well as sleeping itself.

EPICOL'IC, from *επι*, and *κολων*, 'the colon.' The *epicolic regions*, *regio'nes epicol'icæ*, are those parts of the abdomen which correspond to the colon.

EPICOLLESIS, *Agglutination*.

EPICON'DYLE, *Epicon'dylus*, from *επι*, and *κονδύλος*, 'a condyle.' A name given, by Chausier, to an eminence at the outer part of the inferior extremity of the os humeri; so called, because it is seated above the condyle. The epicondyle gives attachment to the outer lateral ligament of the elbow joint, and to a very strong tendon to which several of the muscles of the posterior part of the forearm are attached.

EPICONDYLO-CUBITALIS, see *Anconæus*—e. *Radial*, *Supinator radii brevis*—e. *Sus-métacarp'ien*, see *Radialis*—e. *Sus-phalangettien commun*, *Extensor digitorum communis*—e. *Sus-phalangettien du petit doigt*, *Extensor proprius minimi digiti*.

EPICOPHO'SIS, from *επι*, and *κωφωσις*, 'deafness.' Deafness supervening on another disease.

EPICRA'NIUM, from *επι*, and *κρανιον*, 'the cranium.' A name given to different parts seated on the cranium. The skin, aponeurosis between

the occipital and frontal muscles, and the whole of the scalp, have been so called. Portal includes under this name the occipito-frontalis muscle, pyramidalis nasi, and superior and anterior auricular muscles. The pericranium.

EPICRANIUS, Occipito-frontalis.

EPICRA'SIS, *Contempcratio*, from *ἐπι* *κραννυμι*, (ἐπι, and *κραννυμι*, 'I mix,') 'I temper.' A term used by the humourists for an amelioration of the humours. They called *Cure by Epicrasis*—*per epicrasin*, a mode of treatment by imagined alternatives, which they supposed to be possessed of the power of gradually correcting the vitiated humours.

EPICROUSIS, Percussion.

EPICTENIUM, Pubes.

EPICTETI MORBI, Acquired diseases.

EPICYEMA, Superfetation.

EPICYESIS, Superfetation.

EPICYSTOTOMIA, see Lithotomy.

EPIDESIS, Deligation.

EPIDEM'IC, *Epidem'icus*, *Epidem'icus*, (F.) *Epidémique*. Same etymon as Epidemy. Appertaining to an epidemy. An epidemy.

EPIDEMIC CONSTITUTION, Epidemy—*c.* Disease, Epidemy—*c.* Influence, Epidemy.

EPIDEMICITÉ, see Epidemy.

EPIDEMICUS, Epidemy.

EPIDEMIOG' RAPHY, *Epidemiograph'ia*, from *ἐπι*, 'upon,' *ἄνθρωπος*, 'the people,' and *γραφη*, 'a description.' A history or description of epidemics.

EPIDEMIOLOG'Y, *Epidemiolog'ia*, from *epidemy*, and *logos*, 'a description.' The doctrine of epidemics.

EPIDÉMIQUE, Epidemy.

EPIDEMIUS, Epidemy.

EPIDÉMY, *Epidem'ic*, *Epidem'ia*, *Morbus publicus seu popularis seu epidem'icus seu epidem'ius*, *Epidem'ic disease*, *E. influence*, *E. constitution*; from *ἐπι*, and *ἄνθρωπος*, 'the people.' (F.) *Epidémie*. A disease which attacks at the same time a number of individuals, and which is referred to some particular *constitutio aeris*, or condition of the atmosphere, with which we are utterly ignorant. It differs from *endemic*, the latter being owing to locality; but it is obvious, that should a particular epidemic constitution of the air—*epidemic'ity*, (F.) *épidémicité*—exist along with a favouring endemic condition, these combined influences may act in the causation of several of those serious and fatal complaints, which at times visit a district, and are never afterwards met with, or at least not until after the lapse of a considerable period.

EPIDENDRUM VANILLA, see Vanilla.

EPID'ERIS, *Epider'rhia*, *Epider'rhion*, *Hypod'eris*, *Hypoderm'is*, *Hypod'er'rhia*, from *ἐπι*, and *δερμα*, 'a skin.' This word, with some, means the nymphæ; with others, the clitoris; with others, again, the prepuce of the clitoris.

EPIDERMA, Epidermis.

EPIDERMATIS, Epidermis.

EPIDERM'IC, *Epiderm'icus*, *Epiderm'al*, *Epiderma'lis*, (F.) *Épidémique*: same etymon as epidermis. Belonging to the epidermis,—as

EPIDERMIC CELLS, see Cell, epidermic.

EPIDERMIDON'OSI, *Epidermidon'oi*, from *ἐπι* *δερμα*, 'epidermis,' and *ὄσος*, 'disease.' Diseases of the epidermis.

EPIDERMION, Epidermis.

EPIDER'MIS, *Epider'mion*, *Epider'matis*, *Epider'ma*, *Epicho'rion*, *Sum'mitas cutis*, *Cutic'ula*, *C. ex'tima seu extre'ma*, *Cutis ex'tima seu summa seu ult'ima*, *Pellis summa*, *Pellio'ula summa seu super'ior*, *Lam'ina prima cutis*, *Oper'culum cutis*, from *ἐπι*, 'upon,' and *δερμα*, 'the true

skin.' *Scarf-skin*, *Cuticle*. (F.) *Surpeau*, *Cuticule*. A transparent, dry, thin membrane, devoid of nerves and vessels, which covers all the surface of the body, except the parts that correspond to the nails. It appears to consist of minute scales, placed one above the other. Chaussier considers it to be formed and reproduced by an excretory action of the true skin; to act like a dry varnish, which prevents the immediate contact of bodies with the nervous papillæ; and consequently to deaden tactile impressions, which, without its intervention, might be painful. The *Epider'mis*, *Epider'meous*, or *Epider'moid* (as Bichat called it) *Systém*, in general anatomy, comprises three parts. 1. External Epidermis. 2. Epidermis spread over the mucous membranes. 3. The Nails and Hair.

EPIDERMIS LINGUÆ, see Tongue.

EPIDERMOID, *Epidermoi'des*, from *epider'mis*, and *εἶδος*, 'resemblance.' Resembling the epidermis.

EPIDERRHION, Epideris.

EPIDERRHIS, Epideris.

EPIDERRHITIS, Nymphitis.

EPIDESIS, see Bandage—*c.* Hæmostasia, Ligature.

EPIDESMIS, Epidesmus.

EPIDESMUM, Epidesmus.

EPIDES'MUS, *Epides'mis*, *Epides'mum*, *Superliga'men*, from *ἐπι*, and *δω*, 'I bind.' A fasciæ, bandage or ligature, by which dressings are secured.—Hippocrates.

EPIDID'YMIS, from *ἐπι*, 'upon,' and *διδυμος*, 'a testicle.' *Epidid'ymus*, *Did'ymis*, *Paras'tate*, *Testic'ulus accessorius*, *Caput testis*, *Corpus varicosum seu varicifor'me testis*, *Spermatid'ia*, (F.) *Corpus variqueux ou variciforme*. That which is placed upon the testicle. A small, oblong, vermiform, grayish body, lying along the superior margin of the testicle. The Epididymis is a canal formed by the union of all the seminiferous vessels folded several times upon themselves, after having traversed the *Corpus Highmoria'num*. Its lower portion or tail, *Cauda seu globus seu caput minor*, is curved upwards, and is continuous with the vas deferens; the opposite extremity is called the head, *globus seu caput major*. The length of this canal, folded as it is upon itself and describing numerous curvatures, is, according to Munro, 32 feet.

EPIDIDYMITIS, Parastatitis—*c.* Blennorrhagic. Hernia humoralis.

EPID'OSIS, from *ἐπι*, 'upon,' and *δίδωμι*, 'to give.' Augmentation, increase. A word applied to the natural increase of the body, or to the increase of a disease.

EPID'ROME, *Epidrom'ia*, *Epiph'ora*, from *ἐπι* *δρῶμαι*, (ἐπι, and *δρῶμαι*, 'I run,') 'I run upon.' An afflux or congestion of humours.—Galen.

EPIDROMIA, Epidrome.

EPIFAGUS AMERICANUS, Orobanche Virginiana—*c.* Virginianus, Orobanche Virginiana.

EPIGÆ'A (properly EPIGÆ'A) REPENS, (ἐπι, 'upon,' and γη, 'the earth,' from its trailing growth.) *Trailing Arbutus*, *Ground Laurel*, *Mayflower*, called, by the Shakers, *Gravelplant*. A small, trailing plant, of the Family Ericaceæ, which grows in sandy woods, or in rocky soil, especially in the shade of pines. Its flowers appear in early spring, and exhale a rich, spicy fragrance. The leaves and stems are prepared and used like *uva ursi*.

EPIGASTRAL'GIA, from epigastrium, and αλγος, 'pain.' Pain at the epigastrium.

EPIGAS'TRIC, *Epigas'tricus*, *Epigas'trius*, from *ἐπι*, 'upon,' and *γαστρ*, 'the stomach.' This name has been given to several parts.

EPIGASTRIC AR'TERY, *Arteria epigas'trica*, *Artère sus-pubienne*, (Ch.) It arises from the

external iliac, on a level with the crural arch; sometimes by a trunk proper to it, at others by one common to it and the *Obturator artery*. It ascends upwards and inwards, behind the spermatic cord, follows the outer edge of the rectus abdominis muscle, and anastomoses, towards the umbilicus, with the internal mammary. The epigastric artery anastomoses, also, with the obturator, spermatic, lumbar, and intercostal vessels.

EPIGASTRIC REGION, *Regio epigas'trica* seu *cardi'aca* seu *stomach'ica* seu *stomacha'lis*, is the superior region of the abdomen, comprised between the false ribs on each side, and extending from the lower surface of the diaphragm to two fingers' breadth above the umbilicus. It is divided into three parts, one *middle*, the *epigas'trium*,—and two *lateral*, the *hypochond'ria*.

EPIGASTRIC VEIN follows nearly the same course as the artery.

EPIGASTRICUS, Epigastric.

EPIGAS'TRIUM. Same etymon. The belly; the epigastric region.

EPIGASTRICUS, Epigastric.

EPIGASTROCE'LE, from *επι*, *γαστρο*, 'the stomach,' and *κηλη*, 'a tumour.' Hernia, formed by the stomach; *Gas'troce'le*. This name has been given more especially to hernia, which occurs towards the upper part of the linea alba or in the epigastric region, whether formed or not by the stomach.

EPIGEA, Epigæa.

EPIGENEMA, Epigenesis.

EPIGEN'ESIS, from *επι*, and *γενεσις*, 'generation.' A theory of conception, according to which the new being is created entirely anew; and receives at once from each parent the materials necessary for its formation. Also, a new formation.

EPIGN'ESIS, as well as *Epigene'ma*, or *Epigenema*, is, also, applied to any symptom occurring during a disease, without changing its nature. An accessory symptom. *Epiginom'enos*, *Superveniens*, has a similar acceptation.

EPIGINOMENOS, see Epigenesis.

EPIGLOT'TIC, *Epiglot'ticus*: same etymon as Epiglottis. Relating to the epiglottis.

EPIGLOTTIC GLAND, *Periglot'tis*, *Caro glandulosa*. A collection of small, glandular granula, situate in the adipose, areolar texture at the base of the anterior surface of the epiglottis, in a triangular space, bounded anteriorly by the thyro-hyoid membrane and thyroid cartilage. It pours out an unctuous and mucous fluid, which lubricates the epiglottis, keeps it supple and movable, and prevents the larynx from being irritated by the constant passage of the air, in the act of respiration.

EPIGLOT'TIS, from *επι*, and *γλωττις*, 'the glottis.' *Epiglot'tic cartilage*, *Lig'ula*, *Operculum laryn'gis*, *Lingua Ex'igua*, *Ling'ula fist'ula*, *Sublingu'm*, *Superlig'ula*. A fibro-cartilage, situate at the upper part of the larynx, behind the base of the tongue. Its form is oval; texture elastic; thickness greater below than above, and greater in the middle than at the sides. By its smaller extremity, which is the lower, it is attached to the thyroid cartilage; its two surfaces are covered by the mucous membrane of the pharynx and larynx. The special use of the epiglottis would seem to be to cover the glottis accurately at the moment of deglutition, and, thus, to assist in opposing the passage of alimentary substances into the air tubes.

EPIGLOTTITIS, *Inflamma'tio Epiglot'tidis*, *Angi'na epiglottide'a*, (F.) *Inflammation de l'Epiglote*. Inflammation of the epiglottis.

EPIGLOT'TUM. An instrument, mentioned

by Paracelsus, for raising the eyelids. It resembled the epiglottis in shape.

EPIGLOU'TIS, from *επι*, and *γλωττις*, 'the buttocks.' The superior region of the nates.—Gorræus.

EPIGNA'THUS, from *επι*, 'upon,' and *γναθος*, 'the jaw.' A double monster, in which an incomplete fœtus is rooted, with its blood vessels in the palate of one more complete.

EPIGONĒ, *Gonē*, *Hypoph'ysis*, *Progen'ies*, *Proles*, *Sub'oles*, *Sub'oles*, from *επι*, and *γενν*, 'sperm.' Offspring. Progeny. Also, Superfecundation.

EPIGONION, Fœtus.

EPIGONIS, Patella.

EPIGONON, Superfecundation.

EPIGOUNIS, Patella.

EPIGUNIS, Patella.

ÉPILATOIRE, Depilatory.

EPILATORIUM, Depilatory.

EPILEMP'SIS, Epilepsy.

EPILENTIA, Epilepsy.

EPILEPSIA, Epilepsy—e. *Acuta* *nŕntum*, *Eclampsia*—e. *Algetica*, see *Algos*—e. *Febrilis* infantum, *Eclampsia*—e. *Nocturna*, *Incubus*—e. *Puerilis*, *Eclampsia* infantum—e. *Saltatoria*, *Chorea*—e. *Uterina*, *Lorind* *matricis*.

ÉPILEPSIE, Epilepsy—e. *Utérine*, *Hysteria*—e. *Vertige*, see *Epilepsia*.

EP'ILEPSY, *ἑπληψία* *vouscos*, *Epilep'sia*, *Epilep'sis*, *Epilemp'sis*, *Epilen'tia*, *Syp'sia* *Epilepsia*, *Catalen'tia*, *Decident'tia*, *Pas'sio cad'i'va*, *Heracle'us morbus*, *Hieran'osus*, *Deprehen'sio*, *Clonos epilep'sia*, *Prehen'sio*, *Perdit'io*, *Epilep'tica pas'sio*, *Morbus comit'ialis* seu *cadu'cus* seu *Hercule'us* seu *sacer* seu *astro'lis* seu *sidera'tus* seu *demoni'acus* seu *Sancti Johan'nis* seu *fœdus* seu *pueri'lis* seu *magnus* seu *major* seu *interlu'nus* seu *divi'nus* seu *dæmo'nus* seu *son'ticus* seu *seleni'acus* seu *lunat'icus* seu *menal'is* seu *deif'icus* seu *acces'tus* seu *vitriola'tus* seu *viridel'us* seu *Sancti Valenti'ni*, *Mulum cadu'cum*, *Analep'sia*, *Catapto'sis*, *Insa'nia cad'i'va*, *Apoplex'ia parva*, *Cadu'ca pas'sio*, *Lues divi'na*, *Vit'ium cadu'cum* seu *divi'nus* seu *Hercule'us*, *Falling sickness*, *Falling-down*, (Old E.) *Fallandevyl*, (F.) *Mal caduc* ou *divin* ou *Saint-Jean* ou *de Terre*, *Hautmal*, *Goutte caduque*. The word is derived from *επιλαμβάνω*, 'future,' *ἐπιληψω*, (*επι*, and *λαμβάνω*, 'I seize,') 'I seize upon.' It is a disease of the brain, which may either be idiopathic or symptomatic, spontaneous or accidental, and which occurs in paroxysms, with uncertain intervals between. These paroxysms are characterized by loss of consciousness and by convulsive motions of the muscles. Frequently, the fit attacks suddenly; at other times, it is preceded by indisposition, vertigo, and stupor. At times, before the loss of consciousness occurs, a sensation of a cold vapour is felt, hence called *aura epilep'tica*. This appears to rise in some part of the body, proceeds towards the head; and as soon as it has reached the brain the patient falls down. The ordinary duration of a fit is from 5 to 20 minutes. Sometimes it goes off in a few seconds; at others, it is protracted for hours. In all cases, there is a loss of sensation, sudden falling down, distortion of the eyes and face; countenance of a red, purple or violet colour; grinding of the teeth; foaming at the mouth; convulsions of the limbs; difficult respiration, generally stertorous; with, sometimes, involuntary discharge of feces and urine. After the fit, the patient retains not the least recollection of what has passed, but remains, for some time, affected with head-ach, stupor, and lassitude.

The disease is in the brain, and is generally

organic; but it may be functional and symptomatic of irritation in other parts, as in the stomach, bowels, &c. The prognosis, as to ultimate recovery, is unfavourable. It does not, however, frequently destroy life, but is apt to lead to mental imbecility. Dissection has not thrown light on its pathology.

To the attacks of epilepsy which are unaccompanied by convulsions, as is sometimes the case, the French give the name *Petit-mal*, and *Épilepsie Vertige, Cerebral Epilepsy*. Fully formed epilepsy is the *Grand-mal* of the French, *Spinal Epilepsy*. When furious mania succeeds to a paroxysm, it is termed *Mania epileptica* and *Epileptic Delirium*.

In the treatment, the cause must be sought after, and if possible removed. In the paroxysm, but little can be done, but as the tongue is liable to be injured by the teeth, the jaws may be kept open by putting a cork or piece of wood between them. If the fit has been brought on by indigestible food, the stomach must be cleared. It is between the paroxysms that the great effort must be made. Generally, there is considerable irritability and debility of the nervous system, and hence tonics have been found the best remedies. Of these, perhaps the most powerful, in epilepsy, is the *argenti nitras*, given regularly and continued for months, if necessary. Preparations of iron, copper, and zinc, have also been used, and vegetable tonics and antispasmodics in general. Counter irritants, as blisters, moxa, &c., may be employed, if necessary, along with this course.

Unfortunately, in many cases, these means are found insufficient, and all that can be done is to palliate, removing carefully the exciting causes; such as the use of spirituous liquors, strong emotions, violent exercise, &c.; and regulating the diet.

EPILEPSY, CEREBRAL, see Epilepsy—c. Spinal, see Epilepsy.

EPILEPTIC, *Epilepticus, Epileptus, Lunaticus, Canticans*; same etymon. One affected with epilepsy. Any thing relating to epilepsy; as an *epileptic individual*; an *epileptic symptom*. Some authors also speak of *epileptic remedies*.

EPILEPTOID, *Epileptoides*; same etymon. Resembling epilepsy—as *epileptoid symptoms*.

EPILOBIUM ANGIUSTIFOLIUM, (from *επι, λοβου, and ιον*, 'a violet on a pod,') *Great Willow herb*. An indigenous plant, *Order, Onagraceæ*, which flowers in July. The root is emollient and slightly astringent.

EPILOBIUM COLORATUM, *Purple-veined Willow herb*, has similar properties.

EPILOBIUM VILLOSUM, a South African plant, is used at the Cape of Good Hope as a domestic remedy for cleansing foul ulcers.

EPIMANES, from *επιμαίνομαι, (επι, and μαίνομαι, 'I am in a rage at,')* A name given to a maniac when in a paroxysm.

EPIMELE'TAL, from *επιμελομαι, (επι, and μελω, 'I take care,')* 'I take care of.' A name by which the ancient Persians called the cures of wounds, who followed their armies.

EPIMOR'IOS, 'unequal;' from *επι, and μετροω, 'I divide.'* An epithet applied to the pulse when unequal.—Galen.

EPINARD, Spinacia—*é. Sauvage, Chenopodium bonus Henricus*.

ÉPINE, Spine—*é. de Cerf, Rhamnus—é. du Dos, Vertebral column—é. Gutturale, Nasal spine, inferior and posterior—é. Sphénoïdale, Sphenoidal spine—é. Vinette, Oxyacantha Galeni*.

EPINEMESIS, Dispensation.

EPINEPH'ELOS, *Nubilo'vus*. Presenting a cloud. Applied to the urine.—Hippocrates.

ÉPINE'US, Spinous—*é. du Dos, grand, Spinialis dorsi major*: see *Interspinales dorsi et lumborum—é. du Dos, petit*, see *Interspinales dorsi et lumborum*.

ÉPINGLE, Pin.

ÉPINIÈRE, Spinal.

EPINOTION, Scapula.

EPINYCTIDES, from *επι, and νυξ, νυκτος, 'the night.'* Eruptions which appear on the skin in the night, and disappear in the day. *Pustula livens et noctibus inquit'ans*.—Pliny.

EPINYCTIS PRURIGINOSA, Urticaria.

ÉPIONE, Decidua.

EPIPAROXYS'MUS. A paroxysm super-added to an ordinary paroxysm.

EPIASTUM, Catapasma.

EPIPE'CHU, *επιπηνυ, from επι, and πηνυς, 'the elbow.'* The upper part of the elbow. The part above the elbow.—Gorræus.

EPIPEPHYCOS, Conjunctiva.

EPIPHENOM'ENON, *Post appa'rens, Epiphenomenon*; same etymon as the next. Any kind of adventitious symptom which occurs during the progress of an affection; and which is not as intimately connected with the existence of the disease as the symptoms properly so called.

EPIPHAN'IA, from *επι, and φαινω, 'I exhibit.'* The external appearance of the body.—Castelli.

EPIPEHGUS VIRGINIANA, *Orobanche Virginiana*.

EPIPHENOMENON, *Epiphenomenon*.

EPIPH'LEBUS, from *επι, and φλεψ, φλεβε, 'a vein.'* An epithet given to those whose veins are very apparent.—Hippoc., Aretæus.

EPIPHLEGIA, Inflammation.

EPIPHLOGISMA, *Erysipelatous inflammation*.

EPIPHLOGOSIS, *Hyperphlogosis*.

EPIPH'ORA, *Dacryoma, Lacrymatio, Dacrymatid, Ilacrymatid, Stillicid'ium lacrymat'rium, Oc'ulus lacrymans*; from *επιφω, (επι, and φω, 'I carry to,')* Weeping. (F.) *Larmoiement*. The watery eye;—an involuntary and constant flow of tears upon the cheek. It is almost always symptomatic of some disease of the lachrymal passages, and occurs when the tears cannot pass into the *ductus ad naum*. Occasionally, it is owing to the tears being secreted in too great quantity; as in certain cases of ophthalmia. The treatment must of course be directed to the removal of the cause.

EPIPHORA, Epidrome, Ophthalmia—c. Alvi, Diarrhœa—c. Ptyalismus, Salivation.

EPIPHYMATA, Cutaneous diseases.

EPIPH'YSIS, *Additamentum*, from *επι, and φω, 'I arise.'* Any portion of a bone, separated from the body of the bone by a cartilage, which becomes converted into bone by age. The *epiphysis* of the fœtus becomes the *apophysis* of the adult.

EPIPHYSIS CRURIS LONGIORIS INCUDIS, Os orbiculare.

EPIPHYTE, *Epiph'yton*, (pl. *Epiph'yta*), from *επι, and φυρον, 'a plant.'* *Phyto-parasite*. A parasite derived from the vegetable kingdom. Epiphytes are divided by Vogel into, 1. Vegetations in the human fluids, as the *tor'ula cerevis'ia* in vomited fluids, and faecal evacuations, and the *sar'cina ventriculi*. 2. Vegetations on the external skin and its appendages, as in *tinea favosa*, *mentagra*, *herpes tonsurans* and *plica polonica*; and, 3. Vegetations on the mucous membrane,—as in the aphthæ of children; in the cicatrices of the mucous membrane after typhus, &c. By some, the term *epiphytes* is restricted to those

vegetable parasites which grow on the surface of the body.

EPIPLASMA, from *επιπλασσω*, (*επι*, and *πλασσω*, 'I smear,') 'I plaster.' Cataplasma. Galen uses it especially for an application of wheaten flour, boiled in *hydrelaum*, and applied to wounds.

EPIPLEGIA, from *επι*, and *πληγη*, 'a stroke.' Paralysis of the upper extremities.

EPIPLEROSIS, from *επι*, and *πληρωσις*, 'repletion.' Excessive repletion; distension.

EPIPLOCĒ, from *επι*, and *πλεκω*, 'I twine.' Coition. Entanglement.

EPIPLOCĒ INTESTINALIS, *Conglomeratio intestino-rum*. Conglomeration of the intestines.

EPIPOCELE, from *επιπloon*, and *κηλη*, 'tumour.' *Epiploic hernia*, *Zirbalis hernia*, *Hernia omentalis*, *Omental hernia*, (F.) *Hernie de l'épiploon*. Omental hernia is recognised—but at times with difficulty—by a soft, unequal, pasty tumour, in which no gurgling sound is heard on pressing or reducing it. It is less dangerous than hernia of the intestines.

EPIPO-ENTEROCĒLĒ, from *επιπloon*, *εντερον*, 'intestine,' and *κηλη*, 'tumour.' Hernia, formed by the epiploon and a portion of intestine. It is more commonly called *Entero-epiplocele*.

EPIPOĒTEROÖSCHEOCĒLĒ, *Epiplo-enteroschocele*, from *επιπloon*, *εντερον*, 'intestine,' *σχων*, 'scrotum,' and *κηλη*, 'rupture.' Hernia with omentum and intestine in the scrotum.

EPIP'LOIC, *Epiploicus*. Relating to the epiploon.

EPIPOIC APPENDAGES, *Appendiculæ pinguedinosæ*.

EPIPOIC ARTERIES; *Arteria Epiploica*. The arterial branches, which are distributed to the epiploon, and which are given off by the gastro-epiploica.

EPIPO-ISCIOCEĒLĒ, from *επιπloon*, *ισχιον*, 'the ischium,' and *κηλη*, 'a tumour.' Hernia formed by the epiploon through the ischiatic notch.

EPIPLOITIS, from *επιπloon*, and *itis*, denoting inflammation. Inflammation of the omentum; *Omentitis*, *Empresma peritonitis omentalis*, *Omenti inflammatio*, *Enteritis epiploitis*, *Omentitis*. A form of partial peritonitis.

EPIPLOMEROCEĒLĒ, from *επιπloon*, *μηρος*, 'the thigh,' and *κηλη*, 'a tumour.' Femoral hernia, formed by the epiploon.

EPIPLOMPHALOCEĒLĒ, *Epiplophalon*.

EPIPLOMPHALON, *Epiploōm'phalon*, *Epiplomphalocele*, *Hernia umbilicalis*, from *επιπloon*, and *ομφαλος*, 'the navel.' Umbilical hernia, formed by the epiploon.

EPIPLOMPHRASIS, from *επιπloon*, and *εμψρασω*, 'I obstruct.' Induration of the epiploon.

EPIPLOCOMISTES, from *επιπloon*, and *κομιζω*, 'I carry.' One who has the omentum morbidly large. Also, one labouring under epiplocele.

EPIP'LOON, *Epiploom*, from *επι*, 'above,' and *πλωω*, 'I swim or float.' *Omentum*, *Retē*, *Reticulum*; *Dertron*; *Zirbus*, *Gan'ganē*, *Gangannum*, *Operimentum intestino-rum*, *Saccus epiploicus*, *Sage'na*, *The Caul*, (Prov.) *Flick*, *Web*. A prolongation of the peritoneum, which floats above a portion of the intestines, and is formed of two membranous layers, with vessels and fatty bands distributed through it. Anatomists have described several epiploons. The uses of the omentum are but little known. The chief one seems to be, to retain the viscera *in situ*, and to give passage to vessels.

EPIPLOON, COLIC, *Colic Omentum*, *Omentum colicum*, *Third Epiploon* or *Omentum*, is a duplicature of the peritoneum, situate along the as-

cending portion of the colon, as far as its junction with the transverse portion. It is behind the great epiploon, and fills up the angle formed by the junction of the ascending with the transverse colon. Its two laminae are separated by colic arteries and veins.

EPIPLOON, GASTRO-COLIC, *Great Omentum*, *Omentum gastro-colicum seu majus*, *Peritonæum duplicatum*, *Retē majus*, *Zirbus adiposus*, consists of an extensive duplicature, which is free and floating on the folds of the intestines. It is quadrilateral, and longer on the left side than on the right:—its base being fixed, anteriorly, to the great curvature of the stomach; and, posteriorly, to the arch of the colon. It is formed of two laminae, each consisting of two others. In this epiploon a number of vessels is distributed, and there is much fat.

EPIPLOON, GASTRO-HEPATIC, *Lesser Omentum*, *Omentum hepato-gastricum seu minus*, *Membrana macilentior*, is a duplicature of the peritoneum, which extends transversely from the right side of the cardia to the corresponding extremity of the fissure of the liver, and downwards from this fissure to the lesser curvature of the stomach, the pylorus and duodenum. Below it is the *foramen* or *hiatus* of WINSLOW; and between its laminae are lodged the biliary and hepatic vessels. It contains but little fat. If air be blown in at the foramen of Winslow, the cavity or sac of the omentum, *Sacculus Peritonæi retroventriculæris*, *Bursa omentalis*, will be rendered perceptible.

EPIPLOON, GASTRO-SPLENIC, *Gastro-Splenic Omentum*, *Omentum Gastro-splenicum*. A reflection of the peritoneum, which passes between the concave surface of the spleen and the stomach, from the cardiac orifice to near its great curvature, and which lodges the vasa brevia and splenic vessels between its laminae.

EPIPOSAKCOMPHALUS, from *επιπloon*, *σαρξ*, *sarkos*, 'flesh,' and *ομφαλος*, 'the navel.' Umbilical hernia, formed of indurated omentum.

EPIPOSCHEOCĒLĒ, *Epiploschocele*, from *επιπloon*, *σχων*, 'the scrotum,' and *κηλη*, 'tumour.' Scrotal hernia, formed by a displacement of the epiploon.

EPIPLOUM, Epiploon.

EPIPNOIA, Afflatus.

EPIPOROMA, from *επιπρωω*, (*επι*, and *πρωω*), 'I harden.' A tophaceous concretion which forms in the joints. The callus of fracture.

EPIR'HOĒ, *Epirrhœa*, from *επι*, and *ρρω*, 'I flow.' Afflux or congestion of humours.—Hippocrates.

EPIRRHŒA, *Epirrhœe*.

EPISARCIDIUM, *Anasarca*.

EPISCHESIS, Retention; from *επιχωω*, 'I restrain.' A suppression of excretions.

EPISCHET'ICUS, same etymon. *Sistens*, *Rep'rimens*, *Co'hibens*. 'Restraining.' An agent that diminishes—secretion, for example.

EPISCHION, Abdomen, Pubes, Pubis os.

EPISCOPALES VALVULÆ, Mitral valves.

EPISEION, Pubis (os), Vulva, Labium Pudendi, Tressoria.

EPISEMA'SIA, from *επισημαινω*, (*επι*, and *σημα*, 'a sign,') 'I afford some sign.' *Sema'sia*. It has been used synonymously with *sign*, and also with the invasion of an attack of fever. See Annotatio.

EPISTOCLE, see Hernia, vulvar.

EPISTŒDE'MA, from *επισιον*, (modern,) 'the labium pudendi,' and *οιδημα*, 'œdema.' Œdema of the labia pudendi.

EPISIOT'IS, from *επισιον*, 'the labium pudendi,' and *itis*, denoting 'inflammation.' Inflammation of the labia pudendi.

EPISION, Episeion.

EPISION'CUS, *Epiophy'ma*, from *επισιον*,

(modern,) 'the labium pudendi,' and *ογκος*, 'a tumour.' *Tumor labio'rum*. A swelling or tumour of the labia pudendi.

EPISIOPHYMA, Episioncus.

EPISIORRHAG'IA, from *επισιον*, 'the labium pudendi,' and *ρυσση*, 'rupture.' Hemorrhage from the labia pudendi.

EPISIORRHAPHY, *Episiorrhaph'ia*, from *επισιον*, 'the labium pudendi,' 'the vulva,' and *ραφη*, 'suture.' An operation practised in cases of prolapsus uteri, which consists in paring the opposing surfaces of the labia pudendi, bringing them together and uniting them by suture, so as to diminish the outlet of the vulva.

EPISPA'DIA; *Hyperspa'dia*; same etymon as the next. The condition of an Epispadias.

EPISPA'DIAS, *Epispadiæ'us*, *Anaspadiæ'us*, *Anaspadiæ'us*, *Hyperspadiæ'us*, *Hyperspadiæ'us*, from *επι*, 'above,' and *σπασω*, 'I draw.' The opposite to Hypospadias. One who has a preternatural opening of the urethra at the upper part of the penis.

EPISPASTICS, *Epispas'tica*, *Ves'icants*, *Helc'tica*, (F.) *Épispastiques*; same etymon. An epithet for every medicinal substance which, when applied to the skin, excites pain, heat, and more or less redness, followed by separation of the epidermis, which is raised up by effused serum. Cantharides and mustard are epispastics. See Blistar, and Derivative.

EPISPASTICUM, Blister.

ÉPISPASTIQUES, Epispastics.

EPISPASTUM, Catapasma.

EPISPHÆRIA, from *επι*, and *σφαيرا*, 'a sphere.' Some physiologists have applied this term to the different convolutions and sinuosities presented by the external surface of the brain.

EPISPAGISMOS, Signature.

EPISPLETIS, see Splenitis.

EPISTACTIS'CHESIS, from *επισταξις*, and *οσχευ*, 'to restrain.' Arrest of bleeding from the nose.

EPISTAPHYLINI, see Azygos Muscles.

EPISTASIS, *Insident'ia*, from *επι*, and *σπασω*, 'I rest.' A substance which swims at the surface of urine. It is opposed to the *hypostasis* or sediment. Epistasis has also been employed synonymously with suppression.

EPISTAX'IS, from *επι*, and *σταξω*, 'I flow drop by drop.' *Hæmorrhag'ia acti'va na'rrium*, *H. nari'ea seu na'rrium*, *Epistar'is junio'rum seu nari'ea seu hæmorrhin'ia*, *Hæmorrhin'ia*, *Hæmorrhag'ia*, *Hæmatis mus*, *Chæmorrhag'ia*, *Rhinorrhag'ia*, *Stillicid'ium seu Stillactio Sang'uinis e nari'ibus*, *Hæmorrhage from the pituitary membrane*, *Bleeding at the nose*, *Nosebleed*, (F.) *Hémorrhagie nasale*, *Saignement du nez*, *Hémorrhinie*. This is one of the most common varieties of hemorrhage; the organization of the Schneiderian membrane being favourable to it, as the blood-vessels are but slightly supported. It does not generally flow from both nostrils, and is very apt to recur. Puberty is the period at which it is most common. Its exciting causes are:—any thing that will induce local congestions, as running, coughing, blowing the nose, &c., provided there be a predisposition to it. A common cause of epistaxis requires but little treatment, especially if in a young person. Light diet and a dose or two of the sulphate of magnesia will be sufficient. In more severe attacks, cold and astringent washes of alum, sulphate of zinc, weak sulphuric acid, or creosote, may be used, and the nostrils be plugged anteriorly; but if the bleeding goes on posteriorly, the posterior nares must be plugged likewise,—the patient being kept with the head raised. The flow of blood has been arrested by directing the patient to stand up with his head

elevated, compressing the nostril whence the blood flows with the finger, raising the corresponding arm perpendicularly, and holding it in this position for about two minutes. A less vigorous circulation through the carotids appears to result from the increased force required to carry on the circulation through the upper extremities when raised.—Négrier. To prevent the recurrence, strict diet must be inculcated.

EPISTER'NAL, *Episterna'lis*, from *επι*, and *στερνον*, 'the sternum.' An epithet applied to two bones which form part of the sternum, and are situated upon its superior and lateral part. In the young subject, they are attached to the sternum by a synovial membrane, and a fibrous capsule. They have somewhat the appearance of the pisiform bones, but are of a greater size.

According to Professor Owen, the piece of a segment of an articulate animal, which is immediately above the middle inferior piece or sternum.

EPISTHOTONOS, from *επισθεν*, 'forwards,' and *τείνω*, 'I extend.' A spasm of the muscles, drawing the body forwards. The word *emprosthotonos* is more commonly employed.

EPISTROPHE, Relapse.

EPISTROPHEUS, Axis.

EPISTROPHUS, Axis.

EPISYNAN'CHÆ, (*επι*, and *synanche* or *cynanche*), *Episynangi'na*. A Greek word, used by a modern writer for *oedema* of the pharynx, by which deglutition is prevented, and the solid and liquid food driven back towards the mouth or nasal fossæ.

EPISYNTHET'IC, *Episynthet'icus*, from *επισυνθημι*, (*επι*, *συν*, and *τιθημι*), 'to collect, accumulate.' The name of a medical sect, whose object it was to reconcile the principles of the Methodists with those of the Empirics and Dogmatics. Leonides of Alexandria seems to have been one of the first partisans of this sect; of which we know little.

EPIT'ASIS, from *επι*, and *τείνω*, 'to extend.' The period of violence of a fever, (*Febris intensio*), attack or paroxysm; sometimes it is used in the same sense as epistasis, for 'suppression.'

EPITELIUM, Epithelium.

EPITHE'LIAL, *Epitheliæ'lis*; same etymon as *epithelium*. Appertaining or relating to the epithelium,—as *epithelial cells*, and *scales*.

EPITHE'LIAL CANCER, *Canceroid of the skin*, *Epithelioma*, *Carcinoma epitheliodes*, is a morbid growth of epithelial cells; forming an *epithelial tumour*, which, by some, is regarded as *carcinoma*. See *Canceroid*.

EPITHELIOMA, Epithelial cancer.

EPITHE'LIUM, *Epithelium*, *Epithel'ia*, *Epithel'ia*, *Endepiderm'is*, from *επι*, and *θηλη*, 'a nipple.' The thin layer of epidermis, which covers parts deprived of *derma*, properly so called,—as the nipple, mucous membranes, lips, &c. Modern histological researches have shown that it exists, in different forms, *pavement*, *cylindrical*, and *vibratile or ciliated epithelium*. *Tessellated*, *pavement*, *scaly*, *lamellar*, *tubular*, or *flattened epithelium*, *Epithelium polyed'ricum*, and *E. lamellosum*, (F.) *Épithélium pavimenteux*, covers the serous and synovial membranes, the lining of blood-vessels, and the mucous membranes, except where cylinder epithelium exists. It is spread over the mouth, pharynx and œsophagus, conjunctiva, vagina, and entrance of the female urethra. *Cylinder*, *Columnar* or *Prismatic epithelium*, *Epithelium cylind'ricum*, (F.) *Épithélium cylindrique* is found in the intestinal canal, beyond the cardiac orifice of the stomach; in the larger ducts of the salivary glands; in the ductus communis choledochus, prostate, Cowper's glands, vesiculæ seminales, vas deferens, tubuli uriniferi,

and urethra of the male; and lines the urinary passages of the female, from the orifice of the urethra to the beginning of the tubuli uriniferi of the kidneys. In all these situations, it is continuous with tessellated epithelium, which lines the more delicate ducts of the various glands. For the uses of the *ciliated epithelium*, *Epithelium vibrans*, see Cilia.

Mr. Bowman has applied the term *spheroidal* to a variety of epithelium, the cells of which, for the most part, retain their primitive roundness; or, being flattened where they touch, acquire a polyhedral figure. At times, these show a disposition to pass into the columnar, or the tessellated variety; and, hence, this form has been called the *transit'ional*. The spheroidal epithelium is found in the urinary passages, where it succeeds the columnar epithelium of the urethra at the internal orifice of that canal, and lines the whole of the bladder, ureters and pelvis of the kidneys. It is found, also, in the excretory ducts of the mammary, perspiratory, and of many mucous, glands; and a modification of it lines the inmost secreting cavities or commencing ducts of glands generally.

EPITHEM, *Epith'ema* or *Epithe'ma*, from *ἐπι*, and *τιθημι*, 'I put.' This term is applied to every topical application, which cannot be classed either under ointments or plasters. Three sorts of epithems have been usually distinguished,—the *liquid*, *dry*, and *soft*, which comprise *fomentations*, *bags filled with dry substances*, and *cataplasmata*.

EPITHESES. Same etymon. The rectification of crooked limbs by means of instruments.

EPITHYM'IA, from *ἐπι*, and *θυμος*, 'desire.' Morbid desires or longings.

EPITHYMIAMA, Fumigation.

EPITHYMIUM, *Cuscuta epithymum*—e. *Officinarium*, *Cuscuta Europæa*.

EPITROCHLEA, from *ἐπι*, and *τροχάλια*, 'a pulley.' (F.) *Epitrochlée*. That which is situate above a pulley. Chaussier gives this name to the unequal, round protuberance situate on the inside of the inferior extremity of the humerus above its articular trochlea. It is the part usually called the *inner* or *lesser condyle* of the humerus. From it originate several muscles of the forearm, to which it has given part of their names, according to the system of Chaussier.

ÉPITROCHLO-CARPI-PALMAIRE, *Palmaris longus*—*ε. Cubito-carpien*, see Cubital muscles—*ε. Métacarpien*, *Palmaris magnus*—*ε. Phalanginien commun*, *Flexor sublimis perforatus*—*ε. Radial*, *Pronator radii teres*.

EPIZO'A, (sing. *Epizo'on*), (F.) *Epizootères*, *Ectoparasites*, from *ἐπι*, and *ζωον*, 'an animal.' Parasitic animals, which infest the surface of the body, or the common integument.

EPIZOÔ'TIA, *Epizooty*, (F.) *Epizootie*. Same etymon. A disease which reigns amongst animals. It corresponds, in the veterinary art, to epidemic in medicine.

EPIZOÔT'IC, *Epizoot'icus*, same etymon. Relating or appertaining to an epizooty,—as an 'epizootic aphtha.'

EP'OCHE, from *ἐπι*, and *χω*, 'to have or to hold.' Doubt, suspension of judgment—*Suspens'ia centent'ia*. Sometimes employed in the same sense as *epicheia* or suppression.

EPOCHETEUSIS, Derivation.

EPODE, Incantation.

EPO'MIS, *Superhumeral'is*, from *ἐπι*, and *ωμος*, 'the shoulder.' The acromion. The upper part of the shoulder.—Hippocrates, Galen.

EPOMPHALICUM, *Emphalium*.

EPOMPHALION, Uterus.

EPOMPHA'LIUM, *Emphal'icum*, from *ἐπι*,

and *ομφαλος*, 'the navel.' A medicine, which, when placed upon the umbilicus, moves the belly.—Paulus of Ægina.

ÉPONGE, Spongia—*ε. d'Eglantier*, *Bedegar*.

ÉPOQUES, Menses.

EPOSTOMA, Exostosis.

EPOSTOSIS, Exostosis.

ÉPREINTES, Tenesmus.

EPEMA, Decoction.

EPESIS, Decoction, Elixatio.

EPSOM, MINERAL WATERS OF. Epsom is a town in Surrey, about 15 miles from London. The water is a simple saline;—consisting chiefly of sulphate of magnesia, which has consequently received the name of *Epsom salts*, though no longer prepared from the Epsom water.

ÉPUISEMENT, Exhaustion.

EPULIS, from *ἐπι*, and *ουλον*, 'the gum.' *Odon'tia exores'cens*, *Sarco'ma epul'is*, *Ulon'cus*, *Eccrescent'ia gingiv'ce*. (F.) *Épulide*, *Épulis*. An excrescence on the gum, sometimes ending in cancer.

EPULOSIS, Cicatrization.

EPULOTICA, Cicatrifiantia.

ÉPURGE, Euphorbia lathyris

EQUAL, *Æqual'is*. An epithet applied particularly to the pulse and to respiration. The pulse and respiration are equal, when the pulsations and inspirations which succeed each other are alike in every respect.

EQUILIBRIUM, from *æquus*, 'equal,' and *librare*, 'to weigh.' In medicine, this word is sometimes used to designate that state of organs, fluids and forces, which constitutes health.

EQUINA CAUDA, see Cauda Equina.

EQUI'NIA, from *equinus*, 'belonging to a horse.' *Glanders*, (F.) *Morve*. A dangerous contagious disorder, accompanied by a pustular eruption, which arises from inoculation with certain diseased fluids generated in the horse, the ass and the mule. Two forms are met with,—*E. mitis*, contracted from horses affected with grease, and *E. glandulo'sa*, a dangerous and commonly fatal disease communicated to man, either in the acute or chronic form, from the glandered horse. The veterinary surgeons make two varieties of the disease in the horse—*Glanders*, *Mal'ia*, *Malias'mus*, *Mal'lie*, *Malis*, *Malleus*; and *Farcy* *glanders*, *Farcino'ma*, *Farci'men*, *Mal'leus farcimio'sus*, *Morbus farcimino'sus*, *Cachex'ia lymphat'ica farcimino'sa*, (F.) *Morve farcineuse*, *Farcin*;—the former affecting the pituitary membrane and occasioning a profuse discharge from the nostrils, with pustular eruptions or small tumours, which soon suppurate and ulcerate, being attended by symptoms of malignant fever and by gangrene of various parts:—the latter being the same disease, but appearing in the shape of small tumours about the legs, lips, face, neck, &c. of the horse; sometimes very painful, suppurating and degenerating into foul ulcers. They are often seen together.

The mild cases require little treatment but rest; the more severe generally resist all remedies.

EQUSETUM ARVENSE, &c., see Hippuris vulgaris—e. *Minus*, *Hippuris vulgaris*.

EQUITA'TION, *Equita'tio*, *Hippe'ia*, *Hippa'sia*, *Hippe'usis*, from *equitare*, *equitatum*, 'to ride on horseback.' *Horseback exercise*. A kind of exercise, advisable in many diseases, although improper in others—in uterine affections, for instance. It has been much recommended in phthisis pulmonalis. It is less fatiguing than walking, and consequently more available in disease. But to prevent disease, where much exercise is required, the latter is preferable.

EQUIVOROUS, *Equi'vorus*, *Hippoph'agous*.

from *equus*, 'a horse,' and *voro*, 'I devour.' Feeding or subsisting on horseflesh.

EQUIVOCAL, *Equivocus*, from *æquus*, 'equal,' and *vox, vocis*, 'voice.' Those symptoms are occasionally so denominated which belong to several diseases. Generation is said by some to be 'equivocal,' when it is, or seems to be, spontaneous. See Generation.

EQUUS, *Hippus*.

ÉRABLE, *Acer saccharinum*.

ERADICA'TION, *Eradicatio*, from *e*, 'from,' and *radix, radicis*, 'a root.' The act of rooting out, or completely removing a disease.

ERAD'ICATIVE, *Eradi'cans*: same etymon. Any thing possessed of the power of completely rooting out a disease.

ÉRAILLEMENT DES PAUPIÈRES, *Ectropion*.

ERCLE, *Blister*.

EREBINTHUS, *Acer arietinum*.

ERECTEUR DU PÉNIS, *Ischio-cavernosus*.

ERECTHITES HIERACIFO'LIA, *Senecio hieracifolius*, *Fireweed*. An indigenous plant, *Order Compositæ*, which flowers from July to September. It is said to be an acrid tonic, and astringent, and in large doses emetic.

ERECTILE TISSUE, *Tela seu Textus erectilis*, from *erigere, erectum*, (*e*, and *rigere*), 'to erect,' 'to become erect.' A tissue, whose principal character is, to be susceptible of dilatation, i. e. of turgescence and increase of size. It is formed of a collection of arteries and veins, intermixed with nervous filaments; which form, by their numerous anastomoses, a spongy substance, whose areolæ communicate with each other. Smooth muscular fibres have been detected in it. This tissue exists in the corpora cavernosa of the penis and clitoris, at the lower and inner surface of the vagina, in the spongy part of the urethra, the lips, nipples, iris, &c. Sometimes it is developed accidentally, and constitutes a kind of organic transformation.

ERECTION, *Erectio, Arrectio*; same etymon. The state of a part, in which, from having been soft, it becomes stiff, hard, and swollen by the accumulation of blood in the areolæ of its tissue, as in the case of the penis—*Styma, Styxis*.

ERECTOR CLITORIDIS, *Ischio-cavernosus*—*c. Penis, Ischio-cavernosus*.

EREMACAU'SIS, (*F.*) *Érémacausie*, from *ερμος*, 'waste,' and *καυσis*, 'combustion.' A term applied by Liebig to the slow combustion, oxidation, or decay of organic matters in the air.

EREMIA, *Acinesia*.

ÉRÉSIPLELE, *Erysipelas*.

ERETHILYTICUM HÆMATOLYTICUM, *Spanæmic*.

ER'ETHISM, *Erethis'mus, Erethis'sin, Irritamentum, Irritatio*, (*F.*) *Érêthisme*, from *ερεθίζω*, 'I irritate.' Irritation. Augmentation of the vital phenomena in any organ or tissue. *Orgasm*. Under this name, Mr. Pearson has described a state of the constitution produced by mercury acting on it as a poison. He calls it *Mercu'rial Er'ethism* or *Erethis'mus*. It is characterized by great depression of strength; anxiety about the præcordia; irregular action of the heart; frequent sighing; tremors; small, quick, sometimes intermitting pulse; occasional vomiting; pale, contracted countenance, and sense of coldness; but the tongue is seldom furred, nor are the vital and natural functions much disturbed. In this state any sudden exertion may prove fatal.

ERETHISMA, *Rubefacient*.

ERETHISMUS, *Erethism, Irritation*—*e. Ebriorum, Delirium tremens*—*e. Hydrophobia, Hydrophobia*—*e. Oneirodynia, Incubus, Paroniria*—*a. Simplex, Fidgets*.

ERETHISTICUS, *Erethiticus, Rubefacient*.

ERETHIT'IC, *Erethit'icus, Erethis'ticus*. Belonging or relating to erethism—as "erethitic phenomena," or phenomena of irritation.

ÉRÉTISME, *Erethism*.

ERUGMOS, *Ereuctio*.

ERUXIS, *Ereuctio, Flatulence*.

ERGASMA, *Myrrha*.

ERGASTERION SPIRITUS, *Pulmo*.

ERGASTERIUM, *Laboratory*.

ERGOT, (*F.*) *Ergot*, 'a spur.' *Er'gota, Seca'le cornutum* seu *mater'mum* seu *tur'gidum* seu *temulentum* seu *lucubr'ians* seu *clavatum*, *Clavus temulentus* seu *lucubr'ians* seu *Silig'vinis*, *Spermoë'dia clavus*, *Mater seca'lis*, *Calcar*, *Seca'lis mater*, (*S-tile'go*, *Frumentum cornutum* seu *corniculatum* seu *lucubr'ians* seu *tur'gidum* seu *temulentum*, *Brizoc'eras* [*'*], *Melanophy'ma*, *Spurred or Horned Rye*, *Spur*, *Hornseed*, *Cocks'pur Rye*, *Cock'spur*, (*F.*) *Seigle ergoté* ou *cornu* ou *corrompu* ou *à Éperon* ou *Ergotisé*, *Blé cornu*, *Ergot* ou *clou* ou *mère de seigle*, *Bled avorté* ou *girouche*, *Faus seigle*. *Ergot* is generally considered to be the result of a disease in rye, occurring most frequently when a hot summer succeeds a rainy spring. Decandolle, however, regards it as a parasitic fungus, and calls it *Sclero'tium clavus*; whilst Leveillé esteems it to be a fungus giving a coating to the diseased grain; the medical virtues residing in the coating. This parasitic fungus he calls *Sphace'lia seg'etum*. More recently, it has been maintained, that it is a diseased state of the grain occasioned by the growth of a fungus not previously detected, to which the names *Ergotæ'tia abortans* seu *abortifaciens*, *Oidium abortifaciens*, *Ergot-mould*, have been given. It is found projecting from among the leaves of the spike or ear, and is a long, crooked excrescence, resembling the spur of a cock, pointed at its extremities, of a dark brown colour externally, and white within.

This substance has been long used in Germany to act on the uterus, as its names *Mutterkorn* and *Gebärpulver* ('womb-grain,' *pulvis parturiens, pulvis ad partum*) testify. Upwards of forty years ago, it was recommended in this country, by Dr. John Stearns, of Saratoga County, New York, for accelerating parturition; and since that period, numerous testimonials have been offered in its favour. Half a drachm of the powder is gently boiled in half a pint of water, and one-third part given every 20 minutes, until proper pains begin. Some recommend the clear watery infusion; others advise the powder; others the oil; and others the wine. It is obvious that, in many cases, the uterine efforts would return in the period which would necessarily elapse in the administration of ergot; so that several of the cases, at least, of reputed efficacy, may not have been dependent upon the assigned agent. *Ergot* also possesses, it is affirmed, narcotic virtues, which have rendered it useful in hemorrhagic and other affections of excitement.

Bread, made of spurred rye, has been attended with the effects described under *ERGOTISM*.

ERGOT, *Hippocampus minor*—*e. de Seigle*, *Ergot*.

ERGOT-MOULD, see *Ergot*.

ERGOTA, *Ergot*.

ERGOTÆTIA ABORTANS, see *Ergot*—*e. Abortifaciens*, see *Ergot*.

ER'GOTIN, *Extractum Er'gotæ*, *Extract of Ergot*, (*F.*) *Ergotine*, *Extrait hémotatique de Bonjean*. The extract of ergot has been found an excellent hemastatic in the dose of two grains several times a day. It has also been used externally.

ERGOTINE, *Ergotin*.

ERGOTISM, *Ergotismus*, from *ergota*, 'ergot.' *Morbis cerealis*. Poisoning by ergot. An affection produced by the use of spurred rye. At times, the symptoms are limited to vertigo, spasms, and convulsions. See *Convulsio cerealis*. Most commonly, there is torpor with numbness of the hands and feet, which waste away, lose sensation and the power of motion, and separate from the body by dry gangrene.—*Ergotismus Sphacelosus*, *Sphacelus cerealis*, *Necrosis cerealis*, *Gangræna ustilaginea*, *Necrosis ustilaginea*, *Ergot* or *Mildeu* mortification, (F.) *Ergotisme gangrèneux*, *Gangrène des Solnois*, *Mal de Solagne*.

ERGOTISME CONVULSIVE, *Convulsio cerealis*.

ERGOTISMUS, *Ergotismus*—e. *Spasmodicus*, *Convulsio cerealis*—e. *Sphacelosus*, *Ergotismus*.

ERICA VULGARIS, *Calluna vulgaris* seu *Erica*, *Common Heath*, (Sc.) *Heather*, (F.) *Brugère vulgaire*. This has been regarded as diuretic and diaphoretic.

ERIGERON AMBIGUUM, *E. Philadelphicum*—e. *Annum*, see *E. Canadense*.

ERIGERON BELLIDIFOLIUM, *Robin's plantain*, *Rosy Betty*, (*nyctepur*, from *ny*, 'spring,' and *pur*, 'an old man';) owing to the hoary appearance of some of the vernal species.) An indigenous plant of the Composite family; flowering in May. It is said to possess properties like those of *E. Philadelphicum*.

ERIGERON CANADENSE, *Canada Fleabane*, *Prideweed*, *Horseweed*, *Butterweed*, is considered to be allied in properties to *E. Philadelphicum*; and the same may be said of *Erigeron heterophyllum*. *E. annuum*, *Sweet Scabious* or *various-leaved Fleabane*; all of which are in the secondary list of the Pharmacopœia of the United States.

ERIGERON HETEROPHYLLUM, see *E. Canadense*—e. *Integrifolium*, *E. Philadelphicum*.

ERIGERON PHILADELPHICUM, *E. strigosum* seu *ambiguum* seu *integrifolium*, *Scabious*, *Skevich*, *Philadelphica* or *Skevich* or *Narrow-leaved Flea-Bane*, *Daisy*, *Cocash*, *Frostweed*, *Fieldweed*, *Squaw-weed*, (F.) *Vergerette de Philadelphie*. The plant has been used in decoction or infusion for gouty and gravelly complaints, and is said to operate powerfully as a diuretic and sudorific. The essential oil has been given in uterine hemorrhage. Dose, five drops.

ERIGERON STRIGOSUM, *E. Philadelphicum*.

ERIGERUM, *Senecio*.

ÉRIGNE, *Hook*.

ÉRINE, *Hook*.

ERIOCEPHALUS UMBELLULATUS, *Wild Rose-mary*. A South African shrub, *Ord*. Composite, which is diuretic, and used by the farmers and Hottentots in various forms of dropsy.

ERIOSPERMUM LATIFOLIUM. A South African plant, the tuber of which is muculent, and used externally in abrasions of the skin, and in superficial ulcers. It is also employed by the Mohammedans, in decoction, in amenorrhœa.

ERIX, *epix*. This word signifies, in Galen, the upper part of the liver.

ERODENS, *Caustic*.

ERODIUM MOSCHATUM, *Geranium moschatum*.

EROMANIA, *Erotomania*.

ERONOSUS, *Eronus*, from *ny*, 'spring,' and *nosos*, or *vovos*, 'disease.' A disease which appears chiefly in spring.

EROS, *Love*.

EROSIO, *Corrosion*—e. et *Perforatio spontanea Ventriculi*, *Gastromalacia*.

EROSION, *Erosio*, *Diabrosis*, *Anabrosis*, *Corrosion*, from *erodere*, *erosum*, (e, and *rodere*,) 'to eat away.' The action of a corrosive sub-

stance, or the gradual destruction of a part by a substance of that kind. It is often employed in the same sense as ulceration,—*spontaneous erosion*.

EROSION, *GANGRENOUS*, OF THE CHEEK, *Cancer aquaticus*.

EROTIC, *Eroticus*, from *eros*, *ερωτας* 'love' That which is produced by love—as *Erosic melancholy*, *E. delirium*, &c.

EROTICOMANIA, *Erotomania*.

EROTION, *Melissa*.

EROTOMANIA, *Eromania*, *Mania erotica*, from *eros*, *ερωτας*, 'love,' and *mania*, 'mania.' A species of mental alienation caused by love. *Erotic melancholy* is *Love melancholy*. Some authors write it *Erotico-mania*. Also, *Nymphomania*.

ERPES, *Herpes*.

ERRABUNDUS, *Planetes*.

ERRATIC, *Erraticus*, *Ataotos*, *Atypus*; from *errare*, *erratum*, 'to err.' Wandering, irregular. That which disappears and recurs at irregular intervals. This name is given to intermittents, *Febres erraticæ*, which observe no type. Most commonly, it is applied to pains or to any diseased manifestations which are not fixed, but move from one part to another, as in gout, rheumatism, erysipelas, &c.

ERRERE DE LIEU, *Error loci*.

ERRHINE, *Errhinum* or *Errhinum*, *Sternutatorium*, *Sternutatory*, *Ptaricum*, *Apophlegmatizans per nares*, *Nasalé*, from *er*, 'in,' and *rhin*, 'the nose.' A remedy whose action is exerted on the Schneiderian membrane, exciting sneezing and increased discharge. Errhines have been mainly used as local stimulants in head affections. The chief are: *Asari Folia*, *Euphorbiæ gummi-resina*, *Hydrargyri sulphas flavus*, *Tabacum*, and *Veratrum album*.

ERRHINUM, *Sternutatory*.

ERRHYSIS, from *er*, 'in,' and *rho*, 'I flow.' With some, a draining of blood. A trifling hemorrhage.

ERROR LOCI, *Apoplane'sis*, (F.) *Erreur de Lieu*. A term, frequently used by Boerhaave to express deviation of fluids, when they enter vessels not destined for them. Boerhaave admitted several orders of capillary vessels, the diameters of which went on decreasing. The largest received red blood, those which came next received only white, others only lymph, whilst the last of all were destined for yet more subtle fluids—for a kind of vapour. When the red globules entered vessels destined for the white, or the white those intended for the thinner fluids, there was, according to the theory of Boerhaave, an *error loci*, the proximate cause of inflammation and other mischiefs. See *Aberration*.

ERUCA, *Brassica eruca*, *Sinapis*—e. *Barbarea*, *Erysimum barbarea*—e. *Fœtida*, *Brassica eruca*—e. *Sativa*, *Brassica eruca*.

ERUCTATIO, *Eructatio*, *Erugismus*, *Erugge*, *Eructatio*, *Ructus*, *Ructa'men*, *Ructuatio*, *Eruxio*, *Restagnatio*, (Old Eng.) *Bolke*, *Bouffe*, *Belke*, (Sc.) *Bocking*, *Bokking*, *Rucking*, *Rift*, (Prov.) *Kelking*, *Rasping*, *Roeeping*, *Riving*, *Ruft* (N. of England); from *eructare*, *eructatum*, (e, and *ructare*, 'to belch.') *Belching*, (F.) *Rot*, *Rapport*. A sonorous emission, by the mouth, of flatus proceeding from the stomach. When so frequent as to occasion a diseased condition, this is termed *Ructuositas*, *Morbis ructuosus*.

ERUGATORIA, *Tetanothra*.

ÉRUGINEUX, *Eruginus*.

ERUPTIO, *Eruption*—e. *Exanthematis*, *Epanastasis*—e. *Sanguinis*, *Hæmorrhagia*.

ERUPTION, *Eruptio*, from *erumpere*, *eruptum*, (e, and *rumpere*,) 'to issue with violence,' 'to break out.' This word has several meanings.

1. The sudden and copious evacuation of any fluid from a canal or cavity—of serum, blood, pus, &c.,

for example; 2. The breaking out of an exanthem; and, 3. the exanthem itself, whether simply in the form of a rash, or of pustules, vesicles, &c.

ÉRUPTION ANOMALE, *Roseolæ*—*e. Roseolæ*, *Roseolæ*—*e. Violet*, *Ionthus*.

ERUPTIVE, *Eruptivus*. Same etymon as Eruption. That which is accompanied by an eruption. Thus small-pox, measles, scarlet fever, miliaria, &c., are *eruptive fevers*. The term *eruptive disease* is nearly synonymous with cutaneous disease.

ERVA DE SANCTA MARIA, *Arum Dracunculæ*—*e. do Rato*, *Palicourea Maragraavii*.

ERVALENTA, see *Ervum*.

ERVUM, *E. Ervilia*, *E. plicatum*, *Vicia ervilia*, *Orobus*, (F.) *Lentille* *Ers* ou *Orobe*. In times of scarcity, the seeds have been made into bread, but it is said not to be wholesome. The meal was once much used in poultices: it was formerly called *Orobion*, *orobion*. Under the name *Ervalemta*, a secret dietetic preparation has been introduced. It is a vegetable powder, which, when mixed with milk and soup and taken night and morning, is said to have succeeded in removing old and obstinate obstructions! Buchner thinks it is bean-meal, *Ditterich*, the meal of the seed of *Ervum Ervilia*.—*Aschenbrenner*.

A specimen of Warton's *Ervalemta*, obtained at the dépôt, and analyzed by Dr. Hassall, was found to contain a mixture of the French and German lentil, reduced to powder, including portions of the shells or husks, and of a substance very closely resembling, in its microscopic characters, *corn meal*. A sample of Du Barry's *Revalenta Arabica* was found to consist of a mixture of the Egyptian or Arabian lentil and barley meal.

ERVUM ERVILIA, *Ervum*.

ERVUM LENS, *Cicer Lens*, *Lens esculenta*, *Phacos*, the *Lentil*, (F.) *Lentille*. The seed, *Phacé*, *Phacée*, *Phacus*, is eaten in many places like peas: they are flatulent and difficult of digestion.

ERVUM PLECATUM, *Ervum*.

ERYGE, *Eruetation*.

ERYNGIUM, *Ηρυγγιον*, *Ηρυγγος*, see *E. Aquaticum*.

ERYNGIUM AQUATICUM, *E. yuccaefolium*, *Water Eryngo*, *Button Snake Root*. Order, Umbelliferae. This root, *Eryngium*, (Ph. U. S.) is nearly allied to the contrayerva of the shops, and acts more especially as a sudorific. It is a secondary article in the Pharm. U. S.

ERYNGIUM CAMPESTRIS, *E. vulgare*, *Centum Capita*, *Lyrin*gum, *Aster inquina*lis, *Capitulum Martis*, *Acus Ven'eri*s, (F.) *Panicaut*, *Char-don-Roland*, is sometimes used for *E. maritimum*.

ERYNGIUM FETIDUM, indigenous, is said to possess properties like those of valerian and contrayerva.

ERYNGIUM MARITIMUM seu MARI'NUM. The *Sea Holly* or *Eryngo*, *Æthe*rea *Herba*, (F.) *Panicaut Maritime*. *Eryngo* root has a slightly aromatic odour, and sweetish and warm taste. It is considered to be diuretic and expectorant, but its powers are so weak, that it is scarcely ever used.

ERYNGIUM VULGARE, *E. Campestre*—*e. Yucca-folium*, *E. aquaticum*.

ERYNGO, SEA, *Eryngium maritimum*—*e. Water*, *Eryngium aquaticum*.

ERYSIMUM, *E. officinale*, from *ερυσμα*, 'to draw,' from its power of raising blisters. *Sisymbrium officinale*, *Chama*epion. *Hedge-mustard*, (F.) *Vélar*, *Tortelle*, *Herbe aux Chantres*. This was, formerly, much used for its expectorant and diuretic qualities. The seeds greatly resemble mustard.

ERYSIMUM ALLIARIA, *Alliaria*.

ERYSIMUM BARBARE'A, *E. lyratum*, *Arabis barbare'a*, *Sisymbrium barbare'a*, *Eruca barbare'a*, *Barbare'a*, *B. stric*ta, (F.) *Herbe de Sainte*

Barbe, has been ranked amongst the antiscorbutic plants. See *Alliaria*.

ERYSIMUM CORNIPOLIUM, *Alliaria*—*e. Lyratum*, *E. barbarea*.

ERYSIPELA'CEA. Same etymon as the next. A family of eruptive diseases, comprising erysipelas, variola, rubella, and scarlatina.

ERYSIPELAS, from *ερυσ*, 'I draw in,' and *πelas*, 'near,' from its tendency to spread. *Febriis erysipelato'sa* seu *erysipela'cea*, *Erythrap'ura*, *Em'phlysis Erysi'elas*, *St. Anthony's fire*, *Wild-fire*, *Ignis Sancti Anto'nii* seu *Columellæ* seu *Sacer*, *Hieroppyr*, *Icteric*ia *rubra*, *Rose*, *Rubea icteric*ia, *Vicinitra'ha*, *Vicinitra'ctus*, *Brunus*, *Luglus*, (F.) *Erysipèle*, *Érysipèle*, *Feu St. Antoine*, *Feu sacré*, *Mal Saint Antoine*. A disease, so called because it generally extends gradually to the neighbouring parts. Superficial inflammation of the skin, with general fever, tension and swelling of the part; pain and heat more or less acrid; redness diffused, but more or less circumscribed, and disappearing when pressed upon by the finger, but returning as soon as the pressure is removed. Frequently, small vesicles appear upon the inflamed part, which dry up and fall off, under the form of branny scales. Erysipelas is, generally, an acute affection: its medium duration being from 10 to 14 days. It yields, commonly, to general refrigerant remedies. Topical applications are rarely serviceable. At times, when the disease approaches the phlegmonous character, copious bleeding and other evacuations may be required, as in many cases of erysipelas of the face; but this is not commonly necessary. In most cases, indeed, the general action seems deficient, and it becomes necessary to give tonics.

When erysipelas is of a highly inflammatory character, and invades the parts beneath, it is termed *Erysi'elas phlegmono'des*, (F.) *Érysipèle phlegmoneux*: when accompanied with phlyctenæ, and the inflammation terminates in gangrene, *Erysi'elas gangræno'sum*, (F.) *Érysipèle gangréneux*; and when associated with infiltration of serum, *Erysi'elas œdemato'sum*, (F.) *Érysipèle œdémateux*.

At times, the surface in erysipelas has a bronzed rather than a red hue—*bronzed erysipelas*.

ERYSIPELAS, BRONZED, see *Erysipelas*—*e. Gangrænoso*, see *Erysipelas*—*e. Œdematosum*, see *Erysipelas*—*e. Phlegmonodes*, see *Erysipelas*—*e. Periodica nervosa chronica*, *Pellagra*—*e. Phlyctenoides*, *Herpes zoster*—*e. Pustulosæ*, *Herpes zoster*—*e. Zoster*, *Herpes zoster*.

ERYSIPELATODES, *Erysipelatous*.

ERYSIPELATOUS, *Erysipelato'des*, *Erysipelato'sus*. Belonging to erysipelas; as an *erysi-pel'atous affection*.

ERYSIPEL'ATOUS INFLAMMATION, *Inflammatio erysi-pelato'sa*, *Epiphlogis'ma*, *Cuti'tis*, *Dermati'tis*, is the character of inflammation which distinguishes erysipelas, *Erythe'ma erysi-pelato'sum*.

ÉRYSIPE'LE, *Erysipelas*—*é. Ambulant*, see *Ambulant*—*é. Gangréneux*, see *Erysipelas*—*é. Œdemateux*, see *Erysipelas*—*é. Phlegmoneux*, see *Erysipelas*—*é. Serpiginieux*, see *Serpiginous*—*é. Pustuleux*, *Herpes zoster*.

ERYSOS, *Erythema*.

ERYTHE'MA, from *ερυθραιος*, *ερυθρος*, 'red, rose-coloured,' *ερυθραιω*, 'toadden.' *Inflammatory blush*, *Erythre'ma*, (F.) *Érythème*. This name is, sometimes, given to erysipelas, especially when of a local character. It is, also, applied to the morbid redness on the cheeks of the hectic, and on the skin covering bubo, phlegmon, &c. It forms the 6th genus of the 3d order, *Exanthemata*, of Bateman's classification of cutaneous diseases; and is defined, 'a nearly continuous redness of some portion of the skin, attended with disorder of the constitution,

but not contagious.' Many varieties are described by dermatologists,—for example, *E. intertrigo*, *E. fugax*, *E. papulatum*, *E. tuberculatum*, *E. nodosum*, *E. læve*, *E. centrifugum*, *E. marginatum*, and *E. acrodynamicum*.

ERYTHEMA ACRODYNIA, Acrodynamicum—e. Ambustio, Burn—e. Anthrax, Anthrax.

ERYTHEMA CENTRIFUGUM. A variety of erythema attacking the face, commencing with a small red spot, and spreading around, so as, at times, to affect the whole face.

ERYTHEMA ENDEMICUM, Pellagra—e. à Frigore, Chilblain.

ERYTHEMA FUGAX. Patches of erythema, that sometimes appear on the body in febrile diseases, teething children, &c., and which are of brief duration.

ERYTHEMA GANGRENOsum, Anthrax—e. Ichorosum, Eczema mercuriale—e. Intertrigo, Chafing.

ERYTHEMA LEVE. (F.) *Érythème léger*. Slight erythema affecting the skin, as in oedema.

ERYTHEMA MARGINATUM, (F.) *Érythème marginal*. A form in which the prominent patches of erythema are distinctly separated from the skin at their margins.

ERYTHEMA MERCURIALE, Eczema mercuriale.

ERYTHEMA NODOSUM, (F.) *Érythème noueux*. A more severe form of Erythema tuberculatum.

ERYTHEMA PAPULATUM, (F.) *Érythème papuleux*. Patches of erythema, which are at first papulated, appearing on the face, neck, breast, &c., of females, and young persons more especially.

ERYTHEMA PELLAGRUM, Pellagra—e. Pernio, Chilblain.

ERYTHEMA SOLARE. A name given by the Italians to Pellagra, owing to its being attended with an eruption of small red spots or patches on the parts that are exposed to the sun.

ERYTHEMA TUBERCULATUM, *E. tuberculatum*, (F.) *Érythème tuberculeux*. A form of erythema in which the patches are in small lumps.

ERYTHEMATOUS, *Erythemat'ic*, *Erythematous*, *Erythemat'icus*, (F.) *Erythémateux*. Appertaining or relating to Erythema.

ÉRYTHÈME, Erythema—*é. Léger*, Erythema læve—*é. Marginal*, Erythema marginatum—*é. Noueux*, Erythema nodosum—*é. Papuleux*, Erythema papulatum—*é. Tuberculeux*, Erythema tuberculatum.

ERYTHREA CENTAURIUM, Chironia centaurium—e. Chilensis, Canchalagua, Chironia Chilensis.

ERYTHRÆMIA, Plethora.

ERYTHRAN'CHE. *Angina erysipelato'sa*, from *ερυθρος*, 'red,' and *αγγω*, 'I suffocate.' Erysipelas or erythematous sore throat.

ERYTHREMA, Erythema.

ERYTHRINA MONOSPERMA, Butea frondosa.

ERYTHROCHÆRAS, Cynanche parotidæa.

ERYTHRODANUM, Rubia.

ER'YTHROID, *Erythro'des*, *Erythro'id*. Reddish.

ERYTHROID COAT, *Tunica erythro'des*. Vaginal coat of the testis.

ER'YTHROID VES'ICLE, *Vesic'ula Erythro'id*. A vesicle of the foetus described by Pockels and others. It is pyriform, and much longer than, although of the same breadth as, the umbilical vesicle. Velpeau, Weber, and others, doubt its being a physiological condition.

ERYTHRONIUM, E. Americanum.

ERYTHRONIUM AMERICANUM, *E. flavum* seu *dens canis* seu *lincolnatum* seu *longifolium*, Common Erythronium, (from *ερυθρος*, 'red,' which does not, however, apply to the American species,) Yellow Snake-leaf, Yellow Adder's tongue, Adder-leaf, Dog violet, Dog's tooth violet, Rattlesnake

violet, Lamb's tongue, Scroph'ula root, Yellow Snoudrop. Ord. Liliacæa. This plant is possessed of emetic properties, but is rarely, if ever, used. The root and herb, *Erythronium*, (Ph. U. S.), are in the secondary list of the Pharmacopæia of the United States.

ERYTHRONIUM DENS CANIS, E. Americanum—e. Flavum, E. Americanum—s. Lanceolatum, E. Americanum—e. Longifolium, E. Americanum.

ERYTHROPLEUM JUDICIALE, Sassy bark tree.

ERYTHROPYRA, Erysipelas.

ERYTHRORRHIZA ROTUNDIFOLIA, Galax aphylla.

ERYTHROSIS, Plethora.

ES'APHE, from *εσάφω*, (*εσ*, and *αφω*, 'I feel,') 'I feel inside,' (F.) *Le Toucher*, *Le Toucher vaginal*. The introduction of a finger or fingers into the vagina, for the purpose of examining the condition of the uterus. It is employed to verify the existence or non-existence of pregnancy and its different stages; to detect certain affections or faults of conformation in the genital organs and pelvis; and, during the process of accouchement, to examine into the progress of labour, &c.

ESCA, Aliment.

ESCARA, Eschar.

ESCARGOT, Helix pomatia, Limax.

ESCHALOTTE, Échalotte.

ESCHAR, *Es'chara*, *εσχα*, (Old Eng.) *Asker*, *a. slough*, *a. crust* or *scab*. The crust or disorganized portion, arising from the mortification of a part, and which is distinguishable from the living parts by its colour, consistence, and other physical properties. The inflammation which it excites in the contiguous parts, gives occasion to a secretion of pus between the living and dead parts, which ultimately completely separates the latter.

ESCHARA, Vulva.

ESCHAROT'IC, *Escharot'icum*, from *εσχα*, 'eschar.' Any substance, which, when applied upon a living part, gives rise to an eschar, such as *caustic potassa*: the concentrated mineral acids, &c.

ESCHAROTIC POISON, see Poison.

ESCHELLOHE, MINERAL WATERS OF. This water rises at the foot of a mountain near the convent of Etal in Bavaria. It is a sulphuro-saline.

ESCHID'NA OCELLATA. A very poisonous viper, which inhabits the sugar-cane fields of Peru. Its bite is almost instantaneously fatal.

ESCORZONERA, Scorzonera.

ESCREVISSE, Crab.

ESCULAPIAN, Medical.

ES'CULENT, *Esculen'tus*, *Ed'ible*, *Eat'able*, *Ciba'rious*, from *εσκα*, 'food.' (F.) *Comestibile*. Such plants or such parts of plants or animals as may be eaten for food.

ESÈRE, Ordeal nut.

ESKIN, Singultus.

ESO, *εσω*, 'within.' A prefix which signifies an affection of an internal part; as *Esoenteritis*, inflammation of the inner membrane of the intestines:—*Esogastritis*, inflammation of the inner membrane of the stomach, &c.

ESOCHAS, Esoche.

ES'OCHE, *Es'ochas*. A tumour—as a hemorrhoid—within the anus.

ESOCOLITIS, Dysentery.

ESODIC, Afferent.

ESOENTERITIS, see Enteritis.

ESOGASTRITIS, Endogastritis.

ESOTER'IC, *εσωτερος*, 'interior,' from *εσω*, 'within.' Private; interior; in contradistinction to *Exoter'ic*, public, exterior. Hence, we speak of 'esoteric and exoteric causes of disease.'

ESOTERISM, MED'ICAL. Same etymon as Esoterie. A term used by M. Simon, in his *Dé-ontologie médicale*, for that esotery, or mystery and secrecy, with which the practitioner performs his daily duties, and which, he conceives, he is compelled to adopt by the prejudices and ignorance of his patients.

ESPACES INTERCOSTAUX. see Costa.

ESPÈCES. Species, see Pulvis — *e. Pulvéru-lentes.* see Pulvis.

ESPÈCES VULNÉRAIRES. Falltranck.

ESPILYSIS. from *es, eis*, 'inwards,' and *φλασσω*. 'I break, bruise,' &c. A fracture of the skull by comminution, the fragments being depressed. See Depression, and Enthlasia.

ESPRIT. Spirit — *e. d'Ammoniaque*, Spiritus ammoniæ — *e. d'Anisi*, Spiritus anisi — *e. de Bois*, Spiritus Pyroxylicus — *e. de Camphre*, Spiritus camphoræ — *e. de Cannelle*, Spiritus cinnamomi — *e. de Carvi*, Spiritus carui — *e. d'Éther Sulfurique*, Spiritus ætheris sulphurici compositus — *e. de Genièvre composé*, Spiritus juniperi compositus — *e. d'Ileæ*, Achillea moschata — *e. de Lavande*, Spiritus lavandulæ — *e. de Lavande composé*, Spiritus lavandulæ compositus — *e. de Menthe poivrée*, Spiritus menthæ piperitæ — *e. de Muscade*, Spiritus myristicæ — *e. de Poudiot*, Spiritus pulegii — *e. Pyroxylic*, Spiritus pyroxylicus — *e. de Rai-fort composé*, Spiritus armoraciæ compositus — *e. de Rosmarin*, Spiritus rosmarini — *e. de Vin délayé*, Spiritus tenuior — *e. de Vin rectifié*, Spiritus rectificatus.

ESPRITS ANIMAUX. Nervous fluid.

ESQUILLE. Splinter.

ESQUINANCIE. Cynanche, Cynanche tonsillar.

ESQUILLINE. Smilax Chinn.

ESSENCE. *Essentia*, corrupted from *existentia*, 'standing out,' [?] or, from *esse*, 'to be.' By some, used synonymously with *volatile oil*; by others, with *simple tincture*. The oil of pepper-mint is the oil obtained by distillation; the *essence*, the oil diluted with spirit. See Tinctura olei menthæ.

ESSENCE OF ANISE. *Essentia anisi* — *e.* of Bardana, Hill's, Tinctura guaiaci ammoniata — *e.* of Caraway, *Essentia Carui* — *e.* of Cinnamon, *Essentia Cinnamomi* — *e.* of Coltsfoot, see Tinctura benzoini composita — *e.* of Fennel, *Essentia Feniculi* — *e.* of Musk, Tinctura moschi — *e.* of Mustard Pills, Whitehead's, see Sinapis — *e.* of Mustard, Whitehead's, see Sinapis — *e.* of Pennyroyal, *Essentia menthæ pulegii* — *e.* of Peppermint, Tinctura olei menthæ piperitæ — *e.* of Pimento, *Essentia Pimentæ* — *e.* of Rosemary, *Essentia Rosmarini* — *e.* of Roses, see Rosa centifolia — *e.* of Spearmint, Tinctura olei menthæ viridis — *e.* of Spruce, see Pinus Abies.

ESSENCES. Olea volatilata.

ESSENTIA. Essence, Tincture — *e.* Abietis, see Pinus abies — *e.* Aloes, Tinctura aloes — *e.* Absinthii amara, Tinctura A. composita — *e.* Absinthii composita, Tinctura A. composita.

ESSENTIA ANISI. *Essence of Anise.* (Ol. Anisi, f3j; Sp. rectif. f3ix. M. Ph. D.) Dose, ℥xx to f3j, as a carminative.

ESSENTIA ANTIMONII SEU STIBII. Vinum antimonii tartarizati — *e.* Antiseptica Huxhami, Tinctura cinchonæ composita — *e.* Aromaticæ, Tinctura cinnamomi composita.

ESSENTIA CARUI. *Essence of Caraway.* (Ol. Carui f3j; Sp. rectif. f3ix. Ph. D.) Dose, gtt. xx to f3j, as a carminative.

ESSENTIA CASTOREI. Tinctura castorei — *e.* Catholica purgans Rothii, Tinctura jalapii composita — *e.* de Cedro, Oleum cedrinum — *e.* Chinæ, Tinctura cinchonæ composita — *e.* Cinnamomi, Tinctura cinnamomi.

ESSENTIA CINNAMOMI. *Essence of Cinnamon.* (Ol. Cinnam. f3j; Sp. rect. f3ix. M.) Dose, gtt. x to xx.

ESSENTIA FENICULI. *Essence of Fennel.* (Ol. Fenic. f3j; Alcohol. f3ix. M. Ph. D.) Dose, gtt. xx to f3j.

ESSENTIA MENTHÆ PIPERITÆ. Tinctura olei menthæ piperitæ.

ESSENTIA MENTHÆ PULEGII. *Essence of Pennyroyal.* (Ol. Puleg. f3j; Sp. rectif. f3ix. Ph. D.) Dose, gtt. xv to xxx.

ESSENTIA MENTHÆ VIRIDIS. Tinctura olei menthæ viridis.

ESSENTIA MYRISTICÆ MOSCHATÆ. *Essence of Nutmeg.* (Ol. Myrist. f3j; Sp. rectif. f3ix. M. Ph. D.) Dose, gtt. xx—xxx.

ESSENTIA NEROLI. see Citrus aurantium — *e.* Corticis Peruviani antiseptica Huxhami, Tinctura cinchonæ composita — *e.* Corticis Peruviani composita, Tinctura cinchonæ amara.

ESSENTIA PIMENTÆ. *Essence of Pimento.* (Ol. Piment. f3j; Sp. rectif. f3ix. M. Ph. D.) Dose, gtt. xxx ad lx.

ESSENTIA ROSMARINI. *Essence of Rosemary.* (Ol. Rosmarini f3j; Sp. rectif. f3ix. M. Ph. D.) Dose, gtt. xxx ad gtt. lx.

ESSENTIAL. *Essentialis.* The word *essentia* has been applied to the immediate or active principles of vegetables, which were believed to be endowed with the properties of the plants from which they were extracted. *Essential oil* was so called because it was regarded as the only immediate principle which was *essential*. This expression is retained. *Essential salts* are true salts or extracts which exist, ready formed, in vegetables; and which are obtained by distillation, incineration, or some other process.

An *essential disease* is synonymous with a general disease; that is, one not depending on any local affection, — not symptomatic. See Entity.

ES'SERA. *Es'serè, Sora, Saire, Sare, Morbus porci'vus, Rosa saltans, Urticaria porcell'vina,* (F.) *Ampoules, Porcelaine.* Of Arabic derivation. A species of cutaneous eruption, consisting of small, reddish tubercles over the whole body, accompanied by a troublesome itching. It seems to be a variety of lichen or urticaria. See Lichen tropicus.

ESSIEU. Axis.

ESSOUFFLEMENT. Anhelatio.

ESTHEMA. Vestitus.

ESTHESIS. Vestitus.

ESTHIOMÈNE. Esthiomenus.

ESTHIOMÈNUS. *Estiom'enus*, from *estiv*, 'I eat.' *Es'edens.* (F.) *Esthiomène.* That which devours or eats away. Certain ulcers and eruptions — *Dartres rongeantes* — are so called.

ESTIVAL. *Estivus*; from *æstus, æstatis*, 'summer.' Happening in summer, belonging to summer. This epithet is given to *summer diseases*, so called because they reign at that season, and appear to depend on the influence exerted by it on the animal economy. In the United States, the term *summer disease* or complaint means disorder of the bowels; as *diarrhæa*, *cholera morbus*, &c.

ESTOMAC. Stomach.

ESTRAGON. Artemisia dracunculus.

ESULA CYPARISSIAS. Euphorbia cyparissias — *e.* Major, Euphorbia palustris — *e.* Minor, Euphorbia cyparissias.

ESULE, GRANDE. Euphorbia lathyris.

ESURIES. Hunger.

ESURIGO. Hunger.

ESURITIO. Hunger.

ESYCHIA. Acinesia.

ÉTAGE. Stage.

ÉTAÏN, Tin.—*é. de Glace*, Bismuth.—*é. Gris*, Bismuth.

ÉTAÏT, Acme.—*é. Granuleux du Rein*, Kidney, Bright's Disease of the.

ÉTERNUEMENT, Sneezing.

ESTICULATIO, Castration.

ETHER, Æther, Æther sulphuricus—*e. Acetic*, see Æther—*e. Acétique*, see Æther sulphuricus—*e. Azoteux*, see Æther sulphuricus—*e. Azoteux alcoolisé*, Spiritus ætheris nitrici.

ETHER, CHLORIC, see Chloroform. Under the names concentrated chloric ether and strong chloric ether—a compound of pure chloroform and nearly absolute alcohol, in the proportion of one-third of the former to two-thirds of the latter—has been used as an anæsthetic by inhalation, in the same cases as sulphuric ether and chloroform. It is properly an alcoholic solution or tincture of chloroform.

ETHER, CHLOROHYDRIC, CHLORINATED, see Æther muriaticus.—*é. Chlorohydrique*, see Æther sulphuricus.—*é. Chlorohydrique chloré*, see Æther muriaticus.

ETHER, COMPOUND. A preparation consisting of a solution of chloroform in sulphuric ether, which has been used as an anæsthetic by inhalation, in the same cases as chloroform.

ÉTHER, HYDRATIQUE. Æther sulphuricus—*e. Hydriodic*, Æthér hydriodici.—*é. Hydrique alcoolisé*, Spiritus ætheris sulphurici compositus—*e. Hydrochloric*, see Æther—*e. Hydrochloric*, Spirit of, Spiritus ætheris muriatici—*e. Hydrocyanic*, Æther hydrocyanicus—*e. Iodhydric*, Æther hydriodici.—*e. Marine*, Æther muriaticus—*e. Muriatic*, Æther muriaticus—*e. Muriatic*, spirit of, Spiritus ætheris muriatici.—*é. Nitreux*, see Æther sulphuricus—*e. Nitric*, see Æther—*e. Nitrique*, see Æther sulphuricus—*e. Nitrous*, see Æther—*e. Phosphoric*, see Æther—*e. Prussic*, Æther hydrocyanicus—*e. Rectified*, Æther sulphuricus.—*é. Sulfurique*, Æther sulphuricus.—*é. Sulfurique alcoolisé*, Spiritus ætheris sulphurici compositus—*e. Sulphuric*, Æther sulphuricus—*e. Terebinthinated*, Æther terebinthinatus.—*é. Vitriolique*, Æther sulphuricus,

ÉTHEREAL, Æthereal.

ÉTHERÉ, Æthereal.

ÉTHEREOUS, Æthereal.

ETHERINE, HYDROCYANATE OF. Æther hydrocyanicus—*e. Muriate of*, Æther muriaticus.

ETHERISM. The aggregate of phenomena induced by the inhalation of ether—*Ætherismus*.

ETHERIZA'TION, *Ætherisa'tio*. The administration of ether by inhalation, as an anæsthetic.

ETHERIZED, *Ætherisa'tus*. Presenting the phenomena induced by the inhalation of ether.

ÉTÉROLAT, *Éthérat*. The product of the distillation of ether on aromatic substances.

ÉTÉROLATURE, see Tincture.

ÉTÉROLÉ. A liquid medicine formed of ether and medicinal principles united with it by direct solution or simple mixture.—*Béral*.

Éthérolé d'essence de térébinthine. See Liquor anodynus terebinthinatus.

ÉTÉROLIQUE, *Etherol'ic*. A medicine which has ether for excipient.—*Béral*.

ÉTÉROLOTIF. An etherolic medicine which is exclusively employed externally.

ETHICS, MEDICAL, Deontology, medical.

ETHIK, Hectic.

ETHIOPIAN, see Homo.

ETHIOPS, see Æthiops.

ETHISIS, Filtration.

ETHISMUS, Filtration.

ETHMOCÉPHALE, *Ethm-cep'h'alus*; from *ἔθμος*, 'a sieve,' 'the ethmoidal portion of the

nose,' and *κεφαλή*, 'head.' A genus of monsters, with eyes close together, but distinct; the nasal apparatus atrophied and rudimental.—*I. G. St. Hilaire*.

ETH'MOID, *Ethmo'des*, *Ethmoï'des*, *Ethmoï'deus*, from *ἔθμος*, 'a sieve,' and *εἶδος*, 'form.' Shaped like a sieve.

ETHMOID BONE, *Os ethmoidaleum seu multifor'me seu spongio'sum seu spongioi'des seu cribro'sum seu cribri'for'me seu cu'bicum seu crista'tum seu foraminulen'tum seu coloi'f'me seu colato'rium*, (F.) *Os cribreux*. One of the eight bones which compose the cranium; so called, because its upper plate is pierced by a considerable number of holes. The ethmoid bone is situate at the anterior, inferior, and middle part of the base of the cranium. It seems to be composed of a multitude of thin, fragile, semi-transparent laminae, which form more or less spacious cells, called the *Ethmoidal labyrinth* or *cells*, distinguished into *anterior* and *posterior*. These communicate with the nose, and are lined by a prolongation of the pituitary membrane. The ethmoid bone is constituted of compact tissue, and is surrounded by the *Ethmoid suture*. It is articulated with the *frontal* and *sphenoidal bones*, the *cornua sphenoidalia*, the *superior maxillary bones*, the *palate bones*, the *ossa turbinata inferiora*, the *vomer*, the *proper bones of the nose*, and the *lacrimal bones*.

ETHMOID'AL, *Ethmoidal'is*. Belonging to the ethmoid bone; as, *Ethmoidal cells*, *E. suture*, &c. The *Ethmoidal apoph'ysis* or *process* is the advanced part of the sphenoid bone, which articulates with the ethmoid. The *ethmoidal arteries* are two in number, the anterior of which arises from the ophthalmic artery. The origin of the other varies. The *Éthmocerue ethmoidale* of the os frontis receives the ethmoid bones. The *Ethmoidal veins* correspond to the arteries.

ETHMOSE, Cellular tissue.

ETHMOSYNE, Habit.

ETHMYPHE, Cellular tissue.

ETHMYPHITIS, *Inflammatio telæ cellulosa*, from *ἔθμος*, 'a sieve,' *ὑψή*, 'texture,' and *itis*, denoting inflammation. Inflammation of the cellular membrane.

ETHMYPHOTYLOSIS, Induration of the cellular tissue.

ETHNOGRAPHY, *Ethnograph'ia*; from *ἔθνος*, 'nation,' and *γραφία*, 'a description.' A history of the races and families of man.

ETHNOLOG'OGY, *Ethnolog'ia*; from *ἔθνος*, 'nation,' and *λογος*, 'a discourse.' The science of the races and families of man.

ETHOS, Habit.

ÉTHUSE, *Æthusa cynapium*—*é. Meum*, *Æthusa meum*.

ETHYLE, CHLORIDE OF, Æther muriaticus—*e. Cyanuret of*, Æther hydrocyanicus.

ETICK, Hectic.

ETIOLATION, *Blanching*, (F.) *Étiollement*. That state of plants which occurs in consequence of privation of light, when they become pale and watery. In *pathology*, it is sometimes used to denote the paleness produced in those persons who have been kept long without light; or a similar paleness, the result of chronic disease.

ÉTIOLEMENT, Etiolation.

ETIOLOGY, *Ætiologia*.

ÉTIQUETTE, MEDICAL. Deontology, medical.

ÉTISE, Hecticis.—*é. Mésentérique*, *Tabes mesenterica*.

ETNAGH-BERRIES, see *Juniperus communis*.

ÉTOILE, *Stella*.

ÉTONNÉE, Stunned.

ÉTOUFFEMENT, Suffocation.

ÉTOUPE, Stupa.

ÉTOURDISSEMENT, Vertigo.

ETRANGLE-LOUP, Paris.

ÉTRANGLEMENT, Strangulation, Hysteria
—*é. des Intestins*, Enteremphaxis.

ÉTRANGUILLON, from *étrangler*, 'to strangle.' An ancient French appellation for a violent angina accompanied by symptoms of imminent asphyxia.

ÉTRIER, Stapes.

ETRON, Abdomen, Hypogastrium.

ETRON'CUS, from *ητρον*, 'the lower belly,' and *ογκος*, 'a tumour.' A tumour of the lower belly.

ETTICK, Hectic fever.

ETTLE, Urtica.

ÉTUI DE L'HIPPOCAMPE. A name given by Viqu d'Azay to the upper part of the sphenoidal portion of the lateral ventricle of the brain, which is bounded by the lateral cornu of the corpus callosum.

ÉTUVE, Stove—*é. Humide*, Bath, vapour.

ÉTUVER, to Foment.

EU, *eu*, 'good, proper,' when prefixed to words. Hence:

EUÉ'MIA, from *eu*, and *αἷμα*, 'blood.' A good condition of the blood.

EUËSTHË'SIA, from *eu*, and *αἰσθησις*, 'perception.' Vigorous perception. A good condition of the perceptive faculties.

EUËSTHË'TOS. Same etymon. One whose senses are in full vigor.

EUANALËP'SIS, from *eu*, and *αναληψις*, 'recovery.' Rapid restoration to strength.

EUANTHEMON, Anthemis nobilis.

EUCALYPTUS MANNIF'ERA, (*eu*, 'well,' and *καλυπτος*, 'covered.') From this Australian species, *Order*, Myrtaceæ, a manna-like substance exudes, which has similar properties to manna.

EUCALYPTUS RESINIFERA, see Kino.

EUCHRË'A, from *eu*, and *χρoια*, 'colour.' A good or healthy colour of the skin. A good appearance of the surface.

EUCHYM'IA, from *eu*, and *χυμος*, 'juice.' A good state of the humours.

EUCRA'SIA, from *eu*, and *κρασις*, 'temperament.' A good temperament.

EUADIAPNEUS'TIA, from *ευαδιαπνευστος*, (*eu*, and *διαπνοη*,—*δια*, and *πνοη*, 'breath'); 'one who breathes well.' Easy transpiration.

EUËCTICA (medicina), Gymnastics.

EUËL'CES, from *eu*, and *ελκος*, 'an ulcer.' One in whom wounds and ulcers are readily healed.

EUËMES, *Euēm'etos*, from *eu*, and *εμεω*, 'I vomit.' That which readily excites vomiting. One who vomits with facility.

EUETHES, Benign.

EUËX'IA, from *eu*, and *εξις*, 'constitution.' A good constitution.

EUFRASE, *Euphrasia officinalis*.

EUGENIA CARYOPHYLLA'TA, called after Prince Eugene. *Garryophyllus*, *Caryophyllus aromaticus*, *Myrtus caryophyllus*. The Indian tree which affords the clove. *Order*, Myrtaceæ. The Clove, *Caryophyllus*, *C. aromaticus*, is the unexpanded bud, (F.) *Girofle*, *Gérofle*, *Gyrophle*, *Clou de Girofle*. Its odour is strong, aromatic, and peculiar; taste, pungent and acrid. Like all substances whose virtue depends upon essential oil, it is stimulant and carminative. It is generally used as a corrigent to other remedies, and in cases where substances containing the essential oils are demanded. The oil—(F.) *Huile de Girofle*; *Oleum caryophylli*, Oil of Cloves—has the properties of the cloves.

EUGENIA PIMENTA, *Myrtus pimenta*.

EUGION, Hymen.

EULË, *εὐλη*, 'a worm.' *Eula*, *εὐλαι*, with Hippocrates, meant especially worms bred in a wound or ulcer.

EULOPHIA, see Salep.

EUNUCH, *Eunuchus*, from *ευνη*, 'the bed,' and *εχω*, 'I keep.' *Hemian'drus*, *Hemianor*, *Hemianthro'pus*, *Seminas*, *Seminascutus*, *Semimacula'tus*, *Semivir*, *Thla'dias*, *Thla'sias*, *Thlib'ias*, *Castrat'us*, *Gallus*, *Eneastrat'us*, *Evira'tus*, *Emacula'tus*, *Spado*, *Exsec'tus*, *Esteticula'tus*, *Euton'us*. (F.) *Eunuche*. One whose organs of generation have been removed, or so altered, that he is rendered incapable of reproducing his species, or of exercising the act of venery. Eunuchs were common with the ancient Romans. In Italy, this horrible mutilation still takes place to improve the voice; and in the East, eunuchs have the surveillance of the seraglio.

EUNUCHISMUS, Castration.

EUNUCHIUM MECONIS, Lactuca.

EUNUCHUS, Castratus, Eunuch.

EUNUQUE, Eunuch.

EUODES, Benevolentia.

EUONYMUS, Quassia simarouba.

EUONYMUS AMERICA'NUS, *Strawberry bush*, *Strawberry tree*, *Burning bush*, *Indian arrow-wood*. A shrub of the *Ord. Celastraceæ*, *Ses. Syst.* Pentandria Monogynia, which, like the next, is found throughout the United States and Canada; flowering from May to June.

EUONYMUS ATROPURP'UREUS, *Burning bush*, *Spindle tree*, *Indian Arrow-wood*, flowers from June to July.

From these varieties of *Euonymus* the *Wahoo* or *Whahoo bark* is said to be obtained. It is diuretic, antiperiodic, tonic, and a hydragogue cathartic, and has been used in dropsy in infusion, in the proportion of an ounce to a pint of water.

EUPATH'IA, from *eu*, and *παθος*, 'suffering.' A disposition for being affected by pain easily. Also, a good state of health—*Euphor'ia*.

EUPATOIRE D'AVICENNE, *Eupatorium cannabinum*—*c. Percefeuille*, *Eupatorium perfoliatum*.

EUPATORIUM, (called after Mithridates Eupator,) *Eupatorium perfoliatum*.

EUPATORIUM AGERATOI'DES, *E. Urticifolium*, *Ageratum altiss'imum*, *White Snakeroot*; indigenous; flowering in August and September; has the same properties as *Eupatorium perfoliatum*.

EUPATORIUM AROMAT'ICUM, *Aromatic Eupatorium* or *Hempweed*, has similar properties.

EUPATORIUM CANNAB'INUM, *Eupatorium*, *E. Arab'icum* seu *Japon'icum* seu *trifoliatum*, *E. of Avicenna*, *Cannab'ina aquatica*, *Trifolium cervi'num*, *Orig'anum aquaticum*, *Hemp ag'rimony*, *Herb of Saint Cuneconde*, (F.) *Eupatoire d'Avicenne*. *Ord. Compositæ. Ses. Syst.* Syngenesia æqualis. The juice of this plant proves violently emetic and purgative, if taken in sufficient quantity. It has been considered diuretic, cathartic, and emetic.

EUPATORIUM CONNATUM, *E. perfoliatum*—*e. Guaco*, *Guaco*—*e. Japonicum*, *E. Cannabinum*—*e. Mesues*, *Achillea ageratum*.

EUPATORIUM PERFOLIA'TUM, *E. Connatum*, *Boneset*, *Thor'oughwort*, *Thor'oughstem*, *Thor'oughwaax*, *Veg'table ant'imony*, *Crosswort*, *Ague-weed*, *Feverwort*, *Indian sage*, *Joe-pye*, *Tazel*, *Sweating plant*, (F.) *Eupatoire percefeuille*, is a plant which grows in low, wet meadows throughout the United States. The tops and leaves—*Eupatorium*, (Ph. U. S.)—are considered to be stimulating, tonic, emetic, purgative, diuretic, and sudorific. Dose, gr. xv of the powder, as a gentle purgative.

EUPATORIUM, PURPLE-STALKED, *E. purpuricum*.

EUPATORIUM PURPUREUM, Mohawk tassel, Purple-stalked *Eupatorium*, Trumpet weed, Joe Pye-weed, Gravelroot, is used in similar cases, as well as

EUPATORIUM ROTUNDIFOLIUM, Round-leaved Hempweed, Wild horehound;

EUPATORIUM SESSILIFOLIUM, Upland Boneset, and

EUPATORIUM TEUCRIFOLIUM, Wild horehound, German-leaved horehound, Rough boneset.

EUPATORIUM TRIFOLIATUM, *E. Cannabinum* — *e. Urticifolium*, *E. ageratoides*.

EUPEPSIA, from *eu*, and *πεψω*, 'I digest.' A good digestion. The antithesis to *dyspepsia*.

EUPEPTIC, *Eupepticus*, *Eupepticus*. Same etymon. Relating to a good digestion. One endowed with a good digestion.

EUPHLO'GIA, from *eu*, and *φλεγω*, 'I burn.' Mild inflammation. Variola.

EU'PHONY, *Euphonia*, from *eu*, and *φωνη*, 'voice.' An agreeable or regular voice.

EUPHORBE CYPRESS, *E. cyparissias* — *e. des Marais*, *E. palustris* — *e. Vomitive*, *E. ipecacuanha*.

EUPHORBIA CAPITATA, *Caecica*, *Ord.* *Euphorbiaceæ*. Called after Euphorbus, physician to king Juba. A Brazilian plant, which is strongly astringent and not poisonous. It is considered to be one of the best remedies for the bites of serpents. It is, also, called *Columbina Lusitanica*.

EUPHORBIA CAPUT MEDUSÆ. A South African plant, with which, according to Bruce, the wild inhabitants poison their arrows. The Ethiopians use, for the same purpose, *Euphorbia heptagona*, *E. virosa*, and *E. cereiformis*; the savages, in the most southern part of America, the sap of *E. cotinifolia*.

EUPHORBIA CEREIFORMIS, see *Euphorbia caput Medusæ*.

EUPHORBIA COROLLATA, *Large flowering spurge*, *Milkweed*, *Snakes' milk*, *Ipecacuanha*, *Hippo*, *Picac*, *Ip'ecac*, *Milk purslane*, *Purge root*, *Emet'ic root*, *Bowman's root*, *Apple root*, *Indian Physic*; indigenous; has similar properties to the last. The dose of the root is the same.

EUPHORBIA COTINIFOLIA, see *Euphorbia caput Medusæ*.

EUPHORBIA CYPARIS'IAS, *E. Cypressina*, *Esula minor* seu *Cyparissias*, *Tithym'alus Cyparissias*, *Cypress spurge*, *Welcome to our house*, (*F.*) *Euphorbe cypres*. This, like most of the spurges, is very acrimonious. Amongst the rustics, it was formerly called *poor man's rhubarb*, and was, consequently, a laxative. It is not used.

EUPHORBIA CYPRESSINA, *E. Cyparissias*.

EUPHORBIA HELIOSCOP'IA, *E. obturata*, *Sun spurge*, *Wartwort spurge*; indigenous. The juice has been given in syphilis. It is said to be cathartic.

EUPHORBIA HEPTAGONA, see *Euphorbia caput Medusæ*.

EUPHORBIA HYPERICIFOL'IA, *Large spotted spurge*, *Eyebright*. A native of the United States. It is astringent and tonic. Used in infusion — f3ss to Oj of boiling water. Dose, a tablespoonful. *EUPHORBIA MACULATA*, *Milk Purslane*, *Spotted spurge*, has similar properties.

EUPHORBIA IPECACUAN'HA, *Anisophyllum Ipecacuanha*, *Ipecacuanha spurge*, *Wild ip'ecac*. (*F.*) *Euphorbe vomitive*. This species of spurge is common in the southern and middle parts of the United States. The root is a powerful emetic, in the dose of from five to fifteen grains: twenty grains act as a cathartic likewise. In large doses, it acts as a narcotico-acrid.

EUPHORBIA LATH'YRIS. The systematic name of the plant which affords the lesser *catapulta* seeds, *Catapulta minor*, *Caper Spurge*, *Lath'yris*, *Garden spurge*, *Caper plant*, *Mole-plant*, *Tithym'alus latifolius* seu *lath'yris*, *Galurhaeus*

Lath'yris, (*F.*) *Épurga ou grande Ésule*. The seeds possess cathartic properties, and an expressed oil of the seeds — *O'leum Euphorbiæ Lath'yridis*, (*F.*) *Huile d'épurga* — has been given as a cathartic in the dose of six to twelve drops.

EUPHORBIA MACULATA, see *E. hypericifolia* — *e. Myrtifolia*, *Pedilanthus tithymaloides* — *e. Obturata*, *E. helioscopia*.

EUPHORBIA OFFICINARUM. The systematic name of the plant which affords the *Euphorbium*, *Euphorbiæ gummi-resi'na*, *Gummi euphorbiæ*, in greatest abundance. The euphorbium is an inodorous gum-resin, in yellow tears, which have the appearance of being worm-eaten. It enters into the composition of some plasters, and has been used as an errhine.

EUPHORBIA PALUS'TRIS, *E. paniculata*, *Greater spurge*, *Esula major*, *Tithym'alus palustris*, *Galurhaeus palustris*, *Marsh spurge*, *Wretweed*, (*F.*) *Euphorbe des marais*. The juice is given, in Russia, as a common purge. It is used, too, as an irritant in tinea, warts, &c.


EUPHORBIA PALUSTRIS and *E. VILLOSA*, or *E. PILOSA* have been brought forward as preventives of hydrophobia — the bitten part being washed with a decoction, and, at the same time, the decoction being taken internally.

EUPHORBIA PANICULATA, *E. palustris*.

EUPHORBIA PARAL'IAS, *Tithym'alus paralias*, *Sea spurge*. This is violently cathartic and irritating, and is seldom used.

EUPHORBIA PEPLUS, *Petty spurge*; indigenous. The powdered root is actively cathartic. It has been given in dropsy.

EUPHORBIA PILOSA, see *Euphorbia palustris* — *e. Tithymaloides*, *Pedilanthus tithymaloides* — *e. Villosa*, see *Euphorbia palustris* — *e. Virosa*, see *Euphorbia caput Medusæ*.

 All the spurges are vesicant and rubefacient, when applied externally.

EUPHORBUM, *Euphorbia officinarum*.

EUPHORIA, *Eupathia*.

EUPHRAGIA, *Euphrasia officinalis*.

EUPHRA'SIA (εὐφρασία, cheerfulness; so called from its effects), *E. Officin'alis* seu *Min'ima* seu *Imbricata*, *Ocularia*, *Euphra'gia*, *Ophthalmica*, *Eyebright*, (*F.*) *Enfraise*, *Euphrase*, *Casse-Lunette*. It has been recommended in diseases of the eye, but is unworthy of notice.

EUPLAST'IC, *Euplast'icus*, (*F.*) *Euplastique*: from *eu*, and *πλασσω*, 'I form.' Having the capacity of becoming organizable in a high degree, — as in false membranes resulting from acute inflammation in a healthy person.

EUPLEURUM SERRULATUM, *Diosma cre-nata*.

EUPNŒ'IA, from *eu*, and *πνέω*, 'I respire.' Freedom or facility of respiration.

EURHYTH'MIA, from *eu*, and *ρυθμος*, 'rhythm.' Regularity of the pulse.

EURIBALI, *Juribali*.

EURODES, *Carius*.

EURODON'TICUS, from *eupos*, 'caries,' and *ὀδοντος*, 'a tooth.' Suffering from carious teeth.

EURUS, *eurus*, 'mould, putrefaction,' *Mucor*, *Caries*. Corruption of the humours.

EURYCHORIA, *Sinus*.

EURYCLES, *Eugastri-myth*.

EURYCLITUS, *Engastri-myth*.

EURYSMUS, *Dilatation*.

EURYTH'MIA, *Euryth'mus*, from *eu*, and *ρυθμος*, 'rhythm.' Regularity of pulse, both as regards quantity and quality.

EUSAR'CUS, from *eu*, and *σαρξ*, *σαρκος*, 'flesh.' One who is fleshy, robust, muscular.

EUSEM'IA, (*F.*) *Eusémie*, from *eu*, and *σημειο*, 'a sign.' A collection of good signs.

EUSOMPHALIEN, from *eu*, and *σφαλος*,

'navel.' A double monster in which each being has a separate navel, and is able to execute, independently, almost all the vital functions.

EUSPLANCHNIA, from *eu*, and *σπλᾶγχνον*, 'a viscous.' A healthy state of the viscera.

EUSTACHIAN TUBE, *Tuba Eustachia'na* seu *Aristotel'ica*, *Syrin'ga*, *Syrinx*, *Med'tus cæcus*, *Ductus auris palati'nus*, *Iter a Pala'to ad Aures*, (F.) *Trompe ou Conduit d'Eustache*, *Conduit naturel de l'oreille*.—Ch. So called from its discoverer, Eustachius. It is partly bony and partly cartilaginous: extending from the cavity of the tympanum to the upper part of the pharynx. Its length is about two inches; the bony portion which belongs to the temporal bone, is about three-fourths of an inch long. It is lined, internally, by a prolongation of the lining membrane of the pharynx. Its nerves are furnished by the palatine branches of the ganglion of Meckel, and its vessels proceed from those of the pharynx and velum pendulum. The use of the tube seems to be, to permit the renewal of air in the cavity of the tympanum.

EUSTACHIAN VALVE, *Valve of Eusta'chius*, *Valvule d'Eustachi ou d'Eustache*. A membranous, semilunar fold, which corresponds to the opening of the vena cava inferior into the right auricle of the heart.

EUSTHENIA, *Vigor*, *Exuberan'tia*, from *eu*, and *σθενος*, 'strength.' Flourishing, exuberant health.—Hippocrates.

EUSTOMACHUS, from *eu*, and *στομαχος*, 'stomach.' Digesting rapidly. Having a good stomach.

EUSTRONGYLUS GIGAS, *Strongylus gigas*.

EUTAXIA, from *eu*, and *ταξις*, 'I arrange.'

Euthe'sia. A well-ordered constitution, in which every part has its proper relation. The ready return of a rupture, or of a luxated bone.

EUTHANASIA, from *eu*, and *θάνατος*, 'death.' An easy death. The opposite to this is *Cacothanasia*.

EUTHESIA, *Eutaxia*.

EUTHYENTERON, Rectum.

EUTHYENTEROSTENOMA, Stricture of the Rectum.

EUTHYMIA, *An'imi tranquill'itas*, from *eu*, and *θυμος*, 'mind.' Tranquillity of mind. A good state of the mental faculties.

EUTHYPNOE, Orthopnoea.

EUTHYPNOEA, Orthopnoea.

EUTOCIA, from *eu*, and *τοκος*, 'delivery.' An easy labour. Fecundity.

EUTROPHIA, from *eu*, and *τροφή*, 'nourishment.' A good state of nutrition.

EUTROPHIC, *Eutroph'icus*, same etymon. A term introduced into medical terminology, by the author, for an agent whose action is exerted on the system of nutrition, without necessarily occasioning manifest increase of any of the secretions.

The chief eutrophics are,—*mercurials*, the preparations of iodine, bromine, cod liver oil, the preparations of gold and silver, sulphur, sugar, and *sarsaparilla*.

EUZOODYNAMIA, *Sanitas*.

EVACUANTS, *Evacuan'tia*, from *e*, and *vacuare*, 'to empty.' (F.) *Evacuatif*. Medicines are so called which occasion a discharge by some emunctory; such as purgatives, emetics, &c.

EVACUATIFS, Evacuants.

EVACUATIO, Evacuation, Excretion.

EVACUATION. Same etymon. *Lapar'is*. The discharge of any matter whatever by the natural passages or by an artificial opening.

ÉVANOUISSMENT, Syncope.

EVAN ROOT, Geum Virginianum.

EVAPORATION, *Evapora'tio*, *Vapora'tio*,

Exhala'tio, from *e*, and *vaporare*, *vaporatum*, 'to emit a vapour.' Transformation of a liquid into vapour, in order to obtain the fixed matters contained in it dry and separate from the liquid. When the vapour is received into a proper vessel and condensed, the process is called distillation. Evaporation produces cold, and this is one of the processes by which the body is cooled, through the evaporation of the perspiratory fluid.

ÉVAUX, MINERAL WATERS OF. Evaux is situate in Auvergne, in France. The waters are hydrosulphurous and thermal.

EVENTRATION, *Eventra'tio*, *Hypogastro'rix'is*, from *e*, 'out of,' and *venter*, 'the belly.' A tumour, formed by a general relaxation of the parietes of the abdomen, and containing a great part of the abdominal viscera. Also, *ventral hernia*, or that which occurs in any other way than through the natural openings of the abdominal parietes. Lastly, any very extensive wound of the abdominal parietes, with issue of the greater part of the intestines.

EVENTUS, from *e*, 'out of,' and *venire*, *ventum*, 'to come.' *Apobat'ion*, *Apobesom'enon*, *Ec-besom'enon*, *Terminat'io morbi*. The event or issue of a disease, either favourable or unfavourable.

EVERLASTING, **DICECIOUS**, *Antennaria dioica*.

EVERRICULUM, *Speci'llum*, from *everrere*, 'to clean out.' A sort of sound or scoop, used for extracting sand or fragments of stone or clots of blood from the bladder, after or during the operation of lithotomy.

EVERSIO PALPEBRÆ, Ectropion.

ÉVIGILATION, see Sleep.

EVIL, THE, Scrofula—e. King's, Scrofula.

EVIRATIO, Castration.

EVIRATUS, Castratus, Eunuch.

EVISCERATION, Exenterismus.

EVOLUTIO, Development—e. Spontanea, Evolution, spontaneous.

EVOLUTION OF GERMS, see Generation.

EVOLUTIO, **SPONTANEOUS**, *Ver'sio seu Evolut'io sponta'nea*, from *e*, and *volvere*, *volutum*, 'to roll.' *Sponta'neous ver'sion*, *Sponta'neous expul'sion*, *Duplicat'io of the fœtus*. A term, applied, by Dr. Denman, to what he considered to be a spontaneous turning of the fœtus in utero, in an arm presentation, in consequence of powerful uterine contractions forcing out the breech and feet, whilst the arm recedes. It is now usually considered to be a doubling of the fœtus, so that the arm changes its position but little, whilst the breech is forcibly expelled before the upper extremity;—the case becoming similar to a breech presentation.

EVOLUTION, **VESICLE OF**, Vesicle of development.

EVOMITIO, Vomiting.

EVONYMUS, Euonymus.

EVULSION, *Avulsion*, *Evul'sio*, from *evellere*, *evulsus*, (e, and *vellere*), 'to pluck out.' *Apot'il'mos*, *Ect'il'mos*, *Til'mos*. The action of plucking out; forcible extraction. (F.) *Arrachement*. This word is oftener used by the French than by the English surgeons, for the act of extracting certain parts, the presence of which is injurious,—as the teeth, &c.

EW-GOWAN, Bellis.

EX MARIBUS, Castratus.

EXACERBATIO, Exacerbation, Paroxysm.

EXACERBATION, *Exacerba'tio*, *Exaspera'tio*, *Exrescen'tia*, from *exacerbare*, *exacerbatum*, (ex, and *acerbus*), 'to provoke.' (F.) *Rédoublement*. An increase in the symptoms of a disorder. Often used synonymously with paroxysm.

EXACUM SPICATUM, *Coutouba alba*.

EXEMATOSIS, *Hæmatosis*.

EXÆMIA, Anæmia.

EXÆMON, Exanguis.

EXÆRESIS, from *ἐξ*, 'out of,' and *αἶμα*, 'I take away.' A surgical operation, which consists in drawing, extracting, or removing, from the human body, every thing that is useless, noxious, or extraneous. It is a generic term, which includes extraction, evulsion, evacuation, excision, ablation, amputation, &c.

EXÆRESIS, Extraction.

EXAGGERATED, *Exaggeratus*, (F.) *Exagéré*, from *exaggero*, (ex, and *aggero*, 'I heap up,') 'I magnify.' Heightened. Increased by expression. An epithet for sounds heard on auscultation and percussion, when much increased—*Hypereche'sis*, *Hypereche'ma*.

EXAGION. The sixth part of an ounce:—four scruples.—Actuarius.

EXALMA, from *ἐξ*, 'out of,' and *αλλομαι*, 'I leap.' *Ecalis*. Hippocrates calls thus the displacement of the vertebræ.

EXALSIS, Exalma, Leap.

EXALTATION OF THE VITAL FORCES, *Exaltatio vi'rium*, from *exalto*, (ex, and *alto*,) 'I raise.' This expression has been used, by modern pathologists, to designate a morbid increase in the action of organs, and partly that which occurs in an inflamed organ. Some use *exaltation of the vital forces*, and *inflammation*, synonymously.

EXAMBLOMA, Abortion.

EXAMBLOSIS, Abortion.

EXAMEN RIGOROSUM. 'Rigorous examination.' An examination of a Candidate for the degree of Doctor of Medicine in the Prussian Universities, which, like the *Tentamen medicum*, is conducted in Latin, and takes place before the medical faculty on all branches of medicine.

EXANASTOMOSIS, Anastomosis.

EXANASTROPHE, Convalescence.

EXANGIA, from *ἐξαγγίζω*, (*ἐξ*, and *αγγος*, 'a vessel,') 'I evacuate from a vessel.' *Exangei'a*. An enlargement or rupture of a blood-vessel, without external opening. A genus in the order *Dystheticæ*, class *Hæmaticæ* of Good. It comprises *aneurism* and *varix*.

EXANGIA ANEURISMA, Aneurism.

EXANGUIOUS, *Exsangui'ne*, *Exsanguineus*, *Exæ'mos*, *Exsang'uis*, (F.) *Exsangu*, from *ex*, 'out of,' and *sanguis*, 'blood.' Seeming to be bloodless; as a female, who has suffered largely from uterine hemorrhage. See *Anæmic*.

EXANIA, Protocele.

EXANIMATION, *Exanimatio*, from *ex*, 'out of,' and *anima*, 'the spirit.' This word has two acceptations. Sometimes, it means real death, corresponding with the Greek *Savatos*, *mors*. At others, it signifies apparent death, corresponding with the Greek *αἰψυχία*, *εκψυχία*, *εκλσις*, *Ant'imi dell'quim*.

EXANTHEM, *Exanthē'ma*, *Anthē'ma eruptio*, *Epanthē'ma*, *Efflorescent'ia*, *Exanth'is'ma*, *Ethyl'ma*, *Epanaste'ma*, *Epanthē'ma*, *Epanthis'ma*, *Effloratio*, from *ἐξανθεω*, (*ἐξ*, and *ανθεω*,) 'I flourish.'

A rash. (F.) *Exanthème*, *Élevure*. Under this term, is comprehended, by some, every kind of eruption, of which the skin is the seat. Others comprehend by it those eruptions that are accompanied by fever, *Febres exanthemat'icæ*: including, under the head of the *major exanthemata*, those which attack a person but once in his life, and which are communicated by specific contagion; and, under the *minor exanthemata*, those which are not marked by these characteristics. Small-pox, measles, cow-pox, &c., belong to the major:—chicken-pox, herpes, lichen, &c., to the minor. The general acceptation of Exanthem is, however, a more or less vivid, circumscribed, or diffuse redness of the skin, which diminishes, or

disappears transiently, under the pressure of the finger.

EXANTHEM, CARBUNCULAR, Anthracia—e. Ichorous, Emphyysis—e. Pustulous, Empyysis—e. Rash, Eanthesis.

EXANTHEMA, Eminence—e. Internum, Eisanthema.

EXANTHĒMA IUP'ICUM. An eruption of dark-red definite spots, of various sizes, spreading over the whole body, without the formation of scales, and disappearing only after a long time, which seems to be produced occasionally by the use of iodine.

EXANTHEMA MILIARIA, Miliary fever—e. Pestis, Plague—e. Antivariololum, *Vaccina*—e. Serosum, Pemphigus—e. Strophulus, Strophulus—e. Urticatum, Urticaria—e. Vaccina, Vaccina—e. Varicella, Varicella.

EXANTHEMATICA. Same etymon. *Eruptive fevers*. Cutaneous eruptions, essentially accompanied with fever;—the third order in the class *Hæmaticæ* of Good.

EXANTHEMATIS'CHESIS, *Exanthematos'chesis*, from *ἐξανθημα*, 'exanthem,' and *ἰσχειν*, 'to withhold.' Suppression of a cutaneous eruption.

EXANTHEMATOLOG'Y, *Exanthematolog'ia*, from *ἐξανθημα*, 'exanthem,' and *λογος*, 'a discourse.' The doctrine of cutaneous eruptions.

EXANTHEMATOPHTHALMIA, *Ophthalm'ia exanthemat'ica*, from *ἐξανθημα*, 'exanthem,' and *οφθαλμία*, 'inflammation of the eye.' Ophthalmia in the course of, or succeeding to, a cutaneous eruption.

EXANTHEMATOUS, *Exanthemat'ous*, *Exanthemat'icus*, (F.) *Exanthématique*. Relating or appertaining to exanthem, as "an exanthematous eruption." (F.) *Éruption exanthématique*.

EXANTHÈME INTESTINALE, see Typhus.

EXANTHESIS, see Efflorescence—e. Arthrosis, Dengue—e. Roseola, Roseola—e. Rubella, Rubella.

EXANTHISMA, Exanthem.

EXANTHROPIA, Misanthropia.

EXAPSYS, Inflammation.

EXARAG'MA, from *ἐξαρασσω*, 'I tear away,' 'I break.' Collision, violent fracture or friction.

EXARCHIATER, *Exarchiat'ros*. Chief of the archiatri, or chief of physicians, a title, like that of *Comes Archiatro'rum*, given to the chief physician of an emperor or king. Archiater appears to have meant, at times, the same thing.

EXARMA, Swelling.

EXAR'SIO. A burning heat.

EXARTEMA, Amuletum.

EXARTERITIS, *Exarterit'is*, from *ex*, 'out of,' and *arteria*, 'an artery.' Inflammation of the outer coat of an artery.

EXARTHREMA, Luxation, Sprain.

EXARTHROMA, Luxation.

EXARTHROSIS, Luxatio, Sprain—e. Paretica, see Pareticus.

EXARTICULATIO, Amputation, joint; Luxatio.

EXARYSIS, Exhaustion.

EXASPERATIO, Exacerbation.

EXCÆCARIA AGALLOCHA, Agallochum.

EXCALEFACTIO, Échauffement.

EXCARNA'TION, *Excarna'tio*, from *ex*, and *caro*, *carnis*, 'flesh.' A mode of making anatomical preparations, which consists in separating injected vessels from the parts in which they are situated. This is done by means of corrosion by an acid or by putrefaction.

EXCASTRATIO, Castration, Eunuch.

EXCATHISMA, Bath, half; Semicupium.

EXCAVATOR, Dental, see Dental.

EXCEREBRATIO, Eccephalosis.

EXCEREBRATUS, Delirious.

EXCERNENT, Secreting.

EXCIDENTIA, Ecliptoma.

EXCIPIENT, *Excipiens*, from *excipere*, (ex, and *capere*,) 'to receive.' (F.) *Intermède*. A substance, which, in a medicinal prescription, gives form and consistence to it, and serves as a *vehicle* or *medium* for the exhibition of the other ingredients.

EXCIPULA, Cup.

EXCISIO, *Entaille*.

EXCIS'ION, *Excisio*, from *excidere*, *excisum*, (ex, and *cadere*,) 'to cut off.' *Ec'tomè*. A surgical operation, by which parts of a small size are removed with a cutting instrument.

EXCITABIL'ITY *Excitabil'itas*. Irritability. From *excitare*, (ex, and *citare*, *citatum*,) 'to excite.' The faculty, possessed by living beings, of being sensible to the action of excitants. The doctrine of excitability forms an important part of the Brunonian system.

EXCITANT, Stimulant.

EXCITA'TION, *Excitatio*, *Excite'ment*; same etymon. The act of exciting; the state of an organ or organs excited. Excitement is, sometimes, used synonymously with augmented arterial action. The effect of the exciting powers acting on the excitability, according to Brown, constitutes *excitement*. Cullen used the term to express the restoration of the energy and action of the brain, which had been interrupted by sleep or some debilitating cause,—a state opposite to that of *collapse*. Not unfrequently, it is employed in the sense of excessive action,—*Super-excitatio*, (F.) *Sur-excitation*.

EXCITED DISEASES, Feigned diseases.

EXCITO-MOTION, see *Excito-motory*.

EXCITO-MO'TORY. (F.) *Excito-moteur*. An epithet applied by Dr. Marshall Hall to a division of the nervous system—comprising the gray matter of the spinal marrow, with the afferent and efferent nerves connected with it;—all of which are concerned in *reflex* actions; or those by which impressions are transmitted to a centre, and reflected so as to produce muscular contraction without sensation or volition. They constitute the *reflex system of nerves*. See *Nerves*.

The term *excito-motion* has also been employed to signify motion, no matter how excited, by the reflex nerves or by volition.—C. J. B. Williams.

EXCORIATIO, *Ecdora*.

EXCORIA'TION, *Excoriatio*, *Excoriatio'ra*, *Am'yché*, from ex, and *corium*, 'skin.' (F.) *Écorchure*. A slight wound, which removes only the skin.

EXCORIATURA, Excoriation.

EXCREA'TION, *Excreatio*, *Exscreatio*, *Serectus*, *Rasca'tio*, from ex, and *screare*, 'to spit.' Act of spitting. (F.) *Crachement*. See *Exspuitio*.

EXCREMENT, *Excremen'tum*, *Retrimen'tum*, *Excretum*, *Excretio*, *Perit'oma*, *Diachore'ma*, *Aph'odos*, *Aphoden'ma*, *Apoch'reon*, *Apoc'risis*, *Ar'das*, *Ar'dalos*, from *excernere*, *excretum*, (ex, and *cernere*,) 'to separate,' 'cleanse.' Everything, which is evacuated from the body of an animal by the natural emunctories, as superfluous; such as the faecal matters, the urine, perspiration, nasal mucus, &c. Generally, however, the term is restricted to the faecal evacuations;—*Purgamenta*, *Hedra*, *Sedes*, *Faeces*, *Stercus*, *Cacæ*, *Spat'ile*, (especially when liquid,) *Dejectio alvi'na* *O'nera alvi*, *Sordes ventris*, *Hypocho're'ma*, *Cæ'lia*, *Hypocho're'is*, *Merda*, *Merdus*, *Catarrhex'is*, *Copros*, *Scor*.

EXCREMENT, HUMAN, *Stercus humanum*.

EXCREMENTIT'IAL, *Excrementit'ius*, *Excrementit'ius*, (F.) *Excrémentieux*, *Excrémentiel*.

That which is similar to excrement, and forms part of it. *Excrementitial humours* or *parts* are those destined to be evacuated as incapable of administering to the nutrition of the body.

EXCREMENTO-RECREMENTIT'IAL, *Excremento-recrementit'ius*. Animal fluids, intended to be partly absorbed and partly rejected.

EXCRESCENCE, *Exerescen'tia*, *Exphy'ma*, *Phymatosis*, *Hypersarcosis*, *Sarcophy'ia*, *Carc'eres'cens*, from *exerescere*, (ex, and *erescere*,) 'to grow outwards.' (Sc.) *Su'alme*. (F.) *Excroissance*. A tumour, which forms at the surface of organs, and especially on the skin, mucous membranes, or ulcerated surfaces. Excrecences differ in size, shape, cause, &c., and each requires its own treatment. *Warts*, *condylomata*, *polyp'ii*, *hemorrhoids*, belong to this head.

EXCRESCENTIA, Exacerbation, Protuberance, Tumour—e. *Carnosa*, *Sarcoma*—e. *Fungosa*, *Fungosity*—e. *Gingivæ*, *Epulis*—e. *Vesicæ urina'riæ carnosæ*, *Cysthypersarcosis*.

EXCRETA, see *Excretion*.

EXCRÉTEUR, Excretory.

EXCRETIO, *Excretion*, *Excrement*—e. *Alvina*, *Defecation*—e. *Fæcum alvinarum*, *Defecation*—e. *Urinæ involuntaria*, *Enuresis*.

EXCRETION, *Excretio*, *Ec'crisis*, *Evacuatio*, *Ejectio*, *Eapul'sio*, *Eges'tio*, *Diachore'sis*, from *excernere*, *excretum*, (ex, and *cernere*,) 'to separate.' The separation or throwing off of those matters, *Excret'a*, *Eges'ta*, *Ion'ta*, *Apion'ta*, from the body of an animal, which are supposed to be useless, as the urine, perspiration, and faeces.

EXCRÉTOIRE, Excretory.

EXCRÉTO'RY, *Excretor'ius*, (F.) *Excréteur* ou *Excrétoire*; same etymon. An *Excretory vessel* or *duct*, *Ductus excretor'ius*, is one which transmits the fluid secreted by a gland either externally or into the reservoirs into which it has to be deposited. The existence of an excretory duct was regarded as a distinctive character of the glands properly so called.

EXCRETORY ORGAN means any one charged with the office of excreting; thus, the skin is said to be an excretory organ, because through it the perspiration takes place.

EXCRETUM, Excrement.

EXCROISSANCE, Excrecence.

EXCUTIA VENTRIC'ULI, *Stomach Brush*. An instrument, composed of iron or brass wire, at one of the extremities of which is a pencil of bristles. Some ancient authors proposed this to extract foreign bodies from the œsophagus, as well as to cleanse the stomach of viscid and tenacious matters adhering to it.

EXECHEBRONCHUS, *Bronchocele*.

EXECHEGLUTI, *Exischio*.

EXEDENS, see *Herpes exedens*.

EXELCOSIS, see *Ulceration*.

EXELCYS'MOS, from ελ, 'from,' and εκωω, 'I draw.' *Extraction*. Also the act of breaking out into ulcers.

EXELCYSMUS, *Extraction*.

EXENCEPHALE, from ελ, 'out of,' and εγκεφαλον, 'the encephalon.' A genus of monsters, in which the encephalon is situate in a great measure out of the cranium, the upper portion of which is almost wholly wanting. *Isid. G. St. Hilaire*.

EXENTERATION, *Exenterismus*.

EXENTERIS'MUS, *Exenter'isus*, *Exenteratio*, *Exenteratio*, *Evisceratio*, *Devisceratio*, *Unbow'elling*, *Embow'elling* or *Imbow'elling*, *Visceratio*, from ελ, 'out of,' and εντερον, 'an intestine.' The operation of taking out the bowels.

EXERA'MA, from εξεραω, 'I throw out.' Any thing cast out. Vomiting; or the matter vomited.—*Hippocrates*.

EXERCISE, *Exercita'tio, Exercit'ium, Asce'sis, Gymna'sion*, from *exercere, exercitum*, 'to work.' Every motion of the body arising from the contraction of muscles subjected to the will. Also, the action of any organ whatever. Exercise may be *active* or *passive*. The passive are referred to, under the head of Gestation. The chief active exercises are:—walking, running, dancing, hunting, fencing, playing at ball, cricket, racket, quoits, swimming, declamation, and singing. Exercise is an important prophylactic, particularly for those disposed to be plethoric. It improves the digestion; augments the secretions; and, when used in moderation, gives strength to the body; but when carried to excess, produces debility and disease.

EXERRHO'SIS, from *ἐξ*, 'out of,' and *ῥεω*, 'I flow.' The discharge which takes place by insensible perspiration.

EXFETATION, see Pregnancy.

EXFOLIATIF, *Exfoliative*.

EXFOLIATION, *Enfolia'tio, Desquama'tio, Eclep'isis, Eclep'isis*, from *ex*, and *folium*, 'a leaf.' By this is meant the separation of the dead portions of a bone, tendon, aponeurosis, or cartilage, under the form of lamellæ or small scales. Exfoliation is accomplished by the instinctive action of the parts, and its object is to detach the dead portion from those subjacent, which are still alive. For this purpose the latter throw out fleshy granulations, and a more or less abundant suppuration occurs, which tends to separate the exfoliated part,—now become an extraneous body. The ancients distinguished exfoliation into *sensible* and *insensible*, according as the dead portions of bone were detached in fragments of greater or less size, or in very thin pieces, and in an almost insensible manner. When the dead part embraces all or almost all the substance of a bone, it takes the name *Sequestrum*.

EXFO'LIATIVE, *Exfoliativ'us, Desquamato'rius*, (F.) *Exfoliatif*. That which takes away by leaves or scales. The term has been applied to certain medicines, which were regarded as proper to hasten exfoliation, such as alcohol, oil of turpentine, tincture of myrrh, &c.

EXFO'LIATIVE TREPAN, *Eclep'isitrep'anum*. An ancient raspatory, or instrument for scraping exfoliating portions of bone.

EXHALATION, *Exhalation*.

EXHA'LANT, *Exha'lent, Exha'lans*, from *exhalare, exhalatum*, (*ex*, and *halare*, 'to breathe,') 'to exhale,' 'throw out.'

EXHALANT VESSELS, *Vasa exhalan'tia*, are very minute, and rise from the arterial capillary system. They are situate in every tissue of the body, and on the surface of the mucous and serous membranes and skin; on which each pours its particular fluid. Bichat distinguished three sets. 1. The *external*, terminating on the mucous and external dermoid system, where they pour the matter of perspiration. 2. The *internal*, comprising those of the areolar and medullary tissues, and of synovial surfaces; and, 3. The *nutritive exhalants*, which vary in each organ where they are found, and preside over the phenomena of composition and increase of every part of the body. The exhalants are the antagonists of the absorbents. They are imaginary vessels, inasmuch as they cannot be detected.

EXHALATIO, *Evaporation*.

EXHALATION, *Exhala'tio*; same etymon. *Anathymi'asis, Apopneus'is*, (F.) *Exhalaison*. A function, by virtue of which certain fluids, obtained from the blood, are spread, in the form of dew, in the areolæ of the different textures, or at the surface of membranes; either for the sake of being thrown out of the body, or to serve certain purposes. The sweat is a *liquid, excrementitious*

exhalation; the serous fluid of the pleura, a *liquid recrementitious exhalation*.

Exhalation is, also, applied to that which ex-hales from any body whatever, organic or inorganic, dead or living.

EXHALATION, PULMONARY, see Perspiration.

EXHAUSTION, *Exaur'ysis, Vires exhaust'a*, from *exaurire, exhaustum*, (*ex*, and *haurire*,) 'to draw out.' (F.) *Épuisement*. Loss of strength, occasioned by excessive evacuations, great fatigue or privation of food, or by disease.

EXHIL'ARANT, *Exhil'arans*, from *ex*, and *hilaro*, 'I make merry.' An agent that exhilarates or enlivens.

EXHUMA'TION, *Exhuma'tio, Effos'sio*, from *ex*, and *humus*, 'the ground.' The disinterment of a corpse. The circumstances which render this necessary are:—1. Judicial investigations relative to the body of the person inhumed. 2. The removal of a body from one cemetery to another; and, 3. The evacuation of cemeteries or sepulchral vaults. The operation is attended with much unpleasant smell and annoyance, and requires the use of disinfecting agents, of which the most powerful is chlorinated lime. See Disinfection. The putrid effluvia from animal substances are not, however, found to excite endemic disease.

EXIDIA AURICULA JUDÆ, *Peziza auricula*.

EXINANIT'ION, *Exinanit'io*, from *ex*, 'out of,' and *inanis*, 'empty.' Extreme exhaustion; complete evacuation.

EXISCH'TUM. Same etymon as the next. Prominence of the hips.

EXIS'CHIUUS, *Exis'chus*, from *ἐξ*, 'out of,' and *ἰσχίον*, 'the ischium.' A luxation of the os femoris. Those with large nates, and prominent hips, were formerly called *Exis'chioi* and *Exeche-gluti*.

EXITU'RA, from *exire, exitum*, (*ex*, and *eo*,) 'to go out.' According to some, any abscess which discharges. Paracelsus calls thus every kind of putrid excrement.

EXITUS. Same etymon. The outer termination or exit of a canal. The termination of a disease.

EXOARTERITIS, see Arteritis.

EXOCARDIAC, see Endocardiac.

EXOCARDIAL, see Endocardiac.

EXOCARDITIS, Pericarditis.

EX'OCHAS, *Ex'ochè*, from *ἐξέχω*, (*ἐξ*, and *ἐχω*,) 'I project.' A soft tumour—as a hemorrhoid—outside of the anus. An outward pile.

EXOCHÈ, Eminence, Exochas.

EXOCHORION, see Chorion.

EXOCOLITIS, see Colitis.

EXOCULA'TIO, from *ex*, 'out of,' and *oculus*, 'an eye.' Want of eyes. Want of vision. Blindness.

EXOCYSTÈ, *Exocys'tis*, from *ἐξ*, 'out of,' and *κύστις*, 'the bladder.' A prolapsus of the bladder into the urethra. Also called *Prolapsus vesicæ*, *Ædopto'sis vesicæ*, (F.) *Renversement de la vessie*.

EXOCYS'TE NOBILIA'NA. Protrusion of the inner membrane of the bladder. So called from M. Noel, who first accurately described it.

EXOCYS'TE SOLINGENIA'NA. Protrusion of the neck of the bladder. Called after M. Solingen, who first accurately described it.

EXCEDESIS, Swelling.

EXODIC, Eferent.

EXOG'GENOUS, from *ἐξ*, 'out of,' and *γενναω*, 'I engender.' A term first applied to plants—hence called *Ex'ogens*—in which the wood increases by annual additions to the outside.

In animal anatomy, processes which shoot out from any part are termed *exogenous*. An *exoge-*

non aneurism is one caused by division, from without, of the arterial walls.

• **EXOGENS**, see *Exogenous*.

EXOLOGIE, Extraction.

EXOLOGNUM PURGA, *Convolvulus jalapa*.

EXOGONYAN'CON, from *ἐξω*, 'outwards,' *γωνία*, 'the knee,' and *ἄγκυρα*, 'an elbow.' Bowing of the knees outwards.

EXOINE, *Excuse*, from *ex*, 'out of,' and *idoneus*, 'fit,' or rather from *excusare*, 'to exonerate' [?]. In France, a certificate of excuse, exemption, or dispensation, given to those summoned to appear before a court of justice, and who are unable to do so.

EXOLCE, Extraction.

EXOLUTION, Syncope.

EXOMETRA, *Prolapsus uteri*.

EXOMOTIOSIS, Assimilation.

EXOMPHALOCLE, *Exomphalus*.

EXOM'PHALUS, *Exumbilical'itis*, *Exomphalocel'e*, *Om'phalocel'e*, *Hernia umbilical'is*, *Om'phalocel'ic*, *Om'phaloprop'itis*, *Prolapsus umbil'icis*, (F.) *Hernie ombilicale*, *H. du nombril*, *Umbilical hernia*, from *ἐξ*, 'out of,' and *ὀμφαλός*, 'the navel.' Hernia occurring at the navel. This affection happens more frequently in infants, and takes place by the umbilical ring. In adults, it occurs more commonly in females than in males; and, when it does so, the sac passes in the vicinity of the umbilicus. The organs, found in this kind of hernia, are particularly, — the epiploon, the jejunum, the arch of the colon, and sometimes the stomach. The tumour is, in general, round, and presents all the characters of hernia. It is, commonly, really reducible, and not subject to strangulation. It must be reduced, and retained by an elastic bandage, made in the form of a girdle, and furnished with a pad at its middle part. When strangulated, the stricture may be divided upwards and towards the left side.

EXONCOMA, *Exoncosis*, from *ἐξ*, and *ογκος*, 'a tumour.' A large, prominent tumour. Used, by Galen, for protuberance of the vertebrae after luxation.

EXONCOSIS, *Exoncoma* — e. *Linguae*, *Glossoneus*.

EXONEIROGMUS, Pollution, nocturnal.

EXONEIROSIS, Pollution, nocturnal.

EXONEURISM, Magnetism, animal.

EXOPHTHALMIA, *Ptoxis* seu *Prolapsus* seu *Procidencia Oculi* seu *Bulbi Oculi*, *Exorbitismus*, *Ophthalmocel'e*, *Ophthalmoprop'itis*, *Oculi totius prominencia*, from *ἐξ*, 'out of,' and *ὀφθαλμος*, 'eye.' (F.) *Providence de l'œil*. A protrusion of the eye from its orbit, occasioned by an abscess or tumour in the areolar texture of the orbit; by exostosis of the parietes of the orbit, &c. In exophthalmia, the eye is pressed forwards; the eyelids are raised and separated, so that they can no longer cover the eye and defend it from the action of extraneous bodies; it becomes inflamed, and the sight is disturbed or destroyed. The treatment of course depends upon the cause.

EXOPHTHALMIA FUNGOSA, *Sarcosis bulbi* — e. *Sarcomatica*, *Sarcosis bulbi*.

EXOPHTHALMUS, same etymon. One whose eyes are very prominent. The opposite to *Ceolophthalmus*.

EXORBITISMUS, *Exophthalmia*.

EXORMIA, *Papula* — e. *Lichen*, *Lichen* — e. *Prurigo*, *Prurigo* — e. *Strophulus*, *Strophulus*.

EXORTUS UNGUIUM, see *Nail*.

EXOSIS, Luxation.

EXO-SKELETON, see *Skeleton*.

EXOSMA, from *ἐξ*, 'out of,' and *σμά*, 'to move.' A luxated or dislocated limb or organ.

EXOSMIC, *Exosmotic*.

EXOSMOSE, *Exosmosis*, *Transudation*, from

ἐξ, 'out of,' and *σμος*, 'impulse.' The opposite to *Endosmose*. The act by which substances transude from within to without an animal or other membrane.

EXOSMOTIC, *Exosmot'icus*, *Exos'mic*, *Exos'micus*; same etymon. Belonging to *Exosmose*: — as an *exosmotic* current.

EXOSSATIO, *Exostosis*.

EXOSTEMMA CARIBÆA, *Cinchona caribæa* cortex.

EXOSTOMA, *Exostosis*.

EXOSTONE, *Exostosis* — e. *des Dents*, *Exostosis dentium* — e. *Sous-angéale*, see *Subungual*.

EXOSTOSIS, *Hyperostosis*, *Emphy'ma exostosis*, *Exosto'ma*, *Eposto'ma*, *Eposto'sis*, *Osteo'ma*, *Osto'ma*, *Osteoph'ytia*, *Exosena'tio*, *Ossis Eminen'tia*, (F.) *Exostose*, *Osteophyte*; from *ἐξ*, 'out of,' and *οστέον*, 'a bone.' An osseous tumour, which forms at the surface of bones, or in their cavities. Various kinds have been enumerated. *Ivory Exostosis*; (F.) *E. eburnée*; — that which has the appearance and consistence of ivory. *Laminar Exostosis*; (F.) *E. laminaire*; — that which is formed of laminae in superposition, or of distinct filaments. The *Spongy Exostosis* is that whose structure is analogous to the spongy tissue of bones. *Exostoses* are sometimes distinguished into the *true*, which seem to be a projection of the osseous substance, and which have the same organization and hardness as that substance; and the *false* or *osteosarcoma*. *Exostosis* may depend on syphilis, scrofula, rickets, gout, &c. In such cases, it is important to get rid of the primary disease.

Those *exostoses*, which occur within the bones, have, by some, been called *Enostoses*.

Exostosis, *Spina ventosa*.

EXOSTOSIS DENTIUM, (F.) *Exostose des Dents*. *Exostosis* of the teeth.

EXOSTOSIS STRATOMATODES, *Osteosteotoma* — e. *Subungual*, see *Subungual*.

EXOTERIC, see *Esoteric*.

EXOTIC, *Exoticus*, from *ἐξω*, 'without.' That which comes from abroad. Plants or drugs which are procured from abroad are so called. It is opposed to *indigenus*.

EXOTICADEN'IA, from *exotic*, and *αἰμα*, 'I dislike.' Aversion for exotic drugs.

EXOTICHEMATOSIS, Transfusion.

EXOTICOMANIA, The opposite to *exoticadenia*. Fondness for exotic remedies.

EXOTICOSYMPHYISIS, from *ἐξωτικός*, 'foreign,' and *συνφυσις*, 'a growing together.' A union or growing together of foreign bodies, as of a foreign body with the human.

EXPANSIO, *Expansion* — e. *Musculosa*, *Platysma myoides*.

EXPAN'SION, *Expan'sio*, from *expandere*, *expansum*, (ex, and *pandere*, 'to open,') 'to spread out.' A prolongation or spreading out, presented by certain organs. Thus, we say an *aponeurotic expansion*, &c.

EXPATRATIO, Ejaculation (of Sperm.)

EXPECTANT, *Expectatus*, from *expectare*, *expectatum*, (ex, and *spectare*, 'to look,') 'to wait.' That which waits: — as *Expectant Medicine*. — *La Médecine expectante*. See *Expectation*.

EXPECTATION, *Expecta'tio*; same etymon. The word *expectation* has been applied, in medicine, to that method, which consists in observing the progress of diseases, and removing deranging influences, without prescribing active medicines, unless such shall be imperiously required. It consists, in fact, in leaving the disease almost wholly to the efforts of nature, and has been termed the *art of curing diseases by expectation* or *waiting*. — *Ars sanandi cum expecta'tione*.

EXPECTORANT, *Expect'orative*, *Expect'orans*, *Anacathar'ticus*, *Sputato'rius*, *Antide'chicus*,

Phys' magogue, from *ex*, 'out of,' and *pectus*, *pectoris*, 'the breast.' A medicine capable of facilitating or provoking expectoration. There is probably no such thing as a *direct* expectorant. They all act through the system, or by impressions made on parts at a distance, which, through the medium of general, continuous, or contiguous sympathy, excite the secretory vessels of the air-passages into action. The following are the chief reputed expectorants:—*Ammoniacum*; *Asafoetida*; *Galbanum*; *Ipecacuanha*; *Myroxylon*; *Myrrha*; *Inhalations* of *Iodine*, *Stramonium*, *Tar*, *Burning Wool*, *Tobacco*, &c.; *Scilla*; *Senega*, and *Toluianum*.

EXPECTORATIO, Expectation — e. *Sanguinis*, *Hæmoptysis* — e. *Solida*, *Cynanche trachealis*.

EXPECTORATION, *Expectatio*, *Ecchelysis*, *Ex h'umida*, *Anaptyxis*, *Prop'ptysis*, *Stathocathar'sis*, *Anacathar'sis*, *Anabes'sis*, *Raising*. Same etymon. The act of expelling from the chest matters or secretions there collected or existing. It is, likewise, used for the expectorated matter.

EXPECTORATION DE SANG, *Hæmoptysis*.

EXPECTORATION, PRUNE-JUICE, see *Prune-juice*.

EXPECTORATIVE, Expectorant.

EXPELLENS, Expulsive.

EXPERIENCE, *Experientia*, *Peira*, *Empeira*, from *εξ*, and *πειρα*, 'a trial.' A knowledge of things acquired by observation. In medicine, this knowledge can be obtained both by the practitioner's own experience, and by that obtained from tradition and from books. To profit by experience requires a mind capable of appreciating the proper relations between cause and effect; and hence it happens, that *false experience*, *Experientia fallax*, is extremely common; and that a man had better, in many instances, trust to that which he has learned from others, than to his own fallacious observation.

The union of accurate observation by the physician with that handed down by medical writers constitutes perfect experience, so far as it is attainable in any individual case.

EXPÉRIENCE, Experiment — e. *de Mariotte*, *Mariotte*, experiment of.

EXPERIMENT, *Experimentum*; same etymon. (F.) *Expérience*. A trial, made on the bodies of men or animals, for the purpose of detecting the effect of a remedy, or of becoming better acquainted with their structure, functions, or peculiarities. In a more general sense, it means any trial instituted with the intent of becoming better acquainted with any thing. By experiments on living animals, we have obtained much valuable information in the various departments of medicine; but particularly in physiology and toxicology.

EXPERIMENT OF MARIOTTE, see *Mariotte*.

EXPERIMENT OF WEPFER, *Experimentum Wepferianum*. An experiment attributed to *Wepfer*, which consisted in injecting air into the blood-vessels of animals, with the view of observing its effects on the economy.

EXPERTS NUPTIARUM, *Virgin*.

EXPERT (F.), *Expertus*, from *ex*, and *peritus*, 'skilled.' Skillful or of good experience. A physician, charged with the duty of making a report, (F.) *Expertine*, upon any case of legal medicine.

EXPERTINE, see *Expert*.

EXPIRATION, *Expiratio*, *Espiratio*, *Ec'pnoë*, *Epepneumatō'sis*, *Epepneū'sis*, *Apopneū'sis*, *Apop'noë*, *Apopneū'a*, from *exspirare*, (*ex*, and *spirare*, *spiratum*), 'to breathe out.' The act of expelling from the chest, the air received during inspiration.

EXPIRATORY, *Expiratio'ni inser'viens*. Same etymon. Relating or appertaining to expiration. The expiratory muscles are all those which contribute to diminish the cavity of the chest, for the purpose of expelling the air contained in the lungs, or of producing expiration. These muscles are, chiefly, the intercostals, triangularis sterni, quadratus lumborum, serratus posticus inferior, the oblique and recti muscles of the abdomen, and the sacro-lumbalis.

EXPLORATIO, Exploration — e. *Abdominis*, *Abdominoscopia*.

EXPLORATION, *Exploratio*, *Recognit'io*, from *explorare*, *exploratum*, 'to search into.' The act of observing and attentively examining or investigating every thing connected with a case of disease. The word is chiefly used in this sense by the French practitioners.

EXPLORATOR, CHEST. An instrument, proposed by *Dr. B. Babington* for exploring the chest in cases of empyema. It consists of a needle, contained in the smallest sized canula. This is passed between the ribs into the chest. The needle is then withdrawn, and the escape of fluid indicates the nature of the case.

EXPLORATORIUM, Sound, Specillum.

EXPRES'SION, *Expres'sio*, *Ecpies'mos*, *Echthip'sis*, from *ex*, 'out of,' and *premere*, *pressum*, 'to press.' The act of compressing a substance, for the purpose of separating from it the fluids which it contains. Also, the manner in which impressions made upon us are depieted; especially in the traits of the countenance.

EXPRESSION OF SWEAT, (F.) *Sueur d'expression*, is a term given to the passive perspiration observable in very debilitated individuals.

EXPUITION, Exspuition.

EXPULSION, Excretion — e. *Fæcum*, *Defecation* — e. *Fœtus*, *Parturition*.

EXPULSIF, Expulsive.

EXPULSION, SPONTANEOUS, *Evolution*, spontaneous.

EXPULSIVE, *Expellens*, *Expulso'rius*, (F.) *Expulsif*, from *expellere*, *expulsum*, (*ex*, and *pellere*), 'to drive away.'

An *expulsive bandage*, (F.) *Bandage expulsif*, is one constructed with the view of compressing a part, from which we are desirous of expelling pus, serum, &c. Certain medicines were formerly called *expulsiues*, which were believed to have the power of driving the humours towards the skin; — as diaphoretics, and sudorifics.

EXPULTRIX, see *Vis expultrix*.

EXSANGUE, Exanguious.

EXSANGUINE, Exanguious.

EXSANGUINITY, *Anæmia*.

EXSANGUIS, Exanguious. *Exsanguis* is used by *Ausonius* for one exhausted by venery; — as *sanguis* meant sperm as well as blood. See *Sperm*.

EXSARCOMA, *Sarcoma*.

EXSCREATIO, Excretion.

EXSECTIO VIRILIIUM, *Castration*.

EXSECTUS, *Castratus*, *Eunuch*.

EXSICCATIO, *Drying*.

EXSICCATIVA, *Desiccativa*.

EXSOLUTIO, *Eclysis*.

EXSPIRATIO, *Exspiration*.

EXSPUIT'ION, *Exspuit'io*, *Spuil'io*, *Sputatio*, *Anachremp'sis*, *Apochemp'sis*, *Chremp'sis*, *Pty'sis*, *Anacine'ma*, *Anacine'sis*, (F.) *Exspuition*, *Sputation*, from *ex*, 'out of,' and *spuo*, 'I spit.' Rejection of the matters accumulated in the pharynx and larynx; spitting.

EXSTASIS, *Eccstasis*.

EXSTIRPATIO, *Extirpation*.

EXSTROPH'IA, *Ex'strophē*, *Ec'estrophē*, from *εξ*, 'out of,' and *στροφή*, 'turning.' *Extrover'sio*, *Extrover'sion*. Eversion or turning out of a part

-as of the eyelids. A term used by M. Chausier for certain displacements of organs, and especially of the urinary bladder.

EXSTROPH'IA or EX'STROPHY OF THE BLADDER. (F.) *Renversement de la Vessie*, is a faulty conformation, in which the organ opens above the pubes; so that in the hypogastric region there is a red, mucous surface, formed by the inner coat of the bladder; on which two prominences are distinguishable, corresponding to the openings of the ureters.

EXSUCCATIO. Ecchymoma.

EXSUC'TIO, *Suc'tio*, *Ecmyze'sis*, from *ex*, and *sugere*, *nectum*, 'to suck.' The action of sucking.

EXSUDATIO. Ephidrosis.

EXSUFFLATIO, Ecphysexis.

EXTA, see Entrails.

EXTASE. Ecstasis.

EXTEMPORA'NEOUS, *Extempora'neus*, *Extemporalis*, (F.) *Extemporane*, from *ex*, and *tempus*, *temporis*, 'out of time.' Those prescriptions are called 'extemporaneous,' or 'magistral,' which are made on the spot, and composed according to the prescription of the physician.

EXTENSEUR COMMUN DES DOIGTS, Extensor digitorum communis — *e. Commun desorteils*, Extensor communis digitorum pedis — *e. Court du ponce*, Extensor pollicis brevis — *e. Long du ponce*, Extensor longus pollicis — *e. Petit desorteils*, Extensor brevis digitorum pedis — *e. Propre du petit doigt*, Extensor proprius minimi digiti — *e. Propre de l'Index*, Extensor proprius indicis.

EXTENSIBILITY, *Extensibil'itas*. A property, possessed by certain bodies, of being capable of extension or elongation.

EXTENSIO, Extension, Tetanus.

EXTEN'SION, *Extens'io*, *Tasis*, *Et'asis*, *Cat'asis*, *Anat'asis*, from *extendere*, (*ex*, and *tendere*, *tensum*), 'to stretch out.' An operation in surgery, in which either with the hands alone, or by straps, a fractured or luxated limb is pulled strongly, to restore it to its natural position. It is the opposite of Counter-extension.

EXTENSOR. Same etymon. (F.) *Extenseur*. A muscle, whose office is to extend certain parts.

EXTENSOR BREVIS DIGITO'RUM PEDIS, *Ped'icus*, (F.) *Calcaneo-phalanginien commun*, of DUMAS; *Muscle pédieux ou petit extenseur desorteils*. *Calcaneo-sus-phalangien commun* — (Ch.), *Carré du pied*; *Short Extensor of the Toes*. A muscle, situate on the dorsal region of the foot. It arises from the external surface of the calcaneum, and at the anterior edge of a ligament, which unites that bone to the astragalus. Anteriorly, each of its divisions terminates by a small tendon, which is fixed successively, at the superior part of the posterior extremity of the first phalanx of the great toe, and to the second and last phalanges of the next three toes. Its use is to extend the first four toes, and to direct them a little outwards.

EXTENSOR CARPI RADIALIS BREVIOR, see Radialis — *e. Carpi radialis longior*, see Radialis — *e. Carpi ulnaris*, see Cubital muscles.

EXTENSOR COMMUNIS DIGITO'RUM PEDIS, *E. Longus Digitorum Pedis*, *E. Digitorum longus*, *Cnemodactyla'us*, (F.) *Péronéo-tibi-sus-phalangien commun*, *Péronéo-sus-phalangien commun*, *Extenseur commun des Orteils*. This muscle, also, is situate at the anterior part of the leg. It is long, thin, flattened, simple, and fleshy above, and divided into four tendons below. It arises from the outer tuberosity of the tibia and the anterior surface of the fibula, and is inserted into the superior part of the posterior extremity of the second and third phalanges of the last four toes. It extends the three phalanges of these toes.

EXTENSOR DIGITO'RUM COMMUNIS, *E. digito-*

rum communis manus cum extenso're proprio auricularis, *Digitorum Tensor*; (F.) *Épicondylo-sus-phalangien commun* — (Ch.) — *Extenseur commun des Doigts*, *Common Extensor of the Fingers*. A long, flattened muscle; simple above, and divided into four portions inferiorly. It is situate at the posterior part of the forearm; arises from the external tuberosity of the humerus; from the aponeurosis of the forearm, and from the aponeurotic septa situate between it and the neighbouring muscles, and is inserted at the posterior surface of the second and third phalanges of the last four fingers. This muscle extends the phalanges of the last four fingers upon each other, and upon the metacarpal bone. It can, also, extend the hand on the forearm.

EXTENSOR DIGITORUM LONGUS, *E. communis digitorum pedis* — *e. Longus digitorum pedis*, *E. communis digitorum pedis* — *e. Proprius hallucis*, *E. proprius pollicis pedis* — *e. Indicis*, *E. proprius indicis* — *e. Internodii ossis pollicis*, *E. longus pollicis* — *e. Primi internodii*, *Abductor longus pollicis*.

EXTENSOR POL'ICIS BREVIS, *E. minor pollicis manus seu primi internodii seu pollicis primus seu secundus internodii* (DOUGLAS.) *seu secundus internodii ossis pollicis*; (F.) *Cubito-sus-phalangien du ponce*, — (Ch.), *Court extenseur du ponce*. Seated at the posterior and inferior part of the forearm. It is thin, long, and broader at its middle than at the extremities. It arises from the posterior surface of the ulna and the interosseous ligament, and is inserted behind the superior extremity of the first phalanx of the thumb. It extends the thumb, and aids in supination.

EXTENSOR POL'ICIS LONGUS; *E. major pollicis manus seu secundus internodii seu pollicis secundus seu tertius internodii* (DOUGLAS.) *seu internodii ossis pollicis*, (F.) *Cubito-sus-phalangien du ponce*, — (Ch.), *Muscle long extenseur du ponce*. This muscle is long, flat, and fusiform; and is seated at the posterior part of the forearm. It arises from the posterior surface of the ulna and the interosseous ligament, and is inserted at the posterior part of the superior extremity of the first phalanx of the thumb. It extends the last phalanx of the thumb upon the first.

EXTENSOR PRO'PRIUS INDICIS, *E. proprius primi dig'iti manus*, *E. Indicis*, *Indicator seu secundus internodii indicis proprius*, (F.) *Cubito-sus-phalangien de l'Index*, — (Ch.), *Extenseur propre de l'Index*. This muscle is long and thin; broader in the middle than at the extremities; and is situate at the posterior part of the forearm. It arises from the posterior surface of the ulna, and is inserted at the posterior part of the upper extremity of the second and third phalanges of the index-finger. It extends the three phalanges of the index-finger; and has, besides, the same uses as the other extremities of the fingers.

EXTENSOR PRO'PRIUS MINIMI DIG'ITI, (F.) *Épicondylo-sus-phalangien du petit doigt*, — (Ch.), *Extenseur propre du petit doigt*. Situate on the inside of the *Extensor communis digitorum*. It arises from the external condyle of the os humeri and the aponeurotic septa seated between it, the extensor communis digitorum, and the extensor carpi ulnaris; and is inserted into the last two phalanges of the little finger. Its use is to extend the little finger, and even the hand upon the forearm.

EXTENSOR PRO'PRIUS POL'ICIS PEDIS, *E. proprius Hallucis seu Longus* (DOUGLAS.) *seu pollicis longus*, *Péronéo-sus-phalangien du ponce*, *Péronéo-sus-phalangien du ponce*, — (Ch.) This muscle is situate at the anterior part of the leg. It is

fleshy, broad, flat above; small and tendinous below. It arises from the anterior part of the middle third of the fibula, and is inserted into the posterior part of the superior extremity of the last phalanx of the great toe. It extends the last phalanx of the great toe upon the first, and the first upon the first metacarpal bone.

EXTENSOR PRIMI INTERODII, *E. pollicis brevis*—*e. Secundi internodii*, *E. pollicis brevis*, *Extensor longus pollicis*—*e. Secundi internodii indicis proprius*, *E. proprius indicis*—*e. Tertii internodii*, *E. longus pollicis*—*e. Tertii internodii indicis*, *Prior annularis*—*e. Tertii internodii minimi digiti*, *Abductor minimi digiti*—*e. Ossis metacarpi pollicis manus*, *Abductor longus pollicis*—*e. Pollicis longus*, *E. proprius pollicis pedis*—*e. Pollicis secundus*, *E. longus pollicis*—*e. Minor pollicis manus*, *E. pollicis brevis*—*e. Primus pollicis*, *Abductor longus pollicis*.

EXTENSOR TARSII MAGNUS, *E. Tarsi suralis*. A name given, by some anatomists, to the gastrocnemius and soleus combined.

EXTENSOR TARSII MINOR, *Plantar muscle*.

EXTENUATIO, Emaciation.

EXTERGENATIO, Detergents.

EXTERNAL DISEASES, *Extrinsic Diseases*, *Morbi extrinseci*, *Surgical diseases*, (*F.*) *Maladies externes*. Those diseases which occupy the surface of the body, and form the object of surgical pathology, requiring, generally, external means, or surgical operations.

EXTERNAT. The post or office of an *externe*.

EXTERNE, see *House-surgeon*.

EXTERNUS AURIS, *Laxator tympani*—*e. Mallei*, *Laxator tympani*.

EXTESTICULATUS, *Castratus*, *Eunuch*.

EXTINCTIO, *Death*—*e. Hydrargyri*, *Extinction of Mercury*—*e. Mercurii*, *Extinction of Mercury*.

EXTINCTIO VOICIS, (*F.*) *Extinction de voix*. The French use this term for cases in which the voice is not wholly suppressed, but produces only feeble sounds:—*Incomplete aphonia*.

EXTINCTION OF MERCURY, *Extinctio Mercurii* *sen Hydrargyri*. Trituration of mercury with lard or other substance, until the metallic globules disappear. The mercury is then so divided, that it forms a black powder, generally considered to be a protoxide of mercury, but, perhaps, erroneously.

EXTIRPATIO, *Extirpation*—*e. Linguae*, *Glossosteresis*—*e. Testiculorum*, *Castratio*.

EXTIRPATION, *Extirpatio*, *Exstirpatio*, *Ablatio*, *Apharesis*, from *extirpare*, (*ex*, and *stirpe*, 'a root,') 'to root out.' The complete removal or destruction of any part, either by cutting instruments or the action of caustics. Thus, we speak of the extirpation of cancer, polypus, encysted tumour, &c.

EXTOMIUS, *Eunuch*.

EXTOZOA, *Ectozoa*.

EXTOZOAIRES, *Ectozoa*.

EXTOZOARIA, *Ectozoa*.

EXTRA-PELVIO-PUBI-TROCHANTÉRIEN. *Obturator externus*.

EXTRACT, *Extractum*—*e. of Aconite*, *Extractum aconiti*—*e. of Aconite*, *alcoholic*, *Extractum aconiti alcoholicum*—*e. Alcoholic*, see *Extractum*—*e. of Aloes*, *Extractum aloes*—*e. of Bark*, *Extractum cinchonæ*—*e. of Bark*, *resinous*, *Extractum cinchonæ resinosum*—*e. of Belladonna*, *Extractum belladonnæ*—*e. of Belladonna*, *alcoholic*, *Extractum belladonnæ alcoholicum*—*e. of Bittersweet*, *Extractum dulcamaræ*—*e. of Broom-tops*, *Extractum cacinum genistæ*—*e. of Butternut*, *Extractum juglandis*—*e. of Cascarrilla*, *resinous*, *Extractum cascarrillæ resinosum*—*e. of Chamomile*, *E. anthemidis*—*e. of Cinchona*, *Extractum cinchonæ*—*e. of Colchicum*, *Extract*

of Colchicum—*e. of Colchicum*, *acetic*, *Extractum colchici aceticum*—*e. of Colocynth*, *Extractum colocynthidis*—*e. of Colocynth*, *compound*, *Extractum colocynthidis compositum*—*e. of Cubebs*, *fluid*, *Extractum cubebæ fluidum*—*e. of Dandelion*, *Extractum taraxaci*—*e. of Elaterium*, *Extractum elaterii*—*e. of Ergot*, *Ergotin*—*e. Ethereal*, see *Extractum*—*e. of Foxglove*, *Extractum digitalis*—*e. of Gentian*, *Extractum gentianæ*—*e. of Hemlock*, *Extractum cicuta*—*e. of Hellebore*, *black*, *Extractum hellebori*—*e. of Hemlock*, *alcoholic*, *Extractum conii alcoholicum*—*e. of Henbane*, *Extractum hyoscyami*—*e. of Henbane*, *alcoholic*, *Extractum hyoscyami alcoholicum*—*e. of Hops*, *Extractum humuli*, *E. lupuli*—*e. of Jalap*, *Extractum jalapæ*—*e. of Lettuce*, *Extractum lactucæ*—*e. of Liquorice*, *Extractum glycyrrhizæ*—*e. of Logwood*, *Extractum hæmatoxyli*—*e. of Mayapple*, *Extractum podophylli*—*e. of Meat*, *saponaceous*, *Osmazome*—*e. of Nux Vomica*, *Extractum nucis vomicæ*—*e. of Oak-bark*, *Extractum corticis quercus*—*e. of Pareira Brava*, *Extractum pareiræ*—*e. of White Poppy*, *Extractum papaveris*—*e. of Quassia*, *Extractum quassiae*—*e. of Quinia*, *Extractum quiniæ*—*e. of Rhatany*, *Extractum kramerizæ*—*e. of Rhubarb*, *Extractum rhei*—*e. of Rhubarb*, *fluid*, *Extractum rhei fluidum*—*e. of Rue*, *Extractum rutæ*—*e. of Sarsaparilla*, *Extractum sarsaparillæ*—*e. of Sarsaparilla*, *fluid*, *Extractum sarsaparillæ fluidum*—*e. of Savine*, *Extractum foliorum sabinæ*—*e. of Senna*, *fluid*, *Extractum sennæ fluidum*—*e. of Spigelia* and *Senna*, *Extractum spigeliæ et sennæ*—*e. of Stramonium*, *Extractum stramonii*—*e. of Uva Ursi*, *Extractum uvæ ursi*—*e. of Valerian*, *Extractum valerianæ*—*e. of Wormwood*, *Extractum cacuminum absinthii*.

EXTRACTA, see *Extractum*—*e. Fluida*, see *Extractum*.

EXTRACTIF, *Extractive*.

EXTRACTION, *Extractio*, *Ecæresis*, *Exogogê*, *Exolcê*, *Exelcysmus*, *Enchyloësis*, from *extrahere*, *extractum*, (*ex*, and *trahere*,) 'to draw out.' The act of removing an extraneous substance from any part of the body. Thus, a splinter is said to be extracted. It is also applied to the removal of certain parts. The cataract is said to be *extracted*; a tooth is *extracted*, when carious, &c.

EXTRACTIVE, (*F.*) *Extractif*. Same etymon. A peculiar, immediate principle, which has been admitted in extracts. Thus, *bitter extractive* is the immediate principle of bitter vegetables, &c.

EXTRACTS, see *Extractum*—*e. Fluid*, see *Extractum*.

EXTRACTUM, *Extract*, *Ecchyloëma*, (*F.*) *Extrait*, *Apostolé*. Same etymon. An extract is prepared by evaporating vegetable solutions, till a tenacious mass is obtained. When prepared from an infusion or decoction, it is called a *watery*; from ether, an *etheral*; and from alcohol, an *alcoholic* or a *spirituous extract*. Both kinds contain all the principles of the vegetable that are soluble in the menstrua with which they are prepared; but the volatile parts are dissipated, and some of the fixed parts are decomposed; the proper extractive is oxygenized, and the virtues of the vegetable substance consequently altered or destroyed. Extracts are *hard*, *soft*, or *fluid*: the consistence of the soft being such as to retain the pilular form without the addition of a powder. A patent was taken out, many years ago, by a Mr. Barry, of London, for preparing them in vacuo; and, as the temperature is much lower than in the ordinary method, the virtues of the plant are less altered, and the extracts are generally green. Extracts are also prepared by dis-

placement or percolation. They have, likewise, received different names, according to their predominant principle. The *gummy* or *mucous*, or *mucilaginous*, are those which are mainly composed of gum or mucilage. Gum tragacanth may be considered a pure gummy extract. *Gelatinous extracts* are those composed especially of gelatin: *resinous extracts*, those of a resinous character; *extracto-resinous*, those composed of extractive or colouring matter and resin; *gum-resinous*, those containing gum and resin; and *saponaceous* or *saponeous saline*, those containing a notable quantity of saline substances and a resinous matter, so combined with mucus and other soluble substances, that they cannot be separated. Under the name *Fluid Extracts*—*Extracta Fluida*—a class of preparations has been introduced into the United States Pharmacopœia (1851), the distinctive character of which is, the concentration of the active ingredients of medicinal substances into a small bulk, in the liquid form. *Extractum sarsaparillæ fluidum* is one of these.

EXTRACTUM ABSINTHII CACUMINUM, *Extract of Wormwood*. (A decoction defecated and evaporated.) The flavour is dissipated along with the essential oil. It is a bitter tonic. Dose, gr. x to ℥j, in pill.

EXTRACTUM ACONITI, *Extract of Aconite, Succus spissatus aconiti napelli*, (from the inspissated juice without defecation.) It is esteemed to be narcotic and diuretic; and has been given in the cases referred to under Aconitum. Dose, gr. j, gradually increasing it.

EXTRACTUM ACONITI ALCOHOLICUM, *Alcoholic Extract of Aconite*. (Aconit. in pulv. crass. ℥j; Alcohol, dilut. Oiv. Moisten the aconite with half a pint of diluted alcohol: let it stand for 24 hours; transfer it to a displacement apparatus, and gradually add the remainder of the diluted alcohol. When the last portion of this has penetrated the aconite, pour in from time to time water sufficient to keep the powder covered. Stop the filtration when the liquid which passes begins to produce a precipitate, as it falls, in that which has already passed. Distil off the alcohol, and evaporate to a proper consistence.—Ph. U. S.) Dose, half a grain or a grain.

EXTRACTUM ALLOES, *Extract of Aloes, E. aloës aquosum seu purificatum*, *Purified extract of aloes*. (The gummy part extracted by boiling water, defecated and inspissated.) Dose, gr. v to gr. xv.

EXTRACTUM ALOES AQUOSUM, *E. aloës*—*E. Aloës purificatum*, *E. aloës*.

EXTRACTUM ANTHEMIDIS, *E. anthemidis nobilis seu chamæmeli seu florum chamæmeli*, *Extract of Chamemile*. The volatile oil is dissipated in this preparation. It is a pure, grateful bitter, and is tonic and stomachic. Dose, gr. x to gr. xx, in pills.

EXTRACTUM ASPARAGI, see Asparagus.

EXTRACTUM BELLADONNÆ, *Succus spissatus atropæ belladonnæ*, *Extract of Belladonna* (an expressed juice inspissated). Properties same as those of the plant. Dose, gr. ¼, gradually increased. It dilates the pupil when applied to the eye.

EXTRACTUM BELLADONNÆ ALCOHOLICUM, *Alcoholia extract of Belladonna*. (Prepared like the extractum aconiti alcoholicum.—Ph. U. S.) Dose, half a grain or a grain.

EXTRACTUM CANNABIS, see Banguæ.

EXTRACTUM CARNIS, *Extract of Flesh*. An extract formed by evaporating beef tea in a water-bath. It can be kept for sale. An ounce of it may represent thirty-two ounces of meat.

EXTRACTUM CASCARILLÆ RESINOSUM, *Resinous extract of cascarrilla*. (Cort. cascarrillæ, in

pulv. crass. ℥j; Sp. vini rect. ℔iv. Digest for four days; then decant and strain: boil the residuum in ten pints of water to two: filter and evaporate the decoction, and distil the tincture in a retort, till both are thickened: then mix and evaporate to a pilular consistence.) Dose, gr. x to gr. xx, in pills.

EXTRACTUM CATECHU, *Catechu*—*E. Catharticum*, *E. Colocyntithidis compositum*.

EXTRACTUM CATHOLICUM, (F.) *Extrait Catholique*. This epithet is given to pills composed of aloes, black hellebore, and colocynth, resin of jalap, and scammony. See, also, *Extractum colocyntithidis compositum*.

EXTRACTUM CHAMÆMELI, *E. Anthemidis*—*E. Cicutæ*, *E. Conii*.

EXTRACTUM CINCHONÆ, *E. Corticis Peruviani seu Cinchonæ Molle*, *Extract of Cinchona or of Bark*. (A decoction evaporated.) The active principles are similar to those of the bark in substance; but it is not so effectual, owing to the chymical change induced in the drug during the boiling. When reduced, by drying, to a state fit for being powdered, it is called the *Hard Extract of Bark*, *Extractum Corticis Peruviani durum*, *E. Cinchonæ durum*. Dose, gr. x to ℥ss.

EXTRACTUM CINCHONÆ FLAVÆ, of the Pharmacopœia of the United States (1842), is directed to be prepared as follows: *Yellow Bark*, in coarse powder, ℔j; Alcohol, Oiv; Water, a sufficient quantity. Macerate the Peruvian bark with the alcohol for four days; then filter by a displacement apparatus, and when the liquid ceases to pass, pour gradually on the bark water sufficient to keep its surface covered. When the filtered tincture measures four pints, set it aside, and proceed with the filtration until six pints of infusion are obtained. Distil off the alcohol from the tincture and evaporate the infusion till the liquids are respectively brought to the consistence of thin honey; then mix and evaporate to form an extract.—Ph. U. S.

EXTRACTUM CINCHONÆ PALMIDÆ (Ph. L & E.) May be prepared in the same manner.

EXTRACTUM CINCHONÆ RESINOSUM, *E. Cinchonæ lanceifolia seu Cinchonæ Rubra resinolum*, *Resinous Extract of Bark, E. Cinchonæ Resinæ*. The aqueo-spirituous extract contains both the extractive and resin of the bark. Dose, gr. x to xxx.

EXTRACTUM CINCHONÆ RUBRÆ (Ph. U. S.), may be prepared in the same manner as *Extractum cinchonæ flavæ*.

EXTRACTUM COLCHICI, *Extract of Colchicum*, (Ph. L.) is prepared in the same manner as *Extractum Aconiti*. Dose, one or two grains.

EXTRACTUM COLCHICI ACETICUM, *Aceticum or Acetic Extract of Colchicum*. (Colchic. rad. in pulv. crass. ℥j, Acid. acet. f℥iv, Aquæ q. s. To the acid add a pint of water, and mix this with the root. Put the mixture in a percolator, and pour on water until the liquid that passes has little or no taste. Evaporate to a proper consistence. Ph. U. S.) Dose, gr. j to gr. iij.

EXTRACTUM COLOCYNTHIDIS, *Extract of Colocyntith*. (Ph. L & E.) A cathartic, in the dose of from gr. v to ℥ss.

EXTRACTUM COLOCYNTHIDIS COMPOSITUM, *E. Catharticum seu Catholicum seu Quercetani*, *Compound Extract of Colocyntith*. (Colocyntith. pulp. concis. ℥vj, Aloes pulv. ℥xij, Scammon. pulv. ℥iv, Cardamom. pulv. ℥j, Saponis ℥ij, Alcohol, dilut. cong. Macerate the pulp in the spirit at a gentle heat for four days; strain; add the aloes and scammony; then distil off the spirit and mix in the cardamom seeds. Ph. U. S.) It is an active cathartic, and is used in obstinate visceral obstructions, &c. Dose, gr. vj to ℥ss.

EXTRACTUM CONII, *E. Cicutæ*, *Succus cicutæ*

episcatus, *S. spissatus conii maculati*. *Extract of Hemlock.* (Expressed juice inspissated without defecation.) Employed in the same cases as the conium. Dose, gr. ij to ℥j.

EXTRACTUM CONII ALCOHOLICUM, *Alcoholic Extract of Hemlock.* (Prepared like the Extractum aconiti alcoholicum.—Ph. U. S.) Dose, two or three grains.

EXTRACTUM CONVULVULI JALAPÆ, *E. Jalapæ* —e. *Corticis Peruviani*, *E. Cinchona*.

EXTRACTUM CUBEBÆ FLUIDUM, *Fluid Extract of Cubebs.* (Cubebs exhausted by ether through percolation, and the solution evaporated. Ph. U. S.) Dose, from 5 to 30 minims.

EXTRACTUM DIGITALIS, *Extract of Foxglove.* Prepared like Extractum conii. Dose, from half a grain to two grains.

EXTRACTUM DULCAMARÆ, *Extract of Bittersweet.* (Prepared by displacement from bittersweet in coarse powder.—Ph. U. S.)

EXTRACTUM ELATERII, *Elaterium, Extract of Elaterium.* (The fecula of the expressed juice.) It is violently cathartic, hydragogue, and sometimes emetic. Dose, gr. ss every hour till it operates.

EXTRACTUM ERGOTÆ, *Ergotin.*

EXTRACTUM GENISTÆ CACUMINUM, *Extract of Broom Tops.* Diuretic and stomachic. Dose, ʒss to ʒj.

EXTRACTUM GENTIANÆ, *E. Gentianæ luteæ seu Radicis Gentianæ, Extract of Gentian.* (The evaporated decoction.) Prepared also by displacement.—Ph. U. S.) Properties like those of Gentian. Dose, gr. x to ʒss.

EXTRACTUM GLYCYRRHIZÆ, *Succus Glycyrrhizæ inspissatus seu Liquiritiæ, Spanish Juice, Italian Juice, Extract of Liquorice.* (The evaporated decoction.) It is demulcent, taken ad libitum.

Refined Liquorice, which is sold in the form of cylinders, is made by gently evaporating a solution of the pure extract of liquorice with half its weight of gum Arabic, rolling the mass and cutting it into lengths, and then polishing, by rolling them together in a box.

EXTRACTUM GRAMINIS, *Extract of Grass.* An extract prepared from *Triticum repens*. It is considered by the Germans to be a mild tonic; and is greatly used, especially in convalescence from fever. It is probably devoid of all injurious properties, and as probably totally inefficacious.

EXTRACTUM HÆMATOXYLI, *E. Hæmatoxyli Campechiani seu Scobis Hæmatoxyli, Extract of Logwood.* (The evaporated decoction.) It is astringent. Dose, gr. x to ʒj.

EXTRACTUM HELLEBORI, *E. Hellebori nigri, Extract of Black Hellebore.* (The evaporated decoction.) In large doses, this is cathartic; in smaller, diuretic, resolvent, [?] and emmenagogue. [?] Dose, as a cathartic, gr. x to ℥j; as an emmenagogue, gr. ij to gr. x.

EXTRACTUM HELLEBORI, of the Pharmacopœia of the United States (1842), is prepared from Black Hellebore, in coarse powder, like the Extractum aconiti alcoholicum.

EXTRACTUM HUMULI, *E. Lu'puli, Extract of hops.* (The evaporated decoction.) It is tonic, anodyne, [?] diuretic. [?] Dose, gr. v to ʒi.

EXTRACTUM HYOSCYAMI, *Succus spissatus Hyoscyami, Extract of Henbane.* (The expressed juice, inspissated without defecation.) Its virtues are narcotic. Dose, gr. v to ℥ss.

EXTRACTUM HYOSCYAMI ALCOHOLICUM, *Alcoholic Extract of Henbane.* (Prepared from leaves of Hyoscyamus, in coarse powder, like the Extractum aconiti alcoholicum.—Ph. U. S.) Dose, one or two grains.

EXTRACTUM JALAPÆ, *E. Convolvuli Jalapæ seu Jala'pii seu Jala'pi, Extract of Jalap.* (A

spirituous tincture distilled; and an aqueous decoction evaporated; the residua being mixed together: kept both soft and hard.) It is cathartic and hydragogue. Dose, gr. x to ℥j.

EXTRACTUM JALAPÆ of the Ph. U. S. is prepared like Extractum cinchona. Ph. U. S.

EXTRACTUM JALAPÆ RESINOSUM, *Resina Jala'pæ, Resinous Extract of Jalap.* This is cathartic.

EXTRACTUM JUGLANS'DIS, *Extract of Butternut.* (Prepared by displacement from butternut, in coarse powder.—Ph. U. S.)

EXTRACTUM KRAMERIE, *Extract of Rhatany* (Prepared by displacement from rhatany, in coarse powder.—Ph. U. S.)

EXTRACTUM LACTU'CÆ, *Succus spissatus Lactuce sativæ, Extract of Lettuce.* (Leaves of fresh lettuce ℥ij; beat them in a stone mortar, sprinkling them with water; then express the juice and evaporate, without allowing it to subside until it acquires a proper degree of consistence.) It is said to be narcotic and diaphoretic. Dose, gr. ij to gr. x.

An extract is, sometimes, made from the juice of the wild lettuce, *Lactuca viro'sa*, which is regarded as diuretic.

EXTRACTUM MARTIS ACETICI, *Ferri Acetas*—e. *Nucis Vomiceæ*, see *Strychnos nux vomica*.

EXTRACTUM NUCIS VOMICÆ, *Extract of Nux Vomica* (*Nucis Vomice*, ℥j; Alcohol, q. s. Expose the nux vomica to steam until it is softened, slice, dry and grind to powder. Put it into a percolator, and pour alcohol upon it gradually, until the liquor passes without bitterness. Filter, distil off the alcohol, and evaporate. Ph. U. S.) Dose, from half a grain to two grains.

EXTRACTUM OPII, *E. Opii gummosum seu Thebaicum seu Opii gummosum, Extract of Opium, Laudanum opiatum seu simplex, Opium colatum seu depuratum.* (A watery solution defecated and evaporated.) Dose, gr. ss to gr. v,—about half that of opium.

EXTRACTUM PANCHYMACO'GUM. A drastic medicine, composed of colocynth, bruised with its seeds; senna bruised; black hellebore root, Agaric, Scammony, in powder, Extract of Aloe, and Powder of Diarrhodon.

EXTRACTUM PAPAV'ERIS, *E. Papav'eris somniferi seu Papav'eris albi, Extract of white poppy.* (The decoction evaporated.) It possesses nearly the same virtues as opium, but is much weaker. Dose, gr. ij to ℥j.

EXTRACTUM PAREIRÆ, *Extract of Pareira Brava.* Prepared like Extractum Hæmatoxyli. Dose, ten grains to half a drachm.

EXTRACTUM PIP'ERIS FLUIDUM, *Fluid Extract of Black Pepper.* (Black pepper exhausted by ether through percolation, the solution evaporated, and the piperin in crystals separated by expression. Ph. U. S.) Dose, one or two minims.

EXTRACTUM PODOPHYLLI, *Extract of May-apple.* (Prepared from podophyllum, in coarse powder, in the same manner as the Extract of cinchona.—Ph. U. S.) Dose, gr. v to gr. xv, as a cathartic.

EXTRACTUM PURGANS, see *Hedera helix*—e. *Quercetani*, *E. Colocythidis compositum*.

EXTRACTUM QUASSIÆ, *Extract of Quassia.* (Prepared by displacement from Quassia rasped.—Ph. U. S.) Dose, as a tonic, five grains.

EXTRACTUM CORTICIS QUERCUS, *Extract of oak bark.* (The decoction evaporated.) It is astringent and tonic.

EXTRACTUM QUINIE, *Quiniæ sulphas impuratus.* This is made by evaporating the liquor poured off the crystals of sulphate of quinia to the consistence of a pilular mass. Twenty-four grains will generally arrest an intermittent.

EXTRACTUM RHEI, *Extract of Rhubarb.* (A

olution in diluted alcohol evaporated.) Uses like those of the powdered root. Dose, gr. x. to ʒss.

EXTRACTUM RHEI FLUIDUM, *Fluid Extract of Rhubarb.* (Rhef in pulv. crass. ʒviij, Sacchar. ʒv. Tinct. Zingib. fʒss, Ol. fœnicul., Ol. anis. aa ʒiv; Alcohol. dilut. q. s. Digest the rhubarb, mixed with an equal bulk of coarse sand, with ʒxij of the diluted alcohol for 24 hours. Put the mass into the percolator, and pour on diluted alcohol until the liquid that passes has little odour or taste of rhubarb; evaporate to fʒv; dissolve it in the sugar, and mix the tincture of ginger and oils. Ph. U. S.) Dose, fʒj.

EXTRACTUM RU'DII, *Extract of Rudius, (F.) Extrait de Rudius.* Pills made of colocynth, agaric, scammony, roots of black hellebore and jalap, socotrine aloes, cinnamon, mace, cloves, and alcohol.

EXTRACTUM RUTÆ GRAVEOLENTIS, *E. Ruta seu foliorum Rutæ, Extract of Rue.* (A decoction evaporated.) Tonic, stomachic. The volatile oil being dissipated in the boiling, this is not a good preparation. Dose, gr. x to ʒj.

EXTRACTUM FOLIORUM SABINÆ, *Extract of Savine.* (A decoction evaporated.) Tonic. The same remarks may be made on this preparation as on the last. Dose, gr. x to ʒss.

EXTRACTUM SANGUINIS BOVINI, see Blood.

EXTRACTUM SARSAPARILLÆ, *Extract of Sarsaparilla.* (A strained decoction evaporated.) Virtues the same as those of the powdered root. Dose, gr. x to ʒj. *Extractum Sarsaparillæ* of the United States Pharmacopœia is prepared from Sarsaparilla, in coarse powder, like the *Extractum aconiti alcoholicum*.

EXTRACTUM SARSAPARILLÆ FLUIDUM, *E. Sarsæ Liq'idum, E. Sarsæ Fluidum, Fluid Extract of Sarsaparilla.* (Sarsaparill. concis. et contus. ʒxxvj, Glycyrrhiz. contus., Sassafr. rad. contus., aa ʒij, Alceæci concis. ʒvj, Sacchar. ʒxij, Alcohol. dilut. Oviij. Macerate, with the exception of the sugar, for 14 days; express and filter; evaporate to fʒxij; and add the sugar. Ph. U. S.) Dose, a fluidrachm.

EXTRACTUM SATURNI, GOULARD'S, Liqueur plumbi subacetatis — e. Scobis hæmatoxyli, E. hæmatoxyli.

EXTRACTUM SENNÆ FLUIDUM, *Fluid Extract of Senna.* (Senna in pulv. crass. ʒijss; Sacchar. ʒxx; Ol. Fœnicul. fʒj; Sp. Ether. compos. fʒij; Alcohol. dilut. Oiv. Mix the senna and diluted alcohol; let the mixture stand for 24 hours; put it into a percolator, and gradually pour on water mixed with one-third its bulk of alcohol, until a gallon and a half of liquid shall have passed; evaporate to fʒxx; add the sugar, and, when it is dissolved, the compound spirit of ether holding the oil in solution. Dose, fʒss.

EXTRACTUM SPIGELIÆ ET SENNÆ FLUIDUM, *Fluid Extract of Spigelia and Senna.* Spigel. in pulv. crass. ʒij; Senna in pulv. crass. ʒvj; Sacchar. Biss; Potass. carbon. ʒvj; Ol. Carui, Ol. Anisi, aa fʒss; Alcohol. dilut. q. s. Pour on the spigelia and senna Oij of diluted alcohol; let it stand for 48 hours; place it in a percolator, and pour on gradually diluted alcohol until half a gallon has passed: evaporate to a pint; add the carbonate of potassa; and afterwards the sugar, previously triturated with the oils, and dissolve. Ph. U. S.)

EXTRACTUM STRAMONII, *E. Stramonii foliorum, (Ph. U. S.), Extract of Stramonium or of Stramonium leaves.* (The expressed juice inspissated.) The *Extractum Stramonii Sem'inis, Extract of Stramonium Seed*, is made from the powdered seed by means of diluted alcohol, and with the aid of the percolator; — the solution being evaporated. Ph. U. S.) Used as a narcotic in

asthma and other spasmodic affections. Dose, gr. ij to gr. x.

EXTRACTUM STYRACIS, see STYRAX.

EXTRACTUM TARAXACI, *E. Herbae et Radicis Taraxaci, Extract of Dandelion.* (The strained juice evaporated.) It has been considered deobstruent, laxative, and diuretic. Dose, gr. x to ʒj.

EXTRACTUM THEBAICUM, E. Opii.

EXTRACTUM UVÆ URSI, *Extract of Uva Ursi.* (The Decoction evaporated.) Dose, gr. v to gr. xxx.

EXTRACTUM VALERIANÆ, *Extract of Valerian.* (An expressed decoction evaporated.) The virtues of the valerian being dependent upon its essential oil, this is an objectionable preparation. Dose, gr. x to ʒj.

A fluid extract of Valerian, *Extractum Valerianæ fluidum*, has been introduced into the last edition of the Pharmacopœia U. S. (1851). It is prepared by exhausting the valerian by ether and alcohol, through the percolator, and evaporating. Dose, fʒj or fʒij.

EXTRAIT, *Extract* — e. *Alcoolique de noix vomique*, see *Strychnos nux vomica* — e. *de Kino*, *Kino* — e. *des Fruits*, *Rob* — e. *Mémostatique de Bonjean*, *Ergotin* — e. *de Viande*, *Osmazome*.

EXTRA'NEOUS BODY, *Foreign body*, from extra, 'without.' *Corpus extraneum*, *C. externum*, *C. alienum*, (F.) *Corps étranger*. Any solid, liquid, or gaseous substance, inanimate or animate, proceeding from without, or formed in the body; and which constitutes no part of the body, but occupies, in the substance of the textures, or some of the cavities, a place foreign to it.

EXTRAVASATION, *Extravasatio*, *Extravasation*, from extra, 'out of,' and vasa, 'vessels.' Escape of a fluid — *extravasatum* — from the vessel containing it, and infiltration or effusion of the fluid into the surrounding textures.

EXTRAVASATUM, see *Extravasatio*.

EXTRAVASION, *Extravasation*.

EXTREMITY, *Extrem'itas*, (F.) *Extrémité*; from *extremus*, 'the outermost;' the end or termination of a thing. The limbs, *acrotéria*, have been so called, — as the upper and lower extremities. See *Membrum*. It has been, also, used to express the last moments of life; as when we say, a patient is in 'extremity,' (F.) *le malade est à l'extrémité, à toute extrémité*.

EXTRIN'SIC, *Extrin'secus*. 'From without.' That which comes from without. This term has been used for muscles, which surround certain organs and attach them to the neighbouring parts; in order to distinguish them from other muscles, which enter into the intimate composition of those organs, and which have been named *intrinsic*. Thus, there are extrinsic and intrinsic muscles of the tongue, ear, &c.

EXTROVERSI'ON, *Extrophia*.

EXTUBERANTIA, *Protuberance*.

EXTUBERATIO, *Protuberance*.

EXTUMEFAC'TIO, *Swelling*.

EXTUS'SIO, from *ex*, and *tussio*, 'a cough.' 'I cough with expectoration.'

EXU'BER, from *ex*, 'out of,' 'devoid of,' and *ubera*, 'breasts,' *Apogalactos*. 'A child which has been weaned.'

EXUDATION CORPUSCLES, see *Corpuscles*, *exudation*.

EXULCERATIO, *Eethlimma*, *Ulceration* — e. *Uteri*, *Hysterelcosis* — e. *Ventriculi*, *Gastrelcosis*.

EXUMBILICATIO, *Exomphalos*.

EXUSTIO, *Cauterization*.

EXUTORIUM, *Fonticulus*.

EXUTORY, *Fonticulus*.

EXUVIÆ, see *Exuviation*.

EXUVIATION, *Exuviat'io*, from *exuere*, 'to put off.' The shedding or casting off of parts, as

the deciduous teeth, the skins of serpents, &c., which are hence called *Ecu'vixæ*.

EYE, Sax. *eag*, Teuton. *Auge*, *Oculus*, *Ops*, *Omma*, *Ophthalmos*, *Illos*, *Op'tilos* (Doric,) *Vigilium* *Org'uanum*, (Sc.) *Ee*, plural *Een*, (F.) *Œil*. The eye is the immediate organ of vision. It is seated in the orbit, while its dependencies, called by Haller *Tutamina Oculi*, occupy the circumference of the cavity, and are composed of the eyebrows, the eyelids, cilia, glands of Meibomius, &c. The *Ball*, *Globe*, or *Bulb* of the *Eye*, *Bulbus Oculi*, is covered anteriorly by the tunica conjunctiva; is moved by six muscles, four straight, two oblique, and is constituted of membranes, as the *sclerotic*, *cornea*, *choroid*, *tunica Jacobi*, *retina*, *iris*, *hyaloid*, and, in the fetus, the *membrana pupillaris*; and of fluids, called *Humours*, or *Media*—the *aqueous*, *crystalline*, and *vitreous*. The eyeball is invested with a membranous tunic, which separates it from the other structures of the orbit, and forms a smooth, hollow surface, by which its motions are facilitated. This investment has been called *cell'ular capsule of the eye*, *oc'ular capsule*, *tunica vagina'lis oculi*, *vag'inul coat*, and *submuc'ular fascia of the eye*. The vessels of the eye proceed from the ophthalmic artery. The nerves, except the optic, are chiefly furnished from the ophthalmic ganglion. The following are the dimensions, &c., of the organ, on the authority of Petit, Young, Gordon, and Brewster:

Length of the antero-posterior diameter of the eye.....	0.91
Vertical chord of the cornea.....	0.45
Versed sine of the cornea.....	0.11
Horizontal chord of the cornea.....	0.47
Size of pupil seen through the cornea, 0.27 to	0.13
Size of pupil diminished by magnifying power of cornea to, from.....	0.25 to 0.12
Radius of the anterior surface of the crystalline.....	0.30
Radius of posterior surface.....	0.22
Principal focal distance of lens.....	1.73
Distance of the centre of the optic nerve from the foramen centrale of Sömmerring...	0.11
Distance of the iris from the cornea.....	0.10

Distance of the iris from the anterior surface of the crystalline.....	0.02
Field of vision above a horizontal line, 50° } 120°	
Field of vision below a horizontal line, 70° }	
Field of vision in a horizontal plane.....	150°
Diameter of the crystalline in a woman above fifty years of age.....	0.373
Diameter of the cornea.....	0.400
Thickness of the crystalline.....	0.172
Thickness of the cornea.....	0.042

EYE, APPLE, see Melum—e. Apple of the, Pupil.

EYE BALM, *Hydrastis Canadensis*.

EYE-BREEK, Palpebra.

EYE-BREEN, see Supercilium.

EYE BRIGHT, *Euphrasia officinalis*, *Lobelia Veronica*—e. Brow, Supercilium.

EYE, CAT'S, Amaurotic, see Amaurotic—e. Cellular capsule of the, see Eye.

EYE, CYLIND'RICAL. An eye is so called in which the rays of light, on entering the organ, are refracted to a nearer focus in a vertical than in a horizontal plane.

EYE DROP, Tear.

EYE GLASS, *Scaphium oculare*, see Spectacles. Also, a glass adapted for the application of collyria to the eye.

EYE, GUM OF THE, *Chassie*.

EYE LASHES, Cilia—e. Lid, Palpebra—e. Lid, granular, Trachoma—e. Melon, see Melum—e. Purulent, Ophthalmia, purulent, of infants.

EYE SALVE, SINGLETON'S, Unguentum Hydragryi nitrico-oxydi; Eye-salve, Smellome's, see Cupri subacetat.

EYE, SIGHT OF THE, Pupil.

EYE STONE. The shelly operculum of small turbinidæ. Used at Guernsey to get things out of the eyes. Being put into the inner corner of the eye, under the eyelid, it works its way out at the outer corner, and brings out any foreign substance with it.—Gray.

EYE TEETH, Canine teeth—e. of Typhon, Scilla—e. Water, Collyrium—e. Water, common, Liquor zinci sulphatis cum camphorâ—e. Water, blue, Liquor cupri ammoniati—e. Watery, Epiphora.

EYE, WHITE OF THE, see Sclerotic.

F.

FABA, Phaseolus, *Vicia faba*—f. *Ægyptiaca*, *Nymphæa nelumbo*—f. *Cathartica*, *Jatropha curcas*—f. *Crassa*, *Sedum telephium*—f. *Febri-fuga*, *Ignatia amara*—f. *Græca*, *Diospyros lotus*—f. *Indica*, *Ignatia amara*—f. *Major*, *Vicia faba*—f. *Peckurei*, *Pichurim Beans*—f. *Pichurim*, *Pichurim Beans*—f. *Purgatrix*, *Ricinus communis*—f. *Sancti Ignatii*, *Ignatia amara*—f. *Squilla*, *Hyoxyeanus*—f. *Vulgaris*, *Vicia faba*.

FABÆ, *Onisci aselli*.

FABAGELLE, *Zygophyllum fagabo*.

FABARIA CRASSULA, *Sedum telephium*.

FABRICA ANDROGYNA, Hermaphrodeity.

FACE, *Facies*, *Vultus*, *Vultus*, *Proso'pon*, (F.) *Fare*. The face is the anterior part of the head.

It is formed of 13 bones, viz., the two superior maxillary, the two malar, the two ossa nasi, the two ossa unguis, the vomer, the two ossa spongiosa inferiora, the two palate bones, and the inferior maxillary, without including the frontal portion of the os frontis, and the 32 teeth, which may be considered to form part of it. Its numerous

muscles are chiefly destined for the organs of sight, hearing, taste, and smell. Its arteries proceed from the external carotid: its veins end in the jugular, and its nerves draw their origin immediately from the brain.

The face experiences alterations in disease, which it is important to attend to. It is yellow in jaundice, pale and puffy in dropsy; and its expression is very different, according to the seat of irritation, so that, in infants, by an attention to medical physiognomy, we can often detect the seat of disease.

Hippocrates has well depicted the change which it experiences in one exhausted by long sickness, by great evacuations, excessive hunger, watchfulness, &c., threatening dissolution. Hence this state has been called *Facies Hippocrat'ica*, *Facies Cadaver'ica*, *F. Tortiva'lis*. In this, the nose is pinched; the eyes are sunk; the temples hollow; the ears cold, and retracted; the skin of the forehead tense, and dry; the complexion livid; the lips pendent, relaxed, and cold, &c.

The term *Face* (F.) is likewise given to one of the aspects of an organ; thus, we say, the *superior face of the stomach*.

FACE AGUE, Neuralgia, facial.

FACE INJECTÉE, see *Vultueux*—*f. Vultueux*, see *Vultueux*.

FACET, (F.) *Facette*. Diminutive of *Face*. A small face. A small, circumscribed portion of the surface of a bone, as the *articular facette of a bone*.

FACHINGEN, MINERAL WATERS OF. These springs are at no great distance from those of Geilenau, and two miles north of Wisbaden. They contain free carbonic acid: carbonate, sulphate, and phosphate of soda; chloride of sodium, carbonate of lime, magnesia, and iron.

FACIAL, *Facialis*, from *facies*, 'the face.' Belonging to, or connected with, the face.

FACIAL ANGLE, see *Angle, facial*.

FACIAL ARTERY, *La'bial artery*, *Angular* or *external maxillary artery*, *A. palato-labial*—(Ch.), is a branch of the external carotid, which rises beneath the digastricus, and is distributed to almost every part of the face. It furnishes the *inferior palatal, submental, superior labial, inferior labial, and dorsal nasi*.

FACIAL LINE, see *Angle, facial*.

FACIAL NERVE, *Ramus du'rrior septimæ conjugatio'nis*, *Sympatheticus minor*, *Par septimum seu faciale*, *Comunicans faciei nervus*, *Portio dura of the 1th pair*, *Respiratory nerve of the face*. This nerve arises from the inferior and lateral part of the tuber annulare, in the groove which separates it from the medulla oblongata, external to the corpora olivaria, and by the side of the auditory nerve. It issues from the cranium by the *meatus auditorius internus*; enters the aqueduct of Fallopius; receives a branch of the Vidian nerve; forms a gangliform swelling—*Intumescentia gangliformis*;—sends off filaments to the internal muscles of the malleus and stapes; furnishes, according to many anatomists, that called *Chorda Tympani*; makes its exit at the foramen stylo-mastoideum, and divides into two branches—the *temporo-facial*, and *cervico-facial*. On the face it is termed *Pes auseri'nis*, *Plexus nervorum auseri'nis*. See *Portio Wisbergii*.

FACIAL VEIN, *Palato-labial*—(Ch.), arises between the skin and frontalis muscle, and bears the name *Vena Frontalis*, (F.) *Veine frontale* ou *V. Préaryate*. It then descends, vertically, towards the greater angle of the eye, where it is called *Angularis*; and afterwards descends, obliquely, on the face, to open into the internal jugular, after having received branches, which correspond with those of the facial artery. It is only in the latter part of its course that it is called *Facial Vein*. Chaussier calls the external carotid artery, *Facial Artery*.

FACIENT, *faciens*, 'making,' from *facio*, 'I make.' A suffix, as in *Calefacient*, *Rubefacient*, &c., 'warm making,' 'red making.'

FACIES, *Face*—*f. Cadaverica*, see *Face*—*f. Concaua pedis*, *Sole*—*f. Hippocratica*, see *Face*—*f. Inferior pedis*, *Sole*—*f. Tortualls*, see *Face*.

FACTICE, *Factitious*.

FACITIV'IOUS, *Facitiv'ius*, (F.) *Factice*, from *facere*, *factum*, 'to make.' Artificial. That which is made by art, in opposition to what is natural, or found already existing in nature. Thus, we say, *factitious mineral waters*, for artificial mineral waters.

FACULTAS, *Faculty*—*f. Auctrix*, *Plastic force*—*f. Formatrix*, *Plastic force*—*f. Nutrix*, *Plastic force*—*f. Vegetativa*, *Plastic force*—*f. Zotica*, *Vis vitalis*.

FACULTATES NATURALES, see *Function*.

FACULTY, *Facultas*, *Dynamis*, *Power*, *Virtue*, (F.) *Faculté*, from *facilis*, (itself from *facere*, 'to make,') 'what can be made or done.' The power of executing any function or act. The collection of the intellectual faculties constitutes the understanding. We say, also, *vital faculties for vital properties*, &c. *Faculty* likewise means the whole body of the medical profession, and, also, a body of medical or other professors.

FÆCAL, *Stercoraceus*—*f. Retention*, *Constipation*.

FÆCES, Plural of *Fæx*; *Feces*, *Chersæ*, 'the dregs of any thing.' *Feculence*, (F.) *Fèces*. The alvine evacuations are so called; (F.) *Garderobes*; the excrements, *Impuritates alvine*, *Fæcal matter*. See *Excrement*.

FÆCES INDURATE, *Seybala*.

FÆCOSITAS, *Feculence*.

FÆCULA, *Fecula*.

FÆCULENTIA, *Feculence*.

FÆCUNDATIO, *Fecundatio*.

FÆCUNDITAS, *Fecundity*.

FÆX, *Feculence*.

FAGARA OCTAN'DRA, *Xanthoxylum octan'dra*, *Elaph'rium tomento'sum*, *Am'yris tomento'sum*, from *fagus*, 'the beech,' which it resembles. *Ord. Rutaceæ*. The systematic name of the plant, which affords *Tacamaha'ca*, a resinous substance, that exudes from the tree *Tacamahaca*, which has a fragrant, delightful smell, was formerly in high estimation, as an ingredient in warm, stimulating plasters, and was given internally, like the balsams generally. The *East India Tacamahac*, *Balsamum Vir'idæ seu Cal'oba seu Mariæ*, *O'leum Mariæ*, (F.) *Baume de Calaba*, *Baume vert*, *B. Marie*, is yielded by *Calophyllum inophyllum* seu *Balsamaria inophyllum*.

The name *Tacamahac* is also given to a resin furnished by *Populus balsamifera* seu *tacamahaca*, (F.) *Peuplier baumier* ou *de la Caroline*, which grows in the northern parts of America and Siberia.

FAGARA PIPERITA, (F.) *Fagarier poivré*; a native of Japan, possesses the qualities of pepper, and is used as such by the Japanese. It is, also, employed as a rubefacient cataplasm.

FAGARAS'TRUM CAPEN'SÈ. *Nat. Ord. Xanthoxylæ*. A South African plant, the fruit of which is known to the Colonists as *wild Cardamom*; and, on account of its aromatic qualities, is prescribed in flatulency and paralysis.

FAGARIER POIVRÉ, *Fagara octandra*.

FAGOPYRUM, *Polygonum fagopyrum*.

FAGUS, *F. sylvatica*.

FAGUS CASTANEA. The systematic name of the *Chestnut Tree*; *Castanea*, *C. vulg'aris* seu *vesca*, *Lo'pina*, *Mota*, *Glans Jovis THEOPHRASTI*, *Ju'piter's* or *Sardin'ian Acorn*; the Common Chestnut, (F.) *Châtaignier commun*, of the Oak Family. *Ord. Cupuliferæ*. *Ser. Syst. Monœcia Polyandria*. The Chestnut, *Castanea nux*, (F.) *Châtaigne*, is farinaceous and nutritious, but not easy of digestion.

FAGUS CASTANEA PU'MILA. The *Chin'capin* or *Chinquapin*, *Castanea Pumila*, (F.) *Châtaignier nain*. The nut of this American tree is eaten like the chestnut. The bark, *Castanea* (Ph. U. S.), has been used in intermittents.

FAGUS PURPUREA, *F. sylvatica*.

FAGUS SYLVATICA. The systematic name of the *Beech*, *Fagus*, *F. sylvæ'tris* seu *purpu'rea*, *Oxya*, *Balan'da*, *Valan'ida*; the *Beech Tree*, (F.) *Hêtre*. The *Beech-nut* or *Beech-mast*, (F.) *Faine*, affords an oil, by expression, which is of a palatable character, and is eaten in some places

instead of butter. It has been supposed to be a good vernifuge, but it is no better than any mild oil.

FAGUS SYLVESTRIS, *F. sylvatica*.

FAIBLESSE, Debility.

FAIM, Hunger—*f. Canine*, *Boulimia*—*f. de Loup*, *Fomes lupina*.

FAINE, see *Fagus sylvatica*.

FAINT, see *Syncope*.

FAINTING, *Syncope*.

FAINTING-FIT, *Syncope*.

FAINTISHNESS, see *Syncope*.

FAINTNESS, *Languor*, *Syncope*.

FAIRBURN, MINERAL WATERS OF. The mineral waters at this place, which is in the county of Ross, in Scotland, are sulphureous, and frequented.

FAIRNTICKLED, see *Ephelides*.

FAISANDÉES (*VIANDES*), see *Hyposaprus*.

FAISCEAU, *Fasciculus*—*f. Intermédiaire de Wrisberg*, *Portio Wrisbergii*—*f. Petit*, *Fasciculus*.

FAIX, *Fœtus*.

FALCADINA, see *Scherlievo*.

FALCES, see *Falx*.

FAL'CIFORM, *Falciformis*, *Drepanoïdes*, from *falx*, *falcis*, 'a scythe,' and *forma*, 'shape.' Having the shape of a scythe. This term has been applied to different parts. See *Falx*, and *Sinus*.

FALCIFORM EXPANSION OF THE FASCIA LATA, is the scythe-shaped reflection of the fascia lata, which forms, outwards and upwards, the opening for the vena saphæna, and is attached to the crural arch by its superior extremity, forming the anterior paries of the canal of the same name.

FALLACIA, *Hallucination*.

FALLACIA OPTICA. An optic illusion.

FALLAND-EVYL, *Epilepsy*.

FALLING-DOWN, *Epilepsy*.

FALLING SICKNESS, *Epilepsy*.

FALLOPIAN TUBE, see *Tube*, *Fallopian*.

FALLPOISON, *Amianthum muscætoxicum*.

FALLTRANCK, *Faltranck* (G.), literally, *a drink against falls*. A vulnerary. It is a mixture of several aromatic and slightly astringent plants, which grow chiefly in the Swiss Alps, and hence the name—*Vulnéraire Suisse*—given to such dried plants cut into fragments. They are called, also, *Espèces Vulnéraires*, and *Thé Suisse*. Within the present century, in England, a kind of vulnerary beer was often prescribed, in country practice, in all cases of inward bruises. It bore the name *Cerevis'ia nigra*, or *black beer*, and was formed by infusing certain reputed vulnerary herbs in beer or ale.

The infusion of the *Falltranck* is aromatic, and slightly agreeable, but of no use in the cases for which it has been particularly recommended.

FALMOUTH, CLIMATE OF. The climate of Falmouth, in Cornwall, England, resembles that of Penzance: and, like it, is in many respects, a favourable retreat for the phthical during the winter months.

FALSA VIA, *False passage*.

FALSE, *Falsus*, *Nothus*, *Pseudo*, *Spu'rious*, *Bastard*, (F.) *Faux*, *Fausse*. This epithet has been frequently added to peripneumony, pleurisy, &c., to designate a disease similar to these, but less severe. Most commonly, a severe catarrh or pleurodynia has received the name. See *Peripneumonia notha*.

FALSE PASSAGE, *Falsa Via*, (F.) *Fausse Route*. An accidental passage, made in surgical operations, and particularly in introducing the catheter. The catheter is sometimes passed through into the rectum.

FALSE WATERS, *Hydrallan'tis*, *False Deliv'ery*. Water, which sometimes collects between the amnion and chorion, and is commonly discharged before the birth of the child.

We say, also, *False Ribs*, *False Rhubarb*, &c.

FALSETTO VOICE, see *Voice*.

FALSIFICA'TION, *Adultera'tio*, from *falsus*, 'false,' and *facere*, 'to make.' A fraudulent imitation or alteration of an aliment or medicine by different admixtures. *Manga'nium*, *Manganis'a'tio*. It is synonymous with *adulteration* and *sophistication*.

TABLE OF COMMON FALSIFICATIONS OF SOME OF THE MOST USEFUL DRUGS, &c.

MEDICINES.	ADULTERATIONS.	MODE OF DETECTION.
ACACIE GUMMI.	<i>Gum Senegal</i>	G. S. is clammy and tenacious. The A. G. is perfectly soluble in water, and its solution limpid.
ACETUM DESTILLATUM.	<i>Sulphuric Acid</i>	Acetate of barytes causes a white precipitate.
	<i>Nitric Acid</i>	By evaporating it, the residuum deflagrates, when thrown on burning charcoal.
	<i>Copper</i>	Supersaturate with ammonia—a blue colour is produced.
	<i>Lead</i>	Sulphuretted hydrogen causes a dark precipitate.
ACIDUM MURIATICUM.	<i>Sulphuric Acid</i>	Deposites by evaporation the salts it may contain; precipitates with solution of hydrochlorate of baryta if it contains sulphuric acid.
— NITRICUM.	<i>Muriatic and Sulphuric Acids</i> .	The presence of chlorine is indicated by a precipitate with nitrate of silver: that of sulphuric acid by the same result with hydrochlorate of baryta.
— SULPHURICUM.	<i>Muriatic and Nitric Acids</i> .	The presence of muriatic and nitric acid is indicated by the smell, when the acid tested is strongly heated.
— CITRICUM.	<i>Tartaric and Oxalic Acids</i> .	Their presence is indicated by forming a granular sediment in a concentrated solution of a neutral salt of potassa.

MEDICINES.	ADULTERATIONS.	MODE OF DETECTION.
ETHER RECTIFICATUS.	<i>Too dilute.</i> <i>Sulphuric Acid</i>	The S. G. detects this. By acetate of baryta. Precipitate white.
	<i>Alcohol</i>	With phosphorus a milky instead of limpid solution is formed.
AQUA AMMONIÆ.		A precipitation occurs on adding a solution of muriate of lime.
AMMONIÆ CARRO-	<i>Carbonic Acid</i>	It should be capable of complete volatilization by heat.
NAS.		The <i>Gutta Ammoniaci</i> are white, clear, and dry.
AMMONIACUM.....		The <i>lump Ammoniacum, lapis Ammoniaci</i> , is often adulterated with common resin.
	It ought to be entirely volatilized by a red heat.
AMMONII SULPHURE-	<i>Lead.</i>	Imparts a foliated texture, and is not vaporizable.
TUM.	<i>Arsenic</i>	A smell of garlic is emitted when thrown on live coals, &c.
	<i>Manganese and Iron.</i>	Are not vaporizable.
	<i>Copper</i>	The solution assumes a blue colour, when super-saturated with ammonia. It may be suspected when the salt deliquesces.
ARGENTI NITRAS.		The adulteration with nitrate of potassa is easily recognized by the fracture of a stick of it, which is radiated when pure; and granular if adulterated; or by precipitating a solution of the salts with a sufficient quantity of muriatic acid, and evaporating the clear liquor: the nitrate of potassa or other salts will remain.
ASBENICUM ALBUM.	<i>Chalk, Sulphate of Lime, Sulphate of Baryta.</i>	Not volatilizable by heat.
BALSAMUM PERUVIANUM.	<i>A mixture of Resin and some Volatile Oil, with Benzoin.</i>	Not easily detected.
	<i>Chloride of Sodium</i>	This disposes it to deliquesce.
CAPSICUM.		Digest in acetic acid, and add a solution of sulphuret of ammonia—a dark-coloured precipitate will be produced.
<i>Cayenne Pepper.</i>	<i>Red Lead</i>	
	<i>A mixture of dried blood, gum ammoniac, and a little real castor, stuffed into the scrotum of a goat.</i>	Smell and taste will generally detect the fraud.
CASTOREUM.		
<i>Castor.</i>	<i>Earth or Peasmeal</i>	May be suspected when the cake is brittle and colour grayish.
	<i>Resin</i>	Put it in cold alcohol, which will dissolve the resin, without acting on the wax.
CERA FLAVA.		Is known by the greater softness and unctuousity, and its smell when melted. Turmeric is generally added in this case to obviate the paleness.
<i>Yellow Wax.</i>	<i>Tallow</i>	Melt the wax, the oxide will subside.
		The cake has not its ordinary translucency.
CERA ALBA.	<i>White Lead</i>	
<i>White Wax.</i>	<i>Tallow</i>	
CINCHONA.	<i>This is variously adulterated, but generally with the Carthagena and other inferior barks.</i>	Can only be detected by practice, and examining into the quantity of quinia or cinchonia it contains.
<i>Bark.</i>		
COCCUS.	<i>Pieces of dough formed in moulds, and coloured with cochineal.</i>	Throw it into water, the adulteration will appear.
<i>Cochineal.</i>		
COLOMBA.	The true is distinguishable from the false Colomba by adding to an infusion of the root, a few drops of solution of sulph. iron, which gives to the infusion of the false Colomba a greenish black colour; but produces no change in the other.
		If it does not retain its spherical form when dropped into water, its adulteration may be inferred.
COPAIBA.		Mix one part of strong liquid ammonia of 22°, with three parts of copaiba. If pure, the mixture will, in a few minutes, become transparent; if not, it will remain opaque.
<i>Balsam of Copaiba.</i>	<i>Oil</i>	
	<i>Fibres of smoked Beef</i> ..	Affords an unpleasant odour when thrown on live coals.
CROCUS.	<i>Petals of the Calendula officinalis, and Carthamus Tinctorius.</i> ..	Infuse the specimen in hot water, and the difference will be perceptible.
<i>Saffron.</i>		

MEDICINES.	ADULTERATIONS.	MODE OF DETECTION.
CUBEBA. <i>Cubebs.</i>	<i>Turkey Yellow Berries, or the dried fruit of the Rhamnus Catharticus.</i>	Detected by attentive examination.
CUSPARIÆ CORTEX. <i>Angustura Bark.</i>	<i>False Angustura sometimes sold for it.</i>	The epidermis of the true Cusparia is characterized by being covered with a matter resembling the rust of iron.
GUAIACI RESINA. <i>Resin of Guaiacum.</i>	<i>Common Resin</i>	Detected by the turpentine smell emitted when thrown upon hot coals.
	<i>Manchineel Gum</i>	Add to the tincture a few drops of spirit of nitre, and dilute with water; the guaiacum is precipitated—the adulteration floats in the white striae.
HYDRARGYRI CHLORIDUM MITE. <i>Calomel.</i>	<i>Corrosive Sublimate, and Subnitrate of Bismuth.</i>	A precipitation will be produced by the carbonate of potash, from a solution made by boiling the suspected sample with a small portion of muriate of ammonia in distilled water; or, the presence of deuto-chloride of mercury is indicated, by warming gently a small quantity of calomel in alcohol, filtering and adding to the clear liquor some lime-water, by which a reddish yellow precipitate is afforded. When calomel is rubbed with a fixed alkali, it ought to become intensely black, and not exhibit any orange hue.
HYDRARGYRI OXIDUM RUBRUM. <i>Red Precipitate.</i>	<i>Red Lead</i>	Digest in acetic acid: add sulphuret of ammonia, which will produce a dark coloured precipitate. It should be totally volatilized by heat.
HYDRARGYRI SULPHURETUM RUBRUM	<i>Red Lead</i>	Digest in acetic acid, and add sulphuret of ammonia—a black precipitate will be produced.
HYDRARGYRI SULPHURETUM NIGRUM.	<i>Ivory Black</i>	Throw a suspected portion on hot coals—the residuum will detect the fraud.
IODINE.	<i>Plumbago, Charcoal, and Oxide of Manganese.</i>	The tests of its purity are—that it is perfectly soluble in ether. Heated on a piece of glass or porcelain, it sublimes without residuum.
JALAPÆ RADIX. <i>Jalap Root.</i>	<i>Bryony Root, spurious or false Jalap Root, and Liquorice Root.</i>	Bryony root is of a paler colour, and less compact texture, and does not easily burn at the flame of a candle. Liquorice is detected by the taste.
MAGNESIA.	<i>Lime</i>	Detected by the solution in dilute sulphuric acid affording a precipitate with oxalate of ammonia.
MAGNESIÆ SUBCARBONAS. <i>Carbonate of Magnesia.</i>	<i>Sulphuret of Lime</i>	Gives off when moistened, the smell of sulphuretted hydrogen.
	<i>Chalk</i>	Detected by adding dilute sulphuric acid to the suspected substance, when, if chalk be present, there will be a white insoluble precipitate.
	<i>Gypsum</i>	Boil in distilled water, and test the solution by a barytic and oxalic reagent.
MANNA.	<i>A factitious article, consisting of honey or sugar, mixed with scammony, is sometimes sold for it.</i>	The colour, weight, transparency and taste detect it.
MORPHIA ET EJUS SALES. <i>Morphia and its Salts.</i>	Morphia and its salts, when placed in contact with nitric acid, are coloured red; with persalts of iron, blue. They are perfectly soluble in warm alcohol, and acidulated warm water. When morphia is mixed with narcotina, the adulteration is ascertained by mixing them with sulphuric ether, which dissolves the narcotina, without sensibly affecting the morphia.
MOSCHUS. <i>Musk.</i>	<i>Dried Blood</i>	The bag must not appear to have been opened. This may be suspected, if it emits a fetid smoke when inflamed.
	<i>Asphaltum</i>	Discovered by its melting and running, before it inflames.
	<i>Fine particles of Lead ..</i>	Rub with water. The metallic particles will subside.
OLEA DESTILLATA. <i>Essential Oils.</i>	<i>Fixed Oils</i>	Touch writing paper with it, and hold it before the fire: fixed oil leaves a stain of grease.
	<i>Alcohol</i>	Add water. A milkiness and increase of temperature occurs.
OLEUM RICINI. <i>Castor Oil.</i>	<i>Olive or Almond or Poppy Oil.</i>	Alcohol S. G. .820 will mix with any proportion of castor oil, whilst it dissolves very little of the others.

MEDICINES.	ADULTERATIONS.	MODE OF DETECTION.
OPIMUM.	<i>Extract of Liquorice, Bullets and Stones sometimes in it; Extract of Poppy, of Chelidonium majus; G. Arabic, G. Tragacanth, Linseed Oil, Cow's Dung.</i>	The best opium is covered with leaves and the reddish capsules of a species of <i>Rumex</i> . The inferior kinds have capsules adherent. It is bad when soft and friable, when intensely black or mixed with many impurities, and when sweet. The quantity of morphia affords the best test.
POTASSII IODIDUM. <i>Iodide of Potassium.</i>	<i>Chlorides of Potassium and Sodium, Nitrate of Potassa.</i>	The adulteration is ascertained by precipitating a solution of the salt with nitrate of silver, and treating the precipitate with ammonia, which dissolves the chloride of silver, without acting upon the iodide of this metal.
QUINIE SULPHAS. <i>Sulphate of Quinia.</i>	<i>Mannite.....</i>	Leaves no residue when submitted to calcination: is perfectly soluble in warm alcohol, and in water slightly acidulated with sulphuric acid.
STRYCHNIE ET EJUS SALES. <i>Strychnia and its Salts.</i>	<i>Brucia.....</i>	They are free from brucia when no colour is produced by contact with nitric acid.
ZINCI OXYDUM. <i>Flowers of Zinc.</i>	<i>Chalk.....</i> <i>White Lead.....</i>	Sulphuric acid excites an effervescence. Sulphuric acid forms an insoluble sulphate of lead.

FALTRANCK, Faltranck.

FALX. Anatomists have given this name to several membranous reflections having the shape of a falx or scythe.

FALX CEREBELLI, *Falx minor, Septum Cerebelli seu parvum occipitale, Processus falcatiformis Cerebelli*, (F.) *Faux du cervelet, Septum médian du cervelet*, (Ch.), is a triangular process of the dura mater opposite the internal occipital protuberance. Its base is attached to the middle of the tentorium, and its top or apex bifurcates, to proceed to the sides of the foramen magnum. Its convex surface is towards the cranium, and its concave in the fissure or groove, which separates the two lobes of the cerebellum.

FALX CEREBRI, *F. major, Vertical superior longitudinal process, Mediastinum cerebri, Septum Cerebri*, (F.) *Faux du cerveau, Repli longitudinal de la méninge*, (Ch.), *Processus falcatiformis duræ matris*. The greatest process of the dura mater. It extends from the fore to the hind part of the skull, on the median line; is broad behind and narrow before, and is lodged in the groove which separates the hemispheres from each other—the *interlobular fissure*. At its superior part is situated the longitudinal sinus (*superior*), and at its lower, corresponding to the edge of the scythe, the inferior longitudinal sinus. Its anterior extremity is attached to the crista galli; its posterior is continuous with the tentorium cerebelli, and contains the straight sinus.

FALX MAJOR, *Falx cerebri*—f. Minor, *Falx cerebelli*.

FALX OF THE PERITONEUM, GREAT, *F. peritonei major*, (F.) *Grande faux du péritoine, Faux de la Veine Omphalique, Falx of the umbilical vein*, is a reflection of the peritoneum, which ascends from the umbilicus to the anterior and inferior surface of the liver.

FALCES OF THE PERITONEUM, LESSER, *Falces Peritonei minores*, (F.) *Petites faux du péritoine*, are the lateral ligaments of the liver and the reflections which the peritoneum forms, raised up by the umbilical arteries.

FALX OF THE UMBILICAL VEIN, *Falx*, great, of the Peritoneum.

FAME, Lancet.

FAMELICA FEBRIS, from *fames*, 'hunger.' Fever accompanied with insatiable hunger.—Sylvius.

FAMELICUS, Hungry.

FAMES, Hunger—f. Bovina, Boulimia—f. Canina, Boulimia.

FAMES LUPINA, *Lycorex'is*, (F.) *Faim de Loup*. Authors have described, under this name, a kind of boulimia, or depravation of the digestive function, in which the patient eats voraciously, and passes his food, almost immediately afterwards, *per anum*.

FAMEX, Contusio.

FAMIGERATISSIMUM EMPLASTRUM, from *fama*, 'fame,' and *gero*, 'I wear.' A plaster, extolled in ague, and made of aromatic, irritating substances. It was applied to the wrist.

FAMILIARICA SELLA, Close stool.

FAMILY DISEASES, see Hereditary.

FAMIS, Contusio.

FAMIX, Contusio.

FANCULUM, Anethum.

FANCY MARK, Nævus.

FANFECKLES, Ephelides.

FANG, Radix.

FANON (F.) from (G.) *Fahne*, 'a banner,' 'ensign,' 'standard.' *Ferula, Lectulus stramineus, Thorulus stramineus*. A splint of a particular shape, employed in fractures of the thigh and leg to keep the bones in contact.

The Fanons were divided into *true* and *false*.

The *true* consists of a cylinder of straw, strongly surrounded with a cord or riband, in the centre of which a stick is usually placed to ensure its solidity. The *false* consists of a thick piece of linen, made flat like a compress, and folded at the extremities. It was placed between the fractured limb and the *true fanon*. The *Drap-fanon* is a large piece of common cloth placed between the fractured limb, in which the fanon or lateral splints are rolled.

FANTICKLES, Ephelides.

FANTOM, *Phantom, Phantoma*, from *φάντασμα*, 'a spectre.' (F.) *Phantôme, Fantôme*. This word has two acceptations. It means the spectres and images which the imagination presents to the sick, when asleep or awake; and, also, the figure called, also, *Man'ikin, Man'ikin*, and *Man'akin*, (F.) *Mannequin*, on which surgeons practice the application of bandages, or the accoucheur the manual part of midwifery,—*Phantoma obstetricium*.

Fantom or *Phantom Tumour* is one that forms, in the abdomen for example, simulating organic disease, and occasionally exceedingly deceptive.

FARADISATION, *Faradism*; from *Faraday*, the celebrated physicist. A term proposed by

Duchenne, of Boulogne, for electricity by induction.

FARCIMEN, see Equinia.

FARCINALLIS MEMBRANA SEU TUNICA. Allantois.

FARCIN, see Equinia.

FARCINOMA, Equinia.

FARCTURA, *Fractura*; from *farcire*, *farcum*, 'to stuff.' The operation of introducing medicinal substances into the cavities of animals or of fruits, which have been previously emptied.

FARCTUS, Emphraxia.

FARCY GLANDERS, see Equinia.

FAR'D. Paint.

FARDEAU, Mola.

FARFARA, Tussilago—f. Bechium, Tussilago.

FARIGOLE, Thymus.

FARINA, *Al'phiton*, *Crimmon*, *Al'eton*, *Alen'ton*, *Al'ema*, from *far*, 'corn,' or grain of any kind, of which it is made. *Meal* or *flour*. The powder, obtained by grinding the seeds of the gramineous, leguminous, and cucurbitaceous plants in particular. It is highly nutritious and much used, dietetically as well as medicinally. In the pharmacopœias of London, Edinburgh, and Dublin, *Fari'na* means wheat flour—*Farina Tritici*.

Leath's Alimentary Far'na, or *Homœopathic Farina'ceous Food*, is said to consist principally of wheat flour, slightly baked, and sweetened with sugar, together with potato flour and a very small quantity of Indian corn meal and tapioca.

FARINA AMYGDALARUM, see Amygdala.

FARINA, COMPOUNDED, BASTER'S, is said to consist of wheat flour, sweetened with sugar.

FARINA, NUTRITIOUS, MAIDMAN'S, is said to consist of potato flour, artificially coloured of a pink or rosy hue, the colouring matter being probably rose pink.

FARINA TRITICI, wheaten flour; *F. Secal'is*, Rye flour or meal; *F. Hor'dei*, Barley meal; *F. Avena'cea*, Oat meal, &c. See Amylum.

FARINE RESOLUTIVES, (F.) *Farines Résolutives*. This name was formerly given to a mixture of the farina of four different plants; the lupine, *Lupinus albus*, the *Erum Ervilia*, the *Vicia faba*, and the Barley, *Hordeum distichum*. They were recommended to form cataplasms.

FARINACEOUS, *Farina'ceus*, *Farino'us*, (F.) *Farinacé*, *Farineux*, *Mealy*. Having the appearance or nature of farina. A term given to all articles of food which contain farina. The term *Farinacea* includes all those substances, called *cerealia*, *legumina*, &c., which contain farina, and are employed as nutriment.

Hard's farinaceous food is fine wheat flour, which has been subjected to some heating process. *Braden's farinaceous food* is said to be wheat flour, baked.

In *Pathology*, the epithet *farinaceous*, (F.) *farineux*, is applied to certain eruptions, in which the epidermis exfoliates in small particles, similar to farina.

FARINACEOUS FOOD, BRADEN'S, see Farinaceous—f. Food, Hard's, see Farinaceous—f. Food, homœopathic, see Farinaceous.

FARINACEOUS FOOD, PLUMBE'S, is said to consist principally of bean or pea flour, most probably the former, with a little Tacca arrowroot, some potato flour, and a very little Maranta arrowroot.

FARINACEOUS FOOD, PRINCE ARTHUR'S, for infants and invalids of all ages, is said to consist entirely of wheat flour, slightly baked.

FARINARIUM, Alia.

FARINES RÉSOLUTIVES, *Farinæ resolutives*.

FARINEUX, Farinaceous.

FARINOSUS, Farinaceous.

FARMERY, Hospital.

FARNTICKLES, Ephelides.

FARRIER, Hippiatre.

FARSIGHTEDNESS, Presbytia.

FART, Sax. *fart*, from Teut. *fahren*, 'to go.'

—*fart*, 'a voyage.' (G.) *Farz*, *Bellus*, *Bellus*, *Bilegg'mia*, *Bilegg'mus*, *Bombus*, *Trulla*, *Pocda*, *Physa*, *Physé*, *Flatus*, *Crepitus*, *C. ventris*, (F.) *Pet*. A sonorous or other discharge of wind from behind. A low word, but of respectable parentage.

FARTURA, Fartura.

FARTUS, Emphraxia.

FAS'CIA, from *fascis*, 'a bundle.' *Ligatio*, *Ligatura*, *Alligatura*, *Anadesmos*, *Vincula*, *Spar'ganon*, *Epidesmos*, *Vinctura*. A bandage, gilet, roller, ligature. The aponeurotic expansions of muscles, which bind parts together, are, likewise, termed *Fasciæ*. See *Tania*.

FASCIA APONETROTICA FEMORIS, *Fascia lata aponeurosis*—f. *Capitalis*, *Bonnet d'Hippocrate*, *Capelina*—f. *Capitis*, *Diadema*—f. *Cooperi*, *F. Transversalis*.

FASCIA CRIBRIFORM, *F. Cribrifor'mis*. The sieve-like portion of the fascia lata; so called from its being pierced by numerous openings for the passage of lymphatic vessels.

FASCIA DEXTATA, *F. D. Tarsi'ni*, *Corps godronné* of *Vicq d'Azyr*. A band of gray matter seen beneath the *tania hippecampi* on raising it up, which runs along the inner border of the *cornu ammonis*. It is, as it were, crenated by transverse furrows.

FASCIA DIGITALIS, *Gantlet*—f. *Diopthalmica*, *Binoculus*—f. *Dividens*, *Dividing bandage*—f. *Heliodori*, *T bandage*.

FASCIA ILIACA, *Iliac aponeurosis*. An aponeurosis which proceeds from the tendon of the *psaos minor*, or which arises from the anterior surface of the *psaos magnus*, when the former muscle does not exist. It is attached, externally, to the inner edge of the crest of the ilium, below, and anteriorly—on one side, to the crural arch, sending an expansion to the fascia transversalis; and on the other, continuous with the deep-seated lamina of the fascia lata, which forms the posterior paries of the crural canal. Within and behind, the fascia iliaca is attached to the brim of the pelvis, and is continuous with the aponeurosis, which M. Jules Cloquet has called *Pelvicum*. The iliac aponeurosis covers the iliac and *psaos* muscles, which it separates from the peritoneum.

FASCIA INTENDITIFORMIS, a funnel-shaped offset from the fascia transversalis, which is prolonged as a sheath upon the *vas deferens* and spermatic vessels as they pass out of the abdomen.

FASCIA INGUINALIS, *Spica*.

FASCIA LATA. A name given by anatomists to an aponeurosis, and to a muscle.

FASCIA LATA APONETROSIS, *F. aponeuroticæ fem'oris*, *Vagi'na fem'oris*, *Crural* or *Fem'ral Aponeurosis*, is the most extensive in the body, and envelopes all the muscles of the thigh. Above, it is attached to the outer edge of the ilia; before, it arises from the crural arch by two distinct laminae, separated by the femoral vessels, and becoming confounded a little below the part where the great vena saphæna opens into the crural vein. Of these two laminae, the one is more anterior and thicker than the other, and may be considered as a prolongation of the aponeurosis of the external oblique. It is intimately united to *Poupart's* ligament. The other, which is thinner, is behind, and deeper seated, and, after its union with the former, proceeds to be inserted into the pubis. Inferiorly, the fascia lata becomes confounded with the tendon of the triceps, and is attached to the external tuberosity

of the tibia. The use of the fascia lata, like that of other aponeuroses, is to strengthen the action of the muscles, &c.

FASCIA LATA MUSCLE, *Tensor vaginæ femoralis*, *Fascia lita*, *Membrano-sus*, *Musculus aponeurosis seu fascia lata seu femoralis membrano-sus*, (F.) *Ilio-aponeurosi-femoralis*, *Ilio-aponeuroti-femoralis*—(Ch.), *Tenseur de l'aponeurose fémorale*. A muscle, situate at the upper and outer part of the thigh. It arises, *above*, from the outer part of the anterior and superior spine of the ilium; and is inserted, *below*, between the two laminae of the fascia lata, which it stretches and raises when it contracts.

FASCIA, OBTURATOR, see *Pelvic aponeuroses*—f. *Pelvic*, *Internal*, see *Pelvic aponeuroses*—f. *Pelvic*, *Lateral*, see *Pelvic aponeuroses*—f. *Pelvic*, *Superior*, see *Pelvic aponeuroses*.

FASCIA PROPRIA. A layer of areolar tissue derived from the sheath of the femoral vessels,—or according to some from the cribriform fascia. It is one of the coverings of femoral hernia, and is generally pretty dense about the neck of the hernia; but thin or even wanting on its fundus.

FASCIA REPENS, *Spica*—f. *Scapularis*, *Scapularis*—f. *Sculpti*, *Bandage of separate strips*—f. *Semicircularis*, *Tænia semicircularis*—f. *Spiralis*, *Ascia*—f. *Stellata*, *Stella*—f. *Submuscular*, see *Vaginal*, (of the eye.)

FASCIA, SUPERITONEAL, *Superitoneal aponeurosis*. A thin tendinous layer on the outer surface of the peritoneum.

FASCIA SUPERFICIALIS, *Superficial aponeurosis of the abdomen and thigh*, (F.) *Aponérose superficielle de l'abdomen et de la cuisse*. A very thin aponeurosis, which covers the muscles and aponeuroses of the abdomen; passes before the crural arch, to which it adheres with some degree of force; sends a membranous sheath, which surrounds the spermatic cord; and is continuous with the dartos, which it assists in forming. The fascia superficialis presents, beneath the crural arch, very distinct fibres, whose direction is parallel to the fold of the thigh. It is applied over the fascia lata aponeurosis, and is attached, internally, to the ascending ramus of the ischium, near the root of the corpus cavernosum. Before the descent of the testicle from the abdomen, the fascia superficialis is very manifestly continuous with the *Gubernaculum testis*.

FASCIA TFORMIS, *T. bandage*—f. *Tortilis*, *Tourniquet*.

FASCIA TRANSVERSALIS, *F. Cooperi*. An aponeurosis, which separates the transversalis muscle from the peritoneum in the inguinal region. It arises above the posterior edge of the crural arch, where it seems to be continuous with the aponeurosis of the greater oblique muscle. *Above*, it is lost in the areolar tissue at the internal surface of the transversalis abdominis. *Within*, it is continuous with the outer edge of the tendon of the rectus muscle and Gimbernat's ligament; *below*, it is continuous with the aponeurosis of the greater oblique, and receives an expansion from the *Fascia lita*. Towards its middle and a little above the crural arch, the fascia transversalis has the wide orifice of a canal, which is occupied, in the female, by the round ligament of the uterus; and, in man, furnishes an expansion, that serves as a sheath to the spermatic vessels.

FASCIÆ, *Swathing clothes*—f. *Coli*, see *Colon*—f. *Ligamentosa Coli*, see *Colon*.

FASCIALIS, *Fascia lata muscle*, *Sartorius*—f. *Longus*, *Sartorius*—f. *Sutorius*, *Sartorius*.

FASCIARUM APPLICATIO, *Deligation*.

FASCIATIO, *Deligation*, see *Bandage*—f. *Cucullata*, *Couvrechef*.

FASCICULI CORTICALES, *Ferrein*, *pyramids of*—f. *Graciles*, see *Corpora restiformia*.

FASCICULI INNOMINATI. Two large bundles of fibres in the interior of the medulla oblongata, behind the corpora olivaria, and more or less apparent between those bodies and the corpora restiformia. They ascend, and become apparent in the fourth ventricle, under the name *Fasciculi seu Processus teretes*.

FASCICULI, MEDIAN POSTERIOR, OF THE MEDULLA OBLONGATA, *Funiculi graciles*—f. *Musculorum*, see *Muscular fibre*—f. *Pyramidales*, *Ferrein*, *Pyramids of*—f. *Teretes*, *Processus teretes*, see *Fasciculi innominati*—f. *Teretes cordis*, *Columnæ carnea*.

FASCICULUS, *Phacellus*, *Phacelus*, *Fascicle*, diminutive of *fascia*, 'a bundle'; *Desmedion*, 'a small bundle'. In *Anatomy*, it is employed in this sense; as "*a fasciculus of fibres*." (F.) *Fascicule*, *Faisceau ou Petit Faisceau*, *Troussseau*. In *Pharmacy*, it means *manipulus*, *Chetivolethes*, *χαρτοληθης*, 'a handful'. Musa Brasavolus says,—as much as can be held in two fingers.

FASCICULUS CUNEATUS, *Reinforcement*, *fasciculus of*—f. of *Reinforcement*, *Reinforcement*, F. of.

FASCICULUS UNCINATUS. A white bundle, seen on the lateral aspect of the cerebral hemispheres, passing across the bottom of the fissure of Sylvius, and connecting the anterior with the middle and posterior lobes. The fibres of this fasciculus expand at each extremity, and the superficial portions of them curve or *hook* sharply between the contiguous parts of the anterior and middle lobes, whence the epithet '*uncinatus*.'

FASCINOSUS, *Membrosus*.

FASCINUM, *Penis*.

FASCIOLA, *Bandelette*—f. *Cinerea*, *Tuberculum cinereum*—f. *Hepatica*, *Distoma hepaticum*—f. *Humana*, *Distoma hepaticum*—f. *Lanceolata*, *Distoma hepaticum*.

FASELUS, *Phaseolus vulgaris*.

FASEOLE, *Phaseolus*.

FASTIDIUM, abridged from *fastis tedium*. *Fastidiousness*, *Squeamishness*, or the condition of a stomach that is really affected with nausea.

FASTIDIUM CIBI, *Asitia*, *Disgust*—f. *Potdis*, *Loathing of drink*.

FASTIGIUM, *Acme*.

FASTING, from Sax. *fæstan*, *Limo'sis experta*, *protracta*, *Anorexia mirabilis*, *Inedia*, *Jejunium*. Loss or want of appetite, without any other apparent affection of the stomach; so that the system can sustain almost total abstinence for a long time without faintness. Some wonderful cases of this kind are on record. See *Abstinence*.

FAT, *Pingue'do*, *Pim'elē*, *Piar*, *Pie'ron*, *Lipos*, *Stear*, *Adeps*, *Sevum*, *Sebum*, *Corpus adiposum*, *Axungia*, *Fat*, (F.) *Graisse*: from (G.) *Fett*, (Sax.) *fat*. A soft, white, animal substance; inodorous; insipid; oily; inflammable, easy to melt; spoiling in the air, and becoming rancid by union with oxygen: almost insoluble in alcohol; insoluble in water; soluble in fixed oils. Fat is formed of the immediate principles, *stearin*, *margarin*, and *olein* or *clain*, all of which are regarded as salts composed of stearic, margaric and oleic acids, and a common base, to which, from its sweetish taste, the name *Glycerin* *Glycerina*, *Glycerinum*, (F.) *Glycérine* has been given. To these are, almost always, joined an odorous and a colouring principle. **GLYCERIN**, *Glycerina*, has been introduced into the last edition of the *Pharmacopœia* of the U.S. (1851), in which it is directed to be prepared as follows:—*Lead plaster*, recently prepared and yet fluid; *boiling water*, of each a gallon; mix: stir briskly for 15 minutes; allow it to cool and pour off the liquid. Evaporate until it has the s. g. 1.15, and pass slowly through it a current of sulpho-hydric acid

until a black precipitate is no longer thrown down. Filter and boil until the sulpho-hydric acid is driven off, and evaporate the liquid until it ceases to lose weight. Glycerin is a colourless or straw-coloured syrupy fluid; s. g. 1.25. It is soluble in water and in alcohol, but not in ether. It is used in the form of lotion, composed of half an ounce to ten fluidounces of water, in cutaneous diseases, as psoriasis, pityriasis, lepra and ichthyosis. It forms a kind of varnish, and might be useful in cases of burns.

Fat is found in a number of animal tissues, and is very abundant in the neighborhood of the kidneys and in the epiploon. It is generally fluid in the cetacea; soft, and of a strong smell in the carnivora; solid, and inodorous in the ruminating animal; white, and abundant in young animals; and yellowish in old. It generally forms about a twentieth part of the weight of the human body. The fat, considered physiologically, has, for its function, to protect the organs; maintain their temperature; and to serve for nutrition in case of need; as is observed in torpid animals.

FAT, Corpulent—f. Cells, Fatty vesicles.

FAT, MACKAW, see *Cocos butyracea*.

FATIGATIO, Copos.

FATIGUE, Copos.

FATTY, *Adiposus*, *Adipatus*, *Pimel'icus*, *Pimelo'des*, *Liparo'des*, *Lip'arus*, Sax. *fætt*; past participle of *fætan*, to feed, *Ad'ipous*, *Pinguid*, *Pinguid'inous*. Relating to fat. Resembling or containing fat. The cellular membrane has been called *fatty* or *adipous*; from an opinion that, in its areolæ, the fat is deposited. The areolar membrane, however, merely lodges, between its lamellæ and filaments, the vesicles in which the fat is contained.

FATTY LIGAMENT, *Ad'ipous ligament*. This name has been given to a reflection of the synovial membrane of the knee joint, which passes from the ligamentum patellæ towards the cavity that separates the condyles of the femur.

FATTY MEMBRANE, *Ad'ipous membrane*, *Adipous tissue*. The subcutaneous areolar tissue, or that containing the fatty or adipous vesicles.

FATTY VESICLES, *Sacculi adiposi*, *Ad'ipous vesicles*, *Fat cells*. This name is given to small bursæ or membranous vesicles which enclose the fat, and are found situate in the areolæ of the areolar tissue. These vesicles vary much in size. Generally, they are round and globular; and, in certain subjects, receive vessels which are very apparent. They form so many small sacs without apertures, in the interior of which are filaments arranged like septa. In fatty subjects, the adipous vesicles are very perceptible, being attached to the areolar tissue and neighbouring parts by a vascular pedicle. Raspail affirms that there is the most striking analogy between the nature of the adipose granules and that of the amylaceous grains.

FATTY VESSELS, *Ad'ipous vessels*. The vessels connected with the fat. Some anatomists have called *Adipous canals*, (F.) *Conduits adipæus*, the vessels to which they attribute the secretion of fat.

FATTY DEGENERATION OF THE LIVER, *Adiposis hepatica*—f. Liver, *Adiposis hepatica*.

FATUISME, *Fatuitas*.

FATUITAS, *Moro'sis*, *Stultit'ia*, *Stupor mentis*, *Amen'tia*; (F.) *Fatuisme*, *Fatuité*, from *fatuus*, 'foolish.' Mental imbecility. Idiotism. Dementia. One affected with fatuity is said to be *fat'uous*, (Sc.) *Daupit*.

FATUITAS ALPICOLARUM, Cretinism—f. *Convallinus*, Cretinism.

FATUOUS, see *Fatuity*.

FAUCES, Isthmus, Pharynx, Throat.

FAUCETTE VOICE, see *Voice*.

FAUNORUM LUDIB'RIA. *The sports of*

the Fauni. Some authors have called thus the incubus; others, epilepsy.

FAUQUIER SPRINGS, see *Virginia*, Mineral Waters of.

FAUSSE, False—f. *Conception*, *Conception*, false.

FAUSSE COUCHE (F.), *Vanum partu'cium*. Some authors have used this term for the expulsion of different bodies constituting false conceptions; such as moles, hydatids, clots of blood, &c. Most accoucheurs use the term synonymously with abortion.

FAUSSE GROSSESSE, *Pregnancy*, false—f. *Membrane*, *Membrane*, false—f. *Pleur'cite*, *Pleurisy*, false—f. *Pneumonie*, *Peripneumonitis* notha.

FAUSSE POSITION (F.), *False position*. The French use this term, in vulgar language, to indicate any attitude in which torpor, tingling, and loss of power over the motion of a part, are produced by too strong contraction or painful compression.

FAUSSE RHABARBE, *Thaliectron*—f. *Route*, False passage—f. *Variole*, *Varicella*.

FAUSTI'NI PASTIL'LI, *Faustinus's Lozenges*. These were once celebrated. They were composed of burnt paper, quicklime, oxide of arsenic, sandarach, lentils, &c.

FALX, False—f. *du Cerveau*, *Falx cerebri*—f. *du Cervelet*, *Falx cerebelli*—f. *Germe*, *Conception*, false—f. *Grande du péritoine*, *Falx*, great, of the peritoneum—f. *Petite du péritoine*, *Falx*, lesser, of the peritoneum—f. *de la Veine ombilicale*, *Falx*, great, of the peritoneum.

FAVEUX, *Favosus*.

FAVIFORMIS, *Favosus*.

FAVOSUS. Similar to a honeycomb. *Faviformis*, *Favosus*, from *favus*, *cerion*, 'a honeycomb.' (F.) *Faveux*. An epithet given to a species of porrigo. *Cerion*, *Kerion*, *Favus*; means also a state of ulceration, resembling a honeycomb.

FAVULUS; diminutive of *favus*, 'a honeycomb.' *Favuli* is used by Dr. Morton, of Philadelphia, for the honeycomb-like depressions in the lining membrane of the stomach—the *stomach-cells* of Messrs. Todd and Bowman.

FAVUS, *Porrigo*, *Porrigo favosa*, see *Favosus*—f. *Confertus*, *Porrigo scutulata*—f. *Dispersus*, *Porrigo lupinosa*—f. *Disseminé*, *Porrigo lupinosa*—f. *en Groupes*, *Porrigo scutulata*.

FAXWAX, see *Nucha*.

FEABERRY, *Ribes uva crispa*.

FEATHERFEW, *Matricaria*.

FEBRICITANS, *Feb'riens*, *Emp'yros*, (F.) *Fébricitant*. One attacked with fever; from *fébricitare*, 'to have a fever.'

FEBRICITATIO, *Feverishness*.

FEBRICOSUS, *Feverish*.

FEBRIC'ULA, *Fe'veret*, (F.) *Fébricule*. Diminutive of *febris*, 'fever.' A term employed to express a slight degree of fever. *Ephemera*.

FEBRICULOSITY, *Feverishness*.

FEBRIENS, *Feb'ricitans*, *Feverish*.

FEBRIFACIENT, see *Feverish*.

FEBRIFEROUS, *Feb'rifer*, from *febris*, 'fever,' and *fero*, 'I carry.' Fever-bearing, as a *febriferous* locality.

FEBRIFIC, see *Feverish*.

FEBRIFUGA, *Leucopyr'eticus*, *Lecopyr'etus*, *Pyret'icus*, *Aleucopyr'eticus*, *Antifeb'r'is*, *Antipyret'ic*, *Feb'rifu'gus*, from *febris*, 'a fever,' and *fugare*, 'to drive away.' A medicine which possesses the property of abating or driving away fever.

FEBRIFUGUM LIGNUM, *Quassia*.

FE'BRILE, *Feb'r'is*, *Pyret'icus*. Relating to fever, as *febrile* movement, *febrile* pulse, &c.

FEBRIS, Fever—f. *Acnastic*, *Synocha*—f.

Acuta, Synocha.—f. Acuta continua, Synocha.—f. Adeno-meningea, Fever, adeno-meningeal.—f. Adeno-nervosa, Plague.—f. Africana, Fever, African.—f. Agrypnodes, see Agrypnodes and Agrypnos.—f. Albu, Chlorosis.—f. Algidæ, see Algidus.—f. Amatoria, Chlorosis, Hæctic fever.—f. Americana, Fever, yellow.—f. Amphemera, Quotidian.—f. Amphimerina hæctica, Hæctic fever.—f. Amphimerina laticæ, Laticæ (febris).—f. Ampullosa, Pemphigus.—f. Anabatica, Continued fever.—f. Angiotenica, Synocha.—f. Anginosa, Angina.—f. Annua, see Annual diseases.—f. Anomala, Fever, anomalous.—f. Aphonica, Fever, aphonic.—f. Aphthosa, Aphtha.—f. Apoplectica, Fever, apoplectic.—f. Ardens, Synocha.—f. Arte promota, Fever, artificial.—f. Arthritica, Gout.—f. Asodes, Fever, bilious, see Asodes.—f. Assiduæ, Continued fever.—f. Asthenica, Fever, asthenic, Typhus.—f. Asthmatica, Fever, asthmatic.—f. Ataxo-ady-namica, Fever, ataxo-ady-namic.—f. Azodes, see Asodes.—f. Biliosa, Fever, bilious.—f. Bullosa, Pemphigus.—f. Cardialgia, Fever, cardialgic.—f. Catarrhalis, Catarrh.—f. Catarrhalis epidemica, Influenza.—f. Cautatodes, Synocha.—f. Causodes, Synocha.—f. Cephalalgica, Fever, cephalalgic.—f. Cephalica, Fever, cephalic.—f. Choleryctica, Fever, bilious.—f. Cholericæ, Fever, bilious, Fever, choleric.—f. Chronica, Fever, chronic.—f. Coäl-terna, see Coalternæ febres.—f. Colliquativa, Fever, colliquative.—f. Comatodes, Fever, apoplectic.—f. Communicans, see Subintrantes F.—f. Confusa, see Confusæ febres.—f. Continens, Typhus.—f. Continens non putrida, Synocha.—f. Continens putrida, Typhus.—f. Continua inflammatoria, Empresma.—f. Continua putrida, Synochus.—f. Continua putrida icterodes Caroliniensis, Fever, yellow.—f. Continua non putris, Synocha.—f. Continua sanguinea, Synocha.—f. Convulsiva, Fever, convulsive.—f. Crymodes, see Crymodes, and Algid fever.—f. Culicularis, Miliary fever.—f. cum Delirio, Fever, delirious.—f. Depuratoria, Fever, depuratory.—f. Deurens, Synocha.—f. Diaphoretica, Fever, diaphoretic.—f. Diaria, Ephemera.—f. Duodecimana, Fever, duodecimane.—f. Dysenterica, Dysentery.—f. Elodes, see Elodes.—f. Elodes icterodes, Fever, yellow.—f. Enterica, see Typhus.—f. Enteromesenterica, Entero-mesenteric.—f. Epacnastica, see Epacnasticos.—f. Ephemera, Ephemera.—f. Epidemica cum angina, Cynanche maligna.—f. Epileptica, Fever, epileptic.—f. Erotica, Fever, erotic.—f. Erratica, Fever, anomalous, Fever, erratic, see Erraticæ and Planetes.—f. Erroneæ, see Planetes.—f. Erysipelacea, Erysipelas.—f. Erysipelatosa, Erysipelas.—f. Esserosa, Miliary fever.—f. Exquisita, Fever, regular.—f. Exanthematica articularis, Dengue.—f. Famelica, see Famelica Febris.—f. Flava, Fever, yellow.—f. Flava Americanorum, Fever, yellow.—f. Gangræ-nodes, Fever, gangrenous.—f. Gastrica, Fever, bilious, Fever, gastric.—f. Gastrico-biliosa, Fever, gastric.—f. Gastro-ady-namica, Fever, gastro-ady-namic.—f. Hæmoptica, Fever, hæmoptoic.—f. Hebdomadana, Octana.—f. Hæctica, Hæctic fever.—f. Hæctica infantum, Tabes mesenterica.—f. Hæctica maligna nervosa, Typhus mitior.—f. Hemeresia, Quotidian.—f. Hemitritæ, Hemitritæ.—f. Hepatica, Fever, bilious.—f. Hepatica inflammatoria, Hepatitis.—f. Horrida, see Algidus.—f. Humoralis, Fever, humoral.—f. Hungarica, Fever, Hungaric.—f. Hydrocephalica, Hydrocephalus internus.—f. Hydrophobica, Fever, hydrophobic.—f. Hysteretica, see Postpositio.—f. Hysterica, Fever, hysteric.—f. Ictericæ, Fever, icteric.—f. Iliaca inflammatoria, Enteritis.—f. Infantum remittens, Fever, infantile remittent.—f. Inflammatoria, Synocha.—f. Intensis, Epitasis.—f. Intermittens, Intermittent fever.—f. Intermittens cephalica larvata, Cephalalgia periodica.—f. Intestinalis ulcerosa, see Typhus—

f. Intestinorum, Enteritis.—f. Irregularis, Fever, anomalous.—f. Lactea, Fever, milk.—f. Larvata, Fever, masked.—f. Lenta, Fever, infantile remittent, Hæctic fever, Synochus.—f. Lenta nervosa, Typhus mitior, Fever, nervous.—f. Lenticularis, Miliary fever.—f. Lethargica, Fever, apoplectic.—f. Lochialis, Fever, lochial.—f. Lygmodes, Fever, singultous.—f. Lyngodes, Fever, singultous.—f. Maligna, Fever, malignant.—f. Maligna biliosa America, Fever, yellow.—f. Maligna cum Sopore, Typhus.—f. Maligna flava Indiæ occidentalis, Fever, yellow.—f. Marasmodes, Hæctic fever, Marasmpyrra.—f. Meningo-gastricus, Fever, gastric.—f. Mesenterica, Fever, adeno-meningeal, Fever, mesenteric.—f. Methemerina, Quotidian.—f. Miliaris, Miliary fever.—f. Minuta, Fever, syncopal.—f. Morbillosa, Rubella.—f. Mucosa, Fever, adeno-meningeal.—f. Mucosa verminosa, Fever, infantile remittent.—f. Nautica pestilentialis, Typhus gravior.—f. Nephritica, Fever, nephritic.—f. Nervosa, Fever, nervous.—f. Nervosa enterica, see Typhus.—f. Nervosa epidemica, Typhus.—f. Nervosa exanthematica, Typhus.—f. Nervosa gastrica, see Typhus.—f. Nervosa mesenterica, see Typhus.—f. Nervosa petechialis, Typhus.—f. Neurodes, Fever, nervous.—f. Nocturnus, see Nocturnal.—f. Nonana, Fever, nonane.—f. Nosocomium, Typhus gravior.—f. Nycterinus, see Nocturnal.—f. Octana, Fever, octane.—f. Oscitans, Oscitant fever.—f. Paludosa, see Elodes.—f. Panonica, Fever, Hungary.—f. Pemphigodes, Pemphigus.—f. Pemphingodes, Pemphigus.—f. Pempta, Quintan.—f. Periodica, Fever, periodic.—f. Perniciosa, Fever, pernicious.—f. Pestilens, Plague.—f. Pestilens maligna, Typhus gravior.—f. Pestilentialis, Fever, pestilential.—f. Pestilentialis Europæ, Typhus gravior.—f. Petechialis, Typhus gravior.—f. Phthisica, Hæctic fever.—f. Planetes, see Planetes.—f. Pleuritica, Pleuritis.—f. Podagrica, Gout.—f. Polycholica, Fever, bilious.—f. Pneumonica, Fever, pneumonic, Pneumonia.—f. Puerperalis biliosa, Metrocholis.—f. Puerperarum, Puerperal fever.—f. Puncticularis, Miliary fever, Typhus gravior.—f. Puerperalis, Fever, puerperal.—f. Purpurata rubra et alba miliaris, Miliary fever.—f. Puerulenta, Fever, purulent.—f. Putrida, Typhus gravior.—f. Putrida nervosa, Typhus mitior.—f. Quartana, Quartan.—f. Quæ-quæ, see Algidus.—f. Quinta, Quintan.—f. Quintana, Fever, quintan, Quintan.—f. Quotidiana, Fever, quotidian, Quotidian.—f. Regularis, Fever, regular.—f. Remittens, Remittent fever.—f. Remittens infantum, Fever, infantile remittent.—f. Rheumatica inflammatoria, Rheumatism, acute.—f. Rubra, Scarlatina.—f. Rubra pruriginosa, Urticaria.—f. Sanguinea, Synocha.—f. Sapropyya, Typhus gravior.—f. Scarlatinosa, Scarlatina.—f. Scorbutica, Fever, scorbutic.—f. Semitertiana, Hemitritæ.—f. Septana, Fever, septan.—f. Sesquialtera, Hemitritæ.—f. Sextana, Fever, sextan.—f. Simplex, Ephemera, Fever, simple.—f. Singultosa, Fever, singultous.—f. Soporosa, Fever, apoplectic.—f. Stercoralis, Fever, stercoral.—f. Sthenica, Synocha.—f. Stomachica inflammatoria, Gastritis.—f. Subintrans, Fever, subintrant.—f. Sudatoria, Fever, diaphoretic, Hydropyretus, Sudor anglicus.—f. Syncopalis, Fever, syncopal.—f. Syphilitica, Fever, syphilitic.—f. Tabida, Fever, colliquative, Hæctic fever.—f. Tertia, Fever, tertian, Tertian fever.—f. Tonica, Synocha.—f. Topica, Neuralgia, facial.—f. Toxicæ, Fever, yellow.—f. Tragica, Fever, tragic.—f. Traumatica, Fever, traumatic.—f. Tropica, Fever, yellow.—f. Typhodes, Typhus.—f. Urticata, Urticaria.—f. Uterina, Metritis.—f. Vaga, Fever, anomalous, see Planetes.—f. Variolosa, Variola.—f. Verminosa, Fever, infantile remittent, F. verminous, Helminthopyrra.—f. Vernalis, Fever, vernal.—f. Vesicularis, Miliary Fever, Pemphigus.—f. Virginum, Chlorosis.

FEBRUA, from *februus*, 'I purge.' In ancient mythology, a goddess who presided over menstruation.

FECAL, Stercoraceous.

FÆCES, Fæces.

FÉCONDATION, Fecundation.

FÉCONDITÉ, Fecundity.

FECULA, *Fæcula*, diminutive of *fæx*, *fæcis*, 'lee.' An immediate principle of vegetables, composed of hydrogen, oxygen, and carbon. It exists in several plants, and has different names, according to that which furnishes it. (F.) *Fécule*, *Amygdol.* When extracted from wheat or barley, it is called *starch*, *Amylum*. When from *Cycas circinalis*, *Sago*:—from *Orchis mol'ris*, *Salep*. We say, also, *Fecula* of the *Potato*, *Bryony*, *Arum*, *Manioc*, &c.

FECULA, GREEN. This name is given to a green, solid matter, of variable character, which is believed to be resinous, and which renders turbid several kinds of juices, extracted from vegetables. It is also called *Chlorophyll*.

FECULA AMYLACEA, *Amylum*—f. *Marantæ*, *Arrow-root*—f. *Tapioca*, see *Jatropha manihot*.

FÉCULE, *Fecula*.

FECULENCE, *Fæculentia*, *Fæcositas*, *Fæx*, plural *Fæces*, *Lemma*, *Lee*, *Depos't*, *Dregs*. In *Pharmacy*, feculent, aluminous, or other substances, which are deposited from turbid fluids.

FECULENT, *Fæculentus*, *Hypodrygus*, *Trygo'des*. 'Foul, dreggy, excrementitious;' as a feculent fluid, feculent evacuations, &c.

FECUND, *Fecundus*, *Fæcundus*, (F.) *Fécond*. Same etymon as the next. Fruitful, prolific.

FECUNDATION, *Fæcundatio*, *Impregnation*, *Impregnatio*, *Ingravidatio*, *Prægnatio*, *Gravidatio*, *Prægnatus*, *Fruetificatio*, *Cyc'sis*, *Encyc'sis*, *Encymo'sis*, *Procreatio*, (F.) *Fécondation*. The act by which, in organized beings, the material furnished by the generative organs of the female, unites with that prepared by those of the male, so that a new being results.

FECUNDITY, *Eutoc'ia*, *Fæcunditas*, *Productivitas*, (F.) *Fécondité*. The faculty of reproduction, possessed by organized bodies.

It has been estimated that, throughout a country, taking one marriage with another, not more than 4 children are the result; and in towns only 35 children to 10 marriages.

FEE, Psychorages, Sostrum.

FEET, BURNING OF THE. A singular cachectic disease, described by Mr. Malcolmson as occurring in India, the prominent symptom of which was a sense of burning in the feet.

FÉGARITE, Cancer aquaticus, Stomatitis, pseudo-membranous.

FEIGNED DISEASES, *Morbi dissimulati* seu *simulati* seu *celati* seu *institi* seu *pseudalei*, *Sim'ulated* or *Pretend'ed diseases*, (F.) *Maladies dissimulées* ou *simulées* ou *feintes* ou *supposées*. The tricks employed, by impostors, to induce a belief that they are attacked with diseases when they are not. These are generally assumed by beggars to obtain alms; by criminals to escape punishment; and by soldiers to be exempt from duty.

The following table exhibits the chief feigned diseases, with the means of detection:

A TABLE OF FEIGNED, PRETENDED, SIMULATED, OR EXCITED DISEASES OR DISQUALIFICATIONS.

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
1. ABORTION.	By staining the clothes and body with borrowed blood.	
2. ABSTINENCE.	By constant and minute attention.
3. AMAUROTIC BLINDNESS.	By applying the extract of belladonna or datura stramonium to the eye.	Amaurosis is characterized by dilated pupil. Where these substances have been applied, the effects will go off in ten days or a fortnight.
4. APOPLEXY.	By falling down as if deprived of sensation and consciousness.	By powerful stimulants; an electric shock; application of hot water, sternutatories, actual cautery, &c.
5. CACHEXIA, ANÆMIA, AND DEBILITY.	Using substances to make the face appear pale and livid. Indulging freely in wine, and privation of sleep prior to examination.	By examining if the pulse be strong, and the skin hot, and whether there be loss of appetite or of strength, or swelling of the limbs.
6. EXCRETION OF CALCULI.	Putting sand, pebbles, &c., into the urine.	By the aid of chymistry. We are acquainted with the chymical composition of urinary calculi.
7. CANCEROUS ULCER.	By gluing on a portion of a spleen with the smooth side to the skin, leaving on the outside the appearance of an ulcerated surface.	By noticing whether there be signs of cachexia, and by attentive examination of the part.
8. CATALEPSY.	By seeming to be suddenly motionless, the joints remaining flexible, and external objects making no impression.	By powerful stimulants, as recommended under apoplexy. Letting fall a drop of boiling water on the back. Proposing to use the actual cautery, and seeing whether the pulse rises.
9. CHOREA.	By assuming the convulsive motions of a part which characterize chorea.	By examining the patient whilst he may imagine himself unobserved, and seeing whether the convulsive motions go on. By anæsthetics. (?)
10. CONTRACTION OF JOINTS IN GENERAL.	Mode of discrimination sometimes so obscure as to deceive the most practised and attentive.

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
11. CONTRACTION OF THE FINGERS.	Introduce a cord between the fingers and the palm of the hand, and gradually apply weights so as to expand the fingers. Confine him so that he cannot obtain his food without using his clenched hand.
12. CONVULSIONS.	When feigned, they do not present the rigidity of muscles or the rapidity of action which characterize the real. The mode of detection must be the same as in epilepsy.
13. OPAKE CORNEA.	Produced by the application of a strong acid, by acrid powders, as quicklime, &c. Some articles of diet will bring on <i>urticaria</i> or <i>nettle-rash</i> , in particular individuals, as shell-fish, bitter almonds, &c. By acrids, acids, or any irritants applied to the surface. An ointment of tartarized antimony causes a painful pustular eruption. See Porriego, in this list.	The existence of the opacity can be detected by attentive observation.
14. CUTANEOUS DISEASES.		By careful examination on the part of practitioner and nurse.
15. DEAF-DUMBNESS.	The really deaf and dumb acquire an expression of countenance and gestures which it is difficult to assume.
16. DEAFNESS.	It may be assumed or excited by putting a pea in the ear, or by inserting irritants, so as to induce inflammation and temporary loss of function.	Make a noise when not expected, and see if the countenance varies or pulse rises. Put to sleep by opium, and then fire a pistol close to the ear, when he may be thrown off his guard. Examine the ear to see if any trick has been played there.
17. DEATH.	Some persons possess the power of suspending or moderating the action of the heart.	If suspected, the plan recommended under apoplexy will be found most efficacious.
18. DEBILITY, see CACHEXIA.		
19. DEFORMITY.	Examine the part and its articulation, naked, and compare it with the opposite.
20. DELIVERY.	After enlargement produced artificially, a subsidence of the tumefaction; the parts being moistened by borrowed blood, and the child of another substituted as the female's own.	Can only be positively detected by examination <i>per vaginam</i> . Soon after delivery, the vagina will be relaxed and the lochial discharge be flowing in greater abundance, the shorter the time that may have elapsed since delivery.
21. DIARRHŒA.	Said to have been caused by a mixture of vinegar and burnt cork. (?) May be occasioned by the use of any of the purgative roots, &c.	When diarrhœa is feigned by the lower classes, inspect the linen; if clean, the bowels are probably not much out of order. Let every individual have a close stool of his own; and inspect the evacuations, taking care that one suffering under the disease does not lend his evacuations to another.
22. DROPSY.	May be feigned, like pregnancy, by wearing pads. The anasarous condition of the lower limbs has been caused by applying a ligature round them. By inflating the cellular membrane of the abdomen.	Can be detected by attentive examination. There will be a want of that leucophlegmatic habit which accompanies and characterizes dropsy.
23. DYSENTERY.	May be feigned, like diarrhœa, by adding a little blood to the evacuations, or by introducing a soap or some more irritating suppository.	Same rules as under diarrhœa.

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
24. EPILEPSY.	The foaming of the mouth has been produced by keeping a piece of soap in it.	Sensation in epilepsy is totally abolished. If any remain, disease probably feigned. Incontractility of pupil, which occurs in epilepsy, cannot be feigned. Same means to be used as in feigned apoplexy.
25. FEVER.	By various stimulants, as wine, brandy, pepper; swallowing a small quantity of tobacco, or introducing it into the anus. Flour or chalk used to whiten the tongue. Redness of skin, caused by friction with a hard brush.	This deceit is generally developed by a day or two's examination. Where flour or chalk has been used to whiten the tongue, the line of demarcation between the whitened part and the clean, healthy margin of the tongue, is too well marked to escape observation.
26. FISTULA IN ANO.	By making an incision near the verge of the anus, and introducing into it an acrid tent, such as the root of white hellebore, &c.	By careful examination.
27. FRACTURES.	There is generally nothing but the man's own testimony. He complains of pain in the part; if fracture of the skull be feigned, he states, perhaps, that he becomes deranged on tasting liquor.	By attentive examination.
28. HÆMATEMESIS.	By drinking the blood of some animal, or using some coloured liquid, and then throwing it up.	By cutting off the supply of the fluid and careful examination.
29. HÆMOPTYSIS.	By secreting bullock's blood for the purpose of colouring the saliva; making small incisions in the mouth; using bole armeniac or paint of vermillion.	Blood from the lungs is frothy and light-coloured. Mouth and fauces must be carefully inspected, and the individual be observed.
30. HÆMORRHOIDS.	By introducing bladders of rats or of small fish partly into the rectum. The linen has also been stained with borrowed blood.	The means are obvious.
31. HEART, DISEASES OF, see PALPITATION.		
32. HEPATITIS.	Unless the person be a well-educated impostor, acute inflammation of the liver will be detected by the absence of marks of strong inflammatory action. <i>Chronic liver disease</i> is, frequently, not characterized by well-marked symptoms, and hence, when assumed, is difficult of detection.
33. HERNIA.	In the same manner as hydrocele; — by inflation.	The detection is easy.
34. HYDROCELE.	By puncturing the skin of the scrotum, and inflating the cellular membrane.	Do.
35. HYDROCEPHALUS.	By opening the integuments of the head, near the vertex, e. g. and blowing in air.	Do.
36. HYSTERIA.	Does not easily resist the application of strong sternutatories to the nostrils. Attentive examination necessary.
37. INSANITY.	The expression of countenance cannot easily be feigned. Nor can the affection be kept up so long as in real mental alienation. The individual cannot do so long without food, sleep, &c.
38. JAUNDICE.	By colouring the skin with an infusion of turmeric or tincture of rhubarb. <i>Clay-coloured stools</i> produced by taking daily a small quantity of muriatic acid. <i>High-coloured urine</i> by rhubarb taken internally.	The eyes cannot be coloured, although smoke has been used for this purpose. The skin must be washed to remove the colouring matter if any exist, and the supply of acid and rhubarb be prevented.

DISEASES, &c.	HOW FEIGNED.	HOW DETECTED.
39. LAMENESS.	By keeping the limb in a contracted state, and resisting any efforts to move it.	By two persons taking hold of the individual and moving rapidly along with him; and when they are tired, having relays. The impostor will generally give in.
40. MENSTRUATION.	By staining the clothes and body with borrowed blood.	By cutting off the supply.
41. MYOPIA. <i>Short-Sightedness.</i>	Present an open book, and apply the leaves close to the nose. If it cannot be read distinctly, when thus placed, or when glasses proper for short-sightedness are used, the disease is feigned.
42. OPHTHALMIA.	Excited by a variety of acrid and corrosive substances applied to the eye; as lime, &c. A portion of black muslin, spread over the cornea. The eyelashes are sometimes extracted, and caustic applied to excite disease in the palpebræ.	When ophthalmia is thus excited, its progress is ordinarily very rapid, arriving at its height within a few hours.
43. OZENA.	By impregnating a piece of sponge with some offensive juices or oils, mixed with decayed cheese, and putting the imbued sponge into the nostrils.	
44. PALPITATION.	White hellebore, given in the dose of 10 or 12 grains, and repeated, will occasion general indisposition, and undue action of the heart.	Cut off the supply.
45. PAINS.	The detection is here often difficult. The non-existence of pain cannot be proved, and great pain may be present without any appearance externally.
46. SHAKING PALSY.	The imposition is more frequently detected by inconsistencies and contradictions in the patient's history of the case, than in any other manner.
47. PARALYSIS.	May be suspected, if the person be in an ordinary state of vigour. Try violent remedies and means, recommended under Chorea.
48. PHTHISIS PULMONALIS.	Individuals with long necks and contracted shoulders have simulated phthisis, by covering the chest with blisters, cicatrices of issues, &c., and by taking drugs which cause paleness.	Violent remedies are here required. Cold affusion, actual cautery, electric shocks, &c.
49. POLYPUS NASI.	By introducing the testicle of a young cock, or the kidney of a rabbit, into the nostril, and retaining it there by means of a sponge fastened to it.	By attentive examination of the symptoms.
50. PORRIGO. <i>Scalped Head.</i>	By applying nitric acid to the head, after protecting the face with fatty substances: but the chronic state is imitated by the use of depilatories of different kinds applied sometimes in patches, so as to resemble the <i>Porrigio decalvans</i> .	
51. PREGNANCY.	By wearing pads, and assuming the longing after particular articles of diet, &c.	By the absence of the areola; the presence of a pad; and, if necessary, by examination <i>per vaginam</i> .
52. PROLAPSUS ANI.	By a portion of the intestine of the ox, in which a sponge filled with a mixture of blood and milk is placed.	
53. PROLAPSUS UTERI.	By a similar fraud.	

DISEASES, &c.	How FEIGNED.	How DETECTED.
54. PULSE, WEAKNESS OR DEFECT OF.	By ligatures applied to the corresponding arm.	By examining whether the arteries of the two arms beat alike; and if a ligature be placed on the arm.
55. CHRONIC RHEUMATISM. See Pain.		
56. SCROFULA.	By exciting ulcers below the angles of the jaw.	By examining the general habit, and observing whether the ulcerations be glandular, and the discharge of a scrofulous character.
57. SCURVY.	By covering the teeth with wax, and then applying acid, corrosive substances to the gums.	By examining the general habit: whether debilitated, cachectic, and possessing the usual general signs of scorbutus.
58. STAMMERING.	Simulators of this defect generally state, that it is connate, or ascribe it to a fit of apoplexy or severe fever. Where the organs of speech were perfect, and the moral evidence of the previous existence of the infirmity was not satisfactory, the French authorities used to confine the soldiers, and not supply them with food, until they called for it without stammering.
59. STRICTURE OF THE URETHRA.	By passing a bougie.
60. SWELLING OF THE LEGS.	By ligatures round the thighs.	Examine the limbs uncovered.
61. SYNCOPE.	Ligatures are sometimes used to prevent the pulse being felt. By applying lotions to the face to make it pale.	By using sternutatories. By the absence of some of the symptoms of syncope. Examine the naked arms. Wash the face.
62. TYMPANY.	Persons have possessed the power of swallowing air, so as to distend the stomach, and simulate tympany. By introducing a tent, imbued with blistering plaster into the ear, and repeating the application, until the tube becomes ulcerated, and a discharge of puriform matter is established. The fetid smell is imitated, by dropping into the ear a mixture of empyreumatic oil, asafetida, and old cheese. Also, by introducing a little honey into the meatus.	By careful examination.
63. ULCER OF THE EAR.	By corrosives, or irritants. Sometimes by abrasion, by rubbing sand on the shin-bone. At others, they are pretended, by gluing on a piece of spleen or the skin of a frog.	
64. ULCERS OF THE LEGS, &c.	The fruit of the Indian Fig (<i>Cactus opuntia</i>) colours the urine as red as blood. Cantharides will cause it. Blood may also be procured and mixed with the urine.	Artificial ulcers have, usually, a more distinct margin, and are more readily healed than others; the latter being generally indicative of an impaired constitution.
65. BLOODY URINE.		By making the patient pass his urine in the presence of the physician, and examining the vessel before and after. By cutting off the supply of any substance, which could cause the appearance.
66. INCONTINENCE OF URINE.	Difficult, at times, of detection. Give the person a full dose of opium, and introduce the catheter when he is asleep. If there be urine, the incontinence is feigned.
67. VARICOSE VEINS.	By a ligature, placed tightly round the limb. They may be excited in this manner, or aggravated if already existing.	By examining the limb.

FEL, Bile.—*f. Anguillæ*, see Bile.—*f. Bovinum*, Bile of the Ox.—*f. Bovis*, see Bile.—*f. Naturæ*, Aloes.—*f. Tauri*, see Bile.—*f. Terræ*, Ludus Helmontii.—*f. Ursi*, see Bile.—*f. Vitri*, see Vitrum.

FELDSCHEREN, Bathers.

FELINEUS, Bilious.

FELLEUS, Biliary, Bilious.

FELLIDUCUS, Chologogue.

FELLIS OBSTRUCTIO, Icterus.—*f. Superfusio*, Icterus.—*f. Suffusio*, Icterus.

FELON, Paronychia.

FELTING, from Anglo-Saxon *felt*, cloth or stuff made without weaving. *Tangling*. A term applied to the hair when inextricably interlaced, as occurs occasionally in women from inattention.

FÉLURE DE GLASER, Fissure, glenoid.

FELWORT, Gentiana lutea.

FEMALE, *Fœmina*, *Fœmina*, *Gynē*, (F.) *Femelle*, (*Femme*, 'woman.') In animals, the one that engenders and bears the young. It is, also, used adjectively;—as, the *Female Sex*, &c.

FEMALE PARTS, Vulva.

FEMELLE, Female.

FEMEN, Thigh (inner part), Vulva.

FEMINA, Female.

FEMINESCENCE, *Fæminescen'tia*, from *fœmina*, 'a female.' The possession or assumption of certain male characteristics by the female.—*Mehlis*.

FEMME EN COUCHE, Puerpera.

FEM'ORAL, *Femora'lis*, from *femur*, *femoris*, 'the thigh-bone.' Belonging or relating to the thigh; as *Femoral artery*, *Femoral hernia*, &c. See *Crural*.

FEMORALIS, Triceps cruris.

FÉMORO-CALCANIEN PETIT, Plantar muscle.—*f. Popliteal*, great, Sciatic nerve, great.—*f. Popliti-tibial*, Popliteus muscle.

FEM'ORO-TIBIAL, *Femoro-tibia'lis*. Belonging to the femur and tibia. The *Femoro-tibial articulation* is the knee-joint.

FEMUR, *Merus*, *Me'rium*. The thigh (q. v.) Also, the thigh bone, *Os fem'oris*, (F.) *L'os de la Cuisse*. The strongest and longest of all the bones of the body, extending from the pelvis to the tibia, and forming the solid part of the thigh. The femur is cylindrical, slightly curved anteriorly, unsymmetrical and oblique downwards and inwards. The body of the bone is prismatic, and has, behind, a prominent ridge, the *linea aspera*.

PROCESSES.	Upper or Pelvic Extremity.	Head supported on a column or neck.
		Great trochanter.
	Inferior or Tibial Extremity.	Lesser trochanter.
		External condyle.
		Internal condyle.
		External tuberosity.
		Internal tuberosity.

The femur ossifies from five points:—one on each process of the pelvic extremity; one in the body of the bone; and two on the condyles. It is articulated with the pelvis, tibia, and patella. See Thigh.

FÉMUR COL DU, Collum femoris.—*f. Moventium septimus*, Iliacus internus.—*f. Moventium sextus*, Psoas magnus.—*f. Summum*, Vulva.

FENES'TRA, *A window*, (akin to *φαῖνον*, 'I show,' and to *Sanscrit fānu*.) (F.) *Fenêtre*. Anatomists have given this name to two apertures, situate in the inner paries of the cavity of the tympanum. The one of these is the *Fenest'ra ovalis seu vestibularis*, *Fora'men ova'le*. It is oval-shaped; situate at the base of the stapes, and corresponds with the cavity of the vestibule. The other is the *Fenest'ra rotun'da seu cochlea'ris seu triquetra*, *Fora'men rotun'dum*. It is closed by a fine, transparent membrane, called

Membr'na Tympani secun'daria, *Tympanum minus seu secunda'rium*, *Membr'na fenest'ra rotun'da*, and corresponds to the inner scale of the cochlea.

FENESTRA OCULI, Pupil.

FENES'TRAL, *Fenest'ra'tus*, from *fenestra*, 'a window.' (F.) *Fenêtré* et *Fenêtré*. Bandages, compresses, or plasters with small perforations or openings, are so called. The openings prevent the detention of the discharge.

FENESTRATUS, Fenestral.

FENESTRÉ, Fenestral.

FEN'TRE, Fenestra.

FENÊTRÉ, Fenestral.

FENNEL, Anethum.—*f. Dogs'*, Anthemis cotula.—*f. Flower*, Nigella.—*f. Hog's*, Peucedanum.—*f. Sweet*, Anethum.—*f. Water*, Phellandrium aquaticum.

FENOUIL, Anethum.—*f. d'Eau*, Phellandrium aquaticum.—*f. Marin*, Crithlum maritimum.—*f. de Pore*, Peucedanum.—*f. Puant*, Anethum graveolens.

FENTE, Fissure.—*f. Capillaire*, see *Pilatio*.—*f. Glénoidale*, Fissure, glenoid.—*f. Orbitaire*, Orbital fissure.—*f. Orbitaire inférieure*, Sphenomaxillary fissure.—*f. Sphéno-maxillaire*, Sphenomaxillary fissure.—*f. Sphénoïdale*, Sphenoidal fissure.

FÉNUGREC, Trigonella fœnum.

FENUGREEK, Trigonella fœnum.

FER, Ferrum.—*f. Acétate* de, Ferri acetate.—*f. Brômure* de, see Bromine.—*f. Carbonate* de, Ferri protocarbonas.—*f. Carbure* de, Graphites.—*f. Chaud*, Pyrosis.—*f. Hydrate* de, tritoxide de, *Élites*.—*f. Hydriodate* de, see Ferri iodidum.—*f. Iodure* de, Ferri iodidum, see Iodine.—*f. Lactate* de, Ferri Lactas.—*f. Limaille* de, Ferri limatur.—*f. Perchlorure* de, Ferri chloridum.—*f. Peroxide* de, Ferri subcarbonas.—*f. et de Potasse*, tartrate de, Ferrum tartarizatum.—*f. et de Potassium*, cyanure de, Potassii Ferrocyanuretum.—*f. et de Potassium*, protocyanure de, Potassii Ferrocyanuretum.—*f. Réduit par l'hydrogène*, Ferri pulvis.—*f. Sulphate* de, Ferri Sulphas.—*f. Sulphure* de, Ferri Sulphuretum.—*f. Tannate* de, Ferri tannas.—*f. Trichlorure* de, Ferri chloridum.—*f. Valériante* de, Ferri Valerianas.

FERALS PEDICULUS, see Pediculus.

FERINE, *Fer'inus*; 'savage, brutal,' from *fera*, 'a wild beast.' *Therio'des*. A term, applied to any malignant or noxious disease. In France it is used only when joined to *Toux*; *Toux férine*, a dry, obstinate, and painful cough.

FERMENT, *Fermen'tum*, (quasi *fervimentum*, from *ferveo*, 'I burn, I ferment,') *Zyma*, *Zymo'ma*, *Leven*, (F.) *Lévain*. The Iatrochemists applied this name to imaginary substances, *Fermen'ta mor'bi*, to which they attributed the power of producing disease, by exciting a fermentation in the humours.

FERMENTA'TION, *Fermenta'tio*, *Zymo'sis*, *Æstua'tio*, *Causis*, *Brasmos*. Same etymon. An intestinal movement, occurring spontaneously in a liquid; whence result certain substances, that did not previously exist in it.

The chemical physicians attributed all diseases to an imaginary fermentation of the humours.

FERMENTATION, PUTREFACTIVE, Putrefaction.

FERMENTUM CEREVISIE, Yest.—*f. Morbi*, Ferment.—*f. Ventriculi*, Gastric juice.

FERN, BUSH, Comptonia asplenifolia.—*f. Cinnamon*, Osmunda cinnamomea.—*f. Eagle*, Pteris aquilina.—*f. Female*, Asplenium filix femina, Pteris aquilina.—*f. Flowering*, Osmunda regalis.—*f. Gale*, Comptonia asplenifolia.—*f. Male*, Polypodium filix mas.—*f. Meadow*, Comptonia asplenifolia.—*f. Rock*, Adiantum pedatum.—*f. Root*, Polypodium vulgare.—*f. Sweet*, Adiantum peda-

tum, *Comptonia asplenifolia*.—f. Sweet, shrubby, *Comptonia asplenifolia*.

FERNAMBUCO WOOD, *Cæsalpinia echinata*.

FERNFRECKLED, *Ephelides*.

FERNITICKLES, *Ephelides*.

FERO'NIA ELEPHANTUM, (from the goddess *Feronia*.) A tree of India, Order Aurantiaceæ, which yields a gum closely resembling Gum Arabic. Dr. Pereira thinks it not improbable, that a part of the *East India gum* taken to England may be the produce of this tree. The leaves are aromatic, of an odour resembling anise, and are used as a carminative.

FERRAMENTUM, *Siderion, Instrumentum ferreum*, from *ferreum*, 'iron.' Any surgical instrument made of iron. By the vulgar, in France, the word *ferrements* means the instruments used in difficult labours.

FERRARIA, *Scrophularia aquatica*.

FERRATUS, *Chalybeatus*.

FERREIN, CANAL OF, *Ductus Ferrei'ni*. A triangular channel, which Ferrein supposed to result from the approximation of the free edges of the eyelids applied to the globe of the eye; and which he considered adapted for directing the tears towards the puncta lacrymalia, during sleep. The canal is, probably, imaginary.

The same name is likewise given to the *cortical canals*,—the first portions of the uriniferous ducts, whilst still in the cortical substance of the kidney.

FERREIN, PYR'AMIDS OF, *Pyramides Ferrei'ni*. Each of the papillæ of the kidney, according to Ferrein, consists of, at least, 700 subordinate cones or pyramids. To these last the names 'pyramids of Ferrein,' *Pyramides renales Ferrei'ni*, *Fasciculi pyramidales*, *Fasciculi corticales*, *Lobuli renum*, have been given.

FERRI ACETAS, *Ectæticum martis aceticum*, *Aceticum martiale*, *Acetate of Iron*, *Iron Liquor*, (F.) *Acétate de fer*. A preparation of the Dublin Pharmacopœia, made by digesting 1 part of carbonate of iron in 6 parts of acetic acid for three days, and filtering. Dose, as a tonic and astringent. ℞v to ℞xx, in water.

FERRI ERUGO, F. Subcarbonas—f. et Ammoniac murias, *Ferrum ammoniatum*—f. Ammonio-chloridum, *Ferrum ammoniatum*.

FERRI ET ALUMINE SULPHAS, *Sulphate of Iron and Alumina*, is made by treating bicarbonated solution of soft iron and carbonated solution of pure washed alumina, with sulphuric acid. It has been recommended as a valuable astringent. Dose, from five to ten grains.

FERRI AMMONIO-CITRAS, *Ammonia ferro-citras*, *Ferrum ammoniaco-citricum*, *Ammonio-citrate of iron*, *Citrate of ammonia and iron*. Prepared by adding ammonia to citrate of iron, so as to neutralize the excess of acid. Dose, gr. v. to gr. viij. A POTASSIO-CITRATE and a SODIO-CITRATE OF IRON have also been introduced; and a CITRATE OF THE MAGNETIC OXIDE OF IRON prepared by combining the magnetic oxide with citric acid.

FERRI AMMONIO-TARTRAS, *Ammonio-tartrate of Iron*. This salt is best made by dissolving to saturation freshly precipitated hydrated oxide of iron in a solution of bitartrate of ammonia, and evaporating to dryness. It is very soluble. The dose is five grains or more in pill or solution.

FERRI ARSENIAS, *Arseniate of Iron*—f. *Borusias*, Prussian blue—f. *Bromidum*, see *Bromine*—f. *Carbonas*, F. subcarbonas—f. *Carbonas Saccharatum*, see F. Protocarbonas—f. *Carbonas cum Saccharo*, see F. Protocarbonas—f. *Carbonatum*, Graphites—f. *Carburetum*, Graphites.

FERRI CHLORIDUM, F. *Sesquichloridum*, (F.) *Trichlorure on Perchlorure de fer*, *Chloride*, *Sesquichloride*, *Perchloride* or *Muriate of Iron*, is obtained by dissolving sesqui-chloride of iron in

chlorohydric acid. It is generally prescribed in the form of *Tinctura Ferri Chloridi*. A concentrated solution of the perchloride has been used as an injection for the radical cure of aneurism.

FERRI CITRAS, *Ferrum citricum*, *Citrate of Iron*. Two citrates of iron are prepared—the *Sesquicitrate* or *Citrate of the Sesquioxide*; and the *Citrate of the Protoxide*. The former is official in the Ph. U. S. They resemble, in their medical properties, the tartrate and the lactate of the metal. Dose, five grains or more.

FERRI CYANURETUM, Prussian blue—f. *Deuto-carbonas fuscus*, F. subcarbonas—f. *Deutoxydum nigrum*, *Æthiops martial*—f. *Ferrocyanas*, Prussian blue—f. *Ferrocyanuretum*, Prussian blue.

FERRI FILUM. The pharmacopœial name of iron wire.

FERRI HYDRIOBAS, F. *Iodidum*—f. *Hydrocyanas*, Prussian blue—f. *Hypercarburetum*, Graphites.

FERRI IODIDUM, F. *Ioduretum*, *Ferrum iodatum*, *Iodide*, or *Ioduret of Iron*, (F.) *Iodure de fer*; in solution, *Hydroiodate of Iron*, *Ferri hydroiodas*, *Ferrum hydroiodatum*, (F.) *Hydroiodate de fer*, (Iodin. ℥ij, *Ferri rament*. ℥i. *Sp. destillat*. Oiss.) Mix the iodine with a pint of the distilled water, in a porcelain or glass vessel, and gradually add the iron filings, stirring constantly. Heat the mixture gently until the liquid acquires a light greenish colour; then filter; and, after the liquid has passed, pour upon the filter half a pint of the distilled water, boiling hot. When this has passed, evaporate the filtered liquor, at a temperature not exceeding 212°, in an iron vessel, to dryness; keep the dry iodide in a closely stopped bottle. (Ph. U. S.) Dose, gr. iij to gr. x or more.

FERRI IODURETUM, F. *Iodidum*.

FERRI LACTAS, *Ferrum lacticum*, *Lactate of Iron*, L. of *Protoxide of Iron*, (F.) *Lactate de Fer*. Prepared by digesting in a sand-bath, at a low temperature, diluted lactic acid with iron filings. It is employed in the same cases as the precipitated subcarbonate of iron, and especially in chlorosis. Twelve grains of the lactate may be given in the 24 hours, in the form of lozenges.

FERRI LIMAT'RA PURIFICATA, *Purified Iron Filings*:—purified by means of the magnet, for internal use. The filings are, also, called *Ferri Nodis*, F. *Ramentis et Filis*, *Martis Limat'ra*, *Speculum Indicum*, *Ferri in pulverem resolutio*, (F.) *Limaille de Fer*. They are considered to possess the general properties of iron:—the iron becoming oxidized.

FERRI ET MAGNE'SIE CITRAS, *Citrate of iron and magnesia*, is prepared by dissolving hydrated oxide of iron in a solution of citric acid, saturated with carbonate of magnesia, and evaporating to dryness. It is soluble in water, and does not constipate. Dose, from four grains to fifteen, in solution.

FERRI NITRAS, see *Liquor Ferri nitratis*—f. *Nitratis Liquor*, see *Liquor Ferri nitratis*—f. *Oxidum fuscum*, F. subcarbonas—f. *Oxidum hydratum*, *Ferrum oxydatum hydratum*—f. *Oxidum Magneticum*, *Oxydum Ferri nigrum*—f. *Oxidum Nigrum*, *Oxydum Ferri nigrum*—f. *Oxidum rubrum*, *Colecthar*, *Ferri subcarbonas*—f. *Oxidum hydratum*, *Ferrum oxydum hydratum*—f. *Percyanidum*, Prussian blue—f. *Peroxydum hydratum*, *Ferrum oxydatum hydratum*—f. *Persequintras*, see *Liquor Ferri nitratis*.

FERRI PHOSPHAS, *Phosphate of Iron*, (F.) *Sulphat*. ℥v, *Soda Phosphat*. ℥vi, *Aqua*, cong. Dissolve the sulphate of iron and phosphate of soda, severally, in four pints of the water; then mix the solutions, and set the mixture by, that the powder may subside; lastly, having poured off the supernatant liquor, wash the phosphate

of iron with water, and dry it with a gentle heat.—Ph. U. S.) Dose, 5 to 10 grains as a chalybeate. Rarely used.

FERRI PILA, *F. limatura*—*f. et Potassæ tartras*, Ferrum tartarizatum—*f. Potassio-citras*, see Ferri ammonio-citras—*f. Potassio-tartras*, Ferrum tartarizatum.

FERRI PROTOCARBONAS, *Proto-carbonate of Iron*, (F.) *Carbonate de Fer*. The protocarbonate of iron, thrown down from a solution of the sulphate of iron by the carbonate of soda, readily attracts oxygen and becomes converted into the sesquioxide. To prevent this, it may be associated with honey and sugar, and the mixture be reduced by evaporation to a pilular consistence. The mass constitutes the *Ferruginous Pills*, *Pilules ferrugineas*, of Vallet, *Vallet's Pills*. See *Pilule ferri Carbonatis*. The Edinburgh Pharmacopœia contains the *Ferri Car'bonas Saccharatum*, *Ferri Car'bonas cum Saccharo*, *Saccharine Car'bonate of Iron*, prepared in this manner. It is the *Ferrum Carbonicum Saccharatum* of Klauer.

The protocarbonate is given in the same diseases as the lactate of iron; ten or fifteen grains in the course of the twenty-four hours.

FERRI PROTOSULPHAS VIRIDIS, *F. Sulphas*—*f. Prussias*, Prussian blue.

FERRI PULVIS, *Fer réduit*, *F. réduit par l'hydrogène*, *Ferrum metallicum*, Powder of iron, *Réd'ced Iron*, *Quevonne's Iron*, prepared by passing a stream of hydrogen gas over the sesquioxide of iron, contained in an iron or porcelain tube heated to low redness. It is very liable to become oxidated, and must be kept in a dry, well-stopped bottle. It has been prescribed in anæmic, and especially in chlorotic cases. The ordinary dose is three to six grains, or more, three times a day, in pill made with sugar and gum. A formula for its preparation is given in the last edition of the Pharmacopœia of the United States (1851).

FERRI ET QUINIE CITRAS, *Ferrum et Chininum citricum*, *Citrate of Iron and Quinia*, formed by the union of four parts of citrate of iron and one part of citrate of quinia, has been prescribed in cases where a combination of these tonics is indicated.

FERRI RAMENTA, *F. limatura*—*f. Rubigo*, *F. Subcarbonas*—*f. Scobs*, *F. limatura*—*f. Sesquichloridum*, *Ferri chloridum*—*f. Sesquioxidum*, *F. subcarbonas præcipitatus*.

FERRI SESQUINITRATIS, *Liquor*, see *Tinctura Ferri muriatis*—*f. Sodio-tartras*, see *Ferri et ammonio-citras*.

FERRI SUBCARBONAS, *F. Car'bonas*, *F. Rubigo*, *Ferrum præcipitatum*, *Chalybis Rub'go præparata*. (The last two terms, as well as *Ferr'go*, are applied to the subcarbonate or rust, *Æru'go Ferri*, *Cacæferri*, formed by the action of moist air on metallic iron.) A protoxide of iron, oxidized by the decomposition of water; the carbonic acid being attracted from the air. Dose, gr. v to ʒss and more.

Ferri Subcarbonas may be precipitated from sulphate of iron by carbonate of soda. The following is the formula of the Pharmacopœia of the United States:—*Ferri sulph. ʒviij; sodæ carb. ʒix; aquæ bullient. cong.* Dissolve the sulphate of iron and carbonate of soda severally in four pints of the water, then mix the solutions, and, having stirred the mixture, set it by that the powder may subside: having poured off the liquor, wash the subcarbonate with hot water, wrap it in bibulous paper, and dry with a gentle heat. It is, also, called *Ferri sesquioxidum*, *Oxidum ferri fuscum*, *O. ferri rubrum*, *Deuto-car'bonas Ferri fuscum*, *Crocus martis ap'riens*,

Sesquioxide or peroxide of iron, (F.) *Peroxide de fer*, *Safran de Mars ap'ritif*.

FERRI SULPHAS, *Sal Martis*, *Vitriolum Martis* seu *Ferri*, *Ferrum Vitriolatum*, *Sulphus* seu *Protosulphus Ferri viridis*, *Calcud'num*, *Cal'catar*, *Cal'catar*, *Chalcant'um*, *Calcit'a*, *Atramentum aut'rium*, *Vit'riol*, *Vitriolum viride*, *An'ima Hep'atis*, *Sulphate of Iron*, *Green Vitriol*, *Cop'peras*, (F.) *Sulfate de fer*. The Pharmacopœia of the United States directs it to be made by the action of sulphuric acid ʒxviij, on iron wire, cut in pieces, ʒxij—water, a gallon—evaporating, crystallizing, and drying the crystals on bibulous paper. This salt is inodorous, and of a strong styptic taste. The crystals are light green, transparent, rhomboidal; and soluble in two parts of water. It is tonic and anthelmintic. Dose, gr. j. to vj and more.

FERRI SULPHAS CALCINATUM, *Colcothar*.
FERRI SULPHAS EXSICCATUM, *Ferri Sulphas siccatum*, of the Pharmacopœias, is *Sulphate of Iron* exposed to a moderate heat, until it is capable of being reduced to powder. In this form it can be made into pills. Three grains may be esteemed equivalent to five of the crystallized salt.

FERRI SULPHURETUM, *Sulphuret of Iron*, *Iron pyrites*, (F.) *Sulfure de fer*; may be made by heating one part of sublimed sulphur, over three parts of iron filings in a crucible until the mass begins to glow. It is employed as a ready means for obtaining hydrosulphuric acid gas by the addition of sulphuric or chlorohydric acid.

FERRI TANNAS, *Ferrum tannicum*, *Tannate of Iron*, (F.) *Tannate de fer*. This salt is usually obtained by adding a solution of a salt of sesquioxide of iron, as the persulphate, to a decoction of nut-galls. It is possessed of tonic and astringent properties, and has been extolled, especially in chlorosis.

FERRI TARTARUM, *Ferrum tartarizatum*—*f. Tritoxydum*, *Colcothar*.

FERRI VALERIA'NAS, *Ferrum Valerianicum*, *Vale'rianate of Iron*, (F.) *Valérianate de fer*. Formed by the action of valerianic acid on oxide of iron. It is a dark brick-red powder, insoluble in water, and has been given in hysteria complicating chlorosis. Dose, from two to four grains. The Dublin Pharmacopœia of 1850 has a formula for this salt.

FERRICUS HYDRAS, *Ferrum oxydum hydratum*.

FERRO-MAN'GANIC, *Ferro-mangan'icus*. Composed of iron and manganese—as a “ferro-manganic preparation.”

FERRUGINEOUS, Chalybeate.

FERRUGINOUS, Chalybeate.

FERRUGO, *Ferri subcarbonas*, *Ferrum oxydatum hydratum*.

FERRUM, *Mars*, *Side'ros*, *Metal'um hæmatopœticum*, *Iron*, (F.) *Fer*. A metal of a bluish-gray colour; fibrous texture; brilliant and fine-grained fracture. Specific gravity 7·600 to 7·800; hard, ductile, malleable, and magnetic. The medicinal virtues of iron are tonic; producing fetid eructations, when it takes effect, owing to its meeting with acid in the stomach, which oxidizes it, and causes the evolution of hydrogen gas. When given in the metallic state, the filings are chiefly used; but the oxides and salts are most commonly employed.

FERRUM AMMONIACALE, *F. ammoniatum*—*f. Ammoniao-citricum*, *Ferri ammonio-citras*.

FERRUM AMMONIATUM, *F. ammoniacal'e*, *Murias Ammon'ia et Ferri*, *Ferri ammonio-chloridum*, *Flores martiales seu salis ammon'iaci martiales*, *Ens martis seu Ven'eris Boy'lei*, *Sal martis murial'icum sublimatum*, *S. ammon'acum martial'e*, *Ammonium murial'icum martia'tum*

sea martialis, *Ammoni'acum hydrochlorat'um ferat'um*, *Aroph. Paracel'si*, *Calendulae mineral'es*, *Ammoni'ated Iron*, *Ammoni'o-chloride of Iron*, (F.) *Muriate d'ammoniaque et de fer*. A mixture of the hydro-chlorates of ammonia and iron. [?] (*Ferri Subcarb.* ʒiij; *acid. muriat.* f ʒx; *ammon. muriat.* lbjss; *aq. destillat.* Oij. Mix the subcarbonate with the acid in a glass vessel and digest for two hours; then add the muriate previously dissolved in distilled water, and having filtered the liquor, evaporate to dryness. Rub to powder.—Ph. U. S.) Dose, gr. iij. to gr. xv.

FERRUM ARSENIATUM, Arseniate of iron—f. *Arsenicicium oxydulatum*, Arseniate of iron—f. *Borussicum*, Prussian blue—f. *Bromatum*, see Bromine—f. *Carbonicum Saccharatum*, Ferri protocarbonas—f. et *Chininum citricum*, Ferri citras—f. *Citricum*, Ferri citras—f. *Cyanogenatum*, Prussian blue—f. *Hæmatites*, Hæmatites—f. *Hydriodatum*, see Ferri iodidum—f. *Hydrobromicum oxydatum*, see Bromine—f. *Lacticum*, see Ferri lactas—f. *Magnes attractorium*, Magnet—f. *Metallicum*, Ferri pulvis.

FERRUM OXYDATUM HYDRATUM, *Ferri Oxidum Hydratum* (Ph. U. S.) seu *Peroxidum Hydratum*, *Ferrug'o*, *Hydras Fer'ricus*, *Hydro-oxide of Iron*, *Hydrated peroxide of Iron*, *Hydrated Trizoxide of Iron*. It may be prepared by taking a solution of sulphate of iron, increasing its dose of oxygen by heating it with nitric acid, and precipitating the oxide by adding pure ammonia in excess, washing the precipitate, and keeping it moist.

The following is the formula adopted by the Pharmacopœia of the United States: *Ferri Sulph.* ʒiv; *acid. sulphuric.* f ʒiiss; *acid. nitric.* f ʒvi, vel q. s., *Liq. ammoniæ* q. s., *aquæ* Oij. Dissolve the sulphate of iron in the water, and having added the sulphuric acid boil the solution: then add the nitric acid in small portions, boiling the liquid for a minute or two after each addition, until the acid ceases to produce a dark colour. Filter the liquid, allow it to cool, and add the liquor ammoniæ in excess, stirring the mixture briskly; wash the precipitate with water, until the washings cease to yield a precipitate with chloride of barium, and keep it close in bottles with water sufficient to cover it.

It has been brought forward as an antidote to arsenic; and many cases of its efficacy have been published.

From 10 to 20 parts of the hydrated oxide would seem to be more than sufficient to convert 1 part of arsenious acid into the basic salt of iron.

FERRUM OXYDULATUM HYDROCRANICUM, Prussian blue—f. *Potabile*, *F. tartarizatum*—f. *Præcipitatum*, Ferri subcarbonas.

FERRUM TANNICUM, Ferri tannas.

FERRUM TARTARIZATUM, *Tartaras Potassæ et Ferri*, *Tartarum Ferri*, *Ferri et Potassæ Tartaras*, (Ph. U. S.) *Tartarus chalybeatus seu martialis*, *Mars solubilis*, *Chalybs tartarizatus*, *Ferri potass-i-tartaras*, *Tartaras kalico-fer'ricus* seu *Potassæ ferrugineus*, *Ferrum potabile*, *Globus martialis*, *Globuli Tartari martiales seu martiales soluti* seu *martiali seu tartaratis ferri et liciis*, *Boli Martis*, *Pyri martiales*, *Tartarized Iron*, *Tartarate of Potassa and Iron*, (F.) *Tartarate de potasse et de fer*, *Boule de Mars ou de Nancy ou de Molsheim ou d'Acier*. (*Ferri subcarb.* ʒiij, *Acid. muriat.* f ʒx; *Liquor Potassæ*, Ovss; *Potassæ Bitart.* ʒvijs, *Aquæ destillat.* cong. iss. Mix the subcarbonate of iron and the muriatic acid, and digest for two hours; pour the solution into a gallon of the distilled water; set aside for an hour, and pour off the supernatant liquor. To this add the liquor potassæ: wash the precipitate formed frequently with water, and while yet moist, mix it with the bitartrate of potassa and

half a gallon of the distilled water. Keep the mixture at the temperature of 140° for 30 hours, frequently stirring; filter the solution, and evaporate by means of a water bath, at the same temperature, to dryness.—(Ph. U. S.) It is one of the mildest of the salts of iron, and not unpalatable. Dose, gr. x. to ʒss.

The *Tinctura Martis Aperiti'va*, *Tincture of Ludwig*; *Al'cohol cum Sulphat'e Ferri tartariadatus*; *Tinctura Martis Glauberi* is, essentially, a solution of this salt. It is also called *Eau de Boule*, and is used in contusions.

Helvetius's Styptic was composed of the filings of iron and tartar, mixed to a proper consistence with French brandy. It was called in England *Eaton's Styptic*; but this is now formed of Sulphate of Iron.

FERRUM VALERIANICUM, Ferri valerianas—f. *Vitriolatum*, Ferri sulphas—f. *Vitriolatum ustum*, Colcothar—f. *Zooticum*, Prussian blue.

FERSÆ, Rubecula.

FERTILITY, Fecondity.

FER'ULA, *Fanon*, *Palette*, *Splint*—f. *Africana*, *Bubon galbanum*—f. *Asafœtida*, see *Asafœtida*—f. *Graveolens*, *Anethum graveolens*—f. *Opopanax*, *Pastinaca opopanax*—f. *Persica*, see *Sagapenum*—f. *Tingitana*, see *Ammoniac*, gum.

FERUS, Homicidal.

FESSIAN, Physician.

FESSES, Nates.

FESSIER, Gluteal—f. *Grand*, *Glutæus maximus*—f. *Moyen*, *Glutæus medius*—f. *Petit*, *Glutæus minimus*.

FESTER. A word of unknown etymology, which signifies to corrupt or rancle; and, when applied to a sore, means to suppurate, or rather to discharge a thin fluid as an evidence of its being irritated.

FETID, *Fætidus*, *Dysœdes*, *Cacoœdes*, *Bromoœsus*, *Gravœolens*. Having a bad smell.

FETUS, Fœtus, Pregnant.

FEU ACTUEL, Cauterium—f. *Persigue*, *Herpes zoster*, see *Anthrax*—f. *Potential*, see *Cauterium*—f. *Sacré*, *Erysipelas*—f. *St. Antoine*, *Erysipelas*—f. *Sauvage*, *Ignis sylvaticus*—f. *Volage*, *Ignis sylvaticus*.

FEUILLET, Omasum; see *Tache embryonnaire*.

FEUX DE DENTS, Strophulus—f. *Volages*, *Porrigo larvalis*.

FÈVE, *Vicia faba*—f. *de Carthagène*, *Habilla de Carthagena*—f. *à Cochon*, *Hyoscyamus*—f. *Épaisse*, *Sedum telephium*—f. *des Marais*, *Vicia faba*—f. *de Sainte Ignace*, *Ignatia amara*—f. *Purgatif*, *Ricinus communis*.

FEVER, *Febris*, from *feritas*, 'wildness,' or from *feror*, 'heat,' or from *februo*, 'I purify.' *Pyr*, *Pyr'etos*, *xuperos*, (F.) *Fievre*. One of the most frequent and dangerous affections to which the body is liable. A person has an attack of fever, when he is affected with rigors, followed by increased heat of skin, quick pulse, languor, and lassitude. Rigors, increased heat, and frequency of pulse have each been assumed as the essential character of fever. It is not characterized, however, by any one, but depends upon the coexistence of many symptoms. Fevers have been usually divided into *idiopathic* or *essential*, and into *symptomatic*. The idiopathic arise without any obvious local cause. The symptomatic are dependent upon local irritation. Idiopathic fevers may be divided into three classes: 1. Those attended with distinct paroxysms:—intermittents. 2. Remittent and continued fevers:—ad. 3. Fevers complicated with eruptions, or the exanthematous. These divisions admit of great variety, owing to climate, season, soil, age, &c. All ages and climates are liable to fever; and its exciting

causes are very numerous. These causes may be common; as irritations in the intestines; external injuries, stimulants, &c.; or they may be specific; as miasmata, contagion, &c. The greatest diversity has prevailed regarding the theory of fever. Its primary seat has been placed in the brain, mucous membrane of the stomach and intestines, skin, nerves, blood-vessels, liver, vena cava, pancreas, &c. It would seem, however, that although, in fever, the whole of the functions are morbidly impressed, the arguments in favour of the impression being first made on the nervous system and the system of nutrition are the strongest. The exciting cause of fever, whatever it may be, produces an irregular action in the system of nutrition, which is soon conveyed to the rest of the system, owing to the extensive sympathy which exists between every part of the body; and it is probable, that all those local inflammations and congestions are the consequence, rather than the cause, of this disordered condition of the system. The general character of fever is clearly shown by examination of the blood. When fever is devoid of inflammatory complication, the quantity of fibrin is in no case augmented. It frequently remains in the healthy proportion, and at times diminishes to an extent not met with in any other acute disease. The alteration of the blood in fevers, which consists generally in a diminution of the fibrinous element, is the reverse of what occurs in inflammation.

Many phenomena of fever are influenced by that periodicity, which we notice in the execution of several of the functions of the body. The types of intermittents are strong evidences of such an influence.

In the treatment of fever, it is important to bear in mind, 1. Its tendency, particularly in the case of the exanthemata, to run a definite course, and terminate in restoration to health. 2. The disposition to local determination or hyperæmiæ:—the most frequent cause of the fatal termination of fever; a circumstance requiring the vigilant attention of the physician. 3. That the symptoms must be attentively studied, in order to deduce, as far as possible from them, the indications of cure. Lastly, attention must be paid to the prevalent epidemic. There are particular seasons in which fevers are very malignant; and others in which they are as mild; circumstances which necessarily have an effect upon the treatment.

FEVER AND AGUE, Intermittent Fever—f. Acclimating, F. strangers.

FEVER, ADE'NO-MENING'EAL, *Febris ade'no-mening'eal* seu *mesenter'ica* (BAGLIVI) seu *muc'o'sa*, *Morbus muc'o'sus*, *Gastro-duodenop'gra*, *Mucous* or *Pitu'itous* or *Catar'rhal* Fever, *Phlegmap'gra*, *Phlegmap'gra*, *Phlegmatop'gra*, *Gastro-Bronchi'tis*. Fever, accompanied with considerable mucous secretion; especially from the digestive tube.

FEVER, ADE'NO-NERVOUS, Plague.

FEVER, ADYNAM'IC, *Febris adynam'ica*, *Asthenicop'gra*, *Asthenicop'gr'etus*, *Asthenop'ora*, *Asthenop'gr'etus*. Fever attended with great prostration of the vital powers—as Typhoid and Typhus fever.

FEVER, AF'RICAN, *Febris Africa'na*. The malignant bilious remittent fever, which prevails on the western coast of Africa,

FEVER, ALGID, see *Algidus*.

FEVER, ANOM'ALOUS, *Febris Anom'ala* seu *erru'tica* seu *irregula'ris* seu *vaga*. A fever, whose progress and type are irregular.

FEVER, APHON'IC, *Febris apho'nica*. A variety of intermittent, in which the voice is lost during the paroxysm.

FEVER, APOPLEC'TIC, *Febris apoplec'tica* seu *comato'des* seu *lethar'gica* seu *soporo'sa*, *Com'a-*

tose Fever. An intermittent or continued fever attended with apoplectic symptoms.

FEVER, ARTICULAR ERUPTIVE, Dengue—f. Articular rheumatic, Dengue.

FEVER, ARTIFIC'IAL, *Febris artificialis* seu *arte promo'ta*. Fever produced designedly by the internal or external use of stimulants.

FEVER, ASTHEN'IC, *Febris asthen'ica*. Fever accompanied with debility. It may include every variety of fever under certain circumstances, but is generally appropriated to typhus.

FEVER, ASTHMAT'IC, *Febris asthma'tica*. A pernicious intermittent, accompanied with symptoms of asthma.

FEVER, ATAXO-ADYNAM'IC, *Febris atax'o-ady-nam'ica*. Fever characterized by symptoms of ataxia and adynamia.

FEVER, BARCELONA, Fever, yellow—f. Bastard, Illegitimate fever.

FEVER, BIL'IOUS, *Febris bilio'sa* seu *polychol'ica* seu *as'o'des* seu *choleric'a* seu *gastro'ica* seu *hepat'ica* seu *cholopyret'ica*, *Cholop'gra*, *Cholopyr'etus*, *Hepato'gastro'choloid'osis*, *Enterop'gra bilio'sa*, *Choloz'e'mia febril'is*, *Syn'ochus bilio'sa*. The common remittent fever of summer and autumn; generally supposed to be owing to, or connected with, derangement of the biliary system.

FEVER, BILIOUS REMITTING. YELLOW, Fever, yellow—f. Bladdery, Pemphigus.

FEVER BLISTER. A term, vaguely applied to vesicular or pustular eruptions, which appear particularly about the mouth, during or after febrile disturbance.

FEVER, BOA VISTA. A malignant bilious remittent fever, greatly resembling yellow fever, which was very fatal at Fernando Po, and in ships in its waters, in the year 1845.

FEVER, BONA. A malignant paludal fever, which prevailed amongst the troops of the garrison at Bona in Algeria, from 1832 to 1835.

FEVER, BONE, see Inflammation—f. Bouquet, Dengue—f. Brain, F. cerebral, Phrenitis—f. Brain, water, Hydrocephalus internus—f. Breakbone, Dengue—f. Bucket, Dengue—f. Bulam, Fever, yellow—f. Camp, Typhus gravior.

FEVER, CARDIAL'GIC, *Febris Cardial'gica*. A variety of pernicious intermittent, accompanied with violent cardialgia during the paroxysm.

FEVER, CARDIT'IC. Intermittent fever, accompanied with pain at the heart.

FEVER, CATARRHAL, F. adeno-meningeal.

FEVER, CEPHAL'GIC, *Febris cephal'gica*. A pernicious intermittent, accompanied with intense pain of the head. Also, intermittent cephalalgia.

FEVER, CEPHAL'IC, *Febris cephal'ica*. A febrile affection of children—intermediate between the acute form of active cerebral congestion and the chronic form—which is attended by fever, pain in the head, disorder, or more generally constipation of the bowels, and a train of phenomena often supposed to be premonitory of an attack of hydrocephalus.—Mauthner.

FEVER, CER'EBRAL, Brain fever. Fever, generally of an ataxic character, in which the brain is considerably affected.

FEVER, CHAGRES, Panama' Fever. A severe form of endemic fever, common on the isthmus of Panama.

FEVER, CHILDBED, Fever, puerperal, Puerperal Fever—f. Childbed, Low, see Peritonitis.

FEVER, CHOL'ERIC, *Febris choleric'a*. A variety of pernicious intermittent, accompanied with symptoms of cholera morbus.

FEVER, CHOLERIC, OF INFANTS, Cholera infantum.

FEVER, CHRONIC, *Febris chron'ica*. Some authors apply this name to protracted fevers; others to hectic fever.

FEVER, COLLIQ'UATIVE, *Febris tab'ida seu colliquat'iva*. Fever, characterized by rapid emaciation, copious evacuations, and rapid prostration of strength.

FEVER, COMATOSE, *F. apoplectic*.

FEVER, CONGESTIVE, *Febris congest'iva*. Fever accompanied by obscure symptoms; or by great oppression and depression; in which it is difficult—and often impossible—to induce reaction. Congestive fevers occurs in various parts of this country, especially in the fall; and they are very common in India. The term congestive fever is often used in some parts of the south of the United States very indefinitely—to include winter typhus, and typhoid fevers, typhoid pneumonia, as well as intermittents and autumnal remittents.—Dickson.

FEVER, CONTINENT, see Continent and Continued fever.—*f. Continual*, see Continued fever.—*f. Continued*, common, Synochus.

FEVER, CONVULSIVE, *Febris convuls'iva*. A pernicious intermittent or remittent, accompanied by convulsions.

FEVER, COUNTRY. Fever occasioned by exposure to the miasms of a highly malarious soil. A term much used in the Southern States.

FEVER, DELIRIOUS, *Febris cum delir'io*, (*F.*) *Fèvre délirante*. A pernicious intermittent, characterized by delirium in the paroxysms.

FEVER, DEPURATORY, *Febris depurato'ria*. A fever, to which was attributed the property of purifying the blood; or which indicated, that such a supposed depuration had occurred.

FEVER, DIAPHORETIC, *Febris diaphoret'ica*, *Febris sudato'ria*. A pernicious intermittent, with excessive sweating during the fit.

FEVER, DIARY, Ephemera.

FEVER, DIGESTIVE. The chilliness, followed by increased heat and quickness of pulse, which frequently accompanies digestion.

FEVER, DOUBLE, (*F.*) *Fèvre double ou doublée*. An intermittent, which has two paroxysms in a given time, instead of one.

FEVER, DOUBLE-QUARTAN. A fever, whose paroxysms occur two days in succession, and fail the third day; the first paroxysm resembling the 4th, and the second the 5th.

FEVER, DOUBLE-QUOTIDIAN. An intermittent, whose paroxysms return twice every day at corresponding hours.

FEVER, DOUBLE-TERTIAN. An intermittent, whose paroxysms return every day; the first corresponding with the 3d, the second with the 4th, and so on.

FEVER, DUODECIM'IANE, *Febris duodecima'na*. A supposititious intermittent, whose paroxysms recur on the 12th day, or every 11 days.

FEVER, DYNAMIC, Synocha.—*f. Endemial*, Remittent *F.*—*f. Endemic*, Remittent *F.*—*f. Endemial*, Remittent *F.*—*f. Enteric*, see Typhus—*f. Ephemeral*, Ephemera.

FEVER, EPILEPTIC, *Febris epilept'ica*. A variety of pernicious intermittent, accompanied with attacks of epilepsy.

FEVER, EROTIC, *Febris erot'ica*, (*F.*) *Fèvre d'amour*. A chronic fever, occasioned by unpropitious love.

FEVER, ERRATIC, see Erratic.—*f. Eruptive*, Exanthematica.—*f. Eruptive*, articular, Dengue.—*f. Exacerbating*, Remittent Fever.

FEVER, FAINTING, OF PERSIA. A singular and fatal epidemic, presenting some points of analogy with cholera, which prevailed at Teheran in the autumn of 1842.

FEVER, FIVE DAY, see Relapse.

FEVER, GANGRENOUS, *Febris gangræno'des*. Fever, accompanied by gangrene of various parts, and especially of the limbs and genitals. Sénac describes an intermittent of this kind.

FEVER, GASTRAL'GIC. An intermittent accompanied with acute burning lacerating pain at the stomach.

FEVER, GASTRIC, *Febris gas'trica seu gas'trica bilio'sa*, *Gastrop'yra*, *Gastropyretus*, *Gastrocholo'sis*, *Hepatogastrocholo'sis*, *Stomach'ic fever*, *Syn'ochus Bilio'sa*. A name given by some to bilious fever, which has appeared to them to be dependent on an affection of the stomach. Also, called *Menin'go-gastric fever*, *Febris meningo-gas'trica*, *Meningo-gastrit'is*, *Gastro-meningit'is*, *Harvest fever*, (*F.*) *Fèvre de la Moisson*, *F. Méningo-gastrique*.

FEVER, GASTRO-ADYNAM'IC, *Febris gastroadynam'ica*. A fever, in which the symptoms of bilious fever are joined with those of adynamic fever.

FEVER, GASTRO-ANGIOTEN'IC. A fever, in which the symptoms of bilious are united with those of inflammatory fever.

FEVER, GASTRO-ATAX'IC. A fever, in which the symptoms of bilious fever are united with those of ataxic fever.

FEVER, GIBRALTAR, Fever, yellow.—*f. Harvest*, *F. Gastric*—*f. Hæmagastric*, *F. yellow*.

FEVER, HÆMOP'TOIC, *Febris Hæmopto'ica*. A variety of masked intermittent, in which periodical hæmoptysis is the chief symptom.

FEVER, HAY, *Catarr'hus æstivus*, *Summer Catarrh*, *Hay Asthma*, *Rose Catarrh*, *Summer Bronchit'is*. A catarrh to which certain persons are subject in summer, and which has been ascribed in England to the effluvium of hay, but this is not the probable cause. It is a catarrh with sneezing, headach, weeping, snuffling and cough, with, at times, fever and general discomfort. It is not uncommon in this country. It disappears spontaneously,—to recur on subsequent years about the same period.

FEVER, HEBDOM'ADAL. A supposititious fever, whose paroxysms return weekly, and on the same day.

FEVER, HECTIC, see Consumption, and Hectic Fever.

FEVER, HEPAT'IC or *HEPATAL'GIC*. A pernicious intermittent, with violent pain in the right hypochondrium.

FEVER, HILL. A modification of remittent, occurring in the hilly districts of India.

FEVER, HOSPITAL, Typhus gravior.

FEVER, HU'MORAL, *Febris humora'lis*. Fever in which an alteration or deterioration of the humours is suspected.

FEVER, HUNGARY, *Febris Hungar'ica seu Pannon'ica*, *Lues Panno'nia*, *Morbus Ungar'icus*, *Cephalon'osus*, *Vermis cer'ebri*, (*F.*) *Fèvre Hongroise ou de Hongrie*. An epidemic typhus, common amongst the soldiers in barracks, in Hungary.

FEVER, HYDROPHOB'IC, *Febris hydrophob'ica*. Pernicious intermittent, with dread of liquids.

FEVER, HYSTER'IC, *Febris hyster'ica*. Hysteria, accompanied by fever. Hysteria, occurring with each paroxysm of an intermittent.

FEVER, ICTERIC, *Febris icter'ica*. Fever, accompanied with jaundice. Some intermittents exhibit this complication at each paroxysm.

FEVER, ICTERIC, REMITTING, see Relapse.

FEVER, INFANTILE REMIT'TENT, *Febris infantum remittens*, *F. vermino'sa*, (of many), *F. mucosa vermino'sa*, *F. lenta*, *Spurious worm fever*, *Remitting fever of children*, *Hæctic infant'ilis*. A fever occurring in childhood, which often assumes many of the characters of hydrocephalus. It appears generally to be dependent upon a morbid condition of the stomach and bowels.

FEVER, INFLAMMATORY, Synocha.

FEVER, INSIDIOUS. Fever, which, at first, seems devoid of danger, but subsequently becomes of a more or less malignant character.

FEVER, INTERMITTENT, see Intermittent fever —f. Jail, Typhus gravior —f. Jungle, see Jungle fever.

FEVER, LAKE. Fever produced by exposure to the malaria in the neighbourhood of the northern lakes of this country.

FEVER, LENTICULAR, Miliary fever —f. Irritative, Irritation, morbid.

FEVER, LO'CHIAL, *Febris lochia'lis*. That acceleration of the circulation which sometimes occurs during the discharge of the lochia.

FEVER, LUNG, Catarrh, Pneumonia.

FEVER, MALIGNANT, *Febris maligna*. Fever which makes its approaches insidiously, and subsequently becomes formidable. Any fever which exhibits a very dangerous aspect. *Typhus gravior*.

FEVER, MALIGNANT PESTILENTIAL, Fever, yellow—f. Marsh, Elodes (febris).

FEVER, MASKED, *Febris larvata*, (F.) *Fièvre larvée*, Dead or Dumb Ague. An anomalous intermittent, the paroxysms of which have not the regular stages, or are masked.

FEVER, MENINGO-GASTRIC, Fever gastric.

FEVER, MESENTERIC, *Febris mesenterica*. A name given, by Baglivi, to a species of fever which appears to have belonged either to the mucous or bilious.

FEVER, MILK, *Febris lactea*, (F.) *Fièvre de lait ou lactée*; *Galactopyra*, *Galactopyretus*. The fever which precedes or accompanies the secretion of milk in women recently delivered. It comes on generally about the third day after delivery, and is characterized by quick pulse; increased heat; redness of face; diminution or temporary suspension of the lochial discharge; tumefaction and tension of the breasts. It commonly terminates in twenty-four hours, and often with profuse perspiration. It requires the use of antiphlogistics, with dry diet.

FEVER, MIXED, Synochus—f. Mucous, F. adeno-meningeal.

FEVER, NEPHRITIC, *Febris nephritica*. Intermittent fever, accompanied with nephritic pain during the paroxysm.

FEVER, NERVOUS, *Febris nervosa* seu *neurodes* seu *lenta nervosa*, *Neuropyra*, *Neuropyrætus*. A variety of Typhus; the *Typhus mitior* of Cullen. By many, however, it is esteemed a distinct disease. See Typhoid fever.

FEVER, NERVOUS, WITH EXANTHEMATOUS ERUPTION, Typhus.

FEVER, NEURALGIC, Dengue.

FEVER, NIGER. A malignant fever, of the bilious remittent kind, which proved fatal to many in the expeditions sent out by the British government to explore the Niger, in the years 1841-42, and previously.

FEVER, NON'ANE, *Febris nona'na*. A supposititious fever, whose paroxysms recur every ninth day, or every eight days.

FEVER, OCT'ANE, *Febris octa'na*. An intermittent, whose paroxysms recur every eighth day.

FEVER, PALUDAL, Intermittent —f. Panama, Fever, Chagres—f. Paroxysmal, Remittent fever —f. Periodic, Intermittent.

FEVER, PERIODICAL, *Febris periodica*. An intermittent or remittent fever.

FEVER, PERNICIOUS, *Febris pernicio'sa*. Intermittent fever, when attended with great danger, and which destroys the majority of those affected by it in the first four or five paroxysms; sometimes in the very first.

FEVER, PESTILENTIAL, *Febris pestilentialis*. The Plague. Also a severe case of typhus. The yellow fever and sweating sickness have, likewise, been thus designated.

FEVER, PESTILENTIAL, OF CATTLE, Murr —f. Putrid, F. adeno-meningeal.

FEVER, PLEURITIC. An intermittent or remittent, accompanied with inflammation of the pleura.

FEVER, PNEUMONIC, *Febris pneumonica*. An intermittent, accompanied with inflammation of the lungs. Also, pneumonia.

FEVER POISON, see Poison —f. Pseudo, Irritation, morbid.

FEVER, PUERPERAL, *Febris puerperalis*, *Child-bed fever*, (F.) *Fièvre puerpérale*. This name has been given to several acute diseases, supervening on delivery. It means, generally, a malignant variety of peritonitis, which runs its course very rapidly, and passes into a typhoid condition, unless met, at the very onset, by the most active depleting measures. By the generality of practitioners, it is esteemed to be eminently contagious; some, however, deny that it is so. See Peritonitis, and Puerperal fever.

FEVER, PUERPERAL DYNAMIC OR MALIGNANT, see Peritonitis—f. Puking, Milk sickness.

FEVER, PU'ULENT, *Febris purulenta*. Fever, which accompanies suppuration.

FEVER, PUTRID, Typhus gravior.

FEVER, QUINTAN, *Febris quinta'na*. A fever, whose paroxysms return every fifth day. It is seen rarely, or never.

FEVER, QUOTIDIAN, see Quotidian —f. Red Tongue, see Typhus.

FEVER, REGULAR, *Febris regularis* seu *exquisita*. An intermittent whose paroxysms follow a determinate type. It is opposed to *atypic*. Sometimes opposed to *anomalous*.

FEVER, RELAPSING, see Relapse.

FEVER, REMITTENT, see Remittent Fever —f. Remittent, Bilious, see Relapse—f. Remittent, infantile, see Fever, infantile remittent—f. Remittent of children, F. infantile remittent—f. Remitting icteric, see Relapse—f. Rheumatic, Rheumatism, acute—f. Root, *Triosteum perfoliatum*.

FEVER, SCORBU'IC, *Febris scorbutica*. The febrile movement, which sometimes accompanies scorbutus or scurvy.

FEVER, SEASONING, F., strangers'.

FEVER, SEC'ONDARY. A febrile condition which recurs in certain affections after having ceased; such as the secondary fever, which comes on at the time of the maturation of the various pustules, or as the eruption of scarlatina, &c., disappears.

FEVER, SEPTAN, *Febris septa'na*. An intermittent, whose paroxysms recur every six days, and consequently on the seventh.

FEVER, SEVEN DAY, see Relapse.

FEVER, SEXTAN, *Febris sexta'na*. A fever, which recurs every five days, and consequently on the sixth.

FEVER, SHIP, see Typhus.

FEVER, SIMPLE, *Febris Simplex*. Simple fever is that which has no predominant character—bilious, inflammatory, or nervous; and which is unaccompanied by any local determination, hyperæmia, or complication. It may be continued, remittent, or intermittent.

FEVER, SIMPLE CONTINUED. This is the most favourable form of continued fever, and has a tendency to wear itself out, provided only the *lædientia* be avoided. The prognosis is consequently favourable, and the treatment simple, consisting in perfect repose of body and mind, abstinence, and relieving the thirst by cold drinks.

FEVER, SINGUL'TOUS, *Febris singulto'sa* seu *lygmo'des* seu *lyngo'des*. Fever, accompanied with singultus or hiccough.

FEVER, SOLAR, Dengue—f. Spotted, Typhus gravior.

FEVER, STE'CORAL, *Febris stercora'lis*. Fever,

produced by an accumulation of fæces in the intestines.

FEVER, STOMACHIC, Gastric Fever.

FEVER, STRANGERS', *Acclimating* or *Seasoning fever*. Yellow, or remittent fever, which is endemic in certain places, and to which strangers are especially liable.

FEVER, SUBCONTINUAL, Remittent Fever.

FEVER, SUBINTRANS, *Febris subintrans*. An intermittent, in which one paroxysm is scarcely finished before the other begins.

FEVER, SUN, Dengue—f. Sweating, Sudor Anglicus.

FEVER, SYN'OPAL, *Febris synopalis* seu *mitis*, (F.) *Fidre synopale*. A variety of pernicious intermittent, in which there is, in every paroxysm, one or more faintings.

FEVER, SYNCHOID, Synchus.

FEVER, SYPHILITIC, *Febris syphilitica*. Fever, accompanying syphilis, or supposed to be owing to a syphilitic taint.

FEVER, TERTIAN, *Febris tertia*. A fever, whose paroxysm returns on the third day, and consequently every two days.

FEVER, THIRD DAY, Tertian.

FEVER, TRAG'IC, *Febris Tragica*. A low fever, in which the patient declaims like an actor during the delirium.

FEVER, TRAUMAT'IC, *Febris traumatica*. The fever, which supervenes on wounds or great surgical operations.

FEVER, TREMBLING, Intermittent fever—f. Typhoid, see Typhus—f. Typhoid, of India, Cholera—f. Typhous, Typhus—f. Varicellar, see Varioloid—f. Varioloid, see Varioloid.

FEVER, VERMINOUS, *Febris verminosa*, *Helmintophora*, Worm fever. Fever, produced by the presence of worms in the digestive tube, or accompanied by their expulsion.

FEVER, VERNAL, *Febris vernalis*. An intermittent or other fever occurring in the spring. Vernal intermittents were formerly considered salubrious.

"An ague in the spring
Is physic for a king."

FEVER, VESICULAR, Pemphigus.

FEVER, WALCHEREN, *Gall-sickness*. The remittents and intermittents to which the British troops were exposed, who were attached to the expedition to Walcheren, in 1809.

FEVER, WATER BRAIN, Hydrocephalus interus—f. Winter, see Tongue, black—f. Worm, Verminous F.—f. Worm, spurious, Fever, infantile remittent.

FEVER, YELLOW, *Febris flava*, *F.* seu *Pestis America*'na, *F. flava Americano*'rum, *F. tropica*, *F. continua putrida ictero*'des *Carolin*'ensis, *F. clo*'des *ictero*'des, *F. maligna bilio*'sa *Americ*'æ, *F. maligna flava Indiæ Occident*'alis, *F. toxica*, *Cholo*'sis *America*'na, *Ochro*'typhus, *Loimocholo*'sis, *Pestilen*'tia hæmagastrica, *Pestis occident*'alis seu *intertropica*, *Vomitus niger*, *Epan*'ctus *malignus flavus*, *Remittens ictero*'des, *Tritaph*'ya *America*'na, *Typhus ictero*'des, *Ochro*'typha, *Syn*'ochus *ictero*'des, *Endem*'ical *caus*'us of the *West Indies*, *Causus tropicus endem*'icus; Bilious remitting yellow fever, Malignant pestilential fever, *Fidre jaune d'Amérique*, *F. mato*'te, *F. gastro*'dynamique, *F. de la Barbade*, *F. de Siam*, *F. Jé*'rique, *F. Gastro*'hépatique, *Typhus mias*'matique *atarique putride* *jaune*, *T. jaune*, *T. d'Amé*'rique, *Vomito prieto*, *Vomito negro*, *Mal de Siam*, *Hæmagas*'tric *Fever* or *Pestilence*, *Black vomit*, *Bulam*, *Gibral*'tar and *Barcelo*'na *Fever*,—vulg., in the *West Indies*, *Mula*'to *Jack* and *Yellow Jack*. A very acute and dangerous febrile affection; so called, because complicated, in its second stage, with a jaundiced appearance, accompanied by vomiting of black matter. Sauvages, Cullen, and

others regard it as a variety of typhus; and Pinel, as a species of gastro-adyamic, or bilious putrid fever. It occurs, endemically, only within the tropics; but it has been met with epidemically in the temperate regions. Broussais regards the disease as gastro-enteritis, exasperated by atmospheric heat; so that it runs through its stages with much greater rapidity than the gastro-enteritis of our climates. The yellow colour of the skin, according to him, is owing to the inflammation of the small intestine,—and especially of the duodenum,—augmenting the secretion of the liver, and at the same time preventing its discharge into the duodenum. The pathology of this affection, as well as its origin, is still unsettled. The treatment must generally be of the most active nature at the onset; consisting in bleeding largely, and exhibiting mercury, so as to excite a new action, if possible:—the other symptoms being combated according to general principles. It must vary, however, according to the epidemic.

FEVER, YELLOW, MILD, see Relapse.

FEVERBUSH, *Laurus benzoin*, *Prinos*.

FEVER DROPS, WARBURG'S, see Bebeeru.

FEVERET, *Febricula*.

FEVERFEW, *Matricaria*.

FEVERFOUILLIE, *Matricaria*.

FE'VERISH, *Fe'verous*, *Fe'briens*, *Febrico*'us, (F.) *Fidreus*, from *febris*, 'fever.' That which causes fever or is febrif'icent or febrif'ic; as *feverish food*, *feverish diathesis*, &c. Also, the state of one labouring under fever, *Feversick*. If affected with intermittent fever, he is said to be a'garified in Suffolk, England.

FE'VERISHNESS, *Febricitas*, *Febriculos*'-ity. The state of having fever. A slight febrile disorder.

FEVEROUS, *Feverish*.

FEVERROOT, *Pterospara andromeda*.

FEVERSICK, see *Feverish*.

FEVERTREE, *Pinckneya pubens*.

FEVERWOOD, *Laurus benzoin*.

FEVERWORT, *Eupatorium perfoliatum*, *Triosteum*.

FEWS, *Sempervivum tectorum*.

FEY, *Psychorages*.

FEYK, *Fidgets*.

FIBER, *Castor fiber*.

FIBRA, *Fibre*—f. *Aurea*, *Coptis*—f. *Nervea*,

Nerve—fibre—f. *Sanguinis*, *Fibrin*.

FIBRÆ ARCIFORMES, see *Arciform*.

FIBRE, *Fibra*, *Is*, *Cledon*, *Filum*. An organic filament, of a solid consistence, and more or less extensible, which enters into the composition of every animal and vegetable texture. The simple or elementary fibre of the ancients, from a particular assemblage and arrangement of which every texture of the body was conceived to be constituted, seems entirely ideal. The moderns usually admit, with Haller and Blumenbach, three elementary fibres or tissues. 1. The cellular or laminated, formed chiefly of thin plates, of a whitish colour and extensible, which seems to consist of concrete gelatin. 2. The nervous, pulpy, or medullary, formed of a soft substance, contained in a cellular sheath, and consisting of albumen united to a fatty matter. 3. The muscular, composed of round filaments, of a grayish or reddish colour, and formed of fibrin. Chaussier has added to these the albuginous fibre, but it seems to differ from the cellular fibre only in greater condensation of the molecules. See *Fibrous*.

A very small or ultimate fibre is called a *Fibril*, *Fibrilla*.

FIBRE, ALBUGINEOUS, see *Albuginea*.

FIBRÆ LISSE, see *Muscular fibre*—f. *Striæ*, see *Muscular fibre*.

FIBRES, CONVERG'ING. Nervous fibres, whose office it is to associate different portions

of the nervous centres with each other. They form the *Commissures*.

FIBRES, DIVERG'ING. The fibres composing the columns of the medulla oblongata, which separate in their progress to the periphery of the cerebrum and cerebellum.—Gall and Spurzheim.

FIBRES OF REMAK. Fibres described by Remak as peculiar to the sympathetic nerve, but which Valentin considers to be neurilemma, and to consist of fibro-cellular bundles.

FIBREUX, Fibrous.

FIBRIL, see Fibre.

FIBRILLA, Fibril—f. Muscularis, Muscular fibre.

FYBRILLAR, *Fibrilla'ris*. Relating or appertaining to fibrils.

FIBRILLATION, *Fibrilla'tio*. The formation of fibrils.

FIBRIN or *Fib'rin*, *Fibrine*, *Fibri'na*, *Fibri'nē*, *Fibra sanguinis*, *Mate'ria fibro'sa*, *Lympha plastica*, *Matière fibreuse du sang*. An immediate animal principle—solid, white, and inodorous; insipid; heavier than water; without action on the vegetable blues; elastic, when moist; hard and brittle when dry. It enters into the composition of the chyle and the blood, and forms the chief part of the muscles of red-blooded animals. Muscular fibrin, *Syntonin*, *Musculin*, has been shown, however, to be different from that of the blood. In certain diseased actions, *Fibrin* or *Coagulable lymph*, *gluten*, is separated from the blood, and is found in considerable quantity on the surfaces of membranes, and in the cavities of the body. See *Liquor Sanguinis*.

Fibrin is likewise a proximate principle of vegetables, and differs but little in chemical composition from animal fibrin; nor does it differ much from albumen and casein.

Fibrin is very nutritious.

FIBRINATION, *Fibrina'tio*. The act of adding fibrin to the blood. The opposite to *defibrination*.

FIBRINOUS, *Fibrino'sus*, (F.) *Fibrineux*. That which is composed of fibrin, or has the appearance of fibrin.

FIBRO. In composition, appertaining or relating to fibrous tissue.

FIBRO-AREOLAR. Fibro-cellular.

FIBRO-BRONCHITIS. Fibrous or rheumatic inflammation of the bronchial tubes.

FIBRO-CARTILAGE, *Fibro-cartila'go*. An organic tissue, partaking of the nature of the fibrous tissue, and of that of cartilage. It is dense, resisting, elastic, firm, supple, and flexible. Fibro-cartilages are distinguished into.—1. *Membriform*, or those which serve as moulds to certain parts, as the *alæ nasi* and eyelids. 2. *Vaginiform*, or those which form sheaths for the sliding of tendons. 3. *Interarticular*, those which are met with in the movable articulations. 4. *Uniting*, (F.) *Fibro-cartilages d'union*, which form a junction between two bones, as the symphysis pubis. Fibro-cartilages are sometimes formed adventitiously, as the result of a morbid process in different organs.

FIBRO-CARTILAGES, TAPSAI, see Tarsus.

FIBRO-CARTILAGINES INTERVERTEBRALES, Intervertebral cartilages.

FIBRO-CELLULAR, *Fibro-cellula'ris*, *Fibro-areolar*, *Fibro-areola'ris*. That which partakes of the fibrous and of the cellular tissue, as the fascia transversalis.

FIBRO-CELLULAR TUMOUR, Tumour, fibro-cellular.

FIBROMA, Tumour, fibrous.

FIBRO-MUCOUS, *Fibro-muco'sus*, (F.) *Fibromucous*. Possessing the nature of fibrous and of mucous membranes. A term applied to fibrous membranes, which are intimately united

with other membranes of a mucous nature, as the pituitary membrane, the membrane of the urethra, &c.

FIBRO-PLASTIC, *Fibro-plast'icus*, (F.) *Fibroplastique*, from *fibra*, 'fibre,' and *πλασσω*, 'I form.' An epithet for a morbid formation, constituted of the elements of areolar tissue transformed, in part, into fibre.—Lebert.

FIBRO-PLASTIC TUMOUR, Sarcoma.

FIBRO-SEROTIS, *Fibro-sero'sus*, (F.) *Fibrosérous*. Possessing the nature of fibrous and serous membranes. Membranes, composed of a fibrous, and a serous sheet, intimately united:—as the *Dura Mater*, *Pericardium*, *Tunica albuginea testis*, &c.

FIBROID, Fibrous, Tumour, fibrous.

FIBROUS, *Fibro'sus*, *Fibroid*, (F.) *Fibreux*. Composed of fibres. Certain membranes, as the dura mater, periosteum, ligamentous capsules of the joints, &c., are fibrous. The fibrous system of Bichat includes the system of organs formed by the *albugineous fibre* of Ghaussier. It comprises, particularly, the periosteum and perichondrium; the articular capsules and ligaments; the tendons; the dura mater, pericardium, tunica sclerotica, tunica albuginea testis, outer membrane of the spleen, &c. Under *simple fibrous tissues*, *Tela fibro'sa*, certain writers have classed the *white* and *yellow fibrous tissues*, and areolar tissue. Both the yellow and the white may be detected in the areolar tissue. The *white* is said to exist alone in ligaments, tendons, fibrous membranes, aponeuroses, &c. The *yellow*, (F.) *Tissu jaune*, exists separately in the middle coat of the arteries, the chordæ vocales, ligamentum nuchæ of quadrupeds, &c. It differs from the white in possessing a high degree of elasticity, owing to the presence of a distinct principle, called by MM. Robin and Verdeil, *élastine*.

FIBROUS GROWTH, Tumour, fibrous—f. Matter of the Brain, see Cerebrum—f. Membranes, see Membranes, fibrous.

FIBULA, (for *Figibula* [?] from *figo*, 'to fix.') *Cruris radialis*, *Canna minor*, *Os peronæ*, *Peronæum*, *Facile minus*, *Sura*, *Arat'do minor*, *Fistula Cruris*, *Tibia min'ima*, *Oa tib'æ minus*, *Spellbone*, 'a clasp.' The splinter bone of the leg, (F.) *Péroné*. The long, small bone, situate at the outer part of the leg. The superior or tibial extremity of the fibula is rounded and forms the *caput* or *head*. It is articulated with the tibia. Its inferior or tarsal extremity is broader than the superior. It is articulated with the tibia and astragalus, and forms the *malleolus externus* or *outer ankle* by means of its *coronoid process*. The body of the bone has three faces, having more or less prominent edges. It is separated from the tibia by the interosseous space, and is developed by three points of ossification; one at the body, and one at each extremity. It prevents the foot from turning outwards.

FIBULAD, see Fibular aspect.

FIBULAR, Peroneal.

FIBULAR ASPECT. An aspect towards the side on which the fibula is situated.—Barclay, *Fibulad* is used by the same writer adverbially, to signify 'towards the fibular aspect.'

FIBULATIO, Infibulation.

FIC, Ficus.

FICAIRE, Ranunculus ficaria.

FICARIA, *Scrophularia aquatica*—f. *Communis*, *Ranunculus ficaria*—f. *Ranunculoides*, *Ranunculus ficaria*—f. *Verna*, *Ranunculus ficaria*.

FICATIO, Ficus.

FICITION, Physician.

FICOSA EMINENTIA, Ficus.

FICUS, *Sycē*, *Syl'cea*, *Sycum*, *Sycō'sis*, *Sycō'ma*, *Ficatio*, *Fico'sus Tumor*, *Fico'sa eminentia*, *Mari'sca*, (F.) *Fic*. A fleshy excrescence, often

soft and reddish, sometimes hard and scirrhous, hanging by a peduncle, or formed like a fig; occurring on the eyelids, chin, tongue, anus, or organs of generation. The fig seated on the last-mentioned parts are generally of a syphilitic character.

FICUS, F. Carica.

FICUS CARICA, *F. vulgaris* seu *communis*. The systematic name of the fig tree; (F.) *Figuer*. Ord. *Urticacæ*. *Carica*, *Ficus*, *Sycè*, *ovun*, (F.) *Figue*: the *fig*—is a pleasant fruit when ripe; as well as when dried in the state in which it is found in the shops—*Ficus*, (Ph. U. S.) It is used, at times, in place of a cataplasm; especially in gum-boils.

FICUS COMMUNIS, F. Carica—f. *Indiæ grana*, *Coccus cacti*—f. *Indica*, *Musa paradisiaca*, see *Caoutchouc*, *Carica papaya*, and *Lacca*—f. *Religiosa*, see *Lacca*.

FIDERIS, MINERAL WATERS OF. These waters, in the Canton of the Grisons, are strong, acidulous, and alkaline, and possess all the virtues of the class.

FIDGETS. Of doubtful etymology. *Dysphoria simplex*, *D. nervosa*, *Erethismus simplex*, *Tibutio*, *Creepers*, (Sc.) *Feyk*, (F.) *Fréttillement*. General restlessness and troublesome uneasiness of the nerves and muscles, with increased sensibility, and inability of fixing the attention, accompanied with a perpetual desire of changing the position. See *Agacement des Nerfs*.

FIDICINALES, *Lumbricales manus*.

FIEB, *Psychorages*.

FIEL, Bile—f. *de Bœuf*, Bile of the ox—f. *de Terre*, *Fumaria*.

FIELDWEED, *Anthemis cotula*, *Erigeron Philadelphicum*.

FIENTE, *Fimus*.

FIÈVRE, Fever—f. *d'Accès*, *Intermittent fever*—f. *Adynamique*, *Typhus*—f. *Algide*, see *Algidus*—f. *d'Amour*, *Fever, erotic*—f. *Angioténique*, *Synocha*—f. *Annuelle*, see *Annual Diseases*—f. *Ardent*, *Ardent fever*, *Synocha*—f. *Ataxique*, *Typhus*—f. *de la Barbade*, *Fever, yellow*—f. *Bulleuse*, *Pemphigus*—f. *des Camps*, *Typhus gravior*—f. *Catarrhale*, *Catarrh*—f. *Catarrhale épidémique*, *Influenza*—f. *Cérébrale*, *Cerebral fever*—f. *Cérébrale des Enfants*, *Hydrocephalus internus*—f. *Cérébro-spinal*, *Meningitis, cerebro-spinal*—f. *Cholérique*, *Choleric fever*—f. *de Concomption*, see *Consumption*—f. *Contingente inflammatoire*, *Synocha*—f. *Délirante*, *Fever, delirious*—f. *Double*, *Fever, double*—f. *Doublee*, *Fever, double*—f. *Entéro-mésentérique*, *Typhoid fever*—f. *Éphémère*, *Ephemera*—f.

Étique, *Hectic fever*—f. *Gastro-adyynamique*, *Fever, yellow*—f. *Gastro-hépatique*, *Fever, yellow*—f. *Hectique*, *Hectic fever*—f. *de Hongrie*, *Fever, Hungaric*—f. *d'Hôpital*, *Typhus gravior*—f. *Isotérique*, *Fever, yellow*—f. *Inflammatoire*, *Synocha*—f. *Intermittente*, *Intermittent fever*—f. *Intermittente paludéenne*, see *Elodes*—f. *Irritative*, *Synocha*—f. *Jaune d'Amérique*, *Fever, yellow*—f. *Lactée*, *Fever, milk*—f. *de Lait*, *Fever, milk*—f. *Laitieuse*, *Fever, milk*—f. *Larvée*, *Fever, masked*—f. *Lenticulaire*, *Typhus gravior*—f. *du Levant*, *Plague*—f. *des Marais*, *Intermittent fever*—f. *Maremmatique*, see *Elodes*—f. *Matelote*, *Fever, yellow*—f. *Méningo-gastrique*, *Fever, gastric*—f. *Mésentérique*, see *Typhus*—f. *de la Moisson*, *Fever, gastric*—f. *Morbilleuse*, *Rubeola*—f. *Nerveuse*, *Typhus mitior*—f. *Noxocomiale*, *Typhus gravior*—f. *Ortiée*, *Urticaria*—f. *Oscitante*, *Oscitant fever*—f. *Paludéenne*, see *Elodes*—f. *Périodique*, *Intermittent fever*—f. *Péripneumonique*, *Pneumonia*—f. *Pernicieuse délirante*, see *Delirious*—f. *Pleurétique*, *Pleurisy*—f. *des Prisons*, *Typhus gravior*—f. *Pneumonique*, *Pneumonia*—

f. *Pourprée*, *Scarlatina*—f. *Puerpérale*, *Puerperal*, see *Peritonitis*—f. *Quarte*, *Quartan*—f. *Quotidienne*, *Quotidian*—f. *à Reclute*, see *Relapse*—f. *Rémittente*, *Remittent fever*—f. *Rhumatisme*, *Rheumatism*, *acute*—f. *Rouge*, *Roseola*, *Scarlatina*—f. *Sanguine*, *Synocha*—f. *Semitierce*, *Hemitritea*—f. *de Siam*, *Fever, yellow*—f. *Suante*, *Sudor Picardicus*—f. *Syncope*, see *Fever synoptal*, and *Syncope*—f. *Tierce*, *Tertian fever*—f. *Typhoïde*, see *Typhus*—f. *Vésiculaire*, *Pemphigus*.

FIÈVREUX, *Feverish*.

FIG, see *Ficus carica*—f. *Indian*, *Cactus opuntia*.

FIGUE, see *Ficus carica*.

FIGULIER, *Ficus carica*—f. *d'Inde*, *Cactus opuntia*.

FIGURA VENOSA, *Circulus venosus*.

FIGURATIO, *Imagination*.

FIGURE, *Countenance*.

FIGURÉ, (F.) An epithet for a compressive bandage, applied over the head after bleeding from the frontal vein. It has also been called *bandage royal*.

FIGWORT, *Scrophularia nodosa*—f. *Water, greater*, *Scrophularia aquatica*.

FILACEOUS, *Filamentous*.

FILA NERVEA, *Nerve-fibres*—f. *Spermatica*, *Spermatozoa*.

FILAIRE DE MÉDINE, *Dracunculus*.

FIL'AMENT, *Filum*, *Filamentum*, from *filum*, 'a thread.' This word is used synonymously with *fibril*; thus, we say, a *nervous* or *cellular filament* or *fibril*. Also, the glairy, thread-like substance which forms in the urine in some diseases, and which depends on a particular secretion from the mucous membrane of the urinary passages.

FILAMEN'TOUS, *Filamento'sus*. *Filaceous*; threadlike; filiform. Same etymon. Containing threadlike substances, as the *tunica filamentosa* or decidua:—*Filamentous urine*; urine containing threadlike substances.

FILAMENTUM, *Frænum*.

FILARIA GUINEENSIS, *Dracunculus*—f. *Hominis bronchialis*, see *Worms*—f. *Medinensis*, *Dracunculus*—f. *Oculi*, see *Worms*.

FILBERT, *Corylus avellana*.

FILE, *DENTAL*, *Lima dentaria*.

FILELLUM, *Frænum*.

FILET (DE LA LANGUE), *Frænum*—f. *de la Verge*, *Frænum penis*—f. *Opération du*, see *Frænum*.

FILETUM, *Frænum linguae*.

FILICULA, *Polypodium filix mas*—f. *Dulcis*, *Polypodium vulgare*.

FILIFORM, *Filiformis*, from *filum*, 'a thread,' and *forma*, 'form.' Having the shape of a thread; as the "filiform papillæ of the tongue;" a "filiform pulse." See *Papillæ*.

FILING, *Limatio*.

FILIPENDULA, *Spiræa filipendula*.

FILIUS ANTE PATREM, *Tussilago*.

FILIX FEMINEA, *Pteris aquilina*—f. *Florida*, *Osmunda regalis*—f. *Mas*, *Polypodium filix mas*—f. *Non ramosa dentata*, *Polypodium filix mas*—f. *Nymphæa*, *Pteris aquilina*—f. *Pinnata*, *Polypodium filix mas*—f. *Veneris*, *Adiantum pedatum*.

FILLE, *Girl*.

FILLET, *Fascia*, *Laqueus*—f. of the *Corpus Callosum*, *Gyrus fornicatus*.

FILLING. A term employed in dentistry for the operation of stuffing the cavity of a carious tooth with some indestructible substance, to prevent the access of extraneous matters—*Odontoplero'sis*. See *Plugging*.

FILTRA'TION, *Filtra'tio*, *Percola'tio*, *E'this'sis*, *E'this'mus*. Same etymon as the next. A

pharmaceutical operation, which consists in passing a fluid through a filter or strainer, for the purpose of clarifying it. In ancient physiology, it meant the action by which the different humours of the body are separated from the mass of the blood.

FILTRUM. A filter. (F.) *Filtre*;—from *fentre*, 'felt or fulled cloth,' which was used to form it. Any porous material; such as sand, some kinds of freestone, powdered charcoal, pounded glass, flannel, unsized paper, &c., through which a fluid is passed for the purpose of separating it from the matters suspended in it.

FILUM. Filament—f. Muscular, Muscular fibre—f. *Tæniaforme*, *Tæniola*.

FILUM TERMINALE. A slender ligament, prolonged from the nervous sheath, formed by the spinal pia mater, which descends through the centre of the cauda equina, and is attached to the dura mater, lining the canal of the coccyx.

FIMBRIA, Pavar'na. A band, a fringe; as the fimbria or fimbriated or fringed extremity of the Fallopian tube.

FIMBRIÆ CARNOSÆ COLI. Appendiculae pinguedinosæ—f. *Tubarum Fallopii*, see Tube, Fallopian.

FIMBRIATED, see Fimbria.

FIMUS, Bor'borus, Onthus, Stercus. (F.) *Fiente*. Dung, excrement.

FIMUS seu STERCUS AN'SERIS, Goose-dung, was applied as a poultice to the feet in malignant fever. See *Chenocoprus*.

FIMUS EQU'NI, Stercus equi non castrati. *Stone horse-dung* was once thought antiepileptic.

FIMUS VACCÆ, Cow-dung, was employed as a cataplasm, especially in gout.

* **PINCKLE,** Anethum.

FINGER, Digitus—f. Middle, *Medius digitus*—f. Ring, *Annular finger*—f. Stall, *Digitale*.

PINIS ASPERÆ ARTERIÆ, Larynx.

PIOLE, Phiala.

PIPPLE, Lip, under.

PIR, BALSAM, Pinus balsamea.

PIR-MOSS, UPRIGHT, Lycopodium selago.

PIR, SCOTCH, Pinus sylvestris—f. Spruce, *Norway*. *Pinus abies*.

PIR TREE, SILVER, EUROPEAN, Pinus picea.

PIR, YEW-LEAVED, Pinus abies.

FIRE DAMP, Hydrogen, carburetted—f. Persian, *Anthracion*—f. St. Anthony's, *Erysipelas*.

FIRE WEED, Erechtites hieracifolia, Senecio.

FIRING, Cauterization.

FIRST INTENTION, see Intention.

FISHSKIN, Ichthyosis.

FISH-TONGUE. (F.) Langue de Carpe. An instrument—so called from its shape—used by some dentists for the removal of the dentes sapientiae.

FISSICULA'TIO, from fissiculare, 'to cut off,' 'open,' 'make incisions.' An old word for an opening made with a scalpel.

FISSIPARITE, see Generation.

FISSIPAROUS, see Generation.

FISSURA, Fissure—f. *Canalis Fallopii*, see *Aqueductus Fallopii*—f. *Capillorum, Distrix*—f. *Contrajacens, Contra-fissura*—f. *Cerebri longitudinalis, Fissure, longitudinal, of the Brain*—f. *Interlobularis, Interlobular fissure*—f. *Magna vulvae, see Rima*—f. *Palpebrarum, see Palpebra*—f. *Pilaris, Trichismus*.

FISSURA LONGITUDINALIS, ANTE'RIOR ET POSTERIOR. Two vertical fissures in the median line, in front of, and behind, the medulla oblongata, which divide it superficially into two symmetrical lateral columns.

FISSURA TRANSVERSA MAGNA CEREBRI, Fissure, transverse, of the Brain.

FISSURE, Fissu'ra, Schisma, Scissu'ra, Rhegē, Rhegma, Rhegnus, from findere, fissum, 'to

cleave,' a long and narrow cleft or opening in a bone—*Rhagē, ραγν, Cœasma, κτασπα.* (F.) *Fissure, Fente*.

FISSURE has various acceptations: 1. A fracture, *Catag'ma fissu'ra*, in which the bone is cracked, not separated, as in fracture; 2. A narrow, long, and superficial solution of continuity, around the external openings of the mucous membranes. A sort of chaps—(F.) *Gercure*—observed on the hands, particularly on the callous hands, of workmen, in certain mechanical employments; 3. Small chapped ulcerations, sometimes noticed in young children, owing to the contact of the feces and urine with the fine, delicate skin of the thighs, nates, and genital organs; 4. Clefts of a more or less deep nature, occurring on the genital organs in the vicinity of the anus, in those labouring under syphilis. These are usually called *rhagades*. See *Monster*.

FISSURE OF BICHAT, Fissure, transverse, of the Brain—f. *Capillary, see Pilatio*.

FISSURE, CENTRAL. The aggregate of the cavities or ventricles of the brain. Meckel considers them but one cavity, in the form of a cross.

FISSURE OF GLASER, Fissure, glenoid—f. of *Glaserius, F. Glenoid*.

FISSURE, GLENOID, Fissure of Glaser or Glase'rius, (F.) Fissure ou Scissure de Glaser, Fente glenoïdale ou Fêlure de Glaser, divides the glenoid cavity of the temporal bone into two parts, and gives passage to the chorda tympani. &c.

FISSURE OF THE HELIX. A small vertical fissure of the helix of the ear, a little above the tubercle for the attachment of the *atrahens aurem* muscle.

FISSURE, INFRAORBITAR, Suborbital fissure.

FISSURE, LONGITUDINAL OF THE BRAIN, Fissu'ra cerebri longitudinalis. The space which separates the two hemispheres of the brain.

FISSURE, ORBITAR, see Orbital fissure—f. *Orbital, inferior, Spheno-maxillary fissure*—f. *Orbital, superior, Sphenoidal fissure*—f. *Portal, see Liver*.

FISSURE OF ROLAN'DO. A transverse fissure placed between two superior cerebral convolutions, which are met with above the fissure of Sylvius.

FISSURE, SEMILUNAR. A notch at the anterior edge of the cerebellum, where it receives fibres which connect it to the cerebellum and mesocephalon.

FISSURE OF SYLVIVS, Fissu'ra seu Fossa Magna Syl'vii. A deep, narrow sulcus, which ascends obliquely backwards from the temporal ala of the sphenoid bone, near to the middle of the parietal bone, and which parts the anterior and middle lobes of the cerebrum on each side.

FISSURE OF THE TRAGUS. A fissure on the anterior surface of the tragus of the ear.

FISSURE, TRANSVERSE, OF THE BRAIN, Great transverse fissure, Fissure of Bichat, Fissu'ra seu Rima transver'sa magna cerebri. A fissure, which passes beneath and behind the edge of the middle lobe of the brain, and extends beneath the hemisphere of one side to the same point of the opposite side.

FIST, Sax. fýst, πνγν, Pygmē, Pugnus, (F.) Poing. The clenched hand.

FISTULA, 'a pipe or reed,' Syrinx, Syrin'ga, Aulos—when of a small size, Auti'cos. A solution of continuity, of greater or less depth and sinuosity; the opening of which is narrow, and the disease kept up by an altered texture of parts, so that it is not disposed to heal. A fistula is incomplete or blind, when it has but one opening; and complete when there are two, the one communicating with an internal cavity, the other externally. It is lined, in its whole course, by a membrane which seems analogous to mucous

membranes. *Incomplete fistula* may be *internal* or *external*. The former are those which open *internally*; the latter those which open *externally*. *External incomplete fistula* are kept up by caries or necrosis of bones, by extraneous bodies in any of the living textures, or by purulent cavities, the walls of which have not become united. *Internal incomplete fistula* generally soon become complete, since the discharge that escapes from them into the cavities into which they open has a constant tendency to make its way outwardly, and soon occasions ulceration of the integuments. *Fistulae* have received different names, according to the discharge which they afford, and the organs in which they are seated,—as *lachrymal*, *biliary*, *salivary*, *synovial*, *urinary*—*Fistula lachrymalis*, *Urticaria*. The great object of treatment, in fistulous sores, is to bring on an altered condition of the parietes of the canal, by astringent or stimulating injections, caustics, the knife, pressure, &c. Those which are dependent on diseased bone, cartilage, tendon, &c., do not heal until after the exfoliation of the diseased part. *Fistulae* of excretory ducts are produced either by an injury of the duct itself or by the retention and accumulation of the fluids to which they have to give passage. Thus, *Fistula lacrymalis*, *Dacryosyrinx*, *Emphragma lacrymale*, *Hydrops sacci lacrymalis*, *Dropsy of the lachrymal sac*, commonly proceeds from the obliteration of the nasal ducts, or from atony of the lachrymal sac; which circumstances prevent the tears from passing into the nostrils.

FISTULA IN ANO, *Archosyrinx*, generally occurs from some mechanical pressure or impediment. The principal indication in the treatment of these fistulae of the excretory canals being to put a stop to the constant discharge of the secretions, &c., through the preternatural channel, the fistulous passage is at times laid open, and a communication established with the natural excretory canal; at others, strong pressure is employed to procure its obliteration.

FISTULA BELLINIANA, Uriniferous tube—*f. Cibalus*, *Oesophagus*—*f. Cruris*, *Fibula*—*f. Duræ matris*, *Sinus of the dura mater*—*f. Lacrymalis*, see *Fistula*—*f. Mammæ*, *Mastosyrinx*—*f. Nervorum*, *Neurilemma*—*f. Sacra*, *Medulla spinalis*, *Vertebral column*—*f. Spiritualis*, *Trachea*—*f. Urinaria*, *Urethra*—*f. Ureterum renium*, see *Calix*—*f. Urinæ*, see *Fistula*—*f. Ventriculi*, *Oesophagus*.

FISTULES STERCORAIRES, see *Stercoraceous*.

FISTULOUS, *Fistulosus*, *Syringicus*, *Syringoides*, (F.) *Fistulens*. Relating to, or resembling, a fistula; as 'a fistulous opening.'

FIT, *Pes*.

FIT-ROOT, *Monotropa uniflora*.

FITS, NINE DAY, *Trismus nascentium*.

FIVE FINGERS, *Panax quinquefolium*.

FIXATIO MONONCEA, *Melancholey*.

FIXED, *Fixus*, from *figere*, *fixum*, 'to fasten.'

A body not capable of being volatilized by fire is said to be fixed. Thus, we say *fixed oils*, in contradistinction to *volatile oils*.

FIXEN, MINERAL WATERS OF. This spring is four leagues from Waldsassen, in Bavaria. It contains carbonic acid, holding in solution carbonates of lime and magnesia, chlorides of lime and magnesia, carbonate of soda and silica. In Bavaria, it replaces the Seltzer water.

FIXI DENTES. The teeth of the second dentition.

FLABELLATION, *Flabella'tio*, from *flabellum*, *flabellatum*, (from *flabrum*, 'a blast of wind,') 'to agitate the air.' An operation recommended by Ambrose Paré, which consists in keeping fractured limbs cool, as well as the

dressings surrounding them, by the renewal of the air around them, either by the use of a fan, or the repeated change of position of the parts affected.

FLABELLUM ET VENTILABRUM *CORDIS*, *Pulmo*.

FLACCIDITY, *Flacciditas*, from *flaccidus*, 'flabby,' 'soft.' Softness of a part, so as to offer little resistance on pressure.

FLAG, BLUE, *Iris versicolor*—*f. Blue*, slender, *Iris Virginica*—*f. Dwarf*, *Iris lacustris*—*f. Myrtle*, *Acorus calamus*—*f. Root*, *Acorus calamus*.

FLAGELLATIO, *Mastigosis*.

FLAMBE, *Iris Germanica*—*f. Bâtard*, *Iris pseudacorus*.

FLAMBOISE, *Ignis sylvaticus*.

FLAME, VITAL, *Vital principle*.

FLAMED, *Inflamed*.

FLAMMA, *Fleam*—*f. Cordis*, *Biolychnium*—*f. Vitalis*, *Biolychnium*.

FLAMME, *Fleam*—*f. Vitale*, *Vital principle*.

FLAMMETTE, *Fleam*.

FLAMMON, *Lachesis rhombeata*.

FLAMMULA, *Fleam*, *Ranunculus ficaria*—*f. Cordis*, *Biolychnium*—*f. Jovis*, *Clematis recta*—*f. Vitalis*, *Animal heat*, *Biolychnium*, *Vital principle*.

FLANCS, *Flanks*.

FLANKS, *l'ia*, *l'ea*, *La'gones*, *Lap'ara*, *Cenoe'nes*, (Prov.) *Liste*, (F.) *Flancs*, *Les Iles*. The regions of the body which extend, on the sides, from the inferior margin of the chest to the *crista ili*.

FLAP, (F.) *Lambeau*. A portion of the soft parts of the body separated from those beneath, but still attached by the base. Hence there may be 'flap wounds,' (F.) *Plaies à lambeaux*, and 'flap operations,' (F.) *Opérations à lambeaux*. When the flap is taken from one part, and applied to another, in plastic operations, the operation is said to be by migration of the flap, (F.) *Migration du Lambeau*. At times, the flap is made to slide over the part to which it has to adhere:—the operation is then said to be by the sliding of the flap, (F.) *Glissement du Lambeau*. The flap is also, at times, rolled, inverted, &c., (F.) *Roulement ou Inversion du Lambeau*.

FLAP-DOCK, *Digitalis*.

FLAP OPERATION OF AMPUTATION, *Amputation à lambeau*.

FLAT TOP, *Vernonia Novboracensis*.

FLATUARIUS, *Alchymist*.

FLATUEUX, *Windy*.

FLATULENCE, *Flatulentia*, *Flatus*, *Flatus-itis*, *Aërifluxus*, *Pneumato'sis*, *Pneumatosia ventriculi et enterica*, *Bdes'ma*, *Hyperpneus'tia*, *Clydon*, *Physa*, *Polyphys'ia*, *Limo'sis* *Flatus*, *Ereux'ia*, *Bombus*, *Flatuos'ity*, *Wind*, *Windiness*, *Ventos'ity*, *Vapour*, (F.) *Ventosité*, *Flatuosité*, *Vents*; from *flō*, *flatum*, 'to blow.' Wind emitted from, or accumulated in, the digestive tube especially.

FLATULENT, *Windy*.

FLATULENTIA, *Flatulence*.

FLATUOSITÉ, *Flatulence*.

FLATUOSITY, *Flatulence*.

FLATUS, *Crepitation*, *Flatulence*—*f. Furiosus*, *Ambulo-flatulens*—*f. Spinae*, *Spina ventosa*.

FLAVEDO CORTICUM CITRI, see *Citrus medica*.

FLAVOUR, from (F.) *flairer*, 'to smell.' The quality of a sapid body, which is appreciated by the taste and smell combined, and more especially by the latter. Some physiologists consider that flavour is effected through the smell alone.

FLAVUS, *Yellow*.

FLAX, COMMON, *Linum usitatissimum*—*f. Purging*, *Linum catharticum*—*f. Seed*, see *Linum*

usitatissimum—*f.* Seed tea, Infusum lini compositum—*f.* Toad, Antirrhinum linaria.

FLEA, COMMON, Pulex irritans.

FLEABANE, GREAT, *Conyza squarrosa*—*f.* Various-leaved, *Erigeron heterophyllum*—*f.* Canada, *Erigeron Canadense*—*f.* Narrow-leaved, *Erigeron Philadelphicum*—*f.* Philadelphia, *Erigeron Philadelphicum*—*f.* Skevish, *Erigeron Philadelphicum*.

FLAM, *Flamma, Flam'mula, Schaste'rium, Fosso'rium, Phlebot'omum*. A surgical instrument used for the operation of phlebotomy. (*F.*) *Flamme, Flammette*. It consists of a small metallic box, containing a spear-pointed cutting instrument, which, by means of a spring, can be forced into the vein. It is much used in Germany and some other European countries, and is not unfrequently employed in America; but is scarcely ever seen in France or Great Britain.

FLEAWORT, *Plantago psyllium*.

FLÉCHISSEUR, *Flexor*—*f.* *Court commun desorteils, Flexor brevis digitorum pedis*—*f.* *Court du petit doigt, Flexor parvus minimi digiti*—*f.* *Court du petit orteil, Flexor brevis minimi digiti pedis*—*f.* *Profond des doigts, Flexor profundus perforans*—*f.* *Sublime des doigts, Flexor sublimis perforatus*—*f.* *Superficiel des doigts, Flexor sublimis perforatus*—*f.* *Court du gros orteil, Flexor brevis pollicis pedis*—*f.* *Grand commun desorteils, Flexor longus digitorum pedis profundus perforans*—*f.* *Long commun desorteils, Flexor longus digitorum pedis profundus perforans*—*f.* *Long du gros orteil, Flexor longus pollicis pedis*—*f.* *Court du ponce, Flexor brevis pollicis manûs*.

FLECK, Abrasion.

FLECTENS PAR LUMBORUM, *Quadratus lumborum*.

FLEGMEN, Flemen.

FLEMEN, *Flegmen*. A tumour about the ankles. Also, a chap on the feet and hands.

FLERECIN, Gout.

FLESH, *Caro, Sarc, (F.) Chair*. Every soft part of an animal is so named; but more particularly the muscles, which are called *muscular flesh*.

FLESH, PROUD, Fungosity.

FLESHY, Corpulent.

FLETUS, Lacrymatio.

FLEUR DE COUCOU, *Primula veris*—*f.* *Sainte-Catherine, Nigella*—*f.* *de Sainte-Jacques, Thlaspi bursa*—*f.* *de Tous les mois, Calendula officinalis*.

FLEURS, Flowers, Menses—*f.* *de Benjoin, Benjamin*—*f.* *Blanches, Leucorrhœa*—*f.* *de Muscade, Mace*—*f.* *de Soufre, Sulphur sublimatum*.

FLEXIBILITY, *Flexibilit'as*, from *flectere, flexum*, 'to bend.' Capability of being bent. A physical property of the tissues, which varies greatly according to the structure. The tendons exhibit this property in a marked manner.

FLEXIO, *Campsis, Flexion*.

FLEXION, *Flexio, Campé*. Same etymon. The state of being bent. The action of a flexor muscle.

FLEXOR. Same etymon. (*F.*) *Fléchisseur*. A muscle, whose office it is to bend certain parts.

FLEXOR BREVIS DIGITORUM PEDIS PERFORATUS, *F. Sublimis* seu *brevis seu digito'rum brevis seu perforatus pedis, Perforatus seu flexor secun'di interno'dii digito'rum pedis, (F.) Calcanéo-sous-phalangien commun, —Calcanéo-sous-phalangien commun, —(Ch.), Muscle court fléchisseur commun desorteils*. A muscle, placed at the middle of the sole of the foot. It is narrower and thicker behind than before, where it is divided into four portions. It arises from the posterior part of the inferior surface of the os calcis, and is inserted at the inferior surface of the second phalanx of the last four toes. It bends

the second phalanges of the toes on the first, and the first on the metatarsal bones; in this manner augmenting the concavity of the vault of the foot.

FLEXOR BREVIS MINIMI DIGITI, *F. Parvus minimi digiti*.

FLEXOR BREVIS MINIMI DIGITI PEDIS, *Para'flexor minor, (F.) Court fléchisseur du petit orteil, Tarso-sous-phalangien du petit orteil—(Ch.)*. A muscle, situate at the anterior and outer part of the sole of the foot. It arises from the posterior extremity of the fifth metatarsal bone, and is inserted into the posterior part of the first phalanx of the little toe, which it bends.

FLEXOR BREVIS POL'ICIS MANÛS, *F. secun'di interno'dii, F. primi et secun'di ossis pol'icis, Thenar, (F.) Court fléchisseur du ponce, Carpo-phalangien du ponce—(Ch.), Demi-interosseux du ponce*. A muscle, situate at the outer part of the palm of the hand. It is divided into two portions by the tendon of the *Flexor longus pollicis*. It arises from the os magnum, the anterior annular ligament of the carpus and the third metacarpal bone; and is inserted into the superior part of the first phalanx of the thumb, and into the two ossa sesamoidea at the articulation of the first phalanx with the first metacarpal bone. Its use is to bend the first phalanx of the thumb on the first metacarpal bone, and the latter upon the trapezium.

FLEXOR BREVIS POL'ICIS PEDIS, *F. brevis seu hallucis seu brevis pol'icis, (F.) Tarso-phalangien du ponce, Court fléchisseur du gros orteil, Tarso-sous-phalangien du premier orteil—(Ch.)*. It is situate at the anterior and inner part of the sole of the foot; is thin and narrow behind, thick and divided into two portions before. It arises from the inferior part of the os calcis and the last two cuneiform bones, and is inserted at the inferior part of the base of the first phalanx of the great toe, and into the two sesamoid bones of the corresponding metatarso-phalangeal articulation. It bends the first phalanx of the great toe on the first metacarpal bone.

FLEXOR CARPI RADIALIS, *Palmaris magnus*—*f.* *Carpi ulnaris*, see *Cubital (muscles)*—*f.* *Hallucis, F. brevis pollicis pedis*—*f.* *Hallucis longus, F. longus pollicis pedis*.

FLEXOR LONGUS DIGITORUM PEDIS PROFUNDUS PERFORANS; *Perforans seu flexor profundus, Perodactyleus, Peronodactylus, Peronodactylæus, Peronodactylus, Flexor digitorum longus seu Perforans pedis, Perforans seu flexor tertii interno'dii digito'rum pedis; (F.) Tibio-phalangien—(Ch.), Grand ou long fléchisseur commun desorteils*. A muscle, situate at the posterior and deep-seated part of the leg. It is broader at its middle than at its extremities, the inferior of which is divided into four portions. It arises from the posterior surface of the tibia, and its tendons are attached to the posterior part of the lower surface of the three phalanges of the last four toes. It bends the three phalanges on each other, and the toes on the metatarsus, and extends the foot on the leg.

The *Accessorius Flexoris Longi Digito'rum Pedis, Carpo quadrata Syl'vii, C. accessorius, Massa car'nea Jacob'i Syl'vii, Planta'ria verus, (F.) Accessoire du long fléchisseur commun desorteils, Carrée (Chair)*, is a small muscle of the sole of the foot, which passes obliquely from the os calcis to the outer edge of the flexor longus, whose force it augments, and corrects its obliquity.

FLEXOR LONGUS POL'ICIS MANÛS, *F. longus pollicis, F. tertii interno'dii seu longissimus pol'icis; (F.) Radio-phalangien du ponce—(Ch.)*. Situate at the anterior and profound part of the forearm. It arises from the upper three quarters of the anterior surface of the radius and interos-

aeous ligament, and is inserted, by a tendon, into the anterior surface of the last phalanx of the thumb. It bends the second phalanx of the thumb on the first; the first on the corresponding metacarpal bone, and this upon the radius. It can, also, bend the hand on the forearm.

FLEXOR LONGUS POLICIS PEDIS, *F. Hallucis seu Pollicis longus*, (F.) *Péronéo-phalangien du gros orteil*, *Long fléchisseur du gros orteil*, *Péronéo-sous-Phalangien du ponce*,—(Ch.) It is situate at the posterior and profound part of the leg. It arises from the posterior surface of the fibula and the interosseous ligament, and is inserted, by means of a long tendon, into the inferior part of the first phalanx of the great toe. It bends the third phalanx on the first, and this upon the corresponding metatarsal bone. It augments the concavity of the sole of the foot, and extends the foot on the leg.

FLEXOR PARVUS MINIMI DIGITI, *F. brevis minimi digiti manus*, *Abductor minimi digiti*, *Hypophenar Riola'ni seu minimi digiti*, (F.) *Carpophalangien du petit doigt*,—(Ch.) *Court fléchisseur du petit doigt*. It arises from the anterior annular ligament of the carpus and the process of the os unguiforme, and is inserted at the inner side of the superior extremity of the first phalanx of the little finger. It bends the first phalanx of the little finger.

FLEXOR PERFORANS, *F. profundus perforans*—*f. Perforatus*, *F. sublimis perforatus*—*f. Primi internodii*, *Opponens pollicis*—*f. Primi internodii digitorum manus*, *Lumbicalis manus*—*f. Primi et secundi ossis pollicis*, *F. brevis pollicis manus*.

FLEXOR PROFUNDUS PERFORANS, *F. Profundus seu Perforans seu tertii internodii digitorum manus*, *Perforans manus*; (F.) *Cubito-phalangien commun*—(Ch.) *Fléchisseur profond des doigts*. A thick, flat, long muscle, seated beneath the *Flexor sublimis perforatus*. Its upper extremity is simple, and arises from the anterior surface of the ulna and from the interosseous ligament. Its inferior extremity terminates by four tendons, which, after having passed through slits in the *sublimis*, are inserted into the anterior surface of the last phalanges of the four fingers. It bends the third phalanges on the second, and, in other respects has the same use as the *flexor sublimis perforatus*.

FLEXOR SECUNDI INTERNODII DIGITORUM PEDIS, *F. brevis digitorum pedis*—*f. Tertii internodii*, *F. longus pollicis manus*—*f. Tertii internodii digitorum manus*, *F. profundus perforans*—*f. Tertii internodii digitorum pedis*, *F. longus digitorum pedis profundus perforans*—*f. Ossis metacarpi pollicis*, *Opponens pollicis*—*f. Perforatus pedis*, *F. brevis digitorum pedis*—*f. Sublimis*, *F. brevis digitorum pedis*.

FLEXOR SUBLIMIS PERFORATUS, *F. Perforatus*, (F.) *Fléchisseur sublime ou superficiel des doigts*, *Epitrochlo-phalangien commun*,—(Ch.) It is a thick, flat, muscle, seated at the anterior part of the forearm. Its upper extremity, which is simple, arises from the internal condyle of the os humeri—from the coronoid process of the ulna, and from the anterior edge of the radius. Its lower extremity divides into four tendons, which slide under the anterior annular ligament of the carpus, and are inserted into the second phalanges of the last four fingers, after having been slit to allow the tendons of the *flexor profundus* to pass through them. This muscle bends the second phalanges on the first; these on the carpal bones, and the hand on the forearm.

FLEXURA, *Curvature*—*f. Iliaca*, *Sigmoid flexura*—*f. Sigmoidea*, *Sigmoid flexure*.

FLICK, *Epiploon*.

FLIXWEED, *Sisymbrium sophia*.

FLOATING, *Hæmorrhagia*.

FLOCCI, see *Villous membranes*—*f. Volitantes*, see *Metamorphopsia*.

FLOCCILATION, *Carphologia*.

FLOCCILEGIUM, *Carphologia*.

FLOCCITATION, *Carphologia*.

FLOCCORUM VENATIO, *Carphologia*.

FLOCCULI, see *Villous membranes*.

FLOC'ULUS; diminutive of *flocus*, 'a lock of wool'—*Pneumogast'ric lob'ule*, *Lob'ulus pneumogast'ricus*. A long and slender prominence, extending from the side of the vallecula around the corpus testiforme to the crus cerebelli, lying behind the filaments of the pneumogastric nerves.

FLOR DE MISTELA, see *Mistura*.

FLORENCE, **CLIMATE OF**. This agreeable Italian city is by no means a favourable residence for the phthisical invalid. Sir James Clark affirms, indeed, that he does not know any class of invalids for whom Florence offers a favourable residence. It is subject to sudden vicissitudes of temperature, and to cold, piercing winds during the winter and spring.

FLORES BENZOËS, Benjamin, flowers of—*f. Boracis*, *Boracic acid*—*f. Cosso*, see *Hagenia Abyssinica*—*f. Macidis*, see *Myristica moschata* *f. Macis*, *Mace*—*f. Martiales*, *Ferrum ammoniatum*—*f. Salis ammoniaci martiales*, *Ferrum ammoniatum*.

FLORIDA, **CLIMATE OF**, see *Saint Augustine*.

FLORIDA, **WATERS OF**. Near Long Lake, in Florida, United States, which communicates with St. John's River by a small creek, there is a vast fountain of hot mineral water, issuing from a bank of the river. From its odour it would seem to be sulphureous.

FLORION, *Influenza*.

FLOS, *Anthos*. A flower. Also, the finest and noblest part of the body, and virginity.

FLOS, *Jovis*, *Crocus*—*f. Lactis*, *Cremor lactis*—*f. Salis Soda*, *subcarbonate of*—*f. Sanguineus monardi*, *Tropeolum majus*—*f. Trinitatis*, *Viola tricolor*—*f. Virginitatis*, *Hymen*.

FLOUR, *COLD*, *Pinoli*.

FLOUR, **PATENT**, *JONES'S*. A farinaceous preparation, which is said to consist of wheat-flour, with tartaric acid and carbonate of soda.

FLOUR, **POTATO**, see *Solanum tuberosum*.

FLOWER DE LUCE, *Iris Germanica*.

FLOWERS, *Flores*, (F.) *Fleurs*. The ancient chymists gave this name to different solid and volatile substances obtained by sublimation. The term is not yet entirely banished from chymical and medical language, as *Flowers of Benjamin*, *Flowers of Sulphur*, &c.

FLOWERS, *Menses*.

FLOWERS, **FOUR CARMIN'ATIVE**, *Quatuor flores carminat'ivi*, were chamomile, dill, fever-few, and melilot.

FLOWERS, **FOUR COR'DIAL**, *Quatuor flores cordiales*, were formerly, borage, bugloss, roses, and violets.

FLUCTUATIO, *Fluctuation*—*f. Aurium*, *Tinnitus aurium*.

FLUCTUA'TION, *Fluctuatio*, from *fluctuare*, *fluctuatum*, (*fluctus*, 'a wave,') 'to undulate.' *Undula'tio*, (F.) *Ondulation*. The undulation of a fluid collected in any natural or artificial cavity, which is felt by pressure or by percussion, properly practised—*periph'erie fluctuation*. In ascites, the fluctuation is felt by one of the hands being applied to one side of the abdomen whilst the other side is struck with the other hand. In abscesses, fluctuation is perceived by pressing on the tumour, with one or two fingers alternately, on opposite points.

FLUCTUATIO, **PERIPHERIC**, see *Fluctuation*—*f. Rhonchal*, see *Rhonchal*—*f. by Succussion*, see *Succussion*.

FLUELLEN, *Antirrhinum elatine*.

FLUELLIN, Veronica.
FLUEURS, Meuses—*f. Blanches*, Leucorrhœa.

FLUID, *Fluidus*, from *fluere*, 'to flow.' The human body is chiefly composed of fluids. If one, weighing 120 pounds, be thoroughly dried in an oven, the remains will be found not to weigh more than 12 or 13 pounds; so that the proportion of fluids to liquids in the body is about 9 or 10 to 1.

TABLE OF FLUIDS OF THE HUMAN BODY.

1. *Blood.*
2. *Lymph.*
3. *Exhaled or Perspiratory.* { Transpiration of the mucous, serous, and synovial membranes; of the areolar membrane; of the adipous cells; of the medullary membrane; of the interior of the thyroid gland; of the thymus; suprarenal capsules; eye; ear; vertebral canal, &c.
4. *Follicular.* { Sebaceous humour of the skin; cerumen; gum of the eye; mucus of the mucous glands and follicles; that of the tonsils, of the glands of the cardia, the environs of the anus, the prostate, &c.
5. *Glandular.* { Tears; saliva; pancreatic fluid; bile; cutaneous transpiration; urine; fluid of the glands of Cowper; sperm; milk; of the testes and mammae of the new-born child.

FLUID, BURNETT'S DISINFECTING, see Burnett—*f. Cephalo-rachidian*, Cephalo-spinal fluid—*f. Cephalo-spinal*, Cephalo-spinal fluid—*f. Cerebro-spinal*, Cephalo-spinal fluid—*f. Deodorizing*, Ellerman's, see Ellerman—*f. Ledoyen's* Disinfecting, see Ledoyen—*f. of Scarpa*, *Vitrine auditive*—*f. Subarachnoidæan*, Cephalo-spinal fluid.

FLUIDE SÉMINAL, Sperm.

FLUIDUM NERVEUM, Nervous fluid—*f. Cerebro-spinale*, Cephalo-spinal fluid.

FLUKE, *Distoma hepaticum*—*f. Liver*, *Distoma hepaticum*.

FLUMEN DYSENTERICUM, Dysentery.

FLUMMERY, (Scotch) *Sovens*. A preparation of oatmeal, which forms a light article of food during convalescence. It may be made as follows:—Take of oatmeal or groats, a quart. Rub with two quarts of hot water, and let the mixture stand until it becomes sour; then add another quart of hot water, and strain through a hair sieve. Let it stand till a white sediment is deposited; decant, and wash the sediment with cold water. Boil this with fresh water till it forms a mucilage, stirring the whole time.

FLUOR, Flux—*f. Albus Intestinorum*, *Coeliac flux*—*f. Albus malignus*, *Gonorrhœa impura*—*f. Muliebris*, *Leucorrhœa*—*f. Muliebris non Gallicus*, *Leucorrhœa*—*f. Sanguinis pulmonum*, *Hæmoptysis*—*f. Sanguinis vesicæ*, *Cystorrhœa*.

FLUSH, *Flushing*, (F.) *Rougeurs*, *Bouffé de Chaleur*; from (G.) *Fliesen*, 'to flow.' The temporary redness and heat produced by accumulation of blood in the capillaries of the face; as the sudden 'flush' or 'blush of emotion;' the 'flush of hectic.'

FLUX, *Fluxus*, *Profluviu*, *Fluor*, from *fluere*, 'to flow.' A discharge. *Rhysis*. In nosology, it comprises a series of affections the principal symptom of which is the discharge of a fluid. Generally it is employed for dysentery.

FLUX, BILIOUS, *Fluxus bilio'sus*. A discharge

of bile, either by vomiting or by stool, or by both, as in cholera.

FLUX, BLOODY, Dysentery—*f. de Bouche*, *Salivation*—*f. Bronchique*, *Bronchorrhœa*—*f. Dysenterique*, Dysentery—*f. Hemorrhoidal*, see *Hæmorrhœis*—*f. Hépatique*, *Hepatirrhœa*—*f. Menstrual*, *Menstris*—*f. Muqueux*, *Catarrh*—*f. Muqueux de l'estomac*, *Gastrorrhœa*—*f. Muqueux de la vessie*, *Cystirrhœa*—*f. Root*, *Asclepias tuberosa*—*f. Salivaire*, *Salivation*—*f. de Sang*, *Hæmorrhagia*, *Dysentery*—*f. Sebaceous*, *Stearrhœa*—*f. de Sperme*, *Spermatorrhœa*—*f. de Sucre*, *Ephidrosis*—*f. Weed*, *Sysimbrium sophia*—*f. d'Urine*, *Diabetes*—*f. de Ventre*, *Diarrhœa*.

FLUXIO, Fluxion—*f. Alba*, *Leucorrhœa*—*f. Arthritica*, *Gout*—*f. Vulvæ*, *Leucorrhœa*.

FLUXION, *Flux'io*, *Aflux'us*. A flow of blood or other humour towards any organ with greater force than natural. A determination. Thus we say, in those disposed to apoplexy, there is a *fluxion* or *determination* of blood to the head.

FLUXION CATARRHALE, *Catarrh*—*f. sur les Dents*, *Odontalgia*.

FLUXION DE POITRINE, (F.) By this name, the French often understand *acute pulmonary catarrh*, or *pleurisy*, but most commonly *peripneumony*.

FLUXUS, Discharge—*f. Alvinus*, *Diarrhœa*—*f. Capillorum*, *Alopecia*—*f. Chylosus*, *Coeliac flux*—*f. Coeliacus*, *Coeliac flux*—*f. Coeliacus per Renes*, *Chyluria*—*f. Cruentus cum tenesmo*, *Dysentery*—*f. Dysentericus*, *Dysentery*—*f. Hepaticus*, *Hepatirrhœa*, see *Hepateros*—*f. Lientericus*, *Lientery*—*f. Lunaris*, *Menstris*—*f. Matricis*, *Leucorrhœa*—*f. Menstrualis*, *Menstruation*—*f. Menstruus*, *Menstris*—*f. Muliebris*, *Leucorrhœa*—*f. Salivæ*, *Salivation*—*f. Sebaceous*, *Stearrhœa*—*f. Splenicus*, *Malena*—*f. Venereus*, *Gonorrhœa impura*—*f. Ventriculi*, *Gastrorrhœa*.

FLY-GOLDING, *Coccinella septem punctata*.

FLY-POLSON, *Amianthum muscetoxicum*.

FLYTRAP, *Apocynum androsaemifolium*, *Saracenia*.

FOALFOOT, *Tussilago*.

FOC'ILE. The name was formerly given to the bones of the leg, as well as to those of the forearm.

FOCILE MAJUS, *Tibia*—*f. Inferius seu majus*, *Ulna*—*f. Minus*, *Fibula*—*f. Minus seu superius*, *Radius*.

FOCUS APOPLECTICUS, *Apoplectic cell*.

FŒCUNDATIO, *Fecundation*.

FŒCUNDITAS, *Fecundity*.

FŒCUNDUS, *Fecund*.

FŒDI COLORES, *Chlorosis*.

FŒMEN, *Perinæum*.

FŒMINA, *Female*, *Woman*.

FŒMINESCENTIA, *Feminescence*.

FŒNICULUM, *Anethum*—*f. Aquaticum*, *Phellandrium aquaticum*—*f. Erraticum*, *Peucedanum silaus*—*f. Marinum*, *Crithum maritimum*—*f. Officinale*, *Anethum*—*f. Porcinum*, *Peucedanum*—*f. Vulgare*, *Anethum*.

FŒNUGREEK, *Trigonella fœnum*.

FŒNUM CAMELORUM, *Juncus odoratus*—*f. Græcum*, *Trigonella fœnum*.

FŒTAB'ULUM, An encysted abscess.—*Marcus Aurelius Severinus*.

FŒTAL, *Fætalis*. Relating to the foetus. A name given to the parts connected with the foetus. Thus we say—the *fœtal surface* of the placenta, in contradistinction to the *uterine* or *maternal surface*.

FŒTAL CIRCULATION differs from that of the adult in several respects. Commencing with the placenta, where it probably undergoes some change analogous to what occurs in the lungs in extra-

uterine existence, the blood proceeds by the umbilical vein as far as the liver, where a part of it is poured into the vena porta; the other proceeds into the vena cava inferior; the latter, having received the suprahepatic veins, pours its blood into the right auricle. From the right auricle, a part of the blood is sent into the right ventricle; the rest passes directly through into the left auricle, by the foramen ovale. When the right ventricle contracts, the blood is sent into the pulmonary artery; but as the function of respiration is not going on, no more blood passes to the lungs than is necessary for their nutrition; the remainder goes directly through the ductus arteriosus into the aorta. The blood, received by the left auricle from the lungs, as well as that which passed through the foramen ovale, is transmitted into the left ventricle; by the contraction of which it is sent into the aorta, and by means of the umbilical arteries, which arise from the hypogastric, it is returned to the placenta.

FETAL HEAD. The diameters of this at the full period are as follows:—1. The *Biparietal* or *transverse*, extending from one parietal protuberance to the other, and measuring $3\frac{1}{2}$ inches. 2. The *Temporal*, from one temple to another, 3 inches. 3. The *Occipito-mental*, from the occiput to the chin; the greatest of all, 5 inches. 4. The *Occipito-frontal* or *antero-posterior*, $4\frac{1}{2}$ or $4\frac{1}{2}$ inches. 5. The *Cervico-bregmatic*, from the nape of the neck to the centre of the anterior fontanelle. 6. The *Fronto-mental*, from the forehead to the chin, about $3\frac{1}{2}$ inches. 7. The *Trachelo-bregmatic*, from the front of the neck to the anterior fontanelle, $3\frac{1}{2}$ inches. 8. The *Vertical diameter*, from the vertex to the base of the cranium.

FETAL NUTRITION, Cyotrophy.

FETATION, Pregnancy.

FETICIDE, Feticid'ium, from *fœtus*, and *cædere*, 'to kill.' **Aborticid'ium.** Criminal abortion.

FETIDUS, Fetid.

FETOR, Dysodia.

FETOR ALARUM. The disagreeable rank odour exhaled from the axillæ.

FETOR ORIS, Breath, offensive.

FETUS, Fetus, Cyc'na, Onus ventris, Sar'cina. 'The young of any creature.' The unborn child. (F.) *Fœtus, Foiz.* By *κυννα, Cyc'na, Hippocrates* meant the fecundated, but still imperfect, germ. It corresponded with the term *embryo*, as now used; whilst *εμβρυον*, 'embryo,' signified the fetus at a more advanced stage of the utero-gestation. The majority of anatomists apply to the germ the name *embryo*, which it retains until the third month of gestation, and with some until the period of quickening; whilst *fœtus* is applied to it in its latter stages. The terms are, however, often used indiscriminately. When the ovule has been fecundated in the ovarium, it proceeds slowly towards, and enters the uterus, with which it becomes ultimately connected by means of the placenta. When first seen, the fetus has the form of a gelatinous flake, which some have compared to an ant, a grain of barley, a worm curved upon itself, &c. The fetal increment is very rapid in the first, third, fourth, and sixth months of its formation, and at the end of nine months it has attained its full dimensions—*Enfant à terme*. Generally, there is but one fetus in utero; sometimes, there are two; rarely three. The fetus presents considerable difference in its shape, weight, length, situation in the womb, proportion of its various parts to each other, arrangement and texture of its organs, state of its functions at different periods of gestation, &c. All these differences are important in an obstetrical and medico-legal point of view. The following table exhibits the length and weight of the fetus at different periods of gestation, on the authority of different observers. Their discordance is striking. It is proper to remark, that the Paris pound—*Poid de Marc*—of 16 ounces, contains 9216 Paris grains, whilst the avoirdupois contains, only 8532.5 Paris grains, and that the Paris inch is 1.065977 English inch.

	BECK.	MAYGRIER.	GRANVILLE.	BECK.	MAYGRIER.	GRANVILLE.
	Length.			Weight.		
At 30 days,	3 to 5 lines	10 to 12 lines			9 or 10 grains	
2 months,	2 inches	4 inches	1 inch	2 ounces	5 drachms	20 grains
3 "	$3\frac{1}{2}$ "	6 "	3 inches	2 to 3 "	$2\frac{1}{2}$ ounces	$1\frac{1}{2}$ ounces
4 "	5 to 6 "	8 "		4 to 5 "	7 or 8 "	
5 "	7 to 9 "	10 "		9 to 10 "	16 "	
6 "	9 to 12 "	12 "	9 inches	1 to 2 lbs.	2 pounds	1 pound
7 "	12 to 14 "	14 "	12 "	2 to 3 "	3 "	2 to 4 "
8 "	16 "	16 "	17 "	3 to 4 "	4 "	4 to 5 "

FÆTUS, Pregnant—f. in *Fœta*, *Cryptodidymus*.

FÆTUS SEPTIMES'TRIS. A seven months' fetus.

FÆTUS ZEPHYRIUS, Mole.

FOIE, Liver—f. d'*Antimoine*, *Oxidum stibii sulphuretum*—f. de *Soufre*, *Potassæ sulphuretum*.

FOIN DE CHAMEAU, Juncus odoratus.

FOIROLE, Mercurialis.

FOLIA APALACHINES, see *Ilex Paraguensis*—f. *Peraguæ*, *Ilex Paraguensis*.

FOLIACEUM ORNAMENTUM, see *Tuba Fallopiana*.

FOIE, Insanity—f. *Alcoolique*, *Delirium tremens*—f. des *ivrognes*, *Delirium tremens*.

FOLLETTE, Atriplex hortensis, Influenza.

FOLLICLE, Foll'culus, Cyatis, diminutive of *Follis*, a bag. See *Crypta*.

FOLLICLES, CILIARY, Meibomius, glands of—f. of *De Graaf*, *Folliculi Graafiani*—f. *Hair*, see

Hair—f. of *Lieberkühn*, see *Intestine*—f. *Nail*, see *Nail*—f. *Palpebral*, *Meibomius, glands of*—f. *Solitary*, *Brunner's glands*—f. *Synovial*, *Bursæ mucosæ*.

FOLLICULAR, Foll'culus, Foll'culose, Folliculo'sus, (F.) Foll'culeux, from 'follicle.' Relating or appertaining to a follicle,—as 'follicular inflammation,' inflammation affecting crypts or follicles, *Follicul'itis*.

FOLLICULAR ELEVATIONS, Tubercles, pearly.

FOLLICULE, Crypta—f. *Ciliaire*, *Meibomius, gland of*—f. *Ceruminæuse*, *Ceruminous glands*—f. *Dentaire*, *Dental follicle*—f. *Palpebral*, *Meibomius, glands of*.

FOLLICULES DE GRAAF, Folliculi Graafiani.

FOLLICULI GRAAFIANI, F. Ova'rii, Ova seu *O'vula Graafiana, Follicles or Vesicles of De Graaf, Graafian or Ova'rian vesicles, (F.) Fol-*

Vesicles ou Vésicules de Graaf. Small spherical vesicles in the stroma of the ovary, which have at least two coats: the outer termed *ovicapsule* and *tunic* of the *ovisac*, *tu'nica fibro'sa* and *theca folliculi*; the latter, *ovisac* and *membrana propria*; *vésicule ovulifère* of M. Pouchet. See *Membrana granulosa*. They exist in the fetus. The ovum—*ovule* of some—is contained in, and formed by, them.

Kölliker terms the follicles themselves, *O'visac*, *O'visac'ci*. The interior of the follicles contains a clear, light yellowish fluid—*liquor folliculi*—of the density of the serum of the blood, in which are, almost always, isolated granules, nuclei and cells, which appear to be detached portions of the *membrana granulosa*.

FOLLICULI OVARI, Folliculi Graafiani—f. Rottundi et Oblongi cervicis uteri, Nabothi glandulæ—f. Sanguinis, Globules of the blood—f. Sebacei, Sebaceous glands—f. Sebacei coronæ glandis, see Sebaceous glands.

FOLLICULITIS, Follicular inflammation.

FOLLICULOSE GLAND, Crypta.

FOLLICULUS, Vulva.

FOLLICULUS AERIS, *Air-chamber*. A space at the larger end of the bird's egg, formed by a separation of the two layers of the shell membrane, which is inservient to the respiration of the young being.

FOLLICULUS DENTIS, Dental Follicle—f. Fellis, Gall-bladder—f. Genitalis, Scrotum—f. Mucosus, see Crypta—f. Pili, see Hair.

FOMENT, *Fove're*, (F.) *Éluyer*, *Bassiner*. To apply a fomentation to a part.

FOMENTATION, *Fomentatio*, *Fotus*, *Py'ria*, *Thermas'ma*, *Chlidas'ma*, *Æone'sis*, *Perfu'sio*, *Asper'sio*, *Fomen'tum*, (quasi *fovementum*), from *fovere*, 'to bathe.' (F.) *Hydrolé*, *Hydrolitif*. A sort of partial bathing, by the application of cloths which have been previously dipped in hot water, or in some medicated decoction. They act, chiefly, by virtue of their warmth and moisture, except in the case of narcotic fomentations, where some additional effect is obtained.

A dry fomentation, *Fomen'tum siccum*, *Lee'tulus medicatus*, is a warm, dry application to a part;—as a hot brick, wrapped in flannel;—a bag, half filled with chamomile flowers made hot, &c.

FOMENTATION HERBS, *Herba pro fotu*. The herbs, ordinarily sold under this title by the English apothecary, are—*southernwood*, *tops of sea wormwood*, *chamomile flowers*, each two parts; *bay leaves*, one part. Zijss of these to Ovj of water.

FOMENTUM, Fomentation—f. Siccum, see Fomentation.

FOMES MORBI, *F. mali*. The *focus* or seat of any disease. (F.) *Foyer*.

FOMES VENTRICULI, Hypochondriasis, Spleen.

FOMITES, plural of *fomes*, *fomitis*, 'fuel, any thing which retains heat.' *Enaus'ma*, *Zop'yon*. A term applied to substances which are supposed to retain contagious effluvia; as woollen goods, feathers, cotton, &c.

FONCTION, Function.

FUNCTIONNEL, Functional.

FOND, Fundus.

FONDAMENTAL, Fundamental.

FONDANT, Solvent.

FONDEMENT, Anus.

FONDLING, Idiot.

FONGIFORME, Fungoid.

FONGOÏDE, Fungoid.

FONGOSITÉ, Fungosity.

FONGUEUX, Fungous.

FONGUS, Fungus—f. *Médullaire*, Encephaloid.

FONS, Fontanella—f. Lacrymarum, see Canthus—f. Medicatus, Water, mineral—f. Pulsans, Fontanella—f. Pulsatilis, Fontanella—f. Salutaris, Water, mineral—f. Soterius, Water, mineral—f. Vitalis, Centrum vitale.

FONSANGE, MINERAL WATERS OF. Fonsange is situated near Nismes in France. The water is sulphuretted.

FONTAINE DE LA TÊTE, see Fontanella.

FONTA'NA, CANAL OF. A canal of a triangular shape, at the inner side of the ciliary circle; partly formed by the groove at the inner edges of the cornea and sclerotic.

FONTANEL'LA. A *fontanel*:—diminutive of *fons*, *fontis*, 'a fountain.' (F.) *Fontanelle*. *Fons*, *F. pulsat'ilis* seu *pulsans*, *Vertex pallipitans*, *Bregma*, *Fontic'ulus*, *Lacu'na*, *Mould*. The opening of the head. A name, given to a space occupied by a cartilaginous membrane, in the fetus and new-born child, and situate at the union of the angles of the bones of the cranium. There are six fontanels. 1. The great or *incip'ital* or *anterior*, *fontic'ulus quadrangula'ris*, *Fontaine de la Tête*, situate at the junction of the sagittal and coronal sutures. 2. The small or *bregmat'ic* or *posterior*, situate at the part where the posterior and superior angles of the parietal bones unite with the upper part of the os occipitis. 3. The two *sphenoidal*, in the temporal fossæ; and, 4. The two *mastoïd*, or of *Casseri*, *Fonticuli mastoïdes* seu *Casse'rii*, at the union of the parietal, occipital, and temporal bones.

FORTE, Colligation.

FONTES, see Fons—f. *Medicati Plumbarii*, *Plombières*, mineral waters of—f. *Sulphurei calidi*, Waters, mineral, sulphureous.

FONTICULE À POIS, see Fonticulus.

FONTICULUS, *Fontanel'la*, *Exuto'rium*, *Exuto'ry*, *Issue*, (F.) *Fonticle*, *Cautére*, *Exutoire*. Diminutive of *fons*, *fontis*, 'a fountain.' A small ulcer produced by art, either by the aid of caustics or of cutting instruments; the discharge from which is kept up with a view to fulfil certain therapeutical indications. The *Pea issue*, (F.) *Fonticule à pois*, is kept up by means of a pea placed in it. This *pea*, (F.) *Pois à cautère*, is sometimes formed of wax; at others, the young, blasted fruit of the orange is employed. The common dried garden pea answers the purpose. The seton is also an issue.

FONTICULUS CASSERII, see Fontanella—f. Mastoïdeus, see Fontanella—f. Quadrangularis, see Fontanella.

FOOD, Aliment—f. Farinaceous, see Farinaeous—f. of the Gods, *Asnæctida*.

FOOD, PRINCE OF WALES'S. A farinaceous preparation, which is used in the same cases as arrow-root, is said to consist entirely of potato-flour.

FOOL'S STONES, MALE, *Orchis mascula*.

FOOT, Pes—f. Flat, see Kylosis—f. Griffo'n's, Gryphus pes.

FOOTLING CASE. A presentation of the foot or feet in parturition.

FORA'MEN, *Trema*, from *foro*, 'I pierce.' *Aulos*, (F.) *Trou*. Any cavity, pierced through and through. Also, the orifice of a canal.

FORAMEN ALVEOLARE ANTERIUS, see Palatine canals—f. Alveolare posterius, see Palatine canals—f. *Amplum pelvis*, *Obturatorium foramen*—f. *Anonymum Ferreini*, see *Aquæductus Fallopii*—f. *Aquæductus Fallopii*, *F. stylomastoïdeum*—f. *Auditorium externum*, see *Auditory canal*, external—f. *Auditorium internum*, see *Auditory canal*, internal—f. of Bichat, see Canal, arachnoid—f. of Botal, see Botal foramen—f. *Cæcum ossis maxillaris superioris*, see Palatine canals.

FORAMEN CÆCUM OF THE MEDULLA OBLON-

GATA or of Vico. d'AZyr. (Fr.) *Trou borgne*. A tolerably deep fossa at the point where the median furrow at the anterior surface of the medulla oblongata meets the pons.

FORAMEN CÆCUM OF THE TONGUE, see Cæcum foramen, and Tongue—f. Carotid, see Carotica foramina.

FORAMEN CENTRALE ET LIMBUS LUTÆUS RETINÆ. The central foramen and yellow spot of the retina; discovered by Sömmerring; *Macula lutea, flava seu aurea*, (F.) *Tache jaune*. It is situate about two lines to the outside of the optic nerve, and in the direction of the axis of the eye.

FORAMEN COMMUNE ANTERIUS, Vulva—f. Commune posterius, see Anus—f. Condyloid, see Condyloid—f. Conjugationis, see Conjugation—f. Ethmoideum, Orbital foramen, internal—f. Incisivum, see Palatine canal—f. Infraorbitarium, see Sub-orbital canal—f. Infrapubianum, Obturator foramen—f. Jugulare, Lacerum posterius foramen—f. Lacerum in basi cranii, Lacerum posterius foramen—f. Lacerum inferius, Sphenomaxillary fissure—f. Lacerum superius, Sphenoidal fissure—f. Magnum, see Occipital bone—f. Mastoid, see Mastoid foramen—f. Mental, see Mental foramen.

FORAMEN OF MONRO, *Foramen Monroianum*. An opening behind the anterior pillar of the fornx, somewhat above the anterior commissure, by which the third ventricle communicates with the lateral ventricle; so called after the second Monro.

FORAMEN OF MORGAGNI, see Cæcum foramen and Tongue—f. Obturatorium, Obturator foramen—f. Oculi, Pupil—f. Œsophageum (of the diaphragm), see Œsophagus, and Œsophageal aperture of the diaphragm—f. Oodes, Ovale foramen—f. Opticum, Optic foramen—f. Opticum Chorioideæ, see Optic nerves—f. Orbitarium internum, Orbital foramen, internal—f. Orbitarium superius, Orbital foramen, superior, F. supra-orbitarium—f. Ovale, Botal foramen, Fenestra ovalis, Obturatorium foramen, Ovale foramen—f. Ovale, patency of the, see Cyanopathy—f. Palatinum anterius, see Palatine canals—f. Palatinum posterius, see Palatine canals—f. Palato-maxillare, see Palatine canals—f. Rotundum, Fenestra rotunda—f. Spheno-spinosum, Spinal foramen—f. Spinale, Spinal foramen—f. Spinosum, Spinal foramen—f. Tarini, see Aqueductus Fallopii—f. Thyroideum, Obturatorium foramen—f. of Winslow, Hiatus of Winslow.

FORAMINA CRIBROSA, see Cribra.

FORAMINA THEBÆSII. Openings resembling vascular orifices, found below the orifice of the vena cava superior in the right auricle, which are supposed to be the openings of veins.

FORATIO, Trepanning.

FORCE, from *fortis*, 'strong,' *Vis*, *Potentia*, *Energia*, *Dynamis*, *Cratos*. Any power which produces an action. These powers which are inherent in organization are called *vital forces*. We say, also, *organic force*, and *muscular force*, to designate that of the organs in general, or of the muscles in particular. To the latter the word *Dynamis*, *dynamis*, corresponds; and the absence of this force is termed *adynamia*. The *vital forces* have to be carefully studied by the pathologist. The doctrine of diseases is greatly dependent on their augmentation or diminution; freedom or oppression, &c.

FORCE OF ASSIMILATION, Plastic force—f. Catalytic, see Catalysis—f. Cell, see Cell force—f. of Formation, Plastic force—f. Germ, Plastic force—f. *Médicatrice*, *Vis medicatrix*—f. Metabolic, see Metabolic force—f. of Nutrition, Plastic force—f. Osmotic, see Osmotic force—f. of Vegetation, Plastic force—f. Vital, *Vis vitalis*—f. *Vitale*, *Vis vitalis*.

FORCE-REAL, MINERAL WATERS OF.

The name of a mountain, situate four leagues from Perpignan in France. The water is chalybeate.

FORCEPS, quasi, *ferriiceps*, from *ferrum*, 'iron,' and *capio*, 'I take.' *Pincers*, *Lobis*, *Volsell'er*. An instrument, for removing bodies which it would be inconvenient or impracticable to seize with the fingers. (F.) *Pince*, *Pincette*. There are various kinds of forceps, 1. The ordinary kind, contained in every dressing-case, for removing lint, &c. from wounds or ulcers. (F.) *Pince à pansement ou à anaux*. 2. *Dissecting or Ligature or Arterial Forceps*, (F.) *Pince à dissection*, *P. à ligature*, to lay hold of delicate parts. 3. *Polypus Forceps*, *Tooth Forceps*, *Forceps of Muscæ* for laying hold of the tonsils or other parts to be removed. 4. *The Bullet Forceps*, *Strombul'cus*, (F.) *Tireballe*. 5. The *Lithotomy Forceps*, *Lithot'abon*, *Lithago'gon*, *Tenac'ula*, *Volsell'a*, which resembles the *Craniotomy Forceps*, (F.) *Tenettes*. 6. The *Cataract Forceps*, (F.) *Pince à Cataracte*, resembling the dissecting forceps, but much finer. 7. The *Pince à guîne*, the branches of which can be brought together, and retained by a movable sheath.

Forceps is also an instrument used by obstetrical practitioners to embrace the head, and bring it through the pelvis. It consists of two branches, blades or levers; one of which, in the case of the *short forceps*, is passed over the ear of the child, and the other opposite the former, so that the blades may lock. When the head is securely included between the blades, the operation of extraction can be commenced. See Parturition.

FORCEPS, see *Corpus callosum*—f. Anterior, see *Corpus callosum*—f. Arterial, see *Forceps*—f. Bullet, see *Forceps*—f. Craniotomy, see *Forceps*—f. Deceptoris, see *Forceps*—f. Lithotomy, see *Forceps*—f. of Muscæ, see *Forceps*—f. Polypus, see *Forceps*—f. Tooth, see *Forceps*.

FORD'S BALSAM OF HOREHOUND, see Balsam of horehound.

FOREARM, *Antibrah'chium*, *Pygmæ*, *Pars inferior bra'chii*, *Cu'bitus*, (F.) *Avant-bras*. The part of the upper extremity, comprised between the arm and the hand. It is composed of two bones—*radius* and *ulna*—and 20 muscles.

FOREFRONT, Front.

FOREHEAD, Front.

FORE-MILK, Colostrum.

FOREIGN BODY, Extraneous body.

FORENSIC ANATOMY, see *Anatomy*—f. Medicine, Medicine, legal.

FORES, Genital organs.

FORESKIN, Prepuce.

FORETOP, Front.

FORFEX, *Ciseau*, *Scissors*—f. *Dentaria*, *Dentagra*.

FORFICULA AURICULARIA, *Earwig*. An insect of the Order Orthoptera, which occasionally enters the meatus auditorius externus, and excites intense pain. It may be destroyed by tobacco-smoke, or by oil poured into the meatus.

FORGES, MINERAL WATERS OF. Forges is situate four leagues from Gournay, in the department of Seine Inférieure, France. There are three springs, which are acidulous chalybeates. These are called *Royal*, *Reinette*, and *Cardinal*, in honor of Louis XIII., Queen Anne of Austria, and Cardinal Richelieu, who used them.

FORMATIVE, Plastic.

FORMICA, *Myrmex*. The ant or pismire. (F.) *Fourmi*. It contains an acid juice and gross oil, which were formerly extolled as aphrodisiacs. The chrysalides of the animal are said to be diuretic and carminative; and have been used in dropsy. 2. Also the name of a black wart, *verruca formica'ria*, with a broad base and cleft surface; so called because the pain attending it

resembles the biting of an ant, *μυρμικία*, *myrmecia*. — Forestus. 3. A varicose tumour on the anus and glans penis. 4. Miliary herpes.

FORMICA AMBULATORIA, *Herpes circinatus* — f. *Corrosiva*, *Herpes exedens*.

FORMICANT, *Formicæans*, *Myrmecizon*, from *formica*, 'an ant' (F.) *Fourmillant*. An epithet given to the pulse, *Pulsus formicæans*, when extremely small, scarcely perceptible, unequal, and communicating a sensation like that of the motion of an ant through a thin texture.

FORMICATION, *Formicatio*, *Myrmeciasimus*, *Myrmeciasis*, *Myrmecismus*, *Myrmecosis*, *Stupor formicæans*. Same etymon. (F.) *Fourmillément*. A pain, compared with that which would be caused by a number of ants creeping on a part.

FORMIX, *Herpes esthiomenus*.

FORMULA, Prescription — f. *Extemporanea*, see *Magistral* — f. *Magistralis*, see *Magistral* — f. *Medica*, Prescription — f. *Officinalis*, see *Magistral*.

FORMULARY, *Formulaarium*, *Narthecia*, *Narthecium*, *Narther*. From *formularius*, (diminutive of *forma*, 'form'). One skilled in preparing writs or forms of process. — Quintilian. A collection of medical formulæ or receipts.

FORMULE, Prescription.

FORMYL, PERCHLORIDE OF. Chloroform.

FORMYLE, IODURE DE, Iodoform.

FORNIX, 'an arch or vault.' F. *cerëbri* seu *trilatëris* seu *tricuspidatilis*, *Pæalis*, *Pæalidium*, *Can'eca*, *Testat'lo cerëbri*, *Arcus medullaris*, (F.) *Trigone cerëbral* — (Ch.) *Voûte à trois piliers*, *Triangle Médullaire*. A medullary body in the brain, below the corpus callosum and above the middle ventricle, on the median line. This body, which is curved upon itself, terminates anteriorly by a prolongation, which constitutes its *anterior pillar*, *Crus* seu *Colum'na forni'cia*, (F.) *Pilier antérieur*, and posteriorly by two similar prolongations, called *posterior pillars* or *crura*. See *Achicolum*, and *Vault*.

FORNIX CEREBRI, Fornix.

FORNPECKLES, *Ephelides*.

FORPEX, *Scissors*.

FORRET, *Front*.

FORSTERANIA DIFFORM'IS, *Echites difform'is* seu *puberula*; indigenous; Ord. *Apocynaceæ*; is said to be used as a wash with milk to remove freckles. The juice destroys warts.

FORTIFIANT, *Corroborant*, *Tonic*.

FORTRAITURE, *Hysteralgia*.

FORTYKNOT, *Achyranthes repens*.

FOSSA, *Fo'vea*, from *fodio*, *fossam*, 'I dig.' *Scamma*. A cavity of greater or less depth, the entrance to which is always larger than the base. The fossæ of bones have been called *simple*, when they belong to one bone only, as the *parietal fossæ*; and *compound*, (F.) *Fossæ composæ*, when several concur in their formation, as the *orbital fossæ*, *temporal fossæ*, &c.

FOSSA, AMYG'DALOID, *Amygdaloid excavatio*. The space between the anterior and posterior pillars of the fauces, which is occupied by the tonsils.

FOSSA AMYN'TÆ. A kind of bandage, used in fractures of the nose; so called, by Galen, from Amyntas of Rhodes, its inventor. It consisted of a long band, applied round the head, the turns of which crossed at the root of the nose.

FOSSA CANINA, *Canine fossa*.

FOSSA CEREBEL'LI, (F.) *Fossæ cerëbelleuse*. The inferior occipital fossa, which lodges the corresponding portion of the cerebellum.

FOSSA CORONA'NIS, *Coronal* or *frontal fossa*.

A depression on the orbital plate of the frontal

or coronal bone, which supports the anterior lobe of the brain.

FOSSA CORONOID'EA, *Coronoid fossa*. A cavity before the inferior extremity of the humerus, in which the coronoid process of the ulna is engaged during the flexion of the forearm.

FOSSA COTYLOIDEA, see *Cotylid*.

FOSSA DIGAS'TRIC, *Fossa digastrica*. A deep groove on the mastoid portion of the temporal bone, which gives origin to the digastric muscle.

FOSSA ETIMOID'ALIS, *Ethmoid fossa*. A shallow gutter on the upper surface of the cribriform plate of the ethmoid bone, in which is lodged the expanded portion of the olfactory nerves.

FOSSA GENU, *Popies*.

FOSSA GUTTURA'NIS, *Guttural fossa*. The depression which forms the guttural region of the base of the cranium, between the foramen magnum and posterior nares.

FOSSA HYALOIDEA, see *Hyaloid* (Fossa) — f. *Iliac*, see *Iliac fossæ* — f. *Infra-orbital*, *Canine fossa* — f. *Infra-spinous*, see *Infra-spinata fossa*.

FOSSA INNOMINA'TA. The space between the helix and antihelix of the ear.

FOSSA, ISCHIO-RECTAL, *Perineal fossa* — f. *Jugularis*, *Jugular fossa* — f. *Lacrymalis*, *Lacrymal fossa* — f. *Lenticularis*, see *Hyaloid membrane* — f. *Magna Muliebris*, *Vulva* — f. *Magna Sylvi*, *Fissura Sylvi*.

FOSSA, MENTAL, *Fossa mentalis*. A small depression on each side of the symphysis on the anterior surface of the body of the maxilla inferior, for the attachment of muscles.

FOSSA NAVICULARIS, *Navicular fossa*.

FOSSA OCCIPITA'NIS, *Occipital fossa*. The occipital fossæ are four in number: the *superior* or *cerebral*, and the *inferior* or *cerebellous*. They are separated by a groove, which lodges the lateral sinus.

FOSSA OVALIS, see *Ovalis fossa* — f. *Palatina*, *Palate* — f. *Patellaris*, see *Hyaloid membrane* — f. *Perinæi*, *Perineal fossa* — f. *Pituitaria*, *Sella Turcica*.

FOSSA POPLITE'A, *Poplite'al fossa*. The hollow of the ham; — the popliteal region.

FOSSA, PORTAL, see *Liver* — f. *Scaphoides*, *Navicularis fossa*.

FOSSA SIGMOIDEA, *Sulcus sinus lateralis*. A broad deep groove on the inner surface of the mastoid portion of the temporal bone, which supports part of the lateral sinus.

FOSSA, SUBORBITAR, *Canine fossa*.

FOSSA, SUPRYPYRAMIDAL, *F. suppyramidalis*. A deep fossa under the pyramid and behind the fenestra rotunda in the middle ear, remarkable for its constancy, and pierced by several foramina at the bottom.

FOSSA SUPRA-SPHENOIDALIS, *Pituitaria fossa* — f. of *Sylvius*, *Fissure* of S. — f. *Temporalis*, *Temporal fossa* — f. *Transversa*, see *Liver* — f. *Umbilicalis*, see *Liver*.

FOSSÆ CEREBRA'LES, *Cerëbral fossæ*. Fossæ or excavations at the base of the cranium. They are nine in number: three occupy the median line, and three are placed at each side. They are distinguished into *anterior*, *middle*, and *posterior*.

FOSSÆ DIGITALES, see *Impression*.

FONNE BASILAIRE, *Basiliary fossa* — f. *Sous-épineuse*, *Infra-spinata fossa* — f. *Sus-épineuse*, *Fossa supra-spinata*.

FOSSES CONDYLOIDIENNES, *Condylodea foramina* — f. *Nasales*, *Ouvertures postérieures des Nares*, posterior.

FOSSETTE (F). Diminutive of *fossa*. *Scrobiculus*, *Both'ion*. Several depressions are so called. A *dimpled chin*, *Fossette du menton*, consists in a slight depression, which certain persons have on the chin. 2. A *dimple of the cheek*, (F.)

Fossette des joues, a depression which occurs on the cheeks of certain persons when they laugh. *Scrobiculus cordis*, *Anticardium*, *Præcordium*, (F.) *Fossette du cœur*. The depression observed on a level with the xiphoid cartilage at the anterior and inferior part of the chest. It is, also, called *pit of the stomach*, (F.) *Cœur de l'estomac*.

FOSSETTE, *Fos'sula*, *Annulus*, *Bulbium*, is also a small ulcer of the transparent cornea, the centre of which is deep.

FOSSETTE ANGULAIRE DU QUATRIÈME VENTRICULE, *Calamus scriptorius*—*f. du Cœur*, *Scrobiculus cordis*.

FOSSORIUM, Fleam.

FOSSULA, *Argema*, *Fossa*, *Fossette*, *Fovea*.

FOTHERGILL'S PILLS, see *Pilulæ aloes et colocynthis*.

FOTUS, Fomentation—*f. Communis*, *Decoctum papaveris*.

FOU, Insane.

FOUETS, *Sempervivum tectorum*.

FOUGÈRE DOUCE, *Polypodium*—*f. Femelle*, *Asplenium filix femina*, *Pteris aquilina*—*f. Grande*, *Pteris aquilina*—*f. Mâle*, *Polypodium filix mas*.

FOUL DISEASE, Syphilis.

FOULURE, Sprain.

FOUNDLING HOSPITAL, *Brephotropeum*.

FOURCHE, (F.), *fourché*, 'cleft;' from *furca*, 'a fork.' *Apostema Phalangium*. A French provincial term for small abscesses which form on the fingers and hands of working people. Also, an instrument, invented by M. J. L. Petit, for compressing the ramine artery in cases of hemorrhage from that vessel.

FOURCHETTE', diminutive of *fourche*, *Furci'la*, a little fork, *Fur'cula*. A surgical instrument used for raising and supporting the tongue, during the operation of dividing the frænum.

Fourchette, in anatomy, is, 1. The posterior commissure of the labia majora, called also, *Frænum*, *Frænulum pudendi*, *Fur'cula Labiorum*; 2. The cartilago ensiformis; so called from its being sometimes cleft like a fork. Also, the semilunar notch at the superior or clavicular extremity of the sternum.

FOURMI, Formica.

FOURMILLANT, Formicant

FOURMILLEMENT, Formication.

FOUSEL OIL, see Oil, fusel.

FOVEA,—diminutive, *Fove'ola*,—from *fodio*, 'I dig.' [?] *Bothros*. A slight depression. *Fos'cula*. The *pudendum muliebre*; see *Valva*. The *fossa navicularis*. A vapour-bath for the lower extremities.

FOVEA AXILLARIS, *Axilla*—*f. Elliptica*, see *Crista Vestibuli*—*f. Hemisphærica*, see *Crista Vestibuli*—*f. Lacrymalis*, *Lachrymal fossa*—*f. Oculi*, *Orbit*—*f. Ovalis*, *Ovalis fossa*—*f. Semielliptica*, see *Crista Vestibuli*—*f. Sulciformis*, see *Crista Vestibuli*.

FOVEOLA, *Fovea*.

POWER, Syncope.

FOX-BERRY, *Arbutus uva ursi*.

FOX-GLOVE, *Digitalis*.

FOYER, *Fomes morbi*.

FRACTIONNEMENT, Segmentation.

FRACTURA, Fracture—*f. Dentis*, *Odontoclasia*.

FRACTURE, *Fractu'ra*, from *frangere*, *fractum*, 'to break,' 'bruise.' *Catag'ma*, *C. Fractu'ra*, *Catag'is*, *Clasis*, *Olasma*, *Agma*, *Aigma*, *Diar'rhagē*. A solution of continuity in a bone, *Osteoclasia*. A simple fracture is when the bone only is divided. A compound fracture is a division of the bone with a wound of the integuments communicating with the bone,—the bone, indeed,

generally protruding. In a *com'minuted fracture*, *Alphite'don*, *αλφίτηδον*, *Cerge'don* *Catag'ma*, *καργήδον κατάρμα*, the bone is broken into several pieces; and in a *complicated fracture* there is, in addition to the injury done to the bone, a lesion of some considerable vessel, nervous trunk, &c. Fractures are also termed *transverse*, *oblique*, &c., according to their direction. The treatment of fractures consists, in general, in reducing the fragments when displaced; maintaining them when reduced; preventing the symptoms which may be likely to arise; and combating them when they occur. The reduction of fractures must be effected by extension, counter-extension, and coaptation. The parts are kept in apposition by position, rest, and an appropriate apparatus. The position must vary according to the kind of fracture. Commonly, the fractured limb is placed on a horizontal or slightly inclined plane, in a state of extension; or rather in a middle state between extension and flexion, according to the case.

FRACTURE, CAMERATED, Camerosis—*f. en Rave*, *Raphanodon*.

FRACTURE OF THE RADIUS, *BARTON'S*, *Barton's fracture*. A term applied to a fracture of the lower extremity of the radius, which commences at the articular surface, and extends upwards for an inch or more, to terminate on the dorsal aspect. Owing to the extensor muscles drawing up the separated portion of the bone, and with it the carpus, a deformity results, which has been confounded with simple dislocation. In consequence of the fracture having been well described by Dr. John Rea Barton, of Philadelphia, it is often called after him.

FRACTURES PAR RÉSONNANCE, see *Contrafractura*.

FRÆNA MORGAGNII, F. of the Valve of Bauhin.

FRÆNA OF THE VALVE OF BAUHIN, (F.) *Freins de la valvule de Bauhin*. A name given by Morgagni to the projecting lines formed by the junction of the extremities of the two lips of the ileo-cæcal valve. They are also called *Frænd* and *Retinacula Morgagnii*.

FRÆNULUM, see *Frænum*, *Bride*—*f. Clitoridis*, *Frænum clitoridis*—*f. Epiglottidis*, *Glossopiglottic ligament*—*f. Labiorum*, *Fourchette*—*f. Novum*, *Tænia semicircularis*—*f. Pudendi*, *Fourchette*.

FRÆNULUM VELI MEDULLÆ RIS ANTERIORIS A narrow slip, given off by the commissure of the encephalic testes, which strengthens the junction of the testes with the valve of Vieussens.

FRÆNUM, *Fre'num*, (F.) *Frein*, 'a bridle.' *Frænulum*, *File'tum*, *Filamentum*. 'A small bridle.' A bridle. Names given to several membranous folds, which bridle and retain certain organs.

FRÆNUM CLITORIDIS, *Frænulum Clitoridis*, (F.) *Frein du Clitoris*. A slight duplicature formed by the union of the internal portions of the upper extremity of the nymphæ.

FRÆNUM EPIGLOTTIDIS, *Glossopiglottic ligament*—*f. Glandis*, *F. Penis*.

FRÆNUM LABIORUM, (F.) *Frein des lèvres*. There are two of these; one for the upper, the other for the lower lip. They unite these parts to the maxillary bone, and are formed by the mucous membrane of the mouth. Also, the *Fourchette*.

FRÆNUM LINGUÆ, *Frænulum seu Vin'culum Lingue*, *Glossodesmus*, *File'tum*, (F.) *Filet on Frein de la langue*, is a triangular reflection, formed by the mucous membrane of the mouth, and situate between the inferior paries of that cavity and the inferior surface of the tongue.

When the frænum extends as far as the extremity of the tongue, it cramps its movements, interferes with sucking, &c. This inconvenience is remedied by carefully snipping it with a pair of scissors. The French call this *l'opération du filet*.

FRÆNUM PENIS, *F.* seu *Frænulum* seu *Vinculum Præputii*, *F. Glandis*, *Cynodesmion*, *Cynodesmus*, *Vinculum Caninum*, (*F.*) *Filet* ou *Frein de la verge*, is a membranous reflection which fixes the prepuce to the lower part of the glans. When too short, it prevents the prepuce from sliding over the glans.

FRÆNUM PRÆPUTII, *F. Penis*.

FRAGARIA, from *frago*, 'I smell sweetly.' The Strawberry, *Frageria vesca* seu *vulgaris* seu *sempervirens*, *Chamaebatos*, (*F.*) *Fraisier*. *Ord.* Rosaceæ. The fruit is agreeable and wholesome, and the roots have been used as tonic and slightly diuretic. The fruit is the *Fragrum*, *κοναρον*, of the ancients; (*F.*) *Fraise*.

FRAGARIA ANSERINA, *Potentilla anserina*—*f.* *Pentaphyllum*, *Potentilla reptans*—*f.* *Tormentilla officinalis*, *Tormentilla*.

FRAGARIA VIRGINIANA, *Wild Strawberry*. An indigenous plant, which has astringent leaves.

FRAGILE VITREUM, *Fragilitas ossium*.

FRAGILITAS, *Ruptibilitas*, from *frango*, 'I break.' Fragility, Brittleness. The state of being easily broken or torn.

FRAGILITAS OSSIIUM, *Osteopathyrosis*, *Parostia fragilis*, Brittleness of the bones, Friability of the bones, *Fragile vitreum*. Pathologists have given this name to the extreme facility with which bones break in certain diseases of the osseous texture. It is owing to a deficiency of the animal matter.

FRAGMENT, *Fragmentum*, *Fragmen*, *Ramentum*, from *frangere*, *fractum*, 'to break.' The French use this term for the two portions of a fractured bone; thus, they speak of the *superior* and the *inferior fragment*.

FRAGMENTS, *PRECIOUS*. A name formerly given, in *Pharmacy*, to the garnet, hyacinth, emerald, sapphire and topaz. The Arabs falsely attributed to them cordial and alixiterial properties.

FRAGON, *Ruscus*.

FRAGUM, see *Fragaria*.

FRAISE, see *Fragaria*.

FRAISIER, *Fragaria*.

FRAMBŒSIA, *Frambœsia*, *Lepra fungifera*, from *framboise*, (*F.*) 'A raspberry.' *Syphilis Indica*, *Anthraxia rubula*, *Thymiosis*, *Th.* seu *Lues Indica*, *Variola Amboinensis*, *Lepra fungifera*, *Scrophula Moluccana*. The *Yaws*, *Epian*, *Pian*. A disease of the Antilles and of Africa, characterized by tumours, of a contagious character, which resemble strawberries, raspberries, or champignons; ulcerate, and are accompanied by emaciation, The *Pian*, for so the Indians call it, differs somewhat in America and Africa.

Pian of America, *Frambœsia Americana*, *Anthraxia Rubula Americana*, occurs under similar circumstances with the next, and seems to be transmitted by copulation. The tumours have a similar form, and are greater in proportion to their paucity. In some cases they are mixed with ulcers.

Pian of Guinea, *Frambœsia Guineensis*, *Anthraxia Rubula Guineensis*, is common amongst the negroes, especially in childhood and youth. It begins by small spots, which appear on different parts, and especially on the organs of generation and around the anus; these spots disappear, and are transformed into an eschar, to which an excrescence succeeds, that grows slowly, and has the shape above described.

The treatment is nearly the same in the two varieties. The tumours will yield to mercurial friction, when small. When large, they must be destroyed by caustic. In both cases, mercury must be given to prevent a recurrence.

An endemic disease resembling yaws was observed in the Feejee Islands by the medical officers of the United States' Exploring Expedition. It is called by the natives *Dhoko*.

FRAMBŒSIA ILLYRICA, *Scherlievo*—*f.* *Scotica*, *Sibbens*.

FRAMBOISE, *Rubus idæus*.

FRANCOLIN, *Attagen*.

FRANGES SYNOVIALES, *Synovial glands*.

FRANGIPANE. An article of food, prepared by evaporating to dryness, in a water-bath, milk mixed with almonds and sugar.

FRANGULA ALNUS, *Rhamnus frangula*.

FRANKINCENSE, *COMMON*, see *Pinus abies*—*f.* *African Daniellia thurifera*—*f.* *Tree*, *Daniellia thurifera*—*f.* *True*, *Juniperus lycia*.

FRANZENSBAD, **FRANZBAD**, or **FRANZENBRUNN**, **MINERAL WATERS OF**. A celebrated water at Eger, in Bohemia, which contains sulphate of soda, carbonate of iron, and carbonic acid gas. The springs are called *Franzenbrunnen*.

FRAPPER, *Percuss*.

FRASERA, *Calumba*, *American*—*f.* *Carolinensis*, see *Calumba*—*f.* *Officinalis*, see *Calumba*—*f.* *Verticillata*, *Calumba American*—*f.* *Walteri*, see *Calumba*.

FRATER UTERINUS, see *Uterinus frater*.

FRATERINITAS, *Adelphixia*.

FRATRATIO, *Adelphixia*.

FRAXINELLA DICTAMNUS, *Dictamnus albus*—*f.* *White*, *Dictamnus albus*.

FRAXINUS AMERICANA, *White Ash*; and **FRAXINUS QUADRANGULATA**, *Blue Ash*, indigenous; *Ord.* *Oleaceæ*; have bitter and astringent barks, and have been used as antiperiodics.

FRAXINUS APETALA, *F. excelsior*—*f.* *Aurea*, *F. excelsior*—*f.* *Crispa*, *F. excelsior*—*f.* *Excelsa*, *F. excelsior*.

FRAXINUS EXCELSIOR. The systematic name of the *Ash tree*, *Fraxinus Ornus* seu *Fraxinus sylvestris*, *F. apetal* seu *aur* seu *crispa* seu *excelsa* seu *pendula* seu *verruco* seu *Bume* seu *Macedonica* *Fraxinus*, *Bume*, (*F.*) *Frêne*. The fresh bark has a moderately strong, bitterish taste. It has been said to possess resolvent and diuretic qualities, and has been given in intermittents, and in gout and rheumatism, and as an antidote to serpent bites. The seeds, called *Birds' tongue*, *Lingua avis*, *Ornithoglossa*, have been exhibited as diuretics, in the dose of a drachm. Its sap has been extolled against deafness.

FRAXINUS FLORIFERA, *F. ornus*—*f.* *Macedonica*, *F. excelsior*.

FRAXINUS ORNUS, *F. melia* seu *paniculata* seu *florifera*, *Ornus mannifera* seu *rotundifolia*. The systematic name of the tree whence *manna* flows. This substance is also called *Manna Calabrin*, *Ros Calabrinus*, *Aëromeli*, *Drosomeli*, *Drysomeli*, *Mel ærium*, *Succus orn* concretus. In Sicily, not only the *Fraxinus Ornus*, but also the *F. rotundifolia* and *F. excelsior* are regularly cultivated for the purpose of procuring *manna*, which is their condensed juice. In the Ph. U. S. it is assigned to *Ornus Europæa*. *Manna* is inodorous, sweetish, with a very slight degree of bitterness; in friable flakes, of a whitish or pale yellow colour; opaque, and soluble in water and alcohol. It is laxative, and is used as a purgative for children, who take it readily on account of its sweetness. More generally it is employed as an adjunct to other purgatives. Dose, ʒss to ʒij. Its immediate principle is called

Mannite or **Mannin**. This has been recommended by Magendie as a substitute for manna. Dose, $\frac{3}{4}$ ij. for children.

FRAXINUS PANICULATA, F. Ornus—f. Quadrangulata, see F. Americana—f. Rotundifolia, F. ornus.

FRAYEUR NOCTURNE, Panophobia.

FRECKENS, Ephelides.

FRECKLED, see Ephelides.

FRECKLES, Ephelides.

FREEMAN'S BATHING SPIRITS, see Limentum saponis compositum.

FREIN, Frænum—f. du Clitoris, Frænum clitoridis—f. de la Langue, Frænum linguae—f. des Lèvres, Frænum labiorum—f. de la Verge, Frænum penis.

FREINS DE LA VALVULE DE BAUHIN, Fræna of the valve of Bauhin.

FREINWALDE, MINERAL WATERS OF. These springs are in Brandenburg, twelve leagues from Berlin. They contain chloride of sodium, sulphate of magnesia, chloride of magnesium, sulphate of lime, carbonates of lime and magnesia, iron, &c., and are used in asthenic diseases.

FRÉMISSEMENT, Shuddering.

FRÉMISSEMENT CATAIRE (F.). 'Cat's purr.' Purring Tremor. Laënnec has given this name to the agitation which is sensible to the hand, when applied on the præcordial region, and which he considers a sign of ossification or other contraction of the auriculo-ventricular openings. The name was chosen by him from the analogy of the sound to the purring of a cat.

FRÉMITUS, Bruissement, Frémissement, Shuddering.

The **PECTORAL** or **VOCAL FRÉMITUS**, *Pectoral Vibration*, *Tactile vibration*, is an obscure diffused resonance of the voice, which is felt when the hand is applied to the chest.

A **FRIC'TION FRÉMITUS** is perceptible to the hands in many cases of pleurisy, when the surfaces of the pleura have become roughened from deposition; as well as in rare cases of pericarditis.

FRENA, see Alveolus.

FRENCH CRUST, Syphilis—f. Distemper, Syphilis—f. Nut, Juglans regia—f. Pox, Syphilis.

FRÈNE, Fraxinus excelsior—f. Épineux, Xanthoxylum clava Herculis—f. Sauvage, Sorbus aucuparia.

FRENGA, see Scherlievo.

FRENULUM, see Frænum.

FRENUM, Frænum.

FRET, Chafing, Herpes.

FRETILLEMENT, Fidgets.

FRETTING, Tormina.

FRICATIO, Friction.

FRICATORIUM, Liniment.

FRICE, *Fricum*, *Friconium*. A medicine which the ancients employed under the form of friction. They distinguish the *Fricæ sicca* and *F. mollæ*.—Gaubius.

FRICONIUM, Fricæ.

FRICTA, Colophonia.

FRICTIO, Friction—f. Humida, see Friction—f. Sicca, see Friction.

FRIC'TION, *Fric'tio*, *Frica'tio*, *Anat'ribæ*, *Anat'rip'sia*, *Trypsia*, *Chirap'sia*, from *fricare*, *frictum*, 'to rub.' The action of rubbing a part of the surface of the body more or less forcibly, with the hands, a brush, flannel, &c., constituting *Xerotrib'ia*, *Xerotrip'sis*, *Fric'tio sicca* or *dry friction*; or with ointments, liniments, tinctures, &c., constituting *moist friction*, *Fric'tio hu'mida*. It is a useful means for exciting the action of the skin.

FRIC'TION FRÉMITUS, see *Frémitus*—f. Sound, *Bruit de frottement*.

FRICTRIX, Tribas,

FRICTUM, Liniment.

FRICUM, Fricæ.

FRIGEFACIENTIA, Refrigerants.

FRIGID, *Frig'idus*, (F.) *Froid*; from *frigeo*, 'to be cold.' Cold. Not easily moved to sexual desire; *Imbell'is ad ven'erem*. Impotent.

FRIGIDARIUM, Bath, cold.

FRIGID'ITY, *Frigid'itas*. Same etymon. A sensation of cold. Also, impotence, and sterility. *Frigidity of the stomach* is a state of debility of that organ, imputed to excessive venery,—the *Anorexia exhaustivum* of Sauvages.

FRIGIDUS, Frigid.

FRIGORIFIC, from *frigus*, 'cold,' and *fio*, 'I become.' That which has the power of producing cold. The best **FRIGORIFIC MIXTURES** are the following. Their effects are owing to the rapid absorption of heat when solids pass into the liquid state.

FRIGORIFIC MIXTURES WITH SNOW.

Mixtures.	Therm. falls.
Snow, or pounded ice, two parts by weight;	From any temperature. to—5°
Chloride of Sodium..... 1.	
Snow or pounded ice..... 5.	
Chloride of Sodium..... 2.	to—12°
Mur. of Ammonia..... 1.	
Snow or pounded ice..... 24.	
Chloride of Sodium..... 10.	to—18°
Muriate of Ammonia..... 5.	
Nitrate of Potash..... 5.	
Snow or pounded ice..... 12.	to—25°
Chloride of Sodium..... 5.	
Nitrate of Ammonia..... 5.	
Dilut. Sulph. Acid..... 2.	from +32°
Snow..... 3.	to—23°
Concentr. Mur. Acid..... 5.	from +32°
Snow..... 8.	to—27°
Concentr. Nitrous Acid..... 4.	from +32°
Snow..... 7.	to—30°
Chloride of Calcium..... 5.	from +32°
Snow..... 4.	to—40°
Crystall. Chloride of Calcium 3.	from +32°
Snow..... 2.	to—50°
Fused Potash..... 4.	from +32°
Snow..... 3.	to—51°

Frigorific Mixtures may also be made by the rapid solution of salts, without the use of snow or ice. The salts must be finely powdered and dry.

FRIGORIFIC MIXTURES WITHOUT SNOW.

Mixtures.	Therm. falls.
Mur. of Ammonia..... 5.	from +50°
Nitrate of Potash..... 5.	
Water..... 16.	to—10°
Mur. of Ammonia..... 5.	from +50°
Nitrate of Potash..... 5.	
Sulphate of Soda..... 8.	to—10°
Water..... 16.	
Nitrate of Ammonia..... 1.	from +50°
Water..... 1.	to— 4°
Nitrate of Ammonia..... 1.	from +50°
Carbonate of Soda..... 1.	to— 7°
Water..... 1.	
Sulphate of Soda..... 3.	from +50°
Dilut. Nitrous Acid..... 2.	to— 3°
Sulphate of Soda..... 6.	
Mur. of Ammonia..... 4.	from +50°
Nitrate of Potash..... 2.	to—10°
Dilut. Nitrous Acid..... 4.	
Sulphate of Soda..... 6.	from +50°
Nitrate of Ammonia..... 5.	to—14°
Dilut. Nitrous Acid..... 4.	
Phosphate of Soda..... 9.	from +50°
Dilut. Nitrous Acid..... 4.	to—12°

<i>Mixtures.</i>		<i>Therm. falls.</i>
Phosphate of Soda.....	9.	from +50° to—21°
Nitrate of Ammonia	6.	
Dilut. Nitrous Acid.....	4.	from +50° to 0°
Sulphate of Soda.....	3.	
Muriatic Acid.....	5.	from +50° to—3°
Sulphate of Soda.....	5.	
Dilut. Sulphuric Acid.....	4.	

FRIGUS, Cold—*f. Tenue*, see Rigor.

FRINGE TREE, *Chionanthus Virginica*.

FRISSE, Rigor.

FRISSEMENT, Horripilation.

FROGLEAF, *Brasenia Hydropeltis*.

FROG TONGUE, *Ranula*.

FROGS' SPAWN, *Sperma ranarum*.

FROID, Cold, Frigid—*f. Glacial*, Ice cold.

FROISSEMENT (F.), 'Rubbing, bruising.'

FROISSEMENT PULMONAIRE, *Bruit de froissement pulmonaire*, Pulmonary crumpling sound. A name given by M. Fournet to a respiratory sound, which communicates to the ear the sensation of the rubbing (*froissement*) of a texture compressed against a hard body. It is by no means well defined.

FRÔLEMENT, (F.) 'Grazing or touching lightly.'

FRÔLEMENT PÉRICARDIQUE, *Bruit de frôlement péricardique*. Rustling noise of the pericardium. A sound resembling that produced by the crumpling of a piece of parchment or of thick silken stuff, accompanying the systole and diastole of the heart. It indicates roughness of the pericardium induced by disease.

FROMAGE, Cheese.

FROMENT, Triticum.

FRONCEMENT, Corrugation.

FRONCLE, Furunculus.

FRONDE, Funda.

FRONT, *Frons*, *Metopon*, Forehead, Fore-front, Fore-top, Brow, (Sc.) Forret. That part of the visage, which extends from one temple to the other, and is comprised in a vertical direction, between the roots of the hair and the superciliary ridges.

FRONTAL, *Frontalis*. From *frons*, *frontis*, 'the front.' Relating or belonging to the front. This name has been given to several parts. Winslow, Sömmering, and others, call the anterior part of the occipito-frontals—the *frontal muscle* or *frontalis*, *Musculo'sa Frontis Cutem movens subcutia* Par (Vesalius).

FRONTAL ARTERY, *Arteria supra-orbitalis*, is given off by the ophthalmic, which is itself a branch of the internal carotid. It makes its exit from the skull at the upper part of the base of the orbit, and ascends the forehead between the bone and the orbicularis palpebrarum; dividing into three or four branches, which are distributed to the neighbouring muscles.

FRONTAL BONE, *Os frontis* seu *coronallē* seu *invarecul'dum* seu *puppis* seu *syncephalis* seu *proræ* seu *ratio'nis*, *Metopon*. A double bone in the fetus, single in the adult, situate at the base of the cranium, and at the superior part of the face. It forms the vault of the orbit; lodges the ethmoid bone in a notch at its middle part; and is articulated, besides, with the sphenoid, parietal, and nasal bones, the ossa unguis, superior maxillary, and malar bones.

FRONTAL FURROW extends upwards from the frontal spine, and becomes gradually larger in its course, to lodge the upper part of the superior longitudinal sinus, and to give attachment to the falx cerebri.

FRONTAL NERVE, *Palpebro-frontal*—(Ch.), is the greatest of the three branches of the ophthalmic nerve,—the first division of the 5th pair.

It proceeds along the superior paries of the orbit, and divides into two branches:—the one, *internal*, which makes its exit from the orbital fossa, passing beneath the pulley of the oblique muscle:—the other, *external*, issuing from the same cavity by the foramen orbitarium superius.

FRONTAL PROTUBERANCE, *F. tuberosa* ity. *Tuber frontallē*, *Suggrun'dium superciliorum*. The protuberance of the frontal bone above the superciliary arch.

FRONTAL SINUSES, *Frontales Sinus*, *Metopop'tra*, *Prosopop'tra*, *Sinus Supercilia'res* seu *pituitarii frontis*, *Cavernæ frontis*, are two deep cavities in the substance of the frontal bone, separated from each other by a median septum, and opening, below, into the anterior cells of the ethmoid bone.

FRONTAL SPINE, *Crista inter'na*, is situate in the middle of the under part of the bone, and is formed by the coalescence of the inner tables for the attachment of the falx cerebri.

Surgeons have given the name *FRONTALIS* to a bandage or topical application to the forehead. Such have, also, been called *τρομετωπία* and *αναπολληρατι*.

FRONTAL TUBEROSITY, Frontal protuberance.

FRONTAL'LE, from *frons*, 'the forehead.' A medicine applied to the forehead.

FRONTALIS ET OCCIPITALIS, Occipito-frontalis.

FRONTALIS VERUS, Corrugator supercilii.

FRONTODYMIA, see Cephalodymia.

FRONTO-ETHMOID FORAMEN, Cæcum foramen—*f. Nasal*, Pyramidalis nasi—*f. Sourcilier*, Corrugator supercilii.

FROSTBITE, Congelation.

FROSTPLANT, *Helianthemum Canadense*.

FROSTWEED, *Erigeron Philadelphicum*, *Helianthemum Canadense*.

FROSTWORT, *Helianthemum Canadense*.

FROTH'Y, from Gr. *αφρός*, 'froth.' [?] *Spu'mous*, (F.) *Spumeux*, *Écumeux*, *Mousseux*. An epithet given to the faeces or sputa, when mixed with air.

FROTTEMENT, see *Bruit de frottement*.

FROTTEMENT GLOBULAIRE. A name given by M. Simonnet, to the pulse in aortic regurgitation, when it is jerking; and, in well marked cases, appears as if the blood consisted of several little masses, which passed in succession under the finger applied to the artery.

FRUCTIFICATION, Fecundation.

FRUCTUS HORÆI, Fruit, (summer)—*f. Immaturus*, Abortion.

FRUGIVOROUS, *Frugivorus*, from *fruges*, 'fruits,' and *voro*, 'I eat.' One that eats fruits.

FRUIT, *Fructus*, from *fruo*, *fructus*, 'to enjoy.' *Carpus*. In botany, the seed with its enclosing pericarp. In a medical sense it may be defined to be,—that part of a plant which is taken as food. The effects of fruits on the body, in a medical as well as a dietetical point of view, are various. They may be distinguished into classes: for, whilst the *Cerealia*, for example, afford fruits, which are highly nutritious; the *Summer Fruits* (*Fructus Horæi*), which include strawberries, cherries, currents, mulberries, raspberries, figs, grapes, &c., are refrigerant and grateful, but afford little nourishment.

FRUIT-SUGAR, Glucose.

FRUMENT, *Fru'menty*, *Fur'menty*, from *frumentum*, 'wheat' or 'grain,' quasi *frugimentum*; from *fruges*, 'fruit.' Pottage made of wheat. Food made of wheat boiled in milk.

FRUMENTUM. Same etymon. *Sitos*. Any kind of grain from which bread was made;—especially wheat.

FRUMENTUM, Triticum—*f. Corniculatum*, *Ergot*

—f. Cornutum, Ergot—f. Luxurians, Ergot—f. Temulentum, Ergot—f. Turgidum, Ergot.

FRUSTRATOIRE, (F.) Any liquor, taken a short time after eating, for the purpose of assisting digestion when difficult. Sugared water, eau sucrée, or water with the addition of a little brandy, or some aromatic substance, is commonly used for this purpose.

FRUTEX BACCIFER BRAZILIENSIS, Caa-guayo—f. Indicus spinosus, Cara schulli.

FUCUS, F. vesiculosus, Paint.

FUCUS AMYLACUS, Jaffna, Edible or Ceylon Moss, *Plocaria candida*, *Gracilaria* seu *Sphaerococcus* seu *Gigartina* seu *Fucus lichenoides*, Marine Moss. This moss belongs to the natural order Algae. It was introduced some years ago into England, from India. It is white, filiform and fibrous, and has the usual odour of sea-weeds. Its medical properties are similar to those of Irish moss.

FUCUS, BLADDER, F. vesiculosus.

FUCUS CRISPUS, F. *Irlandicus*, Lichen *Carra-geen*, *Chondrus* (Ph. U. S.), *Ch. crispus* seu *polymorphus*, *Sphaerococcus crispus*, *Ulva crispus*, Irish Pearl, *Carraheen* or *Corrigreen* moss, (F.) *Mousse d'Irlande* ou *perle*. This Fucus is found on the coasts of Scotland, Ireland, Western France, Spain, and Portugal, and as far as the tropics. It is also a native of the United States. In Ireland, it is used by the poor as an article of diet. As met with in America, it is of a light yellow colour, and resembles plates of horn, crisped and translucent. An agreeable jelly is obtained from it by boiling it in water or milk, which forms a good article of diet in consumptive cases. Its properties are indeed exactly like those of the Iceland Moss.

FUCUS HELMINTHOCORTON, *Corallina Corsicana*—f. *Infatus*, F. vesiculosus—f. *Irlandicus*, *Fucus crispus*—f. *Lichenoides*, F. *amylaceus*—f. *Saccharine*, *Rhodomela palmata*.

FUCUS VESICULOSUS, *Fucus*, F. *inflatus*, *Hallidrys vesiculosa*, *Quercus Marina*, Bladder Fucus, Sea Oak, Sea Wrack, Yellow Bladder Wrack, (F.) *Varec vésiculeux*, *Chêne marin*. It has been said to be a useful assistant to sea-water, in the cure of disorders of the glands. When the wrack, in fruit, is dried, cleaned, exposed to a red-heat in a crucible with a perforated lid, and is reduced to powder, it forms the *Æthiops vegetabilis*—the *Pulvis Quercus Marinae* of the Dublin Pharmacopœia—which is used, like the burnt sponge, in bronchocele and other scrofulous swellings. Its efficacy depends on the iodine it contains. Dose, gr. x to ʒij, mixed in molasses or honey. See Soda.

FUGA DÆMONUM, *Hypericum perforatum*.

FUGACIOUS, *Fugax*, *fugacis*, (F.) *Fugace*, from *fugere*, 'to fly.' An epithet given to certain symptoms, which appear and disappear almost immediately afterwards; as a *fugacious redness*,—a *fugacious swelling*, *Tumor fugax*, &c.

FUGAX, *Fugacious*.

FUGE, from *fugo*, 'I expel,' 'an expeller.' A common suffix. Hence, *Febrifuge*, *Vermifuge*, &c.

FUGILĒ, *Fugilla*. This term has several acceptations. It means, 1. The cerumen of the ear. 2. The nebulous suspension in, or deposition from, the urine. 3. An abscess near the ear.—Ruland and Johnson. 4. Abscess in general.

FUGILLA, *Fugile*.

FULGUR, *Astrape*.

FULIGINOUS, *Fuliginosus*, *Lignyoides*, (F.) *Fuliginacea*, from *fuligo*, *fuliginis*, 'soot.' Having the colour of soot. An epithet given to certain parts, as the lips, teeth, or tongue, when they assume a brownish colour, or rather are covered with a coat of that colour. *Fuliginosity*.

FULIGO, genitive *fulig'iniæ*, *Lignys*, *Soot*, (F.) *Suie*, &c., *Woodsoot*, *Fuligo Ligni*, consists of volatile alkaline salt, empyreumatic oil, fixed alkali, &c. A tincture, *Tinctura Fulig'iniæ*, prepared from it, has been recommended as a powerful antispasmodic in hysterical cases. (*Fulig. lign.* ʒij; *potass. subcarb.* lbss; *ammon. muriat.* ʒi; *aqua fluvial.* Oijj. Digest for three days.) This tincture bears the name *Soot drops* and *Fit drops*. An ointment of soot has been used in various cutaneous diseases.

FULIGO ALBA PHILOSOPHUM, *Ammonia murias*.

FULIGO KALI, from *fuligo*, 'soot,' and *kali*, 'potassa.' This is an analogous preparation to *anthrakali*; soot being used in the place of coal. It is employed in the same diseases. What might be regarded as a weak solution of *fuligo-kali* has been used for many years in Philadelphia, under the names *medical lye*, *soot tea*, *alkaline solution*, *dyspeptic lye*, and *Physick's lye tea*. It is made by mixing a quart of *hickory ashes*; half a pint of *soot*; and a gallon of *boiling water*; allowing the mixture to stand for twenty-four hours, and decanting.

FULLERS' EARTH, *Cimolia purpureoens*.

FULMEN, *Astrape*.

FULMICOTON, see *Collodion*.

FULNESS, *Repletio*, *Plenitudo*, *Plethora*, *Repletio*. The state of being filled. Also, a feeling of weight or distension in the stomach or other part of the system.

FUMARIA, *Fumaria officinalis* seu *medica*, *Fumus terra*, *Capnos*, *Herba melancholifuga*, *Solamen Scabiosorum*, *Fumitory*, *Common Fumitory*, *Fumiterra*, (F.) *Fumeterre*, *Fiel de terre*, *Pisse-sang*. Ord. *Fumariaceæ*. The leaves are extremely succulent, and have a bitter, somewhat saline, taste. The infusion of the dried leaves and the expressed juice of the fresh plant have been extolled for their property of clearing the skin of many disorders of the leprosy kind.

FUMARIA BULBOSA, *F. cava* seu *major*, *Borchhansenia* seu *Capnoi-des cava*, *Aristolochia fabacea* seu *cava* seu *vulgaris rotunda*, *Corydalis bulbo* seu *cava* seu *tuberosa*, *Capnorhis*. The root of this plant was formerly given as an emmenagogue and anthelmintic. (F.) *Fumeterre bulbeuse*.

FUMARIA CAVA, *F. bulbosa*—f. *Major*, *F. bulbosa*—f. *Media*, *Fumaria*—f. *Officinalis*, *Fumaria*.

FUMETERRE, *Fumaria*.

FUMIGATIO, *Fumigation*—f. *Antiloinica* *Gaubii*, see *Disinfection*—f. *Guytoniensis*, see *Disinfection*—f. *Smythiana*, see *Disinfection*.

FUMIGATION, *Fumigatio*, from *fumus*, 'smoke.' *Suffitus*, *Suffitio*, *Suffuminatio*, *Suffumigatio*, *Suffimen*, *tum*, *Apocynus mus*, *Thymidum*, *Epithymidum*, *Hypothymidum*, *Hypothymiasis*, *Thymiasis*, *Capnis mus*, *Hypercynus mus*, *Hypatmus*, *Hypatmis mus*, *Anathymiasis*. An operation, the object of which is to fill a circumscribed space with gas or vapour, with the intention either of purifying the air, of perfuming it, or of charging it with a substance proper for acting upon a part of the surface of the human body. See *Sulphurous acid*. Hence, *fumigations* have been distinguished into *aqueous*, *aromatic*, *sulphureous*, *mercurial*, *disinfecting*, *Guytonian*, &c. Benzoin generally constitutes the chief ingredient in the *Fumigating Pastilles*, to which any variety of odoriferous substances may be added. The following are formulæ.

R. Benzoin. ʒi; *cascarilla*. ʒss; *myrrh*. ʒj; *ol. myrist.*, *ol. caryoph.* ʒā gr. x; *potassæ nitrat.* ʒss; *carbon. lign.* ʒvj; *mucil. trag.* q. s.—or

R. Benzoin. p. xvj; *Balsam. Tolut.* p. iv; *Santal. alb.* p. iv; *Labdan.* p. j; *Carbon. Lign.* p. xlvij; *Potass. nitrat.* p. ij; *Tragacanth.* p. j;

Acacia, p. ij; *Aq. cinnam.* p. xij. Reduce the solid ingredients to powder, and mix the whole into a plastic mass, which must be formed into cones, flattened at the base, and dried, first in the air, and afterwards in a stove.

FUMIGATION, *Chloirine*, see Disinfection — *f.* Nitrous, see Disinfection — *f.* Oxy muriatic, see Disinfection.

FUMITERRA, *Fumaria*.

FUMITORY, *Fumaria*.

FUMUS ALBUS, *Hydrargyrum* — *f.* Citrinus, Sulphur — *f.* Terre, *Fumaria*.

FUNAMBULATIO, from *funis*, 'a cord,' and *ambulare*, 'to walk.' An exercise with the ancients, which consisted in scaling ropes.

FUNCTION, *Functio*, *Actio*, (*F.*) *Fonction*; from *fungor*, *functus*, 'I act,' 'perform.' The action of an organ or system of organs. Any act necessary for accomplishing a vital phenomenon. A function is a special office in the animal economy, which has, as its instrument, an organ or apparatus of organs. Thus respiration is a function. Its object is the conversion of venous into arterial blood, and its instrument is the lungs. The ancient physiologists divided the functions into *vital*, *animal*, and *natural*. They called *vital functions* those which are essential to life, as innervation, circulation, respiration; *animal functions*, those which belong to the encephalon; viz.: the functions of the intellect, the affections of the mind, and the voluntary motions; and *natural functions*, *Facultates* seu *Actiões naturales*, those relating to assimilation, such as the actions of the abdominal viscera, of the absorbent and exhalant vessels, &c. Bichat divided the functions into those which relate to the preservation of the individual, and those that relate to the preservation of the species. The former he subdivided into *animal* and *organic*. The *animal functions* or *functions of relation* are those of the intellect, sensation, locomotion, and voice. The *organic functions* include digestion, absorption, respiration, circulation, secretion, nutrition, and calorification. The *functions*, whose object is the preservation of the species—the *organic, nutritive*, or *vegetative functions*—are all those that relate to generation;—such as conception, gestation, accouchement, &c. Each of these admits of numerous subdivisions in a complete course of *Physiology*;—for so the doctrine of the functions is called.

FUNCTIONAL, *Functionalis*, (*F.*) *Fonctionnel*. Relating or belonging to the function of an organ, or to the functions in general.

Functional or *vital phenomena* are those produced by some modification in the action of an organ or organs, in contradistinction to those of a material or physical character;—thus we distinguish the *functional phenomena* of phthisis from those indicated by physical diagnosis—the *physical phenomena*.

FUNCTIONAL DISEASES, see Organic Diseases.

FUNDA, 'a sling.' *Sphenodon*, (*F.*) *Fronde*. A bandage, composed of a fillet or long compress, cleft at its extremities to within about two inches of its middle. It is used in diseases of the nose and chin, and especially in cases of fracture of the lower jaw. In such case it has, also, been called *Mentonnière*, because placed beneath the chin; from (*F.*) *Menton*, 'the chin.'

FUNDAMENT, Anus—*f.* Falling down of the, Proctocele.

FUNDAMENTAL, from *fundare*, 'to found; to lay the bottom of any thing.' (*F.*) *Fondamental*. Some anatomists have called the sacrum *Os Fundamentale*, because it seems to serve as a base to the vertebral column. The *sphenoid bone* has likewise been so denominated, from its being situate at the base of the cranium.

FUNDUS, 'the bottom.' (*F.*) *Fond*. The base of any organ which ends in a neck, or has an external aperture;—as the *Fundus vesicæ*, *F. uteri*, &c. Also, the *Vulva*.

FUNDUS VAGINÆ, *Laquear vaginæ*.

FUNES CORDIS, *Columnæ carnae*—*f.* Semicirculares, Semicircular canals.

FUNGIFORM PAPILLÆ, see Papillæ of the Tongue.

FUNGOID, *Fungoides*, *Mycoides*, *Fongiformis*, *Fun'giform*, (*F.*) *Fonguide*, *Fongiforme*, from *fungus*, 'a mushroom,' and *idos*, 'resemblance.' That which has the shape of, or grows in some measure like a mushroom, as the *fungoid* or *fungiform papillæ* of the tongue.

FUNGOID DISEASE, Encephaloid.

FUNGOSITY, *Fungositas*, *Caro laceriana* seu *fungosa*, *Escarco'ma*, *Hyperescarco'ma*, *Hyperescarco'is*, *Proud Flesh*, (*F.*) *Fonguité*. The quality of that which is fungous:—fungous excrecence, *Exerescencia fungosa*. The fungosities which arise in wounds or ulcers are easily repressed by gentle compression, dry lint, the sulphas cupri, or other gentle caustics. At times, the more powerful are necessary, and sometimes excision is required.

FUNGOUS, *Fungosus*, (*F.*) *Fongueux*. Having the characters of fungosity or fungus—as a *fungous* wound.

FUNGUS, *Mycæ*, (*σφογγος*), (*F.*) *Fungus*, *Champignon*. The mushroom order of plants; class Cryptogamia, in the Linnean system. In *Pathology*, the word is commonly used synonymously with fungosity, *mycosis*. M. Breschet has proposed to restrict the term *fungosity* to vegetations which arise on denuded surfaces, and to apply the term *fungus* to the tumours which form in the substance of the textures, without any external ulceration. Fici and warts, for example, would be fungi of the skin.

FUNGUS ALBUS SALICINUS, *Dædalea suaveolens*—*f.* *Articuli*, *Spina ventosa*—*f.* Bleeding, *Hæmatodes fungus*—*f.* *Cancerosus hæmatodes*, *Hæmatodes F.*—*f.* *Cancerosus medullaris*, see Encephaloid—*f.* *Cerebralis*, see Encephaloid—*f.* *Cerebri*, *Encephalocele*—*f.* *Chirurgorum*, *Lycoperdon*—*f.* *Cynosati*, *Bedeguar*—*f.* *Faginosus*, *Morchella esculenta*—*f.* *Hæmatodes*, *Hæmatodes fungus*—*f.* *Igniarius*, *Boletus igniarius*—*f.* *Laricis*, *Boletus laricis*—*f.* of Malta, *Cynomorion coccineum*—*f.* *Medullaris*, see Encephaloid—*f.* *Melanoides*, *Cancer*, *melanotic*—*f.* *Melitensis*, *Cynomorion coccineum*—*f.* *Petræus marinus*, *Umbilicus marinus*—*f.* *Quercinus*, *Boletus igniarius*—*f.* *Rosarum*, *Bedeguar*—*f.* *Salicis*, *Dædalea suaveolens*—*f.* *Sambucinus*, *Peziza auricula*.

FUNIC BELLOWS' SOUND, see Bellows' Sound, *funic*.

FUNICULI GRACILES, (diminutive of *funis*, 'a cord,') *Posterior Median Columnæ* or *Funiculi of the medulla oblongata*. Along the posterior border of each corpus restiforme, and separated from it by a groove, is a narrow white cord, separated from its fellow by the *fissura longitudinalis posterior*. The pair of cords are the *funiculi graciles*. Each funiculus forms an enlargement—*processus clavatus*—at its upper end, and is then lost in the corpus restiforme.

FUNICULI SILIQÆ. Longitudinal fibres seen in the groove which separates the corpus olivare from the corpus pyramidale and corpus restiforme. They enclose the base of the corpus olivare,—those which lie on its inner side forming the *funiculus internus*; and those on its outer side the *funiculus externus*.

FUNICULUS, Cord—*f.* *Externus*, see *Funiculi siliquæ*—*f.* *Internus*, see *Funiculi siliquæ*—*f.* *Spermaticus*, *Spermatic cord*—*f.* *Tympani*, *Chorda tympani*.

FUNICULUS UMBILICALIS, *Funis umbilicalis*, *Intestinulum*, *Vinculum umbilicale*, *Umbilical cord*, *Navel string*, diminutive of *Fanis*, 'a cord.' (F.) *Cordon ombilicale*. A cord-like substance, which extends from the placenta to the umbilicus of the foetus. It is composed of the chorion, amnion, an albuminous secretion called the *Jelly of the Cord*, cellular substance, an umbilical vein, and two umbilical arteries. The former conveys the blood from the placenta to the foetus—the latter return it. All these parts are surrounded by a sheath—*Investitura seu Vagina funiculi umbilicalis*. Its usual length is from 16 to 22 inches.

FUNICULUS VARICOSUS, *Cirsocele*.

FUNIS, Cord, Laqueus—f. *Argentens*, *Medulla spinalis*—f. *Hippocristis*, *Achillis tendo*—f. *Umbilicalis*, *Funiculus umbilicalis*.

FUNK, *Boletus igniarius*.

FUNNEL, see *Infundibulum*.

FUR, *Enduit*.

FURCELLA, *Furcula*; diminutive of *furca*, 'a fork.' The upper part of the sternum, the clavicle. The *Fourchette*.

FURCELLA INFERIOR, Xiphoid cartilage.

FURCHMUEHL, MINERAL WATERS OF. These Bavarian springs contain carbonic acid, sulphuretted hydrogen, carbonates of lime and soda; chlorides of lime and magnesium, oxides of iron and magnesium, &c.

FURCIFERA, *Penis*.

FURCILLA, *Fourchette*.

FURCULA, *Furcella*, *Clavicle*, *Fourchette*.

FUREUR UTÉRINE, *Nymphomania*.

FURFUR, (from *far*, 'bran,') *Bran*, *Pit'gron*, *Ach'gron*, *Apobras'na*, *Lemna*, *Canabru'no*, (F.) *Son*. The decoction is sometimes employed as an emollient.

FURFURA, *Scarf*.

FURFURACEOUS, from *furfur*, 'bran.' *Scurfy*, *Canicaceous*, *Pithyri'nus*, *Pityroides*, *Pityro'des*, (F.) *Furfuracé*. Resembling bran. A name given to eruptions, in which the epidermis is detached in small scales resembling bran. Also, a bran-like sediment observed at times in the urine;—*Urina furfuracea*, *Sedimentum Urinæ pityroides*.

FURFURAMIDE, see *Furfurine*.

FURFURATIO, *Desquamation*, *Porrigo*, *Pityriasis*.

FUR/FURINE, *Furfuri'na*, from *furfur*, 'bran.' By the action of dilute sulphuric acid on corn meal or bran an oil is obtained—*furfurol* or *furfurole*. By the action of ammonia on this, *furfuramide* or *furfurolamide* results; and by the action of dilute potassa on this the alkaloid *furfurine* is obtained. This was found by Professor Simpson, of Edinburgh, to possess tonic, if not antiperiodic, properties.

FURFURISCA, *Pityriasis*.

FURFUROL, see *Furfurine*.

FURFUROLAMIDE, see *Furfurine*.

FURIA INFERNALIS. A kind of vermiform insect, scarcely two lines long, common in Sweden, which flies about and stings both man and animals, exciting the most excruciating torture.

FURIBUNDUS, *Maniodes*.

FURIOSUS, *Maniodes*.

FURIOUS, *Maniodes*.

FURMAGE, *Cheese*.

FURMENTY, *Frument*.

FURNAS, MINERAL WATERS OF. A

thermal chalybeate water in St. Michael's, Azores, which contains carbonic acid, and carbonate of iron.

FURUNCLE, *Furunculus*.

FURUNCLE GUÉPIER. A malignant boil, *Wasp's nest boil*, which generally attacks the nape and region of the neck, and rarely others than old people. Hence it has been called *Old People's boil*.

FURUNCULEUX, *Furuncular*.

FUROR, *Mania*—f. *Brevis*, *Rage*—f. *Mania*, *Mania*—f. *Uterinus*, *Nymphomania*.

FURRED, see *Enduit*.

FURROW, **MENTO-LABIAL**, see *Mento-labial furrow*.

FURUNCULAR, *Furunculous*, *Furunculous*, (F.) *Furunculenz*. Relating or appertaining to furunculus,—as 'a furuncular epidemic.'

FURUNCULI ATONICI, *Ecthyma*—f. *Ventriculus*, see *Furunculus*.

FURUNCULOID, *Furunculoïdes*. A hybrid term, from *furunculus*, and *αἶος*, 'resemblance.' Resembling furunculus,—as 'a furunculoid epidemic.'

FURUNCULUS, from *furere*, 'to be mad.' *Chi'adus*, *Chi'oli*, *Doth'ien*, *Dothion*, *Furunculus suppuratorius*, *F. Versus*, *F. benignus*, *Phyma furuncululus*, *Abscessus nucleatus*, a *furuncle*, a *boil*, a *bile*, (Sc.) *Bruick*, *Bruck*, (Prov.) *Pinsaveal*, (F.) *Furonce*, *Fronce*, *Clou*. A small phlegmon, which appears under the form of a conical, hard, circumscribed tumour, having its seat in the dermoid texture. At the end of an uncertain period, it becomes pointed, white or yellow, and gives exit to pus mixed with blood. When it breaks, a small, grayish, fibrous mass sometimes appears, which consists of dead areolar tissue. This is called the *Core*, *Setfast*, *Ventriculus* seu *Nucleus Furuncululi*, (F.) *Bourbillon*. The abscess does not heal until after its separation. The indications of treatment are,—to discuss by the application of leeches and warm fomentations,—or, if this cannot be done, to encourage suppuration by warm, emollient cataplasms. When suppuration is entirely established, the part may be opened or suffered to break, according to circumstances.

The term *blind boil* is given to an indolent, imperfectly suppurating, phlegmonous tumour, of the kind described above, which is often seated in a sebaceous follicle, as in *Acne indurata*.

FURUNCULUS GANGRÆNOSUS, *Anthrax*—f. *Malignus*, *Anthrax*.

FUSARIA VERMICULARIS, *Ascaris vermicularis*—f. *Visceralis et renalis*, *Strongylus gigas*.

FUSÉE PURULENTE (F.). The long and sinuous route which pus takes, in certain cases, in making its way to the surface. These *Fusées* almost always form beneath the skin between the muscles; or along aponeuroses, bones, tendons, &c.

FUSEL OIL, see *Oil, fusel*.

FUSIBLE, see *Fusion*.

FUSIBILITY, see *Fusion*.

FUSION, *Fu'sio*, *Melting*, *Liquefaction*; from *fundere*, *fusum*, 'to melt.' In chemistry, the transition of a solid body into a liquid by the aid of heat. Substances capable of such transition are said to be *fusible*; or to be possessed of *fusibility*.

FUSTIC TREE, *Cladastrix tinctoria*.

FUTUTRIX, *Tribas*.

G.

THE Greek G, Γ, with the ancient Greek physicians, signified an ounce.—Rhod. ad Scribonium.

GAB, Mouth.

GABALLA, Cabal.

GABELLA, Mesophryon.

GABIR'EA, γαβίρα. A fatty kind of myrrh, mentioned by Dioscorides.

GADUS ÆGLIFINUS, see *Oleum jecoris aselli*—*g.* Callarius, see *Oleum jecoris aselli*—*g.* Carbonarius, see *Oleum jecoris aselli*—*g.* Lota, see *Oleum jecoris aselli*—*g.* Merluccius, see *Oleum jecoris aselli*—*g.* Morrhuæ, see *Oleum jecoris aselli*—*g.* Pollachius, see *Oleum jecoris aselli*.

GEOPHAGIA, Geophagism.

GEOPHAGUS, from γαῖα, 'earth,' and φάγω, 'I eat.' One who eats earth.

GAGEL, Myrica gale.

GAGUE SANG, Cague-sang.

GAJET, Coyot.

GAÏAC, Guaiacum.

GAILET ACCROCHANT, Galium aparine—*g.* Crochant, Galium aparine—*g.* Jaune, Galium verum—*g.* Vrai, Galium verum.

GAINE, Vagina or sheath—*g.* de l'Apophyse styloïde, Vaginal process of the temporal bone—*g.* de la Veine porte, Vagina or sheath of the vena porta.

GAITLING, Infans.

GALA, γάλα, genuine γαλακτος, milk; hence:

GALACTACRA'SIA, from γάλα, and ακρασία, 'imperfect mixture.' A morbid mixture or constitution of the milk.

GALACTACRATIA, Galactia.

GALACTE'MIA, from γάλα, and αἷμα, 'blood.' A condition of the blood in which it contains milk.

GALACTAGOGA, Galactopoetica.

GALACTAPOSTEMA, Mastodynia apostematosa.

GALACTIA, Galactorrhœa, Galactorrhœa, from γάλα, Lactis redundantia, Polygalactia, Galactozemia. A redundant flow of milk, either in a female who is suckling, or in one who is not. It may occur without being provoked by suckling. When to a great extent, it sometimes causes wasting; *Tubes lactea*, *T. nutritivum*. Dr. Good uses *Galactia*, in his Nosology, for 'morbid flow or deficiency of milk,' *Galactocratia*.

GALACTICUS, Lactic.

GALACTIDRO'SIS, from γάλα, and ἰδρωσις, 'sweating.' Sweating of milk.

GALACTIFER, Galactophorous.

GALACTINE, Casein.

GALACTINUS, Lactic.

GALACTIRRHOEA, Galactia.

GALACTIS, Galaxias.

GALACTISCHESIS, Galactoschesis, Lactis retentio, from γάλα, and ἰσχειν, 'to restrain.' Retention or suppression of milk.

GALACTITES, Galaxias.

GALACTOCATARACTA, Cataract, milky.

GALACTOCE'LE, Galactoncus, Lactoce'le, from γάλα, and κηλη, 'a tumour.' *Tumor lacteus*. Distension of one or more of the galactophorous sinuses by milk. Lactiferous engorgement. The French use the term *Poile* for such engorgement; a relic of an idea, as old as Aristotle, that a hair swallowed may have obstructed one of the milk tubes, which must be got rid of by sucking.

The name has, also, been given by Vidal de Cassis to a tumour produced by an effusion of a

white liquid into the tunica vaginalis, which he considered to be true milk.

GALACTODEN'DRON U'TILE, from γάλα, and δένδρον, 'a tree.' *Palo de Vacca*, *Arbol de Leche*, *Coco tree* of South America. *Family*, Urticaceæ. When an incision is made into the trunk, a fluid similar to milk flows out, which is agreeable and nutritious.

GALACTO'DES, from γάλα, and εἶδος, 'resemblance.' In Hippocrates, the term signifies milk-warm, and likewise a milky colour, as of the urine—*uri'na galactodes*.

GALACTODIÆ'TA, *Diæt'a lactea*, from γάλα, and διαίτα, 'diet.' A milk diet.

GALACTODIARRHOEA, Cœlic Flux.

GALACTOGANGLION, Milk knot.

GALACTOHE'MIA, *Galacthœ'mia*, *Galæ'mia*, from γάλα, and αἷμα, 'blood.' *Lactis sanguinolenti Excretio*. The secretion of bloody or bloodlike milk.

GALACTOMASTOPARECTOMA, Mastodynia apostematosa.

GALACTOMETER, *Lactom'eter*, from γάλα, and μετρον, 'measure.' An instrument for appreciating the quantity of cream in milk. It is a kind of graduated separatory or *éprouvette*—the degrees on the scale indicating the thickness of the layer of cream that forms on the surface of the milk.

GALACTONCUS, Galactocele.

GALACTOPHAGOUS, *Galactoph'agus*, *Lactiv'orus*, sometimes used substantively; from γάλα, and φάγω, 'I eat.' That which feeds on milk. A name given to certain people, with whom milk appears to constitute the chief nourishment.

GALACTOPHORA, Galactopoetica.

GALACTOPHORITIS. Inflammation of the galactophorous ducts. It has been inaccurately used for ulceration of the top of the nipple towards their orifices.

GALACTOPHO'ROUS, Lactiferous—*g.* Ducts, Lactiferous ducts—*g.* Receptacle, see Lactiferous vessels—*g.* Sac, see Lactiferous vessels—*g.* Sinuses, see Lactiferous vessels.

GALACTOPH'ORUS. Some accoucheurs have given this name to an instrument intended to facilitate sucking, when the faulty conformation of the nipple prevents the child from laying hold of it.

GALACTOPH'YGUS, from γάλα, and φεύγειν, 'to shun.' That which arrests or disperses the secretion of milk. Hence *Galactoph'ygæ Medicament'a*.

GALACTOPLA'NIA, *Metastasis* seu *Aberratio* seu *Via extraridinalis lactis*, *Galactorrhœa erronea*, from γάλα, and πλανή, 'wandering.' Extravasation of milk into the areolar membrane. Secretion of milk elsewhere than from the breasts.

GALACTOPLERO'SIS, from γάλα, and πληρωσις, 'repletion.' Redundance of milk.

GALACTOPOEA, Galactopoetica.

GALACTOPOESIS, Galactosis.

GALACTOPOET'ICA, *Galactoph'ora*, *Galactag'ga*, *Galactopœ'a*, from γάλα, and ποιέω, 'I make.' Substances to which has been attributed the property of favouring the secretion of milk and augmenting its quantity.

GALACTOPOIESIS, Galactosis.

GALACTOPO'SIA, from γάλα, and ποσις,

'drink.' The drinking of milk. Treatment of a disease by means of milk.

GALACTOPOTES, *Galactop'otus*, *Lactip'otor*. Same etymon. A drinker of milk. One subjected to a milk diet.

GALACTOPYRA, Fever, milk.

GALACTOPYRETUS, Fever, milk.

GALACTORRHOEA, *Galaetia*—g. Erronea, *Galactop'lania*—g. *Saccharata*, *Saccharorrhoea lactea*.

GALACTOSACCHARUM, *Saccharum lactis*.

GALACTOSCHESIS, *Galactischesis*.

GALACTO'SIS, *Galactopoe'is*, *Galactopoe'is*, *Secretio lactis*; from *γαλα*. The secretion or formation of milk.

GALACTOTROPHE, *Galactotroph'ia*, from *γαλα*, and *τροφή*, 'nourishment.' Nourishment by means of milk.

GALACTOZE'MIA, from *γαλα*, and *ζημία*, 'loss.' Loss of milk. Also, *Galaetia*.

GALACTU'CHOS, from *γαλα*, and *εχειν*, 'to have.' Suckling. Giving milk.

GALACTURIA, *Chyluria*.

GALÆMIA, *Galactohæmia*.

GALANGA, *Maranta galanga*.

GALANGALE, *Cyperus longus*.

GALARHÆUS LATHYRIS, *Euphorbia lathyris*—g. *Palustris*, *Euphorbia palustris*.

GALARIPS, *Allamanda*.

GALAX APHYLLA, *G. rotundifolia*, *Erythrorrhiza rotundifolia*; *Carpenter's leaf*, *Beetleweed*; indigenous; Order *Ericaceæ*. The root is astringent, and the leaves, in domestic practice, are applied to cuts and wounds.

GALAXIA, Thoracic duct.

GALAX'IAS, *Galaet'ies*, *Galaet'is*; from *γαλα*. A milk stone. A stone supposed to be capable of promoting the secretion of milk.

GALBANUM, see *Bubon galbanum*—g. *Long-leaved*, *Bubon galbanum*.

GAL'BULUS, from *galbus*, 'yellow.' A kind of congenital jaundice, in which the yellow colour continues through life. It is rather a defect in colour than a disease.—Vogel.

GALE, *Myrica gale*—g. *Odorant*, *Myrica gale*—g. *Scotch*, *Myrica gale*—g. *Sweet*, *Myrica gale*.

GALE, *Psora*—g. *Canine*, *Psoriasis*—g. *Épidermique*, *Eczema*—g. *Miliaire*, *Psoriasis*—g. *Sèche*, *Lichen*, *Psoriasis*.

GA'LEA. A helmet, from *γαλεα*, 'a cat;' of the skin of which it was formerly made. [?] A name given to the *amios*, and also to the *bandage of Galen*. In *Pathology*, it indicates a headache affecting the whole head. See, also, *Caul*.

GALEA APONEUROTICA CAP'ITIS, *Ga'lea tendinea Santorini*, *Ga'lea cap'itis*, *Membra'na epierania*. The tendinous expansion which unites the frontal and occipital portions of the occipito-frontalis muscle.

GALEAMAUROSIS, Amaurotic cat's eye.

GALEAN'CON, *Galian'con*, from *γαλεα*, 'a cat,' 'a weasel,' and *αγκων*, 'an elbow.' *Mustela'neus*. One who has two short arms.

GALEAN'THROPY, *Galeanthro'pia*, from *γαλεα*, 'a cat,' and *ανθρωπος*, 'a man.' A variety of melancholy in which the patient believes himself changed into a cat. An affection similar to lycanthropy and cynanthropy.

GALE'GA, *G. officinalis* seu *gal'aris* seu *Per'sica*, *Ruta capra'ria*, *Goat's Rue*, (F.) *Rue de chèvre*, *Fane Indigo*. Ord. *Leguminosæ*. It is slightly aromatic, and was once used as a sudorific and alexiterial in malignant fevers, &c.

GALEGA APOLLINEA, *Tephrosia apollinea*—g. *Persica*, *Galega*—g. *Purpurea*, *Tephrosia apollinea*—g. *Toxicaria*, *Tephrosia toxicaria*.

GALEGA VIRGINIA'NA, *Tephrosia Virginia'na*, *Turkey Pea*, *Hoary Pea*, *Devil's shoestrings*, *Vir-*

ginia Goat's rue or *cat-gut*, is used in some parts of the United States as an anthelmintic. The decoction of the root is given.

GALEGA VULGARIS, *Galega*.

GALENE, *Graphites*.

GALENEA, *Graphites*.

GALEN'IC, *Galen'ical*, *Galen'icus*, *Gale'nus*, (F.) *Galénique*, from *Galenus*. That which relates to the doctrine of Galen or to Galenism. Used, substantively, for drugs that are not chymical.

GALENIC MEDICINE, *Galenism*.

GA'LENISM, *Galen'ic med'icine*. The doctrine of Galen.

GA'LENIST, *Galenis'ta*, *Galenis'tes*. A follower of the doctrine of Galen.

GALENIUS, *Galenic*.

GALEODOLON, *Galeopsis*.

GALEOPDOLON, *Galeopsis*.

GALEOP'SIDE, *Galeopsis grandiflora*.

GALEOPSIS, *Lamium album*.

GALEOP'SIS, *Galiop'sis*, *Galeob'dolon*, *Galeop'dolon*, *Le'mium rubrum*, *Urti'ca iners* magna *fec-tidis'sima*, *Stachys fœ'tida*, *Hedge nettle*, (F.) *Ortie morte des bois*. Ord. *Labiatae*. From *γαλεα*, 'a weasel,' and *οψis*, 'appearance;' the corolla having been supposed to resemble the head of the weasel. This plant was formerly reckoned a vulnerary and anodyne.

GALEOPSIS ANGUSTIFOLIA, *G. grandiflora*—g. *Dubia*, *G. grandiflora*.

GALEOP'SIS GRANDIFLO'RA, *G. Ochroleuca* seu *la'danum* seu *angustifolia* seu *du'bia* seu *prostra'ta* seu *villo'sa*, *Tetrahit longiflorum*, *G. Seg'tetum*, *Herba Sideri'tidis*, (F.) *Galeopsiside*, *Chanvre bâtarde*. This plant is regarded in Germany as a bitter resolvent. It is the basis, also, of a celebrated nostrum, the *Blankenheimer Tea*, called likewise *Lieber's pectoral* and *phthisical herbs* (*Liebersche Brust oder Auszehrungs-Krauter*), which has enjoyed great repute in pectoral complaints. The tops of the plant are given in decoction (℥i), boiled in a pint of water for a quarter of an hour. This quantity to be taken in a day.

GALEOPSIS LADANUM, *G. grandiflora*—g. *Ochroleuca*, *G. grandiflora*—g. *Prostrata*, *G. grandiflora*—g. *Segetum*, *G. grandiflora*.

GALEOP'SIS VERSICOLOR, is possessed of the same virtues.

GALEOPSIS VILLOSA, *G. grandiflora*.

GALEROP'IA, *Oxyopia*.

GALEUX, *Psoric*.

GA'LIA. An ancient composition, in which galls were an ingredient; the *Galia pura*. There was also a *Galia aromatica*, *moscata* seu *musca'ta*, which consisted of a mixture of several perfumes, such as musk.

GANIANCON, *Galeancon*.

GALIOPSIS, *Galeopsis*.

GALIPEA FEBRIFUGA, *Cusparia febrifuga*—g. *Officialis*, see *Cusparia febrifuga*.

GALIPOT, see *Pinus sylvestris*.

GALIUM, *Gall'ium*, (from *γαλα*, 'milk,' because some species curdle milk.) *G. verum*.

GALIUM ALBUM, *G. Mollugo*.

GALIUM APARIN'E, *G. infestum* seu *aparinoi'des* seu *brachycarp'on* seu *scaber'rimum*, *Valant'ia aparin'e*, *Aparin'e*, *A. his'pida*, *Lappa*, *Philanthropus*, *Ampelocarp'us*, *Omphalocarp'us*, *Ilex*, *Asphari'nè*, *Aseper'ula*, *Goose-grass*, *Cleaver's bees*, *Cleavers*, *Goose-shave*, *Robin-run-the-hedge*, *Haygriff*, (Old Eng.) *Clitheren*, (Prov.) *Clider*, *Cliver*; Order, *Rubiaceæ*; Ser. *Syst.* *Tetrandria Monogynia*. (F.) *Gaillet cœcrochant*, *G. crochant*, *Grat-turon*. The expressed juice has been given as an aperient diuretic in incipient dropsies; also in cancer; and the decoction in *Lepra* and *Psoriasis*.

GALIUM APARANOIDES, *G. aparine*.

GALIUM ASPREL'UM, *Rough bed-straw*, *Rough*

caules' bed-straw; indigenous; has the diuretic properties of most of its genus.

GALUUM BRACHYCARPON, *G. aparine*—*G. Caucasicum*, *G. verum*.

GALUUM CIRCEAZANS, *Wild Liquorice*, *Master of the Woods*. An indigenous plant, which flowers from June to August. It is demulcent and diuretic, and is a popular domestic remedy.

GALUUM INFESTE, *G. aparine*—*G. Luteum*, *G. verum*.

GALUUM MOLLU'GO, *Galium album* seu *Tyrolense*, *Alys'sum Plin'ii*, *Greater ladies' bed-straw*, (*F.*) *Caillelait blanc*. The herb and flowers have been used medicinally in epilepsy.

GALUUM ODORATUM, *Asperula odorata*—*G. Scaberrimum*, *G. aparine*.

GALUUM TINCTORUM, an American species, closely allied in properties to *G. verum*.

GALUUM TYROLENSE, *G. mollugo*—*G. Tuberculatum*, *G. verum*.

GALUUM VERUM, *Galium*, *G. luteum* seu *Caucasicum* seu *tuberculatum*, *Ladies' bed-straw*, *Cheese-remet*, *Bed-straw*, *Cleavewort*, *Goose-grass*, *Savoyan*, *Clabber-grass*, *Milkweet*, *Poor Robin*, *Gravel-grass*, (*F.*) *Gaillet jaune* ou *vrai*, *Vrai Caillelait*. The tops were used in the cure of epilepsy. The leaves and flowers possess the property of curdling milk.

GALL, Bile, see *Chafing*, and *Eczema impetiginodes*, *Quercus infectoria*, and *Vitrum*—*G.* of the Earth, *Prenanthes*, *P. alba*—*G. Nut*, see *Quercus infectoria*—*G.* of the Ox, see *Bile*—*G.* of the Skin, *Chafing*—*G. Turkey*, see *Quercus infectoria*.

GALL-BLADDER, *Vesicula fellea*, *Cholecyst*, *Cholecystitis*, *Folliculus felleus*, *Cystis fellea* seu *choledochus*, *Vesicula fellea* seu *biliaria*, *Vesicula bilis*, *Folliculus felleus*, (*F.*) *Vésicule du fiel* ou *biliaire*, *Réservoir de la bile*. A membranous, pyriform reservoir, lodged in a superficial depression at the inferior surface of the right lobe of the liver. It receives, by the hepatic and cystic ducts, a portion of the bile secreted by the liver, when the stomach is empty, which becomes in it more acrid, bitter, and thick. It receives an artery, called the *cystic*. Its veins empty into the vena porta. Its nerves come from the hepatic plexus, and its lymphatic vessels join those of the liver.

GALLA, see *Quercus infectoria*—*G. Maxima Orbiculata*, see *Quercus infectoria*.

GALLÆ QUERCŪS, see *Quercus infectoria*—*G. Tinctoria*, see *Quercus infectoria*—*G. Turcica*, see *Quercus infectoria*.

GALLATURA, Molecule.

GALLÉ DE CHÊNE, see *Quercus infectoria*.

GALLI GALLINACEI CAPUT, *Gallinaginis caput*.

GALLINAG'INIS CAPUT, *Galli gallina'cei Caput*, *Caput gallina'ceum*, *Colliculus semina'lis*, *Caruncula semina'lis*, *Vernu montanum*, *Crista urethralis* seu *ure'thræ*, *Crête uréthrale*.—(*Ch.*) from *gallinago*, *gallinaginis*, 'a woodcock.' An oblong, rounded projection, formed by the mucous membrane in the spongy portion of the urethra, at the sides of which the ejaculatory ducts open.

GAL'LIPOT, (*Prov.*) *Glumpot*. Perhaps from *gala*, 'finery.' [?] A pot painted and glazed, or merely glazed, and commonly used to hold medicines.

GALLITRICHUM, *Salvia sclarea*.

GALLIUM, *Galium*.

GALLS, see *Quercus infectoria*—*G. Bussorah*, see *Quercus infectoria*—*G. Mecca*, see *Quercus infectoria*.

GALLSICKNESS, Fever, Walcheren.

GALLSTONES, Calculi, biliary.

GALLUS, Eunuch.

GALREDA, Gelatin.

GALVANIA VELLOZII, *Palicourea Maro-gravii*.

GALVANIC CAUTERY, see *Moxa*.

GALVANISATION, *Galvanization*.

GALVANISM, *Galvanismus*, *Electric'itas animalis* seu *Galvanica* seu *metall'ica*, *Irritamentum metallorum* seu *metall'icum*, *Voltaism*, *Volta'ic* or *Chemical* or *Contact Electricity*. Called after Galvani. A series of phenomena, consisting in sensible movements, executed by animal parts, which are endowed with irritability, when placed in connexion with two metallic plates of different nature, between which a communication is established by direct contact or by means of a metallic wire. Galvanism has been employed medicinally in the same cases as electricity, and especially in neuralgic affections. It is often applied in the form of plates,—"*Mansford's plates*." In asthma, for example, a small blister, the size of a dollar, may be placed on the neck over the course of the phrenic and pneumogastric nerves, and another on the side, in the region of the diaphragm. One metal is placed medially or immediately over the vesicated surface on the neck, and another over that in the side. They are then connected by means of a wire. The new nervous impression, in this way induced, is often signally beneficial.

GALVANIZATION, *Galvanisatio*, (*F.*) *Galvanisation*. The act of affecting with galvanism.

GALVANIZATION, *Lo'calized*. (*F.*) *Galvanisation localisée*. A mode of employing galvanism, proposed by Duchenne, of Boulogne, by which, he conceives, the electric influence may be arrested, at will, in the skin; or, without any incision or puncture, the skin may be traversed, and the electric influence be limited to the organs which it covers—to the nerves, muscles, and even the bones.

GALVANOPUNCTURE, *Electropuncture*.

GAMBA, *Patella*.

GAMBARUS, Crab.

GAMBIER, see *Catechu* and *Nauclea gambir*.

GAMBIR, see *Catechu* and *Nauclea gambir*.

GAMBOGIA, *Cambogia*.

GAMBOIDIA, *Cambogia*.

GAMMARUS, Crab.

GAM'MATA FERRAMEN'TA. *Cauteries*, having the shape of the Greek letter Γ , *gamma*, which were used for cauterizing hernie.

GAMMAUT, *Gammot*. The Italians, according to Scultetus, gave this name to a kind of crooked bistouri, used for opening abscesses.

GAMMISMUS, *Psammismus*.

GAMPHE, *Gena*.

GAMPHELE, *Gena*, *Maxillary Bone*.

GANGAME, *Epiploon*.

GANGAMUM, *Epiploon*.

GANGLIA CEREBRI POSTICA, *Thalami nervorum optico'rum*—*G. Formative*, see *Ganglion*—*G. Hemispherical*, *Hemispheres of the brain*—*G.* of Increase, see *Ganglion*—*G. Nervorum*, *Ganglions*, *nervous*, see *Ganglion*—*G. Respiratory*, see *Encephalon*—*G. Sensory*, see *Sensory ganglia*—*G. Stomato-gastric*, see *Encephalon*.

GANGLIAR, *Ganglionic*.

GANGLIATED, *Ganglionic*.

GANG'LIFORM, *Gang'lioform*, *Gangli'form'is*. Having the shape of a ganglion. From *ganglion*, and *forma*, 'form.'

GANGLIITIS, *Ganglionitis*.

GANGL'OLUM, *Diminutive of ganglion*. A small ganglion.

GANGLIO'MA. A tumour of the glands or of the lymphatic ganglions, and especially epithelioma of them.—*Nysten*.

GANG'LION, *Gang'lium*. In anatomy, a name generally given to a knot-like enlargement in the course of a nerve. It is applied, however, to organs differing considerably from each other in

size, colour, texture, functions, &c. They are divided into *glandiform*, *lymphatic*, and *nervous*.

1. *Glandiform ganglia*, called also *adenoid*, *vascular*, and *sanguineous ganglia*; *blind*, *aporic*, *ductless*, *blood*, and *vascular glands*; *glandula spuria*, &c., are organs of whose functions we are, in general, ignorant; and which have the appearance of glands. They are formed of agglomerated globules, pervaded by blood-vessels surrounded by areolar membrane, and contain a milky or yellowish fluid. To this class belong the spleen, thymus, thyroid, and supra-renal glands. They are, doubtless, concerned in sanguification.

2. *Lymphatic ganglia*. See *Conglobata*. 3. *Nervous ganglia*, *Ganglia seu Ganglio'nes seu Nodi seu Noduli Nervo'rum*, *Tumores seu Plexus ganglioform'es*, *Plexus glandifo'rm'es*, *Tuber'cula nodo'sa Nervo'rum*, *Diverticula spirituum animalium*, *Ganglia of increase*, *Form'ative ganglia*. Enlargements or knots in the course of a nerve. They belong, in general, to the system of the great sympathetic. One exists on the posterior root of every spinal nerve, and on one cerebral,—the 5th. Bichat regarded them as so many small brains, or centres of nervous action, independent of the encephalon, and intended exclusively for organic life. Being formed by the union of the cerebral and spinal nerves, they may send out the influence of both these nervous centres to the parts to which the nerves proceeding from them are distributed. Ganglia are chiefly composed of vesicular neurine; and appear to be concerned in the formation and dispensation of nerve power.

GANGLION, (*γαγγλιον*, *Ætius*.) Same etymon. *Emphy'ma encystis ganglion*. A globular, hard, indolent tumour, without change in the colour of the skin; of a size varying from that of a pen, to that of an egg, and always situate in the course of a tendon. The tumour is formed of a viscid, albuminous fluid, contained in a cyst of greater or less thickness. The cyst is sometimes loose; but in the majority of cases it communicates, by a narrow footstalk, with the sheath of a tendon, or even with the synovial capsule of a neighbouring articulation. The causes are generally unknown. The treatment consists in compression, percussion, the use of discutients, extirpation, or incision.

GANGLION ABDOMINALE, *G. semilunare*—*g. Adenoid*, *G. glandiform*—*g. of Andersch*, *Petrous ganglion*—*g. Annular*, see *Ciliary ligament*—*g. of Arnold*, *Oticum ganglion*—*g. Auricular*, *Oticum G.*—*g. Azygous*, see *Trisplanchnic nerve*—*g. Cardiac*, *Cardiac ganglion*—*g. Carotic* or *Carotid*, see *Carotid* or *Carotic nerve*—*g. Cavernous*, see *Carotid* or *Carotic nerve*—*g. Cerebelli*, *Corpus dentatum*—*g. Cerebral*, anterior, *Corpora striata*—*g. Cerebri Anterior*, *Corpora striata*—*g. Cerebri, inferior*, *grand*, *Thalami nervorum optico'rum*—*g. Cerebri, posterior*, *Thalami nervorum optico'rum*—*g. du Cervelet*, *Corpus dentatum*—*g. Ciliare*, *Corpus dentatum*—*g. Ciliary*, *Ophthalmic ganglion*—*g. Corpuscles*, see *Neurine*.

GANGLION OF EHRENRITTER, *Ganglion nervi glosso-pharynge'i superius*, *G. jugulare superius seu Ehrenritteri seu Mulleri*. A reddish-gray mass on the glosso-pharyngeal nerve in the foramen lacerum, above the ganglion of Andersch.

GANGLION OF THE FIFTH NERVE, *G. of Gasser*.

GANGLION OF GASSER, *Ganglion seu Ganglion Gass'eri seu Gasserii'nium seu semilunare*, *Moles gangliiform'is*, *Intumescen'cia gangliiform'is seu semilunaris*, *Ta'nia nervo'sa Hall'eri*, *Ganglion of the fifth nerve*. A semicircular knot on the 5th pair of nerves, before its division into three branches.

GANGLION, GLANDIFORM, see *Ganglion*—*G. Globules*, see *Neurine*—*g. Impar*, see *Trisplanchnic*

nerve—*g. Jugulare superius*, *G. of Ehrenritter*—*g. Latteur*, *Milk-knot*—*g. of Laumonier*, see *Carotid* or *Carotic nerve*—*g. Lenticular*, *Ophthalmic G.*—*g. Lymphatic*, *Conglobate gland*—*g. Maxillo-tympanique*, *Oticum G.*—*g. of Meckel*, *Sphenopalatine G.*—*g. Mulleri*, *G. of Ehrenritter*—*g. Nasopalatine*, see *Nasopalatine ganglion*—*g. Nervi glosso-pharyngei superius*, *G. of Ehrenritter*—*g. Nervous*, *Neuroma*—*g. Ophthalmic*, see *Ophthalmic ganglion*—*g. Optic*, *Quadrigenina tubercula*—*g. Orbital*, *G. ophthalmic*—*g. Oticum*, *Oticum G.*—*g. Petrosal*, see *Petrous ganglion*.

GANGLION OF THE PNEUMOGASTRIC. A ganglionic structure in the pneumogastric as it passes through the foramen lacerum posterius.

GANGLION OF RIBES. A nervous ganglion upon the anterior communicating artery of the brain, and to be found at the point of junction of the right and left trunks of the sympathetic.

GANGLION, SANGUINEOUS, *G. glandiform*—*g. Semilunare*, *G. of Gasser*, *Ophthalmic ganglion*—*g. Sensory*, see *Sensory ganglia*—*g. Solare*, *G. semilunare*—*g. Sphenoidal*, *Sphenopalatine ganglion*—*g. Spinal*, see *Encephalon*—*g. Splanchnicum*, *G. semilunare*—*g. Superieur du cerveau (grand)*, *Corpora striata*—*g. of the Superior Laryngeal Branch*, see *Pneumo-gastric nerves*—*g. Surre'nal*, *G. semilunare*—*g. Thyroid*, see *Trisplanchnic nerve*—*g. Transversum*, *G. semilunare*—*g. Vascular*, *G. glandiform*—*g. Vertebral*, see *Trisplanchnic nerve*—*g. de Vieussens*, *Cœliac plexus*.

GANGLIONARY, *Ganglionic*.

GANGLIONES NERVORUM, *Ganglions*, *nervous*.

GANGLIONIC, *Ganglion'icous*, *Gan'gionary*, *Gan'gliar*, *Gan'gliated*, from *ganglion*, (*F.*) *ganglionnaire*. Relating to ganglia. Nerves are so called in the course of which ganglions are met with; as the greater part of the branches of the great sympathetic or trisplanchnic, the posterior roots of the spinal nerves, &c. *Ganglionics*, according to Dr. Pereira, are agents, which affect the ganglionic or great sympathetic system of nerves, [?]—as stimulants and sedatives.

GANGLIONIC NERVE, *Trisplanchnic nerve*—*g. Nervous System*, see *Trisplanchnic nerve*.

GANGLIONITIS, *Ganglii'tis*, from *γαγγλιον*, 'a ganglion,' and *itis*, denoting inflammation. Inflammation of a nervous ganglion. Sometimes used for inflammation of a lymphatic ganglion.

GANGLIONITIS PERIPHERICA ET MEDULLARIS, *Cholera*.

GANGLIONNAIRE, *Ganglionic*.

GANGLIUM, *Ganglion*—*g. Gasseri*, *Ganglion of Gasser*.

GANGRÆNA, *Gangrene*—*g. Alopecia*, *Alopecia*—*g. Caries*, *Caries*—*g. Nosocomiorum*, *Hospital gangrene*—*g. Oris*, *Cancer aquaticus*—*g. Ossis*, *Spina ventosa*—*g. Ossium*, *Caries*—*g. Pottii*, see *Gangrene*, and *Gangrene of old people*—*g. Pulmonum*, *Necropneumonia*—*g. Senilis*, *Gangrene of old people*—*g. Sphacelus*, *Sphacelus*—*g. Tonsillarum*, *Cynanche maligna*—*g. Ustilaginea*, *Ergotism*—*g. Vagine*, *Colpocœce*.

GANGRENESENTIA, *Gangranosis*.

GANGRENICUS, *Gangrenous*.

GANGRENODES, *Gangrenous*.

GANGRENOPTIS, from *γαγγραινα*, 'gangrene,' and *οτις*, 'the countenance.' *Cancer aquaticus*; also, gangrenous inflammation of the eyelids; *Blepharitis gangranosa*.—Siebenhaar.

GANGRENO'SIS, *Gangrenescen'cia*, from *γαγγραινα*, 'gangrene.' The state of being gangrenous or of becoming gangrenous.

GANGRENOSUS, *Gangrenous*.

GANGRENE, *Gangræ'na*, *Cancere'no*, *Hot mortification*, (*F.*) *Gangrène*, *G. Chaude*, *Asphyxie des parties*, from *γρᾶν*, *γρᾶν*, 'I eat.' Privation of

life or partial death of an organ. Authors have generally distinguished mortification into two stages: naming the first *incipient* or *gangrene*. It is attended with a sudden diminution of feeling in the part affected; livid discoloration; detachment of the cuticle, under which a turbid fluid is effused; with crepitation, owing to the disengagement of air into the areolar texture. When the part has become quite black, and incapable of all feeling, circulation, and life, it constitutes the *second stage*, or *mortification*, and is called *sphac'elus*. Gangrene, however, is frequently used synonymously with mortification.—*local asphyxia* being the term employed for that condition, in which the parts are in a state of suspended animation, and, consequently, susceptible of resuscitation. When the part is filled with fluid entering into putrefaction, the affection is called *humid gangrene*, (F.) *Gangrène humide*: on the other hand, when it is dry and shrivelled, it constitutes *dry gangrene*; (F.) *Gangrène sèche*. To this class belongs the *gangræ'na seni'lis*, *G. Pottii*, *Presbyosphac'elus*, or *spontaneous gangrene* of old people, which rarely admits of cure. Whatever may be the kind of gangrene, it may be caused by violent inflammation, contusion, a burn, congelation, the ligature of a large arterial trunk, or by some inappreciable internal cause.

The treatment, both of external and internal gangrene, varies according to the causes which produce it. Gangrene from excessive inflammation is obviated by antiphlogistics; and that from intense cold by cautiously restoring the circulation by cold frictions, &c. When the gangrene has become developed, the separation of the eschars must be encouraged by emollient applications, if there be considerable reaction; or by tonics and stimulants, if the reaction be insufficient.

GANGRENE, HOSPITAL, see Hospital gangrene—*g.* of the Lungs, Necropneumonia.

GANGRÈNE DE LA BOUCHE, Cancer aquaticus—*g.* *Chande*, Gangrene—*g.* *Froide*, Sphacelus—*g.* *Humide*, see Gangrene—*g.* *Sèche*, see Gangrene—*g.* *Hôpital*, Hospital gangrene—*g.* *du Panon*, Necropneumonia—*g.* *des Solonois*, Ergotism.

GANGRENOUS, *Gangræ'nicus*, *Gangræno'sus*, *Gangræno'des*, (F.) *Gangrèneux*. Affected with or relating to gangrene.

GANGRENOUS' EROSION OF THE CHEEK, Cancer aquatics.

GANJAH, Gunjah.

GANTELET (F.), *Chirothe'ca*, *Fas'cia digi'talis*, *Gauntlet*; from (F.) *gant*, 'a glove.' A sort of bandage which envelops the hand and fingers like a glove. It is made with a long roller, about an inch broad: and is applied so that the fingers are covered to their tips, when it is called *Gantelet entier ou complet*. The *Demi-gantelet* includes only the hand and base of the fingers. Both bandages are used in fractures and luxations of the fingers, burns of the hand, &c. See Chirotheca.

GANTS DES DAMES, Condom—*g.* *de Notre Dame*, Digitalis.

GAOL CACHEXIA, see Cachexia, gaol.

GAPING, Yawning.

GARANÇE, Rubia.

GARCIN'IA CAMBO'GIA, *G.* seu *Cambogia gutta*, *Mangost'a'na Cambogia*. *Ord.* Guttifera. A tree of Ceylon, *Family*, Guttifera, which affords a concrete juice similar to Gamboge.

GARCINIA GUTTA, *G.* *Cambogia*.

GARCIN'IA MANGOSTA'NA, *Mangost'a'na Garcin'ia*, from *Garcin*, who accurately described it. The systematic name of the *Mangostan'*, *Mangoustan'*, *Mangostine'* or *Mangosteen' tree*, *Man-*

gosta'na. It grows in great abundance in Java and the Molucca islands. The fruit, which is about the size of an orange, is delicious, and is eaten in almost every disorder. The dried bark is used medicinally in dysentery and tenesmus; and a strong decoction has been much esteemed in ulcerated sore throat.

GARCIN'IA PURPUREA. From this species a concrete oil of *Mangosteen* is obtained, which is called, in India, *Kokum butter*, *Kokum* being the Indian name of *Mangosteen*. The oil is extracted from the fruit by boiling. It is used in India in cases of chaps.

GARDE-MALADE, Nurse.

GARDENIA DUMETORUM, *Randia dumetorum*—*g.* *Spinosa*, *G.* *dumetorum*.

GARDEROBE, *Artemisia abrotanum*.

GARDEROBES, Faeces.

GARDINER'S ALIMENTARY PREPARATION, see Oryza.

GARDOUCHES, *Vesiculae seminales*.

GARETUM, Poples.

GARGALISMUS, Gargalus.

GARGALUS, *Gar'gale*, *Gargalis'mus*, *Titilla'tio*, *Irrita'tio*, *Pruri'tus*. Titillation, irritation, itching. Also, masturbation; and, rarely, animal magnetism.

GARGAREON, Uvula.

GARGARISATIO, Gargarism.

GARGARISM, *Gargaris'mus*, *Gargaris'ma*, *Anagargalic'ton*, *Gargaris'mum*, *Collu'tio*, *Dia-cly's'ma*, *Anagargaris'ton*, *Anagargaris'mus*, *Ana-conchylis'mus*, *Ana-conchylis'mus*, *Titillamen'tum*, from γαργαρίζω, 'I wash the mouth,' 'I gargle.' A gargle. Any liquid medicine, intended to be retained in the mouth, for a certain time, and to be thrown in contact with the uvula, velum pendulum, tonsils, &c. For this purpose, the liquid is agitated by the air issuing from the larynx, the head being thrown back. Gargles are employed in cynanche tonsillar and other diseases of the fauces, and are made of stimulants, sedatives, astringents, refrigerants, &c., according to circumstances.

The process is termed *gargling*, *gargarisa'tio*.

The term *collu'torium* or *collutatorium oris* is generally restricted to a wash for the mouth.

GARGET, *Phytolacca decandra*.

GARGLE, Gargarism.

GARGLING, see Gargarism.

GARGOUILLEMENT, Borborygmus, Gurgling. See Râle muqueux.

GARLIC, *Allium*—*g.* *Hedge*, *Alliaria*—*g.* *Meadow*, *Allium Canadense*.

GARLOCK, *Sinapis arvensis*.

GAROSMUM, *Chenopodium vulvaria*.

GAROU BARK, *Daphne gnidium*.

GARRETUM, Poples.

GARROPHYLUS, *Eugenia caryophyllata*.

GARROT (F.), *garrot*, from *garrotter*, 'to tie fast.' A compressing bandage, tightened by twisting a small cylinder of wood, by which the arteries of a limb are compressed, for the purpose of suspending the flow of blood in cases of hemorrhage, aneurism, amputation, &c.

GARROTILLO, *Cynanche maligna*.

GARRULITAS, Loquacity.

GARULEUM BIPINNATUM. A South African plant, *Ord.* Compositæ; known under the name *Snakeroot*, from its reputed effects as an antidote to the bites of venomous serpents. The root is a great favourite with the Boers in chest diseases—as asthma—and in affections in which a free secretion from the mucous membrane of the bronchia is indicated. It has diaphoretic properties, and acts as a diuretic in gout and dropsy. It is given in decoction or tincture. Dr. Pappe thinks the root ought to have a place in the *Materia Medica*.

GARUM. γαρων, *Licul'men, Liqua'men.* The ancient Romans gave this name to a kind of pickle made by collecting the liquor which flowed from salted and half-putrefied fish. It was used as a condiment. — *The Geoponics*, Humelberg on Apicius, Martial, &c.

GARVANCE, Cicer arietinum.

GAS. see Gaz — g. Ammoniacale, Ammonia — g. Animale sanguinis, Gaz sanguinis — g. Azoticum, Azote — g. Azoticum oxygenatum, Nitrogen, gaseous oxide of — g. Hepaticum, Hydrogen, sulphuretted — g. Hydrogenium sulphuretum, Hydrogen, sulphuretted — g. Intoxicating, Nitrogen, gaseous oxide of — g. Laughing, Nitrogen, gaseous oxide of — g. of the Lungs, Gaz, pulmonary — g. Nitrous, dephlogisticated, Nitrogen, gaseous oxide of — g. Oxygenated muriatic acid, Chlorine — g. Oxymuriatic acid, Chlorine — g. Palustre, Miasm, marsh — g. Paradise, Nitrogen, gaseous oxide of — g. Sulphuris, Sulphurous acid.

GASCARILLA, Croton cascarilla.

GASPING, Anhelatio.

GASTEIN, MINERAL WATERS OF. Gastein or Gasteiner Wildbad is in the Noric Alps, Austria. The waters are thermal. Temp. 106° to 118° Fah. They contain sulphate of soda, chloride of sodium, chloride of potassium, carbonate of soda, carbonate of lime, magnesia, manganese, iron, &c.

GASTER, γαστρ. The abdomen. At times, but rarely, the uterus. Also, the stomach, in particular; Hence:

GASTERALGIA, Gastralgia.

GASTERANAX, from γαστρ, 'the abdomen,' and ἀναξ, 'a king.' A name given by Doletus to a hypothetical vital principle, corresponding to the Archæus of Van Helmont, the sent of which he placed in the lower belly. See Bithnimalca.

GASTERANGEMPHRAXIS, Gasterangiemphraxis.

GASTERANGIEMPHRAXIS, *Gasterangiemphrax'is, Gasteremphrax'is.* Obstruction of the pylorus; from γαστρ, 'stomach,' ἀγγω, 'I strangle,' and ἐμφρασσω, 'I obstruct.' — Vogel. Also, and properly, obstruction or congestion of the vessels of the stomach.

GASTÉRISE, Pepsin.

GASTERASTHENIA, *Imbecill'itas seu Aton'ia seu Lax'itas ventriculi;* from γαστρ, 'stomach,' and ἀσθενεία, 'debility.' Debility of the stomach.

GASTERATAXIA, Embarras gastrique.

GASTERECHÉMA, *Gastreche'ma, Son'itus stomach'icus;* from γαστρ, 'the stomach,' and ἤχημα, 'sound.' Sound heard on auscultating or percussing the region of the stomach, and referred to that organ.

GASTEREMPHRAXIS, Gasterangiemphraxis.

GASTERHYSTEROTOMY, Cæsarean section.

GASTRÉMIA, from γαστρ, 'the stomach,' and αἷμα, 'blood.' Hyperæmia of the stomach.

GASTRALGIA, Cardialgia.

GASTRANEURYSMA, Gastrectasis.

GASTRATROPHIA, from γαστρ, 'the stomach,' and τροφία, 'wasting.' Atrophy of the stomach.

GASTRECHEMA, Gastrechema.

GASTRECTASIS, *Gastrecta'sia, Gastroneury'sma, Dilata'tio ventriculi;* from γαστρ, 'the stomach,' and εκτασις, 'dilatation.' Dilatation of the stomach.

GASTRELCOBROSIS, Gastrobrosis ulceroza.

GASTRELCO'SIS, *Ulcera'tio seu Eru'deera'tio seu Ul'cera ventriculi;* from γαστρ, 'the stomach,' and ἐλκος, 'ulceration.' Ulceration of the stomach. When accompanied with hectic, it constitutes *Gastrophth'isis, Gastroph'thoë.*

GASTRELYTROTOMIA, see Cæsarean Section.

GASTRENCEPHALOMA, *Gastromyelo'ma, Gastromyelo'sis, Gastroencephalo'sis;* from γαστρ, 'the stomach,' and ἐγκεφαλος, 'the brain.' Encephaloid of the stomach.

GASTRENCHYTA, Stomach pump.

GASTRENTERIC, Gastroentericus.

GASTRENTERTIC, Gastroenteritic.

GASTRENTEROMALACIA, *Gastroenteromalac'ia,* from γαστρ, 'stomach,' εντερον 'intestine,' and μαλακία, 'softening.' Softening of the stomach and intestines.

GASTREPATICUS, Gastrohepatic.

GASTREPATITIS, *Inflamma'tio ventriculi et hep'atis;* from γαστρ, 'stomach,' and ἥπαρ, 'liver.' Inflammation of the stomach and liver.

GASTREPIPOLOICUS, Gastroepiploic.

GASTRETHIIS'IA, from γαστρ, 'stomach,' and ἐρεθίζω, 'I irritate.' Irritation of the stomach.

GASTRIC, Gastricus; from γαστρ, 'the stomach.' Belonging or relating to the stomach.

GASTRIC ARTERIES are three in number, *Arteria gastro-epiplo'ica dextra, A. gastro-epiplo'ica sinistra,* and *A. coronaria ventriculi.*

GASTRIC JUICE, Succus gastricus, Men'strum seu Fermentum Ventriculi, (F.) Suc Gastrigue, Gastric Acid. A fluid, secreted from the mucous membrane of the stomach. As met with, it is a mixture of the fluids secreted by that organ with those of the supra-diaphragmatic portion of the alimentary canal. Owing to such admixture, the most contrary properties have been assigned to it. That such a fluid is secreted, which concurs powerfully in digestion, is evident from many considerations, and has been positively proved by the author and numerous others. It was found by him to contain, in man chlorohydric and acetic acids. The gastric fluid in cases of sudden death sometimes corrodes and perforates the stomach; giving rise to interesting questions in medical jurisprudence.

GASTRIC NERVES. The two cords by which the pneumogastric nerves terminate, and which descend on the two surfaces of the stomach; as well as the filaments of the great sympathetic, which accompany the gastric vessels.

GASTRIC PLEXUS, Plexus coronarius ventriculi. A nervous net-work, formed by the solar plexus. It accompanies the *Arteria coronaria ventriculi*, and passes along the lesser curvature of the stomach, to which it gives branches.

GASTRIC VEINS follow the same distribution as the arteries, and open into the *Vena porta abdominalis.*

GASTRICISM, Gastricis'mus, from γαστρ, 'the stomach.' A name by which is designated the medical theory, that refers all, or almost all, diseases to an accumulation of saburæ in the digestive passages.

GASTRICITÉ, Embarras gastrique.

GASTRILOQUE, Engastrimyth.

GASTRILOQUIST, Engastrimyth.

GASTRILOQUUS, Engastrimyth.

GASTRIMARGUS, Glutton.

GASTRINUM, Potash.

GASTRISMUS, Gluttony, see Saburra.

GASTRITIC, Gastrit'icus; same etymon as the rest. Relating to gastritis.

GASTRITIS, from γαστρ, 'the stomach,' and itis, denoting inflammation. *Ventriculi inflammatio, Causa seu Empres'ma seu Cardial'gia inflammato'ria, Febris stomach'ica inflammato'ria, Inflammat'io gastritis, I. ventriculi seu stom'achi, Phlegmonè ventriculi, Inflammation of the stomach, (F.) Inflammation de l'estomac, Gastrite, Catarrhe gastrique.* A disease, characterized by pyrexia; great anxiety; heat and pain in the epigastrium, increased by taking any thing into the stomach; vomiting and hiccup. Gastritis may either be seated in the peritoneal or mucous

coat. It is most frequently in the latter—*Eso-gastritis*, *Endogastritis*, *Gastromycoderitis*, *Gastritis mucosa*—being excited directly by acrid ingesta. It requires the most active treatment:—bleeding, blistering, fomentations, diluents, &c. Some degree of inflammation of the mucous coat of the stomach was considered by the followers of Broussais to be present in almost all fevers; and the various forms of dyspepsia have been supposed by some to be nothing more than chronic *endogastritis*.

GASTRITIS ARTHRITICA, *Cœliagra*.

GASTRO, from *γαστρον*, 'stomach.' In composition, relating or appertaining to the stomach, or belly.

GASTRO-ARTHRITIS, Gout—*g.* Ataxia, Dyspepsia—*g.* Atonia, Dyspepsia—*g.* Bronchitis, Fever, adenomeningeal—*g.* *Entérite intense*, Synocha—*g.* *Entérite*, with nervous affection of the brain, see Typhus—*g.* Enteritis, follicular, Dotheritis; see Typhus.

GASTROBROSIS, *Perforatio ventriculi*, *Gastrostomachia*, (F.) *Perforation de l'Estomac*; from *gastro*, and *βρωσις*, 'the act of gnawing,' Corrosion and perforation of the stomach.

GASTROBROSIS ULCEROSA, *Gastrocœbro'sis*. Destruction and perforation of the coats of the stomach by ulceration.

GASTROCELE, from *gastro*, and *κῆλη*, 'a tumour,' *Hernia of the stomach*, *Hernia ventriculi*, (F.) *Hernie de l'Estomac*. *Hernia* formed by the stomach through the upper part of the linea alba; a disease, the existence of which has been doubted by many. See *Epigastrocele*.

GASTRO-CEPHALITIS: from *gastro*, *κεφαλή*, 'head,' and *itis*, denoting inflammation. Inflammation of the stomach and head,—a not uncommon concomitant of certain malignant fevers.

GASTROCHOLOSIS, Fever, gastric.

GASTROCNEME, Sura.

GASTROCNEMIA, Sura.

GASTROCNEMII, from *gastro*, and *κνήμη*, 'the leg.' The name of the two fleshy masses which occupy the posterior and superficial part of the leg, *Genelli*: (F.) *Gastrocnémien*, *Summeux de la jambe*, *Bifemoro-calcaneus*—(Ch.) the two constituting the *Gastrocnemius externus*, of English anatomists. These muscles are distinguished into *internal* and *external*, which are distinct above, but united at their inferior extremity. They are long, flat, and thick; and arise—the former from the posterior part of the outer condyle of the femur; the latter, from the posterior part of the inner condyle of the same bone. The aponeurosis, which unites these muscles below, joins with that of the solaris, and forms with it, a large tendon, which, under the name *Tendo Achillis*, is inserted at the posterior part of the calcaneum. These muscles extend the foot on the leg, and the leg on the foot. They can, also, bend the leg and the thigh reciprocally on each other. For the *Gastrocnemius internus*, see *Soleus*.

GASTROCNEMIUM, Sura.

GASTROCELIACUS, *Cœliac*.

GASTROCELICUS, *Cœliac*.

GASTROCOLIC, see *Epiploon*, *gastrocolic*.

GASTROCOLICA, *Cardialgia*.

GASTROCOLITIS, from *gastro*, and *κώλον*, 'colon.' Inflammation of the stomach and colon. *GASTROCOLPOTOMIA*, see *Cæsarean* section.

GASTRODIDYMUS, *Gastrodyms*, *Didymus*, *Symphogastrius*, *Pseudymus*; from *gastro* and *δίδυμος*, 'a twin.' A monstrosity in which twins are united by the abdomen.

GASTRODUODENAL, *Gastroduodenalis*; from *gastro*, and *duodenum*. Relating to the stomach and duodenum.

GASTRO-DUODENITIS, (F.) *Gastro-duodénite*. Inflammation of the stomach and bowels, an affection, which figured largely in the system of Broussais. See *Gastro-enteritis*.

GASTRODUODENOPYRA, Fever, adenomeningeal.

GASTRODYNE, *Cardialgia*.

GASTRODYNIA, *Cardialgia*—*g.* *Flatulenta*, *Colica flatulenta*.

GASTRO-ELYTROTOMIA, see *Cæsarean* section.

GASTRO-ENTERALGIA, from *gastro*, *εντερον*, 'intestine,' and *αλγος*, 'pain.' Pain in the stomach and intestines.

GASTROENTERIC, *Gastroentericus*, *Gastroenteric*, *Gastroentericus*, *Gastro-intestinal*, from *gastro*, and *εντερον*, 'intestine.' Relating to the stomach and intestine.

GASTROENTERITE, *Gastroenteritis*.

GASTROENTERITIC, *Gastroenteritic*, *Gastroenteritis*, *Gastroenteritic*, *Gastroenteritic*; same etymon as the next. Relating to *gastroenteritis*.

GASTROENTERITIS, *Inflammatio ventriculi et Intestini*, (F.) *Gastro-entérite*, *Gastro-enteric disease*, from *gastro*, *εντερον*, 'an intestine,' and *itis*, a suffix denoting inflammation. Inflammation of the stomach and small intestine. According to Broussais, the essential fevers of authors are *gastro-enteritis*, simple, or complicated.

GASTROEPIPLOIC, *Gastro-epiploicus*, *Gastroepiploicus*, from *gastro* and *επιπλοον*, 'the epiploon.' That which relates to the stomach and epiploon.

GASTROEPIPLOIC ARTERIES, *Gastric inferior arteries*, are two in number, and distinguished into *right* and *left*. The *right*, also called *Gastro-hepatic*, *Gastrica inferior dextra*, *Gastro-epiploica dextra*, is furnished by the hepatic artery. It descends behind the pylorus, and passes from right to left, along the great curvature of the stomach. It gives branches to the pancreas, duodenum, stomach, omentum majus, and terminates by anastomosing with the *Gastro-epiploica sinistra*, *Gastrica sinistra seu inferior sinistra*. This—the *left*—arises from the splenic artery. It is of considerable magnitude, and passes from left to right, along the great curvature of the stomach, distributing its branches more particularly to the stomach and omentum majus. It terminates by joining the right *gastro-epiploic*.

GASTROEPIPLOIC GLANDS are the lymphatic ganglions or glands, situate towards the great curvature of the stomach, between the two anterior laminae of the omentum majus.

GASTROEPIPLOIC VEINS are distinguished, like the arteries, into *right* and *left*. They empty themselves:—the former, into the superior mesenteric; the latter, into the splenic vein.

GASTROHEMORRHAGIA, *Hæmatemesis*.

GASTROHEPATIC, *Gastro-hepaticus*, *Hepato-gastricus*, *Gastrohepaticus*; from *gastro*, and *ήπαρ, ηπατος*, 'the liver.' Relating to the stomach and liver. This name has been given to several organs. See *Epiploon*, *gastrohepatic*, &c.

GASTRO-HEPATITIS, *Simultaneous inflammation of the stomach and liver*.

GASTROHYSTEROTOMY, *Cæsarean* section.

GASTRO-INTESTINAL, *Gastro-enteric*.

GASTROLIENALIS, *Gastro-splenicus*.

GASTROLITHIASIS, from *gastro*, and *λίθαις*, 'formation of stone.' The formation of concretions, *gastrolithi*, in the stomach.

GASTROMALACIA, *Gastromalaxia*.

GASTROMALAXIA, *Gastro-malacia*, *Gastromalacia*, *Malacogaster*, *Malacia ventriculi*, *Dissolutio ventriculi*, *Emollitio ventriculi*, *Pseudophlogosis ventriculi* *resolutiva et colli-*

quat'ra, *Metamorphosis ventriculi gelatinifor-mia*, *Erosio et perforatio spontanea ventriculi*, *Resolutio et diabrosis ventriculi*, (F.) *Ramollissement de l'Estomac*, from *gastro*, and *μαλαξίς*, 'softening.' Softening of the stomach, induced at times by the gastric secretions after death,—*Resolutio ventriculi autopeptica*.

GASTROMANTIS, Engastrium.

GASTROMELES, from *gastro*, and *μελος*, 'a limb.' A monster with one or more supernumerary limbs on the abdomen.

GASTROME'NIA, from *gastro*, and *μην*, 'a month.' Vicarious menstruation from the stomach.

GASTRO-MENINGITIS, Fever, gastric.

GASTROMETROTOMIA, Cæsarean section.

GASTROMYCODERIS, see Stomach.

GASTROMYCODERITIS, see Gastritis.

GASTROMYELOMA, Gastroencephaloma.

GASTRONOSUS, Gastropathy.

GASTROPARALYSIS, *Gastroplegia*, *Paralysis ventriculi*; from *gastro*, and *παράλυσις*, 'paralysis.' Paralysis of the stomach.

GASTROP'ATHY, *Gastropathia*, *Gastron'-osis*, *Gastron'-us*, from *gastro*, and *πάθος*, 'disease.' Disease of the stomach.

GASTROPERIODYN'IA; *Sool* (India.) A violent periodical neuralgic pain at the pit of the stomach, not uncommon in Hindoostan.

GASTROPHREN'IC, *Gastro-phren'icus*, from *gastro*, and *φρενής*, 'the diaphragm.' Belonging to the stomach and diaphragm.

GASTROPHRENIC LIGAMENT is a reflection of the peritoneum, which descends from the inferior surface of the diaphragm to the cardia.

GASTROPHTHISIS, Gastrectosis.

GASTROPHTHOE, Gastrectosis.

GASTROPLEGIA, Gastroparalysis.

GASTROPYRA, Fever, gastric.

GASTROPYRETUS, Fever, gastric.

GASTROR'APHY, *Gastrorrhaphia*, *Gastror'-rhaphe*, *Sutura abdominalis*, from *gastro*, and *ραφή*, 'a suture.' The suture used for uniting wounds penetrating the abdomen, when they are too extensive or too unequal to be kept in contact by position, adhesive plaster, or appropriate bandages. The interrupted and quilted sutures are those chiefly employed.

GASTRORRHAGIA, Hæmatemesis.

GASTRORRHÆXIS, Gastrobrosis.

GASTRORRHÆA, from *gastro*, and *ρρω*, 'I flow.' *Blenorrhœa seu Fluxus ventriculi*, (F.) *Flux muqueux de l'estomac*, *Catarrh stomachi*. A morbid condition of the stomach, which consists in the secretion of an excessive quantity of mucus from the lining membrane. Also, Coeliac flux.

GASTROSCIR'RHUS, *Induratio ventriculi scirrhosa*, *Scirrhus seu carcinoma ventriculi*. Scirrhus induration or cancer of the stomach.

GASTROSCOPIA, Abdominoscopy.

GASTRO'SES. A generic name for diseases which are seated in the stomach.—*Alibert*.

GASTROSPLE'NIC, *Gastrosplicus*, *Gastro-liena'lis*, from *gastro*, and *σπλην*, 'the spleen.' Relating to stomach and spleen.

GASTROSTENOSIS, Stricture of the stomach.

GASTROSTENOSIS CARDIACA et **PYLORICA**; from *gastro*, and *στενός*, 'narrow.' Narrowness of the cardiac and pyloric orifices of the stomach from cancer of that organ.

GASTROT'OMY, from *gastro*, and *τομή*, 'incision.' Several different operations have been so called. 1. The Cæsarean Section. 2. An incision made into the abdomen for the purpose of removing some internal strangulation or volvulus, or to reduce hernia, *Laparotomy*: and, 3. The opening made in the stomach, to remove a

foreign body which has passed into it through the œsophagus.

GASTRYPERNEUR'IA, from *gastro*, *ὑπερ*, 'above,' and *νεῦρον*, 'a nerve.' Morbidly increased activity of the nerves of the stomach.

GÂTEAU FÉBRILE, Ague cake.

GÂTEUX (F.), from *gâter*, 'to spoil.' A name given, in the insane hospitals, and in the hospitals for incurables and the aged, in France, to persons who are incapable of attending to them selves, and have lost all sense of cleanliness.

GATHERING, Abscess, Suppuration.

GATTILIER, Vitex.

GAUDIA FEDA, Masturbation.

GAULTHER'IA, *Gaultheria seu Gualtheria procumbens*, *Gaultheria repens*, Mountain Tea, Partridge Berry, Berried Tea, Grouseberry, Deerberry, Spice berry, Tea berry, Red berry, Wintergreen, Red berry Tea, Ground berry, Ground ivy, Ground holly, Hill berry, Box berry, Chequer berry, (F.) *Gaulthérie*, *Palommier*, *Thé du Canada*, called after Dr. Gauthier, of Quebec. Ord. Ericaceæ. An American plant, which is one of the principal articles of the materia medica of some Indian tribes. The infusion of the leaves is stimulant and anodyne, and is said to have been used, with advantage, in asthma. The oil—*Oleum Gaultheriæ*, Ph. U. S.,—is used, chiefly on account of its pleasant flavour, to cover the taste of other medicines.

GAUNTLET, *Gantelet*.

GAQUAVA, Smilax China.

GAUTIERA REPENS, *Gaultheria*.

GAY FEATHER, *Liatriis spicata*.

GA'YAC, *Guaiacum*.

GAYLUSSAC'IA RESINO'SA, *Vaccinium resinosum*, *Black Huckleberry*. Order, Ericaceæ, (called after the chemist, *Gay-Lussac*.) An indigenous plant, whose fruit is sweet and agreeable. The leaves and root, as well as those of *GAYLUSSACIA DUMO'SA*, *Dwarf huckleberry*, and of *G. FRONDO'SA*, *Blue tangle*, indigenous, are used in decoction as astringents in diarrhœa, sore throat, &c.

GAZ, *Gas*, *Air*. Van Helmont first designated by this name,—the etymology of which is unknown,—the carbonic acid developed in the vinous fermentation. Afterwards, the term was appropriated to every permanently elastic fluid; that is, which preserves its æriform state at all temperatures; and ultimately it was extended to all æriform bodies;—which were divided into permanent and non-permanent gases. The latter are generally termed vapours:—they return to the liquid state, when a portion of their caloric is abstracted. The permanent gases, or *gazes* properly so called, are numerous, and may be divided into three sections with regard to their effects on the animal economy.

- | | |
|----------------------------------|---|
| 1. Irrespirable gases. | { Carbonic acid, ammoniacal gas, chlorohydric acid gas, deutoxide of azote, nitrous acid gas, and chlorine. |
| 2. Negatively deleterious gases. | |
| 3. Positively deleterious gases. | |
- Hydrogen, azote.
Oxygen, protoxide of azote, carburetted hydrogen, carbonic oxide, sulphuretted hydrogen, and arseniuretted hydrogen.

It is proper to remark that the term *respirable* has been very differently employed by different writers. Sometimes it has meant the power of supporting life when applied to the blood in the lungs. At others, all gases have been deemed irrespirable, which are incapable of being intro-

duced into the lungs by voluntary efforts,—without any relation to their power of maintaining vitality; and this is perhaps the best sense. The gazes were, at one time, employed in medicine, under great expectations,—especially by the enthusiastic Beddoes; but they are now scarcely ever had recourse to. They differ, considerably, in their effects on the animal economy. Some, as oxygen, are exciting; others, as azote, depressing; whilst others, again, as the *Protoxide of azote* or *laughing gas*, produce the most singular effects.

GAZ DES BALLONS, Hydrogen.

GAZ, PULMONARY, *Gaz of the lungs*. A name given to the expired air; which contains—besides common air, an increase of carbonic acid, water, and some animal matter.

GAZ SANGUINIS, *Gaz anima'le san'guinis*, *Halitus seu Aura seu Vapor san'guinis*, *Spiritus roriferus*, *Hæmatismus*. The halitus, or vapour, given off by freshly-drawn blood.

GAZEL, *Ribes nigrum*.

GAZELLE, Antelope.

GÉANT, Giant.

GÉANTISME, see Giant.

GEBARPULVER, Ergot.

GEDE'OLA. The convex part of the liver.—Du Cange.

GEILNAU, MINERAL WATERS OF. Geilnau is a village in the grand duchy of Nassau, at no great distance from Frankfort. The waters contain carbonic acid, carbonate, sulphate, and phosphate of soda, chloride of sodium, carbonate of lime, magnesia, and iron.

GEISMAR, MINERAL WATERS OF. The mineral waters of Geismar, in Bavaria, are acidulous chalybeates.

GEISUM, γεῖσον, *Geison*. The part of the frontal bone over the eyes.

GELAPPIUM, *Convolvulus jalapa*.

GELASINI DENTES, Incisive teeth.

GELASMUS, Canine laugh.

GELATIG'ENOUS. A hybrid term, from *gelatin*, and γεῖω or γεῖνω, 'to engender.' That which forms gelatinous structure. Dr. Carpenter restricts the term to substances yielding gelatin. The propriety of this may well be questioned.

GEL'ATIN, *Gel'atine*, *Gelat'ina*, *Galreda*, (F.) *Gélatine*, from (F.) *Gélee*, 'gelly or jelly.' From (L.) *Gelus*, 'ice.' An immediate animal principle. It is semitransparent, insipid, inodorous, insoluble in cold water, very soluble in hot, which it thickens, and transforms into jelly on cooling. Gelatin is a nutritious substance; and, when dissolved in a considerable quantity of water, forms an emollient fluid, much used in therapeutics, but not the most easy of digestion. Two forms of gelatin are admitted—the one *glutin* or gelatin proper: the other *chondrin*.

GELATIN OF WHARTON, *Gelat'ina Whartonia'na*, *Jelly of the Cord*. A soft, dense, fluid, gelatinous substance, which envelops the umbilical cord, and is conceived, by some, to be inservient to the nutrition of the fœtus.

GELATINA AQUATICA, *Brasenia hydropeltis*.

GÉLATINE DES CARTILAGES, Chondrin.

GELATINIFORM MATTER OF THE INTESTINE, Albuminose.

GELAT'INOUS, *Gelatino'sus*, (F.) *Gélatineux*. Relating, or appertaining to, or having the character or appearance of gelatin or jelly. Viscid.

GELATINOUS NERVE-FIBRE, see Nerve fibres.

GELATIO, Congelation.

GELDING, Castration.

GELÉE, Gelly—*g. c ntre le Goître*, Balsam, 'aduretted.

GELLY, *Jelly*, *Jus gelatum seu coagulatum*, *Gelu*, (F.) *Gélee*. Same etymon. A substance of a soft consistence, tremulous, and transparent, which is obtained, by an appropriate treatment, from animal and vegetable matters: hence the distinction into *animal* and *vegetable jelly*. The former is merely a concentrated solution of gelatin, left to cool.

Vegetable Jelly is found in the juice of the currant, mulberry, and of almost all acid fruits, when ripe. It is of itself colourless, but almost always retains a little colouring matter of the fruit which has furnished it. It has an agreeable taste; is scarcely soluble in cold water, but boiling water dissolves it readily: the jelly is, however, almost all deposited on cooling. If this aqueous solution be boiled for a long time, it becomes analogous to muciilage, and loses the property of being jellied on cooling.

GELSEMINUM NIT'IDUM, *G. Sempervirens*, *Bigno'nia sempervirens*, *Yellow Jes'samine*. The flowers, root, &c., of this shrub of the Southern States, *Sex. Syst.* Pentandria Digynia, *Nat. Ord.* Apocynæ, are narcotic, and the effluvia from the former are said sometimes to induce stupor.

GELU, Gelly.

GELUS, Risus.

GEMELLI, *Gastrocnemii*, *Ischio-trochanterianus*, Testicles.

GEMEL'US, *Gem'inus*, 'a twin,' *Did'yms*, (F.) *Jumeau*, *Jumelle*, *Besson*, *Bessonne*. One of two children, twins, born at the same accouchement, or gestation. Also, relating to twins, as 'a twin conception.'

GEMELLUS MUS'culus. Cowper applies this name to the long portion of the triceps brachialis united to the inner portion.

GEMINI, *Ischio-trochanterianus*, Testicles.

GEMINUM CENTRUM SEMICIRCULARE, *Tænia semicircularis*.

GEMINUS, *Gemellus*.

GEMIPOMA, *Mamma*.

GÉMISSEMENT, see Moaning.

GEMMA, Granulation—*g. Oculi*, Crystalline.

GEMMATION, see Generation.

GEMMATION, GENERATION BY, see Generation.

GEMMIPARITÉ, see Generation.

GEMMIPAROUS, see Generation.

GEMUR'SA, a corn; also a name given by the ancients to a disease seated between the toes;—the nature of which is unknown to us.

GEN, *Gen'esis*, 'generation,' from γεῖνω, 'I make.' Hence Hydrogen, Osteogeny, &c.

GENA. The Cheek, *Genys*, *Pare'ia*, *Gamp'hé*, *Gamp'hé'le*, *Gnathos*, *Gnathmus*, *Malas*, (Sc.) *Wongé*, (F.) *Joue*. The cheeks form the lateral parts of the mouth. *Externally*, they have no precise limits: they are continuous, *above*, with the lower eyelid; *below*, they descend as far as the base of the jaw; *before*, they terminate at the *alæ nasi*, and at the commissures of the lips; and *behind*, at the ear. Their thickness varies, according to the degree of fatness of the individual. They are formed of three layers;—one dermoid, another muscular, and the third mucous.

GENCIVES, *Gingivæ*.

GENEANTHROPY, *Anthropogenia*.

GENE'IAS, *Lanugo prima*, *Probar'bium*. The downy hairs, which first cover the cheek. Also, a bandage which passes under the chin.

GENEION, Beard, Mentum.

GENERAL ANATOMY, see Anatomy—*g.* Practitioner, Surgeon-apothecary.

GÉNÉRALE, *Influenza*.

GENERATIO, Generation—*g.* *Æquivoca*, see Generation—*g.* *Calculi*, *Lithia*—*g.* *Homogenea*,

see Generation—*g.* *Prinigena*, see Generation—*g.* *Primitiva*, see Generation—*g.* *Originaria*, see Generation—*g.* *Spontanica*, see Generation.

GENERATION, *Genera'tio*. *Gen'esie*, *Genne'is*, *Genê*, *Gonus*, *Procrea'tio*, *Procreation*, *Breeding*, from (*Lu.*) *geno*, *genere*, *genitum*, *γενω*, or *γενωω*, 'to engender.' Under this name physiologists comprehend the aggregate of functions, which concur, in organized beings, towards the production of their kind. The act of generation means the union of the sexes. See COITION. The writers of antiquity believed, that all organized bodies are produced either by what is termed *univocal* or *regular generation*, *Homogen'esie*, *genera'tio homogen'ea*, *propaga'tio*, which applies to the upper classes of animals and vegetables, or by *spontaneous generation*, *Autogen'ia*, *heterogen'esie*, *generatio heterogen'ea* seu *aquiv'oca* seu *primiti'va* seu *primig'ena* seu *origina'ria* seu *sponta'nea*, *sponteparité* (*Dugès*), which they considered applicable to the very lowest classes only, as the mushroom, the worm, the frog, &c. There are still many distinguished naturalists who consider that beings, low in the scale of animality, are produced in the latter way. Spontaneous generation and *equivocal generation* have been regarded by many to be synonymous. Others, however, mean by spontaneous generation, the production of a new being from the mere combination of inorganic elements; whilst by *equivocal generation* they understand the evolution of a new being from organized beings dissimilar to themselves, through some irregularity in their functions, or through the incipient decay or degeneration of their tissues. As to the mode in which regular generation is accomplished, there have been many views. According to the doctrine of Hippocrates, and of the ancient philosophers, the ovaries of the female furnish a prolific fluid, similar to that of the male; and the fetus results from the mixture of the two seeds in copulation. Steno and others conceived, that the ovaries contain ova, which are not developed until vivified by the male sperm. Bonnet and Spallanzani believed in the pre-existence of germs, created since the origin of the world, but *encased* in each other, and becoming developed in succession; whence it would follow that the ovary of the first female must have contained the germs of all subsequent generations: and that the number of these germs must go on always diminishing, until ultimately extinct. This was the system of the *evolution of germs*. According to Leeuwenhoek, the ovaries do not contain eggs, but vesicles destined to receive animalcules; which, in his view, live in the sperm. Thousands of these animalcules are thrown into the uterus during copulation, and the most expeditious and vigorous reaches the ovary, after having scattered and destroyed its competitors. Buffon—admitting the hypothesis of the two seeds—supposed that they were formed of molecules proceeding from every part of the body of each parent; and that, by a kind of elective affinity, those which were furnished by the head, the trunk, or the extremities of the male parent, could only unite with those proceeding from the same parts of the female. Before him, Maupertuis, admitting, with many of the ancient philosophers, the system of *Epigenesis*, and adopting, as regarded the composition of the sperm, a theory analogous to that of Buffon, had supposed that the molecules, capable of being organized, were attracted towards a centre; that the nose attracts the two eyes; the body, the arms; the arms, the hands, &c., nearly as the particles of a salt, dissolved in a liquid, arrange themselves in regular crystals around the same nucleus. These and various other systems have been successively proposed

and abandoned, and the mystery of generation remains impenetrable.

The simplest kind of reproduction does not require sexual organs. The animal separates into several fragments, which form so many new individuals. This is *Fissip'arous generation*, *Fissip'arism*, *G. from fission*, (*F.*) *Fissiparité*, *Scissiparité*, *Scission*, *Cloisonnement*, *Reproduction ou Multiplication mérismatique*. *Gemmip'arous generation*, (*F.*) *Gemmiparité*, *Génération par Gemmation*, *Surculation* ou *Bourgeonnement*, consists in the formation of buds, sporules or germs on some part of the body, which at a particular period drop off and form as many new individuals. These kinds of reproduction require but one parent—*Monog'eny*. In *Ovip'arous generation*, (*F.*) *Oviparité*, the egg is hatched out of the body. In *ovovivip'arous generation*, the new being is hatched in the excretory passages. In *vivip'arous generation*, the new individual is born under its appropriate form; and in *mar'sup'ial* or *mar'sup'iate generation*, the young being, born at a very early stage of development, is received and nourished in a *mar'supium* or pouch. In *alternate generation*, (*F.*) *Génération alternante*, the young not only do not resemble the parent at birth, but remain dissimilar during their whole life, so that their relationship is not apparent until a succeeding generation. Thus, the cercaria undergoes a change into the distoma. See METAGENESIS. All these kinds of reproduction require the union of sexes—*Dig'eny*.

All the acts comprising the function of generation in man may be referred to five great heads. 1. *Copulation*. 2. *Conception* or *fecundation*. 3. *Gestation* or *Pregnancy*. 4. *Delivery* or *Accouchement*; and, 5. *Lactation*.

GENERATION, ACT OF, see Generation—*g.* *Alternate*, see Generation, and Metagenesis—*g.* *Equivocal*, see Generation—*g.* by *Fission*, see Generation—*g.* *Fissiparous*, see Generation—*g.* by *Gemmation*, see Generation—*g.* *Gemmiparous*, see Generation—*g.* *Marsupial*, see Generation—*g.* *Organs of*, female, see *Vulva*—*g.* *Oviparous*, see Generation—*g.* *Regular*, see Generation—*g.* *Spontaneous*, see Generation—*g.* *Univocal*, see Generation—*g.* *Viviparous*, see Generation.

GEN'EROUS, *Genero'sus*. From *genus*, *genc'ris*, 'a race'; *generosus*, 'of a noble race, noble.' A name given to wines which contain a great quantity of alcohol.

GENESIAL, Genital.

GENESIOLOGY, *Genesiolog'ia*, from *γενεσις*, 'generation,' and *λογος*, 'a discourse.' The doctrine of generation.

GENÉSQUE, *LE*, Appetite, venereal.

GENESIS, Generation.

GENÉT, *Spartium scoparium*—*g.* *à Balai*, *Spartium scoparium*—*g.* *des Teinturiers*, *Genista tinctoria*.

GENETHLIACUS, from *γενεθλιος*, 'natal,' 'pertaining to nativity.' A name given by the ancients to certain astrologers, who, from the state of the heavens at the time of the birth of an individual, predicted his future character and the events of his life.

GENETICA, from *γενεσις*, 'generation.' Diseases of the sexual functions; the 5th class in Good's Nosology. Also, agents that act on the sexual organs.—Pereira.

GENETICUS, Genital.

GENEVA, Gin, see Spirit.

GENÉVRIER, *Juniperus communis*—*g.* *Oxycedre*, *Juniperus oxycedrus*.

GENIVITE, *Ulitis*.

GÉNI, Genian.

GEN'IAN, *Genia'nus*, *Ge'nial*, from *γενειον*, 'the chin.' The *Genian apoph'ysis* or *Process*,

(F.) *Apophyse génienne* ou *géné*, is situate at the posterior part of the symphysis menti, and is formed of four small tabercles.

GENICULATUM, see Corpora geniculata.

GÉNIE, *Genius*. The French sometimes apply this term to diseases nearly synonymously with *nature*; as *Génie inflammatoire*, *G. bilieux*, *G. adynamique*. Some use it in the same sense as type; *Génie intermittent*. The unwonted predominance of any mental faculty is also so called.

GÉNIEN, Genian.

GENIÈVRE, Gin, Juniperus communis (the berry.)

GENIO, from γενιον, 'chin.' In composition, means relating or appertaining to the chin.

GENIOGLOSSUS, *Genio-hyoglossus*, from γενιον, 'the chin,' and γλωσσα, 'the tongue.' Mesoglossus, Mesoglossus, Nonus lingua musculus. The name of a flat, triangular muscle, which extends from the genian apophysis to the inferior surface of the os hyoides and tongue, which last it carries forward.

GENIOHYODES, Geniohyoideus.

GENIOHYOGLOSSUS, Genioglossus.

GENIOHYOIDES, Geniohyoideus.

GENIOHYOIDEUS, *Geniohyoideus*, from γενιον, 'the chin,' and ὡς, 'the os hyoides.' A muscle which arises from the genian apophysis, and is inserted at the anterior part of the body of the os hyoides. Its use is to raise the os hyoides, and carry it forwards. It may, also, contribute to depress the lower jaw in contracting towards the os hyoides.

GENIO-PHARYNGEUS, from γενιον, 'the chin,' and φάρυγξ, 'the pharynx.' A name given, by some anatomists, to a bundle of fibres which passes from the lower jaw to the sides of the pharynx, and forms part of the constrictor pharyngis superior.

GENIOPLASTY, *Genioplasty*, from γενιον, 'the chin,' and πλασσω, 'I form.' The operation for restoring the chin. At times used instead of *Genioplasty*, for the operation for restoring the cheek.

GENIPA OBLONGIFOLIA, *Huito*. A plant of Peru, with the juice of which the Indians paint their legs, to protect them against the stings of insects.

GENIPI ALBUM. *Achillea moschata*, *Artemisia rupestris* — *g. Blanc*, *Artemisia rupestris* — *g. Verum*, *Achillea atrata*.

GENISTA, *Spartium scoparium* — *g. Canariensis*, see *Rhodium lignum* — *g. Hirsuta*, *Spartium scoparium* — *g. Scoparia*, *Spartium scoparium*.

GENISTA SPINOSA IN DICA, *Bahel Schulli*. *Ord. Leguminosæ*. An oriental tree, a decoction of the roots of which is diuretic. The leaves boiled in vinegar have the same effect. — Ray.

GENISTA TINCTORIA, *Genistoïdés tinctoria*, *Spartium tinctorium*, *Dyers' broom* or *weed*, *Green weed*, *Wood wazen*, *Woodweez*, (F.) *Genêt des Teinturiers*. A shrub cultivated in this country and in Europe. The flowering tops and seed have been used in medicine. It has the same properties as *Spartium scoparium*.

GENISTOIDES TINCTORIA, *Genista tinctoria*.

GENITAL, *Genita'lis*, *Genet'icus*, *Generatio'ni inser'viens*, *Gene'sial*, *Gennet'icus*. Same etymon as *Generation*. That which belongs or relates to generation.

GENITAL ORGANS, *Sex'ual Organs*, *Puden'da*, *Natura'lia*, *Nat'ura*, *Eda'ca*, *Me'zea*, *Me'ra*, *Mo'rion*; *Genital*, *Noble*, *Nat'ural* or *Private parts*; *Privities*, *Privy parts*, *Privy Members*, *the Parts*, *Pars*, *Pars corporis seu obsec'na*, *Fores*, *Partes genita'les seu generatio'ni inservien'tes seu obsec'*

na, *Me'dea*, *Veren'da*, *Pedes*, *Inguen*, *Genita'lia*, *Gennet'ica*, *Gymna*, *Membra puden'da*, *Organa generatio'ni inservien'tia*, *O. sexua'lia seu genita'lia*, (F.) *Organes génitales*, *Parties génitales* ou *honteuses* ou *génétoires* ou *nobles* ou *sexuelles* ou *naturelles*, *Les Parties*, *Génétoires*. The parts that are inservient to the reproduction of the species. These are very different in the male and female. In man, they are numerous; some secreting the sperm, as the testicles and their appendages; others retaining it, as the vesiculæ seminales; and another for carrying it into the organs of the female, — the penis. In the female, the parts of generation form an apparatus, perhaps more complicated than that of the male. Some are inservient to copulation, as the vulva, vagina, &c., others to conception and the preservation of the product for a determinate time, as the uterus and its appendages; whilst others concur in the alimentation of the infant after birth, as the mammae.

GENITALE, Sperm — *g. Caput*, *Glans*.

GENITALIA, Genital organs — *g. Viri*, *Pudibilia*.

GENITO-CRURAL NERVE, *Nervus genito-cruralis*, *Subpu'bial* or *Inter'nal inguinal* nerve. A branch of the second lumbar nerve, which passes through the psoas muscle, and, approaching the femoral arch, divides into two branches, — an internal, scrotal, or gen'ital, *nervus spermaticus seu puden'dus exter'nus*, and an external or femoral cutaneous branch, *lumbo-inguinalis*.

GENITO-URINARY, see Uro-genital.

GÉNITOIRES, Genital Organs.

GENITURA. That which is fecundated or engendered in the maternal womb. This word has been used synonymously with *embryo*, *fœtus*, and *infant*. Also, the sperm; and the penis. See GONÉ.

GENIUM, Mentum.

GENNESIS, Generation.

GENNETICA, Genital Organs.

GENNETICOCNESMUS, from γεννητικός, 'genital,' and κνησμος, 'itching.' Itching of the genital organs.

GENNETICON'OSI, *Genneticon'osi* from γεννητικός, 'genital,' and νοσος, 'disease.' Diseases of the genitals.

GENNET'ICUS, Genital.

GEN'OA, CLIMATE OF. The climate of this Italian city and its vicinity has been often selected as favourable for the phthisical valetudinarian during the winter; but it is now admitted to be decidedly improper for pulmonary affections, being subject to frequent and rapid changes of temperature, and to dry cold winds from the north, alternately with warm moist winds from the south-east. — Sir James Clark.

GENONU'SI, *Morbi sexus*, from γυνος, 'sex,' and νοσος, 'disease.' Sexual diseases.

GENOPLASTIC, Meloplastic.

GENOS, Sex.

GENOU, Genn.

GÉNOUILLES, (Corpa,) Corpora geniculata.

GÉNOUILLET, *Convallaria polygonatum*.

GENRE, Genus.

GENSANG, *Panax quinquefolium*.

GENTIA, *Gentianina*.

GENTIAN, (from *Gentius*, King of Illyria, who used some species medicinally), *Gentiana lutea*, *Triosteum* — *g. Blue*, *Gentiana catesbaei* — *g. Catesbian*, *Gentiana catesbaei* — *g. Horse*, *Triosteum* — *g. White*, *Laserpitium latifolium* — *g. Southern*, *Gentiana catesbaei* — *g. White*, *Triosteum* — *g. Yellow*, *Gentiana lutea*, see *Calumba*.

GENTIANA, *g. lutea* — *g. Alba*, *Laserpitium latifolium* — *g. Amarylloides*, *G. quinqueflora* — *g. Cachenlagen*, *Chironia chilensis* — *g. Centaurium*, *Chironia centaurium*.

GENTIANA CATESBÆI; *G. Catesbia'na*, *Blue Gentian*, Ph. U. S., *Catesbian* or *Southern Gentian*, *Bluebelle*, *Bitterroot*. Ord. *Gentianacæ*. It is a pure and simple bitter, and the root may be used wherever that of the *Gentiana lutea* is proper.

GENTIANA CHIRAYTA, *G. Chirayi'ta*, *Henrice'a Pharmacæ'cha*, *Swer'tia*, *Agath'ies chirayi'ta*, *Ophelia chirai'ta*, *Chire'ta*, *Chirayi'ta*, *Chiræ'ta*. A native of India, which has been much employed in that country, in dyspepsia, and as an antiperiodic in intermittents. It is preferred by some to sarsaparilla, where the latter is considered to be indicated,—as after large quantities of mercury have been taken, or where profuse salivation has been induced. It has also been advised in atonic leucorrhœa. It yields its virtues to alcohol and water.

GENTIANA CRINITA, see *G. quinqueflora*—*g. Gerardi*, *Chironia centaurium*.

GENTIANA LUTEA. The systematic name of the official gentian; *Gentia'na*, *G. major* seu *vet'ernum seu rubra*, *Swer'tia lutea*, *Asterias lutea*, *Yellow Gentian*, *Felwort*. Ord. *Gentianæ*. (*F.*) *Gentiane jaune*. This is a plant common in the mountains of Europe. The root is almost odorless, extremely bitter, and yields its virtues to ether, alcohol, and water. It is tonic and stomachic; and in large doses, aperient. Dose, gr. x to ℥ij. It is most frequently, however, used in infusion or tincture.

GENTIANA MAJOR, *G. lutea*—*g. Ochroleuca*, see *G. saponaria*—*g. Peruviana*, *Chironia Chilensis*.

GENTIANA QUINQUEFLO'RA, *G. amaryll'oides*, *Five-flowered Gentian*; and **GENTIANA CRINITA**, *Fringed Gentian*, indigenous, are possessed of like virtues.

GENTIANA RUBRA, *G. lutea*.

GENTIANA SAPON'RIA, *Soapwort Gentian*, and *G. ochroleuca*, *Sampson's Shakeroot*; indigenous, have the properties of *Gentiana lutea*; and are said also to be diaphoretic.

GENTIANA VETERUM, *G. lutea*.

GENTIANE JAUNE, *Gentiana lutea*.

GENTIANI'NA, *Gentianine*, *Gentianin*, *Gentia'nia*, *Gent'ia*. A supposed neutral substance, obtained from gentian, and, by some, presumed to be its active principle. It is not so.

GENTILITIUS, *Hereditary*.

GENU, *Gony*, γονυ, (*F.*) *Genou*. The articulation of the leg with the thigh;—the *Femoro-tibial* or *knee-joint*, the *knee*. It is the most complicated in the body, and is formed by the inferior extremity of the femur, the superior extremity of the tibia, and the rotula. The articular surfaces of the bones are covered by layers of cartilage—more or less thick—and by the synovial membrane of the articulation. The soft parts of this joint are,—the *ligamentum patellæ*, two lateral ligaments, distinguished by the names *internal* and *external*; a posterior ligament, *Ligamentum post'cum Winslow'ii*; two crucial ligaments,—the one anterior, and the other posterior; two interarticular fibro-cartilages; some albugineous fibres, which form an imperfect capsule, &c. The knee receives its arteries from the femoral and popliteal. They bear the name *articular*. Its veins have the same distribution as the arteries, and discharge their blood into the saphena and crural. Its nerves are furnished by the sciatic, popliteal, and crural. The joint is protected by the tendons and muscles which surround it.

The French use the term *Articulation en genou* for a joint, in which the head of a bone is received into a bony cavity of another, where it rolls and moves in all directions.

GENU CORPORIS CALLOSI, see *Corpus callosum*—*g. Varum*, *Entogonyancon*.

GENUGRA, *Gonagra*.

GENUINUS, *Legitimate*.

GENUS, (*F.*) *Genre*. A collection or group of species, analogous to each other, and which can be united by common characters. When a species cannot be referred to a known genus, it constitutes a distinct one.

GENUS CURATIONIS, *Ratio medendi*.

GENY, *Genys*.

GENYANTRAL'GIA, from γένυς, 'the maxilla,' αντρον, 'the antrum,' and αλγος, 'pain.' Pain in the antrum of Highmore.

GENYANTRI'TIS, from γένυς, 'the maxilla,' αντρον, 'the antrum,' and itis, denoting inflammation. Inflammation of the antrum of Highmore.

GENYANTRUM, Antrum of Highmore.

GENYOXYNANCHE, *Cynanche parotidæa*.

GENYOPLASTY, see *Genioplasty*.

GENYS, γένυς, *Genu*, γένυ, 'the jaw;' also the chin. See *Gena*.

GEOFFRÆA INER'MIS, *Geoffræ'a*, *G. racem'osa* seu *Piso'nis* seu *Jamaicæ'nsis*, *Vouacap'oua America'na*, *Geoffroy'a*, *Cabbag'ium*, *Andi'ra inerm'is* seu *racem'osa*, *Cabbage Tree*, *Cabbage Bark Tree*, *Worm-bark Tree*. Ord. *Leguminosæ*. *Sex. Syst.* *Diadelphia Decandria*. The odour of the bark is very unpleasant. It is antelmintic and cathartic. Dose of the powder, ℥j to ℥ij.

GEOFFRÆA JAMAICENSIS, *G. inermis*—*g. Piso'nis*, *G. inermis*—*g. Racemosa*, *G. inermis*. **GEOFFRÆA SURINAMENSIS**, *Andi'ra Surinamen'sis*, has similar properties.

Hutteneschmidt has separated their active principles, to which he has given the names *Jamaicine* and *Surinamine*. They are all alkaline.

GEOFFRÆA VERMIF'UGA, *Andira ibai*, *Arriba*, *Skolemo'ra Fernambucen'sis*; a South American plant, has a fruit, the almond of which, called *angelina*, has a reputation at Rio Janeiro as a vermifuge. Dose, a grain to fifteen; or it may be given in infusion.

GEOFFROYA, *Geoffræa inermis*—*g. Jamaicensis*, *G. inermis*.

GEOGRAPHY, MEDICAL, *Geographia Med'ica*, *Nosogeography*, *Nosogeograph'ia*. The description of the surface of the globe as regards the influence of situation on the health, vital functions, and diseases of its inhabitants, vegetable and animal; but principally on those of man.

GEOPH'AGISM, *Geophag'ia*, *Geophag'ia*, from γη, 'earth,' and γαγν, 'I eat.' The act or practice of eating earth. See *Chthonophagia*.

GEORGIA, MINERAL WATERS OF. The water of *Indian Springs*, in Butts County, is sulphureous. The *Warm Springs*, in Meriwether County, have a temperature of 90° Fahr. *Madison's Springs*, in Madison County, are a chalybeate; and so are *Rowland's Springs*, in Cass County. All these are frequented by invalids.

GERÆOLOG'IA, *Geratolog'ia*, from γρηας, γρηας, 'old age,' and λογος, 'a discourse.' The doctrine, or a description, of old age.

GERÆLOURIS, *Dianthus caryophyllus*.

GER'ANIS, from γρηας, 'a crane,' which it resembled. A bandage, used by the ancients in cases of fractured clavicle. Some authors attribute it to Hippocrates; others, to Perigenes.

GERANIUM, same etymon, because its pistil is long, like the bill of the crane. Ord. *Geraniacæ*. The *Crane's Bill*.

GERANIUM CAROLINIANUM, see *G. Robertianum*—*g. Fœtidum*, *G. Robertianum*—*g. Maculatum*, see *Geranium*, and *G. Moschatum*.

GERANIUM MOSCHA'TUM, *Erod'ium Moschatum*, (*F.*) *Bec de grue musqué*. A European plant, esteemed to be excitant and diaphoretic.

GERANIUM NOVEBORACENSE, *G. Maculatum*—*g. Purpureum*, *G. Robertianum*.

GERANIUM ROBERTIA'NUM, *G. foetidum* seu *purpureum*, *Stinking Crane's Bill*, *Herb Robert*, (F.) *Herbe à Robert*, *Bec de Grue Robertin*. This plant was, at one time, used as an antispasmodic and slight stimulant, as well as for an external application in various painful sores and inflammations.

GERANIUM CAROLINIA'NUM, has similar properties.

Most of the species of geranium have been used as astringents. In some of the northwestern parts of the United States, the root of *Geranium maculatum*—*Geranium*, (Ph. U. S.)—*G. Noveboracense*, *Spotted Crane's Bill*, *Crowfoot*, *Alum root*, *Tormentil*, *Stork-bill*—is called *Racine à Becquet*, after a person of that name. It is highly extolled by the Western Indians as an antisyphilitic.

GERAS, *Senectus*.

GERCURE, *Fissure*.

GERM, *Germen*, *Blasē*, *Blasē'ma*, (F.) *Germe*. The rudiment of a new being, not yet developed, or which is still adherent to the mother.

GERM FORCE, *Plastic force*.

GERMANDER, **COMMON**, *Teucrium chamædrys*—*g. Creeping*, *Teucrium chamædrys*—*g. Marum*, *Teucrium marum*—*g. Small*, *Teucrium chamædrys*—*g. Water*, *Teucrium scordium*.

GERMANDRÉE AQUATIQUE, *Teucrium scordium*—*g. Maritime*, *Teucrium marum*—*g. Officinale*, *Teucrium chamædrys*—*g. Scordone*, *Teucrium scordium*.

GERME FAUX, *Conception*, *false*.

GERMEN, *Germ*, *Sperm*—*g. Dentale*, *Dental Pulp*—*g. Falsum*, *Mole*—*g. Spurium*, *Conception*, *false*.

GERMINAL, *Germinative*, *Germinà'lis*, *Germinat'vus*, (F.) *Germinatif*; from *germen*, a germ. Relating or belonging to germ; as "*germinal cell*."

GERMINAL CELL, see *Cytoblast*—*g. Membrane and Vesicle*, see *Molecule*—*g. Nucleus*, see *Molecule*—*g. Spot*, see *Molecule*—*g. Vesicle*, see *Molecule*.

GERMS, **DISSEMINATION OF**, *Panspermia*—*g. Pre-existence of*, see *Preformation*.

GEROBOS'CIA, *Gerontobos'cia*, from *γρηας*, 'old age,' and *βοσκν*, 'food.' Nourishment or maintenance proper for the aged.

GEROCOMEUM, *Gerocomium*.

GEROCO'MIA, *Gerocom'icē*, *Gerontocom'icē*, from *γερων*, *γερωντος*, 'an aged person,' and *κομειν*, 'to take care of.' The part of medicine whose object is the preservation of the health of the aged:—the hygiene of old people.

GEROCOMICE, *Gerocomia*.

GEROCOMI'UM, *Gerocom'eum*, *Presbydochi'um*, *Gerontocom'ium*, *Gerotrophe'um*. Same etymon. An hospital for the aged.

GÉROFLE, see *Eugenia caryophyllata*.

GÉROFLEE JAUNE, *Cheiranthus cheiri*.

GERONTATROPHIA, *Marasmus senilis*.

GERONTOBOS'CIA, *Geroboscia*.

GERONTOCOMICE, *Gerocomia*.

GERONTOCOMIUM, *Gerocomium*.

GERONTO'PIA, from *γερων*, *γερωντος*, 'an old person,' and *ωπ*, 'the eye.' Weakness of sight of the aged.

GERONTOTOXON, *Gerotoxon*.

GERONTOXON, *Gerotoxon*.

GEROTOX'ON, *Gerontox'on*, *Gerontotox'on*, *Leuco'ma Gerontotox'on*, *Mac'ula cor'neæ arcuà'ta*, *Arcus seni'lis*; from *γερων*, *γερωντος*, 'an old person,' and *τοξον*, 'a bow.' A bow-shaped obscenity at the under margin of the cornea, common to old people.

GEROTROPHEUM, *Gerocomium*.

GEROTROPHIA, *Geroboscia*.

GERSA, *Plumbi subcarbonas*—*g. Serpentinae*, see *Arum maculatum*.

GÉSIER, *Ventriculus callosus*.

GESTA, 'things done,' from *gerere*, *gestum*, 'to do,' 'carry.' *Acta*. A Latin term, introduced by Hallé into medical language, to designate, among the objects which belong to hygiene, the functions which consist in the voluntary movements of muscles and organs. In the class *Gesta* are found *sleep*, the *waking state*, *movements* or *locomotion*, and *rest*.

GESTA'TION, *Gesta'tio*, *Phora*, from *gestare*, *gestatum*, 'to carry.' The time during which a female who has conceived carries the embryo in her uterus. See *Pregnancy*. Gestation, likewise, signifies the bearing or carrying of an individual; a kind of exercise easier than that in which he moves by virtue of his own powers. Thus, we speak of *gestation on horseback*, in a *carriage*, &c.

GESTATION, **PROTRACT'ED**. *Pregnancy* protracted beyond the usual period. See *Pregnancy*.

GESTICULA'TION, *Gesticula'tio*, same etymon. The act of making many gestures; a symptom in disease, which indicates great encephalic erethism. It is met with in numerous affections.

GETHYL'IS SPIRA' LIS. A South African plant, *Nat. Ord. Amaryllidæ*, the orange-coloured fruit of which has a peculiar fragrance. An infusion in spirit or tincture is used in flatulence and colic.

GEUM, *G. rivale*—*g. Caryophyllum*, *G. urbanum*—*g. Nutans*, *G. rivale*—*g. Palustre*, *G. rivale*.

GEUM RIVALE, *G. palus'trē* seu *nutans*, *Caryophylla'ta aquat'ica* seu *nutans*, *Benedic'ta sylvestris*, *Water or Purple avens*, *Choc'olate root*, (F.) *Benôte aquatique* ou *des Ruisseaux*. *Ord. Rosacæ*. *Ser. Syst. Icosandria Polygynia*. The root of this plant—*Geum* (Ph. U. S.)—is astringent. It has been much extolled in the cure of intermittents, diarrhoea, hemorrhage, &c.

GEUM URBA'NUM, *G. caryophyllatum*, *Caryophylla'ta*, *G. vulgà'ris* seu *urba'na*, *Caryophyl'lus vulgà'ris*, *Sanamun'da*, *Logophthal'mus*, *Caryophyll'a*, *Janamun'da*, *Herba benedic'ta*, *Common avens*, *Herb Bennet*, *Yellow avens*, *Star of the North*, *Blessed herb*, (F.) *Benôte*. The root of this plant has a smell not unlike that of cloves. Taste bitterish, austere; virtues yielded to water and alcohol. It has been used in intermittents, dysentery, chronic diarrhoea, debility, &c. Dose, ʒss to ʒj of the powder.

GEUM VERNUM, *Western early avens*, has the same properties as

GEUM VIRGINIA'NUM, *White avens*, *Avens*, *Evan root*, *Choc'olate root*, *Bennet*, *Throat root*, *Cure-all*, (F.) *Benôte de Virginie*, is common from Maine to Carolina and Kentucky, flowering in June and July. It has the same medical properties as *Geum rivale*.

GEUMA, *Taste*.

GEUSION'OSI, from *γεωσις*, 'taste,' and *ωσος*, 'disease.' *Geusion'osi*. Diseases of the organ or sense of taste.

GEUSIS, *Taste*. Rarely, the root of the tongue—*Radix lingua*.

GEUTHMOS, *Taste*.

GEZIB, see *Pastinaca opoponax*.

GÉZIT, *Cagot*.

GÉZITAIN, *Cagot*.

GHERKIN, see *Cucumis sativus*.

GHITTA JEMOCO, *Cambogia*.

GIALAPPA, *Convolvulus Jalapi*.

GIANT, *Gigas*, (F.) *Géant*. One much above the ordinary stature; of excessive growth—*Macrosoma'tia*, *Macrosomat'ia*, *Magnitu'do gigante'a*, (F.) *Géantisme*.

GIBBA, Hump.
 GIBBER, Hump.
 GIBBEROSITAS, Gibbositas.
 GIBBOSITAS, *Gibberositas*, *Cyphosis*, *Rhachio-cyphosis*, from (F.) *Gibbeus*, (L.) *Gibbus*, *gibbus*, 'something arched or vaulted; prominent,' *Gibbosity*, *Gibbousness*, *Curvature of the spine*, *Hydro-ma*. A phenomenon which occurs in different diseases, particularly in rickets and caries of the vertebrae. See Hump.

GIBBOSITAS CARIOSA, Vertebral disease.

GIBBOUSNESS, Gibbositas.

GIBBUS, Hump—g. *Pottii*, Vertebral disease.

GIBLETS. According to Minsheu, from *Gobbet*, 'a mouthful' [?]; but, according to Junius, from (F.) *Gibier*, 'game.' The word seems to be the old (F.) *Gibelez*, *Gibelet*, &c., i. e. *Gibier*, (L.) *Cidarium*, 'food,' (F.) *Abattis*. It means, generally, the parts which are cut off from a goose before it is roasted. Also, the extremities of fowls, such as the head, wings, feet; to which are sometimes added the liver, gizzard, &c. Soup, made from these, is moderately nutritious.

GIDDINESS, Vertigo.

GIDDY, Vertiginous.

GIGANTEUS, Gigantic.

GIGANTESQUE, Gigantic.

GIGANTIC, *Giganteus*, *Giganto-des*, (F.) *Gigantesque*. Relating to one much above the ordinary stature.

GIGANTODES, Gigantic.

GIGARTINA HELMINTHOCHORTON, *Corallina Corsicana*—g. *Lichenoides*, *Fucus amylaceus*.

GIGARUS SERPENTARIA, *Arum dracunculus*.

GIGAS, Giant.

GILARUM, *Thymus serpyllum*.

GILEAD, BALM OF, see *Amyris opobalsamum*.

GILET DE FORCE, Waistcoat, strait.

GILL, *Glechoma hederacea*.

GILL-GO-BY-GROUND, *Glechoma hederacea*.

GILLA THEOPHRASTI, *Zinci sulphas*—g. *Vitrioli*, *Zinci sulphas*.

GILLE'NIA STIPULA'CEA, *Small-flowered Indian Physic* has the same properties as GILLE'NIA TRIFOLIA'TA, *Spiraea trifoliata*, *Common Gille'nia*, *Indian Physic*, *Western Dropwort*, *Indian Hippo*, *Ipæaca*, *Bowman's root*, *Meadow sweet*, *Beaumont root*, (F.) *Gillénie*. *Ord. Rosaceae*. *See. Syst. Icosandria Pentagynia*. The root of this shrub, and of *G. stipulacea*—*Gille'nia*, (Ph. U. S.)—which grows plentifully in the United States, is a safe and efficacious emetic, in the dose of about 30 grains. It resembles *ipeacuanha* in its properties.

GILLÉNIE, *Gillenia trifoliata*.

GILLIFLOWER, *Dianthus caryophyllus*.

GILLON, *Viscum album*.

GIMBERNAT'S LIGAMENT. A fibrous, triangular expansion, which is detached from the posterior and inner part of the crural arch, and is inserted into the crest of the pubis. This ligament forms the inner part of the superior aperture of the crural canal. It is one of the most frequent causes of strangulation in crural hernia.

GIN, *Hollands*, *Gene'va*, (F.) *Genièvre*. This spirit, which is distilled from corn and juniper berries, or from some substitute for them, is largely used in Great Britain, and is extremely detrimental, to the lower classes particularly. It possesses the properties of other spirituous liquors, but is diuretic, and, in popular medicine, is more used than other varieties, in cases of colic or intestinal pain of any kind. *Schiedam Schnapps* is understood to be *Hollands* of good quality. See Spirit.

GIN DRINKER'S LIVER, Liver, nutmeg—g. Liver, Liver, nutmeg.

GINGEMBRE, *Amomum zingiber*.

GINGER, *Amomum zingiber*.

GINGER BEER POWDER, see *Amomum zingiber*.

GINGER, GARDEN, *Capsicum*—g. *Indian*, *Asarum Canadense*—g. *Jamaica*, concentrated essence of, see *Amomum zingiber*—g. *Preserved*, see *Amomum zingiber*—g. *Wild*, *Asarum Canadense*.

GINGIBRACHIUM, from *gingivæ*, 'the gums,' and *brachium*, 'the arm.' A name given to the scurvy, because the gums and arms are chiefly affected by it. It has also been called *Gingipe'dium*, because the lower limbs are in many cases the seat of scorbutic spots. See *Porphyra nautica*.

GINGIPEDIUM, see *Gingibrachium*, *Porphyra nautica*.

GINGIVALIS, from *gignere*, 'to beget,' because the teeth are, as it were, begotten in them. [?] The gums, *oua*, *U'ta*, *Carnicula*, (F.) *Gencives*. The portion of the mucous membrane of the mouth which covers the maxillary bones to the level of the alveolar arches. The gums are formed of a red tissue, more or less solid, and of a fibro-mucous nature, which adheres strongly to the necks of the teeth, and transmits, between the roots and their alveoli, a very thin expansion,—the *alveolo-dental periosteum*. The gums fix the teeth and contribute greatly to their solidity. In the aged, after the loss of the teeth, they become fibrous and very solid, and are inservient to mastication.

GINGIVALIS, *Uleticus*, *U'licus*; from *gingivæ*, 'the gums.' Relating to the gums.

GINGLYMOID, *Ginglymoïdeus*, *Ginglymoïdes*, from *γινγλυμος*, 'a ginglymus or hinge,' and *αἶδος*, 'resemblance.' Resembling a ginglymus or hinge. An epithet applied to joints which resemble a hinge, as a *ginglymoid joint*.

GIN'GLYMUS, *Cardinamentum*, *Cardo*, 'a hinge.' (F.) *Charnière*, *Articulation en charnière*. A species of diarthrodial articulation, which only admits of motion in two directions, like a hinge,—as the knee-joint or elbow-joint.

GINSENG, *Panax quinquefolium*—g. *Asiatic*, *Panax Schinseng*—g. *Blue*, *Caulophyllum thalictroides*—g. *Yellow*, *Caulophyllum thalictroides*—g. *Horse*, *Triosteum*—g. *White*, *Triosteum*.

GIRAFFE, *Dengue*.

GIRARD-ROUSSIN, *Asarum*.

GIRDLE, *Cingulum*.

GIRL. This seems, formerly, to have been an appellation common to both sexes. Many etymologists deduce the word from the Su. Goth. *Karl*, 'a man.' It means a young female, (L.) *Fil'ia*, from *φιλεῖν*, 'to love.' (F.) *Fille*.

GIROFLE, see *Eugenia caryophyllata*.

GIROFLÉE JAUNE, *Cheiranthus cheiri*.

GIROLE, *Sium sisarum*.

GIRON, *Arum maculatum*.

GIRSLE, *Cartilage*.

GISN, *Anhelatio*.

GISSA, *Cynanche parotidea*.

GIZZARD, *Ventriculus callosus*.

GIZZEN, see *Parturition*.

GLABELLA, *Mesophryon*.

GLABELLAD, see *Glabella*.

GLABEL'LAR, from *glabella*, 'the space between the eyebrows.' An epithet for an aspect towards the glabella.—Barclay. *Glabbellad* is used adverbially by the same writer to signify 'towards the glabellar aspect.'

GLABRITIES, *Calvities*.

GLACE, *Ice*.

GLACIALE, *Mesembryanthemum crystallinum*.

GLACIALIS HUMOR, *Crystalline*.

GLACIES, *Ice*.

GLADY'OLUS, diminutive of *gladius*, 'a sword,' from the sword-like shape of its leaf. *Machæroides*—g. *Cæruleus*, *Iris Germanica*.

GLADI'OLUS COMMUNIS, *G. vulgaris*, *Victoria's rotunda* seu *feminea*, *Cornplav*, (F.) *Glacéul. Ord. Iridiæ*. The root of this plant has been considered aphrodisiac. Applied in cataplasms, it has been extolled against scrofulous tumours.

GLADIOLUS LUTEUS, *Iris pseudacorus*—*g. Vulgaris*, *G. communis*.

GLADIUS, Penis.

GLAMA, Lippitudo, *Chassie*.

GLAND, *Glândula*, *Gran'dula*, (diminutive of *glands*, 'an acorn, a kernel.') *Aden*. The ancient anatomists gave this name to a number of organs of a texture generally soft, and a shape more or less globular, but differing greatly in their nature and functions. They applied it, for instance, 1. To those organs which separate from the blood, any fluid whatever. When such organs were composed of several lobules, united by common vessels, they received the name *conglomerate glands*, as the parotid, pancreas, &c. 2. To the reddish and spongy, knot-like bodies, which are met with in the course of the lymphatics. These they called *conglobate glands*;—see *Ganglion*, (*lymphatic*;) and 3dly and lastly, to various other organs, whose intimate texture and functions are still unknown, as the *Pineal gland*, *Pituitary gland*, *Glands of Pacchioni*, *Thyroid gland*, *Thymus gland*, *Supra-renal glands*;—see *Chaussier* restricts the word *gland* to those softish, granular, lobated organs, composed of vessels, and a particular texture, of which there are in the human body, the *lacrimal, salivary, and mammary*, the *testicles*, the *liver, pancreas*, and *kidneys*. These *permanent glands*, or *glands with permanent ducts*, are all destined to draw from the blood the molecules necessary for the formation of new fluids, and to convey these fluids externally, by means of one or more excretory ducts. Several glands, besides their excretory ducts, have especial reservoirs, in which the fluids, secreted by them, collect, remain for a greater or less space of time, and undergo slight modifications before being evacuated;—such are, the gall-bladder for the liver, the urinary bladder for the kidneys, &c. Each gland has an organization peculiar to it, but we know not the intimate nature of the glandular texture.—Malpighi believed that the vessels terminate in small, solid masses, to which he gave the name—*glandular grains* or *acini*. In these, he considered, the excretory ducts originate. Ruysch thought that the glands are entirely vascular, and that the excretory ducts are immediately continuous with the *vasa afferentia*, &c. The best view, perhaps, is, that the exhaling or secreting vessel is distributed on the animal membrane, which forms the blind extremity of the excretory duct, and that the secretion is effected through it by means of cells.

The term *glande* (F.) is sometimes appropriated to the tumour formed by inflammation or engorgement of a lymphatic ganglion.

GLAND, *Glands*—*g. Accessory*, of the Parotid, see *Parotid*—*g. Anteprostatic*, see *Cowper's glands*—*g. Blood*, see *Ganglion*—*g. Globate*, *Conglobate gland*—*g. Prostate*, *Prostate*—*g. Salivary*, *abdominal*, *Pancreas*.

GLANDAGE, *Adenophyma*.

GLANDE, *Gland*—*g. Thyroide*, see *Thyroid gland*—*g. Vulvaire conglomerée*, *Cowper's glands* in the female—*g. Vulvo-vaginale*, *Cowper's glands* in the female.

GLANDERS, *Equinia*—*g. Farcy*, see *Equinia*.

GLANDES BRONCHIQVES, *Bronchial glands*—*g. Cérumineuses*, *Ceruminous glands*—*g. Conglobées*, *Conglobate glands*—*g. de Cowper*, *Cowper's glands* in the female—*g. de Croissance*, *Waxing kernels*—*g. de Litré*, see *Sebaceous glands*—*g. de Meibomius*, *Meibomius glands* of—

g. de Méry, *Cowper's glands* in the female—*g. de Pacchioni*, *Pacchioni glands* of.

GLANDIFORM CORPUSCLE, *Acinus*—*g. ganglion*, see *Ganglion*.

GLANDIUM, *Thymus*.

GLANDS, see *Quercus Alba*—*g. Accessory*, *Cowper's glands*—*g. Aggregate*, *Peyer's glandulæ*—*g. Agminated*, *Peyer's glands*—*g. Aporic*, *Ganglions*, *glandiform*—*g. Blind*, *Ganglions*, *glandiform*—*g. of Bartholinus*, *Cowper's glands* in the female—*g. Brunner's*, see *Brunner's glands*—*g. Cowper's*, see *Cowper's glands*—*g. Diapnogenous*, see *Perspiration*—*g. Ductless*, see *Ganglion*—*g. of Duverney*, *Cowper's glands* in the female—*g. Havers's*, *Synovial glands*—*g. Lenticular*, *Lenticulares glandulæ*—*g. Lieberkühn's*, *Lieberkühn's glands*, see *Intestine*—*g. Military*, *Sebaceous glands*—*g. of Naboth*, *Nabothi Glandulæ*—*g. Oil*, *Sebaceous glands*—*g. of Pacchioni*, *Glandulæ Pacchioni*—*g. Permanent*, see *Gland*—*g. Peyer's*, *Peyer's glandulæ*—*g. Renal*, *Capsules*, *renal*—*g. Sebaceous*, see *Sebaceous glands*—*g. Solitary*, *Brunner's glands*—*g. Sudoriparous*, see *Perspiration*—*g. Sweat*, see *Perspiration*.

GLANDS, *TEMPORARY*, *Glands without permanent orifices*. Glands, that consist of a single primary vesicle or sacculus, which, having elaborated a secretion in its interior, bursts, discharges it, and disappears. The Graafian vesicles and the gland cells afford examples of these.

GLANDS OF TYSON, *Sebaceous glands* of *Tyson*—*g. Vascular*, see *Ganglion*—*g. of Vesalius*, *Bronchial glands*—*g. of Willis*, *Albicantia corpora*, *Mammillary tubercles*.

GLANDULA, *Gland*—*g. ad Aures*, *Parotid*—*g. Avicennæ*, *Encystis*—*g. Bartholiniana*, *Sublingual gland*—*g. Basilaris*, *Pituitary gland*—*g. Colli*, *Tonsil*—*g. Innominate*, *Galen's Lacrymal gland*—*g. Lacrymalis*, *Lacrimal gland*—*g. Lactifera*, *Mammary gland*—*g. Mucosa*, *Conglobate gland*—*g. Pinealis*, *Pineal gland*—*g. Pituitosa*, *Pituitary gland*—*g. Riviniana*, *Sublingual gland*—*g. Salivaris abdominalis*, *Pancreas*—*g. Socia Parotidis*, see *Parotid*—*g. Thymus*, *Thymus*—*g. Thyroidea*, *Thyroid gland*.

GLANDULÆ ACINOSÆ, see *Acinus*—*g. Agminatæ*, *Peyer's glandulæ*—*g. Articulares*, *Synovial glands*—*g. Assistentes*, *Prostate*—*g. Brunner's*, *Brunner's glands*—*g. Cervicis uteri*, *Nabothi glandulæ*—*g. Coronæ penis*, see *Sebaceous glands*—*g. Cowper's*, *Cowper's glands*—*g. Duræ matris*, *g. Pacchioni*—*g. Duræ meningis*, *G. Pacchioni*—*g. Agmen congregatæ intestinorum*, *Peyer's glandulæ*—*g. Intestinales*, *Peyer's glandulæ*—*g. Lieberkühniannæ*, *Lieberkühn's glands*—*g. Littrianæ*, see *Sebaceous glands*—*g. Meibomianæ*, *Meibomius glands* of—*g. Mucosæ congregate intestinorum*, *Peyer's glandulæ*—*g. Muciparæ racematum congestæ intestinorum*, *Peyer's glandulæ*—*g. Myrtiformes*, *Carunculæ myrtiformes*—*g. Odoriferæ*, see *Sebaceous glands*—*g. Odoriferæ penis*, see *Sebaceous glands*—*g. Odoriferæ Tysoni*, see *Sebaceous glands*—*g. Pacchioni*, *Pacchioni glands* of—*g. Peyerianæ*, *Peyer's glandulæ*—*g. Plexiformes*, *Peyer's G.*—*g. Prostatæ mulierum*, see *Corpus glandulosum mulierum*—*g. Sebaceæ ciliares*, *Meibomius glands* of—*g. Sebaceæ glandis*, see *Sebaceous glands*—*g. Solitaria*, *Brunner's glands*—*g. Spuriæ*, *Peyer's glandulæ*, *Ganglions*, *glandiform*—*g. Sudoriferæ*, see *Perspiration*—*g. Suprarenales*, *Capsules*, *renal*—*g. Tubulosæ*, see *Tubular*—*g. Tysoni*, *Sebaceous glands* of *Tyson*—*g. Uterinæ*, *Utricular glands*—*g. Utriculares*, *Utricular Glands*—*g. Vasculosæ*, *Conglomerato glands*—*g. Vesaliamæ*, *Bronchial glands*.

GLANDULAIRE, *Glandular*.

GLAND'ULAR, *Glandularis*, *Glanduleux*, (F.) *Glandulaire*, *Glanduleux*. Having the ap-

pearance, form, or texture of *Glands*; as a *glandular* body, a *glandular* texture, &c.

GLANDULAR SUBSTANCE OF THE KIDNEY, see Kidney.

GLANDULE OF MALPIGHI, see Corpora Malpighiana.

GLANDULEUX, Glandular.

GLANDULOUSO-CARNEUS. Ruysch gives this epithet to fleshy excrescences which he found in the bladder.

GLANDULOSUS, Glandular.

GLANS, genitive *glandis*, ('an acorn.') *Balanus*, *Cyt'taros*, *Cuspis*, *Caput*, *C. Penis*, *Genita'le caput*, (F.) *Gland*. The extremity of the penis and of the clitoris. The *glans penis* is of a conical, slightly flattened shape. It is continuous with the urethra, which opens at its apex; and is circumscribed by a projecting edge, called the *Coro'na glandis*, behind which is a depression. *Collum seu Sulcus retroglandula'ris*. It is covered by a thin mucous membrane, is furnished, at its base, with sebaceous follicles, called *glan'dulae odoriferae Tyso'ni*, the secretion from which is termed *Smegma preputii*; and can, almost always, be covered by the reflection of the skin, called the *prepuce*. Lastly, it is formed of a spongy texture, susceptible of being thrown into erection.

GLANS, Bronchocele, Pessary, Suppository.

GLANS CLITORIDIS is smaller. It is imperforate, and likewise covered with a sort of prepuce formed by the mucous membrane of the vulva.

GLANS JOVIS THEOPHRASTI, *Fagus castanea*—*g. Ulnæ*, *Olecranon*—*g. Unguentaria*, *Guilandina moringa*, *Myrobalanus*.

GLAREA, Gravel.

GLASS, Vitrum.

GLAUBER'S SALTS, Soda, sulphate of.

GLAUCEO, Glaucoma.

GLAUCO'MA, from *γλαυκος*, 'sea-green.' *Glaucosis*, *Glauc'do*, *Cataracta glauca*, *Oculus cæsius*, *Cæsius*, *Phtharma glaucoma*, *Paropsis glaucosis*, *Apoglaucosis*. Amongst the older pathologists, this word was used synonymously with cataract. It is now ordinarily applied to opacity of the vitreous humour or of the tunica hyaloidea, which manifests itself by a grayish, or greenish spot, apparent through the pupil. The diagnosis is generally difficult; and the disease is almost always incurable.

GLAUCOMA WOUHOUZI, Cataract.

GLAUCOSIS, Glaucoma.

GLAYEUL, *Gladiolus vulgaris*—*g. Puant*, *Iris foetidissima*.

GLECHO'MA HEDERA'CEA, *G. hirsuta*, *Nep'tea glecho'ma*, *Heder'ula*, *Chamæcle'ma*, *C. hederacea*, *Calamin'tha hederacea*, *Calamin'tha humilior*, *Chamæcis'sus*, *Hed'era terrestris*, *Nep'tea glecho'ma*, *Panacea pectoris*, *Ground-ivy*, *Gill*, *Gill-go-by-ground*, *Alehoof*, *Robin runaway*, (F.) *Lierre terrestre*, *Terrette*, *Rondote*, *Herbe Saint-Jean*. This plant—*Ord. Labiata*—has a strong smell, and a bitterish, somewhat aromatic, taste. It has been considered expectorant and tonic.

GLECHOMA HIRSUTA, *G. hederacea*.

GLECHON, *Mentha pulegium*.

GLECHONITIS. Wine, impregnated with the Glechon, *γλῆχων*, *mentha pulegium* or pennyroyal.

GLEEING, Strabismus.

GLEET, see Gonorrhœa.

GLEME, *Chassie*, Lippitudo.

GLENĒ, γλῆνῃ. The pupil. The anterior part of the eye. The eyeball. The eye. According to some, the crystalline lens. Also, a glenoid cavity.

GLENGORE, Syphilis.

GLENGOUR, Syphilis.

GLENITIS, Phacitis.

GLENN'S SPRINGS, see South Carolina, Mineral waters of.

GLENOID, Glene.

GLENOID, *Glenoid'al*, *Glenoid'al'is*, *Glenoi'des*, *Gleno'des*, *Glenoi'deus*, from γλῆνῃ, and αἶδος, 'resemblance.' (F.) *Glénôide*, *Glénôidale*, *Glénôidien*. Any shallow, articular cavity, γλῆνῃ, *glenê*, which receives the head of a bone; such as, 1. The glenoid cavity or fossa of the scapula, *Fossa glenoidæa*, *Omocol'yte*, *Acetabulum hu'meri*, *Oavi'tas hu'meri glenoidæa*, *Entypo'sis*, is situate at the anterior angle of the scapula; and is articulated with the head of the humerus. 2. The glenoid cavity or fossa of the temporal bone. It is seated between the two roots of the zygomatic process, and receives the condyle of the lower jaw.

GLENOID LIG'AMENT, (F.) *Ligament Glénôidien*, is a fibro-cartilaginous ring or *bouvrelet*, which seems formed by the expansion of the tendon of the long head of the biceps brachialis, and surrounds the glenoid cavity of the scapula, the depth of which it increases.

GLÉNOÏDALE, Glenoid.

GLEYED, see Strabismus.

GLIADINE, Hæmatin.

GLISCHRAS'MA, *Glischrotes*, from γλίσχραινο, (γλῖα, 'glue,') 'I become glutinous.' *Leni-*, viscosity.—Hippocrates.

GLISCHROCH'OLUS, from γλίσχρος, 'viscid,' and χολη, 'bile.' An epithet for excrement which is glutinous and bilious.

GLISCHROTES, Glischrasma.

GLISOMARGO, Creta.

GLISSEMENT DU LAMBEAU, see Flap.

GLISTER, Clyster.

GLOBE, Bandage (head)—*g. of the Eye*, see Eye—*g. Flower*, *Cephalanthus occidentalis*, *Trollius laxus*.

GLOBULAIRE PURGATIVE, Globularia alypum.

GLOBULARIA ALYPUM, *Globularia*, *Montpell'ier Turbith*, (F.) *Globulaire purgative*, *Turbith blanc*. *Ord. Globulariæ*. The leaves of this plant are bitter, and have been used in intermittents and in constipation. See Alypon.

GLOB'ULE, *Glob'ulus*, *Sphæ'rion*, *Sphæ'rula*. A small globe.

GLOBULES OF THE BLOOD, *Blood globules*, *Blood-corpuscles*, *Blood-disks*, *Blood-vesicles*, *Glob'uli seu Vesic'ulæ seu Sphæ'rulæ seu Follic'uli seu Corpus'cula seu Cell'ulæ seu Partic'ulæ seu Gran'ulæ san'guinis*, (F.) *Globules du sang*, *Disques sanguins*, are small, circular bodies, which are particularly observable when the transparent parts of cold-blooded animals are examined by the aid of the microscope; and are met with in the blood of all animals. They are circular in the mammalia, and elliptical in birds and cold-blooded animals; are flat in all animals, and generally composed of a central nucleus enclosed in a membranous sac. Chemically, they consist of hæmatin and globulin—*hæmato-globulin*. The chemical composition of the nucleus, where it exists,—it does not in man,—has been by some regarded as fibrin; by others, as fat; and by others, again, as a species of horn, to which the name *nuclein* has been given.

SIZE OF THE GLOBULES.

Sir E. Home and Bauer, with colouring matter	1.1700th part of an inch.
Eller.....	1.1930
Sir E. Home and Bauer, without colouring matter.....	1.2000
Müller.....	1.2300 to 1.3500
Mandl.....	1.2625 to 1.3150

Hodgkin, Lister, and Rudolphi, 1.3000	
Sprenzel,.....	1.3000 to 1.3500
Cavallo.....	1.3000 to 1.4000
Donné.....	1.3150 to 1.3280
Jurin and Gulliver.....	1.3240
Blumenbach and Sénac.....	1.3330
Tabor.....	1.3600
Milne Edwards.....	1.3900
Wagner.....	1.4000
Kater.....	1.4000 to 1.6000
Prévost and Dumas.....	1.4056
Haller, Wollaston, and Weber...	1.5000
Young.....	1.6060

Besides these, the blood contains *white globules*, (F.) *Globules blancs*, *Leucocytes*, which are probably lymph and chyle corpuscles in process of development into red corpuscles.

Dried blood globules of the ox have been given internally in cases of *anæmia*.

GLOBULE D'ARANTIUS, see Sigmoid valves—*g. du Sang*, Globule of the blood.

GLOBULES BLANCS, see Globules of the blood—*g. Ganglion*, see Neurine—*g. Chyle*, see Chyle—*g. Lymph*, see Lymph—*g. Milk*, see Milk—*g. Mucous*, see Mucus—*g. Pus*, see Pus—*g. Pyoid*, see Pus.

GLOBULI ARTERIARUM TERMINI, see Acinus—*g. Mammillares*, see Mammillary eminence—*g. Sanguinis*, Globules of the blood—*g. Tartari martialis*, Ferrum tartarizatum—*g. Tartatis ferri et lixivie*, Ferrum tartarizatum.

GLOBULIN, *Glob'uline*, *Blood ca'sein*. The colourless substance that remains after the abstraction of the colouring matter of the blood-corpuscle. It is a peculiar albuminous principle. The globulin of Berzelius consists of the envelopes of the blood globules, and of the part of their contents that remains after the extraction of the hæmatin. Lecanu regards it as identical with albumen; and, according to Mulder, it belongs to the combinations of protein. Robin and Verdeil consider it to be *albuminose*.

The term *globulin* is likewise given by M. Donné to small granulations appertaining to the chyle, which are observable in the blood with the microscope. They are small, white, roundish, isolated or irregularly-agglomerated grains; of about the 1-300 of a millimetre in diameter, and are regarded by M. Donné as the first elements of the blood globules. They are the *white granulated corpuscles* of Mandl.

GLOBULISM, Homœopathy.

GLOBULUS ARANTII, see Sigmoid valves—*g. Nasi*, see Nasus—*g. Sanguineus*, Punctum saliens—*g. Stapedius Ossis*, Os orbiculare.

GLOBUS HYSTERICUS, Angone—*g. Major*, see Epididymis—*g. Minor*, see Epididymis—*g. Martialis*, Ferrum tartarizatum.

GLOBUS UTERI'NUS. A term applied by accoucheurs to the round tumour, formed by the uterus in the lower part of the abdomen immediately after delivery.

GLOMERATE, Conglomerate.

GLOMERULE, *Glomer'ulus*; from *glomys*, 'a slew of thread.' A ball or clew, formed by an agglomeration of vessels; as *Glomerule* of Malpighi; see Corpora Malpighiana.

GLOMERULI RENALES, Corpora Malpighiana.

GLOSSA, *Glotta*, 'the tongue.' The power of speech. Speech. Hence:

GLOSSAGRA, Glossalgia.

GLOSSAL'GIA, *Gloss'agra*, *Glottal'gia*, *Glott'agra*, from *γλωσσα*, 'the tongue,' and *αλγος*, 'pain.' Pain in the tongue.

GLOSSANIS'CHUM, *Glossanoch'eus*, *Glossan'*

ochum, from *γλωσσα*, 'tongue,' and *αρχειν*, 'to hold up.' An instrument for holding up the tongue.

GLOSSANOCHEUS, Glossanischum.

GLOSSANOCCHUM, Glossanischum.

GLOSS'ANTHRAX, *Pestis gloss'anthrax*, (F.) *Boussole*, from *γλωσσα*, 'the tongue,' and *ανθραξ*, 'a carbuncle.' Carbuncle of the Tongue. A disease more common in cattle than in man.

GLOSSEPIGLOTTIC, *Glossepiglott'icus*. Relating to the tongue and epiglottis, as *Ligamentum glossepiglott'icum*.

GLOSSIANUS, Lingual muscle.

GLOSSITIS, from *γλωσσα*, 'the tongue,' and *itis*, a suffix denoting inflammation. *Glosson'cus inflammato'rius*, *Angina lingua'ria seu lingua'lis*, *Inflammat'io Lingua*, Inflammation of the tongue, (F.) *Inflammation de la Langue*. When confined to the mucous membrane, it is of slight importance. That which affects the whole of the tongue is a serious disease, and requires the vigorous use of antiphlogistics. It is rare.

GLOSSO, from *γλωσσα*, 'the tongue.' In composition, relating or appertaining to the tongue.

GLOSSOC'ACE, from *glosso*, and *κακος*, 'evil.' Ulceration of the tongue, with symptoms of adynamic fever.

GLOSSOCARCINO'MA, *Glossoscir'rhus*, *Carcino'ma Lingua*, from *glosso*, and *καρκινωμα*, 'cancer.' Cancer of the tongue.

GLOSSOCAT'CHUS, *Glossocat'ochēz*, *Catagloss'eum*, from *γλωσσα*, 'the tongue,' and *κατεχυ*, 'I arrest.' *Lingua Deten'tor*, (F.) *Abaisseur de la langue*. An instrument, the invention of which is attributed to Paulus of Aegina, and which was employed to depress the tongue, in order to examine diseases of the fauces. It was composed of two branches; one of which had, at its extremity, a plate for depressing the tongue; whilst the other, shaped like a horse-shoe, was applied under the chin. The finger, or the handle of a spoon, or a spatula, is generally used in similar cases. See Cataglossum.

GLOSSOCE'LE, from *glosso*, and *κηλη*, 'hernia,' 'tumour.' *Hernia of the Tongue*, *Paragloss'eē*, *Prolap'sus lingua*, *Glossomegis'tus*, *Glossop'tosis*. Projection of the tongue from the mouth. It depends, generally, on an inflammatory swelling of the organ. At times, however, a chronic glossocele, or sort of oedematous engorgement, is met with; which proceeds to a great length, and deforms the dental arches, the lips, &c. Inflammatory glossocele must be combated by antiphlogistics. In the oedematous kind, such as is sometimes caused by excessive salivation, the infiltrated fluid may be pressed back by the hand of the practitioner, to get the tongue behind the teeth; and it may be kept there by a piece of gauze tied over the mouth. The chronic, elongated kind sometimes requires amputation of a portion of the organ.

GLOSSOCOMA, Glossospasmus.

GLOSSOC'OMON, *Glossoc'omum*, *Glossocomi'on*, from *glosso*, and *κομειν*, 'to guard.' The ancients gave this name to a small case for holding the tongues of their wind-instruments. By extension, it was applied to the box or cradle in which fractured limbs were kept. We find, in the ancient writers, a *Glossocomon* of Hippocrates, of Nymphodorus, Galen, &c.

GLOSSODESMUS, Frænum lingua.

GLOSSO-EPIGLOTTIC, *Glosso-epiglott'icus*. That which belongs to the tongue and epiglottis. Some anatomists have so denominated certain fleshy fibres, which pass from the base of the tongue towards the epiglottis. These muscles are more evident in some of the mammalia than in man; and their use seems to be,—to raise the

epiglottis, and to remove it farther from the glottis. Santorini, who described them after Eustachius, calls them *Retractor es Epiglottidis*.

GLOSSO-EPIGLOT'TIC LIGAMENT, *Frænum seu Frænum epiglottidis, Ligamentum membranaecum, Velamentum linguæ*. Three folds of mucous membrane, which connect the anterior surface of the epiglottis with the root of the tongue.

GLOSSOGRAPHY, *Glossographia*, from *glossō*, and *γραφη*, 'a description.' An anatomical description of the tongue.

GLOSSO-HYAL. A name given, by Geoffroy Saint-Hilaire, to the posterior cornua of the os hyoides.

GLOSSOL'OGY, *Glossolog'ia*, from *glossō*, and *λογος*, 'a treatise,' 'a discourse.' A treatise on the tongue. Nomenclature.

GLOSSOLYSIS, *Glossople'gia, Paral'ysis linguæ, P. Nervi hypoglossi*, from *glossō*, and *λυσis*, 'solution.' Paralysis of the tongue.

GLOSSOMANT'IA, *Prognos'is ex linguâ*, from *glossō*, and *μαντεia*, 'divination.' Prognosis from the state of the tongue.

GLOSSOMEGISTUS, *Glossocèle, Paraglossa*.

GLOSSON'CUS, from *glossō*, and *ογκος*, 'tumour.' *Exoncos'is linguæ*. Swelling of the tongue.

GLOSSONCUS INFLAMMATORIUS, *Glossitis*.

GLOSSO-PALATINUS, *Glossostaphylinus*—g. Pharyngeal, *Glossopharyngeus*, *Pharyngoglossal*—g. p. Nerve, *Pharyngo-glossal nerve*.

GLOSSO-PHARYNGE'US, *Glossopharynge'al*, from *glossō*, and *φαρυγξ*, 'the pharynx.' Belonging to the tongue and pharynx. Some anatomists thus designate certain fleshy bundles, which arise from the lateral parts of the base of the tongue, and are inserted into the parietes of the pharynx. They form part of the constrictor pharyngis superior.

GLOSSOPLEGIA, *Glossolysis*.

GLOSSOPTOSIS, *Glossocèle*.

GLOSSOSCIRRHUS, *Glossocarcinoma*.

GLOSSOSCOP'IA, from *glossō*, and *σκοπεω*, 'I view.' Inspection of the tongue as an index of disease.

GLOSSOSPAS'MUS, *Glossoco'na, Spasmus linguæ*, from *glossō*, and *σπασμος*, 'spasm.' Cramp or spasm of the tongue.

GLOSSOSPAT'HA, *Spat'ula pro orê*, from *glossō*, and *σπαθη*, 'spatula.' A spatula for pressing down the tongue to enable the fauces to be examined.

GLOSSO-STAPHYLINUS, from *glossō*, and *σταφυλη*, 'the uvula.' *Glossopalatinus, Palato-glossus, Constrictor Isthmi Fau'cium*. A small, thin, narrow, and long muscle, which arises from the base of the tongue, and is inserted at the inferior and lateral part of the velum palati, in the anterior pillar of which it is situate. Its use is to contract the isthmus faucium, by depressing the velum palati, and raising the base of the tongue.

GLOSSOSTERE'SIS, *Linguæ extirpa'tio*, from *glossō*, and *στερησις*, 'privation.' Extirpation of the tongue.

GLOSSOSTROPH'IA, from *glossō*, and *στροφω*, 'I turn.' Doubling of the point of the tongue upwards and backwards,—said to have been a mode of suicide. [?]

GLOSSOT'OMY, *Glossotom'ia*, from *glossō*, and *τεμνειν*, 'to cut.' Dissection of the tongue. Amputation of the tongue.

GLOSSYPERTROPH'IA, from *glossō*, and *υπερ*, 'over,' and *τροφειν*, 'to nourish.' Hypertrophy or supernutrition of the tongue.

GLOTTA, *Glossa, Tongue*.

GLOTTAGRA, *Glossalgia*.

GLOTTALGIA, *Glossalgia*.

GLOTTIS, *γλωττις*, ('the mouth-piece of a flute,') *Lig'ula*. A small oblong aperture, in the larynx, comprised between the *chordæ vocales*. It is narrow, anteriorly; wider, posteriorly; and is capable of being modified by muscular contraction, as may be required by the voice. It is by the *chordæ vocales*, that voice is produced. The glottis is nearly an inch long in the adult male; less in the female and child.

Glottis is, by some, used synonymously with ventricle of the larynx: with others, it includes the whole of the larynx.

GLOTTIS, LIPS OF THE, *Thyreo-arytenoid ligaments*—g. Spasm of the, *Asthma thymicum*.

GLOUGLOU D'UNE BOUTEILLE, *Gurguling*.

GLOUTERON, *Arctium Lappa*—g. *Petit, Xanthium*.

GLOUTIUS, *Gluteal*—g. *Maximus et extimus, Glutæus maximus*—g. *Secundus et medius, Glutæus medius*—g. *Tertius et intimus, Glutæus minimus*.

GLOUTON, *Glutton*.

GLOWWORM, *Cicindela*.

GLUANT, *Glutinous*.

GLUCOGENIA, *Glycogeny*.

GLUCOGENESIS, *Glycogeny*.

GLUCOSE, *Glycose*, from *γλυκός*, 'sweet.'

Grape, Fruit, Starch, and Honey sugar. A variety of sugar, that occurs naturally in many vegetable juices, and in honey. Compared with cane sugar, it is much less soluble in water, and less disposed to crystallize; and, when injected into the blood-vessels, does not pass off to the like extent by the kidneys. *Diabetic, Urinary, and Hepatic sugar* appear to have the same chemical composition.

GLUCOSURIA, *Diabetes mellitus*.

GLUE BONE, *Osteocolla*.

GLUE FISH, *Ichthyocolla*.

GLUMPOT, *Gallipot*.

GLUTÆUS MAGNUS, *G. major*—g. *Major, G. maximus*.

GLUTÆUS MAX'IMUS, *G. major seu magnus, Maximus et ext'imus gloutius*, (F.) *Muscle grand fessier, Ilio-sacro-fémoral; Sacro-fémoral*, (Ch.), from *γλωττος*, 'the buttock,' 'the breech.' This muscle is situate at the posterior part of the pelvis, and at the upper and posterior part of the thigh. It is large, thick, and quadrilateral; and is attached, above, to the posterior part of the crista ilii, to the part of the ilium comprised between the crista and the upper curved line, to the posterior surface of the sacrum, coccyx, and great sacro-sciatic ligament; and below, it terminates by a broad aponeurosis, which is inserted into the rugged surface that descends from the trochanter major to the linea aspera of the femur. This muscle extends the thigh on the pelvis, and rotates the thigh outwards. It is greatly concerned in station and progression.

GLUTÆUS M'ED'US, *Gloutius Secun'dus et M'edius*, (F.) *Ilio-trochantérien: Grand Ilio-trochantérien*, (Ch.), *Moyen Fessier*. This muscle is situate in part beneath the preceding; it is broad, very thick, radiated, and triangular; attached, above, to the crista ilii, and to the part of the outer surface of that bone comprised between the three anterior fourths of its crista, its upper curved line, and its lower; and below, it ends by a tendon, inserted at the upper edge of the great trochanter. It is an abductor of the thigh; but can turn the thigh outwards or inwards, according as its posterior or inferior fibres are thrown separately into contraction.

GLUTÆUS M'IN'IMUS, *G. minor, Ter'tius et Int'imus Gloutius*, (F.) *Petit Fessier; — Ilio-ischii-trochantérien, Petit Ilio-trochantérien* (Ch.)

This muscle, which is situate beneath the preceding, is flat, triangular, and with radiated fibres. It is attached, *above*, to the external surface of the os ilii, from the inferior curved line to the acetabulum; and, *below*, is inserted into the anterior part of the great trochanter. It has the same uses as the preceding.

GLUTEUS MINOR, G. minimus.

GLUTEAL, *Gloutinus*, *Glutæus*. Same etymon. (F.) *Fessier*. That which belongs or relates to the nates. This name has been given to many parts which compose the nates.

GLUTEAL APONEUROSIS. The upper and back part of the femoral fascia. In it is a remarkable opening, called the *gluteal arch*, for the passage of the gluteal vessels and nerves.

GLUTEAL ARCH, see Gluteal aponeurosis.

GLUTEAL ARTERY, *Posterior Iliac Artery*, (F.) *Artère fessière*, is one of the largest branches of the hypogastric. It makes its exit from the pelvis at the upper part of the superior sciatic foramen; gains the posterior part of the pelvis, and divides into two branches;—the one *superficial*, the other *deep-seated*. The last subdivides into three secondary branches, whose ramifications are distributed particularly to the *Glutæi*, *Longissimus Dorsi*, *Sacro-lumbalis*, &c., and anastomose with the sciatic and internal circumflex arteries.

GLUTEAL NERVE, (F.) *Nerv Fessier*, is a large branch, furnished by the 5th pair of lumbar nerves. It is chiefly distributed to the glutæi muscles.

GLUTEAL VEIN, (F.) *Veine fessière*, follows the same march as the artery of the same name.

GLUTEN, *Glutininum*, *Lentor*, 'glue, paste.' *Veg'etabile Gluten*, *Veg'etabile Casein*. An immediate principle of vegetables. It is soft, of a grayish white, viscid consistence, and very elastic. Exposed to the air, it becomes hard, brown, and fragile; and, in moist air, putrefies. Water and alcohol do not dissolve it. It is soluble in vegetable, and in weak mineral, acids, at a high temperature. The farinæ, in which it is found, are those preferred for the preparation of bread; on account of the property it has of making the paste rise. It is a compound of protein, and hence has been ranged amongst the "*proteinaceous alimentary principles*" by Dr. Pereira. By washing wheaten dough with a stream of water, the gum, sugar, starch and vegetable albumen are removed: the ductile, tenacious, elastic, gray mass left is the gluten, *common gluten*, *Beccaria's gluten*. *Pure gluten* is the soluble portion on boiling common gluten in alcohol. *Granulated gluten*, *Gluten granule*, is a paste made by adding wheat gluten to ordinary wheat. It is said to form an agreeable and nutritious aliment.

GLUTEN ARTICULORUM, Synovia—g. *Beccaria's*, see Gluten—g. Bread, see Bread, gluten—g. Common, see Gluten—g. Granulated, see Gluten—g. Granule, see Gluten—g. Pure, see Gluten.

GLUTI, Nates.

GLUTIA, Nates, Quadrigemina corpora.

GLUTIN, see Gelatin.

GLUTINANS, Agglutinant.

GLUTINATIF, Agglutinant.

GLUTINATIO, Agglutination.

GLUTINEUX, Glutinous.

GLUTINOUS, *Glutinosus*, *Collo'des*, from *gluten*, 'paste, glue.' (F.) *Glutineux*, *Gluant*. An epithet given to substances taken from the animal or vegetable kingdom, and endowed with unusual viscosity. The decoctions of marsh-mallows, and figs, and the jelly of hartshorn, are said to be glutinous.

GLUTINUM, Gluten.

GLUTITIS, from *γλουρι*, 'the nates,' and

itis, denoting inflammation. Inflammation of the nates.

GLUTOI, Nates.

GLUTOPYOSIS, from *γλουρι*, 'the nates,' and *πωσις*, 'suppuration.' Suppuration of the nates.

GLUTTON, same etymon as the next. *Gastrimargus*, *Gnathon*, *Hel'uo*, *Mando*, *Gulo'sus*, *Lurco*, *Bellygod*, (F.) *Glouton*, *Gourmand*, *Goulou*. An excessive eater.

GLUTTONY, from *glutio*, 'I swallow,' *gluttus*, 'the gullet.' *Limo'sis* *Helluo'num*, *Gastrim'us*, (F.) *Gourmandise*. Excessive appetite, owing often to habitual indulgence.

GLUTTUPATENS, Stomach.

GLUTTUS, Œsophagus.

GLUTUS, Trochanter major.

GLYCANSIS, Edulcoration.

GLYCAS'MA, from *γλυκς*, 'sweet.' A sweet wine, prepared from must.—Linden.

GLYCERATON, Glycerrhiza.

GLYCERIN, see Fat—g. Butyrate of, see Butter.

GLYCIPICROS, Solanum dulcamara.

GLYCISIDE, Pæonia.

GLYCOG'ENY, *Glycogen'ia*, *Glycogen'esie*, *Glucogen'ia*, *Glucogen'esie*, (F.) *Glycogénie*, from *γλυκς*, 'sweet,' and *γενσις*, 'generation.' Generation or formation of sugar,—as in the liver.

GLYCOSE, Glucose.

GLYCOPHYTON, Glycerrhiza.

GLYCYRRH'IZA, from *γλυκς*, 'sweet,' and *ρίζα*, 'a root.' *G. Glabra* seu *Lævis*, *Liquorit'ia* *Scyth'ica*, *Glycyra'don*, *Glycyph'yton*, *Adip'son*, *Alcacas*, *Al'mos*, *Liq'orice*, *Lic'orice*, (F.) *Réglisse*, *Bois doux*. Ord. Leguminosæ. Sex. Syst. Diadelphia Decandria. The root of this southern European plant is inodorous; has a sweet taste; is mucilaginous; and leaves, when unpeeled, a degree of bitterness in the mouth. It is used as a demulcent, and chiefly in catarrh. The extract, made from it and sold in the shops, is known under the name *Spanish Liquorice* or *Liquorice Juice*, (F.) *Jus de Réglisse*.

Pectoral Balm of Liquorice—a quack preparation—is said by Dr. Paris to consist chiefly of *Paregoric Elixir*, strongly impregnated with *Oil of Aniseed*.

GLYCYRRHŒ'A, from *γλυκς*, 'sweet,' and *ρεω*, 'I flow.' A discharge of saccharine fluid from the system.

GLYCYRRHŒA URINOSA, Diabetes mellitus.

GLYSTER HERBS, *Herbæ pro Enem'atē*. The herbs ordinarily sold by the English apothecary under this title, are:—mallow leaves, one part; chamomile flowers, one part. (3iss to Oj of water).

GNAMPISIS, Curvature.

GNAPHALUM DIOICUM, *Antennaria dioica*—g. *Plantaginifolium* seu *Plantagineum*, *Antennaria plantaginifolia*.

GNAPHALUM MARGARITACEUM, *Antennaria Margaritacea*, *Cudweed*, *Pearly Life* everlasting, from *γναφαλον*, 'a lock of wool,' owing to the floccose down of the leaves. An indigenous plant, growing in woods and fields, and flowering in August. Its virtues are not defined, and the same may be said of

GNAPHALUM POLYCEPH'ALUM; *Sweet-scented Life everlasting*, *Catfoot*. They appear to be mild astringents.

GNATHALGIA, Neuralgia maxillarum.

GNATHANCYLO'SIS, from *γναθος*, 'the jaw,' and *αγκυλωσις*, 'stiffness of joint.' Ankylosis of the lower jaw.

GNATHANTROPOLYPUS, from *gnathantrum*, 'the antrum of Highmore,' and *polypus*. Polypus of the antrum of Highmore.

GNATHITIS, *Inflamma'tio genæ*, from *γναθος*,

'the cheek, the jaw.' Inflammation of the cheek or upper jaw.

GNATHMUS, Gnathus.

GNATHOCEPHALUS, from *γναθος*, 'the jaw,' and *κεφαλή*, 'head.' A monster which has no head visible externally, but exhibits voluminous jaws.

—G. St. Hilaire.

GNATHOCYNANCHE, Cynanche parotidæa.

GNATHON, Glutton.

GNATHONEURALGIA, Neuralgia maxillaris.

GNATHOPLASTY, *Gnathoplasticæ*, from *γναθος*, 'cheek,' and *πλαστικός*, 'formative.' The formation of an artificial cheek.

GNATHOPLEGIA, *Gnathoparalysis*, from *γναθος*, and *πληγη*, 'a stroke.' Paralysis of the cheek. *Gnathoparalysis* is employed by Fuchs to signify paralysis of the lesser portion of the trifacial nerve, which supplies the muscles of mastication.

GNATHORRHAGIA, from *γναθος*, and *ρηνυμι*, 'to burst forth.' Hemorrhage from the internal surface of the cheeks.

GNATHOSPASMUS, Trismus.

GNATHUS, *Gnathus*, from *κναι*, 'I scrape, rub.' The cheek, the jaw. Also, the part of the jaws in which the teeth are fixed.—Hippocrates, *Œsius*. See *Bucca*, *Gena*, and *Maxillary Bone*.

GNESIOS, Legitimate.

GNOME, Intellect.

GNOSIS, *γνῶσις*, 'knowledge.' A common suffix, as in *Diagnosis*, *Prognosis*, &c.

GOACONAX, see *Toluifera balsamum*.

GOADBY'S SOLUTION, see *Solution*, *Goadby's*.

GOATS' BEARD, COMMON, *Tragopogon*.

GOATS' MILK, see *Milk*, *goats'*—g. *Milk*, *artificial*, see *Milk*, *goats'*—g. *Thorn*, *Astragalus verus*.

GOB, Mouth.

GOBELET ÉMÉTIQUE, *Goblet*, *emetie*.

GOBLET ÉMÉTIC, *Poculum emeticum*, *Calix vomitoria*, (F.) *Goblet émetique*. A vessel, made by pouring melted antimony into a mould. By putting wine into this and allowing it to stand some time, it acquires the property of producing vomiting. This kind of emetic has been long rejected, as the practitioner could never be certain of the dose he exhibited.

GODESBERG, MINERAL WATERS OF. These waters, at Godesberg, a German mile from Bonn, are an efficacious, acidulous chalybeate, formerly known by the name, *Draitsch Water*. They contain chloride of sodium, carbonate of soda, carbonate of lime, carbonate of magnesia, and carbonate of iron.

GODFREY'S COR'DIAL. A celebrated nostrum, for which Dr. Paris has given the following formula. Infuse $\frac{3}{4}$ of *sassafras* and of the seeds of *caraway*, *coriander*, and *anise*, each $\frac{3}{4}$, in six pints of water. Simmer the mixture until reduced to $\frac{1}{2}$ iv; then add $\frac{1}{2}$ vj of *treacle*, and boil the whole for a few minutes. When cold, add $\frac{3}{4}$ ij of *tincture of opium*. The following form is recommended by a committee of the Philadelphia college of Pharmacy. *Tinct. Opii*, Oiss; *Syrupi Nigri*, Oxxvj; *Alcoholis*, Oij; *Aquæ*, Oxxvj; *Carbonatis Potassæ*, $\frac{3}{4}$ ijss; *Olei Sassafras*, $\frac{1}{4}$ iv, M. It is anodyne.

GODRONNÉ, (CANAL), *Canalis Petitiæ* *nus* *seu Petiti*, *Couronne ciliaire*, *Canal de PETIT* ou *goudronné*, *Canal* or *Bulbular Canal* of *Petit*. *Petit* gave this name (from F.) *godron*, 'a plait or fold,' to the semicircular canal, formed by the tunica hyaloidea around the edge of the crystalline; because it appears, as it were, plaited or festooned.

GOGGLE-EYED, (Prov.) *Grozet-eyed*. Having prominent, staring and rolling eyes.

GOITRE, Bronchocele—g. Leaf, see *Laminaria*.

GOITRE STICKS. In South America the stems of a seaweed are so called, because they are chewed by the inhabitants where goitre prevails.—Royle.

GOITROUS, (F.) *Goitreux*. Relating or appertaining to goitre. One affected by goitre or Bronchocele, —*Goitred*.

GOLD, *Aurum*, *Chryso*, *Sol*, *Res metallosum*, (F.) Or. (Sax.) *golo*. A solid, yellow, very brilliant, hard, very ductile, malleable, tenacious, and heavy metal; found in nature, either in its native state, or combined with a little silver, copper, or iron. S. g. 19.25.

Muriate or *Chloride of Gold*, *Auri Chloridum* *seu Terchloridum* *seu Murias* *seu Chloruretum*, *Aurum Muriatum* *seu Chloratum* *seu Oxydulatum* *seu muriatum* *seu Salatum*, (F.) *Chlorure* ou *Muriate* ou *Hydro-chlorate d'or* has been admitted into the Pharmacopœia of the United States, and into that of Paris, &c. The formulæ, however, differ. That of the United States is a muriate with two bases; and is prepared, according to the form of Dr. Christien, by dissolving the gold in a mixture of nitric and muriatic acids, and adding chloride of sodium to the residuum after evaporation; then redissolving and evaporating slowly to dryness. The Parisian formula for the *Muriate d'or*, *Murias* *seu Chloruretum Auri*, consists in simply dissolving the gold in the acids, and evaporating to dryness. It has been recommended as an antisyphilitic in old, rebellious, venereal affections, exostoses, and in venereal, scrofulous or cancerous glandular enlargements. Dose, gr. 1–8th to gr. ss, rubbed on the tongue or gums. Internally, one-sixteenth of a grain, in pills.

Various other preparations, as the *Cy'anide* or *Tercy'anide*, (*Auri Cy'anidum* *seu Cyanuretum* *seu Tercyanidum*, (F.) *Cyanure d'or*;) the metallic gold in a state of division (*Aurum metallicum*, *Pulvis Auri*, (F.) *Or divisé* ou *métallique*,) obtained by amalgamating gold with mercury and driving the latter off by heat; and in the form of filings (*Aurum limatum*); the *Chloride of Gold* and *Sodium*, (*Aurum muriatum* *natronatum* *seu muriatum* *seu chloratum* *natronatum*, *Chloreum Auricum* *Chlore* *natrui*, *Murias Aurico-natricum*, *Chloruretum auri* *et natrui*, *Sol'dii auro-terchloridum*, *Hydrochlorate* *ou* *muriate of Gold* and *Soda*, (F.) *Chlorure d'or* *et de Soda*, *Hydrochlorate* *ou* *muriate d'or* *et de Soude*); the *Nitromuriate of Gold*, (*Aurum Nitro-muriatatum*, *Auri nitromurias*, (F.) *Nitromuriate d'or*;) the *Oxide of Gold*, (*Auri Oxidum*, *Aurum Oxidatum*, *Auri teroxidum*, *Peroxide of gold*, *Auric acid*, (F.) *Oxide d'or*;) and the *Iodide of Gold*, (*Auri Iodidum* *seu Ioduretum*, (F.) *Iodure d'or*,) have been employed in the like affections, and with similar results.

GOLD-BEATER'S SKIN. The intestina recta of the ox, which have been beaten quite smooth, for the manufacture of gold leaf. Used as a defensive dressing for slight cuts, &c.

GOLD, CHLORIDE OF, see *Gold*—g. *Cyanide* of, see *Gold*—g. *Hydrochlorate* of, see *Gold*—g. *Iodide* of, see *Gold*—g. *Muriate* of, see *Gold*—g. *Nitro-muriate* of, see *Gold*—g. *Oxide* of, see *Gold*—g. *Peroxide* of, see *Gold*—g. *Tercy'anide* of, see *Gold*—g. and *Sodium*, *Chloride* of *oxide* of, see *Gold*—g. and *Soda*, *hydrochlorate* of, see *Gold*—g. and *Soda*, *muriate* of, see *Gold*.

GOLD LEAF, *Gold Foil*, *Aurum foliatum*, *Aurum in libellis*. Used to gild pills and to plug carious teeth. The gold leaf or foil, used by the dentist for filling teeth, is much thicker than ordinary gold leaf.

GOLD THREAD, *Coptis*.

GOLDEN BUG, *Coccinella septempunctata*—*g.* Rod. *Knop*, *Coccinella septempunctata*—*g.* Rod. *Solidago virgaurea*—*g.* Rod. fragrant. *Solidago odora*—*g.* Rod. rigid, *Solidago rigida*—*g.* Rod. sweet-scented, *Solidago odora*—*g.* Seal, see *Calumba*, *Hydrastis Canadensis*.

GOLDENS, *Chrysanthemum leucanthemum*.

GOLDING, *Calendula officinalis*.

GOLDSHRUB, *Palicourea speciosa*.

GOLDWASSER, see Spirit.

GOLFE, Sinus—*g. de la Veine jugulaire*, see Jugular veins.

GOLUNCHA, *Menispermum cordifolium*.

GOMBO, *Hibiscus esculentus*.

GOMME, Gumma, Gummi—*g.* *Adragant*, *Tragacanth*—*g.* *Ammoniacque*, *Ammoniac gum*—*g.* *Arabique*, *Acacia gummi*—*g.* *Astringente de Gambie*, see *Butea frondosa*—*g.* *Carayne*, *Caranna*—*g.* *Carane*, *Caranna*—*g.* *Elastique*, *Caoutchouc*—*g.* *de Gaïac*, *Guaiacum*—*g.* *de Générrier*, *Sandarac*—*g.* *Gettania*, *Gutta Percha*—*g.* *Gutte*, *Cambogia*—*g.* *de Kino*, *Kino*—*g.* *Laque*, *Lacca*—*g.* *de Lierre*, see *Hedera helix*—*g.* *du Pays*, *Gummi nostras*—*g.* *Séraphique*, *Sagapenum*.

GOMPHIASIS, *Gomphias'mus*, from *γομφος*, 'a nail.' A disease of the teeth, and particularly of the molares; looseness of the teeth in their sockets. *Agomphia'sis*. Pain in the teeth. *Odontalgia*.

GOMPHIASMUS, *Gomphiasis*.

GOMPHIOL, Molar teeth.

GOMPHOCARPUS CRISPUS. A South African plant, the root of which, formerly known to the Dutch apothecaries as *Radix Asclepi'adis crispæ*, is extremely bitter and acrid; and, on account of its diuretic virtues, a decoction or infusion of it has been advised in various kinds of dropsy. A tincture of it is said to be valuable in colic.

GOMPHOMA, *Gomphosis*.

GOMPHO'SIS, *Cardinamen'tum*, *Clava'tio*, *Gompho'ma*, *Coagmenta'tio*, *Inclava'tio*, *Incuneat'tio*, from *γομφος*, 'a nail.' *Engompho'sis*. An immovable articulation, in which one bone is received into another, like a nail or peg into its hole. *Gomphosis* is only met with in the articulations of the teeth with the alveoli. It is, also, called *Articulation par implantation*.

GOMPHUS, *Clavus*.

GOMUTO PALM, see *Toddy*.

GOMUTOS GOMUTO, see *Toddy*.

GONACRASIA, *Spermatorrhœa*.

GONACRATIA, *Spermatorrhœa*.

GON'AGRA, *Gon'yagra*, from *γονν*, 'the knee,' and *agya*, 'a prey.' "That which attacks the knees." Gout or rheumatism in the knees. *Gongorrheu'na*. Paracelsus calls it *Gen'ugra*.

GONAL'GIA, from *γονν*, 'the knee,' and *αλγος*, 'pain.' Pain in the knee. *Gonyal'gia*. This is almost always produced by gout. It may, however, depend on some other disease, either of the knee or of another part—particularly of the hip-joint.

GONARTHRI'TIS, from *γονν*, 'the knee,' *αρθρον*, 'joint,' and *itis*, denoting inflammation. Inflammation of the knee-joint.

GONARTHROCACE, *Gonocace*.

GONAURA, see *Sperm*.

GONDOLE OCULAIRE, *Scaphium oculare*.

GONDRET'S AMMONIACAL CAUSTIC, *Pommade de Gondret*—*g.* Counter-irritant, *Pommade de Gondret*.

GONE, *Gonos*, *Genitu'ra*. The semen; (hence, *gonorrhœa*)—the uterus, offspring.—*Hippocrates*. See *Epigone*, and *Generation*.

GONECYSTIDES, *Vesiculæ seminales*.

GONECYSTITIS, *Inflammat'io vesiculæ'rum semina'tium*, from *γονν*, 'sperm,' *κυστις*, 'bladder,'

and *itis*, denoting inflammation. Inflammation of the vesiculæ seminales.

GONEPEUS, *Spermatopeus*.

GONEPOETICUS, *Spermatopeus*.

GONFLEMENT, Swelling.

GONGRONA, *Bronchocele*.

GONGROPHTHISIS, *Phthisis pulmonalis*.

GONGYLIDIUM, *Pilula*.

GONGYLION, *Pilula*.

GONGYLIUS, *Pilula*.

GONIOCATH'ETER, from *γωνια*, 'an angle,' and *καθετηρ*, 'a catheter.' An angular or crooked catheter.

GONIOME'LE, from *γωνια*, 'an angle,' and *μηλη*, 'a probe.' An angular or crooked probe or sound.

GONIOM'ETER, *Goniom'etrum*, from *γωνια*, 'an angle,' and *μετρον*, 'a measure.' An instrument for measuring angles.

A 'FACIAL GONIOMETER' has been invented by Mr. Turpenney, of Philadelphia, which is well adapted for measuring the facial angle.

GONOBOLIA, Ejaculation, spermatic.

GONOC'ACE, *Gonarthritis'acis*, from *γονν*, 'the knee,' and *κακος*, 'evil.' *Tumor genui albus*. White swelling of the knee. *Hydrarthrus*.

GONOCELE, *Spermatocele*.

GONOIDES, from *γονν*, 'sperm,' and *ειδος*, 'appearance.' *Genitu'ra sim'ilitis*. Similar to sperm. *Sperm'atoid*, *Spermatoï'des*, *Spermato'des*. A term appropriated to any substance which resembles sperm.

GONOL'OBUS MACROPHYL'LUS, (*γωνια*, 'an angle,' and *λοφος*, 'a pod'); indigenous. *Nat. Ord.* *Asclepiadaceæ*. The root is cathartic. It is said to furnish a juice with which the Indians poison their arrows. [?]

GONOPOETICUS, *Spermatopeus*.

GONORRHÉE BATARDE, *Gonorrhœa spuria*.

GONORRHO-BLEPHARRHŒA, *Ophthalmia*, *gonorrhœal*.

GONORRHO-PROSTATITIS, see *Prostatitis*.

GONORRHŒA. Erroneously called from *γονν*, 'sperm,' and *ρœω*, 'I flow,' because the older writers believed it to be a flux of semen. *Blennorrhag'ia*, *Blennorrhœa*, *B. urethral'is*, *Blennure'thria*, *Phallorrhœa*, *Medorrhœa*, *M. viri'tis*, *Catar'rhus gonorrhœa* seu *ure'thræ*, *Ure'thritis*, *Inflammat'io ure'thræ*, *Urethral'gia*, *Proflu'vium mucosum ure'thræ*, *Catarrhus urethral'is*, (*F.*) *Écoulement*, *Urétrite*, *Urétrite*. An inflammatory discharge of mucus from the membrane of the urethra in both sexes; and from that of the prepuce in man, and the vagina in woman. It may be excited spontaneously, or by irritants applied directly to the membrane; but is usually produced by impure connexion. Two great varieties have been generally reckoned.—1. *GONORRHŒA PURA* SEU *BENIGNA*. That which does not follow an impure connexion; (*F.*) *Échauffement*, *Blennorrhag'ia benigna*, *Caulorrhœa benigna*, *Catar'rhus Ure'thræ*, *Gonorrhœa catarrhal'is* seu non contagio'sa:—and, 2. *GONORRHŒA IMPURA*, *maligna*, *contagio'sa*, *syphilit'ica*, *et virulenta*; *Fluor albus malignus*, *Fluor venereus*, *Blennorrhœa luo'des*, *Baptorrhœa*, *Myxio'sis*, *Lues gonorrhœica*, *Seroph'ulæ gonorrhœica*, *Tuber'cula gonorrhœica*, *Clap*, (*Old E.*) *Chawdpye*, (*F.*) *Chaudepisse*; that which is the result of impure commerce. The French, also, distinguish the *Chaudepisse sèche*, or that unaccompanied with discharge; and the *Chaudepisse chordée*, *Gonorrhœa cordata*, or that accompanied with chordee, and which, of course, occurs only in the male. It is the kind that most frequently engages the attention of the practitioner, and is characterized by mucous discharge from

the urethra or vagina, intermixed with specific matter, and accompanied with burning pain on micturition. It is decidedly infectious. It is, however, a distinct disease from syphilis, and never produces it. Its duration is various, but the inflammatory symptoms usually subside in four or five weeks; leaving generally behind more or less of the gonorrhœa mucosa or gleet. Gonorrhœa of every kind, attended with any inflammatory symptoms, is best treated by the antiphlogistic regimen; avoiding every kind of irritation, and keeping the body cool by small doses of salts, and the urine diluted by the mildest fluids. After the inflammatory symptoms have subsided, cubebs, or the balsam of copaiba, exhibited in the dose of a teaspoonful, three times a day, will be found effectual: indeed, during the existence of the inflammatory symptoms, it often affords decided relief. Injections are rarely required.

Sometimes, gonorrhœa affects the glans; when it is called *Gonorrhœa Spuria* seu *Balani*, *Balanoblennorrhœa*, *Balanorrhœa*, *Balani* 'tis, *Blennorrhag* 'ia *spuria* seu *notha*, (F.) *Blennorrhagie du gland*, *Gonorrhée bâtarde*, *Fausse Blennorrhagie*. It requires only cleanliness and cooling lotions. This is the simple form of balanitis. It may, however, be of the impure kind—*Balanorrhag* 'ia, (F.) *Balanite contagieuse*.

Some other varieties of gonorrhœa have been enumerated, but they are of little moment.

In consequence of repeated attacks of gonorrhœa, or of the debility induced by a single attack, it not unfrequently happens, as already remarked, that a constant, small discharge occurs, or remains behind, after all danger of infection is removed. The great difference between it and gonorrhœa is, that it is uninfected. The discharge consists of globular particles, contained in a slimy mucus, and is generally devoid of that yellow colour which characterizes the discharge of gonorrhœa virulenta. It is unattended with pain, scalding, &c. To this state the names *Gleet*, *Gonorrhœa mucosa*, *Blennorrhœa chronica*, *Blennorrhœa*, &c., have been given. It is commonly a disease of some duration, and demands the use of the copaiba, astringent injection; and, if obstinate, the introduction of the bougie.

GONORRHEA BALANI, *G. spuria*—*g.* Benigna, Leucorrhœa—*g.* Catarrhalis, see *Gonorrhœa*—*g.* Chordata, Chordee—*g.* Contagiosa, see *Gonorrhœa*.

GONORRHEA DORMIENTIUM, *G. Oneirogonos*. The seminal discharge which occurs during sleep, and is occasioned by libidinous dreams. See *Pollution*.

GONORRHEA IMPURA, see *Gonorrhœa*.

GONORRHEA LAXORUM, *G. libidinosæ*, *Spermatorrhœa atonica*, consists of a pellucid discharge from the urethra, whilst awake, without erection of the penis, but with venereal thoughts.

GONORRHEA LAXORUM, *Pollution*, *G. libidinosa*—*g.* Maligna, see *Gonorrhœa*—*g.* Mucosa (gleet), see *Gonorrhœa*—*g.* Noncontagiosa, *G. pura*—*g.* Notha inveterata, *Leucorrhœa*—*g.* Oneirogonos, *G. dormientium*, *Pollution*—*g.* Spuria, see *Gonorrhœa*—*g.* Syphilitica, see *Gonorrhœa*—*g.* Vera, *Pollution*, *Spermatorrhœa*—*g.* Virulenta, see *Gonorrhœa*.

GONOS, *Gone*.

GONOSTROMA, *Proligerous disc*.

GONOZEMIA, *Spermatorrhœa*.

GONY, *yovv*, *Genu*, 'the knee;' hence:

GONYAGRA, *Gonagra*.

GONYALGIA, *Gonalgia*.

GONYANCON, *Gonycampsis*, from *yovv*, 'the knee,' and *αγκων*, 'a curvature.' A curvature or bending of the knee. A bent knee.

GONYCAMPISIS, *Gonyancœon*.

GONYCROTUS, from *γόνυ*, 'the knee,' and *κρῶν*, 'I strike.' One who is knock-kneed, or in-kneed. See *Entogonyancœon*.

GONYONCUS, from *yovv*, 'the knee,' and *ογκος*, 'a tumour.' A swelling of the knee.

GONYORRHEUMA, *Gonagra*.

GOOD KING HENRY, *Chenopodium bonus Henricus*.

GOODYE'RA PUBES'CENS, *Tussa'ca reticulata*, *Satyrium*, *Neottia*, *Rattlesnake leaf*, *Rattlesnake Plantain*, *Networt*, *Nettleaf*, *Scrofula weed*, *Cancer weed*. Ord. *Orchidaceæ*. An indigenous plant, used empirically in scrofula—the fresh leaves being applied to the sores. It is employed by the Indians.

GOOSEBERRY, *Ribes uva crispa*.

GOOSEFOOT, *Chenopodium anthelminticum*—*g.* Angular-leaved, *Chenopodium bonus Henricus*—*g.* Mercury, *Chenopodium bonus Henricus*

—*g.* Stinking, *Chenopodium vulvaria*.

GOOSEGRASS, *Galium aparine*, *G. verum*.

GOOSESHARE, *Galium aparine*.

GOOSESKIN, *Horrida cutis*.

GORBELLY, *Physconia*.

GORDIUS MEDINENSIS, *Dracunculus*.

GORGE, *Ingluvies*, *Pharynx*, *Throat*—*g.* *Grosse*, *Bronchocele*—*g.* *Mal de*, *Cynanche*.

GORGET, *Gorget*—*g.* *Lithotome*, *Gorget*, *lithotomy*—*g.* *à Repoussoir*, see *Gorget*.

GORGET, from (F.) *gorge*, 'the throat.' *Canalis canaliculatus*, *Ductor canaliculatus*, (F.) *Gorget*. An instrument representing a long gutter, in the shape of a throat, which is especially employed in the operations of lithotomy and fistula in ano.

GORGET, CUTTING, see *Gorget*, and *Lithotomy*.

GORGET, LITHOTOMY, (F.) *Gorget Lithotome*, *Cutting Gorget*, is the one used in the operation for the stone, for the purpose of dividing the prostate and the neck of the bladder, so as to enable the surgeon to introduce the forceps and extract the stone. At the end of this gorget is a crest or beak, which fits the groove of the staff, and admits of the gorget being passed along it into the bladder. Besides cutting, there are also blunt gorgets, intended to be introduced into the wound—their concavity serving as a guide for the forceps into the bladder.

The chief modifications in the gorget have been made by Andouillet, Bell, Blicke, Bromfield, Cline, Desault, Foubert, Hawkins, Larrey, Lefèvre, Michaelis, Thomas, &c.

GORGET FOR FISTULA IN ANO consists of a semi-cylindrical wooden staff, four inches long, without including the handle, and furnished with a wide groove. This is introduced into the rectum, to prevent the point of the bistoury from injuring the intestine, when the internal orifice of the fistula is deeply situate, and it is desirable to perform the operation by incision. This instrument, invented by Marchetti, has been modified by Percy, Runge, &c.

Desault invented an instrument for conducting the wire by the anus, in the operation for fistula by ligature. He called it *Gorget à repoussoir*.

GORGONEI FONTES. Fountains described by Libavius as containing water which possessed a petrifying property; probably, water holding in solution supercarbonate of lime.

GORGOSSET, *Pyrosis*.

GOSIER, *Pharynx*, *Throat*.

GOSSUM, *Bronchocele*.

GOSSYP'IIUM, *Gossypium Herba'ceum*, *Gossipion* *Xylon*, *Xylum*, *Bombac*, *Cotton*, (F.) *Coton*. Family, *Malvaceæ*. Sex. Syst. *Monadelphia Polyandria*. The seeds of the Cotton Tree, *Gossypium arbor'ceum*, have been administered in coughs, on account of the mucilage they contain, and a decoction of a pint of the seeds to a quart

of water has been given, in the Southern States, in intermittents. A decoction of the root has been prescribed as an emmenagogue and parturient. The cotton wool is used in medicine for making moxas, &c., and is applied in the treatment of recent burns and scalds.

GOTCHBELLY, Physconia.

GOTIUM, Bronchocele.

GOUDRON, see Pinus sylvestris.

GOUFF, Arum maculatum.

GOÛTRE, Bronchocele.

GOULARD'S LOTION, see Lotion, Goulard's.

GOULARD WATER, Liquor plumbi subacetatis dilutus.

GOULU, Glutton.

GROUND OF THE EYE, Chassie.

GOULD, Cucurbita—g. Bitter, Cucumis coloyanthis—g. Bottle, Cucurbita pepo.

GOULD WORM, Distoma hepaticum.

GOURLANDISE, Gluttony.

GOURME, Porrigio larvalis. Vulgarly, in France, any cutaneous eruption.

GOUSSE, Legumen.

GOUT, Arthrit'is, Arth'ragra, Arthral'gia, Morbus domino'rum seu articu'laris, Malum articulo'rum, Gutta, Arthro'ia Pod'agra, Podal'gia, Pod'agra, Arthrit'icus verus, Arthrit'is Podagra, Podagra Arthrit'is, Flux'io arthrit'ica, Febris arthrit'ica seu Podag'rica, Arthrodyn'ia podag'rica, Cauma podag'ricum, Flerecin, Gastroarthrit'is, Misopto'chos, (Sc.) Gat, (F.) Goutte, Agrie.—Piorry. The gout was formerly regarded as a catarrh, and received its name from (F.) goutte, (L.) gutta, 'a drop,' because it was believed to be produced by a liquid, which distilled, goutte à goutte, 'drop by drop,' on the diseased part. This name, which seems to have been first used about the year 1270, has been admitted into the different languages of Europe. Gout is an inflammation of the fibrous and ligamentous parts of the joints. It almost always attacks, first, the great toe; whence it passes to the other smaller joints, after having produced, or been attended with, various sympathetic phenomena, particularly in the digestive organs: after this, it may attack the greater articulations. It is an affection which is extremely fugitive, and variable in its recurrence. It may be acquired or hereditary. In the former case, it rarely appears before the age of thirty-five; in the latter, it is frequently observed earlier. It is often difficult to distinguish it from rheumatism. A combination is, indeed, supposed to exist sometimes; hence called *Rheumatic gout*. During the paroxysm or fit, a burning, lancinating pain is experienced in the affected joint, attended with tumefaction, tension, and redness. One or more joints may be attacked, either at the same time or in succession; and, in either case, the attack terminates by resolution in a few days. This is the Arthrit'is acu'ta seu inflammato'ria seu regula'ris, Regular gout, Arthro'ia pod'agra regula'ris, Arth'ragra legit'ima seu vera seu genu'i'na seu norma'lis, (F.) Goutte régulière chaude. At other times, pains in the joints exist, of more or less acute character; the swelling being without redness. These pains persist, augment, and diminish irregularly, without exhibiting intermission, and, consequently, without having distinct paroxysms. The disease is then called *aton'ic*, *asthen'ic*, *imperfect* or *irregular* gout, Chronic G., Arthrit'is aton'ica seu asthen'ica, Arthro'ia Podagra larva'ta, Dyarthrit'is. It is, also, commonly called in France *Goutte froide*, *Goutte blanche*. It may appear primarily, or succeed attacks of regular gout.

Gout does not always confine itself to the joints. It may attack the internal organs: when it is called *Arthrit'is aber'rans* seu *errat'ica* seu *planot'ica*, Arth'ragra anom'ala, Pod'agra aber'rans,

Vare'ni, Wandering, misplaced, or anomalous gout, (F.) Goutte vague.

Ret'rograde gout, Arthritis retrograda, Podagra retrograda, Arthro'ia Podagra complicata, Rec'e'dent or m'iepla'ced gout, (F.) Goutte remontée, G. malplacée, G. rentrée, is when it leaves the joints suddenly and attacks some internal organ, as the stomach, intestines, lungs, brain, &c.

Gout is also called, according to the part it may affect, *Podagra*, *Gonagra*, *Chiragra*, &c. It may be acute or chronic, and may give rise to concretions, which are chiefly composed of urate of soda. See Calculus, (arthritis.) It may, also, give occasion to nodosities, when it is called *Arthritis nodo'sa*, (F.) Goutte nouée.

The treatment is of the antiphlogistic kind, and the local disorder should be but little interfered with. Colchicum seems to have great power over the disease. It forms the basis of the *Eau médicinale d'Husson*, a celebrated French gout-remedy. The bowels must be kept regular by rhubarb and magnesia; and a recurrence of the disease be prevented by abstemious habits.

GOUT, DIAPHRAGMATIC, Angina pectoris.

GOUT PAPER, so called, Charta antiarthrit'ica, Charta antirheumat'ica, is made by spreading a very thin layer of a mixture of an ethereal or spirituous extract of the bark of mezereum root, with wax, spermaceti, and oil, over the surface of paper.

GOUT, RHEUMATIC, see Rheumatism, acute—g. Weed, Ligusticum podagraria.

GOUT STONES, Calculi, arthritic.

GOUT, Taste.

GOUTTE, Gout, Gutta—g. Blanche, Gou' (atonic)—g. Caduque, Epilepsy—g. Froide, Gout (atonic)—g. Malplacée, Gout (retrograde)—g. Nouée, Gout (with nodosities)—g. Régulière chaude, Gout (regular)—g. Remontée, Gout (retrograde)—g. Rentrée, Gout (retrograde)—g. Rose, Gutta rosea—g. Sciatique, Neuralgia femoro-poplitea—g. Sereine, Amaurosis—g. Vague, Gout (wandering).

GOUTTEUX, Arthritic, Podagric.

GOUTTIÈRE (F.), Collic'ie. A gutter in a bone, like that used for carrying off rain. Some of these cavities are intended to facilitate the sliding of tendons, such as the *Gouttière Bicipitale* or *Bicipital groove*. Others, as the *Gouttière sagittale* or *Sagittal groove*, lodge blood-vessels and especially veins. Others, again, are merely intended for the support of certain organs; as the *Gouttière basilaire* or *Basillary fossa*, which supports the medulla oblongata.

GOUTTIÈRE BASILAIRE, see Gouttière—g. Bicipitale; Bicipital groove—g. Lacrymale, Lachrymal groove—g. Sacré, Sacral groove—g. Sagittale, see Gouttière.

GOUTY, Arthritic, Podagric—g. Rheumatism, see Rheumatism, acute.

GOVERNAIL DU TESTICULE, Gubernaculum testis.

GOWAN, EWE, Bellis—g. Yellow, Ranunculus.

GOWLAND'S LOTION, see Lotion, Gowland's.

GOWN, RED, Strophulus—g. Yellow, Icterus infantum.

GOWNE OF THE EYE, Chassie.

GRACE-WIFE, Midwife.

GRACILARIA LICHENOÏDES, Fucus amy-laceus.

GRAC'ILIS, Macer, Macilent'us. Slender, lean. Also, the slender Rectus int'rior fem'oris seu Grac'ilis int'ior, (F.) Sous-pubio-crétitibial, Sous-pubio-prétibial (Ch.), Droit ou grêle interne de la cuisse. This muscle is situate at the inside of the thigh. It is thin and very long; and

arises from the descending ramus of the pubis. to be inserted at the inner and inferior part of the tuberosity of the tibia. It bends the leg and causes adduction of the thigh. See *Grêle*.

GRACILIS ANTERIOR, Rectus femoris.

GRADUATE, *Graduatus*, from *gradus*, 'a step,' 'a degree.' In medicine, one who has attained a degree,—usually, the degree of doctor.

GRÆA, *γραια*. The pellicle which forms on milk. The folds of skin round the umbilicus. An old woman.

GRAIN, *Granium*; the 60th part of a Troy, and the 72d part of a *Poids de marc* drachm.

GRAIN OIL, Oil, fusel —g. Oily, Sesamum orientale.

GRAINE D'ASPIG, see Phalaris Canariensis —g. d'Écarlate, Coccus cacti—g. *Musc*, Hibiscus abelmoschus —g. de Paradis, Amomum granum paradisi —g. de Perroquet, Carthamus tinctorius (seed) —g. de Santé, see *Pilulæ aloes et kinæ* —g. de Turquie, Zea mays.

GRAINS, GUINEA, see Pepper, Malagueta.

GRAINS OF PARADISE, Amomum granum Paradisi.

GRAISSE, Pinguedo —g. de Mouton, Sebum —g. d'Oie, Adeps asnerina —g. *Oxygénée*, Unguentum acidi nitrosi —g. de Porc, Adeps præparatus.

GRAMEN ÆGYPTIACUM, G. Crucis cyprioidis—g. Caninum, Triticum repens.

GRAMEN CRUCIS CYPRIODIS, *Gramen Ægyptiacum*, *Ægyptian Cock's foot grass*. Ord. Graminæ. The roots and plants possess the virtues of the *Triticum repens*, and have been recommended in the earlier stages of dropsy. They were, formerly, considered to possess many other properties.

GRAMEN DIOSCORIDIS, Triticum repens—g. Major, Sarsaparilla Germanica—g. Orientale, Juncus odoratus —g. Repens, Triticum repens —g. Rubrum, Sarsaparilla Germanica.

GRAMIA, CHASSIE, Lippitudo.

GRAMINIVOROUS, *Graminivorus*, from *gramen*, 'grass,' and *voro*, 'I eat.' Feeding or subsisting on grass.

GRAMMARUM, Scruple.

GRAMME, *γρᾶμμα*. An ancient weight, equivalent to the 24th part of an ounce, or to 24 grains, or a scruple, avoirdupois. At the present day, the gramme is equal in weight to a cubed centimètre of water; or to 18 grains, poids de marc—15.434 grains, Troy.

GRAMME, Iris, Line.

GRAMMITIS AUREA, Asplenium ceterach—g. Ceterach, Asplenium ceterach.

GRANA, Hemicrania—g. Guineensis, Pepper, malagueta—g. Molucca, Croton tiglium—g. Moschi, Hibiscus abelmoschus —g. Orientis, see *Menispermum cocculus*—g. Paradisi, Amomum granum Paradisi —g. Tiglii seu Tiglia, see *Croton tiglium*—g. Tili, Croton tiglium—g. Triteæ, see *Hyoides os*.

GRANADILLA, APPLE-SHAPED, *Passiflora maliformis*.

GRANATI RADICIS CORTEX, see *Punica granatum*.

GRAND DORSAL, Latissimus dorsi.

GRANDEB'ALÆ, The hair which grows in the arm-pits.

GRANDGORE, Syphilis.

GRANDINOSUM OS, Cuboid.

GRANDO, Chazaza.

GRANDULA, Gland.

GRANIVOROUS, *Granivorus*, from *granum*, 'a grain,' and *voro*, 'I eat.' Feeding or subsisting on grain or seeds.

GRANTRISTUM, Anthrax.

GRANULA SANGUINIS, Globules of the blood—g. Seminis, see *Sperm*.

GRANULAR, *Granulosus*, *Granulosus*, (F.) *Granulé*, *Granuleux*, from *granulum*, 'a small grain.' Composed of or containing granules or small particles,—as 'granular eyelid,' 'membrana granulosa,' &c., &c.

GRANULAR DEGENERATION or DISORGANIZATION OF THE KIDNEY, Kidney, Bright's disease of the —g. Conjunctiva, Trachoma—g. Eyelid, Trachoma—g. Layer, Membrana granulosa —g. Liver, Cirrhosis—g. Tin, see Tin.

GRANULATED LIVER, Cirrhosis.

GRANULATION, *Granulatio*, from *granum*, 'a grain.' Gemma, (F.) *Bourgeon*, *B. charnu*. Granulations are the reddish, conical, flesh-like shoots, which form at the surface of suppurating wounds and ulcers. They are the product of inflammatory excitement, and may be produced in indolent ulcers, by exciting the parts by proper stimulants. They form the basis of the cicatrix.

GRANULATION is, likewise, a name given by the modern French physicians to an organic lesion, consisting in the formation of small, round, firm, shining, semi-transparent tumours, of the size and shape of millet-seed, or of a pea; which are met with in the lungs particularly, and in considerable quantity; often without materially interfering with their functions.

In pharmacy, granulation is a process by which a metal is reduced to fine grains, by melting it, and causing it, whilst liquid, to pass through a kind of sieve into a vessel of water,—as in the making of shot:—or by shaking or rubbing the melted metal in an appropriate box or vessel,—as in the formation of granular tin or granulated zinc.

GRANULATIONS CÉRÉBRALES, Glandulæ Pacchioni.

GRANULATIONS MILITARY, or *Miliary tubercles*, are the small, transparent grains, of variable size, from that of a millet-seed to that of a grain of hemp, which are presumed to be the primitive state of tubercles.

GRANULE, *Granulum*; diminutive of *granum*, 'a grain.' A small grain; a small compact particle; a cytoblast.

GRANULES SEMINAL, *Granula seminis*. Minute, rounded, granulated bodies, observable in the semen, which are, in all cases, much less numerous than the spermatozooids. See *Sperm*.

GRANULEUX, Granular.

GRANULOUS, Granular.

GRANVILLE'S LOTION, see Lotion, Granville's counter-irritant.

GRAPE, see *Vitis vinifera*—g. Sea-side, see Kino.

GRAPE-CURE, (F.) *Cure de raisin*, (G.) Traubenkur, Weintrau. A mode of medication in Germany, which consists in the use of the grape for both meat and drink; nothing more at the farthest being allowed than a piece of dry bread. This diet is continued for weeks. Its effects are altogether revellent, and resemble in many respects those of hydropathy.

GRAPES, DRIED, *Uvæ passæ*.

GRAPHIDOIDES, Styloid.

GRAPHIOIDES, Styloid.

GRAPHIOIDES, Styloid.

GRAPHIS'CUS, *Graphis'cus Di'oclis*. An instrument invented by Diocles for extracting darts [?]. Said, in many of the Dictionaries, but erroneously, to be described by Celsus. He speaks only of the *Cyathis'cus Di'oclis*, *Κυαθισκων*. See *Cyathiscus*.

GRAPHITES, *Plumbago*, *Supercarbure'tum seu Carbure'tum Ferri*, *Ferri Carbonat'um seu Supercarbure'tum*, *Carbo minéralis*, *Galène*, *Gale'ne*, *Plumbum nigrum*, *Cerussa nigra*, *Graphite*, *Carburet of iron*, *Black lead*, *Wad*, (F.)

Crayon noir, Plombagine, Charbon minéral, Carburé de fer, from γράφω, 'I write.' This substance has been esteemed slightly astringent and desiccative. It has been advised by Weinhold in the cure of herpes.

GRAPHOIDES, Styloid.

GRAPHOSPASMUS, Cramp, writers'.

GRAS DES CADAVRES, Adipocire—*g. de Jambe*, Sura—*g. des Cimetières*, Adipocire—*g. de Jambe*, Sura.

GRAS FONDURE (F.), *Diarrhœa adiposa*, literally, *molten grease*. A species of diarrhœa, referred to by old writers; accompanied with great emaciation, and in which the evacuations contain fat-like matter. According to Sauvages, the *Gras-fondure* differs from colliquative diarrhœa in not being attended with hectic fever.

GRASS, Asparagus—*g. Bear's*, *Yucca filamentosa*—*g. Bermuda*, *Cynodon dactylon*—*g. Bitter*, *Aletris farinosa*—*g. Blue-eyed*, *Sisyrinchium Bermudianum*—*g. Brome*, *Bromus ciliatus*—*g. Brome*, soft, *Bromus ciliatus*—*g. Canary*, cultivated, *Phalaris Canariensis*—*g. Couch*, *Triticum repens*—*g. Dog*, *Triticum repens*—*g. Egyptian cock's foot*, *Gramen crucis cyperioïdes*—*g. Goat's*, *Scorzonera*—*g. Knot*, *Polygonum aviculare*—*g. Lily*, *Sisyrinchium Bermudianum*.

GRASS OIL OF NEMAU, *Oleum graminis Indici*.

GRASS, PHYSIC, *Sisyrinchium Bermudianum*—*g. Rib*, *Plantago lanceolata*—*g. Scurvy*, *Sisyrinchium Bermudianum*—*g. Silk*, *Yucca filamentosa*—*g. Sweet*, *Acorus Calamus*.

GRASS TREE, *Xanthorrhœa*.

GRASS, VIPERS', *Scorzonera*—*g. Yellow-eyed*, *Xyris bulbosa*.

GRASSET (F.). The anterior region of the thigh, bounded below by the patella.

GRASSEYEMENT (F.), *Sonus blæsus*, *Rotacismus*, from (F.) *gras*, 'thick.' 'Speaking thick.' According to Sauvages, a vicious pronunciation of the letter *r*. They who speak thick, like the inhabitants of Newcastle, in England, or of Havre, in France, have difficulty in pronouncing the *r*, and they frequently substitute for it the letter *l*; but this does not properly constitute *Grasseyement*. It consists in this: that, in words in which the letter *r* is joined to another consonant, a sort of *burrowing* or guttural rolling is heard, nearly like that produced by gargling. See Rotacismus.

GRASUS, Cinabra.

GRATELLE, Psoriasis.

GRATIA DEI, *Gratiola officinalis*.

GRATIOLE CENTAURIODES, *G. officinalis*.

GRATIOLE OFFICINALE, *Digitalis minima*, *Gratia Dei*, *Gratiola Centaurioides*, *Hedge hyscop*, *Herb of Grace*, from *gratia*, 'grace or favour,' owing to its presumed excellent medical virtues. It is a native of the south of Europe. (F.) *Gratiolæ, Séné des prés, Petite digitale, Herbe au pauvre homme*. The plant is inodorous; taste strong, bitter, nauseous. It is possessed of anthelmintic, purgative, emetic, and diuretic properties. Dose, ten grains.

GRATIOLE, *Gratiola officinalis*.

GRATTERON, *Galium aparine*.

GRATTOIR, Raspatorium.

GRAVATIF, Heavy.

GRAVE, Serious—*g. Plant*, *Datura sanguinea*.

GRAVEDO, Catarrh, Coryza—*g. Neonatorum*, Sunifles.

GRAVEL, *Lithiâ renâlis areno'sa*, *Lithiâasis nephritica seu renâlis*, (F.) *Gravelle*. A disease occasioned by small concretions, similar to sand or gravel, *Gla'rea*, (F.) *Gravies*, which form in the kidneys, pass along the ureters to the bladder, and are expelled with the urine. These concretions, which are commonly composed of uric acid and an animal matter, are deposited at the

bottom of the vessel, immediately after the excretion of the urine; and, by their hardness and resistance under the finger, differ considerably from the ordinary sediment of that liquid. A vegetable diet and alkaline drinks are the best prophylactics. See Calculi, urinary. *A fit of the Gravel, Nephral'gia calculosa seu areno'sa, Col'lica nephrit'ica*, is the excruciating suffering induced by the passage of gravel from the kidney to the bladder. It can only be relieved by anæsthetics, opiates, the warm bath, &c.

When the deposit is in fine particles, it is termed *Sand*, *Are'na*, *Are'nula*, *Psam'ma*, *Psammus*.

GRAVEL GRASS, *Galium verum*.

GRAVEL, PILEOUS or HAIRY, (F.) *Gravelle pileuse*. A species of gravel containing hairs, phosphate of lime, ammoniaco-magnesian phosphate, and a little uric acid.—Magendie.

GRAVEL PLANT, *Epigæa repens*—*g. Root*, *Eupatorium purpureum*.

GRAVELEUX, Calculous.

GRAVELLE, Chalaza, Gravel—*g. Pileuse*, Gravel, pileous.

GRAVEOLENS, Fetid.

GRAVID, Pregnant.

GRAVIDIN; from *gravidus*, 'pregnant,' *gravis*, 'heavy.' A sediment in the urine of pregnant women, which by its decomposition gives rise to the pellicle kysteine. It differs from albumen, casein and gelatin.—Stark.

GRAVIDITAS, Fecondation, Pregnancy—*g. Abdominalis*, Pregnancy, abdominal—*g. Extra-uterina*, see Pregnancy—*g. Extra-uterina in Ovario*, Pregnancy, ovarian—*g. Extra-uterina Secundaria*, Metacyesis—*g. Interstitialis*, Pregnancy, interstitial—*g. Molaris*, Mole—*g. Ovarii*, Pregnancy, ovarian—*g. Spuria*, Pregnancy, false—*g. Tubaria*, Pregnancy, tubal, Salpingo-cyesis—*g. Uteri substantia*, Pregnancy, interstitial—*g. Uterina*, Pregnancy.

GRAVIER, Gravel.

GRAVIMETER, Aerometer.

GRAVIS, Heavy.

GRAVITY, SPECIFIC, *Gravitas specific'a*, (F.) *Pesanteur spécifique*. The relation between the weight of a body and its bulk: thus, supposing four bodies to be of the same size, but to weigh, one four, another three, another two, and the fourth one; the specific gravity of the first will be four times greater than that of the last. The specific gravities of different bodies are, therefore, as the weights, bulk for bulk. For solids, and liquids, water is taken as the unit; atmospheric air for the gases. Thus, water is 1.000; mercury, sat the common temperature, 13.58. Whence, we conclude mercury is between thirteen and fourteen times heavier than water.

GRAYHAIRD, see Poliosis.

GRAY MATTER OF THE BRAIN, Cortex cerebri, see Neurine.

GREASE, from (F.) *Graisse*, 'fat.' A specific inflammation, affecting the skin of the heels of the horse, which is especially interesting from the circumstance, that the matter, if inserted under the cuticle of an unprotected individual, may give rise to an affection—*grease-pox, variolâ equinâ*—which preserves the person from small-pox. [?]

GREASE, BARROW'S, *Adeps suillus*—*g. Goose*, *Adeps anserinus*—*g. Molten*, *Gras fondure*.

GREASE POX, see Grease.

GREENHEART, see Bebeera.

GREENHOW'S TINCTURE FOR THE TEETH, *Spiritus armoracæ compositus*.

GREEN SICKNESS, Chlorosis.

GREENWEED, *Genista tinctoria*.

GRÈLE (F.), *Grac'itis*, 'long and thin.' This

epithet is given by the French to various parts, as the

Apophyse Grêle du Marteau, the slender apophysis or process of the malleus. A long process situate at the anterior part of the neck of the malleus, which passes out by the fissure of Glaserius. It is also called the *Apophysis of Rau*, although it was already known to Fabricius ab Aquapendente and to Cæcilus Follius.

GRÊLE, Chalaza—*g. Interne de la Cuisse*, Gracilis.

GRÊMIL OFFICINAL, *Lithospermum officinale*.

GREMIUM, Vulva.

GRENADÉ, Influenza.

GRENADIER, *Punica granatum*.

GRENADIN, see *Punica granatum*.

GRENIERS, *Vesiculæ seminales*.

GRENOUILLE, *Rana esculenta*.

GRENOUILLETTE, *Ranula*.

GRÉOULX, MINERAL WATERS OF. Sulphuretted springs in the department of Basses-Alpes, France.

GRESSURA, Perinæum.

GRESSUS, Walking.

GREVEURE, Hernia.

GRIELUM, *Apium petroselinum*, *Smyrniolum asutrum*.

GRIFF, see *Mulatto*.

GRIFFO, see *Mulatto*.

GRINCEMENT DES DENTS, *Brygmus*.

GRINCOMES, Syphilis.

GRINDERS, Molar teeth—*g. Asthma*, see *Asthma*, grinders'—*g. Rot*, *Asthma*, grinders'.

GRINDING MILL, see *Pulverization*.

GRIPE, CUTTING ON THE, see *Lithotomy*.

GRIPEs, *Tormina*, *Colic*.

GRIPEs, WATERY. A popular name for a dangerous disease of infancy, common in England, which does not differ essentially from the cholera infantum of this country.

GRIPHOSIS, *Onychogryphosis*.

GRIPPE (F.), from *gripper*, 'to gripe,' 'catch hold of.' A vulgar name for several catarrhal diseases, which have reigned epidemically; as the influenza.

GRIPPE, Pinched.

GRISOU, see *Hydrogen*, carburetted.

GRISTLE, *Cartilage*.

GRITS, *Groats*.

GRIT GRUEL, *Water gruel*. This is made as follows:—Take three ounces of grits; wash them well in cold water, and, having poured off the fluid, put them into four pints of fresh water, and boil slowly until the water is reduced one-half; then strain through a sieve. It is a good demulcent, and is employed also as a vehicle for clysters.

GROAN, see *Suspirium*.

GROATS, German *Grütze*; *Grutum*, *Avena excorticata*; (F.) *Gruau*, *Oatmeal*, (*Yorkshire*.) Oats, hulled, but unground, (*Lancashire*.) Hulled oats, half ground. Oats that have the hulls taken off; *Grita*, (Prov.) *Culina*. When crushed, they are termed *Emden groats*. In America, fine hominy is called *Grita*, and wheat prepared in the same way is likewise so designated. It is also called *wheaten hominy*.

GROATS, CRACOW, *Semolina*.

GROCERS' ITCH, see *Eczema Impetiginodes*, *Psoriasis*.

GROG-BLOSSOMS, *Gutta rosea*.

GROG-ROSES, *Gutta rosea*.

GROMWELL, *Lithospermum officinale*.

GROOVE, *Furrow*, *Sulcus*, (F.) *Rainure*. Icelandic, *grafa*, Sax. *grafan*, 'to dig.' A channel or gutter in a bone or surgical instrument. See *Coulisse*.

GROOVE, PRIMITIVE, *Primitive streak or trace*, *Nota primitiva*. A bright streak in the long axis of the pellucid part of the area germinativa, after it presents a central pellucid and a peripheral opaque part, and passes from the round to the pear shape.

GROOVED. Same etymon. *Sulcatus*, *Striatus*, *Canaliculatus*, (F.) *Cannelé* ou *Canelé* ou *Canaliculé*; *Canaliculated*. Having a small channel or gutter.

GROS, *Drachm*—*g. Cou*, *Bronchocele*.

GROSEILLES, *Ribes rubrum*—*g. à Maque-reau*, *Ribes uva crispa*.

GROSELLER NOIR, *Ribes nigrum*—*g. Rouge*, *Ribes rubrum*.

GROSSE GORGE, *Bronchocele*.

GROSSESSE, *Pregnancy*—*g. Abdominale*, *Pregnancy*, abdominal—*g. Afœtale*, *Pregnancy*, afœtal—*g. Bigéminale*, *Pregnancy*, bigeminal—*g. Complexe*, *Pregnancy*, complex—*g. Composée*, *Pregnancy*, compound—*g. Contre-nature*, *Pregnancy*, extra-uterine—*g. Fausse* ou *apparente*, *Pregnancy*, false—*g. Fœtale*, *Pregnancy*, foetal—*g. Gazo-hystérique*, *Pregnancy*, gazo-hysteric—*g. Hémato-hystérique*, *Pregnancy*, hemato-hysteric—*g. Hydro-hystérique*, *Pregnancy*, hydro-hysteric—*g. Inter-extra-utérine*, *Pregnancy*, complex—*g. Ovarienne*, *Pregnancy*, ovarian—*g. Sarco-hystérique*, *Pregnancy*, sarco-hysteric—*g. Sarcœfœtale*, *Pregnancy*, sarcœfœtal—*g. Simple*, *Pregnancy*, solitary—*g. Solitaire*, *Pregnancy*, solitary—*g. Trigéminale*, *Pregnancy*, trigeminal—*g. Triple*, *Pregnancy*, trigeminal—*g. Tubaire*, *Pregnancy*, tubal—*g. Utéro-abdominale*, *Pregnancy*, utero-abdominal—*g. Utéro-ovarienne*, *Pregnancy*, utero-ovarian—*g. Utéro-tubaire*, *Pregnancy*, utero-tubal.

GROSSULARIA, *Ribes uva crispa*—*g. Nigra*, *Ribes nigrum*—*g. Non spinosa*, *Ribes nigrum*—*g. Rubra*, *Ribes rubrum*.

GROTTO DEI SERPI (I.) 'Grotto of serpents.' A grotto near Braccano in Italy, filled with warm water, to which invalids, affected with skin diseases, resort.

GROUILLEMENT D'ENTRAILLES, *Borborygmus*.

GROUND BERRY, *Gaultheria*—*g. Holly*, *Gaultheria*—*g. Ivy*, *Gaultheria*, *Glechoma hederacea*—*g. Nut*, *Arachis hypogæa*, *Pignut*—*g. Pine*, *Teucrium chamæpitys*—*g. Pine*, *French*, *Teucrium iva*.

GROUNDIE SWALLOW, *Senecio*.

GROUNDSEL, *Senecio*—*g. Tree*, *Baccharis halimifolia*.

GROUSEBERRY, *Gaultheria*.

GROWTH, from Dutch *groyen*, 'to grow,' *Incrementum*, *Crescentia*, *Anaploxis*, *Anaptyxis*, (F.) *Croissance*, *Crue*. The development of the body; particularly in the direction of its height. Also, any adventitious tissue; thus, we speak of a morbid growth or formation. See *Development*.

GROZET-EYED, *Goggle-eyed*.

GRUAU, *Groats*.

GRUB, *Larve*, see *Ectozoa*.

GRUBS, *Acne punctata*.

GRUEL, GRIT, see *Grits*—*g. Water*, see *Avo-na*, and *Grits*.

GRUFFE, from Teutonic *ge*, and *ruh*, 'rough.' In pharmacy, the coarse residue, which will not pass through the sieve in pulverization.

GRUMEAU, *Coagulum*.

GRUMOUS, *Grumous*, (F.) *Grumelé*, *Grumeleux*, from *grumus*, 'a clot.' Clotted.

GRUMUS, *Coagulum*.

GRUTUM. 'Groats.' *Grutum Milvium*, *Milvium*. A hard white tubercle of the skin, resembling, in size and appearance, a millet-seed. It is confined to the face. See, also, *Groats*.

GRYPHIUS PES. The *Griffon's foot*, (F.)

Pied de Griffon, An instrument of which Ambrose Paré speaks, which was used for extracting moles from the uterus.

GRYPHOSIS, Onychogryphosis.

GRYPPOSIS, from *γῡρῡσῡ*, 'I incurvate.' *Incurvatio*. Curvature or crookedness in general. Crookedness or incurvation of the nails. See *Onychogryphosis*.

GRYPOTES, see Grypus.

GRYPUS. One who has a crooked or aquiline nose. The condition is termed *Grypotes*.

GUACO, *Huaco*. The name of a plant, *Eupatorium Guaco*, described by Humboldt and Bonpland under the name *Mikania Guaco*, *Ord.* Compositæ, which grows in the valleys of Madalena, Rio-Cauca, &c., in South America. The negroes use the juice against the bites of poisonous reptiles;—both in the way of prevention and cure. It has been, of late, brought forward as a remedy in cholera.

GUAIAC, see Guaiacum.

GUAIACI LIGNUM, see Guaiacum—*g. Resina*, see Guaiacum.

GUAIACINE, see Guaiacum.

GUAIACUM, *G. officinale* seu *America-num*, *Lignum vitæ* seu *sanctum* seu *benedictum* seu *Indicum*, *Palus sanctus*, *Hagia-lylun*, (F.) *Gayac*, *Gaiac*. *Ord.* Zygophylles. The resin—*Guaiaci Resina*, *Guaiac*, (F.) *Résine ou Gomme de Gaiac*—and the wood—*Guaiaci lignum*—are both used in medicine. Their odour is slightly fragrant; taste warm and bitter, of the resin more so than of the wood. The resin is concrete, brittle; colour, externally, greenish; internally grayish. Water dissolves about one-tenth; alcohol 95 parts. It is soluble, also, in *liquor potassæ* 15 parts, *liquor ammoniæ* 38 parts. The powder is whitish, but changes green in the air. The base of the guaiacum is a peculiar resin, called *Guaiacine*.

Guaiacum is stimulant and diaphoretic; and in large doses, purgative. It is administered in chronic rheumatism, gout, cutaneous diseases, and the sequelæ of syphilis. Dose of resin, gr. v to xx:—to purge, gr. xx to xl.

Guaiacum sanctum, and *G. angustifolium* have the same virtues as *G. officinale*.

GUALTHERIA, *Gaultheria*.

GUANO,—according to Tschudi, properly *Huanu*,—is formed of the excrements of different kinds of marine birds—mews, divers, sheerbreaks, &c., but especially of the *Sula variegata*. It is found in enormous layers in the South American islands of the Pacific, and is used as manure.

GUARANA, *Paullinia*.

GUARANHEM, *Monesia*.

GUARANIN, see *Coffea Arabica*.

GUARAPO. A fermented liquor made, in Peru, of sugar-cane pulp and water. It is a very favourite beverage of the negroes.—Tschudi.

GUARD (for a bed), *Alde*.

GUARERBA ORBA, *Momordica elaterium*.

GUAVA APPLE, *Psidium pomiferum*.

GUAYAVA, *Psidium pomiferum*.

GUAZA, see *Bangue*.

GUERNACULUM DENTIS, (*gubernaculum*, 'a rudder.') A cord, which passes from the follicle of the permanent tooth along a small long canal behind the alveolus of the milk tooth, and becomes continuous with the gum. The gubernaculum has been supposed to direct the permanent tooth outwards. The canal has been termed *iter dentis*.

GUERNACULUM TESTIS, *Gubernaculum testis*, *Hunteri*, *G. Hunteri*, *Hunteri Ligamentum suspensorium Testis*, (F.) *Gouvernail du testicule*. A triangular, fibro-cellular cord; which, in the foetus, arises from the ramus of the ischium and the skin of the scrotum, and proceeds to the posterior part of the testicle, before this organ issues from the

abdomen. It has been supposed to be a continuation of the fascia superficialis with muscular fibres from the internal oblique muscle, which pass upwards to the testis when in the abdomen; and by their contraction draw the testis down, and ultimately form the cremaster muscle.

GUÊPE, Wasp.

GUÉRISON, Cure.

GUÉRISABLE, Curable.

GUI, *Viscum album—g. de Chêne*, *Viscum album*.

GUIDER, Tendon.

GUILANDINA MORINGA, *Hyperanthea moringa*. A plant, *Ord.* Leguminosæ, which affords the *Ben nut*, and the *lignum nephriticum*. It is also called *Moringa Oleifera* seu *Zeylanica* seu *Nux ben*. Under the name *Guilandina moringa*, Linnæus is said to have confounded *Moringa pterygosperma* and *M. aptera*. *Hyperanthea moringa* is a synonyme of the former species. The nut *Ben*, *Ben nux*, *Glans unguenta*, *ria*, *Balanus Myrepisia*, *San'dalum cernu*, *leum*, *Oily Acorn* or *Ben nut*, is a West India nut which furnishes an oil, *O'leum Balani-num*, *Oil of Ben*, that does not become rancid by age, and is hence used by perfumers. It is purgative.

The wood of the Guilandina is called *Lignum Nephriticum*, and has been used in decoction, in affections of the urinary organs.

GUILLOTINE, TONSIL, *Amygdalotome*.

GUILTY CUPS, *Ranunculus acris*.

GUIMAUVE, *Althæa—g. Veloutée*, *Hibiscus abelmoschus*.

GULA, *Œsophagus*, *Pharynx*.

GULÉ IMBECILLITAS, *Pharyngoplegia—g. Principium*, *Pharynx*.

GULLET, *Œsophagus*.

GULLION, *Colic*.

GULOSUS, *Glutton*.

GULSCHUCH, *Icterus*.

GUM, ACAROID, see *Xanthorrhœa—g. Anime*, *Anime—g. Arabic*, *Acaciæ gummi—g. Artificial*, *Dextrin—g. Bassora*, *Bassora gum—g. Black-boy*, see *Xanthorrhœa*.

GUM BOIL, *Parulis*.

GUM, BRITISH, *Dextrin—g. Butea*, see *Butea frondosa—g. Caranna*, *Caranna—g. Dragon*, *Tragacantha—g. East India*, see *Feronia elephantum—g. Elastic*, *Caoutchouc*.

GUM OF THE EYE, *Chassie—g. Falling away of the*, *Ulatrophia—g. Hemlock*, see *Pinus Canadensis—g. Indigenous*, *Gummi nostras—g. Ivy*, see *Hedera helix—g. Juniper*, *Sandarac*.

GUM-LANCET, *Denticulipium*, *Odontoglyphon*, (F.) *Déchaussoir*. An instrument for separating the gum from the cervix of the tooth, prior to extraction. It is formed much like a fleam. The operation itself is called *Lancing the gums*, *Denticulipium*, (F.) *Déchaussement*.

GUM, ÖRENBURG, see *Pinus larix—g. Red*, *Strophulus*.

GUM RESIN, *Gummi resina*.

GUM, SANDARACH, *Sandarac—g. Sassa*, *Sassa—g. Seneca*, *Acaciæ gummi—g. Senega*, *Acaciæ gummi—g. Shrinking of the*, *Ulatrophia—g. Sweet*, *Liquidambar styraciflua—g. Tragacanth*, *Tragacanth*.

GUM TREE, BROWN, see *Kino*.

GUM, WHITE, *Liquidambar styraciflua*, *Strophulus—g. Yellow*, *Icterus infantum*.

GUMMA, (F.) *Gomme*. An elastic tumour formed in the periosteum occupying particularly the cranium and sternum, and produced by the syphilitic virus, when it has been long in the constitution. It is so called, because, when opened, it contains a matter like gum.

GUMMI, *Commis*, *копм*, (F.) *Gomme*. An immediate principle of vegetables. It is a solid, uncrystallizable, inodorous substance, of a mawk-

fish taste, unchangeable in the air, insoluble in alcohol, but soluble in water, with which it forms a mucilage. It is obtained from various species of *minosa* and *prunus*; and consequently there are many varieties of gum. They are used in medicines as demulcents, emollients, and relaxants, particularly in catarrh, intestinal irritations, &c.; and in *Pharmacy*, they are employed in the formation of emulsions, pills, &c.

GUMMI ACACIE ARABICÆ, *Acaciæ gummi*—*g. Acanthium*, *Acaciæ gummi*—*g. Acaroides*, see *Xanthorrhœa*—*g. Adstringens* Fothergilli, Kino—*g. Ammoniacum*, *Ammoniac*—*g. Anime*, *Anime*—*g. Arabicum*, *Acaciæ gummi*—*g. Astragali* *Tragacanthæ*, *Tragacantha*—*g. Bogia*, *Cambogia*—*g. Brelisis*, *Caranna*—*g. Copalimum*, *Copal*—*g. Elasticum*, *Caoutchouc*—*g. Euphorbia*, see *Euphorbia officinarum*—*g. Gamandra*, *Cambogia*—*g. Gambiense*, *Kino*—*g. de Goa*, *Cambogia*—*g. Gutta*, *Cambogia*—*g. Hederæ*, see *Hedera helix*—*g. de Jemu*, *Cambogia*—*g. Juniperi*, *Sandarac*—*g. Laccæ*, *Lacca*—*g. Ladanium*, see *Cistus creticus*—*g. Lamac*, *Acaciæ gummi*—*g. Laricis*, see *Pinus larix*—*g. Leucum*, *Acaciæ gummi*.

GUMMI NOSTRAS, (F.) *Gomme du Pays*; *Indig'neuse Gum*. These generic names are given to several species of gum, which flow spontaneously from certain indigenous fruit trees, such as the almond, cherry, peach, apricot, &c. The indigenous gums have nearly the same properties as gum Arabic; but they are inferior to it.

GUMMI ORENBURGENSE, see *Pinus larix*—*g. Parnacis*, see *Pastinaca opoponax*—*g. ad Podagram*, *Cambogia*.

GUMMI-RES'INA, *Gum-resin*. A milky juice, obtained by making incisions into the branches, stalks and roots of certain vegetables. Gum-resins are compounds of resin, gum, often with essential oil, and different other vegetable matters. They are solid, opaque, brittle, of a strong odour, acid taste, variable colour, and are heavier than water. Water dissolves a part of them, and alcohol another; hence proof spirit is the proper menstruum. The generality of the gum-resins are powerful stimulants to the whole or to parts of the economy. The chief are *asafoetida*, *gum ammoniac*, *euphorbium*, *galbanum*, *cambooge*, *myrrh*, *olibanum*, *opoponax*, *scammony*, *aloes*, &c.

GUMMI RUBRUM ADSTRINGENS GAMBIESE, *Kino*, see *Butea Frondosa*—*g. Seneca*, *Senegal gum*—*g. Senega*, *Senegal gum*—*g. Senegalense*, *Senegal gum*—*g. Senica*, *Senegal gum*—*g. Serapionis*, *Acaciæ gummi*—*g. Thebaicum*, *Acaciæ gummi*—*g. Tragacantha*, *Tragacantha*—*g. Uralense*, see *Pinus larix*.

GUMMIDODES, Mucilaginous.

GUMMIODES, Mucilaginous.

GUMMOSUS, Mucilaginous.

GUMS, *Gingivæ*.

GUN-COTTON, see *Collodion*—*g. c. Ethereal solution of*, *Collodion*.

GUNJAH, see *Bangue*.

GUNNERA PERPEN'SA. A South African plant, *Nat. Ord.* *Urticacæ*; the decoction of which is taken as a domestic remedy by the farmers, as a tonic in dyspepsia. A tincture has been used in gravel. An infusion of the leaves is demulcent, and is employed in pulmonary affections. The leaves are applied fresh, to cure wounds and ulcers.

GURGES, *Pharynx*.

GURGITELLO, MINERAL WATERS OF.

A thermal spring in the isle of Ischia. Temp. at its source, 176° Fah. It contains carbonic acid, carbonates of lime, magnesia, iron and soda, sulphates of lime and soda, chloride of sodium and silica.

GURGLING. 'Gushing with noise,' as water from a bottle. (F.) *Gargouillement*, same etymon

as gargle. The rhonchus or *râle* heard on auscultation when there is a cavity in the lungs containing pus. It is the 'cavernous rattle or rhonchus,' *Râle cavernæ*. The size of the bubbles heard varies, and hence the rhonchus has been called *cavernous* and *cavernulous*, (F.) *Râle cavernuleux*. If the cavern be large, this *râle* will nearly resemble the gurgling of a bottle (*gorglous d'une bouteille*;) if, on the contrary, the cavern be small, it will not differ from the *râle muqueux*.

GURGULIO, *Penis*, *Uvula*.

GUSTATIF, (Nerf,) see *Lingual Nerve*.

GUSTATION, Degustation, Taste.

GUSTATORY NERVE, see *Lingual Nerve*.

GUSTUS, Taste—*g. Depravatus*, *Parageusia*.

GUT, Gout, Intestine—*g. Blind*, *Cæcum*.

GUTS, SLIPPERINESS OF THE, *Lientery*.

GUTTA, Apoplexy, *Cambogia*, *Gout*.

GUTTA. A Drop, *Stalagma*, *Alun'sel*, *Stilla* (F.) *Goutte*. A quantity of liquid, generally valued, in pharmacy, at the weight of a grain. The weight, however, varies according to different circumstances, as the degree of tenacity of the fluid, and the extent of moist surface to which the suspended drop is attached before it falls; and it was found by Mr. Alsop to be influenced by the size of the bottle, and the angle of inclination at which it was held during the operation of dropping. The following are some of his results as to the number of drops required to measure a fluidrachm, when dropped from a large and a small bottle.

	From a large bottle.	From a small bottle.
(f3) Diluted sulphuric acid.....	24 drops	84 drops
Scheele's hydrocyanic acid 35.....	50	74
Distilled water.....	31.....	54
Solution of ammonia.....	40.....	48
Tincture of opium.....	84.....	135
Rectified spirits.....	100.....	130
Tincture of chloroide of iron.....	100.....	150

GUTTA GAMANDRÆ, *Cambogia*—*g. Gamba*, *Cambogia*—*g. Opaca*, *Cataract*.

GUTTA PERCHA, *Gutta tuban*, *Gomme getta'nia*. The concrete juice of a tree—*Isanan'dra gutta*—which is indigenous in Singapore and its vicinity, and belongs to the *Natural Order* *Sapotacææ*. Plunged in boiling water it softens, when it may be moulded like caoutchouc to any form, which it retains on cooling. Splints and other instruments have been made of it.

A solution of *gutta percha* in *chloroform*—*Traumaticine*—has been used with advantage, topically, in various cutaneous affections, serofulous and indolent ulcers, and as an ecrotic in small-pox.

GUTTA ROSACEA SYPHILITICA, Crown of Venus.

GUTTA RO'SEA, *G. Rosa'cea*, *Ion'thus corymbifer*, *Crusta serpiginosa seu pruriginosa*, *Anê rose'cea*, *Rose'ola acn'ea*, *Thylaci'itia*, *Bacchia*, *Butiga*, *Cav'buncled face*, *Rosy Drop* or *Wheel*, *Copper-nose*, *Bottle-nose*, *Grog-blossoms*, *Grog-roses*, *Brandy-face*, (F.) *Couperose*, *Goutte Rose*, *Bourgeons*, *Dartre pustuleuse couperose*. An eruption of small, suppurating tubercles, with shining redness, and an irregular granular appearance of the skin of the part of the face which is affected. The redness commonly appears first at the end of the nose, and then spreads on both sides. It is often produced by hard drinking. Its cure must be attempted by regular regimen, and cooling means internally: weak spirituous or saturnine lotions externally. The affection is usually very obstinate.

GUTTA SERENA, Amaurosis—*g. Tuban*, *Gutta percha*.

GUTTÆ ABBATIS ROUSSEAU, *Laudanum abbatiss Rousseau*—*g. Ammoniaci*, see *Ammoniac*.

gum—*g.* Acidæ tonicæ, Elixir Acidum Halleri—*g.* Nervinæ, Alcohol sulfurico-æthereus ferri.

GUTTÆ NIGRÆ, *Ace'tum o'pii*, Common Black Drop. (*Opit* ζ'viii, *acetii* distillat. bij. Infuse.)

The celebrated Black Drop, Lan'caster or Quaker's Black Drop may be made as follows. Take half a pound of opium sliced; three pints of good verjuice, (juice of the wild crab,) one and a half ounce of nutmegs, and half an ounce of saffron. Boil to a proper thickness, and add a quarter of a pound of sugar, and two spoonfuls of yeast. Set the whole in a warm place, near the fire, for six or eight weeks; then place it in the open air until it becomes a syrup. Lastly, decant, filter, and bottle it up, adding a little sugar to each bottle. One drop is equal to three of laudanum; and it is nearly devoid of all the unpleasant exciting effects of the latter. An analogous formula is contained in the Pharmacopœia of the United States under the name *Ace'tum opii*, *Vinegar of Opium*.

GUTTALIS CARTILAGO, Arytenoid Cartilage.

GUTTERIA, Bronchocele.

GUTTUR. The throat; the larynx; the trachea.

GUTTUR GLOBOSUM, Bronchocele—*g.* Tumidum, Bronchocele.

GUTTURAL, *Guttura'lis*, from *guttur*, 'the throat.' Relating or belonging to the throat.

The Superior Thyroideal Artery is sometimes called *Guttural Artery*.

A *Guttural Cough* is one occasioned by irritation of the larynx or trachea.

A *Guttural Sound* is one produced, as it were, in the throat.

GUTTURIS OS, Hyoides os.

GUTTURNIA, Arytenoid cartilages.

GYMNA, Genital Organs.

GYMNASION, Exercise.

GYMNASIUM, from *gymnos*, 'naked.' *Palæstra*. An establishment amongst the ancients, intended for bodily exercises, as wrestling, running, &c.; a term now used, in Germany more especially, for an academy or higher school.

GYMNAST, *Gymnas'tes*, *Gymnas'ta*; same etymon. The manager of a gymnasium. One, whose profession it is to prevent or cure diseases by gymnastics.

GYMNASTICS, *Gymnas'tica*, *Medici'na gymnast'ica* seu *euect'ica*, *Somascét'ics*, (F.) *Gymnastique*, *Somascétique*. Same etymon. That part of hygienic medicine which treats of bodily exercises. It is called *Med'ical Gymnastics*. The ancients had also *Athlet'ic Gymnastics*, and *Military Gymnastics*. Herodotus, of Selivriæ, first proposed gymnastics for the cure of disease.

GYMNASTIQUE, Gymnastics.

GYMNEURA LACTIFERUM, *Ceylon cow-tree*, *Kiriaghuma*. A tree, of the Family Apocynaceæ, the juice of which is used by the Cingalese as milk.

GYMNOCLADUS CANADENSIS, *Coffee Tree*, *Kentucky Coffee-bean Tree*, *Mahogany*, *Nickar Tree*, *Bondue*. Order, Leguminosæ; from *gymnos*, 'naked,' and *κλαδος*, 'a branch.' An indigenous tree, which grows from Ohio to Louisiana. The leaves are cathartic, and said to contain cytisin. The seeds are a good substitute for coffee.

GYMNOGRAMME CETERACH, *Asplenium ceterach*.

GYMNOPTERIS CETERACH, *Asplenium ceterach*.

GYMNOSIS, Denudation.

GYNE/CANER, from *gyn*, *gynaikos*, 'a woman,' and *ανω*, 'a man.' *Vir effemina'tus*. An effeminate man.

GYNECANTHE, *Tamus communis*.

GYNECATOP'TRON, from *gyn*, *gynaikos*, 'a woman,' and *κατοπτρον*, 'a speculum.' A speculum for investigating the female organs of generation.

GYNECEA, *Gynæceia*.

GYNECEIA, *Gynæci'a*, *Gynæce'a*, from *gyn*, *gynaikos*, 'a woman.' The catamenia;—the lochia. The diseases of women in general.—Hippocrates, Galen, Fœsius.

GYNECEUM, Antimonium, Vulva.

GYNECE'US, from *gyn*, *gynaikos*, 'a woman.' Belonging to women. Female. Feminine.

GYNECIA, *Gynæcia*, Menses.

GYNECOLOG'IA, *Gynecol'ogy*, from *gyn*, *gynaikos*, 'a woman,' and *λογος*, 'a description.' The doctrine of the nature, diseases, &c. of women.

GYNECOMA'NIA, from *gyn*, *gynaikos*, 'woman,' and *μανια*, 'mania,' 'rage.' That species of insanity, which arises from love for women. Some have used the word synonymously with nymphomania.

GYNECOMAZIA, see *Gynæcomastus*.

GYNECOMAS'TUS, *Gynæcomas'thus*, from *gyn*, *gynaikos*, 'woman,' and *μαστος*, 'a breast.' A man whose breasts are as large as those of a woman.—Galen, Ingrassias. The condition is called *Gynæcoma'zia*.

A considerable enlargement of the breasts of a female was formerly called *Gynæcomas'ton*.

GYNECOMYS'TAX, from *gyn*, *gynaikos*, 'woman,' and *μοσταξ*, 'the beard.' The hair on the pubes of women.—Rolfink.

GYNECOPHO'NUS, from *gyn*, *gynaikos*, 'woman,' and *φωνη*, 'voice.' A man who has an effeminate voice.

GYNANDRIA, Hermaphrodeity.

GYNANDROP'SIS PENTAPHYL'LA, *Cleo'nē pentaphyl'la*. Ord. Capparidaceæ; an indigenous plant, is possessed of diaphoretic properties. The juice alone, or mixed with oil, has been used in earache.

GYNANDRUS, *Gynanthropus*.

GYNANTHRO'PUS, *Gynan'drus*. An hermaphrodite who belongs more to the male than to the female sex.

GYNATRE'SIA, from *gyn*, 'a woman,' and *απντρος*, 'imperforate.' Closure or imperforation of the external parts of generation of the female.

GYNE, Female, Woman.

GYNECOLOGY, *Gynæcologia*.

GYNIDA, Hermaphrodite.

GYNOARIUM, Ovarium.

GYNOCARDIA ODORATA, *Chaulmoogra*.

GYNOPLAS'TIC, *Gynoplas'ticus*, from *gyn*, 'a woman,' and *πλασσω*, 'I form.' The gynoplastic operation is employed for opening or dilating the closed or contracted genital openings of the female.

GYPSY-WEED, *Lycopus sinuatus*, and *L. Virginicus*.

GYRI CEREBRI, Convolutions (cerebral).—*g.* Cochleæ, 'Scale of the Cochlea'—*g.* Intestinales, see *Anse*.

GYROMIA VIRGINICA, *Medeola Virginica*.

GYROPHLE, see *Eugenia caryophyllata*.

GYROPHORA, see *Tripe de Roche*.

GYRUS, Convolution—*g.* Intestinalis, *Anse intestinale*.

GYRUS FORNICA'TUS, 'Arched convolution.' *Fillet of the corpus callosum*. A large convolution of the brain, which lies horizontally on the corpus callosum, and may be traced forwards and backwards to the base of the brain, terminating by each extremity at the fissure of Sylvius. The surface of the hemisphere, where it comes in contact with the corpus callosum, is bounded by it.

GYTLING, Infans.

H.

HABBERING, Balbuties.

HAB-EL-KALIMBAT, *Pistacia terebinthus*.

HABBI, Hagenia Abyssinica.

HABBLING, Balbuties.

HABE'NA. *A Bridle, Tel'amon*. A bandage for uniting the lips of wounds; which, in many instances, replaced the suture.—Galen.

HABENÆ OF THE PINEAL GLAND, see Pineal Gland.

HABENULÆ OF THE PINEAL GLAND, see Pineal Gland.

HABIL'LA DE CARTHAGE'NA, *Bejuco, Carthage'na Bean*, (F.) *Fève de Carthagène*. A kind of bean of South America, famed as an effectual antidote for the poison of all serpents, if a small quantity be eaten immediately.

HABIT, *Habitu'do*, from *habere, habitum*, 'to have or to hold,' *Assuetu'do, Mos, Usus, Consuetu'do, Ethos, ἔθις, Hexis, Ethmos'ynē*, (F.) *Habitude, Accoutumance*. Habit is the aptitude for repeating certain acts:—or, a facility, which results from the frequent repetition of the same act. It is, according to vulgar expression, 'a second nature.' Habit may predispose to certain diseases, or it may protect against them. It ought not to be lost sight of, in attending to the progress of disease, or of its treatment.

HABIT OF BODY, *Constitu'tio, Hab'itus, H. Cor'poris, Catas'tasis, Hexis, Epiphan'a*, (F.) *Habitude extérieure, Habitude du corps*. The aggregate of the physical qualities of the human body.

HABITUDE, Habit—*h. du Corps*, Habit of body—*h. Extérieure*, Habit of body.

HABITUDO, Habit.

HABITUS, Habit of body—*h. Apoplectic*, Apoplectic habit—*h. Corporis*, Habit of body—*h. Quadratus*, see Apoplectic—*h. Torosus*, see Apoplectic.

HABROMA'NIA, from *ἄβρος, 'gay,' and μανία, 'mania'*. Insanity in which the delusions are of a gay character.

HACHICH, see Bangué.

HACHISCH, Bangué.

HACKBERRY, *Celtis occidentalis, Prunus padus*.

HADDOCK, see *Oleum jecoris aselli*.

HÆMA, *αἷμα, 'aiματος, 'blood.'*

HÆMACHROINE, Hæmatine.

HÆMACHRO'SES, ((G.) *Hämachrosen*), from *hæma*, and *χρῶσις, 'coloration.'* A family of diseases in which the blood has its colour different from usual, as in purpura and cyanosis.—Fuchs.

HÆMACY'ANIN, from *hæma*, and *κρῶσις, 'blue'*. A blue colouring matter, detected by Sanson in healthy blood, and in bile by some chemists, but not by others.

HÆMADON'OSUS, from *hæma*, and *νοσος, 'a disease.'* *Hæmatangion'osus, Hæmatangionu'sus, Hæmatangio'is*. Disease of the blood-vessels.

HÆMADOSTO'SIS, from *hæma*, and *ὄστωις, 'a bony tumour.'* Ossification of the blood-vessels.

HÆMADROMOMETER, Hæmodromometer.

HÆMADYNAMETER, Hæmadynamometer.

HÆMADYNAM'ICS, (F.) *Hémadynamique*, from *hæma*, and *δύναμις, 'power.'* The mechanical theory of the circulation of the blood.

HÆMADYNAMOM'ETER, *Hæmatodynam'eter, Hæmadynam'eter, Hæmom'eter, Hæmatom'eter*, (F.) *Hémadynamètre, Hémomètre*; from *hæma, δύναμις, 'power,' and μέτρον, 'a measure.'* An instrument for measuring the force of the blood in the vessels. It consists of a bent glass tube, the lower bent part of which is filled with mercury. A brass head is fitted into the artery, and a solution of carbonate of soda is interposed between the mercury and the blood, which is allowed to enter the tube for the purpose of preventing its coagulation. The pressure of the blood on the mercury in the descending portion of the bent tube causes the metal to rise in the ascending portion; and the degree to which it rises indicates the pressure under which the blood moves.

HÆMAGASTER, Hæmatogaster.

HÆMAGASTRIC PESTILENCE, Fever, yellow.

HÆMAGOGUM, Pæonia.

HÆMAGO'GUS, from *hæma*, and *αἷω, 'I drive off.'* A medicine which promotes the menstrual and hemorrhoidal discharges.

HÆMAL, from *hæma*. Relating to the blood or blood-vessels.

HÆMAL ARCH. The arch formed by the projections anteriorly of the ribs and sternum from the bodies of the vertebræ. It encloses the great blood-vessels.—Owen.

HÆMAL AXIS, Aorta—*h. Spine, Sternum*.

HÆMALOPIA, Hemophthalmia, Hæmalopis.

HÆMALOPIS, Hæmophthalmia.

HÆMALOPS, Hemophthalmia.

HÆMAN'THUS COCCIN'EUS. A beautiful South African plant, *Nat. Ord. Amaryllidæ*; the bulb of which is employed as a diuretic. It is given as an oxymer in asthma and dropsy. The fresh leaves are antiseptic, and applied to foul, flabby ulcers, and in anthrax.

HÆMAPERITONIRRHA'G'IA, (F.) *Hémapéritonirrhagie*; from *hæma, περίτονον, 'peritoneum,' and γαγν, 'a violent rupture.'* An exhalation of blood into the peritoneum.

HÆMAPHÆ'IN, *Hæmatophæ'um, Hæmophæ'um*, from *hæma*, and *φαῖος, 'of a dusky colour.'* A term applied by Simon to the brown colouring matter of the blood, supposed by some to be nothing more than hæmatin modified by an alkali.

HÆMAPH'OBUS, *Hæmoph'obus*, from *hæma*, and *φοβος, 'dread.'* One who has a dread of blood—who cannot look at it without fainting.

HÆMAPOPHYSES, Costal cartilages, see Costa.

HÆMAPOR'IA, *Hæmatapor'ia, Hæmatopor'ia, Oligohæ'mia, Oligæ'mia*, from *hæma*, and *ἀπορῆς, 'poor.'* Paucity of blood. See Anæmia.

HÆMAPTYSIS, Hæmoptysis.

HÆMAS, *αἷμας, gen. αἱματος*. A blood-vessel.

HÆMASTATICA, Hæmatostatica.

HÆMASTAT'ICS, *Hæmatat'ics*, from *hæma*, and *στατική, 'statics.'* The statics of the blood and blood-vessels; *Hæmatat'ics*.

HÆMATANAGOGÉ, Hæmatemesis.

HÆMATAL'LOSCOPY, *Hæmatallonscop'ia*, from *hæma, αλλοίος, 'different,' and σκοπεῖν, 'to view.'* The mode of distinguishing different kinds of blood from each other.—Toddei.

HÆMATANGIONOSUS, Hæmadonosus

HÆMATANGIOSIS, *Hæmadonosus*.

HÆMATAPORIA, *Hæmaporia*.

HÆMATAPORRHOSIS, *Cholera*.

HÆMATAPOSTEMA, from *hæma*, and *αποστήμα*, 'an abscess.' An abscess which contains blood.

HÆMATULICA, Vascular system.

HÆMATECLYSIS, *Hæmotex'ia*, *Hæmotex'is*, *Hæmatol'ysis*, *Hæmatosep'sis*, *San'guinis dissolutio*; from *hæma*, and *εκλυσίς*, 'loosening.' Dissolution of the blood.

HÆMATELÆUM, see Blood.

HÆMATEMESIS, from *hæma*, and *εμεω*, 'I vomit.' *Hæmorrhæ'gia* *Hæmatem'esis*, *Vom'itus cruent'us*, *Vom'itus seu Vomit'io San'guinis*, *Gastro-hæmorrhag'ia*, *Gastro-hæmorrhag'ia*, *Hæmorrhag'ia ventriculi*, *Hæmorrhæ'a ventriculi*, *Hæmatanag'o'gê*, *Vomiting of Blood*, (F.) *Hématémèse*, *Vomissement de sang*. Hæmatemesis is generally preceded by a feeling of oppression, weight, and dull or pungent pain in the epigastric and in the hypochondriac regions; by anxiety, and, occasionally, by syncope. Blood is then passed by vomiting, and sometimes, also, by stool,—the blood being generally of a grumous aspect. Hæmatemesis may be active or passive, acute or chronic. The blood effused proceeds from rupture of vessel, or from a sanguineous exhalation at the surface of the mucous membrane of the stomach. It is often observed in females whose menstrual secretion is irregularly performed. It is not of much danger, except when connected with disease of some of the solid viscera of the abdomen. On dissection of those who have died from protracted hæmatemesis,—for the acute kind is comparatively devoid of danger,—the mucous membrane of the stomach is found red and inflamed, or black, and the vessels considerably dilated. Complete abstinence from food; rest; the horizontal posture; bleeding, if the hæmorrhage be active; cold, acidulous drinks, &c., constitute the usual treatment.

HÆMATENCEPHALUM, Apoplexy.

HÆMATEPAGOGÉ, Congestion.

HÆMATERA, Hepatirrhœa.

HÆMATERÛS, Sanguine.

HÆMATERYTHRUM, Hæmatin.

HÆMATEXOSTOSIS, Osteosarcoma.

HÆMATHIDROSIS, *Sudor cruentus*.

HÆMATHORAX, Hæmatothorax.

HÆMATIO, Sanguine—h. Crystals, Hæmatoidin.

HÆMATIASIS, Hæmatonosus.

HÆMATICA, from *hæma*. Diseases of the sanguineous function:—the third class in the nosology of Good. Also, medicines that act on the blood.—Pereira.

HÆMATICA DOCTRINA, Hæmatology.

HÆMATICUS, Sanguine.

HÆMATIDROSIS, *Sudor cruentus*.

HÆMATIN, *Hæmat'ina*, *Hem'atine*, *Hæ'matoin*, *Hem'atoin*, *Hæmater'ythrum*, *Hæmer'ythrum*, *Zoohem'atin*, *Hemachro'in*, *Hematoero'ine*, *Gl'i'adine*, *Phæ'nodine*, *Phæ'nicine*, *Hæmatochro'ite*, *Cru'orin*, *Rubrin*, *Glob'ulin* of some; *Rouge du Sang*; from *hæma*. The red colouring matter of the blood. It resides in distinct particles or globules, and, in the opinion of some observers, in the envelope of the globules. It appears to be of a peculiar character, and one that has not yet been determined by the chemist. That the colour of the blood is not owing to the peroxide of iron which it contains is shown by the fact mentioned by Scherer, that he removed the iron by acids, and yet a deep red tincture was formed when alcohol was added to the residuum.

HÆMATIN'IC, *Hæmatin'icus*: from *hæmatin*, 'the red colouring matter of the blood.' An agent that augments the number of red corpuscles of the blood.—Pereira.

HÆMATINIC CRYSTALS, Hæmatoidin.

HÆMATINUS, Sanguine.

HÆMATIS'CHESIS, *Hæmis'chesis*, *Hæmocyph'ia*, from *hæma*, and *σχυν*, 'to suppress.' The retention or suppression of a natural or artificial flow of blood.

HÆMATISMUS, Epistaxis, Hæmorrhagia.

HÆMATISTIMUS, from *hæma*, and *ισμος*, 'pharynx.' Hemorrhage into the pharynx.

HÆMATYTES, *Hæmat'y'tis*, from *αἷμα*, 'blood,' Resembling blood. *Lapis Hæmat'y'tes*. A beautiful ore of iron, called also, *Bloodstone*, *O'chrea rubra*, *Oxydum fer'ricum crystallizatum nativum*, *Ferrum Hæmatites*. When finely levigated, and freed from the grosser parts, by frequent washings with water, it has been long recommended in hæmorrhage, fluxes, uterine obstructions, &c., in doses of from one scruple to three or four. Also, a vessel that contains blood, *αἱματ'ις φλεψ*.

HÆMATMUS, *Gaz sanguinis*.

HÆMATOCATHARTICA, from *hæma*, and *καθάρσις*, 'purification or purgation.' Remedies for purifying the blood.

HÆMATOCE'LE, from *αἷμα*, *αἷματος*, 'blood,' and *κηλη*, 'tumour.' A tumour formed by blood. By some, this term has been applied to a tumour formed by blood, effused into the areolar texture of the scrotum. Others have used it for tumours arising from effusion of blood into the tunica vaginalis:—hæmatocele, according to them, differing from hydrocele, only in the character of the effusion.—Heister. Others, again, have applied it to effusions of blood into the interior of the tunica albuginea itself.—Richter. The first is the usual acceptance. It is most commonly caused by wounds or contusions; and requires the use of antiphlogistics, discentients, &c. Sometimes it is necessary to evacuate the effused blood.

HÆMATOCELE ARTERIOSA, Aneurism.

HÆMATOCEPH'ALUS, (F.) *Hæmatoceph'ale*. A monster in which an effusion of blood on the brain has given occasion to more or less singular deformities.

HÆMATO-CEREBRIFORM DISEASE, Encephaloid.

HÆMATOCHE'ZIA, from *hæma*, and *χεῖω*, 'I go to stool.' *Sedes cruent'a*, *Catarrh'es vero*, *Enterorrhag'ia simplex*, *Diarrhæ'a cruenta seu sanguinolenta*, *Hæ'mato-diarrhæ'a*, *Hæmorrhag'ia intestino'rum*, *Enter'o-hæmorrhag'ia*, (F.) *Hæmorrhagie des intestins*, *Écoulement de sang par l'Intestin*. Discharge of blood by stool. See Melæna.

HÆMATOCHROÏTE, Hæmatin.

HÆMATOCHYSIS, Hæmorrhagia.

HÆMATOCE'LIA, from *hæma*, and *κοιλία*, 'the cavity of the abdomen.' Effusion of blood into the abdomen.

HÆMATOCOL'PUS, from *hæma*, and *κολπος*, 'vagina.' Effusion of blood into the vagina. Accumulation of blood, owing to occlusion of the vagina.

HÆMATOCRYSTALLIN, Hæmatoidin.

HÆMATOCYANOPATHIA, Cyanopathy.

HÆMATOCYANOSIS, Cyanopathy.

HÆMATOCYSTE, Hæmatoma saccatum.

HÆMATOCYST'IS, from *hæma*, and *κυστίς*, 'bladder.' Hemorrhage into the bladder.

HÆMATO'DES, *Hæmatoi'des*, *Sanguin'eus*, *Hæ'matoid*, *Hem'atoid*, from *hæma*, and *εἶδος*, 'appearance.' That which contains blood, or has the character of blood.

HÆMATODES FUNGUS, *Hæmatomyces*, *Fungus cancro'sus hæmato'des*, *Mela'na fungo'sa carci'no-des*, *Angidiospon'gus*, *Angiomy'ces*, *Hæmatospon'gus*, *Tumor fungo'sus sanguin'eus seu anom'alus*, *Carcino'ma Hæmato'des*, *Carcino'ma Spongio'sum*, *Spon'goid inflammation*, *Pulpy* or *Med'ul-lary Sarco'ma*, *Bleeding Fungus*, *Soft Cancer*, (F.) *Carcinôme sanglant*, *Hématoncie fongoi'de*. An extremely alarming carcinomatous affection, which was first described, with accuracy, by Mr. John Burns, of Glasgow. It consists in the development of cancerous tumours, in which the inflammation is accompanied with violent heat and pain, and with fungus and bleeding excrescences. Even when the diseased part is extirpated at a very early period, recovery rarely follows; other organs being generally implicated at the same time. Fungus hæmatodes was the term first applied to the disease by Mr. Hey of Leeds. Mr. J. Burns called it *Spongioid inflammation*, from the spongy, elastic feel, which peculiarly characterizes it, and continues even after ulceration has taken place. The disease has, most frequently, been met with in the eyeball, the upper and lower extremities, testicle and mamma; but it occurs in the uterus, ovary, liver, spleen, brain, lungs, thyroid gland, and in the hip and shoulder-joint.

Some French surgeons designate, by this name, those tumours which were formerly termed *anormales*, *cavernæuses*, *variqueuses*, called *Erectiles* by Dupuytren, *Hématoncies*, by Alibert, and *Telangiectasia* by Gräfe.

HÆMATODIARRHŒA, *Hæmatochezia*.

HÆMATODYNAMOMETER, *Hæmadynamometer*.

HÆMATOGASTER, from *hæma*, and *γαστήρ*, 'stomach.' Effusion of blood into the stomach.

HÆMATOGENETICA, *Hæmatopoetica*.

HÆMATOGLOBULIN, see Globules of the blood.

HÆMATOGRA'PHIA, *Hæmatog'raphy*, from *hæma*, and *γραφη*, 'a description.' A description of the blood.

HÆMATOID, *Hæmatodes*.

HÆMATOID'IN, *Hæmatocry'stallin*, *Hæmatin'ic*, *hæmatic*, *hæmin* or *blood crystals*, (F.) *Hématoidine*, *Crystaux d'Hématique*. A substance found in the midst, or in the vicinity, of effusions of blood, which is either in amorphous masses, or, more frequently, in minute rhomboidal crystals, or in fine needles. It is considered to be hæmatosin separated from the globules of the blood, and passed from the state of a non-crystallizable organic substance to one of a crystalline chemical compound.

HÆMATOL'OGY, *Hæmatol'ogy*, *Hæmatolog'ia*, *Hæmat'ica doctri'na*, (F.) *Hématologie*, from *hæma*, and *λογία*, 'a discourse.' That part of medicine which treats of the blood.

HÆMATOLOGY, **PATHOLOG'ICAL**, (F.) *Hématologie pathologique*. Observation of the blood to detect its varying characters in disease.

HÆMATOLYSES, (G.) *Hæmatolysen*, from *hæma*, and *λυσις*, 'solution.' An order of diseases in which there is diminished coagulability of the blood.—Fuchs.

HÆMATOLYSIS, *Hæmateclysis*.

HÆMATOLYTICUM, *Spanæmic*.

HÆMATO'MA, *Thrombus*, *Tumor sanguin'eus*. A bloody tumour, especially of the scalp of the new-born.

HÆMATO'MA SACCA'TUM. An encysted tumour containing blood.—*Hæmatocyst'ic*.

HÆMATOMANT'IA, from *hæma*, and *μαντεία*, 'divination.' Judgment of disease from the appearance of the blood.

HÆMATOMATRA, *Metrorrhagia*.

HÆMATOMETACHYSIS, *Transfusio san-guinis*.

HÆMATOMETRA, *Metrorrhagia*.

HÆMATOMMA, *Hæmophthalmia*.

HÆMATOMPHALOC'E'LE, *Hæmatom'phalum*, *Hæmatompha'lus*, from *hæma*, and *ομφαλος*, 'the navel,' and *κληη*, 'a tumour.' Umbilical hernia, the sac of which encloses a bloody serum, or which has, at its surface, a number of varicose veins; constituting *Varicom'phalus*.

HÆMATOMPHALUM, *Hæmatomphalocèle*.

HÆMATOMYCES, *Hæmatodes fungus*.

HÆMATON'CUS, (F.) *Hématoncie*, from *hæma*, and *ογκος*, 'a tumour.' Alibert has given this name to *Nevi mater'ni*, *Varicose tumours*. He admits three varieties:—the *H. fongoi'de*, *H. framboisée*, and *H. tubéreuse*.

HÆMATON'OSUS, *Hæmat'osis*, *Hæmatopath'ia*, *Hæmopath'ia*, from *hæma*, and *νοσος*, 'disease.' A disease of the blood. A *hæmic* or *blood-disease*.

HÆMATOPATHIA, *Hæmatonosus*.

HÆMATOPEDESIS, see Diapedesis.

HÆMATOPERICAR'DIUM, from *hæma*, and *περικαρδιον*, 'pericardium.' Effusion of blood into the pericardium.

HÆMATOPHŒUM, *Hæmaphæin*.

HÆMATOPHIL'IA, *Hæmophil'ia*, *Hæmorrhophil'ia*, *Idiosyncra'sia hæmorrhag'ica*, from *hæma*, and *φιλεω*, 'I love.' A hæmorrhagic diathesis.

HÆMATOPHLEBES'TASIS. Sudden suppression of a hæmorrhage;—from *στασις* 'αίματος φλεβων', 'suppression of the blood of the veins.'—Galen.

HÆMATOPHOB'IA, *Hæmophob'ia*, from *hæma*, and *φοβος*, 'dread.' Dread or horror at the sight of blood, producing syncope, &c.

HÆMATOPHTHALMIA, *Hæmophthalmia*.

HÆMATOPH'THORES, (G.) *Hæmatoph'thoren*; from *hæma*, and *φθορα*, 'corruption.' An order of diseases in the classification of Fuchs, in which the blood is materially altered in composition, as in typhus.

HÆMATOP'ISIS, (F.) *Hématopisie*. M. Capuron, of Paris, has applied the term *Hématopisie utérine*, from analogy with *Hydropisie utérine*, to a collection of blood which sometimes takes place in the uterus, when, owing to faulty conformation, the exit of the menstrual flux is prevented.

HÆMATOPLA'NIA, *Hæmatoplane'sis*, from *hæma*, and *πλάνη*, 'wandering.' A vicarious hæmorrhage.

HÆMATOPLANIA MENSTRUALIS, *Menstruation*, vicarious.

HÆMATOPLASMA, *Liquor sanguinis*.

HÆMATOPLETHORA, *Plethora*.

HÆMATOPŒA, *Hæmatopoetica*.

HÆMATOPOESIS, *Hæmatosis*.

HÆMATOPOËT'ICA, *Hæmatopœ'a*, *Hæmatogenet'ica*, 'bloodmakers,' from *hæma*, and *ποιεω*, 'I make.' Agents that favour hæmatosis. Iron was called by the ancients *met'al'um hæmatopœt'icum*.

HÆMATOPOIESIS, *Hæmatosis*.

HÆMATOPORIA, *Hæmaporia*.

HÆMATOPS, *Hæmophthalmia*.

HÆMATOPSIA, *Hæmophthalmia*.

HÆMATOPTYSIS, *Hæmoptysis*.

HÆMATOPTYSIS, *Hæmoptysis*.

HÆMATORRHACHIS, *Apoplexia myelitia*.

HÆMATORRHAGIA, *Hæmorrhagia*.

HÆMATORRHŒA, *Hæmorrhagia*.

of a florid red. In such active hemorrhages, the great indications of treatment will be, to diminish plethora where it exists, and to lessen the heart's action. Bleeding, purgatives, and cold, will be the chief agents.

Passive Hemorrhage, Hæmorrhag'ia passiva seu *veno'sa, Profusio, P. hæmorrhag'ica*, occurs in cases of weak constitution; or who have been debilitated by protracted disease, poor diet, long watching, excessive evacuations, &c.

The direct causes may be:—previous active hemorrhage; scorbutus, or any thing capable of inducing atony or asthenic hyperæmia of the small vessels. These hemorrhages are not preceded by excitement or by any signs of local determination. They are usually accompanied by paleness of the countenance; feeble pulse; fainting, &c. The indications of treatment will be:—to restore the action of the small vessels and the general tone of the system; hence the utility of styptics and cold externally; and of tonics and astringents, creasote, mineral acids, &c., internally. Hemorrhage also occurs from mechanical hyperæmia, as when hæmoptysis is produced by tubercles in the lungs; hæmatemesis by disease of some of the solid viscera of the abdomen, &c.

In *Traumatic Hemorrhages*, or those which are the consequences of wounds of arterial or venous trunks, the blood is of a florid red colour, and issues by jets and pulses, if it proceed from an artery; whilst it is of a deeper red, issues slowly and by a continuous flow, if from a vein. If the capillary vessels be alone divided, the blood is merely effused at the surface of the wound. Of the means used for arresting these traumatic hemorrhages, some act mechanically as *absorbents, ligature, and compression*; others chymically, as *fire, caustics, creasote, astringents, &c.*

HÆMORRHAGIA ACTIVA NARIUM, Epistaxis—*h. Bronchica, Hæmoptysis*—*h. Cerebri, Apoplexy*—*h. per Cutem, Sudor cruentus*—*h. per Diæresin, see Diæresis*—*h. per Diapedesin, Diapedesis*—*h. Faucium, Stomatorrhagia*—*h. Gingivarum, Ulorrhagia*—*h. Hæmatemesis, Hæmatemesis*—*h. Hæmaturia, Hæmaturia*—*h. Hæmoptysis, Hæmoptysis*—*h. Hepatica, Hepathæmorrhagia*—*h. Hepatis, Hepatorrhagia*—*h. Intestinorum, Hæmatochezia*—*h. Mucosa, see Hæmorrhoids*—*h. Nabothi, see Parturition*—*h. Narinae, Epistaxis*—*h. Narium, Epistaxis*—*h. Oris, Stomatorrhagia*—*h. Penis, Stimatosis*—*h. Pulmonis, Hæmoptysis*—*h. Renum, Nephorrhagia*—*h. Universalis, Purpura hæmorrhagica*—*h. Uterina, Metorrhagia*—*h. Ventriculi, Hæmatemesis*—*h. Vesicæ, Cystorrhagia.*

HÆMORRHINIA, Epistaxis.

HÆMORRHINORRHAGIA, Epistaxis.

HÆMORRHŒ'A, from *hæma*, and *rho*, 'I flow.' *Hæmorrhag'ia*, Loss of blood. Some writers have proposed to restrict this name to passive hemorrhages.

HÆMORRHŒA PETECHIALIS, Purpura hæmorrhagica—*h. Pulmonalis, Hæmoptysis*—*h. Uterina, Metorrhagia*—*h. Vasorum hæmorrhoidalium, Hæmorrhoids*—*h. Ventriculi, Hæmatemesis*—*h. Viarum urinarium, Hæmaturia.*

HÆMORRHŒDAL, HæmorrhŒdal, HæmorrhŒdal'is, HæmorrhŒdew. Relating to hemorrhoids; as *hemorrhoidal flux, hemorrhoidal tumours, &c.*

HÆMORRHOIDAL ARTERIES have been distinguished into *superior, middle, and inferior*. 1. The first is the termination of the inferior mesenteric artery, which assumes the name *superior hemorrhoidal*, when it reaches the upper and posterior part of the rectum. 2. The *middle hemorrhoidal* is furnished by the hypogastric or internal pudic. It ramifies on the inferior and anterior part of the rectum. 3. The *inferior hemor-*

rhoidal arteries are branches of the internal pudic, furnished to the inferior part of the rectum and to the muscles of the anus.

HÆMORRHOIDAL NERVES. These emanate from the sciatic and hypogastric plexuses; and cover the rectum with their numerous filaments.

HÆMORRHOIDAL VEINS follow the same distribution, and generally empty themselves into the lesser mesenteric. Some of them assist in the formation of the hypogastric vein.

HÆMORRHOIDAL VESSELS, Vasa sed'alia, are those vessels which are distributed to the rectum—the seat of hemorrhoids.

HÆMORRHOIDALES NODI, Hæmorrhoids.

HÆMORRHIDES, see *Hæmorrhoids*—*h. Cæcæ*, see *Hæmorrhoids*—*h. Fluentes, Hæmorrhoids*—*h. Furentes*, see *Hæmorrhoids*—*h. Mariscosæ, Hæmorrhoids*—*h. non Fluentes, Hæmorrhoids*—*h. Oris, Stomatorrhagia.*

HÆMORRHŒDRO'SIS, from *ἡμorrhῶσις*, 'hemorrhoids,' and *ῥῶσις*, 'sweating.' Sweating of hemorrhoidal blood. [?]—Bock.

HÆMORRHOIS, Aïmor'rhois, Asclepias'mus, from *hæma*, and *rho*, 'I flow.' *Aïmor'rhois, Proctica Maris'ca, Maris'ca, Proctul'gia HæmorrhŒdal'is, Morbus HæmorrhŒdal'is, Piles, Em'erods, (F.) Hæmorrhoides.* The essential symptoms of this affection are:—Livid and painful tubercles or excrescences, (*HæmorrhŒdal Tumours*), usually attended with a discharge of mucus or blood, (*HæmorrhŒdal flux, Proctorrhœ'a, Hæmoproctia, Hæmorrhœ'a vasorum hæmorrhŒdalium, (F.) Hæmoproctie.*) The most common causes of piles are a sedentary life; accumulation of feces in the rectum; violent efforts at stool; pregnancy, &c. The precursory symptoms are:—pains in the loins; stupor of the lower limbs; and uneasiness in the abdomen and rectum, with more or less gastric, cerebral, and indeed general disorder;—constituting the *Diathesis HæmorrhŒdal'is, Motus seu Turba HæmorrhŒdal'is*, and *Molimen HæmorrhŒdal'is*, of most of the writers of Continental Europe. To these symptoms follow one or more round, smooth, renitent, painful, pulsating, and erectile tumours, around the margin of the anus, or within the anus; some pouring out blood occasionally. After having remained, for a time, tense and painful, they gradually shrink and disappear. The chief symptoms, occasioned by hemorrhoidal tumours, when much inflamed, are:—constant pain, liable, however, to exacerbations, and to become augmented by the least pressure, or by the passage of the feces, and obliging the patient to preserve the horizontal posture.

Hæmorrhoids have generally been distinguished into—*Hæmorrhoides Fluentes seu Mariscosæ, Proctica Maris'ca cruent'a, Bleeding or Open Piles*; and into *H. non fluent'es, Proctica Maris'ca cæca, HæmorrhŒdes cæcæ seu furent'es, HæmorrhŒdal'es nodi, Shut or blind piles.* They have, also, been divided into *internal or occult*, and *external*, according to their situation; and into *accidental or constitutional.*

Hæmorrhoidal Tumours are extremely troublesome, by their disposition to frequent recurrence; and they are apt to induce fistula; otherwise, they are devoid of danger. When anatomically examined, they are found not to consist in a varicose dilatation of the veins of the rectum; but to be formed of a very close, spongy, texture; similar to that which surrounds the orifice of the vagina; and to be erectile, like it. They are surrounded by a delicate membrane, and have no internal cavity. The treatment, in mild cases of hemorrhoidal tumours, is simple. Rest; the horizontal posture; the use of mild laxatives, as sulphur, castor oil, and emollient gylsters, will

be sufficient. If they be much inflamed, leeches may be applied; and warm cataplasms or cold lotions, according to circumstances, be prescribed, with abstinence, and cooling drinks. Afterwards, an ointment, composed of powdered galls and opium, may afford relief. It is in the relaxed kind, that such ointment, and the internal use of *Ward's Paste*, can alone be expected to afford much benefit. If, after repeated attacks, the tumours remain hard and painful, and threaten fistula, they may be removed.

By *Hæmorrhoidal Flux*, *Fluxus hæmorrhoidalis*, *Proctorrhag'ia*, is meant the hæmorrhage which takes place from the rectum, owing to hæmorrhoids. It is a common affection. The quantity of blood discharged is various: at times, it is very trifling; at others, sufficient to induce great debility, and even death. It is announced and accompanied by the same symptoms as precede and attend hæmorrhoidal tumours. Like other hæmorrhages it may be *active* or *passive*; *accidental* or *constitutional*. The prognosis is rarely unfavourable. The affection may, almost always, be relieved by properly adapted means. These resemble such as are necessary in hæmorrhages in general. Perfect quietude,—mental and corporeal; light diet; cooling drinks; bleeding if the symptoms indicate it; astringents, (if the disease be protracted and passive), such as the *Tinctura Ferri Chloridi*; aspersions of cold water on the anus: astringent injections; plugging and compression. Such will be the principal remedial agents. When the hæmorrhage has become habitual, or is vicarious, some caution may be required in checking it; and, if inconvenience arise from a sudden suppression, its return may be solicited by the semicupium, sitting over warm water, aloetic purgatives, glysters, irritating suppositories, &c.; or leeches may be applied to the anus.

To the internal bleeding pile, a soft, red, strawberry-like elevation of the mucous membrane, Dr. Houston, of Dublin, gives the name *vascular tumour*. For its removal he recommends the application of nitric acid, so as to produce sloughing of its surface.

Some authors have described a species of *Leucorrhœa Ana'tis* or whitish discharge from the anus, which often attends ordinary hæmorrhoids. This they have called *Proctica maris'ca mucosa*, *Hæmor'rhois alba*, *Hæmorrhag'ia mucosa*, *Leucor'rhois*, &c. It requires no special mention.

The term *Hæmorrhoid* is applied at times to varicose veins developed elsewhere, as at the orifice of the uterus. See *Metrorrhœides*.

HÆMORRHOIS AB EXANIÀ, Proctocœle—h. Proccœdes, Proctocœle.

HÆMORRHOÏS'CHESIS, from *ἄιρρῳπος*, *hæmorrhôis*, *pew*, 'I flow,' and *ἄρξω*, 'I restrain.' *Reten'tio fluxus hæmorrhoidalis*. Suppression or retention of the hæmorrhoidal flux.

HÆMORRHOPHE'SIS, *Absorp'tio sang'uinis*, from *hæma*, and *ρῶσσω*, 'I sip up.' Absorption of blood.

HÆMORRHOPHILIA, Hæmatophilia.

HÆMORRHOSCOPIA, Hæmatoscopia.

HÆMORRHYSIS, Hæmorrhagia.

HÆMOSCHEOCELE, Hæmatoschœcele.

HÆMOSCOPIA, Hæmatoscopia.

HÆMOSPASIA, see Hæmospastic.

HÆMOSPAS'TIC, (F.) *Hémospasique*, from *hæma*, and *σπασω*, 'I draw.' An agent which draws or attracts blood to a part; as a cupping-glass. The operation is termed *Hæmospa'sia*, (F.) *Hémospasie*. It is generally applied to a process by which the air is exhausted over a considerable surface, as over one or more of the extremities, by an appropriate pneumatic apparatus.

HÆMOSTA'SIA, *Hæmostasis*, *Epid'esis*, *San'guinis stagna'tio*, (F.) *Hémostasie*, from *hæma*, and *στασις*, 'stagnation.' Stagnation of blood. This name has, also, been given to any operation, the object of which is to arrest the flow of blood.

HÆMOSTATIC, PAGLIA'RI'S, *Pagliari's Styptic*, (F.) *Eau de Pagliari*. A styptic liquor, said to be of great power, discovered by M. Pagliari, a pharmacien at Rome. It is composed as follows:—Eight ounces of tincture of benzoin, one pound of *alum*, and ten pounds of water are boiled together for six hours in a glazed earthen vessel, the vaporized water being constantly replaced by hot water, so as not to interrupt the ebullition, and the resinous mass being kept stirred round. The fluid is then filtered, and kept in stoppered bottles. A drop of the fluid, poured into a glass containing human blood, produces an instantaneous magma.

HÆMOSTATICS, Hæmostatica.

HÆMOTELANGIOSIS, Telangiectasia.

HÆMOTEXIA, Hæmatecylis.

HÆMOTEXIS, Hæmatecylis.

HÆMOTHORAX, Hæmathorax.

HÆMOTISMUS, Hæmoptysis.

HÆMOTROPHY, *Hæmotroph'ia*; from *hæma*, and *τροφή*, 'nourishment.' Excess of sanguineous nourishment.—Prout.

HÆMURIA, Hæmaturia.

HÆMURESIS, Hæmaturia.

HÆMYDOR, Serum of the blood.

HÆMYDRIA, Hydroæmia.

HÆREDITARIUS, Hereditary.

HÆSITATIO, Balbuties.

HÆVEA GUANENSIS, see Caoutchouc.

HAG, Abdomen.

HAGARD, Haggard.

HAGBERRY, *Prunus padus*.

HAG'NIA ABYSSIN'ICA, *Bray'era anthelmintica*, *Bank'sia Abyssin'ica*. An Abyssinian tree; Order, Rosacæ; Sexual System, Icosandria Digynia, which the natives plant round their habitations, as an ornament. The infusion or decoction of its flowers, *Flores cosso*, is employed by them as an anthelmintic, especially in cases of tape-worm. It is called, there, *Cusso* or *Cosso*, *Kosso*, *Kousse*, *Kosso*, *Kwoso*, *Cotz*, *Cabota*, and *Habbi*.

HAGGARD, (F.) *Hagard*. The French use the term *Air hagard*, *Œil hagard*, 'Haggard air,' 'Haggard eye,' for a physiognomy, in which there is at once an expression of madness and terror.

HAGIOSPERMUM, *Artemisia santonica*.

HAGIOXYLUM, Guaiacum.

HAHNEMANNISM, Hæmœopathy.

HAIL, Chazala.

HAIR, Sax. *hær*, *Crinis*, *Pilus*, *Thrix*, *Pile*, (F.) *Poil*. A conical, corneous substance, the free portion or *shaft*, *scapus*, of which issues to a greater or less distance from the skin, to the tissue of which it adheres by a bulb, *Bulbus pili*, seated in a *hair follicle*—*follic'ulus pili*—made by an inversion of the integument, the epidermis of which forms a "root sheath," *vagi'na pili*. At the base of the hair follicle, there is a small papilla, well supplied with bloodvessels and nerves, *Papilla pili*; at times called, but improperly, *Pulpa seu Bliste'ma pili*. The hair receives various names in different parts—as *Beard*, *Cilia*, *Eyebrows*, *Hair of the head*, (*Capilli*), &c.

HAIR DYE, TWIGGS'S. A preparation much used to give colour to the hair. It may be formed of *precipitated sulphur* and *acetate of lead*, each ʒi; *rose water*, ℥iv; triturated together in a mortar. It should be applied twice a day, until the desired effect is produced.

HAIR, FALLING OFF OF THE, Alopecia.

HAIR FOLLICLE, see Hair.

HAIR, MATTED, Plica—h. Plaited, Plica—h. Trichomatose, Plica—h. Worm, Seta equina, see Ectozoa—h. Worm, muscular, Dracunculus.

HAIRY, Pileous. Also, covered with hair.

HAKE, see Oleum jecoris aselli.

HAL, Salt.

HALA'TION, *Hala'tium*; diminutive of 'als, 'salt.' A pungent remedy, with the ancients, which contained salts.

HALCHEMIA. The alchemists so called the art of fusing salts: from 'als, 'salt,' and $\chi\epsilon\mu\alpha$, 'I pour out.'—Libavius.

HALCYON, Alcyon.

HALCYONIUM ROTUNDUM, *Pila marina*.

HALE, Ephelides.

HALEINE, Breath—h. *Courte*, Dyspnoea.

HALELE'ON, *Halele'um*, from 'als, 'salt,' and $\epsilon\lambda\epsilon\omega$, 'oil.' *Oleum Sali mixtum*. A mixture of oil and salt, for removing swellings of the joints.—Galen.

HALÉTÉRATION, Alteration.

HALF-CASTE, *Half-Cast*, see Caste. A term applied, in India, to the offspring of a Hindoo and a European. Since, extended to the offspring of mixed races. The subjoined list from Tschudi's Travels in Peru, 1838–1842, shows the parentage of the different varieties of half-casts, and also the proper designations of the latter, as observed in South America.

PARENTS.	CHILDREN.
White Father and Negro Mother,	Mulatto.
White Father and Indian Mother,	Meztizo.
Indian Father and Negro Mother,	Chino.
White Father and Mulatta Mother,	Quarteron.
White Father and Meztiza Mother	Creole, (only distinguished from the white by a pale brownish complexion.)
White Father and China Mother,	Chino-blanco.
White Father and Quarterona Mother.....	Quintero.
White Father and Quintera Mother.....	White.
Negro Father and Mulatta Mother,	Zambo-negro.
Negro Father and Meztiza Mother,	Mulatto oscuro.
Negro Father and China Mother,	Zambo chino.
Negro Father and Zamba Mother,	Zambo-negro (perfectly black.)
Negro Father and Quarterona or Quintera Mother.....	Mulatto (rather dark.)
Indian Father and Mulatta Mother.....	China-oscuro.
Indian Father and Meztiza Mother.....	Mestizo-claro (frequently very beautiful.)
Indian Father and China Mother,	Chino-cholo.
Indian Father and Zamba Mother.....	Zambo-claro.
Indian Father with China-chola Mother.....	Indian (with rather short frizzy hair.)
Indian Father and Quarterona or Quintera Mother.....	Meztizo (rather brown.)
Mulatto Father and Zamba Mother.....	Zambo (a miserable race.)
Mulatto Father and Meztiza Mother.....	Chino (of rather clear complexion.)
Mulatto Father and China Mother,	Chino (rather dark.)

See Mulatto.

HALICA, Alica.

HALICACALUM, *Physalis*.

HALICES, Paudiulation.

HALIDRYS VESICULOSA, *Fucus vesiculosus*.

HALINA'TRUM, *Halini'trum*, *Haloni'trum*; from 'als, 'salt,' and *natrum* or *natron*. A name given by the ancients to subcarbonate of soda, containing a little subcarbonate of ammonia, which is found, ready-formed, on the plaster of damp walls, in places inhabited by man or animals. Also, Potassæ nitras.

HALINITRUM, *Halinatrum*.

HALINUS, Saline.

HALITUOUS, *Halituous*, from *halitus*, 'vapour.' (F.) *Halitueus*. The skin is said to be *halitueus*, when covered with a gentle moisture. The vapour, exhaled in all the cavities of the body, so long as the blood is warm, is called *Hal'itus*. The odorous vapour, exhaled by the blood itself whilst warm, is called *Hal'itus Sanguinis*.

HALITUS, Breath—h. *Oris Fœtidus*, Breath, offensive—h. *Sanguinis*, *Gaz sanguinis*. See *Halituous*.

HALL, MINERAL WATERS OF. The springs of Hall, in Upper Austria, contain iodine.

HALLUCINATIO HYPOCHONDRIASIS, Hypochondriasis—h. *Vertigo*, *Vertigo*.

HALLUCINATION, *Hallucina'tio*, *Halucina'tio*, *Falla'cia*, *Illu'sto*, *I. sensus*, *Socor'dia*, *Allu'sia*, *Allucina'tio*, *Parora'sis*, *Somnia'tio* in statu vig'ili, *Ido'lum*, *Waking dream*, *Phantasm*, from *allucinari* or *hallucinari*, *allucinatus*, 'to be deceived.' A morbid error in one or more of the senses. Perception of objects, which do not in fact exert any impression on the external senses. Hallucination or *delusion* almost always, if not always, depends on disorder of the brain, but is not an index of insanity, unless the patient believes in the existence of the subject of the hallucination.

HALLUS, Pollex pedis.

HALLUX, Pollex pedis.

HALME, Muria.

HALMYRIS, see Muria.

HALMYRO'DES, *Salsugino'sus*, *Salen'ginous*, from $\alpha\lambda\mu\upsilon\pi\iota\varsigma$, ($\alpha\lambda\mu\eta$, 'sea-water,' 'als, 'salt,') 'a salt liquor,' saltish. An epithet given to any affection, in which the heat feels pungent under the finger of the physician.—Hippocrates.

HALMYRUS, Saline.

HALO, Areola.

HALO SIGNA'TUS. The impression made by the ciliary processes on the anterior surface of the vitreous humour. So called from its consisting of a circle of indentations.—Sir C. Bell. Called by Haller, *Stria retinæ subjectæ ligamento cilia'ri*; by Winslow, *Sulci cilia'ria*.

HALODERE'UM, from 'als, 'salt,' and $\delta\epsilon\rho\alpha\iota\omega$, 'a collar,' ($\delta\epsilon\rho\eta$, 'neck,') *Colla'rium sali'num*. A collar of salt applied to the neck, as in cases of croup.

HALOGENE, Chlorine.

HALODUM OXYGENATUM, Potassæ murias hyperoxygenatus.

HALONITIS, Induration of the cellular tissue.

HALONITRUM, *Halinatrum*.

HALOPE'GÆ, from 'als, 'alos, 'salt,' and $\pi\eta\eta\eta$, 'a spring.' Mineral waters whose chief ingredient is common salt—chloride of sodium.

HALOS, Areola.

HALOTYLE, *Cul'cita sali'na*, from 'als, 'salt,' and $\tau\upsilon\lambda\eta$, 'a bolster.' A bolster or pillow of salt, recommended in croup.

HALSBANE, Clavicle.

HALS, 'als, 'salt,' hence *Halogene*, *Haloid*, &c.

HALSE, Throat.

HALTERES. Pieces of lead held in the hands by the ancients to assist them in leaping.

HALUCINATIO, Hallucination.

HALYCODES, Saline.

HALYSIS LATA, *Bothriocephalus latus*—h. *Membranacea*, *Bothriocephalus latus*—h. *Solium*, *Tenia solium*.

HAM, Poples.

HAMAME'LIS VIRGINIA'NA, *H. Virgin'ica* seu *macrophy'la*, *Witch hazel*, *Winter witch hazel*, *Snapping hazelnut*, *Winter bloom*. Ord. *Hamelaceæ*. The bark of this tree, which is

a native of the United States, is somewhat bitter and sensibly astringent; but it has not been much used. A cataplasm of the inner rind of the bark is said to have been found efficacious in painful inflammation of the eyes.

HAMARTHRITIS, *Holarthrit'is*, *Catholarthrit'is*, *Arthrit'is universal'is*, from *ἄμα*, 'at once,' and *ἄρθριτις*, 'gout.' Gout in all the joints. Universal gout.

HAMCH, Coxofemoral articulation.

HAM'MA, *Nodus*, 'a tie.' A knot, used for retaining bandages on any part. — Hippocrates. A truss.

HAMMERING, Balbuties.

HAMPSTEAD, MINERAL WATERS OF. These waters, situate in the neighbourhood of London, are a good chalybeate.

HAMSTRING, To, see Hamstrings.

HAMSTRINGS, (F.) *Jarretiers*. The strings or tendons of the ham.

The **HAMSTRING MUSCLES** are the biceps femoris, whose tendon forms the *outer hamstring*; and the semimembranosus, semitendinosus, gracilis and sartorius, whose tendons form the *inner hamstring*.

'To hamstring' means to cut the strings or tendons of the ham.

HAMULAR PROCESS, see Hamulus.

HAM'ULUS. Diminutive of *hamus*, 'a hook.' A hook or crook: *Ancis'tron*. Also, any hook-like process; as the *hamulus* or *hamular process* of the pterygoid process of the sphenoid bone, which projects from the termination of the internal pterygoid plate.

HAMULUS LAMINÆ SPIRALIS, see Lamina spiralis.

HAMUS, see Hamulus.

HANCHE, Haunch.

HANCLETH, Malleolus.

HAND, Manus.

HANDWOMAN, Midwife.

HANDS, DROPPED, *Hand-drop*, *Wrist-drop*. A popular term for the paralysis of the hand, induced by the action of lead.

HANDWRIST, Carpus.

HANGERS, Crusta genu equinæ.

HANGNAIL, (old Eng.) *Agnail*, (Prov.) *Ang-nail*, *Stepmother's blessing*, *Wirt-springs*, *Nail-spring*, *Nang-nail*, (F.) *Envie*. A portion of epidermis, detached so as to tear the integument in the vicinity of the finger nails.

HANNEBANE, Hyoscyamus.

HAPANTIS'MUS. Badly formed from *πας*, or *πας*, 'all.' *Oblitera'tio comple'ta*. The matted or growing together of organic parts.

HAPHÈ, *ἅψη*, 'feeling, touch.' Hence:

HAPHON'OSI, *Haphon'usi*, from *ἅψη*, 'the touch.' *Morbi tactūs*. Diseases of the sense of touch.

HAPLOACNE, Aene simplex.

HAPLOPATH'IA, from *ἁπλος*, 'simple,' and *πάθος*, 'disease.' *Morbus simplex*. A simple or uncomplicated affection.

HAPLOTOM'IA, *Simplex rec'tio*, *Incis'io simplex*, *Opera'tio simplex*, from *ἁπλος*, 'simple,' and *τομή*, 'incision.' A simple incision.

HAPSIS, Touch.

HAPTODYSPHOR'IA, *Tactus dolorificus*; from *ἅψη*, 'touch,' and *δυσφορος*, 'difficult to be borne.' Painful to the touch.

HAPTOGEN, Haptogenous.

HAPTOGENÈ, Haptogenous.

HAPTOG'ENOUS, *Haptog'ennus*, *Hapt'ogen*, (F.) *Haptogène*, from *ἅπτω*, 'I attach,' and *γεννῶ*, 'I engender.' Ascherson calls the pellicle, which forms around fatty matter when albumen is brought in contact with it, *Haptogen membrane*, (F.) *Membrane haptogène*.

HAPTOT'ICA, *Haptot'icè*, from *ἅψη*, 'touch.' The doctrine of the phenomena of touch.

HARD, (G.) *hart*, *Durus*, *Scleros*, (F.) *Dur*. That which offers much resistance. In anatomy, the *hard parts* are those which compose the osseous basis of the body; in other words, the skeleton. See Pulse, hard.

HARD'S FARINACEOUS FOOD, see Farinaceous.

HARDESIA, Hibernicus lapis.

HARDHACK, *Spiraea tomentosa*.

HARE-LIP, *Lagochi'lus*, *Lagos'toma*, *Lagon'tomum*, *Lagen'tomum*, *La'bium Lepor'num*, *Olopho'nia Labii Loba'ta*, *Lepor'num rostrum*, (Sc.) *Hareshaw*, (Prov.) *Hayscale*, (F.) *Bec de Lièvre*. A fissure or perpendicular division of one or both lips. It has been so called, in consequence of the upper lip of the hare being thus divided. Hare-lip is generally congenital; at other times it is *accidental*, or produced by a wound, the edges of which have not been brought into contact, and have healed separately. It is *simple*, when there is but one division; *double*, when there are two; and *complicated*, when there is, at the same time, a division or cleft of the superior maxillary bone and of the palate — *Cleft* or *fissured palate*, *Wolf's jaw*, *Palat'um fissum*, *Rictus lupi'nus*; or a projection of the teeth into the separation of the lip.

In the *Hare-lip operation*, there are two indications to be fulfilled. First, to pare, with the knife or scissors, the edges of the cleft, and, afterwards, to preserve them in contact; in order to cause adhesion. This last object is accomplished by means of pins, passed through the edges of the division; in other words, by the twisted suture. The projecting teeth must, of course, be previously removed. If there be separation of the palate, it will become less and less after the union of the lip, or the operation of staphyloplasty may be performed upon it.

HARE'S EAR, *Eupleurum rotundifolium* — *h.* Eye, *Lagophthalmia*.

HARESHAW, Harelip.

HARGNE, *Hernia* — *h.* *Anévrismale*, *Cirsomphalus*.

HARICOT, *Phaseolus vulgaris* — *h.* *Grand de Perou*, *Jatropha curcas*.

HARMALIA, Aliment.

HARM'ONY, *Harmo'nia*, *Harmos*, primarily from *ἁρμ*, 'I adjust.' Anatomists have called *Suture by Harmony*, or simply *Harmony*, *False* or *superficial suture*, *Sutura os'sium spu'ria*, *Compa'ges os'sium per lin'eam simp'licem*, an immovable articulation, in which the depressions and eminences, presented by the bony surfaces, are but slightly marked; so that it might be presumed that the junction of the bones took place by simple apposition of their surfaces. An instance of harmony occurs in the union of the superior maxillary bones with each other.

HARMUS, Articulation, Harmony.

HARNPAN, Cranium.

HARRODSBURG SPRINGS, see Kentu ky, Mineral Waters of.

HAR'ROWGATE, MINERAL WATERS OF. The villages of High and Low Harrowgate are situate in the centre of the county of York, near Knaresborough, twenty miles from York, and fifteen from Leeds. Here are several valuable sulphurous and chalybeate springs. The *sulphureous springs* contain chloride of sodium, chloride of calcium, chloride of magnesium, bicarbonate of soda, sulphohydric acid, carbonic acid, carburetted hydrogen, and azote. The *chalybeate springs* contain protoxide of iron, chloride of sodium, sulphate of soda, chloride of calcium, chloride of magnesium, carbonic acid, azote, and oxy-

gen. For the former, Harrowgate is celebrated and frequented.

HARROWGATE SALTS, ARTIFICIAL, are much employed, and not unfrequently by those who drink the genuine water, for the purpose of increasing its aperient power. They may be made as follows:—*Sulph. Potass. cum Sulph.* 3vj; *Potass. bitart.* 3j; *Magnes. Sulph.* in pulv. 3vj. M. The usual dose is a teaspoonful, in a small tumblersful of tepid water, early in the morning.

HARROWGATE WATER, ARTIFICIAL, may be formed of common salt, 3v; water, Oij; impregnated with the gas from sulphuret of potass, and sulphuric acid, aa 3iv. The following form has also been recommended. *R. Sulphat. Potass. cum sulph.* (Ph. Ed.) 3j; *Potass. bitart.* 3ss; *Magnes. sulphat.* 3vj; *Aqua destillat.* Oij. One-half to be taken for a dose.

HARTFELL, MINERAL WATERS OF. Hartfell Spa is about five miles from Moffat, in Scotland. The water is a chalybeate, and is much used.

HARTSHORN, Cervus, Cornu cervi, Liquor cornu cervi—h. Red, Spiritus lavandulæ compositus—h. and Oil, Linimentum ammoniæ carbonatis.

HARTS' TONGUE, Asplenium scolopendrium.

HARTS' TRUFFLES, Elaphomyces granulatus.

HARTWORT OF MARSEILLES, Seseli tortuosum.

HASCHE, Pain.

HASCHISCH, see Bangué.

HASCHISCHINE, see Bangué.

HASTA, Penis—h. Nuptialis, Penis—h. Virilis, Penis.

HASTECLA, Splint.

HASTINGS, CLIMATE OF. This place has the reputation of being one of the mildest and most sheltered winter residences on the south coast of England. Owing to its low situation, and the height of the neighbouring cliffs, it is protected in a great degree from all northerly winds; and hence is found a favourable residence generally for invalids labouring under diseases of the chest.

HASTULA REGIS, Asphodelus ramosus.

HATFIELD'S TINCTURE, see Tinctura Guaiaci ammoniata.

HATTREL, Abscess.

HAUNCH, *Coxa*, *Coxen'dix*, of the Latins; *αἰχμή*, or *ισχίον*, of the Greeks; *Ancha*, *Ischion*, *Osphys*, *Hip*, *Cossa*, (Prov.) *Hubbon*, *Shoup*, (F.) *Hauche*. The region of the trunk which is formed by the lateral parts of the pelvis and the hip-joint, including the soft parts. In women, on account of the greater width of the pelvis, the haunches are more marked and prominent than in men.

HAUSE, Collum, Throat.

HAUSTEL/LATE, *Haustella'tus*, from *haurio*, *haustum*, 'I drink up.' An epithet for the structure of mouth, which is adapted for drinking or pumping up liquids. Insects, which possess that kind of mouth, are so named. Ehrenberg refers the fancied spermatozoon to the haustellate entozoa.

HAUSTRA COLI, see Colon.

HAUSTUS, *Potio*, *Potiu'n'cula*. A Draught. A liquid medicine, which can be taken at a draught.

HAUSTUS NIGER, see Infusum sennæ.

HAUT MAL, Epilepsy.

HAVAN'A or HAVAN'NAH, CLIMATE OF. The climate of Cuba is often selected for the phthisical invalid during the winter months, and so far as regards elevation and comparative equality of temperature, it is more favourable for those of weak lungs than that of the United States. The mean annual temperature is high

(78°), but the difference between the mean temperature of the warmest and coldest months is twice as great as at Madeira.

HAVERSIAN CANALS, Canals, nutritive, of Bones.

HAW, Hungry.

HAW, BLACK, *Viburnum prunifolium*.

HAWKNUT, *Bunium bulbocastanum*.

HAWKWEED, VEINY, *Hieracium venosum*.

HAWSE, Throat.

HAWTHORN, WHITE, *Mespilus oxyacantha*.

HAY ASTHMA, Fever, Hay.

HAY, CAMEL'S, *Juncus odoratus*.

HAYRIFF, *Galium aparine*.

HAYSCALE, Harelip.

HAZEL, BEAKED, *Corylus rostrata*.

HAZEL CROTCHES, *Lichen pulmonarius*—h.

Nut, snapping, *Hamamelis Virginiana*—h. Nut tree, *Corylus avellana*.

HAZEL, WITCH, *Hamamelis Virginiana*, *Sorbus aucuparia*.

HAZELRAW, *Lichen pulmonarius*.

HEAD, Sax. *heapoc*, *hearp*, *heaved*; the past participle of *hearan*, 'to heave up.' *Caput*, *Ceph'alê*, *Cy'bê*, *Poll*, (Prov.) *Hede*, *Pow*, *Scop*, (F.) *Tête*. The head forms the upper extremity of the body, and tops the skeleton. It consists of the cranium and face. The first, which comprises all the superior and posterior part, has the encephalon in its cavity: the latter forms only the anterior part, and serves as the receptacle for the greater part of the organs of the senses.

HEAD, WATER IN THE, Hydrocephalus.

HEAD-ACHE, *Cephalæa*, *Cephalalgia*—h. Ache, intermittent, *Cephalalgia periodica*—h. Ache, sick, *Cephalæa spasmodica*.

HEADSWOMAN, Midwife.

HEADY, same etymon as Head. (F.) *Capiteux*. That which inebriates readily. An epithet, applied to wines which possess this quality.

HEAL, *Sanitas*.

HEAL-ALL, *Collinsonia Canadensis*, *Prunella vulgaris*, *Scrophularia nodosa*.

HEALING, Curative—h. Art, Medicine.

HEALTH, *Sanitas*.

HEALTHY, Salutory.

HEARING, HARDNESS OF, Deafness—h. Perverse, *Paracusis perversa*.

HEART, Sax. *heort*, Germ. Herz, Cor, Cœur, Cœr, *Card'ia*, *καρ*, *κρ*, *καρδία*, (F.) *Cœur*. An azygous muscle, of an irregularly pyramidal shape; situate obliquely and a little to the left side, in the chest;—resting on the diaphragm by one of its surfaces:—suspended by its base from the great vessels; free and movable in the rest of its extent, and surrounded by the pericardium. The right side of the body of the heart is thin and sharp, and is called *Margo acut'us*: the left is thick and round, and termed *Margo obtu'us*. It is hollow within, and contains four cavities; two of which, with thinner and less fleshy walls, receive the blood from the lungs and the rest of the body, and pour it into two others, with thick and very fleshy parietes, which send it to the lungs and to every part of the body. Of these cavities, the former are called *auricles*, the latter *ventricles*. The right auricle and right ventricle form the *Pulmonic* or *right or anterior heart*, *Cor veno'sum seu pulmona'le*, (F.) *Cœur du poumon*, *C. du sang noir*, *C. droit*, *C. antérieur*: and the left auricle and ventricle, the *systemic, corporeal, left, or aortic heart*, *Cor arterio'sum seu aor'ticum*, (F.) *Cœur du corps*, *C. gauche*, *C. aortique* ou *C. rouge*. In the adult, these are totally distinct from each other, being separated by a partition;—the *septum cordis*. Into the right auricle, the *venæ cavæ*,—superior and inferior,—and the coronary vein, open. The pulmonary artery arises from the right ventricle; the four pulmo-

nary veins open into the left auricle, and the aorta arises from the left ventricle.

The mean weight of the heart, in the adult, from the twenty-fifth to the sixtieth year, is, according to Bouillaud, from eight to nine ounces. The dimensions, according to Lobstein and Bouillaud, are as follows:—Length, from base to apex, five inches six lines; breadth, at the base, three inches; thickness of the walls of the left ventricle, seven lines; at a finger's breadth above the apex, four lines; thickness of the walls of the right ventricle, two and a quarter lines; at the apex, half a line; thickness of right auricle, one line; of the left auricle, half a line. The heart is covered, externally, by a very thin, membranous reflection from the pericardium. The muscular structure of which it is constituted is much thicker in the parietes of the ventricle than in those of the auricles. Its cavities are lined by a very delicate membrane, the *endocardium*, which is continuous with the inner membrane of the arteries, as regards the left cavities, and with that of the veins, as regards the right. Its arteries—the *coronary*—arise from the commencement of the aorta. Its nerves proceed, chiefly, from the pneumogastric and the cervical ganglions of the great sympathetic. The heart is the great agent in the circulation. By its contraction, the blood is sent over every part of the body. Its action does not seem to be *directly* owing to nervous influence received from the brain or spinal marrow, or from both. The circulation may, indeed, be kept up, for some time, if both brain and spinal marrow be destroyed.

When the ear is applied to the chest, a dull, lengthened sound is heard, which is synchronous with the arterial pulse. This is instantly succeeded by a sharp, quick sound, like that of the valve of a bellows or the lapping of a dog, and this is followed by a period of repose. The first sound appears to be mainly produced by the contraction of the ventricles; the second, by the reflux of the blood against the semilunar valves. These are what are called the *Sounds of the Heart*, (F.) *Bruits du Cœur*. Dr. C. J. B. Williams thinks that the word *lub-dup* conveys a notion of the two sounds. The *Beating or Impulse of the heart*, *Heart-stroke*, *Pulsus* seu *Ictus cordis*, (F.) *Impulsion du Cœur*, against the parietes of the chest, is mainly caused, perhaps, by the systole of the heart, which tends to project it forwards. It is doubted, however, by some, whether the impulsion be produced by the dilatation or the contraction of the ventricles.

The following table exhibits the different actions of the heart, and their coincidence with its sounds and impulse. It presumes, that the period from the commencement of one pulsation to that of another is divided into eight parts; and if the case of a person, whose pulse beats, sixty times in a minute, be taken, each of these parts will represent the eighth of a second.

EIGHTHS OF A SECOND.

Last part of the pause....1...	Auricles contracting; ventricles distended.
First sound and impulse...4...	Ventricles contracting; auricles dilating.
Second sound.....2...	Ventricles dilating; auricles dilating.
Pause.....1...	Ventricles dilating; auricles distended.

The heart is subject to different organic diseases: the chief of which are *aneurism*, *contraction of the apertures*, and *rupture of its parietes*.

HEART, AT'ROPHY, OF THE, *Atroph'ia* seu *Ari-*

du'ra seu *Phthisis Cordis*, *Cardi'atroph'ia*, *Acardi'atroph'ia*. A condition of the organ in which there is diminution in the thickness of the parietes of the whole organ.

HEART, BEATING OF THE, see Heart—h. Dilatation of the, see Aneurism—h. Displacement of the, *Ectopia Cordis*—h. Fatty, *Steatosis cordis*.

HEART, HY'PERTROPHY OF THE, *Hypertroph'ia Cordis*, *Hypercor'dia*, *Hyperar'co'sis Cordis*, *Hypereardi'atroph'ia*, *Cardiaux'e*, *Aneurys'ma cordis acti'vum* (Corvisart), *Cor bovi'num* seu *tauri'num*, *Enor'mitas cordis*, (F.) *Hypertrophie du Cœur*. Supernutrition of the muscular parietes of the heart, which are thicker than usual; the cavities being generally diminished. The *physical signs* which indicate it are the following. In cases of long standing, the præcordial region is generally prominent; the pulsations of the heart are visible over a greater extent than natural; and a marked vibration is communicated to the hand when placed on the cardiac region. The dull sound on percussion is more extensive than natural; and on auscultation there is a permanent increase of the force and extent of the heart's action; there is no increase, however, of frequency, and the rhythm is regular. The pulse is generally strong, full, and hard.

HEART, HYPERTROPHY WITH DILATATION OF THE, *Active an'eurism*, *Eccen'tric hy'p'trophy*. In this affection, the pulsations can be seen and felt over a larger space, and the apex is more to the left and lower down than natural. The impulse is less steady, but at times more violent than that which accompanies simple hypertrophy. Percussion gives more distinct evidence of the enlargement, the sound being more extensively dull. On auscultation, the impulse is often violent, but irregular: in extreme cases, it produces the sensation of a large mass of flesh rolling or revolving beneath the ear. The pulse is strong, full, and vibratory. The shock of the heart's action is often transmitted to the whole person, and to the bed on which the patient is lying.

HEART, CONCENTRIC HYPERTROPHY OF THE, (F.) *Hypertrophie concentrique*, *Hypertrophie centripète du Cœur*, *Retrait ventriculaire*, *Cardiac'tie*, is when the parietes augment at the expense of the cavities.

HEART, IMPULSE OF THE, see Heart—h. Neuralgia of the, *Angina Pectoris*—h. Rupture of the, *Cardiorrh'xis*—h. Sounds of the, see Heart—h. Stroke, see Heart.

HEART'S EASE, *Polygonum persicaria*, *Viola tricolor*.

HEARTS, LYMPH, Lymphatic hearts—h. Lymphatic, Lymphatic hearts.

HEARTAXES, Ardor ventriculi.

HEARTBURN, Ardor ventriculi, Cardialgia.

HEARTSCALD, Ardor ventriculi.

HEART-SCIRTS, Diaphragm.

HEARTWORT, *Laserpitium album*.

HEAT, past participle of Sax. *hætan*, 'to make warm.' *Calor*, *Therma*, *Σεμν*. (F.) *Chaleur*. The material cause, which produces the sensation—or the particular sensation itself—produced by a body of an elevated temperature, on our organs, especially on the organs of touch. Modern chymists have given the name *Caloric* to the principle, whatever may be its nature, which is the cause of heat. When we touch a body of a temperature superior to our own, a portion of caloric passes from the body to the hand, and produces the sensation of *heat*. If, on the contrary, we touch a body of a temperature inferior to our own, we communicate a portion of our caloric to it, and experience the sensation of *cold*. Our own sensations are but imperfect indexes of temperature. Two men meeting at the middle of a mountain,—the one ascending, the other de-

ascending,—will experience different sensations. The one ascending, passes from a warmer to a colder atmosphere;—the one descending from a colder to a warmer.

The chief instrument for measuring heat, used

in medicine, is the thermometer. Of this there are three kinds:—that of Fahrenheit, that of Réaumur, and that of Celsius or the *Centigrade*. The following Table exhibits the correspondence of these different thermometric scales.

Fah.	Reau.	Centig.	Fah.	Reau.	Centig.	Fah.	Reau.	Centig.	Fah.	Reau.	Centig.
0	0	0	143	51.55	64.44	85	23.55	20.44	22	— 4.44	— 5.55
212 _u	80.00	100.00	147	51.11	63.88	84	23.11	20.88	21	— 4.88	— 6.11
210 _b	79.55	99.44	146	50.66	63.33	83	22.66	21.33	20 _u	— 5.33	— 6.66
209	79.11	98.88	145	50.22	62.77	82	22.22	21.77	19	— 5.77	— 7.22
208	78.66	98.33	144	49.77	62.22	81 _o	21.77	22.22	18	— 6.22	— 7.77
207	78.22	97.77	143	49.33	61.66	80	21.33	22.66	17	— 6.66	— 8.33
206	77.77	97.22	142 _g	48.88	61.11	79	20.88	23.11	16	— 7.11	— 8.88
205	77.33	96.66	141	48.44	60.55	78	20.44	23.55	15	— 7.55	— 9.44
204	76.88	96.11	140	48.00	60.00	77	20.00	24.00	14	— 8.00	— 10.00
203	76.44	95.55	139	47.55	59.44	76 _p	19.55	24.44	13	— 8.44	— 10.55
202	76.00	95.00	138	47.11	58.88	75	19.11	24.88	12	— 8.88	— 11.11
201	75.55	94.44	137	46.66	58.33	74	18.66	25.33	11	— 9.33	— 11.66
200	75.11	93.88	136	46.22	57.77	73	18.22	25.77	10	— 9.77	— 12.22
199	74.66	93.33	135	45.77	57.22	72	17.77	26.22	9	— 10.22	— 12.77
198	74.22	92.77	134	45.33	56.66	71	17.33	26.66	8	— 10.66	— 13.33
197	73.77	92.22	133 _h	44.88	56.11	70	16.88	27.11	7 _v	— 11.11	— 13.88
196 _c	73.33	91.66	132	44.44	55.55	69	16.44	27.55	6	— 11.55	— 14.44
195	72.88	91.11	131	44.00	55.00	68	16.00	28.00	5	— 12.00	— 15.00
194	72.44	90.55	130	43.55	54.44	67	15.55	28.44	4	— 12.44	— 15.55
193	72.00	90.00	129	43.11	53.88	66	15.11	28.88	3	— 12.88	— 16.11
192	71.55	89.44	128	42.66	53.33	65	14.66	29.33	2	— 13.33	— 16.66
191	71.11	88.88	127	42.22	52.77	64	14.22	29.77	1	— 13.77	— 17.22
190	70.66	88.33	126	41.77	52.22	63	13.77	30.22	0	— 14.22	— 17.77
189	70.22	87.77	125	41.33	51.66	62	13.33	30.66	— 1	— 14.66	— 18.33
188 _q	69.77	87.22	124	40.88	51.11	61	12.88	31.11	— 2 _w	— 15.11	— 18.88
187	69.33	86.66	123	40.44	50.55	60	12.44	31.55	— 3	— 15.55	— 19.44
186	68.88	86.11	122	40.00	50.00	59	12.00	32.00	— 4	— 16.00	— 20.00
185	68.44	85.55	121	39.55	49.44	58	11.55	32.44	— 5	— 16.44	— 20.55
184	68.00	85.00	120	39.11	48.88	57	11.11	32.88	— 6	— 16.88	— 21.11
183	67.55	84.44	119	38.66	48.33	56	10.66	33.33	— 7 _x	— 17.33	— 21.66
182	67.11	83.88	118	38.22	47.77	55	10.22	33.77	— 8	— 17.77	— 22.22
181	66.66	83.33	117	37.77	47.22	54	9.77	34.22	— 9	— 18.22	— 22.77
180	66.22	82.77	116	37.33	46.66	53	9.33	34.66	— 10	— 18.66	— 23.33
179	65.77	82.22	115	36.88	46.11	52	8.88	35.11	— 11 _y	— 19.11	— 23.88
178 _d	65.33	81.66	114	36.44	45.55	51	8.44	35.55	— 12	— 19.55	— 24.44
177	64.88	81.11	113	36.00	45.00	50 _g	8.00	36.00	— 13	— 20.00	— 25.00
176	64.44	80.55	112	35.55	44.44	49	7.55	36.44	— 14	— 20.44	— 25.55
175	64.00	80.00	111	35.11	43.88	48	7.11	36.88	— 15	— 20.88	— 26.11
174 _e	63.55	79.44	110 _i	34.66	43.33	47	6.66	37.33	— 16	— 21.33	— 26.66
173	63.11	78.88	109	34.22	42.77	46	6.22	37.77	— 17	— 21.77	— 27.22
172	62.66	78.33	108	33.77	42.22	45	5.77	38.22	— 18	— 22.22	— 27.77
171	62.22	77.77	107	33.33	41.66	44	5.33	38.66	— 19	— 22.66	— 28.33
170	61.77	77.22	106 _j	32.88	41.11	43	4.88	39.11	— 20	— 23.11	— 28.88
169	61.33	76.66	105	32.44	40.55	42	4.44	39.55	— 21	— 23.55	— 29.44
168	60.88	76.11	104 _k	32.00	40.00	41	4.00	40.00	— 22	— 24.00	— 30.00
167 _f	60.44	75.55	103	31.55	39.44	40	3.55	40.44	— 23	— 24.44	— 30.55
166	60.00	75.00	102 _l	31.11	38.88	39	3.11	40.88	— 24	— 24.88	— 31.11
165	59.55	74.44	101	30.66	38.33	38	2.66	41.33	— 25	— 25.33	— 31.66
164	59.11	73.88	100 _m	30.22	37.77	37	2.22	41.77	— 26	— 25.77	— 32.22
163	58.66	73.33	99	29.77	37.22	36	1.77	42.22	— 27	— 26.22	— 32.77
162	58.22	72.77	98 _n	29.33	36.66	35	1.33	42.66	— 28	— 26.66	— 33.33
161	57.77	72.22	97	28.88	36.11	34	0.88	43.11	— 29	— 27.11	— 33.88
160	57.33	71.66	96	28.44	35.55	33	0.44	43.55	— 30	— 27.55	— 34.44
159	56.88	71.11	95	28.00	35.00	32 _r	0.00	44.00	— 31	— 28.00	— 35.00
158	56.44	70.55	94	27.55	34.44	31	— 0.44	44.44	— 32	— 28.44	— 35.55
157	56.00	70.00	93	27.11	33.88	30 _s	— 0.88	44.88	— 33	— 28.88	— 36.11
156	55.55	69.44	92	26.66	33.33	29	— 1.33	45.33	— 34	— 29.33	— 36.66
155	55.11	68.88	91	26.22	32.77	28 _t	— 1.77	45.77	— 35	— 29.77	— 37.22
154	54.66	68.33	90	25.77	32.22	27	— 2.22	46.22	— 36	— 30.22	— 37.77
153	54.22	67.77	89	25.33	31.66	26	— 2.66	46.66	— 37	— 30.66	— 38.33
152	53.77	67.22	88	24.88	31.11	25	— 3.11	47.11	— 38	— 31.11	— 38.88
151	53.33	66.66	87	24.44	30.55	24	— 3.55	47.55	— 39 _z	— 31.55	— 39.44
150	52.88	66.11	86	24.00	30.00	23	— 4.00	48.00	— 40	— 32.00	— 40.00
149	52.44	65.55									
148	52.00	65.00									

a Water boils (*Barom.* 30 inches.)
b Heat of a stove, borne by Dr. Solander.

c Heat of a stove, borne for 10 minutes by Sir Joseph Banks and Dr. Solander.

d Water simmers.

e Alcohol boils.

f Very pure ether distills.

g Bees' wax melts.

h Spermaceti melts.

i Temperature at which liquids are often drunk.

j Heat observed in Scarlatina.

k Temperature of the common hen.

l Temperature of arterial blood. (?)

m Temperature of venous blood.

Phosphorus melts.

n Ether boils. *o* Nitric ether boils.

p Muriatic ether boils.

q Medium temperature of the globe.

r Ice melts.

s Milk freezes.

t Vinegar freezes.

u Strong wine freezes.

v A mixture of one part of alcohol

and three of water freezes.

w Cold at the battle of Eylau, 1807

x A mixture of equal parts of alcohol and water freezes.


y A mixture of two parts of alcohol and one of water freezes.

z Melting point of quicksilver.

To reduce Centigrade degrees to those of Fahrenheit, multiply by 9, divide by 5, and add 32: thus, 40° Cent. $\times 9 \div 5 + 32 = 104^\circ$ Fahr. To reduce Fahrenheit's degrees to those of Centigrade, subtract 32, multiply by 5, and divide by 9: thus,

104° Fahr. $- 32 \times 5 \div 9 = 40^\circ$ Cent. To reduce Réaumur's degrees to those of Fahrenheit, multiply by 9, divide by 4, and add 32: thus, 32° Réaumur. $\times 9 \div 4 + 32 = 104^\circ$ Fahr. To reduce Fahrenheit's degrees to those of Réaumur, subtract

32, multiply by 4, and divide by 9: thus, 104° Fahr. — $32 \times 4 \div 9 = 32$ Réaumur. To reduce Réaumur's degrees to those of Centigrade, multiply by 5, and divide by 4: thus 32° Réaumur. $\times 5 \div 4 = 40^\circ$ Cent.; and, lastly, to reduce Centigrade degrees to those of Réaumur, multiply by 4, and divide by 5: thus, 40° Cent. $\times 4 \div 5 = 32^\circ$ Réaumur.



Thermometers.

The human body can bear a high degree of heat diffused in the atmosphere. There are cases on record, where air of 400° and upwards, of Fahrenheit's scale, has been breathed with impunity for a short time. It can likewise withstand very severe cold. In the expedition of Capt. Back to the Arctic regions, the thermometer was as low as — 70° of Fahr., and Sir George Simpson affirms, that it has fallen in Siberia to — 83°, or 115° below the freezing point. Dr. Kane, in the Arctic regions, in 1853–54, observed the range of eleven spirit thermometers, selected as standards, to vary from — 60° to — 75°. The mean annual temperature was 5°·2—the lowest ever registered.

Excessive heat disposes the body to gastric and intestinal diseases, and particularly to inflammation and enlargement of the liver; hence, the frequency of such affections within the torrid zone. Heat is often used therapeutically: the actual cautery, at a white heat, disorganizes the parts to which it is applied: a lesser degree occasions violent inflammation. Heat higher than that of the human body is excitant; of a lower degree, sedative. Excessive cold acts as a powerful sedative—inducing sleep, the tendency to which, after long exposure, becomes irresistible. See Cold. Many of the topical applications—as cataplasms—act wholly by virtue of their warmth and moisture.

HEAT, *Ardor venereus*, *Pruritus*, (F.) *Chaleurs*, *Chaleurs des Animaux*, *Rut*, is the periodical sexual desire experienced by animals,—(*Estruation*). It is supposed by some to be owing to the periodical maturation and discharge of ova,—(*Ovulation*).

HEAT, ACRID, see *Acrid*—h. Animal, see *Animal Heat*.

HEAT APOPLEXY, see *Coup-de-soleil*.

HEAT, INTERNAL, (F.) *Chaleur interne*, is a sensation of heat felt by the patient, but not sensible to the touch. *External heat*, (F.) *Chaleur extérieure*, that which can be felt by others. *Heat*, *Ardor*, is called moist, (F.) *haliteuse*, when

accompanied with moisture, like that felt after bathing; dry, (F.) *sèche*, when the skin has not its ordinary softness. It is called *acid* and *pungent*, (F.) *âcre et mordicante*, when it conveys a disagreeable tingling to the fingers. The French employ the terms *nervous heat* and *erratic heat*, *Chaleur nerveuse* and *Chaleur erratique*, for that which comes by flushes, alternating with chills, and which moves rapidly from one part to another. M. Double has used the term *septic heat*, (F.) *Chaleur septique*, for that which produces a pungent sensation on the hand, similar to that of the acid heat, but milder and more uniform, and which is accompanied with feebleness and frequency of pulse, &c.

HEAT, PRICKLY, *Lichen tropicus*.

HEATH, COMMON, *Erica vulgaris*.

HEATHER, *Erica vulgaris*.

HEATHER-CLU, *Malleolus*.

HEAUTOPHONICS, see *Autophonia*.

HEAVINESS, *Somnolency*.

HEAVING, *Vomiturition*.

HEAVY, *Gravis*, (F.) *Gravatif*, from Sax. *heapan*, 'to heave.' An epithet given to any pain which consists in a sensation of weight or heaviness, or is accompanied by such sensation.

HEBDOMADAL CYCLE, see *Cycle*.

HEBDOMADARIA, *Octana*.

HEBE, ἡβη, *Juventa*, *Juventus*, *Juventus*, *Hora*. In antiquity, the goddess of puberty. The word has been employed to designate, 1. The first hair that grows on the pubes, 2. The pubic region, and 3. Puberty.

HEBETES, *Adolescens*.

HEBETOR, *Adolescens*.

HEBETUDO ANIMI, *Imbecility*—h. *Dentium*, *Hamodia*—h. *Visus*, *Amblyopia*, *Asthenopia*, *Caligo*.

HEBRADENDRON CAMBOGIODES, see *Cambogia*.

HEBREWS, MEDICINE OF THE. Medicine seems to have been at a very low ebb with the ancient Hebrews. Of anatomy they knew nothing. Their physiology was imperfect and filled with superstitions; and their therapeutics unsatisfactory. Hygiene appears to have been most attended to. Of the other departments of medicine we cannot judge of their knowledge.

HECATOMA PALUSTRIS, *Ranunculus sceleratus*.

HECATOMPHYLLUM, *Rosa centifolia*.

HECH, *Anhelatio*.

HECKBERRY, *Prunus padus*.

HECTEUS, ἑκτεως. A Greek measure, containing about 72 chopines or pints.

HECTIC FEVER, *Febris hectica* seu *phthisica* seu *marasmodica* seu *tabida* seu *lenta* seu *amphimerina* *hectica* seu *amato'ria*, *Hecticopy'ra*, *Hecticopy'retos*, *Hectica*, *Amphimerina* *hectica*, *Syntecopy'ra*, *Syntectopy'ra*, *Marasmodopy'ra*, *Leucopy'ria*, *Epan'eus* *hectica*, *Chlorosis* *amato'ria*, *Ettick*, (Sc.) *Ethick*, *Ettick*, (F.) *Fièvre Hectique* on *Étique*, from ἥξις, 'habit of body,' because in this disease every part of the body is emaciated; or, perhaps, from ἐκτρεω, 'I consume,' 'I am exhausted.' The name of a slow, continued, or remittent fever, which generally accompanies the end of organic affections, and has been esteemed idiopathic, although it is probably always symptomatic. It is the fever of irritation and debility; and is characterized by progressive emaciation, frequent pulse, hot skin,—especially of the palms of the hands and soles of the feet,—and, towards the end, colliquative sweats and diarrhoea. Being symptomatic, it can only be removed by getting rid of the original affection. This is generally difficult, and almost hopeless in the disease which it most commonly accompanies,—consumption.

HECTICA, Hectic Fever—*h. Infantilis*, Fever, Infantile remittent.

HECTICOPYRA, Hectic fever.

HECTICOPYRETOS, Hectic fever.

HECTITIS, (*F.*) *Étisie*. Same etymon. The state of those who have hectic fever.

HECTOGRAMME, *Hectogramma*, from *ἑκατον*, 'a hundred,' and *γραμμα*. A measure of 100 grammes, i. e., 3 ounces, 1 drachm, and 44 grains, Troy.

HECTOLITRE. A measure containing 100 litres or 26.42 wine pints.

HECUSIUS, Voluntary.

HEDEO'MA, *Hedeo'ma pulegioides*, *Cuni'la pulegioides*, *Melis'sa pulegioides*, *Pennyroyal*, *Tickweed*, *Stinking Balm*, *Squawmint*. *Sec. Syst.* *Diandria Monogynia*; *Nat. Ord.* *Labiatae*. From *ἡδύς*, 'agreeable.' An indigenous plant, common in all parts of the United States, and which, where it is abundant, perfumes the air for a considerable distance. It is employed in the same cases as the mints and the English pennyroyal. In popular practice it is used as an emmenagogue.

The *O'leum Hedeo'mæ* (*Ph. U. S.*) or *Oil of Pennyroyal* is used as a stimulating carminative, dropped on sugar. Dose, 2 to 6 drops.

HERERA ARBOREA, *H. Helix*.

HERERA HELIX, *Hed'era arbo'rea*, *Bac'ehica*, *Cissos*, *κισσός*, *Cittos*, *κίττος*, *Ev'era*, *Corymb'etra*, *Corym'bos*, (*Sc.*) *Bindwood*, *Ivy*, (*F.*) *Lierre*, *L. Commun* ou *grimpant*. *Ord.* *Araliacæ*. The taste of *ivy leaves* is bitter, styptic, and nauseous. They are not used in medicine. According to Haller, they were recommended in Germany against the atrophy of children; and the common people of England sometimes apply them to running sores and to keep issues open. The berries were supposed, by the ancients, to have an emetic and purgative quality; and a watery extract was made from them, called by *Quercetanus* *Extractum purgans*. From the stalk of the tree a resinous juice exudes, in warm climates, called *Cummi Hed'erae*, *Ivy Gum*, (*F.*) *Gomme de lierre*, *Résine de lierre*. It is possessed of tonic and astringent properties, but is not used.

HERERA TERRESTRIS, *Glechoma hederacea*.

HERERULA, *Glechoma hederacea*.

HEDE, Head.

HEDEVERK, *Cephalalgia*.

HEDEWARK, *Cephalalgia*.

HEDISARUM ALHAGI, *Agul*.

HEDRA, *ἑδρα*, *ἑδρη*, 'a vestige,' 'a seat.' A seat. A fracture of the bones of the cranium, in which the trace of the fracturing instrument is still perceptible. It was, also, used by the ancients, for the anus, the breech, excrement, a privy, a night-chair, and for the bottom of an abscess.

HEDROCELE, *Proctocèle*.

HEDYCH'ROUM, from *ἡδύς*, 'agreeable,' and *χρῶς*, 'colour.' A remedy of a pleasant colour. Applied to certain trochs, the chief constituent of which was *theriac*.

HEDYPHO'NIA, from *ἡδύς*, 'sweet,' and *φωνη*, 'voice.' Sweetness of voice.

HEDYPNEUSTUS, *Hedyp'nous*, from *ἡδύς*, 'sweet,' and *πνεύω*, 'I breathe.' Breathing sweetly or softly. Smelling sweetly.

HEDYPNOSI TARAXACUM, *Leontodon taraxacum*.

HEDYSARUM ALHAGI, *Agul*.

HEDYSMA, *Condiment*.

HEEL, see *Calcaneum*—*h. Clubfoot*, see *Kyl'osis*.

HEGH, *Anhelatio*.

HEIL, *Sanitas*.

HELCENTERITIS, *Dothinenteritis*.

HELCOC'ACĒ, *Cachelco'ma*, from *ἑλκος*,

'ulcer,' and *κακή*, 'badness.' A malignant ulcer.

HELCODES, *Ulcerated*, *Ulcerous*.

HELCOLO'OGY, *Helcology'ia*, from *ἑλκος*, 'ulcer,' and *λογος*, 'discourse.' The doctrine of, or a treatise on, ulcers.

HELCOMA, *Ulcer*.

HELCOMENIA, *Menelcosis*.

HELCPH'THAL'MIA, *Ophthal'mia ulcero'sa*. Ophthalmia with ulceration.

HELCOPLASTY, *Hel'koplasty*, *El'koplasty*, *El'coplasty*; from *ἑλκος*, 'an ulcer,' and *πλασσω*, 'I form.' An operation which consists in grafting on an ulcer a piece of skin from the opposite limb, or from the limb of another person, when the destruction of skin is too great to permit the healing process.

HEL'COS, from *ἑλκος*, 'an ulcer.' Hence:

HEL'COSIS, *Eclcosis*, *Ulceration*—*h. Cerebri*, *Encephalopoyosis*—*h. Laryngis*, *Phthisis laryngea*—*h. Pulmonalis*, *Phthisis pulmonalis*—*h. Renalis*, *Nephrelosis*—*h. Uteri*, see *Metrophthisis*—*h. Vesicæ*, *Cystophthisis*.

HEL'COSTAPHYLO'MA, from *ἑλκος*, 'ulcer,' and *σταφυλωμα*, 'staphylo'ma.' Staphylo'ma ending in ulceration.

HELCOXERO'SIS, from *ἑλκος*, 'ulcer,' and *ξηρωσις*, 'dryness.' The drying of an ulcer.

HELC'TICA, *Epispastics*.

HELCUS, *Ulcer*.

HEL'CYD'RION, *Helcyd'rion*, *ἑλκιδιον*, *Ulcusculum*, a small ulcer, a superficial ulceration of the cornea.—*Galen*, *Paulus*, *Foësius*.

HEL'CYSTER, *ἑλκυστηρ*, from *ἑλκω*, 'I draw.' An iron hook or crochet for extracting the fœtus. See *Crochet*.

HELENIUM, *Inula helenium*.

HELENIUM AUTUMNA'LE, *Falsæ Sunflower*, *Sneeze'wort*, *Sneeze'weed*, *Swamp Sunflower*, *Yellow star*, *Ox-eye*. *Ord.* *Compositæ*. An indigenous herb, with large golden-yellow compound flowers, which appear in August. All its parts are bitter and somewhat acrid, and when snuffed up the nostrils in powder are powerful sternutatories.

HELIANTHEMUM CANADEN'SĒ, *Cistus Canaden'sis*, *Frostwort*, *Frostweed*, *Frost plant*, *Rock rose*. *Ord.* *Cistacææ*. An herbaceous plant, *Helianthemum* (*Ph. U. S.*), having large yellow flowers, which grows in all parts of the United States, and flowers, in the Middle States, in June. It has an astringent, slightly aromatic, and bitterish taste. It has been prescribed in scrofula, but probably is nothing more than an aromatic tonic.

HELIANTHEMUM CORYNDO'SUM, *Rock rose*, an indigenous plant, is used in the same cases.

HELIASIS, *Astrabolismus*, *Insolation*.

HELICHRYSUM NUDIFOLIUM, *Caff'tea*. A South African plant, *Nat. Ord.* *Compositæ*, which is demulcent, and, in the form of infusion, is recommended in catarrh, phthisis, and other pulmonary affections.

HELICHRYSUM SERPYLLIFOLIUM, *Hottentot's tea*, and **HELICHRYSUM AURICULATUM** have similar virtues.

HELICIA, *Age*.

HEL'ICINE, *Helic'inus*, *Helicoi'des*, *Helico'des*, from *helix*, *helicia*, 'the tendrill of the vine.' Resembling in form the tendrill of the vine.

HELICINE ARTERIES of the penis, as described by J. Müller, are short vessels given off from the larger branches, as well as from the finest twigs of the artery of the organ; most of those come off at a right angle, and project into the cavity of the spongy substance, either terminating abruptly or swelling out into a club-like process without again subdividing. Almost all these vessels are bent like a horn, so that the end de-

scribes half a circle or somewhat more. They have a great resemblance to the tendrils of the vine, whence their name. A minute examination of them, either with the lens or the microscope, shows that, although they at all times project into the venous cavities of the corpora cavernosa, they are not entirely naked, but are covered with a delicate membrane, which, under the microscope, appears granular.

HELICIS MAJOR. A muscle of the ear, which originates from the anterior, acute part of the helix, upon which it ascends and is inserted into the helix. It pulls the part into which it is inserted a little downwards and forwards.

HELICIS MINOR. This muscle originates from the under and fore part of the helix, and is inserted into the helix, near the fissure in the cartilage, opposite the concha. Its use is to contract the fissure.

HELICOIDES, Helicine.

HELICOTREMA, from *ἑλῖς*, 'helix, cochlea,' and *τρημα*, 'a foramen.' The hole by which the two scale of the cochlea communicate at the apex.

HELIENCEPHALITIS, see *Coup de Soleil*.

HELIGMUS, Convolution.

HELIKIA, Age.

HELIONOSIS, Insolation.

HELIOSCIADIUM NODIFLORUM, *Sium nodiflorum*.

HELIOSIS, Astrabolism, Insolation.

HELIOTROPE, *Heliotropium Europæum*.

HELIOTROPION, *Cichorium intybus*.

HELIOTROPIMUM EUROPEUM, *H. erectum seu canescens seu supinum, Vernucaria.* The *Heliotrope*, (F.) *Tournesol, Herbe aux verres.* Ord. Boraginacæ. This plant is considered to possess apert properties; and to be capable of destroying cutaneous excrescences; hence one of its names.

HELIOTROPION INDICUM, an East India plant, naturalized in some of the Western and Southern States, probably has similar virtues.

HELIX, *ἑλῖς, Capreaulus*, from *ελεῖν*, 'to envelop,' 'to roll.' The fold is thus called, which forms the outer circumference or ring of the external ear.

HELIX, Limax.

HELIX POMATIA, (F.) *Limax, Colimaçon, Escargot.* A large kind of snail, transported from the south of Europe to England by Sir Kenelm Digby, for his lady when in a decline. It was considered highly restorative.

HELLA, Incubus.

HELLEBORASTER, Helleborus foetidus—h. Foetidus, Helleborus foetidus.

HELLEBORE, Dracontium foetidum—h. American, Veratrum viride—h. Black, Helleborus niger—h. Blanc, Veratrum album—h. Green, Helleborus viridis—h. Noir, Helleborus niger—h. Stinking, Helleborus foetidus—h. Swamp, Veratrum viride—h. White, Veratrum album.

HELLEBORIS'MUS, Elleboris'mus. The method of treating disease, amongst the ancients, by hellebore. This comprised not only the choice, preparation, and administration of the medicine, but, likewise, the knowledge and employment of preliminary precautions and remedies proper for aiding its action, and preventing the pernicious effects which it might occasion.

HELLEBORUS, H. niger—h. Albus, Veratrum album—h. Grandiflorus, H. niger—h. Trifolius, Coptis.

HELLEBORUS FOETIDUS, Helleboraster, Elleboraster, Stinking Hellebore or Bear's foot, Setterwart, (F.) Hellebore ou Ellebore fétide, Pied de Griffon. Ord. Ranunculacæ. The leaves of this plant are said to be anthelmintic. The smell of the fresh plant is extremely fetid, and

the taste bitter and acrid. It usually acts as a cathartic.

HELLEBORUS NIGER, H. grandiflorus, Elleb'orus niger, Melampo'dium, Melanorrh'izum, Ec'tomon, Black Hellebore, Melampode, Christmas Rose, Clove-tongue, (F.) Hellebore noir. The root of this European plant—*Helleborus*, (Ph. U. S.)—has a disagreeable odour, and bitter, acrid taste. It is possessed of cathartic properties, and has been recommended as an emmenagogue. It has been given in mania, melancholia, dropsy, suppressed menses, &c. Dose, gr. x to ʒj, as a cathartic.

HELLEBORUS VIRIDIS, Green Hellebore. A European plant, naturalized on Long Island, has similar properties to the last.

HELLECEBRA, Sedum.

HELLENIA GRANDIFLORA, Costus.

HELMET-FLOWER, YELLOW, Aconitum Anthora.

HELMET POD, Jeffersonia Bartoni.

HELMINS, genitive, *Helminthos*, plur. *Helmin'thes* seu *Elmin'thes*, from *ἑλμιν*, 'a worm.' A worm; an entozoon:—hence:

HELMINTHAGOGUE, Anthelmintic.

HELMINTHI, Worms.

HELMINTHIA, Helminthiasis—h. Alvi, Worms.

HELMINTHIA ERRATICA, Worms, introduced by accident and without finding a proper habitation in the stomach or intestines; producing spasmodic colic, with severe tormina, and occasionally vomiting or ejections of blood; the 12th genus of the order *Enterica*, class *Cœliaca*, of Good.

HELMINTHIA PODICIS, Worms.

HELMINTHIASIS, Helminthogen'esis. A generic name for the condition which gives occasion to the presence of intestinal worms. It is, also, called *Helmin'thia, Vermina'tio, Morbus vermino'sus, Status vermino'sus, Sabur'ra vermino'sa, Scolec'asis, Scolece'asis, Entozoögen'esis, Parasit'ismus intestinal'is; Worm disease, Invermin'ation.* See Worms.

HELMINTHIC, Anthelmintic.

HELMINTHOCHORTUM, Corallina Corsicana.

HELMINTHOCOLICA, Colica verminosa.

HELMINTHOCORTON, Corallina Corsicana.

HELMINTHOGENESIS, Helminthiasis.

HELMINTHOL'OGY, Helmintholog'ia, from *helmins, helminthos*, and *λογος*, 'a description.' A treatise on worms.

HELMINTHONCUS, Malis—h. Medinensis, Dracunculus.

HELMINTHOP'YRA, Helminthopyr'etos, Febris vermino'sa, from *helmins, helminthos*, and *πυρ*, 'a fever.' Fever occasioned by worms. See Fever, verminous.

HELMINTHOPYRETOS, Helminthopyra.

HELMINTHUS GORDII, Seta equina.

HELODES, Elodes.

HELONIAS DIOICA, Chamælerium luteum—h. Erythrosperma, Amianthum muscætoxicum—h. Lutea, Chamælerium luteum—h. Officinalis, see Veratrina.

HELOPYRA, Elodes (febris).

HELOPYRETUS, Elodes (febris).

HELOS, Clavus, Marsh.

HELO'SIS, Helo'tis: from *ἑλω*, 'I turn.' Eversion of the eyelids, and convulsions of the muscles of the eyes. *Plica Polonica. Stabismus.*

HELOTIS, Helosis, Plica.

HELUS, Clavus, Helos.

HELVELLA ESCULENTA, Morchella esculenta.

HELXINE, Parietaria.

HEMACHROIN, Hæmatin.

HÉMADYNAMÈTRE, *Hæmadynamometer*.

HÉMADYNAMIQUE, *Hæmadynamics*.

HÉMAPÉRITONIRRHAGIE, *Hæmaperitonirrhag'ia*; from *αἷμα*, 'blood,' *peritonæum*, and *πρὺναι*, 'I break forth.' Hemorrhage into the peritoneum.

HÉMAPROCTIE, *Hæmorrhoids*.

HÉMASTATIQUES, *Hæmostatica*.

HÉMATIDROSE, *Sudor eruentus*.

HÉMATIN, *Hæmatin*.

HÉMATO-ENCÉPHALIE, *Apoplexy*.

HÉMATOID, *Hæmatodes*.

HÉMATOÏDINE, *Hæmatoidin*.

HÉMATOLOGIE, *Hæmatology*.

HÉMATOLOGY, *Hæmatology*.

HÉMATOMYÉLIE, *Hæmorrhagie de la Moëlle Épinière*.

HÉMATONCIE, *Hæmatoncus*, see *Hæmatodes fungus*—*h. Fungoides*, *Hæmatodes fungus*.

HÉMATOPISIE, *Hæmatops*.

HÉMATORRHACHIS, *Apoplexia myelitica*.

HÉMATOSIN, *Hæmatin*.

HÉMATOSTATICS, *Hæmostatica*.

HÉMATOSTATIQUES, *Hæmostatica*.

HÉMATOZOAIRES, *Hæmatozoa*.

HÉMENDOCARDE, *Hæmendocardium*.

HÉMERA, *ἡμέρα*, 'a day.' *Dies*. Hence:

HEMERALOPIA, see *Nyctalopia*.

HEMERAL'PIA, from *ἡμέρα*, 'the day,' and *εἶσθαι*, 'I see.' *Hæmeralops*, *Dysop'ia* seu *Caligo tenebrarum*, *Parop'sis* *Noct'ia*, *Visus diurnus*, *Nyctalop'ia* (of some), *Nyctophlo'sis*, *Anblyop'ia crepuscularis*, *Cæcitas crepuscularis* seu *nocturna*, *A'cies diurna*, *Day-sight*, *Day-vision*, *Hen blindness*, *Night blindness*, *Nocturnal Anauro'sis*, (F.) *Vue diurne*, *Avenglement de Nuit*. A disease, in which the eyes enjoy the faculty of seeing, whilst the sun is above the horizon, but are incapable of seeing by the aid of artificial light. Its causes are not evident. The eye, when carefully examined, presents no alteration, either in its membranes or humours.

HEMERALOPS, *Do'rea*. One labouring under hemeralopia. Also, *Hemeralopia*.

HEMERATYPHLOSIS, *Nyctalopia*.

HEMERODROMA, *Ephemera*.

HEMEROPATH'IA; from *ἡμέρα*, 'a day,' and *πάθος*, 'an affection.' A disease, which continues only a day; or, which is only observed during the day.

HEMEROTYPHLOSIS, *Nyctalopia*.

HEMI, *ἡμι*, 'half,' 'semi.' Hence:

HEMIACEPH'ALUS, (F.) *Hemiacephale*, from *hemi*, *α*, privative, and *κεφαλή*, 'head.' A monster whose head is represented by a shapeless tumour, with some appendages or cutaneous folds anteriorly, the thoracic extremities existing.

HEMIAMAUROSIS, *Hemiopia*.

HEMIANDRUS, *Eunuch*.

HEMIANOR, *Eunuch*.

HEMIANTHROPIA, *Mania*.

HEMIANTHROPUS, *Eunuch*, *Maniac*.

HEMIAZYGA (*Vena*), see *Azygos vein*.

HEMICEPHAL'EA, *Hæmicrania*.

HEMICEPHALÆUM, *Sinciput*.

HEMICEPHALUM, *Sinciput*.

HEMICEPH'ALUS, *Hemienceph'alus*, *Semiceph'alus*, from *hemi*, 'half,' and *κεφαλή*, 'head.' One who has half a head.

HEMICRANIA, *Hemicephal'ea*, *Migra'na*, *Crania*, *Hemip'gia*, *Hemip'gia*, *Heterocrania*, *Monop'gia*, *Monop'gia*, *Cephal'ea* *Hemicrania*, *Hemipath'ia*, *Neural'gia cerebri*, *Megrim*, from *hemi*, 'half,' and *κρανιον*, 'cranium.' (F.) *Migraine*. Pain, confined to one half the head.

It is almost always of an intermittent character;—at times, continuing only as long as the sun is above the horizon; and hence sometimes called *Sun-pain*,—and is cured by cinchona, arsenic, and the remedies adapted for intermittents.

HEMICRANIA IDIOPATHICA, *Neuralgia facialis*.

HEMIDESMUS IN'DICUS, *Perip'laea Indica*, *Asclepias Pseudosarsa*, *Sarsaparilla Indica*, *Nannari*, *Indian Sarsaparilla*. Ord. *Asclepiadaceæ*. A Hindoostance plant, the root of which, *Hemidesm'us* (Ph. D.), has a peculiar aromatic odour, and a bitterish taste. It is used in India as a substitute for sarsaparilla.

HEMIDIALYSIS, *Hemiplegia*.

HEMIDIAPHORE'SIS, *Transpiratio Unilateralis*; from *hemi*, 'half,' and *διαφωρεω*, 'perspiration.' Perspiration of one half the body.

HEMIECTON, *Hemiecton*, *Semisextum*. A vessel capable of containing 36 chopines or pints, and in which fumigations were made in diseases of the uterus and vagina.—*Hippocrates*.

HEMIENCEPH'ALUS, (F.) *Hemiencephale*, from *hemi*, 'one half,' and *κεφαλος*, 'the cephalon.' A monster, which has no trace of the organs of the senses, and yet has a brain nearly natural. See *Hemicephalus*.

HEMIM'ILES, (F.) *Hémimèle*, from *hemi*, 'one half,' and *μῦλος*, 'a limb.' A genus of monsters, in which the upper or lower extremities are very defective—mere stumps,—and the fingers and toes are entirely wanting or very imperfect.—*I. G. St. Hilaire*.

HEMIMER'ION, from *hemi*, and *μοῖρα*, 'a part.' 'One half.'—*Foësius*. Also, half a drachm.—*Eratian*.

HEM'INA. A Greek measure, answering to the *Cotyle*, *κοτύλη*, i. e. one half the sextarius, or about half a pint, English.

HEMIOBOL'ION, *Hemiobolon*. Half the obolus. A weight of about five grains.—*Goræus*.

HEMIO'LION. A weight of 12 drachms, or oz. 1½.—*Galen*. See *Sesuncia*.

HEMIO'PIA, *Hemiop'sis*, *Hemiop'asia*, *Suf'fusio dimidia*, *Visus dimidia'tus*, *Marmor'gæ Hippocratis*, *Hemiamauro'sis*, *Anauro'sis dimidia'ta*, (F.) *Hémiopie*, from *hemi*, 'one half,' and *οπτομαί*, 'I see.' Depraved vision, in which the person sees only one half of an object.

HEMIOPIASIS, *Hemiopia*.

HEMIOPSIS, *Hemiopia*.

HEMIPAG'ES, *Dicor'yphus dihypogastrius*, *Ocotopus synapheoceph'alus*, from *hemi*, 'half,' and *πηγνυμι*, 'I fasten.' A monstrosity, in which twins are united from the navel to the vertex.—*I. G. Saint Hilaire*.

HEMIPAGIA, *Hemicrania*.

HEMIPATHIA, *Hemicrania*.

HEMIEPIGIA, *Hemicrania*.

HEMIPHO'NIA, from *ἡμι*, 'half,' and *φωνή*, 'voice.' Great weakness of voice. The husky voice of those affected with cholera.

HEMIPLE'GIA, *Hemiplex'ia*, *Hemidial'ysis*, *Paral'ysis Hemiplegia*, *Semiple'gia*, *Semi-sideratio*, from *hemi*, 'one half,' and *πλῆσσω*, or *πληττω*, 'I strike.' Paralysis of one side of the body. See *Paralysis*. One so palsied is said to be *hemipleg'ic*, *semi-sideratus*.

HEMIPLEGIA FACIALIS, *Paralysis*, *Bell's*.

HEMIPLEXIA, *Hemiplegia*.

HEMIPROSOPLEGIA, *Paralysis*, *Bell's*.

HEMISPHERÆ CEREBRI, *Hemispheres of the brain*.

HEMISPHERE, *Hemisphæ'ra*, *Hemisphæ'rium*, from *hemi*, 'one half,' and *σφαῖρα*, 'a sphere.' One half of a sphere or of a body having a spheroidal shape.

HEMISPHERES OF THE BRAIN, *Hemisphæ'ræ cerebri*, *Hemispherical ganglia*, are the upper

spheroidal portions of the brain, separated from each other by the falx cerebri.

HEMISPHERICAL GANGLIA, Hemispheres of the brain.

HÉMITE, *Hæmitis*.

HÉMITEUTÉA (FEBRIS), *Febria semi-tertiana seu æsqualitèra*, (F.) *Demi-tierce*, *Fèvre demitierce*, from *hemi*, 'half,' and *tertius*, 'tertiary.' A semi-tertian fever, so called because it seems to possess both the characters of the tertian and quotidian intermittent. — Galen, *Spigelius*.

HEMIUNCIA, *Hemiuncion*.

HEMIUNCION, *Hemiuncia*, *Seminuncia*, *Seminuncia*. Half an ounce.

HEMLOCK, *Pinus Canadensis*, *Conium maculatum* — h. American, *Cicuta maculata* — h. Bastard, *Cherophyllum sylvestre* — h. Common, *Conium maculatum*.

HEMLOCK DROPPORT, *Eranthe*.

HEMLOCK GROUND, *Taxus Canadensis*.

HEMLOCK GUM, see *Pinus Canadensis* — h. Pitch, see *Pinus Canadensis* — h. Poison, *Conium* — h. Spruce, *Pinus Canadensis*.

HEMLOCK, WATER, AMERICAN, *Cicuta maculata*, *Cicuta aquatica* — h. Water, fine-leaved, *Phellandrium aquaticum* — h. Wild, *Cicuta maculata*.

HÉMO-ANGIAIRRHAGIE, *Hæmoptysis*.

HÉMOCARDE, *Hæmocardiorrhagia*.

HÉMOCARDIOPLASTIES, see *Polypus*.

HÉMODIA, *Hæmodia*.

HÉMO-ENCÉPHALORRHAGIE, *Apoplexy*.

HEMOHÉPATORRHAGIE, *Hepatorrhagia*.

HÉMONÈTRE, *Hæmodynamometer*.

HÉMONYÉLORRHAGIE, *Apoplexy*, spinal.

HÉNOPÉRICARDE, *Hæmopericardium*.

HÉMOPTYSIE, *Hæmoptysis* — h. *Foudroyante*, see *Hæmoptysis*.

HEMORRHAGE, *Hæmorrhagia* — h. Accidental, see *Hæmorrhagia* — h. Active, see *Hæmorrhagia* — h. from the Bladder, *Cystirrhagia* — h. Constitutional, see *Hæmorrhagia* — h. Critical, see *Hæmorrhagia* — h. by Exhalation, see *Hæmorrhagia* — h. External, see *Hæmorrhagia* — h. General, see *Hæmorrhagia* — h. from the Intestines, *Melæna* — h. Internal, see *Hæmorrhagia* — h. Interstitial, *Apoplexy* — h. Local, see *Hæmorrhagia* — h. of the Mucous Membranes, see *Hæmorrhagia* — h. Passive, see *Hæmorrhagia* — h. from the Pituitary Membrane, *Epistaxis* — h. Pleural, *Hæmatothorax* — h. Secondary, see *Hæmorrhagia* — h. from the Skin, *Sudor cruentus* — h. Spinal, *Apoplexy*, spinal — h. Spontaneous, see *Hæmorrhagia* — h. Supplementary, see *Hæmorrhagia* — h. Symptomatic, see *Hæmorrhagia* — h. of the Tissues, see *Hæmorrhagia* — h. Traumatic, see *Hæmorrhagia* — h. Uterine, *Metrorrhagia*.

HEMORRHAGIC PLEURISY, *Hæmatothorax*.

HÉMORRHAGIE BUCCALE, *Stomatorrhagia* — h. *Cerebrale*, *Apoplexy* — h. *du Cœur*, *Hæmocardiorrhagia* — h. *du Foie*, *Hepatorrhagia* — h. *Interstitielle*, *Apoplexy* — h. *Interstitielle du Poupon*, *Hæmoptysis* — h. *des Intestins*, *Hæmatochezia* — h. *de la Matrice*, *Metrorrhagia* — h. *Méningée*, *Apoplexy meningéale* — h. *de la Moëlle épinière*, *Apoplexia myelitica* — h. *Nasale*, *Epistaxis* — h. *de la Vessie*, *Cystirrhagia*.

HEMORRHAGIP'AROUS, (F.) *Hémorrhagipare*, from *hæmorrhagia*, 'hemorrhage,' and *parire*, 'to bring forth.' That which gives occasion to hemorrhage: thus, softening of the neurine may be *hæmorrhagiparous*.

HÉMORRHINIE, *Epistaxis*.

HÉMORRHOÏDAIRE (F.). One who is subject to hemorrhoids.

HÉMORRHOÏDES, *Hæmorrhoids* — h. *Aveugles*, *Cæca hæmorrhoides*. See *Hæmorrhoids*.

HEMORRHOSCOPIE, *Hæmatoscopia*.

HÉMOSPASIE, *Hæmospasia*.

HÉMOSPASIQUE, *Hæmospastic*.

HÉMOSTASIE, *Hæmostasia*.

HÉMOSTATICS, *Hæmatostatica*.

HÉMOSTATIQUES, *Hæmatostatica*.

HEMP, INDIAN, *Apocynum cannabinum*, *Bangue* — h. Wild, *Ambrosia trifida*.

HEMPWEED, AROMATIC, *Eupatorium aromaticum* — h. Roundleaved, *Eupatorium rotundifolium*.

HEMPSEED, see *Cannabis sativa*.

HENBANE, *Hyoscyamus*.

HENBIT, *Lamium amplexicaule*.

HEN-BLINDNESS, *Hemeralopia*.

HENNÉ, *Lawsonia inermis*.

HENRICEA PHARMACEARCHA, *Gentiana chirayita*.

HENRICUS RUBENS, *Coleothar*.

HEPAR, 'ἥπαρ, 'genitive,' 'ἥπατος, 'liver.' A name for substances resembling liver in appearance. The ancient name for the liver of sulphur, *Hepar sulphuris*; which is sometimes a compound of sulphur and potassium; at others, of sulphur and potassa. See *Potassæ sulphuretum*.

HEPAR ADIPOSUM, *Adiposus hepaticus* — h. *Adulterinum*, *Spleen*.

HEPAR ANTIMONIA'TUM is a compound of a sulphuret of antimony and an alkali. See *Oxydum stibii sulphuratum*.

HEPAR MARTIA'LE. A compound of sulphuret of potass and an oxide of iron.

HEPAR SANGUINIS, see *Blood* — h. *Sinistrum*, *Spleen*.

HEPAR SUCCENTURIA'TUM. An additional liver or liver lobe; a case of malformation.

HEPAR SULPHURIS SALINUM, *Potassæ sulphuretum* — h. *Sulphuris volatile*, *Ammoniac sulphuretum* — h. *Uterinum*, *Placenta* — h. *Variegatum*, *Liver*, *nutmeg*.

HEPATAL'GIA, *Hepatodyn'ia*, *Neural'gia hepatis*, *Col'ica hepatic'a*, from 'ἥπαρ, 'ἥπατος, 'liver,' and *αλγος*, 'pain.' Pain in the liver. *Neuralgia of the liver*, (F.) *Néuralgie du Foie*.

HEPATALGIA CALCULOSA, *Colica hepatica* — h. *Petitiana*, *Turgescencia vesiculæ fellæ* — h. *Phlegmonoides*, *Hepatitis*.

HEPATAPOSTE'MA, from 'ἥπαρ, 'ἥπατος, and *ἀποστήμα*, 'an abscess.' Abscess of the liver.

HEPATARIUS, *Hepaticus*.

HEPATATROPH'IA, *Atroph'ia seu Aridit'ra hepatis*, from 'ἥπαρ, 'ἥπατος, and *atrophia*, 'atrophy.' Atrophy of the liver: — a general concomitant of *Cirrho'sis hepatis*.

HEPATAUXÉ, *Hypertroph'ia hepatis*, (F.) *Hyperhépatotrophie*, *Hypertrophie du foie*. Hypertrophy of the liver.

HEPATECHE'MA, *Son'tus hepaticus*, from 'ἥπαρ, 'ἥπατος, and 'ἤχημα, 'sound.' Sound rendered by the liver on percussion.

HEPATEMPHRAX'IS, from 'ἥπαρ, 'ἥπατος, and *ἐμφρασις*, 'I obstruct.' *Emphrax'is hepatis*. Hepatic obstruction. — *Plouquet*.

HEPATENCEPHALO'OMA, *Hepatomyelo'ma*, *Fungus hepatis medullaris*, from 'ἥπαρ, 'ἥπατος, and *ἐγκεφαλος*, 'encephalon.' *Encephaloid of the liver*.

HEPATER'US, *Hepaticus*, *Jecora'rius*. A variety of diarrhoea, *Fluvius hepaticus*. — *Gorræus*.

HEPATI'HEMORRHAG'IA, *Hæmorrhag'ia hepatic'a*, *Hepatorrhag'ia*, *Apoplex'ia hepatic'a*, (F.) *Hémorrhagie du foie*, *Hémohépatorrhagie*, from 'ἥπαρ, 'ἥπατος, and *αἱμορραγία*, 'hemorrhage.' Hemorrhage from the liver.

HEPAT'IC, *Hepat'icus*, *Hepata'rius*, *Hepate'rus*, *Hepato'i'des*, *Hepato'des*, *Jecora'rius*, *Jec'oral*, (F.) *Hépatique Jécoraire*, from *ἥπαρ*, *ἥπατος*. Belonging or relating to or resembling liver.

HEPATIC AR'TERY, *Arte'ria Hepat'ica*. One of the three branches given off by the coeliac. It passes towards the inferior surface of the liver; where it divides into two branches, a *right* and a *left*, which proceed towards the corresponding parts of that organ. The right branch gives off the cystic artery. Before dividing, the hepatic artery sends off two considerable branches, the *A. pylorica* and *Gastro-epiploica dextra*.

HEPATIC DUCT, *Ductus hepatic'us*, (F.) *Canal hépatique*, is about three fingers' breadth in length, and of the size of a quill. It is formed by the union of the biliary ducts, and joins the cystic duct at a very acute angle, to form the ductus choledochus. Its function is to convey the bile from the liver towards the duodenum.

HEPATIC ISLETS, see Liver—h. Lobules, see Liver.

HEPATIC PLEXUS, *Plexus hepatic'us*, consists of nervous filaments, sent by the coeliac plexus to the liver, which accompany the hepatic artery.

HEPATIC VEINS, *Supra-hepatic Veins*, *Vena cava hepatic'a*, (F.) *Veines sus-hépatiques*, *Intra-lobular veins*, do not follow the course of the arteries of the same name. They arise in the substance of the liver; converge towards the posterior margin of that viscus, and open into the vena cava inferior. They convey away the blood carried to the liver by the hepatic artery and vena porta.

HEPATICA. Medicines believed to be capable of affecting the liver.

HEPATICA, *H. triloba*—h. *Americana*, *H. triloba*—h. *Fontana*, *Marchantia polymorpha*—h. *Nobilis*, *H. triloba*—h. *Stellata*, *Asperula odorata*.

HEPATICA TRI'L'OB'A, *H. America'na*, *Anemo'né hepatic'a* seu *nobil'is*, *Herba tri'nit'is*, *Hepatic'a*, *Hepatic'us flos*, *Trifolium hepatic'um* seu *an'reum*, *Liverwort*, *Liverweed*, *Trefoil*, *Noble Liverwort*, *Herb Trinity*, (F.) *Hépatique des jardins*. The leaves of this plant—*Hepatic'a* (Ph. U. S.)—are a simple astringent, but not possessed of much virtue.

HEPATICULA, *Hepatitis*, chronic.

HEPATICUS, *Hepateros*—h. *Flos*, *Hepatica triloba*.

HEPATIFICATIO, *Hepatization*.

HÉPATIQUE, *Hepatic*—h. *Étoilée*, *Asperula odorata*—h. *des Fontaines*, *Marchantia polymorpha*—h. *des Jardins*, *Hepatica triloba*.

HEPATIRRHŒA, *Fluxus hepatic'us*, *Dysenter'ia hepatic'a*, *Diarrhœa hepatic'a*, *Hepatorrhœa*, *Hepatocholorrhœa*, *Hepatodysenter'ia*, *Hepatorrhag'ia*, *Hæmat'ia*, *Aimat'ia*, (F.) *Flux hépatique*, from *ἥπαρ*, *ἥπατος*, and *ρῶω*, 'I flow.' A species of diarrhœa in which the excreted matters seem to come from the liver, or are much mixed with bile.

HEPATIS EMUNCTORIA, *Inguen*—h. *Suspensorium*, *Ligament*, *suspensory*, of the liver.

HEPATISATIO, *Hepatization*—h. *Pulmonum*, *Hepatization of the Lungs*.

HÉPATISATION, *Hepatization*—h. *Grisee*, see *Hepatization*—h. *Rouge*, see *Hepatization*.

HEPATITES VENA, *Cava vena*.

HEPATIT'IS, *Empres'ma hepatic'is*, *Cauma* seu *Inflamma'tio hepatic'a*, *I. Jecin'oris*, *Febris hepatic'a inflammato'ria*, *Hepatic'gia phlegmonoi'des*, *Morbus jecin'oris*, *Hepatophlegmonë*, *Inflammation of the liver*, (F.) *Hépatite*, *inflammation du foie*, *Pièce*, (Provincial); from *ἥπαρ*, *ἥπατος*, 'liver,' and *itis* denoting inflammation.

It may be seated either in the peritoneal covering, *Sero-hepatit'is*, *Hepatic'is velamento'sa* seu *extern'a*, *Peritonit'is hepatic'a*, *Perihepatit'is*, or in the substance of the liver, *Hepatic'is parenchymato'sa*, or in both, *Puro-hepatit'is*, and may be acute or chronic. The peculiar symptoms are:—pain in the right hypochondrium, shooting to the back and right shoulder, and increased on pressure; difficulty of lying on the left side; sometimes jaundice with cough, and synocha. Its termination is generally by resolution:—in tropical climates it often runs on to suppuration, *Jec'oris vomica*, *Hepatic'is apostemato'sa*, the abscess breaking either externally, or forming a communication with the intestines or chest, or breaking into the cavity of the abdomen. The causes are those of inflammation in general. Heat predisposes to it; hence its greater frequency in hot climates. On dissection of those who have died of it, the liver has been found hard and enlarged; colour of a deep purple; or the membranes have been more or less vascular; or adhesions, or tubercles, or hydatids, or abscesses, or biliary calculi may be met with. The treatment must be bold. Bleeding, general and local, fomentations, blisters, purgatives, and the antiphlogistic regimen. In hot climates especially, a new action must be excited by mercury as early as possible.

HEPATITIS APOSTEMATOSA, see *Hepatitis*.

HEPATITIS, CHRONIC, *Hepatic'is chronica*, *Inflamma'tio hepatic'a lenta*, *Hepatitis ocul'ia*, *Hepatic'ula*, *Subinflammato'ria hepatic'a*, (F.) *Chronic'hépatite*, *Chronic liver disease*, is not as common as is believed. It may be suspected from the existence of the symptoms above mentioned, when in a minor degree; enlargement, constant dull pain in the region of the liver; sallow countenance; high-coloured urine; clay-coloured fæces, &c. The great object of treatment is to excite a new action by mercury and counter-irritants, and to keep the liver free by cathartics.

HEPATITIS CYSTICA, *Cholecystitis*—h. *Externa*, see *Hepatitis*—h. *Occulta*, *H. chronic*—h. *Velamentosa*, see *Hepatitis*—h. *Parenchymatosa*, see *Hepatitis*.

HEPATIZATION, *Hepatisa'tio*, *Hepaticifa'tio*, from *ἥπαρ*, *ἥπατος*. Conversion into a liver-like substance. Applied to the lungs when gorged with effused matters, so that they are no longer pervious to the air—*Hepatisa'tio pulmo'num*, *Carnifica'tio pulmo'num*. In such state, they are said to be *hepatized*.

HEPATIZATION, RED, (F.) *Hépatisation rouge*, *Endurcissement rouge*, *Ramollissement rouge*, characterizes the first stage of consolidation of the lungs in pneumonia.

HEPATIZATION, GRAY, (F.) *Hépatisation grise*, *Induration grise*, *Ramollissement gris*, *Infiltration purulente*, characterizes the third stage, or stage of purulent infiltration.

HEPATIZON, *Chloasma*.

HEPATOC'ACÉ, from *ἥπαρ*, *ἥπατος*, and *κακος*, 'evil,' *Gangrene of the liver*.

HEPATOCE'LE, from *ἥπαρ*, *ἥπατος*, and *κῆλη*, 'a tumour,' *Hernia hepatic'a*; *Hernia of the liver*, (F.) *Hernie du foie*. The liver has never been found entirely out of the abdominal cavity. Increase of its bulk, or injuries of the parietes of the abdomen, have been the sole cause of the protrusions which have been occasionally met with, especially in infants, in whom the upper part of the linea alba is very weak, and indeed scarcely seems to exist. Sauvages has distinguished two species of hepatocèle:—the *ventral* (in the linea alba), and the *umbilical* or *hepatophthalmum*.

HEPATOCHOLORRHEA, *Hepatirrhœa*.

HEPATOCOLICUM. A ligament of the liver, described by Haller, as passing from the gall-bladder and contiguous sinus portarum, across the duodenum, to the colon. Another, termed *Hepato-renal*, descends from the root of the liver to the kidney. They are both peritoneal.

HEPATOCYSTIC, *Hepatocysticus*, *Cystidopaticus*, from *ἥπαρ*, *ἥπατος*, and *κύστις*, 'bladder.' Relating to the liver and gall-bladder.

HEPATOCYSTIC DUCT, *Ductus hepatocysticus*. The choledoch duct.

HEPATODYME, *Hepatodymus*, from *ἥπαρ*, *ἥπατος*, and *δίδυμος*, 'a twin.' A monster united at the liver, and double above.

HEPATODYNIA, *Hepatalgia*.

HEPATODYSENTERIA, *Hepatirrhœa*.

HEPATOGASTRIC, *Gastrohepatic*.

HEPATOGASTROCHOLOSIS, Fever, bilious, Fever, gastric.

HEPATOGRAPHY, *Hepatographia*; from *ἥπαρ*, *ἥπατος*, and *γραφία*, 'a description.' The part of anatomy which describes the liver.

HEPATOHÆMIA, *Hyperhæmia hepatis*, *Hepatic Engorgement*, (F.) *Hyperémie du Foie*, from *ἥπαρ*, *ἥπατος*, and *αἷμα*, 'blood.' Sanguineous congestion of the liver.

HEPATODES, *Hepatic*.

HEPATOIDES, *Hepatic*.

HEPATOLITHIASIS, from *ἥπαρ*, *ἥπατος*, and *λίθιασις*, 'formation of stone.' The formation of concretions, *Hepatolithia*, in the liver.

HEPATOLGY, *Hepatology*, from *ἥπαρ*, *ἥπατος*, and *λόγος*, 'a discourse,' 'treatise.' A treatise on the liver.

HEPATOMALACIA, *Malacoësis hepatis*, (F.) *Ramollissement du Foie*. Softening of the liver.

HEPATOMPHALUS, from *ἥπαρ*, *ἥπατος*, 'liver,' and *ομφαλός*, 'navel.' Hernia of the liver at the umbilicus.

HEPATOMYELOMA, *Hepatencephaloma*.

HEPATONCUS, from *ἥπαρ*, *ἥπατος*, and *ὄγκος*, 'a tumour.' Tumefaction of the liver.

HEPATOPAREC'TAMA, from *ἥπαρ*, *ἥπατος*, and *παρεκταμα*, 'considerable extension.' Excessive enlargement of the liver.

HEPATOPATHIA, from *ἥπαρ*, *ἥπατος*, and *πάθος*, 'suffering.' *Liver-disease*. Disease of the liver.

HÉPATOPATHIE CANCÉREUSE, *Hepatoscirrhus*—*h. Tuberculeuse*, *Hépatostromosie*.

HEPATOPHLEGMONÉ, *Hepatitis*.

HEPATOPHTHISIS, *Phthisis hepatica*, from *ἥπαρ*, *ἥπατος*, and *φθίω*, 'I consume.' Consumption from suppuration of the liver.

HEPATORRHAGIA, *Hepatirrhœa*, *Hepathæmorrhagia*.

HEPATORRHEX'IS, from *ἥπαρ*, *ἥπατος*, and *ῥήξις*, 'rupture.' Rupture of the liver.

HEPATORRHŒA, *Hepatirrhœa*.

HÉPATOSARCOMIE, *Hepatoscirrhus*.

HEPATOSCIRRHUS, from *ἥπαρ*, *ἥπατος*, and *σκίρρος*, 'cancerous induration.' *Scirrhus hepatis*, *Encephalosis* or *Carcinoma* of the liver, (F.) *Hépatosarcomie*, *Hépatopathie cancéreuse*, *Cancer du Foie*. *Scirrhus* or *Cancer* of the liver.

HÉPATOSTRUMOSIE, *Tubercula hepatis*, (F.) *Tubercules du Foie*, *Hépatopathie tuberculeuse*; from *ἥπαρ*, *ἥπατος*, and *stroma*, 'a tumour,' 'a scrofulous tumour.' Tubercle of the liver.

HEPATOT'OMY, from *ἥπαρ*, *ἥπατος*, and *τέμνω*, 'I cut.' Dissection of the liver.

HEPIALOS, *Epialos*.

HEPS, see *Rosa canina*.

HEPSEMA, *Decoction*.

HEPSESIS, *Decoction*.

HEPTAL CYCLE, see *Cycle*.

HEPTAL'ON GRAV'EOLENS, *Hogwort*, *Bears' fright*. An indigenous plant, which has a fetid porcine smell; and is said to be used by the Indians as a diaphoretic, cathartic, &c.

HEPTAPHAR'MACUM, from *επτα*, 'seven,' and *φάρμακον*, 'a remedy.' A medicine composed of seven substances; cerusse, litharge, pitch, wax, colophony, frankincense, and bullock's fat. It was regarded as laxative, suppurating and healing.

HEPTAPHYLLUM, *Tormentilla*.

HEPTAPLEURON, *Plantago major*.

HERACLEUM, see *H. lanatum*—*h. Branca*, *H. spondylium*.

HERACLE'UM SPONDYL'UM, *H. Branca*, *Branca ursina* seu *Germanica*, *Spondylium*, *Spondylium*, *seu Parsnep*, *All-heal*, *Cone-mumble*. (Sc.) *Buneward*, (Prov.) *Cushia*, (F.) *Berce*, *Brancursine bâtarde*, *Fausse Acanthe*. Order, *Umbelliferae*. *Sex. Syst.* Pentandria Digynia. Dedicated to *Ἡρακλῆς*, 'Hercules.' The root of this plant has a strong, rank smell; and a pungent, almost caustic taste. It has been given as a tonic, stomachic and carminative; both in powder and in decoction. The Russians, Lithuanians, and Poles obtain from its seeds and leaves, by fermentation, a very intoxicating spirituous liquor, which they call *Parst*.

The root of *Heracle'um Lana'tum*, *Masterwort*, *Cow parsnep*,—*Heracle'um* (Ph. U. S.)—is in the secondary list of the Pharmacopœia of the United States.

HERACLEUS MORBUS, *Epilepsy*.

HERB, *Herba*, *Bot'anê*. Any ligneous plant, which loses its stalk during the winter.

HERB, BLESSED, *Geum urbanum*—*h. Christopher*, *Actæa spicata*.

HERB-DOCTOR, *Botanical doctor* or *physic'ian*. One who treats diseases altogether by herbs; as the—so called—"Thompsonians."

HERB OF GRACE, *Gratiola officinalis*—*h. Mastich*, common, *Thymus mastichina*—*h. Mastich*, *Syrian*, *Teucrium marum*—*h. Peter*, *Primula veris*—*h. Robert*, *Geranium Robertianum*—*h. of Saint Cunegonde*, *Eupatorium cannabinum*—*h. Sophia*, *Sisymbrium sophia*—*h. Trinity*, *Hepatica triloba*.

HERBA, *Herb*—*h. Alexandrina*, *Smyrniolum olusatrum*—*h. Althæa*, *Pelargonium cucullatum*—*h. Anthos*, *Rosmarinus*—*h. Benedicta*, *Geum urbanum*—*h. Britannica*, *Rumex hydroslaphium*—*h. Canni*, *Artemisia santonica*—*h. Cardiaca*, *Leonurus cardiaca*—*h. Cardui veneris*, *Dipsacus fullonum*—*h. Dorea*, *Solidago virgaurea*—*h. Felis*, *Nepeta*—*h. Genipi*, *Achillea atrata*—*h. Ignis*, *Lichen pyxidatus*—*h. Melancholifuga*, *Fumaria*—*h. Militaris*, *Achillea millefolium*—*h. Papillaris*, *Lapsana*—*h. Paralyseos*, *Primula veris*—*h. Paralytica*, *Primula veris*—*h. Patæ lapinæ*, *Leonurus cardiaca*—*h. Pulicaris*, *Plantago psyllium*—*h. Quercini*, *Lichen plicatus*—*h. Sacra*, *Verbena officinalis*—*h. Salivaris*, *Anthemis pyrethrum*—*h. Sardonica*, *Ranunculus sceleratus*—*h. Scelerata*, *Ranunculus scleratus*—*h. Sideritidis*, *Galeopsis grandiflora*—*h. Tabaci*, *Nicotiana tabacum*—*h. Trinitatis*, *Anemone hepatica*, *Viola tricolor*—*h. Veneris*, *Adiantum pedatum*—*h. Ventis*, *Anemone pulsatilla*—*h. Vitri*, see *Salsola kali*—*h. Zazarhendi*, *Origanum*.

HERBÆ PRO ENEMATE, *Glyster herbs*—*h. pro fotu*, *Fomentation herbs*.

HERBALIST, *Herborist*.

HERBARIUM, from *herba*, a plant. (F.)

Herbar. A collection of plants. Generally applied to a collection of dried plants — *Hortus sicca.* In Pharmacy, a plant that is used entire.

HERBARIUS, Herborist.

HERBE. Herb — *h. aux Abeilles*, *Spiræa ulmaria* — *h. au Cancer*, *Plumbago Europæa* — *h. au Centaure*, *Chironia centaurium* — *h. aux Chantres*, *Erysimum* — *h. aux Choqueriers*, *Achillea millefolium*, *Justicia pectoralis* — *h. aux Chauts*, *Nepeta*, *Tenarium marum* — *h. au Coq*, *Tanacetum balsamita* — *h. au Cuillers*, *Cochlearia officinalis* — *h. aux Écorcelles*, *Scrophularia nodosa* — *h. aux Écus*, *Lysimachia nummularia* — *h. à l'Éternuer*, *Achillea ptarmica* — *h. à la Fidère*, *Chironia centaurium* — *h. aux Femmes battues*, *Tamus communis* — *h. de Grace*, *Ruta* — *h. aux Gueux*, *Clematis vitalba* — *h. aux Hémorrhoides*, *Ranunculus ficaria* — *h. à l'Hirondelle*, *Chelidonium majus* — *h. à la Houette*, *Asclepias Syriaca* — *h. d'Irroque*, *Lolium temulentum* — *h. des Juifs*, *Solidago virginæa* — *h. aux Lades*, *Veronica* — *h. aux Mamelles*, *Lapsana* — *h. aux Monches*, *Conyza squarrosa* — *h. aux Murrailles*, *Parietaria* — *h. des Nones*, *Parietaria* — *h. aux Oies*, *Potentilla reptans* — *h. à Paris*, *Paris* — *h. à la Paralysie*, *Primula veris* — *h. au Pauvre homme*, *Gratiola officinalis* — *h. aux Perles*, *Lithospermum officinale* — *h. à Pisser*, *Pyrola umbellata* — *h. à la Poudre de Chypre*, *Hibiscus abelmoschus* — *h. aux Poux*, *Delphinium staphisagria* — *h. à la Puce*, *see Apocynum hypericifolium* — *h. aux Puces commune*, *Plantago psyllium* — *h. à Robert*, *Geranium Robertianum* — *h. à la Roste*, *Drosera rotundifolia* — *h. de Sainte Barbe*, *Erysimum barbarea* — *h. Sainte Christophe*, *Actæa spicata* — *h. de Sainte Étienne*, *Circæa luteana* — *h. Saint Fiacre*, *Verbascum thapsus* — *h. de Saint Jean*, *Artemisia vulgaris* — *h. Saint Jean*, *Glechoma hederacea* — *h. Saint Quirin*, *Tussilago* — *h. de Saint Roch*, *Inula dysenterica* — *h. Sainte Rose*, *Pæonia* — *h. aux Sorciers*, *Circæa luteana*, *Datura stramonium* — *h. aux Toiguesux*, *Arctium lappa* — *h. à Tous les maux*, *Verbena officinalis* — *h. Turque*, *Herniaria glabra* — *h. aux Vexes*, *Tanacetum vulgare* — *h. aux Verrues*, *Heliotropium Europæum*.

HERBIER, Herbarium, Ingluvies.

HERBIVOROUS, *Herbiv'orus*, from *herba*, 'grass,' and *voro*, 'I eat.' An epithet applied to animals which feed on herbs.

HERBORIST, *Herba'rius*. One who deals in useful plants. An *Herb'alist* or *Herb'arist*.

HERBORIZATION, *Herba'rum inquisit'io*. An excursion, made with the view of collecting plants. Such excursions are directed by the Apothecaries' Company of London, for the use of their apprentices, &c.

HERBS, FIVE CAP'ILLARY, *Quinque herbæ capilla'res*, were, anciently, hart's tongue; black, white, and golden maiden-hair, and spleenwort.

HERBS, FIVE EMOL'IENT, *Quinque herbæ emol'ientes*, were, anciently, beet, mallow, marsh-mallow, French mercury, and violet.

HERCULES ALLHEAL, *Pastinaca opoponax*. *HERCULES MORBUS*, Epilepsy.

HERÉDITAIRE, Hereditary.

HERÉDITARY, *Hæreditarius*, *Hæreditarius*, *Gentili'tius*, *Syn'phytus*, *Syn'genes*, (F.) *Héréditaire*, from *hæris*, *hæredis*, 'an heir.' An epithet given to diseases, communicated from progenitors. Such diseases may exist at birth; or they may supervene at a more or less advanced period of existence. *Hereditary diseases*, *Morbi hereditarii*, (F.) *Maladies héréditaires*, often prevail amongst several members of a family, or are family diseases or complaints.

HERMAPHRODITE, *Hermaphrodis'ia*, *Hermaphrodit'us*, *Hermaphrodis'mus*, *Fab'rica androg'yna*, *Androgyn'ia*, *Gynand'ria*, *Hermaph'*

rodism, *Hermaph'roditism*; from *Ἑρμης* 'Mercury,' and *Ἀφροδίτη*, 'Venus.' Appertaining to Mercury and Venus. Union of the two sexes in the same individual.

HERMAPHRODISIA, Hermaphrodeity.

HERMAPHRODITE. Same etymon. *Hermaphrodit'us*, *G'y'nida*, *Androg'ynus*, (Sc.) *Searcht*, (Prov.) *Scrat*. One who possesses the attributes of male and female: who unites in himself the two sexes. A term, applied to an animal or plant which is, at the same time, both male and female. True hermaphrodites are only met with in the lower degrees of the animal scale, amongst the zoophytes, mollusca, or gasteropoda. The individuals of the human species, regarded as hermaphrodites, owe this appearance to a vicious conformation of the genital organs: a kind of monstrosity, which renders them unfit for generation, although an attentive examination may exhibit the true sex. Hermaphrodites have, likewise, been described, which, instead of uniting the attributes of both sexes, cannot be considered male or female. These have been called *neutral hermaphrodites*.

HERMAPHRODITISMUS, Hermaphrodeity.

HERMAPHRODITUS, Hermaphrodite.

HERMETICA DOCTRINA, *Hermetica ars*, (F.) *Hermétique*. The doctrine of Hermes, a celebrated Egyptian philosopher, who is considered the father of alchemy. That part of chymistry, whose object was the pretended transmutation of the metals.

HERMODACTYLUS, *Dactyle'tus*, *An'ima articulo'rum*, from *Ἑρμης*, 'Mercury,' and *δάκτυλος*, 'a finger;' or rather from *Hermus*, a river in Asia, upon whose banks it grows, and *δάκτυλος*, 'a date;' or from *Ἑρμης*, 'Mercury,' and *δάκτυλος*, 'a date;' (F.) *Hermodactyle*, *Hermodactæ* or *Hermodate*. The root of the *Hermodactyl* was formerly used as a cathartic. By some, it is supposed to be identical with the *Iris tuberosa*. The best testimony seems to be in favour of its being a variety of the colchicum. — *Col'chicum Illy'ricum*. Under the name *Hermodactyle*, roots or bulbs of an unknown plant, growing in the countries about the eastern extremity of the Mediterranean, are sold in the shops of Europe.

HERMODATE, *Hermodactylus*.

HERMOPH'ILUS, from *Ἑρμης*, 'Mercury,' and *φίλος*, 'a lover.' One who is fond of mercury as a medicine.

HERNIA, *Ramez*, *Ruptu'ra*, *R. hernio'sa*, *Crepitu'ra*, *Ectop'ia hernio'sa*, *Celè*, *Rupture*, *Burst*, (F.) *Hargne*, *Descente*, *Effort*, *Greveure*, *Rompure*. From *ερπος*, 'a shoot' [?]. Any tumour, formed by the displacement of a viscus or a portion of a viscus, which has escaped from its natural cavity by some aperture, and projects externally. Herniæ have been divided into, — 1. *Hernia of the Brain*; *Encephalocele*; 2. *Hernia of the Thorax*; *Pneumocèle*; 3. *Hernia of the Abdomen*.

Abdom'inal Herniæ are remarkable for their frequency, variety, and the danger attending them. They are produced by the protrusion of the viscera, contained in the abdomen, through the natural or accidental apertures in the parietes of that cavity. The organs, which form them most frequently, are the intestines and the epiploon. These herniæ have been divided, according to the apertures by which they escape, into:

1. *In'guinal* or *Supra-Pu'b'ian Hernia*. These issue by the inguinal canal: they are called *Bubonoce'lè*, when small; and *Scro'tal Hernia* or *Ochoceol'è*, in man, when they descend into the scrotum: — *Vulvar Hernia* or *Puden'dal* or *La'b'ial Hernia*, *Episioce'lè*, in women, when they extend to the labia majora. 2. *Crural* or *Fem'oral Hernia*, *Me'roce'lè*, when they issue by the

crural canal. 3. *Infra-Pubian Hernia*, (F.) *Hernie sous-pubienne*, *Ovidocœlæ*, *Opeocœlæ*, *Opodocœlæ*, *Hernia foraminis ovalis*, when the viscera escape through the opening, which gives passage to the infra-pubian vessels. 4. *Ischiatic Hernia*; when it takes place through the sacro-sciatic notch. 5. *Umbilical Hernia*, *Exomphalos*, *Omphalocœlæ*; when it occurs at the umbilicus or near it. 6. *Epigastric Hernia*;—occurring through the linea alba, above the umbilicus. 7. *Hypogastric or Infra-umbilical Hernia*, *Celiacœlæ*, *Hypogastricœlæ*,—when it occurs through the linea alba below the umbilicus. 8. *Perineal Hernia*, *Mesocœlæ*, *Hernia perinæi*, *Perinæocœlæ*, *Perineocœlæ*—when it takes place through the levator ani, and appears at the perineum. 9. *Vaginal Hernia*, *Coleocœlæ* seu *Elytracœlæ*—through the parieties of the vagina. 10. *Diaphragmatic Hernia*, *Diaphragmatocœlæ*; when it passes through the diaphragm.

Hernia are likewise distinguished,—according to the viscera forming them,—into *Enterocœlæ*, *Epiplocœlæ*, *En'tero-epiplocœlæ*, *Gastrocœlæ*, *Cystocœlæ*, *Hepatocœlæ*, *Splenocœlæ*, &c.

When a hernia can be restored to its natural cavity, by the aid of pressure, &c., properly applied, it is said to be *reducible*. It is, on the contrary, *irreducible*, when adhesion, bulk, &c., oppose its return. When the aperture, which has given passage to the hernia, occasions more or less constriction on the protruded portion, the hernia is said to be *incarcerated or strangulated*; and, if the constriction be not removed, constipation, hiccough, vomiting, and all the signs of violent inflammation, followed by gangrene, supervene, with alteration of the features, small pulse, cold extremities, and death.

The therapeutical indications are,—1. *As regards reducible hernia*:—to replace the viscera in the abdomen by the taxis; and to retain them there by the use of a *truss*, which, if properly adapted, may effect a radical cure. 2. *As regards irreducible hernia*:—to support the tumour by an appropriate suspensory bandage. 3. *As regards strangulated hernia*:—to have recourse to the taxis; blood-letting; warm bath; tobacco gylsters; ice to the tumour; and, if these should not succeed, to perform an operation, which consists in dividing the covering of the hernia, and cutting the aponeurotic ring, which causes the strangulation;—reducing the displaced viscera, unless their diseased condition should require them to be retained without;—dressing the wound appropriately:—restoring the course of the fæces by means of gentle gylsters—preventing or combating inflammation of the abdominal viscera;—conducting the wound to cicatrization, by appropriate means; and afterwards supporting the cicatrix by a bandage.

The word *Hernia* was also used, of old, for the scrotum, and, not unfrequently, for the testicle.

HERNIA, ANEURISMAL, *Cirsonphalus*—h. *Arteriarum*, see *Aneurism*—h. of the Bladder, *Cystocœlæ*—h. *Bronchialis*, *Bronchocœlæ*—h. *Carnosa*, *Sarcocœlæ*—h. of the Cerebellum, *Paracephalocœlæ*—h. *Cerebri*, *Encephalocœlæ*.

HERNIA, CONGENITAL, *Hernia congenita*, is a protrusion of some of the contents of the abdomen into the *Tunica vaginalis testis*, owing to a want of adhesion between its sides, after the descent of the testicle.

HERNIA CORNEÆ, *Ceratocœlæ*—h. *Crural*, *Merocele*—h. *Epiploicæ*, *Epiplocœlæ*—h. *Femoral*, *Merocele*—h. *Foraminis Ovalis*, see *Hernia*—h. *Gutturis*, *Bronchocœlæ*—h. *Hepatica*, *Hepatocœlæ*.

HERNIA HUMORALIS, *Empres'ma Orchitis*, *Dydymitis*, *Orchitis*, *Orchiditis*, *Inflammatio testium*, *Inflammation of the Testicle*, *Swell'd Testi-*

tle, *Hernia Ven'eriæ*, *Orchiocele*, *Orchidocœlæ*, *Orchidioncus*, (F.) *Inflammation du testicule*, *Orchite*. Swelling and inflammation of the testicle is a common symptom of gonorrhœa, (F.) *Chancroëpisée tombée dans les Bourses*; but it may arise from external injuries, or from other causes. It is a disease which cannot be mistaken, and the treatment must obviously be strongly antiphlogistic, supporting the testicle during the treatment, and for some time afterwards: methodical compression has also been found useful. The disease is not generally of a serious character, going off as suddenly as it comes on. As it affects the epididymis more especially, when supervening on gonorrhœa or blennorrhœa, it is sometimes termed *blennorrhag'ic epididymitis*.

HERNIA INCARCERATA, *Entero-peristole*—h. *Inguinalis*, *Bubonocœlæ*—h. *Intestinalis*, *Enterocœlæ*—h. *Intestinalis omentalis*, *Entero-epiplocœlæ*—h. *Iridis*, *Ptoxis Iridis*—h. *Ischiatica*, *Ischiocele*—h. *Lienalis*, *Splenocœlæ*.

HERNIA LITT'ICA, *H. e. Diverticulo Intestini*. *Hernia* first described by Littre, in which the intestinal canal proper is not included in the hernial sac, the protruded portion of intestine consisting of a digital prolongation of the ileum, which Littre concluded was formed by the gradual extension of a knuckle of the bowel, that had been engaged in the inguinal canal.

HERNIA OF THE LIVER, *Hepatocœlæ*—h. of the Lung, *Pneumocœlæ*.

HERNIA, MESOCOLIC, *Hernia Mesocolica*. *Hernia*, caused by the bowels gliding between the layers of the mesocolon.—Sir A. Cooper.

HERNIA, OMENTAL, *Epiplocœlæ*—h. *Parorchido-enterica*, *Parorchido-enterocœlæ*—h. *Perinæi*, see *Hernia*—h. *Pharyngis*, *Pharyngocœlæ*—h. *Phrenicæ*, *Diaphragmaticæ* *hernia*—h. *Pinguedinosæ* *Scroti*, *Liparocœlæ*—h. of the Pleura, *Pleurocœlæ*—h. *Pleurica et pulmonalis*, *Pleurocœlæ*—h. *Putendalis*, *Putendal hernia*—h. *Sacci lacrymalis*, *Lachrymal hernia*, *Mucocœlæ*—h. *Scrotalis*, *Scrotocœlæ*—h. *Seminalis* *scroti*, *Spermatocele*—h. of the Stomach, *Gastrocœlæ*—h. *Suprapubian*, *H. inguinal*—h. of the Tongue, *Glossocœlæ*—h. *Umbilicalis*, *Epiplophalon*, *Exomphalos*—h. *Umbilicæ* *Aquosa*, *Hydromphalum*—h. *Utracæ*, *Uromphalus*—h. *Uteri*, *Hysterocœlæ*—h. *Varicose*, *Cirsecœlæ*—h. *Varicosa*, *Varicocœlæ*—h. *Venarum*, *Varix*—h. *Veneris*, *Hernia humoralis*—h. *Ventosa*, *Physocœlæ*—h. *Ventral*, see *Ventral*—h. *Ventriculi*, *Gastrocœlæ*—h. *Vesicæ* *Urinariæ*, *Cystocœlæ*—h. *Zirbalis*, *Epiplocœlæ*.

HERNIAIRE, *Hernial*, *Herniaria glabra*.

HERNIAL, *Herniarius*, (F.) *Herniaire*. Belonging to, or concerning hernia:—as *Hernial Bandage*, *Hernial Sac*, &c.

The (F.) *Hernié*, is applied to a part enveloped in a hernial sac. (F.) *Hernieux*, *Cele'ta*, *Celo'tes*, *Ruptured*, *Burst*, *Bursten*, means one affected with hernia; and *Chirurgien herniaire*, one who devotes himself to the treatment of hernia.

HERNARIA GLABRA, *H. vulgaris* seu *hirsuta* seu *alpes'tris* seu *anmia* seu *cine'ra* seu *fruticosa*, *Millig'ra*, *En'petrum*, *Rupturewort*, from *hernia*, 'rupture'. (F.) *Herniaire*, *Turquette*, *Herniale*, *Herbe Turque*. Ord. *Paronychieæ*. This plant, which as its name imports, was formerly considered efficacious in the cure of hernia, seems destitute of all virtues.

HERNIARIUS, *Hernial*.

HERNIE CHARNUE, *Sarcocœlæ*—h. *de l'Épiploon*, *Epiplocœlæ*—h. *de l'Estomac*, *Gastrocœlæ*—h. *du Foie*, *Hepatocœlæ*—h. *Inguinale*, *Bubonocœlæ*—h. *du Nombril*, *Exomphalos*—h. *Ombilicæ*, *Exomphalos*—h. *du Poupon*, *Pneumocœlæ*—h. *de la Vessie*, *Cystocœlæ*.

HERNIÉ, *Hernial*.

HERNIEMPHRAGMUS, *Herniempfrax'is*, from *hernia*, and *εμπαγμος*, 'obstruction.' The mechanical obstruction of a hernial canal for the radical cure of the hernia.

HERNIEUX, Hernial.

HERNIOLE, *Herniaria glabra*.

HERNIOTOMY, *Celotomia*.

HERO'IC, *Hero'icus*, *Hero'ius*, *Hero'us*, from *ἥρως*, 'a hero.' An epithet applied to remedies or practice of a violent character.

HERPEDON, *Herpes*.

HERPEN, *Herpes*.

HERPES, *Erpes*, *Herpe'don*, *Herpen*, *Serpens*, from *ἑρπω*, 'I creep,' because it creeps and spreads about the skin; *Echph'ysis Herpes*, *Cytis'ma Herpes*, *Tetter*, *Fret*, (F.) *Dartre*, *Olophlyctide*. A vesicular disease, which, in most of its forms, passes through a regular course of increase, maturation, decline, and termination, in from 10 to 14 days. The vesicles arise in distinct, but irregular clusters, which commonly appear in quick succession, and near together, on an inflamed base; generally attended with heat, pain, and considerable constitutional disorder. The term, like *and* others which refer to cutaneous diseases, has not been accurately defined. The ancients had three varieties: the military, *κρηταίος*; vesicular, *φλυκταίνων*, and eroding, *εσθιονέος*. Bateman has the following varieties:

1. HERPES PHLYCTÆNODES, *Herpes militaris*, *Echph'ysis Herpes Militaris*, *Nirles*, (F.) *Dartre phlyctænoides*, in which the vesicles are millet-sized; pellucid; clusters commencing on an uncertain part of the body, and being progressively strewed over the rest of the surface; succeeded by fresh crops. 2. HERPES ZOSTER, *Zoster*, *Zona ignea* seu *serpiginosa*, *Ignis Per'sicus*, *Cinzill'a*, *Cin'gulum*, *Sacer ignis*, *Echph'ysis Herpes zoster*, *Herpes peris'celis*, *Erysip'elas zoster* seu *phlyctænoides* seu *pustulosa*, *Zona*, *Circ'cinus*, *Perizo'ma*, (F.) *Ceinture*, *C. dartreusae*, *Feu Persique*, *Erysipèle pustuleux*, *Shingles*, (Prov.) *Cruels*, in which the vesicles are pearl-sized; the clusters spreading round the body like a girdle; at times confluent, and occasionally preceded by constitutional irritation. 3. HERPES CIRCINATUS, *Formica ambulatoria*, *An'nulus re'pens*, *Herpes Serpi'go*, *Serpi'go*, *Echph'ysis Herpes Circinatus*, *Ringworm*, *Vesic'ular Ringworm*, consisting of vesicles with a reddish base, uniting in rings; the area of the rings slightly discoloured, often followed by fresh crops. 4. HERPES LABIALIS, and 5. HERPES PRÆPUTIALIS, *Aphthæ præputii*, *Ulcus'cula præputii*, appearing, respectively, on the lips and prepuce. 6. HERPES IRIS, *Iris*, *Rainbow-worm*, occurring in small circular patches, each of which is composed of concentric rings of different colours. To these may be added, HERPES EX'EDENS, *H. eethiomi'enus* seu *depassens* seu *ferus* seu *eethiomi'enus* seu *ferox*, *Echph'ysis Herpes ex'edens*, *Darta exoriati'va* seu *maligna*, *Lupus non exedens*, *Formica corrosi'va*, *Formica*, *Pap'ula fera*, *Ul'cerative Ringworm*, *Ag'ria*, (F.) *Dartre rongeante*, in which the vesicles are hard; clusters thronged; fluid dense, yellow or reddish, hot, acrid, corroding the subjacent skin, and spreading in serpentine trails.

All the varieties demand simply an antiphlogistic treatment, when attended with febrile irritation. The *herpes circinatus*, alone, requires the use of astringent applications which have the power of repressing the eruption.

HERPES DEPASSENS, *H. exedens*—*h. Eethiomenus*, *H. exedens*—*h. Estiomenus*, *H. exedens*—*h. Farinosus*, *Pityriasis*—*h. Ferox*, *H. exedens*—*h. Ferus*, *H. exedens*—*h. Furfuraceus*, *Pityriasis*—*h. Furfuraceus circinatus*, *Lep'ra*, *H. exedens*—*h. Miliaris*, *H. phlyctænoides*—*h. Peris-*

celis, *H. zoster*—*h. Serpigo*, *H. circinatus*—*h. Tonsurans*, *Porri'go decalvans*.

HERPET'IC, *Herpet'icus*, (F.) *Herpétique*, *Dartreux*. Possessing the nature of herpes.

HERPETOGRAPHY, *Herpetograph'ia*, from *ἑρπης*, 'herpes,' and *γραφία*, 'a description.' A description of the different forms of herpes, as *Herpetology*, *Herpetolog'ia* is a treatise on the same, and also on reptiles, from *ἑρπετος*, 'a reptile.'

HERPETON, *Herpet'ion*, from *ἑρπετν*, 'to creep.' A creeping eruption or ulcer.—*Hippocrates*.

HERPYLOS, *Thymus serpyllum*.

HERRENSCHWAND'S SPECIFIC, see Specific of Herrenschwand.

HERIN, *Urine*.

HERSKET, *Ardor ventriculi*.

HERPERIS ALLIARIA, *Alliaria*.

HERTERADELPH'IA, *Heterodidym'ia*, *Heterodym'ia*, from *ἑτερος*, 'other,' and *ἀδελφος*, 'a brother.' A double monstrosity, in which the components of the double bodies are very unequal, and of which one portion may be regarded as the stem or trunk, to which another organized part, or even a whole body, less developed than itself, is affixed like a parasite.

HERTERADEN'IC, *Heteraden'icus*, (F.) *Hétéradénique*, from *ἑτερος*, 'other,' and *ἀδην*, 'a gland.' An accidental tissue is so termed, which has a glandular structure, but occurs in parts devoid of glands.

HETEROCHRON'ICUS, *Heteroch'ronus*, from *ἑτερος*, and *χρονος*, 'time.' Relating to difference of time.

Pulsus heterochron'icus. A pulse of varying rhythm. An irregular or intermittent pulse.

HETEROCHYMEU'SIS, from *ἑτερος*, and *χυμεισις*, 'mixture.' A state of the blood in which it contains other matters than in health, as urea, bile, &c.

HETEROCOLITE, see Homology, and Tissues

HETEROCRANIA, *Hemicrania*.

HETEROCRIN'IA, from *ἑτερος*, and *κρινω*, 'I separate.' Modification in the situation of secretions.—*Andral*.

HETERODIDYMIA, *Heteradelphia*.

HETERODYMIA, *Heteradelphia*.

HETEROGENESIS, see Generation. Also, any anomaly in organic formation.

HETEROLALIA, *Heterophonia*.

HETEROLOGOUS TISSUES, see Tissues.

HETEROMORPHISM, *Heteromorphi'mus*, from *ἑτερος*, and *μορφή*, 'shape.' A deviation from the natural shape of parts.

HETEROMORPHOUS, see Homology.

HETEROPA'GIA; from *ἑτερος*, and *παγω*, 'I unite together.' A double monstrosity, in which the accessory subject, which is very small and imperfect, but provided with a distinct head, and at least rudimental lower limbs, has its body implanted on the anterior surface of the body of the principal subject.

HETEROPATHIC, *Allopathic*.

HETEROPHON'IA, from *ἑτερος*, and *φωνή*, 'voice.' A cracked or broken voice. A change of the voice or speech—*Heterolalia*.

HETEROPHTHAL'MIA, from *ἑτερος*, and *ὀφθαλμος*, 'eye.' A difference in the two eyes,—as when one squints, or is of a different colour.

HETEROPLASM'US, from *ἑτερος*, and *πλασμα*, 'formation.' A morbid formation foreign to the economy, and which is composed of elements different from those that are found in it in the normal condition.

HETEROPLAS'TY, *Heteroplas'ticē*, *Heteroplas'tia*, from *ἑτερος*, and *πλασσω*, 'I form.' Irregular plastic or formative actions or operations, that do not admit of exact classification.

HETEROPROSOP'US, from *ἑτερος*, and *προ-*

ὄμων, 'countenance.' A monster having two faces.—Gurlt.

HETEROPUS, from *ἑτερος*, and *πους*, 'foot.' One who has one foot different from the other.

HETERORENNA, Malacia.

HETERORRHYTHMUS, from *ἑτερος*, and *ῥυθμος*, 'rhythm.' Having another rhythm. An epithet given to the pulse, when it is such, in any individual, as is usually felt at a different age.

HETEROSARCOSES, from *ἑτερος*, and *σὰρξ*, 'flesh.' A class of diseases which consist in the formation of accidental tissues.—Gendrin.

HETEROTAXIA, (F.) *Hétérotaxie*, from *ἑτερος*, and *τάξις*, 'order.' A malformation, which consists in the general transposition of organs. A change in the relation of organs.

HETEROTOPIA, (F.) *Hétérotopie*, from *ἑτερος*, and *τοπος*, 'place.' A deviation from the natural position of parts.

HÊTRE. *Fagus sylvatica*.

HEUCHERA, see *H. cortusa*—*h. Acerifolia*, *H. cortusa*.

HEUCHERA CORTUSA, *H. America'na* seu *Acerifolia* seu *Vis'cida*, *Cortu'sa America'na*, *Alum Root*, *American San'icle*, *Ground-maple*, *Cliff-weed*, *Split-rock*. Ord. Saxifragaceæ. The root—*Heuchera* (Ph. U. S.)—is a powerful astringent, and is the basis of a *cancer powder*. The American Indians apply the powdered root to wounds, ulcers, and cancers. It is said to have been sold for colchicum.

HEUCHERA RICHARDSONII, which abounds on the rocks of Churchill River, is used medicinally by the Cree Indians, who chew its astringent root, and apply it as a vulnerary to wounds and sores.

HEUCHERA VISCIDA, *H. cortusa*.

HEUCK-BANE, *Ischion*.

HEUDELOTIA AFRICANA, see *Bdellium*.

HEUKS, *Singulus*.

HEVEA GUIANENSIS, see *Caoutchouc*.

HEXAGIUM, *Sextula*.

HEXAGONE ARTÉRIELLE, see *Circulus Willisii*.

HEXATHYRIDIVM VENARUM, see *Worms*.

HEXIS, *ἔξις*. Habit, habit of body, constitution. Hence, *hectic*, *cachectic*, &c.

HEYLE, *Sanitas*.

HIA TSAO TONG TCHONG, *Sphæria Sinensis*.

HIA'TUS, from *hiare*, *hiatum*, 'to gape,' 'to open.' A foramen or aperture. Mouth. The vulva. Also, yawning.

HIATUS CANALIS FALLOPII, see *Aquæductus Fallopii*.

HIATUS DIAPHRAGMATICUS AÖR'TICUS, *Semicirculus exculp'tus*. The opening in the diaphragm for the passage of the abdominal aorta.

HIATUS FALLOPII, see *Aquæductus Fallopii*—*h. Occipito-pétræus*, *Lacerum posterius foramen*—*h. Sphæno-pétræus*, *Lacerum anterius foramen*.

HIATUS OF WINSLOW, *Foramen Winslowii*, *Foramen of Winslow*. An opening—situate behind the lesser omentum, and behind the vessels and nerves of the liver—which forms a communication between the peritoneal cavity and that of the omenta.

HIBERNICUS LAPIS, *Teg'ula* seu *Arde'sia Hibernica*, *Harde'sia*, *Irish Slate*. A kind of slate or very hard stone, found in different parts of Ireland, in masses of a bluish-black colour, which stains the hands. It has been taken, powdered, in spruce-beer, against inward constitutions.

HIBISCUS, *Althæa*.

HIBISCUS ABELMOSCHUS, *Abelmoschus moschatus*. Ord. Malvaceæ. The name of the plant, whose seeds are called *Grana Moschi* or

Muskseed. It is the *Belmoschus*, *Abelmoschus*, *Granum Moschi*, *Ket'mia Egyptiaca*, *Moschus Arabum*, *Egyptia moschata*, *Bamia moschata*, *Al'cea*, *A. Ind'ica* seu *Egyptiaca Villo'sa*, *Abelmosch*, *Abelmosk*, *Muskmalloe*, (F.) *Graine de Mue*, *Herbe à la poudre de Chypre*, *Ambrette*, *Guimauve veloutée*. It is indigenous in Egypt and the Indies. The seeds are chiefly used as perfumes; and especially in the formation of *Cyprus Powder*.

HIBISCUS ESCULENTUS, *Okra*, *Okro*, *Combo*. A plant, the green pods of which abound in mucilage, and are much used in the West Indies for soups.

HIBISCUS PUPLEUS, *Balimba'go*. A small Molucca tree. The fruit is full of a juice similar to camboge. The root is emetic. It is used in chronic diarrhœa, colic, dyspepsia, &c.

HIBRIDA, *Hybrid*.

HICCOUGH, *Singulus*.

HICCUP, *Singulus*.

HICK'ORY. The name of several American trees of the genus *Carya*; Order, *Juglandaceæ*. The leaves are usually aromatic; and are reputed to be antispasmodic [?]. The bark of those species that have bitter nuts, as *Carya amara* and *C. porc'ina*, is somewhat astringent. Some of them bear fruit that is much esteemed, as *Carya oliviformis*, *Pecan'* or *Peccan'* nut, and *C. vulcata*, *Shellbark*.

HIDDEN SEIZURES. An expression employed by Dr. Marshall Hall for obscure encephalic and spinal attacks, of an epileptoid character for example, which may be immediately owing to trachelismus.

HIDE-BOUND, Induration of the cellular tissue.

HIDRISCHEISIS, *Hidroschesis*.

HIDROA, *Desudatio*, *Echaubouures*, *Hydron*, *Sudamina*.

HIDRON'OSUS, *Hidronus'us*, from *ἰδρως*, 'sweat,' and *νοσος*, 'a disease.' A disease accompanied by violent sweats. *Sudor Anglicus*.

HIDRONUSUS, *Hidronosus*.

HIDROPEDESIS, *Ephidrosis*.

HIDROPHOROS, *Sudoriferous*.

HIDROPOETICUM, *Sudorific*.

HIDROPYRA, *Sudor Anglicus*.

HIDROPYRETOS, *Sudor Anglicus*.

HIDRORRHŒA, *Ephidrosis*.

HIDROS, *ἰδρως*, *ἰδρωτος*, *Sudor*, 'sweat.' Hence—*Hidropyra*, *Hidrosis*, &c.

HIDROSCHESIS, *Hidris'chesis*, *Reten'tio sudoris*, from *hidros*, and *σχέσις*, 'retention.' Suppression of perspiration.

HIDROSIS, *Ephidrosis*.

HIDROTERIUM, *Acicolum*.

HIDROTERIUM, *Sudorific*.

HIDROTIUM, *Sudorific*.

HIDRO'TIUM, Diminutive of *hidros*. A gentle sweat or perspiration.

HIDROTOPEUM, *Sudorific*.

HIEBLE, *Sambucus ebulus*.

HIERA DIACOLOCYNTHIDOS, from *ἱερος*, 'holy.' *Hiera of Colocynth*. An electuary, composed of 10 parts of *colocynth*, as much *agarie*, *germander*, *white horehound*, *stechas*:—5 parts of *opoponax*, as much *sagapenum*, *parsley*, *round birthwort root*, and *white pepper*:—4 parts of *spikenard*, *cinnamon*, *myrrh*, and *saffron*; and 3 pounds, 3 ounces, and 5 drachms of *honey*.

HIERA OF COLOCYNTH, *H. Diacolocynthidos*—*h. Logadii*, *Hiera picra*.

HIERA PICRA, from *ἱερος*, 'holy,' and *πικρος*, 'bitter.' *Holy bitter*, *Pulvis aloë'icus*, formerly called *Hiera logadii* when made into an electuary with honey. It is now kept in the form of dry powder;—prepared by mixing *socotrine aloes*,

one pound, with 3 ounces of *canella alba*. See *Pulvis aloes cum canella*.

HIERA SYRINX, Vertebral column, Epilepsy.

HIERACITES, *ἱεράκεις*, from *ἱερά*, 'precious,' and *αἶμα*, 'blood.' The ancient name of a precious stone, believed to be capable of arresting the hemorrhoidal flux.—Pliny, Galen, Paulus.

HIERACIUM LACHENALII, *H. murorum*.

HIERACIUM MURORUM, *H. Lachenalii*, *Polmonaria Gallica*, *Auricula muris major*, (F.) *Épave des murailles*, *Palmatoire des Français*. Ord. Compositæ. A European plant, which is a slight tonic.

HIERACIUM OLERACEUM, *Sonchus oleraceus*.

HIERACIUM PILOSELLA, The systematic name of the *Auricula Muris*, *Pilosella*, *P. Alpina*, *Mousse-ear*, (F.) *Piloselle*, *Oreille de Souris*. This plant contains a bitter, lactescent juice, which has a slight degree of astringency. The roots are more powerful than the leaves.

HIERACIUM VENSOSUM, *Rattlesnake-weed*, *Tring Hackweed*, *Bloodwort*, indigenous, has similar properties.

HIERANOSUS, Chorea, Convulsion, Epilepsy.

HIERAX, Accipiter.

HIEROBOTANE, *Verbena officinalis*.

HIEROGLYPHICA, from *ἱερός*, 'holy,' and *γλύφω*, 'I carve.' A name given to the signs employed in medicine; and, also, to the folds in the hands, feet, and forehead, which afford chiromancy its pretended oracles.

HIEROPYR, Erysipelas.

HILIX PIKRY, *Pulvis aloes et Canellæ*.

HILLS BALSAM OF HONEY, see *Balsam of Honey*, *Hill's*—h. Essence of Bardana, *Tinctura gaultheriæ*.

HILLBERRY, *Gaultheria*.

HILL FEVER, see *Fever*, *hill*.

HILON (F.), *Hilon*, improperly *Hilus*, 'the black spot on the top of a bean.' A name given by some writers to a small blackish tumour, formed by the protrusion of the iris through an opening in the transparent cornea, so called from its comparison with the *hile* or black mark presented by the *vicia faba* at one of its extremities. In Botany, the *Hile* or *Hilon* is the cicatrícula of a seed, which indicates the place by which it was attached in the cavity of the pericarp. It is the umbilicus of the seed. The fissure of the spleen, kidney, lung, ovary, &c., is, also, called *Hilus*. See *Porta*.

HILUS, *Hilon*.

HILUS LIENALIS, *Incisura lienalis*. The concave part of the spleen.

HILUS RENALIS, *Incisura renal*, *Hilus* of the kidney. The concave part of the kidney.

HILWORT, *Mentha pulegium*.

HIMANTOMA, see *Himas*.

HIMANTOSIS, *Himas*.

HIMAS, *ἱμας*, *ἱμάντος*, 'a thong of leather.' The uvula; likewise, elongation, and extenuation of the uvula. It is also called *Himantosis*, *Himantoma*. See *Staphylœdema*.

HIMEROS, Libido.

HINDBERRIES, see *Rubus idæus*.

HINDHEEL, *Tanacetum*.

HIP, Haunch, *Hypochondriasis*—h. Bone, Ischium—h. Disease, *Coxarum morbus*—h. Joint, *Coxofemoral articulation*—h. Tree, *Rosa canina*.

HIPPAË, *ἵππα*, from *ἵππος*, 'a horse.' *Cheese Equinus*. A cheese prepared from mare's milk.

HIPPANTHROPICIA, from *ἵππος*, 'a horse,' and *άνθρωπος*, 'a man.' A variety of melancholy, in which the patient believes himself changed to a horse. The Greek word *ἵππανθρωπος*, means the fabulous Centaur.

HIPPASIA, Equitation.

HIPEED, *Hypochondriac*.

HIPPEIA, Equitation.

HIPPEUSIS, Equitation.

HIPPIA'TER, *Hippiatros*, *Medicus equarius*, from *ἵππος*, 'a horse,' and *αἶμα*, 'a physician.' A farrier. A horse doctor. Used also for one who treats the diseases of other domestic animals; *Veterinarius*, *Malomedicus*, *Medicus Veterinarius*, *Zoëtros*.

HIPPIATRIA, *Hippiatrica*, *Hippiatrics*, *Medicina equaria*, (F.) *Hippiatrique*. A science, whose object is the knowledge of the diseases of the horse. It is sometimes made to include other domestic animals. See *Veterinary art*.

HIPPIATRQUE, *Hippiatria*.

HIPPIATRUS, *Hippiater*.

HIPPO, *Ipecacuanha*. A corruption, in Ireland, of *Ipecacuanha*; and hence applied, at times, in the United States, to the American plant, *Euphorbia Corollata*, which is, also, called *Ipecacuanha*.

HIPPO, *INDIAN*, *Gillenia trifoliata*.

HIPPOCAMPE GRANDE, *Cornu ammonis*.

HIPPOCAMPUS MAJOR, *Cornu ammonis*.

HIPPOCAMPUS MINOR, *Pes hippocampi minor*, (*ἵπποκαμπος*, 'the sea horse.') *Unguis*, *U. Acis* seu *Halleri*, *O'crea*, *Colliculus cavæ posterioris ventriculi cum laterali lino*, *Calcar acis*, *Utriciform Em'luence*, (F.) *Ergot*, *Éperon*. A medullary tubercle or projection, observed in the posterior cornu of the lateral ventricle of the brain.

HIPPOCENTAUREA CENTAURIUM, *Chironia centaureum*.

HIPPOCRAS, *Claret*.

HIPPOCRATES, *CAP OF*, *Bonnet d'Hippocrate*—h. Sleeve, *Chausse*.

HIPPOCRATIC, *Hippocraticus*. Relating to Hippocrates, or concerning his doctrine,—as *Hippocratic doctrine*, *Hippocratic juice*, &c.

HIPPOCRATISM, *Hippocratismus*. The doctrine of Hippocrates, which consisted in observing and following nature, as indicated by her presumed efforts, and the crises set up by her.

HIPPOCRATIST. A partisan of the Hippocratic doctrine.

HIPPOGONYOLEPUS, *Crusta genu equinae*.

HIPPOLAPATHIUM, *Rumex patientia*.

HIPPOLITHUS, Bezoard of the horse.

HIPPOMANÉ MANCINELLA, (*ἵππος*, 'a horse,' and *mania*, 'madness,' 'horse, or violent madness,') *Manchineel*, (F.) *Mancevillier*. Family, *Euphorbiaceæ*. *Sex. Syst.* Monœcia Monadelphica. A poisonous tree of South America, with the milky juice of which the Indians poison their arrows. The fruit, also, is highly poisonous.

HIPPOMARATHRUM, *Peucedanum silaus*.

HIPPOPATHOLOG'Y, *Hippopathologia*, from *ἵππος*, 'a horse,' *πάθος*, 'a disease,' and *λογία*, 'a discourse.' The science of the diseases of the horse. Pathology of the horse.

HIPPOPHAGOUS, Equivorous.

HIPPOPUS, see *Kyllosis*.

HIPPOS, *Equus*.

HIPPOSELINUM, *Smyrniolum olusatrum*.

HIPPOSTEOLOGY, *Hipposteologia*, from *ἵππος*, 'a horse,' *οστέον*, 'a bone,' and *λογία*, 'a discourse.' Osteology of the horse.

HIPPOTOMY, *Hippotomia*, from *ἵππος*, 'a horse,' and *τεμννν*, 'to cut.' Anatomy of the horse.

HIPPURIA, from *ἵππος*, 'a horse,' and *ουρον*, 'urine,' because the urine contains hippuric acid, which is found in the urine of the horse. A pathological condition, in which there is an excess of hippuric acid in the urine.

HIPPURIC ACID, see *Acid*, *hippuric*.

HIPPURIS VULGARIS, from *ἵππος*, 'a

horse,' and *cupa*, 'a tail.' The systematic name of the *Horse's Tail*, *Equisetum minus*, *Equisetum*, (F.) *Prêle*, *Presle*, *Asprêle*, *Queue de cheval*. It is an astringent, and frequently used, as tea, by the vulgar, in diarrhœa and hemorrhage. The same virtues are attributed to the *Equisetum arvense*, *fluviatile*, *limosum*, &c.

HIPPUS, *Equus*, *Nictatio*, from 'ἵππος, 'a horse.' A disease of the eyes, in which, from birth, they perpetually twinkle, like those of a man on horseback. [?] Also, a tremulous condition of the iris, which occasions repeated alternations of contraction and dilatation of the pupil; *Iridodromus*, *Iridodone'sis*, *Iridoplasia*, *Tremor Pridis*.

HIPS, Sax. *heopa*. The fruit of the dog-rose, *Rosa canina*: chiefly used as a confection. See *Confectio Rosæ Caninæ*.

HIRCIS/MUS, *Hircus*, *Hirquus*, from *hircus*, 'a goat.' Stinking like a goat:—applied especially to the odour of the secretions of the axilla.

HIRCUS, *Canthus* (greater), *Hircismus*, *Tragus*—h. *Alarum*, *Cinabra*.

HIRPLING, *Claudication*.

HIRQUITAL/TAS, from *hircus*, 'a goat,' *Paraphonia puerum*. Goat's voice. See *Egophony*.

HIRQUUS, *Hircus*, *Canthus* (greater), *Hircismus*, *Tragus*.

HIRSUTIES, *Dasya*, *Dasytes*, *Trichosis*, *Hirsuties*, *Hairiness*. Growth of hairs on extraneous parts, or, superfluous growth on parts; as in cases of bearded women.—Good. When congenital, it is termed *Hirsuties adnata*.

HIRUDO, *The Leech*, *Sanguisuga*, *Bdella*: from *haurio*, 'I draw out,' or from *hæreo*, 'I stick.' [?] In medicine, *Hirudo Medicinalis*, *Bdella Medicinalis* or *Medicinal Leech*, (F.) *Sanguis*, is employed. In the United States, *H. decora* is used. The leech lives in fresh water, and feeds on the blood of animals, which it sucks, after having pierced the skin with its three sharp teeth. This habit has been taken advantage of, to produce local blood-letting. In applying the leech, the part must be wiped dry; and if there be difficulty in making it suck, a little milk or cream may be applied. When satiated, it will drop off, and by applying a little salt or vinegar to its head it will disgorge the blood. A good English leech will take about half an ounce of blood, including that which flows by fomenting the part subsequently. The American takes less.

HIRUDO ARTIFICIALIS, *Antlia sanguisuga*—h. *Decora*, see *Hirudo*—h. *Medicinalis*, *Hirudo*.

HIRUNDINARIA, *Asclepias vincetoxicum*, *Lysimachia nummularia*.

HIRUN'DO, *Chelidon*, from *hærendo*, 'sticking,' because it sticks its nests against the houses [?]. *The Swallow*. The nests of the swallow were once employed as rubefacients, boiled in vinegar.

HISPANICUM VIRIDE, *Cupri subacetatis*.

HISPIDITAS, *Dystecchiasis*, *Phalangosis*.

HISPIDULA, *Antennaria dioicum*.

HISSING RESPIRATION, see *Râle sibilant*.

HISTIOLOGICAL, *Histological*.

HISTIOLOGY, *Histology*.

HISTOCHEMIA, see *Chymistry*.

HISTOCHEMY, see *Chymistry*.

HISTOCHEMISTRY, see *Chymistry*.

HISTODIALYSIS, from 'ἵστος, 'organic texture,' and *διαλυσις*, 'dissolution.' A morbid dissolution of the tissues.

HISTOGENETIC, *Histogeneticus*, 'Tissue making.' Same etymon as the next. Relating, or appertaining to *histogeny* or the formation and development of the organic textures.

HISTOGENETIC ELEMENTS OF NUTRITION, see *Plastic Element*.

HISTOGENY, *Histogenia*, *Histogenesis*,

Tissue formation, from 'ἵστος, 'the organic texture,' and *γενεσις*, 'generation.' The formation and development of the organic textures.

HISTOLOGIA, *Histology*.

HISTOLOGICAL, *Histologicus*, *Histologicus*, *Histologicus*. Same etymon as the next. Relating to histology. Applied, also, at times, to the natural transformations that occur in the tissues in the embryo, in contradistinction to *morphological*, which applies to the alterations in the form of the several parts of the embryo.

HISTOL'OGY, *Histolog'ia*, *Histology*, *Histolog'ia*, from 'ἵστος, 'the organic texture,' and *λογος*, 'a description.' Anatomy (general). The term is, also, more particularly appropriated to the minute anatomy of the tissues. See *Anatomy*.

Patholog'ical Histology, *Microscopic patholog'ical Anat'omy*, is the morbid anatomy of the tissues.

HISTOLYSIS, from 'ἵστος, 'the organic texture,' and *λυσις*, 'solution.' The decay and dissolution of the organic tissues, and of the blood. It includes the various forms of retrograde metamorphosis and degeneration.

HISTON'OMY, *Histonomia*, from 'ἵστος, 'the organic texture,' and *νομος*, 'law.' The aggregate of laws, which preside over the formation and arrangement of the organic tissues.

HISTORY, MEDICAL, *Historia Medicinæ*. A narration of the chief circumstances, and the persons connected with them, in the progress of medicine.

HISTOS, 'ἵστος, 'a web or tissue,' 'the organic texture'; *Textura seu Tela organica*.

HISTOT'OMY, *Histotomia*, from 'ἵστος, 'organic texture,' and *τομή*, 'incision.' Dissection of the tissues.

HIVE, *Hives*.

HIVE SYRUP, *Syrupus scillæ compositus*.

HIVES, *Cynanche trachealis*, *Urticaria*, *Variella*. In Scotland, according to Dr. Jameson, *Hives* or *Hives* means any eruption of the skin, proceeding from an internal cause; and, in Lothian, it is used to denote both the red and the yellow gum. In the United States, it is vaguely employed: most frequently, perhaps, for *Urticaria*.

HIVES, *BOLD*, *Cynanche trachealis*, *Urticaria*—h. *Eating*, *Rupia escharotica*.

HOARSENESS, *Raucedo*.

HOAST, *Tussis*.

HOBBIL, *Idiot*.

HOB-NAIL LIVER, *Cirrhosis of the liver*, *Liver*, *nutmeg*.

HOC, *Alcea rosea*.

HOCK, *Poples*.

HOCKS, *Alcea rosea*.

HOD, *Crick in the neck*.

HOFF, *Poples*.

HOG-LICE, *Onisci aselli*.

HOGWORT, *Heptallon graveolens*.

HOIST, *Tussis*.

HOLANENCEPHALIA, from 'ολος, 'entire,' and *anencephalia*, 'absence of brain.' Entire absence of brain,—the same as *Anencephalia*.—I. G. St. Hilaire.

HOLARTHRTIS, *Hamarthritis*.

HOLCE, 'ολκη. A drachm.—Galen.

HOL'CIMOS, 'ολκιμος, from 'ολκη, 'a weight.'

A tumour of the liver.

HOLCUS SORGHUM, *Panicum Italicum*.

HOLENE, *Ilex aquifolium*.

HOLERA, *Cholera*.

HOLES, BURN'T, *Rupia escharotica*.

HOLLANDS, *Gin*.

HOLLEN, *Ilex aquifolium*.

HOLLY, AMERICAN, *Ilex opaca*—h. *Common*, *Ilex aquifolium*—h. *Dahcon*, *Ilex vomitoris*

—h. Ground, *Pyrola maculata*, *Pyrola umbellata*.

—h. Sea, *Eryngium maritimum*.

HOLLYHOCK, COMMON, *Alcea rosea*.

HOLMES WEED, *Scrophularia nodosa*.

HOLMICOS, Alveolus.

HOLMOS, Mortar.

HOLOCYRON, *Teucrium chamæpitys*.

HOLOGANGLITIS, Cholera.

HOLONARCO'SIS, from *ὅλος*, 'whole,' and *νάρκωσις*, 'stupor.' Narcosis of the whole body. *Torpiditas universalis*.

HOLOPHYCTIDES, Phlyctæna.

HIOLOSTEUM ALSINE, *Alsine media*.

HIOLOSTEUS, *Osteocolla*.

HIOLOTETANUS, see Tetanus.

HIOLOTONIA, *Holotoniceus*, Tetanus.

HIOLOTON'ICUS, *ὅλος*, 'the whole,' and *τεινω*, 'I stretch.' *Holoton'ia*. A spasm of the whole body:—a variety of tetanus.—Sauvages.

HOLYN, *Ilex aquifolium*.

HOLYWELL, MINERAL WATERS OF. Holywell is a town in Wales, and takes its name from the famous well of St. Winifred. It is a simple cold water, remarkable for its purity.

HOMAGRA, Omagra.

HOMEOPATHY, *Homœopathy*.

HOMERDA, *Stercus humanum*.

HOMERIA COLLI'NA, *Cape Tulip*. A poisonous South African plant, *Nat. Ord.* Iridææ, the bulb of which acts as a violent acro-narcotic, producing fatal results very speedily.

HOMESICKNESS, Nostalgia.

HOMICID'AL, *Truculentus*, *Ferus*, from *homo*, 'man,' and *cædo*, 'I kill.' Pertaining or relating to homicide or the killing of man.

HOMICIDAL INSAN'ITY, *Homicid'ali Monomani'a*, (F.) *Monomanië homicide*. Insanity, with an irresistible impulse to destroy life.

HOMILIA, Coition.

HOM'INY. A word of Indian derivation. Maize or Indian corn hulled and coarsely broken. It is prepared for food by being mixed with water and boiled.

HOMINY, WHEATEN, see Groats.

HOMIOSIS, Homiosis.

HOMME, Homo.

HOMO, (F.) *Homme*, Man,—the chief and most perfect of the mammalia; in Greek, *ανθρωπος*, *Anthro'pos*, from *ana*, 'upwards,' and *τερω*, 'I turn,' because man, alone, of all animals, possesses the natural power of standing erect. He is, also, the only animal whose incisor teeth, wedged in a projecting jaw, are absolutely vertical. Man is especially distinguished from other mammalia by the faculty, which he possesses, of classing his ideas; comparing them with each other; and connecting, representing, and transmitting them by signs and articulate sounds. He possesses, in the highest degree, all the attributes of intelligence,—memory, judgment, and imagination. He inhabits all countries,—the burning regions of the torrid zone, and the chilling atmosphere of the polar climes. In different situations, he presents, in his figure, colour, and stature, differences which have caused mankind to be divided by naturalists into races or varieties. The number of such races can only be approximated. Blumenbach admits five, the *Caucasian*, *Ethiopian*, *Mongolian*, *Malay*, and *American*. Every division must necessarily be arbitrary, and the individuals composing each variety are far from being alike.

HOMO ALATUS, see Alatus.

HOMO CAUDA'TUS, 'Tailed man.' A fabulous, tailed variety of the human species, 'incola orbis antarctici,' admitted by Linnæus, although he is uncertain whether to rank them with men or apes!

HOMO FATUUS, Idiot.

HOMOED'RUS, from *ἄμα*, 'together,' and *ἔδρα*, 'seat.' Having the same seat. *Morbi homœdri*:—diseases that have the same seat.

HOMŒOMORPHOUS, see Homology.

HOMŒOPATHI, *Homœopathist*.

HOMŒOPATHI'IC, *Homœopath'icus*, *Homœo-*

opathes. Relating to homœopathy.

HOMŒOPATHIE, *Homœopathy*.

HOMŒOPATHIST, *Homœopath*, *Homœopath'icus*, *Homœopathis'ta*, *Homœopathis'tes*. One who believes in homœopathy.

HOMŒOP'ATHY, *Homœopath'i'a*, *Arx homœopath'ica*, *Homœop'athy*, *Hahnemannism*, *Ugma-mop'athy*, (F.) *Homœopathie*, *Homœothérapeutique*, *Homœosympathie*, *Doctrine homœo-organique*, from *ὅμοιος*, 'like,' and *πάθος*, 'affection.' A fanciful doctrine, which maintains, that disordered actions in the human body are to be cured by inducing other disordered actions of a like kind, and this to be accomplished by infinitesimally small doses, often of apparently inert agents; the decillionth part of a grain of charcoal, for example, is an authorized dose. It has also been called *Globulism*.

HOMŒOSIS, Homiosis.

HOMŒOSYMPATHIE, *Homœopathy*.

HOMŒOTHÉRAPEUTIQUE, *Homœo-*

opathy. HOMŒOZ'YGY, from *ὅμοιος*, 'like,' and *ζυγω*, 'I join together.' (F.) *Sot-pour-soi*. The law of association of organs, by which like parts adhere to like parts.—Serres.

HOMOETHNIA, Sympathy.

HOMOGENESIS, see Generation.

HOMOPATHIA, Sympathy.

HOMOIOS'IS, *Homœo'sis*, *Homio'sis*, from *ὅμοιος*, 'I resemble,' 'I assimilate.' An elaboration of the nutritious juice, by which it becomes proper for assimilation.

HOMOLINON, see Apolinos.

HOMOLOGOUS, see Homology.

HOMOLOGUE, *Homol'ogue*, same etymon as homology. A term applied to the same organ in different animals under every variety of form and function.—Owen.

HOMOL'OGY, from *ὅμοιος*, 'like,' and *λογος*, 'a description.' The doctrine of similarity of structure. Thus, the two sides of the body are said to be 'homologous.' *Homol'ogous*, *homomorph'ous*, or *homœomorph'ous* tissues are those that resemble others; in opposition to *heterol'ogous*, *heterocritic*, or *heteromorph'ous*, which are new formations. Homology seems now to be accepted as the designation of the doctrine or study, the subject of which is the relations of the parts of animal bodies.—Owen.

HOMOMORPHOUS, see Homology.

HOMONOPAGIA, Cephalalgia.

HOMOPH'AGUS, from *ὥμος*, 'raw,' and *φαγω*, 'I eat.' One who eats raw flesh.

HOMOPLAS'TY, *Homœopla'sia*, from *ὅμοιος*, 'like,' and *πλασσω*, 'I form.' The formation of homologous tissues.

HOMOPLATA, Scapula.

HOMOT'ONOS, *Equa'tis*, from *ὅμος*, 'equal,' and *τονος*, 'tone.' That which has the same tone. A continued fever, whose symptoms have an equal intensity during the whole course of the disease, has been so called. See *Acnasticos*, and *Synocha*.

HONESTY, *Lunaria rediviva*.

HONEWORT, FIELD, *Sison amomum*.

HONEY, Mel—h. Balsam of, Hill's, see Mel—h. Bloom, *Apocynum androsæmifolium*—h. of Borax, Mel boracis—h. Clarified, Mel despumatum—h. Prepared, Mel præparatum—h. of Roses, Mel rosæ—h. of Squill, compound, *Syrupus scillæ compositus*.

HONEYCOMB BAG, *Reticulum*.

HONEYSUCKLE, *Lonicera periclymenum* — h. Bush, *Diervilla trifida* — h. Scarlet, *Lonicera sempervirens* — h. Trumpet, *Lonicera sempervirens*.

HONEY SUGAR, Glucose.

HONGLANE, *Coptis teeta*.

HONOR CAPITIS, see *Capillus*.

HONORAIRE, *Sostrum*.

HONORARIUM, *Sostrum*.

HONTEUX, *Padie*.

HOODWORT, *Scutellaria lateriflora*.

HOOK, Sax. *hoco*, *hooce*, Dutch, *hoeck*, *Un-*

cus, *Unc'ius*, *Anc'gra*, (F.) *Érigne*, *Airigne* ou

Érine. An instrument, consisting of a steel wire,

flattened at the middle, and having the extremi-

ties crooked and pointed. Some hooks are fur-

nished with a handle at one extremity—the other

having one or two hooks;—constituting the *single*

or *double hook*. The hook is used by anatomists

and surgeons to lay hold of, and raise up, certain

parts, the dissection of which is delicate, or which

would slip from the fingers.

The *Tenaculum* is a variety of the hook.

HOOK, BLUNT, *Ich'thya*, *Ich'thye*. *Ongis*, *Unguis*

ferreus. An instrument which is passed over

the flexures of the joints to assist in bringing

down the fetus in parturition.

HOOP TREE, *Melia azedarach*.

HOOPER'S PILLS, *Pilule Aloes* et *Myrrhæ*.

HOOPING-COUGH, *Pertussis* — h. *Roche's*

Embrocation, see *Roche*.

HÔPITAL, *Hospital* — h. *des Fous*, *Morotro-*

phium.

HOPLOCHRIS'MA, from *ὅπλον*, 'a weapon,'

and *χρῆμα*, 'salve.' *Unguentum armarium*,

Armatory Unguent. A salve which was sup-

posed to cure wounds by sympathy,—the instru-

ment with which the wound was inflicted being

anointed with it.

HOPLOMOCIL'ION, *ὅπλομοχλῖον*, from *ὅπλον*,

'a weapon,' and *μοχλός*, 'a lever.' The name of

an iron machine or apparatus, which embraced

the whole body like armour. A figure of it is

given by Fabricius ab Aquapendente.

HOP, *Humulus lupulus* — h. *Pillow*, see *Humu-*

lus lupulus — h. *Plant*, *Humulus lupulus*.

HOPS, see *Humulus lupulus*.

HOQU'ET, *Singultus*.

HORA, *Hebe*.

HORÆA, *Menses*.

HORÆOTES, *Maturity*.

HORDEI MALTUM, *Malt*.

HORDE'OLUM, diminutive of *Hordeum*, 'bar-

ley.' *Orde'olum*, *Pos'thia*, *Sclerophthal'mia*, *Cri-*

thæ, *Crieth'id'ion*, *Crythæ*, *Stye*, *Styan*, *Stine*, (F.)

Orgelet, *Orgeolet*. A small, inflammatory tu-

mour, of the nature of a boil, which exhibits

itself near the free edge of the eyelids, particu-

larly near the inner angle of the eye.

HOR'DEUM, *Or'deum*, *Crithæ*. The seeds of

Hordeum vulgare or *Scotch Barley*, *Ord*. *Gra-*

mineæ, are ranked amongst the *Cerealia*. (F.)

Orge. They afford a mucilaginous decoction,

which is employed as a diluent and antiphlo-

gistic. The seeds of the *Hordeum distichon* seu

antivum, *Zoe'riton dietichum*, and *H. hexas-*

tichon possess similar properties. Barley is freed

from its shells in mills, forming the *Hor'deum*

mundatum seu *decortica'tum* seu *excortica'tum*,

Hor'deum of the *Pharmacopœias*; and, at times,

is rubbed into small, round grains, somewhat

like pearls, when it is called *Hordeum perlatum*,

Pearl Barley, (F.) *Orge perlé*, and forms the

Hordeum denudatum seu *perlatum*, the *Hordei*

sem'ina tubicis munda'ta, of the *Pharmacopœias*,

—*Hordeum* (Ph. U. S.).

HORDEUM CAUSTICUM, *Veratrum sabadilla* — h.

Decortica'tum, see *Hordeum* — h. *Denudatum*, see

Hordeum — h. *Excortica'tum*, see *Hordeum* — h.

Galacticum, *Oryza* — h. *Mundatum*, see *Hordeum*

— h. *Perlatum*, see *Hordeum*.

HOREHOUND, *Marrubium* — h. *Black*, *Bal-*

lota fetida — h. *Germander-leaved*, *Eupatorium*

teucrifolium — h. *Stinking*, *Ballota fetida* — h.

Water, *Lycopus sinuatus*, *Lycopus Virginicus* —

h. Wild, *Eupatorium rotundifolium*, *E. teucri-*

folium.

HORME, *Instinct*.

HORMINUM, *Salvia horminum* — h. *Colora-*

tum, *Salvia horminum* — h. *Sativum*, *Salvia hor-*

minum.

HORMON, *Enormon*.

HORN, *Cornu*.

HORN POX, see *Varicella*.

HORNSEED, *Ergot*.

HORN, *Corneus*.

HORN EXCRESCENCES, *Lepido'sis*, *Ichthy'i'a-*

sis cornig'era, *Cor'nea cuta'nea*, (F.) *Cornes* de

la Peau. Certain excrescences, which occasion-

ally form on some part of the skin, and resemble,

in shape, the horns of an animal.

HORN SUBSTANCE, see *Tooth*.

HORRENTIA, *Horripilation*.

HORRIDA CUTIS, *Goose-skin*, *Cutis anse-*

ri'na, *Dermatospasmus*, *Spasmus periph'ericus*.

A state of the skin produced by cold or fear, and

accompanying the rigor of an intermittent.

HORRIPILA'TION, *Horripila'tio*, *Horror*,

Horren'tia, *Phric'e*, *Phric'us*, *Phric'us*, *Phric'us*,

Phri'cia, from *horreo*, 'to bristle up,' and *pilus*,

'hair.' (F.) *Horripilation*, *Frisonnement*. Ge-

neral chilliness, preceding fever, and accompanied

with bristling of the hairs over the body.

HORROR, *Horripilation* — h. *Ciborum*, *Disgust*

for food.

HORSE BALM, *Collinsonia Canadensis* — h.

Cane, *Ambrosia trifida* — h. *Chestnut*, *Aesculus*

hippocastanum — h. *Crust*, *Crusta genu equinae*.

HORSEFLY WEED, *Sophora tinctoria*.

HORSEFOOT, see *Kyllosis*.

HORSEMINT, *Ambrosia trifida*, *Monarda coc-*

cinea, *M. fistulosa*, and *M. punctata* — h. *Sweet*,

Canilla mariana.

HORSE NETTLE, *Solanum Carolinense*.

HORSE RADISH, *Cochlearia armoracia*.

HORSE'S TAIL, *Hippuris vulgaris*.

HORSE WEED, *Ambrosia trifida*, *Collinsonia*

Canadensis, *Erigeron Canadense*.

HORTULUS CUPIDINIS, *Vulva*.

HORTUS, *Vulva* — h. *Siccus*, *Herbarium*.

HOSE, *Collum*, *Throat*.

HOSEFISH, *Sepia*.

HOSPITAL, primarily from *hospes*, *hospitis*,

'a guest.' *Infirmary*, (Old Eng.) *Farm'ery*,

Spit'al, *Nosocomi'um*, *Adynatoconi'um*, *Adynato-*

dochi'um, *Xenodoche'um*, *Xenodoche'um*, *Infirma-*

rium, *Infirmary'um*, *Nosodochi'um*, *Valetudi-*

na'rium, (F.) *Hôpital*. An establishment for the

reception of the sick, in which they are main-

tained and treated medically. Hospitals were

first instituted about the end of the 4th century;

a period at which the word *νοσοκομειον* was em-

ployed, for the first time, by St. Jerome. They

may be general, receiving all cases; or special,

admitting only the subjects of certain diseases.

HOSPITAL GANGRENE, *Phagedæ'na gangra-*

no'sa, *Putrid* or *Malignant Ulcer*, *Gangræ'na*

Nosocomi'um seu *Nosocomia'tis* seu *Contagio'sa*,

Sphac'elus nosocomia'tis, *Hospital Sore*, *Putre-*

do, (F.) *Pourriture ou Gangrène d'hôpital*. *Gan-*

grene, occurring in wounds or ulcers, in hos-

pitals, the air of which has been vitiated by the

accumulation of patients, or some other circum-

stance. Hospital gangrene—many different va-

rieties of which are met with, and always accom-

panied or preceded by fever—commonly commences with suppression of the suppuration of the wound, which becomes covered with a grayish and tenacious sanies. The gangrene then manifests itself. It extends from the centre of the ulcerated surface towards the edges; these become swollen, painful, and everted; and the patient dies with all the signs of typhus. The treatment must be varied according to circumstances. Sometimes, it requires the use of stimulating, acid, caustic, and antiseptic applications; with, occasionally, the actual cautery, aided by the exhibition of tonics, internally:—at others, the antiphlogistic regimen and emollient applications may be necessary.

HOSPITAL, INSANE, Morotrophium—h. Leper, *Ladrière*.

HOST, Tussis.

HOSTIARIUS, Pylorus.

HOT SPRINGS, see Virginia, mineral waters of.

HOUBLON, *Humulus lupulus*.

HOUGH, Poples, Sickly.

HOUNDS' TONGUE, *Cynoglossum*.

HOUPPE NERVEUSE, see Papilla—h. du Menton, Depressor labii inferioris, Levator labii inferioris.

HOURLASS CONTRACTION OF THE UTERUS, see *Chaton*.

HOUSELEEK, *Sempervivum tectorum*—h. Small, *Sedum*.

HOUSEMAID'S KNEE, see Knee, housemaid's.

HOUSE-PUPIL, see House-Surgeon.

HOUSE-SURGEON, *Res'ident Surgeon*. Usually a senior house-pupil or graduate, who resides in an hospital, and attends to every accident and disease, in the absence of the attending physician or surgeon. It answers, in the British hospitals, to the *Élève interne* or *Interne* of the French. The *Élève externe* or *Externe* is a less advanced pupil; from whom *Internes* are chosen. In ordinary schools, *Élève externe* means a day-scholar; whilst *Élève interne* means a boarder.

HOUX, *Ilex aquifolium*—h. *Apalachine*, *Ilex vomitoria*—h. *Petit*, *Ruscus*.

HOWDY, Midwife.

HOWDY-WIFE, Midwife.

HUACACACMU, *Datura sanguinea*.

HUACO, Guaco.

HUANU, Guano.

HUBBON, Haunch.

HUCKLEBERRY, see *Gaylussacia*, and *Vaccinium*—h. Dwarf, *Gaylussacia humosa*.

HUCKLEBONE, *Ischion*.

HUCKSHOULDERED, see *Hump*.

HUDSON'S PRESERVATIVE FOR THE TEETH AND GUMS, see *Tinctura Myrrhæ*.

HUILE, Oil—h. *d'Absinthe*, *Artemisia absinthium* (oil of)—h. *d'Acajou*, see *Anacardium occidentale*.

HUILE ACOUSTIQUE (F.), *O'leum acusticum*, Acoustic Oil. An oil for deafness, prepared of olive oil, $\frac{3}{4}$ ij; garlic, ox-gall, and bay leaves, each $\frac{3}{4}$ ij; boiled for a quarter of an hour, and strained.

HUILE D'AMANDES, *Oleum amygdalarum*—h. *d'Aneth*, see *Anethum graveolens*—h. *Animale*, *Oleum animale*—h. *Animale de Dippel*, *Oleum animale Dippelii*—h. *Animalisée par infusion*, *Oleum animalizatum per infusionem*—h. *d'Avis*, see *Pimpinella anisum*—h. *Anthelmin-tique de Chabert*, *Oleum animale empyreumaticum Chaberti*—h. *Aromatique*, *Oleum animalizatum per infusionem*—h. *d'Aurone*, *Artemisia abrotanum* (oil of)—h. *d'Cacao*, Butter of Cacao—h. *de Cade*, see *Juniperus oxycedrus*—h. *de Carvi*, *Carum* (oil)—h. *de Cédrat*, *Oleum cedrinum*—h.

de petits Chiens, *Oleum animalizatum per infusionem*—h. *de Corne de Cerf*, *Oleum animale Dippelii*—h. *d'Épurga*, *Euphorbia lathyris*—h. *de Foie de Morue*, *Oleum jecoris aselli*—h. *de Glabian*, *Petroleum*—h. *de Gérofle*, see *Eugenia caryophyllata*—h. *de Laurier*, *Unguentum laurinum*—h. *de Lin*, see *Linum usitatissimum*—h. *de Lis*, see *Lilium candidum*—h. *de Morelle*, see *Solanum*—h. *de Morue*, *Oleum jecoris aselli*—h. *de Noir*, see *Juglans cinerea*—h. *d'Éillette*, *Papaver* (oil)—h. *de Papier*, *Pyrothionide*—h. *de Pieds de Bœuf*, *Oleum bubulum*—h. *de Riein*, see *Ricinus communis*—h. *de Succin*, see *Succinum*—h. *de Turtre par défaillance*, *Liquor potassæ subcarbonatis*—h. *Verte*, Balm, green, of Metz—h. *de Vin douce*, *Oleum Etherium*.

HUILES ANIMALES, *Olea animalia*—h. *Distillées*, *Olea volatilia*—h. *Empyreumatiques*, *Olea empyreumatica*—h. *Essentielles*, *Olea volatilia*—h. *Éthérées*, *Olea volatilia*—h. *Élixes*, ou *Grasses*, *Olea fixa*—h. *Fugaces*, *Olea fugacia*—h. *Médicinales*, *Olea medicinalia*—h. *Pyrogénées*, *Olea empyreumatica*—h. *Volatiles*, *Olea volatilia*.

HUIT DE CHIFFRE. Figure of 8. A bandage of which the turns are crossed in the form of the figure 8. Such is the bandage used after bleeding from the arm.

HUITO, *Genipa oblongifolia*.

HUITRE, *Ostrea*.

HUKEBANE, *Ischion*.

HULCUS, Ulcer.

HULGIE-BACKED, see *Hump*.

HULLUPING, Vomiting.

HUM, VENOUS, *Bruit de diable*.

HUMBLE, Rectus inferior oculi.

HUMECTANTIA. From *humectare*, 'to moisten.' A name formerly given to drinks, which appeared to possess the property of augmenting the fluidity of the blood.

HUMERAL, *Humeralia*. That which belongs to, or is connected with, the arm or humerus.

HUMERAL ARTERY, *Arteria humeralis*, see *Brachial artery*.

HUMÉRO-CUBITAL, *Brachialis anterior*—h. *Sus-métacarpien*, see *Radialis*—h. *Sus-radial*, *Supinator radii longus*.

HUMERUS, *Scapula*, *Omo*, *Armus*, (F.) *Épau*, *Shoulder*, (Prov.) *Spaul*. The most elevated part of the arm. The bones, which concur in forming it, are—the scapula, head of the humerus, and the clavicle, united together by strong ligaments, and covered by numerous muscles.

HUMERUS, *Os humeri* seu *brachii* seu *adjutorium sui brachia'le*, *Brachium*, *Lacertus*, *Shoulderbone*, (Prov.) *Spaul-bone*, *Spaul-bone*, *Shullbane*, is the cylindrical, irregular bone of the arm; the upper extremity of which has a hemispherical head connected with the scapula; and two tuberosities or tubercles, a greater and a lesser, for the attachment of muscles, between which is the *Bicipital groove* or *Fossa*. At the inferior extremity may be remarked—the *inner condyle*, the *outer condyle*; the *small head*, which is articulated with the radius; the *trochlea* articulated with the ulna, &c. The humerus is developed by seven points of ossification:—one for the body; one for the head; one for the greater tuberosity; one for the trochlea; one for the epitrochlea; one for the epicondyle; and another for the lesser head.

HUMERUS SUMMUS, Acromion.

HUMEUR AQUEUSE, Aqueous humour—h. *Crystalline*, *Crystalline*—h. *Hyaloidé*, *Corpus vitreum*—h. *de Morgagni*, *Morgagni*, humour of.

HUMEURS FROIDES, *Serofula*.

HUMIDE RADICALE, *Humidum radicale*.

HUMIDUM NATIVUM, *H. radicale*—h. *Na-*

tivum Articulorum, Synovia—h. Primigenium, H. radiale.

HUMIDUM RADICALE, *Hu'midum primigenium seu nativum seu seminale*, Radical Moisture, (F.) *Humide radicale*. Names formerly given to the liquid which was conceived to give flexibility and proper consistence to the different organic textures.

HUMIDUM SEMINALE, H. radiale.

HUMILIS, Rectus inferior oculi.

HUMLOIK, Conium.

HUMOR, Humour—h. Albugineous, Aqueous humour—h. Articularis, Synovia—h. Ceruminous, Cerumen—h. Doridis, Water, sea—h. Genitalis, Sperm—h. Glacialis, Crystalline, Corpus vitreum—h. Hyalinus seu Hyaloides, Corpus vitreum—h. Lacteus, Milk—h. Lacrymalis, Tear—h. Melancholicus, see Mercurialis—h. Mercurialis, see Mercurialis—h. Morgagnianus, Morgagni, humor of—h. Ovatus, Aqueous humour—h. Oviformis, Aqueous humour—h. Pericardii, see Pericardium—h. Purulentus, Pus—h. Seminalis, Sperm—h. Venerous, Sperm—h. Vitreus, Corpus vitreum.

HUMORAL, *Humoralis*, from *humere*, 'to moisten.' Proceeding from, or connected with, the humours.

HUMORES INQUILINI, Recrementitial Secretions.

HUMORIQUE, see *Bruit humorique*.

HUMORISM, *Hu'moral Pathology*, *Pathologia humeralis*. A medical theory, founded exclusively on the parts which the humours were considered to play in the production of disease. Although traces of this system may be found in the most remote antiquity, the creation, or, at all events, the arrangement of it may be attributed to Galen, who enveloped it in metaphysical subtleties relating to the union between the elements and the four cardinal humours.

HUMORISTS. The Galenic and other physicians, who attributed all diseases to the depraved state of the humours, or to vicious juices collected in the body.

HUMOUR, *Humor*, *Hygra'sia*, *Hygredon*, *Hygum*. Every fluid substance of an organized body;—as the blood, chyle, lymph, &c. The *Humours*, *χυμοι*, *Chymi*, *Humores*, differ considerably as to number and quality in the different species of organized beings; and even in the same species, according to the state of health or disease. The ancients reduced them to four; which they called *cardinal humours*:—the blood, phlegm, yellow bile, and atrabilis or black bile. A modern classification of the humours is given under Fluid.

HUMP. Perhaps from *umbo*, 'the boss of a buckler.' *Hunch*, *Gibber*, *Gibbus*, *Gibba*, *Tuber*, (F.) *Bosse*. A prominence, formed by a deviation of the bones of the trunk. Commonly, it is formed by the spine or sternum, and is seated at the posterior or anterior part of the trunk. It may, also, be produced by deviation of the ribs or pelvis. The spine may be curved in three principal directions. 1. *Backwards*, the most common case; this the ancients called *κνυσσις*, *Cynphosis*, *Gibbositas*. 2. *Forwards*, *λοφωσις*, *Lophosis*, *Recurvatio*; and 3. *Laterally*, *σκολιωσις*, *Scoliosis*, *Obstipatio*. Most curvatures occur at a very early age, and are caused by scrofula, rickets, &c.; and, not unfrequently, they are accompanied by caries of the vertebrae. See Vertebral disease. One who is affected with a hump, or is *hump-backed*, *Humpty*, *Huckshounded*, *Shrumph-shouldered*, (Sc.) *Hugliebacked*, *Trudgebacked*, (Prov.) *Croopbacked*, is called, by the French, *bossu*.

HUMPBACED, see *Hump*.

HUMPTY, see *Hump*.

HUMULUS, *Humulus lupulus*.

HUMULUS *Lu'PULUS*, *Lu'pulus*, *L. scandens* seu *communis* seu *salicetinus*, *Humulus*, (from *humus*, 'the ground on which it creeps,') *Convolverulus perennis*, the *Hop-plant*. Nat. Ord. Urticaceae. (F.) *Houblon*, *Vigne du nord*. Its cones or strobiles, *Hu'muli strobili* (Ph. L.), *Humulus* (Ph. U. S.), have a fragrant odour; and a bitter, aromatic taste, depending on a peculiar principle, named *Lu'pulin*, extractive and essential oil, which may be extracted, equally, by water and spirit, from the dried strobiles. The hop is employed as a tonic and hypnotic, and enters into the composition of ale and beer.

The *Hop pillow*, *Pulv'nar Hu'muli*, has long been used for producing sleep.

HUNCH, *Hump*.

HUNGARICA FEBRIS, Fever, Hungary.

HUNGER, Anglo-Saxon, *hunger*, *Fames*, *Limos*, *Peine*, *Peina*, *Esu'ries*, *Jeju'nium*, *Jeju'ntas*, *Esuri'tio*, *Esuri'go*, (F.) *Faim*. The necessity for taking food. Hunger is an internal sensation, which some authors have attributed to the friction between the sides of the stomach in its empty state; others, to the compression of the nerves, when the organ is contracted; others, to the action of the gastric juice, &c. It is dictated by the wants of the system:—farther we know not. See Appetite.

HUNGER-CURE, *Limotherapeia*.

HUNGRY, *Famelicus*, *Limicus*, *Limo'des*; (Prov.) *Haw*, (vulg.) *Peekish*; same etymon. Affected with hunger.

HURA BRASILIENSIS, *Assaon*, *Assaëd*, *Ussacü*. A Brazilian tree, of the Nat. Ord. Euphorbiaceae, which, in form of the extract of the bark, is esteemed a specific in leprosy. It is, also, given in elephantiasis, and as an anthelmintic.

HURDIES, *Nates*.

HURTLEBERRY, *Vaccinium myrtillus*.

HYA-HYA, *Tabernaemontana utilis*.

HYACINTH, WILD, *Scilla esculenta*.

HYACINTHUS MUSCARI, *Bulbus vomitorius*.

HYÆNAN/CHE GLOBO'SA. An arborescent shrub of South Africa, Nat. Ord. Euphorbiaceae, the fruit of which, pounded, is used to destroy hyænas and other beasts of prey, and seems to contain strychnia.

HYALEUS, *Hyaline*.

HY'ALINE, *Hyalinus*, *Hyal'eus*, *Vit'reus*, *Vit'reous*, (F.) *Vitreux*, *vitré*. Glassy. Resembling glass.

HYALINE SUBSTANCE, *Cytoblastema*.

HYALITIS, *Inflammatio tunica hyaloïdeæ*, *Hyaloiditis*, *Vit'reo-capsulitis*, from *hyaloid*, and *itis*, denoting inflammation. Inflammation of the hyaloid membrane of the eye.

HY'ALOID, *Hyaloides*, *Hyaloi'des*, from *υαλος*, 'glass,' and *αδος*, 'resemblance.' Vitreiform; resembling glass.

HYALOID CANAL, see *Hyaloid membrane*.

HYALOID FOSSA, *Fossa Hyaloïde'a*, is a cup-like excavation in the vitreous humour, in which the crystalline is imbedded.

HY'ALOID MEMBRANE, *Tunica Hyaloïde'a* seu *Vit'rea*, *Membrana Arachnoïde'a*, is the extremely delicate membrane, which forms the exterior covering of the vitreous humour, and transmits within it prolongations, which divide it into cells. Fallopius discovered this membrane, and gave it the name *Hyaloid*. On a level with the entrance of the optic nerve into the eye, the hyaloid membrane has been described as forming, by reflection, a cylindrical canal, which pierces the vitreous humour from behind to before, as far as the posterior part of the crystalline. See Canal, hyaloid. At its anterior part is a depression—

Fossa patella'ris seu *lenticula'ris* for the reception of the crystalline lens.

HYALOIDITIS, Hyalitis.

HYALONIXIS, see Cataract.

HYALONYXIS, see Cataract.

HYALOS, Vitrum.

HYANCHE, *Cynanche tonsillaris*.

HYBOMA, *Gibbositas*.

HYBRID, *Hí'brida*, *Hy'brida*, (F.) *Hybride*, from the Greek *ὑβρις*, *ὑβριδος*, 'mongrel or mongrel.' (F.) *Métis*, *Mulet*. A being born of two different species,—as the mule. The term is applied to plants as well as to animals. The result is termed *Hybridity*. Hybrid is often, also, used to designate words which are formed from two different languages—as *uter-itis*, for inflammation of the uterus, in place of *metritis*.

HYBRIDITY, see Hybrid.

HYDARTHROSIS, *Hydrarthrus*.

HYDARTHROS, *Hydrarthrus*.

HYDAS, *Hydor*.

HYDATID, *Hy'datis*, *Bulla*, *Aqu'ula*, *Hydro'a*, *Hydrocyst'is*, *Hygrocyst'is*, *Tania hydatid'ena*, *Echinococ'cus humanus*, from *ὑδωρ*, *ὑδατος*, 'water.' This name was long given to every encysted tumour which contained an aqueous and transparent fluid. Many pathologists subsequently applied it to vesicles, softer than the tissue of membranes, more or less transparent, which are developed within organs, but without adhering to their tissues. It is by no means clear that these formations are really entozoa. They have been found in various parts of the body; sometimes in the uterus, occasioning signs nearly similar to those of pregnancy, but being sooner or later expelled. The expulsion is generally attended with more or less hemorrhage. See *Acephalocystis*.

Hydatis, *Aqu'ula*, *Phlycten'ula*, *Verru'ca Palpebrarum*, *Mil'ium*, also, meant a small, transparent tumour of the eyelids.—Galen, C. Hoffmann.

HYDATIDES CERVICIS UTERI, *Nabothi glandulæ*.

HYDATIDOCELE, *Hydatoc'elē*, from *ὑδατις*, 'hydatid,' and *κύηλη*, 'a tumour.' Osechocele containing hydatids; the *Osechocele hydatido'ea*, *Hydatidoschocele'le* of Sauvages.

HYDATIDOÏDES, *Hydatoid*.

HYDATIDOMA, from *ὑδατις*, 'hydatid.' A tumour caused by hydatids.

HYDATIDOSCHEOCELE, *Hydatidocele*.

HYDATIFORM, *Hydatiform'is*. A hybrid word from *hydatid* and *forma*, 'form.' Resembling an hydatid.

HYDATINUS, *Hydatoid*.

HYDATIS FINNA, *Cysticereus cellulose*.

HYDATIS'MUS, from *ὑδωρ*, *ὑδατος*, 'water.' The noise caused by the fluctuation of pus contained in an abscess.—Aurelian, *Poësius*.

HYDATOCELE, *Hydatidocele*, *Hydrocele*.

HYDATOCH'OLOS, from *ὑδωρ*, *ὑδατος*, 'water,' and *χολή*, 'bile.' *Aquo'eo-bil'iosus*. An epithet given to evacuated matters when mixed with water and bile.—Hippocrates, *Poësius*.

HYDATODES, *Aqueous*.

HYDATOID, *Hydatoid'is*, *A'queous*, *Aquo'us*, *Hydatoid'is*, *Hydat'ina*, *Hydatido'is*, *Hydatido'is*, from *ὑδωρ*, *ὑδατος*, 'water,' and *ειδός*, 'resemblance.' Watery. Resembling water. This name has been given to the membrane of the aqueous humour; and, also, to the aqueous humour itself. *Vinum hydatoid'is*; wine and water.

HYDATONCUS, *Anasarca*, *Edema*.

HYDATOPO'SIA, from *ὑδωρ*, *ὑδατος*, 'water,' and *ποσις*, 'drinking.' Water-drinking;—hence HYDATOPOTES. A water-drinker.

HYDATOTHERAPEIA, *Hydrosudotherapeia*.

HYDERICUS, *Hydropic*.

HYDERODES, *Hydropic*.

HYDERONCUS, *Anasarca*, *Edema*.

HYDEROS, *Anasarca*, *Hydrops*.

HYDOR, *hy'datos*, *ὑδωρ*, *ὑδατος*, and *Hydas*, *hydatos*, *ὑδας*, *ὑδατος*, 'water.' Hence:

HYDRACHNIS, see *Varicella*.

HYDRADEN, *Conglobate gland*.

HYDRADENT'ITIS, *Inflammat'io glandularum lymphaticarum*, from *Hydraden*, 'a lymphatic gland,' and *itis*, denoting inflammation. Inflammation of a lymphatic gland.

HYDRÆDUS, from *ὑδωρ*, 'water,' and *αἰδοία*, 'genital organs.' Edema of the female organs.

HYDRÆMIA, *Hydroæmia*.

HYDRAGOGA, *Hydragogues*.

HYDRAGOGIA, *Hydragogues*.

HYDRAGOGICA, *Hydragogues*.

HYDRAGOGUES, *Hydrago'ga*, *Hydrago'gin*, *Hydrago'gica*, *Hydroph'ica*, *Hydroph'ica*, *Aquid'ica*, from *ὑδωρ*, 'water,' and *αἶμα*, 'I expel.' Medicines believed to be capable of expelling serum effused into any part of the body. These are generally cathartics or diuretics.

HYDRAGOGUM BOYLEI, *Argenti nitras*.

HYDRALLANTE, *False Waters*.

HYDRALLAS, from *ὑδωρ*, 'water,' and *αλλας*, 'a sausage,'—see *Allantois*. A morbid increase in the quantity of the fluid of the allantois.

HYDRALMÆ, *Waters*, *mineral* (saline).

HYDRAMNIOS, from *ὑδωρ*, 'water,' and 'amnios.' An excessive quantity of the liquor amnii.

HYDRAN'GEA ARBORESCENS, *H. vulgari's*, *Sorbarbark*, *Wild Hydran'gea*, *Bissum*. *Ord.* *Hydrangeæ*. An indigenous plant, which flowers in July. The leaves are said to be tonic, sialagogue, cathartic, diuretic, and antilithic. [?]

HYDRANGEITIS, *Angioloecitis*.

HYDRANGIA, *Lymphatic vessels*.

HYDRANGIOGRAPHIA, *Angiohydrography*.

HYDPANGIOTOMIA, *Angiohydrotomy*.

HYDRAGYRANATRIP'SIS, *Hydragyrentrip'sis*, from *ὑδραργυρος*, 'quicksilver,' and *ατραπις*, 'rubbing in.' The rubbing in of a preparation of quicksilver:—*Hydragyrotrip'sis*.

HYDRARGYRENTRIPSIS, *Hydragyranatripsis*.

HYDRARGYRI ACETAS, *Hydrargyrus acetatus*—*h.* *Bichloridum*, *H. oxymurias*—*h.* *Bicyanidum*, *H. cyanuretum*—*h.* *Biniodidum*, *H. iodium rubrum*—*h.* *Binoxidum*, *H. oxydum rubrum*—*h.* *Bisulphuretum*, *H. sulphuretum rubrum*—*h.* *Borussias*, *H. cyanuretum*—*h.* *Bromidum*, see *Bromine*—*h.* *Calx alba*, *Hydrargyrum præcipitatum*—*h.* *Chloridum*, *H. submurias*—*h.* *Chloridum corrosivum*, *H. Oxymurias*—*h.* *Chloridum mite*, *H. submurias*.

HYDRARGYRI CYANURETUM, *H. Cyan'idum* seu *Borus'sias* seu *Bicyan'idum*, *Hydrargyrum Cyanogenat'um* seu *Hydrocyan'icum* seu *Borus'sicum* seu *Zoot'icum*, *Prussias Hydrargyri*, *Cyanuret* or *Prussiate of Mercury*, (F.) *Cyanure de Mercure*. (Ferri Ferro-cyanuret. \mathfrak{z} iv; *Hydrarg. oxid. rubr.* \mathfrak{z} ij, vel q. s.; *Aquæ destillat.* Oij. Put the ferro-cyanuret and three ounces of the oxide of mercury, previously powdered and thoroughly mixed together, into a glass vessel, and pour on two pints of the distilled water. Boil the mixture, stirring constantly; and if, at the end of half an hour, the blue colour remains, add small portions of the oxide of mercury, continuing the ebullition until the mixture becomes of a yellowish colour; then filter through paper. Wash the residue in a pint of the distilled water, and filter. Mix the solution and evaporate till a pellicle appears, and set the liquor aside, that crystals may form. To purify the crystals, subject it to re-

solution, evaporation, and crystallization.—Ph. U. S.) This preparation has been strongly recommended as a powerful antisyphilitic, and is admitted into the Parisian codex. Twelve to twenty-four grains may be dissolved in a quart of distilled water, and three or four spoonfuls of the solution be taken daily in a glass of any appropriate liquid.

HYDRARGYRI DEUTO-IODIDUM, H. Iodidum rubrum—h. Deuto-ioduretum, see Iodine—h. Deutonitras, see II. nitras—h. Dipernitras, see II. nitras—h. Hyperoxodes, Hydrargyri nitrico-oxydum.

HYDRARGYRI IODIDUM, H. Protoiodidum seu Proto-ioduretum seu Subiodidum, Hydrargyrum Iodidum flaccum seu Iodidulum, Iodidum seu Ioduretum hydrargyrosum, Protoioduretum mercurii, Iodide or Protoiodide of Mercury, (F.) Protoiodure or Protoiodure of Mercure, (Hydrarg. 5j, Iodini ʒv, Alcohol q. s. Rub the mercury and iodine together, adding sufficient alcohol to form a soft paste, and continue the trituration till the globules disappear. Dry the iodide in the dark, with a gentle heat, and keep it in a well-stopped bottle, the light excluded.—Ph. U. S.) Dose, one grain, gradually raised to three or four.

HYDRARGYRI IODIDUM CHLORIDI, Mercury, iodide of chloride of.

HYDRARGYRI IODIDUM RUBRUM, H. Biniodidum seu Deuto-iodidum seu Deuto-ioduretum seu Periodidum, Hydrargyrum Iodidum rubrum seu Biiodidum seu Periodatum, Iodidum seu Iodidum Hydrargyricum, Deuto-ioduretum mercurii, Red Iodide, Biniodide, Deutiodide and Periodide of Mercury, (F.) Deutiodure or Biiodure or Periodure of Mercure, (Hydrarg. corros. chlorid. 5j, Potassii Iodid. ʒx. Aquæ destillat. Oij. Dissolve the chloride in a pint and a half, and the iodide of potassium in half a pint of distilled water, and mix the solutions. Collect the precipitate on a filter, and, having washed it with distilled water, dry it with a moderate heat, and keep it in a well-stopped bottle.—Ph. U. S.) Dose, one-sixteenth to one-fourth of a grain.

HYDRARGYRI MURIAS BASI OXYDI IMPERFECTI, H. oxyurias—h. Muriar corrosivum, H. oxyurias—h. Muriar dulcis sublimatus, H. submuriar—h. Oxygenatus, H. Oxyurias—h. Muriar spirituosus liquidus, Liquor hydrargyri oxyuriatis—h. Muriar suboxygenatus præcipitatione paratus, Hydrargyrum præcipitatum.

HYDRARGYRI NITRAS, Nitrus Hydrargyri in crystallis concretus, Hydrargyrum nitricum, Mercurius nitrosus frigore paratus, Nitras hydrargyrosum, Azotas hydrargyrosum, Nitrate of Mercury, (F.) Azotate de mercure, Protozotate de mercure, Nitrate mercurienx. It is employed in syphilis; and, externally, in fungous, obstinate ulcers.

It is used in the formation of the Soluble Mercury of Hahnemann.

An acid nitrate of mercury, Hydrargyri deuto-nitras seu dipernitras seu supernitras, Mercurius nitrosus calidè paratus, Nitras hydrargyricus, Azotas hydrargyricus liquidus, Liquor Hydrargyri supernitatis, Hydrargyri pernitatis Liquor, (Ph. D.) Solution of supernitrate of mercury, (F.) Deutozotate acide liquide de mercure, Nitrate acide de mercure, made by dissolving four parts of mercury in eight of nitric acid, and evaporating the solution to nine parts, has been used as a caustic in malignant ulcerations and cancerous affections.

Ward's White Drops,—a once celebrated antisyphilitic nostrum,—were prepared by dissolving mercury in nitric acid, and adding a solution of carbonate of ammonia; or, frequently, they con-

sisted of a solution of sublimate with carbonate of ammonia.

HYDRARGYRI NITRICO-OXYDUM, Hydrargyri ox'idum rubrum (Ph. U. S.), Hydrargyrum nitratum rubrum, Mercurius corrosivus ruber seu præcipitatus corrosivus seu præcipitatus ruber seu corallinus, Arcanum corallinum, Pulvis præcipis, Præcipitatus ruber, Ox'idum hydrargyri completum seu hydrargyricum, Panacea mercurii rubra, Pulvis Joannis de Vigo, Ox'idus hydrargyri rubrum, Hyperoxodes hydrargyri, Ox'idum hydrargyri nitricum seu hydrargyri rubrum per ac'idum nitricum, Nitric oxide of mercury, Red præcipitate, (F.) Oxide nitrique de mercure, Præcipité rouge ou per re, Mercure corallin, Nitrate de mercure rouge. (Hydrarg. ʒxxxvj; Acid. nitric. f ʒxviij; Aquæ Oij. Dissolve the mercury with a gentle heat, in the acid and water previously mixed, and evaporate to dryness. Rub into powder, and heat in a very shallow vessel till red vapours cease to rise.—Ph. U. S.) It is a stimulant and escharotic, and used as such in foul ulcers, being sprinkled on the part in fine powder, or united with lard into an ointment.

HYDRARGYRI OXODES RUBRUM, Hydrargyri nitrico-oxydum—h. Oxydi murias ammoniacalis, Hydrargyrum præcipitatum—h. Oxydulum nigrum, H. Oxydum cinereum.

HYDRARGYRI OXYDUM CINEREUM, Ethiops per se, Mercurius niger Moscati seu præcipitatus niger, Oxydum hydrargyri nigrum seu hydrargyrosum seu hydrargyricum præcipitatum, Oxydulum hydrargyri nigrum, Pulvis mercurialis cinereus, Mercurius cinereus, Turpe'ntum nigrum, Gray or Black oxide of Mercury, (F.) Oxide de mercure cendré, Oxide gris ou noir de mercure, Protoxide de mercure. This oxide is made in various ways. It may be formed by boiling sublimate of mercury in lime water. The dose of this Pulvis Hydrargyri cinereus is from two to ten grains. There are four other preparations of it in estimation, viz:—*Plenk's solution*, made by rubbing mercury with mucilage. 2. By rubbing equal parts of sugar and mercury together. 3. A compound of honey or liquorice and purified mercury. 4. The blue pill, and ointment. All these possess the usual properties of mercury.

The Hydrargyri Oxidum Nigrum of the Ph. U. S. is made as follows:—*Hydrarg. Chlorid. nit. Potassa. aa ʒiv, Aquæ Oj.* Dissolve the potassa in the water, allow the dregs to subside, and pour off the clear solution. To this add the chloride, and stir constantly till the black oxide is formed. Pour off the supernatant liquor, wash the black oxide with distilled water, and dry with a gentle heat. Dose, one or two grains.

The Mercurius solubilis of Hahnemann is formed from a black oxide of mercury. It is the Mercurius solubilis Hahnemannii, Oxydum hydrargyri nigri median'te ammo'niâ ex protonitrate hydrargyri præcipitatum, Azotas hydrargyroso-ammoniæ, (F.) Azotate de mercure et d'ammoniaque, Oxide gris ou noir de Mercure, Præcipité noir, Turbith noir, which has been esteemed, in Germany and France, as an antisyphilitic. Dose gr. 1-5th to gr. i. It is used in the same cases as the Hydrargyri oxydum cinereum.

HYDRARGYRI OXYDUM NIGRUM, H. oxydum cinereum—h. Oxydum nigrum mediante ammo'niâ et protonitrate hydrargyri præcipitatum, see H. oxydum cinereum—h. Oxydum nitricum, Hydrargyri nitrico-oxydum.

HYDRARGYRI OXYDUM RUBRUM, H. Binoxydum, Mercurius calcinatus, Hydrargyrum calcinatus, (F.) Oxide de Mercure rouge, Red oxide of mercury. (Made by precipitation from a solution of bichloride of mercury by solution of potassa.) See Hydrargyri nitrico-oxydum.

It is stimulant and escharotic; and, in large

doses, emetic. Owing to the violence of its operation, it is seldom given internally.

HYDRARGYRI OXYDUM RUBRUM PER ACIDUM NITRICUM, Hydrargyri nitrico-oxylum—h. Oxylum saccharatum, Hydrargyrum saccharatum—h. Oxylum sulphuricum, Hydrargyrus vitriolatus.

HYDRARGYRI OXYMURIAS, *H. Chloridum Corrosivum* (Ph. U. S.), seu *Bichloridum* seu *Pernuriar*, *Hydrargyrus mercurialis*, *Murias hydrargyri corrosivus* seu *oxygenatus* seu *hydrargyri basi oxydi imperfecti* seu *hydrargyri corrosivum*, *Sublimatus corrosivus*, *Sublimatum corrosivum*, *Mercurius corrosivus* seu *corrosivus sublimatus*, *Supermurias hydrargyri*, (F.) *Deutochlorure* ou *Bichloride de mercure*, *Muriate oxygéné de mercure*, *Sublimé corrosif*; *Bichloride* or *Oxymuriate* or *Corrosive muriate of mercury*, *Corrosive sublimate*. (Hydrarg. ℞ij, Acid. Sulphur. ℞ijj, Sodii Chlorid. ℥ss. Boil the mercury with the sulphuric acid until the sulphate of mercury is left dry. Rub this, when cold, with the chloride of sodium, in an earthenware mortar; then sublime with a gradually increasing heat.—Ph. U. S.) It is used as an antisyphilitic eutrophic in venereal complaints, old cutaneous affections, &c. Gr. iij to ℥j of water is a good gargle in venereal sore-throats, or an injection in gonorrhoea. Externally, it is applied in cases of tetter, and to destroy fungus, or stimulate old ulcers. Dose, gr. 1-16 to gr. 1-8, in pill, once in twenty-four hours. White of egg is the best antidote to it, when taken in an overdose.

The **YELLOW WASH**, *Aqua phagedænica*, *Lo'tio Hydrargyri flava*, *Phagedænic Wash*, is made by adding *corrosive sublimate* to *lime-water*, usually in the proportion of fifteen grains to half a pint. It is a common application to syphilitic sores.

HYDRARGYRI PERIODIDUM, *H. Iodidum rubrum*—h. *Pernurias*, *Hydrargyri oxyurias*—h. *Proto-iodidum*, *H. Iodidum*—h. *Proto-ioduretum*, *H. Iodidum*—h. *Proto-tartras*, *H. tartras*—h. *Prusias*, *H. cyanuretum*—h. *Saccharum vernifugum*, *Hydrargyrum saccharatum*—h. *Subchloridum*, *H. Submurias*—h. *Subiodidum*, *H. Iodidum*.

HYDRARGYRI SUBMURIAS, *H. Chloridum* seu *Subchloridum* seu *Chloridum mitè* (Ph. U. S.), *Calomelas*, *Calomelanos Torqueti*, *Calomeli*, *Hydrargyrum mercurialeum mitè*, *Draco mitigatus*, *Submurias hydrargyri mitis*, *Submurias Hydrargyri sublimatum*, *Mercurius dulcis* seu *dulcis sublimatus*;—when precipitated, *M. dulcis precipitatus*, *M. sublimatus dulcis*, *M. Zoticus Hartmanni*,—*Panacea Mercurialis* (when nine times sublimed), *Murias hydrargyri dulcis sublimatus*, *Aquila*, *Manna Metallorum*, *Panchymagogum mineralis*, *P. Quercetanus*, *mild Chloride*, *protochloride*, *submuriate*, *subchloride*, or *mild Muriate of Mercury*, *Calomel*, (F.) *Mercur doux*, *Protochlorure de mercure*, *Mild chloride of mercury* is thus directed to be prepared in the Pharmacopœia of the United States:—*Mercury*, ℞iv; *Sulphuric Acid*, ℞ij; *Chloride of Sodium*, ℥ss; *Distilled water*, a sufficient quantity. Boil two pounds of the mercury with the sulphuric acid, until the sulphate of mercury is left dry. Rub this, when cold, with the remainder of the mercury, in an earthenware mortar, until they are thoroughly mixed. Then add the chloride of sodium, and rub it with the other ingredients till all the globules disappear: afterwards sublime. Reduce the sublimed matter to a very fine powder, and wash it frequently with boiling distilled water, till the washings afford no precipitate upon the addition of liquid ammonia; then dry it. *Properties*. Antisyphilitic and sialagogue: in large doses, purgative. *Dose*: one or two grains given at night gradually excite ptialism. Gr. v to xx, purge. Children bear larger doses than adults.

The **BLACK WASH**, *Lo'tio Hydrargyri nigra*, is formed of *calomel*, ℥ij; *Lime-water*, ℥j. Used for syphilitic sores.

HYDRARGYRI SUBMURIAS AMMONIATUM, *Hydrargyrum præcipitatum*—h. *Subsulphas flavus*, *Hydrargyrus vitriolatus*—h. *Subsulphas peroxidati*, *Hydrargyrus vitriolatus*—h. *Sulphas*, *Hydrargyrus vitriolatus*, *Hydrargyri sulphas flavus*.

HYDRARGYRI SULPHURETUM NIGRUM, *H. sulphuretum cum sulphure*, *Hydrargyrus seu Mercurius cum sulphure*, *H. e sulphure*, *Ethiops mineralis*, *Pulvis hyppocratis*, *Ethiops narcoticus*, *Calomelas*, (F.) *Sulfure de mercure noir*, *Black sulphuret of mercury*, *Sulphuret of mercury with sulphur*, *Ethiops mineral*. (Hydrarg., Sulphur, aa ℔j. Rub together till the globules disappear, Ph. U. S.) Used chiefly in scrofulous and cutaneous affections. Dose, gr. x to ℥ss.

HYDRARGYRI SULPHURETUM RUBRUM, *H. Bisulphuretum*, *Hydrargyrus sulphuratus ruber*, *Milium purum seu Græcorum*, *Magnes epileptica*, *Ammon*, *Purpuriscum*, *Mercurius Cinnabarinus*, *Cinnabaris*, *Cinnabaris*, *Cinnabarium*, *Bisulphuret* or *Red Sulphuret of Mercury*, *Cinnabar*, *Cinoper*, *Vermilion*, (F.) *Sulfure de Mercure rouge*, *Cinnabre*, *Cinnabre*. (Hydrarg. ℥xl; Sulphur, ℥viii. Mix the mercury with the sulphur melted over the fire; and as soon as the mass begins to swell remove the vessel from the fire, and cover it with considerable force to prevent combustion. Rub the mass into powder and sublime. (Ph. U. S.)

It is an antisyphilitic, but is chiefly used in fumigation against venereal ulcers of the nose, mouth, and throat;—℥ss being thrown on a red-hot iron. This preparation is the basis of a nostrum, called *Boerhaave's Red Pill*.

HYDRARGYRI SUPERMURIAS, *H. oxyurias*—h. *Supernitras*, see *Hydrargyri nitras*.

HYDRARGYRI TARTRAS, *H. Proto-tartras*, *Tartrate of Mercury*. Antisyphilitic. Dose, one or two grains twice a day.

HYDRARGYRI ET ARSENICI IODIDUM, *Arsenicum* and *Mercury*, iodide of.

HYDRARGYRI ET QUINÆ PROTO-CHLORIDUM, *Protochloride of Mercury* and *Quinia*. A combination of mild chloride of mercury and quinia, administered in obstinate cutaneous diseases.

HYDRARGYRIA, *Eczema mercuriale*.

HYDRARGYRIASIS, *Hydrargyrosis*, from *ὕδαρ*, 'water,' 'mercury.' A disease induced by the use of mercury; *Merbus Mercurialis*, (F.) *Maladie Mercurielle*. Poisoning by mercury. *Eczema mercuriale*. Also, the phenomena produced by mercury on the organism, *Mercurialism*, *Mercurialisimus*.

HYDRARGYRICUM, *Mercurial*.

HYDRARGYRIUM, *Mercurial*.

HYDRARGYROSIS, *Eczema mercuriale*, *Hydrargyriasis*.

HYDRARGYRO-STOMATITIS, see *Salivation*, *mercurial*, and *Stomatitis*, *mercurial*.

HYDRARGYROTRIPSIS, *Hydrargyriatripsis*.

HYDRARGYRUM, *Hydrargyrus*, from *ὕδαρ*, 'water,' and *αργυρος*, 'silver,' *Mercurius*, *Argentum vivum* seu *mobile* seu *fusum* seu *fugitivum* seu *liquidum*, *Missadan*, *Fumus albus*, *Lorvus*, *Arca arcæcorum*, *Da'dalus*, *Mater metallorum*, *Mercury*, *Quick'silver*, formerly *Argent-vive*, (F.) *Mercur*, *M. cru*, *Vif Argent*. A fluid, brilliant metal; of a slightly bluish white colour; fluid above—39° of Fahr. and under 656°. S. g., when liquid, 13.568 (Cavendish); easily oxydized. Metallic quicksilver does not act on the body, even when taken into the stomach. When oxydized, and combined with acids, it acts powerfully. It has been exhibited in cases of constriction of the bowels and in intussusception, from a notion

that it must certainly pass through the bowels by its gravity. The water, in which mercury has been boiled, has been recommended as a vermifuge; but it probably enjoys no such property, as chemical tests do not exhibit the presence of the metal. When the crude metal is distilled in an iron retort, it forms the *Hydrargyrum purificatum*.

HYDRARGYRUM AMMONIATO-MURIATICUM, H. præcipitatum—h. Ammoniatum, H. præcipitatum—h. Biotatum, Hydrargyri iodidum rubrum—h. Biotatum cum kalio iodato, Potassii hydrargyriodidum—h. Borussicum, Hydrargyri cyanuretum—h. Cyanogenatum, Hydrargyri cyanuretum.

HYDRARGYRUM CUM CRETÂ (Ph. U. S.), *Hydrargyrum cum cretâ*, *Mercurius alkalisatus*, *Mercury with chalk*, *Æthiops alcalisatus*, *Gray powder*, (F.) *Mercure avec la craie*. (Hydrarg. 3iij; Cretæ præparat. 3v. Rub them together till the globules disappear. Ph. U. S.) A protoxide of mercury, formed by trituration with carbonate of lime. It is somewhat uncertain; and consequently not much employed as a mercurial. It possesses the properties of the black oxide of mercury, and may be advantageously exhibited in cases of diarrhoea in children, dependent upon acidity and vitiated secretions. Dose, gr. v to 3ss, twice a day, in any viscid substance.

HYDRARGYRUM CUM MAGNEŒIÂ of the Dublin Pharmacopœia resembles it in properties.

HYDRARGYRUM HYDROCYANICUM, Hydrargyri cyanuretum—h. Iodatum, Hydrargyri iodidum—h. Iodatum cum chlorido Mercurii, Mercury, iodide of chloride of—h. Iodatum flavum, Hydrargyri iodidum—h. Iodatum rubrum, Hydrargyri iodidum rubrum—h. Iodidulatum, Hydrargyri iodidum—h. Muriaticum Mite, Hydrargyri submuriatis—h. Nitricum, Hydrargyri nitris—h. Periodatum, Hydrargyri iodidum rubrum.

HYDRARGYRUM PRÆCIPITATUM, H. ammoniatum (Ph. U. S.), H. ammoniato-muriaticum, Hydrargyri ammonio-chloridum, *Mercurius cosmecticus*, *Murius oxidii hydrargyri ammoniacalis*, *M. hydrargyri sub-oxygenatus præcipitatio-neparatus*, *Submurius ammoniaco-hydrargyricus*, S. Hydrargyri ammoniatum seu Præcipitatum, Calc Hydrargyri alba, Præcipitatum album, Ammonio-chloride of Mercury, White precipitate of Mercury, White precipitate, Calcinetum majus Poterii, (F.) *Sousmuriate de mercure précipité ou Précipité blanc*. (Hydrarg. chlorid. corros. 3vj; aquæ destillat. cong., Liquor ammonia, f3viij. Dissolve the chloride in the water, with the aid of heat, and to the solution, when cold, add the solution of ammonia, frequently stirring. Wash the precipitate till it is tasteless, and dry it—Ph. U. S.)

A peroxide, combined with muriatic acid and ammonia, forming a triple salt. It is used in powder, to destroy vermin; and, united with lard, for the same purpose, as well as in scabies and some other cutaneous affections.

HYDRARGYRUM SACCHARATUM, *Æthiops saccharatus*, *Mercurius saccharatus*, *Oxidum hydrargyri saccharatum*, *Saccharum hydrargyri vermicugum*. A mild mercurial formula in several of the Pharmacopœias of continental Europe; formed by triturating one part of mercury with two of white sugar. It is used in the venereal affections of children.

HYDRARGYRUM ZOOTICUM, Hydrargyri cyanuretum.

HYDRARGYRUS, Hydrargyrum.

HYDRARGYRUS ACETATUS, *Sperma mercurii*, *Terra foliata mercurii*, *Mercurius acetatus*, Hydrargyri Acetatus, *Acetas seu Proto-acetas Hydrargyri*, Acetate of mercury. This was the basis of *Keyser's pills*, and was once much celebrated in the cure of the venereal disease. The

dose is from three to five grains, but it is not much used.

The formula for *Keyser's anti-venereal pills* was as follows:—Hydrarg. Acet. 3iv; Manna, 3xxx; Amyg. 3ij; Mac. G. Trag. q. s. To be made into pills of gr. vj each. Dose, two pills.

HYDRARGYRUS CALCINATUS, Hydrargyri oxidum rubrum—h. cum Cretâ, Hydrargyrum cum cretâ—h. Muriatus, Hydrargyri oxyurias—h. Nitratu ruber, Hydrargyri nitrico-oxidum.

HYDRARGYRUS PHOSPHORATUS, *Phosphuret mercury*. This preparation has been recommended in cases of inveterate venereal ulcers, but is now scarcely used.

HYDRARGYRUS SULPHURETUS RUBER, Hydrargyri sulphuretum rubrum—h. cum Sulphure, Hydrargyri sulphuretum nigrum—h. e Sulphure, Hydrargyri sulphuretum nigrum.

HYDRARGYRUS VITRIOLATUS, *Turpe'tum mineralis*, *Mercurius emet'icus flavus seu canst'icus flavus seu luteus*, Hydrargyri sulphus seu sulphus flavus (Ph. U. S.), *Subsulphus Hydrargyri flavus*, *Oxidum hydrargyri sulphuricum*, *Subsulphus hydrargyri peroxidati*, *Calc mercurii vitriolata*, *Turbith mineral*, (F.) *Sous-sulfate de mercure ou turbith mineral*. (Hydrarg. 3iv; Acid. Sulph. 3vj. Mix in a glass vessel, and boil in a sand-bath till a dry, white mass remains. Rub this into powder, and throw it into boiling water. Pour off the liquor, and wash the yellow precipitated powder repeatedly with hot water: then dry it—Ph. U. S.) Two grains of this mercurial act on the stomach violently. It is sometimes recommended as an emetic in amaurosis.

HYDRARTHROS, Hydrarthrus.

HYDRARTHROSIS, Hydrarthrosis.

HYDRARTHIROSIS, Hydrarthrosis.

HYDRARTHROSIS, Hydrarthrosis, H. synovia'lis, *Hydrops articuli*, *Hydrarthrosis*, *Hydrarthros*, *Meliceria*, *Spina ventosa* of Rhazes and Avicenna, *Arthritis Hydrarthros*, *Hydrarthrosis*, *Hydrarthros*, *Emmyxium articuli*, *Tumor albus*, White swelling; from ὑδωρ, 'water,' and αρθρον, 'a joint.' (F.) *Tumeur blanche*, T. lymphatique des articulations. The French surgeons apply the term Hydrarthrosis to dropsy of the articulations. White swelling is an extremely formidable disease. It may attack any one of the joints; but is most commonly met with in the knee, the haunch, the foot, the elbow, and generally occurs in scrofulous children. It consists, at times, in tumefaction, and softening of the soft parts and ligaments, which surround the joints; at others, in swelling and caries of the articular extremities of bones; or both these states may exist at the same time. The treatment consists in the employment of counter-irritants; the use of iodine internally and externally, &c. Also, Synovia.

HYDRASPIS EXPANSA, *Great freshwater Tortoise*. On the sandy banks of rivers in Peru this animal buries its eggs, from which the Indians extract oil. Its flesh supplies well-flavoured food.—Tschudi.

HYDRASTIS, H. Canadensis.

HYDRASTIS CANADENSIS, *Hydrastis*, (perhaps from ὑδωρ, 'water,' and δραω, 'I am active,') *Warnera Canadensis*, *Hydrophyllum verum*, *Canad'ian Hydrastis*, *Yellow Root*, *Orange Root*, *Yellow Puccoon*, *Ground Rasp'berry*, *Yellow Paint*, *Golden Seal*, *Indian Paint*, *Eye-balm*. Nat. Ord. Ranunculaceæ. It is used in Kentucky as a 'mouth water,' and as an outward application in wounds and local inflammations.

HYDRATION, *Hydratio*, from ὑδωρ, 'water.' Impregnation or moistening with water: as *Hydraté* means impregnated or moistened with water.

HYDRELÆON, *Hydrolæum*, from ὑδωρ,

'water,' and *ελαιον*, 'oil.' A mixture of water and oil.

HYDRELYTRON, see Hydrocele.

HYDREMA, Œdema.

HYDRENCEPHALITIS, Hydrocephalus internus.

HYDRENCEPHALIUM, Hydrocephalus internus.

HYDRENCEPHALOCÉLĒ, *Hydrocephalocélē*, from 'ὕδωρ,' 'water,' *εγκέφαλος*, 'the encephalon,' and *κλήη*, 'rupture, protrusion.' A monstrosity in which there is a fissure of the cranium, the integument of the head being present, and forming a hernial sac in which the brain lies outside the skull—the sac containing a large quantity of serous fluid. Also, Hydrocephalus chronicus.

HYDRENCEPHALON, see Hydrocephalus chronicus.

HYDRENCEPH'ALOID, from 'ὕδωρ,' 'water,' *εγκέφαλος*, 'the brain,' and *εἶδος*, 'resemblance.' Resembling hydrocephalus. *Hydroceph'aloïd disease*, *Spir'ious hydroceph'alus*, *Pseudo-encephalit'is*. Disorders of the bowels, and exhaustion in children, are at times attended with hydrocephaloid symptoms.

HYDRENCEPHALUS, Hydrocephalus internus.

HYDRETEROCÉLĒ, from 'ὕδωρ,' 'water,' *εντερον*, 'intestine,' and *κλήη*, 'a tumour.' Intestinal hernia, the sac of which encloses fluid.

HYDRETEROMPHALOCÉLĒ, *Hydrenteromphalocélē*, from 'ὕδωρ,' 'water,' *εντερον*, 'intestine,' *ομφαλος*, 'umbilicus,' and *κλήη*, 'rupture.' Umbilical hernia with intestine and water in the sac.

HYDRETEROMPHALUS, Hydrenteromphalocèle.

HYDREPIGASTRIUM, see Ascites.

HYDREPIPOCÉLĒ, from 'ὕδωρ,' 'water,' *επιπλοον*, 'omentum,' and *κλήη*, 'rupture.' Omental hernia, with water in the sac.

HYDREPIPIOMPHALUS, *Hydrepipiomphalocélē*, from 'ὕδωρ,' 'water,' *επιπλοον*, 'omentum,' and *ομφαλος*, 'umbilicus.' Umbilical hernia, with omentum and water in the sac.

HYDRETNUM, Ascites.

HYDRIASIS, Hydrosudotherapeia.

HYDRIATER, see Hydropathic.

HYDRIATRIA, Hydrosudotherapeia.

HYDRIATRICA ARS, Hydrosudotherapeia.

HYDRIATRICUS, see Hydropathic.

HYDRIATRUS, see Hydropathic.

HYDRIODAS KALICUS, see Potassæ hydriodas.

HYDRIODIC ACID, see Acid, hydriodic.

HYDRO'A, *Hydro'a*, *Aqu'ula*, *Boa*, *Planta noctis*, from 'ὕδωρ,' 'water.' An affection, which consists in an accumulation of water or serous fluid under the epidermis. Some have used *hydro'a* synonymously with *sudamina*; others with *pemphigus*. In the first case, it has generally, however, been written *hidro'a*, from 'ἵδρως,' 'sweat,' and in the latter *hydro'a*. See Hydatid.

HYDROÆ'MIA, *Hydræ'mia*, *Hæmyd'ria*; from 'ὕδωρ,' 'water,' and 'αἷμα,' 'blood.' Anæmia. The state of the blood in which the watery constituents are in excess.

HYDROAËROPLEURIE, Hydropneumothorax.

HYDROA'RION, *Hydroöph'oron*, *Hydroöphor'ia*, *Hydroö'rion*, *Hydroöva'rion*, *Hy'drops ovar'ii*, *Ascit'es ovar'ii*, *A. acca'tus*, (F.) *Hydropisie de l'ovaire*, from 'ὕδωρ,' 'water,' and *ωαριον*, 'ovarium.' Dropsy of the ovary.

HYDROATA, Sudamina.

HYDROBLEPH'ARON, *Hy'drops Pal'pebræ*,

Blepharocœ'dema aquo'sum, *Œdema pal'pebrarum*. An œdema or watery swelling of the eyelids; from 'ὕδωρ,' 'water,' and *βλεφαρον*, 'eyelid.'

HYDROCACHEXIA, Leucoplegmatia.

HYDROCARDIA, Hydropericardium, see Pericardium.

HYDROCATARRHOPHE'SIS, *Hydrocatarrhop'h'ia*, from 'ὕδωρ,' 'water,' and *καταρροφειν*, 'to sip up.' Absorption of water from without.

HYDROCE'LE, *Hydrops Scroti seu testiculor'um*, *Hydroscœ'celē*, *Hydatocē'le*, *Hydroscœ'chem*, *Hydror'chis*, *Hydrochœ'nie*, (Alibert) from 'ὕδωρ,' 'water,' and *κλήη*, 'a tumour.' A term generally applied to a collection of serous fluid in the areolar texture of the scrotum or in some of the coverings, either of the testicle or spermatic cord. To the first of these varieties the names—*External Hydrocele*, *H. œdematodes*, (F.) *H. par infiltration* have been given; and to the second, those of *Hydrocē'le interna*, *H. tunica vaginal'is testis*, *Hydatocē'lytron*, (F.) *H. par épanchement*. When the collection occurs in the envelope of the testicle, it is called *H. of the tunica vaginalis*; and the epithet *congenital* is added, when the interior of the membrane, in which it is situate, still communicates freely with the cavity of the abdomen. When it exists in the spermatic chord, it is called *encysted*, or *diffused Hydrocele of the spermatic cord*, as the case may be. The tumour of the distended scrotum is oblong: greater below than above; indolent and semi-transparent. When it becomes inconveniently large, the fluid may be evacuated by puncturing with a trocar, but, as it collects again, this operation can only be considered palliative. The radical cure consists, usually, in injecting, through the canula of the trocar, which has been left in, after puncturing, some irritating liquid, as wine. This is kept in the tunica vaginalis for a few minutes, and then withdrawn. The coat inflames; adhesion takes place, and the cavity is obliterated.

HYDROCELE OF THE NECK. A tumour, filled with a watery fluid, occupying some portion of the neck.

HYDROCELE PERITONÆI, Ascites—h. Spinalis, Hydrorachis.

HYDROCENO'SIS, from 'ὕδωρ,' 'water,' and *κνωσις*, 'evacuation.' The evacuation of water morbidly accumulated in the body.

HYDROCEPHALE, Hydrocephalus.

HYDROCÉPHALE AIGUË, Hydrocephalus internus.

HYDROCEPHALITIS, Hydrocephalus internus.

HYDROCEPHALIUM, Hydrocephalus.

HYDROCEPHALOCÉLE, Hydrencephalocèle.

HYDROCÉPHALOËCTASIE, Hydrocephalus chronicus.

HYDROCEPH'ALUS, *Hydroceph'alum*, *Hydrocra'nia*, *Hydrocra'nium*, *Hydroceph'alium*, *Hydroceph'ale*, *Hydrops Cap'itis seu Cer'e'br'i*, from 'ὕδωρ,' 'water,' and *κεφαλή*, 'the head.' Water in the head. Dropsy of the head. Dropsy of the brain. A collection of water within the head. It may be internal or external.

HYDROCEPHALUS ACUTUS, H. internus—h. Acutus senum, Apoplexy, serous—h. Adnatus, see H. chronicus—h. Congenitus, see H. chronicus—h. Externus, H. chronicus—h. Meningeus, H. internus.

HYDROCEPHALUS CHRONICUS, *Hydrencephalocē'le*, *Hydrops Cap'itis seu arachno'idææ*, *Hydrocephalus externus*, (F.) *Hydroceph'ale*, *Hydrocéphaloëctasie* of some, may exist at birth. *Hydrenceph'alon*, *Hydrocephalus congenitus* seu *adna'tus* commonly commences at an early period

of existence, and the accumulation of fluid gradually produces distension of the brain, and of the skull, with separation of the sutures. It commonly proves fatal before puberty.

HYDROCEPHALUS EXTERNUS, (*Eidema capitis*, *Cephalædema* of some, is a mere infiltration into the subcutaneous cellular tissue of the cranium.

HYDROCEPHALUS INTERNUS, *Hydrocephalus acutus* seu *meningæus*, *Hydrops cerebri*, *Encephalalgia hydrocephalica*, *Encephalitis exsudatoria*, *Encephalitis seu Meningitis Infantum* seu *hydrocephalica*, *Morbus cerebitalis Whyllii*, *Enterocephalopygia Infantum*, *Hydrophlogosis Ventriculorum cerebri*, *Encephalodachrysis*, *Phrenicula hydrocephalica*, *Hydrencephalus*, *Hydrencephalitis*, *Hydrencephalium*, *Hydrocephalitis*, *Hydromeningitis*, *Febris* seu *Apoplexia hydrocephalica*, *Curus hydrocephalica*, *Water Brain Fever*, *Whyll's Disease*, (F.) *Hydrocéphale aiguë*, *Fiebre cérébrale des Enfants*, is generally seated, according to modern observers, in the meninges and surface of the encephalon, and is a tubercular meningitis, *Meningitis tuberculosa*, *Encephalotrombosis*, *Granular* or *Serophulous meningitis*. (F.) *Méningite tuberculeuse ou granuleuse*, *Padencephalie*.—Piorry. It is observed particularly in childhood. Its march is extremely acute and often very rapid; admitting, generally, however, of division into three stages. The symptoms of the first stage are those of general febrile irritation, with head-ache, intolerance of light and sound, delirium, &c. Those of the second, which generally denote that the inflammation has ended in effusion, are, great slowness of pulse, crying out as if in distress, moaning, dilated pupil, squinting, &c.; and lastly, in the third stage—profound stupor, paralysis, convulsions, involuntary evacuations, quick pulse, and frequently death. The disease is of uncertain duration; sometimes, destroying in two or three days; at others, extending to two or three weeks. The prognosis is unfavourable. The treatment must be active during the stage of excitement,—precisely that which is necessary in phrenitis. In the second stage, the indication is;—to promote the absorption of the effused fluid. This must be done by counter-irritants, and mercury, chiefly. On dissection, water is generally found in the ventricles, or at the base of the brain; or there are evidences of previous vascular excitement, as effusions of coagulable lymph, &c.

HYDROCEPHALUS SPURIUS, *Hydrencephaloid*, (disease.)

HYDROCHAMAIMELUM, *Infusum anthemidis*.

HYDROCHEZIA, *Diarrhœa serosa*.

HYDROCHLORATE D'OR, see Gold.

HYDROCHLORINAS NATRICUS, Soda, muriate of.

HYDROCHOLECYSTIS, *Turgescencia vesiculæ fellæ*.

HYDROCHYSES, (G.) *Hydrochysen*, from *ὕδωρ*, 'water,' and *χυσίς*, 'effusion.' A family of diseases, according to the classification of Fuchs, in which there is a sudden effusion of serous fluid, as in serous apoplexy, hydrocephalus, &c.

HYDROCISSOCELE, *Hydrocissocèle*, *Hydrocœle*, from *ὕδωρ*, 'water,' *κισσός*, 'varix,' and *κύλη*, 'tumour.' A tumour, formed by the varicose distension of the veins of the spermatic cord, and by the accumulation of serous fluid in the areolar texture of the scrotum.

HYDROCELEA, *Ascites*.

HYDROCONIUM, Bath, shower.

HYDROCOTYLE ASIATICA, *Codagam*, *Pescuinus*, *Asiatie Pen'nyport*. Order, *Umbellifera*. A plant of India, Cape of Good Hope,

and Jamaica, where it is supposed to be possessed of astringent and diuretic properties. The leaves are applied to bruised parts to prevent inflammation. The *Hydrocotyle* has recently been extolled, in India, as a remedy for leprosy.

HYDROCOTYLE CENTE'LA. A South African plant, the roots and stalks of which are astringent; and used in diarrhœa and dysentery.

HYDROCOTYLE UMBELLATUM, *Acanthoides*.

HYDROCRANIA, *Hydrocephalus*.

HYDROCRANIUM, *Hydrocœlatus*.

HYDROCRITHE, *Decoctum Hordei*.

HYDROCYNAN'IC ACID, *Ac'idum Hydrocyanicum*; from *ὕδωρ*, 'water,' and *κυανός*, 'blue.' *Prussic Acid*, *Ac'idum Prussicum* seu *Berussicum* seu *Zoët'icum* seu *Zoët'icum*, *Cyanohydric* or *Cyanhydric Acid*, (F.) *Acide Hydrocyanique* ou *Prussique*. This acid exists in a great variety of native combinations in the vegetable kingdom, and imparts to them certain properties, which have been long known and esteemed; as in the bitter almond, cherry laurel, leaves of the peach tree, kernels of fruit, pips of apples, &c. When concentrated, it is liquid, colourless, of a strong smell and taste, at first cool, afterwards burning. Its s.g. at 7° Centigrade, is 0.7058. It is very volatile, and enters into ebullition at 80° Fahr. It speedily undergoes decomposition, sometimes in less than an hour, and consists of a peculiar gaseous and highly inflammable compound of carbon and azote, to which the name *Cyanogen* has been assigned; and of hydrogen, which acts as the acidifying principle; hence its name *Hydrocyanic acid*. In the Pharmacopœia of the United States, (1851,) two formulæ for the preparation of the *Ac'idum hydrocyanicum dilutum*—*Ac'idum hydrocyanicum*, Ph. U. S. of 1842—are given; the one from the *Ferro-cyanuret of Potassium*; the other from the *Cyanuret of Silver*. According to Magendie, the acid, prepared after Scheele's method—the one in common use—is of irregular medicinal power; he, therefore, recommends Gay Lussac's acid, diluted with 6 times its volume, or 8.5 times its weight of distilled water, for medicinal purposes, and this he calls *Medicinal Prussic Acid*. Dr. Ure has proposed, that the specific gravity should indicate that which is proper for medicinal exhibition; and, after comparative experiments of the gravity of the acids, obtained by different processes, he states, that the acid, usually prescribed, is of s. g. 0.996 or 0.997. Great caution is, however, necessary. One drop of pure prussic acid may instantly destroy, and the animal show hardly any traces of irritability, a few moments after death. It has been advised in laryngeal phthisis, in pulmonary phthisis, pulmonary inflammation and irritation, dyspepsia, uterine affections, hectic cough, cancer, chronic rheumatism, and mania, and as a local remedy in impetiginous affections; but, although possessed of powerful sedative properties, it is so unmanageable and the preparation so uncertain, that it is not much used. The *Dose* of Scheele's Acid, or of the *Medicinal Prussic Acid*, is from a quarter of a drop to two drops.

HYDROCYSTIS, from *ὕδωρ*, 'water,' and *κύστις*, 'a bladder.' A cyst containing a watery or serous fluid. An hydatid. Also, sacculated ascites.

HYDRODERMA, *Anasarca*.

HYDRODES, *Aqueus*.

HYDRODIARRHŒA, *Diarrhœa*, serous.

HYDROEDEMA, *Edema*.

HYDRO-ENCÉPHALORRHÉE, *Apoplexy*, serous.

HYDRO-ENTERO-EPIPOCE'LE, *Hydroepiplo-enterocœle*; from *ὕδωρ*, 'water,' *εντέρον*, 'intestine,' and *επιπλουν*, 'omentum.' *Entero-*

epiplocele, the sac of which contains a serous fluid.

HYDRO-ENTERO-EPIPLOMPHALUM, from 'ὕδωρ, 'water,' *εντερων*, 'an intestine,' *επιπλον*, 'the caul,' and *ομφαλος*, 'the navel.' Umbilical hernia, the sac of which contains intestine, epiploon, and serum.

HYDRO-ENTEROMPHALUM, *Hydrenteromphalum*; from 'ὕδωρ, 'water,' *εντερων*, 'an intestine,' and *ομφαλος*, 'the umbilicus.' Hernia umbilicalis, the sac of which contains intestine and serum.

HYDRO-EPIPOCELE, from 'ὕδωρ, 'water,' *επιπλον*, 'omentum,' and *κηλη*, 'a tumour.' Hernia, formed by omentum, the sac of which contains serum.

HYDRO-EPILO ENTEROCELE, *Hydro-entero-epiplocele*.

HYDRO-EPIPLOMPHALUM, from 'ὕδωρ, 'water,' *επιπλον*, 'the omentum,' and *ομφαλος*, 'the umbilicus.' Umbilical hernia, the sac of which contains epiploon and serum.

HYDROMELA, from 'ὕδωρ, 'water,' and *γαλα*, 'milk.' A mixture of water and milk.

HYDROGASTER, *Ascites*.

HYDROGEN, *Hydrogenium*, *Influmabile air*, *Phlogiston*, *Principium hydrogneticum* seu *hydroticum*, *Mephitis inflammabilis*, (F.) *Hydrogène*, *Air inflammable*, *Gaz des ballons*, from 'ὕδωρ, 'water,' and *γεννω*, 'I produce.' This gas, when breathed, proves fatal from containing no oxygen. When diluted with two-thirds of atmospheric air, it occasions some diminution of muscular power and sensibility, and a reduction of the force of the circulation. It has been resorted to in catarrh, hæmoptysis, and phthisis.

HYDROGEN, *Carburetted*, *Influmabile air*, *Fire damp* of miners, (F.) *Hydrogène carboné* ou *proto-carboné*, *Grisou*—obtained by passing the vapour of water over charcoal, at the temperature of ignition, in an iron tube—has been found possessed of similar properties, when diluted, and has been used in like cases.

HYDROGEN, *Protoxide of Water*.

HYDROGEN, *Sulphuretted*, *Hydrosulphuric*, *Sulphuretted*, *Sulphohydric* or *Hydrothionic acid*, *Hydrothion*, *Gaz hepaticum* seu *hydrogenium mephuratum*, *Mephitis hepatica*, (F.) *Hydrogène sulfuré*, *Acide hydrosulfurique*, *Air purant* ou *légitique*, may be disengaged from any of the sulphurets by the addition of a strong acid. It is a violent poison, but has been recommended to be inhaled, diluted, to allay the increased irritability which occasionally exists after diseases of the lungs. See *Hydro-sulphuretted Water*.

HYDROGENATION, see *Hydrogénèses*.

HYDROGÈNE, *Hydrogen*—*h*. *Carboné* ou *proto-carboné*, *Hydrogen*, *carburetted*—*h*. *Sulfuré*, *Hydrogen*, *sulphuretted*.

HYDROGÉNÈSES. Baumes gives this name to diseases which he fancifully considers to depend upon disturbed *hydrogenation*. In it he includes intermittent and remittent fevers.

HYDROGENO-SULPHURETUM AMMONIACÆ LIQUIDUM, *Ammoniacæ sulphuretum*.

HYDROGLOSSA, *Ranula*.

HYDROGRAPHY, see *Hydrology*.

HYDROHÉMIE, *Anæmia*, *Hydroæmia*.

HYDROHYMENITIS, *Orrhohymenitis*, from 'ὕδωρ, 'water,' *ὑμην*, 'a membrane,' and *itis*, denoting inflammation. Inflammation of a serous membrane.

HYDROHYSTERA, *Hydrometra*.

HYDROLÆUM, *Hydrelæon*.

HYDROLAPATHUM, *Rumex hydrolapathum*.

HYDROLATA, *Aquæ destillatæ*.

HYDROLATS, *Aquæ destillatæ*, *Water*, distilled.

HYDROLÉ, *Decoction*, *Fomentation*, *Lotion*, *Mucilage*, *Potion*, *Tisane*. See *Hydrolia*.

HYDROLICA, (F.) *Hydroliques*, from 'ὕδωρ, 'water.' Watery solutions of the active principles of medicinal agents. Those prepared by solution or admixture are termed, by some of the modern French pharmacists, *Hydrolés*; those by distillation, *Hydrolats*.

HYDROLIQUES, *Hydrolia*.

HYDROLITE, *Decoction*, *Mucilage*, *Potion*.

HYDROLOGY, *Hydrologia*, from 'ὕδωρ, 'water,' and *λογος*, 'a discourse.' A treatise on water. By the term *Medical Hydrology* is meant that part of physics, whose object is the study of water, considered as it respects medicine; and, consequently, embracing that of mineral waters. *Medical Hydrography* comprises the study of the influence exerted by the sea or by navigation on the health of man.

HYDROLOTIE, *Fomentation*, *Lotion*.

HYDROMANIA, from 'ὕδωρ, 'water,' and *μανια*, 'mania.' A name given by Strambi to pellagra, in which the patient has a strong propensity to drown himself.

HYDRO-MEDIASTINUM, *Hydrops mediastini*, Effusion of serous fluid into the mediastinum.

HYDROMEL, *Hydromeli*, from 'ὕδωρ, 'water,' and *μελι*, 'honey.' *Aqua mada*, *Melitis*, *Mulsam*, *Melicratum*, *Mellivatum*, *Braggart*, *Medo*. A liquid medicine, prepared with an ounce and a half of honey and a pint of tepid water. It is used as a demulcent and laxative, and is generally known under the names *Simple hydromel*, *Vinous hydromel*, *Mead*, *Hydromeli vinosum*, *Enomel*, is a drink made by fermenting honey and water, and is much used in some countries. As a common drink, mead is often made of honey and some alcoholic liquor well beaten together. See *Mahogany*.

HYDROMELLÉ. A preparation formed like the syrups, using hydromel in place of sugar.—Béral.

HYDROMENINGITIS, *Aquo-membranitis*, *Hydrocephalus internus*.

HYDROMETER, *Areometer*.

HYDROMETRA, *Hydrometria*, *Hydrohysteria*, *Hydrometrectasia*, *Hysterites*, *Hysterocelema*, *Hydrometra uteri*, from 'ὕδωρ, 'water,' and *μετρα*, 'the womb.' *Dropsy of the womb*. A disease characterized by circumscribed protuberance in the hypogastrium,—with obscure fluctuation, progressively enlarging, without ischury or pregnancy. If it ever occur, it must be a rare disease.

HYDROMETRECTASIA, *Hydrometra*.

HYDROMPHALUM, *Hydrops umbilicalis*, *Exomphalus aquosus*, *Her'nia umbilici aquosa*, from 'ὕδωρ, 'water,' and *ομφαλος*, 'the navel.' A tumour, formed by the accumulation of serum in the sac of umbilical hernia: or simply by distension of the navel in cases of ascites.

HYDROMYRINGA, *Hydromyrinx*, *Hydrops tympani*, from 'ὕδωρ, 'water,' and *myrinxa* or *myrinx*, 'the membrana tympani.' *Dropsy of the drum of the ear*; giving rise to difficulty of hearing.—*Dysacousia hydroptica*.

HYDROMYRINX, *Hydromyringa*.

HYDRONCUS, *Anasarca*, *Edema*.

HYDRONEPHROSIS, (F.) *Hydronephrose*, *Hydroneurde distension*, from 'ὕδωρ, 'water,' and *νεφρος*, 'kidney.' An accumulation in the kidney, owing to the obstruction of the tubes of the papillæ.—Rayer.

HYDRONOSUS, *Hydrops*.

HYDRONUSUS, *Hydrops*.

HYDROOARIUM, *Hydrooarion*.

HYDROOPHORON, *Hydrooarion*.

HYDROOVARIIUM, *Hydrooarion*.

HYDROPATHIC, *Hydropathicus*, *Hydriaticus*, from 'ὕδωρ, 'water,' and *παθος*, 'disease.'

Relating to hydrophaty or the water-cure,—as a hydrophatic physician, *Hydriū'ter, Hydriū'trus*.

HYDROPATHY, *Hydrosudotherapeia*.

HYDROPEDE'SIS, from *ὑδωρ*, 'water,' and *πῆδω*, 'I break out.' *Ephidro'sis*. Excessive sweating.

HYDROPEGE, Water, spring.

HYDROPELTIS PURPUREA, *Brasenia hydropeletis*.

HYDROPERICARDIA, *Hydropericardium*.

HYDROPERICARDITIS, *Hydropericardium*.

HYDROPERICARDIUM, *Hydropericardium*. *Hydropericardi'tis*, from *ὑδωρ*, 'water,' and *pericardium*; *Hydrops Pericardii, Hydrocar'dia*. Dropsy of the pericardium, (F.) *Hydropisie du Péricarde*. This is not a common disease. Palpitations; irregular or intermitting pulse; excessive dyspnoea, amounting often to orthopnoea, and dulness over a large space on percussion, will cause the pericardium to be suspected. The treatment is that of dropsies in general. It is, usually, however, of the active kind.

HYDROPÉRIONE, from *ὑδωρ*, 'water,' *περί*, 'around,' and *ων*, 'an egg, or ovum.' The sero-albuminous substance, secreted by the lining of the uterus prior to the arrival of the impregnated ovum in that cavity.—Breschet.

HYDROPERITONEUM, *Ascites*.

HYDROPÉRTONIE, *Ascites*.

HYDROPHAL'LUS, from *ὑδωρ*, 'water,' and *φαλλος*, 'the male organ.' Edema of the male organ.

HYDROPHIMOSIS, *Phimos oedematodes*.

HYDROPHLEGMASIA TEXTŪS CELLULARIS, *Phlegmasia alba*.

HYDROPHLOGOSIS VENTRICULORUM CEREBRI, *Hydrocephalus internus*.

HYDROPHOB'IA, *Paraphob'ia, Parophob'ia, Phobodip'son, Pheulgydron, Pheigophob'ia, Cynolys'sa, Cynolys'sum, Morbus hydrophob'icus seu hydroph'obus, Lycan'che, Lycan'chis, Aërophob'ia, Phren'itis latrans, Lytta, Lyssa, L. can'ina, Panto'hob'ia, Rabies can'ina, Erethis'mus hydrophob'us, Cynos hydrophob'ia, Hydrophobia, Aqua metus, Canine madness, Water fright, (F.) Rage; from ὑδωρ, 'water,' and *φοβος*, 'dread.'*

The term *Rabies* is more appropriate for the aggregate of symptoms resulting from the bite of rabid animals. *Hydrophobia* literally signifies a 'dread of water,' and, consequently, ought to be applied to one of the symptoms of rabies, rather than to the disease itself. It is a symptom which appears occasionally in other nervous affections. Rabies is susceptible of spontaneous development in the dog, wolf, cat, and fox, which can thence transmit it to other quadrupeds or to man; but it has not been proved that it can supervene,—without their having been previously bitten,—in animals of other species; or that the latter can, when bitten, communicate it to others. Many facts induce the belief, that the saliva and bronchial mucus are the sole vehicles of the rabid virus; the effects of which upon the economy sometimes appear almost immediately after the bite, and are, at others, apparently dormant for a considerable period. The chief symptoms are—a sense of dryness and constriction of the throat; excessive thirst; difficult deglutition; aversion for, and horror at, the sight of liquids as well as of brilliant objects; red, animated countenance; great nervous irritability; frothy saliva; grinding of the teeth, &c. Death most commonly happens before the fifth day. *Hydrophobia* has hitherto resisted all therapeutical means. Those which allay irritation are obviously most called for. In the way of prevention, the bitten part should always be excised, where practicable; and cauterized.

In some cases, symptoms like those which follow the bite of a rabid animal are said to have come on spontaneously. This affection has been termed *nervous* or *spontaneous hydrophobia*.

HYDROPHOBUS, *Lyssodectus*.

HYDROPH'THAL'MIA, from *ὑδωρ*, 'water,' and *οφθαλμος*, 'the eye.' *Hydrophthal'mus, Dropsy of the eye, Hydrops Oculi, Bupthal'mus, Zoophthal'mus, Oculus Bovi'nus seu Bul'bus seu Elephant'inus, Ophthalmop'tis, Parop'tis Staphylo'ma simplex, (F.) Hydrophthal'mie, Hydropisie de l'œil*. This affection is caused, at times, by an increase in the quantity of the aqueous, at others, of the vitreous, humour. In the former case, the iris is concave anteriorly, and pushed backwards:—in the latter, it is convex, and pushed forwards. Most commonly, the disease seems to depend on both humours at the same time. *Hydrophthal'mia* sometimes affects both eyes; at others, only one. Children are more exposed to it than adults or old persons. The treatment must vary according to the cause; its longer or shorter duration; greater or less extent, &c. Hence, according to circumstances, hydragogue medicines, purgatives, general and local blood-letting, blisters, setons, moxa, cupping-glasses, fomentations, collyria, and fumigations of different kinds, have been employed. When all means fail, and the disease continues to make progress, the fluid may be evacuated, by a puncture made with a cataract needle at the lower part of the transparent cornea.

HYDROPH'THAL'MIA, CONICAL, *Staphyloma of the cornea*.

HYDROPH'THAL'MION. Samt etymon. An oedematous swelling of the conjunctiva in hydroptic persons.

HYDROPH'THALMUS, *Hydrophthal'mia*—h. Cruentus, *Hæmophthal'mia*.

HYDROPHYLLUM VERUM, *Hydrastis Canadensis*.

HYDROPHYSOCE'LE, *Hydropneumatoce'le*, from *ὑδωρ*, 'water,' *φύσα*, 'wind,' and *κληη*, 'a tumour.' *Hernia*, which contains a serous fluid and gas.

HYDROPHYSOME'TRA, from *ὑδωρ*, 'water,' *φύσσω*, 'I inflate,' and *μήτρα*, 'the womb.' A morbid condition of the womb, in which both fluid and air are contained in it.

HYDROPTIC, *Hydropticus, Hy'phydros, Hyder'icus, Hydero'des, Drop'sical, (Sc.) Edroppit, (F.) Hydropique*. One labouring under dropsy. *Drop'sical*. Relating to dropsy. Also, an anti-hydrotic, and a hydragogue.

HYDROPIPEK, *Polygonum hypopiper*.

HYDROPI'SIA, *Hydros*.

HYDROPI'SIE, *Hydros*—h. du Bas-ventre, *Ascites*—h. Cardiaque, *Dropsy, cardiac*—h. Cérébrale suraiguë, *Apoplexy, serous*—h. Hépatique, *Dropsy, hepatic*—h. de l'Œil, *Hydrophthal'mia*—h. de l'Ovaire, *Hydronarion*—h. du Péricarde, *Hydropericardium*—h. du Péritoine, *Ascites*—h. des Plèvres, *Hydrothorax*—h. de Poitrine, *Hydrothorax*—h. Rénale, *Dropsy, renal*—h. de la Vésicule du Fiel, *Turgescencia vesicæ felleæ*.

HYDROPI'SIS, *Hydros*—h. Vera, *Anasarca*.

HYDROPI'SMUS, *Hydros*.

HYDROPLET'RIE, *Hydrothorax*.

HYDRO-PNEUMATOCELE, *Hydrophysocèle*.

HYDRO-PNEUMON, *Hydropneumonia*.

HYDRO-PNEUMON'IA, *Hydropneum'ion, Œde'ma pulmonum chron'icum, Hydrops pulmonum seu pulmonum cellulosus, Anasar'ca pulmonum*, from *ὑδωρ*, 'water,' and *πνευμων*, 'the lung.' Dropsical infiltration of the lungs. See *Œdema of the lungs*.

HYDRO-PNEUMOSAR'CA, from *ὑδωρ*, 'water,' *πνευμα*, 'wind, air,' and *σαρξ*, 'flesh.' An ab-

acess, containing water, air, and matters similar to flesh.—M. A. Severinus.

HYDROPNEUMOTHORAX, *Hydroaëropleurie*, from *ὕδωρ*, 'water,' *πνεῦμα*, 'the lung,' and *θώραξ*, 'the chest.' Pneumothorax with effusion of blood into the chest. See Pneumothorax.

HYDROPOIDES, from *ὕδωρ*, 'water,' and *ποιεῖν*, 'I make.' An epithet for watery excretions, such as sometimes take place in hydropsies.

HYDROPOTA, *Hydropotes*, *Potator Aquæ*, from *ὕδωρ*, 'water,' and *ποτῆς*, 'a drinker.' A water drinker. One who drinks only water, or drinks it in an extraordinary quantity.

HYDROPS, from *ὕδωρ*, 'water,' *Phlegmatia*, *Hydros*, *Affectus hydroides*, *Hydrophisis*, *Hydrophisia*, *Hydrophismus*, *Hydrophisia*, *Hydrophisa*, *Hydrophoeis*, *Hydrophoeis*, *Hydrophoeis*, *Polyhydria*, *Hydrophoeis*, *Hydrophoeis*, *Dropsy*, *Hydropsy*, (Old Eng.) *Dropsy*, (F.) *Hydropsie*. A preternatural collection of a serous fluid in any cavity of the body, or in the areolar texture. When the areolar texture of the whole body is more or less filled with fluid, the disease is called *Anasarca* or *Leucophlegmatia*;—and when this variety is local or partial, it is called *Œdema*. The chief dropsies, designated from their seat, are:—*Anasarca*, *Hydrocephalus*, *Hydrorachitis*, *Hydrothorax*, *Hydropericardium*, *Ascites*, *Hydrometra*, *Hydrocele*, &c.

Encysted Dropsy, *Hydrops sacculus*, *incarceratus* seu *cysticus*, is that variety in which the fluid is enclosed in a sac or cyst; so that it has no communication with the surrounding parts. *Dropsy of the Ovary*, *Hydrops* seu *Ascites Ovarii*, *Ascites sacculus*, is an instance of this variety.

Dropsy may be active or passive. The former consists in an increased action of the exhalants, so that those vessels pour out much more fluid than is absorbed: the latter arises from a state of atony of the absorbent vessels, which allows of an accumulation of fluid. It may also be mechanical, or produced by obstructions to the circulation, as in cases of diseased liver. Active dropsy, occurring accidentally in a sound individual, generally ends favourably. That which supervenes on other diseases, or is symptomatic of some internal affection, is rarely curable.

The treatment consists in the use of all those remedies which act on the various secretions: so that, the demand being increased, the supply will have to be increased accordingly; and in this manner some of the collected fluid may be taken up by the absorbents. To this end bleeding, if the dropsy be very active; purgatives, diuretics, sudorifics, sialogogues, &c., are the remedies chiefly depended upon.

HYDROPS ABDOMINIS, *Ascites*—h. *Abdominis acutus*, *Tympanites*—h. *Abdominis sacculus*, see *Ascites*—h. *Anasarca*, *Anasarca*—h. *Anasarca acutus*, see *Anasarca*—h. *Arachnoidea*, *Hydrocephalus chronicus*—h. *Articulorum*, *Hydrarthrosis*—h. *Ascites*, *Ascites*—h. *Capitis*, *Hydrocephalus*, *Hydrocephalus chronicus*—h. *Cardiacus*, *Dropsy*, *cardiac*—h. *Cavitatis columnæ vertebralis*, *Hydrorachis*—h. *Cellularis artuum*, *Œdema*—h. *Cellularis totius corporis*, *Anasarca*—h. *Cellulosus*, *Anasarca*—h. *Cerebri*, *Hydrocephalus*, *Hydrocephalus internus*—h. *Cutaneus*, *Anasarca*—h. *Cysticus*, see *Ascites*, and *Hydrops*—h. *Glottidis*, *Œdema of the Glottis*—h. *Hepaticus*, *Dropsy*, *hepatic*—h. *Incarceratus*, see *Hydrops*—h. *Intercus*, *Anasarca*—h. *Leucophlegmatia*, *Leucophlegmatia*—h. *ad Matulum*, *Diahetes*—h. *Mediastini*, *Hydromediastinum*—h. *Medullæ spinalis*, *Hydrorachis*—h. *Matellæ*, *Diahetes*—h. *Oculi*, *Hydrophthalmia*—h. *Ovarii*, *Hydrooion*, see *Hydrops*—h. *Palpebræ*, *Hydroblepharon*—h. *Pectoris*, *Hydrothorax*—h. *Pericardii*,

Hydropericardium—h. *Pleura*, *Hydrothorax*—h. *Pulmonis*, *Hydrothorax*—h. *Pulmonum*, *Hydropneumonia*, *Œdema of the Lungs*—h. *Pulmonum cellulosus*, *Hydropneumonia*—h. *Renalis*, *Dropsy*, *renal*—h. *Sacculus*, see *Hydrops*—h. *Sacci lacrymalis*, *Fistula lacrymalis*—h. *Serotii*, *Hydrocele*—h. *Siccus et flatulentus*, *Tympanites*—h. *Spinæ*, *Hydrorachis*—h. *Spinæ vertebralis*, *Hydrorachis*—h. *Subcutaneus*, *Anasarca*—h. *Tela cellulosa*, *Anasarca*—h. *Testiculorum*, *Hydrocele*—h. *Thoracis*, *Hydrothorax*—h. *Tubarium Falcipii*, *Hydrosalpinx*—h. *Tympani*, *Hydromyrrina*—h. *Tympanites*, *Tympanites*—h. *Umbilicatis*, *Hydromphalum*—h. *Uteri*, *Hydrometra*—h. *Vesicæ fellæ*, *Turgescencia vesiculæ fellæ*.

HYDROPSIA, *Hydrops*.

HYDROPSY, *Hydrops*.

HYDROPYRETOS, from *ὕδωρ*, 'water,' and *πυρετος*, 'fever.' *Febbris sudatoria*. Fever with sweating. *Hydropyretos* would be more proper; from *ἵδρω*, 'sweat.' See *Sudor Anglicus*.

HYDRORACHIS, from *ἵδωρ*, 'water,' and *ραχίς*, 'the spine.' *Hydrorachis*, *Hydrorachitis*, *Hydrorachia*, *Myelochyria*, *Hydrops Cavitas Columnæ Vertebralis* seu *Spinæ vertebralis* seu *medullæ spinalis* seu *spinæ*, *Hydrocellæ spinalis*, *Hydrorachitis spinosa*. An effusion of serum, often owing to inflammation of the spinal membranes—*myelitis exudativa*—and forming a soft, frequently transparent, tumour, constituted of the membranes of the spinal marrow, which are distended and projecting backwards from the vertebral canal, the posterior paries of which, when the affection is congenital, is wanting to a certain extent—*Spina bifida*, *Atelorachidia*, *Hydrorachis dehiscentis* seu *congenita*, *Schistorrhachis*, *Spina*. The disease is often accompanied by paralysis of the lower extremities. It is congenital, and situate in the lumbar or sacral regions. It is almost always fatal:—the tumour rupturing, and death occurring instantaneously. On dissection, a simple separation or complete absence of the spinous processes of the vertebrae is perceived, with, at times, destruction or absence of spinal marrow. The treatment is the same as in *hydrocephalus chronicus*; and, as in it, advantage seems occasionally to have been derived by puncturing with a fine needle.

HYDRORACHIS DEMISCENS, see *Hydrorachis*.

HYDRORACHITIS, *Hydrorachis*.

HYDRORACHIS, *Hydrocele*.

HYDRORÉNALE DISTENSION, *Hydro-nephrosis*.

HYDRORRHACHIA, *Hydrorachis*.

HYDRORRHACHIS, *Hydrorachis*—h. *Congenita*, see *Hydrorachis*—h. *Dehiscentis*, see *Hydrorachis*.

HYDRORRHAGIE, *Apoplexy*, *serous*.

HYDRORRHŒ, *Hydrops*.

HYDRORRHŒA, *Hydrops*. Also, a copious watery discharge,—as from the vagina.

HYDRORRHŒUS, *Hydrops*.

HYDRORTHOPNŒA, from *ὕδωρ*, 'water,' and *orthopnea*, 'difficulty of breathing, except in the erect posture.' *Orthopnea*, owing to a collection of water in the chest.

HYDROSACCHARUM, *Aqua saccharata*, (F.) *Eau sucrée*. Sugared water.

HYDROSALP'INX, *Hydrops tubarum Fallopii*; from *ὕδωρ*, 'water,' and *σαλπῆξ*, 'a tube.' Dropsy of the Fallopian tube.

HYDROSAR'CA, from *ὕδωρ*, 'water,' and *σάρξ*, 'flesh.' A tumour containing a fluid, as well as portions of flesh. Also, *Anasarca*.

HYDROSARCOCELE, from *ὕδωρ*, 'water,' *σάρξ*, 'flesh,' and *κύλη*, 'a tumour.' *Sarcocystocèle*. A tumour, formed by a sarcocele, complicated with dropsy of the tunica vaginalis.

HYDROSCHEOCELE, *Oscheocelē aquosa*, from 'ὕδωρ, 'water,' σκῆλον, 'the scrotum,' and κηλη, 'rupture.' A collection of water in the scrotum. Hydrocele.

HYDROSCHEONIE, Hydrocele.

HYDROSCHEUM, Hydrocele.

HYDROSIS, Hidrosis.

HYDROSTATIC TEST OF INFANTICIDE, see Doctinasia—h. Bed, Water bed.

HYDROSUDOPATHY, Hydrosudotherapeia.

HYDROSUDOTHERAPEIA, *Hydrops'athy, Hydropathia, Hydrosudop'athy, Hydriasis, Ars hydriatrica, Hydriatri'a, Hydatotherapeia, Hydrotherapeia, Water cure*, (G.) Wasserkur, from 'ὕδωρ, 'water,' sudo, 'I sweat,' and θεραπεύω, 'I remedy.' A badly compounded word, formed to express the mode of treating diseases systematically by cold water, sweating, &c.

HYDROSULPHURET'ED WATER, *Aqua hydrosulphurata simplex, A. hepatica*, (F.) Eau hydrosulphurée simple. (Sulphuret of iron 1000 parts, sulphuric acid 2000 parts, distilled water 4000 parts: add the water to the acid, and put the sulphuret of iron into a retort, to which a Wolf's apparatus of five or six vessels is adapted: the last containing about an ounce of potassa dissolved in a quart of water. Pour the diluted acid gradually on the sulphuret, and, ultimately, throw away the water in the last vessel. Ph. P.) It is stimulant, diaphoretic, and deobstruent [?]. and is used in rheumatism, diseases of the skin, &c.

It has been, also, called *Ac'idum Hydrothion-icum liq'idum*.

HYDROSULPHURETUM AMMONIACALE AQUOSUM, Ammoniac sulphuretum—h. Ammoniacum, Ammoniac sulphuretum.

HYDROTHERAPIA, Hydrosudotherapeia.

HYDROTHION, Hydrogen, sulphuretted.

HYDROTHORAX, from 'ὕδωρ, 'water,' and θώραξ, 'the chest.' *Hydrops Thoracis seu pectoris seu pulmonis seu pleuræ, Stethochylisis, Pleurorrhœa lymphatica seu serosa, Dyspnœa et Orthopnœa hydrothoracica*, (F.) *Hydropisie de Poitrine ou des Plèvres, Dropsy of the Chest*. Idiopathic hydrothorax, termed, by Laënnec, *Hydropisie des plèvres, Dropsy of the Pleuræ*,—by Piorry, *Hydropleurie*,—is a rare disease, and difficult of diagnosis. It generally exists only on one side, which, if the fluid effused be considerable, projects more than the other. Dyspnœa, and fluctuation perceptible to the ear, are characteristic symptoms. When the chest is examined with the stethoscope, respiration is found to be wanting every where, except at the root of the lung. The sound is also dull on percussion.

Effusion into the chest, as a result of inflammation of some thoracic viscus, is as common as the other is rare. It is usually a fatal symptom. It has been called *symptomatic hydrothorax*.

In hydrothorax, the course of treatment proper in dropsies in general must be adopted. Diuretics seem, here, to be especially useful: probably on account of the great activity of pulmonary absorption. Paracentesis can rarely be serviceable.

HYDROTHORAX CHYLOSUS, Chylothorax—h. Purulentus, Empyema.

HYDROTICA, Hydragogues.

HYDROTIS, from 'ὕδωρ, 'water,' and otis, gen. ωτος, 'the ear.' Dropsy of the ear. Properly, an accumulation of mucous or muco-purulent matter in the middle ear.

HYDRURESIS, Diabetes.

HYDRURIA, Diabetes, see Urine.

HYÈRES. This small town, agreeably situated on the declivity of a hill, about two miles from the Mediterranean, and twelve from Toulon, is

the least exceptionable residence in Provence for the pulmonary invalid. It is in some measure protected from the northerly winds; but not sufficiently so from the *mistral* to render it a very desirable residence for the phthisical.—Sir James Clark.

HYGEA, Hygiene, Sanitas.

HYGEIA, Sanitas.

HYGEISMUS, Hygiene.

HYGEOLOGY, Hygiene.

HYGIANISIS, Sanitas.

HYGIASIS, Sanitas.

HYGIASMA, Medicament.

HYGIAS'TICA DOCTRINA. The doctrine of health. The doctrine of the restoration of health.

HYGIASTICUS, Salutory.

HYGIEA, Sanitas.

HYGIEIA, Hygiene, Sanitas.

HYGIENUS, Salutory.

HYGIEIOLOGIA, Hygiene.

HYGIÈNE (F.), (generally Anglicised, *Hygiene*, and pronounced *hygeene*), from 'ὑγιεινᾶ, 'health.' *Hygeis'mus, Hygie'nē, Hygie'sis, Hygieia; Hygie'a, Hygie'a, Hygie'a, Hygiene, Hygien'ics, Medicina conservat'ica seu diastatica, Hygeolog'ia, Hygienolog'ia, Hygeol'ogy, Hygiol'ogy*; from 'ὑγιος, 'healthy.' The part of medicine whose object is the preservation of health. It embraces a knowledge of healthy man, both in society and individually, as well as of the objects used and employed by him, with their influence on his constitution and organs. See Regimen.

HYGIEN'IC, (F.) *Hygiénique*. Same etymon. Relating to Hygiene—as 'hygienic precautions, hygienic rules,' &c. &c.

HYGIENICS, Hygiene.

HYGIÉNIQUE, Hygienic.

HYGIEN'IST. One who understands the principles of hygiene.

HYGIERUS, Salutory.

HYGIESIS, Hygiene.

HYGIOLOGY, Hygiene.

HYGRA, from 'ὕδωρ, 'water,' or 'ὑγρος, 'humid.' Liquid plasters;—*Hygremplastræ*.

HYGRASIA, Humour.

HYGRECHEMA, from 'ὑγρος, 'humid,' and χῆμα, 'sound.' *Sonitus fluitidi*. The sound of fluid, heard by auscultation, or otherwise.

HYGREDON, Humour.

HYGREEMPLASTRA, Hygra.

HYGROBLEPHARICI, from 'ὑγρος, 'humid,' and βλεφαρον, 'eyelid.' *Hygrophthalmici*. The excretory ducts of the lachrymal gland have been so called.

HYGROBRONCHIORRHONCHUS, *Râle muqueux*.

HYGROCATARACTA; from 'ὑγρος, 'humid,' and καταρακτῆς, 'cataract.' *Cataracta liquida*. Liquid or fluid cataract.

HYGROCELE, Hydrocirsocele.

HYGROCOLLYRIA, see Collyrium.

HYGROCYSTIS, Hydatid.

HYGROLOGY, *Hygrol'ogy*, from 'ὑγρος, 'humid,' and λογος, 'a discourse.' The anatomy of the fluids of the body.

HYGRO'MA, from 'ὑγρος, 'humid.' *Tumor cysticus serosus, Cys'tis serosa*. Dropsy of the bursæ mucosæ.

HYGROMET'RIC, *Hygromet'ricus*, (F.) *Hygrométrique*. Same etymon as the next. Relating or appertaining to hygrometry.

HYGROMETRIC PROPERTY of an organized tissue, (F.) *Hygrométrieité*, is that by which it gives off to the air, and receives from it moisture under varying conditions of that medium.

HYGROMÉTRICITÉ, Hygrometric property.

HYGROMETRY, *Hygromet'ria*, *Hygrosco'p-ia*, from *υγρος*, 'humid,' and *μετρον*, 'measure.' The part of physics which concerns the measurement of the dryness or humidity of the atmosphere. It is probable, that diseases are as frequently caused by the varying moisture of the atmosphere or by changes in its weight or temperature. The *hygrometer* ought, consequently, to form part of every apparatus for medical meteorological observations.

HYGRON, Liquor.

HYGROPHOBIA, *Hydrophobia*.

HYGROPHTHAL'MIA, from *υγρος*, 'humid,' and *οφθαλμία*, 'inflammation of the eye.' Ophthalmia with much lachrymation.

HYGROPHTHALMICI, *Hydrophorarii*.

HYGROPISSOS, see *Pinus sylvestris*.

HYGROSCOPIA, *Hygrometry*.

HYGROTES, Humour, Liquor.

HYGRUM, Humour.

HY'L'E, *υλη*, *Materia*, 'Matter. Wood.' *Materia Medica*; also, the Philosopher's stone.

HYLE, *ΙΥΡΙΞ*, *Materia Medica*.

HYLISIS, *Colatio*.

HYLISMUS, *Colatio*.

HYLISTER, *Colatorium*.

HYLOPH'AGOUS, from *υλη*, 'wood,' and *φαγω*, 'I eat.' One that feeds upon the young shoots of trees, roots, &c. *Hylophagous* tribes yet exist in some parts of Africa.

HYMEN, *υμην*, which signifies 'marriage,' 'nuptial song,' 'membrane or pellicle.' *Claustrum seu Sigi'llum seu Cunctid'ia seu Colum'na seu Zona virginita'tis*, *Flos virgin'a'lis*, *Virgin'ia*, *Cer'entus membran'osus*, *Buclon*, *Interceptum virgin'a'le*, *Cento virgin'a'lis*, *Argumentum Integritat'is*, *Munimen'tum seu Zona Castita'tis*, *Panniculus hymen'aeus seu virgin'a'lis*, *Eugian*, *Vul'vula seu Membran'ula lunata vagin'ae*, *Vir'ginal membrane*. The semilunar, parabolic, or circular fold, situate at the outer orifice of the vagina in virgins, especially during youth, and prior to menstruation. This membrane is ordinarily ruptured by the first venereal act, and is effaced by accouchement; some irregular flaps remaining, to which the name *Caruncula Myrtiformis* has been given by reason of their resemblance to the leaves of the myrtle. Many circumstances of an innocent character may occasion a rupture or destruction of this membrane. It is often, indeed, found absent in children soon after birth; whilst it may remain entire after copulation. Hence the presence of the hymen does not absolutely prove virginity; nor does its absence prove incontinence; although its presence would be *primâ facie* evidence of continence.

HYMEN, Membrane—*h*. *Diaphragmaton*, *Mediastinum*.

HYMENÆA, see *Copal*—*h*. *Courbaril*, see *Animo*.

HYMEN'ICA AMENORRHŒ'A. *Amenorrhœa* occasioned by closure of the hymen.

HYMEN'ICUS, *Hymen'odes*, from *υμην*, 'hymen,' &c. Relating to the hymen. Also, membranous.

HYMENITIS, from *υμην*, 'a membrane,' and *itis*, denoting inflammation. Membranous inflammation. Inflammation of an internal membrane.

HYMEN'NIUM, *Membran'ula*, diminutive of *υμην*, 'a membrane.' A fine, delicate membrane.

HYMENODES, *Hymenicus*.

HYMENOGANGLIITIS, *Cholera*.

HYMENOG'ENY, *Hymenogen'ia*, (*F.*) *Hyménogénie*, from *υμην*, 'a membrane,' and *γεννω*, 'I engender.' The production of membranes by the effect of simple contact of two liquids, as albumen and fat, when the former gives a coating to the globules of the latter.—*Ascherson*.

HYMENOG'RAPIHY, *Hymenogra'phia*, from *υμην*, 'a membrane,' and *γραφω*, 'I describe.' That part of anatomy whose object is the description of the different membranes.

HYMENOL'OGY, *Hymenolog'ia*, from *υμην*, 'a membrane,' and *λογος*, 'a description.' A treatise on the membranes.

HYMENOPHTHALMIA, *Ophthalmia membranarum*.

HYMENOR'RHAPHY, *Hymenorrh'a'phia*, from *υμην*, 'the hymen,' and *ραφω*, 'a suture.' A form of elytrorrhaphy, in which the operation is performed in the natural situation of the hymen.

HYMENOT'OMY, *Hymenotom'ia*, from *υμην*, 'a membrane,' and *τεμνω*, 'I cut,' 'I dissect.' The part of anatomy which treats of the dissection of the membranes. The term has also been applied to the incision of the hymen, practised in certain cases of imperforation of the vagina, in order to give exit to the blood retained and accumulated in the cavity of the uterus.

HYMNIUM, *Amnios*.

HYO: in composition, an abridgment of *Hyoides os*.

HYOBASTOGLOSSUS, *Basioglossus*.

HYOCHONDROGLOSSUS, *Hyoglossus*.

HYODEGLOSSUS, *Hyoglossus*.

HYODEOTHYREODES, *Thyreohyoideus*.

HYODES, *Hyoïdes*.

HYODEPIGLOTTICUS, *Hyo-epiglotticus*.

HYO-EPIGLOT'TICUS, *Hyo-epiglotticus*.

Belonging to the os hyoides and epiglottis. Some anatomists have given the name *Hyo-epiglottic ligament* to a bundle of condensed areolar tissue, which passes from the posterior part of the body of the hyoid bone to the base of the epiglottic fibro-cartilage.

HYO-GLOSSO-BASI-PHARYNGIEN, *Constrictor pharyngis*.

HYOGLOSSUS, *Hyo'deo-glossus*, *Hyo-chondro-glossus*, *Hyopiglossus*, *Cer'ato-glossus* of Douglass and Cowper: *Basio-Cerato-Chondro-glossus*. A large, thin, quadrilateral muscle, situate at the anterior and superior part of the neck. Its insertions at three different points of the os hyoides permit it to be divided into three portions:—the first, (*Cerato-glossus* of Albinus) is attached to the great cornu of the os hyoides: the second, (*Basio-glossus* of Albinus,) arises from the superior part of the body of the same bone; and the third, (*Chondro-glossus* of Albinus,) arises from the lesser cornu and the cartilage, situate between the body and the greater cornu. The fibres of these three bundles are inserted into the lateral and inferior parts of the tongue. This muscle depresses the base of the tongue, or raises the os hyoides, when the tongue is fixed.

HYO-THYREOÏDES, *Thyreohyoideus*.

HYO-THYROID, *Thyreohyoïd*.

HYOID BONE, *Hyoïdes os*.

HYOÏDES OS, *Oa Bico'r'nē seu hypsela'dēs seu Lambdoï'des seu Gnat'uris seu Gnat'ula seu Lingua seu Lingua'lē*, *Upsilonoïdes*, *Ypsiloïdes*. The *Hyoid Bone*; from the Greek *υ*, and *ειδος*, 'shape.' *Hyoïdens*, *Hyoïdes*. The hyoid bone is a very movable, osseous arch; of a parabolic shape; convex before, and suspended horizontally in the substance of the soft parts of the neck, between the base of the tongue and the larynx. This bone, separated entirely from the rest of the skeleton, is composed of five distinct portions, susceptible of motion on each other. The first, and most central, is the body of the hyoid, *Ossi'dium me'dium Hyoï'dis*, which affords attachment to several muscles; the two others are lateral, and bear the name of *branches* or *greater cornua*. The last two are smaller, situate above the other, and are known under the name *lesser cornua* and *styloid cornua*, the *Ossa pisi'formia lingua'lia*

of Sömmerring, *Ossa seu Grana seu Ossicula seu Cor'pora seu Corpuscula tritic'ea*. The os hyoides is ossified from five points.

HYOIDES PRIMUS, Sternoc-hyoideus.

HYOÏDEUS, Hyoïdes.

HYOÏDIS QUARTUS MUSCULUS, Omo-hyoïdeus.

HYOSCY'AMUS, from *ὅς*, 'a swine,' and *κνῆπος*, 'a bean.' *Faba sni'lla*, Bengi, *Jusqui'amus*, *Hyoscyamus niger* *seu agre'stis*, *Apollin'aris*, *Alter'cum*, *Agloné*, *Altercan'yennu*, *Henbane*, *Poison Tobacco*, *Stinking nightshade*, (F.) *Jusqui'amus*, *Fève a Cochon*, *Hannebane*, *Patelée*, *Porcelot*. Ord. Solanaceæ. The leaves, *Hyoscy'ami Fol'ia* (Ph. U. S.), and seeds, *Hyoscy'ami Semen* (Ph. U. S.), are the parts used in medicine. Their odour is narcotic and peculiar; taste insipid and mucilaginous. The virtues are yielded to proof spirit. *Hyoscyamus* is narcotic, anodyne, antispasmodic, and slightly stimulant. It is used as a substitute for opium, where the latter disagrees; and is applied, externally, as a cataplasm in cancer and glandular swellings. Dose, gr. iij to x of the powder. An alkaloid was obtained from it by Brandes, *Hyoscyamine*, *Jusqui'amine*.

HYOSCYAMUS AGRESTIS, *Hyoscyamus*.

HYOSCYAMUS ALBUS, *White Henbane*, possesses similar virtues.

HYOSCYAMUS LUTEUS, *Nicotiana rustica* — h. Niger, *Hyoscyamus* — h. Peruvianus, *Nicotiana tabacum*.

HYOSCYAMUS SCOPOL'IA, *Scopol'ia atropoi'des*. The herb and root of this plant, which grows in Illyria, Hungary, Croatia, Gallizia, and Bavaria, are used in the same cases as belladonna. Dose of the powder, half a grain.

HYPACTICUS, Cathartic.

HYPÆMIA, from *ὑπο*, 'beneath,' and *αἷμα*, 'blood.' *Hypæ'mia*, *Hypohæ'mia*, *Oligæ'mia*, *Oligohæ'mia*, *Anæ'mia*. Deficiency of blood. — Andral. Also, extravasation of blood.

HYPÆMIA OCULI, *Hæmophthalmia*.

HYPAGOGÆ, Dejection.

HYPAGOGUS, Laxative.

HYPALEIM'MA, *Hypalim'ma*, *Hypaleip'tum*, *Hypalip'tum*, from *ὑπαλειψω*, 'I anoint.' An ointment or liniment to be rubbed or spread on a part.

HYPALEIPTRIS, *Hypaleiptrum*.

HYPALEIP'TRON, *Hypaleip'trum*, *Hypaleip'tris*, *Hypalip'trum*, *Speci'um*, *Spatha*. A sort of spatula for spreading ointments. — Hippocrates.

HYPALEIPTRUM, *Hypaleiptron*.

HYPALEIPTUM, *Hypaleimma*.

HYPALIMMA, *Hypaleimma*.

HYPALIPTRUM, *Hypaleiptron*.

HYPAMAURO'SIS, from *ὑπο*, 'under,' and *amaurosis*; *Amauro'sis imperfecta*. Imperfect amaurosis; *Morcamau'ro'sis*.

HYPANCO'NIUM, from *ὑπο*, 'beneath,' and *αγκων*, 'the elbow.' A pillow for supporting the elbow in surgical cases.

HYPAPOPLEX'IA, from *ὑπο*, 'under,' and *apoplexia*. An incomplete attack of apoplexy.

HYPATMISMUS, Fumigation.

HYPATMUS, Fumigation.

HYPAUCHE'NIUM, from *ὑπο*, 'under,' and *αυχην*, 'the neck.' A pillow or cushion for the neck.

HYPECCHORESIS, Dejection.

HYPECCHORETICUS, Laxative.

HYPELATUS, Cathartic, Laxative.

HYPENANTIOMA, Allopathy.

HYPENANTIOSIS, Allopathy.

HYPE'NĒ, *ὕπην*. The beard which grows under the chin, according to some. Also, the upper lip. — Vesalius. See *Mystax*.

HYPENETES, Adolescens.

HYPER, *ὑπερ*, 'above,' 'in excess.' Hence: HYPERACU'SIS, *Organo'ia*, *Hyperac'ia*, *Phthongodypho'ria*, from *hyper*, and *ακον*, 'audition.' Excessive sensibility of the organ of hearing.

HYPERADENO'SIS, *Hyperaden'oma*, *Hyper'troph'ia glandula'rum*, from *hyper*, and *αδην*, 'a gland.' Hypertrophy of a gland.

HYPERÆMIA, *Hyperhæ'mia*, *Hæmorrhæ'sis*, (F.) *Hyperhæ'mia*, *Angiohæ'mia*, from *hyper*, and *αἷμα*, 'blood.' Prenatural-accumulation of blood in the capillary vessels, more especially local plethora — *pletho'ra partiæ'lis*: congestion. — Andral. Various forms of hyperæmia are admitted by pathologists. — for example, the *active* or *sthenic*; as in the phlegmasiæ, — the *asthenic* or *passive*, from weakness of vessels; the *cadaveric*, or that which forms immediately before or after death; and the *hypostatic*, which occurs in depending parts.

HYPEREMIA ABDOMINIS, *Cæliæmia* — h. Activa, Inflammation — h. Capitis, *Cephalæmia* — h. of the Brain, *Stethæmia* — h. Cerebri, *Cephalæmia* — h. Pectoris, *Stethæmia* — h. Pulmonum, *Stethæmia* — h. of the Lungs, *Stethæmia* — h. Hepatic, *Hepatohæmia*.

HYPERÆMIC, *Hyperæ'micus*, *Hyperæ'mial*, *Conges'ted*, *Bloodshot*. Affected with hyperæmia.

HYPERÆSTHESIA, *Hyperæsthesis* — h. Lingue, *Hypergeusia* — h. Olfactoria, *Hyperosphresia* — h. Plexus cardiæci, *Angina pectoris*.

HYPERÆSTHES'IS, *Hyperæsthæ'sia*, *Oxyæsthæ'sia*, (F.) *Hypæcæsthesis*, from *hyper*, and *αἰσθησις*, 'the faculty of feeling.' Excessive sensibility, impressibility, or passibility.

HYPERÆSTHETICA, same etymon. Agents that are conceived to augment general sensibility — as strychnia, brucia, &c. Pereira.

HYPERANARRHOPHESIS, from *hyper*, and *αναρροήσις*, 'absorption.' Excessive absorption.

HYPERANTHERA MORINGA, *Guilandina moringa*.

HYPERANTHRAXIS, see *Cholera*.

HYPERAPH'IA, *Oxyaph'ia*, *Oxyaph'ia*, from *hyper*, and *αφν*, 'touch.' Excessive acuteness of touch.

HYPERAPHRODIS'IA, from *hyper*, and *Αφροδιτη*, 'Venus.' Excessive venereal desire.

HYPERASTHEN'IA, from *hyper*, and *ασθενεια*, 'weakness.' Excessive debility.

HYPERAUXE'SIS, from *hyper*, and *αυξησις*, 'augmentation.' *Hyperepid'osis*. Excessive increase or enlargement of a part, as *Hyperæuc'sis Iridis*, an excessive enlargement of the iris, so as to stop up the pupil.

HYPERBOL'IC (attitude), from *hyper*, and *βαλλω*, 'I throw.' 'Excessive.' Galen, by this term, designates certain extraordinary attitudes, in which the limbs and vertebral column are in a state of complete extension or flexion.

HYPERBO'REAN, from *hyper*, and *βορæας*, 'the north wind.' A race of men found at the extreme north of the two continents, in the vicinity of the polar circle. It includes the Thibetans, Ostiaks, Kamtschadales, Laplanders, Samoiedes, Esquimaux, &c.

HYPERBUL'IA, from *hyper*, and *βουλη*, 'will.' Ungovernable will or volition.

HYPERCARDIA, Heart, hypertrophy of the. HYPERCARDIOTROPH'IA, Heart, hypertrophy of the.

HYPERCATAPINO'SIS, from *hyper*, and *καταπινειν*, 'to sip up.' Excessive activity of absorption.

HYPERCATHAR'SIS, *Hyperine'sis*, *Hyperinos*, *Superpurgatio*, *Effrenitatio*, from *hyper*, and *καθαρσις*, 'purgation.' Superpurgation.

HYPERCENO'SIS, from *hyper*, and *κενωσις*,

'evacuation.' Excessive evacuation, as of blood, bile, &c.

HYPERCERASIS, Staphyloma of the cornea.
HYPERCERATOSIS, Staphyloma of the cornea.

HYPERCHOLIA, Polychochia.

HYPERCHROMA, from *hyper*, and *χρῶμα*, 'colour.' A red fleshy excrecence at the inner angle of the eye, near the caruncula lacrymalis, which sometimes grows very large, so as to separate the eyelids and interfere with their movements.

HYPERCHROMATOPSIA, *Hyperchromatopsia*, from *hyper*, *χρῶμα*, *χρωματος*, 'colour,' and *ὥψις*, 'vision.' A defect of vision, owing to which ideas of colour are attached to objects, which convey no such coloured impressions to a healthy eye. It is the antithesis to *achromatopsia*.

HYPERCINESIA, *Hypercinesia*, from *hyper*, and *κίνησις* (*kineu*, 'I move,') 'motion. Excessive motion.' Under the term *hypercinesis*, Romberg includes the spasmodic neuroses.

HYPERCINESIA NERVOSA, see Irritable—h. Uterina, Hysteria.

HYPERCINESIA GASTRICA, Hypochondriasis.

HYPERCONJUNCTIVITE, see Ophthalmia.

HYPERCORYPHOSIS, from *hyper*, and *κρυφή*, 'the vortex,' the extreme point of anything. The lobes of the liver and lungs. — Hippocrates.

HYPERCRINIA, *Hyperdiacrisis*, *Chymoze-mia*, from *hyper*, and *κρῖνω*, 'I separate.' A morbid increase in the quantity of the secretions.

HYPERCRISIS, same etymon. *Superexcretion*, *Superexcretion*. An excessive crisis, or evacuation; a flux.—A very violent, critical effort, or too copious critical evacuations.—Galen.

HYPERCYEMA, Superfetation.

HYPERCYESIS, Superfetation.

HYPERDERMATOSIS, *Hyperdermatoma*, *Hyperdermo-sis*, *Hyperdermo-ma*, from *hyper*, and *δέρμα*, 'skin.' Hypertrophy of the skin.

HYPERDIACRISIS, Hypercrinia.

HYPERDIURESIS, Diabetes.

HYPERDYNAMIA, Hypersthenia.

HYPERDYNAMIC, *Hyperdynamiceus*, from *hyper*, and *δυναμις*, 'strength.' Appertaining to or having the characters of hyperdynamia, or excessive strength—of the vital powers more especially.

HYPERECHEMA, see Exaggerated.

HYPERECHESIS, see Exaggerated.

HYPEREMESIS, *Hyperemesia*, *Vomitus profusus*; from *hyper*, and *εμεω*, 'I vomit.' Excessive vomiting.

HYPERÉMIE, Hyperæmia — h. du Cerveau, Cephaloæmia — h. Cérébrale, Cephaloæmia — h. du Foie, Hepatoæmia — h. de la Moëlle épinière, Hypermyeloæmia — h. des Poumons, Stethæmia.

HYPERENCÉPHALOTROPHIE; from *hyper*, *εγκεφαλον*, 'the encephalon,' and *τροφή*, 'nourishment.' Hypertrophy of the encephalon. — Piorry.

HYPERENCEPHALUS, (F.) *Hyperencephale*, from *hyper*, and *κεφαλή*, 'the head.' A monster whose excessive brain is situate in the skull.

HYPERENDOSMOSE, Inflammation.

HYPERENERGIA, from *hyper*, and *ενεργεια*, 'activity.' Excessive activity, as of the nervous system.

HYPEREPHIDROSIS, Ephidrosis.

HYPEREPIDOSIS, Hyperauresis.

HYPERERETHISIA, from *hyper*, and *ερεθίζω*, 'I excite.' Excessive irritability.

HYPERESIA, 'vaseoria,' 'a ministry.' This

word is sometimes applied to the organs;—when it means function.

HYPERESTHÉSIE, Hyperæsthesia.

HYPERETRIA, Midwife.

HYPERFIBRINATION, Hyperinosis.

HYPERGENESIS, from *hyper*, and *γενεσις*, 'generation,' (F.) *Hypergénésie*. The excess of formative power, which gives occasion to monstrosities by excess of parts. See Monster.

HYPERGEUSTIA, *Hypergustia*, *Hyperæsthesia* (*lingue*, *Oxygustia*, from *hyper*, and *γεύω*, 'taste.' Excessive sensibility of the organ of taste.

HYPERHÆMATOSIS, Inflammation.

HYPERHÆMIA, Hyperæmia.

HYPERHÉPATOTROPHIE, Hepatauxia.

HYPERHIDROSIS, Ephidrosis.

HYPERHO'RA, from *hyper*, and *ώρα*, 'time.' Premature development of the body, or of some part.

HYPERICUM BACCIFERUM, *Arbuscula gummifera* *Brazilien'sis*, *Cau-opia*. Ord. Hypericaceæ. A Brazilian tree, whose bark emits a juice, when wounded, which resembles gamboge. Hypericum is considered by Linnaeus to be derived from *υπερ*, 'over,' and *εικων*, 'an image or spectre,' from its having been fancied to possess the power of driving away evil spirits.

HYPERICUM OFFICINALE, II. perforatum — h. Officinarium. II. perforatum.

HYPERICUM PERFORATUM, *Hypericum*, II. officinale seu officinarum seu vulgare seu Virginicum, *Fuga Dæmonum*, *Androsæmum*, *Colerin*, *Perforated* or *Common St. John's Wort*, (F.) *Milepertuis ordinaire*. It is aromatic and astringent, and enters into a number of aromatic preparations; and, amongst others, into the *Fall-trancka*. It was, at one time, much used in dysentery, hemorrhages, &c. The juice is acrid. The Oil of St. John's Wort, Red oil, *Linimentum seu Oleum hyperici*, *Balsamum hyperici simplex*, is made by infusing 5iv. of the flowers in a quart of olive oil. It is applied to recent bruises.

HYPERICUM SAROTHRA, *Orange grass*, *Pine-weed*, indigenous, is applied in domestic practice, boiled, to contusions and sprains.

HYPERICUM VIRGINICUM, II. perforatum — h. Vulgare, II. perforatum.

HYPERIDROSIS, Ephidrosis.

HYPERINESIS, Hypercatarsis.

HYPERINOS, Hypercatarsis.

HYPERINO'SIS, *Ino'sis*, *Hyperplasma*, from *hyper*, and *ινωσις*, 'fibre.' The condition of the blood in which it contains an increase in the proportion of fibrin, *Hyper-fibrination*, *Super-fibrination*, a decrease of the corpuscles in proportion to the excess of fibrin, and an increase of the fat,—as in inflammation. In proportion to the increase of the fibrin and fat, and the decrease of the corpuscles, the whole solid residue will be diminished. Also, morbidly-increased muscular activity.—Siebenhaar.

HYPERLYMPHIA, (F.) *Hyperlymphie*; from *hyper* and *lymphica*. Excessive formation or accumulation of lymph.

HYPERMÉTROHÉMIE, Metro hæmia.

HYPERMNE'SIA, from *hyper*, and *μνησμαι*, 'I recollect.' Excessive memory.

HYPERMYELOHÆMIA, (F.) *Hypermie de la Moëlle Épinrière*, *Congestion sanguine rachidienne*, from *hyper*, *μυελος*, 'marrow,' and *αἷμα*, 'blood.' Hyperæmia of the spinal marrow.

HYPERNÉPHROTROPHIE, from *hyper*, *νεφρος*, 'kidney,' and *τροφή*, 'nourishment.' Hypertrophy of the kidney.

HYPERNEURIA, from *hyper*, and *νευρον*, 'a nerve.' Excessive nervous activity.

HYPERNEURO'MA, same etymon. Morbid development of the neurine or nervous masses.

HYPERO-PHARYNGEUS, Palato-pharyngeus.

HYPERO'A, from *hyper*, and *ων*, 'a high place.' The palatine arch,—the base of the cranium; the palate.

HYPEROITIS, *Inflammatio palati*, *Angina Palati'na*, (F.) *Inflammation du Palais*: from *hyperos*, 'the palate,' and *itis*, a suffix denoting inflammation of the velum palati.

HYPEROÖCHAS'MA, from *ὑπερω*, 'the palate,' and *χασμα*, 'an opening;' *Lycostoma*, *Palatum fissum*. Fissure of the palate.

HYPEROPSIA, Oxyopia.

HYPEROS, Pilon.

HYPEROSMIA, Hyperosphresia.

HYPEROSPHERE'SIA, *Hyperos'mia*, *Hyperæsthesia olfactoria*, *Olfactus acutus*, from *hyper*, and *σφρησις*, 'smell.' Excessive acuteness of smell.

HYPEROSTOSIS, Exostosis.

HYPERPATHIA, from *hyper*, and *παθος*, 'suffering.' Excessive sensibility in disease.

HYPERPHLEBO'SIS, from *hyper*, and *φλεβ*, 'a vein.' Too great development of the venous system; predominant venosity.

HYPERPHLEGMASIA, Hyperphlogosis.

HYPERPHLOGO'SIS, *Epiphlogo'sis*, *Hyperphlegmasia*, *Inflammatio peracuta*; from *hyper*, and *φλογωσις*, 'inflammation.' A high degree of inflammation.

HYPERPIRE'NIE, Mania.

HYPERPIMELE, see Polysarcia.

HYPERPLASMA, Hyperinosis.

HYPERPLEXIE, Ecstasis.

HYPERPNEUSTIA, Flatulence.

HYPERPRESBTIA, Presbytia.

HYPERSARCHIDIOS, Physconia.

HYPERSARCO'MA, *Hyperarco'sis*, *Hyperarctis*, from *hyper*, and *σας*, 'flesh.' A soft fungous excrescence, especially such as appears upon ulcerated parts.—A fungosity.

HYPERSARCOSIS, Excrecence, Fungosity, —h. Cordis, Heart, hypertrophy of the.

HYPERSARXIS, Hypersarcoma.

HYPERSPADIAS, Epispadias.

HYPERSPASMIA, Convulsion.

HYPERSPLÉNOTROPHIE, Splenoncus.

HYPERSPONGIA, Spina ventosa.

HYPERSTHENIA, *Hyperdyna'mia*, *Hyperzoödyna'mia*, *Status inflammatorius verus*; from *hyper*, and *σθενος*, 'strength.' Super-excitement. A morbid condition, characterized by over-excitement of all the vital phenomena.

HYPERSTHENIC, Stimulant.

HYPERSTHENICUS, Active, Stimulant.

HYPERTONTA, *hyper*, and *τονος*, 'tone.' Excess of tone in parts. It is opposed to atony.

HYPERTRICHOSIS: from *hyper*, and *τριξ*, *τριχος*, 'hair.' Excessive production of hair.

HYPERTROPHÆ'MIA, from *hyper*, *τροφη*, 'nourishment,' and *αἷμα*, 'blood.' A state in which the plastic powers of the blood are increased.

HYPERTROPHIA, Hypertrophy —h. Cerebri, Phrenauxe —h. Cordis, Heart, hypertrophy of the —h. Glandularum, Hyperadenosis —h. Hepatis, Hepatauxa —h. Intestinorum, Enterhypertrophia —h. Lienis, Splenoncus —h. Splenis, Spleoncus —h. Uteri, Metrauxa —h. Unguium, Onychauxa —h. Vesicæ urinariæ, Cystauxa.

HYPERTROPHIE, see Hypertrophy.

HYPERTROPHIE DU CŒUR, Heart, hypertrophy of the —h. *Centripète du Cœur*, Heart,

hypertrophy of the —h. *Concentrique du Cœur*, Heart, hypertrophy of the —h. *du Foie*, Hepatauxa —h. *de la Rate*, Splenoncus —h. *du Corps Thyroïde*, Bronchocele.

HYPERTROPHIED, see Hypertrophy.

HYPERTROPHIEN, see Hypertrophy.

HYPERTROPHOUS, see Hypertrophy.

HYPERTROPHY, *Hypertroph'ia*, from *hyper* and *τροφη*, 'nourishment.' The state of a part in which the nutrition is performed with greater activity; and which, on that account, at length acquires unusual bulk. The part thus affected is said to be *hypertrophied* or *hypertrophous*. *Hypertrophie* is used by Neligan, as *Hypertrophien* is by Simon, for an order of cutaneous diseases, characterized by hypertrophy of the derma, or epidermis, or both, or of the hair follicles.

HYPERURESIS, Enuresis —h. *Aquosa*, Diabetes.

HYPERURORRHÉE, Diabetes —h. *Saccharine*, Diabetes (mellitus.)

HYPERZOODYNAMIA, Hypersthenia.

HYPEX'ODOS, *ὑπεξόδος*, from *ὑπο*, 'beneath,' and *εξόδος*, 'passing out.' An alvine discharge or flux. —Hippocrates.

HYEZOCUS, Diaphragm, Pleura. See Hypozoma.

HYPHA, Texture.

HYPHEMA, Ecchymoma.

HYPHEMATO'SIS, *Hypohæmato'sis*, from *ὑπο*, 'under,' and *αἷματος*, 'sanguification.' Morbidly diminished hæmatosis. —*Sanguificatio debili'or*.

HYPHEMOS, Subcruentus.

HYPHE, Texture.

HYPHYDROS, Hydropic.

HYPINO'SIS, *Hypoplas'ma*, from *ὑπο*, 'under,' and *ισ*, *ινος*, 'fibre.' The condition of the blood in which the quantity of fibrin is frequently less than in health, or if it amounts to the usual quantity, its proportion to the blood corpuscles is less than in health: the quantity of corpuscles is either absolutely increased or their proportion to the fibrin is larger than in healthy blood; the quantity of solid constituents is also frequently larger than in health. Such is the condition of the blood in fevers, hemorrhages, and polyæmia, —Simon.

HYPNÆSTHESIS, Somnolency.

HYPNAGO'GIC, *Hypnagogicus*, from *ὑπνος*, 'sleep,' and *αγωγος*, 'a leader.' Leading to sleep. *Hypnagogic hallucinations* are those visions which occur when we are half asleep.

HYPNIA'TER, (F.) *Hypniatre*, from *ὑπνος*, 'sleep,' and *ιατρος*, 'a physician.' A name given to deluded or designing persons who have affirmed that they were able, during their 'magnetic sleep,' to diagnose disease and its appropriate treatment.

HYPNIC, *Hyp'nicus*, from *ὑπνος*, 'sleep.' An agent that affects sleep. —Pereira.

HYPNOBADES, Somnambulist.

HYPNOBADISIS, Somnambulism.

HYPNOBATASIS, Somnambulism.

HYPNOBATES, Somnambulist.

HYPNOBATESIS, Somnambulism.

HYPNOBATIA, Somnambulism.

HYPNO'DES, *ὑπνωδης*, from *ὑπο*, 'under,' and *υπνος*, 'sleep.' One in a state of slumber or somnolency.

HYPNODIA, Somnolency.

HYPNOLOG'ICA, *Hypnolog'icæ*. The part of hygiene which treats of sleep.

HYPNOL'OGY, *Hypnology*, from *ὑπνος*, and *λογος*, 'a discourse.' Same etymon. A treatise on sleep. The doctrine of sleep.

HYPNONERGIA, Somnambulism.

HYPNOPHOBIA, from *ὑπνος*, 'sleep,' and *φοβος*, 'dread.' Dread of sleep. Incubus.
HYPNOPŒUS, Somniferous.
HYPNOS, Sleep.
HYPNOSIS, from *ὑπνος*, 'sleep.' The suppression or production of sleep.
HYPNOSIS BIOMAGNETICA, Sleep, magnetic.
HYPNOTIC, Somniferous.
HYPNOTISM, see Magnetism, animal.
HYPNOTIZED, see Mesmerized.
HYPŌ, *ὑπο*, 'sub,' 'under,' in composition.
HYPŌ, Hypochondriasis.
HYPŌŒMA, Ecchymoma.
HYPŌŒMIA, Ecchymoma, Hypæmia.
HYPOBLEPH'ARUM, from *ὑπο*, 'under,' and *βλεφαρον*, 'eyelid.' Tamefaction under one or both eyelids. Also, an artificial eye, placed under the eyelids.
HYPOCAPNISMA, Suffimentum.
HYPOCAPNISMUS, Fumigation.
HYPOCARO'DES, *Subsoporatus*, from *ὑπο*, 'under,' and *καρος*, 'a heavy sleep.' One who is in a state approaching carus.—Hippocrates.
HYPOCATHAR'SIS, from *ὑπο*, 'under,' and *καθαρσις*, 'purgation.' Too feeble purgation. A word opposed to hypercatharsis.
HYPOCATHARTICUS, Laxative.
HYPOCAUSTUM, from *ὑπο*, 'under,' and *κατω*, 'I burn.' A name given to a stove, or any such contrivance, to sweat in. Also, a furnace in any subterraneous place, used for heating baths:—*ὑποκαυστον*, *Balnearium*, *Vaporarium*.
HYPOCEPHALĒ'UM, from *ὑπο*, 'under,' and *κεφαλη*, 'head.' A pillow for the head.
HYPOCERCHA'LEON, from *ὑπο*, and *κερχαλος*, 'hoarse.' Roughness of the fauces affecting the voice.—Hippocrates.
HYPOCHLORETUM SULPHUROSUM, Sulphur, chloride of.
HYPOCHLOROM'ELAS, *sub-pallidæ nigricans*, from *ὑπο*, *χλωρος*, 'green,' and *μελας*, 'black.' A term applied to one whose skin is pale, with a blackish hue.—Hippocrates, Galen.
HYPOCHOILION, Abdomen.
HYPOCHONDRE, *Hypochondrium*, *Subcartilagineum*, *Regio Hypochondriaca*, *Hypochondriac Region*, from *ὑπο*, 'under,' and *χονδρος*, 'a cartilage.' Each lateral and superior region of the abdomen is so called, because it is bounded by the cartilaginous margin of the false ribs, which forms the base of the chest. There is a right and a left hypochondrium.
HYPOCHONDRIAC, *Hypochondriacus*, *Hypochondriacal*, *Vapoury*, *Vapourish*, (F.) *Hypochondriacque*. Same etymon. Belonging to hypochondriasis. One labouring under hypochondriasis—hipped.
HYPOCHONDRIACISMUS, Hypochondriasis.
HYPOCHONDRIALGIA, Hypochondriasis.
HYPOCHONDRIAQUE, Hypochondriac.
HYPOCHONDRIASIS. Same etymon. *Alusis* seu *Hallucinatio hypochondriasis*, *Morbis hypochondriacus* seu *Resicicatorius* seu *Ructuosus*, *Malum hypochondriacum*, *Hypochondriacismus*, *Hypochondriacismus*, *Dyspepsia hypochondriasis*, *Pas'sio* seu *Affectio hypochondriaca*, *Anathymiasis*, *Hypercine'sis gastrica*, *Splenes*, *Melancholia nervæ* seu *statio'sa* seu *hypochondriaca*, *Suffocatio hypochondriaca*, *Morbis stituosus* seu *eruditiorum*, *Fomes ventriculi*, *Hypochondrism*, *Hypo*, *Hip*, *Spleen*, *Vapours English*, *Malady*, *Low Spirits*, (F.) *Hypochondrie*, *Maladie imaginaire* ou *Anglaise*, *Affectio vaporeuse*, *Vapours*. This disease is probably so called, from the circumstance of some hypochondriacs having felt an uneasy sensation in the hypochondriac regions. The disease seems really to be, as Pinel has classed it, a species of neurosis, and of mental alienation, which is observed in persons who in

other respects are of sound judgment, but who reason erroneously on whatever concerns their own health. Hypochondriasis is characterized by disordered digestion, without fever or local lesion; flatulence; horborygmi; extreme increase of sensibility; palpitations; illusions of the senses; a succession of morbid feelings, which appear to simulate the greater part of diseases; panics; exaggerated uneasiness of various kinds; chiefly in what regards the health, &c. Indigestion has usually been considered the cause of hypochondriasis. They are, unquestionably, much connected with each other: but there is every reason to believe, that the seat of the affection is really, though functionally, in the brain. The disease almost always appears at the adult age, most commonly in irritable individuals; and in those exhausted, or rather in the habit of being exhausted by mental labour, overwhelmed with domestic or public affairs, &c.

The treatment is almost entirely moral. The condition of the digestive function must, however, be accurately attended to.

HYPOCHONDRICTISMUS, Hypochondriasis.

HYPOCHONDRISM, Hypochondriasis.

HYPOCHOREMA, Excrement.

HYPOCHORESIS, Dejection, Defecation, Excrement.

HYPOCHORETICUS, Cathartic.

HYPOCHYMA, Cataract.

HYPOCHYROSIS, Barycoia.

HYPOCHYSIS HÆMATODES, Hæmophthalmia.

HYPOCISTIS, Cytinus.

HYPOCLEP'TICUM, from *ὑπο*, 'beneath,' and *κλεπτω*, 'I steal.' A chymical vessel, formerly used for separating oil from water.

HYPOCŒLIS, Palpebra inferior.

HYPOCŒLIUM, Abdomen.

HYPOCŒLUM, Hypocœilon, Palpebra inferior.

HYPOCOI'LON, *Hypocœlon*, *Hypocœlum*, *Hypocœylum*, from *ὑπο*, 'under,' and *κοilon*, 'a cavity.' A cavity situate under the lower eyelid. The lower eyelid itself.

HYPOCOPHOSIS, Barycoia, Deafness.

HYPOCRANI'UM (*ΑΠΟΣΤΕΜΑ*), from *ὑπο*, 'under,' and *κρανιον*, 'the cranium.' A collection of pus between the cranium and dura mater.

HYPOCRATES, Infirm.

HYPOCYLUM, Hypocœilon, Palpebra inferior.

HYPOCYSTEOTOMIA, see Lithotomy.

HYPODERIS, Epideris.

HYPODERMAT'OMY, *Hypodermatomia*, from *ὑπο*, 'under,' *derma*, 'the skin,' and *τομη*, 'incision.' The section of subcutaneous parts, as of tendons and muscles.

HYPODERMIC, *Hypodermicus*, (F.) *Hypodermique*, from *ὑπο*, 'under,' and *derma*, 'skin.' That which is under the skin,—as '*hypodermic medication*,' (F.) *Médication hypodermique*; that which consists in the application of remedies—as morphia—under the skin.

HYPODERMIS, Clitoris, Epideris.

HYPODERRHIS, Epideris.

HYPODESMA, Bandage.

HYPODYNAMIC, Adynamic.

HYPODYN'IA, from *ὑπο*, 'under,' and *δυνη*, 'pain.' *Dolor mitis seu lenis*. A slight pain.

HYPOGALA, Hypopyon.

HYPOGASTRAL'GIA, from *ὑπογαστριον*, 'the hypogastrium,' and *αλγος*, 'pain.' Pain in the hypogastrium.

HYPOGASTRIC, *Hypogastriacus*. Relating or belonging to the hypogastrium.

HYPOGASTRIC ARTERY, *A. Ili'aca inter'na*, *A. Ili'aca post'rior*, *A. pelvienne*—(Ch.), is the more internal of the two branches into which the primary iliac divides. It descends into the cavity

of the pelvis, and gives off a considerable number of branches, which arise, at times, separately; at others, by common trunks. These branches are, 1. The *posterior*, i. e. the ilio-lumbar arteries, lateral, sacral, and gluteal. 2. The *anterior*, i. e. the umbilical, vesical, and obturator. 3. The *internal*,—the middle hemorrhoidal arteries, uterine, and vaginal in women. 4. The *inferior*, i. e. the *ischiatric arteries*, and internal pudic.

HYPOGASTRIC GANGLION. A large nervous ganglion, described by Dr. Robt. Lee as seated on each side of the cervix uteri, immediately behind the uterus; which receives the greater number of the nerves of the hypogastric and sacral plexuses, and distributes branches to the uterus, vagina, bladder, and rectum.

HYPOGASTRIC OPERATION OF LITHOTOMY, (F.) *Taille hypogastrique.* The high operation, or that practised above the pubes.

HYPOGASTRIC PLEXUS, Plexus sous-mésentérique of Winslow, is situate at the lateral and posterior parts of the rectum and the *bas fond* of the bladder. It is formed by the sacral nerves and the inferior mesenteric plexus, and gives off numerous filaments, which accompany the arteries that pass to the rectum and genital organs.

HYPOGASTRIC VEIN furnishes nearly the same branches.

HYPOGASTRION, Abdomen, Hypogastrium.

HYPOGASTRIUM, Hypogastrium, from 'υπο, 'under,' and γαστήρ, 'the stomach or belly;' *Εγρον, πτρον, Venter imus, V. parvus, Aquiliculus, Simen, Rumen;* the lower part of the abdomen. The *Hypogastric region, Regio hypogastrica seu hypogastria*, which extends as high as three fingers' breadth beneath the umbilicus, is divided into three secondary regions—one middle or *pubic*, and two lateral or *inguinal*.

HYPOGASTROCELĒ, from 'υπο, 'under,' γαστήρ, 'the stomach or belly,' and κηλη, 'a tumour.' Hernia in the hypogastric region, occurring through the separated fibres of the lower part of the linea alba. See *Hernia, hypogastric*.

HYPOGASTRODIDYMIUS, Didymus Symphyohypogastrius, Ichiopoda'ges, from 'υπο, 'under,' γαστήρ, 'the belly,' and διδύμος, 'a twin.' A monstrosity in which twins are united by the hypogastrium.—Gurlt.

HYPOGASTRORIXIS, Eventration.

HYPOGLOSSA, Hypoglossides.

HYPOGLOSSAL NERVE, Hypoglossus.

HYPOGLOSSIA, Hypoglossides.

HYPOGLOSSIADENITIS, from 'υπο, 'under,' γλωσσα, 'tongue,' αδην, 'a gland,' and itis, denoting inflammation. *Inflammatio glandularum sublingualium.* Inflammation of the sublingual gland.

HYPOGLOSSIDIA, Hypoglossides.

HYPOGLOSSIS, Hypoglossium, Hypoglossis, Hypoglossia, from 'υπο, 'under,' and γλωσσα, 'tongue.' The under surface of the tongue; anything under the tongue; and hence, *Ranula*.

HYPOGLOSSIUM, Hypoglossis, Ranula.

HYPOGLOSSUM, Ruscus hypoglossum, Ranula.

HYPOGLOS'SUS, from 'υπο, 'under,' and γλωσσα, 'the tongue.' That which is under the tongue.

HYPOGLOSSUS, Lingualis Me'dius, Hypoglossal or Lingual or Gustatory Nerve, (F.) Nef Hypoglossæ ou Grand Hypoglosse, Hypoglossien (Ch.), is the ninth pair of nerves, *Par nomen, Nervus hypoglossus seu loquens*, of many anatomists. It arises by ten or twelve very fine filaments from the grooves, which separate the corpora pyramidalia from the C. olivaria; issues from the cranium by the foramen condyloideum anterius; and divides, near the angle of the jaw, into two

branches; the one, the *cervicalis descendens* or *descendens noni*. It forms, with the cervical plexus, a large anastomotic arch, and furnishes branches to several of the muscles of the neck. The other, the *lingual branch*, is the continuation of the principal trunk, and gives its numerous filaments to the muscles of the tongue and pharynx. The ninth pair communicates motion to the muscles to which it is distributed.

HYPOGLOTTIA, Hypoglossis, Hypoglossides, HYPOGLOTTIDES, (PILULÆ,) Hypoglossia, Hypoglossia, Hypoglossis, Hypoglossis, Hypoglossis, Pilule sublinguales. Pills placed under the tongue to dissolve there.

HYPOGLOTTIS, Hypoglossis, Ranula.

HYPOGLUTIS, from 'υπο, 'under,' and γλουτος, 'the nates.' The lower and projecting part of the nates.—Gorræus.

HYPOGNATHADEN, Submaxillary gland.

HYPOGNATHADENITIS, Hypoglossadenitis, from *hypognathaden*, the submaxillary gland, and *itis*, denoting inflammation. Inflammation of the submaxillary gland.

HYPOHÆMA, from 'υπο, 'under,' and αιμα, 'blood.' Effusion of blood into the chambers of the eye. It is generally absorbed in time.

HYPOHÆMATOSIS, Hypohæmatosis.

HYPOHÆMIA, Hypæmia.

HYPOLEPSIOMANIA, Melancholy.

HYPO'MIA, from 'υπο, 'under,' and ωμος, 'the shoulder.' That which lies under the shoulder, as the axilla.

HYPOMORIA, from 'υπο, 'under,' and μωρια, 'folly.' Slight imbecility of mind, or slight delirium.

HYPONARTHÉCIE, Hyponarthecia, from 'υπο, 'under,' and αρθής, 'a splint.' A term used by M. Mayor for his mode of treating fractures by position only,—the limb resting upon a properly cushioned board or splint.

HYPONEU'RIA, from 'υπο, 'under,' and νευρον, 'a nerve.' Morbidly diminished nervous energy.

HYPON'OMOS, Ulcus subtus depas'cens, from 'υπο, 'under,' and νεμω, 'I feed.' A deep fistula or ulcer.

HYPON'YCHON, from 'υπο, 'under,' and ονχ, 'the nail.' Ecchymo'ma Hyponychon. Effusion of blood under a nail.

HYPOPATHIA, Subaffectio, from 'υπο, 'under,' and παθος, 'disease.' A disease of a slight character.

HYPOPATUS, Dejectio.

HYPOPE'DIUM, from 'υπο, 'under,' and πους, 'the foot.' A cataplasm for the sole of the foot.

HYPOPH'ASIS, from 'υπο, 'under,' and φαινω, 'I appear.' The state of the eyes in which the white only is seen through the opening of the eyelids.—Hippocrates.

HYPOPHLEGMASIA, Subinflammatio.

HYPOPH'ORA, from 'υπο, 'under,' and φερω, 'I carry;' Ulcus sinu'eum seu fistulo'eum. A fistulous ulcer.—Galen. A dejection.

HYPOPTHALMIA, Hypopyon.

HYPOPTHAL'MION, from 'υπο, 'under,' and οφθαλμος, 'the eye.' That part under the eye where oedema generally commences in chronic diseases and in cachexia.—Hippocrates.

HYPOPHYSIS, Cataract, Epigone—h. Cerebri, Pituitary gland.

HYPOPITYS LANUGINOSA, from 'υπο, 'under,' and πυρς, 'a pine tree,' from the place of its growth. American Pine-sap, False Beechdrops, Birds' Nest. Indigenous; flowering from June to August. Order, Ericacææ. Used as a nervine in the form of the powdered root.

HYPOPIUM, from 'υπο, 'under,' and πιω,

'eye.' The part of the face under the eye:—a black eye.

HYPOPIUM OS, Male os.

HYPOPLASMA, Hypinosis.

HYPOPLASTE'MA, from 'υπο, 'under,' πλαστικός, 'formative,' and 'αἷμα, 'blood.' Diminished plasticity of the blood.

HYPOPLEURIUS, Pleura.

HYPOPO'DIA, *Supplanta'lia*, *Suppeda'nea*, from 'υπο, 'under,' and ποῦς, 'the foot.' Remedies, as sinapisms, which are applied under the foot.

HYPOP'YON, *Hypop'yum*, *Hypophthal'mia*, *Pyophthal'mia*, *Emphy'sis* seu *Diapye'sis* seu *Abscessus Oculi*, *Oculus purulent'us*, *Linell'a*, *Hypopy'ala*, *Hypopyum lacteum*, *Pys'sis*, *Parop'sis* *Staphylo'ma purulentum*; from 'υπο, 'under,' and πύον, 'pus,' because the pus is under the cornea. This name has been given to small abscesses between the laminae of the cornea, as well as to different purulent collections in the chambers of the eye; hence, some pathologists have distinguished *Hypop'yon* of the chambers from *Hypop'yon* of the cor'nea. In abscesses of the chambers, the purulent matter is mixed with the aqueous humour, which it renders turbid; and is deposited particularly at the lower part of the eye; forming behind the cornea a kind of whitish crescent, that rises more or less before the pupil, and closes it entirely or in part. It requires the use of local and general antiphlogistics, and sorbefacients. At times, it is necessary to puncture the cornea and evacuate the pus.

Hypopion was used by Galen to signify "an effusion of blood under the eyelids and tunica conjunctiva, or under the lower eyelid."—Blancard.

HYPOPYUM LACTEUM, Hypopyon.

HYPORIN'TON, *Hyporrhini'um*, from 'υπο, 'under,' and ριν, 'the nose.' That part of the beard which grows beneath the nose. The mustaches. Also, the upper lip.

HYPORRHYSIS, Defluxion, Prolapsus.

HYPOSA'PRUS, *Subputris*, *Putres'cens*. Growing putrid; slightly putrid. Game, of various kinds. (F.) *Viandes faisandées*, is, at times, preferred in an incipient state of decomposition, which certainly renders it more digestible.

HYPOSAR'CA, from 'υπο, 'under,' and σαρξ, 'flesh,' *Hyposarcid'ius*, *Anasarca*. In the Nosologies of Linnæus and Cullen, it is synonymous with *Physconia*.

HYPOSARCIDIDIUS, *Anasarca*, *Hyposarca*.

HYPOSARCO'SIS, from 'υπο, 'under,' and σαρκωσις, 'a fleshy growth.' A small, fleshy growth:—a wart.

HYPOSTIAGON, Maxillary bone, inferior.

HYPOSTIAGONARTHRI'TIS, from 'υπο, 'under,' σιαγών, 'the jawbone,' ('lower jawbone,') αρθρον, 'a joint,' and ιτις, denoting inflammation; *Inflammatio articuli maxillæ inferioris*. Inflammation of the joint of the lower jaw.

HYPOSIALADEN, Submaxillary gland.

HYPOSIALADENTIS, Hypogathadenitis.

HYPOSPA'DIA, *Hypospadi'asis*, from 'υπο, 'under,' and σπας, 'I draw.' A malformation, in which the canal of the urethra, instead of opening at the apex of the glans, terminates at its base, or beneath the penis, at a greater or less distance from the symphysis pubis. When the orifice of the urethra is very near the root of the penis, the scrotum is divided, as it were, into two great labia; and this malformation has often been taken for a case of hermaphroditism. Hypospadia is ordinarily incurable; and an idea has been entertained that it is the cause of impotence. It is not exactly so; but it renders impregnation less probable.

HYPOSPA'DIAS, *Hypospadi'us*, *Hypospad'icus*. One affected with hypospadia.

HYPOSPADIASIS, Hypospadia.

HYPOSPATHIS'MUS, from 'υπο, 'under,' and σπαθιον, 'a ladle or spatula.' A surgical operation, with the ancients, in cases of severe 'defluxion to the eyes,' which consisted in incisions made in the scalp down to the bone, and the insertion of an instrument called a *spath'ion* or *hypospathi'ster* to enable the intermediate parts to be divided. The precise character of the operation is not well described, and it is no longer known.—Paulus of Ægina.

HYPOSPATHISTER, see Hypospathismus.

HYPOSPHAG'MA, from 'υπο, 'under,' and σφαζω, 'I kill.' The coagulated blood, which is collected when an animal is killed and used for food. Also, an effusion of blood, especially under the conjunctiva.

HYPOSPHAGMA, Hemophthalmia.

HYPOSTAPHYLE, Staphyloedema.

HYPOSTAPHYLITIS, see Staphylitis.

HYPOS'TASIS, from 'υπο, 'under,' and στας, 'the act of placing.' A morbid deposition in the body. Sediment.

HYPOSTAT'IC, *Hypostat'icus*, from 'υπο, 'under,' and στας, 'stagnation.' Relating to hypostases, sediments or depositions.

HYPOSTATIC HYPE'RMIA. A congestion of blood in the vessels of a part, caused by its depending position.

HYPOSTEMA, Sediment.

HYPOSTHENIA, Debility.

HYPOSTHENIC, Contrastimulant.

HYPOSTROPHE, 'υποστροφή, 'change of position,' from 'υπο, and στρέφω, 'I turn.' Act of a patient turning himself. Also, a relapse or return of a disease.—Hippocrates, Foësius.

HYPOSTYP'TIC, *Hypostyp'ticus*, from 'υπο, 'under,' and στυψις, 'constriction.' A feeble astringent.

HYPOSYPHILIS, from 'υπο, 'under,' and *Syphilis*. A mild form of syphilis.

HYPOTH'ENAR, *Sub'vola*, from 'υπο, 'under,' and *Sevap*, 'the palm of the hand or sole of the foot.' *Hypothenar Em'ince*. The fleshy projection of the palmar surface of the hand, which corresponds with the little finger, and is supported by the fifth metacarpal bone. This eminence is formed of four muscles: the *Palmaris brevis*, *Adductor seu Flexor brevis minimi digiti*, and *Opponens minimi digiti*. The name *Hypothenar* has also been given to different muscles of the hand. The *Hypothenar minimi digiti* of Riolan comprehended the *Adductor*, *Flexor brevis* and *Opponens minimi digiti*; and his muscle, *Hypothenar pollicis*, corresponded to the *Adductor* and a portion of the *Flexor brevis pollicis*. Winslow called the *Adductor minimi digiti*, muscle *petit hypothenar* ou *hypothenar du petit doigt*.

HYPOTHENAR MINIMI DIGITI, *Flexor parvus minimi digiti*—h. Minor metacarpeus, *Abductor minimi digiti*—h. Riolani, *Flexor parvus minimi digiti*.

HYPOTHETON, Suppository.

HYPOTHYMIAMA, Fumigation.

HYPOTHYMIASIS, Fumigation.

HYPO'TIUM, (*Emplastum*;) from 'υπο, 'under,' and ος, 'the ear.' *Emplastum auriculæ*; a plaster applied behind or under the ear.

HYPOTROPHE, Relapse.

HYPOTROPH'IA, *Hypot'rophy*, from 'υπο, 'under,' and τροφη, 'nourishment.' Scanty nourishment or nutrition. Atrophy.

HYPOTROPIASMUS, Relapse.

HYPOTRYGUS, Feulent.

HYPOUCANNA, Ipecacuanha.

HYPOUTRION, Abdomen.

HYPOXÉMIE, Asphyxia.

HYPOXYS ERECTA, *Stargrass*; from 'υπο, 'beneath,' and ὄξυς, 'sharp,' because the pod is acute at the base [?]; indigenous; Order, Amarillidacæ. The root is eaten, and has been used as a vulnerary; and in chronic ulcers and agues.

HYPOZO'MA, from 'υπο, 'under,' and ζωννυμι, 'I bind round,' *Membrana succinigena*. A membrane or septum, as the mediastinum, diaphragm, &c.

HYPOCRAS, Claret.

HYPSELOGLOSSUS, Basioglossus.

HYPSILODES, OS, Hyoides os.

HYPSILOGLOSSUS, Hyoglossus.

HYPSONOSUS, Puna.

HYPSONPHONUS, from ὑψος, 'high,' and φωνή, 'voice.' One who has a clear high voice.

HYPSSOSIS, Sublimation.

HYPTIASMA, Supination.

HYPTIASMOS, from ὑπτιάζω, 'I lie with the face upwards.' Lying in a supine posture. Also, inversion of the stomach, as in nausea, regurgitation, or vomiting.—Hippocrates, Galen.

HYPULUS, from 'υπο, and οὐλη, 'cicatrix.' Imperfectly cicatrized.

Ulcera hypu'la. Ulcers healed at the top, but not at the bottom.

HYRA'CEUM. A substance found in the Cape Colony, which Thunberg and other travellers mistook for a kind of bitumen; but, according to Dr. Pappe, it is obtained from the urine of the *Klipdas* or *Hyrax Capensis*, which, when passed, is thick and of a glutinous nature. The animal is in the habit of evacuating the urine at one spot, where its aqueous parts evaporate in the sun—the more tenacious adhering to the rock and hardening.

In smell, and medical properties, it most resembles castor, which, according to Dr. Pappe, it may replace. It is used by the Cape farmers in nervous and spasmodic affections.

HYRAX CAPENSIS, see Hyraceum.

HYRTOCHEILIDES, Labia pudendi.

HYSOPE, Hyssopus.

HYSSOP, Hyssopus—h. Hedge, Gratiola officinalis.

HYSSOPTES. Ancient name of a wine, of which hyssop was the chief ingredient, and which Dioscorides extolled in chronic inflammation of the chest. It was regarded as diuretic and emmenagogue.

HYSSO'PUS, from the Hebr. Azob, *Cassia alba*, *Hyssopus officinalis*, Common hyssop, (F.) *Hyssope*. It has been chiefly used as an aromatic, stimulant, and pectoral, in the form of infusion.

HYSTERA, 'vutera, 'vutera, 'the uterus.' Hence:

HYSTERA, Secundinas.

HYSTERALGIA, *Hysterodyn'ia*, (F.) *Fortraiture*, from 'vutera, 'the uterus,' and αλγος, 'pain.' Pain in the uterus. Irritable uterus, (F.) *Névrologie de l'utérus*. Hippocrates uses the epithet *Hysteralgæ*, *vuteralgyn*, for any thing that excites uterine pain; and, especially, [?] for vinegar.

HYSTERALGIA CATARRHALIS, Metrorrhœma—h. Galactica, Phlegmatia alba dolens—h. Lochialis, Dyslochia—h. Rheumatica, Metrorrhœma.

HYSTERAPOPNIXIS, Angone.

HYSTERATRE'PSIA, *Atréto-me'tria*, from 'vutera, 'the uterus,' and ἀτρητος, 'imperforate.' Imperforation of the os uteri.

HYSTERELCO'SIS, from 'vutera, 'uterus,' and ἔλκος, 'an ulcer.' *Uteri ezulceratio*, *Uteri ulcus*. Ulceration of the uterus.

HYSTERELOSIS, Hysteroloxia.

HYSTEREMPHYSEMA, Physometra.

HYSTERIA, *H. vaga*, *Hystericismus*, *Hysteria-*

mus, *Hysterias*, *Hysteri'asis*, *Hysteropath'ia*, *Hypocine'sia uteri'na* seu *hyste'ria*, *Uteri adscensus*, *Suffocatio hyster'ica* seu *uteri'na* seu *mulierum*, *Asthma uteri*, *Præfocatio materi'cis* seu *uteri'na*, *Synap'sia hysterica*, *Malum hyster'icun* seu *hysterico-hypochondri'acum*, *Vapo' res uteri'ni*, *Affectio* seu *Passio hyster'ica*, *Morbus hyster'icus*, *Strangulatio uteri'na* seu *Vulva*, *Vapours*, *Hyster'ics*, *Hyster'ic fit*, *Mother*, (F.) *Hystérie*, *Mot de Mère*, *Maladie imaginaire*, *Passion hystérique*, *Suffocation utérine*, *Étranglement*, *Épilepsie utérine*, *Vapeurs*, *Maux de Nerfs*, from 'vutera, 'the uterus.' A species of neurosis, classed amongst the spasmi by Sauvages and Cullen, and in the *Névroses de la génération*, by Pinel. It received the name of *hysteria*, because it was reputed to have its seat in the uterus. It generally occurs in paroxysms; the principal characters of which consist in alternate fits of laughing and crying, with a sensation as if a ball set out from the uterus and ascended towards the stomach, chest, and neck, producing a sense of strangulation. If the attack be violent, there is, sometimes, loss of consciousness (although the presence of consciousness generally distinguishes it from epilepsy) and convulsions. The duration of the attacks is very variable. It appears to be dependent upon irregularity of nervous distribution in very impressible persons, and is not confined to the female; for well-marked cases of hysteria are occasionally met with in men. During the fit,—dashing cold water on the face, stimulants applied to the nose, or exhibited internally, and antispasmodics, form the therapeutical agents. Exercise, tranquillity of mind, amusing and agreeable occupations constitute the prophylactics. See Mania, dancing.

HYSTERIA CATALEPTICA, Catalepsy—h. *Vaga*, *Hysteria*.

HYSTERIAS, *Hysteria*.

HYSTERIASIS, *Hysteria*.

HYSTERICA, see *Hysterical*.

HYSTERICAL, *Hyster'icus*, *Va'pourish*, *Va'pouring*, (F.) *Hystérique*. Same etymon as hysterical. Relating to or affected with hysteria.

The word *hysterica* was used by Martial for a female affected with nymphomania or with strong sexual desires.

HYSTERICISMUS, *Hysteria*.

HYSTERICUS, *Hysteria*.

HYSTÉRIE, *Hysteria*.

HYSTERISMUS, *Hysteria*.

HYSTERITES, Hydrometra.

HYSTERITIS, Metritis, Hydrometra.

HYSTEROCARCINOMA, Metrocarcinoma.

HYSTEROCE'LE, from 'vutera, 'the womb,' and κηλη, 'hernia.' *Metroce'le*, *Her'nia uteri*, *Hernia of the womb*. This is a rare disease. The womb may protrude through the inguinal or the crural canal, or through the lower part of the linea alba.

HYSTEROCELE NUDA, Prolapsus uteri.

HYSTEROCOLICA, Colica uterina.

HYSTEROCNESMUS, from 'vutera, 'uterus,' and κνησμος, 'itching.' Pruritus of the uterus or genitals.

HYSTEROCRANUM, Occiput.

HYSTEROCYESIS, Pregnancy.

HYSTERO-CYSTIC, *Hysterocys'ticus*, from 'vutera, 'the uterus,' and κυστις, 'the bladder.' Relating to the uterus and bladder. Some authors have called *Hystero-cystic Retention of urine*, that which is caused by the compression of the bladder by the uterus, during pregnancy.

HYSTERO-CYSTOCE'LE, from 'vutera, 'the womb,' κυστις, 'the bladder,' and κηλη, 'a tumour.' Hernia of the uterus complicated with displacement of the bladder.

HYSTERODYNIA, Hysteralgia.

HYSTERŒDEMA, Hydrometra.

HYSTEROID, *Hystero'des*, *Hysteroi'des*, from *hysteria*, and *eidōs*, 'resemblance.' Resembling hysteria:—as a hysteroid disease, symptom, &c.

HYSTEROLŒGY, *Hysterolog'ia*, from *ὑστέρα*, 'the uterus,' and *λόγος*, 'a description.' A treatise on the sound and morbid uterus.

HYSTEROLOX'IA, *Hysterelo'sis*, *Obliquitas seu Situs obliquus uteri*, *Flexio seu Versio incomple'ta seu Inclina'tio uteri*, *Metrolox'ia*, *Metrocamp'sis*, *Metroscolio'ma*, *Uterus obliquus seu Inclina'tus*; from *ὑστέρα*, 'the uterus,' and *λόγος*, 'oblique.' An oblique position of the uterus, occurring during pregnancy. Anteversion of the uterus, *Hysterolox'ia anterior*, *Anteversio seu Prona'tio uteri*, *Venter propendens*; and Retroversion of the uterus, *Hysterolox'ia posterior*, *Retrover'sio uteri*, *Reflex'io uteri completa*, are varieties.

HYSTEROMALA'CIA, *Hysteromalaco'sis*, *Hysteromalaco'ma*, *Metromalaco'sis*, *Metromalaco'ma*, *Mulaco'sis uteri*, *Emolli'ties uteri morbose*, *Putrescent'ia u'teri grav'idi*, from *ὑστέρα*, 'the uterus,' and *μαλακία*, 'softness.' Softness of the uterus during pregnancy, which renders it liable to rupture in labour.

HYSTEROMALACOMA, *Hysteromalacia*.

HYSTEROMANIA, *Nymphomania*.

HYSTEROMETER, see Sound, uterine.

HYSTEROMOCHLIUM, *Lever*.

HYSTERON'CUS, from *ὑστέρα*, 'the uterus,' and *ογκος*, 'tumour.' *Metron'cus*, *Hysterophy'ma*, *Metrophy'ma*, *Tumor uteri*. A tumour of the uterus.

HYSTERO-PARALYSIS, *Hysterople'gia*, from *ὑστέρα*, 'the uterus,' and *παράλυσις*, 'paralysis.' Paralysis of the uterus.

HYSTEROPATH'IA, from *ὑστέρα*, 'the uterus,' and *πάθος*, 'suffering.' Disease or suffering in the uterus. Also, hysteria, and deutero-pathia.

HYSTEROPHYMA, *Hysteroncus*.

HYSTEROPHYSE, *Physometra*.

HYSTEROPHYSIS, from *ὑστέρα*, 'the uterus,' and *φύση*, 'wind.' Distension of the uterus with air: *Emphyse'ma u'teri*.

HYSTEROPLEGIA, *Hysteroparalysis*.

HYSTEROPNIX, *Angene*.

HYSTEROPOLYPUS, *Metrotypus*.

HYSTEROPHOBIA, *Physometra*.

HYSTEROPTOSIS, from *ὑστέρα*, 'the womb,' and *πτωσις*, 'fall,' *Prolapsus uteri*. Also, *Inversio*

uteri, *Anastrophē uteri*. In a general sense, a protrusion of any part of the genital organs or of excrescences from them into the genital passages; *Adoptosis*.

HYSTEROPTOSIS, *Prolapsus uteri*—h. *Vagina prolapsus*, *Prolapsus vaginae*.

HYSTERRORRHAGIA SANGUINEA, *Metrorrhagia*.

HYSTERRORRHESIS, *Uterus, rupture of the*. **HYSTERRORRHŒA**, *Metrorrhagia*—h. *Mucosa*, *Leucorrhœa*.

HYSTERSALPINX, *Tube*, *Fallopian*.

HYSTEROSCIRRHUS, *Metroscirrhus*.

HYSTEROSCOPE, from *ὑστέρα*, 'the uterus,' and *σκοπεω*, 'I view.' A metallic mirror, used in inspecting the state of the os uteri for throwing the rays of a taper to the bottom of the speculum uteri.—Colombat de l'Isère.

HYSTEROSPASM'US, from *ὑστέρα*, 'the womb,' and *σπασμος*, 'spasm.' Spasm of the uterus. Also, hysterical spasm.

HYSTEROSTOMA, *Os uteri*.

HYSTEROSTOMAT'OMUS, from *ὑστέρα*, 'the womb,' *στομα*, 'orifice,' and *τεμνω*, 'to cut.' An instrument invented by Coutouly for dividing the os uteri, when it is important to deliver immediately, as in cases of convulsions.

HYSTEROSTOMIUM, *Os uteri*.

HYSTERO SYNZISIS, *Metrosynzisis*.

HYSTEROTOME, *Hysterotomus*.

HYSTEROTOM'IA, *Hysterotomy*. Same etymon as *Hysterotomus*. Cæsarean section. Also, dissection of the uterus.

HYSTEROTOMOTOCIA, *Cæsarean section*.

HYSTEROT'OMUS, *Hysterotome*, *Metrot'omus*, *Me'trotome*, from *ὑστέρα*, 'the womb,' and *τεμνω*, 'to cut.' An instrument for dividing the womb through the vagina. It is a kind of *Bistouri caché*, and is intended to divide the cervix and os uteri.—*Hysterostomat'omus*.

HYSTEROTRISM'US, from *ὑστέρα*, 'the womb,' and *τριζω*, 'I gnash.' Spasmodic contraction of the uterus, especially of its mouth.

HYSTREMPHYSEMA, *Physometra*.

HYSTRIASIS, *Hystriasis*.

HYSTRIC'ASIS, *Hystri'asis*, *Hystrix*, *Hystriasis'mus*, *Cerostro'sis*, *Tricho'sis seto'sa*, from *ὑστρίς*, *ὑστρίγος*, 'a hedgehog or porcupine.' A disease of the hairs, in which they stand erect like the quills of the porcupine.

HYSTRICISMUS, *Hystriasis*.

HYSTRIX, *Hystriasis*.

HYVES, *Hives*.

I.

IAMA, *Medicament*.

IAMATOLOGY, *Materia medica*.

IAMATOSYNTAXIOLOGIA, see *Prescription*.

IAMATOSYNTAXIS, see *Composition*.

IAMATOTAXIOLOGIA, see *Prescription*.

IASIMOS, *Curable*.

IASIS, *Curation*.

IATERIA, *Medicina*.

IATERIUS, *Medicinal*.

IATRALEIP'TICE, *Iatralipticē (Ars)*, *Iatraleptic* or *Iatralip'tic method*; from *ιατρος*, 'a physician,' and *αλειψω*, 'I anoint.' The method of treating diseases adopted by the *Iatraleptes*,—that is, by friction chiefly.

IATRALEP'TES, *Iatralaip'tes*, *Iatralip'ta*,

Iatraleip'tes, *Med'icus Unguenta'rius*. One who treats diseases by unguents, frictions, and by external means generally. *Prodicus*, a disciple of *Æsculapius*, was the chief of the *Iatraleptes*.

IATRALIPTES, *Iatraleptes*.

IATRALIPTICE, *Iatraleptice*.

IATREUSIOLOG'IA, from *ιατρειός*, 'the exercise of the art of healing,' and *λόγος*, 'a description.' The doctrine of the exercise of the healing art.—*Reil*.

IATREUSIS. The exercise of the healing art. **IATRIA**, *Medicina*.

IATRICE, *Medicina*.

IATRICUS, *Medical*; see *Digitus*.

IATRINĒ, *Medicina*. Also, a female practitioner of medicine; a midwife.

IATRION, *ιατρίον*, *Iatre'on*, *Tà'tron*, *Iet'ri'on*, *Medicatriona*. The house or office of a physician or surgeon. Also, the physician's fee or honorarium.

IATROCHEMIA, Chymiatría.

IATROCHYMIA, Chymiatría.

IATROGNOMICA, *Iatrognom'icé*; from *ιατρος*, 'a physician,' and *γινωσκω*, 'I know.' A knowledge of medical objects.—Hufeland.

IATROLEPTES, *Iatroleptes*.

IATROLOG'IA, *Iatrol'ogy*, from *ιατρος*, 'a physician,' and *λογος*, 'a description.' A treatise on physio and physicians.

IATROMANTIS, from *ιατρος*, 'a physician,' and *μαντεια*, 'divination.' A physician who is a soothsayer.

IATRO-MATHEMATICAL PHYSICIANS, *Iatromathem'ici*. Mechanical physicians.

IATROMECHANICI, Mechanical physicians.

IATRON, *Iat'ron*.

IATROPHYSICS, Physics, medical.

IATROS, *Iat'ros*.

IATROSOPHIST'IA, from *ιατρος*, 'a physician,' and *σοφιστης*, 'one skilled in an art or science.' A learned, or thoroughly educated physician.

IATROTECH'NA, *Iatrotech'nes*; from *ιατρος*, 'a physician,' and *τεχνη*, 'art.' A practical physician or surgeon.

IATROTECHNICE, *Medicina*, Therapeutics.

IATRUS, *Iat'ros*, Physician; also, a surgeon.

IBERIS, *Lepidium iberis*.

IBERIS AMA'RA, *Bitter candytist*, (F.) *Passerage*, from *Iberia*, where it is indigenous. A small herbaceous plant, *Ord.* Cruciferae, indigenous in Europe, which was employed, of old, in gout, rheumatism, &c. The seeds have been used, in the dose of one to three grains, in asthma, bronchitis, dropsy, and hypertrophy of the heart. It is said to possess acro-narcotic properties; but it is not much used.

IBERIS BURSA PASTORIS, *Thlaspi bursa*—i. *Campestris*, *Thlaspi campestre*—i. *Sophia*, *Cardamine pratensis*.

IBICUIBA, *Becuibá*, *Becuibá nux.* A species of nut from Brazil, the emulsive kernel of which is ranked amongst balsamic remedies.

IBIS, *ibis*. A bird held sacred by the Egyptians. When sick, it is asserted that it was wont to inject the water of the Nile into its fundament: whence, according to Langius, was learned the use of glysters.

IBISCHA MISMALVA, *Althæa*.

IBISCUS, *Althæa*.

IBIXUMA, *Saponaria*.

ICE, *Sax. ír*, *Gla'cie*, *Frozen water*, (F.) *Glacé*. Iced water is much used internally, as the best refrigerant in fever. It is, also, applied externally, in cases of external inflammation, as well as in phrenitic and hernial affections, &c., in the form of the ice-cap and ice-cataplasm or ice-poultice; formed by half filling a bladder with pounded ice.

ICE-CAP, see Ice.

ICE-COLD, *Icy cold*, (F.) *Froid glacial*. A very strong morbid sensation of cold, compared by the patient to that which would be produced by the application of pieces of ice.

ICE-POULTICE, see Ice.

ICELAND, MINERAL WATERS OF. Hot springs are found in every part of Iceland. The most noted of these is one called Geyser, two days' journey from Hecla, and near Skalholt. The diameter of the basin is 59 feet; and the height to which the water is thrown is often more than 100. The heat of the water is 212°.

ICE-PLANT, *Mesembryanthemum crystallinum*, *Monotropa uniflora*.

ICHNOS, *ichnos*. The foot. The sole of the foot. The heel.

ICHOR, *ιχωρ*, *Ichos*. The serum of the blood, *Sarax*, *Sorides*, *Vicus*, *Pus malignum*, *Tubum*. A thin, aqueous, and acid discharge.

ICHOROIDES, *Ichorous*.

ICH'OROUS, *Ichorous*, *Ichoroides*, *Sarionous*, *Sa'nious*, (F.) *Ichoroux*, *Sarionux*. Belonging to or resembling ichor.

ICHOS, *Ichor*.

ICHTHYA, Hook, blunt.

ICHTHYOCOL'LA, from *ιχθυος*, *ichthos*, 'a fish,' and *κόλλα*, 'glue.' *Colla Piscium*, *Piscinassa*, *Fish-glue*, (F.) *Ichthyocolle*, *Colle de Poisson*. A name given to the dried swimming bladder or sound of *Acipenser huso*, and other species of acipenser, which is almost wholly composed of gelatin, and is employed in medicine in the formation of nutritive jellies. It is, also, occasionally used in anatomical injections. The *English Court Plaster* is made with it. In Russia, the *Acipenser* is not the only genus from which isinglass is procured; it is also obtained from *Silurus Glanis*, which Dr. Royle suggests to be the source of the *Samory* or *Samory Isinglass* of commerce. The sounds of the cod, ling, hake, &c., yield it. From these it is made in New England; and, in England, is termed *New York Isinglass*.

ICHTHYOPHAGISTS, *Ichthyoph'agi*, from *ιχθυος*, *ichthos*, 'a fish,' and *φαγω*, 'I eat.' People who feed habitually on fish;—generally the most uncivilized of mankind.

ICHTHYOPHAGOUS, *Piscivor* as.

ICHTHYOSE, *Ichthyosis*.

ICHTHYO'SIS, from *ιχθυος*, *ichthos*, 'a fish,' from the resemblance of the scales to those of a fish. *Lepido'sis*, *Ichthy'o'sis*, *L. Ichthy'o'sis*, *Lep'ra Ichthy'o'sis*, *Alvaras nigra* (Arab.), *Impetigo excorticat'ra*, (F.) *Ichthyose*; *Fisheskin*, *Porcupine Disease*. A permanently harsh, dry, scaly, and, in some cases, almost horny texture of the integuments of the body, unconnected with internal disorder. Willan and Bateman have two varieties, *I. simplex* and *I. cornea*. Alibert has three, the *I. nacrée* or *pearly*, the *I. corne*, and the *I. pellagre* or *Pellagra*.

ICHTHYOSIS PELLAGRA, *Pellagra*.

ICHTHYOSIS SEBA'CEA, *Sebaceous Ichthy'o'sis*. A morbid incrustation of a concrete subaceous substance upon the surface of the epidermis, confounded, according to Mr. E. Wilson, with ichthyosis, to which it bears a close resemblance. In many cases there is neither redness nor heat, nor is the affection often accompanied by constitutional symptoms.

ICHTHYOTOX'ICUM, from *ιχθυος*, *ichthos*, 'a fish,' and *τοξικον*, 'a poison,' *Ich'thyus*, fish poison.

ICHTHYUS, *Ichthyotoxicum*.

ICICA ARACOUCHI'NI, *Aracouchini* is a balsam, extracted by incision from this tree in Guyana. The Galibis use it for healing wounds.

ICICA ICICARIBA, see *Amyris elemifera*.

ICTA, *Vulva*.

ICTÈRE, *Icterus*—i. *Bleu*, *Cyanopathy*—i. *des Nouveaux-nés*, *Icterus infantum*—i. *Noire*, *Melæna*.

ICTERIC, *Ictericus*.

ICTERIC FEVER, *REMITTING*, see *Relapse*.

ICTÉRICIE, *Icterus*.

ICTERICUS, *Antiicteric*. Also, relating to, or resembling *Icterus*; *Ic'teric*, *Icteroï'des*, *Ictero'-des*, (F.) *Ictérique*.

ICTERITIA ALBA, *Chlorosis*—i. *Flava*, *Icterus*—i. *Rubea*, *Erysipelas*—i. *Rubra*, *Erysipelas*.

ICTERODES, *Ictericus*.

ICTERUS, *I. flavus* seu *verus*, from *ικτερος*, a species of weasel, whose eyes are yellow [?]. (*Icterus* meant, with the ancients, a yellow bird, perhaps the Oriole;) *Morbus arcuatus* seu *arguatus*, *Leuco-*

oli morbus (Paracelsus), *Auri'go*, *Auri'go*, *Morbus regius* seu *lute'olus*, *Cholelith'ia* *icterus*, *Cholelith'ia*, *Cholel'mia*, *Cholelith'ia*, *Cholel'sis*, *Dermatocholo'sis*, *Suffu'sio* *aurigino'sa* seu *fellis* seu *bilis*, *Icteric'ia*, *Icteric'ia*, *Icteric'ia* seu *icteroides*, *Cachex'ia* *icter'ica*, *Fellis* *suffu'sio* seu *obstructio* seu *Superf'u'sio*, *Bill'ous* *Dys'crasy*, *Jaundice*, *Yellow's*, *Jandere*, *Jaundere*, (Sc.) *Gulschoch*, (Prov.) *Jonas*, (F.) *Ictère*, *Ictérie*, *Jaunisse*, *Bile répandue*. A disease, the principal symptom of which is yellowness of the skin and eyes, with white feces and high-coloured urine. It admits of various causes; in fact, anything which can directly or indirectly obstruct the course of the bile, so that it is taken into the mass of blood and produces the yellowness of surface;—the bile being separated by the kidneys, causes yellowness of urine, and its being prevented from reaching the intestine occasions the pale-coloured feces. The prognosis, in ordinary cases, is favourable;—when complicated with hepatic disease, unfavourable. The treatment is simple:—an emetic or purgative, given occasionally so as to elicit the return of the bile to its ordinary channels; light tonics; unirritating diet; cheerful company, &c.

ICTERUS ALBUS, Chlorosis.

ICTERUS INFANTUM, *I. Neonatorum*, *Neotoc'ic'erus*, *Padiet'erus*, *Auri'go* *neophyto'rum*, *Yellow gum*, *Yellow govin*, (F.) *Ictère des nouveau-nés*, is a common affection, and frequently dependent upon obstruction of the choledoch duct by the meconium. It requires time; and castor oil, occasionally.

ICTERUS MELAS, *Melæna*—*i. Neonatorum*, *I. infantum*—*i. Niger*, *Melæna*.

ICTERUS SATURNI'US, *Lead jaundice*. The earthy-yellow hue in saturnine cachexy.

ICTERUS VIRIDIS, *Green jaundice*. *Icterus* when the skin is of a green or olive hue.

ICTODES FÆTIDUS, *Dracontium foetidum*.

ICTUS. 'A stroke or blow.' *Plegè*, *Plaga*, *Ictus solis*, a stroke of the sun. See *Coup de soleil*. *Ictus*, also, means the pulsation of an artery, and the sting of a bee or other insect.

ICTUS, Blow—*i. Cordis*, see *Heart*—*i. Cordis* et *arteriarum*, *Pulse*—*i. Sanguinis*, *Apoplexy*—*i. Solis*, *Coup de soleil*.

IDEA, *Id'ea*, *Ide'olum*, *Ideach* [? Paracelsus], (F.) *Idée*, from *ideo*, 'I see.' The image or representation of an object in the mind.

IDEA, *FIXED* or *PREDOMINANT*, *Id'ea fixa*, *Ideopeg'ma*. Tension of the mind on one notion; often observed in insanity.

Id'ea MORBI. Knowledge or idea of a disease.

IDEACH, *Idea*.

IDEAGENOUS, *Sentiment*.

IDEAL, *Idea'lis*. Mental, notional, fancied. *Morbi ideales*. Ideal diseases. Diseases of the imagination.

IDEA'TION, *Idea'tio*, from 'idea.' The establishment of a distinct mental representation or idea of an object.

IDÉE, *Idea*.

IDEN'TITY (PERSONAL), *Iden'titas*, from *idem*, 'the same.' *Sameness*. It is sometimes a question in legal medicine to decide upon personal identity: that is, whether an individual be the same he represents himself to be. Physical marks form the great criteria.

IDEOL'OGY, *Ideolog'ia*, from *ideo*, 'I see,' and *logos*, 'a discourse.' The science of ideas. Intellectual philosophy.

IDEO-MOTOR. A name given by Dr. Carpenter to muscular movements, the result of complete engrossment by an idea, which he regards as automatic, although originating in the cerebrum.

IDEOPEGMA, *Idea*, fixed.

IDÉOSYNCHYSIE, *Delirium*.

IDIAN'EA, *Idiotism*.

IDIAPOCAUSIS, *Combustion*, *human*.

IDIEMPRESIS, *Combustion*, *human*.

IDIOC'RASIS, *Idiosyncrasy*.

IDIOCTONIA, *Suicide*.

IDIOCY, *Idiotism*.

IDIO-MIASMATA, see *Miasm*.

IDIOPATH'IA, *Idiopath'ia*, *Protopath'ia*, *Pro'prius affectus*, *Morbus idiopath'icus* seu *pro'prius seu prima'rius seu protopath'icus seu origina'lis*, *Malum prima'rium*, from *idios*, 'peculiar, proper,' and *pathos*, 'an affection.' A primary disease; one not depending on any other.

IDIOPATH'IC, *Pro'prio hab'itu* seu *Constitutio'nè pro'pria pendens*, *Idiopath'icus*, *Idiop'athes*. Primary affections and their symptoms are so denominated.

IDIOPTCY, *Achromatopsia*.

IDIOPTS, see *Achromatopsia*.

IDIOSYNCRASIA, *Idiosyncrasy*—*i. Hæmorrhagica*, *Hæmatophilia*—*i. Olfactoria*, *Parosmia*.

IDIOSYNCRASY, *Idio'crasy*, *Idiosyncrasis*, *Idiosyncrasis*, *Idiotrop'ia*, *Idio'crasia*, *Idiosyncrasis*, from *idios*, 'peculiar,' *syn*, 'with,' and *crasis*, 'temperament.' A peculiarity of constitution, in which one person is affected by an agent which, in numerous others, would produce no effect. Thus, shell-fish, bitter almonds, produce urticaria in some, by virtue of their idiosyncrasies. Others faint at the sight of blood, &c.

ID'OT, *Idio'ta*, 'foolish, stupid, ignorant.' Now used for one who is fatuous, or who does not possess sufficient intellectual faculties for the social condition, and for preserving himself from danger,—*Homo fat'uus*, (Prov.) *Daleop*, *Pondling*, *Hobbit*. In law, one who has been without understanding from his birth, and whom the law presumes to be never likely to attain any.

IDIOTIA, *Idiotism*—*i. Endermica*, *Cretinism*.

IDIOTIE, *Idiotism*.

ID'IOTISM, *Idiot'ismus*, *Idioti'a*, *Idian'cia*, *Imbecil'itas mentis*; *Mo'ria demens*, *Ano'cia*, *Mo'ria*, *Mo'ro'sis*, *Meio'sis*, *Fatu'itas*, *Amen'tia*, *Demen'tia natura'lis*, *Stupid'itas*, *Vecor'dia*, *Imbecil'itas Ingen'ii*, *Id'ocy*, *Id'iotcy*, *Fatu'ity*, (F.) *Démence innée*, *Idiotisme*, *Idiotie*, *Apsychisme*—*Piorry*. Same etymon. A species of unsound mind, characterized by more or less complete obliteration of the intellectual and moral faculties. It may supervene on mania and melancholia, when it is termed *Demen'tia*, but more commonly it depends upon original conformation. It may also, be symptomatic of organic disease of the brain, which has come on after birth. *Idiotism* exists in various degrees. Some idiots are mere automata, exhibiting scarcely any sensibility to external impressions; others are capable of articulating a few words, and possess certain mental emotions to a limited extent. The physiognomy is usually vacant, step unsteady, and articulation imperfect or broken. The affection is almost always incurable; but it may often be palliated.

IDIOTISME, *Idiotism*.

IDIOTISMUS, *Idiotism*—*i. Endemicus*, *Cretinism*.

IDIOTROPIA, *Idiosyncrasy*.

IDOLUM, *Hallucination*, *Idea*.

IDROSIS, *Ephidrosis*.

IETRION, *Iatrimon*.

IF, *Taxus baccata*.

IFFIDES, *Plumbi subcarbonas*.

IGDE, *Mortar*.

IGDIS, *Mortar*.

IGNA'TIA AMA'RA, *Strychnos Ignat'ii*, *Ignatia'na Philippin'ica*. *Ord. Apocynaceæ*. The systematic name of the plant which affords *St. Ignatius's Bean*, *Faba Indica* seu *Sancti Ignat'ii* seu *febrif'uga*, (F.) *Ignatie*, *Fève de Saint*

Ignace, Noix d'Ignacur. The seeds are bitter and poisonous, containing strychnia, and only appear to differ from nux vomica in containing a larger proportion of that alkaloid.

IGNATIE, *Ignatia amara.*

IGNAVIA, *Inertia*—i. *seu Ignavitas partium genitalium*, Impotence.

IGNIS, 'fire.' Hence:

IGNIS ACTUALIS, *Cauterium*—i. *Animalis*, Animal heat.

IGNIS CALIDUS. 'A hot fire.' A violent inflammation, about to degenerate into gangrene.

IGNIS COLUMELLÆ, *Erysipelas.*

IGNIS FRIGIDUS. 'A cold fire.' *Sphacelus.*

IGNIS NATURALIS, *Animal heat*—i. *Persicus*, Anthrax, Herpes zoster—i. *Philosophicus*, Phosphorus—i. *Potentialis*, *seu Cauterium*—i. *Sacer*, *Erysipelas*, Herpes zoster—i. *Sancti Antonii*, *Erysipelas.*

IGNIS SAPIENTUM. The ancient name for the heat of horses' dung.

IGNIS SYLVATICUS, *I. sylvæstris seu volatilis seu volagrus*, *Strophilurus sylvæstris seu volatilis*, Wildfire rash, (F.) *Feu sauvage ou volage*, *Flamboie*. Probably, the *Porri'go larvælis* or *Crusta lactea* of infants. Also, a transient redness of the face and neck, sometimes observed in hysterical and chlorotic females.

IGNIS SYLVESTRIS, *I. sylvaticus*—i. *Vitalis*, Animal heat—i. *Volagrus*, *I. sylvaticus*—i. *Volaticus*, *I. sylvaticus.*

IGNIVOROUS, *Pyrophagus.*

IGNYE, *Poples.*

IGNYS, *Poples.*

ILAPHIS, *Arctium lappa.*

ILEA, *Flanks.*

ILEAC PASSION, *Ileus.*

ILEADELPH'US, from *ileum*, 'the ileon,' and *adelphos*, 'brother.' A double monster, united at the pelvis, and below.

ILECH, *Y'tech*, *Ilei'as*, *Ilias'ter*, *Ylia'ter*, *Eli-as'ter*, *Ilias'trum*, *Ilei'ados*, *Ilei'dos*, *Ilei'adum*, *Ili'adus*. Terms used by Paracelsus to designate the first matter:—the beginning of every thing.

ILEIADOS, *Ilech.*

ILEIADUM, *Ilech.*

ILEIAS, *Ilech.*

ILEIDOS, *Ilech.*

ILEI'TIS, from *ileum*, 'the intestine ileum,' and *itis*, a suffix denoting inflammation. Inflammation of the ileum.

ILEO. In composition, relating or appertaining to the ileum.

ILEO-CHOLEOSIS, *Diarrhœa*, bilious.

ILEO-COLIC, *Ileo-co'licus*. Relating to the ileum and colon:—as the ileo-colic valve or valve of Bauhin.

ILEO-COLITIS, *Enteritis.*

ILEO-DICLIDITE, *see Typhus.*

ILEO-LUMBAR, *Ileo-lumba'lis*, *Ilio-lumba'ris*. Belonging to the ilium and lumbar region.

ILEO-LUMBAR ARTERY, *Ilio-lumbar artery*, *Iliaco-muscular*, (Ch.) is given off by the hypogastric, opposite the base of the sacrum. It ascends behind the psoas muscle, and divides into two branches;—an ascending and a transverse, which give off numerous ramifications to the neighbouring parts.

ILEO-LUMBAR LIGAMENT, *Ilio-lumbar* or *Ilio-lumbo-ver'tebral ligament*, *Vertebro-iliae ligament*, is a broad, membraniform, triangular ligament, extending horizontally from the transverse process of the 5th lumbar vertebra to the upper and posterior part of the iliac crest. It unites the vertebral column with the pelvis.

ILEON, *I'l'eum*, *Ei'leon*, *I'ium*, *Intesti'num circumvolu'tum*, from *ilaiv*, 'to turn,' 'to twist.'

Anatomists have given this name to the longest portion of the small intestine, which extends from the jejunum to the cæcum. It was so called from its forming a considerable number of convolutions.

ILEOPYRA, *see Typhus.*

ILEOSIS, *Ileus.*

ILEO-TYPHUS, *see Typhus.*

ILES, *Flanks.*

ILEUS, *Ei'leus*, from *ilaiv*, 'I twist or contract.' *Col'ica Ileus seu spasmod'ica*, *Enterel'e'sin*, *Enterel'o'sis*, *Ileus spasmod'icus*, *Chordap'sia*, *Passio Ili'aca*, *Iliac* or *Ileac Passion*, *Voll'rulus*, *Ileo'sis*, *Misere'rê mei*, *Convul'culus*, *Tormentum*, *Intercept'io Intestino'rum*, *Enterosten'o'sis volvulus*, (F.) *Colique de Miséricorde*, *C. de Miséréré*. A disease, characterized by deep-seated pain of the abdomen, stercoraceous vomiting, and obstinate constipation. It is occasioned by hernia or other obstruction to the passage of the fæces through a part of the intestinal canal, *Enterosten'o'sis*. The term *Ileus* has been applied to various affections—to simple nervous colic, intussusception, and to strangulation of the small intestine, &c. Various remedies have been employed;—the majority for the purpose of procuring alvine evacuations,—as purgatives, in draught, pill, or glyster; suppositories, tobacco glysters; pure mercury; leaden bullets; antispasmodics and narcotics; blisters to the epigastrium; ice by the mouth, or injected into the rectum. Some of these have, occasionally, succeeded,—especially the tobacco glyster, and ice to the tumour, where the disease has been occasioned by strangulated hernia. It is very dangerous.

ILEUS FLAVUS, *Icterus*—i. *Icteroides*, *Icterus*—i. *Inflammatorius*, *Enteritis.*

ILEX AQUIFO'LIUM. The systematic name of the Common Holly, *Aquifo'lium*, *Agri'folium*, *πρινος* of the ancients [?], (Prov.) *Hollen*, (Sc.) *Holyn*, *Holene*, (F.) *Houx*, *H. commun* ou *Chêne vert*. Ord. Aquifoliaceæ. The leaves of this plant have been recommended as tonic, astringent, and antiseptic, and have been prescribed in atonic gout, intermittents, dyspepsia, &c. *Al'icine*, the active principle, has also been advised. It is obtained by dissolving the alcoholic extract of the leaves of the holly in water, and successively treating it with the subacetate of lead, sulphuric acid, and carbonate of lime. The filtered and evaporated product is then dissolved in alcohol; and the mixture filtered and evaporated in shallow vessels.

ILEX CASSINE, *Ilex vomitoria.*

ILEX MAJOR. From the berries of this tree, called by the Spaniards *Bellotas*, a juice may be expressed, which forms a slightly astringent emulsion with water, and has been recommended by some Spanish physicians in humid cough, hæmoptysis, &c.

ILEX MATE, *Ilex paraguensis.*

ILEX OPA'CA, *American Holly*, grows throughout the Atlantic portion of the United States. It is said to possess the same properties as the European variety.

ILEX PARAGUEN'SIS, *Viburnum lavigatum seu Cassinoi'des*, *Cassi'nê Pera'gua seu Carolini'na*, *I. Maté*; a native of Paraguay. It affords the celebrated *Mattee*, or *Matê*, *Folia Apalachines seu Pera'gua*, which is drunk in place of the Chinese tea by the people of Paraguay. By M. Lenoble, of Montevideo, the maté is referred to *Peara'lea glandulo'sa*.

ILEX VOMITO'RIA, *I. Cassi'ne*, *Cassi'na*, *Dahoon holly*, *Apalach'inê Gallie*, *South-Sea Tea*, *Evergreen Cassi'ne*, *Casse'na*, *Yanpon*, *Yopon*, (F.) *Thé des Apalaches*, *Houx Apalachine*, *Apalachine*, *Thé de la Mer du Sud*. A tree, indigenous in the southern parts of the United States.

The leaves, when dried, are aromatic, stimulant, stomachic, and expectorant, and are used as a tea. When fresh, they are emetic and cathartic.

ILIA. Flanks.

ILIAC, *Il'acus*, *Il'al*, (F.) *Iliaque*, from *Ilia*, 'the flanks.' Relating to, or connected with, the flanks.

ILIA. ARTERIES, *Arte'ria Il'acæ*. This term has been given to several arteries. The *Primary Iliaca*, (F.) *A. Iliacæ primitives*,—*Pelvicurales*, (Ch.) arise from the bifurcation of the aorta, opposite the body of the 4th lumbar vertebra, and descend, in a divergent manner; until, opposite the sacro-iliac symphysis, they divide into two considerable trunks,—the *internal Iliac* or hypogastric (which see), and the *external Iliac*—*Portio Iliacæ de la crurale*, (Ch.) This proceeds from the sacro-iliac juncture as far as the crural arch, when it assumes the name *Femoral Artery*. Before passing under the arch it gives off two pretty considerable branches,—the *Epigastric*, and the *Circumflexa Ilii*. This last, which is called, also, *anterior Iliac* by some anatomists,—*Circonflexe de l'Ilium* (Ch.); leaves the external iliac at the crural arch. It ascends along the outer edge of the iliacus muscle, and divides into two branches—an *internal* and an *external*.

The *Iliaca Inter'na Minor* seu *Ilio lumbal'is* is a small artery, which sometimes arises from the hypogastric; at others, from the beginning of the gluteal. It is sent, chiefly, to the psoas and iliacus internus muscles.

ILIAC CREST, *Crista Il'ii*, is the upper margin of the ilium. (F.) *Crête Iliacæ*. It is very thick, and curved like the *Italica S.* It affords attachment to the broad muscles of the abdomen.

ILIAC FOSSE are two in number; the *internal*, the depression presented by the ilium on its interior, and at the upper part in which the *Iliacus internus* is lodged;—and the *external*, an excavation on the outer surface of the same bone, occupied by the *Glutæi muscles*.

ILIAC MUSCLE, INTERNAL, *Il'acus internus Muscæ*, *Iliacus*, *Iliaco-trochantérien* (Ch.); *Femur moven'tium sept'imus*, *Iliac muscle*, is situate in the fossa iliacæ, and at the anterior and superior part of the thigh. It is broad, radiated, triangular; and is attached, *above*, to the two upper thirds of the fossa iliacæ, and to the internal part of the iliac crest;—*below*, it terminates by a tendon, which is common to it and the psoas magnus, and is fixed into the lesser trochanter. When this muscle contracts, it bends the pelvis on the thigh, and conversely. It is also a rotator of the thigh outwards, and prevents the trunk from falling backwards.

ILIAC PASSION, Ileus.

ILIAC REGIONS, *Regio'nes il'iacæ*, *Inan'ia*, are the sides of the abdomen between the ribs and the hips.

ILIAC SPINES, *Spinous Pro'd'esses of the Ilium*, are four in number. They are distinguished into *anterior* and *posterior*. Of the *anterior*, one is *superior*. It bounds the crista ili anteriorly, and affords origin to the sartorius muscle. The other is *inferior*, and receives the tendinous origin of the rectus femoris.

The two *Posterior Iliac Spines* are divided, like the anterior, into *superior* and *inferior*. They afford insertion to strong ligaments, which unite the ilium with the sacrum.

ILIACO-TROCHANTÉRIEN, *Iliacus internus*.

ILIACUS, I. internus—i. Externus, Pyramidalis.

ILIADUS, Ilech.

ILIAL, Iliac.

ILIASTER, Ilech.

ILIASTRUM, Ilech.

ILINGOS, Vertigo.

ILIO. In composition, relating or appertaining to the ilium.

ILIO-ABDOMINAL, *Obliquus internus abdominis*—i. *Aponévrosi-fémoral*, *Fascia lata muscle*—i. *Costal*, *Quadratus lumborum*—i. *Cresti-tibial*, *Sartorius*—i. *Femoral*, *Coxo-femoral*—i. *Iliopsoas* (nervus), *Musculo-cutaneous nerve*, *superior*—i. *Inguinal*, *Musculo-cutaneous nerve*, *inferior*—i. *Ischii-trochantérien*, *Glutæus minimus*—i. *Lombo-costo-abdominal*, *Obliquus internus abdominis*—i. *Lumbalis*, *Ileo-lumbar*, see *Iliac arteries*—i. *Lumbi-costal*, *Quadratus lumborum*—i. *Lumbo-vertebral ligament*, *Ileo-lumbar ligament*.

ILIO-PECTINE'A, *Ileo-pectineal*, belonging to the ilium and to the pecten, or pubis.

ILIO-PECTINEAL EMINENCE is so called from being formed by the junction of the ramus of the ilium with that of the pubis. It affords attachment to the psoas minor.

ILIO-PECTINEA LIN'E'A, *Linea innominata*, is the projecting line or ridge of the ilium and pubis, which forms part of the brim of the pelvis.

ILIO-PRÉTIBIAL, *Sartorius*.

ILIO-PSOÏTIS, from *Ilium*, 'the ilion,' *psout*, 'the loins,' and *itis*, denoting inflammation. Inflammation of the ilium and psoas muscle.

ILIO-PUBO-COSTO-ABDOMINAL, *Obliquus externus abdominis*—i. *Rotulien*, *Rectus femoris*—i. *Sacral*, *Sacro-iliac*—i. *Sacral articulation*, *Sacro-iliac articulation*—i. *Sacro-fémoral*, *Glutæus major*—i. *Scrotal nerve*, *Musculo-cutaneous*, *superior*—i. *Trochantérien*, *Glutæus medius*—i. *Trochantérien petit*, *Glutæus minimus*.

IL'ION, *Il'ium*, *I'leum*, *Haunch bone*. The largest of the three bones which constitute the os innominatum in the fetus and child. It was probably so called from its seeming to support the intestine ilion; or, perhaps, because its crest is curved upon itself, from *ilew*, 'I twist,' 'I roll.' This portion of the os innominatum is usually so called in the adult. The posterior surface is called *Dorsum*, the internal *Venter*. The upper semicircular edge is the *Crista* or *Spine*, at the anterior part of which is the *anterior* and *superior spinous process*; and below, the *anterior* and *inferior spinous process*. At the back part of the spine are two *spinous processes*, the *posterior* and *superior*, and *posterior* and *inferior*. See *Iliac*.

ILITHY'IA, *Ειλαθυια*, *Luci'na*, *Juno Lucina*. The goddess who presided over parturient females, with the Greeks and Romans.

ILIU, Ileum, Ilium.

IL'KESTON, MINERAL WATERS OF. Ilkeston is in Derbyshire, England, about eight miles from Nottingham. The water contains carbonates of lime and soda, chloride of calcium, sulphate of magnesia, sulphate of soda, carbonic acid, sulphohydric acid, and a little iron.

ILL, Sick.

ILLACRYMATIO, Epiphora.

ILLECEBRA, *Sedum*—i. Major, *Sedum telephium*.

ILLECEBRUM POLYGONOIDES, *Achyranthes repens*—i. *Vermiculare*, *Sedum*.

ILLEGITIMATE, *Illegit'imus*, from *il*, *in*, 'negation or opposition,' and *legitimus*, 'legitimate,' from *lex*, *legis*, 'law.' *Bastard*. That which is contrary to law: which has not the conditions required by law, as an illegitimate birth—one out of wedlock; *base-born*, *love-begot*, *love-child*.

ILLEGITIMATE or BASTARD is also applied to diseases—as fevers, whose progress is anomalous. See *False*.

ILLI, see *Strabismus*.

ILLIC'IIUM ANISA'TUM, from *illicere*, 'to allure.' *Yellow-flowered Anise, Aniseed Tree, Star Anise, Ani'mum stellatum seu Sinen'se, Semen Badian, (F.) Anis de la Chine, A. étoilé, Badiane.* Ord. Magnoliaceæ. *Sc. Syst.* Polyandria Polygynia. The seeds are used like the aniseed. The same tree is supposed to furnish the aromatic bark, called *Cortex Ani'si Stella'ti, Cortex Lavalæ*.

Illicium Anisatum is said to furnish much of the so called *Oil of Anise*, used in the United States.

ILLICIUM FLORIDA'NUM, *Florida Anise Tree, Star Anise, Sweet Laurel.* An evergreen shrub, the bark, leaves, &c., of which have a spicy odour like anise, and might be used for it.

ILLICIUM PARVIFLO'rum, a shrub of the hilly regions of Georgia and Carolina, has a flavour closely resembling that of sassafras root.

ILLIDES, see Strabismus.

ILLINITIO, Inunction.

ILLISIO, Enthlasia.

ILLITIO, Inunction.

ILLITUS, Inunction.

ILLNESS, Disease—i. Monthly, see Menses.

ILLOS, Eye, see Strabismus.

ILLO'SIS, *Pl'o'sis*, from *ἄλσος*, 'squinting.'

Distortion of the eyes. Strabismus.

ILLUSIO SENSUS, Hallucination.

ILLUSION, Hallucination.

ILLUTAMENTUM, see Illutatio.

ILLUTA'TIO, from *il*, *in*, 'upon,' and *lutum*, 'mud.' A word used, by the ancients, for the act of covering any part of the body with mud,—*illutamentum*—with therapeutical views.

ILOSIS, Illosis.

IM. In composition, generally means the same as *in*; occasionally, however, it has the privative sense of *un*.

IMAGINA'TION, *Imagina'tio, Figura'tio, Phanta'sia, Dianoi'ma*, from *imago*, *imago*, 'image.' The faculty of forming in the mind an assemblage of images and combinations of ideas which are not always in connexion with external objects.

IMBALSAMATIO, Embalming.

IMBECIL'ITY, *Imbecillitas, Stupiditas, Hebetudo ani'mi, Oligopsychia*, (F.) *Imbecillité*. Weakness, especially of the intellect; incoherency: *Imbecillitas mentis*.

IMBECILLIS, Infirm.

IMBECILLITAS INGENII, Idiotism, Imbecility—i. Mentis, Idiotism, Imbecility—i. Ventriculi, Gasterasthenia.

IMBELLIS AD VENEREM, Frigid.

IMBER'BUS, *Imber'bus, Agenci'os, Apo'gon*, from *im*, *in*, 'negation,' and *barba*, 'beard.' One devoid of beard.

IMBITIO, Absorption, Imbibition.

IMBITI'ON, *Imbitio, Emp'osis, Aspiratio*, from *imbibere* (*in*, and *bibere*), 'to drink, to imbibe.' The action, by which a body becomes penetrated by a liquid. See Endosmose. Many of the phenomena of absorption are owing to imbibition.

IMBIBITION, DOUBLE, Endosmose.

IMBOWELLING, Exenterismus.

IMBEX NARIUM, Septum narium.

IMBRICARIA SAXATILIS, Lichen saxatilis.

IMMERSUS, Subscapularis muscle.

IM'MINENCE, *Imminen'tia*, from *immineo* (*im*, and *minuo*), 'to hang over.' Staying over, or upon; impending. Some authors have designated, by this term, the period which precedes the invasion of a disease; when certain indications foretell its approach.

IMMINENCE MORBIDE, Diathesis, Predisposition.

IMMISSIO CATHETERIS, Catheterismus.

IMMISSOR, Catheter.

IMMOBILITAS, Acinesia—i. Pupillæ, Amaurosis.

IMMODERANTIA, Intemperance.

IMMODERATIO, Intemperance.

IMMOVABLE APPARATUS, see Apparatus, immovable.

IMMUTANS, Alternative.

IMPACTED, Wedged.

IMPACT'ION, *Impac'tio*, from *impingere* (*in*, and *pangere*, *paetum*, 'to drive in'), 'to strike against,' 'a collision.' A fracture of the cranium, ribs, or sternum, with depression of some of the fragments and projection of others externally. See Epiesma.

IMPAT'IENTS BALSAM'INA, *Balsam weed, Touch-me-not.* Ord. Balsaminaceæ. This probably resembles the other species in its properties.

IMPAT'IENTS FULVA and **I. PAL'LIDA**, *Touch-me-not, Jewel weed, Balsam weed, Slippers, Cel'andine, Quick-in-the-hand, Weath'eroocks, Snap-weed*; called from the sudden bursting of the pods when touched. Indigenous plants, having tender, juicy, almost transparent stems, and yellow flowers, which appear in July and August. It is found in low, moist ground, in every part of the Union. The properties are probably the same as those of

IMPAT'IENTS NOLI-ME-TAN'GERE of Europe, which has an acrid taste, and acts as an emetic, cathartic, and diuretic.

IMPATIENTIA VESICÆ, Cysterethismus.

IMPEDIMENTUM, Emphragma.

IMPÉ'ROIRE, Imperatoria.

IMPERATO'RIA, *I. Ostruthium, Selinum ostruthium seu Imperato'ria, Angelica officinalis, Astruthium, Ostruthium, Astran'tia, Mugistrantia, Masterwort*, (old F.) *Austruche, Impératoire.* Ord. Umbelliferae. The roots of this plant were formerly considered *divinum remedium*. They are merely aromatic and bitter, and are not used.

IMPERATORIA SYLVESTRIS, *Angelica sylvestris.*

IMPERFORATE, Atretus.

IMPERFORATIO, Imperforation—i. Ani, Atrisia ani'adnata, Proctatresia—i. Pupillæ, Synexizis—i. Uteri, Metratresia.

IMPERFORATIO, from *im*, 'privative,' *per*, 'through,' and *fora*, *e*, 'to bore.' *Imperforatio, Atrisia, Atrétismus, Cap'tetus*. Absence of a natural aperture; as, of the mouth, anus, vulva, nostrils, &c. It is congenital, being dependent upon faulty conformation.

IMPERFORATION DE L'ANUS, Atrisia ani adnata.

IMPE'RIAL. A pleasant, cooling drink, formed of bitartrate of potassa, ʒss; one lemon, cut into slices; white sugar, lbss; and water, Oijj. Let the mixture stand for half an hour, and strain.

IMPETIGINOSITAS, Impetigo.

IMPETIG'INOUS, *Impetiginosus*, (F.) *Impétigineux*. Same etymon as the next. Having the nature of, or relating to, impetigo.

IMPETI'GO, from *impeto*, 'I invade.' *Impetiginositas, Dart, Der'bia, In'petus, Peti'go, Epye'sis seu Phlysis impetigo, Lepa Squamosa, Running Scall or Tetter, Crusted, Pustular or Humid Tetter or Scall, Cowrap*, (F.) *Dartre, D. crustacée ou croûteuse, Lèpre humide, Mili-tagre*. A word used in various acceptations. With some writers it is synonymous with itch. In others, it means a variety of herpes. Sauvages employs it as a generic term, under which he comprises syphilis, scorbutus, rachitis, elephantiasis, the itch, tinea, scrofula, &c. It forms, also, a genus in the class *Cachexiæ* of Cullen. In Bateman, it is the first genus of the 5th order, *Pustula*, and is defined—the humid or running tetter; consisting of small pustules, denominated

Pydracia. It is unaccompanied by fever; not contagious, or communicable by inoculation. He has five species:—the *I. figura'ta*, *I. sparsa*, *I. erysipelato'des*, *I. scab'ida*, and *I. rodens*. See *I'sorialis*.

IMPETIGO EXCORTICATIVA, Ichthyosis—i. Figurat'a, Porrigo lupinosa—i. Pellagra, Pellagra—i. Ulcerata, Zerna.

IMPETUM FACIENS, Enormum.

IMPETUS, Impetigo, Paroxysm.

IMPLACENTAL, see Placenta.

IMPLICATIONES RETICULARES NERVORUM, Plexus nervorum.

IMPLIC'ITI MORBI, *Implica'ti* seu *complica'ti* Morbi, from *im*, 'in,' and *plico*, 'to fold.' Diseases, which exist in an organ; and produce, concurrently, disorders in other organs.

IMPLUVIUM, Embrocation, Bath, shower.

IMPOSTHUME, Abscess.

IMPOTENCE, *Acrati'a*, *Impoten'tia*, *I. generandi*, *Agene'sia*, *Agene'sis*, *Ignavia* seu *Ignavia's* partium genitalium, (F.) *Impuissance*; from *im*, 'privative,' and *potens*, 'able.' Loss of power over one or more of the members. Commonly, it means want of sexual vigour; incapacity for copulation; and chiefly on the part of the male. *Astys'ia*, *Astyph'ia*, *Aeyno'dia*, *Adynamia viri'tis*. It has, also, been used synonymously with sterility. Impotence may be absolute or relative, constitutional or local, direct or indirect, permanent or temporary.

IMPOTENTIA, Adynamia, Impotence—i. Generandi, Impotence.

IMPOVERISHED, *Effic'tus*, *Depanperatus*, (F.) *Appauvri*. 'Having become poor;' originally from (L.) *pauper*, (F.) *pauvre*, 'poor.' The Humorists applied this epithet to a humour, deprived of a part of its constituents, and particularly to the blood. This fluid was considered to be impoverished when it was pale, without the proper consistence, and abounding in serum. It was, on the contrary, rich, in their opinion, when of a scarlet colour; possessing consistence; when it coagulated promptly, and the quantity of serum, compared with that of the clot, was by no means considerable.

IMPREGNATIO, Feundation, Pregnancy.

IMPREGNATION, Feundation, Pregnancy.

IMPRESSIBILITY, see Sensibility—i. Great, *Hyperæsthesis*.

IMPRES'SION, *Impres'sio* (in, and *premere*, *pressum*, 'to press'), *Pros'bole*, (F.) *Empreinte*. A more or less deep indentation which certain organs seem to make on others. Inequalities observable on the bones, which appear to be made by the subjacent organs.

IMPRES'SIONS DIG'ITAL, *Impressio'nes digita'tæ cran'ii*, *Fossæ digita'les*, are the depressions of various forms, observable at the inner surface of the cranium, which look, at first view, as if they were made with the fingers.

IMPRESSIONABILITÉ, see Sensibility.

IMPU'BER, *Impu'bes*, *Impu'bis*, *Ane'bus*, *Capilla'tus*, from *in*, 'negation,' and *puber*, 'at the age of puberty.' (F.) *Impubère*. One who has not attained the age of puberty.

IMPUBES, Impuber.

IMPUDICUS, see Digitus.

IMPUSSANCE, Impotence.

IMPULSE, DIAS'TOLIC, *Impul'sus diastol'icus*, *Back stroke of the heart*. A jog or stroke which has been termed the 'back stroke,' felt at the end of each pulsation, and which would seem to be owing to the refilling of the ventricles.

IMPULSE OF THE HEART, see Heart.

IMPULSION DU CŒUR, see Heart.

IMPURITATES ALVINÆ, Fæces.

INANIA, Iliac regions.

INANITIATED, see Inanitation.

INANITATION, *Inanitia'tio*, same etymon as the next. The act of being exhausted for want of nourishment. One so exhausted is said to be *inaniti'tated*.

INANIT'ION, *Inaniti'tio*, *Inan'itas*, from *inane*, *inanitum*, 'to empty.' *Ceno'ria*. Exhaustion for want of nourishment. To die from inanition is to die from exhaustion.

INAPPETENTIA, Anorexia, Dysorexia.

INARTICULATIO, Enarthrosis.

INAURA'TION, *Inauratio*, from *in*, and *aurum*, 'gold.' The gilding of pills or boluses.

INCANTAMENTUM, Charm.

INCANTATION, *Incanta'tio*, *Incantamen'tum*, *Ep'o'de*, from *in*, and *cantare*, *cantatum*, 'to sing,'—for example, a magical song. A mode of curing diseases by charms, &c., defended by Paracelsus, Van Helmont, and others.

INCARCERA'TIO, *Chatoniement*, Incarceration—i. *Intestinorum interna*, *Enteremphraxia*—i. *Intestinorum*, *Enteropéristole*.

INCARCERATION, *Incarcera'tio*, from *in*, 'in,' and *carcere*, 'prison.' Hernia is said to be 'incarcerated,' *Hernia incarceration'a*, when, owing to constriction about the neck of the hernial sac or elsewhere, it cannot be reduced with facility. Incarceration is sometimes used in the same sense as strangulation.

INCARNANS, *Incarnat'ivus*, *Sarco'ticus*, *Stal'ticus*, *Plero'ticus*, *Anaplero'ticus*, *Sarcot'ic*, from *in*, and *caro*, *carnis*, 'flesh.' Medicines, which were fancied to promote the regeneration of the flesh. Certain bandages and sutures have, also, been so called.

INCARNATIO, Incarnation—i. Unguim, *Onychogryphosis*.

INCARNATION, *Incarna'tio*, same etymon. Growth of flesh or granulations.

INCARNATUS, Carneous.

INCENDIUM, from *incendere* (in, and *candere*), 'to burn.' *Pyra'ca*, *νυκται*, *Causis*, *Phlogosis*. A burning fever or any burning heat.—*Incen'dium febrile*, *Incen'sio*.—Willis. Inflammation, Phlegmon.

INCENDIUM SPONTANEUM, Combustion, human.

INCENSIO, Incendium.

INCENTIVUM, Stimulant.

INCERA'TION, *Incera'tio*, *Encero'sis*, from *in*, and *cera*, 'wax.' The act of incorporating wax with some other body; or, rather, the operation whose object is to communicate to a dry substance the consistence of wax.

INCERNIC'ULUM, from *incernere* (in, and *cernere*), 'to sift.' A strainer or sieve.

INCESSIO, Bath, half, Semicupium.

INCESSUS, Walking.

INCIDENT'IA, from *incidere*, *incisum* (in, and *cædere*), 'to cut.' (F.) *Incisifs*. This name was formerly given to medicines to which was attributed the property of cutting thick or coagulated humours. The fixed alkalies, hydrosulphurets of alkalies, sulphurous water, &c., were considered to belong to this class.

INCINERA'TION, *Incinera'tio*, *Cinera'tion*, *Cinera'tio*, from *in*, and *cinis*, *cineris*, 'ashes.' In pharmacy, a process by which animal or vegetable substances are reduced to ashes.

INCINCTA, Pregnant.

INCISIFS, Incidentia.

INCISIO, Cut, Incision—i. Simplex, Haplotomia.

INCIS'ION, *Incis'io*, *En'copè*, *Incis'ura*, *Incisus*, *Tomè*, (F.) *Taillade*. Same etymon as Incidentia. A methodical division of soft parts with a cutting instrument.

INCISIVE, *Incisi'vus*, *Incisori'us*, *Fossa incis'iva* seu *myrtifor'mis*, is a depression in the superior maxillary bone, above the incisor teeth.

INCISIVE BONE, Intermaxillary bone—i. Canal, see Palatine canals.

INCISIVE NERVE, *Nervus incisivus*. A branch of the inferior dental nerve, which supplies the incisor teeth.

INCISIVE TEETH, *Incisor Teeth*, *Dentes incisives* seu *tomici* seu *ctenes* seu *dichastres* seu *gelastini* seu *primores* seu *pasorii*. The teeth which occupy the anterior part of the upper and lower jaws are so called, because they are used for cutting the food in the manner of cutting instruments. The two middle incisors of the upper jaw are, at times, called *Butter teeth*.

INCISIVUS (MUSCULUS), Levator labii superioris proprius — i. Inferior, Levator labii inferioris — i. Lateralis et pyramidalis, Levator labii superioris alaeque nasi — i. Medius, Depressor alae nasi.

INCISOR TEETH, Incisive teeth.

INCISORIUM, *Tomion*. An operating table. A scalpel.

INCISORIUS, Incisive.

INCISURA, *Échancreure*, Incision — i. Crucialis, Crucial Incision — i. Ischiadica Major, Sciatic notch, greater — i. Ischiadica Minor, Sciatic notch, lesser — i. Lienalis, Hilus lienalis — i. Renalis, Hilus renalis — i. Scapularis, Notch, scapular — i. Septi, see Ventricles of the brain.

INCISURA TRAGICA, Notch of the Concha. A wide, deep and rounded notch, which separates the tragus from the antitragus.

INCISURES DE SANTORINI, Santorini, fissures of.

INCISUS, Incision.

INCITABILITY, Irritability.

INCITAMENTUM, Stimulus.

INCITANS, Stimulant.

INCITATIO, Stimulation.

INCITO-MOTOR, (F.) *Incito-moteur*. An epithet, applied to an action the reverse of excitomotor; as in the case of muscular motion, which commences in the nervous centres and excites the muscles to contraction.

INCLAVATIO, Gomphosis, see Wedged.

INCLINATIO, Decline — i. Colli, Climate.

INCLUSION, MONSTROSITY BY, *Emboîtement*.

INCLUSUS, see Wedged.

INCOCTUS, see Crudity.

INCOHERENCE, *Incoherence*, (F.) *Anacoluthe*, *Réasserie*; from *in*, 'negation, co, con, cum, 'with,' and *herere*, 'to stick.' Want of dependence of one part on another. The condition of the mental manifestations in dementia, &c.

INCOMPATIBLE, from *in*, 'negation,' and *competere*, 'to agree.' A substance, which cannot be prescribed with another, without interfering with its chemical composition or medicinal activity.

INCONTINENTIA, from *in*, 'negation,' and *contineo*, 'I contain,' *Anepischesis*. 'Incontinence,' 'Inability to retain the natural evacuation.' Abuse of the pleasures of love.

INCONTINENTIA ALVI, Diarrhoea — i. Urinae, Enuresis. Scorracria.

INCORPORATION, *Incorporatio*, *Corporeatio*, *Ensomato'sis*, *Metensomato'sis*, from *in*, and *corpus*, *corporis*, 'a body.' An operation, by which medicines are mixed with soft or liquid bodies, in order to give them a certain consistence. Also, the thorough admixture of various substances.

INCRASSANTIA, *Incrassantia*, *Inviscantia*, *Spissantia*, *Epepytica*, *Condensantia*, *Pachynotica*, *Pycnotica*, *Pycnemasica*, from *in*, and *crassus*, 'thick.' Medicines which were formerly believed to possess the power of thickening the humours, when too thin. All mucilaginous substances were so regarded.

INCRASSATIVA, Incrassantia.

INCREASE, *Incrementum*, *Auris*, *Aure'sis*, from *in*, and *crecere*, 'to grow.' (F.) *Accroissement*. Augmentation of the size or weight of a body, by the application of new molecules around those which already exist.

INCREMENTUM, Augmentation, Growth, Increase.

INCRUCIATIO, Decussation.

INCUSTATION, *Incrustatio*, from *in*, and *crusta*, 'a crust,' *Con'asis*. The act of forming a crust on the surface of a body, as well as the crust itself. Also, the calcareous deposits or cartilaginous plates, which sometimes form in organs.

INCUBATIO, Incubation.

INCUBATION, *Incubatio*, *Incubatio*, *Incu-bitus*, from *incubare*, *incubitus*, (*in*, and *cubare*), 'to lie upon.' This word, which is used in natural history for the sitting of birds upon their eggs, is employed, figuratively, in medicine, for the period that elapses between the introduction of a morbid principle into the animal economy and the invasion of the disease.

INCUBITIO, Incubation.

INCUBITUS, Incubation.

INCUBO, Incubus.

INCUBUS. Same etymon. *In'cubo*, *Epi'al'tes*, *Ephial'tes*, *I. nocturnus*, *Hypnophob'ia*, *Epilep'sia nocturna*, *Asthma nocturnum*, *Prigal'ion*, *Sue'cubus*, *Oneirodyn'ia gravis*, *Erethis'mus oneirodyn'ia*, *Noctur'na oppres'sio*, *Epi'b'ole*, *Babuzica'rius*, *Nightmare*, *Wildmare*, *Riding-hag*, (Prov.) *Bitch-daughter*, *Hella*, (F.) *Cauchemar*, *Cauchevieille*, *Cochemar*, *Oneirodynie gravative*. Same etymon. A sensation of a distressing weight at the epigastrium during sleep, and of impossibility of motion, speech, or respiration; the patient at length awaking in terror, after extreme anxiety. Nightmare is often the effect of difficult digestion or of an uneasy position of the body. At other times, it occurs in consequence of severe emotions. The sensation of suffocation was formerly ascribed to the person's being possessed, and the male spirits were called *incubes* — the female *succubes*. The disease requires no particular treatment. The causes must be avoided.

INCUBUS VIGILANT'UM, *Ephial'tes vigilan'tium*, *E. hypochondri'aca*, *Daymare*. This is produced during wakefulness; the sense of pressure being severe and extending over the abdomen; respiration frequent, laborious, and constricted; eyes fixed; sighing, deep and violent; intellect, undisturbed.

INCUNABULA, Swathing clothes.

INCUNEATIO, Gomphosis, see Wedged.

INCURABLE, *Incurabil'is*, *Rem'ediless*, *Irre-me'diable*, *Insanab'ilis*, *Ana'thes*, *Anaces'tos*, *Atherapeu'tus*, from *in*, negative, and *cura*, cure. Not susceptible of cure; — applied to both patients and diseases.

INCURIA, Acedia.

INCURSUS ARTERIARUM, Pulse.

INCURVATIO, Gynopsis.

INCUS, genitive, *Incu'dis*, (*in*, and *cudo*, 'I hammer,') 'an anvil.' *Aemon*, *Os incu'di sim'ile*, *Ossic'ulum Incu'di* seu *mola'ri denti compara'tum*, (F.) *Enclume*. One of the small bones of the ear, so called from its fancied resemblance to a smith's anvil. It is situate in the cavity of the tympanum, between the malleus and orbiculaire. Its body is articulated with the malleus; its horizontal ramus, which is the shortest, corresponds with the entrance of the mastoid cells; and its vertical ramus is articulated with the os orbiculaire. The incus is composed of a compact tissue, and is developed from a single point of ossification.

INDEX, *Dig'itus index* seu *secun'dus* seu *salu-*

tator'ius, Demonstrator, Indica'tor, Indicato'rius, Lichanos, Deic'ticos, from indicare, 'to point out.' The forefinger, index finger. See Digitus.

INDIAN ARROW WOOD, see *Euonymus*—*i. Paint, Hydrastis Canadensis*—*i. Physic, Gillenia trifoliata*—*i. Physic, small-flowered, Gillenia stipulacea.*

INDIAN SPRINGS, see *Georgia, Mineral waters of.*

INDIA'NA, MINERAL WATERS OF. A medicinal spring, near Jeffersonville, is much frequented. Its waters are strongly impregnated with sulphur and iron.

INDICANT, *In'dicans, Endeic'ticos, (F.) Indictif.* Same etymon as *index*. Any thing which, in the course of a disease or in what precedes or accompanies it, concurs in pointing out the means to be used for its cure.

INDICATIVE, *Indicant.*

INDICATING DAYS, *Dies In'dices seu Indicato'rii seu Indican'tes seu Contemplan'tes, (F.) Jours indicateurs.* Hippocrates and others have thus called the middle day of the septenary; because they imagined, that indications were sometimes given then of the crisis which would occur in the last day of the septenary.

INDICATION, *Indica'tio, Accusa'tio, Dela'tio, Endeix'is, Endex'is, Boethemat'icum Seme'ion.* The object proposed in the employment of any means which are had recourse to for the cure of disease. It may also be defined—the manifestation afforded by the disease itself of what is proper to be done for its removal.

INDICATOR, *Extensor proprius indicis,* see *Digitus, and Index.*

INDICUM, *Indigo.*

INDICUS COLOR, *Indigo.*

INDIGENCE, *Want.*

INDIG'ENOUS, *Indig'ena, 'Inborn.* From *indū*, obsolete for 'in,' and *geno*, obsolete for *gigno*, 'I generate.' Whatever is native in a country, in opposition to *exotic*; as an *indigenous remedy, indigenous disease, &c.*

INDIGESTION, *Dyspepsia*—*i. Alkaline,* see *Dyspepsia*—*i. Neutral,* see *Dyspepsia.*

INDIGITATIO, *Intussusceptio.*

INDIGNABUNDUS, *Rectus externus oculi.*

INDIGNATORIUS, *Rectus externus oculi.*

INDIGO, *Indigum, Ind'icum, Indicus color, Pignētum Ind'icum.* A dye-stuff, in small solid masses, of a deep azure blue colour, and devoid of smell and taste. It is obtained, by a fermentative process, from *Indigofera anil, (I. suffruticosa), I. argentea, (I. articulata seu tinctoria seu glauca seu colorata), and I. tinctoria seu Ind'ica,* and is sometimes used by the apothecary for colouring certain preparations. It has likewise been administered internally, of late, in spasmodic diseases, especially in epilepsy. The dose may be at first grains, but it may be elevated to drachms.

INDIGO, FALSE, (TALL WHITE,) see *Sophora tinctoria.*

INDIGO, FAUX, *Galega officinalis*—*i. Weed, Sophora tinctoria.*

INDIGO, WILD, *Sophora tinctoria*—*i. Yellow, Sophora tinctoria.*

INDIGOFE'RA, see *Indigo, Sophora tinctoria.*

INDIGUM, *Indigo.*

INDISPOSIT'ION, *Mala disposit'io, Dyspho'ria, (F.) Malaise.* From *in*, 'negation,' and *disponere, dispositum*, 'to put in order.' A slight functional disturbance, which may scarcely bear the name of disease. A feeling of sickness—*sensus agri'tu' dinis*; vulgarly an *all-overish* feeling.

INDOLENT, *Indolens, from in*, privative, and *dolere, 'to be in pain.'* Exhibiting little or no pain. An epithet particularly applied to certain tumours.

INDOLENTIA, *Anodynia.*

INDOLES, A natural disposition or character.

INDOLES AN'IMI. The natural disposition or character of mind.

INDOLES MORBI. The nature or character of a disease.

INDOSYNCLONUS, *Beriberi.*

INDUC'TIO, *Apogogē, from inducere, (in, and ducere, ductum, 'to lead.')* A word used especially for the action of extending a plaster upon linen.

INDUCULA, *Waistcoat, strait.*

INDUMENTUM CORDIS, *Pericardium*—*i. Nervorum, Neurilemma*—*i. Ventriculorum, Ependyma ventriculorum.*

INDURANS, (*in* and *durus*, 'hard.')

Sclerot'icus, Sclerut'icus, Sclerypticus. A medicine which hardens the parts to which is applied.

INDURATIO, *Induration*—*i. Intestinorum, Enteropathia cancerosa*—*i. Maligna, Scirrhus*—*i. Renum, Nephroscleria*—*i. Telæ cellulosa neonatorum,* see *Induration*—*i. Ventriculi scirrhusa, Gastroscirrhus.*

INDURA'TION, *Indura'tio, Sclerys'ma, Sclerysmus, Callos'itas, Indurescent'ia, Poro'ma, Poro'sis, (F.) Endurcissement.* The hardness which supervenes, occasionally, in an inflamed part. It is one of the terminations of inflammation, and is owing to a change in the nutrition of the part.

INDURATION OF THE AREOLAR or CELLULAR TISSUE, *Indura'tio cellulosa, Induratio seu Opila'tio telæ cellulosa neonato'rum, Ethniphytolo'sis, Edema neonato'rum, Compact Edema of Infants, Oat'ochus Infantum, Scleremia, Sclerema, Sclero'ma, Scleroder'ma, Cutis tensa chronica, Dermatoperiarterio'sis, Dermatoperisclerismus, Halonit'is, Phlegmasia cellularis, Scirrhus neonato'rum, Scler'asis neonato'rum, Sclero'sis, Stipa'tio telæ cellulosa Infantum, Skin-bound Disease, Skin-blind, Skin-tight, Hæm-bound, (F.) Endurcissement du tissu cellulaire; Œdème du tissu cellulaire des nouveau-nés, Asphyxie lente des nouveau-nés,* is a disease which attacks infants a few days after birth, and which Chaussier proposed to call *Sclérème*, from *σκληρος, 'hard.'*

The Induration of the areolar tissue of the adult, *Sclerosteno'sis cutanea,* has been regarded as the result of inflammation of the corium, *Chorionit'is.*

INDURATION OF THE BRAIN, *Sclerencephalia*—*i. Grise,* see *Hepatitis.*

INDURESCENTIA, *Induration.*

INDU'SIUM, *Chitonis'cos.* Strictly, 'a shirt,' 'a small tunic.' Some have so called the amnion.

INE'BRIANT, *Inebrians, Phantas'ticus;* from *in* and *ebrio*, 'I intoxicate.' Intoxicating. An agent that intoxicates.

INEBRIATION, *Temulentia.*

INÉBRIATION QUINIQUE, *Quinism.*

INEDIA, *Fasting.*

INER'TIA, from *in*, 'negation,' and *ars, artis, 'art, Ignavia, Ener'vitas.* Sluggishness, inactivity.

INER'TIA OF THE WOMB, (F.) *Inertie de la matrice.* The diminution and even total cessation of the contractions of the uterus during labour; as well as the species of languor into which it sometimes falls after the expulsion of the fetus.

INERTIE PAR ÉPUISEMENT, see *Parturition*—*i. de la Matrice, Inertia of the womb.*

INESIS, *Cenosis.*

INETHMOS, *Cenosis.*

IN EXTREMIS, *Psychorages.*

IN'FANCY, *Infan'tia, from in*, 'negation,' and *fans, from fari, 'to speak.'* Early childhood. Childhood; *Ne'piotes, Paid'a, (F.) Enfance.* It generally includes the age from birth till the seventh year. See *Age.*

INFANS, *Paid'ion, Ne'pion, (Sc.) Gaitling,*

Simple accumulation of blood in the capillaries of the part is *hyperæmia*, not inflammation. The latter requires, in addition, altered nutrition either of the vessels themselves, or of the surrounding tissues, or of both.

The inflammations of the areolar and serous membranes greatly agree;—and those of the mucous and skin; the former being more active, and constituting the *phlegmonous* variety:—the latter, the *erythematic* or *erysipelatous*. Of this variety is the *diffusive inflammation* produced by morbid poisons,—as during dissection, where solutions of continuity exist on the fingers of the operator. It is seen, too, in workers in bone, and hence has been called *bone fever*. Preparatory to the turning of bones, it is customary to macerate them in water. The fluid soon becomes putrid, and if the hands be kept in it diffusive inflammation results. Inflammation may end by resolution, suppuration, gangrene, adhesion, effusion, or induration. Each of the inflammations of internal organs has received a name according to the organ affected;—as, *gastritis*, *cephalitis*, *enteritis*, *hepatitis*, &c. Besides the above inflammations, it has been considered, that there is an instinctive kind established for the union of parts which have been divided, whether the union takes place *immediately*, or by the aid of an intermediate body. This is the *adhesive inflammation* of some; but it may well be questioned whether inflammation is ever desirable in such cases. See Adhesion, Callus, Cicatrix. Broussais considered that the term *inflammation* should include every local exaltation of the organic movements which is sufficiently great to disturb the harmony of the functions, and disorganize the texture in which it is situate. He farther extended the name *inflammation* to irritations which do not induce disorganization of the textures, and which had been previously, and are still, called *fevers*.

Examination of the blood drawn always exhibits an increase of the fibrinous elements—the average proportion of which, in healthy blood, is about three in the thousand. In inflammation, it at times rises as high as ten. In fevers unaccompanied with inflammation, the proportion is natural, or below the average; but whenever inflammation supervenes, it immediately rises.

External inflammation is easily detected by the characters already mentioned:—*internal*, by disturbance of function and pain upon pressure; but the last sign is often not available. Both forms require the removal of all irritation, and the reduction of vascular excitement and nervous irritability; hence, blood-letting—local and general—sedatives, refrigerants, and counter-irritants become valuable remedies in almost all cases of inflammation.

Inflammation is not always of the active kind. There is a form, *Inflammatio debilis*, which occurs in low conditions of the system, and in broken down constitutions, which, like many forms of chronic inflammation, requires a stimulant, rather than a sedative, mode of treatment.

INFLAMMATION DES AMYGDALES. Cynanche tonsillaris—*i. des Artères*, Arteritis—*i. of the Bladder*, Cystitis—*i. de la Bouche*, Stomatitis—*i. des Bronches*, Bronchitis—*i. du Cæcum*, Typhlo-enteritis—*i. du Cerveau* et *du Cervelet*, Phrenitis—*i. du Cœur*, Carditis—*i. du Colon*, Colitis—*i. de la Conjonctive*, see Ophthalmia—*i. of the Diaphragm*, Diaphragmitis—*i. Diffusive*, see Inflammation—*i. Diphtheritic*, Diphtheritis—*i. of the Ear*, Otitis—*i. de l'Épiglotte*, Epiglottitis—*i. de l'Estomac*, Gastritis—*i. of the Eye*, Ophthalmia—*i. du Foie*, Hepatitis—*i. des Gencives*, Ulitis—*i. General*, Synocha—*i. of the Internal Membrane*

of the Heart, Endocarditis—*i. Internal*, Empyema—*i. des Intestins*, Enteritis—*i. of the Iris*, Iritis—*i. of the Kidney*, Nephritis—*i. de la Langue*, Glossitis—*i. of the Larynx*, Laryngitis—*i. of the Liver*, Hepatitis—*i. de la Luette*, Uvulitis—*i. of the Lungs*, Pneumonia—*i. of the Malpighian Bodies*, Kidney, Bright's disease of the—*i. des Mamelles*, Mastitis—*i. de la Matrice*, Metritis—*i. de la Membrane ulcéolo-dentaire*, Periodontitis—*i. de la Membrane séreuse céphalo-rachidienne*, Meningitis—*i. of the Mesentery*, Mesenteritis—*i. de la Moelle épinière* ou *rachidienne*, Myelitis—*i. of the Mouth*, pseudo-membranous, Stomatitis, pseudo-membranous—*i. of the Mouth*, pultaceous, Aphthæ—*i. des Muscles*, Myositis—*i. des Nerfs*, Neuritis—*i. de l'Œil*, Ophthalmia—*i. de l'Oreille*, Otitis—*i. de l'Os*, Oaritis—*i. du Palais*, Hypoeritis—*i. du Parachyme pulmonaire*, Pneumonia—*i. de la Parotide*, Cynanche parotidæa—*i. Pellicular*, Diphtheritis—*i. du Péricarde*, Pericarditis—*i. of the Pericardium*, Pericarditis—*i. du Péritoneum*, Peritonitis—*i. of the Peritonæum*, Peritonitis—*i. of the Pleura*, Pleuritis—*i. de la Plèvre*, Pleuritis—*i. des Poumons*, Pneumonia—*i. du Muscle psoas*, Psositis—*i. de la Rate*, Splenitis—*i. des Reins*, Nephritis—*i. Spongoid*, Hamatodes Fungus, see also Encephaloid—*i. of the Stomach*, Gastritis—*i. de la Testicule*, Hernia humoralis—*i. of the Testicle*, Hernia humoralis—*i. des Tissus blancs*, Angioleucitis—*i. of the Tongue*, Glossitis—*i. des Vaisseaux Lymphatiques*, Angioleucitis—*i. des Veines*, Phlebitis—*i. de la Vésicule du Fiel*, Cholecystitis—*i. de la Vessie*, Cystitis—*i. of the Womb*, Metritis.

INFLAMMATIUM' CULA, *Subinflammatio*. A superficial and often insignificant inflammation of the skin, as in many cutaneous affections.

INFLAMMATORY, *Inflammato'rius*; *Phlog'ic*, *Phlogo'us*, *Phlogis'ticus*, *Phlogis'tic*, belonging to inflammation;—as, *inflammatory tumour*, *inflammatory fever*, &c. The blood is said to be *inflammatory* when cupped or buffed. See *Corium phlogisticum*.

INFLAMMATORY BLUSH, Erythema.

INFLAMMATUS, Inflamed.

INFLATIO, Emphysema, Puffiness, Colica flatulenta—*i. Parotidum*, Cynanche parotidæa—*i. Uteri*, Physometra.

INFLEXIO, Campsis.

INFLUENCE, Influenza.

INFLUENTIA, Influenza.

INFLUENZA. The Italian for 'Influence.' *Influenza Europæa*, *Influentia*, *Catar'hus epidemicus*, *C. a contagio*, *Febris catarrh'al'is epidem'ica*, *Rheuma epidemicum*, *Morbus Vervecinus* seu *Catarrh'al'is* seu *Arie'tis*, *Syn'ochus* seu *Deflus'io catarrh'al'is*, *Cephalalg'ia contagiosa* (epidemics of the 16th and 17th centuries): (F.) Tac, Quinte, Florion, Coqueluche, Barquette, Générale, Grippe, Follette, Grenade, Coquette, Cocote, Petite Poste, Petit Courier, Alture, Ladendo, Fièvre catarrh'ale épidémique; Influence, Epidemic catarrh. A severe form of catarrh occurring epidemically, and generally affecting a number of persons in a community. See Catarrh, epidemic. Gluge, from his investigations considers that the following is the chronological order of the return of the influenza:—14th century, 1323, 1326—15th century, 1410, 1411, 1414—16th century, 1510, 1537, 1562, 1574, 1580, and 1593—17th century, 1658, 1669, 1675, 1693—18th century, 1708, 1712, 1729, 1732, 1733, 1742, 1743, 1761, 1762, and 1775—19th century, 1800, 1803, 1831, and 1833. To these may be added 1837, and 1843.

INFLUENZA EUROPÆA, Influenza.

INFLUENZOID, *Influenzoid'es*. An expressive but hybridous compound: from *influenza*, and

cedos, 'resemblance.' Resembling influenza.—Dr. T. Thompson.

INFORMITAS, Deformation.

INFORTUNUM, Contrafissura.

INFRA-ATLOIDÆUS, Sub-atloideus.

INFRA-AOXIDÆUS, Sub-axoideus.

INFRA-COSTALES, see Intercostal muscles.

INFRA-MAXILLARIS, Sub-maxillary.

INFRAMAXILLOSTERNODYMIA, Cephalostomatodymia.

INFRA-ORBITAR, Sub-orbital.

INFRAPUBIAN LIGAMENT, Triangular ligament.

INFRA-SCAPULARIS, Subscapularis.

INFRA-SPINALIS, Infra-spinatus.

INFRA-SPINATUS, *Infra-spinatus*, from *infra*, 'beneath,' and *spina*, 'spine.' Situate beneath the spine of the scapula.

INFRA-SPINATA FOSSA, (F.) *Fosse sous-épineuse*. A large excavation on the posterior surface of the scapula, beneath its spine. It is filled by the

INFRA-SPINATUS Muscle, *Grand Scapulo-trachitærius*, *Superseapularis inferior* (Ch.), (F.) *Sous-épineuse*, which is broad, flat, and triangular. It is attached, by its base, to the three inner quarters of the fossa; and is inserted, by a long tendon, into the middle part of the great tuberosity of the os humeri (*Trochiter*). It turns the arm outwards, and, when the arm is elevated, carries it backwards.

INFRINGENS, Corrigent.

INFUNDIBULUM, (in, and *fundere*, *fusum*, 'to pour out.'). A Funnel, *Trito-rium*, *Chod-ne*, *Chanos*, (F.) *Entomoir*,—of which various kinds are employed in pharmaceutical operations. A name, given to many parts which, more or less, resemble a funnel. See Proventriculus. It is particularly appropriated to the following organs:—

INFUNDIBULUM OF THE BRAIN, *Infundibulum seu Pelvis seu Chod-na seu Cyathus seu Scyphus seu Concha seu Lacin-na seu Embolus seu Aqueductus seu Labrum seu Concretus emchulæris seu Processus orbicularis Cerebri*, (F.) *Entomoir du ventricule moyen du cerveau*, *Tige Pituitaire*, *Tige sus-phénoïdale*. A depression in the inferior paries of the middle ventricle, above the pituitary gland. It was, anciently, regarded as a canal by which the fluid, collected in the ventricles of the brain, was evacuated, and poured into the nasal fosse.

INFUNDIBULUM CEREBRI, I. of the Brain.

INFUNDIBULUM OF THE COCHLEA, *I. Cochleæ*, *Scyphus Vienssenii seu auditorius*, *Cucullus*, *Caulis Scalærum communis*. This, with the modiolus, forms the nucleus around which the gyri of the cochlea pass. It is an imperfect funnel, the apex of which is common with that of the modiolus; and the base is covered with the apex of the cochlea, termed *Cupola*.

In *Surgery*, infundibula or funnels are used to direct steam or vapours; to conduct the actual cautery to certain morbid parts, &c.

INFUNDIBULUM OF THE ETMOID BONE, or OF THE NASAL FOSSÆ. One of the anterior cells of that bone, which is broad and expanded above, and narrow below; opening, above, into the frontal sinus; below, into the anterior part of the middle meatus of the nasal fossæ.

INFUNDIBULUM OF THE HEART, *Conus arteriosus*—I. of the Kidney, see *Calix*—I. Lacrymale, Lacrymal Sac—I. Tubarum Fallopii, see *Tube*, Fallopiæ—I. Ventriculi, *Œsophagus*.

INFUSË, Infusion.

INFUSION, *Infusio*, *Enchysis*, from *infundere* (in, and *fundere*, *fusum*), 'to pour in,' 'to introduce.' A pharmaceutical operation, which consists in pouring a hot or cold fluid upon a substance whose medical virtues it is desired to extract. *Infusion* is, also, used for the product of this

operation. (F.) *Infusion*, *Infusé*. In *Surgery*, infusion—*Chirurgia infusoria*, *Ars clysmatica nova*—is the act of introducing into the veins medicinal substances, by aid of an instrument called *Infusor*. See *Infusoir*. This mode of introducing medicines was called *Ars infusoria*.

INFUSION OF ANGUSTURA, Infusum Cuspariæ—I. of Bark, Infusum cinchonæ—I. of Buchu, Infusum Diosmæ—I. of Calumba, Infusum calumbæ—I. of Cascarella, Infusum cascarillæ—I. of Catechu, Infusum catechu compositum—I. of Cayenne pepper, Infusum capsici—I. of Chamomile, Infusum anthemidis—I. of Cinchona, Infusum cinchonæ—I. of Cloves, Infusum caryophyllorum—I. of Cusparia, Infusum cuspariæ—I. of Dandelion, Infusum Taraxaci—I. of Foxglove, Infusum digitalis—I. of Gentian, compound, Infusum gentianæ compositum—I. of Ginger, Infusum Zingiberis—I. of Hops, Infusum humuli—I. of Horseradish, compound, Infusum armoraciæ compositum—I. of Linseed, Infusum lini compositum—I. of Matico, Infusum matico—I. of Mint, compound, Infusum menthæ compositum—I. of Orange-peel, compound, Infusum aurantii compositum—I. of Pareira Brava, Infusum Pareiræ—I. of Pinkroot, Infusum spigeliæ—I. of Quassia, Infusum quassia—I. of Rhatany, Infusum Krameriæ—I. of Rhubarb, Infusum rhei—I. of the Rose, Infusum rosæ compositum—I. of Sarsaparilla, Infusum sarsaparillæ—I. of Sassafras pith, Infusum sassafras medullæ—I. of Seneca, Infusum Senegæ—I. of Senna, Infusum sennæ compositum—I. of Simaruba, Infusum simarubæ—I. of Slippery elm, Infusum ulmi—I. of Thoroughwort, Infusum eupatorii—I. of Tobacco, Infusum tabaci—I. of Valerian, Infusum valerianæ—I. of Virginia snakeroot, Infusum serpentariæ—I. of Wild cherry, Infusum pruni Virginianæ.

INFUSOIR (F.), *Infusor*. An instrument for injecting medicinal substances into the veins. It was a kind of funnel, the elongated apex of which was stopped by a metallic rod, which could be withdrawn when the apex was introduced into a vein.

INFUSUM, *Enchyma*. The product of an infusion.

INFUSUM ACACIÆ CATECHU, I. catechu compositum—I. Amarum vinosum, Vinum gentianæ compositum—I. of Angustura, I. cuspariæ.

INFUSUM ANTHEMIDIS, *Hydrochamaimeum*, Infusion of Chamomile. (*Anthemid. flor.* 3ss; *aq. bullient.* Oj. Macerate for ten minutes in a covered vessel, and strain.) Dose, f3j to f3iiss.

INFUSUM ARMORACIÆ COMPOSITUM, I. Armoraciæ (Ph. U. S.), Compound Infusion of Horseradish. (*Armorac. concis.*, *sinapis cont.*, sing. 3j; *aqua bullient.* Oj. Macerate for two hours, and strain.) Dose, f3j to f3iij.

INFUSUM AURANTII COMPOSITUM, Compound Infusion of Orange Peel. (*Aurant. cort. sicc.* 3ij; *limon. cort. recent.* 3j; *caryoph.* cont. 3ss; *aq. fervent.* Oss. Macerate for fifteen minutes, and strain.—Ph. L.) Dose, f3iiss to f3ij.

INFUSUM BRASII, Wort—I. Buchu, Infusum Diosmæ—I. Byones, Wort.

INFUSUM CALUMBÆ, I. Calumbæ seu Colombæ, Infusion of Calumba. (*Calumb. rad. concis.* 3ss; *aq. fervent.* Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f3iiss to f3ij.

INFUSUM CAPSICI, Infusion of Cayenne Pepper. (*Capsic. in. pulv. crass.* 3ss; *aq. bullient.* Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f3iiss.

INFUSUM CARYOPHYLLI, Infusion of Cloves. (*Caryoph. contus.* 3ij; *aq. bullient.* Oj. Macerate for two hours, and strain.) Dose, f3iiss to f3ij.

INFUSUM CASCARILLÆ, Infusion of Cascarella. (*Cascarill. contus.* 3j; *aq. bullient.* Oj. Macerate for two hours, and strain.) Dose, f3iiss to f3ij.

rate for two hours, and strain.) Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ij.

INFUSUM CASSIÆ SENNÆ, *I. sennæ compositum*.

INFUSUM CAT'ECU COMPOS'ITUM, *I. Catechu seu Aca'cia Catechu, Infusion of Catechu*. (Catechu, in pulv. $\frac{3}{4}$ ss; cinnam. cont. $\frac{3}{4}$; aq. bullient. Oj. Macerate for an hour, and strain.) Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ij.

INFUSUM CINCHO'NÆ, *I. Cinchonæ lancifoliae, Infusion of Cinchonæ*. (Cinch. contus. $\frac{3}{4}$ ij; aq. fervent. Oj. Macerate for two hours, and strain.) This infusion may also be made from the same quantity of bark in coarse powder by the process of displacement with hot or cold water. Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij. The Pharmacopœia of the United States has an *Infusum Cinchonæ flavæ, Infusion of yellow bark*, and an *Infusum Cinchonæ rubræ, Infusion of red bark*, which are prepared in the same manner. It has also an *Infusum Cinchonæ Compositum*, which is made as follows: (Cinchon. rubr. in pulv. $\frac{3}{4}$ ij; Acid. Sulph. aromat. f $\frac{3}{4}$ ij; aquæ, Oj. Macerate for twelve hours, occasionally shaking, and strain.) Dose, same as the last.

INFUSUM CINCHONÆ COMPOSITUM, see *Infusum Cinchonæ*—i. *Cinchonæ flavæ*, see *Infusum Cinchonæ*—i. *Colombæ*, i. *calumbæ*.

INFUSUM CUSPARIÆ, *I. Angusturæ, Infusion of Cusparia*. (Cuspar. cort. contus. $\frac{3}{4}$ ss; aq. bullient. Oj. Macerate for two hours, and strain.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM DIGITA'LIS, *I. Digitalis purpureæ, Infusion of Foxglove*. (Digital. $\frac{3}{4}$ ij; aq. bullient. Oss; tinct. cinnamomi, f $\frac{3}{4}$ ij. Macerate for two hours, and add sp. cinnam. f $\frac{3}{4}$ ij.) Dose, f $\frac{3}{4}$ ij.

INFUSUM DIOS'MÆ, Ph. U. S., 1842, *I. Buchu*, Ph. U. S., 1851, *Infusion of Buchu*. (Diosm. $\frac{3}{4}$ ij; aq. bullient. Oj. Macerate for two hours in a covered vessel, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss.

INFUSUM EUPATO'RII, *Infusion of Thoroughwort*. (Eupator. $\frac{3}{4}$ ij; aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij.

INFUSUM GENTIANÆ COMPOS'ITUM, *Compound Infusion of Gentian*. (Gentian. cont. $\frac{3}{4}$ ss; Aurant. cort., Coriandr. contus. $\frac{3}{4}$ ss; alcohol. dilut. f $\frac{3}{4}$ iv; aquæ, f $\frac{3}{4}$ xij. First pour on the alcohol, and three hours afterwards, the water; then macerate for twelve hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

Marsden's *Anticorbutic Drops*, an empirical preparation, consist of a solution of corrosive sublimate in an infusion of gentian.

INFUSUM HUMULI, *I. Lu'puli, Infusion of Hops*. (Humul. $\frac{3}{4}$ ss; aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ijij.

INFUSUM JUNIP'ERI, *Infusion of Juniper*. The common mode of making this infusion is by an ounce of *Juniper Berries* to a pint of boiling water. The whole may be taken in the twenty-four hours.

INFUSUM KRAMERIÆ, *Infusion of Rhat'any*. (Kramer. contus. $\frac{3}{4}$ ij; aq. bullient. Oj. Macerate for four hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss.

INFUSUM LINI, *I. L. compositum*.

INFUSUM LINI COMPOS'ITUM (Ph. U. S., 1851), *I. Lini* (Ph. U. S., 1842), *I. Lini usitatissimi, Infusion of Linseed, Flaxseed Tea*. (Lini sem. cont. $\frac{3}{4}$ ss; glycyrrh. rad. cont. $\frac{3}{4}$ ij; aquæ bullient. Oj. Macerate for two hours, and strain.) Dose, a teacupful, ad libitum.

INFUSUM LUPULI, *Infusum Humuli*—i. *Malti Wort*.

INFUSUM MATICO, *Infusion of Matico*. (Matico, $\frac{3}{4}$ ss (avoirdupois); aq. bullient. Oss (Imp. meas.). Infuse for an hour, and strain.—Ph. D.) Dose, f $\frac{3}{4}$ ss to f $\frac{3}{4}$ ij.

INFUSUM MENTHÆ VIRIDIS, *Infusion of Spearmint, Mint Tea*. (Menthæ virid. $\frac{3}{4}$ ij; aq. bullient. Oss. Infuse for an hour, and strain.—Ph. D.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM PAREIRÆ, *Infusion of Paraira Brava*. (Pareir. $\frac{3}{4}$ ij; aq. bullient. Oj (Imp. meas.). Infuse for two hours, and strain.—Ph. E.) Dose, f $\frac{3}{4}$ ij or f $\frac{3}{4}$ ijij.

INFUSUM PICIS EMPYREUMATICÆ LIQUIDÆ, see *Pinus sylvestris*—i. *Picis liquidum*, see *Pinus sylvestris*—i. *Polygalæ, Infusum Senegæ*.

INFUSUM PRUNI VIRGINIÆ, *Infusion of Wild Cherry Bark*. (Prun. Virginian. cont. $\frac{3}{4}$ ss; aquæ, Oj. Macerate for 24 hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM QUASSIÆ, *I. Quassia excel'sæ, Infusion of Quassia*. (Quassia lign. cont. $\frac{3}{4}$ ij; aq. (frigid.) Oj. Macerate for twelve hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM RHEI, *Antima Rhei, Infusion of Rhubarb*. (Rhei, cont. $\frac{3}{4}$ ij; aq. ferv. Oss. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM ROSÆ COMPOS'ITUM, *I. Rosæ Gallicæ, Infusion of the Rose*. (Ros. Gallic. $\frac{3}{4}$ iv; aq. bullient. Ojss; acid. sulph. d. f $\frac{3}{4}$ ijij; succh. purif. $\frac{3}{4}$ ss. Add the water, and afterwards the acid;—macerate for half an hour; strain, and add the sugar.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss to Oss.

INFUSUM SARSAPARILLÆ, *Infusion of Sarsaparilla*. (Sarsaparill. contus. $\frac{3}{4}$ ij; aquæ bullient. Oj. Digest for two hours in a covered vessel, and strain.—Ph. U. S.) It may also be prepared by displacement. Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM SASS'AFRAS, *I. Sassafras medullæ, Infusion of Sassafras Pith, Mucilage of Sassafras Pith*. (Sassafras medull. $\frac{3}{4}$ ij; aquæ (frigid.) Oj. Macerate for three hours, and strain.—Ph. U. S.) An emollient collyrium, and a demulcent drink.

INFUSUM SENEGÆ, *Infusum Polygalæ, Infusion of Senega*. (Seneg. $\frac{3}{4}$ x; aq. bullient. (Imp. meas.). Infuse for four hours, and strain.—Ph. E.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM SENNÆ, *I. S. Compositum, I. Cassiæ Sennæ seu Senna simplex, Infusion of Senna*. (Sennæ, $\frac{3}{4}$ ij; coriandr. cont. $\frac{3}{4}$ ij; aq. bullient. Oj. Macerate for an hour, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

The *Black Draught, Black Dose, Black bottle, Haustus niger*, is usually formed of this infusion. It may be made of infus. sennæ, f $\frac{3}{4}$ v; aq. cinnam. f $\frac{3}{4}$ ij; mannæ, $\frac{3}{4}$ v; magnes. sulph. $\frac{3}{4}$ v. Dose, a wineglassful.

Schwab's *Prepared Essence of Senna* is a concentrated infusion of the leaves in combination with an alkali.

INFUSUM SENNÆ COMPOSITUM, *I. sennæ*.

INFUSUM SERPENTARIÆ, *Infusion of Virginia Snakeroot*. (Serpentar. $\frac{3}{4}$ ss; aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ ss.

INFUSUM SIMARUBÆ, *Infusion of Simaruba*. (Simarub. cort. cont. $\frac{3}{4}$ ij; aq. fervent. Oj. Macerate for two hours, and strain.—Ph. E.) Dose, f $\frac{3}{4}$ ij to f $\frac{3}{4}$ ijij.

INFUSUM SPIGELIÆ, *Infusion of Pinkroot*. (Spigel. $\frac{3}{4}$ ss; aq. bullient. Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f $\frac{3}{4}$ iv to Oss.

INFUSUM TAB'ACI, *Infusion of Tobacco*. (Tabaci fol. $\frac{3}{4}$ ij; aq. ferv. Oj. Macerate for an hour, and strain.—Ph. U. S.) Half of this per anum in strangulated hernia.

INFUSUM TARAX'ACI, *Infusion of Dan'delion*.

(*Taraxac. contus.* 3ij; *aq. bullient.* Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f3iiss, as a diuretic, &c.

INFUSUM ULMI, *Infusion of Slip'pery Elm, Slippery Elm Tea.* This preparation, in the Pharmacopœia of the United States, is made by infusing one ounce of *slippery elm bark* in a pint of boiling water.

INFUSUM VALERIANÆ, *Infusion of Valerian.* (*Rad. valerian.* in crass. pulv. 3ss; *aq. bullient.* Oj. Macerate for an hour, and strain.—Ph. U. S.) Dose, f3iiss to 3ij.

INFUSUM ZINGIBERIS, *Infusion of Ginger, Ginger Tea.* (*Zingib. contus.* 3ss; *aq. bullient.* Oj. Macerate for two hours, and strain.—Ph. U. S.) Dose, f3iiss to f3ij, as a carminative.

INGA. A genus of plants, *Ord. Leguminosæ*, belonging to South America, many of the species of which are possessed of astringent properties. *Inga bark* is used in South America as a tonic astringent in diarrhoea, gonorrhœa, &c. An extract, prepared from it, possesses all the properties of rhatany.

INGENIUM (*Morbi*). The nature or genius of a disease, from *in*, and *geno*, for *gigno*, 'to generate.' This word is employed, especially by the French, synonymously with *nature*. They speak, for instance, of *Génie inflammatoire, bilieux*, &c. Some, also, use *génie* in the place of *type* of an intermittent.

INGESTA, from *in*, and *gerere, gestum*, 'to hear or carry into.' Substances, introduced into the body by the digestive passages; as food, condiments, drinks, &c.

INGLUVIES, *Apletia, Victis intemperantia*.—Gluttony, Insatiableness; also, the *Crop or Gorge of Birds*, (*F. Jabot*; and the *first stomach or pouch of ruminant animals*—*Pen'ula, Rumex, Venter magnus*, (*F. Panse, Herber, Double*. Also, the Pharynx.

INGRASCIAS, APOPHYSES OR. The lesser alæ of the sphenoid bone.

INGRAVIDATIO, *Fecundation, Pregnancy.*

INGRAVIDATION, *Fecundation, Pregnancy.*

INGRESSUS SUPERIOR, *Cardia.*

INGUEN, *Bubon, Bubo*, 'the groin,' (*Prov. Lank, Laet, Lisk, Lesk; Ædæon, Hepatis emunctoria, (F.) Aine.* The oblique fold or depression which separates the abdomen from the thigh. It is only, properly speaking, a line that extends from the anterior and superior spinous process of the ilium to the middle part of the horizontal ramus of the pubis. Also, the genital organs.

INGUINAL, *Inguinalis*, from *inguen, inguinis*, 'the groin.' Belonging or relating to the groin. This epithet has been given to various parts met with in the region of the groin or inguinal region.

INGUINAL ARTERY is that portion of the femoral artery situate immediately beneath the crural arch in the inguinal region.

INGUINAL CANAL is a canal, about two inches in length, proceeding obliquely downwards, inwards and forwards at the lower part of the abdomen; through which passes the spermatic cord, in men, and the round ligament of the uterus in women. This canal is formed, inferiorly and anteriorly, by the aponeurosis of the greater oblique muscle; posteriorly by the *fascia transversalis*, which is joined to the preceding aponeurosis, and forms with it a deep channel, into which are received the lesser oblique and transversalis muscles. The inguinal canal has two apertures; the one, the *lower and inner*, is called the *inguinal or abdominal ring*. It is bounded by two strong tendinous pillars which fix it—the innermost to the symphysis, the outermost to the spine of the pubis. The upper and outer aperture is formed

by the fascia transversalis. From the edge of this aperture arises a membranous funnel,—a prolongation of the fascia transversalis,—which receives the spermatic vessels; forms their proper sheath, and accompanies them as far as the testicle. On its inside lies the epigastric artery. Above it, is the lower edge of the transversalis muscle: and, below, it is bounded by the channel of the greater oblique. By following the oblique direction of this canal, and passing, consequently, on the outside of the epigastric artery, the viscera are displaced, so as to constitute internal inguinal hernia.

INGUINAL RE'GION, *Re'gio inguinalis, Bubo, Inguen.* The region of the groin.

INGUINAL RING, *Abdominal Ring, An'nulus abdominis*, (*F. Anneau Inguinal*), is the inferior aperture of the inguinal canal.

INGUINO-CUTA'NEUS. A name given by Professor Chaussier to the middle ramus of the anterior branch of the first lumbar nerve, because it sends its numerous filaments to the groin, scrotum, and to the skin of the superior part of the thigh.

INHÆRENS, *Inherent.*

INHALATIO, *Absorption, Inhalation*—i. *Cutis*, see *Absorption*.

INHALATION, *Inhalatio*; from *in*, and *halare*, 'to breathe.' The act of drawing air or vapour into the lungs—*Inhalatio pulmonalis*. Also, *absorption*.

INHA'LER. Same etymon. An apparatus for inhalation. *Mudge's Inhaler* is an apparatus for inhaling the steam of hot water, in affections of the air-passages. It consists of a pewter tankard provided with a lid, into which a flexible tube is inserted. Through this the vapour is inhaled.

INHE'RENT, *Inhærens*, (*in*, and *harco*, 'I stick.') That which adheres, or which is joined or united to any thing.

INHERENT CAUTERY, (*F. Cautére inhérent*, is the actual cautery, left in contact with a part until it is reduced to the state of a deep eschar.

INHUMATION, *Inhumatio*, from *inhumo, inhumatus*, (*in*, and *humus*), 'I put into the ground.' The sepulture of the dead. This belongs to the subject of medical police. Also, digestion of substances in earth or dung. See *Arcuatio*.

INIAD, see *Inial*.

IN'IAL, from *iniov*, 'the nape of the neck.' An epithet, proposed by Dr. Barclay, for an aspect towards the plane of the ridge of the occiput. *Iniad* is employed by him adverbially to signify 'towards the inial aspect.'

INIENCEPH'ALUS, (*F. Iniencephale*, from *iniov*, 'occiput,' and *εγκεφαλος*, 'encephalon.' A monster whose encephalon is in great part in the cranium, and in part out of it, behind, and a little beneath the cranium, which is open in its occipital portion.—I. G. St. Hilaire.

INIODYMUS, *Diprosopus*.

INION, *iniov*. Some of the Greek physicians give this name to the occiput, or the ridge of the occiput; others to the back part of the neck, and the muscles of the occiput. Blancard says it is the commencement of the spinal marrow.

INIOPS, *Iniopæ*, from *iniov*, 'occiput,' and *ωψ*, 'the eye, face.' A monster, double above the umbilicus, which has two ears very near each other, or only one median ear, and above it, at a greater or less distance, a median orbit and eye, more or less imperfect. The rudimental face appearing to be directly opposite the principal face suggested the name.—I. G. St. Hilaire.

INI'TIS, *Inohymenitis*, from *is*, gen. *ivos*, 'a fibre,' and *itis*, a suffix denoting inflammation. Fibrous inflammation.

INITIUM, Arche.—i. *Asperæ arteriæ*, Larynx
—i. *Extubercans Coli*, Cæcum.

INIUM, *inior*. The nucha. Also, the muscles
at the back of the neck. See Inion.

INJACULA'TIO, *in*, and *jaculor*, *jaculatus*,
'to shoot.') A term employed by Van Helmont
to designate an acute pain of the stomach, with
rigidity and immobility of the body.

INJECT'ED, *Injec'tus*, from *injacere*, *injec'tum*,
(*in*, and *jacere*, 'to throw into.') The face and
other parts are said to be *injected*, when the accu-
mulation of blood in the capillary vessels gives
them an evident red colour. A subject or part
of a subject is also said to be *injected*, when its
vessels have been filled, by the anatomist, with
an appropriate composition.

INJECTION, *Injec'tio*, *Eis'bolê*: same etym-
on. The act of introducing, by means of a sy-
ringe or other instrument, a liquid into a cavity
of the body. The liquid injected is also called an
injection. Anatomists use injections—*Injec'tio-
nes anatom'icæ*—for filling the cavities of ves-
sels, in order that they may be rendered more
apparent, and their dissection be facilitated. For
this purpose, they employ syringes of different
dimensions, and various materials. The most
common injections are made of soot, wax, and
turpentine, coloured with lamp-black, vermilion,
&c. There are three kinds chiefly used by ana-
tomists,—the *coarse*, the *fine*, and the *minute*.
The following are formulæ for each.

COARSE INJECTION.

No. 1.

Pure yellow wax, oz. xvj.
Bleached rosin, oz. viij.
Turpentine varnish, by measure, oz. vj.

No. 2.

Yellow rosin, lb. ij.
Yellow wax, lb. j.
Turpentine varnish, a sufficient quantity to
make the mixture flexible when cold.

No. 3.

Tallow, lb. ij.
White wax, oz. x.
Common oil, oz. vj.
Venice turpentine, oz. iv.
Mix and liquefy over a slow fire or over boiling
water.

To make any of these mixtures.

Red—add vermilion, oz. ij.
Yellow—King's yellow, oz. iiss.
White—best flake white, oz. vss.
Pale-blue { best flake white, oz. iiiss.
 { fine blue smalt, oz. iiiss.
Dark-blue—blue verditer, oz. xss.
Black—lamp-black, oz. j.
Green { powdered verdigris, oz. ivss.
 { best flake white, oz. iiss.
 { powdered gamboge, oz. j.

FINE INJECTION.

Brown spirit varnish, oz. iv.
White spirit varnish, oz. iv.
Turpentine varnish, oz. j.

To make this mixture,

Red—add vermilion, oz. j.
Yellow—King's yellow, oz. j̄.
White—best flake white, oz. ij.
Light-blue { fine blue smalt, oz. iiss.
 { best flake white, oz. j̄.
Dark-blue—blue verditer, oz. iv.
Black—lamp-black, oz. ss.

MINUTE INJECTION.

Take of transparent size, broken to pieces, or
Isinglass, oz. viij.
Water, lb. iiss. Dissolve.

To make this mixture,

Red—add vermilion, oz. v.
Yellow—King's yellow, oz. iv.
White—best flake white, oz. v.
Blue—fine blue smalt, oz. viij.
Green { powdered verdigris, oz. iij.
 { best flake white, dr. ij.
 { powdered gamboge, dr. j.
Black—lamp-black, oz. j.

Beautiful injections, are made with ether as
the menstruum.

COLD INJECTION.

White lead and red lead, each, oz. iv; linseed
oil, enough to form a thick paste when they
are rubbed well together. Liquefy this paste
with turpentine varnish, oz. viij.

The advantage of this mixture is, that the sub-
ject need not be heated.

In order to inject the arteries, the injection
must be forced from the great trunks towards
their ultimate ramifications. To inject the veins,
on the contrary, it is indispensable, on account of
their valves, to send the injection from the smaller
divisions towards the greater. The lymphatics
are usually injected with mercury. The practi-
tioner injects, by forcing with a syringe, liquids,
such as emollient, narcotic, stimulant, and other
decoctions or infusions, into different hollow or-
gans, as the rectum, vagina, nasal fossæ, urethra,
tunica vaginalis, auditory canal, &c., to fulfil va-
rious therapeutical indications.

The following injection has been strongly re-
commended by Dr. Horner to preserve the dead
body.

Take of Liverpool, St. Ubes,
or Turk's Island Salt, oz. 36 avoidr.
Nitrate of potassa, " 19
Carbonate of Soda, " 8
Molasses (Sugar-house), " 4 by meas.
Water, six pints.

The saline constituents to be dissolved first of
all in boiling hot water; the molasses to be af-
terwards stirred in; the starch to be mixed well with
half a pint of cold water, and then to be stirred
in with the other articles. As soon as it begins
to boil, the whole mass swells up, when it must
be removed from the fire. On the proper reduc-
tion of temperature it is fit for use.

INJECTION, MATTHEW'S, see *Tinctura cantha-
ridis*.

INJECTIONES ANATOMICÆ, see Injection.

INK, Atramentum.

INKBERRY, *Prinos glaber*.

IN-KNEED, *Entogonyancon*.

IN'NATE, from *in*, and *nascor*, *natus*, 'I am
born.' Inborn.

INNATE DISEASES, *Morbi conna'ti seu congen-
iti*, (F.) *Maladies innées*. Diseases with which
the infant is born. They are not always heredi-
tary, as hereditary diseases are not always innate.

INNERLEITH'EN, MINERAL WATERS
OF. These springs, situate near the Tweed, and
supposed to be the scene of "St. Ronan's Well,"
contain chlorides of sodium and calcium, and
carbonate of magnesia.

INNERVA'TION, *Innerva'tio*, from *in*, 'in,'
and *nervus*, 'a nerve.' By this term is meant—
the nervous influence, necessary for the mainte-
nance of life and the functions of the various or-
gans;—an influence of whose character and source

we are ignorant. It seems to resemble the galvanic or electric agencies. See Nerves.

INNOMINATUM, *Anonymum*, from *in*, 'negation,' and *nomen*, 'a name.' (F.) *Anonyme*. Having no name.

INNOMINATA ARTERIA, *Brachio-cephalic artery*, *Arteria anonyma*, *Right Subclavian*, (F.) *Artère innominée*, *Artère brachio-céphalique* (Ch.), is the trunk common to the right primitive carotid and to the subclavian. It arises from the anterior part of the arch of the aorta, ascends obliquely to the right, along the trachea; and after a course of about an inch in length, divides into two trunks, which go off at right angles. The one is *external*—the *right subclavian proper*; the other *superior*—the *primitive carotid*, of the same side.

INNOMINATA CARTILAGO, *Cricoid*.

INNOMINATA CAVITAS. A cavity of the outer ear, between the *helix* and *anthelix*.

INNOMINATUM FORAMEN. A foramen, near the middle of the anterior surface of the pars petrosa of the temporal bone, leading backwards for the passage of the Vidian nerve, reflected from the 2d branch of the 5th to the portio dura of the 7th pair.

INNOMINATA FOSSA, see Fossa—i. Linea, see *Lio-pectinea Linea*.

INNOMINATI seu ANONYMI NERVI. Some anatomists have thus called the nerves of the fifth pair.

INNOMINATUM OS, *Os Corendicis* seu *Coxæ* seu *anonymum* seu *pelvis lateralis*, (F.) *Os innominé*, *Os Coxal*, *Os anonyme*. A very large, flat bone, which forms the anterior and lateral paries of the pelvis. It is curved upon itself in two opposite directions. In the first periods of life, it is composed of three portions;—the *ilium*, *ischium*, and *pubis*, which join each other in the acetabulum. It is articulated *before* with its fellow, —*behind* with the sacrum; and *laterally* with the femur.

INNOMINATA MINORA OSSA, *Lesser Ossa innominata*, *Ossicula innominata*. Some anatomists have given this name to the three cuneiform bones of the tarsus.

INNOMINATA OSSICULA, *Innominata minora ossa*—i. *Tunica Oculi*, *Sclerotic*.

INNOMINATÆ VENÆ, of VIEUSSENS. Vieussens has given this name to two or three veins, which arise on the anterior surface and right margin of the heart, and open into the auricle towards its right margin. The term VENÆ INNOMINATÆ of MECKEL is given to the *brachiocephalic* veins, which are generally included in the description of the subclavian vein, and correspond to the *arteria innominata*, being formed by the union of the internal jugular vein and the subclavian properly so called, which correspond to the common carotid and subclavian arteries.

INNOMINATUS, *Anonymous*.

INNUTRITIO OSSIUM, *Rachitis*.

INOCULABLE. Same etymon as the next. Capable of inoculation, or of being communicated by inoculation.

INOCULATION, *Inoculatio*, *Insit'io*, *Insit'io variolarum*, from *inoculare*, (*in*, and *oculus*, 'an eye,') 'to ingraft.' Any operation by which small-pox, for example, may be artificially communicated, by introducing the virus of the particular disease into the economy, by means of a puncture or scratch made in the skin. When the word inoculation is used alone, it usually means that for the small-pox,—*Variolation*. See *Syphilization*.

INOCULATION, *Cowpox*, *Vaccination*—i. *Jennian*. *Vaccination*.

INOCULATOR, same etymor. *In'sitor*. One who practises inoculation.

INODULAR TISSUE, see *Tissæ*, *inodular*.

INODULE, *Tissue*, *inodular*.

INOHYLOMA, *Tumour*, *fibrous*.

INOHYMENT'IS: from *is*, gen. *ivos*, 'a fibre;' *μην*, 'a membrane,' and *itis*, denoting inflammation. *Inflammatio telæ fibro-æ*. Inflammation of the fibrous tissue.

INOMA, *Tumour*, *fibrous*.*

INONCTION, *Inunction*.

INOPLYTUS, see *Polypus*.

INORGAN'IC, *Inorgan'icus*, *Unorganiz'ed*, (F.) *Inorganique*; from *in*, 'negation,' and *organum*, 'an organ.' A term applied to bodies which have no organs;—such as minerals. At the present day, naturalists admit of but two classes of bodies,—the *organized* and *inorganic*. Parts of the body which, like the epidermis, are devoid of blood-vessels and nerves, have been called *anorganic*.

INOSCLERO'MA, from *is*, gen. *ivos*, 'a fibre,' and *σκληρωμα*, 'induration.' Induration of the fibrous tissue.

INOSCULATIO, *Anastomosis*.

INOSIS, *Hyperinosis*.

IN'OSITE, *In'osit*, *Muscle sugar*; from *is*, *ivos*, 'a fibre.' An unfermentable sugar, found only in the juice of the cardiac muscle.—Lehmann.

INOSTEATO'MA, from *is*, *ivos*, 'a fibre,' and *stentoma*. A tumour of a mixed fibrous and steatomatous character.

INQUIES, *Inquietude*.

INQUIETATIO, *Inquietude*.

INQUIETUDE, *Inquietudo*, *In'quies*, *Inquietatio*, *Jactitatio*, from *in*, 'negation,' and *quies*, 'rest.' Agitation or trouble, caused by indisposition. Restlessness.

INQUINAMENTUM, *Miasm*.

INQUISITIO MEDICO-LEGALIS, see *Medico-legal*.

INSALIVATION, *Insalivatio*, from *in*, and *saliva*. The mixture of the food with the saliva, and other secretions of the mouth.

INSALUBRIOUS, *Insalu'bris*, *Nose'ros*, *No'sodes*. That which is unhealthy,—which injures the health.

INSANABILIS, *Incurable*.

INSANE, *Insanus*; from *in*, 'negation,' and *sanus*, 'sound;' *Alienatus*, *Crazy*, *Mad*, *Nonsane*, *Demented*, *Deranged*, *Bedlamite*, (Sc.) *Daft*, *Wod*, (F.) *Aliéné*, *Fou*, *Insensé*. One affected with mental aberration, or of unsound mind.

INSANTA, *Mania*—i. *Cadiva*, *Epilepsy*—i. *Lupina*, *Lycanthropia*—i. *Puerperarum*, *Mania*, *puerperal*.

INSANITAS, *Insanity*.

INSANITY, *Insania*, *Insan'itas*. Same etymon. *Mental alienation*, *Abalienatio* seu *Alienatio Mentis*, *Arreptio*, *Unsound Mind*, *Derangement*, *Deranged intellect*, *Craziness*, (Sc.) *Duffin*, *Duffing*, *Wodnes*; *Aphros'yne*, *Ephiro'nia*, *Eephros'yne*, *Paral'lagé*, *Parallax'is*, *Delir'ia*, *Vesania*, *Delir'ium* (Crichton), *Insuper'entia*, (F.) *Folie*, *Égarement d'Esprit*, *Paraphrénie*. This term includes all the varieties of unsound mind,—*Mania*, *Melancholia*, *Moral Insanity*, *Dementia*, and *Idiocy*. A slight degree of insanity is sometimes popularly called "a kink in the head;" in Scotland, "a bee in the bonnet."

INSANITY, *HOMICIDAL*, see *Homicidal*—i. *Incoherent*, *Dementia*—i. *Moral*, *Pathomania*—i. *Puerperal*, *Mania*, *puerperal*—i. *Religious*, *Sebatomania*—i. *Senile*, *Delirium senile*—i. *Suicidal*, see *Suicide*.

INSANUS, *Insane*.

INSCRIPTION, see *Matriculate*.

INSCRIPTIO'NES TENDIN'Æ MUSCULORUM, *Intersectio'nes seu Éneratio'nes tendin'æ musculorum*, *Intermedia ligamenta'lia seu nerv'ea*. The tendinous portions which cross

several muscles, and especially the straight muscles of the abdomen.

INSEMINATION, see Ejaculation.

INSENESCENCIA, Agerasia.

INSENSE, Insano.

INSENSIBILITY, *Anæsthesia, Insensibilitas*. Loss or absence of sensibility. It is very common in cerebral affections, and may extend to every part, or be limited to one or more. Some organs are much more sensible than others. The bones, cartilages, ligaments, &c., are insensible in health, but acutely sensible in disease.

INSENSIBLE, *Sensibilitas carens*. That which is devoid of sensibility. This word is applied, also, to phenomena which cease to be appreciable to the senses. Thus, we say, the pulse becomes insensible.

INSERTION, *Inser'tio, Symphysis, Eno'sis*, from *inserere, insertum*, (in, and *serere*, to join or knit,) 'to ingraft.' (F.) *Attache*. The attachment of one part to another. Insertions occur chiefly on bones, cartilages, and fibrous organs: thus, we speak of the insertion of muscular fibres into a tendon or aponeurosis; the insertion of a tendon, aponeurosis, or ligament, into a cartilage or bone. The word *insertion* has likewise been used by pathologists for the act of inoculating or introducing a virus into the body.

INSES'SIO, from *insidere, insessum*, (in, and *sedere*,) 'to sit in.' This term is, sometimes, applied to a vapour bath, the person being seated in a perforated chair, beneath which a vessel, filled with hot water, or the hot decoction of some plant, is placed. See *Semcupium*.

INSESSUS, Bath, half.

INSICIUM, Isicium.

INSIDENTIA, *Epistaxis*. Any thing which swims on or in—the urine, for example. It is opposed to the *Hypostaxis* or *subsidentia*.

INSIPIENTIA, Insanity.

INSITIO, Inoculation—i. *Ciliorum, Blepharoplasticæ*—i. *Dentis, Transplantatio Dentis*—i. *Variolarum, Inoculation*.

INSOLATION, *Insolutio*, from *in*, and *sol*, the sun; *Apriectio, Heliosis, Heliasis, Heliorosis, Stris'asis*. Exposure to the sun. Exposure of a patient to the rays of the sun is, sometimes, had recourse to, with the view of rousing the vital forces when languishing, or of producing irritation of the skin. Insolation is occasionally used in the same sense as *coup de soleil*.

In Pharmacy, *insolation* means the drying of chemical and pharmaceutical substances.

INSOLAZIONE DE PRIMIVERA, Pel-lagra.

INSOMNIA, *Insomnitas, Sahara, Zavra, Perrigillum, Pernocatio, Apynia, Agrypnia, Apynia, Anpynia, Typhnodia, Sleeplessness, Vig'rance*, from *in*, 'negation,' and *somnus*, 'sleep,' 'absence of sleep.' This may exist alone, and constitute a true disease; or it may be connected with another affection. It is an unequivocal sign of suffering in some organ; even when the patient experiences no pain.

INSOMNIUM, Somnium.

INSPECTIO MEDICO-LEGALIS, see *Medico-legal*.

INSPIRATEUR, Inspiratory.

INSPIRATION, *Inspiratio, Empneumato'sis, Esp'noë, Adspiratio, Aspira'tio*, from *in*, 'in,' and *spiro*, 'I breathe.' The action by which the air penetrates into the interior of the lungs. A movement opposed to that of expiration. As regards the average quantity of air received into the lungs at each inspiration, there is much discrepancy amongst observers. The following table sufficiently exhibits this:—

Cubic inches at each inspiration.

Reil.....	42 to 100	
Menzies,	}	40
Sauvages,		
Hales,		
Haller,		
Ellis,		
Sprengel,		
Sömmering,		
Thomson,	}	35 to 38
Bostock,		
Jurin.....	35 to 38	
Fontana.....	35	
Richerand.....	30 to 40	
Dalton.....	30	
Jeffreys.....	26	
Herbst.....	21 to 30	
Herholdt.....	20 to 29	
Jurine and Coathupe.....	20	
Allen and Pepys.....	16½	
J. Borelli.....	15 to 40	
Goodwyn.....	14	
Sir H. Davy.....	13 to 17	
Abernethy and Mojon.....	12	
Keutsch.....	6 to 12	

INSPIRATION OF VENOUS BLOOD. By this is meant the aspiration of blood towards the heart, occasioned by the approach to a vacuum produced by the dilatation of the thorax during inspiration.

INSPIRATORY. Same etymon. *Inspiratio ni inser'viens*, (F.) *Inspirateur*. A name given to muscles, which, by their contraction, augment the size of the chest, and thus produce inspiration. The diaphragm and intercostal muscles are the chief agents of inspiration. In cases where deep inspirations are necessary, the action of these muscles is aided by the contraction of the pectoralis major and pectoralis minor, subclavius, serratus major anticus, scaleni, serratus posticus superior, &c. Most of these muscles become inspiratory, by taking their fixed point at the part which they ordinarily move, and elevating the ribs.

INSPISSANT, *Insipissans*. Same etymon as the next. Any remedial agent, that renders the blood thicker, directly, or indirectly. The opposite to diluent.

INSPISSATIO, *Pycno'sis*, from *in*, and *spissare, spissatum*, 'to thicken.' The act of rendering thick; as in the formation of an extract—*Succus Insipissatus*.

INSTEP, *Collum pedis, Neck-of-the-foot*, (F.) *Conde-pied, Cou-de-pied*. The projection at the upper part of the foot, near its articulation with the leg—the *tarsus*.

INSTILLATIO, Instillation.

INSTILLATION, *Enstalar'is, Instillo'tio*, from *in*, 'into,' and *stilla*, 'a drop.' The act of pouring a liquid drop by drop.

INSTINCT, (L.) *Instinctus*, (in, and *stinguo, stinctum*, 'to sting,' 'inwardly moved.' *Brutia, Hormæ*. The action of the living principle, whenever manifestly directing its operations to the health, preservation, or reproduction of a living frame or any part of such frame—*Mollimen nature salutarium*. The law of instinct is, consequently, the law of the living principle, and instinctive actions are the actions of the living principle. Instinct is natural. Reason is acquired.

INSTINCTIVE, *Instinctus*. Same etymon. Relating to or caused by instinct; as *instinctive* actions. See *Emotional*. Those instinctive actions of animals which are owing to impressions made on the sensory ganglia, exciting respondent motor influences that are propagated to the various muscles of the body, are termed *consequential*.

Consensual movements which accompany voluntary efforts without our consciousness are termed *associated*. (F.) *Associés*.

INSTITUTES OF MEDICINE, see Theory of medicine.

INSTITUTUM ORTHOPÆDICUM, Orthopedic institution.

INSTRUMENT, *Instrumen'tum*, *Or'ganum*. A tool, an agent.

INSTRUMENTA, Pudibilia.

INSTRUMENTUM CHIRURGICUM. A surgical tool or instrument.

INSTRUMENTUM DIGESTIONIS. The digestive apparatus.

INSTRUMENTUM INSTRUMENTORUM, Manus.

INSUFFICIENCY, (F.) *Insuffisance*; from *in*, and *sufficient*. Inadequateness to any end or purpose, — as *Insufficiency of the valves of the heart*, (F.) *Insuffisance des valvules du Cœur*; — a condition in which they are not adapted, as in health, to properly close the apertures.

INSUFFISANCE, Insufficiency.

INSUFFISANCE DES VALVULES DU CŒUR, Insufficiency of the valves of the heart.

INSUFFLATION, *Inufla'tio*, from *in*, *sub*, and *flare*, 'to blow.' The act of blowing a gas or vapour into some cavity of the body; as when tobacco smoke is injected into the rectum; or when air is blown into the mouths of new-born children to excite the respiratory functions.

INSULA, *Insula cerebri*.

INSULA CEREBRI, *Island* or *In'sula* of Reil. The intermediate lobe of the brain, *Lobus intermedius cerebri*. A remarkable group of convolutions within the fissure of Sylvius. It is called, by Cruveilhier, *Lobule of the Fissure of Sylvius*, *Lobule of the corpus striatum*. The 'island' of Reil, with the substantia perforata, forms the base of the corpus striatum.

INSULA SANGUINIS, see Blood.

INSULÆ, *Maculæ albæ* — i. *Pulmonales*, see Pulmo.

INSULTUS, Attack, Paroxysm.

INSURANCE OF LIFE, see Life Insurance.

INTEGRITAS, Sanitas, Virginity.

INTEGUMENT, *Integumen'tum*, *Tegumen'tum*, *Teg'men*, *Teg'men*, *Involutum* seu *Velamen'tum corporis communē*; *Velamen'tum* seu *Velamentum nativum*, from *in*, and *tegere*, *tectum*, 'to cover.' (F.) *Tégument*. Any thing which serves to cover, to envelop. The *skin*, including the cuticle, rete mucosum, and cutis vera, is the common integument or tegument of the body.

INTEGUMENTA FŒTUS. Membranes of the foetus.

INTEGUMENTUM, Integument — i. Common, Integument.

INTELLECT, *Intellectus*, *Nous*, *Mens*, *Gnomē*, *Noos*, *Nūs*, *Noc'is*, *Syn'esis*, from *intelligere*, *intellectum*, (*inter*, 'between,' and *legere*, 'to choose') 'to understand,' 'conceive,' 'know.' (F.) *Entendement*, *Intelligence*. The aggregate of the intellectual faculties — perception, formation of ideas, memory, and judgment.

INTELLECT, DERANGED, Insanity.

INTELLIGENCE, Intellect.

INTEMPERANCE, *Intemperan'tia*; *Immoderantia*, *Immoderatio*, *Acras'ia*, *Acolas'ia*, *Ples'monē*, *Ames'ria*, *Aples'ia*, from *in*, 'negation,' and *temperare*, 'to temper.' Immoderate use of food and drink, especially the latter; — a fruitful source of disease.

INTEMPERIES. Same etymon. *Dyscras'ia*. Derangement in the constitution of the atmosphere and of the seasons; bad constitution; derangement or disorder in the humours of the body.

INTENSIVUS, Entaticus.

INTENTION, *Inten'tio*, from *in*, and *tendere*,

tensum seu *tentum*, 'to stretch.' *Propositum*. The object which one proposes. In surgery, a wound is said to heal by the *first intention*, *Réunion per primam intentio'nem*, when cicatrization occurs without suppuration; union by the *second intention*, *Réunion per secundam intentio'nem*, being that which does not occur until the surfaces have suppurated. To obtain union by the first intention, the edges of a recent wound must be brought in apposition and kept together by means of adhesive plasters and a proper bandage. Delpech has substituted for those expressions, *Réunion primitive*, and *Réunion secondaire*.

INTERANEA, Entrails.

INTERARTICULAR, *Interarticula'ris*; from *inter*, 'between,' and *articulus*, 'a joint.' Parts situate between the articulations are so called; as *interarticular cartilages*, *interarticular ligaments*, &c.

INTERCADCENCE, *Intercaden'tia*, *Interciden'tia*, from *inter*, 'between,' and *cadere*, 'to fall.' Disorder of the arterial pulsations, so that, every now and then, a supernumerary pulsation occurs. The pulse, in such case, is said to be *intercurrent*.

INTERCALARY, *Intercala'ris*, *Inter'cidens*, from *intercalare*, 'to insert.' *Embol'imos*, *Interpol'atus*, *Provocato'rius*. The days which occur between those that are critical. The term has, also, been applied to the days of apyrexia in intermittent fevers.

INTERCELLULAR, *Intercellula'ris*, (F.) *Intercellulaire*; from *inter*, 'between,' and *cellula*, 'a small cell, a cellule.' That which is between cells or cellules.

INTERCELLULAR FLUID, Liqueur Sanguinis.

INTERCELLULAR PASSAGES. A term given by Mr. Rainey to irregular passages through the substance of the lung, which form the terminations of the bronchial tubes, are clustered with air-cells, and not lined by mucous membrane.

INTERCELLULAR SUBSTANCE, see Cytoblastema.

INTERCEPTIO, from *inter*, 'between,' and *capere*, 'to take.' A bandage, by the aid of which the ancients proposed to arrest the progress of the material cause of gout and rheumatism; and which consisted in covering the affected limbs with carded wool; surrounding them, afterwards, with broad bandages, applied from the fingers to the axilla, or from the toes to the groin.

INTERCEPTIO INTESTINORUM, Ileus.

INTERCERVICAUX, Interspinales colli.

INTERCIDENS, Intercalary.

INTERCIDENTIA, Intercadence.

INTERCILIUM, Mesophryon.

INTERCLAVICULAR, *Interclavica'ris*, from *inter*, 'between,' and *clavicula*, 'a clavicle.' That which is placed between the clavicles.

INTERCLAVICULAR LIGAMENT is a fibrous bundle, placed transversely above the extremity of the sternum, between the heads of the two clavicles. This ligament is flat. Its fibres, which are always longer above than below, are separated by small apertures, which are traversed by vessels. It prevents the separation of the two clavicles in the forced depression of the shoulder.

INTERCOSTAL, *Intercosta'lis*, from *inter*, 'between,' and *costa*, 'a rib.' *Mesopleu'rus*, *Mesopleu'rius*. That which is situate between the ribs. (F.) *Sous-costal*.

INTERCOSTAL ARTERIES vary in number. There is constantly, however, a *superior*, *Arteria*, *Intercosta'lis superior* seu *Intercosta'lis subclavia*, which is given off from the posterior part of the subclavian, and which sends branches into the first two or three intercostal spaces; and, generally, eight or nine *inferior* or *aortic intercostals*. These arise from the lateral and posterior parts

of the pectoral aorta, and ascend obliquely in front of the vertebral column, to gain the intercostal spaces, where they divide into a *dorsal* branch and an *intercostal*, properly so called.

INTERCOSTAL MUSCLES are distinguished into *internal*, — *inter pleurocostalis* of Dumas, — and *external*. The former are inserted into the inner lip, the latter into the outer lip of the edge of the ribs. The fibres of the *external intercostals* are directed obliquely downwards and forwards; and those of the *internal* downwards and backwards. Both are inspiratory or expiratory muscles, according as they take their origin on the upper or lower rib. Some small, fleshy fibres, seen occasionally at the inner surface of the thorax, descending obliquely from one rib to another, have been called *Infracostales*.

INTERCOSTAL NERVE, Trisplanchnic nerve.

INTERCOSTAL NERVES, *Branches souscostales* (Ch.), *Costal* or *Dorsal nerves*, proceed from the anterior branches of the dorsal nerves. They are twelve in number, and are distributed especially to the muscles of the parietes of the chest and abdomen.

INTERCOSTAL SPACE, *Intervallum seu Interstitium intercostale*, *Mesoplexurum*, *Mesoplexurium*, is the interval which separates one rib from that immediately above or below it.

INTERCOSTAL VEINS are distinguished like the arteries. The right superior intercostal vein is often wanting. When it exists, it opens into the back part of the subclavian. The same vein of the left side is very large. It communicates with the demi-azygos, receives the left bronchial vein, and opens into the corresponding subclavian. The right inferior intercostal veins open into the vena azygos; and those of the left into the demiazygos.

INTERCOSTO-HUMERAL NERVES. So called from their origin and distribution. They are the cutaneous branches of the second and third intercostal nerves.

INTERCURRENT, *Intercurrens*, from *inter*, 'between,' and *currere*, 'to run.' Diseases are so called which supervene at different seasons of the year, or which cannot be considered as belonging to any particular season. — Sydenham. A disease is likewise so termed which occurs in the course of another disease, as *Intercurrent Pneumonia*.

INTERCUTANEOUS, Subcutaneous.

INTERDENT'ITIUM, from *inter*, 'between,' and *dens*, *dentis*, 'a tooth.' The interval between teeth of the same order. — Linden.

INTERDIGITAIRE, Interdigital.

INTERDIG'ITAL, *Interdigita'lis*, (F.) *Interdigitaire*; from *inter*, 'between,' and *digitus*, 'a finger.' That which relates to the spaces between the fingers. *Interdigital space*, (F.) *Espace interdigitaire*, is used, also, for the commissure between the fingers.

INTERDIGIT'IUM. Same etymon as the last. A corn or wart, which grows between the fingers and toes, especially between the latter. — Pliny.

INTERÉPINEUX CERVICAUX, Interspinales colli.

INTERFEMINEUM, Perinæum, Vulva.

INTERFINIUM NARIUM, Septum narium.

INTERFEMIN'EUM, (*inter*, 'between,' and *femen*, *femina*, the inner part of the thigh,) Perinæum, Vulva.

INTERFORAMINEUM, Perinæum.

INTERGANGLION'IC, *Interganglion'icus*, from *inter*, 'between,' and *ganglion*, 'a knot.' An epithet for nervous cords, placed between ganglia, which they connect together.

INTERITUS, Death.

INTERLO'BAR, *Interloba'ris*, (F.) *Interlobaire*, from *inter*, 'between,' and *lobus*, 'a lobe.' That which is between lobes, — as of the lungs.

INTERLOB'ULAR, *Interlobula'ris*, from *inter*, 'between,' and *lobulus*, 'a small lobe.' That which is between lobules, — as of the lungs.

INTERLOBULAR FISSURES. The intervals between the lobules of the liver. The narrower spaces are called *Fissuræ interlobula'res*, the wider *spatia interlobula'res*. — Källiker. See Liver.

INTERLOBULAR PLEXUS OF THE BILIARY DUCTS. See Liver.

INTERLOBULAR SPACES. The angular interstices formed in the liver by the apposition of several lobules.

INTERLOB'ULAR TISSUE, *Tectum interlobula're*, *Ligamenta interlobula'ria pulmonum*. The cellular tissue between the pulmonary lobules.

INTERLOBULAR VEINS, see Liver.

INTERMAX'ILLARY, *Intermaxilla'ris*, from *inter*, 'between,' and *maxilla*, 'a jaw.' That which is situate between the maxillary bones.

INTERMAXILLARY BONE, *Inci'sive*, *Pal'atine*, or *Labial bone*, is a bony portion, wedged in between the two superior maxillary bones, which supports the upper incisors. This bone is found in the mammalia; and, also, in the human foetus.

INTERMÈDE, Excipient.

INTERMEDIA LIGAMENTALIA seu **NERVEA**, *Inscriptiones tendineæ musculorum*.

INTERMÉDIAIRE DE WRISBERG, Portio Wisbergii.

INTERMEDIATE VASCULAR SYSTEM, Capillary system.

INTERMENT, PREMATURE, Zoothapsis.

INTERMIS'SION, *Intermis'sio*, *Dialeip'sis*, *Dialeip'sis*, *Tempus intercala're*, *Interval'lum*, from *inter*, 'between,' and *mittere*, *missum*, 'to put or send.' (F.) *Intermittence*. The interval which occurs between two paroxysms of an intermittent or other disease — during which the patient is almost in his natural state. There is said to be *intermission* of the pulse, when, in a given number of pulsations, one or more may be wanting.

INTERMITTENCE, Intermision.

INTERMITTENT, *Intermittens*. Same etymon. That which has intermissions.

INTERMITTENT FEVER, *Febris intermittens*, *Dialeip'gra*, *Intermittens*, *An'etus*, *Pyretotypos'sis*, *Aque*, *Aque and Fever*, *Fever and Aque*, *Paludal* or *Periodic Fever*, (Prov.) *Agah*, Cold-chill, *Shaking Shakes*, (Old Eng.) *Aksis*, *Auggeres*, *Axes*, (F.) *Fièvre intermittente*, *F. d'Accès*, *F. des marais*, *F. Périodique*, is a fever consisting of paroxysms, with a complete state of apyrexia in the intervals. The chief types are the *Quotidian*, *Tertian*, and *Quartan*. The symptoms of intermittents are those of a decided and completely marked cold stage: (F.) *Stade de froid*, attended with paleness; collapse; impaired sensibility; and coldness, more or less diffused, followed by general rigors. After this occurs the *hot stage*; *Stade de la Chaleur*, the heat returning partially and irregularly, and at length becoming universal, and much above the standard of health. The pulse is now hard and strong; tongue white; urine high-coloured; thirst considerable. At length, the *sweating stage*, (F.) *Stade de Sueur*, makes its appearance: the moisture usually beginning on the forehead, face, and neck, and soon extending universally; the heat abating; thirst ceasing; the urine throwing down a sediment, and the functions being gradually restored to their wonted state. The tertian type is the most common, and the quartan the most severe. The quotidian more readily changes into a remittent and continued fever. The quartan has, generally, the longest cold stage, the tertian the longest hot. The chief exciting cause is marsh miasmata. Aque, also, occurs in districts where there are no marshy emanations. Such districts are, gene-

rally, of a volcanic nature; further we know not. When the disease has once attacked an individual, it is apt to recur. The *prognosis* is, in general, favourable, as far as regards life; but long protracted intermittents are apt to induce visceral obstructions, and engorgements, which may end in dropsy. In some countries, the disease is of a very pernicious character. The indications of treatment are, 1. To abate the violence of the paroxysm; and, 2. To prevent its return. The first indication demands the adoption of the general principles and remedies required in ordinary fever. The *second* is the most important. The period of apyrexia is that for action. The means for fulfilling this indication are:—the use of emetics, purgatives, cinchona, quinia, &c., arsenic, and forcible impressions made on the mind of the patient.

INTERMUSCULAR, *Intermuscularis*, from *inter*, 'between,' and *musculus*, 'a muscle.' That which is placed between muscles.

INTERMUSCULAR APONEUROSIS are aponeurotic laminae or septa, situate between muscles, to which they often give attachment.

INTERNAL, *Imer*, *Internus*. That which is placed on the inside. This epithet is given to parts that look towards an imaginary central plane, which divides the body into two equal and symmetrical portions, as well as to those which correspond with a cavity. Thus, we say—the *internal* or *inner surface* of the arm or thigh, skull, &c.

INTERNAL DISEASES, *Morbi interni*, are those which occupy the inner parts of the body. Their investigation belongs to the physician; *external* diseases falling under the management of the surgeon.

INTERNAT. The post or office of an *interne*.

INTERNE, see House-Surgeon.

INTERNODIA DIGITORUM MANUS, Phalanges of the fingers—i. *Digitum pedis*, Phalanges of the toes.

INTERNO'DIUM, from *inter*, 'between,' and *nodus*, 'a knot.' The part of the fingers between the joints. A phalanx.

INTERNUNCIAL, *Internuncia'lis*, from *inter*, 'between,' and *nuncius*, 'a messenger.' Relating or belonging to a messenger between parties. A term applied by Mr. Hunter to the functions of the nervous system.

INTERNUNTII DIES, Critical days.

INTERNUS AURIS, Tensor tympani. i. Mallo, Tensor tympani.

INTEROSSEOUS, *Interosseus*. That which is situate between the bones; from *inter*, 'between,' and *os*, 'a bone.'

INTEROSSEOUS ARTERIES OF THE FOREARM AND HAND. Of these there are several. The common *interosseous artery* arises from the posterior part of the ulnar, a little below the bicipital tuberosity of the radius; passes backwards and divides into two branches: the one called *anterior interosseous* descends vertically, anterior to the interosseous ligament; the other, called *posterior interosseous*, passes above the ligament, appears at its posterior part and divides into two great branches,—the *posterior recurrent radial* and the *posterior interosseous*, properly so called. In the hand,—1. The *dorsal metacarpal interosseous arteries* are given off by the *dorsalis carpi*, a division of the radial artery. 2. The *palmar interosseous arteries*, which arise from the convexity of the deep palmar arch, and give off the *middle interosseous arteries*. 3. The *dorsal interosseous of the index*, proceeding directly from the radial artery. In the foot, are distinguished:—1. The *dorsal interosseous arteries*, three in number, which arise from the artery of the metatarsus, a branch of the *dorsalis tarsi*. 2. The *plantar*

interosseous arteries, which are, also, three in number, and arise from the plantar arch.

INTEROSSEOUS LIGAMENTS. Ligaments seated between certain bones, which they unite; such are the ligaments between the radius and ulna, and between the tibia and fibula.

INTEROSSEOUS MUSCLES. These occupy the spaces between the bones of the metacarpus and metatarsus; and, consequently, belong,—some to the hand, others to the foot.

INTEROSSEOUS NERVE is a branch, given off by the median nerve, which descends before the interosseous ligament, accompanying the artery of the same name.

INTEROSSEOUS VEINS have the same arrangement as the arteries.

INTEROSSEUS MANUS, (F.) *Muscle interosseus de la main*.—*Métacarpo-phalangien-lateral sup-palmar* et *métacarpo-phalangien latéral*, (Ch.) These muscles are seven in number; two for each of the three middle fingers, and one for the little finger. Four are situate on the back of the hand, and three only in the palm. They are inserted into the metacarpal bones, and send a tendon to the tendon of the extensor communis. According to their office, they are, to each finger, an *adductor* and an *abductor*. The *index* has a dorsal abductor and a palmar one. The *middle finger* has two dorsal muscles for adductor and abductor; the *ring finger* has a dorsal adductor and a palmar abductor; and the *little finger* has only one interosseous abductor, which is palmar. These muscles produce abduction and adduction of the fingers, which they can also extend, owing to their connexion with the extensor tendons.

INTEROSSEUS PEDIS, *Métatarso-phalangien-lateral*, (Ch.) The number, arrangement, shape, and uses of these are the same as in the case of the preceding muscles. Four are *dorsal*, and three *plantar*: six belong to the three middle toes and one to the little toe. The great toe is devoid of them. As in the hand, they are distinguished, in each toe, into abductor and adductor.

INTEROSSEUS SECUNDUS, *Prior medii digiti*—i. *Quartus*, *Prior annularis*.

INTEROSSEUX DE LA MAIN, *Interosseus manus*.

INTERPARIETAL BONE, *Os Interparietale*. A bone found in the skulls of the children of the Peruvian races. It lies in the situation of the upper angle of the occipital bone, where the parietal bones separate from each other; and is the analogue of the interparietal bone of ruminants and carnivora.—Tschudi.

INTERPEDUNCULAR, *Interpeduncularis*; from *inter*, 'between,' and *pedunculi*, 'peduncles.' A term, applied to that which is placed between the peduncles of the encephalon,—as *Interpeduncular space*; a lozenge-shaped interval, behind the diverging peduncles or crura of the cerebrum, in which are found the posterior perforated space, corpora albicantia, tuber cinereum, and infundibulum, and pituitary body.

INTERPELLATUS, (*Morbus*), from *interpello*, (*inter*, and *pello*), 'I interrupt.' A term, by which Paracelsus designated those diseases whose progress is unequal, and paroxysms irregular.

INTERPLEVRO-COSTAUX, *Intercostal* muscles.

INTERPOLATUS, *Intercalary*.

INTERRUPTIO MENSTRUATIONIS, *Amenorrhœa*.

INTERSCAPULAR, *Interscapularis*, (*inter*, and *scapula*.) That which is between the shoulders; as the *interscapular region*. The *interscapular cavities*, *Cavitates interscapulares*, are the depressions between the scapulæ and the spinous processes of the vertebrae.

INTERSCAPULIUM. The spine of the scapula.

INTERSECTIO, Decussation, Intersection.

INTERSECTION, Intersection, Diac'ap'se, from *inter*, 'between,' and *seco, sectum*, 'to cut.' The point where two lines meet and cut each other. The name *aponeurotic intersection* is given to fibrous bands, which certain muscles present in their length, and by which they seem interrupted. Aponeurotic intersections are found in the recti muscles of the abdomen; in the semi-membranosus, complexus, sterno-thyroides, &c.

INTERSECTIONS TENDINEÆ MUSCULORUM. Inscriptions tendineæ musculorum.

INTERSEPIMENTUM THORACIS, Mediastinum.

INTERSEPTA HORIZONTALIA PACCHIONI. Tentorium.

INTERSEPTUM, from *inter*, 'between,' and *septum*, 'a partition.' *Cion, κίων.* The uvula; also, the septum narium. The diaphragm.

INTERSEPTUM NARIUM, Septum narium—i. Virginal, Hyman.

INTERSPINAL, Interspinalis, from *inter*, 'between,' and *spina*, 'the spine.' That which is seated between the spinous processes.

INTERSPINAL LIGAMENT, Membra'na interspinalis, (F.) *Ligament interépineux.* These occupy the intervals between the spinous processes in the back and loins. In the neck, they are replaced by the muscles of the same name. They prevent the too great flexion of the spine, and keep the spinous processes *in situ*. The outermost thickened portions of these are, sometimes, termed *Ligamenta d'apicium*.

INTERSPINALES COLLI, Spinal'es Colli mino'ra, Spinal'ti, (F.) *Intérépineux-cervicaux, Intercervicaux,* (Ch.) These are twelve in number, and occupy, in two parallel rows, the intervals between the spinous processes of the cervical vertebræ, from that of the atlas and vertebra dentata, to that between the last cervical and first dorsal vertebra. They are flat, thin, and quadrilateral. These muscles contribute to the extension of the neck and to throwing the head backwards.

INTERSPINALES DORSI ET LUMBORUM; portions of the *Transversaire-épineux* of Boyer, and the *sacro-spinal* of Chaussier. These muscles are of two kinds. The one (*Muscle grand épineux du dos*, of Winslow) representing fleshy bundles of different lengths, applied upon the lateral surfaces of the spinous processes, from the third dorsal vertebra to the second lumbar. The other (*Muscle petit épineux du dos*, Winslow) covered by the preceding. They are situate on each side of the interspinal ligament, in the form of small, short flat bundles, which pass from one spinous process to the second, third or fourth above it. These muscles aid in extending the vertebral column; and incline it a little to one side, when they act on one side only.

INTERSTICE, Interstit'ium, Aræo'ma, from *inter*, 'between,' and *sto, ἵστημι*, 'I stand.' *Dia'stéma, Intervallum.* Anatomists have given this name to the intervals between organs. The *iliac crest, crista ili*, for example, has two *lips* and an interstice between them, which affords attachment to the lesser oblique muscle of the abdomen.

INTERSTICE, Pore.

INTERSTIT'IAL, Interstitia'lis, same etymon. Applied to that which occurs in the interstices of an organ,—as 'interstitial absorption,' 'interstitial pregnancy,' &c.

INTERSTITIUM, Interstices—i. Ciliare, Ciliary ligament—i. Interco'stale, Interco'stal space—i. Jugulare, Throat.

INTERTINCTUS, Discreet.

INTERTRACHÉLIENS, Intertransversales colli.

INTERTRANSVERSAIRES DES LOMBES, Intertransversales lumborum.

INTERTRANSVERSALIS, Intertransversa'rius. That which is placed between the transverse processes of the vertebra.

INTERTRANSVERSALES COLLI, (F.) Intertransversaires cervicaux, Intertrachéliens, (Ch.) These are small, muscular bundles; quadrilateral; thin and flat; situate, in pairs, in the intervals between the transverse processes of the neck, except between the first and second, where there is only one. They are distinguished into *anterior* and *posterior*. The former are six in number; the latter five. These muscles bring the transverse processes of the neck nearer each other, and contribute to the lateral flexion of the neck.

INTERTRANSVERSALES LUMBORUM, Musculi intertransversi lumborum, (F.) Intertransversaires des lombes. These are almost entirely fleshy, and ten in number; five on each side. They are similar to the preceding in general arrangement, except that they are more marked, and not placed in two rows. Each intertransverse space contains only one. They are quadrilateral, and flat. The first occupies the space between the transverse processes of the first lumbar and the last dorsal vertebræ; and the last is between that of the fourth and fifth lumbar vertebræ. These muscles incline the lumbar region laterally; and straighten it when inclined to one side.

INTERTRIGO, Chafing—i. Podicis, Chafing.

INTERVAL, LUCID, see Lucid.

INTERVALLUM, Intermission, Interstice—i. Interco'stale, Interco'stal space.

INTERVENTUM, Mesophlebium.

INTERVERTEBRAL, Intervertebra'lis, (inter, and vertebra.) That which is situate between the vertebræ.

INTERVERTEBRAL CARTILAGES, Fibro-cartilages or discs, Ligamenta intervertebra'lia, Fibro-cartilagin'ines intervertebra'les. These organs are of a fibro-cartilaginous nature; sections of a cylinder; flexible; whitish; resisting; and situate between the bodies of the vertebræ,—from the space between the second and third as far as that between the last vertebra and sacrum. Their form is accommodated to that of the vertebra with which they are in connexion; so that, in the neck and loins, they are oval, whilst in the dorsal region they are nearly circular.

INTERVERTEBRAL DISCS, Intervertebral cartilages—i. Fibro-cartilages, Intervertebral cartilages.

INTESTABILIS, Castratus.

INTESTATUS, Castratus.

INTESTIN, Intestine.

INTESTINAL, Intestina'lis, Enter'icus, Enter'ic, (F.) Entérique, from *intus*, 'within.' That which belongs to the intestines,—as *intestinal canal*, &c. Borden uses the term *Pouls intestinal* for a pulse, which he conceived to announce an approaching crisis by the intestines.

INTESTINAL JUICE, Succus entericus.

INTESTINAL TUBE OR TRACT, Cana'lis seu Tractus intestino'rum. The canal formed by the intestines from the pyloric orifice of the stomach to the anus.

INTESTINE, Intesti'num, En'teron, Chorda, Gut, Nedyia (pl.), *Pan'tices* (pl.), *Boel'li* (pl.), *Bowel, (F.) Intestin, Boyan.* Same etymon. A musculo-membranous canal, variously convoluted, which extends from the stomach to the anus, and is situate in the abdominal cavity; the greater part of which it fills. In man, its length is six or eight times that of the body. It is divided into two principal portions, called *small intestine* and *large intestine*. The former, *Intesti'num*

*ten'ui, I. grac'ile, (F.) Intestin grêle, constituting nearly four-fifths of the whole length, begins at the stomach and terminates in the right iliac region. It is divided into duodenum, jejunum and ileum. Some anatomists give the name small intestine to the last two only; which are kept in place by the mesentery, and form a large paquet, occupying the umbilical and hypogastric regions, a part of the flanks, of the iliac regions, and of the cavity of the pelvis. It is composed of, 1. A serous membrane, which is peritoneal. 2. Of a muscular coat, whose fibres are very pale, and are placed, in part, longitudinally; but the greater part transversely. 3. Of a whitish, mucous membrane; villous, and forming folds or valves—*valvulae conniventes*—at its inner surface, and furnished with a number of mucous follicles, called *glands of Lieberkühn, (Crypts or Follicles of Lieberkühn),* and in the duodenum with the *glands of Brunner*. The arteries of the small intestine proceed from the superior mesenteric; its veins open into the vena porta. Its nerves proceed from the superior mesenteric plexus. The large intestine, *Intestinum crassum, Megalocol'ia, (F.) Gros intestin,* forms a sequence to the small. It is much shorter, and is solidly attached in the regions of the abdomen which it occupies. It begins in the right iliac region; ascends along the right flank, till beneath the liver, when it crosses the upper part of the abdomen, descends into the left iliac fossa, and plunges into the pelvic cavity, to gain the anus. The great intestine is usually divided into three portions,—the *cæcum, colon and rectum*. It receives its arteries from the superior and inferior mesenteries. Its veins open into the vena porta. Its nerves are furnished by the mesenteric plexuses. Its lymphatic vessels, which are much less numerous than those of the small intestine, pass into the ganglions or glands seated between the different reflections of the peritoneum, which fix it to the abdominal parietes. The use of the intestines is,—in the upper part, to effect the chyliification of the food and the absorption of the chyle;—in the lower, to serve as a reservoir, where the excrementitious portion of the food collects; and, also, as an excretory duct, which effects its expulsion.*

* **INTESTINORUM LÆVITAS, Lientery.**

INTESTINULA CEREBRI, Convolutions (cerebral)—i. Meibomii, Meibomius, glands of

INTESTINULUM, Funiculus umbilicalis.

INTESTINUM CELLULATUM, Colon—i. Circumvolutum, Ileon—i. Crassum, Colon, see Intestine—i. Gracile, see Intestine—i. Grande, Colon—i. Laxum, Colon—i. Majus, Colon—i. Medium, Mesentery—i. Plenum, Colon—i. Rectum, Rectum—i. Tenue, see Intestine.

INTIMUM UNGUIS, Nail, root of the.

INTONATIO INTESTINALIS, Borborygmi.

INTORSIO, Distortion.

INTOXICATIO, Poisoning—i. Arsenicalis, Arsenicisms—i. Opiaca, Meconism—i. Saturnina, Saturnisms.

INTOXICATION, Poisoning, Temulentia—i. Antimonial, Stibialism—i. des Marais, see Miasm.

INTOXICATION SATURNINE. Lead poisoning; saturnine cachexy. The aggregate of symptoms which present themselves prior to an attack of lead colic.—*Tanquerel des Planches.*

INTRALINGUAL SALIVARY GLANDS, see Salivary glands.

INTRALOBULAR, Intralobula'ris; from intra, 'within,' and lobulus, 'a lobule.' Relating to the space within a lobule:—as the "intralobular veins," *vena seu ven'ula intralobula'ris seu central'es, of the liver:—veins which communi-*

cate with the interlobular veins, and are the radicles of the hepatic veins.

INTRA-PELVIO TROCHANTÉRIEN, Ob-turator internus.

INTRA-UTERINE, Intrauteri'nus; from intra, 'within,' and uterus, 'the womb' That which takes place within the womb—as 'intra-uterine life.'

INTRICATURA, Decussation.

INTRIN'SIC, Intrin'secus; from intra, 'within,' and secus, 'towards.' Applied to the internal muscles of certain organs; as those of the ear, tongue, and larynx. Linnaeus gave the name *Intrin'seci* to internal diseases.

INTROITUS PELVIS, see Pelvis—i. Vaginae, Rima vulvæ.

INTROMIS'SION, Intromis'sio; from intro, 'within,' and mitto, missum, 'I send.' The act of introducing one body into another, as *Intromissio Penis, (F.) Intromission de la Vierge.*

INTUSSUSCEPTIO, Intussusceptio—i. Entropium, Entropion.

INTUBUM, Cichorium endivia—i. Erraticum, Cichorium intybus.

INTUMESCENTIA, Intumescen'tia, Onco'sis, Dionco'sis, from intumescere, (in, and tumescere, 'to swell.') Augmentation of size in a part or in the whole of the body. Sauvages uses the word *Intumescen'tia* for an order of diseases, in which he comprehends polysarcia, pneumotosis, anasarca, oedema, phlyctonia, and pregnancy.

INTUMESCENTIA GANGLIFORMIS, Ganglion of Gasser, see Facial nerve—i. Lactea Mammarum, Sparganosis—i. Lienis, Splenoncus—i. Semilunaris, Ganglion of Gasser.

INTUS INVERSUS, Transposition of the viscera.

INTUSSUSCEPTIO, from intus, 'within,' and suscipio, susceptum, 'I receive.' Introsuscept'io. In physiology:—the mode of increase peculiar to organized bodies. In pathology, like *Convolvulus, Volvulus* seu *Suscept'io* seu *Parasago'gê* intestino'rum, *Chordap'us, Enteroparisa'gogê, Indigita'tio, Tormen'tum, and Incagina'tio,* it means the introduction of one part of the intestinal canal—*intussuscept'um*—into another, which serves it as a sort of vagina or sheath. Generally, it is the upper part of the small intestine, which is received into the lower—*intussuscept'icus*—when the intussusception is said to be *progres'sive*. At times, however, it is *retrograde*. As the disease cannot be positively detected by the symptoms, it must be treated upon general principles. At times, the invaginated portion has separated and been voided per anum,—the patient recovering. The disease is, however, of a very dangerous character.

INTUSSUSCEPTUM, see Intussusceptio.

INTUSSUSCIPiens, see Intussusceptio.

INTYBUM, Cichorium endivia.

INTYBUS HORTENSIS, Cichorium endivia.

INULA, see Inula helenium—i. Britannica, I. dysenterica—i. Common, I. helenium—i. Conyzæa, I. dysenterica.

IN'ULA DYSENTERICA. The systematic name of the *Lesser Inula, I. Britan'nica* seu *Conyzæa, Aster dysenter'icus* seu *undulatus, Pulcra'ria dysenter'ica, Cony'za, C. media, Ar'nica Sueden'sis* seu *spu'ria, Cun'ila mas'cula* seu *me'dia, Cun'ila'go, (Prov.) Mare's fat, (F.) Aunée antidysentérique, Herbe de Saint Roch. Nat. Ord. Composites. This plant was once considered to possess great antidysenteric virtues. The whole plant is acrid and somewhat aromatic.*

INULA HELEN'UM. The systematic name of the *El'campane, En'ula, E. camp'na, Elen'ium, Necta'rium, Helenium, Aster helen'ium* seu *offici'nalis, Corvisea'tia helen'ium, Common Inula, (F.) Aunée ou Aunée officinale. The root In'ula*

(Ph. U. S.) was formerly in high esteem in dyspepsia, cachexia, pulmonary affections, &c. It is now scarcely used.

INULA, Lesser, *Inula dysenterica*—i. Squarrosa, *Conyza squarrosa*.

INUNCTIO, Inunction, Liniment.

INUNCTIO, *Inunctio*, *Pernunctio*, *Militio*, *Militus*, *Militio*, *Oblinitio*, *Chirisis*, *Catachirisis*, *Diachirisis*, *Epichirisis*, *Unctio*, from *ungere*, *inunctum*, (*in*, and *ungere*, *unctum*), 'to anoint.' The act of rubbing in an ointment or a liniment, *unction*, (F.) *Unction*, *Inunction*. Inunction of the whole body with a piece of bacon, lard, or with oil, has been recommended in scarlatina, measles, and in scrofulous or tuberculous cachexia.

INUNDATIO, Depuration.

INUSTIO, Cauterization, Cauterium.

INUSTORIUM, Cauterium.

INVAGINATED, *Invagina'tus*, from *in*, and *vagina*, 'a sheath.' Applied to a part which is received into another, as into a sheath.

The *invaginated* or *slit* and *tail bandage* is one in which strips or tails pass through appropriate slits or button-holes.

INVAGINATIO, Intussusceptio.

INVAGINATION, *Invagina'tio*. Intussusceptio. Also, an operation, which consists in introducing one end of a divided intestine into the other, with the view of restoring the continuity of the intestinal canal.

The name *Invagination* is, likewise, given to an operation for the radical cure of crural hernia, which consists in introducing into the crural canal a thick tent of charpie, the effect of which is, by compression and adhesive inflammation, to obliterate the neck of the hernial sac.—Nysten.

INVALETUDO, Infirmary.

INVALID, Valetudinary.

INVALIDING, (F.) *Réforme*. The removal from the army of a soldier, who has become, from infirmity, unfit for service.

INVALIDUS, Valetudinary.

INVASIO, Arche, Attack, Paroxysm.

INVERMINATION, Helminthiasis.

INVERSIO PALPEBRARUM, Entropion—i. Uteri, Hysteroptosis, Uterus, inversion of the.

INVERSION DE LA MATRICE, Uterus, inversion of the—i. *du Lambeau*, see Flap.

INVERTENTA, Absorbents, Antacids.

INVESTITURA FUNICULI UMBILICALIS, see Funiculus umbilicalis.

INVETERATUS, Chronic.

INVISCONTIA, Incrassantia.

INVISCONTIO OCULI, from *in*, and *viscum*, 'bird lime.' A morbid adhesion of the eyelids to each other, or to the globe of the eye. Also, gluing together of the eyelids by a viscid secretion.

INVOLUCRUM, *Involumen'tum*, from *involvere*, (*in*, and *volvere*, *volutum*), 'to fold in.' A covering: hence *Involucra cer'e'ri*, the membranes of the brain. *Involucra nervo'rum*, the sheaths of the nerves.

INVOLUCRUM CORDIS, Pericardium—i. *Corporis commune*, Integument—i. *Linguae*, see Tongue—i. *Membranaceum*, *Decidua reflexa*—i. *Nervorum*, *Neurilemma*—i. *Reti comparatum*, *Retina*.

INVOLUMENTUM, Involverum.

INVOLVENTIA, Demulcents.

IODE, Iodine.

IODES, Æruginous.

IODETUM HYDRARGYRICUM, Hydrargyri iodidum rubrum.

IODHYDRARGYRITE DE CHLORURE MERCUREUX, Mercury, iodide of chloride of.

IOD'IC, *Iod'icus*, (F.) *Iodique*. Same etymon as Iodine. Containing iodine.

IODIDUM HYDRARGYRI CHLORIDI,

Mercury, iodide of chloride of—i. Hydrargyricum, Hydrargyri iodidum rubrum—i. Hydrargyrosium, Hydrargyri iodidum.

IODINE, *Iod'ina*, *Iod'ium* (Ph. U. S., 1842), *Iod'in'ium* (Ph. U. S., 1851), *Iod'in'eum*, *Iod'um*, *Iod'ium*, *Iod'ium*, *Iod'um*, (F.) *Iode*; from *ιώδης*, *viol'aceus*, 'of a violet colour,' so called from the violet vapour it exhales when volatilized. It is contained in the mother waters of certain fuci, and is obtained by pouring an excess of concentrated sulphuric acid on the water obtained by burning different fuci, lixiviating the ashes and concentrating the liquor. The mixture is placed in a retort to which a receiver is attached, and is boiled. The iodine passes over and is condensed. It is solid, in the form of plates; of a bluish gray colour, of a metallic brightness, and smell similar to that of the chloride of sulphur. Its s. g. is 4.946. When heated, it becomes volatilized, and affords the vapour which characterizes it. With oxygen it forms *Iodic* acid, and with hydrogen *Hydriodic acid*. The tincture of iodine and the iodides have been employed with great success in the treatment of goitre and of some scrofulous affections. It must be administered in a very small dose and for a long period. It is said to be apt, however, to induce cholera morbus, signs of great nervous irritability, *Iodome'thē*, (F.) *Ivresse iodique*, and emaciation of the mammae. When these symptoms, collectively termed *Iodism*, *Iodosis*, and *Iod'in'ia*, (F.) *Cachexie iodée*, are urgent, the dose may be diminished, or it may be wholly discontinued, and afterwards resumed.

Various preparations of iodine are employed in medicine.

IODINED, Iodized.

I'ODIZED, *I'odined*; *Iodisa'tus*. Impregnated, or affected with, iodine, or its preparations.

IODINIA, see Iodine.

IODINIUM, Iodine.

IODINUM, Iodine.

IODIQUE, Iodic.

IODISM, see Iodine.

IODIUM, Iodine.

I'ODOFORM, *Iodofo'r'mum*, (F.) *Iodoforme*, *Iodure de formyle*; so called from its analogy to chloroform. It is obtained by the reaction on each other of iodine, bicarbonate of potassa, water, and alcohol. It is in crystalline plates, of a beautiful citrine colour; and may be given in the same cases as the other preparations of iodine. Dose, one grain, three times a day. It is the *sesqui-iodide of carbon*. See Carbonis sesqui-iodidum.

IODOGNOSIE, Iodognosis.

IODOGNO'SIS, (F.) *Iodognosie*; from *iodine*, and *γνῶσις*, 'knowledge.' A knowledge of iodine in its various relations.—Dorvault.

IODOHYDRARGYRATE OF POTASSIUM, see Iodine.

IODOMETHE, see Iodine.

IODOPE'GÆ, from *iodum*, 'iodine,' and *πῆγη*, 'a spring.' Mineral waters containing iodine.

IODOPHTHISIE, from *iodine*, and *φθίσις*, 'phthisis.' Loss of flesh and strength produced by iodine.

IODOSIS, see Iodine.

IODUM, Iodine.

IODURE DE FORMYLE, Iodoform—i. *de Mercure*, Hydrargyri iodidum—i. *de Plomb*, Plumbi iodidum—i. *Plombique*, Plumbi iodidum—i. *de Potassium*, Potassæ hydriodas—i. *de Quinine*, Quinæ hydriodas—i. *de Soufre*, Sulphuris iodidum—i. *de Zinc*, Zinci iodidum.

IODURETUM AMMONIÆ, see Iodine—i. Amyli, see Iodine—i. Hydrargyrosium, Hydrargyri iodidum—i. Sulphuris, see Iodine.

IODYMUS, Dicephalus.

ION, Viola.

IONIA, *Tencrium chamæpitya*.

IONIDIUM MARCUCCI, *Cinchunchulli*—a South American plant. *Sex. Syst.* Pentandria Monogynia; *Nat. Ord.* Violariæ; said to be extremely efficacious in the Mal de San Lazaro, of Colombia.

IONIUM, Iodine.

IONTA, see Excretion.

ION'THUS, *Varnu*, *Violet Eruption*, from *ion*, 'the violet,' and *avdos*, 'a flower,' or *ovdos*, 'foulness' [?]. An unsuppurative, tubercular tumour; stationary; chiefly on the face. A genus in the class *Hæmatica*, order *Phlogotica* of Good.

IONTHUS CORYMBIFER, *Gutta rosea*—i. Varus, Aene.

IONUM, Iodine.

IOTACISMUS, from the Greek letter *Iota*. Defective articulation,—the patient not being able to pronounce the palatals *j* and *g* soft.

IPECAC, Apocynum androsæmifolium, Euphorbia corollata, Gillenia trifoliata, Ipecacuanha, Triosteum—i. American, Apocynum androsæmifolium—i. Wild, Euphorbia ipecacuanha.

IPECACUAN'HA. In common parlance, often abridged to *Ipecac*, and in Ireland to *Hippo*. The pharmacopoeial name of the root of *Cephaelis* seu *Calliocoeca ipecacuan'ha*, *Ipecacuan'ha*, *Cogosanga*, *Hypocuan'na*, *Hippocuan'na*, (F.) *Ipecacuanha* *anellé*. *Nat. Ord.* Cinchonaceæ. It is also obtained from the *Psychotria emetica* of Peru. The odour of the root, *Ipecacuan'ha* root, *Ipecacuan'ha* radior. *Radic. Brazilien'sis*,—*Ipecacuanha* (Ph. U. S.), (F.) *Racine Brésilienne* ou *de Brésil*,—is faint and peculiar; taste bitter, subacid, and mucilaginous; both water and alcohol extract its virtues, which depend on a peculiar principle, called *Emetia*. It is emetic in large doses; sudorific in smaller. *Dose*, as an emetic, gr. xx to xxx,—alone, or united with gr. i. to ij of tartarized antimony.

IPECACUANHA, Euphorbia corollata, Pedilanthus tithymaloides—i. *Anellé*, *Ipecacuanha*—i. Bastard, *Asclepias curassavica*, Triosteum perfoliatum—i. *Blanc de l'île de France*, *Tylophora asthmatica*—i. *Blanc de St. Domingue*, *Asclepias curassavica*—i. of the Isle of France, *Tylophora asthmatica*.

IPECACUAN'HA, WHITE, is obtained from different species of *Richardsonia* and *Ionidium*. *Viola ipecacuan'ha* seu *Solea ipecacuan'ha* seu *Pomblea ipecacuan'ha* also affords it. It is weaker than the gray.

IPO, Upas.

IPOMÆA BATATOÏDES, see *Convolvulus jalapa*—i. *Jalapa*, *Convolvulus jalapa*—i. *Macrorrhiza*, *Convolvulus jalapa*—i. *Mestitlanica*, see *Convolvulus jalapa*—i. *Nil*, *Convolvulus nil*—i. *Operculata*, see *Batata* de purga—i. *Orizabensis*, see *Convolvulus jalapa*—i. *Purga*, see *Convolvulus jalapa*—i. *Schiedeana*, see *Convolvulus jalapa*.

IRA, Rage.

IRACUNDUS, *Rectus externus oculi*.

IRAL'GIA; *Ireal'gia*, *Iridal'gia*, from *iptis*, 'the iris,' and *alges*, 'pain.' Pain in the iris.

IRASCIBLE, *Rectus externus oculi*.

IREALGIA, *Iralgia*.

IREONCION, *Iriancistron*.

I'RIAN, *Iria'nus*, (F.) *Irien*. Belonging to the iris.

IRIANCIS'TRON, *Iridancistron*, *Ireon'cion*, from *iptis*, and *αγκιστρον*, 'a hook.' A hooked instrument, used by Schlagintweit in the formation of an artificial pupil.

IRIDÆ'A EDU'LIS. One of the Algae, eaten in Scotland and the Southwest of England.

IRIDALGIA, *Iralgia*.

IRIDANCISTRON, *Iriancistron*.

IRIDAUXE'SIS, from *Iris*, *iridis* and *αὐξήσις*, 'augmentation;' *Staphylo'ma Iridis*, *S. Uveæ*,

Iridonco'sis, *Lymph'neus Iridis*. Exudation of fibrin into the tissue of the iris.

IRIDECTOMEDIALYSIS, see *Coretomia*.

IRIDECTOMIA, see *Coretomia*.

IRIDENCELEISIS, see *Coretomia*.

IRIDENCEISMUS, see *Coretomia*.

IRIDERE'MIA, from *Iris*, *iridis*, and *εἰρηνο*, 'deprived of;' *Iridostere'sis*. Absence of iris, either apparent or real.

IRIDITIS, *Iritis*.

IRIDOCELE, *Ptoxis iridis*.

IRIDOCOLOBOMA, *Coloboma iridis*.

IRIDODIALYSIS, *Coretomia*.

IRIDODONE'SIS, from *Iris*, and *δονεω*, 'I shake.' Tremulousness of the Iris. See *Hippus*.

IRIDOMALACIA, from *Iris*, *iridis*, and *μαλακία*, 'softness.' Mollescence or softening of the iris.

IRIDOMELANO'MA, *Iridomelano'sis*, from *Iris*, and *μελανν*, 'to colour black.' The deposition of black matter on the tissue of the iris.

IRIDONCO'SIS, *Iridon'cus*, *Ironco'sis*, *Hyperronco'sis Iridis*, from *Iris*, and *ογκος*, 'a swelling.' Tumefaction or thickening of the iris. Also *Iridauxesis*.

IRIDONCUS, *Iridoncosis*.

IRIDOPERIPHACITIS, *Iridoperiphakitis*, from *Iris*, *περι*, 'around,' and *phacitis*, inflammation of the lens. Inflammation of the anterior hemisphere of the capsule of the lens.

IRIDOPHACIA, *Hippus*.

IRIDOPTOSIS, *Ptoxis iridis*.

IRIDORRHAGAS, *Iridorrhog'e*, *Iridor'rhox*, *Iridoschia'ma*, *Fissura Iridis*, from *Iris*, and *ρῆγας*, 'a fissure.' Fissure of the iris. See *Coloboma*.

IRIDORRHAGE, *Iridorrhagas*.

IRIDORRHOX, *Iridorrhagas*.

IRIDOSCHISMA, *Coloboma Iridis*, *Iridor'rhagas*.

IRIDOSTERESIS, *Irideremia*.

IRIDOTENCELEISIS, *Coretomia*.

IRIDOTOMEDIALYSIS, see *Coretomia*.

IRIDOTOMTA, see *Coretomia*.

IRIDOTROMUS, *Hippus*.

I'RIEN, *Irian*.

IRIS, (*gen. Iridis*) *Grammè*. So called from its resembling the rainbow in a variety of colours. A membrane, stretched vertically at the anterior part of the eye, in the midst of the aqueous humour, in which it forms a kind of circular, flat partition, separating the anterior from the posterior chamber. It is perforated by a circular opening called the *pupil*, which is constantly varying its dimensions, owing to the varying contractions of the fibres of the iris. Its posterior surface has been called *uvea*, from the thick, black varnish which covers it. The pigmentary stratum, on its free surface, appears to be bounded by a delicate, but sharply defined, line, which has been described as a special membrane—*Membrana pigmenti seu lim'itane seu lim'itane Poci'ni* seu *Jaco'bi Arnold*. The greater circumference of the iris is adherent to the ciliary processes and circle. It has an external plane of radiated fibres and an internal one of circular fibres, which serve—the one to dilate, the other to contract the aperture of the pupil. The iris receives the *irian* nerves. Its arteries are furnished by the long ciliary arteries, which form two circles by their anastomoses; the one very broad, near the great circumference; the other, smaller, and seated around the circumference of the pupil. The veins of the iris empty themselves into the *Vasa vorticosa*, and into the long ciliary veins. The use of the iris seems to be,—to regulate by its dilatation or contraction, the quantity of luminous rays necessary for distinct vision. The different colours of the iris occasion the variety in the colours of the human eye.

IRIS, see Herpes Iris — i. Common, I. Germanica — i. Common, I. Germanica — i. de Florence, I. Florentina.

IRIS FLORENTINA, *Florentine iris* or *orris*, (F.) *Iris de Florence*. Order, Iridaceæ. The rhizoma of this plant is extremely acrid in its recent state; and, when chewed, excites a pungent heat in the mouth, that continues for several hours. When dried, the acrimony is lost, or nearly so, and the smell is very agreeable. It is indebted to its agreeable flavour for its retention in the pharmacopœias, although it is ranked as an expectorant. The *orris* root of the shops is said to consist of the rhizomes of three species—*I. Florentina*, *I. paludosa*, and *I. Germanica*.

IRIS FÆTIDIS'SIMA, *I. fœtida*, *Spath'ula fœtida*, *Xyris*, *Stinking iris*, (F.) *Iris puant*, *Glayeul puant*. The root has been esteemed antispasmodic and narcotic.

IRIS GERMANICA. The systematic name of the *Flower-de-Luce*, *Iris nostras* seu *vulgaris*, *Aier'-en*, Common *iris* or *orris*, *Gladi'olis cœruleus*, (F.) *Iris commun*, *Flambe*. The fresh roots have a disagreeable smell, and an acrid, nauseous taste. They are powerfully cathartic, and are given in dropsies, where such remedies are indicated.

IRIS HEXAGONA, I. versicolor.

IRIS LACUS'TRIS, *Dwarf-flag*, *Dwarf lake-iris*; has the properties of I. versicolor.

IRIS, LAKE, DWARF, I. lacustris — i. Lutea, I. pseudacorus — i. des Marais, I. pseudacorus — i. Nostras, I. Germanica — i. Palustris, I. pseudacorus.

IRIS PSEUDACORUS. The systematic name of the *Yellow water-flag*, *Iris palustris* seu *lutæa*, *Gladi'olis luteus*, *Ac'orus vulgaris* seu *adulterinus* seu *palustris*, *Pseudacorus*, *Buttoman*, (F.) *Iris des marais*, *Faux acore*, *Flambe bâtar*, *Acore bâtar*. The root has an acrid, styptic taste. It is an errhine, sialogogue, and acrid astringent. The expressed juice is diuretic, and said to be a useful application to serpiginous and other cutaneous affections. Rubbed on the gums, or chewed, it is said to cure toothache.

IRIS PUANT, I. fetidissima — i. Stinking, I. fetidissima.

IRIS VERSICOLOR, *I. hexago'na*, *Blue flag*. The rhizoma of this is an active cathartic, and has been much used, as such, by the American Indians. It is reputed to be diuretic.

IRIS VIRGINICA, *Slender Blue flag*; and *Iris verna*, are said to have virtues like those of I. versicolor.

IRIS VULGARIS, I. Germanica.

IRITIS, Iritis.

IRITIS, *Inflammation of the iris*, *Inflamma'tio Iridis*, *Irisi'tis*, *Iridi'tis*, from *Iris*, and *itis*, denoting inflammation, *Uvei'tis* of some. The chief symptoms are;—change in the colour of the iris; fibres less movable; tooth-like processes shooting into the pupil; pupil irregularly contracted, with the ordinary signs of inflammation of the eye. If the inflammation does not yield, supuration takes place; and although the matter may be absorbed, the iris remains immovable. It is often caused by syphilis. The general principles of treatment are, to deplete largely and exhibit mercury freely: along with attention to other means advisable in ophthalmia. The free use of quinia is sometimes serviceable. When the inflammation is seated in the serous covering of the iris, it is termed *Iritis serosa*.

IRON, Ferrum — i. Acetate of, Ferri acetat — i. Alumina, sulphate of, Ferri et aluminis sulphas — i. Ammoniated, Ferrum ammoniatum — i. Ammonio-citrate of, Ferri ammonio-citras — i. Ammonio-chloride of, Ferrum ammoniatum — i. Ammonio-tartrate of, Ferri ammonio-tartas — i. and Ammonia, citrate of, Ferri ammonio-citras — i. and Quinia, Hydriodate of, see Iodine.

IRON BARK, see Kino.

IRON, Bromide of, see Bromine — i. Carbonate of, saccharine, Ferri proto-carbonas — i. Carburet of, Graphites — i. Black oxide of, Oxydum ferri nigrum — i. Chloride of, Ferri chloridum — i. Citrate of, Ferri citras — i. Citrate of the magnetic oxide of, see Ferri ammonio-citras — i. Ferrocyanuret of, Prussian blue — i. Ferropussiate of, Prussian blue — i. Filings, purified, Ferri limatura purificata — i. Hydrated oxide of, Ferrum oxydatum hydratum — i. Hydrated peroxide of, Ferrum oxydatum hydratum — i. Hydrated tritoxide of, Ferri iodidum — i. Hydrobromate of, see Bromine — i. Hydro-oxide of, Ferrum oxydatum hydratum — i. Iodide of, see Iodine, and Ferri iodidum — i. Ioduret of, Ferri iodidum — i. Lactate of, Ferri lactas — i. Liquor, Ferri acetat — i. and Magnesia, citrate of, Ferri et Magnesiæ citras — i. Magnetic Oxide of, Oxydum ferri nigrum — i. Muriate of, Ferri chloridum — i. Nitrate of, solution of, see Tinctura Ferri muriatis — i. Perchloride of, Ferri chloridum — i. Peroxide of, Ferri subcarbonas — i. Phosphate of, Ferri phosphas — i. Pills of, compound, Pilule ferri composita — i. Potassio-citrate of, see Ferri ammonio-citras — i. Protocarbonate of, Ferri protocarbonas — i. Protoxide of, lactate of Ferri lactas.

IRON PYRITES, Ferri sulphuretum.

IRON, QUEVENNE'S, Ferri pulvis.

IRON AND QUINIA, citrate of, Ferri et Quiniæ citras — i. and Quinia, hydriodate of, see Quinia, iodide of iodhydrate of.

IRON REDUCED, Ferri Pulvis.

IRON, SESQUICHLORIDE OF, Ferri chloridum — i. Sesquinitrate of, Solution of, see Tinctura ferri muriatis — i. Sesquioxide of, Ferri subcarbonas — i. Sodio-citrate of, see Ferri ammonio-citras — i. Sulphate of, Ferri sulphas — i. Sulphuret of, Ferri sulphuretum — i. Tannate of, Ferri tannas — i. Tartarized, Ferrum tartarizatum — i. Ternitrate of, solution of, see Tinctura ferri muriatis — i. Tritohydro-ferrocyanate of, Prussian blue.

IRON AND POTASS, Tartrate of, Ferrum tartarizatum.

IRON, VALERIANATE OF, Ferri valerianas.

IRON WEE, Vernonia noveboracensis.

IRONCOSIS, Iridoncosis.

IRRADIATION, *Irradiatio* (ir and radius). Shooting or proceeding from a centre — as an 'irradiating pain.'

IRREDUCIBLE, (F.) *Irréductible*. An epithet given to fractures, luxations, herniæ, &c., when they cannot be reduced.

IRREDUCTIBLE, Irreducible.

IRREGULAR, *Irregula'ris*, *Anom'alous*; from ir, negation, and regula, 'a rule.' A term chiefly applied to the types of a disease; and also, to the pulse when its beats are separated by unequal intervals.

IRREGULAR PRACTITIONER. One who does not practise his profession according to rules sanctioned by law or custom.

IREMEDIABLE, Incurable.

IREPTIO, Attack.

IRRIGATION, *Irrigatio*, *Embroc'atio*, *Irrora'tio*: 'the act of watering or moistening,' from ir, and rigo, 'I water.' The methodical application of water to an affected part, to keep it constantly wet.

IRRITABILITAS, Irritability — i. Halleriana, Irritability — i. Morbosa, see Subsultus tendinum — i. Vesice, Cysterethismus.

IRRITABILITY, *Irritabil'itas*, from irrita, (in, 'negation,' and ritus, 'the usual manner,') 'I provoke.' *Via irritabilit'is*, *V. Vita*, *V. insita* of Haller, *V. vitalis* of Gorter, *Oscillatio* of Boerhaave, *Tonic power* of Stahl, *Vita propria*, *Irritabilitas Halleriana*, *Inherent power*, *Con-*

tractility, Excitability, &c. (F.) *Irritabilité, I. Hallérienne*. A power, possessed by all living, organized bodies of being acted upon by certain stimuli, and of moving responsive to stimulation. It is the ultimate vital property.

IR'RITABLE, *Irritabilis*. That which is endowed with irritability. Every living organized tissue is irritable; that is, capable of feeling an appropriate stimulus, and of moving responsive to such stimulus. Irritable is often used in the same sense as *impressible*, as when we speak of an irritable person, or habit, or temper. This last condition has been variously termed—*Debilitas nervosa*, *D. Erethis'ica*, *Hypercinesia nervosa*, *Neurosthenia*, *Neurosthenia*, *Sensibilitas anomala*, *S. Morbosa*, *S. aucta*.

IRRITAMEN, Irritant, Stimulus.

IRRITAMENTUM, Erethism, Irritant, Stimulus—i. Metallicum, Galvanism.

IR'RITANT, *Irritans, Irritamen, Irritamentum*. Same etymon. That which causes irritation or pain, heat and tension; either mechanically, as puncture, acupuncture, or scarification; chemically, as the alkalies and acids; or in a specific manner, as cantharides.

Irritants are of great use in the treatment of disease.

IRRITATING POISON, see Poison.

IRRITATIO, Erethism, Gargale, Irritation.

IRRITATION, *Irritatio, Erethis'mus*. Same etymon. The state of a tissue or organ, in which there is excess of vital movement; commonly manifested by increase of the circulation and sensibility. Broussais defines irritation to be;—the condition of an organ, the excitation of which is carried to so high a degree, that the equilibrium resulting from the balance of all the functions is broken. In this signification, he also uses the word *surirritation*, which he considered as a higher degree, and as the essential cause of fever. Irritation is the precursor of inflammation.

IRRITATION, MORBID, *Constitutional irritation, Irritative fever, Pseudo-fever*, is that excitement which occurs after injuries done to the body, or to any part thereof;—constituting cases of diseased sympathy.

IRRITATIVE FEVER, Irritation, morbid.

IRRORATIO, Irrigation.

IS, Fibre.

ISADELPH'IA, from *isos*, 'equal,' and *adelphos*, 'brother.' A double monstrosity, in which both bodies are equally and perfectly developed, and possessed of independent vital organs; and which are united by parts of little significance.—Gurlt.

ISATIS TINCTORIA, *Woad, Pastel*. Ord. Capparidaceae. A European plant, whose leaves have a fugitive pungent smell, and an acrid durable taste. They are not used, however, in medicine at the present day, but are the source of the dye-stuff, *woad*.

ISCA, *Boletus igniarius*.

ISCHÆ'MIA, from *ισχω*, 'I retain,' and *αἷμα*, 'blood.' Morbid retention or suppression of an habitual flux of blood, as of the hemorrhoidal or menstrual flux or of epistaxis. See *Hæmatostatica*.

ISCHÆ'MON, *Ischa'mum*. Same etymon. A medicine which restrains or stops bleeding.

IS'CHESIS, from *ισχειν*, 'to retain.' Suppression or retention of a discharge or secretion.

IS'CHIA, MINERAL WATERS OF. In this volcanic isle, five miles from Naples, there are several thermal waters, one of which—that of Gurgitello—raises the thermometer of Fahr. to 167°.

ISCHIACUS, Ischiatic.

ISCHIADELPHUS, Ischiopages.

ISCHIADIUS MORBUS, Neuralgia femoro-poplitea—i. Nervus, Sciatic nerve.

ISCHIAOCELE, Ischiocele.

ISCHI'AGRA, *Is'chias, Ischul'gia*, from *ισχιον*, 'the haunch,' and *αγρα*, 'a seizure.' A name given to ischiatic gout. *Femoro-popliteal neuralgia* has, also, been so called.

IS'CHIAL, *Ischia'lis*. Relating or belonging to the ischion.

ISCHIALGIA, Ischiagra, Neuralgia femoro-poplitea.

ISCHIAS, Ischiagra—i. *Nervosa Antica*, Neuralgia femoro-prætibialis—i. *Nervosa Catumbii*, Neuralgia femoro-poplitea—i. *Nervosa digitalis*, Neuralgia cubito-digitalis—i. *Nervosa postica*, Neuralgia femoro-poplitea—i. à Sparganoesi, Phlegmatia dolens.

ISCHIATIC, *Ischiad'ic, Ischiat'icus, Ischiad'icus, Isch'iscus, Sciatic'ic*, (F.) *Ischiatique*, from *ισχιον*, 'the haunch;' whence the word *Sciatic*. An epithet given to parts connected with the ischium. The SCIATIC NOTCHES, (F.) *Échan-crures Ischiatiques*, are formed by this bone; and the ISCHIATIC SPINE belongs to it, and gives attachment to the small sacro-sciatic ligament. The TUBEROSITY OF THE ISCHIUM, *Os ischiadicum, Tuber Ischii*, (F.) *Tubérosité ischiatique*, is formed by it. It receives the insertions of different muscles of the thigh, and forms the projection on which the body rests when seated.

ISCHIATIC ARTERY, *Sciatic Artery*, (F.) *Artère Femoro-Poplitée* (Ch.) arises singly from the hypogastric, or with the gluteal; and seems to be really a continuation of the trunk of the hypogastric. It issues from the pelvis, at the lower part of the great sciatic notch; and, afterwards, divides into a considerable number of branches, which are distributed particularly to the posterior and superior region of the thigh.

ISCHIATIC REGION, *Sciatic region; Regio ischiad'ica*. The region of the hip.

ISCHIATIC VEIN presents the same arrangement as the artery.

ISCHIATICUS, Sciatic.

ISCHIAOCELE, Ischiocele.

ISCHIDRO'SIS, *Sudor'is suppressio, Oligid'ria*, from *ισχω*, 'I restrain,' and *ἰδρωσ*, 'sweat.' Suppression of perspiration.

ISCHIO. In composition, belonging or relating to the ischium.

ISCHIOBLEN'NIA, *Ischoblen'nia*, from *ισχω*, 'I restrain,' and *βλεννα*, 'mucus.' The suppression of a morbid but habitual discharge of mucus.

ISCHIO-CAVERNO'SUS. Belonging to the ischium and corpus cavernosum.

ISCHIO-CAVERNOSUS muscle, *Collateralis, Director seu Erector seu Sustentator Penis*, (F.) *Érecteur du Pénis, Ischio-wéthral*, (Ch.) is a small, long, flat muscle, which surrounds the origin of the corpus cavernosum. It is fixed below to the inner side of the tuberosity of the ischium, and above to the root of the penis, where it is confounded with the fibrous membrane of the corpus cavernosum. It draws the root of the penis downwards and backwards.

ISCHIO-CAVERNOSUS of the female, *Erector seu Sustentator seu Superior rotundus clitoridis*, (F.) *Ischio-clitoriden, Ischio sous-clitorien*, (Ch.), is arranged nearly as in the male, but is less bulky. It arises, by aponeurosis, from the tuberosity of the ischium, and terminates by embracing the corpus cavernosum of the clitoris, to the erection of which it appears to contribute.

ISCHIOCE'LE, *Ischiadoc'le, Ischiadoc'le, Hernia ischiatic'a*; from *ισχιον*, 'the ischium,' and *κηλη*, 'tumour.' Ischiatic hernia. One in which the viscera issue by the great sciatic notch. It is a rare disease. The protruded parts must be reduced—the patient being placed in a favourable position, and they must be retained by a bandage.

ISCHIO-CLITORIANUS. That which is connected with the ischium and clitoris.

ISCHIO-CLITORIAN ARTERY of Chaussier is a division of the internal pudic, which furnishes the two arteries of the clitoris—the *superficial* and *deep-seated*.

ISCHIO-CLITORIAN NERVE, of the same professor, is the superior branch of the pudic nerve, which is distributed to the clitoris.

ISCHIO-CLITORIDIEN. Ischio-cavernous—i. Cocegyeus, Cocegyeus—i. Crèti-tibial, Semi-tendinosus—i. Femoralis, Adductor magnus—i. *Fémoro-péronier*, Biceps flexor cruris—i. *Périnéal*, Transversus perinæi—i. *Popliti-fémoral*, Semi-membranosus—i. *Popliti-tibial*, Semi-membranosus—i. *Prétibial*, Semi-tendinosus—i. *Sous-clitorien*, Ischio-cavernosus—i. *Sous-trochanterien*, Quadratus femoris—i. *Spino-trochanterien*, Ischio-trochanterianus—i. *Uréthral*, Ischio-cavernosus.

ISCHIODYMIA, see Somatodymia.

ISCHION, *Ischium*, Os *ischii*, Os *coxen'dicis*, Hip-bone, Huckle-bone, Seat-bone, (Sc.) *Heuck-bone*, *Huckebane*, *Lunyjebane*, (Prov.) *Chancebone*, *Pinbone*, (F.) *Os de l'assiette*. The lowermost of the three portions which compose the os innominatum in the foetus and young individual. The inferior region of the same bone has, also, been called *ischium*, in the adult. According to Hesychius, the ancients designated by the word *ischion* the capsular ligament of the coxo-femoral articulation, as well as the articulation itself. Some derive the word from *ισχς*, the lumbar region; others from the verb *ισχω*, 'I arrest,' 'I retain'—because that bone serves as a base or support for the trunk, when we are seated.

ISCHION, Haunch.

ISCHIO-NEURALGIA, Neuralgia femoro poplitea.

ISCHIOPA'GES, *Ischiadelphus*, from *Ischion*, and *παω*, 'I fasten.' A monstrous union of two fetuses, in which they are attached to each other by the ischia.—Geoffroy St. Hilaire. See Hypogastrodidymus.

ISCHIOPHTHISIS, *Tubes coxa'ria* seu *ischiad'ica*, *Phthisis ischiad'ica*, from *ισχον*, 'ischium,' and *phthisis*. Phthisis in consequence of suppuration of the hip-joint. See Coxarum morbus.

ISCHIO-PROSTATICUS. Winslow, Sancto-rini, Albinus, and Sömmerring, have given this name to the fibres of the transversus perinæi muscle, which go towards the prostate.

ISCHIO-RECTAL FOSSA, Perineal fossa.

ISCHIOSIS, Neuralgia femoro poplitea.

ISCHIO-TROCHANTERIANUS, *Gem'ini*, *Gemel'li*. Part of the *Marsupialis* of Cowper; *Car'neus Marsup'ium*, *Ischio-spini-trochanterius*, *Secundus et tertius quadrigen'imus*, (F.) *Muscle cannelé*, *Accessoire à l'obturateur interne*. Two small, fleshy bundles, long and flat, which arise,—the *superior* on the outside of the sciatic spine; the *inferior* behind the tuberosity of the ischium. Both pass horizontally outwards, and are attached to the tendon of the obturator internus, which they accompany into the fossa of the trochanter. These muscles are rotators of the lower limb outwards. They can, also, turn the pelvis on the femur, in standing on one foot.

ISCHIUM, Ischion.

ISCHL, MINERAL WATERS OF. These springs, which are in Upper Austria, contain iodine and bromine.

ISCHNOPHO'NIA, from *ισχυος*, 'slender,' and *φωνη*, 'voice.' Slenderness of voice.—Hippocrates, Galen.

ISCHNOTES, Emaciation.

ISCHO, *ισχω*, 'I keep back,' 'I restrain,' 'I hold firm.' Hence:

ISCHOBLENNIA, Ischioblenia.

ISCHOCENO'SIS, from *ισχω*, 'I arrest,' and *κνωσις*, 'evacuation.' Retention or suppression of a natural evacuation—as of the menses.

ISCHOCOL'IA, from *ισχω*, 'I arrest,' and *χολη*, 'bile.' Retention of the secretion of the biliary secretion.

ISCHOCOLIA, Constipation.

ISCHOCOPRIA, Constipation.

ISCHOGALAC'TIA, *Reten'tio lactis*, from *ισχω*, and *γαλα*, 'milk.' Want of milk in the mammae.

ISCHOLOCH'IA or *Ischolochi'a*, *Lochios'chesis*, *Suppres'sio lochii* rum, from *ισχω*, 'I restrain,' and *λοχεια*, 'the lochial discharge.' Retention of lochiorum. Suppression of the lochial discharge.

ISCHOMENIA, Amenorrhœa.

ISCHONEURALGIA, Neuralgia femoro-poplitæa.

ISCHOPHONIA, Balbuties.

ISCHOSPERR'MIA, from *ισχω*, 'I retain,' and *σπερμα*, 'sperm'; *Sem'inis reten'tio*. Retention or suppression of the spermatic secretion.

ISCHURET'IC, *Ischuræticum*. Same etymon as the next. A medicine for relieving suppression of urine. A diuretic.

ISCHUR'IA, *Urinæ suppres'sio*, *Stoppage of urine*, *Suppres'sio lot'ii*, *Paru'ria reten'tio'nis*; from *ισχω*, 'I arrest,' 'I retain,' and *ουρον*, 'urine.' Retention of, or impossibility of discharging, the urine, *Anurorrhœe*—Piorry. Often restricted to suppression of the secretion, *Renal Ischuria*, *Anu'ria*, *Anure'sis*, *Nephra'uria*, *Anurocrinie*—Piorry.

Ischuria Vera is that in which the urine, having accumulated in the bladder, the patient is unable to pass it, notwithstanding the inclination which constantly distresses him. In false *ischuria*, *Paru'ria inops*, *P. reten'tio'nis rena'lis*, *Ischuria notha* seu *spu'ria*, of some, (F.) *Suppression d'Urine*, owing to some disease of the kidney or ureters, the urine cannot reach the bladder.

Ischuria has likewise received various other names, according to the seat and character of the obstacle which opposes the exit of the urine: hence the expressions—*Renal*, *Ureteric*, *Vesical*, *Uréthral*, and *Calculus Ischuria*. *Vesical Ischuria* is generally used synonymously with Retention of Urine, which see.

ISCHURIA CYSTO-PHEGEMATICA, Cystirrhœa—i. *Phimosica*, see *Phimosicus*—i. *Spasmodica*, *Cystospasmus*—i. *Spuria*, see *Ischuria*—i. *Uréthralis* a phimosi, see *Phimosicus*—i. *Vera*, see *Ischuria*—i. *Vesicalis*, Retention of urine.

ISIC'IUM, *Paicos*, *Isic'ium*, *Tomac'ulum*, *Tom'actum*, *Tomac'inum*. A kind of sausage, formed of mixed meat and spices.

ISINGLASS, Ichthyocolla.

ISIR, Elixir.

ISIS NOBILIS, see Coral.

ISLAND OF REIL, *Insula cerebri*.

ISLE OF WIGHT, CLIMATE OF. This beautiful island is a favourable summer retreat for invalids. Undercliff is the situation chosen as a winter residence for phthisical valetudinarians. Cowes, Niton, Sandown, Shanklin, and Ryde, are delightful summer residences.

ISNARDIA PALUSTRIS, *Ludwigia palustris*.

ISOCH'RONOUS, *Isch'ronus*, *Ischron'ius*, *Isod'romus*, from *ισος*, 'equal,' and *χρονος*, 'time.' That which takes place in the same time, or in equal times. The pulsations of the arteries, in various parts of the body, are nearly isochronous.

ISOCH'RYSON, *Auro compar*; from *ισος*, 'equal,' and *χρυσος*, 'gold.' A collyrium, described by Galen as worth its weight in gold. Libavius has also given this name to an amalgam, made with equal parts of antimony and mercury.

ISOCRATES, from *ισος*, 'equal,' and *κρατος*, 'strength.' A mixture of equal parts of wine and water.—Hippocrates.

ISODROMUS, Isochronous.

ISOLUSINE, see *Polygala senega*.

ISOMERIC, *Isomer'icus*, *Isom'erus*, *Isomce'rus*, *Isom'orus*, from *isos*, 'equal,' and *μερος*, 'part.' An epithet applied to different bodies which agree in composition, but differ in properties. The condition is termed *Isomerism*.

ISOMERISM, see *Isomeria*.

ISOMORPHISM, see *Isomorphous*.

ISOMORPHOUS, *Isomor'phus*, *Isomor'phicus*, from *isos*, 'equal,' and *μορφη*, 'form.' An epithet applied to different bodies which have the same crystalline form. The condition is called *Isomorphism*.

ISONANDRA GUTTA, see *Gutta percha*.

ISOPATHIA, *Isop'athy*, from *isos*, 'equal,' and *παθος*, 'disease.' This term has been used by some of the German writers to signify the cure of diseases by the disease itself, or its products;—under the hypothesis, that every contagious disease contains in its contagious matter the means for its cure;—thus, that variola may be cured by homœopathic doses of variolous matter; syphilis, with venereal matter, &c., &c. Others have given to isopathy another form;—maintaining, that every diseased organ has its remedy in the same organ,—that eating liver, for example, will remove disease of the liver!

Isopathy has been used by an American writer—Dr. J. M. B. Harden, of Georgia—to mean "*Parallelism of Diseases*;" "the disposition of diseases to 'anastomose with each other,' or to wear each other's livery."

ISSUE, *Fonticulus*—i. *Peas*, *Aurantia curasaventia*, see *Fonticulus*.

ISTHME DU GOSIER, *Isthmus* of the fauces.

ISTHMION, *Isthmus*, *Pharynx*.

ISTHMITIS, *Inflammat'io Fau'cium* seu *Pal'oti*, *Angi'na simplex* seu *mitis* seu *fau'cium*, *Angor fau'cium*, *Cynan'che simplex*, (F.) *Angine simple*, from *isthmus*, and *itis*, denoting inflammation. Inflammation of the *isthmus faucium*, and of the fauces. See *Cynanche pharyngea*, and *Angina*.

ISTHMIUM, *Isthmus*.

ISTHMOCATAR'RHUS; from *ισθμος*, 'the fauces,' and *καταρροος*, 'catarrh.' *Catarrh* of the fauces.

ISTHMODYN'IA, from *ισθμος*, 'the fauces,' and *δυνη*, 'pain.' Pain in the fauces.

ISTHMOPL'GIA; from *ισθμος*, 'the fauces,' and *πληγη*, 'a stroke.' Paralysis of the fauces.

ISTHMOS, *Pharynx*.

ISTHMYPHUS, *Cynanche maligna*.

ISTHMUS, *Isth'mion*, *Isth'mium*, (F.) *Isthme*. A tongue of land joining a peninsula to a continent, or which separates two seas. Anatomists have given the name *Isthmus of the Fauces*, *I. Fau'cium*, *Clastrum Gut'turis*, (F.) *Isthme du gosier*, to the strait which separates the mouth from the pharynx. It is formed above by the velum palati and uvula; at the sides, by the pillars of the fauces and the tonsils; and below, by the base of the tongue. Also, the *Fauces*.

ISTHMUS OF THE ENCEPHALON. The narrowed and constricted portion of the encephalic mass, situate between the cerebrum, the cerebellum and the medulla oblongata, which corresponds to the free margin of the tentorium cerebelli, and comprises the pons varolii, and middle peduncles of the cerebellum, the peduncles of the cerebrum, the tubercula quadrigemina, the superior peduncles of the cerebellum, and the valve of Vieussens.—*Cruevilhier*.

ISTHMUS OF THE FOSSA OVA' LIS: *Isthmus Vieussen'is*, *Striga cartilagino'sa cordis*, is the prominent arch formed above the fossa ovalis by the union of the two pillars which bound the cavity.

ISTHMUS OF THE THYROID GLAND is a narrow band that unites the two chief lobules composing the thyroid gland.

ISTHMUS HEPATIS, see *Lobulus anonymus*—i. *Urethra*, see *Urethra*—i. *Uteri*, see *Uterus*—i. *Vieussenii*, *Isthmus* of the fossa ovalis, see *Ovalis fossa*.

ITALIAN JUICE, *Extractum glycyrrhizæ*.

ITALICUS MORBUS, *Syphilis*.

ITCH, *Psora*—i. *Animalcule*, see *Psora*—i. of *Animals*, *Scabies ferina*.

ITCH, *BAKERS'*, see *Psoriasis*—i. *Barbers'*, *Sycosis*—i. *Bricklayers'*, *Eczema impetiginodes*.

ITCH, *COWPOX*. A cutaneous eruption, observed and described by Gölis, which appeared after the fourteenth day from vaccination, and consisted of isolated vesicles, often filled with a puriform fluid.

ITCH, *DRUNKARDS'*, *Psora ebriorum*—i. *Grocers'*, see *Psoriasis*.

ITCH INSECT, see *Psora*—i. *Mite*, see *Psora*.

ITCH, *VENEREAL*, *Scherlievo*.

ITCH WEED, *Veratrum viride*.

ITCHING, *Parap'sis pruri'tus*, *Antal'gia pruriginosa*, *Pruri'tus*, *Pruri'go*, *Cnesmos*, *Cnesma*, (Sc.) *Youk*, *Yuke*, (F.) *Prurit*. *Démangeaison*. A sensation, more inconvenient than painful, seated especially at the surface of the body, which provokes the patient to scratch the part. It may be either an external or an internal sensation: that is, produced by an external body, or by some modification in the organic actions of the part to which it is referred.

ITEA, *Salix*.

ITER A TERTIO AD QUARTUM VENTRICULUM, *Aqueductus Sylvii*—i. *Dentis*, see *Gubernaculum dentis*—i. *Femineum*, *Perinaeum*—i. *ad Infundibulum*, *Vulva*—i. *a Palato* *ad Arem*, *Eustachian tube*—i. *ad Quartum ventriculum*, *Aqueductus Sylvii*—i. *ad Tertium ventriculum*, *Vulva*—i. *Urinæ*, *Urethra*—i. *Urinarium*, *Urethra*.

ITHYPHAL'LUS, from *εὐς*, 'straight,' and *φαλλος*, 'penis.' An anulet, in the form of an erect penis, anciently worn round the neck, to which were attributed alexiterial properties.

ITINERARIUM, *Conductor*, *Director*.

ITIS, from *ιτης*, 'bold,' 'rash,' [?]. A suffix denoting inflammation:—as *encephalitis*, inflammation of the encephalon;—*Pleuritis*, inflammation of the pleura. According to some, as *Kraus*, *encephalitis*, for example, was, with the ancients, the adjective agreeing with *εγκεφαλος*, 'a disease,' understood to express a disease of the encephalon, and it was only in modern times, that the suffix was used to denote inflammation.

IVA, *Teucrium iva*—i. *Arthritica*, *Teucrium chamæpitys*—i. *Moschata Monspeliensium*, *Teucrium iva*—i. *Pecania*, *Smilax sarsaparilla*.

IVETTE MUSQUÉE, *Teucrium iva*—i. *Pelite*, *Teucrium chamæpitys*.

IVOIRE, *Ivory*.

IVORY, *Ebur*, *El'ephaz*, (F.) *Ivoire*. The tusk of the elephant. It is chiefly composed of phosphate of lime, and is used for the fabrication of pessaries, artificial teeth, handles of instruments, &c. Formerly, when calcined to whiteness, it entered into some pharmaceutical preparations. It was regarded as astringent and anthelmintic, and was called *Spo'dium*.

The *dentine* is the ivory of the human tooth. See *Tooth*.

IVORY BLACK, see *Carbo animalis*—i. *Vegetable*, *Phytelephas macrocarpa*.

IVRAIE, *Lolium temulentum*.

IVRESSE, *Temulentia*—i. *Iodique*, see *Iodine*—i. *Quinique*, *Quininism*.

IVROGNERIE, see *Temulentia*.

IVY. *Hedera helix*, *Kalmia angustifolia*—i. American, *Ampelopsis quinquefolia*—i. Big, *Kalmia latifolia*—i. Fine-leaved, *Ampelopsis quinquefolia*—i. Ground, *Glechoma hederacea*.

IXIA, *Varix*, *Viscum album*.

IXINE, *Atractylus gummifera*.

IXIR, Elixir.

IXOS, *Viscum album*.

IXUS, *Galium aparine*.

IXYOMYELI'TIS, from *ixus*, 'the lumbar region,' *μυελος*, 'marrow,' and *itis*, denoting inflammation. Inflammation of the spinal marrow in the lumbar region.

IXYS, *ixus*, *Ix'ya*, *Ix'ya*. Used by different authors for the ilia, flanks, and loins; most frequently for the last.

J.

JABOT, Inguivies.

JACA INDICA, *Thymus mastichina*.

JACEA, Tricolor, *Viola Tricolor*—j. Ramosissima, *Centaurea calcitrapa*.

JACENS, Sick.

JACK OF THE BUTTERY, *Sedum acre*—j. in the Hedge, *Alliaria*.

JACKET, BARK. A jacket of flannel or cotton in which powdered bark is quilted. It is occasionally worn as an antiperiodic and tonic.

JACKSON'S BATHING SPIRITS, *Linimentum saponis compositum*.

JACOBÆA, *Senecio Jacobæa*.

JACOBÉE, *Senecio Jacobæa*.

JACOB'S COAT, *Tunica Jacobi*—j.'s Ladder, false, *Polemonium reptans*—j.'s Membrane, *Tunica Jacobi*—j.'s Ulcer, *Lupus*.

JACOBSON, VENOUS SYSTEM OF, see Venous system.

JACOBSON'S ANASTOMOSIS, see Petrosal ganglion.

JACTA'TION, *Jactita'tion*, *Jacta'tio*, *Jacta'tus*, *Rhaptas'mos*, *Jactita'tio*, from *jactare*, 'to toss about.' Extreme anxiety; excessive restlessness;—a symptom observed in serious diseases.

JACTITATIO, Inquietude, Jactation.

JADE NÉPHRITE ou ORIENTALE, Nephreticus lapis.

JAGGERY, Spirit Arrack, see Toddy.

JAGRE, see Tari.

JALAP, *Convolvulus jalapa*, *Phytolacca decandra*—j. *Blanc*, *Convolvulus mechoacan*—j. Cancer root, *Phytolacca decandra*—j. Fusiform, light male, see *Convolvulus jalapa*—j. Wild, *Convolvulus panduratus*.

JALAPA, *Convolvulus jalapa*.

JALAPINE, see *Convolvulus jalapa*.

JALAPIUM, see *Convolvulus jalapa*.

JALAPPA, *Convolvulus jalapa*—j. Alba, *Convolvulus mechoacan*.

JALEYRAC, MINERAL WATERS OF. These waters are situate two leagues from Mauriac in France, on the road from Clermont in Auvergne. The waters contain lime and carbonate of soda, and are esteemed to be tonic, aperient, &c.

JALOPA, *Convolvulus jalapa*.

JAMAICA, see West Indies—j. Bark tree, *Bursera gummifera*.

JAMAICINE, see *Geoffræa inermis*.

JAMBES DE BARBADE, see Elephantiasis—j. du Cervelet, *Corpora testiformia*.

JAMBIER, Tibial—j. Antérieur, Tibialis anticus—j. Grêle, Plantar muscle—j. Postérieur, Tibialis posticus.

JAMES'S ANALEPTIC PILLS, see Analptica.

JAMESTOWN WEED, *Datura stramonium*.

JANAMUNDA, *Geum urbanum*.

JANDERS, Icterus.

JAN'NICEPS, from *Janus*, a divinity with two faces, and *caput*, 'head.' A double monster, the two bodies intimately united above a common umbilicus; and a double head, with the faces looking in opposite directions.—I. G. St. Hilaire.

JANIPHA MANIHOT, *Jatropha manihot*.

JANITOR, Pylorus.

JANITRIX, Porta vena.

JANON-TARENTISME, Tarantismus.

JANUS, Dicephalus.

JAQUIER, *Artocarpus*.

JARRET, Poples.

JARRETIER, *Popliteus muscle*, Hamstring.

JARRETIÈRE (F.), *Peris'celis*, a garter, from *garetum* or *garretum*, in low Latin, the ham. A kind of furfuraceous herpes, which occupies the part of the leg where the garter is worn.

JASMIN, *Jasminum officinale*.

JASMINUM ARABICUM, *Coffea Arabum*.

JASMINUM OFFICINALE, *Jasminum*, *Jesem'ianum*, the *Jessamine*, (F.) *Jasmin*. Ord. *Jasmineæ*. The flowers of this beautiful plant have a very fragrant smell, and a bitter taste. They afford, by distillation, an essential oil; which is much esteemed in Italy, for rubbing paralytic limbs, and in the cure of rheumatic pains.

JASPER, *Jaspis*.

JASPI, *Jasper*. A precious stone, supposed by the ancients to be capable of arresting hemorrhage when worn as an amulet.

JATAMANSI, Sumbul.

JATROPHA CURCAS, (from *iaopat*, 'I heal,' and *ρεφα*, 'I nourish.') *Ric'inius major*, *Ricinioides*, *P'ineus* seu *Curcas purgans*, *Pinho'nes* *Indici*, *Faba cathartica*, *Avella'na cathartica*, *Nux cathartica*, *Americ'na seu Barbadesis*, *Physic Nut*, (F.) *Pignon d'Inde*, *Médecinier cathartique*, *Grand haricot de Pérou*, *Pignon de Barbarie*, *Noix cathartique*, *N. Américaine* ou *des Barbades*. Nat. Order, *Euphorbiaceæ*. The seeds of this plant afford a quantity of oil, given, in many places, like the castor oil; to which, indeed, it is nearly allied. They contain a peculiar acid, the *Jatrophie* or *Igasurie*.

The seeds of *JATROPHA MULTIFIDA* are called *Purging Nuts*, and give out a similar oil.

JATROPHA ELASTICA, see *Caoutchouc*.

JATROPHA MANIHOT, *Jan'ipha Man'ihot*, *Manihot utilis'ima*, *Bitter Cassa'va*. The plant affording the *Cassa'da* or *Cassa'va* Root, *Caca'vi*, *Cassa'cê*, *Cazabi*, *Pain de Madagascar*, *Ric'innus Minor*, *Man'ioc*, *Mandioc*, *Magnoc*, *Maniot*, *Yucca*, *Maniibar*, *Aipi*, *Aipima cozera*, *Aipipoca*, *Jani'pha*. The juice of the root is extremely acrid and poisonous. What remains, after expressing it, is made into cakes or meal; of which the cassada or cassava bread is formed. This bread constitutes a principal food of the inhabitants of those parts where it grows.

The fecula of the root forms *Tapioca*, *Cipipe*, *Fec'ula Tapi'o'ka*, *Am'yum manihot'icum*, which

is very nutritious. It may be prepared in the same manner as sago.

A factitious Tapioca is met with in the shops, which is in very small, smooth, spherical grains, and is supposed to be prepared from potato starch. It is sold under the name *pearl tapioca*.

JATROPHA URENS. A most deadly plant. The touch of its five bristly stings had nearly cost Mr. Smith, the curator of Kew gardens, his life.

JAUNDERS, Icterus.

JAUNDICE, Icterus—*j.* Black, Melæna—*j.* Green, Icterus viridis—*j.* Lead, Icterus saturninus—*j.* Red, Phenigmus.

JAUNE, Yellow—*j.* d'Œuf, see Ovum.

JAUNISSE, Icterus.

JAW BONE, Maxillary Bone.

JAW-DISEASE, PHOSPHORUS, see Phosphorus.

JEAN-DE-GLAINES, ST., MINERAL WATERS OF. These waters are situate two leagues from Billom in Auvergne. They contain chloride of calcium, and carbonate of magnesia; dissolved in an excess of carbonic acid.

JEANNETTE, Narcissus pseudonarcissus.

JECINUS, Liver.

JÉCORAIRE, Hepatic.

JECORAL, Hepatic.

JECORARIA, Marchantia polymorpha.

JECORARIUS, Hepateros, Hepatic.

JECORIS VOMICA, see Hepatitis.

JECTIGATIO. A word used by Van Helmont for a species of epilepsy or convulsion.

JECUR, Liver—*j.* Uterinum, Placenta.

JEFFERSONIA BARTONI, J. Diphyl'la, Common Twinleaf, Yellow Root, Rheumatism Root, Helmet Pod, Ground Squirrel Pea. An indigenous plant, belonging to *Nat. Ord.* Berberideæ. *Ser. Syst.* Octandria Monogynia; which possesses medical properties analogous to those of hydrastis.

JEJUNITAS, Hunger.

JEJUNITIS; a term of hybrid formation,—from *jejunum*, 'the intestine jejunum,' and *itis*, a suffix denoting inflammation. Inflammation of the jejunum.

JEJUNIUM, Hunger, Fasting.

JEJUNUM, from *jejunus*, 'empty,' *Nestis, vovis, Nestia*. The part of the small intestine comprised between the duodenum and ileum. It has been so called, because it is almost always found empty in the dead body.

JELLY, Gelly—*j.* of the Cord, Gelatin of Wharton—*j.* Rice, see *Oryza*—*j.* Water, Brasenia hydropeltis.

JENEPERE, Juniperus communis.

JENKINSONIA ANTIDYSENTERICA, Pelargonium antidysentericum.

JERGON, see Arrow-poison.

JERKING RESPIRATION, Respiration, jerking.

JERKS, see Mania, dancing.

JEROFFLERIS, Dianthus caryophyllus.

JERSEY, CLIMATE OF. Jersey is the largest of the islands of the British Channel, and is most frequented by invalids. Its climate closely resembles that of the south-west coast of England, and especially of Penzance, and it is adapted to the same class of invalids.

JERUSALEM OAK OF AMERICA, Chenopodium anthelminticum.

JESEMINUM, Jasminum officinale.

JESSAMINE, Jasminum officinale—*j.* Yellow, Gelseminum nitidum.

JETICUCU, Batata de purga.

JEUNESSE, Adolescence.

JEWBSH, Pedicularis tithymaloïdes.

JEWEL WEED, Impatiens balsamina.

JEW'S EAR, Peziza auricula—*j.* Harp, Trilium latifolium.

JIGGER, Chique.

JIMPSON WEED, Datura stramonium.

JIMSTON WEED, Datura stramonium.

JIZZEN-BED, see Parturition.

JOANNESIA PRINCEPS, Anda.

JOANNETTE, MINERAL WATERS OF.

Several springs are found at Joannette, about five leagues from Angers, in France. The waters are both cold and warm, and contain sulphate of lime; subcarbonate of soda; chloride of sodium; some iron; chloride of calcium, and, sometimes a kind of saponaceous matter. They are chiefly used as a tonic in chlorosis, leucorrhœa, &c.

JOE PYE, Eupatorium perfoliatum, and E. purpureum.

JOHNE, MINERAL WATERS OF. Johnne is a village near Dol in Franche-Comté, France, where is a mineral spring, which contains subcarbonate of soda, and subcarbonates of magnesia and iron. It is used chiefly as a bath in certain diseases of the skin, and in atonic affections.

JOINT, Articulation—*j.* Ball and Socket, Enarthrosis—*j.* Dove-tail, Suture—*j.* Stiff, Ankylosis.

JONAS, Icterus.

JONAS, MINERAL WATERS OF. This spring is situate to the south-west of Bourbon-l'Archambault, in France. The waters contain chloride of calcium and sulphate of lime; chloride of sodium, and sulphate of soda; carbonate of iron, and carbonic acid gas. They are tonic and aperient.

JORDAN'S WHITE SULPHUR SPRINGS. These are in Virginia, five or six miles from Winchester. They are used in the same cases as sulphureous springs in general.

JOT-GUT, Rectum.

JOUAN, ST., MINERAL WATERS OF. St. Jouan is a village, near Saint-Malo, in France. The water is a cold chalybeate.

JOUBARBE ACRE, Sedum—*j. des Toits*, Sempervivum tectorum—*j. des Vignes*, Sedum telephium.

JOVE, Genu.

JOULOS, Julus.

JOUR, Dies.

JOUS CANICULAIRES, Dog days—*j.* Critiques, Critical days—*j.* Indicateurs, Indicating days—*j.* Paroxyntiques, see Paroxysmic.

JOVIS FLOS, Crocus—*j.* Glans, Juglans.

JUCATO CALLELOE, Phytolacca decandra.

JUDE'US, Juda'icus seu Phœnic'ites *Lapis, Tecol'ithos*, (F.) *Pierre Judaïque*. A stone, found in Judæa, Palestine, &c. Called, also, *Lapis Syriacus*. It was formerly esteemed to be diuretic and lithontriptic.

JUDGMENT, *Judic'ium*, (F.) *Jugement*. The faculty of the intellect, by which ideas are compared with each other, and their relative worth appreciated.

JUDICATORII DIES, Critical days.

JUDICIUM, Judgment.

JUGA ALVEOLARIA, see Alveolar arches.

JUGA CEREBRALIA. Prominences on the inner surface of the bones of the cranium, between the digital impressions.

JUGAL, Zygomatic—*j.* Process, Zygomatic process.

JUGAL REGION, *Re'gio Jugal'is*. The region of the cheek-bone.

JUGALE OS, Mala os.

JUGALIS, Zygomatic.

JUGALIS SUTURA, from *zygon, jugum*, (from *ungere, junctum*, 'to join,') 'a yoke.' The suture which unites the *Os Mala* or *Os Jugal'e*

with the superior maxillary bone. Also, the sagittal suture.

JUGEMENT. Judgment.

JUGOLINE. Sesamum orientale.

JUGLANS, BLACK. *J. nigra*.

JUGLANS CINEREA (*Ju-glans*, quasi *Jovis glans*), *Butter or Oil Nut, White Walnut.* *Ord.* Juglandaceæ. The inner bark of the root of this tree, *Juglans* (Ph. U. S.), which is abundant in the United States, is used in medicine, in the form of extract, as an efficacious and mild laxative, in doses of from 10 to 20 grains. It is in the secondary list of the Pharmacopœia of the United States.

JUGLANS NIGRA, Black Juglans, Black Walnut; indigenous. The rind of the unripe fruit is said to remove ringworm and tetter. A decoction has been given as an anthelmintic.

JUGLANS RE'GIA, Juglans, Nux Juglans, Carya basili'ca, kapva, Carya. The Walnut, French nut, (F.) *Noyer*, the tree; *Noix*, the nut, *kapvov, Car'yon*. The unripe fruit, in the state in which it is pickled, was formerly esteemed to be anthelmintic. The putamen or green rind of the walnut has been celebrated as a powerful antisyphilitic, and used as a sort of diet drink. The kernel is an agreeable article of dessert; but, like all nuts, is difficult of digestion. The expressed oil, *Huile de Noix*, is used in France as an aliment, and, like other fixed oils, is laxative.

Various preparations of the leaves have been recommended in scrofulous affections.

JUG'ULAR, Jugula'ris, from *jugulum*, (itself from *jugum*, because the 'yoke' was attached there,) 'the throat.' Relating to the throat.

JUGULAR FOSSA, Fossa Jugula'ris, is a cavity in the petro-occipital suture. It is formed by the petrous portion of the temporal bone, and by the occipital bone, and lodges the origin of the internal jugular vein. It is, sometimes, called *thimble-like cavity*.

JUGULAR VEINS, Venæ Jugula'res seu sphagit'ides seu apoplecticæ seu sopora'les. These are situate at the lateral and anterior parts of the neck. They are two on each side; one external, the other internal. 1. The External Jugular Vein, *Trachélo-sous-cutanéë*, (Ch.) is of less size than the internal. It descends, almost vertically, along the anterior and lateral part of the neck, from the cervix of the lower jaw to the subclavian vein, into which it opens, a little above the internal jugular. It is formed by the internal maxillary, superficial temporal, and posterior auricular veins. It is this vein which is commonly opened in bleeding in the neck. 2. The Internal Jugular Vein, *V. Céphalique* (Ch.), *Vena apoplecticæ*, is much larger, and more deeply seated than the preceding. It descends, vertically, along the anterior and lateral part of the neck, from the posterior part of the foramen lacerum posterius as far as the subclavian vein. It commences at the sinus of the jugular vein, (F.) *Golfe de la veine jugulaire*, and receives the blood, which returns by the sinus of the dura mater, and that of the facial, lingual, pharyngeal, superior thyroid, occipital, and diploic veins.

JUGULUM, Clavicle, Throat—j. Uteri, Collum uteri.

JUGUM PENIS, Presse-urèthre.

JUICE, EXPRESSED, see Succus.

JU'JUBE, Ju'juba, Ziz'yphum, Baccæ jujubæ, B. Ziz'yphi. A word of Arabic origin. The fruit of *Rhamnus Ziz'yphus* seu *Ziz'yphus vulga'ris.* *Ord.* Rhamnææ. A native of the south of Europe. It was formerly ranked amongst the pectoral fruits. It has an agreeable sweet taste. The fruits of two other species of *Zizyphus*—*Z. jujuba, Rhamnus jujuba, Manksana arbo'rea*, a native of the East Indies, and *Z. lotus, Rhamnus*

lotus, Zizyphus nit'ida seu *sati'va* seu *sylices'tris*, growing in North Africa, possess properties similar to those of *Zizyphus vulgaris*.

JUJUBE PASTE, Paste of jujubes.

JULAPIUM, Julep—j. Camphoræ, Mistura Camphoræ.

JULEB, Julep.

JULEP, Jula'pium, Jule'pus, Zula'pium, Juleb of the Persians. A sweet drink. A demulcent, acidulous or mucilaginous mixture.

JULEP. CAMPHOR, Mistura camphoræ—j. Camphræ, Mistura camphoræ—j. Chalk, Mistura cretæ—j. Mint, see Mint julep.

JULUS, Julos, Ioulos, ιουλος. The first down that appears upon the chin.—Rufus of Ephesus.

JUM, Lolium temulentum.

JUMBLE BEADS, see Abrus precatorius.

JUMEAU, Gemellus.

JUMEAUX DE LA JAMBE, Gastrocnemii.

JUMELLE, Gemellus.

JUMENTOUS URINE, Urine jumentæus.

JUMPERS, see Convulsionnaire.

JUNCTURA, Articulation.

JUNCUS ODORA'TUS, from jungere, junctum, 'to join,' in allusion to the use of rushes for bands. [?] *Andropogon schœnan'thus* seu *bicor'nis* seu *citra'tus* seu *citridod'rus*, *Cymbopogon schœnan'thus*, *Fœnum camelo'rum*, *Juncus aroma'ticus*, *Camel's hay*, *Sweet rush*, *Schœnan'thus*, *Schœnan'thum*, *Scœnan'thum*, *Cal'amus odora'tus*, *Squinan'thus*, *Gramen orienta'le*, (F.) *Jonc odorant*, *Foin de chameau.* *Ord.* Graminææ. The dried plant, which is generally procured from Turkey and Arabia, has an agreeable smell, and a warm, bitterish taste. It was formerly used as a stomachic and deobstruent.

JUNGLE FEVER. A variety of remittent occurring in the jungle districts of India.

JUNIPER TREE, Juniperus communis.

JUNIPERUM VINUM. Wine impregnated with juniper berries.

JUNIPERUS COMMUNIS, J. Suec'ica, Arceu'thoa, Ju'niper tree; Akat'alis, Akat'era, (Sc.) *Jenepere.* Family, Coniferae. Sex. Syst. Diœcia Monadelphica. The berries, *Juniper'us* (Ph. U. S.), (Sc.) *Eatin'-berries, Etnagh-berries*, are ordered in the pharmacopœias. Their odour is strong, but not unpleasant. Taste warm, pungent. Properties dependent upon essential oil, which they yield to both water and alcohol. Dose, ℥j to ʒss, rubbed with sugar. In Holland, juniper berries are used for flavouring gin. The oil, *O'leum junip'eri*, possesses the virtues of the plant. It is called, by Ruland, *Targar.* (F.) *Genévrier* (the plant), *Genêdre* (the berry).

JUNIPERUS LYCIA. This plant—*Thu'rea, Th. virga. Arbor thurifera*—has been supposed to afford the true frankincense, *Cedros, Olib'anum, Thus Libano'tos, Libano'tum, Lib'anos, Thus mas'culum, Thus verum,* (F.) *Encens.* By some, however, it is supposed to be the produce of an *Am'yris*, and by others of *Boswel'lia serrat'a*, or *B. thurifera*. The odour of olibanum is peculiar and aromatic; taste bitterish and slightly pungent; partly soluble in alcohol, and forming a milky emulsion, when triturated with water. It was formerly used in dysentery and hæmoptysis, but is now never employed except as a perfume in a sick room.

JUNIPERUS OXYCE'DRUS, (F.) Genévrier Oxyèdre, Cade, Ozicèdre, Oxyèdre, grows in the south of Europe, Siberia, &c. By combustion of the wood, a liquid tar, *O'leum ca'dinum, Junip'eri o'leum empyreuma'ticum, Oil of Cade, (F.) Huile de Cade ou de Genévrier.* is obtained, which is employed externally in various chronic cutaneous and other diseases.

JUNIPERUS SABI'NA, Sabi'na, S. ster'ilis, Savi'

na, *Bruta*, *Cedrus baccifera*, Common or barren *savin*; *Brathu*, *Brathys*, *Bo'rathron*, *Bo'rathron*, (F.) *Sabine*, *Savinier*. The odour of *savin* tops, *Sabi'na* of the pharmacopœias, is strong and disagreeable. Taste hot, acrid, and bitter, depending on an essential oil. Their operation is stimulant, emmenagogue, anthelmintic, and abortive; externally, escharotic. Dose, internally, gr. v to x of the powder. As an escharotic, they are applied in powder or formed into a cerate. The essential oil, *O'leum Sabi'nae* (Ph. U. S.), has the virtues of the *savine*. Dose, two to five drops.

JUNIPERUS SUECICA, *J. communis*.

JUNIPERUS VIRGINIANA, *Red cedar*. This tree is known throughout the United States by the name of *savine*, and is often used for the same purposes. The tops and leaves are officinal in the Pharmacopœia of the United States.

JUNK, Pad.

JUNO LUCINA, *Ilithyia*.

JUNO'S TEARS, *Verbena officinalis*.

JUPITER, Tin.

JUPITER'S BEARD, *Sempervivum tectorum*.

JURIBALI, *Euribali*. A tree in the forests of Pomeronia. Fam. Meliaceæ. Class, Octandria. Nat. Order, Monogynia. The bark is febrifuge, and may be given in powder or infusion, (3j ad aq. bullient. Oij.)

JURISPRUDENCE, MEDICAL, from *jus*, *juris*, 'law,' and *prudentia*, 'knowledge.' *Jurisprudentia Medica*. This word is often used synonymously with *Legal Medicine*. It is at times employed for the embodied laws and regulations that relate to the teaching and practice of medicine.

JURY OF MATRONS. A jury formed of women empanelled under a writ *de ventre inspicendo*, to try the question, whether a woman is with child or not.

JUS, *Zomos*, *Zomid'ium*, *Sorbi't'io*, *Sor'bitum*. Soup, broth, *Jus'culum* or *bouillon*.

JUS ROVINUM. Beef-tea—*j. Coagulatum*. Gelly. *JUS D'HERBES* (F.). The juice of certain vegetables administered as depuratives; as that of fumitory, burdock, water trefoil, &c.

JUS JELATUM. Gelly—*j. de Réglisse*, see *Glycyrrhiza*—*j. de Viande*, see *Bouillon*.

JUSCULUM, see *Jus*.

JUSÉE (F.). An aqueous *maceration* of tan, coarsely powdered, in which the skins of animals have been steeped for a time, in the preparation of leather. An extract has been prepared from this, which has been prescribed in syrup and in pill; each pill containing about a grain and a half of the extract; and two being taken daily in phthisis, and in general debility connected with scrofula and rickets.—Barruel.

JUSQUIAMINE, Hyoscyamus.

JUSQUIAMUS, Hyoscyamus.

JUSQUIAUME, Hyoscyamus.

JUSTICIA BIVALVIS, *Adulasso*—*j. Adhatoda*, *Adhatoda*.

JUSTICIA ECROLIUM, *Carra curini*. A Malabar plant, the root of which, and the leaves, in decoction, are considered in the country to be lithontriptic.

JUSTICIA PECTORALIS. A West India plant, which is slightly astringent. (F.) *Carmentine*, *C. Pectorale*, *Herbe aux Charpentiers*.

JUVANS, pl. *Juvantia*, *Aucill'ium*, *Remed'ium*. A medicine or substance of any kind, which relieves a disorder. An *Adjuvant*.

JUVENIS, Adolescents.

JUVENTA, Adolescence, Hebe.

JUVENTUS, Adolescence, Hebe.

JUVENTUS, Adolescence, Hebe.

JUXTAPOSIT'ION, *Juxtaposit'io*, from *juxta*, 'near to,' and *ponere*, *positum*, 'to place.' The mode of increase proper to minerals; which consists in the successive application of new molecules upon those that form the primitive nucleus. It is opposed to *intussusception*.

K.

N. B.—Most of the terms under *K*, derived from the Greek, are found under the letter *C*.

K. This letter was formerly used to designate a compound of gold.

KAATH, Catechu.

KAAWY. Ancient name of an Indian drink, prepared from maize.

KABALA. Cabal.

KÆMPFERIA GALANGA, see *Maranta galanga*.

KÆMPFERIA ROTUNDA. Called after Kæmpfer, the naturalist. *Zedoaria*, *Z. rotunda*, *Amomum zedoaria*, *Col'chicum Zeylan'icum*, *Cureuma aromatica*, *Zedoary*, (F.) *Zédoaire rond*. Ord. Zingiberaceæ. The roots of this Ceylonese plant have a fragrant smell, and warm, bitterish, aromatic taste. They are in wrinkled, gray, ash-coloured, heavy, firm, short pieces; of a brownish-red colour within; and are stimulant and carminative. Dose, ʒj to ʒj of the powder.

According to some, the round *zedoary* is furnished by *Cureuma zerumbet*; the long, *Zedoaria longa*, by *Cureuma zedoaria*.

KAHINCÆ RADIX, *Caincæ radix*.

KAIB, *Kayl*. A word employed by the alchemists for sour and coagulated milk.

KAJEPUT, Cajeput.

KAKERLAKISM, see *Albino*.

KAKOCHYMIA, *Cacochymia*.

KALI, Potash, Potassa—*k. Acetas*, Potassæ *acetas*—*k. Aeratum*, Potassæ *carbonas*—*k. Bichromicum*, Potassæ *bichromas*—*k. Causticum*, Potassa *fusa*—*k. Causticum cum calce*, Potassa *cum calce*—*k. Chloricum*, Potassæ *urias hyperoxygenatus*—*k. Chromicum acidum*, Potassæ *bichromas*—*k. Chromicum flavum*, Potassæ *bichromas*—*k. Chromicum rubrum*, Potassæ *bichromas*—*k. Chromicum neutrale*, Potassæ *chromas*—*k. Hydriodicum*, see Potassæ *hydriodas*—*k. Inermis*, see *Salsola kali*—*k. Nitricum*, Potassæ *nitras*—*k. Oxalicum acidulum*, Potassa, *oxalate of*—*k. Præparatum*, Potassæ *subcarbonas*—*k. Præparatum e tartaro*, see Potash—*k. Purum*, Potassa *fusa*—*k. Soda*, see *Salsola kali*—*k. Spinosum cochleatum*, *Salsola kali*—*k. Subcarbonas*, Potassæ *subcarbonas*—*k. Sulphas*, Potassæ *sulphas*—*k. Sulphuricum*, Potassæ *sulphas*—*k. Sulphuretum*, Potassæ *sulphuretum*—*k. Tartarizatum*, Potassæ *tartras*—*k. e Tartaro*, see Potash—*k. Vitriolatum*, Potassæ *sulphas*.

KALICUM HYDRAS, Potassa *fusa*.

KALIUM IODATUM, see Potassæ *hydriodas*.

—*k. Iodatum Hydrargyrum, Potassii hydrargyro-iodidum.*

KALMIA ANGUSTIFOLIA. Called after Kalm, the botanist. *Ivy, Narrow-leaved Kalmia* or *Laurel, Dwarf Laurel, Sheep Laurel.* Ord. Ericaceae. This plant has the same virtues as *K. latifolia*. So also have *KALMIA CUNEATA* and *KALMIA HIRSUTA*.

KALMIA, BROAD LEAVED, K. latifolia.

KALMIA GLAUCA, Swamp Laurel, Pale Laurel.

KALMIA LATIFOLIA, Broad-leaved Kalmia, Calico bush, Laurel, Mountain or Rose or Broad-leaved Laurel, Big Ivy, Spoonwood, Lambkill, Sheep-poison. (F.) *Laurier des Montagnes*. This plant kills sheep and other animals. The Indians use it as a poison. The powdered leaves have been applied successfully in tinea capitis; and a decoction of it has been used for the itch. The powder, mixed with lard, has been applied in herpes.

KALMIA, NARROW-LEAVED, K. angustifolia.

KALO, Arum esculentum.

KAMEYLA, see Rottlera tinctoria.

KAMPHUR, Camphor.

KANKROID, Canceroid.

KAPHUR, Camphor.

KARABE, Asphatium.

KARABITUS, Phrenitis.

KARCINOSE, Carcinosis.

KARENA, Carena.

KARKINOMA, Carcinoma.

KARSON-KHAYI, Swietenia Senegalensis.

KASSADER, Convolvulus panduratus.

KASSAUDER, Convolvulus panduratus.

KATASARCA, Anasarca.

KATASTALTICA, Astringents.

KAVA, Ava.

KAVIAC, Caviare.

KAWING, Anhelatio.

KAYL, Kaib.

KECKCORN, Trachea.

KECKER, Trachea.

KEEN-BITE, see Congelation.

KEER, Sorbus aucuparia.

KEITA, Monsonia ovata.

KELIS, see Canceroid.

KEKING, Eruption.

KELOID, Canceroid.

KELOTOMIA, Celotomia.

KELP, Varec. The impure mineral alkali obtained by burning certain marine plants. See Soda.

KENCH, Sprain.

KENNELWORT, Scrophularia nodosa.

KENTUCKY, MINERAL WATERS OF. The *Olympian Springs, Mudlick*, in this State, are near the sources of Licking river. There are three different kinds, in the space of half a mile. One of them is saline, impregnated with sulphur;—another is chalybeate, and a third a sulphureous spring. In various parts of Kentucky, there are saline waters, which are frequented by invalids. The *Salines* at Bigbone, formerly employed in the manufacture of salt, are now resorted to. A spring, near Harrodsburg, in Mercer County, is strongly impregnated with sulphate of magnesia, and is much visited. At the *Blue Licks* on both banks of the Licking river, there are sulphureous saline waters.

KERATIASIS, Ceratiasis.

KERATITIS, Ceratitis.

KERATOCELE, Ceratocele.

KERATODEITIS, Ceratitis.

KERATO-GLOSSUS, Cerato-glossus.

KERATO-IRITIS, Aquo-membranitis.

KERATO-MALACIA, Cerato-malacia.

KERATONYXIS, Ceratonyxis.

KERATO-PHARYNGEUS, Cerato-pharyngus.

KERATOPLASTY, Ceratoplastic.

KERATO-STAPHYLINUS, Cerato-staphylinus.

KERATOTOMUS, Ceratotomus, Knife, cataract.

KERECTOMY, Cerectomy.

KERION, see Favosus.

KERMES, Chermes, Alkerm'es. One of the species of the genus *kermes* lives on a green oak, and is called *Coccus ilicis, Kermes animal, Cocci granum, Coccum, C. buphicium seu infer-torium seu tinctorium seu scarlatinum, &c.* The oak, to which allusion has been made, is known by botanists under the name *Quercus coccifera*, and grows abundantly in the uncultivated lands of southern France, Spain, and in the islands of the Grecian Archipelago. The *kermes* inhabiting it has the appearance of a small, spherical, inanimate shell. Its colour is reddish-brown, and it is covered with a slightly ash-coloured dust. This is the *kermes* of the shops. It is now only used in dyeing; but was formerly reputed to possess aphrodisiac, analeptic, anti-abortive, and other virtues.

KERMES MINERAL, Antimonii sulphuretum precipitatum and A. sulphuretum rubrum.

KERNEL, Tubercle.

KERNELS, WAXING, see Waxing kernels.

KERN-MILK, Buttermilk.

KERUA, Ricinus communis.

KETCHUP, Catchup. A pickle prepared from the liquor of the mushroom, walnut, tomato, &c.

KETMIA ÆGYPTIACA, Hibiscus abelmoschus.

KEY, Sax. cæg, Clavis, Oleis, Clavis Angliæ, (F.) Clef; Olef de Garangeot. An instrument, used for extracting teeth. It consists of a firm handle, with a claw at right angles to it, and moving upon a pivot. This claw embraces the tooth. It has undergone several modifications, and hence various instruments are used under this denomination. The French have the *Clef à pompe, Clef à pivot, and Clef à noir.*

KEYRI CHEIRI, Cheiranthus cheiri.

KHALA MIMUC, Bit noben.

KHAYA SENEGALENSIS, Swietenia Senegalensis.

KHUS-KHUS, Andropogon muricatus.

KIAS'TER, Chiaster, from χιαστω, 'to cross.' A species of bandage, having the form of the Greek letter χ , which the ancients used for approximating, and maintaining in contact, the fragments of the patella, in cases of fracture of that bone. It was applied in the form of the figure 8.

KIBE, Chilblain.

KIBISTITOME, Cystitome.

KIDNEY. Its etymology is uncertain. Sere-nius derives it fancifully from *Su. Goth. qued*, the belly; and *nigh*, (quasi, *quidney*.) *Ren, Nephros, Protme'sia, (Sc.) Neir, Nere, (F.) Rein.* The kidneys or reins are the secretory organs of the urine. They are two glands, situate deeply,—the one on the right, and the other on the left side—in the hypochondres: at the sides of the lumbar vertebræ; behind the peritoneum; and in the midst of an abundant, fatty areolar tissue, *Tunica seu Capsula adiposa*. The kidney is of a reddish-brown colour; oval form; and flattened on two surfaces. It has, at its internal margin, a deep fissure, by which the renal vessels and nerves enter or quit the organ, and the ureter issues. It resembles, pretty accurately, the haricot or kidney-bean. Two substances are readily distinguishable in it;—the outer, *secreting, cortical, glandular or vascular, Substantia corticalis seu glandulosa*, which secretes the urine; and the inner, *tubular, medullary, uriferous, conical or radiated, Substantia Medullaris seu Tubulosa seu Fibrosa*, which appears

under the form of small cones or unequal papillæ or mamillæ, each resulting from the union of small capillary tubes, adherent by one of their extremities to the cortical substance; and opening, by the other, at the summit of the cone, into calices, a species of membranous tubes, more or less numerous, which transmit the urine of the papillæ to the pelvis. By the pelvis is meant a small, membranous sac, of an irregularly oval shape, at the base of which are the orifices of the calices, and the other extremity of which is continuous with the ureter. The kidney is surrounded by a fibrous membrane proper to it, *Perinephrus*, *Tunica propria* seu *abuginæa*. It has been shown by Mr. Bowman and others that the renal artery is distributed to the corpora Malpighiana, forming a pellet of convolutions, which is received into a flask-like dilatation of the ureter—*Bowman's Capsule*, and through which—it is conceived—the watery portion of the urine is separated. The blood then becomes venous, and is distributed by different veins—*portal veins* of the kidney—to the convoluted tubes through which the proper urine is secreted. Hence the blood passes into the renal vein. The intermediate vessels between the Malpighian bodies and the convoluted tubes have been termed the *Portal System of the Kidney*.

KIDNEY, BRIGHT'S DISEASE OF THE, *Morbus Brightii* seu *albuminæsis* seu *granulæus Renum*, *Nephritis albuminosa* seu *sociata*, *Nephritis*, *Nephrocalasosis*, *Nephrococcosis*, *Urozeimia albuminosa*, *Cachectic* or *Associated nephritis*, *Inflammation of the Malpighian corpuscles*, *Disease of Bright*, *Granular Degeneration or Disorganization of the Kidney*, *Granular Kidney of Bright*, *Cachectic nephritis*, *Chalazonephritis*, (F.) *Maladie de Bright*, *Néphrite albumineuse*, *État Granuleux ou Dégénérescence granulee du Rein*, *Albuminurorrhée*. A granular disease of the cortical part of the kidney, which gives occasion to the secretion of urine that contains albumen, and is of less specific gravity than natural, and which destroys by inducing other diseases. It was first described by Dr. Bright, of London.

KIDNEY, GRANULAR, OF BRIGHT, Kidney, Bright's disease of the.

KIESTEINE, *Ki'estein*, properly *Ky'esteine*, *Ky'estein*, *Kystein*, *Cy'esteine* or *Cy'estein*, *Kiesthein*,—(Lehmann), from *κειν*, 'to be pregnant,' and *εθος*, 'a garment or pellicle.' A peculiar albuminoid pellicle, which forms on the urine of a pregnant female when allowed to stand for a few days. It is whitish, opalescent, slightly granular, and may be compared to the fatty substance that swims on the surface of soups, after they have been allowed to cool. When taken in conjunction with other phenomena, it is a valuable aid in the diagnosis of early pregnancy.

KIESTINIC, *Kiestinicus*; same etymon. Relating or appertaining to kiesteine; as '*kiestinic urine*.'

KILBURN, MINERAL WATERS OF. These springs contain carbonic acid, sulphohydric acid; carbonates of lime, magnesia, and iron; sulphates of soda, lime, and magnesia, and chloride of sodium.

KILE, Ulcer.

KILL LAMB, *Andromeda mariana*.

KILOGRAMME, *Chiliogramma*, from *χιλοι*, 'a thousand,' and *γραμμα*, 'a gramme.' The weight of a thousand grammes;—two pounds, eight ounces, one drachm, and twenty-four grains, Troy.

KILOLITRE, from *χιλοι*, 'a thousand,' and *λιτρον*, 'a litre.' A measure containing a thousand litres.

KINA KINA, *Cinchona*.

KINANCIE, *Cynanche*.

KINCOUGH, *Pertussis*.

KINDCOUGH, *Pertussis*.

KINESIP'ATHY, *Kinesitherapia*, *Cinesip'athy*, *Movement cure*, *Motor'pathy*, *Medicinal mechanica*, from *κινω*, 'motion,' and *παθω*, 'disease.' A mode of treating disease by gymnastics or appropriate movements proposed by Ling, a Swede, and hence called *Lingism*.

KINETIC, *Motory*.

KINGCURE, *Pyrola maculata*.

KING'S EVIL, *Scrofula*.

KINGSHOOD, *Reticulum*.

KINICI ACETAS, *Quinia*, acetate of.

KININUM, *Quinine*.

KINIUM, *Quinine*.

KINK, *Quinte*.

KINK IN THE HEAD, see *Insanity*.

KINKHAUST, *Pertussis*.

KINKHOST, *Pertussis*.

KINKINA, *Cinchona*.

KINO, *Gummi Gambien'se* seu *rubrum adstringens Gambien'se* seu *adstringens Fothergilli*, *African*, *East India* or *Amboy'na kino*, (F.) *Goman*, *suc* ou *extrait de kino*. The trees, whence one variety of this resin is obtained, are not botanically ascertained. The London college ascribe it to *Pterocarpus erinacea*; the Edinburgh to *Eucalyptus resinifera* seu *gummifera*, *Iron bark*; and the Dublin to *Butea frondosa*. The Pharmacopœia of the United States, (1842 and 1851), defines it to be "an extract obtained from an uncertain plant;" that of 1851 states it to be the inspissated juice of *Pterocarpus marsupium* (De Candolle) and of other plants. On wounding the bark, the kino flows drop by drop. A West India variety is said to be derived from *Coccoloba wuifera* or *Sea-side Grape*; and a *Botany Bay kino* is said to be the concrete juice of *Eucalyptus resinifera* or *brown gum-tree* of New Holland. *Sez. Syst. Icosandria Monogynia. Nat. Ord. Myrtacæ*. Kino consists chiefly of tannic and gallic acids, oxide of iron, and colouring matter. It is inodorous; the taste a sweetish bitter; and it is sometimes gritty between the teeth. It comes to us in fragments of a dark ruby red colour, and is easily pulverized. Its properties are powerfully astringent. Dose, gr. x to gr. xx in powder.

KIONORRHAPHIA, *Staphyloraphy*.

KI'OTOME, *Ki'otomus*, from *κτω*, 'a pillar,' 'support,' and *τεμνω*, 'to cut.' (F.) *Coupebride*. An instrument invented by Desault to cut any accidental brides or filaments in the rectum and bladder; and which he afterwards used for the removal of the tonsils. It is composed of a flat, silver sheath, open at one edge. This sheath is provided with a cutting blade, which can be forced through the opening, and thus all the parts can be divided with which it comes in contact.

KIPPERNUT, *Bunium bulbocastanum*.

KIRATE. A weight of four grains, according to Blancard.

KIRIAGHUMA, *Gymneura lactiferum*.

KIRKLAND'S NEUTRAL CERATE, see *Cerate*, *Kirkland's Neutral*.

KIRNMILK, *Buttermilk*.

KIRRHONOSIS, *Cirrhosis*.

KIRRHOSIS, *Cirrhosis*.

KIRSCHWASSER, (G.) *Kirsch* and *Wasser*, 'Cherry water.' An alcoholic liquor, obtained from cherries bruised with their stones, by subjecting them to distillation, after having caused them to ferment. See *Spirit*.

KISS'INGEN, MINERAL WATERS OF. These Bavarian springs have been long frequented. There are three,—two chalybeate, and one alkaline and acidulous.

KIST. A weight of 14 grains.—*Paracelsus*.

KITILLING, Tickling.

KITTLING, Tickling.

KLAPROTHII SULPHAS, Cadmii sulphas.

KLAPROTHIUM SULPHURICUM, see Cadmii sulphas.

KLEPTOMANIA, *Cleptoma'nia*, *Klopema'nia*, *Cleptomania*; from κλέπτω, 'I steal,' and mania. Insanity, with an irresistible propensity to steal.

KLINOCEPHALIA, Clinoccephalia.

KLIPDAS, see Myrax Capensis.

KLOPEMANIA, Kleptomania.

KNAP-KNEED, see Entogonyancon.

KNARESBOROUGH, MINERAL WATERS OF, see Harrogate, mineral waters of.

KNEADING, Shampooing.

KNEE, Genu.

KNEE, HOUSEMAID'S. An inflammation of the bursa, which in most individuals is in front of the patella, and is apt to inflame and enlarge from effusion in those in whom it is subjected to much pressure. Hence its name. It is a form of capsular rheumatism.

KNEE-HOLLY, Ruscus.

KNEE-JOINT, Genu.

KNEE SCAB, Crusta genu equinæ—k. Encircled hoof. Crusta genu equinæ.

KNIFE, Sax. cnif. Swed. Knif. Dan. Kniv. (F.) Canif. *Culter*, *Smilæ*, *Cutell'us*, *Tome'ion*, *Tome'us*, *Macha'ra*, *Macha'rium*, *Macha'rie*, (F.) *Couteau*. A cutting instrument, used in surgery to divide the soft parts, and which only differs from the bistouri or scalpel in being usually larger.

The most common knives are the following:

KNIFE, AMPUTATION, (F.) *Couteau à amputation*, *C. droit*, *Culter rectus*. This is the largest of the knives used in surgery. Formerly, they were curved; now they are straight, and provided with one or two edges.

KNIFE, CATARACT, *Cerat'omus*, *Kerat'omus*, (F.) *Couteau à cataracte*. The cataract knives of Richter, Wenzel, Ward, and others, being intended to perform the section of the transparent cornea, are so shaped as to exactly fill the small wound made by them; and thus to prevent the discharge of the aqueous humour, until the section is completed. The blade of the knives of Wenzel and Ward resembles a very narrow lancet, blunt in the posterior five-sixths of one of its edges. The blade of that of Richter is pyramidal, cutting through the whole length of its inferior edge, and also blunt in the five-sixths of the upper.

KNIFE, CHES'ELDEN'S. A knife with a fixed handle; very convex on its edge, concave on the back, which was used by Cheselden in *lithotomy*.

KNIFE, CROOKED, *Culter falcatus*, *Culter curvus*, (F.) *Couteau courbe*. A knife, which is crooked and concave on its cutting edge. It was formerly employed in amputation of the limbs.

KNIFE, CROTCHET, *Cutell'us uncus*, (F.) *Couteau à crochet*. A steel instrument, composed of a round staff, furnished with a handle at one extremity, and at the other with a curved knife. It was formerly used to cut to pieces monstrous fetuses in utero, and to open the head when necessary.

KNIFE, DOUBLE-EDGED, *Anceps cutell'us*, *Amphis'mela*, *Amphis'milæ*, *Catling*, (F.) *Couteau à deux tranchans ou désarticulateur ou interosseux*. A knife, the blade of which is straight and sharp on both sides. It is used for disarticulating bones; and for cutting the soft parts situate between the bones, in amputation of the leg and forearm.

KNIFE EN SERPETTE, (F.) *Couteau en serpette*. A sort of knife, of the shape of a *serpette* or pruning-knife, invented by Desault for

dividing the bony paries of the maxillary sinus, for the purpose of extracting fungi from it.

KNIFE, LENTICULAR, *Culter lenticularis*, (F.) *Couteau lenticulaire*. An instrument, used in the operation of trepanning, for removing inequalities in the inner table of the skull, which may have been left by the crown of the trephine around the opening made by it. It is formed of a lenticular button, fixed at the extremity of an iron staff, which is convex on one side, flat on the other; sharp at both edges, and mounted on an ebony handle.

KNIFE, LITHOTOMY, (F.) *Couteau lithotome*. A name, given by Foubert to a large knife, the narrow blade of which, four and a half inches in length, was sharp in its whole extent, and made an obtuse angle with the handle. He used it in the lateral operation.

KNIFE, POLYPUS, Polypotome.

KNIFE, ROOT-CUTTING. In Pharmacy, a knife moving on a joint at its pointed extremity, by which roots and other ligneous matters are divided in pharmaceutical processes.

KNIFE'SMAN, see Surgery, salvation.

KNIT-BACK, Symphytum.

KNOBWEED, Collinsonia.

KNOCK-KNEED, Entogonyancon.

KNOT, Tubercle.

KNOT, PACKER'S, (F.) *Nœud d'emballleur*. A compressive bandage, used for arresting hemorrhage from the temporal artery or its branches. The *nœud d'emballleur* is made with a double-headed roller, five ells long. A graduated compress is placed over the opening in the artery, and the bandage is applied over it; the balls of the roller being carried horizontally round to the opposite temple, where they are crossed obliquely and carried back to the part where the compress is situate. The hands are then changed with the rollers, crossing them so as to form a knot, and taking one above the head; the other beneath the chin. They are then crossed again, so as to form several knots, one above the other. This bandage is called, by some surgeons, *solar* or *oblique chevestre* or *capisternum*.

KNOT, SURGEON'S, *Nodus chirurgicus*, (F.) *Nœud du chirurgien*. A double knot made by passing the thread twice through the same noose. This knot is used frequently in the ligature of arteries, the umbilical cord, &c.

KNOTBERRIES, *Rubus chamæmorus*.

KNOTGRASS, *Polygonum aviculare*.

KNOTROOT, *Collinsonia Canadensis*.

KNOTWEED, *Collinsonia Canadensis*, *Polygonum aviculare*.

KNOWLTONIA VESICATORIA, *Nat. Order*, Ranunculaceæ. An acrid plant of South Africa, used by the Cape colonists as a blister in rheumatism.

KOINO-MIASMATA, see Miasm.

KOKUM, *Garcinia purpurea*—k. Butter, see *Garcinia purpurea*.

KOLERUS, a name given by Paracelsus to a dry ulcer.

KOLPORRAPHY, Elytrorrhaphy.

KOLTO, Plica.

KOOCHLA TREE, *Strychnos nux vomica*.

KOOSSO, *Hagenia Abyssinica*.

KOPYOPIA, see Copyopia.

KORIS, Cimex.

KOSSO, *Hagenia Abyssinica*.

KOUMIS, Kumyss.

KOUSSO, *Hagenia Abyssinica*.

KRAMERIA, *Krameria trian'dra*, *Crame'ria*, *Ratan'hia*, *Rhatan'ia*, *R'at'ahy*, (F.) *Ratan'hie*, —*Sec. Syst.* Tetrandria Monogynia, *Nat. Ord.* Polygalææ. — is a native of Java. The root, *Krame'ria* (Ph. U. S.), has a bitter taste; and is

astringent, diuretic, and detergent. Dose, \mathfrak{Dj} to \mathfrak{ss} .

KRAMERIA IXT'NA or *Ratanhy* of the Antilles has similar virtues.

KREASOTON, Creasote.

KREATIC NAUSEA, see Nausea.

KREATINE, Creatine.

KREATININE, see Creatine.

KREATOPHAGOUS, Carnivorous.

KREOSOTON, Creasote.

KRETINISMUS, Kretinism.

KRIEBELKRANKHEIT, Convulsio cerealis.

KRONE, Crown.

KRYSTALLINE, see Crystalline.

KUA KAH, *Curcuma longa*.

KUMYSS, *Koumyss*, *Koumie*. A beverage used in families by the people of Yakutz. It resembles sour buttermilk, without being greasy. According to Sir George Simpson, it is prepared in a very simple way from mare's milk, which is merely allowed to stand for some days in a leather churn till it becomes sour. It is then bottled for use. This drink is rather nutritious than exhilarating; but from the same material the Burats and the Kirghez prepare an intoxicating spirit in which they indulge to excess.

KUSSAUER, *Convolvulus panduratus*.

KUTKULEJA, *Cæsalpinia bonducella*.

KUTOOKURUNJA, *Cæsalpinia bonducella*.

KUTUBUTH. An Arabic name for a species of melancholy in which the patient is never quiet at any one place, but wanders about here and there, see Cuttubuth. Also, the name of an insect, which lives at the surface of stagnant waters, and is in a constant state of agitation. Some lexicographers imagine that it is on account of this last circumstance, that the name of the insect has been given to the disease.

KWOSO, *Hagenia Abyssinica*.

KYANOSIS, Cyanopathy.

KYAPUTTY, Caieput.

KYESTEINE, Kiesteine.

KYESTINIC, Kiestinic.

KYLLOPODIA, Kyllosis

KYLLO'SIS, from *κυλλος*, 'crooked,' 'lame.' *Cyllo'sis*, *Kyllopod'ia*. Professor Chaussier so calls congenital distortion of the feet, *Clubfoot*, *Poltfoot*, *Stumpfoot*, (Prov.) *Pummel-foot*, *Tal'ipes*, (F.) *Pied bot*. Of this there are many varieties. In one, the foot, instead of resting on the soil, by the whole plantar surface, touches it only with the metatarso-phalangan articulations. It seems as if turned backwards and broken upon the leg, (*Pes seu Tal'ipes Equinus*, *Hip'popus*, *Oxy'pod'ia*, *Tip-foot*, *Horsefoot*.) In other cases the foot is twisted inwards, (*Varus*, *Tal'ipes varus*, *Blasop'odes*, *Bl'a'opus*, *Cyllo'pus*, *Cyllop'oda*, *Cyllopod'ion*, *Lo'ripes*, *Crossfoot*, (F.) *Cagneux*,) so that it rests only on the ground on its

outer edge; or it may be twisted outwards, (*Valgus*, *Tal'ipes Valgus*, *Out-bow-foot*,) or rest only on its inner edge, or on the back part of the heel, *Talipes calcaneus*, *Heel clubfoot*. In the *flatfoot* or *splayfoot*, *Duck-foot*, *Splat-foot*, *Sar'apus*, *Plat'ypus*, *Platypod'ia*, (F.) *Pied plat*, the plantar surface of the foot is flattened instead of being concave.

These deformities are rarely accidental. They are almost always congenital, and may be rectified, at an early period, by proper mechanical means to strengthen the foot gradually and restore it to its proper shape and direction; and if these means fail, the tendons and muscles concerned in the deformity may be divided.

KYMOGRAPH'ION, *Cymograph'ion*; from *κύμα*, 'wave,' and *γράφω*, 'I describe.' A wave describer or measurer.' An instrument invented by Ludwig, which is self-registering, and exhibits the relation between the waves of the pulse and the undulations produced by respiration.

KYNA, *Pastinaca opoponax*.

KYPHOSIS, see Cyphosis—k. *Inflammatoria*, *Vertebral disease*—k. *Paralytica*, *Vertebral disease*.

KYST, *Cyst*, *Kystie*, *Cystis*, from *κυστις*, 'a bladder,' 'pouch.' (F.) *Kyste*. This term is generally applied to a pouch or sac, — *Cysto'ma*, — without opening, and commonly of a membranous nature, which is accidentally developed in one of the natural cavities, or in the substance of organs. Many theories have been successively emitted to explain the formation of cysts, but none are entirely satisfactory. Some are formed by a thin, translucent membrane, having scarcely the thickness of the arachnoid; others of a whitish, fibro-cellular membrane, more or less thick. Some contain cartilaginous or bony flakes. The greater part have but one cavity; others, on the contrary, have several, separated by complete or imperfect septa, as is frequently seen in those developed in the ovaries. The matter contained in cysts is sometimes limpid, serous, yellowish white, reddish, and, at others, more or less thick, albuminous, adipous, or caseous. The tumour, formed by them, is called Encysted.

KYSTE, Kyst — k. *Anévrismal*, *Aneurismal sac*.

KYSTEIN, Kiesteine.

KYSTH'ITIS, *Cyathi'tis*, from *κυσθός*, 'the vagina.' Inflammation of the vulva and of the mucous membrane of the vagina.

KYSTITOME, *Cystitome*.

KYSTHOPTO'SIS, from *κυσθός*, 'the vagina,' and *πτωω*, 'I fall.' Prolapsus or inversion of the vagina.

KYTE, Abdomen, Stomach.

KYTTARRHAG'IA, from *κυσταριον*, 'an alveolus,' and *ρῥαγω*, or *ρρηγνυμι*, 'I break forth.' Discharge of blood from an alveolus.

L.

LABARIUM, from *labi*, 'to fall.' Looseness of the teeth.

LABDACISMUS, Lallation.

LABDAMEN, *Cistus creticus*.

LABDANUM, see *Cistus creticus*.

LABE, from *λαβω*, *λαμβάνω*, 'I seize,' 'I take,' 'the act of grasping.' Invasion. Also, em-

ployed to denote the first paroxysm of fevers. — Galen, Hippocrates.

LABEO, Chilon.

LABES, Chilon, Macula.

LABIA CUNNI, Lips of the vulva—l. *Interna* seu *Minora*, *Nymphæ*—l. *Majora*, Lips of the vulva—l. *Pubendi*, Lips of the vulva—l. *Pubendi minora*, *Nymphæ*.

LABIAL. *Labialis*, from *labium*, 'a lip.'

LABIAL. Orbicularis oris.

LABIAL ARTERY. Haller and Sabatier call thus the *facial artery* of the majority of anatomists. The *labial arteries*, properly so called, *coronary arteries of the lips*, (F.) *Coronaires des lèvres*, are two in number. The *superior* arises from the facial, above, and very close to, the commissure of the lips. It is large and tortuous, and is distributed to the upper lip. The *lower* arises from the facial, at a considerable distance from the commissure, and proceeds, in a serpentine course, into the substance of the lower lip, to which it is distributed.

LABIAL GLANDS. This name is given to a multitude of muciparous follicles, of some size, round, prominent, and separate from each other, which are found on the inner surface of the lips, below the mucous membrane.

LABIAL VEINS are distinguished, like the arteries, into *superior* and *inferior*. They open into the facial vein;—a division of the internal jugular.

LABIALIS. Orbicularis oris.

LABIDOMETER. Labimeter.

LABIMETER. (F.) *Labimètre* ou *Labidomètre*, from *λαβίς*, *labidos*, 'forceps,' and *μετρον*, 'measure.' A scale adapted to the handles of the forceps, which indicates the distance of the blades from each other, when applied to the head of the child in *utero*.

LABIS. Forceps.

LABIUM. Lip.—1. *Leporinum*, Harelip.—1. *Uteri*, Amphideum.

LABLAB. *Dolichos lablab*.

LABORANS. Sick.

LABORATOIRE. Laboratory.

LABORATORIUM. Laboratory.—1. *Chymicum* seu *pharmaceuticum*, see *Pharmacopœia*.

LABORATORY. *Laboratorium*, *Ergasterion*, (F.) *Laboratoire*, from *laborare*, *laboratum*, 'to work.' A work-shop. A place for preparing chemical or pharmaceutical products, &c.

LABORIOSUS. Sick.

LABORIOUS. Delivery is said to be *laborious*, *Partus laboriosus*, *Mogostocia*, (F.) *Accouchement laborieux*, when attended with more difficulty and suffering than usual. With some, *laborious labour* means one that requires the use of instruments. See *Dystocia*.

LABOUR. Parturition.

LABOUR CHAIR. *Obstetric chair*. A chair, in which a parturient woman is placed during delivery.

LABOUR, COMPLICATED, see *Parturition*—1. Difficult, *Dystocia*—1. Dry, *Partus siccus*—1. Forced, see *Parturition*—1. Impracticable, see *Parturition*—1. Morbid, *Dystocia*.

LABOUR PAINS, see *Pains*.

LABOUR, POWERLESS, see *Parturition*—1. Premature, *Parturition*, (premature)—1. *Preternatural*, *Metatocia*, see *Parturition*.

LABOUR SHOW, see *Parturition*.

LABRISULCIUM. Cheiloeace, Stomaceae.

LABRUM. Lip.—1. *Cerebri*, *Infundibulum* of the brain.

LABRUSCA. *Bryonia alba*.

LABYRINTH. *Labyrinthus*, *Antrum buccinum*. A place, full of turnings, the exit of which is not easily discoverable. Anatomists have given this name to the aggregate of parts, constituting the internal ear, *Labyrinthus auris internæ*, *Intima pars organi auditûs*, *Labyrinthic cavity of the ear*. The Labyrinth is situate between the tympanum and meatus auditorius internus. It is composed of several cavities, which communicate with each other in the dried bone; as the *vestibule*, *cochlea*, *semicircular canals*, &c. It is lined by periotestum, and also by a pulpy membrane, constituting the membra-

nous *labyrinth*, on which the auditory nerve is regularly dispersed. This membrane forms two sacks in the vestibule, called *sacculus vestibuli* and *sacculus*, respectively, which resemble in shape that of the bony cavities containing them. Each sac contains calcareous matter, constituting the *Otoliths* and *Otoconies*. When the sac is laid open, upon the upper and outer part, a partition appears, partaking of the nature of the sac, and called by Meckel, *Septum vestibuli nervi semembranaceum*.

Labyrinth is also applied to the lateral portions of the ethmoid bone, which are formed of large irregular cells.

LABYRINTH, MEMBRANOUS, see *Labyrinth*.

LABYRINTHIC. *Labyrinthicus*, (F.) *Labyrinthique*. Relating or appertaining to the labyrinth.

LABYRINTHIC CAVITY OF THE EAR. *Labyrinth*.

LABYRINTHUS. *Labyrinth*—1. *Auris Intima*, see *Labyrinth*.

LAC. *Lacca*, *Laqueus*, Milk—1. *Ammoniacy*, *Mistura ammoniaci*—1. *Amygdales*, *Emulsio amygdalæ*—1. *Asafoetida*, *Mistura asafoetida*—1. *Avis*, *Albumen ovi*—1. *Guaiacy*, *Mistura Guaiacy*.

LAC, GUM. *Lacca*.

LAC LUNE. *Marga candida*—1. *Maris*, *Sperm*—1. *Pressum*, *Curd*—1. *Primum Puerperæ*, *Colostrum*.

LAC SEED. *Lacca*—1. *Shell*, *Lacca*—1. *Stick*, *Lacca*.

LAC SULPHURIS. *Sulphur præcipitatum*—1. *Terra*, *Magnesia carbonas*—1. *Virginia*, *Virgin's milk*.

LACAPHTHON. Supposed to be the bark of some fir or other tree. It entered into the composition of the *Cyphi*, used in fumigations—Paulus of Ægina. Supposed to be the *Narcaphthum* or *Nascaphthum*.—Ruellius.

LACCA. from *lakah*, Arab. *Gummi lacce*, *Stick*, *Gum*, *Seed*, or *Shell*, *lac*, (F.) *Laque*, *Gomme Laque*. Lac is a substance formed by an insect, and deposited on different species of trees, chiefly in the East Indies, for example, on *Croton lacciferum*, and two species of *Ficus*,—*Ficus religiosa*, and *F. Indica*. The various kinds, distinguished in commerce, are *stick-lac*, which is the substance in its natural state, investing the small twigs of the tree; and *seed-lac*, which is the same broken off. When melted, it is called *shell-lac*.

Lac was, at one time, used in the form of tincture, as a tonic and astringent; and it still forms part of particular dentifrices.

LACERATION. *Laceratio*, *Rhegê*, *Rhegmus*, *Rhegma*, *Ruptura*, *Rupture*, from *lacerare*, *laceratum*, 'to tear'—*Dilaceratio*, *Sparagma*, (F.) *Arrachement*, *Déchirement*, *Dilacération*, *Broiement*. The act of tearing or rending. The breach made by tearing or rending; as a *lacerated wound*, *Tresis vulnus laceratum*, *Laceratum*, *Vulnus laceratum*.

LACERATURA. see *Laceration*.

LACERTA. *Lizard*—1. *Aquatica*, see *Ectozaa*.

LACERTI CORDIS. *Columnæ carneæ*—1. *Musculorum*, see *Muscular fibre*.

LACERTOSUS. *Muscular*.

LACERTULI CORDIS. *Columnæ carneæ*.

LACERTUS. *Brachium*, *Muscle*, see *Muscular fibre*.

LACERUM. Same etymon as *Laceration*. Any thing torn, or appearing as if torn.

LACERUM FORAMEN ANTERIUS. (F.) *Trou déchiré antérieur*, *Hiatus sphæno-pétreux*, (Ch.) is an irregular opening, formed by the sphenoid, occipital and petrous portion of the temporal bone. It is closed by cartilage, and forms a kind of fontanelle, situated at the termination of the petro-occipital suture.

LACERUM FORAMEN POSTERIUS. *Foramen jugulârê*, *F. lacerum* in *Basi Cranii*, (F.) *Trou*

Osseus postérieur, Hiatus occipito-pétrous, (Ch.) is formed by the occipital bone, and the inferior edge of the petrous portion of the temporal bone. Through it, the internal jugular vein, the eighth pair of nerves, and accessory nerve pass out of the cranium.

LACERUM FORAMEN SUPERIUS, Sphenoidal fissure.

LACH'ANON, from *λαχανον*, 'I dig.' A pot herb; *Olus*, pl. *Olera*, a vegetable used as food.

LACHESIS PICTA, see Arrow-poison.

LACHESIS RHOMBEA'TA, *Flagmon*. A poisonous serpent common in the lower forests of Peru.

LACHNANTH'ES TINCTO'RIA, *Dila'tris tincto'ria*, *Redroot*; indigenous. Ord. *Hæmodoraceæ*, (from *λαχν*, 'wool,' and *ανθος*, 'a flower,' owing to its woolly flowers.) The root is astringent and tonic.

LACHRYMA, Tear; see *Vitis vinifera*.

LACHRYMAL, *Lacryma'lis*, from *lacryma*, 'a tear.' Belonging to the tears. This epithet is given to various parts.

LACHRYMAL APPARATUS, see *Lachrymal Passages*.

LACHRYMAL ARTERY proceeds from the ophthalmic; and distributes its principal branches to the lachrymal gland.

LACHRYMAL CANAL OR DUCT, *Nasal Canal* or *duct*, *Canalis lacryma'lis* seu *orbitæ nasalis* seu *Sacci lacryma'lis*, *Ductus ad Nasum* seu *Nasalis orbitæ*, is formed by the superior maxillary bone, os unguis, and os turbinatum inferius; and is seated in the outer paries of the nasal fossæ. It is lined by a prolongation of the mucous membrane of the lachrymal sac; and its inferior orifice is furnished with a valvular duplicature. This duct transmits the tears, which have been absorbed at the great angle of the eye by the puncta lacrymalia, into the nasal fossæ.

LACHRYMAL CARUNCLE, see *Caruncula lacrymalis*.

LACHRYMAL DUCT, *Lachrymal Canal*.

LACHRYMAL FOSSA, *Fossa* seu *Fovea lacryma'lis*, is a slight depression at the upper part of the orbit, which lodges the lachrymal gland.

LACHRYMAL GLAND, *Glandula lacryma'lis* seu *innomina'ta Gale'ni*, is seated in a depression of the frontal bone at the upper, anterior, and outer part of the orbit. It is of about the size of an almond; and of an oval shape, flattened above and below:—its great diameter being the antero-posterior. It is composed of several small lobules, united by areolar tissue, and separated by it as well as by vessels and nerves which creep in the intervals. This gland has seven or eight excretory ducts, which open behind the upper eyelid. Its use is to secrete the tears, and pour them on the globe of the eye by the excretory ducts.

LACHRYMAL GROOVE, (F.) *Gouttière lacrymale*, is the bony channel, which lodges the lachrymal sac. It is seated at the anterior and inner part of the orbit, and is formed by the os unguis and the ascending process of the upper jaw bone.

LACHRYMAL HERNIA, *Lachrymal Tumour*, *Hernia Sacci Lacryma'lis*, is when the tears enter the puncta, but cannot pass to the nose, and accumulate. By Anel, this was called *Dropsy of the Lacrymal sac*.

LACHRYMAL NERVE is the smallest of the three branches formed by the ophthalmic nerve. It is distributed, particularly, to the lachrymal gland and to the upper eyelid. In its course it gives off a *spheno-maxillary* and a *malar* filament.

LACHRYMAL PAPILLA, see *Lachrymal Puncta*.

LACHRYMAL PASSAGES, *Via lacryma'les*, (F.) *Voies lacrymales*. The organs concerned in the secretion of tears, in spreading them over the eye, and taking them up again to transmit them into the nasal fossæ. The lachrymal passages

are composed of the lachrymal gland, *caruncle*; *puncta*, *ducts*, *lachrymal sac*, and *nasal duct*, which, together, form the *lachrymal apparatus*, (F.) *Appareil lacrymal*.

LACHRYMAL PUNCTA, *Puncta Lacryma'lia*, *Spiram'ina Palpebra'rum*, (F.) *Points lacrymaux*, are two small, round, and contractile openings, situate in the centre of a tubercle or papilla, *Papilla lacryma'lis*, *Tuberculum lacryma'le*, about a line and a half distant from the inner commissure of the eyelids, and continuous with the lachrymal ducts. These ducts, *Lacryma'les Canaliculi*, *Canales seu Ductus lacryma'les*, *Ductus lacrymales laterales* seu *puncto'rum lacryma'tium*, *Cor'nea lacryma'lia* seu *limacum*, *Collic'ia puncto'rum lacryma'tium*, *canaliculi limacum*, (F.) *Conduits lacrymaux*, are two in number—a superior and an inferior— which extend from the puncta to the lachrymal sac. They seem formed by a very delicate prolongation of the conjunctiva, which is continuous with the mucous membrane of the lachrymal sac, (F.) *Reservoir des larmes*:—the *Saccus* seu *Sinus* seu *Lacus lacryma'lis*, *Dacryocystis*, *Infundibulum lacryma'le*.

LACHRYMAL TUBERCLE, see *Lachrymal puncta*.

LACHRYMAL VEINS accompany the artery of the same name, and open into the ophthalmic and palpebral veins.

LACHRYMATION, *Lacrymatio*—1. Sanguineous, *Dacryhæmorrhysis*.

LACYNIA, 'a hem or fringe.'

LACINLÆ TUBARUM FALLOPII, see *Tube*, *Fallopian*.

LACIS, *Plexus*.

LACMUS, *Lichen roccella*.

LACMUS TINCTORIUS, *Lichen roccella*.

LACONICUM, *Vaporarium*, see *Stove*.

LACQ, *Laqueus*.

LACRIMA, Tear.

LACRUMA, Tear.

LACRYMA, Tear.

LACRYMALIS, *Lachrymal*.

LACRYMA'TIO, *Lachryma'tion*, *Epiphora*. Also, a profuse secretion of tears from any cause:—weeping, *Fletus*, *Plora'tio*, *Plora'tus*.

LACRYMINE, see *Tears*.

LACTALBUMEN, *Casein*.

LACTANS, *Nurse*.

LACTA'RIUM, *Lacticin'ium*; from *lac*, 'milk.' Food prepared from milk. Milk diet or food.

LACTATE DE FER, *Ferri lactas*—*l. de Quinine*, *Quinæ lactas*.

LACTA'TION, *Lacta'tio*, *Thela'sis*, *Thela'smus*, from *lacto*, *lactatum*, (*lac*, 'milk,') 'I suckle,' 'I give milk.' *Suckling*:—the giving of suck, (F.) *Allaitement*. The French make four varieties of lactation. 1. *Allaitement maternal*—*Maternal Lactation*, when the mother suckles the child. 2. *A. étranger mercenaire*,—when another suckles it. 3. *A. artificiel*, when the child is brought up by hand. 4. *A. animal*, when the child is suckled by an animal.

LACTÉ, *Lactic*.

LACTEAL, *Lactic*—1. *Sac*, see *Lactiferous vessels*.

LACTEALS, *Chyliferous vessels*.

LACTEIN, *Milk*, solidified.

LACTENS, *Sucking child*.

LACTES, *Mesentery*, *Pancreas*.

LACTEUS, *Lactic*.

LACTIC, *Lac'teus*, *Galac'ticus*, *Galac'tinus*, *lacteal*, *milky*, (F.) *Lacté*, *Laiteux*, from *lac*, *lactis*, 'milk.' Appertaining to, or resembling milk.

LACTIC ACID, *Ac'idum Lac'teum* seu *Lactis* seu *Lact'ium*, *Acid of milk*, *Galac'tic*, *Nan'ceic* or *Ca'seic Acid*, (F.) *Acide Lactique*. This has been recommended as a therapeutical agent in atonic dyspepsia, owing to its being presumed to be one

of the gastric acids secreted in health. It is given either in the form of lemonade or of lozenges. The acid is obtained either from milk or from the juice of the red beet.

LACTICA, Typhoid.

LACTICANS, Nurse.

LACTICINIA, Parotid.

LACTICINIUM, Lactarium.

LACTIFÈRE, Lactiferous.

LACTIFÈROUS, *Galactoph'orous*, *Lact'ifer*, (F.) *Lactifère*, from *lac*, *lactis*, 'milk,' and *fero*, 'I carry.' Milk-conveying.

LACTIFEROUS VESSELS, *Lactiferous* or *Galactoph'orous Ducts*, *Tubuli lactiferi*, *Ductus lactiferi* seu *lac'tei*, (F.) *Vaisseaux ou conduits lactifères*, are the excretory ducts of the mammary gland, each of which, running towards the nipple, dilates beneath the areola into an elongated sacculus—the *lacteal* or *galactophorous sac*, *receptacle* or *sinus*—*sacculus* seu *sinus lactiferus*.

LACTIFEROUS OF LACTEAL SWELLING. A tumefaction of the breast, supposed by Sir Astley Cooper to arise from a large collection of milk in one of the lactiferous tubes, the result of chronic inflammation of the tube near the nipple, with closure of its aperture, and obliteration of the canal for an inch or more. The tube requires to be punctured.

LACTIFUGA, *Lact'ifuge*, from *lac*, *lactis*, 'milk,' and *fugo*, 'I drive away.' Medicines which dispel milk.

LACTIGO, Porrigo larvalis.

LACTINE, Sugar of milk.

LACTIN'IA, from *lac*, *lactis*, 'milk.' Food prepared with milk.

LACTIPOTOR, *Galactopotes*.

LACTIS CONCRETIONES, *Infarctus Mam-mæ lacteus*.

LACTIS REDUNDANTIA, *Galactia*—I. Retentio, *Galactischesis*—I. *Sanguinolenti Excretio*, *Galactohæmia*.

LACTISUGIUM, *Antlia lactea*.

LACTIVORUS, *Galactophages*.

LACTOCELE, *Galactocele*.

LACTOLINE, Milk, solidified.

LACTOMETER, *Galactometer*.

LACTOSE, Sugar of milk.

LACTUCA, from *lac*, *lactis*, 'milk,' so called, from its milky juice. *Lactuca Sati'va*, *Lettuce*, *Garden Lettuce*, *Eunn'chium Meco'nis*, *Thridax*, *Cherbas*, (F.) *Laitue ordinaire*. Ord. *Compositæ*, is used as a wholesome salad. The seeds possess a quantity of oil, which, when formed into an emulsion, has been advised in *ardor urinæ*, &c.

The inspissated juice, *Lactuca'rium*, *Thrid'acé*, *Lettuce opium*, resembles, in odour and appearance, that of opium, and is, like it, narcotic, but uncertain. Dose, gr. j to x, and more.

LACTUCA ELONGATA, see *L. virosa*—I. *Graveolens*, *L. virosa*—I. *Floridana*, *Mulgedium Floridanum*.

LACTUCA SCARIOLA, *L. Sylves'tris*, *Scariola*, (F.) *Laitue Scariole* ou *Sauvage*, possesses a stronger degree of bitterness than *L. sativa*. It has similar virtues with *Z. virosa*.

LACTUCA SYLVESTRIS, *L. scariola*—I. *Villosa*, *Mulgedium acuminatum*.

LACTUCA VIRO'SA, *L. grav'olens*, *Strong-scented Lettuce*, (F.) *Laitue vireuse*. The odour of this plant, the leaves of which are used in medicine, is strongly narcotic, and the taste bitter. They are narcotic, diuretic, and aperient; and have been used in dropsies. *Lactuca elong'ata*, *Wild Lettuce* of the United States, has been employed for *L. virosa*.

LACTUCARIUM, see *Lactuca*.

LACTUCIMEN, *Aphthæ*.

LACTUCININA, *Aphthæ*.

LACTUMEN, *Porrigo larvalis*.

LACTUMINA, *Aphthæ*.

LACUNA, *Canaliculus*, diminutive of *lacus*, 'a lake or deep ditch.' A *Fossa* or *Ditch*. A small cavity in a mucous membrane, the parietes of which secrete a viscid humour. It is used synonymously with *crypt*. Also, the external opening of such cavity.

LACUNA, *Crypta*, *Fontanella*—I. *Cerebri*, *Infundibulum* of the brain, *Pituitary gland*.

LACUNA seu SULCUS seu SULCULUS LABII SUPERIORIS, *Amabile*, *Anato'rium*, *Philtrum*, *Phile'trum*. The hollow of the upper lip under the nose.

LACUNA MAGNA, see *Urethra*.

LACUNÆ, see *Urethra*.

LACUNÆ OF BONE. Certain dark stellate spots, with thread-like lines radiating from them, seen under a high magnifying power. These were at first believed to be solid *osseous* or *bone corpuscles* or *cells*, *Corpuscula os'sium*, *Corpuscles* of *Parkinje*, (F.) *Cellules des os ou osseuses*, *Corpuscules des os ou noirs ou ramifiés ou calcaires*, *Ostéoplastes*; but are now regarded as excavations in the bone—*Sacculi chalicoph'ori*—with minute tubes or *canaliculi* proceeding from them, and communicating with the *Haversian canals*. The *lacunæ* and *canaliculi* are fibres concentrated in the transit of nutrient fluid through the osseous tissue.

LACUNÆ GRAAFIA'NÆ, *L. mucosæ vulvæ*. The mucous follicles of the vagina.

LACUNÆ MUCOSÆ VULVÆ, *L. Graafianæ*—I. *Palpebrarum*, *Meibomius*, glands of.

LACUNAR, *La'quear*, 'an arched roof.' The roof of a chamber. Hence,

LACUNAR OR'BITÆ. The roof of the orbit.

LACUNAR VENTRICULI QUARTI SUPERIOR, *Valvula Vieussenii*.

LACUNE DE LA LANGUE, *Cæcum foramen*.

LACUS LACRYMALIS, *Lachrymal sac*.

LACUS LACRYMAR'UM. A small space in the inner angle of the eye between the lids, towards which the tears flow, and at which the triangular canal formed between the closed lids terminates.

LADA, *Piper nigrum*.

LADANUM, see *Cistus creticus*.

LADENDO, *Influenza*.

LADIES MANTLE, *Alchemilla*.

LADIES' SLIPPER, SHOWY, *Cypripedium spectabile*—I. *Slipper*, stemless, *Cypripedium acaule*—I. *Slipper*, yellow, *Cypripedium luteum*.

LADIES' SMOCK, *Cardamine pratensis*.

LADRE, *Leprosus*.

LADRERIE (F.), from *ladre*, 'a leper.' (F.) *Léproserie*, *Maladrerie*. A vulgar name for elephantiasis, or lepra. Also, an hospital for the reception of the leprous, *Leprosarium*, *Domus leprosa'ria*, *Leper hospital*.

LADYBIRD, *Coccinella septempunctata*.

LADYBUG, *Coccinella septempunctata*.

LADYCOW, *Coccinella septempunctata*.

LADY CRESPIGNY'S PILLS, see *Pilulæ Aloes et Kinæ Kinæ*.

LADY HESKETH'S PILLS, see *Pilulæ Aloes et Kinæ Kinæ*.

LADY WEBSTER'S PILLS, see *Pilulæ Aloes et Kinæ Kinæ*.

LADY'S THUMB, *Polygonum persicaria*.

LÆDENS, pl. *Læden'tia*, from *lædo*, 'to hurt.' Any thing hurtful or injurious, as to the economy.

LÆMOPARALYSIS, *Cæphalogiplegia*.

LÆMOS, *Pharynx*.

LÆMOSCIR'RHUS, *Cancer pharyn'gia et œsoph'agi*, *Læmostenosis* seu *Dysph'agia scirrhus'sa*; from *læmos*, 'the pharynx or œsophagus.' *Cancer* of the pharynx or œsophagus.

LÆMOSTENOSIS, *Dysphagia constricta*—I. *Scirrhus*, *Læmoscirrhus*.

LÆSIO, Lesion — 1. Continui, see Continuity, solution of.

LÆTIFICAN'TIA, from *lætifico* (*lætus*, and *facio*). 'I make glad.' Medicines formerly used as cordials, in depression of spirits, &c.

LÆVIGATIO, Levigation.

LÆVITAS INTESTINORUM, Lientery.

LAGENTOMUM, Hærelip.

LAGNEA, Coition, Satyriasis, Sperm.

LAGNEIA, Coition, Satyriasis, Sperm.

LAGNESIS, Furor Femininus, Nymphomania — 1. Furor masculinus, Satyriasis — 1. Salacitas, &c. yriasis.

LAGNEUMA, Coition, Sperm.

LAGNIA, Satyriasis.

LAGNOSIS, Satyriasis.

LAGOCHEILUS, Hærelip.

LAGONES, Flanks.

LAGONOPONOS, Pleurodynia.

LAGONTOMUM, Hærelip.

LAGOPHTHAL'MIA, from *lagos*, 'a hare,' and *ophthalmos*, 'an eye.' *Lagophthalmus*, *Leporinus ocululus*, *Hare's Eye*; (F.) *Ceil de Lièvre*. A vicious arrangement of the upper eyelid, which is so retracted that it cannot cover the globe of the eye during sleep. It has been asserted that this condition of the eye is natural in the hare when asleep.

LAGOPHTHALMUS, *Lagophthalmia*, Geum urbanum.

LAGOSTOMA, Hærelip.

LAG TEETH, Molar teeth.

LAICHE, *Sarsaparilla Germanica* — 1. *des Saibles*, *Sarsaparilla Germanica*.

LAIT, Milk — 1. *Adonissant*, Loach ex ovo — 1. *d'Anesse*, Milk, asses' — 1. *d'Amandes*, Emulsion amygdalæ — 1. *d'Asafœtida*, Mixture asafœtidæ — 1. *de Beurre*, Buttermilk — 1. *de Brebis*, Milk, ewes' — 1. *Caillé*, Curds — 1. *de Chèvre*, Milk, goats' — 1. *Épanché*, *L. répandu* — 1. *de Femme*, Milk, human — 1. *de Jument*, Milk, mares' — 1. *de Poule*, see Ovum.

LAIT RÉPANDU, (F.) *L. épanché*. A popular expression in France, under which is comprehended every kind of disease (and particularly vague pains) occurring after delivery; all being ascribed to diffusion or deposition of milk.

LAIT DE TERRE, *Magnesiæ carbonas* — 1. *de Vache*, Milk of the cow — 1. *Virginal*, Virgin's milk.

LAITEUX, Lactic.

LAITERON DOUX, *Sonchus oleraceus*.

LAITIAT, (F.) Sour whey, in which different wild fruits have been macerated. Said to be much used in the Jura as a refreshing drink.

LAITUE ORDINAIRE, *Lettuce* — 1. *Sauvage*, *Lactuca scariola* — 1. *Scariola*, *Lactuca scariola* — 1. *Vireuse*, *Lactuca virosa*.

LA'ITY, *La'ici*; from *laos*, 'the people.' Formerly, and generally at the present day, the people as distinguished from the clergy. Frequently, however, employed in the sense of the people as distinguished from those of the other learned professions. A *layman* is one of the laity.

LAKEWEED, *Polygonum hydropiper*.

LALIA, Voice, articulated.

LALLA'TION, *Lalla'tio*, *Lambdacismus*, *Labdacismus*, *Lu'laby speech*. Sauvages uses this term for a vicious pronunciation, in which the letter L is improperly doubled, or softened, or substituted for R.

LAMAC, *Acaciæ gummi*.

LAMBDAÇISMUS, Lallation.

LAMBDOID, *Lambdoid'al*, *Lambdo's des*, *Lambdo's des*, *Lambdoideus*, from the Greek letter Λ , $\lambda\alpha\beta\delta\alpha$, and *eidos*, 'shape,' 'resemblance.' Anatomists have given the name **LAMBDOIDAL SUTURE**, *Sutura Lambdoidalis* seu *lambdoïdes* seu *lamb-*

doïdea seu *Proræ*, to the suture, formed by the parietal bones and the occipital, because it resembles the letter Λ , *lambda*, of the Greeks. It is the *Occipito-parietal suture*, *Suture occipitale*, (Ch.) In this suture, the ossa Wormiana are most frequently met with; and the denticulations are most distinctly marked.

LAMBEAU, Flap.

LAMBIVIVUM, Eclectos.

LAMBKILL, *Kalmia latifolia*.

LAMBKILLER, *Andromeda mariana*.

LAME, *Lamina* — 1. *Cornée*, *Tænia semicircularis* — 1. *Ruyschienne*, *Ruyschiana tunica*.

LAMELLA, *Lamina* — 1. *Plana*, *Planum os*.

LAMEL'AR, *Lamello'sus*, *Lam'inated*, (F.) *Lamelleux*, *Lamineux*, composed of thin laminae or leaves — as the *Lamellar* or *laminated tissue*; i. e. the *areolar tissue*.

LAMELLEUX, *Lamellar*.

LAMENESS, Claudication.

LAMIER, *Lamium album*.

LAM'INA, *Lamell'a*, (F.) *Lame*. A thin, flat part of a bone, a plate or table, as the cribriform lamina or plate of the ethmoid bone. *Lamina* and *Lamella* are generally used synonymously; although the latter is properly a diminutive of the former.

LAMINA CILIARIS, Ciliary zone.

LAM'INA CINER'EA. A thin layer of gray substance, which forms the anterior part of the inferior boundary of the third ventricle of the brain.

LAM'INA COR'NEA, *Tæ'nia Tari'ni*. A yellowish band or a thickening of the lining membrane of the ventricle, by which the vena corporis striati is overlaid in the lateral ventricle of the brain.

LAMINA CRIBROSA, *Cribriform lamella* — 1. *Cribrrosa ossis ethmoidæ*, see *Criblé* — 1. *Fusca*, see *Sclerotic* — 1. *Medullaris triangularis cerebri*, *Lyra* — 1. *Papyracea*, *Planum os* — 1. *Perforata anterior*, *Locus perforatus anticus* — 1. *Perforata posterior*, see *Tarini pons*.

LAMINA PERITONÆI EXTERNA. The outer lamina or fold of the peritonæum.

LAMINA PRIMA CUTIS, *Epidermis*.

LAMINA SPIR'ALIS, *Septum scalæ seu coch'leæ audit'riæ*. A partition between the scalæ of the cochlea. The largest part of this, next the modiolus, is formed of bone. The remainder, or that part next the opposite side of the scalæ, is composed of a cartilaginous membrane, called, by Valsalva, *Zona seu Zo'nula Coch'leæ*. By some anatomists, the lamina is divided into a *Zona os'sea* and *Z. mollis*. By others, it is considered to consist of four laminae, when examined with a strong glass: a *Zona os'sea*, next to the modiolus — a *Zona cori'acea*, on the outer side of this: a *Zona vesicul'aris* — and a *Zona membrana'cea*, which is, perhaps, the lining membrane of the cochlea. At the apex of the cochlea, the lamina spiralis terminates by a pointed hook-shaped process, *ham'ulus lam'ina spir'al'is*.

LAMINA OF A VERTEBRA, see *Vertebra*.

LAM'INÆ DORSAL'ES, *Dorsal laminae*. Two oval masses on each side of the primitive groove of the embryo, which approach so as to form a groove, in which are lodged the future brain and spinal marrow.

LAMINÆ SPONGIOSÆ NASI, *Turbinated bones*.

LAM'INÆ VENTRA'LES, *L. viscera'les*. Thickened prolongations of the serous layer of the germinal membrane, which, by their union, form the anterior wall of the trunk of the new being.

LAMINARIA DIGITATA, *Tangle*. One of the Algæ eaten in Scotland, and hawked about the streets with the Pepper-dulse.

The leaf of a sea-weed — a species of *Laminaria* — is employed in the Himalayas under the

name of *goitre leaf*; so called because chewed by the inhabitants, where goitre prevails.

LAMINATED, Lamellar.

LAMINEUX, Lamellar.

LA'MIUM ALBUM, *L. folio'sum*, *Urti'ca mor'tua*, *Galeop'sis Archangel'ica*, *Dead or White Arch'angel Nettle*, (F.) *Ortie blanche ou morte*, (S.) *Daynettle*, *Lamier*. *Ord. Labiate*. Infusions of this plant have been recommended in uterine hemorrhage, and leucorrhœa. It is not used.

LA'MIUM AMPLEXICAUL'E, *Dead Nettle*, *Henbit*; naturalized; flowering from May to October; is regarded as tonic, diaphoretic, and laxative.

LAMIUM FOLIOSUM, *L. album*—*L. Montanum*, *Melittis melissophyllum*—*L. Plinii*, *Melittis melissophyllum*—*L. Rubrum*, *Galeopsis*.

LAMOTTE, MINERAL WATERS OF. These thermal springs are in the department of Isère, France. Temperature, 184°.

LAMPOURDE, *Xanthium*.

LAMPROPH'ONUS, from *λαμπος*, 'clear,' and *φωνη*, 'voice.' One who has a clear voice.

LAMPSANA, *Lapsana*.

LAMPYRIS NOCTILUCA, *Cicindela*.

LANA PHILOSOPHORUM, *Zinci oxydum*.

LANARIA, *Saponaria*, *Verbascum nigrum*.

LANCE DE MAURICEAU, (F.) An instrument invented by Mauriceau for perforating the head of the fœtus. A perforator.

LANCELET, *Lancet*.

LANCEOLA, *Lancet*.

LANCET, *Lance'ola*, *Lancet'ta*, *Schaste'ron*, *Scalpm chirurg'icum*, *Lancet'let*, (Prov.) *Fame*, (F.) *Lancette*,—diminutive of *lancea*, 'a lance.' A surgical instrument, used in the operation of phlebotomy. It is composed of two parts, the handle, (F.) *Chasse*, and the blade, (F.) *Lame*. The former is made of two small plates of ivory, bone, or shell, movable on the blade for whose preservation they are intended. The blade is formed of well-polished steel. Lancets are made of different shapes; some being broad-shouldered—others, *spear-pointed*. The French distinguish three kinds: 1. *Lancette à grain d'orge*, which, on account of the almost oval shape of its point, makes a large opening. 2. The *L. à langue de serpent*, which is very narrow towards the point; and, 3. *L. à grain d'avoine*, which holds a medium station between the two former, and is generally preferred.

The *Abscess Lancet* is merely a large lancet for the purpose of opening abscesses.

The employment of a lancet, for opening abscesses, &c., is called *Lancing*.

LANCETTA, *Lancet*.

LANCETTE, *Lancet*.

LANCETTIER, (F.) A lancet-case.

LANCINANT, *Lancing*.

LANCINATING, *Lan'cinans*, from *lancinare*, *lancinatum* (*lancea*, 'a lance,') 'to strike or thrust through.' (F.) *Lancinant*, (substantive, *Élancement*.) A species of pain, which consists in lacerations or shootings, similar to those that would be produced by the introduction of a sharp instrument into the suffering part. It is especially in cancer that this kind of pain is felt.

LANCING, see *Lancet*—*L. the Gums*, *Dé-choussement*.

LAND'S END, CLIMATE OF. The climate of the Land's End, in England, resembles that of the south of Devonshire, but is more relaxing. It is considered to be most likely to prove beneficial in consumptive cases, in which the disease is accompanied by an irritated state of the pulmonary mucous membrane, producing a dry cough. Where the system is relaxed, and the secretion from the lungs considerable, the climate, it is conceived, will generally prove injurious.

As a brumal retreat, the southern coast of Devonshire would seem to be preferable to it.

LANGÉAC, MINERAL WATERS OF. Acidulous, mineral waters at Langeac, in the department of Haute-Loire, France. They are employed as refrigerant, aperient, and diuretic. They contain carbonic acid, carbonates of soda and magnesia, and a little iron.

LANGDEBEFF, *Ancusa officinalis*.

LANGENSCHWALBACH, *Schwalbach*.

LANG'II AQUA EPILEP'TICA, *Epilep'tic Water of Langius*, formerly employed against epilepsy. It was composed of the flowers of *convallaria* and *lavender*, *Spanish wine*, *cinnamon*, *nutmeg*, *mistletoe*, *peony* and *dittany roots*, *long pepper*, *cubeba*, and *rosemary flowers*.

LANGUE, Tongue—*L. Abaisseur de la*, *Glossocotichus*—*L. de Carpe*, *Fish-tongue*, see *Lever*—*L. de Cérif*, *Asplenium scolopendrium*—*L. de Cheval*, *Ruscus hypoglossum*—*L. de Chien*, *Cynoglossum*—*L. de Serpent*, *Ophioglossum vulgatum*.

LANGUEUR, *Langueur*.

LANGUOR, *Aph'esis*, *Faintness*, (F.) *Langueur*. A species of atony, depression, or debility, which generally comes on slowly.

LANGWORT, *Veratrum album*.

LANK, *Inguen*.

LAN'TANA, *Sage Tree*, *Blueberry*, (F.) *Cail-leau*. The leaves of this indig-nous plant form a fine-scented tea, like *L. Camara* or *Bahama Tea*, and *L. Pseudothe'a* or *Brazil Tea*. The tea is used as a diaphoretic.

LANU'GO, *Pili cutis*, *Pluma*, from *lana*, 'wool.' The soft, fine hair on different parts of the body, especially of the young.

LANUGO PRIMA, *Genecia*—*L. Pudendorum*, *Pubes*—*L. Siliquæ hirsutæ*, see *Dolichos pruriens*.

LANUVIUM, *Vulva*.

LAÏON'ICA CURA'TIO seu CURA, from *λαος*, 'the people.' A popular mode of treating the gout in the middle ages, which consisted in the employment of local applications, proper for evacuating the morbid matter. [?]

LAOS, *Tin*.

LAPACTICUS, *Cathartic*, *Laxative*.

LAPARA, *Abdomen*, *Flanks*, *Lumbi*.

LAPARENTEROTOMIA, *Laparo enterotomy*.

LAPAROC'ELE, from *λαπαρά*, 'the lumbar region,' and *κηλη*, 'rupture.' *Lumbar Hernia*, through a separation of the fibres of the quadratus lumborum, and a protrusion of the aponeurosis of the transverse muscle on the outside of the mass common to the sacro-lumbalis and longissimus dorsi.—*Cloquet*.

LAPAROCYSTOTOMIA, see *Lithotomy*.

LAPARO-ELYTROTOMIA, *Cæsarean section*.

LAPARO-ENTEROT'OMY, *Lap'aro-enterotom'ia*, *Laparenterotom'ia*, from *λαπαρά*, 'the lumbar region,' the 'abdomen,' *εντερον*, 'intestine,' and *τομή*, 'incision.' The operation of opening the abdomen and intestinal canal, for the removal of disease.

LAPAROSCOPIA, *Abdominoscopia*.

LAPAROTOMY. *Gastrostomy*, especially when executed in the lumbar region. See *Gastrostomy*.

LAPATHOS, *Rumex acutus*—*L. Aquaticum*, *Rumex hydrocephalum*—*L. Chinense*, *Rheum*—*L. Orientale*, *Rheum*—*L. Sanguineum*, *Rumex sanguineus*—*L. Unctuosum*, *Chenopodium bonus Henricus*.

LAPATHUM, *Rumex acutus*—*L. Acutum*, *Rumex acutus*—*L. Hortense*, *Rumex patientia*—*L. Pratense*, *Rumex acetosus*—*L. Scutatum*, *Rumex scutatus*—*L. Sylvestre*, *Rumex acutus*, *R. obtusifolius*.

LAPAXIS, *Evacuation*.

LAPE, *Mucus*.

LAPIDIL'LUM, from *lapis*, 'a stone.' *Blasius*

has given this name to a kind of scoop, used for extracting stone from the bladder.

LAPILLATIO, Lithia.

LAPILLI GLANDULÆ PINEALIS, see Pineal gland.

LAPILLUS, diminutive of *lapis*, 'a stone.' A small stone; gravel; a grain of sand.

LAPIS, Calculus.—*l. Accipitrum*, Hieracites.—*l. Aerous*, Calamina.—*l. Ammoniaci*, see Ammoniac gum.—*l. Animalis*, Blood.—*l. Armenius*, Melochites.—*l. Aureus*, Urine.—*l. Bezoardicus*, Bezoar.—*l. Cæruleus*, Cupri sulphas.—*l. Calaminaris*, Calamina.—*l. Carneolus*, Cornelian.—*l. Cansticus*, Potassa cum calce, Potassa fusa.—*l. Collymus*, Ætites.—*l. Contraryervæ*, Pulvis contraryervæ compositus.—*l. Cyanus*, *l. lazuli*.

LAPIS DIVINUS, *L. Ophthalmicus* seu *Ophthalmicus Sti. Ivesii*, *Coprum aluminatum*, (*F.*) *Pierre divine*, (*Cupri sulphat.*, *Alumini*, *Potass. nitrat.* æ 3j. Melt together, adding at the end *Camphor.* ʒss.) Employed to make an eye-water, ʒij ad aqua ʒiv.

LAPIS FULMINEUS, Ceraunion.—*l. Hematites*, *Hæmatites*—*l. Heracleus*, Magnet.—*l. Infernalis*, *Argenti nitrus*—*l. Infernalis alkalinus*, Potassa fusa.—*l. Infernalis sive septicus*, Potassa cum calce.—*l. Judaicus*, Judæus (lapis).

LAPIS LAZULI, *L. Cyannus*, *Asulei*, *Lazulite*, (*F.*) *Pierre d'azur*, *Outremer*. A stone, of a beautiful blue colour; opaque; and close-grained; fracture, dull. It is composed of silex; alumina; carbonate and sulphate of lime; oxide of iron, and water. It was formerly looked upon as a purgative and emetic, and given in epilepsy.

LAPIS MALUCENSIS, Bezoard of the Indian porcupine.—*l. Nauticus*, Magnet.—*l. Ophthalmicus*, *L. Divinus*—*l. Ophthalmicus Sti. Ivesii*, *L. Divinus*—*l. Phœnicites*, Judæus (lapis)—*l. Porcinus*, Bezoard of the Indian porcupine.—*l. Prunellæ*, Potassæ nitrus fusus sulphatis pucillo mixtus.—*l. Sardius*, Cornelian.—*l. Septicus*, Causticum commune, Potassa fusa.—*l. Specularis*, *S. lucidum*—*l. Syderitis*, Magnet.—*l. Syriacus*, Judæus (lapis)—*l. Vini*, Potassæ supertartas impurus.

LAPPA, *Arctium lappa*, *Galium aparine*, *Lip-pitudo*—*l. Minor*, *Xanthium*.

LAPPULA HEPATICA, Agrimony.

LAPSAÑA, *Lampisana*, *Nâpium*, *Papillaris herba*, *Dock-creases*, *Nipple-Wort*, (*Sc.*) *Bolgan leaves*, (*F.*) *Lampasane*, *Herbe aux Mamelles*. This plant is a lactescent bitter. It has been chiefly employed, however, as an external application to sore nipples, &c.

LAPSUS PILORUM, Alopecia.

LAQ, Laqueus.

LAQUE, Laccæ.

LAQUEAR, Lacunar.

LAQUEAR VAGINÆ, *Fundus Vaginæ* (laquear, 'the inward roof of a house'). The part of the vagina in which the cervix uteri terminates.

LAQUEUS, *Funnis*. A cord, ligature or bandage, with running knots:—a Noose, a loop. A *fillet*, *Brochos*, *Pach'eton*, (*F.*) *Lac*, *Laq* ou *Lacq*. The term is applied to a bandage or fillet of any kind, attached by means of a loop upon any part, with the view of fixing it; as in certain cases of labour, where a hand or foot presents; or to facilitate extension in luxations and fractures.

Also:—A prominent band in the brain, *Lemniscus*, behind the brachium posterius of the corpora quadrigemina, which marks the course of the superior division of the fasciculus olivaris.

LAQUEUS GUTTURIS, 'Noose of the throat.' Violent inflammation of the tonsils, in which the patient appears as if suffocated by a noose. According to some, gangrenous cynanche.

LARCH, *Pinus larix*.

LARD, Adeps.

LARDACÆ, Lardaceous.

LARDA'CEOUS, *Larda'ceus*, *Lar'deus*, *Lard'iform*, *Lardiformis*, (*F.*) *Lardacæ*; from *lardum*, 'lard,' the fat of bacon. An epithet given to certain organic alterations in the textures, whose aspect and consistence resemble lard. (*F.*) *Tissus lardacæ*. They constitute what has been termed the *Larda'ceous* or *Chol'esterine Disease*. From their waxy appearance the pathological condition has been called *Cero'ma*.

LARDEUS, Lardaceous.

LARDIFORM, Lardaceous.

LARDUM, Adeps.

LARGE, Broad.

LARIDUM, Adeps.

LARIX, *Pinus larix*—*l. Communis*, *Pinus larix*—*l. Decidua*, *Pinus larix*—*l. Europæa*, *Pinus larix*—*l. Pyramidalis*, *Pinus larix*.

LARKSPUR, BRANCHING, *Delphinium consolida*.

LARME, Tear.

LARMOIEMENT, Epiphora.

LAROCHE POSAY, MINERAL WATERS OF. Simple sulphurous waters in the department of Vienne, France.

LAR'REA MEXICA'NA, *L. glutinosum*, *Crescote plant*. Order, *Zygophyllacæ*. A shrub, growing in Arkansas, Southern Texas, New Mexico, and Northern Mexico, which abounds in a strong-scented resinous matter; and is used, externally and internally, in rheumatism, and in syphilitic, and other pains.

LARVA, Mask. Also the larve, grub, or vermiform condition of an insect: the first change it experiences after leaving the ovum. Larves of insects are occasionally developed in the intestinal canal from ova swallowed. See *Ectozaa*.

LARVE, Mask.

LARVÉ, Masked.

LARYNGÉ, Laryngeal.

LARYNGE'AL, *Laryng'eus*, (*F.*) *Laryngé*, *Laryngien*. Same etymon as *Larynx*. That which belongs to the larynx.

LARYNGEAL ARTERIES are given off from the thyroid arteries.

LARYNGEAL NERVES, (*F.*) *Nervi Laryngés*, are two in number:—a superior and an inferior. The superior laryngeal nerve is given off from the trunk of the pneumogastric, at the upper and deep part of the neck. It passes downwards and inwards, behind the internal carotid artery, and divides into two secondary branches; the one, *external*, which distributes its filaments, on the outside of the larynx, to the sterno-thyroid, hyo-thyroid, constrictor inferior, crico-thyroid muscles, &c. The other, the *internal*, which crosses the thyro-hyoid membrane, and gives filaments to the epiglottis, the mucous membranes of the pharynx and larynx, to the arytenoid gland, the arytenoid and crico-thyroid muscles, and ultimately anastomoses with the inferior laryngeal nerve. The inferior laryngeal nerves or *recurrents*, *Nervi reversi'vi*, *Rameaux Trachæaux* (*Chaus.*) arise from the trunk of the pneumogastric within the thorax. They ascend in the furrow, which separates the trachea from the œsophagus, to be distributed on the neck, after having been reflected;—the left around the arch of the aorta; the right, around the corresponding subclavian. They send off filaments to the cardiac plexuses; to the pareties of the œsophagus, and trachea; to the thyroid gland; to the inferior constrictor of the pharynx; the posterior and lateral crico-arytenoid and thyro-arytenoid muscles; and to the mucous membrane of the pharynx and larynx.

LARYNGEAL PHTHISIS, Phthisis, laryngeal.

LARYNGEAL POUCHES. Small *culs-de-sac*, which lead from the anterior part of each ventricle of the larynx upwards, for the space of half an inch, between the superior ligaments on the inner side, and the thyroid cartilage on the outer, reaching as high as the upper border of that cartilage at the sides of the epiglottis. Numerous small glands open into them.

LARYNGEAL SOUND, Laryngeche.

LARYNGEAL VEINS open into the internal jugular. Winslow gives the name *laryngeal* to the superior thyroid artery.

LARYNGEAL VOICE, see Voice.

LARYNGE'CHE, from *larynx*, and *ηχη*, or *ηχος*, 'sound.' The laryngeal sound heard by the stethoscope during breathing and speaking.

LARYNGEMPHRAX'IS, from *larynx*, and *εμφραξις*, 'obstruction.' Obstruction of the larynx from any cause.

LARYNGIEN, Laryngeal.

LARYNGISM'US, from *larynx*. Spasm of glottis, giving occasion to contraction or closure of the opening.

LARYNGISMUS STRIDULUS, Asthma thymicum.

LARYNGITE, Laryngitis—*l. Muqueuse*, Laryngitis (simple)—*l. Edémateuse*, Edema of the glottis—*l. Edémateuse et séro-purulente*, Edema of the glottis—*l. avec Production de Faussees membranes*, Cynanche trachealis—*l. Pseudo-membraneuse*, Cynanche trachealis—*l. avec Sécrétion de Pus*, Phthisis laryngea—*l. Sus-glottique*, Edema of the glottis—*l. Striduleuse*, Asthma thymicum—*l. Sous-muqueuse*, Edema of the glottis.

LARYNGIT'IS, from *larynx*, and *itis*, a suffix denoting inflammation; *Inflammat'io Laryng'ie*, *Cynanchè* seu *Angi'na laryng'e'a*, (F.) *Laryngite*, *Catarrhe laryngien*, *Angine laryngé*, *Inflammation of the Larynx*. This disease, in some measure, resembles croup; but is usually devoid of that peculiar sonorous inspiration, which attends the latter. There is, also, pain upon pressing the larynx; and, whilst laryngitis is a disease of more advanced life, croup attacks children. The membraniform exudation is, also, absent; probably, because the inflammation, being seated above the glottis, the coagulable lymph is readily expectorated. It requires the most active treatment.

Simple Laryngitis is called by some *mucous Laryngitis*, *Laryngitis acuta* seu *mucosa acuta*, (F.) *Laryngite muqueuse*, to distinguish it from *submucous Laryngitis* or Edema of the glottis.

Chronic Laryngitis is generally regarded as synonymous with laryngeal phthisis; but it may exist independently.

LARYNGITIS ACUTA, Laryngitis—*l. Chronic*, see Phthisis laryngea—*l. Exsudativa*, Cynanche trachealis—*l. Membranacea*, Cynanche trachealis—*l. Mucosa acuta*, Laryngitis—*l. Mucous*, Laryngitis (simple)—*l. Edematous*, Edema of the glottis—*l. Polyposa*, Cynanche trachealis—*l. Seropurulenta*, Edema of the glottis—*l. Submucous*, Edema of the glottis—*l. et Tracheitis chronica*, see Phthisis laryngea—*l. et Tracheitis infantilis*, Cynanche trachealis.

LARYNGOCACE, Cynanche trachealis.

LARYNGO-CATAR'RHUS, *Catarrhus Laryngis et tracheal'is*. Catarrh affecting the larynx and trachea more especially, as indicated by alteration of the voice—hoarseness—itching and sensation of burning in those parts; short cough and expectoration, &c.

LARYNGO-ET-TRACHEO-PHTHISIS, Phthisis laryngea.

LARYNGO'GRAPHY, *Laryngogra'phia*; from *larynx*, and *γραφη*, 'a description.' An anatomical description of the larynx.

LARYNGOL'OGY, from *larynx*, and *λογος*, 'treatise.' A treatise on the larynx.

LARYNGOPARALYSIS, see Aphonia.

LARYNGOPH'ONY, *Laryngopho'nia*, *Tracheoph'ony*, *Larynge'al voice*, *Tracheal voice*: from *larynx*, and *φωνη*, 'voice.' The sound heard in health, when the stethoscope is placed over the larynx or trachea at the time a person speaks. The voice appears to pass immediately up to the ear of the auscultator. A similar physical sign exists when there is a cavity in the lungs, and the instrument is placed over it whilst the patient speaks. See Pectoriloquy.

LARYNGOPHTHISIS, Phthisis laryngea.

LARYNGOSPASMUS, Asthma thymicum.

LARYNGOSTASIS, Cynanche trachealis.

LARYNGOSTENO'SIS, from *larynx*, and *στενωσις*, 'contraction.' Contraction or narrowing of the larynx.

LARYNGOT'OMY, *Laryngotom'ia*, from *larynx*, and *τεμνειν*, 'to cut.' A surgical operation, which consists in opening the larynx, either to extract a foreign body, or to remedy an obstruction of the glottis. The operation is, sometimes, erroneously called *Bronchotomy*, and *Trachelotomy*.

LARYNGO-TRACHEITIS, Cynanche trachealis—*l. Tracheitis* with Diphtheritic exudation, Cynanche trachealis.

LAR'YNX. (gen. *Laryng'is*,) λαρυγξ, λαρυγγος. *Caput* seu *Operculum* seu *Init'ium* seu *Finis superior* seu *Terminus superior* seu *Partis primæ asperæ arte'riæ*. The apparatus of voice is situated at the superior and anterior part of the neck; and at the top of the trachea, with which it communicates. It is composed of four cartilages,—the thyroid, cricoid, and two arytenoid; is moved by a number of muscles, and lined by a mucous membrane, having certain membranous reflexions, constituting the superior ligaments of the glottis, &c.

PARTS COMPOSING THE LARYNX.

- | | | |
|---------------------|------------|--------------------------------------|
| | | Thyroid. |
| | | Cricoid. |
| 1. Cartilages..... | | Two arytenoid. |
| | | Epiglottis. |
| | Extrinsic, | Sterno-thyroid. |
| | | Constrictors of the pharynx. |
| | | All the muscles of the hyoid region. |
| 2. Muscles. | | Crico-thyroid. |
| | | Crico-arytenoid, posterior. |
| | Intrinsic. | Crico-arytenoid, lateral. |
| | | Thyro-arytenoid. |
| | | Arytenoid. |
| 3. Mucous Membrane. | | |
| | | Epiglottic. |
| 4. Glands..... | | Arytenoid. |
| | | Thyroid. |
| 5. Membranes..... | | Thyro-hyoid. |
| | | Crico-thyroid. |
| 6. Ligaments..... | | Crico-arytenoid. |
| | | Thyro-arytenoid. |

The vessels and nerves of the larynx are called *Laryngeal*.

The larynx is destined to give passage to the air, in the act of respiration, and to impress upon it certain modifications, which constitute voice. Its dimensions vary in different individuals. In men, it is always larger, and situate lower, than in women.

LARYNX, PELLICULAR OR PLASTIC INFLAMMATION OF THE, Cynanche trachealis.

LAS'ANUM, *Sella familiar'ica*, (F.) *Chaise percée*. A close stool.

LASCIVIA, Satyriasis.

LASCIVITAS, Satyriasis.

LASCIVUS, Libidinous.

LASER, *Laserpitium*—*l. à larges feuilles*, *Laserpitium latifolium*.

LASERPIT'IUM, *Laser*, *Sil'phium*, *σικερίον*. A term applied, anciently, both to a plant and its juice, regarding the nature of neither of which we possess any precise information. Bentley, Laurence, Geoffroi, &c., regard it to have been the same as *asafoetida*:—Theophrastus, Dioscorides, and the ancient scholiast of Aristophanes, however, by assigning a sweet and agreeable flavour to the *laserpitium*, discountenance the idea. From whatever plant obtained, it was so rare, and consequently so costly, that the Romans deposited it in the public treasury. It was obtained from Cyrene—*Succus Cyren'dicus*—and likewise from Persia—the latter being the most valuable. The *Laserpitium* is called by *Ariceana*, *Altihit*.

LASERPITIUM ASPERUM, *L. latifolium*.

LASERPITIUM LATIFOLIUM, *L. as'perum*, *Gentiana alba*, *White Gentian*, *Cervaria alba*; (F.) *Laser à larges feuilles*. The root of this plant is bitter and tonic.

LASERPITIUM MONTANUM, *L. siler*.

LASERPITIUM SILER, *L. trifoliatum* seu *montanum*, *Ses'eli*, *Seselia*, *Siler montanum* seu *lan-cifolium*, *Heart-wort*, *Sermountain*. The seeds and roots are possessed of aromatic properties.

LASERPITIUM TRIFOLIATUM, *L. siler*.

LASSITUDE OCULAIRE, *Coryopia*.

LASSITUDO, *Copos*.

LAST, *Inguen*.

LATENS IN ORE, *Pterygoideus internus*.

LATENT, *Latens*, from *latere*, 'to lie hid.' 'Lying hid,' 'concealed.' An epithet applied to certain diseases or states of disease, in which the symptoms are so concealed and obscure, *morbi occult'i*, as to escape the observation of the physician. Thus, we say *latent inflammation*, *latent period of small-pox*.

LATERAD, from *latus*, *lateris*, 'the side.' A term used adverbially by Dr. Barclay to signify 'towards the lateral aspect.'

LATERAL ASPECT, see *Mesial*.

LATERIT'IOUS, *Laterit'ius*, *Lateric'ius*, (F.) *Briqueté*, from 'later', *lateris*, 'a brick.' An epithet applied to the brick-like sediment, occasionally deposited in the urine of people afflicted with fever; *Sedimentum urinae lateric'ium*;—(F.) *Urine briqueté*.

LATEROFLEX'ION, *Lat'ero-flex'io*. A bending to one side,—as of the uterus.

LATÉRO-VERSION, *Lat'ero-ver'sio*. A turning or version to one side,—as of the uterus.

LATESCENTIS CHORDÆ (*Musculus*), *Palmaris longus*.

LATEX NIVEUS, *Milk*.

LATHYRIS, *Euphorbia lathyris*.

LATIB'ULUM, from *lateo*, 'I lie hid.' The foyer of a febrile poison; whence it spreads to every part to induce a paroxysm. See *Clapier*.

LATICA. Same etymon. *Amphimer'ina lat'ica* of *Sauvages*. A species of quotidian remittent, whose paroxysms are very long, and which is accompanied with *latent* or slightly developed fever, whence its name.

LATICES LACTEI, *Receptaculum chyli*.

LATISSIMUS COLLI, *Platysma myoides*.

LATIS'SIMUS DORSI, (superlative of *latus*, 'broad.') *Aniscal'ptor*, *Brachium movens quartus*, (F.) *Lombo-huméral* (Ch.), *Dorsi-lombo-sacro-huméral*, *Muscle grand dorsal*, *M. très large du dos*. A flat muscle; broad, especially below; thin; quadrilateral; and situate at the posterior, lateral, and

inferior region of the trunk. It is attached to the posterior half of the outer lip of the crest of the ilium; to the posterior surface of the sacrum; to the spinous processes of the six or seven last dorsal vertebrae, to all those of the loins, and to the last four false ribs, and is inserted by a strong tendon at the posterior edge of the bicipital groove of the humerus.⁶⁶ Its upper fibres are almost horizontal; the middle very long and oblique upwards and outwards; and the anterior almost vertical. This muscle carries the arm backwards, depressing it, and making it turn on its axis. It also draws backwards and downwards the prominence of the shoulder. When, suspended by an arm, we make an effort to raise ourselves, it draws the trunk towards the arm. It can, also, raise the ribs by assuming its fixed point on the humerus, and become an inspiratory muscle.

LATITUDO HUMERI, *Scapula*.

LATTICE WORK, *Cancelli*.

LATUS, Broad—*L. Ani*, *Levator ani*.

LAU, see *Spirit*.

LAUCA'NIA, *Leuca'nia*, *Lau'chané*. The fauces and œsophagus. Also, the chin.—*Gorræus*.

LAUCHANE, *Laucania*.

LAUD'ANUM or **LAUDA'NUM**. Perhaps, from *laus*, *laudis*, 'praise'; *lauda'tum*, 'praised.' Every preparation of opium, solid or liquid, but more particularly the extract and tincture, and especially the latter.

LAUDANUM ABBATIS ROUSSEAU, *Gutta Abbatis Rousseau*, *Vinum opiatum fermentati'vè parat'um*, *Abbe Rousseau's Drops*. (*Mel. Narbonnensis*, f3xij; *aqua calidâ*, Oijj. Set in a warm place, and, as soon as the mixture ferments, add *opium*, ℥iv, dissolved in *water*, f3xij. Let it ferment for a month, and evaporate to f3x; strain, and add *rectified spirit of wine*, f3ivss.

LAUDANUM, *FOR'D's*, *Vinum opii*—*L. Liquid*, *Tinctura opii*—*L. Liquidum Hoffmanni*, *Vinum opii*—*L. Liquidum Sydenhami*, *Vinum opii*—*L. Opiatum*, *Extractum opii*—*L. Simplex*, *Extractum opii*—*L. Sydenham's*, *Vinum opii*.

LAUGH, *Risus*—*L. Sardonic*, *Canine laugh*.

LAUGHING, *Risus*.

LAUGHTER, *Risus*.

LAUREL, *Kalmia latifolia*, *Magnolia macrophylla*—*L. Broad-leaved*, *Kalmia latifolia*—*L. Cherry*, *Prunus lauro-cerasus*—*L. Common*, *Prunus lauro-cerasus*—*L. Dwarf*, *Kalmia angustifolia*—*L. Great*, *Rhododendron maximum*—*L. Ground*, *Epigæa repens*—*L. Mountain*, *Kalmia latifolia*, *Rhododendron*—*L. Narrow-leaved*, *Kalmia angustifolia*—*L. Pale*, *Kalmia glauca*—*L. Poison*, *Prunus lauro-cerasus*—*L. Rose*, *Kalmia latifolia*—*L. Sheep*, *Kalmia angustifolia*—*L. Swamp*, *Kalmia glauca*—*L. Sweet*, *Illicium floridanum*—*L. Water*, see *Prunus lauro-cerasus*—*L. White*, *Magnolia glauca*.

LAURENT, *SAINT*, *MINERAL WATERS OF*. A thermal spring, five leagues from Joyeuse in France. Temp. 127° Fahr.

LAUREN'TIA PINNATIFIDA, *Pepperdulse*. One of the Algae, eaten in Scotland, and hawked about the streets of Edinburgh along with *Lamina'ria digit'ata* or *Tangle*.

LAUREOLA, *Daphne laureola*.

LURIER, *Laurus*—*L. Alexandrin*, *Ruscus hypoglossum*—*L. Amandier*, *Prunus lauro-cerasus*—*L. d'Apollon*, *Laurus*—*L. Cerise*, *Prunus lauro-cerasus*—*L. de Ceylon*, *Laurus cinnamomum*—*L. Commun*, *Laurus*—*L. des Cuisines*, *Laurus*—*L. des Montagnes*, *Kalmia latifolia*—*L. Noble*, *Laurus*—*L. Rose*, *Nerium oleander*—*L. de Trébisonde*, *Prunus lauro-cerasus*.

LAURO-CERASUS, *Prunus lauro-cerasus*.

LAUROSE, *Nerium oleander*.

LAURUS, *Laurus nobilis*, *Daphne*, *Sweet Bay*. *Nat. Ord.* Lauracæ. (F.) *Lawrier*, *L. noble* ou *des cuisines* on *commun* ou *d'Apollon*. The leaves and berries, *Laurus* (Ph. L.), have a sweet, fragrant smell, and an aromatic, astringent taste. Sweet bay has been advised as a stomachic and carminative, but is now rarely used. It is, sometimes, employed as a fomentation and in glysters; and the berries are an ingredient in the *Emplastrum Cuscutæ*.

LAURUS ÆSTIVALIS, *L. benzoin*—*L. Alexandrina angustifolia*, *Ruscus hypoglossum*.

LAURUS BENZOIN, *Benzoin odoriferum*, *Laurus Pseudo-benzoin* seu *æstivalis*, *Spice wood*, *Spice bush*, *Benjamin bush*, *Allspice bush*, *Wild allspice*, *Spice berry*, *Fever wood*, *Fever bush*. An indigenous shrub, growing in moist, shady places, in all parts of the United States; flowering early in spring. All parts of the shrub have a spicy, agreeable flavour, which is strongest in the bark and berries. An infusion or decoction of the small branches is used in popular practice as a vermifuge, and an agreeable drink in low fevers. The bark has been used in intermittents; the berries, dried and powdered, for allspice. The oil of the berries is used as an excitant.

LAURUS CAMPHORA, see *Camphor*—*L. Canella*, *L. Cassia*.

LAURUS CAS'SIA, *L. Canella*, *Persea cassia*. The species of *laurus* which yields the *Cassia*, *Cassia*, *Canella*, *Cassia lignea* seu *cinnamomea*, *C. lignea Malabarica*, *Cortex Canellæ Malabarica*, *Cortex cras'sior*, *Xylo-cassia*, *Canella Malabarica* et *Javen'sis*, *C. Cuba'na*, *Arbor Judæa*, *Canellifera Malabarica*, *Cinnamomum Malabaricum* seu *Indicum* seu *Sinense*, *Culiac'ha canella*, *Wild Cinnamon*, *Malabar Cinnamon*, *Cassia*, (F.) *Cannelle de Malabar* ou *de Java* ou *de la Chine* ou *des Indes* ou *de Coromandel*. *C. fausse*, *C. matte*, *Casse en bois* ou *aromatique*. The bark and leaves abound with the flavour of cinnamon, for which they may be substituted; but they are much weaker. The unopened flower-buds are used in the same manner. **AQUA CASSIÆ**, *Water of Cassia*, prepared by distillation, is official in the Edinburgh Pharmacopœia.

LAURUS CINNAMOMOIDES, *Nectandra cinnamomoides*.

LAURUS CINNAMOMUM, *Persea cinnamomum*, *Cinnamomum*, *C. Zeylanicum*, *Darseni*, *Darsi'ni*, *Cinnamon*, *Xylo-cinnamomum*, *Cuirdo Canella*, (Sc.) *Cannel*, (F.) *Cannelle*, *Baume de Cannelle*, *Cannelle officinale* ou *Ceylon*. *Cinnamon bark*, which is obtained, also, from the *Cinnamomum aromaticum*, is stimulant and carminative, and is employed, chiefly, as a grateful aromatic, to cover the taste of nauseous remedies. Dose, gr. x to ℥j. *The Flowers*, called *Cassia Flores* in the shops, possess aromatic and astringent virtues, and may be used wherever cinnamon is required. The volatile oil of the bark—*O'leum Cinnamomi*—*Oil of Cinnamon*, is official in the Pharm. U. S.

LAURUS CUBEBA, *Piper cubeba*—*L. Culilaban*, *L. Culilawan*.

LAURUS CULILAWAN, *L. Oulilaban* seu *Caryophyllus*, *Cinnamomum Culilawan*. The tree that affords the *Cortex Culilawan* seu *Culilaban*, *Culilawan*, *Cortex caryophylloides* seu *Caryophylloides Amboinensis*. This bark resembles the *sassafras* in appearance and properties, and is used in Java as a condiment.

LAURUS MALABATHRUM, see *Malabathrum*—*L. Nobilis*, *Laurus*—*L. Persea*, *Persea gratissima*—*L. Pseudobenzoin*, *L. Benzoin*.

LAURUS SASSAFRAS, *Persea sassafras*, *Sassafras*, *Cornus mas odorata*, *Anhuiba*, *Aque-free*, *Aque-tree*, (F.) *Pavanne*. Indigenous in the United States. *Sassafras* wood and root, and especially the bark of the root, *Sassafras Radicis Cortex* (Ph. U. S.), have been considered

stimulant, sudorific, and diuretic. The virtues depend upon essential oil, *Oleum Sassafras*, the odour of which is not unlike that of fennel. It has been used in cutaneous diseases, chronic rheumatism, &c.

The pith of the stems, *Sassafras medulla* (Ph. U. S.), abounds in gummy matter, which it readily imparts to water, forming a limpid mucilage, which is much employed as a collyrium in ophthalmia, and as a drink in dysentery, catarrh, &c. (one drachm of the pith to a pint of boiling water).

LAUTIS'SIMA VINA. (Superlative of *lautus*, 'elegant.') Wines were formerly so called, which were strongly impregnated with myrrh.

LAVAMENTUM, *Clyster*, *Enema*.

LAVANDE, *Lavendula*—*L. de Mer*, *Statice limonium*—*L. Trieste*, *Statice limonium*.

LAVAN'DULA, *Laven'dula*, from *lavo*, 'I wash'; so called from being used in baths. *Lavan'dula spica* (Linnaeus) seu *angustifolia* seu *officinalis* seu *vera*, *Common Lavender*, (F.) *Lavande*. *Ord.* Labiatae. The odour of lavender flowers, *Lavand'ula* (Ph. U. S.), is fragrant and agreeable; taste warm and bitterish—depending upon an essential oil. It has been used as a stimulant; particularly in the form of the oil,—*O'leum lavandulæ*. The dried leaves have been employed as an emmenagogue.

The French use, also, *LAVANDULA STÆCHAS*, *Stæchas* et *Stichas Arabica*, *French Lavender*, of which they have a compound syrup, *Syrupus de stæchade compos'itus*: given as a pectoral.

LAVANDULA LATIFOLIA, *Lavandula*.

LAVANDULA SPICA (Linnaeus), *L. Latifolia Spica*, *French Lavender*, (F.) *Lavande*, *Aspic*, *Spic*, yields the *Oil of Spike*, *O'leum spicæ*, (F.) *Huile d'Aspic* ou *de Spic*, which is used to make varnishes, and as a liniment in paralysis.

LAVEMENT, *Clyster*, *Enema*—*L. Antispasmodique*, *Enema fœtidum*.

LAVENDER, *COMMON*, *Lavendula*—*L. French*, *Lavandula spica*—*L. Sea*, *Statice limonium*, *Statice Caroliniana*.

LAVENDULA, *Lavandula*.

LAVER, *Uva latissima*—*L. Broad green*, *Uva latissima*—*L. Germanicus*, *Veronica beccabunga*—*L. Laciniated*, *purple*, *Uva umbilicalis*—*L. Odoratum*, *Sisymbrium nasturtium*.

LAVIPEDIUM, *Pedilivrum*.

LAW MEDICINE, *Medicine*, *legal*.

LAWSONIA ALBA, *L. Inermis*.

LAWSONIA INERMIS, *L. alba*, *Alican'a vera* seu *Orientalis*, *Cyperus antiquorum*, *Ligustrum Egyptianum*, *Smooth Lawsonia*, (F.) *Henné*, *Al-hanne*. An East Indian and African plant, the root of which is slightly astringent.

In India, the root of the *Lawsonia spinosa* is employed in lepra and other cutaneous affections.

LAX, *Diarrhoea*.

LAXANS, *Laxative*, *Relaxant*.

LAXATIVE, *Laxativus*, *Lazans*, *Le'niens*, from *laxare*, *laxatum*, 'to loosen'; *Minorati'vus*, *Soluti'vus*, *Alvid'ucus*, *Hypochoreticus*, *Hypog'us*, *Hypel'atus*, *Hypocatharticus*, *Lapac'ticus*. A medicine which gently opens the bowels; such as tamarinds, manna, &c.

LAXATIVUS INDICUS, *Cambogia*.

LAXATOR AURIS INTERNUS, *L. tympani*.

LAXATOR TYMPANI, *L. major tympani*, *Exter'vus* seu *Anterior mall'ei*, *Oblit'vus* seu *Externus auris*, *Lazator auris inter'nus*, *Eustachii musculus*, (F.) *Antérieur du marteau*, *Acoustico-malléen*, *Spheni-malpingo-mallien*. A muscle which arises from the spine of the sphenoid bone and from the cartilage of the Eustachian tube, and is inserted, by a tendon, into the apophysis of Rau. It relaxes the *membrana tympani*.

LAXATOR TYMPANI MINOR. A very small muscle which extends from the upper part of the external auditory canal, and is inserted at the inferior part of the process of the handle of

the malleus. Its existence is denied by most anatomists.

LAXITAS, Atony.—1. Alvi, Diarrhœa.—1. Ingestorum, Lientery.—1. Intestinorum, Lientery.—1. Scroti, Rachosis.—1. Ventriculi, Gastrasthenia.

LAX'ITY, *Lax'itas*, *Lazness*. Same etymon. Condition of a tissue when loose or relaxed: or of one which wants tone. We say *laxity of fibre*, *laxity of skin*, to express that those parts have lost some of the tenseness proper to them.

LAYER, **ANIMAL**, see *Tache embryonnaire*—1. Mucous, see *Tache embryonnaire*—1. Serous, see *Tache embryonnaire*—1. Vascular, see *Tache embryonnaire*—1. Vegetative, see *Tache embryonnaire*.

LAYERS OF THE BLASTODERMA, see *Tache embryonnaire*.

LAYMAN, see *Laity*.

LAZARETTO, *Lazaret*, *Lazar-house*, from (I.) *lazzero*, 'a leper.' A solitary edifice in most seaports of magnitude, intended for the disinfection of men and goods proceeding from places where contagious diseases are prevailing.

LAZULITE, *Lapis lazuli*.

LEAD, **Plumbum**—1. Black, Graphites.—1. Chloride of, *Plumbi chloridum*—1. Colic, see *Colica metallica*—1. Iodide of, *Plumbi iodidum*—1. Nitrate of, *Plumbi nitras*—1. Oxyd of, semi-vitrified, *Plumbi oxydum semivitreum*.

LEAD PARALYSIS, see *Palsy*, lead.

LEAD POISONING, *Molybdo'sis*, *Molybdon'o-ous*, *Morbus plumb'eus*, *Cacochym'ia*, *plumbea*, (F.) *Intoxication saturnine*. Morbid phenomena induced by lead received into the system.

LEAD, RED, *Plumbi oxydum rubrum*.

LEAD RHEUMATISM, *Lead Neural'gia*, *Arthral'gia* of M. Tanquerel. The neuralgic and spasmodic pains caused by the poison of lead.

LEAD, SUCARBONATE OF, *Plumbi superacetas*—1. Tannate of, see *Tannin*.

LEAD WATER, *Liquor Plumbi subacetatis dilutus*.

LEAD, WHITE, *Plumbi subcarbonas*.

LEADER, *Tendon*.

LEADWORT, *Plumbago Europæa*.

LEAF, SOUR, *Andromeda arborea*.

LEAM'INGTON, MINERAL WATERS OF. Saline waters at Leamington, about two miles east of Warwick, England, which contain chloride of sodium, sulphate of soda, and chlorides of calcium and magnesium.

LEANNESS, *Emaciation*.

LEAP, *Sax. hlæpan*, *Saltus*, *Salit'io*, *Exaltis*, (F.) *Saut*; Bound, Jump,—the act of leaping. Muscular movement or movements, by which the body is detached from the soil by the forcible and sudden extension of the lower limbs, previously flexed upon the pelvis.

LEAPERY, *Lepira*.

LEAPING AGUE. This disease is said by the Scotch writers to be characterized by increased efficiency, but depraved direction, of the will, producing an irresistible propensity to dance, tumble, and move about in a fantastic manner, and often with far more than the natural vigour, activity, and precision! See *Mania*, dancing.

LEATHER FLOWER, *Clematis viorna*—1. Wood, *Dicra palustris*.

LEB'ANON, MINERAL WATERS OF. The spring at Lebanon, 26 miles east of Albany, New York, is a simple thermal; temperature 73° Fahr.

LECAINE, *Pelvis*.

LECHE, *Physician*.

LECHENEION, *Torcular Herophilii*.

LECHEROUS, *Libidinous*.

ECHO, *Puerpera*.

LECHOPYRA, *Puerperal fever*.

LECTHODES, *Vitelline*.

LECTITHOS, see *Ovum*.

LE CRAN, *Cochlearia armoracia*.

LECO, *Penis*.

LECONTIA VIRGINICA, *Peltandra Virginea*.

LECTISTERNIUM, from *lectus*, 'a bed,' and *sternere*, 'to spread.' The arrangement of a bed so as to adapt it to a particular disease. Also, a supplication, with the Romans, in times of public danger, when beds or couches were spread for the gods, as if they were about to feast, and their images were taken down from their pedestals and placed upon these couches around the altars. The lectisternium was first introduced in the time of a pestilence.—*Livy*.

LECTUAL'IS, from *lectus*, 'a bed.' An epithet applied to a protracted disease.

LECTULUS MEDICATUS, see *Fomentation*—1. *Stramineus*, *Fanon*.

LEDE SAUVAGE, *Ledum sylvestre*.

LEDOYEN'S DISINFECTING LIQUID or **FLUID**. A solution of nitrate of lead, (*Plumb. nitrat. 3j ad aquæ f 3j*), used as an antiseptic and antibromic.

LEDUM LATIFOLIUM, *Labrador' Tea*, *Ord. Ericaceæ*, grows in damp places, in Canada and the United States. The leaves have a pleasant odour and taste, and have been used as tea. They have also been esteemed pectoral and tonic.

LEDUM PALUS'TRÆ, *Rosmar'inus seu Anthos sylvest'ris*, *Marsh Tea*, (F.) *Lède* ou *Rosmarin sauvage*, *Olivier de Bohême*. *Nat. Ord. Ericinæ*. *Sex. Syst. Decandria Monogynia*. This plant has a bitter, substringent taste, and was formerly used in Switzerland in place of hops: the virtues are equivocal. The leaves are chewed by the Cree Indians, and applied to burns, which are said to heal rapidly under their influence. The cake of chewed leaves is left adhering to the sore until it falls off.—*Sir J. Richardson*.

LEE, *Feculence*, *Lixivium*, *Ley*, *Lye*, *Urine*.

LEECH, *Hirudo*, see *Ectozaa*, *Physician*.

LEECHCRAFT, *Medicina*.

LEEK, *Allium porrum*.

LEES, *SOAP*, *Liquor potassæ*.

LEG, (*Da. Lag.*) *Crus*, *Scelos*, *Onemē*. The portion of the lower extremity, which extends from the knee to the foot. It consists of three bones, *Tibia*, *Fibula*, and *Patella*, and also of a great number of muscles, vessels, and nerves. The projection, formed by the muscles at the back part of the leg, has received the name of *Calf of the leg*. It is the special attribute of man, and proves that he is destined to be biped.

LEG, SWELLED, *Phlegmatia dolens*—1. *Cochin*, see *Elephantiasis*.

LEGITIMATE, *Legit'imus*, from *lex*, *legis*, 'law,' *Genu'nus*, *Gne'sios*. An epithet applied to things which are according to rule. *A legitimate child*, (F.) *Enfant légitime*, is one conceived or born during marriage *Legitimate diseases*, (F.) *Maladies légitimes*, are those which follow a regular march.

LEGNA, from *λεγνν*, *λεγνον*, 'a fringed edge,' 'the border of a garment.' The orifice of the pudendum mulieb're, or of the uterus.

LÉGUME, *Legumen*.

LEGU'MEN, from *lego*, 'I gather,' [?] akin to *λαχαρον*, 'pulse,' (F.) *Légume*, *Gousse*. So called because it is usually gathered by the hand, instead of being reaped. All kinds of *pulse*, as peas, beans, &c., are thus termed.

LEGUMIN, *Casein*.

LEICHEN, *Lichen*.

LEIMANTHIUM VIRGINICUM, *Melan-thium Virginicum*.

LEIMAX, *Limax*.

LEIOPUS, *Li'opus*, *Planus*, *Plantus*, *Plotus*, from *λεω*, 'smooth,' and *πους*, 'a foot.' One who is affected with *flat-footedness*, *splay-footedness*. *Lecopodia*, *Liopodia*. One, the soles of whose feet are flat, instead of having the concavity which they commonly present.

LEIOSIS, Communion.

LEIPHÆMIA, *Liphæ'mia*: same etymon as *Leiphæmos*. Poverty or paucity of blood.

LEIPHÆMOI, Achroi.

LEIPHÆMOS, *Liphæ'mos*, (F.) *Leiphème*, from *λεω*, 'I want,' and *αἷμα*, 'blood.' A word sometimes used adjectively; at others, substantively, either for a vicious state of the blood—or rather for a sort of anemia—or for the patient who labours under this condition. See Achroi.

LEIPHÈME, *Leiphæmos*.

LEIPO, *Lipo*.

LEIPODERMIA, *Aposthia*.

LEIPODERMOS, *Lipoder'mos*, *Leipoderm'a-tus*, from *λεω*, 'I want,' and *δέρμα*, 'skin.' One who wants a part of his skin. It is especially applied to one who wants the prepuce. See *Apella*, and *Aposthia*.

LEIPOMERIA, *Lipomeria*.

LEIPOPSYCHIA, *Syncope*.

LEIPOTHYMIA, *Syncope*.

LEIPYRIAS, from *λεω*, 'I want,' and *πυρ*, 'fire,' or 'heat.' A species of continued fever, referred to by the Greek physicians, in which there is burning heat of the internal parts and coldness of the extremities. Avicenna described, under this name, a kind of hemitritea.

LEMA, *Chassie*.

LEME, *Lippitudo*.

LEMITHOCORTON, *Corallina Corsicana*.

LEMMA, *Cortex*, *Feculence*, *Furfur*, *Sedimentum*.

LEMNISCUS, *Pessary*, *Tent*. See *Laqueus*.

LEMON, *GROUND*, *Podophyllum montanum*—1. Juice, see *Citrus medica*—1. Juice, artificial, see *Citrus medica*—1. Peel, see *Citrus medica*—1. Tree, *Citrus medica*.

LEMON, *WILD*, *Podophyllum montanum*.

LEMONADE, *Limonada*, (F.) *Limonade*, *Citronnade*. Lemon juice diluted with water and sweetened. See *Citrus medica*.

LEMONADE, DRY, (F.) *Limonade sèche*. Citric or tartaric acid reduced to powder and mixed with sugar. *Lemonade powders* may be made as follows:—Pound $\frac{3}{4}$ of citric acid with a few drops of essence of lemon-peel and $\frac{3}{4}$ or more of lump sugar. Divide into six papers, each of which will make a glass of lemonade. See *Citric acid*.

Limonade Gazeuse (F.), is an agreeable drink prepared by adding syrup of lemons, raspberry, &c., to water saturated with carbonic acid.

LEMONADE, MAGNESIAN, *Magnesia citras*.

LEMONADE, NITRIC. Nitric acid considerably diluted with water, and sweetened.

LEMONADE, SULPHURIC, and **LEMONADE, TARTARIC**, are made with the sulphuric and tartaric acids.

LEMOSITAS, *Chassie*, *Lippitudo*.

LENDE, *Lumbi*.

LENDIS, *Lumbi*, *Nates*.

LENIENS, *Laxative*, *Lenitive*.

LENIS, *Lenitive*.

LENITIVE, *Lenitivus*, *Lenis*, *Leniens*, *Epiad-ma*, *Mitigans*, from *lenio*, *lenitum*, 'to assuage.' A medicine, which allays irritation or palliates disease; also, a laxative medicine. A *lenitive electuary* is one that purges gently.

LENOS, *Torcular Herophili*.

LENS, *Ervum lens*—1. *Crystalline*, *Crystalline*—1. *Esculenta*, *Ervum lens*.

LENTICULA. Dim. of *lens*, *lentic*, 'a lentil.' A freckle. Also, the eruption of lenticular fever. See *Ephelides*.

LENTICULAR GANGLION, *Ophthalmie G.* **LENTICULARES GLANDULÆ**, *Lenticular glands*. Mucous follicles, having the shape of a lentil, which are observed especially towards the base of the tongue.

LENTIGO, *Ephelides*.

LENTIL, *Ervum lens*.

LENTILLE, *Ervum lens*—1. *Crystalline*, *Crystalline*—1. *Ers*, *Ervum ervilia*.

LENTIS/CINUM VINUM. Wine impregnated with mastich; from *Lenticus*, 'the mastich tree.'

LENTISCUS VULGARIS, *Pistacia lentiscus*.

LENTITIA, *Lentor*.

LENTITIS, *Phacitis*.

LENTITUDO, *Lentor*.

LENTOR, *Lentit'ia*, *Lentitudo*, from *lentus*, 'clammy.' A viscosity or sizziness of any fluid. See *Gluten*.

LENT ROSE, *Narcissus pseudonarcissus*.

LENUM, *Torcular Herophili*.

LEONOTIS LEONURUS. *Ord. Labiatae*, A South African plant, which has a peculiar smell and nauseous taste, and is said to produce narcotic effects if incautiously used. It is employed in decoction in chronic cutaneous diseases. The Hottentots smoke it like tobacco, and take a decoction of its leaves as a strong cathartic. It is also given as an emenagogue. In the eastern districts of the Cape Colony, *Leonotis ovalis*, (Prov.) *Crowparenep*, *Male*, is used for the same purpose.

LEONOTIS OVATA, see *Leonotis leonurus*.

LEONTIASIS, *Leontion*, from *λεων*, *leontos*, 'a lion.' A name given to lepra of the face, from some fancied resemblance between the countenance of those labouring under it and that of the lion. To this kind of lepra the epithets *le'onine* and *le'ontine* have been given.

LEONTICE THALICTROIDES, *Caulophyllum thalictroides*.

LEONTION, *Leontiasis*.

LEONTODON TARAXACUM, *L. officinale* seu *vulgaris*, *Taraxacum officinale*, *Dens Leo'nis*, *Hedys'nois taraxacum*, *Urina'ria*, *Caput Mon'achi*, *Dan'delion*, *Piss-a-bed*, (Prov.) *Shitabed*, *Puffball*, (Sc.) *Dentiloun*, (F.) *Pissenlit*, *Liondent*, *Dent de lion*, *Couronne de Moine*, from *λεων*, *leontos*, 'a lion,' and *odon*, *odonos*, 'a tooth.' *Order*, *Compositae*. The young leaves are sometimes eaten as salad. The roots are, also, roasted and used as a substitute for coffee. The root, *Taraxacum*, (Ph. U. S.) is, moreover, reputed to be aperient and diuretic; hence its vulgar names. Its efficacy is doubtful.

LEONTOPODIUM, *Alchemilla*.

LEONU'RUS, from *λεων*, 'a lion,' and *ουρα*, 'a tail.' *Lion's Tail*.

LEONURUS CARDIACA, *Agripal'ma Gallis*, *Marrubium*, *Cardiaca crispia seu trilobata seu vulgaria*, *Herba cardiaca seu Patà lapi'næ*, *Motherwort*, *Throatwort*, (F.) *Agripaume*. *Ord. Labiatae*. Its properties are those of a nauseous bitter; and hence it has been used in hysteria and other nervous affections.

LEONURUS LANATUS, *Ballota lanata*.

LEOPARD'S BANE, *ROMAN*, *Arnica montana*, *Doronicum pardalianches*.

LEPAS, *Lepis*, *lepas*, gen. *λεπας*; *λεπς*, gen. *λεπας*, 'a scale.'

LEPER, see *Leprons*.

LEPER HOSPITAL, *Ludverie*.

LEPIA CAMPESTRIS, *Thlaspi campestre*—1. *Sativa*, *Lepidium Iberis*.

LEPIDIUM, from *λεπς*, *λεπας*, 'a scale;' so called from its supposed usefulness in cleansing the skin from scales and impurities. *Pepper-wort*. *Ord. Cruciferae*.

LEPIDIUM CAMPESTRE, *Thlaspi campestre*.

LEPIDIUM IBE'RI'S, *Ibe'ris*, *Cardaman'tica*, *Sci-at'ica crensca*. This plant possesses a warm, penetrating, pungent taste, like other cresses, and is recommended as an antiscorbutic, antiseptic, and stomachic.

LEPIDIUM SATI'VUM, *Lep'ia sati'va*, *Thlaspi nasturtium* seu *sati'vum*, *Nasturtium horten'se* seu *sati'vum*. Garden cress, *Dittander*, *Green mustard*, (F.) *Cresson alénois*, *Cresson des Jardins*. This plant possesses warm, stimulating properties, and is used like the last.

LEPIDIUM SQUAMATUM, *Cochlearia coronopus*.

LEPIDODES, Squamous.

LEPIDOIDES, Squamous.

LEPIDOSARCO'MA, *Tumor squamifor'mis carno'sus*. A fleshy tumour, covered with scales; from *λεπς*, *lepidos*, 'a scale,' and *σarkoma*, 'a fleshy tumour.' Marcus Aurelius Severinus describes tumours of this kind in the interior of the mouth.

LEPIDOSIS, Scaly disease.

LEPIDOSIS ICHTHYIASIS, *Ichthyosis*—1. *Ichthyiasis cornigera*, Horny excrescences—1. *Lepriasis*, *Lepra*—1. *Pityriasis*, *Pityriasis*—1. *Psoriasis*, *Psoriasis*.

LEPIDOTIS CLAVATA, *Lycopodium clavatum*.

LEPIRA, *Lepra*.

LEPIS, Scale.

LEPISMA, Squamous portion.

LEPORINUM LABIUM, *Harelip*—1. *Ros-trum*, *Harelip*.

LEPORINUS OCULUS, *Lagophthalmia*.

LEPOSTEOPHYTON; from *λεπς*, 'a scale,' *οστέον*, 'a bone,' and *φυτον*, 'a plant or growth.' A thin scale of bone, the result of a morbid growth.

LEPRA, *Lep'ra*, from *λεπρος*, 'scaly.' *Lepido'sis Lepri'asis*, *Lepro'sis*, *Lepro'sitas*, *Vitili'go*, *Lepra Græcorum*, *Herpes furfuraceus circin'atus*, *Leprosy*, *Leap'ery*, (Sc.) *Lepyr*, *Lipper*, (F.) *Lèpre*, *Mal Saint-Main*, *Dartre furfuracée arandonie*. This term has been applied to various affections, very different in character. 1. To the *Leprosy of the Jews*, *Leucé*, *Lepido'sis Lepriasis canescens*, *Lepra Moa'ica* seu *Hebræorum*,—a variety of the *Alphos* or *Lepra alphi'des*. The leucæ was, generally, not scaly, but consisted of smooth, shining patches, on which the hair turned white and silky, and the skin, with the muscular flesh, lost its sensibility. It was incurable. 2. To the *Elephantiasis* or *Lepra of the Arabs*, see *Elephantiasis*; and 3. To the *Lepra of the Greeks*, which includes all the varieties met with at the present day. It is characterized by scaly patches of different sizes, but having always nearly a circular form. Bateman and Willan describe three chief varieties of this lepra.

1. *Lepra alphi'des*, *Lepido'sis Lepri'asis al-bida*, *Alphos*, *Morphæ'a alba*, *Vitili'go albus*, *Albaras alba*, *Albaræ*, *Albaros*, *Lèpre écailluse* of Alibert, *White leprosy*. An affection, characterized by white patches, surrounded by a rose-coloured areola, which appears here and there on the surface; depressed in the middle.

2. *Lepra nigricans* seu *melas seu maculo'sa nigra*, *Vitili'go melas seu nigra*, *Morphæ'a seu Albaras nigra*, *Melas*, *Lepido'sis lepri'asis nigricans*, *Black leprosy*; in which the scales are livid; the size of half a dollar; and diffused over the body, but less widely than in the *Alphi'des*. The French pathologists usually admit three species of lepra, to which they give the epithets *scaly*, (*squameuse*), *crustaceous*, (*crustacée*), and *tubercular*, (*tuberculeuse*), according as the skin may be covered with scales, crusts, or tubercles.

3. *Lepra vulga'ris*, *Lepido'sis lepri'asis vulga'ris*, *Dartre furfuracée arandonie*, of Alibert, characterized as follows: scales glabrous, whitish,

size of a crown piece; preceded by smaller reddish and glossy elevations of the skin, encircled by a dry, red, and slightly elevated border; often confluent; sometimes covering the whole of the body except the face.

Lepra appears to be endemic in Egypt, in Java, and certain parts of Norway and Sweden. Imperfect and faulty nutriment appears to contribute to its development. The means, best adapted for its removal, are:—a mild, unirritating diet, emollient fomentations—sulphureous baths, fumigations, &c; but, often, all remedial agents will be found ineffectual.

LEPRA ARABUM, *Elephantiasis Arabica*—1. *As-turica*, see *Pellagra*—1. *Borealis*, *Radzyge*—1. *Fungifera*, *Frambosia*—1. *Gangrænosa*, *Ngeren-gere*—1. *Græcorum*, *Lepra*—1. *Hebræorum*, see *Lepra*—1. *Ichthyosis*, *Ichthyosis*—1. *Lombardica*, *Pellagra*—1. *Maculosa nigra*, *L. Nigricans*—1. *Mediolanensis*, *Pellagra*—1. *Melas*, *Lepra nigricans*—1. *Mercurial*, *Eczema mercuriale*—1. *Mosaica*, see *Lepra*—1. *Norvegica*, *Radzyge*—1. *Squamosa*, *Impetigo*—1. *Taurica*, *Mal de Crimée*.

LÈPRE, *Lepra*—1. *des Cosaques*, *Mal de Crimée*—1. *Écailluse*, *Lepra alphi'des*—1. *Humide*, *Impetigo*.

LÉPREUX, *Leprous*.

LEPRICUS, *Leprous*.

LEPROSARIUM, *Ladrerie*.

LÉPROSERIE, *Ladrerie*.

LEPROSIS, *Lepra*.

LEPROSITAS, *Lepra*.

LEPROSUS, *Leprous*.

LEPROSY, *Lepra*—1. *Black*, *Lepra nigricans*—1. *Norwegian*, see *Radzyge*—1. *White*, *Lepra alphi'des*.

LEPROUS, *Lepro'sus*, *Lepro'ticus*, *Lep'ricus*, *Lepro'des*, (Sc.) *Lipper*, (F.) *Lépreux*, *Ladre*, same etymon. Relating to or resembling or affected with leprosy; a *leper*.

LEPSIS, *Attack*.

LEPTANDRA VIRGINICA, *Veroni'ca* seu *Padero'ta Virginica*, *Culver's physic*. *Ord. Scrophulariaceæ*. An indigenous plant, which grows throughout the United States, and flowers in August. The flowers are white, and terminate the stem in a long spike. A variety with purple flowers has been described, as

LEPTANDRA PURPUREA, *Physic root*, *Black root*, *Whorlywort*, *Culver's root*, *Brinton root*, *Bowman root*. The root is bitter and nauseous, and when fresh is emetic and cathartic. In the dried state it is more uncertain. Dose of the powder, gr. xx to ʒj.

LEPTO, **LEPTOS**, *λεπτος*, 'thin,' 'light.'

Hence:

LEPTOCEPHA'LIA, from *λεπτος*, 'thin, fine,' and *κεφαλή*, 'head.' Monstrosity, in which the head is abnormally small.

LEPTOCHRO'A, *Leptochros*, from *λεπτος*, 'thin, fine,' and *χρoα*, *χρoα*, 'the colour of the skin: the skin.' Fineness, thinness of skin.

LEPTOCHYM'IA, from *λεπτος*, 'thin,' and *χυμος*, 'a juice.' Morbid thinness of the juices.

LEPTOHYME'NIA, from *λεπτος*, 'thin,' and *μνη*, 'a membrane.' Thinness, delicacy of membrane.

LEPTOMER'IA, from *λεπτος*, 'thin, fine,' and *μερος*, 'a part.' Fineness, delicacy of bodily formation.

LEPTONTHIQUES, *Attenuans*.

LEPTOPHONIA, *Oxyphonia*.

LEPTOSPERMUM LEUCADENDRUM, *Melaleuca cajuputi*.

LEPTOTHRIX, *Leptotrichus*.

LEPTOTRICH'IA; same etymon as the next. Fineness of hair.

LEPTOTRICHUS, *Leptothrix*, from *λεπτος*,

'fine,' and *δοξ, τριχος*, 'hair.' One who has fine hair.

LEPTOTROPHIA, *Microtroph'ia*, *Microtra-ph'ia*, from *λεπτος*, 'light,' and *τροφη*, 'nourishment.' Light nutrition.

LEPTYNSIS, Emaciation.

LEPTYNTICA, Attenuants.

LEPTYSMUS, Emaciation.

LEPYR, Lepra.

LEREMA, see Dementia.

LERESIS, see Dementia.

LERUS, Delirium, see Dementia.

LESEOLI MORBUS, Icterus.

LESEOLUS. According to Paracelsus, a perfectly transparent salt, which cured jaundice: hence *Leseoli morbus*.

LESION, *La'sio*, from *laedere*, *laesum*, 'to injure.' Derangement, disorder; any morbid change, either in the exercise of functions or in the texture of organs. *Organic lesion* is synonymous with *organic disease*.

LESION OF CONTINUITY, see Continuity, solution of.

LESK, Inguen.

LESSIVE, Lixivium.

LESSIVE DES SAVONNIERS, Liquor potassæ. Also, a solution of caustic soda in water, containing about 3 parts of soda to 8 of water.

LESSIVE DE TARTRE, Liquor potassæ subcarbonatis.

LETALITAS, Mortality.

LETHALIS, Lethiferous, Mortal.

LETHALITAS, Mortality.

LETHARGIC, *Lethargicus*, *Veterno'us*, (F.)

Léthargique. Relating to lethargy; affected with lethargy.

LÉTHARGIQUE, Lethargic.

LETH'ARGY, *Lethar'gia*, *Lethar'gus*, *Carnus lethargus*, *Veter'nus*, *Obliv'io iners*, from *ληθη*, 'oblivion,' and *αργος*, 'active.' A constant state of stupor from which it is almost impossible to arouse the individual; and, if aroused, he speedily relapses into his former condition.

LETHEA, Papaver.

LE'THEON, from *ληθη*, 'oblivion.' A name given by some to sulphuric ether, when inhaled as an anæsthetic agent.

LETHIFEROUS, *Le'thifer*, *Letha'lis*, *Lethif'icus*, *Mort'ifer*, *Mortif'erous*, from *ληθη*, 'death,' and *φερω*, 'I bear.' Death-bearing; deadly.

LETHUM, Death.

LETTUCE, *Lactuca*—1. Blue, *Mulgedium acuminatum*—1. False, *Mulgedium Floridanum*—1. Indian, see Calumba.

LETTUCE OPIUM, see *Lactuca*.

LETTUCE, STRONG-SCENTED, *Lactuca virosa*—1. White, *Nabulus albus*—1. Wild, *Lactuca elongata*.

LETUM, Death.

LEUCADES, see Sclerotic.

LEUCÆMIA, *Leukæ'mia*, *Leuchæ'mia*, from *λευκος*, 'white,' and *αἷμα*, 'blood,' (F.) *Leucæmie*. A condition of the blood in which it is deficient in colouring matter. See *Leucocythæmia*.

LEUCÆTHIOPIA, see Albino.

LEUCÆTHIOPS, Albino.

LEUCANIA, *Laucania*.

LEUCANTHEMUM, *Anthemis nobilis*, *Matri-caria chamomilla*—1. Vulgare, *Chrysanthemum eucanthemum*.

LEUCE, Lepra (of the Jews.)

LEUCELECTRUM, *Succinum (album)*.

LEUCÉMIE, *Leucæmia*.

LEUCÆMIA, *Leucæmia*.

LEUCITIS, Sclerotitis.

LEUCOCYTE, see Globules of the blood.

LEUCOCYTHÆMIA, 'white cell blood,' from *λευκος*, 'white,' *κυτος*, 'cell,' and *αἷμα*, 'blood.' A condition of the blood, which consists in a su-

perabundant development of the white corpuscles, a disease which has been observed at times to be accompanied by enlargement of the spleen and liver, and at others by increased size of the lymphatic glands. See *Leucæmia*.

LEUCODENDRON, *Melaleuca cajuputi*.

LEUCENUS, see Wine.

LEUCOIUM, *Lunaria rediviva*—1. *Luteum*, *Cheiranthus cheiri*.

LEUCOLACHANUM, *Valeriana*.

LEUCOLEIN, *Leukoleinum*.

LEUCOMA, from *λευκος*, 'white.' *Oculo'rum albu'go*, *Leucom'ma*, *Leuco'sis*, *Albu'go*, *Al'bulu*, *Ceratoleucoma*, *Ephel'otes*. *Leucoma* and *Albugo* are often used synonymously to denote a white opacity of the cornea. Both are essentially different from nebula of the cornea; nebula being usually the result of chronic ophthalmia and an effusion of a milky serum into the texture of the delicate continuation of the conjunctiva over the cornea;—the others are the result of violent, acute ophthalmia. In this state, a thick, coagulable lymph is extravasated from the arteries, sometimes superficially, at other times deeply into the substance of the cornea. On other occasions, the disease consists of a firm, callous cicatrix on this membrane,—the effect of a wound or ulcer with loss of substance. The affection is more difficult of cure in proportion to its duration and to the age of the individual; the activity of the absorbents being greater in youth. If inflammation still exist, antiphlogistics must be persevered in, and, afterwards, gentle stimulants be used to excite absorption; along with the internal use of mercury or iodine.

LEUCOMA, Albumen—1. *Gerontotoxon*, *Gero-tonox*—1. *Margaritaceum*, see *Margaritaceus*.

LEUCOMMA, *Leucoma*.

LEUCOMORIA, *Cuttubuth*.

LEUCONECROSIS, from *λευκος*, 'white,' and *νεκρωσις*, 'death.' A form of dry gangrene,—the opposite in appearance to anthraconecrosis.

LEUCONYMPHÆA, *Nymphæa alba*.

LEUCOPATHIA, see Albino, Chlorosis—1. *Partialis acquisita*, *Achroma*.

LEUCOPHAGIUM, *Blanc-manger*.

LEUCOPHLEGMASIA, *Leucophlegmatia*—1. *Dolens puerperarum*, *Plegmatia dolens*.

LEUCOPHLEGMATIA, *Leucophlegmatia*, *Hydrops leucophlegmatia*, *Tumescen'tiu piti-t'o'sa*, from *λευκος*, 'white,' and *φλεγμα*, *φλεγματος*, 'phlegm.' A dropsical habit, *Hydrocoachex'ia*. Some writers use the word synonymously with *anasarca* and *oedema*; others with *emphysema*.

LEUCOPHLEGMATIA ÆTHIOPUM, *Chthonophagia*.

LEUCOPIPER, *Piper album*.

LEUCOPYRIA, *Hætic fever*.

LEUCORRHÆE, *Leucorrhœa*.

LEUCORRHŒA, *Fluxus seu Fluor mulie'bris*, *Proflu'vium mulie'brè*, *Cursus matric'is*, *Fluxus matric'is*, *Elytr'itis*, *Coloos'itis*, *Colp'itis*, *Destilla'tio uteri*, *Fluxio alba seu mul'va*, *Uleus uteri*, *Catar'hus genita'lium*, *Hysterorrhœa mucosa*, *Cutame'nia alba*, *Menses albi*, *Men'strua alba*, *Menorrhag'ia alba*, *Fluor mulie'bris non Gal'licus*, *Blennelytr'ia*, *Gonorrhœa benigna notha inveterata*, *Purgatio mulie'bris alba*, *Alba purgan'tia*, *Cachex'ia uterina*, *Rheuma uteri*, *Uteri Cory'za*, *Medorrhœa femina'rum insons*, *Blennorrhœa seu Blennorrhag'ia genita'lium*, *Edvoblenorrhœa seu Medoblenorrhœa femina'rum*; *The whites*; (F.) *Fleurs ou Fluens blanches*, *Pertes blanches*, *Écoulement blanc*, *Catarrhe utérin*, *Perte utérine blanche*, from *λευκος*, 'white,' and *ρρω*, 'I flow.' A more or less abundant discharge of a white, yellowish, or greenish mucus; resulting from acute or chronic inflammation, or from irritation of the membrane lining the genital

organs of the female. *Vaginal Leucorrhœa* has been termed *Blennorrhœa seu Fluor albus vaginæ*, *Leucorrhœa*, *Medorrhœa vagina*, *Vaginitis*, *Elythroblennorrhœa*, *Colporrhœa*.—*Uterine Leucorrhœa* has received the names *Fluor albus uteri*, *Leucorrhœa seu Medorrhœa uteri*, *Metroblennorrhœa*, *Blennometritis*, *Metroblennorrhœa*, *Metrocystitis*, *Metroleucorrhœa*, (F.) *Leucorrhœa uterine*, *Catarrhe utérin*. It is often attended with pain and a sense of heaviness in the loins, abdomen and thighs; disordered digestive functions, &c., so that, at times, the health suffers largely, although there are few females who are not occasionally subject to moderate leucorrhœa. Attention to the general health, change of air, keeping up a perspirable state of the surface by flannel worn next the skin, the horizontal posture, &c., do more than pharmaceutical agents; which are almost entirely confined to astringent injections. These may be employed, when the discharge is so great as to require them.

LEUCORRHEA ANALIS, see Hæmorrhoids — 1.

Nabothi, see Parturition.

LEUCORRHOIS, see Hæmorrhoids.

LEUCOSSES; from *λευκος*, 'white.' In the nosology of M. Alibert, all the diseases of the lymphatic apparatus. The 7th family in his *Nosologie*. In that of Fuchs, it is a family of diseases, (G.) *Leukosen*, which includes the various forms of anæmia.

Leucosis, Leucoma, Paleness. See Albino.

LEUK or LOCHEE, MINERAL WATERS OF. Saline, chalybeate waters, about six leagues distant from Sion. They contain chloride of sodium, with a little sulphate of magnesia; sulphate of lime, carbonate of magnesia, carbonic acid, and protoxide of iron.

LEUKÆMIA, Leucæmia.

LEUKOLEIN'UM, *Chinolein'um*, *Leukol*, *Leu'colein*, *Leu'colein*, *Chi'nolein*, *Chi'nolin*. This substance is the product of the dry distillation of coal; mixed with picolin, anilin and other substances, in mineral tar. It is procured, also, by heating *quinia*, *cinchonia* and *strychnia*, with as concentrated a ley of *potassa* as can be made. Its specific gravity is 1.081; and it is slightly soluble in water, and miscible in all proportions with alcohol, ether and essential oils.

LEVAIN, Ferment.

LEVATOR, Elevator.

LEVATOR AN'GULI ORIS, *Abducens labiorum*, *Eleva'tor labiorum commun'is*, *Caninus*, (F.) *Sus maxillo-labial*, *Petit sus-maxillo-labial* (Ch.), *Muscle canin*. A small, flat, long, quadrilateral muscle, which arises from the fossa canina, and is inserted at the commissure of the lips, where it is confounded with the triangularis. It raises the corner of the mouth, and draws it towards the ear.

LEVATOR ANI, *Levator magnus seu internus*, *Latus ani*, *Eleva'tor ani*, *Sedem attolens*, (F.) *Pubo-coccygien annulaire*, *Sous pubio-coccygien* (Ch.), *Releveur de l'anus*. A muscle, situate at the lower part of the pelvis. It is broad, flat, quadrilateral, and broader above than below. It represents a kind of membranous partition, which closes the outlet of the pelvis, and the upper concavity of which is opposed to that of the diaphragm. It is attached, above, to the posterior surface of the body of the pubis, to the upper part of the obturator foramen, and to the spine of the ischium; and is inserted into the coccyx, into an aponeurotic line common to it and its fellow, and into the lateral parts of the rectum. This muscle supports the rectum; raises it, and carries it upwards during the evacuation of the excrement. It can, also, compress the bladder and vesiculæ seminales, and thus favour the expulsion of the urine and sperm.

LEVATOR ANI PARVUS. Transversus perinæi — 1. *Auris*, *Attollens aurem* — 1. *Coccygis*, *Coccygeus*.

LEVATOR GLAN'DULÆ THYROIDEÆ. A muscle occasionally found connected with the upper border or isthmus of the thyroid gland; and attached superiorly to the body of the os hyoides, or to the thyroid cartilage.

LEVATOR MENTI, *Levator labii inferioris* — 1. *Oculi*, *Rectus superior oculi* — 1. *Proprius scapulæ*, *L. scapulæ*.

LEVATOR LABII INFERIO'NIS, *L. menti*, *Elevator labii inferioris seu labii inferioris proprius*, *Incisivus inferior*, *Musculus penicillatus*, (F.) *Houppé du menton*, *Releveur de la lèvre inférieure ou du menton*. A portion of the mento-labial of Chaussier. A small muscle situate before the symphysis menti. It is thick, conical; and attached by its apex to a fossette at the side of the symphysis in the inferior maxillary bone. Its fibres proceed diverging and vanishing in the manner of a tuft, (F.) *Houppé*, on the skin of the chin. This muscle raises the chin, and pushes upwards the lower lip.

LEVATOR LABII SUPERIO'NIS ALÆQUE NASI, *Incisivus lateralis et pyramidalis*, (F.) *Grand sus-maxillo-labial* (Ch.), *Élévateur commun de l'aile du nez et de la lèvre supérieure*. This muscle is a fleshy, thin, triangular bundle, situate at the sides of the nose. It arises from the ascending process of the superior maxillary bone: thence its fibres descend in a diverging manner, a part being inserted into the ala of the nose, and a part losing themselves in the upper lip. This muscle raises the upper lip and ala nasi, which it draws a little outwards.

LEVATOR LABII SUPERIORIS PROPRIUS, *Musculus incisivus*, *Elevator labii superioris proprius*, (F.) *Moyen sus-maxillo-labial*, (Ch.), *Orbito-maxillo-labial*. This thin, flat, quadrilateral muscle is situate at the middle and inner part of the face. It arises from the os malæ and the os maxillare superius, and is inserted into the upper lip, which it raises, at the same time carrying it a little outwards.

LEVATOR PALA'TI, *L. Palati Mollis*, *Petrosalpin'go-staphylinus*, *Salpin'go-staphylinus internus*, *Salpingo-staphylinus*, *Pter'gyo-staphylinus externus*, *Spheno-staphylinus*, *Spheno-palati-nus*, *Peristaphylinus internus superior*, *Pétro-staphylin*, (Ch.). This muscle is long, narrow, and almost round above; broader and flatter inferiorly. It arises from the lower surface of the *pars petrosa*; from the cartilage of the Eustachian-tube; and is inserted into the substance of the velum palati. Its use is to raise the velum palati.

LEVATOR PAL'PEBRÆ SUPERIO'NIS, *Palpebræ superioris primus*, *Aperitor Oculi*, *Aperiens Palpebrarum rectus*, *Recl'isor palpebrarum*, (F.) *Orbito-palpebral* (Ch.), *Orbito-sus-palpebral*, *Élévateur de la paupière supérieure*. A long, small, thin muscle, situate at the upper part of the orbital cavity. By its posterior extremity it is inserted into the little ala of the sphenoid bone, immediately in front of the foramen opticum, and, by its anterior extremity, which is expanded, it terminates at the upper margin of the tarsal cartilage of the eyelid. This muscle raises the upper eyelid, draws it backwards, and sinks it into the orbit.

LEVATOR PROS'TATÆ, (F.) *Releveur de la prostate*. Santorini has given this name to the anterior fibres of the levator ani, which embrace the prostate.

LEVATOR SCAP'ULÆ, *Eleva'tor scapulæ*, *E. seu Musculus Potent'ia*, *Angula'ris*, *Levator proprius scapulæ*, — (F.) *Trachélo-scapulaire* (Ch.), *Angulaire de l'omoplate*, *Releveur de l'omoplate*.

This muscle is so called, because it is attached to the upper and internal angle of the scapula. It is situated at the posterior and lateral part of the neck, and at the upper part of the back. It is long, flat, and broader above than below. In this latter direction, it is inserted into the superior internal angle of the scapula; and, in the former, into the tops of the transverse processes of the first four cervical vertebrae. It depresses the prominence of the shoulder, by raising the posterior angle of the scapula, on which it impresses a kind of rotary motion. It can, also, draw the head to one side.

LEVATOR URETHRAE, (F.) *Releveur de l'Urethre*. Santorini describes under this name a portion of the transversus perinaei muscle.

LEVATORES COSTARUM, Supracostales.

LEVEN, Ferment.

LEVER, from *levare*, 'to lift up.' *Vectis*, *Mochlus*, *Porrec'tum*. An inflexible rod, turning round a fixed point, and used for moving bodies, bearing burdens, or raising them. The point on which the lever moves is called the *Fulcrum*, *Hypomochlion*. The force which moves the lever is called the *power*; and the weight to be moved, the *resistance*. There are three kinds of levers. A lever of the first kind has the fulcrum between the power and resistance. A lever of the second kind has the resistance between the fulcrum and power; whilst a lever of the third kind has the power between the fulcrum and resistance. In the locomotive system of the human body, we have examples of all the three kinds. The bones represent *levers*; the muscles of locomotion are *powers*; the weight of parts to be moved constitutes the *resistance*. The *fulcra*, are, at times, the joints; at others, the ground, &c. The head moves or the neck, as a lever of the first kind; the first cervical vertebra forming the fulcrum. We rise on tip-toe by a lever of the second kind, the fulcrum being the ground under the toes; and we have examples of a lever of the third kind in the flexion of the fore-arm on the arm, in the elevation of the arm, &c.

LEVER, *Hystero-mochlium*, *Mochlis'eus*, *Vectis obstetric' ius*, *Vectis*, (F.) *Levier*, is an instrument curved at the extremity, and having a fenestra. It is used to assist the extraction of the child's head, when instrumental aid is necessary. Levers are, also, used by the dentist for extracting stumps, &c. The *Levier de l'Écluse*, *Langue de Carpe*, *Trivelin* or *Punch*, is employed for extracting the molar teeth.

LEVIATHAN PENIS, *Pria'pus Ceti*, *Bale'nae*. The penis of the whale. This singular medicine was, at one time, given in powder, in cases of dysentery and leucorrhœa.

LEVIER, Lever—*l. de l'Écluse*, see Lever.

LEVIGATION, *Laviga'tio*, from *lavigare*, *laviga'tum*, (*lavis*, 'smooth,') 'to polish.' *Porphyrazia*. An operation, by which bodies are reduced to very fine powder. It is performed by putting substances, already pulverized, into water; the coarser parts are not long in being deposited, whilst the finer molecules remain suspended in the water. The liquor is decanted into another vessel, and suffered to remain at rest, until the fine particles are collected at the bottom. The fluid part is then separated by decantation.

LEVISTICUM, *Ligusticum levisticum*.

LÈVRE, Lip.

LÈVRES GRANDES, *Labia pudendi*—*l. Petites*, *Nymphæ*—*l. de la Vulve*, *Labia pudendi*.

LEVURE, *Yeast*—*l. de la Bière*, *Yest*.

LEXIPHARMACUS, *Alexipharmic*.

LEXIPYRETICUS, *Febrifuge*.

LEY, *Lixivium*, *Lye*—*l. Soap*, *Liquor potassæ*.

LEYS'SERA GNAPHALOIDES. A South

African plant, *Nat. Ord.* *Compositæ*, which is emollient, and highly recommended at the Cape, in catarrh, cough, and even in phthisis.

LEZARD, Lizard.

LIAN À RÉGLISSE, *Abrus precatorius*.

LIA'TRIS SPICATA, *Gayfeather*, *Button snakeroot*. *Ord.* *Compositæ*. An indigenous plant, growing in meadows and moist grounds in the Middle and Southern States. Its beautiful purple compound flowers are in a spike. They appear in August. The root has been considered diuretic.

LIA'TRIS SCARIO'SA and *L. Squarro'sa*, *Throatwort*, *Sow-wort*, *Backache root*, *Devilsbite*, *Blazing star*, *Prairie pines*, *Rough root*, are called from their reputed powers in bites from the rattlesnake, *Rattlesnake's master*. The roots, bruised, are applied to the wound, and the decoction, in milk, is given internally.

LIBANOTIS ANNUA, *Athamanta cretensis*—*l. Coronaria*, *Rosmarinus*—*l. Cretensis*, *Athamanta cretensis*—*l. Hirsuta*, *Athamanta cretensis*.

LIBANOTUS, see *Juniperus lycia*.

LIBANUS, *Juniperus lycia*.

LIBER PHARMACEUTICUS, *Dispensatorium*.

LIBIDINIS SEDES, *Clitoris*.

LIBIDINOSUS, *Libidinous*.

LIBIDINOUS, *Libidinosus*, *Lascivus*, *Veneriv'agus*; from *libido*, 'lust,' *Lecher'ous*, *Lick'orous*. *Lewd*: *lustful*.

LIBIDO. Desire, necessity. Authors speak of *Libido urinae*, *Libido intestinali*. Some employ it synonymously with *Prurigo*; others, with *Salac'itas*, *Lubi'do*, *Himeros*.

LIBOS, from *λεῖω*, 'to drop.' A defuxion from the eyes.—*Galen*.

LIBRA, *Pound*.

LICHANOS, *Index*, see *Digitus*.

LICHE, *Body*.

LICHEN, *λεῖχην* or *λεχην*, (pronounced *li'ken*), *Exor'mia Lichen*, *Leichen*, *Serpi'go*, *Volat'ica*, *Pap'ulae*, *P. sicca*, *Peti'go*, *Pustula sicca*, *Scab'ies sicca* seu *a'gria*, *Lichen'asis adult'orum*, *Li'chenous rash*, (F.) *Gale sèche*, *Dartre furfuracée volante*, *Poussée*. The cutaneous affection described under this name by the Greek writers, is not clearly defined. Some have believed it to be *Impetigo*, but this is doubtful. The name is, now, generally applied to a diffuse eruption of red pimples, accompanied by a troublesome sense of tingling or pricking. Drs. Willan and Bateman define it,—"an extensive eruption of papulae affecting adults, connected with internal disorder, usually terminating in scurf; recurrent, not contagious." One of their varieties, however, the *Lichen Tropicus*, does not accord well with this definition; for it affects children as well as adults, and is unconnected with internal disorder.

LICHEN A'GRISUS, *L. A'gria*, *Exor'mia Lichen ferus*, *Lichen ferus*, *Pap'ula a'gria*, *Wild Lichen*, is distinguished by pimples in clusters or patches, surrounded by a red halo; the cuticle growing gradually harsh, thickened, and chappy, often preceded by general irritation. In addition to antiphlogistics, a cooling ointment may be used, to allay itching.

LICHEN BARBATUS PLICATUS, *L. plicatus*.

LICHEN CIRCUMSCRITUS is characterized by clusters or patches of papulae, having a well defined margin, and an irregularly circular form; continuing for six or eight weeks. These varieties require but little medical treatment. The antiphlogistic plan is all that is necessary.

LICHEN FERUS, *Lichen agrius*.

LICHEN LIVIDUS. The papulae have a dark red or livid hue, without any fever. They are

more permanent in this variety. It requires the mineral acids and bark.

LICHEN PILARIS, *Exor'mia Lichen pilaris*, is merely a modification of the preceding; the papule appearing at the roots of the hair.

LICHEN SIMPLEX, *Exor'mia Lichen simplex*, consists of an eruption of red papule, appearing first on the face or arms, and afterwards extending over the body; preceded for a few days by slight febrile irritation, which usually ceases when the eruption appears,—with an unpleasant sense of tingling during the night. It generally dies away in ten days or a fortnight.

LICHEN TROPICUS, *Exor'mia Lichen tropicus*, *Ecce'sma*, *Es'sera* [?], *Sudam'ina* [?], *Prickly Heat*, *Summer Rash*. The pimples are bright red, and of the size of a small pin's head; with heat, itching, and pricking, as if by needles. It is local; produced by excessive heat; and disappears when the weather becomes cooler, or the individual is inured to the climate.

LICHEN URTICATUS, *L. Urtico'sus*, *Exor'mia Lichen urtico'sus*, is another variety. The Nettle Lichen consists of papule, accompanied by wheals like those of nettle rash. A variety of this is said to be called, in America, *blights*.

LICHEN, WILD, *Lichen agrius*.

LICHEN APHTHOSUS, *Pelti'dea aphtho'sa*; *Ord. Lichenaceæ*; is said to be cathartic and anthelmintic.

LICHEN ARBORUM, *Lichen pulmonarius*—*l. Barbatus plicatus*, *Lichen plicatus*—*l. Blanc de Neige*, *L. caninus*—*l. Boëtier*, *L. pyxidatus*.

LICHEN CANINUS seu *spu'rius* seu *terres'tris* seu *veno'sus* seu *ciner'eus terres'tris*, *Muscus can'inus*, *Phys'cia niva'lis*, *Peltig'era can'ina*, *Pelti'dea can'ina* seu *leucor'rhiza* seu *mala'cea* seu *amplis'sima* seu *spu'ria*, *Ash-coloured Ground Liverwort*, (F.) *Lichen contre-rage*, *Lichen blanc de neige*. This cryptogamous plant was, for a long time, considered capable of preventing and curing *Rabies canina*. It has, also, been used in mania and in spasmodic asthma.

LICHEN CINEREUS TERRESTRIS, *L. caninus*—*l. Carrageen*, *Fucus crispus*—*l. Cocciiferus*, *L. pyxidatus*—*l. contre Rage*, *L. caninus*—*l. Entonnoir*, *L. pyxidatus*—*l. Eryngifolius*, *L. islandicus*—*l. Floridus hirtus*, *L. plicatus*—*l. Hirtus*, *L. plicatus*.

LICHEN ISLANDICUS, *L. eryngifo'lius*, *Lichenoides Island'icum*, *Loba'ria Island'ica*, *Muscus Islandicus* seu *cathart'icus*, *Clado'nia* seu *Phys'cia* seu *Cetra'ria* seu *Parme'lia Island'ica*, *Ice-land Lichen* or *Liverwort* or *Moss*, (F.) *Lichen d'Islande*, *Mousse marine perlée*, *Mousse d'Islande*. This plant is inodorous, with a bitter and mucilaginous taste. It is esteemed to be tonic, demulcent, and nutrient. Dose, ʒj to ʒiv, being first steeped in water holding in solution some carbonate of potassa to extract the bitter, and then boiled in milk. A bitter principle has been extracted from it, termed *Cetrarin*, *Cetrari'num*, *Cetra'rium*, which has been given in intermittents.

LICHEN LACINIATUS, *L. saxatilis*—*l. Parietinus*, *Parmelia parietina*.

LICHEN PLICATUS, *L. hirtus* seu *barba'tus plicatus* seu *Floridus hirtus*, *Parme'lia plica'ta*, *Usne'a plica'ta*, *U. Flor'idia hirta* seu *hirta*, *Muscus arbo'reus* seu *albus* seu *quernus*, *Querci'ni Herba*. This plant is applied, by the Laplanders, as an astringent, to bleeding vessels; and to parts which are excoriated after long journeys.

LICHEN PULMONARIUS, *L. arborum* seu *reticulatus*, *Parme'lia pulmona'cea*, *Reti'cula'ria officina'lis*, *Muscus pulmonari'us querci'ni*, *Pulmona'ria arbo'rea*, *Loba'ria pulmonaria*, *Sticta pulmona'cea*, *Onk Lungs*, *Tree Lungwort*, *Hazel Crotches*, *Hazeltree*, (F.) *Lichen pulmonaire*, *Pulmo'saire en arbre* ou *de chêne*. This plant is sub-

astringent, and rather acid. It was once in high repute for curing diseases of the lungs.

LICHEN PYXIDATUS, *Muscus Pyxida'tus*, *Seyphoph'orus pyxida'tus*, *Mus'culus pyxo'ides ter'es'tris*, *Lichen pyxida'tus major* seu *coccif'erus*, *Herba Ignis*, *Cup Moss*, (F.) *Lichen entonnoir*, *Lichen Boëtier*, *L. Pyxidé*. This plant is sub-astringent, and has been used in decoction in whooping-cough.

LICHEN RETICULATUS, *L. pulmonarius*.

LICHEN ROCCEL'LA, *Parme'lia roccel'la*, *Roccella*, *R. tincto'ria*, *Seta'ria roccella*, *Litmus*, *Lacmus*, *L. tincto'rius*, *Orchill*, *Canary Archell*, *Chinney Weed*, *Herb Archell*, (F.) *Orseille*. The chief use of this plant is as a blue dye. It has been employed to allay cough in phthisis, &c.

LICHEN SAXATILIS, *L. tincto'rius* seu *laciniatus*, *Parme'lia saxat'ilis*, *Imbricia'ria saxat'ilis*, *Loba'ria saxat'ilis*, *Muscus Cra'nii huma'ni*, *Usne'a*. This moss, when found growing on a human skull, was formerly in high estimation against head affections, &c.

LICHEN SPURIUS, *L. caninus*—*l. Stellatus*, *Marchantia polymorpha*—*l. Tinctorius*, *L. Saxatilis*—*l. Wall*, yellow, *Parmelia parietina*.

LICHENIASIS ADULTORUM, *Lichen*—*l. Strophulus*, *Strophulus*.

LICHENOIDES ISLANDICUM, *Lichen Islandicus*.

LICKOROUS, Libidinous.

LICINIUM, Tent.

LICORICE, Glycyrrhiza.

LICULMEN, Garum.

LIEBENSTEIN, MINERAL WATERS OF. These waters, situated in the duchy of Saxe-Meiningen, are amongst the strongest acidulous chalybeates in Germany. They contain sulphate of soda, chloride of sodium, sulphate of lime, chloride of calcium, carbonate of lime, chloride of magnesium, and carbonate of iron.

LIEBERKÜHN'S GLANDS or FOLLICLES, *Glandulae* seu *Cryptae Lieberkühnia'nae*, so called from their first describer, are fine, capillary, blind sacs, the openings of which are from 1-20th to 1-30th of a line in diameter, so closely placed over the whole of the small intestine as to give the mucous membrane a general sieve-like or perforated appearance. They secrete the *succus entericus*.

LIEBERSCHES AUSZEHRUNG'S KRAUTER, see *Galeopsis grandiflora*—*l. Brustkrauter*, see *Galeopsis grandiflora*.

LIEBWERDA, MINERAL WATERS OF. Liebwerda is a Bohemian village, near the Silesian frontier. The springs are much frequented. There are four, all of which are rich in carbonic acid, and contain but little saline matter.

LIÈGE, Suber.

LIEN (F.), *Vin'culum*. A band, strap or garter, used in certain operations; as to tie patients during the operation of lithotomy; to fix the apparatus in fracture, &c. Also, the spleen.

LIEN ACCESSORIUS, *Lienculus*—*l. Ingens Splenoncus*—*l. Succenturiatus*, *Lienculus*.

LIEN'ULUS, diminutive of *lien*, 'spleen,' *Lien succenturiatus* seu *accessorius*. A supernumerary spleen.

LIENITIS, Splenitis.

LIENOSUS, Splenic.

LIENTERY, *Lienter'ia*, from *λειος*, 'smooth,' 'slippery,' and *εντερον*, 'intestine' *La'vitas* seu *Lax'itas* *intestino'rum*, *Diarrhoe'a* *lienter'ia*, *Fluxus* *Lienter'icus*, *Chymorrhoe'a* seu *La'vitas* *ingesto'rum*, *Bromatodœ'crisis*, *Diarrhoe'a* *Dyspep'tica* seu *cum Apepsiâ* seu *Ingesto'rum*, *Slip'periness of the Guts*. Frequent liquid evacuations, the food only half digested. This condition is always symptomatic of great irritation in the

Intestinal canal, the sensibility of which is so much augmented that it cannot bear the sojourn of the food in it.

LIERRE, *Hedera helix*—*l. Grimpant*, *Hedera helix*—*l. Terrestris*, *Glechoma hederacea*.

LIFE, *Sax. lif*, *lyf*, *Vita*, *Bios*, *Bi'otē*, *Bi'otos*, *Zōē*, *Pneuma*, *Spir'itus*, (F.) *Vie*. The state of organized beings, during which, owing to the union of an unknown force with matter, they are capable of performing functions different from those that regulate other natural bodies; all of which functions, however numerous and diversified, work to one end. Life has only a limited duration; beyond which,—the organic functions ceasing to be executed,—the body is given up to the agency of chemical affinity. Hence Bichat has defined life to be—the aggregate of the functions which resist death. On account of the difference that exists among the vital functions, he applied the term *Organic Life* to the functions inservient to composition and decomposition;—as digestion, respiration, circulation, calorification, absorption, secretion, and nutrition; and *Animal Life*, to the functions which connect man and animals with external bodies; as the understanding, sensations, locomotion and voice.

LIFE-BLOOD. A term, commonly used for a twitching or quivering of the eyelids—a kind of involuntary or muscular tic. See *Cillo*.

LIFE, DURATION OF, see *Longevity*.

LIFE EVERLASTING, *Gnaphalium margaritaceum*—*l. Everlasting*, plantain, *Antennaria plantaginifolia*—*l. Everlasting*, sweet-scented, *Gnaphalium polycephalum*.

LIFE, EXPECTATION OF, see *Longevity*.

LIFE-INSURANCE. A contract entered into, usually by an *insurance company*, to pay a certain sum of money on a person's death, on the condition of his paying an annual premium during his life. The medical practitioner may be applied to to certify that the life of the insured is one that is insurable according to the rules of the company.

LIFE ROOT, *Senecio aureus*.

LIFE, VALUE OF, see *Longevity*, and *Mortality*.

LIFE'S BLOOD, *Life-blood*, *Cillo*.

LIGAMENT, *Ligamen'tum*; from *ligare*, *ligatum*, 'to bind'; *Desmos*, *Syndes'mos*, *Colliga'men*, *Cop'ula*. A name given to fibrous structures, which serve to unite bones, and to form articulations; hence the division into *interosseous* and *articular* ligaments. They are of a white, close texture; are but little extensible, and difficult to break. The name *ligament* has, also, been given to any membranous fold, which retains an organ in its situation.

TABLE OF THE PRINCIPAL LIGAMENTS.

1. Of the Lower Jaw.	Capsular ligament. Suspensory ligament of the stylo-glossus. Lateral ligament.
2. Connecting the Head with the first and second Vertebrae, and these with each other.	Two capsular ligaments between atlas and head. Circular ligament. Two capsular between atlas and axis. Perpendicular ligament. Two lateral or moderator ligaments. Transverse ligament and its appendices. Anterior common. Crucial intervertebral. Ligaments running from the edge of the bony arch and spinous process of one vertebra to that of the next.
3. Of the other Vertebrae.	Interspinous ligament. Ligamentum nuchæ. Intertransverse. Capsular. Posterior or internal common.
4. Of the Ribs, Sternum, &c.	Capsular of the head of ribs. Capsular of the tubercles. Ligamenta transversaria interna. Ligamenta transversaria externa. Ligamenta cervicis costarum externa. Ligamentous fibres running from the margins of the extremities of the ribs to the corresponding cartilages. Radiated ligaments from cartilage of ribs to the sternum. Capsular ligaments of the cartilages of the ribs. Proper membrane of sternum. L. of cartilago-ensiformis. Tendinous expansions over the intercostales, &c.
5. Of the Pelvis.....	Two transverse,—one superior, one inferior. Ilio-sacral. Capsular of the sacro-iliac synchondrosis. Two sacro-ischiatic,—posterior and anterior, with the superior and inferior appendices. Longitudinal of os coccygis. Inguinal ligament. Capsular of symphysis pubis. Ligament of foramen thyroideum.
6. Of the Clavicle.....	Radiated ligament. Capsular. Interclavicular. Ligamentum rhomboideum. Claviculo-acromial. Conoid. Trapezoid.
7. Of the Scapula.....	Anterior triangular Proper posterior.
8. Of the Shoulder-Joint.	Capsular.
9. Of the Elbow-Joint.	Capsular. Brachio-ulnar. Brachio-radial. Coronary of the radius. Anterior and posterior accessory. Intermuscular of the os humeri.
10. Carpal Extremity of Radius and Ulna, and between those bones.	Interosseous ligament. Oblique or chorda transversalis cubiti. Capsular.
11. Between Fore-arm and Wrist.	Capsular. Two lateral. Mucous.
12. Of the Carpus.....	Annular. Capsular. Short ligaments.—oblique, transverse, capsular, and proper.
13. Between Carpal and Metacarpal bones.	Articular lateral. Straight, perpendicular, &c.
14. Between the extremities of the Metacarpal bones.	Interosseous, at the bases and heads,—dorsal, lateral, palmar.
15. At the base of the Metacarpal bone of the Thumb, and at the first joint of the Fingers.	Capsular. Lateral.
16. Of the first and second joints of the Thumb, and second and third joints of the Fingers.	Capsular. Lateral.
17. Retaining the Tendons of the Muscles of the Hand and Fingers in situ.	Annular. Vaginal or flexor tendons. Vaginal or crucial of the phalangen. Accessory of the flexor tendons. Posterior annular. Vaginal of extensors. Transverse of extensors.

18. *Connecting the Os Femoris with the Os innominatum.* { Capsular and accessory slips.
Round or teres ligament.
Cartilaginous ligament.
Double cartilaginous ligament.
Ligamenta mucosa.
19. *Of the Knee-Joint.* { Lateral,—internal and external.
External short lateral ligament.
Posterior ligament.
Ligament of the patella.
Capsular.
Ligamentum alare,—majus et minus.
Ligamentum mucosum.
Two crucial,—anterior and posterior.
Transverse.
20. *Connecting Fibula and Tibia.* { Capsular.
Interosseous.
Anterior superior.
Posterior superior.
21. *Connecting the Tarsal with the Leg Bones.* { Anterior ligament of the fibula.
Posterior of fibula.
Deltoides of tibia.
Capsular.
22. *Of the Tarsus.....* { Capsular.
Short ligaments.
Capsular, broad superior, and lateral ligaments, connecting astragalus and naviculare.
Superior, lateral, and inferior, fixing os calcis to os cuboides.
Long, oblique, and rhomboid, forming the inferior ligaments.
Superior superficial, interosseous and inferior transverse ligaments, fixing the os naviculare and os cuboides.
Superior lateral, and plantar, which fix the os naviculare and cuneiform.
Superior superficial and plantar, connecting the os cuboides, and os cuneiforme externum.
Dorsal and plantar, uniting the ossa cuneiformia.
The proper capsular of each bone.
23. *Between Tarsus and Metatarsus.* { Capsular.
Dorsal, plantar, lateral, straight, oblique, and transverse.
24. *Connecting the Metatarsal Bones.* { Dorsal, plantar, and lateral, connecting the metatarsal bones.
Transverse ligaments.
25. *Of the Phalanges of the Toes.* { Capsular.
Lateral.
Annular.
Vaginal of the tendons of the peronei.
Laciniated.
Vaginal of the tendon of the flexor longus pollicis.
Vaginal and crucial of the tendons of the flexors of the toes.
Accessory of the flexor tendons of the toes.
Transverse of the extensor tendons.
26. *Retaining the Tendons of the Foot and Toes in situ.* {

LIGAMENT, ANTERIOR, OF THE BLADDER. A name given by the older anatomists to a portion of the superior pelvic aponeurosis, which becomes attached to the front of the neck of the bladder.

LIGAMENT ARTÉRIEL, Arterial ligament—*l.* Camper's, Perineal fascia—*l.* Cervical supraspinal, see Supraspinosa ligamenta—*l.* Ciliaire, Ciliary ligament.

LIGAMENT, CORONARY, OF THE LIVER, is a reflection formed by the peritoneum, between the posterior margin of the liver, and the lower surface of the diaphragm. See Falx.

LIGAMENT, COSTO-XIPHOID, Xiphoid ligament—*l.* Dentelé, Denticulatum ligamentum—*l.* Denticulé, Denticulatum ligamentum—*l.* Dors-

lumbo-supraspinal, see Supra-spinosa ligamenta—*l.* of Fallopius, Crural arch—*l.* Gimbernati's, see Gimbernati's ligament—*l.* Glossepiglottic, see Glossepiglottic—*l.* Infra-pubian, Triangular ligament—*l.* Interépineux, Interspinal ligament—*l.* of the Ovary, see Ovarium—*l.* Palmar inferior, Metacarpal ligament—*l.* Poupart's, Crural arch—*l.* Rond, Teres ligamentum—*l.* Sur-épineux cervical, see Supra-spinosa ligamenta—*l.* Sur-épineux-dorso-lombaire, see Supra-spinosa ligamenta—*l.* Suspensory, of the liver, see Suspensory—*l.* Triangular, Perineal fascia.

LIGAMENTA ALARIA, Odontoid ligaments—*l.* Apicum, see Interspinal ligament—*l.* Coli, see Colon—*l.* Interlobularia pulmonum, Interlobular tissue—*l.* Intervertebralia, Intervertebral cartilages—*l.* Lata uteri, see Uterus—*l.* Mauthart, Odontoid ligaments—*l.* Rotunda uteri, Round ligaments of the uterus—*l.* Tarsæ lata, see Tarsæ lata (ligamenta).

LIGAMENTEUX, Ligamentous.

LIGAMENTOSUS, Ligamentous.

LIGAMENT'OUS, Ligament'ous, Desmons, Desmo'sus, (F.) Ligamenteux, Desmeux. Having the character of, or relating to, a ligament.

LIGAMENTS, BROAD, OF THE UTERUS, see Uterus—*l.* Croisés, Crucial ligaments—*l.* Cruciformes, Crucial ligaments—*l.* Jaunes, Yellow ligaments—*l.* Grandes de l'utérus, see Uterus—*l.* of the Larynx, inferior, Thyreo-arytenoid ligaments—*l.* Ronds de l'utérus, Round ligaments of the uterus.

LIGAMENTUM, Ligament—*l.* Arteriosum, see Arterial duct—*l.* Capsulare, Capsular ligament—*l.* Ciliare, Ciliary ligament—*l.* Conoides, Conoid ligament—*l.* Dentatum, Denticulatum ligamentum—*l.* Fallopii, Crural arch—*l.* Iridis, Ciliary ligament—*l.* Latum, Suspensory ligament of the liver—*l.* Membranaceum, Glosso-epiglottic ligament—*l.* Nuchæ, Cervical ligament—*l.* Posticum Winslowii, see Genu—*l.* Suspensorium dentis, see Odontoid ligaments—*l.* Suspensorium hepatis, Suspensory ligament of the liver—*l.* Suspensorium testis, Gubernaculum testis—*l.* Teres, Teres ligamentum—*l.* Trapezoides, Trapezoid ligament—*l.* Triangulare, Triangular ligament.

LIGATE, see Ligature.

LIGATIO, Fascia, Ligature—*l.* Linguae, Aphonia.

LIGATION, see Ligature.

LIGATURA, Ligature, Fascia—*l.* Glandis, Phimosis.

LIG'ATURE, Ligatu'ra, Liga'tio, Alligatu'ra, Deligat'io, Vincitu'ra, Fascia, Epid'esia. This word has various acceptations. It means, 1. The thread with which an artery or vein is tied, to prevent or arrest hemorrhage. 2. The cord, or thread, or wire, used for removing tumours, &c. 3. The bandage used for phlebotomy. *Ligature* is, also, sometimes applied to the act of tying an artery or considerable vessel—*Liga'tion*, and occasionally we meet with the terms *ligature* and *to ligate*, used unnecessarily in this sense, in place of to take up or tie a vessel. When the artery alone is tied, the ligature is said to be *immediate*; when any of the surrounding parts are included, it is said to be *mediate*. The ligature occasions obliteration or adhesion of the arterial parietes, by cutting through the middle and internal coats; the adhesion being favoured by the formation of a coagulum, which acts, in some degree, as a barrier against the impulse of the blood, and subsequently disappears by absorption.

LIGHT, Sax. lecht, lih't, (G.) Licht, Lur Lumen, Phos, (F.) Lumière. An extremely rare fluid; diffused over the universe; emanating from the sun and fixed stars; traversing more than four millions of leagues in a minute; passing through transparent bodies, which refract it ac-

ording to their density and combustibility; and arrested by opaque bodies, by which it is reflected at an angle equal to the angle of incidence. It is the cause of colour in all bodies, being entirely reflected by white surfaces and absorbed by black. It is decomposed in passing through a transparent prism into seven rays—red, orange, yellow, green, blue, purple, and violet.

Light acts upon the body as a gentle and salutary stimulus. It urges to exercise, whilst privation of it induces sleep and inactivity, and disposes to obesity. Hence it is, that, in rural economy, animals which are undergoing the process of fattening are kept in obscurity. When vegetables are deprived of light, their nutrition is interfered with, and they become *etiolated*. To a certain extent this applies to animals, and there is every reason to believe, that want of light prevents the due development of organized bodies. It has been found that, when tadpoles were deprived of light, they did not undergo the perfect metamorphosis into the frog, but that monstrous from arrest of development were induced.

LIGHTNING, *Astrape*.

LIGHTS, *Pulmo*—1. Rising of the, *Cynanche trachealis*.

LIGNE, *Line*—1. *Apre*, *Linea aspera*—1. *Blanche*, *Linea alba*—1. *Courbe*, *Curved line*—1. *Médiane de l'abdomen*, *Linea alba*—1. *Sous-trochantérienne*, *Linea aspera*.

LIGNEOUS, *Ligno'ous*, (F.) *Ligneux*. Woody. Having the character or appearance of wood; as '*Ligno'ous Scirrhus*,' or S. of the mamma, which has for its principal characteristics, hardness, the extensibility of wood, and no defined limits.—*Velpeau*. A form of this, in plates, or diffused over the mammary integuments, is termed, by M. *Velpeau*, *Squirrel en cuirasse*.

LIGNIN, from *lignum*, 'wood.' Ligneous or woody fibre; the fibrous structure of vegetable substances.

LIGNUM ALOES, *Agallochum*—1. *Aspalathi*, *Agallochum*—1. *Benedictum*, *Guaiacum*—1. *Brasilianum rubrum*, *Hæmatoxylon campechianum*—1. *Cæruleum*, *Hæmatoxylon campechianum*—1. *Campechense*, *Hæmatoxylon campechianum*—1. *Campechianum*, *Hæmatoxylon campechianum*—1. *Campechanum*, *Hæmatoxylon campechianum*—1. *Colubrinum*, see *Strychnos*—1. *Febrifugum*, *Quassia*—1. *Hæmatoxyl*, *Hæmatoxylon campechianum*—1. *Indicum*, *Guaiacum*, *Hæmatoxylon campechianum*—1. *Infelix*, *Sambucus*—1. *Nephriticum*, *Guilandina Moringa* (the wood)—1. *Pavane*, *Croton tiglium*—1. *Quassia*, see *Quassia*—1. *Sanctum*, *Guaiacum*—1. *Sandalinum*, see *Pterocarpus santalinus*—1. *Sappan*, *Hæmatoxylon campechianum*—1. *Serpentum*, *Ophioxylum serpentinum*—1. *Vitæ*, *Guaiacum*.

LIGNYODES, *Fulgionis*.

LIGNYS, *Fuligo*.

LIG'ULA, *Lin'gula*, 'a small tongue, a tongue of land; the tongue of a shoe.' The clavicle; also, the glottis and epiglottis. A measure containing 3 drachms and a scruple, or about half an ounce. Also, a species of bandage or ligature.—*Scribonius*.

LIGUSTICUM, *L. Levisticum*.

LIGUSTICUM ACTEIPOLIUM, *Thas'pium actæipolium*, *Angelica brœida*, *Angelica Nendo*, *White root*, indigenous; *Ord.* Umbellifera. The root is aromatic and carminative.

LIGUSTICUM CAPILLACRUM, *Ethusa meum*—1. *Carvi*, *Carum*—1. *Fœniculum*, *Anethum*.

LIGUSTICUM LEVISTICUM, from *Λιγυστικός*, pertaining to Liguria. *Levis'ticum*, *Laeripitium* *Germanicum*, *Ligusticum*, *Angelica levis'ticum*, seu *palustri'folia*, *Lavage*. (F.) *Liveche*, *Ache des montagnes*. *Ord.* Umbellifera. The pro-

perties of this plant are said to be stimulant, carminative, emmenagogue, &c.

LIGUSTICUM MEUM, *Ethusa meum*—1. *Phellandrium*, *Phellandrium aquaticum*.

LIGUSTICUM PODAGRARIA, *Podagra'ria ægopodinum*, *Ægopodinum* seu *Sison podagra'ria*, *Pimpinella angelica'folia*, *Sium vulgare*, *Tragoselinum angelica*, *Ses'eli ægopodinum*, *Angelica sylvestris*, *Gout-weed*. A British plant, once considered useful in cases of gout.

LIGUSTICUM SILAUS, *Peucedanum silaus*.

LIGUSTRUM ÆGYPTIACUM, *Lawsonia inermis*.

LIGUSTRUM VULGARE, *Privet*, *Privy*, *Prim*, (F.) *Troëne*. *Ord.* Oleaceæ. A shrub, which grows wild both in Europe and in the United States, usually in hedges. The leaves are astrigent and bitter; and the flowers, which are snow-white, and of an agreeable odour, have been employed in decoction, in sore throat, and ulcerous stomatitis. The berries are said to be cathartic.

LIGYSMA, *Distortion*.

LILAC, COMMON, *Syringa vulgaris*—1. *Vulgaris*, *Syringa vulgaris*.

LILL, *Lilium Paracelsi*.

LILIA'GO. Dim. of *Lil'ium*, the *lily*; *Spi-derwort*; *Lilias'trum*. This plant was formerly said to be alexipharmic and carminative.

LILIASTRUM, *Liliago*.

LIL' IUM CANDIDUM, *L. album*, *Crinon*. The *white lily*. (F.) *Lis blanc*. *Ord.* Liliacæ. The infusion of the flowers of the lily in olive oil is emollient, and often applied externally, under the name of *Lily oil*, (F.) *Huile de lis*. The scales of the bulb, roasted, are sometimes employed as maturatives.

LIL' IUM CONVALLIUM, *Convallaria maialis*.

LIL' IUM PARACEL'SI, *Tinctu'ra Metallo'rum*, *Lili*. A medicine employed by Paracelsus. It was prepared by melting in a crucible four ounces of each of the following alloys: *Antimony and iron*, *antimony and tin*, *antimony and copper*, previously mixed with eighteen ounces of nitrate of potassa and as much salt of tartar. The melted mass, when pulverized, was treated with rectified alcohol, which really dissolved only the potassa set at liberty by the decomposition experienced by the nitre and salt of tartar. The *Lilium Paracelsi* was used as a cordial. It entered into the composition of the theriacal elixir, &c.

LILY, GROUND, *Trillium latifolium*—1. *May*, *Convallaria maialis*—1. *Pond*, *Nelumbium luteum*—1. *Pond*, *White*, *Nymphaea odorata*—1. *Pond*, *yellow*, *Nuphar advena*—1. *Toad*, *Nymphaea odorata*—1. of the Valley, *Convallaria maialis*—1. *Water*, *little*, *Brasenia hydropeltis*—1. *Water*, *sweet*, *Nymphaea odorata*—1. *Water*, *sweet-scented*, *Nymphaea odorata*—1. *Water*, *white*, *Nymphaea alba*—1. *Water*, *yellow*, *Nymphaea lutea*—1. *White*, *Lilium candidum*, *Nelumbium luteum*.

LIMA, CLIMATE OF. The climate of Peru does not appear to be favourable to the generation of consumption: and Lima would seem to be a good residence for the phthisical valetudinarian. Many have been benefited by a residence there: but when they have gone farther south, as to Chili, the effect, according to Dr. M. Burroughs, has generally been fatal.

LIMA DENTARIA, *Scalprum dentarium*.

LIMAÇIEN (F.), from *limaçon*, (*limax*, *limacis*, 'the snail,' 'the cochlea of the ear.' A branch of the acoustic or labyrinthic nerve sent to the cochlea.

LIMAÇON, *Cochlea*. *Helix pomatia*, *Limax*—1. *Rampes du*, see *Cochlea*.

LIMACUM CORNUA, see *Lachrymal puncta*.

LIMANCHIA, *Abstinence*.

LIMA'TIO, from *limo*, *limatum*, 'to file.' Filing: an operation employed by the dentist more

especially to prevent immediate contact of the teeth with each other.

LIMATURA FERRI, Ferri Limatura.

LIMAX, λιμαξ, λιμακος, *Helix*, *Slug* or *Snail*, (F.) *Limaçon*, *Colimaçon*, *Escargot*. A syrup has been prepared from these animals, which has been given in phthisis, &c. The raw snails have also been taken in consumption. They have been used as food.

LIMB, Membrum.

LIMBUS ALVEOLARIS, Alveolar border—1. Fossa Ovalis, see Ovalis fossa—1. Luteus retinæ, see Foramen centrale—1. Posterior corporis striati, Tænia semicircularis.

LIME. A fruit like a small lemon, the juice of which is strongly acid, and is much used for making punch. It is, also, used in long voyages as an antiscorbutic, &c. It is a species of lemon, the fruit of *Citrus acida*.

LIME, Calx—1. Carbonate of, Creta—1. Chloride of, Calcis chloridum—1. Chlorite of, Calcis chloridum—1. Chloruret of, Calcis chloridum—1. Hydrate of, see Calx—1. Hydrosulphate of, Calcis sulphuretum—1. Hypochlorite of, Calcis chloridum—1. Muriate of, solution of, see Calcis murias—1. Oxymuriate of, Calcis chloridum—1. Phosphate of, see Cornu ustum—1. Bone phosphate of, see Cornu ustum—1. Slaked, see Calx—1. Sulphuret of, Calcis sulphuretum.

LIME TREE, Tilia—1. Water, Liquor calcis—1. Water, compound, Liquor calcis compositus.

LIMICUS, Hungry.

LIMITROPHES, see Trisplanchnic nerve.

LIMNE, Marsh.

LIMNE'MIC, *Limnæ'mic*, *Limnhæ'mic*, *Limnæ'micus*, *Limnhæ'micus*, from λιμνη, 'a marsh,' and 'αἷμα,' 'blood.' An epithet given to affections induced by paludal emanations; (F.) *Affections limnhémiques*.

LIMOCTON'IA, Abstinence. Abstinence to death. Death from hunger; from λιμος, 'hunger,' and κτος, 'death.' Suicide by hunger.

LIMODES, Hungry.

LIMON, see Citrus medica.

LIMONADA, Lemonade.

LIMONADE, Lemonade—1. Gazeuse, see Lemonade—1. Sèche, Lemonade, dry; see Citric acid.

LIMONIUM, Statice limonium—1. Malum, see Citrus medica.

LIMONUM BACCA, see Citrus medica.

LIMO-PHTHISIS, from λιμος, 'hunger,' and φθις, 'wasting.' Wasting from privation of food.

LIMOS, λιμος, 'hunger.' Hence:

LIMO'SIS, Stomach disease, Morbid appetite. A genus in the class Celiacæ, order Enterica, of Good.

LIMOSIS CARDIALGIA MORDENS, Cardialgia—1. Cardialgia sputatoria, Pyrosis—1. Dyspepsia, Dyspepsia—1. Expers, Anorexia—1. Expers protracta, Fasting—1. Flatus, Flatulence—1. Hellonium, Gluttony—1. Pica, Malacia.

LIMOTHEAPEIA, *Nestitherapeia*, *Nentotherapeia*, *Nestiatri'a*, *Peinotherapi'a*, *Cura'tio morbi per ine'diam*, from limos, and *Separæia*, 'treatment.' Hunger-cure. Cure by fasting.

LIMUS, Limos,

LIN, Linum usitatissimum—1. Graines de, see Linum usitatissimum—1. Purgatif, Linum catharticum.

LIN'NAIRE, Antirrhinum linaria.

LINAMENTUM, Lintum.

LINARIA, Antirrhinum linaria—1. Cymbalaria, Antirrhinum linaria—1. Elatine, Antirrhinum elatine—1. Vulgaris, Antirrhinum linaria.

LINCTUARIUM, Electos.

LINCTUS, Electos—1. Albus, Looch album—1. Amygdalinus, Looch album—1. ad Aphthas,

Mel boracis—1. de Borace, Mel boracis—1. Communis, Looch album.

LINDEN TREE, Tilia, Tilia Americana.

LINE, Lin'ea, Grammè, (F.) *Ligne*. Extent in length, considered without regard to breadth or thickness. As a measure, it means the 12th part of an inch.

MEDIAN LINE OF THE BODY is an imaginary line supposed to set out from the top of the head and to fall between the feet, so as to divide the body vertically into two equal and symmetrical parts.

LINEA, Line.

LINEA ALBA, *Linea alba Abdominis*, (F.) *Ligne blanche ou médiane de l'abdomen* (Ch.), *Candidula abdominis lin'ea*, *Lin'ea centralis*. A tendinous, strong, and highly resisting cord; extending from the ensiform cartilage of the sternum to the symphysis pubis, with the umbilicus near its middle. The linea alba is formed by the decussation of the aponeurosis of the abdominal muscles; and its use is to limit the movement of the chest backwards; to prevent it from separating too far from the pelvis, and to furnish a fixed point for the muscles of the abdomen in their contraction.

LINEA AS'PERA, *Ragged ridge*, (F.) *Ligne âpre*, *Ligne sous-trochantérienne*, (Ch.) A rough projection at the posterior surface of the femur, which gives attachment to muscles.

LINEA CANDIDULA ABDOMINIS, L. alba—1. Centralis, L. alba—1. Innominata, Ilio-pectinea Linea.

LINEÆ ARCUATÆ, Curved lines—1. Semicirculares externæ, Curved lines.

LINEÆ SEMILUNARES are the lines, which bound the outer margin of the recti muscles of the abdomen. They are formed by the union of the abdominal tendons. The lines which cross these muscles are called *Lineæ transversæ*.

LINEÆ TIBIÆ, *Anguli tib'ie*. Sharp lines on the tibia.

LINEÆ TRANSVERSÆ, see L. semilunares, and Processus teretes.

LINÉAIRE, Linear.

LINÉAMENT, *Lineamentum*, from *linea*, 'a line,' *lineare*, 'to mark by lines.' A delicate trait observed on the countenance, which constitutes its special character, enables us to preserve its image, and is the cause of resemblance to others. A feature.

Bonnet gave the name *Linéament* to the first traces of organization in the embryo of man and animals.

LIN'EAR, *Linea'ris*, *Linea'rius*, from *linea*, 'a line.' (F.) *Linéaire*. Pathologists apply the epithet *linear* to fractures which are very narrow, and in which the fragments are scarcely separated.

LINE'OLA; a diminutive of *linea*, 'a line.' A small line.

LINE'OLE MAMMA'RUM. The white lines on the breasts.

LINGISM, Kinesipathy.

LINGUA, Tongue—1. Avis, see Fraxinus excelsior—1. Bovis, *Anchusa officinalis*—1. Canina, *Cynoglossum*—1. Cervina, *Asplenium scolopendrium*—1. Exigua, *Epiglottis*—1. Prognosis ex, *Glossomantia*—1. Serpentina, *Ophioglossum vulgatum*.

LINGUÆ DETENTOR, *Glossocatochus*—1. Exoncosis, *Glossoncus*—1. Scalpinus, *Cure-langue*.

LIN'GUAL, *Lingua'lis*, from *lingua*, 'the tongue.' Relating or belonging to the tongue.

LINGUAL ARTERY, *Arteria lingua'lis*, arises from the external carotid; and, after several tortuosities, reaches the base of the tongue, becomes horizontal, and, under the name *Ranina*, advances

to the tip, where it anastomoses with its fellow. In its course, it gives off the *Dorsalis linguae* and *sublingual*.

LINGUAL MUSCLE, *Lingualis*, *Basio-glossus* (Copper), *Glossia'nus*, is a small, long, fasciculus of fibres, hidden beneath the sides of the tongue, between the hyoglossus and styloglossus muscles, which are on the outside, and the genioglossus, within. This muscle passes from the base to the tip of the tongue; and, at its sides, is confounded with the muscles just referred to. It shortens the tongue, depresses its point, and can carry it to the right or left side.

LINGUAL NERVE is a name which has been given to the ninth pair or hypoglossus. It is, also, a term applied to a branch given off from the *Inferior maxillary*, or third branch of the fifth pair. Near its origin it anastomoses with, or simply runs close to, the chorda tympani. It afterwards gives off a considerable number of filaments, which are distributed to the tongue, and some of which are said to have been even traced as far as the papillae;—a distribution which has occasioned it to be regarded as the *Gustatory nerve*, (F.) *Nervus gustatif*.

LINGUAL SALIVARY GLANDS, see *Salivary glands*.

LINGUAL VEIN follows nearly the same distribution as the artery. It opens into the internal jugular.

LINGUALIS, Lingual muscle.

LINGUETTA LAMINOSA, Lingula.

LINGULA, Ligula.

LINGULA, *Linguet'ta lamino'sa*. A thin, transversely grooved lobule of gray substance, derived from the anterior border of the cerebellum, which, for a short distance, lies over the velum medullare arterius.

LINGULA FISTULÆ, Epiglottis.

LINI USITATISSIMI SEMINA, see *Linum usitatissimum*.

LINIMENT, *Linimen'tum*, *Litus*, *Enchris'ton*, *Catechris'ton*, *Perich'risis*, *Perichris'ton*, *Aleiph'a*, *Aleim'ma*, *Enchris'ta*, *Frictum*, *Fricato'rium*, *Inunctio*, from *linire*, *lini'tum*, 'to anoint gently,' 'to anoint.' An unctuous medicine, containing usually oil or lard, which is used externally in the form of friction.

LINIMENT OF AMMONIA, **COMPOUND**, see *Lotion*, *Graville's*.

LINIMENT AMMONIACAL, *Linimentum ammoniacum*—*l. de Carbonate d'Ammoniaque*, *Linimentum ammoniac carbonatis*.

LINIMENT, *Anodyne*, *Linimentum opii*.

LINIMENT ANTIHÉMORRHOÏDALE DE SIEUR ANDRY, (F.) This is formed of *Norbonne honey*, *Olive oil*, and *Turpentine*.

LINIMENT ANTIPARALYTIQUE, (F.) Composed of *subcarbonate of ammonia*, *alcoholized oil*, *black soap*, and *oil of rosemary*.

LINIMENT, ANTISCROFULOUS, OF *HUFELAND*. Composed of *fresh ox-gall*, *White soap*, *Unguentum althææ*, *Volatile oil of petroleum*, *Carbonate of ammonia*, and *Camphor*.

LINIMENT CALCAIRE, *Linimentum aquæ calcis*—*l. Camphor*, *Linimentum camphoræ*—*l. Camphré*, *Linimentum camphoræ*—*l. of Cantharides*, *camphorated*, *Linimentum e cantharidibus camphoratum*—*l. Cantharide camphré*, *Linimentum e cantharidibus camphoratum*—*l. Croton oil*, *Linimentum crotonis*—*l. d'Eau de chaux*, *Linimentum aquæ calcis*—*l. of Limewater*, *Linimentum aquæ calcis*—*l. de Mercure*, *Linimentum hydrargyri*—*l. Mercurial*, *Linimentum hydrargyri*—*l. Oléo-calcaire*, *Linimentum aquæ calcis*—*l. of Opium*, *Linimentum opii*.

LINIMENT RÉSOLUTIF DE POTT, com-

posed of *oil of turpentine* and *muritic acid*. Used in rheumatism, swellings, &c.

LINIMENT, SAINT JOHN LONG'S. A liniment used by a celebrated empiric at the commencement of the second quarter of the 19th century. It is said to have consisted of *oil of turpentine* and *acetic acid*, held in suspension by yolk of egg. It was a powerful counter-irritant.

LINIMENT DE SAVON, *Linimentum saponis compositum*—*l. de Savon opiacé*, *Linimentum saponis et opii*.

LINIMENT SAVONNEUX HYDROSULFURÉ DE JADELLOT. Composed of *sulphuret of potass*, $\mathfrak{Z}\text{vj}$; *white soap*, liij ; *oil of poppy*, liiv ; *oil of thyme*, $\mathfrak{Z}\text{ij}$. Used in itch.

LINIMENT, SIMPLE, *Linimentum simplex*—*l. Soap*, *Linimentum saponis*—*l. Soap and opium*, *Linimentum saponis et opii*—*l. Soap, compound*, *Linimentum saponis compos.*—*l. Turpentine*, *Linimentum terebinthinæ*—*l. of Verdigris*, *Linimentum æruginis*—*l. de Vert-de-gris*, *Linimentum æruginis*—*l. Volatil*, *Linimentum ammoniac fortius*.

LINIMENTUM, *Liniment*.

LINIMENTUM ÆRUGINIS, *Ocymel Æruginis*, *Unguentum Ægyptiacum*, *Mellitum de aceta'te cupri*, *Liniment of verdigris*, (F.) *Liniment de Vert-de-gris*, *Miel d'acétate de cuivre*. (*Ærugin*, cont. $\mathfrak{Z}\text{ij}$; *acet.* $\mathfrak{Z}\text{vij}$; *in illis de'sum.* pond. $\mathfrak{Z}\text{xiv}$. Liquefied, strained, and inspissated by boiling.—Ph. L.) It is used as an escharotic and detergent;—diluted, as a gargle in venereal ulcerations and in foul ulcers.

LINIMENTUM ALBUM, *Ceratum cetacei*, *Unguentum cetacei*—*l. ad Ambustiones*, *L. aquæ calcis*.

LINIMENTUM AMMONIÆ, *Sapo ammoniacalis*, *Linimentum ammoniacum* seu *ammoniacum* seu *Anglicanum*, *Sapo ammoniac oleaceus*, *Unguentum album resolvens*, *Oleum ammoniacum*, *Strong liniment of ammonia*, (F.) *Liniment volatil ou ammoniacal*, *Savon ammoniacal*. (*Liq. ammon.* $\mathfrak{f}\mathfrak{Z}\text{ij}$; *olei olivæ*, $\mathfrak{f}\mathfrak{Z}\text{ij}$. Mix.—Ph. U. S.) A stimulating and rubefacient soap.

LINIMENTUM AMMONIÆ CARBONATIS, *Liniment of subcarbonate of ammonia*, *Linimentum ammoniac sesquicarbonatis*, *L. volatilè*, *Hartshorn and oil*, (F.) *Liniment de carbonate d'ammoniaque*. (*Solut. subcarb. ammon.* $\mathfrak{f}\mathfrak{Z}\text{ij}$; *olei olivæ*, $\mathfrak{f}\mathfrak{Z}\text{ij}$. Shake till they unite.—Ph. L.) A stimulating liniment, mostly used to relieve rheumatic pains, bruises, &c.

LINIMENTUM AMMONIÆ COMPOSITUM, see *Graville's Lotion*—*l. Ammoniac Sesquicarbonatis*, *Linimentum Ammoniac Carbonatis*—*l. Ammoniacum* seu *Ammonicum*, *L. Ammoniac fortis*—*l. Anglicanum*, *L. Ammoniac fortis*—*l. ad Aphthas*, *Mel boracis*.

LINIMENTUM AQUÆ CALCIS, *L. Calcis* (Ph. U. S.), *L. ad ambustiones*, *Oleum lini cum calcē*, *Sapo calcarius*, *Liniment of limewater*, *Carron oil*, (because much used in the Carron iron works in Scotland), (F.) *Liniment d'eau de chaux*, *Savon calcaire*, *Liniment calcaire ou oléo-calcaire*. (*Olei lini*, *aquæ calcis*, \mathfrak{aa} $\mathfrak{f}\mathfrak{Z}\text{ij}$. Misce.) A cooling and emollient application to burns and scalds.

LINIMENTUM ARCEI, *Unguentum elemi compositum*—*l. de Borace*, *Mel boracis*—*l. Calcis*, *L. aquæ calcis*.

LINIMENTUM CAMPHORÆ, *Oleum Camphoratum*, *Solutio camphoræ oleo'sa*, *Camphor liniment*, (F.) *Liniment Camphré*, (*Camphoræ*, $\mathfrak{Z}\text{iv}$; *olei olivæ*, $\mathfrak{f}\mathfrak{Z}\text{ij}$. Dissolve.—Ph. U. S.) It is used as a stimulant and discutient.

LINIMENTUM CAMPHORÆ COMPOSITUM, *Ward's essence for the headache*, *Compound camphor liniment*. (*Camphor*, $\mathfrak{Z}\text{iiiss}$; *ol. Lavand.* $\mathfrak{f}\mathfrak{Z}\text{ij}$; *Sp. rectificat.* $\mathfrak{f}\mathfrak{Z}\text{xvij}$; *Liquor ammon. fortior.* $\mathfrak{f}\mathfrak{Z}\text{ij}$. Dissolve the camphor and oil in the spirit; add

the ammonia, and shake till they are mixed. Ph. L.) It is stimulant and discutient.

LINIMENTUM CANTHARIDIS, *Liniment of Spanish Flies*. (Cantharid. in pulv. $\frac{3j}{i}$; Ol. Terebinth. Oss. Digest for three hours by means of a water-bath, and strain.—Ph. U. S.) Used as an excitant liniment in typhus, &c.

LINIMENTUM E CANTHARIDIBUS CAMPHORATUM, *Camphorated liniment of cantharides*, (F.) *L. cantharide camphrè*. (Tinct. cantharid. $\frac{3ss}{i}$; Ol. amygd. dulc. $\frac{3iv}{i}$; sapon. amygd. $\frac{3j}{i}$; camphor. $\frac{3ss}{i}$. Dissolve the camphor in the oil, and add the mixture to the tincture and soap.—Ph. P.) Rubefacient, and discutient.

LINIMENTUM CROTONIS, *Croton oil Liniment*, (Ol. Croton, $\frac{f3j}{i}$; Ol. Terebinth, $\frac{f3vij}{i}$, M. Ph. D.) Rubefacient, causing an eruption when continued.

LINIMENTUM HYDRARGYRI, *L. Hydragryri compos'itum*, *Mercurial liniment*, (F.) *L. de Mercure*. (Ung. Hyd. fort., adip. præp. $\frac{aa}{i}$, $\frac{3iv}{i}$; camphora, $\frac{3j}{i}$; sp. rect. gtt. xv; liquor ammon. $\frac{f3iv}{i}$. Rub the camphor with the spirit; add the ointment and lard; and, lastly, gradually add the solution.—Ph. L.) It is used as a stimulant and discutient to venereal swellings, &c.

LINIMENTUM HYPERICI, see *Hypericum perforatum*.

LINIMENTUM OPII, *Liniment of Opium*, *Anodyne Liniment*. (Sapon. $\frac{3vj}{i}$; Opii. $\frac{3iss}{i}$; Camphor. $\frac{3iij}{i}$; Ol. Rosmarin. $\frac{f3vj}{i}$; Spirit rectificat. Oij. (Imp. meas.) Macerate the soap and opium in the spirit for three hours; filter, and add the oil and camphor.—Ph. L.) Anodyne and slightly rubefacient.

LINIMENTUM PLUMBATUM, *Unguentum plumbi superacetatis*—1. Saponaceum opiatum, L. Saponis et opii.

LINIMENTUM SAPONACEUM HYDROSULPHURATUM, (*Common Soap*, 500 p. Liquefy in a water bath in an earthen vessel; and add white poppy oil, 250 p. Mix intimately, and add dry sulphuret of potass, 100 p. Beat together, and add oil of poppy seed, 750 p.—Ph. P.) In cutaneous affections, as psora, herpes, &c.

LINIMENTUM SAPONATO-CAMPHORATUM, L. Saponis camphoratum.

LINIMENTUM SAPONIS, *Tinctura saponis camphorata* (Ph. U. S.), *Camphorated Tincture of Soap*, *Soap Liniment*. (Saponis concis. $\frac{3iv}{i}$; camphor. $\frac{3ij}{i}$; Ol. Rosmarin. $\frac{f3ss}{i}$; alcohol. Oij. Digest the soap and alcohol in a water-bath, until the former is dissolved; filter, and add the camphor and oil.—Ph. U. S.) Used in sprains, bruises, and as an embrocation.

LINIMENTUM SAPONIS CAMPHORATUM, L. S. compos'itum, L. Saponato-camphoratum, *Balsamum opodel'doc*, *Tinctura saponis camphorata* (Ph. U. S.), *Opodel'doch*, *Steer's opodel'doch*, *Compound Soap Liniment*, (F.) *L. de Savon*. (Sapon. rasur. $\frac{3iv}{i}$; Camphor. $\frac{3ij}{i}$; Ol. Rosmarin. $\frac{f3ss}{i}$; Aquæ, $\frac{f3iv}{i}$; Alcohol. Oij. Mix the alcohol and water; digest the soap in the mixture by means of a water-bath, until it is dissolved; filter, and add the camphor and oil.—(Ph. U. S.) Stimulant and anodyne; in bruises, local pains, &c.

Freeman's Bathing Spirits consist of lin. sapon comp. coloured with *Daffy's elixir*.

Jackson's Bathing Spirits differ from *Freeman's* in the addition of some essential oils.

LINIMENTUM SAPONIS ET OPII, L. *saponaceum opiatum*, *Soap and opium liniment*, *Bates's anodyne balsam*, *Balsamum anodynum*, *Tinctura saponis et opii*, (F.) *L. de Savon opiacé*. (Saponis duri. $\frac{3ij}{i}$; opii. $\frac{3j}{i}$; camphora, $\frac{3ij}{i}$; olei rosmarin. $\frac{f3ss}{i}$; alcohol. Oij.—Ph. E.) Anodyne; in chronic rheumatism and local pains in general.

LINIMENTUM SIMPLEX, *Simple Liniment*. (Ol. oliv. 4 p.; ceræ alba. 1 p. fiat linimentum.—Ph. E.) Emollient; used in chaps, &c.

LINIMENTUM SIMPLEX, *Unguentum ceræ*.

LINIMENTUM TEREBINTHINÆ, L. *Terebinthina-tum*, *Turpentine liniment*. (Cerat. resinæ, $\frac{ij}{i}$; ol. terebinth. Oss. Melt the cerate and stir in the oil.) A stimulant; applied to burns, &c.

LINIMENTUM VOLATILE, L. *ammoniac carbonatis*.

LINN, *Tilia Americana*.

LINNÆA, L. *Borea'lis*. Ord. Caprifoliaceæ.

This plant, called after Linnæus, has a bitter, sub-astringent taste; and is used, in some places, in the form of fomentation, in rheumatic pains. An infusion in milk is used in Switzerland for the cure of sciatica.

LINOSPERMUM, see *Linum usitatissimum*.

LINSEED, see *Linum usitatissimum*.

LINT, Carbasus, Linteum.

LINTAMEN, Linteum, Pledget.

LINTEUM, 'a linen cloth; from *linum*, 'flax, linen,' L. carptum seu rasum, *Linte'men*, *Oth'onè*, *Othon'ion*, *Carbasus*, *Car'basæ*, *Car'pia*, *Motos*, *Motè*, *Motum*, *Linamentum*, *Tilma*, *Xystos*, *Xysma*, *Achnè*, *Lint*, (F.) *Charpie*. A soft, flocculent substance, made by scraping old linen cloth, (F.) *Charpie râpée*, or by unravelling old linen cut into small pieces—(F.) *Charpie brute*; and employed in surgery as a dressing to wounds, ulcers, &c., either simply or covered with ointment.

LINTEUM CARPTUM, see *Linteum*—1. Rasum, see *Linteum*—1. Scissum, *Splitcloth* (residue).

LINUM, see *Linum usitatissimum*—1. Arvense, L. usitatissimum.

LINUM CATHARTICUM, L. *min'imum*, *Chamælinum*, *Purging flax*, *Mill mountain*, (F.) *Lin purgatif*. Ord. Linaceæ. This plant is possessed of cathartic properties, and has a bitterish, disagreeable taste. Dose, $\frac{3j$, in substance.

LINUM CRUDUM, see *Apolinosis*—1. Minimum, L. Catharticum.

LINUM USITATISSIMUM, L. *arven'se*, *Common flax*, (F.) *Lin*. The seed, *Linum*, (Ph. U. S.), *Sem'ina lini usitatissimi*, *Linosp'er'mum*, *Linseed*, *Flaxseed*, (F.) *Grains de lin*, are inodorous, and almost tasteless; yielding mucilage to warm water, and oil by expression. They are demulcent and emollient. Linseed, when ground into powder, forms a good emollient poultice. It is only necessary to stir the powder into boiling water. The oil, *Oleum lini*, *Flaxseed oil*, (F.) *Huile de Lin*, is emollient and demulcent.

LIONDENT, *Leontodon taraxacum*.

LION'S FOOT, *Nabalus albus*, *Prenanthes*—1. Tall, *Leonurus*.

LIOPODIA, see *Leopopus*.

LIOPUS, *Leiopus*.

LIP, Sax. and Germ. *Lippe*, *Cheilos*, *La'bium*, *Labrum*, (F.) *Lèvre*. [In Entomology, *labium* means the lower lip, —*labrum*, the upper.] The lips are composed of different muscular fasciculi, nerves, and vessels, covered by the skin and mucous membrane of the mouth. They circumscribe the anterior aperture of that cavity; and are inservient to mastication, pronunciation, &c. They are distinguished into *upper* and *lower*—*Anochei'lon*; and *Catochei'lon*, (Prov.) *Fipple*—and are placed in front of each jaw, forming between them the anterior aperture of the mouth. They unite at each side, and form what are called the *angles* or *commissures* of the mouth—*Chal'ini*. Their free edge is covered with a mucous membrane, of a more or less livid red, according to the individual. They receive their arteries from the external carotid. Their veins open into the two jugulars. Their lymphatic vessels descend into the ganglions situate beneath the chin. Their nerves are derived from the infra-orbital, mental, and facial.

LIPS, *La'bia*, (F.) *Lèvres de la vulve*, are folds belonging to the genital organs of the female,

and distinguished into—1. *Labia pudendi* seu *Cunni* seu *majo'ra*, *Episi'a*, *Alæ pudendi muliebris*, *Hytrochei'tides*, *Rupes*, *Alæ majo'res*, *Crem'noi*, (F.) *Grandes Lèvres*. These are two membranous folds, thicker above than below, which limit the vulva laterally, and extend from the inferior part of the mons veneris to the perinaeum. They unite anteriorly and posteriorly, forming commissures; the posterior of which is called *Fourchette*. Their outer surface is convex; formed of skin and covered with hair. The inner surface is white, and covered by a mucous membrane, continuous with that lining the other parts of the vulva. The space between the skin and mucous membrane is filled with a fatty tissue and fibrous bands, some fibres of the constrictor vaginae muscle, vessels, and nerves.

We speak, also, of the *lips of a wound, ulcer, &c.*, when alluding to the edges of these solutions of continuity.

LIPA, λιπα, fat; also, Lippitudo.

LIP'ARA, from λιπαρος, 'fatty.' Plasters, containing much oil or fat.

LIPARIA, see Polysarcia.

LIPAROCE'LE, *Lipoce'le*, from λιπαρος, 'fatty,' and κηλη, 'tumour.' *Lipo'ma*, or fatty tumour of the scrotum, *Her'nia pinguedino'sa scroti*.

LIPARODES, Fatty.

LIPAROIDÉ, Cerate.

LIPAROLE, Pomatum, *Pommade*.

LIPAROTES, see Polysarcia.

LIPAROTRICH'IA, from λιπαρος, 'fatty,' and σπῆς, τριχος, 'hair.' Too great oiliness of the hair.

LIPARUS, Fatty.

LIPASMA, see Polysarcia.

LIPEMANIA, Melancholy, see Lypemania.

LIPHÆMIA, Leiphæmia.

LIPHÆMOS, Leiphæmos.

LIPO or LEIPO, λιπω, 'I leave,' 'I forsake.' Hence, *Leipopsychia*, *Leipothymia*.

LIPOCELE, Liparocèle.

LIPODERMIA, Aposthia.

LIPODERMOS, Leipodermos.

LIPO'MA, *Lypo'ma*, from λιπα, 'fat.' A fatty tumour of an encysted or other character.

LIPOMATODES, Lipomatous.

LIPOM'ATOUS, *Lipomato'us*, *Lipomato'des*, (F.) *Lipomateuz*. Having the nature of lipoma—as a 'lipomatous mass.'

LIPOME'RIA, *Leipome'ria*, from λιπω, 'I leave,' and μερος, a 'part.' Monstrosity from arrest of development, or from defect. Deficiency of one or more of the parts of the body; for example, in one who has only four fingers.

LIPOPSYCHIA, Syncope.

LIPOS, Fat.

LIPOSIS, see Polysarcia.

LIPOSPHYXIA, Syncope.

LIPOTHYMIA, Syncope.

LIPPA, Chasie.

LIPPER, Lepra, Leprous.

LIPPIA, Adali.

LIPPITU'DO, *Ophthal'mia chron'ica* of some, *Lippitude*, *Blar-eye*, *Blar-eyedness*, *Blar'edness*, *Lappa*, *Lipa*, *Xerophthal'mia*, *Lema*, *Lemè*, *Lemos'ias*, *Glemè*, *Glama*, *Gra'mia*. A copious secretion of the sebaceous humour of the eyelids, which renders them gummy. It is owing to a state of chronic inflammation of the tarsal margins; the eyelids being generally red, tumefied, and painful.

LIPPITUDO NEONATORUM, see Ophthalmia.

LIPPUS, one affected with blar eye; *blar-eyed*; (F.) *Chasieux*.

LIPSIS ANIMI, Syncope.

LIPSOTRICHIA, Alopecia, Calvities.

LIQUAMEN, Garum.

LIQUAMU'MIA, *Adeps humu'nus*. *Humenz* fat.—Ruland and Johnson.

LIQUARUM, Syrupus simplex.

LIQUATIO, Liquefaction.

LIQUEFA'CIENT, *Liquefa'ciens*, from *liquidus*, 'liquid,' and *facere*, 'to make.' A medicinal agent, which seems to have the power of liquefying solid dispositions. To this class mercury, iodine, &c., have been referred by some.

LIQUEFAC'TION, *Liqua'tio*, *Liquefac'tio*; same etymon. Transformation of a solid substance into a liquid. It is used particularly in speaking of metals and fatty bodies, see Fusion.

LIQUEUR DE L'AMNIOS, *Liquor amni'i*—*l. d'Alumine Composee*, *Liquor aluminis compos.*—*l. Anodine d'Hoffman*, *Spiritus ætheris sulphurici compositus*—*l. Anodine nitreuse*, *Spiritus ætheris nitrici*—*l. Arénicale*, *Liquor arsenicalis*—*l. Fumante de Boyle*, *Ammonia sulphuretum*, *Liquor fumans Boylii*—*l. Volatile de Corne de cerf*, *Liquor volatilis cornu cervi*—*l. de Cuivre ammoniacal*, *Liquor cupri ammoniaci*—*l. de Fer alcaline*, *Liquor ferri alkali*—*l. de Sous-acétate de plomb*, *Liquor plumbi subacetatis*—*l. de Potasse*, *Liquor potassæ*—*l. du Sung*, *Liquor sanguinis*—*l. de Sous-carbonate de potasse*, *Liquor potassæ subcarbonatis*.

LIQUID, ADHESIVE, MAYNARD'S, Collo-dion—*l. Disinfecting*, *Burnett's*, see *Burnett's disinfecting liquid*—*l. Disinfecting*, *Labarraque's*, *Liquor sodæ chlorinatæ*—*l. Disinfecting*, *Ledoyen's*, see *Ledoyen's disinfecting liquid*.

LIQUIDAMBAR ASPLENIFOLIA, *Comptonia asplenifolia*—*l. Officialis*, see *Styrax*—*l. Perregina*, *Comptonia asplenifolia*.

LIQUIDAM'BAR STYRACIF'LUA, *Liquidam'bra*, *Sweet gum*, *White gum*. *Ord. Balsamineæ*. The name of the tree which affords the *liquid amber* and *Storax liquida*, *Styrax liquida* or *Liquid storax*. *Liquid amber* is a resinous juice of a yellow colour, inclining to red; at first of about the consistence of turpentine; by age, hardening into a solid, brittle mass. It is not used medicinally.

Styrax Liquida is obtained from this plant by boiling. There are two sorts,—one more pure than the other. It is used occasionally, as a stomachic, in the form of plaster.

LIQUIDE AMNIOTIQUE, *Liquor Amni'i*—*l. Céphalo-rachidienne*, *Cephalo-spinal fluid*.

LIQUIDUM NERVEUM, *Nervous fluid*.

LIQUIRITIA, *Glycyrrhiza*.

LIQUOR, (*liqueur*, 'I am liquid,') *Liqua'men*, *Hyggon*, *Hy'grotes*, (F.) *Liqueur*. A name given to many compound liquids, and especially to those the bases of which are water and alcohol.

LIQUOR ACIDUS HALLERI, *Elixir acidum Halleri*—*l. Æthereus*, *Ether*—*l. Æthereus sulphuricus*, *Spiritus ætheris sulphurici*.

LIQUOR ALU'MINIS COMPOS'ITUS, *Aqua alu'minis compos'ita* seu *alumin'o'sa Bates'na*, *Bates's alum water*, *Compound solution of Alum*, (F.) *Liqueur d'alumine composée*. (*Alumina*, *zinci sulphat.* sing. \mathfrak{z} ss, *aqua fervent.* Oij. Dissolve and filter through paper.) Detergent and stimulant. Used as a collyrium, when properly diluted, in ophthalmia; as an injection in gleet, leucorrhæa, &c.

LIQUOR EX AMMONIÀ ET OLEO SUCCINI, *Spiritus ammoniæ succinatus*.

LIQUOR AMMO'NIÆ, *L. ammoniæ puræ*, *Al'cali ammoniacum flu'idum*, *Ammoni'a liquida* seu *caustica liquida* seu *pura liquida*, *Lixiv'ium ammoniacal'e*, *Spiritus salis ammoniaci aquo'us*, *Aqua ammonia* seu *ammonia caust'ica*, *Liquid ammonia*, *Solution of ammonia*, (F.) *Ammoniaque liquide*. Directed to be made in the Pharm. U. S. from *muriate of Ammonia*, in fine powder; and *Lime*, each a pound; *distilled water* a pint; *water*,

nine fluid ounces. The water is employed to slake the lime; this is mixed with the muriate of ammonia and put into a glass retort in a sand-bath. Heat is applied so as to drive off the ammonia, which is made to pass into a quart bottle containing the distilled water. To every ounce of the product three and a half fluidrachms of distilled water are added, or as much as may be necessary to raise its s. g. to 0.96. This Liquor Ammonia may also be made by mixing one part of liquor ammonia fortior with two parts of distilled water. (Ph. U. S. 1842.) Its s. g. is 0.960. LIQUOR AMMONIAE FORTIOR, *Stronger solution of ammonia* (Ph. U. S.) is an aqueous solution of ammonia of the s. g. 0.882. Liquor ammonia is stimulant, antacid, and rubefacient. Dose, gtt. x to xx, in water or milk.

LIQUOR AMMONIAE ACETA'TIS, *Al'cali ammoniacum acetatum, A. volat'ile acetatum, Aqua acetatis ammoniacæ seu ammoniacæ acetatæ, Solution of acetate of ammonia, Acet'as ammoniacæ, Spirit of Mindere'rus, Sal ammoniacum vegetabile, Spiritus ophthalmicus Mindere'ri, Sal acetosus ammoniacalis.* (Acid. acetic. dilut. Oj, Ammon. carbonat. in pulv. Add the salt to the acid until it is saturated. Ph. U. S. 1851.) A sudorific; externally, cooling. Dose, fʒij to fʒiiss.

LIQUOR AMMONIAE SESQUICARBONATIS, *Liquor Ammonia Subcarbonatis.*

LIQUOR AMMONIAE CITRA'TIS, *Solution of Citrate of Ammonia.* This preparation is official in the Dublin Pharmacopoeia. It is made by saturating a solution of citric acid with sesquicarbonate of ammonia. It is given in cases of adynamic fever.

LIQUOR AMMONIAE SESQUICARBONATIS, *Liquor Ammonia Sesquicarbonatis, Solutio subcarbonatis ammoniacæ, Aqua carbonatis ammoniacæ, Solution of subcarbonate of ammonia, Water of carbonate of ammonia.* (Ammonia carbon. ʒiv, aquæ destillat. Oj. Dissolve and filter through paper.) Use; — the same as that of the carbonate of ammonia.

LIQUOR AMMONII HYDROTHIODIS, *Ammonia sulphuretum*—*l. Ammonii vinosus, Spiritus ammoniacæ.*

LIQUOR AMNII, *Aqua amni.* The liquor of the amnios. (F.) *Eau ou Liqueur de l'amnios, Liqueur amniotique.* The fluid exhaled by the amnios, and which envelops the fetus during the whole period of utero-gestation. It is often simply called the waters, (F.) *Les Eaux.* Its relative quantity diminishes as pregnancy advances, although its absolute quantity continues to increase till the period of delivery. In some women, only five or six ounces are met with; in others, it amounts to pints. It is limpid, yellowish, or whitish; exhales a faint smell, and has a slightly saline taste. It contains water in considerable quantity; albumen; chloride of sodium; phosphate of lime; an alkaline substance; and a particular acid. It facilitates the dilatation of the uterus, and aids delivery by acting as a soft wedge enclosed in its membranes, *Bag of waters, Poche des Eaux, &c.* It is probably inservient to useful purposes in the nutrition of the fetus.

LIQUOR AMNII, FALSE. The fluid contained between the amnion and chorion in the early periods of foetal existence.

LIQUOR ANODYNE, HOFFMANN'S, *Spiritus ætheris sulphurici compositus*—*l. Anodynus martialis, Alcohol sulphurico-æthereus ferri.*

LIQUOR ANODYNUS TEREBINTHINATUS. A formula prescribed by Rademacher in cases of gallstone, and of obstructions and indurations of the liver and spleen. It was composed of Hoffmann's anodyne liquor ʒi; rectified oil of turpentine ʒij. Dose, 5 to 10 drops. It resembles the Remède de Durand, *Mixture lithontriptique de Durand, M. de Whitt, Éthérol d'essence de térébenthine*, which

is formed of Sulphuric ether, 30 parts: *Essence of turpentine*, 15 parts. It is used in the same cases.

LIQUOR ARSENICALIS, *L. Potassæ arsenitæ* (Ph. U. S.), *Solutio arsenicalis seu arsenicata seu arsenitæ kalicæ, Arsenical solution, Mineral solvent, Arsenic potassæ liquidus seu potassæ aquosus, Fowler's solution of arsenic, Solvent minéral, Italian poison, Acqua seu Aqua Toffana, Acqua della Toffana, A. di Napoli, Acquetta* [?], *Tasteless aque drop, (F.) Liqueur arsenicale.* (Acid. Arsenios. in frustulis, potassæ carbonatis pur., sing. gr. lixiv, aquæ destillat. q. s. Boil together the arsenious acid and carbonate of potassa with twelve fluidounces of distilled water, in a glass vessel, until the arsenic is dissolved. When the solution is cold, add Spirit. lavand. o. fʒiv, and as much distilled water as will make the whole one pint. Ph. U. S.) fʒj contains gr. ss of the arsenious acid. Dose, gtt. x.

For Pearson's Solution, see Arseniate of Soda.

LIQUOR ARSENICI CHLORIDI, *Solution of Chloride of Arsenic, of the London Pharmacopoeia of 1851,* is made by boiling Arsenious acid ʒss, in hydrochloric acid fʒiiss (Imp. meas.), mixed with a fluidounce of distilled water, until it is dissolved; then adding water to make the quantity a pint (Imp. meas.) Each fluidounce contains a grain and a half of arsenious acid, or an equivalent quantity of terchloride. Properties the same as those of Liquor Arsenicallis. It is a substitute for, and imitation of *De Valangin's Arsenical Solution, Solutio solven'tis mineralis*, much used by many British practitioners. Dose, gtt. v-x.

LIQUOR ARSENICI ET HYDRARGYRI IODIDI, see Arsenic and Mercury, iodide of—*l. Barii chloridi, Baryta, muriate of, solution of*—*l. Bellotii, L. Hydrargyri nitrici*—*l. Calci chloridi, see Calcis murias.*

LIQUOR CALCIS, *Solutio calcis, Aqua calcis seu benedicta seu Calcaria usta, Calcaria pura liquidæ, Solution of Lime, Lime Water, (F.) Eau de chaux. (Calcis ʒiv, eq. destill. cong. Pour the water on the lime, and stir. Let it stand in a covered vessel three hours; bottle the lime and water in stopped bottles, and use the clear solution.)* It is astringent, tonic, and antacid; and is used in diarrhœa, diabetes, heartburn, &c., and as a lotion to foul and cancerous ulcers, &c. Dose, fʒij to Oss, in milk.

LIQUOR CALCIS CHLORINATÆ, *Solution of Chlorinated Lime* (Ph. D.), is made by triturating half a pound of chlorinated lime with half a gallon of water; shaking the mixture; filtering; and keeping in a well-stopped bottle. Used in the same cases as chlorinated lime. Dose, fʒss to fʒj.

LIQUOR CALCIS COMPOSITUS, *Aqua Calcis composita seu benedicta composita, Compound lime water.* (F.) *Eau de chaux compositæ. (Liqu. quaiac. ras. lbss, rad. glycyrrh. ʒi, cort. saussfras, ʒss; semin. coriand. ʒij, liquor calcis, Ovj. Macerate for two days, and filter.)* It is stimulant, diaphoretic, and astringent, and is used in cutaneous affections.

LIQUOR CALCIS MURIATIS, see Calcis murias—*l. Cereris, Cerevisia*—*l. Chloreti natri, L. sodæ chlorinatæ*—*l. Chlorini, see Chlorine*—*l. Chlorureti natri, L. Sodæ chlorinatæ*—*l. Chlorureti sodæ, L. sodæ chlorinatæ.*

LIQUOR CHYLI, see Chyle.

LIQUOR CUPRI AMMONIATII, *Liquor Cupri ammonio-sulphatis, Aqua seu solutio cupri ammoniacæ seu sappharina, Blue eyewater, Solution of ammoniated copper; (F.) Liqueur ou Eau de cuivre ammoniacale. (Cupri ammoniat. ʒi, aquæ destill. Oj. Dissolve and filter the solution through paper. Ph. L.) Corrosive and detergent. Used externally to foul ulcers; and diluted with an equal part of distilled water, it is applied by means*

—a hair pencil to specks and films on the eye.

LIQUOR CUPRI AMMONIO-SULPHATIS, Liquor Cupri ammoniati.

LIQUOR CUPRI SULPHATIS COMPOSITUS, *Aqua cupri vitrioli et composita*. (Cupri sulphat., alumin. sulphat. āā ʒij , aqua purā Oij , acid sulph. ʒij . Boil the salts in the water until they are dissolved; then filter the liquor, and add the acid.) Used as an astringent in epistaxis, &c. It was also called *Aqua Styptica*.

LIQUOR CYRENIACUS, Benjamin—1. Entericus, Succus entericus—1. Excitans, Spiritus ammonis succinatus.

LIQUOR FERRI ALKALINI, *Solution of Alkaline Iron*, (F.) *Liqueur de fer alcaline*. (Ferri ʒiiss , acid. nitric. ʒij , aqua destillat. fʒxj , liq. potass. eubcarb. fʒvj . To the acid and water mixed, add the iron; and, after the effervescence, add the clear solution, gradually, to the liq. potassæ eubcarb.; shaking it occasionally till it assumes a deep brown-red colour, and the effervescence stops. After six hours' settling, pour off the clear solution. Ph. L.) It is tonic like other preparations of iron. Dose, fʒss to fʒiss .

LIQUOR FERRI IODIDI, *Solution of Iodide of Iron*, *Syrupus Ferri iodidi*, *Syrup of Iodide of Iron*. (Iodin. ʒij , Ferri rament. ʒj , Sacchar. pulv. ʒxij , Aqua destillat. q. s. Mix the iodine with fʒx of the distilled water, in a porcelain or glass vessel, and gradually add the iron filings, constantly stirring. Heat the mixture gently until the liquor acquires a light greenish colour; then, having added the sugar, continue the heat a short time, and filter. Lastly, pour distilled water upon the filter, and allow it to pass until the whole of the filtered liquor measures twenty fluidounces. Keep the solution in closely stopped bottles.—Ph. U. S.) Dose, 20 or 30 drops.

LIQUOR FERRI MURIATIS, *Tinctura ferri muratis*.

LIQUOR FERRI NITRATIS, *L. Ferri Sesquinitratæ seu ternitratæ*, *Solutio Ferri nitratæ*. *Solution of nitrate, of ternitrate of sesquioxide, or of sesquinitrate of iron*, has been recommended in chronic diarrhoea and dysentery. Its virtues exactly resemble those of chloride of iron. It is prepared as follows:—Ferri ʒij , incis. ʒj , Acid. nitric. fʒij , Aq. destillat. q. s. Mix the acid and a pint of distilled water, until gas ceases to be given off; filter, and add distilled water to make fʒxxx .—Ph. U. S.) Dose, 10 to 20 drops.

LIQUOR FOLLICULI, see Folliculi Graafiani.

LIQUOR FUMANS BOYLEI, *Sulphuretum Ammoniacæ Hydrogenatum*, *Hydrosulphuretum Ammoniacæ*, *Tinctura Sulphuris Volatilis*, *Aqua Sulphureti Ammoniacæ*, *Boyle's Fuming liquor*, (F.) *Liqueur fumante de Boyle*. It is possessed of nauseating and emetic properties, and has been given in diabetes and diseases of excitement as a deoxygenizer!

LIQUOR FUMING, *BOYLE'S*, *Liquor fumans Boylei*—1. Fuming, of Libavius. Tin, muriate of—1. Genital, Sperm—1. of Hartshorn, volatile, *Liquor volatilis cornu cervi*—1. Hydrargyri Bichloridi, *Liquor hydrargyri oxymercuriatis*—1. Hydrargyri chloridi corrosivi, *L. hydrargyri oxymercuriatis*—1. Hydrargyri et arsenici iodidi, see Arsenic and Mercury, iodide of.

LIQUOR HYDRARGYRI NITRICI, *L. Bellotti*, (F.) *Eau mercurielle ou contre la gangrène*, *Remède du Duc d'Antin*, *R. du Capucin*, (Hydrarg. 120 p., acid. nitric. (33°) 150 p. Dissolve, and add to the solution distilled water, 900 p. Ph. P.) Dose, two or three drops in a glass of water. Not much used.

LIQUOR HYDRARGYRI OXYMURIATIS, *L. Hydrargyri Chloridi corrosivi seu Hydrargyri bichloridi*, *Solution of Oxymercuriate of Mercury*,

Liquor Sicietenis seu Syphiliticus Turneri. *Muriæ hydrargyri spiritalis liquidus*, *Solutio Muriæ hydrargyri oxygenati*, *Liquor or Solution of Corrosive Sublimate or of Van Sieten*. (Hyd. oxyg., ammon. muriat. āā gr. x, aquæ destill. Oj , (Imp. meas.) Dissolve in the water and add the spirit. Ph. P.) Dose, fʒss , or fʒj .

Norton's Drops, *Green's Drops*, and *Solomon's Anti-impetiginæ*—all nostrums—seem to be disguised solutions of Corrosive Sublimate.

LIQUOR HYDRARGYRI PERNITRATIS, see Hydrargyri nitras—1. Hydrargyri Supernitratæ, see Hydrargyri nitras—1. Hydriodatis Arsenici et Hydrargyri, Arsenic and mercury, iodide of—1. Intestinalis, Succus entericus.

LIQUOR IODINI COMPOSITUS, (Ph. U. S. 1842,) *L. Iodini compositus*, (Ph. U. S. 1851,) *Solutio Potassii Iodidi Ioduræta*, *Compound Solution of Iodine*, *Lugol's Solution*. (Iodin. ʒvj , Potass. iodid. ʒiss , Aqua destillat. Oj . Dissolve the iodine and iodide of potassium in the water.—Ph. U. S.) Dose, gtt. vj , ad xij , in sugared water.

LIQUOR LITHARGYRI SUBACETATIS, *L. plumbi subacetatis*—1. Lithargyri subacetatis compositus, *Liquor plumbi subacetatis dilutus*.

LIQUOR LYMPE, see Lymph.

LIQUOR OF MONRO, *Solution of Monro*. A solution, used by Monro for the preservation of anatomical preparations. It was composed of alcohol at 22° or 24°, with a drachm of nitric acid to each pint.

LIQUOR MORGAGNI. The small quantity of fluid contained within the capsule of the crystalline lens.

LIQUOR MORPHÆ ACETATIS, *Solution of Acetate of Morphia*, *Anodyne Drops*. (Acetate of morphia, ʒiv , distilled water, Oj . (Imp. meas.), Spirit. tenuior. Oss. (Imp. meas.) Ph. L.) *acetic acid*, mxxv . Dose, from six to twenty-four drops.

LIQUOR MORPHÆ MURIATIS, *Solution of Muriate of Morphia*. (Morphiæ muriat. ʒiv , aq. destillat. Oj . (Imp. meas.) Sp. tenuior, Oss. (Imp. meas.) Ph. L.) Dose, from 6 to 20 minims.

LIQUOR MORPHÆ SULPHATIS, *Solution of Sulphate of Morphia*. (Morphiæ sulphat. gr. viij , aqua destillat. Oss. Dissolve the sulphate of morphia in the water.—Ph. U. S.) Dose, fʒj to fʒij —containing from an eighth to a quarter of a grain.

LIQUOR MUCI, see Mucus—1. Natri Oxymuriatici, *L. Sodæ chlorinatæ*—1. Nervinus Bangii, *Tinctura æthereæ camphorata*—1. Oleosus Sylvii, *Spiritus ammoniacæ aromaticus*—1. Opii sedativus, (Haden's), see Tinctura opii.

LIQUOR OPII SEDATIVUS. An empirical preparation by a London druggist of the name of Battley. It is said to be an aqueous solution of opium, evaporated to dryness to get rid of the acid resin, re-dissolved in water, and a small portion of alcohol added to give it permanence.—Redwood. It is devoid of many of the narcotic effects of opium.

LIQUOR OVI ALBUS, *Albumen ovi*—1. Pancreaticus, see Pancreas—1. Pericardii, see Pericardium—1. Plumbi acetatis, *L. P. subacetatis*—1. Plumbi diacetatis, *Liquor Plumbi subacetatis*.

LIQUOR PLUMBI SUBACETATIS, *L. Subacetatis Lithargyri seu Plumbi Acetatis seu Plumbi Diacetatis*, *Gould's Extrac'tum Saturni*, *Lithargyri Acetum*, *Acetum plumbicum seu saturninum*, *Solution of Subacetate of Lead*, (F.) *Liqueur de sous-acétate de Plomb*. (Plumb. acet. ʒxvj , Plumb. oxid. semivitr. in pulv. subtil. ʒixsa , aq. destillat. Oiv . Boil together in a glass or porcelain vessel, for half an hour, occasionally adding distilled water, so as to preserve the measure. Filter through paper, and keep the solution in closely-stopped bottles.—Ph. U. S.) It is

used externally as a cooling astringent, and discontinue, when diluted with distilled water.

LIQUOR PLUMBI SUBACETATIS DILUTUS, *L. Subacetatis Lithargyri Compositus*, *L. Plumbi acetatis dilutus*, *L. Plumbi Diacetatis dilutus*, *Acetas Plumbi dilutus alcoholici*, *Diluted Solution of Subacetate of Lead*, *Aqua Saturni seu regl'eto-mineralis seu Plumbi seu Goulard's seu Lithargyri Acetati composita*, *Tinctura Plumbosa*, *Lead water*, *Goulard water*, (F.) *Eau 'anche*, *Eau de Goulard*, *Eau végétominérale*, *White Wash*, *Royal Preventive*. (*Liq. plumbi subacet. f3ij*, *aquæ destillat. Oj*. Ph. U. S.) Properties the same as the last, but feebler.

LIQUOR POTASSÆ, *Aqua Potassæ seu Kali Caustici seu Kali puri*, *Solution of Potash or of Potassa*, *Lixivium magistralis seu Saponarium*, *Soap Lees*, *Soap Ley*, *Lixivium causticum*, *Potassæ liquida*, (F.) *Eau, solution ou liqueur de Potasse*, *Potasse liquide*, *Lessive des Savonniers*. (*Potassæ carb. lbj*, *calcis. lbss*, *aquæ destill. fervent. congium*. Dissolve the alkali in Oij of the water, and add the remainder of the lime. Mix the whole: set aside in a close vessel, and, when cold, filter through calico.) It is antilithic in cases of uric acid calculi, and antacid. Externally, stimulant and escharotic. Dose, gtt. x to xx.

LIQUOR POTASSÆ ARSENITIS, *L. arsenicalis* — *l. Potassæ Carbonatis*, *L. P. Subcarbonatis*.

LIQUOR POTASSÆ CITRATIS, *Solution of Citrate of Potassa*, *Neutral Mixture*, *Saline Mixture*. (*Succ. Limon. Oss. Potassæ Bicarbonat. q. s.*) saturate by the carbonate of potassa, and filter; or, *Acid. Citric. 3ss*; *Ol. Limon. M. ij*; *Aquæ Oss. Potassæ Bicarbonat. q. s.*; dissolve, saturate by the carbonate of potassa, and filter. Ph. U. S.) Used in fever, but probably of little or no efficacy.

LIQUOR POTASSÆ SUBCARBONATIS, *L. Potassæ Carbonatis* (Ph. U. S.), *Aqua Kali seu Kali præparati seu Subcarbonatis Kali*, *Oleum Tartari per deliquium*, *Lixivium Tartari*, *Oil of Tartar*, *Saline oil of Tartar*, *Solution of Subcarbonate of Potassa*, (F.) *Liqueur de sous-carbonate de Potasse*, *Lessive de Tartre*, *Huile de Tartre par défaillance*. (*Potassæ subcar. lbj*, *aquæ destillat. f3xij*. Dissolve and filter. Ph. U. S.) Dose, gtt. x to xxx.

LIQUOR POTASSII IODIDI COMPOSITUS. *Compound Solution of Iodide of Potassium*. (*Potassii iodid. gr. x*, *Iodin. gr. v*, *aquæ destillat. Oj*. (Imp. meas.) Ph. L.) Dose, f3j.

LIQUOR PROPAGATORY, *Sperm—l. Prostaticus*, *Prostatic liquor*, see *Prostate—l. Puris*, see *Pus*.

LIQUOR SANGVINIS. *Blood liquor*, (F.) *Liqueur du Sang*. A term given by Dr. B. Babin to one of the constituents of the blood, the other being the red particles. It is the *Hæmatoplasma*, the *Plasma*, of Schultz, *Lymph*, *Coagulable or plastic Lymph*, the *Mucago* or *Mucilage* of Harvey, *Hewson* and others, and the *intercellular fluid* of Lehmann, (F.) *Lympe plastique ou coagulable*, *Suc nourricier*, and is the effused material from which the cells obtain the constituents of the different tissues and secretions. See *Blood*.

LIQUOR OF SCARPA, *Vitrine auditive*.

LIQUOR SEMINIS. The homogeneous, transparent fluid, in which the spermatozooids and seminal granules are contained. See *Sperm*.

LIQUOR SODÆ, *L. Sodæ Causticæ*, *Solution of Soda*, *Solution of Caustic Soda*. (Prepared, like *Liquor Potassæ*, from Carbonate of Soda.)

LIQUOR SODÆ CARBONATIS, *Solution of Carbonate of Soda*. (*Sodæ carbon. 3iss*, (avoirdupois); *Aquæ destillat. Oj*. (Imp. meas.) Ph. D.) Dose, as an antacid, one to two tablespoonfuls, diluted with water.

LIQUOR SODÆ CHLORIDI, *L. sodæ chlorinatæ*.

LIQUOR SODÆ CHLORINATÆ, *L. sodæ chlo'ridi*

seu sodæ oxymuriat'icæ seu chlore'ti natri seu chlorure'ti natri seu chlore'ti sodæ seu chlorure'ti sodæ, *seu natri oxymuriat'ici*, *Natrum chloratum liq'idum*, *Aqua natri oxymuriat'ici*, *Labarraque's Disinfecting Liquid*, *Solution of Chlorinated Soda*. (*Calcis Chlorinat. lbj*; *Sodæ Carbonat. lbj*; *Aquæ cong. iss*. Dissolve the carbonate of soda in three pints of the water, with the aid of heat. To the remainder of the water add, by small portions at a time, the chlorinated lime, previously well triturated, stirring the mixture after each addition. Set the mixture by for several hours, that the dregs may subside; decant the clear liquid, and mix it with the solution of carbonate of soda. Lastly, decant the clear liquor from the precipitated carbonate of lime, pass it through a linen cloth, and keep it in bottles secluded from the light.—Ph. U. S.) Used in the same cases as the chloride of lime. Internally, 10 drops to a fluidrachm, for a dose. Diluted with water, it is an exciting and disinfectant in various morbi externi.

LIQUOR SODÆ EFFERVESCENS, *Acidulous water*, simple—*l. Sodæ Oxymuriatæ*, *L. sodæ chlorinatæ*—*l. Stypticus Ruspini*, *Styptic*, *Ruspini's*—*l. Sulphuricus Alcolisatus*, *Spiritus ætheris sulphurici*—*l. Swietenis*, *L. hydrargyri oxymuriatis*—*l. Syphiliticus Turneri*, *L. hydrargyri oxymuriatis*—*l. Tartari emetici*, *Vinum antimonii tartarizati*—*l. of Van Swieten*, *L. hydrargyri oxymuriatis*.

LIQUOR VOLATILIS CORNU CERVI, *L. volatilis Cornu Cervi*, *Volatilis osseum*, *Volatile Liquor of Hartshorn*, *Spiritus Lumbricorum seu Milipedarum seu Cornu Cervi*, *Hartshorn*, *Spirit of Hartshorn*, *Bone Spirit*, (F.) *Liqueur volatile de Corne de cerf*. This is a solution of subcarbonate of ammonia, impregnated with empyreumatic oil. It possesses the same virtues as the subcarbonate of ammonia. It is in common use to smell at, in faintings, &c.

LIQUOR VOLATILIS OSSIIUM, *L. volatilis cornu cervi*—*l. Zinci Chloridi*, see *Burnett's Disinfecting Liquid*.

LIQUOR ZINCI SULPHATIS CUM CAMPHORÆ, *Aqua Zinci vitriolat'i cum Camphorâ seu vitriol'ica camphorata seu ophthalmica*, *Common Eye Water*. (*Zinci sulph. 3ss*, *camphor. ʒij*, *aq. bullient. Oij*; dissolve and filter.) Used as a lotion for ulcers; or, diluted with water, as a collyrium.

LIQUORICE, *Glycyrrhiza*—*l. Bush*, *Abrus precatorius*—*l. Juice*, see *Glycyrrhiza*.

LIQUORICE, *REFINED*, *Extractum glycyrrhizæ*—*l. Spanish*, see *Glycyrrhiza*—*l. Wild*, *Aralia nudicaulis*, *Galium circeazans*.

LIQUORITIA, *Glycyrrhiza*.

LIRIODENDRON, from *λίρον*, 'lily or tulip,' and *δένδρον*, 'tree.' *L. tulipif'era*, *Tulipif'era Liriodendron*, *Old wife's shirt*, *Tulip* or *Poplar Tree*, *Tulip-bearing* or *American Poplar*, *White Wood*, *Cypress Tree*, (New England.) (F.) *Tulipier*. The bark—*Liriodendron* (Ph. U. S.)—especially of the root, of this noble forest tree, which is indigenous in the United States, is a strong aromatic bitter, and has been employed advantageously as a tonic. An active principle was separated from it by Professor J. P. Emmet of the University of Virginia, and has been called *Lirioden'drin*. It is not used in medicine.

LIS BLANC, *Lilium candidum*—*l. Asphodellæ*, *Asphodelus ramosus*—*l. des Étangs*, *Nymphæa alba*.

LISERON, *GRAND*, *Convolvulus sepium*—*l. des Haies*, *Convolvulus sepium*—*l. Mechameek*, *Convolvulus panduratus*.

LISK, *Inguen*.

LISTS, *Flanks*.

LISTON'S ISINGLASS PLASTER, see Spadurapum adhasivum.

LITE, λιτῆρ. A plaster, formerly made of verdigris, wax, and resin.—Galen.

LITHAGOGECTASIA, Lithectasy.

LITHAGO'GUM, from λιθος, 'a stone,' and γυω, 'I expel.' A remedy which was supposed to possess the power of expelling calculi. Also, a lithotomy forceps.

LITHANTHRAX, Carbo fossilis.

LITHANTHROKOKALI, Anthrakokali.

LITHARGE, Plumbi oxydum semivitreum—I. of Gold, see Plumbi oxydum semivitreum—I. of Silver, see Plumbi oxydum semivitreum.

LITHARGYRI ACETUM, Liquor plumbi subacetatis.

LITHARGYRUM, Plumbi oxydum semivitreum.

LITHARGYRUS, Plumbi oxydum semivitreum.

LITHAS, Urate.

LITHATE, Urate—I. of Soda, Urate of soda.

LITHECT'ASIA, Lithectasia, Lithagogaectasia, Lithodectasia, from λιθος, 'a stone,' and εκτasis, 'dilatation,' Cystectasy. An operation which consists in extracting stone from the bladder by dilating the neck of the organ, after making an incision in the perineum, and opening the membranous portion of the urethra.

LITH'IA, Lithiasis, Lithogenia, Uriasis, Urolithiasis, Catech'ia calculosa, Calculi Morbus, Lapillatio, Generatio calculi, from λιθος, 'a stone.' The formation of stone, gravel, or concretions in the human body. Also, an affection in which the eyelids are edged with small, hard, and stone-like concretions.

LITH'IA, CARBONATE OF, Lithia Carbonas, Lithium Carbonicum, (F.) Carbonate de Lithine. A salt found in certain mineral waters, which have been serviceable in lithuria. Hence, it has been suggested in that morbid condition.

LITHIA RENALIS ARENOSA, Gravel—I. Renalis, Nephrolithiasis—I. Vesicalis, Calculi, vesical.

LITHIAE CARBONAS, Lithia, carbonate of.

LITHIASIS, Lithia—I. Cystica, Calculi, vesical—I. Nephritica, Gravel, Nephrolithiasis.

LITHIASIS PULMONUM, Pulmo'nes tartarizati. The formation of concretions in the lungs, occasioning at times Litho-phthiasis, Phthisis calculosa, Phthisis calculeuse, of Bayle.

LITHIASIS RENALIS ARENOSA, Gravel—I. Renalis, Nephrolithiasis—I. Vesicalis, Calculi, vesical.

LITHIC, Lithicus. Belonging to lithic or uric acid, or to stone: hence Lithic Diathesis. Also, an antilithic.

LITHIC ACID, Uric acid—I. Acid diathesis, Lithuria—I. Diathesis, Lithuria—I. Sediments, see Lithuria.

LITHINE, CARBONATE DE, Lithia, carbonate of.

LITHIUM CARBONICUM, Lithia, carbonate of.

LITHIURIA, Lithuria.

LITHOCENOSIS, Lithotripsy.

LITHOCYSTOTOMY, Lithotomy.

LITHOCLAST, see Lithotripsy.

LITHOCLASTIA, Lithotripsy.

LITHODECTASIA, Lithectasy.

LITHODIALYSIS, from λιθος, 'a stone,' and διαλυω, 'to dissolve.' The dissolving of a stone in the bladder.

LITHODRAS'SIC, Lithodras'sicus, (F.) Lithodras'sique, from λιθος, 'a stone,' and δρασσειν, 'to seize hold of.' An epithet given to a form of stone forceps—Pince Lithodras'sique—used in the operation of lithotripsy, by MM. Meirieu and Tanchou.

LITHOGENIA, Lithia.

LITHOID, Lithoides, Lithoïdes; from λιθος, 'stone,' and εidos, 'resemblance.' Of the nature of stone, or resembling stone: as

LITHOIDES OS, see Temporal bone.

LITHOLABE (F.), Litholabum, from λιθος, 'a stone,' and λαβω, λαβανω, 'I seize hold of.' An instrument, employed for laying hold of a stone in the bladder, and keeping it fixed, so that lithotritic instruments can act upon it.

LITHOLABON, Forceps, (Lithotomy.)

LITHOLABUM, from λιθος, 'a stone,' and λαβανω, 'I seize.' An instrument concerned in extracting stone from the bladder. It had various shapes.—Fabricius ab Aquapendente, Hildanus.

LITH'OLYTE, from λιθος, 'a stone,' and λυω, 'I dissolve.' An instrument, of the catheter kind, for conveying solvents of calculi into the bladder.

LITHOLYTIC, Lithotriptic.

LITHOME'TRA, from λιθος, 'a stone,' and μετρα, 'the uterus.' Osseous, or other concretions of the uterus.

LITHONEPHRITIS, see Nephritis.

LITHONLYTIC, Lithotriptic.

LITHONTHRYPTIC, Lithotriptic.

LITHONTRIP'TIC, Lithontrip'ticus, Lithon-thryp'tic, Litholytic, Lithonlytic, Calculifragus, Sacifragus, from λιθος, 'a stone,' and θρυπτω, 'I break in pieces.' A remedy believed to be capable of dissolving calculi in the urinary passages. There is not much reliance to be placed upon such remedies. By antilithics, exhibited according to the chemical character of the calculus (see Calculi, urinary,) the disease may be prevented from increasing; but most of the vaunted lithontrip'tics for dissolving the calculus already formed have been found unworthy of the high encomiums which have accompanied their introduction.

LITHOP'EDION, Infans lapideus, Osteop'edion, from λιθος, 'a stone,' and παις, παίδος, 'a child.' A foetus, petrified in the body of the mother.

LITHOPH'AGUS, from λιθος, 'a stone,' and φάγω, 'I eat.' A stone-eater. Cases are on record of persons who have swallowed stones for purposes of exhibition.

LITHOPHTHISIS, see Calculi, pulmonary, and Lithiasis pulmonum.

LITHOPRINE, Lithotripsy.

LITHOPRIONE, from λιθος, 'a stone,' and πριων, 'a saw.' An instrument proposed by M. Leroy for preventing the fragments of a calculus, when subjected to lithotripsy, from falling into the bladder. It is a variety of litholabe.

LITHOPRISIE, Lithotripsy.

LITHORINEUR, from λιθος, 'a stone,' and ρινειν, 'to file.' An instrument, proposed by MM. Meirieu and Tanchou for filing down calculi in the bladder.

LITHOS, Calculus.

LITHOSPERMUM OFFICINALE, Milium Solis, Egon'ychon, Gramvell, Bastard Alkanet, Little-wale, (F.) Grémil officinal, Herbe aux Perles. Ord. Boraginæ. The seeds of this plant were formerly supposed, from their stony hardness, (λιθος, 'a stone,' and σπέρμα, 'seed,') to be efficacious in calculous affections. They have, also, been considered diuretic.

LITHOSPERMUM VILLOSUM, Anchusa tinctoria.

LITHOTERE'THRUM, from λιθος, 'stone,' and τερεν, 'to rub.' A lithotritor.

LITHOTHLIBIE (F.), from λιθος, 'stone,' and θλιβω, 'I crush.' The crushing of a friable calculus between a finger, passed by the rectum or vagina as far as the bas-fond of the bladder, and a catheter introduced into the bladder.

LITHOTHRYPsis, Lithotripsy.

LITHOTHRYPTORS, see Lithotrixy.

LITHOTOME, *Lithotomus*, from *λίζω*, 'a stone,' *τέμνω*, 'I cut,' and *τομή*, 'an incision.' This name has been given to a number of instruments of different shapes and sizes, which are used in the operation for the stone, to cut the neck or body of the bladder. They ought, with more propriety, to be called *Cystotomes*.

The *Lithotome Caché* of Frère Côme is the most known, and is still occasionally used. It is composed of a handle, and a flattened sheath, slightly curved: in this there is a cutting blade, which can be forced out, by pressing upon a *bascule* or lever, to any extent that may be wished by the operator.

A *Double Lithotome* was used by Dupuytren in his bilateral operation. See Lithotomy.

LITHOTOMIST. Same etymon. *Lithotomus*. One who devotes himself entirely to operating for the stone. One who practises lithotomy.

LITHOTOMY, *Lithotomia*, *Cystotomia*, *Urolithotomia*, *Sectio vesicalis*, *Lithocystotomy*. Same etymon. (F.) *Taille*. The operation by which a stone is extracted from the bladder. The different methods, according to which this operation may be practised, are reducible to five principal; each of which has experienced numerous modifications.

1. *The Method of Celsus*, *Methodus Celsiana*, *Cystotomia cum apparatu parvo*, *Apparatus Minor*, *Cutting on the Gripe*. This consisted in cutting upon the stone, after having made it project at the perinæum by means of the fingers introduced into the rectum. This method was attended with several inconveniences; such as the difficulty of dividing the parts neatly, injury done to the bladder, as well as the impossibility of drawing down the stone in many persons. It is sometimes, also, called *Methodus Guytoniana*; from Guy de Chauliac having endeavoured to remove from it the discredit into which it had fallen in his time. It was termed *Apparatus Minor*, (F.) *Le petit appareil*, from the small number of instruments required in it.

2. *Apparatus Major*. This method was invented, in 1520, by John de Romani, a surgeon of Cremona, and communicated by him to Mariano-Santo-di-Barletta, whence it was long called *Mariano's Method*, *Sectio Mariana*. It was called, also, *Apparatus Major*, and *Cystotomia seu Methodus cum apparatu magno*, (F.) *Le grand appareil*, from the number of instruments required in it. An incision was made on the median line; but the neck of the bladder was not comprehended in it. It was merely dilated. The greater apparatus was liable to many inconveniences, such as ecchymoses; contusion; inflammation of the neck of the bladder; abscesses; urinary fistulæ; incontinence of urine; impotence, &c.

3. *The High Operation*, *Apparatus altus*, *Cystotomia cum apparatu alto*, *C. Hypogastrica*, *Epicystotomia*, *Laparocystotomia*, *Sectio seu Methodus Franconiana*, *S. Hypogastrica seu alta*, (F.) *Haut appareil*, *Taille Hypogastrique*, *Taille sus-pubienne*, was first practised by Peter Franco, about the middle of the 16th century. It consisted in pushing the stone above the pubis by the fingers introduced into the rectum. Rousset afterwards proposed to make the bladder rise above the pubis by injecting it. The method had fallen into discredit, when Frère Côme revived it. It is used when the calculus is very large. It was practised by opening first the membranous part of the urethra upon the catheter passed into the canal. Through this incision, the *Sonde à dard*—a species of catheter, having a spear-pointed stilet—was introduced into the bladder.

An incision was then made into the *linea alba*, above the symphysis pubis, of about four or five fingers' breadth, and the peritoneum detached to avoid wounding it. The stilet was pushed through the bladder, and used as a director for the knife, with which the bladder was divided anteriorly, as far as the neck; and the stone extracted. It was performed in England by Douglass, in 1719, and since by others, with various modifications.

4. *The Lateral Operation*, *Hypocysteotomia*, *Cystotomia lateralis*, *Cystauchenotomia*, *Cystotrachelotomia*, *Urethrocystauchenotomia*, *Urethrocysteotrachelotomia*, *Sectio lateralis*, *Apparatus lateralis*, (F.) *Appareil latéralisé*, so named from the prostate gland and neck of the bladder being cut laterally, was probably invented by Peter Franco. It was introduced into France by Frère Jacques de Beaulieu. He performed it with rude instruments, invented by himself, and improved by the suggestions of some of the Parisian surgeons. In England, it received its earliest and most important improvements from the celebrated Cheselden. It is the method practised at the present day, according to different modes of procedure. In this method, the patient is placed upon a table; his legs and thighs are bent and separated; the hands being tied to the feet. The perinæum is then shaved, and a staff is introduced into the bladder: the handle being turned towards the right groin of the patient. An oblique incision is now made from the raphe to the middle of a line drawn from the anus to the tuberosity of the ischium of the left side; and, taking the staff for a guide, the integuments, areolar tissue of the perinæum, membranous portion of the urethra, transversus perinæi muscle, bulbo-cavernosus, some fibres of the levator ani, the prostate and neck of the bladder, are successively divided. For this latter part of the operation, the knife, the beaked bistoury, *Bistouri* ou *Lithotome Caché*, cutting gorget, &c., is used, according to the particular preference. The forceps are now introduced into the bladder, and the stone extracted. In the operation, care must be taken not to injure the rectum, or the great arterial vessels, distributed to the perinæum.

A variety of the *Lateral Apparatus*, called by the French *Appareil latéral*, consisted in cutting into the *bas-fond* of the bladder, without touching the neck of that organ; but it was soon abandoned, on account of its inconveniences.

The method of Le Cat and of Pajola—*Urethrocysteo-aneurysmatotomia*—consists in dividing the prostate in part only,—the enlargement of the wound being effected by a peculiar dilator.

The *Bilateral Operation* is founded on that of Celsus. It consists in making an incision posterior to the bulb of the urethra, and anterior to the anus, involving both sides of the perinæum by crossing the raphe at right angles: an incision is then made through the membranous part of the urethra, and the prostate may be cut bilaterally, either with the double lithotome of Dupuytren, or the prostatic bisector of Dr. Stevens, of New York.

5. *Lithotomy by the Rectum*, *Proctocystotomia*, *Sectio recto-vesicalis*, (F.) *Taille par la Rectum*, *Taille postérieure*, *T. Recto-vésicale*. This was proposed by Vegetius in the 16th century; but it was never noticed until M. Sanson, in the year 1817, attracted attention to it; since which time it has been successfully performed in many instances. It consists in penetrating the bladder through the paries corresponding with the rectum, by first cutting the sphincter ani and rectum about the root of the penis, and penetrating the

bladder by the neck of that organ, dividing the prostate,—or by its *bas-fond*.

Lithotomy in women, from the shortness of the urethra, is a comparatively insignificant operation.

LITHOTOMY BY THE RECTUM, see Lithotomy—1. by the Vagina, see Lithotomy.

LITHOTRESIS, Lithotrity.

LITHOTRIPSIS, Lithotrity.

LITHOTRIPSY, Lithotrity.

LITHOTRIPATORS, see Lithotrity.

LITHOTRITES, see Lithotrity.

LITHOTRITEURS, see Lithotrity.

LITHOTRITOR, see Lithotrity.

LITHOTRITY, *Lithotritia*, *Lithotryp'sy*, *Lithotripsy*, *Lithotrip'sy*, *Lithotrypsis*, *Lithotres'sis*, *Lithotripsis*, *Lithocoeno'sis*, *Lithoclastia*, *Lithoprinie*, *Lithoprisie*, (F.) *Broyement de la Pierre*, from *lithos*, 'a stone,' and *tripein*, 'I break.' The operation of breaking, bruising or sawing the stone in the bladder. It has been performed, of late years, with success, by French, and, after them, by English and American surgeons. The instruments employed for this purpose are called, in the abstract, *Lithotrites*, *Lithotriteurs*, *Lithotritors*, *Lithotriptors*, *Lithotryptors*, *Lithotrypetæ* and *Lithoclasts*. The most celebrated are those of Civiale, Jacobson, Heurteloup and Weiss. See *Brise-Pierre articulé*; and *Percuteur à Marteau*.

LITHOTRYPETÆ, see Lithotrity.

LITHOXIDURIA, from *lithos*, 'a stone,' *oxide*, and *urine*, 'urine.' The discharge of urine containing lithic or xanthic oxide.

LITHURIA, *Lithuria*, *Lithure'sis*, *Lithourrhée* (Piorry); from *lithos*, 'a stone,' and *urine*, 'urine.' *Lithic*, or *Lithic Acid*, *Diathesis*. The condition of the system and of the urine in which deposits of lithic acid and the lithates—*Lithic sediments* or *concretions*—take place in the urine. See *Urine*.

LITHUS, Calculus.

LITMUS, Lichen roccella.

LITRA, Pound.

LITRE, *Litra*. A measure containing a cubed decimetre, which is equal nearly to 2.1135 pints. The ancients gave the name *litra*, *litra*, to a measure capable of containing 16 ounces of liquid.

LITSÆA CUBEBA, Piper cubeba—1. Pipe-rita, Piper cubeba.

LITTLE-WALE, *Lithospermum officinale*.

LITUS, Liniment.

LIVÊCHE, *Ligusticum levisticum*.

LIVER, Sax. *lisen*, *Hepar*, *Jecur*, *Jec'inus*, (F.) *Foie*. The liver is the largest gland in the body. It is an azymous organ; unsymmetrical; very heavy; and of a brownish-red colour; occupying the whole of the right hypochondrium, and a part of the epigastrium. Above, it corresponds to the diaphragm; below, to the stomach, transverse colon, and right kidney; behind, to the vertebral column, aorta, and vena cava; and before, to the base of the chest. Its upper surface is convex; the lower, irregularly convex and concave, so that anatomists have divided the organ into three lobes,—a large or right or colic lobe;—a lesser lobe, lobule, or inferior lobe, the *Lobulus Spigelii*,—and a middle or left lobe. At its inferior surface, are observed:—1. A Sulcus or Furrow or Fissure, called horizontal or longitudinal or great fissure, *Fossa Umbilicalis*, (F.) *Sillon horizontal*, *longitudinal* ou *de la veine ombilicale*, *Sulcus antero-posterior Jecoris* seu *horizontalis Jecoris* seu *longitudinalis Jecoris* seu *sinister Jecoris* seu *Umbilicalis*, which lodges, in the fœtus, the umbilical vein and ductus venosus. 2. The Principal Fissure, termed *Sulcus Transversus*, *Fossa transversa*, *Transverse fossa*, *Sinus Porta'rum*, *Porta*, *P. seu Manus Jecoris* seu *He'patis Fossa* or *Fissure of the Vena porta*, *Portal*

Fissure or *Fossa*, (F.) *Sillon transversal* ou *de la veine porte*, which receives the sinus of the vena porta. 3. The Fissure of the Vena Cava inferior, *Sillon de la veine cave inférieure*, situated at the posterior margin of the organ, and lodging the vena cava inferior. 4. The *Lobulus Spigelii*, or posterior portal eminence. 5. The anterior portal eminence, *Auriga* seu *Lobulus anon'ymus*. 6. Depressions corresponding to the upper surface of the stomach, gall-bladder, arch of the colon, right kidney, &c. Continued from the fossa umbilicalis is a small fossa, called *Fossa Ductus Venosi*, between the left lobe and *Lobulus Spigelii*. The posterior margin of the liver is very thick; much more so than the anterior. The liver is surrounded by a serous or peritoneal covering, which forms for it a suspensory or broad ligament and two lateral and triangular ligaments. See *Falx*. The blood-vessels of the liver are very numerous. The hepatic artery and vena porta furnish it with the blood necessary for its nutrition and the secretion of bile. The hepatic veins convey away the blood, which has served those purposes. The lymphatic vessels are very numerous; some being superficial; others deep-seated. The nerves are, also, numerous, and proceed from the pneumogastric, diaphragmatic, and from the hepatic plexuses.

The intimate structure of the parenchyma of the liver has been well studied. When cut, it presents a porous appearance, owing to the division of a multitude of small vessels. When torn, it seems formed of granulations;—the intimate structure of which has given rise to many hypotheses. In these granulations are contained the radicles of the excretory ducts of the bile; the union of which constitutes the hepatic duct. According to Mr. Kiernan, the intimate structure consists of a number of lobules—*hepatic lobules*, *hepatic islets*—composed of *intralobular* or hepatic veins, which convey the blood back that has been inservient to the secretion of bile. The *interlobular* plexus of veins is formed by branches of the vena porta, which contain both the blood of the vena porta and of the hepatic artery; both of which, according to Mr. Kiernan, furnish the pabulum of the biliary secretion. The biliary ducts form likewise an *interlobular plexus*, having an arrangement similar to that of the interlobular veins. Mr. Kiernan's views are embraced by many anatomists; but are denied by some.

The liver is the only organ, which, independently of the red blood carried to it by the hepatic artery, receives black blood by the vena porta. The general opinion is, that the vena porta is the fluid which furnishes bile, whilst that of the artery affords blood for the nutrition of the liver. It is probable, however, that bile is secreted from the blood of the latter vessel. Besides bile the liver forms sugar, and is a great assimilating organ.

The liver is liable to a number of diseases. The principal are—*Hepatitis* or inflammation, cancer, *biliary calculi*, *encysted* and other tumours or tubercles, *hydatids*, &c.; and it has, at times, been the fashion to refer to it as the cause of symptoms with which it is no wise connected.

LIVER, *Hepar*. Under this name the ancients designated several substances, having a brownish colour analogous to that of the liver; and composed of sulphur and some other body. See *Potassæ Sulphuretum*, *Liver of Sulphur*.

LIVER of Antimony is the semi-vitreous sulphuret, &c.

LIVER DISEASE, *Hepatopathia*—1. Fatty, *Adiposis hepatica*—1. *Gin*, *L. nutmeg*—1. *Gin-drinkers'*, *L. nutmeg*—1. Granulated, *Cirrhosis of the liver*.

LIVER-GROWN, *Tu'mido jecoræ præditus*. Having a large liver.

LIVER, HORNAIL, Cirrhosis of the liver—**L. Mammillated, Cirrhosis of the liver.**

LIVER, NUTMEG, *Hepar variegatum, Tubiform liver.* An appearance of the liver when cut across, resembling that of the section of a nutmeg, *Myristicatio hepatis*, supposed by some to be the result of intemperance in the use of alcoholic drinks; but occurring under other causes. The terms *whisky liver, gin-drinkers' liver*, and *gin liver*, occasionally applied to it, are, consequently, not distinctive.

LIVER SPOT, Chloasma.

LIVER, TUBERCULATED, Cirrhosis of the liver—**L. Tubiform, L. nutmeg.**

LIVER WEED, Hepatica triloba.

LIVER, WHISKY, Liver, nutmeg.

LIVER WORT, Hepatica triloba, Marchantia polymorpha—**L. Wort, ground, ash-coloured, Lichen caninus**—**L. Wort, Iceland, Lichen Islandicus**—**L. Wort, noble, Hepatica triloba.**

LIVIDUS MUSCULUS, Pectinalis.

LIVOR, Suggillation—**L. Emortualis, see Suggillation**—**L. Sanguineus, see Ecchymoma.**

LIVRE, Pound.

LIVRET, Omasum.

LIXIVIA TARTARIZATA, Potassæ tartras—**L. Vitriolata, Potassæ sulphas**—**L. Vitriolata sulphurea, Potassæ sulphas cum sulphure.**

LIXIVIA seu KALICUM ACETAS, Potassæ acetas.

LIXIVIAL, *Lixiviosus*, from *lixivium*, 'lee.' (F.) *Lixivial, Lixivieux*. An ancient term for salts obtained by washing vegetable ashes,—such as the fixed alkalis.

LIXIVIA'TION, *Elizivia'tion, Lixivia'tio*, Same etymon. An operation which consists in washing wood-ashes with water, so as to dissolve the soluble parts. The filtered liquor is the *lee*.

LIXIVIUM, *Lixivia, Con'ia, Lee, Ley, Lye*, (F.) *Lensive*. Any solution containing potassa or soda—*Sal lixiviosum*—in excess; from *lix*, 'lye.'

LIXIVIUM AMMONIACALE, Liquor ammoniæ—**L. Ammoniacale aromaticum, Spiritus ammoniæ aromaticus**—**L. Causticum, Liquor potassæ**—**L. Magistrale, Liquor potassæ**—**L. Saponarium, Liquor potassæ**—**L. Tartari, Liquor potassæ subcarbonatis.**

LIXIVUS CINIS, Potash of commerce.

LIZARD, *Lacer'ta, Lacer'tus*, said to be so called in consequence of its limbs resembling the arms (*lacerti*) of man [?]. *Saura, Sauros*, (F.) *Léopard*. Lizards were formerly employed in medicine as sudorifics; and were, at one time, extolled in syphilis, cutaneous affections, and in cancer.

LIZARD'S TAIL, *Saurus cernuus*.

LOADED, see Enduit.

LOADSTONE, Magnet.

LOATHING, Disgust.

LOBARIA ISLANDICA, Lichen Islandicus—**L. Pulmonaria, Lichen pulmonarius**—**L. Saxatilis, Lichen Saxatilis.**

LOBE, *Lobus*. A round, projecting part of an organ. The liver, lungs, and brain, for example, have lobes.

LOBE OF THE EAR, *Lob'ule of the Ear*, is a soft, rounded prominence, which terminates the circumference of the pavilion inferiorly, and which is pierced in those who wear rings.

The under surface of the brain is divided into two anterior, two lateral, two posterior, and two intermediate lobes or processes. These Chaussier calls *lobules of the brain*: the cerebral hemispheres he terms *lobes*.

LOBE, BIVEN'TRAL. A wedge-shaped lobe of the cerebellum, situate behind the amygdala.

LOBE DOUDÉNAL, Lobulus Spigelii—**L. Pancréatique, Lobulus Spigelii**—**L. Petit du foie, Lobulus Spigelii**—**L. de Spigel, Lobulus Spigelii.**

LOBELIA, BLUE, L. syphilitica.

LOBELIA CARDINALIS, *Lobelia coccin'ea, Trachelium Americanum, Cardinal Plant, Cardinal Flower, Scarlet Lobelia.* Ord. Lobeliaceæ. This species is indigenous in the United States. It blooms in autumn, having beautiful carmine flowers. The root is a reputed anthelmintic with the Indians.

LOBELIA COCCINEA, L. Cardinalis.

LOBELIA INFILATA, *Indian or Wild Tobacco, Puke Weed, Asthma Weed, Eyebright, Emet'ic Weed, Lobelia* (Ph. U. S.). The prominent virtues of this American plant are those of an emetic. In smaller doses it is sedative, and has been given as a pectoral in croup, asthma, &c. It is, also, sudorific and cathartic, and is an acro-narcotic poison. Twenty grains act as an emetic.

LOBELIA, PALESPIKED, L. spicata.

LOBELIA PINIFOLIA. A South African plant, the root of which is excitant and diaphoretic. A decoction of it is sometimes used in the Cape Colony as a domestic remedy in cutaneous affections, chronic rheumatism, and gout.

LOBELIA, SCARLET, L. Cardinalis.

LOBELIA SPICATA, *Palespiked Lobelia*; indigenous: is a diuretic.

LOBELIA SYPHILITICA, L. reflex'a, *Ranunculus Virginianus, Ranunculum Syphiliticum, Blue Lobelia, Blue Cardinal Flower.* The root of this plant, which is indigenous in the United States, is an emetic and drastic cathartic. It has been used in syphilis; hence its name. The mode of preparing it is to boil 3ss of the dried root in Oxiij of water, until the fluid is reduced to Oviij. Dose, Oss.

LOBES BIJUMEAUX, Quadrigenita tubercula—**L. Cerebral, see Lobe.**

LOBES OF THE LIVER, *Pinnæ seu Lobi seu Pin-nulæ Hepatis.* See Lobule.

LOBES, OPTIC, Quadrigenita tubercula—**L. Optiques, Quadrigenita tubercula.**

LOBI HEPATIS, Lobes of the liver—**L. Pulmonum, see Pulmo.**

LOBULAR, *Lobularis*. Same etymon as Lobule. Relating to or belonging to a lobule:—as *lobular pneumonia*, (F.) *Pneumonia lobulaire, P. mamelonée, P. disséminée*. Pneumonia anatomically characterized by nuclei of red or gray hepatization disseminated in variable numbers in one or both lungs.

LOBULAR BILIARY PLEXUS. The plexus formed of lobular hepatic ducts, which are derived chiefly from the interlobular. This plexus forms the principal part of the substance of the lobule.

LOBULAR VENOUS PLEXUS. The plexus interposed between the interlobular portal veins, and the intralobular hepatic vein.

LOB'ULE, *Lob'ulus*, diminutive of lobus, 'a lobe.' A little lobe. Mr. Kiernan uses the term *lobule*, for the *acinus* of the liver of many anatomists.

LOBULE OF THE CORPUS STRIATUM, *Insula cerebri*—**L. of the Ear, Lobe of the Ear**—**L. of the Fissure of Sylvius, *Insula cerebri***—**L. du Foie, Lobulus Spigelii**—**L. of the Liver, see Liver**—**L. of the Lung, see Pulmo**—**L. Pneumogastric, Flœculus.**

LOBULI, see Lobulus—**L. Pulmonales, see Pulmo.**

LOBULUS ACCESSORIUS ANTERIOR, QUADRATUS, L. anonymus.

LOBULUS seu LOBUS ANONYMUS, L. accessorius ante'rior quadra'tus, L. quadra'tus, (F.) *Éminence porte antérieure*. This is situate in the liver between the passage for the round ligament and the gall-bladder, and is less prominent, but broader, than the *Lobulus caudatus*. From the lobulus anonymus a bridge runs across the passage for the round ligament. It is called *Pons seu Isthmus hepatis*.

LOBULUS seu **LOBUS CAUDA'TUS**, *Processus caudatus*. This is merely the root or one of the angles of the lobulus Spigelii, advancing towards the middle of the lower side of the great lobe, and representing a kind of tail. Also, the termination of the helix and anthelix of the ear, which is separated from the concha by an extensive fissure.

LOB'ULUS CERTRA'LIS. A small lobule or prominence of the superior vermiform process of the cerebellum, situate in the incisura anterior.

LOBULUS NASI, see **NASUS**—1. Pneumogasticus, Flocculus—1. Posterior, L. Spigelii—1. Posterior papillatus, L. Spigelii—1. Quadratus, L. anonymus—1. Renum, Ferrein, pyramid of.

LOBULUS seu **LOBUS SPIGE'LII**, *L. poste'rior seu poste'rius papilla'tus*, (F.) *Éminence porie' postérieure*, *Lobule* ou *Petit lobe du foie*, *Lobe de Spigel* ou *duodénal* ou *pancréatique*, is situate near the spine, upon the left side of the great lobe of the liver, and is of a pyramidal shape, projecting, like a nipple, between the cardia and vena cava, at the small curvature of the stomach.

LOBUS, *Lobe*, see **Lobulus**—1. Quadrangularis, see **Cerebellum**—1. Semilunaris, see **Cerebellum**—1. Superior anterior, see **Cerebellum**—1. Superior posterior, see **Cerebellum**.

LOCAL, *Local'is*, *Top'icus*, *Mer'icus*, *Partia'lis*, *Top'ical*, (F.) *Locale*, *Topique*. An affection is called local—*Morb'us Local'is*,—when confined to a part, without implicating the general system; or, at all events, only secondarily. *Local* is thus opposed to *general*. A *local* or *topical* application is one used externally. See **Topical**.

LOCH, *Looch*.

LOCHADES, see **Sclerotic**.

LOCH'IA or **LO'CHIA**, *Purgamen'ta Puerperi'rii* seu *Uteri*, *Purga'tio puerperi'rii*, *Lyma*, from *λογος*, 'a woman in childbed,' *λογεια*, 'parturition;' (F.) *Suites de couches*, *Vidanges*. The *cleansings*. A sero-sanguineous discharge following delivery. During the first two or three days, it is bloody; but afterwards becomes green-coloured,—*green waters*—and exhales a disagreeable and peculiar odour. The lochia, in different stages, have received the names *Lochia cruen'ta*, *L. sero'sa* and *L. alba* seu *muc'o'sa* seu *lac'tea*. The duration, quantity, and character of the discharge vary according to numerous circumstances. It flows from the part of the uterus which formed a medium of communication between the mother and foetus, and continues, usually, from 14 to 21 days. See **Parturition**.

LOCHIOCELIITIS, *Puerperal fever*.

LOCHIODOCHIUM, *Lochodochium*.

LOCHIOPIRYA, *Puerperal fever*.

LOCHIORRHAG'IA, from *lochia*, and *ρρννρη*. 'I make an irruption.' An immoderate flow of the lochia. Hemorrhage from the uterus in the child-bed state.

LOCHIORRH'E'A, from *lochia*, and *ρρω*, 'I flow.' Discharge or immoderate discharge of the lochia.

LOCHIORUM RETENTIO, *Ischlochchia*.

LOCHIOSCHESIS, *Ischlochchia*.

LOCHIOCH, *Looch*.

LOCHODOCHI'UM, *Lochiodochi'um*, from *λογος*, 'a female in childbed,' and *δοχωμ*, 'I receive.' An institution for the reception of pregnant and childbed females. A *Lying-in-hospital*.

LOCHOMETRITIS, see **Metritis**.

LOCHOMETROPHEBITIS, see **Phlebitis**.

LOCHOOPHORITIS, see **Oaritis**.

LOCHOPERITONITIS, see **Peritonitis**.

LOCHOS, *Puerpera*.

LOCI, *Uterus*—1. Muliebres, *Uterus*, *Vulva*.

LOCK HOSPITAL. A term of uncertain origin, generally given to venereal hospitals in Great Britain.

LOCKED JAW, *Trismus*.

LOCKING, *Coxofemoral articulation*.

LOCOMOTILITY, see **Locomotion**.

LOCOMO'TION, *Locomoti'o*, from *locus*, 'a place,' and *mo'vere*, *motum*, 'to move.' An action peculiar to animal bodies, by which they transport themselves from place to place. It, as well as *muscula'tion*, has also been used for the function of animal movements. The faculty is sometimes called *Locomotiv'ity* and *Locomotiv'ity*.

LOCOMOTION OF AN ARTERY, is the movement produced in the vessel with a curvature, by the impulse of the blood sent from the heart, which tends to straighten the artery, and causes the movement in question.

LOCOMOTIVITY, see **Locomotion**.

LOCULAMENTA COLLI, see **Colon**.

LOCUS CINEREUS, *Substantia ferruginea*.

LOCUS NIGER, see *Peduncles of the Brain*.

LOCUS PERFORA'TUS ANTI'CUS seu **ANTE'RIOR**. A triangular flat surface of the brain, which corresponds to the posterior extremity of each of factory process in front of the optic tract, and near the entrance of the fissure of Sylvius. *Substantia seu Lam'ina perforata ante'rior*.

LOCUS PERFORA'TUS POSTICUS, see **Tarini pons**.

LOCUST, **BLACK**, *Robinia Pseudo-acacia*. **LOCUST EATERS**, *Acridophagi*—1. *Plant*, *Cassia Marilandica*—1. *Tree*, *Robinia Pseudo-acacia*.

LOCUST, **YELLOW**, *Cladrastis tinctoria*.

LODGED, *Wedged*.

LODOICEA, see *Coco of the Maldives*.

LOECHE, **MINERAL WATERS OF**, *Leuck*, *mineral waters of*.

LOEME, *Plague*.

LOEMIA, *Plague*.

LOEMICUM, see **Læmology**.

LOEMOCHOLYSIS, *Fever, yellow*.

LOEMOCOMI'UM, from *λοιμος*, 'plague,' and *κομειν*, 'I take care of.' An hospital for those affected with plague.

LOEMOGRAPHY, *Loimography*.

LOEMOLOGIUM, see **Læmology**.

LOEMOL'OGY, *Læmolog'ia*, from *λοιμος*, 'plague,' and *λογος*, 'a description.' The doctrine of plague and pestilential diseases. A treatise on the same,—*Lo'micium*, *Læmolog'ium*.

LOEMOPHTHALMIA, see **Ophthalmia**.

LOEMPYRA, *Plague*.

LOEMOS, *Plague*.

LOGADES, *Sclerotic*.

LOGADITIS, *Sclerotitis*.

LOGIA'TROS, *Logia'ter*; from *λογος*, 'a word,' and *ιατρος*, 'a physician.' In the bad sense, a physician without experience; a mere theorist. In the good sense, a rational physician; one who treats disease according to theoretical or scientific principles.

LOGONEURO'SES, from *λογος*, 'mind, reason,' and *νευρον*, 'a nerve.' Diseases of the mental activity.—*Romberg*.

LOGOS, *Reason*.

LOGWOOD, *Hæmatoxylon Campechianum*.

LOG'Y, *λογος*, 'a description.' A suffix denoting 'a treatise or description.' Hence, *Angiology* and *Neurology*, &c.

LOHOCH, *Looch*.

LOIMOCHOLYSIS, *Fever, yellow*.

LOIMOG'RAPHY, *Loimograph'ia*, *Læmog'raphy*, from *λοιμος*, 'plague,' and *γραφω*, 'I describe.' A description of the plague and pestilential diseases.

LOIMOLOGY, *Læmology*.

LOIMOS, *Plague*.

LOINS, *Lumbi*.

LOISELEUR'IA PROCUMBENS, *Azalea procumbens*; indigenous; *Order*, *Ericaceæ*; has some reputation as an astringent.

LOLIACEUM RADICE REPENTE, *Triticum repens*.

LOLIUM ANNULUM, *L. temulentum*.

LOLIUM TEMULENTUM, *L. annuum*, *Crepedia temulentata*, *Bromus temulentus*, *Darnel*, (Prov.) *Cockgrass*, *Jum*, (F.) *Herbe d'Ivrogne*. A species of the genus *Lolium*, (F.) *Ivraie*; *Ord.* Gramineæ; *Gen.* *Syst.* Triandria digynia, which has decidedly poisonous properties; occasioning, when mixed in bread or beer, intoxication, vertigo, nausea, and vomiting.

LOMBAIRE, *Lumbar*.

LOMBES, *Lumbi*.

LOMBO-ABDOMINAL, *Transversalis abdominalis*—*l. Costal*, *Serratus posticus inferior*—*l. Costo-trachélien*, *Sacro-lumbalis*—*l. Dorso-spinal*, *Transversalis dorsi*—*l. Dorso-trachélien*, *Longissimus dorsi*—*l. Huméral*, *Latissimus dorsi*—*l. Sacré*, *Lumbo-sacral*.

LOMBRIC, *Ascaris lumbricoides*.

LOMBRICAUX, *Lumbricales*.

LOMBRICOÏDE, *Ascaris lumbricoides*.

LONCHADES, see *Sclerotic*.

LONCHADITIS, *Sclerotitis*.

LONCHITIS, *Polypodium filix mas*.

LONG, *Longus*, *Macros*. That which is much greater in length than in breadth; as the *long bones*. The epithet is, also, applied to several muscles, to distinguish them from others of similar function, when the latter are shorter. We say, for instance, *long flexors*, and *long extensors*, in opposition to *short flexors*, and *short extensors*.

LONG DU COU, *Longus colli*—*l. du Dos*, *Longissimus dorsi*.

LONGÆVUS, *Macrobiotic*.

LONGANON, *Rectum*.

LONGAON, *Rectum*.

LONGAS, *Rectum*.

LONGEVITY, *Longævitas*, from *longus*, 'long,' and *ævum*, 'age,' *Macrobiosis*, *Macrobiosis*. The prolongation of existence to an advanced age. Haller collected examples of more than one thousand centenarians. He had knowledge of sixty-two persons aged from 110 to 120 years; of twenty-nine, from 120 to 130 years; and of fifteen, who had attained from 130 to 140 years. Beyond this advanced age, examples of longevity are much more rare and less sufficiently attested.

The following list of instances of very advanced ages has been given:

	<i>Lived.</i>	<i>Age.</i>
Apollonius of Tyana.....	A. D. 99...	130
St. Patrick.....	491...	122
Attila.....	500...	124
Llywarch Hên.....	500...	150
St. Coemgene.....	613...	120
Piastus, King of Poland.....	861...	120
Thomas Parr.....	1635...	152
Henry Jenkins.....	1670...	169
Countess of Desmond.....	1612...	145
Thomas Damme.....	1643...	154
Peter Torten.....	1724...	185
Margaret Patten.....	1739...	137
John Rovin and wife.....	1741...	172 and 164
St. Monagh or Kentigen.....	1781...	185

Longevity also means *length or duration of life*, (F.) *Durée de la vie*. The mean age at death, (F.) *Vie moyenne*, of different classes and professions enables an estimate to be formed of the *expectation or value of life* in each.

LONGING, *Malacia*.

LONGISSIMUS, superlative of *longus*, 'long.' Longest. A term applied to parts, which are very long in their relation to others.

LONGISIMUS DORSI, *Semi-spina'tus*, (F.) *Lombo-dorso-trachélien*, *Portion costo-trachélienne du sacro-spinal* (Ch.) *Long dorsal*, *Long du dos*,

is situate vertically at the posterior part of the trunk, and fills, in a great measure, the vertebral furrows. It is thick and almost square below; thin and pointed above. It is attached to the posterior surface of the sacrum, to the transverse processes of all the lumbar and dorsal vertebrae, and to the inferior margin of the last 7 or 8 ribs. It maintains the vertebral column in a straight position; straightens it when bent forwards, and can even carry it back. It also assists in the rotatory motion of the trunk.

LONGISSIMUS FEMORIS, *Sartorius*—*l. Oculi*, *Obliquus superior oculi*.

LONG-SIGHTEDNESS, *Presbytia*.

LONGUS, *Long*.

LONGUS COLLI, (F.) *Pré-dorso-cervical*, *Pré-dorso-atloïdien*, (Ch.) *Long du cou*. This muscle is situate at the anterior and superior part of the vertebral column. It is long, flat, and broader at its middle than at its extremities, which are pointed. It is attached to the anterior surface of the bodies of the first three dorsal and last six cervical vertebrae; to the intervertebral ligaments; to the anterior edge of the transverse processes of the last five cervical vertebrae; and to the tubercle on the anterior arch of the first. This muscle bends the cervical vertebrae upon each other and upon the dorsal vertebrae. If the upper portion acts on one side only, it occasions the rotation of the atlas on the vertebra dentata; and, consequently, of the head on the neck.

LONGICERA GERMANICA, *L. Periclymenum*—*l. Marilandica*, *Spigelia Marilandica*.

LONGICERA PERICLYMENUM, *L. Germanica*, *Periclymenum*, *P. vulgare*, *Caprifolium*, *C. Periclymenum* seu *sylvaticum* seu *distinctum*, *Common Woodbine*, *Honeysuckle*, (Prov.) *Suckling*, (F.) *Chèvre-feuille*. This common plant is slightly astringent and tonic, and was formerly much used in gargles.

LONGICERA SEMPERVIRENS, *Caprifolium sempervirens*, *Scarlet* or *Trumpet Honeysuckle*. *Ord.* *Caprifoliaceæ*; indigenous. A syrup of the leaves is said to have been given in asthma and in amygdalitis. [?]

LONK, *Coxofemoral articulation*.

LOOCH, *Lochoch*, *Logh*, *Lochoch*, *Looch*, of Arabic extraction. A linctus. See *Eclegma* and *Eclectos*.

LOOCH ABSQUE EMULSIONE PARATUM, *Look préparé sans émulsion*; *Looch prepared without emulsion*. (*Pulv. g. trag. gr. xvj*—*gr. xxx*, *ol. amygd. dulc. ℥ss*, *sacchar. ℥ij*, *aquæ ℥iij*, *aquæ flor. aurant. ℥ij*. Mix by rubbing in a marble mortar.) *Demulcent*.

LOOCH ALBUM, *L. amygdalinum*, *Linctus albus* seu *amygdalinus* seu *communis*, *Eclegma album* seu *gummo-so-oleosum*, (F.) *Looch blanc*, *L. b. amygdalin*, *L. b. pectoral*. (*Amygd. dulc. ℥ss*, *amygd. amar. No. ij*, *sacchar. alb. ℥iv*. Make an emulsion by gradually adding *℥iv* of water. Then take *pulv. tragacanth. gr. xvj. ol. amygd. dulc. recent. ℥ss*, *sacch. ℥ij*. Add the almond milk gradually to this, and afterwards *aq. flor. aurant. ℥ij. Ph. P.*) It is demulcent and pectoral.

LOOCH AMYGDALINUM, *L. album*—*l. Blanc*, *L. album*—*l. cum Croco* et *pistaciis*, *L. viride*—*l. of Egg*, *L. ex Ovo*.

LOOCH EX OVO, *L. pectoralē leniens*, *Potio seu emul'sio seu mist'ra leniens seu oleo'sa*, *Looch of Egg*, (F.) *Look d'œuf*, *Looch rouge*, *Emulsion huileuse*, *Mixture calmante*, *Potion pectorale*, *Lait adoucissant*. (*Vitell. ovi. recent. ℥ss*, *ol. amygd. dulc. ℥iss*, *syrup. althææ. ℥j*. Rub in a mortar, and add by degrees, *aq. flor. aurant. ℥j*, *aq. papav. rhæad. ℥ij. Ph. P.*) Virtues the same as the preceding.

LOOCH ROUGE, *L. ex Ovo*—*l. Vert*, *L. viride*.

LOOCH VIR'IDE, *L. cum crocæ et pista'ciis*, (F.) *Looch vert.* (*Synop. Violar.* 3j, *tinct. croci* gtt. xx., *aquæ* 3iv. Mix, and add *pistaciæ semin. sicc.* 3vj. *Ph. P.*) Virtues like the last.

LOOK, *Looch*—*l. d'Æuf*, *Looch ex ovo*—*l. Préparé sans émulsion*, *Looch absque emulsione paratum*.

LOOSE STRIFE, CREEPING, *Lysimachia nummularia*—*l. s.* Four-leaved, *Lysimachia quadrifolia*.

LOOSENESS, *Diarrhœa*—*l. of the Teeth*, *Odontoseisis*.

LOPEZ RADIX, *Radix lopesia'na seu Indica lopezia'na*. The root of an unknown Indian tree, not possessed of any remarkable smell or taste, or of any appearance of resinous matter. It has been extolled, notwithstanding, in cases of colliquative diarrhœa. Gaubius compares its action to that of simarouba, but thinks it more efficacious.

LOPHADIA, *Lophia*.

LOPHIA, *Lopha'dia*, *λοφία*. The upper part of the neck. The first vertebra of the back—*Vertebra dorsi prima*—Gorraus.

LOPIMA, *Fagus castanea*.

LOQUAC'ITY, *Garrulitas*; from *loquor*, 'I speak. (F.) *Babillement*. The act of speaking with volubility. It is sometimes a symptom of disease, and is observable in hysteria, &c.

LOQUELA, Voice, articulated—*l. Abolita*, *Aphonia*—*l. Blassa*, *Balbuties*—*l. Impedita*, *Barryphonia*.

LORDO'SIS, *Lordo'ma*, from *lordôs*, 'curved,' 'bent.' *Repanda'tio*, *Repan'ditas*. A name given to curvatures of the bones in general; and particularly to that of the vertebral column forwards; *Spina dorsi intro'sum flexa*, *Emprostocyrto'ma*. This gives rise to the projection of the sternum called chicken-breasted or pigeon-breasted. *Pectus carina'tum*, *Pig'leon breast*, narrow chest.

LORIND MATRICIS, *Epilepsia uteri'na*, *Convulsivus ute'ri morbus*. A barbarous name given to a pretended epilepsy of the womb.

LORIPES, see *Kyllosis*.

LORUS, *Hydrargyrum*.

LOT, *Urine*.

LOTA MOLVA, see *Oleum jecoris aselli*—*l.* *Vulgaris*, see *Oleum jecoris aselli*.

LOTIO, *Enema*, *Lotion*—*l. Cupri sulphatis camphorata*, *Lotion*, camphorated, of Bates—*l. Rubra*, *Lotion*, camphorated, of Bates.

LOTIO RUBRA, *Red wash*, (*Hydrarg. chlorid. corrosiv.* gr. xviii; *H. sulphureti rubr.* gr. ix; *Creasot.* ℥ vi; *Aquæ* f3viij. May be used in the proportion of f3j; to from f3i to f3iiij of water; like the yellow wash.

LOTIO SAPONACEA, see *Sapo*.

LOT'ION, *Lo'tio*, *Lotu'ra*, from *lavare*, *lotum*, 'to wash.' (F.) *Hydroloti*, *Hydro'le*. A fluid external application. Lotions are ordinarily applied by wetting linen in them and keeping it on the part affected.

LOTION, *BARLOW'S*, *Lotion of Sulph'uret of potass'um*. (*Potassii sulphur.* 3ij; *Sapon.* 3iss; *Aq. Calcis*, f3viiss; *alcohol. dilut.* f3ij. M.) Used in various chronic cutaneous diseases.

LOTION, *CAMP'ORATED*, OF *BATES*, *Bates's camph'orated water*, *Lotio cupri sulphat'is camphorata*, *Lotio rubra*, *Red wash*. (*Cupri sulphat.* 3ij; *Bol. rubr.* 3ij; *Camphor.* 3ss; *Aq. bullient.* Oij. Strain through linen. f3ij of this, added to f3iv of water, forms *Bates's Collyrium*, *Collyrium Batea'num*, which may be made extemporaneously as follows:—*Cupri sulphat.* gr. viij; *Boli*, gr. viij; *Camphor.* gr. ij; *Aq. bullient.* f3viij. Mix and filter.

LOTION, *FEBURE'S*. A cancer remedy, which consisted of *arsenic.* gr. x; *water*, Oj; *extract of*

hemlock, 3j; *Liquor Plumbi subacetatis*, f3iiij; *laudanum*, f3j.

LOTION, *GROWLAND'S*. An empirical preparation. (*Bitter almonds*, 3j; *sugar*, 3ij; *distilled water*, lbj. Grind together, strain, and add *corrosive sublimate*, ʒij, previously ground with *sp. vini rect.* 3ij.) Used in obstinate eruptions.

LOTION, *GRANVILLE'S COUNTER-IRRITANT*, *Granville's Lotion*, *Granville's antid'ynous lotion*. Of this lotion, Dr. Granville gives two forms—a milder, and a stronger. The milder is made as follows: *Liq. ammon. fort.* f3j; *Sp. Rosmarin.* f3vj; *Tinct. camphor.* f3ij, M.

The stronger is made as follows: *Liq. ammon. fort.* f3x; *Spiri. Rosmar.* f3ss; *Tinct. camphor.* f3ij, M.

The stronger lotion vesicates rapidly. A piece of cotton or linen folded six or seven times, or a piece of thick flannel may be imbued with them, and laid for a few minutes on the part to be irritated.

The Dublin Pharmacopœia has a *Linimen'tum Ammoniac Compo'situm*, *Compound Liniment of Ammonia*, which is a close imitation of *Granville's Lotion*.

LOTION, *HANNAY'S*, *Preven'tive wash*. This famous nostrum for the prevention of venereal infection, was nothing more than a solution of caustic potass.

LOTION, *HYDROCYAN'IC*, *Lotio Ac'idi Hydrocyan'ici*. (*Hydrocyanic acid*, f3iv; *rectified spirit of wine*, f3j; *distilled water*, f3xss.) Used with much success in impetigo, &c.

LOTION, *STRUVE'S*, FOR *HOOPING-COUGH*. (*Antim.* et *Potass.* tart. 3j; *Aquæ*, 3ij. Add *tinct. cantharid.* 3j.)

LOTIUM, *Urine*.

LOTURA, *Lotion*.

LOTUS SYLVESTRIS, *Trifolium melilotus*—

l. Virginiana, *Diospyros Virginiana*.

LOUCHEMENT, *Strabismus*.

LOUCHES, see *Strabismus*.

LOUPE, *Wen*.

LOUSE, *Pediculus*.

LOUSINESS, *Phtheiriasis*.

LOUSY DISEASE, *Phtheiriasis*.

LOUTRON, *Bath*.

LOVAGE, *Ligusticum levisticum*.

LOVE, *Eros*, *Amor*, from *Sax. lufian*, (G.) *lieben*, 'to love.' (F.) *Amour*. A tender and elevated feeling, which attracts one sex to the other. Love is occasionally a cause of disease, especially of insanity.

LOVE APPLE PLANT, *Solanum lycopersicum*—*l. Begot*, *Illegitimate*—*l. Child*, *Illegitimate*—*l. Pea*, *Abrus precatorius*.

LOW SPIRITS, *Hypochondriasis*.

LOWER, TUBERCLE OF, *Tuberculum Loweri*. Anatomists have given this name to a small projection, the existence of which is by no means constant, and which is found in the sinus venosus, between the superior and inferior cava.

LOXARTHROS, *Loxarthrum*, from *loxos*, 'oblique,' and *arthron*, 'articulation.' (F.) *Perversion de la tête des os et des muscles*. A vicious deviation or direction of the joints, without spasm or luxation,—as in *clubfoot*.

LOXIAS, *Torticollis*.

LOXOPHTHALMUS, *Strabismus*.

LOZANGIA, *Lozenge*.

LOZENGE, *Tabella*.

LOZENGES, *BARK*, *Tabellæ cinchonæ*—*l.* of *Catechu* and *magnesia*, *Tabellæ antimoniales Kunckelii*—*l. Faustinus's*, *Faustini pastilli*—*l.* for the *Heart-burn*, *Trochisci carbonatis calcis*—*l. Magnesia*, *Tabellæ de magnesiâ*—*l.* of *Marsh-mallows*, *Tabellæ de althæâ*—*l.* of *Oxalic acid*, *Tabellæ acidî oxalicî*—*l. Pectoral*, *black*, *Trochisci glycyrrhizæ glabræ*—*l. Pectoral*, of *eme-*

tine, Trochisci emetinæ pectorales—1. Rhubarb, Tabellæ de rheo—1. of Scammony and senna, compound, Tabellæ de scammonio et sennâ—1. Spittæ's, see Trochisci glycyrrhizæ cum opio—1. Steel, Tabellæ de ferro—1. Steel, aromatic, see Tabellæ de ferro—1. Sulphur, simple, Tabellæ de sulphure simplicibus—1. Sulphur, compound, Tabellæ de sulphure compositæ—1. of Sulphuret of antimony, Tabellæ antimoniales Kunkelii—1. Wistar's, Trochisci glycyrrhizæ cum opio—1. Worm, Ching's, see Worm lozenges, (Ching's.)

LUBIDO, Libido—1. Intestini, Voluntas desedendi.

LUBRICANTIA, Demulcents, Emollients.

LUBRICUM CAPUT, Penis.

LUCCA, MINERAL WATERS AND CLIMATE OF. The baths and waters, near this ancient Italian city, have been long celebrated. They are thermal, and resemble, in properties, those of Plombières in France, and of Bath England. They contain carbonic acid, sulphates of alumina, soda, magnesia, and iron. There are ten different sources, the temperature of which varies from 94° to 130° Fahr.

Luca is much frequented in summer; partly on account of its mineral waters, but more on account of the coolness of the situation.

LUCID, *Lu'cidus*, from *luceo*, 'I shine.' In medicine, the word *lucid* is particularly applied to the intervals, *Interval'la lu'cida*, of apparent reason, which occur in mental alienation.

LUCIFER-MATCH-MAKERS' DISEASE, see Phosphorus.

LUCIFUGUS, from *lux*, *lucis*, 'light,' and *fugere*, 'to shun.' *Photoph'obus*, *Photoph'ob'icus*. Dreading or avoiding the light.

LUCINA, Lithyia.

LUCOMANIA, Lycanthropia.

LUCTUEUX (F.), *Luctu'ous*, 'mournful.' *Plaintive*. An epithet applied to the respiration—(F.) *Respiration luctueuse*—when accompanied by moaning.

LUCUMA, (S.) A fruit which grows in the southern provinces of the coast of Peru, and the north of Chili. It is round; and the gray-brown husk encloses a fibrous, dry, yellow-coloured fruit with its kernel.—Tschudi.

LUCUMORIA'NUS, probably from *lux*, 'light,' and *morari*, 'to tarry.' Continuing for several days: hence, *Dormit'io lucumoria'na*. A morbid sleep persisting for several days.

LUDUS HELMONTII, *L. Paracel'si, Fel terræ*. A calcareous stone, the precise nature not known, which was used by the ancients in calculous affections. The term was also applied to every species of calculous concretion occurring in the animal body.

LUDUS PARACELSI, L. Helmontii.

LUDWIG'IA PALUSTRIS, *L. nit'ida, Ienar'dia palus'tris, Water Purlaine, Phthiis Weed*. Order, Onagraceæ; indigenous. An infusion of the dried plant has been used as a pectoral in asthma and chronic cough.

LUES, Disease, Plague, Syphilis—1. Divina, Epilepsy—1. Dysenterica, Dysentery—1. Gonorrhoeica, Gonorrhoea impura—1. Gutturis epidemica, Cynanche maligna—1. India, Framboesia—1. Inguinaria, Plague—1. Pannonie, Fever, Hungary—1. Polonica, Plica—1. Sarmatica, Plica—1. Syphilis, Syphilis—1. Syphilodes, Syphilis pseudo-syphilis—1. Trichomatica, Plica—1. Venerea, Syphilis—1. Scorbatica, see Purpura.

LUETTE, Uvula—1. *Vésicale*, see Urinary bladder.

LUG, Ear.

LUGDUS, Erysipelas.

LUJULA, Oxalis acetosella.

LULLABY SPEECH, Lallation.

LUMBA'GO, from *lumbi*, 'the loins.' *Arthro'*

sia lumb'o'rum, Lumbago seu Nephral'gia rheumat'ica, Rachir'rheu'ma, Rachiorrheu'ma, Rheumat'ismus dorsi'lis, Osphyrrheu'ma, Rheumatism affecting the lumbar region.

Lumba'go psoad'lica, L. apostemato'ea, L. ab arthro'acæ. Pain in the loins from abscess. See Psotitis.

LUMBAGO A NISU, see Effort.

LUMBAR, *Lumba'ris seu Lumba'lis*, (F.) *Lombaire*. From *lumbi*, 'the loins.' Belonging or having reference to the loins.

LUMBAR ABSCESS, *Psoas abscess, Aposte'ma psoad'icum, Absces'sus lumb'o'rum, Morbus psoad'icus*. This abscess is so called from the matter being found on the side of the psoas muscle, or betwixt that and the iliacus internus. Between these muscles is a quantity of loose, areolar substance; and when an abscess takes place there, it can find no outlet except by a distant course. Generally, it descends along the psoas muscle, forming a swelling immediately beneath Poupart's ligament; at times, however, it extends down the thigh under the fascia. Severe hectic follows the bursting of the abscess, and often death. Its causes are, — scrofula, injury to the loins, &c.

When, from the pain of the back continuing for some time, with other symptoms, the disease is suspected, caustics, or the moxa, applied opposite the transverse processes of the lumbar vertebrae, may be recommended; and, when the abscess is ready for opening, it may be emptied by repeated discharges of the matter, through a small opening, made with a lancet, or small, lancet-pointed trocar. The medical treatment consists in supporting the system, under the great restorative efforts required of it.

LUMBAR ARTERIES are four or five in number on each side. They arise from the sides of the abdominal aorta, and pass behind the muscles situate in front of the lumbar portion of the spine, to gain the broad muscles of the abdomen. They give off, 1. Spinal branches. 2. Anterior, posterior, and external muscular branches.

LUMBAR NERVES are five in number, and issue from the vertebral column by the spinal foramina of the loins. The first lumbar nerve gives off three branches:—the external or *ilio-sciatic*; the middle or *inguino-cutaneous*, and the internal or *infra-pubic*. Along with the three pairs below it, it forms the lumbar plexus.

LUMBAR PLEXUS, *Portion lombaire du plexus crural* (Ch.), *Plexus Lombo-abdominal*. This plexus is formed by the union of the *Rami communican'tes* of the anterior branches of the first four lumbar nerves. It is situate behind the psoas muscle, and before the transverse processes of the lumbar vertebrae. It furnishes, besides the branches which proceed from the first pair, several filaments, that are distributed to the psoas muscle, to the iliacus, the integuments, and glands of the groin; and three great branches—the *crural*, *obturator*, and *lumbo-sacral*.

LUMBAR REGION, Lumbi.

LUMBAR VEINS have an arrangement analogous to that of the arteries of the same name. They communicate with the vertebral sinuses, azygous veins, &c., and pass into the vena cava inferior.

LUMBARIS EXTERNUS, *Quadratus lumborum*—1. Internus, *Psoas magnus*.

LUMBI, *The loin or loins, Lumbar region*, (Old Eng.) *Lende*, (Sc.) *Lendis*; *Re'gio lumba'lis, Lap'ara, Psca*, plural *Pscae, Osphyæ, Reins*, (F.) *Lombes*. The posterior regions of the abdomen, composed between the base of the chest and the pelvis. The parts which enter into the formation of the lumbar region are,—the skin; a considerable quantity of areolar texture; broad and strong aponeuroses:—the *Latissimus Dorsi*,

Obliquus externus, and *Obliquus internus abdominis*, *Transversalis abdominis*, *Quadratus lumborum*, and the mass common to the *Sacro-lumbalis*, *Longissimus dorsi*, and *Multifidus spinæ*. These muscles surround the lumbar region of the vertebral column. The vessels, nerves, &c., of the loins, are called *lumbar*.

LUMBO-SACRAL. Belonging to the lumbar and sacral regions. (F.) *Lombo-sacré*. Bichat calls this a very large nerve, given off from the anterior branch of the fifth lumbar pair, which descends into the pelvis before the sacrum to join the sciatic plexus.

LUMBRICALIS, *Vermicularis*. Resembling a *lumbricus*, or 'earthworm.' A name given to small muscles, (F.) *Lombricaux*, met with in the palm of the hand and sole of the foot.

LUMBRICALES MANUS, *Fidicinales*, *Flexor primi internodii digitorum manus seu perforatus lumbricalis*, (F.) *Annuli-tendino-phalangiens*, *Palmi-phalangiens*. Four small, fleshy, thin, round, long, fusiform fasciculi, situate in the palm of the hand, and distinguished into first, second, third, and fourth, counting from without to within. They arise from the tendons of the flexor communis digitorum, and are inserted at the outer and posterior side of the superior extremity of the first phalanges of the last four fingers. These muscles bend the fingers on the metacarpus, and fix the tendons of the flexor digitorum communis.

LUMBRICALES PEDIS, (F.) *Planti-tendino-phalangiens*, *Planti-sous-phalangiens*, (Ch.) They are analogous to those of the hand in form, number, and arrangement. They increase the flexion of the toes, and draw them inwards.

LUMBRICUS, *Ascaris lumbricoides*—1. *Latus*, *Bothriocephalus latus*, *Tænia solium*—1. *Teres hominis*, *Ascaris lumbricoides*.

LUMBUS VENERIS, *Achillea millefolium*.

LUMEN, Light, Pupil—1. *Constans*, *Phosphorus*.

LUMIÈRE, Light.

LUMINOSITY, Phosphorescence.

LUNA, Argentum, Moon—1. *Albini*, *Sciatic notch*, lesser—1. *Imperfecta*, *Bismuth*—1. *Potabilis*, *Argenti nitras*.

LUNACY, see *Lunatic*.

LUNAR, *Lunatic*.

LUNA'RE OS, *Os semilunare*. The second bone in the upper row of the carpus.

LUNARIA, *Menses*.

LUNA'RIA REDIVIVA, *Leucoïum*, *Bulbonach*, *Satin*, *Honesty*. Said, by Ray, to be a warm diuretic.

LUNARIS, *Lunatic*.

LUNATIC, *Lunaticus*, *Luna'ris*, *Luna'rius*, *Lunar*, from *luna*, 'the moon.' Relating to the moon. An epithet given to diseases which are supposed to appear at certain phases of the moon, or to those who are affected by them. The term *lunatic* (*Moon'ed*, *Moonstruck*, *Selenoble'tus*), is restricted to one labouring under lunacy or mental alienation, *Dementia accidentalis seu adventitia*, that is, under any form of unsoundness of mind, except idiocy. In law, a *lunatic* is one who has had an understanding, but by disease, grief, or other accident, has lost the use of his reason, and become *non compos mentis* or *non compos*.

LUNATICUS, *Epileptic*, *Lunatic*, *Somnambulist*.

LUNATISMUS, *Somnambulism*.

LUNE, Moon—1. *d'Eau*, *Nymphæa alba*.

LUNELLA, *Hypopyon*.

LUNES, *Menses*.

LUNETTES ORDINAIRES, *Spectacles*.

LUNG, *Pulmo*—1. *Black*, of coal miners, *Anthracosis*—1. *Cancer* of the, *Phthisis*, *cancerous*.

LUNG FEVER, *Catarrh*, *Pneumonia*.

LUNG, *PERFORATING ABSCESS* OF THE. A purulent collection, which forms exterior to the lung, and afterwards perforates its tissue, so that it is evacuated through the bronchial tubes.

LUNG PROOF, *Docimasia pulmonum*—1. *Wort*, *Pulmonaria*—1. *Wort*, cow's, *Verbascum nigrum*—1. *Wort*, tree, *Lichen pulmonarius*.

LUNULA, *Onyx*—1. *Unguicium*, see *Nail*—1. *Scapula*, *Notch*, *scapular*.

LUNYIE, *Lumbi*.

LUNYIE-BANE, *Ischion*.

LUPIA, *Encystis*, *Wen*—1. *Junctura*, *Spina ventosa*.

LUPINUS. Under this term the *white lupin*, *Lupinus albus seu sativus*, *Ord.* *Leguminosæ*, is meant, in some pharmacopœias. The seeds, which were much eaten in the days of Pliny and of Galen, are now neglected. The meal is, however, occasionally used as an anthelmintic, and as a cataplasm.

LUPOID, *Lupoïdes*, badly compounded from *lupus*, and *eidos*, 'resemblance.' Resembling, or having the character of, *lupus*.

LUPULINA, *Lupulin*, *Lupulinic glands*. A substance which exists in *Humulus lupulus* or hop. It is in the form of small, shining, yellowish grains, which cover the base of the scales of the hop; is pulverulent, and of an aromatic odour. When analyzed, it is found to contain resin, volatile oil in small quantity, and a bitter principle—*Lupulite*. It is aromatic and tonic, and—according to some—narcotic.

LUPULINIC GLANDS, *Lupulina*.

LUPULITE, see *Lupulina*.

LUPULUS, *Humulus lupulus*—1. *Communis*, *Humulus lupulus*—1. *Salictarius*, *Humulus lupulus*—1. *Scandens*, *Humulus lupulus*.

LUPUS, 'the wolf.' So named from its rapacity. *Lupus exedens seu vorax seu devorans*, *Jacob's Ulcer*, *Ulcus Tuberculo'sum*, *Cancer seu Phymato'sis lupus*, *Noli me tang'ere*. Tubercular excrescences, with ragged, spreading ulcerations, chiefly about the *alæ nasi*, where they destroy the skin, &c., for some depth. Sometimes they appear in the cheek, circularly, or in the shape of a sort of ringworm, destroying the substance, and leaving a deep and deformed cicatrix. The knife or caustic should be used to separate the sound from the morbid parts. Arsenic has been given internally with advantage. See, also, *Herpes exedens*, and *Lycoides*.

LUPUS CANCRO'SUS, *Cancer*—1. *Devorans*, see *Lupus*—1. *Exedens*, *Lupus*—1. *Varicosus*, *Nævi*—1. *Vorax*, *Herpes exedens*, see *Lupus*.

LURCO, *Glutton*.

LURID, *Luridus*. Pale, yellow, sallow,—applied to the complexion.

LURKEY-DISH, *Mentha pulegium*.

LUROR, *Paleness*.

LUS, *Sacrum*.

LUSCIOSTAS, *Luscitās*, *Myopia*.

LUSCIOSUS, *Borgne*, *Myops*.

LUS'CITAS, *Luscio'sitas*, *Luscit'ies*, *Luscit'ion*, from *lucius*, 'a blinkard,' *lusciosus*, 'dim-sighted.' *Strabismus*. The term has also been given to any case of obliquity in which the eye is fixed in an unnatural position.—Beer. The obliquity termed *Cock-eye* is a form of *Luscitās*.

LUSCITIES, *Luscitās*.

LUSCUS, *Borgne*.

LUSS, *Pityriasis*.

LUSTRAMENTUM, *Cathartic*.

LUT, *Lute*.

LUTE, *Lutum* ('mud'), *Cæmentum*. (F.) *Lut*. A composition employed either for closing apertures in a pharmaceutical instrument, or for covering the surface of retorts, tubes, &c., which have to support a great degree of heat. *Lutes*

are composed differently, according to the object to be accomplished. Commonly, they are made of linseed meal and starch. The *fat lute* is formed of clay and drying oil. Sometimes, the white of egg and lime are used; and that which is employed for covering vessels, intended to be strongly heated, is made of clay, sifted sand, and water.

LUTEOLA, *Reseda luteola*.

LUTEUS, Yellow.

LUTRON, λουτρον. A bath. Also, an ophthalmic medicine.—Galen.

LUX, Light.

LUXATIO, Luxation—I. Imperfecta, Sprain.

LUXATION, *Eluxation*, from *luxare*, *luxatum*, 'to put out of place.' *Dislocation*, *Olisthema*, *Apopedasia*, *Luxatio*, *Dislocatio*, *Ecclisis*, *Stremma*, *Luxaturia*, *Etetopia*, *Eptosis*, *Elapsus*, *Lygis mus*, *Emotio*, *Ecstasis*, *Exarthrema*, *E. luxatio*, *Exarthroma*, *Exarthrosis*, *Excoxis*, *Exotheis*, *Paratopia*, *Streblosis*, *Delocatio*, *Exarticulatio*, (F.) *Luxation*, *Dislocation*, *Déboitement*. A displacement of a part from its proper situation. A putting out of joint. A displacement of two or more bones, whose articular surfaces have lost, wholly, or in part, their natural connexion; either owing to external violence, (*accidental luxation*), or to disease of some of the parts about the joint (*spontaneous luxation*.) *Luxation* is complete when the bones have entirely lost their natural connexion; *incomplete*, when they partly preserve it; and *compound*, when a wound communicates with the luxated joint. The general indications of treatment, are;—1. To reduce the protruded bone to its original place. 2. To retain it *in situ*. 3. To obviate any attendant or consequent symptoms.

To reduce requires extension, counter-extension, and coaptation.

LUXEUIL, MINERAL WATERS OF. Saline waters, at the town of Luxeuil, at the foot of the Vosges, in the department of Haute Saône. Five springs are thermal, and two cold. They seem to contain carbonic acid, carbonates of iron and lime, and chloride of sodium; and are employed as aperient, tonic, and stimulant.

LUZ, Sacrum.

LYCAN'CHE, *Lycan'chis*, from λυκος, 'a wolf,' and αγκω, 'I strangle.' *Wolf quinsy*, *Wolf choak*;—as *Cynanche* means *Dog choak*. Also, hydrophobia. See *Lycoides*.

LYCANCHIS, *Lycanche*.

LYCANTHROPE, see *Lycanthropia*.

LYCANTHROPIA, from λυκος, 'a wolf,' and ανθρωπος, 'a man.' *Lyc'oon*, *Ina'nia lupi'na*, *Lycoman'ia*. A variety of melancholy, in which the person believes himself to be changed into a wolf, and imitates the voice and habits of that animal. One so circumstanced is called a *Lycanthrope*, *Lycanthropus*.

LYCANTHROPUS, see *Lycanthropia*.

LYCAON, *Lycanthropia*.

LYCHNIDIUM, *Biolychnium*.

LYCHNIS GITHA'GO, *Agrostem'ma githa'go*, *Corn cockle*, *Rose campion*. A naturalized European plant. *Ord.* Caryophyllaceæ. The seeds are somewhat acrid, but not poisonous. They were, at one time, employed in cutaneous diseases, and in hemorrhage, and as a topical application to ulcers and fistule.

LYCHNIS OFFICINALIS, *Saponaria*—*L. Sylvestris*, *Saponaria*.

LYCHNIUM, *Biolychnium*.

LYCIUM, λυκίον. A medicinal substance, anciently obtained from the roots and branches of a thorny shrub growing in Lycia. It was astringent, and used in dysentery, ulcers of the gums, &c. By some, it has been considered to be the catechu.

LYCOÏDES, *Lycoides*, from λυκος, 'a wolf,' and αἰδης, 'form.' *Lupo sim'ilis*. An epithet for a species of cynanche, called, also, *Lupus*, and *Strangulator*, and absurdly attributed to excessive retention of sperm in its reservoirs and its passage into the blood [?].—Galen.

LYCOMANIA, *Lycanthropia*.

LYCOPE DE VIRGINIE, *Lycopus Virginicus*.

LYCOPER'DON, from λυκος, 'a wolf,' and πῦρ, 'I break wind.' *L. Bovis'ta* seu *Arrhizon* seu *Globosum*, *Bovis'ta*, *Crep'tus Lupi*, *Fungus Chirurgo'rum*, *Puff-ball*, *Bull Fists*, *Moltipuffs*, (Prov.) *Puff*, (Sc.) *Deil's snuff-box*, (F.) *Vesseloup*. *Ord.* Fungi. The puff-ball dries into a very fine, light brownish dust, which is sometimes used as a mechanical styptic to arrest hemorrhage.

LYCOPERDON ARRIZON, *Lycoperdon*—*L. Cervinum*, *Elaphomyces granulatus*—*L. Globosum*, *Lycoperdon*—*L. Gulosorum*, *L. Tuber*.

LYCOPERDON PRO'TEUS, *Common puff-ball*. The fumes of this substance have been found to produce anæsthetization in animals.

LYCOPER'DON TUBER, *L. Gulosorum*, *Tuber Agriocastanum* seu *Cibarium* seu *Gulosorum* seu *Nigrum*, *Tubera Terræ*, *Trubs*, *Truffle*, (pronounced *troofle*), (F.) *Truffe*. A fleshy, firm body, unknown in its mode of reproduction, which is found under ground, especially in different parts of France, Italy, &c., and is much esteemed as an aliment. It seems to belong to the champignons, and has a particular perfume. Aphrodisiac virtues have been ascribed to it.

A Virginia truffle is called *Tuckaho*.

LYCOPERSICUM ESCULENTUM, *Solanum lycopersicum*—*L. Pomum amoris*, *Solanum lycopersicum*—*L. Tuberosum*, *Solanum tuberosum*.

LYCOPODE, *Lycopodium*.

LYCOPODIUM, from λυκος, 'a wolf,' and πους, 'a foot.' *L. Clavatum* seu *Officinale*, *Lepidotis clavata*, *Cingularia*, *Muscus clavatus*, *Club-moss*, *Wolf's-claw*, *Witch-meal*. *Ord.* *Lycopodiaceæ*. This is the largest of the European mosses. The dust, which fills the capsules of its spikes, is very inflammable, and hence has been called *Vegetable Sulphur*;—and, also, *Farina* seu *Pulvis seu Semen seu Sulphur Lycopo'dii*, (F.) *Lycopode*, *Soufre végétal*. It is used in France and in the United States to roll pills and boluses in; and in medicine is a desiccative, in the excoaration to which infants are liable. It is collected chiefly in Switzerland and Germany for commerce.

LYCOPODIUM CLAVATUM, *Lycopodium*.

LYCOPODIUM COMPLANATUM, *Common Club-moss*, *Ground Pine*; indigenous; has the same properties as *Lycopodium selago*.

LYCOPODIUM OFFICINALE, *L. clavatum*—*L. Recurvum*, *L. Selago*.

LYCOPODIUM SELA'GO, *L. Recurvum*, *Planan'thus fatigatus* seu *selago*, *Muscus erectus* seu *Catharticus*, *Sela'go*, *Upright Club-moss*, *Fir Club-moss*, *Upright Fir-moss*. A decoction of this plant acts violently as an emetic and purgative; and was formerly, on this account, used as an abortive. It is also employed to kill vermin.

LYCOPUS, *L. Virginicus* seu *uniflorus* seu *pumila*, *Bugleweed*, *Water Bugle*, *Water Horehound*, *Gypsy Weed*, *Paul's Bet'ony*, (F.) *Lycopo de Virginie*. This indigenous herb, *Sec. Syst.* *Diandria Monogynia*, *Nat. Ord.* *Labiata*, grows throughout the greater part of the United States. The whole herb is said to be slightly narcotic. It is given in infusion, (*Lycop. ʒj; aq. fervent. Oj.*)

LYCOPUS PUMILA, *Lycopus*.

LYCOPUS SINUA'TUS, *Water Horehound*, *Gypsy Weed*, *Paul's Bet'ony*, indigenous, has similar properties.

LYCOPUS UNIFLORUS, *L. pumila*—*L. Virginicus*, *Lycopus*.

LYCOREXIS. *Fames lupina*.

LYCOSA TARENTULA, see Tarantula.

LYCOSTOMA, *Hyperoöchasma*.

LYE, Lixivium, Ley — *l.* Dyspeptic, see Fuligokali — *l.* Medical, see Fuligokali — *l.* Tea, Physick's, see Fuligokali.

LYGISMUS, Luxation.

LYGMODES, Singultus.

LYGMUS, Singultus.

LYGODES, Singultus.

LYING-IN STATE, Parturient state.

LYING-DOWN STATE, Parturient state.

LYMA, Lochia; from *λυω*, 'to wash,' also, filth or sordes, (to be washed away.) *λυμα*, *Lymē*.

LYME, Lyma.

LYMPH, *Lympha*, from *νυμφη*, 'water,' by changing *ν* into *λ*. *White blood*, *Lympha Nutritiva*. A name given to the fluid contained in the lymphatic vessels and thoracic duct of animals, which have been made to fast for 24 hours. According to Chevreul, the lymph of the dog contains water, fibrin, albumen, common salt, subcarbonate of soda, phosphates of lime and magnesia, and carbonate of lime. The properties and composition of lymph vary somewhat according to the part whence the lymphatic vessels obtain it. Generally, it is under the form of a transparent, slightly alkaline fluid, sometimes of a madder-red or yellowish colour, — of a spermatic odour, and saline taste; soluble in water, — the solution becoming turbid, when mixed with alcohol. When left to itself, it coagulates. The clot or solid portion becomes of a scarlet red, if put in contact with oxygen; and of a purple red, if placed in carbonic acid.

Like the blood, the lymph consists of a fluid — *Liquor lymphæ* — in which *lymph corpuscles* or *globules* are suspended.

The lymph is probably the product of internal absorption in different parts of the body: it then flows along the lymphatic vessels, uniting with the chyle, and is poured with it into the veins; thus becoming one of the materials of the blood. According to others, the blood, when it reaches the arterial radicles, is divided into two portions, — the one red, which is carried to the heart, — the other serous or white, which is absorbed by the lymphatic vessels, and constitutes the lymph. By others, again, the lymphatics are considered to be the vessels of return for the white blood sent to certain tissues. White blood, however, has probably no existence.

The word *lymph* is sometimes used unhappily by the surgeon to signify *liquor sanguinis*.

LYMPH, COAGULABLE, Fibrin, *Liquor sanguinis*.

LYMPH CORPUSCLES, see Lymph — *l.* Globules, see Lymph — *l.* Hearts, Lymphatic hearts.

LYMPH, PLASTIC, *Liquor sanguinis*.

LYMPHA, Lymph — *l.* Arborum, see Sap — *l.* Muculenta narium, Nasal mucus — *l.* Nutritia, Lymph — *l.* Pancreatis, see Pancreas — *l.* Pericardii, see Pericardium — *l.* Plastica, Fibrin.

LYMPHADENITIS: from *lymph*, and *adenitis*, 'inflammation of a gland.' *Inflammatio glandularum lymphaticarum*, *Adenitis lymphatica*, (F.) *Adénite lymphatique*. Inflammation of a lymphatic gland or ganglion.

LYMPHÆDUCTUS, Lymphatic vessels.

LYMPHANGIITIS, Angioloecitis.

LYMPHANGIA, Lymphatic vessels.

LYMPHANGIECTASIS, *Lymphœury'sma*; from *lymph*, *αγγειον*, 'a vessel,' and *εκτασις*, 'dilatation.' Dilatation of lymphatic vessels.

LYMPHANGIOGRAPHY, *Lymphangiographia*; from *lymph*, *αγγειον*, 'a vessel,' and *γραφη*, 'a description.' A description of the lymphatic vessels.

LYMPHANGIOITIS, Angioloecitis.

LYMPHANGIOLOGY, from *Lymphangion*,

'a lymphatic,' and *λογος*, 'a description.' A treatise on the lymphatics.

LYMPHANGIOTOMY, *Lymphangiotoμία*; from *lymph*, *αγγειον*, 'a vessel,' and *τομή*, 'incision.' The anatomy or dissection of the lymphatic vessels.

LYMPHANGITIS, Angioloecitis.

LYMPHONCUS, *Lymphoncus*; from *lymph*, *αγγειον*, 'a vessel,' and *ογκος*, 'a tumour.' Tumefaction of the lymphatics.

LYMPHATIC, *Lymphaticus*. That which relates to lymph; also, abounding in lymph. See Phlegmatic.

LYMPHATIC HEARTS, *Lymph Hearts*. The frog and several other animals are provided with large receptacles for the lymph immediately underneath the skin, which exhibit distinct and regular pulsations like the sanguiferous heart. Their use appears to be — to propel the lymph.

LYMPHATIC SYSTEM is that particular system of organs which is inservient to the formation and circulation of lymph. These organs are:

1. LYMPHATIC GLANDLIONS OR GLANDS, see Conglobate.

2. LYMPHATIC VESSELS, *Lymphæductus*, *Lymphangi'a*, *Lymphang'e'a*, *Vena lymphatica*, *Ductus serosi*, *Vasa lymphatica seu resorbentia seu hydragoga*, *Hydrangi'a*. These are very numerous. Arising at the surface of membranes and in the tissue of the organs, they carry into the veins the lymph from those parts. Lymphatic vessels are found in every part of the body. Wherever they are met with, however, they form two orders, — one *superficial*, the other *deep-seated*; — the two orders frequently communicating with each other. Lymphatic vessels are generally smaller than arteries and veins. They are very thin, diaphanous, and cylindrical; but present, here and there, more or less considerable dilatations, caused by valves in their interior. They are slightly tortuous in their course; their anastomoses are very numerous, and they often cross each other, forming successive plexuses. Of the arrangement of the extreme radicles we are ignorant. All the branches, before joining the principal trunks, pass through lymphatic ganglions, in which they are convoluted, or subdivided almost *ad infinitum*. They are formed of an outer areolar membrane and an internal coat, similar to that of the veins; of the latter, the valves are formed. All the lymphatics of the body ultimately discharge themselves at the junction of the subclavian and internal jugular veins. Two of these trunks are considerably larger than the others, — the *thoracic duct*, and the great trunk of the right side, (F.) *La grande veine lymphatique droite*. The former receives the lymphatics of the abdomen, of the lower extremities, the left side of the thorax, the left upper extremity, and the corresponding side of the head and neck; the latter receives those of the right upper extremity, and of the right side of the head, neck and thorax.

LYMPHATITIS, Angioloecitis.

LYMPHE COAGULABLE, *Liquor sanguinis* — *l.* de Cotugno, Cotunnus, liquor of — *l.* Plastique, *Liquor sanguinis*.

LYMPHÆURYSMA, Lymphangiectasis.

LYMPHITIS, Angioloecitis.

LYMPHIZATION. A term used by Professor Gross to signify effusion of coagulable lymph. It is not a happy word, inasmuch as lymph has another meaning, whilst the term 'coagulable lymph' is now almost abandoned.

LYMPHOCHÆZIA, from *lymph*, and *χρησ*, 'I go to stool.' Serous diarrhœa.

LYMPHONCUS, Lymphangoncus — *l.* Iridis, Iridæxesis.

LYMPHOPYRA, Fever, adeno-meningeal.

LYMPHORRHAG'IA, from *lymph*, and *ρρηνυμι*, 'I break forth.' Discharge of lymph from a lymphatic vessel, owing to a lesion of its coats.

LYMPHO'SIS. The preparation or elaboration of lymph.

LYMPHOT'OMY, from *lymph*, 'lymph,' and *τεμνω*, 'I cut.' Dissection of the lymphatics.

LYNCH'S EMBROCA'TION. An emollient nostrum, formed of olive oil impregnated with bergamot and other essences, and coloured with alkanet root.

LYNGODES, Singultous — 1. Febris, Fever, singultous.

LYNGYODES, Singultous.

LYNN WAHOO, Ulmus alata.

LYNX, Singultus.

LYPE, Athymia.

LYPEMANIA, Melancholy.

LYPE'RIA CRO'CEA. A South African plant, *Nat. Ord.* Scrophulariaceæ, the flowers of which closely resemble saffron in smell and taste, and possess similar medical virtues.

LYPEROPHRE'NIE, Melancholy.

LYPOMA, Lipoma.

LYPOTHYM'IA, from *λυπη*, 'sadness,' and *ψυχο*, 'the mind.' Very great sadness or despondency.

LYRA, *λυρα*, 'the lyre.' *Psalter*, *Psalterium*, *Corpus Psalloi'des*, *Lyra Da'vidis*, *Lam'ina medulla'ris triangularis cer'ebr'i*, *Spatium trigonum*. The under surface of the posterior part of the body of the fornix is impressed with numerous transverse and oblique lines, which have been so called from some resemblance they bear to the ancient lyre.

LYRINGIUM, *Eryngium campestre*.

LYSIA, Lysis.

LYSIMA'CHIA NUMMULA'RIA, *L. nem'orum*, *Nummularia*, *Hirundinaria*, *Centimor'bia*, *Creeping Loose Strife*, *Money Wort*, (F.) *Herbe aux œufs*. *Ord.* Primulacæ. This plant was formerly accounted vulnerary. It has been considered to possess antiscorbutic and astringent qualities, and has been used in leucorrhœa.

LYSIMACHIA PURPUREA, *Lythrum salicaria*.

LYSIMA'CHIA QUADRIFO'LIA, *Four-Leaved Loose Strife*, *Crosswort*, an indigenous plant, is astringent; and has been used as a stomachic and antiperiodic.

LYSIS, *Λυσις*, *λυσις*, from *λυω*, 'I dissolve.' Solution. A common suffix; also, a name given to solutions or terminations of disease, which are operated insensibly; that is, gradually and without critical symptoms.

LYSSA, Hydrophobia — 1. Canina, Hydrophobia.

LYSSAS, Maniodes.

LYSSETER, Maniodes.

LYSSODEC'TUS, from *λυσσα*, 'canine madness,' and *δηκω*, or *δακνω*, 'I bite.' *Hydroph'obus*; *Canē rab'ido morens*. One who has been bitten by a mad dog, or is actually labouring under hydrophobia.

LYSSODEG'MA, *Lyssodeg'mus*, *Lyssodec'is*, same etymon. The bite of a mad dog.

LYSSODEXIS, *Lyssodegma*.

LYTE'RIOS, *λυτηριος*, (from *λυω*, 'I dissolve,') 'solving.' An epithet given to those signs which announce the solution of a very violent disease.

LYTHRUM ALATUM, see *Lythrum salicaria* — 1. *Hyssopifolia*, *Lythrum salicaria*.

LYTHRUM SALICA'RIA, from *λυθρον*, 'blood, and dust,' on account of the colour of the flowers of some species, *Lysima'chia purpu'rea*, *Salica'ria vulga'ris seu epica'ta*; — the Common or Purple Willow Herb. *Ord.* Lythracæ. The herb, root, and flowers possess a considerable degree of astringency, and are used, occasionally, in the cure of diarrhœa and dysentery, leucorrhœa, hæmoptysis, &c.

LYTHRUM HYSSOPIFO'LIA, and *L. ALA'TUM*; indigenous, have similar properties.

LYTTA CÆRULEA, see *Cantharis* — 1. *Gigas mas*, see *Cantharis* — 1. *Ruficeps*, see *Cantharis*.

LYTTA RU'PIES. *Class*, Insecta; *Ord.* Coleoptera. A variety of *Lytta*, peculiar to Chili, which is more active as a vesicant than the *cantharis*, or the *Lytta vittata*.

LYTTA SEGETUM, see *Cantharis* — 1. *Vesicatoria*, *Cantharis*.

LYTTA VITTA'TA, *Cantharis vittata*, *Pota'to Fly*, (F.) *Cantharide tachetée*. Four species of meloë that blister, are found in the United States. The *lytta vittata* feeds principally upon the potato plant, and, at the proper season of the year, may be collected in immense numbers. The potato fly resembles the *cantharides* in every property, and is fully equal to them.

M.

M. This letter signifies, in prescriptions, *manipulus*, 'a handful.' Also, *misce*, 'mix.' See Abbreviation.

MACAPATLI, *Smilax sarsaparilla*.

MACARO'NI. An alimentary paste, moulded of a cylindrical shape, and formed of rice or wheaten flour. It is eaten — when boiled — in soup; prepared with cheese, &c. See *Turandæ Italica*.

Also, a name formerly given to a pulverulent compound of sugar and glass of antimony, carried into France by the Italian monks, and employed at the hospital *La Charité* in the treatment of painters' colic.

MACARONS, Biscuits, medicinal.

MACE, see *Myristica moschata* — m. Reed, *Typha latifolia*.

MACEDONISIUM, *Smyrniolum olusatrum*.

MACER, *Gracilis*.

MACERA'TION, *Macera'tio*, from *macere*, *maceratum*, 'to soften by water.' An operation which consists in infusing, usually without heat, a solid substance in a liquid, so as to extract its virtues.

MACERONA, *Smyrniolum olusatrum*.

MACES, see *Myristica moschata*.

MACESCERE, Emaciate.

MACHÆRA, Culter, Knife, Penis.

MACHERIDION, *Machærian*.

MACHÆ'RION, *Machæ'rius*, *Machærid'ion*, *Gladi'olus*, from *μαχαίρα*, 'a sword,' *μαχρ*, 'a fight.' A knife. An amputating knife. Rufus of Ephesus asserts, that the Aruspices gave this name to a part of the liver of animals.

MACHÆRIS, Knife, Novacula.

MACHA'ON, from *μαχαιω*, 'I desire to fight.' The son of Æsculapius, and a celebrated physician.

MACHAÖNIA seu **MACHAÖN'ICA** ARS. Medicine. The *Healing Art*:—so called after Machaon, the son of Æsculapius.

MACH'INAL, *Mechan'icus*. This epithet is added especially by French writers to the word *movement*, to express that the will takes no part in it.

MACHINE', *Mach'ina*, *Machinamen'tum*, *Mech'anê*, *Mechan'e'ma*. A more or less compound instrument or arrangement for putting a body in motion, or for producing any action whatever. Physiologists sometimes use it for the *animal body*;—as the *machine* or *animal machine*.

MACHLOSZYNE, *Nymphomania*.

MACHOIRE, Maxillary bone—*m. Diacranienne*, Maxillary bone, lower—*m. Syncratiennne*, Maxillary bone, superior.

MACIES, Atrophy, Emaciation—*m. Infantum*, *Tabes mesenterica*.

MACILENTUS, *Gracilis*.

MACIS, see *Myristica moschata*.

MACRAU'CHEN, from *macro*, and *αυχην*, 'the neck.' *Longo collo præ'ditus*. One who has a long neck.—Galen.

MACRE FLOTTANTE, *Trapa natans*.

MACRITUDO, Emaciation.

MACROBIOSIS, Longevity.

MACROBIOTES, Longevity.

MACROBIOT'IC, *Macrobiot'icus*, *Macrobi'otus*, *Macro'bius*, *Longæ'vus*, from *macro*, and *βίος*, 'life.' That which lives a long time. The *macrobiotic art*, *Macrobiot'ics*, (F.) *Macrobiotique*, is the art of living a long time.

MACROBIOTUS, *Macrobiotic*.

MACROBIUS, *Macrobiotic*.

MACROCEPHALIA, see *Capitones*.

MACROCEPH'ALUS, *Makroceph'alus*, (F.) *Macrocéphale*; from *macro*, and *κεφαλή*, 'head.' *Qui magnum habet caput*. 'One who has a large head.' This epithet is given to children born with heads so large that they seem to be hydrocephalic; but in which the unusual development is owing to a large size of the brain. Such are supposed to be more than ordinarily liable to convulsions. [?] The term has also been applied by Hippocrates to certain Asiatics who had long heads. See *Capitones*.

MACROCHI'RIA, from *macro*, and *χείρ*, 'hand.' Monstrosity, characterized by excessive development of the hands.

MACROCO'LIA, from *macro*, and *κῶλον*, 'a limb.' Great length of limbs in general, and of the lower limbs in particular.

MAC'ROCOSM, *Macrocosm'us*, from *macro*, and *κόσμος*, 'world.' The universe, in contradistinction to *Microcosm*, which means man.

MACRODACTYLUS, from *macro*, and *δάκτυλος*, 'a finger.' Having long fingers.

MACROGASTER PLATYPUS, *Acarus folliculorum*.

MACROGLOSSA, see *Macroglossus*.

MACROGLOSSUS, from *macro*, and *γλῶσσα*, 'tongue.' One who has a very large or prolapsed tongue. See *Paraglossa*.

MACROMEL'IA, from *macro*, and *μέλος*, 'limb.' Monstrosity, which consists in the excessive size of some limb.

MACRONOSIÆ, Chronic diseases.

MACROPHAL'LUS, from *macro*, and *φαλλος*, 'the male organ.' A large size of the male organ.

MACROPHO'NUS, from *macro*, and *φωνή*, 'voice.' One who has a strong voice.

MACROPHTHAL'MUS, from *macro*, and *ὀφθαλμος*, 'eye.' Having large eyes.

MACROPHYSOCEPH'ALUS, from *macro*, *φύσα*, 'air,' and *κεφαλή*, 'head.' A word used by Ambrose Paré to designate an augmentation of

the head of the fœtus, produced by a sort of emphysema [?], which retards delivery.

MACROP'IPER, *Piper longum*.

MACROP'NCE'A, from *macro*, and *πνεύω*, 'I breathe.' A long and deep respiration.

MACROP'NUS, *Macrop'nois*. Same etymon. Long breathed. One who breathes slowly:—a word met with in some authors.—Hippocrates.

MACROP'ODUS, *Mac'ropus*, *Megal'opus*, from *macro*, and *πους*, *πόδος*, 'foot.' One who has a large foot.

MACROPROSO'PIA, from *macro*, and *προσωπον*, 'face.' Monstrosity, characterized by excessive size of face.

MACROR, Emaciation.

MACROR'RHIS, from *macro*, and *ῥις*, or *ριν*, 'nose.' One who has a long nose.

MACROS, *μακρός*, 'great.' Long.

MACROS'CELES, *Cyura longa habens*; from *macro*, and *σκελος*, 'the leg.' One who has long legs.

MACROSIÆ, Chronic diseases.

MACROSOMATIA, see *Giant*.

MACROSOMIA, see *Giant*.

MACRO'TES, from *macro*, and *ους*, *ωτος*, 'an ear.' One who has long ears.

MACROT'RY, *RACEMOSA*, see *Actæa racemosa*.

MAC'ULA. A spot. *Dyschro'a*, *Celia*, *Labea*, (F.) *Tache*. A permanent discoloration of some portion of the skin, often with a change of its texture. *Ephelis*, *Nævus*, *Spilus*, &c., belong to *Maculæ*.

MACULA, Molecule—*m. Aurea*, see *Foramen centrale retinae*—*m. Cornæa*, *Caligo*—*m. Cornæa arcuata*, *Gerotoxon*—*m. Cornæa margaritacea*, *Paralampsis*—*m. Cribrosa*, see *Auditory canal*, internal—*m. Emortualis*, see *Suggillation*—*m. Flava*, see *Foramen centrale retinae*—*m. Fusca*, *Epelides*—*m. Germinativa*, see *Molecule*—*m. Hepatica*, *Chloasma materna*, *Nævus*—*m. Lenticularis*, *Ephelides*—*m. Lutea retinae*, see *Foramen centrale*—*m. Matricialis*, *Nævus*—*m. Matricis*, *Nævus*—*m. Nigra*, see *Optic nerve*—*m. Solaris*, *Ephelides*.

MAC'ULÆ ALBÆ, *M. Lac'tea*, *In'sulæ*, *Milk-spots*. White spots, seen on serous membranes, especially on that investing the heart, and which appear to be the result of previous inflammatory action.

MACULÆ ANTE OCULOS VOLITANTES, *Metamorphopsia*—*m. Lactea*, *M. Albæ*.

MACULOSUS, *Sablé*.

MAD, Insane—*m. Doctor*, see *Psychiatria*.

MADAR, *Mudar*.

MADARO'SIS, from *μαδρος*, 'bald.' *Madaro'ma*, *Made'sis*, *Madar'otes*, *Made'ma*, *Mad'isis*, *Deplumatio*, *Milphæ*, *Milpho'sis*, *Pilo'sis*, *Calvi'ties*. Loss of the hair, particularly of the eyelashes.

MADAROTES, *Madarosis*.

MADDER, *DYERS'*, *Rubia*.

MADEIRA, CLIMATE OF. This island is much frequented by pulmonary invalids, on account of the mildness and equability of its climate. Owing, indeed, to the mildness of the winter, and the coolness of the summer, together with the remarkable equality of the temperature during day and night, as well as throughout the year, it has been considered that the climate of Madeira is the finest in the northern hemisphere. Sir James Clark is of opinion that there is no place on the continent of Europe, with which he is acquainted, where the pulmonary invalid could reside with so much advantage, during the whole year, as in Madeira.

MADELEON, *Bellium*.

MADEMA, *Madarosis*.

MADESIS, *Depilation*, *Madarosis*.

MADISIS, Depilation, Madarosis.

MADISON SPRINGS, see Georgia, Mineral waters of.

MADISTERIUM, μαδιστηριον, *Trichola'bium*, *Trichol'ubia*, *Voelsell'a*. Tweezers. An instrument for extracting hairs.

MADJOUND, *Madjoun*. A mixture of powdered hemp and honey, used as a narcotic agent, like tobacco, by the Algerines.

MADNESS, CANINE, Hydrophobia — *m*. Raving or furious, Mania.

MADOR, Moisture. A cold sweat.

MADORIUS, Mudar.

MADREPORA OCOLATA, see Coral.

MADWEED, *Scutellaria lateriflora*.

MÆA, Midwife.

MÆEIA, Obstetrics.

MÆEUMA, see Parturition.

MÆIA, Obstetrics.

MAESA PICTA, Saoria.

MÆUSIS, Parturition.

MÆUTRIA, Midwife.

MAGDALIA, *Magda'leon*, *Magda'lis*. 'Crumb of bread.' Any medicine, as a pill, formed of crumb of bread. A roll of plaster.

MAGEIRICE, Culinary art.

MAGGOT PIMPLE, see Acne.

MAGIS, *payis*, 'a cake.' A sort of cake, composed of cloves, garlic and cheese, beaten together. — Hippocrates.

MAGISTERIUM, Magistery — *m*. Bismuthi, Bismuth, subnitrate of — *m*. Jalapæ, Resin of Jalap — *m*. Marcasitæ, Bismuth, subnitrate of — *m*. Plumbi, Plumbi subcarbonas — *m*. Sulphuris, Sulphur lotum — *m*. Tartari purgans, Potassæ acetas.

MAG'ISTERY, *Magist'rium*, from *magister*, 'a master.' Certain precipitates from saline solutions were formerly so called; as well as other medicines, the preparation of which was kept secret.

MAGISTERY OF BISMUTH, Bismuth, subnitrate of.

MAG'ISTRAL, *Magistra'lis*. Same etymon. *Extempora'neus*. A prescription is so called — *Form'ula magistra'lis seu extempora'nea* — which is prepared extemporaneously; an official prescription — *Form'ula officina'lis* — being such as has been prepared for some time before it is prescribed.

MAGISTRANTIA, Imperatoria.

MAGMA, *payma*, (F.) *Marc*. The thick residuum, obtained after expressing certain substances to extract the fluid parts from them. The grounds which remain after treating a substance with water, alcohol, or any other menstruum. Also, a salve of a certain consistence.

MAGMA or MARC OF OLIVES is the residuum after the greatest possible quantity of oil has been extracted from olives by making them ferment. It was formerly employed as a stimulant, under the form of a bath — to which the name *Bain de Marc* was given by the French.

MAGMA or MARC OF GRAPES, *Bry'tia*, was once employed for the same purposes.

MAGMA RÉTICULÉ, 'reticulated magma.' The gelatiniform substance found between the chorion and amnion in the early period of embryonic existence.

MAGNES, Magnet.

MAGNES ARSENICA'LIS. (*Sulphur, white arsenic*, and common *antimony*, of each equal parts. Mix by fusion.) It is corrosive. See Magnetic plaster.

MAGNES EPILEPSIÆ, *Hydrargyri sulphuretum rubrum*.

MAGNE'SIA, *Abarna'as*, *Chambar*, *Terra amara*, *Magnesia terra*, *Talc earth*; from *magne's*, 'the magnet.' An alkaline earth whose me-

tallic base is *magnesium*. The term *Magnesia* was employed, of old, for any substance that seemed to have the power of attracting some principle from the air. At present it is restricted to —

MAGNESIA, *M. usta seu calcina'ta*, *Cal'cined Magnesia*, *Oxide of magn'esium*, (F.) *Magnésie brûlée*, *Magnésie*, *M. Caustique*. This is obtained by exposing carbonate of magnesia to a strong heat. It is inodorous; taste very slightly bitter; in the form of a white, light, spongy, soft powder. S. g. 2.3; requiring 2000 times its weight of water for its solution. It is antacid, and laxative when it meets with acid in the stomach. Dose, gr. x to ʒi in water or milk.

Henry's, *Husband's*, and *Ellis's Magnesia*, are dense forms of this preparation.

MAGNESIA, ACETATE OF, *Magnesia acetas* — *m*. *Aerata*, *Magnesia carbonas* — *m*. *Alba*, *M. carbonas* — *m*. *Calcinata*, *M. usta* — *m*. *Citrate* of, *Magnesia citras* — *m*. *Dinneford's*, see *Magnesia*, fluid — *m*. *Edinburgensis*, *M. carbonas*.

MAGNESIA, EFFERVESCING, *MOXON'S*. (*Magnes. carb.*; *M. sulphat.*; *Soda bicarbon.*, *Acid tartaric.* *aa partes æquales*; to be pulverized, well dried, mixed, and enclosed in bottles hermetically sealed.) Dose, a teaspoonful in half a tumbler of water, drunk in a state of effervescence.

MAGNESIA, ELLIS'S, see *Magnesia*.

MAGNESIA, FLUID. Under this name a preparation is designated, which consists of a solution of carbonate of magnesia in carbonated water. It is also termed *carbonated magnesia water*, *aërated magnesia water*, and *condensed solution of magnesia*, (F.) *Eau magnésienne*. *Dinneford's Magnesia* is a solution of this kind.

MAGNESIA, HENRY'S, *Magnesia* — *m*. *Husband's*, see *Magnesia* — *m*. *Mitis*, *Magnesia carbonas* — *m*. *Muriate* of, *Magnesi chloridum* — *m*. *Nigra*, *Manganese*, black oxide of.

MAGNESIA OPALI'NA. A name given by Léméry to a mixture of equal parts of *antimony, nitrate of potassa*, and *chloride of sodium*, (decrepitated.) It has emetic properties, but is not used.

MAGNESIA SALIS AMARI, *Magnesia carbonas* — *m*. *Salis Ebsdamiensis*, *Magnesia carbonas* — *m*. *Saturni*, *Antimonium* — *m*. *Solution* of, *condensed*, *M. fluid* — *m*. *Subcarbonate* of, *Magnesia carbonas* — *m*. *Subcarbonate* of, *hydrated*, *Magnesia carbonas* — *m*. *Terra*, *Magnesia* — *m*. *Vitriolata*, *Magnesia sulphas* — *m*. and *Soda*, *sulphate* of, see *Soda*, *sulphate* of — *m*. *Usta*, *Magnesia*.

MAGNE'SIÆ ACETAS, *Ac'etate of Magn'esia*. This salt is prepared by saturating carbonate of magnesia with *acetic acid*. It is of a syrupy consistence, and about four ounces is a dose. It acts like citrate of magnesia.

MAGNE'SIÆ CAR'ONAS seu *Subcarbonas*, *Magnesia* (Dublin), *M. aëra'ta seu carbon'ica seu alba*, *M. Subcarbon'ica seu Mitis* seu *Edinburgensis* seu *Salis Ebsdamiensis* seu *Salis ama'ri*, *Carbonas magn'esicum*, *Lac terra*, *Hypocarbonas magn'esia*, *Terra amara aëra'ta seu absorbens minera'lis seu talco'sa oxyanthraci'des*, *Subcarbonate of Magnesia*, *Hydrated Subcarbonate of Magnesia*, (F.) *Sous-carbonate ou carbonate de Magnésie*, *Magnésie aérée ou blanche ou erayuse ou douce ou effervescente ou moyenne ou Anglaise*, *Craie ou Terre magnésienne*, *Lait de Terre*, *Panacée Anglaise*, *Poudre de Sentinelli* ou de *Valentini* ou du *Comte de Palme* ou de *Zwinger*. Prepared from sulphate of magnesia by subcarbonate of potassa. It is inodorous; insipid; light; white; spongy; opaque; effervescing with acids; insoluble in water. Properties the same as the last; but the carbonic acid, when set free, sometimes causes unpleasant distension.

MAGNE'SIÆ CITRAS, *Citrate of Magnesia*, (F.)

Citrate de Magnésie. A saline preparation, formed by saturating a solution of citric acid with either magnesia or its carbonate. Dose, an ounce. It is devoid of the bitter taste of the magnesian salts.

A solution in water, or in mineral water, sweetened with syrup, and acidulated with citric acid, makes an agreeable purgative. A simple solution in water has been called *magnesian lemonade*. In the effervescent state, it is the *effervescent magnesian lemonade*. The Pharmacopœia of the United States (1851) has a form for the LIQUOR MAGNESIÆ CITRATIS, *Solution of Citrate of Magnesia*. Take of Carbonate of magnesia, ℥v; citric acid, ℥viiss; Syrup of citric acid, f℥ij; Water, a sufficient quantity. Dissolve the citric acid in f℥iv of water, and add ℥iv of the carbonate of magnesia, previously rubbed with f℥ij of water. When the reaction has ceased, filter into a strong f℥xij glass bottle, into which the syrup of citric acid has been previously introduced. Rub the remaining carbonate of magnesia with f℥ij of water, and pour the mixture into the bottle, which must be well corked, and secured with twine; and shake the mixture occasionally until it becomes transparent. As a cathartic the whole of this may be taken.

MAGNESIÆ HYPOCARBONAS, *M. carbonas*—m. Subcarbonas, *M. carbonas*—m. Vitriolicum, *Magnesiæ sulphas*.

MAGNESIÆ SULPHAS, *Sulphas Magnesiæ purificata*, *Magnesia vitriolata*, *Sal catharticus amarus* seu *catharticum amarum* seu *amarum* seu *Anglicanum* seu *Epomœnis* seu *catharticum Anglicanum* seu *Sedlicen'sis* seu *Ebsdamen'se* seu *Schützen'se*. Terra amara sulphurica, *Vitriolicum Magnesiæ*, *Sulphate of Magnesia*, *Epsom Salt*, *Bitter purging Salt*, (F.) *Sulfate de magnésie*, *Sel admirable de Lémery*, *Sel d'Egra*. Generally obtained from sea-water. Its taste is bitter and disagreeable. It is soluble in an equal quantity of water at 60°. It is purgative and diuretic. Dose, as a cathartic, ℥ss to ℥ij.

MAGNESIAN LEMONADE, *Magnesiæ citras*.

MAGNÉSIE AÉRÉE, *Magnesiæ carbonas*—m. *Anglaise*, *Magnesiæ carbonas*—m. *Blanche*, *Magnesiæ carbonas*—m. *Brûlée*, *Magnesia usta*—m. *Carbonate de Magnesiæ carbonas*—m. *Caus-tique*, *Magnesia usta*—m. *Citrate de Magnesiæ citras*—m. *Crayeuse*, *Magnesiæ carbonas*—m. *Douce*, *Magnesiæ carbonas*—m. *Effervescente*, *Magnesiæ carbonas*—m. *Moyenne*, *Magnesiæ carbonas*—m. *Souscarbonate de Magnesiæ carbonas*—m. *Sulfate de Magnesiæ sulphas*.

MAGNÉSII CHLORIDUM, *Chloride of Magnesium*, *Muriate of Magnesia*. This bitter deliquescent salt has been given as a mild and effective cholagogue cathartic, in the dose of half an ounce to the adult. Being deliquescent, it may be kept dissolved in its weight of water.

MAGNESIUM, see *Magnesia*—m. *Chloride of Magnesiæ chloridum*—m. *Oxide of Magnesia usta*.

MAGNET, *Magnes*, *Magne'tes*, *Ferrum magnes attractorium*, *Sideri'tes*, *Sideri'tis*, *Lapis heracleus* seu *Syderi'tis* seu *nauteicus*, *Magni'tis*, so called from *Magnes*, its discoverer, or from *Magnesia*, where it is said to have been first found. (F.) *Aimant*, *Pierre d'Aimant*. The magnet or loadstone. An amorphous, oxydulated ore of iron, which exerts an attraction on unmagnetized iron, and has the property of exhibiting poles; that is, of pointing by one of its extremities to the north. This ore, by constant or long rubbing, communicates its properties to iron; and thus artificial magnets are formed. Magnetic ore is found in many countries, and particularly in the island of Elba. The magnet is sometimes

used to extract spicula of iron from the eye or from wounds. It has been employed as an antispasmodic; but acts only through the imagination. The powder has been given as a tonic. In Pharmacy, it is used to purify iron filings. It attracts the iron, and the impurities remain behind. It formerly entered, as an ingredient, into several plasters, to draw bullets and heads of arrows from the body—as the *Emplastrum divinum Nicolai*, the *Emplastrum nigrum* of Augsburg, the *Opodeldoch*, and *Attractivum* of Paracelsus, &c.

MAGNETES, *Magnet*.

MAGNETIC, *Magne'ticus*, (F.) *Magnétique*. Same etymon. That which belongs or relates to magnetism;—mineral or animal.

MAGNETIC FLUID. A name given to the imponderable fluid to which the magnet owes its virtues. By analogy it is applied to a particular principle, supposed to be the source of organic actions, which affects, it is conceived, the nervous system principally, and is susceptible of being transmitted from one living body to another, by contact or simple approximation, and especially under the influence of fixed volition. See *Magnetism*, *Animal*.

MAGNETIC PLASTER. A plaster, at present, not used. It has for its base a mixture, called *Magnes arsenicalis*; formed of equal parts of antimony, sulphur and arsenic melted together in a glass cucurbit. The name *Magnetic plaster* was, likewise, given to such as contained powdered magnet.

MAGNETINUS, *Potassæ supertartaras impurus*.

MAGNETISM, ANIMAL, *Mesmerism*, *Path-etism*, *Neurog'mia*, *Bioga'mia*, *Biomagnetis'mus*, *Zoëmagnetis'mus*, *Ezoon'erism* (proposed by Mr. H. Mayo), *Telluris'mus*, *Anthropomagnetis'mus*, *Garg'alis'mus*, *Garg'alia*. Properties attributed to the influence of a particular principle, which has been compared to that which characterizes the magnet. It is supposed to be transmitted from one person to another, and to impress peculiar modifications on organic action, especially on that of the nerves. The discussions, to which this strange belief has given rise, are by no means terminated. There is no evidence whatever of the existence of such a fluid. Highly impressible persons can be thrown into a kind of hysteric or 'magnetic' sleep and somnambulism, (designated by Mr. Braid, *hypnotism*, *neuro-hypnotism*, and *nervous sleep*); but farther than this, the efforts of the magnetizer cannot reach. It is a mode of action upon the nerves through the medium of the senses.

MAGNETIZATION, *Mesmerization*.

MAGNETIZED, *Mesmerized*.

MAGNETIZER, *Mesmerizer*.

MAGNETIS, *Magnet*.

MAGNITUDO CORPORIS, *Stature*—m. *Gigante*, see *Giant*.

MAGNOC, *Jatropha manihot*.

MAGNOLIA FRAGRANS, *M. glauca*.

MAGNOLIA GLAUCA, *Magnolia*, *M. fragrans*, *Small Magnolia*, *Swamp Sweetgum*, *Elk Bark*, *Indian Bark*, *White Laurel*, *Sweet Bay*, *Beaver Wood*, *Beaver Tree*, *White Bay*, *Cinchona of Virginia*, *Castor Bay*, *Sweet or Laurel magnolia*, (F.) *Magnolia glauca*. Ord. *Magnoliaceæ*. The bark is possessed of tonic properties, resembling those of cascarilla, canella, &c. The same may be said of the *Magnolia tripet'ala* seu *Umbrella*, *Umbrel'la tree*; the *M. acuminata* or *Cucumber tree*, the *M. grandiflora* and *M. macrophylla*, *Laurel*, *Elk wood*, *Silverleaf*, *Big leaf*, *White Bay*, *Beaver Tree*, *Elk bark*, *Big bloom*. *Magnolia* (Ph. U. S.) is the bark of *Magnolia glauca*, *M. acuminata*, and *M. tripet'ala*.

MAGNOLIER GLAUQUE, *Magnolia glauca*.

MAGNUM DEI DONUM, *Cinchona*.

MAGNUM OS. The third bone of the lower row of the carpus, reckoning from the thumb. It is the largest bone of the carpus; and is, also, called *Os capitatum*; (F.) *Grand Os*.

MAGRUMS. A popular name in the State of New York for a singular convulsive affection, which resembles chorea. It rarely, however, occurs before the adult age; never ceases spontaneously, and, when fully developed, is devoid of any paroxysmal character.

MAGUEY, *Agave Americana*.

MAHMOUDY, *Convolvulus scammonia*.

MAHOGAN FÉBRIFUGE, *Swietenia febrifuga*.

MAHOGANY, *Gynocladus Canadensis*. Also, a drink in use among the Cornish fishermen of England, which is made of two parts gin and one part treacle, well beaten together. An analogous drink—a mixture of whisky and honey, (see Mead, under *Hydromel*), is known in the Highlands of Scotland by the name *Atol porridge*.

MAHOGANY MOUNTAIN, *Betula lenta*—m. Tree, *Swietenia mahogani*.

MAIANTHEMUM, *Convallaria maialis*.

MAIDENHAIR, *Adiantum capillus veneris*—m. American, *Adiantum pedatum*—m. Canada, *Adiantum pedatum*—m. Common, *Asplenium trichomanoides*—m. Golden, *Polytrichum*—m. White, *Asplenium ruta muraria*.

MAIDENHEAD, *Virginity*.

MAIDENHOOD, *Virginity*.

MAIEIA, *Obstetrics*.

MAIEUMA, see *Parturition*.

MAIEUSIS, *Parturition*.

MAIEUTA, see *Parturition*.

MAIEUTER, *Accoucheur*.

MAIEUTES, *Accoucheur*.

MAIEUTICA ARS, *Obstetrics*.

MAIGREUR, *Emaciation*.

MAILLET, *Mallet*.

MAILLOT, *Swathing clothes*.

MAIN, *Manus*.

MAIRANIA UVA URSI, *Arbutus urva ursi*.

MAÏS, *Zea mays*.

MAISON PETITE, *Morotrophium*.

MAJOR HELICIS. A narrow band of muscular fibres situate upon the anterior border of the helix of the ear, just above the tragus.

MAJORANA, *Origanum majorana*—m. Hortensis, *Origanum majorana*—m. Syriaca, *Teucrium marum*.

MAKROCEPHALUS, *Macrocephalus*.

MAL D'AMOUR, *Odontalgia*.

MAL DES ARDENS. A name given to a species of pestilential erysipelas or *Saint Anthony's fire*, which reigned epidemically in France, in 1130.

MAL D'AVENTURE, *Paronychia*—m. di Breno, *Scherlievo*—m. Caduc, *Epilepsy*—m. de Cerf, *Tetanus*.

MAL DE CRIMÉE (F.), *Lèpre des Cosaques*, *Lepra Taurica*. A variety of lepra in the Crimea.

MAL DE DENT, *Odontalgia*—m. d'Estomac, *Chthonophagia*—m. Divin, *Epilepsy*—m. d'Enfant, *Pains (Labour)*—m. di Fiume, *Scherlievo*—m. Française, *Syphilis*—m. de Gorge, *Cynanche*—m. Grand, see *Epilepsy*—m. Haut, *Epilepsy*—m. de Mâchoire, *Trismus*—m. de Mer, *Nausea marina*—m. de Mère, *Hysteria*—m. de Mière, *Pellagra*—m. de Montagne, *Puna*—m. de Naples, *Syphilis*—m. Petit, see *Epilepsy*—m. des Pieds et des mains, *Acrodynia*—m. Régionale, *Endemy*—m. de la Rosa, *Pellagra*—m. du Roi, *Serofula*—m. Rouge de Cayenne, *Elephantiasis of Cayenne*—m. Saint Antoine, *Erysipelas*—m. Saint Jean, *Epi-*

lepsy—m. Saint Main, *Lepra*, *Psora*—m. di Scherlievo, *Scherlievo*—m. de Stiam, *Fever*, *yellow*.

MAL DE SAN LAZARO, *Cocobay*. A leprous disease, common in Colombia, S. America.

MAL DEL SOLE, *Pellagra*—m. de Solagne, *Ergotism*—m. de Terre, *Epilepsy*—m. à Tête, *Cephalalgia*—m. del Valle, *Proctocæ*—m. Vat, see *Anthrax*.

MALA, Gena—m. Aurea, see *Citrus aurantium*.

MALABATH'RINUM. Ancient name of an ointment and a wine, into which the *malabathrus* entered.

MALABA'THRUM, *Cadeji-Indi*. The leaves of a tree of the East Indies. These leaves entered into the theriac, mithridate, and other ancient electuaries. They are believed to be from a species of laurel—*Laurus Cassia*; but, according to others, from *Laurus Malabathrum*. The *O'leum Malabathri* is obtained from it.

MALACCÆ RADIX, *Sagittarium alexipharmacum*.

MALACHE, *Malva rotundifolia*.

MALACIA, from *μαλακία*, 'softness.' A depravation of taste; in which an almost universal loathing is combined with an exclusive longing for some particular article of food. If the patient desires substances that are noxious or not eatable, it constitutes *Pica*, *Pisso'sia*, *Pitto'sia*, *Heterorez'ia*, *Heterorrhæz'ia*, *Cissa*, *Citto'sia*, *Citta*, *Limo'sia Pica*, *Allotriophagia*, *Picacismus*, *Pica'tio*, *Depraved appetite*, (F.) *Envie*. These symptoms accompany several nervous affections,—those of females in particular. In pregnancy it is common, and is termed *Longing*.

MALACIA AFRICANORUM, *Chthonophagia*—m. Cordis, *Cardiomalacia*.

MALACISMUS, *Mollities*.

MALACOGASTER, *Gastromalacia*.

MALACOPHONUS, from *μαλακία*, 'softness,' and *φωνη*, 'voice.' One who has a soft voice.

MALACOPEA, *Emollients*.

MALACORIUM, see *Punica granatum*.

MALACOSARCO'SIS, from *μαλακος*, 'soft,' and *σαρξ*, *sarcos*, 'flesh.' State of softness of the muscular system.

MALACOSIS, *Mollities*—m. Cerebri, *Mollities cerebri*—m. Cordis, *Cardiomalacia*—m. Hepatis, *Hepatomalacia*—m. Uteri, *Hysteromalacia*.

MALACOSTEON, *Mollities ossium*.

MALACOSTEOSIS, *Mollities ossium*.

MALACTICA, *Emollients*.

MALACTICUM, *Relaxant*.

MALADE, *Sick*.

MALADIE. Disease—m. Anglaise, *Hypochondriasis*—m. Bleue, *Cyanopathy*—m. de Bright, *Kidney*, *Bright's disease of the*—m. Bulleuse, see *Bullous*, *Pemphigus*—m. de Cruveilhier, see *Brash*, *weaning*—m. Cuculaire, *Pertussis*—m. Glandulaire, *Elephantiasis Arabica*—m. Imaginaire, *Hypochondriasis*, *Hysteria*—m. Mercenaire, *Hydrargyriasis*.

MALADIE DES MINEURS. *Anæmia* occurring in the workers in mines.

MALADIE NOIRE, *Melæna*—m. du Pays, *Nostalgia*—m. Pédiculaire, *Phthieriasis*—m. de Pott, *Vertebral disease*—m. Typhoïde, see *Typhus*—m. Vénérienne, *Syphilis*—m. de Vénus, *Syphilis*—m. de Werthof, *Purpura hæmorrhagica*.

MALADIES ACQUISES, *Acquired diseases*—m. Annuelles, *Annual diseases*—m. Chroniques, *Chronic diseases*—m. Connées, *Connate diseases*—m. Dissimulées, *Feigned diseases*—m. Externes, *External diseases*—m. Feintes, *Feigned diseases*—m. Hérititaires, *Hereditary diseases*—m. Innées, *Innate diseases*—m. Légitimes, see *Legitimate*—m. Nervieuses, *Nervous diseases*—m. Religieuses, *Religious diseases*—m. Simulées, *Feigned diseases*—

m. Sporadiques, see Sporadic — *m. Supposées*, Feigned diseases — *m. Ventuses*, Pneumatosis.

MALADIF, Sickly.

MALADRERIE, *Ladrerie*.

MALADY, ENGLISH, Hypochondriasis.

MALÆ OS, from *malum*, 'an apple,' so called from its roundness. *Zygo'ma*, *Os mala'rè* seu *Jugulè* seu *Jugamen'tum* seu *Genæ* seu *Zygomat'ium* seu *Hypo'pium* seu *Subocula'rè* seu *Pudi'um*, (F.) *Os Malaire* ou *Zygomatique* ou *de la Pommette*. The cheek or malar bone. This bone is situate at the lateral and superior part of the face; and constitutes the zygomatic region of the cheek. It is irregularly quadrilateral. Its outer surface is convex, covered by muscles and skin, and pierced with canals, called *malar*, through which vessels and nerves pass. Its upper surface is concave, and forms part of the orbit. Its posterior surface is concave, and enters into the composition of the temporal fossa. This bone is thick and cellular. It is articulated with the frontal, temporal, sphenoid, and superior maxillary bones, and is developed by a single point of ossification.

The part of the face rendered prominent by it, the French call *Pommette*.

MALAG'MA, from *μαλασσω*, 'I soften.' An emollient cataplasm, and, in general, every local application which enjoys the property of softening organic tissues.

MALAIRE, *Malair*.

MALAISE, Indisposition.

MALAKIEN, Mollities.

MALAMBO BARK, *Matias*.

MALANDERS, *Malandria*.

MALAN'DRIA. A species of lepra or elephantiasis. — *Marcellus Empiricus*. *Mal'anders* or *Mall'enders*, is a disease of the horse, which consists of a scurfy eruption on the inside of the hook, or a little below it, as well as at the bend of the knee. It is called *Mallenders* in the foreleg, and *Sallenders* in the hind leg. — *Youatt*.

MALANDRIO'SUS, *Leprous*. Affected with *malandria*.

MALAR, *Mala'ris*, (F.) *Malaire*, from *mala*, 'the cheek,' Belonging to the cheek, — as the *malar* bone.

MALAR PROCESS, Zygomatic process.

MALARE OS, *Malæ os*.

MALARIA, *Miasm*.

MALARIOUS, *Mal'arial*. Owing to, or connected with *Malaria*, — as a *malarious soil*, *malarious disease*, &c.

MALASSIMILA'TION, *Malassimila'tio*; from *mala*, 'bad,' and *assimilatio*, 'assimilation.' Imperfect or morbid assimilation or nutrition.

MALAXATE, *Moll'itè*, *Subig'erè*, *Malacissu'rè*, (F.) *Malaxer*, from *μαλασσω*, 'I soften.' To produce softening of drugs, by kneading them. The process is called *Malaxation*, *Malaxa'tio*.

MALAXATION, see *Malaxate*.

MALAXIA VENTRICULI, *Gastromalaxia*.

MALAXIS, Mollities — *m. Cordis*, *Cardioma-lacia* — *m. Hepatica*, *Adiposis hepatica*.

MALAY, see *Homo*.

MALAZISSA'TUS, *Malacissu'tus*, same etymon as *Malaxate*. One in whom the testicles have not descended. It has, also, been used synonymously with *emascula'tus* and *muliera'tus*. — *Castelli*.

MALCE, *Chûblain*.

MALE, *Mas*, *Mas'culus*. Of the sex that begets young. Not female. What belongs to the male sex: as the *male organs of generation*.

MALE, *Axilla*.

MALE ORGAN, *Penis*.

MALE DI BRENO, see *Scherlievo* — *m. Scabioso venereo*, *Scherlievo*.

MALEFICIUM, *Poisoning*.

MALFORMATION, *malforma'tio*, *malconforma'tio*; from *mal*, and *forma*, 'form.' A wrong formation; or irregularity in the structure of parts. See *Monster*.

MALHERBE, *Plumbago Europæa*.

MALIA, see *Equinia*.

MALIASMUS, *Malis*, see *Equinia*.

MALICE, *Althæa*.

MALICHORIUM, see *Punica granatum*.

MALICORIUM, see *Punica granatum*.

MALIE, *Equinia*.

MALIG'NANT, *Malig'nus*, (F.) *Malin*. A term applied to any disease whose symptoms are so aggravated as to threaten the destruction of the patient. A disease of a very serious character, although it may be mild in appearance; — *Morbus malignus*.

MALIN, *Malignant*.

MALING'ERER; from (F.) *malingre*, 'sickly.' A simulator of disease, so termed in the British military service.

MALIS, *Malias'mos*, *Cuta'neous vermina'tion*, *Helminth'on'cus*. The cuticle or skin infested with animalcules. — *Phthiri'asis*, *Parasiti'mus superfic'iei*. In Persia, this affection is produced by the Guinea worm; in South America, by the Chigre; and in Europe, occasionally by the Louse. See *Equinia*, and *Phtheiriasis*.

MALIS DRACUNCULUS, *Dracunculus* — *m. Gordii*, *Dracunculus* — *m. Pediculi*, *Phtheiriasis*.

MALLENDERS, *Malandria*.

MALLE'OLAR, *Malleola'ria*, from *malleolus*, 'the ankle.' (F.) *Malléolaire*. Belonging or relating to the ankles.

MALLEOLAR ARTERIES are two branches furnished by the anterior tibial about the instep; the one — the *internal* — passes transversely behind the tendon of the tibialis anticus, to be distributed in the vicinity of the malleolus internus; — the other — the *external* — glides behind the tendons of the *extensor communis digitorum pedis* and the *peroneus brevis*, and sends its branches to the parts which surround the outer ankle, as well as to the outer region of the tarsus.

MALLE'OLUS. Diminutive of *malleus*, 'a mallet, or hammer;' *Race'ta*, *Raste'ta*, *Rascha*, *Rasetta*, *Rase'ta*, *Sphyræ*, *Talus*, *Diab'ebos*, *Tale'olus*, *Peza*, *Ankle*, (Prov.) *Ancler*, *Ancler*, *Ancliff*, *Anclor*, *Ankley*, *Riek*, (Sc.) *Cnte*, *Coot*, *Hancloth*, *Heather-clu*, (F.) *Malléole*, *Cherville du Pied*. The two projections formed by the bones of the leg at their inferior part. The inner belongs to the tibia; the outer to the fibula. The ankles afford attachment to ligaments: and each has a sort of gutter, in which certain tendons slide. See *Malleus*.

MALLET, *Malle'olus*, (F.) *Maillet*. A kind of hammer, used with a gouge for removing or cutting bones, in certain surgical and anatomical operations.

MAL'LEUS, *Malle'olus*, *Ossic'ulum Malleolo assimila'tum*, (F.) *Marteau*. The longest and outermost of the four small bones of the ear. It is situate at the outer part of the tympanum, and is united to the *membrana tympani*. It has, 1. An ovoid head, which is articulated behind with the *incus*, and is supported by a narrow part called the neck: this has, anteriorly, a *long apophysis*, which is engaged in the glenoid fissure, and is called the *Apophysis* or *Processus Processus grac'ilis*, of *Rau*. It affords attachment to the anterior seu *externus mallei* muscle. 2. A *handle*, which forms an obtuse angle with the neck, and corresponds to the membrane of the tympanum, which it seems to draw inwards. It is furnished at its upper extremity with a process — the *processus brevis*, to which the *internus mallei* is attached. This bone is developed by a single point of ossification.

MALLEUS, Equinia—*M. farciminosus*, see Equinia—*m.* Slender Process of the, *Grêle apophyse du marteau*.

MALLOW, COMMON, *Malva*—*m.* Compound decoction of, *Decoctio malvæ compositum*—*m.* Indian, *Abutilon Avicennæ*—*m.* Yellow, *Abutilon Avicennæ*.

MALLS, *Rubeola*.

MAL'MEDY, MINERAL WATERS OF. *Mal-medey* is a town in Rhenish Prussia, between Spa and Coblenz. In its immediate vicinity are several acidulous chalybeate springs.

MALO DI SCARLIEVO, *Scherlievo*.

MALOGRANATUM, *Punica granatum*.

MALPIGHI, ACINI OF, *Corpora Malpighiana*.

MALPIG'HIA MOUREL'LA, (F.) *Mourel-liee*, *Simarouba fauz*. The bark of this shrub—a native of Cayenne—is reputed to be febrifuge, and useful in diarrhœa.

MALPIGHIAN BODIES, *Corpora Malpighiana*—*m. b.* Inflammation of the Kidney, Bright's disease of the—*m. b.* of the Spleen, see Spleen.

MALPRAX'IS, *Mala praxis*, *Malum reg'imen*. Bad management or treatment.

MALT, Sax. *mealt*, Dutch, *mout*, Teut. *malt*; from *μαλάρω*, 'I soften;' (?) *Bynē*, *Mal-tum*, *Hor'dei maltum*, *Brasium*, (F.) *Drêche*. Barley made to germinate, for the purpose of forming beer. It has been recommended in medicine, as antiscorbutic, antiscrofulous, &c.

MALT SPIRIT. A spirit distilled from malt. It is the basis of most of the spirituous cordials.

MALTA, CLIMATE OF. The climate of Malta is pretty equable, the range of temperature in the twenty-four hours being rarely more than 6°. The air is almost always dry and clear. The most disagreeable wind is the sirocco, which is the source of more or less suffering to the pulmonary invalid. The winter climate is favourable. Dr. Liddell thinks that no place which he has seen in the south of Europe can compete with Malta, for a mild, dry, bracing air in November, December, and part of January; and during the other winter and spring months, he thinks it is equal to any of them.

MALTHA, *Malthē*, from *μαλάρω*, 'I soften.' Wax, particularly soft wax.

MALTHACTICA, Emollients.

MALTHAXIS, Mollities.

MALUM, Disease, Melum—*m.* *Articulorum*, Gout—*m.* *Caducum*, Epilepsy—*m.* *Caducum*, pulmonum, Asthma—*m.* *Canum*, see *Pyrus cydonia*—*m.* *Coense*, *Averrhoa carambola*—*m.* *Cotoncum*, see *Pyrus cydonia*—*m.* *Hypochondriacum*, *Hypochondriasis*—*m.* *Hystericum*, *Hysteria*—*m.* *Insaanum*, see *Solanum Melongena*—*m.* *Ischiadicum*, *Neuralgia femoro-poplitea*—*m.* *Lazari*, *Elephantiasis*—*m.* *Lycopersicum*, *Solanum lycopersicum*.

MALUM MOR'TUUM. A species of lepra, in which the affected portions of skin seem to be struck with death.

MALUM PILARE, *Trichosis*—*m.* *Pottii*, *Vertebral disease*—*m.* *Primarium*, *Idiopathia*—*m.* *Regimen*, *Malpraxis*—*m.* *Spinosum*, *Datura stramonium*—*m.* *Terrestre*, *Atropa Mandragora*—*m.* *Venerium*, *Syphilis*.

MALUS, *Pyrus malus*—*m.* *Aurantia major*, *Citrus aurantium*—*m.* *Communis*, *Pyrus malus*—*m.* *Dasyphylla*, *Pyrus malus*.

MALUS INDICA, *Bilumbi biting-bing* of Bon-tius. *Ord.* *Rosaceæ*. The juice of this East India tree is cooling, and is drunk as a cure for fevers. The leaves, boiled and made into a cataplasm with rice, are famed in all sorts of tumours. The juice, mixed with arrack, is drunk for the cure of diarrhœa. The ripe fruit is eaten as a delicacy; and the unripe is made into a pickle for the use of the table.

MALUS LIMONIA ACIDA, see *Citrus medica*—*m.* *Medica*, see *Citrus medica*—*m.* *Sylvestris*, *Pyrus malus*.

MALVA, *Malva sylvestris seu vulgaris*, *Commor Mallo*, (F.) *Mauve sauvage*. *Ord.* *Malvaceæ*. The leaves and flowers are chiefly used in fomentations, cataplasms, and emollient enemas. Its properties are demulcent.

Malva rotundifolia, *Mal'achē*, *Mal'ochē*, has like virtues; as well as the other varieties.

MALVAISCUM, *Althæa*.

MALVERN, WATERS OF. The village of Great Malvern, (pronounced *Mal'vern*), in Worcestershire, England, has for many years been celebrated for a spring of remarkable purity, which has acquired the name of the *Holy well*. It is a carbonated water; containing carbonates of soda and iron, sulphate of soda, and chloride of sodium; and is chiefly used externally, in cutaneous affections.

MAMA-PIAN. (F.) *Maman-pian*. An ulcer of a bad aspect, which is the commencement of the pian; and which, after having destroyed the flesh, extends to the bones. It is also called the *Mother of Pians*;—*La mère des pians*.

MAMEI, *Mamoe*, *Momin* or *Toddy tree*. From incisions made in the branches of this West Indian tree, a copious discharge of pellucid liquor occurs, which is called *momin* or *Toddy wine*. It is very diuretic, and is esteemed to be a good antilithic and lithontriptic.

MAMELLE, *Mamma*.

MAMELON, Nipple.

MAMELONNÉ, Mammillated.

MAMELONS DU REIN, *Papillæ* of the kidney.

MAMILLA, *Mammula*, see *Mamma*.

MAMMA, from *mamma*, one of the earliest cries of the infant, ascribed to a desire for food. *Masthos*, *Mastus*, *Mazos*, *Thelē*, *Tithos*, *Titty*, *Rama*, *Uber*, *ουδαν*, *Nutrix*, *Gemip'ona*. The female breast, *Pap*, *Pappe*, (*Mammilla*, *Mamilla* being the male breast); (F.) *Mamelle*. A glandular organ, proper to a class of animals—the *mammalia*—and intended for the secretion of milk. The *mammæ* exist in both sexes, but they acquire a much greater size in the female; especially during pregnancy and lactation. In women, before the age of puberty, the breasts are but little developed. At this period, however, towards the central part of each breast, the skin suddenly changes colour, and assumes a rosy tint. It is of a reddish brown in women who have suckled several children. This circle has a rugous appearance, owing to the presence of sebaceous glands, and is called *Are'ola* or *Aure'ola*. These glands—*Tubercles of the Areola*, of Sir Astley Cooper—furnish an unctuous fluid for defending the nipple from the action of the saliva of the sucking infant. In the midst of the *aure'ola* is the nipple, a conoidal eminence, of a rosy tint, susceptible of erection, and at the surface of which the galactophorous ducts open. Besides the skin covering them, the breasts are, also, composed of a layer of fatty areolar tissue, more or less thick; of a large gland; excretory ducts; vessels, nerves, &c. See *Mammary*.

In the animal, the mamma is called the *Udder*, *ουδαν*, *Uber*, the *Bag*.

The breasts, plus the space between, are called the *bosom*, *sinus*, *colpos* (κολπος). At times, these terms are restricted to the space between the breasts. (F.) *Sein*.

Mamma also means a nurse.

MAMMAIRE, *Mammary*.

MAMMAL, plural *Mamma'lia*, *Mam'mifer*, *mammiferous animal*; from *mamma*, 'a breast.' An animal that suckles its young.

MAMMARY, *Mamma'rius*, (F.) *Mammaire*.

6 m *mamma*, 'the breast.' Relating to the breasts.

MAMMARY ABSCESS, *Mastodynia apostematosa*.

MAMMARY ARTERIES are three in number. They are distinguished into—1. The *Internal Mammary*, *Arteria sternalis*, *A. Sous-sternal* (Ch.), *Internal thorac'ic*. It arises from the subclavian, and descends obliquely inwards, from its origin to the cartilage of the third rib. Below the diaphragm it divides into two branches; the one *external*, the other *internal*. From its origin until its bifurcation, it gives branches to the muscles and glands of the neck, to the thymus, mediastinum, pericardium, and oesophagus. In each intercostal space it gives off *internal* and *external* musculo-cutaneous branches, and also, on each side, the *superior diaphragmatic*. Its two ultimate branches are distributed on the parietes of the abdomen, and anastomose with the external mammary, intercostal, lumbar, circumflexa ili, and epigastric arteries. 2. The *External Mammary Arteries* are two in number, and are distinguished into *superior* and *inferior*. The *superior external mammary*, *First of the thoracics* (Ch.), *Superior external thoracic*, *Superior thoracic*, is furnished by the axillary artery. It descends obliquely forwards between the pectoralis major and pectoralis minor, to which it is distributed by a considerable number of branches. The *inferior external mammary*, the *second of the thoracics* (Ch.), *Long or inferior thoracic*, arises from the axillary artery, a little below the preceding. It descends vertically over the lateral part of the thorax; curves, afterwards, inwards; becomes subcutaneous and divides into a number of branches, which surround the breast. It gives branches to the pectoralis major, serratus major anticus, the intercostal muscles, the glands of the axilla, and the integuments of the breast.

MAMMARY GLAND, *Glandula lactifera*, is the secretory organ of the milk. It is situated in the substance of the breast, to which it gives shape. The tissue of this gland results from the assemblage of lobes of different size, united intimately by a dense areolar tissue. Each of these is composed of several lobules, formed of round granulations, of a rosy white colour, and of the size of a poppy seed. The glandular grains give rise to the radicles of the excretory canals of the mamma, which are called *galactophorous* or *lactiferous*. These excretory vessels unite in ramusculi, rami, and in trunks of greater or less size; collect towards the centre of the gland; are tortuous, very extensible, and semi-transparent. All terminate in sinuses, situate near the base of the nipple, which are commonly from 15 to 18 in number. These sinuses are very short, conical, and united by areolar tissue. From their summits, a fasciculus of new ducts sets out, which occupy the centre of the nipple and open separately at its surface. The arteries of the mammary gland come from the thoracic, axillary, intercostal, and internal mammary. The veins accompany the arteries; the nerves are furnished by the intercostals, and brachial plexus; the lymphatic vessels are very numerous, and form two layers. They communicate with those of the thorax, and pass into the axillary ganglions.

MAMMARY SARCO'MA, *Mastoid sarcoma* of Abernethy, *Emphy'ma sarcoma mamma'rum*. A tumour, of the colour and texture of the mammary gland; dense and whitish; sometimes softer and brownish; often producing, on extirpation, a malignant ulcer with indurated edges. Found in various parts of the body and limbs.

MAMMARY VEINS follow the same course as the arteries, and have received the same denominations. The *internal mammary vein*, of the right

side, opens into the superior cava; that of the left, into the corresponding subclavian vein. The *external mammary veins* open into the axillary vein.

MAMM'E'A AMERICA'NA, (F.) *Abricot sauvage*. Ord. *Guttifera*. The systematic name of the tree on which the *mammee* or *mamei* fruit grows. This fruit has a delightful flavour when ripe; and is much cultivated in Jamaica, where it is generally sold in the markets as one of the best fruits of the island.

MAMM'E'TA, *Mammo'ea*, from *mamma*, 'the breast.' One who has large breasts.

MAMMELLA, Nipple.

MAMMIFER, Mammal.

MAMMIFEROUS ANIMAL, Mammal.

MAMMIFORM, Mastoid.

MAMMILLA, *Mamma* (male), Nipple.

MAMMILLÆ MEDULLARES, see *Mammillary*.

MAMMILLARIS, Mastoid.

MAM'MILLARY, *Mammilla'ris*, from *Mammilla*, 'a small breast, a nipple.' See *Mastoid*.

MAMMILLARY EM'INENCE is a name given, 1. To more or less marked prominences on the inner surface of the bones of the cranium, which correspond to the anfractuositities of the cranium. 2. To white, round, medullary tubercles, of the size of a pea, situate at the base of the brain, behind the gray substance from which the *Tige pituitaire*, of the French anatomists, arises. These *Mammillary Tubercles*, *Cor'pora albican'tia* seu *Candican'tia* seu *Mammilla'ria* seu *Pisiformia*, *Bulbi for'nicis*, *Mammilla'rie* seu *Globuli medulla'rie*, *Prominentia albican'tes*, *Processus mammilla'rie* cer'e'br'i, *Protensio'nes glandula'rie*, *Eminen'tiæ candican'tes*, *Priorum crurum for'nicis bulbi*, *Willis's Glands*, (F.) *Bulbes de la voûte à trois piliers*, *Tubercles pisiformes* (Ch.), are united to each other by a small grayish band, which corresponds with the third ventricle. They receive the anterior prolongations of the fornix.

Some ancient anatomists, taking the nervous trunks, to which Willis first gave the name of *olfactory nerves*, for simple appendages of the brain, called them, an account of their shape, *Crurunculae mammilla'rie*. Vesalius, Fallopius, Columbus, and several others, termed them *Processus mammilla'rie cer'e'br'i ad nares*. They have also been called *Trigo'na olfacto'ria*.

MAM'MILLATED, (F.) *Mamelonné*, from *mamma*, 'the female breast.' That which has mammiform projections on its surface.

MAMMILLATED LIVER, *Cirrhosis*.

MAMMITE, *Mastitis*.

MAMMITIS, *Mastitis*.

MAMMOSA, *Mammeata*.

MAM'MULA; diminutive of *Mamma*, *Mamula*. A small breast.

MAMOE, *Mamei*.

MAN, *Aner*, *Anthropos*, *Homo*.

MANAKIN, *Fantom*.

MAN-IN-THE-GROUND, *Convolvulus panduratus* — m. of the Earth, *Convolvulus panduratus*

MANCENILLIER, *Hippomane mancinella*.

JANCHE D'HIPPOCRATE, *Chausse*.

MANCHINEEL, *Hippomane mancinella*.

MANCURANA, *Origanum*.

MANDIBULA, *Maxillary bone*.

MANDIBULARIS MUSCULUS, *Masseter*.

MANDIOC, *Jatropha manihot*.

MANDO, *Glutton*.

MANDRAGORA, *Atropa mandragora* — m. *Acaulis*, *Atropa mandragora* — m. *Officinalis*, *Atropa mandragora* — m. *Vernalis*, *Atropa mandragora*.

MANDRAGORITES, from *μανδραγόρας*, the *Atropa mandragora* or mandrake. Wine in which the roots of mandrake have been infused.

MANDRAKE, *Atropa mandragora*, *Podophyllum peltatum*, *P. montanum*.

MANDRIN A VIRGULE, see *Brise-coque*.

MANDUCATIO, Mastication — *m. Difficilis*, *Bradymanesis*.

MAN'DUCATORY, *Manducatorius*; from *Manducatio*, 'mastication.' Appertaining or relating to mastication; — as,

MANDUCATORY NERVE, see *Trigemini*.

MANŒUVRE, *Manœuvre*.

MANGANESE, *Manganesium*, *Manga'nium*. A metal, which is pretty abundant in nature in an oxidized state. It is of a grayish-white colour, resembling some varieties of cast iron, is hard and brittle, and destitute of magnetic properties. Specific gravity about 8.

MANGANESE, BLACK OXIDE OF, *Mangane'sii oxydum* seu *Binox'idum*, *Tetrox'ide* of *manganese*, *Magne'sia nigra*, *Mangane'sium vitrario'rum* seu *oxyda'tum nativum* seu *nigrum* seu *ochraceum nigrum* seu *oxyda'tum nativum*, *Man'ganum oxyda'tum nativum*, *Molybde'num magne'sii*, *Oxo'des man'gani nativa*, *Peroxy'dum mangane'sii nigrum nativum*, *Superoxy'dum mangan'icum*, (*F.*) *Oxyde noir de manganese*. This oxide is not much used in medicine. It has been advised to dust the affected parts, in tinea capitis, with the powder. It is official in the Edinburgh pharmacopœia, and is used in preparing chlorine, solution of chlorine, and solution of chlorinated soda.

MANGANESE, SALTS OF, see *Manganese*, sulphate of.

MANGANESE OXYDE NOIR DE, *Manganese*, black oxide of.

MANGANESE, SULPHATE OF, *Mangane'sii Sulphas* seu *Protox'idi Sulphas*, *Sulphate of Protox'ide of Man'gane*. A rose-coloured and very soluble salt, isomorphous with sulphate of magnesia. It is prepared on a large scale for the use of the dyer, by heating, in a close vessel, peroxide of manganese and coal, and dissolving the impure protoxide, thus obtained, in sulphuric acid, with the addition of a little cholehydric acid towards the end of the process. The solution is evaporated to dryness, and again exposed to a red heat, by which the persulphate of iron is decomposed. Water then dissolves the pure sulphate of manganese, leaving the oxide of iron behind.

This salt has been recommended as a cholagogue, in doses of a drachm or two.

The salts of manganese have been recommended in chlorosis and amenorrhœa, and as substitutes for chalybeates generally. The subcarbonate and the oxide have been chiefly employed.

MANGANESE, TETROXIDE OF, *M. Black oxide of*. **MANGANESII BINOXIDUM**, *Manganese*, black oxide of — *m. Oxidum*, *Manganese*, black oxide of — *m. Peroxydum nigrum nativum*, *Manganese*, black oxide of — *m. Protoxidi sulphas*, *Manganese*, sulphate of — *m. Sulphas*, *Manganese*, sulphate of.

MANGANESIUM, *Manganese*.

MANGANESIU OCHRACEUM NIGRUM, *Manganese*, black oxide of — *m. Oxydatum nativum* seu *nigrum*, *Manganese*, black oxide of — *m. Vitroriorum*, *Manganese*, black oxide of.

MANGANI OXODES NATIVA, *Manganese*, black oxide of.

MANGANICUM SUPEROXIDUM, *Manganese*, black oxide of.

MANGANIU, *Manganese*.

MANGANUM OXYDATUM NATIVUM, *Manganese*, black oxide of.

MANGE, *Scabies ferina*.

MANGIFERA INDICA, *M. domes'tica*, from *mango*, the fruit, and *fero*, 'I bear.' The *Mango tree*, (*F.*) *Manguier*. *Ord.* Terebinthaceæ. A tree cultivated over Asia, and in South America. Mangos, when ripe, are juicy, of a good flavour,

and so fragrant as to perfume the air to a considerable distance. They are eaten, either raw or preserved with sugar. From the expressed juice a wine is prepared; and the remainder of the kernel can be reduced to an excellent flour for bread.

MANGO TREE, *Mangifera Indica*.

MANGONISATIO, Falsification.

MANGONIUM, Falsification.

MANGOSTAN, *Garcinia mangostana*.

MANGOSTANA, *Garcinia mangostana* — *m.*

Cambogia, *Garcinia cambogia* — *m. Garcinia*, *Garcinia mangostana*.

MANGOUSTAN, *Garcinia mangostana*.

MANGOSTEEN, *Garcinia mangostana*.

MANGUIER, *Mangifera Indica*.

MANHOOD, Adult age.

MANI, *Arachis hypogæa*.

MAN'IA, *Furor*, *Hemianthro'pia*, *Furor* seu *Ephro'nia* seu *Delir'ium mania*, *Insania*, *Delir'ium mani'acum* seu *furio'sum*, *Mania universalis*, *Vesania mania*, (*F.*) *Manie*, *Hyperphrénie*, *Raving* or *furious madness*, (*Sc.*) *Wodnes*; from *μωωπα*, 'I am furious.' With some, it means insanity. Disorder of the intellect, in which there is erroneous judgment or hallucination, which impels to acts of fury. If the raving be not directed to a single object, it is mania properly so called; if to one object, it constitutes *monomania*, which term is, however, usually given to melancholy. Mania attacks adults chiefly; and women more frequently than men. The prognosis is unfavourable. About one-third never recover; and they who do are apt to relapse. Separation is one of the most effective means of treatment, with attention to the corporeal condition and every thing that can add to the mental comfort of the patient, and turn his thoughts away from the subjects of his delusion. In the violence of the paroxysms, recourse must be had to the strait waistcoat, the shower bath, &c. Separation should be continued for some weeks during convalescence, with the view of preventing a relapse.

MANIA, DANCING, *Dancing plague*. A form of convulsion, which has appeared, at various times, epidemically under the form of *St. Vitus's dance*, *St. John's dance*, *Tarantism*, *Hysteria*, *Tigretier* (in Abyssinia), and diseased sympathy; and which has been fully described by Hecker in his 'Epidemics of the Middle Ages.' See *Convulsionnaire*.

MANIA EPILEPTICA, see *Epilepsy* — *m. Erotica*, *Erotomania* — *m. Lactea*, *m. puerperal* — *m. Melancholica*, *Melancholy* — *m. a Pathemata*, *Empathemata* — *m. a Potu*, *Delirium tremens*.

MANIA, PUERPERAL, *M. puerperarum acuta* seu *puerperalis* seu *lactea*, *Insania puerperarum*, *Encephalopathia puerperalis*, *Puerperal Insanity*. Mania which supervenes in the child-bed state.

MANIA PUERPERARUM ACUTA, *M. puerperal* — *m. sine Delirio*, *Pathomania* — *m. a Temulentia*, *Delirium tremens* — *m. Pellagria*, *Pellagra*.

MANIACAL, *Maniodes*.

MANIACUS, *Maniodes*.

MANICA HIPPOCRATIS, *Chaussée*.

MANICOCOMIUM, *Morotrophium*.

MANIE, *Mania* — *m. sans Délire*, *Empathema*.

MANIGUETTA, *Amomum grana paradisi*.

MANIHOT UTILISSIMA, *Jatropha manihot*.

MANIIBAR, *Jatropha manihot*.

MANIKIN, *Fantom*.

MANILUVIUM, *Manulu'vium*, from *manus*, 'the hand,' and *lavo*, 'I wash.' A bath for the hands. It may be rendered stimulating, by means of muriatic acid, mustard, &c.

MANIOC, *Jatropha manihot*.

MANIO'DES, *Mani'acus*, *Mani'acal*. One labouring under mania; *Hemianthro'pus*, *Furio'sus*, *Furibundus*, *Lyssas*, *Lysses'ter*.

MANIPULATION, from *manus*, 'a hand.' Mode of working, as in pharmacy.

MANIPULUS, (F.) *Poignée*. The quantity of a substance capable of filling the hand. A handful. See Fasciculus.

MAN-MIDWIFE, Accoucheur.

MANNA, see *Fraxinus ornus*—m. Briançon, see *Pinus larix*.

MANNA BRIGANTINA, see *Pinus larix*—m. Calabrina, see *Fraxinus ornus*—m. Croup, Semolina—m. Laricea, see *Pinus larix*—m. Metallorum, *Hydrargyri submurias*.

MANNEQUIN, Fantom.

MANNIKIN, Fantom.

MANNIN, see *Fraxinus ornus*.

MANNITE, see *Fraxinus ornus*.

MANŒUVRE (F.), pronounced and anglicized *Manœuvre*; from *main*, 'the hand,' and *œuvre*, 'work.' A dexterous movement. Applied in France to the practice of surgical or obstetrical operations on the dead body or phantom; *Opératio chirurgica seu obstetrica*.

MANSANA ARBOREA, see Jujube.

MANSFORD'S PLATES, see Galvanism.

MANSORIUS, Buccinator.

MANSTUPRATIO, Masturbation.

MANTELE, Bandage (body).

MANTI'A, *Man'tica*, *Man'ticē*, in English, *mancy*; a common suffix, from the Greek, denoting 'divination'; *Divina'tio*, *Prædivinatio*, *Præcægium*.

MANTILE, Bandage (body).

MANTLE, Panniculus carnosus.

MANU'BRIUM, from *manus*, 'a hand.' The handle of any thing:—as *manu'brium mal'lei*, *petiolus mal'lei*, 'the handle of the mallets.'

MANUBRIUM, *Chasse*—m. *Manûs*, Radius.

MANUBRIUM STERNI. The uppermost broad part of the sternum.

MANULUVIUM, Bath, hand, Maniluvium.

MANUS, *paw*, 'I feel,' *παῖπα*, 'I take,' *Choir*, *Instrumentum Instrumentorum*, *Hand*, *Paw*, (F.) *Main*. The part which terminates the upper extremity in man, and which is inservient to prehension and touch. It extends from the fold of the wrist to the extremity of the fingers. The hand is sustained by a bony skeleton, composed of a number of pieces, movable on each other; of muscles, tendons, cartilages, ligaments, vessels, nerves, &c. It is divided into three parts—the *carpus* or wrist, the *metacarpus*, and *fingers*. Its concave surface is called the *palm*; the convex surface the *back of the hand*. The facility of being able to oppose the thumb to the fingers in order to seize objects forms one of the distinctive characters of the human hand.

MANUS CHRISTI PERLA'TA. A name anciently given to troches, prepared of pearls and sugar of roses. They were called *Manus Christi simplices*, when pearls were not employed.

MANUS DEL. An ancient plaster, prepared of *rac*, *myrrh*, *frankincense*, *mastic*, *gum ammoniac*, *galbanum*, *oil*, &c. See Opium.

MANUS HEPATIS, see Liver—m. *Jecoris*, see Liver—m. *Parva majori adjutrix*, see *Digitus*.

MANSTUPRATIO, Masturbation.

MANSTUPRATOR, Masturbator.

MANYPLIES, Omasum.

MAPLE, *Acer saccharinum*—m. Ground, *Heuchera cortusa*—m. Red, *Acer rubrum*—m. Striped, *Acer pennsylvanicum*.

MARAI'S, Marsh.

MARANSIS, Atrophy.

MARANTA, Arrow-root—m. *Arundinacea*, Arrow-root.

MARANT'A GALAN'GA, *Alpinia galan'ga*, *Amomum galanga*, *Galanga*. Ord. *Zingiberaceæ*. The smaller *galangal*. Two kinds of galangal are mentioned in the pharmacopœias; the greater,

obtained from *Kæmpferia galanga*, and the smaller, from the root of *Maranta galanga*. The dried root is brought from China, in pieces from one to two inches in length, but scarcely half as thick; branched; full of knots and joints, with several circular rings, of a reddish brown colour, on the outside, and brownish within. It was formerly much used as a warm stomachic bitter, and generally ordered in bitter infusions.

MARASCHINO, see Spirit.

MARASMO'PYRA, *Febbris marasmo'des*, from *μαρasmus*, 'marasmus,' and *πυρ*, 'fever.' Fever of emaciation in general. Hectic fever.

MARASMUS, Atrophy—m. *Lactantium*, *Pædatrophia*—m. *Phthisis*, *Phthisis pulmonalis*.

MARASMUS SEN'ILIS, *Tabes senum*, *Gerontotroph'ia*. Progressive atrophy of the aged.

MARASMUS TABES, *Tabes*—m. *Tabes dorsalis*, *Tabes dorsalis*.

MARATHRI'TES, from *μαραθρον*, 'fennel.' Wine impregnated with fennel.

MARATHROPHYLLUM, *Peucedanum*.

MARATHRUM, *Anethum*—m. *Sylvestre*, *Peucedanum*.

MARAUGIA, *Metamorphopsia*.

MARBLE, Marmor.

MARBRE, Marmor.

MARÇ, Magma.

MARCASITA, Bismuth—m. *Alba*, Bismuth, subnitrate of—m. *Plumbea*, Antimonium.

MARCASTÆ MAGISTERIUM, Bismuth, subnitrate of.

MARCHAN'TIA POLYMOR'PHA, *M. stellata* seu *umbellata*, *Hepatica fontana*, *Lichen stellatus*, *Jecora'ria*, *Liverwort*, (F.) *Hépatique des fontaines*. Ord. *Hepaticæ*. This plant is mildly pungent and bitter. It is recommended as aperient, solvent, and antiscorbutic; and is used in diseased liver. &c.

MARCHE (LA), Walking.

MARCHIONIS PULVIS, *Powder of the Marquis*. A powder, formerly considered to be antiepileptic; and composed of *Male peony root*, *Mistletoe*, *Ivory shavings*, *Horn of the hoof of the stag*, *Spodium*, *Tooth of the monodon*, *coral*, &c.

MARCORES, Atrophy, Emaciation.

MARCORY, *Stillingia*.

MARÉCAGEUX, Elodes.

MARENMATIQUE, see Elodes.

MAREO, Puna.

MARE'S FAT, *Inula dysenterica*—m. *Tail*, *Hippuris vulgaris*.

MARGA CAN'DIDA, *Lac luna*. An ancient name for a variety of spongy, white, friable marl, which was employed as an astringent and refrigerant.

MARGARETIZZA, *Scherlievo*.

MARGARITA, Pearl.

MARGARITACEOUS, *Margaritaceus*, *Nacreous*, (F.) *Nacré*; from *margarita*, 'pearl, mother of pearl.' Resembling, or of the nature of, mother of pearl,—as *Leuco'ma Margaritaceum*; Pearl-like leucoma.

MARGARON, Pearl.

MARGELIS, Pearl.

MARGELLUM, Pearl.

MARGINI-SUS-SCAPULO-TROCHITÉRIEN, *Teres minor*.

MARGO, *Bord*—m. *Dentatus*, see *Retina*—m. *Orbitalis*, see *Orbit*.

MARGUERITE PETITE, *Bellis*—m. *des Prés*, *grande*, *Chrysanthemum leucanthemum*.

MARIENBAD, MINERAL WATERS OF. Celebrated springs in Bohemia. The *Kreuzbrunn* contains sulphate of soda, carbonate of iron, and carbonic acid.

MARIGOLD, DIAMOND FIG, *Mesembryanthemum crystallinum*—m. *Garden*, *Calendula officinalis*—m. *Marsh*, *Caltha palustris*—m. *Sin-*

gle, *Calendula officinalis*—m. Wild, *Calendula arvensis*.

MARIOTTE', EXPERIMENT OF, (F.) *Expérience de Mariotte*. A celebrated experiment of the Abbé Mariotte, which consists in placing two small round spots on a wall at some distance from each other, standing opposite the left-hand object, and looking at it with the right eye, the left being closed. By walking backwards, until the distance from the object is about five times as great as the distance between the two objects, the latter will be found to disappear. Mariotte and, after him, many ophthalmologists, inferred that the optic nerve, on which the ray doubtless falls in this experiment, is insensible; and hence that the choroid may be the seat of vision,—not the retina. The inference is illogical; for it doubtless falls on the part of the optic nerve where the central artery enters, and the central vein leaves the eyeball, and where there is necessarily no neurine.

MARIS, mās. Ancient name of a measure, containing 83 pints and 4 ounces.

MARISCA, Ficus, Hæmorrhoids.

MARJOLAINE, *Origanum majorana*.

MARJORAM, COMMON, *Origanum*—m. Wild. *Origanum*—m. Sweet, *Origanum majorana*.

MARJORANA, *Origanum majorana*—m. Mancurana, *Origanum*.

MARMALADE, Marmelade.

MARMARYGE, see *Metamorphopsia*.

MARMARYG'DES, μαρμαρυγῶδες, 'brilliant.'

An epithet, joined particularly to the word *οφθαλμος*, to indicate a brilliant eye, a flashing eye. An eye which transmits the image of imaginary objects. See *Metamorphopsia*.

MAR'MELADE, *Marmela'da*, *Marmela'ta*, *Marmalade*, *Miva*. Parts of vegetables, confectioned with sugar, and reduced to a pulaceous consistence.

MARMELADE OF APRICOTS. A marmelade, prepared with two parts of ripe apricots deprived of their stones, and one part of white sugar.

MARMELADE OF FERNEL, M. of Tronchin.

MARMELADE OF TRONCHIN, or OF FERNEL. A kind of thick looch, of an agreeable taste, prepared with two ounces of oil of sweet almonds, as much syrup of violets, manna in tears, very fresh pulp of cassia, 16 grains of gum tragacanth, and two drachms of orange flower water. It is used as a laxative, demulcent, and pectoral.

MARMELATA, Marmelade.

MARMOR, μαρμωρος, Marble, Calcis Car'bonas durus, (F.) *Marbre*. White granular carbonate of lime. Used in pharmacy for the preparation of a pure lime, and the disengagement of carbonic acid.

MARMORATA AURIUM, Cerumen.

MARMOROS, Marmor.

MARMORYGE, see *Metamorphopsia*—m. Hippocratis, Hemiopia.

MAROUTE, *Anthemis cotula*.

MARRIAGEABLE, Nubile.

MAR'RIOTT, DRY VOMIT OF. This once celebrated emetic, called *dry*, from its being exhibited without drink, consisted of equal portions of tartarized antimony, and sulphate of copper.

MARRONIER D'INDE, *Æsculus Hippocastanum*.

MARROW, *My'elos*, *Medu'lla*, *M. ossium*, *Medullary Juice*, *Azun'gia de Mum'ia*, *Mumia de Medullis*, *Sax. mers*, (F.) *Moëlle*, *Suc médullaire*. The oily, inflammable, whitish or yellowish juice, which fills the medullary canal of the long bones, the cancellated structure at the extremities of those bones, the diploë of flat bones, and the interior of short bones. The marrow is furnished by the exhalation of the medullary membrane. It is fluid during life, and appears under the form

of small points or brilliant grains after death. It is enveloped in the medullary membrane.

MARROW, SPINAL, *Medulla spinalis*—m. Vertebral, *Medulla spinalis*.

MARRUBE BLANC, *Marrubium*—m. *Noir*, *Ballota fetida*.

MARRUBIUM, *M. vulga'rē seu album* seu *German'icum* seu *apulum*, *Præ'sium*, *Phrasum*, *Horehound*, (F.) *Marrube blanc*. Ord. Labiatae. The leaves have a moderately strong, aromatic smell; and a very bitter, penetrating, diffusive, and durable taste. It has often been given in coughs and asthmas, united with sugar. Dose, ʒss to ʒj, in infusion; dose of extract, gr. x to ʒss.

FORD'S BALSAM OF HOREHOUND, is made as follows:—*horehound, liquorice root*, aa lbijj and ʒviij; *water*, q. s. to strain lbvj. Infuse. To the strained liquor add:—*proof spirit or brandy*, lb12; *camphor*, ʒj and ʒij; *opium and benjamin*, aa ʒj; *dried squills*, ʒij; *oil of aniseed*, ʒi; *honey*, lbijj and ʒviij.—Gray. It is pectoral.

MARRUBIUM, *Leonurus cardiaca*—m. *Album*, *Marrubium*—m. *Apulum*, *Marrubium*—m. *Germanicum*, *Marrubium*—m. *Nigrum*, *Ballota fetida*—m. *Vulgare*, *Marrubium*.

MARS, Ferrum—m. *Solubilis*, *Ferrum tartarizatum*.

MARSEILLES, (CLIMATE OF) The remarks made upon the climate of Montpellier apply even in greater force to that of Marseilles. It possesses all the objectionable qualities of the climate of southeastern France.

MARSH, Linné, Helos, Palus, Sax. merye, (F.) *Marais*. Marshy districts give off emanations, which are the fruitful source of disease and the cause of great insalubrity in many countries. The chief disease, occasioned by the malaria or miasm, is intermittent fever. Hence it becomes important to drain such regions, if practicable. Some marshy countries are not so liable to phthisis pulmonalis, and it has been found, that where intermittents have been got rid of by draining, consumption has, at times, become frequent. The most unhealthy periods for residence in a marshy district are during the existence of the summer and autumnal heats; at which times the water becomes evaporated, and the marshy bottom is more or less exposed to the sun's rays. This postulatium seems necessary for the production of the miasmata: for whilst the marsh is well covered with water, no miasm is given off.

MARSH POISON, Miasm (marsh)—m. *Mallow*, *Althæa*—m. *Root*, *Statice Caroliniana*—m. *Tea*, *Ledum palustre*.

MARSHALL'S CERATE, see *Cerate*, *Marshall's*.

MARSHY, Elodes.

MARSIPIUM, *Marsupion*.

MARSUM, *Mar'sium*, *Mar'sicum*. An ancient wine of Marsia, in Italy, which was used as an astringent in certain diseases of the mouth.

MARSUPIAL, see *Marsupion*.

MARSUPIALIS, *Ischio-trochanterianus*, *Obturator internus*.

MARSUPIATE, see *Marsupion*.

MARSUPIUM, *Marsupeion*, *Marsyp'ion*, *Mar'sip'ion*, *Mar'sip'ium*, *Mars'icium*, *Sac'culus*, (pap'ovion, 'a small sack, a purse.') A sack or bag, with which any part is fomented. Also, the abdominal pouch in the kangaroo, opossum, &c., into which the young, born at a very early stage of development, are received and nourished with milk secreted from glands which open into the pouches. Such animals are termed *Mar'sup'ial*, *Mar'sup'iate*, *Marsup'ialia*. See *Generation*.

MARSUPIUM, *Scrotum*—m. *Musculosum*, *Dartos*.

MARSYPION, *Marsupion*.

MARTEAU, Malleus.

MARTIAL, Chalybeate.

MARTIALIS, Chalybeate.

MARTIANA POMA, see Citrus aurantium.

MARTIA'TUM UNGUENTUM, (F.) *Onguent de Soldat, Soldier's ointment.* This was composed of bay berries, rue, marjoram, mint, sage, wormwood, basil, olive oil, yellow wax, and Malaga wine. It was invented by Martian; and was employed by soldiers as a preservative against injuries from cold.

MARTIS LIMATURA, Ferri limatura.

MARUM CORTUSI, Teucrium marum—m. Creticum, Teucrium marum—m. Syriacum, Teucrium marum—m. Verum, Teucrium marum—m. Vulgare, Thymus mastichina.

MARUTA COTULA, Anthemis cotula.

MAS, Male, Modiolus.

MASCARPIO, Masturbator.

MASCHALADENITIS, from *μασχαλη*, 'the axilla,' and *adenitis*, 'inflammation of a gland.' Inflammation of the glands of the axilla.

MASCHALE, Axilla.

MASCHALIBUS, Axillary.

MASCHALIATR'IA, from *μασχαλη*, 'the axilla,' and *ia-tris*, 'healing,' Treatment of disease by applications made to the axilla.

MASCHALIS, Axilla.

MASCHALISTER, Axis.

MASCHALONCUS, *Maschalop'anus*, from *μασχαλη*, 'the axilla,' and *ογκος*, 'a tumour.' A tumour or bubo or swelling in the axilla.

MASCHALOPANUS, Maschalonicus.

MASCULA, Tribade.

MASCULUS, Male.

MASESIS, Mastication.

MASHUA, (S.) A tuberous root, of a flat, pyramidal shape, which is cultivated and cooked like the potato by the Serranos of Peru. It is watery and insipid, but nevertheless is much eaten by them. The Indians use the mashua as a medicine in dropsy, dyspepsia, and dysentery. The plant is unknown to botanists.—Tschudi.

MASK, Larva, (F.) *Masque, Larve.* A bandage applied over the face, as a sort of mask, in cases of burns, scalds, or erysipelas. It serves to preserve the parts from the contact of air, and to retain topical applications *in situ*. It is made of a piece of linen, of the size of the face, in which apertures are made corresponding to the eyes, nose, and mouth, and which is fixed by means of strings stitched to the four angles.

MASKED, Larva'tus, (F.) *Larvé.* Disguised, —as a "masked fever," or one which is not clearly detected. See Fever, masked.

MASLACH, *Moslich, Am'phion, An'fion.* A medicine much used by the Turks, and into the composition of which opium enters. It is excitant.

MASQUE, Mask.

MASS, *μαζα, Maza, Massa*, from *μασσω*, 'I mix.' The compound, from which pills have to be formed.

MASSA, Mass —m. Carnea Jacobi Sylvi, see Flexor longus digitorum pedis profundus perforans—m. de Dactylis, Paste, date —m. de Extracto glycyrrhizæ, Pasta glycyrrhizæ, &c.—m. de Gummi Arabico, Paste, marshmallow —m. de Zizyphorum fructu, Paste of jujubes.

MASSAGE, Shampooing.

MASSE D'EAU, Typha latifolia.

MASSEMA, Mastication.

MASSEMENT, Shampooing.

MASSEPAINS, Biscuits, medicinal.

MASSES APOPHYSAIRES, see Vertebrae.

MASESIS, Mastication.

MASSETER, from *μασσαι*, 'I eat,' 'I chew.' *Musculus mandibula'ris*, (F.) *Zygomo-maxillaire.* A muscle situate at the posterior part of

the cheek, and lying upon the ramus of the lower jaw-bone. It is long, quadrilateral, and is attached, *above*, to the inferior edge and to the inner surface of the zygomatic arch; *below*, it terminates at the angle of the jaw, and at the outer surface and inferior margin of the ramus of that bone. It is composed of an intermixture of fleshy and aponeurotic fibres. Its office is to raise the lower jaw, and to act in mastication.

MASSETER INTERNUS, Pterygoideus internus.

MASSETER'IC, *Mas'seterine, Masse'tericus, Masseter'i'cus*, (F.) *Massétérin, Massétérique.* Relating or belonging to the masseter muscle.

MASSETERINE, Masseteric.

MASSETERINE ARTERY arises from the trunk of the internal maxillary or temporalis profunda posterior, and is distributed to the masseter muscle, after having passed, horizontally, through the sigmoid notch of the lower jaw-bone.

MASSETERINE NERVE is given off from the inferior maxillary branch of the fifth pair. It passes through the sigmoid notch, and is distributed on the inner surface of the masseter. In luxation of the lower jaw, this nerve is strongly stretched, and considerable pain, consequently, produced.

MASSETERINE VEIN has the same distribution as the artery. It opens into the internal maxillary vein.

MASSETERINUS, Masseteric.

MASETTE, Typha latifolia.

MASSICOT, Plumbi oxydum semivitreum.

MASSING, Shampooing.

MASSULA, Molecule.

MASTADENTITIS, Mastitis.

MASTALGIA, Mastodynia.

MASTAX, Mystax.

MASTEMA, Masticatory.

MASTER OF THE WOODS, Galium circæans.

MASTERWORT, Angelica atropurpurea, Heracleum lanatum, Imperatoria.

MASTESIS, Mastication.

MASTHELICOSIS, Mastodynia apostematosa.

MASTHOS, Mamma.

MASTICA'TION, *Mastica'tio, Mase'sis, Mase'sis Masse'ma, Maste'sis, Manduca'tio, Com-manduca'tio, Manduca'tion, Chewing, Chawing*, from *μαστιγναι*, 'I gnash the teeth.' The action of bruising food in the mouth, to prepare it for the digestion it has to undergo in the stomach. This is executed by the joint action of the tongue, cheeks, and lips, which push the alimentary substance between the teeth; and by the motions of the lower jaw it is cut, torn, or bruised.

MASTICATORY, *Masticato'rium, Maste'ma, Diamaste'ma*, (F.) *Masticatoire.* Same etymon. Chewing. Relating or appertaining to mastication or chewing. Also, a substance, chewed with the intention of exciting the secretion of saliva.

MASTICATORY NERVE, see Trigemini.

MASTICH, see Pistacia lentiscus—m. Herb, common, Thymus mastichina—m. Herb, Syrian, Teucrium marum—m. Tree, Pistacia lentiscus.

MASTICHELEON, see Pistacia lentiscus.

MASTICHINA GALLORUM, Thymus mastichina.

MASTIGODES HOMINIS, Trichocephalus.

MASTIGO'SIS, *Flagella'tio.* Flagellation, scourging; employed by the ancients as a remedy in many diseases.

MASTI'TIS, *Mastoi'tis, Mastoi'd'itis, Mammi'tis, Mazoi'tis, Inflammat'io Mammae, Mastadeni'tis*, (F.) *Inflammation des Mamelles, Mammite*, from *μαστος*, 'the breast,' and *itis*, denoting inflammation. Inflammation of the breast. Inflammation of the mammary gland of the pregnant or parturient female is vulgarly called a *weid* or *weed*, and a *weed* in the breast.

MASTITIS APOSTEMATOSA, Mastodynia apostematosa.

MASTITIS PUERPERA'LIS. Inflammation of the mamma in the childbed woman.

MASTIX, see Pistacia lentiscus.

MASTO-CARCINO'MA, from *μαστος*, 'the breast,' and *carcinoma*. Carcinoma or cancer of the breast.

MASTO-CHONDRO'SIS, from *μαστος*, 'the breast,' and *χόνδρος*, 'a cartilage.' A cartilaginous formation in the breast.

MASTODES, Mastoid.

MASTODYN'IA, *Mastal'gia*, *Mazodyn'ia*, from *μαστος*, 'the breast,' and *ὀδυνή*, 'pain.' Pain in the breasts; a form of neuralgia. See Neuralgia mammae.

MASTODYN'IA APOSTEMATO'SA, *Phleg'monē Mammæ*, *Masthelco'sis*, *Mastit'is apostemato'sa*, *Galactapostema*, *Galactomastoparecto'ma*, *Abscessus Mammæ seu lacteus*, *Phleg'monē Mastodyn'ia*, *Abscess of the Breast*, *Mam'mary Abscess*, *Milk-abscess*. Phlegmonous inflammation of the breasts, running on to suppuration, generally in the childbed female. It is one of the best examples of acute phlegmonous inflammation, and requires the active use of appropriate treatment.

MASTODYNIA POLYGALA, Sparganosis—*m. Polygalæ*, *Mastospargosis*.

MASTOID, *Mastoi'des*, *Masto'des*, *Mastoidæus*, *Mammilla'ris*, *Papil'li seu mammilli-formis*, *Mammiform'is*, *Mam'miform*, (F.) *Mastoidien*, from *μαστος*, 'a breast,' and *εἶδος*, 'form, resemblance.' Having the form of a nipple. Also, that which relates to the mastoid process, *Mastoidæus*.

MASTOID APERTURE—the opening of communication between the cavity of the tympanum and the mastoid cells.

MASTOID CANCER. A kind of firm carcinomatous growth, the section of which is thought to resemble the boiled udder of the cow.

MASTOID CELLS, *Cell'ula Mastoidæa*, *Antrum mastoidæum*, *Mastoid Si'nuses*. These are situated in the mastoid process; communicate with each other, and open into the cavity of the tympanum. Their use seems to be to increase the intensity of sound.

MASTOID FORAMEN is situated behind the mastoid process, and gives passage to a small artery of the dura mater, as well as to a vein which opens into the lateral sinus.

MASTOID FOSSA, (F.) *Gouttière mastoïdienne*, is a depression at the inner surface of the mastoid portion of the temporal bone, which forms part of the lateral sinus.

MASTOID OR DIGAS'TRIC GROOVE, (F.) *Rainure mastoïdienne ou digastrique*, is a groove, situated at the inner side of the mastoid process, which affords attachment to the posterior belly of the digastric muscle.

MASTOID MUSCLE, POSTERIOR, *Splenius*.

MASTOID OR MAM'MIFORM OR MAM'MILLARY PROCESS, *Processus mastoidæus*, *Apoph'ysis mammilla'ris*, *Pars mastoidæa*, is situated at the inferior and posterior part of the temporal bone, and gives attachment to the digastric and mastoid muscles.

MASTOIDEUS, *Sterno-cleido-mastoidæus*—*m. Lateralis*, *Complexus minor*.

MASTOIDITIS, Mastitis.

MASTOÏDO-CONCHINIEN, *Retrahens auris*—*m. Génien*, *Digastricus*—*m. Hyogénien*, *Digastrius*—*m. Oriculaire*, *Retrahens auris*.

MASTOME'NIA, from *μαστος*, 'the breast,' and *μήν*, 'a month.' Vicarious menstruation from the mamma.

MASTON'CUS, *Thelon'cus*, from *μαστος*, 'the breast,' and *ογκος*, 'a tumour.' A tumefaction of the nipple, or of the breast itself. When the

tumour is of an adipous character it is termed *Mastopion'cus* or *Mastopimelon'cus*.

MASTONCUS POLYGALACTICUS, *Mastospargosis*. **MASTOPAREC**'TAMA, from *μαστος*, 'the breast,' and *παρεκταμα*, 'excessive distension.' Immoderate distension of the mammae.

MASTOPATHI'A, from *μαστος*, 'the breast,' and *πάθος*, 'disease.' An affection of the breast.

MASTOPIMELONCUS, see Mastoncus.

MASTORRHAG'IA, from *μαστος*, 'the breast,' and *ρρρρρρ*, 'to flow.' An unusual flow of milk.

MASTOS, Mamma.

MASTOSCIR'RHUS, from *μαστος*, 'the breast,' and *σκιρπος*, 'a hard tumour.' Scirrhus, or hard cancer of the breast.

MASTOSPARGO'SIS, *Mastodyn'ia polyg'ala*, *Maston'cus polygalacticus*, from *μαστος*, 'the breast,' and *σπαργαω*, 'I am full to bursting.' Fulness of the breasts with milk, so that they are ready to burst.

MASTOSYRINX, *Fis'tula mammæ*, from *μαστος*, 'the breast,' and *συριγξ*, 'a pipe.' Fistula of the mamma.

MASTURBA'TION, *Cinæ'dia*, *Gar'galé*, *Gar'galus*, *Gargali'smus*, *Mastupra'tio*, *Manustupra'tio*, *Manustupra'tio*, from *manus*, 'the hand,' and *ετρω*, 'I ravish,' *Ona'nia*, *On'anism*, *Ædæo-gargari'smus*, *Ædæo-gar'galus*, *Anaplas'mus*, *Gau'dia fœda*, *Duct'io præputii*, *Vol'untary Pollu'tion*, *Self Pollution*, *Self-abuse*, (F.) *Abus de soi-même*, *Attonchement*. Excitement of the genital organs by the hand.

MASTURBA'TOR, *Mastupra'tor*, *Manustupra'tor*, *Mascar'pio*: same etymon. One given to masturbation.

MASTUS, Mamma.

MAT, (F.) Dull.

MAT SON. A dull or flat sound. The obscure noise, (F.) *Matité*, afforded in certain diseases when any part, as the chest, is percussed. It has been compared to that produced when the thigh is struck.' It is opposed to the *Son clair*, or 'clear sound.'

MATE, see *Ilex Paraguensis*.

MATECÓ, Matico.

MATELLA, Urinal.

MATER, Uterus—*m. Dura*, *Dura mater*—*m. Herbarum*, *Artemisia vulgaris*—*m. Metallorum*, *Hydrargyrum*—*m. Mollis*, *Pia mater*—*m. Peralarum*, see Pearl—*m. Pia*, *Pia mater*—*m. Scalis*, *Ergot*—*m. Tenuis*, *Pia mater*.

MATERIA, Matter—*m. Fibrosa*, *Fibrin*.

MATE'RIA MED'ICA, *Pharmacolog'ia*, *Pharmacology*, *Acol'ogy* or *Akol'ogy* or *Aceolog'ia*, *Hylé Iat'ricé*, *Iamatolog'ia*, *Iamatology*, (F.) *Matière Médicale*. The division of medical science which treats of the knowledge of medicines; their action on the animal economy, and mode of administration. The study of the *Materia Medica* is one of great importance;—it is a study of the tools with which the practitioner has to work in the cure of disease. Much labour has been spent in contriving classifications of the *Materia Medica*. Some have arranged the articles according to their natural resemblances: others according to their real or presumed virtues: others, according to their active constituent principles. The Pharmacopœias place them alphabetically. Perhaps the best classification would be one founded on the agency exerted by the articles on the different tissues; but this arrangement, in the present state of science, is by no means easy; and, moreover, ideas in regard to the action of medicines are so associated with certain terms,—as narcotics, tonics, sedatives, &c., employed to denote certain operations, which they are esteemed capable of producing, that, to abandon them, would be to throw obstacles in the way of the student, with-

out the ultimate advantage accruing to him of possessing a better knowledge of the *modus operandi* of medicines than when a classification, somewhat resembling those usually embraced, is adopted.

The following is the classification adopted by the Author, in his "General Therapeutics and Materia Medica, 6th edit., Philad., 1857:—

- | | |
|---|---|
| 1. Agents that affect prominently the alimentary canal or its contents, | { Emetics,
Cathartics,
Anthelmintics. |
| 2. Agents that affect prominently the respiratory organs, | { Expectorants. |
| 3. Agents that affect prominently the follicular or glandular organs, | { Errhines,
Sialogogues,
Diuretics,
Antilithics,
Diaphoretics. |
| 4. Agents that affect prominently the nervous system, | { Narcotics,
Tetanics,
Antispasmodics. |
| 5. Agents that affect prominently the organs of reproduction, | { Emmenagogues,
Parturificants, |
| 6. Agents that affect various organs, | { Excitants,
Tonics,
Astringents,
Sedatives,
Refrigerants,
Revelents,
Eutrophics. |
| 7. Agents whose action is prominently chemical, | { Antacids,
Antalkalics,
Disinfectants. |
| 8. Agents whose action is prominently mechanical, | { Demulcents,
Dilutents. |

Of old, the *Materia Medica* consisted of more articles than at present. The tendency, indeed, is, and must be, to diminish it still further; to get rid of those articles which possess no advantages over others equally common, or whose properties are doubtful. In a dictionary, it becomes necessary to insert all that have been reputed to possess virtues; but the majority are unnecessary. The catalogue might be largely reduced with impunity.

MATERIA MEDICA, DYNAMICAL, Pharmacodynamias.

MATERIA MORBO'SA, *M. Peccans, Materies Morbi*, Morbid matter. The matter or material which is the cause of disease.

MATERIA OSSEA, Terra Ossea—*m. Peccans, M. Morbosa*—*m. Sialina*, see *Saliva*—*m. Scytophica*, Tannin—*m. Testacea dentium*, see *Tooth*—*m. Urinosa, Urea*.

MATERIALISTA, Druggist.

MATÉRIAUX IMMÉDIATS, Principles, immediate.

MATERIES, Matter—*m. Morbi, Materia morbosa*.

MATHEW'S PILLS, *Pilulæ ex helleboro et myrrha*.

MATHIEU'S VERMIFUGE, see *Vermifuge*.

MATIAS. The bark of a South American tree not yet determined. It is used in its native country in intermittents, and as a tonic generally. Its principal characteristic constituent is a bitter resinous matter. It is probably the same as the *Malambo bark*.

MATY'CO, (pronounced *matee'co*), *Motecó, Matica, Yerba del Soldado, Soldier's weed* or herb. A South American herb—*Piper angustifolium* seu *longatum, Stephens'ia elongata, Artanthe elongata, Ord.* Piperaceæ—which is possessed of astringent virtues, and is used both internally and externally. It is given in *infusion* made of one ounce of the leaves to a pint of boiling water,

of which the dose is f $\frac{3}{4}$ ss, or in *tincture*, made of $\frac{3}{4}$ ss of the leaves to a pint of dilute alcohol.

MATÈRE, Matter—*m. Caséuse, Casein*—*m. Cérébriforme, Encéphaloid*—*m. Extractive du Bouillon, Osmazome*—*m. Fibreuse du Sang, Fibrin*—*m. Médicale, Materia Medica*—*m. de la Mélanose*, see *Choroidea tunica*—*m. Noire des Pouxons, Anthracosis*—*m. Perte de Kerkring, Antimonium diaphoreticum*—*m. Pulmonaire noire*, see *Pulmo*.

MATITÉ, see *Mat son*.

MATLOCK, MINERAL WATERS OF. Matlock is a village in Derbyshire, England, at which there is a mineral spring of the acidulous class. Temperature 66°. It differs but little, except in temperature, from good spring-water. It is generally used as a tepid bath.

MATONIA CARDAMOMUM, Amomum cardamomum.

MATORIUM, Ammoniac, gum.

MATRACIUM, Matrass, Urinal.

MATRASS, *Matra-cium*. A glass vessel with a long neck; and a round, and sometimes oval, body. It may be furnished with tubulures, or not. It is used in *Pharmacy* for distillation, digestion, &c.

MATRES CEREBRI, Meninges.

MATRICIAIRE, Matricaria.

MATRICA'LIS, *Matrica'rius*, from *matrix*, *matricis*, 'the uterus.' Relating to the uterus.

Matricaria are remedies for diseases of the uterus.

MATRICA'RIA, from *matrix, matricis*, 'the womb;' so called from its reputed virtues in affections of that organ. *M. Parthe'nium, Parthe'nium febrif'ugum, Py'rethrum* seu *Chrysanth'enum parthe'nium, Aryg'rocha'ta, Chrysoc'alis, Fever-few, Feather-few, Mother-wort, Coothwort, (Sc.) Fever-fouillie, (F.) Matricaire. Nat. Ord. Compositæ*. The flowers resemble, in their properties, chamomile and tansy; and, like them, have been esteemed tonic, stomachic, resolvent, emmenagogue, vermifuge, &c. They are not much used.

MATRICARIA, see *Matricaria chamomilla*.

MATRICA'RIA CHAMOMIL'LA, *M. Suaveolens, An'thenis vulga'ris, Chamæ'melum Vulga're, Chamomil'la nostras, Leucan'themum* of Dioscorides, *Wild Corn, Dog's Cham'omile, German Chamomile, (F.) Camomille vulgaire*. The flowers, *Matrica'ria* (Ph. U. S.), resemble *Matricaria* in properties.

MATRICA'RIA GLABRA'TA. A South African plant, known at the Cape as *Wild Chamomile*, has the same properties as the other species of *matricaria*.

MATRICARIA LEUCANTHEMUM, Chrysanthemum Leucanthemum—*m. Suaveolens, M. Chamomilla*.

MATRICE, Uterus—*m. Col de la, Collum uteri*—*m. de l'Ongle*, see *Nail*.

MATRICULATE, from *matricula*, diminutive of *matrix, matricis*, 'a roll,' originally 'an army roll or register.' One who is admitted into a university or college, by enrolling or having his name enrolled on the register of the institution. In France, *prendre inscription* means 'to matriculate,' and, in the university regulations of that country, it is required, that the *inscription* shall be made every three months, until the termination of the prescribed period of study: the student having to take his inscription within the first fortnight of each *trimestre* or of every three months, and to present himself within the last fortnight of the *trimestre* to establish the fact of his attendance.

In this country, it is only necessary to matriculate at the commencement of each session.

One who has thus enrolled himself in an institution is called a *Matriculate*.

MATRISYLVA, *Asperula odorata*.

MATRIX, *Cytoblastema*, Uterus—*m.* Unguis, see Nail.

MATRONA, Midwife.

MATRONALIS, Viola.

MATTEE, see *Ilex Paraguensis*.

MATTER, *Mat'ria*, *Hyle*, *Mat'ries*, (F.) *Matière*. Any substance which enters into the composition of a body. In *Medicine*, it is sometimes applied to the substance of evacuations; and is also used synonymously with pus.

MATTING OF PARTS, *Hapantismus*.

MATULA, Urinal.

MATURATION, *Matura'tio*, from *maturare*, *matura'tum*, 'to ripen.' *Pepas'mos*, *Pepan'sis*. Progression of an abscess towards maturity. The state of maturity. Coction.

MATURATIF, *Maturative*.

MATURATIVE, *Matu'rans*, *Pepanticos*, (F.) *Maturatif*. Same etymon. A medicine which favours the maturation of an inflammatory tumour.

MATURATUS, Concocted.

MATURITY, *Matu'ritas*, *Horæ'otes*. Same etymon. The state of fruits and seeds, when comparatively developed. State of an abscess, in which the pus is completely formed.

MATURITY, *Preecociu's*, *Preecocity*.

MAUDLIN, *Achillea ageratum*—*m.* Tansey, *Achillea ageratum*—*m.* Wort, *Chrysanthemum leucanthemum*.

MAULES, *Rubeola*.

MAUVE SAUVAGE, *Malva*.

MAUX DE NERFS, *Hysteria*.

MAW, *Stomach*—*m.* Bound, *Constipated*—*m.* Worm, *Ascaris*.

MAXILLA, *Maxillary bone*.

MAXILLARY, *Maxilla'ris*, (F.) *Maxillaire*, from *maxilla*, 'a jaw.' Relating or belonging to the jaws; from *massaw*, 'I chew.'

MAXILLARY ARTERIES are three in number.

1. *External maxillary*. See *Facial*. 2. *Internal maxillary*—*A. Gutturæ-maxillaire* (Ch.). This arises from the external carotid with the temporal. It is remarkable for its complex course, and for the number of branches which it transmits to the deep-seated parts of the face. Immediately after its origin, it buries itself under the neck of the lower jaw, curving inwards and downwards. It then advances directly inwards; proceeding in the space between the two pterygoid muscles towards the *maxillary tuberosity*. It turns again, becomes vertical, and ascends into the bottom of the zygomatic fossa, until, having arrived at the floor of the orbit, it takes a horizontal and transverse direction; enters the sphenomaxillary fossa, and divides into several branches. See *Artery*, (table.)

MAXILLARY BONE, *Maxil'la*, *Mandib'ula*, *Gamphela*, *Gnathus*, *Mola*, *Singon*, *Jawbone*, (Prov.) *Chaffbone*, *Chall*, *Chaule*, *Chavel*, *Chawbone*, *Gill*, (F.) *Mâchoire*. A name given to two bones, which support the teeth, and, by means of them, are inservient to the cutting, bruising, and tearing of alimentary substances.

MAXILLARY BONE, INFERIOR, *Lower jaw-bone*, *Os Maxilla'rè inferius*, *Maxill'a inferior*, *Hypo'gon*, (F.) *Os maxillaire* (Ch.), *Mâchoire d'cranienne*, is a symmetrical, nearly parabolic bone, the middle portion of which is horizontal, and called the *body*; and the posterior is vertical, the angular portions being termed *Rami* or *branches*. These have behind a *parotidian edge*, which forms—by uniting with the base—the *angle of the jaw*. The branches terminate above by two processes, separated by the *sigmoid notch* or *fossa*; the anterior of which is called the *coronoid*; the posterior, the *condyloid process* or *maxillary condyle*, supported by a *Cervix*, *Collum*,

or *Neck*. The chief parts observed on the lower jaw are—1. *Externally*,—the *Symphysis menti*, *Apophysis menti*, *Mental foramen*, and the *external oblique line*. 2. *Internally*,—the *Genian apophysis*, the *internal* or *Myloid oblique lines*, and the *entrance of the dental canal*. The lower jaw has, also, an *alveolar edge*, which contains alveoli for the reception of the teeth. The lower jaw-bone is developed by two points of ossification, which unite at the *symphysis of the chin*. It is articulated with the temporal bone and with the teeth.

MAXILLARY BONE, SUPERIOR, *Upper jaw-bone*, *Maxil'la superior* seu *syncrania*, *Os maxilla'rè superius*, (F.) *Os sus-maxillaire* (Ch.), *Mâchoire syncranienne*. The upper jaw-bones are to the face what the sphenoid bone is to the cranium. They are articulated with all the portions composing it. They determine, almost alone, the shape of the face, and give it solidity. Their size is considerable; form unequal. They occupy the middle and anterior part of the face; and enter into the composition of the nasal fossæ, orbit, and mouth. The chief parts observable in the upper jaw are—1. *Externally*,—the *Nasal process*, *Processus nasalis* seu *frontalis* seu *ascendens*, (F.) *Apophyse montante*, the *superior* or *orbital surface*, *Superficie'ies orbitalis*, *Planum orbitale*, which forms almost the entire floor of the orbit, the *Foramen infra-orbitarium*, *Zygomatic process*, *Canine* or *Myrtiform fossa*. 2. *Internally*,—the *Palatine process*, *Anterior palatine canal*, and the *Antrum of Highmore*. Its circumference is very unequal, and it has behind a round, unequal prominence, called the *Maxillary tuberosity*, which is pierced by the *posterior dental canal*. Anteriorly, there is a notch, which forms part of the anterior aperture of the nasal fossæ, and beneath, an eminence, called the *anterior nasal spine*. The lower part of this circumference forms the *alveolar margin*. Each superior maxillary bone is articulated with the ethmoid, frontal, nasal, lacrymal, palatine, inferior spongy, vomer, its fellow, the teeth of the upper jaw, and sometimes the sphenoid bone. It is developed by four or five points of ossification.

MAXILLARY NERVES are two in number, and formed of the second and third branches of the fifth pair. The SUPERIOR MAXILLARY NERVE, *Nerv sus-maxillaire* (Ch.), arises from the middle of the gangliform enlargement of the fifth pair; passes forwards, and issues from the cranium through the foramen rotundum of the sphenoid bone; enters the sphenomaxillary fossa, which it crosses horizontally; passes into the infra-orbital canal, which it traverses; and makes its exit to vanish on the cheek. It gives off the following branches,—the *orbital*, a branch which goes to the *spheno-palatine ganglion*; *posterior dental branches*; the *anterior dental*,—and terminates in the *infra-orbital* nerves, which are divided into *superior*, *inferior*, and *internal*. The *inferior maxillary nerve*, *Nerv maxillaire* (Ch.), is the largest of the three branches furnished by the fifth pair. It issues from the cranium by the foramen ovale of the sphenoid. Having reached the zygomatic fossa, it divides into two trunks; the one *superior* and *external*, which gives off the *temporales profundi*, *masseterine*, *buccal*, and *ptyergoids*;—the other—*inferior* and *internal*—the larger of the two, which furnishes the *inferior dental*, *lingual*, and *auricular*.

MAXILLARY VEINS present the same arrangement as the arteries they accompany.

MAXILLO-ALVÉOLI-NASAL. *Depressor alæ nasi*—*m.* *Labial*, *Depressor anguli oris*—*m.* *Narinal*, *Compressor naris*—*m.* *Palpébral*, *Orbicularis palpebrarum*—*m.* *Scleroticæ*, *Obliquus inferior oculi*.

MAY APPLE, *Podophyllum peltatum*—m. a. Mountain, *Podophyllum montanum*—m. Flower, *Anthemis cotula*—m. Weed, *Anthemis cotula*.

MAYELLA, *Curcuma longa*.

MAYNARD'S ADHESIVE LIQUID, Colloidion.

MAYS, *Zea mays*—m. Americana, *Zea mays*—m. *Zea*, *Zea mays*.

MAYWORM, TRUE, *Meloe majalis*.

MAYZ, *Zea mays*.

MAZA, Mass, Placenta.

MAZALYSIS, Mazolysis.

MA'ZICUS, from *maza*, 'placenta.' Relating to the placenta.

MAZISCH'ESIS, from *maza*, 'the placenta,' and *σχέσις*, 'holding,' 'retention.' Retention of the placenta.

MAZODYNIA, Mastodynia.

MAZOITIS, Mastitis.

MAZOLYSIS, *Mazal'ysis*, from *maza*, 'placenta,' and *λυσίς*, 'solution.' Separation of the placenta.

MAZOPATHIA, from *maza*, 'placenta,' and *πάθος*, 'disease.' A disease of the placenta. One originating from the placenta.

MAZOS, Mamma.

MEAD, Hydromeli, Melizomum.

MEADOW BLOOM, *Ranunculus acris*—m. Fern, *Comptonia asplenifolia*—m. Pride, see *Calumbā*—m. Rue, *Thalictrum*—m. Saffron, Vinegar of, *Aacetum colchici*—m. Sweet, *Gillenia trifoliata*, *Spiraea ulmaria*—m. Sweet, American, *Spiraea lobata*—m. Sweet, red, *Spiraea tomentosa*.


MEALY TREE, *Viburnum dentatum*.

MEASLES, Rubella—m. Black, see Rubella—m. False, Roseola—m. French, Roseola.

MEASLINGS, Rubella.

MEASLY, Morbillous.

MEASURE, GRADUATED. A glass measure, used in pharmacy, which is so divided as to indicate accurately the quantities in dispensing liquid medicines.



Four-ounce Glass Measure.

MEASUREMENT, Mensuration.

MEASURES, see Weights and Measures.

MEASURING, MEDICAL, Mensuration.

MEAT BISCUIT, see Biscuit, meat.

MEAT, EXTRACTIVE OF, Osmazome.

MÉAT, Meatus.

MEA'TUS, *Ductus*, *Canalis*, *Porus*, (F.) *Méat*, from *meo*, *meatum*, 'to go.' A passage or canal.

MEATUS AUDITORIUS EXTERNUS, Auditory canal, external—m. Auditorius internus, Auditory canal, internal—m. Cæcus, Eustachian tube—m. Cuticulae, Pori cutanei—m. Cysticus, Cystic duct—m. Narium, see Nasal fossæ—m. Seminales nteri, Tubæ Fallopiæ—m. Seminarius, Corpus Highmori—m. Urinarius, Urethra.

MECHAMECK, *Convolvulus panduratus*—m. Bindweed, *Convolvulus panduratus*.

MECHANE, Machine.

MECHANEMA, Machine.

MECHANICAL, *Mechan'icus*, from *μηχανή*, 'a machine.' An epithet given to irritating bodies, which do not act chymically—as a mechanical irritant.

MECHANICAL or IATRO-MATHEMATICAL PHYSICIANS, *lū'tro-mechan'ici* seu *lū'tromathemat'ici*, are such as refer every function, healthy or morbid, to mechanical or mathematical principles;—*Medic'ina mechan'ica*.

MECHANICS, ANIMAL. That part of physiology whose objects are to investigate the laws of equilibrium and motion of the animal body.

MECHANISM, *Mechanism'us*. The structure of a body; the collection or aggregate of the parts of a machine; the mode in which forces produce any effect, &c.

MÊCHE, Tent. This term is usually applied in French surgery to a collection of threads of charpie, cotton or raw silk, united together, which are used for detaching sinuous or fistulous ulcers; or to keep open or enlarge natural or artificial apertures. They are generally applied by means of an instrument called *Porte-mêche*.

MECHOACANA NIGRA, *Convolvulus jalapa*—m. Nigricans, *Convolvulus jalapa*.

MECHOACANÆ RADIX, *Convolvulus mechoacan*.

MÊCHOACAN DU CANADA, *Phytolacca decandra*.

MECOMETER, (F.) *Mécomètre*; from *μεκος*, 'length,' and *μετρον*, 'measure.' A kind of graduated compass,—*compas de proportion*,—used at the Hospice de Maternité of Paris, to measure the length of new-born infants.

MECON, Meconium, Opium, Papaver.

MECONICUM, Opiate.

MECONIS'MUS, *Intoxica'tio opia'ca*; from *μηκων*, 'the poppy.' Poisoning by opium.

MECO'NIUM, same etymon. Poppy juice, *Papaver'culum*, *Purgamen'ta Infantis*, *Mecon*. The excrement passed by the infant a short time after birth, which had accumulated in the intestines during pregnancy. It is of a greenish or deep black colour, and very viscid. It seems formed of the mucous secretions of the intestines mixed with bile.

MECONOLOG'IA, *Opiolog'ia*; from *μηκων*, 'a poppy,' and *λογος*, 'a description.' A treatise on opium.

MECONOPSIS DIPHYLLA, *Stylophorum diphyllum*.

MEDICINARE, Doctor.

MEDEA, Genital organs.

MÉDECIN, Physician—m. *Consultant*, Consulting physician—m. *de Fous*, see *Psychiatria*.

MÉDECINE, *Medicina*—m. *Expectante*, Expectation—m. *Légale*, Medicine, legal—m. *Militaire*, Military medicine—m. *Opératoire*, Surgery, operative—m. *Perturbatrice*, Perturbatrix (*Medicina*).

MEDELA, Curation.

MEDENL A term used by Paracelsus for cutaneous ulcerations like those of boils.

MEDEOLA VERTICILLIFOLIA, *M. Virginica*.

MEDÉOLA VIRGINICA, *M. verticillifolia*, *Gyro'mia Virginica*, *Indian cucumber*. Order, Smilacæ. An indigenous herb, growing in every part of the United States, the root of which resembles a small cucumber. It has been thought to be diuretic.

MEDIAN, *Medi'anus*; from *medium*, 'the middle.' That which is situate in the middle.

MEDIAN LINE. A vertical line, supposed to divide a body longitudinally into two equal parts; the one right, the other left. Chaussier calls the *linea alba* the *ligne médiane* of the abdomen.

MEDIAN NERVE, *Median digital*, (Ch.) This nerve arises chiefly from the anterior branches of

the last two cervical nerves and first dorsal. The fifth and sixth cervical pairs also send it a branch, which separates from the musculo-cutaneous nerve. The median nerve descends the inner part of the arm along the biceps muscle. Opposite the elbow joint it buries itself behind the aponeurosis of that muscle, and engages itself between the two fasciuli of the pronator teres. Lower down, it is situate between the flexors—sublimis and profundus—and passes, with their tendons, under the anterior annular ligament; it then divides into five branches, which are distributed to the muscles of the thenar eminence, to the lumbricales, the integuments of the thumb, the index, middle finger, and outer part of the ring finger. This nerve gives no branches to the arm. In the forearm, it furnishes filaments to all the pronator and flexor muscles; and one of them accompanies the anterior interosseous artery. It also gives off a filament to the integuments of the palm of the hand.

MEDIAN VEINS. Three of the superficial veins of the forearm are so called. The *median basilic*—the *median cephalic* and *common median* or *funis brachii*. See *Basilic* and *Cephalic*.

MEDIANUM, Mediastinum, Mesentery.

MEDIAS'TINAL, Mediastin'alis, (F.) Médiastin. Relating to the Mediastinum.

MEDIAS'TINAL ARTERIES are very delicate arterial branches, distributed in the areolar texture of the mediastinum. They are distinguished, according to their situation, into anterior and posterior.

MEDIASTINITIS, Mesodmitis.

MEDIASTINUM, Mediastin'us, quasi, in medio stans, as being in the middle; *Mesod'mē, Medi'num, Mesotē'chium, Mesotē'chum, Septum thorac'is, Hymen Diaphrag'aton, Membra'na Diaphrag'aton seu thorac'em intersep'iens, Intersepim'entum thorac'is, Diribito'rium, Dissipim'entum thorac'is, (F.) Médiastin.* A membranous space formed by the approximation of the pleuræ, dividing the chest into two parts, the one right, the other left. The mediastinum, formed by a double reflection of the pleura, extends from the spine to the posterior surface of the sternum. Its anterior part, called *Anterior mediastinum, Mediastinum pectoral'e*, lodges, at its upper part, the thymus gland in the fœtus, and is filled below with fatty, areolar tissue. Its posterior part, parallel to the spine, is occupied by the œsophagus, vena azygos, thoracic duct, the lower part of the windpipe, the origin of the bronchia and a number of lymphatic glands. This part is called the *posterior mediastinum—Mediastinum dors'al'e*.

By some anatomists, Mediastinum is defined to be the pleural walls of the membranous space, whilst the space itself is termed *Cavum mediastini*.

MEDIASTINUM AURIS, see *Tympanum*—*m. Cerebri, Falx cerebri, Septum lucidum*—*m. Testis, Corpus Highmori*.

MEDIASTINUS, Mediastinum.

MEDICABILIS, Curable.

MEDICABLE, Curable.

MED'ICÆ. Sworn midwives, whose duty it was, of old, to inspect women in cases of suspected pregnancy.

MED'ICAL, Medica'lis, Iūt'ricus, Pæon'ius, Pæon'icus, Physic'al (of old), *Eculā'pian, (F.) Médical.* Same etymon as *Medicament*. Appertaining or relating to medicine or to medicines.

MEDICAL JURISPRUDENCE, Medicine, legal—m. Man, Surgeon-Apothecary.

MEDICALIS, Medical.

MED'ICAMENT, Medicamen'tum, Remed'ium, Ace'sis, Ace'sma, Ace's'tium, Ace's'tys, Althos, Hygias'ma, Ia'ma, Aeos, Medica'men, Phar'macum, Pharmaceu'ma, Alkar, Auxil'ium, Boëthe'ma,

*Med'icine, (old E.) Medsine; Physic, Rem'edy, Rem'etic, (F.) Remède; from medicor, medicatus, 'to cure or heal.' A medicine. Any substance exhibited with the view of curing or allaying morbid action. Medicines are obtained from the three kingdoms of nature, and are divided into *internal* and *external*, according as they are administered internally or applied externally.*

In common language, medicine is often called *Stuff, Doctors' stuff*, and, in the Southern States, *Truck*.

MEDICAMENTAL, Medicinal.

MEDICAMENTO'SUS LAPIS, Medie'inal stone, (F.) Pierre médicamēteuse. A name formerly given to a mixture of *peroxyd of iron, litharge, alum, nitre, sal ammoniac, and vinegar*; evaporated and calcined at a red heat for an hour. The product was regarded as eminently astringent.

MEDICAMENTUM, Drug—m. ex Palmulis, Diaphœnicon.

MEDICASTER, Medicastra. An ignorant practitioner. A charlatan.

MEDICA'TION, Medica'tio. Same etymon as *Medicament*. The change in the animal economy produced by the operation of remedies. Treatment by medicine. The act of impregnating with some medicine.

MÉDICATION HYPODERMIQUE, see Hypodermic.

MEDICATRINA, Iatrimon.

MEDICA'TRIX, same etymon. 'Having the power of healing.' A female practitioner.

MEDIC'NA, Ars medica seu Machaō'n'ia seu Machaō'n'ia, Res medica, Iat'rice, Iate'ria, Iūt'ria, Iat'ri'nē, Pæon'yñē, Pæon'icē, Iatrotech'nice, Acesto'ria, Scien'tia medica seu medendi, Sote'ria doctri'na, Med'icine, (old E.) Medsine, The healing'art, Leech craft, Physic, (F.) Médecine. Same etymon. A science, the object of which is the cure of disease and the preservation of health. Occasionally, it is used to comprehend all the branches of the healing art; at others to comprise one great division, in contradistinction to *surgery and obstetrics*. Medicine, in this sense, includes many branches;—the chief of which are, *Anatomy, Physiology, Pathology, Therapeutics, Hygiene, Materia Medica, and Pharmacy*.

MEDICINA CONSERVATIVA, Hygiene—m. Diætetica, Dietetics—m. Diasostica, Hygiene—m. Eclectica, see Eclectic—m. Efficax, Surgery—m. Equaria, Hippia'tria—m. Euectica, Gymnastics—m. Forensis, Medicine, legal—m. Gymnastica, Gymnastics—m. Hermetica, see Spagyrist's—m. Judiciaria, Medicine, legal—m. Mechanica, Kinesipathy, see Mechanical—m. Methodica, see Methodists—m. Militaris, Military medicine—m. Operativa, Surgery—m. Paracelsistica, see Spagyrist's—m. Perturbatrix, see Perturbatrix—m. Politica, Police, medical—m. Politico-forensis, Medicine, state—m. Publica, Medicine, state—m. Sinica, Chinese medicine—m. Spagyrica, Chymiatria; see Spagyrist's—m. State, Police, medical—m. Tristitia, Crocus—m. Veterinaria, Veterinary art.

MEDICINABLE, Medicinal.

MEDIC'INAL, Medicina'lis, Medicamen'ti, Medie'inal, Med'ical, Iūt'er'ius, Reme'dial, Reme'diate, (F.) Médicinal, Médicamēteux. Same etymon. Having a remedial power;—as *medicinal plants, &c.* Relating to medicine, as

MEDIC'INAL DAYS, Medice'nales Dies. Days on which the ancients considered that remedies might be administered; and especially evacuates. Such days were not esteemed critical.

MEDICINAL HOURS. Times at which—the stomach being freed from its digestive actions—medicine, it was thought, could be most advantageously administered, as in the morning fast-

ing, an hour before dinner, some hours after it, and at bedtime.

MEDICINE, Medicament, Medicina.

MEDICINE is, also, used in the same sense as Medicament, and for a purging potion.

To MEDICINE was formerly used for "to restore or cure by medicine."

MEDICINE CHEST, Pharmacotheca.

MEDICINE, CLINICAL, see Clinical — m. Eclectic, see Eclectic — m. Empirical, Arcanum — m. Galenic, Galenism.

MEDICINE, LEGAL, *Medical jurisprudence* of some, *Law or Forensic medicine*, *Medicina forensis* seu *judicialis*, *Diatricia*, (F.) *Médecine légale*. The application of medical knowledge to the solution of every question connected with the preservation of the species, and the administration of justice.

MEDICINE, PATENT, see Patent medicine — m. Political, Police, medical — m. Quack, Arcanum.

MEDICINE STAMP, Cachet.

MEDICINE, STATE, *Medicina publica* seu *politico-forensis*. The medical knowledge, brought to bear on state objects, as on public hygiene, and matters appertaining to medical jurisprudence.

MEDICINAR, Doctor.

MEDICINIER CATHARTIQUE, *Jatrophacurcas*.

MEDICO-CHIRURGICAL, *Med'ico-Ohirurgicalis*. Relating or appertaining to medicine and surgery: — as ' *medico-chirurgical society*.'

MEDICO-CHIRURGICAL ANATOMY, see Anatomy.

MEDICO-LEGAL, *Med'ico-legalis*. Relating to legal medicine; as 'a *medico-legal inquiry*, *Inquisitio medico-legalis* — 'a *medico-legal inspection*, *Inspectio medico-legalis*.

MEDICO-STATISTICAL, see Statistics, medical.

MEDICUS, Doctor, Physician — m. Equarius, Hippiastr — m. Ocularis, Oculist — m. Ophthalmicus, Oculist — m. Unguentarius, Iatralaptes — m. Veterinarius, Hippiastr — m. Vulnerarius, see Deligation — m. Vulnerum, Surgeon.

MEDIMNUS, *Medimnos, pebiuvos*. An ancient measure, capable of containing about 4 pecks and 6 pints.

MEDINOGRDIUS, *Dracunculus*.

MEDITULLIUM, Diploë.

MEDIUM MUSCULI, see Muscle — m. Ventris, Umbilicus.

MEDIUS, Middle, Median.

Medius Dig'itus, *Middle Finger*, (F.) *Doigt du Milieu*, is between the index and ring finger.

MEDLAR, *Mespilus*.

MEDSINE, Medicament, Medicina.

MEDO, Hydromeli.

MEDOBLENNORRHEA, *Leucorrhœa*.

MEDORRHEA, *Gonorrhœa* — m. *Feminarum* insons, *Leucorrhœa* — m. *Virilis*, *Gonorrhœa*.

MEDULLA, Marrow — m. *Cerebri*, see Cerebrum — m. *Dorsalis*, *M. Spinalis* — m. *Dorsualis*, *M. Spinalis* — m. *Nervorum*, *Neurine*.

MEDULLA OBLONGATA, *Oblong medulla*, *Cerebral protuberance*, *Nervous system of the senses*, (Gall.) *Cerebrum elongatum*, (F.) *Mésocephale*, *Moëlle allongée*, *Méencéphale*, (Ch.) The medullary substance that lies within the cranium upon the basilar process of the occipital bone. The anterior surface which rests in the basilar groove, is impressed by the basilar artery. At the upper extremity, and on its posterior surface are the *Tubercula quadrigemina*. At the same extremity, the medulla gives rise to two prolongations, the *peduncles of the brain*, separated from each other by the mammillary eminences, and becoming lost in the optic thalami. The lower extremity is called the *tail* or *Rachidian bulb*, *Bulbus medullæ spinalis* seu *rachidicus*, (F.)

Bulbe rachidien, and is continuous with the medulla spinalis. It is to this part only that some anatomists apply the name *Medulla oblongata*. From the posterior angles two other prolongations arise, called *Peduncles of the cerebellum*. The medulla oblongata has several eminences — the *Pons varolii*, *Corpora pyramidalia*, *C. olivaria*, &c. — and it has a longitudinal fissure before, and another behind, called the *anterior* and *posterior median fissures*. The vesicular neurine in the centre of the medulla is the nervous centre of respiration and deglutition: hence it has been called *centrum vitæ*.

MEDULLA OSSIUM, Marrow — m. *Spinæ*, *M. spinalis*.

MEDULLA SPINALIS seu *Spinæ* seu *dorsualis* seu *vertebralis* seu *dorsalis*, *Corda spinalis*, *Funis argenteus*, *Rachetrum*, *My'elus*, *M. rachit'es* seu *noticus* seu *dianche'nus* seu *pao'tes*, *My'elon*, *Notomy'elus*, *Rhachit'es*, *Eon*, *Proces'sus rachidianus*, *Sacra fistula*, *Spinal prolongation*, *Vertebral marrow*, *Spinal cord* or *marrow*, *Nervous system of voluntary motion and tactile impression* (Gall.) (F.) *Moëlle épinière*, *Moëlle vertébrale*, called by Ch., *Prolongement rachidien* of the encephalon, is the continuation of the medulla oblongata. It commences at the foramen magnum of the occipital bone, and descends in the vertebral canal as low as the 2d lumbar vertebra, without filling it; presenting, in its course, several evident enlargements. It is grooved on both its anterior and posterior surfaces by a furrow, which divides it, in its whole length, into two great nervous cords, intimately united with each other. It terminates by an oval tubercle, whence a number of nerves set out called *Cauda equina*, (F.) *Queue de cheval*, from its resemblance to a horse's tail. The spinal marrow has no analogy, as regards its structure, with the marrow of long bones. It is formed of two substances; one white, the other gray, presenting an inverse arrangement to that which they have in the brain, the white being external, the cineritious at the centre, *Substantia grisea centralis*. It is enveloped by a yellowish, fibrous membrane, very resisting, which seems to be continued insensibly from the pia mater, and by two other membranes, which are merely prolongations of the arachnoid and dura mater. See Nerve.

MEDULLA VERTEBRALIS, *M. spinalis*.

MEDULLÆ, *Sperm*.

MEDULLARY, *Medullaris*, (F.) *Médullaire*, from *medulla*, 'marrow.' Relating to the marrow, or analogous to marrow.

MEDULLARY ARTERIES. The arteries, which enter bones and pass to the marrow.

MEDULLARY CANAL, see Canal, medullary — m. Cells, see Medullary Membrane — m. Juice, Marrow — m. Matter of the brain, see Cerebrum.

MEDULLARY MEMBRANE, *Perio'teum*, *Inter'neum*, *Endo'teum*. A vascular, areolar web of extreme tenuity, which envelops the marrow and lines the inner surface of the medullary canal of the long bones. This membrane has been considered as a species of internal periosteum of those bones. It has numerous vessels, which bury themselves in the thickness of the marrow; and others which nourish the innermost plates of the bone. The cells formed by it are termed *Cellulæ* seu *Sacculi medullares*.

MEDULLARY NEURINE, see Neurine — m. Sarcoma, Encephaloid, Hæmatodes fungus.

MEDULLARY SUBSTANCE OF THE KIDNEY, see Kidney — m. s. of Schwann, see Nerve-fibre.

MEDULLARY SYSTEM. Bichat gives this name to the marrow and its membranes. He distinguishes two species of medullary systems; the one occupies the cellular tissue at the extremities of the long bones and that of the flat and short

bones; the other is found merely in the central canal of long bones.

MEDULLARY TUMOUR OF THE LUNGS, Phthisis, cancerous.

MEDULLITIS, Myelitis. M. Gerdy uses *MéduUite* for inflammation of the marrow of bones, *Osteomyelitis*. — *Myelitis* being usually appropriated to inflammation of the spinal marrow.

MEERAN, *Daucus carota*.

MEGALANTHROPOGENESIS, from *μεγας*, *μεγαλη*, 'great,' *ανθρωπος*, 'man,' and *γενεσις*, 'birth.' A term used by a French physician, named Robert, to designate the art of procreating great men; men of mind; men of genius. He considered that they may be perpetuated by always taking care to have talented men united to clever women!

MEGALOCARDIA, from *μεγας*, *μεγαλη*, 'great,' and *καρδια*, 'heart.' The state of having a very large heart.

MEGALOCÆLIA, Intestine, great, Megalop-splanchnia.

MEGALOMELTA, from *μεγας*, *μεγαλη*, 'great,' and *μελος*, 'a limb.' A monstrosity characterized by very large limbs.

MEGALOPHONIA, from *μεγας*, *μεγαλη*, 'great,' and *φωνη*, 'voice.' The condition of having a full, strong voice.

MEGALOPHTHALMUS, from *μεγας*, *μεγαλη*, 'great,' and *οφθαλμος*, 'eye.' A congenital deformity, in which the eye is inordinately large.

MEGALOPUS, Macropodus.

MEGALOSPLANCHNUS, from *μεγας*, *μεγαλη*, 'great,' and *σπλγχνον*, 'a viscus.' *Megalocælia*. Hippocrates applies the epithet to those in whom a viscus, or the viscera, are tumid. Some have used the substantive *Megalosplanchnia* for the tumour itself.

MEGALOSPLENIA, Splenoncus.

MEGETHOS, Stature.

MEGRIM, Hemicrania.

MEIBOMIUS, GLANDS OF, *Follicles of M.*, *Palpebral* or *Ciliary Follicles*, *Glandulæ sebaceæ ciliares* seu *Meibomia-næ*. *Intestinula Meibomii*, *Lacuna palpebrarum*, (F.) *Follicules palpébraux* ou *ciliaires*, *Glandes de Meibomius*. Small, sebaceous follicles, called after Henry Meibomius, although known long before his description of them. They are situate in special grooves in the tarsal cartilages; and are ranged by the side of each other, in the form of yellowish, parallel, and vertical lines; sometimes straight, at others, tortuous. They may be seen at the inner surface of the eyelids, through the conjunctiva, and secrete a sebaceous humour, called *Lippitudo*, Gum, (F.) *Chassie*.

MEIOSIS, *Meosis*, *Miosis*, *Imminutio*, *Minutio*, *Minutio*, from *μειωω*, 'I lessen.' The period of a disease, in which the symptoms begin to diminish; — the decline. Remission. Also, idiotism.

MEIUIROS, Decurtatus, Myurus.

MEKILWORT, *Atropa belladonna*.

MEL, *Meli*, *Nilivum*, *Honey*, (F.) *Miel*. A substance of a muco-saccharine nature, prepared by the *Apis mellifica*, or common bee, which collects it from the nectaries of flowers. Honey is employed as aliment, condiment, and medicine. It is demulcent and aperient; and is prescribed as an adjunct to gargles in cynanche tonsillaris, &c. It is, at times, used as a detergent to foul ulcers. *Virgin Honey*, *Mel virginum* seu *virginum*, (F.) *Miel vierge*, is that which flows from the wax spontaneously.

HILL'S BALSAM OF HONEY is formed of *balsam of tolu*, ℥j, *honey*, ℥j, *rectified spirit*, one gallon. It was long a celebrated empirical pectoral.

HONEY WATER is a mixture of essences coloured with saffron. A little honey is added to

communicate a clamminess; the effect of which is to make it retain the scent the longer. It is used as a scent.

MEL ACETATUM, Oxy-mel — m. *Ægyptiacum*, *Ægyptiacum* — m. *Aërium*, *Fraxinus ornus* — m. *Arundinaceum*, *Saccharum*.

MEL BORA'CIS, *M. subbora'cis* seu *bora'atum*, *Colluto'rium adstrin'gens*, *Linimentum de Bora'cè* seu *ad aphthas*, *Linctus de Bora'cè* seu *ad aphthas*, *Honey of Borax*, *Mellite de borax*, *Miel boratè*. (*Boracis contrit. ʒj. mellis despumati ʒij*, Ph. L.) Detergent. Applied to the mouth in aphthous affections.

MEL CANNÆ, *Saccharum* (non purificatum) — m. *Coctum*, *Pelicide*.

MEL DESPUMA'TUM, *M. Depura'tum*, *Clar'ified Honey*, *Anaphrom'eli*, (F.) *Miel Clarifié* ou *despumé* ou *dépuré*, *Mellite simple*, *Sirap de miel*. Melt the honey in a water-bath, and remove the scum. Use the same as honey.

MEL ROSÆ, *M. rosa'tum*, *Mellitum de Rosis*, *Rhodom'eli*, *Rhodostacton*, *Honey of Roses*, (F.) *Miel rosat*, *Mellite de Rosas*. (*Rosæ gallic. ʒij*, *aqua bullient. f ʒij*, *mellis despum. f ʒxx*. Macerate the roses in f ʒvij of boiling water for four hours; press out as much fluid as possible and set aside. Macerate the residue in f ʒiv of boiling water for half an hour, and again express. Reserve f ʒiv of the first infusion; mix the remainder with the infusion last obtained; add the honey, and evaporate to a pint. Lastly, add the reserved infusion, and strain. Ph. U. S.) Astringent and detergent. Used chiefly in gargles and washes for aphthæ, &c.

MEL SCILLÆ, Oxy-mel scillæ — m. *Scillæ compositum*, *Syrupus scillæ compositus*.

MEL VINOSUM, *Ænomet* — m. *Virginum*, see Mel.

MELÆ'NA, *Melanorrhag'ia*, *Melanorrhag'ia*, *Melæ'no-diarrhœ'a*, *Icterus niger*, *Melauchlos'ia*, *Melanict'erus*, *Melas Icterus*, *Melanism'us*, *Morb'us niger seu niger Hippoc'ratie*, *Enterorrhœ'a*, *Black Jaundice*, (F.) *Mélène*, *Mêlène*, *Mélasictère*, *Maladie noire*, *Mêlène*, *Ictère noir*; from *μελας*, *melanos*, 'black.' A name given to vomiting of black matter, ordinarily succeeded by evacuations of the same character. It seems to be often a variety of hæmatemesis. The *Black Vomit* in yellow fever is owing to a morbid secretion from the lining membrane of the stomach and small intestines mixed with blood. *Melæna* also signifies hemorrhage from the intestines; *Fluxus sple'nicus*, *Dysenter'ia sple'nica*, *Defec'tiones nigre*, *Seces'sus niger*, *Hemorrhage from the Intes'tines*.

MELÆNA FUNGOSA CARCINODES, *Hæmatodes fungus*.

MELÆNODIARRHÆA, *Melæna*.

MELÆNOGOGUS, from *μελας*, *melanos*, 'black,' and *αγω*, 'I drive away.' A medicine believed to be capable of expelling atrabilis or 'black bile.'

MELÆNORRHAGIA, *Melæna*.

MELLAINE, see *Choroidea tunica*.

MELALBU'CA CAJAPUTI, *M. Minor* seu *Leucoden'dron*, *Myrtus Leucaden'dron*, *Leptospermum Leucaden'drum*. Ord. *Myrtaceæ*. This plant affords the *Cajaput Oil*, (see *Cajuput*.) The leaves are esteemed diuretic, stomachic, and emmenagogue.

MELALBU'CA LEUCODENDRON, *M. Cajaputi* — m. *Minor*, *M. Cajaputi*.

MELAMPHONUS, from *μελας*, 'obscure,' and *φωνη*, 'voice.' Having a hoarse or indistinct voice.

MELAMPHYLLUM, *Acanthus mollis*.

MELAMPODE, *Helleborus niger*.

MELAMPODIUM, *Helleborus niger*.

MELANÆMA, Suffocation.

MELANÆMIA, Venosity.

MELANAGOGUE, *Melanago'gus*, from *μελας*,

melanos, 'black,' and *αγω*, 'I expel.' A medicine which the ancients believed adapted for expelling black bile or melancholy.

MELANCHLO'RUS, from *μελας*, *μελανος*, 'black,' and *χλωρος*, 'green.' The ancients gave this name to certain dark-coloured topical remedies. See *Melæna*.

MELANCHOLE, *Atrabilis*.

MELANCHOLIA, *Melancho'ly* — *m. Autochrica*, Suicide — *m. Errabunda*, *Cuttubuth*, *Kutubuth* — *m. Flatuosa*, *Hypochondriasis* — *m. Hypochondriaca*, *Hypochondriasis* — *m. Nervæa*, *Hypochondriasis* — *m. Pleonectica*, see *Pleonectica* — *m. Saltans*, *Chorea* — *m. Suicidium*, *Suicide* — *m. Uterina*, *Nymphomania* — *m. Zoanthropia*, *Melancho'ly*.

MELANCHOLIC, *Melancho'licus*, *Melan'cho'lus*, *Melancho'les*, (Sc.) *Dram*, *Drum*, (F.) *Mélanco'lique*, from *μελας*, *μελανος*, 'black,' and *χολη*, 'bile.' One labouring under melancholy. That which belongs or relates to melancholy. In popular language, one of a gloomy disposition.

MELANCHOLINESS, *Melancho'ly*.

MELANCHOLODES, *Melancho'lic*.

MELANCHOLUS, *Melancho'lic*.

MEL'ANCHOLY, *Melancho'ly*, same etymon. *Lypemania*, *Euphro'nia*, *Melancho'lia*, *Mania*, *Melancho'lica* seu *Melancho'lia*, *Tristema'nia*, *Baryth'mia*, *Hypoleptoma'nia*, *Ano'sia adstricta*, *Fixatio nominum*, *Mel'ancholiness*, (Sc.) *Dismal*, (F.) *Mélan'colie*, *Lypophrénie*. Same etymon. A disease supposed, by the ancients, to be caused by black bile. A variety of mental alienation, characterized by excessive gloom, mistrust, and depression, generally, with insanity on one particular subject or train of ideas, *Monomania*, *Monoma'nia*, *Panophobia*, *Demonomania*, *Erotomania*, *Nostalgia*, &c., may be referred to this head.

Melancho'ly is also used for unusual gloominess of disposition.

MELANCHOLY, **EROTIC**, *Erotomania* — *m. Love*, *Erotomania*.

MELANCHRUS, *Melan'ochrus*, from *μελας*, *μελανος*, 'black,' and *χρως*, 'a colour.' One attacked with black jaundice. Supposed, by some, to be an abbreviation of *Melanchlorus*.

MÉLANCOLIE, *Melancho'ly*.

MÉLANCOLIQUE, *Melancho'lic*.

MELANEPHRIDOSIS, *Sudor Anglicus niger*.

MÉLANGE PECTORAL, *Mistura acidi hydrocyanici*.

MELAN'IC, *Melan'icus*, *Melanot'ic*, *Melanot'icus*. (F.) *Mélané*; from *μελας*, *μελανος*, 'black.' Of or belonging to *Melanosis*; — as *Melan'ic deposit*, a black colouring matter deposited from the blood under special circumstances; — see *Melanosis*.

MELANIC DEPOSIT, *Melanosis*.

MELANICTERUS, *Melæna*.

MÉLANIENNES, (*Tachee*), *Nævus*.

MÉLANINE, see *Choroidea tunica*.

MÉLANISME, *Melanismus*.

MELANISMUS, *Melæna*. M. I. G. St. Hilaire employs the term *Mélanisme*, for the change of coloration of the skin in which the pigment is of a deeper hue and in greater quantity than natural.

MELANOCARCINOMA, *Cancer*, *melanotic*.

MELANOCHRUS, *Melanochrus*.

MELANOMA, *Melanosis* — *m. Pulmonum*, see *Melanos*.

MELANOMYCES, *Melanospongus*.

MELANOPATHI'A, from *μελας*, *μελανος*, 'black,' and *παθος*, 'affection.' *Nigritism*, *Nigrit'ies*. A disease of the skin, which consists

in augmentation of black pigment; generally in patches.

MELANOPHYMA, *Ergot*.

MELANOPIPER, *Piper nigrum*.

MELANOPNEUMON, see *Melanos*.

MELANORRHAGIA, *Melæna*.

MELANORRHIZUM, *Helleborus niger*.

MÉLANOSE, *Melan'ois*.

MELANO'SIS, *Melan'ois*, *Melano'ma*, *Nigritu'do*, (F.) *Dégénérescence noire*, *Mélanose*; from *μελας*, *μελανος*, 'black.' An organic affection, in which the tissue of the parts is converted, owing to a *melan'ic deposit*, into a black, hard, homogeneous substance, near which ulcers or cavities may form; — owing to the softening, either of the substance itself, or of some other morbid tissue, — of tubercles especially. This morbid change affects the lungs particularly; when it is called *Melano'ois* seu *Melano'ma* seu *Anthraco'ois Pulmo'nium*, *Melanopneu'mon*, *Nigritu'do melanot'ica*, *Melan'is Pulmonum*, *Pneumomonelano'ois*, *Black tubercle*, (F.) *Mélanose Pulmonaire*. It is, also, met with in the liver and areolar texture. Its causes are very obscure. *Melanos* of the lungs constitutes one of the species of phthisis of Bayle; but it is impossible to distinguish it from the other species during life.

In the classification of Fuchs, *Melanosen* (G.) forms a family of diseases.

The deposition of melan'ic matter in the tissues is not necessarily connected with malignant disease. See *Cancer*, *melanotic*.

MELANOSIS PULMONUM, see *Melanos* — *m. Universalis*, *Venosity*.

MELANOSMEGMA, see *Sapo*.

MELANOSPONG'US, *Melanomy'ces*, from *μελας*, *μελανος*, 'black,' and *σπογγος*, 'sponge.' The tubiform variety of *melanos* of the lungs.

MELANOTIC, *Melan'ic* — *m. Cancer*, *Cancer*, *melanotic*.

MELANOTHRIX, *Atricapill'us*, (F.) *Mélanotrique*, from *μελας*, 'black,' and *τριξ*, 'hair.' Having black hair.

MELANSIS, *Melanos* — *m. Pulmonum*, *Melanos* *pulmonum*.

MELANTHIUM, *Nigella*, *M. Virginicum* — *m. Muscætoxicum*, *Amiantum muscætoxicum*.

MELANTHIUM VIRGIN'ICUM, *Melanthium*, *Leiman'thium Virgin'icum*, *Quasdil*; indigenous; Order, *Melanthaceæ*, flowering in July, is said to be a sure but severe remedy for itch.

MELAS, *Lepra nigricans* — *m. Icterus*, *Melæna*.

MELAS'MA, *Echymo'ma Melas'ma*, *Nigror*, from *μελας*, 'black.' A black spot or ecchymosis, occurring on the lower extremities of old people especially. Also a cutaneous affection analogous to *chloasma*, differing from it only in the dark colour of the morbid pigment. It is also called *Pityri'asis nigra*.

MELAS'SES, *Molas'ses*, *Melus'tum*, *Theriaca communi's*, *Syr'upus empyreumaticus* seu *niger*, *Sac'chari Fæx*, *Treacle*, (Old Eng.) *Triacle*; from *μελι*, 'honey,' or from *μελας*, 'black.' The syrup, which remains after the juice of the sugar-cane has been subjected to all the operations for extracting sugar from it. In the United States, the syrup, made in the process of forming common sugar, is called *melasses*; that remaining after the refinement of sugar being termed *sugar-house melasses* or *treacle*.

MELASSES SPIRIT, *Rum*.

MELATROPH'IA, from *μελος*, 'a limb,' and *ατροφια*, 'wasting.' Wasting of the limbs.

MELCA, *μελκα*. Food made of milk, coagulated by boiling vinegar.

MELE, *Specillum*.

MELEA, *Pyrus malus*.

MELEGUETTA, *Amomum grana paradisi*.

MELEI'OS, *μηλεις*, *Meli'us*, *Melias*. A species of alum found in the island of Melos.

MÉLÉNA, *Melæna*.

MÉLÈNE, *Melæna*.

MÉLÈZE, *Pinus larix*.

MELI, *Mel*.

MELIA, *Fraxinus ornus*.

ME'LIA AZED'ARACH, *Azedàra'cha amœ'na*, *Poison Berry Tree*, *Azed'arach*, *Pride of India* or of *China*, *Pride tree*, *Hop tree*, *Bead tree*. *Nat. Ord.* Meliaceæ; *Sec. Syst.* Decandria Monogynia. The bark of the root—called in India *Neem Bark*, *Azedarach*, (*Ph. U. S.*), is usually given in decoction, in the proportion of three or four ounces of the bark of the fresh root to a quart of water, boiled down to a pint. The dose is one or two table-spoonfuls, every two or three hours, till purging is induced. It is given in this manner as an anthelmintic. It is sometimes formed into an ointment, and used in tinea capitis. The tree is a native of Syria, Persia, and the north of India, and is abundant in the southern states of the Union.

MELIA TERRA, *Creta*.

MELIAN'THUS MAJOR, *Nat. Ord.* Zygophyllæ. A south African plant, a decoction of whose leaves is a good external remedy in tinea capitis, crusta serpigiosa, necrosis and foul ulcers. It is also useful as a gargle and lotion in sore throat and diseases of the gums. The bruised leaves applied to ulcers promote granulation.

MELICERA, *Porrigio favosa*.

MELICERIA, *Hydrarthrus*, *Porrigio favosa*.

MELICE'RI'S, *Melifa'vium*, from *μελι*, 'honey,' and *κρυος*, 'wax.' *Emphy'ma encys'tis melice'ris*. An encysted tumour filled with a substance resembling honey. Also, *Porrigio favosa*.

MELIC'EROUS, *Melicer'itos*: same etymon. Having the characters of meliceris,—as a melicerous tumour.

MELICERUM, *Porrigio favosa*.

MELICHEI'UM, from *μελι*, 'honey,' and *χew*, 'I pour out.' A honey-like discharge from an ulcer.

MELICRATUM, *Hydromeli*.

MELIFAVIUM, *Meliceris*, *Porrigio favosa*.

MELIGE'TON, from *μελι*, 'honey.' A fetid humour of the consistence of honey, discharged from ulcers, accompanied with caries of a bone.

MÉLILOT, *Trifolium melilotus*.

MELILOTUS, *Trifolium melilotus*.

MELIME'LUM, from *μελι*, 'honey,' and *μηλον*, 'an apple.' A name given to two compounds:—one of honey and quince, the other of honey and apples.

MÉLINE, *Melæna*.

MELINI SULPHAS, *Cadmii sulphas*.

MELI'NUM, *μηλινον*. The ancient name of an ointment, and of several plasters, described by Dioscorides and Galen: so called from their resembling the *μηλον* or quince in colour.

MELINUM SULPHURICUM, *Cadmii sulphas*.

MELINUS, *Meleios*.

MÉLIOLE, *Mellitum*.

MELIPHYLLUM, *Melissa*.

MELIS'SA, from *μελισσα*, 'a bee,' because bees gather honey from it. *M. officina'lis* seu *Romana* seu *hirsu'ta* seu *cit'rina* seu *cit'ra'ta* seu *horten'sia*, *Citra'go*, *Citra'ria*, *Melite'a*, *Melissob'otos*, *Melite'na*, *Melissobot'anum*, *Melissophyll'um*, *Meliphyllum*, *Mentha cit'ra'ta*, *Melit'tia*, *Cedronel'la*, *Apiastrum*, *Ero'tion*, *Balm*, (*F.*) *Melisse*, *La Citronelle*, *Céline*; *Nat. Ord.* Labiatae. The leaves of balm, *Melissa* (*Ph. U. S.*), have a pleasant odour; and an austere and aromatic taste. Balm was formerly much used in nervous diseases, but is now only em-

ployed when made into tea, as a diuretic, in febrile affections. It is the basis of a celebrated preparation,—the *Eau des Carmes*.

MELISSA CALAMIN'THA, *Calamin'tha*, *C. vulga'ris* seu *officina'rum* seu *grandiflo'ra*, *Thymus calamin'tha*, *Cal'amint*, *Nespite*, (*F.*) *Calament*. This plant smells like wild mint, though more agreeably. It is used, popularly, as a tea in dyspepsia; flatulent colic; hysteria; uterine obstructions, &c.

MELISSA CANARIENSIS, *Dracocephalum Canariense*—m. *Citrata*, *Melissa*—m. *Citrina*, *Melissa*—m. *Clinopodium*, *Clinopodium vulgare*.

MELISSA GRANDIFLO'RA, *Thymus glandiflorus*, *Calamin'tha magno flore*, *C. mont'na*, *Mountain Cal'amint*. This plant resembles the last in virtues.

MELISSA HIRSUTA, *Melissa*—m. *Hortensis*, *Melissa*—m. *Humilis*, *Melittis melissophyllum*.

MELISSA NEP'ETA, *Thymus nep'eta* seu *multiflorus*, *Calamin'tha Anglica* seu *Pulegi'i odorè* seu *nep'eta* seu *parviflo'ra* seu *trichot'oma*, *Nep'eta agrest'is*, *Field Cal'amint*, *Spotted Cal'amint*. Formerly used as an aromatic.

MELISSA OFFICINALIS, *Melissa*—m. *Pulegioides*, *Hedeoma*—m. *Romana*, *Melissa*—m. *Tragi*, *Melittis melissophyllum*—m. *Turcica*, *Dracocephalum Canariense*.

MÉLISSE DES BOIS, *Melittis melissophyllum*—m. *de Moldavie*, *Dracocephalum Canariense*.

MELISSOBOTANUM, *Melissa*.

MELISSOBOTOS, *Melissa*.

MELISSOPHYLLUM, *Melissa*, *Melittis melissophyllum*.

MÉLISSOT, *Melittis melissophyllum*.

MELITÆ'MIA, *Melithæ'mia*, from *μελι*, 'honey,' and *αἷμα*, 'blood.' A condition of the circulating fluid, in which it contains an unusual quantity of saccharine matter.

MELITÆNA, *Melissa*.

MÉLITAGRE, *Impetigo*.

MELITEIA, *Melissa*.

MELITHÆMIA, *Melitæmia*.

MELITHOCORTON, *Corallina Corsicana*.

MELITHYPERURIA, *Diabetes*.

MELI'TIS, *Inflammat'io genæ*; from *μηλον*, 'the cheek,' and *itis*, denoting inflammation. Inflammation of the cheek.

MELITITES, *Hydromeli*.

MELITTA, *Bee*.

MELITTIS, *Melissa*.

MELI'TIS MELISSOPHYLL'UM, *La'mium montanum*, *Melissa humilis* seu *sylyest'ris* seu *Tragi*, *La'mium Plin'ii*, *Melissophyll'um*, *Bastard Balm*, (*F.*) *Mélisse des Bois*, *Mélistot*. *Ord.* Labiatae. This plant was formerly employed, like the balm, in uterine obstructions.

MELITTIS SYLVESTRIS, *M. Melissophyllum*.

MELITURIA, *Urine*, diabetic, see *Diabetes*.

MELI'TUS, *Mell'itus*, from *mel*, 'honey.' Appertaining to honey. Of the nature of honey.

MELIZO'MUM, from *μελι*, 'honey,' and *ζωμος*, 'broth'; *Mead*, *Melli'na*, *Metheg'lin*, (*Prov.*) *Mulin*. A drink prepared with honey.

MELLA'GO, from *mel*, 'honey.' Any medicine having the consistence of honey. A fluid extract.

MELLICRATUM, *Hydromeli*.

MELLIFAVIUM, *Porrigio favosa*.

MELLINA, *Melizomum*.

MELLITE DE BORAX, *Mel boracis*—m. *de Roses*, *Mel rose*—m. *Simple*, *Mel despumatum*.

MELLITUM, from *mel*, 'honey.' (*F.*) *Miel me'dicinal*, *Hydromellé*, *Saccharolé liquide*, *Meliolé*. A prepared honey:—a medicated honey.

MELLITUM DE ACETATE CUPRI, *Linimentum Æuginis*.

MELLITUM DE MERCURIA'LI COMPOS'ITUM, (F.)
Miel de mercuriale composé. Syrup de longue vie.
 (Succ. purif. mercurialis ℥ij, boraginis, anchusæ officinalis, āā ℥ss, iris pseudacori radicis recent. ℥ij, rad. nicc. gentianæ 3j, mellis ℥ij, vini albi 3xij.) Macerate the bruised roots in the wine for 24 hours; strain; mix the juices and honey; boil slightly, and filter; then add the two liquors, and boil to the consistence of syrup. (Ph. P.) Cathartic, stomachic, &c.

MELLITUM DE ROSIS, Mel rosæ.

MELLITUM SIMPLEX, Syr'upus de Mellē, Syrup of Honey. (Mellis ℔vj, aquæ ℔ss, carbonat. calcis. præparat. 3ij.) Mix the honey, water, and carbonate of lime in a silver vessel: boil the mixture, stirring, at the same time, for 2 or 3 minutes: then add prepared animal charcoal 3vj, two whites of eggs, mixed in a pint of water. Mix all, and boil to the consistence of syrup: remove from the fire; let the syrup rest for 15 minutes, and pass through a cloth. (Ph. P.) Demulcent.

MELLITUS, Melitus.

MELOÆMIA, from μέλας, 'black,' and αἷμα, 'blood.' A state of blood, characterized by its incoagulability, black colour, and septic properties.

MELOCHITIS, Lapis Arme'nus, Armeni'tes, Arne'nian Stone. A variety of blue carbonate of copper. It is found in Germany, the Tyrol, and especially in Armenia. It was formerly employed as a cardiac, and as proper for purging away melancholy.

MELODIDYMUS, from μέλος, 'a limb,' and ἴδιος, 'double.' A double member or limb.

MELOE MAJALIS, see Cantharis.

MELOE NIGER, M. Pennsylvan'icus; Class, Insecta; Ord. Coleoptera. A blistering fly, native of the United States. It feeds upon *Prunella vulgaris* or *Self-heal*, and *Ambrosia trifida* or *Strick-weed*. These flies resemble the Spanish flies in properties.

MELOE PENNSYLVANICUS, M. niger.

MELOE PROSCARABÆUS, Cantarel'lus, was anciently used as a diuretic and anti-hydrophobic. See *Cantharis*.

MELOE VESICATORIUS, Cantharis.

MELOMÆLES, from μέλος, 'a member,' and μέλος, 'unprofitable.' A monster with supernumerary limbs.

MELON, μῆλον, 'an apple.' A disorder of the eye, in which it protrudes out of the socket. See *Exophthalmia*.—Castelli.

MELON, Cucumis melo — m. d'Eau, Cucurbita citrullus — m. Plant, Cucumis melo — m. Water, Cucurbita citrullus.

MELONGENA, Solanum melongena.

MELOPLACUNTUM, Meloplacus.

MELOPLA'CUS, Meloplacuntum, from μῆλον, 'an apple,' 'a quince.' A compound obtained by boiling wine, honey, quince, pepper, &c., together.—Galen.

MELOPLAS'TIC, Meloplas'ticus, Genoplas'tic, from μῆλον, 'the cheek,' and πλασσω, 'I form.' (F.) Meloplastie, an epithet for the operation for forming a new cheek when any part of it has been lost.

MELOS, μέλος, Membrum. A member. A limb. An organized part, composed of other parts.—Castelli.

MELO'SIS, Catheteris'mus, Cenembate'ois, from μῆλη, 'a probe.' The act of probing a wound, ulcer, &c.—Hippocrates. Catheterism.

MELOTHRUM, Bryonia alba.

MELO'TIS, Melo'tris, Specill'um minus. From μῆλη, 'a probe.' A small probe,—a probe for the ear. See *Apyromele*.

MELOTRIS, Melotis.

MELT, Spleen. Also, the sperm of fish,—Milt.

MELTING, Fusion.

MELUM, μήλον, Malum. An apple. Fruit in general. Also, a round, firm female breast. The cheek. The apple-eye or melon-eye is an apple-formed projection of the eye from the orbit.

MELUM ARMENIACUM, Prunus armeniaca — m. Cydonium, see Pyrus cydonia.

MELUSTUM, Melasses.

MEMBRA PUDENDA, Genital organs.

MEMBRANA, Membrane — m. Abdominis, Peritonæum — m. Adiposa, Cellular membrane — m. Adventitia, Decidua reflexa — m. Agnina, Amnios — m. Amphiblestodes, Retina — m. Arachnoidea, Hyaloid membrane — m. Capsularis testis, Albuginea — m. Carnosa, Dartos — m. Cellulosa, Cellular membrane — m. Cellulosa, Decidua — m. Cerebri tenuis, Pia mater — m. Cerebri mollis, Pia mater — m. Cerebri propria, Pia mater — m. Chorion capillaris, see Choroida tunica — m. Circumossalis, Periosteum — m. Communis, Decidua — m. Cor circumplexa, Pericardium — m. Cordis, Pericardium — m. Costalis, Pleura — m. Costas succingens, Pleura — m. Crassa, Decidua reflexa — m. Cribrosa, Decidua — m. Cuticularis, Dura mater — m. Decidua serotina, see Decidua — m. Demuriana, see Aqueous humour, and Cornea — m. Dentata, Denticulatum ligamentum — m. Dermatodes, Dura mater — m. Descemetii, see Aqueous humour, and Cornea — m. Diaphragmaton, Mediastinum — m. Dura cerebrum ambiens, Dura mater — m. Epicrania, Galea aponeurotica capitis — m. Externa dentium, Tapetum alveoli — m. Farcimialis, Allantois — m. Fenestra rotunda, see Fenestra — m. Filamentosa, Decidua reflexa — m. Flocculenta, Decidua — m. Fœtum involvens, Amnios — m. Fusca, see Sclerotic.

MEMBRANA GRANULO'SA, Gran'ular membrane or layer, (F.) Membrane granuleuse, Couche celluleuse (Coste). A layer of yellow, granular matter, which lines the inner layer of the Graafian follicle, and is regarded by Kölliker as epithelium.

MEMBRANA HUMORIS AQUEI, see Aqueous humour — m. Interspinalis, Interspinal ligament — m. Intestinalis, Allantois — m. Jacobi, Tunica Jacobi — m. Jacobi Arnold, see Iris — m. Limitans, see Iris — m. Limitans Pacini, see Iris — m. Linguae, see Tongue — m. Macilentior, Epiploon, gastrohepatic — m. Mucosa, Decidua, see Membrane — m. Nervorum, Neurilemma — m. Nictitans, Valvula semilunaris — m. Olfactoria, Pituitary membrane — m. Ossis, Periosteum — m. Ovi materna, Decidua — m. Ovuli corticalis, Oöcylomma — m. Pellucida, Amnios — m. Pigmenti, see Iris — m. Pinguedinea, see Cellular membrane — m. Pinguis intestinorum, Mesentery — m. Pituitaria, Pituitary membrane, see Membrane — m. Pituitosa, see Membrane — m. Pleuretica, Pleura — m. Præexistens, Decidua — m. Propria, Basement membrane, and Folliculi Graafiani, see Cortex ovi — m. Pupillar, Pupillary membrane.

MEMBRANA PUTAMINIS, (putamen, 'a shell or husk.') Pellic'ula Ovi. The membrane adherent to the inner surface of the eggshell. — Putamen seu Testa Ovi.

MEMBRANA RETIFORMIS CHORII, Decidua reflexa — m. Ruychiana, Choroid (inner layer.)

MEMBRANA SACCIFORM'IS. A separate synovial membrane for the inferior radio-cubital articulation, which covers the upper surface of the triangular ligament, and the sort of incomplete ring which circumscribes the head of the ulna.

MEMBRANA SEMIPELLUCIDA. Caligo — m. Serosa, Membrane (serous) — m. Sinuosa, Decidua — m. Subcostalis, Pleura — m. Succingens, Diaphragm, Pleura.

MEMBRANA TESTE, Shell membrane. The

membrane that lines the shell of the bird's egg:—a simple membrane.

MEMBRANA THORACEM INTERSEPIENS, Mediastinum—*m.* Tympani, see Tympanum—*m.* Tympani secundaria, see Fenestra—*m.* Urinaria, Alantoides—*m.* Uteri interna evoluta, Decidua—*m.* Uvea, Uvea—*m.* Vasculosa cerebri, Pia mater—*m.* Vasorum Communis, Endangium—*m.* Verricularis, Retina.

MEMBRANA VERSICOLOR. A peculiar membrane—according to Mr. Fielding, of Hull—situate immediately behind the retina, and in connexion with it. It is separable into distinct layers from the choroid, and is supplied with blood-vessels.

MEMBRANA WACHENDORFIANA, Pupillary membrane.

MEMBRANACEOUS, Membranous.

MEMBRANE, *Membrana*, *Hymen*, *Meninx*, *Ependyma*, *Pannicle*. A name given to different thin organs, representing a species of supple, more or less elastic, webs; varying in their structure and vital properties, and intended, in general, to absorb or secrete certain fluids; and to separate, envelop, and form other organs. Bichat has divided the membranes into simple and compound.

Simple membranes comprise three orders. 1. A Mucous membrane, *Membrana mucosa* seu *pituitosa* seu *pituitaria*, *Phlegmymen*, *Phlegmat'hymen*, *Membrane folliculeuse* (Ch.), is so called, on account of the mucous fluid by which it is constantly lubricated. Mucous membranes line the canals, cavities, and hollow organs, which communicate externally by different apertures on the skin. Bichat refers them to two great divisions—the *gastro-pulmonary*, and the *genito-urinary*. The mucous membranes have a striking analogy with the cutaneous tissue, in organization, functions, and diseases. They are composed of chorion, papillæ, and epidermis; and are furnished with a multitude of follicles, which secrete a viscid humour,—mucus. They receive a quantity of arterial vessels, veins, lymphatics, and nerves. 2. The Serous membranes, *Membrana serosa*, *M. villosa* simple (Ch.), are transparent, thin, and composed of one lamina. One surface adheres to other textures; the other is smooth, polished, and moistened by a serous fluid. They are arranged—in the form of sacs without apertures—as great, intermediate reservoirs for the exhalant and absorbent systems, in which the serous fluid, in passing from one system, carries some time before it enters the other. The serous membranes resemble the areolar membrane in structure and diseases. They facilitate the motion on each other of the organs which they envelop. They may be divided into (a) *Serous membranes*, properly so called; as the arachnoid, pleura, peritoneum, and tunica vaginalis. (b) *Synovial membranes or capsules*; which belong to joints, tendons, aponeuroses, &c. These membranes—mucous and serous—are constituted of similar layers—epithelium, basement membrane, condensed areolar tissue; and a looser form of areolar tissue, termed *submucous* in one case,—*subserous* in the other. 3. *Fibrous membranes*, *Membranes albugineuses* (Ch.) These are almost all continuous, and terminate at the periosteum—their common centre. They have been divided into two sections. 1. *Enveloping aponeuroses*, *Aponeuroses of insertion*, *Fibrous capsules of the joints*, and *Fibrous sheath of tendons*. 2. The *Periosteum*, *Dura mater*, *Sclerotica*, *Tunica albuginea testis*, *Fibrous membrane of the spleen*, &c. The fibrous membranes are not free or moistened by any particular fluid. They adhere by both surfaces to the neighbouring parts; are firm, resisting, but slightly elastic, and of a white colour;

sometimes pearly and glistening. Their vessels are numerous, in some, as in the dura mater and periosteum; in others, scarcely perceptible, as in the aponeuroses. The presence of nerves has never been proved, although several circumstances, regarding their sensibility, render their existence probable. The fibrous membranes serve, in general, to augment the solidity of the organs which they envelop; to retain the muscles in their respective positions; to favour the motion of the limbs, and that of the muscles and skin; to form canals and rings for the passage of different organs, &c.

Bichat admits three species of Compound Membranes. 1. The *Sero-fibrous*, formed of a serous and fibrous lamina, intimately adherent to each other;—as the pericardium, dura mater, and tunica albuginea. 2. The *Sero-mucous*, formed of a serous and mucous lamina;—as the gull-bladder at its lower part. 3. The *Fibro-mucous*, constituted of the union of a fibrous and mucous membrane; as the mucous membrane of the nasal fossæ, gums, &c. Chaussier admits six kinds of membranes. 1. The *laminated*. See Cellular Tissue. 2. The *serous or simple villous*. 3. The *follicular or complicated villous*. 4. The *muscular or fleshy*. 5. The *albugineous*. 6. The *albuminous*.

MEMBRANE, ACCIDENTAL, *M. false*—*m.* Accidentelle, *M. false*—*m.* Adnée, Conjunctiva—*m.* Adventitious, *M. false*—*m.* Anhistous, Decidua.

MEMBRANE, BASEMENT, *Primary or liminary membrane*, *Membrana propria*. A delicate, structureless lamella of membrane found beneath the epidermis or epithelium, on all the free surfaces of the body.

MEMBRANE CADUQUE, Decidua—*m.* Caduque réfléchi, Decidua reflexa—*m.* de Demours, see Aqueous Humour, and Cornea—*m.* de Desceemet, see Aqueous Humour, and Cornea.

MEMBRANE, FALSE, *Accidental or Adventitious membrane*, *Pseudo-membrane*, *Pseud'hymen*, *Pseudomeninx*, (F.) *Fausse membrane*, *M. accidentelle*. Membranous productions, which form on all the free natural surfaces, and on every free accidental surface, are so called: They are, in general, produced by the exudation of a fibrinous matter, susceptible of organization, which takes place in consequence of inflammation of the various tissues. These accidental membranes occur on the skin after the application of a blister; on mucous surfaces, as in croup; on the parietes of inflamed veins and arteries, &c. The cicatrices of wounds are formed of them.

Of late, it has been proposed to give the name *pseudo-membrane* to fibrinous exudations, which are devoid of organization, whilst *neo-membrane* may be applied to such fresh productions as have become supplied with vessels and nerves.

MEMBRANE FAUSSE, Membrane, false—*m.* Germinal, see Molecule—*m.* Granular, *Membrana granulosa*—*m.* Granulose, *Membrana granulosa*—*m.* Haptogen, see Haptogenous—*m.* Primary, Membrane, basement—*m.* Pseudo, *M. false*—*m.* Hyaloid, see Hyaloid membrane—*m.* Jacob's, *Tunica Jacobi*—*m.* Limitary, Membrane, basement—*m.* Nasnyth's, see Enamel—*m.* Ruyschienne, Choroid (inner layer), *Ruyschiana tunica*—*m.* Shell, *Membrana testæ*—*m.* du Tympan, see Tympanum—*m.* Virginal, Hymen—*m.* Vitellary, *Zona pellucida*—*m.* Vitelline, *Zona pellucida*.

MEMBRANES ALBUGINEUSES, Membranes, fibrous, see Albuginea—*m.* Folliculeuses, Membranes, mucous—*m.* Veloutées, Villous membranes—*m.* Villoses simples, Membranes, serous.

MEMBRANES OF THE FETUS, *Membranae foetus involvrentes*, *Velamenta infantis*, *Integumenta foetus*, *The membranes*. The membranes which immediately envelop the fetus in the cavity of the uterus and the rupture of which gives rise

to the discharge of the liquor amnii. These membranes are the *decidua*, *chorion*, and *amnion*.

MEMBRANES, THE, Membranes of the Fœtus.

MEMBRANEUX, Membranous.

MEMBRANIFORM, *Membraniformis*, from *membrana*, and *forma*—resembling a membrane. A name given to thin and flat parts, which resemble membranes.

MEMBRANIFORM MUSCLES. Very broad and thin muscles, as the *platysma myoides*, *obliquus abdominis*, &c.

MEMBRANOSUS, *Fascia lata* muscle.

MEMBRANOUS, *Membranosus*, *Membrana'-ceus*, *Hymenoides*, (F.) *Membraneus*. Having the nature of membrane. Formed of membrane.

MEMBRANULA, *Hymenium*—m. *Coronæ ciliaris*, see *Ciliary*—m. *Lunata Vaginæ*, *Hymen*—m. *Nervorum*, *Neurilemma*—m. *Semilunaris conjunctivæ*, *Valvula semilunaris*.

MEMBRANULE, *Membran'ula*, diminutive of *membrana*, 'a membrane.' A small membrane.

MEMBRE, *Membrum*—m. *Viril*, *Penis*.

MEMBROSUS, *Membro'sior*, *Mentula'tus*, *Mutonia'tus*, *Nasa'tus*, *valde Mentula'tus*, *Fascino'-eus*, *Peolon*, (F.) *Membru*, from *membrum*, 'the male organ.' One whose penis is very large.

MEMBRU, *Membrosus*.

MEMBRUM, *Artus*, *Melos*, *Colon*, a limb, a member, (F.) *Membre*. The limbs or extremities or members of animals are certain parts exterior to the body, which are more or less apparent, long, and moveable. It is by means of their limbs, that animals transport themselves from one place to another; and that they defend themselves, or attack others. The limbs are arranged on each side of the trunk, with which they are articulated. In man, they are four in number:—two upper or thoracic, and two lower, pelvic or abdominal. Also, the male organ.

MEMBRUM GENITALE VIRORUM, *Penis*—m. *Mutieb're*, *Clitoris*—m. *Seminalis*, *Penis*—m. *Virile*, *Penis*.

MEMOIRE, *Memory*—m. *Perte de*, *Amnesia*.

MEMORIA DELETA, *Amnesia*.

MEMORY, *Memo'ria*, *Mnemē*, *Recordatio*, *Recorda'tio*, (F.) *Mémoire*. The cerebral faculty, by virtue of which past impressions are recalled to the mind.

MEMPHITES LAPIS. A sort of stone, found anciently in the environs of Memphis; which was esteemed narcotic.

MENAGOGUES, *Emmenagogues*.

MENDESION, *Ægyptiacum*.

MENE, *Moon*.

MENECRATIS ACCIPITER, *Accipiter*.

MENELCO'SIS, *Helcome'nia*, from *μην*, 'a month,' and *ἐλκωσις*, 'ulceration.' Vicarious menstruation from an ulcer.

MENES, *Menses*.

MÉNESPAUSIE, see *Menses*.

MENIDROSIS, from *μην*, 'a month,' and *ἰδρωσις*, 'sweating.' A sweating of menstrual blood.—Bock.

MENFRIGE, see *Pistacia lentiscus*.

MENINGARTHROCAÏE, *Inflammat'io membrana'rum articulatio'nis*, *Arthromeningi'tis*, from *μηνίγξ*, *μηνίγγος*, 'membrane,' *αρθρον*, 'a joint,' and *κακος*, 'disease.' Inflammation of the membranes of a joint.

MÉNINGE, *Dura mater*.

MENINGÉ, *Meningeal*.

MENINGEAL, *Menin'geus*, from *μηνίγξ*, *μηνίγγος*, 'a membrane.' (F.) *Méningé*. Relating to the meninges, or merely to the dura mater.

MENINGEAL APOPLEXY, see *Apoplexy*.

MENINGEAL ARTERY, MIDDLE, *Arteria menin'-gea media* seu *Sphæno-spino'sa* seu *Dura Matris media* max'ima seu *Sphæno-spinalis*, *Spiral Ar-*

tery, (F.) *Artère méningée moyenne* ou *Sphéno-épineuse*, ou *Épineuse*, is the largest of the branches given off by the internal maxillary artery. It enters the cranium by the foramen spinale, and distributes its branches chiefly to the dura mater.

MENINGEAL VESSELS, *Vasa menin'gea*. The vessels of the membranes of the brain.

MENIN'GES, *Eilan'tides*, *Matres seu Involu'-era Cer'ebr'i*, *Omen'ta seu Velamen'ta cerebra'lia*, from *μηνίγξ*, *μηνίγγος*, 'a membrane.' The three membranes which envelop the brain—*Dura mater*, *Arachnoid*, and *Pia mater*.

MÉNINGETTE, *Pia mater*.

MENINGITIS, *Meningitis*.

MENINGI'NA, (F.) *Méninge*. Same etymon. A name given by Chaussier to the pia mater, united to the cerebral layer of the arachnoid.

MENINGINA, INNER LAMINA OF THE, *Pia mater*.

MÉNINGINE, *Meningina*.

MÉNINGITE, *CÉRÉBRO-SPINAL*, *Meningitis*, *cerebro-spinal*.

MENINGINITIS, from *meningina*, and *itis*, denoting inflammation. Inflammation of the meningina.

MENINGION, *Arachnoid membrane*.

MENINGIS CUSTOS, *Meningophylax*.

MÉNINGITE, *Meningitis*—m. *Granulose*, *Hydrocephalus internus*—m. *Rachidienne*, *Meningitis*, *spinal*—m. *Tuberculeuse*, *Hydrocephalus internus*.

MENINGITIS, *Meningi'tis*, *Encephali'tis peripher'ica seu membrano'sa*, (F.) *Inflammation de la membrane séreuse céphalo-rachidienne*, *Méningite*, from *meninges*, and *itis*, denoting inflammation. Inflammation of the meninges or membranes of the brain. See *Phrenitis*.

MENINGITIS ARTHRITICA, *Cephalagra*.

MENINGITIS, *CÉRÉBRO-SPINAL*, *Myeli'tis*, *petechia'lis*, *Cer'ebral* or *cerebro-spinal typhus*, (F.) *Méningite cérébro-spinal*, *Fièvre cérébro-spinal*, *Typhus cérébro-spinal*. An alarmingly fatal form of epidemic fever, which prevailed in different countries of Europe, and, also, in certain portions of the United States, near the middle of the 19th century. It is considered, by Boudin, to constitute the cerebro-spinal form of the nosological group of typhus.

MENINGI'TIS GRANULAR, *Hydrocephalus internus*—m. *Hydrocephalica*, *Hydrocephalus internus*—m. *Infantum*, *Hydrocephalus internus*—m. *Membranosa*, *M. spinal*.

MENINGITIS MESENCOPHALICA, *Inflammation of the meninges of the medulla oblongata and pons varoli more especially*.

MENINGITIS PERIPHERICA, *M. spinal*—m. *Potatorum*, *Delirium tremens*—m. *Rachidian*, *M. spinal*—m. *Scrophulous*, *Hydrocephalus internus*.

MENINGITIS, *SPINAL*, *M. Rachid'ian* seu *Spina'lis seu peripher'ica seu membrano'sa*, *Perimyeli'tis*, *Myelo-meningi'tis*, (F.) *Méningite rachidienne*. Inflammation of the meninges of the spinal marrow.

MENINGITIS, TUBERCULAR, see *Hydrocephalus internus*—m. *Tuberculosa*, *Hydrocephalus internus*.

MENIN'GIUM. Diminutive of *μηνίγξ*, *μηνίγγος*, 'a membrane.' A fine, delicate membrane. The tunica arachnoidea of the brain.

MENINGO-CEPHALI'TIS, *Meningo-encephali'tis*, *Ceph'alo-meningi'tis*, *Enceph'alo-meningi'tis*, from *μηνίγξ*, *μηνίγγος*, 'a membrane,' *κεφαλη*, 'head,' and *itis*, a suffix denoting inflammation. Inflammation of the membranes and brain.

MENINGO-GASTRIC FEVER, *Fever, gastric*.

MENINGOGASTRITIS, *Fever, gastric*.

MENINGOMALACIA, from *μηνίγξ*, *μηνίγγος*,

'membrane,' and *μαλακία*, 'softening.' Softening of membranes in general.

MENINGOPHYLAX, *Custos mening'ia*, *Depressorium*, (F.) *Dépresseoire*, from *μηνίξ*, *μηνίγος*, 'a membrane,' the dura mater; and *φύλαξ*, 'a preserver.' An instrument for depressing the dura mater, and guarding it from injury, whilst the bone is cut or rasped.

MENINGORRHŒA, from *μηνίξ*, *μηνίγγος*, 'membrane,' and *ῥεω*, 'I flow.' Effusion of blood upon or between the membranes of the brain.

MENINGO'SIS. Same etymon. The union of bones by means of membrane; e. g. the articulation of the bones of the cranium in the fœtus. Meningosis is a variety of Syndesmosis.

MENINGOSYMPHYSIS; *Meningosyn'ia*, from *μηνίξ*, *μηνίγγος*, 'membrane,' and *συνφύσις*, 'growing together.' Adhesion of or by membrane. Adhesion of the membranes of the brain to each other, or to the brain.

MENINGOSYZIA, *Meningosymphysis*.

MENINGURIA; from *μηνίξ*, 'a membrane,' and *ουρον*, 'urine.' The passage of urine containing membranous shreds.

MENINX, Membrane—m. Choroides, Pia mater—m. Crassa, Dura mater—m. Dura, Dura mater—m. Exterior, Dura mater—m. Interior, Pia mater—m. Media, Arachnoid membrane—m. Pacheia, Dura mater—m. Sclera, Dura mater—m. Serosa, Arachnoid membrane—m. Vasculosa, Pia mater.

MENISCHEMIS, see *Amenorrhœa*.

MENISCI, Cartilages, interarticular.

MENISPERMUM ANGULATUM, *M. Canadense*.

MENISPERMUM CANADENSE, *M. angulatum*, Ord. *Menispermaceæ*, from *μην*, 'the moon,' and *σπερμα*, 'seed,' in allusion to the crescentic shape of the seed. A climbing plant, growing in various parts of the United States. Said to be used in Virginia as a substitute for sarsaparilla in scrofula, and to be called, in the West, *Sarsaparilla*. It is an excitant tonic.

MENISPERMUM COCCULUS, *M. glaucum*, *Anamirta Cocculus* seu *paniculata*, *Cocculus suberosus*. The systematic name of the plant, the berries of which are well known by the name of *Cocculus Indicus* seu *suberosus*, *Indian Berries*, *Bacca Piscatoria*, *Indian Cockles*, *Coccus Indicus*, *Coccula officinalis*, *Cocci orientales*, *Grana Orientis*, (F.) *Cogue du Levant*, *C. levant*. These berries are remarkable for their inebriating and destructive quality to fish. The deleterious principle appears to reside in the kernel, and in the active principle called *Picrotoxine*. *Cocculus Indicus* has been used in decoction to kill vermin. It has, at times, been added to beer, by fraudulent tradesmen, to render it more inebriating.

MENISPERMUM GLAUCUM, *M. cocculus*—m. *Palatum*, *Calumba*.

MENISPERMUM TUBERCULATUM, *M. Verrucosum*, *Cocculus crispus*. A Chinese plant, an extract from the root of which is tonic to the stomach and bowels, and possesses some astringency. Dose, 5 to 10 grains.

The same may be said of *Menispermum Cordifolium*—the *Golmucha*, of Bengal.

MENISPERMUM VERRUCOSUM, *M. tuberculatum*.

MENOCRYPHIA, *Amenorrhœa*.

MENOLIPSIS, see *Menses*.

MENOPAUSIS, see *Menses*.

MENOPHANIA, see *Menses*.

MENOPLANIA, Menstruation, vicarious.

MENORRHAG'IA, from *μην*, *μηνος*, 'a month,' and *ρρῡνναι*, 'to flow fiercely.' Flow of the menses. Frequently, the word is used synonymously with uterine hemorrhage or *metrorrhagia*, or for immoderate flow of the menses—*profuse menstrua-*

tion, *Parame'nia superflua* seu *profusa*, *Menorrhag'ia rubra*, *Catamenio'rum fluxus immodicus*, *Menstrua immod'ica* seu *superflua*, *Menorrh'ia*.

MENORRHAGIA, Menstruation, *Metrorrhagia*—m. Alba, Leucorrhœa—m. Difficilis, Dysmenorrhœa—m. Erronea, Menstruation, erronea—m. Stillatitia, Dysmenorrhœa.

MENORRHŒA, from *μην*, *μηνος*, 'a month,' and *ῥεω*, 'I flow.' The ordinary flow of the menses. Also, too long continuing, or too often returning, menstruation. *Menorrhagia*.

MENOSCHESIS, see *Amenorrhœa*.

MENOSTASIA, *Amenorrhœa*.

MENOSTASIS, from *μην*, *μηνος*, 'month,' and *στασις*, 'stasis,' 'stagnation.' This word, according to some, signifies the retention of the menses and their accumulation in the uterus. According to others, it means the acute pain which, in some females, precedes each appearance of the menses: a pain which has been presumed to proceed from the stasis of blood in the capillary vessels of the uterus.

MENOXEN'IA, *Menoxeno'sis*, from *μην*, *μηνος*, 'a month,' and *ξενος*, 'foreign.' Irregular menstruation.

MENOXENOSIS, *Menoxenia*.

MENS, Anima, Intellect.

MENSA, *Trop'æa*, 'a table.' The upper superficial part or corona of the jaw teeth.

MENSES, (pl. of *menstris*, 'a month;') *M. Mulieb'ria*, *Menses*, *Fluvius men'struis* seu *luna'ris*, *Profluvium mulieb're* seu *genita'le mulieb're*, *Consuetudo menstrua*, *Catame'nia*, *Men'strua*, *Emme'nia*, *Tributum luna're* seu *men'struum*, *Purgatio'nes*, *P. menstrua*, *Gynæcei'a*, *Aphedri'a*, *Cursus men'struis*, *Hor'a*, *Luna'ria*, *Courses*, *Menstrual flux*, *Monthly courses or periods*, *Flowers*, *Turns*, *Terms*, the *Reds*, *Troubles*, (F.) *Mois*, *Règles*, *Lunes*, *Flueurs*, *Affaires*, *Époques*, *Ordinaires*, *Purgations*, *Fleurs*, *Menstrues*. The sanguineous evacuation from the uterus, *Sanguis men'struis*, *Men'strual Blood*, the monthly occurrence of which constitutes menstruation or monthly illness. The first appearance of the menses—*menophania*—is usually preceded by the discharge of a fluid whitish matter from the vagina; by nervous excitement, and by vague pains and heaviness in the loins and thighs; numbness of the limbs, tumefaction and hardness of the breasts, &c. More or less indisposition and irritability also precede each successive recurrence of the *menstrual flux*. In temperate climates, each period, *men'strual period*, *periodus luna'ris*, ordinarily continues from three to six days; and the quantity lost varies from four to eight ounces. The menses continue to flow from the period of puberty till the age of 45 or 50. At the term of its natural cessation, *Menolip'sis*, *Ménopause* (Gardanne), *Menopau'sis*, the flux becomes irregular; and this irregularity is occasionally accompanied with symptoms of dropsy, glandular tumours, &c., constituting the *Parame'nia cessatio'nis* of Good, and what is called the *critical time* or *turn* or *change of life*; yet it does not appear that the mortality is increased by it. With the immediate causes of menstruation we are unacquainted. We express only our ignorance, when we assert it to depend upon periodicity; the discharge comes from the vessels of the uterus and vagina, and differs from ordinary blood by its peculiar odour, and by its not coagulating. It is evidently connected with the condition of the ovaries, and—it has been supposed—with the periodical discharge of ova from them. It is arrested, as a general principle, during pregnancy and lactation. In warm climates, women usually begin to menstruate early, and cease sooner than in the temperate regions. The quan-

tity lost is also greater. In the colder regions, the reverse of this holds as a general rule.

Dried menstrual blood, especially of a woman in her first menstruation—*Sanguis virginis primum menstruale*—was formerly given internally as an anti-epileptic.

MENSES ALBI, Leucorrhœa—m. Anomalæ, Paramenia—m. Devii, Menstruation, vicarious—m. Dolorifici, Dysmenorrhœa—m. Retention of the, see Amenorrhœa—m. Suppression of the, Amenorrhœa.

MENSIIUM RETENTIO, see Amenorrhœa.

MENTRUA, Menses—m. Alba, Leucorrhœa—m. Difficilia, Dysmenorrhœa—m. Dolorosa, Dysmenorrhœa—m. Immodica, Menorrhagia—m. Superflua, Menorrhagia.

MENTRUAL, Catamenial—m. Flux, Menses—m. Period, see Menses.

MENTRUANT, *Menstruans*, *Menstruat*, from *menstrua*, 'the catamenia.' One subject to the catamenia.

MENTRUSTATIO ANOMALA, Paramenia—m. Difficilis, Dysmenorrhœa—m. Dolorifica, Dysmenorrhœa—m. Impedita, see Amenorrhœa—m. Per insolitas Vias, Menstruation, vicarious.

MENTRUSTATIO PRÆCOX, Premature menstruation. Menstruation occurring before the usual age or time.

MENTRUSTATIO RECIDI'VA. Menstruation, when protracted beyond the usual age.

MENTRUSTATIO RETENTA, Amenorrhœa, Paramenia—m. Suppressa, Amenorrhœa, Paramenia.

MENTRUSTATION, *Menstruat*, *Chronogynia*, *Menorrhagia*, *Fluxus menstrualis*. The flow of the menses.

MENTRUSTATION DIFFICILE, Dysmenorrhœa—m. Difficult, Dysmenorrhœa—m. Laborious, Dysmenorrhœa—m. Premature, Menstrustatio præcox—m. Profuse, Menorrhagia.

MENTRUSTATION, VICA'RIOUS, is that which occurs from other parts than the uterus. It is called *Paramenia erronea*, *Menorrhagia erronea*, *Mensum per aliæna loca excretio*, *M. devii*, *Aberratio mensium seu menstruum*, *Menstrustatio per insolitas vias*, *Hæmatoplasia seu Hæmatoplasia menstrualis*, *Menoplasia*, (F.) *Déviation des Règles*, *Règles déviées*. At times, the secreted fluid has all the characters of the menstrual secretion; at others, it appears to be mere blood.

MENTRUSTATUS, Menstruant.

MENTRUEL, Catamenial.

MENTRUES, Menses.

MENTRUSTUOUS, Catamenial. Also, one affected with the menses or catamenia.

MENTRUM, (F.) *Menstrue*. Same etymon;—the menstruum being, of old, usually continued in action for a month. The monthly discharge of the female. The plural, *menstrua*, is now, however, always used for the menses. The name was formerly given to every substance which possesses the property of dissolving others slowly and with the aid of heat. At present, it is used synonymously with solvent. Thus—water, alcohol, ether, acids, oils, &c., are menstrua.

MENTRUM AURI, Nitro-muriatic acid—m. Ventriculi, Gastric juice.

MENSURA MEDICA. A measure of 48 ounces.

MENSURATIO, *Meta*, from *mensura*, 'measure.' *Measurement*, *Medical measuring*. One of the means used for exploring the state of the thoracic and other cavities. It consists in a comparative measurement of each side of the chest, by means of a riband extended from the median line of the sternum to the spine. When effusion exists on one side, the measurement is usually greater than on the other. When, on the

other hand, a portion of a lung has ceased, for some time, to exert its respiratory functions, the corresponding side of the chest becomes smaller, in consequence of the contraction of the pleura, retraction of the pulmonary tissue, and greater approximation of the ribs.

MENTA, *Mentha*, Penis.

MENTAGRA, Mentulagra, Sycosis—m. Infantum, Porriga lupinosa.

MENTAGRAPHYTE, *Men'tagrophyte*, *Microsporon mentagrophytes*, from *mentagra*, and *groy*, 'a plant.' A name proposed by M. Gruby, of Vienna, for a cryptogamous plant, which he considers to produce mentagra.

MENTAL, *Menta*, from *mens*, *mentis*, 'mind'; belonging or relating to the mind, as *mental phenomena*, *mental diseases* (*Morbi mentales*), &c.

In Anatomy, Mental, *Menta*, (F.) *Mentonnier*, signifies that which relates to the chin; from *mentum*, 'the chin.'

MENTAL ARTERY, (F.) *Artère mentonnière*. This is given off by the inferior dental, and issues at the mental foramen to be distributed on the lower lip.

MENTAL FORAMEN, (F.) *Trou mentonnier*, *Anterior maxillary F.*, is the outer orifice of the inferior dental canal. It is situated on the outer surface of the lower jaw-bone, opposite the second incisor, or the canine tooth, and gives passage to the vessels and nerves.

MENTAL FOSSA, see Fossa, mental.

MENTAL NERVE, (F.) *Nerf mentonnier*, is furnished by the inferior dental nerve. It issues by the mental foramen, and is distributed, in numerous filaments, to the muscles of the lower lip.

MENTAL REGION, *Re'gio mentalis*. The region of the chin.

MENTHA seu MENTA AQUATICA, (μνθρ, μνθα, μνθος, 'mint.') *M. palustris* seu *hirsuta* seu *rotundifolia palustris*, *Menthastrium*, *Sisymbrium menthastrium*, *Balsamum palustre*, *Watermint*, (F.) *Menthe aquatique*, *M. rouge*. *Nat. Ord.* Labiatae. This is less agreeable than the *Mentha viridis*, and more bitter and pungent. It is used like spearmint.

MENTHA BALSAMEA, *M. piperita*.

MENTHA CAPEN'SIS, which grows at the Cape of Good Hope, possesses the medical properties of the mints.

MENTHA CERVINA, *Hart's pennyroyal*, *Hyssop-leaved mint*, *Pulegium cervinum*, (F.) *Menthe cervine*. Possesses the properties of pennyroyal, but is very unpleasant.

MENTHA CITRATA, *Melissa*.

MENTHA CRISPA, *M. hircynica*, *Colymbifera minor*, *Curled Mint*, (F.) *Menthe frisée* on *crêpe*, *Baume d'eau à feuilles ridées*. Possesses the properties of peppermint.

MENTHA GENTILIS, *M. viridis*—m. *Hercynica*, *M. crispa*—m. *Hirsuta*, *M. Aquatica*—m. *Lævigata*, *M. viridis*—m. *Officinalis*, *M. piperita*—m. *Palustris*, *M. Aquatica*.

MENTHA PIPERITA, *M. Piperitis* seu *officinalis* seu *balsamea*, *Peppermint*, (F.) *Menthe poivrée*. The odour of this variety is strong and agreeable. Taste, pungent, aromatic, and producing a sensation of coldness in the mouth. Virtues depend upon essential oil and camphor. It is stomachic and carminative. It is chiefly used in the form of essential oil—the *oleum menthae piperitæ*.

Essence of Peppermint consists of *ol. menthae pip.* 3ij, *sp. vin. rectif.*, coloured with *spinach leaves*, Oij. See *Tinctura olei menthae piperitæ*.

MENTHA PULEGIUM, *Pulegium*, *P. regale* seu *latifolium*, *Glechon*, *Pennyroyal*, *Pudding grass*, (Prov.) *Hilwort*, *Lurkey-dink*, *Organ*, (F.) *Pouliot*. The oil—*O'leum pulegii*—possesses the virtues of the plant. It resembles the mints in

general in properties. Amongst the vulgar, it is esteemed an emmenagogue. *AQUA PULEGII*, *Aqua Mentha pulegii*, *Pennyroyal Water*, is official in the Edinburgh and Dublin Pharmacopœias.

MENTHA ROMANA, *Balsamita suaveolens*, *Tanacetum balsamita*—*m. Rotundifolia palustris*, *M. aquatica*—*m. Saracenicæ*, *Balsamita suaveolens*, *Tanacetum balsamita*—*m. Sativa*, *M. viridis*—*m. Spicata*, *M. viridis*.

MENTHA VIRIDIS, *M. Sati'va seu vulgari's seu gentilis* seu *spicata* seu *lævigata*, *Spearmint*, (*F.*) *Mentha sauvage* ou *verte* ou *des jardins*, *Baume des jardins*. Odour strong, aromatic; taste warm, austere, bitterish. Virtues the same as those of the peppermint. The *Oleum mentha viridis* is obtained from it.

MENTHA VULGARIS, *M. viridis*.

MENTHASTRUM, *Mentha aquatica*.

MENTHA AQUATIQUE, *Mentha aquatica*—*m. Cervine*, *Mentha cervina*—*m. Cog*, *Tanacetum balsamita*—*m. Crêpe*, *Mentha crispa*—*m. Frisée*, *Mentha crispa—m. des Jardins*, *Mentha viridis*—*m. Poirée*, *Mentha piperita*—*m. Romaine*, *Balsamita suaveolens—m. Rouge*, *Mentha aquatica—m. Sauvage*, *Mentha viridis—m. Verte*, *Mentha viridis*.

MENTIGO, *Porrigio larvalis*, *Sycosis*.

MENTISME, from *mens, mentis*, 'mind.' Any irregular movement of the mind, whether from emotion or a vivid imagination.—*Baumes*.

MENTOBICORNEUS, *Geniobhyoides*.

MENTO-LABIAL, *Mento-labialis*, from *mentum*, 'the chin,' and *labium*, 'a lip.' Under this name Chaussier has united the Levator labii inferioris, and Quadratus muscles. See Depressor labii inferioris.

MENTO-LABIAL FURROW: a transverse depression situate between the lower lip and the chin, which is remarkable for the perpendicular direction of the hairs growing upon it.

MENTON, *Mentum*.

MENTONNIER, *Mental*—*m. Labial*, *Depressor labii inferioris*.

MENTONNIÈRE, see *Funda*.

MENTULA, *Penis*—*m. Muliebris*, *Cereosis*, *Clitoris*—*m. Parva*, *Pipinna*.

MENTULAGRA, *Men'tagra*. A hybrid word, from *mentula*, 'the penis,' and *aggra*, 'a seizure.' A convulsive erection of the mentula or penis; such as is said to be sometimes observed in eunuchs. See *Pudendagra*.

MENTULATUS, *Membrosus*.

MENTUM, *Geni'um*, *Geni'um*, *Anthe'reon*, 'the chin.' (*F.*) *Menton*. The inferior and middle part of the face, situate below the lower lip. The chin is formed of skin, areolar tissue, muscles, vessels, nerves, and the os maxillare inferius.

MENYAN'THES TRIFOLIA'TA, *Menyan'thes*, *Minyan'thes*, from *μνν*, 'a month,' and *ανθος*, 'a flower,' because it keeps in flower about a month [?]. *Trifolium paludosum* seu *aquaticum* seu *palustris* seu *fibris'um*, *Water trefoil*, *Buckbean*, *Bogbean*. *Nat. Ord.* *Gentianacæ*. (*F.*) *Trèfle d'eau* ou *des Marais* ou *de Castor*. The taste is intensely bitter, and is extracted by water. It is tonic, anthelmintic, diuretic, cathartic, and, in large doses, emetic. In some countries, it is used as a substitute for hops in making beer.

MENYAN'THES Verna, *American buckbean*, *Marsh trefoil*, *Water shamrock*, *Bitter root*, has similar properties.

MEOSIS, *Meiosis*.

MEPHITIC, *Mephitis'icus*, (*F.*) *Méphitique*; from a Syriac word, signifying 'to blow, to breathe.' Any thing possessed of an unwholesome property; chiefly applied to exhalations.

MEPHITIC ACID, *Carbonic acid*—*m. Air*, *Carbonic acid*.

MEPHITIS, see *Mephitism*—*m. Hepatica*, *Hydrogen sulphuretted—m. Indammabilis*, *Hydrogen—m. Urinosa*, *Ammonia*.

MEPHITISM, *Mephitis'm*, (*F.*) *Méphitisme*. Same etymon. Any pernicious exhalation. *Mephitis* and *Mophe'ta*, (*F.*) *Mofette* ou *Moufette*, are old names for azote; and, in general, for all exhalations and gases that are unfit for respiration.

MERACUS, *Merus*.

MERAMAUROSIS, *Hypamaurosis*.

MERANÆSTHESIS, *Meranæsthes'ia*, from *μερος*, 'a part,' *av*, 'privative,' and *αισθησις*, 'sensation.' Insensibility of a part of the body.

MERATROPHIA, from *μερος*, 'a part,' and *αρροφια*, 'want of nourishment.' Atrophy of some part of the body.

MERCURE, *Hydrargyrum*—*m. Bichlorure de*, *Hydrargyri oxyurias—m. Biiodure de*, *Hydrargyri iodium rubrum—m. Bromures de*, see *Bromine—m. Coralli*, *Hydrargyri nitrico-oxylum—m. avec la Craie*, *Hydrargyrum cum cretâ—m. Cru*, *Hydrargyrum—m. Cyanure de*, *Hydrargyri cyanuretum—m. Deutiodure de*, *Hydrargyri iodium rubrum—m. Deuto-chlorure de*, *Hydrargyri oxyurias—m. Doux*, *Hydrargyri suburias—m. Gommeux de Plenck*, *Syrupus de mercurio mediante gummi—m. Muriate oxygénée de*, *Hydrargyri oxyurias—m. Oxide de, cendré*, *Hydrargyri oxydum cinereum—m. Oxide gris de*, *Hydrargyri oxydum cinereum—m. Oxide nitrique de*, *Hydrargyri nitrico-oxylum—m. Oxide noir de*, *Hydrargyri oxydum cinereum—m. Oxide rouge de*, *Hydrargyri oxydum rubrum—m. Périodure de*, *Hydrargyri iodium rubrum—m. Protiodure de*, *Hydrargyri iodium—m. Protochlorure de*, *Hydrargyri suburias—m. Protoxide de*, *Hydrargyri oxydum cinereum—m. Sous-muriate de*, *précipité*, *Hydrargyrum præcipitatum—m. Sous-sulfate de*, *Hydrargyrum vitriolatus—m. Sulfure de, noir*, *Hydrargyri sulphuretum nigrum—m. Sulfure de, rouge*, *Hydrargyri sulphuretum rubrum*.

MERCURIAL, *Mercurialis*, (*F.*) *Mércuriel*; from *mercurius*, 'mercury.' That which contains or relates to mercury. A preparation of mercury—*Hydrargyricum*, *Hydrargyrium*.

MERCURIAL ERETHISM, see *Erethism*—*m. Rash*, *Eczema mercuriale*.

MERCURIALE ANNUELLE. *Mercurialis—m. des Bois*, *Mercurialis perennis—m. de Chien*, *Mercurialis perennis—m. Vivace*, *Mercurialis perennis*.

MERCURIALIS, *Mercurialis*, (*F.*) *Mercuriale annuelle*, *Foivole*, *Ramberge*, *Vignoble*; *Class.* *Diœcia*; *Order*, *Enneandria*. It has been esteemed cathartic, hypnotic and cosmetic, and has been chiefly used by way of lavement. See *Chenopodium Bonus Henricus*.

MERCURIALIS HUMOR, *Humor melancholicus*. A supposititious humour with the older physicians, out of which they presumed that morbi melancholici seu mercuriales arose.

MERCURIALIS PERENNIS, *Cynocrambæ*, *Canina bras'ica*, *M. montana*, *M. Sylvestris*, *Dog's mercury*, (*F.*) *Mercuriale vivace*, *M. de chien*, *M. des bois*. This plant is possessed of acro-narcotic properties.

MERCURIALISMUS, *Hydrargyriasis*.

MERCURIALIST, *Mercurialis'ta*, from *mercurius*, 'mercury.' One inordinately addicted to prescribing mercury.

MERCURIALIZATION. Same etymon. The state of being affected by mercury. One under the influence of mercury is said to be mercurialized.

MERCURII CALX VITRIOLATA, *Hydrargyrum vitriolatus—m. Deuto-ioduretum*, *Hydra*

gyri iodium rubrum—m. Proto-ioduretum, Hydrargyri iodium—m. Sperma, Hydrargyrus acetatus—m. Terra foliata, Hydrargyrus acetatus.

MERCURIUS, Hydrargyrum—m. Acetatus, Hydrargyrus acetatus—m. Alkalisatus, Hydrargyrum cum cretâ—m. Calcinatus, Hydrargyri oxydum rubrum—m. Causticus flavus, Hydrargyrus vitriolatus—m. Cinereus, Hydrargyri oxydum cinereum—m. Cinnabarinus, Hydrargyri sulphuretum rubrum—m. Corallinus, Hydrargyri nitricco-oxydum—m. Corrosivus, Hydrargyri oxymurias—m. Corrosivus ruber, Hydrargyri nitricco-oxydum—m. Corrosivus sublimatus, Hydrargyri oxymurias—m. Cosmeticus, Hydrargyrum præcipitatum—m. Dulcis, Hydrargyri submurias—m. Emeticus flavus, Hydrargyrus vitriolatus—m. Gummosus Plenkii, Syrupus de mercurio mediante gummi—m. Luteus, Hydrargyrus vitriolatus—m. Mortis, Algaroth—m. Niger Moscati, Hydrargyri oxydum cinereum—m. Nitrosus calide paratus, see Hydrargyri nitras—m. Nitrosus frigore paratus, Hydrargyri nitras—m. Præcipitatus corrosivus, Hydrargyri nitricco-oxydum—m. Præcipitatus niger, Hydrargyri oxydum cinereum—m. Saccharatus, Hydrargyrum saccharatum—m. Solubilis of Hahnemann, see Hydrargyri oxydum cinereum—m. Sublimatus dulcis, Hydrargyri submurias—m. cum Sulphure, Hydrargyri sulphuretum nigrum—m. Terrestris, Polygonum Hydropiper—m. Vitæ, Algaroth—m. Zoticus Hartmanni, Hydrargyri submurias.

MERCURY, Hydrargyrum—m. Acetate of, Hydrargyrus acetatus—m. Acid nitrate of, see Hydrargyri nitras—m. Ammoniated, Hydrargyrum præcipitatum—m. Ammonio-chloride of, Hydrargyrum præcipitatum—m. Bichloride of, Hydrargyri oxymurias—m. Biniodide of, Hydrargyri iodium rubrum, see Iodine—m. Bisulphuret of, Hydrargyri sulphuretum rubrum—m. Bromide of, see Bromine.

MERCURY WITH CHALK, Hydrargyrum cum cretâ.

MERCURY, CHLORIDE OF, MILD, Hydrargyri submurias—m. Compound cerate of, Ceratum Hydrargyri compositum—m. Cyanuret of, Hydrargyri cyanuretum—m. Deuto-iodide of, see Iodine—m. Dog's, Mercurialis perennis—m. English, Chenopodium bonus Henricus—m. French, Mercurial—m. Iodide of, Hydrargyri Iodium—m. Iodide of, red, Hydrargyri Iodium rubrum.

MERCURY, IODIDE OF CHLORIDE OF, *Iod'idum Hydrar'gyri chlo'ridi*, *Hydrar'gyrum ioda'tum cum chlorido mercur'ii*, (F.) *Iodhydrargyrite de chlorure mercur'is*. Made by the reaction of iodine on mild chloride of mercury. Two iodides may be prepared—the iodide and the biniodide. Both are violent irritants: the biniodide, especially, is a powerful caustic. Both have been given in scrofula. The biniodide is only used externally like nitrate of silver in scrofulous and certain syphilitic ulcerations. An ointment of the iodide (gr. xv ad adipis ʒij) is rubbed on scrofulous tumefactions.

MERCURY, IODO-ARSENITE OF, Arsenic and Mercury, iodide of—m. Muriate of, corrosive, Hydrargyri oxymurias—m. Muriate of, mild, Hydrargyri submurias—m. Nitrate of, Hydrargyri nitras—m. Nitric oxyd of, Hydrargyri nitricco-oxydum—m. Oxide of, black or gray, Hydrargyri oxydum cinereum—m. Oxide of, red, Hydrargyri oxydum rubrum—m. Oxymuriate of, Hydrargyri oxymurias—m. Periodide of, Hydrargyri iodium rubrum—m. Phosphuretted, Hydrargyrus phosphoratus—m. Precipitate of, white, Hydrargyrum præcipitatum—m. Protochloride of, Hydrargyri submurias—m. Protoiodide of, see Hydrargyri iodium, and Iodine—m.

Prussiate of, Hydrargyri cyanuretum—m. Soluble, Hydrargyri nitras—m. Subchloride of, Hydrargyri submurias—m. Submuriate of, Hydrargyri submurias—m. Sulphuret of, black, Hydrargyri sulphuretum nigrum—m. Sulphuret of, red, Hydrargyri sulphuretum rebrum, Realgar—m. Sulphuret of, with Sulphur, Hydrargyri Sulphuretum nigrum—m. Supernitrate of, Solution of, see Hydrargyri nitras—m. Tartrate of, Hydrargyri tarttras—m. Three-seeded, Acalypha Virginica.

MERCURY AND ARSENIC. IODIDE OF, Arsenic and Mercury, Iodide of—m. and Quinia, Protochloride of, Hydrargyri et Quiniæ Protochloridum.

MERDA, Excrement.

MERDIVOMUS, Copremetus.

MERDUS, Excrement

MÈRE DES PIANs, Pian—m. de Seigle, Ergot.

MEREMPHRAX'IS, from *μερος*, 'a part,' and *σπραγξίς*, 'obstruction.' Obstruction or infarction of an organ.

MERICUS, Local.

MERIDRO'SIS, *Sudor parti'alis seu loca'lis*; from *μερος*, 'a part,' and *ἰδρωσις*, 'sweating.' A partial perspiration.

MERIMNA, Cura.

MERIMNE, Cura.

MERISIER, Prunus avium.

MERISMOPEDIA VENTRICULI, Sarcina ventriculi.

MEROBALANEUM, Bath, partial.

MEROBALNEUM, Bath, partial.

MEROCELE, *Mirocel'is*, *Meror'is*, *Meror-rher'is*, from *μπος*, 'the thigh,' and *κῆλη*, 'tumour,' *Hernia crura'lis seu femora'lis*, *Fem'or'al* or *crural hernia*, (F.) *Hernie crurale ou fémorale*. In this hernia, the viscera issue from the abdomen through the crural canal; or through an opening immediately on the outer side of Gimbernat's ligament, which gives passage to lymphatic vessels. This affection is more frequent in the female than in the male, and especially in those who have had children. The tumour, formed by merocele, is generally small, round, and more or less difficult of reduction. In other respects, it possesses all the characters of hernia. The neck of the sac has, close on the outside, the epigastric artery; above, the spermatic cord and spermatic artery in the male—the round ligament in the female; on the inside, Gimbernat's ligament; and below, the pubes. When the obturator artery arises from the epigastric, it generally passes on the outside of and below the orifice of the sac; sometimes, however, it takes a turn above, and then to the inside of the opening. J. Cloquet asserts, that of 134 cases, in one only did he find the epigastric artery on the inside of the orifice of the sac.

The operation, required in strangulation of this variety, may be practised on the aponeurotic opening, by cutting downwards and inwards on the side of Gimbernat's ligament.

MEROCOXYALGIA, Coxalgia.

MEROL'OGY, *Merolog'ia*, (F.) *Mérologie*, from *μερος*, 'a part,' and *λογος*, 'a discourse.' The branch of anatomy, which treats of the elementary constituents of the tissues and humours of the organism.—Robin and Verdeil.

MEROPIA, from *μερος*, 'a part,' and *ορασις*, 'vision.' A partial obscurity of vision.

MERORIXIS, Merocele.

MERORRHEU'MA, *Rheumatismus parti'alis seu loca'lis seu Membro'rum singulo'rum*; from *μερος*, 'a part,' and *ρευμα*, 'defluxion, rheumatism.' Rheumatism affecting a part. Topical or local rheumatism.

MERORRHESIS, Merocele.

MEROS, Femur, Thigh.

MERRY-ANDREW. An itinerant quack, who exposes his nostrums for sale at fairs and markets. See CHARLATAN.—So called from Dr. Andrew Boorde, who lived in the reigns of Henry VIII., Edward VI., and Queen Mary, and who was in the habit of frequenting fairs and markets, at which he harangued the populace.

MERTEN'SIA VIRGINICA, *Pulmonaria Virginica*, *Virginian Cowslip*, *Smooth Lungwort*; indigenous. Family, Boraginacæ. The plant is mucilaginous, and may be used as a demulcent.

MERULIUS, *Morchella Esculenta*—m. Auricula, *Peziza auricula*.

MERUS, *Mera'cus*, *A'cratos*, 'pure, genuine,' as *Vinum merum*, unmixed wine.

MERUS, Femur, Thigh.

MERVINE, MALE, *Cypripedium luteum*.

MERYCISMUS, Ruminantion.

MÉRYCOLE, see Ruminantion.

MERYCOLOG'IA, from *μῆρυκα*, 'I ruminate,' and *λογος*, 'a description.' Any work on ruminantion may be so termed.

MERYPERTROPH'IA, from *μερος*, 'a part,' and *hypertrophia*, 'supernutrition.' Hypertrophy of a part.

MESA, Genital organs.

MESARÆUM, Mesentery.

MESARAIC, Mesenteric.

MESEMAR, Mismar.

MESEMBRYANTHEMUM COPTICUM, see Soda.

MESEMBRYANTHEMUM CRYSTALLINUM, *M. fico'ides* (*μεμβρῖανον*, 'to do at noon,' because its flowers expand at midday.) *Di' amond fig-marygold*, *Ice-plant*, (F.) *Glaciale*, *Crystalline*. Nat. Ord. Ficoideæ: a plant common in the neighbourhood of Cape Town. It has been recommended in dysuria, ischuria, and some other affections of the urinary organs.

MESEMBRYANTHEMUM EDULÈ, and *M. acinaciforme*, South African plants, are much used as domestic remedies at the Cape. The expressed juice of the leaves acts as an astringent in dysentery, and as a mild diuretic. It is also used as an antiseptic gargle in malignant sore throat, violent salivation and aphthæ, and as a lotion to burns and scalds.

MESEMBRYANTHEMUM TORTUOSUM, also a South African plant, is said to possess narcotic properties.

MÉSENCÉPHALE, Medulla oblongata, Pons Varolii.

MES'ENTERIC, *Mesara'ic*, *Mesenter'icus*, *Mesara'icus*, *Mesara'icus*, (F.) *Mésentérique*, *Mésaraïque*; from *μεσεντεριον*, or *μεσαραιον*, 'the mesentery.' That which relates or belongs to the mesentery.

MESENTERIC ARTERIES are two in number, and distinguished into, 1. The *Superior mesenteric*, which arises from the anterior part of the abdominal aorta, below the coeliac. It immediately descends to the left, forwards, behind the pancreas, and in front of the third portion of the duodenum,—to gain the superior extremity of the mesentery, where it makes a long curvature, the convexity of which is turned forwards and to the left. Towards the termination of the ileum, it ends by anastomosing with a branch of the *A. colica dextra inferior*. At its concavity, it gives off the three *Arteria colica dextra*, which belong to the great intestine; and, at its convexity, it sends off fifteen or twenty branches, which are distributed to the small intestine, after having formed numerous arches by their anastomoses.

2. The *Inferior Mesenteric Artery* arises from the anterior part of the abdominal aorta, an inch

and a half before its termination. It descends, on the left side, behind the peritoneum; engages itself in the substance of the iliac mesocolon, forming a considerable curvature, whose convexity faces the left side. When it reaches the brim of the pelvis, it passes along the posterior separation of the mesorectum, and attains the neighbourhood of the anus, under the name *Superior hemorrhoidal artery*. It gives off no branch at its concave part; but, from the convex, the three *Arteria colica sinistra* arise.

MESENTERIC DISEASE, *Tabes mesenterica*.

MESENTERIC GLANDS are the lymphatic ganglions of the mesentery. Through them, the chyliferous vessels pass to the thoracic duct. Their uses are unknown. When diseased, nutrition is interfered with, and atrophy produced.

MESENTERIC HERNIA. If one of the layers of the mesentery be torn by a blow, whilst the other remains in its natural state, the intestines may insinuate themselves into the aperture, and form a kind of hernia. It is not known during life.

MESENTERIC PLEXUSES are furnished by the solar plexus, and have been distinguished into *superior* and *inferior*; like the mesenteric arteries which they accompany.

MESENTERIC or MESARAIC VEINS are two in number, and belong to the vena porta. They are distinguished into, 1. The *Superior mesenteric* or *mesaraic* or *great mesaraic*. This trunk receives, from above to below and on the right side, the three *vena colica dextra* and the *Gastro-epiploica dextra*. Into its left side, the veins of the small intestine open. It passes in front of the transverse portion of the duodenum; and, behind the pancreas, unites with the splenic vein to concur in the formation of the vena porta.

The *inferior* or *lesser mesenteric vein* corresponds to the artery of the same name, and opens into the splenic, near the union of that vein with the superior mesenteric, and behind the pancreas.

MÉSÉNTÉRIQUE, Mesenteric.

MESENTERIITIS, Mesenteritis.

MESENTERITIS, *Mesenterii'tis*, *Empres'ma periton'itis mesenter'ica*, *Enteri'tis mesenter'ica*, *Inflammation of the mesentery*; from *μεσεντεριον*, 'mesentery,' and *itis*, denoting inflammation. The pain is here deeper seated and more immediately in the mesenteric region. The external tenderness is less than in some of the other varieties of peritonitis.

MESENTERIUM, Mesentery—m. Crassum, Mesocolon.

MESENTERON, Mesentery.

MESENTEROPHTHISIS, *Tabes mesenterica*.

MES'ENTRY, *Mesenter'ium*, *Mesara'ion*, *Mesara'um*, *Media'num*, *Membra'na pinguis intestino'rum*, *Lactes*, *Mesen'teron*, *Mesera'um*, *Mesora'um*, *Medium intesti'num*, *Epichor'dis*; from *μεσος*, 'in the middle,' and *εντερον*, 'intestine,' (F.) *Mésentère*. A term in anatomy, applied to several duplicatures of the peritoneum, which maintain the different portions of the intestinal canal in their respective situations; allowing, however, more or less motion. They are formed of two laminae, between which are contained the corresponding portion of intestine and the vessels that pass to it. One only of these duplicatures has received the name *mesentery*, properly so called. This belongs to the small intestine, which it suspends and retains *in situ*. Its posterior margin, which is the smallest, is straight, and descends obliquely from the left side of the body of the second lumbar vertebra to the right iliac fossa. Its anterior margin is curved, undulating, plaited, and corresponds to the whole length of the small intestine. The mesentery contains, between the two laminae

which form it, a number of lymphatic ganglions; the trunks and branches of the mesenteric vessels; the nervous plexuses accompanying them, and many lacteals and lymphatics.

MESERA. Tutia.

MESERÆUM, Mesentery.

MESERAION, Mesentery.

MESIAL, from *μεσος*, 'in the middle.' Relating or appertaining to the middle.

MESIAL PLANE, an imaginary plane, dividing the head, neck, and trunk into similar halves, towards right and left. Every aspect towards this plane is *mesial*; and every aspect towards right or left is *lateral*; every lateral aspect being *dextral* or *sinistral*.

MESIAMUM, Aniceton.

MESMER'IC, *Mesmer'icus*; from Anthony Mesmer, the great promulgator of animal magnetism. Relating to mesmerism or animal magnetism:—as the '*mesmeric state*,' '*mesmeric sleep*,' &c. &c.

MESMERISM, Magnetism, animal.

MESMERIST. A practiser of, or believer in, mesmerism.

MESMERIZA'TION, *Magnetiza'tion*. The act of mesmerizing. The state of being mesmerized.

MES'MERIZED, *Mesmeriza'tus*, *Mag'netized*. Affected with mesmerism or animal magnetism. When the person is in a state of '*magnetic sleep*,' he is said to be *hypnotized*.

MES'MERIZER, *Animal Magneti'zer*, *Mag-neti'zer*. One who practises mesmerism.

MESMERO-PHRENOLOGY, *Phreno-magnetism*.

MESO, *μεσος*, 'in the middle.' A prefix to certain words.

MESOBREGMATE. see Cranium.

MESOCÆ'CUM, (*meso* and *cæcum*). A name given to a duplicature of the peritoneum, (in some persons only), at the posterior part of the cæcum.

MESOCÉPHALE, *Medulla oblongata*, Pons Varolii.

MESOCHON'DRIAC, *Mesochondri'acus*; from *meso*, and *χονδρος*, 'cartilage.' A name given by Boerhaave to fleshy fibres situate between the cartilaginous rings of the trachea.

MESOCOL'IC, *Mesocol'icus*. Same etymon as the next. Belonging or relating to the mesocolon.

MESOCO'LON, *Mesocol'um*, *Mesenter'ium cras-um*, from *meso*, and *κωλον*, 'the colon.' A name given to the duplicatures of the peritoneum, which fix the different parts of the colon to the abdominal parietes. It has received different names, according to its situation. The *right lumbar mesocolon* fixes the ascending colon to the corresponding lumbar region. The *transverse mesocolon* arises from the concave arch of the colon, and forms a septum between the epigastric and umbilical regions. Its inferior portion is continuous with the mesentery. The *left lumbar mesocolon*, which contains the ascending colon, is continuous below with the *Riæc mesocolon*. The last includes between its layers the sigmoid flexure of the colon, and ends in the mesorectum. Under the right kidney, it is narrow and firm, and forms the *right ligament of the colon*: at the under end of the left kidney, it forms the *left ligament of the colon*.

MESOCRANIUM, Sinciput.

MESOCRANUM, Sinciput.

MESODERMUM, *Corpus mucosum*.

MESODME, Mediastinum.

MESODMI'TIS, from *μεσodem*, (*μεσος*, 'middle,' and *δωμος*, 'a building,') 'the mediastinum,' and *itis*, 'inflammation.' *Mesotechi'tis*, *Mediastinitis*. *Inflammatio mediastini*. Inflammation of the mediastinum.

MESOËPIDID'YMIS, from *μεσος*, 'middle,' and *ἐπιδιδυμς*, 'epididymis.' A double serous fold, like the mesocolon, formed by the tunica vaginalis passing from the body of the epididymis, and uniting it to the testicle.

MESOGASTRIUM, Umbilical region.

MESOGASTRIUM MÜLLERI. A portion of mesentery, which passes from the vertebral column to the left curvature of the stomach.

MESOGLOSSUS, Genioglossus.

MESOGLOTTUS, Genioglossus.

MESOGONIUM, Internodium.

MESOLO'BAR. Belonging or relating to the *Mésolobe* or *Corpus callosum*.

MESOLOBAR ARTERIES, *Arteriae mesolob'icæ*, *A. corporis callo'si cerebri*, are the arteries of the corpus callosum.

MÉSOLLOBE, *Corpus callosum*.

MESOMER'IA, from *meso*, and *μερος*, 'the thigh.' The parts of the body situate between the thighs and hips.

MESOMERION, Perinæum.

MESOMPHALUM, Umbilicus.

MESOPHLEB'IUM, from *meso*, and *φλεψ*, 'a vein;' *Intervenium*. The space between two veins.

MESOPH'RYON, *Glabella*, *Gabell'a*, *Intercil'ium*, *Metop'ium*, *Nasal eminence*, from *meso*, and *ὄφρυς*, 'the eyebrow.' The part between the eyebrows.

MESOPLEURIUM, Intercostal space.

MESOPLEURIUS, Intercostal.

MESORÆUM, Mesentery.

MESOR'CHIUM, from *meso*, and *ρχης*, 'a testicle.' A duplicature of the peritoneum, which supports the testicle in its passage from the abdomen into the scrotum.—Seiler.

MESORECT'UM. A hybrid word; from *meso*, and *rectum*, 'the intestine rectum.' A triangular reflection, formed by the peritoneum, between the posterior surface of the rectum and the anterior surface of the sacrum. Between the two layers of which the meso rectum is composed are found much areolar tissue, and the termination of the inferior mesenteric vessels.

MESOS, *Meso*.

MESOSCEOCLE. Hernia, perineal.

MESOSCEOLON, Perinæum.

MESOSCELOPHY'MA, from *Mesoscelon*, (*meso*, and *σκελος*, 'leg,') 'perinæum,' and *φυμα*, 'swelling.' A tumour of the perinæum.

MESOTH'ENAR, from *μεσος*, and *Sevap*, 'the thenar,' 'the palm of the hand.' A muscle, which carries the thumb towards the palm. Winslow applied the term to the abductor pollicis, united to the deep-seated portion of the flexor brevis pollicis.

MESOT'ICA, from *μεσος*, 'middle.' Diseases affecting the parenchyma. Pravity in the quantity or quality of the intermediate or connecting substance of organs; without inflammation, fever, or other derangement of the general health. The first Order, class *Ecritica* of Good.

MESOTECHITIS, Mesodmitis.

MESOTECHIUM, Mediastinum.

MESOTECHIUM, Mediastinum.

MES'PILUS, *Mespilus Germanica* seu *domestica*. Ord. Rosaceæ. The medlar. (F.) *Néflier*. The fruit (F.) *Nefle*, and seeds of the medlar have both been used medicinally:—the immature fruit as an astringent, and the seeds in nephritic diseases. See *Cratægus Aria* and *Sorbus aucuparia*.

MESPILUS ARIA, *Cratægus aria*—*m. Domestica*, *Mespilus*, *Sorbus domestica*—*m. Germanica*, *Mespilus*—*m. Intermedia*, *Mespilus oxyacantha*—*m. Levigata*, *Mespilus oxyacantha*.

MESPILUS OXYACANTHA. *M. Oxyacanthoides* seu *intermedia* seu *levigata*. *Oxyacantha*, *Spina alba*, *White Hawthorn*. (F.) *Aulepine*. The

flowers of this uncommon European plant are sometimes used in infusion as a pectoral.

MESPILUS OXYACANTHOIDES, *Mespilus oxyacantha*.

META, *μετα*, *Meth'*, *μεθ'*, 'with,' 'together with,' 'after,' 'change of form and place.' A common prefix to words. Hence:

METABASIS, *Transitus*, *Metab'ole*, *Metabol'ia*, *Metall'age*, *Metallax'ia*, from *μεταβαίνω*, (*meta*, and *βαίνω*, 'I go,') 'I digress.' A change of remedy, practice, &c.—Hippocrates.

METABOLE, *Metabasis*, Transformation.

METABOLELOGY, *Metabolelog'ia*; from *μεταβολή*, (*meta*, and *βύλλω*), 'change,' and *λογος*, 'a description.' A description of the changes which supervene in the course of a disease.

METABOLIA, *Metabasis*.

METABOLIC, *Metabol'icus*, from *μεταβολή*, 'change.' Appertaining to change or transformation.

METABOLIC FORCE. *Vis metabol'ica*. A term employed by Schwann for the power possessed by living cells of changing the character of the substances brought in contact with them.

METACARPAL, *Metacarpia'nus*, (F.) *Métacarpien*. From *meta*, and *καρπος*, 'wrist.' Relating or belonging to the metacarpus.

METACARPAL ARTERY, *Arte'ria dorsa'lis metacarp'i*, arises from the radial, at the moment it engages itself in the upper extremity of the abductor indicis. It descends obliquely upon the back of the hand. Its branches are distributed to the abductor indicis, and the integuments: some communicate with the dorsalis carpi.

METACARPAL ARTICULATIONS are those by which the last four metacarpal bones are united together at their upper extremity.

METACARPAL BONES, or bones which compose the metacarpus, are five in number; and distinguished into *first*, *second*, *third*, &c., beginning from the outer or radial side. They are articulated by their superior extremity with the bones of the second range of the carpus; and by the lower with the first phalanges.

METACARPAL LIGAMENT, *Inferior palmar Lig'ament*, is a fibrous band, stretched transversely before the inferior extremities of the last four metacarpal bones, which it keeps in their respective positions.

METACARPAL PHALAN'GES are the first phalanges of the fingers; so called, because they are articulated with the bones of the metacarpus.

METACARPAL RANGE or row of the carpal bones, (F.) *Rangée métacarpienne du carpe*, is the lower row of carpal bones; so called because they are articulated with the bones of the metacarpus. It is composed of the trapezium, trapezoides, magnum, and unciniforme.

METACARPEUS, Abductor metacarpi minimi digiti.

MÉTACARPIEN, Metacarpal — *m. du Petit doigt*, Opponens minimi digiti — *m. du Pouce*, Opponens pollicis.

METACARPION, Metacarpus.

METACARPO-PHALANGÆUS POLLICIS, Adductor pollicis manus.

METACARPO-PHALAN'GIAN, *Metacarpophalangia'nus*, *Metacarpophal'angal*, (F.) *Métacarpophalangien*. That which belongs to the metacarpus and phalanges.

METACARPO-PHALANGIAN or METACARPO-PHALANGAL ARTICULATIONS are formed by the bones of the metacarpus and the corresponding phalanges. In these articulations, the bony surfaces are incrustated with cartilage, covered by a synovial membrane, and kept in connexion by means of an anterior and two lateral ligaments.

MÉTACARPO-PHALANGIEN LATÉRAL SUS-PALMAIRE, ET MÉTACARPO-PHALANGIEN LATÉRAL, *Interosseus manus*.

METACARPUS, same etymon as Metacarpal. *Metacarp'ion*, *Postbrachia'le*, *Postcarp'ium*, *Torsus manûs*, (F.) *Métacarpe*. The part of the hand comprised between the carpus and fingers. It is composed of five parallel bones, called *metacarpal*; forming the back of the hand, external; and the palm internally.

METACERASMA, *Cerasma*.

METACHEIRIXIS, Surgery.

METACHIRISIS, Surgery.

METACHIRISMUS, Surgery.

METACHORESIS, Metastasis.

METACHYSIS, Transfusion.

METACINEMA, Metastasis.

METACINESIS, Metastasis.

METACONDYLE, see Phalanx.

METACYE'SIS, from *meta*, and *κύησις*, 'pregnancy'; *Graviditas extra-uteri'na secunda'ria*. Extra-uterine pregnancy in which the foetus is at first in the uterus, but subsequently in some other place.

METADERMATO'SIS, from *meta*, and *δερμα*, 'skin.' A morbid development of the epidermis or epithelium.

METAGEN'ESIS, (F.) *Metagénèse*, *Altern'ate genera'tion*, from *μετα*, and *γενεσις*, 'generation.' A term, proposed by Professor Owen, to express the changes of form, which one animal species undergoes in a series of successively produced individuals, extending from the one developed from the ovum to that which becomes the imago or last perfected individual. This term he employs in contradistinction to *metamorphosis*, which he restricts to the changes of form undergone by one and the same individual. See Generation, alternate.

METAL, *Meta'llum*. From *μεταλλω*, 'I seek after.' A class of simple, combustible bodies; distinguished from others by considerable specific gravity; a particular splendour; almost total opacity; insolubility in water; and the property they have of ringing when struck. Metals have no effect, except of a mechanical nature, when taken into the stomach; unless they have already undergone, or undergo in the stomach, oxidation or union with an acid; when, at times, deleterious compounds may be formed. Copper cents; half-pence; quicksilver; lead, have frequently been swallowed in the metallic state with impunity. Tin and mercury are the only metals prescribed for a mechanical effect; the former as an anthelmintic,—the latter, idly enough, in cases of fancied intussusception.

MÉTAL DES CLOCHES, Bell-metal.

METALLAGE, Metabasis.

METALLAXIS, Metallage.

METALLIC VOICE, *Tintement métallique*.

METALLODYN'IA, from *μεταλλον*, 'a metal,' and *δύνη*, 'pain.' Pain owing to the injurious influence of a metal—as lead, quicksilver, &c.

METALLUM, Metal — *m. Hæmatopoeticum*, ferrum.

METAMORPHOP'SIA, *Phantasmascop'ia*, *Phantasmascop'ia*, from *μεταμορφω*, (*μετα* and *μορφη*, 'form,') 'I transform, and *οψις*, 'vision.' *Suffu'sio*, *S. Oculo'rum seu metamorphop'sis seu myo'des*, *Imagination* (*Maître-Jean*); *Crup'ia*, *Marmar'ygis*, *Marmor'ygis*, *Mac'ula ante oc'ulos volitan'tes*, *Marmar'gia*, *Oc'uli marmar'ygides*, *Vis'io phantas'matum*, *Virus figura'tus*, *False sight*, (F.) *Berlue*. Aberration of the sense of sight, which transmits the image of imaginary objects. This affection sometimes depends on a slight opacity in the transparent parts of the eye; the cornea, crystalline, or vitreous humour, when

it is symptomatic and of no consequence. At other times, it appears to be idiopathic; and occurs particularly in those who have been in the habit of constantly fixing their eyes on very brilliant or small bodies. The objects fancied to be seen are various. They are sometimes circular, perpetually moving; or shining, or black spots, cobwebs, insects, or pieces of wool—when they are often termed *Muscae seu Flocci volitan'tes*, (F.) *Monches volantes*, and the condition *Visus muscarum*, *Scotom'ata*, *Suffusio Myo'des*, *Myiodesopsia*, *Myiodopsia*, *Myodeopsia*, *Myodesopsia*, *Visus muscarum*. These appearances sometimes continue for a few days; being dependent on the state of the nerves of the individual at the time; or they may exist for life, and ultimately impair the sight. Certain of them change their position, and appear to be seated in the humours of the eye, and—it has been supposed—in the vitreous humour more especially: hence the term *enthy'aloid muscae* applied to them.

If the affection be symptomatic, it is of but little moment. If idiopathic, and connected with any excitement of the brain, which is not often the case, attention will have to be paid to that organ.

METAMORPHOSIS, see Metagenesis.

METAMORPHOSIS VENTRICULI GELATINIFORMIS, Gastromalacia.

METANGISMOS, Decantation.

METAPEDIUM, Metatarsus.

METAPHRENON, Dorsum.

METAPHYTEIA, Transplantation.

METAPODIUM, Metatarsus.

METAPOROPH'EA, *Metaporopoe'sis*, from *meta*, πορος, 'a pore,' and ποιεω, 'to make.' The change produced in the minute pores, in the capillary extremities of vessels,—when they pass from the morbid to the healthy condition.—Galen.

METAPTO'SIS, *Metaschematis'mus*, *Metaschemat'isia*. Mutation, change; from μεταπιπρω, (μετα, and πιπρω, 'I fall,' 'I digress.' Any change in the form or seat of a disease. Transformation.

METARRHŒ'EA, *Metarrhysis*, from *meta*, and ρεω, 'I flow.' Reflux. The transfer of a disease from without to within, or from one part to another.

METARRHYSIS, Metarrhœa.

METASCHEMATISIS, Metaptosis.

METASCHEMATISMUS, Metaptosis.

METAS'TASIS, *Metachore'sis*, *Metacine'ma*, *Metacine'sis*, from μεταστημι, (μετα, and στημι, 'I place,') 'I change place.' Displacement, Translation. A change in the seat of a disease; attributed, by the Humorists, to the translation of the morbid matter to a part different from that which it had previously occupied: and by the Solidists, to the displacement of the irritation. It has also been used in the same extensive sense as Metaptosis. Disputes have often been indulged, whether a case of metastasis ought not rather to be esteemed one of extension of the disease. The phenomena of gout and acute rheumatism are in favour of metastasis occasionally supervening.

METASTASIS LACTIS, Galactoplasia, Phlegmatia dolens.

METASTAT'IC, *Metastat'icus*, (F.) *Métastatique*. Belonging or relating to metastasis. A *metastatic crisis* is one produced by metastasis:—a *metastatic affection*, one caused by metastasis, &c. See Abscess, metastatic.

METASYN'CRISIS, from *meta*, and συγκρινω, (συγ, and κρινω, 'I choose,' 'I judge.') *Recomposition*. A word employed by some disciples of Asclepiades. This physician supposed, that all animals are formed by the union or assemblage of atoms. He designated all bodies by the

word συγκριματα, which signifies 'assemblage.' Συγκρινομαι, 'to be assembled,' was, with him, synonymous with 'to exist,' and διακρινομαι, 'to separate,' was, with him, synonymous with 'to dissolve,' 'to cease to exist.' The word *Metasyn-crisis* was invented to express the recombination of bodies after their momentary dissolution. Some have rendered the word by the Latin *Re-corporatio*.

METASYNCRIT'IC, *Metasyncrit'icus*, *Re-corporat'ivus*. Belonging or relating to metasyn-crisis. A name formerly given to medicines to which was attributed the virtue of producing the metasyn-crisis or regeneration of the body, or some of its parts.

METASYNCRITIC CYCLE meant a determinate series of remedies employed for this purpose.

METATARS'AL, *Metatar'ceus*, (F.) *Métatarsien*. Same etymon as metatarsus. Relating or belonging to the metatarsus.

METATARSAL ARTERY, *Artère sus-métatarsienne* of Chaussier, arises from the *Arteria dorsalis Tarsi*; passes transversely over the back of the foot; and furnishes, at its convexity, which is anterior, three branches, called *A. interos'seæ dorsales Pedis*. These branches are distributed in the interosseous spaces.

METATARSAL ARTICULA'TIONS are those resulting from the junction of the metatarsal bones with each other. They are strengthened, behind, by dorsal and plantar ligaments: and before, by a transverse metatarsal ligament, which is plantar, and has the greatest analogy to the inferior transverse metacarpal ligament.

METATARSAL BONES, (F.) *Os métatarsiens*, are five in number, and distinguished by their number; first, second, third, &c., reckoning from the outer side.

METATARSAL PHALANGES are the first phalanges of the toes; so called because they are united to the metatarsus.

METATARSAL ROW—(F.) *Rangée métatarsienne*—of the bones of the tarsus, is the second row, or that contiguous to the metatarsus; comprehending the cuboides and three cuneiform bones: some add the scaphoides.

METATARSEUS, see Abductor minimi digiti pedis.

MÉTATARSIE'N, Metatarsal.

METATARSO-PHALAN'GIAN, *Metatarso-phalangia'nus*, *Metatarso-phalangal*. Relating to the metatarsus and phalanges.

METATARSO-PHALANGIAN or METATARSO-PHALANGAL ARTICULATIONS are formed by the bones of the metatarsus and the corresponding phalanges. They bear the greatest analogy to the metacarpophalangian articulations.

MÉTATARSO-PHALANGIEN-LATÉRAL, see Interosseal pedis—*m. Sous-phalangien transversal du premier orteil*, Transversus pedis.

METATAR'SUS, *Metatar'sium*, *Metape'dium*, *Metapo'dium*, *Præcor'dium* seu *So'ium pedis*, *Vestig'ium pedis*, (F.) *Métatarsee*, from μετα, 'after,' and ραπος, 'the tarsus.' That part of the foot which is situate between the tarsus and toes, corresponding with the metacarpus. It is composed of five parallel bones; one to each toe. Like the metacarpal bones, they are developed from two points of ossification.

METATH'ESIS, from μεταστημι, (μετα, and στημι, 'to place,') 'I change place.' *Transposit'io*. An operation, by which a morbid agent is removed from one place to another, where it may produce less disturbance in the exercise of the functions:—as, for example, in the operation of depressing cataract, or when calculus in the urethra is pushed back into the bladder. Also, Derivation.

METATIO, Mensuration.

METATOC'IA, from *meta*, and *tokos*, 'birth.' Parturition in a preternatural manner. Preternatural labour.

METATOPTOSIS, Diadexis.

METENSOMATOSIS, Incorporation.

METEORISMUS, Sublimation, Tympanites.

METEOROL'OGY, *Meteorolog'ia*, from *μετεωρος*, (*meta*, and *αιωρω*, 'I lift up,') 'a meteor,' and *λογος*, 'a discourse.' The science, whose object is a knowledge of the origin, formation, appearance, &c. of meteors. The state of the atmosphere has a most important bearing upon the health of animals. Every epidemic affection has its causes seated there. Meteorological conditions have, hence, ever been attended to by the physician for the purpose of detecting the precise character of any particular epidemic influence. The barometer, thermometer, and hygrometer are the instruments used with this intent—to detect, as well the varying weight or pressure, and the temperature, as the moisture. Perhaps, of the three conditions, the last exerts more influence in the production of disease than either of the others. Our knowledge, however, of this part of physics is extremely limited and unsatisfactory.

METER, *μετρον*, 'a measure.' A suffix to words denoting 'a measure,' as in *Barometer*, *Pleximeter*, &c.

METH. A prefix to terms. Same signification as *Meta*.

METHÆMACHYmia, Transfusion.

METHÆMATA; from *meth*, and *αιμα*, 'blood.' The capillary or intermediate system of vessels in which the blood undergoes the change from venous to arterial, and conversely.—Marshall Hall. See Capillary vessels.

METHE, Temulencia.

METHEGLIN, Melizomum.

METHEMERINUS, Quotidian.

METHOD, *Meth'odus*, from *meth*, and *οδος*, 'way.' This word has different acceptations in the sciences. In medicine, *curative method*, *meth'odus medendi*, is the methodical treatment of disease.

METHOD OF COUNTING, Method, numerical—*m.* Mariano's, see Lithotomy.

METHOD, NUMER'ICAL, *Method of Observation or of Counting*, of Louis, (F.) *Méthode numérique*, consists in observing every case and every symptom of a case numerically, so as to ensure, as far as practicable, accuracy of observation; and to enable us, by the analysis and collation of such facts, to deduce general laws and conclusions. Also, the application of numbers to the study of disease. See Statistics, medical.

MÉTHODE NUMÉRIQUE, Method, numerical, Statistics, medical—*m.* *Perturbatrice*, *Perturbatrix* (Medicina)—*m.* *Stibio-dermique*, see Unguentum antimonii.

METHODICS, Methodists.

MÉTHODIQUES, Methodists.

METH'ODISTS, *Method'ical sect*, *Method'ics*, (F.) *Méthodistes*, *Méthodiques*. Same etymon. A sect of physicians whose doctrine was in vogue after that of the Empirics and Dogmatists, towards the end of the first century. According to the Methodists, of whom Themison was the chief, almost every disease is dependent on contraction or relaxation—*strictum* or *laxum*. To these two causes, they added a third—*mixed* or *compound*—to include those affections which partook of the two characters. The doctrine, *medici'na method'ica*, resembled, in some respects, that of Brown.

METHODOL'OGY (MEDICAL), *Methodolog'ia Med'ica*, from *μεθόδος*, 'method,' and *λογος*,

'a discourse.' A word used, by the French more especially, to signify *method* applied to the study of any science. *Medical Methodology*, consequently, means *method* applied to the study of medicine.

METHODUS CATALEPTICA, see Cataleptic—*m.* Celsiana, see Lithotomy—*m.* Curatoria, Therapeutics.

METHODUS DERIVATO'RIA. The derivative or revellent system of treatment.

METHODUS ENDERMATICA seu ENDERMICA, see Endermic—*m.* Franconiana, see Lithotomy—*m.* Guytoniana, see Lithotomy—*m.* Medendi, Method of cure, Therapeutics.

METHOMANIA, from *μεθην*, 'drunkenness,' and *μανια*, 'mania.' An irresistible desire for intoxicating substances. Temulencia.

METHU, see Temulencia, Wine.

METHYLE, HYDRATED OXIDE OF, Spiritus pyroxylicus.

METHYLENE, BIHYDRATE OF, Spiritus pyroxylicus.

METHYSIS, Temulencia.

METHYSMUS, Temulencia.

METHYS'TICUS, from *μεθην*, 'drunkenness.' That which causes, or pertains to, drunkenness.

METHYSTOPHYLLUM GLAUCUM. A South African plant, *Nat. Ord.* Amyridaceæ, an infusion of whose leaves is pleasant to the taste, and is used in bronchitis, asthma, and other thoracic diseases. With the Bushmen and others, it is a favourite beverage, and is called by them *Boschjeemanthee*, 'Bushman's tea.' It is, also, chewed.

MÉTIS, Hybrid.

MÉTISAGE, Crossing.

METODONTIASIS, from *meta*, and *οδοντιας*, 'dentition.' Faulty development of the teeth.

METOPAGES, Symphycephalus.

METOPANTRA, Frontal sinuses.

METOPANTRAL'GIA, from *μετωπον* (*meta*, and *ωψ*, 'the eye') 'the forehead,' *αντρον*, 'a cavity,' and *αλγος*, 'pain.' Pain in the frontal sinuses.

METOPANTRITIS, *Inflammat'io sinuum frontali'um*, from *metopantra*, and *itis*, denoting inflammation. Inflammation of the frontal sinuses.

METOPION, Bubon galbanum.

METOPIMUM, Mesopryon.

METOPODYNIA, Neuralgia frontalis.

METOPON, Front, Frontal bone.

METOPOS/COPY, *Metoposcop'ia*, from *μετωπον*, (*meta*, and *ωψ*, 'the eye,') 'the forehead,' and *σκοπειν*, 'to view.' The art of knowing the temperament and character of a person by inspecting the traits of his forehead or face. See Physiognomy.

METRA, *μητρα*, Uterus.

METRÆ, Secundines.

METRÆMORRHAGIA, Metrorrhagia.

METRÆMORRHŌIDES, from *metra*, and, *αιμορροειδης*, 'hemorrhoids.' Hemorrhoids or varicose veins at the orifice of the uterus.

METRAL'GIA, *Metrodyn'ia*, from *metra*, and *αλγος*, 'pain.' Pain in the uterus.

METRANÆMIA, from *metra*, and *anæmia*. An anæmic condition of the uterus.

METRANASTROPHE, Uterus, inversion of the.

METRANEURYSMUS, Metreuryisma.

METRATRE'SIA, *Imperforat'io uteri*, from *metra*, and *αρρηα*, 'imperforation.' An unnatural closure of the uterus.

METRATROPH'IA, from *metra*, a, 'privative,' and *τροφην*, 'nourishment.' Atrophy or want of development of the uterus.

METRAUXË, *Hypertroph'ia seu Sarcœsia*

u'teri, from *metra*, and *αὔξω*, 'increase.' Hypertrophy of the uterus.

METRECTASIA, from *metra*, and *εκτασις*, 'dilatation.' Dilatation of the womb.

METRELCO'SIS, *Metrelco'sis*, from *metra*, and *ἔλκος*, 'an ulcer.' Ulceration of the uterus.

METREMPHRAX'IS, from *metra*, and *εμφρασσω*, 'I obstruct.' Obstruction of the womb or of the vessels of the womb—*Infarc'tus u'teri*.

A name under which some authors have confounded chronic inflammation of that viscus, and the different degenerations to which it is exposed.

METREMPHYSEMA, *Physometra*.

METREN'CHYTA, *Metren'chytes*, from *metra*, and *εγχύω*, 'I inject.' Injection of the uterus. Substances injected into the uterus are called *Metren'chyta*.

METREURYS'MA, *Metraneurys'mus*, from *metra*, and *εὔρω*, 'dilated.' A morbid dilatation of the womb.

METRHELCO'SIS, *Metrelcosis*.

METRHYMENTIS, see *Metritis*.

METRIOPATHI'A, from *μετρός*, 'tempered,' and *πάθος*, 'affection.' State of an individual whose passions are temperate.

METRI'TIS, from *metra*, and *itis*, denoting inflammation. *Febris uteri'na*, *Hysteri'tis*, *Empres'ma Hysteri'tis*, *Inflammat'io U'teri*, *Metrophlogo'sis*, *Inflammation of the Uterus or Womb*, (F.) *Inflammation de la matrice*. The characteristic symptoms of this affection are:—pain, swelling, and tenderness in the hypogastric region; with heat, pain, and tenderness of the os uteri; vomiting, smallness, and frequency of pulse. It occurs most frequently after delivery, *Lochometri'tis*, when there is generally suppression of the lochial discharge. The treatment must be vigorous,—bleeding early, so as to make a decided impression; followed by a full dose of opium, fomentations, blisters, &c.

Acute inflammation of the womb, seated in its internal membrane, *Endo-metri'tis*, *Metrhymen'i'tis*, *Enmetri'tis*, has been called *Inflammat'io catarrhalis u'teri* or *Acute catarrh*, (F.) *Catarrhe aigu*, of that viscus. It is known by the discharge of a clear, stringy fluid *per vaginam*; preceded by pains, which, from the hypogastric region, shoot to the thighs, groins, &c., with more or less fever. It requires the antiphlogistic treatment. *Chronic metritis* sometimes succeeds the acute. To it must be referred the indurations, observed in the uterus, and many of the leucorrhœal discharges to which females are subject.

METRITIS RHEUMATICA, *Metrorrhœuma*—m. Septica, *Metrocace*—m. Venosa, see *Phlebitis*.

METROBLENNORRHŒA, *Leucorrhœa uteri*.

METROBLENNOZEMIA, *Leucorrhœa uteri*.

METROBOTRYTES, Cauliflower excrescence of the uterus.

METROC'ACE, *Metri'tis sept'ica*, from *metra*, and *κακος*, 'evil.' Putrescency or gangrene of the uterus.

METROCAMPSIS, *Hysteroloxia*.

METROCARCINO'MA, *Hystero-carcino'ma*, *U'teri carcino'ma*, *Cancer u'teri*, from *metra*, and *καρκίνωμα*, 'cancer.' Cancer of the uterus.

METROCATARRHUS, *Leucorrhœa uteri*.

METROCELE, *Hysterocele*.

METROCELIDES, *Nævus*.

METROCHOL'SIS, from *metra*, and *χολος*, 'bile.' *Febris puerpera'lis bilio'sa*.

METRODYNIA, *Metralgia*.

METROHÆMIA, *Hypermetrohémie*, from *metra*, and *αἷμα*, 'blood.' Hyperæmia or congestion of blood in the uterus.

METROHEMORRHAGE, *Metrorrhagia*.

METROLEUCORRHŒA, *Leucorrhœa uteri*.

METROLOXIA, *Hysteroloxia*.

METROMALACOMA, *Hysteromalacia*.

METROMALACOSIS, *Hysteromalacia*.

METROMANIA, *Nymphomania*.

METROMY'ZEUS, from *metra*, and *μυζω*, 'I suck.' An exhausting syringe, to be applied to the os uteri, for exhausting the womb in cases of amenorrhœa.

METRONCUS, *Hysteroncus*.

METROPARAL'YSIS, from *metra*, and *παράλυσις*, 'paralysis.' Paralysis of the uterus.

METROPATHI'A, from *metra*, and *πάθος*, 'affection.' An affection of the womb.

METRO-PÉRITONITE PUERPÉRALE, see *Peritonitis*.

METROPERITONITIS, from *metra*, and 'peritonitis.' *Inflammat'io u'teri et peritonæi*. Inflammation of the uterus and peritoneum. Puerperal Fever. See *Peritonitis*.

METROPHLEBITIS, from *metra*, *φλεψ*, 'a vein,' and *itis*, denoting inflammation. Inflammation of the veins of the uterus.

METROPHLEBITIS PUERPÉRALIS, see *Phlebitis*.

METROPHLOGOSIS, *Metritis*.

METROPHTHISIS, *Phthi'sis uteri'na*, from *metra*, and *φθίω*, 'I consume.' Consumption from ulceration of the uterus—*Ul'cera seu Helco'sis u'teri*.

METROPHYMA, *Hysteroncus*.

METROPLETHO'RA, *Metrupe'ra'mia*, from *metra*, and *πληθωρα*, 'fulness.' Congestion or hyperæmia of the uterus.

METROPOLYPUS, *Hys'tero-pol'ypus*, *Polypus U'teri*, from *metra*, and *polypus*. Polypus of the uterus.

METROPROPTOSIS, *Prolapsus uteri*.

METROPTOSIS, *Prolapsus uteri*.

METRORRHAG'IA, *Metæmorrhag'ia*, *Hæmorrhagia uteri'na* seu *u'teri*, *Sanguiflux'us uteri'nus*, *Sanguinis stillicid'ium ab U'tero*, *Fluor uteri'ni sanguinis*, *Profluv'ium Sanguinis ex u'tero*, *Hystero'rrhag'ia sanguin'ea*, *Hystero'rrhœa*, *Hæmorrhœa uteri'na* (of some), *Menorrhag'ia* (of some), *Hæmatometra*, *Hæmatomatra*, *Uterine Hemorrhage*, *Hemorrhage from the Womb*, *Metrohæmorrhage*, (F.) *Hémorrhagie de la Matrice*, *Pertes*, *Pertes utérines rouges*, *P. de sang*; from *metra*, and *ρρῑνναι*, 'I break forth.' An effusion of blood from the inner surface of the uterus, either at the menstrual or other periods; but in a greater quantity than proper. Uterine hemorrhage may be caused by those influences which produce hemorrhage in general. It happens, however, more frequently during pregnancy, and during or after delivery, when the vascular system of the uterus is so circumstanced as to favour its occurrence more than at other periods. The termination of metrorrhagia is usually favourable. Should it, however, be very copious, or frequently recur; or should it happen to a great extent after delivery, death may take place very speedily; and, in some cases, without the discharge being perceptible; constituting *internal hemorrhage*.

Uterine hemorrhage may be *active* or *passive*; requiring obviously a different treatment. The main management is similar to that of hemorrhage in general;—the horizontal posture; acid drinks; free admission of cool air; cold applications to the loins, thighs, and abdomen; injection of cold water, even of iced water, into the vagina; plugging the vagina, so as to prevent the discharge *per vaginam*, and thus induce a coagulum in the mouths of the bleeding vessels. Such will be the *special* plan adopted where the hemorrhage has occurred in one not recently delivered. In uterine hemorrhage after delivery, the same cooling plan must be followed; but, as the flow of blood is owing to the uterus not contracting so

as to constrict its vessels, pressure must be made on the abdomen to aid this: and, if necessary, the hand must be introduced into the uterus to stimulate it to contraction. Should the female be excessively reduced, so as to render the accoucheur apprehensive that she may expire from loss of blood, brandy may be exhibited. The profuse exhibition of opium in such cases is, at least, a doubtful plan.

Transfusion has, at times, been practised as a last resource.

METORRRHEUMA, *Rheumatis'mus Uteri*, *Hyster'al'gia rheumat'ica seu catarrh'al'is*, *Metri'tis rheumat'ica*, (F.) *Rhumatisme de l'utérus*; from *metra*, and *ρῆμα*, 'defluxion,' 'rheumatism.' Rheumatism of the uterus.

METORRRHEXIS, Uterus, rupture of the.

METORRRHŒA, from *metra*, and *ρῆω*, 'I flow.' A protracted discharge of any fluid from the uterus.

METROSALPINX, Tube, Fallopian.

METROSCIRRHUS, *Hysteroscir'rhus*, *Scirr'hus seu Carcin'o'ma uteri*, from *metra*, and *σκίρρ'ος*, 'scirrhus.' Scirrhus of the uterus.

METROSCOLIOMA, Hysteroloxia.

METROSCOPE, *Metroscop'ium*, from *metra*, and *σκοπεῖν*, 'to view.' An instrument, invented by M. Nauche, for listening to the sounds of the heart of the fœtus, in utero-gestation, when the sounds and movements are imperceptible through the parietes of the abdomen. The extremity of the instrument—the first notion of which was given to M. Nauche by the stethoscope of Laënnec—is introduced into the vagina and applied against the neck of the uterus.

METROSIDEROUS GUMMIFERA, see Kino.

METROSTERE'SIS, from *metra*, and *στερεσις*, 'privation.' Extirpation of the uterus. Want of uterus.

METROSYNIZE'SIS, *Hysterosynize'sis*, from *metra*, and *συνίζω*, 'running together.' Morbid union of the uterus to adjacent parts.

MÉTROTOME, Hysterotomus.

METROTOMIA, Cæsarean section.

METRYPERŒMIA, Metroplethora.

METRYPERŒSTHESIS, see Uterus, irritable.

MEU, *Æthusa meum*.

MEULIÈRE, Molar.

MEUM, *Æthusa meum*—m. *Anethifolium*, *Æthusa meum*—m. *Athamanticum*, *Æthusa meum*.

MEURTRISSURE, Confusion.

MEVIUM, Syphilis.

MEZEA, Genital organs.

MEZEREON, *Daphne mezereum*.

MEZEREUM ACETATUM, see *Daphne mezereum*.

MIAMMA, Miasma.

MIANSIS, Miasma.

MIARIA, Miasma.

MIASMA, *Miasm*, *μιασμα*, 'a stain,' from *μιαίνω*. 'I contaminate.' *Miasm'ma*, *Mian'sis*, *Mia'ria*, *Mias'mus*, *Inquinamen'tum*, *Molyn'sis*, *Cont'gium*. The word *miasm* has, by some, been employed synonymously with contagion. It is now used more definitely for any emanation, either from the bodies of the sick, or from animal and vegetable substances, or from the earth, especially in marshy districts, (*Marsh poison*, *Efflu'vium palus'tre*, *Gas palus'tre*, (F.) *Intoxication des Marais*), which may exert a morbid influence on those who are exposed to its action. To these terrestrial emanations—the *Koino-mias'mata* of Dr. E. Miller, of New York—the Italians give the name *aria cattiva*, but, more commonly, *mal'aria* ('bad air'); a word which has been adopted into other languages. The deleterious effluvia, originating from the decomposition of matter de-

rived from the human body, have been called by Dr. Miller, *Idiomias'mata*; the epithets *Koino* and *Idio* being derived respectively from *κοινος*, 'common,' and *ιδιος*, 'personal.' Of the miasms which arise either from the animal body or from the most unhealthy situations, we know, chemically, nothing. All that we do know is, that, under such circumstances, emanations take place, capable of causing disease in many of those who are exposed to their action.

MIASMAT'IC, *Miasmaticus*. Belonging or relating to miasmata. Sauvages, in his classification of diseases according to their etiology, has a class under the name *Morbi miasmatici*. A fever that arises from marshy miasms, is styled *malaria*, *palud'al*, *Helop'ya*, *Helopyr'etus*, &c.

MIASMUS, Miasma.

MIAULANTES, see *Convulsivonaire*.

MICÆ PANIS, see *Triticum*.

MICATIO CORDIS, Systole.

MICHELIA CHAMPACA, *Champaca*, *Michelia Tsjampaca*. Ord. Magnoliaceæ. An Oriental tree, much prized for the odour and beauty of its flowers. The oily infusion of the flowers is employed in the Moluccas in headache.

MICL'TA. A medicine used by Mesue for arresting hemorrhage; perhaps, according to Siebenhaar, from the Arabic, *micnata*, "proved by experience."

MICOSIS, Framboesia.

MICRANATOMIA, see Anatomy.

MICRANGIA, Capillary vessels.

MICRENCEPHALIUM, Cerebellum.

MICRENCEPHALON, Cretinism.

MICRENCEPHALUM, Cerebellum.

MICROCEPHALIA, see Microcephalus.

MICROCEPHALUS, *Mikroceph'alus*, *Micro-cra'nus*, from *micros*, and *κεφαλή*, 'head.' One who has a small head. A monster having a small imperfect head, or a small imperfect cranium. The condition is called *Microceph'al'ia*, *Cranio'stero'sis*. With some it means Idiostism.

MICROCORTIA, Myosis.

MICROCOSM, *Microcos'mus*, from *micros*, and *κοσμος*, 'world.' A little world. Some philosophers have given this appellation to man, whom they consider as the epitome of all that is admirable in the world. The world they call *Ma'crocosm*.

MICROCOSMETOR, Vital principle.

MICROCOSM'ICA MACH'INA. The organism of man.

MICROCOSMICA SCIENTIA, Physiology.

MICROCOSMICUM SAL, with the ancients meant the salts of the urine,—*Sal uri'næ*.

MICROCOSMOGRAPHIA, Physiology.

MICROCOUST'IC, *Microcous'ticus*, from *micros*, and *ακουω*, 'I hear.' This word, as well as *Microph'onous*, from *μικρος*, 'small,' and *φωνη*, 'voice,' means any thing that contributes to increase the intensity of sound—as the speaking trumpet—by collecting the sonorous rays.

MICROCRANIUM, Microcephalus.

MICROGLOS'SIA, from *micros*, and *γλωσσα*, 'tongue.' Original smallness of tongue.

MICROLEUCONYMPHŒA, *Nymphæa alba*.

MICROLOGY, see Microscopy.

MICROM'ELUS, *Hemim'eles*, from *micros*, and *μελος*, 'a limb.' A monster having imperfectly developed extremities.

MICROMMATUS, Microphthalmus.

MICROPHONOUS, Microacoustic.

MICROPH'THAL'MUS, *Microm'matus*, from *micros*, and *ὀφθαλμος*, 'eye.' One who has small eyes. A monster with two small, or imperfectly developed eyes. A small eye,—*Ophthalmid'ium*, *Ophthalm'mum*.

MICROPODIA, Micropus.

MICROPROSOPUS, *Aprosopus*, from *micros*, and *πρῶσῶπ*, 'face.' A monster in which a part of the face is absent.

MICROPUS, from *micros*, and *πους*, 'foot.' One who has small feet. The condition is called *Micropodia*.

MICROPYLE, *Micropylum*, from *micros*, and *πύλη*, 'gate.' An opening in the episperm of seeds, through which the fecundating pollen enters the ovule to fecundate it. It has been supposed, that such a micropyle exists in the human ovule.

MICROCHIDES, from *micros*, and *ορχίς*, 'a testicle.' They who have very small testicles.

MICROS, *μικρός*, 'little,' as in

MICROSCOPY, *Microscopia*; from *micros*, and *σκοπή*, 'a view.' Observation by the microscope: an important agency in the examination of the healthy and morbid tissues. A description of the results of such observation is *Microraphy*. *Micrology* is often used with the same signification, and for a treatise on the microscope and the results of microscopy.

MICROSOMIA, *Nanosomia*.

MICROSPHYCTUS. Same etymon as the next. One who has a small pulse.

MICROSPHYXIA, from *micros*, and *σφυγμος*, 'pulse.' Smallness or weakness of pulse.

MICROSPORON AUDOUNI, see *Porrigio decalyans*—*m. Furfur*, see *Chloasma*—*m. Mentagrophytes*, *Mentagrophyte*.

MICROSOTOMUS; from *micros*, and *στομα*, 'mouth.' One who has a small mouth.

MICROTESIA, *Partium organica'rum parvitas morbo'sa*; from *μικρός*, 'smallness.' Morbid smallness of organic parts.

MICROTRAPEZIA, *Leptotrophia*.

MICROTRICHIA; from *micros*, and *τριχ*, 'hair.' Fineness or shortness of hair.

MICROTROPHIA, *Leptotrophia*.

MICTIO, Micturition — *m. Cruenta*, *Hæmaturia*—*m. Inopportuna*, *Enuresis*—*m. Involuntaria*, *Enuresis*—*m. Sanguinea*, *Hæmaturia*.

MICTION, Micturition.

MICTURITION, *Mictio*, *Mictus*, *Ure'sis*, *Uris'sis*, *Omiche'sis*, (F.) *Miction*, *Action de pisser*, from *micturio*, and *mingere*, *mictum*, 'I make water.' The act of making water—*Urination*. Also, morbid frequency of passing the water.

MICTUS, Micturition — *m. Cruentus*, *Hæmaturia*.

MIDRIFF, *Diaphragm*.

MIDWIFE, from *mid*, 'with,' and *pir*, 'wife,' or, from *midz*, 'meed,' 'recompense,' and *pir*, 'wife.' *Matro'na*, *Obstet'rix*, *Hyper'tria*, *Mæa*, *Mæas*, *Mæu'tria*, *Aces'toria*, *Aces'tria*, *Aces'tria*, (Sc.) *Howdy*, (Prov.) *Grace-wife*, *Handwoman*, *Headswoman*, *Howdy-wife*, (F.) *Sage femme*, *Accoucheuse*. A female who practises obstetrics.

MIDWIFERY, *Obstetrics*.

MIEL, *Mel*—*m. d'Acétate de Cuivre*, *Linimentum æruginis*—*m. Boratè*, *Mel boracis*—*m. Dépuré*, *Mel despumatum*—*m. Despuné*, *Mel despumatum*—*m. Médicinal*, *Mellitum*—*m. Mercuriale composé*, *Mellitum de mercuriali compositum*.

MIEUTER, *Accoucheur*.

MIGMA, *Mistura*.

MIGRAINE, *Hemicrania*.

MIGRANA, *Hemicrania*.

MIGRATION DU LAMBEAU, see *Flap*.

MIKANIA GUACO, *Guaco*.

MICROCEPHALUS, *Microcephalus*.

MIL, *Panicum miliaceum*.

MILFOIL, *Achillea millefolium*.

MILIAIRE, *Miliary*, *Miliary fever*.

MILIARIA, *Miliary fever*—*m. Sudans*, *Miliary fever*—*m. Sudatoria*, *Miliary fever*.

MILIARIS SUDATORIA, *Sudor anglicus*, *S* *picardicus*.

MILIARY, *Miliaris*, (F.) *Miliaire*; from *milium*, 'millet.' Resembling millet seed—hence, *miliary fever*, *miliary glands*, &c.

MILIARY FEVER, *Em'phlysis miliaris*, *Miliaris*, *Miliaria*, *M. sudans* seu *sudatoria*, *Sudor*, *S. miliaris*, *Pap'ula miliaris*, *Exanthema miliaria*, *Syn'ochus miliaria*, *Apre'do milia'cea*, *Feb'ris miliaris* seu *essero'sa* seu *puncticula'ris* seu *culicula'ris* seu *vesicula'ris* seu *lenticula'ris* seu *purpura'te rubra* et *alba miliaris*, *Pur'pura puerpera'rum*, *P. miliaris*, *Pap'ule sudo'ris* *Millet seed rash*, (F.) *Miliaire*, *Milloy*, *Pourpre blanc*, *Millet*. It is so called from the eruption resembling the seed of the *milium* or *millet*. Fever, accompanied by an eruption of small, red, isolated pimples, rarely confluent, but almost always very numerous, slightly raised above the skin, and presenting, at the end of 24 hours, a small vesicle filled with a white transparent fluid; which quickly dries up, and separates in the form of scales. Miliary fever is now rare, in consequence of the cooling practice in fevers and other states of the system. It is almost always brought on by external heat, and hence the prevention is obvious. The treatment is simple. It requires merely the antiphlogistic regimen, in ordinary cases.

MILIARY GLANDS, *Sebaceous glands*—*m. Granulations* or *Tubercles*, see *Granulation*.

MILII SEMINA, *Panicum Italicum*.

MILIOLOM. Diminutive of *milium*, 'millet.' *Cenchr'id'ion*. A small tumour on the eyelids, resembling, in size, a millet-seed.

MILITARIS HERBA, *Achillea millefolium*.

MILITARY, *Militaris*, from *miles*, *militis*, (F.) *Militaire*, 'a soldier.' Pertaining to a soldier or an army—as '*Military medicine*,' *Médecine militaire*, (F.) *Médecine militaire*; and *Military surgery*, *Chirur'gia militaris*, (F.) *Chirurgie militaire*. Medicine and surgery applicable to the soldier.

MILIUM, *Grutum*, *Panicum miliaceum*, see *Hydatid*—*m. Esculentum*, *Panicum miliaceum*—*m. Indicum*, *Panicum Italicum*—*m. Panicum*, *Panicum miliaceum*—*m. Solis*, *Lithospermum officinale*.

MILK, *Sax. melc*, *Lac*, *Gala*, *Humor lac'teus*, *Latex ni'veus*, (F.) *Lait*. A fluid secreted by the mammary glands of the females of the mammalia. The skimmed milk of the cow, (F.) *Lait de Vache*, contains water, caseous matter, traces of butter, sugar of milk, chloride of sodium, phosphate, and acetate of potassa, lactic acid, lactate of iron, and earthy phosphate. The cream is formed of butter, casein, and whey, in which there is sugar of milk and salts. When examined by the microscope, milk is seen to contain a large number of particles, of irregular size and shape, varying from $\frac{1}{125}$ to $\frac{1}{300}$ of an inch in diameter. They consist of oily matter, surrounded by a delicate pellicle, and are the *milk globules*.

Cow's milk is employed for the preparation of cream, butter, cheese, whey, sugar of milk, and frangipane. It is useful in a number of cases of poisoning; either by acting as a demulcent, or by decomposing certain poisons, or by combining with others so as to neutralize them. It is constantly employed as aliment, and may be regarded as an emulsion in which butter and casein are found in suspension. When taken into the stomach, it is coagulated by the gastric fluids, and the coagulum is digested like any other solid. The watery parts are absorbed.

Between milk, flour, and blood, there is great similarity of composition. The following table is given by Dr. Robert Dundas Thomson:

Milk.	Flour.	Blood.
Curd or Casein	Fibrin,	Fibrin,
	Albumen,	Albumen,
	Casein,	Casein,
	Gluten,	Colouring matter.
Butter.....	Oil.	Fat.
Sugar.....	Sugar,	Sugar.
	Starch,	
Chloride of potassium, sodium,	Do.	Do.
Phosphate of soda,		
lime,		
magnesia, iron,		

From a considerable number of experiments, Messrs. Deyeux and Parmentier class the six kinds of milk, which they examined, according to the following table, as regards the relative quantity of materials they contain.

Casein.	Butter.	Sugar of Milk.	Serum.
Goat	Sheep	Woman	Ass
Sheep	Cow	Ass	Woman
Cow	Goat	Mare	Mare
Ass	Woman	Cow	Cow
Woman	Ass	Goat	Goat
Mare	Mare	Sheep	Sheep

MILK ABSCESS, Mastodynia apostematosa.

MILK, ALMOND, Emulsio amygdalæ.

MILK AND SODA WATER. An agreeable mode of taking milk in cases where it lies heavily on the stomach. Heat, nearly to boiling, a teacupful of milk; dissolve in it a teaspoonful of sugar, put it into a large tumbler, and pour over it two-thirds of a bottle of soda water.

MILK, ASSES', (F.) *Lait d'Anesse*, considerably resembles human milk, of which it has the consistency, smell, and taste: but it contains a little less cream, and more soft, caseous matter. It is often used by those labouring under pulmonary affections.

Artificial Asses' Milk may be made by taking gelatin 3ss; dissolving it, by the aid of heat, in barley water Oij; adding refined sugar 3j; pouring into the mixture new milk Oj; and beating the whole with a whisk.

It may also be prepared by dissolving sugar of milk 3ij in tepid skimmed cow's milk Oj.

MILK BLITCH, Porrigo larvalis — m. Crust, Porrigo larvalis.

MILK, EWES', (F.) *Lait de Brebis*. It affords more cream than cows' milk; but the butter is softer. The caseum, on the contrary, is fatter, and more viscid. It contains less serum than cow's milk. The Roquefort cheeses are made from it.

MILK FEVER, see Fever, milk — m. Glass, Breast glass — m. Globules, see Milk.

MILK, GOATS', (F.) *Lait de Chèvre*, resembles cows' milk: the butyraceous matter, however, which enters into its composition, is more solid than that of the cow.

Artificial Goat's Milk may be made by taking fresh sweet 3j; cutting it into small pieces; tying it in a muslin bag, large enough to leave the morsels free from compression; and boiling in a quart of cow's milk, sweetened with a quarter of an ounce of white sugar candy.

Used as a diet in scrofulous cases, and also in phthisis.

MILK, GREEN, Colostrum.

MILK, HUMAN, (F.) *Lait de femme*,—contains

more sugar, milk, and cream, and less caseum, than cows' milk. Its composition differs according to the distance of time from delivery.

MILK KNOT, *Galacto-ganglion*; (F.) *Ganglion lacteux*. The knots often observed in the breast after inflammation of the organ, or for some time after the suppression of the secretion. They generally end by resolution.

MILK LEG, Phlegmatia dolens.

MILK, MALES', Sperm.

MILK, MARES', (F.) *Lait de Jument*,—contains only a small quantity of fluid butyraceous matter; a little caseum, softer than that of cows' milk, and more serum.

MILK SCALL, Porrigo larvalis.

MILK SICKNESS, *Sick stomach, Swamp sickness, Tires, Slows, Stiff joints, Puking fever, River sickness*. A disease occasionally observed in the States of Alabama, Indiana, and Kentucky, which affects both man and cattle, but chiefly the latter. It is attributed in cattle to something eaten or drunk by them; and in man to the eating of the flesh of animals labouring under the disease. Owing to the tremors that characterize it in animals, it is called the *Trembles*. It is endemic.

The symptoms of the disease are such as are produced by the acro-narcotic class of poisons—vomiting, purging, extreme nervous agitation, &c.: and the approved indications of treatment appear to be—gentle emetics and laxatives, with quiet, and mucilaginous drinks.

MILK, SNAKES', Euphorbia corollata.

MILK, SOLIDIFIED. Milk, to which little less than an equal weight of white sugar has been added, evaporated to dryness. When re-dissolved in water, it has all the qualities of fresh milk, with the additional quantity of sugar.

To milk, evaporated to dryness, MM. Grimaud and Galais have given the names *Lactoline* and *Lacteine*.

MILK SPOTS, *Maculæ albæ*, Strophulus.

MILK STONE, Morochthus.

MILK, SUGAR OF, Sugar of milk.

MILK OF SULPHUR, Sulphur præcipitatum—m. Sweet, Galium verum.

MILK Teeth, see Dentition—m. Thrush, Aphthæ—m. Vetch, Astragalus verus—m. Vetch, stemless, Astragalus exscapus—m. Weed, Apocynum androsaemifolium, Asclepias Syriaca, Euphorbia corollata—m. Weed, long-leaved, green, Acerates longifolia—m. Weed, smooth, Asclepias Sullivantii—m. Wort, bitter, Polygala amara—m. Wort, common, Polygala—m. Wort, dwarf, Polygala paucifolia—m. Wort, rattlesnake, Polygala.

MILKWOMAN, Nurse, wet.

MILKY, Lactic.

MILL MOUNTAIN, Linum catharticum.

MILLEFEUILLE, Achillea millefolium—m. Aquatique, Phellandrium aquaticum.

MILLEFEUILLET, Osmorus.

MILLEFOLIUM, Achillea millefolium.

MILMEMORBIA SCROPHULARIA, Scrophularia nodosa.

MILLEPEDES, Onisci aselli.

MILLEPERTUIS ORDINAIRE, Hypericum perforatum.

MILLET, Aphthæ, Miliary fever—m. Barbadoes, Panicum Italicum—m. Common, Panicum miliaceum—m. Indian, Panicum Italicum—m. des Oiseaux, Panicum Italicum—m. Ordinaire, Panicum miliaceum.

MILLET-SEED RASH, Miliary fever.

MILLIGRAMME, from mille, 'a thousand, and γραμμα, 'a gramme.' The thousandth part of a gramme;—about 0.0154 Troy grain.

MILLIGRANA, Herniaria glabra.

MILLIMÈTRE, Millim'eter, from mille, 'a thousand,' and μετρον, 'measure.' The thou-

ndth part of the metre;—equal to 0.03937 English inch, or about two-fifths of a line.

MILLOT, Miliary fever.

MILPHÆ, Madarosis.

MILPHOSIS, Madarosis.

MILT, Melt, Spleen.

MILT-LIKE. An epithet for a form of cancer, characterized by a soft pale growth, containing scarcely any blood;—so called from its resemblance to the milt of a fish.

MILT-LIKE TUMOUR, Enecephaloid.

MILTUS, Plumbi oxydum rubrum.

MILTWASTE, Asplenium.

MIMOSA CATECHU, Catechu—*m. Cochliacarpa*, see *Cortex adstringens Brasiliensis*—*m. Leucophea*, see *Spirit (Arrack)*—*m. Nilotica*, see *Acacia gummi*—*m. Scandens*, *Cachang-parang*—*m. Senegal*, see *Senegal, gum*—*m. Virginalis*, see *Cortex adstringens Brasiliensis*.

MIND, **ABSENCE OF**, *Aphelxia socors*—*m. Abstraction of*, *Aphelxia intentia*—*m. Unsound*, *Insanity*.

MINERALIUM, Antimonium.

MINER'S ELBOW. An enlargement of a bursa over the olecranon, occurring in such as are in the habit of leaning much upon it; and, therefore, often seen in those who work on the side in low-roofed mines.

MIN'IM, *Min'imum*, 'the least part or portion.' The 60th part of a fluidrachm. This measure has been introduced by the London College of Physicians, in consequence of the uncertainty of the size of the drop, (see *Gutta*.) The subdivision of the wine pint has, accordingly, been extended to the 60th part of the fluidrachm; and glass measures, called "minim-measures," have been adopted by the London College. The proportion between the minims and the drops of various fluid preparations is exhibited in the following table. The results were obtained by Mr. Durand, a skilful *pharmacien* of Philadelphia, under circumstances, as regards the different articles, as nearly identical as possible.

TABLE OF THE NUMBER OF DROPS OF DIFFERENT LIQUIDS EQUIVALENT TO A FLUIDRACHM.

Acid Acetic, crystallizable.....	120
Acid Hydrocyanic (medicinal).....	45
— Muriatic.....	54
— Nitric.....	84
— — diluted (1 to 7).....	51
— Sulphuric.....	90
— — aromatic.....	120
— — diluted (1 to 7).....	51
Alcohol (rectified Spirit).....	138
Alcohol, diluted (proof Spirit).....	120
Arsenite of Potassa, solution of.....	57
Ether, Sulphuric.....	150
Oils of Aniseed, Cinnamon, Cloves, Peppermint, Sweet Almonds, Olives.....	120
Tinctures of Asafoetida, Foxglove, Guaiac, Opium.....	120
Tincture of Chloride of Iron.....	132
Vinegar, distilled.....	78
— of Colchicum.....	78
— of Opium (black drop).....	78
— of Squill.....	78
Water, distilled.....	45
Water of Ammonia (strong).....	54
Do. do. (weak).....	45
Wine (Teneriffe).....	78
— Antimonial.....	72
— of Colchicum.....	75
— of Opium.....	78

MINIM'ETER; badly compounded from *minim* and *perpo*, 'measure.' An instrument for measuring minims, invented by Mr. Alsop. It consists of a glass tube, graduated from the con-

ical point into minims; and having a piston, by the elevation of which, fluid may be drawn into the tube, and by its depression be forced from it.

MINIM MEASURE. A measure for the divisions of a fluidrachm.

MINIMUM, Minim.

MINISTER GYMNASTÆ, *Pædotribes*.

MINITHOSIS, Meiosis.

MINIUM, Plumbi oxydum rubrum—*m. Græcorum*, *Hydrargyri sulphuretum rubrum*—*m. Purum*, *Hydrargyri sulphuretum rubrum*.

MINOR HEL'ICIS. A muscle situate on the posterior border of the helix of the ear, at its commencement in the fossa of the concha.

MINORATIVUS, Laxative.

MINT, **COCK**, *Tanacetum balsamita*—*m. Curled*, *Mentha crispa*—*m. Cat*, *Nepeta*—*m. Horse*, *Monarda coccinea* and *M. punctata*—*m. Horse*, hairy, *Blephilia hirsuta*—*m. Horse*, Ohio, *Blephilia hirsuta*—*m. Horse*, sweet, *Cunila Mariana*—*m. Hyssop-leaved*, *Mentha cervina*.

MINT JULEP. A drink, consisting of brandy, sugar, and pounded ice, flavoured by sprigs of mint. It is an agreeable alcoholic excitant.

MINT MOUNTAIN, *Monarda coccinea*—*m. Mountain*, common, *Pycnanthemum incanum*—*m. Pepper*, *Mentha piperita*—*m. Spear*, *Mentha viridis*—*m. Squaw*, *Hedeoma*—*m. Stone*, *Cunila Mariana*.

MINT TEA, *Infusum Menthæ viridis*.

MINT WATER, *Mentha aquatica*.

MINTHA, Mint.

MINUTHESIS, Meiosis.

MINYANTHES, *Menyanthes*.

MIOSIS, Meiosis.

MIRLES, *Rubeola*.

MIROCELE, *Merocele*.

MIRROT, *Daucus Carota*.

MISADIR, *Ammonia murias*.

MISANTHRO'PIA, *Misan'thropy*, *Exanthro'pia*, *Phyganthro'pia*, from *μῖσος*, 'hatred,' and *ἄνθρωπος*, 'man.' Aversion to man and society;—a symptom of melancholy, and hypochondriasis.

MISCARRIAGE, Abortion.

MISERERE MEI, *Ileus*.

MISHME TEETA, *Coptis teeta*.

MISMAR, *Mesemar*. A name given by Avicenna to a kind of nodus, which forms on the toes as the sequel of contusion or inflammation of those parts.

MISOPTO'CHUS, from *μῖσος*, 'hatred,' and *πῶχος*, 'poor.' That which has hatred for the poor. The gout has been so called by some, because it commonly affects the rich.

MISSADAN, *Hydrargyrum*.

MISSIO SANGUINIS, Blood-letting.

MISSISSIP'PI, **MINERAL WATERS OF**. *Cooper's Well*, in Hinds County, four miles from Raymond, and thirteen from Jackson, is a sulphuretted saline, the chief constituents being the sulphates of soda and magnesia. Of the solid ingredients upwards of 3 per cent. are of peroxide of iron. The waters are chiefly used in chronic gastro-enteric affections. *Ocean Springs*, in Jackson County, is an acidulo-sulphureous chalybeate.

MISTIO, *Mistura*.

MISTLETOE, *Viscum album*—*m. Yellowish*, *Viscum flavesens*.

MISTURA, *μῖξις*, *Propotisma*, *Migma*, *Mis'tio*, *Mix'tio*, *Mixtu'ra*, *Mixture*; (F.) *Potion*, from *miscere*, *mixtum*, 'to mix.' A mingled compound, in which different ingredients are contained in



Minim measure.

the fluid state; suspended or not by means of mucilaginous or saccharine matter. In this sense, it is synonymous with the French *Potion*. In France, however, the word *mixture* is more frequently understood to mean a liquid medicine, which contains very active substances, and can only be administered by drops. A mixture, in other words, in the French sense, may be regarded as a *potion* deprived of watery vehicle.

MISTURA, (S.) A fragrant yellow-coloured water, used as a perfume by the ladies of Peru. It is prepared from gillyflower, jasmine, and *flor de mistela* (*Talinum umbellatum*). See *Campomanesia lineatifolia*.

MISTURA ACACIÆ, *Mucilago Acaciæ*.

MISTURA AC'IDI HYDROCYANICI, *Mixture of Prussic acid, Mélange pectoral* (Magendie). (*Medicinal prussic acid*, ʒij, distilled water, ʒxiv, pure sugar, ʒss.) A dessert-spoonful every morning and evening, as a pectoral, &c.

MISTURA AMMONIACI, *Lac ammoniaci*, *Mixture of ammo'niac*. (*Ammoniac*, ʒij, aqua Oss. Rub the ammoniacum: adding the water gradually until they are perfectly mixed. Ph. U. S.) Dose, fʒss to ʒj.

MISTURA AMYGDALÆ, *Emulsio amygdalæ*.

MISTURA ASAFCETIDÆ, *Lac asafetidæ*, *Emulsio antihyster'ica*, *Mixture of asafetida*. (F.) *Lait d'asafetida*. (*Asafetida* ʒij, aqua Oss. Rub together, adding the water by degrees. Ph. U. S.) Used where pills cannot be swallowed, and as a glyster in irritations during dentition, and in ascariæ. Dose, fʒss to fʒss.

MISTURA ASAFCETIDÆ PRO CLYSMATE, *Enema fetidum*.

MISTURA CAM'PHORÆ, *Aqua camphoræ* (Ph. U. S.), *Camphor mixture* or *julep*, *Mistura camphorata*, *Julep'ium e' camphoræ seu camphoratum*, (F.) *Julep camphré*. (*Camphor*, ʒij, alcohol, ℥xl, magnes. carb. ʒiv, aqua Oij. Rub the camphor with the alcohol, and afterwards with the magnesia, add the water, and filter. Ph. U. S.) Virtues like those of camphor. See *Emulsio camphoræ*.

MISTURA CORNU USTI, *Decoctum cornu cervini seu album*, *Decoction of burnt hartshorn*, *Mixture of burnt hartshorn*. (*Cornuum ust.* ʒij, acaciæ gum. cont. ʒij, aqua Oij. Boil to Oij, constantly stirring, and strain. Ph. L.)

MISTURA CREASOTI, *Creasote mixture*. (*Creasot.*, Acid. acetic. aa ℥xv, Sp. Junip. comp., Syrup., aa fʒj; Aqua fʒxiv. Mix the creasote and the acid; add gradually the water, and then the syrup and spirit. Ph. E.) A fluidounce contains a minim of creasote.

MISTURA CRETÆ, *Potio carbonat'is calcis*, *Mistura creta'cea*, *Mucilago cret'ica*, *Chalk* or *Creta'ceous mixture*, *Chalk julep*, (F.) *Mixture de Craie*. (*Creta* pp. ʒss, sacchar., acaciæ gum. in pulv., aa ʒij, aqua cinnam., aqua, aa fʒiv. Mix. Ph. U. S.) Antacid and absorbent. Dose, fʒj to fʒij.

MISTURA DIABOLICA. A mixture under this name is kept in military hospitals for maligners. It is made of sundry nauseous ingredients, as aloes, asafetida, castor, &c., and is given so as to keep up a disagreeable impression on the gustatory nerves.

MISTURA EMETI'NÆ VOMITO'RIA, *Emetic mixture of emetine*. (*Emetine* 4 gr., orange flower water ʒij, syrup ʒss; M.) A dessert-spoonful every half hour till it acts.

MISTURA FERRI COMPOSITA, *Compound mixture of iron*, (F.) *Mixture de fer composée*, *Griffith's mixture*. (*Myrrh. cont.*, ʒj, potassæ carb. gr. xxv, aqua rosæ fʒviiss, ferri sulph. in pulv., ʒj, sp. lavand. ʒss, sacchar. ʒi. Rub together the myrrh. subcarbonate of potassa, and sugar: then add, while triturating, the rose-water, spirit of nutmeg; and, lastly, the sulphate of iron.

Pour the mixture directly into a glass bottle, and stop it close. Ph. U. S.) It is tonic and emmenagogue, and is useful wherever iron is indicated.

MISTURA GLYCYRRHIZÆ COMPOSITA, *Compound mixture of Licorice*, *Brown Mixture*, (*Extract. Glycyrrhiz. pulv.*, Acaciæ pulv., Sacchar., aa ʒss; Tinct. opii camphorat. fʒij; Vin. antimon. fʒj; Spirit. æther. nitric. fʒss; Aqua fʒxij. Rub the liquorice, gum arabic, and sugar with the water, gradually poured upon them; add the other ingredients and mix. Ph. U. S.) A popular cough medicine, but not deserving of being made official. Dose, a tablespoonful or fʒss.

MISTURA GUAIACI, *Mixture of guai'ac*, *Lac guaiaci*, *Emulsio guaiaci'na*, *Mistura guaiaci gummosa seu gummi guaiaci Ber'geri*, *Solutio guaiaci gummosa seu resina' guaiaci aquosa*, (F.) *Mixture de Guyac*. (*Guaiaac. g. resin.* ʒiss, sacch. pur. ʒij, muc. acaciæ gum. ʒij, aq. cinnam. fʒviij. Rub the guaiacum with the sugar; then with the mucilage. Add, gradually, the cinnamon water. Ph. L.) Dose, fʒss to fʒij.

MISTURA GUMMI GUAIACI BERGERI, *Mistura Guaiaci*—m. Hordei, *Decoctum Hordei compositum*—m. Leniens, Looch ex ovo.

MISTURA MOSCHI, *m. moscha'ta*, *Musk mixture*. (*Moschi, acaciæ gum. contus.*, sacch. purif., sing. ʒj, aq. rosæ fʒvj. Rub the musk with the sugar; add the gum, and, by degrees, the rose-water. Ph. L.) Dose, fʒss to fʒij.

MISTURA SCAMMONII, *Scammony Mixture*. *Scammon. gr. viij, Lactis fʒij*. Triturate to form an emulsion. Ph. E.) The whole is a cathartic dose.

MISTURA SPIRITUS VINI GALLICI, *Brandy Mixture*. (*Sp. Vin. Gallic.*, Aq. cinnam. aa fʒij; Vitel. ovo. ʒj; Sacchar. ʒss; Ol. cinnam. ℥ij. M. Ph. L.) An excitant draught for the low stages of fever.

MISTURA STRYCHNINÆ, *Mixture of strychn'ia*. (*Distilled water*, ʒij, very pure strychnina, 1 gr., white sugar, ʒij; M.) Dose, a dessert-spoonful.

MISTURA SULPHURICO-ACIDA, *Elixir acidum Halleri*—m. *Vulneraria acida*, *Arquebusade*, Eau d'.

MITCHELLA REPENS, *Partridge berry*. A pretty little indigenous trailing evergreen, of the *Cinchona family*—*Cinchonaceæ*,—which flowers from June to July. It has been considered an expectorant, emmenagogue, and diuretic; and has been prescribed in dropsy and gout.

MITELLA, Sling.

MITHRIDATE, *Mithridat'ium*, *Mithridat'icum medicamentum*, *Antidotum Mithridat'ium*, *Diascin'ci antidotus*, *Confectio Damoc'ratia*. A very compound electuary, in which entered—Myrrh of Arabia, Saffron, Agaric, Ginger, Cinnamon, Frankincense, Garlic, Mustard, Birthwort, Galbanum, Castor, Long pepper, Opoponax, Beldium, Gum Arabic, Opium, Gentian, Orris, Sagenum, Valerian, Acacia, Hypericum, Canary wine, Honey, &c. It was invented by Mithridates, king of Pontus and Bithynia, and was formerly regarded as alexipharmic. It is little used at the present day, and, from its heterogeneous nature, should be wholly abandoned.

MITHRIDATICUM MEDICAMENTUM, *Mithridate*.

MITIGANS, Lenitive.

MITRA HIPPOCRATICA, *Bonnet d'Hippocrate*.

MITRAL, *Mitra'lis*, from *μῆτρον*, *mitra*, 'a headdress.' Having the form of a mitre; resembling a bishop's mitre.

MITRAL REGURGITATION, see *Regurgitation*.

MITRAL VALVES, *Valvula mitral'is seu Cordis mitral'is seu episcopales seu bicuspidales seu bicuspidatæ*. Two triangular valves at the open-

ing of communication between the left auricle of the heart and the corresponding ventricle. These valves are formed by the inner membrane of the left cavities of the heart; and are retained on the side of the ventricle by tendinous cords, proceeding from the columnæ carneæ. They form a species of valve, which permit the blood to pass from the auricle into the ventricle and oppose its return.

MITRIUM, Sling.

MITTE (F.), *Effluvium Latrina'rium*. An emanation exhaled from privies, which strongly irritates the eyes. It consists of ammonia, united to the carbonic and hydro-sulphuric acids.

MIUAMARU, see Arrow poison.

MIVA, Marmelade.

MIXIS, Coition, Mistura.

MIXTIO, Mistura.

MIXTURA, Mistura.

MIXTURE, Mistura—m. Almond, Emulsio Amygdalæ—m. of Ammoniac, Mistura ammoniaci—m. of Asafetida, Mistura asafetidæ.

MIXTURE, BRANDY, Mistura spiritus vini Gallici—m. Brown, Mistura glycyrrhizæ composita—m. Calnante, Looch ex ovo.

MIXTURE, CAMPHOR, Mistura camphoræ—m. Chalk, Mistura cretæ.

MIXTURE DE CRAIE, Mistura cretæ—m. of Creasote, Mistura creasoti.

MIXTURE, CRETACEOUS, Mistura cretæ.

MIXTURE OF EMETINE, (emetic,) Mistura emetinæ vomitoria—m. de Fer composée, Mistura ferri composita—m. de Gayac, Mistura Guaiaci.

MIXTURE, GRIFFITH'S, Mistura ferri composita.

MIXTURE OF GUAIAC, Mistura Guaiaci—m. of Gum Arabic, see Emulsio Acaciæ—m. of Harts-horn, burnt, Mistura cornu usti—m. of Iron, compound, Mistura ferri composita.

MIXTURE, HOPPE'S. A mixture recommended by Mr. Hope in diarrhœa and dysentery. It is essentially a nitrate of morphia. Although the proposer employed the acidum nitrosum of the Edinburgh Pharmacopœia, nitric acid is generally used. (*Acid. nitros. f 3j; Aque camphor. f 3viij; Tinct. opii gutt. xl.; M.*) Dose, a fourth part every three or four hours.

MIXTURE OF LIQUORICE, COMPOUND, Mistura glycyrrhizæ composita—m. *Lithontriptique de Durande*, see Liquor anodynus terebinthinatus—m. of Marsh mallows, Decoctum althææ.

MIXTURE, MUSK, Mistura moschi—m. Neutral, Liquor potassæ citratis—m. of Prussic acid, Mistura acidi hydrocyanici.

MIXTURE OF SCAMMONY, Mistura scammoni.

MIXTURE, SCUDAMORE'S. A mixture recommended by Sir C. Scudamore in gouty and rheumatic affections, and much prescribed in the United States. *Magnes. sulph. 3i—3ij; Aqua mentha f 3x; Aceti colchic. f 3j—f 3iiss; Syrup croci f 3j; M.* Dose, one, two, or three tablespoonfuls, repeated every two hours in a paroxysm of gout, until from four to six evacuations are produced in the twenty-four hours.

MIXTURE OF STRYCHNIA, Mistura strychniæ—m. de Whitt, see Liquor anodynus terebinthinatus.

MNEMĒ, Memory.

MNEME CEPHALICUM BAL'SAMUM. A very compound medicine, into which entered Balm, Lily, Rosemary, Lavender, Borage, Broom, Roses, Violet, Saffron, Thyme, Storax, Galbanum, &c.

MNEMONEU'TICE, Mnemonics.

MNEMON'ICS, *Mnemou'icē, Mnemoneu'ticē, Mnemotechnē, μνημη, 'memory' μνημονικος, 'appertaining to memory,'* from μνασθαι, 'I recollect.' The art of aiding the memory; and of forming, in some sort, an artificial memory.

MNEMOTECHNE, Mnemonics.

MOANING, from Sax. *mænan*, 'to grieve.'

Respiratio luctuosæ, Mychthis'mos. A plaintive

respiration, in which the patient utters audible groans,—*moans*, (F.) *Gémissements*.

MOBILE, see Saint Augustine.

MOBILITY, *Mobil'itas*, from *mob'ilis*, contraction of *movibilis*, 'that can be moved.' The power of being moved. In *physiology*, great nervous susceptibility, often joined to a disposition to convulsion. Greatly developed excitability.

MOCCASIN FLOWER, *Cypripedium acaule*, *C. luteum*—m. Plant, *Cypripedium luteum*.

MOCH'LIA. Reduction of a luxated bone; from *μολος*, a lever.

MOCHLICUS, Purgative.

MOCHLISCUS, Lever.

MOCHLUS, Lever.

MOCHTHUS, Agony.

MOCK-KNEES, *Crusta genu equinæ*.

MOD'ELLING PROCESS. A term proposed by Dr. Macartney, of Dublin, to signify the mode in which wounds are healed without inflammation or suppuration, by a deposit of plastic matter from the surface of the wound, by which the gap is more rapidly filled,—portion being laid upon portion, without waste, after the manner of clay in the hands of the sculptor;—hence the term.

MOD'OLUS, 'the nave of a wheel.' *Pyr'amis seu Axis seu Columella coch'leæ*. A hollow cone in the cochlea of the ear, forming a nucleus, axis, or central pillar, round which the gyri of the cochlea pass. The modiolus forms the inner and larger portion of the central pillar, and is the cavity seen at the bottom of the meatus auditorius internus. It lodges a branch of the auditory nerve. The central portion of the modiolus contains a number of minute canals, and is called in consequence *Tractus spiralis foraminulosus*. Into these the nerves of the cochlea enter, and pass out at right angles between the bony plates forming the zona ossea of the lamina spiralis, to be expanded on the membranous portion of the lamina.

MODIOLUS, *Mas*, also means the crown of the trepan.

MODIRA CANIRAM, *Strychnos colubrina*.

MO'DIUS. A dry measure, the third of an amphora, equal to the bushel.—Varro.

MODUS OPERAN'DI, 'mode of operating.' This term is applied to the general principles upon which remedies act in morbid states of the body;—a subject of much interest, although involved in considerable obscurity.

The following classification will convey some idea of the ways in which different organs may be excited into action.

1. By actual or immediate contact of the remedy, and by absorption or mediate contact.
2. By an impulse conveyed by the nerves, through an impression made on the stomach or elsewhere.
3. By contiguous or continuous sympathy, or by mere proximity or continuity of parts.

MOËLLE, Marrow—m. *Allongée*, Medulla oblongata—m. *Épinière*, Medulla spinalis—m. *Vértébrale*, Medulla spinalis.

MENIA DENTIMUM, Alveoli dentium.

MEROR, Athysia.

MOFETTE, Azote, see Mephitism.

MOFFAT, MINERAL WATERS OF. Moffat is a village situate about 56 miles S. W. of Edinburgh. The water is a cold sulphureous; containing, in a wine gallon, 36 grains of chloride of sodium, 5 cubic inches of carbonic acid, 4 of azote, and 10 of sulphohydric acid. It resembles Harrogate water in its properties.

A strong chalybeate spring was discovered there about the year 1828, which contains sesquisulphate of peroxide of iron, sulphate of alumina, and uncombined sulphuric acid.

MOGIGRAPHIA, Cramp, writers'.

MOGILA LIA, *Parulid'ia*, from *μῆνις*, 'with difficulty,' and *λαλέω*, 'to speak.' This word is inaccurately written by Paul Zacchias,—*Mola'ia*, and *Molila'ia*. Difficult or defective articulation. Impediment of speech.. See Balbuties.

MOGOSTOCIA, Dystocia, Laborious labour.

MOGUS, Agony.

MOHAWK TASSEL, Eupatorium purpureum.

MOHRIA THURIFRAGA. A South African fern. *Nat. Ord.* Filices, the dry leaves of which, pulverized and made into an ointment with fat, are serviceable in burns and scalds.

MOIGNON, Stump.

MOIS, Menses.

MOISTURE, *Mador*, (F.) *Moiteur*, anciently *moisteur*. Simple humidity of the skin. The skin is said to be *moist* or in a state of moisture, when there is slight perspiration.

MOITEUR, Moisture.

MOLA, Maxillary bone, Molar tooth, Mole, Patella.

MOLAGOCODI, Piper nigrum.

MOLAIRE, Molar.

MOLALIA, Mogilalia.

MOLAR, *Mola'ris*, *My'licus*, (F.) *Molaire*, *Meulière*. That which bruises or grinds; from *molaris*, 'a grindstone,' or *mola*, 'a millstone,' (*mola*, 'I grind.')

MOLAR GLANDS. Two small bodies formed by a collection of mucous crypts; seated in the substance of the cheeks, between the masseter and buccinator muscles, and whose excretory ducts open into the mouth opposite the last molar tooth.

MOLAR TEETH, *Grinders*, *Jaw Teeth*, *Mola'res permanentes dentes*, *Dentes molares seu maxillares seu gom'phii seu clava'les*, *Molz*, *Gomphioi*, *My'lacri*, *Mylodontes*, *My'lodi*, *Momis'ci*, *Lag-teeth*, *Wall teeth*, (Prov.) *Azzle teeth*, *Azle teeth*, *Cheek teeth*, *Wang teeth*, (F.) *Dents molaires ou mâchelières*, occupy the farther part of each alveolar arch. Their coronæ are broader than they are high; unequal; tuberculated; and the roots are more or less subdivided. They are 20 in number, 10 in each jaw. The first two pairs of molar teeth in each jaw have been called *lesser molares*, *premo'lar* or *bicuspid*, *Dentes bicuspidati seu bucca'les*. The coronæ have two tubercles; the outer being more prominent and larger than the inner. The root is commonly simple; sometimes bifurcated. The other three pairs have been termed, *Great molares*—*Dentes multicuspidées* (Ch.) Their coronæ are cubical, rounded, and have, at the upper surface, four or five tubercles, separated by deep furrows. The root is divided into two, three, four, and five *fangs*, which are more or less curved. The third great molaris appears a long time after the others, and hence has been called *Dens sapient'ie* or *Wisdom tooth*.

MOLASSES, Melasses.

MOLE, *Mola*, from *moles*, 'mass.' *Mylē*, *Zephyr'ius fœtus*, *Germen falsum*, *Pseudoceph'is molaris*, *Gravid'itis molaris*, *Mooncalf*, old French, *Furdeau*. A fleshy, insensible, at times, soft—at others, hard—mass; of variable and determinate size, which forms in the uterus, and is slightly united by vessels to that organ; from which its feeble vitality is derived. It has been conceived by some to be always owing to imperfect conception; but moles may form in the undoubted virgin. They seem to be owing to a morbid process; and certainly are generally connected with conception. At times, they contain parts of the fœtus; but commonly do not. At very different periods, in different women, the diseased mass is expelled from the uterus, with ordinary symptoms of abortion; and the case requires similar

management. See, also, *Nævus*, and *Conception*, (false.)

MOLE PLANT, Euphorbia lathyris.

MOLECULAR, *Molecula'ris*. Of or belonging to molecules or minute portions of any thing. Hence *molecular death*, in contradistinction to *somatic death*.

MOL'ECULE, *Molec'ula*, *Mass'ula*; diminutive of *moles*, 'a mass.' A minute portion of any body. Also, the *cicatric'ula*, *ma'cula*, *gallatu'ra*, *gelatinous molecule*, *tread of the cock*, or embryo part of the impregnated ovum, observable by the microscope before the ovum has left the ovarium of the hen. It lies under the epidermic coats of the yolk, and upon its proper coat. If the ovum, according to Valentin, be lacerated and its contents minutely examined, the cicatricula is found like a grayish white disk, which in its whole periphery is dense, granulous, and opaque; but in the centre presents a clear, nongranulous, and perfectly diaphanous point. Purkinje found, that when he removed the dark granulous mass by suction with a small tube, there remained a perfectly transparent vesicle filled with a pellucid lymph, which had a decidedly spherical form, but, being extremely delicate, was easily lacerated, and then its fluid escaped. As he found this, which later naturalists have named—after its discoverer—the *Purkin'jean ves'icle*, in the ova of the ovary, but could not see it in ova, which had already entered the oviduct, he gave it the name *germinal vesicle*, or *germ cell*, *vesic'ula prolifera seu germinati'va*, (F.) *Vesicula germinative*. Besides a perfectly colourless fluid, this contains one or more dark corpuscles, which appear as a nucleus through the including membrane in the shape of opaque spots—the *germinal spot*, *mac'ula germinati'va seu germinans*, *nucleus germinativus*, *Blasto-car'dia*, *Cor'culum germinis*, (F.) *Tache germinative*. The granulous membrane—its thickened portion, the so-called 'cicatricula,'—and the germinal vesicle, constitute those parts of the ovum which pass immediately into the original foundation of the embryo, the *blastoderm* or *germinal membrane*, *vesic'ula blastodermatica*, *blastoder'mic vesicle*.

MOLECULE, GELATINOUS, Molecule.

MOLÈNE, *Verbascum nigrum*.

MOLES GANGLIFORMIS, Ganglion of Gasser.

MOLIBDUS, Plumbum.

MOLILALIA, Mogilalia.

MOLIMEN, pl. MOLIM'INA, from *moior*, 'to move or stir.' An attempt, a struggle. Hence:

MOLIMEN CRIT'ICUM. An impulsion towards a sudden solution or crisis of a disease.

MOLIMEN HÆMORRHAG'ICUM. The hemorrhagic diathesis or impulsion.

MOLIMEN HÆMORRHOIDA'LE. The hemorrhoidal diathesis or impulsion. See *Hæmorrhoids*.

MOLIMEN MENSTRU'ALE. The menstrual diathesis or impulsion.

MOLIMEN NATURÆ SALUTARIUM, Instinct.

MOLKENKUR, Whey cure.

MOLLESCENCE, Mollities.

MOLLET, Sura.

MOLLIPUFFS, Lycoperdon.

MOLLIS MATER, Pia mater.

MOLLIT'IES, *Molles'cence*, *Malacis'mus*, *Mollit'ia*, *Mollitu'do*, *Emollit'ies*, *E. morbo'sa*, *Mollitio*, *Malax'is*, *Malaco'is*, *Malthar'is*, (F.) *Rammolissement*; (G.) *Malakien* (Fuchs); from *mollis*, 'soft.' Preternatural softness of an organ or part of an organ.

MOLLITIES CER'E'BRI, *Malaco'is Cer'e'bri*, *Encephalomala'cia*, *Encephalomalax'is*, *Necrenceph'alus* (W. Farr), *Cerebromala'cia*, *Encephalosep'is*, *Molles'cence* or *softening* or a kind of lique-

faction of the cerebral substance; the remainder preserving its ordinary consistency. (F.) *Ramollissement du cerveau*. The neurine often contains small clots of blood, giving rise to what has been termed *Apoplexie capillaire*, (Cruveilhier.) The symptoms denoting it are equivocal.

MOLLITIES CORDIS, Cardiomalacia.

MOLLITIES MEDULLÆ SPINALIS, *Myelomalacia*, (F.) *Ramollissement de la Moëlle Épinrière*. Softening of the spinal marrow.

MOLLITIES OSSIS, *Malacosteo-sis*, *Malacosteon*, *Emollit'ies ossium*, *Paros'tia flex'ilis*, *Osteomalaco'sis*, *Osteomala'cia*, *Rachi'tis seu Rachi'tis adultorum*, *Spina vento'sa* (of some), *Softening of the bones*, (F.) *Ostéo-malacie*, *Ostéo-malaxie*, *Ostéo-malakie*, *Ramollissement des os*. A rare affection, in which the bones are deprived of their salts, particularly of the phosphate of lime, and consist only, or mainly, of gelatin; hence they acquire a degree of suppleness which renders them unfit for the performance of their functions. The disease generally affects all the bones; but it is especially remarkable during life in the long bones, which assume any curvature that may be wished. Very violent pain is often experienced in them; and the urine frequently contains an enormous proportion of calcareous phosphate. The patient is compelled to remain in the horizontal posture; the bones no longer being fixed points for the muscles to act upon. The disease has, hitherto, always terminated fatally, and dissection has exhibited the gelatinous nature of the bones; which, by desiccation, have become transparent, as if they had been macerated in acid.

Experience has afforded no means of relief in this dreadful affliction. The alkaline salts, earthy phosphates, &c. are of little or no use.

MOLLITIO. Mollities.

MOLLITUDO. Mollities.

MOLLUSCUM, *Athero'ma*, *Ochthi'asis*. A cutaneous affection, so called in consequence of its resemblance to certain molluscous animals. It consists of numerous tumours, varying in size from that of a pea to that of a pigeon's egg, filled with an atheromatous matter, which are developed in the substance of the derma, and are of various shapes, some having a large base, — others adherent by means of a pedicle.

Molluscum contagiosum, (F.) *Acné molluscoïde*, a singular variety, is characterized by the presence of hard, round, tubercles, which are smooth and transparent, and when pressed, pour out from an orifice in their summits a little opaque or milky fluid.

The disease is probably seated in the sebaceous follicles.

If internal treatment be adopted at all, it must consist of eutrophics, as arsenic, iodine, &c. The external treatment consists in the employment of measures calculated to excite the tubercles to inflammation, as by touching them with potassa.

MOLOCHE, *Malva rotundifolia*.

MOLOPES, *Vibices*.

MOLYBDÆNUM MAGNESII, Manganese, black oxide of.

MOLYBDOCACHEXIA, Saturnismus.

MOLYBDOCOLICA, *Colica metallica*.

MOLYBDONOSUS, Lead-poisoning.

MOLYBDOPARESIS, Palsy, lead.

MOLYBDOS, Plumbum.

MOLYBIDOSIS, Lead-poisoning.

MOLYNE, Anus.

MOLYNSIS, Miasma.

MOMIE, Mummy.

MOMIN, Manei.

MOMISCI, Molar teeth.

MOMORDICA, *M. Balsami'na*, from *mordeo*, *monordi*, *morsum*, 'to bite,' *Balsami'na*, *Neuro-*

sper'ma cuspidata, *Balm-apple*, *Balsam apple*, (F.) *Balsamine*, *Pomme de merveille*. Ord. Cucurbitaceæ. The fruit of this oriental tree, *Pomum mirab'ile seu hicrocolymita'num*, was formerly considered vulnerary.

MOMORDICA ASPERA, *M. Elaterium*.

MOMORDICA ELATERIUM: *M. as'pera*, *Elate'rium*, *Echa'lia Elate'rium*, *Eebatium officina'rum seu agrestè seu cordifo'tium*, *Cucumis agrest'is seu asini'nus seu sylvest'ris*, *Charan'tia*, *Bouba'tios*, *Guarbera orba*, *Eebol'ium elate'rium*, *Wild*, *Squaring*, or *Asse's cucumber*, (F.) *Concombre Sauvage*, *Momordique*. Nat. Order, Cucurbitaceæ. The dried sediment from the juice is the elaterium of the shops. (See *Extractum Elaterii*.) It is a most powerful cathartic, and, as such, has been used in dropsies. Its active principle is the *Elatin*, *Elaterin* or *Elaterium*. Dose, gr. $\frac{1}{2}$ to gr. j. until it operates.

MOMORDIQUE, *Momordica elaterium*.

MONAD, from *monas*, *μονάδος*, 'unity.' A simple particle, or atom, or unit. — Leibnitz. The smallest of all visible animalcules. A primary cell or germ.

MONÆ, Nates.

MONARDA, *M. did'yma seu coccinea seu purpurea seu Kalmia'na*, *Scarlet Rose balm*, *Mountain mint*, *Oncel'go Tea*, *Mountain Balm*, *Horsemint*, *Square stalk*, *Red balm*; a beautiful indigenous plant, *Sex. Syst.* *Diandria Monogynia*. Nat. Ord. *Labiata*, having the excitant properties of *Monarda punctata*.

MONARDA ALLOPHYLLA, *M. fistulosa* — *m. Clinopodia*, *M. fistulosa* — *m. Coccinea*, *Monarda* — *m. Didyma*, *Monarda*.

MONARDA FISTULOSA, *M. rugosa* and *oblongata*; *M. clinopodia seu allophyl'la*. *Wild bergamot*, *Horsemint*. The leaves have a fragrant smell; and an aromatic, somewhat bitter taste. They are reputed to be nervine, stomachic, and deobstruent [?].

MONARDA KALMIANA, *Monarda* — *m. Oblongata*, *M. fistulosa*.

MONARDA PUNCTATA, *Horsemint*. Indigenous in the United States. Stimulant and carminative. The *Oleum monardæ* is officinal in the United States.

MONARDA PURPUREA, *Monarda* — *m. Rugosa*, *M. fistulosa*.

MONDER (F.), from *mundus*, 'cleanly.' To render clean or pure. In *Pharmacy*, it means to separate any substance from its impure or useless portion. In surgery, *monder* on *mondifier une plaie*, is to clean or debride a wound.

MONDIFICATIFS, Detergents.

MONDIFIER UNE PLAIE, see *Monder*.

MONE'MERON, *Monoe'meron*, *Monohemer'on*; from *monos*, 'one,' and *μερον*, 'a day.' A name given to several collyria, which were supposed to be capable of curing diseases of the eyes in a day.

MONENTERUM, Colon.

MONE'SIA. A vegetable extract imported into Europe from South America, in hard, thick cakes. It is prepared from the bark of a tree, whose botanical name is uncertain — probably *Chrysophyll'um glycyphall'un seu Buranhem*, *Buranhem*, *Guaranhem*; Ord. *Sapotaceæ*. It is very friable, and its fracture very much resembles that of a well-torrefied coconut. It is wholly soluble in water; and its taste — which is at first sweet like liquorice — sometimes becomes astringent. It is on account of its astringent properties that it has been prescribed in chronic bronchitis, hæmoptysis, diarrhœa, leucorrhœa, uterine hemorrhage, &c. It has been applied locally, in the form of ointment, (*Mones*, p. 1, *Adipis*, p. vij).

The dose of *Monesia* is from 12 to 40 grains a day.

MONEYWORT, *Lysimachia nummularia*.

MONGO'LIAN. Anthropologists give the name *Mongolian race* to a variety of the human species, spread over a great part of the north of Asia, in China, India, Thibet, the Moluccas, &c. The individuals composing it have the skin of a brown red, forehead flat, nose broad, cheeks prominent, and lips large. See Homo.

MONGREL, Hybrid.

MONIPLIES, Omasum.

MONISM, *Monis'mus*; from *μονας*, 'unity.' The doctrine, which declares matter and mind to be identical.

MONK PHYSICIANS AND SURGEONS. A class of practitioners of whom Frère Cosme and Jacques Beaulieu in France, and Pravetz in Germany, were the most distinguished.

MONKSHOOD, *Aconitum*—m. Common, *Aconitum napellus*—m. Salutory, *Aconitum anthora*—m. Wild, *Aconitum ucinatum*.

MONNI'NA POLYSTACHIA. A beautiful South American plant. *Nat. Ord.* Polygalæ. *Sex. Syst.* Diadelphia Octandria. The bark of the root is a powerful astringent, and much used in South America in diseases of the bowels.

MONNYPLIES, Omasum.

MONOBLEP'SIS, from *monos*, and *βλεψις*, 'sight.' An affection in which vision is confused, imperfect, and indistinct, when both eyes are employed; but perfect or nearly so, when either eye is used singly.

MONOCAR'PÆ, from *monos*, and *καρπος*, 'fruit.' A division of cutaneous affections, of the family of *Eczematosen*, of Fuchs, which includes strophulus, psyrdracia, and ecthyma.

MONOCEPH'ALUS, from *monos*, and *κεφαλη*, 'head.' A compound monster having two bodies with a single head.

MONOCOLON, Cæcum, Rectum.

MONOCRA'NUS, from *monos*, and *κρανον*, *κρανιον*, 'cranium.' A monster with one cranium, but with the face in part double.—Gurlt.

MONOCULUM, Cæcum.

MONOC'ULUS, *Mon'ocle*, *Monophthal'mus*, *Unio'culus*, *Unoc'ulus*, from *monos*, and *oculus*, 'an eye.' A bandage employed to maintain topical applications over one of the eyes. This bandage, called by some surgeons *Simplex oculi*, (F.) *Œil simple*, is made of a roller three or four ells long, rolled into a single ball. See Cyclops.

MONODIPLO'PIA, from *διπλous*, 'double,' and *ωψ*, 'eye.' Double vision with one eye.

MONOEMERON, Monemerom.

MONOGAS'TRIC, *Monogas'tricus*, from *monos*, and *γαστρικ*, 'stomach.' That which has but one stomach. Man is *monogastric*; ruminating animals, on the contrary, are *polygastric*.

MONOGENY, see Generation.

MONOGRAPH, MEDICAL, *Monograph'ia medica*, from *monos*, and *γραφη*, 'description.' An *ex professo* treatise on a single class of diseases, or on a single disease.

MONOHEMERON, Monemerom.

MONOMACHON, Cæcum.

MONOMACUM, Cæcum.

MONOMANIA, see Melancholy—m. Homicidal, Homicidal insanity—m. Incendiary, Pyromania.

MONOMA'NIAC, *Monomani'acus*, (F.) *Monomane*, *Monomaniacque*, from *monos*, and *μανια*, 'mania.' One who labours under monomania, or insanity on one subject.

MONOMANIE, Melancholy—m. Homicide, Homicidal insanity.

MONOMMATOS, Cyclops.

MONOMORIA, see Melancholy.

MONOM'PHALUS, from *monos*, and *μφαλος*, 'a navel.' A monster produced by the union of two bodies having a common umbilicus. Such a

double monster is called by I. G. St. Hilaire *Monomphal'ian*, (F.) *Monomphalien*.

MONOPAGIA, *Clavus hystericus*, Hemicrania.

MONOP'ATHY, *Monopath'ia*, from *monos*, and *παθος*, 'disorder.' A state in which one organ or function is disordered. Applied to melancholy or monomania; which is said to be a *monopathic* affection.

MONOPEGIA, *Clavus hystericus*, Hemicrania.

MONOPHTHALMUS, Cyclops, Monoculus.

MONOPLAS'TIC, *Monoplas'ticus*, from *monos*, and *πλασσω*, 'I form.' That which has one form. A *monoplastic element*, in histology is one which retains its primary form.—Gerber.

MONOPODIA, Symphodia.

MONOPS, Cyclops.

MONOPSIA, Cyclopia.

MONOPUS, Symmeles.

MONOR'CHIS, (F.) *Monorchide*, from *monos*, and *ορχις*, *ορχιδος*, 'testicle.' One who has only one testicle.

MONOS, *μονος*, one alone; as in

MONOS'CELUS, from *monos*, and *σκελος*, 'a leg.' One legged.

MONOSI'TIA, from *monos*, and *σιτος*, 'food,' 'repast.' The habit of taking only one meal in the day.

MONOSO'MIAN, (F.) *Monosomien*, from *monos*, and *σωμα*, 'a body.' An epithet applied to a double monster, which seems, in the first instance, to have had but one body.

MONOTROPA UNIFLO'RA, (*monos*, and *τροπος*, 'a turn,' the solitary flower turned to one side,) *Indian pipe*, *Icelandic pipe-plant*, *Nest root*, *Fit root*. *Ord.* Ericacæ. An indigenous plant, whose juice, mixed with water, has been extolled by the Indians in ophthalmia. The dried root in powder has been given in epilepsy and nervous diseases.

MONS VEN'ERIS, *Montic'ulus Ven'erie*, ('Mount of Venus,') *Ephebe'um*, (F.) *Mont de Vénus*, *Pénit*, *Motte*, *Eminence sus-pubienne*. The more or less projecting eminence, situate at the base of the hypogastrium, above the vulva and in front of the os pubis. At the period of puberty it becomes covered with hair, formerly called *Tresso'ria*.

MONSO'NIA OVA'TA, *Keita*, of the Hottentots. A plant of the *Nat. Ord.* Geraniacæ, which grows at the Cape of Good Hope. The root and herb are very astringent, and are used successfully in dysentery.

MONSTER, *Monstrum*, ('anything strange,') *Porten'tum*, *Teras*, *Pelor*, *Pelo'ria*, *Pelo'rimum*, *Pelo'rurum*, (F.) *Monstre*. Any organized being, having an extraordinary vice of conformation, or a preternatural perversion of every part, or of certain parts only.

The following classification embraces the main varieties of malformations—*Vitia primæ conformationis*.—1. Those in which certain parts of the normal body are absent or defective—*monstra defectiva*, constituting *Ectrog'eny* (Serres),—*Monstrosity by defect*, (F.) *Anomalie ou Monstruosité par défaut*, *Agénésie*. 2. Those produced by fusion or coalition of organs—*coalitio partium*, *sympphysis*. 3. Those, in which parts, united in the normal state, are separated from each other—*clefths*, *fissures*. 4. Those in which normal openings are occluded—*atresia*. 5. Those by excess, or in which certain parts have a disproportionate size—*monstra abundantia*, *Monstrosity by excess*, (F.) *Monstruosité par excès*, *Hypergénésie*. 6. Those, in which one or more parts have an abnormal position—*situs mutatus*. 7. Those affecting the sexual organs—*hermaphroditism*; and to these "true malformations" Vogel adds 8. Diseases of the fœtus, and abnormal states of its envelopes.

Amongst the numerous hypotheses entertained on the origin or cause of monsters, or of monstrosity—*teratosis*, *monstrositas*, (F.) *monstruosité*, as the state has been called—three only are worth mentioning. They have been attributed, 1. To the influence of the maternal imagination on the fœtus in utero. 2. To accidental changes, experienced by the fœtus at some period of its uterine existence: and 3. To a primitive defect in the germs. The second seems to be the only one that is philosophical.

As a medico-legal question, monsters, if capable of action as individuals, have the same rights as other persons.

MONSTERS, COMPOUND, see *Unitaire*—m. Double, see *Diplogenesi*—m. Triplet, see *Triplet*.

MONSTRA ABUNDANTIA, see *Monster*—m. Deficientia, see *Monster*—m. Trigemina, see *Triplet*—m. Triplica, see *Triplet*.

MONSTRE, *Monster*—m. *Composé*, see *Unitaire*—m. *Unitaire*, see *Unitaire*.

MONSTROSITAS, see *Monster*.

MONSTROSITY, see *Monster*—m. by Defect, see *Monster*—m. by Excess, see *Monster*.

MONSTRUM, *Monster*.

MONSTRUOSITÉ, see *Monster*—m. *par Défaut*, see *Monster*—m. *par Excès*, see *Monster*.

MONT DE VÉNUS, *Mons Veneris*.

MONTBRISON, MINERAL WATERS OF. Montbrison is a town in France, in the department of the Loire, fifteen leagues from Lyons. There are three cold springs here, which are highly acidulous, and used as refrigerants and aperients.

MONT-DE-MARSAN, MINERAL WATERS OF. Mont-de-Marsan is a small French town, 10 leagues from Dax; where there is a chalybeate spring, somewhat frequented.

MONT-D'OR, MINERAL WATERS OF. Mont-d'Or is a village seven leagues from Clermont, (Puy-de-Dôme,) at which are several mineral springs, cold and thermal. Those of the Fountain *La Madelaine*, and of the *Great Bath*, are thermal, temp. 112° Fahr., and contains carbonic acid, subcarbonate and sulphate of soda, chloride of sodium, subcarbonate of lime and magnesia, alumine and oxide of iron. Those of *St. Margaret* are cold, and contain much carbonic acid.

MONTECATINI, MINERAL WATERS OF. This town is situate near Borgo-Buggiano, in Tuscany. The springs are thermal; ranging from 78° to 118° Fahrenheit.

MONTHLY COURSES, *Menses*—m. *Illness*, see *Menses*—m. *Periods*, *Menses*.

MONICULUS CEREBELLI, see *Vermiform* processes of the Cerebellum—m. *Veneris*, *Mons veneris*.

MONTPELLIER, CLIMATE OF. This seat of a celebrated medical school in the S. E. of France was at one time renowned for its climate, and especially in regard to its adaptation to consumptive cases. This is now exploded, and it is at present esteemed an unfit residence for them. The climate possesses the general characters of that of south-eastern France—being dry, hot, and irritating; subject to frequent vicissitudes, and especially to blasts of keen, cold, northerly winds. It is, indeed, most injurious in pulmonary diseases.

MOON, *Luna*, *Sel'nē*, *Menē*, Sax. *Mona*, (G.) *Mond*, (F.) *Lune*. The moon has been supposed to exert considerable influence over the human body, in health and disease. Such influence has been grossly exaggerated. Not many years ago, it would have been heretical to doubt the exacerbation of mania at the full of the moon; yet it is now satisfactorily shown, that if the light be

excluded at this period, the *lunatic* is not excited more than ordinarily.

MOONED, *Lunatic*.

MOONSTRUCK, *Lunatic*.

MOONCALE, *Mole*.

MOORBERRY, *Vaccinium oxycoccos*.

MOORGRASS, *Potentilla anserina*.

MOORSHEAD, *Alembic*.

MOORWORT, BROAD-LEAVED, *Anaromeda mariana*.

MOOSEWOOD, *Dirca palustris*.

MOPHETA, see *Mephitism*.

MORBEUX, *Morbid*.

MORBI, see *Morbus*—m. *Acquisiti*, *Acquired diseases*—m. *Acuti*, see *Acute*—m. *Adventitii*, *Acquired diseases*—m. *Anniversarii*, *Annual diseases*—m. *Annui*, *Annual diseases*—m. *Asthenici*, *Adynamia*—m. *Celati*, *Feigned diseases*—m. *Chronici*, *Chronic diseases*—m. *Cognati*, *Innate diseases*—m. *Complicati*, see *Complication*, *Impliciti morbi*—m. *Congeniti*, *Innate diseases*—m. *Connati*, *Connate diseases*—m. *Connexi*, *Conjuncti morbi*—m. *Consequentes*, *Conjuncti morbi*—m. *Constrictorii*, see *Spasmodici*—m. *Dermatodes*, see *Dermatoid*—m. *Dispersi*, *Sporadic diseases*—m. *Dissimulati*, *Feigned diseases*—m. *Epicteti*, *Acquired diseases*—m. *Evolutionis*, see *Development*—m. *Hereditarii*, *Hereditary diseases*—m. *Homoedri*, see *Homoëdri*—m. *Infiati*, *Feigned diseases*—m. *Interni*, *Internal diseases*—m. *Melancholici*, see *Mercurial*—m. *Mercuriales*, see *Mercurial*—m. *Nervosi*, *Nervous diseases*—m. *Oculti*, see *Latent*—m. *Olfactūs*, *Osmonosi*—m. *Perplexi*, see *Complication*—m. *Pseudalei*, *Feigned diseases*—m. *Recidiva*, *Relapse*—m. *Recursus*, *Palindromia*—m. *Sexuales*, see *Sexual*—m. *Sexūs*, *Genonosi*—m. *Simulati*, *Feigned diseases*—m. *Sparsi*, *Sporadic diseases*—m. *Spasmodici*, see *Spasmodicus*—m. *Subacute*, see *Acute*—m. *Tactūs*, *Haphonosi*—m. *Tropici*, *Troponosi*.

MORBID, *Morbidus*, *Morbo'sus*, *Pathic*, *Pathical*, *Path'icus*, (F.) *Morbeux*, *Morbide*; from *morbus*, 'a disease.' Diseased, or relating to disease.

MORBID ANATOMY, see *Anatomy*.

MORBIFIC, *Morbif'icus*, (F.) *Morbifique*; from *morbus*, 'a disease,' and *facere*, 'to make.' Causing or producing disease—as *morbific emanations*.

MORBILITY, *Disease*.

MORBILLEUX, *Morbillous*.

MORBILLI, *Rubeola*—m. *Confluentes*, *Scarlatina*—m. *Ignei*, *Scarlatina*.

MORBILLOUS, *Morbillous*.

MORBILLOUS, *Morbillo'sus*, *Measly*, (F.) *Morbilleux*, from *morbilli*, 'rubeola or measles.' Affected or connected with measles.

MORBOSUS, *Sickly*.

MORBULENT, *Sickly*.

MORBUS, *Disease*, see *Morbi*—m. *Ampullaceus*, *Pemphigus*—m. *Anglicus*, *Phthisis*, *Rachitis*—m. *Aphrodisius*, *Syphilis*—m. *Arqueatus*, *Icterus*—m. *Arieti*, *Influenza*—m. *Arquatus*, *Icterus*—m. *Articularis*, *Gout*—m. *Astralis*, *Epilepsy*—m. *Attonitus*, *Apoplexy*—m. a. *Celsi*, *Catalepsy*—m. *Brightii*, *Kidney*, *Bright's disease* of the—m. *Bullosus*, *Pemphigus*—m. *Caducus*, *Epilepsy*—m. *Cæruleus*, *Cyanopathy*—m. *Calculi*, *Lithia*—m. *Cardiacus*, *Cardialgia*—m. *Catarrhalis*, *Influenza*—m. *Catoxys*, see *Catoxys*—m. *Cerealis*, *Ergotism*—m. *Cerebralis* *Whyttii*, *Hydrocephalus Internus*—m. *Cirrhorum*, *Plica*—m. *Cœliacus*, *Cœliac flux*—m. *Comitialis*, *Epilepsy*—m. *Cordis*, *Cardionosus*—m. *Costalis*, *Pleuritis*—m. *Coxarius*, *Coxarum morbus*—m. *Croatus*, *Scherlievo*—m. *Cucullaris*, *Pertussis*—m. *Cucullis*, *Pertussis*—m. *Cystopneumaticus*, *Cystirrheæ*—m. *Dæmoniacus*, *Epilepsy*—m. *Dæmonius*, *Epilepsy*—m. *Deificus*, *Epilepsy*—m.

Dissolutus, Dysentery—*m.* Divinus, Epilepsy—*m.* Dominorum, Gout—*m.* Endemius, Endemic—*m.* Epidemicus, Epidem'y—*m.* Epidemicus gutturi Foresti, Scarlatina—*m.* Eruditorum, Hypochondriasis—*m.* Extrinsicus, External disease—*m.* Farciniosus, see Equinia—*m.* Febrilis, Pyreticosis—*m.* Fellidus, Cholera—*m.* Flatusus, Hypochondriasis—*m.* Fluminensis, Scherlievo—*m.* Fædus, Epilepsy—*m.* Frigidus, Atrophy—*m.* Gallicus, Syphilis—*m.* Gesticulatorius, Chorea—*m.* Granulosus renum, Kidney, Bright's disease of the—*m.* Gravis, see Serious—*m.* Hæmorrhoidalis, Hæmorrhoids—*m.* Heracleus, Epilepsy—*m.* Herculeus, Epilepsy—*m.* Hispanicus, Syphilis—*m.* Hydrophobicus, Hydrophobia—*m.* Hypochondriacus, Hypochondriasis—*m.* Hystericus, Hysteria—*m.* Idiopathicus, Idiopathia—*m.* Indicus, Syphilis—*m.* Interlunus, Epilepsy—*m.* Ischiadicus, Neuralgia femoro-poplitea—*m.* Italicus, Syphilis—*m.* Jecinoris, Hepatitis—*m.* Lascivus, Chorea—*m.* Lateralis, Pleuritis—*m.* Lazari, Elephantiasis—*m.* Leseoli, Icterus—*m.* Lunaticus, Epilepsy—*m.* Luteolus, Icterus—*m.* Maculosus hæmorrhagicus Werlhofii, Purpura hæmorrhagica—*m.* Magnus, Epilepsy—*m.* Major, Epilepsy—*m.* Mensalis, Epilepsy—*m.* Mercurialis, Eczema mercuriale, Hydrargyriasis—*m.* Metallicus, Colic, metallic—*m.* Mucosus, Fever, adenomenigeal—*m.* Nauticus, Nausea marina—*m.* Neapolitanus, Syphilis—*m.* Niger, Melena—*m.* Originalis, Idiopathia—*m.* Oesophagi, Oesophagopathia—*m.* Oryzeus, Cholera—*m.* Pallidus, Chlorosis—*m.* Panconus, Pandemic—*m.* Parthenius, Chlorosis—*m.* Pandemius, Pandemic—*m.* Pedicularis, Phthiriasis—*m.* Peracutus, see Catoxys—*m.* Phœniceus, Elephantiasis—*m.* Pilaris, Trichiasis—*m.* Pleuriticus, Pleuritis—*m.* Plumbeus, Lead-poisoning—*m.* Porcinus, Essera—*m.* Popularis, Epilepsy—*m.* Primarius, Idiopathia—*m.* Proprius, Idiopathia—*m.* Protopathicus, Idiopathia—*m.* Proteiformis, see Proteiformis—*m.* Psodacis, Lumbar abscess—*m.* Publicus, Epidem'y—*m.* Puerilis, Epilepsy—*m.* Pulicularis, Typhus gravior—*m.* Pustulosus Finnicus, see Anthrax—*m.* Recidivus, Relapse—*m.* Recidivus, Palindromia—*m.* Regionalis, Endemic—*m.* Regius, Icterus—*m.* Resecatorius, Hypochondriasis—*m.* Ructuosus, Hypochondriasis—*m.* Ructuosus, see Eructation—*m.* Sacer, Epilepsy—*m.* Saluatorius, Chorea—*m.* Sancti Joannis, Epilepsy—*m.* Sancti Valentini, Epilepsy—*m.* Scarlatinus, Scarlatina—*m.* Scelestus, Epilepsy—*m.* Scropholus, Scrofula—*m.* Secundarius, Deuteropathia—*m.* Seleniacus, Epilepsy—*m.* Sideratus, Epilepsy—*m.* Silesiacus, Raphania—*m.* Simplex, Haplopathia—*m.* Solstitialis, *Coup-de-Soleil*—*m.* Sonticus, Epilepsy—*m.* Spasmodicus malignus seu popularis, Raphania—*m.* Strangulatorius, Cyanache trachealis—*m.* Sudatorius, Sudor Anglicus—*m.* Truculentus infantom, Cyanache trachealis—*m.* Tuberculosus, see Tubercle and Tubercular cachexia—*m.* Ungaricus, Fever, Hungarie—*m.* Verminosus, Helminthiasis—*m.* Vernaculus, Endemic—*m.* Vervecinus, Influenza—*m.* Vesicularis, Pemphigus—*m.* Virgineus, Chlorosis—*m.* Viridellus, Epilepsy—*m.* Vitriolatus, Epilepsy—*m.* Vocis, Phononosis—*m.* Vulpis, Alopecia.

MORCEAU DU DIABLE, see Tuba Fallopiana—*m.* *Frangé*, see Tuba Fallopiana.

MORCELLEMENT (F.), 'Dividing into pieces.' A mode adopted by *m.* Maisonneuve for removing successfully a large fibrous tumour of the neck.

MORCHELLA ESCULENTA, *Phallus esculentus*, *Helvella esculenta*, *Fungus fagino-us*, *Merulius*, Common Morell, (F.) *Morelle*. *Nat. Order*, Fungi. A fungus, which is esteemed a luxury, entering into ragouts and other dishes,

or cooked itself between slices of bacon. It has been considered to be aphrodisiac.

MOR'DICANT, *Mor'dicans*, *Calor mor'dicans*, from *mordeo*, 'I bite.' A morbid heat, causing a disagreeable pungent sensation in the fingers of the physician.

MORDICES, Teeth: see Tooth.

MOREA, *Morus nigra*.

MORELL, COMMON, *Morehella esculenta*.

MORELLE, *Boletus esculentus*—*m.* *d. Fruit noir*, *Solanum*—*m.* *Furieuse*, *Atropa belladonna*—*m.* *en Grappes*, *Phytolacca decandra*—*m.* *Grim-pante*, *Solanum dulcamara*.

MORETARIUM, Mortar.

MORE'TUM. A kind of salad, with the ancient Romans, made of garlic, parsley, vinegar, oil, &c.

MORE'TUS, from *morum*, 'the mulberry.' Ancient name of a cordial julep, into the composition of which the syrup of mulberries entered.—Schröder.

MORGAGN'I, HUMOUR OF, *Humor Morgagnianus*, (F.) *Humeur de Morgagni*. A peculiar, transparent, slightly-viscid fluid, found between the crystalline and its capsule.

MORGE LINE, Alsine media.

MORQUE (F.). A dead-house, in France, wherein persons, found dead, are exposed with the view of being recognised by their friends—*Necrocom'um*, *Necrodochi'um*, *Nec-ro's'toa*.

MORI SUCCUS, see *Morus nigra*.

MORIA, Idiôtism, *m.* Demens, Idiôtism, Dementia—*m.* Demens lerema, see Dementia—*m.* Imbecillis amnesia, Amnesia.

MORIBUND, Psychorages.

MORILLE, *Morchella esculenta*.

MORINGA APTERA, see *Guilandina moringa*—*m.* *Nux Ben*, *Guilandina moringa*—*m.* *Oleifera*, *Guilandina moringa*—*m.* *Pterygosperma*, *Guilandina moringa*—*m.* *Zeylanica*, *Guilandina moringa*.

MORION, Genital organ, Penis.

MORIOPLASTICŒ, *Chirurgia curto'rum seu Anaplastica*, *Restitutio organ'ica seu partium deperditarum*, *Transplantatio*, *Autoplas'tica*, *Chirurgia plas'tica*, *Anaplas'tic Surgery*, *Autoplas'ty*, *Autoplastic or Plastic Surgery*, from *μοιον*, 'a part,' and *πλαστικός*, 'forming.' The restoration of lost parts. The operations for this purpose have various names, according to the part concerned, as *Cheiloplastice*, *Oceratoplastice*, *Rhinoplastice*, &c.

MORNING GLORY, BLUE, *Convolvulus nil*.

MORO, from *morum*, 'a mulberry'—*Morum*, (F.) *Mure*. A small abscess resembling a mulberry. A small tumour, of a similar appearance, particularly on the genital organs after impure coition.

MOROCH'THUS, *Moroch'tus*, *Morochi'tes*, *Merochi'tes*, *Meroc'tes*, *Galax'ia*, *Leucograph'ia*, (F.) *Pierre au lait*, Milk stone. An Egyptian stone, used by the ancients as an application to the intertrigo of children: probably a variety of fuller's earth, inasmuch as it appears to have been used by fullers.

MOROCCHITES, *Morochthus*.

MOROCHTUS, *Morochthus*.

MOROCOMIUM, *Morotrophium*.

MORODOCHIUM, *Morotrophium*.

MOROSIS, *Fatuitas*, Idiôtism.

MOROSITATES, (*moros'us*, 'morose,' 'peevish,' 'fretful,') from *μωρία*, and *μωρωσις*, 'stupidity.' A generic name given by Linnaeus to an order of Venærie, in which he included pica, bulimia, polydipsia, antipathia, nostalgia, panophobia, satyriasis, nymphomania, tarentismus, and hydrophobia.

MOROTROPH'TUM, *Morocom'um*, *Morodo-*

ch'um, *Manicocomi'um*, from *μωρος*, 'fatuous,' and *τροφή*, 'support.' An insane establishment, lunatic asylum, mad-house, Bedlam, (F.) *Hôpital des fous*, *Maison petite*.

MORPHÆA, *Morphew*—m. Alba, *Lepra alba*—m. Nigra, *Lepra nigricans*.

MORPHÉE, *Morphew*.

MORPHEW, *Morphæa*, (F.) *Morphée*. A term vaguely applied to scurfy eruptions on the face. *Morphæa* was formerly applied to squamous diseases in general.

MORPHIA, *Morphina*—m. Acetate of, *Morphinæ acetas*—m. and Zinc, double iodide of, Zinc and Morphia, double iodide of—m. Citrate of, see *Morphina*—m. Muriate of, see *Morphina*—m. Sulphate of, *Morphinæ sulphas*.

MORPHIÆ ACETAS, *Morphinæ acetas*—m. Citras, see *Morphinæ sulphas*—m. Murias, see *Morphinæ sulphas*—m. Sulphas, *Morphinæ sulphas*.

MORPHICA, from *μωρφή*, 'shape.' Monstrosities of birth.' The 3d order, class *Typhica* of Good.

MORPHINA, *Morphia*, *Morphium*, *Morphin'ne*, *Morphine*, from *Μορφeus*, *Morpheus*, the 'god of sleep.' A solid, colourless alkali; crystallizable in beautiful pyramids, truncated and transparent; soluble in alcohol, and slightly so in boiling water. It exists, combined with meconic acid, in opium. It may be obtained by decomposing, by means of ammonia or calcined magnesia, an aqueous solution of opium made in the cold.

The following is the process of the Pharm. U. S. *Opium*, sliced, ℥j; *distilled water*, alcohol, animal charcoal, each a sufficient quantity; *solution of ammonia*, f℥vj. Macerate the opium with four pints of distilled water for 24 hours, and having worked it with the hand, digest for 24 hours, and strain. Macerate the residue twice successively with distilled water, and strain. Mix the infusions; evaporate to six pints, and filter; then add first five pints of alcohol, and afterwards three fluidounces of the solution of ammonia, previously mixed with half a pint of alcohol. After 24 hours, pour in the remainder of the solution of ammonia, mixed as before with half a pint of alcohol; and set the liquor aside for 24 hours, that crystals may form. To purify these, boil them with two pints of alcohol till they are dissolved; filter the solution, while hot, through animal charcoal, and set it aside to crystallize.

Morphia dissolves perfectly in the acids, which it saturates; and with which it forms very crystallizable salts, with the exception of the acetate. All these salts have a bitter taste, and act upon the animal economy in the same manner as opium, but more powerfully. The acetate is the salt commonly employed in medicine. It was, at one time, supposed that Morphia is the purely sedative part of opium, divested of its irritating properties; but experience has not confirmed this. On the contrary, it will generally disagree where opium does. Dose of morphia, $\frac{1}{4}$ to $\frac{1}{2}$ of a grain.

MORPHIÆ ACETAS, *Morphinæ Acetas*, *Morphium aceticum*, *Ac'etate of Morphine*, *Acetate of Morphia*. A salt formed by saturating morphia with acetic acid. (*Morphia*, in powder, freed from narcotina by boiling with sulphuric ether, ℥j; *distilled water*, Oss; *acetic acid*, a sufficient quantity. Mix the morphia with the water; then carefully drop in the acid, constantly stirring until the morphia is saturated and dissolved. Evaporate the solution in a water-bath to the consistence of syrup. Lastly, dry the acetate with a gentle heat and rub it into powder.—Ph. U. S.) Dose, from $\frac{1}{4}$ of a grain to half a grain.

MORPHIÆ SULPHAS, *Morphiæ Sulphas*, *Sulphate of Morphia*. A salt, formed by saturating morphia with sulphuric acid, evaporating, and crystallizing. Dose, the same as of the acetate.

Muriate, and *Citrate of Morphia*, *Morphia citras*, have likewise been used; but they possess no advantages over the other salts. The sulphate and muriate are official in the Ph. U. S. The muriate—*MORPHIÆ MURIAS*—is directed to be prepared as follows. (*Morphiæ*, in pulv., ℥j; *Aquæ destillatæ*, Oss; *Acid. muriat.* q. s. Mix the morphia with the water; carefully drop in the acid, till the morphia is saturated and dissolved. Evaporate and crystallize.) One-sixth of a grain is equal to a grain of opium.

MORPHIUM, *Morphina*—m. Aceticum, *Morphinæ acetas*.

MORPHOLOG'ICAL, *Morpholog'icus*, from *μωρφή*, 'shape,' and *λογος*, 'a description.' That which has relation to the anatomical conformation of parts. Applied, at times, to the alterations in the form of the several parts of the embryo, in contradistinction to *histological*, which is applied to the transformation by which the tissues are gradually generated. In comparative anatomy, it is applied to the history of the modifications of forms, which the same organ undergoes in different animals.

MORPHOLOGY, Anatomy.

MORPHOLYSIS, from *μωρφή*, 'shape,' and *λυσις*, 'solution.' Destruction of organization. An agent, that occasions such destruction, is called a *morpholytic*.

MORPHOLYTIC, see *Morpholysis*.

MORPHON'OMY, *Morphonom'ia*: from *μωρφή*, 'shape,' and *νομος*, 'a law.' The laws of organic formation. The department of anatomical science which teaches the laws of organic configuration.

MORPHOTOMY, see *Anatomy*.

MORPIO, see *Pediculus*.

MORPION, Crab-louse, see *Pediculus*.

MORRHUA AMERICANA, see *Oleum jecoris aselli*.

MORS, Death—m. Apparens, *Asphyxia*—m. Putativa, *Asphyxia*—m. Repentina, *Demotivus lapsus*—m. *du Diable*, *Scabiosa succisa*.

MORSELLUS, *Tabella*.

MORSULI ANTIMONIALES, *Tabellæ antimoniales Kunckellii*—m. Stibii, *Tabellæ antimoniales Kunckellii*.

MORSULUS, *Tabella*.

MORSURE, *Degmos*.

MORSUS, *Degmos*—m. Diaboli, *Scabiosa succisa*, see *Tuba Fallopiana*—m. Gallinæ, *Alsine media*—m. Stomachi, *Cardialgia*—m. Ventriculi, *Cardialgia*.

MORT, *Deaps*.

MORT, Dead, Death—m. Apparente, *Asphyxia*—m. *du Chien*, *Cholera*—m. *aux Chiens*, *Colicum autumnale*—m. *Né*, Still-born—m. *aux Panthères*, *Doronicum pardalianches*.

MORTA, *Pemphigus*.

MORTAL, *Mortal'is*, from *mors*, *mortis*, 'death.' *Lethal'is*, *Leta'lis*, *Thanato'ides*, *Thanas'imus*, (F.) *Mortel*. That which is subject to death;—that which causes death:—as, *man is mortal*; a disease is mortal. Of old, it was the custom to have a division of mortal or deadly wounds, which gave rise to many errors in medico-legal investigations, as the mortality of wounds depends upon various circumstances; and it is often a matter of extreme difficulty to pronounce whether or not a wound is necessarily mortal.

MORTAL'ITY, *Mortal'itas*, *Lethal'itas*, *Leta'litas*, (F.) *Mortalité*. Same etymon. This word, taken in an extended sense, expresses the condition of all organized bodies,—of being subject to

the cessation of life. In the sense in which it is most frequently employed, it signifies,—the proportional quantity of individuals who, in a certain population, die in a given time. If we assume the population of the earth to be one thousand millions, and a generation to last thirty-three years; in that space of time the one thousand millions must all die, and, consequently, the number of deaths will be, by approximation,

Each year.....	30,000,000
Each day	82,109
Each hour	3,421
Each minute	57
Each second.....	1 nearly.

If, on the other hand, as has been supposed, the number of deaths is to that of the births as TEN to TWELVE: there will be born,

Each year.....	36,000,000
Each day.....	98,356
Each hour	4,098
Each minute	68
Each second.....	1 & 2-15ths.

It has been estimated that the average mortality of the Pays du Vaud, is 1 in 49; of Sweden and Holland, 1 in 48; of Russia, 1 in 41; of France, 1 in 40; of Austria, 1 in 38; of Prussia and Naples, 1 in 33 to 35; of England, 1 in 45; and of South America, 1 in 30. The same rate of mortality has been given to the United States as to France; but the statistical details on all this matter have been inadequate, and—it is not improbable—inaccurate. The following has been given as the annual mortality of some of the chief cities of this country and Europe: (See the author's Human Health, p. 101: Philadelphia, 1844.) Philadelphia, 1 in 45-68; Glasgow, 1 in 44; Manchester, 1 in 44; Geneva, 1 in 43; Boston, 1 in 41-26; Baltimore, 1 in 41; London, 1 in 40; New York, 1 in 37-83; St. Petersburg, 1 in 37; Charleston, 1 in 36-50; Leghorn, 1 in 35; Berlin, 1 in 34; Paris, Lyons, Strasburg, and Barcelona, 1 in 32; Nice and Palermo, 1 in 31; Madrid, 1 in 29; Naples, 1 in 28; Brussels, 1 in 26; Rome, 1 in 25; Amsterdam, 1 in 24; and Vienna, 1 in 22½.

In the cities, the mortality under two years of age bears a large ratio. This is exhibited by the following table:

AGES.		London.	Philad.	Baltimore.
Under 1 year,	}	28-52	22-7	24-11
From 1 to 2,				
2 to 5,		9-97	7-3	11-18
5 to 10,		4-33	4-	5-
10 to 20,		4-03	5-	6-3
20 to 30,		6-64	12-	9-87
30 to 40,		8-08	12-	10-58
40 to 50,		8-89	10-	8-88
50 to 60,		8-89	7-2	5-78
60 to 70,		9-15	5-	4-5
70 to 80,		7-83	3-5	3-
80 to 90,		3-18	1-9	1-67
90 to 100,		0-40	0-5	0-26
100 to 120,		—	0-09	0-18
110 to 120,		—	0-013	

It would not be proper to regard the value of life in different countries, or in different periods in the same country, to be indicated by the average mortality; inasmuch as in one case a greater mortality may occur amongst children, and in another amongst adults.

MORTUARY, BILLS OF, *Bills of Necrology*, *Mortuary Registers*, (F.) *Tableaux de Mortalité*. Tables, instituted for the purpose of exhibiting the number of deaths, &c., in a given time. Well-kept bills of mortality are of great use to the physician and political economist. Those of London

were proverbial for their inaccuracy and insufficiency; especially as regards the complaints of which the persons have died. They were formerly made out by the parish clerks, and the information was conveyed to them by two old women, who were appointed in each parish, and were called *Searchers*;—their duty being to see that the deceased had died a natural death. This miserable system has, however, been abolished; and the Registrar-General's annual reports of births, deaths, and marriages, exhibit how admirably vital statistics are now collected in Great Britain.

MORTAR, *Morta'rium*, *Moreta'rium*, *Morto'rium*, *Piso*, *Ac'onē*, *Pila*, *Holmos*, *Igdē*, *Igdus*, (F.) *Mortier*. A vessel for reducing to powder different solid substances, and for making certain mixtures. Various substances are employed in the construction of mortars;—iron, marble, glass, wedgewood ware, &c.

MORTARIOLUM, Alveolus.

MORTELL, Mortal.

MORTIER, Mortar.

MORTIFER, Lethiferous.

MORTIFICATIO, Mortification—m. Pulmonum, Necropneumonia.

MORTIFICATION, *Mortifica'tio*, *Necro'sis*, *Sphacela'tion*, from *mors*, *mortis*, 'death,' and *fit*, 'I become.' The loss of vitality in a part of the body. The incipient stage of mortification, when the case is still recoverable, is called Gangrene; when totally destroyed, Sphacelus. Mortification of a bone is called Necrosis.

MORTIFICATION, COLD, Sphacelus—m. Ergot, Ergotism—m. Hot, Gangrene—m. Mildew, Ergotism.

MORTUOTH, Cupri sulphas.

MORTUARY REGISTERS, Mortality, bills of.

MORTUUS, Dead.

MORUM, see *Morus nigra*—m. *Palpebræ interne*, *Trachoma carunculolum*.

MORUS NIGRA, *M. lacinia'ta*, *Mul'berry Tree*, *More'a*, *Sycami'nos*, (F.) *Murier Noir*. Ord. Urticaceæ. The fruit of the mulberry tree, *Morum*, *μῑρον*, *Sycami'num*, (F.) *Mûre*, is pleasant, sweet, subacid, and abounds with a deep violet-coloured juice, *Mori succus* (Ph. L.). A syrup is directed in the London Pharmacopœia. The bark of the root has been regarded as an anthelmintic.

MORUS RUBRA, *Red Mulberry*, *Wild Mulberry*; indigenous. The root has been used to destroy tænia.

MORVE, Mucus, Nasal mucus; see *Equinia*—m. *Farcineuse*, see *Equinia*.

MOS, Habit.

MOS MORBI BENIG'NUS seu MALIG'NUS. The benign, or malignant character of a disease.

MOSCHARDINA, *Muscer'da*, from *μῑχος*, 'musk.' A lozenge for the mouth, composed of musk, ambergris, and other aromatics.

MOSCHELÆ'ON, *Moscolæ'a*. A compound aromatic oil, containing musk.

MOSCHUS, Musk—m. Arabum, *Hibiscus abelmoschus*—m. *Facitiosus*, Musk, artificial.

MOSCOLÆA, *Moschelæon*.

MOSLICH, Maslach.

MOSQUITA, see Mosquito.

MOSQUITO (S.), Diminutive of (S.) *mosca*, 'a fly.' *Musquito*. A very troublesome insect in warm, moist situations. Curtains, called *Mosquito curtains* or *bars*, are used by way of protection. The entrance of mosquitoes into the bedchamber may also be prevented, to a certain extent, by keeping the doors and windows closed until the lights are extinguished. The pain and itching from the bites are alleviated by washing them with hartshorn.

MOSQUITA is a name given, by the Portuguese, to small, red pimples on the skin, resembling the bites of the mosquito.

MOSS BERRY, *Vaccinium oxycoccus*—*m.* Ceylon, *Fucus amylaceus*—*m.* Club, *Lycopodium*—*m.* Cup, *Lichen pyxidatus*—*m.* Edible, *Fucus amylaceus*—*m.* Haircap, *Polytrichum Juniperinum*—*m.* Iceland, *Lichen Islandicus*—*m.* Irish, *Fucus crispus*—*m.* Jaffna, *Fucus amylaceus*—*m.* Marine, *Fucus amylaceus*—*m.* Pearl, *Chondrus crispus*—*m.* Wall, common yellow, *Parmelia parietina*.

MOTA, *Fagus castanea*.

MOTAMEN, Motion.

MOTATIO, Motion.

MOTE, *Carbasus*, Linteum.

MOTEUR, Motory.

MOTHER, *Hysteria*, Phlegm.

MOTHER'S MARKS, *Nævus*—*m.* Spots, *Nævus*.

MOTHERWORT, *Leonurus cardiaca*, *Matricaria*.

MOTILITY, *Motilitas*, from *motus*, 'movement.' (F.) *Motilité*, *Motricité*. Faculty of moving; moving power; contractility.

MOTION, *Motus*, *Mo'tio*, *Mota'tio*, *Mota'men*, *Commo'tio*, *Permo'tio*, (*move*, *motum*, 'to move.') *Cine'sis*, *Movement*, (F.) *Mouvement*. The act of changing place. The various motions may be divided into,—*First*, the *voluntary* or those that are executed under the influence of the brain. *Secondly*, the *involuntary*, which may be subdivided into, 1. The *excited*, of the *reflex function* of Dr. Marshall Hall and others,—as the closure of the larynx on the contact of acrid vapours, of the pharynx on that of the food,—a function of the spinal marrow; and, 2. Those that are executed under the organic and other nerves of involuntary function. It is probable, too, that every living tissue is capable of moving responsive to its appropriate irritant. See Irritability.

MOTION, CILIARY, see *Cilia*—*m.* Involuntary, see *Automatic*—*m.* Vibratory, see *Ciliary*—*m.* Voluntary, *Autocinesis*.

MOTIVE, Motory.

MOTOR, Motory.

MOTOR OCULI EXTERNUS, *Nervus tim'idus*, (F.) *Nerf oculo-musculaire externe* (Ch.), *Nerf moteur oculaire externe*, Sixth pair of nerves, arises from the furrow which separates the pons Varolii from the medulla oblongata. It enters the cavernous sinus; anastomoses on the outside of the carotid artery with two filaments furnished by the superior cervical ganglion; enters the orbit by the sphenoidal fissure, and is lost on the rectus externus oculi. This pair is, also, called *Abducentes*.

MOTOR ES OCULORUM, *Moto'rii Oculorum*, *Oculares communes*, *Ophthalmici externi*, *Common Oculo-muscular Nerves*, (F.) *Nerfs moteurs oculaires communs* (Ch.), The third pair of nerves, *Par tertium*. This nerve arises from a depression at the inner side of the peduncles of the brain, between the tuber annulare and corpora mamillaria. It enters the outer part of the cavernous sinus, and passes into the orbital cavity. Behind the sphenoidal fissure, it divides into two branches; the one *superior* and the other *inferior*. The former is lost on the rectus superior oculi and the levator palpebræ superioris; the latter—the larger of the two—divides into three branches, which pass to the rectus internus, rectus inferior, and lesser oblique. The last furnishes a filament, which passes to the ophthalmic ganglion.

MOTORIAL, Motory.

MOTORPATHY, Kinesipathy.

MOTORY, *Motive*, *Motor*, *Moto'rial*, *Motrix*, *Kinet'ic* or *Cinet'ic*, (F.) *Moteur*. Same etymon.

That which moves or causes movement, as *motor nerves*, in contradistinction to *sensory*.

MOTOS, Linteum, Tent.

MOTO'SIS, *mosis*, 'lint.' The application of lint to a wound.

MOTRICITÉ, Motility.

MOTRIX, Motory.

MOTTE, *Mons veneris*.

MOTUM, *Carbasus*, Linteum.

MOTUS, Motion—*m.* Assimilationis, Plastic force—*m.* Automaticus, see *Automatic*—*m.* Compressorius, *Peristole*—*m.* Cordis diastalticus, *Diastole*—*m.* Hamorrhoidalis, *Hamorrhoids*—*m.* Intestinorum, *Borborygmus*—*m.* Involuntarius, see *Autocinesis*—*m.* Peristalticus, *Peristole*—*m.* Testudineus, *Peristole*—*m.* Vermicularis, *Peristole*—*m.* Voluntarius, *Autocinesis*.

MOUCHE, *Cantharis*—*m.* d'Espagne, *Cantharis*.

MOUCHER L'ACTION DE, Munctio:

MOUCHES, see *Pains*, labour—*m.* *Volantes*, see *Metamorphopsia*.

MOUCHETURE, see *Scarification*.

MOUCHOIR EN TRIANGLE, *Couvrechef*, see *Bandeau*.

MOUFETTE, see *Mephitism*.

MOULD, Fontanella.

MOULE, *Mytilus edulis*.

MOULEUR, Muller.

MOULTING, Desquamation.

MOUNTAIN ASH, *Sorbus acuparia*—*m.* Ash, American, *Sorbus Americana*—*m.* Sickness, *Puna*.

MOUREILLER, *Malpighia mourella*.

MOURON DES OISEAUX, *Alsine media*—*m.* *Rouge*, *Anagallis*.

MOUSE-EAR, *Alsine media*, *Hieracium pilosella*.

MOUSE-SIGHT, *Myopia*.

MOUSSE DE CORSE, *Corallina Corsicana*—*m.* d'Islande, *Lichen Islandicus*—*m.* d'Irlande, *Fucus crispus*—*m.* *Marine perlée*, *Lichen Islandicus*—*m.* *Perlée*, *Fucus crispus*.

MOUSSERON, see *Agaric*.

MOUSSEUX, Frothy.

MOUSTACHE, Mystax.

MOÛT DE LA BIÈRE, Wort.

MOÛTARDE, *Sinapis*—*m.* des Allemands, *Cochlearia armoracia*—*m.* des Moines, *Cochlearia armoracia*.

MOUTH, Sax. *muð*, *Os*, *Cavum Oris*, *Stoma*, *Gab*, (Sc.) *Mou*, (Prov.) *Gob*, (F.) *Bouche*. This word sometimes signifies the cavity situate between the jaws, and containing the tongue, &c.;—at others, the outer orifice of that cavity. The mouth, in the first acceptation, *Cavitas seu Spatium Oris*, is the cavity; bounded, above, by the palatine arch; below, by the tongue; before, by the lips; and behind, by the velum palati and pharynx. The sides of the mouth and the organs it contains are lined by a mucous membrane. The anterior aperture of the mouth is, sometimes, called *facial*—the posterior, *pharyngeal*. In the mouth are the teeth, gums, alveolar margins, tongue; the excretory ducts of the salivary glands, and those of a number of mucous follicles, &c. It is in this cavity that the food is cut, torn, or bruised by the teeth; is impregnated with saliva, and formed into a mass or *bolus*, which is then subjected to the act of deglutition. The mouth contains the organs of taste; and it serves in respiration, articulation, expectoration, suction, &c.

The condition of the mouth requires to be attended to in *Semeiology*. The state of the mucous membrane, like that of the tongue, indicates the condition of the membrane of the alimentary canal generally.

MOUTH, Os, Apertu'ra, Orific'ium, Hia'tus, Peristom'ium, is, also, applied to the open extremities of vessels or other canals.

MOUTH ROOT, Coptis.

MOUTH, SORE, Aphthæ.

MOUTH WATERING, see Salivation.

MOUVEMENT, Motion—*m. Péristaltique*, see Peristole.

MOVEMENT, Motion—*m. Cure, Kinesipathy. MOW*, Mouth.

MOXA, Moxib'rium. A word by which the Chinese and Japanese designate a cottony substance, which they prepare by beating the dried leaves of the *Artemisia moxa*,—a kind of mugwort. With this down they form a cone, which is placed upon the part intended to be cauterized, and is set fire to at the top. The heat and pain gradually increase, in proportion as the combustion proceeds downwards, until ultimately an eschar may be formed. In Europe and this country, the same operation is usually practised with a cylinder of cotton-wool, or with one formed from the pith of the greater sun-flower. This mode of cauterization is employed as a powerful counter-irritant; which it assuredly is. Sloughing may be in some measure prevented, according to Baron Larrey, by the application of liquid ammonia to the burnt part.

The term moxa has been extended to any substance, which by gradual combustion on or near the skin is employed as a counter-irritant,—as the *electric or galvanic moxa* or *cautery*, in which voltaic electricity is employed as the cauterizing agent.

MOXIBURIUM, Moxa.

MOXIBUSTION, *Moxibustio, Moxocausis, Encausis*. Mode of cauterization by means of moxa.

MOXOCAUSIS, Moxibustion.

MOXON'S EFFERVESCING MAGNESIA, see Magnesia, effervescing, Moxon's.

MUCAGO, Liquor sanguinis, Mucilage.

MUCARUM, Mucharum.

MUCCINIUM, Mucilage.

MUCEDO, Mucilage.

MUCHA'RUM, Muca'rum. A barbarous term, formerly used for an aqueous infusion of roses, sweetened, and evaporated to the consistence of syrup.

MUCIFIC, Blennogenous.

MUCIFORM, *Muciformis, Myxo'des, Blenno'des, Blenno'des, Blenno'des*, from *mucus*, and *forma*, 'form.' Resembling mucus in character or appearance.

MUCILAGE, Mucila'go, Muca'go, Muce'do, Muccin'um, from the Lat. *mucus*,—itself presumed to be from *mungere, munctum*, 'to wipe the nose.' A mixture of gum and a small quantity of matter analogous to mucus, which is found in abundance in linseed, quince-seed, &c. It is obtained by heating in water the parts, or products, of plants which contain it. It is much used in the preparation of emollient cataplasms and the greater part of the demulcent *tisanes*, (F.) *Hydrolés, Hydrolites*.

MUCILAGE, see Liquor sanguinis—*m. Adragant, Mucilago astragali tragacanthæ*—*m. de gomme Adragante, Mucilago astragali tragacanthæ*—*m. de gomme Arabique, Mucilago acaciæ*.

MUCILAGE, ANIMAL, Mucus.

MUCILAGE ARABIQUE, Mucilago acaciæ—*m. of Barley, Mucilago hordei*—*m. of Quince-seeds, Decoctum cydoniæ*—*m. of Rice, see Oryza*—*m. of Sassafras Pith, Infusum sassafras medullæ*.

MUCILAGE, STARCH, Mucilago amyli.

MUCILAGINEUX, Mucilaginous.

MUCILAG'INOUS, *Mucilaginosus, Gummo'sus, Gummio'des, Gummido'des*, (F.) *Mucilagineux*. Gummy. Resembling gum. That which possesses the character of mucilage.

MUCILAGO, Mucilage, Mucilago Acaciæ, Synovia.

MUCILAGO ACA'CLE, M. Acaciæ Arabicæ see *Gummi Arabicæ, Mucila'go, Mistu'ra Aca'ciæ, Mucilage of acaciæ or of gum Arabic*, (F.) *Mucilage Arabique ou de gomme Arabique*. (Acaciæ gum. contus. \mathfrak{z} iv; aq. bullient. Oss. Rub the gum with the water gradually added.) Demulcent. To allay cough; but, chiefly, to suspend insoluble matters in water, &c.

MUCILAGO AM'YLI, Starch mucilage, Decoctum Amyli. (Amyli, \mathfrak{z} ij; aq. \mathfrak{o} j. Rub the starch with the water gradually added; then boil till it forms a mucilage.—Ph. L.) Demulcent. Generally given *per anum*.

MUCILAGO ARTICULORUM seu JUNCTURARUM, Synovia.

MUCILAGO ASTRAG'ALI TRAGACAN'THÆ, M. tragacanthæ (Ph. U. S.) seu *gummi tragacanthæ, Mucilage of tragacanth, Mucilage Adragant ou de gomme Adragante*. (Gummi tragacanth. \mathfrak{z} ij; aq. bullient. \mathfrak{o} j. Macerate for 24 hours; triturate till the gum is dissolved, and press through linen.) Used for pharmaceutical purposes.

MUCILAGO CRETICA, Mistura cretæ—*m. Gummi Arabici, M. acaciæ*.

MUCILA'GO HOR'DEI, Mu'cilage of Barley. (Hordei, \mathfrak{z} ss, (avoirdupois) Aqua, f \mathfrak{z} xvi. Triturate well; and boil. Ph. D.) A vehicle for substances given in enemata.

MUCILAGO SEMINIS CYDONII MALI, Decoctum cydoniæ—*m. Gummi tragacanthæ, M. astragali tragacanthæ*—*m. Tragacanthæ, M. astragali tragacanthæ*.

MUCIN, Mucus.

MUCIP'AROUS, Mucip'arus, (F.) *Mucipare*, from *mucus*, and *pario*, 'I bring forth.' Mucus-producing; as *muciparous glands* or *follicles*.

MUCYTIS, Mycodermitis, from *mucus*, and *itis*, denoting inflammation. A term, badly formed, for inflammation of a mucous membrane.

MUCKSWEAT, Desudatio.

MUCOCE'LE, Hernia sacci lacrymalis, from *mucus*, and *κλῆν*, 'rupture.' An enlargement or protrusion of the mucous membrane of the lacrymal passages, giving occasion to fistula lacrymalis. Also, dropsy of the lacrymal sac.

MUCO-ENTERITIS, see Enteritis.

MUCO-PURULENT, Mucopurulentus, Pyoblen'neus, from *mucus*, and *pus*. Having the character or appearance of mucus and pus.

MUCOR, Euros, Mucus—*m. Narium, Nasal mucus*.

MUCOSIN, see Mucus.

MUCOSITAS, Mucosity, Mucus.

MUCOSITY, Mucos'itas, (F.) *Mucosité*. A fluid, which resembles mucus, or contains a certain quantity of it.

MUCOUS, Mucos'us, (F.) *Muqueux*. An epithet for all bodies containing mucilage or mucus. It is, also, sometimes used synonymously with gummy. In pathology, it is occasionally employed to express the seat of a disease, as *mucous disease, mucous phlegmasia*; that is, having its seat in a mucous membrane.

MUCOUS CORPUSCLES, see Mucus—*m. Layer*, see *Tache embryonnaire*—*m. Membranes*, see *Membranes*—*m. Web, Corpus mucosum*.

MUCRO, Apex.

MICRO seu *Cor'yp'hê* seu *Fundus* seu *Vertex Cordis*. The apex or point of the heart.

MUCRO HUMERI, Acromion.

MUCRONATA CARTILAGO, Xiphoid cartilage.

MUCRONATUM OS, Xiphoid cartilage.

MUCULENT, *Muculen'tus*, from *mucus*. Slimy, viscid.

MUCUNA, see *Dolichos pruriens*—*m. Pruriens*, *Dolichos pruriens*—*m. Prurita*, see *Dolichos pruriens*.

MUCUS, *Mucor*, *Muco'sus humor*, *Mucos'itas*, *Myra*, *Mycus*, *Lapê*, *Pituita*, *Blenna*, *Zoömy'cus*, *Zoömyx'a*, *Pit'uite*, *Animal mucus*, *Animal mucilage*, (*F.*) *Muqueux animal*, *Morve*, *Mucilage animal*, presumed to be from *mungere*, 'to wipe the nose.' A substance, analogous to vegetable mucilage; from which, however, it differs, by affording subcarbonate of ammonia on distillation. Mucus exudes through the skin, in a state of combination with a peculiar oily matter; and, drying, forms the epidermis. It constitutes, in part, the different epidermeous productions, as the hair, nails, wool, and horn of animals, feathers of birds, and scales of fish. It is found at the surface of the mucous membranes, and presents some difference in its composition and properties, according to the particular mucous membrane from which it is obtained. Its chief organic constituent is an albuminoid compound,—*mu'cin*, *mu'cosin*. Mucus preserves the membranes moist, and in a state best fitted for the performance of their functions. The French give the term *glair* to the thick, stringy mucus, secreted by the mucous membranes when in a state of disease.

When mucus is examined with the microscope it is found to contain numerous epithelial scales or flattened cells; together with round, granular bodies, commonly termed *mucous corpuscles* or *mucous globules*, which are contained in a fluid portion—*liquor muci*.

MUCUS CARPHODES, see *Carphodes*—*m. Catharticus*, *Lichen Islandicus*—*m. Malpighii*, *Corpus mucosum*—*m. Narium*, *Nasal mucus*.

MUD-APPLE PLANT, *Solanum melongena*.

MUDAR, *Madar*. The *Calotropis mudarii* seu *madarii*, *C. gigante'a*, *Ascle'pias gigante'a*, *Mad'o'rius*. *Ord.* *Asclepiadaceæ*. In the Hindo practice of physic, the bark of the root, as well as the concrete juice of the plant, enters into various compound formulæ for the cure of elephantiasis and many other disorders. Experiments instituted by Dr. Duncan, Jr., exhibit its properties to be like those of ipecacuanha. Mixed with oil it is used in gout in Malabar.

MUDGE'S INHALER, see *Inhaler*.

MUDLICK, see *Kentucky*, *Mineral waters of*.

MUGUET, *Aphtha*, *Convallaria maialis*—*m. des Bois*, *Asperula odorata*—*m. de Mai*, *Convallaria maialis*.

MUGWORT, *Artemisia vulgaris*.

MULATRE, *Mulatto*.

MULATTO, (*F.*) *Mulâtre*, (*S.*) *Mulata*. An individual of the human species engendered of a white and black race; from *mulus*, 'a mule.'

The following table exhibits the proportion of white blood in the various castes,—arising from the hybridous admixture of white and black,—according to the principles sanctioned by usage.

PARENTS.	OFFSPRING.	DEGREE OF MIXTURE.
Negro and White,.....	Mulatto,.....	1-2 White, 1-2 Black.
White and Mulatto,.....	Terceron,.....	3-4 — 1-4 —
Negro and Mulatto,.....	{ Griffo, Griff. or Zambo, } or Black Terceron, }	1-4 — 3-4 —
White and Terceron,.....	Quarteron or Quadroon,.....	7-8 — 1-8 —
Negro and Terceron,.....	Black Quarteron or Quadroon....	1-8 — 7-8 —
White and Quarteron,.....	Quinteron,.....	15-16 — 1-16 —
Negro and Black Quarteron,.....	Black Quinteron.....	1-16 — 15-16 —

The last two were considered to be respectively white and black, in the British West India Islands; and the former, prior to modern changes, were white by law, and consequently free. See *Half-caste*.

MULATTO JACK, Fever, yellow.

MULBERRY CALCULUS, see *Calculi, urinary*.

MULBERRY MASS. The mulberry-like appearance, presented by the embryo, at a very early stage, when the different cells have drawn around them a portion of the vitellus or yolk.

MULBERRY RASH, see *Typhus*—*m. Red*, *Morus rubra*—*m. Tree*, *Morus nigra*—*m. Wild*, *Morus rubra*.

MULES (*F.*), *Mulæ*. Chilblains on the heels.

MULET, Hybrid.

MULGEDIUM ACUMINATUM, (*mulgere*, *mulsum*, 'to milk,') *Lactu'ca villo'sa*, *Blue Lettuce*; and

MULGEDIUM FLORIDANUM, *Lactu'ca Florida'na*, *Sonchus Florida'nus*, *False Lettuce*; indigenous plants; *Order*, *Compositæ*; have the reputation of curing the bites of rattlesnakes.

MULIEBRIA, *Menses*, *Vulva*.

MULIEBRITY, *Mulieritas*.

MULIEBROSITAS, *Phylogyn'ia*, from *mulier*, 'a woman,' *muliebris*, 'appertaining to women.' Fondness for women. Hence, *Muliebro'sus*, *Muliera'rius*, and *Muliero'sus*; one who is fond of women.

MULIEBROSUS, see *Muliebro'sitas*.

MULIERARIUS, see *Muliebro'sitas*.

MULIERITAS, *Mulieb'rity*, *Womanhood*, from *mulier*, 'a woman.' The state of puberty in the female.

MULIEROSUS, see *Muliebro'sitas*.

MULIER PLENA, *Pregnant*.

MULLED WINE, see *Wine*.

MULLEIN, *BLACK*, *Verbascum nigrum*—*m. Broad-leaved*, *great*, *Verbascum nigrum*—*m. Yellow*, *Verbascum nigrum*.

MULLER, (*F.*) *Mouleur*. A moulder. A stone held in the hand, with which any powder is ground upon a horizontal slab. It is sometimes called *mullet*, (*F.*) *Mollette*.

MULLIGRUBS, *Tormina*, *Colic*.

MULLIN, *Melizomum*.

MULOMEDICINA, *Veterinary art*.

MULOMEDICUS, see *Hippiater*.

MULSA ACIDA, *Oxyglycus*.

MULSUM, *Hydromeli*—*m. Vinosum*, *CEnomal*.

MULTIFIDUS SPINÆ, *Transversalis dorsi*.

MULTIFŒTATION, *Multifœta'tio*, from *multus*, 'many,' and *fœtus*. Pregnancy with more than two fœtuses.

MULTILOCULAR, *Multilocula'ris*, from *multus*, 'many,' and *loculus*, 'a cell.' Having many cells or small cavities.

MULTIAMMÆ, (F.) *Multimammæ*, from *multus*, 'many,' and *mamma*, 'a breast.' A variety of hypergenæsis, in which there are supernumerary mammae.

MULTIP'AROUS, *Multip'arus*, (F.) *Multipare*, from *multus*, 'many,' and *parire*, 'to bring forth.' One that brings forth several young at the same time. Such a birth is called *plural*.

MULTIPLICATION MÉRISMATIQUE, see Generation.

MULTITUDO, *Plethora*.

MULTIVORANTIA, *Polyphagia*.

MULTIVOROUS, *Polyphagous*.

MUMIA, *Mummy*—*m. de Medullis*, Marrow.

MUMISATIO, *Mummification*.

MUMMIFICATION, *Mumisa'tio*, from *mumy*, and *fio*, 'to be made.' *Sceletu'sis*. The mode of preparing a mummy.

MUMMY, *Mumia*, Arab. *Moumya*, from *mum*, 'wax.' *Rebolea*, *Rebona*, (F.) *Momie*. A dead body simply dried, or dried after having been embalmed. The latter acceptation is the most common. Formerly, the Egyptian mummy was extolled as useful in contusions. It was presumed, also, to have healing, tonic, and resolvent properties. It is now only regarded as an archæal curiosity.

MUMPS, *Cynanche parotidea*.

MUNCHATOË, *Mystax*.

MUNC'TIO, *Apomex'is*, (F.) *l'Action de moucher*, from *mungere*, *muncium*, 'to blow the nose.' The act of blowing the nose.

MUNDIFICANTIA, *Detergents*.

MUNDIFICATIVA, *Detergents*.

MUND'TIA SPINO'SA. A plant of the *Nat. Ord. Polygalæ*, which grows in Southern Africa, and a decoction of whose branches is used in atrophy, phthisis, &c.

MUNGOS RADIX, *Ophiorrhiza mungos*.

MUNGREL, *Hybrid*.

MUNIMENTUM CASTITATIS, *Hymen*.

MUQUEUX ANIMAL, *Mucus*.

MUR, see *Coryza*.

MURAL, *Mura'lis*, from *murus*, 'a wall.' Vesical calculi are so called, when rugous and covered with tubercles or asperities. They are composed of oxalate of lime.

MURE, *Moro*, see *Moros nigra*.

MURFLES, *Ephelides*, *Papulae*.

MUR'IA, *Halmē*, *Salsila'go*, *Brine*, (F.) *Saumure*, ('*ἀλμυρ*,' 'salt water.') This was formerly used as a glyster in dysentery, in certain inveterate neuralgia, &c.

MUR'IA, *Soda*, *muriate of*.

MURIAS AURICO-NATRICUM, see *Gold*.

MURIATE D'AMMONIAQUE, *Ammoniac murias*—*m. d'Ammoniaque et de fer*, Ferrum ammoniatum—*m. d'Or*, *Gold*, *muriate of*, see *Gold*.

MURIAT'IC AC'ID, from *muria*; *Ac'idum salis seu mari'nium concentra'tum seu salis culina'ris seu salis mari'ni*, *Spir'itus salis mari'ni seu salis Glaube'ri seu salis ac'idus seu humans*, *Acidum hydrochlor'icum seu muriat'icum*, *Spirit of salt*, (F.) *Acide hydrochlorique ou muriatique*. An aqueous solution of chlorohydric acid gas of s. g. 1.16. The odour of muriatic acid is suffocating; taste very acid and caustic. It is nearly

colourless when pure; but commonly of a pale yellow; volatile; the fumes visible. Muriatic acid is possessed of tonic and antiseptic properties. It is used in typhus; cutaneous eruptions; in gargles for inflammatory and putrid sore throats, (gtt. xxx to fʒvj of water,) &c. When added to a pediluvium, it renders it stimulating.

ACIDUM MURIAT'ICUM DILU'TUM of the Ph. U. S. contains four fluidounces of muriatic acid, to twelve fluidounces of distilled water.

MURIDE, *Bromine*.

MURIER NOIR, *Morus nigra*.

MURIGENE, *Chlorine*.

MURINA, *Bromine*.

MURMUR, (F.) *Murmure*. A word existing both in the Romanic and Teutonic languages, and probably a variety of onomatopœia;—*mur mur* expressing the kind of sound which the word indicates. A low continued or continuously repeated sound, as that of flame, or of a stream running over a stony bottom.

MURMUR AURIUM, *Tinnitus Aurium*—*m. Intestinales seu intestinorum seu Ventris*, *Borborygmus*.

MURMUR, RES'PIRATORY. (F.) *Bruit respiratoire*. The noise heard during inspiration and expiration, especially the former. It is produced by the passage of the air through the bronchial tubes and into the air-cells. It has been also called *murmur of the expansion of the lungs*, (F.) *Bruit d'expansion pulmonaire*; and, when distinctly vesicular, *Respiration of the cells or vesicular respiration*, (F.) *Respiration ou murmure vésiculaire*. Vesicular respiration is of course absent when the cells of the lungs have been obliterated from any cause. We may then have the *Respiration nulle*, *Absence du bruit respiratoire*, *Silence*, and *Respiration silencieuse* of the French writers.

At times, it is rude during inspiration or expiration, or both—the *Respiration rude* or *R. râpeuse* of the French. At others, there is a blowing sound, (F.) *Souffle*, *Respiration soufflante*, as if some one were blowing into the auscultator's ear through a tube. This is heard in the healthy state over the larynx, trachea, (F.) *Souffle trachéal*, and about the bifurcation of the bronchia, (F.) *Souffle bronchique*; but when it proceeds from the lungs it denotes disease. It may be *tubular* or *diffused*. In the former, the *whiffing* murmurs appear to occur in a space limited to the immediate neighbourhood of the part examined. In the latter, they are produced with but moderate intensity, and sometimes at a distance from the ear, over a tolerably extended space.

The respiration, perceived over the trachea and bronchia in health, is called *tracheal* or *bronchial* or *tubal*, (F.) *Respiration bronchique*, *Souffle tubaire*, according to the situation in which it is heard.

MURMUR, UTERINE, *Bruit placentaire*—*m. Utero-placental*, *Bruit placentaire*.

MURMUR, WHIFFING, see *Murmur, respiratory*.

MURR, *Murray*, from A. S. *mýrrþan*, 'to destroy,' or from (L.) *mori*, 'to die.' (F.) *Claveau*, *Clavelle*, *Pestilential Fever*. An epizootic, perhaps contagious, disease, having some resemblance to small-pox, which affects cattle—especially sheep; and is said to have been transferred to man.

MURRAIN, see *Coryza*.

MURREN, *Coryza*.

MUSA, *M. Paradisiaca*—*m. Cliffortiana*, *M. Paradisiaca*—*m. Mensaria*, *M. Paradisiaca*.

MUSA PARADIST'ACA, *Musa. M. Mensa'ria seu Cliffortia'na*, *Annona*, *Palma hu'milis*, *Ficus In-*

dica, *Bata*, *Plantains*, *Plantain tree*. *Family*, *Musaceæ*. *See*, *Nyct.* *Hexandria Monogynia*. It grows in many parts of India and South America. The fruit, which is largely eaten for bread, consists of a mealy substance, *Plantain meal*, *Conquin-tay*. It is clammy: has a sweetish taste, and will dissolve in the mouth without chewing. The whole spike of fruit often weighs forty or fifty pounds. The leaves of the tree serve the Indians for tablecloths and napkins. Being smooth and soft, they are also employed as dressings for blisters. The water from the soft trunk is astringent, and sometimes used in diarrhoea.

MUSA SAPIENTUM. The *Banana tree*, *Baco'ba*, (F.) *Bananiar*, *Bacove*. This differs somewhat from the last. The fruit, *Banana*, (S.) *Platano*, is shorter, straighter, and rounder: the pulp softer and of a more luscious taste. When ripe, it is very agreeable; it is eaten like the plantain, and relished by all ranks of people in the West Indies. Both varieties are natives of Guinea.

MUSC, Musk.

MUSCA HISPANICA, *Cantharis*.

MUSCÆ ENTO-HYALOID, *see* *Metamorphopsia*—*m*. *Volitantes*, *see* *Metamorphopsia*.

MUSCADE, *see* *Myristica moschata*.

MUSCADIER, *Myristica moschata*.

MUSCERDA, *Moscharina*.

MUSCLE, (old Eng.) *Brawn*, *Musculus*, *Torus*, *Lozer'us*, *Mys*, *Myon*, from *mys*, 'a mouse or rat:' because, say some etymologists, the ancients compared the muscles to flayed mice or rats. According to Diemerbroeck, Douglass, Chaussier, &c., *mys* comes rather from *muo*, 'to close,' 'to move,' &c., a function proper to muscles. This etymon is the more probable. Muscles have been divided into those of *Animal life* or of the *life of relation*—*voluntary muscles*—which execute movements under the influence of the will; as the muscles

of the limbs, head, trunk, &c., and into those of *organic life*—*involuntary muscles*—which contract under the influence of certain special stimuli; as the heart, fleshy fibres of the stomach, &c. *Mixed muscles* are those which belong partly to each of these divisions;—as the muscles of respiration; the sphincters, &c. Muscles that act in opposition to each other are called *antagonists*; thus, every extensor has a flexor for an antagonist, and conversely. Muscles that concur in the same action are termed *congenerous*. The muscles present numerous varieties in form, size, situation, use, &c., and have been divided, by some, into *long*, *broad*, and *short*. Each of these divisions comprises *simple* and *compound* muscles. *Simple* or *rectilinear muscles* have all their fibres in a similar direction, and only one body—as the *Sartorius*, *Pronator quadratus*, &c. *Compound muscles* are those which have only one belly and several tendons, as the flexors of the fingers and toes; or several bellies and several tendons,—as the biceps flexor cubiti, sacro-lumbalis, &c. To the compound muscles belong, also, the *radiated muscles*. Their fibres set out from a common centre, and are arranged like the radii of a circle;—such are the diaphragm, iliacus, temporal, &c. *Pennate* or *Penniform Muscles*. Their fibres are arranged in two rows, which are united at a median line, at greater or less angles; nearly as the feathers are inserted into a quill. The *palmaris longus* is one of these. *Semi-penniform muscles*: their fibres are oblique, as in the last case; but they are inserted only on one side of the tendon. *Hollow Muscles* are,—the heart, intestines, urinary bladder, &c.

Much difference has existed in the enumeration of muscles. Some authors reckon them upwards of 400. Chaussier admits only 100. The greater part of them are in *pairs*. Very few are *azygous*.

TABLE OF THE MUSCLES, ARRANGED AFTER THE MANNER OF DR. BARCLAY, ACCORDING TO THEIR ACTIONS.

THE HEAD IS MOVED

Forwards by	Backwards by	To either side by
Platysma myoides, Sterno-mastoideus, Rectus anticus major, " " minor,	Part of trapezius, Splenius capitis, Complexus, Trachelo-mastoideus, Rectus posticus major, " " minor, Obliquus capitis superior.	Platysma myoides, Sterno-mastoideus, Part of trapezius, Splenius capitis, " colli, Trachelo-mastoideus, Complexus.
Assisted (when the lower jaw is fixed, by Mylo-hyoideus, Genio-hyoideus, Genio-hyo-glossus, Digastrici.		

THE NECK IS MOVED

Forwards by	Backwards by	Laterally by
Platysma myoides, Sterno-mastoideus, Digastricus, Mylo-hyoideus, Genio-hyoideus, Genio-hyo-glossus Omohyoidei, Sterno-hyoidei, Thyro-hyoidei, Rectus anticus minor, Longus colli.	Part of trapezius, Rhomboides minor, Serratus posticus superior, Splenius capitis, " colli, Complexus, Trachelo-mastoideus, Transversalis colli, Inter-spinales colli, Semi-spinales colli, Rectus posticus major, " " minor, Obliquus capitis superior, " inferior, Scaleni postici, Levator scapulae.	Various combinations of those muscles which separately move it: forwards and backwards, assisted by the scaleni, intertransversales, and recti laterales.

THE TRUNK IS MOVED

Forwards by

Rectus abdominis,
Pyramidalis.
Obliquus externus abdominis,
Obliquus internus,
Psoas magnus,
" parvus.

Assisted (when the arms are carried forwards) by

Pectoralis major,
" minor,
Serratus magnus.

Backwards by

Trapezius,
Rhomboides major,
Latissimus dorsi,
Serratus posticus superior,
" inferior,
Sacro-lumbalis,
Longissimus dorsi,
Spinales dorsi,
Semi-spinales dorsi,
Multifidus spine.
Inter-transversales dorsi et lumborum.

Laterally by

Obliquus externus,
internus,
Quadratus lumborum,
Longissimus dorsi,
Sacro lumbalis,
Serrati postici,
Latissimus dorsi.

THE SCAPULA IS MOVED

Upwards by

Trapezius,
Levator scapulae,
Rhomboides.

Downwards by

Lower part of trapezius,
Teres major,
Pectoralis minor.

Forwards by

Pectoralis minor,
Serratus magnus.

Backwards by

Part of trapezius,
Rhomboides,
Latissimus dorsi.

THE HUMERUS IS MOVED

Forwards by

Part of deltoid,
Part of pectoralis major,

Assisted in some circumstances by

Biceps,
Coraco-brachialis.

Backwards by

Part of deltoid,
Teres major,
" minor,
Long head of triceps,
Latissimus dorsi.

Inwards by

Part of pectoralis major,
Latissimus dorsi.

Rotated inwards by

Subscapularis,

Assisted occasionally by
Pectoralis major,
Latissimus and teres major.

Outwards by

Supra-spinatus,
Infra-spinatus,
Teres minor.

THE FORE-ARM IS MOVED

Forwards by

Biceps,
Brachialis anticus,
Pronator teres,

Assisted by

Flexor carpi radialis,
" sublimis,
" ulnaris,
Supinator longus.

Backwards by

Triceps,
Anconeus.

Rotated inwards by

Pronator teres,
Flexor carpi radialis,
Palmaris longus,
Flexor sublimis,
Pronator quadratus,

Outwards by

Biceps,
Supinator brevis,
Extensor secundi internodii.

THE CARPUS IS MOVED

Forwards by

Flexor carpi radialis,
Palmaris longus,
Flexor sublimis,
" carpi ulnaris,
" profundus,
" longus pollicis.

Backwards by

Extensor carpi radialis longior,
Extensor carpi radialis brevior,
Extensor secundi internodii,
Indicator,
Extensor communis digitorum,
Extensor proprius pollicis.

Outwards by

Flexor carpi radialis,
Extensor carpi radialis longior,
Extensor carpi radialis brevior,
Extensor ossis metacarpi,
Extensor primi internodii.

Inwards by

Flexor sublimis,
" carpi ulnaris,
" profundus,
Extensor communis digitorum,
Extensor minimi digiti,
Extensor carpi ulnaris.

THE THUMB IS MOVED

Inwards and forwards across the palm, by

Opponens pollicis,
Flexor brevis,
" longus

Outwards and backwards by

Extensor ossis metacarpi pollicis,
Extensor primi internodii,
Extensor secundi internodii.

Upwards and forwards, away from the other fingers, by

Abductor,
Assisted by part of the
Flexor brevis.

Backwards and inwards, to the other fingers, by

Adductor,
Extensor primi internodii,
Extensor secundi internodii.

THE FINGERS ARE MOVED

<i>Forwards, or flexed, by</i>	<i>Backwards, or extended, by</i>	<i>Outwards, to radial border, by</i>	<i>Inwards by</i>
Flexor sublimis.	Extensor communis.	Abductor indicis.	Abductor digiti minimi,
" profundus.	" minimi digiti,	" digiti minimi,	Interossei.
Lumbricales,	Indicator.	Interossei.	
Interossei.			
Flexor brevis digiti minimi,			
Abductor digiti minimi.			

THE THIGH IS MOVED

<i>Forwards by</i>	<i>Backwards by</i>	<i>Inwards by</i>	<i>Outwards by</i>
Psoas magnus,	Gluteus maximus.	Psoas magnus,	Tensor vaginæ femoris
Iliacus,	Part of gluteus medius,	Iliacus.	Gluteus maximus,
Tensor vaginæ femoris,	Piriformis.	Pectineus.	" medius,
Pectineus.	Obturator internus.	Gracilis.	" minimus,
Adductor longus,	Part of adductor magnus,	Adductor longus,	Piriformis.
brevis.	Long head of biceps,	" brevis,	
	Semi-tendinosus,	" magnus,	
	Semi-membranosus.	Obturator externus,	
		Quadratus femoris.	

THE THIGH IS ROTATED

<i>Inwards by</i>	<i>Outwards by</i>
Tensor vaginæ femoris,	Gluteus maximus,
Part of gluteus medius,	Part of gluteus medius,
	Piriformis.
<i>And, when the leg is extended, by</i>	Gemellus superior,
Sartorius.	Obturator internus,
Semi-tendinosus.	Gemellus inferior,
	Quadratus femoris,
	Obturator externus,
	Psoas magnus,
	Iliacus.
	Adductor longus,
	" brevis,
	" magnus,
	Biceps cruris, slightly.

THE LEG IS MOVED

<i>Backwards, or flexed, by</i>	<i>Extended by</i>
Semi-tendinosus,	Rectus.
Biceps,	Crureus.
Semi-membranosus,	Vastus externus,
Gracilis,	" internus.
Sartorius.	
Popliteus.	

THE FOOT IS MOVED

<i>Forwards, or flexed, by</i>	<i>Backwards, or extended by</i>	<i>Inclined inwards by</i>	<i>Outwards by</i>
Tibialis anticus.	Gastrocnemius,	Extensor proprius pollicis,	Peroneus longus,
Extensor proprius pollicis,	Plantaris,	Flexor longus digitorum,	" brevis.
Extensor longus digitorum,	Soleus.	" longus pollicis,	Extensor longus digitorum,
Peroneus tertius.	Flexor longus digitorum,	Tibialis posticus.	Peroneus tertius.
	" longus pollicis,		
	Tibialis posticus,		
	Peroneus longus,		
	" brevis.		

THE TOES ARE MOVED

<i>Backwards, or flexed, by</i>	<i>Forwards, or extended, by</i>	<i>Inclined inwards by</i>	<i>Outwards by</i>
Abductor pollicis.	Extensor longus digitorum,	Abductor pollicis,	Adductor pollicis.
Flexor brevis digitorum,	Extensor proprius pollicis,	Interossei.	" digiti minimi,
Abductor minimi digiti	" brevis digitorum.		Interossei.
Flexor longus pollicis,			
" digitorum.			
" accessorius,			
Lumbricales,			
Flexor brevis pollicis,			
Adductor pollicis			
Flexor brevis minimi digiti.			
Interossei.			

Muscles have been variously named. 1. *According to their uses*, as diaphragm, buccinator, extensors, flexors, adductors, abductors, levators, depressors, &c. 2. *According to their position*, as interspinales, interossei, subclavius, popliteus, anconeus, cubitalis, iliacus, temporalis, &c. 3. *According to their shape*, as trapezius, splenius, lumbricalis, serratus, digastric, deltoid, scalenus, rhomboides, &c. 4. *According to their dimensions*, as pectoralis major, rectus capitis anticus major, pectoralis minor, glutæus maximus, medius, and minimus. 5. *According to their direction*, as obliquus abdominis, transversalis abdominis, rectus femoris, rectus abdominis, &c. 6. *According to their composition*, as semi-membranosus, semi-tendinosus, complexus, &c. 7. *According to their attachments*, or the different points of the skeleton with which they are connected by means of tendons or aponeuroses; as sterno-clavido-mastoideus, sterno-hyoideus, &c. On this is grounded the nomenclature of M. Dumas, and that of Chaussier.

The end of the muscle, which adheres to the most fixed part, is usually called the *origin* or *head*, (F.) *Tête*; and that which adheres to the more moveable part, the *insertion* or *tail*, (F.) *Queue*; the intervening part or *body* of the muscle being called the *venter* or *belly*, *Venter musculi*, *Medium musculi*, (F.) *Ventre*: hence the names gastrocnemii, digastricus, biceps, and triceps; according as they have two bellies, two or three heads, &c.

Muscles are formed,—1. Essentially of the *muscular* or *fleshy fibre*, (see Muscular Fibre.) 2. Of *Areolar tissue*, which unites together the fibres. This areolar tissue is not very visible between the fine and loose fibres; but becomes more so, when they unite in more considerable fasciculi. It forms, moreover, to each muscle, an external envelope, which unites it to the neighbouring parts, and admits of its motion. This envelope was formerly called *Tunica propria musculorum*. 3. Of *Arteries*. These proceed from neighbouring trunks, and are, generally, very large. Their size and number are always in proportion to the bulk of the muscle. With the exception of some viscera, as the lungs and the kidneys, there are few organs that receive as much blood as the muscles. 4. Of *Veins*. They follow the same course in the muscles as the arteries. Bichat asserts that they are generally devoid of valves. 5. Of *Lymphatics*. Of these we know little, and cannot easily follow them between the fleshy fibres. 6. Of *Nerves*. These are numerous, and of different sizes. They, almost all, proceed from the encephalon; some, however, issue from ganglions, and accompany the arteries. In general, they penetrate the fleshy tissue along with the vessels, with which they are closely united. After they have entered the muscles, they divide and subdivide until they are lost sight of.

MUSCLE CANIN, Levator anguli oris—m. Cannelé, Ischio-trochanterianus.

MUSCLE of GAVARD. The oblique muscular fibres of the stomach.

MUSCLE GRAND FESSIER, Glutæus major—m. of Guthrie, Compressor Urethræ—m. of Horner, Tensor Tarsi—m. of Houston, Compressor venæ dorsalis penis—m. très Large du dos, Latissimus dorsi—m. of Wilson, Compressor urethræ.

MUSCULAR, *Muscula'ris*, *Musculo'sus*, *Toro'sus*, (F.) *Musculaire*, *Musculeux*. That which belongs or relates to the muscles. Well furnished with muscles, *Lacerto'sus*.

MUSCULAR ARTERIES. Arteries that are distributed to the muscles. The name, *Muscular Arteries* of the eye, has been especially given to

two branches sent off by the ophthalmic artery:—the one, *inferior*, which furnishes branches to the rectus inferior oculi, lachrymal sac, &c., and some of the anterior ciliary arteries: the other—the *superior*, which is sometimes wanting; but, when it exists, gives branches to the rectus superior oculi and levator palpebræ superioris.

MUSCULAR CONTRACTION. The exertion of the power, possessed by muscles, of shortening themselves, or of contracting to produce motion:—*muscular motion* being the change in the situation and relation of organs, induced by muscular contraction. When a muscle contracts, its fibres assume more the zigzag direction, and the extremities approximate; but the bulk of the whole muscle is not augmented. This contraction takes place, at times, with extreme velocity: a single thrill, in the letter R, can be pronounced in the 1-30,000th part of a minute. The *force* of contraction, *Myody'namis*, depends upon the healthy physical condition of the muscle, combined with due energy of the brain. The *duration*, in voluntary motion, is for a certain time dependent upon the will:—contractions, excited involuntarily, cannot be so long maintained.

MUSCULAR CURRENT. A current, which, according to Matteucci, exists in every living muscle, whose tendinous extremities are not equally disposed—one, for example, forming a cord; the other broad and ribbon-like—and which circulates in the muscle from the tendinous extremity to the muscular surface.

MUSCULAR FIBRE, *Fleshy fibre*, *Filum muscula'rè*, *Fibrilla muscula'ris*. A name given to the filaments, which, by their union, form the muscles. This fibre is flat, soft, downy, linear, but little elastic, and more or less red. It is firmer in adults than in the young or the aged; of the same size in the great and small muscles; and runs its course without bifurcation or ramification. It is but slightly resisting in the dead body, tearing readily; but during life it supports very great efforts without laceration. It is, itself, composed of a considerable number of fibrils, similar to each other, and subdividing almost *ad infinitum*. The ultimate filaments into which the fibre can be decomposed by mechanical means seem to be hollow or tubular. The fibre exists under two forms, the *striated* or *striped*, (F.) *Fibre striée*, and the *non-striated*, *smooth*, or *unstriated*, (F.) *Fibre lisse*; the muscles composed of the former ministering, as a general rule, to the *animal functions*,—the latter always, perhaps, to the *organic*. The colour of the muscular fibres is red in man, and white in several animals. A greater or less number,—united in fasciculi, *Fasciculi* seu *Lacer'ti musculorum*, approximated to each other, and forming a distinct mass, of very variable size and shape, the extremities being attached to bones by means of tendons,—constitutes a muscle. In this are included areolar membrane, vessels and nerves. See Muscle.

MUSCULAR NERVES. The nerves distributed to the muscles are so named. Winslow calls the 4th pair of nerves—*Nerf musculaire oblique supérieur*.

MUSCULAR SYSTEM. A term given to the aggregate of the muscles of the body.

MUSCULAR VEINS. These bring back the blood, which has been carried to the muscles by the muscular arteries.

MUSCULARIS, Corroborant.

MUSCULATION, See Locomotion.

MUSCULEUX, Muscular.

MUSCULI ACCESSORII AD SACRO-LUMBALUM, see Sacro-lumbalis—m. Papillares, Columnæ carneæ—m. Pectinati, Pectinated muscles. **MUSCULINE**, see Fibrin.

MUSCULO-CUTANEOUS, *Musculo-cutaneus*, from *musculus*, 'a muscle,' and *cutis*, 'skin.' That which appertains to muscles and skin.

MUSCULO-CUTANEOUS NERVE. This name is especially given to two nerves;—the one, the *External cutaneous nerve*, furnished by the brachial plexus (see *Cutaneous*:—) the other, given off from the popliteus externus, (*F.*) *Nerv. Sciatique*, *Poplite externe*, *Prétibio digital* (*Ch.*). It descends on the anterior and outer part of the leg,—at first, hid among the muscles; becomes superficial about the middle of the leg; and divides into two branches, which pass superficially on the back of the foot. Two musculo-cutaneous nerves—superior and inferior—proceed from the first lumbar. The *superior musculo-cutaneous—il'io-scrotal*, *il'io-hypogast'ricus*—which divides into two branches,—abdominal and scrotal. The *inferior musculo-cutaneous* is smaller than the superior, and is distributed as its name, *il'io-in'guinal*, indicates.

MUSCULO-RACHIDÆ'US. Belonging or relating to muscles and to the spine. A name given to branches, furnished behind by the intercostal, lumbar, and sacral arteries, which are distributed to the spine and to the muscles of the neighbourhood.

MUSCULOSA EXPANSIO, *Platysma myoides*.

MUSCULO-SPIRAL NERVE, *Radial nerve*.

MUSCULOSUS, *Muscular*.

MUSCULUS ACCLIVIS, *Obliquus internus abdominis*—*m. Auxiliarius*, *Pyramidalis abdominis*—*m. Constrictorius*, *Sphincter*—*m. Cutaneus*, *Platysma myoides*—*m. Eustachii*, *Laxator tympani*—*m. Fallopii*, *Pyramidalis abdominis*—*m. Femoris membranous*, *Fascia lata*—*m. Patientiæ*, *Levator scapulæ*—*m. Penicillatus*, *Levator labii inferioris*—*m. Pyxoides terrestris*, *Lichen pyxidatus*—*m. Scandularius*, *Parathenar*—*m. Subcutaneus*, *Platysma myoides*—*m. Succenturiatus*, *Pyramidalis abdominis*—*m. Succingens*, *Diaphragm*—*m. Supercilii*, *Corrugator supercilii*—*m. Testicondus*, *Cremaster*—*m. Testis*, *Cremaster*—*m. Tubæ novæ*, *Circumflexus*.

MUSCUS ARBOREUS, *Lichen plicatus*—*m. Caninus*, *Lichen caninus*—*m. Catharticus*, *Lycopodium selago*—*m. Clavatus*, *Lycopodium*—*m. Corallinus*, *Corallina Corsicana*—*m. Cranii humani*, *Lichen saxatilis*—*m. Erectus*, *Lycopodium selago*—*m. Helminthocortos*, *Corallina Corsicana*—*m. Islandicus*, *Lichen islandicus*—*m. Marinus*, *Corallina Corsicana*—*m. Maritimus*, *Corallina*—*m. Pulmonarius quercinus*, *Lichen pulmonarius*—*m. Pyxidatus*, *Lichen pyxidatus*—*m. Villosus*, *Peristroma*.

MUSEAU DE TANCHE, *Os uteri*.

MUSEUM ANATOMI'UM, *Supel'lex anatom'ica*. An anatomical museum.

MUSHROOM, *Fungus*.

MUSIC, *Mu'sica*, (*F.*) *Musique*, from *muca*, 'a song.' The art of producing harmonious and cadenced sounds; an art, which has, at times, been beneficially used in diseases, particularly in those of the mind; or on which the mind could act in a salutary manner.

MUSICOMANIA, *Musomania*, from *music*, and *mania*. A variety of monomania in which the passion for music is carried to such an extent as to derange the intellectual faculties.

MUSING, *LISTLESS*, *Aphelxia otiosa*.

MUSIQUE, *Musie*.

MUSK, *Moschus*, *Mosch* (*Arab.*), (*F.*) *Musc*. A peculiar concrete animal substance, of a very diffusible odour, bitter taste, and a deep brown colour: solid, and enclosed in the preputial follicles of *Moschus moschiferus*, *Musk Deer*, (*F.*) *Chevrotain porte-musc*. It is possessed of stimu-

lant and antispasmodic properties. Dose, gr. v to ʒj in bolus.

MUSK, ARTIFIC'IAL, *Moschus factit'ius*, *Resin'na Succini*. (*Ol. succini rectif.* 1 part, *acid. nitric.* 4 parts. Digest:—a black matter will be deposited, which must be well washed with water.)

MUSKGRAPE FLOWER, *Bulbus vomitorius*.

MUSKMALLOW, *Hibiscus abelmoschus*.

MUSKROOT, *Sumbul*.

MUSKSEED, see *Hibiscus abelmoschus*.

MUSKWOOD, *Thymiana*.

MUSOMANIA, *Musicomania*.

MUSQUITO, *Mosquito*.

MUSSÆN'DA FROND'OA, *Belil'la*, *Bele'son*.

A decoction of this Ceylonese plant has been esteemed refrigerant.

MUSSEL, *Mytilus edulis*.

MUSSELLED, see *Mytilus edulis*.

MUSSITATIO, *Musaitation*, from *musitare*, *musstatum*, itself from *muscare*, 'to murmur.' A condition, in which the tongue and lips move, as in the act of speaking, but without sounds being produced. This sort of murmuring is an unfavourable sign in disease, as indicating great cerebral debility.

MUSTA'CÆUM: from *Mustacea*, a kind of laurus used in making it. A kind of wedding-cake used by the ancients, which consisted of meal, aniseed, cummin, and several other aromatics; its object—it has been conceived—being to prevent or remove the indigestion occasioned by too great indulgence at the marriage feast. *Laureolum in mustaceo quævere*—"to seek for glory in pitiful performances"—Cicero.

MUSTACHE, *Mystax*.

MUSTARD, BLACK, *Sinapis*—*m. Clammy*, *Polanisia graveolens*—*m. Essence of*, *Whitehead's*, see *Sinapis*—*m. False*, *Polanisia graveolens*—*m. Green*, *Lepidium sativum*—*m. Hedge*, *Erysimum*—*m. Mithridate*, *Thlaspi campestre*—*m. Stinking hedge*, *Alliaria*—*m. Treacle*, *Thlaspi arvense*—*m. White*, *Sinapis alba*—*m. Wild*, *Sinapis arvensis*.

MUSTELANEUS, *Galeancon*.

MUSTUM, *Wort*.

MUTACIS'MUS, *Mutacism*, from *mutakiso*, 'I use the letter *m*, *uv*, too frequently.' A vicious pronunciation; consisting, according to Sauvages, in the frequent repetition of the letters *m*, *b*, and *p*, which are substituted for others.

MUTATIO SEXÛS. 'A change or conversion of sex.' A notion prevailed in antiquity that such a conversion was possible.

MUTEOS'IS, from *mutus*, 'dumb.' Under this name is comprehended every phenomenon of expression, voluntary and involuntary, which impresses the sight or touch.

MUTILATION, *Mutatio*, *Maiming*, from *mutilus*, 'broken.' The removal or privation of a limb, or of some other external part of the body. In all countries, this crime has been punished with severity.

MUTINUS, *Penis*.

MUTISME, *Mutitas*.

MUTITAS, (*mutus*, (*F.*) *Muet*, 'dumb.') *Obtumescent'ia*, *Ala'lia*, *Anau'dia*, *Dumbness*, *Speechlessness*, (*F.*) *Mutisme*, *Mutité*. Impossibility of articulating sounds; although they can be elicited. Dumbness is often congenital, and united with deafness, of which it is an effect. In some cases it is accidental.

MUTITAS SURDORUM, *Apho'nia Surdorum*, *Surdumut'itas*, *Deaf-dumbness*, (*F.*) *Surdmutité*. Speechlessness from deafness; congenital, or produced during infancy. The subjects of this affection are called *deaf-dumb*, (*F.*) *Sourde-muets*.

MUTITÉ, *Mutitas*.

MUTO, Penis.

MUTONIATUS, Membrosus.

MUTTERKORN, Ergot.

MUTTON TEA. Prepared from a pound of mutton, freed from the fat and cut into thin slices; and a pint and a half of boiling soft water poured over them, as in the case of beef tea; except that it requires to be boiled for half an hour after the maceration, before it is strained through a sieve.

MUTUNNUS, Priapus.

MYACANTHA, Ruscus.

MYALGIA, Cramp.

MYASTHENIA, from μυς, 'a muscle,' and ἀσθενία, 'debility.' Muscular debility.

MYCE, Occlusion.

MYCES, Fungus.

MYCHMUS, Suspirium.

MYCHTHISMUS, Moaning, Suspirium.

MYCODERMIA, *My'coderm*; from μυκη, 'a mushroom,' and δερμα, 'skin.' A cryptogamous growth, which constitutes the crusts of favus. See Porrigio favosa.

MYCODERMA CEREVISIÆ, *Torula cerevisiæ*.

MYCODERMA PLICÆ, *Trichomaphyton*. A cryptogamous growth, said to be the essence of Plica.

MYCODERMITIS, Mucitis.

MYCODES, Fungoid.

MYCODIARRHŒA, see Diarrhœa.

MYCODYSENTERIA, see Dysentery.

MYCOPHTHESIS, see Bronchorrhœa.

MYCORTHOPNŒA; from μυκος, 'mucus,' and ὀρθοπνοία, 'difficulty of breathing except when in the erect attitude.' Orthopnea from excessive secretion of mucus in the air-passages — *Orthopnœa pituitosa*.

MYCOSIS, Fungus.

MYTERES, Myxæ, Nares.

MYTEROPHONIA, Rhinophonia.

MYTEROXE/ROTES, *Mycterocero'sis*, from μυκτηρες, 'the nares,' and ξηρότης, 'dryness.' *Narium siccatas*. Dryness of the interior of the nose.

MYCUS, Mucus.

MYDE/SIS, from μυδω, 'I abound with moisture,' 'I am putrid.' Moisture. In its most general sense, it means *corruption*. By some, it is used for a mucous discharge from inflamed eyelids.

MYDON, μυδων. Same etymon. Fungus or putrid flesh in certain fistulous ulcers.—Pollux. Also, flesh putrid from mouldiness.

MYDRIA/SIS, *Platyco'ria*, *Platyco'ria'sis*, *Amydri'asis*. A name given by several writers to morbid dilatation of the pupil; and, by others, to weakness of sight, produced by hydropthalmia, or superabundance of humours, from μυδω, 'moisture.'

MYDRIAT/ICUS, *Mydriat'ic*. Relating to or causing dilatation of the pupil.—Pereira.

MYELAL/GIA; from μυελος, 'the spinal marrow,' and αλγος, 'pain.' Pain in the spinal marrow.

MYELANALOSIS, *Tabes dorsalis*.

MYELAPOPLEXIA, Apoplexia myelitica.

MYELATEL/A; from μυελος, 'the spinal marrow,' and ἀτελεια, (α, privative, and τελος, 'end,') 'want of end or finish.' Incompleteness of the spinal marrow.

MYELATROPHIA, *Tabes dorsalis*.

MYÉLITE, Myelitis.

MYELI/TIS, *M. parenchymatosa seu spina'lis*, *Inflammat'io medullæ spina'lis*, *Spini'tis*, *Spino-dorai'tis*, *Notomyeli'tis*, *Rachionomyeli'tis*, *Rachial-gi'tis* (of some), (F.) *Myélite*, *Inflammation de la moëlle épinière*, *l. rachidienne*, from μυελος, 'the marrow,' and ἴτις, denoting inflammation. Inflammation of the spinal marrow or its membranes;—indicated by deep-seated burning pain in the spine, with various nervous and vascular

irregularities of function. It is not common. Dr. Marshall Hall proposes to call inflammation of the membranes of the brain *Meningitis*; that of the substance of the brain *Myelitis*.

MYELITIS PTECHIALIS, Meningitis, œrebro-spinal — *m. Spinalis*, *Myelitis* — *m. Exsudativa*, *Hydrorachis* — *m. Parenchymatosa*, *Myelitis*.

MYELOCHYSIS, *Hydrorachis*.

MY'ELOCYTES, from μυελος, 'marrow,' and κύτος, a mass, body or cell. Granules of the brain. Nuclei of the cells of the gray substance.

MYELOGANGLIITIS, *Cholera* (sporadic).

MY'ELOÏD, *Myelo'i'des*, *Myelo'des*; from μυελος, 'marrow,' and εἶδος, 'resemblance.' Marrow-like.

MYELOID TUMOR, *Fibroplastic tumour*. A sarcomatous tumour, the more characteristic constituents of which are — peculiar many-nucleated corpuscles, which have been recognised as constituents of the marrow and diploe of bones, especially in the fœtus, and in early life.—Paget.

MYELOMA, see Encephaloid.

MYELOMALACIA, *Mollities medullæ spinalis*.

MYELOMENINGITIS, Meningitis, spinal.

MYELOMYCES, see Encephaloid.

MYELON, Medulla spinalis.

MYELOPARALYSIS, Paraplegia.

MYELOPHTHISIS, *Tabes dorsalis* — *m. Sicca*, *Tabes mesenterica*.

MYELOPHY/MATA, *Tuber'cula medullæ spina'lis*, from μυελος, 'marrow,' and φυμα, 'a tubercle.' Tubercles of the spinal marrow.

MYELOPLAXE, from μυελος, 'marrow,' and πλαξ, 'a plate or lamella.' Plates or lamellæ with numerous nuclei in the marrow of bones.—Robin.

MYELORRHAGIA, Apoplexia myelitica.

MYELOSPONGUS, see Encephaloid.

MYELUS, Marrow, Medulla spinalis — *m. Dicauchenius*, *Medulla spinalis* — *m. Dorsites*, *Medulla spinalis* — *m. Notizæus*, *Medulla spinalis* — *m. Psoides*, *Medulla spinalis*.

MYENERGIA, from μυς, 'a muscle,' εν, and εργον, 'work.' Muscular strength.

MYGMUS, Suspirium.

MYIOCEPHALUM, Staphyloma.

MYIODOPSTA, Metamorphopsia.

MYITIS, Myositis, Rheumatism, acute.

MYLABRIS CICHORII, see Cantharis — *m. Punctata*, see Cantharis — *m. Pustulata*, see Cantharis.

MYLACEPHALUS, from μυλη, 'a mole,' α, privative, and κεφαλη, 'head.' A monster with only traces of viscera, and without head or extremities.

MYLACRI, Molar teeth.

MYLACRIS, Patella.

MYLE, Patella, Mole.

MYLICUS, Molar.

MYLO, in composition, from μυλη, 'the jaw.' Relating or belonging to the jaw.

MYLODI, Molar teeth.

MYLODONTES, Molar teeth.

MYLO-GLOSSUS, from μυλο, and γλωσσα, 'the tongue.' Winslow has given this name to muscular fibres, which pass from the posterior part of the myloid line of the lower jaw, and from the sides of the base of the tongue to the parietes of the pharynx. These fibres belong to the constrictor superior pharyngis.

MYLO-HYOID LINE, see Mylo-hyoideus.

MYLO-HYOID FURROW or GROOVE. A furrow which passes from the superior orifice of the inferior dental canal of the lower jaw in the same direction as the canal, and lodges the *mylo-hyoid nerve*, a branch of the inferior-dental.

MYLO-HYOID NERVE, see Mylo-hyoid furrow.

MYLO-HYOÏDEUS, from μυλο, and ὀείδης, 'the os hyoides.' *Mylo'des*, (F.) *Mylo-hyoïdien*.

The *Mylo-pharyngeus* of Morgagni and Santorini. This muscle is situate at the upper and anterior part of the neck, behind the lower jaw. It is broad, flat, and has the form of a truncated triangle. It arises from the internal oblique, *internal maxillary or mylo-hyoid line* of the lower jaw bone, and is inserted at the fore part of the body of the os hyoides. Its innermost fibres unite with those of the opposite side to form a raphe on the median line. The mylo-hyoideus raises the os hyoides and carries it forward, or it depresses the lower jaw.

MYLO-PHARYNGEUS, Constrictor pharyngis, Mylo-hyoideus.

MYLUODES, Mylo-hyoideus.

MYO, in composition, from *μῦον*, 'a muscle.' Appertaining or relating to a muscle.

MYOCARDITIS, Carditis.

MYOCEPHALITIS, *Myi'tis cephal'ica*; from *μῦο*, *κεφαλή*, 'head,' and *itis*, denoting inflammation. Inflammation of the muscles of the head.

MYOCEPHALON, Staphyloma.

MYOCELIAL'GIA, from *μῦο*, *κοιλία*, 'abdomen,' and *ἀλγῶ*, 'I suffer.' Pain in the muscles of the abdomen.

MYOCELI'TIS, *Myoceli'tis*, *Inflammatio musculorum abdominalium*, from *μῦο*, *κοιλία*, 'lower belly,' and *itis*, denoting inflammation. Inflammation of the muscles of the abdomen.

MYODEOPSIA, Metamorphopsia.

MYO'DES, from *μῦο*, and *εἶδος*, 'resemblance.' Like unto muscle. Muscular.

MYODESOPIA, Metamorphopsia.

MYODESOPSIA, Metamorphopsia.

MYODYNAMIA, *Vis muscularis*; from *μῦο*, and *δύναμις*, 'power.' The force or power of a muscle.

MYODYNAMICS, see Muscular contraction.

MYODYNAMIOMETER, Dynamometer.

MYODYNOMETER, Dynamometer.

MYODYN'IA, from *μῦο*, and *ὄδυνη*, 'pain.' Pain in the muscles,—rheumatismal or other. Rheumatism.

MYODYNIA INFLAMMATORIA, Myositis.

MYOGRAPHY, *Myogra'phia*, from *μῦο*, and *γράφειν*, 'to describe.' An anatomical description of the muscles.

MYOID, *Myoi'des*, from *μῦς*, 'muscle,' and *εἶδος*, 'resemblance.' An epithet given to tumours composed of fibre cells or muscular fibres of organic life.

MYOLEM'MA, *Myole'ma*, from *μῦο*, and *λεμμα*, 'a coat.' The membranous tube of each muscular fibre. It has also the same signification as *Sarcolemma*.

MYOLINE, from *μῦον*, 'a muscle.' A name given by Mr. Erasmus Wilson to a transparent substance that fills the cells, which, he conceives, by their juxtaposition form the ultimate muscular fibril.

MYOL'OGY, *Myolog'ia*, *Sarcolog'ia*, *Myoiol'og'ia*, from *μῦο*, and *λόγος*, 'a discourse.' That part of anatomy which treats of the muscles.

MYON, Muscle.

MYONARCO'SIS, from *μῦο*, and *νάρκη*, 'stupor.' Numbness of the muscles.

MYONITIS, Myositis.

MYON'OSUS, *Myoni'us*, *Myopathi'a*, from *μῦο*, and *νόσος*, 'disease.' A disease of the muscles.

MYOPALMUS, Subsultus tendinum.

MYOPARALYSIS, see Paralysis.

MYOPATHIA, Myonosis.

MYOPE, Myops.

MYOPHO'NIA, from *μῦο*, and *φωνή*, 'voice.' The sound of muscular contraction,—as that of the ventricles of the heart, during the systole of 'bat organ.

MYO'PIA, *Myopi'asis*, *Luscios'itas*, *Myo'sis*,

Parop'sis propin'qua, *Amblyo'pia dissito'rum*, *Dyso'pia dissito'rum*, *Dyspho'tia*, *Visus ju'venum*, from *μῦο*, 'I close,' or from *μῦς*, 'a mouse,' and *ὤψ*, 'the eye;' *My'opy*, *Short-sightedness*, *Near-sightedness*, *Mouse-sight*, *Purblindness*, (F.) *Vue courte*. V. basse. Persons who can only see objects very near. The defect is owing to the too great convexity of the eye, or too great density of the humours, and is palliated by wearing concave glasses.

MYOP'IC, *Myop'icus*; same etymon. *Short-sighted*, *Poreblind*, *Purblind*, *Spurblind*. Relating or appertaining to myopia.

MYOPODIORTHO'TICON, from *μῦοψ*, 'one that is short-sighted,' and *διορθωτικόν*, 'having power to correct.' An apparatus for the cure of short-sightedness.—A. A. Berthold.

MYOPS, *Lucio'us*, *Myo'pus*, *My'ope*. Same etymon. One affected with myopia or short-sightedness.

MYOPY, Myopia.

MYORRHEX'IS, from *μῦο*, and *ρῥῆξις*, 'rupture.' Rupture of a muscle.

MYOSARCO'MA, from *μῦον*, 'a muscle,' and *sarcoma*. A sarcomatous tumour in which muscular fibres are found.

MYO'SIS, from *μῦο*, 'I close.' *Microcor'ia*. Smallness of the pupil. *Phthisis pupillaris*. Permanent contraction of the pupil. It is usually caused by iritis, and is extremely difficult to cure. When it exists to such an extent as to obliterate the pupil, it is called *Synizesis*. Also, Myopia.

MYOSITE, Myositis.

MYOSIT'IC, *Myosit'icus*, from *Myosis*. Causing contraction of the pupil,—as opium.—Pereira.

MYOSIT'IS, *Myi'tis*, *Mysi'tis*, *Myoni'tis*, *Sarci'tis*, *Rheumatis'mus phlegmoni'des*, *Myodyn'ia inflammato'ria*, *Phlegma'sia myoica*, *Inflammatio seu Phleg'monē Musculo'rum*, (F.) *Myosite*, *Inflammation des muscles*, from *μῦον*, 'a muscle.' A name proposed by Sagar for inflammation of the muscles. Also, Rheumatism.

MYOSOTIS, *Hieracium pilosella*, Rheumatism, Rheumatism, acute.

MYOSPASMUS, Cramp.

MYOTIL'ITY, *Myotil'itas*, (F.) *Myotilité*, from *μῦον*, 'a muscle.' Muscular contractility. —Chaussier.

MYOT'OMY, *Myotom'ia*, from *μῦο*, and *τεμνειν*, 'to cut.' The part of practical anatomy which treats of the dissection of the muscles. Also, the surgical operation of the division of muscles to remove deformity.

MYOTYRBE, Choreia.

MYRAC'OPUM, from *μυρον*, 'an ointment,' *a*, 'privative,' and *κοπος*, 'fatigue.' An ointment used by the ancients in cases of fatigue.

MYREPSUS, Unguentarius.

MYR'IAGRAMME, from *μυρία*, '10,000,' and *γράμμα*, 'gramme.' A weight equal to 10,000 grammes, or to 26 pounds, 9 ounces, and 6 drachms Troy.

MYRIAMÈTRE. A measure of 10,000 metres; equal to 16 miles, 1 furlong, 156 yards, and 14 inches.

MYRICA ASPLENIFOLIA, *Comptonia asplenifolia*—m. Cerifera, see *Cera flava et alba*, and Wax, myrtle.

MYRICA GALE, *M. palustris*, *Myrtus Braban'tica* seu *Ang'lica*, *Myrtifolia Belg'ica*, *Gale*, *Gagel*, *Rhus sylvest'ris*, *Ac'aron*, *Elæagnus*, *E. Cordo*, *Chamaelag'nus*, *Dodonæ'a*, Dutch myrtle, Sweet gale, Sweet willow, Candleberry myrtle, (Sc.) Scotch gale. (F.) *Piment Royal*, *Thé de Simon Pauli*, *Gale odorant*. Ord. Myricaceæ. Sex. Syst. Dicoica Tetrandria. The leaves, flowers, and seeds have a strong, fragrant smell, and a bitter taste. They are used for destroying moths

and cutaneous insects. The infusion is given internally as a stomachic and vernifuge.

MYRICA PALUSTRIS, M. Gale.

MYRINGA, see Tympanum.

MYRINGITIS, *Inflammatio tympani*, from *Myringa*, 'the membrana tympani'; and *itis*, denoting inflammation. Inflammation of the membrana tympani; and, also, of the tympanum.

MYRINX, see Tympanum.

MYRIS, *μυρίς*, *Myrothe'ce*, *Narthe'cia*, *Narthe'cium*, *Narthes*, from *μυρον*, 'a perfumed oil or ointment.' A perfumed oil jar or ointment box.

MYRISTICA, *M. Moscha'ta* seu *aromat'ica* seu *officina'lis* seu *fragens*, *Comacon*. *Nat. Ord.* Myristicæ. The tree which produces the nutmeg and mace. (F.) *Muscadier*. The *Nutmeg*, *Myristica*, (Ph. U. S.) *Myristicæ* seu *Myristicæ moscha'tæ nucleus*, *Nux moscha'ta* seu *unguenta'ria*, *Nucis'ta*, *Nux myrist'ica* seu *aromat'ica*, *Chryso'balanus Gale'ni*, *Unguenta'ria*, *As'sola*, (F.) *Muscade*, is the seed or kernel. It has a fragrant, aromatic odour; an agreeable pungent taste, and is much used for culinary purposes. Alcohol extracts its active matter. It has the properties of aromatics in general; being stimulant and stomachic. The oil—*O'leum myristicæ* seu *nucis'tæ æther'eum*—possesses the virtues of the nutmeg. Dose of the nutmeg, gr. v. to ʒj; —of the oil gtt. iij. to gtt. vj.

Mace, *Macis*, *Maces*, *Flores macis* seu *ma'cidos*, (F.) *Fleurs de muscade*, is the involucrem of the fruit. It is membranous, with the odour and taste of the nutmeg, and is possessed of similar qualities. The *O'leum macis* seu *ma'cidis* is a fragrant, sebaceous substance, expressed in the East Indies. It is only used externally. It is the *Adepe myrist'ica*, *Myristicæ o'leum express'um*, *O'leum* seu *Bal'samum nucis'tæ*, *Butter of nutmegs*, *Express'ed oil of Mace*.

MYRISTICÆ NUCLEUS, see *Myristica moschata*.

MYRISTICATIO HEPATIS, Liver, nutmeg.

MYRMECIA, Formica.

MYRMECIASIS, *Convulsio cerealis*, Formication.

MYRMECIASMUS, *Convulsio cerealis*, Formication.

MYRMECISMUS, *Convulsio cerealis*, Formication.

MYRMECIZON, Formicant.

MYRMECOSIS, *Convulsio cerealis*, Formication.

MYRMEX, Formica.

MYROBALANUS, from *μυρον*, 'an ointment,' and *βαλανος*, 'a nut'; so called, because formerly used in ointment; *Angeloc'acos*, *Myro'bolan*, *Glans unguenta'ria*, *Palma unguentario'rum*. A dried Indian fruit, of the plum kind, of different species of *Terminalia*. *Ord.* *Combretacæ*. Of this there are several varieties—the *M. Bellir'ica* or *Belliric myrobalan*, *Bellegu*, *Bellerigi*, *Belniley*;—the *M. cheb'ula* or *Chebule myro'balan*; the *M. cit'rina* or *Yellow myrobalan*, *Ara'ra*; the *M. Emblica* or *emblic myrobalan*, and the *M. Indica*, *As'uar*, *Indian* or *Black myrobalan*. All the myrobalans have an unpleasant, bitterish, very austere taste; and strike an inky blackness with a solution of steel. They are said to possess laxative as well as astringent properties.

MYROLÉS, see *Olea volatilia*.

MYRON, Myrum, Unguentum.

MYROISSOCERON. A topical application in alopecia, referred to by Galen; from *μυρον*, 'ointment,' *ρ'icca*, 'pitch,' and *κνος*, 'wax.'

MYROPEUS, Unguentarius.

MYROPOLES, Apothecary, Unguentarius.

MYROSPERMUM FRUTESCENS, *Myroxylon Peruiferum*—m. *Peruiferum*, see *Myroxylon Peruiferum*—m. of *Sonsonate*, see *Myroxylon*

Peruiferum—m. *Toluiferum*. see *Toluifera Balsamum*.

MYROXYLON, see *M. Peruiferum*.

MYROXYLON PERUIFERUM, *Myrospermum frutescens* seu *Peruiferum*, *Quinquino*, from *μυρον*, 'an ointment,' and *ξύλον*, wood; *Caburei'ba*. *Nat. Ord.* *Leguminosæ*. Supposed to be the tree which affords the *Peruvian balsam*, *Bal'samum Peruianum* (Ph. U. S.) seu *Peruianum*, *Putzochill*, *Myrox'yl' Peruiferi* *Bal'samum*, *Myrox'ylon*, *Cabureiciba*, *Indian*, *Mexican*, or *American balsam*, (F.) *Baume de Pérou* ou *des Indes*. This balsam consists of benzoic acid, resin, and essential oil. Its odour is fragrant and aromatic; taste hot and bitter: it is soluble in alcohol, and miscible in water by the aid of mucilage. It is stimulant and tonic, and considered to be expectorant: as such, it has been employed in paralysis, chronic asthma, chronic bronchitis and rheumatism, gleet, leucorrhœa, &c., and externally, for cleansing and stimulating foul indolent ulcers. Dose gtt. v. to gtt. xxx.

White Balsam of Peru, *Natural balsam*, *Bal'samum album*, *Syrax alba*, *Balsamelæon*, is obtained by incision from *Myrospermum peruiferum*.

The *Myroxylon of Sonsonate* in Central America—*Myrospermum Pereira* of Dr. Royle, has been supposed to furnish balsam of Peru.

MYROXYLON OF SONSONATE. see *M. Peruiferum*—m. *Toluiferum*, *Toluifera Balsamum*.

MYRRHA, Heb. *מוֹר*; *Bola*, *Stactê*, *Ergas'ma*, *Myrrh*, *Calo'nia*, *Smyrna*, *Myrrha rubra*, (F.) *Myrrhe*. The exudation, *μυρον*, of an unknown plant of Abyssinia or Arabia Felix, said to be the *Balsumaden'dron myrrha*. This gum-resin has a fragrant, peculiar odour; and bitter aromatic taste. It is in reddish-yellow, light, brittle, irregular tears; partially soluble in distilled water when aided by friction. S. G. 1.360. It is stimulant; and has been used in cachectic affections, humoral asthma, chronic bronchitis, &c. Dose, gr. x. to ʒj.

MYRRHA IMPERFECTA, *Bdellium*.

MYRRHINE, Myrtus.

MYRRHIS ANNUA, *Athamanta Cretensis*—m. *Major*, *Cherophyllum odoratum*—m. *Odorata*, *Cherophyllum odoratum*.

MYRSINA AFRICANA, *Tatzé*.

MYRSINE, Myrtus.

MYRSINELÆON, from *μυρτιν*, 'the myrtle,' and *ελαον*, 'oil.' Oil of myrtle.—*Dioscorides*.

MYRSINITES, *μυρτινιτης*. Wine in which branches of myrtle have been macerated.

MYRTACANTHA, *Ruscus*.

MYRTE COMMUN, Myrtus.

MYRTIDANON, *μυρτιδανον*. An excrescence, growing on the trunk of the myrtle, and used as an astringent. With Hippocrates, either the myrtus caryophyllata or myrtus pimenta. [?] Also, a wine—*Vinum Myrtid'anum*—made from wild myrtle berries.

MYRTIFOLIA BELGICA, *Myrica gale*.

MYRTIFORM, *Myrtiform'is*, *Myrto'des*, from *myrtus*, 'a myrtle,' and *forma*, 'shape.' Having the shape of a leaf of myrtle. A name given to a muscle (depressor alæ nasi), to the fossa incisiva; and to certain crurales, &c.

MYRTIFORMIS, *Compressor naris*, *Depressor alæ nasi*.

MYRTITES, from *μυρτιν*, the 'myrtle berry.' A name given to a medicine prepared with honey and myrtle berries.

MYRTLE, Myrtus—m. *Berry*, *Vaccinium myrtillus*—m. *Candleberry*, *Myrica gale*—m. *Dutch*, *Myrica gale*—m. *Wild*, *Ruscus*.

MYRTOCHEILIDES, *Nymphæ*.

MYRTOCHILA, *Nymphæ*.

MYRTODES, *Myrtiform*.

MYRTON, Clitoris.

MYRTUS, *Myrtus communis* seu *communis Italica*, *myrtos*, *Myrsi'nē*, *Myrrh'i'nē*, *The Myrtle*, (F.) *Myrte Commun.* Ord. *Myrtacæ*. The berries of this plant have been recommended in alvine and uterine fluxes and other disorders of relaxation and debility. They are moderately astringent and somewhat aromatic.

MYRTUS ANGLICA, *Myrica gale*—m. *Brabantica*, *Myrica gale*.

MYRTUS CARYOPHYLLA'TA, *Calyptanthus caryophyllata*, *Cassia* seu *Canel'la caryophyllata*, (F.) *Capelet*, *Cannelle giroflée*. The tree which is considered to afford the *Clove bark*; *Cortex caryophyllatus* seu *caryophyllata*. This bark is a warm aromatic; resembling clove with an admixture of cinnamon. It may be used with the same views as cloves or cinnamon.

MYRTUS CARYOPHYLLUS, *Eugenia caryophyllata*—m. *Leucodendron*, *Melaleuca cajuputi*.

MYRTUS PIMENTA, *Eugenia Pimenta*. The tree which bears the *Jamaica pepper*, *Pimenta buccæ*, *Pimento berries*, *Pimento*, *Piper caryophyllatum* seu *chia'pæ* seu *odora'tum Jamaicensē* seu *Jamaicensē* seu *tabac'um*, *Coc'culi Indi aromati'ci*, *Ano'mum pimenta*, *Fructus pimenti*, *Carive*, *Caryophyllus America'nus* seu *Pimenta*, *Allspice*, (F.) *Poire de Jamaïque*, *Toute épice*, *Assourou*. The unripe berries—*Pimenta* (Ph. U. S.)—have an aromatic odour; resembling a mixture of cinnamon, nutmeg, and cloves; the taste is pungent, but mixed, like the odour. Like other peppers, this is stimulant and carminative. The oil—*O'leum Pimentæ*—possesses the virtues of the berries. The powdered fruit has been called *Quatre épices* or *Four spices*. Dose, gr. v. to ʒij.

Aqua Pimentæ, *Pimento water*, is official in the London, Edinburgh, and Dublin Pharmacopœias. It is carminative. Dose, fʒi or fʒij.

MYRUM, *Myron*, *myron*. A perfumed oil or ointment. A liquid perfume. The spontaneously exuding juice of many plants, especially of that from which myrrh is obtained.

MYS, Muscle.

MYSIOLOGIA, Myology.

MYSTITIS, Myositis.

MYSTAX, *Mastax*, *μυσταξ*, *μυσταξες*. The hair growing on each side of the upper lip in men. The *mustache* or *mustachiu*, (Old Eng.) *Munchat'oe*, (F.) *Moustache*. Also, the upper lip.

MYSTERION, *μυστήριον*, 'a mystery.' An arcanum, nostrum, or secret preparation in general. Also, an antidote referred to by Galen and others.

MYSTRON, *μυστρον*. A Greek measure, which held about three drachms.

MYTILUS EDU'LIS. The common mussel, (F.) *Moule*. A bivalve, the flesh of which, when at all in a state of decomposition, is highly poisonous. When fresh, it is an agreeable, but not very digestible article of diet, producing, in some persons, all the symptoms of urticaria. One affected with such phenomena is said, occasionally, to be *musselled*.

MYU'RUS, *Meiurus*. A pulse is so called when it sinks progressively and becomes smaller and smaller like a rat's tail; from *μῦς*, 'rat,' and *οὐρα*, 'tail.' *Pulsus myurus reciprocus*, a pulse, which, after having become gradually weaker resumen, by degrees, its former character.

MYXA; Mucus, Sebestina.

MYXÆ, Mycteres.

MYXIOSIS, Gonorrhœa impura.

MYXODES, Muciform.

MYXORRHOOS, *μυξορροος*, from *μῦξα*, 'mucus,' and *ρῶω*, 'I flow.' One who is subject to mucous discharge. Applied to an infant, that discharges a considerable quantity of mucus and saliva.

MYXOSARCO'MA, from *μῦξα*, 'mucus,' and *σαρξ*, *σαρκος*, 'flesh.' A tumour which is 'mucocarcinous'; partly mucous and partly fleshy.

MYXOTER, Nasus.

MYZESIS, Sucking.

N.

N. This letter, in prescriptions, is an abridgement of *Numero*, 'by number.'

NAB'ALUS ALBUS, *Prenanthes alba* seu *serpenta'ria*, *White Lettuce*, *Lion's Foot*, *Rattlesnake's Master*, *Rattlesnake root*, *Gall of the Earth*, from *ναβλα*, 'a harp,' owing to the lyrate leaves. An indigenous plant, of the order *Compositæ*, which, with several other species of the genus, is reputed to possess the power of curing the bites of serpents. The root has been used in dysentery.

NABALUS FRASERI has similar properties.

NABOT, *Saccharum candidum*.

NABOT'HI GLAN'DULÆ, *O'vula* seu *Ova Nabothi* seu *Nabothia'na*, *Ova'rium Nabothi*, *Folliculi rotundi et oblongi*. *Vesicula* seu *Bulla rotunda cervicis uteri*, *V. seminales mulierum*, *Cervicis seu Corpus'cula globosa*, *Glandula* seu *Hydatides cervicis uteri*, *Glands of Naboth*. Small, mucous crypts or follicles situate in the interstices of the duplicatures of the lining membrane of the cervix uteri. Naboth, from noticing them in a morbid condition, mistook them for ova; and hence they received the name *Ovula Nabothi*.

NACRA, *Nakra*, *Nasa*. A kind of influenza common in the East Indies.

NACRÆ, Margaritaceus.

NACREOUS, Margaritaceus.

NÆVI, see *Nævus*—n. Pigmentary, see *Nævus*.

NÆVUS, Plural *Nævi*; *Nævus mater'nus* seu *Sigil'lum*, *Nota mater'na*, *Nævus*, *Mac'ula matricis* seu *mater'na* seu *matrica'les*, *Notæ infan'tum*, *Lupus varico'sus*, *Sig'mata*, *Metrocel'ides*, *Congen'itæ notæ*, *Mother's* or *Fancy marks*, *Mother's spots*, (F.) *Envie*, *Tache de Naissance*. Spots of various kinds on the skin of children when born, which have been attributed to the influence of the maternal imagination on the fœtus in utero. They are of various appearances, some much better supplied with blood than others. Some are merely superficial or stain-like spots, *Pigmentary nævi*; others are prominent, *Vascular tumours*, and often have long, irregular hairs growing from them. These have usually been called *Moles*, *Spili*, *Spilo'ma*, *Epichro'ses spili*, &c. The red, rosy, violet or bluish spots are called by I. G. St. Hilaire, *Taches Sanguines*;—those of an intermediate hue, between the normal colour and black, *Taches melaniennes*. See *Telangiectasia*.

When *nævi* are superficial, without any disposition to enlarge or spread, they need not be

meddled with: but all those that partake of the character of aneurism by anastomosis had better be removed, where practicable.

NAEVUS SIGILLUM, Nævus.

NAFDA, Naphtha.

NAIL, Sax. næsl, (G.) Nagel, *Unguis*, *Onyx*, (F.) *Ongle*. A whitish transparent substance, similar to horn, which covers the dorsal extremity of the fingers. Three portions are distinguished in it; the *extremity*, which is free, at the end of the finger;—the *body* or middle portion adherent by its inner surface; and the *root*, *Radix* seu *Matrix unguis*, *In'timum unguis*, (F.) *Matrice de l'ongle*. The last presents two distinct parts; the one, terminated by a thin, serrated edge, is buried in a duplicature of the skin, *Vallecula unguis*, which acts the part of a *nail-follicle*; the other, called *Lu'nula* seu *Semilu'nula* seu *Sele'nē* seu *Arcus* seu *Exor'tus* seu *Anat'olē* seu *Albe'do un'guis*, is whitish and of a semilunar shape, and is situate above the part where the epidermis terminates. The nails are composed of a horny tissue, of the same nature as that which forms the hoofs, horns, and scales of different animals. When the nail has been torn off, the papillæ of the skin become covered by a soft, whitish lamina, whose consistence gradually augments. New laminæ are then formed underneath, and give the nail the thickness it ought to possess. The corneous substance, being thus constantly produced at the extremity of each of these laminæ, the whole of the nail is pushed forwards, and it would grow indefinitely, were it not cut or worn by friction.

The nails protect and support the extremities of the fingers against the impression of hard bodies. They are, also, useful in laying hold of small bodies; and dividing those that have but little consistence.

NAIL-FOLLICLE, see Nail.

NAIL SPRING, Hangnail.

NA'V. Nanus.

NAISSANCE, see Parturition.

NAKRA, Nacra.

NANGNAIL, Hangnail.

NANISME, see Nanus

NANNARI, Hemidesmus Indicus.

NANNYBERRY, Viburnum lentago.

NANOCEPHALIA, see Nancecephalus.

NANOCEPH'ALUS, from *νανός*, 'a dwarf,' and *κεφαλή*, 'head.' One of a monstrous state—*Nanoccephalia*—in which the whole head or certain of its parts are too small, whilst the trunk and extremities are normal.

NANOCORMIA, see Nanocormus.

NANOCOR'MUS, from *νανός*, 'a dwarf,' and *κορμός*, 'a trunk.' One of a monstrous condition—*Nanocormia*—in which the trunk is too small, whilst the head possesses its normal size.

NANOMELIA, see Nanomelus.

NANOM'ELUS, from *νανός*, 'a dwarf,' and *μελος*, 'a limb.' One of a monstrous condition—*Nanomelia*—in which some part of an extremity is too small, and the whole limb too short.

NANOSO'MIA, from *νανός*, 'a dwarf,' and *σώμα*, 'body.' *Microsomia*. Dwarf'ishness, (F.) *Nanisme*. A state in which the entire body with all its parts is smaller than common.

NANOSOMUS, Nanus.

NANUS, *navos*, *Pumil'io*, *Pu'milo*, *Pusil'ius*, 'a dwarf.' *Nanosomus*, (Prov.) *Crut*, *Rope Spinick*, (Sc.) *Croil*, *Crok*, *Dearch*, *Derch*, *Droich*, *Wratack*. (*Pu'milus*, *Pumil'ius*, 'dwarfish.') (F.) *Nain*. One who is much below the usual stature. The term is applicable to all organized beings from man to the vegetable. Trees have their dwarfs, as the human species have theirs. Some curious authenticated instances of human dwarfs are on record. Bebe, the dwarf of Stanislaus,

king of Poland, was 33 inches (French) long, and well proportioned. The Polish nobleman, Borwlaski, who was well made, clever, and a good linguist, measured 28 Paris inches. He had a brother, 34 inches high, and a sister 21. A Friesland peasant, at 26 years of age, had attained 29 Amsterdam inches. C. H. Stöberin, of Nürnberg, was under three feet high at 20, yet he was well proportioned and possessed of talents. General Tom Thumb, so called, was seen by the Author in 1847. He was said to be 15 years old; measured 28 inches in height, and when weighed at the mint was found to weigh 20 pounds and 2 ounces. Two remarkable Indian dwarfs were exhibited in the United States and in Europe, about the middle of the 19th century, under the name of 'Azeo children,'—the boy, from 7 to 8 years of age, was 33½ inches in height, and weighed 20½ pounds:—the girl, from 4 to 6 years of age, was 29½ inches high, and weighed 17 pounds. They were very deficient in intellect. See Pygmy.

NAPE OF THE NECK, Nucha.

NAPELLUS VERUS, Aconitum napellus.

NAPHÆ FLORES, see Citrus aurantium.

NAPHTHA, Acetone.

NAPHTHA, *ναφθα*, *Nafda*, *Napta*, *Napta'lius*, *O'leum petræ album*; (F.) *Naphte*; from a Chaldaic and Syriac word signifying *bitumen*. A bituminous substance, found in Persia, Calabria, Sicily, &c. It is liquid, limpid, of a yellowish-white colour, a smell slightly resembling that of oil of turpentine, and lighter than water. It resembles petroleum in its properties, and has been chiefly used as an external application; although occasionally, as an anthelmintic, and in inhalation in phthisis pulmonalis. Also, *Æther*.

NAPHTHA ACETI, *Æther aceticus*—n. Coal Tar, Light, see *Anæsthetic*—n. Sulphurica, *Æther sulphuricus*—n. Vitrioli, *Æther sulphuricus*—n. Vitrioli camphorata, *Tinctura ætherea camphorata*—n. Vitrioli martialis, *Tinctura seu Alcohol sulphurico-æthereus ferri*—n. Wood, Acetone.

NAPH'THALINE, *Naphthal'ina*, *Naph'thalin*. When coal tar is subjected to distillation, naphthaline passes over after coal naphtha. It is a white, shining, concrete, crystalline substance, fusible at 176° and boiling at 423°. It is soluble in alcohol, ether, naphtha and the oils, but insoluble in water. It has been used as an excitant expectorant, in the dose of 8 to 30 grains, in emulsion or syrup, and repeated. It has also been used as an anthelmintic; and, when made into an ointment, in psoriasis, lepra vulgaris, &c.

NA'PIFORM, *Napiform'is*, *Bu'noid*, *Bunioi'des*, from *napus*, 'a turnip,' and *forma*, 'shape.' An epithet for a form of scirrhus tumour whose cut surface presents concentric lines resembling those seen in the interior of a turnip.

NAPIUM, Lapsana, Sinapis.

NAPLES, (CLIMATE OF). The climate of Naples in its general characters resembles that of Nice, but it is more changeable: the sirocco too, which is little known in Nice, is severely felt at Naples. It is not a good residence for the phthisical invalid: Sir James Clark, indeed, considers it altogether unsuitable.

NAPLES, MINERAL WATERS OF. In the Quarter Santa Lucia, near the coast, is a cold spring, rich in sulphuretted hydrogen and carbonic acid. It is much used as an aperient tonic, and in cutaneous affections.

NAPLES WATER, FACTIT'IOUS, *Aqua Neapolitana* seu *acid'ula hydrosulphurata*, (F.) *Eau de Naples*. (*Acidulous water*, containing four times its bulk of carbonic acid, $\frac{3}{4}$ xv, and $\frac{3}{4}$ ij; *hydrosulphuretted water*, $\frac{3}{4}$ ix; *carbonate of soda*, gr. viij; *carbonate of magnesia*, gr. x. M. Ph. F.)

NAPTA, Nanus, Nata.

NAPTALIUS, Nanus.

NAPUS, Sinapis — n. Leucosinapis, Sinapis alba — n. Sylvestris, Brassica napus.

NAPY, Sinapis.

NARCAPHTE, Thymiamia.

NARCAPH'THON, *Nascaph'thon*. The bark of an aromatic tree formerly brought from India. By some, supposed to be that of the tree which affords the olibanum. It was used in fumigation, in diseases of the lungs. See Lacaphthon.

NARCE, Narcosis.

NARCEMA, Narcosis.

NARCESIS, Narcosis.

NARCISSE D'AUTOMNE, Colchicum autumnale — n. Fauz, Narcissus pseudo-narcissus — n. des Prés, Narcissus pseudo-narcissus — n. Sauvage, Narcissus pseudo-narcissus.

NARCIS'SUS PSEUDO-NARCIS'SUS, *N. festalis* seu *glauca* seu *grandiflorus* seu *hispanicus* seu *major* seu *serotus* seu *sylvestris*, *Bulbocodium*, *Pseudo-narcissus*, *Daffodil*, (Old Eng.) *Aff'dil*, *Belle-Blome*, (Prov.) *Affrodile*, *Bellflower*, *Butter and Eggs*, *Lent-rose*. Ord. Amaryllidaceæ. (F.) *Narcisse sauvage*, *Narcisse des prés*, *Faux narcissus*, *Pörrillon*, *Aiault*, *Zouzinette*, *Jeannette*. The root is emetic and cathartic, in the dose of ʒij. The flowers are antispasmodic. Dose, 24 grains.

NARCODES, Narcotized,

NARCO'SIS, *Narcotism*, *Narcē*, *Narco*, *Narce'ma*, *Narce'sis*, from *ναρκω*, 'I benumb.' *Torpe'do*, *Torpor*, *Stupor*, *Stupefactio*. The aggregate effects produced by narcotic substances. At times, narcotism is confined to a state of more or less profound stupor; and constitutes, in certain cases, a useful remedial condition; at others, it is a true poisoning, characterized by vertigo, nausea, a state of intoxication or apoplexy, constant delirium, convulsive motions, &c. Emetics in strong doses, and not much diluted with water; purgatives and glysters are the first means to be used in this condition. The stupor may afterwards be combated by the use of exciting and stimulating drinks.

NARCOSIS FOLLICULO'RUM. A state of the scalp, which Mr. Erasmus Wilson conceives to be dependent upon torpid action of the hair follicles, and in which the scalp and hair are found covered with a yellowish, dirty-looking powder, composed of an admixture of granular particles and furfuraceous scales.

NARCOSPASMUS, from *ναρκ*, 'stupor,' and *σπασμος*, 'spasm.' Stupor combined with spasm.

NARCOTIA, Narcotine.

NARCOTIC POISON, see Poison.

NARCOTICO-ACRID, see Poison.

NARCOTICS, *Narcotica*, *Carotica*, *Obstupefacientia*, *Stupefacientia*, *Stupeficientes*, (F.) *Stupefactifs*, *Stupefians*. Same etymon. Substances, which have the property of stupefying; — as opium, stramonium, hyoscyamus, belladonna, &c. They are used in medicine as soothing agents; exerting their special influence on the brain and tubular matter of the spinal marrow. In small doses, as a general rule, narcotics stimulate: in large, they act as sedatives. The following is a list of the chief narcotics: — Aconitum, Atheraea, Belladonna, Camphora, Cannabis, Conium, Digitalis, Humulus, Hyoscyamus, Lactucarium, Opium and Morphia, Stramonii Folia, Stramonii Semina, Mental Narcotics, (Appropriate Music, Monotonous sounds, or any succession of monotonous impressions.)

NAR'COTINE, *Narcoti'na*, *Narco'tia*, *Narco'tin*, *Anarco'ti'na*, *Opia'nium*; same etymon; (F.) *Sel de Déroïne ou d'opium*, *Opiane*, *Principe cristallisable de Déroïne*. A solid, white, inodorous, and insipid substance; by some considered

to be alkaloid; by others neuter; crystallizable in straight prisms with a rhomboidal base; fusible like the fats; soluble in boiling alcohol and ether, and scarcely soluble in water. Narcotine produces all the unpleasant effects of opium; but, at the same time, throws the animal into a state of stupor. It is not used in medicine, on account of these objections. The salts, which are very bitter, have been used successfully in India for the cure of intermittents.

NARCOTISM, Narcosis.

NAR'COTIZED, *Tor'pidus*, *Narco'des*. Affected with stupor, as from the use of a narcotic.

NARD, CELTIC, Valeriana Celtica — n. Indica, Nardus Indica — n. Indien, Nardus Indica — n. Indique, Nardus Indica — n. Petit, Aralia nudicaulis — n. Sauvage, Asarum.

NARDOSTACHYS JATAMANSI, Nardus Indica — n. Patrinia, Nardus Indica.

NARDUM GALLICUM, Valeriana Celtica.

NARDUS AMERICANUS, Aralia nudicaulis — n. Celtica, Valeriana Celtica — n. Celtica altera, Arnica montana — n. Montana. Asarum.

NARDUS IN'DICA, *Spica nardi* seu *In'dica*, *Andropogon nardus* seu *citriodorus*, Ord. Gramineæ [?], or *Nardostachys* seu *Valeria'na* seu *Patrin'ia Jataman'si*, *Valeria'na spica*; Order, Valerianaceæ; *Indian nard*, *Spikenard*. The root of this plant is one of the ingredients in the mithridate and theriaca. It is moderately warm and pungent, and has a flavour by no means disagreeable. It is used by the Orientals as a spice: — (F.) *Nard Indien ou Indique*. An ointment was formerly used, called *Unguentum nardi'num*. It was prepared of *Nard*, *Malobathrum* leaves, *Oil of worm*, *Costus*, *Anomum*, *Myrrh*, &c., and was used as a detergent.

NARDUS RUSTICA, Asarum.

NARES, *Rhines*, *Cavitas* seu *Cavum* seu *Ca'væa* seu *Caver'na na'rium*, *Nasus inter'nus*, *Mycte'res*, *Na'rium ad'itus*, *Ocheteu'mata*, *The nostrils*, (Prov.) *Snuirrels*, *Snurles*, (F.) *Narines*. Two elliptical apertures, situate beneath the nose, and separated from each other by the cartilaginous *septum* seu *colum'na nasi*. These apertures are continually open, and give passage to the air we breathe, and to the mucous fluids secreted in the nasal fossæ.

NARES INTERNE, Nasal fossæ.

NARES POSTERIOR, *Nares postre'mæ*, *Extre'mæ nares*, *Os'tia posteriora* seu *Cho'na'na'rium*, *Na'rium Ex'itus*, (F.) *Arrières narines*, *Ouvertures postérieures des fosses nasales*, are the posterior apertures of the nasal cavities, which establish a communication between those cavities and the pharynx. They are bounded, *above*, by the body of the sphenoid bone; *below*, by the palate bone, and the base of the velum pendulum; and, on the *outside*, by the internal ala of the pterygoid process. They are separated from each other by a septum, of which the vomer is the bony part.

NARIFUSO'RIA, from *nares*, 'the nostrils,' and *funderæ*, *fusum*, 'to pour.' Medicine dropt into the nostrils.

NARINES, Nares.

NARIUM ADITUS, Nares — n. Siccitas, Mycteroxerotes.

NARROW CHEST, see Lordosis.

NARTHECIA, Formulary, Myris.

NARTHECIUM, Formulary, Myris.

NARTHEX, Formulary, Myris, Splint — n. Asafoetida, see Asafoetida.

NASA, Nara, Nata.

NASAL, *Nasa'tis*, from *nasus*, 'the nose,' That which relates to the nose.

NASAL ARTERY. This is the largest of the two branches in which the ophthalmic artery terminates. It issues from the orbit, above the tendon

of the orbicularis palpebrarum, passes above the side of the root of the nose, and anastomoses with the last extremity of the facial. Haller gave the name *nasal* to the sphenopalatine. He also called the dorsals *nasi*, furnished by the external maxillary, *Nasa'les laterales*.

NASAL BONES, *Ossa nasi seu nasa'lia*, *Ossicula nasi*, *Ossa maxillæ superioris quarta seu quinta seu secunda*, (F.) *Ossa nasaux*, *Osses propres du nez*. These bones are situate beneath the nasal notch of the os frontis, and occupy the space between the nasal or angular processes of the superior maxillary bone. Their shape is nearly quadrilateral. They have an external or cutaneous surface, an internal or nasal, and four margins. Each is articulated with its fellow, with the os frontis, ethmoid, and superior maxillary bones. They ossify from a single point.

NASAL CARTILAGE, *Cartilago triangularis Nasi*. A cartilage formed of three portions, which unite at the dorsum nasi, and are distinguished into the cartilage of the septum, and the lateral cartilages. This cartilage is continuous, above, with the ossa nasi, and, inferiorly, with membranous fibro-cartilages, which form the supple and movable part of the nostrils. They are two in number—the one before, the other behind.

NASAL DUCT, *Lachrymal duct*—n. Eminence, Mesophryon.

NASAL FOSSÆ, *Cavi narium*, *Nares interiores*. Two large, anfractuous cavities, situate between the orbits below the cranium, and lined by the pituitary or Schneiderian membrane. These cavities have no communication with each other; but the various sinuses in the neighbouring bones—the ethmoidal, sphenoidal, superior maxillary, &c.—all communicate with them. The general cavity of each nostril is divided by the ossa spongiosa into three *meatus* or *passages*, which run from before backwards. 1. The *Meatus narium superior*, placed at the upper, inner, and back part of the superior spongy bone. 2. The *Meatus medius*, situate between the superior and inferior spongy bones; and 3. The *Meatus inferior*, situate between the inferior spongy bone and the bottom of the nose. The Schneiderian membrane receives the first pair of nerves, and various branches from the fifth pair. The arteries are furnished by the branches of the internal maxillary, known under the names of sphenopalatine, infra-orbital, superior alveolar, palatine, pterygo-palatine; by the supra-orbital, and ethmoidal branches of the ophthalmic artery; by the internal carotid, superior labial, and dorsals nasi. Its veins are little known, and generally follow the course of the arteries. The lymphatics are almost unknown. The nasal fossæ are the seat of smell; they aid, also, in respiration and phonation.

NASAL MEATUS, see Nasal fossæ.

NASAL MUCUS, *Mucus narium*, *Phlegma narium crassum*, *Mucor seu Pituita seu Blenna seu Lympha muculenta narium*, *Apomyxia*, vulgarly called *Snot*, *Snob*, *Snoth*, (F.) *Morve*, is the mucus secreted by the Schneiderian membrane.

NASAL NERVE, *Naso-palpebralis* (Ch.), *Naso-oculalis* (Sömmering), *Naso-ciliaris*. One of the three branches of the ophthalmic nerve of Willis. It enters the orbit by the sphenoidal fissure, passes along the inner paries of that cavity, and divides, opposite the internal and anterior orbital foramen, into two branches. 1. The internal and posterior, which passes into the *Foramen orbitarium internum anterius*, enters the cranium beneath the dura mater, and passes into the nasal fossæ, through an aperture at the side of the *Crista galli*. It afterwards divides into several filaments: one of them—the *Naso-lobar*, of

Chaussier—is very small, and descends on the posterior surface of the os nasi, and ramifies on the integuments of the ala nasi. A second terminates near the septum; others descend along the outer paries of the nasal fossæ. 2. The other is external and anterior, and is called the external nasal nerve. It is distributed to the outside of the orbit. Before dividing, the nasal nerve communicates with the ophthalmic ganglion, and gives off two or three ciliary nerves. Sömmering has given the name *Posterior Nasal Nerves* to the Nerves which arise from the internal part of the sphenopalatine ganglion.

NASAL NOTCH, (F.) *Échancrure nasale*. A semicircular notch, situate between the nasal prominence of the frontal bone, and articulated with the nasal bones and the nasal processes of the superior maxillary bones.

NASAL PROCESS, Maxillary bone, superior.

NASAL PROMINENCE, (F.) *Bosse nasale*. A prominence, situate on the median line, at the anterior surface of the os frontis, between the two superciliary arches.

NASAL REGION, *Regio nasa'lis*. The region of the nose.

NASAL SPINES. These are three in number.

1. The *Superior nasal spine* of the os frontis, occupying the middle of its nasal notch, and articulated before with the nasal bones, behind with the ethmoid. 2. The *Inferior and anterior nasal spine*, situate at the inferior part of the anterior opening of the nasal fossæ. It is formed by the two superior maxillary bones; and 3. The *Inferior and posterior nasal spine*, *Spine gutturale* (Ch.), a process, formed on the median line by the two palate bones at the posterior part of the palatine arch.

NASALE, Errhine.

NASALIS, Compressor naris.

NASA'LIS LABII SUPERIORIS. A small muscular slip, which runs up from the middle of the orbicularis and the lip to the tip of the nose. It lies exactly in the furrow, and is occasionally a levator of the upper lip, or a depressor of the tip of the nose.

NASAS, Nata.

NASATUS, Membrosus.

NAS'CALÈ. A kind of pessary, made of wool, or cotton, which was formerly introduced into the vagina, after being impregnated with oil, ointment, or some other proper medicament.

NASCAPHTHON, Narscarphthon.

NASDA, Nata.

NASI, see Oryza.

NASITAS, Rhinophonia.

NASITIS, Rhinitis—n. Postica, Angina nasalis.

NASO, *Nasutus*, from *nasus*, 'the nose.' One who has a long nose. In composition, appertaining or relating to the nose.

NASOCILIARIS (Nervus), Nasal nerve.

NASO-LABIAL, *Naso-labia'lis*; from *naso*, and *labium*, 'a lip.' Relating to the nose and lip.

NASO-LABIAL LINE. A line or furrow, which separates the lip from the cheek, and commences at the ala nasi.

NASO-LABIA'LIS. A muscular fasciculus described by Albinus, which arises from the anterior extremity of the septum nasi, and terminates in the orbicularis oris.

NASO-PALATINE, *Naso-palati'nus*. That which belongs to the nose and velum palati.

NASO-PALATINE GANGLION is situate in the foramen palatinum anterius. Its greater extremity receives the two naso-palatine branches.

whilst the smaller gives off two or three filaments, which reach the palatine vault, where they ramify on the membrane of the same name, anastomosing with filaments of the great palatine nerve.

NASO-PALATINE NERVE, *Nerve of Cotunnus*, is furnished by the sphenopalatine, which proceeds from the ganglion of Meckel. It traverses the vault of the nasal fossæ, and proceeds upon the septum between the two layers of the pituitary membrane. It enters the anterior palatine canal, and terminates at the superior angles of the nasopalatine ganglion, without attaining the mouth.

NASO-PALPÉBRAL, *Orbicularis palpebrarum*.

NASTA, *Nata*.

NASTURTIIUM AMPHIBIUM, see *Sisymbrium* — *n. Aquaticum*, *Cardamine pratensis*, *Sisymbrium nasturtium* — *n. Bursa pastoris*, *Thlaspi bursa-n*. *Hortense*, *Lepidium sativum* — *n. Indicum*, *Tropæolum majus* — *n. Officinale*, *Sisymbrium nasturtium* — *n. Palustre*, see *Sisymbrium* — *n. Peruvianum*, *Tropæolum majus* — *n. Pratense*, *Cardamine pratensis* — *n. Sativum*, *Lepidium sativum*.

NASUM DILATANS, *Pyramidalis nasi*.

NASUS, *Nose*, *Rhin*, *Rhis*, *Organon olfactus seu odoratus seu olfactorium*, *Promontorium faciei*, *Myroter*, *Emunctorium cerebri*, *Snout*, (*Sc.*) *Neis*, *Nes*, (*F.*) *Nez*. The nose is a pyramidal eminence, situate above the anterior apertures of the nasal fossæ, which it covers; and, consequently, occupying the middle and upper part of the face, between the forehead and upper lip, the orbits and the cheeks. Its lateral surfaces form, by uniting angularly, a more or less prominent line, called *Dorsum* seu *Rhachis* seu *Spina Nasi*, (*F.*) *Dos du nez*. This line terminates, anteriorly, by the lobe, — *Lobulus*. The sides are called *Alæ Nasi*, *Pinnæ Naris*, (*F.*) *Ailes du nez*. The column is the inferior part of the partition. Its apex or tip has been called *Globulus nasi*.

The chief varieties of the nose are the *aqutline*, the *flat nose*, (*F.*) *Nes camarat ou épate*, and the *snub nose*, (*F.*) *Nes retroussé*. The nose is formed, besides its bones, of fibro-cartilage, cartilage, muscles, vessels, and nerves; and its use seems to be to direct odours to the upper part of the nasal fossæ.

NASUS INTERNUS, *Nares*.

NASUTUS, *Naso*.

NATA, *Natta*, *Nasa*, *Nasda*, *Nasta*, *Nasas*, *Napta*. A fleshy, indolent excrescence, having the shape of the nates.

NATARON, *Natron*.

NATA'TION, *Nata'tio*, from *natare*, *natatum*, itself from *nare*, 'to swim.' *Swimming*. The action of swimming, or of supporting one's self, or moving upon the water. Swimming resembles the horizontal leap in its physiology — the medium being water instead of air. The difference between the specific gravity of the human body and that of water is not great; so that but little exertion is required to keep a part of the body above water. Swimming is a healthy gymnastic exercise, combining the advantages of bathing.

NATES, *Ephed'rana*, *Scaphia*, *Podex*, *Clunes*, *Glutoi*, *Gluti*, *Glut'ia*, *Sedi'tia*, *Pyge*, *Monæ*, *Bottom*, *Backside*, *Posteriori*, *Buttocks*, (*Prov.*) *Tout*, (*Sc.*) *Doup*, *Doepp*, *Dolp*, *Droddum*, *Hurdies*, *Lendie*, &c., (*F.*) *Fesses*. Two round projections, at the inferior and posterior part of the trunk, on which we sit. Amongst the mammalia, man alone has the nates prominent and round. They are formed chiefly by the skin, and a thick layer of areolar tissue, which covers the three glutæi muscles.

NATES CEREBRI, *Quadrigenina corpora* — *n. et Testes*, *Quadrigenina corpora*.

NATRIUM, *Sodium*.

NATROCRENÆ, *Natropegæ*.

NATRON, *Natrum*, *Nat'aron*, *Nitrum*, *Anatrum*, *Aphronitrum*, from *Natron*, a lake in Judea. A saline compound, very abundant in Egypt, which is almost wholly formed of subcarbonate of soda.

NATRONIUM, *Sodium*.

NATROPE'GÆ, *Natrocre'næ*, from *natron*, and *πηγη*, 'a spring.' Soda springs.

NATRUM, *Natron* — *n. Chloratum liquidum*, *Liquor sodæ chlorinatæ* — *n. Muraticum*, *Soda*, *muriate* of — *n. Muriatum*, *Soda*, *muriate* of — *n. Præparatum*, *Soda*, *subcarbonate* of — *n. Tartarizatum*, *Soda*, *tartrate* of — *n. Vitriolatum*, *Soda*, *sulphate* of — *n. Nitricum*, *Soda*, *nitrate* of — *n. Oxymuriaticum*, *Soda*, *chloride* of — *n. Oxyphosphorodes*, *Soda*, *phosphate* of — *n. Sulphuricum*, *Soda*, *sulphate* of.

NATTA, *Nata*.

NATULÆ, *Quadrigenina tubercula*.

NATU'RA, from *nasci*, 'to be born or arise.' *Physis*. *Nature*. Also, the genital organs.

NATU'RA MORBI. The essence or condition of a disease.

NATURAL PARTS, *Genital organs*.

NATURALIA, *Genital organs*.

NAT'URISM, *Nat'uralism*. A view which attributes every thing to nature, as a sage, prescient, and sanative entity. — *Nysten*. See *Expectation*.

NAT'URIST. A physician who scrupulously investigates, interprets, and follows the indications presented by nature in the treatment of disease.

NAU'CLEA GAMBIR, *Uncaria gambir*. A plant of the *Nat. Order*, *Rubiaceæ*, native of the Malayan Peninsula and Indian Archipelago, which yields large quantities of the kind of Catechu known by the names *Terra Japonica* and *Square Catechu*, and which, in Indian commerce, is called *Gambier* or *Gambir*. It is a powerful astringent, much used in tanning, and in medicine. See *Catechu*.

NAU'SEA, *Nau'sia*, *Nausi'asia*, *Nausio'sis*, *Nau'tia*, *Queas'iness*, *Squeas'iness*, (*F.*) *Nausee*, *Envie de vomir*; from *vau*, *navis*, 'a ship;' because those unaccustomed to sailing are so affected. *Sickness*. *Inclination to vomit*.

NAUSEA, *KREAT'IC*, (*xpeas*, *xpeatos*, 'flesh.') The sickness and vomiting, excited, in some nervous patients, by the smallest portion of animal food.

NAUSEA MARI'NA, *Morbus nau'ticus*, *Vomitus navigantium* seu *mari'nus*, *Sea-sickness*, (*F.*) *Mal de mer*. The sickness, vomiting, &c., experienced at sea by those unaccustomed to a sea-life; and from which those who are accustomed are not always exempt. It generally ceases when the person becomes habituated to the motion of the vessel, and not till then.

NAUSEABOND, *Nauseant*.

NAU'SEANT, *Nau'seans*, (*F.*) *Nauseabond*, *Nauseans*. An agent that excites nausea, which is a state of diminished action. *Nauseants* are, hence, valuable remedies in diseases of excitement.

NAUSIA, *Nausea*.

NAUSIASIS, *Nausea*.

NAUSIO'SIS. This word, besides being synonymous with nausea, has been used to express the state of venous hemorrhage, when blood is discharged by jets.

NAUTIA, Nausea.

NAVEL, Umbilicus—*n.* String, Funiculus umbilicalis—*n.* Wort, Cotyledon umbilicus—*n.* Wort, Venus's, Cotyledon umbilicus.

NAVET, Brassica rapa—*n.* du Diable, Bryonia alba—*n.* Galant, Bryonia alba.

NAVETTE, Brassica rapa.

NAVIC'ULAR, *Navicula'ris*, *Navifor'mis*, (F.) *Naviculaire*, from *navicula*, 'a little ship.'

NAVIC'ULAR FOSSA, *Fossa Navicula'ris* seu *Scaphoi'des*, *Navic'ula*, *Scaph'ula*. See Scaphoid. This name has been given, 1. To a small depression between the entrance of the vagina and the posterior commissure of the labia major or fourchette. 2. To a perceptible dilatation, presented by the urethra in man, near the base of the glans. 3. To the superficial depression which separates the two roots of the helix. This is also called *Scapha*.

NAVICULARE OS, Scaphoides os.

NAVIFORMIS, Navicular.

NAVIS, Vulva.

NEAR-SIGHTEDNESS, Myopia.

NEARTHRO'SIS, from *neos*, 'young or new,' and *arthron*, 'a joint.' A new joint.

NEBLE, Nipple.

NEB'ULA, *Nubes*, *Nubec'ula*, *Nephos*, *νεφελή*, *Neph'elē*, *Nephe'lon*, *Vitus nebulo'sus*, (F.) *Nuage*, *Ombage*. A slight speck on the cornea. A mist or cloud suspended in the urine. See Caligo.

NEBULOUS, Nepheloid.

NEC'ESSARY, *Necessa'rius*, (*ne*, and *cessare*.) The *Necessaries of Life*, *Vita necessita'tes*, (F.) *Besoins de la vie*, include every thing requisite for the maintenance of life, and particularly food.

NECK, Collum—*n.* Derbyshire, Bronchocele.

NECK OF THE FOOT, Instep.

NECK, SWELLED, Bronchocele—*n.* Stiff, Torticollis—*n.* Wry, Torticollis.

NECKLACE, AN'ODYNE, (F.) *Collier anodin*. These are formed of the roots of hyoscyamus, Job's tears, allspice steeped in brandy, or the seeds of the wild liquorice vine, to suit the fancy of the prescriber. They are employed to facilitate dentition in children, and to procure sleep in fever. (!)

NECKWEED, *Veronica beccabunga*, *V. peregrina*.

NECRÆ'MIA, from *nekros*, 'death,' and *aima*, 'blood.' Death of the blood. Death beginning with the blood.

NECENCEPHALUS, Mollities cerebri.

NECROCEDIA, Embalming.

NECROCOMIUM, *Morgue*.

NECRODES, Cadaverous.

NECRODOCHIUM, *Morgue*.

NECROLOGY, BILLS OF, Mortality, bills of.

NEC'ROMANCY, *Necromanti'a*, *Negromanti'a*, *Necyomanti'a*, *Nigromanti'a*, from *nekros*, 'death,' and *μαντεία*, 'divination.' Divination by the dead.

NECROMANTIA, Necromancy.

NECRON, Cadaver.

NECRONARCEMA, Rigor mortis.

NECROPHOB'IA, from *nekros*, 'death,' and *φοβος*, 'fear.' Exaggerated fear of death. This symptom occurs in patients where the disease is not mortal; as in hypochondriasis. In fevers, it is not a good symptom.

NECROPNEUMON'IA, *Pneumo'nia gangræno'sa* seu *typho'sa*, *Gangræ'na* seu *Mortifica'tio* seu *Anthrax* seu *Carbunculus Pulmo'nium*, *Pneumosep'tis*, *Pneumocac'hē*, (F.) *Gangrène du Poumon*, *Pneumonécrose*, — Piorry; from *nekros*,

'death,' and *pneumonia*. Gangrenous inflammation of the lungs. This may be *diffused* or *circumscribed*. The only pathognomonic symptom is the extraordinary and repulsive odour of the breath and expectoration. The treatment consists in the use of the chlorides internally, or of chlorine by inhalation; allaying irritation by opium, and supporting the patient by wine-whey, and nourishing diet.

NECROPSIA, Autopsia cadaverica.

NECROPSY, Autopsia cadaverica.

NECROSCOPIA, Autopsia cadaverica.

NECROSCOPY, Autopsia cadaverica.

NECROSED, see Necrosis.

NECROSEMIOTICE, from *νεκρος*, 'death,' and *σημειον*, 'a sign.' The doctrine of the signs of death.

NECRO'SIS, from *νεκρω*, 'I kill.' Mortification. State of a bone or of a portion of a bone deprived of life. *Osteogangræ'na*, *Osteonecros'is*. The bone, in such case, is said to be *necrosed*. Necrosis may take place without the surrounding soft parts being struck with gangrene. It is to the bones what gangrene is to the soft parts. The part of the bone affected with necrosis becomes a foreign body, similar to the gangrenous eschar, and its separation must be accomplished by the efforts of nature, or by art. When necrosis occurs in the centre of long bones, it never extends to their articular extremities. The exterior layers of bone form a canal round the dead portion or *sequestrum*; between these swollen layers and the sequestrum, suppuration takes place; the matter presses against the bony canal; perforates it, and is discharged by apertures, which become fistulous.

In the treatment, the exit of the sequestrum must be facilitated by proper incisions, by the application of the trepan to the bone, &c.

NECROSIS CERVALIS, Ergotism — *n.* Dentium, Dental gangrene — *n.* Ustilaginea, Ergotism.

NECROSTOA, *Morgue*.

NECROTOMY, Dissection.

NEC'TANDRA CINNAMOMOI'DES, *Laurus cinnamomoi'des*, *Cinnamomum sylvestrē Americānum*: Order, Lauraceæ; native of New Grenada, where it is called *Canela*; has the odour and taste of cinnamon, and is used as such.

NECTANDRA CYMBARUM, *Ocote'a cymbarum* seu *ama'tra*, grows in Oronoko, where it is called *Sass'afra's*. The bark is aromatic, bitter, and stomachic; and Von Martius thinks it is an ingredient in the curare poison.

NECTANDRA PUCHURY, see Pichurim beans — *n.* Rodiei, see Bebeuru.

NECTAR, *νεκταρ*, from *νη*, 'a particle of negation,' and *κταω*, 'to kill.' A pleasant liquor, feigned by the poets to have been the drink of the gods, and to have rendered immortal those who partook of it. A name given, by the ancients, to many drinks; and particularly to one made with wine, evaporated, and sweetened with honey.

NECTARIUM, *Inula helenium*.

NECUSIA, see Wound.

NECYOMANTIA, Necromancy.

NEDYIA, Intestines.

NEDYS, *νῆδος*. The belly, abdomen, stomach, uterus.

NEEDLE, Sax. *neol*, *næol*, from Teut. *neten*, 'to sew.' *Acus*, *Bel'onē*, *Rophis*, *Raph'ion*, *Aces'tra*, (F.) *Aiguille*. A steel instrument, used in many professions. In *Surgery*, a steel, gold, silver, or platina instrument, that may be round, flat, or triangular, straight, or curved, supported or not by a handle, but having *always* a point, by

means of which it penetrates the textures; and often having, either near the point, or more commonly, near the other extremity, an aperture or eye for the reception of a thread or tape, which it introduces into the parts. The *Aiguille à appareil* of the French is the ordinary sewing needle used in the making of bandages, &c.

NEEDLE, ACUPUNCTURE, (F.) *Aiguille à Acupuncture*. An inflexible gold or silver needle; conical, very delicate, four inches long, furnished with a handle, and, at times, with a canula shorter than it by about half an inch. An ordinary needle, waxed at the head, will answer as a substitute for this. See *Acupuncture*.

NEEDLE, CATARACT, *Acus ophthalmica*, (F.) *Aiguille à cataracte*. Needles of gold, silver, and steel have been used; the latter, alone, at the present day. The cataract needle is employed to depress or tear the crystalline when opaque. This needle is usually made from 15 to 24 lines long; and is attached to a fine handle. The extremity may be, as in Scarpa's and Langenbeck's, pointed, prismatic, triangular, and curved; in Dupuytren's and Walther's, flat, curved, and sharp-edged; in Hey's, flat, with a semicircular and sharp end; or, as in Beer's, Siebold's, Schmidt's, Himly's, Von Gräfe's, &c., straight and spear-pointed. A mark is generally placed upon the handle to inform the operator,—when the instrument is engaged in the eye,—what side corresponds to the crystalline.

NEEDLE FOR A COUNTER-OPENING, *Acus invaginata*, (F.) *Aiguille à Contre-ouverture ou en gaine ou à gaine*,—a long, narrow instrument of steel; the point of which is fine and sharp on both sides;—the heel (*talon*) having an aperture to receive a thread, tape, &c., provided with a flat, silver sheath, shorter than the blade, the point of which it covers when passing through parts that have to be respected.

NEEDLE, DESCHAMP'S, *Paupé's needle*, (F.) *Aiguille de Deschamps, A. à manche*, &c., is the last described needle, fixed to a handle. The eye is placed near the point. It is employed in the ligation of deep-seated arteries.

NEEDLE, FIS'TULA, (F.) *Aiguille à Fistule*. A long, flat, flexible, silver instrument; having an aperture near one extremity; blunt at the other. This was formerly used for passing a seton into fistulous ulcers. On one of its sides was a groove for guiding a bistouri in case of necessity.

Also, a long, steel instrument, terminated by a point like that of a trocar, which Desault employed for penetrating the rectum, when operating for fistula that had no internal aperture.

NEEDLE, HARE-LIP, (F.) *Aiguille à Bec de Lièvre*. A small, silver canula, to which is attached a spear-point, that can be readily withdrawn. This needle, armed with the point, is introduced at one side of the fissure in the lip, and through the other. The twisted suture is then applied, and the pin withdrawn.

NEEDLE, LIG'ATURE, (F.) *Aiguille à ligation*. A long, steel instrument, sharp towards one extremity, with an eye near the other, which was formerly used for suspending the circulation of blood prior to amputation, by being passed through the limb, so as to include the principal artery and a part of the muscles and integuments.

Also, a steel instrument of various dimensions, round towards one of its extremities, which was straight, and furnished with an aperture; curved and flat towards the opposite, which was pointed, and had a slight ridge on its concave side. At the commencement of the last century, this instrument was used to pass ligatures around vessels.

Also, a steel instrument of various dimensions,

flat, regularly curved in the form of a semicircle, with a sharp or lance point, and a long eye, used with advantage in place of the last.

NEEDLE, PAUPÉ'S, *Needle, Deschamps*.

NEEDLE, SETON, (F.) *Aiguille à Séton*. A long, narrow, steel blade; pointed and sharp at one extremity; pierced at the other by an aperture. The *Aiguille à contre-ouverture* may be used for the same purpose.

NEEDLE, SUTURE, (F.) *Aiguille à Suture*. For the twisted suture, the hare-lip needle is used; for the others, the straight or curved needle: the straight needle is preferable for stitching up the abdomen, &c., in dissection. In the suture of the tendons, a curved needle has been used; first on both sides, and cutting only at the concave edge, in order that the instrument may pass between the tendinous fibres without dividing them.

NEEDLE-BEARER, *Porte-aiguille*.

NEEDLE-CARRIER, *Porte-aiguille*.

NEEM BARK, see *Melia azedarach*.

NEESING, Sneezing.

NÉFLE, *Mespilus* (the fruit).

NÉFLIER, *Mespilus*.

NEFREN'DES, properly, 'sucking pigs,'—Varro. *Nodoi, Eden'tuli* (vn, privative, and *odonos*, 'a tooth.') Persons devoid of teeth. Young children, for instance, who have not cut them; or aged persons, who have lost them. This state is called *Nefrendia, Nudo'sia, Odon'tia eden'tula*.

NEGOTIUM PARTURITIONIS, Parturition.

NÈGRE BLANC, Albino.

NEGRETIA PRURIENS, *Dolichos pruriens*.

NEGRO, *Nigrita*, (F.) *Nègre*. One of the Ethiopian race. See *Homo* and *Mulatto*.

NEGROMANTIA, Necromancy.

NEIERA, Abdomen.

NEIGE, Snow.

NEIR, Kidney.

NEIRA, Abdomen.

NEIS, Nasus.

NELUMB'BIUM LUTEUM, *Yellow nelumbo, Yellow water lily, Pond lily, Water shield, Water nuts, Water chin'capin, Rattle nut, Sacred bean*. A beautiful water plant, common in the United States, and belonging to *Nat. Ord. Nymphaeaceae*; *Sex. Syst. Polyandria Polygynia*. The leaves are cooling and emollient when applied to the surface. The roots, leaves, and nuts are eaten. The last are called by the Indians and others *water chincapins*.

NELUMBO, YELLOW, *Nelumbium luteum*.

NENDO, *Angelica lucida*.

NENNDORF, MINERAL WATERS OF. Nenndorf is a village three and a half German miles from Hanover. Its cold sulphureous spring is much celebrated.

NENUPHAR, *Nymphaea alba*—*n. Blanc*, *Nymphaea alba*—*n. Jaune*, *Nymphaea lutea*—*n. Lutea*, *Nymphaea lutea*—*n. Odorata*, *Nymphaea odorata*.

NEOARTHROSIS, from *neos*, 'new,' and *arthron*, 'a joint.' A new joint; an artificial joint.

NEOG'ALA, from *neos*, 'new,' and *gala*, 'milk.' Milk secreted immediately after the colostrum. Also, the colostrum.

NEOGENES, *Nouveau-né*.

NEOGILUS, *Nouveau-né*.

NEOGNUS, *Nouveau-né*.

NEOGONUS, *Nouveau-né*.

NEO-MEMBRANE, see *Membrane*, false.

NEONATUS, *Nouveau-né*.

NEOPLASTY, *Neoplas'ticē*, from *neos*, 'new,'

and πλασσω, πλασσω, 'I form.' An operative process for the formation of new parts. It includes autoplasty, cicatrization of wounds, and the formation of adhesions.—Burdach.

NEOSCYTECHUS, see *Craquement de Cuir neuf*.

NEOTOCICTERUS, Icterus infantum.

NEOTIA, Goodyera pubescens.

NEP, Nepeta.

NEPEN'THA DESTILLATO'RIA, *Bandu'ra*. A Ceylonese plant, the root of which is astringent.

NEPEN'THES, from *νη*, negative particle, and *πενθος*, 'grief.' A remedy much extolled by the ancients against sadness and melancholy.

The women of Thebes, according to Diodorus Siceliotus, alone possessed the secret of its composition; and, according to Homer, Helen introduced it from Egypt. Some suppose it to have been opium.

NEPENTHES, Bangué.—n. Opiatum, Pilulæ opiatæ.

NEP'ETA, *N. Cata'ria seu vulga'ris*, *Cata'ria vulga'ris*, *Herba felis*, *Nep*, *Catnep*, *Catmint*, (F.) *Herbe aux Chats*, *Cataire*,—so called, because cats are fond of it. *Ord.* Labiatae. The leaves, *Cata'ria*, (Ph. U. S.), have a smell and taste like those of an admixture of spearmint and pennyroyal. It has been recommended in uterine disorders, dyspepsia, flatulency, &c., like pennyroyal; and is much used in domestic medicine, on the American continent, in flatulencies, &c., of children.

NEPETA AGRESTIS, Melissa nepeta.—n. Glechoma, Glechoma hederacea.—n. Vulgaris, Nepeta.

NEPHALIOTES, Temperance.

NEPHELE, Enæorema, Nebula.

NEPHELION, Nebula.

NEPH'ELOID, *Nepheloides*, *Nepheloides*, *Neb'ulous*, *Nubilosus*. From *νεφελή*, 'a cloud,' and *εἶδος*, 'resemblance.' An epithet applied to urine when it is cloudy.—*Uri'na nepheloides*.

NEPHOS, Nebula.

NEPHRÆMORRHAGIA, see Hæmaturia.

NEPHRAL'GIA, *Dolor Nephreticus*, *Neural'gia Renum*, (F.) *Néuralgie des Reins*, *Dysnéphro-nervie*, from *νεφρός*, 'a kidney,' and *αλγος*, 'pain.' Pain and neuralgia in the kidney.

NEPHRALGIA ARENOSA, see Gravel.—n. Calculosa, see Gravel.—n. Rheumatica, Lumbago.

NEPHRANURIA, Icturia renalis.

NEPHRAPOSTASIS, *Nephropyosis*, *Abscess'ous renalis*, from *νεφρός*, 'kidney,' and *αποστασις*, 'abscess.' Renal abscess.

NEPHRATON'IA, from *νεφρός*, 'kidney,' and *ατωνα*, 'want of tone,' *Renum aton'ia seu paral'ysis*. Atony of the kidney.

NEPHRAUX'E, from *νεφρός*, 'a kidney,' and *αὖξω*, 'to increase.' Enlargement of the kidney.

NEPHRELCO'SIS, *Nephropyosis*, *Helco'sis renalis*, from *νεφρός*, 'kidney,' and *ἐλκωσις*, 'ulceration.' Ulceration of the kidney.

NEPHRELMIN'TIC, *Nephrelminticus*, from *νεφρός*, 'a kidney,' and *ελμινς*, *ελμινθος*, 'a worm.' That which is owing to the presence of worms in the kidney.

NEPHREMPHRAX'IS, from *νεφρός*, 'a kidney,' and *εμφρασσω*, 'I obstruct.' A name given by Plouquet to obstruction of the kidneys.

NEPHRET'IC, *Nephrid'ius*, *Nephrit'ic*, *Nephrit'iculus*, from *νεφρός*, 'a kidney.' That which relates to the kidney. Applied, especially, to pain, &c., seated in the kidney.

NEPHRETICUM, in *Materia Medica*, means

a medicine employed for the cure of diseases of the kidney.

NEPHRETICUM LIGNUM, *Guilandina moringa*.

NEPHRET'ICUS LAPIS, *Talcum nephrit'icum*, (F.) *Pierre néphrétique*, *Jade néphrite ou oriental*. A green, fatty kind of stone,—once used as an amulet against epilepsy,—an absurd name, as there can be no such remedy.

NEPHRIA, Kidney, Bright's disease of the.

NEPHRID'ION, *Pingue'do renalis*. From *νεφρός*, 'the kidney.' The fat which surrounds the kidneys.—Hippocrates.

NEPHRIDIDIUM, Capsule, renal.

NEPHRIDIDIUS, Nephretic.

NEPHRINE, Oxyd, cystic, Urea.

NÉPHRITE, Nephritis.—n. *Albumineuse*, Kidney, Bright's disease of the.

NEPHRITES, Asphalitics.

NEPHRITIC, Nephretic.

NEPHRITICUM LIGNUM, *Guilandina moringa*.

NEPHRITIS, Asphalitics.

NEPHRIT'IS, *Empres'na Nephritis*, from *νεφρός*, 'kidney,' and *itis*, 'denoting inflammation,' *Renum inflammatio*, *Nephro-phlegmoné*,—*Inflammation of the Kidney*, (F.) *Néphrite*, *Inflammation des Reins*, is characterized by acute pain; burning heat, and a sensation of weight in the region of one or both kidneys; suppression or diminution of urine; fever; dysuria; ischuria; constipation, more or less obstinate; retraction of the testicle, and numbness of the thigh of the same side. It may be distinguished into *simple* and *calculous nephritis*—*Lithonephritis*. In the latter, the urine often contains small particles of uric acid or of urate of ammonia. The most common causes of nephritis are,—excess in irritating and alcoholic drinks; abuse of diuretics; blows or falls on the region of the kidneys; the presence of renal calculi, &c. It may be distinguished from lumbago by the pain which attends the latter on the slightest motion, &c. It usually terminates by resolution in from one week to two or three. It may, however, end in suppuration—*pyonephritis*; or may become chronic—*chrononephritis*, (Piorry.) In the treatment, antiphlogistics, as bleeding, baths, &c., are required to the full extent; with the use of diluents, opiates, &c.

NEPHRITIS ALBUMINENSIS, Kidney, Bright's disease of the.—n. *Albuminosa*, Kidney, Bright's disease of the.—n. *Associated*, Kidney, Bright's disease of the.—n. *Cachectic*, Kidney, Bright's disease of the.—n. *Calculous*, see Nephritis.—n. *Sociata*, Kidney, Bright's disease of the.

NEPHROCHALAZOSIS, Kidney, Bright's disease of the.

NEPHROCOCOSIS, Kidney, Bright's disease of the.

NEPHROCYSTO'SIS, from *νεφρός*, 'a kidney,' and *κυστις*, 'a cyst or bladder.' A morbid condition, which consists in the formation of cysts in the bladder.

NEPHROCOLICA, Colica nephretica.

NEPHRODES, Nephroid.

NEPHRODIUM FILIX MAS, *Polypodium filix mas*.

NEPHROG'RAPHY, from *νεφρός*, 'a kidney,' and *γραφη*, 'a description.' An anatomical description of the kidney.

NEPHROHE'MIA, from *νεφρός*, 'kidney,' and *αἷμα*, 'blood.' Hyperæmia or congestion of the kidney.

NEPH'ROID, *Nephroides*, *Nephroides*, *Reniform'is*, from *νεφρός*, 'kidney,' and *εἶδος*, 'form,'

resemblance.' Reniform. Having a resemblance to a kidney—as '*nephroid* cancer,' so called because the morbid growth resembles the kidney in structure.

NEPHROLITHIASIS, from νεφρος, 'a kidney,' and λίθος, 'a stone.' The disease of calculus in the kidney; *Lithi'a renalis*, *Lithi'asis nephritica*, *L. Renalis*. See Gravel.

NEPHROLITHIC, same etymon. Belonging to calculi in the kidneys. This epithet has been applied to ischuria, occasioned by calculi formed in the kidneys.

NEPHRO-LITHOTOMY, Nephrotomy.

NEPHROLOGY, *Nephrolog'ia*, from νεφρος, 'kidney,' and λογος, 'a treatise.' A treatise on the kidneys. Dissertation on the kidneys and their functions.

NEPHROMALACIA, from νεφρος, 'kidney,' and μαλακία, 'softness.' Softening of the kidney.

NEPHROMETRÆ, Psom.

NEPHRONCUS, *Tumor renalis*, from νεφρος, 'kidney,' and ογκος, 'a tumour.' A tumefaction of the kidney.

NEPHROPHLEGMATIC, *Nephro-phlegmaticus*, from νεφρος, 'kidney,' and φλεγμα, φλεγματος, 'phlegm.' A name given, by some authors, to ischuria produced by mucus contained in the urine.

NEPHROPHLEGMONE, Nephritis.

NEPHROPHTHISIS, from νεφρος, 'kidney,' and φθισις, 'consumption.' Phthisis from suppuration of the kidney.

NEPHROPLETHORIC, *Nephro-plethoricus*, from νεφρος, 'a kidney,' and πληθωρα, 'plethora.' Belonging to plethora of the kidneys. An epithet given to ischuria dependent upon this cause.

NEPHROPYIC, *Nephro-py'icus*, from νεφρος, 'a kidney,' and πυρ, 'pus.' Belonging to supuration of the kidney.

NEPHROPYOSIS, Nephropostasis, Nephrelcosis.

NEPHRORRHAGIA, *Hæmatu'ria renalis*, *Hæmorrhag'ia renum*, from νεφρος, 'kidney,' and ραγη, 'rupture;' *Proflu'vium sanguinis renibus*. Hemorrhage from the kidney.

NEPHROS, Kidney.

NEPHROSCLERIA, from νεφρος, 'kidney,' and σκληρία, 'hardness;' *Induratio renum*. Induration of the kidneys.

NEPHROSPASTIC, *Nephrospas'ticus*, from νεφρος, 'a kidney,' and σπασω, 'I draw.' That which depends upon spasm of the kidney. An epithet given to a variety of ischuria.

NEPHROTHROMBOID, *Nephro-thromboïdes*, from νεφρος, 'a kidney,' and θρομβος, 'a clot.' That which depends upon clots of blood, contained in the kidneys or their ducts. An epithet for a species of ischuria.

NEPHROTOMY, *Nephro-lithot'omy*, *Nephrotom'ia*, *Nephro-lithotom'ia*, *Sectio renalis*, *S. renis*, from νεφρος, 'a kidney,' and τεμνω, 'to cut.' Dissection of the kidney. Also, an operation proposed with the view of extracting calculi formed in the kidney, by means of an incision into the tissue of that organ.

NEPHRUS, Kidney.

NEPIOS, Infans.

NEPIOTES, Infancy.

NEPTA, Asphaltum.

NERA, Toddy.

NERANTIA, see *Citrus aurantium*.

NERE, Kidney.

NERF, Nerve.—*n. Circumflexe*, Axillary nerve —*n. Dentaire*, Dental nerve —*n. Fessier*, Gluteal nerve —*n. Glosso-pharyngien*, Pharyngo-glossal nerve —*n. Gustatif*, see Lingual nerve —*n. Gustatif innominé*, Trigemini —*n. Gutturel*, Palatine (middle) nerve —*n. Gutturel-palatin*, Palatine nerve —*n. Honteux*, Pudic nerve —*n. Irien*, Ciliiary nerve —*n. Ischio-clitorien*, Pudic nerve —*n. Ischio-pénien*, Pudic nerve —*n. Labyrinthique*, Auditory nerve —*n. Mentonier*, Mental nerve —*n. Moteur oculaire externe*, Motor oculi externus —*n. Oculo-musculaire externe*, Motor oculi externus —*n. Oculo-musculaire interne*, Patheticus nervus —*n. Pathétique*, Patheticus nervus —*n. Péronière*, branch, Popliteal nerve, external —*n. Pharyngo-glossien*, Pharyngo-glossal nerve —*n. Pneumogastrique*, Pneumogastria nerve —*n. Première paire trachélienne*, Occipital nerve —*n. Prétipto-digital*, Musculo-cutaneous nerve, see Popliteal nerve —*n. Prétipto-auphalangéaire*, see Popliteal nerve —*n. Prétipto susplantaire*, Tibial nerve, anterior —*n. Radio-digital*, Radial nerve —*n. Sciatique grand*, Sciatic nerve, great —*n. Sciatique petit*, Sciatic nerve, lesser —*n. Sciatique poplitée externe*, Musculo-cutaneous nerve —*n. Sous-occipital*, Occipital nerve —*n. Sous-pubio-fémoral*, Obturator nerve —*n. Spino-cranio-trapézien*, Spinal nerve —*n. Susmaxillaire*, Maxillary superior nerve —*n. Suspubein*, Supra-pubian nerve —*n. Sympathique*, Trisplanchnic nerve —*n. grand Sympathique*, Trisplanchnic nerve —*n. Tibiale*, branch, Popliteal internal —*n. Trachéal*, Laryngeal inferior nerve —*n. Trachélo-dorsal*, Spinal nerve —*n. Tri-jumeux*, Trigemini —*n. à Trois cordes*, Trigemini —*n. Tympanique*, Chorda tympani —*n. Vague*, Pneumogastria nerve —*n. Vertébro-digital*, Musculo-cutaneous nerve.

NERFS BRONCHIQUES, Bronchial nerves —*n. Ciliaires*, Ciliary nerves —*n. Craniens*, Cerebral nerves —*n. Moteurs oculaires communs*, Motores oculorum —*n. Rachidiens*, Vertebral nerves —*n. Sacrés*, Sacral nerves —*n. Sous-costales*, Intercoastal nerves.

NÉRION, Nerium oleander.

NÉRIS, MINERAL WATERS OF. Nérís is on the high road from Moulins to Limoges, eighty leagues from Paris. There are four springs, the water of which is clear, inodorous, tasteless, and has an unctuous feel. It contains carbonic acid gas, carbonate of soda, and sulphate of soda, chloride of sodium, silex, and an animal matter to which its 'oleaginous' property is probably owing. The waters are generally used in the form of thermal baths.

NE'RIMUM, from νηρος, 'humid,' because growing in moist places. *N. oleander*.

NE'RIMUM ANTIDYSENTERICUM, *Wright'ia antidysenterica*. The tree which affords the *Codagapala Bark*, *Cones'si cortex*, *Tillicher'ry cortex*, *Cortex proflu'vii*. Family, Apocynæe. *Nar. Syst.* Pentandria Monogynia. (F.) *Codagapale*, *Cropale*. The bark of this Malabar tree is of a black colour, externally, and is generally covered with a white moss or scurf. It has an anstere, bitter taste, and has been recommended as an astringent in diarrhoea, dysentery, &c.

NE'RIMUM OLEAN'DER, *Ne'rium*, *Rhododaph'nia*, *Rosa'go*, (F.) *Laurier rose*, *Laurrose*, *Nérion*, *Rosage*, *Rhododaphné*. The leaves are reputed to be narcotic. The infusion is employed internally, in herpetic affections, and the powder, incorporated with lard, is used in the itch.

NERONIA'NA seu **NERO'NICA** (PHLEBOTOM'IA). An epithet given, for some cause, to phlebotomy, when more than one vein was opened

on the same day, or to bleeding *coup sur coup*. Termed *Neroniana*, because the philosopher Seneca was bled to death under Nero [?].

NERPRUN PURGATIF, Rhannus.

NERVE, *Nervus*, *Neuron*, *νευρον*, 'a string.' (F.) *Nerf*. *Neuron* and *Nervus* meant also, with the ancients, a tendon and a ligament, *Partes nervosæ*; and hence the different acceptations of 'nervous':—'a man of nerve—a strong, nervous man; and a weak, nervous woman.' The nerves are tubular cords of the same substance as that which composes the encephalon and spinal marrow. They extend from one or other of the nervous centres to every part of the body, communicating, frequently, with each other; forming *plexuses*, and, occasionally, *ganglions*; and being, at length, lost in the parenchyma of organs. There are 42 pairs, and, according to their origin, they are termed *Cranial* or *Encephalic*, and *Spinal*. Each nerve is composed of several filaments or cords placed alongside each other, and is surrounded by a neurilemma. The encephalic nerves, in general, have only one root in the brain, whilst the spinal arise from the marrow by two roots; the one from an anterior fasciculus of filaments, the other from a posterior, separated from each other by the *Ligamentum denticulatum*; uniting outside this ligament, and presenting, near the intervertebral foramen, a ganglion formed only by the posterior root. The two roots make,

afterwards, but one nerve; and, like the encephalic nerves, proceed to their destination, subdividing into rami and ramusculi, until they are finally lost in the texture of the organs. The trunks first formed are commonly round, and proceed alone, or accompany the great vessels, being placed in the areolar spaces which separate the organs, and are thus protected from injury. Their manner of termination we are not acquainted with; whether the nervous pulp, for instance, as is more probably the fact, be distributed or lost in a membrane, as seems to be the case with the nerves of sight, hearing, and smell,—or are looped. Certain it is, that there is considerable difference in the organs, as respects the quantity of nerves that terminate in them; and the particular arrangement of the nervous extremities. Some organs have numerous nerves; others seem to have none: a circumstance which influences considerably the sensibility of parts.

The *Encephalic Nerves* arise from the encephalon, or are inserted into it; (according as we consider the brain the origin or termination of the nerves;) and make their exit by foramina at the base of the skull. They are 12 in number. The spinal nerves are 31 in number, 8 *cervical*, 12 *dorsal*, 5 *lumbar*, and 6 *sacral*: the four inferior cervical being much larger than the superior, because they furnish the nerves of the upper extremities.

SYNOPTICAL TABLE OF THE NERVES.

I. Cranial or Encephalic Nerves.		
1. <i>Olfactory</i> (1st pair.)	Divided into internal, external, and middle branches, which are distributed on the Schneiderian membrane.	6. <i>Abducentes</i> (6th pair.)
2. <i>Optic</i> (2d pair.)	Terminate in the retina.	To the rectus externus oculi.
3. <i>Motores Oculorum</i> . (3d pair.)	<i>Superior Branch.</i> To the rectus superior oculi and levator palpebræ superioris. <i>Inferior Branch.</i> To the rectus internus, rectus inferior and lesser oblique muscles; a filament which goes to the ophthalmic ganglion.	1. At its exit from the cranium, the <i>posterior auricular</i> , <i>stylohyoid</i> , and <i>infra-mastoid</i> , to the pavilion of the ear, the mastoid process, the digastricus, and the muscles attached to the styloid process.
4. <i>Pathetici</i> (4th pair.)	To the greater oblique muscle of the eye.	2. Near the parotid gland, the <i>temporal</i> , <i>malar</i> , <i>buccal</i> , <i>supra-maxillary</i> , and <i>infra-maxillary</i> to the whole superficies of the face.
	<i>Ophthalmic Branch.</i> Divided into three branches. 1. The <i>lacrimal</i> branch, to the lacrimal gland and upper eyelid. 2. <i>Frontal branch</i> , to the forehead and upper eyelid. 3. <i>Nasal branch</i> , to the eyelids, nasal fosse, and nose.	7. <i>Facial</i> (Portio dura of the 7th pair.)
	<i>Superior Maxillary Branch.</i> 1. The <i>orbital branch</i> , to the orbit. 2. The <i>posterior and superior dental</i> , to the last three molar teeth and gums. 3. The <i>anterior dental</i> , to the incisor, canine, and two lesser molars. 4. <i>Infra-orbital</i> , to the upper lip, cheek, and nose.	8. <i>Auditory</i> (Portio mollis of the 7th pair.)
	<i>Inferior Maxillary Branch.</i> 1. <i>Temporal</i> profound branches, to the temporal muscle. 2. <i>Masseterine</i> branch, to the masseter muscle. 3. <i>Buccal</i> , to the inner surface of the cheek. 4. <i>Pterygoid</i> , to the internal pterygoid muscle. 5. <i>Lingual</i> , to the mucous membrane of the tongue. 6. <i>Inferior dental</i> , to the teeth of the lower jaw, and to the lower lip. 7. <i>Auricular branch</i> , to the pavilion of the ear and forehead.	9. <i>Glosso-pharyngeal</i> . (Portion of the 8th pair.)
5. <i>1st Trigeminum</i> . (5th pair.)		To the vestibule, semi-circular canals, and cochlea.
		To the base of the tongue and pharynx.
		1. <i>IN THE NECK</i> ,—a <i>pharyngeal</i> branch to the pharynx; <i>superior laryngeal</i> branch to the larynx, and to some muscles of the inferior hyoid region. <i>Cardiac</i> branches to the cardiac plexus.
		2. <i>IN THE CHEST</i> ,—the <i>inferior laryngeal</i> branch to the larynx— <i>pulmonary</i> branches, which form the plexus of the same name: <i>oesophageal</i> branches to the oesophagus.
		3. <i>IN THE ABDOMEN</i> ,— <i>gastric</i> branches to the plexures of the stomach: <i>filaments</i> which go to the neighbouring plexures.
		At its exit from the cranium—the <i>accessory of the pneumo-gastric nerve</i> anastomosing with this nerve. In the neck—filaments to the trapezius muscle.
		<i>Cervicæ descendens</i> , to the muscles of the inferior hyoid region, and to the cervical nerves. Filaments to the muscles of the tongue.
		10. <i>Par Vagus</i> (8th pair.)
		11. <i>Spinal</i> or <i>Accessory of Willis</i>
		12. <i>Hypoglossal</i> (9th pair.)

II. Spinal or Vertebral Nerves.

1. Cervical Nerves.

- 1st Cervical Pair..... { Anterior Branch. Anastomosing *par arcade* with the second pair.
- 2d Cervical Pair..... { Anterior Branch. A branch anastomosing with the first pair: a branch which goes to the cervical plexus
- 3d and 4th Cervical Pairs..... { Anterior Branch. Concurring in the cervical plexus.
- The Posterior Branch of these four pairs ramifies on the occiput and muscles of the superficial and deep-seated cervico-occipital regions.

DESCENDING BRANCHES.—1. *Internal descending*, anastomosing with a branch of the hypoglossus. 2. *Plexenic branch*, to the diaphragm. 3. *External descending branches*, dividing into supra-clavicular, supra-acromial, infra-clavicular, and cervicales profundi, to the muscles and integuments of the upper part of the chest and shoulder, to the trapezius, levator scapulae, rhomboid-eus, &c.

Cervical Plexus.....

ASCENDING BRANCHES.—1. *Mastoid*, to the posterior and lateral part of the head and the inner surface of the pavilion of the ear. 2. *Auricular*, to the parotid gland and pavilion of the ear.

Superficial Cervical Branches. (2.) To the platysma myoides, digastricus, the integuments of the neck, &c.

Anterior Branches. Assisting in the formation of the brachial plexus.

5th, 6th, and 7th Cervical Pairs.....

Posterior Branches. To the muscles and integuments of the posterior part of the neck and upper part of the back.

1. *Thoracic Branches*, (2.) to the anterior and lateral parts of the chest. 2. *Supra-scapular* branch to the muscles of the posterior scapular region. 3. *Infra-scapular* branches (3.) to the subscapularis, teres major, teres minor, and latissimus dorsi. 4. The *brachial interna cutaneous*, to the integuments of the palmar and dorsal surfaces of the forearm, near the ulnar margin. 5. The *brachial external cutaneous*, principally to the integuments of the palmar and dorsal surfaces of the forearm, near the radial margin. 6. The *median nerve*, to the forearm, hand, and palmar surface of all the fingers. 7. The *cubital nerve*, to the last two fingers. 8. The *radial*, to the first three. 9. The *axillary*, around the shoulder-joint, and to the neighbouring muscles.

Brachial Plexus.....

2. Dorsal Nerves, (Nervi dorsales.)

- 1st Dorsal Pair..... { Anterior Branch. To the brachial plexus.
- 2d and 3d Dorsal Pairs..... { Anterior Branches. An intercostal and brachial branch.
- 4th, 5th, 6th, and 7th Dorsal Pairs..... { Anterior Branches. Internal branches, to the intercostals, triangularis sterni, pectoralis major, and to the skin. External branches to the

4th, 5th, 6th, and 7th Dorsal Pairs.....

8th, 9th, 10th, and 11th Dorsal Pairs.....

12th Dorsal Pair.....

integuments of the chest, the obliquus externus abdominis, and the skin of the abdomen.

Anterior Branches. Internal branches, to the transversalis, obliquus internus, and rectus muscles, and to the skin of the abdomen. External branches, to the integuments of the chest, and to the muscles and skin of the abdomen.

Anterior Branch. To the first lumbar nerve, and to the muscles and skin of the abdomen as far as the iliac crest.

The Posterior Branches of the dorsal nerves are distributed to the muscles and integuments of the back and loins.

3. Lumbar Nerves.

- 1st, 2d, 3d, and 4th Lumbar Pairs..... { Anterior Branches. Concurring to form the lumbar plexus.

5th Pair..... { Anterior Branch. Aiding in forming the sciatic plexus.

The Posterior Branches of the lumbar nerves are distributed to the loins, sacrum, and nates.

Lumbar Plexus.....

1. *Musculo-cutaneous Branches*, to the number of three. One superior, to the muscles of the abdomen, to the fold of the groin, and the scrotum: a middle, to the integuments and muscles of the abdomen: and an inferior, to the skin of the thigh. 2. A *genito-crural* branch to the integuments of the scrotum, the groin, and the thigh. 3. The *crural* nerve, to the integuments and muscles of the thigh, skin of the leg, and foot. 4. The *obturator* nerve, to the muscles at the inner part of the thigh. 5. The *lumbo-sacral*, to the sciatic plexus. It gives off the *gluteal* nerve to the glutei muscles.

4. Sacral Nerves.

- 1st, 2d, 3d, and 4th Sacral Pairs..... { Anterior Branches. They form by their union the sciatic plexus.

5th and 6th Pairs..... { Anterior Branches. To the parts in the vicinity of the coccyx.

The Posterior Branches ramify on the muscles and integuments of the nates.

Sciatic Plexus.....

1. *Hemorrhoidal nerves*, to the rectum. 2. *Vesical*, to the bladder. 3. *Uterine and vaginal*, to the vagina and uterus. 4. *Inferior gluteal*, to the gluteal muscles, perinaeum, and integuments of the posterior part of the thigh. 5. *Pudic*, to the perinaeum, penis, or vulva. 6. *Sciatic*, divided into the *external popliteal*, which ramifies on the integuments and muscles of the external side of the leg, on the dorsum of the foot, and the dorsal surface of the toes; and into the *internal popliteal*, distributed on the dorsal surface of the two last toes, to the muscles of the foot, and on the plantar surface of all the toes.

Classifications of the nerves have been recommended according to their uses, in preference to the ordinary anatomical arrangement. It has been remarked that the encephalic nerves have generally one root; the spinal two. Experiments and pathological facts have proved, that the anterior column of the marrow and the anterior roots of the spinal nerves are inservient to volition or voluntary motion: and that the posterior column and roots are destined for sensibility. Hence the spinal nerves, which have two roots, must be the conductors both of motion and feeling; whilst the encephalic, which, with but few exceptions, have but one, can possess but one of these properties:—they must be either *sensitive* or *motor*, according as they arise from the posterior or anterior column of the medulla: and, consequently, three classes of nerves may be distinguished.

- | | |
|-----------------------------------|---|
| 1. <i>Sensory</i> | { Arising, by a single root, from the posterior column of the medulla oblongata or spinal marrow. |
| 2. <i>Motor</i> | { Arising, by a single root, from the anterior column of the same parts. |
| 3. <i>Sensory and Motor</i> | { Which have two roots: one from the anterior, and one from the posterior column. |

According to Sir Charles Bell, the medulla oblongata is composed of three columns on each side; an *anterior*, a *middle*, and a *posterior*. Whilst the anterior and posterior columns produce the nerves of motion and sensation respectively, the middle, according to Sir Charles, gives rise to a third set of nerves—the *respiratory*. To this order belong:—

1. The accessory nerve of Willis, or *superior respiratory*.
2. The *par vagum*.
3. The *glosso-pharyngeal*.
4. The *facial* or *respiratory of the face*.
5. The *phrenic*.
6. A nerve which has the same origin as the phrenic;—the *external respiratory*.

When a horse has been hard-riden, every one of these nerves is in action.

This division is now, however, generally abandoned, and there does not seem to be a third column, especially destined for respiration.

Sir C. Bell, again, reduced the system of nerves to two great classes. 1. Those that are *regular*, *primitive*, *symmetrical*, and common to all animals, from the worm to man; which have double roots, and preside over sensibility and motion: and, 2. The *irregular* or *superadded*, which are added to the preceding, in proportion as the organization of animals offers new or more complicated organs. To the first class belong all

the spinal nerves and at least one encephalic—the 5th pair;—to the second, the rest of the nervous system.

Dr. Marshall Hall has proposed a division of the nervous system, which is calculated to explain many of the anomalous circumstances so frequently witnessed. He proposes to divide all the nerves into, 1. The *cerebral* or the sentient and voluntary. 2. The *true spinal* or excitomotor. 3. The *ganglionic* or *cyclo-ganglionic*,—the nutrient and secretory. If the sentient and voluntary functions be destroyed by a blow upon the head, the sphincter muscles will still contract when irritated, because the irritation is conveyed to the spine, and the reflex action takes place to the muscle so as to throw it into contraction. But if the spinal marrow be now destroyed, the sphincters remain entirely motionless, because the centre of the system is destroyed. Dr. Hall thinks that a peculiar set of nerves constitutes, with the vesicular centre of the marrow as their axis, the second subdivision of the nervous system; and as those of the first subdivision are distinguished into sentient and voluntary, these may be distinguished into the *excitor* and *motory*. The *first*, or the excitator nerves, pursue their course principally from internal surfaces, characterized by peculiar excitabilities, to the centre of the medulla oblongata and m. spinalis; the *second*, or the motor nerves, pursue a reflex course from the medulla to the muscles, having peculiar actions concerned principally in ingestion and egestion. The motions connected with the first or cerebral subdivision are sometimes, indeed frequently, *spontaneous*; those connected with the true spinal are, he believes, *always excited*. Dr. Hall thinks, too, that there is good reason for viewing the fifth, and posterior spinal nerves as constituting an external ganglionic system for the nutrition of the external organs; and he proposes to divide the *ganglionic* subdivision of the nervous system into, 1. The *internal ganglionic*, which includes that usually denominated the sympathetic, and probably filaments of the pneumogastric; and, 2. The *external ganglionic*, embracing the fifth and posterior spinal nerves. To the *cerebral* system he assigns all diseases of sensation, perception, judgment, and volition—therefore all painful, mental, and comatose, and some paralytic diseases. To the *true spinal*, *excito-motory*, *reflex*, or *diastaltic nervous system*, belong all spasmodic and certain paralytic diseases. He properly adds, that these two parts of the nervous system influence each other both in health and disease, as they both influence the ganglionic system.

The main views of Dr. Hall on the excitomotor function have been generally embraced.

The following tabular view of the arrangement and connexions of the nerves and nervous centres has been given by Dr. Carpenter. It affords a good general view, although some of the details might admit of modification.

TABULAR VIEW OF THE NERVOUS CENTRES.

Afferent fibres derived from Sensory Ganglia; efferent fibres transmitted to motor centres.	CEREBRAL GANGLIA, the seat of the formation of Ideas, and the instrument of the Reasoning processes and Will; participating also with the Sensory Ganglia in the formation of the Emotions; and thus the original source of <i>Voluntary</i> and <i>Emotional</i> movements.	Afferent fibres derived from Sensory Ganglia; efferent fibres transmitted to motor centres.
Afferent fibres derived from posterior column of spinal cord; efferent fibres transmitted into posterior column.	CEREBELLIC GANGLIA, for harmonization of muscular actions; including also the ganglionic centre of the sexual sense (?).	Afferent fibres derived from posterior column of spinal cord; efferent fibres transmitted into motor column.
Radiating fibres to Cerebral Ganglia:—Nerves of Common and Special Sensation:—Motor nerves forming part of general motor system.	CRANIO-SPINAL AXIS, or centre of Automatic actions; including— SENSORY GANGLIA, the seat of Sensation, and centre of <i>Consensual</i> (or Instinctive) movements, or of Automatic actions involving sensation.	Radiating fibres to Cerebral Ganglia:—Nerves of Common and Special Sensation:—Motor nerves forming part of general motor system.
Afferent and motor nerves of Respiration, Deglutition, &c.	RESPIRATORY and STOMATO-GASTRIC GANGLIA, forming the true centres of the <i>Medulla Oblongata</i> ; instruments of <i>Reflex</i> movements or automatic actions independent of sensation.	Afferent and motor nerves of Respiration, Deglutition, &c.
Afferent and motor fibres, forming Trunks of Spinal Nerves.	SPINAL GANGLION, or <i>True Spinal Cord</i> , consisting of a coalesced series of segmental ganglia, the instruments of <i>Reflex</i> operations, or Automatic actions independent of Sensation.	Afferent and motor fibres, forming Trunks of Spinal Nerves.

The nerves are sheathed, and are united to the neighbouring parts, by an abundant layer of fatty areolar texture, which sends, inwards, prolongations that separate the nervous cords and filaments from each other. The arterial trunks, distributed to them, transmit branches into their interior. The veins follow the same course as the arteries. Absorbents are not easily traced even on the greatest trunks.

NERVE, ACCESSORY, OF WILLIS, Spinal nerve—n. Articular, Axillary nerve—n. Buccinator, Buccal nerve.

NERVE CELLS, see Neurine—n. of Cotunnus, Nasopalatine nerve.

NERVE, FEMORAL CUTANEOUS, see Genito-crural nerve—n. Femoro-popliteal, lesser, Sciatic nerve, lesser.

NERVE FIBRES, *Fila ner'væ*, *Fibræ ner'væ*. In the nerves of the higher animals, two forms of nerve fibre exist—the *tubular*, and the *gelatinous*. In the minute fibrils or tubules, *Tubuli ner'væ*, full of nervous matter, which constitute the former, there is a difference between the central and peripheral portion:—the former has been called the *axis cylinder* and *primitive band*; the latter, the *medullary* or *white substance* of Schwann,—being that to which the peculiarly white aspect of cerebrospinal nerves is principally due. In the gelatinous fibre, no tubular envelope can be distinguished, and the white substance of Schwann seems to be wanting. It contains numerous cell-nuclei. See Neurine.

NERVE, GENITAL, see Genito-crural nerve—n. Genito-crural, see Genito-crural nerve—n. Gluteal of the Sacral plexus, Sciatic nerve, lesser—n. Gustatory, Hypoglossal nerve—n. Inguinal, internal, Genito-crural nerve—n. of Jacobson, see Petrosal ganglion—n. Lateral nasal, Sphenopalatine nerve—n. Lingual, Hypoglossal nerve—n. Naso-ocularis, Nasal nerve—n. Naso-palpebral, Nasal nerve—n. Orbito-frontal, Ophthalmic nerve—n. Ocular, Optic nerve—n. of Organic life, Trisplanchnic nerve—n. Palpebro-frontal, Frontal nerve—n. Popliteal internal, Tibial nerve, posterior.

NERVE POWER, Nervous power.

NERVE, RESPIRATORY, OF THE FACE, Facial nerve—n. Respiratory, superior, Spinal nerve—n. Scapulo-humeral, Axillary nerve—n. Scrotal, see Genito-crural nerve—n. Spinal, Radial nerve—n. Spinal, muscular, Radial nerve—n. Sub-occipital, Occipital nerve—n. Sub-pubic, Genito-crural nerve—n. Superficial temporal, Auricular nerve—n. Sympathetic, Trisplanchnic nerve—n. Temporal cutaneous, Auricular nerve—n. Ulnar, Cubital nerve—n. Vidian, Pterygoid nerve—n. of Wisberg, see Cutaneous nerves—n. Zygomatico-auricular, Auricular nerve.

NERVES, COSTAL, Intercostal nerves—n. Dorsal, Intercostal nerves—n. Eighth pair of, Pneumogastric—n. Encephalic, first pair, Olfactory nerves—n. Encephalic, eleventh pair, Spinal nerves—n. Ethmoidal, Olfactory nerves—n. Fifth pair, Trigemini—n. Fourth pair of, Patheticus ner-

vus—n. Infra-orbital, Sub-orbital nerves—n. Ninth pair of, Hypoglossal nerve—n. Oculo-muscular, common, Motores oculorum—n. Phrenic, Diaphragmatic nerves—n. Recurrent, Laryngeal inferior nerves—n. Sixth pair of, Motor oculi externus—n. Spinal, Vertebral nerves—n. Tenth pair of, Pneumogastric—n. Third pair of, Motores oculorum. NERVEUX. Nervous.

NERVI ACROMIALES, Acromial nerves—n. Anonymi, Innominati nervi—n. Claviculares, Clavicular nerves—n. Cutanei clunium inferiores, see Sciatic nerve, lesser—n. Divisi, Trigemini—n. Entobænonotes, Afferent nerves—n. Gustatorii, Trigemini.

NERVI MOLLES. 'Soft nerves.' The anterior branches of the superior cervical ganglion of the great sympathetic, which accompany the carotid artery and its branches, around which they form intricate plexuses, and, here and there, small ganglia. They are called *molles* from their softness, and *subrufi*, from their reddish hue.

NERVI ODORatorii, Olfactory nerves—n. Reversivi, Laryngeal inferior nerves—n. Subrufi, Nervi molles—n. Vulneratio, Neurotroxis.

NERVIMOTION, *Nervimotio*, from *nervus*, 'nerve,' and *motio*, 'motion.' A name given, by M. Dutrochet, to the movement which is caused in the organs of the senses by external agents, and transmitted to the muscles by the nerves.

NERVIMOTOR, (F.) *Nervimoteur*. An epithet given, by M. Dutrochet, to external agents, which are capable of causing nervimotion.

NERVINE, *Nervinus*, *Neurotic*, *Neurit'ic*, *Antineurotic*, *Antineuropath'ic*, (F.) *Névrolique*, from *nervus*, 'a nerve.' A medicine which acts on the nervous system.

NERVOUS, *Nervosus*, *Neurodes*, *Nerveus*, *Neuricus*, (F.) *Nerveux*. Same etymon. Relating or belonging to the nerves. Strong. Also, weak, irritable.

NERVOUS ATTACK, (F.) *Attaque des nerfs*, is an affection accompanied with spasm, pain, and different nervous symptoms, to which impressible individuals are liable.

NERVOUS CENTRES, see Centres, nervous.

NERVOUS DIATHESIS is termed *Atax'ia spirituum*, *Anomalia nervorum*, *Status nervosus* seu *erethiticus*, *Diathesis* seu *Constitutio nervosa*, *Nervousness*. Medically, *nervousness* means unusual impressibility of the nervous system. Formerly, it signified strength, force, and vigour. Recently, it has been applied to a hypochondriacal condition verging upon insanity, occasionally occurring in those in whom the brain has been unduly tasked;—a condition termed by some *cerebrop'athy*; by others, *brain-fag*.

NERVOUS DISEASES, *Morbi nervosi*, (F.) *Maladies nerveuses*. Affections seated in the nervous system. To purely functional disease of the nerves, Dr. Laycock has given the name *neuræmia*, and to the class of diseases he applies the epithet *neuræmic*.

NERVOUS FLUID, *Nervous principle*, *Fluidum* seu *Liquidum nervinum*, *Sucus nervus*, *Spiritus vitalis*. The fluid which is supposed to circulate through the nerves, and which has been regarded as the agent of sensation and motion. Of this fluid we know nothing, except that it resembles, in many respects, the electric or galvanic. It was formerly called *Animal spirits*, *Spiritus animal'es*, (F.) *Esprits animaux*.

NERVOUS PAIN, Neuralgia.

NERVOUS POWER, *Nerve power*, *Vis nervosa* seu *nervorum*, *Neurodynamia*, *Neurodynamia*. The power of the nerves as exhibited in the living organism.

NERVOUS PRINCIPLE, Nervous fluid.

NERVOUS SYSTEM. The nerves and nervous centres considered collectively. See Nerves.

NERVOUS SYSTEM OF THE AUTOMATIC FUNCTIONS, Trisplanchnic nerve—n. System, ganglionic, see Trisplanchnic nerve—n. System, organic, see Trisplanchnic nerve—n. System of the senses, Medulla oblongata—n. System of voluntary motion, &c., Medulla spinalis.

NERVOUSNESS, Nervous diathesis.

NERVUS, see Nerve, Penis—n. Acusticus, Auditory nerve—n. Ambulatorius, Pneumogastric nerve—n. Anonymus, Trigemini—n. Auditorius, Auditory nerve—n. Caroticus, Carotid nerve—n. Cervico-facialis, Cervico-facial nerve—n. Communicans faciei, Facial nerve—n. Cutaneus internus, see Saphenous nerves—n. Cutaneus internus longus, see Saphenous nerves—n. Decimus, Pneumogastric—n. Ganglionicus, Trisplanchnic nerve—n. Genito-cruralis, Genito-crural nerve—n. Hypoglossus, Hypoglossus.

NERVUS IMPAR. A prolongation of the neurilemma below the lower extremity of the spinal cord, as a fibrous filament, which is inserted into the base of the coccyx. It was formerly regarded as a nerve;—hence its name.

NERVUS INCISIVUS, Incisive nerve—n. Innominatus, Trigemini—n. Ischiadicus, Sciatic nerve—n. Juvenilis, Penis—n. Latus, Achilles tendon—n. Lingualis lateralis, Pharyngo-glossal nerve—n. Lingualis medius, Hypoglossal nerve—n. Loquens, Hypoglossus—n. Lumbo-inguinalis, Genito-crural nerve—n. Mixtus, Trigemini—n. Musculi pterygoidei, Pterygoid nerve—n. Nasociliaris, Nasal nerve—n. Opticus, Optic nerve—n. Pneumogastricus, Pneumogastric—n. Pudendus externus, see Genito-crural nerve—n. Quintus, Trigemini—n. Spermaticus, see Genito-crural nerve—n. Subcutaneous male, Orbital nerve—n. Sympatheticus medius, Trigemini, Pneumogastric nerve—n. Sympatheticus magnus, Trisplanchnic nerve—n. Sympatheticus minor, Facial nerve—n. Sympathicus, Trisplanchnic nerve—n. Sympathicus medius, Trigemini—n. Temporo-facialis, Temporo-facial nerve—n. Timidus, Motor oculi externus—n. Tremellus, Trigemini—n. Trisplanchnicus, Trisplanchnic nerve—n. Trochlearis, Patheticus nervus—n. Vagus, Pneumogastric nerve—n. Vagus cum accessorio, Spinal nerve—n. Vaso-motorius, Trisplanchnic nerve.

NES, Nesus.

NESPIE, Melissa calamintha.

NESTIA, Jejunum.

NESTIATRIA, Limotherapeia.

NESTIS, Jejunum.

NESTITHERAPEIA, Limotherapeia.

NESTOTHERAPEIA, Limotherapeia.

NESTROOT, Monotropa uniflora.

NETLEAF, *Goodyera pubescens*.

NETTING, Urine.

NETTLE, ARCHANGEL, WHITE, *Lamium album*—n. Day, *Lamium album*—n. Dead, *Lamium album*, L. amplexicaule—n. Dwarf, *Urtica urens*—n. Hedge, *Galeopsis*—n. Horse, *Solanum Carolinense*—n. Pill-bearing, *Urtica pilulifera*.

NETTLE RASH, *Urticaria*—n. Springe, *Urticaria*.

NETTLE, STINGING, COMMON, *Urtica*.

NETTLE TREE, *Celtis occidentalis*.

NETWORT, *Goodyera pubescens*.

NEURADYNAMIA, Neurasthenia.

NEURÆMIA, see Nervous diseases.

NEURÆMIC, see Nervous diseases.

NEURAL; from *νευρον*, 'a nerve.' Relating to a nerve or to the nervous system.

NEURAL ARCH. The arch formed by the posterior projections connected with the body of the vertebra, which protect the medulla.

NEURAL AXIS, see Encephalon—n. Spines, Spinous processes.

NEURALGIA, from *νευρον*, 'a nerve,' and *αλγος*, 'pain.' *Rheumatismus spinarius nervosus*, *Neurodyn'ia*, (F.) *Néuralgie*, *Néurodynie*, *Ner-*

vous pain. A generic name for a certain number of diseases, the chief symptom of which is a very acute pain, exacerbating or intermitting, which follows the course of a nervous branch, extends to its ramifications, and seems, therefore, to be seated in the nerve. The principal neuralgias have been distinguished by the names *facial* (of which the *infra-orbital*, *maxillary*, and *frontal* are but divisions)—the *ilio-scrotal*, *femoro-popliteal*, *femoro-pretibial*, *plantar*, and *cubito-digital*. A division of anomalous neuralgiae has likewise been admitted.

All varieties of neuralgia are obstinate, and the greatest diversity of means has been made use of:—bleeding, general and local,—emetics, purgatives, rubefacients, vesicants, actual cautery, narcotics, mercurial frictions, electricity; destruction of a portion of the nerve, &c. The most successful remedy, perhaps, is the carbonate of iron, given in doses of some magnitude; as, for instance, ℞ss or ℥ij, twice or thrice a day, in molasses. This plan of treatment, continued for a month or two, will often relieve, and ultimately remove this much dreaded affection. The mode in which it acts is by no means clear: but it is almost as certain as any other remedy used in disease in producing its salutary effects. The bowels must be kept free; and all inflammatory symptoms removed during its administration.

NEURALGIA, ANOMALOUS. Under this name Cbaussier has included different neuroses, some of which are characterized by acute pains circumscribed within a short compass, or extending by irradiations, but not having their seat in the course of a nerve; and others which are occasioned by tumours in the course of a nerve, or which succeed contusions or incomplete divisions of nerves.

NEURALGIA BRACHIALIS, Brachialgia—*n.* Brachio-thoracica, Angina pectoris—*n.* Cerebralis, Hemisrania—*n.* Cœliaca, Neuralgia of the solar plexus (Rombert)—*n.* Cruralis, *N. femoro-tibialis*.

NEURALGIA CU'BITO-DIGITALIS, Is'chias nervo'sa digita'lis. In this variety the pain extends from the part where the nerve passes under the inner condyle to the back of the hand and to its cubital edge.

NEURALGIA DENTALIS, Odontalgia nervosa.

NEURALGIA, FA'CIAL, Neuralgia fac'iei, Trismus maxilla'ris seu *Dolorificus* seu *Clon'icus, Ophthalmia, Dolor cr'icæ faciei* seu *faciei period'icus, Hemisrania idiopath'ica, Autal'gia doloro'sa, Tic douloureux, Dolor faciei seu faciei typ'ico caracte'rè* seu *faciei Fothergill'i, Rheumatism'us canero'sus, Rheumatism'us larva'tus, Prosopod'gia, Prosopodyn'ia, Febris top'ica, Ophthalmodyn'ia period'ica, Tortura Oris, Affectus spasmodico-convulsiv'us Labiorum, Face* *ague*, is characterized by acute lancinating pains, returning at intervals; and by twinges in certain parts of the face, producing convulsive twitches in the corresponding muscles. It may be seated in the frontal nerve, in the infra-orbital, or in the maxillary branch of the fifth pair.

Metopodyn'ia, Brow-ague, Neuralgia fronta'lis, Ophthalmodyn'ia, Tic douloureux, Dolor period'icus, &c., commences at the superciliary foramen and extends along the ramifications of the frontal nerve that are distributed on the forehead, upper eyelid, caruncula lacrymalis, and nasal angle of the eyelids. Sometimes it is felt particularly in the orbit.

Neuralgia infra-orbita'ria, Dolor faciei atrox, Rheumatism'us canero'sus, Trismus clon'icus, Prosopod'gia, Febris top'ica, Odontal'gia remittens et intermit'tens, Hemisrania æva, Infra-orbital neuralgia, Tic douloureux, is chiefly felt in the

infra-orbital foramen, whence it passes to the cheek, upper lip, ala nasi, lower eyelid, &c.

In *Neuralgia maxilla'ris, Gnathal'gia, Gnathoneural'gia, Tic douloureux, Trismus catarrh'al'is maxilla'ris*, the pain usually sets out from the mental foramen and passes to the chin, lips, temple, teeth, and tongue.

NEURALGIA, FALSE. A term assigned to pains along a nerve or its ramifications, produced by some body compressing it,—those pains terminating with the removal of the compressing cause.

NEURALGIA FEM'ORO-POPLITE'A, N. Ischiad'ica, Sciatic'a, Schias, Malum ischiad'icum, Morbus ischiad'icus, Passio ischiad'ica, Isch'iagra, Ischial'gia, Ischio'sis, Ischioneural'gia, Coxal'gia, Dolor Ischiad'icus nervo'sus, Is'chias, I. nervo'sa Coturni' seu Coturnii, I. nervo'sa post'ica seu rheumat'ica, Neuris'chias, Cox'agra, Ischial'gia, Ischioneural'gia, (Prov. Devonshire) Bone-shave, (F.) Sciaticque, Goutte sciaticque. This is characterized by pain following the great sciatic nerve from the ischiatic notch to the ham, and along the peroneal surface of the leg to the sole of the foot.

NEURALGIA FEM'ORO-PRÆTIBIA' LIS, N. crural, Ischias nervo'sa anti'ca, Scelal'gia anti'ca. In this, the pain, setting out from the groin, spreads along the fore part of the thigh, and passes down, chiefly on the inner side of the leg, to the inner ankle and back of the foot.

NEURALGIA OF THE HEART, Angina pectoris—*n.* Hepatica, Hepatalgia.

NEURALGIA ILIO-SCROTAL has been rarely observed. It is characterized by a very acute pain, in the course of the branches of the first lumbar pair; this pain follows the crista ili and accompanies the spermatic vessels to the testicle, which is often painfully retracted.

NEURALGIA, LEAD, see Lead rheumatism—*n.* Lienes, Splenalgia.

NEURALGIA MAMMÆ, Ir'ritable breast. An exceedingly painful affection of the female mamma, unaccompanied by inflammation.

NEURALGIA PLANTA' RIS. This is rare; and the pain is confined to the course of the plantar nerves.

NEURALGIA RENUM, Nephralgia—*n.* Spinalis, Spinal irritation—*n.* Testis, Orchidalgia.

NEURANAGEN'ESIS, Neuranoph'ysis, νευρον, 'a nerve,' and αναγενναω, 'to regenerate.' Regeneration or renewal of a nerve.

NEURANAPHYSIS, Neuranagenesis.

NEURAPOPHYSIS, see Vertebræ.

NEURASTHENIA, Neuradyna'mia, Debilitas nervo'sa. Debility or impaired activity of the nerves; from νευρον, 'a nerve,' and ἀσθένεια, 'debility.'—See Irritable.

NEURECTOM'IA, from νευρον, 'a nerve,' and εκτομη, 'excision.' Excision of a nerve.

NEURICUS, Nervous.

NEURILEM'MA, Neurile'ma, Neurily'ma, Neurymen, Perineurion, Fis'tula seu Tu'bulus seu Cap'sula seu Involucrum nervo'rum, Membra'na seu Membran'ula seu Tu'nica seu Indumen'tum nervo'rum, Vagi'na nervo'rum; (F.) Névrlème, Périnère; from νευρον, 'a nerve,' and λεπτα, 'a coat.' The fine transparent, and apparently fibrous membrane that surrounds the nerves—to every filament of which it forms a true canal.

Robin restricts the term *périnère* to the structure that invests the primitive bundles of nerves; and every nervous filament. It differs from the neurilemma, in its ordinary acceptation, which forms a coat of some thickness in the larger nerves, and supports the nutrient vessels, whilst the *périnère* exhibits no capillaries.

NEURILEMMATITIS, Neurilemmitis.

NEURILEMMITIS, Neurilemmat'itis, Neuronemmat'itis, Neurilit'is, (F.) Névrlémite, from

neurilemma, and *itis*, denoting inflammation. Same etymon. Inflammation of the neurilemma.

NEURILITIS, Neurilemmitis.

NEURILYMA, Neurilemma.

NEURINE, *Medulla nervo'rum*, (F.) *Névrine*, *Albumine cérébrale ou du cerveau*, from *νευρον*, 'a nerve.' The substance of which the nervous system is composed. It consists chiefly of albumen and a peculiar fatty matter, associated with phosphorus. There are two kinds of neurine—the one *vesicular* or consisting essentially of *nerve vesicles* or *nerve cells* or *corpuscles*, or, as they are sometimes called, from their prevailing in the ganglia—*ganglion corpuscles* and *ganglion globules*; the other, the *tubular*, formed—as the word imports—of tubules. The former is the *cineritious* or *cortical* nervous matter of the older anatomists;—the latter, the *white* or *medullary*. The *vesicular* neurine has appeared to be restricted to the production and distribution of nerve-power; the *tubular*, to its conduction; yet the experiments of Dr. Brown-Séquard show, that the former conveys sensitive impressions from the posterior spinal roots to the brain.

NEURISCHIAS, Neuralgia femoro-poplitea.

NEURITIC, Nervine.

NEURITIS, *Inflammatio nervo'rum*, *Neurophlogosis*, *Neurophlegmonē*, (F.) *Inflammation des Nerfs*, *Névrite*, from *νευρον*, 'a nerve,' and *itis*, a suffix denoting inflammation. Inflammation of a nerve.

NEUROBLASTA, from *νευρον*, 'a nerve,' and *βλακεια*, 'stump.' Insensibility of the nerves.

NEUROCHONDROIDES, from *νευρον*, 'a sinew,' *χονδρος*, 'a cartilage,' and *ειδος*, 'resemblance.' A hard substance between a sinew and a cartilage.

NEURODES, Nervous.

NEURODYNAMIA, Nervous power.

NEURODYNAMIS, Nervous power.

NEURODYNIA, Neuralgia.

NEUROGAMIA, Magnetism, animal.

NEUROGRAPHY, *Neurographia*; from *νευρον*, 'a nerve,' and *γραφειν*, 'a description.' The part of anatomy which describes the nerves.

NEUROHYPNOLOGY, *Neurohypnologia*, from *νευρον*, 'a nerve,' *υπνος*, 'sleep,' and *λογος*, 'a treatise.' A name given by Mr. Braid to his theory of magnetic sleep. See Magnetism, animal.

NEURO-HYPNOTISM, see Magnetism, animal.

NEUROLEMMATITIS, Neurilemmitis.

NEUROLOGY, *Neurologia*, (F.) *Névrologie*, from *νευρον*, 'a nerve,' and *λογος*, 'a discourse.' The part of anatomy which treats of the nerves.

NEUROMA, (F.) *Névrome*, from *νευρον*, 'a nerve.' A morbid enlargement of a nerve. Applied to subcutaneous, circumscribed, and highly painful tumours formed on the tissue of the nerves; and likewise to a small, hard, grayish tumour, *Neuromatation*, *Nervous ganglion*, *painful tubercle*, of the size of a pea, which is observed in the course of nerves, and appears to be formed from the neurilemma.

NEUROMALACIA; from *νευρον*, 'a nerve,' and *μαλακεια*, 'softening.' Softening of nerves.

NEUROMATION, Neuroma.

NEUROMETERES, Psomē.

NEUROMYELITIS, from *νευρον*, 'a nerve,' *μυελος*, 'marrow,' and *itis*, denoting inflammation. Inflammation of the medullary matter of the nerves.

NEURON, see Nerve.

NEURONOSOS, *Neuronosios*, from *νευρον*, 'a nerve,' and *νοσος*, 'a disease.' A disease of the nerves.

NEURONYGME, *Neuronygmus*, *Neuronyxia*,

from *νευρον*, 'a nerve,' and *συνη*, 'puncture,' *Punctio nervi*. Puncture of a nerve.

NEUROPATHIC, *Neuropathicus*, from *νευρον*, 'a nerve,' and *παθος*, 'a disease.' Belonging to disease of the nerves, to *Neuropathia* or *Neuronosia*.

NEUROPHLEGMONE, Neuritis.

NEUROPHLOGOSIS, Neuritis.

NEUROPLASTY, *Neuroplastikē*, (F.) *Névroplastie*, from *νευρον*, 'a nerve,' and *πλαστικός*, 'forming.' M. Serres applies this term to a ganglionic alteration of the peripheral nerves.

NEUROPYRA, Fever, nervous, Typhus mitior.

NEUROPYRETUS, Fever, nervous, Typhus mitior.

NEUROSES, (F.) *Névroses*, from *νευρον*, 'a nerve.' A generic name for diseases supposed to have their seat in the nervous system, and which are indicated by disordered sensation, volition, or mental manifestation; without any evident lesion in the structure of the parts, and without any material agent producing them. Such is the usual definition. Broussais attributes them to a state of irritation of the brain and spinal marrow.

NEURO-SKELETON, see Skeleton.

NEUROSPASMI, see Spasm.

NEUROSTHENIA, (F.) *Névrosthénie*, from *νευρον*, 'a nerve,' and *σθενεια*, 'strength.' Excess of nervous irritation. Nervous irritation.

NEUROTHELIC APPARATUS, Corpus papillare.

NEUROTIC, Nervine.

NEUROTICA. Diseases of the nervous function. The 4th class of Good's Nosology.

NEUROTOME, *Nérotome*.

NEUROTOMY, *Neurotomia*, (F.) *Névrotomie*. Same etymon. Dissection of nerves. Also, an incised wound of a nerve.

NEUROTROSIS, *Neurotrosmus*, from *νευρον*, 'a nerve,' and *τραωσις*, 'wounding.' *Nervi vulneratio*. Wound or wounding of a nerve.

NEUROTROSMUS, Neurotrosis.

NEURYMEN, Neurilemma.

NEUTA, *Neutha*. A membrane, according to Paracelsus, which covers the eyes and ears of the foetus in utero.

NEUTRAL, from *neuter*, 'neither.' Belonging to neither in particular.

NEUTRAL SALTS, *Secondary salts*, (F.) *Sels neutres*. Salts, composed of two primitive saline substances, combined together, and not possessing the characters of acid or of base.

NEVIS, MINERAL WATERS OF. Thermal springs in the department of Allier, France. Temperature, 136° to 148° Fahrenheit.

NÉVRALGIE, Neuralgia—*n. du Foie*, *Hepatalgia*—*n. des Reins*, *Nephralgia*—*n. de l'Utérus*, *Hysteralgia*.

NÉVRILÈME, Neurilemma.

NÉVRILÈMITE, Neurilemmitis.

NÉVRINE, Neurine.

NÉVRITE, Neuritis.

NÉVRODYNIE, Neuralgia.

NÉVROLOGIE, Neurology.

NÉVROME, Neuroma.

NÉVROSE DU CŒUR, Angina pectoris.

NEVROSES, Neuroses.

NÉVROSTHÉNIE, Neurosthenia.

NÉVROTÈLE, from *νευρον*, 'nerve,' and *σηλη*, 'papilla.' Concerning the papillae of the skin.

APPAREIL NÉVROTÈLE. The papillary apparatus, or arrangement of the skin in the aggregate.—Breschet.

NÉVROTOME, *Neurotome*, *Neurotomus*; from

νευρον, 'a nerve,' and τεμνω, 'I cut,' 'I dissect.' An instrument used by anatomists to dissect the nerves. It is a long and very narrow scalpel, having two edges.

NEVROTOMIE, Neurotomy.

NEVUS, Nævus.

NEW JERSEY, MINERAL WATERS OF.

In the upper part of Morris county and in the county of Hunterdon, near the top of Musconetcong mountain, there are chalybeate springs, which are resorted to. See Schooley's Mountain.

NEW YORK, MINERAL WATERS OF.

The chief waters are those of Saratoga, Ballston, Sharon, and Avon.

NEXUS STAMINEUS OCULI, Ciliary body.

NEZ, Nasus.

NGERENGERE. A singular disease, with the New Zealanders, which Dr. Arthur S. Thomson considers to be a variety of such diseases as the Elephantiasis Græcorum, Lepra of the Cosacks, &c., and which he proposes to call *Lepra gugerengensis*.

NIANDRIA ANOMALA, *Anisodus luridus*.

NICARAGUA WOOD, see *Cæsalpinia*.

NICCOLI SALES, *Salts of Nickel*, (F.) *Sels de Nickel*. These salts are gentle tonics, and have been given with advantage in periodic headache, and in chlorosis and amenorrhœa. The dose of the sulphate is gr. ss or gr. j, three times a day.

NICCOLUM, Nickel.

NICE, CLIMATE OF. The climate of Nice possesses some advantages over the neighbouring climates of Provence and Italy, being free from the sirocco of the latter, and protected from the mistral of the former. Spring is the most unfavourable season. The climate is very dry. It has been a great winter retreat for the consumptive; but does not deserve the encomiums that have been passed upon it.

NICKAR TREE, *Gymnocladus Canadensis*.

NICKEL, *Niccolum*, from (G.) Nickel, 'a prostitute.' A term of detraction, having been applied by the old German writers to what was regarded as a kind of false copper ore. A white malleable metal, s. g. 8.8; less oxidizable than iron. The sulphate has been given as a tonic in the dose of half a grain to a grain.

NICODEMI O'LEUM, *Oil of Nicodemus*.

An oil made by digesting, for some time, — in a mixture of white wine and olive oil, — old turpentine, litharge, aloes, saffron, oxyd of zinc, &c.

NICOTIANA, *N. tabacum* — *n. Minor*, *N. rustica*.

NICOTIANA RUSTICA, called after Nicot, who carried it to Europe; *N. minor*, *Priapeia*, *Hyoscyamus luteus*, *English* or *Wild Tobacco*. *Ord. Solanaceæ*. The leaves possess the properties of tobacco, but are milder.

NICOTIANA TABACUM, *Nicotia'na*, *Herba tabaci*, *Tabacum*, *Petum*, *Petun*, *Hyoscyamus Peruvia'nus*, *Tobacco*, *Virgin'ia tobacco*, *Drunkwort*, (F.) *Tabac*, *Nicotiane*. Tobacco is a violent acro-narcotic; its properties seeming to depend upon a peculiar principle, *Nicotin* or *Nicotianin*. It is narcotic, sedative, diuretic, emetic, cathartic, and errhine. In incarcerated hernia, it is injected, in the form of smoke or infusion, but requires great caution. It is extensively and habitually used as an errhine and sialogogue. The infusion is used to kill vermin, and in some cutaneous eruptions.

NICOTIANE, *Nicotiana tabacum*.

NICOTIANIN, see *Nicotiana tabacum*.

NICOTIN, see *Nicotiana tabacum*.

NICTATIO, *Hippus*, *Nictatio*.

NICTATION, *Nicta'tio*, *Nictita'tio*, from *nictare*, *nictatum*, 'to wink.' *Clonus nictita'tio*, *Ble-*

pharium'us, *Palpebra'tio*, *Twinkling of the eye*, (F.) *Clignotement*, *Cillement*. A rapid and repeated movement of the eyelids, which open and shut alternately. As occasionally performed, it is physiological; if repeatedly, a disease. It seems to be executed chiefly by the motor 7th pair of nerves; but it is necessary that the extensor 5th pair should likewise be in a state of integrity.

NIDAMENTUM, *Decidua*.

NIDDICK, *Nucha*.

NIDOREUX, *Nidorous*.

NIDOROUS, *Nidoro'sus*, (F.) *Nidoreux*, from *Nidor*, *Cnissa*, 'the smell of any thing burnt,' &c. Having the smell of burnt or corrupt animal matter.

NIDUS, *Nidus hirundinis*.

NIDUS HIRUNDINIS, *Nidus*, *Swallows' Nest*. A deep fossa in the cerebellum — so called from its fancied resemblance — situate between the verum medullare posterius in front, and the nodulus and uvula behind.

NIEDERBRONN, MINERAL WATERS OF. Saline waters in the department of Bas-Rhin, France, which contain chloride of sodium, sulphate of magnesia, and carbonic acid.

NIELLE, *Nigella*.

NIEPA BARK, see *Samadera Indica*.

NIGELLA, *Coptis*.

NIGELLA, *N. Sati'va*; *Ord. Ranunculaceæ*; (*nigrella*, from *niger*, 'black.' *Melan'thium*, *Cuminum Nigrum*, *Fennel flower*, *Nutmeg flower*, *Devil in a bush*, (F.) *Nigelle*, *Nielle*, *Faux cumin*, *Fleur Sainte-Catherine*. This small southern European and Syrian plant was formerly used medicinally as an expectorant and deobstruent, errhine, sialogogue, &c.

NIGELLE, *Nigella*.

NIGHT-BLINDNESS, *Hemeralopia*.

NIGHT EYES, *Crusta genu equinæ* — *n. Mare*, *Incubus*.

NIGHTSHADE, AMERICAN, *Phytolacca decandra* — *n. Bittersweet*, *Solanum dulcamara* — *n. Common*, *Solanum* — *n. Deadly*, *Atropa belladonna* — *n. Enchanter's*, *Circea Luteitana* — *n. Garden*, *Solanum* — *n. Palestine*, *Solanum sanctum* — *n. Stinking*, *Hyoscyamus niger* — *n. Vine*, *Solanum dulcamara* — *n. Woody*, *Solanum dulcamara*.

NIGREDO À SOLE, *Ephelides*.

NIGRITA, *Negro*.

NIGRITIES, *Melanopathia* — *n. Ossium*, *Caries*.

NIGRITISM, *Melanopathia*.

NIGRITUDO, *Melanosis* — *n. Pulmonum*, *Melanosis pulmonum*.

NIGROMANTIA, *Necromancy*.

NIGROR, *Melasma*.

NIGRUM OCULI, *Pupil*.

NIHIL ALBUM, *Zinci oxydum* — *n. Griseum*, *Zinci oxydum*.

NILIACUM, *Mel*.

NIMA QUASSIÓ'DES, *Sima'ba quassioides*, a native of Nepal and the Himalaya mountains, is as bitter as quassia, and is used like it.

NINDSIN, *Sium ninsi*.

NINE BARK, *Spiræa opulifolia*.

NINE-DAY FITS, *Trismus nascentium*.

NINSI, *Sium ninsi*.

NINSING, *Panax quinquefolium*.

NINZIN, *Sium ninsi*.

NIPPLE, *Sax. nypele*. The *Teat*, *Tit*, *Tet*, *Dug*, *Pap*, (Old Eng.) *Nebbe*, *Titty*; *Acromæ'tium*, *Uber*, *Staph'ylis*, *Staphyl'ium*, *Eubona*, *Mammil'la*, *Mammel'la*, *Papill'a*, *P. mamma*, *U'beris apex*, *Thela*, *Tit'hê*, *Titthos*, *Tit'thion*, *Tit'this*, (F.) *Mamelon*, *Bout de sein*. The conical tubercle, situate at the centre of the breast. Towards the central part of each breast the skin changes colour, and assumes a rose tint in young females.

or a reddish brown in those who have suckled several children. The circle is called the *Areola* or *Aureola* of the nipple. The nipple is capable of erection on being excited.

NIPPLE, ARTIFICIAL, *Bout de Sein*.

NIPPLEWORT, *Lapsana*.

NIRLES, *Herpes phlyctænodes*.

NIRMALLEES, *Strychnos potatorum*.

NIRMILLIES, *Strychnos potatorum*.

NISUS, *Nixus*, *Peira*, *Straining*, from *Lat. nitor, nisus*, and *nixus*, 'to endeavour.' A voluntary retention of the breath, so as to force down the diaphragm; the abdominal muscles being at the same time contracted forcibly. In this manner the contents of the abdomen are compressed; and the evacuation of the feces, urine, &c., is effected.

NISUS, Effort—*n. Formativus*, Plastic force.

NISUS PARTURIENTIIUM. The efforts or forcing during parturition.

NITEDULA, *Cicindela*.

NITON, MINERAL WATERS OF. These springs in the Isle of Wight contain iron, and sulphate of alumina and potassa.

NITRAS, Nitrate—*n. Hydrargyricus*, see *Hydrargyri nitras*—*n. Hydrargyrosus*, *Hydrargyri nitras*—*n. Kalicus*, *Potassæ nitras*—*n. Lixivæ*, *Potassæ nitras*—*n. Natrium*, *Soda*, nitrate of—*n. Plumbicus*, *Plumbi nitras*—*n. Potassæ cum sulphure fusus*, *Potassæ nitras fusus sulphatis paucillo mixtus*—*n. Sub-bismuthicum*, *Bismuth*, sub-nitrate of.

NITRATE, *Nitras*, *Azotate*. A salt, formed of a base and nitric acid. Several nitrates are employed in medicine.

NITRATE ACIDE DE MERCURE, see *Hydrargyri nitras*—*n. d'Ammoniaque*, *Ammonia nitras*—*n. d'Argent*, *Argenti nitras*—*n. Mercureux*, *Hydrargyri nitras*.

NITRE, *Potassæ nitras*—*n. Cubic*, *Soda*, nitrate of.

NITRIC, *Nitricus*, (F.) *Nitrique*, from *nitrum*, 'nitre.' Appertaining or relating to nitre.

NITRIC ACID, *Acidum nitricum* seu *azoticum* seu *septicum* seu *nitri*, *Aqua fortis*, *Nitrous acid*, *Spiritus nitri duplex seu nitri fumans seu nitri Glauberi* seu *nitri ac'idus*, *Azotic acid*, *Dephlogisticated nitrous acid*, *Oxyseptonic acid*, (F.) *Acide nitrique ou azotique*. Nitric acid is obtained from nitre—*Nitrate of potassa*. Its odour is suffocating; taste very acid and caustic. It is corrosive, liquid, colourless, and transparent. S. g. 1.500.

Strong nitric acid is rarely used except as an application to foul, indolent ulcers, or to warts. When given internally, it is in the form of the *ACIDUM NITRICUM DILUTUM*, *Spiritus nitri simplex* seu *nitri vulgaris*—*Diluted nitric acid*, which, in the Pharmacopœia of the United States, consists of *nitric acid*, f $\frac{3}{4}$ ij, *water*, f $\frac{3}{4}$ ij by measure. Diluted largely with water, it is used, as a drink in fevers of the typhoid kind; in chronic affections of the liver, syphilis, &c.; but in the latter affections, it is not to be depended upon.

NITROGEN, from *νιτρον*, 'nitre,' and *γεννω*, 'I generate.' Azote.

NITROGEN, GA'SEOUS OXIDE OF, *Nitrous oxide*, *Protaxide* of nitrogen or of azote, *Paradise*, *Intoxicating* or *Laughing gas*, *Dephlogisticated nitrous gas*, *Gas azoticum oxygenatum*. This gas, when respired, produces singular effects; great mental and corporeal excitement; and, generally, so much exhilaration as to render the appellation, "laughing gas" by no means inappropriate. It has not been much used in medicine, although recommended in paralysis. Its effects are too violent and too transient to render it a valuable remedial agent; and, in the delicate, it has been

productive of unpleasant effects, inducing palpitation, fainting, and convulsions.

NITROGEN, PROTOXIDE OF, Nitrogen, gaseous oxide of.

NITROGENE, Azote.

NITROGENIUM, Azote.

NITROGENIZED, *Azoted*, *Azotized*, (F.) *Azoté*. Containing nitrogen or azote i—as a nitrogenized, azoted, or azotized aliment.

NITRO-MURIATIC ACID, *Ac'idum nitro-muriaticum* seu *muriaticum nitro'so-oxygens'tum*, *Mens'trum auri*, *Nitro-hydrochloric* or *Hydrochloro-nitric acid*, *Aqua regia* seu *styg'ia* seu *regis Chrysalea*, (F.) *Eau régale*. A mixture of the nitric and muriatic acids, has been used in diseases of the liver, in the form of a bath for the feet and legs, made sharply acidulous. It has, also, been employed, and with more uniform results, in cutaneous affections. Whatever advantage it may possess in internal diseases—and these advantages are doubtful—they are probably dependent upon the chlorine formed by the mixture, or upon the properties possessed by the mineral acids in general. See *Chlorine*.

The *Acidum nitro-muriaticum* of the Pharmacopœia of the United States is formed by mixing four fluidounces of nitric acid with eight of muriatic acid.

NITROUS OXIDE, Nitrogen, gaseous oxide of.

NITRUM, *Natron*, *Potassæ nitras*—*n. Antiquorum*, *Soda*—*n. Cubicum*, *Soda*, nitrate of—*n. Facitium*, *Borax*—*n. Flammanis*, *Ammonia nitras*—*n. Lunare*, *Argenti nitras*—*n. Prismaticum*, *Potassæ nitras*—*n. Rhomboidale*, *Soda*, nitrate of—*n. Saturninum*, *Plumbi nitras*—*n. Vitriolatum*, *Schroederi*, *Potassæ sulphas*—*n. Tabulatum*, *Potassæ nitras fusus sulphatis paucillo mixtus*—*n. Vitriolatum*, *Potassæ sulphas*.

NIX, *Snow*—*n. Fumans*, *Calx viva*—*n. Zinci*, *Zinci oxydum*.

NIXUS, *Nisus*—*n. Parturientium*, *Nisus parturientium*.

NOAH'S ARK, *Cypripedium luteum*.

NOBLE, *Nobilis*. Principal, Essential.

NOBLE PARTS, *Partes essential'es*. Some anatomists have given this name to parts, without which life cannot exist; such as the heart, liver, lungs, brain, &c. The organs of generation have, likewise, been so called.

NOCAR, *νωκαρ*, *Torpor Soporificus*. Lethargic torpor.

NOCTAMBULATIO, *Somnambulism*.

NOCTAMBULISMUS, *Somnambulism*.

NOCTAMBULUS, *Somnambulist*.

NOCTISURGIIUM, *Somnambulism*.

NOCTUINI OC'ULI. Gray or blue eyes.—*Castelli*.

NOCTURNA BELLA, *Coitus*.

NOCTURNAL, *Nocturnus*, *Nycter'inus*, from *nox*, *noctis*, 'night.' Relating to night, as *Febris nocturna* seu *nycter'ina*. A fever occurring in the night.

NODDING, *Annuatio*, *Nutatio*.

NODDLEPOX, *Syphilomania*.

NODE, *Nodus*, *Emphy'ma exosto sis perios'tea*, (F.) *Nodosité*, *Nœud*; from *Hebr. גו', 'a heap.'* A hard concretion or incrustation, which forms around joints attacked with rheumatism or gout. Some include, under this name, exostoses, articular calculi, ganglions, and even the chronic swellings of the joints known under the name of white swellings.

NODES, *Edentulus*.

NODI DIGITORUM MANUS, *Phalanges of the fingers*—*n. Lactæ*, *Infarctus mammae lacteæ*—*n. Nervorum*. Ganglions, nervous—*n. Hæmorrhoidales*, see *Hæmorrhoids*.

NODOI, *Nefrendes*.

NODOSIA, see *Nefrendes*.

NODOSITÉ, Node.

NODULI ARANTII, *Noduli Morgagni*, *Cor'pora Aran'tii*. The small sesamoid bodies situate on the periphery of the semilunar valves of the aorta and pulmonary artery for the better occlusion of the artery.

NODULI MORGAGNII, *Noduli Arantii* — *n. Nervorum*, Ganglions, nervous.

NOD'ULUS, diminutive of *nodus*, 'a knot.' A small knot. A small prominence or lobule in the portion of the cerebellum, which forms the posterior boundary of the fourth ventricle. The nodulus is on the median line, and before the uvula. See Vermiform process, inferior.

NODULUS UTERINUS, Pessary.

NODUS, Articulation, Edentulus, Encystis, Hamma — *n. Cerebri*, *Pons Varolii* — *n. Chirurgicus*, Knot, surgeon's — *n. Encephali*, *Nœud de l'Encéphale*, *Pons Varolii* — *n. Hystericus*, Angone — *n. Vitæ*, Centrum vitale.

NOESIS, Intellect.

NEUD, Node — *n. du Chirurgien*, Knot, surgeon's — *n. d'Emballleur*, Knot, packer's.

NEUD DE L'ENCÉPHALE (F.), *Nodus Enceph'ali*. M. Cruveilhier, under this name, includes the pons Varolii, peduncles of the cerebrum and cerebellum, and the tubercula quadrigemina.

NEUD VITAL, Centrum vitale.

NEUDS, Calculi, arthritic.

NOIR ANIMAL, Carbo animalis — *n. d'Os*, Carbo animalis.

NOISETIER, Corylus avellana.

NOISETTE, Corylus avellana (nut).

NOIX, *Juglans regia* (nut) — *n. Américaine*, *Jatropha curcas* — *n. des Barbades*, *Jatropha curcas* — *n. Cathartique*, *Jatropha curcas* — *n. d'Eau*, *Trapa natans* — *n. de Galle*, see *Quercus infectoria* — *n. d'Igasur*, *Ignatia amara* — *n. de Serpent*, see *Thevetia Ahouai* — *n. Vomique*, see *Strychnos nux vomica*.

NOLA CULINARIA, *Anemone pulsatilla*.

NOLI ME TANGERE, *Lupus*, see *Chancreux (Bouton)*.

NOMA, Cancer aquaticus — *n. Pudendorum*, Colpocæcæ infantilis.

NOMAD, *Nomade*, *Nomas*, (*vouas*, *vouados*), from *voun*, 'pasturage.' An epithet given to people who have no fixed habitation, and who travel, with their flocks, from country to country, for pasturage. Such are the Tartars. By analogy, the word *Nomad'ic* has been applied to a spreading ulcer.

NOMBRIL, Umbilicus — *n. de Vénus*, *Cotyledon umbilicus*.

• **NOME**, Cancer aquaticus, Phagedenic ulcer.

NOMENCLATURE, *Nomenclat'io*, *Nomenclat'ura*, *Onomatolog'ia*, *Onomatocle'sis*, *Terminol'ogy*, *Terminol'ogy*, *Glossol'ogy*, *Orismol'ogy*, from *onoma*, 'name,' and *kalos*, 'I call.' A collection of terms or words peculiar to a science or art. In all sciences, nomenclature is an object of importance; and each term should convey to the student a definite meaning. The Lavoisierian nomenclature was a valuable gift to chemistry; and anatomy has derived advantage from the labours of Barclay, Dumas, and Chaussier, who have given names to parts indicative of their situation. See Muscle. The nomenclature of pathology has required the greatest attention; and although repeated attempts have been made to improve it, the barbarous terms that disgrace it are still frequently adopted. It consists of Hebrew and Arabic terms; Greek and Latin, French, Italian, Spanish, German, English, and even Indian, African, and Mexican; often barbarously and illegitimately compounded. A want of principle in founding the technical terms of medicine is every where observable. They have

been formed:—1. From colour; as *Melana*, *Melas*, *Atrabilis*, *Leuce*, *Alphos*, *Chlorosis*, *Rubeola*, *Scarlatina*, *Purpura*, &c. 2. From duration; as *ephemeral*, *quotidian*, *tertian*, and *quartan*, continued, and intermittent, &c. 3. From Birds, Beasts, Fishes, Insects, and Plants; as *Rabies canina*, *Cynanche*, *Boutimira*, *Pica*, *Hippopoyon*, *Elephantiasis*, *Urticaria*, *Lichen*, *Ichthyos'is*, &c. 4. From Persons or Places; as *Morbus Herculeus*, *Facies Hippocratica*, *Lep'ra Arabum*, *Plica Polonica*, *Sudor Anglicus*, *Morbus Gallicus*, *Ignis Sancti Antonii*, *Chorea Sancti Viti*, &c.

NOMUS, Cancer aquaticus.

NON-COMPOS, Lunatic.

NON-COMPOS MENTIS, Lunatic.

NON-NAT'URALS, *Non natural'ia*. Under this term the ancient physicians comprehended air, meat, and drink, sleep and watching, motion and rest, the retentions and excretions, and the affections of the mind. They were so called, because they affect man without entering into his composition, or constituting his nature; but yet are so necessary that he cannot live without them.

NONPLACENTAL, see Placental.

NON-SANE, Insane.

NON-STRIATED MUSCULAR FIBRE, see Muscular fibre.

NONUS HUMERI PLACENTINI, *Teres minor* — *n. Linguae musculus*, *Genio-glossus* — *n. Vesalii*, *Peronæus tertius*.

NOOS, Intellect.

NORICE, Nurse.

NORMA VERTICALIS, see Normal.

NORMAL, *Norma'lis*, from *norma*, 'a perpendicular,' 'a rule.' According to rule; perpendicular.

The normal line, *norma vertical'is*, of Blumenbach, is a vertical line let fall from the prominence of the frontal bone and shaving the superior maxillary, so as to mark the projection of the latter bone beyond the arch of the forehead.

NORRIS'S DROPS, see *Antimonium tartarizatum*.

NORTH CAROLINA, MINERAL WATERS OF. The chief springs are the *Warm* and *Hot Springs of Buncombe*. Temperatures varying from 94° to 104° Fahr. They have not much mineral impregnation, and have the properties of the thermal waters in general.

NORTON'S DROPS, *Liquor hydrargyri oxy-muriatis*.

NORYSS, Nurse.

NOSACERUS, Sickly.

NOSE, *Nasus* — *n. Bleed*, *Epistaxis*.

NOSE, RUNNING AT THE, *Coryza*.

NOSELIA, *νοσληα*, *Nosocom'e'sis*, *Nosocom'i'a*, from *nosos*, 'disease.' Care of the sick.

NOSEMA, Disease.

NOSENCEPH'ALUS, (F.) *Nosencéphale*, from *nosos*, 'disease,' and *εγκεφαλος*, 'brain.' A monster whose skull is open only on the frontal and parietal regions, the posterior fontanelle being distinctly present.

NOSEROS, Insalubrious, Sick.

NOSEUMA, Disease.

NOSOCOMESIS, *Noselia*.

NOSOCOMIA, *Noselia*.

NOSOCOMIAL, *Nosocomia'lis*, from *νοσος*, 'a disease,' and *κομω*, 'I take care of,' 'an hospital.' Relating to an hospital, — as 'nosocomial or hospital fever.'

NOSOCOMIUM, Hospital.

NOSOCOMUS, *Infirmier*.

NOSOCRATIC, Specific.

NOSODES, Insalubrious, Sick, Sickly.

NOSODOCHIUM, Hospital.

NOSOGENESIS, Pathogeny

NOSOGENIA, Pathogeny.

NOSOGENY, Pathogeny.

NOSOGEOGRAPHY, Geography, medical.

NOSOG'RAPHY, *Nosograph'ia*, from *nosos*, 'a disease,' and *γραφω*, 'I describe.' A description of diseases.

NOSOL'OGY, *Nosolog'ia*, from *nosos*, 'a disease,' and *λογος*, 'a discourse.' A name given to that part of medicine whose object is the classification of diseases. The most celebrated nosological systems have been those of Sauvages (1763), Linnaeus (1763), Vogel (1764), Sagar (1776), Macbride (1772), Cullen (1772), Darwin (1796), Selle, Crichton (1804), Parr (1809), Swediaur (1812), Pinel (1813), Young (1813), Good (1817), Hosack (1818), &c. Besides these general nosographies, others have been published on *Surgery* exclusively, none of which are particularly worthy of enumeration amongst nosological systems. Nosological arrangements have, also, been formed of single families or groups of diseases. Plenck, of Baden, is the author of two different treatises of this kind: the one, a methodical arrangement of the diseases of the eyes, and the other, of cutaneous diseases. Dr. Willan published an arrangement of cutaneous diseases, which was completed by Dr. Bateman, and adopted into the Nosology of Dr. Hosack. Mr. Abernethy, also, published a methodical classification of tumours, and many other partial nosological classifications might be enumerated. Also, Pathology.

NOSON'OMY, *Nosonom'ia*, from *nosos*, 'a disease,' and *ονομα*, 'name.' The nomenclature of diseases.

NOSOPHYTA, from *nosos*, 'disease,' and *φυτον*, 'a plant.' A disease supposed to be produced by, or to consist in the development of parasitic plants—as *porrigio*, *mentagra*, &c.—Gruby.

NOSOP'EUS, *Nosopœt'icus*, from *nosos*, 'disease,' and *ποιω*, 'I make.' That which causes disease.

NOSOS, Disease.

NOSOTAX'Y, *Nosotax'ia*, from *nosos*, 'a disease,' and *ταξις*, 'arrangement.' The distribution and classification of diseases.

NOSOTHEO'RIA, from *nosos*, 'disease,' and *θεωρια*, 'doctrine.' The doctrine or theory of disease.

NOSTAL'GIA, from *nostros*, 'return,' 'a journey home,' and *αλγος*, 'pain.' *Nostoma'nia*, *Nostros'ia*, *Apodemial'gia*, *Pathopatridal'gia*, *Patopatridal'gia*, *Philopatridal'gia*, *Philopatridoma'nia*, *Ademon'ia*, *Ademos'ynê*, Home-sickness, (F.) *Nostalgie*, *Maladie du pays*. An affection produced by the desire of returning to one's country. It is commonly attended by slow wasting, and sometimes by hectic, which may speedily induce death. M. Pinel properly regards it as a variety of melancholy.

NOSTOMANIA, Nostalgia.

NOSTRASSIA, Nostalgia.

NOSTRILS, Nares.

NOSTRUM, Arcanum.

NOSTRUM, CHITTICK'S. An empirical remedy for stone in the bladder, said to be a solution of alkali in veal-broth.

NOTA MATERNA, *Nævus*—n. Primitiva, Groove, primitive.

NOTÆ INFANTUM, *Nævi*.

NOTAL, Dorsal.

NOTAL'GIA, *Notial'gia*, (*Nostalgia*, improperly, of Köchlin and others,) from *notos*, 'the back,' and *αλγος*, 'pain.' Pain in the back. Spinal irritation.

NOTANENCEPHALIA, see *Notencephalus*.

NOTCH, Teut. Nocke, Ital. *Nocchia*, *Emarginatio*, *Emarginatura*, (F.) *Échancrure*. A depression or indentation of different shape and size, observed on the circumference or edges of certain bones.

NOTCH OF THE CONCHA, *Incisura tragica*.

NOTCH, ETHMOID'AL, (F.) *Échancrure ethmoïdale*, is situate on the frontal bone, and joins the ethmoid.

NOTCHES, ISCHIAT'IC, (F.) *Échancrures Ischiatiques*, are two in number:—the greater and the less. The former is large, situate at the inferior part of the pelvis, and formed by the sacrum and ilium. It gives passage to the sciatic nerve, pyramidalis muscle, and to the superior gluteal vessels and nerves. The latter is much smaller than the other, from which it is separated by the sciatic spine. It gives passage to the tendon of the obturator internus, and to the internal pudic vessels and nerves.

NOTCH, PAROT'ID, (F.) *Échancrure parotidienne*, is the triangular space comprised between the parotid edge of the inferior maxillary bone and the mastoid process; so called, because it lodges the parotid gland. The notches in soft parts are generally called *Fissures*.

NOTCH, SCAP'ULAR, *Incisura scapularis*, *Lu'mula scap'ulæ*, (F.) *Échancrure scapulaire*. The notch on the superior edge or *costa* of the scapula, which is converted into a foramen by means of a ligament, and gives passage to the suprascapular nerve.

NOTCH, SEMILUNAR OF THE STERNUM, *Fourchette*.

NOTENCEPHALIA, see *Notencephalus*.

NOTENCEPH'ALUS, from *notos*, 'the back,' and *εγκεφαλος*, 'the brain.' A monster whose head, with the brain, is on the back. The condition is termed *Notencephalia* and *Notanencephalia*.—I. G. St. Hilaire.

NOTHEUSIS, Degeneration.

NOTHIA, Degeneration.

NOTHROTÉS, Torpor.

NOTHUS, False.

NOTIÆUS, Dorsal—n. *Myelus*, *Medulla spinalis*.

NOTIALGIA, *Notalgia*.

NOTOM'ELES, from *notos*, 'the back,' and *μελεος*, 'unprofitable.' A monster with superfluous flesh on the back.

NOTOMYELITIS, *Myelitis*.

NOTOMYELUS, *Medulla spinalis*.

NOTON, Dorsum.

NOTOS, Dorsum, Vertebral column

NOUAGE DE LA CORNÉE, Caligo.

NOUÉ, (*Bandage*) (F.); 'knotted,' from *nodus*, 'a knot.' A bandage which has a considerable number of knots placed over each other. It is made with a roller, 6 or 7 ells long, rolled into two balls, and is used to compress the parotid region, after the extirpation of the parotid gland.

Also, an epithet applied to children in whom the disease of rickets has swollen the articulations.

It is, likewise, applied to the gout when it has caused nodes on the joints.

NOUET (F.), *Nod'ulus*. A bag filled with medicinal substances, and infused in a liquid to communicate their properties to it.

NOUFFER'S, MADAME, REMEDY, *Poly-podium filix mas*.

NOURRICE, Nurse.

NOURRICIER, Nutritious.

NOURRITURE, Aliment.

NOUS, Intellect.

NOUURE, Rachitis.

NOUVEAU-NÉ (F.), ('new-born,') *Neonatus*, *nuperrimè Natus*, *Neog'enes*, *Neog'iles*, *Neog'ilus*, *Neog'onius*, *Neog'nus*, *Infans recens natus*. That which has been just born. A new-born infant.

NOVACULA, Razor.

NOW, Vertex.

NOYAU, see Cytoblast.
NOYAU CENTRAL DES PÉDONCULES
DU CERVELET, Corpus dentatum.

NOYER, *Juglans regia*—n. de Ceylon, Adhatoda.

NOYRIS, Nurse.

NUAGE, Nebula.

NUBECULA, Enaërema, Nebula.

NUBES, Enaërema, Nebula.

NUBILE, *Nu'bilis*, 'marriageable,' 'fit to marry.' Generally, the period of puberty is considered to be the age at which both sexes are nubile. They are truly nubile, however, only when they are fitted to procreate healthy and vigorous children, and are competent to discharge their duties as parents.

NUBILITY, *Nubil'itas*, (F.) *Nubilité*; same etymon. The state of being nubile or marriageable.

NUBILOSUS, Nepheloid.

NUCES AQUATICÆ, see *Trapa natans*—n. *Quercus*, see *Quercus alba*.

NUCHA, *In'tum*, *Nucha cap'itis*, (F.) *Nuque*. The nuche, hinder part or nape of the neck, (Prov.) *Niddick*. The part where the spinal marrow begins.

Ligamentum Nuchæ. A strong ligament from the neck, proceeding from one spinous process to another, and inserted into the occipital bone. It is very strong in quadrupeds. It is called in them *Pazywazy*, *Paxwaz*, *Packwaz*, formerly, *Faxwaz*, and *Tarwaz*.

NUCHAL REGION, *Re'gio nuchæ* seu *nuchæ* *lis* seu *occipitalis inferior*, from *nucha*. The region of the nucha or nape of the neck.

NUCISTA, see *Myristica moschata*.

NUCK, CANAL OF. A small prolongation of the peritoneum often sent into the inguinal canal of the female fetus. So called from Nuck, who first described it.

NUCLEATED, *Nuclea'tus*, from *nucleus*, 'a kernel.' Having a nucleus or central particle. Applied to the elementary cells of organized tissues; the vital properties of which are seated in the nucleus. See Cytoblast.

NUCLEATED CELL, see Cytoblast.

NUCLEI CEMBRÆ, see *Pinus Cembra*—n. Ossei, Ossification, points of—n. *Pinæ*, see *Pinus pinea*.

NUCLEIN, see Globules of the blood.

NUCLEOLE, see Cytoblast.

NUCLEOLULE, see Cytoblast.

NUCLEOLUS, see Cytoblast.

NUCLEUS, see Cytoblast—n. *Blastodermatis*, *Tache embryonnaire*—n. *Centralis*, *Corpus dentatum*—n. *Cicatriculæ*, *Tache embryonnaire*—n. *Dentatus*, *Corpus dentatum*—n. *Dentis*, *Dental pulp*—n. *Encased*, *Cytoblast*—n. *Fimbriatus*, *Corpus dentatum*—n. *Furunculi*, see *Furunculus*—n. *Germinal*, see *Molecule*—n. *Germinativus*, see *Molecule*—n. *Olivæ*, *Corpus dentatum*—n. *Ossificationis*, *Ossification*, point of—n. *Rhomboidalis*, see *Corpus dentatum*.

NUCULA TERRESTRIS, *Bunium bulbocastanum*.

NUKE, Nucha.

NULE-KNEED, *Entogonyancon*.

NUMERICAL METHOD, see Method, numerical.

NUM'MULAR, *Nummular'is*. Relating to money, from *nummus*, 'money.' An epithet applied to the sputa in phthisis, when they flatten at the bottom of the vessel, like a piece of money, and to the mode of aggregation of the red corpuscles of the blood—'nummular rolls.'

NUMMULARIA, *Lysimachia nummularia*.

NUMPOST, Abscess.

NUPHAR AD'VENA, *Nymphæa ad'vena*, Yellow Pond Lily, Spatterdock. Order, Nym-

phæaceæ; indigenous in Canada and the United States. The root is a popular tonic, and the fresh root bruised is applied as a cataplasm.

NUPHAR LUTEUM, *Nymphæa lutea*.

NUQUE, Nucha.

NURICE, Nurse.

NURSE, Sax. *noþice*, *Nutrix*, (from *nourish*, itself from *nutrire*), *Tithe'nē*, *Trephou'sa*, *Lactans*, *Lacticans*, *Nutrients*, *Thelactaria*, *Mamma*, (L.) *Nutrice*, (F.) *Nourrice*, (Sc.) *Noyris*, *Nourys*, *Nurice*, *Norice*. One who suckles her own child or another's. Also, one that has the care of a sick person, *Care-taker*, (F.) *Garde-malade*.

NURSE, DRY. One who gives every care to a child, but does not suckle it.

NURSE, MONTHLY. One who attends a woman during the month of her parturient state.

NURSE, WET, (Sc.) *Milkwoman*. A female, who suckles the child of another.

To 'nurse artificially,' is to bring up a child by the hand.

NŪS, Intellect.

NUSUS, Disease.

NUT, BRAZILIAN, see *Pichurim* beans—n. *Butter*, *Juglans cinerea*—n. *Cembros*, see *Pinus cembra*—n. *Clearing*, *Strychnos potatorum*—n. *French*, *Juglans regia*—n. *Para*, see *Pichurim* beans—n. *Physic*, *Jatropha curcas*—n. *Pine*, see *Pinus picea*—n. *Pistachio*, see *Pistacia vera*—n. *Poison*, see *Strychnos nux vomica*—n. *Purging*, see *Jatropha* and *Croton tiglium*—n. *Rattle*, *Nelumbium luteum*—n. *Sassafras*, see *Pichurim* beans—n. *Soap*, see *Sapindus saponaria*.

NUT TREE, MALABAR, *Adhatoda*.

NUT, VOMIC, see *Strychnos nux vomica*—n. *Zirbel*, see *Pinus picea*.

NUTATION, *Nuta'tio*, from *nutare*, *nutatum*, 'to nod.' Nodding. Also, a constant oscillation of the head, by which it moves involuntarily in one or more directions.

NUTATOR CAPITIS, *Sterno-cleido-mastoidæus*.

NUTMEG, see *Myristica moschata*—n. *Butter*, see *Myristica*—n. *Flower*, *Nigella*—n. *Liver*, *Liver*, *nutmeg*.

NUTMEGS. Testes.

NUTRICATIO, Nutrition.

NUTRICIER, Nutritious.

NUTRICIUS, Nutritious.

NUTRIENS, Aliment, Nurse.

NUTRIMEN, Aliment.

NUTRIMEN'TAL, *Nutrimen'talis*, *Aliment'al*, *Alimen'tary*, *Alimenta'tis*, (F.) *Alimenteux*, from *nutrimen*, 'aliment.' Having the qualities of food or nutriment.

NUTRIMENTUM, Aliment, Pabulum.

NUTRIT'ION, *Nutrit'io*, *Nutrica'tio*, *Nutritus*, *Alit'ura*, *Threpsis*, from *nutrire*, *nutritum*, 'to nourish.' Nutrition is that function by which the nutritive matter already elaborated by the various organic actions, loses its own nature, and assumes that of the different living tissues,—to repair their losses and maintain their strength. Sometimes the word is used in a more extended signification, to express the whole series of actions by which the two constant movements of composition and decomposition are accomplished, in organized bodies. Nutrition, then, would comprehend digestion, absorption, respiration, circulation, and assimilation; the latter being *nutrition*, properly so called, and being effected in the intermediate system over the whole of the body,—the cells of the tissues attracting from the blood the elements necessary for their repair.

NUTRITION, FORCE OF, Plastic force.

NUTRIT'IOUS, *Nutrit'ius*, *Nutritive*, *Alit'ilis*, *Troph'imos*, *Troph'ides*, (F.) *Nourricier*, *Nutricier*, *Alibile*. Same etymon. Having the

quality of nourishing: as nutritious food, nutritive lymph, &c.

NUTRITIOUS or NUTRITIVE ARTERIES, (F.) *Arteres nutritivæ*. Arterial branches which enter the *foramina nutriticia* of long bones, and penetrate to the medullary membrane.

NUTRITIVE, Nutritious, (F.) *Nutritif*. Also, relating to nutrition: hence the 'nutritive functions,' or those that are concerned in nutrition.

NUTRITUS, Aliment, Nutrition.

NUTRIX, Mamma, Nurse.

NUTS, WATER, Nelumium luteum.

NUX AROMATICA, see *Myristica moschata* —n. *Avellana*, *Corylus avellana* (nut) —n. *Barbadiensis*, *Jatropha curcas* —n. *Becuba*, *Ibicuiba* —n. *Cathartica Americana*, *Jatropha curcas* —n. *Gallia*, see *Quercus cerris* —n. *Juglans*, *Juglans regia* —n. *Medica*, *Coco* of the *Maldives* —n. *Metella*, *Strychnos nux vomica* —n. *Methel*, *Datura stramonium* —n. *Moschata*, see *Myristica moschata* —n. *Myristica*, see *Myristica moschata* —n. *Pistacia*, see *Pistacia vera* —n. *Unguentaria*, see *Myristica moschata* —n. *Vomica*, *Strychnos nux vomica*.

NYCTALOPE, see *Nyctalopia*.

NYCTALOPEX, see *Nyctalopia*.

NYCTALOP'IA, from νύξ, νυκτος, 'night,' and οπτομαι, 'I see'; *Parop'sis Lucif'uga*, *Nyctalop'iasis*, *Cæcitatis diurna*, *N. Vius nocturnus*, *Oxyo'pia*, *Hemeralo'pia* (moderns), *Amblyo'pia meridiana*, *Hemeratophlo'sis*, *Hemerotophlo'sis*, *Photophob'ia*, *Photophophthal'mia*, *Dyso'pia l'uminis*, *Vius a'crior*, *Nyctalopy*, *Dayblindness*, *Diurnal amauro'sis*, (F.) *Vue nocturne*, *Aveuglement de Jour*. The faculty of seeing during the night, with privation of the faculty during the day. It affects both eyes at once, when idiopathic. Its duration is uncertain, and treatment very obscure. It is, however, a disease of nervous irritability, and one of excitement of the visual nerve in particular. The indications of cure will consequently be—to allay direct irritation in every way; to excite counter-irritation by blisters; and to gradually accustom the eye to the impression of light.

One labouring under this affection is called a *Nyctalope*, *Nyctalops*, *Nyctalopez*.

Nyctalopia has also been used for night blindness. See *Hemeralopia*.

NYCTALOPS, see *Nyctalopia*.

NYCTALOPY, *Nyctalopia*.

NYCTERINUS, *Nocturnal*.

NYCTHEMERON, see *Dies*.

NYCTHEMERUM, from νύξ, νυκτος, 'night,' and ημερα, 'day.' The space of 24 hours, or of a day and night. Certain complaints continue only so long.

NYCTOBADIA, *Somnambulism*.

NYCTOBASIS, *Somnambulism*.

NYCTOBATESIS, *Somnambulism*.

NYCTOBATIA, *Somnambulism*.

NYCTOTYPHLOSIS, *Hemeralopia*.

NYGMA, *Wound, punctured*.

NYGMATICUM EMPLASTRUM, *Emplastrum resinae*.

NYMPHA, Clitoris.

NYMPHÆ, from νυμφη, 'a water nymph.' *Alæ minores seu internae minores clitoridis seu muliebres minores*, *Carunculæ cuticulæ res*, *Cristæ clitoridis*, *Collieuli vaginæ*, *Myrtoschilla*, *Myrtoscheilides*, *Labia minora seu interna seu pudendi minora*, (F.) *Nymphes*, *Petites lèvres*. Two membranous folds, which arise from the lateral parts of the prepuce of the clitoris, and descend on the inner surface of the labia majora; terminating, by becoming gradually thinner, about the middle of the circumference of the orifice of the vagina. They are formed each of two folds of the mucous membrane of the vulva; and

contain, in their substance, a thin layer of spongy erectile tissue. Their use seems to be,—not, as was once supposed, to direct the course of the urine, which notion gave rise to their name, but to favour the elongation and dilatation of the vagina in pregnancy and labour.

The word νυμφη, *Nymphæ*, has also been used synonymously with clitoris by Oribasius, Aëtius, &c.

NYMPHÆA ADVENA, Nuphar advena.

NYMPHÆA ALBA, Leuconymphæa, Nenu'phar, Microleuconymphæa, Castalia speciosa, White Water Lily, (Sc.) Cambie-leaf, (F.) Nénuphar ou Nénufar blanc, Lis des étangs, Lune ou Volant d'Eau. Nat. Ord. Ranunculacææ. Sex. Syst. Polyandria Monogynia. Formerly employed as a demulcent, antaphrodisiac, emollient, and slightly anodyne remedy.

NYMPHÆA INDICA, N. nelumbo —n. *Major lutea*, *N. lutea*.

NYMPHÆA LU'TEA, N. major lutea seu umbilicalis, Nuphar luteum, Nenu'phar lutea, Nymphæanthus vulgaris, Yellow Water Lily, (Sc.) Cambie-leaf, (F.) Nénuphar jaune. Used for the same purposes as *Nymphæa alba*. The flowers are called in Norfolk, England, *Brandy bottles*.

NYMPHÆA NELUM'BO, N. Indica, Faba Egyptiaca, Cy'anus Ægyptiacus, Pontic or Egyptian Bean. The fruit of this is eaten raw in Egypt and some of the neighbouring countries; and is considered to be tonic and astringent.

NYMPHÆA ODORA'TA, Sweet-scented Water Lily, Sweet water or White pond or Toad Lily, Cow or Water Cabbage, (F.) Nénuphar odorant. An indigenous plant, growing in most parts of the United States in fresh water ponds, and on the borders of streams, and having large white, beautiful, sweet-scented flowers. The root is very astringent and bitter. It is sometimes made into a poultice and used as a discutient.

NYMPHÆA UMBILICALIS, N. lutea.

NYMPHE, Clitoris.

NYMPHES, Nymphæ.

NYMPHITIS; from νυμφη, 'the clitoris,' and itis, denoting inflammation. *Epiderrh'itis, Inflammatio Clitoridis*. Inflammation of the clitoris.

NYMPHOCLOUIA, Nymphomania.

NYMPHOMANIA, from νυμφη, 'a bride,' and mania, 'fury'; *Furor uterinus, Uteroromania, Lagne'sis furor femininus, Metromania, Edæogargalis, Edæogargaris'mus, Thelygon'ia, Erotomania, Edæomania, Aidoionomania, Melincho'ia uterina, Nymphoclui'a, Symp'toma surpitudinis, Andromania, Gynæcæomania, Machlos'ynê, Entelipathi'a, Tenti'go vene'reæ, Hysteromania, Sallac'itas vulvæ, Uteri pruritus, Brachu'na, Arascon, Arastum, Estromania, (F.) Nymphomania, Fureur utérine. An irresistible and insatiable desire, in females, for the venereal act. It occurs in those particularly, who possess a nervous temperament, and vivid imagination, especially if excited by improper language, masturbation, &c. Its course, as described, is as follows. In the commencement, the sufferer is a prey to perpetual contests between feelings of modesty and impetuous desires. At an after period, she abandons herself to the latter, seeking no longer to restrain them. In the last stage the obscenity is disgusting; and the mental alienation, for such it is, becomes complete. The treatment consists in the use of the same means as are required in the satyriasis of man. When the mental alienation is complete, solitude is indispensable.*

NYMPHON'CUS, from νυμφη, 'the nymph,' and ογκος, 'a tumour.' A morbid tumefaction of the nymphæ.

NYMPHOTOMY, Nymphotomia, Nymphæctomum Sectio, (F.) Nymphotomie, from νυμφη, and

νεμειν, 'to cut.' An operation, known and practised for a long time, which consists in the excision of the nymphæ. The operation is had recourse to, when they are attacked with scirrhus, cancer, fungus, or gangrene; or when they are so large as to interfere with walking or coition. *Nymphotomy* is the circumcision of the female. It is practised in some countries.

Some authors have used the term *Nymphotomy* for amputation of the clitoris.

NYPHOZANTHUS VULGARIS, *Nymphæa lutea*.

NYSTAGMUS. A partial rotatory movement of the eyeball from side to side. Also, *Coma vigil*.

NYXIS, Puncture.

O.

OAK, BLACK, *Quercus tinctoria*—o. Common, *Quercus robur*—o. Jerusalem, *Chenopodium botrys*.

OAK LUNGS, *Lichen pulmonarius*.

OAK ORCHARD ACID SPRINGS. These springs—eight in number—are in Genesee county, New York, eight miles from Lockport. They are acid salines, consisting of free sulphuric acid, sulphate of protoxide of iron, and sulphates of alumina, magnesia and lime. They are powerfully astringent and tonic.

OAK, POISON, *Rhus toxicodendron*—o. Red, *Quercus rubra montana*—o. Sea, *Fucus vesiculosus*—o. Spanish, *Quercus rubra montana*—o. White, *Quercus alba*.

OARIOCYESIS, Pregnancy, ovarian.

OARION, Ovary.

OARION'CUS, *Ovariophy'ma*, *Ovarium tumidum*, *Tumor Ovari'i*, from *ωαριον*, (diminutive of *ωον*, 'an egg,') 'the ovarium,' and *ογκος*, 'swelling,' Ovarian tumour.

OARIOPAREC'TAMA, *Oophorauz'e*, from *ωαριον*, 'ovarium,' and *παρεκτεινεν*, 'to extend,' Enlargement of the ovary.

OARIORRHEX'IS, from *ωαριον*, 'ovarium,' and *ρηξις*, 'rupture,' *Ruptura Ovari'i*. Rupture of the ovary.

OARIO'OMY, *Oariotom'ia*, *Ovariotom'ia*, *Ovario'omy*, from *ωαριον*, 'the ovarium,' and *τομν*, 'incision,' The operation for removing the ovary.

OARIT'IS, *Oöphoritis*, *Ooritis*, *Inflammatio Ovari'i*, *Ovaritis*, (F.) *Inflammation de l'Ovaire*, from *ωαριον*, 'the ovarium,' and *itis*, the termination denoting inflammation. Inflammation of the ovarium. When occurring in childhood, it is the *Lochoöphoritis* of some.

OARIUM, Ovary.

OARTHROC'ACE, formed by contraction from *omo-artroace*. Said to be used by Rust to designate inflammation of the scapulo-humeral articular surfaces.—Nysten.

OATMEAL, see Avena.

OATS, Avena.

OBAUDITIO, Barycoia.

OBAUDITUS, Barycoia.

OBCÆCATIO, Cæctas.

OB'DORMIT'IO, from *ob*, and *dormio*, *dormitum*, 'to sleep.' The state of the limbs being asleep;—*Stupor artuum*.

OB'DUCTIO LEGALIS, Autopsia cadaverica legalis.

OBELÆA RAPHE, Sagittal suture.

OBSITAS, Polysarcia—o. Colli, Struma adiposa—o. Nimia, Pimelosis—o. Viscerum, Physconia adiposa.

OBESITÊ, Polysarcia.

OBSITY, Polysarcia.

OBSUS, Corpulent.

OB'FUSCATIO, Amaurosis.

OBITUS, Death.

OBJECTIVE CONE. The cone of light pro-

ceeding from an object, the apex of which is on the object, and the base on the cornea.

OBJECTIVE SENSATIONS, see Sensation.

OBLINITO, Inunction.

OBLIQUE', *Obli'quus*, Any thing inclined, or which deviates from the vertical line. Anatomists have given this name to certain muscles, which have an oblique direction as regards the plane that divides the body into two equal and symmetrical halves. These are:—

OBLIQUE MUSCLES OF THE ABDOMEN. They are two in number, and distinguished into: 1. *Obliquus Exter'nus Abdom'inis* seu *descendens exter'nus* seu *descendens* seu *major*, (F.) *Ilio-pubo costo-abdom'inal*, *Costo-abdom'inal* (Ch.), *Grand oblique*, *Oblique externe*. One of the broadest muscles of the body. It is situate at the lateral and anterior part of the abdomen; and is flat and quadrilateral. It is attached, *above*, to the outer surface and lower edge of the 7 or 8 last ribs; *below*, to the anterior third of the external lip of the crista ili: *before*, it terminates at the linea alba by means of a broad and strong aponeurosis, which covers the rectus, and presents towards its inferior part two very solid fibrous fasciculi, which are inserted,—the one at the symphysis, the other at the spine of the pubis,—under the name of *Pillars of the Abdominal Ring*. These pillars leave between them an opening, which forms the inferior orifice of the inguinal canal. The obliquus externus abdominis depresses the ribs, and carries them backwards during a strong expiration. It impresses on the chest a movement of rotation, and bends the thorax upon the pelvis, and conversely. It contracts, also, the abdominal cavity. 2. *Obliquus Inter'nus Abdom'inis* seu *ascendens* seu *minor* seu *internus* seu *ascendens internus*, *Musculus accli'vis*, (F.) *Iliolombo-costo-abdominal*, *Ilio-abdominal* (Ch.), *Muscle petit oblique* ou *oblique interne*, is broad, especially before; thin, and irregularly quadrilateral, like the preceding, beneath which it is situate. It is attached, *above*, to the inferior edge of the cartilages of the 5th, 4th, 3d, and 2d false ribs; *below*, to the anterior two-thirds of the interstice of the crista ili, to the posterior part of the crural arch, and to the pubis; *behind*, to the spinous processes of the last two lumbar vertebræ, and to those of the first two portions of the sacrum; *before*, to the linea alba. Its upper fibres run obliquely upwards and forwards; the middle are horizontal; and the lower pass obliquely downwards and forwards.

These last, in the male, are dragged down through the inguinal ring, when the testicle descends, and form the two fasciculi of the cremaster.

The obliquus internus resembles the O. externus in function.

OBLIQUE MUSCLES OF THE EYE, *Amato'rii*, *Circumagen'tes*, *Rotato'res Oculi*, are two in number.

They are distinguished into: 1. *Obliquus Superior Oculi*, *Anato'rius musculus*, *Trochlea'ris*, *Trochlea'tor*, *Obliquus major*, *Circumductio'nis op'fec.* *Longie'simus oculi*, (F.) *Optico-trochlei scleroti-cien*, *Grand trochléateur* (Ch.), *Grand oblique de l'œil*, *O. supérieur de l'œil*, *Amoureux (Muscle)*, is situate at the inner and upper part of the orbit. It is small, round, fusiform, and reflected upon itself in the middle of its course. Behind, it is attached to the inside of the foramen opticum; and when it arrives opposite the internal orbital process, it becomes a small, round tendon, which slides in a cartilaginous pulley fixed to the os frontis, and is reflected, at an acute angle, to proceed downwards and outwards, and to attach itself to the outer and back part of the globe of the eye. This muscle carries the globe of the eye forwards and inwards; making it experience a movement of rotation, which directs the pupil downwards and inwards. This is conceived to be an involuntary muscle as well as the next. In sleep, according to Sir C. Bell, when the power over the straight or voluntary muscles of the organ is nearly lost, the eye is given up to the oblique muscles, which lodge the transparent cornea under the upper eyelid. At the approach of death, the same thing is observable; hence, the turning up of the eye, at such time, is not an evidence of agony or suffering, but of insensibility. 2. *Obliquus Inferior seu minor oculi*, (F.) *Mozillo-scleroticien*, *Petit Trochléateur* (Ch.), *Petit oblique ou oblique inférieur de l'œil*, is situate at the anterior and inferior part of the orbit. It is flat and attached to the inner and anterior part of the orbital surface of the superior maxillary bone, on the outside of the lachrymal gutter; from thence it passes outwards and backwards, and terminates by an aponeurosis, at the posterior and inner part of the globe of the eye. It carries the globe of the eye inwards and forwards; and directs the pupil upwards and outwards.

OBLIQUE MUSCLES OF THE HEAD. These are two in number. 1. *Obliquus Superior seu minor Cap'itis*, (F.) *Trachélo-atloïdo-occipital*, *Atloïdo-eous-mastoi'dien* (Ch.), *Muscle oblique supérieur ou petit oblique de la tête*. This muscle is situate at the sides of, and behind, the articulation of the head: it is flat and attached, on the one hand, to the top of the transverse process of the atlas; and, on the other, terminates at the occipital bone, beneath the inferior curved line, and sometimes, also, at the mastoid region of the temporal bone. It extends the head,—inclining it to one side. 2. *Obliquus Inferior Capitis*, *Obliquus major*, (F.) *Spini-azoido-trachéli-atloïdien*, *Azoido-atloïdien* (Ch.), *Oblique inférieur ou grand oblique de la tête*, is situate at the posterior part of the neck and head. It is round, fusiform; attached to the spinous process of the axis, and proceeds to terminate behind and below the summit of the transverse process of the atlas. It impresses, on the first vertebra and the head, a movement of rotation, which turns the face to one side.

OBLIQUE PROCESSES, see Vertebrae.

OBLIQUE EXTERNE, *Obliquus externus abdominis*—o. *Grand*, *Obliquus externus abdominis*—o. *Grand de l'œil*, *Obliquus superior oculi*—o. *Grand de la tête*, *Obliquus inferior capitis*—o. *Inferieur de l'œil*, *Obliquus inferior oculi*—o. *Inferieur de la tête*, *Obliquus inferior capitis*—o. *Interne*, *Obliquus internus abdominis*—o. *Petit*, *Obliquus internus abdominis*—o. *Petit de l'œil*, *Obliquus inferior oculi*—o. *Petit de la tête*, *Obliquus superior oculi*—o. *Supérieur de l'œil*, *Obliquus superior oculi*—o. *Supérieur de la tête*, *Obliquus superior capitis*.

OBLIQUITÉ DE LA MATRICE, *Retroversio uteri*.

OBLIQUUS ASCENDENS, *O. internus abdo-*

minis—o. *Auris*, *Laxator tympani*—o. *Descendens externus*, *O. externus abdominis*—o. *Major*, *O. externus abdominis*, *O. inferior capitis*, *O. superior oculi*—o. *Minor*, *O. internus abdominis*—o. *Minor capitis*, *O. superior capitis*—o. *Minor oculi*, *O. inferior oculi*.

OBLITERATED, *Obliteratus*; from *obliterate*, (*ob*, and *litera*), 'to efface.' A vessel or duct is said to be obliterated, when its parietes have approximated and contracted such an adhesion to each other that the cavity has completely disappeared.

OBLITERATIO COMPLETA, *Hapantismus*.

OBLIVIO, *Amnesia*—o. *Iners*, *Lethargy*.

OLOBIUM, *Antilobium*.

OB'OLUS, *Onolosat*. A weight of 9 or 10 grains.

OBSCURÆ PARTES, *Genital organs*.

OBSCURISSEMENT DE LA VUE, *Caligo*.

OBSERVATION, *Observatio*, (from *ob*, and *servare*, *servatum*, 'to keep,' e. g. in sight.) *Tere'sis*, *Symparateresis*. Act of examining a thing by means of the external senses. This word is employed in several acceptations. It expresses—1. The action of observing—2. The aggregate of knowledge, afforded by observation. In French—but not in English—it means the case or history of the phenomena presented by a patient in the course of a disease.

OBSERVATION, *Case*—o. *Method of*, *Numerical method*.

OBSTET'RIC, *Obstet'ricus*, *Obstet'rical*; same etymon as *obstetrics*. Relating or appertaining to *obstetrics*,—as 'obstetric auscultation,' 'obstetric exploration,' &c.

OBSTETRIC CHAIR, *Labour-chair*.

OBSTETRICANS, *Accoucheur*.

OBSTET'RICS, from *Obstetrix*, 'a midwife, (*obstare*, *ob*, and *stare*, 'to stand before.')

Tokol'ogy, *Tocol'ogy*, *Maëi'a*, *Maëi'a*, *Maëi'a*, *Maëi'a*, *Maëi'a*, *Ars obstetric'ia*, *Obstet'ricy*, (F.) *Obstétrique*. The art of midwifery. Midwifery in general.

OBSTETRICY, *Obstetrics*.

OBSTÉTRIC, *Obstetrics*.

OBSTETRIX, *Midwife*.

OBSTIPATIO, *Constipation*—o. *Tenesmus*, *Tenesmus*.

OBSTIPATIO, see *Hump*.

OBSTIPITAS, *Torticollis*—o. *Capitis seu Colli*, *Torticollis*.

OBSTRUCTIO, *Emphraxis*, *Stegnosis*—o. *Alvi*, *Constipation*—o. *Ductus Alimentarii*, *Constipation*—o. *Ductus Stenotiani*, *Stenosis*—o. *Intestinalis*, *Constipation*—o. *Recti Spastica*, *Stricture of the Rectum*, *spasmodic*—o. *Pulmonum pituitosa febrilis*, *Peripneumonia notha*.

OBSTRUCTION OF THE INTESTINES, *Enteremphraxis*.

OB'STRUENS, from *obstruo*; (*ob*, and *struere*, 'to build'), 'I stop up by building against.' *Op'pilativus*. A medicine which closes the crifices of ducts or vessels.

OBSTUPEFACIENTIA, *Narcotics*.

OBTONDANT, *Obtundens*.

OBTURCATIO, *Decollation*.

OBTUMESCENCIA, *Mutitas*.

OBTUNDENS, (F.) *Obtundant*, from *obtundere*, (*ob* and *tundere*, 'to beat,') 'to beat against,' and therefore to blunt the edge. An epithet applied to remedies that were supposed, according to an erroneous theory, to be possessed of the power of blunting the acrimony of the humours. A demulcent is one of these.

OBTURAMENTUM, *Emphragma*.

OBTURATEUR, *Obturator*.

OBTURATEUR DU PALAIS, *Palate*, *artificial*.

OBTURATIO, *Emphraxis*.

OBTURATOR, *Obturatorius*, (F.) *Obturator*,

from *obturare*, (akin to *Supra*, 'a door,' [?]) 'to close,' 'stop up the entrance.' A name given in anatomy, to several parts.

OBTURATOR ARTERY, *Arteria obturatoria seu obturatoria*, (F.) *Sous-pubio-fémorale* (Ch.), *Artère obturatrice*, arises, most commonly, from the hypogastric. It is, however, frequently given off from the epigastric; a matter of importance to be determined in cases of femoral hernia. Of 500 obturator arteries examined by M. J. Cloquet, 340 were furnished by the hypogastric, and 152 by the epigastric or crural. When it arises from the hypogastric, it passes forwards and outwards, and then turns horizontally into the cavity of the pelvis, to issue from this cavity by the opening left at the upper part of the obturator membrane. When, on the contrary, the obturator artery arises from the epigastric or the crural, it descends obliquely inwards, behind the os pubis, to the obturator foramen. At its exit from the pelvis, the artery divides into two branches, a posterior and an anterior, which are distributed to the muscles of the anterior and superior part of the thigh.

OBTURATOR FORAMEN, *Foramen Obturatorium seu obturatum seu infra-pubium seu os ilii seu thyroideum seu Thyroïdes seu Amplum Pelvis*, (F.) *Trou sous-pubien*. A large opening, of an oval or triangular form, in the anterior part of the os innominatum, on the outside of the symphysis pubis and beneath the horizontal ramus of the os pubis. This foramen is closed by a membranous ligament.

OBTURATOR LIGAMENT OR MEMBRANE. *Sub-pubic membrane*, is a fibrous membrane, fixed to the whole circumference of the obturator foramen, except above, where an opening remains for the passage of the vessels and nerves of the same name.

OBTURATOR MUSCLES, *Obturatorres, Rotatores femoris*. These are two in number. They are divided into

a. Obturator Externus, Extra-pelvio-pubi-trochantérien, Sous-pubio-trochantérien externe (Ch.) A muscle, situate at the anterior and inner part of the thigh. It is broad, flat, and triangular; and is attached, on the one hand, to the anterior surface of the os pubis, to that of the ischium, and to the anterior surface of the obturator ligament. Its fleshy fibres converge to the tendon, which proceeds to be inserted at the inferior part of the cavity of the great trochanter. This muscle rotates the thigh outwards.

b. Obturator Internus, Marsupialis, Bursa-lis, Intra-pelvio-trochantérien, Sous-pubio-trochantérien interne (Ch.) is seated, almost entirely, in the pelvis. It arises from the inner surface of the obturator ligament, and from the posterior part of the circumference of the obturator foramen, and is inserted, by means of a strong tendon, running between the two portions of the gemini, into the cavity at the root of the great trochanter; after having turned upon the ischium, which forms for it a kind of pulley. This muscle also rotates the thigh outwards.

OBTURATOR NERVE, *Sous-pubio-fémoral*, (Ch.), proceeds principally from the 2d and 3d lumbar nerves. It descends into the pelvis; gains the obturator foramen; gives branches to the obturator muscles, and divides, behind the abductor primus and pectinalis, into two branches; one anterior, whose branches are distributed to the first two adductors, gracilis, and integuments; the other, posterior, distributing its ramifications to the obturator externus and third adductor.

OBTURATOR VEIN has, ordinarily, the same arrangement as the artery. It is common, however, to find it arising from the epigastric; whilst the

corresponding artery proceeds from the hypogastric, and conversely.

The term *obturator*, (F.) *Obturateur*, is, also, applied to an instrument, adapted for closing an opening caused by a wound or disease.

OBTURATOIRES, Obturator muscles.

OBVOLVENTIA, Demulcents.

OCA, Oxalis tuberosa.

OCCECATIO, Cæcitas.

OCCIPITAL, *Occipitalis*. That which belongs to the occiput.

OCCIPITAL ARTERY. This arises from the posterior part of the external carotid, beneath the parotid. It proceeds backwards, passes between the mastoid process and the transverse process of the atlas; reaches the occipital bone and divides into two branches, which may be called ascending posterior and anterior, and are distributed to the neighbouring muscles and ligaments.

OCCIPITAL BONE, *Os occipitale seu occipitii seu occipitalis seu spheno-basilare seu memorialis seu nervosum seu basilare seu prære seu pyramidalis seu sextum cranii seu lambdoïdes seu lambdæ seu laudæ seu puppis seu nervosum seu fibrosum seu pelvicephalicum*, (F.) *Os occipitalis*, is situate at the posterior and inferior part of the cranium, which it assists in forming. It is flat, symmetrical, and curved upon itself. It presents, 1. An occipital or posterior surface, which is convex, and has, upon the median line, the *basilary surface*, the *foramen magnum*, through which passes the spinal marrow with its membranes and vessels,—the *external occipital crest*, the *external occipital protuberance*; and, at the sides, the *upper curved line*, *large rough arched ridge* or *transverse arch* or *linea semicircularis*, the *lower curved line*, the *posterior condyloid fossæ*, the *condyles* for the articulation of this bone with the atlas; and the *anterior condyloid fossæ*, pierced by a foramen for the passage of the ninth pair of nerves. 2. A *cerebral* or *anterior surface*. On the median line are: the *basilary fossæ*, the *inner orifice* of the foramen magnum, the *internal occipital crest*, the *internal occipital protuberance*, the *cruciform spine*; a *channel*, which lodges the termination of the straight sinus, and on each side, the *occipital fossæ*, distinguished into *superior* or *cerebral*, and *inferior* or *cerebellous*, and separated by a groove which lodges the lateral sinus. 3. The surfaces of the occipital bone are separated by four ridges and four angles. The two superior edges are articulated with the parietal bones; the two lower join the temporal; and the anterior angle, under the name *basilary process*, is united to the sphenoid.

The occipital bone is developed from four points of ossification; and sometimes from a greater number.

OCCIPITAL MUSCLE, *Occipitalis*. Many anatomists have given this name to the posterior fasciculus of the occipito-frontalis.

OCCIPITAL NERVE, *Sub-occipital nerve*, (F.) *Première paire trachélienne* (Ch.), *Nervus occipitalis* or *os occipitalis*. It arises from the upper part of the spinal marrow by eight or ten filaments, united in two fasciculi. Thus formed, it passes between the foramen magnum and the posterior arch of the atlas; and, at this place, forms a long ganglion, afterwards dividing into two branches. Of these, the *anterior*, which is long and small, makes a turn above the transverse process of the atlas and forms an anastomotic noose with a branch of the second cervical nerve. The *posterior* branch, larger and shorter, divides into seven or eight branches, which are distributed to the muscles of the upper and back part of the neck.

OCCIPITAL REGION, Occiput.

OCCIPITAL VEIN. Its roots follow exactly the course of the branches of the artery, and unite

into a single trunk, which opens into the internal jugular vein, and sometimes into the external.

OCCIPITIUM, Occiput.

OCCIPITO-ATLOID, *Occipito-atloïdeus*, (F.) *Occipito-atloïdien*. That which has reference to the occiput and atlas.

OCCIPITO-ATLOID ARTICULATION is the articulation of the condyles of the occipital bone with the superior articular cavities of the atlas. It is strengthened by two ligaments; the one anterior, the other posterior, called *occipito-atloid ligaments*: the one extends from the anterior, the other from the posterior, arch of the atlas, to the corresponding portion of the circumference of the foramen magnum.

OCCIPITO-AX'OID, *Occipito-axoïdeus*, (F.) *Occipito-axoïdien*. That which relates to the occipital bone and the axis or second vertebra.

OCCIPITO-AXOID ARTICULATION is the connection of the occipital bone with the axis or second vertebra, although these bones are not really articulated, but are merely retained in apposition by three strong ligaments, the posterior of which is called the *occipito-axoid*, and the two others *odontoid*.

OCCIPITO-FRONTA'LIS, *Digastricus cranii*, *Epicranius*, *Fronta'lis et occipitalis*. The majority of anatomists call by this name the whole of the fleshy plane, with the epicranial or coronal aponeurosis, (see *Calotte*), which covers the head from the occiput to the forehead. It is attached, by its posterior fasciculus, to the two outer thirds of the upper curved line of the occipital bone, and to the outer surface of the mastoid portion of the temporal; and, by its anterior fasciculus, it terminates at the eyebrow, where it becomes confounded with the superciliaris, pyramidalis nasi, and orbicularis palpebrarum.

The occipito-frontalis, by the contraction of its anterior fasciculus, draws forward a part of the integuments of the cranium. It wrinkles the skin of the forehead transversely, and may, also contribute to open the eye by its decussation with the orbicularis palpebrarum. The posterior fasciculus of the muscle draws backwards a part of the skin of the cranium, and assists in stretching the common aponeurosis.

OCCIPITO-MENINGE'AL. That which belongs to the occipital bone, and to the meninges or dura mater.

OCCIPITO-MENINGEAL ARTERY, in Chaussier's nomenclature, is a branch of the vertebral, given off to the dura mater at its entrance into the cranium.

OCCIPUT, *Occipit'ium* (ob, and caput), *Regio occipita'lis*, *In'ion*, *Opisthocra'nium*, *Opisthoceph'alon*, *Hysteroera'nium*, *Prora*, *Occipit'ium*. The back part of the head, formed by the occipital bone.

OCCIPUT, Soft, Craniotabes.

OCCLU'SION, *Occlu'sio*, *Mycē*, from *occludere*, *occlusum*, 'to shut up.' Sometimes this word signifies, simply, the transient approximation of the edges of a natural opening—the *occlusion of the eyelids*, for example; at others it is synonymous with imperforation, as *occlusion of the pupil*, *vagina*, &c.

OCCULT DISEASES, see Latent.

OCEAN SPRINGS, see Mississippi, Mineral waters of.

OCEMA, Vehicle.

OCHETEUMATA, Nares.

OCHETOS, Canal.

OCHBUS, Scrotum.

OCHLE'SIS, from *ὄχλος*, 'a crowd.' A term, applied by Dr. George Gregory to a morbid condition induced by the crowding together of sick persons under one roof.

OCHREA RUBRA, Hæmatites.

OCHTHIASIS, Paleness.

OCHROMA, Paleness.

OCHROPYRA, Fever, yellow.

OCHROTES, Paleness.

OCHROTYPHUS, Fever, yellow.

OCHTHIASIS, Molluscum.

OCHTHODES, Callous.

OCIMUM ADCSCENDENS, O. Basilicum.

OCIMUM BASIL'ICUM, *O. adscen'dens* seu *pi'o'sum* seu *racemo'sum* seu *citra'tum*, *Beren'daros*, *Basil'icum*, *B. majus* seu *citra'tum*, *Common* or *Citron basil*, (F.) *Basilic commun*; *ἄκρον*, from *ακρῶς*, 'swift,' owing to its swift growth. *Nat. Ord. Labiatae*. *Sex. Syst.* Didynamia Gymnospermia. This herb has a fragrant odour and aromatic taste. It is used as a condiment, and has been supposed to possess nervine properties.

OCIMUM CARYOPHYLLA'TUM, *O. min'imum*, *Small* or *Bush basil*. Possesses properties similar to the former. It is sometimes used as snuff.

OCIMUM CITRATUM, O. Basilicum—o. Pilosum, O. Basilicum—o. Racemosum, O. Basilicum.

OCOTEA AMARA, *Nectandra cymbarum*—o. *Cymbarum*, *Nectandra cymbarum*—o. *Pichurim*, see *Pichurim* beans—o. *Puchury*, see *Pichurim* beans.

OCREA, *Hippocampus minor*, *Shin*.

OCTA'NA, *Hebdomada'ria*, *Febris hebdomada'ria*, from *octo*, 'eight.' A fever whose paroxysms recur every week;—a supposititious case.

OCTA'RUS. A pint. The eighth part of a wine-gallon. Same etymon. It contains sixteen fluidounces (Ph. U. S.),—20 fluidounces imperial measure.

OCTOPUS, *Synapheocephalus*, *Hemipages*.

OCUTUNX, from *octo*, 'eight,' and *uncia*, 'an ounce.' A weight of eight ounces.

OCULAR, *Ocula'ria*, (F.) *Oculaire*, from *oculus*, 'an eye.' Of or belonging to the eye.

OCULAR CONE. The cone formed within the eye by a pencil of rays proceeding from an object; the base of the cone being on the cornea,—the apex on the retina.

OCULARES COMMUNES, *Motors oculorum*—o. *Dentes*. Canine teeth.

OCULARIA, *Euphrasia officinalis*.

OCULI MARMARYGODES, *Metamorphopsia*—o. *Palpebrarum scabies pruriginosa*, *Ophthalmia tarsi*.

OCULIST, *Oculis'ta*, *Ophthalmia'ter*, *Med'icus ocula'rius* seu *ophthal'micus*, from *oculus*, 'an eye.' One who occupies himself, chiefly, with the management of diseases of the eye.

OCULISTIQUE. Ophthalmology.

OCULO-MUSCULAR NERVES, COMMON, *Motors oculorum*.

OCULUM MOVENS PRIMUS, *Rectus internus oculi*—o. *Movens quartus*, *Rectus inferior oculi*—o. *Movens secundus*, *Rectus externus oculi*—o. *Movens tertius*, *Rectus superior oculi*.

OCULUS, Eye—o. *Bovinus*, *Hydrophthalmia*—o. *Bovis*, *Chrysanthemum leucanthemum*—o. *Bubulus*, *Hydrophthalmia*—o. *Cæsius*, *Glaucoma*—o. *Duplex*, *Binoculus*—o. *Elephantinus*, *Hydrophthalmia*—o. *Genu*, *Patella*—o. *Lacrymans*, *Epiphora*—o. *Purulentus*, *Hyppopyon*—o. *Simplex*, see *Monoculus*—o. *Typhonis*, *Scilla*.

OCYMUM, see *Ocimum*—o. *Sylvestre*, *Clino-podium vulgare*.

OCYDINIC, *Ocytocio*.

OCYPH'ONUS, from *ακρῶς*, 'quick,' and *φονος*, 'murder.' An agent that kills speedily.

OCYTOGEUS, *Ocytocio*.

OCYTOTIC, *Oxytocio*, *Ocytoc'eus*, *Ocytoc'ius*, *Ocyt'ocus*, *Ocyodin'ic*, *Odinago'gus*, from *ακρῶς*, 'quick,' and *τοκος*, 'labour.' Any thing that expedites parturition.

OCYTOCIUS, *Ocytocio*.

OCYTOCUS, *Ocytocio*.

OD. A term, employed by Reichenbach to designate an "effective force," which, according to him, not only manifests itself in contact with man, as in the case of magnets, crystals, the hands, &c., but, also, at distances, as from the sun, moon, and stars—indeed from all matter. Such a force is termed by him *odic*.

ODAXIS'MUS, *Odaxes'mus*, *Odontocne'sis*, *Odontocnes'mus*, *οδᾶξιμος*; a sensation of biting or pricking, as that in the gums which precedes the appearance of the teeth.

ODES. A suffix, see Eidos.

ODEUR, Odour.

ODIC, see Od.

ODIN, Pains, (labour.)

ODINAGOGUS, Ocytotic.

ODINOL'YSIS, from *odin*, 'labour pains,' and *λῡσις*, 'solution.' Mitigation of labour pains.

ODINOPÆ'A, from *odin*, 'labour pains,' and *ποιέω*, 'I make.' Agents that encourage labour pains.

ODIS, Pains, labour.

ODME, Odour.

ODONTAGOGON, Dentagra.

ODONT'AGRA, from *odon*, *odontos*, 'a tooth,' and *αγρα*, 'a seizure.' A rheumatic or gouty pain in the teeth. Dentagra.

ODONTAL'GIA, from *odon*, *odontos*, 'a tooth,' and *αλγος*, 'pain.' *Dentium dolor*, Toothach, *Odont'ia*, *O. dolorosa*, *Gomphi'asia*, *Gomphi'asmus*, *Odontodyn'ia*, (F.) *Douleur des dents*, *Fluxion sur les dents*, *Mal de dent* or *d'amour*. A disease dependent upon a variety of causes affecting the cavity of the tooth; but generally owing to caries, which exposes the cavity to the action of the air, and to extraneous matters in general. Hence, the treatment consists in plugging the tooth, or destroying the sensibility of the nerve by powerful stimulants; and, if these means fail, in extracting the tooth.

ODONTALGIA CARIOSA, Dental gangrene—o. Dentitionis, Dentition—o. *Hæmodia*, *Hæmodia*.

ODONTALGIA NERVO'SA, *Neuralgia Denta'lis*. Neuralgia of the teeth. Characterized by periodical pain, shooting with the utmost violence along the branches of the fifth pair distributed to the affected jaw.

ODONTALGIA REMITTENS ET INTERMITTENS, Neuralgia infra-orbitaria.

ODONTALGIC, Anti-odontalgic.

ODONTAMBLYOGMUS, *Hæmodia*.

ODONTHEMÖDIA, *Hæmodia*.

ODONTHARPAGA, Dentagra.

ODONTYPERÆSTHESIS, *Hæmodia*.

ODONTIA, Odontalgia.

ODONT'IA DEFOR'MIS. Deformity of the teeth from error of shape, position, or number.—Good.

ODONTIA DENTITIONIS LACTANTIUM, see Dentition (first)—o. Dentitionis puerilis, see Dentition (second)—o. Edentula, see Nefrendes—o. Excrescens, Epulis—o. Incrustans, Odontolithos—o. Stuporis, *Hæmodia*.

ODONTIASIS, Dentition.

ODONTIATER, Dentist.

ODONTIATRIA, Dentistry.

ODONTIC, Anti-odontalgic.

ODONTI'TIS, *Odontophleg'monē*, from *odon*, *odontos*, 'a tooth,' and *itis*, denoting inflammation. Inflammation of the teeth.

ODONTOBOTHRI'TIS, *Odontophatn'itis*, *Inflam'ma t' alveolo'r'um*, from *odontobothrium*, (*odon*, *odontos*, 'a tooth,' and *βοθρίον*, 'alveolus,') and *itis*, denoting inflammation. Inflammation of the alveoli.

ODONTOBOTHRIUM, Alveolus.

ODONTOCLA'SIS, from *odon*, *odontos*, 'a tooth,' and *κλάσις*, 'fracture; *Fractura dent'is*. Fracture of a tooth.

ODONTOCNESIS, Odaxismus.

ODONTOCNESMUS, Odaxismus.

ODONTODES, Odontoid.

ODONTODYNIA, Odontalgia.

ODONTOG'ENY, *Odontogen'ia*, from *odon*, *odontos*, 'a tooth,' and *γενεσις*, 'generation.' Generation or mode of development of the teeth.

ODONTOLYPHON, Denticulipium, Gum lancet.

ODONTOLYPHUM, Denticulipium, Gum lancet.

ODONTOGRAPHY, *Odontograph'ia*, from *odon*, *odontos*, 'a tooth,' and *γραφη*, 'a description.' A description of the teeth.

ODONTOID, *Odontoi'des*, *Odonto'des*, *Dentiform'is*, *Denta'lis*, *Dentoideus*, *Pyrenoi'des*, from *odon*, *odontos*, 'a tooth,' and *ειδός*, 'shape,' 'resemblance.' An epithet given to the *Processus dentatus* of the second vertebra or axis.

ODONTOID LIG'AMENTS, *Ligament'ia ala'ria* seu *Mauchar'ti*, (F.) *Ligaments odontoi'diens*, are two strong and short conical fasciculi, whose truncated summits embrace the sides of the odontoid process, and whose bases are fixed in the fossæ at the inner side of the condyles of the occipital bone. Their direction is obliquely outwards and slightly upwards. They enter into the composition of the occipito-axoid articulation, strengthen the junction of the head with the vertebral column, and limit the movement of rotation of the atlas on the axis.

Other ligamentous fibres pass from the highest portion of the *processus dentatus* to the anterior margin of the foramen magnum of the occipital bone—constituting the *Ligamentum suspensorium dentis*.

ODONTOLITHOS, (F.) *Odontolithe*, from *odon*, *odontos*, 'a tooth,' and *λίθος*, 'a stone.' A sort of incrustation, of a yellowish colour, which forms at the coronæ of the teeth, and is called *Tartar*, *Tartarus Dentium*, *Odont'ia incrustans*, *Calculus denta'lis*, (F.) *Tartre des dents*, *T. dentaire*. It consists of seventy-nine parts of phosphate of lime; twelve and a half of mucus; one of a particular salivary matter, and seven and a half of animal substance, soluble in chlorohydric acid. A species of infusoria, *Dentic'ola hom'inis*, has been found in it.

ODONTOL'OGY, *Odontolog'ia*, from *odon*, *odontos*, 'a tooth,' and *λογος*, 'a discourse.' An anatomical treatise of the teeth.

ODONTONECROSIS, Dental gangrene.

ODONTOPARALLAX'IS, from *odon*, *odontos*, 'a tooth,' and *παρὰλξις*, 'deviation.' Irregularity and obliquity of the teeth; (Prov.) *Snaggle teeth*.

ODONTOPHATNE, Alveolus.

ODONTOPHATNITIS, Odontobothritis.

ODONTOPLEROSIS, see Filling.

ODONTOPHYIA, Dentition.

ODONTOPRISIS, Brygmus, Stridor dentium.

ODONTOSEY'SIS, *Odontoseis'mus*, *Dentium vacillan'tia*, from *odon*, *odontos*, 'a tooth,' and *σεισις*, 'shaking.' Looseness of the teeth.

ODONTOSEISMUS, Odontoseisis.

ODONTOSMEGMA, Dentifrice.

ODONTOSPHACELISIS, Dental gangrene.

ODONTOSPHACELISMUS, Dental gangrene.

ODONTOSTERE'SIS, from *odon*, *odontos*, 'a tooth,' and *στερησις*, 'privation.' Loss of the teeth.

ODONTOSYNERISM'US, from *odon*, *odontos*, 'a tooth,' and *συνερίζειν*, 'to strike together;' (F.) *Claquement*. Chattering of the teeth.

ODONTOTECHNY, Dentistry.

ODONTOTHERAPIA, Dentistry.

ODONTOTRIMMA, Dentifrice.

ODONTOTRYPY, Odontropy.

ODONT'TRYPY, *Odontot'rupy*, *Odontotrup'ia*, from *odon*, *odontos*, 'a tooth,' and *τρύπω*, 'I per-

forate.' The operation of perforating a tooth to evacuate purulent matter confined in the cavity of the pulp.

ODORAMENTUM, Odoriferum.

ODORANT, see Odour.

ODORAT, Olfaction.

ODORATIO, Olfaction.

ODORATUS, Olfaction—o. Deperditus, Anosmia.

ODORIFERUM, from *odor*, *odoris*, 'odour,' and *fero*, 'I carry;' *Odoramentum*. A medicine that gives odour or flavour. A scent. A perfume.

ODOROUS, see Odour.

ODOS, Way.

O'DOUR, *Odor*, *Odmē*, *Osmē*, (F.) *Odeur*. A smell. Odours are subtle particles, constantly escaping from the surface of certain bodies hence called *odorons*, (F.) *Odorant*. They act, in some manner, by actual contact with the nerves of the Schneiderian membrane, and give occasion to the sense of smell or olfaction.

ODOUS, see Teeth.

OD'YNE, *odynē*, 'pain,' *Dolor*, *Odyne'ma*. A very common suffix to words; as in *Pleurodyne*. ODYNEMA, *Odyne*.

OEIOMA'NIA, *Oikeioma'nia*, from *οικιος*, 'belonging to a household or family,' and *mania*. A form of moral insanity, characterized by domestic perversity.

ECONOMIA, Economy — œ. Animalis, Physiology.

EDALICUS, *Edematous*.

EDALIUS, *Edematous*.

EDE'MA, from *οίδημα*, 'I am swollen.' *Hydron'cus*, *Edemat'ia*, *Hyderon'cus*, *Hydaton'cus*, *Hydrocæ'dema*, *Hydre'dema*, *Hydrops cellula'ris ar'tuum*, *Phlegmat'ia*, *Leucophlegma'tia*, *Ecphy'ma edemat'icum*, (F.) *Edème*, *Edématic*, *Enflure*. Swelling produced by the accumulation of a serous fluid in the interstices of the areolar texture. This swelling is soft; yields under the finger; preserves the impression for some time, and is pale and without pain. It presents the same characters as anasarca, which is general œdema. Its etiology and treatment are also the same. See Anasarca.

EDEMA ACUTUM, see Anasarca.

EDEMA ARSENICA'LIS. The swelling of the eyelids and face, induced by continued use of the preparations of arsenic.

EDEMA CALIDUM, see Anasarca — œ. Capitis, *Hydrocephalus externus*.

EDE'MA CER'E'BRI, (F.) *Edème du cerveau*. Edema of the brain. A condition of the cerebral pulp, in which there is an infiltration of serous fluid into it, so that it appears more moist or watery than common; and, when sliced or pressed, small drops of water are seen to ooze out.

EDEMA, COMPACT, Induration of the cellular tissue — œ. Cruentum, Suggillation — œ. Febrile, see Anasarca — œ. Fugax, Anathymiasis — œ. Hystericum, Anathymiasis.

EDEMA OF THE GLOTTIS, *Æ. glottidis*, *Edem'atous Laryngi'tis* or *angina*, *Laryngi'tis submuco'sa* seu *edemat'ia* seu *sero-purulen'ta*, *Hydrops glott'idis*, *Angi'na aquo'sa* seu *larynge'a edemat'ia*, *Submu'cous Laryngi'tis*, (F.) *Laryngite adémateuse*, *L. adémateuse et séro-purulente ou sus-glottique ou sous-muqueuse*, *Angine laryngée adémateuse*, *Edème de la glotte*. A disease consisting of serous or sero-purulent infiltration into the submucous tissue of the glottis. The symptoms resemble those of croup; but the disease attacks the adult rather than the child. The age is, indeed, a principal means of diagnosis between the two affections.

The disease is almost always fatal. The treatment has to vary according to the accompanying general symptoms.

EDEMA LACTEUM, *Phlegmatia dolens*.

EDEMA OF THE LUNGS, *Æ. pulmo'num*, *Pneumoch'ysis*, *Pneumonœ'dema*, *Hydrops seu anasar'ca Pulmo'num*, *Hydropneumo'nia*, (F.) *Edème du pousmon*. Laënnec has so called the infiltration of serum into the tissue of the lung, carried to such an extent as to diminish its permeability to air. It is not an uncommon sequela of pneumonia, and the major exanthemata. The respiration is laborious; the respiratory murmur scarcely perceptible, although the thorax is largely expanded, and there is a slight *râle crépitant*, particularly at the base and inferior part of the lung. The sound on percussion is clear, and on both sides equally so. The cough is attended with aqueous expectoration. In some cases the respiration becomes purile in a small portion of the summit of the lung.

EDEMA NEONATORUM, Induration of the cellular tissue — œ. Palpebrarum, *Hydroblepharon* — œ. Puerperarum, *Phlegmatia dolens* — œ. Pulmonum, *Edema of the lungs* — œ. Pulmonum chronicum, *Hydropneumonia* — œ. Scroti aquosum, *Oschydrœdema* — œ. Scroti cruentum, *Oschæmatœdema* — œ. Scroti purulentum, *Oscheopyœdema* — œ. Scroti urinosum, *Urocele* — œ. Spasticum, *Anathymiasis* — œ. Uvule, *Staphylœdema*.

EDEMATIA, *Edema*.

EDEMATIE, *Edema*.

EDEMATOSARCA, *Edemosarca*.

EDEMATOSCHEOCE'LE, *Edemoscheoce'le*, *Oscheoce'le* *edemat'ica*. *Oscheocele* with œdema of the scrotum.

EDEMATOUS, *Edemat'ous*, *Edemat'icus*, *Edemat'os*, *Edal'ius*, *Edal'icus*, *Edem'atous*, (F.) *Edémateur*. Affected with œdema, (F.) *Edématique*, of the nature of œdema.

EDE'ME, *Edema* — œ. *du Cerveau*, *Edema cerebri* — œ. *Douloureux des femmes en couche*, *Phlegmatia dolens* — œ. *des Nouvelles accouchées*, *Phlegmatia dolens* — œ. *actif des Nouvelles accouchées*, *Phlegmatia dolens* — œ. *de la Glotte*, *Edema of the glottis* — œ. *de la Glotte*, *Angina edematosa* — œ. *du Tissu cellulaire des nouveaunés*, *Induration of the cellular tissue*.

EDEMOSAR'CA, *Edematosar'ca*. A species of tumour mentioned by M. A. Severinus, which holds a middle place between œdema and sarcoma.

EDEMOSCHEOCELE, *Edemoscheocele*.

EIL, Eye — œ. *de Bonf*, *Anthemis tinctoria* — œ. *Double*, *Binoculus* — œ. *de Lièvre*, *Lagophthalmus* — œ. *Simple*, see *Monoculus*.

EILLET GIROFLÉE, *Dianthus caryophyllus*.

EILLIÈRE, *Scaphium oculare*.

ENAN'THÉ, (*Æ. crocata* seu *chærophy'llis foliis*, *Hemlock dropwort*, *Hemlock water-dropwort*. *Nat. Ord. Umbelliferae*. *Sex. Syst. Pentandria Digynia*, from *οivos*, 'wine,' and *ανθος*, 'a flower;' its flowers smelling like the vine. A violent poison of the acro-narcotic class. Its juice has been recommended in certain cutaneous diseases; but it is scarcely ever used. It is employed in fomentations.

ENANTHE AQUATICA, *Phellandrium aquaticum* — œ. *Phellandrium*, *Phellandrium aquaticum* — œ. *Striata rigida*, *Seseli tortuosum*.

ENELE'ON, from *οivos*, 'wine,' and *ελαιον*, 'oil.' A mixture of oil and wine.—Galen. Also, rectified spirit or alcohol.

ENOG'ALA, *οινογαλα*, from *οivos*, 'wine,' and *γαλα*, 'milk,' — 'a mixture of wine and milk.' According to some, wine as warm as new milk.

ENOG'ARON, *Garum vino mistum*. Wine mixed with garum.—Ætius.

ENOLATURE, see Vinum.

ENOLÉ, see Vinum.

ENOLOTIF (F.). A name given by M

Béral to a medicine intended for external use, whose excipient is wine.

CENOIDES, Vinous.

CENOMANIA, *Cenom'nia*, from *oivos*, 'wine,' and *mania*. Delirium tremens. Properly, wine-mania. An insatiable desire for intoxicating liquors.

CENOMEL, *Cenom'eli*, from *oivos*, 'wine,' and *mel*, 'honey.' *Mel vino'sum*, *Mulsam Vино'sum*. Honey wine. Wine mead. See Hydromel.

CENOPHLYGIA, Temulentia.

CENOPHLYXIS, Temulentia.

CENOS, Wine.

CENOSTAGMA, *Spiritus vini rectificatus*.

CENOTHE'RA BIEN'NIS, *Æ. Mollis'sima seu muricata seu gaurōides*, *On'agra*, Evening Primrose, Tree Primrose, Primrose tree, Cure-all, *Scabish*. An indigenous plant, common on the borders of fields, and in natural hedges. *Ord.* Onagraceæ. *Ser. Syst.* Octandria Monogynia. Its properties are mucilaginous and slightly acid. A decoction has been used in cases of infantile and other eruptions.

CENOTHE'RA GAUROIDES, *O. biennis* — *æ. Muricata*, *O. biennis*.

CESOPHAGE'AL, *Cesophagæ'us*, (F.) *Cesophagien*. Same etymon as *oesophagus*. Relating or belonging to the *oesophagus*.

CESOPHAGEAL AP'ERTURE OF THE DI'APHRAGM, *Fora'men cesophagæ'um*, (F.) *Ouverture cesophagienne du diaphragme*. An opening in the diaphragm for the passage of the *oesophagus*.

CESOPHAGEAL AP'ERTURE OF THE STOMACH. A name given to the superior or cardiac orifice of the stomach, to distinguish it from the inferior or pyloric. See Cardia.

CESOPHAGEAL MUSCLE, *Cesophagæ'us*. Some anatomists have given this name to the transverse muscular fibres which surround the *oesophagus* at its upper extremity.

CESOPHAGEAL TUBE, see Tube, *oesophageal*.

CESOPHAGEURYS'MA, *Cesoph'agus succenturiatus*, from *oesophagos*, 'the *oesophagus*,' and *eurysma*, 'dilatation.' Dilatation of the *oesophagus*.

CESOPHAGIALGIA, *Cesophagismus*.

CESOPHAGIARCTIA, *Dysphagia constricta*.

CESOPHAGIEN, (muscle), *Constrictor cesophagi*.

CESOPHAGIS'MUS, *Cesophagis'mum*. A name given by Vogel to spasm of the *oesophagus*, *Dysphagia Spasmodica seu Spas'tica seu Nervo'sa*, *Strictura cesoph'agi spasmodica*, *Cesophagial'gia*, *Cesophagospas'mus*, *Tenes'mus Gula*, (F.) *Ténésie de l'oesophage*. By some used synonymously with *oesophagitis*, *dysphagia*, and contraction of the *oesophagus*.

CESOPHAGITIS, *Angi'na seu Cyman'chè cesophagæ'a*, *Dysph'gia inflammatoria*, *Inflammatio cesoph'agi seu Gula*, (F.) *Angine cesophagienne*. Inflammation of the *oesophagus*.

CESOPHAGODYN'IA, from *oesophagos*, 'the *oesophagus*,' and *odyn*, 'pain.' Pain in the *oesophagus*.

CESOPHAGOPATHI'A, *Morbus Cesoph'agi*, from *oesophagos*, 'the *oesophagus*,' and *pathos*, 'disease.' A morbid condition of the *oesophagus*.

CESOPHAGOPLE'GIA, *Dysph'gia paralytica*, *Læmoparaly'sis*, from *oesophagos*, 'the *oesophagus*,' and *πληγη*, 'a stroke.' Paralysis of the *oesophagus*.

CESOPHAGORRHAG'IA, from *oesophagos*, 'the *oesophagus*,' and *ραιη*, 'a rupture.' Hemorrhage from the *oesophagus*.

CESOPHAGORRHE'E'A, from *oesophagos*, 'the *oesophagus*,' and *ρρω*, 'I flow.' Discharge of blood or mucus from the *oesophagus*.

CESOPHAGOSPAS'MUS, from *oesophagos*, 'the

oesophagus,' and *σπασμος*, 'spasm.' Spasm of the *oesophagus*.

CESOPHAGOT'OMY, *Cesophagotomia*, from *oesophagos*, 'the *oesophagus*,' and *τεμνω*, 'to cut.' An incision made into the *oesophagus* for the purpose of extracting foreign bodies from it.

CESOPH'AGUS, from *oio*, 'future' *oio*us, 'I carry,' and *φαγω*, 'I eat'; *Gula*, *Fistula cibalis*, *Fistula seu Infundib'ulum Ventriculi*, *Via stom'achi et Ventris*, *Gluttus*. The Gullet, (Prov.) *Wizen*. A musculo-membranous canal, cylindrical, and depressed from before to behind, which extends from the inferior extremity of the pharynx to the upper orifice of the stomach. At its origin, it is situate in the median line; but beneath the larynx, it deviates to the left, and in the chest experiences different reflections. In its cervical portion, it corresponds, behind, with the spine; before, with the larynx and trachea; and, at the sides, it is close to the primitive carotids, internal jugular veins, par vagum, recurrent nerves, &c. In its inferior or thoracic portion, the *oesophagus* is entirely contained in the posterior mediastinum; and enters the abdomen through the *oesophageal* aperture, *Fora'men cesophagæ'um*, of the diaphragm. The *oesophagus* is composed of a very strong muscular layer, sometimes called *Tunica vaginalis gula*; formed, itself, of two sets of fibres, the external being general longitudinal, the internal transverse or annular. 2. Of a mucous membrane which is soft, fine, thin, and white, especially at its lower part. It is continuous, above, with the mucous membrane of the pharynx. The mucous follicles, found beneath it, are not numerous, and have been called *Cesophageal glands*. The arteries of the *oesophagus* proceed, in the neck, from the thyroid; in the chest, from the bronchial arteries and directly from the aorta;—in the abdomen, from the inferior phrenic, and coronaria ventriculi. Its veins empty themselves into the inferior thyroid, the vena cava superior, the internal mammary, azygos, bronchial, phrenic, and coronaria ventriculi. Its lymphatics pass into the ganglia surrounding it. Its nerves are afforded by the pharyngeal and pulmonary plexuses; by the cardiac nerves; the thoracic ganglia, and, especially, by the pneumogastrics and their recurrent branches.

CESOPHAGUS SUCCENTURIATUS, Pharyngocele.

CESTROMANIA, Nymphomania.

CESTRUATION, Orgasm, see Heat.

CESTRUM, Clitoris.

CESTRUM VENERIS, *Æ. vene'reum*, *Cestrus vene'reus*; from *oio*stros, *cestrus*, 'a violent impulse or desire.' A vehement desire for sexual intercourse. Orgasm. With some, *cestrum* signifies Clitoris.

CESTRUS, Clitoris, see Ectozoa—*æ. Venerens*, *Cestrum venerem*.

CE'SYPU, from *ois*, 'a sheep.' The greasy matter of unwashed wool; formerly employed in friction in diseased joints.

CE'US, Ova.

OFFICE, PHYSICIAN'S or SURGEON'S, Iatration.

OFFICINA, Pharmacopolium.

OFFIC'INAL, *Officina'lia*, from *officina*, for *opificina*, (from *opifex* (opus and facio,) 'a workman,') 'a shop.' An epithet for medicines found in the shop of the apothecary, ready prepared—*usua'lia*; in opposition to *magistral* or *extemporaneous*,—those prepared after the prescription of the physician.

OFFIUM, Affion, Opium.

OFFSPRING, Epigone.

OFFUSCATIO, Amaurosis.

OGLA, Oozala.

OGNON, a Corn.—o. *Marin*, Scilla.

OHIO, MINERAL WATERS OF. *Yellow Spring* is a chalybeate, situate in Greene county, 64 miles from Cincinnati. It is somewhat frequented.

OIDIUM ABORTIFACIENS, see Ergot.—o. Albicans, Aphthaphyte.—o. Schoenleini, see Porrigo favosa.

OIE, Sorbus domestica.

OIGNON, Allium cepa.

OIKEIOMANIA, Œciomania.

OIL, *O'leum*, *Elæ'ôn*, *élaov*, (F.) *Huile*; from *êla*, 'the olive.' A collective name, under which two classes of fluids are included, very different from each other: those belonging to the one class, are viscid, mawkish or almost insipid; those of the other are nearly devoid of viscosity, and are caustic and very volatile. The former are called *fat* or *fixed oils*; the latter *volatile* or *essential oils* or *essences*.

OIL, see Ricinus communis.

OIL OF ALMONDS, *Oleum amygdalarum*—o. of bitter Almonds, *Oleum amygdalæ amaræ*—o. of Amber, rectified, see Succinum—o. of Anda, see Anda—o. Animal, *Oleum animale*—o. Animal, of Dippel, *Oleum animale Dippelii*—o. of Bay, *Daphnêlæon*—o. of Bays, *Oleum laurinum*—o. and Beeswax, *Unguentum ceræ*—o. of Ben, see Guilandina moringa—o. of Benjamin or Benzoin, *Oleum benzoini*—o. Benne, see Sesamum orientale—o. of Bergamot, see Bergamot.

OIL, BRITISH. An empirical preparation often used in cases of sprains. A committee of the Philadelphia College of Pharmacy recommended the following form for its preparation.—*Ol. Terebinth.*, *Ol. Lini usitat.* 33 f 3ij. *Ol. Succini*, *Ol. Juniper.* 33 f 3iv, *Petrol. Barbada.* f 3ij, *Petrol. American.* (*Seneca Oil.*) f 3j. M.

OIL OF CACAO, Butter of Cacao—o. of Cade, see Juniperus Oxycedrus—o. Carron, Linimentum aquæ calcis—o. of the Cashew nut, see Anacardium occidentale—o. Castor, see Ricinus communis—o. of Chabert, *Oleum animale empyreumaticum Chaberti*—o. of Cinnamon, see Laurus cinnamomum—o. of Cloves, *Eugenia caryophyllata*—o. Cod, *O. Jecoris aselli*—o. Codliver, *O. Jecoris aselli*—o. of Copiba, *Oleum Copibaæ*—o. Cornspirit, Oil, fusel—o. of Cubebs, see Piper cubeba—o. of Cypress, *Cyprium oleum*—o. of Dill, see Anethum graveolens—o. Dippel's, *Oleum animale Dippelii*—o. of Egg, see Ovum—o. Ethereal, *Oleum Æthereum*—o. Flaxseed, see Linum usitatissimum.

OIL, FUSEL, *Al'cohol amylicum* (Ph. D.), *Fousel oil*, *Pota'to oil*, *Grain oil*, *Cornspirit oil*, *Pota'to spirit oil*, *Oil of potatoes*, *Amylic al'cohol*, *Hydrated oxide of amyle*, ((G.) *Fusel*, an epithet for poor brandy.) An acrid volatile oil, formed in the manufacture of potato brandy, and which is not easily separable from it. Its chemical constitution is analogous to that of alcohol. It exhales a powerful and peculiarly suffocating odour. S. g. 318. In small doses it is highly stimulating,—acting like narcotics in general. In large doses it destroys the mucous membrane of the stomach. It is used by the Dublin College in the preparation of Sodæ valerianas.

OIL, GRAIN, Oil, fusel.

OIL, HAERLEM. An empirical preparation supposed to consist chiefly of petroleum, turpentine, and balsam of sulphur. Used internally in renal and rheumatic affections.

OIL, KRUMHOLTZ, see Pinus mughos—o. of Lemons, see Citrus medica—o. Lily, see Lilium candidum—o. of Mace, expressed, see Myristica—o. of Mastich, see Pistacia Lentiscus—o. of Mucilages, *Oleum e mucilagibus*—o. Neats-foot, *Oleum bubulum*—o. of Nicodemus, Nico-

demi oleum—o. Nut, *Juglans cinerea*—o. of Nutmegs, see Myristica moschata—o. Olive, *Oleum olivæ*—o. Palm, see Cocos butyracea—o. Paper, *Pyrothonide*—o. Parsley, *Apiol*—o. Pennyroyal, see Hedeoma pulegioides—o. of Pike, *Oleum lucii piscis*—o. of Potatoes, Oil, fusel—o. Potato, Oil, fusel—o. Potato spirit, Oil, fusel—o. Rag, *Pyrothonide*—o. Red, see Hypericum perforatum—o. Rock, *Petroleum*—o. of Rose Geranium, *Oleum graminis Indici*—o. of Roses, see Rosa centifolia—o. of Rue, *Peganeleon*, see Ruta—o. Skateler, see *Oleum Jecoris aselli*—o. of Spike, *Oleum terebinthinæ*—o. of St. John's Wort, see Hypericum perforatum—o. Salad, *Oleum olivæ*—o. of Scorpion, see Scorpion—o. Seneca, *Petroleum*, New York—o. of Spike, see Lavandula spica—o. of Spikenard, *Oleum graminis Indici*—o. Sulphuretted, *Balsamum sulphuris simplex*—o. of Tartar, *Liquor potassæ subcarbonatis*—o. of Tobacco, *Oleum tabaci*—o. of Turpentine, *Oleum terebinthinæ*—o. of Turpentine, rectified, *Oleum terebinthinæ rectificatum*—o. of Valerian, *Oleum valerianæ*—o. of Vitriol, *Sulphuric acid*—o. of Wine, *Oleum æthereum*.

OILS, ANIMAL, *Olea animalia*—o. Distilled, *Olea volatilæ*—o. Empyreumatic. *Olea empyreumatica*—o. Essential, *Olea volatilæ*—o. Ethereal, *Olea volatilæ*—o. Expressed, *Olea fixæ*—o. Fatty, *Olea fixæ*—o. Fixed, *Olea fixæ*—o. Fugacious, *Olea fugaciæ*—o. Medicinal, *Olea medicinalia*—o. Volatile, *Olea volatilæ*.

OILY, Oleaginous—o. Grain, *Sesamum orientale*.

OINOMANIA, Œnomania.

OINOS KRITHES, Cerevisia.

OINTMENT, Unguentum—o. Antimonial, *Unguentum antimonii*—o. of Antimony, tartarized, *Unguentum antimonii*.

OINTMENT, ARSENICAL, of Sir Astley Cooper. This is made of *arsenious acid* 3i: sulphur, 3i: spermaceti cerate, 3j. It is spread on lint, and applied to cancerous sores.

OINTMENT, BASILICON, *Unguentum resinæ*—o. Bay, *Unguentum laurinum*—o. of Belladonna, *Unguentum belladonnæ*—o. Blister, *Cerate of cantharides*—o. Blistering, green, *Unguentum lyttæ medicatum*—o. Blistering, milder, *Unguentum lyttæ*—o. Blistering, yellow, *Unguentum lyttæ medicatum aliud*—o. Blue, *Unguentum hydrargyri*—o. Citrine, *Unguentum hydrargyri nitratis*—o. of Cocculus Indicus, *Unguentum cocculi*—o. of Creasote, *Unguentum creasoti*—o. Cucumber, see Cucumis sativus—o. Cyrillo's, *Unguentum muratis hydrargyri oxygenati medicatum*—o. Digestive, simple, *Unguentum digestivum simplex*—o. Edinburgh, see *Unguentum veratri*—o. Elder, *Unguentum sambuci*—o. of Elemi, *Unguentum elemi compositum*—o. for the Eyes, (Smellome's), see *Ceratum resinæ*—o. of Galls, *Unguentum gallæ*—o. of Galls, compound, *Unguentum gallæ compositum*—o. Garlic, see Allium—o. of Geranium, *Oleum graminis Indici*—o. Gingergrass of Nemauro, *Oleum graminis Indici*—o. Golden, *Unguentum hydrargyri nitrato-oxidi*—o. Golden, Singleton's, see Singleton's golden ointment—o. Goulard's, *Ceratum plumbi compositum*—o. Grass of Nemauro, *Oleum graminis Indici*—o. Green, *Unguentum sambuci*—o. Hellebore, white, *Unguentum veratri*—o. of Hemlock, *Unguentum conii*—o. of Hydriodate of Potass, *Unguentum potassæ hydriodatis*—o. of Iodide of Lead, *Unguentum plumbi iodidi*—o. of Iodide of Mercury, *Unguentum hydrargyri iodidi*—o. of red Iodide of Mercury, *Unguentum hydrargyri iodidi rubri*—o. of Iodide of Sulphur, *Unguentum sulphuris iodidi*—o. of Iodine, *Unguentum Iodini*—o. of Iodine, compound, *Unguentum Iodini compositum*—o. Issue, Dr. Physick's, see *Unguentum lyttæ medicatum aliud*—o. Itch,

Unguentum sulphuris compositum—o. Itch, Bailey's, see Unguentum sulphuratum ad scabiem—o. Itch, Bateman's, see Unguentum sulphuratum alcalinum ad scabiem—o. Itch, Helmerick's, Unguentum sulphuratum alcalinum ad scabiem—o. Itch, Jackson's, Unguentum sulphuris compositum—o. Lead, compound, Unguentum plumbi compositum—o. of white oxide of Lead, Unguentum oxidi plumbi albi—o. of Lydia, Bacaris—o. Marshmallow, Unguentum de althæa—o. Mercurial, Unguentum hydrargyri—o. Mercurial, milder, see Unguentum hydrargyri—o. Mercurial, strong, see Unguentum hydrargyri—o. of nitrate of Mercury, Unguentum hydrargyri nitratis—o. of gray oxide of Mercury, Unguentum oxidi hydrargyri cinerei—o. of nitric oxyd of Mercury, Unguentum hydrargyri nitrice-oxydi—o. of Mezeron, Unguentum mezerei—o. of Nitrous acid, Unguentum acidi nitrosi—o. of Pitch, Unguentum resinæ nigræ—o. of Pitch, black, Unguentum resinæ nigræ—o. Plunkett's, for cancer, Plunkett's ointment—o. red Precipitate, Unguentum hydrargyri nitrice-oxydi—o. of white Precipitate, Unguentum hydrargyri præcipitati albi—o. Resin, black, Unguentum resinæ flavæ—o. Resin, black, Unguentum resinæ nigræ—o. of Rosewater, Unguentum aquæ rosæ—o. Soldier's, Martiatum unguentum—o. of Spanish flies, Cerate of Cantharides, Unguentum lyttæ—o. Spermaceti, Unguentum cetacei—o. of Stramonium, Unguentum stramonii—o. of Subacetate of copper, Unguentum subacetatis cupri—o. of Sugar of lead, Unguentum plumbi superacetatis—o. Sulphur, Unguentum sulphuris—o. Sulphur, compound, Unguentum sulphuris compositum—o. Tar, Unguentum picis liquidæ—o. Tartar emetic, Unguentum antimonii—o. Tobacco, Unguentum Tabaci—o. Tutty, Unguentum oxidi zinci impuri—o. Verdigris, Unguentum subacetatis cupri—o. Wax, Unguentum ceræ—o. White, Unguentum oxidi plumbi albi—o. Zinc, Unguentum zinci—o. of impure oxide of Zinc, Unguentum oxidi zinci impuri.

OIOCALYM'MA, *Oiocalyptum*; from *ovov*, 'egg,' and *καλυπτειν*, 'to cover,' *Membra'na O'vuli corticalis*. The membrane of the egg-shell.

OKRA, *Hibiscus esculentus*.

OLD AGE, *Senectus*.

OLD MAN, *Artemisia abrotanum*.

OLD WIFE'S SHIRT, *Liriodendron*.

OLEA ANIMA'LIA, *Animal oils*, (F.) *Huiles animales*. A name given to fixed oils holding in solution the mucilaginous and gelatinous principles of certain animals, as the *Oil of frogs*, *Oil of scorpions*, *Oil of spiders*, &c. Sometimes, also, the term *animal oils* is given to empyreumatic oils, produced during the decomposition of animal substances by heat.

OLEA DESTILLATA, O. Volatilia.

OLEA EMPYREUMAT'ICA, *Empyreumatic oils*, (F.) *Huiles empyreumatiques ou pyrogénées, Pyrélainées*. Oils which have an empyreumatic or burnt smell. They are obtained by treating vegetable or animal matters by heat, in close vessels. They do not exist in organized bodies, but are formed during their decomposition by fire. The animal oil of Dippel is an empyreumatic oil.

OLEA EUROPE'Æ, O. *antiva seu lancifolia seu polymorpha seu Gallica*, *Oli'va*, *Olive tree*, *oliva*, *Ela'ea*, (F.) *Olivier*; (Fruit) *Olive*. Nat. Ord. *Jasminæ*. Sex. Syst. *Diandria Monogynia*. The leaves of the olive are bitter, and an extract prepared from them, and a substance called *olivi'na*, the bitter principle of the leaves, have been given in Italy as antiperiodics. The fruit, when pickled, is extremely grateful to some stomachs. Olives, as met with in the shops, are prepared from the green, unripe fruit, repeatedly

steeped in water. To this, some quicklime or alkaline salt is added, and, afterwards, they are washed and preserved in a pickle of common salt and water. From this fruit is prepared the *Olive oil* or *Salad oil* of the *Pharmacopœias*, which is obtained by grinding and pressing the olives, when thoroughly ripe. The finer and purer oil issues first by gentle pressure, and the inferior sort on heating what is left, and pressing it more strongly. See *Oleum Olivæ*. In Calabria, an odorous resin exudes from its trunk, which is employed as a perfume by the Neapolitans.

A gum flows from certain wild olives, in warm countries, which consists, according to Pelletier, of a resin, a little benzoic acid, and a peculiar substance, called *Olivile*. It is in yellow masses, of a slightly acid taste, and of a vanilla smell. It is called *Olea gummi*, *O. resin'na*, and *O. bals'anum*, (F.) *Gomme olivier*, and is esteemed astringent and detersive.

OLEA FIXA SEU PIN'GUIA, *Expressed, Fixed or Fatty oils*, (F.) *Huiles fixes ou grasses, Oléols*. All the oils obtained from the seeds or pericarps of vegetables, without distillation, and which are viscid, but slightly odorous and sapid; lighter than water, and insoluble in alcohol. The rancidity of oils depends on the absorption of oxygen, and therefore they should be kept in bulk as much as possible, and in narrow-necked bottles, so that a very small surface only can be exposed to the air. All the fixed oils are emollient, and, in a certain dose, they act as purgatives and emetics. They are prepared by expressing the fruit or seed containing them.

OLEA FUGA'CIA, *Fuga'cious oils*, (F.) *Huiles fugaces*. A name given to oils which are so volatile that, in order to obtain them, recourse must be had to a different process from that employed for other essential oils. Such are the oils of jessamine, lily, violet, &c.

OLEA GALLICA, O. *Europæa*—o. *Lancifolia*, O. *Europæa*.

OLEA MEDICINA'LIA, *Medic'inal oils*, (F.) *Huiles Médicinales, Etléols, Solution par les huiles*. A name given to oils prepared by macerating, infusing, or boiling medicinal substances in olive or any other fixed oils. These oils may then be regarded as oily solutions of certain medicinal substances; whence they can never be simple. They have, however, been divided into *simple and compound medicinal oils*. To the former belong the *Oils of St. John's wort*, of the *Solanum nigrum*, &c.; to the other—which have often been called *Oily balsams*, (F.) *Baumes huileux*—the *Balsams of Fioraventi*, Metz, &c. Medicinal oils are, almost always, employed externally.

OLEA POLYMORPHA, O. *Europæa*—o. *Sativa*, O. *Europæa*.

OLEA VOLATIL'IA, O. *destilla'ta*, *Æthero'lea*, *Volatile*, *Ethe'real*, *Essential*, or *Distilled oils*, *Es'sences*, (F.) *Huiles volatiles ou essentielles ou distillées ou éthérées, oléolates, oléoles, oléols*. Oils found in aromatic vegetables, and in every part of them, except in the interior of the seeds. The majority are obtained by distillation; but some by expression. They possess unctuousity, inflammability, and viscosity, like the fixed oils; but they are generally odoriferous, pungent, and acid. The greater part are lighter than water; but some are heavier, and congeal at a moderate temperature. They dissolve, in small quantity, in distilled water, by simple agitation. Almost all are soluble in alcohol. The odour and taste are the usual tests of their goodness. To preserve them, they should be kept in a cool place, in small bottles, quite full and well corked. Volatile oils are possessed of the aromatic properties of the plants whence they are obtained. They

are all, when applied externally, stimulant and rubefacient. Pharmaceutical preparations, in which substances are dissolved in volatile oils, (F.) *Solutions par les huiles volatiles*, are termed by Béal, *Myrolés*.

OLEAGINOUS, *Oleaginosus*, *Oleo'sus*, (F.) *Oléagineux*; from *oleum*, 'oil.' Oily; containing oil,—as 'an oleaginous or oily mixture.'

OLEA'MEN, *Oleamen'tum*. Same etymon. Any soft ointment prepared of oil.—Scribonius.

OLEANDER, *Rhododendron chrysanthemum*.

OLECRANARTHRTIS, from *olecranon*, ὀλκρον, 'joint,' and *itis*, denoting inflammation. Inflammation of the elbow joint.

OLECRANARTHROACÉ, from ὀλκρον, 'the olecranon,' ἄρθρον, 'articulation,' and *kakia*, 'vice or disease.' Inflammation of the articular surfaces of the elbow.—Rust.

OLECRAN'IAL, (F.) *Olécranién*. Relating or belonging to the olecranon.

OLECRANOID CAVITY, see *Ulna*.

OLECRANON, *Olecranon*, *Olecr'num*, *Olecranon mobilis*; from ὀλκρον, 'the elbow,' and κρηνον, 'the head.' *Acrole'nion*, *Additamen'tum nectat'um*, *Awon*, *Proces'sus awone'us seu uncinat'us*, *Glans seu Coro'na seu Coro'na posterior seu Additamen'tum uncat'um Ulnæ*, *Vertex Cu'biti*, *Patel'la fica*, *Rost'rum exter'num seu poste'rius*, *Top of the cubit*. Head or protection of the elbow. A large process at the upper extremity of the ulna, on which we lean. When this process is fractured, it is apt to be drawn up by the triceps, and much care is required to keep the parts in apposition.

OLEFIANT GAS, **CHLORIDE OF**, see *Anæsthetic*.

OLEIN, see *Fat*—o. Codliver, see *Oleum jecoris aselli*.

OLENE, *Ulna*.

OLEO-CERATUM, *Cerate*—o. c. *Aquâ subac'tum*, *Ceratum Galeni*.

OLÉO-CÉROLÉ, *Cerate*.

OLÉO-CÉROLÉ RÉSINEUX, *Unguentum*.

OLÉOLATS, *Olea volatil'ia*.

OLÉOLÉS, *Olea volatil'ia*.

OLÉOLS, *Olea fixa*.

OLÉO-SACCHAROLÉ, *Elæosaccharum*.

OLEOSACCHARUM, *Elæosaccharum*.

OLEOSUS, *Oleaginous*.

OLERA, see *Lachanon*.

OLETTE, **MINERAL WATERS OF**. At Olette, in the department Pyrénées Orientales, is a thermal spring, which raises the thermometer to 190° Fahr. It is the hottest in France.

OLÉULES, *Olea volatil'ia*.

OLEUM, Oil—o. *Abietis*, see *Pinus picea*—o. *Absinthii*, *Artemisia absinthium*, (oil of)—o. *Acusticum*, *Huile acoustique*.

OLEUM ÆTHE'REUM, *Spir'itus æ'theris oleo'sus*, *Æ'ther'cal oil* (formed in the distillation of ether), *Oleum vini*, Oil of wine, *Heavy oil of wine*, *Sweet oil of vit'riol*, (F.) *Huile douce de vin*. A peculiar oleaginous matter, obtained by continuing the distillation, after the whole of the sulphuric ether has passed over in the process for the preparation of the latter. It is used only as an ingredient in the compound spirit of ether. It is official in the Ph. U. S.

OLEUM AMMONIATUM, *Linimentum ammoniac fortius*.

OLEUM AMYGDALÉ AMA'RÆ, *Oil of Bitter Almonds*. An oil, obtained by distilling with water the kernels or the fruit of *Amygdalus communis*—variety, *Amara*. Its action is like that

of hydrocyanic acid—four drops having destroyed a dog. Dose, a quarter of a drop to a drop. A drop to a fluidounce of water forms a wash, which has been used in Prurigo senilis, and other forms of itching.

OLEUM AMYGDALA'RUM, *Oleum amygd'alæ*, *Oil of Almonds*, (F.) *Huile d'amandes*. (Expressed from both sweet and bitter almonds,—*Amygdalus communis*.) 3xvj of almonds yield 3v of oil. It is inodorous, insipid, and of a pale straw colour, and is employed as a demulcent and emollient.

OLEUM ANACARDII, see *Anacardium occidentale*.

OLEUM ANIMAL'É, *Animal oil*, (F.) *Huile animale*. An oil obtained by the decomposition of the immediate principles of animals, subjected to the action of heat. It is fetid, and always contains a certain quantity of subcarbonate of ammonia. See *Olea Empyreumatica*. The name *animal oil* is sometimes also given to the fat contained in the fatty vesicles. The composition of this fat does not, indeed, differ from that of the fixed oil.

OLEUM ANIMAL'É DIPPE'LII, *O. cornu cervi*, *O. C. rectificat'um*, *O. pyro-animal'è depur'atum seu animal'è æ'ther'eum*, *Pyro'leum os'sium rectificat'um*, *Animal oil*, *Animal oil of Dippel*, (F.) *Huile animale de Dippel ou de corne de cerf*, is obtained by distilling animal matters, especially hartshorn, on the naked fire. The subcarbonate of ammonia, which it contains, renders it partly soluble in water, and communicates to it the stimulant properties for which it is used in medicine. It is employed as an antispasmodic.

OLEUM ANIMAL'É EMPYREUMAT'ICUM CHABERTI, *O. Empyreumaticum seu anthelminticum seu contra tæ'niam Chaberti*, *Empyreumatic oil of Chabert*, Oil of Chabert, (F.) *Huile anthelmintique de Chabert*, is made by adding one part of animal oil to three parts of oil of turpentine, leaving them to combine for four days, and then distilling three parts. An effective anthelmintic. Dose, a teaspoonful three times a day.

OLEUM ANIMALIZA'TUM PER INFUSIO'NEM, (F.) *Huile animalisée par infusion*, *H. aromatique ou de petits chiens*. A preparation, formerly esteemed tonic and cephalic. It was obtained by boiling new-born puppies in oil, first depriving them of their blood, skin, and intestines. When the decoction was cold, origanum, thyme, pennyroyal, St. John's wort, and marjoram were added.

OLEUM ANISI, see *Pimpinella anisum*—o. *Anthelminticum Chaberti*, *Oleum animale empyreumaticum Chaberti*—o. *Anthemidis*, see *Anthemis nobilis*—o. *Aurantii*, see *Citrus aurantium*—o. *Balaninum*, *Guilandina moringa* (oleum)—o. *Balsami*, see *Myrris opobalsamum*.

OLEUM BENZO'INI, *Oil of Benzoin or Benjamin*. An oil obtained by heating, in a sand-bath, the matter which remains after benzoic acid has been separated from benzoin by the aid of heat. It has been regarded as balsamic and sudorific.

OLEUM BERGAMI, see *Bergamot*.

OLEUM BU'BULUM, *Neat's-foot oil*, (F.) *Huile de pieds de bœuf*. It is obtained by boiling in water for a long time the feet of the beef—*Bos domesticus*—previously deprived of the hoof. It is introduced into the official list of the Ph. U. S. as an ingredient of the ointment of nitrate of mercury.

OLEUM CACAO SPISSATUM, *Butter of Cacao*—o. *Cadinum*, see *Juniperus oxycedrus*—o. *Cajuputi*, *Caieput* (oil)—o. *Camphoratum*, *Linimentum camphoræ*—o. *Cari seu Carui*, *Carum* (oleum)—o. *Caryophylli*, see *Eugenia caryophyllata*.

OLEUM CEDRINUM, *Essentia de cedro*, (F.) *Huile de cédrat*. The oil of the peel of citron.

obtained in Italy in a particular manner, without distillation.

OLEUM CHABERTI, *O. animale empyreumaticum* Chaberti—*O. Chenopodii*, see *Chenopodium anthelminticum*—*O. Cicinum*, see *Ricinus communis*—*O. Cinnamomi*, see *Laurus cinnamomum*—*O. Contra Tæniam Chaberti*, *Oleum animale empyreumaticum* Chaberti.

OLEUM COPA'IBÆ, *Oil of Copaiba*. (*Copaib. 3liij; Aquæ, cong. iv.* Distil three gallons; separate the oil; return the water to the copaiiba, and again distil three gallons. Separate the oil, and add it to the other. Ph. U. S.) Dose, *gtt. x to xxx.*

OLEUM CORNU CERVI, *O. animale Dippelii*—*O. Cubebæ*, see *Piper cubebæ*—*O. Euphorbiæ lathyridis*, see *Euphorbia lathyris*—*O. Fixum nucum cocos butyraceæ*, see *Cocos butyraceæ*—*O. Fœniculi*, see *Anethum*—*O. Gabianum*, *Petroleum*—*O. Gallinæ*, *Alkalæ*—*O. Gaultheriæ*, see *Gaultheria*—*O. Hedeomæ*, see *Hedeoma pulegioides*—*O. Hyperici*, see *Hypericum perforatum*—*O. Infernale*, *Jatropha curcas* (oil).

OLEUM GRAM'INIS IN'DICI, *Grass-oil of Nemauro*, *Gingergrass oil*, called, by the perfumers, *oil of geranium*, and *oil of rose geranium*, and occasionally sold as *oil of epikenard*, is obtained from *Andropogon calamus aromaticus* of India. It is used as a perfume, and like the essential oils in general.

OLEUM JEC'ORIS ASELI, *O. Morrhue seu Jec'oris Morrhue seu Jecin'oris Aselli*, *Acun'gia Gad'i*, *A. Piscinæ mari'na*, *Codliver oil*, *Cod oil*, (*F.*) *Huile de morue ou de Foie de morue*. The animal oil, which appears under this name in commerce, is obtained from several of the species belonging to the genus *Gadus*. The Cod, *Gadus morrhua*; the Dorse or Dorsch, *Gadus Calla'rias*, *Morrhua America'na*; the Coalfish, *Gadus carbo'narius*; the Burbot, *Gadus lota*, *Lota vulgari's*; the Pollock, *Gadus polla'chius*; the Ling, *Lota molva*; the Torsk, *Bros'mius vulgari's*. There is reason also to believe, that on the American coast it is obtained, also, from the Hake, *Gadus merlu'cius*, and the Haddock, *Gadus ægifi'nus*. The clearest sorts are generally used. An excellent euphotropic, in tuberculous and strumous affections, rheumatism, chronic cutaneous diseases, and tumours of the mammae, &c. The dose for an adult is from *f3ij to f3iiss*.

Cod liver olein, prepared by subjecting codliver oil to a low temperature and separating the olein by pressure, has been used in the same cases as the oil.

Skate liver oil, *Oleum ra'ie*, obtained from *Ra'ia batis* and *R. clava'ta*, is used in the same cases and doses.

OLEUM JECORIS MORRHUE, *Oleum jecoris aselli*—*O. Juniperi*, see *Juniperis communis*—*O. Juniperi empyreumaticum*, see *Juniperus oxycedrus*—*O. de Kervæ*, see *Ricinus communis*—*O. Kervinum*, see *Ricinus communis*.

OLEUM LAURI'NUM, *O. lauri*, *Oil of bays*. An oil obtained from bayberries, and sometimes used in sprains and bruises, unattended with inflammation.

OLEUM LAURINUM, *Daphnelæon*, *Unguentum L.*—*O. Lavandulæ*, see *Lavandula*—*O. Lentiscium*, *Schnelæon*—*O. Limonis*, see *Citrus medica*—*O. Lini*, see *Linum usitatissimum*—*O. Lini cum calce*, *Linimentum aquæ calcis*.

OLEUM LU'CI PISCIS, *Oil of Pike*. From the liver of the *Esox lucius* an oil is spontaneously separated, which is used in some countries to destroy specks on the cornea.

OLEUM MACIDIS, see *Myristica*.

OLEUM MALABATHRI, see *Malabathrum*—*O. Mariæ*, see *Fagara octandra*—*O. Mastichinum*, see *Pistacia lentiscus*—*O. Melaleucæ leucodendri*,

Caieput (oil)—*O. Menthæ piperitæ*, see *Mentha piperita*—*O. Menthæ viridis*, *Mentha viridis*—*O. Monardæ*, *Monarda punctata*—*O. Morrhue*, *O. jecoris aselli*.

OLEUM E MUCILAGINIBUS, *Oil of Mucilages*. (*Rad. althææ rec. 3ss; sem. lini, sem. fœnugraci, 3iij; aquæ, Oij.* Boil for half an hour; add *ol. oliv.* Oiv; continue the boiling till the water is nearly consumed, and pour off the oil.) Emollient.

OLEUM MYRISTICÆ, see *Myristica moschata*—*O. Myristicæ expressum*, see *Myristica*—*O. Neroli*, see *Citrus aurantium*—*O. Nucistæ*, see *Myristica*—*O. Nucistæ æthereum*, see *Myristica*.

OLEUM OLIVÆ, *O. olivæ rum*, *Olives or Salad oil*. An inodorous, insipid, and transparent oil; obtained by expression from the olive, when ripe. It is demulcent and emollient,—possessing the qualities of the fixed oils in general.

An inferior kind, obtained by boiling olives in water, and skimming the oil from the surface, is also used in Pharmacy. See *Olea Europæa*.

OLEUM OMPHACINUM, *Omotribes*—*O. Origani*, see *Origanum*—*O. Oxydi cupri viride*, *Balsam green*, of Metz—*O. de Palmâ Christi*, see *Ricinus communis*—*O. Palmæ*, see *Cocos butyraceæ*—*O. Palmæ liquidum*, see *Ricinus communis*—*O. Petræ*, *Petroleum*—*O. Petræ album*, *Naphtha*—*O. Picinum*, *Brutia*, *Pisselæum*—*O. Pimentæ*, see *Myrtus pimenta*—*O. Pini purissimum*, *O. terebinthinæ rectificatum*—*O. Pulegii*, see *Mentha pulegium*—*O. Pyro-animale*, *O. animale Dippelii*—*O. Raiæ*, see *Oleum jecoris aselli*—*O. Ricini*, see *Ricinus communis*—*O. Rosæ*, see *Rosa centifolia*—*O. Rosarum*, *Rhodolæon*—*O. Rosmarini*, see *Rosmarinus*—*O. Rutæ*, *Peganelæon*—*O. Rutæ*, see *Ruta*—*O. Sabinæ*, see *Juniperus Sabina*—*O. Sassafras*, see *Laurus sassafras*—*O. Sesami*, see *Sesamum orientale*—*O. Sinapis*, see *Sinapis*—*O. Spicæ*, see *Lavandula spica*—*O. Spicæ vulgaris*, *O. terebinthinæ*—*O. Succini*, see *Succinum*—*O. Succini rectificatum*, see *Succinum*.

OLEUM SULPHURATUM, *Balsamum sulphuris simplex*, *Sulphurated oil*. (*Sulphur. lot. 3ij; olivæ olei, 3ij.* Heat the oil in a large iron pot, and throw in the sulphur by degrees; stirring the mixture after each addition till they unite.) It is stimulating, and was formerly much used in coughs, asthma, &c., and, externally, to foul ulcers.

OLEUM TAB'ACI, *Oil of Tobacco*. An empyreumatic oil, obtained from coarsely-powdered tobacco by heating the retort to dull redness.—Ph. U. S.

OLEUM TARTARI PER DELIQUUM, *Liquor potassæ subcarbonatis*—*O. Templinum*, see *Pinus mughos*.

OLEUM TEREBIN'TINÆ, *Oil of Turpentine*. The volatile oil of the juice of *pinus palustris*, and other species of *pinus*.

OLEUM TEREBIN'TINÆ RECTIFICATUM, *O. pini purissimum* *O. T. purificatum* seu *terebinthinæ æthereum*, *Rectified oil of turpentine*, *Sp. of turpentine*. Common oil of turpentine is also called *Common oil of Spike*, *Oleum spicæ vulgaris*. (*Olei terebinth. Oj; aquæ, Oiv.* Distil over the oil.) It is stimulant, diuretic, anthelmintic, and rubefacient. Dose, *gtt. x to f3j*.

Guestonian Embrocation for rheumatism consists of *ol. terebinth.*, *ol. oliv.* 3ss f3iiss; acid sulph. dil. f3ij.

OLEUM THEOBROMÆ CACAO EXPRESSUM, *Butter of cacao*—*O. Tigllii*, see *Croton tiglium*.

OLEUM VALERIANÆ, *Oil of Valerian*. The distilled oil of the root of *Valeriana officinalis*.—Ph. U. S. Dose, four or five drops.

OLEUM VINI, *O. æthereum*—*O. Vitrioli*, *Sulphuric acid*—*O. Vitrioli dulcæ*, *Æther sulphuricus*. OLFACTIF, *Olfactory*.

OLFACTION, *Olfac'tus*, from *olfacere*, *olfac'tum*, (*olere*, 'to smell,' and *facere*, 'to make,') 'to smell;' *Osphe're'sis*, *Osphe'ra'sis*, *Osphe're'sis*, *Osm'e'sis*, *Odora'tio*, *Odora'tus*, *Sensus osometricus*, *Sense of smell*, *Smelling*, (F.) *Odorat*. The sense by which we perceive the impressions made on the olfactory nerves by the odorous particles suspended in the atmosphere. The olfactory nerve or first pair has usually been considered the great nerve of smell; and it is probably the nerve of special sensibility, general sensibility being communicated by the branches of the fifth pair, distributed on the pituitary membrane of the nose and sinuses.

OLFACTIVE, Olfactory.

OLFACTORY, *Olfactive*, *Olfacti'veus*, *Olfact'o-rius*, *Osphe'ran'ticus*, *Osphe'rante'rius*, *Osphe're'ticus*, (F.) *Olfactif*. Same etymon. That which belongs or relates to the sense of smell.

OLFACTORY BULB, see Olfactory nerves.

OLFACTORY FORAM'INA, (F.) *Trous olfactifs*, are the holes in the cribriform plate of the ethmoid bone, through which the olfactory nerve passes.

OLFACTORY LOBE, see Olfactory nerves.

OLFACTORY NERVES, *Ethmoid'al nerves*, *Primum Nervorum cerebri*, *Nervi odoratorii*, *Processus mamillares seu papillares seu mamillares cerebri ad nases seu papilla' res*, *Carunculæ mamillares*,—the first pair of encephalic nerves. This nerve, which probably arises from the medulla oblongata, is observed to leave the brain, opposite the inner part of the fissure of Sylvius, by three roots; which, by their union, form a triangular knot or expansion. When it reaches the ethmoid fossa it expands and forms a triangular ganglion, or grayish, soft bulb,—*Bulbus olfacto'rius*, *Olfactory bulb*, *tubercle*, or *lobe*,—which furnishes, from its inferior surface, the branches that have to be distributed to the nasal fossæ. These filaments are very numerous; they pass through the foramina in the cribriform plate and enter the nasal fossæ. They are distinguished into the *internal*, *external*, and *middle*. The former are distributed over the mucous membrane, covering the outer paries of the nasal fossæ; the second descend upon the septum, and the third are lost, almost immediately, on the portion of the pituitary membrane that lines the vault of the fossæ.

OLFACTORY TUBERCLE, see Olfactory nerves.

OLFACTUS, Olfaction—o. Acutus, Hyperosphresia—o. Amissio, Anosmia—o. Deficiens, Anosmia—o. Depravatus, Parosmia.

OLIBANUM, see Juniperus lycia—o. Sylvestre, see Pinus abies—o. Vulgare, see Pinus abies.

OLIGÆMIA, Anæmia, Hæmaporia, Hypæmia.

OLIGIDRIA, Ischidrosið.

OLIGO, from *oligos*, 'few, little,' in composition has these significations.

OLIGOBLÉN'NIA, from *oligo*, and *βλεννα*, 'mucus.' A deficiency of mucus.

OLIGOCHOL'IA, from *oligo*, and *χολη*, 'bile.' Paucity of bile.

OLIGOCH'YLUS, from *oligo*, and *χυλος*, 'juice,' 'chyle.' An epithet for food which is but little nutritive; which furnishes little chyle.

OLIGOCOP'RIA, from *oligos*, 'few,' and *κοπος*, 'excrement.' Scantiness of alvine evacuations.

OLIGOCYTHÆ'MIA, from *oligo*, *κυτος*, 'a cell,' and *αἷμα*, 'blood.' A condition of the blood in which there is paucity of red corpuscles.

OLIGODAC'RYA, from *oligo*, and *δακρυ*, 'a tear.' Paucity of lachrymal secretion.

OLIGO GALACTIA, Agalactia.

OLIGO GALIA, Agalaxis.

OLIGO HÆMIA, Anæmia, Hæmaporia, Hypæmia.

OLIGOPO'SIA, from *oligo*, and *ποσις*, 'drink.' Diminution in the quantity of drinks.

OLIGOPSYCHIA, Imbecility.

OLIGOSIA'LIA, from *oligo*, and *σάλων*, 'saliva.' Paucity of saliva.

OLIGOSPER'MIA, from *oligo*, and *σπέρμα*, 'sperm.' Paucity of spermatic secretion.

OLIGOSPOND'YLUS, from *oligo*, 'small,' and *σπονδυλος*, 'a vertebra.' A monster with defective vertebrae.—Gurli.

OLIGOTRICH'IA, from *oligo*, and *τριχ*, 'hair.' Want of hair. Paucity of hair.

OLIGOTROPH'IA, *Par'cior nutrit'io*, from *oligo*, and *τρέφω*, 'I nourish.' Deficient nourishment.

OLIGURE'SIA, *Oligure'sis*, *Oligoure'sis*, *Oligoure'sia*, from *oligo*, and *ουρεω*, 'I pass urine.' Morbidity diminished urinary secretion.

OLISTHEMA, Luxation.

OLIVA, Olea Europæa.

OLIVÆ, Corpora olivaria.

OLIVAIRE, Olive-shaped.

Cautère Olivaire, of the French surgeons, is a cautery whose extremity is terminated by a button having the shape of an olive.

OLIVARIS, Olive-shaped.

OLIVARY, Olive-shaped—o. Bodies, Corpora olivaria.

OLIVARY PROC'ESS, *Processus Oliva'ris*, is a small ridge, running transversely between, and a little behind, the roots of the anterior clinoid processes of the sphenoid bone, and by some considered as the fourth clinoid process.

OLIVE, see Olea Europæa—o. Tree, Olea Europæa.

OLIVE-SHAPED, *Oliva'ris*, *Olivifor'mis*, *Ol'ivary*, (F.) *Olivaire*. Resembling an olive. See Corpora olivaria.

OLIVES, PICKLED, Colymbades.

OLIVIER, Olea Europæa—o. de Bohême, Ledum palustre.

OLIVIFORMIS, Olive-shaped.

OLIVINA, see Olea Europæa.

OLLA CAPITIS, Cranium.

OLMITELLO, MINERAL WATERS OF. A thermal spring in the isle of Ischia. Temperature, 100° Fahrenheit.

OLOPHLYCTIDE, Herpes.

OLOPHON'IA, from *ολλω*, *ολλωω*, 'I lose,' and *φωνη*, 'voice.' Congenital misconstruction of the vocal organs.—Good.

OLOPHONTA LABII LOBATA, Harelip—o. Lingua frænata, see Ankyloglossum.

OLSNITIUM, Pucedanum palustre.

OLUS, Lachanon.

OLYMPIAN SPRINGS, see Kentucky, mineral waters of.

OMA, *ωμα*, in the termination of many words, is perhaps derived from *ωμος*, 'crude, wild, imperfect,' as in *sarcoma*, *atheroma*, &c.

OMACEPH'ALUS, (F.) *Omacéphale*, from *ωμος*, 'the shoulder,' and *ακεφαλος*, 'devoid of head.' A genus of paracephalous monsters, without any neck—a slight depression indicating the limits of the head and body.—I. G. St. Hilaire.

OM'AGRA, *Honi'agra*, from *ωμος*, 'the shoulder,' and *αγρα*, 'a seizure.' Gout in the shoulder.

OMAL'GIA, from *ωμος*, 'the shoulder,' and *αλγος*, 'pain.' Pain in the shoulder.

OMARTHRI'TIS, from *ωμος*, 'the shoulder,' *αρθρον*, 'a joint,' and *itis*, denoting inflammation. Inflammation of the shoulder-joint.

OMARTHRO'ACE, from *ωμος*, 'shoulder,' *αρθρον*, 'a joint,' and *κακος*, 'bad.' Arthroace of the shoulder-joint. Caries or suppuration of the shoulder-joint.

OMASUM, *Omasus*, *Manyplies*, (Sc.) *Monipplies*, *Monnyplies*, (F.) *Fenillet*, *Millefenillet*,

Lioret, Psautier. The third stomach of ruminant animals.

OMBILIC, Umbilicus.

OMBILICAL, Umbilical.

OMBRAGE, Nebula.

OMENTA, Meninges.

OMENTESIS, Epiploitis.

OMENTITIS, Epiploitis.

OMENTULA, Appendiculae pinguedinosæ.

OMENTUM, Epiploon—o. Colicum, Epiploon, colic—o. Gastro-colicum, Epiploon, gastro-colic—o. Gastro-splenic, Epiploon, gastro-splenic—o.

Great, Epiploon, gastro-colic—o. Hepato-gastricum, Epiploon, gastro-hepatic—o. Lesser, Epiploon, gastro-hepatic—o. Majus, Epiploon, gastro-colic—o. Minus, Epiploon, gastro-hepatic—o. Ossium, Periosteum—o. Sac of the, see Epiploon, gastro-hepatic—o. Third, Epiploon, colic.

OMICHESES, Micturition.

OMICHA, Urine.

OMITIS, from *omos*, 'the shoulder,' and *itis*, denoting inflammation. Inflammation in or about the shoulder-joint.

OMMA, Eye, Vision.

OMNIPHAGUS, Omnivorous.

OMNIVORE, Omnivorous.

OMNIVOROUS, *Omnivorus*, *Omniphagus*, *Pantophagus*, (F.) *Omnivore*, from *omnis*, 'all,' and *voro*, 'I eat.' An epithet for animals—*Pantophagi*—which eat every kind of food, animal or vegetable.

OMO. In the composition of anatomical terms, an abridgment of *omos* or *omus*, the humerus.

OMO-CLAVICULAR, see Coraco-clavicular.

OMOCOTYLE, see Glene.

OMO-HYOIDEUS, *Cor'aco-hyoïdeus*, *Hyoïdis Quartus Musculus*, (F.) *Scapulo-hyoïdien*, *Omo-hyoïdien*, *Omo-plat-hyoïdien*. This muscle is situate obliquely at the sides and front of the neck. It is slender, long, and flat. It arises from the superior costa of the scapula, near the semi-lunar notch, and from the ligament that runs across it, and is inserted at the sides of the inferior margin of the body of the os hyoides. It consists of two fasciculi, united by a common tendon, and is a true digastric muscle. It depresses the os hyoides, carrying it a little backwards, and to one side, except when it acts with its fellow, when the bone is depressed, and drawn obliquely backwards.

OMOPHAG'IA, *Omosi'ia*, from *omos*, 'raw,' and *phagw*, 'I eat.' Fondness for raw food. Eating of raw food.

OMOPLATE, Scapula—o. *Col de V*, Collum scapulae.

OMOPLAT-HYOÏDIEN, Omo-hyoideus.

OMOS, Crude, Humerus.

OMOSITIA, Omophagia.

OMOTAR'ICHOS, from *omos*, 'the shoulder,' and *tarichos*, 'pickled.' *Salsamentum crudum*; Salted Tunny Fish, in particular. Properly, the neck or shoulder-piece of a salted animal. Once much recommended against the bites of vipers, and in hydrophobia.

OMOTES, Crudity.

OMOTOCIA, Abortion.

OMOTRIBES, *Omphac'inum O'leum*, from *omos*, 'crude,' and *tribw*, 'I bruise.' Oil expressed from unripe olives.

OMPHAC'ION, *omphakion*, from *omphakos*, 'an unripe grape';—*Succus uvæ acer' bæ*. *Omphac'ium*. The juice of unripe grapes. Also, *Verjuice*.

OMPHACITES (VINUM). A name given to wine prepared from the unripe grape. *Omphac'itis* is the name of a small gall.—*Dioscorides*.

OMPHACIUM, Verjuice.

OMPHACOM'ELI, from *omphakos*, 'an unripe

grape,' and *meli*, 'honey.' A sort of oxymel, made of the juice of unripe grapes and honey.

OMPHALECO'SIS, from *omphalos*, 'the navel,' and *'alekos*, 'an ulcer.' Ulceration of the navel.

OMPHALEXOCHE, Exomphalus.

OMPHALOCARPUS, Galium aparine.

OMPHALOCLE, Exomphalos.

OM'PHALO-CRA'NIODID'YMUS, from *omphalos*, 'the navel,' *κρανιον*, 'the cranium,' and *διδυμος*, 'a twin.' A double monster, in which the umbilical cord or the rudiment of one fœtus is rooted in the cranium of another.

OMPHALOMANT'IA, *Om'phalomancy*, from *omphalos*, 'the navel,' and *μαντεια*, 'prophecy.' A species of divination, practised by credulous matrons, who pretend to be able to know the number of children a female will have, by the number of knots in the navel-string of the child.

OM'PHALO-MESENER'IC, *Omphalo-mesenter'icus*, *Om'phalo-mesara'icus*; from *omphalos*, 'the navel,' and *mesenterium*, 'the mesentery.'

OMPHALO-MESENERIC VESSELS, *Vitel'lo-mesenter'ic*, *Vitelline vessels*. Haller gave this name to two very fine vessels, which spread their ramifications on the parietes of the umbilical vesicle. There is an *omphalo-mesenteric artery and vein*. The *omphalo-mesenteric artery* is a branch of the superior mesenteric;—the vein empties itself into the trunk, or into one of the branches of the superior mesenteric. Velpeau affirms, that they insinuate with a branch of the second or third order of those great vessels, with those in particular that are distributed to the cœcum, and he regards them to be the vessels of nutrition of the umbilical vesicle. They are occasionally met with in the fœtus, at the full period, under the form of whitish filaments, which extend from the mesenteric vessels to the umbilicus.

OMPHALO-MESERAIC, *Omphalo-mesenteric*.

OMPHALONCUS, Omphalophyma.

OMPHALONEURORRHEXIS, Omphalorrhæxis.

OMPHALOPHLEBITIS, from *omphalos*, 'the umbilicus,' and *phlebitis*, 'inflammation of a vein.' Inflammation of the umbilical vein.

OMPHALOPHY'MA, *Omphalon'cus*, from *omphalos*, 'the navel,' and *φυμα*, 'tumour.' A tumefaction of the navel.

OMPHALOPROPTOSIS, Exomphalus.

OMPHALORRHAG'IA, *Omphalor'rhagæ*, from *omphalos*, 'the umbilicus,' and *ρηννμι*, 'I break out.' Hemorrhage from the umbilicus, in the new-born in particular.

OMPHALORRHEX'IS, from *omphalos*, 'navel,' and *ρηξις*, 'rupture.' *Omphaloneurorrhæx'is*. Rupture of the navel string.

OMPHALOS, Umbilicus.

OMPH'ALOSITE, from *omphalos*, 'the navel,' and *σιτος*, 'nourishment.' A monster that possesses an imperfect kind of life, which ceases when the umbilical cord is divided.

OMPHALOT'OMY, *Omphalotom'ia*, from *omphalos*, 'the umbilicus,' and *τεμνω*, 'I cut.' The division of the navel string.

OMPHALUS, Umbilicus.

OMUS, Crude, Humerus.

ONAGRA, *E'nothera biennis*.

ONANIA, Masturbation.

ONANISM, Masturbation.

ON-COME. A Scotch term, according to Sir Walter Scott, for "mysterious diseases, which baffle the regular physician."

ONCOS, Tumour.

ONCO'SES, from *ογκος*, 'a tumour.' Tumours, as diseases.

ONCOSTIS, Intumescence.

ONCOT'OMY, *Oncotom'ia*, *Onkotomy*, from *ογκος*, 'a tumour,' and *τομη*, 'incision.' The open-

ing of an abscess with a cutting instrument, or the excision of a tumour.

ONCTION, Inunction.

ONCUS, Swelling, Tumour.

ONDULATION, Fluctuation.

ONEBERRY, Paris.

ONEIROCRITICUS, *Oneirocrites*, from *oneiros*, 'a dream,' and *kritikos*, 'judgment.' One who judges according to dreams.

ONEIRODYN'IA, from *oneiros*, and *odynē*, 'pain.' Painful dreams. *Incubus* and *Somnambulism* are oneirodynia.

ONEIRODYNIA ACTIVA, Paroniria, Somnambulism—o. *Gravans*, Incubus.

ONEIRODYNIE GRAVATIVE, Incubus.

ONEIROG'MUS, *Oneirog'ynē*, *Oneirogopol'esis*, from *oneiros*, 'a dream.' A lascivious dream; pollution; nocturnal pollution.

ONEIROGONORRHEA, Pollution, nocturnal.

ONEIROGONOS, Pollution.

ONEIROLOGY, *Oneirolog'ia*, from *oneiros*, 'a dream,' and *logos*, 'a description.' The doctrine of dreams.

ONEIROMANTIA, from *oneiros*, 'a dream,' and *mantia*, 'divination.' The art of divining by dreams; or of interpreting dreams.

ONEIRON'OSUS, from *oneiros*, and *osos*, 'a disease.' *Somnia'tio morbo'sa*. Morbid, uneasy dreaming.

ONEIROPOLESIS, Oneirogmus.

ONERA ALVI, Excrement.

ONGLADE, Paronychia.

ONGLE, Nail—o. *Incarné*, Onychogryphosis.

ONGLÉE (F.), *Digitorum stupor à gelu*. Painful numbness at the extremities of the fingers, caused by cold.

ONGLET, Pterygion.

ONGUENT, Unguentum—o. *d'Althéa*, Unguentum of Althæa—o. *Blanc*, Ceratum cetacei—o. *de Blanc de baleine*, Unguentum cetacei—o. *Blanc de Rhazes*, Unguentum plumbi subcarbonatis—o. *de Céruse*, Unguentum plumbi subcarbonatis—o. *Oitrin contre la gale*, Unguentum hydrargyri nitras—o. *Digestif simple*, Unguentum digestivum simplex—o. *Gris*, Unguentum oxidi hydrargyri cinereum—o. *de Guimauve*, Unguentum of althæa—o. *de Laurier*, Unguentum laurinum—o. *Mercuriel*, Unguentum hydrargyri—o. *Napolitain*, Unguentum hydrargyri—o. *de Nitrate de mercure*, Unguentum hydrargyri nitratiss—o. *de Poiz et de cire*, Basilicon—o. *de Soldat*, Martiatum unguentum—o. *de Sureau*, Unguentum sambuci—o. *de Tornamira*, Unguentum plumbi subcarbonatis.

ONION, COMMON, Allium cepa—o. Sea, Scilla.

ONIS'CUS, *Asell'us*, *Cut'tio*, *Porcel'tio*, *Porcel'us*, *Por'culus*, diminutive of *ovos*, 'a small ass.' A genus of insects, *Class*, Crustacea, very common in cellars and dark and moist places.

ONISCI ASELLI, *Miller'edes*, *Aselli*, *Miller'edes*, *Faba*, *Wood-lice*, *Slaters*, *Hog-lice*, (F.) *Cloportes ordinaires*, had, at one time, a place in the pharmacopœias. They were considered stimulant and diuretic, and useful in jaundice.

ONITIS, Origanum dictamnus.

ONKOTOMY, Oncotomy.

ONOLOSAT, Obolus.

ONOMATOCLESIS, Nomenclature

ONOMATOLOGIA, Nomenclature.

ONONIS ANTIQUORUM, O. spinosa.

ONONIS ARVEN'SIS, (F.) *Bugrane des champs*, has properties like the next.

ONONIS SPINOSA, *Ano'nis*, *Resta* seu *Arres'ta dovis*, *Rem'ora Ara'tri*, *Ononis antiqu'rum*, *Rest harrow*, (F.) *Arrête-bœuf*, *Bugrande épineuse*, *Bugrane*, *Nat. Ord. Leguminosæ*. *Sec. Syst. Diadelphia Decandria*. The root of this plant was once used as a diuretic.

ONOPORDON ACAN'THIUM, *Onopordum acanthium*, *Carduus tomento'sus*, *Acan'thium*, *Ac'anos*, *A. spina*, *Spina alba*, *Cotton Thistle*, (F.) *Chardon aux Anes*. *Ord. Compositæ*. *Sex. Syst. Syngenesia Polygamia æqualis*. The expressed juice has been recommended as a cure for cancer, applied externally.

ONOPORDUM ACANTHIUM, Onopordon acanthium.

ONTHUS, Fimus.

ONYCHATROPH'IA, *Atroph'ia un'guium*, from *ονυξ*, *ονυχος*, 'a nail,' and *atrophia*, 'want of nourishment.' Atrophy of the nails.

ONYCHAUX'E, *Hypertroph'ia un'guium*, from *ονυξ*, *ονυχος*, 'a nail,' and *αυξη*, 'increase.' Unusual increase of the size and thickness of the nails.

ONYCHEXALLAX'IS, from *ονυξ*, *ονυχος*, 'a nail,' and *ελλαξις*, 'change.' A morbid condition of the nails.

ONYCHIA, Paronychia.

ONYCHITIS, from *ονυξ*, *ονυχος*, 'a nail,' and *itis*, denoting inflammation. Inflammation of a nail. See Paronychia.

ONYCHOC'LASIS, from *ονυξ*, *ονυχος*, 'a nail,' and *κλασις*, 'fracture.' Fracture of a nail.

ONYCHOCRYPTOSIS, Onychogryphosis.

ONYCHOGRYPH'OSIS, *Onychogryppo'sis*, *Onychogryppo'tis*, *Onychocrypto'tis*, *Grypho'sis*, *Grypho'sis*, *Onyx'is*, from *ονυξ*, *ονυχος*, 'a nail,' and *gryposis*, 'crookedness.' Curvature of the nails; such as occurs in hectic individuals. Also, growing in of the nails; *Incarna'tio seu Adunca'tio seu Arctura un'guium*, (F.) *Ongle incarné*.

ONYCHON'OSI, *Onychon'osi*, from *ονυξ*, *ονυχος*, 'a nail,' and *ososis*, 'a disease.' Disease of the nails.

ONYCHOPH'THORA, *Onychophtho'ria*, *Onychoph'thora*, *Onychophtho'ria*, from *ονυξ*, *ονυχος*, 'the nail,' and *φθορα*, 'degeneration.' A degenerate condition or destruction of the nails.

ONYCHOPHY'MA, from *ονυξ*, *ονυχος*, 'the nail,' and *φυμα*, 'a tumour.' A painful degeneration of the nails, which become thick, rough, and crooked. It occurs as a symptom of syphilis and of lepra.

ONYCHOPT'OSIS, *Piptonychia*, *Dia'resis un'guis*, from *ονυξ*, *ονυχος*, 'a nail,' and *πτωσις*, 'falling.' The falling off of the nails.

ONYCOPHTHORA, Onychophthora.

ONYX, Hook (blunt). Nail, Pterygion. Also, a collection of purulent matter between the laminae of the cornea, having the shape of a nail. Unguis Lunula.

ONYXIS, see Onychogryphosis.

ONYXITIS, see Paronychia.

OOCYESIS, Pregnancy, ovarian.

OODEOCELE, see Hernia (foraminis ovalis.)

OODES, Aqueous humour, Oval.

OÖG'ALA, *Oög'la*, *Ogla*, *Puls ex Ovis et Lactis*. Milk of eggs, from *ovon*, 'an egg,' and *gala*, 'milk.' A preparation of eggs and milk.

OÖGLA, Oogala.

OÖIDES, Aqueous humour, Oval.

OÖLEMMA PELLUCIDUM, *Zona pellucida*.

OÖLOG'IA, *Ovolog'ia*, *Ovolog'y*, from *ovon*, 'an ovum or egg,' and *logos*, 'a discourse,' 'a description.' A description of the ovum.

OÖN, Ovum.

OÖNINE, Albumen.

OÖPH'AGOUS, *Oöph'agus*, from *ovon*, 'an egg,' and *φαγω*, 'I eat.' Feeding on eggs.

OÖPHORAUXE, Oarioparectama.

OÖPHORITIS, Oaritis.

OÖPHORON, Ovary.

OÖRITIS, Oaritis.

OPAC'ITY, *Opac'itas*, *Opakeness*; from *opacare*, 'to obscure.' Quality of that which is

opaque. The property possessed by some bodies of not allowing the light to traverse them. It is opposed to transparency. *Opacity of the cornea* constitutes *albugo* or *leucoma*;—opacity of the crystalline causes cataract.

OPAKE', *Opa'cus*, *Opa'cous*. Same etymon. An epithet given to bodies, which do not permit the passage of rays of light. The *opaque cornea* is the sclerotic, in contradistinction to the transparent cornea or true cornea.

OPAKENESS, Opacity.

OP'ALINE. That which resembles the opal. A fluid is said to be opaline, when milky, and when it presents an appearance more or less like that of the opal.

OPEOCLE, see Hernia.

OPERATIO, Action, Operation—o. Chirurgica, see Operation—o. Chymica seu pharmaceutica, Operation, chymical, &c.—o. Simplex, Haplotomia.

OPERATION, *Opera'tio*, from *opus*, *operis*, 'work.' The application of instruments to the human body with the view of curing disease. The object of an operation, *opera'tio chirurgica*, is generally to divide or reunite parts, to extract extraneous or noxious bodies, and to replace organs that are wanting, by different instruments or artificial means. The principal operative methods have been called *Synthesis*, *Diæresis*, *Excresis*, and *Prothesis*. Frequently, the most difficult subject connected with an operation is to decide when it is absolutely called for or advisable, and when improper.

OPÉRATION À LAMBEAU, see Flap.

OPERATION, BILATERAL, see Lithotomy—o. *Cæsarienne*, Cæsarean section.

OPERATION, CHYMICAL or PHARMACEUTICAL, *Opera'tio chymica seu pharmaceutica*, is any process whose object is the preparation of medicines; whose combinations with each other, analysis, and decomposition;—such are, *solution*, *distillation*, *sublimation*, *evaporation*, *digestion*, *maceration*, *infusion*, *decoction*, *calcination*, &c. Some of these operations are mechanical; others, really chymical.

OPERATION, HIGH, see Lithotomy—o. Lateral, see Lithotomy—o. Sigaultian, Symphyseotomy.

OPERATOR. Same etymon. A surgeon who is in the habit of practising the greater operations. One who performs any operation. To be a good operator, the surgeon must be well acquainted with anatomy, and be possessed of strong nerve. See Surgeon.

OPERCULUM, *Opero'trium*, *Operimen'tum*, from *operire*, 'to cover.' A cover or lid. In fishes, the gill cover.

OPERCULUM ASPERÆ ARTERIÆ, Larynx—o. Cutis, Epidermis—o. *Hei*, Bauhin, valve of—o. Laryngis, Epiglottis—o. Oculi, Palpebra.

OPERCULUM PAPILLÆ. A shield for the nipple.

OPERIMENTUM, Operculum—o. Intestinorum, Epiploon—o. Prætensum abdominis, Peritonæum.

OPERITORIUM, Operculum.

OPHELIA CHIRATA, Gentiana chirayta.

OPHIASIS, Calvities, Porrigo, decalvans.

OPHIOGLOS'SUM VULGA'TUM, *O. ophioglossum*, from *ophis*, 'a serpent,' and *γλῶσσα*, 'a tongue,' *Lingum serpentaria*, *Adder's Tongue*, (F.) *Langue de Serpent*. Ord. Filices. *Sex. Syst.* Cryptogamia. This plant was formerly considered to be vulnerary.

OPHIOPH'AGOUS, *Ophioph'agus*, from *ophis*, 'a serpent,' and *φαγω*, 'I eat.' An epithet for animals which feed on serpents. Pliny has called certain African tribes by this name.

OPHIORRHIZA, *O. mungos*, from *ophis*, 'a serpent,' and *ρίζα*, 'root.' Family, Gentianaceæ.

Sec. Syst. Pentandria Digynia. The name of the plant whose root has been called *Rudix Serpentum*, *Mungos Radix*. The bitter root is much esteemed in Java, Sumatra, &c., for preventing the effects that usually follow the bite of the *naja*, a venomous serpent; with which view it is eaten by the natives. It is, also, a reputed antelmintic.

OPHIOXYLON, *Ophioxylum*.

OPHIOXYLUM, from *ophis*, 'a serpent,' and *ξύλον*, 'wood.' *Ophioxylum seu Ophioxylon serpentinum seu trifoliatum*, *Acavéria*, *Lignum serpentum*, Ord. Apocynaceæ, has been recommended in the bites of serpents, and in intermittents. It is said to be very bitter.

OPHIS, Serpent.

OPHITES, from *ophis*, 'a serpent.' *Serpentinus*, *Serpentine* or *Black Porphyry*. This rock was formerly worn as an amulet, to cure diseases of the head.

OPHROSTAPHYLON, *Bryonia alba*.

OPHRYS, Supercilium.

OPHTHALMAL'GIA, from *οφθαλμος*, 'the eye,' and *αλγος*, 'pain.' Pain in the eye. Neuralgia of the eye.

OPHTHALMEMPASMA, Collyrium siccum.

OPHTHAL'MIA, *Ophthalmi'tis*, from *οφθαλμος*, 'the eye.' *Oculo'rum inflammatio*, *Cauma ophthalmi'tis*, *Ophthalmop'onia*, *Inflammation of the Eye*, *Ophthalm'y*, (F.) *Ophthalmie*, *Ophthalmite*, *Inflammation de l'œil*, *Catarrhe oculaire*. Three great varieties of ophthalmia, independently of the acute and chronic (which conditions occur in all inflammations), may be reckoned:—the *ophthalmia membranarum*, *O. purulenta*, and *O. tarai*. The first is characterized by the general symptoms of ophthalmia;—pain and redness of the eye or its appendages; with intolerance of light, and unusual flow of tears; the inflammation being seated chiefly in the coats of the eyeball. It is the *Hymenophthalmia*, *Ophthalmia tarax'is seu hu'mida seu vera seu chemo'sis*, *Conjunctivitis*, *Inflammatio conjunctivæ*, *Symphymen'itis*, *Syndesm'itis*, *Catarr'hal ophthalmia*, *Ophthalmo-conjunctiv'itis*, *Epiph'ora* (Galen), *Conjunctival or Lachrymose Ophthalmia*, *Tarax'is* (Paulus of Aegina), *Chemo'sis* (Aëtius), (F.) *Conjonctivite*, *Inflammation de la conjonctive*. In the second,—*Ophthalmia purulenta seu puriform'is*, *Blennophthalmia*, *Bleph'aroblenorrhœ'a*, *Ophthalmoblenorrhœa*, *Blennorrhœa oculi*—the internal surface of the palpebræ associates in the inflammation of the eye-ball; and there is a copious secretion of a purulent fluid. An epidemic and contagious (?) variety of this is the *Egyptian Ophthalmia*, *O. epidem'ica seu purulen'ta epidem'ica seu contagio'sa seu catarrhal'is bel'lica seu bel'lica seu Asia'tica seu purulenta contagio'sa*, *Blennorrhœ'a Oculi Egyptiaca*, *Lemophthal'mia*, *Conjunctivitis puro-mucosa*, *Contagio'sa seu Egyptiaca*, *Blepharot'is glandular'is contagio'sa*, *Adenit'is palpebrarum contagiosa*, *Suppurative ophthalmia*, *Epidemic contagious ophthalmia*—called *Egyptian*, from its prevalence in Egypt during the British Expedition under Sir Ralph Abercrombie. The inflammation is rapid and destructive; granulations shoot from the tunica conjunctiva, and occasionally there is intolerable pain, often succeeded by delirium. In new-born children, a species of purulent ophthalmia, *O. purulen'ta infantum seu neonato'rum*, *Blepharophthalmia seu Lippitu'do seu Blennorrhœ'a Oculi seu Ophthalmoblenorrhœ'a seu Blepharoblenorrhœ'a seu Blepharopyorrhœ'a seu Pyophthalmia*, *Peorophthalmia neonato'rum*, *Adenoeynchiton'itis*, (F.) *Ophthalmie puriforme des nouveau-nés*, in which the palpebræ are florid and peculiarly tumid, is by no means uncommon. It seems to be caused by acrid discharges from the mother, applied to

the eye of the infant during its exit; or to the stimulus of the light, when the child first opens its eyes. A severe form of purulent ophthalmia—*Hyperconjectivite* (Piorry)—is produced by the application of gonorrhoeal matter to the eye. It is the *Ophthalmia gonorrhoeica*, *Blennorrhoea oculi gonorrhoeica*, *Gonorrhoeblepharhoea*, *Conjunctivitis blennorrhagica* seu *gonorrhoeica*, *Gonorrhoeal Ophthalmia*, (F.) *Ophthalmie blennorrhagique*, *Conjunctivite blennorrhagique*.

Ophthalmia is likewise modified by the condition of the constitution, and hence we have *stru-*

mous, *variolous*, and other inflammations of the conjunctiva.

The third variety—the *Ophthalmia Tarsi* seu *glutinosæ*, *Blepharophthalmia*, *Blepharotis*, *Blepharotitis*, *Blepharitis*, *Blepharadenitis*, *Psorophthalmia*, *Oculi palpebrarum scabies* pruriginosa, *Adenophthalmia*—is seated chiefly in the tarsus; the sebaceous crypts secreting a viscid and acrid fluid, that glues and ulcerates its edges, and irritates the eye.

The different forms of inflammation of the conjunctiva are thus classed by M. Desmarres:

Conjunctivitis.....	1. Pure or phlegmonous.	{	{	{	{
	2. Pustular.				
	3. Granular or catarrhal.				
	4. Purulent.....				
				Contagious.	
				Non-contagious.	
				Miasmatic.	
				Exanthematic.....	Erysipelatous.
					Variolous.
					Morbillous.
					Scarlatinous.
				Of new-born infants.	
				Gonorrhoeal.	
				Egyptian.	

Conjunctivitis, when slight, requires little treatment: the antiphlogistic regimen—and, particularly, avoiding exposure to light,—being all that is necessary. When more severe, along with this, general and local blood-letting, especially by leeches to the side of the septum narium, must be employed so as to make a decided impression on the system; with nauseating doses of antimony, purgatives, blisters (at some distance from the organ), refrigerants; and astringents, cold or warm, according to circumstances, to the eye. In *Purulent Ophthalmia*, these measures must be still more actively employed, otherwise disorganization will speedily follow:—the granulations, which form on the adnata of the eyelids may be removed with the scissors, and the sore be touched with a solution of alum. *Ophthalmia Tarsi* must be treated on general principles, when severe. It usually, however, requires merely the antiphlogistic regimen, with the occasional use, especially at night, of a slightly stimulating ointment, such as the unguentum hydrargyri nitrico-oxydi, considerably reduced with lard. The ointment prevents the eyelids from being glued together during the night, and thus obviates the irritation caused by attempts at separating them.

OPHTHALMIA, *Ægyptian*, see Ophthalmia—o. *Asiatica*, see Ophthalmia—o. *Bellica*, see Ophthalmia—o. *Biliosa*, *Ophthalmocholosis*—o. *Catarrhal*, see Ophthalmia—o. *Catarrhalis bellica*, see Ophthalmia—o. *Chronica*, *Lippitudo*—o. *Conjunctival*, see Ophthalmia—o. *Contagiosa*, see Ophthalmia—o. *Epidemica*, see Ophthalmia—o. *Glutinosæ*, see Ophthalmia—o. *Gonorrhoeica*, see Ophthalmia—o. *Gonorrhoeal*, see Ophthalmia—o. *Humida*, see Ophthalmia—o. *Lachrymose*, see Ophthalmia—o. *Membranarum*, see Ophthalmia—o. *Neonatorum*, see Ophthalmia—o. *Phlyctenular*, see Ophthalmia—o. *Puriformis*, see Ophthalmia—o. *Purulenta*, see Ophthalmia—o. *Purulenta contagiosa*, see Ophthalmia—o. *Purulenta epidemica*, see Ophthalmia—o. *Purulenta infantum*, see Ophthalmia—o. *Rheumatic*, *Scleritis*—o. *Sicca*, *Xerophthalmia*—o. *Suppurative*, see Ophthalmia—o. *Taraxis*, see Ophthalmia—o. *Tarsi*, see Ophthalmia—o. *Ulcerosæ*, *Helcophthalmia*—o. *Varicose*, *Cirsophthalmia*—o. *Vera*, see Ophthalmia.

OPHTHALMIATER, Oculist.

OPHTHALMIATRIA, *Ophthalmiatrotechnica*, *Ophthalmotherapia*, (F.) *Oculistique*; from

ophthalmos, and *iarpos*, 'a physician.' The art of the oculist. Treatment of diseases of the eye.

OPHTHALMIATROTECHNICA, Ophthalmiatria.

OPHTHALMIC, *Ophthalmicus*, from *ophthalmos*. That which relates or belongs to the eye.

OPHTHALMIC ARTERY, *A. orbitaire*, (Ch.) arises from the internal carotid, and issues from the cranium by the foramen opticum. At first, it is situate at the outer and lower side of the optic nerve, but ascends above this nerve, and passes towards the inner angle of the orbit. It furnishes, in its course, the *lachrymal*, *centralis retineæ*, *infra-orbitares*, *ciliaries*, *anterior ciliaries*, *inferior and superior musculares*, *anterior and posterior ethmoidals*, and *superior and inferior palpebrals*. After having given off these, it divides into two branches—the *frontal* and *nasal*, which furnish a number of ramifications,—some superficial, others deep-seated.

OPHTHALMIC GANGLION, *Lenticular* or *Ciliary ganglion*, *Ganglion ophthalmicum* seu *semilunare seu ciliare*, *G. orbitaire* (Ch.), is one of the smallest ganglions of the body, and formed by the 5th and 3d pairs of encephalic nerves. It is situate at the outer side of the optic nerve near the bottom of the orbit; is irregularly quadrilateral and flat; its colour of a reddish gray. Behind, it communicates by its posterior angles—by means of nervous filaments—with the nasal nerve of the ophthalmic, the motor oculi, and offsets from the sympathetic. Each of its anterior angles furnishes a fasciculus of small nerves. These are the ciliary nerves.

OPHTHALMIC NERVE, *Ophthalmic Nerve* of Willis, *Orbito-frontal*, (Ch.) is the smallest and first of the three branches given off by the 5th pair. It proceeds along the external paries of the cavernous sinus, and enters the orbit by the sphenoidal fissure. It divides into three branches, one, *external*—the lachrymal nerve; another, *superior*—frontal nerve; and the last, *internal*—the nasal nerve. See *Lachrymal*, *Frontal*, *Nasal*.

OPHTHALMIC or OPTIC REMEDY, *Ophthalmicum* seu *Opticum*, (F.) *Topique ophthalmique*, is a medicine employed in ophthalmia.

OPHTHALMIC SINUS, see *Cavernous Sinus*.

OPHTHALMIC VEIN has the same arrangement as the artery, which it accompanies in all its divisions. It issues from the orbit at the inner

part of the sphenoidal fissure, and discharges its blood into the cavernous sinus.

The name *Facial Ophthalmica* has been given to the branch by which the facial vein communicates with the ophthalmic.

OPHTHALMICA, *Euphrasia officinalis*.

OPHTHALMICI EXTERNI, *Motores oculorum*.

OPHTHALMICUM, Ophthalmic remedy.

OPHTHALMIDIUM, Microphthalmus.

OPHTHALMIE PURIFORME DES NOUVEAUX-NÉS, see Ophthalmia — *o. Blennorrhagique*, see Ophthalmia.

OPHTHALMITE, Ophthalmia.

OPHTHALMITIS, Ophthalmia. This term is occasionally restricted to inflammation of the globe of the eye, in which, both the external and internal structures are involved.

OPHTHALMIUM, Microphthalmus.

OPHTHALMO-BLENNORRHOEA, Ophthalmia (purulent) — *o. Neonatorum*, see Ophthalmia (purulenta infantum).

OPHTHALMO-CARCINOMA, from *ophthalmos*, and *καρκινωμα*, 'cancer.' Cancer of the eye.

OPHTHALMOCELE, Exophthalmia.

OPHTHALMOCHOLOSIS, from *ophthalmos*, and *χολη*, 'bile;' Ophthalmia biliosa. Ophthalmia from biliary excitement.

OPHTHALMOCHROITES, see Choroidea tunica.

OPHTHALMO-CONJUNCTIVITIS, see Ophthalmia.

OPHTHALMOCOPIA, Copyopia.

OPHTHALMODULIA, from *ophthalmus*, and *δουλεια*, 'servitude.' Eye-service. The title of a book by Bartsch on diseases of the eye.

OPHTHALMODYNIA, from *ophthalmos*, and *δυνη*, 'pain,' especially rheumatic pain of the eye. Plenck has given this name to neuralgia of the frontal nerve, in which the pain radiates particularly towards the bottom of the orbit. See Neuralgia, frontal.

OPHTHALMODYNIA PERIODICA, Neuralgia, facial.

OPHTHALMOEDEMA, from *ophthalmos*, and *οιδημα*, 'oedema.' Oedema of the conjunctiva.

OPHTHALMOGRAPHY, from *ophthalmos*, and *γραφη*, 'a description.' The part of anatomy which gives a description of the eye. An anatomical description of the eye.

OPHTHALMOLOGY, *Ophthalmolog'ia*, *Ophthalmotology*, from *ophthalmos*, and *λογος*, 'a discourse.' The part of anatomy which treats of the eye. An anatomical treatise on the eye. A description of the eye in health and disease.

OPHTHALMOMETER, from *ophthalmos*, and *μετρον*, 'measure.' An instrument of the nature of compasses, invented by F. Petit, for measuring the capacity of the anterior and posterior chambers of the eye in anatomical experiments.

OPHTHALMOMYITIS, *Ophthalmomyositis*, from *ophthalmos*, *μυς*, 'a muscle,' and *itis*, denoting inflammation. Inflammation of the muscles of the eye.

OPHTHALMOPARALYSIS, Ophthalmoplegia.

OPHTHALMOPHLEBOTOMIA, from *ophthalmos*, *φλεψ*, 'a vein,' and *τομη*, 'incision.' Bleeding from the vessels of the conjunctiva.

OPHTHALMOPHTHARISIS, *Ophthalmophthisis*, from *ophthalmos*, and *φθορασις*, 'corruption.' Destruction of the eyeball.

OPHTHALMOPHTHISIS, Ophthalmophtharisis.

OPHTHALMOPLAGIA, *Ophthalmoparaly-sis*, from *ophthalmos*, and *πλησσω*, 'I strike.' Paralysis of one or more of the muscles of the eye.

OPHTHALMOPONIA, Ophthalmia.

OPHTHALMOPTOSIS, from *ophthalmos*, and *πτωσις*, 'a prolapse.' A word employed by some authors in the sense of exophthalmia; by others, in that of hydrophthalmia.

OPHTHALMORRHAGIA, from *ophthalmos*, and *ρρηγμα*, 'I break forth.' Hemorrhage from the tunica conjunctiva. It is rare.

OPHTHALMOS, Eye.

OPHTHALMOSCOPE, see Ophthalmoscopy.

OPHTHALMOSCOPIA, *Ophthalmoscopy*, from *ophthalmos*, and *σκοπεω*, 'I regard attentively.' The art of judging of the temper, &c., of a person by examining his eyes. The art of judging of health or disease by inspection of the eyes. Exploration of the eyes in order to a diagnosis.

An *Ophthalmoscope*, *ophthalmoscopium*, *ophthalmoscopum*, has been invented, which by reflecting the light on the retina enables the condition of the interior of the eye to be appreciated. To a form of the ophthalmoscope the name *Anagnostakis* has been given, after a young Greek physician, who, it was thought, invented it.

OPHTHALMOTATUM (F.) *Ophthalmotat*, from *ophthalmos*, and *στασις*, 'station;' *Speculum Oculi*. An instrument for fixing the eye.

OPHTHALMOTHERAPIA, Ophthalmiatria.

OPHTHALMOTOLOGY, Ophthalmology.

OPHTHALMOTOMY, *Ophthalmotomia*, from *ophthalmos* and *τομη*, 'I cut.' The part of anatomy which treats of the dissection of the eye. It has, also, been applied to extirpation of the eye.

OPHTHALMOXEROSIS, Xerophthalmia.

OPHTHALMOXYSIS, from *ophthalmos*, and *ζωω*, 'I scrape.' A name given to the scarification sometimes practised on the conjunctiva, in cases of ophthalmia.

OPHTHALMOXYSTRUM, *Ophthalmoxyster*, *Ophthalmoxysterium*, *Xystrum ophthalmicum*. Same etymon. An instrument for scraping the eye. Name given to a small brush, with barbs like an ear of barley or rye, intended to scarify the eyelids in certain cases of ophthalmia.

OPHTHALMUS, Eye.

OPHTHALMYMENITIS, *Ophthalmohymenitis*, from *ophthalmos*, *μνην*, 'membrane,' and *itis*, denoting inflammation. Inflammation of one or more of the membranes of the eye.

OPIACE, Opiate.

OPIACUM, Opiatum.

OPIANE, Narcotine.

OPIANINE, *Opiantina*, *Opiania*; from *opium*. An alkaloid discovered in opium, which resembles morphia in its properties, and doses.

OPIANUM, Narcotine.

OPIAT, Opiate — *o. Dentifrice*, Dentifrice.

OPIATE, *Opiatum*, *Opiacum*, *Papaveraceum*, *Mecconicum*, (F.) *Opiat*, *Opiace*, from *οπιον*, (*σπος*, 'juice,') 'opium.' A medicine containing opium. A medicine that procures sleep. An electuary; — formerly, an electuary which contained opium.

OPIATED, *Opiatus*. Impregnated with opium. Affected by opium.

OPIATUM FEBRIFUGUM, *Electuarium de Kina Kina*; *Electuary of Cinchona*. (*Pulv. cinch. 3xviij, ammon. muriat. ʒij, mellis, syrup., absinth. aa ʒij*. Make into an electuary. *Ph. P.*) Given, when cinchona is indicated.

OPIATUM MESENTERICUM, *Electuarium de Aloë*, *Muriatide Hydrargyri, et Ferro*. (*Gum. ammon. ʒss. senecæ ʒij, hydrargyri submuriat., rad. ari., aloës socotrin. aa ʒij, pulv. scammon. comp. (vulg. de tribus), rhej; rad. aa ʒij, ferri limatur. porphyrisat. ʒss*. Bruise and mix together, add of compound syrup of apples double the weight of the other matters, and make into an electuary. *Ph. P.*) Dose, ʒss to ʒij, in obstructions of the liver, mesentery, &c.

OPIATUS, Opiated.

OPION, Opium.

OPISMA, Succus.

OPIS'THENAR, *Dorsum manûs*, from *οπισθε*, 'backwards,' and *θηνα*, 'the flat of the hand.' The back of the hand.

OPISTHOCEPHALON, Occiput.

OPISTHOCRANIUM, Occiput.

OPISTHOCYPHOSIS, Cyphoma.

OPISTHOLOB'IUM, *Opistho'tium*, from *οπισθε*, 'behind,' and *λοβιον*, 'the lobe of the ear.' Any agent applied behind the ear.

OPISTHOTIUM, Opisthobolium.

OPISTHOT'ONOS, *Raptus posterga'neus*, *Tet'anus dorsa'lis seu post'icus seu posterga'neus*, from *οπισθε*, 'backwards,' and *τεινω*, 'I stretch.' A species of tetanus, in which the body is bent backwards.

OPIUM, see Papaver—o. Colatum, Extractum opii—o. Depuratum, Extractum opii.

OPIUM EATERS, Theriaki.

OPIUM, Lettuce, see Lactuca.

OPOBALSAMUM, see Amyris opobalsamum.

OPOCARPSON, Carpasium (juice.)

OPOCEPH'ALUS, (F.) *Opocephale*. A genus of monsters having the two ears approximated or united under the head; the jaws atrophied; and no mouth or nose.—I. G. St. Hilaire.

OPODELDOC, Opodeldoch.

OPODEL'DOCH, *Opodel'toch*, *Opodeldoc*. An unmeaning term, frequently used by Paracelsus. Formerly, it signified a plaster for all external injuries; now, it is applied to the Linimentum Saponis Compositum.

OPODELDOCH, STEER'S. A liniment, called after the inventor. There are many formulæ for its preparation. The following is one. *Sap. alb. ñij, camphor. ʒij, ol. rosmarinum f ʒiv, spiritus vini rectificati Oij*. See Linimentum Saponis Compositum.

OPODEOCELE, see Hernia.

OPODYMUS, Diprosopus.

OPOIDIA GALBANIFERA, see Bubon Galbanum.

OPOLÉ, Succus.

OPOLIDE, Succus.

OPOPIOS, Optic.

OPOPONACUM, Pastinaca opoponax.

OPOPONAX, see Pastinaca opoponax—o. Chironium, Pastinaca opoponax.

OPOPONAXWORT, Pastinaca opoponax.

OPO'RICÉ, from *οπωπα*, 'autumnal fruits.' A medicine, composed of several autumnal fruits, particularly of quinces, pomegranates, &c., and wine. It was formerly administered in dysentery, diseases of the stomach, &c.

OPOS, Succus expressus—o. Silphion, Laserpitium.

OPPIDULUM, Vulva.

OPPILATIO, Emphraxis—o. Telæ Cellulosæ, Induration of the cellular tissue.

OPPILATIVUS, Obscruens.

OPPO'NENS, (F.) *Opponent*, from *ob*, and *ponere*, 'to place.' That which faces or is put in opposition to something. The name has been given to two muscles of the hand.

OPPONENS MIN'IMI DIG'ITII, (F.) *Opposant du petit doigt*, *Carpo-métacarpien du petit doigt* (Ch.), *Métacarpien du petit doigt*. This muscle has the same shape and arrangement as the preceding, but is of less size. It is situate in the hypothenar eminence. Its fibres are inserted into the anterior annular ligament of the carpus, and terminate on the whole length of the inner edge of the 5th metacarpal bone. This muscle carries the 5th metacarpal bone forwards and outwards, and thus augments the concavity of the palm of the hand.

OPPONENS POL'LCIS, *Flexor ossis metacarpi pollicis*, F. *primi intero'dii* (Douglass). *Opponens pollicis manûs*, *Antith'enar sive semi-interosseus pollicis*, (F.) *Carpo-métacarpien du pouce* (Ch.), *Métacarpien du pouce*, *Opposant du Pouce*.

A small, flat, triangular muscle, situate in the substance of the thenar eminence. It is attached, on the one hand, to the anterior annular ligament of the carpus and to the trapezium; and, on the other, to the whole of the outer margin of the first metacarpal bone. This muscle impresses on the first bone of the metacarpus a movement of rotation, which opposes the thumb to the other fingers.

OPPOSANT, Opponens—o. du Petit doigt, Opponens minimi digiti—o. du Pouce, Opponens pollicis.

OPPRESSIO, Catalepsy, Oppression—o. Nocturna, Incubus.

OPPRESSION, *Oppres'sio*, *Thlipsis*, *Cata-thlipsis*, from *opprimere* (*ob*, and *premere*, *pressum*), 'to press against.' A state, in which the patient experiences a sensation of weight in the part affected. When employed abstractedly, it means, particularly, *Oppression of the chest—Oppressio Pec'toris*, (F.) *O. de Poitrine*.

OPPRESSION, *Oppres'sio vir'ium*, (F.) *Oppression des forces*, is, also, used for that condition, at the commencement of fevers, &c., in which the system is oppressed rather than debilitated, and where the vascular action rises, as the obstruction to free circulation is relieved by bleeding, purging, &c.

OPS, Eye.

OPSEONUSI, Opsionusi.

OPSIALGIA, Neuralgia, facial.

OPSIONUS'I, *Opsionu'si*, from *οψis*, 'vision,' and *νοσος*, 'a disease;' *Morbi visus*. Diseases of vision.

OPSIOTOC'IA, from *οψis*, 'too late,' and *τοκος*, 'birth;' *Partus sero'tinus*. Parturition after the usual period. See Pregnancy.

OP SIS, Pupil. Vision.

OPSOMA'NIAC, *Opsom'anes*, (F.) *Opsomane*, from *οψον*, 'aliment,' and *μανια*. One who loves some particular aliment to madness.

OPTESIS, Assatio.

OPTIC, *Opt'icus*, *Op'o'pticus*, (F.) *Optique*, from *οπτοι*, 'I see.' That which relates to vision.

OPTIC CENTRE, *Centrum opt'icum*. The optic centre of the crystalline is the point at which the various rays proceeding from an object cross in their way to the retina.

OPTIC COMMISSURE, see Chiasmus.

OPTIC FORAMEN, *Fora'men opt'icum*, (F.) *Trous optique*. A round opening in the sphenoid bone, near the base of its lesser ala, through which the optic nerve passes.

OPTIC GANGLIA, Quadrigemina tubercula.

OPTIC GROOVE. A transverse groove on the superior surface of the sphenoid bone, on which the commissure of the optic nerve rests, and which is continuous on each side with the optic foramen.

OPTIC LOBES, Quadrigemina tubercula.

OPTIC NERVE, *Nervus opt'icus seu vis'ivus seu vis'o'rius*, *Par secundum seu opt'icum seu vis'o'rium*, *Nerve of the 2d pair*, *Ocular nerve of Chaussier*. The optic nerves are remarkable for their size; for their running a longer course within than without, the cranium; and for their furnishing no branch from their origin to their termination. They do not seem to arise, as was long supposed, from the optic thalami, but from the tubercula quadrigemina. Immediately after their origin, the *optic tracts*, *Tractus opt'ici*, proceed forwards; are, at first, broad and flat; but afterwards become narrower and round. In front of the fossa pituitaria, they unite and decussate, each nerve proceeding through the optic foramen

of the sphenoid bone with the ophthalmic artery, and through the *Foramen opticum chorioideæ*, to form the retina. The nerve passes to the back part of the globe of the eye, becomes narrower, and enters that organ at a flattened circular papilla—*Colliculus* seu *Papilla nervi optici*—to give rise to the nervous expansion called the retina. Between the colliculus and the yellow spot of Sömmering, a small projection or fold of the retina passes—*plica centralis retinæ*. Besides its neurilemma, the optic nerve is surrounded by a sheath, furnished by the dura mater. This accompanies it as far as the eye.

The optic nerve is the nerve of special sensibility of the eye. In the centre of the nerve, there is a *punctum cæcum*, *macula nigra*, where the central artery enters the eye, and the corresponding vein passes out. If a ray of light falls on this spot it is not appreciated.

OPTIC REMEDY, see Ophthalmic remedy—o. Thalami, Thalami nervorum opticorum—o. Tracts, see Optic nerve.

OPTICO-TROCHLÆI-SCLÉROTICIEN, Oblique, superior of the eye.

OPTICUM, Ophthalmic remedy.

OPTILOS, Eye.

OPTOMETER, from *ὀπτομαί*, 'I see,' and *μετρον*, 'a measure.' An instrument for measuring the distance of distinct vision.

OPUNTIA, Cactus opuntia—o. Vulgaris, Cactus opuntia.

OR, Gold—o. Blanc, Platinum—o. Cyanure d', see Gold—o. Divisé, see Gold—o. Hydrochlorate d', see Gold—o. Iodure d', see Gold—o. Métallique, see Gold—o. Nitromuriate d', see Gold—o. Oxide d', see Gold—o. et de Sodium chlorure d', see Gold—o. et de Soude, Hydrochlorate d', see Gold—o. et de Soude, Muriate d', see Gold.

ORA SERRATA, see Ciliary (Body.)

ORACHE, SPREADING, Atriplex patula—o. Stinking, Chenopodium vulvaria.

ORAL, Vocal, (F.) Oral, from *os*, oris, 'a mouth.' Relating to the mouth or to speech.

ORANGE, Citrus aurantium—o. Bitter, see Citrus aurantium—o. Flowers of the, see Citrus aurantium.

ORANGE GRASS, Hypericum sarothra—o. Root, Hydrastis Canadensis—o. Tree, Citrus aurantium.

ORANGEADE. A drink, made with orange juice diluted with water. It is antiphlogistic, and often recommended in acute diseases.

ORANGES, see Citrus aurantium—o. Curasson, Aurantia curassaviana.

ORBES CARTILAGINOSI TRACHEÆ, see Trachea.

ORBITULAIRE DES LÈVRES, Orbicularis oris—o. des Paupières, Orbicularis palpebrarum.

ORBICULAR, Orbicularis, (F.) Orbiculaire, from *orbis*, 'a circle.' Spherical, circular.

ORBICULAR BONE, *Os orbicularè seu lenticula-rè seu Sylvii seu orbicularè Syll'vii*, *Globulus stap'edis ossis*, *Epiph'ysis curvis longioris incis-dis*, *Ossiculum orbicularè seu squamosum seu cochlearè seu quartum*, is the smallest of the four bones of the ear. It is scarcely perceptible, round, convex on two surfaces, and situate between the long ramus of the incus and the head of the stapes.

ORBICULAR MUSCLES are muscles with circular fibres surrounding some natural opening of the body. 1. *Orbicularis seu Constrictor Oris*, *Sphincter Labiorum*, *Semi seu Demi-orbicularis*, *Supra-semi-orbicularis*, *Labialis*, *Osculato-rius*, *Buccinator*, (F.) Labial, Orbiculaire des lèvres. A muscle situate in the substance of the lips, and extending from one commissure to the other. It is formed of two very distinct portions, of a semi-oval shape; one belonging to the upper lip,

the other to the lower. Their extremities cross at the commissures, and are confounded with the other muscles of the parts. The use of this fleshy muscle is to bring the lips together, and to close the aperture of the mouth, by making it represent a sort of *bourrelet* with radiated wrinkles. It is an antagonist to every other muscle of the lips. 2. *Orbicularis Palpebrarum seu oculi seu palpebrarum ciliaris*, *Palpebrarum duo mus-culi*, *Palpebralis*, *Constrictor seu Sphincter Palpebrarum seu Oculi*, (F.) *Naso-palpebral* (Ch.), *Maxillo-palpebral*, *Palpebral*, *Orbiculaire des paupières*. A muscle common to both the eyelids, and seated in their substance. It is broad, thin, transversely oval, and cleft in its great diameter. It is attached to the nasal process of the superior maxillary bone; to the internal angular process of the frontal bone, and to the orbital process of the superior maxillary bone. From these origins the muscle passes upwards, under the skin of the eyelids, surrounding the orbit in a circular manner, extending somewhat beyond it, and covering the upper part of the cheek. It is inserted into the skin of the eyelids, its upper and inner edge being intimately connected with the frontal and corrugator muscles. Its use is to close the eye, by bringing the eyelids together. The part of the orbicularis which covers the cartilages of the eyelids, and which is remarkably thin, is the *Musculus Ciliaris* of some authors.

ORBICULARIS OCULI, Orbicularis palpebrarum, see Orbicular muscles—o. Oris, see Orbicular muscles—o. Palpebrarum, see Orbicular muscles—o. Recti, Sphincter an externus.

ORBICULUS CILIARIS, Ciliary ligament, Ciliary zone.

ORBIS GENU, Patella.

ORBIT, *Or'b'ita*, *O. oculi*, *Troch'ia*, *Conchus seu Cavitas seu Fovea seu Pelvicula oculi*, (F.) *Orbite*, from *orbis*, 'a circle.' The circular cavities are so called, which lodge the organs of sight. The *orbits* or *orbital fossæ* or *cavities*, *conchi*, *κογχαι*, are situate at the upper part of the face, and are composed of seven bones, viz.:—the frontal, above; the palate and superior maxillary, below; the sphenoid and malar, externally; and the ethmoid and lacrymal, internally. The orbit is filled by the globe of the eye, with its different accessory parts—muscles, nerves, vessels, the lacrymal gland, &c. Its margin is termed *margo orbita'lis*.

ORBITA, Orbit—o. Oculi, Orbit.

ORBITAL, Orbital—o. Fissure, inferior, Sphenoidal fissure—o. Fissure, superior, Maxillary fissure.

ORBITAR, *Orbitarius*, *Orbita'lis*, *Or'bital*, (F.) *Orbitaire*. Same etymon. Relating or belonging to the orbit.

ORBITAR APOPH'YSES or PROC'ESSES terminate the two extremities of the orbital arch. The external is much more prominent, and is articulated with the cheek-bone; the internal is thinner, and joins the os unguis.

ORBITAR ARCH, (F.) *Arcade orbitaire*, is the projecting, round margin, which separates the frontal and orbital surfaces of the os frontis, and forms a part of the circumference of the orbit.

ORBITAR FISSURES, (F.) *Fentes orbitaires*. These are situate in the orbit. They are two in number: one superior, the Sphenoidal; the other inferior, the Sphenomaxillary.

ORBITAR FORAM'INA. Of these there are several.—1. The *Foramen orbitarium superius*, (F.) *Trou orbitaire supérieur*, which is situate at the junction of the inner third of the orbital arch with its two outer thirds, and gives passage to the frontal nerve. 2. The *Foram'ina orbitaria interiora seu inter'na*, *Foram'ina Ethmoidea*.

These are situate at the upper part of the internal paries of the orbit, are formed by the union of two notches of the os frontis with two similar ones of the ethmoid; and are distinguished into *anterior* and *posterior*. The former gives passage to the ethmoidal filament of the nasal nerve and to a small artery.

ORBITAR NERVE, *Nervus orbitæ* seu *orbitalis* seu *subcutaneus male*, is a branch given off from the superior maxillary. It enters the orbit by the spheno-maxillary fissure, and divides into two filaments; the one — *molar* — which passes over the cheek-bone, is distributed to the orbicularis palpebrarum, and anastomoses with the facial nerve; the other — *temporal* — which passes to the temporal fossæ, and anastomoses with the inferior maxillary and facial, to become afterwards subcutaneous.

ORBITAR SURFACE, see Maxillary bone (superior).

ORBITO-EXTUS-SCLÉROTICIEN, Rectus externus oculi — *o. Intus-Scléroticien*, Rectus internus oculi — *o. Maxillo-labii*, Levator labii superioris proprius — *o. Palpebræ*, Levator palpebræ superioris — *o. Sus-palpebræ*, Levator palpebræ superioris.

ORCANETTE, Anchusa officinalis.

ORCHALGIA, Orchidalgia.

ORCHEA, Scrotum.

ORCHETIS, *Oschetis*, *Oschetis*, *Inflammati*o *scroti*, from *orchis*, 'the scrotum,' and *itis*, denoting inflammation. Inflammation of the scrotum.

ORCHETIS, Hernia humoralis.

ORCHEOCATABASIS, Orchidocatabasis.

ORCHEOCELE, Orchiocele.

ORCHEOTOMY, Castration.

ORCHESTROMANIA, Chorea.

ORCHIDAL'GIA, *Orchial'gia*, *Orchidodyn'ia*, from *orchis*, *orchidos*, 'a testicle,' and *algos*, 'pain.' Pain of the testicle. Neuralgia of the testicle. Irritable testicle. A form of neuralgia.

ORCHIDATROPH'IA, *Atroph'ia testicul'i*, from *orchis*, *orchidos*, 'a testicle,' and *atrophia*, 'atrophy.' Atrophy or wasting of the testicle.

ORCHIDITIS, Hernia humoralis.

ORCHIDOCARPUM ARIETINUM, see *Carica papaya*.

ORCHIDOCATAB'ASIS, *Orcheocatab'asis*, *Descen'sus testiculo'rum*, from *orchis*, *orchidos*, 'the testis,' and *καταβασις*, 'descent.' The descent of the testes into the scrotum.

ORCHIDOCELE, Hernia humoralis.

ORCHIDODYN'IA, Orchidalgia.

ORCHIDONCUS, Hernia humoralis.

ORCHIDOTOMY, Castration.

ORCHILL, Lichen roccella.

ORCHIOCE'LE, *Orcheoce'le*, from *orchis*, *orchidos*, 'the testicle,' and *κλην*, 'a rupture.' A tumour of the testicle. Scrotal hernia. Hernia humoralis. A name given to several diseases of the testicle and its envelopes, essentially differing from each other.

ORCHIOCE'LE SCIRRHO'SA, *Scirrhoc'e'le*. A malignant tumour of the testis.

ORCHIS, Testicle — *o. Butterfly*, *Orchis bifolia*.

ORCHIS MAS'CVLA, *Saty'rion*, *Testicul'us caninus*, *Cynosor'chis*, *Dogs' stones*, *Male orchis*, *Bal-lock-grass*, *Male Fools' stones*, *Twinroot*. *Ord. Orchidaceæ*. This root had a place in some pharmacopœias, on account of the glutinous, slimy juice it contains. The root of the *Orchis bifolia* or *Butterfly orchis*, has likewise been collected. It has been used as a demulcent. Its fancied aphrodisiac virtues seem to be owing to its resemblance to *orchis*, 'a testicle,' whence its name.

Salap is obtained from this variety of the orchis; — and from the *Orchis morio*, &c. See *Salap*.

ORCHIS, ROUND-LEAVED, LARGE, *Platanthera orbiculata*.

ORCHITE, Hernia humoralis.

ORCHITIS, Hernia humoralis.

ORCHOI, see Tarsus.

ORCHOTOMIA, Castration. Also, removal of the tarsi.

ORDEAL NUT — called *Esère* by the natives of Calabar — of the natural family Leguminosæ, and closely resembling a *Dolichos*, is possessed of virulent acro-narcotic properties.

ORDEOLUM, Hordeolum.

ORDEUM, Hordeum.

ORDINAIRES, Menses.

ORDO MEDICINALIS, Police, medical.

ORDONNANCE, Formula, Prescription.

ORELLE D'ÂNE, Symphytum — *o. d'Homme*, Asarum — *o. de Souris*, *Hieracium pilosella* — *o. d'Ours*, *Primula veris*.

ORELLETTE, Asarum, Auricle of the heart.

O'RELLONS, Parotis, Cynanche parotidæa.

O'RELIA GRANDIFLORA, Allamanda.

O'RELLANA, see Terra Oroleana.

O'REOSELINUM, *Athamanta aureoselinum* — *o. Africanum*, *Bubon galbanum* — *o. Legitimum*, *Athamanta aureoselinum* — *o. Nigrum*, *Athamanta aureoselinum*.

O'REXIS, Appetite, Pyrosis — *o. Cynodes*, *Boulimia*.

ORGAN, *Organum*, *Or'ganon*. From *εργον*, 'work.' 'An instrument.' Part of an organized being, destined to exercise some particular function. Thus — the eyes are organs of sight; the muscles are organs of motion, &c. The collection of organs, which concur in any function, is called *apparatus*.

ORGAN, *Mentha pulegium*.

ORGANA GENERATIONI INSERVIENTIA, Genital organs — *o. Genitalia*, Genital organs — *Sexualia*, Genital organs — *o. Sudoripara*, see Perspiration — *o. Urogenitalia*, Urogenital organs.

ORGANES GÉNITAUX, Genital organs.

ORGAN'IC, *Organ'icus*, (F.) *Organique*. Relating to an organ or organs, and to beings possessed of organs. Hence, **ORGANIC FUNCTIONS** are those possessed by both animals and vegetables. It is also used synonymously with *vital*, in contradistinction to *physical*.

ORGANIC CHEMISTRY, see *Chymistry*.

ORGANIC DISEASES or LESIONS are such as concern the organ itself, in contradistinction to *functional*, *rat'ional*, or *dynam'ic*, which merely concern the function. Of course, *organic* diseases of a part are always more serious than *functional*.

ORGANIC NERVQUS SYSTEM, see *Trisplanchnic nerve*.

OR'GANICISM, *Organicism'us*, from *οργανος*, 'an organ.' The doctrine of the localization of disease, or which refers it always to a material lesion of an organ. A modern term, but not much used.

ORGANISATIO, Organization.

ORGANISATUS, Organized.

OR'GANISM. Same etymon as organ. The living economy. Many physiologists have used this word synonymously with organization; but more especially to designate the aggregate of vital actions in organized beings, animals as well as vegetables. The collection of parts composing an organized body, and the laws which govern it.

ORGANIZABILITY, from *organisatio*, and *habilis*, 'suitable.' Capability of organization or of being formed into living tissue. Substances possessed of such capability — as fibrin — are said to be *organizable*.

ORGANIZABLE, see *Organizability*.

ORGANIZA'TION, *Organisat'io*, from *οργανος*, an organ.' The manner of structure proper to different beings. It is also employed in the sense of the structure of a part; as, the organization of the heart, &c.

ORGANIZATION, COMPOUNDS OF, Principles, immediate.

OR'GANIZED, *Or'ganis instructus*, *Organisatus*, (F.) *Organisé*. That which is composed of organs.

ORGANIZED BODIES, (F.) *Corps organisés*, are those endowed with life; i. e. animals and vegetables: in contradistinction to *inorganic* bodies, which comprise the mineral kingdom. See *Corpus*.

ORGANO-CHEMIA, Chymistry, organic.

ORGANOGEN'Y, *Organogen'ia*, *Organogen'esis*, from *οργανος*, 'an organ,' and *γενεσις*, 'generation.' The doctrine of the formation of different organs.

ORGANOGENO'SIA, from *οργανος*, and *γνωσις*, 'knowledge.' The knowledge and discrimination of organized bodies.

ORGANOGRAPHY, *Organograph'ia*, from *οργανος*, and *γραφη*, 'a description.' A description of the organs of a living body.

ORGANOLEPTIC, Plastic.

ORGANOLOGY, Anatomy, Craniology.

ORGANON OLFACTUS, Nasus.

ORGANONOM'IA, from *οργανος*, and *νομος*, 'a law.' The doctrine of the laws of organic life.

ORGANOZOONOMIA, Zoonomia.

ORGANUM, Instrument.

ORGASM, *Orgas'mus*, *Orgas'is*, from *οργαζω*, 'I desire ardently,' 'I am excited.' *Æstru'dition*. A strong impulse or desire for something. State of excitement and turbulence of any organ whatever. Erethism. Applied particularly to the height of venereal excitement in sexual intercourse.

ORGASMUS, Turgescence.

ORGAS'TICA. Same etymon. The second order of the class *Genetica* of Good. Defined: diseases affecting the orgasm; organic or constitutional infirmity disordering the power or the desire of procreating.

ORGE, Rage.

ORGE, Hordeum—*O. Perlé*, see Hordeum.

ORGEAT, Syrupus amygdalæ.

ORGELET, Hordeolum.

ORGEOLET, Hordeolum.

ORGOSIS, Orgasm.

ORICHALCUM, Brass.

ORICULAR, Auricular.

ORICULE, Auricle, Pavilion of the ear.

ORIFICIUM, Mouth—*O. Dextrum Ventriculi*, Pylorus—*O. Uteri*, Os uteri—*O. Uterinum*, see Uterus—*O. Vaginale*, see Uterus—*O. Ventriculi Sinistrum*, Cardia.

ORIG'ANUM, *O. vulga'rē* seu *heracleoticum*, (*Op'yavon*, from *opos*, 'a mountain,' and *yavos*, 'delight.') *Cuni'lia bu'bula*, *Majora'na mancure'na*, *Zazarhen'di herba*, Wild or Common marjoram, *Mancure'na*, (F.) *Pied de lit*. Ord. Labiatae. *Sex. Syst.* Didynamia Gymnospermia. This plant resembles marjoram in smell; and has a pungent taste, much like thyme. It is aromatic and stimulant; and its oil possesses the properties of the essential oils in general. The oil, *Öleum origani* (Ph. U. S.), is very pungent, and is rarely given internally. It is used in toothache.

ORIGANUM AQUATICUM, Eupatorium cannabium—*O. Creticum*, *O. dictamnus*.

ORIGANUM DICTAM'NUS, *O. Cret'icum*, *Dictam'nus Cret'icus*, *Amar'acius tomento'sus*, *Oni'tis*, *Di'tany* of Crete, (F.) *Dictanne* de Crète. The leaves of this plant were once recommended as emmenagogue and alexipharmic.

ORIGANUM HERACLEOTICUM, Origanum.

ORIGANUM MAJORA'NA, *O. majoranaides*, *Majorana*, *M. horten'sis*, Sweet mar'joram, *Scamp'sucus*, *Agriorig'anum*, *Amar'acius*, (F.) *Marjolaine*. Its odour is strong and fragrant; taste aromatic and bitterish. Its properties are those of a stomachic and errhine. It is chiefly used, however, for culinary purposes, and as a snuff in headache.

ORIGANUM SYRIACUM, Teucrium marum.

ORIGINAIRE (F.), from *origo*, *originis*, 'origin.' This term has been used, by French writers, in the same sense as *congenital*. It is particularly applied to affections depending upon faulty conformation, or to faulty conformation itself.

ORIGO, Arche.

ORISMOLOGY, Terminology.

ORLEANA, see Terra Orleana.

ORME, Appetite.

ORME, Ulmus—*O. Champêtre*, Ulmus—*O. Pyramidale*, Ulmus.

ORMEAU, Ulmus.

ORMSKIRK MED'ICINE. A celebrated antylissic remedy, supposed to consist of *Pulv. Crete*, ʒss; *Bol. Armen.* ʒiij; *Alumin.* gr. x; *Pulv. Inulæ*, ʒj; *Ol. Anisi*, gtt. vj.

This is taken for a dose every morning for six times, in a glass of water, with a small proportion of fresh milk.

ORNAMENTUM FOLIACEUM, see Tube, Fallopian.

ORNITHOG'ALUM ALTIS'SIMUM. (*Orni-thogalon*, a fanciful name, from *ornis*, *ornis*, 'a bird,' and *gala*, 'milk.') A South African plant, *Nat. Ord.* Asphodeleæ, the fleshy bulb of which is diuretic. An oxymel is prepared of it, which is given in catarrh, asthma, phthisis, and hydrothorax. It resembles squill in its medical properties.

ORNITHOGALUM MARITIMUM, Scilla—*O. Squilla*, Scilla.

ORNITHOGLOSSA, see Fraxinus excelsior.

ORNUS EUROPEA, see Fraxinus ornus—*O. Mannifera*, *Fraxinus ornus*—*O. Rotundifolia*, *Fraxinus ornus*—*O. Sylvestris*, *Fraxinus excelsior*.

OROBAN'CHĒ VIRGINIA'NA, (*οροβαγχη*, from *ορος*, 'the orobus,' and *αγχω*, 'I strangle.') *Epiphe'gus* seu *Epifa'gus America'nus* seu *Virginia'nus*, *Virgin'ia broom-rape*, *Beech-drops*, *Cancer-root*. Family, *Orobanchoides*. This parasitic plant is astringent, and a peculiar and extremely nauseous bitter. It is most powerful when fresh. It has been used in dysentery; and, externally, to obstinate ulcers.

OROBAN'CHĒ AMERICA'NA, and *O. UNIFLO'RA*, have the same properties, and are, likewise, called *Cancer-root*, *Earth-club*, and *Clapwort*.

OROBĒ, Ervum ervilia.

OROBION; see Ervum ervilia.

OROBOD'ES, from *ορος*, 'the orobus,' and *ειδος*, 'form.' *Orobo'des*. Having the shape of the orobus. A name given to urine, *Urina orobo'des* seu *orobod'is*, when depositing a substance of a fawn colour, like the meal of the orobus.

OROBUS, Ervum ervilia—*O. Faba*, *Vicia faba*.

ORONGE, see Agarie.

OROPHALL'US, from *opos*, or *oppos*, 'the rump,' and *φαλλος*, 'the male organ.' A monster having a second male organ originating from the rump.—Gurli.

OROS, Crotonion, Serum.

OR'PIMENT, *Auripigmen'tum*, *Orpin*, *Risigallum*, *Risigallum*, *Oloanz*, *Elomē*, *Sira*, *Lempnia*, *Spec'ulum citrinum*, *Yrides*, *Yride*. The native yellow sulphuret of arsenic. It is poisonous, but less so than the oxide. It is also, called *Adarigo*, *Adarnech*, *Ahusal*, *Althana*, *Azarnot*.

The Turks prepare a depilatory with it, which they call *Rusma*.

ORPIN, Orpiment.—*o. Brûlant*, Sedum—*o. Grand*, Sedum telephium—*o. Reprise*, Sedum telephium.

ORPINE, Sedum telephium.

ORRHOCHEDIA, Diarrhoea, serous.

ORRHOCYSTIS, from *oppos*, 'serum,' and *κυστις*, 'a bladder.' An encysted tumour containing a serous fluid.

ORRHOHYMENITIS, Hydrohymenitis.

ORRHOPYG'ION. The inferior extremity of the vertebral column; from *opos*, 'extremity,' and *πυγη*, 'the nates.' The os coccygis. Also, the raphe, extending from the penis to the anus, and separating the scrotum into two parts.

ORRHORRHE'A, from *oppos*, 'serum,' and *ρω*, 'I flow.' A morbidly increased secretion of a serous fluid.

ORRHOS, Croupion, Serum.

ORRIS, Iris Florentina—*o. Common*, Iris Germanica.

ORSEILLE, Lichen roccella.

ORTEIL, Digtus pedis.

ORTHOCOL'UM, from *oppos*, 'straight,' and *κωλον*, 'limb.' Stiffness of the limb, so that it is kept quite straight.

ORTHODON'TIA, from *oppos*, 'straight, right,' and *odontos*, 'a tooth.' The part of dentistry which is concerned in the treatment of irregularity of the teeth.

ORTHODO'RON. A measure with the Greeks, equal to eleven finger-breadths.

ORTHOG'NATHOUS, from *oppos*, 'right, straight,' and *γναθος*, 'jaw.' Having a vertical jaw. A term applied to the form of head in which the facial angle approaches the right angle.

ORTHOMORPHIA, Orthopædia.

ORTHOMORPHOSIS, Orthopædia.

ORTHOTROPIA, Orthopædia.

ORTHOPEDIA, *Orthopæ'dicē*, *Orthomor'phia*, *Orthomorpho'via*, *Orthokotrop'ia*, *Orthopæ'dies*, from *oppos*, 'right,' and *παιδος*, 'a child.' The part of medicine whose object is to prevent and correct deformity in the bodies of children. Often used, however, with a more extensive signification, to embrace the correction or prevention of deformities at all ages. *Orthosomat'ica*, *Orthosomat'icē*, from *oppos*, 'right,' and *σωμα*, 'body,' has been proposed as a preferable term.

ORTHOPE'DIC, *Orthopæ'dicus*; same etymon. Relating to orthopædia,—as *Orthopedic Institution*, *Institutum orthopæ'dicum* seu *orthopæ'dium*: an institution for the correction or prevention of deformities in children.

ORTHOPE'DICS, Orthopædia.

ORTHOPE'DIST, *Ortho'ter*, *Orthopæ'dicus*. Same etymon. One who practises orthopædia.

ORTHOPHO'NIA, from *oppos*, 'right,' and *φωνη*, 'voice.' A good pronunciation.

ORTHOPHREN'IC, *Orthophren'icus*, from *oppos*, 'right,' and *φρον*, 'mind.' Relating to sound mind, and to the cure of insanity.

ORTHOPNE'A, *Suffoca'tio*, *Strangula'tio*, *Præfoca'tio*, *Pnix*, *Pnigmus*, *Euthyp'noē*, *Euthypno'e'a*, *Angor*, *Apno'e'a*, from *oppos*, 'right,' 'straight,' and *πνέω*, 'I respire.' Impracticability of breathing in the horizontal posture. Necessity of being in the erect posture, in order to respire.

ORTHOPNEA CARDIACA, Angina Pectoris—*o. Convulsiva*, Asthma—*o. Cynanchica*, Cynanche trachealis—*o. Hydrothoracica*, Hydrothorax—*o. Hysterica*, Angone—*o. Puituosa*, Mycorthopneæ—*o. Tussiculosa*, Pertussis.

ORTHOSOMATICE, Orthopædia.

ORTHOSOMATIQUE, Orthopædia.

ORTHOTER, Orthopædist.

ORTHYSTEROPTOMA, Prolapsus uteri.

ORTHYSTEROPTOSIS, Prolapsus uteri.

ORTIE, Urtica—*o. Blanche*, *Lanum album*—*o. Brûlante*, *Urtica urens*—*o. Morte*, *Lanum album*—*o. Morte des Bois*, *Galeopsis*—*o. Romaine*, *Urtica pilulifera*.

ORTYX, Tetrao coturnix.

ORUS, Croupion, Serum.

ORVALE, Salvia sclarea.

ORVIETA'NUM. From the Italian *Orvietano*. An electuary regarded as an invaluable antidote, and composed of *Old theriac*, *Dried vipers*, *Scorzonera*, *Carlina*, *Imperatorium*, *Angelica*, *Bistort*, *Rosemary*, *Juniper*, *Cinnamon*, *Cloves*, *Mace*, *Honey*, and a crowd of other substances. Its medical properties resembled, in some respects, those of the theriac. It was so called, according to some, because invented by Orvietano, a celebrated charlatan; or, according to others, from Orvieto, a town in Italy.

ORY'ZA, *Opu'za*, *Ory'za sativa*, *Rice*, (F.) *Riz*. Family, Gramineæ. *Sex. Syst.* Hexandria Monogynia. A plant, almost aquatic, and indigenous in China, one of the richest productions of Egypt, and which prospers in the warmer regions of the four quarters of the globe. The grain inclosed in the husk is called by the Malays *Paddy*, *Padi*, or *Paddie*; when deprived of the husk, *Bras*, and when boiled, *Nasi*. Its amylaceous seeds, *Ory'za*, *Ory'zum*, *Hordeum galactium*, nourish more men than those of wheat and rye together. Besides being alimentary, they are demulcent and emollient. *Gardiner's* alimentary preparation is very finely ground rice-meal.

Mucilage of Rice, *Rice water*; may be formed by taking one ounce of rice, and, having washed it, macerating it for three hours in a quart of tepid soft water, in a pan placed upon a stove; then boiling the whole slowly for another hour, and straining through a sieve.

It may be sweetened and acidulated, or be prepared with milk, like arrow-root, and be used as a demulcent.

Rice Jelly is made by macerating a sufficient quantity of clean rice in as much water as will cover it; boiling slowly, adding water as it evaporates, until the rice is reduced to a pap, sweetening and flavouring with lemon-juice or vanilla, and passing through a fine sieve into a form or mould. It may be eaten alone or with milk; and is a good dietetic preparation for those of weak digestive powers.

Ground rice. Take a tablespoonful of ground rice, a pint and a half of milk, and half an ounce of candied lemon-peel. Rub the rice smooth with the milk: then add the lemon-peel cut into small pieces; boil for half an hour, and strain whilst the milk is hot. It is used in the same cases as rice jelly.

OS, Bone, Mouth, see Ossiculum.—*o. Aeromii*, Aeromion—*o. Adjutorium*, *Humeri os*—*o. Alæforme*, *Sphenoid*—*o. Alagas*, *Coccyx*, *Sacrum*—*o. Alatum*, *Sphenoid bone*—*o. Amphideon*, *Os uteri*—*o. Anonyme*, *Innomminatum os*—*o. Anonymum*, *Innomminatum os*—*o. Anticum*, *Avant bouche*—*o. Arcuale*, *Temporal bone*—*o. Armale*, *Temporal bone*—*o. Asser*, *Sternum*—*o. de l'Assiette*, *Ischion*—*o. Azygos*, *Sphenoid*—*o. Balliste*, *Astragalus*—*o. Basilare*, *Occipital bone*, *Palate bone*, *Sacrum*, *Sphenoid*—*o. Basillare*, *Sphenoid*—*o. Bicornæ*, *Hyoides os*—*o. Brachiale*, *Humeri os*—*o. Brachii*, *Humeri os*—*o. Calcis*, *Calcaneum*—*o. Capitatum*, *Magnum os*—*o. Caudæ*, *Occipital bone*—*o. Clunium*, *Sacrum*—*o. Coccygis*, *Coccyx*—*o. Colatorii*, *Sphenoid bone*—*o. Colatorium*, *Ethmoid bone*—*o. Coliforme*, *Ethmoid*—*o. Convolutum superius*, *Turbinated bone*, *middle*—*o. Coronale*, *Frontal bone*—*o. Coxæ*, *Innomminatum os*—*o. Coxal*, *Innomminatum os*, *Ischion*—*o. Cribuleus*, *Ethmoid*—*c. Cribri*.

forme, Ethmoid — o. Cribrosum, Ethmoid — o. Cristatum, Ethmoid — o. *Crochu*, Os unciforme.

OS CROTAPHALĒ. A name given by M. Bèlard to a bone sometimes met with at the anterior and inferior angle of the parietal bone; from *κροτάφος*, 'the temple.'

Os CROTAPHITICUM, Temporal bone — o. Cubicum, Ethmoid bone — o. ad Cubitale, Radius — o. Cubiti Inferius, Ulna — o. *de la Cuisse*, Femur — o. Cuneiforme, Sphenoid — o. Cuneo comparatum, Sphenoid — o. Cymbiforme, Scaphoid bone — o. Disciforme, Patella — o. Durum, Temporal bone — o. Ensiforme, Sternum.

OS ÉPACTALĒ. A name given to one of the ossa Wormiana; particularly to one met with in the posterior fontanelle, and which has been described by G. Fischer, under the name *Os Épactalē seu Gœthia* num.

OS ÉPACTAUX, Wormiana ossa.

Os EXTERNUM. The entrance into the vagina, in contradistinction to the *Os internum* or *Os uteri*.

Os EXTRA ORDINEM CARPI, Os pisiforme — o. Femoris, Femur — o. Fibrosum, Occipital bone — o. Foraminulentum, Ethmoid — o. Frontis, Frontal bone — o. Genæ, Mala os — o. Gladioli, Sternum — o. *Grand*, Magnum os — o. *Grand*, col du, Collum ossis magni — o. Grandiosum, Cuboid — o. Gutturis, Hyoides os — o. Hamatum, Os unciforme — o. *Hors du rang*, Os pisiforme — o. Hypopium, Mala os — o. Hypseloides, Hyoides os — o. Inconjugatum, Sphenoid — o. Incendi similis, Ineus — o. Inferius ventriculi, Pylorus — o. *Innominé*, Innominatum os — o. *Intercalé*, Wormiana ossa — o. Internum, see Uterus — o. Inverecundum, Frontal bone — o. Ischii, Ischion — o. Interparietale, Interparietal bone — o. Jugale, Mala os — o. Jugamentum, Mala os — o. Juguli, Clavicle — o. Lacrymale, Unguis os — o. Lambda, Occipital bone — o. Lambdoides, Hyoides os, Occipital bone — o. Lapidum, Temporal bone — o. Lapidosum, Temporal bone — o. Latum, Sacrum — o. Latum humeri, Scapula — o. Lenticulare, Os orbiculare, Os pisiforme — o. Lepidoides, Temporal bone — o. Lingue, Hyoides os — o. Linguale, Hyoides os — o. Lithoides, Temporal bone — o. Magnum, neck of the, Collum ossis magni — o. *Malare*, Mala os — o. *Malare*, Mala os — o. Maxillæ superioris undecimum, Vomer — o. *Maxillaire inférieure*, Collum mandibulae — o. Memento mori, Temporal bone — o. Memoria, Occipital bone — o. Mendosum, Temporal bone — o. Morsus Adami, Thyroid cartilage — o. Mucronatum, Xiphoid cartilage — o. Multangulum majus, Trapezium os — o. Multangulum minus, Trapezoides os — o. Multiforme, Ethmoid, Sphenoid — o. *Nasaux*, Nasal bones — o. Navicular, Os scaphoides — o. Nervale, Temporal bone — o. Nervosum, Occipital bone — o. Occipiti, Occipital bone — o. Orbiculare, Os pisiforme — o. *du Palais*, Palate bone — o. Palati, Palate bone — o. *Palatin*, Palate bone — o. Papillare, Sphenoid — o. Parietale inferius, Temporal bone — o. Paxillare, Sphenoid — o. Pectinis, Pubis, os — o. Pectoris, Sternum — o. Pelvi-cephalicum, Occipital bone — o. Pelvis laterale, Innominatum os — o. Perone, Fibula — o. Petrosum, Temporal bone — o. Polymorphon, Sphenoid — o. *de la Pommette*, Mala os — o. Posterum, Pharynx — o. Prærupti rupi assimilatam, Temporal bone — o. Procutibale, Ulna — o. *Propres du nez*, Nasal bones — o. Proræ, Frontal bone, Occipital bone — o. Pterygoideum, Sphenoid — o. Pudicum, Mala os — o. Puppis, Frontal bone, Occipital bone — o. Pyramidale carpi, Trapezoides os — o. Pyxidis, Occipital bone — o. Rationis, Frontal bone — o. *du Rayon*, Radius — o. Sacrum, Sacrum — o. Saxeum, Temporal bone — o. Scutiforme, Patella — o. *de Sèche*, Sepia — o. Sedentarium, see Ischiatic — o. Semilunare, Lunare os — o. Sextum *transi*, Occipital bone — o. *du Sinciput*, Parietal

bones, see Sinciput — o. *Sous-ethmoidal*, Turbinate bone, inferior — o. Spheno-basilar, Occipital bone — o. Sphecoideum, Sphenoid — o. Sphe-noidale, Sphenoid — o. Sphenoides, Sphenoid — o. Spongiosum, Ethmoid — o. Spongiosum inferius, Turbinate bone, inferior — o. Spongiosum medium, Turbinate bone, middle — o. Spongiosum superius, Turbinate bone, middle, Turbinate bone, superior — o. Spongoides, Ethmoid — o. Squamosum, Temporal bone — o. Suboculare, Mala os — o. *Surnuméraires*, Wormiana ossa — o. *Supramaxillaire*, Maxillary (superior) bone — o. Sylvii, Os orbiculare — o. Syncipitis, Frontal bone — o. Temporis, Temporal bone — o. Tesserae, Astragalus, Cuboid — o. Thyroides, Patella — o. Tibia minus, Fibula — o. Tineæ, Os uteri — o. *Triangulaires*, Wormiana ossa — o. Turbinatum, Turbinate bone, middle — o. Turbinatum inferius, Turbinate bone, inferior — o. Uncinatum, Os unciforme — o. Upsilonoides, Hyoides os.

Os UTERI, *Os tinea* seu *internum* seu *amphidæon* seu *amphidæum*, *Protosporus*, *Oscheon*, *Hysterosoma*, *Os culum* seu *orificium uteri*, *Hysterosoma*, *Hysterosoma*, (F.) *Museau de Tanche*. The mouth of the womb, see Uterus.

Os UTERI EXTERNUM, see Uterus — o. Uteri internum, see Uterus — o. Varium, Cuboid — o. Vesiforme, Sphenoid — o. Vomeris, Vomer — o. *Wormiens*, Wormiana ossa — o. Xiphoides, Sternum — o. Ypseloides, Hyoides os — o. Zygomatium, Mala os.

OSCEOD, Yawning.

OSCHÆMATEDE'MA, *Oschæmæde'ma*, *Edema scroti cruentum*; from *σσχον*, 'scrotum,' 'æμα,' 'blood,' and *οίδημα*, 'œdema.' Effusion of blood into the scrotum.

OSCHËMEDEMA, *Oschæmatœdema*.

OSCHE, Scrotum.

OSCHEITIS, Orchitis, Oschitis.

OSCHËOCARCINOMA, Cancer, chimney-sweepers'.

OSCHËOCE/LĒ, *Oschœce'le*, *Scrotoce'le*, from *σσχον*, 'the scrotum,' and *κηλη*, 'tumour'; *Scrotal hernia*, see Bubonocœle. Sauvages uses *oschœcele* in the sense of *Dropsy of the scrotum*, see Hydrocele.

OSCHËOCELE AQUOSA, *Hydroschœcele* — o. Flatulenta, *Physocele* — o. Hydatidosa, *Hydatidocœle* — o. *Œdematica*, *Œdematoschœcele* — o. Seminalis, *Spermatocele* — o. Urinalis, *Urocele* — o. Varicosa, *Varicocele*.

OSCHËOCHALASIE, from *σσχον*, 'the scrotum,' and *χαλασις*, 'relaxation.' A tumour resulting from hypertrophy of the areolar tissue of the scrotum, and the excessive extension of the skin of the part, — an affection which has been confounded with *sarcocele*. — Alibert.

OSCHEON, Scrotum, Os uteri.

OSCHEON'CUS, *Oschon'cus*, *Oschœophy'ma*, *Oschœophy'ma*, from *oscheon*, and *αγκος*, 'a tumour.' A preternatural swelling of the scrotum.

OSCHËOPHYMA, *Oschœoncus*.

OSCHËOPLAS'TIC, *Oschœoplas'ticus*, from *oscheon*, and *πλαστικός*, 'I form.' An epithet given to the operation for restoring the scrotum when lost. *Oscheoplasty*.

OSCHËOPYEDE'MA, *Oschœopyæde'ma*, *Edema scroti purulentum*. Purulent tumefaction of the scrotum.

OSCHEUS, Scrotum.

OS'CHION. The raised margin of the os uteri. According to some, an excrescence from the os uteri.

OSCHI'TIS, *Orchei'tis*, *Oschœi'tis*, *Inflammatio scroti*, from *oscheon*, and *itis*, 'inflammation.' Inflammation of the scrotum.

OSCHOCARCINOMA, Cancer, chimney-sweepers'.

OSCHËOCELE, *Oschœcele*, *Scrotocele*.

OSCHONCUS, *Oschœoncus*.

OSCHOPHYMA, Oscheoncus.

OSCHOPYEDEMA, Oscheopyœdema.

OSCHURŒDE'MA; from *oscheon*, *οσχον*, 'urine,' and *œdema*, 'œdema.' Œdema from effusion of urine into the scrotum.

OSCHUS, Scrotum.

OSCHYDREDEMA, from *oscheon*, 'œdema,' and *œdema*, 'œdema.' Œdema scroti *aquœsum*. Watery œdema of the scrotum.

OSCILLA'TIO, Irritability, Oscillation.

OSCILLA'TION, *Oscilla'tio*; from *oscillum*, 'an image, perhaps of Bacchus, hung on ropes, and swung up and down in the air.' Vibration, *Vibra'men*, *Vibra'tio*, *Vibra'tus*. Also, a partial rotatory movement of the eyeball to and from its antero-posterior axis.

OSCITANT (FEVER), *Febris os'citans*, (F.) *Fèvre oscitante*, from *oscitare*, 'to yawn.' A fever, in which the patient is continually yawning.

OSCITATIO, Yawning.

OSULATORIUS, Orbicularis oris.

OSCULUM UTERI, Os uteri.

OSCUS, Scrotum.

OSEILLE, *Rumex acetosus*—o. *Boucher*, *Rumex scutatus*—o. *Petite*, *Rumex scutatus*—o. *Ronde*, *Rumex scutatus*—o. *Rouge*, *Rumex sanguineus*.

OSIER, *Salix alba*—o. red, *Cornus sericea*.

OS'MAZOME, *Os'mozome*, *Osmazo'ma*, *Osmozo'ma*; *Os'mazone*, *Zomod'mum*, (F.) *Matière extractive du Bouillon, Extrait de viande; Extractive or Saponaceous extract of meat*; from *osmē*, 'smell,' and *ζωπος*, 'soup.' M. Thénard gave this name to an extractive matter, contained in muscular flesh and in the blood of animals, which he considers of a peculiar nature. It has an agreeable smell and taste, and is found in *Bouillons* of meat, in the proportion of one part to seven of gelatin. Vauquelin discovered in it some fungi. It is the substance which gives the flavour of meat to soups, and hence its name. It is a mixture of different crystallizable and uncrystallizable principles with empyreumatic products.

Osmazome has been prescribed in the dose of ʒss to ʒj, in broth or in powder, with aromatics, as a stimulant to the digestive actions and a restorer of appetite in convalescence. An *Osmazome chocolate*, *Ohocolata cum osmazo'ma*, has been introduced as an analeptic article of diet. It consists of chocolate 1bj to osmazome ʒj.

OSME, Odour.

OSMESIS, Olfaction.

OSMIDRO'SIS, from *osmē*, 'odour,' and *ἰδρωσ*, 'sweat.' Perspiration of an unusual odour.

OSMITOPS'IS ASTERISCOY'DES, - *Bellia*.

A plant, of the Cape of Good Hope, which, from its smell and taste, seems to contain camphor; hence its antispasmodic virtues. In infusion, it is often beneficially employed in cough, hoarseness, and diseases of the chest generally, and is said to be serviceable in flatulent colic. Infused in spirit, *spir'itue bell'idie*, it has been used successfully as an external remedy in paralysis.

OSOMETRICUS SENSUS, Olfaction.

OSMON'OSI, *Osmonu'si*, *Morbi olfac'tūs*, from *osmē*, 'odour,' and *voos*, 'a disease.' Diseases of olfaction.

OSMONOSOLOG'IA, from *osmē*, 'odour,' *voos*, 'disease,' and *logos*, 'description.' The doctrine of, or a treatise on, the diseases of the sense of smell.

OSMORRHIZA LONGIS'TYLIS, (*osmē*, 'a scent,' and *ρίζα*, 'a root,') *Sweet Cic'ely*. An indigenous plant, *Order*, Umbellifera, which flowers in May and June. It is an agreeable aromatic, having much of the flavour of anise.

OSMOSE, *Osmo'sis*; from *ωσμος*, 'impulse.' The action by which fluids are impelled through

moist membrane and other porous septa, in endosmotic and exosmotic experiments.

OSMOT'IC, *Osmot'icus*, same etymon. Relating or belonging to osmose; as "osmotic force," the force which induces osmose.

OSMOZOME, Osmazome.

OSMUND, ROYAL, *Osmunda regalis*.

OSMUN'DA CINNAMOM'EAE, *Cinnamom Fern*. Indigenous; *Order*, Filices; is regarded to be demulcent, subastringent, and tonic. Boiled in milk, it yields a fine mucilage, which is useful in diarrhœa.

OSMUN'DA REGA'LIŒ, *O. spectabilis*, *Filix Flor'ida*, *Flow'ering Fern*, *Osmund-royal*. This plant was once thought to possess astringent and emmenagogue virtues.

OSMUNDA SPECTABILIS, *O. regalis*.

OSORON, Opium.

OSPHRANTERICUS, Olfactory.

OSPHRANTICOS, Olfactory.

OSPHRASTA, Olfaction.

OSPHEREIOL'OGY, *Osphresiology'ia*, from *σφρησις*, 'olfaction,' and *λογος*, 'a discourse.' A treatise on olfaction and odours.

OSPHERESIS, Olfaction.

OSPHERETICUS, Olfactory.

OSPHRIS'TICE, from *σφρησις*, 'olfaction.' The doctrine of the phenomena of odours.

OSPHYALGEMA, Coxalgia.

OSPHYALGIA, Coxalgia.

OSPHYARTHRTIS, Osphyitis.

OSPHYARTHROCACE, Coxarum morbus.

OSPHYI'TIS, *Osphy'tis*, *Osphyarthrit'is*, *Inflamma'tio coxæ*, from *σφύς*, 'the hip,' and *itis*, denoting inflammation. Inflammation of the parts about the hip.

OSPHYRRHEUMA, Lumbago.

OSPHYS, Haunch, Lumbi.

OSPHYTIS, Osphyitis.

OSSA BREGMATIS, Parietal bones—o. *Convoluta inferiora*, Turbinate bones, inferior — o. *Digitorum mandis*, Phalanges of the fingers — o. *Digitorum pedis*, Phalanges of the toes—o. *Epactalia*, Wormiana ossa — o. *Maxillæ superioris quarta seu quinta seu secunda*, Nasal bones — o. *Nasalia*, Nasal bones—o. *Nasi*, Nasal bones — o. *Nervalia*, Parietal bones—o. *Pisiformia lingualia*, see *Hyoides*, os—o. *Raphogeminantia*, Wormiana ossa — o. *Sesamoidea*, Sesamoid bones — o. *Sincipitis*, Parietal bones—o. *Spongiosa*, Turbinate bones—o. *Suturarum*, Wormiana ossa—o. *Tertia maxillæ superioris*, Turbinate bones — o. *Tetragona*, Parietal bones—o. *Triangularia*, Sphenoidal cornua — o. *Triangularia Blasii*, Wormiana ossa — o. *Triquetra*, Wormiana ossa—o. *Triticea*, see *Hyoides*, os—o. *Turbinata*, Turbinate bones—o. *Verticalia*, Parietal bones.

OSSEINE, Osteine.

OSSELETS DE BERTIN, Sphenoidalia cornua — o. *de l'Oreille*, *Ossicula auditūs* — o. *de l'Ovie*, *Ossicula auditūs*—o. *du Tympan*, *Ossicula auditūs*.

OSSEOUS, *Os'seus*, *Osto'des*, *Ostoi'des*, *Ostoid*, (F.) *Osseux*; from *os*, 'a bone.' Bony. Resembling bone.

OSSICLES, Ossicula, see Ossiculum.

OSSICULA, see Ossiculum.

OSSICULA AUDIT'US, *O. Auris*, (F.) *Ossicleta de l'Oreille ou de l'ovie ou du tympan*. The small bones or ossicles situate in the cavity of the tympanum, and forming an uninterrupted chain from the membrane of the tympanum to that of the fenestra ovalis. They are four in number; and their series, from without to within, is as follows: *Malleus*, *Incus*, *Os orbiculare*, *Stapes*.

OSSICULA BERTINI, Sphenoidalia cornua — o. *Innomina*, *Innominata minora ossa* — o. *Nasi*, Nasal bones—o. *Sesamoidea*, Sesamoid bones—o. *Triticea*, see *Hyoides*, os.

OSSIC'ULUM, (plural *Ossic'ula*.) *Osteo'rion*, *Os'sicle*. Diminutive of *os*, *ossis*, 'a bone.' A small bone. See *Os*.

OSSICULUM COCHLEARE, Orbicular bone—*o*. Incudi Comparatum, Incus—*o*. Malleolo assimilatum, Malleus—*o*. Molari denti comparatum, Incus—*o*. Orbicular, Orbicular bone—*o*. Quartum, Orbicular bone—*o*. Squamosum, Orbicular bone.

OSSIFICATION, *Ossifica'tio*, *Ossifac'tio*, from *os*, 'a bone,' and *facere*, 'to make.' *Osteogen'ia*, *Osteogen'esis*, *Osteo'sis*, *Osto'sis*. Formation of bone. Development or increase of the osseous system. Ossification takes place in the same manner as the nutrition of other organs. The bones are, at first, mucous, and afterwards cartilaginous; the cartilage, at length, receives the phosphate of lime, and is, at the same time, replaced by a gelatinous parenchyma, when the bone has acquired the whole of its development.

OSSIFICATION, POINTS OF, *Puncta seu Nu'clei ossificatio'nis*, *Nu'clei ossi*, are the points where the ossification of a bone commences, whence it extends to other parts. Almost all the bones present a greater or less number of points of ossification. Besides the natural ossification, which we observe in the fetus and in the first periods of life, there are also *accidental ossifications*, such as those frequently remarked after the inflammation of serous membranes, in the parietes of arteries, and to which the terms *Osteo'ia*, *Incrustation* and *Petrifaction* have, sometimes, been applied.

OSSIFRAGA, *Osteocolla*.

OSSIS SACRI ACUMEN, *Coccyx*.

OSSISANA, *Osteocolla*.

OSSIV'OROUS, from *os*, *ossis*, 'a bone,' and *voro*, 'I devour.' A species of tumour, mentioned by Ruysch, which destroys bone.

OS'TAGRA, *Os'teagra*, from *osteon*, 'a bone,' and *agga*, 'seizure.' A forceps to cut or remove portions of bone.

OSTALGIA, *Osteocopus*.

OSTARIUM, *Ossiculuh*.

OSTARTHTRITIS, *Arthrophlogosis*.

OSTARTHROCACE, *Spina ventosa*.

OSTEAGRA, *Ostagra*.

OSTEALGIA, *Osteocopus*.

OSTÉIDE, *Osteoid*.

OSTEINE, *Osseine*, from *osteon*, 'a bone.' The animal matter of bone.

OSTETES, *Osteocolla*.

OSTEITIS, *Osti'tis*, from *osteon*, 'a bone,' and *itia*. *Inflamma'tio Ossis*. Inflammation of bone; characterized by violent shooting and lancinating pains in a bone.

OSTEMPYE'SIS, *Osteömpye'sis*, *Ostempyo'sis*, from *osteon*, 'a bone,' and *empyois*, (*ev*, and *puon*), 'effusion of pus.' An abscess in the interior of a bone.

OSTEMPYOSIS, *Ostempyosis*.

OSTEOANABROSIS, *Osteonabrosis*.

OSTEOCE'LE, from *osteon*, 'a bone,' and *κηλη*, 'a tumour.' A hernia in which the sac is cartilaginous and bony:—a rare case. Also, osseous induration of one or both testicles.

OSTEOCLASIS, *Fracture*.

OSTEOCOL'LA, from *osteon*, 'a bone,' and *κόλλα*, 'glue.' *Glue-bone*, *Stone* or *Bone binder*; *Ossifraga*, *Holos'teus*, *Ossisa'na*, *Ostri'tes*, *Ostei'tes*, *Amos'teus*, *Osteol'ithos*, *Stelochi'tes*. A name given to petrified carbonate of lime, found in some parts of Germany and other portions of the globe: so called, because it has been supposed to possess the power of favouring the formation of callus in fractures. It is the *Chaux carbonatée concrétionnée incrustante* of Hally. Not used.

OSTEOC'OPUS, *Osteal'gia*, *Ostal'gia*, *Osteo-*

dyn'ia, from *osteon*, 'a bone,' and *κοπος*, 'fatigue.' *Boneach*, *Bone-ague*. Pain in the bones,—as in syphilis.

OSTEOCYST'OID, *Osteocystoi'des*, from *osteon*, 'a bone,' *κύστις*, 'a sack or cyst,' and *ειδος*, 'appearance.' A morbid growth in bone, consisting of the formation of cysts filled with puriform or other fluid.

OSTEODENTINE, see *Tooth*.

OSTEODYNIA, *Osteocopus*.

OSTEOGANGRÆNA, *Necrosis*.

OSTEOGENIA, *Ossification*.

OSTEOG'RAPHY, *Osteogra'phia*, from *osteon*, 'a bone,' and *γραφειν*, 'to describe.' Description of the bones. The part of anatomy which describes the bones.

OSTEOHELCOISIS, *Caries*.

OSTE'OID, *Osteoi'des*, (F.) *Ostéide*, from *osteon*, 'bone,' and *ειδος*, 'appearance.' An accidental bony production, or more frequently a morbid production, which is a calcareous incrustation of the normal tissue or of fibrous tumours.

OSTEOLITHOS, *Osteocolla*.

OSTEOLOG'Y, *Osteolog'ia*, from *osteon*, 'a bone,' and *λογος*, 'a discourse.' The part of anatomy which treats of bones.

OSTEOLYO'SIS, from *osteon*, 'a bone,' and *λυω*, 'I dissolve.' Necrosis or death of the bone, as in hospital gangrene, in which the dead parts are of soft consistence.

OSTEOMA, *Exostosis*.

OSTEOMALACIA, *Mollities ossium*—*o*. Infantum, *Rachitis*.

OSTEOMALACOSIS, *Mollities ossium*.

OSTEOMYELITIS, *Medullitis*.

OSTEON, *Bone*—*o*. Hieron, *Sacrum*.

OSTEONABRO'SIS, *Osteonabro'sis*, from *osteon*, 'a bone,' a priv., and *βρωσις*, 'nutriment.' Atrophy of the osseous texture marked by diminution of volume.

OSTEON'CUS, from *osteon*, 'a bone,' and *ογκος*,

'a tumour.' A bony tumour.

OSTEONEROSIS, *Necrosis*.

OSTEON'OSI, *Osteonu'si*, from *osteon*, 'a bone,' and *νοσος*, 'a disease.' *Mor'bi os'sium*. Diseases of the bones.

OSTEOPÆDION, *Lithopædion*.

OSTEOPALIN'CLASIS, from *osteon*, 'a bone,' *παλιν*, 'again,' and *κλασις*, 'fracture.' The breaking again of a bone which has united unfavourably.

OSTEOPHY'MA, *Osteonecus*. Sometimes used in the sense of *Exostosis*. By Lobstein applied to bony productions originating in the deep-seated layers of the periosteum, in the vicinity of a carious bone, and which seem to be intended to obviate the weakness resulting from the destruction of the bone.

OSTEOPHYTE, *Exostosis*.

OSTEOPHY'TON, *Os'teophyte*, from *osteon*, 'a bone,' and *φυω*, 'I grow.' Under this name, Rokitsansky describes a deposition of bony matter, found by him on the inner surface of the parietes of the skull of pregnant women.

OSTEOPLASTY, (F.) *Osteoplastic*, from *osteon*, 'a bone,' and *πλασσειν*, 'to form.' An operation by which the total or partial loss of a bone is remedied.

OSTEOPLEU'RA, *Pleurosto'sis*, from *osteon*, 'a bone,' and *πλευρον*, 'a rib.' Ossification of the cartilages of the ribs.

OSTEOPORO'SIS, *Osteosclero'sis*, from *osteon*, 'a bone,' and *πωρωσις*, 'induration.' Induration of a bone, from the deposition of too much bony matter. Also, a bony degeneration, which consists in too great porosity with hardening. [?]

OSTEOPSATHYROSIS, *Fragilitas ossium*.

OSTEO-SARCO'MA, *Osteo-sarco'sis*, *Sarcosto'sis*, *Hæmatocysto'sis*, from *osteon*, 'bone,' and *σαρξ*,

σάρκας, 'flesh.' Disease of the bony tissue, which consists in softening of its laminae, and their transformation into a fleshy substance, analogous to that of cancer; accompanied with general symptoms of cancerous affection. The word has often, also, been used synonymously with *spina ventosa*.

OSTEOSARCOSIS, Osteosarcoma.

OSTEOSCLEROSIS, Osteoporosis.

OSTEOSIS, Ossification.

OSTEO-STEATOMA, *Exostosis steatomatodes*, from *οστέον*, 'a bone,' and *στέαρ*, *στεαρός*, 'suet or fat.' A name given to bony tumours or degenerations, which sometimes belong to osteosarcoma; at others, to *spina ventosa*, *exostosis*, &c.

OSTEOTOME; same etymon as Osteotomy. A bone saw, which has the form of a chain.—Nysten.

OSTEOTOMIST, *Osteotomis'ta*. Same etymon as the next. An instrument, invented by Dr. D. Davis, for cutting the bones of the fetal cranium, where it becomes necessary to greatly reduce the size of the head in parturition.

OSTEOTOMY, *Osteotom'ia*, from *οστέον*, 'a bone,' and *τομήν*, 'to cut,' (*τομή*, 'an incision.')

The part of practical anatomy whose object is the dissection of bones.

OSTEOTOPHUS, Tophus.

OSTEOTYLUS, Callus.

OSTEULCUM, Bone-nippers.

OSTEUM, Bone.

OSTHEXIA, see Ossification.

OSTIARIUS, Pylorus.

OSTIOLUM, diminutive of *οστέον*, 'a mouth or opening.' A small door or gate.

OSTIOLA CORDIS, *Valvula cordis, Pellicula cordis*. The valves of the heart.

OSTITIS, Osteitis.

OSTIUM, *οστέον*, *Orific'ium*, a door or gate, — of the heart, for example. The opening — *Ostium atrioventriculārē seu ventriculī venosum* — between the auricle and ventricle of each side. *Ostium ventriculī arteriosum*, is the opening at the origin of the pulmonary artery and aorta.

OSTIUM ABDOMINALE, see Tube, Fallopian — o. Arteriosum, see Ostium — o. Atrioventriculare, see Ostium — o. Duodenale ventriculī, Pylorus — o. Internum, see Ostium uterī — o. Œsophageum, Cardia.

OSTIUM UTERI. The vulva; the vagina. The term *Ostium internum* has been applied to the inferior opening of the uterus, which establishes a free communication between the cavities of the body and neck, and which is often obliterated in old women.—Cruveilhier. The neck of the uterus; the mouth of the uterus. See Uterus.

OSTIUM UTERINUM, see Tube, Fallopian — o. Venosum, see Ostium — o. Ventriculī Duodenale seu Pyloricum seu Dextrum seu Inferius, Pylorus.

OSTODES, Osseous.

OSTOIDEA SUBSTANTIA, see Tooth.

OSTOID, Osseous.

OSTOMA, Exostosis.

OSTOSIS, Ossification.

OSTREA, *Os'treum*, *οστρέον*, *οστρεϊον*, *Os'trea edulis*, (F.) *Huître*. Class, Acephala. The oyster. The oyster is a wholesome article of diet, often advised as analeptic and easy of digestion.

The shells, *Testa Os'trea*, *Testa* (Ph. U. S.), (F.) *Écailles d'huîtres*, have been received into the Pharmacopœias, and used medicinally. Their virtues are similar to those of the carbonate of lime. See Creta.

OSTREUM, Ostrea.

OSTRITES, Osteocolla.

OSTRUTHIUM, Imperatoria.

OSWEGO TEA, *Monarda coccinea*.

OSYRIS, *Antirrhinum linaria*.

OTACOUS'TIC, *Otacus'ticus*, from *οὖς*, *ωτός*,

'the ear,' and *ακούω*, 'I listen.' Any instrument is so called which improves the sense of hearing, — as the different species of trumpets.

OTAGRA, Otagria.

OTAL'GIA, *Ot'agra*, *Otod'ynē*, *Otal'gy*, (F.) *Otalgie*, from *οὖς*, *ωτός*, 'the ear,' and *αλγος*, 'pain.' Pain in the ear; *Dolor aurium*, *Spas'mus aurium*, *Ear-ache*. See Otitis.

OTAL'GIC, *Otal'gicus*, (F.) *Otalgique*. An epithet for remedies used in otalgia.

OTEC'HUS, Tinnitus aurium.

OTEMPLAS'TRUM, *Otoplas'ta*, *Otoplas'trum*, from *οὖς*, *otos*, and *ἐμπλάστρον*, 'a plaster. A plaster put behind the ears.

OTEN'CHYTES, *Oten'chyta*, from *οὖς*, *otos*, *εν*, 'into,' and *χύνω*, 'I pour.' *Sipho auriculāris*. A species of syringe, used for injecting the ear.

OTHELCO'SIS, from *οὖς*, *otos*, and *ἐλκωσις*, 'ulceration.' Ulceration or suppuration of the ear.

OTHONE, Linteam.

OTHONION, Linteam.

OTHYGROTES, Otitrheea.

OTIATER, Aurist.

OTIATRIA, *Otiat'rice*, from *οὖς*, *otos*, and *ιατρεία*, 'medicine.' The business of the aurist. The management of diseases of the ear; *Aural Medicine and Surgery*, *Ear Surgery*.

OTIATRICE, Otiatria.

OTIATRUS, Aurist.

O'TIC, *O'ticus*, (F.) *Otique*, from *οὖς*, *otos*. Belonging or relating to the ear. A medicine employed in diseases of the ear.

OTIC GANGLION, *Otogan'gium*, *Gang'lion auriculārē seu o'ticum*, *Auricular ganglion*, *Ganglion of Arnold*, (F.) *Ganglion maxillo-tympanicum*. This is a small ganglion, although more than double the size of the ophthalmic, ash-coloured and pulpy like the ganglions of the sympathetic. It is situated in advance of the ganglion of Gasser, on the lower surface of the inferior maxillary nerve, at the inner margin of the foramen ovale of the sphenoid bone. See Petrous ganglion.

OTIRRHŒ'A, *Otorrhœ'a*, *Othy'gotes*, *Blennorrhœ'a*, (F.) *Catarrhe de l'Oreille*, from *οὖς*, *otos*, and *ρῶς*, 'I flow.' Discharge of a purulent or puriform liquid, from the meatus auditorius externus, *Pyorrhœ'a aurium*, *Otopyorrhœ'a*, *Pyatorrhœ'a*, *Otorrhœ'a purulenta*. It is a symptom of chronic otitis. Some authors, have, indeed, used the term synonymously with chronic otitis. When it extends to the brain, it is termed *cer'eb'al otorrhœ'a*; and when from the middle or internal ear, it is termed *internal otorrhœa*, *Otorrhœ'a inter'na*, and *Entotorrhœ'a*.

OTIRRHŒA CEREBRAL, see Otitrheea.

OTTES, see Digitus.

OTTITIS, *Otophleg'monē*, *Emp'ree'ma otitis*, *Inflammat'io auris*, *Inflammation of the ear*, (F.) *Otite*, *Inflammation de l'oreille*, from *οὖς*, *otos*, and *itis*, denoting inflammation. Inflammation of the mucous membrane of the ear, characterized by excruciating pain; intolerable humming in the ear, with a discharge of mucus generally from the meatus externus or from the Eustachian tube. When the inflammation is restricted to the lining membrane of the meatus, it is termed *otitis catarrhalis*. Otitis is divided into *external* and *internal*, according as it affects the meatus auditorius, or the cavity of the tympanic or internal ear. It may be *acute* or *chronic*. *Acute otitis* commonly terminates in a few days in a favourable manner, with the aid of antiphlogistics. *Chronic otitis*, *otirrhe'a*, is almost always accompanied by a purulent discharge from the meatus auditorius. It is often incurable, and may ultimately give occasion to disease of the internal ear, and even of the brain.

OTITIS CATARRHALIS, see Otitis.

O'TIUM, *Ota'rium*, from *ous*, *otos*. A small ear. The auricle or pavilion of the ear.

OTOCEPH'ALUS, *Monot'ia*, from *ous*, *otos*, and *κεφαλη*, 'head.' A monster, whose ears are in contact, or united into one.

OTOCONIES, see *Otolithi*.

OTODYNE, *Otagia*.

OTOGLYPHIS, *Earpick*.

OTOGLYPHUM, *Earpick*.

OTOG'RAPHY, *Otogra'phia*, from *ous*, *otos*, and *γραφη*, 'a description.' The part of anatomy which describes the ear.

OT'OLITHS, *Otolithi*, *O'tolites*, (F.) *Otolithes*, from *ous*, *otos*, and *λίθος*, 'a stone.' White calcareous substances contained in the membranous vestibule, and seeming to be suspended in the fluid of the vestibule by means of a number of nervous filaments proceeding from the auditory nerve. Their universal presence in the ear of the mammalia would seem to show that they are inservient to audition. When of a looser consistence, they are called *Otocon'ia*, (F.) *Otoconies*, from *ous*, *ωτος*, 'the ear,' and *κόνια*, 'dust.'

OTOL'OGY, *Otolog'ia*, from *ous*, *otos*, and *λογος*, 'a discourse.' The part of anatomy which treats of the ear. An anatomical treatise on the ear.

OTOPATHEMA, *Otopathy*.

OTOP'ATHY, *Otopathi'a*, *Otopathe'ma*, from *ous*, *otos*, and *πάθος*, 'a disease.' A diseased condition of the ear.

OTOPHLEGMON, *Otitis*.

OTOPLASTA, *Otemplastrum*.

OTOPLAS'TICE, *O'toplasty*, from *ous*, *otos*, and *πλαστικός*, 'forming.' The operation for restoring a lost ear.

OTOPLASTRUM, *Otemplastrum*.

OTOPYORRHŒA, *Otiorrhœa*.

OTORRHAG'IA, from *ous*, *otos*, and *ρην*, 'rupture.' Bleeding from the ear or ears.

OTORRHEUMATIS'MUS, from *ous*, *otos*, and *ρευματισμός*, 'rheumatism;' *Otal'gia rheumat'ica*. Rheumatic ear-ache.

OTORRHŒA, *Otiorrhœa*—o. Cerebral, see *Otiorrhœa*—o. Interna, see *Otiorrhœa*—o. Purulenta, *Otiorrhœa*.

O'TOSCOPE, *Otoscop'ium*, from *ous*, *otos*, and *σκοπη*, 'examination.' An instrument for examining the condition of the ear. An elastic stethoscope, the ends of which are tipped with ivory—one to be inserted into the meatus of the patient, and the other applied to the ear of the examiner.

OTOT'OMY, *Ototom'ia*, from *ous*, *otos*, and *τεμνειν*, 'to cut.' The part of practical anatomy which teaches the mode of dissecting and preparing the ear.

OTTO OF ROSES, see *Rosa centifolia*.

OUACHITA SPRINGS, see *Arkansas*, mineral waters of.

OULE, *Audition*.

OULA, *Gingivæ*.

OULE, *Cicatrix*.

OULITIS, *Ulitis*.

ULORRHAGIA, *Ulorrhagia*.

OUNCE, *Un'cia*. A weight equal to the 16th part of a pound avoirdupois, the 12th of a pound troy. It is composed of 8 drachms. See *Weights and Measures*.

OURAQUE, *Urachus*.

OURARI, *Curare*.

UREMA, *Urine*.

URLES, *Parotis*, *Cynanche parotidæa*.

URON, *Urine*.

OUS, genitive *otos*, *ous*, *ωτος*, *Ear*.

OUTHAR, *Mamma*, *Uterus*.

OUTRE-MER, *Lapis lazuli*.

OVERTURE CADAVERIQUE, *Autopsia cadaverica*.

OVA GRAAFIANA, *Folliculi Graafiani*—o. *Nabothi*, *Nabothi glandulæ*.

OVAIRE, *Ovarium*.

OVAL, *Ova'lis*, *Oð'des*, *Oûi'des*, *Ova'tus*, (F.) *Ovale*, from *ovon*, *ovum*, 'an egg.' Egg-shaped; elliptic. That which is round and oblong like an egg.

OVALÈ FORA'MEN, *Foramen oû'des*, (F.) *Trou ovale*. The foramen obturatorium; also, the foramen in the inner paries of the middle ear, which opens into the vestibule; the foramen in the sphenoid bone, through which the third branch of the fifth pair issues from the cranium; and the aperture, in the foramen, between the auricles of the heart. See *Fotal Foramen*.

OVALIS FOSSA, *Fo'vea ova'lis*, *Val'vula seu Vestig'ium Foram'inis ovalis*, is a depression observed in the right auricle of the heart, in the septum, which separates it from the left auricle. It has thick and strong edges, at its upper and fore part, called *Colum'næ foram'inis ovalis*, *Isthmus seu An'nulus Vieussæni*, *An'nulus seu Limbus fossæ ovalis*: in the fœtus called *An'nulus foram'inis seu fossæ ova'lis*. The fossa ovalis, in the adult, replaces the inter-auricular aperture in the septum auricularum of the fœtus.

OVALBUMEN, *Albumen ovi*.

OVA'RIAN, *Ova'rial*, *Ovaria'nus*, (F.) *Ovarien*, *Ovarique*; from *ovarium*, *ωαριον*, 'the ovary.' Relating or belonging to the ovary—as *Ovarian pregnancy*.

OVARIAN ARTERY, *Spermatic artery*—o. *Nerves*, *Spermatic plexus of nerves*—o. *Pregnancy*, see *Pregnancy*, *ovarian*—o. *Veins*, *Spermatic veins*—o. *Vesicles*, *Folliculi Graafiani*.

OVARIOTOMY, *Oariotomy*.

O'VARISTS, *Ovists*. Same etymon as *Ovarian*. Those physiologists who think that the phenomena of generation, in the human species and in every species of animal, result from the development of the ova or ovula of the female, merely incited by the male. Not many of the physiologists of the present day can be classed among the ovarists.

OVARITIS, *Oaritis*.

OVARIUM, Diminutive of *ovon*, *ovum*, 'an egg.' The *O'vary*, (Sc.) *Eggbed*; *Testis muliebris seu femine'us*, *Oa'rion*, *Ooph'oron*, *Gynoa'rium*, *Vesica'rium*, (F.) *Ovaire*. The ovaries are the organs in which the ova are formed in oviparous animals. By analogy, the name has been given to the organs which the ancients called the *testicles* of the female. They are two ovoid bodies, almost as large as the testicles in men, placed on each side of the uterus, between the Fallopian tube and round ligament, and in the substance of the broad ligament. Their outer extremity gives attachment to one of the projections of the fimbriated extremity of the Fallopian tube; and the internal is fixed to the uterus by a small ligamentous cord, called *Ligamentum rotundum ova'rîi*, *Ligament of the O'vary*. The ovaries are composed of a very close, spongy texture—*stroma*,—and of small vesicles—*Folliculi Graafiani*,—filled with a clear fluid: these vesicles contain ovules, which detach themselves from the ovary before and after fecundation, and are carried into the cavity of the uterus by the Fallopian tube.

OVARIUM TUMIDUM, *Oarioncus*—o. *Nabothi*, *Nabothi glandulæ*.

OVATUS, *Oval*.

OVER-EATING, *COLIC FROM*, *Colica crapulosa*.

OVERO BUTUA, *Pareira brava*.

OVICAPSULE, see *Folliculi Graafiani*.

OVIDUCT, *Tube*, *Fallopian*.

OVIDUCTUS MULIEBRIS, *Tuba Fallopiana*.

OVIG'EROUS, *O'viger*, *Ovig'erus*; from

ovum, 'an egg,' and *gero*, 'I bear.' Containing or producing ova or eggs.

OVIPARITÉ, see *Oviparous*.

OVIP'AROUS, *Ovip'arus*, (F.) *Ovipare*; from *ovum*, 'an egg,' and *pario*, 'I bring forth.' An epithet applied to animals which are hatched from eggs outside of the body. The condition may be termed *Ovip'arousness*, (F.) *Oviparité*. See *Generation*.

OVIPAROUSNESS, see *Oviparous*.

OVISAC, see *Folliculi Graafiani* — o. Tunic of the, see *Folliculi Graafiani*.

OVISTS, *Ovarists*.

OVOLGY, *Oologia*.

OVO-VIVIP'AROUS, (F.) *Ovo-vivipare*, from *ovum*, 'an egg,' *vivus*, 'living,' and *pario*, 'I bring forth.' *Oviparous* animals, whose ova are hatched within the mother,—vipers, for example. See *Generation*.

OVULA GRAAFIANA, *Folliculi Graafiani* — o. *Nabothi*, *Nabothi glandule*.

O'VULAR, *Ovula'ris*; from *ovulum*, 'an ovule.' Relating or belonging to the ovule.

OVULATION; from *ovulum*, diminutive of *ovum*, 'an egg.' The formation of ova in the ovary, and the discharge of the same. Also, the development of the embryo from the egg.

SPONTANEOUS OVULATION, (F.) *Ovulation spontané*, is the spontaneous formation and discharge of ova which takes place in the female of the mammalia. See *Heat*.

OVULE, *Ovulum*.

OVULINE, *Decidua reflexa*.

O'VULUM, *O'vule*; diminutive of *ovum*, 'an egg.' A small egg. That which has the appearance of a small egg. See *Ovum*.

OVUM, *ovv*, *Oön*, an egg. The eggs of poultry are chiefly used as food. The different parts are, likewise, employed in medicine. The shell, *Auran'cum*, *Auran'tum*, (F.) *Coque* ou *Coquille d'œuf*, calcined, is esteemed an absorbent. The Oil of the egg is emollient, and is used, externally, to burns and chaps. The Yolk or Yell of the egg — *Vitel'lum* seu *Vitel'lus ovi*, *Lec'ithos* — renders oils, &c., miscible with water. Eggs, when light boiled, are easy of digestion; but, when very hard-boiled or fried, they are more rebellious.

Egg Brandy is made by taking of Brandy f $\frac{3}{4}$ iv; Cinnamon water, f $\frac{3}{4}$ iv; the yolks of two eggs; Sugar, 3ss; and Oil of cinnamon, gtt. ij. Mix the yolks of the eggs first with the water, the oil and the sugar, stirring constantly. Then add the brandy little and little, until a smooth fluid is formed. It is an agreeable mode of giving brandy in adynamic states.

In Pharmacy, white of egg, which consists chiefly of albumen, is used for clarifying syrups, &c. Yolk of egg, (F.) *Jaune d'œuf*, beaten up with warm water and sugar, and aromatized with orange-flower water, forms an emulsion, which is emollient, and known under the name *Lait de poule* or *chicken's milk*.

Anatomists give the name *Ova*, *O'vula*, *O'vules*, (F.) *Œufs*, to round vesicles, containing a humour similar to the yolk of egg, which are situate in the ovaries of the female, and, when fecundated, constitute the rudiments of the fœtus. During gestation, the embryo and its enveloping membranes retain the name *Ovum*, (F.) *Œuf*. The changes induced in the mammalia after impregnation greatly resemble those in the bird.

OVUM, Testicle — o. *Hystericum*, see *Clavus hystericus*.

OXALAS SUBPOTASSICUS, Potass, oxalate of.

OXALAT'IC, *Oxalat'icus*. Relating to oxalates: hence *Oxalat'ic* or *Oxalic acid Diath'esis*

is the habit of body which favours the formation and deposition of oxalates from the urine.

OXAL'IC ACID, *Acidum oxali'num* seu *sacchari'num* seu *hydro-carbon'icum* seu *acetosel'la*, seu *oxal'icum* seu *sac'chari*, *Acid* of sugar or of sorrel, *Car'bonus acid*, from *ox'alis*, 'sorrel.' An acid found in sorrel. It crystallizes in long, colourless, quadrangular prisms, and is generally found, in nature, united with lime or potassa. It has been proposed, in a diluted state, as a refrigerant, but it is unnecessary; and, besides, in quantity, it is a virulently acrid poison. It is largely employed for cleaning boot-tops; and, as it strongly resembles the sulphate of magnesia, has often been taken for it. Several fatal cases are, indeed, on record. For its antidotes, &c., see *Poison*.

OXALIC ACID DIATHESIS, see *Oxalatic*.

OXALIDE, *Oxalis acetosella*.

OXALIS ACETOSA, *Rumex acetosus*.

OX'ALIS ACETOSEL'LA, *O. America'na*, *Oxy alba*, called from *oxys*, 'acid,' owing to the sharpness of its juice. This systematic name of the Wood-sorrel, Cuckowbread, Sour Tref'oil, White sorrel, Mountain Sorrel, *Trifo'l'ium acetosum*, *Oxitrif'hyllum*, *Lujula*, *Allelu'ia*, *Acetosel'la*, *Oxyphyl'lon*, *Oxytrif'hyll'lon*, *Panis cu'culi*, (F.) *Surelle*, *Oxalide*, *Pain a Coucou*, *Alleluia*. Ord. *Oxalidaceæ*. Sez. *Syst. Decandria Decagynia*. This plant has a gratefully acid taste, and is used in salads. It has been esteemed refrigerant, antiscorbutic, and diuretic.

OXALIS AMERICANA, *O. acetosella*.

OXALIS STRICTA, *Yellow wood sorrel*; indige-nous, has similar virtues.

OXALIS TUBERO'SA, *Oca*. A tuberous plant, whose oval-shaped root is used as food by the Peruvians. It is watery, has a sweetish taste, and is much liked.

OXALIUM, Potass, oxalate of.

OXAL'ME, from *oxys*, 'acid,' and *als*, 'salt.' A mixture of vinegar and salt.

OXALU'RIA, Urine, oxalic.

OXELÆ'ON, from *oxys*, 'acid,' and *elaion*, 'oil.' A mixture of vinegar and oil, used as a condiment.

OX EYE, *Helenium autumnale* — o. Gall, Bile of the ox.

OXÉOLÉ, *Acttolé*.

OXICÉDRE, *Juniperus oxycedrus*.

OXID, *Oxyd*.

OXIDATION, *Oxydation*.

OXIDE, *Oxyd*.

OXIDUM, *Oxyd*, see *Oxydum*.

OXINES, *oxivns*, (*ovos*), from *oxys*, 'acid.' Wine already sour, but not yet vinegar.

OXITRIPHYLLUM, *Oxalis acetosella*.

OXODE, *Oxyd*.

OXODES, *Acidulous*.

OXOIDES, *Acidulous*.

OXOS, *Aceticum acidum*, *Acetum*.

OXTER, *Axilla*.

OXURE, *Oxyd*.

OXYA, *Fagus sylvatica*.

OXYACANTHA, *Mespilus oxyacantha*.

OXYACANTHA GALE'NI, *Berberis*, *B. vulga'ris*, *Spina ac'i'da*, *Creepi'nus*, (F.) *Épine vinette*, *Vinettier*. Family, *Berberidæ*. Sez. *Syst. Hexandria Monogynia*. The *Barberry*. The fruit of the berries, *Berberies*, is gratefully acid, and moderately astringent. By preparing an alcoholic extract of the root and adding water to it, a pulverulent brown substance is thrown down, which, when dissolved in alcohol and evaporated, yields the bitter principle, *Berberin*, which is an excellent stomachic in the dose of 2, 5, or 10 grains.

The American Barberry, *Berberis Canadens'*.

ois, Pipperidge Bush, Souberry, differs slightly from the European. It flourishes on mountains and hilly districts from Canada to Virginia.

OXYÆSTHESIA, Hyperæsthesia.

OXYAPHE, Hyperaphia.

OXYAPHIA, Hyperaphia.

OXYBAPHION, see Actebulum.

OXYBOLIA, from *oxus*, 'quick,' and *βολη*, 'ejection.' Too rapid emission of sperm.

OXYCEDRE, Juniperus oxycedrus.

OXYCEPHALIA, 'Sugar-loaf head,' from *oxus*, 'sharp,' and *κεφαλη*, 'head.' Monstrosity in which the head is peaked like a sugar-loaf.

OXYCOCCOS, Vaccinium oxycoccus — *O. Palustris*, Vaccinium oxycoccus.

OXYCRATE, *Oxyra'tum*, from *oxus*, 'acid,' and *κραω*, 'I mix;' *Posca*, *Phusca*. A mixture of vinegar and water; frequently used in medicine as a refrigerant and antiseptic.

OXYCROCEUM EMPLASTRUM, from *oxus*, 'acid,' and *κροκος*, 'saffron.' A plaster, composed of saffron, pitch, colophony, yellow wax, turpentine, galbanum, gum ammoniac, myrrh, olibanum, and mastic. It was employed as a strengthening plaster.

OXYCUM, Oxygen.

OXYD, *Oxid*, *Oxide*, *Oxyde*, *Ox'ydum*, *Ox'idum*, (F.) *Oxyde*, *Oxode*, *Oxure*, from *oxus*, 'acid.' A combination, not acid, of a simple body with oxygen.

OXYD, CALCULOUS, Oxyd, cystic — *O. Vesical*, Oxyd, cystic.

OXYD, CYSTIC, *Cal'culous* or *Ves'ical oxyd*, *Cystine*, *Nephrine*, *Scor'dosmine*. A species of urinary calculus, having the shape of confused, yellowish, semi-transparent, insipid, very hard crystals. When distilled, it furnishes subcarbonate of ammonia, like all nitrogenized matters. It dissolves in the nitric, sulphuric, phosphoric, and oxalic acids.

OXYDA'TION, *Oxyda'tio*, *Oxida'tion*, *Oxygenation*. The action of oxydizing a body; that is, of combining it with oxygen; a combination from which results an oxyd.

OXYDE, Oxyd.

OXYDE D'ANTIMOINE SULFURÉ, *Oxydum stibii sulphuratum* — *O. Gris ou noir de Mercure*, see Hydrargyri oxydum cinereum.

OXYDER/CIA, *Oxydor'cia*, *Vius a'cies*, from *oxus*, 'quick,' and *δερκα*, 'I see.' Sharpness of vision.

OXYDER/CICUS, *Oxyder'ces*. A medicine which sharpens the sight.

OXYDORCIA, Oxydercia.

OXYDUM, Oxid — *O. Antimonii cum phosphate calcis*, Antimonial powder.

OXYDUM FERRI NIGRUM, *Ferri ox'idum nigrum*, *Black oxyd of iron*. The scales, which fall from iron, when heated, consist of iron, combined with oxygen. They have been used like the chalybeates in general. The Edinburgh Pharmacopœia has a *Ferri ox'idum nigrum*, made by precipitating, by ammonia, a solution of the mixed sulphates of protoxide and sesquioxide of iron. An analogous preparation is in the Dublin Pharmacopœia, under the name *Ferri ox'ydum magneticum*, *Magnetic oxide of iron*, *Martial Ethiops*.

OXYDUM FERRICUM CRYSTALLIZATUM, *Hæmatites* — *O. Hydrargyri completum*, *Hydrargyri nitrico-oxydum* — *O. Hydrargyricum*, *Hydrargyri nitrico-oxydum* — *O. Hydrargyricum præparatum*, *Hydrargyri oxydum cinereum* — *O. Hydrargyrosium*, *Hydrargyri oxydum cinereum*.

OXYDUM STIBII SULPHURATUM, *Hepar antimoni*, *Cr'cus metallo'rum seu antimoni*, (F.) *Oxyde d'antimoine sulfuré*, *Foie d'Antimoine*. This was formerly exhibited in the cure of fevers, but it is now rarely employed.

OXYDUM URICUM, Uric oxide.

OXYECOIA, Hyperacusis.

OXYG'ALA, from *oxus*, 'sour,' and *γαλα*, 'milk.' Sour milk. The *Oxygala equi'num*, *Caracas' mos*, is reckoned amongst the delicacies of the Tartars.

OXYG'ARON, from *oxus*, 'acid,' and *γαρον*, 'garum.' A composition of garum and vinegar.

OXYGEN, *Oxygen'ium*, *Empy'real*, *Pure*,

Vital or *Dephlogis'ticated air*, *Oxyg'enes*, *Ox'yum*, *Princip'ium oxyg'enans* seu *oxygenet'icum* seu *acidificans* seu *ac'idum*, *Elemen'tum acidific'icum*, (F.) *Oxygène*, *Air du feu*; from *oxus*, 'acid,' and *γεννω*, 'I engender,' that is, *generator of acids*; and such it was believed to be, *exclusively*, at the period when the name was given to it. This is now known not to be the case. Oxygen is largely distributed in nature. It exists in the air, in water, in several acids, in all the oxyds, and in vegetable and animal substances, &c. It is obtained by decomposing the peroxyd of manganese or the chlorate of potassa by heat in close vessels. Although oxygen, in the state of admixture in which it is found in the atmosphere, is of vital importance, it cannot be respired in a pure state with impunity. Animals die in it long before the whole of the oxygen is consumed. The properties of oxygen seem to be stimulant. It increases the force and velocity of the pulse, and has, accordingly, been used in cases of chronic debility, chlorosis, asthma, scrofula, dropsy, paralysis, &c. It requires to be diluted with from 10 to 20 parts of atmospheric air; one to two quarts being given during the day.

OXYGENATION, Oxydation.

OXYGÈNE, Oxygen.

OXYGÈNESES. A flame given by Baumes to diseases which he attributed to disordered oxygenation.

OXYGEUSIA, Hypergeusia.

OXYG'LICUS, *Oxyg'lices*, from *oxus*, 'acid,' and *γλυκς*, 'sweet;' *Mulsa acida*, *Acetum mulsum dulcè*. Prepared by macerating and boiling honey-comb (from which the greater part of the honey has been taken) in water, and adding vinegar. An *Oxymel*.

OXY-HYDRO-CARBURETUM EX OLEO PYROXYLICCO PARATUM, *Creasote*.

OXYLAPATHUM, *Rumex acutus*.

OXYMEL, *Apom'eli*, *Mel acetat'um*, (F.) *Acetomel*, from *oxus*, 'acid,' and *μελι*, 'honey.' Honey and vinegar boiled to a syrupy consistence. (*Mel despumat. lbj, acid. acetic. ℥vij, aq. destillat. f℥viij.* Add the acid to the water and mix them with the honey previously heated.—Ph. L.) It is cooling;—externally detergent.

OXYMEL ERUGINIS, Linimentum *Æruginis*.

OXYMEL COL'CHIC' *O. of meadow saffron*, (F.) *Oxymel colchiqué*. (*Rad. colch. recent. in laminas tenues sectæ ℥j, aceti destillati lbj, mellis despumat. pond. lbj.* Macerate in a gentle heat for 48 hours. Press and boil the liquor with the honey to the thickness of a syrup; stirring with a wooden spoon. Ph. D.) It is expectorant and diuretic. Dose, f℥j, in gruel.

OXYMEL COLCHIQUE, *O. colchici* — *O. of Meadow saffron*, *O. colchici*.

OXYMEL SCILLÆ, *O. of squills*, *O. Scillicit'icum*, *Mel scillæ*. (*Mellis despumat. Oiss, aceti scillæ, Oij.* Boil in a glass vessel over a gentle fire to a proper consistence. Ph. U. S.) It is reputed to be expectorant and diuretic. Dose, f℥j to f℥ij.

OXYMEL OF SQUILL, *O. scillæ*.

OXYMYRRHINE, *Ruscus*.

OXYMYRSINE, *Ruscus*.

OXYNGULUM, *Adeps suillus*.

OXYNI'TRON, from *oxus*, 'acid,' and *νιτρον*, 'nitre.' Name of a plaster, composed chiefly of vinegar and nitre, and recommended by Aëtius.

OXYNOSEMA, Acute disease.

OXYNOSOS, Acute disease.

OXYNUSOS, Acute disease.

OXYOPIA, *Galero'pia*/ *Hyperop'sia*, *Suffu'sio excla'rana*, *Acies visus*, from *οξυς*, 'quick,' and *ωπ*, 'sight.' Excessive acuteness of the sense of sight. Instances have occurred where persons could see the stars in the day time. The cause is seated in great sensibility of the retina. The term has been used synonymously with *nyctalopia*.

OXYOSPHRE'SIA, *Oxyosphra'sia*, from *οξυς*, 'acute,' and *οσφρησις*, 'smell.' Excessive acuteness of the sense of smell.

OXYPHLEGMA'SIA, *οξυφλεγμασια*, (*οξυς*, 'acute,' and *φλεγμασια*), *Acu'ta et ve'hemens inflammatio*. A violent inflammation.

OXYPHENICON, *Tamarindus*.

OXYPHO'NIA, *Vox acu'ta*, from *οξυς*, 'sharp,' and *φωνη*, 'voice'; *Parapho'nia clangens* seu *ulcero'sa*, 'Clangor,' *Vox clango'sa*, *Leptopho'nia*. A shrill and squalling voice,—frequently, a symptom of morbid affection of the larynx.

OXYPHYLLON, *Oxalis acetosella*.

OXYPODIA, see *Kyllosis*.

OXYPROTEIN, see *Corium phlogisticum*.

OXYREG'MIA, *Ructus ac'idus*, from *οξυς*, 'sour,' and *ερευνα*, 'I belch.' Acid eructation.

OXYR'IA RENIFORM'IS, *Rumex dig'ynus*, *Boreal sourdock*, *Mountain Sorrel*, *Welsh sorrel*. A plant, *Ord. Polyganaceæ*, *Sex. Syst. Diandria Digynia*, which grows in the northern parts of America and Europe; blossoming in the spring. It possesses the same properties as *Oxalis acetosella*.

OXYRRHOD'INON, *Acetum rosa'tum*, from *οξυς*, 'acid,' and *ροδον*, 'a rose.' A composition of vinegar and roses. (*Ol. rosat.* ζ ij, *acet. rosat.* ζ ij.) Used as a liniment in herpes and erysipelas.

OXYS, Acid, Acute—o. Alba, *Oxalis acetosella*.

OXYSACCHARUM, from *οξυς*, 'acid,' and *σακχαρον*, 'sugar.' A mixture of sugar and vinegar. It was called *Oxysaccharum vomiti'vum*, when it held in solution the glass of antimony; and *Oxysaccharum scillit'icum*, when it contained squills.

OXYSAL DIAPHORET'ICUM AN'GELI SALÆ. A preparation of Angelo Sala. Acetate of potass exposed to the atmosphere; and consequently fallen into a state of deliquium.—Orfila.

OXYTARTARUS, *Potassæ acetat.*

OXYTES, Acidities.

OXYTOC'IA. Same etymon as the next. Rapidity of parturition.

OXYTOC'IC, *Oxyt'ocus*, from *οξυς*, 'quick,' and *τοκω*, 'I bring forth.' A medicine which promotes delivery.

OXYTRIPHYLLON, *Oxalis acetosella*.

OXYURE VERMICULAIRE, *Ascaris vermicularis*.

OXYURIS VERMICULARIS, *Ascaris vermicularis*.

OYSTER GREEN, *Ulva latissima*—o. Root, *Tragopogon*—o. Shells, *Ostreæ testæ*—o. Prepared, *Testa præparata*.

OZÆ'NA, *Coryza enton'ica* seu *purulen'ta* seu *ozæno'sa* seu *ulcero'sa* seu *virulen'ta* seu *malig'na*, *Pyorrhœa nasa'lis*, *Entozæ'na*, *Rhini'tis ulcero'sa*, *Ulcus na'rrium fetens*, from *οζω*, 'I smell of something.' (F.) *Ozène*, *Punaisie*. An affection of the pituitary membrane, which gives occasion to a disagreeable odour similar to that of a crushed bed-bug; hence the name *Punais*, by which the French designate one labouring under *ozæna*. It is, sometimes, owing to caries of the bones; but is, perhaps, most frequently dependent upon syphilitic ulceration of the pituitary membrane, with or without caries of the bones of the nose.

OZENA BENIGNA, *Coryza*, chronic.

OZE, Breath, offensive.

OZEMUN, Albumen.

OZÈNE, *Ozæna*.

OZONE, same etymon as *Ozæna*. The powerfully odorous matter produced when a current of ordinary electricity passes from pointed bodies into the air.—Schönbein. It is presumed to be a peculiar modification of oxygen; and its varying quantity in the atmosphere is supposed to affect the health of man.

Air containing ozone is said to be *ozonized*.

P.

P. A contraction of *Pugillus*.

P. Æ. A contraction of *Partes æquales*.

P. P. A contraction of *Pulvis patrum*.

See Abbréviation.

PAB'ULUM, *Trophē*, *Phorbē*, *Alimen'tum*, *Nutimen'tum*. Food, aliment. The animal heat and animal spirits—unmeaning terms—were formerly considered the *Pabulum vitæ*.

PACAY, *Prosopis dulcis*.

PACCHIO'NI, GLANDS OF, *Pacchio'nian Bodies*, *Glandulæ Duræ Matris* seu *duræ Menin'gis*, *Tuber'cula parva* seu *Cor'pora glandifor'mia duræ matris*, (F.) *Glandes de Pacchioni*, *Graulationes cérébrales*. Small, whitish, or yellowish bodies, sometimes separate, at others united like a bunch of grapes, which are observed in several parts of the dura and pia mater. They receive vessels, but apparently no nerves. Their texture and uses are unknown. A considerable number is observed in the longitudinal sinus, covered by the internal membrane.

PACHÆ'MIA, *Pachyæ'mia*, *Pachæ'ma*, *Pachyæ'ma*, from *παχυς*, 'thick,' and *αἷμα*, 'blood.' Too great thickness of the blood.

PACHEABLEPHARO'SIS, *Pacheableph'ara*, *Pachybleph'aron*, *Pachyblepharo'sis*, *Pach'ytes*, from *παχυς*, 'thick,' and *βλεφαρον*, 'an eyelid.' A name given, by some authors, to a thickening of the eyelids, produced by tubercles or excrescences formed on their margins. It is the same affection as that which Sauvages designates *Caligo* do *Pacheablepharo'sis*.

PACHETOS, see *Laqueus*.

PACHULOSIS, *Pachylosis*.

PACHYÆMIA, *Pachæmia*.

PACHYBLEPHARON, *Pacheablepharosis*.

PACHYCHOL'IA, from *παχυς*, 'thick,' and *χολη*, 'bile.' Morbid thickness of the bile.

PACHYDERMATOCE'LE, from *παχυς*, 'thick,' *δερμα*, *dermatos*, 'skin,' and *κηλη*, 'a tumour.' A disease of the dermoid tissue, apparently congenital, which increases from a small mole or

spot, at times to the size and appearance of a relaxed and very emaciated mamma.—V. Mott.

PACHYLOSIS, *Pachulosis*, from *παχυς*, 'thick.' An inordinate production of the epidermis dependent upon hypertrophy of the papillæ of the skin.—E. Wilson.

PACHYNSIS, Polysarcia adiposa.

PACHYNTICA, Incrassantia.

PACHYSMUS, Polysarcia adiposa.

PACHYTES, Pachæblepharosis.

PACINIAN CORPUSCLES, see Corpuscles.

PACKING, OF THE HYDROPATHISTS; *Emballotage*.

PACKWAX, see Nucha.

PAD, (S.) *Pajado*, from *paja*, 'straw' [?]; *Plumaceolus*, *Pulvillus*, *Pulvinulus*, *Parvum pulvinar*;—a cushion, a little cushion; a junk; (F.) *Coussinet*. A small bag, filled with feathers, wool, hair, bran, chaff, &c. It is used to compress or support parts. In the treatment of certain fractures, soft pads or cushions, called by the French *Remplissages*, are placed between the splints and the fractured limb, in order that the prominent parts may not be chafed by the splints.

PADDIE, see Oryza.

PADDY, see Oryza.

PADNOON, Bit noben.

PAD'UA, MINERAL WATERS OF. Near Padua, in Italy, there are several thermal sulphureous springs, the hottest of which is 188° Fahr. The chief establishments are Abano, Battaglia, and Montegrotto.

PADUS, *Prunus padus* — p. *Avium*, *Prunus padus* — p. *Lauro-cerasus*, *Prunus lauro-cerasus*.

PEDAN'CHONE, from *παῖς*, *παῖδος*, 'a child,' and *αἴχω*, 'I strangle;' *Angina Sicca*. Name of a fatal species of cynanche, mentioned by M. A. Severinus; probably the *Cynanche Trachealis*, or *Cynanche Maligna*.

PEDARTHROCEAE, *Spina ventosa*.

PEDATROPH'IA, *Atrophia infantilis*, *Marasmus lactantium*, from *παῖς*, *παῖδος*, 'a child,' a, privative, and *τροφή*, 'nourishment.' Atrophy of children. *Tabes mesenterica*.

PEDATROPHIA GLANDULOSA, *Scrofula*.

PÆDERAS'TIA, *Pædogogium*, *Pædicatorio*, *Ginædia*, from *παῖς*, *παῖδος*, 'a boy,' and *ἔρως*, 'love.' The love of boys. A disgraceful passion, common in ancient times, but now unknown. These terms are likewise applied to the crime of sodomy.

PÆDEROTA VIRGINICA, *Leptandra virginica*.

PÆDENCÉPHALIE, *Hydrocephalus internus*.

PÆD'IA, *παῖδα*, *Educatio infantium*. The bringing up or education — physical as well as moral — of children.

PÆDIATRI'A, *Pædiatrica*, *Pædiatrics*, from *παῖς*, *παῖδος*, 'a child,' and *ἰατρειά*, 'medical management.' Treatment of the diseases of children.

PÆDICATIO, *Pæderastia*.

PÆDICTERUS, *Icterus infantum*.

PÆDOBAROMACROMETER, *Baromacrometer*.

PÆDOGOGIUM, *Pæderastia*.

PÆDOMETER, *Baromacrometer*.

PÆDON'OSUS, from *παῖς*, *παῖδος*, 'a child,' and *νόσος*, 'disease.' A disease of childhood.

PÆDOPHLEBOTOM'IA, from *παῖς*, *παῖδος*, 'a child,' *φλέψ*, 'a vein,' and *τομή*, 'incision.' *Venæsectio puerorum*. Bleeding performed on children.

PÆDOT'RIBES, *Minister Gymnasæta*, from *παῖς*, *παῖδος*, 'a child,' and *τρίβω*, 'I train.' In ancient gymnastics, the officer acquainted with all the prescribed exercises, and who carried them into effect, according to the direction of the superior.

PÆDOT'ROPHY, *Pædotroph'ia*, (F.) *Pædotrophie*, from *παῖς*, *παῖδος*, 'a child,' and *τροφή*, 'nourishment.' The part of hygiene, which concerns the nourishment of children.

PÆON, Physician.

PÆON'IA, *P. officinalis seu macula seu nemoralis seu femina seu lobata*, *Glycis'idæ*, *Hæmago'gum*, *Pe'ony*, or *Pi'ony*, male and female. (F.) *Pivoine*, *Pione*, *Herbe Sainte-Rose*. From *Παῖων*, 'the god of physic.' The root, flowers, and seeds were long esteemed powerful medicines. They are feebly astringent, and have been considered to be narcotic and antispasmodic. They have been given, particularly, in intermittents and epilepsy, but are not now used. The flowers are called *Rosæ benedictæ*, *R. regizæ*.

PÆONICE, Medicine.

PÆONICUS, Medical.

PÆONIUS, Medical.

PÆOSYNE, Medicine

PAGAPOPLEX'IA, from *παῦν*, 'to nail fast,' and *ἀποπληξία*, 'apoplexy.' Apoplexy from cold.

PAGOS, Peritonæum.

PAIDIA, Infancy.

PAIDION, Infans.

PAIGIL, *Primula veris*.

PAIN, (old F.) *painer*, 'to torment;' Sax. *pinan*, 'to torture.' *Dolor*, *Algos*, *Alge'ma*, *Alge'do*, *Alge'is*, *Od'ynè*, *Ache*. (Sc.) *Hachæ*, (Prov.) *Crazy*, *Wark*, (F.) *Douleur*. A disagreeable sensation, which scarcely admits of definition. It is generally symptomatic, and is called *acute*, (F.) *aiguë*, when very violent, as a *twinge*; *pungent*, (F.) *ponctive*, when it resembles that which would be produced by a sharp instrument run into the part: *heavy*, (F.) *gravative*, when attended with a sensation of weight; *tensive*, when the part seems distended: *lan'cinating*, when occurring in shoots: *lac'erating* or *tearing*, when the part seems to be tearing: *burning*, (F.) *brûlante*, when resembling that produced by a burn, &c.

PAIN, *Son*, *Hemicrania*.

PAIN, see *Triticum* — p. *à Coucou*, *Oxalis acetosella* — p. *de Madagascar*, *Jatropha manihot* — p. *de Pourceau*, *Cyclamen*.

PAINS, *AFTER*, see P. labour.

PAINS, *GNAWING*, *Dolo'res roden'tes*. Pains resembling what might be produced by the gnawing of an animal.

PAINS, *Labour pains*, *Throes*, *Throws*, *Parturien'tis dolo'res*, *Odin*, *Odis*, *Dolo'res ad partum*, *Torm'ina parturien'tium*, (F.) *Mal d'Enfant*, *Douleurs*. The uneasy sensations that accompany labour, and are owing to uterine contraction. The pains that precede actual delivery, and are simply employed in dilating the os uteri, are called *grinding*; those which extrude the child, *forcing*. Those which take place in the first days after delivery, and are employed in forcing away coagula, &c., are termed *After pains*, *Dolo'res seu Torm'ina post partum*, *D. puerperarum*, (F.) *Tranchées utérines*.

"To take a pain." — "To try a pain," — is to make an examination per vaginam, during labour, to discover its progress, character, &c.

The French term the pains, which precede and announce labour, *mouches*; those which accompany it, *douleurs*; and those which occur immediately afterwards, *tranchées* on *coliques*.

PAINT, (from *pingere*, *pinctum*, 'to paint,' (I.) *pinto*, (F.) *peint*;) *Pigmen'tum*, *Fucus*, (F.) *Fard*. A name given to different substances, employed for embellishing the complexion or rendering the skin softer. The substances most commonly used for this purpose are:—the *Subnitrate of Bismuth*, (F.) *Blanc de fard*. A spirituous solution of benzoïn, precipitated by water, and forming *Virgin's milk*, (F.) *Lait virginal*; the red or rouge

of the carthamus; *Spanish white*; the oxides of lead, tin, and mercury, vermilion, red saunders, &c. All paints may be said to be noxious. They injure the skin; obstruct perspiration; and, thus, frequently lay the foundation for cutaneous affections. See Cosmétique.

PAINT, INDIAN, *Hydrastis Canadensis*, *Sanguinaria Canadensis* — p. Yellow, *Hydrastis Canadensis*.

PALAD, Vertex.

PALÆSTRA, Gymnasium.

PALATS, Palate.

PALAME, Palm.

PALATAL, Palatine.

PALATE, *Palatium*, *Hypero'a*, *Hyperö'um*, *Uranis'cos*, *Roof of the mouth*, *Fossa Palati'na*, *Palatum stab'ile*, *Hard Palate*, (F.) *Palais*. The upper part of the cavity of the mouth; a kind of parabolic vault, formed by the two superior maxillary bones, and the two palate bones, covered with a thick and dense mucous membrane: bounded, before and at the sides, by the superior dental arch; and, behind, by the velum palati. A whitish line runs along the middle, from before to behind; at the anterior extremity of which is a tubercle corresponding to the inferior orifice of the anterior palatine canal. The arteries of the palate and corresponding gums proceed from the palatine, alveolar, infra-orbital, labial, and buccal branches. The veins follow a similar course. The nerves are given off from the palatine, facial, infra-orbital, superior dental, and from the naso-palatine ganglion.

PALATE, ARTIFICIAL, (F.) *Obturbateur du Palais*. A plate of tortoise-shell, silver, &c., used to close a deficiency of the palate.

PALATE BONE, *Os Palati seu basilarë*, (F.) *Os Palatin ou du Palais*, is a small bone, of an irregular shape, situate at the posterior part of the nasal fossæ and palate. It seems formed by the union of two bony plates joined at right angles to each other; so that one is *inferior* and *horizontal*; — the other, *superior* and *vertical*. The *horizontal* or *palatine portion*, *Arch of the palate*, is quadrilateral. Its upper surface forms part of the nasal fossæ; the lower forms part of the palatine arch, and contains the inferior orifice of the *posterior palatine canal*. Its anterior edge is articulated with the palatine process of the superior maxillary bone; the posterior is free and sharp. It gives attachment to the velum palati, and presents an eminence, which, when joined with one on the opposite side, forms the posterior nasal spine. Its inner edge is articulated with the corresponding bone; its outer is confounded with the vertical portion. The *vertical* or *ascending portion* has the form of a long square. The inner surface forms part of the outer paries of the nasal fossæ. Its outer surface is unequal, and is articulated with the superior maxillary bone; behind, it presents a vertical groove, which forms part of the posterior palatine canal; and, above, a small free surface, which looks into the zygomatic fossa. Its anterior edge is united with the superior maxillary bone; the posterior is unequal, and rests on the internal ala of the pterygoid process. The angle, which it forms by uniting with the posterior edge of the horizontal portion, presents a very prominent pyramidal eminence, called the *tuberosity of the palate bone*. Its upper margin is surmounted by two eminences; — the one, anterior, called the *orbital process*; — the other posterior, the *sphenoid process*. These two processes are separated by an almost circular notch, which the sphenoid bone forms into a foramen, called the *sphenopalatine*.

This bone appears to be developed from a single point of ossification. It is articulated with

the sphenoid, ethmoid, the cornua sphenoidalia, superior maxillary bone, the inferior spongy bone, the vomer, and its fellow.

PALATE, CLEFT, see Harelip — p. Falling down of the, Uvulitis — p. Fissured, see Harelip — p. Hard, Palate — p. Soft, Velum pendulum palati.

PALATINE, *Pal'atal*, *Palati'nus*, *Pal'atic*, from *palatum*, 'the palate.' Relating or belonging to the palate.

PALATINE or PALATAL ARTERIES are two in number. They are distinguished into, 1. The *superior palatal* or *palatine*, *Palati'na descendens* or *Pal'ato-maxillary*, which leaves the internal maxillary behind the top of the orbit; descends into the pterygo-maxillary fissure, and enters the posterior maxillary canal. Its branches are distributed to the velum palati and the nasal fossæ. 2. The *inferior* or *ascending palatine* or *palatal*, which is very small, and separates from the facial near its origin. It passes along the lateral and superior part of the pharynx, between the pillars of the velum palati, and divides into a number of ramusculi, which are distributed to the pharynx, the corresponding tonsil, the Eustachian tube, and the velum palati.

PALATINE CANALS or DUCTS. These are two in number. The *anterior*, *Ductus incisivus* seu *Naso'lo-palati'nus*, *Incisive canal*, is situate at the anterior part of the palatine arch, and is formed by the two superior maxillary bones. It is single beneath, but opens above by two apertures, one into each nasal fossa. The foramen is called *Fora'men incisivum* seu *Palati'num anterius* seu *Alveola'rë anterius* seu *cævum osis maxilla'ris superio'ris*. The *posterior palatine canal* is situate at the posterior part of the palate. It is formed by the palate and superior maxillary bones, ascends between these two bones, and, in its course, gives rise to two or three small accessory palatine canals, which open above the tuberosity of the palate bone. It transmits the nerves and vessels which bear its name. The foramen is called, *F. Palati'num posterius* seu *Palato-maxilla'rë seu alveola'rë posterius*.

PALATINE MEMBRANE is the fibro-mucous membrane that lines the palate or roof of the mouth.

PALATINE or PALATAL NERVES, *Gutturopalatins* (Ch.), are three in number; and arise from the inferior part of the sphenopalatine ganglion. 1. The *great palatine*, *Palatin inférieur* (Ch.), is situate anteriorly to the two others, and descends into the posterior palatine canal; gives a *nasal filament* and a *staphyline filament*, and is distributed on the palate. 2. The *middle palatine*, *Guttural* (Ch.), which arises behind the preceding, enters one of the accessory palatine canals, and is distributed to the tonsils and velum palati. 3. The *little palatine nerve*, situate still more backward than the last, which loses itself by several filaments in the uvula, tonsils, and follicles of the palatine membrane.

PALATITE, Isthmitis.

PALATITIS, Uranisconitis.

PALATO-GLOSSUS, Glossostaphylinus.

PALATO-PHARYNGEUS, *Pharyngo-staphylinus*, *Staphylin'no-pharyngeus*, *Thyro-pharyngo-staphylin'us*, part of the *Hypero-pharyngeus*, (F.) *Palato-pharyngien*, *Stylo-pharyngien* (Ch.). A membranous muscle, broader at its middle than at its extremities, and situate vertically in the lateral paries of the pharynx and velum palati. It includes the peristaphylo-pharyngeus, pharyngo-staphylinus, and thyro-staphylinus of Winslow. When the two palato-pharyngei contract together, they depress the velum. They elevate and shorten the pharynx, and act principally in deglutition.

PALATO-PHARYNGIEN, Palato-pharyn-

geus — *p. Salpingeus*, *Circumflexus* — *p. Salpingien*, *Circumflexus*.

PALATOSTAPHYLINI, see *Azygos muscle*.

PALATUM FISSUM, *Hyperoochasma*, see *Harelip* — *p. Mobile*, *Velum pendulum palati* — *p. Molle*, *Velum pendulum palati* — *p. Pendulum*, *Velum pendulum palati*.

PALE, *Pollen*.

PALENESS, *Pallor*, *Och'rotes*, *Ochro'ma*, *Och'r'asia*, *Leuco'sia*, *Dealbatio*, *Luror*, (F.) *Pâleur*. Whiteness of complexion. It appears to be owing to a diminution or alteration of the blood circulating in the capillary vessels, and is often a sign of disease.

PALES COULEURS, *Chlorosis*.

PALETTE (F.), *Pal'mula*, *Fer'ula*. An instrument of percussion. It has the shape of a spatula with a long handle; is made of very light, white wood, and has been proposed, by Mr. Percy, for what is called the *Massage*, a kind of pressure or kneading, exerted by the hand on the body and limbs, to excite the tone of the skin and sub-jacent tissues.

Palette also means a cup for bleeding; and the xiphoid cartilage [?].

PALETTE À PANSEMENT. A small splint of thin, scooped wood, having the shape of the hand, and used to support the hand in certain cases—when burnt, for example—in order to prevent the fingers from being deformed by the cicatrices.

PALETTE DE CABANIS. An instrument, invented by Cabanis of Geneva to draw out the extremity of the stylet when passed into the nose, in the operation for fistula lacrymalis. It is composed of two plates of silver, pierced with several holes, and movable on each other. To use it, the instrument must be carried into the nose, so that the holes of the two plates correspond. When the end of the stylet is received into one of these holes, it is pinched, by sliding one of the plates on the other, and is easily withdrawn.

PALEUR, *Paleness*.

PALI PLAGUE, see *Plague*.

PALICOU'REA CROC'EA, *Psycho'tria croc'ea*, *Order*, *Cinchonaceæ*, a native of the West Indies, is possessed, according to Von Martius, of emetic properties.

PALICOU'REA MARGRAA'VII, *Erva do rato*, *Galv'ania Vellozii*, a native of Brazil, is poisonous, and employed there to kill rats and mice.

PALICOU'REA OFFICINA'LIS, a native of Brazil, is reputed to be a powerful diuretic in small doses.

PALICOU'REA SPECIO'SA, *Goldshrub*, a native of New Granada and Brazil. The leaves are considered to be antisiphilitic. The decoction in large doses is poisonous.

PALILLO, *Campomanesia lineatifolia*.

PALIMPISSA, see *Pinus sylvestris*.

PALINCOTESIS, *Recrudescencia*.

PALINDROME, *Palindromia*.

PALINDROM'IA, *Palin'dromê*, from *παλιν*, 'again,' and *δρῶμος*, 'a course.' This word signifies, with some, a reflux of fluids from the exterior towards the interior; and, according to others, the relapse of an affection—the recurrence—*Recursio*, *Repetit'io*—of a paroxysm, or disease,—*Morbus recidi'vus*, *Morbi recur'sus*.

PALINGENESIS, *Regeneration*.

PALLA, *Pila*.

PALLAT, *Vertex*.

PALLI'ATION, *Palliatio*, from *palliare*, *palliatum*, 'to cover, mask;' from *pallium*, 'a mantle.' The act of palliating—that is, of causing the symptoms of a disease to mitigate, without

curing it. The medical art can frequently only moderate the symptoms of a disease, prevent it from making progress, prolong the days of the patient, and diminish his sufferings. This is *palliative treatment*, (F.) *Traitement palliatif*.

PALLIATIVE, *Palliatus*, *Antipath'ic*, *Enantipath'ic*, (F.) *Palliatiif*. A remedy or mode of treatment, which only relieves a disease, without curing it. The *palliative treatment*—*Cura palliativa*, is placed in antithesis to the radical, *Cura radical'is*; see *Radical*.

PALLIDULUS, *Blafard*.

PALLIDUS MORBUS, *Chlorosis*.

PALLOR, *Paleness* — *p. Virginum*, *Chlorosis*.

PALM, *Palma*, *πάλην*, *Vola*, *V. manûs*, *Plate'a*, *Thenar*, *Agos'tus*, *Poc'ulum*, *Pat'era*, *Suppelle'x* seu *Suppelle'ctile* *Diog'enis*, (F.) *Paume* ou *Creux de la main*. The hollow or inside of the hand. It is limited, *without*, by the thenar eminence, which answers to the thumb; *within*, by the hypothenar eminence, which answers to the little finger; *above*, by the wrist-joint; and, *below*, by the four fingers.

PALM, *Gomphro*, see *Toddy*.

PALMA CHRISTI, *Ricinus communis* — *p. Cocos*, *Cocos nucifera* — *p. Dactylifera*, *Date tree* — *p. Humilis*, *Musa paradisiaca* — *p. Indica* *vinaria secunda*, see *Toddy* — *p. Unguentariorum*, *Myrobalanus*.

PALMÆ U'TERI PLICA'TÆ, *Arbor vite uteri'nis*, *A. u'teri viv'ificans*. The arborescent striæ in the interior of the cervix uteri.

PALMAIRE, *Palmar* — *p. Cutané*, *Palmaris brevis* — *p. Grand*, *Palmaris magnus* — *p. Grêle*, *Palmaris longus* — *p. Long*, *Palmaris longus*.

PALMAR, *Palma'ris*, (F.) *Palmaire*, from *palma*, 'the palm of the hand.' That which concerns the palm of the hand. The palm has, sometimes, been divided into three regions, called *palmar*. 1. The *external palmar region* corresponds to the thenar eminence. 2. The *internal palmar region* is formed by the hypothenar eminence; and 3. The *middle palmar region* is the hollow of the hand, (F.) *Creux de la main*.

PALMAR APONEURO'SIS, *Palmar fasc'ia*. A strong triangular aponeurosis, which arises from the inferior tendon of the palmaris brevis, and the anterior annular ligament of the carpus. It covers the whole palm of the hand, and adheres, somewhat strongly, to the skin of the part. The *palmar ligaments* are small, very numerous, fibrous fasciæ, which unite the different bones of the carpus and metacarpus.

PALMAR ARCHES, (F.) *Arcades* ou *Crosses palmaires*. The curved extremities of the radial and cubital arteries in the palm of the hand. There are two palmar arches:—the one, *cubital* or *superficial*,—*grande Arcade palmaire* of Chaussier, furnishes, at its convexity towards the fingers, the collateral arteries of the fingers, and the internal collateral of the thumb; the other, the *radial*, *profound*, or *deep-seated*,—*petite Arcade palmaire* of Chaussier, is deep in the palm of the hand beneath the tendons of the flexor muscles of the fingers. It gives some small branches only to the surrounding parts. The palmar veins accompany the corresponding arteries.

PALMAR ARTERIES, see *Palmar arches* — *p. Fascia*, *P. aponeurosis*.

PALMAR MUSCLES. These are three in number. 1. *Palma'ris Magnus*, *Flexor carpi radialis*, *Radialis intern'us*, (F.) *Épitrochlo-métacarpien* (Ch.), *Palmaire grand*, *Radial antérieur*. A long fusiform muscle, which passes obliquely from the inner tuberosity of the humerus to the upper extremity of the second metacarpal bone. It bends the hand. 2. *Palmaris longus*, *Latecen'tis chorda*, *Ulna'ris grac'ilis*, (F.) *Épitrochlo-carpi-pal-*

maire (Ch.), *Palmaire long*, *P. grêle*. This muscle is situate on the inner side of the last. It is very slender, and passes from the inner tuberosity of the humerus to the anterior annular ligament of the carpus, and to the palmar aponeurosis, which it seems to form by the vanishing of its fibres. It stretches the palmar aponeurosis, and bends the hand. 3. *Palmaris Brevis*, *Caro quadrata*, *Carpia'us*, (F.) *Palmaire cutané*. A small, thin, quadrilateral muscle; situate before the thenar eminence. It passes from the annular ligament of the carpus to the integuments of the palm of the hand, which it wrinkles by its contraction.

PALMAR VEINS, see Palmar arches.

PALMARIS BREVIS, see Palmar muscles—*p. Longus*, see Palmar muscles—*p. Magnus*, see Palmar muscles.

PALMETTO, SAW, *Chamærops serrata*.

PALMI-PHALLANGIENS, *Lumbricales manus*.

PALMISTRY, *Chiromancy*.

PALMOSCOPIA, from *παλμος*, 'palpitation,' and *σκοπεω*, 'I view.' Investigation of the beating of the heart and pulse. Prognosis from the beating of the heart and pulse.

PALMULA, *Date*, *Palette*.

PALMUS, *Palpitation*—*p. Cordis*, *Palpitation*—*p. Plumbarius*, *Colic*, *metallie*—*p. Vomitus*, *Vomiting*.

PALO DE CALENTURAS, *Calenturas*—*p. de Vacca*, *Galactodendron utile*—*p. de Velas*, *Parmentiera cerifera*.

PALOMMIER, *Gaultheria*.

PALOS DE CALENTURA, *Cinchona*.

PALPATION, from *palpare*, *palpatum*, (F.) *palper*, 'to feel.' The sense of touch. It is also used for the mode of exploring diseases by feeling or pressing upon the diseased organ. A delicate and discriminating touch—*tactus eruditus*—is important in the diagnoses of many morbid conditions.

PALPEBRA, supposed to be from *palpitare*, 'to palpitate,' owing to its frequent motion. [?] *Blepharon*, *Ope'culum* seu *Op'erculum oculi*, *Eyelid*, (Prov.) *Eye-breck*, (Sc.) *Eye-winker*, (F.) *Paupière*. The eyelids are two movable vela, situate in front of the eye; which by their separation permit the organ to receive the impression of light, or by their greater or less occlusion protect it from the influence of too great light, or from the action of extraneous bodies. They are separated, above, from the forehead, by the eyebrows; and, below, are confounded with the cheeks. They are distinguished into *upper*, *Palpebra major seu superior*, *Epice'lis*—and *lower*, *Palpebra minor seu inferior*, *Hypoc'e'lis*, *Hypoc'e'lum*, *Hypoc'yllum*. The former is much broader and more movable than the latter. They are separated by a transverse cleft—*Fissura palpebrarum*—and their edges are thick, and furnished with the cilia or eyelashes. Supported by the tarsal cartilages, they unite at the extremities of the transverse diameter of the orbit, forming two angles, the innermost being called the *greater angle of the eye*, or *inner or greater canthus*; the other, the *lesser*, or *external angle or canthus*. They are formed, 1. Of a thin dermoid texture: 2. Of a muscular membrane.—a portion of the *orbicularis*: 3. Of a particular fibrous membrane: 4. Of the tarsi: 5. Of a mucous membrane,—a part of the *tunica conjunctiva*.

Their vessels and nerves are called *Palpebral*. *PALPEBRA INFERIOR EXTRORSUM FLEXA*, *Ectropion*—*p. Tertia*, *Valvula semilunaris*.

PALPEBRÆ SUPERIORIS PRIMUS, *Levator palpebræ superioris*.

PALPEBRAL, *Palpebra'lis*, from *palpebra*, 'the eyelid.' That which belongs or relates to the eyelid.

PALPEBRAL ARTERIES are those distributed to the eyelids. They are two in number; an *upper* and a *lower*. The former arises from the ophthalmic, a little beyond the cartilaginous pulley of the greater oblique muscle, and descends towards the upper eyelid, over which it is distributed. The latter arises at the side of, and a little behind, the preceding, and sends its ramifications to the lower eyelid and the neighbouring parts. The palpebral veins have a similar arrangement; except that some of them open into the temporal and labial veins.

PALPEBRAL FLUX, *Puriform*. A puriform discharge, furnished by the glands of Meibomius, when inflamed; and to which Scarpa attributes *fistula lacrymalis*, in many cases.

PALPEBRAL LIGAMENTS are distinguished into *superior* and *inferior*. Some anatomists, with Winslow, have described, under this name, the fibrous layer which is attached,—on the one hand, to the outer part of the circumference of the orbit; on the other, to the tarsal cartilages.

PALPEBRAL NERVES, *Nerves of the eyelids*, arise, particularly, from the frontal and lacrymal branches; from the ophthalmic nerve of Willis; and from the facial, nasal, and infra-orbital branch of the superior maxillary nerve.

PALPEBRALIS, *Orbicularis palpebrarum*.

PALPEBRARUM APERIENS RECTUS, *Levator palpebræ superioris*—*p. Duo musculi*, *Orbicularis palpebrarum*.

PALPEBRATIO, *Nictation*.

PALPITATIO, *Palpitation*—*p. Arteriarum*, see *Palpitation*—*p. Cordis* *trepidans*, *Cardiotromus*.

PALPITATIO MEMBRORUM. A term used by Gaubius for involuntary shaking of the legs.

PALPITATION, *Palpita'tio*, *Palmus*, *P. seu Tremor Cordis*, *Palpita'tio seu Pula'tio Cordis*, *Vibra'tio Cardi'aca seu Cordis*, *Cardi'mus*, *Cardiopal'mus*, *Subsul'tio*, *Clonus Palpita'tio*, (Sc.) *Dunt*, *Dount*, (F.) *Palpitation*, *P. du Cœur*, from *palpitare*, *palpitatum*, 'to throb.' A name given to the pulsations of the heart, when they become stronger, more extensive, or more sensible to the individual than ordinarily. Palpitations occur in a number of affections. They are, sometimes, owing to organic diseases of the heart—hypertrophy, inflammation, &c.; at others, to a disease in some other viscus; whilst, frequently, they are merely nervous.

PALPITATION is sometimes felt in the arteries, constituting *Palpita'tio Arterio'sa*.

PALPITATION DU CŒUR, *Palpitation*.

PALSTED, *Paralytic*.

PALSY, *Paralysis*—*p. Bell's*, *Paralysis*, *Bell's*—*p. Histrionic*, *Paralysis*, *Bell's*.

PALSY, LEAD, *Paral'ysis saturni'na seu metal'lica*, *Molybdopar'esis*, (F.) *Paralysie saturnine*, *Metal'lic Palsy*, *Painters' Palsy*. The paralysis of the upper extremities especially, induced by the poison of lead. Palsy of the intestines, thus induced, is called, by Piorry, *Anet'roner'vie saturnine*.

PALSY, METALLIC, see *Palsy*, lead—*p. Painters'*, *P. lead*—*p. Shaking*, *Paralysis agitans*—*p. Trembling*, *Paralysis agitans*.

PALSY WORT, *Primula veris*.

PALTA, *Persea gattissima*.

PALUDAL, *Elodes*.

PALUDAPIUM, *Apium graveolens*.

PALUDÉEN, *Elodes*.

PALUDOSUS, *Elodes*.

PALUS, *Marsh*, *Penis*—*p. Sanctus*, *Guaiacum*.

PALUSTRE, *Elodes*.

PAMBIOMA, *Panbioma*.

PAMPOISON, *Syncope*.

PAMPINIFORM PLEXUS, *Corpus pampini-forme*.

PAMPINIFORME CORPUS, see *Corpus pampiniforme*.

PAMPLE/GIA, *Panplegia*, from *πav*, 'all,' and *πληγω*, 'I strike.' General paralysis. Palsy of the whole body.

PANACEA, *panacea*, *Cathol'icon*, *Cathol'icum*, from *πav*, 'all,' and *ακεπα*, 'I cure.' *Universale remedium*, *Panchres'tos*. A pretended universal remedy. There is no such thing in existence.

PANACEA DUCIS HOLSATIÆ, Potassæ sulphas — p. *Duplicata*, Potassæ sulphas — p. *Lapsorum*. *Arnica montana* — p. *Mercurialis*, *Hydrargyri sublimatus* — p. *Mercurii rubri*, *Hydrargyri nitrico-oxidum* — p. *Pectoris*, *Glechoma hederacea* — p. *Vegetabilis*, *Crocus*.

PANACÉE ANGLAISE, *Magnesie carbonas*.

PANADA, diminutive of Ital. *paine*, 'bread.' (Lat. *panis*.) *Pana'ta*, *Panatel'la*. Bread, boiled in water to the consistence of pap. Sometimes administered to the sick.

SIMPLE BREAD PANADA may be made by putting any quantity of grated stale bread into enough water to form a moderately thick pulp; covering it up, and leaving it to soak for an hour; then beating it up with two tablespoonfuls of milk, and a small portion of refined sugar, and boiling the whole for ten minutes, stirring all the time.

PANAIS, *Pastinaca sativa*.

PANARIS, *Paronychia*.

PANARITUM, *Paronychia*.

PANARIUM, *Paronychia*.

PANATA, *Panada*.

PANATELLA, *Panada*.

PANAX, see *Panax quinquefolium* — p. *Costivum*, *Pastinaca opoponax* — p. *Heracleum*, *Pastinaca opoponax* — p. *Pastinacea*, *Pastinaca opoponax*.

PANAX QUINQUEFOLIUM, *Aureliana* seu *Ard'ia Canadensis*, *Ginseng*, *Ninsing*, *Gensang*, *Red-berry*, *Five fingers*. Same etymon as *Panacea*. *Family*, *Araliaceæ*. *Sex. Syst.* *Polyandria Monœcia*. This plant grows in North America. The root — *Panax* (Ph. U. S.) — *Radix ginseng* — is about the thickness of the little finger; an inch or two in length, often dividing into two branches; of a whitish-yellow colour; wrinkled on the surface, and of a compact, almost horny, texture. It has no smell, but a very sweet taste, combined with a slight degree of aromatic bitterness. It is much esteemed by the Chinese, who pay a high price for it. It is a gentle and agreeable stimulant.

PANAX SCHINSENG, *Asiat'ic Ginseng*, a native of Asia, has been usually confounded with the last. The root, *radix ninsi*, has like virtues.

PANBIO'MA, *Pambio'ma*, from *πav*, 'all,' and *βίος*, 'life.' The original animating principle. The general principle of life.

PANCHRESTOS, *Panacea*.

PANCHYMAGO'GUM, *Pantago'gon*, from *πav*, 'all,' *χυμος*, 'juice,' and *αγω*, 'I drive away.' A medicine which the ancients believed capable of purging away all sorts of humours.

PANCHYMAGOGUM MINERALE, *Hydrargyri sublimatus*.

PANCRENOSUS, *Pandemic*.

PANCRENUS (MORBUS), *Pandemic*.

PANCRATIUM, *Pancreas*.

PAN'CREAS, from *πav*, 'all,' and *κρεας*, 'flesh,' i. e. 'quite fleshy.' *Callic'reas*, *Pancrat'ium*, *Panc'reum*, *Lactes*, *Pidui'var ventriculi*, *Totum carnosum*, *Gland'ula salival'is abdom'inis*, *Panc're'nē*, *Sweetbread*. A gland, deeply seated in the abdomen, and lying transversely on the vertebral column, between the three curvatures of the duodenum, behind the stomach, and to the right of the spleen. Its left extremity is called the 'tail of the pancreas'; it is smaller than the

right, which is called the *head*. At its right portion, it has, almost always, a greater or less prolongation; called, by Winslow, and some others, the *lesser pancreas*, (F.) *Petit pancréas*, *Pancreas of Aselli*, *Pancreas minus*. The pancreas resembles the salivary glands in structure, and has been called the *Abdominal Salivary Gland*. It is composed of lobes and granulated lobules, distinct, and united by areolar tissue. From each of the granulations of these lobes arise the radicles of its excretory duct, which are very delicate, and united like veins. The duct itself, *Ductus Pancreat'icus* seu *Wirsung'ianus*, *Pancræat'ic duct*, *Canal or Duct of Wirsung or Wirsungus*, (F.) *Canal pancréatique*, proceeds in a serpentine course through the substance of the organ; and when it has reached behind the second portion of the duodenum, it becomes free, and is of the size of a crow's quill. It opens at an acute angle into the choledoch duct, or proceeds close to it, opening separately into the duodenum. The *Pancræatic arteries* are very numerous and small. They proceed from the celiac, splenic, superior mesenteric, right gastro-epiploic, coronaria ventriculi, and left capsular. Its veins open into the radicles of the vena porta; and, particularly, into the splenic and lesser mesenteric. Its nerves emanate from the solar plexus, and its lymphatics pass into ganglions, to which it gives its name. The pancreas secretes the *Pancræatic juice*, *Succus seu Liquor Pancreat'icus*, *Lympha Pancreat'is*, (F.) *Suc pancréatique*, which resembles the saliva. When this juice is mixed with amylaceous matters, it converts them into dextrin and glucose. Its great use appears, however, to be, to emulsify fatty matters by virtue of a peculiar albuminoid principle — *pancræatin* — coagulable by heat or alcohol, which it contains.

PANCREAS OF ASELLI, see *Pancreas* — p. *Minus*, see *Pancreas* — p. *Petit*, see *Pancreas* — p. *Second*, *Brunner's glands*.

PANCREAT'AL/GIA, from *pancreas*, and *αλγος*, 'pain.' Pain or disease of the pancreas.

PANCREATEMPHRAX'IS, from *pancreas*, and *εμπρασσω*, 'I obstruct.' Obstruction of the pancreas.

PANCREAT'IC, *Pancræat'icus*, (F.) *Pancréatique*. Same etymon as *Pancreas*. Relating to or affecting the pancreas.

PANCREATIC ARTERIES, see *Pancreas* — p. *Duct*, see *Pancreas* — p. *Juice*, see *Pancreas* — p. *Nerves*, see *Pancreas* — p. *Veins*, see *Pancreas*.

PANCREAT'IC SARCO'MA OF ABERNETHY, *Emphy'ma sarco'ma pancræat'icum*. A tumour, in irregular masses, connected by loose fibrous substance, like the irregular masses of the pancreas. Found in the areolar substance: but, more usually, in lymphatic glands, chiefly in the female breast.

PANCREATIN, see *Pancreas*.

PANCREATITIS, *Inflamma'tio pancræat'is*, (F.) *Inflammation du Pancréas*, from *pancreas*, and *itis*, denoting inflammation. Inflammation of the pancreas; — an affection, the phenomena of which are not well known.

PAN'CREATOID, from *pancreas*, and *ειδος*, 'appearance.' *Pancræatoid'es*. A tumour resembling the pancreas in structure.

PANCREATON'CUS, from *pancreas*, and *ογκος*, 'a swelling.' Tumour of the pancreas.

PANCRENE, *Pancreas*.

PANCREUM, *Pancreas*.

PANDA'LEA, *Panda'leum*. A kind of lozenge used by the Arabian physicians.

PANDALEUM, *Pandalea*.

PANDALITIUM, *Paronychia*.

PANDEMIA, *Pandemic*.

PANDEM'IC, *Pan'demy*, *Pande'mia*, *Morbus Pandem'ius* seu *Pande'nus* seu *Panca'nus*, *Pan-*

corno'us, (F.) *Pandémie*; from *παν*, 'all,' and *ἄνθος*, 'people.' An epidemic which attacks the whole, or a great part, of the population.

PANDÉMIE, Pandemic.

PANDEMY, Pandemic.

PANDICULATION, *Pandiculatio*, *Scordine'ma*, *Scordinismus*, *Clonus Pandiculatio*, from *pandiculari*, (*pandere*, 'to stretch out,') 'to stretch as one awaking from sleep.' *Stretching*, *Hal'ices*, (pl.) An action, which consists in elevating the upper limbs, and throwing the head and trunk backwards, at the same time that the lower limbs are extended. This series of motions of the extensor muscles is in some measure independent of the will, and is accompanied with yawning, which is equally involuntary. In the state of health, stretching occurs before and after sleep; especially when we are fatigued. It is also observed at the onset of certain paroxysms of fever, hysteria, &c.

PANDOCE'UM, *Pandoci'um*, *Pandoci'um*, from *παν*, 'all,' and *δεχομαι*, 'to receive.' An inn that receives all classes. A general hospital, or one that admits all cases.

PANDOCCHIUM, *Pandocum*.

PANDOCIUM, *Pandocum*.

PANHIDROSIS, from *παν*, 'all,' and *ἵδρωσις*, 'sweating.' *Sudor universal'is*. Sweating over the whole body.

PANHYPEREMIA, *Plethora*.

PANIC D'ITALIE, *Panicum Italicum*.

PANICAUT, *Eryngium campestre*—*p. Martine*. *Eryngium maritimum*.

PANIC GRASS, *ITALIAN*, *Panicum Italicum*.

PANICUM DACTYLON, *Cynodon dactylon*.

PAN'ICUM ITAL'ICUM, from *panis*, 'bread,' *Italian* *Panic Grass*, *Indian Millet*, *Guinea Corn*, *Barba'does Millet*, *Mil'ium Indicum*, *Sorghum*, *Haleus Sorghum*, (F.) *Millet des oiseaux* ou à grappes, *Panic d'Italie*. *Nat. Ord.* Gramineæ. The seed—*Mil'ii Sem'ina*—are much esteemed in Italy, being a constant ingredient in soups, and made into a variety of forms for the table.

PANICUM MILIA'CEUM, *P. mil'ium*, *Common Millet*, *Milium*, *M. Pan'icium* seu *Esculent'um*, *Cenchrus*, *Cenchrus*, (F.) *Mil'ot ordinaire*, *Mil.* Millet-seed are esteemed a nutritious article of diet, and are often made into puddings.

PANICUM MILIUM, *P. miliaceum*.

PANIS CANICACEUS, see *Canicæ*—*p. Cibar'ius* seu *secundarius*, *Syncomistes*—*p. Cuculi*, *Oxalis acetosella*—*p. Porcinus*, *Cyclamen*.

PANIV'OROUS, *Paniv'orus*, from *panis*, 'bread,' and *voro*, 'I devour.' Subsisting on bread.

PANNI, Swathing clothes.

PANNICLE, Membrane.

PANNICULE CHARNU, *Panniculus carnosus*.

PANNIC'ULUS, diminutive of *pannus*, cloth. A piece of fine cloth. *Pterygion*.

PANNICULUS ADIPOSUS, *Cellular membrane*.

PANNICULUS CARNO'SUS, *Tu'nica carno'sa*, *Muscular Web*, *Mantle*, (F.) *Pannicule charnue*, is a general covering found in the quadruped, and formed by a thin, subcutaneous muscle, which serves to move the skin. It exists only in certain parts of the human body,—as in the forehead, where it is formed by the occipito-frontalis; in the neck, by the platysma myoides; and, in the scrotum, by the cremaster muscle.

PANNICULUS CORDIS, *Pericardium*—*p. Hymenæus*, *Hymen*—*p. Virginalis*, *Hymen*.

PANNIKELL, *Cranium*.

PANNUS. A piece of cloth. This word has several other meanings, many of which are equivocal. It means a tent for a wound, and, also, an irregular nœvus or mark on the skin.—*Castelli*. Modern ophthalmologists apply it to an

opaque and vascular condition of the anterior membrane of the cornea, almost invariably produced by the continuance of purulent or strumous ophthalmia, or by chronic inflammation of the cornea. This part becomes nebulous, covered with red vessels, and opaque. At times, the sclerotic conjunctivæ undergoes a similar change to that which occurs in the corneal portion.

PANNUS HEPATICUS, *Chloasma*—*p. Lenticularis*, *Ephelides*.

PANOCHIA, *Bubo*.

PANOPHO'BIA, *Pantopho'bia*, *Panpho'bia*, *Pavopho'bia*, *Athym'ia*, *Pavor*, (F.) *Frayeur nocturne*, *Souci*; from *Παν*, 'the god Pan,' and *φοβος*, 'fear,' a panic or fear inspired by the god Pan, according to the opinion of the ancients. Others derive it, and with more probability, from *παν*, 'all,' and *φοβος*, 'fear.' This universal and groundless fear is morbid, and is a symptom of melancholy.

PANOPHTHALMITIS, from *παν*, 'all,' and *ophthalmitis*, 'inflammation of the eye.' Inflammation of the whole eye.

PANPHOBIA, *Panophobia*.

PANPLEGIA, *Pamplagia*.

PANSCLERO'SIS, from *παν*, 'all,' and *σκληρωσις*, 'hardness.' Complete induration of a part.

PANSE, *Ingluvies*.

PANSEMENT, *Dressing*.

PANSIES, *Viola tricolor*.

PANSPER'MIA, from *παν*, 'all,' and *σπερμα*, 'sperm.' The theory of *Dissemination of Germs*, according to which, ova or germs are disseminated over all space, undergoing development under favourable circumstances.

PANSY, *Viola tricolor*.

PANTAGOGON, *Panchymagogum*.

PANTANANCEPHA'LIA, from *παν*, *παντος*, 'all,' and *anencepha'lia*, absence of brain. Entire absence of brain—the same as *anencephalia*.—*I. G. St. Hilaire*.

PANTATROPH'IA, from *παν*, *παντος*, 'all,' and *ατροφία*, 'atrophy.' Universal atrophy.

PANTHOD'IC, *Panthod'icus*, from *παν*, *παντος*, 'all,' and *ὅδος*, 'a way.' An epithet applied by Dr. Marshall Hall, to a course of nervous action proceeding from one point in all directions to every other.

PANTICES, *Intestine*.

PANTING, *Anhelatio*.

PANTOGANGLITIS, *Cholera*.

PANTOPHAGISTS, see *Omnivorous*.

PANTOPHAGUS, *Omnivorous*.

PANTOPHOBIA, *Hydrophobia*, *Panophobia*.

PANTOZOOTIA, *Panzootia*.

PANUS, *Pytheglon*—*p. Faucium*, *Stomatopanus*—*p. Inguinalis*, *Bubo*.

PANZOÔT'IA, *Pantozoôtia*; from *παν*, 'all,' and *ζῶον*, 'animal.' An endemic or epidemic disease affecting animals generally.

PAP, *Mamma*, *Nipple*. Also, a soft food made of bread boiled or softened with water, (Sc.) *Breadberry*.

PAP OF THE HASS, *Uvula*—*p. of the Throat*, *Uvula*.

PAPAS AMERICANUS, *Solanum tuberosum*.

PAPA'VER, *P. somniferum*, *Mecon*, *Lethe'a*. The *White Poppy*, *Papaver sati'vum*, (F.) *Pavot des Jardins*. *Ord. Papaveracæ*. *Sez. Syst.* *Polyandria Monogynia*. The *CAPSULE OF THE WHITE POPPY*, *Κόδια*, *κωδια* or *κωδια*, *Scaphion*, *Papaveris Caput* seu *Cap'sula*, *Papaver*, (Ph. U. S.), *Mecon*, *Poppyhead*, is used in fomentation as an anodyne and relaxant, to inflamed or ulcerated parts. The expressed oil of the seeds is called by the French *Huile d'Œillette*. It is from the poppy that *OP'ium*—called, also, *Op'ium Theba'icum*, *Meco'nium*, *Op'ion*, *Manu Dei*, *Asion Aff'ium*, *Amphiam*, *Afium*, *Osoron*, *Lad'anum*,

Mecon, Sopo, (Prov.) Poppy pill, (F.) Opium brut ou cru—is procured. It is the milky juice obtained after the flowering of the plant, by making longitudinal incisions in the capsules and stalks. It is commonly obtained from the Levant or India, where the poppy grows abundantly. Opium is a compound of morphia, narcotina, meconic acid, codeia, narcein, meconin, caoutchouc, fecula, resin, oil, and several other substances. *Turkey Opium* is of a heavy and slightly fetid colour. Its taste is nauseous, bitter, acrid, and warm: it is in flat pieces; solid and tenacious; of a dark brown colour, and yellowish when powdered; making a light brown, interrupted streak on paper. In *East India Opium*, the odour is the same, and empyreumatic; the taste is less bitter, but more nauseous, and the colour darker. The varieties of opium known in commerce are the *Smyrna*, the *Constantinople*, the *Egyptian*, the *Persian*, and the *Indian*. Opium is stimulant in small doses; sedative, in large; narcotic and anodyne; operating through the nervous system. It is a most valuable remedy in all painful affections, and inflammations, when given so as to produce its full, sedative effects, especially in peritonitis. In diarrhoea and dysentery, it acts as an astringent; and in intermittents, it will often prevent the paroxysm. In typhus, in small doses, it acts as a cordial; in larger, it allays irritation and produces sleep. When applied externally, its narcotic effects are developed: and, hence, it is a useful application to irritable sores, painful tumours, &c. Dose, gr. $\frac{1}{4}$ to $\frac{1}{2}$ as a stimulant; gr. $\frac{1}{4}$ to $\frac{1}{2}$ as a sedative.

PAPAVER CEREALE, P. rhœas—p. Corniculatum, Chelidonium majus—p. Erraticum, P. rhœas—p. Luteum, Chelidonium majus.

PAPAVER NIGRUM. This is merely a variety of the white poppy, producing black seeds.

PAPAVER RHÆAS. The systematic and pharmacopœial name of the *Red corn-poppy*, *Corn rose*, *Papaver erraticum seu cereale seu rubrum*, *Copper rose*, *Red-mailes*, (F.) *Coquelicot*. The heads of this species contain a milky juice of a narcotic quality. A syrup, *Syrupus Rhœados*, of its flowers—*Rhœas* (Ph. L. and E.)—has been directed, which has been esteemed diaphoretic and slightly soothing; but is more admired for its colour than for any other property.

PAPAVERACEUM, Opiate.

PAPAVERCULUM, Meconium.

PAPAVERINE, *Papaverina*; from *papaver*, 'the poppy.' An alkaloid contained in opium; but not much investigated.

PAPAVERINUM, Codeine.

PAPAW TREE, *Carica papaya*.

PAPAYER, *Carica papaya*.

PAPER, BLISTERING, see Sparadrapum vesicatorium.

PAPIER ÉPISPASTIQUE, see Sparadrapum vesicatorium.

PAPILLA, (F.) *Papille*—diminutive, perhaps, from *papa*, one of the earliest cries of the infant, ascribed to a craving for food. The end of the breast, the nipple. See Nipple. An eminence similar to a nipple. A name given to small eminences, more or less prominent, at the surface of several parts, particularly of the skin and mucous membranes, in which latter situation they are also called villi. See Villous. They are formed by the ultimate expansion of the vessels and nerves, and are susceptible, in some cases, of a kind of erection. They are also called *Papillæ nervosæ* seu *Nervæ*, *P. Pyramidales*, *Nervous papillæ*, (F.) *Papilles* on *Papilles nerveuses*, *Houpes nerveuses*.

PAPILLA LACRYMALIS, see Lachrymal puncta—p. Mamme, Nipple—p. Nervi optici, see Optic nerve—p. Pili, see Hair.

PAPILLÆ, see Corpuseles, Pacinian—p. Calyciformes, see Papillæ of the tongue—p. Capitatæ, see Papillæ of the tongue—p. Circumvallatæ, see Papillæ of the tongue—p. Conical, see Papillæ of the tongue—p. Conoidal, see Papillæ of the tongue—p. Filiform, see Papillæ of the tongue—p. Fungiform, see Papillæ of the tongue—p. Fungoid, see Papillæ of the tongue.

PAPILLÆ OF THE KIDNEY, (F.) *Mamelons du rein*, are the projections into the pelvis of the kidney of the apices of the cones—*Pyramids of Malpighi*, *Pyramides Malpighiæ*—of the tubular substance of the organ. They are also called *Processus mamillares*, *Curvculæ papillares*, and *Papillæ renum*.

PAPILLÆ, LENTICULAR, see Papillæ of the tongue—p. Maxime, see Papillæ of the tongue—p. Medie, see Papillæ of the tongue—p. Minime, see Papillæ of the tongue—p. Mucosæ, see Papillæ of the tongue—p. Nervæ, Corpuseles, Pacinian—p. Nervous, see Papilla—p. Pyramidal, see Papillæ of the tongue—p. Pyramidales, see Papilla—p. Renum, see Papillæ of the kidney—p. Semilenticulares, see Papillæ of the tongue.

PAPILLÆ OF THE TONGUE are very numerous and marked. They are observed, particularly, at the upper surface of the organ, and are distinguished into—1. The *Lenticular papillæ*, *Papillæ mucosæ seu maxime seu capitatæ seu vallatæ seu circumvallatæ seu truncatæ seu calyciformes*. They are arranged at the base of the tongue in two oblique lines having the shape of the letter V: and united angularly at the foramen cæcum. They have, in general, an ovoid or spherical shape, and have seemed to be mucous follicles similar to those of the palate, lips, &c. 2. The *Fungiform or Fungoid Papillæ*, *Papillæ medice seu semilenticulares*, are of indeterminate number, but always more numerous than the preceding. They are irregularly disseminated near the edges and the point of the tongue; have a rounded and flattened head, supported by a short and narrow footstalk. Their tint is whitish. 3. The *Conical, Conoidal or Pyramidal papillæ*, *P. minime seu villosæ*, are the most numerous of all. Their arrangement is more regular in the forepart of the tongue than behind. They resemble small cones; the base being towards the tongue and the apex free. The anterior are very small, and have been called *Filiform papillæ*. They appear to be produced by the terminations of the filaments of the lingual nerve, and are surrounded by an evident vascular network.

PAPILLÆ TRUNCATÆ, see Papillæ of the tongue—p. Vallatæ, see Papillæ of the tongue—p. Villosæ, see Papillæ of the tongue.

PAPILLARY, *Papillaire*, *Papillous*, (F.) *Papillaire*, from *papilla*. Of or belonging to the nipple or to the papillæ.

PAPILLARY BODY, *Corpus papillare*.

PAPILLI-AUT-MAMMILLI-FORMIS, Mastoid.

PAPILLIFORM, *Papilliformis*, from *papilla*, and *forma*, 'form.' Having the shape or appearance of a papilla.

PAPILLOMA. A tumour, whose specific elements are those of the papillæ of the normal skin; and which consequently is composed of a stem of fibrous or homogenous connective tissue; is club-shaped, cylindrical or acuminated, and covered with a layer of epithelial cells, similar to those of the normal papillæ.—Förster.

PAPILLOUS, Papillary.

PAPOOSE ROOT, *Caulophyllum thalictroides*.

PAPPE, Mamma.

PAPPEA CAPENSIS. A small tree of the *Nat. Ord. Sapindaceæ*, common in Cafferland, the fruit of which is called *Wild Plume*. It

furnishes a vinous beverage, and excellent vinegar. Its kernel contains an oil, which, although eatable, is somewhat cathartic, and is recommended as an external application in tinea capitis, alopecia, &c.

PAPPUS AMERICANUS, *Solanum tuberosum*.

PAP'ULA. Akin to *papilla*. A pimple, *Exor'mia*, (Prov.) *Pouke*, *Puckle*, (Sc.) *Pluke*, *Plouk*, (F.) *Papule*, *Bouton*, *Bourgeon*. A small, acuminate elevation of the cuticle, with an inflamed base; very seldom containing a fluid, or suppurating, and commonly terminating in scurf or desquamation. An eruption of pimples is called, in Devonshire, *Murples*.

Papule form the first order of Willan and Bateman's arrangement of cutaneous diseases.

PAPULA AGRIA, *Lichen agrius* — p. *Fera*, *Herpes exedens* — p. *Miliaris*, *Miliary fever* — p. *Typhosa*, *Spots*, red.

PAPULE, *Lichen* — p. *Siccæ*, *Lichen* — p. *Sudoris*, *Miliary fever* — p. *Sudorales*, *Sudamina*.

PAPULAR, *Papulose*.

PAPULEUX, *Papulose*.

PAP'ULOSE, *Papulo'sus*, *Pap'ulous*, *Papula'tus*, *Pimpled*, *Pap'ular*, *Papula'ria*, (F.) *Papuleux*, from *papula*. Of or belonging to, resembling, or covered with, papule or pimples.

PAQUERETTE VIVACE, *Bellis*.

PAR NERVORUM, *Syz'gyi nervo'rum*. A pair of nerves.

PAR NOKUM, *Hypoglossus* — p. *Opticum Nervorum*, *Optic nerve* — p. *Primum nervorum cerebri*, *Olfactory nerve* — p. *Quartum*, *Pathetic nerve* — p. *Quintum nervorum cerebri*, *Trigeminum* — p. *Secundum nervorum*, *Optic nerve* — p. *Septimum sive faciale*, *Facial nerve* — p. *Tertium*, *Motore oculorum* — p. *Trigeminum*, *Trigeminum* — p. *Trium funiculorum*, *Trigeminum* — p. *Vagus*, *Pneumogastrie* — p. *Visorius*, *Optic nerve*.

PARA, *para*, a common prefix, 'by, near, contrary to, through,' (*per*;) 'above and beyond,' (*ultra*;) 'besides,' (*præ*.) In certain terms, as in *Paracisus*, *Parapsis*, *Parabysma*, &c., used by Dr. Good, *para* means a defective or morbid condition.

PARA NUT, see *Pichurim beans*.

PARABAL'ANUS, from *παρὰβαλλειν*, (*para*, and *βαλλειν*, 'to cast,') 'to expose one-self.' A name given, by the ancients, to a person who devoted himself to the service of the sick in hospitals; and particularly during epidemics and the plague.

PARABYS'MA, from *para*, and *βυσιν*, 'to stuff.' Infarction or engorgement of an organ — as *Parabysma Lie'nis*, *Engorgement of the spleen*.

PARABYSMA MENTERICUM, *Tabes mesenterica*.

PARABYS'TIA, same etymon. Engorgement or infarction of an organ with blood.

PARACELSIS'TA, *Paracel'sist*, *Paracel'sian*; from *Paracelsus*. A follower of *Paracelsus*.

PARACELSISTICA MEDICINA, see *Spagyrists*.

PARACENTERIUM. Same etymon as the next. Woolhouse has given this appellation to a small trocar, employed by Nuck for puncturing the eye in cases of dropsy of that organ.

PARACENTE'SIS, *Perfora'tio*, *Punct'io*, *Punctu'ra*, *Cenembate'sis*, *Cente'sis*, *Compunct'io*, (F.) *Paracentèsis*, from *παράκινναι*, (*para*, and *κινειν*, 'I pierce,') 'I pierce through.' Puncture. The operation of tapping to evacuate the collected fluid in ascites, ovarian dropsy, &c.

PARACENTESIS, *Puncturing* — p. *Thoracis*, *Thoracentesis*.

PARACENTERION, *Trocar*.

PARACENTICION, *Trocar*.

PARACEPH'ALUS, (F.) *Paracéphale*, from

para, 'defective,' and *κεφαλή*, 'head.' A genus of monsters, with the head badly formed, but voluminous; the face distinct, with a mouth; rudimentary sensorial organs, and thoracic members.

PARACHRE'A, from *para*, and *χρῶα*, 'colour.' Morbid change of colour of the surface, especially of the face.

PARACHROMA, *Achromatopsia*.

PARACHROSIS, *Decoloration*.

PARACINE'SES, (G.) *Parakinesien*, from *para*, and *κίνησις*, 'motion.' Diseases of the motor nerves; morbid movements of voluntary muscles.

PARACMASIS, *Decline*.

PARACMAS'TIC, *Paracmas'tions*, from *παράμαζω*, (*para*, and *ακμή*, 'the highest point,') 'I decrease.' Decreasing. An epithet for a distemper, which gradually decreases. Also, according to Galen, the period of life when a person is said to grow old, which he places at from 35 to 49, after which he is said to be old — a definition that would not suit us at the present day. — Gortæus, and Castelli.

PARACMASTICOS, see *Acmaستicos*.

PARACME, *Decline*.

PARACOE, see *Cophosis*, *Paracisus*.

PARAC'OPÉ, *Paraph'ora*, *Paracru'sis*, from *παράκοπτα*, (*para*, and *κοπτα*, 'I strike,') 'I strike falsely.' Delirium. In Hippocrates, it means the slight delirium which sometimes occurs in febrile diseases.

PARACOUSIA, see *Cophosis*.

PARAC'RISES, (G.) *Parakrisien*: from *para*, and *κρίσις*, 'secretion.' Diseases of the secretory function. — Fuchs.

PARACRISIS, *Paracope*.

PARACUSIS, *Cophosis*. Also, *perverted audition*.

PARACU'SIS ACRIIS, *Audi'tus a'crior*, *Paracu'sis oxyco'i'a*, from *para*, and *ακουσις*, 'the act of hearing.' The hearing painfully acute, and intolerant of the lowest sounds; a symptom in otalgia, epilepsy, cephalitis, and febrile affections.

PARACUSIS DUPLICATA, *Double hearing* — p. *Imaginaria*, *Tinnitus aurium* — p. *Imperfecta*, *Double hearing*, *P. perversa* — p. *Obtusa*, *Barycoia* — p. *Oxycoia*, *P. acris*.

PARACUSIS PERVER'SA, *P. Willisiana seu imperfecta*, *Perverse hearing*. The ear only sensible to articulate sounds, when excited by other and louder sounds intermixed with them. — Good.

PARACUSIS SURDITAS, see *Cophosis* — p. *Willisiana*, *Paracisus perversa*.

PARACYCLE'SES, (G.) *Parakyklesen*, from *para*, and *κυκλῆσις*, (*κυκλος*, 'a circle,') 'circulation. Diseases of the blood as regards its distribution and motion. — Fuchs.

PARACYESIS, *Pregnancy*, *morbid*, *Pregnancy*, *preternatural* — p. *Abortus*, *Abortion* — p. *Tubaria*, *Salpingocystis*.

PARACYNAN'CHÉ, (*para*, and *cynanche*.) Inflammation of the external muscles of the larynx. According to others, a slight *cynanche*.

PARÆSTHESIA, *Paresthesia* — p. *Olfactoria*, *Parosmia*.

PARÆSTHESIS, *Paresthesia* — p. *Auditus flacca*, *Proptoma auricularum*.

PARÆTHENAR, *Parathenar*.

PARAGEUSIS, *Ageusia*. Also, *perversion of taste*, *Parageu'sia*, *Paragenu'sia*, *Allotriogenu'sia*, *Gustus deprava'tus*, from *para*, and *γευσis*, 'taste.'

PARAGEUSTIA, *Parageusis*.

PARAGLOSSA, *Paraglos'sis*, from *para*, and *γλῶσσα*, 'the tongue;' *Swelling of the tongue*; *Dyspha'gia Linguo'sa*, *Glossomegi'stus*, *Macroglos'sa*, *Prolap'sus lingue*. Sauvages, under the term *Paraglossa deglutito'ria*, means a pretended

doubling of the tongue into the pharynx, (F.) *Renversement de la langue.* See Glossocoele.

PARAGOGE, Abduction, Coaptation.

PARAGOMPHOSIS, *Paragomphosis cap'itis*, from *para*, and *γομφωσις*, 'a nailing.' Incomplete wedging of the head of the child in the pelvis during labour.

PARAKINESIEN, Paracineses.

PARAKRISIEN, Paracrisis.

PARAKYLESEN, Paracycleses.

PARALALIA, Mogilalia — p. Nasalis, Rhinophonia.

PARALAMP'SIS, *παράλαμψις*, from *παράλαμ-πω*, (*para*, and *λαμπω*, 'I shine,') 'I shine at the side of;' *Mac'ula corneæ margarita'cea*, *Perla*. A cicatrix on the transparent part of the eye.—Hippocrates.

PARALEREMA, Delirium.

PARALLAGE, Insanity.

PARALLAGMA, *Chevauchement*.

PARALLAXIS, *Chevauchement*, Insanity.

PARALLE'LA. A kind of scurfy, cutaneous eruption, affecting the palms of the hands, in parallel lines, and symptomatic of the venereal disease.—Forestus.

PARALLELISM OF DISEASES, Isopathia.

PARALOGIA, Delirium.

PARALOPH'IA, from *para*, and *λοφία*, 'the first vertebra of the back.' The lower and lateral part of the neck, near the vertebræ.—Keil.

PARALYSE, Paralytic.

PARALYSIE, Paralysis—p. *Musculaire atrophique*, *Atrophie musculaire*, progressive—p. *Saturnine*, Palsy, lead.

PARALYSIS, *Par'esis*, *Resolutio membro-rum seu nervo-rum*, *Cerus paralysis*, *Catal'ysis*, (F.) *Paralysis*, *Résolution des membres*; *Aneur'ria*, *Aneur'ria*, *Anervianie*,—Piorry; *Palsy*; from *παράλω*, (*para*, and *λω*, 'I loosen,') 'I relax.' Abolition or great diminution of the voluntary motions, *Myo'paral'ysis*, and sometimes of sensation, in one or more parts of the body.

Palsy is said to be *local*, when it affects only a few muscles. When it extends to half the body, vertically or horizontally, it is called *Hemiplegia* or *Paraplegia*. It is commonly symptomatic of disease of the brain or of the nerves themselves, and is a frequent accompaniment, precursor, or successor of apoplexy,—*Phrenoparal'ysis*. The immediate cause is, generally, pressure, either by blood effused, or by serum, or by vascular turgescence. At its onset, therefore, blood-letting may be advisable; but, afterwards, the disease becomes one of nervous debility, and requires stimulants to the paralyzed parts, or to the mucous membranes,—frictions, sinapisms, blisters, the moxa, baths, electricity, galvanism, purgatives, and the nux vomica, which last has the property of exciting movements where the will has no longer power. The disease generally admits only of palliation, and is extremely apt to recur.

PARALYSIS, Primula veris.

PARALYSIS AG'ITANS, *P. tremula*, *Syn'clonus ballist'us*, *Seclotyr'be festi'ans*, Tremor, Tremos, Shaking Palsy, Trembling Palsy. A variety of tremor, in which the muscles are in a perpetual alternation of contraction and relaxation.

PARALYSIS, BELL'S, *Bell's Palsy*, *Hist'ri'ic Paral'ysis* or *Palsy*, *Paral'ysis facia'lis*, *Proso'paral'ysis*, *Trismus diastrophè Sauvage'sii*, *Diastrophè Gale'ni*, *Hemiplegia facia'lis*, *Oris tortu'ra paralyt'ica Linnæ'i*, *Hemiprosoplegia*. The paralysis of the face induced by a lesion of the portio dura, has been called *Bell's Palsy*, owing to Sir Charles Bell having been one of the first to investigate it minutely.

PARALYSIS BERIBERIA, Beriberi—p. *Extremi-tatum*, *Acro-paralysis*—p. *Facialis*, Paralysis,

Bell's—p. *Hemiplegia*, *Hemiplegia*—p. *Hist'ri'ic*, Paralysis, *Bell's*—p. of the Insane, Dementia paralytica—p. *Intestini recti*, Proctoparalysis—p. *Intestinorum*, Enteroparalysis—p. *Linguae*, Glossolysis—p. *Medullaris*, Paraplegia—p. *Metallica*, Palsy, lead—p. *Nervi hypoglossi*, Glossoplegia—p. *Nervi vagi in parte thoracica*, Asthma—p. *Nervorum thoracis*, &c., Stethoparalysis—p. from Poisons, P. Venenata—p. *Rachialgia*, Colic, metallic—p. *Saturnine*, Palsy, lead—p. *Scurbutica*, Pellagra—p. *Spinalis*, Paraplegia—p. *Tremula*, Paralysis agitans.

PARALYSIS VENENA'TA, *Paralysis e vene'nis*, *Paralysis from poisons*. Paralysis, the result of poisonous substances—as of lead, mercury, arsenic, ergot, &c.

PARALYSIS E VENENIS, P. Venenata—p. *Ventriculi*, Gastroparalysis.

PARALYTIC, *Paralyt'icus*, *Par'alized*, *Pal'sied*, (F.) *Paralysé*, *Paralytique*. Same etymon as Paralysis. Affected with or causing paralysis.

PARALYTIC STROKE, see Stroke, paralytic.

PARALYZED, Paralytic.

PARAMEN'IA, from *para*, and *μην*, 'a month,' *Menstruatio anom'ala*, *Menses anom'ala*, *Mis-menstruation*. Difficult menstruation. Disordered menstruation.

PARAMENIA DIFFICILIS, Dysmenorrhœa—p. *Erroris*, Menstruation, vicarious—p. *Obstructionis*, see Amenorrhœa—p. *Profusa*, Menorrhagia—p. *Superflua*, Menorrhagia.

PARAMERIA, *παράμηνια*, from *para*, and *μηνος*, 'the thigh.' The internal parts of the thigh.

PARAMESOS, Annular finger, see Digitus.

PARAMO DE RUIZ, MINERAL WATERS OF. A remarkable acid spring in New Granada. It contains free sulphuric and chlorohydric acids, and has a temperature of 157° Fahr.

PARANEPHRI'TIS, from *para*, and *nephritis*, 'inflammation of the kidney.' Inflammation of the renal capsules, *Inflammatio renum succenturiato-rum*.

PARANŒA, Delirium, Dementia.

PARANYMPH'US, from *para*, and *νύμφη*, 'a young spouse,' 'a bride.' A name given by the ancients to him who led the bride to the house of her husband. The term was adopted into the ancient schools of medicine to designate the usual discourse pronounced at the end of the prescribed period of study for licentiate, in which the qualifications and merits of the candidate were described.

PARAPATHIA, Pathomania.

PARAPECHYON, Radius.

PARAPHIA, Parapsis.

PARAPHIMOSIS, *Periphimo'sis*, *Circumligatu'ra*, *Phimo'sis circumligu'ta*, *Penis lipoderm'us*, from *para*, and *φίμω*, 'I bridle.' Strangulation of the glans penis, owing to the opening of the prepuce being too narrow. It frequently happens in persons affected with phimosis, when the prepuce has been pushed violently behind the base of the glans, during copulation, &c. In paraphimosis, the narrow and resisting aperture of the prepuce forms, behind the corona glandis, a kind of ligature, which strangulates the part, causes it to swell and inflame, and even to become gangrenous. Attempts must be made to bring back the prepuce. These excite considerable pain, yet they must be persevered in for some time; and should they not succeed, the rings, formed by the tumefied prepuce, must be cut transversely, and the antiphlogistic treatment be adopted.

PARAPHOBIA, Hydrophobia.

PARAPHON'IA, from *para*, and *φωνη*, 'the voice.' Affection of the voice, when its character or timbre is less agreeable.

PARAPHONIA CLANGENS, Oxyphonia — p. Gutturalis, Asaphia — p. Nasalis, Rhinophonia — p. Puberum, Hirquitalitas — p. Rauca, Raucedo — p. Ulerosa, Oxyphonia.

PARAPHORIA, Delirium, Paracope.

PARAPHIRENESIS, Paraphrenitis — p. Diaphragmatica, Diaphragmitis.

PARAPHRENË, Insanity.

PARAPHRENITIS, from *para*, and *φρενες*, 'the diaphragm,' *Paraphrenesis*, (F.) *Paraphrenésis*. A name given to inflammation of the diaphragm, (see Diaphragmitis,) which was supposed always to produce delirium:—as well as to the delirium itself accompanying that affection. See Phrenitis.

PARAPHROSYNE, Delirium.

PARAPHROTES, Delirium.

PARAPLASTIC, *Paraplasticus*, from *para*, and *πλαστικός*, 'formative.' Possessed of depraved formative powers,—as Carcinoma, Encephaloma, Melanosis, &c.

PARAPLEGIA, *Paraplexia*, *Rachio-paraplegia*, *Rhachio-paraplegia*, *Rhachio-plegia*, *Myelo-paraplegia*, *Paralysis spinalis seu Medullaris*, *Cervicis paralysis paraplegia*, (F.) *Paraplégie*, from *para*, and *πλησσω*, 'I strike.' Palsy of the lower half of the body, including the bladder and rectum. It is, generally, owing to disease of some part of the cerebro-spinal axis; especially of the spinal portion. The ancients sometimes used the term for incomplete apoplexy. Some authors have, also, made a distinction between *paraplegia* and *paraplexia*,—using the former for partial apoplexy,—the latter for general or complete apoplexy.

PARAPLEURITIS, Pleurodynia.

PARAPLEXIA, Paraplegia.

PARAPOPHYSES, see Vertebra.

PARAPOPLEXY, *Parapoplexia*, from *para*, and *ἀποπληξία*, 'apoplexy.' False apoplexy; a soporose state, resembling apoplexy; and especially that which occurs in the paroxysms of pernicious intermittents.

PARAPSIS, *Paraphia*, from *para*, and *ἅπτω*, 'I touch.' Sense of touch or general feeling vitiated or lost.—Good.

PARAPSIS ACRI TENERTUDO, Soreness — p. Experts, Torpor, Anæsthesia — p. Illusoria, Pseudæsthesia — p. Pruritus, Itching.

PARARMA, Fimbria.

PARARRHYTHMICUS, Pararrhythmus.

PARARRHYTHMUS, *Pararrhythmicus*, from *para*, and *ῥυθμός*, 'rhythm.' An epithet for the pulse, when it has not the rhythm proper to the age and constitution of the individual.

PARATHREMA, *Pararthroma*, *Parathresia*, *Pararthrosis*, *Parac'tasis*, *Parastrepsis*, from *para*, and *ἄρθρον*, 'a joint,' Incomplete luxation; subluxation.

PARASCEPAS'TRA, from *para*, and *σκεπέω*, 'I cover.' A cap or bandage, which covers the whole head.—Galen.

PARASCEUE, Apparatus.

PARASCHIDES, from *para*, and *σείω*, 'I cleave.' Fragments or splinters of a broken bone.

PARASITE, *Parasitus*, from *παράσιτος*, (*para*, and *σιτος*, 'corn,' 'food.') 'I eat by the side of.' Parasites are plants which attach themselves to other plants; and animals which live in or on the bodies of other animals,—so as to subsist at their expense. The mistletoe is a parasitic plant;—the louse a parasitic animal. The chief parasitic animals are the *Pediculus humanus corporis*, *P. capitis*, *P. pubis*, *Pulex irritans*, *P. penicillans*, and the *Acarus Scabiei*.

PARASTIC, *Parasiticus*, *Parasitoid*, *Parasitoides*. Same etymon. Having the nature of, or resembling a parasite,—as 'parasitic growths,' tubercles, cancer, monster, (F.) *Parasitaire*, &c.

PARASITICIDE, Antiparasitic.

PARASITISM, *Parasitismus*. Same etymon as Parasite. The condition of a parasite, or of an organized body which lives on another organized body.

PARASITISMUS INTESTINALIS, Helminthiasis — p. Superficialis, Malls.

PARASITOG'ENY, *Parasitogenia*, *Parasitogenesis*. The corporeal condition, which favours the development of parasites.

PARASITOID, Parasitic.

PARASPA'DIA, from *para*, and *σπῆω*, 'I draw.' A preternatural opening of the urethra at the side of the penis. One whose urethra opens in this manner is called *Paraspidæus* and *Paraspidicus*.

PARASPADIEUS, see Paraspadia.

PARASPHAGIS, from *para*, and *σφῆγν*, 'the throat.' The part of the neck contiguous to the clavicles.

PARASTATA, *Parastates*, from *παρίστημι*, (*para*, and *ίστημι*), 'I stand near.' The prostate. In Hippocrates, it means the epididymis. The epididymes were called by Herophilus and Galen *Varico'sæ parastata*, to distinguish them from *glandulo'sæ parastata* or *prostate*. Rufus of Ephesus called the Tubæ Fallopianæ *Parastata varico'sæ*.

PARASTATA ADENOIDES, Prostate — p. Glandula, Prostate — p. Glandulosa, Prostate.

PARASTATES, Parastata.

PARASTATITIS, *Epididymitis*; from *parastata*, 'epididymis,' and *-itis*, denoting inflammation. Inflammation of the epididymis.

PARASTREMA, *Parastrephê*, from *παράστροφω*, (*para*, and *στρέφω*, 'I turn,') 'I distort or pervert.' Convulsive distortion, especially of the face, the soft parts being drawn laterally.

PARASTREPSIS, Parastrema.

PARASTROPHE, Parastrema.

PARASYNANCHE, Cynanche parotidæa.

PARASYSTOLE, Acinesia.

PARATH'ENAR, *Parathénar*, from *para*, and *ἔναρ*, 'the sole of the foot.' Winslow called *grand parathénar*, *parathénar major*, a portion of the abductor of the little toe, *musculus scandularius*; and *petit parathénar*, *parathénar minor*, the short flexor of that toe.

PARATHENAR MAJOR, Abductor minimi digiti pedis — p. Minor, Flexor brevis minimi digiti pedis.

PARATHROMA, Elongatio.

PARATONIA, see Spasm.

PARATOPIA, Luxation.

PARATOPIÆ, from *para*, and *τοπος*, 'place.' Diseases in which organs are removed from their proper position. See Ectopia.

PARATRIMMA, Chafing.

PARATROPE, Apotrope.

PARATROPHY, *Paratrophê*, *Paratrophia*, from *para*, and *τροφή*, 'nourishment. Misnutrition.

PARBREAKING, Vomiting.

PARDALIANCHES, Aconitum.

PARDALIANCHUM, Aconitum.

PARRECTAMA, *Parrectasis*, from *πᾶρα*, *εκ*, 'out of,' and *τείνω*, 'I stretch.' Immoderate extension, distension, or dilatation of a part—as of the heart.

PARRECTASIS, Parastrema, Parectama.

PAREGORIC, *Paregoricus*, *Solpiens*, (F.) *Parégorique*, from *παρηγορέω* (*para* and *αγορεύω*), 'I soothe, I calm.' That which soothes or assuages—as paregoric elixir. An anodyne.

PAREIA, Gena.

PAREIRA, see P. brava.

PAREIRA BRAVA, *Cissampelos Pareira*, *Pa-reira*, *Butua*, *Abutua*, *Albutua*, *Boton*, *Cara-pêba*, *Overo butua*, *Wild vine*, *Velvet leaf*. Family

ispermeæ. *Ser. Syst.* Diœcia Monadelphica. root of this plant — *Paraïra* (Ph. U. S.) — a tree of South America and the West Indies, a sweet taste, with considerable bitterness and slight roughness. It has been recommended in nephritic and calculous affections; but it is not much used. By A. de St. Hilaire, Pareira is assigned to *Cissampelos glaberima*.

PARELLE, *Rumex hydrolapathum*, R. patiensia.

PARENCEPHALIS, Cerebellum.

PARENCEPHALITIS, Cerebellitis.

PARENCEPHALOCÈLÈ, from *παρεγκεφαλῖς*, (*para*, and *εγκεφαλός*, 'the brain,') 'the cerebellum,' and *κῆλη*, 'a rupture.' *Hernia of the cerebellum*; a very rare disease. It occurs under the form of a soft, indolent, or slightly painful tumour, without change in the colour of the skin, and is situate in the occipital region. It is constituted of the cerebellum, which escapes through an aperture in the occipital bone. Great attention must be paid not to mistake it for an ordinary tumour, as its extirpation would be attended with fatal consequences.

PARENCEPHALUS, Cerebellum.

PARENCHYMA, *Caro parenchymat'ica seu viscerum seu viscerat'is seu viscerosa*, *Substantia parenchymat'osa*, *Prochyma*, (F.) *Parenchyme*. The texture of glandular and other organs, composed of agglomerated globules united by areolar tissue, and tearing with more or less facility. Such is the texture of the liver, kidneys, &c. *Παρεχυμα* signifies, in Greek, effusion, (*παρά, εν, and χυω*, 'I pour,') and it was believed, that this texture consisted of effused blood or other fluid.

PARENCHYMA TESTIS, *Pulpa testis*.

PARENCHYMAL, *Parenchymat'ous*, *Parenchymous*, *Parenchymat'ia*, *Parenchymat'icus*, (F.) *Parenchymateux*; same etymon. Belonging or relating to the parenchyma of an organ.

PARENCHYMATITIS: from *parenchyma*, and *itis*, denoting inflammation. *Inflammatio parenchymat'ica*. Inflammation of the substance of an organ.

PARENCHYMATOUS, *Parenchymal*.

PARESIS, from *παρεσις*, 'I relax.' (F.) *Parésie*. Paralysis. According to some, a slight, incomplete paralysis, affecting motion, not sensation.

PARESTHESIS, *Paræsthesia*, *Paræsthesia*, *Sensua imperfectus*, from *para*, and *αἰσθησις*, 'sensation.' Congenital misconstruction of the external organs of sense. A genus in the nomenclature of Good. Also, perversion of sensibility.

PARETICA EXARTHROSIS. Luxation from relaxation of the ligaments of a joint.

PARETICUS, *Relaxant*.

PARERYA, *Paraïra brava*.

PARFUM, *Suffimentum*.

PARIERA, *Paraïra brava*.

PARIES, *Table*.

PARIETAL, *Parietalis*, *Pariet'innus*, from *paries*, *parietis*, 'a wall.' Appertaining to the parietes or walls of an organ or to the parietal bones.

PARIETAL ARTERY, see P. Bones.

PARIETAL BONES, *Ossa parietalia seu verticillia seu verticilis seu sincipitis seu bregmatica seu bregmat'ica seu nervalia seu tetragona*, are two fellow-bones, occupying the lateral and upper parts of the cranium, in the formation of which they assist. They have a quadrilateral shape: their external surface is convex, and presents, at its middle part, an eminence called the *Parietal protuberance*, *Tuber parietale*, (F.) *Bosse parietale*. Their inner or cerebral surface, which is concave, has a depression that corresponds with the preceding eminence, and which is called the *Parietal fossa*, *Fossa parietalis*. Above and be-

hind, near the superior edge of these bones, is a foramen, which does not always exist, called the *Parietal foramen*, *Foramen parietale*, through which passes a vein — *parietal vein* — that opens into the superior longitudinal sinus of the dura mater: and an artery — *parietal artery* — a branch of the occipital, which ramifies on the same sinus. Each parietal bone is articulated above, with its fellow: below, with the temporal bone: before, with the frontal; behind, with the occipital, and by the anterior and inferior angle with the great ala of the sphenoid. Each parietal bone is developed from a single point of ossification.

PARIETAL FORAMEN, see P. BONES — p. Fossa, see P. BONES — p. Protuberance, see P. BONES — p. Veins, see P. BONES.

PARIETARIA, *P. officinalis*, *Helz'inz*, *Vit'aria*, *Wall pel'titory*, (F.) *Pariétaire commune*, *Perce-muraille*, *Aumure*, *Herbe aux murailles* ou *des Nones*. Family, Urticæ. *Ser. Syst.* Polygamia Monœcia. This plant grows on old walls, — hence its name, from *paries*, *parietis*, 'a wall,' — and contains much nitrate of potassa. It is diuretic, but is not much used.

PARIETARIA PENNSYLVANICA, *American Pel'titory*, a small weed, flowering from June to August, has been used in juice or decoction as a diuretic, deobstruent [?] and emmenagogue.

PARIETES, Plural of *Paries*, 'a wall,' (F.) *Parois*. A name given to parts which form the enclosures or limits of different cavities of the body, as the *Parietes* of the cranium, chest, &c.

PARIETINUS, *Parietal*.

PARIS, *P. quadrifolia*, *Solanum quadrifolium*, *Uva inrer'ca seu lupina seu vulp'ina*, *Oneberry*, *Trueloce*, (F.) *Parisette*, *Raisin de Renard*, *Herbe à Paris*, *Étrangle-loup*. It possesses narcotic properties, and was formerly considered a powerful love philter. It is emetic, and is sometimes used externally as a resolvent and anodyne.

PARISAGOGE INTESTINORUM, *Intussusceptio*.

PARISETTE, *Paris*.

PARISTH'MIA, *para*, and *ισθμος*, 'the throat.' A word employed to designate the tonsils, and, likewise, inflammation of the tonsils. *Cynanche tonsillaris*, and *Cynanche* in general.

PARISTHMIOTOMUS, from *paristh'mia*, 'the tonsils,' and *τεμνω*, 'I cut.' An instrument with which the tonsils were formerly scarified.

PARISTHMITIS, *Cynanche*, C. tonsillaris.

PARISTHMIUM, *Tonsil*.

PARISWORT, *Trillium latifolium*.

PARMACITY, *Cetaceum* — p. Poor Man's, *Thlaspi bursa*.

PARME'LIA PARIET'INA, *Lichen pariet'innus*, *Yellow wall Lichen*, *Common Yellow Wall Moss*, which grows abundantly on trees and walls. Order, *Lichenacæ*, is used in intermittent fever on account of its bitterness.

PARMELIA ISLANDICA, *Lichen Islandicus* — p. *Plicata*. *Lichen plicatus* — p. *Pulmonacea*, *Lichen pulmonarius* — p. *Roccella*, *Lichen roc-cella* — p. *Saxatilis*, *Lichen saxatilis*.

PARMENTIE'RA CERIFERA, *Pala de velas*, *Candle Tree*. A forest tree in the valley of the Chagres, — *Nat. Ord.* *Crucifera*, — the long cylindrical fruit of which, resembling candles, serves for food for cattle.

PAROARION, *Parovarium*.

PAROCHETEUSIS, *Derivation*.

PARODONTIDES, from *para*, and *οδους*, tooth. 'Tumours on the gums: parulis, &c.'

PARODYNIA, *Dystocia* — p. *Perversa*, *Pre-sentation*, *preternatural*.

PAROIS, *Parietes*.

PAROLE, *Voice*, articulated.

PAROMPHALOCÈLÈ, from *para*, and *μφαλός*,

'the navel,' and κηλη, 'rupture.' Hernia near the navel.

PARONIRIA, from *para*, and ονειρον, 'a dream.' Depraved, disturbed, or morbid dreaming; *Oncirodynia activa*, *Erethismus onciro-dynia*. The voluntary organs connected with the passing train of ideas, overpowered by the force of the imagination during dreaming, and involuntarily excited to their natural or accus-tomary actions, while the other organs remain asleep.—Good.

PARONIRIA AMBULANS, Somnambulism—p. Sa-lax, Pollution.

PARONYCHIA, *Paronychia*, *Paronychitis*, *Pan'aris*, *Redu'via*, *Redu'ia*, *Vermis* seu *Pteryg-i'um dig'iti*, *Panari'tium*, *Digit'ium*, *Pandali'tium*, *Passa*, *Dactylitis*, *Pana'rium*, *Onych'ia*, (F.) *Onglade*, from *para*, and ονυξ, ονυχος, 'the nail.' The ancients gave this name to an inflam-matory tumour.—*Onychitis*, *Onyxi'tis* (Gross),—seated near the nail, and involving the pulp or matrix. Some of the moderns have extended it to inflammation of the hand, and even of the forearm. It usually means every phlegmonous tumour of the fingers or toes, and especially of the first phalanx. Four kinds of paronychia are commonly pointed out: 1. That seated between the epidermis and skin, called in France *Tour-niole*. 2. That seated in the subcutaneous arcu-lar tissue. 3. That occupying the sheath of a tendon; and 4. That considered to be seated be-tween the periosteum and bone. The last three are only different degrees of the same disease, constituting *Whitlow*, (Prov.) *Teak*, (F.) *Mal d'aventure*:—the worst kinds, in English, are called *Felon*. The inflammation generally com-mences in the subcutaneous areolar tissue, and spreads to other parts, occasioning excessive, lancinating pain, owing to the parts being put upon the stretch; hence, the necessity of dividing them freely,—even down to the bone, when there is reason to believe that the affection belongs to the fourth division,—making use of emollient applications, the antiphlogistic treatment, and opiates to relieve pain.

PARONYCHIA, *Asplenium ruta muraria*—p. Di-gitum, Digitum.

PARONYCHIS, *Paronychia*.

PARONYCHITIS, *Paronychia*.

PAROPHOBIA, *Hydrophobia*.

PAROPIA, *Canthus* (lesser.)

PAROPLEXIA, *Paraplegia*.

PAROPSIS, *Dysæsthesia visualis*—p. Amau-rosis, Amaurosis —p. Cataracta, Cataract —p. Glauco-sis, Glaucoma —p. Illusoria, Pseudoblep-sia —p. Lateralis, Dysopia lateralis —p. Longin-qua, Presbytia —p. Lucifuga, Nyctalopia —p. Noctifuga, Hemeralopia —p. Propinqua, Myopia —p. Staphyloma, Staphyloma —p. Staphyloma purulentum, Hypopyon—p. Staphyloma-simplex, Hydrophthalmia —p. Strabismus, Strabismus —p. Synæsis, Synæsisis.

PAROPTESIS, from *para*, dim., and οπταω, 'I roast.' Forcing a sweat by placing a patient before the fire or putting him in a dry bath.

PARORASIS, from *para*, and ορω, 'I see.' Perversion of sight, which prevents the person from judging accurately of the colour of objects. *Achromatopsia*. Hallucination.—Galen.

PARORCHIDIUM, from *para*, 'something wrong,' and ορχης, ορχιδος, 'testicle.' Any position of one or both testicles, different from that which they ought to occupy in the scrotum;—whether they may not have descended through the abdo-minal ring, or are drawn up to the groins by the cremaster muscle.

PAROR'CHIDO-ENTEROCELĒ, *Hernia paror'chido-enter'ica*. Internal hernia compli-cated with displacement of the testicle.—Sau-vages.

PAROSMIA, Anosmia. Also, perversion of smell; *Paræsthesia olfactoria*, *Olfac'tus depræ-vatus*, *Idiosyncrasia olfactoria*, from *para*, and οσμη, 'odour.'

PAROSTIA, from *para*, 'defectively,' and οστειον, 'a bone.' Bones untempered in their sub-stance, and incapable of affording proper sup-port.—Good.

PAROSTIA FLEXILIS, Mollities ossium—p. Fra-gilis, Fragilitas ossium.

PAROTIA, *Cynanche parotidæa*.

PAROTID, *Gland'ula parotide'a* seu *ad Anres*, *Parot'is*, *Animal'as*, *Alimel'as*, *Lacticin'ia*; from *para*, and ος, ως, 'the ear.' The largest of the salivary glands, seated under the ear, and near the angle of the lower jaw. It is composed of many separate lobes, giving rise to excretory ducts, which unite to form one canal, called the *Parotid duct*, *Steno's canal*,—the *Ductus superior* or *Superior salivary canal*, of some. This duct, after having advanced horizontally into the sub-stance of the cheek, proceeds through an opening in the buccinator muscle, and terminates in the mouth opposite the second upper molaris. About the middle of its course, it sometimes receives the excretory duct of a glandular body, situate in its vicinity, and called the *Accessory Gland of the Parotid*, *So'cia Parot'idis*, *Gland'ula so'cia parot'idis*. In the substance of the parotid are found—a number of branches of the facial nerve, of the transverse arteries of the face, and the posterior auricular. It receives, also, some fila-ments from the inferior maxillary nerve, and from the ascending branches of the superficial cervical plexus. Its lymphatic vessels are some-what numerous, and pass into ganglions situate at its surface or behind the angle of the jaw. The parotid secretes saliva, and pours it copiously into the mouth.

PAROTID, ACCESSORY GLAND OF THE, see Pa-rotid.

PAROTID APONEURO'SIS. A sheath of great thickness, continuous, below, with the cervical fascia. It forms a frame-work to the parotid by means of fibrous prolongations from its deep sur-face. Its density accounts for the pain of paroti-tis, and for the difficulty with which pus makes its way to the surface.

PAROTID ARTERIES, see Parotid —p. Duct, see Parotid —p. Veins, see Parotid.

PAROTIDE, Parotis.

PAROTIDE'AN, *Parotidæ'us*, (F.) *Parotidien*. Belonging or relating to the parotid.

PAROTIDITIS, *Cynanche parotidæa*, *Parotis*.

PAROTIDON'CUS, *Tumor parot'idis*, *T. Gland'ula parotide'a*, from *parotis*, and ογκος, 'a tu-mour.' Swelling of the parotid.

PAROTIS, *Phleg'monè parotide'a*, *Inflamma'tio parot'idum*, *Cynan'che parotidæ'a*, *Parotit'is*, *Parotidit'is*, *Dioc'en'ri*, (F.) *Parotide*, is the name of a tumour seated under the ear; which is red-dish, hard, and attended with obtuse pain; the progress to suppuration being slow and difficult. Sometimes it is of a malignant character, slough-ing, and long protracted. Some French patholo-gists restrict the term *parotide* to symptomatic parotitis—as to that which occurs in plague; whilst they apply the terms *Oreillons* or *Ourles*, *Parotoneices*, to *Cynanche parotidæa* or Mumps.

PAROTIS CONTAGIOSA, *Cynanche parotidæa* —p. Epidemica, *Cynanche parotidæa* —p. Seroæ

glutine tumens, Cynanche parotidæa—p. Spuria, Cynanche parotidæa.

PAROTITIS, Cynanche parotidæa, Parotis—p. Epidemica, Cynanche parotidæa—p. Erysipelatosa, Cynanche parotidæa—p. Polymorpha, Cynanche parotidæa—p. Specifica, Cynanche parotidæa.

PAROTONCIES, see Parotis.

PAROVARIUM, properly *Parovarium*; *Corpusculum continent Rosenauilleri*, from *para*, and *ovum*, 'the ovary.' A body closely analogous in structure to the epididymis, which is seated between the Fallopian tube and the ovary in the broad ligament. The Corpus Wolffianum concurs in its formation.—Kobelt.

PAROXYSMATIC, *Paroxysmaticus*, (F.) *Paroxysmatique*, from *παροξυσμός*, 'I excite.' Relating to a paroxysm.

Jours paroxysmatiques are days on which paroxysms take place.

PAROXYSM, *Paroxysmus*, from *παροξυσμός*, (*παρά*, and *οξύω*, 'to render sharp.') 'I irritate.' *Insultus*, *Insultus*, *Insultus*, *Exacerbatio*, *Accessio*, *Accessus*, (Old Eng.) *Axes*, (F.) *Exacerbation*, *Redoublement*, *Access*. The augmentation which supervenes at equal or unequal intervals in the symptoms of acute diseases. Also, a periodical exacerbation or fit of a disease.

PAROXYSMAL, *Paroxysmalis*; same etymon. (F.) *Paroxysmatique*. Of or belonging to a paroxysm:—as a *paroxysmal* day, (F.) *Jour paroxysmatique*; a day on which a paroxysm of disease occurs.

PAROXYSMUS FEBRILIS, Pyrexia.

PAROXYSTIQUE, Paroxysmal.

PARS, Genital organs—p. Corporis, Genital organs—p. Inferior Pelvis, Sole—p. Intermedia, see Corpus spongiosum urethræ—p. Mastoidea of Temporal Bone, Mastoid process—p. Naturalis medicine, Physiology—p. Obscena, Genital organs—p. Petrosa, see Temporal Bone—p. Prima asperæ arteriæ, Larynx—p. Pyramidalis ossis temporis, see Temporal Bone—p. Squamosa, see Temporal Bone—p. Urethræ membranacea, see Urethra—p. Vaga, Pneumogastric—p. Virilis, Penis.

PARSLEY, COMMON, *Apium petroselinum*—p. Fool's, *Æthusa cynapium*—p. Macedonian, *Bubani macedonicum*—p. Mountain, black, *Athamanta aureoselinum*—p. Poison, *Conium maculatum*—p. Spotted, *Conium maculatum*—p. Water, *Cicuta maculata*.

PARSNIP, COW, *Heracleum spondylium*, *Heracleum lanatum*—p. Crow, *Leontodon taraxacum*—p. Garden, *Pastinaca sativa*—p. Meadow, *Thaspium Barbinode*—p. Water, common, *Sium latifolium*.

PART, see Parturition—p. *Légitime et illégitime*, see Parturition.

PARTES GENERATIONI INSERVIENTES, Genital organs—p. Genitales, Genital organs—p. Laterales nasi, Scapulae nasi—p. Nervose, see Nerve—p. Obscena, Genital organs.

PARTHENEIA, Virginity.

PARTHENIUM FEBRIFUGUM, *Matricaria*—p. Mas, *Tanacetum*.

PARTHENIUS (*Morbus*): from *παρθένος*, 'a virgin.' A disease in a young female. Also, chlorosis.

PARTHENOGENESIS, 'virgin birth': from *παρθένος*, 'a virgin.' and *γενεσις*, 'generation.' The successive production of procreating individuals from a single ovum, without any renewal of fertilization.—Owen.

PARTHENOLOGY, *Parthenologia*, from

παρθένος, 'virginity,' and *λογος*, 'a description.' A description or consideration of the state of virginity in health or disease.

PARTHENUS, Virgin.

PARTIALIS, Local.

PARTICLES, PRIMITIVE, OF MUSCLES, Sarcous Elements.

PARTICULÆ SANGUINIS, Globules of the blood.

PARTIES, LES, Genital organs—p. *Génitales*, Genital organs—p. *Montenses*, Genital organs—p. *Génitoires*, Genital organs—p. *Naturæ*, Genital organs—p. *Nobles*, Genital organs—p. *Sexuales*, Genital organs.

PARTIO, Parturition.

PARTIDGE-BERRY, *Gaultheria*, *Mitchella repens*.

PARTS, THE, Genital organs.

PARTURA, Parturition.

PARTURIENT, *Parturiens*, from *parturire*, *parturitus*, itself from *parere*, 'to bring forth.' Bringing forth, or about to bring forth, or having recently brought forth young.

The *Parturient*, *Puerperous*, *Childbed* or *Lying-in* or *Lying-down* state, *Confinement*, *Puerperium*, generally comprises the period from parturition to perfect recovery, which is usually a month. In the north of England this is called the *steg-month*, requires careful management, and is subject to various diseases.

A *Parturient* or *Parturificient* is a medicine which induces or promotes labour; as ergot, and, perhaps borate of soda. See Abortive.

PARTURIFACIENT, Parturient.

PARTURIGO, Parturition.

PARTURITION, *Puerperium*, *Partus*, *Partio*, *Parturitio*, *Parturi'go*, *Partura*, *Tocos*, *Tolctos*, *Tetis*, *Lochi*, *Lochi'sis*, *Mæcul'sis*, *Maient'sis*, *Negotium* seu *Actus parturitio'nis*, *Enic'sis* seu *Enit'io* seu *Expul'sio factus*, *Apoll'ysis*, *Status parturien'di*, *Delir'ery*, *Labour*, *Child-bearing*, *Child-birth*, *Birth*, *Travail*, (Sc.) *Child-ill*, *Crying*, *Crying out*, (F.) *Accouchement*, *Couche*, *Travail d'Enfant*, *Enfement*. Same etymon as Parturient. The act of delivery of the fœtus and its appendages, also the state during and immediately after delivery—*Childbed*, (Sc.) *Gizzen-bed*, *Gizzen*, see Parturient. Labour is the necessary consequence of conception, pregnancy, and the completion of gestation. It does not take place, at the full time, until after nine months of utero-gestation,—the causes producing it being the contraction of the uterus and abdominal muscles. By different authors, the stages of labour have been variously divided. We may, perhaps, admit of four. The *first stage* comprises the precursory signs. One, two, or three days before labour, a mucous discharge, streaked with blood, takes place from the vagina, which is called the *Signum*, *Leucorrhœa Nabothi*, *Hæmorrhag'ia Nabothi*, *Labour-show*, or *Show*; and, along with this, trifling grinding pains, called by the French *Mouches*, are felt in the loins and abdomen. *Second stage*:—Peculiar pains, which extend from the lumbar region towards the uterus; tension and dilatation of the neck and orifice of that organ; protrusion into the vagina of the envelopes of the fœtus, forming the *bag of waters*, (F.) *Poche des eaux*; and rupture or breaking of the waters. *Third stage*:—Contractions of the uterus and abdominal muscles; forcing pains; the head of the fœtus becomes engaged in the pelvis: the occiput being commonly situate above the left acetabulum, passes beneath the arch of the pubis. *Fourth stage*:—the head now presents at the vulva, the perineum being

considerably distended; at length it clears the external parts, and the rest of the body is easily disengaged. The child still continues attached to the mother by the umbilical cord, which has to be divided. In the course of fifteen or twenty minutes, trifling pains occur, and the secundines are expelled. This completes delivery—the *birth*, (F.) *Naissance* of the child. In by far the majority of cases, the head is the presenting part, the occiput directed anteriorly, and the face posteriorly. Of 12,633 children, born at the Hospital *La Maternité*, in Paris, from the 10th of December, 1797, to the 31st of July, 1806, the following were the presentations:

The head, with the face posteriorly...	12,120
The head anteriorly.....	63
The breech.....	193
The feet.....	147
The knees.....	3
Other positions.....	102
	12,633

The following table, drawn up from data furnished by Velpéau, gives an approximate view of the comparative number of presentations, according to the experience of the individuals mentioned:

TABLE, EXHIBITING THE RATIO OF PRESENTATIONS IN 1000 CASES.

	ACCORDING TO							
	Merri-man.	Bland.	Madame Boivin.	Madame Lachapelle.	Nägele.	Lovati.	Hospital of the Faculté.	Boer.
Regular or of the vertex.....	924	944	969	933	933	911	980	
I. Occipito-anterior.....	908		944	910		895		
a. Occipito-cotyloid (left).....			760	717		537		
Do. (right).....			179	209				
b. Occipito pubian.....			0.29					
II. Occipito-posterior.....			9.4	0				
a. Fronto-cotyloid (left).....			5.3	7.3				
Do. (right).....			4.4	2.9				
Face presentation.....	2.2	2.6	3.6	4.6				8.8
Mento-iliac (right).....			2.6					
Of the pelvis.....	36	28	29	36	47			29
Of the foot.....	12.7	9.4		14				10.3
Of the knees.....			0.19	0.40				
Of the breech.....	23	13	18	22				19
Of the trunk.....			4.6	5.3	4.8			
Requiring Forceps.....	6.6	4.7	4.6	3.4	36			5.7
“ Turning.....	16	4.7		7.8	7.2			5.9
“ Cephalotomy.....	3.3	5.2	4.77	0.53	2.4			1.5

Labour has received different names, according to the period of pregnancy at which it occurs, and the mode in which it takes place. It is called *miscarriage* or *abortion*, prior to the seventh month; *premature labour*, *Omotoc'ia*, *Partus præcox* seu *præmaturus*, between this and eight months and a half; and *labour at the full period*, (F.) *Accouchement à terme*, when it happens at the end of nine months. Labours have likewise been variously divided by different authors. Perhaps the one into *Natural*, *Preternatural*, and *Labourious*, is as common as any;—*natural Labour* meaning that which would occur without manual assistance; *preternatural*, requiring the assistance of art, but by the hand solely: and *labourious* or *instrumental*, requiring instrumental assistance; which M. Capuron calls *Accouchement Mécanique*. *Powerless labour*, the *Inertie par*

épuisement of Madame Lachapelle, is that in which the muscular powers from exhaustion are unable to accomplish such delivery.

Forced labour or *parturition*, (F.) *Accouchement ou Avortement provoqué*, is labour or abortion, induced by art. When labour is accompanied with other phenomena of a deranging character, it is said to be *complicated*. When the child, even when reduced in size, cannot pass, it is termed *impracticable*, &c. See *Dystocia*.

The French use the term *Part*, *Partus*, at times, synonymously with *delivery*; at others, with *fœtus*, *Matœu'ma*, *Mœu'ma*. *Suppression de part* means concealment of delivery; *Part légitime*, a fœtus which comes into the world with the condition necessary for its enjoying advantages granted by law; and *Part illégitime*, one that does not fall under those conditions.

TABLE OF THE COURSE AND MANAGEMENT OF DIFFERENT KINDS OF LABOURS.

I. NATURAL LABOURS.	
Or those in which Delivery can be effected without Assistance.	
HEAD PRESENTING. (Four Positions.)	1. Occiput towards the Left Acetabulum. { The head with the chin bent upon the breast clears the brim of the pelvis. Motion of rotation produced by the inclined planes of the pelvis; the occipital protuberance is carried behind the symphysis pubis, and the face towards the hollow of the sacrum.
	2. Occiput towards the Right Acetabulum. { The head attains the outlet: clears it; the occiput being first disengaged, and the face afterwards, which passes successively along the whole of the hollow of the sacrum. The shoulders enter the brim diagonally, and execute the movement of rotation. The one behind first clears the outlet. The same thing occurs with the nates.
	3. Occiput towards the Right Sacro-iliac Symphysis. { The same course is here followed: except that after the rotary movement, the face passes behind the symphysis pubis, and the occiput into the hollow of the sacrum. The occiput is first extruded;—the face next.
	4. Occiput towards the Left Sacro-iliac Symphysis. { When the chin presents and is turned forwards, the face is first expelled; the head is thrown backwards. This presentation is uncommon. If the chin present and be thrown backwards, natural delivery cannot be effected.

NATURAL LABOURS.—CONTINUED.

LOWER EXTREMITY PRESENTING.
(Four Positions of the Feet, Knees, and Breech.)

1. *Heel, Tibia, or Sacrum towards the left Acetabulum.*
2. *Heel, Tibia, or Sacrum towards the Right Acetabulum.*
3. *Heel, Tibia, or Sacrum towards the Right Sacro-iliac Symphysis.*
4. *Heel, Tibia, or Sacrum towards the Left Sacro-iliac Symphysis.*

The nates enter the brim diagonally: movement of rotation: the one passes behind the symphysis pubis; the other towards the hollow of the sacrum. The one behind is first extruded.

The same thing happens to the shoulders. The chin is bent upon the chest to clear the brim: movement of rotation: the occiput passes behind the symphysis pubis, and the face towards the hollow of the sacrum. The face clears the outlet first; the occiput last.

The same progress, except that after the movement of rotation, the face is towards the pubis, and the occiput towards the sacrum. The face is here again delivered first.

II. PRETERNATURAL LABOURS,

Or those which require Manual Assistance only.

I. PRESENTATION OF THE HEAD, UPPER EXTREMITIES, BACK, ABDOMEN, &c.

1. When the labour is without any serious complication, and the head is not in a good position at the brim of the pelvis, endeavours are to be made to place it in its natural position: the hand to be introduced with the fingers united, and the long diameter of the head to be placed diagonally in the brim, if possible. Or the lever may be fixed over the occiput, and, by drawing it down, a closer approximation of the chin and chest may be effected; the position of the head to the pelvis being at the same time attended to.
2. If complicated with syncope, convulsions, hemorrhage, want of uterine action, &c., the feet must be laid hold of, and the child *turned*. For this purpose, empty the rectum and bladder; and turn, if possible, prior to the escape of the liquor amnii. The female may be placed on her back or left side, with the breech over the edge of the bed. Use the right or left hand, according as the feet of the child are to the right or left side of the pelvis; always so introducing the hand, that the child shall be in its palm, and the back of the hand opposed to the inner surface of the uterus. The hand and arm to be introduced during an interval of pain; and they must be flattened and tranquil whilst the uterus is contracting forcibly. If both feet cannot be readily brought down, seize one, and tie a fillet around it.
3. If the toes point to the vertebral column, or to the abdomen of the mother, the head, in its descent, will not enter the pelvis: the chin and occiput will be hitched on the pubis and promontory of the sacrum. In such case, grasp the nates as soon as they have passed the os externum; and, during a pain, endeavour to direct the toes towards either sacro-iliac synchondrosis. When the fingers can be passed along the body of the child, and over the shoulders to the bend of the elbows, draw down the arms successively. When the body is expelled, and the head filling up the brim, the termination of labour must be accelerated by passing two fingers over the shoulder of the child; using moderate and steady extracting power, whilst one finger of the other hand, passed into the mouth, depresses the chin on the sternum. In this way, the head may be extracted.

III. INSTRUMENTAL LABOURS,

Or those requiring the Assistance of appropriate Instruments.

GENERAL OBSERVATIONS.

If the pains become feeble, or inadequate from any cause;—or if the presenting part be firmly wedged in the pelvis; or the woman be becoming exhausted, instrumental assistance may be required. In all such cases, place the female on the back or left side, empty the bladder and rectum; wait till the os uteri is in a fit state; afford assistance during the pains; introduce the instrument during the intervals between the pains; have the patient on her left side (generally); and extract according to the axes of brim and outlet, as the case may be.

1. *Short Forceps*

Apply these to the sides of the head of the child, so that the ears and parietal protuberances may be within the fenestræ. Introduce the left-hand blade first; the index and middle fingers of the right hand guiding it to the ear. With the right hand pass the other in an opposite line, corresponding with the course of the first blade, guided by the fore and middle fingers of the left hand; the third and little fingers being employed to retain the left-hand blade in place; humour the instrument so as to allow the blades to lock. The handles may now be tied gently together. The force used must be from blade to blade, with a gentle, tractile effort.

2. *Long Forceps*

Applicable where the head does not enter the brim. The long forceps must be applied, in most cases, over the occiput and face of the child, so that the convex edges of the blades may be towards the hollow of the sacrum. When used, the power may be exerted from side to side, with moderate traction.

3. *Lever or Vectis*

The lever is applicable to the same cases as the forceps. It must not be used strictly as a lever, but as a hook with which to draw downwards. It may be applied to any part of the head, but is generally hitched on the occiput.

4. *Fillet and Blunt Hook*

In certain cases of breech and knee presentation, where the hand is insufficient, it may be necessary to pass over the joint a fillet, or the blunt hook, with which to exert some force of traction. The operation is very simple.

INSTRUMENTAL LABOURS.—CONTINUED.

EMBRYULEIA

SYMPHYSTOMY AND CÆSAREAN SECTION.

SEPARATION OF THE HEAD OF THE FŒTUS FROM THE BODY.

INDUCTION OF PREMATURE LABOUR.

Disproportionate size of head, or distorted or deformed pelvis, may require the child to be destroyed, and its bulk lessened; or the Cæsarean section or symphysiotomy, or the induction of premature labour may be called for.

The instruments required here, are the *Perforator*, *Crotchet*, and *Craniotomy Forceps*. Make a steady pressure on the abdomen; pass two fingers of the left hand up to the head: feel for a suture or fontanelle; introduce the perforator and bore through, until the progress of the instrument is arrested by its *shoulders*. Open the handles, and turn the instrument in different directions, so that the opening may be large enough to admit the perforator, with which the brain must be broken down.

If, after this, delivery cannot be accomplished without further aid, pass up the craniotomy forceps; open the handles slightly, and introduce the blade without teeth within the cranium. On closing the forceps, a hold is obtained; and a tractive force can now be exerted during the pains, and the extrusion of the child be effected.

Should great difficulty exist at the brim, the bones at the top of the head may have to be removed, until the base alone remains. The chin must then be brought through first.

The crotchet is often used instead of the craniotomy forceps, being passed into the opening made by the perforator, and hooked upon some bony projection in the interior of the skull; but it is not a good instrument. In presentations of the face demanding perforation of the cranium, the perforator should be introduced just above the nose, in the sagittal suture.

When it is necessary to open the head after the lower extremities have been expelled, the perforation must be made behind the ear.

In cases of great narrowness of the pelvis, the operation of *Symphysiotomy* has been adopted by French practitioners,—rarely by British or American; and, in extreme cases, recourse has been had to the *Cæsarean Section*.

If the body of the fœtus has been forcibly separated, and the head left in the uterus, it must be laid hold of with one hand, and the forceps be applied; or the head be opened and delivered, as under embryuleia. Should the head have been brought away, and the body left behind, the feet must be brought down.

In cases of narrowness of the pelvis, where the fœtus, at the full term, cannot be born alive, delivery may be brought about at an earlier period, or as soon as the child is capable of carrying on its functions independently of the mother.

The most advisable plan for accomplishing this object is merely to pass the finger round and round within the os and cervix uteri, so as to detach the decidua;—the membranes being thus left entire, and the life of the child not so much endangered, as where the membranes are punctured, and the waters evacuated. Parturition usually follows within forty-eight hours.

The laying of eggs by animals is termed, in French, *ponte*. The term has been applied to the periodical discharge—*ponte périodique*—of ova, (F.) *Chute des Œufs*, which is supposed to take place from the ovaries at each menstrual period.

PARTURITION, DRY, Partus siccus.

PARTURIUM VANUM, *Fausse couche*.

PARTUS, Parturition—p. Abactio, Abortion—p. Abactus, Abortion—p. Cæsareus, Cæsarean section—p. Difficilis, Dystocia—p. Laboriosus, Dystocia, Laborious labour—p. Præcox, Parturition (premature)—p. Præmaturus, Parturition (premature)—p. Serotinus, Opsitocia.

PARTUS SICCUS, Dry labour or parturition. Labour, which is neither preceded nor attended by a discharge of the liquor amnii.

PARULIS, from *para*, and *ovulov*, 'the gum,' *Apostema parulis* seu *paroulis*, *Phlegmonè Parulis*, Gum Boil, Alveolar abscess. Small abscesses are so called which generally form in the alveoli and involve the gums, sometimes without any known cause, but which often depend upon carious teeth.

PARURIA, from *para*, and *ουρεω*, 'I pass urine,' Morbid secretion or discharge of urine.

PARURIA INCONTINENS, Enuresis—p. Incontinens aquosa, see Diabetes—p. Inops, Ischuria, false—p. Mellita, Diabetes—p. Retentionis, Ischuria—p. Retentionis renalis, Ischuria—p. Retentionis vesicalis, Retention of urine—p. Stillatitia, Strangury—p. Stillatitia mucosa, Cystirrhœa.

PARVITAS MORBOSA PARTIUM ORGANICARUM, Microtesia.

PAR'YGRON, from *para*, and *υγρος*, 'humid,' A liquid or moist preparation for allaying topical inflammation.

PAS D'ÂNE, Tussilago.

PASMA, Catapasma.

PASQUE FLOWER, Anemone pulsatilla.

PASSA, Paronychia.

PASSÆ CORINTHIACÆ, see Vitis Corinthiaca.

PASSAGES, DIGESTIVE, Ways, digestive—p. Second, Ways, second.

PASSAMBIO, Rio vinaigro.

PASSE-PIERRE, Crithmum maritimum.

PASSERAGE, Iberis amara—p. Sauvage, Cardamine pratensis.

PASSIBILITY, GREAT, Hyperæsthesia.

PASSIFLORA LAURIFOLIA, Bay-leaved Passion Flower. Ord. Passifloræ. A native of Surinam. The fruit has a delicious smell and flavour, and quenches thirst, abates heat of the stomach, increases the appetite, &c.

PASSIFLORA MALFORSIS, Apple-shaped Granadilla. The fruit of this species is esteemed a delicacy in the West Indies, where it is served up at table in desserts.

Some of the genus are said to possess narcotic and emetic properties.

PASSIO, Disease, Passion—p. Æmoptoica, Hæmoptysis—p. Cadiva, Epilepsy—p. Cardiaca, Cardialgia—p. Coeliaca, Coeliac flux—p. Choleric, Cholera—p. Pellitica, Cholera—p. Hæmoptoica, Hæmoptysis—p. Hypochondriaca, Hypochondriasis—p. Hysterica, Hysteria—p. Iliaca, Ileus—p. Ischiadica, Neuralgia femoro-poplitea—p. Pleuritica, Pleuritis—p. Stomachica, Dyspepsia—p. Ventriculosa, Coeliac flux—p. Vomica, Phthisis pulmonalis.

PASSION, Pas'sio, An'imi Pathe'ma, Affec'tio, Emo'tio, from *pator*, *passus*, 'to suffer.' An active affection or emotion of the mind; as rage, terror, love, hatred, &c.

PASSION FLOWER, BAY-LEAVED, Passiflora laurifolia—p. Hystérique, Hysteria—p. Iliac, Ileus.

PASSIONES ANIMI, Affections of the mind.

PASSIVE, Passivus. Same etymon as Passion. An epithet for diseases which seem owing

to a greater or less diminution of the strength, or which are without apparent reaction. Thus, passive hemorrhages are such as supervene in debilitated individuals, or under conditions in which increased action of the arteries can scarcely be presumed. The term *passive* has, also, been given to aneurisms of the heart with extension of the parietes; in opposition to *active* aneurisms, with thickness of the parietes—hypertrophy.

PASSULÆ MAJORES, see *Vitis vinifera*—p. *Minores*, see *Vitis Corinthiaca*.

PASSULATUM, (*Uvæ passæ*, 'raisins, dried grapes.') A medicine in which the pulp of raisins is a chief ingredient.

PASSY, MINERAL WATERS OF. This village is in the vicinity of Paris. There are several springs, containing sulphates of lime, iron, and magnesia, alum, common salt, carbonate of iron, carbonic acid, &c. They are employed as a tonic and astringent.

PASTA DE ALTHÆÆ, Paste, marshmallow—p. *Amygdalina*, Confection (almond)—p. *Dactylis*, Paste, date—p. *Emulsiva*, Confection (almond)—p. *de Jujubis*, Paste of jujubes—p. *Regia*, Confection (almond).

PASTA GLYCRRHIZÆ GUMMA'TA ET ANISA'TA seu *Massa de extracto glycyrrhizæ*. A compound of extract of liquorice, gum Senegal, sugar, Florence orris-root, and volatile oil of aniseed. Demulcent.

PASTA ZINCI CHLORIDI, (F.) *Pâte de Canquoin*, Paste of Chloride of Zinc, Canquoin's Caustic Paste. This paste is made of three strengths, consisting, respectively, of chloride of zinc, 1 part; wheaten flour, parts 4, 3 and 2; water sufficient to make a paste. This caustic destroys the part in a direct ratio with the thickness of the layer applied.

PASTE, *Pasta*, (F.) *Pâte*. A viscid, tenacious mixture. Also, a compound medicine, like the pastil, but less consistent, flexible, less saccharine, and more mucilaginous.

A convenient paste for attaching labels to glass, or for other services, is made of gum tragacanth and water, to which alum is added. This the author always keeps by him. Another form is to make a solution of Gum Arabic, 3j, boiling water, f3ij, and glycerin, f3ij; and another is made by mixing powdered Gum Arabic, and powdered tragacanth, of each 3ss, water, f3iss, acetic acid, Mxx.

PASTE, CAGLIARI, see Turundæ Italicæ.

PASTE OF CHLORIDE OF ZINC, *Pasta Zinci chloridi*.

PASTE, DATE, *Massa seu Pasta de dactylis*. Composed of dates, sugar, gum Arabic, water, and orange-flower water.—Ph. P. Demulcent.

PASTE OF ICELAND MOSS, *Pasta Lichenis*, (F.) *Pâte de lichen*. A paste made by boiling lichen Islandicus to a proper consistence, and adding gum Arabic and sugar.

The *Pâte de Lichen opioce* contains opium in addition; about 3d of a grain to an ounce.

PASTE, ITALIAN, see Turundæ Italicæ.

PASTE OF JUJUBES, *Pasta de jujubis*, *Massa de zizyphorum fructu*, (F.) *Pâte de Jujube*. Composed of jujubes, gum Senegal, sugar, water, and mixture of lemon-peel. Demulcent.

PASTE, MARSHMAL'LOW, *Massa de gummi Arabico*, *Pasta de althææ*, (F.) *Pâte de Guimauve*. A compound of gum Arabic, sugar, white of egg, and orange-flower water.—Ph. P. It is demulcent.

PASTE, TOOTH, Dentifricium—p. Vienna, see Powder, Vienna—p. Ward's, Confectio piperis nigri.

PASTEL, *Isatis tinctoria*.

PASTÈQUE, *Cucurbita citrullus*.

PASTIL, *Pastillus*, *Diocres*, *Arvicula Cipria*,

Caude'la fuma'lis, *Phthois*, *Phthois'cus*. A compound of aromatic substances in various shapes, used as a fumigating material when set fire to. See Fumigation.

The French use the term *Pastille*, (L.) *Pastillus*, for a solid, dry, round, oblong, square, or triangular compound, whose base is essential oil, and sugar the excipient. These pastilles are in fact small lozenges, and only differ from the *tablette* in their smaller size. They are termed, also, *Saccharolés solides*.

PASTILLE, see Pastel.

PASTILLES DIGESTIVES D'HAUTE-RIVE-VICHY, see Trochisci Sodæ bicarbonatis—p. Fumigating, see Fumigation—p. *de Vichy*, see Trochisci Sodæ bicarbonatis.

PASTILLI DIGESTIVI DARCETII, see Trochisci Sodæ bicarbonatis—p. Emetina pectorales, Trochisci emetina pectorales.

PASTILLI DE MENTHÆ PIPERITÆ, *Peppermint drops*. (*Succ. alb.*, *aqua menthæ pip.*, *up. destillat.* aa ʒij. Boil to the consistence of an electuary. Take of fine white sugar in powder, ʒiv; volatile oil of peppermint, ʒss. Mix, and add the electuary whilst warm. Drop it then on marble, and dry the drops.—Ph. P.)

PASTILLUS, Pastil.

PASTINACA ALTISSIMA, P. opopanax—p. Anethum, Anethum graveolens—p. Graveolens, Anethum graveolens—p. Opaca, P. sativa.

PASTINACA OPOPANAX. *Ord.* Umbellifere. The name of the plant which furnishes the *Opopanax*. The plant, whence the gum resin is obtained, has, also, been called *Fer'ula opopanax* seu *opopanax*, *Pastinaca altissima*, *Laserpitium* seu *Seli'nium chiro'nium*, *Opopanax'cum*, *Panax heracle'um* seu *cost'num*, *Pastinac'cea*, *Kynu*, *Hercules-all-heal*, *Opopanax'wort*.

Opopanax, *Opopanax'cis gummi-resi'na*, *Gezir*, *Pop'anax*, *Gummi Pana'cis*, exudes from the roots when wounded. Its odour is strong and peculiar; taste, bitter and acrid. It is met with in the shops, in lumps of a reddish-yellow colour, without; white within. It has been recommended as an antispasmodic and emmenagogue; but is seldom used. Dose, gr. x. to ʒj, in powder.

PASTINACA PRATENSIS, P. sativa.

PASTINACA SATI'VA, *Elaphobol'cum*, *Pastinaca sylvest'ris* seu *opa'ca* seu *praten'sis*, *Anethum* seu *Seli'nium pastinac'ca*, *Ban'ica*, *Garden parsnep*, (F.) *Pana'is*. The root is sweet and nutritious, and is in esteem as an article of food. It has been used as a diuretic and demulcent in calculous affections, &c.

PASTINACA SYLVESTRIS, P. sativa—p. *Sylvestris tenuifolia officinarum*, *Daucus carota*.

PATATE, *Convolvulus batatas*.

PATCHOULI, *Pogostemon patchouli*.

PÂTE, Paste—p. *Arsenicale*, Arsenical paste—p. *de Canquoin*, *Pasta Zinci chloridi*—p. *Caustique de Vienna*, Powder, Vienna—p. *de Guimauve*, Paste, marshmallow—p. *de Jujube*, Paste of Jujubes.

PATEL'LA, *Rot'ula*, *R. genu*, *Epigon'atis*, *Epigon'tis*, *Aeromy'le*, *Epigon'nis* seu *Epigon'is* seu *Epig'nis*, *Mola* seu *Concha* seu *Oculus* seu *Oculus* seu *Sentum* genu, *Os scutiforme* seu *discoformis* seu *thyroco'des*, *Supergenu'm'lis*, *Cancroco'des*, *My'lac'is*, *My'le*, *Gamba*, *Olecranon mol'ile*, (Prov.) *Cop bone*, (F.) *Rotule*. Diminutive of *patera*, 'a bowl,' so called from its shape. A sort of sesamoid, flat, short, thick, rounded bone, situate in front of the knee, and enveloped in the substance of the tendon common to the extensor muscles of the leg. The posterior surface of the patella is divided into two *facettes*, united angularly, which are encrusted with cartilage, and each of which is articulated with one of the condyles of

the femur. The patella is developed by a single point of ossification. When the patella is fractured, it is difficult to produce bony union. This can only be accomplished by keeping the fractured extremities closely in apposition.

PATELLA, Cup—*p.* Fixa, Olecranon.

PATENCY OF THE FORAMEN OVALE, see Cyanopathy.

PATENT MEDICINE, *Médecine brevetée* ou *patentée*, *Remède breveté* ou *patenté*. A medicinal compound, for the sale of which the proposer obtains from government an exclusive privilege. A specification of his invention must be registered in the Patent Office, and, hence, it differs from a secret medicine.

PATER OMNIUM VIVENTUM, Penis.

PATERA DIOGENIS, Palm.

PÂTEUX, Clammy.

PATHEMA, Affection, Disease.

PATHEMATOLOGY, Pathology.

PATHE'TIC, *Pathet'icus*, from *παθεω*, 'I feel.' That which moves the passions. A name given to the superior oblique muscle of the eye, and, also, to a nerve.

PATHE'TIC MUSCLE, Oblique superior of the eye.

PATHE'TIC NERVE, *Nervus Pathet'icus* seu *trochlearis* seu *trochleatorius* seu *trochleator*, the 4th pair of nerves, *Par quartum*, (F.) *Oculo-musculaire interne* (Ch.), *Nerf pathétique*, is the smallest encephalic nerve. It arises behind the inferior part of the tubercula quadrigemina (*testes*) at the lateral parts of the valvula Vieussensii. Its roots unite into one thin round cord, which makes a turn upon the peduncles of the brain, and passes into a canal of the dura mater, near the posterior clinoid process. It enters the orbit by the broadest parts of the sphenoidal fissure, and terminates in the superior oblique muscle of the eye, by dividing into several filaments,

PATHE'TISM, Magnetism, animal.

PATHIC, Morbid.

PATHICAL, Morbid.

PATHICUS, Morbid.

PATHNE, Alveolus.

PATHOGENET'IC, *Pathogenet'icus*, *Pathogen'icus*. Same etymon as Pathogeny. Relating to pathogeny.

PATHOGENETIC PHYSIOLOGY, *Physiolog'ia pathogenet'ica*. That part of physiology which relates to the origin of disease.

PATHOGENICUS, Pathogenetic.

PATHOGEN'ENY, *Pathogen'ia*, *Nosogen'esie*, *Nosogen'ia*, from *παθος*, 'a disease,' and *γενεαις*, 'generation.' The branch of pathology which relates to the generation, production and development of disease.

PATHOGNOMON'IC, *Pathognomon'icus*, *Signum conjunctum* seu *characterist'icum* seu *morb essential'e*; from *παθος*, 'an affection,' and *γινωσκω*, 'I know.' A characteristic symptom of a disease.

PATHOGRAPHY, *Pathograph'ia*, from *παθος*, 'disease,' and *γραφω*, 'I describe.' A description of disease.

PATHOLOGIA, Pathology—*p.* Humoralis, Humorism—*p.* Solidaris, Solidism.

PATHOLOG'ICAL, *Patholog'icus*, *Pathol'ogus*, (F.) *Pathologique*. Same etymon as pathology. Relating to pathology. Often used in the sense of pathical.

PATHOLOGICAL ANATOMY, see Anatomy.

PATHOLOGIST, *Patholog'us*. Same etymon. One versed in pathology.

PATHOLOG'Y, *Pathemat'ology*, *Patholog'ia*, *Pathematolog'ia*, *Patholog'ice*, from *παθος*, 'a disease,' and *λογος*, 'a discourse.' The branch of medicine whose object is the knowledge of disease. It has been defined *diseased physiology*,

and *physiology of disease*. It is divided into *general* and *special*. The first considers diseases in common:—the *second*, the particular history of each. It is subdivided into *internal* and *external*, or *medical* and *surgical*.

PATHOLOGY, Humoral, Humorism.

PATHOMA'NIA, *Parapath'ia*, *Moral Insanity*, from *παθος*, 'a disease,' and *μανια*, *Mania sine Delir'io*. A morbid perversion of the natural feelings, affections, inclinations, temper, habits, moral disposition, and natural impulses, without any remarkable disorder or defect of the intellect, or knowing and reasoning faculties, and particularly without any insanity or hallucination.

PATHOMYOTOM'IA; from *παθος*, 'affection,' *μυς* or *μυων*, 'muscle,' and *τομή*, 'incision.' The title of a work by Dr. John Bulwer (1649), being "a dissection of the significative muscles of the affections of the mind."

PATHOPATRIDALGIA, Nostalgia.

PATHOS, Affection, Disease.

PATIENCE, *Rumex patientia*—*p.* *d'Eau*, *Rumex hydrolopathum*—*p.* *Garden*, *Rumex patientia*—*p.* *Rouge*, *Rumex sanguineus*—*p.* *Sauvage*, *Rumex acutus*.

PATIENT, see Sick.

PATIENT OF DEATH, Agony.

PATIENTIA, *Rumex patientia*.

PATIENTIÆ MUSCULUS, Levator scapulæ

PATNE, Alveolus.

PATOPATRIDALGIA, Nostalgia.

PATOR NA'R'IUM. The cavity or the opening of the nostrils.

PATRATIO, Ejaculation, (of sperm.)

PATRINIA JATAMANSI, *Nardus Indica*.

PATTE D'OIE (F.), 'Goose's foot.' An aponeurotic expansion, which covers the internal surface of the tibia at its upper part, and consists of the expansion formed by the tendons of the sartorius, gracilis, and semi-tendinosus muscles.

PATURSA, Syphilis.

PAU, (CLIMATE OF.) Pau is the capital of the department of the Lower Pyrenees, and is about 150 miles from Bordeaux, and 50 from Bayonne. The climate corresponds with that of the south-west of France generally; its great quality is the comparative mildness of its spring, and exemption from cold winds. It is regarded as a favourable winter residence for invalids labouring under chronic affections of the mucous membranes. It is too changeable for the consumptive, and to be avoided by the rheumatic.

PAULLIN'IA. This is an extract from a plant of the same name in Brazil. It is prepared by the Indians, and appears to possess excitant powers. In Brazil, and the neighbouring countries, it is given in the form of tisanæ—the powder being mixed with cocoa—in diarrhoea and dysentery. It has been employed also as a tonic. An extract is prepared in Brazil from *Paullinia sorb'ilis*, Ord. Sapindaceæ, known there under the name of *Guarana*, which is administered in similar pathological cases.

PAUME DE LA MAIN, Palm.

PAUNCH, Ingluivies.

PAUNTA, Sedatives.

PAUPIÈRE, Palpebra.

PAUSIME'NIA, from *παυσις*, 'cessation,' and *μηνες*, 'menses.' Cessation of the menses.

PAVANA WOOD, *Croton tiglium*.

PAVANNE, *Lanrus sassafras*.

PAVIL'ION. The French give the name *pa-villon*, to the expanded extremity of a canal or cavity,—for example, of a catheter, sound, &c.

PAVILION OF THE EAR, *Auric'ula*, *Pinna*, *O'tium*, *Ala*, (F.) *Auricule* ou *Oreille*, *Parillon* de l'oreille, is seated behind the cheeks, beneath

the temple, and anterior to the mastoid process. Its size varies in individuals. It is free above, behind, and below. Anteriorly and within, it is continuous with the neighbouring parts. Its outer surface has several prominences, the *Helix*, *Antihelix*, *Tragus*, *Antitragus*, and *Lobe*; and several cavities—the groove of the *Helix*, the *fossa nuchalis* seu *scaphoides*, and the *Concha*. The skin which covers the pavilion is very fine, and studded with a number of sebaceous follicles. The auricle is fixed to the head by three fibro-articular ligaments;—a *superior*, *anterior*, and *posterior*.

PAVILLON DE L'OREILLE, Pavilion of the ear—*p. de la Trompe*, see *Tuba Fallopiana*.

PAVIMENTUM CEREBRI, Base of the brain.

PAVINA, *Æsculus hippocastanum*.

PAVITATIO, *Tremor*; from *pavitare*, *pavitatum*, 'to tremble from fear.' Trembling from fright.

PAVOPHOBIA, *Panophobia*.

PAVOR (L.), Dread. Great fright. *Panophobia*.

PAVO'RES NOCTUR'NI seu **DORMIENTIUM**. Fear during sleep.

PAVOT, *Papaver*.

PAW, *Manus*.

PAWPAW, see *Carica papaya*.

PAXWAX, see *Nucha*.

PAXYWAXY, see *Nucha*.

PAZHAR, *Bezoar*.

PEA, *Pisum*—*p. Garden*, *Pisum*—*p. Ground Squirrel*, *Jeffersoniana Bartoni*—*p. Hoary*, *Galega Virginiana*—*p. Love*, *Abrus precatorius*—*p. Nut*, *Arachis hypogæa*—*p. Partridge*, *Cassia chamaecrista*—*p. Turkey*, *Galega Virginiana*.

PEACH BRANDY, *Amygdalus Persica*—*p. Tree*, *Amygdalus Persica*—*p. Wood*, see *Cæsalpinia*.

PEAGLE, *Primula* *veris*.

PEAR, see *Pyrus malus*—*p. Avocado*, *Persea gratissima*.

PEARL, *Perla*, *Mar'garon*, *Margari'ta*, *Margellum*, *Margel'is*, *Unio*, (F.) *Perle*. A roundish concretion of a silvery white colour, and brilliant polish, found in several shells, and especially in the *Aric'ula margarit'ifera*, which has, hence, been called *Mater perlarum*, *Margari'ta*, and *Mother of pearl*, and is much used in the fabrication of handles for surgical instruments, &c. Pearls were formerly given in powder as astringents and antacids. They consist of lime and an animal matter.

PEARL ASH, *Polash* of commerce.

PEARL, *Mother of*, see *Pearl*—*p. White*, *Bismuth*, *subnitrate of*.

PEAU, *Cutis*—*p. Divine*, *Condom*.

PEAU'CIER, *Platysma myoides*.

PECAN or **PECCAN NUT**, see *Hickory*.

PECCANT, *Peccans*, from *peccare*, 'to sin.' Morbid, not healthy. An epithet given by the humorists to the humours when erring in quality or quantity. See *Materia Morbosa*.

PECHAN, *Stomach*.

PECHEDEON, *Perinæum*.

PECHEGUERA. A pulmonary affection affecting children three or four months old in South America, which is promptly fatal.

PÉCHER, *Amygdalus Persica*.

PECHLAGRA, *Pechyagra*.

PECHING, *Anhelatio*.

PECHU'RIM seu **PICHU'RIM CORTEX**. A highly aromatic bark, the produce of a species of *laurus*. It is extremely fragrant, like cinnamon, which it greatly resembles in its properties. In Portugal it is used in the cure of dysentery, &c. See *Pichurim beans*.

PECH'YAGRA, *Pech'ingra*, *Ancon'agra*, *Pec-*

tilogra, from *πῆχυς*, 'the elbow,' and *αγρᾶ*, 'a seizure.' Gout affecting the elbow.

PECHYS, *Elbow*.

PECHYTYRBE, *Porphyra nautica*.

PECKISH, *Hungry*.

PECTE, *Cheese*.

PECTEN, *Pubes*. Also, a comb.

PECTEN DENTITUM, see *Sepes*—*p. Veneris*, *Scandix cerefolium*.

PECTIHAGRA, *Pechyagra*.

PECTIN, from *πηκτος*, 'coagulated, curdled.' A name given by *Baconnot* to a principle which forms the basis of vegetable jelly. It is extensively diffused in the juices of pulpy fruits and roots, especially when they are mature; and occasions these juices to coagulate when they are mixed with alcohol or boiled with sugar.

It is the basis of one of the classes of elementary principles of *Dr. Pereira*—the *pectinaceous*.

PECTINACEOUS, *Pectineous*.

PECTINALIS, from *pecten*, *pectinis*, 'the pubes'; *Pectinæ'us*, *Musculus liv'idus*, *Pectinatus*, (F.) *Pubio-fémoral*, *Supubio-fémoral* (Ch.), *Pectiné*. This muscle is situate at the inner and upper part of the thigh. It is long, flat, and triangular; is attached, above, to the space which separates the ileo-pectineal eminence from the spine of the os pubis; and, below, by a flat tendon, to the oblique line which descends from the lesser trochanter to the linea aspera. The pectinalis bends the thigh on the pelvis, and carries it outwards in adduction and rotation. It may also bend the pelvis on the thigh.

PECTINATED, *Pectinatus*, *Pectiniform*, *Pectinate*, (F.) *Pectiné*, from *pecten*, *pectinis*, 'a comb.' Having the shape of the teeth of a comb.

PECTINATED MUSCLES, *Pectinatus musculi*. The fascicular texture observed in the right auricle of the heart:—*Musculi Auriculæ Pectinatus*. Their main use probably is—to prevent the over-dilatation of the auricles.

PECTINATUS, *Pectinalis*.

PECTINÉ, *Pectinalis*, *Pectinated*.

PECTINEUS, *Pectinalis*.

PECTINIFORM, *Pectinated*.

PECTINOUS, *Pectinæ'us*, *Pectino'sus*, *Pectinaceous*, from *pectin*. Of or belonging to pectin or vegetable jelly,—as a *pectinous* or *pectinaceous* vegetable principle.

PECTORAL, from *pectus*, *pectoris*, 'the breast.' Relating to the breast. See *Pectorals*.

PECTORAL GRAND, *Pectoralis major*.

PECTORALE, *Corset*.

PECTORALIS INTERNUS, *Triangularis sterni*.

PECTORALIS MAJOR, from *pectus*, *pectoris*, 'the breast.' *Pectoralis*, *Sterno-cleido-brachialis*, (F.) *Sterno-costo-clavio-humeral*, *Sterno-humeral* (Ch.), *Grand Pectoral*. A large, flat, triangular muscle, situate at the anterior part of the chest, before the axilla. It is attached, on the one hand, by means of aponeurotic fibres, to the inner half of the anterior edge of the clavicle; to the anterior surface of the sternum, and to the cartilages of the first six true ribs;—on the other, by a strong tendon, which is inserted at the anterior edge of the bicipital groove of the humerus. The use of this muscle is to move the arm in various directions. If it be hanging by the side of the body, it carries it inwards and a little forwards. When the arm is raised, it depresses it and carries it inwards. It can, also, move it in rotation inwards. If the humerus be raised and fixed, the pectoralis major draws the trunk towards the upper extremity.

PECTORALIS MINOR, *Serratus anti'cus minor*, (F.) *Costo-coracoidien* (Ch.), *Petit Pectoral*, *Petit*

dentelé antérieur. This muscle is situate beneath the preceding, at the anterior and upper part of the chest. It is flat and triangular; is attached by its base, which appears digitated, to the upper edge and outer surface of the 3d, 4th, and 5th true ribs, and is inserted at the anterior part of the coracoid process. The pectoralis minor draws the scapula forwards and downwards, and makes it execute a rotatory motion, by virtue of which its inferior angle is carried backwards, and the anterior depressed. When the shoulder is fixed, it elevates the ribs to which it is attached, and aids in the dilatation of the chest.

PECTORALS, *Pectoralia*, *Thoracica*. Medicines considered proper for relieving or removing affections of the chest.

PECTORAL DROPS, **BATEMAN'S**, consist chiefly of *tincture of castor*, with some *camphor* and *opium*, flavoured by *aniseed*. Gray gives the following formula: *Castor*, ʒi; *ol. anisi*, ʒi; *camph.* ʒv; *cocci*, ʒss; *opii*, ʒij; *proof spirit*, a gallon.

A form, advised by a committee of the Philadelphia College of Pharmacy, is the following:—*Alcohol. dil. cong.* iv; *Santal. rubr. ras.* ʒi; *digerere per horas* xxiv; *cola*, et adde *Pulv. opii*, *Pulv. catechu*, *Camphoræ*, aa ʒij; *Ol. anisi*, ʒiv. Digest for ten days.

PECTORILOQUE, from *pectus*, *pectoris*, 'the chest,' and *loqui*, 'to speak.' One, according to Laënnec, who presents the phenomenon of *Pectoriloquism*. Also, a stethoscope.

PECTORILOQUE, *Pectoriloquy*—*p. Chérotante*, *Egophony*.

PECTORILOQUOUS BRONCHOPHONY, see *Pectoriloquy*.

PECTORIL'OQUY, *Pectoril'oquism*, *Cav'ernous voice*, *Pectoril'oquia*, (F.) *Pectoriloque*, *Voix cavernueuse*, *V. articulée*. Same etymon. Speech or voice coming from the chest. Laënnec has designated, by this name, the phenomenon often presented by consumptive individuals, when their chests are examined with the stethoscope. The voice seems to issue *directly* from the chest, and to pass through the central canal of the cylinder, — a phenomenon owing to the voice resounding in the anfractuons cavities, produced in the lungs by the suppurative or breaking down of tubercles, which constitute *abscesses* or *ulcers* of the lungs. By some, it is not separated from bronchophony, having been termed, at times, *strong bronchophony*, and *pectoril'oquious bronchophony*. See *Laryngophony*. Whispering *pectoriloquy* is that produced by the whispering of the patient.

PECTORO'SUS, from *pectus*, *pectoris*, 'the chest.' One who has a broad chest.

PECTUS, Sternum, Thorax — *p. Carinatum*, see *Carina* and *Lordosis*.

PECULIUM, Penis.

PEDAGRA, *Potassæ supertartras impurus*.

PEDAL, *Pedal'is*, from *pes*, *pedis*, 'the foot.' Relating to the foot: — as

PEDAL APONEUROSIS, (F.) *Aponévrose pédieuse*. A thin, aponeurotic layer, which invests the extensor brevis digitorum pedis, and separates the tendons of the foot from each other.

PEDEBELLY, *Physconia*.

PEDES, Genital organs.

PEDESIS, Pulsation.

PEDETHIMOS, Pulsation.

PEDAL'GIA, *Pedional'gia*, *Pedioneural'gia*, from *πέδιον*, 'the sole of the foot,' and *άλγος*, 'pain.' Pain in the sole of the foot. Neuralgia of the foot in general. An affection of this kind, under the name *Pédionalgie*, prevailed in 1762 at Savigliano in Piedmont, and in 1806 amongst the military at Padua.

PEDICLE, **VITELLINE**, see *Vesicula umbilicalis*.

PEDICULARIA, *Delphinium staphisagria*.

PEDICULATIO, *Phtheiriasis*.

PEDIC'ULUS, *Phtheir*, *Louse*, (Prov.) *Biôdy*, (F.) *Pou*. Diminutive of *pes*, *pedis*, 'a foot,' from its having many feet. A genus of parasitic insects. The human body is infested with three kinds: the *Body-louse*, or *Clothes-louse*, *Pedic'ulus vestimen'ti*, (F.) *Pou de corps*; the *Head-louse*, *Pedic'ulus cap'itis*, (F.) *Pou de la tête*, which lives in the hair; the *Crab-louse*, *Mor'pio*, *Fera'lis pedic'ulus*, *Plat'ula*, *Phth'rius inguinal'is*, *Pedic'ulus Pubis*, (F.) *Mor'pion*, which infests the hair of the pubes. Infusion of tobacco, or mercurial ointment, or ointment of white precipitate of mercury, or the white or red precipitate of mercury, readily destroys them. The louse occurring in phtheiriasis, *pedic'ulus tabescent'ium*, differs from the common louse.—Vogel.

PEDICULUS CAPITIS, see *Pediculus* — *p. Pubis*, *Crab-louse*—*p. Tabescentium*, see *Pediculus*—*p. Vestimentis*, see *Pediculus*.

PÉDICURE, *Chiropodist*.

PEDICUS, *Extensor brevis digitorum pedis*.

PÉDIEUX (muscle), *Extensor brevis digitorum pedis*.

PEDIGAMBRA, *Ysambra*.

PEDILANTHUS TITHYMALOÏDES, *Euphor'bia tithymaloïdes* seu *myrtifolia*, *Jew bush*, *Order*, *Euphorbiacæ*, grows in various parts of the West Indies. The practitioners of Curacao give a decoction in venereal cases. It is also used in domestic practice, in amenorrhœa. It is, moreover, known and used as *Ipecacuanha*.—Lindley.

PÉDIL'VIUM, from *pedes*, 'the feet,' and *lavo*, 'I wash.' *Lavip'edum*, (F.) *Pédiluve*, *Bain de pied*. A bath for the feet. See *Bath*.

PEDION, *Sole*.

PEDIONALGIA, *Pedialgia*.

PEDIONEURALGIA, *Pedialgia*.

PEDIUM, *Tarsus*.

PÉDONCULE, *Peduncle*.

PÉDONCULES DU CERVEAU, *Peduncles of the brain*—*p. du Cervelet*, *Corpora testiformia* — *p. du Conarium*, *Peduncles of the pineal gland* — *p. de la Glande pinéale*, *Peduncles of the pineal gland*.

PEDORA. The sordes of the eyes, ears, and feet.

PED'UNCLE, *Pedun'culus*, (F.) *Pédoncule*, from *pes*, *pedis*, 'a foot.' This term has been applied to different prolongations or appendices of the encephalon.

PEDUNCLES OF THE BRAIN, *Pedun'culi seu Crura seu Caudex cer'ebri*, called, also, *Crura anteriora medullæ oblongatæ*, (F.) *Pédoncules du Cerveau*, are two white cords, about ʒths of an inch in diameter, on the outside of the corpora albicantia. They arise from the medullary substance of the brain, and gradually approach, till they join the tuber annulare. They are formed, internally, of a mixture of cineritious and medullary matter; the former of which, being of a darker colour at one part than in any other part of the brain, has been called *Locus niger crurum cerebri*, *Substantia nigra pedun'culi*.

PEDUNCLES OF THE CEREBELLUM, *Pedun'culi seu Crura cerebel'li*, called, also, *Crura posteriora*, are six in number, three on each side—a superior, a middle, and an inferior. The superior peduncles are generally known as the *Processus cerebelli ad testes*, *Crura cerebel'li ad corpora quadrigemina*, *Bra'chia copulativa*: the inferior are the *Processus cerebelli ad medullam oblongatam*; and the middle are called, also, *Cerebel'lar peduncles*—*Processus seu Crura cerebel'li ad pontem*, *Bra'chia pontis*.

PEDUNCLES, INFERIOR, OF THE CEREBELLUM,

Corpora restiformia—p. of the Medulla oblongata, Corpora restiformia.

PEDUNCLES OF THE PINEAL GLAND. *Pedunculi conarii*, (F.) *Pédoncules de la glande pinéale*, *P. du Conarium*, are two medullary bands or strips, which seem to issue from the pineal gland, and proceed on each side upon the upper and inner region of the optic thalami. See Pineal Gland.

PEDUNCULI CEREBELLI, Peduncles of the Cerebellum—p. Cerebri, Peduncles of the brain—p. Conarii, Peduncles of the pineal gland, see Pineal gland—p. Medullæ oblongatæ, Corpora restiformia.

PEDUNCULUS, Peduncle.

PEGA, Canthus (lesser).

PEGANELE'ON, *Oleum rutaecum*, *O'leum ruta*, from *pegayon*, 'rué,' and *elaion*, 'oil.' Oil of Rue.

PEGANUM, Ruta.

PEGMIN, Corium phlogisticum.

PEINE, Hunger.

PEINOTHERAPIA, Limotherapeia.

PEIRA, Effort, Experience, Nisus.

PELA'DA, (F.) *Pelade*. A word of uncertain origin, applied to alopecia, especially to that resulting from syphilis. See Alopecia.

PELA'GIA. A kind of scaly erysipelas of the hands—sometimes of the legs, at others, of the face. Pellagra [?].

PELAGRA, Pellagra.

PELARGONIUM ANCEPS, *Peris'terâ anceps*. A South African plant, *Nat. Ord.* Geraniaceæ, the decoction of which is used by the Malays in amenorrhœa, and to promote parturition and abortion.

PELARGONIUM ANTIDYSENTERICUM, *Jenkins'onia antidy'senterica*. This South African plant is used by the natives in dysentery. It is boiled in milk.

PELARGONIUM CUCULLA'TUM. Common along the side of the Table Mountain. It has been recommended in decoction as an enema in colic, nephritis, and suppression of urine. It is an excellent emollient, and was formerly exported to Holland under the name *Herba Althæa*.

PELARGONIUM TRISTE, *Polygac'tium triste*. The root of this plant, which grows at the Cape of Good Hope, is somewhat astringent, and is used there in diarrhœa and dysentery, and as an anthelmintic.

PELICAN, *Pelica'nus*, *Peleca'nus*, *Pel'ecan*, *Pel'ecus*, *πελεκαν*, from *πελεκας*, 'I cleave.' An instrument, curved at the end like the beak of a pelican. Its use is to extract teeth.

PELI'CIDĒ, *Pelu'dē*, *Mel coc'tum*. Boiled honey.—Ruland and Johnson.

PELICOMETER, Pelvimeter.

PELIDNOMA, Eechymoma, Pelioma.

PELIO'MA, *Pelido'ma* from *πελος*, 'black.' A livid eechymosis.—Forestus. See Eechymoma. PÉLIOSE, Purpura.

PELIOSIS, Purpura hæmorrhagica—p. Rheumatica, see Purpura simplex.

PELLA, Cutis, Prepuce.

PELAGRA, *Pel'agra*, *Ichthyo'sis seu Tuber seu Impet'go Pellagra*, *Eryphanti'asis Ital'ica*, *Lep'ra Mediolanen'sis seu Lombard'ica*, *Scorbutus alpi'nus*, *Ma'nia pella'gria*, *Erythe'ma endem'icum seu pell'agrum*, *Insolazione de Primavera*, *Male del Sole*, *Dermat'agra*, *Erysip'elas period'ica nervo'sa chron'ica*, *Paral'ysis scorbut'ica*, *Scurvy of the Alps*, (F.) *Mal de misère*, *Scorbut des Alpes*; from *πελλα*, [?] 'skin,' and *αγρα*, 'seizure.' A disease, particularly noticed among the Milanese, which consists in the skin becoming covered with wrinkles, and assuming a scaly appearance, especially in the parts exposed to the air. The strength diminishes, the intellectual faculties and

sensations become obscure, and cramps, convulsions, and other lesions of muscular contractility supervene. It is a singular endemic affection, and has been supposed to follow everywhere the introduction of Indian corn: hence the name *Rapha'nia maiz'tica*, given to it by some of the Italian physicians.

An affection endemic in the Asturias is said to be a variety of pellagra. It is the *Rosa seu Lep'ra Astu'rica*, *Mal de la Rosa*.

PELLENS, from *pellere*, 'to drive.' Driving; impelling.

PELLENTIA MEDICAMENTA are medicines that promote the occurrence of the menstrual or hæmorrhoidal flux.

PELLICULA OVI, Membrana putaminis—p.

Summa, Epidermis—p. Superior, Epidermis.

PELLICULÆ CORDIS, Ostiola cordis.

PELLIS, Cutis—p. Summa, Epidermis.

PELLITORY, *Xanthoxylum fraxineum*—p. American, *Parietaria Pennsylvanica*—p. Bastard, *Achillea ptarmica*—p. of Spain, *Anthemis pyrethrum*—p. Wall, *Parietaria*.

PELMA, Sole.

PELOHÆ'MIA, (F.) *Pélohémie*, from *πηλος*, 'dirt,' and *αἷμα*, 'blood.' A condition of the blood in which it is thick, syrupy, and of a very dark colour—as in diseases of the blood, gangrene, &c.

PELOR, Monster.

PELORIA, Monster.

PELORIUM, Monster.

PELORUM, Monster.

PELOTE DE MER, Pila marina.

PELTALIS seu PELTATUS CARTILAGO, Thyroid cartilage, Xiphoid cartilage.

PELTAN'DRA VIRGIN'ICA, from *πελτη*, 'a target,' and *ανθη*, 'for anther,' from the shield-shaped stamens. *Arum Virgin'icum*, *Culla seu Lecon'tia seu Rensselaëria Virgin'ica*, *Arrow Arum*, *Taraho*, *Wampee*; indigenous; of the Order Araceæ. The fresh roots and seeds are acrid stimulants, like *Arum*.

PELTIDEA AMPLISSIMA, Lichen caninus—p. Aphthosa, Lichen aphthosus—p. Canina, Lichen caninus—p. Leucorrhiza, Lichen caninus—p. Malacea, Lichen caninus—p. Spuria, Lichen caninus.

PELTIGERA CANINA, Lichen caninus.

PELUDE, Pelicide.

PELVIC, *Pelvi'cus*, *Pelvi'nus*, (F.) *Pelvien*, from *pelvis*. Belonging or relating to the pelvis.

PELVIC APONEUROSIS, (F.) *Aponevrose pelvienne*. A tendinous lamina, given off from the sides of the pelvis, and from the entire circumference of the brim, which passes into and lines the pelvis, and is soon divided into two distinct layers;—one external, the lateral pelvic or obturator fascia, which continues to line the sides of the pelvis, and covers the obturator internus muscle; the other, internal or superior, which passes inwards upon the side of the prostate, bladder, and rectum in the male, and of the bladder, vagina, and rectum in the female, in order to form the floor of the pelvis.

PELVIC CAVITY, (F.) *Cavité pelvienne*. The cavity of the pelvis.

PELVIC MEMBERS. The lower extremities.

PELVIC SURFACE OF THE ILIUM. That which faces the pelvic cavity.

PELVICULA OCULI, Orbit.

PELVIDYMUS, (F.) *Pelvidyme*, from *pelvis*, and *δίδυμος*, 'a twin.' A monster united at the pelvis, and double above.

PELVI-TROCHANTERIAN, *Pelvi-trochanteria'nus*. That which relates to the pelvis and great trochanter. The *Pelvi-trochanterian region* is formed by the muscles—*pyramidalis*, two *obturator*, *geminii*, and *quadratus femoris*—which

pass from the pelvis to the digital cavity of the great trochanter.

PELVIE, Pelvic.

PELVIMETER, *Pelycom'eter*, *Pelyom'eter*, *Pellicom'eter*, *Pyelom'eter*, (F.) *Pelvimètre*, from *pelvis*, and *μετρον*, 'a measure.' A barbarous hybrid. This name has been given to different instruments, invented for measuring the diameters of the pelvis, and particularly the antero-posterior or fore-and-aft diameter of the brim:—*Pelvimeter*, (F.) *Pelvimetrie*. Two have been chiefly employed, especially by French practitioners:—the *Compas d'épaisseur*, and the *Pelvimeter* of Coutouly. 1. The *Compas d'épaisseur* or *Callipers* of Baudelocque, is formed like a pair of compasses—with blunt extremities—the branches of which can be moved at will. One of these is applied to the symphysis pubis, and the other on the sacrum. About three inches must be deducted from this measurement for the thickness of the mons veneris, pubis, and the base of the sacrum. 2. The *Pelvimeter* of Coutouly resembles the instruments used by shoemakers for measuring the length of the foot. The two branches are introduced, in a state of approximation, into the vagina, and then separated, so that one touches the promontory of the sacrum, the other comes behind the os pubis. It is a barbarous contrivance, and its introduction ought never to be attempted on the living subject. It does not, indeed, seem possible to introduce it without mischief.

The finger is, decidedly, the best pelvimeter, and by it we can judge whether the base of the sacrum be unusually prominent. Measurement of the pelvis by the hand has been called *Pelycochirometre'sis*; from *πῶδες*, 'the pelvis,' *χειρ*, 'the hand,' and *μετρον*, 'measure.'

PELVIMETRY, see Pelvimeter.

PELVIS, *Cho'ana*, *Py'elos*, *Lee'ane*, (F.) *Basin*; so called from *πῶδες*, and *πέλλος*, 'a basin,' because fancied to be shaped like an ancient basin. The part of the trunk which bounds the abdomen below. It is a large, bony, irregular, conoidal cavity,—open above and below,—which supports and contains a part of the intestines, and the urinary and genital organs; and serves, at the same time, as a fixed point for the articulation of the lower limbs, the attachment of their muscles, and the execution of their movements. The pelvis supports, behind, the vertebral column, and is sustained, before, by the ossa femorum. It is situate, in the adult, near the middle part of the body, and is composed of four broad, flat, unequally thick bones, differing much in their shape, size, and arrangement, which touch, are articulated at some part of their surface, and intimately united by means of a number of ligamentous fasciæ. Of these bones, two are behind, on the median line,—the sacrum and the coccyx; the two others are before and at the sides,—the ilia. They are fellows, and unite, before, with each other. The most important parts of the pelvis, in an obstetrical point of view, are the brim and the outlet. The BRIM, *Angustia abdominalis*, *Intro'itus*, *Apertu'ra pelvis superior*, *Upper Opening or strait of the Cavity of the Pelvis*, (F.) *Détroit supérieur*, *D. abdominal*, is the narrow part which separates the greater pelvis from the less—the false from the true, *Pelvis vera* seu *minor*. In the well-formed woman it is elliptical, and slightly inclined forwards. Its antero-posterior, sacro-pubic or conjugate diameter, in a standard pelvis, measures 4½ inches, but with the soft parts, 3½ inches; its transverse or iliac or lateral, 5½ inches, but with the soft parts 4 inches; and its oblique, Diameter of Deventer, with the soft parts, 4½ inches. The OUTLET, *Ex'itus*, *Inferior opening*, or *strait*, *Angustia perinæalis*, (F.) *Détroit inférieur*, *D. périnéal*,

forms the lower aperture of the pelvis. The antero-posterior diameter is here, on account of the mobility of the coccyx, 5 inches: the lateral, 4 inches. The AXIS OF THE PELVIS is important to be known in obstetrics. The *Axis of the Brim* is indicated by a straight line drawn from the umbilicus to the apex of the coccyx;—the *Axis of the Outlet* by a line drawn from the first bone of the sacrum to the entrance of the vagina. An imaginary curved line which indicates the direction of the canal of the pelvis, has occasionally been termed the *curve of Carus*; in consequence of its having been pointedly described by the German obstetrician.

PELVIS AURIUM, Cochlea—p. Cerebri, Infundibulum of the brain.

PELVIS OF THE KIDNEY, *Pelvis renalis* seu *renum*, *Sinus*, *Venter* seu *Alvus Renum*, (F.) *Bassinet*. This is a small, membranous pouch, occupying the posterior part of the fissure of the kidney. It is placed behind the renal artery and vein; is elongated from above to below; flattened from before to behind; irregularly oval in figure; and, below, contracts considerably, to be continuous with the ureter. It receives the orifices of the infundibula, which pour the urine secreted in the kidney into its cavity.

PELVIS MINOR, Pelvis—p. Ocularis, Scaphium oculare—p. Renalis, Pelvis of the kidney—p. Renum, Pelvis of the kidney—p. Vera, Pelvis.

PELYCOCHIROMETRESIS, see Pelvimeter.

PELYCOMETER, Pelvimeter.

PELYOMETER, Pelvimeter.

PEM'ICAN or PEM'ICAN. Meat cured, pounded, and mixed with fat. It has been much used as nutriment on long overland journeys.

PEMPHIGODES, *Pemph'igoid*, (F.) *Pemphigode*, from *πῆφις*, 'a blister,' and *εἶδος*, 'form.' An epithet given by the ancients to the fever that accompanies pemphigus.

PEM'PHIGUS, *Emphy'sis Pem'phigus*, *Pompholyx*, from *πῆφις*, *πῆφυος*, 'a blister,' *Febris bullo'sa* seu *vesicul'aris* seu *ampullo'sa* seu *pemphigode's* seu *pemphigode's*, *Exanthema serosum*, *Morta*, *Pem'phigus morta* seu *Helveticus* seu *major seu minor*, *Morbus bullo'sus* seu *vesicul'aris* seu *ampullo'eus*, *Pemph'ix*, *Pemph'ix*, *Pemph'ix*, *Typhus vesicul'aris*, *Vesic'ular Fever*, *Bladdery Fever*, *Water-blebs*, (F.) *Fievre bulleuse* ou *vésiculaire*, *Maladie bulleuse*. A disease, defined to consist of vesicles, scattered over the body; transparent, filbert-sized, with a red, inflamed edge, but without surrounding blush or tumefaction; on breaking, disposed to ulcerate; fluid, pellucid or slightly coloured; fever, typhous. It is doubtful whether any such idiopathic fever have ever existed: the probability is, that the fever and vesications have been an accidental complication. Cullen thinks the *Pemphigus Helveticus* must have been malignant sore throat. If, however, such a distinct disease did exist, it was probably only as an endemico-epidemic.

PEMPHIGUS GANGRENO'SA, *Rupia escharotica*—p. Hungaricus, see Anthrax—p. Infantilis, *Rupia escharotica*—p. Major, *Pemphigus*—p. Minor, *Pemphigus*—p. Morta, *Pemphigus*—p. Variolodes, see Varicella.

PEMPHIX, *Pemphigus*.

PEMPHIX, *Pemphigus*.

PEMPHYX, *Pemphigus*.

PEMPTÆA FEBRIS, Quintan.

PENÆA MUCRONATA, see Sarcocolla—p. Sarcocolla, *Sarcocolla*.

PENCHE, Abdomen.

PENCIL-LIKE PROCESSES, Styloid processes.

PENDULOUS ABDOMEN, *Physeonia*.

PENDULUM PALATI, *Velum pendulum palati*.

PEN'ETRATING, *Pen'etrans*, from *penetrare*, *penetratum*, (*penitus intrare*), 'to go into.' A wound is so called which penetrates one of the great splanchnic cavities—*Vulnus penetrans*.

A medicine is, also, so called, which is supposed to pass through the pores, and stimulate.

PENICILLUM, Compress, *Penicillus*.

PENICULUS, *Penicillum*, *Peniculus*, 'a painter's brush;' diminutive of *penis*, 'a tail.' A tent or plectet. A pessary. The secreting glandiform extremities of the *venæ portæ*. (*Ac'ini bilio'si*) have been so termed, as well as the villous textures.

PENICULUS, Tent.

PENICULUM, Compress.

PENICULUS, *Penicillus*, Tent.

PÉNIDE, *Penidium*.

PENIDES, *Saccharum hordeatum*.

PENID'UM, *Penidium saccharatum*, from *penis*, 'a tail,' from the shape [?], (F.) *Pénide*, *Sucre tors*. A kind of clarified sugar, made up into rolls. *Barley sugar*. It is demulcent; see *Saccharum hordeatum*.

PÉNIL, *Mons veneris*.

PENIS, 'a tail.' *Caulis*, *Coles*, *Caulces*, *Caulos*, *Mentula*, *Menta*, *Phallus*, *Parcifer*, *Posthæ*, *Prosthium*, *Crithæ*, *Cor'ynæ*, *Vomer*, *Pater om'nium viventium*, *Priap'us*, *Virga*, *V. viri'lis*, *V. genital'is*, *Vere'trum*, *Sathæ*, *Membrum viri'le* seu *seminale* seu *genitale viro'rum*, *Morion*, *Cyon*, *Sicula*, *Tentum*, *Hasta*, *H. nuptia'lis*, *H. viri'lis*, *Vas*, *Vas'culum*, *Pecul'ium*, *Vir'gula*, *Vir*, *Thyrus*, *Tennis*, *Clavus*, *Cauda salac*, *Cauda*, *Fas'cinum*, *F. viri'le*, *Muto*, *Nervus*, *N. fistulosus* seu *fistularis* seu *juveni'lis*, *Per'tica per se*, *Scapulis*, *Leco*, *Cucur'lio*, *Venna*, *Contus*, *Telum*, *Gurgulio*, *Sceptrum*, *Arma*, *Gla'dius*, *Lu'bricum*, *Caput*, *Muti'num*, *Palus*, *Pes'sulus*, *Radix*, *Ramus*, *Rutab'ulum*, *Arma Ventris*, *Columna adstans inguin'ibus*, *Columna*, *Pyr'amis*, *Trabs*, *Spina*, *Catagula viri'lis*, *Verpu*, *Macha'ra*, *Taurus*; the *Yard*, *Male organ*, &c. (F.) *Verge*, *Membre viril*. This organ, the use of which is to carry the seminal fluid into the female organs of generation, is cylindroid, long, and erectile, and situate before and beneath the symphysis pubis. In the ordinary state, it is soft and pendent in front of the scrotum. During erection, it becomes elongated, upright, and assumes a triangular shape. Its upper surface is called the *Dorsum penis*, (F.) *Dos de la verge*; and, at its anterior surface, there is a longitudinal projection formed by the canal of the urethra. The two sides of the penis are round, and its posterior extremity or root is attached to the pelvis. Its anterior extremity is free, and presents the glans, prepuce, and orifice of the urethra. The penis is formed of the corpora cavernosa, the principal seat of erection; the corpus spongiosum of the urethra, for the conveyance of the urine and sperm; and of the glans, which terminates the canal. The *Arteries* of the penis are branches of the internal pudic. The *Veins* correspond with the arteries. The *Nerves* are from the internal pudic.

PENIS CEREBRI, Pineal gland—p. *Femineus*, *Clitoris*—p. *Lipodermis*, *Paraphimosis*—p. *Muliebris*. *Clitoris*.

PENNATUS, *Penniform*.

PENNIFORM, *Pennifor'mis*, *Penna'tus*, from *penna*, 'a pen,' and *forma*, 'form.' An epithet for muscles whose fleshy fibres are inserted at the side of a middle tendon, like the feathers of a pen on their common stalk.

PENNSYLVANIA, **MINERAL WATERS** OF. The most celebrated are those of Bedford. The Carlisle, Ephrata, and Yellow Springs are also frequented.

PENNYCRESS, *Thlaspi*.

PENNYR'YAL, *Mentha pulegium*, *Hedeoma pulegioides*—p. *Hart's*, *Mentha cervina*.

PENNYWORT, **ASIATIC**, *Hydrocotyle Asiatica*.

PENSACOLA, see Saint Augustine.

PENSEE, *Viola tricolor*—p. *Sauvage*, *Viola tricolor*.

PENSILIA, *Pudibilia*.

PENSY, *Viola tricolor*.

PENTAM'YRON, from *πεντε*, 'five,' and *μυρον*, 'ointment.' An ancient ointment, consisting of five ingredients. These are said to have been—storax, mastich, wax, opobalsam, and unguentum nardinum.—Paulus, and Aëtius.

PENTAPHAR'MACON, from *πεντε*, 'five,' and *φαρμακον*, 'remedy.' Any medicine consisting of five ingredients.

PENTAPHYLLUM, *Potentilla reptans*.

PENTATEUCH, (SURGICAL,) *Pentateuchus*, from *πεντε*, 'five,' and *τευχος*, 'a book,' which signifies the five books of Moses—Genesis, Exodus, Leviticus, Numbers, and Deuteronomy. By analogy, some surgeons have given the name *Surgical Pentateuch* to the division of external diseases into five classes:—wounds, ulcers, tumours, luxations, and fractures.

PENTATH'ETUM, from *πεντε*, 'five,' and *τιθεσθαι*, 'to place.' An ancient plaster consisting of five ingredients.

PENULA, *Ingluvies*.

PENZANCE, **CLIMATE** OF. This is the chief residence of invalids in Cornwall, England, during the winter. It is situated on Mount's Bay, about ten miles from the Land's End. It is 5½° warmer in winter than London, 2° colder in summer; scarcely 1° warmer in the spring, and only about 2½° warmer in the autumn. It is a very favourable winter residence for the phthisical invalid.

PEONY, *Pæonia*.

PEPANSIS, Coction, *Maturation*.

PEPANTICOS, *Matrative*.

PEPASMOS, Coction, *Maturation*.

PEPAS'TIC, *Pepas'ticus*, from *πεπαισθαι*, 'I concoct.' A medicine supposed to have the power of favouring the concoction of diseases. *Matrative*.

PEPEIRUS, *Concocted*.

PEPERI, *Piper*.

PEPINO (S.). *A cucurbitacea*, which is cultivated in great abundance in the fields of Peru. The pulp or edible part is solid, juicy, and well flavoured; but is apt to disagree.

PEPO, *Cucurbita pepo*—p. *Lagenarius*, *Cucurbita lagenaria*—p. *Vulgaris*, *Cucurbita pepo*.

PEPPER, **BLACK**, *Piper nigrum*—p. *Cayenne*, *Capsicum annum*—p. *Chilly*, *Capsicum annum*—p. *Cubeb*, *Piper cubeba*—p. *Decorticated*, *Fulton's*, see *Piper nigrum*—p. *Guinea*, *Capsicum annum*—p. *Jamaica*, *Myrtus Pimento*—p. *Long*, *Piper longum*.

PEPPER BUSH, *SWEET*, *Clethra alnifolia*.

PEPPER, **MALAGUET'A**. This term has been applied to the fruit or seeds of several zingiberaceous plants, as well as to the pimento. It is usually considered to be synonymous with *Grana Paradi'si*, *Grains of Paradise*, and *Grana Guine'nsia*, *Guinea grains*.—*Pereira*.

PEPPER, **POOR-MAN'S**, *Polygonum hydropiper*—p. *Tailed*, *Piper cubeba*—p. *Turnip*, *Arum triphyllum*—p. *Water*, *Polygonum hydropiper*—p. *Water*, of America, *Polygonum punctatum*—p. *Wall*, *Sedum*—p. *White*, *Piper album*.

PEPPERWORT, *Lepidium*.

PEPSIN, *Pepsine*, *Peps'inum*, *Chymosin*, *Gast'érase*, *Digestive Prin'ciple*, from *πεψω*, 'coction,' 'digestion.' A peculiar organic matter, which, in combination with the gastric acids, is considered to form the proper digestive solvent. Its chemi-

sal constitution is unknown. It would appear that its presence is necessary to induce changes in the elements of the food, which may enable the gastric acids to act upon them so as to form chyme.

PEPSINUM, Pepsin.

PEPSIS, Coction, Digestion.

PEPTIC, *Pep'ticus*, from *πενν*, 'I ripen,' 'I digest.' An agent that promotes digestion, or is digestive. Also, applied adjectively to an article of food that is easy of digestion.

•PEPTIC PERSUADER, see *Pilulæ aloes et Kinæ Kinæ*.

PEPTONE. A presumed definite combination between the solvent fluid in the stomach, and albuminous matters received into the organ as aliments. The same as Albuminose.

PER, 'by or through,' in composition often gives intensity to the signification: as *perussis*, a violent cough.

•PERACUTUS, Catoxys.

PERARTICULATIO, Diarthrosis.

PERATODYNIA, Cardialgia.

PERCE-CRANE, Perforator.

PERCE-FEUILLE, *Bupleurum rotundifolium*.

PERCE-MOUSSE, Polytrichum.

PERCE-MURAILLE, Parietaria.

PERCE-PIERRE, *Crithmum maritimum*.

PERCEP'TA, from *percipere*, *perceptum*, (*per*, and *capio*), 'to perceive,' 'receive.' A word used by some writers on hygiene to indicate, in a general manner, the effects of the nervous action on the animal economy; in other words, the sensations, the functions of the mind, and their deterioration or privation.


PERCEPTIBILITY, see Sensibility.

PERCEPTION, *Percep'tio*. Same etymon. The appreciation which the brain has of an impression made upon an organ of sense.

PERCEPTIV'ITY. Same etymon. The power of perception.

PERCOLATIO, Filtration, Percolation.

PERCOLA'TION, *Percolat'io*, from *percolare*, *percolatum*, (*per*, and *colare*), 'to strain through.'



Displacer or Percolator.

The terms percolation and displacement are applied in pharmacy to an operation which consists in placing any substance, the virtues of which have to be extracted by a menstruum, in a funnel-shaped instrument, having a septum perforated with holes, or its tube stuffed with cotton or tow, and pouring fresh portions of the menstruum upon it until all its virtues have been extracted. The operation is used in the formation of certain infusions, extracts, tinctures, &c.

An instrument used for this purpose is called a *displacer* or *percolator*.

PERCOLATOR, see Percolation.

PERCUSS. *Peren'teræ*, *Percussio*, (*per*, and *quære*, 'to strike,') (F.) *Perceuter*, *Frapper*. Same etymon as the next. To strike upon with the view of appreciating the resulting sound. To practice percussion.

PERCUSSION, *Epicorru'sis*, *Percus'sio*. When immediate or direct percussion is made on the chest or abdomen, the more or less perfect resonance is an index of the state of the contained

organs; and the physician is thus aided in his diagnosis. For this purpose the chest may be struck with the fingers, gathered into a bundle, and their tips placed upon a level. It is better, however, to employ *mediate percussion*, which consists in interposing, between the point of the fingers and the chest, the finger of the other hand, or a *Plexim'eter*, and striking this instead of the naked chest.

PERCUSSION, AUSCULTATORY, see *Acouophonia*.

PERCUTEUR, Percuss.

PERCUTEUR COURBE À MARTEAU, (F.) An instrument used by Baron Heurteloup in the operation of lithotomy, in which a hammer is employed instead of a screw, to force together the blades of the instrument on the foreign body.

PERDONIUM. A medicated wine of herbs.

—Paracelsus.

PERETERIUM, Trepan.

PERETORIUM, Trepan.

PERFORANS, from *perforare*, (*per*, and *forare*,) 'to bore through.' A name given to different muscles, whose tendons pass through intervals between the fibres or tendons of other muscles; thence called *perforated*. See *Flexor Longus Digitorum*, &c. Under the name *Perforating Ar'teries*, are included, 1. *In the hand*—arterial branches, given off by the profound palmar arch, which traverse the muscles and interosseous spaces. 2. *In the thigh*—three or four arteries, furnished by the profunda, which pass through the openings of the abductor magnus. 3. *In the foot*—the anterior and superior branches of the plantar arch.

PERFORANS CASSERI, see *Cutaneous*—p. *Mandis*, *Flexor profundus perforans*—p. *Profundus*, *Flexor longus digitorum pedis profundus perforans*.

PERFORATED SPOT, *Locus perforatus*.

PERFORATING ARTERIES, see *Perforans*.

PERFORATIO, Perforation—p. *Cranii*, see *Perforator*—p. *Intestinorum*, *Enterobrosis*—p. *Ventriculi*, *Gastrobrosis*.

PERFORA'TION, *Perforat'io*, *Anatre'sis*, *Diatre'sis*. Same etymon as *Perforans*. An accidental opening in the continuity of organs, either from an external or internal cause.

PERFORATION, Paracentesis—p. *de l'Estomac*, *Gastrobrosis*—p. *des Intestins*, *Enterobrosis*—p. *Uteri*, *Uterus*, rupture of the.

PERFORATOR, *Perforato'rium*, (F.) *Perce-crane*. An instrument for opening the head of the fœtus in utero, when it is necessary to diminish its size. The operation is called *Cephalotom'ia*, *Perforat'io cranii*.

PERFORATORIUM, Perforator.

PERFORA'TUS. That which is pierced. Anatomists have given this name to muscles whose fibres or tendons separate to suffer other parts to pass through them. Such are the *Flexor brevis digitorum pedis*, *F. sublimis perforatus*, and *Coraco-brachialis*. Also, to parts that are perforated for any purpose—as the *substant'ia perforata* of the brain.

PERFORATUS CASSERI, *Coraco-brachialis*.

PERFRIC'TIO, *Perfrige'rium*, *Catapysx'is*, *Peripsyx'is*, from *perfringere*, *perfrictum*, (*per*, and *frigere*), 'to shiver with cold.' Considerable refrigeration. Great sense of cold—shivering.

PERFRIGERATIO, see *Rigor*.

PERFRIGERIUM, *Perfrictio*.

PERFUSIO, Fomentation—p. *Frigida*, see *Affusion*.

PERI, *peri*, 'about, on all sides, round about.

An augmentative prefix. Hence:

PERIÆRESIS, from *peri*, and *αιρω*, 'I take away.' A sort of circular incision, which the ancients made in the neighbourhood of large ab-

cesses. The periæresis is now only practised in some cases of tumours.

PERIAL'GIA, from *peri*, and *αλγος*, 'pain.' A very violent pain.

PERIAMMA, Amuletum.

PERIAPTON, Amuletum.

PERIBLAST, *Periplast*, (F.) *Périblaste*, from *περι*, 'around,' and *βλαστος*, 'a germ.' The amorphous matter, which surrounds the endoblast, endoplast, (F.) Endoblaste or cell nucleus, and endergoes segmentation.

PERIBLEMA, Catabelma.

PERIBLEP'SIS, from *peri*, and *βλεπω*, 'I look.' The wild look that accompanies delirium.—Fœsius.

PERIB'OLE, from *περιβαλλω*, (*περι*, and *βαλλω*), 'I surround.' The dress of a person. The pericardium. Also, the translation of morbid matters towards the surface of the body.

PERIBRO'SIS, *Pericra'sis*, from *περιβρωσκω*, (*περι*, and *βρωσκω*, 'I eat,') 'I eat around.' Ulceration or erosion at the corners of the eyelids.

PERICAR'DIAC, *Pericardial*, *Pericardi'a-cus*; same etymon as Pericardium. Relating to the pericardium,—as '*pericardial murmur*,' '*pericardial effusion*,' '*pericardial arteries*, veins,' &c., &c.

PERICARDIAL, Pericardiac.

PERICARDI'TIS, from *pericardium*, and *itis*, denoting inflammation; *Inflamma'tio seu Pleuri'tis pericar'dii*, *Cardi'tis exter'na seu sero'sa seu membrano'sa*, *Exocardi'tis*, *Inflammatio of the pericardium*, (F.) *Péricardite*, *Inflammation du Péricarde*. This is probably the proper appellation for most of those cases which have received the names of *Carditis*, *Cardipericardi'tis*, *Cardo-pericardi'tis*, and *Cardipericardi'tis*. Along with signs of pyrexia, the local symptoms resemble those of pneumonia. Those which point out that the pericardium is the seat of disease, are the following:—pain, referred to the region of the heart, or the scrobiculus cordis,—sometimes pungent, at others, dull and heavy: palpitation, accompanied with spasmodic twitchings in the neighbourhood of the heart, shooting up to the left shoulder; pulsation, and sometimes soreness of the carotids, with tinnitus aurium and vertigo; the breathing is by catches; dyspnœa considerable; pulse jarring, jerking, peculiar; the tongue white, covered with a mucous coat, and the skin often bathed in sweat, as in acute rheumatism. The physical signs during the first period are as follows. The action of the heart is generally evident to the eye, and may be felt by the hand. There is soreness to the touch over the intercostal spaces, and over a small surface in the epigastric region, when the pressure is directed upwards towards the pericardium. Percussion is usually natural, but at times there is dulness. On auscultation, the cardiac movements are found to be frequent, abrupt, jerking, and tumultuous; often irregular and intermittent. The pulse presents corresponding characters. When effusion of lymph has occurred, percussion may be negative, or be but slightly affected. On auscultation, in addition to the preceding signs, there may be one or more of the rubbing or friction *bruits* resembling the rustling of parchment, or of a sawing or rasping character. In some cases, the sound is like the creaking of new leather. This has been supposed to be pathognomonic of effused lymph. The most important point in the pathology of pericarditis is its connexion with acute rheumatism; and it forms one of the most dangerous occurrences in the latter disease. It may be *acute* or *chronic*: in either case, it is, of course, formidable. The most active depletion must be used; with large doses of opium, counter-irritants, and all the means re-

quired in the most violent internal inflammations.

PERICARDI'IS EXSUDATORIA SANGUINOLENTA, Hæmopericardium.

PERICAR'DIUM, *Pericar'dion*, *Perib'olē*, *Membr'na Cor circumplex'a*, *Involucrum seu Ar'cula seu Capsa seu Cap'sula seu Cam'era seu Indumentum seu Pannic'ulus seu Membra'na seu Theca seu Sacus seu Sac'culus seu Scrotum seu Vagi'na seu Ves'ca seu Arca seu Thal'amus regalis seu Amphit'es'ma seu Bursa Cordis, Sae' or Capsule of the Heart*, (F.) *Péricarde*; from *peri*, and *καρδία*, 'the heart.' The pericardium is a membranous sac, which envelops the heart, and the arterial and venous trunks that pass from or into it. It is seated in the separation of the mediastinum, above the central aponeurosis of the diaphragm, to which it strongly adheres. Its shape is triangular, like that of the heart, to which it is fitted. The pericardium is composed of two membranes:—the one—*external*—*fibrous*; the other—*internal*—*serous*. The latter, having lined the inner surface of the external layer, is reflected over the heart, and covers it entirely; without, however, having the heart within it; in which arrangement it resembles other serous membranes. The pericardium envelops the heart; retains it in position, and facilitates its movements by means of the serous fluid contained in its cavity—*Cavum pericardii*—and secreted by the membrane—*Liquor seu Aqua seu Humor seu Lympha seu Uri'na Pericar'dii*, *Hydrocar'dia*, which it contains in greater or less quantity.

The arteries of the pericardium are small, and derived from the superior phrenic, anterior mediastinal and bronchial. The veins accompany them, and open into the brachio-cephalic. No nerves have been traced to it. The lymphatics enter the lymphatic glands that surround the vena cava superior.

PERICARPIUM, Epicarpium.

PERICHOLIA, Polycholia.

PERICHONDRITIS, *Inflamma'tio perichon'drii*, from *perichondrium*, and *itis*, denoting inflammation. Inflammation of the perichondrium: hence *Perichondri'tis larynge'a*.—Inflammation of the perichondrium covering the larynx.

PERICHONDRIUM, (F.) *Périchondre*, from *peri*, and *χόνδρος*, 'a cartilage.' A membrane of a fibrous nature, which covers cartilages that are non-articular, and bears considerable analogy to the periosteum in organization and uses.

PERICHRISIS, Circumlitio, Liniment.

PERICHRISTON, Circumlitio, Liniment.

PERICLA'SIS, from *peri*, and *κλω*, 'I break.' A comminuted fracture, with denudation of the bone.—Galen. Fœsius.

PERICLYMENUM, Lonicera periclymenum —p. Vulgare, Lonicera periclymenum.

PERICNE'MIA, from *peri*, and *κνημη*, 'the leg.' The parts surrounding the tibia;—the tibia itself.

PERICOLONI'TIS, *Pericoli'tis*, from *peri*, and *colonitis*. Inflammation of the areolar tissue surrounding the colon.

PERICOLPITIS, from *peri*, *κολπος*, 'vagina,' and *itis*, denoting inflammation. Inflammation of the areolar tissue surrounding the vagina.

PERICRANIŒDE'MA, from *pericranium*, and *οἰδμα*, 'a swelling.' Œdema of the head, owing to fluid effused under the pericranium.

PERICRANIUM, *Pericra'nion*, *Pericra'nia*, (F.) *Péricrane*, from *peri*, and *κρανιον*, 'the cranium.' The periosteum, which covers the cranium externally.

PERICRASIS, Peribrosis.

PERICYSTI'TIS, from *peri*, *κυστις*, 'the bladder,' and *itis*, denoting inflammation. Inflammation.

mation of the areolar tissue separating the bladder from the neighbouring parts.

PERIDERIA, from *peri*, and *δερν*, 'the neck.' Twisting of the cord round the neck of the child.

PERIDESMIC, *Peridesmicus*; from *peri*, and *δεσμος*, 'a ligament.' Some nosologists have given this name to ischuria caused by a ligature round the penis or by stricture of the urethra.

PERIDESMIUM, from *peri*, and *δεσμος*, 'ligament.' The delicate areolar tissue covering a ligament.

PERIDIASTOLE, see Diastole.

PÉRIDIDYME, see Testicle.

PERIDROMUS, from *peri*, and *δρομος*, 'a course.' The extreme circumference of the hairs of the head. — Gorræus.

PERIESTICOS, Salutory.

PERIGLOTTIS, Epiglottic gland: see Tongue.

PERIGRAPHĒ, from *περιγραφω*, (*περι*, and *γραφω*, 'I write,') 'I circumscribe.' The aponeurotic intersections of the rectus abdominis are so called. — Vesalius.

PERIHEPATITIS, see Hepatitis.

PERILARYNGITIS, from *peri*, *larynx*, and *itis*, denoting inflammation. Inflammation of the areolar tissue surrounding the larynx, which not unfrequently ends in *perilaryngeal abscess*.

PERILYMPH, Cotunnus, liquor of.

PERIMYEELITIS, Meningitis, spinal.

PERIMYSIUM, *Fascia*, from *peri*, and *μυς*, 'a muscle.' The areolar membrane or sheath — *vaginata muscula'ris* — that surrounds a muscle, or its fasciculi. The sheath of the whole muscle is termed *Perimysium externum*; of the fasciculi, *Perimysium internum*.

PERIN, *Peris*, from *περα*, 'a sac or pouch.' With some, this means the scrotum; with others, the testicle; with others, the vagina; with others, the peritonæum; and with others, the anus.

PERINÆOCELE, Hernia, perineal.

PERINÆORRHAPHY, *Perinæorrhaphia*, from *περιναϊον*, 'perinæum,' and *ραφή*, 'suture.' Suture of the perinæum, (F.) *Suture du Périnée*, in cases of laceration of the same.

PERINÆUM, *Interfeminæum*, *Interfeminæum*, *Interfeminina*, *Iter feminæum*, *Peris*, *Perin*, *Perineum*, *Gressu'ra*, *Plechas*, *Amphiplex*, *Peche'deum*, *Mesos'celum*, *Mesos'celus*, *Perine'on*, *Perina'os*, *Perine'os*, *Tramis*, *Tauros*, *Interforaminæum*, *Cucho'nē*, *Plichos*, *Mesome'rium*, *Fœmen*, *Re'gio Perinæ'i*, (F.) *Périnée*, (*πρεν*, *πρενις*, 'the scrotum' [?].) The space at the inferior region of the trunk, between the ischiatic tuberosities, anus, and genital organs. It is smaller in the female than in the male; has a triangular shape, and is divided into two equal parts by a median line, called *Raphē*. It is occasionally ruptured in labour. At times, it has been made to extend posteriorly as far as the os coccygis. The part between the pudendum and anus is sometimes called *anterior perinæum*, to distinguish it from that which extends from the anus to the coccyx, called *posterior perinæum*.

PERINÆUS, Perineal.

PERINÆAL, *Perinæus*, *Perinæalis*, *Perinæalis*. That which relates or belongs to the perinæum; as *Perineal artery*, *P. hernia*, &c.

PERINÆAL APONEUROSIS, P. Fascia.

PERINÆAL ARTERY, *Arteria perinæalis*, *Superficial artery of the Perinæum*. A branch of the internal pudic distributed to the perinæum.

PERINÆAL CUTANEOUS NERVE, see Sciatic nerve, lesser.

PERINÆAL FASCIA, *Fascia Perinæ'i*, *Perine'al aponeurosis*. The fasciæ, — *superficial and deep-seated* — *Perine'al ligament* — *Camper's ligament*, *Triangular ligament*, — which belong to the perinæum.

PERINÆAL FOSSA, *Fossa perinæ'i*, *Ischio-rectal fossa*. A conical fossa, the base of which corresponds with the skin; is formed anteriorly by the transversus perinæi muscle; behind, by the inferior border of the glutæus maximus; internally, by the levator ani; and externally, by the tuber ischii. It is filled up with fat and fibrous stræ.

PERINÆAL ISCHURIA, *Ischuria Perinæalis*. A name given by Sauvages to retention of urine, caused by a tumour seated in the perinæum.

PERINÆAL LIGAMENT, see P. fascia.

PERINÆAL NERVE. A branch of the internal pudic; which is mainly distributed to the perinæum and scrotum of the male; and to the vulva and perinæum of the female.

PERINÆAL REGION, Perinæum.

PERINÆAL SECTION, *Sec'tio perinæalis*. An incision into the perinæum for the cure of stricture of the urethra.

PERINÉE, Perinæum.

PERINEPHRITIS, (F.) *Phlegmon périnéphrétique*, from *peri*, *νεφρος*, 'kidney,' and *itis*, denoting inflammation. Inflammation of the external areolar and fibrous membranes of the kidney, or of their investing adipose areolar tissue.

PERINEPHRUS, see Kidney.

PERINEURION, Neurilemma.

PERINÈVRE, Neurilemma.

PERINYC'TIDES, *περι*, and *νυξ*, *νυκτες*, 'night.' An eruption appearing at night and disappearing by day. — Erotian.

PERIOD, *Periódus*, *Periódai'a*, *Periódousis*, *Stad'ium*, *Circu'itus*, (F.) *Période*, from *peri*, and *ódos*, 'way,' *Circuit*. Periods are the different phases or revolutions of a disease, — the epochs which are distinguishable in the course of a disease. Three periods are commonly enumerated.

1. The *augmentation*, *increase*, or *progress*, (*Incrementum*;) 2. The *acme* or height, (F.) *Étut* (*Status*;) and, 3. The *decline* (*Decrementum*.) Some authors reckon only the *invasion* and *termination*.

Period is sometimes used in describing an intermittent, for the time between the commencement of a paroxysm and that of the next, including the fit as well as interval.

PERIODS, MONTHLY, Menses.

PERIODEIA, Period.

PERIODEUSIS, Period.

PERIODEUTES, Charlatan.

PERIODICITY, *Periodic'itas*, *Rhythm*, (F.) *Périodicité*. Same etymon. The aptitude of certain physiological and pathological phenomena, in health or disease, to recur at particular periods, after longer or shorter intervals, during which they cease completely. Diseases, thus recurring, are called *periodical* or *rhythmical* — *Typo'nes*.

PERIODOLOG'Y, *Periodolog'ia*, from *περι*, *ódos*, 'a course or circuit,' and *lógos*, 'a discourse.' The doctrine of periodicity in health and disease.

PERIODONTITIS, (F.) *Périodontite*, *Inflammation de la Membrane alvéolo-dentaire*, from *peri*, and *ódon'tos*, 'a tooth.' Inflammation of the membrane that lines the socket of a tooth.

PERIODONTITIS GINGIVARUM, Ulitis.

PERIDOSCOPE, from *períodos*, 'a period,' and *σκοπεω*, 'I view.' An instrument, proposed by Dr. Tyler Smith, for the ready calculation of the periodical functions of the sex. It consists of a movable circular dial, upon which the months and days are engraved, fixed on a pivot in the centre of a large plate on which are numbered the different conditions of the reproductive system, as conception, abortion, premature labour, hemorrhage, labour, &c. By a knowledge of the date of conception, say November 14th, and fixing

the movable plate opposite the point on the fixed plate which indicates conception, the observer is enabled, at once, to see, that, allowing 280 days for gestation, labour may be expected about the 20th of August. The dial is made of card-board, and is affixed to a small volume, in which its uses and applications are explained.

PERIODUS LUNARIS, see Menses — p. Morbi, Type — p. Sanguinis, Circulation.

PERIODYN'IA, from *peri*, and *όδυν*, 'pain.' A violent and extensive pain.

PÉRIONE, Decidua.

PERIOR'BITA, *Perios'teum or'bitæ*, (F.) *Périorbite*, from *peri*, and *orbita*, 'the orbit.' An appellation used, by some, for the periosteum lining the orbit, which is a continuation of the dura mater. The inflammation of this periosteum is termed *Periorb'itis* or *Priorib'itis*, and *Inflam'matio perios'tei or'bitæ*.

PERIORBITIS, see *Periorbita*.

PERIORBITITIS, see *Periorbita*.

PERIOSTEITIS, see *Periostitis*.

PERIOS'TEOPHYTE, from *periosteum*, and *φύω*, 'I grow.' An osseous formation, commencing in the periosteum.

PERIOS'TEUM, *Perios'teon*, *Perios'teos*, *Périor'stium*, *Circumossat'le*, *Membra'na ossis*, *Omen'tum ossium*, *Circumossat'lis Membra'na*, (F.) *Périorste*, from *peri*, and *οσσω*, 'a bone.' The periosteum is a fibrous, white, resisting medium, which surrounds the bones every where, except the teeth at their coronæ, and the parts of other bones that are covered with cartilage. The external surface is united, in a more or less intimate manner, to the neighbouring parts by areolar tissue. Its inner surface covers the bone, whose depressions it accurately follows. It is united to the bone by small, fibrous prolongations; and, especially, by a prodigious quantity of vessels, which penetrate their substance. The periosteum unites the bones to the neighbouring parts. It assists in their growth, either by furnishing, at its inner surface, as M. Béchard demonstrated, an albuminous exudation, which becomes cartilaginous, and at length ossifies;—or by supporting the vessels, which penetrate them to carry the materials of their nutrition. See Medullary membrane.

PERIOSTEUM INTERNUM, Medullary membrane — p. Orbitæ, Periorbita.

PERIOSTI'TIS, *Perioste'i'tis*, *Inflam'matio perios'tei*, from *periosteum*, and *itis*, denoting inflammation. Inflammation of the periosteum.

PERIOS'TO-MEDULLI'TIS, (F.) *Périosteomédullite*, from *periosteum*, *medulla*, 'marrow,' and *itis*, a suffix denoting inflammation. Simultaneous inflammation of the periosteum and medulla spinalis. — Gerdy.

PERIOS'TO'SIS, (F.) *Périostose*. Tumour of the periosteum. Tumours are so called which are developed on bones, and formed by their outer membrane. These tumours are more rapidly developed, have a less degree of consistence, and sooner disperse than exostoses, which they resemble greatly.

PERIOSTOSTEITIS, (F.) *Périostostéite*, from *periosteum*, *οσσω*, 'a bone,' and *itis*, denoting inflammation. Simultaneous inflammation of the periosteum and bone. — Gerdy.

PERIPHERAD, see *Peripheral* aspect.

PERIPH'ERAL, *Periph'eric*, *Periph'erical*; from *peri*, and *φέρω*, 'I bear.' Relating or appertaining to the periphery or circumference.

PERIPHERAL ASPECT. An aspect towards the circumference of an organ. — Barclay. *Periph'erad* is used by the same writer adverbially to signify 'towards the peripheral aspect.'

PERIPHERAL VASCULAR SYSTEM, Capillary system.

PERIPHIMOSIS, *Paraphimosis*.

PERIPLAST, *Periblast*.

PERIPLEUMONIA, *Peripneumonia*.

PERIPLOCA INDICA, *Hemidesmus Indicus* — p. Mauritiana, see *Scammony*.

PERIP'LYSIS, from *peri*, and *λύσις*, 'the act of washing;' *Pragm'rium*. A copious discharge from any part, especially from the bowels.

PERIPNEUMO'NIA, *Peripleumonia*, *Peripneumoni'tis*, *True peripneumonia*, from *peri*, and *πνευμων*, 'the lung.' Inflammation of the substance of the lungs. See *Pneumonia*.

PERIPNEUMONIA BILIO'SA. Inflammation of the lungs, accompanied with bilious fever.

PERIPNEUMONIA CATARRH'ALIS. Bronchitis or pulmonary catarrh, with pain in some part of the chest. *Peripneumonia notha*.

PERIPNEUMO'NIA NOTHA, *P. spu'ria* seu *pituitosa* seu *catarrhalis*, *Pneumonia notha*, *Pseudo-peripneumonia*, *Pseudo-pneumonia*, *Pseudo-pneumoni'tis*, *Bronchoc'acæ*, *Blennothor'acæ*. *Obstruc'tio pulmonum pituitosa febri'lis*, *Bronchi'tis asthen'ica*, *False* or *Bastard Peripneumonia*, (F.) *Péripneumonie bâtarde* ou *fausse*, *Fausse pneumonie*. An inexact name, under which some affections are comprised that resemble pneumonia; and, especially chronic bronchitis with pleurodynia. See *Bronchitis* (chronic.)

PERIPNEUMONIA PITUITOSA, P. notha.

PERIPNEUMONITIS, *Peripneumonia*.

PERIPNEUMONY, *Pneumonia* — p. *Bastard*, *Peripneumonia notha*.

PERIPNEUMONY, LATENT. *Peripneumony* whose symptoms are so obscure as to be recognised with difficulty.

PERIPROCTI'TIS, from *περι*, 'around,' and *πρόκτος*, 'the anus.' Inflammation of the areolar tissue surrounding the rectum.

PERIPSYXIS, *Catapsyxis*, *Perfrictio*.

PERIPEY'MA, from *peri*, and *πύω*, 'pus. Suppuration around an organ; — a tooth for example.

PERIRRHO'E'A, *Perirr'hoë*, from *peri*, and *ρῶω*, 'I flow.' Afflux of fluids from every point of the body towards an organ which has to remove them from the economy. Also, enuresis.

PERIS, *Perin*.

PERISCCELIS, *Jarretière*.

PERISCYPHISMUS, *Periscythismus*.

PERISCYTHISMUS, *Periscyth'is*, *Periscyth'ismus*, *Periscy'this*, *Periscy'phismus*, *Periscyphismus*, *Scalping*, from *peri*, and *Σκυθ'ης*, 'a Scythian,' that is, 'scalping after the manner of the Scythians.' An operation described by Paulus of Ægina. It consisted of an incision made around the cranium, and was employed in habitual weakness of the eyes, pains of the head, &c.

PERISPAL'SIS, *Circumductio*, from *peri*, and *σπᾶλλω*, 'I move.' A circular motion impressed on a luxated bone, for the purpose of reducing it.

PERISSODACTYLUS, *Polydactylus*.

PERISPLENITIS, see *Splenitis*.

PERISTALTIC, *Systaltic* — p. *Action*, *Peristole*.

PERISTAPHYLINUS EXTERNUS, *Circumflexus musculus* — p. *Inferior*, *Circumflexus musculus* — p. *Internus*, *Levator palati*.

PERISTERA ANCEPS, *Pelargonium anceps*.

PERISTERIUM, *Verbena officinalis*.

PERISTER'NA, from *peri*, and *στέρον*, 'the sternum.' The lateral parts of the chest.

PERISTOLE, *Motus Peristalticus* seu *compressorius seu vermicularis seu testudin'is*, from *peri*, and *στέλω*, 'I contract,' 'I close.' The peristaltic action of the intestines, *Motus peristalt'icus*, (F.) *Mouvement péristaltique*. It consists of a kind of undulation or vermicular movement,

in appearance irregular, but in which the circular fibres of the muscular membrane of the intestine contract successively, from above downwards, in proportion as the chyme advances in the alimentary canal: so that, being compressed above, it is pushed into the next portion of the intestine, whose fibres are in a state of relaxation. When the fibres contract inversely, they occasion an *antiperistaltic action*. The peristaltic action is involuntary, and is not under the immediate influence of either brain or spinal marrow. It continues for some time after death.

PERISTOMA, Peristroma.

PERISTOMIUM, Mouth.

PERISTRO'MA, *Peris'toma, Mus'cus villo'sus, Crusta membrana'cea seu vermicula'ris*, from *peri*, and *σπρῶννυι*, 'to spread.' The villous or mucous coat of the intestines.—Castelli.

PERISYS'TOLE, from *peri*, and *συστολη*, 'contraction.' The interval that exists between the systole and diastole of the heart, which is only perceptible in the dying.—Bartholine.

PERITERION, *Trépan perforatif.*

PERITESTIS, Albuginea.

PÉRITOINE, Peritonæum.

PERITOME, Circumcision.

PERITONÆREX'IS, *Peritonæorex'is*, from *peritonæum*, and *ρνυμι*, 'to break.' Hernia formed by the rupture of the peritonæum.

PERITONÆ'UM, *Peritone'um, Peritonæ'on, Peritonæ'on, Peritonæ'os, Periton'ion, Peritone'os, Velamen'tum abdomina'le, Tu'nica præte'ri'sa seu Operimen'tum præte'rium seu Membra'na abdomi'nis, Pagos, Siphar, Syphar, Zephach, Ziphach, (F.) Péritoine, from peri, and teivo, 'I stretch.'* A serous membrane, which lines the abdominal cavity, *Peritonæ'um parietal'e*, extends over the greater part of the organs contained there, *Peritonæum intestinal'e seu viscera'le*, envelops them wholly or in part, and maintains their respective relations by means of different prolongations and ligamentous folds:—the *mesentery, epiploon, mesocolon*, &c. Like all the serous membranes, the peritoneum, except in the female, is a sort of sac without aperture, which covers the abdominal organs, without containing them within it, and the internal surface of which is smooth, and lubricated by a serous fluid. In the male fœtus, the peritonæum furnishes a prolongation, which accompanies the testicle at the time of its descent, and, in the female fœtus, forms the *Canal of Nuck*. Below the neck of the gall-bladder, the peritonæum has a triangular opening, called the *Foramen or Hiatus of Winslow*, through which it proceeds behind the stomach, to form a sort of secondary cavity, called the *posterior cavity of the peritonæum*, (F.) *Arrière cavité péritonéale ou Cavité des Épipoons*.

PERITONÆUM DUPLICATUM, Epiploon, gastro-
colic

PERITONE'AL, *Peritonæ'us*. That which appertains or relates to the peritoneum.

PÉRITONITE, Peritonitis — *p. Puerpérale*,
see Peritonitis.

PERITONITIS, *Empres'ma seu Cauma Peritonitis*, *Peritonæitis*, *Inflammatio seu Phlegmonæ Peritonæi*, *Phlegmæsia* or *Inflammatio of the Peritonæum*, from *peritonæum*, and *itis*, denoting 'inflammation.' (F.) *Péritonite*, *Inflammation du Péritoine*. The characteristic signs of acute inflammation of the peritonæum are, — violent pain in the abdomen, increased by the slightest pressure, often by the simple weight of the bed-clothes. It generally occurs in the parturient state; *Lochoperitonitis*, and begins on the second or third day after delivery. At times, a malignant epidemic, and perhaps contagious, variety has made its appearance, and destroyed

numbers of females. This has been described under the name *Puerperal Fever, Metropéritonitis, Ene'cia Syn'ochus Puerper'alum*, (F.) *Péritonite puerpérale, Métropéritonite puerpérale, Fièvre puerpérale, Typhus puerpéral, Erythemat'ic or Nonplastic Puerperal Peritonitis, Typho-hæmic Peritonitis, Adynam'ic or Malignant Puerper'al Fever, Low Fever of Childbed*. It is, according to some, a malignant inflammation of the peritoneum and uterus; according to others, a *Uterine Phlebitis*. In any form it requires active treatment, early employed. The appearances on dissection are such as occur whenever a serous membrane has been inflamed, and such inflammation has produced death:—effusion of fluid with flakes of coagulable lymph in it; appearances of vascularity, &c., in the membrane. The treatment—in *active peritonitis*—must consist of the same course as recommended under Enteritis. In the *chronic* kind, symptoms must be combated as they arise, and it may be advisable to excite a new action in the system by means of mercury. In the epidemic and malignant variety, unless bleeding be freely employed at the very outset, it does harm. If used early, it is of decided advantage. Everything depends upon arresting the morbid process at the outset.

PERITONITIS, ERYTHEMATIC, see Peritonitis —
p. Hepatica, see Hepatitis — p. Nonplastic, see
Peritonitis — p. Puerperal, see Peritonitis — p.
Typhohæmic, see Peritonitis.

PERITTOMA, Excrement.

PERITYPHLITIS; from *peri*, and *typhlitis*, inflammation of the cæcum. Inflammation of the areolar substance surrounding the cæcum. See Typhlo-enteritis.

PERI-UTERINE, *Peri-uteri'nus*, badly compounded from *peri*, and *uterus*. That which is situated around the uterus; hence "*periuterine inflammation*" or "*periuterine phlegmon*,"—inflammation of the areolar tissue around the uterus.

PERIWINKLE, LESSER, *Vinca minor*.

PERIZO'MA, from περιζωννυμι, (περι, and ζωννυμι, 'to gird,)' 'to gird around.' A bandage; a girdle. In Fabricius Hildanus, a truss. Herpes zoster. The diaphragm.

PERKINISM. A therapeutical means, first employed by Dr. Elisha Perkins, of Norwich, Connecticut, towards the termination of the last century, and named after him. It consisted in drawing over affected or other parts the pointed extremities of two metallic rods, called *Metallic Tractors*, each made of a different metal. The success obtained, was through the influence of the imagination; and, consequently, as a remedial agent, it ranks as high as animal magnetism, but no higher. The operation was, also, called *Tractoration*.

PERKINIST. A believer in and practiser of Perkinism.

PERKINIS'TIC. Relating or belonging to Perkinism.

PERLA, Pearl, Paralampsis.

PERLE, Péarl.

PERMISTIO, Coition.

PERMIXTIO, Coition.

PERMOTIO, Motion.

PERNICIEUX, Deleterious.

PERNICIOSUS, Deleterious

PERNIO, Chilblain.

PERNOCTATIO, Insomnia.
PEROCEPH'ALUS, from *προς*, 'deficient,'
and *κεφαλη*, 'head.' A monster with a defective
head.

PEROCOR'MUS, from *πηρος*, 'deficient,' and *κορμος*, 'trunk.' A malformation in which the trunk is defective, and too short, from the ab-

sence of one or more vertebrae,—the head and limbs being normal.

PERODACTYLEUS, Flexor longus digitorum pedis profundus perforans.

PERODYNIA, see Cardalgia.

PEROMELTA, from *περος*, 'deficient,' and *μελος*, 'a limb.' Congenital misconstruction, or mutilation of the limbs. A genus in the nosology of Good.

PEROMELUS. A monster possessed of the deformity mentioned above.

PÉRONÉ, Fibula—*p. Col du*, Collum fibulae.

PERONEAL, *Peronealis*, *Fibularis*, *Fibular*, (F.) *Péronier*, from *περονή*, 'the fibula.' Belonging or relating to the fibula.

PERONEAL ARTERY, *Fibular*, *A.*, arises from the popliteal, and is situate deeply in the posterior and outer part of the leg. It gives off branches of little importance. Near the outer ankle, it divides into two branches. 1. The *posterior peroneal*, which descends behind the lower joint of the fibula, on the outer surface of the calcaneum. 2. The *anterior peroneal*, which crosses the inferior extremity of the interosseous ligament, to be distributed on the dorsal surface of the foot.

PERONEAL MUSCLES are three in number. 1. *Peronæus Brevis* seu *Secundus* seu *Anticus* seu *Medius*, *Semi-fibularis*, (F.) *Grand-peronéo-métatarsien*, *Court péronéo-latéral*, *Moyen Péronier*. A muscle, situate beneath the peronæus longus. It is attached above, to the outer surface of the fibula, and terminates, below, at the posterior extremity of the 5th metatarsal bone, after having been reflected beneath the malleolus externus. It extends the foot upon the leg, at the same time raising a little the outer edge. It may also act upon the leg, and extend it upon the foot. 2. *Peronæus Longus* seu *primus* seu *posticus* seu *maximus* seu *posterior*, (F.) *Tibi-péronéo-tarsien*, *Long péronier latéral*, *Péronéo-tarsien*, (Ch.) *Muscle grand péronier*. This muscle is situate at the outer part of the leg. It is long, thick, and triangular above; thin, narrow, and flat below. It is attached above to the outer edge of the upper extremity of the fibula, and to the upper third of the outer surface of that bone. Below, it terminates at the outer part of the posterior extremity of the first metatarsal bone. This muscle extends the foot on the leg, turning its point outward, and raising its outer edge. It acts, also, upon the leg, which it extends on the foot. 3. *Peronæus Tertius*, *Novus Vesalii*, (F.) *Péronier antérieur*, *Petit péronéo-métatarsien* (Ch.), *Petit Péronier*. A muscle, situate at the anterior, outer, and inferior part of the leg. It is long and flat, and is attached, above, to the inferior third of the anterior margin and inner surface of the fibula; terminating, below, at the posterior extremity of the 5th metatarsal bone. This muscle bends the foot on the leg, by raising its outer edge. It can, also, bend the leg on the foot.

PERONEAL NERVE, see Popliteal nerves.

PERONEAL VEIN follows the same course as the artery.

PERONEDACTYLUS, Flexor longus digitorum pedis profundus perforans.

PÉRONÉO-LATÉRAL COURT, *Peronæus brevis*—*p. Phalangien du gros orteil*, Flexor longus pollicis pedis—*p. Sous-phalangien du pouce*, Flexor longus pollicis pedis—*p. Sous-tarsien*, *Peronæus longus*—*p. Sus-métatarsien*, *Peronæus tertius*—*p. Sus-métatarsien grand*, *Peronæus brevis*—*p. Sus-phalangien du pouce*, Extensor proprius pollicis pedis—*p. Tibi-sus-phalangien commun*, Extensor communis digi-

torum pedis—*p. Sus-phalangien du pouce*, Extensor proprius pollicis pedis.

PERONEUM, Fibula.

PÉRONIER, Peroneal.

PÉRONIER, GRAND, *Peronæus longus*—*p. Latéral long*, *Peronæus longus*—*p. Moyen*, *Peronæus brevis*—*p. Petit*, *Peronæus tertius*.

PERONODACTYLÆUS, Flexor longus digitorum pedis profundus perforans.

PERONODACTYLUS, Flexor longus digitorum pedis profundus perforans.

PEROSOMUS, from *περος*, 'deficient,' and *σωμα*, 'body.' A monster, whose whole body is imperfectly developed.—Gurlt.

PEROSPLANCHNICA, from *περος*, 'deficient,' and *σπλῆγχνον*, 'a viscus.' Congenital misconstruction of the viscera. A genus in the nosology of Good.

PERPENDICULUM HEPATIS, Suspensory ligament of the liver.

PERPERACUTUS. 'Extremely acute.' An epithet applied to very violent and rapid diseases.

PERPLEXI MORBI, see Complication.

PERPLICATION, *Perplicatio*; from *per*, 'through,' and *plico*, *plicatum*, 'to fold.' A method of tying arteries, which consists in making a small incision in the side of the artery, near its bleeding orifice, introducing a small pair of forceps, seizing the open extremity, and drawing it backward, through the aperture made in the side of the vessel, so as to form a kind of knot.

PERROSIN, see Pinus abies.

PERRY COUNTY WARM SPRINGS. These are in Pennsylvania, eleven miles from Carlisle, and fourteen from Harrisburg. Temperature 67°; impregnation not much. They are used for bathing.

PERSEA CAMFORA, see Camphor—*p. Cassia*, *Laurus cassia*—*p. Cinnamomum*, *Laurus cinnamomum*—*p. Cubeba*, *Piper cubeba*.

PERSEA GRATISIMA, *Laurus Persea*, *Palta tree*, *Avocado Pear*. A slender, very high tree of Peru, with a small dome-like top, which grows on the eastern declivity of the Andes, and rises, at times, to the height of more than 60 feet. Its fruit—(S.) *Palta*—which is pear-shaped, is generally much liked. It dissolves, like butter, on the tongue, and hence is called, in some of the French colonies, *beurre végétale*. The leaves are reckoned balsamic, pectoral and vulnerary. The seeds are very astringent.

PERSEA PICHURUM, see Pichurim beans—*p. Sassafras*, *Laurus sassafras*.

PERSIAN FIRE, Anthracion.

PERSICA VULGARIS, *Amygdalus Persica*.

PERSICAIRE ÂCRE, *Polygonum hydropiper*—*p. Brulante*, *Polygonum hydropiper*—*p. Douce*, *Persicaria*—*p. Poivrée*, *Polygonum hydropiper*.

PERSICARIA, *Polygonum Persicaria*—*p. Minor*, *Polygonum Persicaria*—*p. Mitis*, *Polygonum Persicaria*—*p. Crens*, *Polygonum hydropiper*—*p. Wandspiked*, *Polygonum Virginianum*—*p. Water*, *Polygonum amphibium*.

PERSIL, *Apium petroselinum*—*p. d'Ane*, *Chærophylum sylvestre*—*p. Faux*, *Æthusa cynapium*—*p. de Macédoine*, *Bubon Macedonicum*—*p. de Montagne*, *Athamanta aureoselinum*.

PERSIMMON, *Diospyros Virginiana*.

PERSOLATA, *Arctium lappa*.

PERSOLLATA, *Arctium lappa*.

PERSOLUTA, *Arctium lappa*.

PERSPICILLUM, see Spectacles.

PERSPIRABLE SANCTORIANUM, *Perspiration*.

PEPSPIRATION, *Perspiratio*, *Transpiratio*, *Diaphatio*, *Diapnoë*, *Diapnoë*, *Diapnoë*, *Perspirabile Sanctorianum*, from *per*, 'through,'

and *spirare, spiratum*, 'to breathe, exhale.' The *insensible transpiration* or exhalation continually going on at the surface of the skin and membranes. Sensible perspiration is called *sweat*. The perspiratory fluid is secreted by an appropriate glandular apparatus, termed by Breschet, *diapnoï'cous*, (F.) *Appareil diapnoïque*; the *sudorip'arous glands*, *Gland'ulæ sudorip'aræ*; *Organa sudorip'ara*, *sweat glands*, consisting of a convoluted tube, situate beneath the true skin, and of spiral ducts, which open obliquely under the scales of the epidermis. A certain portion of the perspired fluid is thrown off by ordinary physical evaporation.

The *Pulmonary exhalation* or *transpiration* is that which takes place into the bronchia, and is sent out with the expired air.

PERSPIRATORY, *Perspirato'rius*, *Transpi'ratory*, *Transpirato'rius*, (F.) *Perspiratoire*, *Transpi'ratoire*. Same etymon as the last. Relating or belonging to, or the product of, perspiration.

PERSUDATIO, Diapedesis.

PERSUTATIO, Diapedesis.

PERTË, *Metrorrhagia*—*p. d'Appetit*, *Anorexia*—*p. de la Mémoire*, *Annesia*—*p. de l'Odo'rat*, *Anosmia*—*p. de Sang*, *Hæmorrhagia*—*p. de la Voiz*, *Aphonia*—*p. de la Vue*, *Cæcitas*.

PERTÉRÉBRANT, *Perter'ebrans*, from *per*, 'through,' and *terebrare*, 'to bore.' The French give this epithet to an acute pain—*Douleur pertérébrante*—which occasions a sensation like that which would be produced by an instrument penetrating and lacerating a part;—the pain from whitlow, for example.

PERTES BLANCHES, *Leucorrhœa*—*p. de Sang*, *Metrorrhagia*—*p. Scémiales*, *Spermatorrhœa*—*p. Utrines*, *Metrorrhagia*—*p. Utrines blanches*, *Leucorrhœa*—*p. Utrines rouges*, *Metrorrhagia*.

PERTICA PER SE, *Penis*.

PERTURBATIONES ANIMI, Affections of the mind.

PERTURBA'TRIX, (*Medici'na*) from *perturbare*, *perturbatum* (*per* and *turbare*) 'to disturb greatly.' *Perturbating treatment*, (F.) *Méthode ou Médecine perturbatrice*; a mode of treatment in which very active means are employed, and such as impress upon diseases a different course from that natural to them. It is the antithesis to the *Medici'na expectans*. Much evil has resulted from such treatment, especially in febrile diseases.

PERTUS'SIS, from *per*, and *tussis*, 'a cough,' 'a violent cough.' *Tussis feri'na* seu *convulsiv'oa* seu *pueros string'ulans* seu *quinta seu stomachal'is* seu *an'hela clamo'sa* seu *stran'gulans* seu *delas'sans* seu *suff'ocans* seu *amphemer'ina* seu *tusiculo'sa* seu *spasmod'ica* seu *asini'na* seu *cani'na*, *Bronchi'tis convulsiv'oa*, *Bronchocephali'tis*, *Pneueis pertus'sis*, *Amphemer'ina*, *Othiopne'oa tussiculo'sa* seu *clango'sa* seu *suffocati'va* seu *clamo'sa* seu *spasmod'ica*, *Morbus Cuen'lius* seu *cuencl'aris*, *Cuen'lius*, *Bechorthopne'oa*, *Bex convulsiv'oa* seu *therio'des*, *Whooping-cough*, *Whooping-cough*, *Chin-cough*, *Kind-cough*, *Kind-coughs*, (Germ. Kind, 'a child,') (Sc.) *Kink-host*, (Prov.) *Kinkhaust*, (F.) *Couqueche*, *Catarhe* ou *Bronchite convulsive*, *Toux convulsive* ou *quintense* ou *bleue*, *Maladie cuculaire*. A violent convulsive cough, returning by fits, (F.) *Quintes*, at longer or shorter intervals; and consisting of several expirations, followed by a sonorous inspiration or *whoop*. The fits of coughing generally recur more frequently during the night, morning, and evening, than in the day. It is esteemed to be contagious, and attacks the young more particularly. It is rare for it to affect an individual for the second time. The duration is various,—six or eight weeks or more. Although the paroxysms are very violent, it is not a dangerous disease. It may, however,

give rise to other affections, as convulsions, pneumonia, &c., when the complication is very dangerous, as the cause cannot be removed. Those children suffer the least, who evacuate the contents of the stomach during the fit. In the treatment, all that can be done is to palliate. It must be borne in mind, that the disease will, in time, wear itself out. If there be such tensive pain of the head, or fever, bleeding may be required, but it is seldom necessary. Narcotics occasionally afford relief, but it is temporary. Gentle emetics, given occasionally, when the paroxysms are long and dry, give decided relief, and aid in the expectoration of the morbid secretions. After the disease has continued for some weeks, and persists in part from habit, change of air is essential, and this, even should the change be to an atmosphere that is less pure.

PERU, see *Lima*.

PERUNCIO, *Inunction*.

PERVENCHE, *Vinea minor*.

PERVERSION, *Perver'sio*, from *per* and *vertere*, *versum*, 'to turn.' *Dias'trophê*, *Diastrem'ma*. One of the four modifications of function in disease: the three others being augmentation, diminution, and abolition. The humourists used this term also to designate disorder or morbid change in the fluids.

PERVERSION DE LA TÊTE DES OS ET DES MUSCLES, *Loxarthrus*.

PERVIGILUM, *Insomnia*, *Coma vigil*.

PES, genitive *Pedis*, *pous*, *podos*, *Ichnus*, (Sc.)

Fit, (F.) *Pied*, 'the foot.' The inferior extremity of the abdominal member, representing a bony arch, an elastic vault, which transmits the weight of the body to the ground. It is divided into three parts;—the *tarsus*, *metatarsus*, and *toes*.

PES ALEXANDRINUS, *Anthemis pyrethrum*—*p. Anserinus*, *Chenopodium bonus Henricus*, *Facial nerve*—*p. Cati*, *Antennaria Dioica*—*p. Equinus*, *Pes equinus*, see *Kyllosis*—*p. Hippocampi major*, *Cornu ammonis*—*p. Hippocampi minor*, *Hippocampus minor*—*p. Hippopotami major*, *Cornu ammonis*—*p. Hippopotami minor*, *Cornu ammonis*—*p. Leonis*, *Alechemilla*.

PESANTEUR, *Weight*—*p. Spécifique*, *Gravity*, *specific*.

PÈSE-ACIDE, *Acidometer*—*p. Liqueur*, *Areometer*—*p. Sirop*, *Saccharometer*.

PESETTE, *Cicer arietinum*.

PESSAIRE, *Pessary*—*p. Intra-utérin*, *Pessary*, *intra-uterine*.

PES'SARY, *Pessa'rium*, from *πεσος*, 'a small stone; a pessary,' *Pessus*, *Perrum*, *Pes'mulus*, *Penicill'us*, *Talus*, *Nod'ulus uteri'nus*, *Suppositorium uteri'num*, *Glans*, *Lemnis'eus*, (F.) *Pessaire*. A solid instrument, composed of cork, ivory, or elastic gum, which is introduced into the vagina to support the uterus, in cases of prolapsus or relaxation of that organ; to keep vaginal hernia reduced, &c. The ancients made use of medicated pessaries, which they distinguished into emollient, astringent, aperient, &c., and they are still used. The form of pessaries is very variable;—round, oval, &c., according to circumstances.

PESSARY, EGYPTIAN, *Ægyptius pessus*.

PESSARY, INTRA-UTERINE, *Stem pessary*, (F.) *Pessaire intra-utérin*, *Redresseur utérin*. An instrument for rectifying uterine displacements—as anteversion and retroversion, consisting of a stem which is introduced into the uterus; this stem is attached to a flattened ovoid bulb or ball, on which the cervix uteri rests.—Connected with this is an external part or wire frame, which is attached, at one extremity, to a flat tubular portion, passing into the vagina, to be fixed to the intra-uterine portion. The wire frame is made to

press upon the pubis, so that the pessary can be kept in position *in utero*. This is essentially the intra-uterine pessary of Professor Simpson of Edinburgh.

PESSARY, *STEM*, Pessary, intra-uterine.

PESSULUS, Pessary, Penis.

PESSUM, Pessary.

PESSUS, Pessary.

PEST, Plague.

PESTE, Plague—*p. Antonine*, Plague of Antoninus—*p. d'Athènes*, Plague of Athens—*p. Noire*, Death, black.

PESTICHIÆ, Peticchiæ.

PESTICLE, Peticchiæ.

PES'TILENCE, *Pestilentia*, *Pestil'itas*, from *pestis*, 'a pest.' A malignant spreading disease;—applied especially to cholera.

PESTILENCE, CHOLERIC, Cholera spasmodica—*p. Glandular*, Plague—*p. Hæmagastric*, Fever, yellow—*p. Septic*, Plague.

PESTILENT-WORT, Tussilago petasites.

PESTILENTIA, Plague—*p. Hæmagastrica*, Fever, yellow.

PESTILENT'IAL, *Pestilentia'lis*, *Pestilentia'rius*, *Pestilentio'sus*, *Pestilen'tus*, (F.) *Pestilential*. Relating or appertaining to a pest or plague. *Pestilential diseases* are those that bear some resemblance to the plague in respect to mode of propagation, symptoms, danger, &c.*

PESTILITAS, Plague.

PESTIS, Plague—*p. Acutissima*, Plague—*p. Adeno-septica*, Plague—*p. Americana*, Fever, yellow—*p. Antoniniana*, Plague of Antoninus—*p. Bellica*, Typhus—*p. Britannica*, Sudor Anglicus—*p. Bubonaria orientalis*, Plague—*p. Contagiosa*, Plague—*p. Glandulosa*, Plague—*p. Glossanthrax*, Plague—*p. Inguinaria*, Plague—*p. Inter-tropica*, Fever, yellow—*p. Nigra*, Death, black—*p. Occidentalis*, Fever, yellow—*p. Orientalis*, Plague—*p. Septica*, Plague—*p. Variolosa*, Variola.

PESTLE, Pilum—*p. Spring*, see Pilum.

PET, Fart.

PETALE, Phtheiriasis.

PETASTES, Tussilago petasites—*p. Hybrida*, Tussilago petasites—*p. Major*, Tussilago petasites—*p. Officialis*, Tussilago petasites—*p. Rubens*, Tussilago petasites—*p. Vulgaris*, Tussilago petasites.

PETAURUM. A seat suspended by ropes, in which a person taking the exercise sat, and was tossed about by assistants.—Juvenal and Martial.

PETE'CHIE, *Petic'ula*, *Pestic'hæ*, *Pestic'ia*, *Punctic'ula*, *Pur'pura maligna*, *Por'phyra Græcorum*, (F.) *Pétéchies*. Small spots, similar in shape and colour to flea-bites, which occur spontaneously upon the skin, in the course of severe fevers, &c. They are, under such circumstances, signs of great prostration.

PETECHIE SINE FERRE, *Purpura simplex*.

PETE'CHIAL, *Petic'hia'lis*; same etymon. Resembling or accompanied by petechiæ.

PETECHIAL SCURVY, *Purpura simplex*.

PETECHIANOSIS, *Purpura hæmorrhagica*.

PETER'S PILLS, *Pilulæ Aloës et Cambogiæ*—*p. Worm lozenges*, see Worm lozenges, Sherman's.

PETICULÆ, Peticchiæ.

PETIGO, Impetigo.

PETINA, Sole.

PETIOLUS, *Pediolus*, diminutive of *pes*, 'a foot. A footstalk or leafstalk of a plant. A petiole.

PETIOLUS EPIGLOTTIDIS. The root of the Epiglottis.

PETIOLUS MALLEI, Manubrium mallei.

PETIT ARGENT, Platinum—*p. Courier*, Influenza—*p. Lait*, Serum lactis—*p. Lait d'Hoffmann*, Serum lactis Hoffmanni—*p. Mal*, see

Epilepsy—*p. Sus-maxillo-labial*, Levator anguli oris.

PETITE-POSTE, Influenza.

PETITS-LAITS, see Serum lactis.

PETRÆLEUM, Petroleum.

PETRÆUM, Solidago virgaurea.

PETRAPTIUM, Bubon Macedonicum.

PÊTRÉOLE, Petroleum.

PÊTREUX, Petrous—*p. Os*, see Temporal bone.

PETRO DEL PORCO, Bezoar of the Indian porcupine.

PÊTROLE, Petroleum.

PETRO'LEUM, *Petrola'um*, *Petræ'leum*, *Petræ'oleum*, *Bitu'men Petrolæum*, *Pissela'um Indicum*, *Rock Oil*, *Petroleum Barbadoes'se*, *Barbadoes Tar*, (F.) *Pétrole* ou *Pétréole*, from *petros*, 'rock,' and *ελαιον*, 'oil.' A bituminous substance flowing spontaneously from the earth, in various parts of the world; of a fetid odour, and bitter, acrid taste. It is semi-liquid, tenacious, semi-transparent; of a reddish-brown colour; insoluble in water and alcohol; combines with fixed and essential oils and sulphur; and is partially soluble in ether. The petroleum found at Gabian, near Béziers, in France, has been called *Oleum Gabianum*, (F.) *Huile de Gabian*, *Petroleum rubrum*. It has been given as an excitant expectorant; and, mixed with tincture of assafoetida, in tapeworm. It is, also, used externally in chilblains, chronic rheumatism, affections of the joints and skin, and paralysis.

New York Petroleum, called *Sen'eca Oil*, is used externally in similar cases.

PETROLEUM BARBADENSE, Petroleum.

PETROLEUM SULPHURATUM, *Balsamum Sulphuris Barbadoes'se*.—(Composed of petrol, *Barbad.* 3xxj, *flor. Sulph.* 3iv) has been used as a pectoral; and as a detergent to ulcers. It has been accounted antispasmodic and sudorific. Dose, grt. x. to xxx. Externally, it is used as a stimulant and discutient.

PETRO-OCIP'ITAL, *Petro-occipitalis*. Belonging to the petrous portion of the temporal bone and to the occipital bone.

PETRO-OCIPITAL SUTURE is formed by the junction of the petrous portion of the temporal with the occipital bone. It is a deep groove separating the bones, which have between them a thin layer of cartilage.

PETRO-SALPIN'GO-PHARYNGEUS. A fleshy fasciculus, which extends from the sphenoid bone, from the petrous portion of the temporal bone, and from the Eustachian tube to the upper part of the pharynx. See Constrictor.

PETRO-SALPINGO-STAPHYLINUS, *Circumflexus*—*p. Salpingo-staphylinus*, *Levator palati*—*p. Staphylin*, *Levator palati*.

PETRO-SPHENOID'AL, *Petro-sphenoida'lis*. Belonging to the petrous portion of the temporal bone and to the sphenoid bone.

PETRO-SPHENOIDAL SUTURE;—a name given to the small suture which is formed by the anterior edge of the petrous portion of the temporal bone and the posterior edge of the sphenoid.

PETROSAL, Petrous.

PETROSELINUM, *Apium petroselinum*—*p. Macedonicum*, *Bubon Macedonicum*.

PETROSUM, OS, see Temporal.

PETROUS, *Petræ'us*, *Petræ'sal*, (F.) *Pétréus*, *Pierreux*, from *petra*, and *petros*, 'a stone.' Resembling stone; having the hardness of stone. Relating or belonging to the petrous portion of the temporal bone.

PETROUS GAN'GLION, *Petræ'sal Ganglion*, *Ganglion of Andersch*, is formed by the glossopharyngeal nerve, shortly after it emerges from the

jugular fossa. From this ganglion, M. Jacobson of Copenhagen traced an anastomosing loop through the cavity of the tympanum, which left branches there, and was thence reflected downwards to join the Vidian nerve.—See Otic Ganglion. This has been called the *Nerve of Jacobson*, *Tympanic branch of the glosso-pharyngeal*, and the *inoculation, Jacobson's Anastomosis*.

PETROUS PORTION OF TEMPORAL BONE, see Temporal Bone—p. Process, see Temporal Bone.

PETROUS SINUSES, *Petrosal Sinuses*, are two venous sinuses of the dura mater, connected with the petrous portion of the temporal bone, and distinguished on each side, into:—1. The *superior petrosal sinus*. It arises from the cavernous sinus; follows the upper edge of the petrous portion which affords it a gutter; passes into the great circumference of the tentorium cerebelli, and opens into the lateral sinus where the latter experiences a bend near the base of the pars petrosa. Its form is triangular. 2. The *inferior petrosal sinus* arises from the cavernous sinus at the same point as the last, with which it communicates at the moment of its origin. It descends between the inferior edge of the pars petrosa and the basilar process; and terminates in the lateral sinus, at the sinus or gulf of the internal jugular vein. It is triangular, and broader at its extremities than at the middle,

PETUM, *Nicotiana tabacum*.

PETUN, *Nicotiana tabacum*.

PEUCE, *Pinus sylvestris*.

PEUCEA BALSAMEA, see *Pinus abies*.

PEUCED'ANUM, from *πευκη*, 'the pine tree,' its leaves resembling those of the pine; *P. officinale* seu *alsaticum* seu *altissimum* seu *paniculatum* seu *ternatum*, *Selinum peuced'anum*, *Mar'athrum sylvest're*, *Marathrophyl'ium*, *Pinastel'ium*, *Poenic'ulum porci'num*, *Bonus Ge'nus*, *Hog's Fenel*, *Sulphurwort*, (F.) *Peucedan*, *Queue de Pourceau*, *Fenouil de Porc*. Ord. Umbelliferae. *Sex. Syst.* Pentandria Digynia. The root has a strong, fetid smell—somewhat resembling that of sulphureous solutions—and an acrid, unctuous, bitterish taste. It has been recommended as antihysterical and nervine.

PEUCEDANUM ALSATICUM, *Peucedanum*—p. *Altissimum*, *Peucedanum*—p. *Officinale*, *Peucedanum*—p. *Oroselinum*, *Athamanta aureoselinum*.

PEUCEDANUM PALUS'TRE, *Thysse'lium* seu *Seli'num palus'tre*, *Selinum sylvest're* seu *thysse'lium* seu *lactescens*, *Thysse'lium*, *Olanit'ium*, *Marsh sel'num*, a European plant, has been recommended as an anti-epileptic. In Courland, it has been a famous remedy in epilepsy.

PEUCEDANUM PANICULATUM, *Peucedanum*—p. *Pratense*, *P. silaus*.

PEUCEDANUM ST'LAUS, *P. pratense*, *Ses'eli pratense*, *Sium* seu *Ligusticium* seu *Cnid'ium oil'aus*, *Saxifraga vulgaris* seu *Anglica*, *Hippomar'athrum*, *Poenic'ulum er'aticum*, *English or Meadow Saxifrage*, (F.) *Seseli des prés*. The roots, leaves, and seeds have been recommended as aperients, diuretics, and carminatives. Rarely used.

PEUCEDANUM TERNATUM, *Peucedanum*.

PEUPLIER, *Populus*—p. *Baumier*, *Fagara octandra*—p. *de la Caroline*, *Fagara octandra*—p. *Franc*, *Populus*—p. *Noir*, *Peucedanum*.

PEXIS, Concretion.

PEY'ERI GLAND'ULÆ, *Pey'eri Plexus glandul'si*, *Glandula Pey'eria'na* seu *mucosa conminata* seu *agminata* seu *mucip'araracematim congesta* intestino'rum seu intestinal'es plexiformes seu *spu'ria*, *Corpuscula glandularum similia*, *Glandulæ in agmen congregatæ intestino'rum*, *Enteradenæ*, *Peyer's Glands*, *Ag'minated or aggregate glands*. Small glands or follicles,

situate beneath the villous coat of the intestines, called after Peyer, who well described them. They are clustered together, (F.) *Plaques de Peyer*, having a honey-comb appearance—*Plaques gaufrées*—and are extremely numerous. They occur most numerous in the lower portion of the ileum, and resemble, in their morphology and functions, the lymphatic ganglions. They are, evidently, concerned in the great function of lymphosis.

PEZA, *πεζα*, Doric and Arcadian for *πους*, 'a foot.' The foot: according to some, the sole of the foot; and to others, the astragalus. The ankle.

PEZI'ZA seu PEZE'ZA AURIC'ULA, from *πεζα*, 'the sole of the foot,' *Tremella auricula Juda*, *Auricula Juda*, *Auricula'ria sambuci* seu *sambucina*, *Eridia auricula Juda*, *Merulius auricula*, *Fungus sambucinus*, *Agaricus Auricula formæ*, *Jew's Ear*. Ord. Fungi. This fungus is said to be astringent; but has chiefly been used as a gargle in relaxed sore throat.

PFEFFERS or PFÄFFERS, MINERAL WATERS OF, *Thermæ Piperi'nae*, *Th. Juba'riae*. These springs, which are in the Swiss Alps, are simply thermal: temperature 100° of Fahrenheit.

PHACĒ, *Φα'cen*, *Phacus*, *φακη*, 'a grain of the lentil,' *Ervum lens*. Also, chloasma; the crystalline lens; and a flask for the application of heat, so called owing to its flattened lens-like shape.

PHACELLUS, *Fasciculus*.

PHACELUS, *Fasciculus*.

PHACHYMENTIS, *Phachomyenitis*.

PHACI, *Ephelides*.

PHAC'ICUS, *Phaco'tus*, *Phaco'des*; from *φακη*, 'lens,' *Lens-shaped*, *lenticular*.

PHACI'TIS, *Phacoid'e'tis*, *Lent'i'tis*, *Crystall'i'tis*, *Cristalloïdi'tis*, *Glen'i'tis*, from *φακος*, 'a lens,' and *itis*, a termination denoting inflammation. Inflammation of the crystalline lens [?]. Inflammation of the capsule of the lens is termed *Capsul'i'tis*, *Capsulit'is*, and *Crystallino-capsulit'is*, *PHACO'DES*, *Phacicus*.

PHACHYMENTIS, *Phakomeni'tis*, *Phachymeni'tis*, *Inflammat'io capsulæ lentis*; from *φακος*, 'a lens,' *φαιν*, 'a membrane,' and *itis*, denoting inflammation. Inflammation of the capsule of the crystalline.

PHACODEITIS, *Phacitis*.

PHACOMALACIA, *Cataract*, soft.

PHAC'OPIS, *Phaco'tus*; from *φακη*, 'lens,' and *κορις*, 'a knife.' *Scalpel'ium lenticular'ræ*. A lenticular-shaped knife.

PHACOPS'IS, from *φακη*, 'lens,' and *οψις*, 'vision.' The state of one who has chloasma or freckles on the face.

PHACOPTIS'ANA, *Phac'ium*; from *φακη*, 'lens,' and *πιναν*, 'drink.' A drink of lentils. A poultice of lentils.

PHACOS, *Ervum lens*.

PHACOSCOTOMA, *Cataract*.

PHACOTUS, *Phacicus*, *Phacopis*.

PHACUS, *Phace*.

PHÆNIGMUS, *Phenigmus*.

PHENOMENOLOGIA, *Semeiology*.

PHÆNOMENON, *Phenomenon*.

PHAGÆNA, *Boulimia*.

PHAGEDÆNA, *Boulimia*, *Phagedenic ulcer* p. *Sloughing*, of the mouth, *Cancer aquaticus*—p. *Gangrenosa*, *Hospital gangrene*.

PHAGEDENIC, *Phagedæ'nicus*, *Depas'cens*, (F.) *Phagédénique*, *Rongeant*, from *φαγω*, 'I eat.' *PHAGEDENIC ULCER*, *Ulcus*, *Phagedæ'na*, *Ulcus depas'cens* seu *obambulans* seu *ferit'um*, (F.) *Ulcère phagédénique ou rongeante*, is one which rapidly eats and corrodes the neighbouring parts. *Phagedenic*, in *Materia Medica*, is a substance

used for destroying fungous granulations in ulcers, &c.

PHAKOHYMENITIS, Phacohymenitis.

PHALACRA, Calvities.

PHALACROMA, Calvities.

PHALACROSIS, Calvities.

PHALACROTIS, Alopecia.

PHALANA PINGUINALIS, see Eetozoa.

PHALATA. A barbarous term, employed by Basil Valentine to designate a universal medicine or panacea.

PHALAN'GAL, *Phalan'gian*, *Phalangia'mus*, from *φάλαγξ*, *phalaggyos*, 'a phalanx.' Having relation to a phalanx.

PHALANGARTHRI'TIS, from *φάλαγξ*, *phalaggyos*, 'a phalanx,' *arthron*, 'a joint,' and *itis*, denoting inflammation. Gout in the fingers. Inflammation of the joints of the fingers.

PHALANGE, Phalanx.

PHALANGES DES DOIGTS, see Phalanx—p. of the Fingers, see Phalanx—p. des Orteils, see Phalanx—p. of the Toes, see Phalanx.

PHALANGETTES, see Phalanx.

PHALANGETTIEN. Professor Chaussier uses this term for any thing relating to the third phalanx of the fingers or toes,—*Phalangien*, for what relates to the first,—and *Phalangiinien*, to the second.

PHALANGIAN, Phalangal.

PHALANGIEN, see Phalangettien.

PHALANGINES, see Phalanx.

PHALANGINIEN, see Phalangettien.

PHALANGIUM ESCULENTUM, Scilla esculenta.

PHALANGO'SIS, *Hispid'itas*, from *φάλαγξ*, *phalaggyos*, 'a phalanx,' 'a row of soldiers.' An affection of the eyelids in which the lashes are arranged in two or three rows. See *Trichiasis*. Also, dropping of the upper eyelid, from paralysis of the levator palpebre superioris muscle.

PHALANX, *Phalan'gē*, *Phalango'ma*. Same etymon. A name given to the small bones which form the fingers and toes, because placed alongside each other like a phalanx. They are distinguished into the *Phalanges of the fingers*, *Ossa seu Ag'mina seu A'cies seu Artic'uli seu Scutic'ula seu Scut'ulae seu Noli seu Interno'dia seu Scy'tal'ides seu Con'dyli Digito'rum Man'is*; and the *Phalanges of the Toes*, *Ossa seu Artic'uli seu Interno'dia Digito'rum Pedis*. I. *Phalanges of the fingers*, (F.) *Phalanges des doigts*. They are, in each hand, fourteen,—three to each finger, and two to the thumb; and are situate one above the other. They are distinguished, numerically, reckoning from the base towards the tip of the finger. 1. The first or metacarpal, *Procon'dyli*, (F.) *Premières phalanges ou Phalanges métacarpiennes*. 2. The second or middle phalanges, (F.) *Secondes phalanges ou Phalanges moyennes*, *Phalangines*, (Ch.) 3. The third phalanges, *Metacon'dyli*, *Phalang'es unguinae*, (F.) *Troisièmes phalanges ou Phalanges unguales ou unguifères ou unguinales*, *Phalanges* (Ch.), which terminate each finger, and support the nail. II. *Phalanges of the toes*, (F.) *Phalanges des Orteils*. They are the same in number as those of the fingers, but much smaller, and of a very different shape.

The phalanges of a finger or toe are united with each other; and the first with the metacarpal or metatarsal bone. They are developed from two points of ossification; one for the superior extremity, and the other for the body and inferior extremity.

PHAL'ARIS CANAREN'SIS, *P. ova'ta*, (*φάλαρος*—from *φαω*, 'I shine,'—'bright, shining,' in application to the seed.) Cultivated Canary Grass, (F.) *Alpiste*. Ord. Graminææ. Sex. Syst. Triandria Dignia. The seed of this grass, *Canary seed*, (F.) *Graines d'aspic*, afford a nou-

rishng meal. The ancients recommended them for allaying pains in the kidney and bladder.

PHALARIS ZIZANOIDES, *Andropogon muricatus*.

PHALLAL'GIA, *Phallody'n'ia*, from *φαλλος*, 'the male organ,' and *αλγος*, 'pain.' Pain in the male organ.

PHALLANCYLOSIS, Chordee.

PHALLITIS, *Priap'itis*, from *φαλλος*, 'the male organ,' and *itis*, a suffix denoting inflammation. Inflammation of the male organ.

PHALLOCAMPSIS, Chordee.

PHALLOCARCINO'MA, from *φαλλος*, 'the male organ,' and *καρκινωμα*, 'cancer.' Carcinoma or cancer of the penis.

PHALLODY'NIA, Phallalgia.

PHALLON'CUS, from *φαλλος*, 'the male organ,' and *ογκος*, 'swelling.' Tumefaction of the male organ.

PHALLOPSOPH'IA, from *φαλλος*, 'the male organ,' and *ψοφος*, 'a noise.' Discharge of wind from the male urethra.

PHALLORRHAG'IA, from *φαλλος*, 'the male organ,' and *ρρηγμα*, 'I break out.' Hemorrhage from the male urethra. Stomatosis.

PHALLORRHOEA, Gonorrhoea.

PHALLORRHOIS'CHESIS, from *φαλλος*, 'the male organ,' *ρροη*, 'discharge,' and *σχησις*, 'to withhold.' Sudden arrest of a gonorrhoeal discharge.

PHALLUS, Penis—p. Esculentus, *Morchella esculenta*.

PHANEROG'AMOUS, *Phanerogam'ic*, *Phanerogam'icus*, (F.) *Phanérogam'ic*, from *φανeros*, (*φαω*, 'I shine,') 'evident,' and *γαμος*, 'marriage.' An epithet used by botanists for plants whose sexual organs are apparent. It is, consequently, the opposite to *cryptogamous*.

PHANTASIA, Delirium, Imagination.

PHANTASM, *Phantas'ma*, *Phanto'ma*, *Phas'ma*, from *φανταζω*, (*φαίνω*, 'I bring to light,') 'I make appear.' A phantom, false appearance, apparition. Hallucination. Derangement of the sense of sight or of the intellectual faculties, causing objects to be perceived, which are not before the eyes.

PHANTASMA, Pseudoblepsia.

PHANTASMATICUS, Phantasticus.

PHANTASMATOSCOPIA, Metamorphopsia.

PHANTASMOSCOPIA, Metamorphopsia.

PHANTASTICUS, Inebriant. Creating phantasms—*phantasmat'icus*.

PHANTOM, Fantom—p. Tumour, see Fantom.

PHANTOMA, Phantasm—p. Obstetricium, Fantom.

PHARMACEIA, Poisoning

PHARMACEUM, Apotheca.

PHARMACEUS, Apothecary.

PHARMACEUTA, Apothecary.

PHARMACEUTIC, *Pharmaceut'icus*, *Pharmaceut'ical*, *Pharmaceut'ialis*, (F.) *Pharmaceutique*, from *φαρμακον*, 'a medicine,' *κατα*, 'beneath,' *γραφη*, 'a writing,' and *λογος*, 'a description.' The art of writing prescriptions.—J. J. Plenck.

PHARMACEUTICE, Pharmacy.

PHARMACEUTIST, Apothecary.

PHARMACIEN, Apothecary.

PHARMACOCATAGRAPHOLOG'IA, *Catagrapholog'ia*, *Ar's for'mulas concinnan'di*, from *φαρμακον*, 'a medicine,' *κατα*, 'beneath,' *γραφη*, 'a writing,' and *λογος*, 'a description.' The art of writing prescriptions.—J. J. Plenck.

PHARMACOCHEMYIA, Chymistry, pharmaceutical.

PHARMACODYNAM'ICS, from *φαρμακον*, 'a medicine,' and *δυναμις*, 'power.' *Pharmacodynami'cæ*, *Dynam'icæ Materia Medica*, *Pharmacodynam'ica doctri'na*, *Pharmacodynamiolog'ia*. A division of pharmacology, which considers the effects and uses of medicines.

PHARMACODYNAMIOLOGIA, *Pharmacodynamies*.

PHARMACOGNOS'IA, *Accognos'ia*, *Acognos'ia*, from *φάρμακον*, 'a medicine,' and *γνῶσις*, 'knowledge.' A division of pharmacology, which treats of simples, or unprepared medicines.

PHARMACOLOGIA, *Materia medica*.

PHARMACOMANIA, from *φάρμακον*, 'a medicine,' and *μανία*, 'mania.' One who has a mania for prescribing or for taking medicine.

PHARMACON, Poison.

PHARMACOPŒ'IA, *Pharmacopœ'a*, *Codex medicamentarius*, (F.) *Pharmacopée*, from *φάρμακον*, 'a medicine,' and *ποιέω*, 'I make.' A work, containing a collection of formulæ for the preparation, &c., of medicines: generally published by authority. Formerly, a chemical laboratory, *Laboratorium chymicum seu pharmacuticum*.

PHARMACOPŒUS, Apothecary, Druggist.

PHARMACOPOLA, Apothecary, Druggist.

PHARMACOPOLE, Apothecary, Druggist.

PHARMACOPOLI'UM, *Apotheca*, *Officina*, *Taberna Medica*. The shop of the druggist or apothecary, (F.) *Apothecairerie*, from *φάρμακον*, 'a medicine,' and *πωλέω*, 'I sell.'

PHARMACOPOS'IA, from *φάρμακον*, 'a medicine,' and *ποσις*, 'a potion.' A liquid medicine, especially one that is cathartic.

PHARMACO'SA MEDICAMEN'TA, from *φάρμακον*, 'a poison.' Medicines that contain poisonous substances. — Fallopius.

PHARMACOTHE'CA, *φαρμακοθήκη*, from *φάρμακον*, 'a medicine,' and *τίθημι*, 'I place.' A case for medicines. A medicine chest.

PHARMACOTHERAPI'A, from *φάρμακον*, and *θεραπεία*, 'treatment.' Cure of diseases by remedies.

PHARMACTER, Apothecary.

PHARMACUM, Medicament, Poison — p. *Ægyptiacum*, *Ægyptiacum* — p. *Anatorium*, *Philitar*.

PHARMACUM AD AURES, *Ægyptium Medicamentum ad Aures*. A name given to a medicine, composed of bitter almonds, white pepper, saffron, myrrh, opium, frankincense, castor, verjuice, sulphate of iron, vinegar, pomegranate-bark, unguentum nardinum, &c. It was believed to be useful for cleansing fetid ulcers in the ears.

PHARMACURGICUS, Apothecary.

PHARMACURGIUS, Apothecary.

PHARMACY, *Pharmacia*, *Pharmaceuticē*, *Pharmaz'ia*, from *φάρμακον*, 'a medicine.' The art which teaches the knowledge, choice, preservation, preparation, and combination of medicines. It was formerly divided into *Galenical* and *Chymical*. The former, called after Galen, embraced the operations undertaken on medicines without chymistry: — the latter, *Chymical Pharmacy* or *Pharmaceutic Chymistry*, *Pharmacochymia*, comprised the preparation of medicines, founded on the chymical action of their principles.

PHARMAXIS, Pharmacy.

PHARNA'CEUM LINEA'RE, called after Pharnaces, King of Pontus. A South African plant, *Nat. Ord.* Paronychiaceæ, employed at the Cape in pulmonary affections. It is generally given in infusion.

PHARUS, Pharynx.

PHARYGATHRON, Pharynx.

PHARYGETHRUM, Pharynx.

PHARYNGE'UM (SAL.), from *pharynx*, *pharyngis*. A compound of cream of tartar, nitrate of potassa, burnt alum, and distilled vinegar. It was formerly employed, united with honey and plantain-water, to make gargles, which were advised in quinsy.

PHARYNGĒ, Pharyngeal.

PHARYNGE'AL, *Pharyngæus*, (F.) *Pharyngé*,

Pharyngien. Same etymon. That which relates to the pharynx. A name given to the muscles, vessels, and nerves of the pharynx.

PHARYNGEAL ARTERIES. These are two in number. The one — *superior* or *pterygo-palatine*, *Ramus Pharyngæus superior* of Haller — arises from the internal maxillary at the base of the zygomatic fossa, passes through the pterygo-palatine canal, and proceeds to be distributed to the pharynx and the corresponding part of the Eustachian tube. The other — the *inferior* — *Pharyngæa ascendens* of Haller — arises from the external carotid, on a level with the facial. It ascends along the lateral and posterior part of the pharynx, and divides into two branches; the one — the *pharyngeal*, properly so called — is distributed to the pharynx; the other — the *meningeal* — gives its branches particularly to the dura mater.

PHARYNGEAL NERVE. A branch sent by the pneumogastric to the pharynx. It descends behind the internal carotid; communicates with the glosso-pharyngeal; and divides, opposite the middle constrictor of the pharynx, into a great number of filaments, which anastomose, in a multitude of different directions, with filaments of the glosso-pharyngeal nerve, of the superior laryngeal branch, and of the first cervical ganglion. These different nervous filaments united constitute the *pharyngeal plexus*, the irregular meshes of which send numerous ramifications to the pharynx.

PHARYNGEAL VOICE, see Voice.

PHARYNGEMPHRAX'IS, from *φάρυγξ*, *φάρυγος*, 'the pharynx,' and *ἐμπραξις*, 'obstruction.' Obstruction of the pharynx.

PHARYNGETHRON, Pharynx.

PHARYNGEURYSMA, Pharyngocœle.

PHARYNGICUS, Pharyngeal.

PHARYNGIEN, Pharyngeal.

PHARYNGITE, Cynanche pharyngea — p. *Tonsillaire*, *Cynanche tonsillaris*.

PHARYNGITIS, Cynanche pharyngea.

PHARYNGITIS APOSTEMATO'SA. Abscess of the pharynx.

PHARYNGI'TIS, DIPHTHERIT'IC, *Angina pseudo-membranacea* seu *membranacea* seu *pellucida* seu *plastica* seu *diphtheritica* seu *maligna* seu *pseudo-membranosa* seu *pestilentialis*, *Pharyngocacē*, *Pseudo-membranous Inflammation of the Throat*, (F.) *Angine couenneuse ou gutturale couenneuse* ou *pseudo-membraneuse* ou *plastique* ou *diphthéritique*. From *pharynx*, and *itis*, denoting inflammation. Diphtheritic inflammation of the pharynx. See Cynanche maligna.

PHARYNGI'TIS, FOLLIC'ULAR, *Follicular Inflammation of the Pharynx*, *Angina folliculosa* of the Pharynx, *Acne of the Throat*. Inflammation and enlargement of the follicles of the pharynx, occasionally extending to the larynx. Sometimes termed *Olergyman's Sorethroat*, *Tubercles of the Larynx and Fauces*, and *Tubercular Sorethroat*.

PHARYNGI'TIS, GANGRENOUS, Cynanche maligna.

PHARYNGOCACE, Pharyngitis, diphtheritic.

PHARYNGOCE'LE, *Pharyngæury'sma*, *Di-verticulum* seu *Hernia* seu *Prolapsus* (*Esoph'agi* seu *pharyng'is*, *Esoph'agus succenturiatus*, from *φάρυγξ*, *φάρυγος*, 'the pharynx,' and *κλήη*, 'a tumour.' An enlargement at the upper part of the gullet, in which the food sometimes rests in considerable quantity.

PHARYNGODYN'IA, from *φάρυγξ*, *φάρυγος*, 'the pharynx,' and *ὄδυν*, 'pain.' Pain in the pharynx.

PHARYNGOGLOSS'AL, *Glossopharyngeal*, *Pharyngoglossia'nus*. That which belongs to the pharynx and tongue.

PHARYNGOGLSSAL NERVE, *Glossopharyngeal*

Nerve, Lingualis lateralis, (F.) Ners Glossopharyngeus, Pharyngoglossus—a portion of the eighth pair—arises from the superior and lateral parts of the spinal marrow, between the facial and pneumogastric nerves, by several filaments, which soon unite into a single cord, that passes out of the cranium by the foramen lacerum anterius; goes downwards and forwards, and gains the posterior and inferior part of the tongue. Immediately after its exit from the cranium, it gives a branch to the meatus auditorius externus, and receives a filament from the facial, and another from the pneumogastric nerve. It then furnishes two other filaments, which descend upon the carotid artery, and unite at the bottom of the neck with branches from the cervical ganglions and cardiac nerves, and give two filaments to the stylo-pharyngeus muscle. The glossopharyngeus, after sending a twig or two to the tonsil, to the upper part of the pharynx and membrane of the epiglottis, divides into many branches which run partly to the margin, and partly to the middle of the root of the tongue; supplying, especially, the papillae majores, and the parts in their neighbourhood.

By some, this nerve is regarded as the controller of the associated motions of the tongue and pharynx. By others, it is esteemed to be the special nerve of gustation.

PHARYNGOG'RAPHY, Pharyngograp'hia, from *φαρυγξ, φαρυγγος*, 'the pharynx,' and *γραφειν*, 'a description.' An anatomical description of the pharynx.

PHARYNGOLOG'Y, Pharyngolog'ia, from *φαρυγξ, φαρυγγος*, 'the pharynx,' and *λογος*, 'a discourse.' The part of anatomy which treats of the pharynx.

PHARYNGOLYSIS, Pharyngoplegia.

PHARYNGOPARALYSIS, Pharyngoplegia.

PHARYNGOPLE'GIA, from *φαρυγξ, φαρυγγος*, 'the pharynx,' and *πλησις*, 'I strike.' *Dysphagia paralytica* seu *torpida* seu *atonia*, *Angina paralytica*, *Asthenia Deglutitionis*, *Pharyngoplegia*, *Pharyngoparaly'sis*, *Paraly'sis Oesophagi*, *Gula Imbecillitas*. Paralysis of the pharynx or oesophagus: commonly a symptom of general paralysis, or of approaching dissolution.

PHARYNGORRHAG'IA; from *φαρυγξ, φαρυγγος*, 'the pharynx,' and *ρρηγν*, 'rupture.' Haemorrhage from the vessels of the pharynx.

PHARYNGOSPASMUS, from *φαρυγξ, φαρυγγος*, 'the pharynx,' and *σπασμος*, 'spasm.' Spasm or cramp of the pharynx.

PHARYNGOSTAPHYLINUS, Palato-pharyngeus.

PHARYNGOTOME, Pharyngotomia, from *φαρυγξ, φαρυγγος*, 'the pharynx,' and *τομω*, 'I cut.' An instrument, invented by J. L. Petit, used to scarify inflamed tonsils, and to open abscesses, which form in the parietes of the pharynx. It is composed of a long, narrow blade with a lancet point, enclosed in a sheath or canula of silver slightly curved. This instrument is carried to the part to be cut, and the point of the blade is made to pass out of the sheath, by pressing on a button at the end of the pharyngotome. A spring causes the blade to re-enter as soon as the pressure is removed.

PHARYNGOTOMY, Pharyngotomia; same etymon. Some authors have used this word synonymously with oesophagotomy. It means, also, scarification of the tonsils, or an incision made for opening abscesses there, &c.

PHARYNX, φαρυγξ, Pharynx, φαρυξ, Pharus, Gula, Læmas, Fauces, Pharyg'ethron, Pharyng'ethron, Pharyng'ethron, Pharyng'ethron, Isthmos, Isth'mion, Ingu'vies, Principium Gulae, Os posterum, Gurgies, (F.) Arrière-bouche, Gasier, Gorge. A species of musculo-membranous, symmetrical

canal, on the median line, irregularly funnel-shaped, and situate between the base of the cranium and the oesophagus, in front of the vertebral column. It is very narrow above; but dilates in the middle, and again contracts below, at its junction with the oesophagus. Into the anterior paries of the pharynx open, successively, from above to below,—the posterior orifices of the nasal fossae; the Eustachian tubes; the posterior aperture of the mouth, and the top of the larynx. The pharynx is formed, externally, of a muscular coat, and, internally, of a mucous membrane, which is continuous, above, with the Schneiderian membrane; and in the middle, with that of the mouth below, with that of the oesophagus; and, at the sides, with that of the Eustachian tubes. This membrane has no villi, and presents only a few inequalities, which are owing to the presence of muciparous follicles. The muscular coat of the pharynx is composed of the constrictor muscles, stylopharyngei, and pharyngostaphylini. The vessels and nerves are called *pharyngeal*. The pharynx serves as a common origin for the digestive and respiratory passages. It gives passage to the air, during respiration; and to the food at the time of deglutition.

PHARYNX, FOLLICULAR INFLAMMATION OF THE, Pharyngitis, follicular.

PHARYX, Pharynx.

PHASELUS, Phaseolus

PHASE'OLUS, φασηλος, φασηλος, φασιολος, P. Vulgaris, Phaseolus, Phase'us, Fase'us, Faseolus, Bona, Buona, Faba; Kidney Bean; French Bean; (F.) Fesole, Haricot. Ord. Leguminosæ. Ser. Syst. Diadelphia Decandria. When young and well boiled, this is easy of digestion, and delicately flavoured. It is less flatulent than the pea.

PHASEOLUS CRETICUS, Cajan, Cayan. A decoction of the leaves, according to Ray, restrains the hemorrhoidal flux when excessive.

PHASEOLUS VULGARIS, Phaseolus.

PHASIA'NUS GALLUS, so called from the river, *Phasis*, in Colchis, near the Black Sea: (F.) *Cog*. The parts of generation of the cock, dried and pulverized, were formerly regarded as proper for increasing the quantity of semen. The fat was used as emollient and resolvent; the brain, in diarrhoea; the gall, in freckles and diseases of the eyes. These phantasies are now abandoned.

PHASIOLUS, Phaseolus.

PHASMA, Phantasm.

PHATNE, Alveolus.

PHATNION, Alveolus.

PHATNORRHAG'IA, Stomatorrhag'ia alveolarum, from *φατιον*, 'an alveolus,' and *ρρηγν*, 'I break forth.' Hemorrhage from the alveoli.

PHAUSIN'GES, φανσιγες, plural of *φανσιγξ*, from *φαιω*, 'I roast.' Spots or blisters on the legs, produced by heat; and, in a more extensive signification, by any other cause. Also, a pustule. — *Foësius*.

PHELLANDRE, Phellandrium aquaticum.

PHELLAN'DRIUM AQUAT'ICUM, Phellandrium, Enan'the phellandrium seu aquat'ica, Ligusticium phellandrium, Fœniculum aquaticum, Cicutaria aquatica, Water Fennel, Fine-leaved Water Hemlock, (F.) Phellandre, Fenouil d'eau, Ciguë d'eau, Millefeuille aquatique. Ord. Umbelliferae. Ser. Syst. Pentandria Digynia. This plant is stimulant and narcotic. It has been used in intermittents and dyspepsia. It has, also, been given in phthisis pulmonalis, and in cases of scrofulous atonic ulcers.

PHENE, Benzole.

PHENIG'MUS, Phœnig'mus, Phœnig'mus, from *φαινει*, 'red.' *Red Jaundice*. A cutaneous affection, consisting of redness diffused over the skin, without fever. *Sauvages* makes it a genus of his

order *Ictericæ*. Also, the act of irritating a part by a stimulating application.

PHÉNOMÈNE, Phenomenon.

PHENOM'ENON, *Phénom'ènon*, (F.) *Phénomène*, from *φαίνωμαι*, 'I appear.' An extraordinary and unexpected event. In *Medicine*, it means any change, appreciable by our senses, which supervenes in an organ or function:—as the *phenomena*, of the circulation, respiration, &c. It is used in pathology, synonymously with symptom.

PHENYLE, **HYDRURET OF**, Benzole.

PHEUGOPHOBIA, Hydrophobia.

PHEUGYDRON, Hydrophobia.

PHI'ALA, A *Phial* or *Vial*, (F.) *Firole*, *φιάλη*, *φιάλη*, 'a vessel with a broad bottom.' A small bottle, in which apothecaries are in the habit of sending out their mixtures, draughts, &c.: hence called *Medicine Vials*, (F.) *Fioles à Médecine*. They are usually sold in grosses, and assorted into vials of oz. 8; oz. 4; oz. 3; oz. 2; oz. 1½; oz. 1; and oz. ½.

PHILADYNAMICA, Debilitants.

PHILANTHROPUS, *Galium aparine*.

PHILETRON, Philtr.

PHILETRUM, *Lacuna labii superioris*.

PHILA'TER, *Philia'tros*, *φιλάτρος*, from *φίλω*, 'I love,' and *τάρην*, 'medicine.' An amateur of medicine. One who studies medicine; *Studio'sus medicinæ*.

PHILOBIO'SIS, (F.) *Philobiosie*: from *φίλω*, 'I love,' and *βίος*, 'life.' The love of life.

PHILÉ'NIA, from *φίλω*, 'I love,' and *οίνος*, 'wine.' Love of wine. Addiction to wine.

PHILOGYNIA, *Muliebrositas*.

PHILONIUM LONDINENSE, *Confectio opii*.

PHILOPATRIDALGIA, *Nostalgia*.

PHILOPATRIDOMANIA, *Nostalgia*.

PHILOSOPHER'S STONE, see *Alchymy*.

PHILOSOPHIA CORPORIS VIVI, *Physiology*—p. *Hermetica*, *Alchymy*—p. *per Ignem*, *Chymistry*.

PHILTER, from *φίλω*, 'I love.' *Philtrum*, *Phil'tron*, *Phile'tron*, *Phar'macum* seu *Poc'ulum amato'rium*, *Amato'rium venefic'ium*, (F.) *Philtre*. A medicine or drink considered capable of causing love.

PHILTRUM, *Lacuna labii superioris*.

PHILYRA, *Tilia*.

PHIMA, *Phyma*.

PHIMOS, *Capistrum*.

PHIMO'SICUS, *Relating to Phimosia*.

Sauvages calls *Ischu'ria phimo'sica* seu *urethra'tis à phimo'si*, the retention of urine which depends upon phimosia.

PHIMO'SIS, *Phymo'sis*, *Capistra'tio*, *Epago'gium*, *Ligatu'ra glandis*, *Strictu'ra præpu'tii*, *Reductio præpu'tii impedi'ta*, from *φίμω*, 'I bind tight.' A disease which consists in a preternatural narrowness of the opening of the prepuce, so that it cannot be carried behind the corona glandis. This affection is most commonly congenital. Sometimes it is accidental, and dependent upon tumefaction of the glans, or of the prepuce, as occurs in syphilis. To remedy this state, the prepuce may be divided by means of a bistoury and director; or circumcision may be practised.

PHIMOSIS ADNATA seu **CONGENITA** seu **PUERILIS**, *Congenital phimosia*—p. *Circumligata*, *Paraphimosia*—p. *Congenita*, *P. adnata*.

PHIMO'SIS ŒDEMATO'DES, *Hydrophimo'sis*. *Phimosia* with œdematous swelling of the prepuce.

PHIMOSIS PUERILIS, *P. adnata*.

PHLASMA, *Contusion*.

PHLEBARTERIODIALYSIS, see *Aneurism*.

PHLEBECTA'SIA, *Phlebec'tasis*, from *φλεψ*, *φλεβος*, 'a vein,' and *εκτασις*, 'dilatation.' Dila-

tion of a vein, or of a portion of a vein.—*Alibert*.

PHLÉBECTIARCTIE. A mongrel term, from *φλεψ*, *φλεβος*, 'a vein,' and *arcto*, 'I contract.' *Contraction of a vein*.

PHLEBES, (*Arteries*). See *Artery*.

PHLEBEURYSMA, *Varix*.

PHLEB'ION, diminutive of *φλεψ*, *φλεβος*, 'a vein.' A vein.—*Hippocrates*. A small vein,—*Ve'nula*.

PHLEBIS'MUS, from *φλεψ*, *φλεβος*, 'a vein.' A term employed by Dr. Marshall Hall for the turgescence of veins, produced by impeded return of the blood. See *Trachelismus*.

PHLEB'I'TIS, *Inflamma'tio venarum*, (F.) *Phlébite*, *Inflammation des veines*, from *φλεψ*, *φλεβος*, 'a vein,' and *itis*, the termination for inflammation. Inflammation of the inner membrane of a vein, *Endo-phlebi'tis*, which sometimes follows blood-letting, and extends from the small wound made in that operation to the neighbouring parts of the venous system. The symptoms are, first, inflammation in the punctured part; and, afterwards, a knotty, tense, painful cord, following the direction of the vessel, and accompanied with more or less fever, according to the extent of the inflammation, the nervous symptoms, &c. *Phlebitis*, also, sometimes supervenes on the ligature or excision of varices; after amputation, gun-shot wounds, delivery (*Uterine Phlebitis*, *Locho-metro-phlebi'tis*, *Metro-phlebi'tis puerperalis*, *Phlebi'tis uteri'na*, *Metri'tis veno'sa*), the ligature of the umbilical cord, &c. It may terminate by resolution, suppuration, ulceration, or gangrene.

PHLEBITIS, *CRURAL*, *Phlegmatia dolens*—p. *Uterina*, see *Peritonitis*, and *Phlebitis*.

PHLEBOG'RAPHY, *Phlebogra'phia*, from *φλεψ*, *φλεβος*, 'a vein,' and *γραφη*, 'a description.' A description of the veins.

PHLEB'OLITE, *Phlebol'ithus*, *Vein stone*, from *φλεψ*, *φλεβος*, 'a vein,' and *λίθος*, 'a stone.' A loose concretion, varying in size from that of a currant to that of a pea, occasionally found in the veins. It would seem to be formed in the blood.

PHLEBOLITHUS, *Phlebolite*.

PHLEBOLOG'Y, *Phlebolog'ia*, from *φλεψ*, *φλεβος*, 'a vein,' and *λογος*, 'a discourse.' The part of anatomy which treats of the veins.

PHLEBOPA'LIA, from *φλεψ*, *φλεβος*, 'a vein,' and *παλλω*, 'I palpitate.' With the ancients, *pulsation*, which they conceived to be produced by the veins. See *Pulsus venarum*.

PHLEBOPHTHALMOTOM'IA, *Ophthalmophlebotom'ia*, from *φλεψ*, *φλεβος*, 'a vein,' *οφθαλμος*, 'the eye,' and *τεμνω*, 'I cut.' Blood-letting in the eye.

PHLEBOPLEROSIS ECPHRACTICA, *Inflammation*.

PHLEBORRHAG'IA, from *φλεψ*, *φλεβος*, 'a vein,' and *ρηννυμι*, 'I break out.' Rupture of the veins; and, in a more extensive sense, venous hemorrhage.

PHLEBORRHEX'IS, from *φλεψ*, *φλεβος*, 'a vein,' and *ρρηξις*, 'rupture.' Rupture of a vein. Also, hemorrhage from a vein, *Phleborrhag'ia*.

PHLEBOSTENO'SIS, from *φλεψ*, *φλεβος*, 'a vein,' and *στενωσις*, 'contraction.' Constriction or contraction of a vein.

PHLEBOT'OMIST, from *φλεψ*, *φλεβος*, 'a vein,' and *τεμνω*, 'I cut.' A bleeder. One who practises particularly the art of phlebotomy. See *Blood-letting*.

PHLEBOTOMUM, *Fleam*.

PHLEBOTOMY, see *Blood-letting*.

PHLEBS, *Vein*.

PHLEDONIA, *Delirium*.

PHLEGM, *Phlegma*, *φlegma*, *Pitui'ta*, *Pit'uite*,

Mucus, Mother, (F.) Phlegme, Pituite. One of the four natural humours of the ancients, which, according to them, was cold and moist, as atrabilis was cold and dry. It predominated, especially, in winter.

Pituita was afterwards applied to every aqueous or excrementitious humour, such as the saliva, nasal and intestinal mucus, serum, &c. The terms *phlegm* and *pituita* are no longer used in physiology,—the different humours having received particular names; but the vulgar still use *phlegm* to designate a stringy mucus, expectorated, or rejected by vomiting. The ancient chymists gave the name 'phlegm' to aqueous, insipid, and inodorous products obtained by subjecting moist vegetable matters to the action of heat.

PHLEGMA NARIUM CRASSUM, Nasal mucus—p. Vitrioli, Sulphuric acid (very dilute.)

PHLEGMAGOGUE, *Phlegmago'gus*; from φλεγμα, 'phlegm,' and αγω, 'I expel.' A medicine, formerly believed capable of 'cutting' or removing phlegm.

PHLEGMAPYRIA, Fever, adeno-meningeal.

PHLEGMASIA, Inflammation, Phlegmon—p. Adenosa, Adenitis—p. Alba dolens puerperarum, Ph. dolens—p. Cellularis, Induration of the cellular tissue—p. Glandulosa, Adenitis—p. Myoica, Myositis—p. of the Peritoneum, Peritonitis.

PHLEGMA'TIA, from φλεγμα, φλεγματος, 'phlegm.' Edema, Anasarca, Hydrops.

PHLEGMA'TIA DOLENS, *P. lac'tea seu dolens seu alba dolens seu Leucophlegmasia dolens seu Spargano'sia seu Scela'l'gia seu (Edema puerperarum, Scelon'eus, Anasar'ca sero'sa, Ecphy'ma oedemat'icum, Echymo'na lymphat'icum, Is'chias a spargano'si, Oede'ma lacteum, Infarc'tus lac'tei extremit'um, Metas'tasis lactis, Hyster'al'gia galact'ica, Hydrophlegmasia testis cellul'osi, Cru'al Phlebi'tis, White leg, Swelled leg, Milk leg, White swelling of lying in women, (F.) Oedème des nouvelles accouchées, Oedème douloureux des femmes en couche, Dépot laiteux sur la cuisse, Oedème acif des nouvelles accouchées, Enflure ou Engorgement des jambes et des cuisses de la femme accouchée, E. laiteux des membres abdominaux.* It occurs, for the most part, in the second or third week after delivery:—is limited to the lower extremity, and chiefly to one side: exhibiting to the touch a feeling of numerous irregular prominences under the skin. It is hot, white, and unyielding; and is accompanied, sooner or later, with febrile excitement. After a few days, the heat, hardness, and sensibility diminish, and the limb remains oedematous for a longer or shorter period. The disease frequently, if not generally, consists in the obstruction of the iliac or other veins. Owing to the pressure of the gravid uterus—the flow of blood being obstructed—the serous part of it is thrown out into the cellular membrane of the limb. Sometimes the vein is found completely obliterated.

The treatment consists in the use of antiphlogistics:—the application of leeches or cupping-glasses near the groin, fomentations to the limb, &c.; and, when the active state has subsided, the use of a roller, applied gently at first, but, afterwards, gradually tightened.

PHLEGMATIA LACTEA, P. dolens.

PHLEGMATIE, see *Empresma*—p. Membranosa et parenchymatosa, *Empresma*.

PHLEGMATIC, *Phlegmat'icus*, from φλεγμα, 'phlegm.' Abounding in phlegm,—as the 'phlegmatic or lymphatic temperament.'

PHLEGMATOCHEMY, see *Chemistry*.

PHLEGMATOPYRA, Fever, adeno-meningeal.

PHLEGMATORRHAG'IA, *Phlegmatorrhœ'a, Catarrh*, from φλεγμα, φλεγματος, 'phlegm,' and

ρῶω, 'I flow.' Stahl and his partisans gave this name to a copious secretion, by the nostrils, of a limpid and thin mucus, without inflammation. It is the *Cory'za Phlegmatorrhag'ia* of Sauvages. See *Coryza*.

PHLEGMHYMENTIS, *Phlegmymentis*.

PHLEGMON, from φλεγω, 'I burn.' *Phlogo'sis phleg'monē, Phleg'monē, Phlegma'sia, Incendium, Inflammat'io, (Prov.) Beal.* Inflammation of the areolar texture accompanied with redness, circumscribed swelling, increased heat and pain; which is, at first, tensile and lancinating; afterwards, pulsatory and heavy. It is apt to terminate in suppuration. It requires the antiphlogistic treatment, until suppuration seems inevitable, when the suppuration must be encouraged as the next favourable termination to resolution.

PHLEGMON PÉRINÉPHRÉTIQUE, *Periphrenitis*—p. Peri-uterine, see *Peri-uterine*.

PHLEGMONÉ, Inflammation—p. Mammæ, Mastodynia apostematosa—p. Mastodynia, Mastodynia apostematosa—p. Parotidæa, Parotis—p. Peritonæi, Peritonitis—p. Parulis, Parulis—p. Musculorum, Myositis—p. Ventriculi, Gastritis.

PHLEGMONÉUX, Phlegmonous.

PHLEGMONODES, Phlegmonous.

PHLEGMONOID, Phlegmonous.

PHLEGMONOUS, *Phlegmono'us, Phleg'monoid, (F.) Phlegmonous, from φλεγμων, (φλεγω, 'I burn,') 'a phlegmon, and εἶδος, 'resemblance.'* That which relates or belongs to or resembles phlegmon.

PHLEGMONOUS INFLAMMATION is the active inflammation of the areolar membrane; in contradistinction to erysipelatous inflammation, or that of the skin.

PHLEGMOPYRA, Fever, adeno-meningeal.

PHLEGMORRHAGIA, *Blennorrhœa*.

PHLEGMORRHAGIE PULMONAIRE, *Bronchorrhœa*.

PHLEGMORRHŒA, *Blennorrhœa*.

PHLEGMYMENITIS, *Phlegmymenitis*, from φλεγμα, 'phlegm,' μῆνις, 'a membrane,' and itis, the termination for inflammation. Inflammation of a mucous membrane.

PHLEGMYMENTIS ENTERICA, see *Enteritis*.

PHLEPS, Vein.

PHLOGIA, Inflammation.

PHLOGICUS, Inflammatory.

PHLOGISTIC, Inflammatory.

PHLOGISTICI, *Empresma*.

PHLOGISTON, Hydrogen.

PHLOGMUS, *Verbasum thapsus*.

PHLOGO'DES, from φλογ, φλογος, (φλεγω, 'I burn,') 'flame,' and εἶδος, 'resemblance.' *Inflamed, Inflammat'us, Inflammat'rius*, Inflammatory. An epithet employed particularly to express the red colour of the face.

PHLOGOGUS, Antiphlogistic.

PHLOGOPYRA, *Synocha*.

PHLOGOSED, Inflamed.

PHLOGO'SIS, *Incendium, Inflammat'ion*, from φλογ, φλογος, 'flame.' Some writers use this word exclusively for external inflammation—phlegmon; others, for that which is superficial or erysipelatous.

PHLOGOT'ICA, *Inflammations*. The second order in the class *Hæmatica* of Good.

PHLOIOS, Cortex.

PHLOMUS, *Verbasum thapsus*.

PHLOOS, Cortex.

PHLORIDZINE, *Phloridzi'na, Phloridzin*, from φλοος, 'inner bark,' and ρίζα, 'root.' The bitter or active principle of the root and trunk of the apple, pear, cherry, and plum tree. It is deposited from the decoction on cooling; crystallizes in silky spicula of a dead white colour when in masses, or in long slender prisms or tables,

when crystallized with care. 1000 parts of water, at a temperature of from 32° to 71°, only dissolve about one part; but at from 70° to 212°, it is dissolved in all proportions. It is very soluble in pure alcohol, but only slightly so in ether; *s. g.* 1.429. It has been administered successfully in intermittents,—10 to 20 grains being generally required to arrest the paroxysm. It may be given either in the form of pill or powder.

PHLOUS, Cortex.

PHLOX, *Verbascom thapsus*.

PHLYCTÆNA, *Phlyctænia*, *Phlyctis*, *Phlyctæcion*, *Ampullæ*, *Holophlyctides*, *Phlyctides*, *Phlysis*, from *φλυζω*, 'I boil.' Tumours formed by the accumulation of a serous fluid under the epidermis.

PHLYCTÆNULA, see Hydatid.

PHLYCTENOID, *Phlyctenoides*, from *phlyctæna*, and *eidos*, 'resemblance.' Resembling phlyctæna. An epithet for a variety of herpes.

PHLYCTENULAR, same etymon as Phlyctæna. Of or belonging to phlyctænula,—as *phlyctenular ophthalmia*; inflammation of the eye, accompanied with phlyctænulæ on the cornea.

PHLYCTIDES, Phlyctæna.

PHLYCTIDIUM, same etymon. A circular or annular spot of inflammation of the corium, encircled by a red ring or zone, within which suppuration takes place. The genuine small-pox pustule, and that produced by the application of tartar emetic ointment, are examples of it.

PHLYCTIS, Phlyctæna.

PHLYSIS, *φλυσις*, same etymon. A cutaneous eruption. An ulcerative, subcutaneous tumour: flat, tense, glabrous: diffused, hot, throbbing; at length fluctuating, with an acrid ichor.—Good.

PHLYSIS, Phlyctæna.—*p.* Ecthyma, Ecthyma.—*p.* Impetigo, Impetigo.—*p.* Porrigo, Porrigo.—*p.* Scabies, Psora.

PHLYZACIA, Ecthyma.

PHLYZACION, Phlyctæna.

PHOBODIPSON, Hydrophobia.

PHOCOMELUS, from *φοκη*, 'a seal,' and *μελος*, 'a limb.' A monster, in which the limbs are shortened; the hands and feet appearing to exist alone, and to be inserted immediately on the trunk, as in the seals and the herbivorous cetacea.—*I. G. St. Hilaire*.

PHENICINE, Hæmatin.

PHENICISMUS, Rubæola.

PHENIGMUS, Phenigmus.—*p.* Petechialis, Purpura simplex.

PHENIX DACTYLIFERA, Date tree.—*p.* Exelsa, Date tree.

PHENODINE, Hæmatin.

PHONA'TION, *Phona'tio*, from *φωνη*, 'voice.' The physiology of the voice.

PHONE, Voice.

PHONICA, *φωνικα*, from *φωνη*, 'voice.' The first order of the class *Pneumatica*, in Good's Nosology. Defined: diseases affecting the vocal avenues:—the passage of the voice impeded, or its power suppressed or depraved.

PHONONOSI, *Phononosi*, from *φωνη*, 'the voice,' and *νοσος*, 'a disease.' *Phonopathi'a*, *Morbi vocis*. Diseases of the voice.

PHONOPATHIÆ, Phononosi.

PHORA, Gestation.

PHORBE, Pabulum.

PHORONOMIA, from *φωω*, 'I move,' and *νομος*, 'law.' The doctrine of muscular motion.

PHOROUS, from *φωω*, 'I carry.' A suffix denoting conveyance. Hence, *Galactophorous*, &c.

PHOS, Light.

PHOSPHAS, *Phosphate*. A salt formed by the union of phosphoric acid with a salifiable base.

PHOSPHAS CALCICUM STIBIATUM. Antimonial powder.—*p.* Calcis stibiatum, Antimonial powder.—*p.* Natrium, Soda, phosphate of.

PHOSPHATE, Phosphas.

PHOSPHAT'IC, *Phosphat'icus*, from *Phosphorus*. Relating to the phosphates; hence the *Phosphat'ic Diathesis* is the habit of body which favours the deposition of the phosphates from the urine.

PHOSPHÈNE, see Photopsia.

PHOSPHORE, Phosphorus.

PHOSPHORENÆSES. M. Baumes unites, under this generic name, diseases which he attributes to disordered phosphorization; i. e. to excess or deficiency of calcareous phosphates or its decomposition. Amongst these affections he ranks rachitis, osteo-malacia, and gout.

PHOSPHORES'CENCE, *Phosphorescentia*, *Luminosity*; from *phosphorus*. The properties possessed by certain bodies of being luminous in obscurity, or under particular circumstances. This phosphorescence has been occasionally witnessed in man, when suffering from some wasting disease, and near death.

PHOSPHORIC ACID, *Ac'idum phosphor'icum*, (F.) *Acide phosphorique*. Same etymon. A solid, colourless compound; highly sapid; very soluble in water, and vitrifiable by fire. It exists in bones combined with lime. It has been employed in medicine in the form of lemonade, in passive hemorrhage, typhus, phthisis, marasmus, &c., and in injection in cases of cancer of the uterus. It does not seem preferable to other acids. An *Acid'um phosphor'icum dilutum* is official in the London pharmacopœia.

PHOSPHORNECROSIS, see Phosphorus.

PHOSPHORURIA, Urine, phosphorescent.

PHOS'PHORUS, *Phos'phorus*, *Autophos'phorus*, *Ignis philosoph'icus*, *Lumen constans*. from *φως*, 'light,' and *φερω*, 'I carry.' (F.) *Phosphore*, 'A light carrier.' An elementary or undecomposed substance; not metallic; solid; transparent or semitransparent; colourless; brilliant; flexible; of a sensible garlic odour; very inflammable, and susceptible of combining with the oxygen of the atmosphere, at even the ordinary temperature. It does not exist pure in nature, but is found frequently united to other bodies. Its fumes in the manufacture of lucifer and Congreve matches have caused, it is said, necrosis of the lower jaw—*Phosphornecro'sis*, *phosphorus jaw disease*, *Lucifer-match-makers' Disease*.

Phosphorus has been used in medicine, dissolved in ether or suspended in the yolk of egg, as a stimulant; but the greatest precautions are necessary in its administration: hence it is rarely employed.

PHOSPHORUS JAW DISEASE, see Phosphorus.

PHOSPHURIA, Urine, phosphorescent. See Urine.

PHOTOCAMPSIS, Refraction.

PHOTOMA'NIA, from *φως*, *φωτος*, 'light,' and *mania*. Insane delirium induced by light.

PHOTOPHO'BIA, from *φως*, *φωτος*, 'light,' and *φοβος*, 'fear.' Aversion to light:—intolerance of light. Nyctalopia.

PHOTOPHOBICUS, *Lucifugus*.

PHOTOPHOBOPHTHALMIA, *Nyctalopia*.

PHOTOPHOBUS, *Lucifugus*.

PHOTOP'SIA, *Visus lu'cidus*, from *φως*, *φωτος*, 'light,' and *ωψ*, 'the eye.' A false perception of light, as of sparks, flashes of fire, &c., occasioned by certain conditions of the retina and brain, and by pressure on the eyeball. When pressure does not induce the luminous appearance, (F.) *Phosphène*, the existence of amaurosis may be inferred. See Metamorphopsia.

PHOTURIA, from *φως*, *φωτος*, 'light,' and *ουρον*, 'urine.' A luminous condition of the urine.

PHRAGMOS, *φραγμος*, 'a fence,' from *φρασσω*, 'I enclose.' A row of teeth.

PHRASIS, Voice, articulated.

PHRASUM, Marrubium.

PHREN, *φρην*, 'the mind.' Also, the diaphragm and scrobiculus cordis.

PHRENAUX'Ē, *Hypertroph'ia cer'ebr'i*, from *φρην*, 'the mind, the organ of mind,' and *αυξη*, 'increase.' Hypertrophy of the brain.

PHRENATROPH'IA, *Atroph'ia cer'ebr'i*, from *φρην*, 'the mind, the organ of mind,' *a*, priv., and *τροφη*, 'nourishment.' Wasting or atrophy of the brain.

PHRENES, Diaphragm.

PHRENESIA, Phrenitis.

PHRĒNĒSIE, Phrenitis.

PHRENESIS, Delirium, Phrenitis.

PHRENETIASIS, Delirium, Phrenitis.

PHRENISMUS, Phrenitis.

PHRENIC, Diaphragmatic.

PHRE'NICA, from *φρην*, 'the mind.' Diseases affecting the intellect; characterized by error, perversion, or debility of one or more of the mental faculties, (*Phrenopath'ia*.) The first order of the class *Neurotica* of Good. Also, remedies that affect the mental faculties—*Phren'ics*.—Pereira.

PHRENICS, see Phrenica.

PHRENICULA HYDROCEPHALICA, Hydrocephalus internus.

PHRENISMUS, Phrenitis.

PHRENTIS, *Phren'ies*, *Phrene'sia*, *Phren'et'iasis*, *Phrenis'mus*, *Phrenet'ismus*, *Cephal'itis*, *Encephal'itis*, *Cerebr'itis*, *Cephal'gia inflammato'ria*, *Karabitus* (Arab.), *Sibore*, *Sphecelis'mus cer'ebr'i*, *Empres'ma cephal'itis*, *Siri'asis*, *Seiri'asis*, *Siderat'io*, *Inflammat'io seu Œdema phren'ie*, *Phren'ey*, *Delirium*, *Brain Fever*, (F.) *Phrénésie*, *Cérébrile*, *Inflammation du cerveau et du cervelet*, from *φρην*, 'the mind,' and *itis*, denoting inflammation. A disease characterized by violent pyrexia, pain in the head, redness of the face and eyes, intolerance of light and sound, watchfulness, and delirium, either furious or muttering. Its causes are the same as those of other inflammations: sometimes, however, it is induced by a *Coup de soleil*; by inebriation; violent mental emotions, &c. Some authors have given the name *Phrenitis* and *Meningitis* to inflammation of the membranes of the brain; and that of *Cephalitis*, *Encephalitis*, or *Cerebritis* to inflammation of the substance of the brain; but there are no pathognomic symptoms, which can lead us to refer the seat of the disease to either of those parts exclusively. Phrenitis is, of course, a very dangerous affection, from the importance of the organ concerned. The most active treatment must necessarily be pursued. Repeated bleeding; purging; the application of ice to the shaved head; a position in which the head is raised; and the avoiding of irritation of every kind;—form the remedial means during the violent stage. After this has yielded, counter-irritants to the nape of the neck or to other parts of the body will be useful adjuvants. When the brain and meninges were affected only symptomatically, as in inflammation of the diaphragm, the disease was formerly called *Paraphrenitis*.

Hydrocephalus acutus is a variety of phrenitis.

PHRENTIS CALENTURA, *Coup-de-soleil*—p. Latrans, Hydrophobia—p. Potatorum, Delirium tremens—p. Vermiosa, Vercoquin.

PHRENO-GLOTTISM, Asthma thymicum.

PHRENOLOGY, Craniology.

PHRENO-MAG'NETISM, *Phreno-mes'merism*, *Mes'mero-phrenology*. A term applied to the condition of a mesmerized person, in which, when any cerebral or phrenological organ—so called—

is touched, the functions of the organ are manifested. It has been established, that no such phenomena can be elicited in children, or in those who have never seen a phrenological cast or picture.

PHRENOPARALYSIS, see Paralysis.

PHRENOPATHIA, see Phrenica.

PHRENOPATH'IA ÆTH'E'RA. The aggregate of encephalic phenomena induced by the inhalation of ether.

PHRENSY, Phrenitis.

PHRICASMUS, Horripilation.

PHRICE, Horripilation.

PHRICIA, Horripilation.

PHRICIASIS, Horripilation.

PHRICO'DES (FEBRIS,) from *φρικν*, 'a shivering;' *Horri'fica*. A fever in which the chill is very intense and prolonged.

PHRYAGMA, Shuddering.

PHRYCTE, Colophonias.

PHRYG'IUS (LAPIS). An earthy matter, found in Phrygia and Cappadocia, and formerly employed as an astrigent. —Gorrieus.

PHTHARMA CALIGO, Caligo—p. Cataracta, Cataract—p. Glaucoma, Glaucoma.

PHTHEIR, Pediculus.

PHTHEIRE'MIA, from *φθειρω*, 'I corrupt,' and *αίμα*, 'blood.' A state in which the plastic powers of the blood are deteriorated.

PHTHEIRIASIS, *Phthiri'asis*, *φθειρίασις*, from *φθειρω*, 'a louse;' *Morbus pediculo'sus seu pedicul'aris*, *Pruri'go pedicul'aris*, *Pedicula'tio*, *Malis pedicul'i*, *Peta'lē*, *Lousiness*, *Lousy disease*, (F.) *Maladie pédiculaire*, *Phthiri'asis*. A disease which consists in the excessive multiplication of lice on the human body, under conditions in which it does not ordinarily take place, and in spite of cleanliness. It has been asserted that the affection has often caused death. Simple and medicated baths have been recommended for its removal; lotions of the decoction or infusion of tobacco and stavesacre, mercurial frictions, &c.

PHTHEIRIUM, Delphinium staphisagria.

PHTHEIROCTONUM, Delphinium staphisagria.

PHTHINAS, Phthisis.

PHTHINODES, Phthisicus.

PHTHIRIASIS, Phthiriasis.

PHTHIRIASIS, Malis, Phthiriasis.

PHTHIRIUM SCABIEI, see Psora.

PHTHIRIUS, Antiparasitic—p. Inguinalis, see Pediculus.

PHTHIROCTONUS, Abortive.

PHTHIROPORON, Autumn.

PHTHISES: same etymon as Phthisis. In the classification of Fuchs, diseases attended with ulceration, hectic fever, and emaciation.

PHTHISIC, Phthisis. Also, a slight tickling cough, (Prov.) *Tissick*.

PHTHISICAL, Phthisicus.

PHTHIS'ICUS, *Phthino'des*, *Phthis'ical*, *Tis'ical*, (F.) *Phthisique*, *Poitrinaire*; same etymon as phthisis. Belonging to phthisis. Also, one labouring under phthisis, or who is consumptive.

PHTHISIE, Phthisis pulmonalis—p. *Aiguë*, Consumption, galloping—p. *Calculense*, Calculi, pulmonary, see Lithiasis pulmonum—p. *Carcinomatense*, Phthisis, cancerous—p. *Galopante*, Consumption, galloping—p. *avec Mélanose*, Anthracosis pulmonum—p. *Mésentérique*, Tabes mesenterica—p. *Trachéale*, see Phthisis laryngea—p. *Tuberculeuse*, Phthisis pulmonalis.

PHTHISIOLOGY, *Phthisiolog'ia*, from *φθισις*, and *λογος*. A treatise on phthisis.

PHTHISIOSIS, Tubercular consumption.

PHTHISI-PNEUMONIA, Phthisis pulmonalis.

PHTHISIQUE, Phthisicus.

PHTHISIS, *Phthoë*, *Phthinas*, *Tabes*, *Te'ce*.

don, Consumption, Decline, Phthisis, Tisic, from φθίω, 'I waste away, I decay.' This word, taken in a general sense, means progressive emaciation of every part of the body. It is usually, however, restricted to phthisis pulmonalis.

PHTHISIS ACUTA, Consumption, galloping — p. Black, see Anthracosis.

PHTHISIS, BRONCHIAL, *Phthisis bronchialis*. Phthisis occasioned by the pressure of enlarged tuberculous bronchial glands, which, not unfrequently, communicate with caverns in the lungs, or with the bronchia.

PHTHISIS CALCULOSA, Lithiasis pulmonum.

PHTHISIS, CANCEROUS, *Scirrhus, Carcinoma et Fungus Pulmonum, Medullary or Encephaloid Tumour of the Lung, (F.) Cancer du Poumon*. Cancer of the lung, which gives rise to carcinomatous phthisis, (F.) *Phthisie carcinomateuse*, of Bayle.

PHTHISIS CONSUMMATA, see Ph. pulmonalis — p. Dorsalis, Tabes Dorsalis — p. Dyspeptic, see Phthisis — p. Exulcerata, Ph. pulmonalis — p. Hepatica, Hepato-phthisis — p. Ischiadica, Ischio-phthisis.

PHTHISIS LARYNGE'A, *Chronic Laryngitis, Laryngitis et Tracheitis chronica, Tuberculosis laryngis et tracheae, Laryngo-et-Tracheophthisis, Laryngophthisis, Ulcera seu Helco'sis Laryngis, (F.) Phthisie laryngée, Ulcère du larynx, Laryngite avec sécrétion de Pus*, is a species of consumption analogous to phthisis pulmonalis, produced by inflammation and ulceration of the interior of the larynx, and sometimes by caries of the cartilages. *Phthisis trachealis, Tracheophthisis, (F.) Phthisie trachéale*, is a similar affection, seated in the trachea. Chronic inflammation of the larynx and trachea may exist, however, without tubercles or phthisis.

PHTHISIS MESERICA, Tabes mesenterica — p. Nodosa, Ph. pulmonalis — p. Mucosa, see Bronchorrhoea — p. Notias, Tabes dorsalis — p. Pituitosa, see Bronchorrhoea.

PHTHISIS PULMONALIS, *Ph. pulmonum seu pulmonaris seu nodo'sa pulmonalis vera seu tuberculo'sa seu scrophulo'sa seu tuberculo-ulcerata seu purulen'ta exulcerata seu exulcerata, Marasmus seu Hectica seu Hamoptysis phthisis, Passio vomitocoflua, Phthisis-pneumonia, Tabes pulmonalis seu pulmo'num, Ulcus pulmonum, Helco'sis pulmonalis, Pneumophthisis, Pneumophthoë, Pneumophthoë, Pulmo'nia, Tuberculosis pulmonum, Gongrophthisis, Morbus Anglicus, Tuberculous disease of the lungs, Pulmonary consumption, Consumption, Decline, Phthisis, Tisic, (Sc.) Teasick, (F.) Phthisie, P. pulmonaire seu tuberculeuse, Crachement de pus*. Occasionally, this formidable disease occurs accidentally: but, usually, it is caused by a conformation obtained from progenitors. This, at least, predisposes to it; and but a slight exciting cause may be required to rouse the predisposition into action. It is less a disease of the torrid zone; and in the temperate regions of the globe, its pathology is often intimately allied with that of scrofula. It consists in the formation of tubercles in the lungs, which sooner or later inflame and break down. In such a constitution, ulcerations of the lungs do not readily heal; and hectic fever — the universal attendant upon irritability and debility — is established. This persists, and gradually wears down the patient, till he ultimately expires, exhausted by the irritative fever. The chief symptoms are the following. The patient feels a sense of uneasiness in the chest, with severe fits of coughing, or a short and dry cough with dyspnoea. This may be considered the incipient stage. The cough then becomes extremely troublesome, especially in the morning; and dyspnoea is increased on the

slightest exertion. The expectoration is evidently purulent, with fever in the evening, and circumscribed redness of the cheeks; wasting and colliquative sweats early in the morning. This must be considered the confirmed stage, — *Phthisis consummata, Pneumophthisis*. In the third, or last stage, the hectic is constant, with small pulse, pungent heat, colliquative sweats, and diarrhoea. The weakness becomes extreme, and the patient exhausted. *Physical Signs*. — 1. In the first stage, whilst there are yet scattered miliary tubercles, — the only appreciable change at the exterior of the chest, and this is not constant, is a want of freedom in the movements under one or both clavicles. Percussion will generally discover a slight diminution of sound corresponding to the summit of one lung. In the same situation, auscultation shows that the inspiratory murmur is either weaker, rougher, or confused. The sound of expiration is more distinct. The voice and cough are rather more resonant, and wheezing or mucous râles are occasionally heard in the same region. 2. When the tubercular deposition has become more abundant, the motion of the chest is more markedly diminished under one or both clavicles. The loss of sound on percussion is more distinct, the inspiration, on auscultation, is bronchial, the expiration louder and more superficial, and the resonance of the voice and cough is increased. The rest of the lung is perhaps healthy, or the respiration may be puerile. 3. When the tubercles are undergoing softening, — in addition to the preceding signs, auscultation exhibits a subcrepitant râle around the apex of the lung, decreasing from above downwards. It is most distinct during or after the cough, and at the close of a deep inspiration. As the disease proceeds, the bubbling becomes successively larger, and ultimately gargouillement is heard. The inspiration and expiration are cavernous or tracheal. The voice and cough indicate unusual resonance and pectoriloquy. The intercostal spaces are often depressed locally, and percussion may become gradually clearer and even tympanitic as the cavern increases.

It is only in the incipient state of this formidable disease, that advantage can be expected from any plan of treatment; and of all means that can be devised, that of removing to a climate in which air and exercise can be daily enjoyed during the autumnal, winter, and spring months is the best. See Climate. If this be impracticable, the patient must be kept in a regulated temperature, taking care that the chamber is duly ventilated; and gestation in the open air be used, whenever the atmosphere, although cold, is dry. The other indications will be: 1. To moderate inflammatory action: 2. To support the strength; and, 3. To palliate urgent symptoms. The whole treatment indeed is palliative. Counter-irritants relieve pain: demulcents and mild opiates alleviate the cough: the mineral acids, united with opium, relieve the colliquative sweats: and chalk and opium the diarrhoea.

Dr. Wilson Philip has pointed out a particular species of phthisis, which is characterized, in the early stage, by a deranged state of the digestive organs; flatulent, irregular bowels; furred tongue; impaired appetite; unnatural faeces; and the epigastric region more or less full and tender on pressure. The affection of the lungs is here secondary. He calls it *Dyspeptic phthisis*.

PHTHISIS PULMONALIS PURULENTA EXULCERATA, *Phthisis pulmonalis* — p. Pulmonalis scrophulosa, *Phthisis pulmonalis* — p. Pulmonalis tuberculosa, *Phthisis pulmonalis* — p. Pulmonalis tuberculosa-ulcerata, *Phthisis pulmonalis* — p. Pulmonalis vera, *Phthisis pulmonalis* — p. Pupillæ, Synechisis — p. Pupillaris, Myosis — p. Scrophu-

Iosa, Phthisis pulmonalis — p. Sicca, Tabes dorsalis — p. Tuberculosa, Phthisis pulmonalis — p. Uterina, Metrophthisis — p. Vesicalis, Cystophthisis.

PHTHISIS WEED, *Ludwigia palustris*.

PHTHISURIA, Diabetes.

PHTHISURIE, Diabetes — p. *Sucrée*, Diabetes mellitus.

PHTHOE, Phthisis.

PHTHOIS, Pastil, Trochiscus.

PHTHOISCOS, Pastil, Trochiscus.

PHTHONGODYSPHORIA, Hyperacusis.

PTHORA, Abortion, Corruption, Plague.

PHTHORIUS, Abortive.

PHTHOROCTONUS, Abortive.

PHUCAGROSTIS MINOR, *Pila marina*.

PHUSCA, Oxyerate.

PHYGANTHROPIA, Misanthropia.

PHYGETH'LO'N, *Panus*, *Phypella*, from *φύω*, 'I grow,' [?]. Inflammation of the superficial

lymphatic glands, not ending in suppuration.

PHYLACTERION, Amuletum.

PHYLACTERIUM, *φυλακτῆριον*, from *φύλασσω*,

'I preserve.' A *Phylactery*, (F.) *Phylactère*.

An amulet: — a prophylactic.

PHYLAXIS, Conservation.

PHYLLANTHUS URINARIA, *Binkohumba*. Ord. Euphorbiaceæ. A plant employed in Ceylon, in decoction, as a diuretic.

PHYLLITIS, Asplenium scolopendrium — p. *Muraria*, Asplenium ruta muraria — p. *Rotundifolia*, Asplenium trichomanoides.

PHYMA, *Phima*, from *φύω*, 'I grow.' With the ancients, this word was used with great latitude: sometimes, in the sense of phlegmon, for tumours larger than a boil, and perfectly suppurative. In the nosology of Sauvages, *Phymata* is an order of diseases, to which he refers erysipelas, oedema, phlegmon, emphysema, scirrhus, &c. In that of Dr. Good it is a genus, including hordeolum, furunculus, syccosis, and anthrax.

PHYMA ANTHRAX, Anthrax — p. Anthrax terminatus, Terminus — p. Syccosis, Syccosis.

PHYMATOID, *Phymatoides*, (F.) *Phymatoides*, from *φύμα*, 'a tubercle,' and *ειδος*, 'resemblance.' A state of cancerous tissue, in which the dull yellow hue and the consistence resemble those of tubercle. — Lebert.

PHYMATOSES, from *φύμα*, 'a hard tumour.' Tuberculous diseases. — Fuchs.

PHYMATOSIS, Excrecence — p. *Cerebri*, Encephalophymata — p. Elephantiasis, Elephantiasis — p. *Lupus*, *Lupus* — p. *Verrucosa*, *Verruca*.

PHYMOSIS, Phimosis.

PHYPELLA, Phygethlon.

PHYSA, Crepitation.

PHYSAGOGA, Carminatives.

PHY'SALIS, from *φύσα*, 'a bladder,' referring to the inflated calyx, *P. Alkekengi* seu *halicacabum*, *Solanum vesicarium*, *Alchachen'ge*, *Alkekengi*, *Halicacabum*, *Winter Cherry*, (F.) *Alkekenge*, *Cogeret*. Ord. Solanaceæ. *Sex. Syst.* Pentandria Monogynia. The berries are recommended as a diuretic, from 6 to 12 for a dose, in dropsical and calculous cases. It has been affirmed, that the whole plant—twigs, leaves, capsules and berries—possesses the antiperiodic virtues of Cinchona.

PHYSALIS ALKEKENGI, *Physalis* — p. *Halicacabum*, *Physalis* — p. Obscura, *Ph. viscosa* — p. *Pennsylvanica*, *Ph. viscosa* — p. *Pubescens*, *Ph. viscosa* — p. *Stramonium*, *Anisodum luridum*.

PHYSALIS VISCO'SA, *P. obscura* seu *Pennsylvanica* seu *pubescens*, *Ground cherry*; indigenous. The berries are said to be diuretic and sedative.

PHYSCE, Abdomen, Colon.

PHYSCIA ISLANDICA, *Lichen Islandica* —

p. *Nivalis*, *Lichen caninus*.

PHYSCONIA, from *φύσκειν*, 'the paunch.'

Ventrositas, *Ecphy'ma physconia*, *Hyposarca*, *Emphractica*, *Visconia*, *Hypersarcid'ius*, *Potbelly*, *Pendulous abdomen*, *Swagbelly*, *Gorbelly*, (Prov.) *Gotchbelly*, *Pedbelly*. A generic term, under which Sauvages has comprised every large tumour developed in the abdomen, that is neither fluctuating nor sonorous. Linnæus calls it *Hyposarca*.

PHYSCONIA ADIPO'SA, *Obesitas* seu *Polyasar'cia viscerum*. Accumulation of fat in the intestines.

PHYSCONIA BILIOSA, *Turgescencia vesicularum*.

PHYSCONIA EMPHYSEMAT'ICA. Accumulation of air in the areolar tissue, or between the coats of particular viscera.

PHYSCONIA LIENIS, Splenoncus — p. *Mesenterica*, *Tabes mesenterica* — p. *Splenica*, *Ague cake*.

PHYSCONIA STRUMO'SA, *Serof'ula abdominalis interna*. Tumid abdomen from serophulous enlargement of the glands.

PHYSCONIE MÉSENTÉRIQUE, *Tabes mesenterica*.

PHYSEMA, Tympanites, Physisis.

PHYSE'SIS, *Physe'ma*, from *φύσσω*, 'I inflate.' An emphysematous tumour. Tympanites.

PHYSETER MACROCEPHALUS, see *Cetaceum*.

PHYSIATRICE, *Vis medicatrix naturæ*.

PHYSIAUTOCRATIA, *Vis medicatrix nature*.

PHYSIC, Medicament, *Medicina* — p. *Indian*, *Euphorbia corollata*, *Gillenia trifoliata* — p. *Root*, *Leptandra purpurea*.

PHYSICAL, Medical — p. *Signs*, see *Symptom*.

PHYSIC'IAN, *Med'icus*, *Iatros*, *Aces'tor*, *Aces'ter*, *Altheus*, *Παῖων*, *Pæon*, (Old Eng.) *Fesic'ian*, *Ficicion*, *Leche*, *Leech*, from *φύσις*, 'nature.' (F.) *Médecin*, *Docteur*. One who has received his degree from an incorporated Institution, as Doctor of Medicine. The French formerly used the word *physicien* in the same sense. It is now appropriated by them to the natural philosopher. *Physicist* has the same signification.

PHYSICIAN, FAMILY. One regarded as the regular attendant on a family.

PHYSICIAN, LEARNED, *Iatrosophista* — p. *Practical*, *Iatrotechna* — p. *Priest*, see *Asclepiada*.

PHYSICIEN, *Physician*.

PHYSICIST, see *Physician*.

PHYSIOS, MED'ICAL, *Iatrophys'ics*. *Physios* directly applied to medicine, — either to the explanation of the vital phenomena, the preservation of individuals, or the treatment of disease.

PHYSIOAUTOCRATIA, *Vis medicatrix nature*.

PHYSIOG'NOMIST, *Physiognomon*, one who busies himself especially with, or pretends to be versed in, physiognomy.

PHYSIOG'NOMY, (Old Eng.) *Vis'onomy*, *Physiognom'ia*, *Physiognom'ica*, *Physiognom'icè*, *Physiognom'ia*, *Physiognomon'ia*, *Physiognom'onè*, *Physiognomons'ynè*, *Physiognomosia*, *Anthroposcop'ia*, *Prosopomanti'ia*, *Prosopolog'ia*, *Semiol'icè fa'ciei*, *Physiognomon'ica*, *Physiognom'on'y*, *Metoposcopy*, *Prosopos'is*, from *φύσις*, 'nature,' and *γνωμν*, 'knowledge.' The art of knowing the dispositions of men from their external appearance; especially from the features. Every inference of this kind must necessarily be extremely fallacious, and cannot be reduced to rule, as was attempted by Lavater and others. In infantile diseases, numerous shades of expression are evident, which experience teaches how to appreciate; and which afford useful guides in understanding the pathology of that period of existence.

PHYSIOLOGIA PATHOGENETICA, see *Pathogenetic*.

PHYSIOLOG'ICAL, *Physiolog'icus*, (F.) *Physiologique*. Relating or belonging to physiology.

PHYSIOLOGICAL ANATOMY, see Anatomy—p. Doctrine, Broussaism.

PHYSIOLOGICE, Physiology.

PHYSIOLOGIST, *Physiologus*. Same etymon as the next. One who occupies himself specially, or is well acquainted, with physiology.

PHYSIOLOGY, *Physiolog'ia*, *Biolog'ia*, *Bion'omy*, *Biol'ica*, *Biophænomenolog'ia*, *Physiolog'icē*, *Æconom'ia animal'is*, *Anatom'ia viva*, *Anat'omē animal'is*, *Anthropolog'ia*, *Microcosmograp'hia*, *Microcos'mica scient'ia*, *Cognit'io physiolog'ica*, *Parē naturālis medic'inæ*, *Philosoph'ia corp'oris vivi*; from *φύσις*, 'nature,' and *λόγος*, 'a description.' Formerly, Physiology meant the same as Physics, in its extensive signification, now does. At the present day, it includes the science which treats of the functions of animals or vegetables; an acquaintance with the phenomena the aggregate of which constitute life. It is the science of life. It is divided into animal—*Zoophysiology*, or *Zoobiolog'ia*; and vegetable—*Phytophysiolog'ia*, or *Phytobiolog'ia*, according as it considers the life of animals or of vegetables singly. Comparative physiology comprises both. Physiology is, also, general or special, according as it treats of life in the abstract or in some particular species. To the latter belongs the *Physiology of Man*,—called also, *Hygienic Physiology* to distinguish it from *Pathological Physiology* or *Pathology*.

PHYSIOLOGY, PATHOGENETIC, see Pathogenetic.

PHYSIONOMIA, *Physiognomy*.

PHYSIONOMIE ÉGARÉE, see Wild.

PHYSIS, *Natura*, *Sex*.

PHYSOCÈLÈ, *Pneumatocel'is*, *Hernia vent'osa*; *Oscheocel'is flatulen'ta*, *Emphyse'ma Scroti*, from *φύσσω*, 'I inflate,' and *κελή*, 'a tumour.' An emphysematous tumour of the scrotum; probably a case of intestinal hernia, containing much flatus.

PHYSOCEPH'ALUS, from *φύσσω*, 'I inflate,' and *κεφαλή*, 'the head.' (F.) *Physocephale*. Emphysematous tumefaction of the head.

PHYSOCÈLIA, *Tympanites*.

PHYSO'DES, from *φύσσω*, 'I inflate,' and *εἶδος*, 'resemblance.' Full of air; apparently full of air—applied to tumours, &c.

PHYSOME'TRA, *Emphyse'ma U'teri seu Uteri'nium*, *Hysteroph'ysē*, *Hysteremph'ysē'ma*, *Hystremph'ysē'ma*, *Metremph'ysē'ma*, *Inf'lat'io U'teri*, *Tym'pany of the Womb*, *U'terine tympani'tes*, from *φύσσω*, 'I inflate,' and *μῆτρα*, 'the womb.' A light, tense, circumscribed protuberance in the hypogastrium, obscurely sonorous, with wind occasionally discharged through the os uteri with noise, *Hysteroph'ysē'ia*, *Ædæopsoph'ia uteri'na*.

PHYSON, *Flatulence*.

PHYSONCUS, *Emphysema*.

PHYSOPSOPI'IA; from *φύσσω*, 'I inflate,' and *ψόφος*, 'a noise.' A discharge of air with noise.

PHYSOSPASMUS, *Colica flatulenta*.

PHYSOTHORAX, *Pneumothorax*.

PHYTEL'EPHAS MACROCAR'PA, from *φύρον*, 'a plant,' and *ελεφας*, 'an elephant.' A tree of South America, called, by Willdenow, *Elephantu'sia*, the dense albumen of whose unt, is the vegetable ivory, which is as durable and nearly as hard as the ivory of the elephant.

PHYTIVOROUS, *Phytophagous*.

PHYTOALOPECIA, *Porrigo decalvans*.

PHYTOBIOLOGIA, see Physiology.

PHYTO-CHEMISTRY, see Chymistry.

PHYTOLAC'CA DECAN'DRA, (P.) *vulgaris*, *Poke Weed*, *American night shade*, *Cancer root*, *Poke*, *Skoke*, *Jucato Calleloe* (?), *Jalap*, *Pigeon Berry*, *Jalap cancer root*, *Sola'nium racem'osum America'num seu magnūm Virginia'nium rubrum*, *Blitum America'num*, *Garget*, (F.) *Raisin d'Amé-*

rique, *Phytolaque à dix étamines*, *Morelle en grappes*, *Méchoacan du Canada*. Ord. Chenopodeace; *Sex. Syst.* Decandria Decagynia; from *φύρον*, 'a plant,' and *lacca*; because the colour of the juice of the berries resembles that of lacca. The leaves, berries, and roots are employed. The young stems, when boiled, are sometimes eaten at table; but when old, they must be cautiously used, as the plant is emetic and cathartic. It is most celebrated as a remedy in chronic rheumatism; and is given in the form of tincture of the ripe berries. It is, also, used in cutaneous eruptions; tinea capitis, itch, cancerous ulcers, &c.

The root is gathered and dried in the fall.

PHYTOLAQUE À DIX ÉTAMINES, *Phytolacca decandra*.

PHYTOLOGIA MEDICA, Botany, medical.

PHYTO-PARASITE, Epiphyte.

PHYTOPH'AGOUS, *Phytoph'agus*; *Phytiv'orous*, *Phytiv'orus*; from *φύρον*, 'a plant,' and *φαγω*, 'I eat.' Feeding or subsisting on vegetables.

PHYTOPHYSIOLOGIA, see Physiology.

PHYTOSPERMATA, (F.) *Phytospermes*; from *φύρον*, 'a plant,' and *σπέρμα*, 'sperm.' Particles similar to the spermatozooids of animals, observed in vegetables, and presumed to possess a similar agency in reproduction.

PHYTOTOMY, see Anatomy, (vegetable.)

PHYTOZOOM, Zoophyte.

PIA MATER, *Pia seu mollis seu ten'uis Mater*, *Pia meninx*, *Meninx interior seu vasculo'sa seu choro'i'des*, *Inner lam'ina of the Méninge*, (Ch.), *Membra'na vasculo'sa Cer'ebr'i*, *M. Cer'ebr'i ten'uis seu mollis seu pro'pria*, (F.) *Pie mère*, *Méningette*. The pia mater is a very delicate membrane, which covers the brain immediately, penetrates into its anfractuosités, and envelops, also, the cerebellum and the spinal prolongation, &c. Bichat does not regard it as a membrane, properly so called, but as a net-work of blood-vessels, united by a lax areolar tissue. The *Pia Mater exterior* is the portion of membrane which covers the exterior of the encephalon, and faces the arachnoid externally. The *Pia Mater interior* is that which lines the cerebral fossæ, into which it penetrates by a cleft situate between the posterior part of the corpus callosum and the fornix; and by two other fissures, situate between the corpora fimbriata and thalami nervorum opticorum. The pia mater contains a great number of granulations similar to the *Glandula Pacchioni*.

PIA MATER TESTIS, *Tunica vasculosa*—p. Meninx, P. Mater.

PIAN, *Frambœsia*.

PIANS, MOTHER OF, *Mama-pian*.

PIAR, *Fat*.

PIARHÈ'MIA, *Piarrhè'mia*, from *πιπ*, 'fat,' and *αἷμα*, 'blood.' Fat in the blood.—Simon.

PICA, *Malacia*—p. Africanorum, Chthonopha'gia.

PICAC, *Euphorbia corollata*.

PICACISM'US, *Pica'tio*, from *piz*, *picis*, 'pitch.' Depilation by means of a pitch plaster. Also, *Malacia*.

PICATIO, *Malacia*, *Picacismus*.

PICEA, *Pinus abies*.

PICERON, *Butter*.

PICHU'RIM BEANS. The seed, *Faba Pichu'rim*, *F. Pechu'rei*, *Para nut*, *Sassafras nut*, *Puziri*, *Brazil'ian nutmeg*, of a tree, supposed to be *Perse'a pichurim*, *Ocote'a pichu'rim* seu *Puchury*, *Tetranthe'ra pichurim*, *Nectan'dra puchury*, or *Puchury* of the Brazilians, Ord. Lauraceæ, which grows in Brazil, Guiana, Venezuela, and other parts of South America. They are of an elliptical shape; flat on one side, convex on the other; of an aromatic odour between that of

nutmegs and saffras, and of a spicy, pungent taste. The medical properties are the same as those of other aromatics. For the bark, see *Pecurium cortex*.

PICK, Emetic.

PICKERELWEED, SHOVEL, *Unisema deltofolia*.

PICOTE, Variola.

PICOTEMENT, Pricking.

PICOTIN, *Arum maculatum*.

PICRA, Pulvis aloës et canellæ.

PICRÆNA EXCELSA, Quassia.

PICRIA, Bitterness.

PICRIUM SPICATUM, *Coutoubea alba*.

PICROCHOLOS, Bilious.

PICROMEL, *Bilin*; from *πικρος*, 'bitter,' and *μελι*, 'honey.' A name given by Thénard to a colourless, soft substance, of an acid, bitter, saccharine taste, nauseous smell, and very soluble in water, which is considered to be a *bilate of soda*. It is not an immediate principle, but a mixture of different substances. It forms part of the bile, and is generally classed amongst the animal immediate principles.

PICROPE'GÆ, *Pikropo'gæ*, from *πικρος*, 'bitter,' and *πνyn*, 'a spring.' Mineral waters; bitter from the presence of chloride of magnesium.

PICROS, Amarus.

PICROTOX'INE, *Picrotox'ina*, *Picrotox'in*, from *πικρος*, 'bitter,' and *toxicum*, 'poison.' A vegetable alkali, discovered by M. Boullay in the *Menispermum Coccus*. It crystallizes in quadrangular, white, brilliant, semitransparent prisms; which are excessively bitter, soluble in three parts of alcohol, and in fifteen of boiling water.

The *Coccus Indicus* owes its poisonous properties to this principle.

PIE MÈRE, *Pia mater*.

PIE-MÉRITE, *Inflammat'io piæ matris*. A miserably compounded term, employed by M. Griolle, to express inflammation of the pia mater:—from (F.) *pie mère*, 'the pia mater,' and *ite, itis*; a termination denoting inflammation.

PIÈCE, Hepatitis.

PIÈCES D'ANATOMIE ARTIFICIELLES, see Artificial.

PIED, Pes—*p. d'Alexandre*, *Anthemis pyrethrum*—*p. d'Alouette des champs*, *Delphinium consolida*—*p. Bot*, see *Kyllosis*—*p. de Chat*, *Antennaria dioica*—*p. de Cheval marin*, *Cornu ammonis*—*p. de Griffon*, *Gryphius pes*, *Helleborus fœtidus*—*p. de Lièvre*, *Arum maculatum*—*p. de Lion*, *Alchemilla*—*p. de Lit*, *Origanum*—*p. d'Ours*, *Acanthus mollis*—*p. Plat*, see *Kyllosis*—*p. de Veau*, *Arum maculatum*—*p. de Veau triphyllæ*, *Arum triphyllum*.

PIERON, Fat.

PIERRE, Calculus—*p. d'Aigle*, *Ætites*—*p. d'Aiment*, *Magnet*—*p. d'Azur*, *Lapis lazuli*—*p. Calaminaire*, *Calamine*—*p. d'Caude*, *Potassa cum calce*—*p. Divine*, *Lapis divinus*—*p. d'Écrevisse*, *Cancerorum chelæ*—*p. Infernale*, *Argent nitras*—*p. Judaïque*, *Judæus (lapis)*—*p. au Lait*, *Morochthus*—*p. Médicamenteuse*, *Medicamentosus lapis*—*p. Néphrétique*, *Nephreticus lapis*—*p. Ponce*, *Pumex*.

PIERRES CRAYEUSES, Calculi, arthritic—*p. au Fiel*, Calculi, biliary—*p. Stercorales*, Calculus of the Stomach and Intestines—*p. Urinaires*, Calculi, urinary.

PIERREUX, Petrous.

PIESTER, Press.

PIESTERION, Press.

PIESTRON, from *πιεω*, 'I press.' An instrument recommended by Hippocrates for breaking the bones of the fetal head when too large to be extracted otherwise.

PIGAMON JAUNÂTRE, *Thalictrum*.

PIGEON BERRY, *Phytolacca decandra*—*p. Breasted*, see *Lordosis*—*p. Tree*, *Aralia spinosa*.

PIGMENT CELLS, see Cell, pigment.

PIGMENTAL, *Pigmentary*, *Pigmenta'rius*, (F.) *Pigmentaire*, from *pigmentum*, (*pingere*, *pic-tum*, 'to paint,') 'paint.' Having relation to pigment or colouring matter.

PIGMENTAL APPARA'TUS, (F.) *Appareil pigmental*. The apparatus concerned in the coloration of the skin of the coloured varieties of man.—*Flourens*.

PIGMENTARIUS, Apothecary, Unguentarius. PIGMENTARY, Pigmental—*p. Nævi*, see *Nævus*.

PIGMENTUM, Paint—*p. Indicum*, *Indigo*—*p. Nigrum*, see *Choroidea tunica*—*p. Uracu*, *Terra Orleana*.

PIGMY, Pygmy.

PIGNEROLE, *Cantaurea calcitrapa*.

PIGNON DE BARBARIE, *Jatropha curcas*—*p. d'Inde*, *Jatropha curcas*—*p. d'Inde, petit*, *Croton tiglium*.

PIGNUT, *Earthnut*, *Groundnut*. The bulbous root of *Bu'nium Bulbocastanum*. *Ord. Umbellifera*. Called *pignuts*, from the fondness of the pig for them. They have been considered to possess a styptic quality; and are deemed serviceable in atonic affections of the kidney.—*Hooper*.

PIGNUT, *Bunium bulbocastanum*.

PIG'S FLARE, *Adeps suillus*.

PIKROPEGÆ, *Picropegæ*.

PIKRY, Pulvis aloës et canellæ.

PILA, *Sphæra*, *Palla*, *παλλα*, *Ball*. A gymnastic exercise, much used by the ancients as well as moderns.—*Galen*. Also, a mortar and a pestle—*Ac'ônê*.

PILA, Mortar, Pillar—*p. Damarum*, *Ægagropila*—*p. Rupicaprarum*, *Ægagropila*.

PILA MAR'NA, *Sphæra Marina* seu *Thalas'sia*, *Halcyonium rotundum*, (F.) *Pelote de Mer*, *Zostère*, *Algæ marine*, *A. commune* ou des *Verriers*. Certain light, round, depressed, or oblong masses—inodorous and insipid—in which rushes, hairs, and the débris of shells are met with. These are found on the sea-shores, and have been regarded by Spielmann as consisting chiefly of the roots of the *Zostera Marina* seu *Marit'ima*, *Alga Mari'na*, (*Big tangles*, *Ware*, *Wair*, *Sea-ware*,) *Phucagros'tis minor*. Iodine is detected in the ashes, which accounts for their success in goitre. They have, also, been given as a vermifuge.

PILAIRE, Pileous.

PILAR, Pileous.

PILARE MALUM, Trichosis.

PILA'TIO, from *pilus*, 'a hair.' Fracture of the cranium in which the fissure is narrow, and has the appearance of a hair—a *capillary fissure*, (F.) *Fente capillaire*.

PILE, Hair.

PILEA PUMILA, from *pilus*, 'a hair,' *pilosus*, 'hairy.' *Richweed*, *Clearweed*, *Coolweed*; indigenous; *Order*, *Urticacæ*; flowering from July to September; has been used as a wash in poisoning by *Rhus*. It has a strong, grateful smell.

PILEOLUM, *Calotte*.

PILEOLUS, Caul, Pileus.

PI'LEOUS, *Pilo'sus*, *Pilose*, *Pilous*, *Pilar*, (F.) *Pileux*, *Pilaire*, from *pilus*, 'a hair.' That which relates to the hair. *Hairy*.

PILEOUS SYSTEM of Bichat is that formed by the hair. It comprises the different species of hair of the human body.

PILES, BLIND, *Cæcæ hæmorrhoides*, see *Hæmorrhoids*.

PILEUS, 'a hat, cap, or bonnet for the head.' *Pile'olus*. A shield. A nipple-like instrument, formed of various materials, and used by females

for protecting the nipple during the child's sucking, when the pain is painful. Also, a caul.

PILEUS, Cucupha—p. Hippocraticus, *Bonnet d'Hippocrate*.

PILEUX, Pileus.

PILEWORT, Ranunculus acris, and R. scaria.

PILI CAPITIS, see Capillus—p. Cutis, Lanugo—p. Palpebrarum, Cilia—p. Pudendorum, Pubes.

PILLER, Pillar.

PILIFEROUS, *Pilifer*, from *pilus*, 'a hair,' and *fero*, 'I bear.' Pile or hair bearing, —as a 'piliferous cyst,'—one containing hair.

PILIMICTIO, see Trichiasis.

PILL, *Pilula*—p. Abernethy's, see *Pilulæ hydrargyri*—p. of Aloes and Camboge, *Pilulæ aloes et cambogiæ*—p. of Aloes and cinchona, *Pilulæ aloes et kinæ kinæ*—p. of Aloes and colocynth, *Pilulæ aloes et colocynthidis*—p. of Aloes and fetid gums, *Pilulæ de aloes et foetidis*—p. of Aloes and ginger, *Pilulæ aloes et zingiberis*—p. of Aloes and iron, *Pilulæ aloes et ferri*—p. Aloes and myrrh, *Pilulæ aloes et myrrhæ*—p. Aloetic, *Pilulæ aloeticæ*—p. Aloetic and assafoetida, *Pilulæ aloes et assafoetidæ*—p. Aloetic, compound, *Pilulæ aloes compositæ*—p. of Ammoniuuret of copper, *Pilulæ ammoniureti cupri*—p. Analeptic, James's, see *Analeptica*—p. Anderson's, see *Pilulæ aloes et jalapæ*—p. Anodyne, *Pilulæ opiatæ*—p. Antibilious, Barclay's, *Pilulæ antibiliosæ*—p. Antibilious, Dixon's, *Pilulæ antibiliosæ*—p. Asiatic, *Pilulæ Asiaticæ*—p. Assafoetida, *Pilulæ Assafoetidæ*—p. Bacher's, *Pilulæ ex helleboro et myrrhæ*—p. Barbarossa's, *Barbarossæ pilulæ*—p. Barthez's, see *Pilulæ aloes et myrrhæ*.

PILL, BLAIR'S. An empirical preparation of colchicum, much used in England in cases of gout.

PILL, BLAUD'S. These pills, proposed by M. Blaud as antichlorotic, are formed as follows:—Take of gum tragacanth, in powder, six grains; water, one drachm. Macerate in a glass or porcelain mortar, until a thick mucilage is formed, and if it be desired to prevent the formation of peroxide of iron, and to make the pill similar to those of Vallet, substitute a drachm of powdered sugar for the mucilage. Add afterwards of sulphate of iron, in powder, half an ounce. Beat well until the mixture is homogeneous, and add subcarbonate of potassa, half an ounce. Beat until the mass becomes of a deep-green colour, and of a soft consistence. Divide into 48 [?] pills. Dose, one a day; gradually increasing to two, and afterwards to three.

PILL, BLUE, *Pilulæ hydrargyri*—p. Brandreth's, *Pilulæ aloes et cambogiæ*—p. Calomel, *Pilulæ hydrargyri chloridi mitis*—p. Calomel, compound, *Pilulæ hydrargyri submuriatis compositis*—p. Calomel and opium, *Pilulæ calomelanos et opii*—p. Camboge, *Pilulæ cambogiæ compos.*—p. Camboge, compound, *Pilulæ cambogiæ compositis*—p. of Carbonate of iron, *Pilulæ ferri carbonatis*—p. Cathartic, compound, *Pilulæ cathartice compositis*—p. of Colocynth and henbane, *Pilulæ colocynthidis et hyoscyami*—p. of Colocynth, compound, *Pilulæ colocynthidis compositis*—p. Common, *Pilulæ aloes et myrrhæ*—p. of ammoniated Copper, *Pilulæ cupri ammoniatis*—p. of Digitalis and squill, *Pilulæ digitalis et scillæ*—p. Dinner, see *Pilulæ aloes et kinæ kinæ*, and *Pilulæ rhej*.

PILL, DINNER, of MR. MAYO. This is composed of P. Rhej, gr. iv; Sodæ Carb. gr. j.

PILL, EVERLAST'ING. A pill once in vogue, and said to be composed of metallic antimony. It was believed to have the property of purging as often as it was swallowed. A lady having swallowed one of these pills, became seriously alarmed at its not passing. "Madam," said her physician, "fear not; it has already passed

through a hundred patients without any difficulty."—Paris.

PILL, FEMALE, *Pilulæ de aloes et foetidis*—p. Ferruginous, of Vallet, see *Ferri protocarbonas*—p. Fothergill's, see *Pilulæ aloes et colocynthidis*—p. Frank's, see *Pilulæ aloes et kinæ kinæ*—p. Fuller's, see *Pilulæ de aloes et foetidis*—p. Galbanum, compound, *Pilulæ galbani compositis*—p. Griffith's, *Pilulæ ferri compositis*—p. Griffith's, see *Pilulæ rhej compositis*—p. Gum, *Pilulæ galbani compositis*—p. Harvey's, *Pilulæ aloes et colocynthidis*—p. of Hellebore and myrrh, *Pilulæ ex helleboro et myrrhæ*—p. of Hemlock, compound, *Pilulæ conii compositis*—p. Hooper's, see *Pilulæ aloes et myrrhæ*—p. of Iodide of iron, *Pilulæ ferri iodidi*—p. of Ipecacuanha and opium, *Pilulæ ipecacuanhæ et opii*—p. of Ipecacuanha and squill, *Pilulæ ipecacuanhæ cum scillâ*—p. of sulphate of Iron, *Pilulæ ferri sulphatis*—p. James's Analeptic, see *Analeptica*—p. Keyser's, see *Hydrargyrum acetatus*—p. Lady Crespigny's, *Pilulæ aloes et kinæ kinæ*—p. Lady Hesketh's, see *Pilulæ aloes et kinæ kinæ*—p. Lady Webster's, *Pilulæ aloes et kinæ kinæ*.

PILL, LARTIGUE'S, *Pilulæ Colchici et Digitalis*, (F.) *Pilules de Lartigue*. (*Ext. Coloc. c. 3ss; Ext. Sem. Colchic. alcoholic., Extract. Digital. alcoholic. 3â gr. iss; f. massa in pilul. x dividend.*) Dose, two, three, or more, in the 24 hours, in gout, until purging is induced.

This is a form that has been employed as a substitute for the *Pilules de Lartigue*, the composition of which is not known. M. Bouchardat recommends for the same purpose a combination of *Extractum colocynthidis compositum*, *Extractum colchici* and *Extractum opii*, which is almost identical with pills prescribed by Sir Henry Holland for the gout; the only difference being, that the latter substitutes *Dover's powder* for the *Extractum opii*.

PILL, OPIATE, of LEAD, *Pilulæ plumbi opiatæ*.

PILL, LEE'S NEW LONDON, is composed of scammony, gamboge, calomel, jalap, aloes, soap and syrup of buckthorn.

PILL, LEE'S WINDHAM, consists of gamboge, aloes, soap and nitrate of potassa.

PILL MACHINE, *Pilulier*—p. Matthews's, see *Pilulæ ex helleboro et myrrhæ*—p. Mercurial, *Pilulæ hydrargyri*—p. of mild chloride of Mercury, *Pilulæ hydrargyri chloridi mitis*—p. Moffat's, *Pilulæ aloes et cambogiæ*—p. Morrison's, *Pilulæ aloes et cambogiæ*, see *Pilulæ cambogiæ compositis*—p. Moseley's, *Pilulæ rhej compositis*—p. Night, common, *Pilulæ opiatæ*—p. Opiate, *Pilulæ opiatæ*—p. Peter's, see *Pilulæ aloes et cambogiæ*—p. Plummer's, *Pilulæ hydrargyri submuriatis compositis*—p. of sulphate of Quinia, *Pilulæ quiniæ sulphatis*—p. Red, *Pilulæ hydrargyri submuriatis compositis*—p. Red, Boerhaave's, see *Hydrargyri sulphuretum rubrum*—p. of Rhubarb, *Pilulæ rhej*—p. of Rhubarb, compound, *Pilulæ rhei compositis*—p. Rudius's, *Pilulæ aloes et colocynthidis*—p. Rufus's, *Pilulæ aloes et myrrhæ*—p. Rush's, see *Pilulæ aloes et cambogiæ*—p. Scotch, *Pilulæ de aloes et foetidis*—p. of Soap and opium, *Pilulæ saponis cum opio*—p. Speediman's, see *Pilulæ aloes et myrrhæ*—p. Squatting, *Pilulæ opiatæ*—p. Squill, compound, *Pilulæ scillæ compositis*—p. Stahl's, see *Pilulæ de aloes et foetidis*—p. Starkey's, see *Pilulæ ex helleboro et myrrhæ*—p. Stomachic, see *Pilulæ aloes et kinæ kinæ*, and *Pilulæ rhej*—p. Storax, *Pilulæ e styrace*—p. of Strychnia, *Pilulæ strychniæ*—p. of Submuriate of mercury, compound, *Pilulæ hydrargyri submuriatis compositis*—p. Tanjore, *Pilulæ Asiaticæ*—p. Turpentine, *Pilulæ de terebinthinâ*—p. Vallet's, see *Ferri protocarbonas*, and *Pilulæ ferri carbonatis*—p. Webster's,

Pilulæ aloes et kinæ kinæ — p. Whytt's, see Pilulæ de aloes et foetidis.

PILLAR, *Pi'la*, *Colum'na*, *Columell'a*, (F.) *Pilier*. A name given to several parts; as the pillars of the velum palati, diaphragm, &c.

PILULA, *Pilula*.

PILNA, MINERAL WATERS OF, see Püllna.

PILON, *Pilum*.

PILOSELLA, *Hieracium pilosella*—v. *Alpina*, *Hieracium pilosella*.

PILOSUS, *Pileosus*.

PILOUS, *Pileosus*.

PIL'ULA, *Pill'ula*, plural *Pil'ulæ*. Diminutive of *pila*, 'a ball.' *Spharid'ion*, *Catapotion*, *Catap'otum*, *Catap'otia*, *Spha'rion*, *Gongyl'ion*, *Gongylis*, *Gongylid'ium*, *Spha'rula Gale'ni*. A pill. (F.) *Pilule*. A simple or compound medicine, weighing from one to five grains, of a firm consistence and spherical form; intended to be swallowed at once, without chewing. The *Pill* differs from the *Bolus*, the latter being softer and larger.

Medicines are often given in the form of pill, on account of their disagreeable taste and odour. *Pilula*, also, means a *Pilular mass*.

PILULÆ EX ALOE, *P. aloes compositæ*—p. de Aloes et cambogiâ, *P. aloes et cambogiæ*.

PILULÆ ALOES, *P. Aloët'icæ* seu *ex Aloë et Sapo'ne*, *Aloët'ic Pill*, (F.) *Pilules d'aloës ou aloëtiques*. (Aloes in pulv.; saponis. sing. $\mathfrak{z}\mathfrak{j}$; syr. q. s. to form 240 pills.—Ph. U. S.) Dose, gr. x. to xx, in habitual costiveness.

PILULÆ ALOES ET ASSAFÆTIDÆ, *Aloetic and Assafetida Pills*. (Aloes, assafetida; saponis. sing. $\mathfrak{z}\mathfrak{ss}$; aquæ. q. s. to form 180 pills.—Ph. U. S.) Purgative and stomacic. Dose, gr. x.

PILULÆ ALOES ET CAMBOGIE, *P. de Aloë et Cambogiâ* seu *Hydrago'gæ Bon'tii*, *Pill of Aloes and Camboge*. (Aloes, cambog., gum. ammoniac., \mathfrak{aa} p. \mathfrak{ss} .) Make into a powder. Dissolve in vinegar; pass through a cloth; express the residuum strongly; and afterwards evaporate, in a water-bath, to a pilular consistence.—Ph. P.) Purgative; used in dropsy. Dose, gr. xij to xxx.

Peter's Pills, —a nostrum—are formed of aloes, jalap, scammony, and camboge, of each equal parts.

The basis of Morrison's Pills, often called *Hygienic Pills*, of Brapdrecht's Pills, and Moffat's Pills, is said to be aloes, with camboge, scammony, &c.

Rush's Pills were formed of calomel, aloes, and camboge, with oil of mint.

PILULÆ ALOES ET COLOCYN'THIDIS, *P. ex Colocynth'idæ* cum aloë seu *Aloes cum Colocynthidæ* seu *coc'cinæ seu coc'chii seu colocyn'thidis compos'itæ* seu *coc'ciæ mino'res*, *Pill of Aloes and Colocynth.*, (F.) *Pilules panchymagogues ou universelles*, *Pilules cochéæ*. (Pulp. coloc. $\mathfrak{z}\mathfrak{ss}$; aloes hepat., scammon. \mathfrak{aa} $\mathfrak{z}\mathfrak{j}$; sapon. $\mathfrak{z}\mathfrak{j}$; ol. caryoph. $\mathfrak{z}\mathfrak{j}$; syrup. q. s.—Ph. D.) Cathartic. Dose, gr. v to x or more.

Fothergill's Pills consist of aloes, scammony, colocynth, and oxide of antimony; and Harvey's Antibilious Pills, and Rudius's Pills do not differ much from them.

PILULÆ ALOES COMPOSITÆ, *P. ex Aloë*, Compound Aloetic Pills. (Aloes epic. extract. cont. $\mathfrak{z}\mathfrak{j}$; ext. gentiana, $\mathfrak{z}\mathfrak{ss}$; olei carui, gtt. xl; syr. q. s.—Ph. L.) These are sometimes, also, called Family pills, Antibilious pills, &c.

PILULÆ ALOES ET FERRI, *Pills of aloes and Iron*. (Ferri sulphat. p. iij; Aloës Barbād. p. $\mathfrak{z}\mathfrak{j}$; Pulv. arom. p. vj; Conserv. Rosæ. p. viij.—Ph. Ed.) In atonic constipation. Dose, one to three pills.

PILULÆ DE ALOË ET FOETIDIS, *P. Benedic'tæ*

Fulleri seu *Ecphrac'ticæ* seu *Aloës martia'les*, *Pills of aloes and fetid gums*, *Fenale pills*, (F.) *Pilules bënites* de Fuller. (Aloes socotr., sennæ, assafetida, galbani, \mathfrak{aa} $\mathfrak{z}\mathfrak{j}$; myrrh. $\mathfrak{z}\mathfrak{j}$; croci, macis, \mathfrak{aa} $\mathfrak{z}\mathfrak{j}$; ferri sulphat. $\mathfrak{z}\mathfrak{ss}$. Reduce to powder separately, and add ol. sucin. gtt. viij; syr. artemisiæ, q. s.—Ph. P.) Purgative, and given in hysteria. Dose, gr. xv to xx.

Whytt's Pills, given in similar cases, consisted of chloride of iron, aloes, extract of horehound, and assafetida.

Stahl's Pills, much used as a tonic aperient in Germany, are made as follows:—Ferri pulv., $\mathfrak{z}\mathfrak{ss}$; Aloes pulv., $\mathfrak{z}\mathfrak{j}$; Ext. colocynth., $\mathfrak{z}\mathfrak{j}$; mucilag. acaciæ q. s. ut fiant pil xl.

PILULÆ ALOES ET JALAP'Æ, commonly called Anderson's pills, Anderson's Scots' pills, Scotch pills. (Aloes Barbād. \mathfrak{ij} ; rad. helleb. nig., rad. jalapæ, potassæ subcarb. \mathfrak{aa} $\mathfrak{z}\mathfrak{j}$; ol. anisi, $\mathfrak{z}\mathfrak{j}$; syrup. simpl. q. s.—Gray.) Dose, gr. x to xxx.

A committee of the Philadelphia College of Pharmacy recommend the following: Aloes Barbādens. $\mathfrak{z}\mathfrak{xxiv}$; saponis. $\mathfrak{z}\mathfrak{j}$; colocynthid. $\mathfrak{z}\mathfrak{j}$; cambogiæ, $\mathfrak{z}\mathfrak{j}$; ol. anis. f. $\mathfrak{z}\mathfrak{ss}$.

PILULÆ ALOES ET KINÆ KINÆ, *P. stomachicæ* seu *ante cibum*, *Pill of aloes and cinchona*, (F.) *Pilules gourmandes*. (Aloes socotr. $\mathfrak{z}\mathfrak{vj}$; ext. cinchon. $\mathfrak{z}\mathfrak{ij}$; cinchon. $\mathfrak{z}\mathfrak{j}$; syrup. absinth. q. s.—Ph. P.) Stomachic and laxative. Dose, gr. x. to xx.

Lady Crespigny's pills, Lady Webster's pills, Lady Hesketh's pills; the *Pilules de Duchesne*, (Querquetanus), the *Grains de Vie* ou *Pilules de Clérambourg*; the *Pilules Indiennes stomachiques de De la Croix*, the *Pilules de Barbier*, and the *Pilulæ stomachicæ Menusès* (Ph. L. 1635), resemble these.

Frank's pills, much employed, also, as 'dinner pills,' or, —to use the language of the eccentric Kitchener—as 'peptic persuaders,' are said to be much used, under the name *Graines de santé*, in various parts of Europe. They consist of aloes, jalap, and rhubarb, with syrup of wormwood as an excipient.

PILULÆ ALOES MARTIALES, *P. de aloes et foetidis*.

PILULÆ ALOES ET MYRRHÆ, *P. aloes cum myrrhâ* seu *de aloë et myrrhâ*, *P. Rufi* seu *Rufi seu commu'nes*, *Rufus's* or *Ruffus's pills*, Common pills, *Pills of Aloes and myrrh*. (Aloes, $\mathfrak{z}\mathfrak{j}$; croci, in pulv. $\mathfrak{z}\mathfrak{ss}$; myrrh. $\mathfrak{z}\mathfrak{j}$; syr. q. s. to form 480 pills.—Ph. U. S.) Cathartic and emmenagogue. Dose, gr. x to xx.

Hooper's pills—possessed of similar properties—according to one of Gray's formulæ are made as follows: Ferri sulph., aquæ, \mathfrak{aa} $\mathfrak{z}\mathfrak{viij}$. Dissolve. Add aloes Barbād. \mathfrak{ij} and $\mathfrak{z}\mathfrak{viij}$; canellæ albæ, $\mathfrak{z}\mathfrak{vj}$; gum. myrrh. $\mathfrak{z}\mathfrak{j}$; opopanax, $\mathfrak{z}\mathfrak{j}$. The committee of the Philadelphia College of Pharmacy recommend the following form: Aloes Barbādensis, $\mathfrak{z}\mathfrak{viij}$; ferri sulphatis exsiccat. $\mathfrak{z}\mathfrak{j}$, and $\mathfrak{z}\mathfrak{ss}$, vel ferri sulph. crystall. $\mathfrak{z}\mathfrak{j}$; ext. helleb. nigr. $\mathfrak{z}\mathfrak{j}$; myrrh. $\mathfrak{z}\mathfrak{j}$; saponis, $\mathfrak{z}\mathfrak{j}$; canellæ, in pulv. tritæ, $\mathfrak{z}\mathfrak{j}$; zingib. in pulv. trit. $\mathfrak{z}\mathfrak{j}$.

Speediman's pills consist of aloes, myrrh, rhubarb, extract of chamomile, and essential oil of chamomile; and the Pills of Barthæz, of aloes, myrrh, musk, camphor, and balsam of Peru.

PILULÆ DE ALOË ET SAPONE, *P. aloeticæ*.

PILULÆ ALOES CUM ZINGIB'ERË, *Pills of aloes and ginger*. (Aloes hepat. $\mathfrak{z}\mathfrak{j}$; rad. zingib. in pulv. $\mathfrak{z}\mathfrak{j}$; sapon. Hispan. $\mathfrak{z}\mathfrak{ss}$; ol. ess. menthæ pip. $\mathfrak{z}\mathfrak{ss}$ —Ph. D.) In habitual costiveness. Dose, gr. x to xx.

PILULÆ AMMONIURE'TI CUPRI, *P. Cupri*, Pills of Ammoniumuret of copper, (F.) *Pilules d'ammonure de cuivre*. (Ammonium. cupri, in pulv. ten. trit., gr. xvj; micæ panis, $\mathfrak{z}\mathfrak{j}$; aquæ carbonatis

ammonia, q. s. Beat into a mass, and divide into xxxij pills.—Ph. E.) Tonic and antispasmodic, in epilepsy, &c. Dose, one pill twice daily; gradually increasing the dose.

PILULÆ ANTE CIBUM, P. aloes et kinæ kinæ.

PILULÆ ANTIBILIO'SÆ, *Barclay's Antibilious pills*. (Extract of colocynth, ʒij; resin of jalap. ʒi; almond soap, ʒiiss; guaiac, ʒij; emetic tart. gr. viij; oils of juniper, carraway, and rosemary, aa iv drops; syrup of buckthorn, q. s. to form 64 pills.) Dose, two or three.

Dixon's Antibilious pills are formed of aloes, scammony, rhubarb, and tartarized antimony.

PILULÆ ANTIHERPETICÆ, *Pilulæ hydrargyri submuriatis comp.*—p. Anthysterica, P. galbani compositæ.

PILULÆ ANTISCROPHULO'SÆ, *P. ex ox'ido stib'ii et hydrargyri sulphure'to nigro compos'itæ*. (P. scammon., hydrarg. sulphur. nigr. aa ʒij; antimoni. oxi'di albi, milleped., sapon. amygdalin. aa ʒij; extract. glycyrrh. ʒv. Make into a mass.—Ph. P.) Dose, gr. iv to xx.

PILULÆ ANTISPASTICÆ, P. galbani compositæ —p. Assafetida compositæ, P. galbani compositæ.

PILULÆ ASIATICÆ, *Asiatic or Tanjore pills*. (Acid. arsenios. gr. lv; piper. nigr. pulv. ʒix; acaciæ, q. s. ut fiant pil. 800. Each pill contains about one-fifteenth of a grain of arsenious acid. These pills are employed in India for the cure of syphilis and elephantiasis.)

PILULÆ ASSAFETIDÆ, *Assafetida pills*. (Assafetid. ʒiiss; saponis, ʒss. Beat with water so as to form a mass to be divided into 240 pills.—Ph. U. S.)

PILULÆ BALSAMICÆ, *P. Docto'ris Morton*. (Milleped. pulverisat. ʒxviij; gum ammon. ʒx; acid. benzoic. ʒvj; croci, balsami Peruv. aa ʒi; balsam. sulph. anteat. ʒvj. Mix.—Ph. P.) Tonic, expectorant, and balsamic.

PILULÆ BENEDICTÆ FULLERI, P. de aloë et foetidis.

PILULÆ CALOMELANOS COMPOSITÆ, P. Hydrargyri submuriatis compositæ.

PILULÆ CALOMEL'ANOS ET OPII, *P. Hydrargyri chlor'idi et op'ii*, *Pills of Calomel and Opium*. (Hydrarg. chlorid. mit. p. iij; Opij p. j; Confect. Ros. q. s. To be divided into pills each containing two grains of calomel.—Ph. Ed.)

PILULÆ CAMBO'GIÆ COMPOS'ITÆ, *P. de guttâ gaman'drâ seu hydrog'gæ seu gummi guttæ aloeticæ*, *Gamboge pills*, *Compound gamboge pills*, (F.) *Pilules de gomme gutte composées*. (Cambogiæ contrita, aloes epicat. extract. contriti., pulv. cinnam. comp. sing. ʒj; saponis, ʒij.—Ph. L.) Cathartic. Dose, gr. x to xx.

Morrison's Pills are said by Dr. Paris to be nothing more than the above, with the addition of a small portion of cream of tartar.

PILULÆ CATHARTICÆ COMPOS'ITÆ, *Compound cathartic pills*. (Ext. colocynth. comp. in pulv. ʒss; ext. jalapæ in pulv. hydrarg. chlorid. mit. aa ʒij; gambog. in pulv. ʒij. M. Form into a mass with water, and make into 180 pills.—Ph. U. S.) Three for a dose.

PILULÆ COCCII, P. aloes et colocynthidis —p. Coccia minores, P. aloes et colocynthidis —p. Coccina, P. aloes et colocynthidis —p. Colechici et digitalis, P. Lartigue's—p. Colocynthidis compositæ, P. aloes et colocynthidis—p. ex Colocynthide cum aloë, P. aloes et colocynthidis.

PILULÆ COLOCYNTHIDIS ET HYOSCY'AMI, *Pills of Colocynth and Henbane*. (Phil. Colocynth. comp. p. ij; Ext. Hyoscyam. p. i.—Ph. E.) Dose, gr. v to gr. xx.

PILULÆ COMMUNES, P. aloes et myrrhæ.

PILULÆ CONII COMPOS'ITÆ, *Compound Pill of Henlock*. (Ext. Coni ʒv; Ipecac. pulv. ʒj; Theriac. q. s.—Ph. L.) Dose, gr. v three times a day.

PILULÆ CUPRI, P. ammoniuretici cupri.

PILULÆ COPA'IBÆ, *Pills of Copaiba*. (Copaib. ʒij; magnesiae, recenté præpar. ʒi. Mix, and set aside till it concretes into a pillular mass, to be divided into 200 pills.—Ph. U. S.) Each pill contains nearly five grains of copaiba. Two to six for a dose.

PILULÆ CUPRI AMMONIA'TI, *Pills of ammoniated Copper*. (Cupri ammon. in pulv. p. j; micæ panis, p. vj; Liq. ammon. carbonat. q. s.—Ph. E.) Dose, one to six pills.

PILULÆ DE CYNOGLOS'SO, *P. de extracto o'pii*. (Pulv. rad. cynogloss., semin. hyoscyam. alb., extract. opii vinos. aa ʒiv; pulv. myrrh. ʒvj; oliban. ʒv; croci, castor., aa ʒiiss; syrup opii, q. s.—Ph. P.) Anodyne. In cough, &c. Dose, gr. iv to gr. viij.

PILULÆ DEPURANTES, P. hydrargyri submuriatis compos.

PILULÆ DIGITA'LIS ET SCILLÆ, *Pills of Digitalis and Squill*. (Digital., Scillæ, aa p. j; Elect. aromat. p. ij.—Ph. E.) One or two pills for a dose.

PILULÆ ECPHRACTICÆ, P. de aloë et foetidis.

PILULÆ FERRI CARBONA'TIS, *Pills of carbonate of iron*, *Vallet's pills*, *Vallet's ferruginous pills*. (Ferri sulph. ʒviij; sodæ carbonat. ʒx; Mel. deepumat. ʒiiij; sacchar. ʒij; aquæ bullientis, Oij; syrup. q. s. Dissolve the sulphate of iron and carbonate of soda, each in a pint of the water, and to each solution add a fluidounce of syrup; then mix the two solutions in a bottle just large enough to contain them; close it accurately with a stopper, and set it by, that the carbonate of iron may subside. Pour off the liquid, and having washed the precipitate with warm water, sweetened with syrup in the proportion of f ʒj of the latter to Oj of the former, until the washings no longer have a saline taste, place it upon a flannel cloth, and express as much water as possible; then mix it with the honey and sugar. Lastly, heat the mixture in a water-bath, until it attains a pillular consistence.—Ph. U. S.)

PILULÆ FERRI COMPOS'ITÆ, *P. ferri cum myrrhâ*, *Compound pills of iron*, *Griffith's Pills*, (F.) *Pilules de fer composées*. (Myrrh. in pulv. ʒij; sodæ subcarb., ferri sulph. aa ʒj; syrup. q. s. Mix, to form 80 pills.—Ph. U. S.) Tonic. Dose, gr. x to xx.

PILULÆ FERRI IOD'IDI, *Pills of Iodide of iron*. (Ferri sulphat. ʒj; Potass. ioidid. ʒiv; tragacanth. pulv. gr. x; sacchar. pulv. ʒss. Beat with syrup into a mass, to form 40 pills.—Ph. U. S.) Dose, one to three where a tonic and eutrophic is indicated.

PILULÆ FERRI CUM MÏRRHÂ, P. Ferri compositæ.

PILULÆ FERRI SULPHA'TIS, *Pills of Sulphate of Iron*. (Ferri sulph. exsic. p. ij; Ext. Taraxac. p. v; Conserv. Rosæ p. ij; Glycyrrhiz. pulv. p. iij.—Ph. E.) Dose, gr. v to gr. xx.

PILULÆ FETIDÆ, P. galbani compositæ.

PILULÆ GAL'BANI COMPOS'ITÆ, *P. assafetida compos'itæ seu foetidæ seu gummo'sæ seu e gummi seu antispas'ticæ seu antihyster'icæ seu myrrhæ compos'itæ*, *Compound galbanum pill*, *Gum pill*. (Galban., myrrhæ, aa ʒvj; assafetida, ʒij; syrup. q. s. to form 240 pills.—Ph. U. S.) Antispasmodic and emmenagogue. Dose, gr. x to xx.

PILULÆ GUMMI, P. galbani compositæ —p. Gummi guttæ aloeticæ, P. cambogiæ compositæ —p. Gummosæ, P. galbani compositæ —p. de Guttâ gamandrâ, P. cambogiæ compositæ.

PILULÆ EX HELLE'ORO ET MYRRHÂ, *P. tonicæ*, *Bacher's pills*, *Tonic pills of Bacher*, *Pills of hellebore and myrrh*. (Ext. helleb. nig., myrrh. aa ʒj; fol. cardui benedict. in pulv. ʒij. M.—Ph. P.) Tonic and emmenagogue.

In dropsical, cutaneous, and verminous affections. Dose, gr. iv to gr. viij.

Matthew's pills, — *Starkey's pills*, are formed of rad. helleb. nigr., rad. glycyrrh., sapon. Castill., rad. curcumæ, opii purif., syrup. croci, 3i; ol. tereb. q. s. ut fiat massa. — Gray.

PILULÆ HYDRAGOGÆ, P. cambogiæ compositæ — p. Hydragogæ Bontii, Pilulæ aloes et cambogiæ.

PILULÆ HYDRARGYRI, P. mercurialis, *Mercurial* or *Blue pill*, (F.) *Pilules de Mercure*. (Hydrag. purif. 3j; conf. rosæ, 3ss; glycyrrh. pulv. 3ss. Rub the quicksilver with the confection, until the globules disappear; then add the liquorice powder, and beat the whole into a uniform mass, to be divided into 480 pills. — Ph. U. S.) Antisyphilitic; mercurial; — in large doses, purgative.

A purgative pill, much prescribed by Mr. Abernethy, and called *Abernethy's pill*, consisted of pil. hydrarg. gr. x; jalap. pulv. gr. xx; syrup. rhamni, q. s. ut fiat pil. vj. Two were given at night, and the next morning a wineglassful of infusum sennæ.

PILULÆ HYDRARGYRI CHLORIDI ET OPII, P. Calomelanos et Opii.

PILULÆ HYDRARGYRI CHLORIDI MITIS, *Pills of mild chloride of mercury*, *Calomel pills*. (Hydrag. chlorid. mit. 3ss; acaciæ, in pulv. 3j; syrup. q. s. ut fiat pil. 240. — Ph. U. S.) Each pill contains one grain.

PILULÆ DE HYDRARGYRO, SCAMMONIO ET ALOE, vulgo *MERCURIAL'ES*, (F.) *Pilules Mercurielles ou de Mercure, Scammonée et Aloes*. (Hydrag. purif. 3j; mellis, 3xj. Rub until the globules disappear; then add aloes Socot., scammon. 3j; mæcis, cinnam. 3j. Make into a mass. — Ph. P.) Uses, same as the last, but they are more purgative. Dose, gr. viij to xxx. These pills are nearly the same as the *Pastilles de Renaudot*.

PILULÆ EX HYDRARGYRO MURIATICO MITI CUM SULPHURE STIBIATO AURIANTIACO, Pilulæ hydrargyri submuriatis compositæ.

PILULÆ HYDRARGYRI SUBMURIATIS COMPOSITÆ, P. ex stibio et hydrargyro seu antihæpeticæ seu ex hydrargyro muriatico miti cum sulphure stibiatō auriantiacō seu resolutæ cum mercurio seu depuratæ seu hydrargyri submuriatis, Pilulæ Hydragyri chloridi compositæ, Pilulæ calomelanos compositæ, Compound Calomel pill, Compound pills of submuriate of mercury, Calomel pills, Plummer's pills; Red pill. (Hydrag. submur., antimon. sulphur. precipitat. sing. 3ij; guaiac. gum resin. contrit. 5iv; spirit rect. 3ss. Beat together. — Ph. L.) Mercurial, and reputed to be diaphoretic and alterative. Dose, gr. v to x.

Leake's pilula salutaria resembles these. It is used in syphilis.

PILULÆ HYDRARGYRI SUBMURIATIS ET OPII, P. Calomelanos et Opii.

PILULÆ IPECACUAN'Æ PULV ET OPII, *Pills of Ipecacuanha and Opium*. (Pulv. Ipecac. et Opii, p. iij; Conserv. Rosæ, p. j. — Ph. E.) One grain of opium in about thirteen grains.

PILULÆ IPECACUAN'Æ CUM SCILLÂ, *Pills of Ipecacuanha with Squill*. (Pulv. Ipecac. comp. 3ij; Scillæ pulv., Ammoniac. in pulv. 3j; Theriac. q. s. — Ph. L.) Dose, gr. v to x.

PILULÆ MERCURIALES, P. hydrargyri, P. de hydrargyro, scammonio et aloë — p. Doctoris Morton, P. balsamicæ — p. Myrrhæ compositæ, P. Galbani composiæ.

PILULÆ OPIATÆ, Pilulæ Opii (Ph. U. S.), P. *Thebæicæ*, *Opiate pills*, *Common night pills*, *Anodyne pills*, (Prov.) *Squatting pills*, *Nepenthes opiatum*, (F.) *Pilules opiacées*. (Opii pulv. 3j; sapon. gr. xij; to be made into 60 pills. — Ph. U. S.) Each pill contains a grain of opium.

PILULÆ DE EXTRACTO OPII, P. de cynoglossio

— p. ex Oxido stibii et hydrargyri sulphureto nigro composiæ, Pilulæ antiscrophulosæ.

PILULÆ PLUMBI OPIATÆ, *Opiate Pills of Lead*. (Plumb. acetat., p. vj; Opii, p. j; Conserv. Rosæ, p. j. To be made into four grain pills. — Ph. E.) Each pill contains three grains of acetate of lead.

PILULÆ RESOLVENTES CUM MERCURIO, P. hydrargyri submuriatis compos. — p. de Rhabarbaro, Pilulæ rhei composiæ.

The stomachic nostrum, sold under the name of *Moseley's pills*, consists merely of rhubarb and ginger.

PILULÆ QUIN'Æ SULPHATIS, *Pills of sulphate of quinia*. (Quin. sulph. 3j; acaciæ, in pulv. 3ij; mel. q. s. ut fiat pilulæ 480. — Ph. U. S.) Each pill contains a grain of the sulphate of quinia.

PILULÆ RHEI, *Pills of Rhubarb*. (Rhej pulv. 3vj; saponis, 3ij. M. et fiat pil. 120. — Ph. U. S.) Each pill contains three grains of rhubarb.

A good dinner or stomachic pill is made of Rhej pulv. 3ij; Capsic. pulv. 3j; make into a mass with molasses, and divide into 60 pills. Dose, two or three before dinner.

PILULÆ RHEI COMPOSITÆ, P. de Rhabarbaro, Compound rhubarb pills. (Rhej, pulv. 3j; aloes, pulv. 3vj; myrrh. pulv. 3iv; ol. menthæ pip. t 3ss. Beat into a mass, with syrup of orange-peel, to form 240 pills. — Ph. U. S.) Laxative and stomachic. Dose, gr. x to xx.

PILULÆ RHEI ET FERRI, *Pills of Rhubarb and Iron*. (Ferri sulph. exsicc. p. iv; Est. Rhej, p. x; Conserv. Rosæ, p. v. Make into five grain pills.) Dose, as a tonic and laxative, two or three pills.

Griffitt's pills, so called after Dr. S. P. Griffitts, of Philadelphia, who prescribed them as a tonic aperient, were formed as follows: — Rhej, pulv. 3ss; ferri sulph. 3ss; saponis, 3ij; aquæ, q. s. ut fiat massa in pil. xl dividenda. Dose, three or four at bedtime.

PILULÆ RUFI, P. aloes et myrrhæ — p. Salutaris, Leake's, see Pilulæ hydrargyri submuriatis composiæ — p. Saponacæ, P. saponis cum opio — p. Saponis composiæ, P. saponis cum opio.

PILULÆ SAPO'NIS COMPOSITÆ, P. *Saponacæ* seu saponis composiæ, *Pills of soap and opium*, (F.) *Pilules de savon et d'opium*. (Opii, 3ss; sapon. 3ij. M. — Ph. U. S.) Five grains contain one of opium.

The Pharmacopœia of the United States has a form for the *Pilulæ opii*, *Opium pills*, as well as for the *Pilulæ saponis composiæ*; the only difference between them being in the proportion of the ingredients. See Pilulæ opiate.

PILULÆ E SCILLÂ, P. scillæ composiæ.

PILULÆ SCILLÆ COMPOSITÆ, P. *scilliticæ* seu *scillæ cum zingib'ere* seu *e scillâ*, Compound squill pill. (Scillæ, pulv. 3j; zingib. pulv., ammoniac. pulv. 3j; saponis, 3ij; syrup. q. s. ut fiat pil. 120. — Ph. U. S.) Dose, two pills. Expectant and diuretic. Dose, gr. x to xx.

PILULÆ SCILLÆ CUM ZINGIBERE, P. scillæ composiæ — p. Scilliticæ, P. scillæ composiæ — p. ex Stibio et hydrargyro, P. hydrargyri submur. compos. — p. Stomachicæ, P. aloes et kinæ kinæ.

PILULÆ STRYCH'NÆ, *Pills of Strychnia*. (Very pure strychnine, 2 gr.; conserve of roses, 3ss. Make into 24 pills.) One or two for a dose.

PILULÆ STY'RACIS, P. S. Composiæ, *Storax Pills*. (Styracis pur. 3vi; opii, croci, sing. 3ij. — Ph. E.) Five grains contain one of opium.

PILULÆ SUBLINGUALES, Hypoglottidæ.

PILULÆ DE TEREBINTHINÂ, *Turpentine Pills*. Made of turpentine boiled in 3 times its weight of water, until, when thrown into cold water, it forms a soft paste, which has the properties of turpentine, and has been administered in gleet, &c.

PILULÆ THEBAICÆ, P. opiatæ — p. Tonicæ, P. ex Helleboro et myrrha.

PIL'ULAR, *Pilula'ris*, (F.) *Pilulaire*, from *pilula*, 'a pill.' Relating to pills, — as 'a *pilular* mass,' 'pilular form,' &c.

PILULES, *Pilulæ* — p. *Agrégatives*, see Aggregate — p. *d'Aloes ou aloétiques*, *Pilulæ aloeticæ* — p. *d'Ammoniaque de cuivre*, *Pilulæ ammoniuretici cupri* — p. *de Barbier*, see *Pilulæ Aloes et kinæ kinæ* — p. *Bénites* of Fuller, *Pilulæ de aloë et fetiditis* — p. *Cochées*, *Pilulæ aloes et colocynthidis* — p. *de Duchesne*, see *Pilulæ Aloes et kinæ kinæ* — p. *de Fer composées*, *Pilulæ ferri compositæ* — p. *Ferrugineuses de Vallet*, see *Ferri protocarbonas* — p. *Gourmandes*, *Pilulæ aloes et kinæ kinæ* — p. *Indiennes stomachiques de Delacroix*, see *Pilulæ aloes et kinæ kinæ* — p. *de Lartigue*, *Pill*, *Lartigue's* — p. *de Mercure*, *Pilulæ hydrargyri* — p. *de Mercure*, &c., *Pilulæ de hydrargyro*, et scammonio et aloë — p. *Mercurielles*, *Pilulæ de hydrargyro*, scammonio et aloë — p. *Opiacées*, *Pilulæ opiatæ* — p. *Panchymagogues*, *Pilulæ aloes et colocynthidis* — p. *de Savon et d'Opium*, *Pilulæ saponis cum opio* — p. *Universelles*, *Pilulæ aloes et colocynthidis*.

PILULIER (F.), *Pill Machine*. An instrument for rolling and dividing pills. Also the earthen pot in which pills are preserved.

PILUM, *Pistillus*, *Pistillum*, *Hypēros*, *Cer'cia*, (F.) *Pilon*. A pestle. A wooden, iron, bell-metal, marble, agate, porcelain, or glass instrument for beating or dividing substances in a mortar.

A *spring pestle* is one attached above to a spring, so as to take off the weight, and thus lessen the labour of pounding.

PILUS, *Capillus*, *Caul*, *Hair*.

PIMELE, *Fat*.

PIMELECHYSIS, *Pimelorrhœa*.

PIMELICUS, *Fatty*.

PIMELI'TIS, from *πυελν*, 'fat,' and *itis*, denoting inflammation. Inflammation of the adipous tissue.

PIMELODES, *Fatty*.

PIMELO'MA, from *πυελν*, 'fat.' A fatty swelling.

PIMELORRHÆ'A, *Pimeleorchysis*, from *πυελν*, 'fat,' and *ρρω*, 'I flow.' A morbid discharge or disappearance of fat.

PIMELORTHOPNÆ'A, *Piorthopnoë'a*; from *πυελν*, 'fat,' *ορθος*, 'upright,' and *πνευω*, 'to breathe.' Dyspnoea, practicable only in the erect attitude, owing to fat.

PIMELO'SIS, *Obesitas nim'ia*, from *πυελν*, 'fat.' Conversion into, or formation of, fat, — as *Pimelo'sis hepatic'a*, 'fatty degeneration of the liver.'

PIMELOSIS HEPATICA, *Adiposis hepatica* — p. *Nimia*, see *Polysarcia*.

PIMENT, *Capsicum annum* — p. *d'Eau*, *Polygonum hydropiper* — p. *Royal*, *Myrica gale*.

PIMENTA, see *Myrtus pimenta*.

PIMENTÆ BACCÆ, see *Myrtus pimenta*.

PIMENTO, see *Myrtus pimenta*.

PIMPERNEL, BLUE, *Scutellaria lateriflora* — p. Red, *Anagallis* — p. Scarlet, *Anagallis arvensis* — p. Water, *Veronica beccabunga*.

PIMPILIM, *Piper longum*.

PIMPINEL, (*bipinnella* or *bipennula*, from the double pennate order of its leaves), see *Pimpinella*.

PIMPINEL ITALIAN, *Sanguisorba officinalis*.

PIMPINELLA ALBA, P. saxifraga — p. *Angelicaefolia*, *Ligusticum podagraria*.

PIMPINEL'LA ANI'SUM, *Anise plant*, *Anisum* (Ph. U. S.), *A. officinalis* seu *vulgaris* *Sison ani'sum*, *A'pium ani'sum*, *Sola'men istiodo'rum*, *Ane'sum*, (F.) *Anis*, *Boucage anis*. Ord. Umbelliferae. Sex. Syst. Pentandria Digynia. A native of Egypt. The seeds, *Ani'sum*, (Ph. U. S.) *Sem'ina*

Ani'si, *Ani'seed*, (Old Eng.) *Aneys*, have an aromatic odour; and a sweetish, warm, grateful taste. They are carminative, and used in dyspepsia and in tormina of infants. Dose, gr. xv to ʒj, bruised. The oil, — OLEUM ANI'SI, (F.) *Huile d'anis* — is official. Dose, gtt. v to xv. Most of the oil used in this country under the name of *Oil of Anise* is said to be obtained from *Illicium Anisatum*.

Aqua ani'si, *Anise water*, (Ph. D.) is made by mixing a fluidounce of essence of anise, with half a gallon (Imp. meas.) of distilled water, and filtering through paper.

PIMPINELLA, GREATER, P. magna — p. *Hircina*, P. Saxifraga — p. *Laterifolia*, *Sion ammi*.

PIMPINEL'LA MAGNA seu *nigra* seu *major* seu *rubra*; Greater *Pimpinell'la*; *Tragoseli'num magnum* seu *majus*, *Daucus*, *Cyan'opus*, (F.) *Boucage majeur*. The root has been extolled as diuretic and resolvent, as well as in erysipelatous ulcerations, tinea capitis, rheumatism, &c.

PIMPINELLA MAJOR, P. magna — p. *Nigra*, P. magna — p. *Nostras*, P. saxifraga — p. *Officinalis*, *Sanguisorba officinalis* — p. *Rubra*, P. magna.

PIMPINELLA SAXIFRAGA, *Saxifraga*, *Burnet saxifraga*, *Poterium sanguisor'ba*, *Sorbastrel'la*, *Tragoseli'num*, *T. saxifraga*, *Pimpinell'la hirci'na* seu *umbellifera* seu *alba* seu *nostras*, (F.) *Boucage mineur*, *Petit B.*, *Pimprenelle*, *Rompt-pierre*. The root has an unpleasant smell, and hot, pungent, bitterish taste. It has been recommended as a stomachic; and as a stimulating gargle in paralysis of the tongue.

PIMPINELLA UMBELLIFERA, P. saxifraga.

PIMPLE, *Papula*.

PIMPLED, *Papulose*.

PIMPRENELLE, *Pimpinella saxifraga* — p. *Noire*, *Sanguisorba officinalis*.

PIN. Perhaps from *pennum*, low Latin; or from *spina* [?], *Spina fer'rea*, *Spi'mula*, *Acus capitata*, (F.) *Épingle*. An iron or brass instrument, generally of a small size, pointed at one extremity, and having a head at the other. It is used, in Surgery, to fix rollers and dressings; and occasionally in sutures.

PIN AND WEB, *Caligo*.

PIN À PIGNON, see *Pinus picea*.

PINASTELLUM, *Peucedanum*.

PINASTER, *Pinus sylvestris*.

PINBONE, *Ischion*.

PINCE, Forceps — p. *d'Anneaux*, see *Forceps* — p. *d'Dissection*, see *Forceps* — p. *d'Ligature*, see *Forceps* — p. *Lithodrassique*, see *Lithodrassic* — p. *Panement*, see *Forceps*.

PINCÉE, *Pinched*.

PINCÉE, *Pugillus*.

PINCERS, *Forceps*.

PINCETTE, *Forceps*.

PINCED, *Contractus*, *Retrac'tus*, (F.) *Pincé*, *Grippé*. An epithet applied to the face, when the features are contracted or shrunken; as in violent abdominal affections, or during intense pain.

PINCKNEY'A PUBENS, P. *Pubescens*, *Georgia Bark*, *Bitter Bark*, *Florida Bark*, *Fever tree*. This bark has been used in Georgia in the cure of intermittents, and successfully. It is a powerful bitter. Dose of the powder, ʒj.

PINDARS, *Arachis hypogæa*.

PINE APPLE, *Bromelia ananas* — p. *Aphernousli*, *Pinus cembra* — p. *Ground*, *Lycopodium complanatum* — p. *Ground*, stinking, *Camphoresma Mospeliaca* — p. *Loblolly*, *Pinus taeda* — p. *Mountain*, see *Pinus mughos* — p. *Mugho*, see *Pinus mughos* — p. *Prince's*, *Pyrola umbellata* — p. *Sap*, *American*, *Hypopitys lanuginosa* — p. *Stone*, *Pinus pinea* — p. *Sugar*, see *Arrow Root* — p. *Yellow*, *Pinus palustris*.

PINEA, *Pinus pinea*.

PINE'AL, *Pineal'is*, from *pinus*, 'a pine.' That which resembles a pine-apple.

PINEAL GLAND, *Glandula pineal'is*, *Cerebral epiph'ysis*, *Cona'rrium*, *Cono'ides corpus*, *Corpus turbinatum*, *Penis* seu *Virga* seu *Turbo cer'e'br*, is a small body, of a conical shape; pale red or grayish colour; and soft consistence; situate between the fornix and the tubercula quadrigemina. It almost always contains sabulous particles, *Sab'ulum cona'rri*, *Brain Sand*, *Pineal Sand*. When these are grouped together over the base of the gland, they form the *Acer'vulus Cer'e'br* of Sömmerring, *Acer'vuli* seu *Lapil'li glan'dulæ pineal'is*. From its anterior parts arise two medullary striae, which proceed over the posterior commissure; coast along the optic thalami, and unite at the anterior pillar of the fornix,—the *Habe'na*, *Haben'ulæ* or *reins of the pineal gland*—*Pedunculi cona'rri*, see *Peduncles of the Pineal Gland*.

The uses of the pineal gland are unknown. Descartes supposed it to be the seat of the soul!

PINEAL SAND, see *Pineal gland*.

PINEL, see *Pinus picea*.

PINEOLI, see *Pinus picea*.

PINEUS PURGANS, *Jatropha curcas*.

PINEWEED, *Hypericum sarothra*.

PINGUEC'ULA, *Pteryg'um pingue* seu *lar-da'ceum*, from *pinguis*, 'fat,' 'fatty.' A small, whitish-yellow tumour in the sclerotic conjunctiva and subjacent areolar tissue, close to the margin of the cornea on its nasal or temporal side; so called from its being supposed, but erroneously, to be of a fatty nature.

PINGUEDO, Fat—*p. Renalis*, *Nephridion*.

PINGUIC'ULA VULGARIS, (*pinguis*, 'fat,') *P. alpi'na*, *Sanic'ula montana* seu *Eboracen'sis*, *V'ola palus'tris*, *Dodecat'hleon Plin'ii*, *Butterwort*, *Yorkshire San'icle*. *Ord. Lentibulacæ*. *Sec. Syst. Lamiæ Monogynia*. The unctuousity of this plant has caused it to be applied to chaps, and as a pomatum to the hair. Decoctions of the leaves, in broths, are used by the common people in Wales as a cathartic.

PINGUID, Fatty.

PINGUIDINOUS, Fatty.

PINGUIN, *Bromelia pinguin*.

PINGUITUDO, *Polysarcia adiposa*.

PINHOLE PUPIL, see *Pupil*, *pinhole*.

PINHONES INDICI, *Jatropha curcas*.

PINK, *CAROLINA*, *Spigelia Marilandica*.

PINK DYE. *Stripped safflower*, $\frac{3}{4}$, *subcarb.* of *potas.* gr. xviii, *spirit of wine*, $\frac{3}{4}$ vi; digest for two hours; add *distilled water*, $\frac{3}{4}$ vi; digest for two hours more, and add *distilled vinegar* or *lemon juice* q. s., to reduce it to a fine rose-colour. Used as a cosmetic.

PINK, *GROUND*, *Silene Virginica*—*p. Indian*, *Spigelia Marilandica*—*p. Wild*, *Silene Virginica*.

PINKROOT, *Spigelia marilandica*—*p. Deme-rara*, *Spigelia anthelmia*.

PINNÄ, *Ala*, *Pavilion of the ear*—*p. Marina*, see *Bissus*.

PINNACULUM FORNICIS GUTTURALIS, *Uvula*.

PINNÆ HEPATIS, *Lobes of the liver*—*p. Naris*, see *Nasus*.

PINNULA, *Aileron*.

PINNULÆ HEPATIS, *Lobes of the liver*.

PINO'LI (S.). A preparation used for subsistence on long journeys in the West. It is parched corn, beaten or ground very fine, and sweetened with sugar, to be used with water, and drunk on the march. It is the cold flour of the Indians, and early Western pioneers.

PINONCILLO TREE, *Castillognia lobata*.

PINSWEAL, *Furunculus*.

PINT, *Octarius*.

PINTA, from (S.) *pintar*, 'to paint.' *Blue stain*: a disease, which is said to prevail in Mexico, and to be a variety of *Pityriasis versicolor*.

PINUS A'BIES, *P. excel'sa* seu *pice'a*, *A'bies*, *A. rubra* seu *excelsa* seu *picea*, *El'atè thelei'a*, *Pice'a*, *Norway Spruce Fir*, *Yew-leaved Fir*. *Nat. Ord. Coniferæ*. *Sec. Syst. Monocia Monadelphica*. The tops are used in making *Spruce Beer*.

Essence of Spruce, *Essen'tia Abie'tis*, is a fluid extract, prepared by decoction from the twigs of this species of fir. From this is made *Spruce Beer*.

This fir affords the Burgundy pitch and common frankincense. 1. *Bur'gundy* or *White pitch*, *Piz Burgun'dica* seu *alba* seu *ar'ida* seu *abiet'i'na* seu *abie'tis* (Ph. U. S.), *Resi'na abie'tis hu'mida*, *R. alba humida*, *Pini abie'tis resina sponte concre'ta*, (F.) *Pois blanche* ou *grasse* ou *jaune* ou *de Bourgogne*. This prepared concrete juice is of a close consistence, rather soft, of a reddish-brown colour, and not unpleasant smell. It is very adhesive to the skin; and, consequently, forms excellent plasters for remaining upon the part for some time;—as in cases of affections of the chest. 2. *Abie'tis Resi'na* (Ph. L. since 1809), *Common Frankincense*, *Perrosin*, *Thus*, *Th. femini'num* seu *vulgar'e*, *Olib'anum vulgare* seu *ayles'trè*, *Resina abie'tis si-ca*, *Resin of the Spruce Fir*. It is solid, dry, brittle; externally, brownish-yellow; internally, whitish. Used in plasters like the last.

PINUS ABIES, see *P. picea*—*p. Alba*, *P. picea*.

PINUS BALSAME'A, *A'bies balsame'a* seu *balsamifera*, *Peu'cea balsame'a*, *Balsam fir*. The tree which affords the *Canada Balsam*, *Balsamum Canaden'se* seu *de Can'ada*, *Resina strobili'na*, *Terebinth'ina Canaden'sis* seu *Pini balsame'a*, *Canada Turpentine* or *Balsam*, *Balsam of Fir*, (F.) *Baume de Canada*, is one of the purest turpentine. It has the common properties of those substances.

PINUS CANADEN'SIS, *Abies Canaden'sis*, *Hemlock*, *Hemlock Spruce*. A tree, which is abundant in Canada, Nova Scotia, and the more northern parts of New England, and is found in the elevated and mountainous regions of the Middle States. The pitch—*Piz Canaden'sis*, *Canada Pitch*, *Hemlock Pitch*—obtained from it is commonly known under the name *Hemlock Gum*.

PINUS CANDICANS, *P. Picea*.

PINUS CEMERA, *P. Monta'na*, *Aphernous'li Pine*. It yields an agreeably scented turpentine:—the *Carpa'thian Balsam*, *Bal'samum Carpa'thicum* seu *Lib'ani*, *Carpa'thicum*, *Briançon Turpentine*. The nuts, *Cembro nuts*, *Nu'clei Cem-bræ*, have an eatable kernel, and yield oil. The shoots yield *Riga Balsam* by distillation.

PINUS DAMAR'RA, *Ag'athis Damarra*, grows in the East India Islands. The juice speedily concretes into a very hard resin,—the *Damarra turpentine*.

PINUS EXCELSA, *P. abies*—*p. Gallica*, *P. picea*.

PINUS LARIX, *A'bies larix*, *Larix*, *L. commu-nis* seu *decid'na* seu *Europæ'a* seu *pyramidal'is*. The *Larch*, (F.) *Mélèze*. From this tree exudes *Or'emberg gum*, *Bri'ançon manna*, *Gummi lar'icis* seu *Oremburgen'se* seu *Uralen'se*, *Manna Briganti'na* seu *larice'a*. It also yields, by boring, *Common Venice Turpentine*, *Resina Lar'icis*, *Terebin-thina Ven'eta* seu *larici'na*, *Pini Lar'icis resina liq'uida*, (F.) *Térébenthine de Vénise* ou *de Mè-lèze*. It is usually thinner than the other sorts of turpentine.

PINUS MUGHOS. The *Mountain* or *Mugho Pine*, *Pinus Pumil'io*. From the extremities of the branches exudes the *Resi'na Strobili'na*, of the Germans, or *Hungarian Balsam*. It is also obtained, by expression, from the cones. By dis-

tillation, the Hungarian balsam affords the *Krumholz oil*, *Oleum Templinum*.

PINUS PALUSTRIS, see *Pinus sylvestris*.

PINUS PICE'A, *P. Abies*, *A'bies*, *A. pice'a* seu *pectinata* seu *Gall'ica* seu *alba* seu *can'dicans* seu *vulgar'is* seu *taxifolia*, *European Silver Fir Tree*, *Él'atè*, (F.) *Sapin commun.* By piercing the tubercles of the bark of this fir, the *Strasburg Turpentine* is obtained:—the *Res'ina Abie'tis* (Ph. L. before 1809), *Oleum Abietis*, *Terebinth'ina Argentonensis*.

PINUS PINASTER, see *P. sylvestris*.

PINUS PI'NEA, *Stone Pine*, *Pinea*, *Pinus*, *P. uber'ima* seu *sati'va*. The nuts, *Zirbel nuts*, *Pine nuts*, *Nu'clei Pi'neæ*, *Pinei*, *Pine'oli*, (F.) *Pin à pignons*, are eaten raw, or preserved like almonds. They are, also, used in emulsions.

PINUS PUMILIO, see *Pinus Mughos*—*p. Sativa*, *Pinea*.

PINUS SYLVES'TRIS, *Pinas'ter*, *Peuœ*, *Scotch Fir*. This pine, as well as *P. marit'ima* (*P. Pinaster*), and other species of *Pinus*, affords common turpentine and its oil, resin, tar, and pitch. 1. *Common Turpentine of Europe*, *Terebinth'ina*, *T. vulgar'is* seu *commu'nis*, *Resina pini*, *Bijon*, *Horse Turpentine*, *Bordeaux Turpentine*, (F.) *Térébithine de Bordeaux*, *Térébithine commune*, is obtained by wounding the tree in hot weather. It is used, chiefly, as a dressing for wounds, &c., in horses, and for the distillation of the oil, (see *Oleum Terebinthinæ*.) The *white Turpentine*, or *common Turpentine of America*—*Terebinth'ina*, Ph. U. S.—is produced chiefly from *Pinus palustris*, *Yellow pine*, and *P. taeda*, *Loblolly pine*, and perhaps from other species inhabiting the Southern States. When the oil is distilled with water, yellow resin, or *Rosin*, (*Res'ina*, Ph. U. S.)—*Resina flava*—is left, which is only used in the formation of ointments and plasters. If without the addition of water, the residuum is *common resin* or *Colophony*. 2. When the cold begins to check the exudation of the common turpentine, part of it concretes in the wounds. This is called, in France, *Gallipot*, *Barras*; and *White Rosin*, *Resina alba*, when hardened after long exposure to the air. 3. When the old trees are subjected to distillation, in a coarse manner, Tar is obtained—*Hygropissos*, *Pissa*, *Pice'cedria* seu *liq'uida*, *Res'ina pini empyreumatica liq'uida*, *Terebinth'ina empyreumatica*, *Alchit'ra*, *Alchit'ra*, *Cedria*, (F.) *Goudron*, *G. végétale*, *Poiz liquide*, *Brat liquide*. *Tar water*, *Aqua Pice'a* seu *Picis*, *Infusum Picis liq'uide* seu *Picis empyreumaticæ liq'uide*, *Potio pice'a*, (F.) *Eau de Goudron*, which is made by mixing two pints of tar with a gallon of water, stirring with a wooden rod for fifteen minutes, and, when the tar has subsided, straining and keeping in well stopped bottles, was, at one time, a fashionable remedy in numerous complaints, and its use has been revived, since its virtues have been presumed to be owing to creasote. It is employed chiefly in pulmonary affections, and the vapour has been recommended in phthisis and other diseases of the lungs. It is used externally as a detergent. A *wine* or *beer* of tar, *Tarbeer*, *Jews'beer*, has been employed in Philadelphia in similar cases. It may be made by adding to tar enough beer to saturate it, mixing, and allowing them to macerate for three days, with occasional agitation. The dose is a table-spoonful. 4. *Common Pitch*, *Pice nigra*, *Black Pitch*, *Burnea*, *Burnia*, *Stone Pitch*, *Pice sicca*, *P. atra*, *P. nava'lis*, *Zopissa*, *Patipiss'ea*, *P. ar'ida* (Ph. L. before 1809), (F.) *Poiz navale*, *P. no'ire*, is obtained by inspissating tar. It is used only as a resolvent in plasters.

PINUS TÆDA, see *P. sylvestris*—*p. Taxifolia*,

P. picea—*p. Uberrima*, *Pinus picea*—*p. Vulgaris*, *P. picea*.

PIONE, *Pæonia*.

PIONY, *Pæonia*.

PIORTHOPNEA, *Pimelorthopneæ*.

PIOLQUES (F.). A kind of sucking-pump, invented by Louis, for extracting water that had entered internal cavities, in cases of drowning. A useless instrument.

PIP, *Syphilis*.

PIPE-PLANT, *Monotropa uniflora*.

PIPER, *peper*, see *Piper nigrum*.

PIPER ALBUM, *Leucop'iper*, *White Pepper*, is black pepper freed from its cuticle.

PIPER ANGUSTIFOLIUM, *Matico*—*p. Aromaticum*, *P. nigrum*—*p. Betel*, *Betel*—*p. Brazilianum*, *Capsicum annuum*—*p. Calecuticum*, *Capsicum annuum*.

PIPER CAPENSE, a South African species, has all the properties of the peppers, and, in appearance and taste, greatly resembles cubeb. It possesses, too, similar virtues.

PIPER CARYOPHYLLATUM, see *Myrtus pimenta*—*p. Caudatum*, *P. cubeba*—*p. Chiapæ*, see *Myrtus pimenta*.

PIPER CUBE'EA, *Perse'a* seu *Laurus cubeba*, *Litæa cubeba* seu *piper'ita*. Ord. *Piperacæ*. A native of Java and Guinea. The odour of these berries—*Cubeba* (Ph. U. S.), *Com'peper*, *Compeba*, *Cubal sini*, *Piper caudat'um*, *Baccæ Piperis glabri*, *Cubeb* or *Tailed Pepper*, *Cuma'mus*, (F.) *Poivre-à-queue*, *Quabebe*—is aromatic; taste at first cooling, afterwards pungent. The active principle is an essential oil—*Oleum Cubebæ*, oil of cubeb—which is official in the Pharm. U. S. The properties of the cubeb are stimulant and purgative. It is used only in gonorrhœa. Dose, from ℥j to ʒj, in powder, three or four times a day; of the volatile oil, 10 or 12 drops.

'Turkey yellow berries,'—the dried fruit of the *Rhamnus Catharticus*—are often substituted for cubeb.

PIPER ELONGATUM, *Matico*.

PIPER GUINEENSE, *Capsicum annuum*—*p. Hispanicum*, *Capsicum*—*p. Indicum*, *Capsicum annuum*—*p. Jamaicense*, see *Myrtus pimenta*.

PIPER LONGUM, *Macrop'iper*, *Acapat'li*, *Catur'ali*, *Pim'pilim*, *Chav'ica Rozburghii*, *Long Pepper*, (F.) *Poivre long*. As we receive the berries, they consist of the unripe fruit dried in the sun; and are small, round grains, disposed spirally on a cylindrical receptacle. They are stimulant, carminative, and tonic, like the other peppers. Dose, gr. v to ʒj.

PIPER LUSITANICUM, *Capsicum annuum*—*p. Murale*, *Sedum*.

PIPER METHISTICUM, see *Ava*.

PIPER NIGRUM, *Melanop'iper*, *Molagoco'di*, *Lada*, *Piper aromaticum*, *Black Pepper*, (F.) *Poivre noir*. Black pepper, *Piper* (Ph. U. S.), is the unripe fruit dried in the sun. *Virtutes*, same as those of the other peppers. When freed from its husk by mechanical trituration it constitutes *Fulton's decoloured pepper*.

PIPER ODORATUM JAMAICENSE, see *Myrtus pimenta*—*p. Tabasum*, see *Myrtus pimenta*—*p. Turcicum*, *Capsicum annuum*.

PIPERINA, *Piperi'num*, *Piper'ine*, *Piperin*, from *piper*, 'pepper.' A resinoid substance, obtained from the *piper nigrum* by M. Erstädt, in which a part of the virtues of the pepper resides. It has been employed, like the quinia, in intermittents, and successfully.

PIPEROID, *Piperoi'dum*, *Zingib'erin*, from *piper*, 'pepper,' and *oidos*, 'resemblance.' A yellow syrupy mass obtained by exhausting ginger root by ether. It is soluble in ether, alcohol, and oils, and may be given in tincture, sugar, lozenges, or syrup, in cases where ginger is indicated.

PIPEROPASTUS, from *πῆπερ*, 'pepper,' and *πασσιν*, 'to strew.' Strewed with pepper,—soaks, for example, where it is desirable to excite the surface of the feet.

PIPETTE. 'A small pipe.' In Pharmacy, a small tube terminating in a perforated point, which is passed into a liquid to be removed—from the surface of a precipitate, for example. The liquid is drawn into the tube by applying the mouth to the free extremity.

PIPIN'NA, *Men'tula parva*. A small penis.

PIPPERIDGE BUSH, see *Oxycantha Galeni*.

PIPSISSEWA, *Pyrola umbellata*—p. Spotted, *Pyrola maculata*—p. White, *Pyrola maculata*.

PIPTONYCHIA, *Onychoptosis*.

PIPTOSTEGIA OPERCULATA, see *Batata de purga*—p. Pisonis, *Convolvulus mechoacan*.

PIQUETÉ, see *Sablé*.

PIQURE, Puncture, Wound, punctured.

PIRTLING, Slaving.

PIRUS, see *Pyrus malus*.

PISA, (CLIMATE OF.) Pisa has long been celebrated as one of the most favourable climates in Italy for the consumptive. It is genial, but somewhat oppressive and damp. It is softer than that of Nice, but not so warm; less soft, but less oppressive than that of Rome.

PISCIARELLI, MINERAL WATERS OF. A spring at Monte Secco, in Italy, between Lake Agnano and Solfatara. It is thermal (100° Fahr.), sulphureous, and carbonated; and is chiefly used in old ulcers, leucorrhœa, chronic diarrhœa, and in cutaneous affections.

PISCID'IA ERYTHRI'NA, *Jamaica Dog-wood*. *Ord. Leguminosæ*. A small branching tree, common in most of the West India Islands; *Diadelphia Decandria* of Linnæus. It is used in the West Indies to poison fish. It appears to be an acro-narcotic, and has been used in the form of a tincture of the bark of the root in toothache.

PISCINA, *Baignoire*.

PISCIV'OROUS, *Piscivorus*, *Ichthyoph'agus*, *Ichthyoph'agus*, from *piscis*, 'a fish,' and *voro*, 'I eat.' Feeding or subsisting on fish.

PIS'IFORM, *Pisiformis*, from *pisum*, 'a pea,' and *forma*, 'shape.' Pealike, pea-shaped.

PISIFORM'E, *Os*, *Os lenticularè* seu *orbicularè* seu *extra ordinem carpi*, (F.) *Os hors du rang*, is the fourth bone of the first row of the carpus. It is round, and gives attachment to the tendon of the cubitalis anterior, and to the anterior transverse ligament of the carpus. It is articulated, behind, with the *os cuneiforme*.

PISO, Mortar.

PISSA, see *Pinus sylvestris*.

PISSASPHALTUS, *Pissasphaltum*, *Mineral Pitch*, from *πίσσα*, 'pitch,' and *ασφαλτος*, 'bitumen.' (F.) *Pois minérale*. A thicker kind of Petroleum or Rock oil; formerly employed as a vulnerary. It is synonymous with *Glu'tinous bitumen*, *Malta bitumen*, and *Mineral tar*.

PISSELEUM, *Oleum Pic'inum*, from *πίσσα*, 'pitch,' and *ελαίον*, 'oil.' An oily matter obtained from boiling pitch, and used for the same purposes as tar.

PISSELEUM INDICUM, Petroleum.

PISSEMENT DE PUS, *Pyuria*—p. *de Sang*, *Hæmaturia*.

PISENENLIT, *Leontodon taraxacum*.

PISSÉ-SANG, Fumarie.

PISSEROS, from *πίσσα*, 'pitch.' Ancient name of a cerate, prepared of olive oil, oil of roses, and dried pitch. It was, also, called *Cera'tum pica'tum*, and *C. tetraphar'macum*.—*Hippocrates*.

PISSITES, *Vinum pica'tum*. Same etymon.

A wine prepared from tar and the must of the grape. It was regarded to be stomachic.

PISSOSIS, Malacia.

PISSWEED, *Anthemis cotula*.

PISTACHIO NUTS, see *Arachis hypogea*, *Pistacia vera*.

PISTACIA CHIA, *P. lentiscus*.

PISTACIA LENTIS'CUS, *P. Chia*, *Terebin'thus lentiscus*, *Lentiscus vulgaris*; the *Mastic Tree*. *Ord. Terebinthaceæ*. The wood of this tree abounds with the resinous principles of mastic; and a tincture made from it has been used in the cure of dyspeptic affections, gout, and dysentery. It yields, by incision, the resin *Mastic*, *Resi'na Lentis'cina*, *Men'frige*, *Mas'tichè*, *Mastix*, *Pistacia Lentis'ci Resi'na*. This resin has an agreeable smell when heated: is almost insipid; in globular, irregular, yellowish, semi-transparent masses; soluble in ether; partially so in alcohol. It is stimulant, and has been used in old coughs, and gleet, and is chewed to sweeten the breath. It is sometimes, also, employed to stop holes in carious teeth.

Oil of Mastic, *Mastichele'on*, *Oleum Mastich'inum*, is mentioned by Dioscorides.

PISTACIA NARBONENSIS, *P. lentiscus*—p. *Reticulata*, *P. lentiscus*.

PISTACIA TEREBIN'THUS, *P. Narbonensis* seu *reticulata* seu *vera*, *Turpentine Tree*, *Hab-el-Kalimat*, *Terebinthus vulgaris*, *Terminthos*. The tree which affords the *Cyprus Turpentine*, *Resi'na Terebin'thi*, *Terebinth'ina vera* seu *Chia* seu *Cyp'ria*, (F.) *Térébinthine de Chio*. This substance is classed amongst the resins. It is procured by wounding the bark of the trunk. The best Chio turpentine is of about the consistence of honey; very tenacious, clear, and almost transparent; of a white colour inclining to yellow, and of a fragrant smell; moderately warm to the taste, but free from acrimony and bitterness. It resembles in virtues the other turpentine.

PISTACIA VERA, *πίσταια*. Name of the tree which affords the *Nux Pistacia*. *Pistachio nuts* have a sweetish, unctuous taste, resembling that of sweet almonds. Like these, they afford an oil, and may be formed into an emulsion.

PISTILLUM, *Pilum*.

PISTOLOCHIA, *Aristolochia pistilochia*.

PISTORIENSIS GLADIUS, *Bistouri*.

PISUM, *πίον*, *P. sativum*, *Pea*, *Garden pea*, (F.) *Pois*, *P. cultivé*. The seeds of the pea, *Ord. Leguminosæ*, *Sex. Syst. Diadelphia Decandria*, are much employed as an aliment, both in the fresh and dried state. They are farinaceous, and, when dried, not easy of digestion.

PIT, Pock mark—p. of the Stomach, *Fossette du cœur*.

PITAY'A BARK. This bark of *Cincho'na pitay'a*, *Ord. Cinchonaceæ*, is preferred in Colombia to any other. It contains a new vegetable alkali, to which the name *Pitayine* has been given.

PITAYINE, see *Pitaya bark*.

PITCAITHLEY, MINERAL WATERS OF. These saline springs, at Pitcaithley, about four miles from Perth, Scotland, contain chlorides of sodium and calcium.

PITCH, Colophonium—p. Black, see *Pinus sylvestris*—p. Burgundy, see *Pinus abies*—p. Canada, see *Pinus Canadensis*.

PITCH CAP, see *Depilatory*.

PITCH, COMMON, *Pinus sylvestris*—p. Hemlock, see *Pinus Canadensis*—p. Jew's, *Asphaltum*—p. Mineral, *Pissasphaltum*.

PITCH PLASTER, see *Depilatory*, *Emplastrum pici*.

PITCH, STONE, see *Pinus sylvestris*—p. White, see *Pinus abies*.

PITYRIASIS, Pityriasis.

PITYRINUS, Acerosus, Furfuraceus.

PITTA-CIUM. A pitch plaster, from *πῖττα*, 'pitch.'

PITTING, Pock mark.

PITTO-SIS, Malacia.

PITTO-TA. Medicines in which pitch is the chief ingredient.

PITTSBURG, MINERAL WATERS OF. A chalybeate mineral spring, situate about four miles south-west of Pittsburgh, in Pennsylvania. It contains chloride of sodium, chloride of magnesium, oxide of iron, sulphate of lime, and carbonic acid.

PITUITA, Mucus, Phlegm—p. Narium, Nasal mucus.

PITU'ITARY, *Pituitarius*, (F.) *Pituitaire*, from *pituita*, 'phlegm.' Concerned in the secretion of mucus or phlegm. A name given to several parts.

PITUITARY BODY, P. Gland—p. Fossa, Sella turcica.

PITU'ITARY GLAND, *Glandula Pituitaria* seu *pituitaria* seu *basilaris*, *Lacu'na* seu *Appendicula cer'ebr'i*, *Senti'na*, *Hypoph'ysis* seu *Appendix cer'ebr'i*, *Corpus pituitar'e*, *Pituitary body*;—*Appendix sus-sphénoïdale du cerveau*, (Ch.) A small, round body; the anterior portion of which is of a cineritious yellow hue, and the posterior, which is smaller, is whitish and pulpy. Its functions are unknown.

PITUITARY MEMBRANE, *Schneide'rian Membrane*, *Membra'na pituitaria* seu *olfactoria*. The mucous membrane which lines the nasal fossæ, and extends to the different cavities communicating with the nose. In the nasal fossæ, the membrane appears formed of two layers, intimately united;—the one in contact with the bone is fibrous;—the other is free at one surface. The latter is a mucous membrane, on which papillæ are not easily distinguishable, but which contains them, as well as a number of very small mucous follicles, that open into the cavity of the nares. On penetrating the maxillary, frontal, sphenoidal, and ethmoidal sinuses, the membrane becomes very thin, transparent, less vascular, and seems reduced to its mucous layer. The pituitary membrane receives the impression of odours, by means of the expansion of the olfactory nerves upon it. It is the seat of smell.

PITUIITE, Mucus.

PITU'ITOUS, *Pituitosus*; from *pituita*, 'mucus or phlegm.' Consisting of, or resembling, or full of mucus or phlegm.

PITYRIASIS, from *πυρρον*, 'bran'; *Lepido'sis* Pityriasis, *Pityris'ma*, *Pithyri'asis*, *Herpes furfuraceus* seu *farinosus*, *Porri'go* (of some), *Tinea furfuracea* seu *porriginosa*, *Furfuratio*, *Furfurica'ca*, *Dandruff*, *Dandruff*, (Sc.) *Lusa*, (Prov.) *Roum*, *Rummet*, (F.) *Teigne*, *Dartre*, *D. furfuracea volante*;—a very superficial affection; characterized by irregular patches of thin scales, which repeatedly exfoliate and recur; but which never form crusts, or are accompanied with exoriations. It is not contagious. It occurs under three or four varieties of form.

PITYRIASIS, Porrigo—p. Nigra, Melasma—p. Versicolor, Chloasma.

PITYRISMA, Pityriasis.

PITYROIDES, Furfuraceus.

PITYRON, Furfur.

PIVOINE, Pæonia.

PIV'OTING. In dentistry, an operation, which consists in fastening in a new crown upon the root of a tooth by means of a piece of gold wire, called 'a pivot.'

PIX, Sperm—p. Abietina, see *Pinus abies*—p. Abietis, see *Pinus abies*—p. Alba, see *Pinus abies*—p. Arida, see *Pinus abies*, and *Pinus syl-*

vestris—p. Atra, see *Pinus sylvestris*—p. Brutia, Brutia—p. Burgundica, see *Pinus abies*—p. Canadensis, see *Pinus Canadensis*—p. Cedria, see *Pinus sylvestris*—p. Græca, Colophonias—p. Li-quida, see *Pinus sylvestris*—p. Navalis, see *Pinus sylvestris*—p. Nigra, see *Pinus sylvestris*—p. Sicca, see *Pinus sylvestris*.

PLACÉBO, 'I will please,' from *placen*. A medicine, prescribed rather to satisfy the patient than with any expectation of its effecting a cure.

PLACEN'TA, *πλάκους*, *πλάκουντρος*. A Latin word, signifying a Cake. *P. uteri* seu *uteri'na* seu *uteri car'nea*, *Hepar* seu *Jecur uteri'num*, *Caro* seu *Affusio orbicularis*, *Maza*. A soft, spongy, vascular body, adherent to the uterus, and connected with the fetus by the umbilical cord. It is not in existence during the first period of the embryo state, but its formation commences perhaps with the arrival of the embryo in the uterus. It is generally considered to have two portions, one fetal, and the other maternal. The fetal consists of highly vascular villi and tufts, which contain the insinuating loops of the umbilical arteries and umbilical vein of the fetus. The maternal portion is considered by Dr. Reid to consist, essentially, of a large sac formed by the inner coat of the vascular system of the mother, into which the maternal blood is poured by the curling arteries of the uterus, and from which it is returned by the utero-placental veins. The tufts and villi of the fetal placenta are ensheathed by this inner coat of the maternal vascular system; and, according to Prof. Goodrip, two distinct sets of nucleated cells—one maternal and the other fetal—separate the maternal and fetal portions where they come in contact, one set of which are probably, he thinks, concerned in the separation of nutritive matter from the maternal blood, and the other destined to convey it into the vessels of the fetus. The placenta is formed of several lobes or cotyledons, which can be readily distinguished from each other on the uterine surface; but towards the fetal surface are confounded into one mass. It is composed of the umbilical vessels, areolar tissue, and whitish, resisting filaments, which are vascular ramifications obliterated. Lymphatic vessels have also been presumed to exist in it; and nerves have been traced, proceeding from the great sympathetic of the fetus.

The main function of the placenta appears to be like that of the lungs in the adult. It admits of the blood of the fetus being shown to that of the mother, and undergoing requisite changes. It may, also, be an organ for nutritive absorption, as observed above.

Animals, which have no placenta—as the Marsupialia—are called *implacental* and *nonplacental*.

PLACEN'TA CRUORIS, see Blood—p. Febrilis, Ague cake—p. Oblata, P. prævia—p. Obvia, P. prævia.

PLACEN'TA PRÆ'VIA, (from *præ*, 'before,' and *via*, 'way,') P. obvia seu *obla'ta*. Presentation of the placenta; a condition which gives rise to unavoidable hemorrhage from the uterus. The safety of the female depends upon speedy delivery.

PLACEN'TA SANGUINIS, see Blood—p. Uteri Succenturiata, Decidua.

PLACEN'TAL, *Placental'is*, (F.) *Placentaire*. Relating, or appertaining to, the placenta. Possessed of a placenta,—as 'a placental mammal.'

PLACEN'TIS, *Placental'is*, *Placoi'tis*, *Inflamma'tio placen'tæ*. A hybrid word—from *placenta*, and *itis*, a suffix denoting inflammation. Inflammation of the placenta.

PLACEN'TULA, 'a little cake,' diminutive of *placenta*, 'a cake.' A rudimental placenta. The placenta of an abortion.

PLACOITIS, Placentitis.

PLACOUNTITIS, Placentitis.

PLADAROMA, Pladarosis.

PLADAROSIS, *Pladaro'ma*, *Pladar'otes*, from *πλαδαγος*, 'soft.' A soft tumour on the eyelid, unaccompanied with redness or pain. Trachoma.

PLADAROTES, Pladarosis.

PLAFOND DES VENTRICULES DU CERVEAU, Corpus callosum.

PLAGA, Ictus, Wound.

PLAGIOBOL'IA, *Planobol'ia*, from *πλαγιος*, 'to one side,' and *βαλλειν*, 'to throw.' Imperfect emission of sperm into the vagina. Emission to one side; sometimes practised to render impregnation less probable.

PLAGUE, from *πληγη*, *plaga*, 'a stroke.' *Pestis*, *P. orientalis* seu *bubona'ria orientalis* seu *inguina'ria* seu *contagio'sa* seu *acutis'sima* seu *septica* seu *glandulo'sa* seu *ade'no-septica* seu *inguinaria*, *Pestil'itis*, *Ephem'era pestilentia'lis*, *E. mortif'era*, *Lues inguina'ria*, *Clades glandu'lar'ia*, *Phthora*, *Pestilen'tia*, *Lues*, *Loimos*, *Febris ade'no-nervo'sa*, *Læmos*, *Læmè*, *Læ'mia*, *Læmop'gra*, *Chao'da*, *Anthra'cia Pestis*, *Anthraco'typhus*, *Febris pest'itens*, *Syn'ochus pestilentia'lis*, *Typhus pestis* seu *pestilentia'lis* seu *gravis'simus* seu *anthra'ci'cus* seu *bubon'icus*, *Ezanth'e'ma pestis*, the *Pest*, *Levant plague*, *Septic* or *glandular pestilence*, *The sickness*, (F.) *Peste*, *Typhus d'Orient*, *Fidère du Levant*, *F. adèno-nervæuse*. An eminently malignant disease; endemic in the Levant; frequently epidemic, and destroying at least two-thirds of those it attacks. It is a fever of the most aggravated kind, with affection of the lymphatic glands of the groins or axillæ, and carbuncles. Its miasmata—it has been conceived—adhere to different organic textures, to woollen goods, clothing and furniture; and may thus be transported to a considerable distance; but this is not certain. The mean duration of the disease is six or seven days; some die in twenty-four hours, others not till ten or twelve days. Pathological anatomy has afforded little light with respect to it. Various means have been used for arresting it, but none have seemed to be pre-eminently distinguished. The great point is, to watch the indications as they develop themselves; and to treat the case, in general, like one of typhus gravior. It is universally agreed that the suppuration of the buboes should be aided as far as practicable. For preventing the importation and spread of the plague, the *Quar'antine Laws* have been instituted; and when the disease has actually appeared, a *cordon sanitaire* has been drawn round the infected district, so as to prevent all communication.

PLAGUE OF ANTONINUS, *Pestis Antoninia'na*, (F.) *Peste Antonine*. A severe febrile disease, which prevailed in the Roman empire, during the reign of Antoninus. It began in Asia, and passed westward. It resembled the Plague of Athens.

PLAGUE OF ATHENS, (F.) *Peste d'Athènes*. A febrile disease, which raged at Athens during the Peloponnesian war, in the 5th century before the Christian era; and a description of which is given in Thucydides. It was characterized by a cutaneous eruption, accompanied by small ulcerations, vomiting, affection of the respiratory organs, and diarrhœa. It began in the far East, and, before reaching Athens, ravaged Egypt, and the greater part of the Persian Empire.

PLAGUE, BLACK, Death, black.

PLAGUE, COLD. A severe form of congestive fever, seen in the Southern States. Bilious pneumonia, in which there is no reaction, has been, also, so called.

PLAGUE MARK, Plague token.

PLAGUE, PALL. A malignant fever, greatly

resembling Plague, which prevailed in Rajpootana after July, 1836.

PLAGUE SPOT, Plague token.

PLAGUE TOKEN, *Plague mark*, *Plague spot*. A mark by which one struck with the plague was known. Perhaps the early stage of carbuncle in unfavourable cases. It is described as a small tubercle — *Bossa* — somewhat resembling a wart, callous and more or less deficient in sensibility, varying in size from that of a millet-seed to that of a bean.

PLAGULA, Compress, Pledget, Splint.

PLAIE, Wound — *p. d'Arme à feu*, Wound, gunshot — *p. d'Arquebuseade*, Wound, gunshot — *p. Contuse*, Wound, contused — *p. Envenimée*, Wound, poisoned — *p. par Arrachement*, Wound, lacerated — *p. à Lambeaux*, see Flap.

PLAINITIVE, *Luctuëux*.

PLANANTHUS FASTIGATUS, Lycopodium selago — *p. Selago*, Lycopodium selago.

PLANARIA LATIUSCULA, Distoma hepaticum.

PLANCHER (F.), 'floor.' In anatomy, the inferior wall or boundary of a cavity.

PLANCHER DU CERVEAU, Tentorium.

PLANCUS, Leiopus.

PLANE, Plexus.

PLANE, MESIAL, see Mesial.

PLANET-STRUCK, see Sideratio.

PLANETTES, *Planeticus*, *Plan'icus*, *Errabundus*, *Errat'ic*, from *πλανν*, 'a wandering about.' An epithet given to diseases, whose returns are irregular — especially to intermittent fever; *Planetes Febris*, *A. Errat'ica*, *F. Erro'nea*, *F. Vaga*.

PLANETICA ARTHRITIS, Gout, wander ing.

PLANICUS, Planetes.

PLANIPEDES, Stéganopodes.

PLANITIES PEDIS, Sole.

PLANOBOLEA, Plagiobolia.

PLANTA NOCTIS, Hydroa, Sudamina — *p. Pedis*, Sole — *p. Prima pedis*, Tarsus.

PLANTA'GO, *P. Major*, *Centiner'via*, *Arnoglossum*, *Septiner'via*, *Polynéu'ron*, *Heptaplex'ron*, *Plant'go latifo'lia* seu *major* seu *vulgar'is*, *Great plantain*, *Waybread*, (Sc.) *Wabran leaves*, (F.) *Plantain commun*. Ord. Plantaginaceæ. *Sex. Syst.* Tetrandria Monogynia. The leaves have been said to be refrigerant, attenuant, sub-styptic, and diuretic.

PLANTAGO AQUATICA, *Alisma plantago*.

PLANTAGO CORDA'TA, *Heart leaved plantain*, *P. LANCEOLA'TA*, *Ribgrass*, and *P. VIRGIN'ICA*, *White plantain*, or *Ribwort*, indigenous; have properties like *Plantago major*.

PLANTAGO CORON'OPUS, *Coronop'dium*, *Cornu cervi'num*, *Stella terra*, *Plantago*, *Buckshorn Plantain*. Properties like the last.

PLANTAGO NITIDA, *P. psyllium*.

PLANTAGO PSYL'LIUM, *P. nit'ida*, *Psyll'ium*, *P. erectum*, *Pulc'aris herba*, *Crystal'lion*, *Cynmoi'a*, *Branching Plantain*, *Fleawort*, (F.) *Herbe aux puces commune*. The seeds have a nauseous, mucilaginous taste, and no remarkable smell. The decoction of the seeds is recommended in hoarseness and roughness of the fauces.

PLANTAGO VULGARIS, *Plantago*.

PLANTAIN DES ALPES, *Arnica montana* — *p. Branching*, *Plantago psyllium* — *p. Buckshorn*, *Plantago coronopus* — *p. Commun*, *Plantago* — *p. d'Eau*, *Alisma plantago* — *p. Great*, *Plantago major* — *p. Heart leaved*, *Plantago cordata*.

PLANTAIN MEAL, see *Musa paradisiaca*.

PLANTAIN, RATTLESNAKE, *Goodyera pubescens* — *p. Robin's*, *Erigeron bellidifolium*.

PLANTAIN TREE, *Musa paradisiaca*.

PLANTAIN, WATER, *Alisma plantago*, *Unisema deltifolia* — *p. White*, *Plantago Virginica*.

PLANTAIRE, *Plantar muscle*—*p. Grêle*, *Plantar muscle*.

PLANTAR, *Plantaris*, (F.) *Plantaire*, from *planta*, 'the sole of the foot.' That which relates or belongs to the sole of the foot. This name has been given to several parts.

PLANTAR APONEUROSIS, *P. fasciæ*. A thick, dense, resisting aponeurosis; of a triangular shape, and superficially situate, which occupies the middle and sides of the sole of the foot, and furnishes insertions to several of the muscles of the region, either directly or through the medium of septa, which it sends between them.

PLANTAR ARCH, see *Plantar arteries*.

PLANTAR ARTERIES. These are two in number, and arise from the extremity of the posterior tibial artery. They are distinguished into:—The *internal plantar artery*, situate above the abductor pollicis pedis. It extends from the termination of the posterior tibial to the trunk of one of the collaterals of that toe.—The *external plantar artery* is larger than the preceding, and ought to be considered as the termination of the posterior, tibial. Situate above the flexor communis digitorum pedis, it extends from beneath the calcaneum to the posterior extremity of the last metatarsal space. When it has reached this part, it turns from without inwards; advances towards the posterior extremity of the first metatarsal bone, and anastomoses with the perforating branch of the *dorsalis tarsi*. In the first part of its course, it furnishes calcaneal, articular, muscular, and adipous branches. The second portion of the artery is called the *Plantar Arch* or *Cross*. It gives branches in all directions. The inferior and posterior have received no name. The superior—three in number—are the *posterior perforating arteries*. The anterior branches are larger, and there are commonly four. They furnish the *anterior perforating*, and the *collaterals* of the last four toes.

PLANTAR CROSS, see *Plantar arteries*—*p. Fasciæ*, *P. aponeurosis*.

PLANTAR LIGAMENTS. A name given to the inferior ligaments, which unite the bones of the tarsus and metatarsus, to distinguish them from the superior, called *dorsal*.

PLANTAR MUSCLE, *Plantaris*, *P. gracilis*, *Tibialis gracilis*, *Extensor tarsi minor*, (F.) *Plantaire*, *P. Grêle*, *Jambier grêle*, *Petit fémorocalcanien*, (Cb.) A muscle, situate at the posterior part of the leg. It is long, thin, and narrow. Above, it is attached to the posterior part of the external condyle of the os femoris: below, it terminates by a long tendon, which is attached to the posterior and inner part of the calcaneum. This muscle extends the leg on the foot, and conversely. It can, also, assist in the flexion of the leg on the thigh.

PLANTAR NERVES. These are two in number, and are furnished by the termination of the posterior tibial nerve. They are distinguished into—1. The *internal plantar nerve*, which extends from the inferior part of the tibial nerve to both sides of the first three toes, and the inner side of the fourth. It furnishes, also, branches to the abductor of the great toe; to the short flexor of the toes; to the accessory; to the short flexor of the great toe, the lumbricales, &c. 2. The *external plantar nerve* proceeds from the termination of the posterior tibial to the posterior extremity of the 5th metatarsal bone, and gives branches to the flexor brevis digitorum pedis, the accessory and the abductor of the little toe. This trunk subsequently divides into two branches;—the one—*profunda*—is distributed to the short flexor of the great toe: the interosseus and abductor pollicis pedis: the other—*superficialis*—sends filaments to the short flexor of the little toe, the

two sides of that toe, the outer side of the 4th, and the last lumbricalis.

PLANTAR REGIONS OR SURFACES are different regions or surfaces of the sole of the foot. Three are generally reckoned,—the *internal*, *middle*, and *external*.

PLANTAR VEINS follow an arrangement similar to that of the arteries.

PLANTARIA, *Dengue*.

PLANTARIS, *Plantar*, *P. muscle*—*p. Superficies pedis*, *Sole*—*p. Verus*, see *Flexor longus digitorum pedis profundus perforans (accessorius)*.

PLANTARIUM, *Pubes*.

PLANTE DU PIED, *Sole*.

PLANTI-SOUS-PHALANGIENS, *Lumbricales pedis*—*p. Tendino-phalangiens*, *Lumbricales pedis*.

PLANUM APONEUROTICUM DIAPHRAGMATIS. The tendinous expansion of the diaphragm.

PLANUM ORBITALE, see *Maxillary bone (superior)*.

PLANUM OS, from *planus*, 'soft, smooth.' The papyraceous or orbital plate of the ethmoid bone—*Lamina papyracea*, *Lamella plana*—was formerly so called.

PLANUM PEDIS, *Sole*.

PLANUM SUPERIORIS LINGUÆ. The upper surface of the tongue.

PLANURIA, from *πλανος*, 'wandering, false,' and *ουρον*, 'urine.' Discharge of urine through unwonted ways.

PLAQUEMINIER D'EUROPE, *Diospyros lotus*.

PLAQUES GAUFRÉES, *Peyeri glandulæ*—*p. Laitenses*, see *Tachæ laticæ*—*p. de Peyer*, *Peyeri glandulæ*.

PLASMA, *Liquor sanguinis*.

PLASMEXHIDROSIS, from *plasma*, the *liquor sanguinis*; *εξ*, 'out of,' and *ἰδρωσις*, 'perspiration.' Exudation of the plasma of the blood from the vessels.

PLASTER, *Emplastrum*—*p. Adhesive*, *Emplastrum resinæ*, *Sparadrapum adhesivum*—*p. Adhesive*, *Baynton's*, see *Emplastrum resinæ*—*p. Adhesive*, of pitch and resins, *Emplastrum de pice et resinis glutinans*—*p. Ammoniacum*, *Emplastrum ammoniaci*—*p. Ammoniacum* with mercury, *Emplastrum ammoniaci cum hydrargyro*—*p. Aromatic*, *Emplastrum aromaticum*—*p. Asafoetida*, *Emplastrum asafoetidæ*—*p. of Belladonna*, *Emplastrum belladonnæ*—*p. Blister*, *Blister*—*p. Blistering*, *Emplastrum lyttæ*—*p. of Blistering fly*, *Emplastrum lyttæ*—*p. Calefacient*, *Emplastrum calefaciens*—*p. of Cantharides*, compound, *Emplastrum lyttæ compositum*—*p. Cephalic*, *Emplastrum picis comp.*—*p. Corn*, *Sparadrapum viride*—*p. Corn*, *Kennedy's*, *Sparadrapum viride*—*p. Court*, *English*, see *Ichthyocolla*, *Sparadrapum adhesivum*—*p. Cummin*, *Emplastrum cumini*—*p. Defensive*, *Sparadrapum Galteri*—*p. Diachylon*, *Emplastrum plumbi*—*p. Fly*, *Emplastrum lyttæ*—*p. Frankincense*, *Emplastrum thuris*—*p. Galbanum*, *Emplastrum galbani*—*p. Galbanum*, compound, *Emplastrum galbani compositum*—*p. Gum*, *Emplastrum gummosum*—*p. of Gum-resins*, *Emplastrum cum gummi-resinis*—*p. Hemlock*, *Emplastrum cicutæ*—*p. Iron*, *Emplastrum ferri*—*p. Isinglass*, see *Sparadrapum adhesivum*—*p. Issue*, *Sparadrapum pro fonticulis*—*p. Lead*, *Emplastrum plumbi*—*p. Machine*, *Sparadrapier*—*p. Mercurial*, *Emplastrum hydrargyri*—*p. of Mercury*, compound, *Emplastrum hydrargyri compositum*—*p. Mustard*, *Cataplasma sinapis*—*p. Nuremberg*, *Emplastrum Norimbergense*—*p. Opium*, *Emplastrum opii*—*p. Pitch*, see *Depilatory*—*p. Pitch*, comp. *Emplastrum picis compositum*—*p. Resin*, *Emplastrum resinæ*—*p. of Red oxide of iron*, *Em-*

plastrum oxidi ferri rubri—p. Soap, Emplastrum zaponis—p. of Spanish flies, compound, Emplastrum lyttæ compositum—p. Sticking, Emplastrum resinæ—p. Strengthening, Emplastrum ferri, Emplastrum oxidi ferri rubri—p. Warm, Emplastrum calefaciens—p. Wax, Emplastrum cereæ.

PLASTIC, *Plas'ticus*, *Formativus*, *Form'ative*, (F.) *Plastique*, from *πλασσω*, 'I form.' That which forms, or serves to form.

PLASTIC ELEMENT. One which bears within it the germs of a higher form.—Gerber. *Plastic* or *histogenetic* or *organoleptic* elements of nutrition are those elements of food which go to the formation of organized tissue.—See Aliment.

PLASTIC FORCE. The generative or formative power in organized bodies. It is also called *Nivus formativus*, *Vis seu Virtus Plastica* seu *formativa seu essentialis* seu *reproductiva*, *Facultas formatrix* seu *nutrix* seu *auctrix* seu *vegetativa*; *Blas alterativum*, *Motus assimilationis*, *Anima vegetativa*, *Plasticismus*, *Virtus formatrix*, *Plasticitas*, *Plasticity*, Force of Nutrition, F. of assimilation, F. of Formation, F. of vegetation, *Puissance du moule intérieur* (Buffon), *Bildungstrieb*, *Germ-force* of the Germans.

PLASTIC LYMPH, *Liquor Sanguinis*—p. Surgery, *Morioplastice*.

PLASTICISMUS, Plastic Force.

PLASTICITAS, Plastic Force.

PLASTICITY, Plastic Force.

PLASTILYTIC, Antiplastic.

PLASTILYTICUM (**HÆMATOLYTICUM**), Spanæmic.

PLATA, Scapula.

PLATANO, *Musa Sapientum*.

PLATANTHE'RA ORBICULA'TA, *Large round-leaved Orchis*; indigenous. Order, *Orchidacæ*; flowering in July. From *πλαρως*, 'wide,' and *ανθη*, for 'anther.' The leaves are large, soft and fleshy, and are used, in certain places, for dressing blisters.

PLATANUS, *Musa paradisiaca*.

PLATEA, *Bothriocephalus latus*.

PLATE-BONE, Scapula.

PLATEIASMA, *Platiasmus*.

PLATES, *MANSFORD'S*, see Galvanism.

PLATIAS'MUS, *Platias'ma*, from *πλατυς*, 'broad.' Excessive development of the lips, causing the articulation to be imperfect.

PLATINA, Platinum.

PLATINUM, *Plat'ina*, from (S.) *plata*, 'silver.' (F.) *Platine*, Or *blanc*, *Petit argent*. The preparations of platinum resemble in their therapeutical properties those of gold. The *BICHLORIDE*, *Plat'ini Bichloridum*, made by dissolving platinum in *aqua regia*, and the *DOUBLE CHLORIDE* of PLATINUM and SODIUM, *So'dii chloroplat'inas*, *Chloroplat'inatæ* of *So'dium*, prepared by dissolving bichloride of platinum and pure chloride of sodium, in proper proportions, in water, evaporating and crystallizing.—are the preparations used. They are not much prescribed.

PLATULA, see *Pediculus*.

PLATYBREGMATE, see *Cranium*.

PLATYCORIA, *Mydriasis*.

PLATYCORIASIS, *Mydriasis*.

PLATYNO'SIS, from *πλαρυς*, 'broad.' *Amplificatio*. Morbid extension of parts.

PLATYOPHTHALMON, *Antimonium*.

PLATYPOLIA, see *Kyllosis*.

PLATYPUS, see *Kyllosis*.

PLATYS'MA, from *πλαρυς*, 'broad.' Anything broadly extended. A spread plaster.

PLATYS'MA MYOÏDES, *P. Myo'des*, *Musculus cutaneus seu subcutaneus* (Winslow), *Quadratus genæ* seu *latis'simus colli*, *Expan'sio musculo'sa*, *Tetragnus*, *Stomomani'con*, (F.) *Thoraco-maxillofacial*, *Thoraco-facial* (Ch.), *Peaucier*. A mus-

cle situate superficially on the lateral parts of the neck. It is flat, broad, and quadrilateral. Its fibres, which are all parallel to each other, pass obliquely upwards, and from without inwards. They arise from the areolar tissue, which covers the anterior and superior part of the chest; pass before the clavicle, and proceed to attach themselves at the interior part of the symphysis menti, at the external oblique line of the jaw, and at the commissure of the lips. The fibres of this last insertion form a distinct fasciculus, called by some anatomists *Musculus Risorius Santorini*. The platysma myoides depresses the commissure of the lips, and carries it outwards. It contributes, also, to the depression of the lower jaw. When it contracts, it wrinkles the skin of the neck transversely.

PLATYSTER'NOS, from *πλατυς*, 'broad,' and *στερνον*, 'the sternum.' One who has a broad, well developed chest.—D. Pechlin.

PLATYS'TOMUS; from *πλατυς*, 'broad,' and *στομα*, 'mouth.' One who has a broad mouth.

PLAUTUS, *Leipus*.

PLECHAS, *Perinæum*.

PLECTANÆ, *Cornua uteri*.

PLECT'ANE, *Plica*, *Plegma*, *Plexus*. From *πλεκω*, 'I twine.' A fold; a plexus. Also, a cornu of the uterus.

PLECT'RUM. The styloid process of the temporal bone. Also, the uvula, and the tongue.—Castelli.

PLEGGET, (D.) *Plagghe*, *Plumaceolus*, *Lintæ'men*, *Plag'ula*, (F.) *Plumaceau*, *Plumaseau*. A small compress or gâteau of lint—(the filaments arranged parallel to each other)—flattened between the hands after the extremities have been cut off or folded down. It is applied over wounds, ulcers, &c., to preserve them from the contact of the air, to retain dressings *in situ*, and to absorb the discharges.

PLEGE, Blow, Ictus.

PLEGMA, *Plectane*, *Plexus*.

PLEGOMETER, *Pleximeter*.

PLEIN, (F.) 'Full.' The middle part of a roller.

PLEIOMASTHUS, *Polymasthus*.

PLEIOMAZIA, see *Polymasthus*.

PLENA MULIER, *Pregnant*.

PLENITUDE, *Fulness*.

PLENTUDO, *Plethora*.

PLENCK'S DEPILOYATORY, see *Depilatory*, *Colley's*.

PLEOMAZIA, see *Polymasthus*.

PLEONEC'TICUS, from *πλεον*, 'more,' and *εχειν*, 'to have.' Covetous. Desirous to acquire: hence *mania pleonectica*.

PLEONEC'TICA ATHYM'IA seu **MELANCHOL'IA**. A form of insanity characterized by a desire for gain or by covetousness.

PLERO'SIS, from *πληρης*, 'full;' also *Repletio*, *Refectio*, *Repletio*. Restoration of flesh after sickness. *Plethora*.

PLEROTICUS, *Incarcans*.

PLES'MONÈ, *πλησμονη*, *Plenitudo*, *Sati'etas*, *Sati'ety*. Intemperance. A condition the opposite to that of hunger.

PLESSER, *Plexor*.

PLESSIMETER, *Pleximeter*.

PLESSIMÈTRE, *Pleximeter*.

PLETHOMER'IA, from *πληθος*, 'many,' and *μερος*, 'a part.' Superabundance of parts, as in the case of six fingers to one hand.

PLETHORA, from *πληθα*, 'I fill.' *Polyæ'mia*, *Polyhyperhæ'mia*, *Panlyhyperæ'mia*, *Polychym'ia sanguinea*, *Augæioprolo'sis*, *Angioprolo'sis*, *Erythræ'mia*, *Erythro'sis*, *Hæmatopletho'ra*, *Hæmopletho'ra*, *Plenitudo*, *Multitudo*, *Redundant'ia*, *Repletio*, *En'chyma*, *Comple'tio*, *Pleth'ory*, *Repletio*, *San'guineus*, *Fulness*, (F.) *Abundance*, *Rédondance*. The word *plethora*, which means

repletion, *Pleuro'sis*, expresses a superabundance of blood in the system, or in some part of it:—hence the division of plethora into *general* and *local*; the latter being called, according to its seat, *cerebral*, *pulmonary*, *uterine*, &c. The principal symptoms of plethora exist in the circulatory system; such as redness of the surface, swelling of the veins, increase in the fulness of the pulse, in the strength of the heart's pulsations, &c., with spontaneous hemorrhages occasionally. With these are usually united general heaviness, torpor, lassitude, vertigo, tinnitus aurium, flushes of heat, &c. The blood of plethoric persons differs from healthy blood in the smaller ratio of water which it contains, and especially in the augmentation of the red corpuscles.

The tendency to plethora, or its actual existence, must be obviated by purgatives, proper diet, exercise, &c.; and, if the danger from it be pressing, by blood-letting. This is, however, a doubtful remedy for general plethora, as it increases hæmatisis.

PLETHORÆ ARTERIO'SA is the form in which the blood is rich in fibrin and red corpuscles—the *arterial constitution*.

PLETHORA, LOCAL, Hyperæmia — p. Partialis, Hyperæmia.

PLETHORIC, *Plêthor'icus*, *Polyæ'mic*, *Polyæ'micus*, *Sanguine*, *Sanguin'eous*, (F.) *Plêthorique*. Same etymon. Full of blood. Relating to or affected with plethora.

PLETHORICUS, Plethoric.

PLETHORY, Plethora.

PLETHOS, Satiety.

PLEUMON, Pulmo.

PLEUMONIA, Pneumonia.

PLEURA, *Pleurum*, *Pleuro'ma*, 'the side, a rib.' *Membra'na pleurit'ica succin'gens* seu *costæ succin'gens*, *M.* seu *Tu'nica costalis* seu *subcosta'lis*, *Hypopleu'rios*, *Hypezo'cus*, (F.) *Plèvre* ou *Pleure*. The pleuræ are two thin, diaphanous, perspirable membranes, which line each side of the chest, and are reflected thence upon each lung. Like all serous membranes—to which class they belong—each represents a sac without aperture. From the junction of the pleura of each side the mediastina are formed. That portion of the pleura which lines the parietes of the chest is called *Pleura Costa'lis*; the portion that covers the lungs, *Pleura Pulmona'lis*. The arteries of the pleuræ are from the intercostals, internal mammaries, phrenic, inferior thyroid, &c. The veins correspond with the arteries. They are, also, supplied with lymphatics, but nerves are not traceable in them.

Pleura is also used for *Cavity of the Pleura*, *Saccus Pleuræ*.

PLEURÆ, The sides of the chest.

PLEURAL, *Pleura'lis*, *Pleur'icus*, *Pleur'ites*. That which concerns the pleura;—as a *pleural fistula*;—*pleural cavity*, *pleural hemorrhage*, &c.

PLEURALGIA, Pleurodyny.

PLEURAPOPHYSES, *Costæ veræ*, see *Costa*.

PLEURAPOSTEMA, Empyema.

PLEURARTHROCA'CE, from *πλευρα*, 'a rib,' *αρθρον*, 'a joint,' and *κακος*, 'bad.' Caries of the ribs.

PLEURE, Pleura.

PLEURESIA, Pleuritis.

PLEURÉSIE, Pleurisy — p. *Fausse*, Pleurodynia.

PLEURESIS, Pleuritis.

PLEURÉTIQUE, Pleuritic.

PLEURICUS, Pleural.

PLEURIS, Pleuritis.

PLEURISY, Pleuritis

PLEURISY, BILIOUS, *Pleurit'is bili'o'sa*. Pleurisy accompanied with bilious symptoms;—the effect of duodenitis or duodeno-hepatitis.

PLEURISY, FALSE, Pleurodynia.

PLEURISY, HEAD. A ridiculous term, occasionally used by the vulgar, in the Southern States especially, for bilious pneumonia before the pneumonic phenomena are developed, and whilst the head is prominently affected.

PLEURISY, HEMORRHAGIC, Hæmatothorax — p. Rheumatic, Pleurodynia — p. Root, Asclepias tuberosa.

PLEURISY, TYPHOID, *Pleurit'is typho'id'es* seu *typho'd'es* seu *typho'sa*. Pleurisy accompanied by adynamic symptoms.

PLEURITES, Pleural.

PLEURITIC, *Pleurit'icus*, *Pleurit'ical*, (F.) *Pleurétique*; same etymon as *pleuritis*. Relating or appertaining to pleuritis — as 'pleuritic pain.'

PLEURITIS, from *πλευρα*, 'the pleura'; *Morbus latera'lis* seu *pleurit'icus* seu *Costa'lis*, *Empres'ma* seu *Cau'ma Pleurit'is*, *Pleuris*, *Pleure'sis*, *Pleure'sia*, *Pleuritis costa'lis* seu *vera*, *Pleurit'ica Febris*, *Passio pleurit'ica*, *Inflamma'tio Pleuræ*, *Pleurisy*, *Inflamma'tion of the Pleura*, (F.) *Pleurée*, *Fèvre pleurétique*, *Inflammation de la Plèvre*, may be acute or chronic. Perhaps the most violent of all internal inflammations is the active form; known by acute, lancinating pain in one side of the chest, increased by inspiration, by coughing, and often by pressure; dyspnoea, the inspirations short and frequent; cough dry, or with little expectoration; and difficulty of lying on the affected side. Symptoms, attending the phlegmasia in general, strongly marked.

Physical Signs. There is seldom any perceptible alteration in the exterior of the chest. When effusion has taken place to a great extent, the signs will be similar to those of chronic pleurisy. The motion of the affected side is diminished, and the vibrations of the voice cannot be detected by the hand. There is more or less dullness of sound on percussion, the dullness diminishing or disappearing by change of position. On auscultation, the inspiration is feeble, distant, or inaudible; but change of position modifies it. Egophony is usually present, when the effusion is in moderate quantity, and is best heard over a portion of the thoracic parietes, represented by a band of three inches, running from below the inferior margins of the scapula in the direction of the ribs to the sternum. Simple pleurisy rarely attacks both sides at once: when such is the case, the presence of tubercles may be suspected.

Formerly, a *true pleurisy* meant one seated in the pleura: a *mediastinal pleurisy*, one situated in the anterior mediastinum; and *false pleurisy* or *rheumatismal pleurisy*, that which occupies the intercostal muscles. The last has received the name *Pleurodyn'ia*; that of *pleurisy* being exclusively appropriated to inflammation of the pleura. Pleurisy has also been distinguished into *dry* and *humid*, according as it is or is not accompanied by expectoration. *Humid* or *moist pleurisy* is now properly regarded as a complication of pleurisy with pulmonary catarrh. It is the *catarrhal pleurisy* of some. The majority of the species of pleurisy, admitted by Sauvages, consist of complications.

The causes of pleurisy are like those of other internal inflammations. It may terminate by resolution, effusion, or suppuration.

CHRONIC PLEURISY, which has been particularly described by Bayle, Broussais, and Laënnec, most commonly terminates by a serous or purulent effusion, which sometimes simulates hydrothorax; at others, phthisis pulmonalis. It

may form slowly, without evident symptoms; or it may succeed acute pleurisy. In the former case, vague pains are felt in the chest, with small, dry cough, oppression at intervals, shivering, irregular febrile symptoms, and hardness of pulse.

Physical Signs. The affected side is smoother, more rounded and motionless; the intercostal spaces are dilated and filled up, or may even protrude slightly. In very chronic cases, however, the affected side may be smaller. The triangular space above the clavicle and the depression immediately above the sternum are often drawn downwards on the diseased side. No vibration is experienced by the hand when the patient speaks. The parietes of the thorax are sometimes œdematous, and fluctuation may be occasionally felt between the ribs. On percussion the sound is dull, or if clear, only so in the upper portions of the chest. On auscultation, there is an absence of the usual sounds over the affected parts, excepting occasionally transmitted sounds. There are no physical signs that can enable us to know whether pus or serum is effused into the chest. The introduction of a grooved needle through the intercostal space, as recommended by Dr. Thomas Davies, would of course settle the question.

On dissection of those who have died of pleurisy, the pleura is found thickened, red, inflamed, and covered with membranous exudations or false membranes. Sometimes it seems cartilaginous and ossified. Its cavity frequently contains serous or sero-purulent effusions.

The acute form of the disease requires the most active treatment. General blood-letting, to such an extent as to make a very decided impression on the system, and repeated if necessary. This is the most important agent. After the violence of the disease has been got over, counter-irritants will be found highly beneficial; as well as when it has passed into the chronic state. The cough may be relieved by demulcents.

Sauvages has given the name **PLEURITIS VENTROSA**, and Pringle that of **PLEURODYNIA VENTROSA**, to a pain behind the false ribs, attributed to the presence of air in the intestines.

PLEURITIS BILIOSA, Pleurisy, bilious—p. Bronchialis, Bronchitis—p. Costalis, Pleuritis—p. Diaphragmatica, Diaphragmitis—p. Humida, Bronchitis—p. Muscularis, Pleurodynia—p. Notha, Pleurodynia—p. Pericardii, Pericarditis—p. Pulmonis, Pleuropneumonia—p. Rheumatica, Pleurodynia—p. Spuria, Pleurodynia—p. Spuria simulata, Pleurodynia—p. Typhoides, Pleurisy, typhoid—p. Vera, Pleuritis.

PLEUROCELE, from *πλευρα*, 'the pleura,' and *κηλη*, 'a tumour.' *Hernia of the pleura*, *Hernia pleurica et pulmonalis*. This name, employed by Sagar, is inaccurate, as the pleura never protrudes alone. It only forms a hernia, when it serves as an envelope to the lung, to tumours, or to purulent collections, which have protruded from the thoracic parietes.

PLEUROCOLLE'SIS, from *πλευρα*, 'the pleura,' and *κολληω*, 'I glue.' An adhesion between the pleura costalis and pleura pulmonalis.

PLEUODYNE, Pleurodynia.

PLEUODYNIA, *Pleuralgia*, *Pseudo-pleuritis*, *Pseudo-pleurisy*, *Pleurodynia*, *Autalgia dolorosa*, *Parapleuritis*, *Stethorrhœuma*, *Rheumatismus pectoris*, *Pleuritis rheumatica seu muscularis seu notha seu spuria seu spuria simulata*, *Pneumonia externa*, *Lagonoponos*, *Thoracodynia*, *Dolor Pectoris externus*. Pain in the side, *Stitch in the side*. *False, rheumatic or rheumatismal pleurisy*, (Old Eng.) *Shott*, (Sc.) *Gaik*, (F.) *Point de côté*, *Douleur de côté*, *Pleur-*

isie finisse; from *πλευρα*, 'the side or pleura,' and *δύνη*, 'pain.' A spasmodic or rheumatic affection, generally seated in the muscles of the chest, and, ordinarily, in the intercostals. It is usually augmented by outward pressure, inspiration, coughing, exertion, moving the thorax, and sometimes even the corresponding arm. It is unaccompanied by fever, cough, or dyspnoea, excepting under the form of a *stitch*, when dyspnoea exists. It is, generally, by no means obstinate, and yields promptly to warm applications and to cupping or leeches. If it still resists, a blister will, almost always, remove it.

PLEUODYNIA VENTOSA, *Pleuritis ventosa*.

PLEUROMA, *Costa*, *Pleura*.

PLEUROPATHIA, from *πλευρα*, 'the pleura,' and *πάθος*, 'a disease.' A disease of the pleura.

PLEUROPERIPNEUMONY, *Pleuroperipneumonia*, *Pleuropneumonia*, *Pneumomopleuritis*, *Pneumopleuritis*, *Pleuritis pulmonis*, *Pneumonia pleuritis*. Inflammation occupying the pleura and the lung at the same time. Pleurisy and peripneumony existing simultaneously. *Physical signs.* A dull sound is yielded by percussion for a variable extent over the more dependent portions of the affected side. The loss of sound is partially modified by change of position. The respiratory murmur on auscultation is very feeble or absent inferiorly; higher up, the crepitation or bronchial respiration of pneumonia may be detected; round the root of the lung, and near the inferior angle of the scapula, egophony is generally present. There is seldom much enlargement of the affected side, or displacement of the adjacent organs, as in the case of simple chronic pleurisy.

PLEUROPNEUMA, *Pneumothorax*.

PLEUROPNEUMONIA, *Pleuroperipneumony*.

PLEUROPNEUMONITIS, *Pleuroperipneumony*.

PLEUROPYESIS, *Empyema*.

PLEUORRHAGIA, *Hæmatothorax*.

PLEUORRHŒA, from *πλευρα*, 'the side,' and *ρῶω*, 'I flow.' Accumulation of fluid in the pleura.

PLEUORRHŒA CHYLOSA, *Chylothorax*—p. *Lymphatica*, *Hydrothorax*—p. *Purulenta*, *Empyema*—p. *Sanguinea*, *Hæmatothorax*—p. *Serosa*, *Hydrothorax*.

PLEUORTHOPNŒA, from *πλευρα*, 'the side,' *ὄρθω*, 'erect,' and *πνέω*, 'I respire.' Pain of the side, which does not permit the patient to breathe, except when in the vertical position.

PLEUROSOMUS, from *πλευρα*, 'side,' and *σῶμα*, 'body.' A malformation in which the fissure is somewhat lateral, with evagination extending chiefly upon the upper part of the abdomen and upon the chest; the upper extremity of the fissured side being more or less atrophied.—Vogel.

PLEUROSTOSIS, from *πλευρα*, 'the pleura,' and *ὀστέον*, 'a bone.' Ossification of the pleura; or in the cavity of the pleura. *Osteopleuria*.

PLEUROTETANUS, *Pleurothotonos*.

PLEUROTOTHONOS, *Pleurothotonos*, *Pleurotetanus*, *Tetanus lateralis*, from *πλευροθεν*, 'laterally,' and *τονος*, 'tension.' A variety of tetanus, in which the body is curved laterally by the stronger contraction of the muscles of one side of the body.

PLEUROTTONOS, *Pleurothotonos*.

PLEURUM, *Costa*, *Pleura*.

PLÈVRE, *Pleura*.

PLEXIMETER, *Plexometer*, *Plegometer*,

Plessim'eter. *Anticooposcopy'ium*, *Anticoop'oscope*. (F.) *Plessimètre*, from πλῆσσω, 'I strike,' and μέτρον, 'measure,' 'a measure of percussion.' An ivory plate of a circular or ovoid shape, from an inch and a half to two inches in diameter, and about one-sixth of an inch in thickness. It has either a raised rim or edge, or projecting handles on its upper side, to permit its being held between the finger and thumb of the left hand, whilst it is struck with the right. It is used in percussion of the chest. A piece of metal, a coin, or the finger of the left hand may be used with equal advantage. — See Percussion, mediate.

PLEXOMETER, Pleximeter.

PLEXOR, from πλῆσσω, 'I strike.' *Plessor*. Any instrument employed in percussion. The ends of the fingers of the right hand brought together are the best *plexor*; as the finger of the left is, perhaps, the best *pleximeter*.

PLEXUS, *Plegma*, *Plec'tanē*, *Planē*, *Retē*, from πλεκο, *plecto*, 'I interlace,' 'I entwine.' (F.) *Lacis*. A network of blood-vessels or of nerves. The nervous plexuses, *Implicatio'nes reticulat'æ* seu *Textus nervorum* belong — some to the system of encephalic nerves — others to that of the great sympathetic; whilst some, as the pharyngeal, seem to be formed of the two sets. The plexuses represent complex networks, with more or less loose meshes, formed by the numerous and diversified anastomoses of the nervous filaments; from which proceed other branches, that are distributed to organs, or to other plexuses.

PLEXUS, CARDIAC, Cardiac plexus — p. Carotid, see Carotid nerve — p. Cavernosus, Bulbus vestibuli — p. Choroides, see Choroid — p. Choroideus Medius, Choroidea tela — p. Ciliaris, Ciliary ligament — p. *Cœliacque*, Cœliac plexus — p. Coronarius ventriculi, Gastric plexus — p. Coronary, see Cardiac plexus, and Coronary plexus — p. Gangliiformis semilunaris, Cœliac plexus — p. Gangliiformes, Ganglions, nervous — p. Glandiformes, Ganglions, nervous — p. Glandulosi Peyerii, Peyerii glandulæ — p. Hederaceus, Corpus pampiniforme — p. Hepaticus, Hepatic plexus — p. Infra-thyroideale, see Thyroideale veins — p. *Lombo-abdominal*, Lumbar plexus — p. Median, Cœliac plexus — p. Mesenterii proprius et maximus, Cœliac plexus — p. Nervorum anserinus, Facial nerve — p. *Opistogastrique*, Cœliac plexus — p. Pampiniformis, Corpus pampiniforme — p. Pharyngeal, see Pharyngeal nerve — p. Pneumogastricus, see Pneumogastric nerve — p. Retiformis, Corpus cavernosum vaginæ — p. Reticularis, Bulbus vestibuli, Choroides plexus — p. Solar, Cœliac plexus — p. *Sous-mésentérique*, Hypogastric plexus — p. Testicularis, Spermatie plexus — p. *Trachélo-sous-côtané*, Cervical plexus — p. Vasculosus funiculi spermatici pampiniformis, Corpus pampiniforme.

PLICA, from *plico*, πλεκο, 'to fold,' *Tricho'sis plica*, *Tricho'ma*, *Trichono'sis plica*, *Tricæ*, *T. Incuborum* seu *Serofo'rum*, *plica Polonica* seu *Polon'ica Juda'ica* seu *Cachect'ica* seu *Saxon'ica* seu *caput Medu'sæ* seu *longicauda lateralis*, *Affectio Sarmat'ica*, *Lues Sarmat'ica* seu *Polon'ica* seu *Trichomat'ica*, *Helo'sis*, *Helo'tis*, *Cir'ragra*, *Cir'rhagra*, *C. Polono'rum*, *Eceph'yma trichoma*, *Morbus Cirrho'rum*, *Coma Casà'rea*, *Kolto*, *Rhopato'sis*, *Plicatio*, *Plicatura*, *Matted hair*, *Plaited hair*, *Trichom'atose hair*, (F.) *Plique*, *P. Polonaise*. A disease endemic in Poland, Lithuania, and other parts of Northern Europe; so called on account of its being characterized by interlacing, twisting, and agglutination or matting of the hair. By some it has been regarded as a real disease; by others, as the want of attention to cleanliness. However this may be, it generally appears upon the hair of the head, but sometimes in that of other parts, as the beard, the

hair on the axilla, pubes, &c. Alibert admits three species of plica. 1: *Plica multiform'is* or *Plica caput Medu'sæ*, in which the hairs are mixed and agglutinated in greater or less masses; and this has been again subdivided into two varieties, according as the meshes are almost straight (*P. C. M. lacinia'ta*, (F.) *Plique en lanières*) or much twisted, (F.) *P. en vrilles*.) 2. *Plica longicauda*, (F.) *Plique solitaire ou à queue*; when the hair is united into a single long mass, and 3. *Plica cæspitosa*, (F.) *Plique en masse*, in which the hairs are matted into one large, shapeless mass. Pinel places this disease amongst the *Cutaneous phlegmasiæ*; but it is far from demonstrated, that it is the result of inflammation of the skin. Certain it is, however, that the tangling of the hair is symptomatic of an affection — *sui generis* — of the dermoid system. The idea that it is entirely owing to want of cleanliness, as has been conceived by some, is not tenable. It has been supposed to be produced by a fungous growth — a *mycoderma* or *trichomaphyte*.

PLICA, Plectane — p. Cachectica, Plica — p. Caput Medusæ, Plica — p. Centralis retinæ, see Optic nerve — p. Longicauda lateralis, Plica — p. Lunata, Valvula semilunaris — p. Polonica, Plica — p. Polonica Judaica, Plica — p. Saxonica, Plica — p. Semilunaris, Valvula semilunaris.

PLICÆ CONNIVENTES, Valvulæ conniventes — p. Semilunares, see Urinary bladder, and Uterus.

PLICATIO, Plica.

PLICATURA, Plica.

PLICATURÆ CEREBRI, Gyri cerebri.

PLICHOS, Perineum.

PLINTHIUM, πλινθιον. A machine invented by one Nileus, which was formerly employed in the reduction of fractures and luxations. Scultetus describes it in his *Armamentarium Chirurgicum*.

PLIQUE, Plica — p. Polonaise, Plica.

PLOCARIA CANDIDA, *Fucus amylaceus* — p. Helminthochorton, *Coralina Corsicana*.

PLOMB, Plumbum.

PLOMB (F.). The sulphuretted gas disengaged from privies, which proves fatal at times to the nightmen, (F.) *Vidangeurs*, engaged in emptying them.

PLOMB, ACÉTATE ACIDE DE, Plumbi superacetat — p. *Blanc*, Plumbi subcarbonat — p. *Blanc* de, Plumbi subcarbonat — p. *Carbonate* de, Plumbi subcarbonat — p. *Chlorure* de, Plumbi chloridum — p. *Iodure* de, Plumbi iodidum — p. *Nitrate* de, Plumbi nitrat — p. *Oxide* de, *fondus*, Plumbi oxydum semivitreum — p. *Oxide rouge* de, Plumbi oxydum rubrum — p. *Oxide semi-vitreux* de, Plumbi oxydum semivitreum.

PLOMBAGE DES DENTS, Plugging.

PLOMBAGINE, Gaphites.

PLOMBIÈRES, MINERAL WATERS OF.

Fontes medica'ti Plumba'rii, *Therma Plumbaria*. Plombières is a town in the department of the Vosges, 17 leagues from Nancy, and two from Remiremont. Here are several sources of thermal water. That which is called *Saponaceous* (F.) *Savonneuse*, is formed of the subcarbonate and sulphate of soda, chloride of sodium, silica, carbonate of lime, and much vegeto-animal matter. This last component, when held in solution by means of the subcarbonate of soda, gives to the water an unctuous character. This is the origin of its name. The waters of Plombières are employed as solvents in engorgements of the abdominal viscera, and as alteratives in chronic diseases of the skin.

PLORATIO, Lacrymatio.

PLORATUS, Lacrymatio.

PLOTUS, *Leipus*.

PLOUK, *Papula*.

PLUGGING, (Sw.) *Plugg*, (F.) *Tamponnement*. The act of stopping a hole. The introduction of a plug or dossil of lint or rag into a wound or natural cavity—the uterus or vagina—to arrest hemorrhage; or of some substance into a carious tooth to prevent toothache, (F.) *Plombage des dents*. See Filling.

PLUKE, *Papula*.

PLUM, *Prunum*, *Prunus domestica*—p. Assyrian, *Sebestina*.

PLUMA, *Lanugo*.

PLUMACEAU, *Pledget*.

PLUMACEOLUS, *Pad*, *Pledget*.

PLUMASSEAU, *Pledget*.

PLUMBAGO, *Graphites*, *Persicaria*.

PLUMBA'GO EUROPE'IA, *P. lapathifolia* seu *undulata*, from *plumbum*, 'lead,' because covered with lead-coloured spots, *Dentaria*, *Crepanella*, *Squama'ria*, *Dentelaria*, *Leadwort*, (F.) *Dentelaire*, *Herbe au Cancer*, *Malherbe*. Ord. *Plumbaginaceæ*. Sex. Syst. *Pentandria Monogynia*. The root was formerly much esteemed as a cure for toothache, like the *Pellitory of Spain*, which is also called *Dentaria*.

PLUMBI ACETAS, *P. superacetas*—p. *Acetas dilutus* alcoholicum, *Liquor plumbi subacetatis dilutus*—p. *Acidulus siccus*, *Plumbi superacetas*—p. *Carbonas*, *P. subcarbonas*.

PLUMBI CHLO'RIDUM, *Chloride of Lead*, (F.) *Chlorure de Plomb*. This salt of lead has been used in the form of lotion and ointment in cancerous ulcerations.

PLUMBI DEUTOXYDUM RUBRUM, *P. Oxydum rubrum*—p. *Hydriodas*, *P. Iodidum*—p. *Iodhydras*, *P. Iodidum*.

PLUMBI IODIDUM, *P. Ioduretum* seu *Hydriodas* seu *Hydroi'odas* seu *Iodhydras*, *Plumbum iodatum* seu *Hydroi'odicum*, *Iodide* or *Ioduret of Lead*, (F.) *Iodure de Plomb*, *I. plombique*. This salt is formed by the double decomposition of *ioidide of potassium* and *nitrate of lead*. It is used in serofulous affections internally and externally; and is official in the *Pharmacopœia* U. S. 1851. Dose, gr. ss to gr. iij or iv.

PLUMBI IODURETUM, *P. Iodidum*.

PLUMBI NITRAS, *Plumbum nitricum*, *Nitras plumbicus*, *Nitrum saturninum*, *Nitrate of lead*, (F.) *Nitrate ou Azotate de Plomb*. A salt formed by the action of *diluted nitric acid on litharge*. Nitrate of lead has the same medical action as the other soluble salts of lead, and is a valuable antihomeric. It has been used, also, as an excellent antiseptic to wounds, ulcers, &c., and is the basis of *Ledoyen's disinfecting liquid*.

PLUMBI NITROSACCHARAS, *Nitrosaccharate of lead*. (*Sacchar*. p. j; *Acid. nitric*. p. ij; *Aqua* p. x.) Beat together in a porcelain dish as long as chemical action continues: dilute the liquor with water, neutralize with chalk, and to the filtered solution add acetate of lead so long as a precipitate is formed. Collect, wash, and dry the precipitate, which is saccharate of lead. Dissolve in dilute nitric acid, (one part acid to nineteen of water,) filter and evaporate to form crystals, which are transparent, of an amber colour, and in regular hexagonal plates or prisms.

Advised as a solvent of phosphatic calculi.

PLUMBI OXYDUM ALBUM, *Plumbi subcarbonas*—p. *Oxydum carbonatum per acidum acetosum*, *Plumbi subcarbonas*—p. *Oxydum fusum*, *Pl. oxydum semivitreum*.

PLUMBI OXYDUM RUBRUM, *Min'ium*, *Miltus*,

Deutoxydum plumbi rubrum, *Superoxydum Plumbosum*, *Plumbum oxydatum rubrum* seu *superoxydatum*, *Sandix*, *Sandys*, *Red lead*, (F.) *Oxide de plomb rouge*. It is sometimes used to kill vermin.

PLUMBI OXYDUM SEMIVITREUM, *Plumbum oxydatum*, *Lithargyrus*, *Lithargyrum*, *Lithargyrium*, *Oxydum plumbi fusum*, *Spuma argenti*, *Semi-vitrified oxyd of lead or litharge*, *Calciteo'sea*, *Cath'mia*, (F.) *Oxide de plomb fondu*, *Oxide de plomb semivitreux*, (a yellow protoxyd of lead, prepared by heat, combined with carbonic acid.) It occurs in scales of a whitish-red colour, and semi-vitrified. It is employed for pharmaceutical purposes, as in the *Liquor plumbi acetatis*, *Emplastrum plumbi*, &c. Formerly, the *Litharge of gold*, *Chrys'tis*, *Mas'sicot*, and *Litharge of silver*, *Almariab*, were distinguished in commerce. These were merely ordinary litharge, resembling gold or silver in colour.

PLUMBI SUBCAR'BNAS, *Car'bonas plumbi* seu *plumbicus*, *Plumbi Carbonas* (Ph. U. S.), *Plumbum carbonicum*, *Pl. oxydatum carbonicum*, *Magisterium plumbi*, *Oxydum plumbi album*, *Cerussa*, *Æru'go plumbi*, *Poimmyth'ion*, *Peyn'mythion*, *Simmitium*, *Flake white*, *Blanca*, *Gersa*, *lf'fides*, *Subcar'bonate of lead*, *Cerussa*, *White lead*, (F.) *Carbonate de plomb*, *Blanc de plomb*, *Plomb blanc*, *Céruse*, *Blanc d'Argent*, *Blanc de Céruse*, *Blanc de Krems*, *B. de Krems*, *B. de Kremnitz*, *Craté de plomb*, (prepared by exposing spiral rolls of lead to the fumes of vinegar, in vessels placed in dung.) It is inodorous; has a sweet taste; is brittle; friable; snow white, and of a minute, scaly texture. It is astringent and sedative, and is sprinkled on parts affected with local inflammation, or to prevent chafing in children. It is also used in the formation of ointments and plasters.

PLUMBI SUPERACETAS, *Cerussa acetata*, *Plumbi acidulus siccus*, *P. acet'as* (Ph. U. S.), *Sal seu saccharum Saturni*, *Plumbum acet'icum*, *Dulcedo Saturni*, *Acetas plumbicum crystallisatum*, *Sal plumbi*, *Sugar of lead*, *Supercac'etate of lead*, *Protoac'etate of lead*, (F.) *Acétate acide de plomb*, *Sel ou sucre de Saturne*. (*Plumb. carb. lbj*, *Acid. acetic. cong. 1½*. Boil the lead with the acid till it is saturated; then strain through paper, and evaporate to a pellicle, that crystals may be formed. Wash the crystals and dry on blotting-paper. Ph. L.) The acetate of lead has a sweet, styptic taste; a very white colour, and silky lustre. It is astringent; and, in weak solution, cooling and sedative; in strong, stimulant. It is given internally, in visceral and other hemorrhages, combined with opium, and is used externally, in solution, in inflammation, burns, bruises, gonorrhœa, &c. Dose, gr. ss to iss.

PLUMBI TANNIN, see *Tannin*.

PLUMBICUM ACETAS CRYSTALLISATUM, *Plumbi superacetas*.

PLUMBOSUM SUPEROXYDUM, *Plumbi oxydum rubrum*.

PLUMBUM, *Molyb'dos*, *Molibdus*, *Saturn'us*, *Capricor'nus*, (F.) *Plomb*. Lead. A solid metal of a bluish white colour; more malleable than ductile; soft enough to be scratched by the nail; very fusible and oxydizable, when heated in the air, and existing, in nature, under a number of different shapes; the principal of which are those of sulphuret and oxyd. S. g. 11.352.

Its use is to form oxides and salts. In its metallic state, it has no action upon the body; but when oxydized or united with an acid, it acts, in large doses, as a powerfully astringent poison.

PLUMBUM ACETICUM, *Plumbi superacetas*—p. *Album*, *Tin*—p. *Candidum*, *Tin*—p. *Carbonicum*,

Plumbi subcarbonas—p. Cinereum, Bismuth, subnitrate of—p. Hydroiodicum, Plumbi Iodidum—p. Iodatum, Plumbi Iodidum—p. Nigrum, Graphites—p. Nitricum, Plumbi nitras—p. Oxydatum, Plumbi oxydum semivitreum—p. Oxydatum carbonicum, Plumbi subcarbonas—p. Oxydatum rubrum, Plumbi oxydum rubrum—p. Superoxydatum, Plumbi oxydum semivitreum.

PLUMIERA ALBA; *Order*, Apocynaceæ; a native of tropical America, has a milky juice which is cathartic.

PLUMIERA DRAS'TICA, is given in Brazil in jaundice, and chronic obstructions, [?] mixed with milk of almonds.

PLUMIERA PHAGEDÆNICA, is in repute in Brazil as an anthelmintic.

PLUMS, SEEDED, see Diospyros Virginiana—p. Winter, see Diospyros Virginiana—p. Yellow, see Diospyros Virginiana.

PLUNGE BATH, see Bath.

PLUNGING SIPHON, see Siphon.

PLUNKETT'S CAUSTIC FOR CANCERS, Plunkett's Ointment.

PLUNKETT'S OINTMENT, *Plunkett's Caustic for Cancers*. This empirical cancer remedy consists of *Arsenious acid*, *Sulphur*, and the powdered flowers of the *Ranunculus flammula* and *Cotula fatida*;—levigated and made into a paste with the white of egg, and applied, on a piece of pig's bladder, to the surface of the cancer.

PLURAL BIRTH, see Multiparous.

PLU'TEA. Avicenna gives this name to duplicatures of the dura mater, such as the *Falx cerebri*.

PLUTONIUM, Barium.

PNEOBIOANTIA, Docimasia pulmonum.

PNEOBIOANTICA, Docimasia pulmonum.

PNEUMA, Air, Life. See Pneumatic Physi- cians.

PNEUMAPOSTE'MA, *Pneumonaposte'ma*, *Ab- scessus pulmo'nis*; from *πνεῦμα*, 'the lung,' and *ἀποστήμα*, 'abscess.' Abscess of the lung.

PNEUMARTHRO'SIS, from *πνεῦμα*, 'air,' and *ἄρθρον*, 'articulation.' Secretion of air into an articular cavity. Seen at times after acute rheumatism, but soon disappearing [?].

PNEUMATELECTASIS, Atelectasis pulmo- num.

PNEUMATHÆMIA, *Pneumatæ'mia*, from *πνεῦμα*, *πνευματος*, 'air,' and *αἷμα*, 'blood.' A morbid condition in which air is developed in the bloodvessels.—Cless.

PNEUMATHORAX, Pneumothorax.

PNEUMATIC, *Pneumat'icus*, *Pneumat'ical*, (F.) *Pneumatique*, from *πνεῦμα*, *πνευματος*, 'Life, air.' Relating to pneuma, air or health.

PNEUMAT'IC PHYSICIANS, *Pneumat'ici*. Name given to a sect of physicians, at the head of whom was Athenæus, who made health and disease to consist in the different proportions of an element—which they called *Pneuma*, *πνεῦμα*—to those of the other elementary principles. This *πνεῦμα* was a fancied spiritual principle, which the Stoics regarded as an element, different from water, air, earth and fire;—the sole elementary principles previously admitted.

PNEUMAT'ICA, from *πνεῦμα*, *πνευματος*, 'air.' Appertaining to the breathing. Diseases of the respiratory function;—the second class in Good's system of nosology. Agents that act on the respiratory organs.—Pereira.

PNEUMATICAL, Pneumatic.

PNEUMATOCELE, Physocèle.

PNEUMATO'DES, from *πνεῦμα*, 'wind,' and

εἶδος, 'resemblance.' One distended with air, or who breathes with difficulty, owing to an accumu- lation of flatus in the digestive tube; or, accord- ing to others, owing to emphysema.

PNEUMATOMETER, Spirometer.

PNEUMATOMPHALOCELE, Pneumatom- phalus.

PNEUMATOM'PHALUS, *Pneumatom- ce'lè*, from *πνεῦμα*, *πνευματος*, 'air,' and *μφαλός*, 'the umbilicus.' Umbilical intestinal hernia, containing a considerable quantity of flatus.

PNEUMATOR'RHACHIS, from *πνεῦμα*, *πνευματος*, 'air,' and *ῥαχίς*, 'the spine.' Accumu- lation of air in the vertebral canal.

PNEUMATOSE, Pneumatosis—p. *Péritonéale*, Tympanites.

PNEUMATO'SIS, *Aëro'sis*, from *πνεῦμα*, *πνευματος*, 'air.' *Flatus*, (F.) *Maladie ventreuse*, *Pneu- matose*. Flatulence. Flatulent distension of the stomach or bowels, or both. Sauvages uses the word synonymously with emphysema. It is now generally employed for every enormous collec- tion of gaseous matters in any part of the body.

PNEUMATOSIS, Emphysema—p. Abdominis, Tympanites—p. Enterica, Colica flatulenta—p. Pericardii, Pneumo-pericardium—p. Pulmonum, Emphysema of the lungs—p. Ventriculi, et Pneumatosis enterica, Flatulence.

PNEUMATOTHORAX, Pneumothorax.

PNEUMECTASIS, Emphysema of the Lungs.

PNEUMOCACE, Necropneumonia.

PNEUMOCE'LE, *Hernia pulmo'nis*, (F.) *Hernie du poulmon*. Hernia of the lung through one of the intercostal spaces.

PNEUMOCHOLOSIS, Pneumonia (bilious).

PNEUMOCHYSIS, Œdema of the lungs.

PNEUMOËCTASIE, Emphysema of the lungs.

PNEUMOGAS'TRIC, *Pneumogas'tricus*, (F.) *Pneumogastrique*, from *πνεῦμα*, 'the lung,' and *γαστήρ*, 'the stomach.' Belonging to the lungs and stomach.

PNEUMOGASTRIC NERVES, *Eighth pair of nerves* (Tenth pair, of some,) *Par vagum*, *Pars vaga*, *Nervus vagus seu ambulatorius seu pneumogas- tricus seu dec'imus seu sympath'icus me'dius*, *Middle sympathetic*, (F.) *Nerfs pneumogastriques ou vagues*; so called because it is distributed espe- cially to the organs contained in the chest and ab- domen. This nerve arises from behind the emi- nentiæ olivariæ and near the corpora restiformia by one or two rows of filaments, regularly arranged. These filaments unite to form a flat cord, which issues from the cranium by the foramen lacerum posterius; behind the glosso-pharyngeal nerve. At its exit from the cranium, the pneumogastric nerve has the appearance of a very compact plexus—*Plexus gangliiform'is*, *Ganglion of the superior laryngeal branch*, of Sir Astley Cooper. It then descends along the neck, deeply seated on the outside of the carotid artery and the inter- nal jugular vein. When it has attained the lower part of the neck, it enters the chest, gliding be- hind the subclavian vein, and passing on the right side, in front of the subclavian artery, and, on the left, before the arch of the aorta. It pro- ceeds, at first, backwards, increasing in size. Afterwards, it passes from behind the bronchia to the œsophagus, which it accompanies through the diaphragm, and terminates on the stomach. In the neck, the pneumogastric nerve anasto- moses with the spinal, glosso-pharyngeal, great hypo-glossal, and great sympathetic nerves; and furnishes, 1, a pharyngeal branch, 2, a superior laryngeal, 3, cardiac branches. In the chest, it

gives off, 1, the inferior laryngeal, or recurrent, 2, the pulmonary plexus, and, 3, the œsophageal branches. On entering the abdomen, the œsophageal branches, which terminate the pneumogastric, are distributed upon the stomach, and give numerous filaments to that viscus, to the hepatic, cœliac, gastro-epiploic, and solar plexuses. Some go, also, to the pancreas, liver, gall-bladder, duodenum, &c.

PNEUMOGRAPHY, *Pneumographia*, from *πνεῦμα*, 'the lung,' and *γραφη*, 'a description.' The part of anatomy that describes the lungs.

PNEUMOLITHIASIS, from *πνεῦμα*, 'lung,' and *λίθιασις*, 'formation of stone.' A morbid state in which calculi form in the lungs.

PNEUMOLGY, *Pneumolog'ia*, from *πνεῦμα*, 'the lung,' and *λογος*, 'a discourse.' An anatomical treatise on the lungs. An anatomical description of the lungs.

PNEUMOMETER, Spirometer.

PNEUMOMETRY, *Pneumomet'ria*, from *πνεῦμα*, 'the lung,' and *μετρον*, 'measure.' Measurement of the capacity of the lungs for air. See Spirometer.

PNEUMON, Pulmo.

PNEUMONALGIA, Angina pectoris.

PNEUMONATELECTASIS, Atelectasis pulmonum.

PNEUMONÉCROSIE, Necro-pneumonia.

PNEUMONECTASIS, Emphysema pulmonum.

PNEUMONEMPHRAXIS, from *πνεῦμα*, 'the lung,' and *εμφραξις*, 'obstruction.' Congestion and infarction of the lungs.

PNEUMONIA, *Peripneumo'nia*, 'Empres'ma pneumoni'cie, *Pleumo'nia*, *Peripneumo'nia*, *Pneumoni'tis*, *Pulmo'nia*, *Pulmonaria*, *Inflammati'o pneumoni'ca*, *I. pectoris acuta seu pulmo'nium*, *Pneumostasis*, *Febri's pneumoni'ca*, *Pneumony*, *Peripneumony*, *Inflammation of the Lungs*, *Lung-fever* (vulgarly), (F.) *Inflammation des poudrons*, *Fiebre pneumonique ou peripneumonique*, *Fluxion de poitrine*, *Inflammation du parenchyme pulmonaire*. From *πνεῦμα*, 'the lung.' The chief symptoms of pneumonia are:—pyrexia, accompanied by pain, sometimes obtuse, at others pungent,—in some part of the thorax; pulse more or less quick and hard, according to the violence and extent of the local disorder; pain, aggravated by the cough, which, with dyspnoea, exists throughout the disease. At first, the expectoration is difficult and painful; but, in the course of a few days, it becomes free, and the oppression of breathing is mitigated. When the inflammation, instead of going off by resolution, passes on to suppuration, rigors are experienced; the respiration becomes more oppressed, but less painful, and a sense of weight is felt in one of the pleuræ. Pneumonia may, also, terminate by gangrene—but this rarely happens,—by induration and by hepatization.

Physical Signs. **FIRST STAGE**, (F.) *Engouement*. Diminution of the sound is generally observed over the affected part on percussion; auscultation exhibits an equal, dry, crepitating *râle*, which is best heard at the close of a deep inspiration, and on coughing. The respiratory murmur is intermingled with the crepitant *râle*, or it may be absent. In the neighbourhood of the affected parts it is natural or puerile. The voice and cough are rather more resonant than usual. **SECOND STAGE**, (F.) *Hépatisation rouge*. When hepatization has occurred, the motion of the affected side is impeded, and immediately above the sternum and in the corresponding triangular space on either side, there is often an evident depression. Percussion is dull over the affected

part in every position of the patient. On auscultation, the crepitation is found to be replaced by bronchial respiration. The respiratory murmur is louder in the other portions of the lung, the voice and cough are more resonant, and the heart's action is more distinct. **THIRD STAGE**, (F.) *Hépatisation grise*. When suppuration supervenes, the same signs persist; but usually the voice and cough are less resonant, and a coarse, mucous *râle* is heard. When gangrenous or suppurative cavities are formed, they present the ordinary signs.

The prognosis must be guarded. It is not a disease which we can always subdue. Great extent of inflammation; very considerable oppression; orthopnoea, and difficult expectoration are unfavourable symptoms. The most active treatment is of course necessary. General bleeding forms the sheet anchor,—employed so as to make a decided impression on the system, and repeated, if necessary; local blood-letting; nauseating doses of antimonials; purgatives; and when the violence of the inflammation has been got under,—counter-irritants, &c. Chronic pneumonia sometimes succeeds the acute, and, at others, occurs accidentally. It must be managed on general principles. Small bleedings, especially locally; and counter-irritants of all kinds are indicated.

PNEUMONIA BILIOSA, P. bilious.

PNEUMONIA BILIOUS, P. bilio'sa, *Pneumochol'oisie*. Inflammation of the lungs, accompanied by gastric fever, and not uncommonly by typhoid symptoms;—*Pneumotyphus*, *Pneumo'nia typhoides seu typhoides seu typho'sa*, *Putrid*, *typhoid*, *adynamic*, or *erysipel'atous pneumo'nia*.

PNEUMONIA ERYSIPELATOUS, P. bilious—p. Gangrenosa, Necropneumonia—p. Hypostatica, Pneumonypostasis—p. Intermittens, Pneumotyposis—p. Lobular, see Lobular—p. Notha, Peripneumonia notha—p. Pleuritis, Pleuropneumonia—p. Putrid, P. bilious—p. Typhodes, P. bilious—p. Typhoid, P. bilious—p. Typhosa, P. bilious, Necropneumonia—p. Vesicular, Bronchitis, vesicular.

PNEUMONIC, Pulmonic.

PNEUMONICA. From *πνεῦμα*, 'the lung.' Diseases affecting the lungs, their membranes or motive power; characterized by irregular, impeded, or painful respiration. The second order of the class Pneumatica of Good.

PNEUMONIC'ULA, diminutive of pneumonia. A slight inflammation of the lung.

PNEUMONICUS, Pulmonic.

PNEUMONIE, Pneumonia—p. *Disséminée*, see Lobular—p. *Lobulaire*, see Lobular—p. *Mamelonnée*, see Lobular.

PNEUMONIE DES AGONISANTS, *Pneumonie hypstatique*. The hyperæmic engorgement, which takes place in the lungs during the last moments of life, has been so termed. It would seem to be hyperæmia rather than inflammation.

PNEUMONITIC, *Pneumonit'icus*. Of, or belonging to pneumonitis, or inflammation of the lungs.

PNEUMONITIS, Pneumonia.

PNEUMONOCARCINOMA, from *πνεῦμα*, 'the lung,' and *καρκινωμα*, 'cancer.' Cancer of the lungs.

PNEUMONOCE'LE, from *πνεῦμα*, 'the lung,' and *κλη*, 'rupture.' Hernia of the lung.

PNEUMONOCE'LE DIAPHRAGMAT'ICA INTER'NA. Hernia of the lung through the diaphragm.

PNEUMONOCELE EXTER'NA seu THORAC'ICA. Hernia through the parietes of the chest.

PNEUMONOCELE THORACICA, Pn. externa.
 PNEUMONŒDEMA, Œdema of the lungs.
 PNEUMONOMELANOSIS, see Melanosis.
 PNEUMONOMETER, Spirometer.
 PNEUMONOPHTHISIS, Phthisis pulmonalis.
 PNEUMONOPHTHOE, Phthisis pulmonalis.
 PNEUMONOPLEURESIS, Pleuroperipneumony.

PNEUMONOPLEURITIS, Pleuroperipneumony.

PNEUMONORRHAGIA, Hæmoptysis.

PNEUMONORRHŒA, Hæmoptysis.

PNEUMONOSCIRRHUS, from πνευμων, 'the lung,' and σκίρρως, 'induration.' Induration of the lungs.

PNEUMONOSES, Pneumoses.

PNEUMONY, Pneumonia.

PNEUMONYPOS'TASIS, *Pneumo'nia hypostat'ica*. Pneumonia in a depending portion of the lung, caused by lying on the back.

PNEUMOPÉRICARDE, Pneumo-pericardium.

PNEUMO-PERICARD'IUM, *Pneumo-pericarditis*, *Pneumato'sis pericard'ii*, (F.) *Pneumo-péricarde*, from πνευμα, 'air,' and περικαρδιον, 'pericardium.' Laënnec designates, under this name, the effusion of air into the cavity of the pericardium.

PNEUMOPHTHISIS, Phthisis pulmonalis.

PNEUMOPHYMATA, Tubercles of the lungs.

PNEUMOPLEURESIS, Pleuroperipneumony.

PNEUMOPHTHOE, Phthisis pulmonalis.

PNEUMORRHAGIA, Hæmoptysis — p. Interna, Hæmatothorax.

PNEUMOSEPSIS, Necropneumonia.

PNEUMOSES, from πνευμων, 'the lung.' A term under which Alibert comprises every morbid affection of the lungs.

PNEUMOTHORAX, *Pneumatho'raz*, *Pneumatho'raz*, *Emphyse'ma pec'toris*, *Asthma ævrium* seu *emphysemat'icum*, *Dyspnoea pneumat'ica* seu *physiothorac'ica*, *Physiotho'raz*, *Aërotho'raz*, *Aëropleurie* (Piorry), from πνευμα, 'air,' and θώραξ, 'the thorax.' An accumulation of air in the cavity of the pleura. A complaint generally sudden in its invasion and fatal in its character. In many instances, along with the air, a liquid is effused, — *Hydropneumothorax*. The disease may be spontaneous and simple, but in a large proportion of cases it is complicated with pleurisy and pulmonary tubercles. The signs vary according as there is, or is not, a communication between the pleura and the bronchia. The affected side gives a hollow, tympanitic sound, even where the thickness of the parietes of the thorax is great. When the effusion is considerable, the affected side is dilated, but there is no rôle of any kind. When a gaseous and a liquid effusion are present at the same time, the sound on percussion is found to be clear at the superior part of the thorax, but dull inferiorly; hence, by varying the position of the patient, and by consequence that of the contained fluids, the seats of the clear and the dull sound will be varied. When the gaseous effusion is owing to a fistulous communication between the pleura and bronchia, the metallic respiration and resonances are heard; and if there be both gaseous and liquid effusion, with a fistulous communication, in addition to these signs there is the *tintement métallique*. The presence of liquid can always be ascertained by succussion. See *Emphysema* of the lungs.

PNEUMOTOMY, *Pneumotom'ia*, from πνευμων,

'the lung,' and τέμνω, 'I cut.' Dissection of the lungs.

PNEUMOTYPHUS, Pneumonia, typhoid.

PNEUMOTYPO'SIS, *Pneumo'nia intermit'tens*, from πνευμων, 'the lung,' and τυπος, 'a stamp, a type.' Pneumonia characterized by periodicity.

PNEUBIOMANTIA, Docimasia pulmonum.

PNEUSIOBIOGNOSIS, Docimasia pulmonum.

PNEUSIOBIOSCOPE, Docimasia pulmonum.

PNEUSIS, Respiration — p. Pertussis, Pertussis — p. Singultus, Singultus — p. Tussis, Tussis.

PNEUSOMETER, Spirometer.

PNIGALION, Incubus.

PNIGMOS, Orthopnoea, Strangulation.

PNIGOPHOBIA, Angina pectoris, Suffocation.

PNIX, Orthopnoea, Strangulation.

PNOE, Aura.

PNŒUM. A nostrum of Dr. Samuel Hahnemann, the founder of the 'Homœopathic System,' which consisted — it is said — of borax only!

PNOCOLYTICUS, Asthmatic.

POAD-MILK, Colostrum.

POCHE DES EAUX, see Liquor Amnii, and Parturition.

POCK, from Tent. ποцца, 'pock or little pouch.' A pustule of smallpox.

POCK-ARR, Pock-mark, see Cicatrix — p. Kine, Vaccina.

POCKFRETTE, see Pockmark.

POCKHOLE, Pockmark.

POCKMARK, *Pockhole*, (Prov.) *Pock-arr*, *Scrobic'ulum* seu *Cica'trix Vari'ola*. The mark or pit or pitting left from a smallpox pustule. One pitted with smallpox is said to be *pockfretten* or *pockfreden*.

POCKY. Infected with, or appertaining to, or resembling, smallpox or syphilis.

POC'ULUM ABSINTHIA'TUM. This was regarded, in remote ages, as a wholesome beverage; the *absinthium*, or 'wormwood,' being supposed to act as an antidote to drunkenness.

POCULUM AMATORIUM, Philiter — p. Diogenis, Palm — p. Emeticum, Gobelet, emetic.

POD'AGRA, *Podarthri'tis*, *Podorrhœu'ma*, *Podal'gia*, from πους, *podos*, 'the foot,' and *agras*, 'seizure.' Pain which attacks the feet. Gout, situate in the articulations of the foot. It has, also, been used in a more extensive signification synonymously with *gout*.

PODAGRA ABERRANS, Gout, (wandering) — p. Arthritis, Gout — p. Retrograda, Gout, (retrograde.)

PODAGRARIA, *Ligusticum podagraria* — p. *Ægopodium*, *Ligusticum podagraria*.

PODAG'RIC, *Podag'ric*, *Podag'ricus*, *Arthrit'icus*, *Arthrit'ic*, *Gouty*, (F.) *Goutteux*. Relating or belonging to gout. Affected with, or liable to gout.

PODALGIA, Gout, Podagra.

PODALYRIA TINCTORIA, *Sophora tinctoria*.

PODANŒCEPHALIA, see Podencephalus.

PODARTHRTIS, Podagra.

PODARTHROC'ACÊ, from πους, *podos*, 'the foot,' *apθpov*, 'an articulation,' and *kakos*, 'bad.' Caries of the articulation of the foot.

PODENCEPHALIA, see Podencephalus.

PODENCEPH'ALUS, *Podanenceph'alus*, from πους, *podos*, 'the foot,' and *κεφαλη*, 'the head.' A monster whose brain is placed outside the skull, and seems to be supported on a pedicle, which traverses the summit of the skull. This state of

monstrosity is termed *Podanencephalia*, or more properly, *Podencephalia*.—I. G. St. Hilaire.

PODEX, Anus, Nates.

PODEDE/MA, from *πους*, *podos*, 'foot,' and *οίδημα*, 'a swelling.' Edema of the foot.

PODOL'OGY, *Podolog'ia*, from *πους*, *podos*, 'the foot,' and *λογος*, 'a discourse.' A description of the foot. A treatise on the foot.

PODOPHYLLUM PELTATUM, *Podophyllum*, *Anapodophyllum Canadense*, *May-apple*, *Mandrake*. Ord. Berberidaceæ. Sez. Syst. Polyandria Monogynia; from *πους*, *podos*, 'a foot,' and *φυλλον*, 'a leaf,' in reference to the shape of the leaf. A common plant throughout North America. The fruit is eatable, and esteemed by many. The leaves are said to be poisonous. The root or rhizoma, *Podophyllum* (Ph. U. S.), is purgative in the dose of 20 grains. It has also been used as an anthelmintic.

PODOPHYLLUM MONTANUM, *Mountain May-apple*, *Mandrake*, *Wild Lemon*, *Ducksfoot*, *Raccoon Berry*, *Yellow Berry*, *Ground Lemon*, has the same properties.

PODORRHEUMA, *Podagra*.

PODOTHE'CA, from *πους*, *podos*, 'a foot,' and *θηκη*, 'a receptacle, a sheath.' The cuticle of the foot. An anatomical preparation. *Chirotheca*, has been used for the cuticle of the hand.

POËLETTE, Cup, *Palette*.

POË'PHAGUS, from *πον*, or *ποα*, 'a plant, an herb,' and *φαγω*, 'I eat.' One who subsists on herbs or vegetables. The act of subsisting or feeding on herbs or vegetables is called *Poëphag'ia*. Hence Strabo calls the Irish *ποηφαγοι*.

POGON, Beard.

POGONI'ASIS, *Pogo'nia*, from *παγων*, 'the beard.' A female beard. Also, great strength or quantity of beard. A female having a beard—*Vira'go*.

POGONIUM, *παγωνιον*, diminutive of *παγων*, 'beard.' A weak or small beard.

POGONOL'OGY, *Pogonolog'ia*, from *παγων*, 'a beard,' and *λογος*, 'a description.' A description of, or treatise on, the beard.

POGOSTEMON INTERMEDIUS, P. Patchouli.

POGOSTEMON PATCHOULI, P. *Suavis seu intermedius*, *Pucha-Pat*, *Patchouli*. Nat. Ord. Labiate. Native of Penang and the Malayan peninsula. The oil is used as a perfume. The distilled oil is used for the same purpose.

POGOSTEMON SUAVIS, P. Patchouli.

POHON ANTIAR, Upas—p. Upas, Upas.

POL A favourite aliment in the Sandwich Islands. It is a sort of paste made from the root of the Kalo—*Arum esculentum*.

POIDS ET MÉSURES, Weights and measures.

POIGNÉE, Manipulus.

POIGNET, Carpus.

POIL, Galactoele, Hair.

POILETTE, Cup.

POILS, see *Typha latifolia*.

POING, Fist.

POINT, Stitch.

POINT, BLISTERING. A name given by Dr. Rush to an indeterminate period in the course of a continued fever, intermediate between the stages of high excitement and collapse, in which blisters will—he thought—usually produce unequivocally good effects.

POINT DE COTÉ, Pleurodynia.

POINTS LACRYMAUX, Lachrymalis puncta.

POIRE, see *Pyrus communis*.

POIREAU, Allium porrum, Verruca.

POIRIER, *Pyrus communis*.

POIS, Pisum—p. à Cautère, see *Fonticulus*—p. Chiche, *Cicer arietinum*—p. Cultivé, Pisum—p. à Gratter, *Dolichos pruriens*—p. Patate, *Dolichos tuberosus*.

POISON, *Phar'macon*, *Phar'macum*, *Tox'icum*, *Vene'num*, *Virus*, *Deleter'ium*, *Venenat'ion*, (F.) *Poison*. A generic name for all substances which, when introduced into the animal economy, either by cutaneous absorption, respiration, or the digestive canal, act in a noxious manner on the vital properties or the texture of organs. Hence we speak of *fever poison*, *cholera poison*, &c. Poisons exist in the three kingdoms of nature; but those which proceed from animals are often called *venom*, as the venom of the viper, scorpion, tarantula, &c.; whilst those that are the products of disease have the name *virus*. In common parlance, therefore, *poison* is restricted to deleterious articles, furnished by the mineral and vegetable kingdoms. Orfila has divided poisons into four classes. 1. ACRID, IR'ITATING, CORRO'SIVE, or ESCHAROT'IC, such as the concentrated acids and alkalies; mercurial, arseniacal, cupreous, and antimonial compounds, cantharides, &c. 2. NARCOT'IC; those that act particularly upon the brain; as hyoscyamus, opium, &c., but without inflaming the organ with which they come in contact. 3. NARCOT'ICO-ACRID or ACRO-NARCOT'IC;—those that act on the brain, but, at the same time, irritate the parts to which they are applied; as aconite, belladonna, &c. 4. SEPTIC or PUTRES'CENT;—those furnished by the animal kingdom. See *Venom* and *Virus*. Various classifications, of a similar character, have been recommended by different toxicologists; but they are liable to the objection, that they throw substances together whose physiological action on the system is very different. It is, indeed, difficult to avoid unnatural compression of matters into places not properly belonging to them, in all such arrangements.

The following table, which exhibits a *coup d'œil* of the chief poisons, with most of the circumstances of importance connected with them, is not free from these objections

TABLE, EXHIBITING THE SYMPTOMS, TREATMENT, AND MODE OF DETECTING THE VARIOUS POISONS, MINERAL, VEGETABLE AND ANIMAL.

N. B. In all cases the Stomach-Pump should be used as soon as possible.

I. INORGANIC POISONS.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
ACIDS.	The acids, generally, are strong corrosive poisons.	The carbonates of soda, potassa, lime and magnesia, are all antidotes to the acids; calcined magnesia also. They are to be used with the following restrictions: for the acetic, citric, muriatic, sulphuric, and tartaric acids, they may be used indiscriminately. For the nitric and oxalic, carbonates of magnesia and lime can alone be employed with safety. In the case of sulphuric acid, water should not be drunk, on account of the great heat which is produced by their mixture. Subsequent inflammation to be treated by ordinary means.	<i>Sulphuric acid</i> is known by a white precipitate with any salt of baryta, insoluble in nitric acid.
<i>Acetic Acid.</i>	Sour, acid taste, burning in the throat, which is increased by pressure, swallowing, or coughing; eructation, and excruciating pain in the stomach; more or less corrugation of the lining membranes of the mouth and primæ viæ; exco-riation about the mouth or such other parts of the skin as the acid may have touched. The matter vomited effervesces with carbonate of lime. The countenance becomes glazed, extremities cold and clammy; convulsions and death. Nitric acid occasions yellow stains, and sulphuric acid, black.	The carbonates of the alkalies and of magnesia and lime have the inconvenience, that a large quantity of gas is extricated in the stomach.	<i>Muriatic acid</i> is known by a white precipitate with nitrate of silver, insoluble in nitric acid, which turns black by the action of light.
<i>Citric Acid.</i>		Ammonia is an antidote; but it should not be employed in a very concentrated form. Liquid chlorine has also been found efficacious. The cold <i>douche</i> to the head has been recommended.	<i>Nitric acid</i> gives rise to orange-coloured fumes, when copper is placed in it, and an ability to dissolve gold when mixed with muriatic acid.
<i>Muriatic Acid.</i>			<i>Acetic acid</i> exhales the odour of vinegar.
<i>Nitric Acid.</i>			<i>Citric acid</i> blackens when heated.
<i>Sulphuric Acid.</i>			<i>Tartaric acid</i> precipitates crystals, with a solution of potassa; and a white precipitate is thrown down by lime-water.
<i>Tartaric Acid.</i>			<i>Oxalic acid</i> gives a white precipitate with lime-water, which, when heated, is converted into carbonate of lime.
<i>Oxalic Acid.</i>			
<i>Prussic Acid.</i>	Is a sedative poison; nausea, giddiness, debility, hurried pulse, weight and pain in the head; eructations having the flavour of the acid; spasms, tetanus, contractile pupil; convulsions, death.		<i>Prussic acid</i> exhales the odour of peach-kernels; forms a white precipitate with nitrate of silver neutralized with an alkali, gives a blue precipitate with the per-salts of iron.
<i>Oil of Bitter Almonds.</i>			
<i>Laurel Water.</i>			
ALKALIES and their SALTS.	Violent, caustic, acid taste; great heat in the throat, with destruction of its lining membrane; difficult and painful deglutition; vomiting of bloody matter, which turns the yellow of turmeric brown; acute pain in the stomach; cold sweats, weakness, hic-cough; violent colic pains, with purging of bloody stools and membranous flakes; death.	The vegetable acids, such as vinegar, lemon juice, citric and tartaric acid in solution are antidotes to the alkalies and their carbonates. The fixed oils, such as castor, linseed, almond and olive, form soaps with the free alkalies, and therefore destroy their caustic effects.	The alkalies are known by their action on turmeric paper: restoring the colour of litmus, which has been reddened by an acid.
<i>AMMONIA, Strong Liquor or Water of.</i>			Carbonates are known by their effervescence with an acid; some of them affect the turmeric paper
<i>Muriate of Ammonia or Sal Ammoniac.</i>			Sal ammoniac gives out the pungent odour of ammonia, when mixed with quicklime.
<i>POTASSA.</i>			
<i>Caustic Potassa, and Liquor Potassæ.</i>			
<i>Carbonate of Potassa, or Pearl Ash, and Salt of Tartar.</i>	The nitrate of potassa will not manifest the effect with turmeric paper. The carbonates, when vomited, will effervesce with acids; and the liver of sulphur will give rise to eructations of sulphuretted hydrogen.	* Poisoning by nitrate of potassa to be treated on general antiphlogistic principles; mucilaginous drinks.	Nitrate of potassa decrepitates and deflagrates when thrown on hot coals.
<i>Nitrate of Potassa or Saltpetre.</i>		Liver of sulphur is said to be decomposed and neutralized by common salt. The liquid chloride of soda will also decompose it.	Liver of sulphur emits the odour of sulphuretted hydrogen when dissolved, or when treated with an acid.
<i>Sulphuret of Potassium or Liver of Sulphur.</i>			
<i>SODA.</i>			
EARTHS and COMPOUNDS.	Analogous to those of the corrosive metals. Violent burning in the stomach, vomiting, gripes, diarrhœa; excessive muscular debility, headache, convulsions, death. Lime differs from baryta in being a pure irritant.	The sulphates of soda and magnesia are prompt and effective antidotes to all the poisonous salts of baryta. Phosphate of soda will also counteract their effects. Lime may be neutralized by dilute acids. Carbonic acid, in soda-water, effervescing draught or yeast, it is supposed, would answer a good purpose. The fixed oils may be employed either for baryta or lime, when not in a compound state.	Baryta and its salts invariably give a white precipitate, insoluble in acid, with a soluble sulphate,
<i>BARYTA.</i>			
<i>Carbonate of Baryta.</i>			Lime, when dissolved, gives a white precipitate with oxalic acid. Also with carbonic acid or a soluble carbonate. The same of baryta.
<i>Chloride of Baryum.</i>			
<i>Nitrate of Baryta.</i>			
<i>LIME.</i>			

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
ALCOHOL. <i>Brandy, Wines, and all Spirituous Liquors.</i>	Intoxication, and when taken very freely, complete insensibility, with apoplexy or paralysis of one side; the countenance swollen and of a dark-red colour; the breathing difficult, and often stertorous, with a peculiar puffing out of the lips; the breath smells of liquor, which will distinguish the symptoms from those of spontaneous apoplexy.	A powerful emetic of white vitriol, or tartar emetic, should be got into the stomach as soon as possible; and if the person has lost the power of swallowing, a flexible catheter or tube should be the means of conveying it thither. The vomiting should be encouraged as much as possible with warm water; and large and active glysters of salt and water should be thrown up. The patient should be placed erect, and if the countenance and other appearances be not improved after these means have been used, the jugular vein may be opened, and cold wet cloths applied to the head, particularly if the body be hotter than natural. If the extremities become cold, warmth and friction should be perseveringly used.	
VOLATILE OILS. <i>Creosote.</i> <i>Dippel's Animal Oil.</i> <i>Oil of Tar.</i> <i>Oil of Tobacco.</i> <i>Oil of Turpentine.</i> <i>Fusel Oil.</i>	General action, that of irritant poisons. Burning pain, vomiting, pungent taste purging, &c. The oils of turpentine and tobacco affect the nervous system, the peculiar odour of each oil will be manifested in the matter vomited.	Creosote is immediately coagulated by albumen. Dippel's animal oil may be counteracted with dilute acids and the fixed oils. The other oils have no particular antidotes, and their effects must, therefore, be counteracted upon general principles.	No better mode of recognizing these substances exists, than that derived from their peculiar odour. Dippel's oil has the pungent odour of ammonia; creosote and oil of tar, a peculiar smell of smoke; the odour of tobacco and turpentine are well known; and fusel oil has a peculiar, offensive, suffocating odour.
GASES. <i>Carbonic Acid, or Fixed Air.</i> <i>Carbonic Oxide.</i> <i>Fumes of Burning Charcoal.</i> <i>Chlorine.</i> <i>Sulphuretted Hydrogen.</i>	Chlorine produces, when inhaled, violent irritation of the organs of respiration; cough, bloody expectoration, inflammation of the lungs, and permanent pulmonary disease. The other gases, although producing some effect on the respiratory organs, act as poisons, in consequence of their sedative agency. The symptoms, therefore, are those of apoplexy, or narcotic poisoning.	The antidotes to chlorine are the cautious inhalation of ammonia, or sulphuretted hydrogen. The inflammatory symptoms from chlorine to be treated on general principles. For the other gases, cold affusions to the head, blood-letting, artificial respiration.	Chlorine is recognized by its peculiar odour and bleaching properties. Sulphuretted hydrogen, by its fetid smell, and the rapid blackening of lead. Carbonic acid, by its occasioning turbidity in lime-water placed in its atmosphere. Carbonic oxide, by the blue colour of its flame.
IODINE. <i>Iodide of Potassium.</i>	Irritant symptoms; burning pain in the throat, lacerating pain in the stomach, and fruitless efforts to vomit; suffusion of the eyes; excessive pain and tenderness of the epigastrium.	Iodine combines with starch, and forms an insoluble compound. The prompt administration of starch, wheat flour, or other vegetable matter containing fecula, beat up in water, is recommended. Iodide of potassium has no antidote. Vomiting should be promoted by draughts of warm water, and inflammation be subdued by general treatment.	Iodine is known by its odour, and the formation of a blue precipitate, when brought in contact with a cold solution of starch. Iodide of potassium gives a crystalline precipitate, with tartaric acid in excess. The supernatant fluid will give the blue colour to starch.
METALS. ANTIMONY. <i>Tartar Emetic.</i> <i>Chloride, or Butyrate of Antimony.</i> <i>Oxide of Antimony.</i>	Vomiting. If vomiting do not occur promptly, violent irritant effects are produced. Burning pain in the pit of the stomach; Purging; colicky pains, sense of tightness in the throat, violent cramps; repeated recurrence of vomiting.	If vomiting have not been produced, it should be brought about by tickling the fauces, and administering copious draughts of warm water. Astringent infusions, such as of galls, oak bark, Peruvian bark, act as antidotes, and should be given promptly; powdered yellow bark may be used, until the infusion is prepared.	Antimony in solution is best distinguished by the peculiar orange-red precipitate, which it forms with sulphuretted hydrogen or solutions of the hydrosulphates. Free alkalis produce white precipitates. The muriate is known by a white precipitate, when it is poured into water. In its solid forms, it may be known by the formation of white fumes, when heated, which redden litmus. It may also be converted into chloride, and then precipitated by adding water.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
ARSENIC. <i>Arsenious Acid, or White Arsenic.</i> <i>Orpiment, or Yellow Sulphuret of Arsenic.</i> <i>King's Yellow.</i> <i>Realgar, or Red Sulphuret of Arsenic.</i> <i>Fly Powder.</i> <i>Fowler's Solution.</i> <i>Arsenical Paste.</i> <i>Arsenical Soap.</i> <i>Arsenite of Copper.</i> <i>Scheele's Green.</i>	<p>Violent burning pain in the region of the stomach and bowels; tenderness on pressure; retching; vomiting; sense of dryness and tightness in the throat; thirst; hoarseness and difficulty of speech; the matter vomited, greenish or yellowish, sometimes streaked with blood; diarrhoea; tenesmus; sometimes excoriation of the anus; urinary organs occasionally affected with violent burning pains and suppression; convulsions and cramps; clammy sweats; lividity of the extremities; countenance collapsed; eyes red and sparkling; delirium; death.</p>	<p>The hydrated peroxide of iron diffused through water; or the precipitated carbonate; or the rubigo ferri, in very fine powder, to be administered every five or ten minutes, until relief is obtained. This is particularly efficacious when the white arsenic has been swallowed. If the arsenic have been taken in the form of Fowler's Solution, lime-water, in copious draughts, may be given. For either of the other forms, emetics of sulphate of zinc; diluents; demulcents, such as flaxseed tea, infusion of slippery elm, &c. Counter-irritants may be used to relieve the spasm, and violent pain in the stomach. Bleeding, as promoting absorption, should not be employed until the stomach is completely evacuated.</p>	<p>If the suspected substance can be obtained free from organic matter, the presence of arsenic may readily be shown by boiling it in distilled water, filtering, and testing with, 1st. Ammoniacal sulphate of copper. 2d. Ammoniacal-nitrate of silver. (<i>Hume's Test</i>) 3d. Sulphuretted hydrogen gas. The first will produce a green precipitate; second, a pale yellow precipitate; third, a bright yellow precipitate, if arsenic be present. A portion of the powder, or one of these precipitates, carefully and thoroughly dried, is then to be mixed with charcoal and exsiccated carbonate of soda, to be placed in a small glass tube, closed at one end, and then to be heated to a red heat in the flame of a lamp. The arsenic will be reduced, and sublime in the upper part of the tube, forming a dark crust, having considerable lustre on its exterior surface, resembling polished steel. By gently heating this crust in a very small flame, crystals of white arsenic will be formed, having a high adamantine lustre.</p> <p>If organic matter be present, it must be entirely destroyed, by heating the substance with nitromuriatic acid. After this has been done, and a clear, transparent solution in distilled water has been obtained, the subsequent process is as above indicated. If arsenious acid be submitted to the action of nascent hydrogen, it is deoxidized, and the metallic arsenic, thereby obtained, combining with hydrogen, forms arsenuretted hydrogen gas. (<i>Morsh's Test.</i>)</p> <p>Add to a suspected solution a few drops of pure chlorohydric acid, and place in it a slip of bright copper; no change occurs until the liquid is brought to the boiling point, when, if arsenic be present, even in small quantity, the copper acquires an iron-gray coating, from the deposit of that metal. Remove the slip of copper; wash it in water, dry it and gradually heat it in a reduction tube, when arsenious acid will be sublimed in minute octohedral crystals. The test succeeds with powdered arsenic, the arsenites, arsenic acid, the arseniates, and orpiment. It will even separate the arsenic from arsenite of copper, and from common lead-shot. <i>Reinsch's Test.</i></p>
BISMUTH. <i>Nitrate of Bismuth.</i> <i>Pearl Powder.</i> <i>Oxide of Bismuth.</i>	<p>Similar to those produced from other irritant poisons. General inflammation of the whole alimentary canal; suppression of urine; hicough, disagreeable metallic taste; vomiting; cramps; delirium; death.</p>	<p>Milk, and sweet mucilaginous drinks are recommended. Leeches, general bleeding; glysters; fomentations to be employed on the general principles of treatment for inflammatory symptoms.</p>	<p>Bismuth is known by the formation of a white precipitate, when its solution in nitric acid is poured into water; and by the formation of a black precipitate with sulphuretted hydrogen gas, and hydrosulphates in solution.</p>
COPPER. <i>Sulphate of Copper</i> <i>Blue Vitriol.</i> <i>Acetate of Copper,</i> <i>Verdigris.</i> <i>Carbonate of Copper,</i> <i>Blue Verditer.</i> <i>Arsenite of Copper</i> <i>Scheele's Green.</i> <i>Food cooked in dirty copper vessels, or pickles made green by copper.</i>	<p>Very similar to those produced by arsenic. Coppery eructations and taste. Fatal cases are generally terminated by convulsions, palsy, insensibility.</p>	<p>Albumen to be administered in either of its forms which can be most readily obtained, as milk or whites of eggs. Vinegar should not be given. The inflammatory symptoms are to be treated on general principles, and so of the nervous.</p>	<p>The presence of copper is readily shown in solutions. By ammonia a beautiful bluish precipitant is thrown down, which dissolves in an excess of the precipitant, and the solution has a rich blue colour. A bright iron bar, introduced into solutions of copper, occasions the separation of the copper, in a metallic form, which deposits itself upon the iron.</p>

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
GOLD. <i>Chloride of Gold.</i> <i>Fulminating Gold.</i>	Very analogous to those of other irritant poisons. This substance communicates a pink stain to the flesh, and patches of that colour may be found about the lips, and inside the mouth.	The salts of gold are decomposed by sulphate of iron; and this has, therefore, been recommended as an antidote. Inflammatory symptoms to be treated on general principles.	Chloride of gold is very readily decomposed, and the gold separated in a metallic state. Proto-sulphate of iron, nitrate of silver and protochloride of tin, all answer this purpose. The precipitated powder will be found insoluble, except in nitro-muriatic acid.
IRON. <i>Sulphate of Iron,</i> <i>Copperas,</i> <i>Green Vitriol.</i> <i>Chloride of Iron.</i>	Symptoms of irritant poisoning; colic pains, constant vomiting and purging. Violent pain in the throat, tension of the epigastrium, coldness of the skin, and feebleness of the pulse.	Carbonate of soda would be a most excellent antidote to either of these substances. Mucilaginous drinks might also be employed; and particular symptoms relieved by general treatment.	Iron is recognised by a bluish-black precipitate, with tincture of galls; rich blue, with prussiate of potassa; and a greenish or reddish precipitate, with the free alkalies or their carbonates.
LEAD. <i>Acetate of Lead.</i> <i>Sugar of Lead.</i> <i>Carbonate of Lead.</i> <i>White Lead.</i> <i>Red Oxide, or Red Lead.</i> <i>Litharge.</i> <i>Wines sweetened by Lead.</i> <i>Water which has been kept in Leadens vessels.</i> <i>Acid food, cooked or left standing in vessels glazed with Lead.</i>	Irritation of the alimentary canal; spasms; nervous symptoms; paralysis, either partial or complete. When taken for some time, in small quantity, violent and obstinate colic; rigidity of abdominal muscles, cramps; remission of pain; obstinate constipation, urine diminished, saliva increased; countenance anxious and gloomy. If relief be not promptly obtained, giddiness, debility, torpor, coma, convulsions and death. The paralysis affects generally the upper extremities.	Sulphate of magnesia and phosphate of soda are both good antidotes for the soluble salts of lead. For the solid forms, dilute sulphuric acid may be drunk. These are applicable to the irritant forms of poisoning by lead. In the chronic form, or colica pictonum, purgatives and anodynes are resorted to; venesection; and external applications to relieve the cramps. The use of strychnia is recommended for the paralysis.	Soluble salts of lead yield a white precipitate with sulphates and carbonates in solution. Salts of chromic acid and iodide of potassium occasion a yellow precipitate. Sulphuretted hydrogen, a black precipitate. If solid, the lead may be converted into a state of solution by nitric acid, and the dilute solution be tested as above.
MERCURY. <i>Corrosive Sublimate.</i> <i>Cyanide of Mercury.</i> <i>Nitrate of Mercury.</i> <i>White Precipitate.</i> <i>Red Oxide, or Red Precipitate.</i> <i>Sulphate, or Turbith Mineral.</i> <i>Vermilion, or Red Sulphuret.</i>	Violent symptoms of irritant poisoning; harsh metallic astringent taste; burning pain in the stomach; vomiting and purging frequently of bloody matter; often irritation of the urinary organs, and sometimes suppression; tightness and burning in the throat, occasionally so great as to prevent speech; countenance not always pale, but sometimes flushed; tendency to doze; stupor, convulsions and death.	Albumen, in some form, must be promptly administered; either white of eggs beaten up with water, milk, or wheat flour beaten up. The inflammatory symptoms to be counteracted by the usual means. Gold, finely mixed in dust, with fine iron filings. (?)	The free alkalies differ in their precipitates from solutions of mercury. Potassa gives, with corrosive sublimate, a yellowish one; ammonia, a white; lime-water, an orange; and sulphuretted hydrogen, a black. If a drop of the solution be placed on a piece of gold, and touched with a knife or key, the mercury will be reduced, which amalgamates with the gold, producing a white spot. The nitrate of mercury gives a black precipitate with the free alkalies. All the solid forms of mercury are volatile, and may be reduced by heating them in a glass tube with charcoal and carbonate of soda, when the metallic mercury will sublime, and may be collected in a globule.
SILVER. <i>Nitrate of Silver,</i> <i>or Lunar Cautic.</i>	Those of other irritant poisons.	Chloride of sodium, or common salt, immediately decomposes this substance, and destroys its activity. Antiphlogistic treatment is to be employed for the inflammatory symptoms.	Nitrate of silver is distinguished by the formation of a white precipitate, insoluble in nitric acid, with chloride of sodium; a gray precipitate with ammonia, which is redissolved in an excess of ammonia; yellow precipitate with phosphate of soda. The precipitate of chloride of silver, obtained by adding muriate of soda to the nitrate of silver, is readily reduced, and the metallic silver obtained, by mixing it with dilute sulphuric acid and granulated zinc.
TIN. <i>Chloride of Tin.</i> <i>Solution of Tin,</i> <i>used by Dyers.</i> <i>Oxide of Tin, or Putty Powder.</i>	The same as those from other irritant poisons, and a peculiar tanned appearance of the villous coat of the stomach.	Milk to be given copiously, and the subsequent treatment to be regulated by the symptoms.	The chloride precipitates vegetable colouring matter; also albumen and gelatin. The white oxide is precipitated from its solution by free alkalies; with sulphuretted hydrogen gas, a brown precipitate.
ZINC. <i>Sulphate of Zinc,</i> <i>White Vitriol.</i> <i>Acetate of Zinc.</i>	Violent vomiting; astringent taste, burning pain in the stomach; pale countenance; cold extremities; dull eyes; fluttering pulse. Death seldom ensues, in consequence of the emetic effects.	The vomiting may be relieved by copious draughts of warm water. Carbonate of soda, administered in solution, will decompose the sulphate of zinc. Milk and albumen also act as antidotes. General principles to be observed in the subsequent treatment.	A white precipitate is thrown down by the free alkalies from the soluble salts of zinc, which becomes yellow when strongly heated—recovering its white colour on cooling. Sulphuretted hydrogen occasions a white precipitate.

POISONS.	SYMPTOMS.	TREATMENT.	TESTS.
PHOSPHORUS.	Symptoms of irritant poisoning; pain in the stomach and bowels; vomiting, diarrhoea, tenderness and tension of the abdomen.	An emetic to be promptly administered; copious draughts containing magnesia in suspension; mucilaginous drinks; general treatment for inflammatory symptoms.	Phosphorus is known by its ready inflammability, luminousness at ordinary temperatures, and alliaceous odour.
GLASS or ENAMEL.	If taken in very coarse powder, it produces irritation and inflammation of the bowels.	Large quantities of crumb of bread should be eaten, to envelop the particles. (?) An emetic of sulphate of zinc should then be given, and vomiting be promoted by demulcent drinks.	

II. ORGANIC POISONS.

I. VEGETABLE POISONS.

IRRITANT POISONS.	SYMPTOMS.	TREATMENT.
<p><i>Anemone pulsatilla</i>. Wind Flower. <i>Arum maculatum</i>. Wake Robin. <i>Bryonia dioica</i>. Bryony. <i>Caladium seguinum</i>. Dumbcane. <i>Calla palustris</i>. Water Arum. <i>Caltha palustris</i>. Marsh Marigold. <i>Chelidonium majus</i>. Celandine. <i>Clematis vitalba</i>. Virgin Bower. <i>Convolvulus jalapa</i>. Jalap. <i>Convolvulus scammonia</i>. Scammony. <i>Croton tiglium</i>. Purging Croton. <i>Cucumis colocynthis</i>. Colocynth. <i>Cyclamen Europæum</i>. Sow Bread. <i>Daphne genkium</i>. Spurge Flax. <i>Daphne mezereum</i>. Mezereon. <i>Doliphinium staphisagria</i>. Stavesacre. <i>Dioica palustris</i>. Swamp leather-wood. <i>Equisetum hyemale</i>. Scourgrass. <i>Euphorbia officinarum</i>. Euphorbium. <i>Spurge</i>. <i>Gratiola officinalis</i>. Hedge Hyssop. <i>Hippomane mancinella</i>. Manchineel. <i>Urtica crepitans</i>. Sand Box. <i>Hydrocotyle vulgaris</i>. Marsh Pennywort. <i>Jatropha curcas</i>. Indian Nut. <i>Jatropha manihot</i>. Cassada. <i>Juniperus sabina</i>. Savin (oil of). <i>Juniperus Virginiana</i>. Red Cedar (oil of). <i>Momordica elaterium</i>. Squinting Cucumber. <i>Narcissus pseudo-narcissus</i>. Daffodil. <i>Pastinax sativa</i>. Common Parsnep. <i>Pedicularis palustris</i>. Marsh Lousewort. <i>Phytolacca decandra</i>. Poke. <i>Piper cubeba</i>. Cubebs. <i>Plumbago Europæa</i>. Toothwort. <i>Ranunculus acris</i>, and other species. <i>Crowfoot</i>. <i>Rhododendron chrysanthemum</i>. Oleander. <i>Ricinus communis</i>. Castor Oil Plant. <i>Sambucus ebulus</i>. Elder. <i>Sedum acre</i>. Stone Crop. <i>Stalagmitis cambogioides</i>. Gamboge. <i>Tanacetum vulgare</i>. Tansy (oil of).</p>	<p>The general effects of this class of vegetable poisons are;—an acrid pungent taste, with more or less of bitterness, excessive heat, great dryness of the mouth and throat, with sense of tightness there; violent vomiting, the efforts being continued even after the stomach is emptied; purging, with great pain in the stomach and bowels; pulse strong, frequent, and regular; breathing often quick and difficult; appearances of intoxication; the pupil of the eye frequently dilated, insensibility resembling death; the pulse becomes slow and loses its force, and death closes the scene.</p> <p>If applied externally, many of them produce violent inflammation of the skin, with vesications or eruptions of pustules.</p>	<p>If vomiting have been occasioned by the poison, and the efforts are still continued, they may be rendered easier by large draughts of warm water, or thin gruel; but if symptoms of insensibility have come on without vomiting, it ought to be immediately excited by the sulphate of zinc, or some other active emetic substance, and after its operation a sharp purgative should be given. After as much as possible of the poison is got rid of, a very strong infusion of coffee, or vinegar diluted with water, may be given with advantage. Camphor mixture with ether may be taken frequently, and if insensibility be considerable, warmth, frictions, and blisters, may be employed. If inflammation or other dangerous consequences have been induced, they are to be treated upon general principles.</p> <p>The fruit of <i>Feuillea Cordifolia</i> has been recommended as a powerful antidote against vegetable poisons; it is to be used in as recent a state as possible. (?)</p> <p>N. B. Bromine, chlorine, and iodine are said to be antidotes to the alkaloïds generally.</p>
ACRO-NARCOTIC.		
<p><i>Ethusa cyapium</i>. Common Fool's Parsley. <i>Aconium napellus</i>. Monkshood. <i>Agaricus</i>, five species. Mushrooms (poisonous). <i>Amanita muscaria</i>. Truffles (poisonous). <i>Anagallis arvensis</i>. Meadow Pimpernel. <i>Aula Gomezi</i>. <i>Apocynum androsaemifolium</i>. Dogsbane. <i>Aristolochia clematidis</i>. Birthwort. <i>Arnica montana</i>. Leopard's bane. <i>Asclepias Syriaca</i>. Swallowwort. <i>Atropa belladonna</i>. Deadly Night Shade. <i>Eschulus Ohioensis</i>. Buckeye. <i>Brucra antidysenterica</i>. False Angustura Bark. <i>Cerbera</i> three species. <i>Cerbera</i>. <i>Cherophyllum sylvestre</i>. Bastard Hemlock.</p>		

IRRITANT POISONS.	SYMPTOMS.	TREATMENT.
<p><i>Chaetothia toriciana</i>. Rathane. <i>Chenopodium murale</i>. Wormseed. <i>Cicuta maculata</i>. American Hemlock. <i>Cicuta virosa</i>. Water Hemlock. <i>Cissus</i>. Cissus. <i>Cocculus Indicus</i>. Fish berries. <i>Calcheum autumnale</i>. Meadow saffron. <i>Conium maculatum</i>. Hemlock. <i>Coriaria myrtifolia</i>. Myrtle-leaved Sumach. <i>Curat</i>. Indian War Poison. <i>Cynanchum erectum</i>. Cynanchum. <i>Cytosus laburnum</i>. Laburnum. <i>Datura stramonium</i>. Thorn Apple. ♀ <i>Digitalis purpurea</i>. Foxglove. <i>Ergot</i>. See <i>Scalce</i>. <i>Eryum cretita</i>. Bitter Vetch. <i>Gaultheria procumbens</i>. Wintergreen (oil of). <i>Hemnanthus toxicarius</i>. <i>Helleborus niger</i>. Black Hellebore. <i>Hypophyllum sanguinaria</i>. Paddock Stool. <i>Iperacuanha</i>. Iperacuanha. <i>Lathyrus cicera</i>. Lathyrus. <i>Laurus camphora</i>. Camphor. <i>Lobelia inflata</i>. Indian Tobacco. <i>Lolium temulentum</i>. Darnel. <i>Melia azederach</i>. Pride of China. <i>Mercurialis perennis</i>. Mountain Mercury. <i>Nerium oleander</i>. Common Oleander. <i>Nicotiana tabacum</i>. Tobacco. <i>Oenanthe crocata</i>. Hemlock Dropwort. <i>Pasiflora quadrangularis</i>. Barbadoe. <i>Piselia erythrina</i>. Jamaica Dogwood. <i>Polysila renouosa</i> (of Java). <i>Rhus radicans</i>. Poison Vine. <i>Rhus toxicodendron</i>. Poison oak, or Sumach. <i>Robinia pseudo-acacia</i>. Locust Tree. <i>Ruta graveolens</i>. Rue. <i>Sanguinaria Canadensis</i>. Blood Root. <i>Scilla maritima</i>. Squill. Sea Onion. <i>Scalce cornutum</i>. Ergot. Spurred Rye. <i>Sium latifolium</i>. Procumbent Water Parsnep. <i>Spigelia Marilandica</i>. Pink Root. <i>Strychnos Ignatii</i>. St. Ignatius's Bean. <i>Strychnos nux vomica</i>. Nux vomica. <i>Scmplecarpus fatida</i>. Stunk Cabbage. <i>Ticunas</i>. Extract of various plants—South America. <i>Tiouté</i>. <i>Upas Tiouté</i>. Tree of Java. <i>Triticum Hybernium</i>. Wheat (diseased.) <i>Upas antiar</i>. Tree in Java. <i>Veratrum album</i>. White Hellebore. <i>Veratrum viride</i>. American Hellebore. <i>Woorara</i>. War Poison of Guiana. <i>Zea mays</i>. Maize (diseased).</p>	<p>The narcotic vegetable poisons, if taken into the stomach or applied to a wound, occasion the following effects:—stupor; numbness; heaviness in the head; desire to vomit, slight at first, but afterwards insupportable; a sort of intoxication, stupid air, pupil of the eye dilated; furious or lively delirium, sometimes pain; convulsions of different parts of the body, or palsy of the limbs. The pulse is variable, but at first generally strong and full; the breathing is quick, and there is great anxiety and dejection, which, if not speedily relieved, soon ends in death.</p>	<p>The stomach to be effectually evacuated by giving four or five grains of tartar emetic, or from ten to twenty of the sulphate of zinc, repeated every quarter of an hour, till the full effect is produced. These means may be assisted by tickling the throat with a feather or the finger. Large and strong glysters of soap dissolved in water, or of salt and gruel, should be speedily administered, to clear the bowels, and assist in getting rid of the poison; and active purgatives may be given after the vomiting has ceased. When as much as possible of the poison has been expelled, the patient may drink, alternately, a tea-cupful of strong hot infusion of coffee, and vinegar diluted with water. If the drowsiness, which is sometimes extreme, and the insensibility bordering on apoplexy, be not remedied by these means, blood may be taken from the jugular vein, blisters may be applied to the neck and legs, and the attention be roused by every means possible. If the heat declines, warmth and frictions must be perseveringly used. Vegetable acids are on no account to be given before the poison is expelled, and it is desirable that but little fluid of any kind should be administered.</p> <p>N. B. Bromine, chlorine, and iodine are said to be antidotes to the alkaloids generally.</p>
<p>NARCOTIC. <i>Actea spicata</i>. Baneberry. <i>Amygdalus communis</i>. Bitter Almond. <i>Amygdalus Persica</i>. Peach. <i>Gelsemium nitidum</i>. Yellow Jessamine. <i>Helonias erythrosperma</i>. Fly Poison. <i>Hyoecyanus albus</i>. White Henbane. <i>Hyoecyanus niger</i>. Black Henbane. <i>Kalmia latifolia</i>. Mountain Ivy. <i>Lactuca virosa</i>. Strong-scented Lettuce. <i>Opium</i>, and its proximate principles. <i>Papaver somniferum</i>. Poppy. <i>Paris quadrifolia</i>. Herb Paris. <i>Prunus Caroliniana</i>. Wild Orange. <i>Prunus lauro-cerasus</i>. Cherry Laurel. <i>Prunus nigra</i>. Black Cherry. <i>Prunus padus</i>. Cluster Cherry. <i>Prunus Virginiana</i>. Wild Cherry. <i>Solanum dulcamara</i>. Bitter-sweet. <i>Sorbus acuparia</i>. Mountain Ash. <i>Taxus baccata</i>. Yew.</p>	<p>Nausea, heat, and pain in the stomach and bowels, with vomiting and purging; thirst; convulsions, and faintings; pulse small and frequent; delirium; dilated</p>	<p>The stomach and bowels to be cleared by an emetic of tartarized antimony, followed by frequent doses of Glimmer's or Epsom salt, and large, stimulating glysters. After the poison is evacuated, ether may be administered, with small</p>
<p>POISONOUS MUSHROOMS. <i>Agaricus muscarius</i>. Fly Agaric. <i>Agaricus piperatus</i>. Pepper Agaric. <i>Agaricus necator</i>. Deadly Agaric. <i>Agaricus bulbosus</i>. Bulbous Agaric. <i>Agaricus chantarellus</i>. Champignon.</p>		

POISONS.	SYMPTOMS.	TREATMENT.
	pupil, and stupor; cold sweats, and death. Poisonous mushrooms may be distinguished from the edible by their botanical characters, and by the following criteria. The former grow in wet, shady places, have a nauseous odour; are softer, more open, and porous; have a dirty-looking surface, sometimes a gaudy colour, or many very distinct hues, particularly if they have been covered with an envelope; they have soft, bulbous stalks, grow rapidly, and corrupt very quickly.	quantities of brandy and water; but if inflammatory symptoms manifest themselves, such stimuli should be omitted, and other appropriate means be had recourse to.

2. ANIMAL POISONS.

POISONOUS FISH.	SYMPTOMS.	TREATMENT.
<p><i>Balistes monoceros</i>. Old wife. <i>Cancer astacus</i>. Crawfish. <i>Cancer rucicolus</i>. Land Crab. <i>Clupea larvissa</i>. Yellow-billed Sprat. <i>Coracius fascus major</i>. Gray Snapper. <i>Coracius minor</i>. Hyne. <i>Coryphæna splendens</i>. Dolphin. <i>Mormyra</i>. Blue Parrot Fish. <i>Muraena major</i>. Conger Eel. <i>Mytilus edulis</i>. Mussel. <i>Ostracion lobellum</i>. Smooth Bottle Fish. <i>Perca major</i>. Barracuda. <i>Perca venenosa</i>. Grooper. <i>Perca venenata</i>. Rock Fish. <i>Physalia</i>. Portuguese Man of War. <i>Scomber caeruleus</i>. Spanish Mackerel. <i>Scomber maximus</i>. King Fish. <i>Scomber thynnus</i>. Bonetta. <i>Sparus chrysops</i>. Porgée. <i>Tetrodon sceleratus</i>. Tunny. <i>Tetrodon ocellatus</i>. Blower.</p>	<p>In an hour or two, or often in a much shorter time, after the fish has been eaten, a weight at the stomach comes on, with slight vertigo and headache, sense of heat about the head and eyes, considerable thirst, and often an eruption of the skin (urticaria), and in many cases death.</p>	<p>An emetic should be speedily administered, or, in the absence of it, vomiting may be excited, by tickling the throat with the finger, and taking large draughts of warm water. After full vomiting, an active purgative should be given, to remove any of the noxious matter that may have found its way into the intestines. Vinegar and water may be drunk after the above remedies have operated, and the body may be sponged with the same. Water, made very sweet with sugar, to which ether may be added, may be drunk freely as a corrective, and a very weak solution of alkali has been recommended to obviate the effects of the poison. If spasm ensue; after evacuations, landanum, in considerable doses, is necessary. If inflammation should occur, the usual means of removing it must be employed.</p>
POISONOUS SERPENTS.		
<p><i>Boa crocoides</i>. Copperhead. <i>Cenchris mockeson</i>. Mockeson. <i>Cerastes nasicornia</i>. Horned Viper of Western Africa. <i>Coleuber berus</i>. Viper. <i>Coleuber prester</i>. Black Viper. <i>Crotalus</i>, (5 species). Rattle Snake <i>Scytale piscivorus</i>. Water Viper.</p>	<p>A sharp pain in the wounded part, which soon extends over the limb or body; great swelling, at first hard and pale, then reddish, livid, and gangrenous in appearance; faintings, vomitings, convulsions, and sometimes jaundice; pulse small, frequent, and irregular; breathing difficult, cold sweats, the sight fails, and the intellectual faculties are deranged. Inflammation, and often extensive suppuration and gangrene, followed by death.</p>	<p>A cupping-glass to be applied over the wound, or a moderately tight ligature above the bites, and the wound left to bleed after being well washed with warm water; the actual cautery, lunar caustic, or huter of antimony, to be then applied freely to it, and afterwards covered with lint, dipped in equal parts of olive oil and spirit of hartshorn. The ligature to be removed if the inflammation be considerable. Warm, diluting drinks, and small doses of ammonia or hartshorn, to cause perspiration; the patient to be well covered in bed, and a little warm wine given occasionally. If gangrene be threatened, wine may be given more freely, and the bark should be had recourse to. Arsenic, the principal ingredient in the Tanjore Pill, has been strongly recommended.</p>
CANTHARIS VESICATORIA.		
<p>Spanish, or Blistering Fly. <i>Lytta vittata</i>. Potato Fly.</p>	<p>Nauseous odour of the breath; acrid taste; burning heat in the throat, stomach, and abdomen; frequent vomitings, often bloody, with copious bloody stools; excruciating pain in the stomach; painful and obstinate priapism, with heat in the bladder, and strangury or retention of urine; frightful convulsions, delirium and death.</p>	<p>Vomiting to be excited by drinking sweet oil, sugar and water, milk, or linseed tea, very freely. Emollient glysters should be administered, and if symptoms of inflammation of the stomach, kidney, or bladder supervene, they must be subdued by appropriate treatment. Camphor dissolved in oil may be rubbed over the belly and the thighs.</p>
VENOMOUS INSECTS		
<p>Tarantula. Scorpio. Scorpion. <i>Vespa crabro</i>. Hornet. <i>Vespa vulgaris</i>. Wasp. <i>Apis mellifica</i>. Bee. <i>Culex pipiens</i>. Gnat. <i>Estrus bovis</i>. Gad-fly.</p>	<p>In general, the sting of these insects occasions only a slight degree of pain and swelling; but occasionally the symptoms are more violent, and sickness and fever are produced by the intensity of the pain.</p>	<p>Hartshorn and oil may be rubbed on the affected part, and a piece of rag, moistened in the same, or in salt and water, may be kept upon it till the pain is removed. A few drops of hartshorn may be given frequently in a little water, and a glass or two of wine may be taken. The sting may, in general, be removed by making strong pressure around it with the barrel of a small watch-key.</p>

	SYMPTOMS.	TREATMENT.
SALIVA OF THE RABID DOG.	At an uncertain interval after the bite, generally, however, between the twentieth day and three or four months, pain or uneasiness occurs in the bitten part, though the wound may have been long healed. Anxiety, uneasiness, languor, spasms, horror, disturbed sleep, difficult respiration succeed, and are soon very much increased; violent convulsions affect the whole body, hideously distorting the muscles of the face; the eyes are red and protruded, the tongue swells and often hangs out, and viscid saliva flows from the mouth; there is pain in the stomach, with bilious vomitings, a horror of fluids, and impossibility of drinking them. All these symptoms are aggravated till the sufferer is relieved by death.	Hydrophobia is more easily prevented than cured; indeed, it is doubted whether it ever has been cured. Mercury, arsenic, opium, musk, camphor, acids, wine, vegetable and mineral alkali, oil, various herbs, and many other remedies whose effects are quite opposite, have been employed, but none can be relied on. Large blood-lettings, the warm and cold bath, and almost every other remedial agent, have been tried without success. The bitten part should be completely cut out, even after it has healed, if the symptoms have not yet come on; the part should then be immersed in warm water, or be washed with it as long as it will bleed, and, after the most persevering ablution, caustic should be applied to every part of the surface, and then the wound covered with a poultice, and suffered to heal by granulation. No milder discipline can ensure safety.

The most energetic poisons are used therapeutically, and often with considerable advantage. They have, of course, to be administered in extremely small doses, to avoid producing poisoning.

POISON, ARROW, see Arrow poison — p. Berry tree, *Melia azedarach* — p. Hemlock, *Conium*.

POISON, ITALIAN, *Liquor arsenicalis* — p. Morbid. Virus.

POISON OAK, *Rhus toxicodendron* — p. Root, *Cicuta maculata*.

POISON, SAUSAGE, *Allantotoximum* — p. Vegetable, *Upas*.

POISON VINE, *Rhus radicans*.

POISONING, *Venefic'ium*, *Toxica'tio*, *Intoxica'tio*, *Entoxia'mus*, *Entoxici'mus*, *Pharmacei'a*, *Malefic'ium*, *Venenat'ion*, (F.) *Empoisonnement*, *Intoxication*. The art of applying a poisonous substance to any of the textures in a dose capable of occasioning more or less serious symptoms. Rigorously speaking, therefore, if a physician, in treating any case, employs a deleterious medicine in a small dose, which, owing to particular circumstances, produces unpleasant symptoms, he may be said to have poisoned. The term is, however, generally restricted to poisoning by design.

Poisoning has also been used, by a few, to denote the series of symptoms resembling those produced by certain poisons, which occasionally follow the use of substances that are generally innocuous. Overloading the stomach with indigestible food, has, for example, given occasion to symptoms resembling those produced by a narcotic poison, and the effect has been termed poisoning.

Secret poisoning was, at one time, practised as an art. It consisted in administering medicines, which should act as slow poisons, and gradually undermine the strength and life. Its agency was far from being so extensive as was imagined.

POISONOUS, *Tox'ical*, *Nox'ius*, (F.) *Toxique*, *Vénéneux*. Having the qualities of a poison.

POISONS, ACRO-NARCOTIC, P. narcotico-acrid — p. Corrosive, P. acrid — p. Escharotic, P. acrid — p. Irritating, P. acrid — p. Putrescent, P. septic. See Poison.

POITRINAIRE, Phthisicus.

POITRINE, Thorax.

POIVRE D'EAU. *Polygonum hydropiper* —

p. de Guinée, *Capsicum annuum* — p. d'Inde, *Capsicum annuum* — p. de Jamaïque, see *Myrtusimenta* — p. Long, *Piper longum* — p. de Moine, *Vitex* — p. Noir, *Piper nigrum* — p. Petit, *Vitex* — p. à Queue, *Piper cubeba* — p. Sauvage, *Vitex*.

POIX BLANCHE, see *Pinus abies* — p. de Bourgogne, see *Pinus abies* — p. Grasse, see *Pinus abies* — p. Jaune, see *Pinus abies* — p. Liqvide, see *Pinus sylvestris* — p. Minérale, *Pissasphaltum* — p. Minérale scorifiée, *Asphaltum* — p. Noale, see *Pinus sylvestris* — p. Noire, see *Pinus sylvestris*.

POKE, *Phytolacca decandra* — p. Indian, *Veratrum viride* — p. Stink, *Dracontium foetidum* — p. Weed, *Phytolacca decandra*, *Veratrum viride*.

POLANIS'IA GRAVEOLENS, *Clammy weed*, *Stinkweed*, *Wormweed*, *Clammy Mustard*, *False mustard*, from *πολις*, 'many,' and *αινεος*, 'unequal,' in reference to its stamens. An indigenous plant, *Ord. Capparidacææ*, *Ses. Syst. Dodecandria Monogynia*, which blossoms in summer, and is found in every part of North America. To the whole plant have been ascribed anthelmintic properties.

POLAR PLANT, *Silphium laciniatum*.

POLECAT WEED, *Dracontium foetidum*.

POLEMONIUM REPTANS, *False Jacob's Ladder*, *Greek Vale'rian*, *Sweat-root*. *Order*, *Polemoniaceæ*; indigenous, flowering in May; has been used as a diaphoretic and expectorant.

POLENTA, Alphon.

POLES, Chalazæ.

POLEY MOUNTAIN, OF CANDY, *Teucrium Creticum* — p. Mountain of Montpellier, *Teucrium capitatum*.

POLIA'TER, from *πολις*, 'a town,' and *ιατρος*, 'a physician.' A physician exercising his profession in a town, by the appointment of the government. — Castelli.

POLICE', MED'ICAL, *Politi'a med'ica*, *Politiobiotherapi'a*, *Medici'na polit'ica*, *Ordo medic'inalis*, *Polit'ical medicine*, (F.) *Police médicale ou sanitaire*. Hygienic rules established, or which ought to be established, for the prevention of disease, and to contribute to the progress of medicine. The salubrity of districts, especially during the existence of spreading disease, is greatly ensured by an efficient medical police.

POLICLIN'ICA, from *πολις*, 'a town,' and *κλινη*, 'a bed.' Treatment of diseases in a town.

POLIOSIS, *Tricho'sis polio'sis*, *Spilo'sis poli-*

o'sis, *Pol'ioties*, *Ami't'ies*, from *πολιος*, 'hoary, gray.' Hair prematurely gray or hoary. The state of becoming gray, *gray-haired*, *Poliot'richos*.

POLIOTES, Poliosis.

POLIOTRICHOS, see Poliosis.

POLITIA MEDICA, Police, medical.

POLITOBIOThERAPIA, Police, medical.

POLIUM CAPITATUM, Teucrium capitatum — p. Creticum, Teucrium Creticum — p. Montanum, Teucrium capitatum.

POLL, Head.

POLLEN, (L.) *Pollen*, *Pollis*, *παλη*, 'dust.' The fecundating dust contained in the anthers of flowers, which is dispersed on the stigma. It is conveyed to the ovary through the stigma by the *Pollen tube*, *Tubus pollin'icis*, (F.) *Boyaupollinique*.

POLLEX, genitive *Pol'licis*, *Ant'icheir*, *Prom'onus*, *Dig'itus primus seu magnus*, *Allus pollex*, *Thumb*, (Prov.) *Thame*, *Tom-toe*, from *pollere*, 'to have much strength.' (F.) *Pouce*. The first of the fingers in size and strength. It is asserted that the term *poltroon* is derived from *poll'ice truncat'o*; — the Romans who were desirous of being exempted from military service being accustomed to cut off the thumb, and thus render themselves unfit to handle arms.

POLLEX, see *Digitus*.

POLLEX PEDIS, *Hollux*, *Hallux*. The great toe.

POLLINCTOR, *Pollincter*, (L.) In antiquity, an anointer, dresser, or embalmer of the dead.

POLLINCTURA, Embalming..

POLLIS, Pollen.

POLLOCK, see *Oleum jecoris aselli*.

POLLOD'IC, *Pollo'd'icis*; from *πολvs*, 'many,' and *odos*, 'a way.' An epithet proposed by Dr. Marshall Hall for a course of nervous action proceeding from one point, in many directions, to every other.

POLLUM, **LITTLE**, *Polygala paucifolia*.

POLLUTION, *Pollut'io*, from *polluo*, *pollutum*, 'to profane.' The excretion of the seminal liquor or sperm, or the production or orgasm in male or female, at other times than during coition. When occasioned by a voluntary act, it is called simply *Pollution* or *Masturbation*, (F.) *Souillure*; when excited during sleep, by lascivious dreams, it takes the name *Nocturnal pollution*, *Exoneiro'sis*, *Oneirog'mos*, *Oneirog'onos*, *Oneirogonorrhoe'a*, *Exoneirog'mus*, *Spermatoclem'ma*, *Spermatoclepsi'sis*, *Spermatolepsi'sis*, *Spermatolip'is*, *Gonorrhoea dormient'ium seu oneirog'onos seu vera seu libidinosa*, *Profluvium sem'inis*, *Spermatorrhoe'a*, *Paronit'ia salax*, *Night pollution*, (F.) *Pollution involontaire*.

POLLUTION INVOLONTAIRE, *Pollution* — p. Nocturnal, *Pollution* — p. Self, *Masturbation* — p. Voluntary, *Masturbation*.

POLONYCH'IA, *Polyonychi'a*, from *πολvs*, 'many,' and *ονυξ*, *ονυχος*, 'a nail.' The condition of having supernumerary nails, — a rare congenital monstrosity.

POLT-FOOT, see *Kyllosis*.

POLY, *Polys*, *πολvs*, 'many, much, full.' Hence:

POLYACTIUM TRISTE, *Pelargonium triste*.

POLYEMIA, *Plethora*.

POLYEMIC, *Plethoric*.

POLYEMICUS, *Plethoric*.

POLYANHEMIA, *Anæmia*.

POLYANHEMIE, *Anæmia*.

POLYANTHES TUBEROSA, *Amaruacachu*. Ord. Liliaceæ. The root is cut into slips and laid upon the bites of serpents by the Peruvian Indians.

POLYBLEN'NIA, from *poly*, and *βλεννα*, 'mucus.' Excessive secretion of mucus.

POLYBREPHIA, *Pregnancy*, complex.

POLYCARP'Æ, from *poly*, and *καρπος*, 'fruit.' A division of cutaneous diseases of the family *Eczematosen*, of Fuchs, including *Psoriasis*, *Lichen*, *Herpes*, and *Impetigo*.

POLY'CERAS, from *poly*, and *κερας*, 'horn.' A monstrous animal having many horns. — Gurlt.

POLYCHOL'IA, *Perichol'ia*, *Hyperchol'ia*, from *poly*, and *χολη*, 'bile.' Superabundance of bile; such as is supposed to exist in bilious fevers.

POLYCHREST'US, from *poly*, and *χρηστος*, 'useful.' Having many virtues. An ancient name for certain medicines considered to be useful in many diseases.

POLYCHRONIUS, *Chronic*.

POLYCHY'LIA, from *poly*, and *χυλος*, 'juice.' *Polychy'mia*. Fulness of juices.

POLYCHYMIA, *Polychylia* — p. *Sanguinea*, *Plethora*.

POLYCLIN'ICA, from *poly*, and *κλινη*, 'bed.' Provided with many beds; — as an hospital.

POLYCOR'IA, from *poly*, and *κορη*, 'the pupil.' The state in which there is more than one pupil. — Von Ammon.

POLYCRINOSSES, see *Profluvium*.

POLYCYE'SIA, *Polycy'e'sis*, from *poly*, and *κυησις*, 'pregnancy.' Frequent or quickly recurring pregnancies: also, compound pregnancy.

POLYCYTHÆMIA, from *poly*, *κυτος*, 'a cell,' and *αίμα*, 'blood.' A condition of the blood in which there is an increase of the red corpuscles.

POLYDACTYLUS, *Perissodactylus*, from *poly*, and *δακτυλος*, 'a finger.' One who has a supernumerary finger or fingers, or toes.

POLYDIP'SIA, from *poly*, and *διψη*, 'thirst;' *Sitis morbo'sa seu inten'sa*, *Polypos'sia*, *Dipso'sis*, *avens*, *Excessive thirst*, (F.) *Soif excessive*. A constant desire for drinking, with a sense of dryness in the mouth and throat. It is mostly symptomatic of fever, inflammation, dropsy, excessive discharges, &c., and is so much a part of diabetes, that it has been called *Polydipsia*.

POLYG'ALA, from *poly*, and *γαλα*, 'milk;' *P. vulgaris*, *Common Milkwort*. Ord. *Polygalaceæ*. *Sez. Syst.* *Diadelphica Octandria*. The leaves are bitter, and are used in the same cases as the *Polygala amara*.

POLYGALA AMA'RA, *P. amarel'la seu vulgaris*, *Bitter Milkwort*. A powerful bitter; and, consequently, tonic. Its reputed virtues are balsamic, demulcent, and corroborant. The *Polygala rubell'a*, or *Bitter Polygala* of the United States, has the same virtues.

POLYGALA AMARELLA, *P. amara* — p. Bitter, see *P. amara* — p. *Grandiflora*, *P. senega*.

POLY'ALA PAUCIFOL'IA, *Triclisper'ma grandiflora*, *Dwarf Milkwort*, *Little Pollom*, *Evergreen Snakeroot*, (F.) *Polygale nuine*, possesses the same properties as *P. Senega*.

POLY'ALA SEN'EGA, *P. Virginica'n'a seu grandiflora*, *Sen'eka*, *Rattlesnake Milkwort*, *Senega* or *Rattlesnake root*, (F.) *Polygalle de Virginie ou de Seneka*. The bark is the active part of the root, *Senega* (Ph. U. S.). Its taste is at first acrid, and afterwards very hot and pungent. It was once esteemed to be specific against the poison of the rattlesnake and in pleurisy, pneumonia, &c., after the inflammatory action had been got under. It has been given also in humoral asthma, chronic rheumatism, dropsy, croup, amenorrhœa, &c. According to Peschier, it contains three particular principles, *Polyg'atine* or *Sen'egine*, *Isol'uine*, and *Polyg'alic Acid*. Dose, gr. 30 to 40 of the powder.

POLYGALA SERPENTA'RIA. A shrub of *Caffra*, the root of which is considered by the natives as a sure antidote against the bites of venomous serpents.

POLYGALA VIRGINIANA, *P. senega* — *p. Vulgaris* *P. amara*.

POLYGALACTIA, *Polygal'ia*, from *poly*, and *γάλα*, *yalaktos*, 'milk.' Superabundant secretion of milk; galactia.

POLYGALE NAINE, *Polygala* paucifolia — *p. de Virginie*, *Polygala Virginiana*.

POLYGALIA, *Polygalactia*.

POLYCALINE, see *Polygala senega*.

POLYGONATUM, *Convallaria polygonatum* — *p. Anceps*, *Convallaria polygonatum* — *p. Caniculatum*, *Convallaria canaliculata* — *p. Latifolium*, *Convallaria canaliculata* — *p. Multiflorum*, *Convallaria multiflora* — *p. Pubescens*, *Convallaria pubescens* — *p. Uniformum*, *Convallaria polygonatum* — *p. Vulgare*, *Convallaria polygonatum*.

POLYGNONUM, *P. aviculare*, from *poly*, and *γων*, 'a knee or joint.' *P. centinod'ium* seu *erectum*, *Centinodia*, *Centummodia*, *Sanguinalis* *Corrigiola*, *Sanguinalis*, *Knotgrass*, *Knotweed*, *Bindweed*, *Doorweed*, (F.) *Renouée*, *Trainasse*, *Centinode*. *Ord. Polygonaceæ*. *Sex. Syst. Octandria Trigynia*. This grass is said to be a feeble astringent.

POLYGONUM ACRE, *P. hydropiper*.

POLYGONUM AMPHIBIUM, *P. coccineum*, *Water Persicaria*, of Europe and this country. The roots have been used like sarsaparilla, especially in diseases of the skin.

POLYGONUM BISTORTA, *P. bistortoides*, *Bistorta*, *Colubrina*, *Officinal Bistort*, *Great Bistort* or *Snakeweed*, (F.) *Bistorte*, *Couleuvre*, *Serpentaire rouge*. The root is a powerful astringent and tonic; and as such, has been used in internal hemorrhage, atonic diarrhoea, ague, &c. *Dose*, gr. xv to ʒi.

POLYGONUM BISTORTOIDES, *P. bistorta* — *p. Centinodium*. *P. aviculare*.

POLYGONUM COCCINEUM, *P. amphibium*.

POLYGONUM DIVARICATUM; *Eastern Buckwheat plant*. The roots, reduced to a coarse meal, are said to be much used as an article of diet by the Siberians.

POLYGONUM ERECTUM, *P. aviculare*.

POLYGONUM FAGOPYRUM, *Fagopyrum*, *F. esculentum*, *Triticum Faginum*, *Buckwheat*, (F.) *Sarrasin*, *Blé noir*. The grain of this plant constitutes a principal article of food with the inhabitants of Russia, Germany, and Switzerland. It is much used in the United States.

POLYGONUM HYDROPIPER, *P. acre* seu *hydropiperoides* (Pursh) seu *punctatum*, *Hydropiper*, *Persicaria wrens*, *Mercurius terrestris*, *Poor Man's Pepper*, *Biting Arse-smart*, *Smartweed*, *Lakeweed*, *Water-pepper*, (F.) *Poire d'eau*, *Persicair poivrée* ou *brûlante* ou *âcre*, *Piment d'eau*, *Cuarge*, *Renouée âcre*. The leaves of this plant have an acrid, burning taste, and resemble those of the arum. They have been recommended as possessing antiseptic, aperient, and diuretic virtues. The fresh leaves have been applied externally as a stimulating cataplasin.

POLYGONUM HYDROPIPEROIDES, *P. Hydropiper*, *P. punctatum*. *Polygonum hydropiperoides* of Michaux, *mild waterpiper*; indigenous; has no acrid properties.

POLYGONUM PERSICARIA, *Persicaria*, *P. minor* seu *mitis*, *Plumbago*, *Lady's Thumb*, *Heart's ease*, (Prov.) *Redshanks*, (F.) *Persicair douce*. This plant is said to possess vulnerary and antiseptic qualities. It has been given in wine to arrest gangrene.

POLYGONUM PUNCTATUM, *P. hydropiperoides*, *Water pepper* of America, *Smartweed*, possesses similar properties to *Polygonum hydropiper*, and is occasionally used internally in gravel, and externally in chronic ulcers. A fluidrachm of the saturated tincture has been extolled in amenorrhœa. See *Polygonum hydropiper*.

POLYGNONUM VIRGINIANUM, *Wand-spiked Persicaria*, *Virgin'ia Bistort*; indigenous; is astringent, diuretic, and tonic.

POLYGROMA, *Staphyloma*.

POLYHIDRIA, *Polyhidria*, *Polyhidrosis*, from *poly*, and *ἵδρω*, 'sweat.' Excessive sweating.

POLYHYDRIA, *Hydrops*.

POLYHYPERHÆMIA, *Plethora*.

POLYIDRIA, *Polyhidria*.

POLYIDROSIS, *Polyhidria*.

POLYLYMPIA, *Anasarca*.

POLYMASTHUS, *Pleoma'sthus*, from *poly*, and *μαστος*, 'breast.' A monster having several breasts. — Gurlt. The condition has been called *Polyma'zia*, *Pleoma'zia*, and *Pleoma'zia*.

POLYMAZIA, see *Polymasthus*.

POLYMELEIAN, (F.) *Polymélien*, from *poly*, and *μελος*, 'a limb.' An epithet given to monsters characterized by the insertion, in a well formed subject, of one or more accessory limbs, accompanied, at times, by the rudiments of certain other parts. — I. G. St. Hilaire.

POLYMERIA, *Polymerismus* — *p. Promiscua* *translativa*, *Transposition of the viscera*.

POLYMERISMUS, *Polymer'ia*, from *poly*, and *μερος*, 'a part.' A monstrosity, which consists in the existence of supernumerary organs; as a sixth finger of the hand.

POLYNEURON, *Plantago*.

POLYONYCHIA, *Polonychia*.

POLYOP'SIA, *Polyopy*, from *poly*, and *οψis*, 'sight.' Vision is so called, when multiple or manifold.

POLYOPY, *Polyopsia*.

POLYOREX'IA, from *poly*, and *ορεξις*, 'appetite.' Excessive appetite or hunger. A genus in the *Gastroses* of M. Alibert.

POLYPE'RIA, from *poly*, and *πηρος*, 'mutilated.' Congenital misconstruction of various parts or organs. — Good.

POLYPHA'GIA, *Multivoran'tia*, from *poly*, and *φαγω*, 'I eat.' Voracity. Faculty of eating or subsisting on many kinds of food. See *Boulimia*.

POLYPHAGOUS, *Polyph'agus*, *Multiv'orous*, (F.) *Polyphage*. Relating or appertaining to polyphagia.

POLYPHARMACIA, *Polypharm'acy*; from *poly*, and *φάρμακον*, 'a medicine.' The prescribing of too many medicines. A prescription, consisting of a number of medicines; hence the name *Polypharm'acus*, (F.) *Polypharmaque*, given to one who is in the habit of prescribing a number of medicines, and whose prescriptions are loaded with ingredients. The term is taken in bad part.

POLYPHARMAQUE, see *Polypharmacia*.

POLYPHY'SIA, *Flatulence*.

POLYPIFORM CONCRETIONS, see *Polypus*.

POLYPIONIA, see *Polysarcia* — *p. Infantum*, see *Polysarcia*.

POLYPIOSIS, see *Polysarcia*.

POLYPLASTIC, *Polyplast'icus*, from *poly*, and *πλασσω*, 'I form.' That which has many shapes.

POLYPLASTIC ELEMENT, in histology, is one which does not retain its primary shape; but undergoes transformation into others. — Gerber.

POLYPODITES. Wine impregnated with polypody.

POLYPO'DIUM, *P. Vulgarē* seu *Virginianum* seu *pinnatifidum*, *Filix'gla dulcis*, *Common Polypody*, *Fern root*, *Rock Brake*, *Brake root*, (F.) *Fougère douce*, *Ord. Filices*. From *poly*, and *πους*, *podos*, 'a foot,' owing to its many roots. This fern grows in the clefts of rocks, walls, &c., in both hemispheres. It was formerly regarded as

a cholagogue, and was given in head diseases. It is not used now.

POLYPODIUM ADIANTIFORME, see *Calagualæ radix* — p. *Ammifolium*, see *Calagualæ radix* — p. *Argenteum*, see *Calagualæ radix* — p. *Cetaræcinum*, *P. incanum* — p. *Calaguala*, *Calagualæ radix* — p. *Coriaceum*, see *Calagualæ radix* — p. *Dentatum*, *Asplenium filix femina* — p. *Filix femina*, *Asplenium filix femina*.

POLYPODIUM FILIX MAS, *Aspidium* seu *Athyrium* seu *Nephrodium* seu *Polystichum Filix Mas*, *A. depauperatum* seu *erectum*, *Filix pinnata*, *Pteris*, *Blechnon*, *Polyptrichon*, *Blancnon* *Oribasit*, *Lonicit*, *Filicula*, *Filix Mas*, *F. non ramo'sa dentata*, *Nephrodium crenatum*, *Polystichum callipteris*, *Male Polypody* or *Fern.* (*F.*) *Fougère mâle*. The rhizoma of this plant, *Filix Mas*, Ph. U. S., has been much extolled as an anthelmintic, and in cases of tænia. It was the basis of a once celebrated remedy—*Madame Nonffer's*—but, perhaps, more of the efficacy was to be attributed to the active purgatives by which it was followed than to the fern. Dose, ʒij to ʒiij of the powdered solid part of the root; followed in the morning by a strong cathartic of camboe or jalap. An ethereal extract has been proposed by Dr. Peschier, of Geneva.

POLYPODIUM INCANUM, *P. cetaræcinum*, *Acrostichum polypodioides*, *Polypody*, *Rock Brake*; indigenous; has been considered demulcent, cathartic and anthelmintic.

POLYPODIUM INCISUM, *Asplenium filix femina* — p. *Molle*, *Asplenium filix femina* — p. *Pennatifidum*, *P. vulgare* — p. *Politum*, see *Calagualæ radix* — p. *Trifidum*, *Asplenium filix femina* — p. *Virginianum*, *P. vulgare* — p. *Vulgare*, *Polypodium*.

POLYPODY, *Polypodium incanum* — p. *Common*, *Polypodium vulgare* — p. *Male*, *Polypodium filix mas*.

POLYPOID, *Polypoides*; from *polypus*, and *oides*, 'form.' Having the shape of, or resembling, a polypus.

POLYPORUS IGNIARIUS, *Boletus ignarius* — p. *Officialis*, *Boletus laricis*.

POLYPO'SIA, *πολυποσία*, from *poly*, and *-osis*, 'drink.' Great thirst—desire for drink. *Polydipsia*.

POLYPO'TOME, *Polypotomus*, *Polypus knife*, from *polypus*, and *τεμνειν*, 'to cut.' An instrument for the removal of polypus by excision.

POLYPOUS, *Polypous*: same etymon as the next. Relating, or belonging to, or resembling polypus.

POLYPUS, *Pseudometamorphosis polyposa*, from *πολύς*, 'many,' and *πους*, *πόδος*, 'a foot.' A name given to tumours, which occur in mucous membranes especially; and which have been compared to certain zoophytes. Polypi may form on every mucous membrane. They are most commonly observed in the nasal fossæ, pharynx, and uterus; more rarely in the stomach, intestines, bronchial tubes, bladder, and meatus auditorius externus. They vary much in size, number, mode of adhesion, and intimate nature. Accordingly, they have received various appellations; *mucous*, *soft*, *vesicular*, when their substance has been soft, spongy, vesicular, and, as it were, gorged with fluid. Others have been called *hard*; and these have been distinguished into *fibrous* or *fleshy*, and into *scirrhous* or *cancerous*. *Fibrous* polypi, *Polypii fibrosi*, *Inopolypii*, are of a dense, compact texture, and whitish colour. They contain few vessels, and do not degenerate into cancer. The *scirrhous* or *cancerinomatous* are true cancerous tumours, painful and bleeding. They present all the pathological changes which belong to cancerous affections, and the prognosis is of course unfavourable.

The mode of treating polypi must differ according to their character. The means used to destroy them have been, 1. To subject them to the action of certain astringent powders or solutions, to obtain their resolution. 2. *Cauterization* or the application of the actual cautery or caustics. 3. *Excision* or ablation with a cutting instrument. 4. *Extirpation* with the fingers or with the polypus forceps. 5. A *ligature*, put round them so as to prevent their nutrition. A seton has, also, been sometimes used to gradually destroy the pedicle.

The term **POLYPY**, *Pseudopolypii*, *Carditis polyposa*, *Polypiform Concretions*, (*F.*) *Concretions polypeuses* ou *polypiformes* ou *sanguines* du Cœur, *Hémocardioplasties*, (*Piorry*), has likewise been applied to collections of blood—fibrinous concretions—found in the heart or great vessels after dissolution. These were once, erroneously, regarded as morbid. The presence of these concretions may be suspected, when in the course of an acute or chronic affection of the heart, or at the close of chronic diseases in general—especially those of the lungs—the heart's action becomes suddenly tumultuous, obscure, and accompanied with a soft bellows' murmur, whilst the general symptoms indicate the effects of obstructed circulation. The right cavities are most frequently affected.

POLYPUS BRONCHIALIS, *Bronchlemmitis*, *Bronchitis membranacea* seu *exsudativa* seu *crouposa*, *B. plastic*, *Bronchial Croup*, *Chronic Croup* [?], *Pseudo-membranous Bronchitis*, (*F.*) *Bronchitis pseudo-membranosa*. A membraniform secretion in the bronchial tubes of a diphtheritic character, having the shape, when unexpecterated, of the bronchia. It is the result of inflammation of the lining membrane of those tubes.

POLYPUS CARNOSUS, *Sarcoma* — p. *Fibrosus*, see *Polypus* — p. *Uteri*, *Metropolypus*.

POLYRRHIZON, *Polypodium filix mas*.

POLYRRHŒA, *Profluvium*.

POLYSAR'CIA, *Polysarcosis*, from *poly*, and *σαρξ*, *σαρκος*, 'flesh.' *Carnositas*, *Polysomatia*, *Adeliparia*, *Corpulentia*, *Stentites*. Excessive corpulency. Authors have distinguished two varieties of this:—**POLYSARCIA CARNO'SA**, *Corpulency*, and **P. ADIPO'SA**, *Polypionia*, *Polypionia*, *Pinelonia nimia*, *Pachymnia*, *Pachymnia*, *Obesitas*, *Hyperpimelia*, *Liparia*, *Adipositas*, *Polysarcia faginata*, *Emphysematum polysarcia*, *Barysomia*, *Barysomatia*, *Liposis*, *Lipomatosis*, *Liposoma*, *Pinguitedo*, *Pinguitedude*, *Obesity*, (*F.*) *Obésité*, *Embonpoint excessif*. A severe regimen and exercise are the best preventives. Children are occasionally met with of unusual fatness, in spite of every care. This morbid disposition has been termed *Polypionia infantum*, and *Chlorosis gigantea*.

POLYSARCIA VISCERUM, *Physeina adiposa*.

POLYSARCOSIS, *Polysarcia*.

POLYSC'EL'IA, from *poly*, and *σκελος*, 'a leg.' A monstrosity, characterized by the presence of one or more supernumerary legs.

POLYSIA'LIA, from *poly*, and *σάλιν*, 'saliva.' Excessive discharge of saliva.

POLYSOMATIA, *Polysarcia*.

POLYSO'MIA, from *poly*, and *σῶμα*, 'body.' A monstrosity characterized by the existence of more bodies than one.

POLYSPAS'TON, from *poly*, and *σπασω*, 'I draw.' A powerful instrument for reducing luxations. — *Gorræus*.

POLYSPERM'IA, from *poly*, and *σπέρμα*, 'sperm.' Richness or fulness of sperm.

POLYSTICUM CALLIPTERIS, *Polypodium filix mas* — p. *Filix Mas*, *Polypodium filix mas*.

POLYSTOMA PINGUICOLA, see *Worms* — p. *Venarum*, see *Worms*.

POLYTRIC, *Asplenium trichomanoides*.

POLYTRICHIA, see *Polytrichus*.

POLYTRICHUM, from *poly*, and *τρίχος*, 'hair,' *P. vulgare*. *Polytrichum*, *Adiantum vulgareum*, *Golden Maiden-hair*, (F.) *Perce-mousse*. *Ord. Musci*. It is possessed of astringent properties, and was formerly given in diseases of the lungs, and in calculous affections.

POLYTRICHUM COMMUNE, *Asplenium trichomanoides*.

POLYTRICHUM JUNIPERINUM, *Hairecap moss*, *Robbin's Rye*, common in New England, is said to possess powerful diuretic properties.

POLYTRICHUS; same etymon as *Polytrichum*. Very hairy. One that has much hair. The condition is termed *Polytrichia*; although this term is, at times, employed for the growth of hair on parts not usually having it.

POLYTROPHIA, from *poly*, and *τροφή*, 'I nourish.' Abundant or excessive nutrition.

POLYURESIS, Diabetes.

POLYURIA, Diabetes.

POLYURICUS, from *poly*, and *ουρον*, 'urine.' Sauvages calls—*Ischuria polyurica*—a kind of ischuria, caused by accumulation of urine in the bladder.

POMA, Drink—*p. Chinensis*, see *Citrus aurantium*.

POMACEUM, Cider.

POMATUM, *Pommatum*, from *poma*, 'apples,' which formerly entered into its composition. (F.) *Pomnade*, *Liparolé*, *Stéarolé*. A pharmaceutical preparation, employed only externally, which differs but little from an ointment, except in having generally less consistence. It is usually scented and coloured to render it more agreeable. Apples formerly entered into the composition of the *pomatum*; hence its name.

Common Pomatum is made of lard, 12lbs; *suet*, 4lbs; *essence*, 3viij.

POMATUM AD LABIA DEMULCENDA. Cerate for the lips—*p. Epispasticum flavum*, *Unguentum lyttæ medicatum aliud*—*p. Epispasticum viride*, &c., *Unguentum lyttæ medicatum*—*p. for the Lips*, Cerate for the lips—*p. Nitricum seu oxygenatum*, *Unguentum acidi nitrosi*—*p. Saturni*, *Unguentum plumbi superacuatiss.*

POMBALEA IPECACUANHA, see *Ipecacuanha*.

POMEGRANATE, *Punica granatum*.

POMILIUS, see *Nanus*.

POMMADE, *Pomatum*—*p. d'Alion*, *Unguentum acidi nitrosi*—*p. Antiporique de Jasser*, *Unguentum sulphuris compositum*—*p. d'Antenrieth*, *Unguentum antimonii tartarizati*—*p. de Cyrrillo*, *Unguentum muriatis hydrargyri oxygenato medicatum*—*p. Épispastique jaune*, *Unguentum lyttæ medicatum aliud*—*p. Épispastique verte*, *Unguentum lyttæ medicatum*—*p. de Garou*, *Unguentum epispasticum de daphne gnidio*.

POMMADE DE GONDRET, *P. ammoniacale de Gondret* (F.), *Gondret's Counterirritant*, *Gondret's ammoniacal Caustic*. (*Liq. ammon. p. ij; Sevi. p. j; Ol. amygdal. p. j; M.*) Used to excite a speedy revulsion in cases of chronic affections of the brain, incipient cataract, amaurosis, &c., as well as to cauterize the integuments deeply.

The above formula, is that of the French Codex, but, according to M. Trousseau, it is much too hard for use, unless the weather is warm. He proposes two formulae, one for summer, the other for winter. In the former he directs three parts of lard, one of *suet*, and four of *liquor ammoniac*; in the latter, equal parts of lard and *liquor ammoniac*.

POMMADE POUR LES LÈVRES, Cerate for the lips—*p. Ophthalmique* (Regent's.) *Unguentum ophthalmicum*—*p. Oxygénée*, *Unguentum*

acidi nitrosi—*p. Stibiée*, *Unguentum antimonii tartarizati*.

POMME, see *Pyrus malus*—*p. d'Adam*, *Pomum Adami*—*p. d'Amour*, *Solanum lycopersicum*—*p. Épineuse*, *Datura stramonium*—*p. de Merveille*, *Momordica balsamica*—*p. Moussueuse*, *Bedegar*—*p. de Terre*, *Solanum tuberosum*.

POMMES HÉMORRHOÏDALES, see *Viscum album*.

POMMETTE, see *Malus os*.

POMMIER, *Pyrus malus*.

POMPE POUR L'ESTOMAC, *Stomach pump*.

POMPHOLYGE'RON, *πομφολυγεον*, from *πομφολυγος*, 'to raise blisters.' The name of a plaster employed by the ancients.—Castelli.

POMPHOLYX, from *πομφος*, 'a bladder.' A small vesicle on the skin. An eruption of *bullæ* or *blebs*, appearing without any inflammation around them, and without fever. It is the *Eczphylia Pompholyx* of Good. See *Pemphigus*.

POMPHOLYX, *Tutia*, *Zinci oxydum*.

POMPHUS, *Wheat*.

POMPION, *Cucurbita pepo*.

POMUM ADA'MI, *Eminentia thyreoidea*, (F.) *Pomme d'Adam*, *Adam's Apple*. The projection formed by the thyroid cartilage in the neck—of males particularly. It is so called from a notion that it was caused by the apple sticking in the throat of our first parent.

POMUM AMORIS, *Testicle*—*p. Hierosolymitanum*, *Momordica balsamina*—*p. Melongena*, see *Solanum melongena*—*p. Mirabile*, *Momordica Balsamina*—*p. Spinosum*, *Datura stramonium*.

PONCTION, *Puncturing*.

POND SHOVEL, *Unisema deltifolia*.

PONDERA ET MENSURÆ, *Weights and measures*.

PONDUS, *Pudibilia*—*p. Pharmaceuticum*, see *Weights and measures*.

PONGITIE, *Pungent*.

PONS CEREBELLI, *Pons Varolii*—*p. Hepatis*, see *Lobulus anonymus*—*p. Sylvii*, *Quadrigemina Tubercula*—*p. Tarini*, *Tarini pons*.

PONS VAROLII, *P. Cerebelli*, *Corpus annulære*, *Processus annularis*, *Prominentia annularis cerebri*, *Eminentia annularis*, *Nodus encephali seu Cerebri*, *Protuberantia basilaris*, *Protuberantia annularis Willisii*, *Varoli's bridge*, *Cerebrum protuberance*, (F.) *Protubérance cérébrale ou annulaire*, *Méencéphale*, *Mésocéphale*, *Pont de Varole*, *Annular protuberance*. An eminence at the upper part of the medulla oblongata, first described by Varolius. It is formed by the union of the crura cerebri and crura cerebelli.

PONS ZYGOMATICUS, *Zygomatic arch*.

PONT DE TARIN, *Tarini pons*—*p. de Varole*, *Pons Varolii*.

PONTE, see *Parturition*—*p. Périodique*, see *Parturition*.

PONTEDERIA CORDATA, *Unisema deltifolia*. **PONTICUM ABSINTHIUM**, *Artemisia Puntica*.

POOK, *Abdomen*.

POPLAR, AMERICAN, *Liriodendron*—*p. Balm of Gilead*, *Populus Candicans*—*p. Black*, *Populus*—*p. Tree*, *Liriodendron*—*p. Tulip-bearing*, *Liriodendron*.

POPLES, *Gare'tum*, *Garretum*, *Ignys*, *Ig'nyē*, *Suffrago*, *Fossa genu*, *Fossa poplitea*, the *Ham*, *Hough*, *Hoff*, *Hock*, (F.) *Jurret*, from *post*, and *plico*, 'I fold.' The posterior part of the knee joint. The ham forms what is called the *popliteal region* or *space*.

POPLESY, *Apoplexy*.

POPLITEUS, *Popliteal muscle*.

POPLITÉ, *Popliteus muscle*.

POPLITEAD, see *Popliteal aspect*.

POPLITE'AL, *Poplitæ'us*, *Poplit'ic*, (F.) *Poplité*, from *poples*, 'the ham.' That which relates to the ham. A name given to various parts.

POPLITEAL ARTERY—*Popliteal portion of the crural*,—(Ch.), is really a continuation of the femoral, which changes its name on passing through the adductor magnus muscle. It descends a little obliquely outwards into the hollow of the ham, and extends from the commencement of the lower third of the thigh to the end of the upper quarter of the leg, where it terminates by dividing into the *posterior tibial and peroneal arteries*. In the ham, it gives off the three *superior articular arteries*; and, at the top of the leg, the arteries of the gemini muscles or *gemini branches*, (F.) *branches jumelles*, the two *inferior articular arteries*, and the *tibialis antica*.

POPLITEAL ASPECT. An aspect towards the side on which the *poples* or ham is situated. (Barclay.) *Poplite'ad* is used by the same writer adverbially, and signifies 'towards the popliteal aspect.'

POPLITEAL FOSSA, *Poples*.

POPLITE'AL GLANDS, *Glan'dulæ poplitæ'æ*. Lymphatic glands, four in number, seated in the popliteal region or space.

POPLITEAL MUSCLE, *Poplitæ'us Muscle*, *Subpoplitæus*, (F.) *Muscle poplité*, *Muscle Jarretier*, *Fémoro-poplitæ-tibial*, (Ch.), is seated in the popliteal region. It is triangular, long, flat; and is inserted, on the one hand, by means of a strong tendon, into the tuberosity at the outer condyle of the femur, terminating below at the upper part of the posterior surface of the tibia. This muscle bends the leg on the thigh, and, reciprocally, the thigh on the leg. When the leg is bent, it causes it to turn, in rotation, inwards.

POPLITEAL NERVES. They result from the bifurcation of the sciatic, and are two in number, viz. 1. The *external popliteal, Peroneal nerve*, *Branche péronière*, (Ch.) It is smaller than the internal; turns to the outside of the fibula at the upper part of the leg, and divides into the *musculo-cutaneous branch, nerf pré-tibio-digital*, (Ch.) and the *anterior tibial, nerf pré-tibio-susphalangé-taire*, (Ch.) 2. The *internal popliteal nerve*, *Branche tibiale* (Ch.), which is larger than the other, and seems to be the real continuation of the sciatic nerve. It descends vertically into the hollow of the ham; and passes beneath the aponeurotic arch of the solaris muscle, beyond which it takes the name *tibial nerve*. It furnishes the *external saphena nerve*, and a number of branches, that are distributed to the posterior muscles of the leg. It then passes to the sole of the foot to be divided into two branches, called *Plantar nerves*.

POPLITEAL REGION, *Poples*—p. Space, *Poples*.

POPLITEAL VEIN has the same arrangement as the artery behind which it is situated.

POPLITIC, *Popliteal*.

POPONAX, see *Pastinaca opoponax*.

POPPY, *CELANDINE*, *Stylophorum diphyl-lum*—p. Corn, red, *Papaver rhœas*.

POPPY HEADS, see *Papaver*.

POPPY HORN, *Stylophorum diphyl-lum*.

POPPY JUICE, *Meconium*—p. Pill, *Opium*.

POPPY, PRICKLY, *Argemone Mexicana*—p. Spotting, *Cucurbalus Behen*—p. Thorn, *Argemone Mexicana*—p. White, *Papaver*—p. White, capsule, see *Papaver*.

POPULUS, *P. nigra*, *Ægei'ros*, *Achero'is*, *Black poplar*, (F.) *Peuplier*, *P. noir* ou *franc*. The young buds, *oculi* or rudiments of the leaves, which appear in the beginning of spring, were formerly used in an official ointment—the *unguentum popule'um*. It was, also, recommended in chronic affections of the lungs.

The bark of *Populus Tremuloi'des*, American

aspen, of *P. TREM'ULA* seu *pen'dula*, *European aspen*, and of *POPULUS CAN'DICANS*, *Balm of Gilead Poplar*, is antiperiodic, and furnishes *salicin* and *populin*, in which the properties probably reside.

POPULUS BALSAMIFERA, see *Fagara octandra*—p. *Candicans*, see *Populus*—p. *Pendula*. *P. tremula*—p. *Tacamahaca*, see *Fagara octandra*—p. *Tremula*, see *Populus*—p. *Tremuloides*, see *Populus*.

PORCELAINE, *Essera*, *Urticaria*.

PORCELET, *Hyoseyamus*.

PORCELIA TRILOBA, see *Carica papaya*.

PORCELLIO, *Oniscus*.

PORCELLUS, *Oniscus*.

PORCULUS, *Oniscus*.

PORCUPINE DISEASE, *Ichthyosis*.

PORCUPINE MEN, *Echinoder'mi*. Persons have been so called in whom the cuticle is produced in the form of pointed prolongations, as if it had been moulded on the papillæ, like the shorter and blunter quills of the porcupine.

PORCUS, *Uvula*.

PORDE, *Crepitation*.

PORE, *Porus*, *In'terstice*, *poros*, 'a passage.'

Anatomists have given this name to the openings at the extremities of vessels at the surface of different membranes. **EXHALANT PORES** have been supposed to exist in the exhalants, to transmit the fluids exhaled.—**ABSORBENT PORES** are employed in taking up parts that have to enter the circulation. Pores exist in the cuticle; yet Humboldt, with a powerful magnifying-glass, was unable to observe them. The pores of the skin have also been called *spira'cula*.

POREBLIND, *Myopic*.

PORED MILK, *Colostrum*.

PORI BILIA'RII, *Ductus bilia'rii* seu *biliferi*, *Canaliculi biliferi*, *Sudor'ici fell'et*. Ducts which receive the bile from the penicilli of the liver, and convey it to the hepatic duct.

PORI CUTA'NEI, *P. Exhalan'tes* seu *Sudoriferi* seu *Resorben'tes*, *Mea'tus cuticula'res*. The pores of the skin.

PORI EXHALANTES, *P. cutanei*—p. *Pulmonum*, *Cellulæ bronchicæ*—p. *Resorben'tes*, *P. cutanei*—p. *Sudoriferi*, *P. cutanei*.

PORNOGRAPHY, *Pornograph'ia*; from *πορνῆ*, 'a prostitute,' and *γραφω*, 'I describe.' A description of prostitutes or of prostitution, as a matter of public hygiene.

POROCE'LE, from *πῶρος*, 'callus,' and *κλῆη*, 'a tumour.' A hard tumour of the testicle or its envelopes.

POROMA, *Induration*.

POROMPHALOCLE, *Poromphalon*.

POROMPHALON, *Poromph'alus*, *Porompha-loce'læ*, from *πῶρος*, 'hard,' and *ομφαλός*, 'the navel.' Umbilical hernia with hard and thickened parietes.

POROS, *Canal*.

POROSIS, *Chalaza*, *Induration*.

POROT'IC, *Porot'icus*, (F.) *Porotique*, from *πῶρος*, 'callus.' A remedy believed to be capable of assisting the formation of callus.

PORPHYRA, *Purpura*—p. *Græcorum*, *Petechiæ*—p. *Hæmorrhagica*, *Purpura hæmorrhagica*—p. *Laciniata*, *Uvula umbilicalis*—p. *Umbilicalis*, *Uvula umbilicalis*.

PORPHYRISMA, *Scarlatina*.

PORPHYRISMUS, *Scarlatina*.

PORPHYRIZATION, *Levigation*.

PORPHYROTYPHUS, see *Typhus*.

PORPHYRU'RIA, *Porphyru'ria*; from *πορφύρα*, 'purple,' and *ουρον*, 'urine.' A state of the urine—*pur'puric urine*—in which it deposits the remarkable colouring matter—*Pur'purine*, *Uro'sacine*, *Rosac'ic acid*, *Uroër'ythin*, *Uro'rhodine*, *Ur'rhodine*—in association with urate of ammonia.

POR'PHYRY, *Porphyri'tes*, from *πορφύρα*,

'purple.' A species of very hard, compound rock, susceptible of considerable polish, and used in the fabrication of mortars and slabs for the apothecary.

PORPHYRY, BLACK. Ophtes.

PORPHYRIA, Porphyria.

PORRACÉ, Porraceous.

PORRACEOUS, *Porraceus*, *Pra'sinus*, from *porra*, 'a leak.' (F.) *Porracé*. Having the green colour of the leak. An epithet given to certain excreted matters, — the sputa, bile and faeces particularly.

PORREAU, *Allium porrum*.

PORRETUM, Liver.

PORRET, *Allium porrum*.

PORRIDGE, ATHOL, see Mahogany.

PORRIGINE, *Porrigio favosa*.

PORRIGINEUX, Porriginous.

PORRIGINOSUS, Porriginous.

PORRIGINOUS, *Porriginosus*, (F.) *Porrigineux*. Appertaining to porrigio.

PORRIGO (L.), 'seurf or scall in the head.' *Tinea*, *Favus*, *Furfuratio*, *Furfurex*, *Ecpye'sis* seu *Phlysis porrigio*, *Trichon'sis*, *Scal'd head* in general, *Scall'd*; from *porrigere*, 'to spread.' (F.) *Teigne*, *Rache*. Some of the varieties are contagious. It is principally characterized by an eruption of the pustules, called favi and achores, unaccompanied by fever.

PORRIGO CONFERTA, *P. scutellata*.

PORRIGO DECALVANS, *Tricho'sis are'a*, *Area*, *Alopec'cia areata* seu *partialis seu circumscripta*, *Phytolepor'cia*, *Trichon'sis furfuracea*, *Tyr'ia*, *Tyro'ma*, *Ophi'asis*, *Tin'ea tonsdens*, *Herpens tonsdens*, *Sporra tonsdens*, *Limited or partial Baldness*, consists of patches of baldness without decay or change of colour in the surrounding hair, the exposed parts of the scalp being glabrous, white, and shining. The morbid condition sometimes spreads, so that the patches coalesce, rendering the baldness extensive. To a vegetable growth, found at the roots of the hair in *Tinea tonsdens*, Gruby gives the name *Microsporon Audouinii*. It is, also, called *Tricophyton*, seu *Trichomyces tonsurans*. The treatment consists in keeping the scalp cleared by constant shaving, and in the steady application of some stimulating liniment or ointment.

PORRIGO FAVOSA, *Ce'ron*, *Favus*, *Ecpye'sis porrigio favosa*, *Tinea favosa*, *Scab'ies cap'itis favosa*, *Melice'ra*, *Melice'rma*, *Melice'ria*, *Mel'ifid'ria*, *Porrigoph'ya*, *Honey-comb Scall* or *Tetter*, (F.) *Teigne favense*, *Porrigine*. This consists in an eruption of pustules common to the head, trunk, and extremities, pea-sized, flattened at the top, in clusters, often united; discharge, fetid; scabs, honey-combed; the cells filled with fluid. It occurs both in early and adult life. Modern researches have satisfied some observers, that the crusts of favus are of vegetable nature, and formed of a parasitic plant, belonging, according to Gruby, to the obsolete genus *mycoderm'a*. MM. J. Müller and Lebert refer it to the genus *Oidium*, and call it *Oidium Schönleini*, whilst Romak assigns it to a new genus under the name *Achorion Schönleini*. The point is not, however, settled. The treatment of this variety must consist in the exhibition of laxatives and the mildest stimulating ointments, such as the *Ung. zinci*, *Ung. hydrargyri præcipitat. alb.*, &c. Eczema of the hairy scalp is often confounded with Porrigio.

PORRIGO FURFURANS, *Ecpye'sis porrigio furfuracea*, *Tinea seu Tricho'sis furfuracea* seu *porriginosa*, *Pityri'asis*, *Teigne furfuracea* (Alibert), commences with an eruption of small achores, the fluid of which soon concretes and separates in innumerable thin, laminated scabs, or scale-like exfoliations. It is attended with a good

deal of itching and some soreness of the scalp, to which the disease is confined. It occurs chiefly in adults. In the treatment, the scalp must be kept shaved. The part must be washed repeatedly with soap and water, and an oil silk cap worn. The nature of the ointments used must vary according to the period of disease and the irritability of the part affected. *Zinc ointment*; — one made with the *Oculus indicus*; — the *Sulphur ointment*; — *Citrine ointment*; *Ung. hydrargyri nitrico-oxidi*; *Tar ointment*, and *Nitric acid ointment* have all been found useful. Should inflammation or discharge arise, these applications must of course be intermitted.

PORRIGO LARVALIS, *P. lupinosa*.

PORRIGO LARVALIS, *Crusta lactea*, *Ecpye'sis Porrigio crustacea*, *Ment'igo*, *Achor*, *Tin'ea lactea* seu *Lactei*, *Lact'men*, *Lact'igo*, *Milky scall* or *Tetter*, *Milkbitch*, (F.) *Crôte de lait* ou *lait-euse*, *Fente volages*, *La Graine*. This is, almost exclusively, a disease of infancy. It commonly appears first on the forehead and cheeks, in an eruption of numerous, minute, whitish achores on a red surface. These break and discharge a viscid fluid, which becomes incrustated in thin, yellowish, or greenish scabs. The patches spread, until the whole face becomes, as it were, enveloped in a mask; hence the epithet *larvalis*. The disease is rather a variety of Eczema. Sometimes the eruption appears on the neck and breast; and, occasionally, on the extremities. In general, the health of the child does not suffer, but the eruption is always attended with much itching and irritation. The treatment consists in mild, tepid ablution, and the application of gently stimulating ointments, as the oxydum zinci. The bowels must be kept in a soluble state by small doses of calomel, or by rhubarb and magnesia.

PORRIGO LUPINOSA, *Tin'ea lupina*, *Favus disper'sus*, *Scabies cap'itis*, *Ecpye'sis porrigio lupinosa*, *Porrigio larva'lis*, *Impeti'go figura'ta*, *Tin'ea granulata* seu *mucosa*, *Crusta lactea* (of some), *Ment'agra infantum*, (F.) *Teigne*, *Dartre crustacée* *flavescente*, *Favus disséminé*, consists of minute pustules in small patches, mostly commencing on the scalp; the patches terminate in dry, delving scabs, resembling lupine seeds; the interstices often covered with a thin, whitish and exfoliating incrustation. It occurs chiefly in early life, and requires the same treatment as the last.

PORRIGO SCUTULATA, *P. conferta*, *Tin'ea granulata* seu *flo'osa*, *Ecpye'sis porrigio galeata*, *Acho'res* seu *Scab'ies cap'itis*, *Favus confertus*, *Ringworm of the Scalp*, *Scal'd head*, (F.) *Teigne annulaire*, *T. en anneau*, *T. nummulaire*, *Favus en groupes*. It commences with clusters of small, light, yellow pustules, which soon scab off; and, if neglected, become hard by accumulation. If the scabs be removed, the surface is left red and shining, but studded with slightly elevated points. If not attended to, it involves the whole head. It occurs in children three or four years of age, and is very unmanageable. The rules laid down under *Porrigio furfurans* must be here still more perseveringly enforced. The whole tribe of stimulating ointments may be used in succession. No one will always succeed, and hence the necessity of change, provided advantage should not seem to follow any particular application. The *Unguentum ad Scabiem* of Bonyer sometimes cures, when others fail.

PORRIGOPHYTA, *Porrigio favosa*. - *Porrigophyte* means the plant of favus or of porrigio favosa.

PORRILLON, *Narcissus pseudonarcissus*.

PORRUM, *Allium porrum*, *Thymion*.

PORRUS, *Sarcoma*, *Thymion*.

PORTA, πύλη, *Pylê*, (F.) *Porte*, 'a gate.' The

part of the liver or other organ—*hilus*—where its vessels enter as by a gate. See Liver. Also, the Vulva.

PORTA HEPATIS, see Liver — p. Jecoris, see Liver — p. Lientis, see Spleen.

PORTA VENA, *Vena porta*, *V. porta-rum* seu *ad portus* seu *portal'is* seu *magna* seu *quæ ad portus est seu stelechiæ'a seu lac'tea seu ostia'ria seu arteria'su seu arteria'lis*, *Roma'lis vena*, *Jan'tiric*, *Alanus Jec'oris seu he'patis*, *Vena ad portas*. The System of the *vena porta* or the Abdominal venous system or Portal system, (F.) *Veine porte*, *Porte veine*, is a vascular apparatus of black blood, situate in the abdomen, and resulting from two distinct orders of vessels, united into one common trunk. 1. One of these trunks, much more extensive than the other, has received the name *Vena porta abdominal'is*, because it arises from all the organs enclosed in the cavity of the abdomen, except the kidneys and bladder; and, in the female, the uterus. The Abdominal *vena porta* has two principal roots, the *splenic vein* and the *superior mesenteric*, which have their ramifications in all the digestive organs. 2. The other trunk, destined solely to the liver, has been called the *Hepatic vena porta*. It appears to be a continuation of the abdominal *vena porta*, and proceeds towards the inferior surface of the liver. Opposite the transverse fissure of that organ, it divides, like an artery, into two branches, which, by their union, seem to form a canal or duct, called by some the *Sinus of the vena porta*, or of the *portal vein*. These branches are distributed in the substance of the liver, whither they are accompanied by a prolongation of the fibrous capsule of that organ, — the capsule of Glisson. The different branches of the *vena porta* are devoid of valves. See Liver.

PORTAL, from *porta*, 'a gate.' Relating to the porta of the liver, — as the Portal system, or system of the *vena porta*. By extension, the term has been applied to an analogous system of vessels in the kidney. See Kidney.

PORTAL FISSURE, see Liver — p. Fossa, see Liver — p. System, see *Porta Vena* — p. System of the Kidney, see Kidney — p. Vein, *Porta Vena* — p. Veins of the Kidney, see Kidney.

PORTE-AIGUILLE, *Acutenaculum*, *Needle-carrier*, *Needle-bearer*, from (F.) *porter*, 'to carry,' and *aiguille*, 'a needle.' An instrument for accurately laying hold of a needle, and giving it greater length, when it is so fine and small that it cannot be held by the fingers. It is of steel or silver, about two inches long, and throughout the whole, almost, of its length, divided into two branches, so as to form a kind of forceps, capable of being closed by means of a sliding ring. A modification of this instrument was devised by Dr. Physick for taking up deep-seated arteries beyond the reach of a tenaculum or common needle. It consists of a forceps so constructed as to hold in its extremity a needle armed with a ligature. The handles of the forceps are fastened together, temporarily, by a spring or catch; and, when the needle is fairly deposited beneath the vessel, it is disengaged from the forceps and drawn out, leaving the ligature behind, which can be tied without difficulty.

PORTE-BOUGIE (F.). A silver canula similar to that of a trochar, except that it is longer. It is used for conducting bougies into the urethra.

PORTE-CAUSTIQUE, *Porte-pierre* — p. *Feuille*, *Subscapularis*.

PORTE-LAQ. An instrument for carrying the laq or fillet over the limb of the foetus whilst in utero, when turning is necessary.

PORTE-LIGATURE, from *porter*, 'to carry,'

and *ligature*. An instrument for conveying a ligature to deep-seated parts.

PORTE-MÈCHE (F.), from *porter*, 'to carry,' and *mèche*, 'a tent.' An instrument for carrying tents or *mèches* deeply into fistulous openings, or sinuses, to the bottom of ulcers, &c. It is a wire of silver or steel, four or five inches long, having a notch at one extremity, and a button at the other.

PORTE-MOXA (F.). A small instrument for keeping the moxa applied to a part to be cauterized. It consists of a metallic ring—separated from the skin by three supports of ebony—to which a handle is attached.

PORTE-NEUD (F.), from *porter*, 'to carry,' and *neud*, 'a knot.' An instrument for carrying a ligature round the base or pedicle of a tumour.

PORTE-PIERRE (F.), *Porte-caustique*, *Caustic bearer*. An instrument similar to a *porte-crayon* for carrying lunar caustic.

PORTE-SONDE (F.). An instrument resembling a *porte-crayon*, employed for holding the style or *sonde*, and inserting it more readily into the nasal duct, during the operation for fistula lacrymalis.

PORTE VEINE, *Porta vena*.

PORTENTUM, Monster.

PORTER, Cerevisia.

PORTIO DURA, Facial nerve — p. Inter duram et mollem, *Portio Wrisbergii* — p. Intermédia Wrisbergii, *Portio Wrisbergii* — p. Media inter portionem duram et portionem mollem, *Portio Wrisbergii*.

PORTIO MOLLIS, Auditory nerve.

PORTIO WRISBERGII, *P. intermédia Wrisbergii* seu *media inter portionem duram et portionem mollem* seu *inter duram et mollem*, (F.) *Faisceau intermédiaire de Wrisberg*, *Intermédiaire de Wrisberg*. A small, white fasciculus, intermediate between the *portio mollis* and *portio dura* of the seventh pair of nerves, which was first described by Wrisberg. The Chorda Tympani is supposed by some to emanate from this nerve.

PORTION COSTO-TRACHÉLIENNE DU SACRO-SPINAL, *Longissimus dorsi* — p. *Écailleux ou squameux du temporal*, *Squamous portion of the temporal* — p. *Lombaire du plexus cervical*, *Lumbar plexus*.

PORTION GODRONNÉE DE LA CORNE D'AMMON. A name given by Vicq d'Azyr to a small, denticulated cord, of a compact tissue and granular appearance, which unites the cornu ammonis to the paries of the lateral ventricle.

PORTONARIUM, Pylorus, Duodenum.

PORTULACA, *P. oleracea*, *Andrach'nè*, *Allium Gallicum*, *Purslane*, (F.) *Pourpier cultivé*. This plant abounds with a watery and somewhat acid juice; and is often put into soups or pickled with spices. It is regarded as antiseptic and aperient.

PORUS, Callus, Meatus — p. Acusticus, Auditory canal, internal.

PORUS OPTICUS. An opening in the centre of the cribriform lamella of the sclerotic, through which the central artery of the retina enters the eyeball.

POSCEA, Oxyerate.

POSE, Coryza.

POSIMOS, Potable.

POSITION, FALSE, False position.

POSOL'OGY, *Posolog'ia*, *Dosilog'ia*, from *posos*, 'quantity,' and *logos*, 'a discourse.' Indication of the doses in which the different articles of the materia medica ought to be exhibited. (F.) *Dosage*.

POSOQUERIA DUMETORUM, *Randia Dumetorum*.

POSSÉDÉ, Possessed.

POSSESS'ED, *Dæmoni'acus*, (F.) *Possédé*. One

affected with epilepsy, or other strange neurotic affection, was formerly supposed to be 'possessed' by a demon. Hence, exorcism and various religious ceremonies, at one time employed to drive off the intruder.

POSSET, *Possetum*, from (L.) *posca*, 'sour wine and water' [?]. Milk curdled with wine, or any acid, or treacle. A preparation of beer and milk; *Zythog'ala*.

POSSET, SACK, see **Sack**—*p.* Sago, see **Sago**.

POSSETTING, Regurgitation (of Infants).

POSTBRACHIALE, *Metacarpus*.

POSTCARPIUM, *Metacarpus*.

POSTERIOR ANNULA'RIS. An external, interosseous muscle, which arises from the corresponding sides of the metacarpal bones of the ring-finger and little finger, and is inserted into the inside of the tendon on the back of the ring-finger. Its use is to draw the ring-finger inwards.

POSTERIOR AURIS, Retrahens auris.

POSTERIOR INDICIS MANŪS. An internal, interosseous muscle, which arises from the inner or ulnar side of the metacarpal bone of the fore-finger, and is inserted into the inside of the tendon on the back of the fore-finger. Its use is to draw the fore-finger inwards.

POSTERIOR INDICIS PEDIS, Abduc'tor Indicis Pedis. An external, interosseous muscle, which arises from the contiguous sides of the metatarsal bones of the fore-toe, and second of the small toes, and is inserted into the outside of the root of the first bone of the fore-toe. Its use is to pull the fore-toe outwards.

POSTERIOR ME'DII DIG'ITI MANŪS. An external, interosseous muscle, which arises from the corresponding sides of the metacarpal bones of the middle and ring-fingers; and is inserted into the inside of the tendon on the back of the middle finger. Its use is to draw the middle finger inwards.

POSTERIOR MEDII DIG'ITI PEDIS, Abduc'tor medii digiti pedis. An external, interosseous muscle, which originates from the contiguous sides of the metatarsal bones of the second and third of the small toes, and is inserted into the outside of the root of the first bone of the second of the small toes. Its use is to pull the toe outwards.

POSTERIOR TER'TII DIG'ITI, Adduc'tor tertii digiti. An external, interosseous muscle, which arises from the contiguous sides of the metatarsal bones of the third and fourth of the small toes, and is inserted into the outside of the root of the first bone of the third of the small toes. Its use is to pull the toe outwards.

POSTHE, *Penis, Prepuce*.

POSTHET'OMIST, Posthet'omus, from *ποσθῆν*, 'prepuce,' and *τομή*, 'incision.' One who performs the operation of circumcision.

POSTHETOMY, Circumcision.

POSTHIA, Hordeolum.

POSTHIOPLAS'TIC, Posthioplas'ticus, from *ποσθῆν*, 'the prepuce,' and *πλασσω*, 'I form.' An epithet applied to the operation for restoring the prepuce.

POSTHIT'TIS, Acrobysti'tis, Acroposthi'tis, from *ποσθῆν*, 'the prepuce, the penis,' and *itis*, the termination for inflammation. Inflammation of the prepuce.

POSTHIUM, Prepuce.

POSTHON'CUS, from *ποσθῆν*, 'the prepuce or penis,' and *ὄγκος*, 'a tumour.' Swelling of the prepuce or penis.

POSTHOPLAS'TICA, from *ποσθῆν*, 'prepuce,' and *πλαστικός*, 'formative.' The art of making a new prepuce.

POST'UMOUS, from *post*, 'after,' and *humus*, 'the ground.' Occurring after death.

POSTHUMOUS CHILD. One born after the death of its father, or taken from the dead body of its mother, as by the Cæsarean section.

POSTIME, Abscess.

POST-MORTEM. 'After death;' as 'a post mortem or necroscopic examination.'

POST-NATUS. 'Born after.' In the old law writers, the second son.

POST-PARTUM. 'After delivery.' As 'post partum hemorrhage.'

POSTPOSIT'IO, (*post*, 'after,' and *ponere*, 'positum,' 'to place,') 'a postponement.' Delay in the return of a paroxysm. It is opposed to *anticipatio*, which means the return of a paroxysm earlier than its accustomed hour.

In this sense, we speak of an *anticipating*, and a *postponing* (*Febris hysteret'ica*) intermittent.

POSTREMA, Uterus.

POTABLE, Potab'ilis, from *potare*, 'to drink.' *Pot'imos*, *Potos*, *Pos'imos*, 'drinkable.' Water is said to be potable, when it contains atmospheric air; is not brackish; and has no mineral impregnation of consequence.

POTASH. So called from the pots or vessels in which it was made;—*Vegetable alkali, Gastrinum*. Also, *Potass*.

POTASH OF COMMERCE, Potassa imp'ura, *Impure potassa, Alu'men cat'ium, Cin'eres Rus'sici, Subcar'bonas potas'sæ imp'uras, Potas'sæ car'bonas impurus, Lix'ivus cinis*, the *Pearlash* of commerce, is prepared from the ashes of land-plants by calcination, solution in water, filtration, and evaporation. It usually consists of subcarbonate of potassa, sulphate of potassa, chloride of potassium, silica, oxide of iron, &c.

From the ashes of lees of wine, and from vine-twigs, a much purer alkali is obtained. It is called *Cinis infecto'rius seu fæcum, Cin'eres clavella'ti seu gravella'ti, Alu'men fæcum*, (F.) *Cendres gravelées ou de sarmen*.

Another variety of the alkali is afforded by burning argol in a crucible; afterwards powdering and calcining it, till it is nearly white; then dissolving in water, filtering, and evaporating. It is called *Salt of Tartar, Sal Tart'ari, Kali prap'aratum e tart'aro, K. e tartaro, Subcar'bonas potas'sæ puris'simus, Potas'sæ subcar'bonas e tart'aro*. Lastly, when wormwood is burnt to ashes, dissolved in water, filtered, and evaporated to dryness, it affords another variety of vegetable alkali, the *Sal absin'thiæ* or *Salt of wormwood*.

Impure potash is used for preparing the subcarbonate for medical use.

POTASS, Potassa fusa, Potash. The protoxide of potassium.

POTASSA, ACETATE OF, Potassæ acetas—*p.* Bichromate of, *Potassæ bichromas*—*p.* Bisulphate of, *Potassæ supersulphas*—*p.* Bitartrate of, *Potassæ supertartras*—*p.* Carbonate of, *Potassæ carbonas*—*p.* Carbonate of, pure, see *Potassæ subcarbonas*—*p.* Caustica cum calce, *Potassa cum calce*—*p.* Chlorate of, *Potassæ murias hyperoxy-genatus*—*p.* Chromate of, *Potassæ chromas*—*p.* Chromate of, neutral, *Potassæ chromas*—*p.* Chromate of, yellow, *Potassæ chromas*.

POTASSA CUM CALCÈ, P. opè calcis para'ta et ignè fusa, Potassa caust'ica cum calcè, Potassæ with lime, Calc' cum kali puro, Caust'icium communè for'tius, Lapis infernali's sive sep'ticus, Kali caust'icium cum calcè, Lapis caust'icus, (F.) *Potassæ préparée avec la chaux, P. à la Chaux, Pierre caust'ique*, (*Potassæ Calcis* ʒʒj. Rub together, and keep in a well-stopped bottle.—Ph. U. S.) Used for the same purposes as the next, but more manageable; being less deliquescent.

POTASSA, EUCHLORATE OF, Potassæ murias hyperoxygenatus—*p.* Ferrocyanate of, *Potassii ferrocyanure'tum*—*p.* Ferropotassiate of, *Potassæ cyanure'tum*.

POTASSA FUSA, *Potassa, Kali causticum seu purum, Fused potass, Alkali vegetabile fixum causticum, Causticum commuñe acer-rimum, C. alkalinum seu commuñe seu potentiale seu Sali-num, Hydras ka'licum, H. potassæ puræ fusæ, Potassæ hydras, Lapis causticus seu causticus chirurgicum seu infernalis alcalinus seu septi-cus, Ruptorium commuñe, Fixed reg'etable alkali, Hydrate of Potassa, (F.) Potasse caustique.* (Prepared by evaporating solution of potass to dryness in an iron vessel.) This preparation is generally run into cylindrical moulds, covered with paper, and kept in well-corked vessels. It is powerfully escharotic, and is used for forming issues, curing strictures, &c.

POTASSA, FUSIDA, *Potassa fusa*—p. Hydriodate of, *Potassæ hydriodas*—p. Hydrobromate of, see Bromine—p. Hydrocyanate of, *Potassii cyanure-tum*—p. Hypermanganate of, *Potassæ hypermanganas*—p. Hyperoxalas, *Potassæ oxalate* of—p. Hypertartaras, *P. super-tartaras*—p. Impura, *Potash of commerce*—p. and Iron, *tartarate of, Ferrum tartarizatum*—p. with Lime, *Potassa cum calce*—p. Liquida, *Liquor potassæ*—p. Mephite of, *Potassæ subcarbonas*—p. Monochromate of, *Potassæ chromas*—p. Muriate of, *Potassæ murias*—p. Nitrate of, *Potassæ nitras*—p. Ope calcis parata et igne fusa, *Potassa cum calce*—p. Oxy-muriate of, *Potassæ murias hyperoxygenatus*—p. Permanganate of, *Potassæ permanganas*—p. Prussiate of, *Potassii ferrocyanuretum*—p. and Soda, *tartarate of, Soda, tartarate of*—p. Subcarbonate of, *Potassæ subcarbonas*—p. Sulphate of, *Potassæ sulphas*—p. Sulphate of, acid, *Potassæ supersulphas*—p. Sulphuret of, *Potassæ sulphure-tum*—p. Supercarbonate of, *Potassæ carbonas*—p. Supersulphate of, *Potassæ supersulphas*—p. Supertartarate of, *Potassæ super-tartaras*—p. Tartrate of, *Potassæ tartras*.

POTASSÆ ACETAS, *Oxytartarus, Acetas kali seu lixivie seu ka'licum, Alkali tartari ace'to saturatum, A. vegetabile cum aceto, Magis-terium tartari purgans, Terra foliata Tartari, Kali acetatum, Sal diureticus seu Senner'ti, Aced'm tartari, Ac'etate of potass, (F.) Acetate de potasse.* (Acid. acetic. Oj, *Potassæ carbon, q. s.* Mix and add, by degrees, enough acetic acid to saturate the alkali; then filter and evaporate till a pellicle is formed, which is to be dried upon bibulous paper; evaporate again in a water-bath, and, removing the pellicles, dry.—Ph. U. S.)

It is cathartic and diuretic. Dose, $\mathfrak{z}\text{ij}$ to $\mathfrak{z}\text{ij}$, or more.

POTASSÆ ET ALUMINÆ SULPHAS, *Alumen*—p. Biantimonias, *Antimonium diaphoreticum*—p. *Bicarbonas, Potassæ carbonas*.

POTASSÆ BICHRONAS, *Kali chro'micum rubrum seu ac'idum seu bichro'micum, Bichro'mate of potassæ*. Procured by adding sulphuric, nitric, or acetic acid to a solution of the neutral chromate, and setting it aside to crystallize. It is a violent irritant poison; and has been used as a caustic in cases of morbid growths, in the proportion of from half a drachm to a drachm of the salt to from an ounce to an ounce and a half of water.

POTASSÆ BISULPHAS, *P. supersulphas*—p. *Bitartaras, P. super-tartaras*.

POTASSÆ CARBONAS, *Kali aëra'tum, Bicar-bonas potassæ, Potassæ bicar'bonas* (Ph. U. S., L., E., & D.), *Deutocarbonas potassii, Carbonate, bicarbonate or supercarbonate of potassa*. This is not caustic, like the subcarbonate of potass, and hence, has been preferred to relieve acidity. (*Potassæ carbon, lbij, Aqua destillat. Ox. Pass carbonic acid through the solution until it is saturated, and filter. Evaporate at a heat not exceeding 160°, and dry the crystals on bibulous paper. Carbonic acid is obtained from marble*

by the addition of dilute sulphuric acid.—Ph. U. S.) Dose, gr. x to $\mathfrak{z}\text{ss}$. It is used in forming the soda powders, as well as for its antacid properties.

Distillers and brewers prepare this salt with great facility by suspending a solution of the purified carbonate of potassa in the fermenting tun. The alkaline salt is thus surrounded by an atmosphere of carbonic acid; and by absorbing it, it crystallizes into bicarbonate. The salt in powder is called *sal aëratum*. Its composition is between that of a carbonate and of a bicarbonate.

POTASSÆ CARBONAS, *P. subcarbonas*—p. *Carbonas e Lixivo Cinere, P. subcarbonas*—p. *Carbonas impurus, P. subcarbonas*—p. *Carbonas purus, Pure carbonate of potassa, see P. subcarbonas*—p. *Chloras, P. murias hyperoxygenatus*—p. *Chloruretum oxydatum, P. murias hyperoxygenatus*.

POTASSÆ CHROMAS, *Kali chro'micum flavum seu neutrale, Neutral yellow Chromate or Monochromate of potassa*. Prepared by igniting a mixture of four parts of native chromite of iron (chrome iron ore) and one part of nitre; dissolving out the chromate of potassa by water, and crystallizing. It has been used as an emetic, expectorant and diaphoretic. Dose, as an emetic, from two to four grains; as an expectorant, from $\frac{1}{16}$ to $\frac{1}{4}$ grain. In the proportion of one drachm to 32 ounces of water, it has been used as an antiseptic wash—called by Jacobson, *Liquor conservatrix*.

POTASSÆ CITRAS, *Citrate of potassa. (Acid. citric. $\mathfrak{z}\text{x}$, Potass. bicarb. $\mathfrak{z}\text{xiv}$, Aq. q. s.* Dissolve the acid in the water; and after effervescence has ceased, strain and evaporate to dryness. The salt is deliquescent, and must be kept in well-stopped bottles. (Ph. U. S.) It possesses the refrigerant properties of the neutral salts, and may be given in fever, when to do little or nothing is the most advisable course. Dose, gr. x to $\mathfrak{z}\text{ss}$.

POTASSÆ ET FERRI TARTRAS, *Ferri tartarizatum*—p. *Hydras, Potassa fusa*.

POTASSÆ HYDRIDODAS, *Potassæ Protohydri'odas, Potassii protox'idi hydri'odas, Kali Hydriodin'icum seu Hydriod'icum, Hydriodas Ka'licus, Hydriodate of potass, (F.) Hydriodate de potasse*. If a solution of potass be poured on iodine, an iodate and hydriodate are formed. These may be separated by means of alcohol, which only dissolves the latter of these salts. The hydriodate, or rather the *Iodide of Potassium, Kalium Iodat'um, Potassii Iodure'tum seu Iod'idum, (F.) Iodure de potassium*, may be obtained by evaporation. When the *Iodide of potassium* is dissolved in water, it becomes the *Hydriodate of Potassa*. The following is the formula in the Ph. U. S. *Potass. $\mathfrak{z}\text{vj}$; Iodin. in pulv. $\mathfrak{z}\text{xvj}$; Carbon. ligni in pulv. subtiliss. $\mathfrak{z}\text{ij}$; Aq. bullient. Oijj.* Dissolve the potassa in the water; add the iodine gradually, stirring until the solution becomes colourless, and continue the additions until the liquid remains slightly coloured from excess of iodine. Evaporate to dryness, stirring in the charcoal towards the close. Rub to powder, and heat to dull redness in an iron crucible, keeping up the temperature for 15 minutes. When cool, dissolve out the saline matter with pure water, filter the solution, evaporate and set aside to crystallize. By evaporating and crystallizing as before, an additional quantity of crystals may be obtained. Virtues,—those of iodine.

POTASSÆ MURIAS, *Sal Digestivus seu febrifugus Syll'vi, Spiritus salis mari'ni coagulatus, Muriate of Potass. Bitter and refrigerant.*

POTASSÆ MURIAS HYPEROXYGENATUS, *Potassæ Chloras seu oxy'murias, Oxy'muriate, Hyperoxy'muriate, Chlorate or Euchlorate of potass,*

Protochlorate of potass'ium, Alkali vegetabilē sal'is-to-dephlogisticat'um, Chloruretum potassæ oxydat'um, Haloid'um oxygend'um, Oxyge'no-Chloruretum potass'ii, Kali chloric'um, Chloras Kal'icus depuratus, (F.) Chlorate de potasse. It has been regarded as antineuropathic, and as an alterative purifier of the blood and humours. It is of doubtful efficacy. Dose, 3 to 6 grains or more.

POTASSÆ NITRAS, Nitrate of Potass, Nitrum, N. primari'um, Sal nitrum, Nitre, Sulp'tre, Sal petra, Salse'do muerum, Sanguin'ia, Berenic'ium, Halinit'rum, Protonit'rate of Potass'ium, Nitrate of protoxide of Potassium, Kali nitricum, Nitras kal'icus seu lixiv'ia, Azot'as potass'icus, (F.) Nitrate on Azotate de potasse. Obtained from the putrefaction of animal matters, in contact with calcareous or alkaline earths, by elixivation; adding, if necessary, wood-ashes to supply the alkaline basis. It is formed, in an impure state, by nature, in warm climates.

It is diuretic and refrigerant, and, in large doses, purgative; externally, cooling and detergent. Dose, gr. x to ʒi. In large doses it is an acrid poison.

POTASSÆ NITRAS FUSUS SULPHATIS PAUCILLO MIXTUS, Sal prunell'ie, Crystal'is minera'lis, Nitras potass'æ cum sulphur' fusus, Nitrum tabulat'um, Lapis Prunell'ie, Aurod'gnum minera'le, Sore-throat salt, Mineral crystal. Nitrate of potass, mixed with a little sulphate of potass and run into moulds. Its properties are those of nitre.

POTASSÆ OXALAS, Ox'ulas Acid'ulum Potass'æ seu Potass'æ acid'ulum seu ba'sicus, Superox'alas kal'icum seu Potassæ, Hyperox'alas Potassæ, Sal Acetosell'ie, Oxal'ium, Kali oxal'icum acid'ulum, Ox'ulas subpotass'icus, Superox'alate of Protoxide of Potass'ium, Oxalate of Potass, Salt of Sorrel, (F.) Sel d'oseille, Sel à détacher. Prepared from the juice of wood sorrel. It is refrigerant, and is used to make lemonade, &c.

POTASSÆ OXALAS ACIDULEM, Potass, oxalate of— p. Oxymurias, P. murias hyperoxygenatus.

POTASSÆ PERMANGANAS, Perman'ganate of potass'ia, Hyperman'ganate of potassa, has been given in cases of diabetes. Dose, from gr. ij to gr. v, three times a day. It has been prescribed, also, in obstinate dyspepsia.

POTASSÆ PROTOHYDRIDAS, P. hydriodas— p. Purio fusæ hydras, Potassa fusa—p. et Sodæ tartras, Soda, tartrate of.

POTASSÆ SUBCARBONAS, P. Car'bonas (Ph. U. S.), P. carbonas e Lixiv'eo Cin'er'e, Subcar'bonas Kali, Car'bonas Potassæ, Kali præparat'um, Sal Herb'arum seu Planta'rum seu Tar'tari seu Absin'thii, Calcareous Tartar, Subcarbonate or Mephite of Potass, &c. (See Potash.) A filtered solution of the impure subcarbonate, Potass'æ car'bonas impur'us (Ph. U. S.), Pearl-ash (see Potash), evaporated to dryness. This deliquescent salt is given in the same cases as the potassæ carbonas, than which it is more caustic and disagreeable.

The Parisian codex has a *Subcar'bonas Potassæ ex tar'taro combus'to, vulgo Sal Tartari or Salt of Tartar—* (it is the *Potassæ carbonas purus, Pure Carbonate of Potassa, Ph. U. S.);* a *Subcarbonas Potassæ ex deflagrant'ibus simul tar'trat'is acid'ulo et Nitrate Potassæ paratus, vulgo Nitrum ex tar'taro fixum;* and a *Subcarbonas potassæ ex deflagrant'ib' cum Carbon'ibus Nitrat'is Potassæ paratus, vulgo Nitrum fixum.* They are all used like the preceding.

POTASSÆ SUBCARBONAS IMPURUS, Potash of commerce.

POTASSÆ SULPHAS, Kali vitriolat'um seu sulphur'icum, Sulphas Kali Lixiv'ia vitriola'ta, Nitrum vitriolatum Schroederi, Sal Sapient'ie seu Tar'tari Sulphurat'um seu Polychres'tus, Specif'icum Paracels'i, Spi'ritus vitrioli coagulatus

Mynsichti, Al'kali vegetabilē vitriolatum, Sal de duobus, Arcanum duplicat'um, Tar'tarum vitriolat'um, Panac'ea Ducis Hobol'tie seu duplicat'a, Sulphate of Potass, Protosulph'ate of Potass'ium, (F.) Sulfate de potasse. (The salt, that remains after the distillation of nitric acid, dissolved; the excess of acid saturated with sugar-carbonate of potass and crystallized.) Purgative and diuretic. Dose, ʒss to ʒj as a laxative. In a large dose it has proved an irritant poison.

POTASSÆ SULPHAS CUM SULPHURE, Sal Polychres'tus Glaseri, Lixiv'ia Vitriola'ta Sulphur'ea, (F.) Formed by deflagrating nitrate of potass with sublimed sulphur. Same virtues as the last. Dose, gr. xv to xxx.

POTASSÆ SULPHURETUM, Potassii sulphuretum— p. Superox'alas, Potassæ oxalas.

POTASSÆ SUPERSULPHAS, P. Bisulphas, Sal eris'mum seu febrif'ugus de Lem'ery, Supersulphate of Potass, Acid Sulphate or Bisulphate of Potass, Super-protosulphate of Potass'ium, (F.) Bisulfate de Potasse. (The salt remaining after the distillation of nitric acid, dissolved, strained, and crystallized.) It is refrigerant and purgative. Dose, ʒss to ʒij.

POTASSÆ SUPERTARTRAS IMPURUS, Tar'tarus crudus, Tar'tarum, Lapis vini, Maquet'ius, Pedagra, Tartar, (F.) Tarte cru ou brut. It is called, also, *Tartarum rubrum* and *Red Argol*, when obtained from red wines; *Tartarum album* or *White Argol*, when from white wines. Tartar is the saline deposit from wines as they become aged; which attaches itself to the sides of the casks and bottles in which they are kept. It consists of a considerable quantity of acidulous tartrate of potass, tartrate of lime, silica, alumen, oxyd of iron, manganese, and a red colouring matter, if the wine be red. It is chiefly used for the preparation of cream of tartar.

POTASSÆ SUPERTARTRAS, P. hypertar'tras, P. Bitar'tras (Ph. U. S.) seu tartras ac'id'a, Crystal'li Tar'tari, Tartras Acid'ulus Potassæ purus, T. superpotass'icus, T. subpotass'icus, Supertar'trate of Protoxide of Potass'ium, Bitartrate or Supertartrate of Potass, Crystals of Tartar, Cremor Tar'tari, Cream of Tartar (when in powder), Supertar'trate of Potass, (F.) Tartrate acide de potasse, Crème de Tartre. (The tartar of wine purified.) It requires 60 parts of water at 60° to dissolve it. It is mildly purgative, refrigerant and diuretic. Dissolved in water, with a small quantity of white wine, some sugar and lemon-peel, it forms an agreeable beverage in febrile diseases, under the name of *Imperial*. Dose, as a diuretic, ʒj to ʒij; as a purgative, ʒss to ʒi.

POTASSÆ TARTRAS, Kali tartarizat'um, Al'cali fixum tartarizat'um, Tar'tarum solu'bile, Tartarus tartarizat'us, Lixiv'ia Tartariza'ta, Sal Panchres'tum, Tartras Kali'us seu lixiv'ia, Sal vegetabil'is, Alkali vegetabilē tartarizatum, Tartrate of Potass, Soluble Tartar, (F.) Tartrate de Potasse. (Potassæ carbon. ʒxxvj; Potassæ Bitar'trat. in pulv. subtiliss. biij, vel q. s., aqua bullientis, cong. Dissolve the carbonate of potassa in the water, add the bitartrate to saturation, and boil; filter, evaporate till a pellicle forms, and set aside to crystallize. Pour off the liquid; dry the crystals on bibulous paper, and keep in closely-stopped bottles.—Ph. U. S.) A saline cathartic in the dose of ʒij to ʒj.

POTASSÆ TARTRAS ACIDA, P. supertartras— p. Tartras acidulus purus, P. supertartras—p. Tartras stibialis seu stibiosus, Antimonium tartarizatum.

POTASSÆ ACÉTATE DE, Potassæ acetas— p. Bisulphate de, Potassæ supersulphas—p. *Camstique, Potassa fusa—* p. *à la Chaux, Potassa cum calce—* p. *Chlorate de, Potassæ murias hyperoxygenatus—* p. *Hydriodate de, Potassæ hy-*

dríodas—*p. Liquide*, Liquor potassæ—*p. Nitrate de*, Potassæ nitræ—*p. Préparée avec la chaux*, Potassæ cum calce—*p. et de Soude*, tartrate de, Soda, tartrate de—*p. Sulfate de*, Potassæ sulphas—*p. Sulfure de*, Potassii sulphuretum—*p. Tartrate de*, Potassæ tartras—*p. Tartrate acide de*, Potassæ supertartras.

POTASSII BROMIDUM, *Potassæ hydrobromas*, Bromide of potassium, (F.) Bromure de potassium. (Bromin, \mathfrak{Zij} ; Ferri filii, \mathfrak{Zj} ; Potassæ carbon. \mathfrak{Zij} and \mathfrak{Zj} , vel q. s.; Aq. destillat., Oiv. Add the iron filings and bromine to a pint and a half of the water; stirring for half an hour. Apply a gentle heat, and when the liquor assumes a greenish colour, add gradually the carbonate of potassa, dissolved in Oiss of the water, until it ceases to cause a precipitate. Continue the heat for half an hour, and filter. Wash the precipitate with the remaining water, boiling hot, and filter. Mix the liquors and crystallize by evaporation.—Ph. U. S.) Virtues:—those of iodide of iron.

POTASSII CYANIDUM, P. Cyanuretum.

POTASSII CYANURETUM, P. Cyan'idum, Cyanuret of Potassium, Hydrocyanate of Potassa. (Potassii Ferrocyanur., in pulv. \mathfrak{Zviij} ; Potassæ Carbonat. exsicc., \mathfrak{Zij} . Mix intimately and throw into a deep iron crucible, heated to redness; keep up the heat till effervescence ceases, and the fused mass concretes, of a pure white colour, upon a warm glass rod dipped into it; then pour out the liquid carefully into a shallow dish to solidify, stopping before the salt becomes contaminated with the precipitated iron. Break up the mass whilst warm, and preserve in well-stopped bottles.—Ph. U. S.) Used in the same cases as the hydrocyanic acid. Dose, a quarter of a grain.

POTASSII DEUTO-CARBONAS, Potassæ carbonas.

POTASSII FERROCYNANURETUM, P. Ferrocyan'idum, Ferrocyanide or Ferrocyanuret of Potassium, Ferrocyanate or Ferropotassium or Prussiate of Potassa, (F.) *Protocyanure jaune de fer et de potassium*, Cyanure de fer et de potassium. The yellow double cyanuret of potassium and iron, the salt from which the cyanuret of potassium is obtained by calcination. It is not much used in medicine. It has been advised as an astringent in the colliquative sweats of phthisis, in leucorrhœa, &c.; but it does not seem to be possessed of much activity. Dose, 10 or 15 grains every four or five hours.

POTASSII HYDRARGYRO-IODIDUM, Potassæ Iodohydrargyris, Hydrargyrum biiodatum cum kalio iodato, Kalium iodatum hydrargyratum, Iodohydrargyrate or Hydrargyro-iodide of Potassium. This salt is made by the combination of iodide of potassium with red iodide of mercury. Its properties are similar to those of the iodides of mercury, but in a less degree.

POTASSII IODIDUM, Potassæ hydriodas—*p. Ioduretum*, Potassæ hydriodas—*p. Oxygeno-chloruretum*, Potassæ murias hyperoxygenatus—*p. Protoxidi hydriodas*, Potassæ hydriodas.

POTASSII SULPHO-CYANURETUM, Sulpho-cyanuret of Potassium. This salt has been prepared as a substitute for hydrocyanic acid, and cyanuret of potassium, on the ground of its possessing their advantages without their inconveniences.

POTASSII SULPHURETUM, Potassæ sulphuretum seu tersulphuretum, Kali sulphuretum, Hepar Sulphuris, H. sulphuris salinum seu sulphuris vulgare, Sulphuretum lixiviale seu kalium, Liver of Sulphur, Sulphuret of Potass, (F.) *Sulfure de potasse*, *Sulfure de potassium sulfaté*, *Foie de Soufre*. (Sulphur. \mathfrak{Zj} . Potassæ carb. \mathfrak{Zij} . Rub them together, and place the mixture in a covered crucible upon the fire until they unite. It has been employed in affections of the chest, as an expectorant, in the dose of from \mathfrak{Zss} to \mathfrak{Zj} ,

largely diluted; but it is chiefly used as a bath or wash in cases of itch, tinea capitis, and other cutaneous affections.)

POTASSII TER-SULPHURETUM, Potassii sulphuretum.

POTASSIUM. Same etymon as potash. The metallic base of potassa.

POTASSIUM, *BROMURE DE*, see Bromine—*p. Cyanuret of*, Potassii cyanuretum—*p. Ferrocyanide of*, Potassii ferrocyanuretum—*p. Ferrocyanuret of*, Potassii ferrocyanuretum—*p. Iodide of*, Potassæ hydriodas—*p. Iodohydrargyrate of*, Potassii hydrargyro-iodidum—*p. Iodure de*, Potassæ hydriodas—*p. Oxide of*, Potassæ—*p. Protoarsenite of*, Arsenite of protoxide of potassium—*p. Protochlorate of*, Potassæ murias hyperoxygenatus—*p. Protinitrate of*, Potassæ nitræ—*p. Protoxide of*, Potassæ—*p. Protoxide of, nitrate of*, Potassæ nitræ—*p. Protoxide of, superoxalate of*, Potassæ oxalate of—*p. Protoxide of, supernitrate of*, Potassæ supertartras—*p. Protosulphate of*, Potassæ sulphas—*p. Sulfure de*, sulfaté, Potassii sulphuretum—*p. Sulphuret of*, Potassii sulphuretum—*p. Superprotosulphate of*, Potassæ supersulphas.

POTATO, Solanum tuberosum.

POTATO FLY, Lytta vittata.

POTATO, HOG, Convolvulus panduratus.

POTATO OIL, Oil, fusel. By distilling potato oil with anhydrous phosphoric acid, a volatile colorless oily liquid is procured, which is lighter than water, and boils at 102°. This is *Amylene* or *Amylen*, (F.) *Amylène*. It is best prepared from fusel oil by the action of chloride of zinc. The vapour has been employed as an anæsthetic.

POTATO, SPANISH, Convolvulus batatas—*p. Sweet*, Convolvulus batatas.

POTATO VINE, WILD, Convolvulus panduratus.

POTATO, WILD, Convolvulus panduratus.

POTBELLY, Physconia.

POTECARY, Apothecary.

POTELÉE, Hyoseyamus.

POTENTIA, Force.

POTENTIA seu POTES'TAS COËUN'DI, Capability of copulation.

POTENTIA GENERAN'DI. The power of procreation.

POTENTIA IRRITANS, Stimulus.

POTENT'IAL, *Potent'ialis*, (F.) *Potentiel*; from *potens*, 'able.' An epithet for remedies which, although energetic, do not act till sometime after their application. The caustic alkalies, for instance, are called *Potentia* canteries in contradistinction to the hot iron, which is termed *actual*.

POTENTILLA, P. reptans.

POTENTILLA ANSERINA, P. argenti'na seu impolita seu neglecta, Argenti'na vulgaria, Fragaria anseri'na, Ar'gentine, Chenopodicon, Argenti'ria, Anseri'na, Silver Weed, Wild Tansy, Moorgrass, (F.) Argentine, Potentille, Herbe aux oies. Ord. Rosacæ. The leaves are mildly astringent and tonic.

POTENTILLA ARGENTINA, P. anserina—*p. Impolita*, P. anserina—*p. Neglecta*, P. anserina.

POTENTILLA NORVEG'ICA, Norway Cinquefoil, an indigenous plant, is possessed of similar properties.

POTENTILLA PALUSTRIS, Comarum palustre.

POTENTILLA REPTANS, Pentaphyllum, Fragaria pentaphyllum, Potenti'la, Quinquefolium, Q. majus, Common cinquefoil, (F.) Potentille rampante, Quintefeuille. The roots are possessed of astringent properties.

POTENTILLA TORMENTILLA, Tormentilla.

POTENTILLE, Potentilla anserina—*p. Ram-pante*, Potentilla reptans.

POTENTIZING, see Dynamic.

POTERIUM SANGUISORBA, Pimpinella saxifraga.

POTESTAS COEUNDI, Potentia coeundi—p. Generandi, Potentia generandi.

POTEX, Anus.

POTHOMORPHA PELTATA, *Pothomorphè peltata* [?], (*potthos*, and *poton*, 'shape,') *Cacaba* [?]. A Brazilian plant, (*Order*, Piperaceæ, which is possessed of diuretic properties, and has been given in strangury.

POTHOMORPHA UMBELLATA, also called *Cacaba* [?], is used in Brazil to modify nutrition, as in glandular affections, in which it is also given externally in the form of decoction.

POTHOS FETIDA, *Dracontium foetidum*—p. Putorii, *Dracontium foetidum*.

POTIMOS, Potable.

POTIO, Drink. Haustus, Potion—p. Calcis carbonatis, *Mistura cretæ*—p. Effervescentis anti-emeticæ, *Potion of Riverius*—p. Leniens, *Looch ex ovo*—p. Oleosa, *Looch ex ovo*—p. Picea, see *Pinus sylvestris*.

POTION (F.), *Potio*; from *poto*, 'I drink.' *Hydrolè*, *Hydrolite*. A liquid medicine, from 4 to 8 ounces or more in quantity, the composition of which varies, and which is administered in divided doses. It corresponds to the English mixture. See *Mistura*.

POTION PECTORALE, *Looch ex ovo*.

POTION OF RIVERIUS, A mixture of *Lemon Juice* and *Subcarbonate of Potass*,—generally given in a state of effervescence. It is the *Potio effervescentis anti-emeticæ dicta Riverii*, of the Parisian codex. The name in the codex indicates its use.

POTIONIS DESIDERIUM, Thirst.

POTIRON, *Cucurbita pepo*.

POTIUNCULA, Haustus.

POTOPARANÆA, *Delirium tremens*.

POTOS, Potable.

POTOTROMOPARANÆA, *Delirium tremens*.

POTTINGAR, Apothecary.

POTT'S CURVATURE, Vertebral disease.

POTT'S DISEASE, Vertebral disease.

POTUS, Drink—p. Hippocraticus, *Claret*.

POU, Pediculus.

POUCE, Pollex, see *Digitus*.

POUCHED, Encysted.

POUCHES, LARYNGEAL, see Laryngeal pouches.

POUDRE, Powder, Pulvis—p. Absorbante et anti-acide, Pulvis de magnesiâ compositus—p. d'Alun composée, Pulvis sulphatis aluminæ compositus—p. Antiarthritica amère, Pulvis de amaris compositus—p. Antiasthmaticque, Pulvis de sulphure et scillâ—p. Antimonial composée ou de James, Antimonial powder—p. Aromatique, Pulvis cinnamomi compositus—p. d'Asarum composée dite sternutatoire, Pulvis asari compositus—p. Balsamique, Pulvis myrrhæ compositus—p. de Birckmann, Pulvis de aro compositus—p. de Blanchement, Calcis chloridum—p. de Cannelle composée ou aromatique, Pulvis cinnamomi compositus—p. Castillon, see Powders, Castillon—p. Cathartique, Pulvis scammonæ compositus—p. Caustique du frère Cosme, Powder, arsenical, of Côme—p. de Chypre, see *Cladonia rangiferina*—p. du Comte de Palme, Magnesiæ carbonas—p. Cornachine, Pulvis cornacini—p. de Corne brûlée avec opium, Pulvis cornu usti cum opio—p. Coton, see Collodion—p. de Craie composée, Pulvis cretæ compositus—p. de Craie composée avec opium, see Pulvis cretæ compositus cum opio—p. de Dover, Pulvis ipecacuanhæ compositus—p. du Duc de Portland, Pulvis Ducis Portlandiæ—p. de Dupuytren, Powder of Dupuytren—p. Gazifère laxative, Powders, Seidlitz—p. Gazifère simple, Powders, soda—p. de Gomme adragant composée, Pulvis tragacanthæ compositus—p. de Gomme gutte, Pulvis e gummi guttâ—p. Hydragogue, Pulvis e gummi guttâ—p. Incisive, Pulvis

de sulphure et scillâ—p. d'Ipecacuanha et opium, Pulvis ipecacuanhæ compositus—p. de Jalap composée, Pulvis jalapæ compositus—p. de Kino composée, Pulvis kino compositus—p. de la Mirandole, Pulvis Ducis Portlandiæ—p. de Pérard, Pulvis de sennâ, scammonæ et lignis—p. de Quercetanus, Pulvis e gummi guttâ—p. de Rousselot, Powder, arsenical, of Côme—p. de Saint Ange, Pulvis asari compositus—p. de Savon végétale, Pulvis gummosus alkalinus—p. de Scammonée composée, Pulvis scammonæ compositus—p. de Seidlitz, Powder, Seidlitz—p. de Sinc composée, Pulvis sennæ comp.—p. de Sentinelli, Magnesiæ carbonas—p. Styptique, Pulvis sulphatis aluminæ compositus.

POUDRE SUBTIL DE DELCROIX. A fashionable depilatory "for removing superfluous hair in less than ten-minutes." It is said to consist of *Quicklime* and *Sulphuret of Arsenic*, with some vegetable powder.—Paris.

POUDRE TEMPÉRANTE DE STAHL, Pulvis de sulphate potassæ compositus—p. de Tennant, Calcis chloridum—p. de Valentini, Magnesiæ carbonas—p. Vermifuge mercurielle, Pulvis e sulphureto hydrargyro nigro et scammonio—p. Vermifuge sans mercure, Pulvis ex helminthocorto compositus.

POUDRE DE ZWINGER, Magnesiæ carbonas.

POUGUES, MINERAL WATERS OF. These alkaline waters are in the department of Nièvre, France. They contain carbonic acid, carbonates of soda, lime, and magnesia; chloride of sodium; alumina, and silica, with protoxide of iron.

POUKE, Papula.

POULAIN, Bubo.

POULET, VIN DE (F.) *Pullet Wine*. A disgusting preparation, made by infusing the excrement of the pullet in white wine. According to Nysten, it is employed by the vulgar in France in a number of diseases, and especially in falls, contusions, &c.

POULIOT, *Mentha pulegium*.

POULS, Pulse—p. Abdominal, Pulsus inferior—p. des Adultes, Pulse, natural—p. Agité, Pulsus fortis—p. Bas, Pulse, low—p. Capital, Pulse, nasal—p. Combiné, Pulse, complex—p. Complicqué, Pulse, complex—p. Composé, Pulse, complex—p. Contracté, Pulse, contracted—p. Défaillant, Pulsus deficient—p. Déprimé, Pulse, depressed—p. Développé, Pulse, developed—p. de la Diarrhée, Pulse, intestinal—p. Dur, Pulse, hard—p. Élevé, Pulse, ardent—p. Embarrassé, Pulsus intricatus—p. de l'Estomac, Pulse, stomachal—p. Étroit, Pulsus strictus—p. de l'Expectoration, Pulse, pectoral—p. Faible, Pulse, feeble—p. du Foie, Pulse, hepatic—p. Grand, Pulse, large—p. Insensible, Pulsus deficient—p. Intercaudent, Pulse, intermittent—p. des Intestins, Pulse, intestinal—p. Languissant, Pulse, languid—p. Lent, Pulse, slow—p. de la Matrice, Pulsus uterinus—p. Moité, Pulse, natural—p. Mou, Pulse, soft—p. de Nez, Pulse, nasal—p. Ondoyant, Pulse, undulating—p. Ondulant, Pulse, undulating—p. Palpitant, Pulse, tremulous—p. Parfait, Pulse, natural—p. Petit, Pulse, small—p. Plein, Pulse, full—p. Précipité, Pulse, frequent—p. Profond, Pulse, deep—p. Prompt, Pulse, sharp—p. Rare, Pulse, slow—p. Rebondissant, Pulsus dicrotus—p. Redoublé, Pulsus dicrotus—p. des Règles, Pulsus uterinus—p. Roide, Pulse, tense—p. Serratile, Pulsus serrinus—p. Serré, Pulse, contracted—p. de la Sueur, Pulsus sudoralis—p. Tardif, Pulse, slow—p. Tendu, Pulse, tense—p. Tremblant, Pulse, tremulous—p. de l'Urine, Pulsus urinialis—p. Véhément, Pulsus fortis—p. Ventral, Pulsus inferior—p. Vide, Pulsus vacuus—p. Vif, Pulse, sharp—p. Vite, Pulse, sharp.

POULTICE, Cataplasm. For different poul-tices, see Cataplasm and Cataplasma.

POUNCE, Sandarac.

POUND, *Libra*, *Litra*, *λίτρα*, *Pondo*, *Ba-ged'ia* (of twelve oz.), *Bard'dia*, (F.) *Livre*, from *pondere*, 'to weigh.' A weight, consisting of twelve ounces Troy,—sixteen ounces avoirdupois. See Weights and Measures.

POURD-MILK, Colostrum.

POURPIER CULTIVÉ, Portulaca.

POURPRE, *Purpura*—*p. Blanc*, Miliary fever—*p. de Cassius*, Purple of Cassius.

POURRIURE, Putrefaction—*p. d'Hôpital*, Hospital gangrene.

POUS, Pes.

POUSSE, Pursiness.

POUSSEE, Lichen.

POUSSIERE TUBERCULEUSE (F.), 'Tuberculous Dust.' A name given by MM. Rilliet and Barthez to fine white or yellow minute points of tuberculous matter occurring in myriads, so as to be likened to dust.

POUSSIF, see Pursiness.

POW, Head.

POWDER, Pulvis.

POWDER OF ETHIOPIA, Guy's. A once celebrated anthelmintic, which consisted of pure *rasped tin*, *mercury*, and *sulphur*, triturated together.

POWDER OF ALOES, COMPOUND, *Pulvis aloes compositus*, *Pulvis sulphatis aluminæ compositus*—*p. of Aloes with canella*, *Pulvis aloes cum canella*—*p. Alum*, compound, *Pulvis sulphatis aluminæ compositus*—*p. Antimonial*, *Chenevix's*, *Antimonial powder*—*p. Aromatic*, *Pulvis cinnamomi compositus*.

POWDER, ARSENICAL, OF CÔME, *Côme's powder*, (F.) *Poudre caustique du frère Cosme ou de Rousselot*—a caustic—is formed of *arsenious acid*, gr. x: *red sulphuret of mercury*, gr. xl: powdered *animal charcoal*, gr. x. It is made into a paste with mucilage.

POWDER, ARSENICAL, OF JUSTAMOND, is made by melting *sulphuret of antimony* 16 parts: *arsenious acid* 1 part, in a crucible. When the mass is cold it must be reduced to powder, and five parts of *extract of opium* be added.

POWDER OF ARUM, compound, *Pulvis de aro compositus*—*p. of Asarabacca*, compound, *Pulvis asari compositus*—*p. of Bladder-wrack*, yellow, *Pulvis quercus marinæ*—*p. Bleaching*, *Calcei chloridum*—*p. Camboge*, *Pulvis e gummi guttâ*—*p. Cardinal del Lugo's*, *Cinchona*—*p. Castillon*, see *Powders*, *Castillon*—*p. of Catechu*, compound, *Pulvis Catechu compositus*—*p. of Chalk*, compound, *Pulvis cretæ compositus*—*p. of Chalk with opium*, compound, *Pulvis cretæ compositus cum opio*—*p. of Cinnamon*, compound, *Pulvis cinnamomi compositus*—*p. Côme's*, *Powder arsenical of Côme*—*p. of Contrayerva*, compound, *Pulvis contrayervæ compositus*—*p. Countess's*, *Cinchona*—*p. Countess of Kent's*, *Cantidus pulvis*—*p. Cyprus*, *Hibiscus abelmoschus*—*p. Dover's*, *Pulvis ipecacuanhæ compositus*.

POWDER OF DUPUYTREN (CAUSTIC) (F.) *Poudre de Dupuytren*, formed of 1 part of *arsenious acid* and 200 parts of mild chloride of mercury. A mild and manageable caustic for causing an eschar on exposed surfaces, and in lupus, &c.

POWDER, EARL OF WARWICK'S, *Pulvis cornachini*—*p. of Ethiopia*, *Powder of Ethiopia*.

POWDER OF FAYNARD. A celebrated powder for arresting hemorrhage, said to be nothing more than the charcoal of beech wood finely pulverized.

POWDER, FEVER, SCHWANBERG'S, *Antimonial powder*.

POWDER-FOLDER, In Pharmacy, an apparatus by which papers containing powders are folded of the desired length.

POWDER, GINGER-BEER, see *Amomum zingiber*—*p. Gout*, *Duke of Portland's*, *Pulvis Ducis Portlandiæ*—*p. Gray*, *Hydrargyrum cum cretâ*—*p. of burnt Hartshorn with opium*, *Pulvis cornu cervi cum opio*—*p. of Ipecacuanha*, compound, *Pulvis ipecacuanhæ compositus*—*p. of Jalap*, compound, *Pulvis jalapæ compositus*—*p. James's*, *factitious*, *Antimonial powder*—*p. Jesuit's*, *Cinchona*—*p. of Kino*, compound, *Pulvis kino compositus*.

POWDER, KNOX'S. A disinfecting preparation, composed of eight parts of *chloride of sodium*, and three parts of *chlorinated lime*. An ounce, dissolved in a tumblerful of water, forms a solution of proper strength.

POWDER OF LEMONADE, see *Lemonade*—*p. of Magnesia*, compound, *Pulvis de magnesiæ compositus*—*p. of the Marquis*, *Marchionis pulvis*—*p. Mead's*, against the bite of a Mad dog, *Pulvis antilyssus*—*p. Myrrh*, compound, *Pulvis myrrhæ compositus*—*p. Portland*, *Pulvis Ducis Portlandiæ*—*p. of Rhubarb*, compound, *Pulvis rhei compositus*—*p. Saline*, compound, *Pulvis salinus compositus*—*p. of the three Sanders*, see *Diatrium*—*p. Scammony and aloes*, *Pulvis scammonii cum aloë*—*p. Scammony*, compound, *Pulvis scammonia compositus*—*p. of Senna*, compound, *Pulvis sen-næ compositus*—*p. Sneezing*, see *Veratrum album*—*p. Spruce beer*, see *Spruce beer*—*p. of Sympathy*, *Sympathetic powder*—*p. Tabor's*, *Cinchona*—*p. Tennant's*, *Calcei chloridum*—*p. Tonquin*, *Pulvis cobbii*—*p. Tooth*, *Dentifrice*—*p. Tra-gacanth*, compound, *Pulvis tragacanthæ compositus*.

POWDER, VIENNA, is composed of equal parts of quicklime and potassa cum calce. In using it, it is made into a paste—*Vienna paste*, (F.) *Caustique de Vienne*, *Pâte Caustique de Vienne*—with spirits of wine, and is applied on the diseased part, previously circumscribed by a hole cut in a sticking plaster.

To be left on from 10 to 20 minutes, according to the depth of the eschar required. Used in cases of lupous ulceration.

By fusing the potass and lime, and raising the heat so as to cause ebullition, the caustic may be run into cylinders:—when it is called *Caustique-Filhos*,* *Causticum Viennense fusum Filhos*, from the name of the author of the process.

POWDERS, CASTILLON, or CASTILHON, so called from the name of the physician who first proposed them. They are formed of *sago*, in powder; *salep*, in powder; *tragacanth*, in powder, each, four parts; *prepared oyster-shells*, one part, *cochineal*, half a part, divided into powders of one drachm each; one of which is given three or four times a day as a demulcent in diarrhoea and dysentery. Dorvault makes *poudre de Castilhon* synonymous with arrow root.

POWDERS, SODA, *Pulveres effervescentes*, (F.) *Poudres gazifères simples*. (*Sodæ carb. 3ss*, in each *blue paper*; *tartaric acid*, gr. xxv, in each *white paper*; for half a pint of water.) A pleasant, cooling drink; taken during the effervescence.

Seidlitz Powders, (F.) *Poudres gazifères laxatives* ou *Poudres de Seidlitz*, contain one or two drachms of tartrate of potassa and soda, added to the alkali in blue paper.

POWER, Faculty—*p. Inherent*, *Irritability*—*p. Sensorial*, see *Sensorial*—*p. Tonic*, *Irritability*.

POX, Syphilis, Variola—*p. Chicken*, *Varicella*.

POX DOCTOR, Syphilidiater.

POX, FRENCH, Syphilis—*p. Grease*, see *Grease*—*p. Horn*, see *Varicella*—*p. Noddy*, *Syphiloma-nia*—*p. Small*, *Variola*—*p. Swine*, see *Varicella*—*p. Water*, see *Varicella*.

POZZUOLI, MINERAL WATERS OF. Pozzuoli is in the neighbourhood of Naples. The springs are, 1. *Aqua della pietra*, Temp. 91° Fah. 2. *Aqua di Cavalcanti*, Temp. 100°. *A. di subævi homini*, Temp. 102°. *A. del cantarello*, Temp. 86 to 89°; and that of the Temple of Serapis in Pozzuoli, Temp. 102 to 107°. The last contains carbonic acid, carbonates of lime, magnesia, alumina, and iron, carbonate of soda, chloride of sodium, sulphate of lime, and silica.

PRACTICE OF PHYSIC, Therapeutics.

PRACTITIONER, GENERAL, Surgeon apothecary—p. Irregular, see Irregular—p. Regular, see Regular—p. Routine, Routinist.

PRÆ. In composition, means 'before.'

PRÆBIUM, Dose.

PRÆCAUTIO, Prophylaxis.

PRÆCIPITANS, Absorbent—p. Magnum, Sepia.

PRÆCIPITATUM ALBUM, Hydrargyrum præcipitatum.

PRÆCIPITATUS RUBER, Hydrargyri nitrico-oxydum.

PRÆCISIO PRÆPUTII, Circumcision.

PRÆCOCIA, Prunus Armeniaca.

PRÆCOCITY, Precocity.

PRÆCOGNITIO, Prognosis.

PRÆCORDIA, from *præ*, and *cor*, *cordis*, 'the heart.' The diaphragm; also, the thoracic viscera, and the epigastrium.

PRÆCORDIAL or PRÆCORDIAL REGION, *Re'gio præcordica* seu *præcordialis*, is the epigastric region. Also, and more properly, the region of the heart. A line drawn from the inferior margins of the third ribs across the sternum, passes over the valves of the pulmonary artery a little to the left of the mesial line, and those of the aorta are behind them, but about half an inch lower down. A vertical line, coinciding with the left margin of the sternum, has about one-third of the heart, comprising the upper portion of the right ventricle, on the right,—and two-thirds, comprising the lower portion of the right ventricle and the whole of the left, on the left. This is the præcordial region.

PRÆCORDIUM, *Fossette du cœur*—p. Pedis, Metatarsus.

PRÆCOX MATURITAS, Precocity.

PRÆCUSTODIA, Prophylaxis.

PRÆDIASTOLICUS, Prediastolic.

PRÆDICTIO, Prognosis.

PRÆDISPOSITIO, Predisposition.

PRÆDIVINATIO, Mantia.

PRÆDOMINIUM, from *præ*, and *domo*, *domitum*, 'I subdue.' Predominance of one thing over another.

PRÆDOMINIUM AQUÆ. A predominance of watery fluid,—*Tenuitas aquosa*.

PRÆDOMINIUM SAN'GUINIS ARTERIO'SI. A predominance of arterial blood, *Arteriositas sanguinis*.

PRÆDOMINIUM SAN'GUINIS VENO'SI, *Venositas sanguinis*. Predominance of venous blood.

PRÆFOCATIO, Orthopnea, Suffocation—p. Faucium, Angone—p. Matricis, Angone—p. Uterina, Angone.

PRÆFOCATION, Strangulatio uterina.

PRÆFORMATIO, Preformation.

PRÆGNANS, Pregnant.

PRÆGNAS, Pregnant.

PRÆGNATIO, Fecundation, Pregnancy.

PRÆGNATUS, Fecundation, Pregnancy.

PRÆGNAX, Pregnant.

PRÆGUSTATIO, *Avant-gout*.

PRÆHENSIO, Prehensio.

PRÆLINGUA. The anterior part or tip of the tongue.

PRÆLIUM, Coition, Press, Tourniquet—p. Arteriale, Tourniquet.

PRÆLUMBARIS, Prelumbar.

PRÆNOTIO, Prognosis.

PRÆNUNTIATIO, Prognosis.

PRÆPARATIO, Preparation—p. Chyli, Chylification.

PRÆPARATUM, Preparation.

PRÆPOTENTIA, from *præ*, and *potens*, 'powerful,' Great power or strength.

PRÆPOTENTIA VIRILIS. Great, or too great procreative power on the part of the male.

PRÆPOTIO, Propositus.

PRÆPUTIALIS, Preputial.

PRÆPUTIUM, Prepuce.

PRÆSAGIUM, Mantia.

PRÆSCIENTIA, Prognosis.

PRÆSCITIO, Prognosis.

PRÆSCRIPTIO, Prescription.

PRÆSCRIPTUM, Prescription.

PRÆSEPIOLUM, Alveolus.

PRÆSEPIUM, Alveolus.

PRÆSERVATIO, Prophylaxis.

PRÆSERVATIVUM, Amuletum.

PRÆSERVATORIA INDICATIO. A preservative indication; one proper for preventing the development of disease.

PRÆSERVATORIUS, Prophylactic.

PRÆSPINALIS, Prespinal.

PRÆTIBIALIS, Pretibial.

PRAIRIE BUSH, STINKING, *Ptelea trifoliata*—p. Pines, *Liatris*.

PRASINUM VIRIDE, *Cupri subacetes*.

PRASINUS, Porraceous.

PRASI'TES, from *prasium*, 'horehound.' Wine in which the leaves of horehound have been infused.

PRASIMUM, Marrubium.

PRASUM, Allium prasum.

PRAXIS, Action.

PRAXIS MEDICA. The practical exercise of the healing art.

PREASER, Rennet.

PRECIPITATE, RED, Hydrargyri nitrico-oxydum—p. White, Hydrargyrum præcipitatum.

PRECIPITATION, *Præcipitatio*, (F.) *Précipité*, from *præcipitare*, *præcipitatum*, (*præ*, and *capit*, 'the head,' 'to throw headlong.' The action by which a body abandons a liquid in which it is dissolved, and becomes deposited at the bottom of the vessel. The matter, so thrown down or precipitated, is called a *precipitate*.

PRÆCIPITATION DE LA MATRICE, Procidencia uteri.

PRÆCIPITE BLANC, Hydrargyrum præcipitatum—p. Noir, see Hydrargyri oxydum cinereum—p. Oxydum per se, Hydrargyri nitrico-oxydum—p. Rouge, Hydrargyri nitrico-oxydum.

PRECOCITY, *Præcociousness*, (F.) *Précocité*, *Præcox maturitas*, *Maturitas præcocious*, from *præ*, 'before,' and *coquere*, 'to cook.' Rapid development before the usual time. See *Præotia*.

PRÆCORDIAL, Præcordial.

PRÆCURSOR, *Præcursorius*, (F.) *Précurseur*, from *præ*, and *curro*, *currere*, 'I run.' That which precedes.

PRÆCURSOR or PREMONITORY SIGNS or SYMPTOMS, (F.) *Signes précurseurs ou prémonitoires*, *S. avant coureurs*, are those which indicate an approaching disease.

PREDIASTOLIC, *Prædiastolicus*, from *præ*, and *diastole*. That which precedes the diastole of the heart,—as 'prediastolic friction sound.'

PREDISPOSANT or PREDISPOSING CAUSES, see Causes, predisponent.

PREDISPOSIT'ION, *Prædispositio*, *Semina morborum* (Gaubius); from *præ*, and *disponere*, *dispositum*, 'to dispose.' (F.) *Imminence morbide*. That constitution or condition of the body, which disposes it to the action of disease

under the application of an exciting cause. Thus, in hereditary diseases, the conformation is such, that but a slight exciting cause may be required to produce them. When the disease arises solely from the predisposition, or under the additional influence of an extremely slight exciting cause, it is, by some, termed a *disposition*.

PREDORSAL, *Prædorsalis*, from *præ*, and *dorsum*, 'the back.' That which is in front of the back.

PREDORSAL REGION of the vertebral column is the anterior surface of the dorsal region.

PRÉ-DORSO-ATLOÏDIEN, *Longus colli*.

PRÉ-DORSO-CERVICAL, *Longus colli*.

PRE-EXISTENCE OF GERMS, see *Preformation*.

PREFORMA'TION, *Præformatio*, from *præ*, 'before,' and *formatio*, 'forming.' An ancient theory of generation, which maintained that the germ of every part exists, but not visibly, before fecundation, and that fecundation occasions their development. It was the doctrine of *pre-existence of germs*.

PREG'NANCY, *Pregnatio*, *Pregnatus*, *Impregnatio*, *Ingravidatio*, from *præ*, and *genere*, 'to beget'; *Hysterocye'sis*, *Graviditas*, *G. uterina*, *Eu'cyon*, *Cye'sis*, *Eucye'sis*, *Eu'cysis*, *Encymo'sis*, *Cyopho'ria*, *Gestatio*, *Fœtatio*, *Ingravidatio*, *Utero-gestatio*; (F.) *Grossesse*. The state of a female who has within her a fecundated germ, which gradually becomes developed in or out of the uterus. When placed in the cavity of the uterus,—by virtue of the vital principle which animates it, it becomes attached to the inner surface of that viscus, and obtains from it indirectly the nutriment necessary for its gradual development, during the space of ten lunar months or two hundred and eighty days—the ordinary *term* or *duration of gestation*, although the period may be protracted to 300 days or upwards,—at the expiration of which parturition occurs. Pregnancy is commonly limited to a single fœtus: at times, the uterus contains two or three.—See *Reproduction*. Occasionally, one or two moles—the degenerated products of conception—alone occupy the uterus:—at other times, a fœtus likewise exists. In certain cases, and owing to accidental circumstances, the fecundated ovum remains in the ovarium, or is arrested in the Fallopian tube. Pregnancy has, accordingly, been distinguished into, 1. *Natural* or *uterine*, when the fœtus is carried in the cavity of the womb; and, 2. *Preternatural* or *extra-uterine*, *Erfœtatio*, *Conceptio vitiosa*, *Paracye'sis* seu *Ecycye'sis*, *Echysterocye'sis*, *Echihysterocye'sis*, *Dysto'cia*, *Dyscye'sis* seu *Graviditas extra-uterina*, (F.) *Grossesse contre-nature*,—when it occurs in the ovarium, tube, or cavity of the abdomen.

During pregnancy, the female experiences remarkable changes in the condition of her organs and functions. The genital apparatus is, of course, most affected. The uterus changes its shape, situation, size, and texture. The usual signs of pregnancy are,—suppression of the catamenia: the abdomen gradually enlarging and forming a hard tumour; the umbilicus prominent in the latter months; the breasts increasing in size; the areola round the nipple becoming darker and the follicles more developed; kystein in the urine from an early period; a bluish tint of the vagina after the fourth week; about the fifth month, the evidence by *ballotement*, the placental *souffle*, and the sound of the foetal heart heard on auscultation; the motion of the fœtus perceptible, and the neck of the uterus diminishing in length at the advanced periods.

PREGNANCY, ABDOM'INAL, *Ecycye'sis abdominalis*, *Graviditas abdominalis*, (F.)

Grossesse abdominale. That in which the ovum has escaped into the cavity of the abdomen.

PREGNANCY, AFET'TAL, *Pregnancy without a fœtus*, (F.) *Grossesse afetale* ou *sans fœtus*. Pregnancy produced by a false or degenerate conception, or any other affection, which has given place to unusual development of the uterus. Madame Boivin includes in this class, *sarco-hysteria*, *hydro-hysteria*, *gazo-hysteria*, and *hemato-hysteria*.

PREGNANCY, BIGEM'INAL, (F.) *Grossesse bigemiale*, *Double pregnancy*. Pregnancy in which there are twins in utero.—Madame Boivin.

PREGNANCY, COMPLEX, (F.) *Grossesse complexe*, *G. inter-extra uterine*, is when the womb, besides a fœtus, contains much water, hydatids, a mole, &c.

PREGNANCY, COMPOUND, *Polybreph'ia*, *Polycye'sis*, *Polycye'sis*, (F.) *Grossesse composée*, is when there are two or three fœtuses in the cavity of the uterus.

PREGNANCY, DOUBLE. Pregnancy, bigeminal—p. Duration of, see *Pregnancy*—p. Extra-uterine, see *Pregnancy*.

PREGNANCY, FALSE, *Graviditas spuria*, (F.) *Grossesse fausse* ou *apparente*, *Pseudocye'sis*. Authors have thus called different affections of a more or less serious character, which resemble pregnancy; so as, at times, to impose upon the best-informed practitioners. These diseases may be seated in the uterus or its dependencies, in the intestines or some other part of the abdomen. They may be:—a mole, hydatids, water, blood, or polypus distending the parietes of the uterus;—chronic enlargement of the uterus; development of tumours, &c., in its substance;—scirrhus; dropsy of the ovaria; ascites; tympanitis, &c.

PREGNANCY, FETAL, (F.) *Grossesse fœtale*. Pregnancy in which there is a fœtus, or more than one, in the uterus or out of it. Madame Boivin puts in this class—the solitary, bigeminal, trigeminal, and sarco-fœtal pregnancies, which form the first order; and the tubal, ovarian, and abdominal, which constitute the second.

PREGNANCY, GAZO-HYSTER'IC, (F.) *Grossesse Gazo-hystérique*. Afœtal pregnancy, caused by development of air or gas in the cavity of the womb.—Madame Boivin.

PREGNANCY, HEMATO-HYSTER'IC, (F.) *Grossesse hémato-hystérique*. Afœtal pregnancy, caused by accumulation of blood in the uterus.—Madame Boivin.

PREGNANCY, HYDRO-HYSTER'IC, (F.) *Grossesse hydro-hystérique*. Afœtal pregnancy, occasioned by the secretion and accumulation of serous fluid in the uterus.

PREGNANCY, INTERSTIT'IAL, *Graviditas in uteri substantiâ seu interstitiâ*. The development of the embryo in the proper substance of the uterus. The case generally terminates fatally at an early period of gestation; the thin external paries of the uterus being ruptured, and the ovum passing into the abdominal cavity.

PREGNANCY, MORBID, *Paracye'sis*, *Dysto'cia*, *Dyscye'sis*. The progress of pregnancy disturbed or endangered by the supervision of general or local disorder.—Good.

PREGNANCY, OVA'RIAN, (F.) *Grossesse ovarienne* ou *de l'ovaire*, *Ecycye'sis ovaria*, *Oâriocye'sis*, *Oûcye'sis*, *Graviditas uteri seu extra-uterina in ovario*. Pregnancy in which the fœtus is developed in the interior of the ovary.

PREGNANCY, PLEA OF. A plea, which a woman, capitally convicted, may set up in arrest of execution until she is delivered. See *Jury of Matrons*, and *De ventre inspicendo*.

PREGNANCY, PRETEND'ED. The criminal act of a female, who, not having been delivered, presents an infant as having been born of her. (F.)

Supposition de part. (This term, in French, likewise includes the case of a female, who, having been delivered of a dead child, substitutes for it a living child which is not hers.)

PREGNANCY, SARCO-FŒTAL, (F.) *Grossesse sarco-fœtale.* Pregnancy in which there are one or more fœtuses, and one or more moles.—Madame Boivin.

PREGNANCY, SARCO-HYSTERIC, (F.) *Grossesse sarco-hystérique.* Afoetal pregnancy formed by moles, hydatids, or a false conception.—Madame Boivin.

PREGNANCY, SOLITARY, (F.) *Grossesse solitaire ou simple.* Pregnancy in which there is a single fœtus. — Madame Boivin.

PREGNANCY, TRIGEMINAL, (F.) *Grossesse trigéminalle ou triple.* Pregnancy in which there are three fœtuses in the cavity of the uterus.—Madame Boivin.

PREGNANCY, TUBAL, (F.) *Grossesse tubaire, Ecty'e'is tuba'lis, Graviditas tuba'ria.* Pregnancy where the fœtus remains and is developed in the Fallopian tube.

PREGNANCY, UTERINE, see Pregnancy.

PREGNANCY, UTERO-ABDOMINAL, (F.) *Grossesse utéro-abdominale.* Pregnancy consisting of two fœtuses; one in the uterus—the other in the abdominal cavity. — Madame Boivin.

PREGNANCY, UTERO-OVARIAL, (F.) *Grossesse utéro-ovarienne.* Pregnancy with two fœtuses, one in the womb, the other in the ovary.

PREGNANCY, UTERO-TUBAL, (F.) *Grossesse utéro-tubaire.* Pregnancy in which there is at the same time a fœtus in the womb and another in the Fallopian tube.

PREGNANT, Prægnans, Prægnas, Prægnax, Fetus, Partus, Encar'por, Gravid'a, Plena mul'ier, Incine'ta, Gravid, Breeding, (Old Eng.) Bealing. Same etymon. (F.) *Enceinte.* One in a state of pregnancy; — one with child.

PREHENSIO, Catalepsy, Epilepsy.

PREHENSION, Prehen'sio, from prehendere, prehensum, 'to lay hold of.' The act of laying hold of.

PREHENSION OF FOOD is the act of seizing and carrying it to the mouth, and introducing it into that cavity. It is, also, called *Assumptio, Prolept'is, Pros'phora.*

PRÊLE, Hippuris vulgaris.

PRÉLOMBAIRE, Prelumbar.

PRÉLOMBO-PUBIEN, Psoas parvus — p. Trochantin, Psoas magnus.

PRELUM, Press — p. Arteriale, Tourniquet.

PRELUMBAR, Prelumbar'is, (F.) *Prélom-baire, from præ, and lumbi, 'the loins.'* That which is placed before the loins. The prelumbar surface of the spinal column is the anterior surface of the lumbar portion.

PREMIÈRES VOIES, Primæ viæ.

PREMOLAR, see Molar teeth.

PREMONITORY, see Precursory.

PRENANTHES, Gall of the Earth, Dewitt Snakeroot, Lion's Foot. There are many species of this indigenous plant, whose root and milky juice are very bitter. They are used in popular practice in dysentery; and in the form of cataplasms to bites of serpents.

PRENANTHES ALBA, Nabalus albus — p. Serpentaria, Nabalus albus.

PRENDRE INSCRIPTION, see Matriculate.

PRÉPARATE (VEINE), see Facial vein.

PRÉPARATION, Prépara'tio, from præ, and parare, paratum, 'to get ready.' The act of making ready any substance whatever that has to be employed for any purpose.

Also, the product of any pharmaceutical operation; — *Prép'aratum.*

Any part of the human body, preserved for the uses of the anatomist, is so called.

PREPUCE, Præputium, Epagoge'ium, Posthæ Pella, Posthium, Putium, Foreskin. The prolongation of the integuments of the penis, which covers the glans. The prepuce is composed of two membranous layers: one external or cutaneous; the other internal or mucous, separated by areolar membrane. The mucous membrane lines the inner surface of the cutaneous layer till beyond the glans, when it is reflected over the latter; forming, behind the corona, a small *cul-de-sac* above the corpora cavernosa, which is interrupted by a triangular duplicature of the mucous membrane, called the *frænum*, fixed in the furrow at the inferior part of the glans, and terminating at a little distance from the orifice of the urethra. The clitoris is covered by a semilunar fold formed by a continuation of the skin of the labia. It is called *Præputium clitor'idis.*

PREPUTIAL, Præputia'lis. Belonging or relating to the prepuce; — as the 'preputial secretion, — or that which takes place from the lining of the prepuce covering the glans.

PRESBYODOCHIUM, Gerocomium.

PRESBYON'OSI, from presbus, 'old,' and vocas, 'disease.' Diseases of old age.

PRESBYOPE, Presby'ter, (F.) Presbyte, from presbus, 'old,' and ωπ, 'the eye.' One who is long-sighted or affected with presbyopia.

PRESBYOPIA, Presbytia.

PRESBYOPIC, Presbyo'picus, Presbyt'ic, Presby'ticus. Relating or belonging to presbyopia or long-sightedness.

PRESBYSPHACELUS, Gangrene of old people.

PRESBY'TIA, Presbyo'pia, Amblyo'pia proximo'rum, Visus sen'ilis, Pres'hypopy, (F.) Vue lanque; from presburns, 'an old person.' A very high degree is termed *Hyperpresby'tia.* A condition of vision, common in old persons, which consists in the circumstance of near objects being confusedly seen, whilst those at a greater distance are clearly distinguished. It is supposed to depend generally on diminution of the convexity of the cornea; whence results an alteration in the convergency of the rays, so that they form a focus behind the retina. This defect is remedied by the use of convex glasses, which augment the convergence of the luminous rays. It is, also, called *Long-sightedness, Far-sightedness, Parop'is longin'qua, Dyso'pia proximo'rum.*

PRESBYTIC, Presbyopic.

PRESCRIPTION, Præscriptio, Anagraphe, from præ, and scribere, scriptum, 'to write; For'mula, F. med'ica, Sched'ula, Præscriptum, Præscriptio med'ica, Recep'tum, (F.) Formulæ, Ordonnance. The formula which the physician writes for the composition of medicines adapted to any case of disease. A prescription should be as simple as possible, and should bear upon its face the evidence of the objects to be fulfilled by it. No article should form part of it, unless adapted for serving some useful purpose. A compound prescription has been divided into four parts: — the *basis* or principal ingredient of the prescription; the *ad'juvans*, or that which is designed to promote the action of the former; the *cor'rigens*, or that intended to correct its operation, or obviate any unpleasant symptom which it may be apt to produce; and the *constit'uens, excip'iens* or *vehic'ulum, Constituent, Excipient* or *ve'hicle*, the substance which gives to the other ingredients consistence or form. All these are seen in the following formula for cathartic pills:

Aloes.....	3jBasis.
Hyd. Chlorid. Mit. gr. x.		Adjuvans.
Olei Carui.....	gtt. v.	Corrigens.
Syrupi.....	q. s.	Constituens.
M. et fiant pilulæ xx. See Symbol.		

It is obvious, however, that most prescriptions are more simple than this. The basis, for example, may require neither adjuvant, corrigent, nor constituent.

Dr. Paris has given the following synopsis of the principles of medicinal combination, *Iamatoxiolog'ia*, *Iamatosyntaxiolog'ia*, *Ars formulæ medicæ concinnand'æ*, which may serve as an instructive guide to the prescriber:—

OBJECT I.

TO PROMOTE THE ACTION OF THE BASIS.

A. By combining the several different forms or preparations of the same substance. *B.* By combining the basis with substances which are of the same nature; i. e., which are individually capable of producing the same effects, with less energy than when in combination with each other. *C.* By combining the basis with substances of a different nature, and which do not exert any chemical influence upon it, but are found by experience, or inferred by analogy, to be capable of rendering the stomach or system more susceptible of its action.

OBJECT II.

TO CORRECT THE OPERATION OF THE BASIS BY OVIATING ANY UNPLEASANT EFFECTS IT MIGHT BE LIKELY TO OCCASION, AND WHICH WOULD PERVERT ITS INTENDED ACTION, AND DEFEAT THE OBJECT OF ITS EXHIBITION.

A. By chymically neutralizing or mechanically separating the offending ingredient. *B.* By adding some substance calculated to guard the stomach or system against its deleterious effects.

OBJECT III.

TO OBTAIN THE JOINT OPERATION OF TWO OR MORE MEDICINES.

A. By uniting those medicines which are calculated to produce the same ultimate results, but by modes of operation totally different. *B.* By combining medicines which have entirely different powers; and which are required to obviate different symptoms, or to answer different indications.

OBJECT IV.

TO OBTAIN A NEW AND ACTIVE REMEDY, NOT AFFORDED BY ANY SINGLE SUBSTANCE.

A. By combining medicines which excite different actions in the stomach and system, in consequence of which new or modified results are produced. *B.* By combining substances which have the property of acting chymically upon each other; the results of which are:—*a.* The formation of new compounds; *b.* The decomposition of the original ingredients, and the development of the more active elements. *C.* By combining substances, between which no other change is induced than a diminution or increase in the solubility of the principles in which their medicinal virtues reside. *a.* By the intervention of substances that act chymically. *b.* By the addition of ingredients whose operation is entirely mechanical.

OBJECT V.

TO AFFORD AN ELIGIBLE FORM.

A. By which the efficacy of the remedy is enhanced. *B.* By which its aspect or flavour is rendered more agreeable, or its mode of administration more convenient. *C.* By which it is preserved from the spontaneous decomposition to which it is liable.

[The vocabulary in the author's "Medical Student" Philad., 1844, will aid the student not

only in translating, but in writing his prescriptions *more solito*.]

PRESENTATION, from (Lat.) *præsens*, (*præ*, 'before,' and *ens*, *entis*, 'being,') The part of a factus which is felt *presenting*, on examination *per vaginam*. When the head presents, and especially the vertex, or the feet, knees, or breech, the presentation is said to be *natural*; when any other part, *preternatural*; and the labour is styled *perverse* or *preternatural*, *Parodynia* seu *Dyslochia perversa*, (*Cross-birth*, (F.) *Accouchement contre nature*. When any part besides the head, feet, knees, or breech presents, the operation of turning becomes necessary. See Parturition.

PRESERVATIVE, Prophylactic.

PRESIS, Swelling.

PRÊSLE, *Hippuris vulgaris*.

PRESMA, Swelling.

PRESPINAL, *Præspinalis*, from *præ*, and *spina*, 'the spine.' That which is situate before the spine. The prespinal surface of the vertebral column is the anterior surface.

PRESS, *Prælium*, *Prælium*, *Pressorium*, *Piester*, *Piesterion*, (F.) *Pressoir*, *Presse*, from *premere*, *pressum*, 'to squeeze.' An instrument for subjecting matters to considerable pressure, for the purpose of separating the liquid from the solid portion.

PRESS OF HEROPHILUS, *Torcular Herophili*.

PRESSE-ARTÈRE, *Serre-nœud de Deschamps*, *Serre-Artère*. An instrument invented by Deschamps for the immediate compression of arteries. It is composed of a plate six or seven lines long, and three broad, and of a rod or stem two inches long, placed perpendicularly above the plate and riveted. At each extremity of the plate is a hole, through which each of the ends of the ligature, to be placed upon the artery, is passed. The ligature is then to be tightened by drawing the ends, and fixing them to the perpendicular rod. This *presse-artère* or *serre-artère*, surrounded with charpie, remains in the wound, until there is no longer danger of hemorrhage. The ligature is then cut, and the instrument withdrawn. It is an unnecessary instrument.

PRESSE-URÈTHRE, *Jugum Penis*. A kind of forceps of elastic iron, the branches of which, padded, may be approximated or separated by means of a screw. The penis is introduced between the branches, one of them being applied to the urethra, and the screw is tightened so as to exercise a degree of pressure sufficient to bring the parietes of the canal in contact. This instrument is sometimes used in incontinence of urine, to prevent the constant discharge of that fluid.

PRESSIO, Pressure.

PRESSIO ABDOMINALE, Pressure, abdominal.

PRESSOIR, Press—*p. d'Hérophile*, *Torcular Herophili*.

PRESSORIUM, Press.

PRESSURE, *Pressio*, from *premere*, *pressum*, 'to press.' The action of pressing.

PRESSURE ABDOMINAL, (F.) *Pressio abdominalis*. A mode of examination by means of which the sensibility, size, suppleness, or hardness of any morbid abdominal viscus may be compared, in these respects, with its healthy condition. Bichat proposed, with the view of exploring the state of the thoracic organs, to exert a pressure from below upwards on the abdominal organs, so as to crowd them upon the diaphragm, and thus to diminish the dimensions of the cavity of the chest, noticing to what extent respiration was affected by this action. No great information can of course be derived from it.

PRESYSTOLIC, *Præystolicus*, from *præ*, 'before,' and *systole*. That which precedes the

contraction of the heart—as “*preystolic friction sound*.”

PRETENDED DISEASES, Feigned diseases.

PRETIBIAL, *Prætibialis*, from *præ*, and *tibia*, ‘the tibia.’ That which is situate before the tibia; as the *ilio-pretibial* and *ischio-pretibial* muscles.

PREVENTIVE, Prophylactic.

PREVENTIVE, ROYAL, from *prævenire*, *præventum*, (*præ*, and *venire*), ‘to come before.’ An empirical preparation, used as a prophylactic against the venereal virus. It is a solution of acetate of lead, — the *Liquor Plumbi Subacetatis dilutus*.

PREVERTEBRAL APONEUROSIS. The aponeurosis which covers the muscles of the prevertebral region.

PREVERTEBRAL ARTERY, (F.) *Artère prévertébrale ou méningienne postérieure*. A ramus of the meningeal branch, and sometimes even of the trunk, of the pharyngeal artery, which is distributed to the prevertebral region.

PREVERTEBRAL MUSCLES, see Cervical region, anterior — p. Region, Cervical region, anterior.

PRIAPEIA, *Nicotiana rustica*.

PRIAPIS/COS. Same etymon as the next. Name of a piece of wood which forms part of the *scamnum Hippocraticum*. Paulus of Ægina calls thus a tent of linen shaped like a penis.

PRIAPISMUS, *Tenti’go*, *T. Penis*, *Pri’apism*, *Horn-colic* (vulg.), *Enta’ria Priapis’mus*, *Rhopa’lismus*, *Satyr’ic’asis*, *Styma*, *Stypsis*, from *Πριαπός*, ‘Priapus.’ Constant and distressing erection, without any voluptuous idea or desire for venery. It is often symptomatic; and sometimes the immediate result of morbid causes, — of the action of cantharides, for example. The application of cold liquids to the penis, and the internal use of opium and camphor, are the means usually had recourse to for its removal.

PRIAPITIS, Phallitis.

PRIAPUS, *Mutunus*. In ancient mythology the deity who presided over gardens and the parts of generation, so called *propter deformitatem et membris virilis magnitudinem*.

PRIAPUS, Penis — p. Cervi, see Cervus — p. Ceti, Leviathan penis.

PRICKET, *Sedum acre*.

PRICKING, Sax. *prickean*, (D.) *pricken*, (F.) *Picotement*. A variety of pain, compared to that which pointed bodies would occasion.

PRIDE OF CHINA, *Melia azedarach* — p. of India, *Melia azedarach* — p. Tree, *Melia azedarach*.

PRIDEWEED, *Erigeron Canadense*.

PRIEST-PHYSICIANS, see *Asclepiades*.

PRIEST’S PINTLE, *Arum maculatum*.

PRIM, *Lignstrum vulgare*.

PRIMA CELLA COLI, Cæcum.

PRIME VLE, (F.) *Premières voies*, ‘The first passages.’ The stomach and intestinal canal: — the lacteals being the *secundæ viæ* or ‘second passages.’ See Ways, digestive.

PRIMARY CELL, see Cell.

PRIMEVÈRE, *Primula veris*.

PRIMP/ARA, *Primiparturiens*, (F.) *Primipare*, from *primus*, ‘first,’ and *parere*, ‘to bring forth.’ A name given to a female who brings forth for the first time.

PRIMP/ARTURIENS, *Primipara*.

PRIM/STERNAL, *Primi-sternalis*, from *primus*, ‘the first,’ and *sternum*, ‘the sternum.’ M. Bôlard has called *os primi-sternal* or *clavi-sternal*, the bony portion of the sternum.

PRIMIT/LÆ, from *primus*, ‘first.’ The first waters, or the waters discharged before the extrusion of the fœtus.

PRIMITIVE BAND, see Nerve fibres — p. Streak, Groove, primitive.

PRIMORDIAL CELL, see Cell.

PRIMORDIUM, *Arche*.

PRIMROSE EVENING, *Oenothera biennis* — p. Tree, *Oenothera biennis*.

PRIMULA OFFICINALIS, *P. veris* — p. *Variabilis*, *P. veris*.

PRIM/ULA VERIS, *P. officinalis* seu *variabilis*, from *primulus*, ‘first.’ So called, because it flowers in the beginning of the spring. *Verbas-culum*, *Paraly’sis*, *Herba paraly’sis* seu *paraly’tica*, the *Cowslip*, *Paigil* or *Peagle*, *Herb Peter*, *Palswort*, (Prov.) *Cow-striplings*, *Cow-stropple*, *Crowel*, (F.) *Primerère*, *Herbe à la paralysie*, *Fleur de coucou*, *Oreille d’ours*, *Brairette*; *Ord.* *Primulaceæ*. See *Syst. Pentandria Monogynia*. The flowers have been considered to be mildly tonic, antispasmodic, and anodyne.

PRIMULA VULGARIS. The leaves and roots of this plant have been used as sternutatories.

PRINCE ARTHUR’S FARINACEOUS FOOD, see Farinaceous.

PRINCEPS, *Rectum*.

PRINCEPS CERV/CIS (Artery) Cervical Artery. A branch of the occipital artery, which descends between the splenius and complexus muscles, and may be followed down to the lower part of the neck.

PRINCEPS POL/LICIS (Arteria), *Arte’ria magna pol’licis*. When the two collateral arteries of the thumb take origin from a common trunk — the trunk is the *Princeps pollicis*.

PRINCEPE CRYSTALLIZABLE DE DÉROSNE, *Narcotine* — p. de la *Mélanose*, see *Choroidea tunica*.

PRINCEPES IMMÉDIATS, *Principles*, immediate.

PRINCIPIUM, *Arche*, *Element* — p. *Acidifcans*, *Oxygen* — p. *Adstringens*, *Tannin* — p. *Byrsodepsicum*, *Tannin* — p. *Coriaceum*, *Tannin* — p. *Hydrogeneticum*, *Hydrogen* — p. *Hydroticum*, *Hydrogen* — p. *Oxygenans*, *Oxygen* — p. *Salivale*, see *Saliva* — p. *Scytodepsicum*, *Tannin* — p. *Urinorum*, *Urea*.

PRIN/CIPLE, *COL/OURING*, OF THE BLOOD. A name, given to the immediate principle in animals, to which the blood owes its red colour. It is solid, inodorous, and insipid. When recently separated from blood, it has a purple-red colour. When distilled, it furnishes, amongst other products, a purple oil. It is insoluble in water; but, if diluted with that liquid, acquires a vinous-red hue. It communicates a red colour to the albumen of the blood, and affords, on analysis, iron, to which the colour has been ascribed. See *Hæmaphæin*, and *Hæmatin*.

PRIN/CIPLE, *DIGESTIVE*, *Pepsin* — p. *Vital*, see *Vital principle*.

PRIN/CIPLES, *IMME/DIATE*, (F.) *Principes immédiats*, *Matériaux immédiats*. A name given to a considerable number of substances, composed of at least three elements, and which are obtained from animals and vegetables without alteration, by simple processes, and, in some measure, immediately.

PRIN/CIPLES, *IMMEDIATE*, OF ANIMALS, *Proximate principles*, *Organic Elements*, *Compounds of Organization*. These are divided into acid principles, fatty principles, and principles which are neither fat nor acid. The first include the *uric*, *rosacic*, *purpuric*, *allantoic*, *caseic*, *butyric*, &c. The fatty principles are: — *stearin*, *olein*, *cholesterin*, &c.; and the last division includes the most important compounds of organization: — *fibrin*, *gelatin*, *albumen*, *cæsin*, *osmazone*, &c.

PRIN/CIPLES, *PROXIMATE*, *P. immediate*.

PRINOS (of the ancients), *Ilex aquifolium*.

PRINOS, *P. verticillatus*, *Alcan’na major latifolia dentata*, *Aquifolium foliis deciduis*, *Prinos Gronovii* seu *padifolius*, *Black Alder*, *Vir-*

ginia or *Whorled Winterberry*, *Feverbush*, (F.) *Apalachine à feuilles de Prunier*. Ord. Aquifoliaceæ. *Ser. Syst. Hexandria Monogynia*. The bark of this common shrub of the United States — *Prinos*, (Ph. U. S.) — is astringent, bitter, and pungent. The berries are, likewise, bitter. The bark has been used as a substitute for the cinchona, in intermittents, and as a tonic.

PRINOS GLABER, see *P. lævigatus* — p. Gronovii, *Prinos*.

PRINOS LÆVIGATUS, *Smooth Winterberry*, and *PRINOS GLABER*, *Inkberry*. Indigenous; have the same properties as

PRINOS PADIFOLIUS, *Prinos*.

PRION, *Saw*.

PRIONODES, from *πριον*, 'a saw,' and *εidos*, 'shape.' *Serra*'ted. Applied to the sutures of the head.

PRIOR, 'first, foremost,' as

PRIOR ANNULARIS, *Interosseus quartus* of Winslow. An internal interosseous muscle which arises from the outside of the metacarpal bone of the ring-finger, and is inserted into the outside of the tendon on the back of the ring-finger. Its use is to draw the ring-finger outwards.

PRIOR INDICIS, *Extensor tertii interni dig'icis*, (F.) *Sous-métacarpo-latéri-phalangien*. It arises from the outer or radial side of the metacarpal bone of the fore-finger, and is inserted into the outside of the tendon on the back of the fore-finger. *Action*: — to draw the finger outwards, towards the thumb.

PRIOR INDICIS PEDIS, *Adductor Indicis pedis*. An external, interosseous muscle of the foot, which arises from the contiguous sides of the metatarsal bones of the great and fore toes, and is inserted into the inside of the root of the first bone of the fore-toe. *Action*: — to pull the fore-toe inwards.

PRIOR MEDII, *Prior medii digiti*.

PRIOR MEDII DIG'IMI, *Prior Medii*, *Second interosseus* of Douglas. An external, interosseous muscle of the hand, which arises from the corresponding sides of the metacarpal bones of the fore and middle fingers, and is inserted into the outside of the tendon on the back of the middle finger. *Action*: — to draw the middle finger outwards.

PRIOR MEDII DIGITI PEDIS, *Adductor medii digiti pedis*. An internal, interosseous muscle of the foot, which arises from the inside of the metatarsal bone of the middle toe, and is inserted into the inside of the root of the first bone of the middle toe. *Use*: — to pull the middle toe inwards, or towards the inner side of the foot.

PRIOR MIN'IMI DIGITI, *Adductor min'imi dig'iti*. An internal, interosseous muscle of the foot, which originates from the inside of the metatarsal bone of the little toe, and is inserted into the inside of the root of the first bone of the little toe. *Action*: — to pull the little toe inwards.

PRIOR TERTII DIGITI PEDIS, *Adductor tertii dig'iti pedis*. An internal, interosseous muscle of the foot, which arises from the inner and under part of the metatarsal bone of the third of the small toes, and is inserted into the inside of the root of the first bone of the third of the small toes. *Action*: — to pull the third of the small toes inwards.

PRISIS, *Brygmus*, *Trepanning*.

PRISMA, *Scobs*.

PRYSMUS, *Brygmus*, *Trepanning*.

PRIVATE PARTS, *Genital organs*.

PRIVET, *Ligustrum vulgare*.

PRIVITIES, *Genital organs*.

PRIVY, *Ligustrum vulgare* — p. *Members*, *Genital organs* — p. *Parts*, *Genital organs*.

PRO, *pro*, 'before.' A common prefix to words, as in *Procatartec*, *Prostate*, &c.

PROAGOREUSIS, *Prognosis*.

PROBANG, *Sponge-probang*, *Detrusorium ex bale'na*. Allied to *probe*. A long slender rod of whalebone, with a piece of sponge at its extremity, intended to push down extraneous bodies, arrested in the œsophagus, into the stomach. A similar instrument is called by J. L. Petit, *Repousseir d'arêtes*.

PROBARBIUM, *Geneias*.

PROBASCANIUM, *Amuletum*.

PROBASCANTIUM, *Amuletum*.

PROBE, *Catheterize*, *Specillum* — p. *Eyed*, see *Specillum*.

PROBING, see *Melosis*.

PROBOLE, *Processus*.

PROCARDIUM, *Scrobiculus cordis*.

PROCÉDÉ, *Process*.

PROCEDURE, *Procedé*.

PROCEPHALI, *Capitones*.

PRO CERUS NASI, *Pyramidalis nasi*.

PROCÈS, *Process* — p. *Ciliaires*, *Ciliary processes*.

PROCESS, *Apophysis*, *Processus*.

Proc'ess, *Ratio*, (F.) *Procedé*, from *procedere*, *processum*, 'to march forwards.' *Procedure*, — method of performing any operation, chemical, pharmaceutical, or surgical.

PROCESS, *Cuneiform*, *Basillary process* — p. *Modelling*, see *Modelling Process* — p. of *Rau*, see *Malleus* — p. *Vertical superior longitudinal*, *Falx cerebri*.

PROCESSUS. A *process*; same etymon; *Projectura*, *Prob'ole*, (F.) *Proclis*. An apophysis or eminence of a bone; see *Apophysis*. Also, any part which seems prolonged beyond others with which it is in connexion; as the *Ciliary processes*.

PROCESSUS ANCONÆUS, *Olecranon* — p. *Annularis*, *Pons Varolii* — p. *Ascendens*, see *Maxillary bone*, (superior) — p. *Belenoides*, see *Belenoid* — p. *Blumenbachii*, *Unciform process* — p. *Caudatus*, *Lobulus caudatus* — p. à *Cerebello ad medullam oblongatam*, *Corpora testiformia* — p. *Cerebelli ad pontem*, *Peduncles of the cerebellum* — p. *Cerebelli ad testes*, *Peduncles of the cerebellum* — p. *Cerebri lateralis*, *Cornu ammonis* — p. *Ciliares*, *Ciliary processes* — p. *Clavatus*, see *Funiculi graciles* — p. *Cochleariformis*, see *Tympanum* — p. *Cuneiformis ossis occipitis*; *Basillary process* — p. *Ensisiformes*, see *Sphenoid bone* — p. *Enteroides cerebri*, *Convulsions (cerebral)* — p. *Falciformis cerebelli*, *Falx cerebelli* — p. *Falciformis duræ matris*, *Falx cerebri* — p. *Frontalis*, see *Maxillary bone*, (superior) — p. *Gracilis* of *Rau*, see *Malleus* — p. *Laterales uteri*, *Tubæ Fallopiæ* — p. *Mamillares*, *Papillæ of the kidney*, *Olfactory nerves* — p. *Mamillares cerebri*, *Mamillary tubercles* — p. *Mastoideus*, *Mastoid process* — p. *Nasalis*, see *Maxillary bone*, (superior) — p. *Olivaris*, *Olivary Process* — p. *Orbicularis cerebri*, *Infundibulum of the brain* — p. *Papillares*, *Olfactory nerves* — p. *Papillarum*, *Olfactory nerves* — p. *Rachidianus*, *Medulla spinalis* — p. *Sterni xiphoides*, *Xiphoid*.

PROCESSUS TER'ETES, *Eminentia seu Fasciculi ter'etes*. Two slightly convex bodies forming the anterior wall or floor of the fourth ventricle of the brain. They are separated by a longitudinal groove, which is continuous, inferiorly, with the sulcus longitudinalis posterior of the spinal cord. They are crossed transversely by several white and gray fasciculi — *Lineæ transversæ, striæ medullares* — the origin of the auditory nerves.

PROCESSUS AD TESTES, see *Valvula Vieussenii* — p. *Transversus duræ matris*, *Tentorium* — p. *Uncinatus*, *Olecranon*, *Unciform process*, see *Coracoid process* — p. *Uvifer*, *Uvula* — p. *Ventriculi*, *Duodenum* — p. *Xiphoides*, *Xiphoid cartilage* —

p. Xipho-sternalis, Xiphoid cartilage—p. Zygomatikus, Zygomatic process.

PROCHEILA, Antelabia.

PROCHEILIDION, Procheilon.

PROCHEILON, *Procheilidion*, *Prolabium*, *Antelabium*, from *pro*, and *cheilos*, 'a lip.' The extreme projecting part or margin of the lips.

PROCHEUMA, Parenchyma.

PROCIDENCE DE L'ŒIL, Exophthalmia.

PROCIDENTIA, Prolapsus—p. Ani, Proctocoele—p. Intestini recti, Proctocoele—p. Iridis, Staphyloma of the Iris—p. Oculi, Exophthalmia—p. Sedis, Proctocoele—p. Uteri, Prolapsus uteri.

PROCNEME, Tibia.

PROCNEMIUM, Tibia.

PROCELIUS, *Procelius*, from *pro*, and *keilia*, 'belly.' One who has a large pendulous abdomen.

PROCELUS, Procelius.

PROCONDYLUS, *prokonduilos*, from *pro*, and *konduilos*, 'a knot.' The first joint of the fingers; the second being called *konduilos*; the third *metakonduilos*.

PROCREATIO, Generation—p. Sanguinis, Hæmatosis.

PROCREATION, Fecundation, Generation.

PROCTAGRA, from *proktos*, 'anus,' and *aggra*, 'seizure.' Gout in the rectum. Proctalgia.

PROCTALGIA, *Proctica simplex*, *Proctagra*, *Dolor ani*, *Proctodyn'ia*, from *proktos*, 'the anus,' and *algos*, 'pain.' Pain in the anus; generally symptomatic of disease, as of hemorrhoids, scirrhus, &c.

PROCTALGIA, Clunisia—p. Hæmorrhoidalis, Hæmorrhoids—p. Inflammatoria, Rectitis—p. Intertriginosa, Chafing—p. Rheumatica, Proctorrheuma.

PROCTATRESIA, *Imperforatio ani*, from *proktos*, 'the anus,' and *atresia*, 'imperforation.' Imperforate state of the anus.

PROCTICA, from *proktos*, 'the anus.' Pain or derangement about the anus, without primary inflammation. A genus in the class *Cœliaca*, order *Enterica*, of Good.

PROCTICA EXANIA, Proctocoele—p. Marisca, Hæmorrhoids—p. Simplex, Proctalgia—p. Tenesmus, Tenesmus.

PROCTISIS, Proctitis.

PROCTITIS, *Proctitis*, *Archi'tis*, *Clun'isia*, *Cyso'tis*, from *proktos*, 'the anus,' and *itis*, denoting inflammation. Inflammation of the anus and rectum. Rectitis.

PROCTITIS GANGRENOSEA, Proctocœce.

PROCTOCÆCE, *Proctitis gangrænosa*, from *proktos*, 'the anus,' and *kakos*, 'evil.' A disease of the rectum, so called by Fuchs, which, according to him, is common in Peru, in the neighbourhood of Quito and Lima, on the Honduras and Mosquito coasts, in Brazil, and on the Gold coast, &c. It is called by the Portuguese, *Bicho*, and *Bicho di Oulo*; by the people of Quito, *Mal del Valle*, from its prevalence in the valleys; and in Africa, *Bitios de Kis*. It is an adynamic, inflammatory condition, frequently ending in gangrene. It has been attributed to bad food, and the use of spices.

PROCTOCELE, from *proktos*, 'the anus,' and *kele*, 'hernia.' *Prolapsus Ani*, *Proctopto'ma*, *Procto'tosis*, *Procto'sis*, *Hedroce'te*, *Archeopto'ma*, *Archeopto'ma*, *Archopto'ma*, *Archeoce'te*, *Archeopto'sis*, *Ecd'nia*, *Hæmorrhoidis procdens* seu *ab ecd'niâ*, *Procdentia Ani* seu *intesti'ni recti* seu *sedis*, *Ecto'pia Ani*, *Sedes procd'ua*, *Falling down of the Fundament*, *Coming down of the Body*, (F.) *Chute du rectum*, *C. du Fondement*, *Renverssement du rectum*. Inversion and prolapse of the mucous coat of the rectum, from relaxation of the sphincter, with more or less swelling. In the treatment, the application of cold water and

astringent lotions must be recommended: and if other means fail, one of the radii of skin, which converge at the anus, may be snipped off. On cicatrizing, the verge of the anus will be contracted, and the cure probably complete. In the way of palliation, the parts may be retained by an appropriate bandage.

PROCTOCYSTOTOMIA, see Lithotomy.

PROCTODYNIA, Proctalgia.

PROCTONCUS, from *proktos*, 'anus,' and *oikos*, 'swelling.' Swelling of the anus.

PROCTOPARALYSIS, *Proctople'gia*, *Paralysis Intesti'ni Recti*, from *proktos*, 'the anus,' and *paral'ysis*, 'paralysis.' Paralysis of the muscles of the rectum.

PROCTOPLEGIA, Proctoparalysis.

PROCTOPTOMA, Proctocoele.

PROCTOPTOSIS, Proctocoele.

PROCTORRHAGIA, Hæmorrhoidal flux.

PROCTORRHEUMA, from *proktos*, 'anus,' and *roma*, 'defluxion.' Rheumatism of the anus.

PROCTORRHEA, from *proktos*, 'the anus,' and *rho*, 'I flow.' A discharge of blood or slime from the anus. See Hæmorrhoids.

PROCTOS, Anus.

PROCTOSIS, Proctocoele.

PROCTOSPASMUS, Tenesmus.

PROCTOSTENOSIS ORGANICA, Stricture of the rectum.

PROCTOTOREUSIS, from *proktos*, 'anus,' and *toruere*, 'to perforate.' Perforation of a closed anus.

PROCTUS, Anus.

PROCURATIO ABORTÛS. The promotion of abortion.

PRODIAGNOSIS, from *pro*, and *diagnosis*. Prognostication of a disease to come,—a diagnosis by anticipation.

PRODROMIC, *Prodrom'icus*, (F.) *Prodromique*. Same etymon as the next. Relating, or belonging, to the prodromus of a disease.

PRODROMUS, *Propath'ia*, from *pro*, and *drōmos*, 'course,' (F.) *Prodrome*. *Stadium opportunitatis* seu *prodromo'rum*; in contagious diseases, *Stadium latentis contagii*. The period immediately preceding an attack of disease; in which the *precursory signs* occur.

PRODUCT, from *producere*, *productum*, (*pro*, and *ducere*, 'to lead,') 'to bring forth.' (F.) *Produit*. The result sought to be attained by any pharmaceutical operation.

PRODUCTIO, Prolapsus—p. Uvulæ à pituitâ, Staphylœdema.

PRODUCTION, *Produce'tio*; same etymon. This word is used, especially by the French, synonymously with *prolongation*. The mesentery is, in this sense, a production of the peritoneum.

PRODUCTION ACCIDENTELLE. An accidental or adventitious structure.

PRODUCTIVITAS, Fecundity.

PRODUIT, Product.

PROGUMENÆ CAUSÆ, Causes, predisponent.

PRÆLIUM, Press, Tourniquet.

PROEMINENT, *Proem'inens*, from *pro*, and *eminere*, 'to project.' Some anatomists call the 7th cervical vertebra the *prominent vertebra*, on account of the length of its spinous process, which passes the level of that of the neighbouring vertebrae.

PROENCEPHALUS, (F.) *Proencephale*; from *pro*, and *encephalos*, 'encephalon.' A monster, whose encephalon is situated, in a great measure, out of, and in advance of, the cranium, which is open in the frontal region.

PRÆOTIA, *Proïotes*, *πρωϊότης* or *προιότης*, from *πρωι*, 'early.' 'Precocity.' *Præcox matur'ritus*, *Præco'cious matur'ity*. Premature development of sexual organization or power.—Good.

PROFLUVII CORTEX, *Nerium antidysentericum*.

PROFLUVIUM, *Discharge, Fluxus, Polyrhœa, Flux*; from *profluo*, (*pro*, and *fluo*,) 'I flow forth.' A term under which—in the plural, *Profluvia*, *Polyrino'es*—some nosologists have comprised all morbid discharges or fluxes:—others, increased excretions attended by fever.

PROFLUVIUM ALVI, *Diarrhœa*—p. *Genitale muliebri*, *Menses*—p. *Mucosum urethræ*, *Gonorrhœa*—p. *Muliebri*, *Leucorrhœa*, *Menses*—p. *Sanguinis*, *Hæmorrhagia*—p. *Sanguinis à Renibus*, *Nephorrhagia*—p. *Sanguinis ex Ore*, *Stomatorrhagia*—p. *Sanguinis ex Utero*, *Metrorrhagia*—p. *Seminis*, *Pollution*—p. *Ventris*, *Diarrhœa*.

PROFOND, *Profundus*.

PROFUNDUS, (*pro*, and *fundus*, 'bottom,') *Altus*, 'having a deep base.' (F.) *Profond*. A name given to different parts, which are seated profoundly as regards others.

PROFUNDA ARTERIA FEMORIS, *A. vasta posterior seu muscularis femoris, Artère-grande musculaire de la cuisse*. (Ch.) This large branch arises commonly from the posterior part of the trural, between the pubis and lesser trochanter. It descends deeply, situates before the adductors; passes through the third adductor above the opening in it for the trunk of the femoral; and terminates in the short portion of the biceps. The *muscularis profunda* gives off, 1. The *external circumflex*; 2. The *internal circumflex*; and, 3. The three *perforantes*.

PROFUNDA ARTERIA HUMERI, *P. superior, Arteria spiralis*, (F.) *Artère humérale profonde, A. collatérale externe, Grand musculaire du bras*, (Ch.) Its origin is variable. When it arises from the brachial artery, it is given off opposite the groove of the humerus destined for the radial nerve, and descends backwards between the three portions of the triceps, accompanied by the nerve; giving branches to that muscle and to the humerus. When it reaches the posterior part of the bone, it divides into two branches, which are distributed to the triceps, supinator longus, &c.

PROFUNDA ARTERIA INFERIOR seu MINOR is often a branch of the last; but, commonly, is given off from the brachial, near the middle of the arm. It gives off branches to the muscles, &c., about the inner side of the os humeri. It is, also, called *large communicating ulnar or profundoulnar*.

PROFUNDA ARTERIA PENIS, *Chaussier* has given this name to the *cavernous artery*, which arises from the artery of the penis, furnished by the internal pudic.

PROFUNDA ARTERIA SUPERIOR, *Profunda humeri*.

Certain muscles are distinguished by the names *profound* or *deep-seated*, and *superficial*. Thus, we speak of the *superficial* and *deep-seated* muscles of the neck, &c.; the *flexor profundus perforans*, &c.

PROFUSIO, *Hæmorrhagia*.

PROFUSIO AQUÆ. The discharge or breaking of the waters at birth.

PROFUSIO SANGUINIS, *Hæmorrhagia*—p. *Seminis*, *Ejaculation*—p. *Subcutanea*, *Purpura simplex*.

PROGASTOR, from *pro*, and *γαστήρ*, 'belly.' One who has a pendulous belly.

PROGENIES, *Epigone*.

PROGENY, *Epigone*.

PROGLOSIS, from *pro*, and *γλῶσσα*, 'the tongue.' *Apex linguae*. The extremity or tip of the tongue.

PROGNA'THOS, from *pro*, and *γναθος*, 'the jaw.' Having a projecting jaw. A term applied to the form of the head in which there is a pro-

longation or forward extension of the jaws, as in the negro.

PROGNOSIS, *Proëgore'sis, Prorrhœ'sis, Præcognit'io, Prognosticè, Præno'tio, Prænum'tia'tio, Præscien'tia, Præscit'io, Pronœ'a, Prop'h'asis, Prædict'io*, (F.) *Prognostic, Pronostic*, from *pro*, and *γινωσκω*, 'knowledge.' A judgment formed by the physician regarding the future progress and termination of any disease.

PROGNOSIS, GENERAL. The opinion formed of a disease in the abstract. Thus, the *General prognosis of Cyanche tonsillaris* is favourable;—of *Phthisis pulmonalis*, unfavourable, &c.

PROGNOSIS, PARTICULAR. The opinion formed of any particular case of disease. The *particular prognosis* of one case of typhus, for example, may be favourable, whilst that of another may be unfavourable.

PROGNOSIS EX LINGUÀ, *Glossomania*.

PROGNOSTIC, *Prognosis*.

PROGNOSTICE, *Prognosis*.

PROGRESSIO, *Augmentation*.

PROGRESSION, *Walking*.

PROGRESSUS, *Augmentation*.

PROIOTES, *Prætia*.

PROJECTURA, *Process*.

PROLABIUM, *Procheilon*.

PROLAPSUS, from *prolabor*, *prolapsus*, (*pro*, and *labor*, *lapsus*, 'to slide,') 'I slip down.' *Prociden'tia*, *Delap'sio*, *Delap'sus*, *Hypor'hysia*, *Propto'ma*, *Propto'sis*, *Productio*, *Propenden'tia*, *Prominen'tia*, *Ptosis*, *Cusus*, (F.) *Chute; a protrusion; a falling down*. A genus of diseases in the class *Locales* and order *Ectopiæ* of Cullen, distinguished by the falling down of a part through the orifice with which it is naturally connected.

PROLAPSUS ANI, *Proctocele*—p. *Bulbi oculi*, *Exophthalmia*—p. *Corneæ*, *Staphyloma* of the cornea—p. *Iridis*, *Ptosis iridis*—p. *Lingue*, *Glossocèle*, *Paraglossa*—p. *Œsophagi*, *Pharyngocele*—p. *Palpebræ*, *Blepharoptosis*—p. *Pharyngis*, *Pharyngocele*—p. *Umbilici*, *Exomphalus*.

PROLAPSUS UTERI, *Exometra, Metropto'sis, Metropropto'sis, Orthysteropto'sis, Orthysteroptoma*, *Falling down of the womb*, *Edopto'sis* seu *Prociden'tia uteri*, *Hysteropto'sis*, *Hysterocele nuda*, (F.) *Abaissement ou Chute ou Descente ou Précipitation de la matrice*. Some use the term '*Procidencia Uteri*' for a minor degree of the affection than *prolapsus*;—*Relaxatio uteri* expressing the slightest state of all. A falling down of the uterus, owing to relaxation of the parts about the utero-vaginal region. In the treatment, the horizontal posture must be insisted upon; the injection of astringent substances *per vaginam* be advised; and if the affection be not remedied by these means, a pessary may be employed as a palliative, or the operation of *Episiorrhaphy* be performed.

PROLAPSUS UVULÆ, *Staphylædema*.

PROLAPSUS VAGINÆ, *Hysteropto'sis* seu *Prolapsus Edoptosis Vaginae*, *Coleopto'sis*, *Colpop'to'sis*, *Elythropto'sis*—*Protrusion of the upper part of the vagina into the lower*. This, like the descent of the uterus, may be *relaxation, procidencia*, *prolapsus*, or complete *inversion*.

PROLAPSUS VESICÆ, *Exocyste*.

PROLECTA'TIO; from *pro*, and *legere*, *lectum*, 'to gather.' Action of separating the finer parts of a body from the grosser.

PROLEPTICS, *Prolepticè*, from *pro*, and *λαμβάνω*, 'I seize hold of.' A term proposed by Dr. Laycock to signify the art and science of predicting in medicine.

PROLEPTICUS, *Anticipating*.

PROLES, *Epigone*.

PROLIDICE, *Prolidicium*, from *proles* (*pro*, and *aliscere*, 'to grow'), 'offspring,' and *cadere*, 'to kill.' Destruction of offspring: a term which includes foeticide as well as infanticide.

PROLIFIC, *Prolificus*, (F.) *Prolifique*, from *proles*, 'offspring,' and *facio*, 'I make.' That which has the faculty of engendering. Applied to man, and animals, and to their sperm or seed; as well as to remedies, which, by strengthening the genital organs, increase the secretion of sperm.

PROLIG'ERUS, *Pro'liger*, *Prolig'erus*, from *proles*, 'offspring,' and *gero*, 'I carry.' That which is connected with carrying the offspring.

PROLIGERUS DISC or **LAYER**, *Discus prolig'erus* seu *vitellinus*, *Stratum prolig'erum*, *Gonotectima*, (F.) *Disque proligrer*. A granular layer, situate generally towards the most prominent part of the ovarian vesicle, in the centre of which the true ovum or ovule exists.—Von Baer.

PROLONGEMENT RACHIDIE, *Medulla spinalis*—p. *Sinus occipital*, Basillary process.

PROMALACTERIUM, from *pro*, and *palasaw*, 'I soften.' The room, in the ancient gymnasium, in which the body was softened and anointed. One of the operations in bathing.

PROMANUS, see *Digitus*, *Pollex*.

PROMETOPIDIA, Frontal bandages.

PROMETOPIS, *Prometopid'ion*, from *pro*, and *μετωπον*, 'the forehead.' The skin of the forehead.

PROMINENTIA, *Prolapsus*, *Protuberance*—p. *Annularis Cerebri*, *Pons Varolii*—p. *Corneæ*, *Ceratoclele*—p. *Ossis Continua*, *Apophysis*.

PROMINENTIA ALBICANTES, *Mammillary tubercles*—p. *Semiövales Medullæ Oblongatæ*, *Corpora olivaria*.

PROMONTIOIRE, *Promontorium*.

PROMONTORIUM, *Promontory*—p. *Faciei*, *Nasus*—p. *Ossis Sacri*, see *Sacrum*.

PROMONTORY, *Promont'rium*, (F.) *Promontioire* (from *pro* and *mons*, *montis*, 'a mountain,') seu *Tuber seu Tuberositas Tympani*. A small projection at the inner paries of the cavity of the tympanum, which corresponds to the external seal of the cochlea, and especially to the outer side of the vestibule.

PROMONTORY OF THE SACRUM, see *Sacrum*.

PRONATEUR CARRÉ, *Pronator radii quadratus*—p. *Grand* ou *roncl*, *Pronator radii teres*—p. *Petit*, *Pronator radii quadratus*.

PRONATION, *Pronatio*, from *pronus*, 'inclined forwards.' Anatomists understand, by pronation, the motion by which the inferior extremity of the radius passes before the ulna, and thus causes the hand to execute a kind of rotation from without inwards.

PRONATOR. That which produces the motion of pronation. This name has been given to two of the muscles of the forearm.

PRONATOR OBLIQUES, *P. radii teres*—p. *Quadratus*, *P. radii quadratus*.

PRONATOR RADII QUADRATUS, *P. quadratus* seu *transversus* (Winslow) seu *radii brevis* seu *radii quadratus* (Cowper), *Quadratus rad'ii*, (F.) *Cubito-radial* (Ch.), *Petit* ou *carré pronateur*. This muscle is situate at the anterior, inferior, and profound part of the forearm. It is flat and square, and is attached, within, to the inferior quarter of the anterior surface of the ulna; and, without, to the inferior quarter of the anterior surface of the radius. It produces the motion of pronation.

PRONATOR RADII TERES, *P. teres* sive *obli'quus* (Winslow), (F.) *Épitrochlo-radial* (Ch.), *Grand* ou *roncl pronateur*. A muscle seated at the upper and anterior part of the forearm. It is long, flat, and larger above than below. It is attached, above, to the inner tuberosity of the humerus and

to the coronoid process of the ulna. Thence it passes obliquely downwards and outwards, and terminates at the middle of the outer surface of the radius. It causes the radius to turn on the ulna, to produce the motion of pronation. It can, also, bend the forearm on the arm, and conversely.

PRONATOR TERES, *P. radii teres*.

PRONAUS, *Vestibulum*.

PRONERVATIO, *Aponeurosis*, *Tendon*.

PRONŒA, *Prognosis*.

PRONONCIATION, *Pronunciation*.

PRONOSTIC, *Prognosis*.

PRONUNCIATION, *Pronuncia'tio*, (F.) *Pronunciation*, from *pro* and *nunciare*, 'to tell.' The mode of articulating or giving accent or utterance to words.

PROPATHIA, *Prodromus*.

PROPENDING, *Prolapsus*.

PROPERTIES, PHYSICAL, OF THE TISSUES. These are *flexibility*, *extensibility*, *elasticity*, *racornissement* and *imbibition*.

PROPERTIES, VITAL. Those which depend upon the vital force, as *contractility*.

PROPHASIS, *Prognosis*.

PROPHYLACTIC, *Prophylacticus*, *Diaphylactic*, *Diasostic*, *Syntereticus*, *Præservatorius*, (F.) *Prophylactique*, from *pro*, and *φύλασσω*, 'I defend.' A preservative or preventive.

PROPHYLACTIC MEDICINE, *Hygiene*.

PROPHYLAX'IS, *Prophyl'acé*, *Prophylact'icé*, *Syntere'sis*. Same etymon. *Præservatio*, *Præcautio*, *Præcustodia*. Preservative or preventive treatment.

PROPOLIS, from *pro*, and *πολις*, 'city,' literally, 'that which is before the city.' *Ceranthemus*, *Commo'sis*, *Bee-bread*. A red, resinous matter, with which the bees cover the bottom of the hive. Its fume has been esteemed antiasmatic.

PROPOMA, from *pro*, and *πομα*, 'a drink.' Ancient name of a medicine, composed of several parts of honey and four of wine. It was drunk before a meal.—Paulus of Ægina.

PROPORTIO, *Symmetry*.

PROPOSITUM, *Intention*.

PROTOTISMA, *Mistura*.

PROPTISMUS, *προποτισμος*. Same etymon. *Prapo'tio*. An ancient name for medicines which were given to the patients before purging them, and, in some sort, to prepare them.

PROPRIUS AURIUS EXTERNÆ, *Retrahens auris*.

PROPTOMA, *Prolapsus*.

PROPTOMA AURICULÆ RUM, *Paræsthe'sis auditus flacca*, *Flap Ear*. Lobe of the ear broad, loose, and pendent from birth.—Good.

PROPTOMA SCROTII, *Rachosis*.

PROPTOSIS, *Prolapsus*—p. *Palpebræ*, *Blepharoptosis*—p. *Uvulæ*, *Staphyloedema*.

PROPTYSIS, *Expectoration*.

PRORA, *Occiput*.

PRORRHESIS, *Prognosis*.

PRORUPTIO SANGUINIS, *Hæmorrhagia*.

PROS, *pros*. In composition, this prefix generally signifies 'in addition to,' 'over and above,' 'besides.'

PROSARTHROSIS, *Diarthrosis*.

PROSBOLE, *Impression*.

PROSCHYSIS, *Affusion*.

PROSCOLLESI, *Adherence*, *Agglutination*.

PROSECTEUR, *Dissector*.

PROSECTIO, *Anatomy*.

PROSECTOR, *Dissector*.

PROSLEPSIS, *Prehension*.

PROSODOS, *Aditus*.

PROSOPALGIA, *Neuralgia*, *facial*.

PROSOPANTRIA, *Frontal sinuses*.

PROSOPANTRITIS, from *prosopatra*, the

frontal sinuses, and *itis*, denoting inflammation. Inflammation of the frontal sinuses.

PROSOPARALYSIS, Palsy, Bell's.

PROSOPIS DULCIS, *Ord. Leguminosæ*. A tree of rather large size, native of Peru. The fruit — *Pacay* — is a pod, from 20 to 24 inches long, containing black seeds imbedded in a white, soft, flaky substance; which last is eaten by the Limenos with pleasure. — Tschudi.

PROSOPODYNIA, Neuralgia, facial.

PROSOPODYSMORPHIA, from *προσωπον*, 'the face,' *δυσ*, 'with difficulty,' and *μορφή*, 'shape,' Malformation or atrophy of the face, — as where the process of nutrition suffers from diminished nervous influence.

PROSOPOLOGIA, Physiognomy.

PROSOPOMANTIA, Physiognomy.

PROSOPON, Face.

PROSOPORRHEUMA, *Rheumatismus faciei*, from *προσωπον*, 'face,' and *ρρευμα*, 'diffusion,' 'rheumatism.' Rheumatism affecting the face.

PROSOPOSIS, Physiognomy.

PROSOPSPASMUS, Canine Laugh.

PROSOPOSTERNODYMIA, see Cephalosomatodymia.

PROSPERITAS VALETUDINIS, Sanitas.

PROSPHORA, Prehension.

PROSPHYSIS, *Adnascen'tia*, *Coalit'io*, 'adhesion, adherence, connexion,' from *pros*, and *φύω*, 'I grow.' In a more limited sense, this word means morbid adhesion of the eyelids, either between themselves, or with the globe of the eye; *Ankylobleph'aron*.

PROSTASIS, *προστασις*, from *προ*, and *στασις*, 'station,' 'superiority, preponderance,' same etymon as *Prostate*. A predominance or abundance of excrementitious humour. — Hippocrates.

PROSTATA, Prostate — *p. Adenoides*, Prostate — *p. Bartholini*, Cowper's glands in the female — *p. Glandula*, Prostate — *p. Muliebris*, Cowper's glands in the female.

PROSTATÆ, Prostate — *p. Adenoides*, Prostate — *p. Glandulosa*, Prostate.

PROSTATAL, Prostatic.

PROSTATAL'GIA, from *προσπαρα*, 'the prostate,' and *αλγος*, 'pain.' Pain in the prostate.

PROSTATAUXE, Prostatoparectasis.

PROSTATE, *προσπαρα*, from *προ*, 'before,' and *στημι*, 'I stand.' — *Prostata*, *P. glandula*, *Prostata*, *P. glandulosa* seu *adenoides*, *Paras'tata* *adenoides* seu *glandulosa*, *Adstans*, *Paras'tata*, *P. gland'ula*, *Testis minor*, *Corpus glandulosum* seu *glandulosum* seu *adeniforme* seu *glandiforme* seu *glandulosum* seu *adenoides*, *Assistentes glandulae*, *Adstites glandulosi*; the Prostate gland. A glandular, cordiform body, of the size of a chestnut, situate before the neck of the bladder, behind the symphysis pubis, and surrounding the first portion of the urethra. Its excretory outlets, to the number of 10 or 12, open into the part of the urethra that traverses it, and pour into it a whitish viscid humour — *Liquor prostat'icus* — intended to lubricate the interior of the urethra, and to serve as a vehicle for the sperm in its ejection.

PROSTATE, ANTERIOR, see Cowper's glands.

PROSTATELCO'SIS, from *προσπαρα*, 'prostate,' and *ελκος*, 'an ulcer.' Ulceration of the prostate.

PROSTATES INFÉRIEURES, Cowper's glands — *p. Petites*, Cowper's glands.

PROSTAT'IC, *Prostat'icus*, *Prostatal*, (F.) *Prostatique*. That which belongs or relates to the prostate. Winslow has named — *Superior Prostatic Muscles* — ligaments which pass from the pubis to the lateral parts of the prostate. He, also, calls — *Inferior prostatic muscles* — the fleshy fibres, which, from the lateral parts of the urethra,

near its membranous portion, pass to be inserted into the pubis.

PROSTATIC BISECTOR. An instrument employed by Dr. Stevens, of New York, in the bilateral operation of lithotomy. In form it resembles a large olive, with a beak at the extremity, cutting edges at the sides parallel to its longest axis, and with a straight handle.

PROSTATIC LIQUOR, *Liquor prostat'icus*. The fluid secreted by the prostate, which mixes with sperm during emission.

PROSTATIC PORTION OF THE URETHRA. The part of the canal of the urethra included by the prostate: it is about 15 lines long.

PROSTATIC SINUS, see Sinus, prostatic.

PROSTATICUS SUPERIOR, Compressor prostate.

PROSTATIQUE, Prostatic.

PROSTATITIS, *Inflammat'io Prostatae*, from *προσπαρα*, 'the prostate,' and *itis*, denoting inflammation. Inflammation of the prostate; when accompanying gonorrhoea, it is the *gonorrhoea prostat'itis*.

PROSTATOCELE, Prostatocencus.

PROSTAT'ICUS, *Prostatocel'e*, *Tumor prostatae*, from *προσπαρα*, 'the prostate,' and *ογκος*, 'a tumour.' Swelling of the prostate.

PROSTATOPARECTASIS, from *προσπαρα*, 'prostate,' and *παρεκτείνω*, 'to extend.' Enlargement of the prostate; also, induration of the same, *Prostatoscirrhus*, *Prostataur'e*.

PROSTATOSCIRRHUS, see Prostatoparectasis.

PROSTERNID'IUM, from *προ*, 'before,' and *στερνον*, 'the sternum.' Any agent applied to the anterior part of the chest; a plaster to the breast.

PROSTHESIS, Prothesis.

PROSTHETA, Subdita.

PROSTHIUM, Penis.

PROSTOM'IA, *Commissu'ra labio'rum*; from *προ*, 'before,' and *στομα*, 'mouth.' The commissure of the lips.

PROSTRATION, *Prostratio vi'rium*, (F.) *Prostration des forces*, *Abatement*. Great depression of strength. Almost total loss of power over the muscles of locomotion.

PROTEA MELIF'ERA. A South African plant, *Nat. Ord. Proteaceæ*, whose involucra and flowers, at the time of inflorescence, are filled with a sweet watery liquid, which contains a great deal of honey. By inspissation it forms a delicious syrup — *Syr'upus Pro'teæ*, which is much used in pulmonary affections.

PROTEA LEPIDOCARPON, and some other Proteæ, supply the same kind of juice.

PROTEAN, *Pro'teiform*, *Proteiform'is*, from *Proteus*, who could assume various shapes. Assuming different shapes.

PROTEIFORMIS, Protean.

PROTEIFORM'IS MORBUS. A disease which assumes various characters.

PROTEIN, *Pro'teine*, *Protei'na*, from *πρωτεινα*, 'I take first rank.' A product of the decomposition of albumen, &c., by potassa. When animal albumen, fibrin or casein is dissolved in a moderately strong solution of caustic potassa, and the solution is exposed for some time to a high temperature, these substances are decomposed. The addition of acetic acid causes, in all three, the separation of a gelatinous translucent precipitate which has exactly the same characters and composition. This is protein. — Mulder.

Protein is the basis of animal and vegetable fibrin, albumen, and casein, and of gluten, which have been, consequently, classed under the "proteinaceous alimentary principle," by Dr. Pereira.

PROTEINACEOUS, Proteinous.

PROTEINOUS, *Proteina'ceous*, *Proteino'sus*,

(F.) *Proteineux*. Of, or belonging to protein; — as "a proteinous alimentary principle."

PROTEINEUX, Proteinous.

PROTENSIONES GLANDULARES, Mamillary eminences.

PROTHESIS, *Prosthesis*, *Adjunctio*, *Adjectio*, *Appositio*, 'addition, application;' from *προς*, 'in addition to,' and *τιθεμι*, 'I put.' That part of surgery whose object is to add to the human body some artificial part, in place of one that may be wanting, as a wooden leg, an artificial eye, &c.

PROTOMESIS, Umbilicus, Kidney.

PROTOAZOTATE DE MERCURE, Hydrargyri nitras.

PROTOGALA, Colostrum.

PROTOIATROS, Archiater.

PROTOMEDICUS, Archiater.

PROTOMUS, Dissector.

PROTOPATHIA, Idiopathia.

PROTOPHYTE, *Protophyton*, from *πρωτος*, 'first,' and *φυτον*, 'a plant.' A vegetable production at the foot of the scale; as a cryptogamous plant, — a fungus.

PROTOPLAST, from *πρωτος*, 'first,' and *πλασσω*, 'I form.' A primary formation. A first formed organized body — the parent of all others.

PROTOSPOROS, Os uteri.

PROTOZÖON, from *πρωτος*, 'first,' and *ζων*, 'an animal.' An animal lowest in the scale; — as a monad.

PROTRAHENS LENTIS, Ciliary muscle.

PROTRUSION, Prolapsus.

PROTUBERANCE, *Protuberantia*, *Extuberantia*, *Prominentia*, *Excrescentia*, *Extuberantia*, from *pro*, 'before,' and *tuber*, 'a projection.' (F.) *Dosse*. A name given to rough, unequal protuberances, seen at the surface of certain organs; the parietal protuberances, occipital protuberances, annular protuberances, &c.

PROTUBERANCE, ANNULAR, Pons Varolii — p. Cerebral, Medulla oblongata, Pons Varolii — p. *Cylindroide*, Cornu ammonis.

PROTUBERANTIA, Eminence — p. Annularis Willisii, Pons Varolii — p. Basilaris, Pons Varolii — p. Cylindrica, Cornu ammonis.

PROTRUSOR, Detrusor urinæ.

PROULMATESIS. A word employed by Forestus for hernia of the stomach [?].

PROUNE, Prunus domestica.

PROVENTRICULUS, *Bulbus glandulosus*, *Infundibulum*, *Secund stomach*; from *προς*, 'before,' and *ventriculus*, 'the stomach.' A bulbous expansion at the termination of the oesophagus, immediately above the gizzard of birds, the walls of which are thickly studded with a layer of glands or follicles which secrete a digestive fluid.

PROVERBS, MEDICAL. A proverbial expression frequently contains sound sense in its directions. In medicine, the greatest collection is in the *Regimen of Health of the School of Salerno*, composed in the 11th century, by John of Milan. Yet, although good sense is often inculcated, we frequently meet with the reverse.

Most of the proverbs are hygienic. The following are a few.

1. *Qu' après la soupe un coupe d'excellent vin
Tire un écu de la poche du médecin.*
2. *Surge quintâ, prauê nonâ, cãna quintâ, dormi nonâ, nec est mortî vita prona.*
3. *Nunquam recte corpus exerceri sine animo,
neque animum sine corpore posse.*
4. *Viande bien machée est à demi digérée.*
5. *Optimum condimentum fames.*
Appetite is the best sauce.
6. *Plures occidit gula quam gladius.*

7. Early to bed and early to rise,
Makes a man healthy, wealthy, and wise.

8. One man's meat's another man's poison.

9. An ounce of prevention is better than a pound of cure.

PROVERBIO, Anteversion.

PROVINS, MINERAL WATERS OF. Chalybeate springs in the department of Seine et Marne, France, which contain carbonic acid and iron.

PROVOCATORIUS, Intercalary.

PROXIMAD, see Proximal aspect.

PROXIMAL, from *proximus*, 'next.' Proximate; nearest; next.

PROXIMAL ASPECT. An aspect towards the trunk, in the course of an extremity. — Barclay. *Prox'mad* is used by the same writer adverbially to signify 'towards the trunk.'

PRUNA, Anthrax.

PRUNE, Prunum.

PRUNE JUICE EXPECTORATION OR SPUTA. The sputa resembling prune juice, which occur in the third or purulent stage of pneumonia.

PRUNEAU, Prunum.

PRUNELLA, Angina pectoris, Aphthæ, Cyanache, Pupil.

PRUNELLA, P. vulgaris seu *officina'lis*, *Brunella*, *Consolida minor*, *Synphytum minus*, *Self-heal*, *Heat-all*, *Bugle*, (F.) *Prunelle*, *Brunelle*; *Ord. Labiate*. *Sex. Syst. Didynamia Gymnospermia*. This plant has been recommended as an astringent, in hemorrhages and fluxes; and, also, in gargles against aphthæ, and in inflammation of the fauces.

PRUNELLE SAL, Potassæ nitras fusus sulphatis puccillo mixtus.

PRUNELLE, Prunella, Pupil.

PRUNELLIER, Prunus spinosa.

PRUNELLOE, Prunum Brignole.

PRUNEOLA SYLVESTRIS, Prunus spinosa.

PRUNES, see Prunum — p. Pulp of, Pruni Pulpa.

PRUNI PULPA, *Pulp of prunes*. Take of prunes a sufficient quantity, soften them in the vapour of boiling water, and having separated the stones, beat the remainder in a marble mortar and press through a hair-sieve. — Ph. U. S.

PRUNIER, Prunus domestica — p. *Sauvage*, Prunus spinosa.

PRUNUM. A dried Plum, a Prune. (F.) *Pruneau*. Three sorts of Prunes have been usually ranked amongst the articles of the materia medica. 1. The *Prunum Brignolen'se*; the *Brignole plum* or *Prunelloe*; from Brignole in Provence. It is of a reddish-yellow colour; and has a very grateful, sweet, subacid taste. 2. The *Prunum Gallicum*, *Prunum* (Ph. U. S.), the *French Prune*; and 3. The *Prunum Damascenum* or *Damson*, *Brab'ylon*. All these possess the same general qualities as the other summer fruits. They are emollient and laxative.

PRUNUM STELLATUM, Averrhoa carambola.

PRUNUS ACACIA, Prunus padus.

PRUNUS ARMENTACA, *Armeniaca vulgaris* seu *epirotica*. The *Apricot*, *Apriocock*, *Bericaucæ*, *Βερικοκκον*, *Præcocia*, (F.) *Abricotier*. *Ord. Rosaceæ*. *Sex. Syst. Icosandria Monogynia*. The apricot, *Melum Armeniacum*, (F.) *Abricot*, when ripe, is easily digested, and considered as a great delicacy.

PRUNUS A'VIUM, *P. cer'anus* seu *sylvestris* seu *macrophylla*, *Cer'anus A'vium* seu *nigra* seu *dulcis*, *Black Cherry Tree*, (F.) *Merisier*. The fruit is eaten; and a gum exudes from the tree, the properties of which are similar to those of gum Arabic.

PRUNUS CAPULIN, (S.) *Capulies*, grows in the

open fields of Peru; and is cultivated in gardens in the towns. The fruit is acid, and not often eaten; but on account of its agreeable odour it is used in making *Pucheros de flores*; and with odoriferous flowers to perfume linen. — Tschudi.

PRUNUS CERASUS, *Cer'asus ac'idula seu hortens'is seu vulgari's seu rubra*. The *Red Cherry Tree*; (F.) *Cerisier*. The fruit *Cer'asum*, *Cera'sion*, has a pleasant, acidulous sweet flavour. It is wholesome. See *Prunus avium*.

PRUNUS DOMESTICA. The *Plum* or *Damson tree*, *Coccyne'lea*, *Proune*, (F.) *Prunier ordinaire*. The damson, when perfectly ripe, affords a wholesome fruit for pies, tarts, &c. It is gently laxative. See *Prunum*.

PRUNUS HORTENSIS, *P. cerasus*.

PRUNUS INSITIA. The *Bullace Plum Tree*. The fruit of this tree is used like the damson.

PRUNUS LAURO-CER'ASUS, *Cer'asus lauro-cerasus*, *Padus lauro-cerasus*, *Poison Laurel*, *Cherry Laurel*, *Common Laurel*, *Lauro-cerasus*, (F.) *Laurier-cerise*, *L. amandier*, *L. de Trébisonde*. The leaves, *Lauro-cer'asus*, (Ph. E. and D.) have a bitter, styptic taste, with the flavour of bitter almonds. The flowers have a similar flavour. The powdered leaves, applied to the nostrils, excite sneezing. The flavour of the leaves has given occasion to their being employed in tarts, custards, &c.; but in large quantities they are poisonous. The poisonous principle is the prussic acid. The distilled water, called *Laurel-water*, is poisonous. (See *Poisons*, Table of.) An *Aqua LAURO-CER'ASI*, *Cherry Laurel-water*, is official in the Edinburgh and Dublin Pharmacopœias. Dose, ℞ xxx to fʒi.

PRUNUS MACROPHYLLA, *P. avium*.

PRUNUS PADUS, *Padus*, *P. avium*, *Cer'asus Padus seu racem'osus sylvest'is*, *Wild cluster* or *Bird Cherry Tree*, (Sc.) *Hagberry*, *Hack-berry*, (Prov.) *Heckberry*, (F.) *Cerisier à grappes*, *Bois puant*. The bark of the tree has a fragrant smell, and a bitter sub-astringent taste, somewhat similar to that of bitter almonds. It has been recommended in intermittents, &c., in the form of decoction. The berries have been used in dysentery.

PRUNUS SEBASTINA, *Sebastina*.

PRUNUS SPINOSA. The *Sloe Tree*, *P. sylvest'ris seu Aca'cia*, *Agriococcine'lea*, *Agru'na*, *Agru'ne'la*, (F.) *Prunellier*, *Prunier sauvage*. The fruit—the *sloe*—is sometimes employed in gargles, and was formerly much used in hemorrhage, owing to the astringent properties. The *Aca'cia nostras seu German'ica seu vulgari's*, *Prune'ola sylvest'ris*, *Succus acaciæ nostrat'is*, *S. Acaciæ Germanicæ inspissat'us* is obtained from this.

PRUNUS SYLVESTRIS, *P. Avium*, *P. Spinosa*.

PRUNUS VIRGINIANA, *Cer'asus scro'tina seu Virginiana*, *Choke cherry*, *Wild Cherry Tree*, (F.) *Cerisier de Virginie*. The bark of this tree has been found useful in intermittents. The leaves are poisonous to certain animals, and the berries intoxicate different kinds of birds. The Indians use the bark in the cure of syphilis. It is bitter and astringent, and possesses some aromatic warmth, and likewise a sedative quality. It is stimulant and tonic. The leaves contain prussic acid. The distilled oil of the bark is very nearly identical with that of the bitter almond. A strong decoction of the bark is anthelmintic.

PRUNUS VULGARIS, *P. Padus*.

PRURIGINOUS, *Prurigin'osus*, (F.) *Prurigenus*. Same etymon as the next. Relating, or belonging to, or resembling prurigo.

PRURIGO, *Pruri'tus*, *Chris'ma*; from *prurio*, *pruritum*, 'to itch.' The word is often used synonymously with *itching*. Willan and Bate-man use the term for a genus of cutaneous diseases, the characteristic symptoms of which are

a severe itching, accompanied by an eruption of papulæ of nearly the same colour as the adjoining cuticle; *Cnemos*, *Scabies papuliform'is*, *Exor'mia pruri'go*. It affects the whole surface of the skin, under three varieties of form;—the *P. mitis*, *P. formicans*, *P. senilis*, &c., as well as some parts of the body, as the *P. podicis*, and *P. pudendi mulie'bris*, *Puden'dagra pruriens*, *Pruri'tus vulvæ*, (F.) *Prurit' de la vulve*.

PRURIGO, Heat—p. *Pedicularis*, *Phthiriasis*.

PRURIT, Itching—p. *de la Vulve*, see *Prurigo*.

PRURITUS, *Gargalus*, *Itching*, *Prurigo*—p. *Vulvæ*, see *Prurigo*.

PRUSSIAN BLUE, *Ferrum Zoö'ticum seu Borussicum seu Cyanogenat'um seu oxydulat'um hydrocyanicum*, *Ferri Ferrocyanas seu Ferrocyanuret'um seu Cyanuret'um seu Perecyanidum*, *Borussicus seu Prussicus seu Hydrocyanus Ferri*, *Cæruleum Borussicum seu Berolinens'e*, *Tritohydro-ferrocyanate* or *Ferrocyanuret* or *Ferropurussiate* of iron, (F.) *Bleu de Prusse*, *B. de Berlin*. In the Pharmacopœia of the United States, the pure salt—*Ferri ferrocyanuretum*—is directed to be made as follows: *Ferri Sulph. ʒiv*, *acid sulph. fʒiiss*, *acid nitric. fʒvj* or *q. s.*, *potassii ferrocyanuret. ʒivss*, *aquæ*, *Oij*. Dissolve the sulphate in a pint of water, and having added the sulphuric acid, boil the solution. Pour into it the nitric acid, in small portions, boiling the liquid for a minute or two after each addition, until it no longer produces a dark colour; then allow the liquid to cool. Dissolve the ferrocyanuret of potassium in the remainder of the water, and add this solution gradually to the first liquid, agitating the mixture after each addition; then pour it upon a filter. Wash the precipitate with boiling water until the washings pass tasteless. Lastly, dry and rub into powder. This salt is chiefly used in the preparation of the hydrocyanic acid and the cyanuret of mercury. It has been advised in the treatment of intermittents, and in epilepsy and scrophulosis. Externally, it has been applied to ill-conditioned ulcers. (ʒj to ʒj, of cerate.) Dose, four to six grains.

PSALIDIUM, *Fornix*.

PSALIS, *Fornix*, *Scissors*.

PSALTERIUM, *Lyra*.

PSAMMA, see *Gravel*.

PSAMMIS'MUS, *Amnis'mus*, *Gammis'mus*, from *ψαμμος*, 'sand.' *Arenat'io*. The application of sand to any part of the body for the cure of disease.—Paulus.

PSAMMODES, *Sabulosus*.

PSAMMUS, see *Gravel*.

PSAUTIER, *Omasum*.

PSELLIS'MUS; *ψελλισμος*, *Psellotes*, *ψελλοτης*, from *ψελλίζω*, 'I stutter.' Imperfect or depraved state of the articulation. A genus in the class *Pneumatica*, order *Phonica*, of Good.

PSELLISMUS NASITAS, *Rhinophonia*—p. *Rhotacismus*, *Rotacism*.

PSELLOTES, *Balbuties*.

PSEUDACACIA ODORATA, *Robinia pseudacacia*.

PSEUDAC'OË, *Pseudacoi'a*, *Pseudecoi'a*, from *ψεύδης*, 'false,' and *ακοη*, 'audition.' *Pseudacui'sis*. False hearing.

PSEUDACORUS, *Iris pseudacorus*.

PSEUDÆSTHES'IA, *Sensus pervers'us*, from *ψεύδης*, 'false,' and *αισθάνομαι*, 'I feel.' *αίσθησις*, 'feeling.' *Paropsis illus'oria*, *Pseudaph'phē*, *Pseudaph'ia*, (F.) *Pseudæsthes'ia*. Depraved feeling. Imaginary sense of touch or general feeling in organs that have no existence. Common to those who have suffered amputation; and to hypochondriacs and the insane.

PSEUDALEI MORBI, Feigned diseases.

PSEUDANGUSTURINUM, *Bruc'ne*.

PSEUDAPHE, *Pseudæsthesia*.

PSEUDARTHROSIS, *Pseudarticulatio*, *Pseudarticulus*, *Articulatio nostra seu artificialis*, from *ψεύδης*, 'false,' and *ἄρθρον*, 'a joint.' A false joint or articulation: *Articulus nostrus seu spurius seu præternaturalis*, (F.) *Articulation enroulée*.

PSEUDARTICULATIO, Pseudarthrosis.

PSEUDECOIA, Pseudæcoe.

PSEUDENCEPHALUS, *Paracephalus*, from *ψεύδης*, 'false,' and *κεφαλον*, 'the encephalon.' A monster, whose cranium is open in its whole extent, from before to behind, its base supporting a vascular tumour. — I. G. St. Hilaire.

PSEUDETHÉSIE, Pseudæsthesia.

PSEUDO, False — p. Asthma, Dyspnoea.

PSEUDOBLEPSIA, *Pseudoblepsia*, *Pseudoræsis*, *Suffusio oculi*, from *ψεύδης*, 'false,' and *βλέψω*, 'sight.' *Pseudopsia*, *Paropsis illusoria*, *Suffusio*, *Phantasma*, *False sight*. A generic name, used by Cullen for perversion of vision. See Metamorphopsia.

PSEUDOBLEPSIS, Paropsis.

PSEUDOCARCINOMA LABII, Cancer aquaticus.

PSEUDOCROMIA, Achromatopsia.

PSEUDOCROUP, Asthma thymicum.

PSEUDOCYESIS, Pregnancy, false — p. *Cycis molaris*, Mole.

PSEUDENCEPHALITIS, Hydrencephaloid disease.

PSEUDO-FEVER, Irritation, morbid.

PSEUDOGEUSIA, *Pseudogeusia*, *Pseudogæstia*, from *ψεύδης*, 'false,' and *γεύω*, 'taste.' *Halucinatio gustus*. False taste.

PSEUDOGEUSTIA, Pseudogeusia.

PSEUDOHELMINTHES, Ectozoa.

PSEUDOLIEN, *Pseudosplen*, from *ψεύδης*, 'false,' and *lien*, 'spleen.' Glands situate in the neighborhood of the spleen; probably, lymphatic glands.

PSEUDOMECHOACANA, *Convolvulus panduratus*.

PSEUDOMEDICUS, Charlatan.

PSEUDOMELANOTIC FORMATIONS, see Anthracosis.

PSEUDOMELANOTIC MENINX, Membrane, false — p. Metamorphosis polyposa, polypus.

PSEUDOMEMBRANA, Membrane, false.

PSEUDOMEMBRANE, Membrane, false.

PSEUDONARCISSUS, *Narcissus pseudonarcissus*.

PSEUDONOMANIA, from *ψεύδης*, 'false,' and *μανία*. A form of insanity, which is characterized by a morbid propensity for lying.

PSEUDO-PARASITES, Ectozoa.

PSEUDOPERIPNEUMONIA, *Peripneumonia notha*.

PSEUDOPHILOGOSIS VENTRICULI RESOLUTIVA ET COLLIQUATIVA, *Gastromalacia*.

PSEUDOPHTHISIS, 'False phthisis.' Emaciation occasioned by other cause than organic lesion of the lungs.

PSEUDOPLASMATA, from *ψεύδης*, 'false,' and *πλάσσω*, 'I form.' A term applied to malignant heterologous tumours. — Vogel.

PSEUDOPLEURITIS, *Pleurodynia* — p. Polyp, see Polyp — p. Pyrethrum, Achillea ptarmica — p. Spleen, Pseudolien.

PSEUDOPNEUMONIA, *Peripneumonia notha*.

PSEUDOPNEUMONITIS, *Peripneumonia notha*.

PSEUDOPSIA, *Pseudoblepsia*.

PSEUDORASIS, *Pseudoblepsia*.

PSEUDOREXIA, *Pseudorexia*, from *ψεύδης*, 'false,' and *ορέξω*, 'appetite.' False appetite.

PSEUDO-RHONCHUS, from *ψεύδης*, 'false,' and *ρῆγχις*, 'rattle.' A sound heard during respiration, which appears to be produced, like the

ordinary rhonchi, in the air-tubes, but is exterior to them—as in the case of *pleural pseudorhonchi*. See Rattle, and Râle.

PSEUDOSCOPIC, *Pseudoscopicus*, from *ψεύδης*, 'false,' and *σκοπεω*, 'I view.' Conveying false impressions to the eye; as where projecting surfaces are seen depressed, and conversely.

PSEUDOSMIA, *Pseudosphresia*, *Pseudosphresis*, from *ψεύδης*, 'false,' and *οσμή*, 'smell.' False sense of smell.

PSEUDOSPIRESIA, Pseudosmia.

PSEUDOSYPHILIS, Syphilis pseudosyphilis.

PSEUDOTHANATOS, Asphyxia.

PSEUDOVARIOLÆ, Varicella.

PSEUDYMEN, Membrane, false.

PSIDIUM POMIFERUM. The Apple *Guava*, *Guayava*, *Ord.* Myrtaceæ.

This plant and *P. Pigriferum* bear fruits; those of the former like apples,—of the latter like pears. The apple kind has an acid flavour, the other is sweet. Of the inner pulp of either, the natives of the Indies make jellies; and, of the outer rind, tarts, marmalades, &c. They are somewhat astringent; this quality, indeed, exists in every part of the tree, and abundantly in the leaf-buds, which are occasionally boiled with barley and liquorice as an excellent drink in diarrhoea. A simple decoction of the leaves is said to be useful in the itch, and most cutaneous eruptions.

PSILOSTIS, Depilation.

PSILOTHRUM, *Bryonia alba*, Depilatory.

PSIMMYTHON, *Plumbi subcarbonas*.

PSOA, *Dysodia*.

PSOÆ, *Psœa*, *Psyæ*, *ψοαί*, see *Lumbi*, 'the loins.' The psœ muscles; *Alopæces*, *Nephrometææ*, *Neuroneures*.

PSOAS MAGNUS, *Psoas seu lumbæris internus*, *Femur moventium sextus*, *Præ-lombo-trochantin*, *Præ-lombo-trochantinien* (Ch.). A muscle, seated on the lateral parts of the lumbar portion of the vertebral column; and passing down to the superior and anterior part of the thigh. It is long, fusiform, and attached, above, to the body and transverse processes of the first four lumbar vertebrae, and last dorsal. Below, it terminates, by a very thick tendon, common to it and the iliacus muscle,—which is inserted into the top of the lesser trochanter. This muscle bends the thigh upon the pelvis, and carries it in rotation outwards. It may, also, bend the pelvis on the thigh, and the loins on the pelvis. It acts considerably in station and progression.

PSOAS PARVUS, *Præ-lombo-pubien*, (Ch.) A muscle situate anteriorly to the last. It is long, thin, flat, and narrow; and is attached, above, to the body of the last dorsal vertebra, and below, to the linea ileo-pectinea, by means of a slender tendon, which furnishes two expansions—one to the *lineæ aponeurosis*, and the other to the *pelvis*. This muscle bends the vertebral column on the pelvis; and, reciprocally, the pelvis on the vertebral column.

PSODYMUS, *Gastrodidymus*.

PSOIE, *Psœa*.

PSOÏTE, *Psotitis*.

PSOÏTIS, *Inflammatio musculi Psœas*, *Lumbægis inflammatoria spæmatica*, (F.) *Psôte*, *Inflammation du muscle Psœas*. From *ψοαί*, 'the loins,' 'the psœ muscles,' and *itis*, denoting inflammation. Inflammation of the psœas muscles and of the parts situate before the lumbar vertebrae: accompanied, from its commencement, by considerable fever, violent pains in the part affected and in the thigh of the same side, which remains commonly half bent. Psotitis may terminate by suppuration, and form a lumbar abscess.

PSOLÊ, *ψωλή*, 'the male organ.' Also, the glands when uncovered.

PSOLON, Membrosus.

PSOLON'CUS, from *psole*, and *ογκος*, 'a tumour.' Swelling of the glans, or of the penis generally.

PSOLUS, from *psole*. One who has the glans uncovered. One who has been circumcised.

PSOPHOMETER, Stethoscope.

PSOPIOS, *ψοφος*, 'a noise.' A loud noise; a hollow sound.

PSORA, from *ψωω*, 'I rub gently.' *Scabies*, *Phlysia* seu *Eccymosis* *Scabies*, *Scabiola*, *Itch*, *Scratch*, *Scotch fiddle*, (Prov.) *Ichet*, *Scrat*, (Sc.) *the Reif*, *Youk*, *Yuke*, (F.) *Gale*, *Mal Saint-Main*, *Rogue*. A contagious eruption of very minute pimples—pustular, vesicular, papular, intermixed, and alternating, itching intolerably and terminating in scabs. It seems to be connected with an insect of the genus *Acarus*;—the *Acarus* seu *Sarcoptes* seu *Sarcoptes* seu *Phthirus* seu *Cheyletus* *Scabiei*, *Acarus* *Ciro*, *Itch-insect*, *Itch-animalcule*, *Itch mite*, (F.) *Ciron*. The itch occupies, particularly, the spaces between the fingers, the back of the hands, wrists, elbows, axillæ, groins, hams, &c., and rarely affects the face. The best applications, for its cure, are,—the *Unguentum Sulphuris* or *Ung. Sulph. comp.* night and morning; but the unpleasant smell of the sulphur has given occasion to the use of other means;—as the *White Hellebore*, *Potass in deliquescence*, *Muriate of Ammonia*, *Sulphuric Acid*, &c.. The repugnance to the smell of sulphur ought not, however, to be regarded; especially as, in the course of four or five days, it will, in almost all cases, produce a perfect cure.

PSORA AGRIA, Psoriasis inveterata.

PSORA EBRIORUM, *Drunkard's Itch*. An eruption, attended by afflicting itching, to which elderly people, who have been much addicted to drinking alcoholic liquors, are liable.—Darwin.

PSORA LEPROSA, Psoriasis—p. *Squamosa*, *Psoriasis*.

PSORALEA, Psoralea pentaphylla.

PSORALEA GLANDULOSA, from *ψωραλεος* 'scurfy or scaly,' in reference to its scurfy glands or dots. *Alraguilla*. Ord. Leguminosæ. Ser. Syst. Diadelphia Decandria. A shrub, used in Chili as a vulnerary. The Paraguay tea is referred to it by M. Lenoble, of Montevideo. See *Ilex Paraguensis*.

PSORALEA PENTAPHYLLA, *Psora'lea*, *Contrayerva noca*, *Maxican Contrayerva*. The root is little, if at all, inferior to the contrayerva in its medical properties; which, by the by, are unimportant.

PSORENTERIA, Cholera.

PSORIA'ASIS, (same etymon as *Psora*,) formerly, the state of being affected with *ψωρα*. *Lepidosis psori'asis*, *Serpe'do*, *Serpi'go*, *Impetigo* (of some), *Scabies sic'ca* seu *fer'na*, *Psora lepro'sa* seu *squamo'sa*, *Scaly Tetter*, *Dry Scale*, *Dry Scall*, (F.) *Dartre squameuse lichenoidé*, *D. écailleuse*, *Gratelle*, *Gale militaire*, *G. canine* ou *sèche*. A cutaneous affection, consisting of patches of rough, amorphous scales; continuous, or of indeterminate outline; skin often chappy. Psoriasis occurs under a considerable variety of forms. The surface, under the scales, is more tender and irritable than in lepra, which psoriasis, in some respects, resembles. Dr. Willan has given names to eleven varieties of psoriasis: the chief of which are the *Psoriasis gutta'ta*, *P. diffu'sa* (one form of which is the *Baker's Itch* and *Grocer's Itch*; and another, the *Washerwoman's Scall*), the *P. gyra'ta*, and *P. invetera'ta* seu *Psora a'grica*. The treatment of irritable psoriasis must be antiphlogistic, with the internal use of the fixed alkalies, sulphur, &c. They do better without any local application. The more local and less inflammatory are ameliorated by emollient and

soothing applications; or by slightly stimulating means, sea-bathing, the *Ung. hydrarg. nitrico-oxyd.*, *Ung. oxidi zinci*, &c.

PSORIC, *Psor'icus*, (F.) *Psorique*, *Galeux*; from *ψωρα*, 'itch.' That which has the nature of itch. A medicine for curing the itch.

PSOROCOMI'UM, from *ψωρα*, 'the itch,' and *κομω*, 'I take care of.' An hospital for those affected with itch.

PSOROPHTHALMIA, Ophthalmia tarsi—p. Neonatorum, see Ophthalmia (purulenta infantum).

PSYÆ. *Psaw*.

PSYCHAGOGA, Psychagogica.

PSYCHAGOGICA, *Psychago'ga*, from *ψυχη*, 'life,' 'soul,' and *αγω*, 'I lead.' Medicines which resuscitate, when life seems extinct; as in syncope and apoplexy.

PSYCHE, Anima.

PSYCHIA'TER. Same etymon as the next. One who treats diseases of the mind—a *Mad-doctor*, (F.) *Médecin de fous*, *Aliéniste*.

PSYCHIATRIA, from *ψυχη*, 'mind,' and *ιατρία*, 'healing.' Treatment of diseases of the mind.

PSY'CHICAL, *Psy'chicus*, from *ψυχη*, 'mind.' That which relates to the mind or mental endowments. *Psycholog'ical* is not unfrequently used with the like signification.

PSYCHOLOGICAL, see Psychical.

PSYCHOLOGY, *Psycholog'ia*, from *ψυχη*, 'mind,' and *λογος*, 'discourse.' A treatise on the intellectual and moral faculties. Also, the intellectual and moral faculties or sphere.

PSYCHONOSOL'OGY, *Psychanosolog'ia*, from *ψυχη*, 'mind,' *νοσος*, 'disease,' and *λογος*, 'a description.' The doctrine of, or a treatise on, diseases of the mind.

PSYCHOPATHIES, Psychoses.

PSYCHORAGES, *Agonizans*, *Moribundus*, *Moribund*, *Fey*, *Fee*, *Fie*, from *ψυχη*, 'mind,' 'life,' and *ραγη*, 'rupture.' Dying; in *Articulo Mortis*, "in the article of death;" in *extre'mis*. *Moribund* means, also, one who is dying.

PSYCHORAGIA, Agony, Death.

PSYCHORRHAGIA, Agony, Death.

PSYCHOS, Cold.

PSYCHO'SES, from *ψυχη*, 'mind.' Neuroses of the intellect. Diseases of the mind. *Psychopathies*.

PSYCHOTHERAPEIA, from *ψυχη*, 'mind,' and *θεραπεω*, 'I attend upon the sick.' Treatment of disease through the mind.

PSYCHOTICA, Analeptica.

PSYCHOTRIA, Analeptica—p. *Crocea*, *Palicourea crocea*—p. *Emetica*, *Ipecacuanha*.

PSYCHOTROPHUM, *Betonica officinalis*.

PSYCHROLUSIA, *Psychrolu'tion*, from *ψυχος*, 'cold,' and *λουω*, 'I wash.' A cold bath.

PSYCHROPHOBIA, from *ψυχρος*, cold, and *φοβος*, 'fear.' Dread of cold, especially of cold water. Impressibility to cold.

PSYCHTICA, Refrigerants.

PSYDRA'CIA, quasi, *ψωρα* 'scabies,' 'cold drops,' from *ψυδραξ*, *ψυδρακος*, 'a small blister or pimple.' The ancients sometimes used this term for *pimples*; at others, for *vesicles*. Willan and Bateman define *Psychracium* to be, a small pustule, often irregularly circumscribed, producing but a slight elevation of the cuticle, and terminating in a laminated scab. Many of the psychracia usually appear together, and become confluent; and, after the discharge of pus, they pour out a thin, watery humour; which frequently forms an irregular incrustation. Frank limits the term to a psoriform eruption, which differs from itch in not originating from an insect, and is not contagious;—perhaps the lichen. Authors have differed regarding its acception.

PSYDRACIA ACNE, Acne.
 PSYLLIUM, Plantago psyllium—p. Erectum, Plantago psyllium.
 PTARMICA, Achillea ptarmica—p. Montana, Arctica montana.
 PTARMICUM, Erhine, Sternutatory.
 PTARMOS, Sneezing.
 PTELEA, Ulnus.
 PTELEA TRIFOLIATA, Shrubby trefoil, Swamp dogwood, Stinking ash, Stinking prairie bush, Wingweed; indigenous. Order, Xanthoxylaceæ. Flowering in June. Is said to be possessed of antiperiodic virtues, and to be anthelmintic.
 PTERIS, Polypodium filix mas.
 PTERIS AQUILINA, (πτερίς, 'a wing,' from the shape of its leaves.) Pteris, Thelypteris, Filix, Nymphaea seu fœminæa, Asplenium aquilinum, Common Brake or Female Fern; Eagle Fern, Bracken, (F.) Fougère femelle, La grande Fougère. Nat. Ord. Filices. The root is considered to be anthelmintic; and as efficacious as that of the male fern in cases of tapeworm.
 PTERIS PALUSTRIS, Asplenium filix fœmina.
 PTERNA, Calcaneum.
 PTERNIUM, Calcaneum.
 PTERNOBATES, Calcigradus.
 PTEROCARPUS ERINACEA, see Kino—p. Marsupium, see Kino—p. Sandalinus, P. Santalinus.

PTEROCARPUS SANTALINUS, P. Santalinus, (πτερον, 'a wing,' and καρπός, 'fruit,') Santalum seu Santalum Rubrum, Red Sanders or Saunders Tree, (F.) Santal rouge. Ord. Leguminosæ. See Syst. Diadelphia Decandria. Red Saunders wood, Santalum, (Ph. U. S.), Lignum santalinum is used only as a colouring matter: it gives to rectified spirit a fine deep red. The juice of this tree, like that of *P. Draco*, affords a species of Dragon's Blood.

PTEROCALON PYCNOSTACHYUM, Cornifera pycnostachya, Black root; indigenous. Order, Compositæ. The root is used as an alternative in the form of decoction. It is said to be possessed of cathartic properties.

PTEROS'PORA ANDROMEDA, (πτερον, 'a wing,' and σπορα, 'seed,' in allusion to the singular wing borne by the seeds,) Scaly Dragon claw, Dragon root, Fever root, Albany beechdrop. An indigenous plant, Nat. Ord. Monotropes, Sec. Syst. Decandria Monogynia—found in the State of New York, which blossoms in July. The root has been regarded as an anthelmintic, diuretic, emmenagogue, &c.: but is not used.

PTERYGION, Pterygium, Ala, from πτερίς, πτερυγός, 'a wing.' A small wing. Web, Onyx, Pygois, Unguis, (F.) Drapeau, Onglet. A varicose excrescence of the conjunctiva, of a triangular shape, and commonly occurring at the inner angle of the eye; whence it extends over the cornea. Sometimes two or three pterygia occur on the same eye, and cover the whole cornea with a thick veil, which totally prevents vision. According to Scarpa, the ancients gave the name Panniculus to this last complication. If the pterygion cannot be dispelled by means of discutient collyria, it must be removed by the scissors.
 PTERYGIUM, Pterygion, Scapula—p. Digniti, Paronychia—p. Lardaceum, Pinguecula—p. Pingue, Pinguecula.

PTERYGO, in the composition of anatomical terms, means relating, or appertaining, to the pterygoid process.

PTERYGO-ANGULI-MAXILLARE, Pterygoideus externus—p. Colli-maxillare, Pterygoideus externus—p. Maxillare, grand, Pterygoideus internus—p. Maxillare, petit, Pterygoideus externus—p. Palatine canal, Pterygoid canal—p. Pharyngeus, Constrictor pharyngis—p. Staphylinus, Circumdex.

PTERYGO-PALATINE, Pterygo-Palatinus. That which belongs to the pterygoid process and palate.

PTERYGO-PALATINE ARTERY or Superior pharyngeal is a very small artery, and is given off by the internal maxillary at the bottom of the zygomatic fossa: after having passed through the pterygo-palatine canal, it is distributed to the superior part of the pharynx, the sphenoid bone, and the Eustachian tube.

PTERYGO-PALATINE CANAL. A small canal, formed by the internal ala of the pterygoid process, and the palate bone. It gives passage to the vessels of the same name.

PTERYGO-STAPHYLINUS INTERNUS, Levator palati—p. Syndesmo-staphylo-pharyngien, Constrictor pharyngis.

PTERYGOIDES, Alatus.

PTERYGOIDE, Pterygoïdes, Pterygoïdeus, Aliformis; from πτερίς, πτερυγός, 'a wing,' and εidos, 'form.' A name given to two processes at the inferior surface of the sphenoid bone; the two laminae which form them having been compared to wings. These alæ or wings or processes, Processus aliformes, have been distinguished, according to their position, into internal (entopterygoïd) and external.

PTERYGOID ARTERY, Vid'ian Artery, Ramulus Ductus Pterygoïde'i, Arteria pharyngea suprema, arises from the internal maxillary artery, at the bottom of the zygomatic fossa, and enters the pterygoid canal with the nerve of the same name, to proceed to be distributed on the Eustachian tube and velum palati. The name, pterygoid arteries, is likewise given to the branches, which the internal maxillary and some of its divisions, as the meningeal media, and posterior temporalis profunda, furnish to the pterygoid muscles, behind the neck of the lower jaw.

PTERYGOID BONE, Sphenoid.

PTERYGOID CANAL, Vid'ian canal, Ductus pterygoïdeus, Canalis Vid'ianus, (F.) Conduit pterygoïdien ou Vidien, Pterygo-palatine canal, is a narrow channel, which traverses, in a direction from before to behind, the base of the pterygoid process, and gives passage to the pterygoid nerve. The foramina of the sphenoid, which terminate the canals, have the same epithets applied to them.

PTERYGOID FOSSA, Fossa Pterygoïde'a, (F.) Fosse pterygoïdienne ou pterygoïde, is the depression which separates the two laminae or alæ of the pterygoid process.

PTERYGOID MUSCLES, Alares musculi, Aliformes musculi, are two in number. 1. The Pterygoideus externus seu minor, (F.) Pterygo-colli-maxillare, Petit Pterygo-maxillare (Ch.), Petit pterygoïdien, Pterygoïdien externe, is a thick, short muscle, which arises, by short aponeuroses, from the outer surface of the external ala of the pterygoid process, and from the outer surface of the great ala of the sphenoid bone; and is inserted at the anterior part of the neck of the lower jaw-bone, and the corresponding part of the inter-articular fibro-cartilage. This muscle draws the condyle of the lower jaw and the inter-articular ligament forwards. It carries the chin forwards, and to the opposite side. If the two external pterygoids act together, the jaw is carried directly forwards. 2. The Pterygoideus internus seu major, Masseter internus, Latens in Oræ, (F.) Grand Pterygo-maxillare, Pterygo-anguli-maxillare, Grand Pterygoïdien ou Pterygoïdien interne. A thick, quadrilateral muscle, which is attached by short and very strong aponeuroses; above, to the pterygoid fossa, whence it proceeds downwards, backwards, and outwards, to be inserted into the inner surface of the ramus of the lower jaw, near its angle. This muscle

raises the lower jaw, and carries it a little forwards. It also performs the grinding motion.

PTERYGOID NERVES. These are two in number. I. The *Pterygoid* or *Vidian Nerve*. It arises from the posterior part of the sphenopalatine ganglion; traverses the Vidian canal of the sphenoid; and, beyond, divides into two filaments. 1. A *superior* or *cranial, Nervus superficialis sive petrosus*, which ascends into the cranium, and enters the hiatus of Fallopius to unite with the facial nerve. 2. An *inferior* or *carotid, Nervus profundus sive major*, which traverses the carotid canal; and unites with the ascending branches of the superior cervical ganglion of the trisplanchnic. II. The *Pterygoid Nerve*, properly so called, *Nervus musculi pterygoidei*, is given off by the inferior maxillary branch of the 5th pair.

PTERYGOID PROCESSES, see *Pterygoid*.

PTERYGOÏDIEN GRAND ou INTERNE, *Pterygoideus internus*—p. *Externe, Pterygoideus externus*—p. *Petit, Pterygoideus externus*.

PTERYGO'MA, from πτερυξ, πτερυγος, 'a wing; a projecting or pendulous body.' A swelling of the vulva, which prevents coition.—M. A. Severinus.

PTERYX, Ala.

PTILOSIIS, Deplumatio, Madarosis.

PTISANA, Tisane—p. *Communis, Decoctum hordei compositum*—p. *Hippocratica, Decoctum hordei*.

PTISSANA, Tisane.

PTOCHIA'TER, *Ptochia'trus*, from πτωχος, 'poor,' and ιατρος, 'a physician.' A physician to the poor, or to an establishment for the poor.

PTOCHIATRI'A, Practice amongst the poor. Also, an institution for the treatment of the diseases of the poor.

PTOCHOCOM'UM, from πτωχος, 'a poor person,' and κομω, 'I take care of,' *Ptochodochi'um, Ptochei'um, Ptochotrophe'um*. An almshouse. An hospital for the poor.

PTOCHODOCHIUM, *Ptochocomium*.

PTOCHOTROPHEUM, *Ptochocomium*.

PTOMA, Cadaver.

PTOSIS, πτωσις, *Prolapsus*—p. *Bulbi oculi, Exophthalmia*.

PTOSIS IRIDIS, Iridocel's, Her'nia seu Prolapsus Iridis, Iridoptosis, from πτωσις, 'a falling.' A prolapsus of the iris through a wound of the cornea. It is known by a blackish tubercle, which projects a little from the cornea in various forms.

PTOSIS PALPEBRÆ, Blepharoptosis.

PTYALOGOGUE, *Sialogogue*.

PTYALIN, see *Saliva*.

PTYALISM, *Salivation*.

PTYALISME, *Salivation*—p. *Mercuriel, Salivation, mercurial*.

PTYALISMUS, *Salivation*—p. *Acutus mellitus*, see *Salivation*—p. *Acutus sympatheticus*, see *Salivation*—p. *Hydrargyratus, Salivation, mercurial*—p. *Iners, Slaving*—p. *Mercurialis, Salivation, mercurial*—p. *Pyrosicus, Pyrosis*.

PTYALOGOGUE, *Sialogogue*.

PTYALON, *Saliva*.

PTYASMAGOGUE, *Expectorant, Sialogogue*.

PTYELON, *Saliva, Sputum*.

PTYSIS, *Exspuition*.

PTYSMA, *Saliva, Sputum*.

PTYSMAGOGUE, *Sialogogue*.

PTYSMATIS'CHESIS, *Ptysmatos'chesis*, from πτυσμα, 'sputum,' and χεω, 'to retain.' Suppression of pulmonary expectoration.

PTYSMATOSCHESIS, *Ptysmatichesis*.

PUANTEUR, *Dysodia*.

PUBENS, *Ephebus*.

PUBER, *Ephebus*.

PUB'ERAL; same etymon as *Puberty*. Re-

lating or appertaining to the age of puberty -- as 'a puberal disease.'

PUBÈRE, *Ephebus*.

PUBERTAS, *Ephebia*.

PUB'ERTY, *Pubertas, Hebē, Pubes, Aphrodis'ia, Ephebi'a, Ephe'bia, Ephe'botes, Ephebo'synē, Pubes'cence*, from *pubescere*, 'to begin to have hair on the chin and sexual organs.' The period of life at which young people of both sexes are *pubes'cent* or *nubile*. According to the civil law, puberty occurs at 12 years in females, and 14 in males. The period varies somewhat in different climates, and in different individuals of the same climate.

PUBES, *Pubis, Episc'h'ion, Oeis, Pecten*; the *Pubic region*. Same etymon. The middle part of the hypogastrio region; so called, because it is covered with hair, in both sexes, at the period of puberty. The hair is also called *Pubes, Lanugo* seu *Pili, Pudendo'rum, Planta'rium*; and, of old, *Epicten'ion* and *pubertas* signified the same. See *Tresoria*.

PUBES, *Puberty*.

PUBESCENCE, *Puberty*.

PUBESCENT, see *Puberty*.

PUBLÆUS, *Pubiē*.

PUBIC, *Pu'b'ian, Pu'b'icus, Pubi'cus, (F.) Pubien*. That which belongs to or concerns the *pubis*.

PUBIC ARCH, *Arch of the Pubis, (F.) Arcades pubienne*. The arch at the anterior part of the inferior circumference of the pelvis. It is more expanded in the female than in the male; to correspond with the additional functions which she has to execute.

PUBIC ARTICULA'TION, *Sym'physis Pubis*, results from the junction of the ossa *pubis*. The medium of union is cartilage.

PUBIC LIG'AMENTS are two very solid, fibrous fasciæ, situate before and beneath the symphysis *pubis*, which they strengthen. The one is called the *anterior pubic ligament*; the other the *inferior* or *subpubic*.

PUBIC REGION, *Re'gio Pubis*. The region of the *pubes*.

PUBIEN, *Pubic*.

PUBIO-COCYGIEN-ANN'LAIRE, *Pubio-coccyge'us annularis*. Dumas gave this name to the levator ani and ischio-coccygeus, which he considers a single muscle.

PUBIO-COCYGIEN-ANNULAIRE, *Levator Ani*—p. *Femoral, Pectinalis*—p. *Femoralis, Adductor longus femoris*—p. *Ombilical, Pyramidalis abdominis*—p. *Prostaticus, Compressor prostatae*—p. *Sous-ombilical, Pyramidalis abdominis*—p. *Sternal, Rectus abdominis*—p. *Urethral, see Compressor Urethrae*.

PUBIS, *Pubes*.

PUBIS OS, Os Pec'tinia, Episc'h'ion, Episc'h'on, Episc'hion, Share Bone, Bar-bone, (Sc.) Ark-bein. The anterior part of the os *innominatum*; so called because it corresponds to the genital organs and *pubic region*. The *angle of the pubis* is formed by the union of the body and rami of each side.

PUB'LICÆ PARTES. 'Public parts.' Those organs are so called, the object of whose functions is the nutrition of all the others. Such are the digestive and respiratory organs.

PUCCOON, Sanguinaria Canadensis—p. *Red, Sanguinaria Canadensis*—p. *Yellow, Hydrastis Canadensis*.

PUCE, *Pulex irritans*—p. *de Bourgogne, see Anthrax*—p. *Maligne, see Anthrax*—p. *Pénetrante, Chigue*.

PUCELAGE, *Virginity*.

PUCELLE, *Virgin*.

PUCHA-PAT, *Pogostemon patchouli*.

PUCHURY, see *Pichurim beans*.

PUCHYRY, see *Pichurim* beans.

PUCKLE. *Papula*.

PUDDING-GRASS, *Mentha pulegium*.

PUDDING-PIPE TREE, *Cassia fistula*.

PUDENDA, Genital organs—p. *Virorum*, *Pudibilia*.

PUDEND'AGRA. A hybrid word; from *pudenda*, ('*pudere*, 'to be ashamed,') 'the genital organs,' the Greek *agrap*, 'a seizure.' According to some, this term means pain in the genital parts: *Dolor pudendorum*, *Mentulagra*, *Ædæodyn'ia*; according to others, it is synonymous with syphilis.

PUDENDAGRA PRURIENS, *Prurigo pudendi muliebris*.

PUDEN'DAL, *Pudenda'lis*. Relating or pertaining to the *pudenda*.

PUDEN'DAL HER'NIA. *Hernia*, which descends between the vagina and ramus ischii, and forms an oblong tumour in the labium; traceable in the pelvis as far as the os uteri.

PUDENDALIS, *Pudendal*.

PUDENDALIS LONGUS INFERIOR, see *Sciatic nerve*, lesser.

PUDENDUM, FEMALE, *Vulva*—p. *Muliebre*, *Vulva*.

PUDENDUS, *Pudic*.

PUDIBIL'IA; from *pudere*, 'to be ashamed.' The male genital organs; *Genita'lia Viri*, *Viril'ia*, *Pensil'ia*, *Instrumenta*, *Puden'da viro'rum*, *Pondas*.

PUDIC, *Pudicus*, *Puden'dus*, (F.) *Honteux*. Same etymon. That which causes shame. A name given to the genital organs as well as to different other parts.

PUDIC ARTERIES are distributed to the organs of generation. They are,—1. The internal *Pudic Arteria* *Pudica*, *A. puden'da communis*, (F.) *Artère honteuse interne*, *A. Sous-pubienne* (Ch.), *Genital Artery*. It is furnished by the hypogastric, and divides into two branches; the one—the inferior or perineal—which gives off the inferior hemorrhoidal and the artery of the septum; the other—the superior, (F.) *Ischiopéniennne*—which furnishes the transverse perineal artery, and divides into two branches; the artery of the *corpus cavernosum*, and the *dorsalis penis*. This latter, in the female, has the name *Dorsalis Clitor'idis*. 2. The external *Pudic Arteries*, (F.) *Artères scrotales ou vulvaires* (Ch.), *A. honteuses externes*, which are two in number. The one is superficial or subcutaneous, and the other deep-seated or subaponeurotic. They are given off from the femoral artery, or some one of its branches; and are distributed to the external parts of generation.

PUDIC NERVE, (F.) *Nerf honteux ou ischio-pénien* ou *Ischio-clitorien*, is a branch furnished by the sacral plexus, which is distributed to the genital organs; and, especially, to the penis in man, and the clitoris in the female.

PUERILE, *Puerilis*, from *puer*, 'a child,' (F.) *Puéril*. Appertaining or relating to childhood.

Respiration puérile ou *exagérée* ou *supplémentaire* ou *hypervésiculaire*:—a term applied to the respiration when it is heard loud through the stethoscope, as in healthy children.

PUERILITAS, *Dementia* of the aged.

PUERTIA, see *Age*.

PUER'PERA, from *puer*, 'a child,' and *parere*, 'to bring forth.' *Tokas*, *Lochos*, *Enix'a*, *Lecho*, *n lying-in woman*, (F.) *Femme en Couchée*, *Accouchée*. A female in child-bed. One recently delivered.

PUER'PERAL, *Puerpera'lis*; from *puerpera*. Relating to parturition and its consequences.

PUERPERAL CONVULSIONS, *Eclampsia gravidarum* et *parturientium*.

PUER'PERAL FEVER. Any fever occurring

during the puerperal state—*Febria puerpera'rum*, *Lechopy'ra*, *Lochioaliti's*, *Lochiopy'ra*, *Lochopy'ra*, *Child-bed fever*. Most commonly restricted to a malignant variety of peritonitis.

PUERPERIUM, *Child-bed state*.

PUERPEROUS, *Parturient*.

PUFF, *Lycoperdon*.

PUFF, VEILED, see *Cavernous respiration*.

PUFF-BALL, *Lycoperdon*, *Leontodon taraxacum*.

PUFFINESS, from (D.) *poffen*, 'to blow.' *Infla'tio*, *Suffla'tio*, *Tumefac'tio mollis*, (F.) *Bouffissure*, *Boursouffure*. A soft intumescence, without redness, which may be partial or general, and be produced by accumulation of serous fluid, extravasation of blood, effusion of air into the subcutaneous areolar tissue, &c. Puffiness is most frequently seen in the face and legs; and sometimes appears around a wound.

PUGIL'US, *Drax*, *Dra'chion*, diminutive of *pugnus*, 'the fist.' *Dragma*, *Dragma*, (F.) *Pincée*. The quantity of any substance, which may be taken with the ends of two or three fingers. The eighth part of a handful.

PUGTEETH, *Canine teeth*.

PUISSANCE DU MOULE INTÉRIEUR, *Plastic force*.

PUKE, *Emetic*.

PUKEWEED, *Lobelia inflata*.

PUKING, *Vomiting*, *Regurgitation*.

PULEGIUM, *Mentha pulegium*—p. *Cervinum*, *Mentha cervina*.

PULEX IRRITANS. The common flea, (F.) *Puce*. A parasitic insect found on the bodies of man and animals.

PULEX PENETRANS, *Chique*.

PULICARIA DYSENTERICA, *Inula dysenterica*.

PULICA'RIS, *Pulicula'ris*, (F.) *Pulicaire*, from *pulex*, *pulicis*, 'a gnat.' An epithet given to cutaneous eruptions resembling gnat-bites;—and, also, to diseases in which such eruptions occur.

PULICARIS HERBA, *Plantago psyllium*.

PULNA, PILNA or BILNA, MINERAL WATERS OF. These well known bitter Bohemian springs contain sulphates of soda and magnesia, and chloride of magnesium.

PULMO, (pl.) *Pulmo'nes*; by transposition of the letter *l* from *pleumon*, *πνευμων*; *Pneumon*, *Pleumon*, *Spiramentum animæ*, *Ergasterion spir'itûs*, *Flabell'um* et *ventila'brum cordis*, *Lights*. The essential organ of respiration, which is double, and occupies the two sides of the chest. The lungs, which are of a spongy, soft, flexible, compressible, and dilatate structure, fill exactly the two cavities of the thorax, and are separated from each other by the mediastinum and the heart. The right lung, which is shorter and broader than the left, is divided, by two oblique fissures, into three unequal lobes, *Lobi seu Ala pulmo'num*. The left has only two lobes, and consequently only one fissure. At the internal surface of those organs, which is slightly concave, there is, about the middle, a pedicle, formed by the bronchia and pulmonary vessels, and called, by anatomists, the *root of the lungs*, (F.) *Racine des Pouxmons*. Essentially, the lungs are composed of prolongations and ramifications of the bronchia, and of the pulmonary arteries and veins,—the divisions of which are supported by a fine, areolar tissue.

When the surface of the lungs is examined in a clear light, we may see, even through the pleura, that their parenchyma is formed by the aggregation of a multitude of small vesicles, of an irregularly spheroid or ovoid shape, full of air, and separated by white and opaque septa, constituting lozenge-shaped spaces, which are called *lobules*—

lobuli seu insula pulmonales—and which are separated by interlobular areolar tissue. These lobules do not communicate with each other. The vesicles are called *Air-cells*, *Spiramenta* seu *Spiramina* seu *Cellulae Pulmonum*. Along the partitions or septa is deposited, in greater or less quantity, *black pulmonary matter*, (F.) *Matière pulmonaire noire*, as it has been called, which seems to be natural. Sometimes it is seen in points; at others, in spots. The colour of the lungs varies according to age and other circumstances. In youth, it is more red; and afterwards, grayish or bluish; often, as if marbled. The pleura pulmonalis is their investing membrane. The air is carried to the lungs by means of the trachea and bronchia. The black venous blood, which requires oxygenation, is conveyed to them from the heart, by the pulmonary artery; and when it has undergone this change, it is returned to the heart by the pulmonary veins. The blood-vessels inservient to the nutrition of the lungs are the bronchial arteries and veins. The pulmonary lymphatics are very numerous. Some are superficial; others, deep-seated. They pass, for the most part, into the bronchial ganglions or glands. The nerves are furnished by the pulmonary plexus.

PULMOMETER, Spirometer.

PULMOMETRY, see Spirometry.

PULMONAIRE, *Pulmonaria*—*p. en Arbre*, *Lichen pulmonaria*—*p. de Chêne*, *Lichen pulmonaria*—*p. des Français*, *Hieracium murorum*.

PULMONARIA, *P. Officinalis* seu *Maculata* seu *Maculosa*, Lungwort, Spotted Lungwort, *Jerusalem Cowslip*, *Jerusalem Sage*, *Spotted Comfrey*, (F.) *Pulmonaire*, *Sauge de Jerusalem*. Ord. Boraginaceae. Sez. Syst. Pentandria Monogynia. The leaves of this plant have been used as demulcents and pectorals; partly from their resemblance to lungs—hence the name, from *pulmo*, *pulmonis*, 'the lung'—and partly because they have a slightly mucilaginous taste. The plant is not worthy of notice.

PULMONARIA, *Pneumonia*—*p. Arborea*, *Lichen pulmonarius*—*p. Gallica*, *Hieracium murorum*—*p. Officinalis*, *Pulmonaria*—*p. Virginica*, *Mertensia Virginica*.

PULMONARY, *Pulmonaris*, *Pulmonalis*, from *pulmo*, *pulmonis*, 'the lung.' That which relates or belongs to the lungs. A name given to various parts.

PULMONARY ARTERY, *Arteria pulmonaris*, *Vena arteriosa*, arises from the upper and left part of the right ventricle of the heart; whence it passes obliquely upwards and to the left side; dividing into two trunks, opposite the second dorsal vertebra—one passing to each lung. At its origin, it is furnished with three sigmoid or semilunar valves. It carries the blood to the lungs for aeration.

PULMONARY MATTER, *Black*, see *Pulmo*.

PULMONARY PLEXUS, *Plexus pulmonaris*. A considerable nervous network, situate behind the bronchia, which is formed by the filaments of the pneumogastric nerve or par vagum, and by other filaments from the inferior cervical ganglion and the first thoracic ganglia. This plexus forms a kind of areolar web, with numerous meshes of different sizes, in which vessels are lodged, surrounded by much areolar tissue. This plexus sends filaments to the bronchia, which they accompany to their extremities.

PULMONARY VEINS, *Venae pulmonares*, *Arteriae venosae*. These arise in the lungs from the final terminations of the arteries of the same name; collect in ramusculi and rami of greater and greater size, and ultimately unite in four trunks; two of which issue from each lung to

open into the left auricle of the heart. The pulmonary veins have the same structure as the other veins of the body; from which they differ, in conveying arterial blood, as the pulmonary artery differs from the others in conveying venous blood. They carry the blood back to the heart after aeration.

PULMONES TARTARIZATI, *Lithiasis pulmonum*.

PULMONIA, *Phthisis pulmonalis*, *Pneumonia*.

PULMONIC, *Pulmonicus*, *Pneumonius*, *Pulmonarius*, *Pneumonice*, (F.) *Pulmonique*, *Pneumonique*. Same etymon as *Pulmonary*. One affected with phthisis—one labouring under diseases of the lungs. Any thing relating or belonging to the lungs. A medicine for the lungs.

PULP, *Pulpa*, *Pulpa'men*, *Sarx*, (F.) *Pulpe*, *Pulpolite*. The soft and parenchymatous parts of vegetables reduced to a paste by the operation of pulping. Applied, also, to parts of the human body, which have the characters of, or resemble a pulp.

PULP OF THE FINGERS, (F.) *Pulpe des doigts*. The fleshy extremity of the fingers.

PULP OF THE TOOTH, *Dental pulp*.

PULPA CASSIÆ, see *Cassia fistula*—*p. Dentis*, *Dental pulp*—*p. Pili*, see *Hair*—*p. Pruni*, see *Pruni pulpa*—*p. Prunorum sylvestrium condita*, *Conserva pruni sylvestris*—*p. Tamarindi*, see *Tamarindi pulpa*.

PULPA TESTIS, *Parent'chyma testis*. The parenchyma of the testis.

PULPAMEN, *Pulp*.

PULPATION, *Pulping*.

PULPE, *Pulp*—*p. Cérébrale*, see *Cerebrum*—*p. Dentarie*, *Dental pulp*—*p. des Doigts*, *Pulp of the fingers*.

PULPEUX, *Pulpy*.

PULPEZIA, *Apoplexy*.

PULPING, *Pulpa'tion*, *Pulpa'tio*. An operation—the object of which is to reduce certain vegetable substances to a pulp. For this purpose, they are bruised in a mortar, and then passed through a hair sieve, by the aid of a wooden spatula, called a *Pulper*, (F.) *Pulpoire*.

PULPOIRE, see *Pulping*.

PULPOLITE, *Pulp*.

PULPY, *Pulposus*, (F.) *Pulpeux*. Full of pulp.

PULQUE, see *Agave Americana*.

PULS EX OVIS et *LACTE*, *Oogala*.

PULSATIF, *Throbbing*.

PULSATILLA NIGRICANS, *Anemone pratensis*—*p. Patens*, *Anemone patens*—*p. Praten-sis*, *Anemone pratensis*—*p. Vulgaris*, *Anemone pulsatilla*.

PULSATILLE NOIRE, *Anemone pratensis*—*p. des Prés*, *Anemone pratensis*.

PULSATIO, *Pulsation*—*p. Cordis*, *Palpitation*.

PULSATION, *Pulsa'tio*, *Sphygmus*, *Sphyxis*, *Pede'sis*, *Pedeth'mos*, (F.) *Battement*. The beating of the heart and arteries. Also, the beating felt in an inflamed part.

PULSATIVUS, *Throbbing*.

PULSATORY, *Throbbing*.

PULSE, from *pulso*, 'I beat;' *Pulsus*, *Pulsa'tio*, *Pulsus* seu *Ictus cordis et arteriarum*, *Incur'tus arteriarum*, *Sphygmus*, *Sphygma*, *Diasphyxis*, (F.) *Pouls*. Beating of the arteries, produced by the afflux of the blood propelled by the heart in its contractions. For the sake of convenience, we choose the radial artery at the wrist to detect the precise character of the pulse. These characters relate to the number of pulsations in a

given time; to their degrees of quickness; hardness and strength; to the equality or inequality either of the pulsations themselves or of their intervals; to the development or fulness of the

artery, and to the different impressions it may produce on the finger.

The following estimates have been made of the pulse at various ages:

Ages.	Number of Beats per minute, according to		
	HEBERDEN.	SÜMMERING.	MÜLLER.
In the embryo			150
At birth.....	130 to 140		Do.
One month.....	120		
One year.....	120 to 108	120	115 to 130
Two years.....	108 to 90	110	100 to 115
Three years.....	90 to 80	90	90 to 100
Seven years.....	72		85 to 90
Twelve years.....	70		
Puberty.....		80	80 to 85
Adult age.....		70	70 to 75
Old age.....		60	50 to 65

It does not appear that the pulse of the aged is less frequent, but the contrary. In general it is more frequent in females and in irritable persons, than in males and those of an opposite temperament. In the healthy state, the pulse, besides having the proper number of pulsations, is neither hard nor unusually soft; it is moderately developed; of a medium strength; and perfectly equal, both as regards the pulsations and their intervals. In diseases it wanders, more or less, from these physiological conditions. The different characters of the pulse that have been recorded, are remarkably numerous. In a dictionary, it is necessary to detail them, although many of them are not now regarded, and some are ridiculous.

PULSE, ARDENT, *Pulsus ardens*, (F.) *Pouls ardent*, *Pouls élevé*. One which seems to raise itself to a point in order to strike the finger.

PULSE, CAPRIZANT, *Pulsus Caprizans* seu *dorcadisans* seu *gazellans* seu *fusalis*, Goat-leap pulse, (F.) *Pouls caprizant*. One which seems to leap, as it were; an imperfect dilatation of the artery being succeeded by a fuller and a stronger one.

PULSE, COMPLEX, (F.) *Pouls combiné* ou *composé*, *P. compliqué*. A pulse which has the character of several critical pulses united.

PULSE, CONTRACTED, (F.) *Pouls serré* ou *concentré*. One whose pulsations are narrow, deep, and slightly hard.

PULSE, CONVULSIVE, *Pulsus convulsivus*, (F.) *Pouls convulsif*. That which is unequally frequent or unequally hard; in short, which resembles the irregular pulsations of a muscular fibre in a state of convulsion.

PULSE, CRITICAL, *Pulsus criticus*. A pulse which, after having possessed the characters of irritation, becomes free, open, soft, &c.

PULSE, DECURTATE, *Pulsus myurus*: see, also, *Myurus*.

PULSE, DEEP, *Pulsus profundus*, (F.) *Pouls profond*. One which cannot be felt without difficulty, and without strong pressure by the finger.

PULSE, DEFICIENT, *Pulsus deficientis* seu *animens* et *circumnuens*, *P. Flickering*, (F.) *Pouls défaillant*, *P. insensible*. One which seems, every instant about to cease.

PULSE, DEPRESSED, *Pulsus depressus*, (F.) *Pouls déprimé*. That which is deep and weak.

PULSE, DEVELOPED, (F.) *Pouls développé*. That which is broad, full, strong, and frequent.

PULSE, DICTOTIC, *Pulsus Dictotus* seu *bicrotus* seu *intestinalis* seu *biferiens* seu *mallearius* seu *interiens* seu *duplic*, (F.) *Pouls dicoté* ou *reboulissant* ou *reboulé*. That in which the finger

is struck twice at each pulsation; once lightly, the other time more strongly.

PULSE, EQUAL, *Pulsus æqualis*. That, in which the pulsations resemble each other, and return at equal intervals.

PULSE, FEBRILE, *Pulsus febrilis*. One very frequent.

PULSE, FEEBLE, *P. Weak*, *Pulsus debilis*, (F.) *Pouls faible*. One which strikes the finger feebly.

PULSE, FILIFORM, *P. thready*. One which resembles a thread slightly vibrating.

PULSE, FLICKERING, *Pulse*, deficient.

PULSE, FORMICANT, *Pulsus formicans*, (F.) *Pouls formicant*. One which resembles, by being small and scarcely perceptible, the motion of an ant.

PULSE, FREQUENT, (F.) *Pouls agité* ou *précipité*. One which strikes more frequently than usual in a given time.

PULSE, FULL, *Pulsus plenus*, (F.) *Pouls plein*. One which gives the notion of great repletion in the vessel.

PULSE, GOAT-LEAP, *P. Caprizant*.

PULSE, GUTTURAL, *Pulsus gutturalis*, (F.) *Pouls guttural*. A pulse which has been supposed to indicate the termination of affections of the throat.

PULSE, HARD, *Pulsus durus*, (F.) *Pouls dur*. That in which the pulp of the finger seems to yield to the pulsation.

PULSE, HECTIC, *Pulsus hecticus*. The weak, frequent pulse observed in hectic fever.

PULSE, HEMORRHOIDAL, (F.) *Pouls des Hémorrhoides*. A critical pulse, announcing the appearance of the hemorrhoidal flux. It is described as unequal;—the pulsations not being alike in force, and still less the intervals.

PULSE, HEPATIC, (F.) *Pouls du Foie*. A pulse imagined to indicate affections of the liver. It is not worthy of description.

PULSE, INFERIOR, (F.) *Pouls inférieur* ou *abdominal* ou *ventral*. One supposed to announce evacuations below the diaphragm.

PULSE, INTERCURRENT, *Pulsus intercurrentis* seu *interiens*. One in which a superfluous pulsation seems to occur from time to time.

PULSE, INTERMITTENT, (F.) *Pouls intercadent*. One in which the pulsations fail from time to time.

PULSE, INTES'TINAL, (F.) *Pouls des intestins*, *P. de la diarrhée*. A critical pulse, presumed to announce an evacuation by the intestines.

PULSE, INTRICATE, *Pulsus intricatus*, (F.) *Pouls embarrassée*. That which is but little developed and unequally slow.

PULSE, IRREGULAR, *Pulsus anormalis*. One

whose pulsations are unequal, and return at unequal intervals.

PULSE OF IRRITATION, *Pulsus ab irritatiōe seu nervinus seu acritivus*. One which is frequent, hard, and contracted.

PULSE, JARRING, *P.* vibrating—*p.* Jerking, Pulse, sharp.

PULSE, LAGUID, (*F.*) *Pulsus languissant*. A slow and feeble pulse.

PULSE, LARGE, *Pulsus crassus seu magnus*, (*F.*) *Pulsus grand*, *P. large*. That which is large and full.

PULSE, LONG. One which strikes the finger to a great extent in length.

PULSE, LOW, *Pulsus humilis*, (*F.*) *Pulsus bas*. One in which the pulsations are scarcely perceptible.

PULSE, NASAL, (*F.*) *Pulsus nasal* ou *capital* ou *du Nez*. One supposed to announce the end of an irritation in the nasal fosse.

PULSE, NATURAL, (*F.*) *Pulsus naturel des adultes* ou *parfait* ou *des adultes* ou *modéré*. One that is equal and regular in strength and frequency; compressible, and devoid of hardness.

PULSE, ORGANIC, (*F.*) *Pulsus organique*. One which indicates the affected organ, or the organ by which an evacuation will take place.

PULSE, PECTORAL, (*F.*) *Pulsus de l'expectoration*. A critical pulse, announcing the termination of some pulmonary irritation.

PULSE, QUICK. One which strikes sharply, but not forcibly, against the finger.

PULSE, REGULAR, *Pulsus normalis seu regularis*. One whose pulsations are equal, and succeed each other at equal intervals.

PULSE, RESISTING, (*F.*) *Pulsus résistant*. One a little tense, a little hard.

PULSE, SHARP, (*F.*) *Pulsus prompt* ou *vite* ou *vif*. One which strikes the finger sharply, rapidly. When to a greater extent, it is called a jerking pulse.

PULSE, SHARP-TAILED, *Pulsus myurus*: see *Myurus*.

PULSE, SIMPLE, *Pulsus simplex*. A critical pulse, indicating an evacuation by one organ only.

PULSE, SLOW, *Pulsus tardus seu rarus*, (*F.*) *Pulsus lent* ou *tardif* ou *rare*. One which strikes the finger slowly.

PULSE, SMALL, *Pulsus parvus*, (*F.*) *Pulsus petit*. One whose pulsations are slender and weak.

PULSE, SOFT, *Pulsus mollis*, (*F.*) *Pulsus mou*. One which yields under the finger.

PULSE, STOMACHAL, (*F.*) *Pulsus stomacal* ou *de l'estomac*. One supposed to announce an evacuation by the stomach.

PULSE, STRONG, *Pulsus fortis*, (*F.*) *Pulsus fort*, *P. véhément*. One which strikes the finger strongly.

PULSE, SUPERIOR, *Pulsus Superior*, (*F.*) *Pulsus supérieur*. A critical pulse, supervening on irritation of organs situate above the diaphragm.

PULSE, SUPPLE, *Pulsus elasticus*, (*F.*) *Pulsus souple*. One in which the pulsations have strength without hardness.

PULSE OF SWEAT, *Pulsus Sudoralis*, (*F.*) *Pulsus Sudoral* ou *de la sueur*. A critical pulse, indicating an approaching perspiration.

PULSE, TENSE, *P. Cordy*, *Pulsus tensus*, (*F.*) *Pulsus tendu*, *P. roide*. One in which the artery resembles a cord fixed at its two extremities. When resembling a wire, it is called wiry.

PULSE, THREADY, *P. filiform*.

PULSE, TREMBULOUS, *Pulsus tremulus*, (*F.*) *Pulsus tremblant* ou *palpitant*. One in which each pulsation oscillates.

PULSE, UNPULATING, *Pulsus fluctuosus seu fluens* ou *incédens Solani*, (*F.*) *Pulsus undulant*

ou *ondoyant*. One which resembles, in its movements, those of the waves.

PULSE, UNEQUAL, *Pulsus inæqualis*, (*F.*) *Pulsus inégal*. One whose pulsations are not alike, or return at unequal intervals.

PULSE, URINAL, *Pulsus urinarius*, (*F.*) *Pulsus de l'Urine*. A critical pulse, fancied to denote an approaching evacuation of urine.

PULSE, UTERINE, *Pulsus uterinus*, (*F.*) *Pulsus Utérine* ou *de la Matrice* ou *des Règles*. One which indicates the flow of the menses, or the occurrence of metrorrhagia.

PULSE, VAGINAL, *Pulsus vaginalis*. The pulsation of the arteria vaginalis, which is increased in pregnancy, during abortion, labour, &c.

PULSE, VENOUS, *Pulsus venarum*.

PULSE, VERMICULAR. That which imitates the movement of a worm.

PULSE, VIBRATING, *Jarring Pulse*. One in which the artery seems to vibrate like a musical string.

PULSE, WIRY, see *Pulse*, tense.

PULSELESS, *Asphyctos*, Devoid of pulsation.

PULSELESSNESS, *Asphyxia*.

PULSEY, *Cataplasma*.

PULSIFIC, *Pulsificus*; from *pulsus*, 'pulse.' That which causes or excites pulsation.

PULSILEGIUM, *Pulsilegium*, *Pulsimeter*, from *pulsus*, 'the pulse,' and *lego*, 'I read,' *Sphygmolegium*, *Sphygmometer*, *Sphygmoscopium*. Any instrument for measuring the quickness or force of the pulse.

PULSIMANTIA, from *pulsus*, and *μαντεια*, 'divination.' The art of predicting or prognosticating by the pulse.

PULSIMETER, *Pulsilegium*.

PULSUS, *Pulse*—*p.* Acriotics, *Pulse* of irritation—*p.* *Equalis*, *Pulse*, equal—*p.* *Annuens* et *circumnuens*, *Pulse*, deficient—*p.* *Anormalis*, *Pulse*, irregular—*p.* *Bicrotus*, *Pulse*, dicrotic—*p.* *Biferiens*, *P.* dicrotic—*p.* *Caprizans*, *Pulse*, caprizant—*p.* *Cordis*, see *Heart*—*p.* *Crassus*, *Pulse*, large—*p.* *Debilis*, *Pulse*, feeble—*p.* *Deficiens*, *Pulse*, deficient—*p.* *Dicrotus*, *Pulse*, dicrotic—*p.* *Dorcadissans*, *Pulse*, caprizant—*p.* *Duplex*, *Pulse*, dicrotic—*p.* *Durus*, *Pulse*, hard—*p.* *Elasticus*, *Pulse*, supple—*p.* *Fluctuosus*, *Pulse*, undulating—*p.* *Fluens*, *Pulse*, undulating—*p.* *Formicans*, *Formicant*, and *Pulse*, formicant—*p.* *Fortis*, *Pulse*, strong—*p.* *Fusalis*, *Pulse*, caprizant—*p.* *Gazellans*, *Pulse*, caprizant—*p.* *Heterochronicus*, see *Heterochronicus*—*p.* *Humilis*, *Pulse*, low—*p.* *Incidens Solani*, *Pulse*, undulating—*p.* *Intercidens*, *Pulse*, intercurrent—*p.* *Mallearis*, *Pulse*, dicrotic—*p.* *Intercisus*, *Pulse*, dicrotic—*p.* *Intercurrens*, *Pulse*, intercurrent—*p.* *Intestinalis*, *Pulse*, dicrotic—*p.* *Intricatus*, *Pulse*, intricate—*p.* *Magnus*, *Pulse*, large—*p.* *Mollis*, *Pulse*, soft.

PULSUS MYURUS, *Pulse* *decurtate* or *sharp-tailed*, (*F.*) *Pulsus myure*. One which goes on diminishing. See *Myurus*.

PULSUS NERVINUS, *Pulse* of irritation—*p.* *Normalis*, *Pulse*, regular—*p.* *Parvus*, *Pulse*, small—*p.* *Plenus*, *Pulse*, full—*p.* *Rarus*, *Pulse*, slow.

PULSUS SERRINUS, (*F.*) *Pulsus serrin* ou *serratile*. One which strikes, sometimes strongly, at others feebly, and recalls the idea of a saw.

PULSUS STRIATUS, (*F.*) *Pulsus striat*. One that strikes the finger to a small extent in breadth.

PULSUS SUDORALIS, *Pulse* of sweat—*p.* *Tardus*, *Pulse*, slow—*p.* *Urinalis*, *Pulse*, urinal—*p.* *Uterinus*, *Pulse*, uterine.

PULSUS VACUUS, (*F.*) *Pulsus vide*. One which gives the idea of emptiness of the artery.

PULSUS VAGINALIS, *Pulse*, vaginal.

PULSUS VENA'RU'U, *Venous Pulse*, *Phlebopalia*. A pulsation occasionally noticed in the jugular veins. It is owing to some of the blood of the right ventricle reflowing into the right auricle,

and communicating an impulse to the venous blood arriving in the auricle through the descending cava. It is often a sign of impediment to the flow of blood along the pulmonary artery.

PULTA'CEOUS, *Pulta'ceus*, (F.) *Pultacé*, from *puls*, *pultis*, 'pap,' or from *pulture*, *pultat*, 'to beat.' Resembling pap. Having a softened, semifluid appearance.

PULTATIO, Atheroma.

PULTISE, Cataplasm.

PULTICULA, *Bouillie*.

PULVERES EFFERVESCENTES, Powders, soda.

PULVERIZABLE, *Pul'verable*. From *pulvis*, *pulveris*, 'dust.' Capable of being reduced to powder.

PULVERIZA'TION, *Pulverisa'tio*, *Pulvera'tio*, *Concotto'is*, (F.) *Pulvérisation*. The operation of dividing substances, so as to reduce them to powder.

Most drugs are reduced to powder by persons who make it a special business, and who are called *drug grinders*. This is done on a large scale in *drug mills*; and the implements used are the *grinding mill* and the *stamping mill*. In the shop of the apothecary, pulverization is effected by the pestle and mortar, and the sieve.

Where substances require the addition of other bodies to facilitate their reduction to powder,—as where camphor needs a few drops of alcohol—the process is called *pulverization by mediation*.

PULVEROLÉS, see *Pulvis*.

PULVERULENT, *Pulverulentus*; from *pulvis*, *pulveris*, 'dust.' Reduced to the state of a fine powder. Covered, as it were, with dust. An epithet applied to the eyes when they appear dirty, and, as it were, sprinkled with dust, by reason of granulations or grayish striae, resulting from thickening of the mucous fluid of the conjunctiva.—Nysten.

PULVILLUS, Pad, *Bourdonnet*—p. e *Lina'mentis confectus*, *Bourdonnet*—p. *Rotundus*, *Bourdonnet*.

PULVINAR, *Pulvinarium*. A pillow.

PULVINAR HUMILI, see *Humulus lupulus*.

PULVINAR MEDICA'TUM seu **MEDICINA'LE**, *Sacculus medicinalis*. A medicated pillow; as of hops.

PULVINAR PARVUM, Pad—p. *Ventriculi*, *Pancræas*.

PULVIS, *Conis*, *Powder*, (F.) *Poudre*. A substance, reduced into small particles by pulverization. *Simple powders* are those of a single substance; *compound*, those which result from a mixture of several different powders. An *impalpable powder* is one which is so fine that it can scarcely be felt under the finger. Compound powders have been called *Species*, (F.) *Especies*, *E. pulvérolentes*, *Pulvérolés*.

PULVIS ABSORBENS, P. de magnesiâ compos.—p. *Adspersorius*, *Catapasma*—p. *Alexiterius*, P. *contrayervæ* comp.—p. *Alexiterius*, P. *ipeacuanhæ* compos.—p. *Algarothi*, *Algarothi*.

PULVIS ALOES COMPOS'ITUS, P. *Aloës cum Guaiaco*, P. *aloëticus cum Guaiaco*, *Compound powder of Aloes*. (*Aloës*, 3ss, *guaiac. res.* 3j, *pulv. cinnam. comp.* 3ss. Rub the aloes and guaiacum separately, then mix the whole. *Ph. L.*) Cathartic, stomachic, and sudorific. Dose, gr. x to ʒj.

PULVIS ALOES CUM CANEL'LA, P. *aloës et canella*, (Ph. U. S.), P. *Aloëticus*, *Powder of Aloes with Canella*, *Species Hi'era pieræ*, *Hi'era pieræ*, vulg. *Hi'kry pi'kry*, *Piera*, and *Pi'kry*. (*Aloës* ʒij, *canell.* ʒijj. Rub separately into a powder, and mix. *Ph. U. S.*) A warm cathartic. Dose, gr. x to ʒj.

PULVIS ALOES CUM GUAIACO, P. A. compos.—p. *Alœticus*, P. *aloës cum canellâ*—p. *Alo-*

eticus, *Hiera pieræ*—p. *Aluminosus*, P. *sulphatis aluminæ* compos.—p. *Aluminis compositus*. P. *sulphatis aluminæ* compos.—p. *Aluminis kinosatus*, see P. *sulphatis aluminæ* compos.

PULVIS DE AMA'RIS COMPOS'ITUS, P. *antiarthriticus amarus*, (F.) *Poudre antiarthritique amère*. (*Gentian.*, *aristoloch. rotund.*, ʒa p. 3, *flor. centaur. min.* p. 4, *fol. chamædr.*, *chamæpith. manipul.*—f. *pulvis*, *Ph. P.*) In gouty affections, dyspepsia, &c. Dose, gr. xx.

PULVIS ANGELICUS, *Algarothi*—p. *Anodynus*, P. *ipeacuanhæ* comp.—p. *Antiacidus*, P. *cretæ* compos.—p. *Antiacidus*, P. de magnesiâ compos.—p. *Antiarthriticus amarus*, P. de amar-
is compos.—p. *Antiarthriticus purgans*, P. de sennâ, *scammonæâ* et lignis—p. *Antiasthmaticus*, P. de sulphure et seillâ.

PULVIS ANTILYS'SUS, *Mead's powder against the bite of a mad dog*. It consisted of ash-coloured liverwort in powder, with an equal quantity of black pepper; a drachm and a half of the two forming a dose for an adult.

PULVIS ANTIMONIALIS, *Antimonial powder*.

PULVIS DE ARO COMPOS'ITUS, *Compound Powder of Arum*, (F.) *Poudre de Birekman*, (*Rud. ari.*, *acori veri*, *pimpinell. saxifrag.* ʒa 40 p., *ocul. caneror.* ppt. 12 p., *cinnam.* 7 p., *potassæ sulph.* 6 p., *amoniac.* muriat. 2 p.—f. *pulv.* *Ph. P.*) Sternutatory.

PULVIS AROMATICUS, P. *cinnamomi* compos.

PULVIS AS'ARI COMPOS'ITUS, P. *sternutatorius* seu *erph'ius* seu *capitalis Sancti Angeli* seu *cephalicus*, *Cephalic sniff*, *Compound powder of Asarabacca*, (F.) *Poudre d'asarum composée dite sternutatoire*, P. de *Saint-Ange*. (*Asar. Europ.* p. 3, *origan. majoranæ*, *lavand.*, sing. p. 1. Rub into powder. *Ph. L.*) Sternutatory.

PULVIS AURI, see Gold—p. e *Bolo* compos.—p. *Bolo* compos.—p. *cretæ* compos.—p. *Bolo* compos.—p. *Cambogiæ* compos.—p. *e. gummi guttâ*—p. *Capitalis Sancti Angeli*, *Pulvis asari* compos.—p. *Carbonatis calcis* compos.—p. *cretæ* compos.—p. *Cardiacus*, P. *contrayervæ* comp.—p. *Carthusianorum*, *Antimonii sulphuretum rubrum*.

PULVIS CAT'ECU COMPOS'ITUS, *Compound Powder of Catechu*. (*Catechu*, kino, ʒa ʒij; *cinnam.*, myrist., ʒa ʒss.) Dose, as an astringent, gr. xv to xxx.

PULVIS CATHARTICUS, P. *scammonææ* comp.—p. *Cephalicus*, P. *asari* compos.

PULVIS CINNAMOMI COMPOS'ITUS, *Compound Powder of Cinnamon*, *Pulvis* seu *Species Imperato'ris*, *Pulvis aromaticus* (Ph. U. S.), P. *vita Imperato'ris* seu *Dianonaton*, *Species diacinnamo'mi* seu *lærificantes Rhazie* seu *Diam'bræ sine odora'tis* seu *aroma'ticæ*, *Tra'gea aroma'ticæ*, *Aromatic powder*, (F.) *Poudre de canelle composée ou aromatique*. (*Cinnam.*, *zingib.* ʒa ʒij, *cardam.*, myrist. ʒa ʒj. Rub to a fine powder. *Ph. U. S.*) Stimulant and carminative. Dose, gr. x to ʒj.

PULVIS COBBI seu **TUNQUINENSIS**. The famous *Tonquin powder*, introduced into England by a Mr. Cobb. It was composed of 16 grains of musk, and 48 grains of cinnamon; and was mixed in a gill of arrack. It was formerly regarded as a specific in hydrophobia.

PULVIS COMITIS WARWICENSIS, P. *Cornachini*—p. *Comitisæ*, *Cinchona*.

PULVIS CONTRAYER'VÆ COMPOS'ITUS, P. *cardiacus* seu *alexiterius* seu *Mantuanus*, *Lapis contrayervæ*, *Contrayerva balla*, *Compound Powder of Contrayerva*. (*Contrayerv. contrit.* ʒv, *test. præp.* bliss. *Mix.* *Ph. L.*) Stimulant and sudorific. Dose, gr. xv to ʒss.

PULVIS CORNACHI'NI, P. *Or'ydî Stib'ti com-*

positus seu *Compositus Warwicen'sis* seu de *Tribus* seu *Scammonii antimonialis*, *Diacetylatus* seu *Paracelsi*, *Earl of Warwick's Powder*, *Cerberus Triplex*, (F.) Poudre Cornuechine. (Scammon., *putass. superlat.*, *antim. diaph.* aa p. æ. f. pulvis, Ph. P.) Cathartic. Used chiefly in cutaneous diseases. Dose, ʒss to ʒi.

PULVIS CORNU USTI CUM OPIO, *P. opiatum*, Powder of Burnt Hartshorn with opium, (F.) Poudre de corne brûlée avec l'opium. (Opium duri cont. ʒi. corn. ust. præp. ʒi, coccor. cont. ʒi. Mix. Ph. L.) Ten grains contain one of opium. Anodyne.

PULVIS CRETACEUS, *P. cretæ compositus*.

PULVIS CRETÆ COMPOSITUS, *P. Carbonatæ Calcis compositus* seu *cretæceus* seu *antac'idus* seu *terres'tris*, *Species e Scor'dio sine opio*, *S. Diacretæ*, Compound Powder of Chalk, *Dioscordium*; (with Bole Armeniac in place of chalk—the Pulvis e Bolo compositus sine opio;) (F.) Poudre de craie composée. (Cretæ ppt. lbss, cinnam. ʒiv, tormentill., acaciæ, sing. ʒiij, pip. long. ʒiv. Reduce separately to a fine powder, and mix. Ph. L.) Antacid, stomachic, absorbent. Dose, gr. xx to ʒij.

PULVIS CRETÆ COMPOSITUS CUM OPIO, Compound Powder of Chalk with opium, *P. e Cretæ compositus cum opio*; (with Bole Armeniac instead of chalk — *P. e Bolo compositus cum opio*;) *Species e Scordio cum opio*, (F.) Poudre de craie composée avec l'opium. (Pulv. cret. c. ʒvss, opii duri cont. ʒiv. Mix. Ph. L.) Contains one grain of opium in 40. Anodyne and absorbent. Dose, ʒj to ʒij.

PULVIS DIAPHORETICUS, *P. ipecacuanhæ comp.* — *p. Diapnoicus*, *P. ipecacuanhæ comp.* — *p. Diaromat.*, *P. cinnamomi compositus* — *p. Diasenæ*, *P. sennæ compos.* — *p. Diaturpethi compositus*, *P. jalapæ compositus*.

PULVIS DUCIS PORTLANDIÆ, Duke of Portland's Gout Powder, (F.) Poudre du Duc de Portland ou de la Mirandole. (Rad. aristoloch., gentianæ, summit. chamædr., centaur. min. aa p. æ.) Dose, ʒss to ʒj.

PULVIS EMMENAGOGUS, *P. myrrhæ compositus* — *p. Epilepticus Marchionum*, *P. de magnesiâ compositus* — *p. Erithrus*, *P. asari compositus* — *p. Galactopæus*, *P. de magnesiâ compos.*

PULVIS E GUMMI GUTTÆ, *P. Cambo'giæ compositus* seu *hydragogus*, *Camboe Powder*, (F.) Poudre hydragogue ou de gomme gutte ou de Quercetanus. (Rad. jalap. 24 p., rad. mechoacan. 12 p., cinnam., rhei. aa 8 p., gum. cambog. 3 p., fol. veldanellæ sic. 6 p., semin. anisi 12 p., fiat pulvis. Ph. P.) A drastic cathartic. In anasarca and in worms. Dose, gr. xv to xxx.

PULVIS GUMMI'S ALKALI'US, *Sapo vegetabilis*, (F.) Poudre de savon végétal. (Guaiac. pulv. 32 p., potass. carb. 4 p. Make into a powder. Ph. P.) Antacid.

PULVIS EX HELMINTHOCORT'O COMPOSITUS, *P. vermifugus absque mercurio*, (F.) Poudre vermifuge sans mercure. (Fuc. helminthocort., sem. contræ., summit. absinth., tanacet. fol. scordii, sennæ, rhei. aa p. æ, fiat pulvis. Ph. P.) Dose, ʒj to ʒi.

PULVIS HYDRAGOGUS, *P. e gummi guttæ* — *p. Hydrargyri cinereus*, see *Hydrargyri oxydum cinereum* — *p. Hypnoticus*, *Hydrargyri sulphuretum nigrum* — *p. Imperatoris*, *P. cinnamomi compositus* — *p. Incisivus*, *P. de sulphure et scillâ* — *p. Pro infantibus*, *P. de magnesiâ compos.* — *p. Infantum*, *P. de magnesiâ compos.* — *p. Infantum antacidus*, *P. de magnesiâ compositus*.

PULVIS IPECACUANHÆ COMPOSITUS, *P. alexiterius* seu *anod'ynus* seu *diaphoreticus* seu *diapnoicus* seu *sudorificus* *Doveri* seu *Ipecacuanhæ et Opium* (Ph. U. S.), Compound Powder of *Ipecacuanha*, *Dover's Powder*, (F.) Poudre de *Dover* ou

d'ipecacuanha et d'opium ou de *Dover*. (*Ipecacuanh. pulv.*, *opii pulv.*, sing. ʒi. *potassæ sulphat.* ʒi. Mix. Diaphoretic. Dose, gr. v to gr. xij.

PULVIS IPECACUANHÆ ET OPII, *P. ipecacuanhæ compositus*.

PULVIS JALAPÆ COMPOSITUS, *P. Diaturpethi compositus* seu *laxans* seu *purgans* seu *laxativus vegetabilis* seu *laxans cum solè* seu *Jalapæ aul'ini* seu *Jalapæ tartar'icus*, *Species Diapalapæ*, Compound Powder of *Jalap*, (F.) Poudre de *Jalap* composée. (Pulv. jalap. ʒi, potass. bitart. ʒij. Mix.) Cathartic. Dose, ʒj to ʒij.

PULVIS JALAPÆ SALINUS, *P. jalapæ compositus* — *p. Jalapæ tartaricus*, *P. jalapæ compositus* — *p. de Jalapâ et scammonio compositus*, *P. scammonæ comp.* — *p. Jamesii*, *Antimonial powder* — *p. Jesuiticus*, *Cinchona* — *p. Joannis de Vigo*, *Hydrargyri nitrico-oxydum*.

PULVIS KINO COMPOSITUS, Compound Powder of *kino*, (F.) Poudre de *kino* composée, (*Kino* ʒxv, cinnam. ʒiv, opii duri ʒi. Rub each to powder, and mix. Ph. L.) Astringent. Dose, gr. v to xx.

PULVIS LAXANS, *P. jalapæ compositus* — *p. Laxans cum sale*, *P. jalapæ compositus* — *p. Laxativus vegetabilis*, *P. jalapæ compositus*.

PULVIS DE MAGNE'SIÂ COMPOSITUS, *P. Galactopæus* seu *Nutri'um* seu *Infan'tium* seu *Infantum antac'idus* seu *pro Infan'tibus* seu *antac'idus* seu *absorbens* seu *epilepticus Marchionum*, Compound Powder of *Magnesiâ*, (F.) Poudre absorbante et anti-acide. (*Magnesiæ*, sacchar. aa p. æ. Mix. Ph. P.)

PULVIS MANTUANUS, *P. contrayervæ comp.* — *p. Marchionis*, *Marchionis Pulvis* — *p. Mercurialis cinereus*, *Hydrargyri oxydum cinereum* — *p. ad Mucum pulmonalem liberius ciendum*, *P. de sulphure et scillâ*.

PULVIS MYRRHÆ COMPOSITUS, *P. e Myrrhâ compositus* seu *ad ul'cera tend'ium* seu *emmenagogus*, Compound *Myrrh Powder*, (F.) Poudre Balsamique. (*Myrrhæ*, sabinæ, rutæ, castor. Russ. aa ʒi.) A stimulant, antispasmodic, and emmenagogue. Dose, gr. xv to xl.

PULVIS NUTRICUM, *P. de magnesiâ compositus* — *p. Opiatus*, *P. cornu usti cum opio* — *p. Oxydi stibii compositus*, *P. cornachini* — *p. ad Partum*, *Ergot* — *p. Parturiens*, *Ergot* — *p. de Phosphate calcis et stibii compositus*, *Antimonial powder* — *p. Principis*, *Hydrargyri nitrico-oxydum* — *p. Purgans*, *P. jalapæ compositus* — *p. Quercetanus*, *Hydrargyri submurias*.

PULVIS QUERCUS MARINÆ, Powder of Yellow Bladder-Wrack. (The wrack in fruit, dried, cleaned, exposed to a red heat in a crucible with a perforated lid, and then reduced to powder. Ph. D.) Virtues dependent on iodine: used, consequently, in bronchocele, &c. Dose, gr. x to xl. See *Fucus vesiculosus*.

PULVIS RHEI COMPOSITUS, Compound Powder of *Rhubarb*. (*Magnes. lbj, zingib. pulv. ʒij, rhei pulv. ʒiv. M. Ph. E.*) An antacid and laxative. Dose, ʒj to ʒij.

PULVIS SALINUS COMPOSITUS, Compound Saline Powder. (*Sodii chlorid.*, *magnesiæ sulphat.*, aa ʒiv, potass. sulphat. ʒij. Ph. E. Dose, as an aperient, two or three drachms dissolved in water.

PULVIS SCAMMONIÆ COMPOSITUS, *P. e Scammonio compositus* seu *Scammonii* seu de *Jalapâ et Scammonio* *compositus* dictus *Catharticus*, Compound Powder of *Scammony*, (F.) Poudre cathartique ou Poudre de *Scammonée* composée. (*Scammon.*, ext. jalap. dur., sing. ʒij, zingib. ʒss. Rub separately to powder, and mix. Ph. L.) Cathartic. Dose, gr. viij to gr. xv.

PULVIS SCAMMONII CUM ALOË, *P. e Scammonio cum Aloë*, *Scammony and Aloë Powder*. (*Scammon.* ʒvj, ext. jalap. dur., aloë Socotr. aa ʒiiss, zing. ʒss.) Cathartic. Dose, gr. x to xv.

PULVIS SCAMMONII ANTIMONIALIS, P. cornachini.

PULVIS SENNÆ COMPOSITUS, P. e Sennâ compositus seu e Senâ compositus seu Diarsenâ, Compound Powder of Senna, (F.) Poudre de séné composée. (Sennâ, potass. superart., sing., ʒij, scammon. ʒss, zingib. ʒij. Ph. L.) Cathartic, hydragogue. Dose, ʒi to ʒij.

PULVIS DE SENNÂ, SCAMMONIÆ ET LIGNIS COMPOSITUS dictus ANTIARTHRITICUS POTASS. (F.) Poudre de Pérad. (Gum Arab., potass. superart., fol. sennâ, chinâ, aa 4 p., scammon., rad. sarsapar., chinâ, guaiac. lign. aa 2 p., fiat pulvis. Ph. P.) Purgative. Used to obviate gout. Dose, ʒi.

PULVIS STANNI, see Tin—p. Sternutatorius, P. asari compositus—p. Stibiatum, Antimonial powder—p. Stypticus, P. Sulphatis aluminæ compositus—p. Sudorificus, P. ipecacuanhæ comp.—p. Sudorificus Doveri, P. ipecacuanhæ compositus.

PULVIS SULPHATIS ALUMINÆ COMPOSITUS, P. aluminosus seu aluminis kinosatus seu stypticus seu aluminis compositus, Alumen kinosatum, Compound Alum Powder, (F.) Poudre styptique ou d'alum composée. (Sulphatis aluminæ, p. iv, kino, p. i. Ph. E.) Astringent. In intestinal hemorrhage and diarrhœa. Dose, gr. x to xv.

PULVIS DE SULPHURÆ ET SCILLÂ, P. antiasthmaticus seu incisivus seu ad mucum pulmonalem liberius ctenidum, (F.) Poudre incisive ou anti-asthmaticque. (Sacchar., p. 3, sulphur. lot. p. 2, scillæ. siccât. p. j. Ph. P.) Dose, gr. v to ʒss.

PULVIS DE SULPHATÆ POTASSÆ COMPOSITUS, P. temperans Stahlii, (F.) Poudre tempérante de Stahl. (Potass. sulph., P. nitrat. aa 9 p., hydrarg. sulph. rubr. 2 p. Mix. Ph. P.) As an antiphlogistic, and in acidity of the first passages. Dose, ʒj to ʒij.

PULVIS E SULPHURETO HYDRARGYRI NIGRO ET SCAMMONIO, dictus VERMIFUGUS MERCURIALIS, (F.) Poudre vermifuge mercurielle. (Pulv. cornachini, hydrarg. sulphuret. nigr. recent. præp., aa p. æ. Mix. Ph. P.) Dose, ʒj to ʒss.

PULVIS SYMPATHETICUS, Sympathetic powder—p. Temperans Stahlii, P. de sulphate potassæ compositus—p. Terrestis, P. cretæ compositus.

PULVIS TRAGACANTHÆ COMPOSITUS, P. e Tragacanthâ compositus, Species Diatrageanthæ frigidiæ, Compound Powder of Tragacanth, (F.) Poudre de gomme adragant composée. (Tragacanth. contus., acaciæ cont., amyli, sing., ʒss, sacch. purif. ʒij. Ph. L.) Demulcent. In coughs, &c. Dose, ʒss to ʒij.

PULVIS DE TRIBUS, P. cornachini—p. Tonquensis, P. Cobbii—p. ad Ulcera tendinum, P. myrrhæ compositus—p. Vermifugus absque mercurio, Pulvis ex helminthocoto comp.—p. Vermifugus mercurialis, P. e sulphureto hydrargyri nigro et scammonio—p. Vitæ Imperatoris, P. cinnamomi compositus.

PUMEX, Lapis pu'micis, Pumice, (F.) Pierre Ponce. This was formerly employed as a desiccative and dentifrice.

PUMICE, Pumex.

PUMILIO, Nanus.

PUMILUS, see Nanus.

PUMMEL-FOOT, Kylosis.

PUMP, BREAST, Antlia lactea—p. Stomach, Stomach pump.

PUMPKIN, Cucurbita pepo.

PUNA, (S.) Veta, Sorocco, Soroche, Mareo; Hipson'ous, Mountain Sickness, (F.) Mal de Montagne. A sickness common in the elevated districts of South America and elsewhere, which attacks those unaccustomed to breathe the rarefied atmosphere. It strongly resembles sea-

sickness, with a most acute and depressing throbbing in the head. It does not often affect the respiration greatly.

PUNAISE, Cimex.

PUNAISE, Ozæna.

PUNCH, Rhi'zagra, from pungere, punctum, 'to prick.' Repulsorium, (F.) Repoussoir. A surgical instrument used for extracting the stumps of teeth. See Lever.

Also, an agreeable alcoholic drink, made of spirit, wine, sugar, lemon juice, and water. Milk is sometimes substituted in part for the water: it then forms *Milk Punch*.

PUNCTA OSSIFICATIONIS, Ossification, points of.

PUNCTÆ MUCOSÆ, see Acne.

PUNCTICULA, Petechiæ.

PUNCTIO, Paracentesis, Puncture—p. Nervi, Neuronygme.

PUNCTUM, Stitch.

PUNCTUM AU'REUM, Punctu'ra au'rea. Formerly, when an intestinal hernia was reduced by an incision made through the integuments down to the upper part of the spermatic vessels, a golden wire was fixed and twisted, so as to prevent the descent of any thing down the tunica vaginalis. Hence the term.

PUNCTUM CÆCUM, see Optic nerve—p. Lacrymale, Lachrymal punctum.

PUNCTUM SALIENS, P. sanguineum saliens, Globulus sanguineus, Punctus ruber saliens. The first moving point which makes its appearance after the fecundation of the germ. It has been generally, but erroneously, applied to the heart of the embryo.

PUNCTURA, Paracentesis—p. Aurea, Punctum aureum.

PUNCTURE, Punctu'ra, Punctio, Cent'esis, Paracente'sis, Nyxis, from pungere, punctum, 'to prick.' Trevis Punctura, (F.) Pique. A wound made by a pointed instrument, as a sword, bayonet, needle, thorn, &c. A puncture is, generally, of trifling consequence; but, sometimes, it becomes important from the parts implicated, and the after consequences.

PUNCTURING. Same etymon. *Compunctio, Paracente'sis, (F.) Ponction.* The operation of opening, by means of a trocar, lancet, &c., any natural or accidental cavity, with the view of evacuating any fluid, effused or accumulated therein. *Paracentesis* is a sort of puncturing.

PUNCTUS RUBER SALIENS, Punctum saliens.

PUNGENT, (F.) Pongitif. Same etymon. Pain is said to be pungent, when it seems as if a pointed instrument were forced into the suffering part.

PUNICA GRANA'TUM. The Pomegranate, *Rhoa, Rheæ, Granatum, Malo-granatum, (F.) Grenadier: Ord. Myrtacæ. Sez. Syst. Icosandria Monogynia.* The rind of the fruit, called *Malicho'rium, Malacorium, Malicorium, Grana'ti Fructus Cortex* (Ph. U. S.); and the flowers—called *Balaustine flowers*—are used in medicine. They have been employed both internally and externally as astringents. The seeds are called *Cocco'nes, κοκκωες*. The portion, however, of the plant, that has attained most celebrity, is the bark of the root, *Grana'ti Radicis Cortex* (Ph. U. S.), which appears to be very efficacious in tænia. It is given in decoction, (*Granat. rad. cort. ʒij; Aquæ Oij. boiled to Oiss.*) Dose, ʒij, every half hour. Three or four doses are usually sufficient to expel the worm.

The bark contains a peculiar matter called *Grenadin*.

PUNK, Boletus igniarius.

PUGENIA, Pyogenia.

PUPIL, Pupill'a, Pu'pula, Pupilio, Vis'io,

Opsis, *Corē*, *Cora*, *Rhoz*, *Fenes'tra* seu *Nigrum* seu *Fora'men* *Oculi*, *Prunel'la*, *Lumen*, *Glēnē*, *Sight* or *Apple* or *Ball* of the eye, (Sc.) *Blak* of the *Eie*, (Prov.) *Candle*, (F.) *Pupille*, *Prunelle*. The aperture of the iris, through which the rays of light pass that have to impress the image of an object on the retina. This aperture can be dilated or contracted so as to allow a greater or smaller quantity of luminous rays to penetrate. The pupil, in man, is round, and by it the anterior and posterior chambers of the eye communicate with each other. In the fetus, during the first six months of gestation, it is closed by the pupillary membrane.

PUPIL, CLOSURE OF THE, *Synezisis*.

PUPIL, PINHOLE. A condition of the pupil in typhus, in which it is so contracted as to resemble a pinhole. It indicates a dangerous form of brain affection.

PUPILLÆ VELUM, Pupillary membrane.

PUPILLARIS HERBA, *Lapsana*.

PUPILLARY, *Pupilla'ris*, (F.) *Pupillaire*. That which belongs to the pupil.

PUPILLARY MEMBRANE, *Membra'na Pupilla'ris*, *Pupill'æ velum*, *Membra'na Wachendorfia'na*. The central aperture of the iris is closed, in the fetus, during a certain period of gestation, by a very thin membrane. It disappears in the seventh month.

PUPPET ROOT, *Veratrum viride*.

PUPPIS OS, Frontal bone.

PUPULA, *Pupil*.

PUPULÆ, *Apices digito'rum*. The pulps or extremities of the fingers.

PURBLIND, *Myopic*.

PURBLINDNESS, *Myopia*.

PURGAMENT, *Purgative*.

PURGAMENTA, Excrement — *p. Alba*, *Leucorrhœa* — *p. Infantis*, *Meconium* — *p. Puerperii*, *Lochia* — *p. Uteri*, *Lochia*.

PURGANS MEDICAMENTUM, *Cathartic*.

PURGATIO, *Catharsis* — *p. Alvi*, *Catharsis* — *p. Muliebris alba*, *Leucorrhœa* — *p. Puerperii*, *Lochia*.

PURGATION, *Catharsis*.

PURGATIONES, *Menses* — *p. Menstruæ*, *Menses*.

PURGATIONS, *Menses*.

PURGATIVE, *Purgati'vus*, *Purgans*, *Moch'licus*, *Catoret'icus*, *Catoret'icus*, *Cathartic*, *Purgament*, *Purge*, (F.) *Purgatif*, from *purgare*, *purgatum*, 'to cleanse.' A medicine which operates more powerfully on the bowels than a laxative, stimulating the muscular, and exciting increased secretion from the mucous coat.

PURGE, *Purgative* — *p. Root*, *Euphorbia corollata*.

PURGING, *Diarrhœa*.

PURIFICANS, *Purif'icus*, from *purificare*, 'to make pure or clean.' Purifying; hence: — PURIFICANTIA, Medicines that cleanse or purify the blood.

PURIFICUS, *Purificans*.

PURIFORM, *Puriform'is*, from *pus*, *puris*, and *forma*. Having the appearance of pus.

PURL. A medicated malt liquor, in which wormwood and aromatics are infused.

PURO-HEPATITIS, see *Hepatitis*.

PURO-MUCOUS, *Puro-muco'vus*, from *pus*, *puris*, and *mucus*. Having the character or appearance of pus and mucus.

PURPLE OF CASSIUS, (F.) *Pourpre de Cassius*. A precipitate, obtained by adding the protochloride of tin to a dilute solution of gold; or by dipping rods of tin in a solution of gold in diluted chlorohydric acid. Used in syphilis.

PURPLES, *Purpura hæmorrhagica*.

PCR/PURA, *Por'phyra*, 'a purple colour.' This word has been employed in many senses.

It generally means, livid spots on the skin from extravasated blood, with languor and loss of muscular strength, pains in the limbs. — (F.) *Pourpre*.

In PURPURA SIMPLEX, *Porph'yra simplex*, *Pete'chiæ* *sinē* *febrē*, *Pelio'is*, *Phœnig'mus* *Pete'chia'lis*, *Profus'io* *subcuta'nea*, *Pete'chial* *Scurvy*, the spots are numerous, but small and flea-bite-shaped; chiefly on the breast, arms, and legs; with paleness of countenance. At times, the affection is accompanied by severe pains in the extremities, constituting *Pelio'is rheumat'ica*.

In PURPURA HÆMORRHAG'ICA, *Porph'yra hæmorrhag'ica*, *Stomac'ace* *universa'lis*, *Hæmorrhagia* *universa'lis*, *Pete'chiano'is*, *Morbus* *maculo'sus* *hæmorrhag'icus* *Werthof'ii*, *Pelio'is* *hæmorrhag'ica*, *Scorbut'us*, *Schorbut'us*, *Hæmorrhœa* *petechia'lis*, *Hæmatospill'ia*, *Land* *Scurvy*, *Purples*, (F.) *Scorbut*, *S. de Terre*, *Maladie* *de Werthof*, *Hémacélinose*, *Pélioë* *hémorrhagique* — the spots are circular, and of different sizes; often in stripes or patches, irregularly scattered over the thighs, arms, and trunk; with occasional hæmorrhage from the mouth, nostrils, or viscera, and great debility and depression of spirits.

In PURPURA NAU'TICA, *Porphyra nautica*, *Scorbut'us*, *Pechtytyrbē*, *Gingibra'chian*, *Gingipe'dium*, *Scelotyr'bē*, *Stomac'acē*, *Scorbut'us* *nauticus*, *Sea* *scurvy*, (Sc.) *Scrubie*, (F.) *Scorbut*, there are spots of different hues, intermixed with the livid, principally at the roots of the hair; the teeth are loose; the gums are spongy and bleeding; the breath is fetid, and the debility universal and extreme. This occurs chiefly at sea, after exposure to a moist, cold, foul atmosphere; with long use of one kind of food and of stagnant water.

All these varieties of purpura considerably resemble each other. In the first two, however, there seems, at times, to be some excitement, which has led to the employment of purgatives, and even of bleeding, with advantage. They are dependent upon a peculiar diathesis, which has been termed *Scorbutic Cachex'ia*, *Lues Scorbut'ica*, and *Cacochyn'ia Scorbut'ica*. Generally speaking, the use of the mineral acids will be indicated; and, in sea scurvy, vegetable acids must be regarded as the best antiscorbutics. Nitre is, also, a good antiscorbutic, and in cases of great prostration it may be united with cinchona or turpentine. Tamarinds and cream of tartar, made into drink, are also serviceable. But the great remedy is change from low to cheerful spirits; from filth to cleanliness; and from a cold and damp, to a temperate and dry climate, with change of food, and especially the use of green vegetables.

The general principles of management are the same in sea, as in land scurvy.

PURPURA, *Scarlatina* — *p. Alba* *benigna* *et* *maligna*, *Miliary* *fever* — *p. Maligna*, *Pete'chiæ* — *p. Miliaris*, *Miliary* *fever* — *p. Puerperarum*, *Miliary* *fever* — *p. Urticata*, *Urticaria*.

PURPURIC URINE, see *Porphyria*.

PURPURINE, see *Porphyria*.

PURPURISSUM, *Hydrargyri sulphuretum rubrum*.

PURRING TREMOR, *Frémissement cataire*.

PURSE, *Scrotum* — *p. Shepherd's*, *Thlaspi bursa*.

PUR'SINESS, *Dyspnœa pinguedino'sa*, (F.) *Pousse*. *Dyspnœa* accompanied with oppressive fatness. The word is, also, applied to *dyspnœa* of every kind. One so affected is said to be *pursey*, or *short-winded*, (Prov.) *Short-of-puff*, *Thick-listed*, (F.) *Poussif*.

PURSLAIN, *Purslane*.

PURSLANE, *Portulaca* — *p. Milk*, *Euphorbia corollata* and *E. maculata* — *p. Speedwell*, *Ve-*

onica peregrina — p. Water, *Ludwigia palustris*, *Veronica beccabunga*.

PURSY, see Pursiness.

PURULENCE, Pus, Suppuration.

PURULENCY, Pus, Suppuration.

PURULENT, *Purulentus*, *Em'pyos*, *Pyo'des*, *P'y'icus*, from *pus*, *puris*. That which has the character of pus or consists of pus; as *purulent matter*, *a purulent collection*, &c.

PURULENT DEPOSIT, Pus.

PURULENTIA, Suppuration.

PURVAIN, *Verbena officinalis*.

PUS, *Humor purulentus*, *Pyon*, *Pyono'ma*, *Matter*, *Pu'rulent depos'it*, *Pu'rulence*, *Pu'rulency*, (Sc.) *Woursaum*, *Worsum*. A secretion from inflamed textures, and especially from the areolar membrane. It is, almost always, of the same nature, whatever may be the part it proceeds from.

Pus of a good quality, — *laudable pus*. *Pus bonum seu laudabil'e seu nutritivum seu verum*, (F.) *Pus louable*, — is of a yellowish-white colour, opaque, inodorous, and of a creamy appearance. Heat, acids, and alcohol coagulate it. When analyzed by Schwilgué, it afforded albumen and water, a particular extractive substance, and a small quantity of soda, phosphate of lime, and other salts. Normal pus consists essentially of two distinct parts, *pus corpuscles* or *pus globules*, *Cytoid corpuscles*, (Henle) — and a colourless, aqueous fluid, *liquor puris*, in which the corpuscles are suspended. A variety of the pus corpuscles is described by M. Lebert under the name *pyoid*. Fully formed pus is aplastic.

PUS BONUM, see Pus — p. Corpuscles, see Pus — p. Corrosivum, Sanies — p. Globules, see Pus — p. Laudable, see Pus — p. *Louable*, see Pus — p. Malignum, Ichor — p. Malignum, Sanies — p. Nutritivum, see Pus — p. Verum, see Pus.

PUSIL, Pustule.

PUSILLA'TUM, *Pusula'tum*, from *pusillum*, 'small' [?]. A coarse powder.

PUSILLUS, Nanus.

PUSKILE, Pustule.

PUSTULA, Pustule — p. Ardens, Eczema — p. Oris, Aphthæ — p. Gangrenosa, see Anthrax — p. Lidaa Estoniæ, see Anthrax.

PUSTULE SICCE, Lichen.

PUSTULAR, *Pustulous*, *Pustula'ris*, *Pustulo'sus*, (F.) *Pustuleux*. Of or belonging to pustules. Affected with pustules.

PUSTULA'TION, *Pustula'tio*, from *pustula*, 'a pustule.' The formation of pustules.

PUSTULE, *Puskile*, *Push*, *Pus'tula*, from *pus*. *Ecpy'e'sis*, *Ecthy'ma*, *Benath*, *Bub'e*, *Epana's'tasis*. An elevation of the cuticle, with an inflamed base, containing pus. Willan and Bateman reckon four varieties of pustules, *Phlyza'cium*, *Paydra'cium*, *Achor*, and *Favus*. The violaceous pustule is often called, by the French, *Grain*; by us, *Pock*.

PUSTULE MALIGNA, see Anthrax.

PUSTULOUS, Pustular.

PUSULATUM, Pusillatum,

PUT TO BED, Deliver.

PUTAMEN OVI, see Membrana putaminis.

PUTIUM, Prepuce.

PUTREDO, Hospital Gangrene, Putrefaction.

PUTREFAC'TION, *Putrefac'tio*, *Putrescen'tia*, *Sepsis*, *Sap'rotes*, *Se'pedon*, *Putrid* or *Putrefac'tive fermentation*, from *putris*, *putre*, 'rotten,' and *facere*, 'to make.' A decomposition, experienced by animal substances, when deprived of life, and placed under special circumstances. The presence of water is indispensable, the temperature most favourable to it, is from 60° to 90° of Fahrenheit. The most common products of putrefaction are, water, carbonic acid, acetic acid, ammonia, carburetted hydrogen, and a semi-putrid substance, which is volatilized, and has an

infected odour. It has been supposed that something like putrefaction may occur in the living body, as in case of putrid fever. To this condition, Piorry has given the name *Typho'hæ'mia*, conceiving it to be owing to alteration of the blood by putrid animal matters. *Putrefaction* signifies the state of becoming putrid: whilst *Putridity*, *Putre'do*, *Putrid'itas*, *Pu'tridness*, (F.) *Pow'rature*, is the state of rottenness.

PUTRESCENT, Hyposaprus — p. Poison, see Poison.

PUTRESCENTIA, Putrefaction — p. Uteri gravidi, Hystero-malacia.

PUTRID, *Pu'tridus*, *Sapros*, *Sepedono'des*, (F.) *Putride*, from *putreo*, 'to rot.' An epithet for some affections, in which the matters excreted and the transpiration itself exhale a smell of putridity. It is particularly applied to typhus.

PUTRIDITAS, see Putrefaction.

PUTRIDITY, see Putrefaction.

PUTRILA'GO, *Putror*, (F.) *Putrilage*. Same etymon. The putrescent matter or slough, which forms in certain gangrenous and other ulcers, and is thrown off.

PUTROR, Putrilage.

PUTZUCHILL, see Myroxylon Peruiferum.

PUXIRI, see Pichurim beans.

PYÆMIA, Pyohæmia.

PYAR'THRUM, from πον, 'pus,' and αρθρον, 'joint.' A suppurated joint.

PYCNANTHEMUM INCA'NUM, (πυχνος, 'dense,' and ανθεμον, 'a blossom.') *Clinopod'ium inca'num*, *Common Mountain Mint*, *Wild Basil*. An indigenous plant of the mint family — *Ord*. Labiata — which flowers in August. It has the aromatic properties of the mints.

PYCNANTHEMUM LINIFO'LIUM, *Virginia Thyme*, has like virtues.

PYCNIMASTICA, Incrassantia.

PYCNOSIS, Inspissatio.

PYCNOTICA, Incrassantia.

PYEC'CHYSIS, from πον, 'pus,' and εκχυσις, 'effusion.' An effusion of pus.

PYELITIS, (F.) *Pyélite*, from πυελος, 'pelvis,' and itis, denoting inflammation. Inflammation of the pelvis and calices of the kidney.

PYELOMETER, Pelvimeter.

PYELONEPHRITIS, from πυελος, 'pelvis,' νεφρος, 'kidney,' and itis, denoting inflammation. Inflammation of the kidney, and of the pelvis and calices,

PYELOS, Pelvis.

PYEMESIS, Pyoemesis.

PYESIS, Suppuration.

PYETIA, Colostrum, Rennet.

PYGE, Nates.

PYGMÆUS, Pygmy.

PYGMIE, Fist, Forearm.

PYGMÉE, Pygmy.

PYGMY, *Pigmy*, *Pygma'us*, (F.) *Pygmée*, from πύγμα, 'the fist; as big as the fist.' A dwarf. The Pygmæi were a fabulous nation of dwarfs in the extreme parts of India; or, according to others, in Ethiopia.

PYGODID'YMUS, *Did'yimus symphyoperina'us*, *Pygop'ages*, from πυγη, 'the nates,' and διδυμος, 'a twin.' A monstrosity, in which twins are united by the sacrum and coccyx.

PYGOM'ELES, (F.) *Pygonèle*, from πυγη, 'the nates,' and μέλος, 'unprofitable.' A monster with excessive growth on the nates or rump.

PYGOPAGES, Pygodidymus.

PYICUS, Purulent.

PYLE, Porta.

PYLEMPHRA'IS, from πύλη, 'porta,' 'a gate,' and εμφρα'ις, 'obstruction.' Obstruction of the vena porta.

PYLEPHLEBITIS, from πύλη, 'porta,' φλεψ,

φλέβος, 'a vein,' and *itis*, denoting inflammation. Inflammation of the vena porta.

PYLORIC, *Pyloricus*, (F.) *Pylorique*. Same etymon as pylorus. That which relates to the pylorus. An epithet given to different parts.

PYLORIC ARTERY, *Arteria Coronaria dextra*, (F.) *Petite artère gastrique droite*, arises from the hepatic, and is distributed to the pylorus and to the lesser curvature of the stomach, anastomosing, particularly, with the A. coronaria ventriculi and A. gastro-epiploica dextra.

PYLORIC VEIN follows the same distribution as the artery.

PYLORIQUE, *Pyloric*.

PYLO'RUS, *Jan'itor*, *Portona'rium*, from *πύλος*, 'a gate,' and *ovos*, 'a guardian.' *Ostium ventriculi duodena'le* seu *pyloricum* seu *dextrum* seu *inferius*, *Orific'ium dextrum* seu *inferius* seu *Egreesus* seu *Ostium inferius* seu *An'nulus* seu *Sphincter ventriculi*, *Ostia'rius*, *Hostia'rius*, (F.) *Pylor*. The lower or right orifice of the stomach—*Orifice intestinal* (Ch.)—is called *Pylorus*, because it closes the entrance into the intestinal canal, and is furnished with a circular, flattened, fibro-mucous ring, which occasions the total closure of the stomach during digestion in that organ. This ring has been called the *Valve of the Pylorus*, *Sphincter Pylo'ri*, *Pylo'rus propriè sic dictus*, (F.) *Valvule du pylore*. It is a fold of the mucous and muscular membranes of the stomach; and is the *Pyloric muscle* of some authors.

PYO, in composition, means 'pus,' from *πύον*, 'pus.'

PYOBLENNICUS, *Mucopurulent*.

PYOBLENNORRHE'A, from *pyo*, and *βλένω*, 'mucus,' and *ρῶω*, 'I flow.' A discharge of puriform mucus.

PYOCENO'SIS, from *pyo*, and *κένωσις*, 'evacuation.' Evacuation of pus.

PYOCHÉZ'IA, *Diarrhœa purulen'ta*, from *pyo*, and *χέζω*, 'I go to stool.' Discharge of pus by stool. *Purulent diarrhœa*.

PYOCÉ'LIA, *Ascites purulen'tus*, from *pyo*, and *κοίλια*, 'the belly.' Pus in the abdominal cavity.

PYOCYSTIS, *Vomica*.

PYODES, *Purulent*.

PYGEDE'MA, from *pyo*, and *οίδημα*, 'swelling.' Tumefaction of the surface owing to effusion of pus.

PYOËM'ESIS, *Pym'esis*, *Vom'itus purulent'us*, from *pyo*, and *εμεσις*, 'the act of vomiting.' Vomiting of pus.

PYOGENESIS, *Pyogenia*.

PYOGEN'IA, *Puogen'ia*, *Pyogen'esis*, *Pyopo'z'esis*, from *pyo*, and *γενεσις*, 'generation.' Generation of pus. The theory or mechanism of the formation of pus. Some have believed that pus is formed by the arterial system, and is deposited, by way of excretion, in the inflamed parts; others, that it is formed by the destruction of solid parts. It seems to be owing to a degeneration of the liquor sanguinis and exudation corpuscles.

PYGENIA CORROSIVA, *Ulceration*—p. Simplex, *Suppuration*.

PYOGEN'IC, *Pyogenicus*. Same etymon. Having relation to the formation of pus.

PYOHE'MIA, *Pyæ'mia*, (F.) *Pyohémie*, from *pyo*, and *αἷμα*, 'blood.' Alteration of the blood by pus, giving occasion to the *Diath'esis seu Infectio purulen'ta*.

PYOID CORPUSCLES or **GLOBULES**, see *Pus*.

PYOMETRA, *P. purulen'ta*, from *pyo*, and *μήτρα*, 'womb.' A collection of pus in the uterus.

PYOMETRA PURULENTA, *Pyometra*.

PYON, *Pus*.

PYONÉPHRITE, see *Nephritis*.

PYONOMA, *Pus*.

PYOO'TORRHŒA, *Otiorrhœa*.

PYOPERICAR'DIA, *Pyopericar'dium*, from *pyo*, *περί*, 'about,' and *καρδία*, 'the heart.' A collection of pus in the pericardium.

PYOPHTHALMIA, *Hypopyon*—p. Neonatorum, see *Ophthalmia (purulenta infantum)*.

PYOPLEURITE, *Empyema*.

PYOPŒSIS, *Pyogenia*.

PYOP'TYSIS, from *pyo*, and *πτύω*, 'I spit.' Spitting of pus.

PYORRHAG'IA, from *pyo*, and *ρᾶγν*, 'violent rupture.' A sudden discharge of a large collection of matter.

PYORRHŒA, from *pyo*, and *ρῶω*, 'I flow.' A discharge of pus.

PYORRHŒA ALVINA, *Pyochezia*—p. Aurium, *Otiorrhœa*—p. Nasalis, *Ozena*—p. Palpebræ, *Blepharopyorrhœa*—p. Urinalis, *Pyuria*—p. Vitarum lacrymalium, *Dacryopyorrhœa*.

PYOSIS, *Hypopyon*, *Pterygion*, *Suppuration*—p. Pectoris, *Empyema*.

PYOTHORAX VERUS, *Empyema*.

PYOTORRHŒE, *Otiorrhœa*.

PYOTURIA, *Pyuria*.

PYR, *Fever*.

PYRA, *Anthrax*—p. Crustumina, *Crustumina*—p. Crustumerina, *Crustumina*.

PYR'A MID, *Pyramis*, *πυραμς*, *πυραμίδος*, *Eminen'tia pyramida'lis tym'pani*, from *πύρ*, 'flame,' which has a conical appearance [?]. A small bony projection in the cavity of the tympanum, which is excavated, to lodge the muscles of the stapes. The *Centre-pin* of the trephine. A small, obtusely pointed eminence of the inferior vermiciform process of the cerebellum.

PYRAMID, see *Calumba*—p. of *Malacarne*, *Vermiform process*, *inferior*.

PYRAMIDAL DE L'ABDOMEN, *Pyramidalis abdominis*—p. de la *Cuisse*, *Pyramidalis*—p. du *Nez*, *Pyramidalis nasi*—p. *Stapédien*, *Stapedius*.

PYRAMIDALE, (OS.) *Cuneiform*, (bone.)

PYRAMIDA' LIS ABDOM'INIS, *Musculus Fallo'pii seu succenturiatus seu auxilia'rius*, (F.) *Pubio-ombilical*, *Pubio-sous-ombilical* (Ch.), *Pyramidal de l'abdomen*. A small, fleshy, pyramidal fasciculus, whose existence is not constant; and which is inserted, by its base, into the upper part of the pubis, and terminates, by its summit, at the inferior part of the linea alba. When it contracts, it stretches the linea alba.

PYRAMIDA' LIS, P. Fem'oris, *Iliacus exter'nus*, *Pyriform'is*, *Piriformis*, *Primus et superior quadrigen'inus*, (F.) *Sacro-trochan'térien* (Ch.), *Pyramidal de la cuisse*. A flat, triangular muscle, situate at the posterior part of the pelvis. By its base, it is attached to the anterior surface of the sacrum; to the corresponding surface of the great sacro-sciatic ligament and to the posterior part of the ilium. It terminates by a tendon, which is attached to the inner surface of the great trochanter. This muscle is a rotator of the thigh outwards, or of the pelvis in an opposite direction.

PYRAMIDALIS FEMORIS, *Pyramidalis*.

PYRAMIDA' LIS NASI, *Procer'us nasi*, *Nasum dilata'tans*, (F.) *Fronto-nasal* (Ch.), *Pyramidal du nez*. A thin, triangular muscle, which has its summit upwards. Above, it is continuous with the occipito-frontalis. It descends vertically before the root of the nose, on which it terminates, becoming confounded with the transversalis. This muscle wrinkles the skin of the root of the nose transversely, and stretches that which covers the tip.

PYRAMIDES, *Corpora pyramidalia*.

PYRAMIDES ANTÉRIEURES, *Corpora pyramidalia*—p. Ferreini, *Ferrein*, *pyramids of*

—p. Malpighianæ, Papillæ of the kidney.—p. *Postérieures du cerveau*, Corpora restiformia.

PYRAMIDS OF MALPIGHI, Papillæ of the kidney.—p. of Ferrein, see Ferrein.—p. Posterior, Corpora restiformia.—p. of Wistar, Sphenoidal cornua.

PYRAMIS, Penis, Pyramid.—p. Cochleæ, Modiolus.—p. Trigona, see Temporal bone.

PYRÆA, Incendium.

PYREC'TICA, 'febrile,' from *πυρετος*, 'fever,' and *εχω*, 'I have.' Fevers: the first order in the class *Hæmatica* of Goo.

PYRÊLAINES, Olea empyreumatica.

PYRENOIDES, Odontoid.

PYRÊTHRE, Anthemis pyrethrum.

PYRETHRUM, Anthemis pyrethrum, Urtica.—p. Parthenium, Matricaria parthenium.—p. Sylvestre, Achillea ptarmica.

PYRETICUS, from *πυρ*, 'fire,' *πυρετος*, 'fever,' *Morbus febrilis*. A febrile affection.

PYRETICUS, Febrifuge, Febrile.

PYRETOGENETIC, Stimulant.

PYRETOGRAPHIA, from *πυρετος*, 'fever,' and *γραφω*, 'a description.' A description of fever.

PYRETOL'OGY, *Pyretology*, from *πυρετος*, 'fever,' and *λογος*, 'a discourse.' *Pyrol'ogy*. A treatise on fevers. A title sometimes given to a monograph on fever.

PYRETOS, Fever.

PYRETOTYPOSIS, Intermittent.

PYREX'IA, *Pyrex'ia*, *πυρεξις*, (*πυρ*, 'fire,') Fever. The febrile condition. Also, a paroxysm of fever.—*Paroxysmus febrilis*.

PYREX'IE, Febrile diseases. The first class of Cullen's Nosology.

PYRI MARTIALES, Ferrum tartarizatum.

PYRIA, Fomentation, Vaporarium.

PYRIATERIUM, Vaporarium.

PYRICAUSTUM, Burn.

PYRIFORMIS, Pyramidalis.

PYRIPH'LEGES, *πυριπλεκεις*, *Pyriphlectos*, *πυριπλεκτος*, from *πυρ*, 'fire,' and *πλεγω*, 'I burn.' One who has a burning fever.

PYRITES, IRON, Ferri sulphuretum.

PYRMONT, MINERAL WATERS OF. A celebrated mineral spring on the Weser, in Westphalia, four leagues from Hamelot. The waters are tonic, and contain chlorides of sodium and magnesium, sulphates of soda and magnesia, and carbonates of iron, lime, and magnesia dissolved in carbonic acid, with some resinous (?) principles. It is a highly carbonated chalybeate.

ARTIFIC'IAL PYRMONT WATER may be formed of Epsom salt, gr. xv; common salt, gr. v; magnesia, gr. x; iron-filings, gr. v; water, Oijj. Impregnated with the gas from marble-powder and sulphuric acid, aa ʒviij.

PYRÆNUS, Spiritus rectificatus.

PYR'OLA, *P. rotundifolia*, *Round-leaved Pyrola* or *Wintergreen*; *Shin leaf*. *Ord. Ericacæ*. Diminutive of *Pyrus*, a pear tree, from a fancied resemblance of the foliage. This plant was once used as a gentle astringent and tonic.

PYR'OLA MACULA'TA, *Chimaph'ila Macula'ta*, *Spotted Pipsi'sewa*, *Spotted Wintergreen*, *Whiteleaf*, *White Pipsi'sewa*, *King cure*, *Ground holly*, *Rheumatism weed*, &c., has similar properties.

PYROLA, ROUNDLEAVED, *Pyrola*.

PYROLA UMBELLA'TA, *Chimaph'ila seu Chimaph'ila umbella'ta*, *C. Corymbo'sa*, *Ground-holly*, *Pipsi'sewa*, *Wintergreen*, *Rheumatism weed*, *Prince's pine*, *Herbe a pisser* (Canada). This common American plant is allied to the *uva ursi*, in botanical as well as medical qualities. It is tonic and diuretic. The bruised leaves irritate the skin. From the decoction or infusion an agreeable beverage may be formed by adding sugar,

ginger to flavour it, and yeast to produce fermentation. Under the name *Pipsi'sewa Beer* it has been used in scrofulous diseases.

PYROLEUM OSSIIUM RECTIFICATUM, Oleum animale Dippelii.

PYROLIG'NEOUS ACID, *Ac'idum pyroaceticum seu pyrolig'neum seu pyrolig'num seu pyroxylicum seu ligni pyro-oleo'sum seu aceticum empyreumaticum seu lig'neum seu ligno-rum empyreumaticum*, *Pyrolig'nic acid*, (F.) *Acide pyroacétique ou pyrolignique ou pyrolique*, *Vinaigre de bois*. An acid, obtained by distillation from wood, coal, &c., which was once believed to be of a peculiar character. It is now known to be formed of acetic acid and an empyreumatic oil, from which it may be freed; and, in this state, is sold in commerce for *Acetum distillatum*, *Acetic acid of commerce*, *Ac'idum aceticum e Ligno vena'te*. It is used for similar purposes as distilled vinegar, and for preserving animal substances. It has also been used as an antiseptic in gangrene and sphacelus, and to fowl ulcers; wherever indeed creasote is indicated.—The dose of the impure pyroligneous acid internally is from five to thirty drops, three or four times a day.

PYROMA'NIA, *Incen'diary monoma'nia*, from *πυρ*, 'fire,' and *mania*. Insanity, with an irresistible desire to destroy by fire.

PYROPH'AGUS, *Igni'vorus*, from *πυρ*, 'fire,' and *φαγω*, 'I eat.' One who eats fire. A juggler, who eats bodies in a state of ignition.

PYROS, Triticum.

PYRO'SIS, from *πυρ*, 'fire,' 'the act of burning,' *Pyralis'mus pyros'icus*, *Limo'sis cardial'is eputato'ria*, *Ardor seu Ebulli'tio stom'achi seu ventriculi*, *Cardial'gia eputato'ria*, *Pyrosis Suec'ica*, *Apoeno'sis vom'itus pyrosis*, *Dyspeps'ia pyrosis*, *Unimoreg'mia*, *Orez'is* (of some), *Soda*, *Waterbrash*, *Waterqualm*, *Waterpang*, *Black-water*, (F.) *Ardur de l'estomac*, *Fer chaud*, *Cremason*, *Gorgosset*. This affection consists of a hot sensation in the stomach, with eructations of an acrid, burning liquid, that causes a distressing sensation in the parts over which it passes. Attention to diet, and the use of bitter absorbents, will usually relieve, when the disease is functional. Occasionally, it is symptomatic of organic disease of the stomach.

Pyrosis also signified, of old, inflammation, gangrene, and a burn.

PYROSIS SUECICA, Pyrosis.

PYROSOPHIA, Chymistry.

PYROTECHNIA, Chymistry.

PYROTECH'NY, *Pyrotech'nè*, *Pyrotech'nia*, from *πυρ*, 'fire,' and *τεχνη*, 'art.' Art of making fire-works, &c. Also, chymistry. M. Percy has used the term *Pyrotechnic Chirurgicale* for the art of applying fire or the actual cautery in surgery.

PYROTHONDE, from *πυρ*, 'fire,' and *θωον*, 'linen.' *Liquor pyro-oleo'sus e linte'o purat'us*. A kind of pyrogenous or empyreumatic oil, produced by the combustion of textures of hemp, linen, or cotton in a copper vessel. The brown product is acid. Its medical properties probably resemble those of creasote. Diluted with three or four times its weight of water, it has been used as a gargle in cynanche tonsillaris, but is not now employed.

It is said, also, to be a popular remedy for toothach and skin diseases. When prepared from rags, it is called *Rag oil*; when from paper, *Paper oil*, (F.) *Huile de Papier*.

PYROTICUS, Caustic.

PYROXYLIN, Gun cotton, see Collodion.

PYRUS AMERICANA, *Sorbus Americana*.

PYRUS CYDO'NIA, *Cydo'nia malifolia seu vulg'aris*, *Sorbus Cydo'nia*, *Cydo'nia*, *Coto'nia*,

Quince Tree, (F.) *Cognassier*, *Cognassier*. Ord. Rosaceæ. Sex. Syst. Icosandria Pentagynia. The fruit is termed *Cydonium malum*, *Malum canum* seu *coto-neum*, *Melum cydonium*, *Quince*, (F.) *Coign*. Quince-seeds—*Cydonium* (Ph. U. S. 1851)—are sometimes used in decoction as a demulcent. The fruit makes a pleasant preserve.

PYRUS DOMESTICA, *Sorbus domestica*.

PYRUS MALUS, *Malus*, *M. dasyphylla* seu *communis* seu *sylvestris*, *Sorbus malus*, *Melea*, the *Apple Tree*, (F.) *Pommier*. The apple—(F.) *Pomme*—is an agreeable fruit when ripe. When unripe, it disagrees, especially with those of weak digestive powers. The same may be said of the pear, (F.) *Poire*; the fruit of *Pyrus communis*, *P. Sorbus* seu *Sativa*, *Pirus*, *Apios*, (F.) *Poirier*.

Apple Brandy is a spirituous liquor obtained from the juice of the apple, and much used in the United States.

PYTIA, *Colostrum*, *Rennet*.

PYTISMA, *Sputum*.

PYUL'CON, *Pyul'eum*, from πύον, 'pus,' and ἔλκω, 'I extract.' (F.) *Tire-pus*. A surgical instrument for extracting pus from any cavity. Many pyulæ have been employed, all of which act as exhausting syringes.

PYURIA, *Pyotu'ria*, *Pyornhæ'a urina'lis*, (F.) *Pyurie*, *Pissement de Pus*, from πύον, 'pus,' and οὖρον, 'urine.' Discharge of purulent urine: a symptom which occurs in many diseases; particularly in cases of renal calculi, and in organic affections of the bladder. It receives various epithets, according to the part whence the pus proceeds:—as *Pyuria urethra'lis*, *P. rena'lis*, and *P. vesica'lis*.

PYURIA CHYLOSA, *Chyluria*—p. *Lactea*, *Chyluria*—p. *Mucosa*, *Cystirrheæ*—p. *Serosa*, *Cystirrheæ*—p. *Viscida*, *Cystirrheæ*.

PYXIS, πύξις. A box; a pill-box. Also, an emollient ointment.—*Aëtius*, and *Paulus*.

PYXIS, *Boîte*, see *Cotyloid*.

Q.

Q., see Abbreviation.

QUABEBE, *Piper cubeba*.

QUACK, *Charlatan*.

QUACK-MEDICINE, *Arcanum*.

QUACKSALVER, *Charlatan*.

QUADRANS, *Quarta pars libræ*. A quarter of a pound. Three ounces (Troy.)

QUADRA'TUS, (F.) *Carré*, from *quadrus*, 'square.' A name for muscles whose figure approaches a square. Depressor Labii inferioris, *Quadratus femoris*.

QUADRATUS DORSI, *Q. lumborum*.

QUADRA'TUS FEM'ORIS, *Quadratus*, *Quartus quadrigem'inus quadratus*, (F.) *Tuber-ischio-trochantérien* (Ch.), *Ischio-sous-trochantérien*, *Carré de la Cuissée*. A muscle situate at the posterior and upper part of the thigh. It is thin, flat, and quadrilateral, and is attached to the outer part of the tuberosity of the ischium; whence it proceeds transversely to the inferior part of the posterior margin of the great trochanter. It rotates the limb, turning the toe outwards.

QUADRATUS GENÆ, *Platysma myoides*—q. *Radii*, *Pronator radii quadratus*.

QUADRATUS LUMBO'RUM, *Quadratus seu Lumbaris externus*, *Q. dorsi*, *Flectens par lumbo'rum*, (F.) *Ilio-costal*, (Ch.) *Ilio-lombi-costal*, *Carré des lombes*. A flat, somewhat thick, irregularly quadrilateral muscle, situate in the loins, at the sides of the vertebral column. It is attached, below, to the crista ili and ilio-lumbar ligament; and above, to the inferior margin of the last rib; and besides, to the base of the transverse processes of the last four lumbar vertebræ. This muscle inclines the loins to its side; depresses the last rib, and is thus a muscle of respiration. It can also raise the hip.

QUADRIGA, *Cataptracta*.

QUADRIGEM'INA TUBER'cula, *Eminentia Quadrigem'ina seu Bigem'ina*, *Optic Lobes*, *Optic Ganglia*, *Pons Syl'vii*, *Corpora quadrigem'ina seu bigem'ina*, *Nates et Testes*, (F.) *Tubercules quadrijumeaux ou bijumeaux*, *Eminences bigeminées*, *Lobes optiques ou bijumeaux*. Four medullary tubercles, situate at the posterior surface of the tuber annulare. They are white, externally; gray, internally; oblong; rounded;

connected by pairs, and separated by two grooves, which divide them crucially. Of these tubercles—the superior called *Nates*, *Na'tulæ*, *Tuber'cula anterio'ra*, *Glu'tia*, are larger and more prominent than the lower, called *Testes*, *Tuber'cula posterio'ra*. The pineal gland corresponds exactly to the point of intersection of the two furrows that separate them.

QUADRIGEMINUS PRIMUS ET SUPERIOR, *Pyramidalis*—q. *Quartus quadratus*, *Quadratus femoris*—q. *Secundus et tertius*, *Ischio-trochanterianus*.

QUADRIJUMEAUX (TUBERCULES), *Quadrigenima Tubercula*.

QUADROON, see *Mulatto*—q. *Black*, see *Mulatto*.

QUAFADIL, *Melanthium Virginicum*.

QUAHUIL, *Cacao*.

QUAIL, *Tetrao coturnix*.

QUAMASH, *EASTERN*, *Scilla esculenta*.

QUANDROS. Ancient name for a precious stone, believed to exist in the brain of the vulture; and to which was attributed the property of augmenting the secretion of milk, and preserving from deadly accidents. No such stone exists.

QUAR'ANTINE, *Quar'entine*, *Quarentaine*, from (L.) *Quaranti'na*, itself from *quar'anta*, 'forty,' (F.) *Quarantaine*. The time during which travellers, returning from a country where the plague or any other pestilential disease exists, are compelled to remain on board ship or in a lazaretto. They, as well as the ship, are said 'to perform quarantine.' Quarantine is sometimes extended beyond forty days, whilst its duration is frequently much less. It is sometimes enforced by land; for example, when contagious diseases require the establishment of *cordons sanitaires*, and when persons are desirous of leaving infected districts. During quarantine, no communication is allowed between the travellers and the inhabitants of the country. Quarantine is a great impediment to commerce; and is, doubtless, very frequently enforced when there is but little danger of contagion. The evils, however, that might arise from the introduction of a widely-spreading contagious disease, are so disastrous, as to induce

legislators to maintain these laws in perhaps an unnecessary degree of rigour.

QUARTAN, from *quartus*, 'the fourth.' *Quartana Febris*, *Anetus Quartanus*, *Tetartæus*, *Quartan Ague*, (F.) *Quarte*. An intermittent, the paroxysms of which recur every fourth day, leaving between them two days' interval. We speak also of the *Quartan Type*. A *Double Quartan*, (F.) *Double quarte*, is one in which one paroxysm occurs on the third and another on the second day, so that there is only one day of apyrexia and two of paroxysms in succession;—the paroxysms of every fourth day resembling each other. A *Triple Quartan* is one in which there is a paroxysm every day; and on every three days the paroxysms correspond. A *redoubtling* or *repeating quartan*, (F.) *Fidère quarte doublée*, is one in which there are two paroxysms every fourth day. See *Intermittent*.

QUARTANUS REMITENS, *Tetartophia*.

QUARTARIUS. An ancient measure, equal to one-fourth of the sextarius. See *Weights and Measures*.

QUARTE, *Quartan*.

QUARTERON, see *Mulatto*—*q.* *Black*, see *Mulatto*.

QUARTIO, *Astragalus*.

QUARTI-STERNAL, *Quadristeralia*. The fourth osseous portion of the sternum, corresponding to the fourth intercostal space.—*Béclard*.

QUASSATIO, *Conquassation*.

QUASSATURA, *Conquassation*.

QUASSIA, *Q. amara* seu *excellentia*, *Picraea excelens*, *Febrifugum lignum*, *Quassy*, *Lusty Bitterwood Tree*, *Bitter-ash*, (F.) *Quassia*, *Bois amer*. *Ord.* *Simarubaceæ*. A West India tree, from which is obtained the *Lignum Quassia*, *Quassia* (Ph. U. S.). It is an intensely durable bitter, without astringency; and has been employed as a tonic and stomachic. It is generally given in the form of infusion. The bitter principle is called *Quassin*.

QUASSIA SIMAROU'BA, *Simarouba*, *Simaru'ba*, *S. officinalis*, *Econ'ymus*, *Bitter Simarouba*, *Mountain Damson*. The bark of the root, *Simaru'ba*, (Ph. U. S.), is bitter, and not unpleasant. Both water and alcohol extract its virtues. It is not astringent. It is exhibited wherever a simple tonic is required.

QUASSIN, see *Quassia*.

QUATERNIO, *Astragalus*.

QUATRE ÉPICES, see *Myrtus pimenta*.

QUATRIO, *Astragalus*.

QUEASINESS, *Nausea*.

QUEEN OF THE MEADOW, *Spiræa lobata*—*q.* of the *Prairie*, *Spiræa ulmaria*.

QUEEN'S DELIGHT, *Stillingia*—*q.* *Root*, *Stillingia*.

QUELLES, *Elixir*.

QUENOUILLE D'EAU, *Typha latifolia*.

QUERCERA, *Epalos*.

QUERCINI HERBA, *Lichen plicatus*.

QUERCULA MINOR, *Teucrium chamædrys*.

QUERCUS ALBA, *White Oak*; QUERCUS ROBUR, *Common Oak*; Q. TINCTORIA, *Black Oak*; Q. RUBRA MONTANA, *Red Oak* or *Spanish Oak*. *Ord.* *Cupuliferæ*. *Sex.* *Syet*. *Monœcia Polyandria*; (F.) *Chêne*. The bark of all these varieties is, probably, alike in medicinal properties. It is powerfully astringent and tonic, and has been used internally in intermittents, and externally in decoction, as an astringent wash, where such is indicated.

Acorns, *Cæces*, *Nuces Quercus*, (F.) *Glands*, were at one time much used as food; and a decoction of them was once recommended in dysentery and diarrhoea, in glandular obstructions, &c. A pessary made of them has been advised in immoderate flow of the menses. They are not now used.

QUERCUS INFECTORIA. A tree of Asia Minor, which affords *Turkey* or *Nut Galls*, *Galla Turcica* seu *Quercina* seu *Tinctoria*, *Nux Galla*, *Galla max'ima orbiculata*, *Cicis*, *Cassanobis*, *Galle*, *Galla*, *Gall-nut*, (F.) *Noix de Galle*, *Galle de chêne*. The *nut-gall*—*Galla* (Ph. U. S.)—is a morbid protuberance or tubercle, produced by the puncture of the *Diplolepis* seu *Cynips Galla Tinctoria*, *Cynips Quercus folii*. It is powerfully astringent and tonic, and has been employed in diarrhoea, intestinal hemorrhage, and intermittents: but is chiefly used, externally, in gargles and injections. The powder, made into an ointment with lard, is used in piles. The ordinary nutgalls of the shops are procured from the Levant. The large *Mecca* or *Bussorah galls*, at times called *Dead Sea apples*, *Mad apples* and *apples of Sodom*, are produced on *Quercus infectoria*.

There are upwards of thirty species of oak in the United States, all of which are probably more or less astringent and tonic.

QUERCUS COCCIFERA, see *Kermes*—*q.* *Marina*, *Fucus vesiculosus*—*q.* *Suber*, see *Suber*.

QUERQUERA, *Epalos*.

QUEUE DE CHEVAL, *Hippuris vulgaris* *Cauda equina*, see *Medulla spinalis*—*q.* *de la Moëlle épinière*, *Cauda equina*—*q.* *de Pourceau*, *Peucedanum*—*q.* *de Rat*, *Sium lunare*.

QUICK-IN-THE-HAND, *Impatiens*.

QUICKEN TREE, *Sorbus acuparia*.

QUICKENING, from *Sax. epcæan*, 'to make alive.' The period of gestation when the motion of the fœtus first becomes perceptible. This usually occurs about the eighteenth week, and was at one time erroneously believed to mark the time at which the fœtus becomes alive—*Animatio fœtus*. It need scarcely be said, that it possesses the principle of life from the moment of the union of the materials furnished by the sexes as a fecundating copulation. When the motion is felt, the female is said to be 'quick with child.'

QUICKENS, *Triticum repens*.

QUICKLIME, *Calx viva*.

QUICKSILVER, *Hydrargyrum*.

QUIES, *Acinesia*.

QUINA, *Quinine*—*q.* *de Campo*, *Strychnos pseudo-china*—*q.* *Quina*, *Cinchona*.

QUINÆ ACETAS, SULPHAS, &c., see *Quinæ Acetas*, &c.

QUINCE, *Scrofula*, see *Pyrus cydonia*—*q.* *Tree*, *Pyrus cydonia*.

QUINIA, *Quinine*—*q.* *Acetate* of, *Quinæ acetat*—*q.* *Amorphous*, see *Quinia*, *extract* of, and *Chinoidine*—*q.* *Arsenate* of, *Quinæ arsenias*—*q.* and *Cinchonia*, *tannate* of, *Quinæ et Cinchonæ tannas*—*q.* *Arsenite* of, *Quinæ arsenis*—*q.* *Citrate* of, *Quinæ citras*.

QUINIA. EXTRACT OF, *Quinæ sulphas impurus*. A preparation which consists in evaporating the liquor poured off the crystals of sulphate of quinia, to the consistence of a pilular mass.

It has the properties of sulphate of quinia, and its active principle appears to be amorphous quinia or chinoidine. Twenty-four grains will generally arrest ague.

QUINIA, FERROCYANATE, OF, *Quinæ ferrocyanas*—*q.* *Hydriodate* of, *Quinæ hydriodas*—*q.* *Impure sulphate* of, *Quinia*, *extract* of—*q.* *Iodhydrate* of, *Quinæ hydriodas*—*q.* *Iodide* of, *Quinæ hydriodas*—*q.* *Iodide* of *iodhydrate* of, see *Quinæ hydriodas*—*q.* *Ioduret* of, *Quinæ hydriodas*—*q.* and *Iron*, *Hydriodate* of, see *Quinia*, *Iodide* of—*q.* *Iodhydrate* of—*q.* *Lactate* of, *Quinæ lactas*—*q.* and *Mercury*, *protochloride* of, *Hydrargyri et Quinæ Protochloridum*—*q.* *Muriate* of, *Quinæ murias*—*q.* *Nitrate* of, *Quinæ nitras*—*q.* *Phosphate* of, *Quinæ phosphas*—*q.* *Sulphate* of, *Quinæ sulphate* of.

QUININÆ ACETAS, *Chini'num acet'icum*, *Acet'as Chini'i* seu *Chini'ni* seu *Chini'nicus* seu *Qui'ninæ* seu *Quinina* seu *Quini'ni* seu *Kini'ni*, *Quina acet'ica*, *Acetate of Quinia* or of *Quinine*. Prepared by saturating *Quinia* with diluted *acetic acid*. Has the properties of the other salts of *Quinia*.

QUININÆ ARSEN'IAS, *Quini'næ* seu *Qui'næ Arsen'ias*, *Chini'num Arsenico'sum* seu *Arsenic'icum*, *Arseniate of Quinia* or of *Quinine*, (F.) *Arséniate de quinine*. Formed by the union of *arsenic acid* and *quinia*. Employed as an antiperiodic in the dose of from three quarters of a grain to a grain and a half in the twenty-four hours.

QUININÆ AR'SENIS, *Q. Diar'senis*, *Ar'senite of Quinia*. A salt resulting from the double decomposition of *arseniate of potassa* and *sulphate of quinia*. It has been used in chronic cutaneous diseases; and as an antiperiodic in ague, neuralgia, &c. Dose, one-third of a grain, three times a day.

QUININÆ CITRAS, *Quinæ* seu *Chini'ni citras*, *Citras chini'nicus* seu *qui'nicus*, *Chini'num* seu *Chini'num Citricum*, *Citrate of Quinia* or of *Quinine*. Formed from the union of *citric acid* and *quinia*. It has the same properties as the sulphate.

QUININÆ ET CINCHO'NIÆ TANNAS, *Chini'num* seu *Chini'num tan'nicum*, *Tannate of Quinia* and *Cincho'nia*. This salt has the same properties as the salts of *quinia* in general.

QUININÆ DIARSENIS, *Quinina Arsenis*.

QUININÆ FERROCY'ANAS, *Chini'num ferrocyanogenat'um* seu *ferrohydrocyan'icum*, *Ferrocyanate* or *hydrocyanoferrate of Quinia* or of *Quinine*. This salt is obtained by the decomposition of *sulphate of quinia* by means of a solution of *ferrocyanuret of potassium*. It has all the properties of *sulphate of quinia*, and, according to some, in a superior degree.

QUININÆ HYDR'ODAS, *Chini'num hydriod'icum*, *Hydriodate* or *Iodhydrate of Quinia*; called, also, *Ioduretum* seu *Iod'idum Qui'næ*, *Iod'uret* or *Iodide of Quinia*, (F.) *Hydriodate* or *Iodhydrate* ou *Iodure de Quinine*. This is formed by precipitating *sulphate of quinia* by means of *iodide of potassium*. Given in scrofulous affections; half a grain to a grain, three times a day, to a child.

An *iodide* and a *biniodide* have been formed; the latter made by mixing double the quantity of the *iodide of potassium* with the *sulphate of quinia*.

An **IODIDE OF IODHY'DRATE OF QUI'NIA**, (F.) *Iodure d'Iodhydrate de Quinine*, is prepared by pouring into an acid solution of *quinia* a solution of *iodide of iron*, containing a slight excess of *iodine*. The precipitate is treated with boiling alcohol: the liquor filtered, and crystals suffered to form. The salt possesses all the properties of the other salts of *quinia*. Under the name, *Hydriodate of iron and Quinia*, a preparation has been introduced by Mr. Battley. It is possessed of tonic and antiperiodic virtues.

QUININÆ IODIDUM, *Quinina hydriodas*—q. *Ioduretum*, *Quinina hydriodas*.

QUININÆ LACTAS, *Quinina lactas*, *Chini'num lac'ticum*, *Lactate of Quinia* or of *Quinine*, (F.) *Lactate de quinine*. Made by the action of *lactic acid* on *quinia*. Used in the same cases as the other salts of *quinia*.

QUININÆ MURIAS, *Chini'num muriat'icum* seu *ant'itum* seu *hydrochlor'icum*, *Muri'as chini'nicus*, *Muriate of Quinia* or of *Quinine*. May be prepared by dissolving *quinia* in dilute *muriatic acid*. Given as an antiperiodic. Dose, half a grain to a grain.

QUININÆ NITRAS, *Chini'num* seu *Chini'num ni'tricum*, *Nitrate of Quinia* or of *Quinine*, (F.) *Nitrate* ou *Azotate de quinine*. May be formed

by the addition of *nitric acid* to *quinia*. Its properties are those of the *sulphate of quinia*.

QUININÆ PHOSPHAS, *Chini'num* seu *Chini'num phosphor'icum*, *Phosphate of Quinia* or of *Quinine*; may be prepared by the addition of dilute *phosphoric acid* to *quinia*. It is thought by many to be next to the *sulphate of quinia* in its remedial powers.

QUININÆ SULPHAS, *Quinine*, *sulphate of*.

QUININÆ VALERIA'NAS, *Chini'num* seu *Chini'num Valerian'icum*, *Valerianate of Quinia*, of *Quina* or of *Quinine*; may be formed by the union of *valerianic acid* with *quinia*. It has the properties of the salts of *quinia*.

QUININICINE, *Quinici'na*, *Quinic'ia*. When a salt of *quinia* is exposed to heat with certain precautions, the alkaloid is changed into another isomeric with it, to which this name has been given.

Quinicine, and its *sulphate*, possess the same medical properties as *quinia* and its salts. Dose, gr. v to gr. xv.

QUINID'IA, *Quinidi'na*, *Qui'nidine*, *Chini'dine*, *Cin'chotin*. An alkaloid of the *cinchonas*—of many of the genuine at least—obtained from them by the same process as that by which *quinia* is obtained from the barks that yield it. Its *sulphate* is more soluble however than the *sulphate of quinia*, and it is left, therefore, in the mother waters.

The salts of *quinidia* are markedly antiperiodic. From gr. x to gr. xxx will usually arrest a paroxysm.

QUININE, *Qui'nia*, *Quina*, *Kini'num*, *Ki'num*, *Sal essential'e cor'ticis Peruvia'ni*, *Chini'num*, *Quinina*. An alkaline, uncrystallizable substance; under the form of a porous, whitish mass; almost insoluble in water, soluble in alcohol and ether. It forms, with acids, salts that are in general soluble. It is obtained from different *cinchonas*, but chiefly from the *yellow*, and is the active principle of those valuable drugs.

QUININE, ACETATE OF, *Quinina acet'as*—q. *Arseniate of Quinina arsenias*—q. *Arsenite of Quinina arsenis*—q. *Azotate of Quinina nitras*—q. *Citrate of Quinina citras*—q. *Ferrocyanate of Quinina ferrocyanas*—q. *Hydriodate de Quinina hydriodas*—q. *Iodhydrate de Quinina hydriodas*—q. *Iodide of Quinina hydriodas*—q. *Iodure de Quinina hydriodas*—q. *Iodure d'iodhydrate de Quinina*, *iodide of iodhydrate of*—q. *Lactate of Quinina lactas*—q. *Muriate of Quinina murias*—q. *Nitrate of Quinina nitras*—q. *Phosphate of Quinina phosphas*.

QUININE, QUINA OR QUINIA, SULPHATE OF, *Quinina* seu *Qui'næ* seu *Quini'næ Sulphas*, *Quina Disul'phas*, *Chini'num* seu *Chini'num Sulphur'icum*, (F.) *Sulfate de Quinine*—the salt usually employed in medicine—occurs in needles of a pearly and satiny appearance. It is employed with great success in the treatment of intermittents; and is available in many cases, where the bark in substance could not be retained, or would be injurious. Dose, as a tonic from 3 to 10 grains in the 24 hours. As an antiperiodic it may be carried much farther. Its action, in a large dose, is decidedly sedative. It is obtained, by treating the *yellow bark* with *sulphuric acid*. The following form is adopted in the *Pharmacopœia of the United States*,—*Cinchon. flav.* in pulv. crass. lbiv; *Acid. muriat.* fʒij; *Calcis*, in pulv. ʒv; *Aqua*, cong. v; *Acid. Sulphur.*, Alcohol; *Carbon. animal.* aa q. s. Boil the bark in one-third of the water mixed with one-third of the *muriatic acid*, and strain through linen. Boil the residue twice successively with the same quantity of acid and water as before, and strain. Mix the decoctions, and, while the liquor is hot, gradually add the lime, previously mixed with two pints of

water, stirring constantly, until the quinia is completely precipitated. Wash the precipitate with distilled water, and having pressed and dried it, digest it in boiling alcohol. Pour off the liquor and repeat the digestion several times, until the alcohol is no longer rendered bitter. Mix the liquors, and distil off the alcohol, until a brown viscid mass remains. Upon this substance, removed from the vessel, pour about half a gallon of distilled water, and having heated the mixture to the boiling point, add as much sulphuric acid as may be necessary to dissolve the impure alkali. Then add an ounce and a half of animal charcoal; boil for two minutes; filter the liquor while hot, and set it aside to crystallize. Should the liquor, before filtration, be entirely neutral, acidulate it very slightly with sulphuric acid. Should it, on the contrary, change the colour of litmus paper to a bright red, add more animal charcoal. Separate the crystals from the liquor, dissolve them in boiling water slightly acidulated with sulphuric acid, add a little animal charcoal, filter and set aside to crystallize. Wrap the crystals in bibulous paper, and dry with a gentle heat. The mother-waters may be made to yield an additional quantity of sulphate of quinia by precipitating the quinia with solution of ammonia, and treating the precipitated alkali with water, sulphuric acid, and animal charcoal, as before.

QUININE, TANNATE OF, Quiniae et cinchonae tannas—q. Valerianate of, Quiniae valerianas.

QUININISM, *Quininismus*, *Quinism*, *Cinchonism*, (F.) *Inebriation ou Ivresse quinique*. The aggregate of encephalic or neuropathic phenomena induced by over-doses of quinia.

QUINOIDINE, Chinioidine.

QUINQUEFOLIUM, *Potentilla reptans*—*Quinquefolium majus*, *Potentilla reptans*.

QUINQUINA, *Cinchona*—q. *Aromatique*, *Croton casearilla*—q. *Bicolor*, *Cinchona* *Caribaea* cortex—q. *Faux de Virginie*, *Cusparia febrifuga*—q. *Gris de Loxa*, *Cinchona* *lancifolia* cortex—

q. *Jaune*, *Cinchona* *cordifolia* cortex—q. *Orange*, *Cinchona* *lancifolia* cortex—q. *Piton*, *Cinchona* *Caribaea* cortex—q. *Rouge*, *Cinchona* *oblongifolia* cortex.

QUINQUINO, *Myroxylon periferum*.

QUINSY, *Cynanche*, *C. tonsillaris*—q. *Nervous*, *Angone*—q. *Wolf*, *Lycanebe*.

QUINTAN, *Febris quinta* seu *pempta* seu *quinta*; from *quintus*, 'a fifth.' A fever whose paroxysms return only every five days inclusively; and between which there are, consequently, three days of apyrexia.

QUINTE, (F.) This word is used synonymously with *paroxysm* or *fit*, when speaking of cough,—as *une Quinte de Tour*—'a fit of coughing,' (Sc.) *Kink*. Also, the influenza.

QUINTEFEUILLE, *Potentilla reptans*.

QUINTERON, see *Mulatto*—q. *black*, see *Mulatto*.

QUINTES, see *Quinte*, and *Pertussis*.

QUINTESSENCE, *Quinta Essentia*. A name formerly given to alcohol, impregnated with the principles of some drug. It was, consequently, often synonymous with *Alcoholic tincture*. The most volatile principles of bodies were, also, called *Quintessences*.

QUINTISTERNAL. The fifth osseous portion of the sternum. It corresponds to the 5th and 6th intercostal spaces.

QUINUA, *Chenopodium quinoa*.

QUOTIDIAN, *Quotidianus*, *Ephe'merus*, (F.) *Quotidien*—itself from *quotidie*, 'daily.' That which takes place every day.

QUOTIDIAN FEVER, *Q. ague*, *Febris quotidiana* seu *hemere'sia* seu *hemere'ria* seu *ephe'mera* seu *metheme'ria* seu *metheme'ria*, *An'etus quotidiana*, *Cathe'merus*, *Cathemer'inus*, *Methemer'inus*, *Amphemer'inus*, *Amphimer'inus*, *Amphemer'inus*, *Quotidianus*, (F.) *Fievre quotidienne*, is an intermittent, the paroxysms of which recur every day.

A *simple*, *double*, or *triple quotidian*, is a quotidian, which has one, two, or three paroxysms in the 24 hours.

R.

R or R is placed at the commencement of a prescription, as a contraction of *Rec'ipē*. (See Abbreviation.) Originally, it was the sign ♃ of Jupiter, and was placed at the top of a formula, to propitiate the king of the gods, that the compound might act favorably.

RABDOIDES, (SUTURA,) Sagittal suture.

RABID, *Rab'idus*, (F.) *Rabique*, *Rabétique*, from *rabies*, (*rabo*, 'I am mad or furious,') 'canine madness.' That which belongs to hydrophobia; as *rabid virus*, &c. Rabid also means affected with hydrophobia, (F.) *Enragé*.

RABIÉQUE, Rabid.

RABIES CANINA, Hydrophobia.

RABIQUE, Rabid.

RACAHOUT, *Amylum quer'neum*. A name given by the Arabs to the starch prepared from an edible acorn obtained from *Quercus flex*, but, according to Plagge, the so called *Racahout* of the Arabs, sold in Paris, is a compound of potatoe starch, chocolate, and aromatics,—as *vanilla*.

RACCOON BERRY, *Podophyllum montanum*.

RACE, from *radice*, abl. of *radix*, 'root,' breed. The union or aggregate of individuals whose con-

formation or particular condition differs perceptibly from those of neighbouring varieties. In the human species, several races are designated. See *Homo*.

RACÉMEUX, *Racemosus*.

RACEMOUS, *Rac'emosus*, *Rac'emose*, (F.) *Racémeux*. In clusters, like grapes.

RAC'EMUS. A cluster—as of grapes.

RACHAMEL'CA, from Hebr. רחם (*rechem*), 'the womb,' and מלך (*molech*), 'a king:' *Rechamel'ca*. A peculiar formative principle, supposed, by Dœtus, to exist in the uterus.

RACHE, *Porrigo*.

RACHIAI, *Rhachitæ*.

RACHIALGIA, *Colic*, *metallie*, *Rhachiodynia*, *Vertebral disease*—r. *Mesenterica*, *Tabes mesenterica*—r. *Pictoniensium*, *Colic*, *metallie*—r. *Pictonum*, *Colic*, *metallie*.

RACHIALGIE MÉSENTÉRIQUE, *Tabes mesenterica*.

RACHIALGITIS, *Rhachialgitis*.

RACHIDIAN, *Spinal*.—r. *Arteries*, *Spinal arteries*—r. *Bulb*, see *Medulla oblongata*—r. *Canal*, *Vertebral canal*.

RACHIDIEN, Spinal.**RACHIOPHYMA**, *Rachiphyma*.**RACHIPHYMA**, *Rhachiphy'ma*, *Rachio'phy-ma*, *Tumor Dorsi*, from *ῥαχίς*, 'the spine,' and *φύμα*, 'a tumour.' A tumour on the spine or on the back.**RACHIS**, Vertebral column.**RACHIS'AGRA**, *Rhachis'agra*, *Rhach'ingra*, *Rhach'engra*, *Rhachid'agra*, *Rhach'itis spinalis*, from *ῥαχίς*, 'the spine,' and *αἴμα*, 'a seizure.' A gouty or rheumatic affection of the spine.**RACHITÆ**, *Rhachitæ*.**RACHIT'IC**, *Rachit'icus*, *Rhachit'icus*, (F.) *Rachitique*. Same etymon as the next. Relating or appertaining to, or affected with rickets. Weak, feeble in the joints; defective in development.**RACHIT'IS**, *Rhachit'is*, from *ῥαχίς*, 'the spine,' and *itis*, a suffix denoting inflammation. Inflammation of the spine. Also, *Cyrtosis Rad'chia*, *Cyrt'on'osis*, *Morbus Ang'licus*, *Osteomala'cia Infant'um*, *Tubes pector'ea*; *Innutritio ossium*, *Spina nodo'sa*, *Rachit'ismus*, *English disease*, *Rickets*; from *ῥαχίς*, 'the spine,' (F.) *Rachitieme*, *Nouure*. A disease characterized by crookedness of the long bones; swelling of their extremities; crooked spine; prominent abdomen; large head; and often precocity of intellect. It is accompanied by leanness, general debility, indigestion; and frequently induces atrophy and hectic. Rickets occurs, particularly, during the first years of existence, in weakly children, born of rickety or scrofulous parents; brought up in damp or confined situations; who are badly nourished, and do not take sufficient exercise. The progress and termination of the disease are very variable. Some children gradually recover their health: others become more and more deformed, and ultimately die of consumption, dropsy, or other organic disease. The treatment is almost wholly *hygienic*. Pure air; a healthy situation; nourishing diet; exercise; sea or common cold bathing, and tonics, afford the best prospect of success.**RACHITIS ADULTORUM**, *Mollities ossium*.**RACHITISME**, *Rachitis*.**RACHITOME**, from *ῥαχίς*, 'spine,' and *τεμνειν*, 'to cut.' An anatomical instrument for opening the medullæ spinalis without injuring the medulla.**RACHO'SIS**, *Rhacho'sis*, *Rako'sis*, from *ῥαχὼς*, or *ῥαχὼς*, 'I wrinkle.' A relaxation of the scrotum. *Propto'ma seu lax'itas Scroti*, *Scrotum pen'dulum*. An excretion of the relaxed scrotum.**RACINE À BECQUET**, *Geranium moschatum*—*r. de Bengal*, *Cassumuniar*—*r. Brésilienne*, *Ipecacuanha*—*r. de Charcis*, *Dorstenia contrayerva*—*r. de Chine*, *Smilax china*—*r. de Dracke*, *Dorstenia contrayerva*—*r. des Philippines*, *Dorstenia contrayerva*—*r. de Safran*, *Curcuma longa*—*r. de Saint Esprit*, *Angelica*—*r. Salivaire*, *Anthemis pyrethrum*—*r. de Turbeth*, *Convolvulus turpethum*.**RACINES DU CERVELET**, *Corpora restiformia*.**RACLURES DES BOYAUX**, *Ramenta intestinorum*.**RACORNISSEMENT** (F.), from *racornir*, (*cornu*, 'horn,') 'to make hard as horn.' *Cor'nification*. A physical property, possessed by animal substances, which consists in a kind of contractility, accompanied with sudden corrugation and curling. It is produced by heat, and by chemical agents, especially by the strong mineral acids.**RADCLIFFE'S ELIXIR**, see *Tinctura aloes composita*.**RADESZYGE**, *Radzyge*.**RADEZYGE**, *Radzyge*.**RADIAD**, see *Radial aspect*.**RA'DIAL**, *Radia'lis*, from *radius*, 'a bone of the forearm.' That which relates to the radius.**RADIAL ANTÉRIEUR**, *Palmaris magnus*.**RADIAL ARTERY**, *A. extern'a Cu'biti*. It arises from the brachial, at the upper and anterior part of the forearm; descends on the outer side as far as the lower part of the radius, and proceeds downwards and backwards towards the proximal extremity of the space which separates the first two metacarpal bones. It then buries itself in the palm of the hand, where it forms the *profund* or *radial* palmar arch. The radial artery gives off the *recurrent radial*, several *radio-muscular* branches, the *transverse palmar radio-carpal artery*, the *external superficial artery of the palm of the hand*, the *external dorsal of the thumb*, the *dorsalis carpi*, *dorsal interosseous artery of the index*, &c.**RADIAL ASPECT**. An aspect towards the side on which the radius is situated.—Barclay. *Radiad* is used by the same writer, adverbially, to signify 'towards the radial aspect.'**RADIAL EXTERNE PREMIER**, see *Radialis*—*r. Externe second*, see *Radialis*—*r. Grand*, see *Radialis*.**RADIAL NERVE**, *Radio-digital* (Ch.), *Spiral Nerve*, *Spiral muscular N.*, *Musculo-spiral nerve*, arises from the four inferior branches of the brachial plexus. It is, at first, situate behind the other nerves of the plexus. It then becomes engaged between the three portions of the triceps brachialis, passes behind the humerus, and descends between the supinator longus and brachialis internus, as far as opposite the upper extremity of the radius. In its course it gives numerous filaments to the triceps, supinator longus, extensor carpi radialis longior, and to the interguments. Opposite the upper extremity of the radius it divides into two branches:—the one *posterior*, which turns backwards into the substance of the supinator brevis, and when it has reached the space between the two layers of muscles on the posterior surface of the forearm, divides into a great number of filaments, which are distributed to those muscles, and to the interguments of the hand. The other branch is *anterior*: it descends before the supinator brevis and the radius; until, near the inferior third of that bone, it engages itself between the tendons of the supinator longus and extensor carpi radialis longior; and, becoming afterwards subcutaneous, divides into two branches, whose filaments are distributed to the interguments of the thumb, index, middle finger, to the outside of the ring finger, and to the first interosseous muscle of the back of the hand.**RADIAL PETIT**, see *Radialis*.**RADIAL VEINS**, *DEEP-SEATED*, follow the same course as the radial artery.**RADIA'LIS EXTER'NUS BRE'VIOR**, *Extensor Carpi Radialis Bre'vior*, *Radialis secundus*, (F.) *Second radial externe*, *Petit radial*, *Épicondylo-sus-métacarpien*, (Ch.) An extensor muscle of the wrist, situate beneath the extensor carpi radialis longior. It is attached, above, to the external tuberosity of the humerus, and terminates below, by a long tendon, inserted into the posterior part of the upper extremity of the third bone of the metacarpus. It has the same uses as the next muscle.**RADIA'LIS EXTER'NUS LON'GIOR**, *Extensor Carpi Radialis Lon'gior*, *Radialis externus primus*, (F.) *Premier radial externe*, *Grand radial*, *Huméro sus-métacarpien*, (Ch.) It is seated at the outer part of the forearm; is thin, but thicker on the outside than on the inside. It is attached, above, to the inferior part of the outer edge, and

to the corresponding tuberosity, of the humerus; and terminates, below, by a long tendon, which is attached to the superior extremity of the second bone of the metacarpus. It extends the hand on the forearm.

RADIALIS EXTERNUS PRIMUS, *R. Externus longior*—*r. Extensor longior*, *R. Externus longior*—*r. Internus*, *Palmaris magnus*—*r. Secundus*, *R. Externus longior*.

RAD'IATED, *Radia'tus*, from *radius*, 'a ray,' (F.) *Rayonné*. That which is arranged in rays; in lines, diverging from a common centre. An epithet given to several ligaments, &c.; as to those which unite the ribs to the sternum; those which unite the tibia and fibula at their inferior extremity, &c.

RADIATED SUBSTANCE OF THE KIDNEY, see *Kidney*.

RAD'ICAL, *Radica'lis*, from *radix*, *radicis*, 'a root.' A radical cure, *cura radica'lis*, is one in which the disease is destroyed, as it were, from the root. It is used in opposition to *palliative cure*.

RADICAL MOISTURE, *Humidum radicale*.

RADICAL VESSELS, *Vascular Rad'icles*, (F.) *Radicules vasculaires*, are the small vessels that take their origin in the tissues, and by their union form larger vessels.

RADICES OSSIS HYOIDEI, *Cornua ossis hyoidei*.

RADICIS'ECA, from *radix*, *radicis*, 'a root,' and *secare*, 'to cut.' One employed in collecting and preparing plants was formerly so called.

RADICULA, *Raphanus hortensis*.

RADICULES VASCULAIRES, *Radical vessels*.

RADII CILIARES, *Ciliary processes*.

RADII FRONTALI. The folds or wrinkles of the forehead.

RADIO, from *radius*. In composition, relating or belonging to the radius, a bone of the forearm.

RADIO-CARPAL, *Radio-carpia'nus*. That which relates to the radius and carpus.

RADIO-CARPAL ARTICULATION is the *wrist joint*, or articulation of the os scaphoides, os semilunare, and os cuneiforme of the carpus with the inferior surface of the radius, and the fibro-cartilage, situated below the ulna. It is strengthened by lateral, anterior, and posterior ligaments.

RADIO-CARPAL TRANSVERSE PALMAR ARTERY, (F.) *Artère Radio-carpienne-transversale-pal-maire*. A transverse branch, given off by the radial artery, opposite the lower edge of the pronator quadratus, which sends several branches to the anterior or palmar surface of the carpus.

RADIO CUTANÉ (NERF), see *Cutaneous*—*r. Phalangien du pouce*, *Flexor longus pollicis manus*.

RADIOLUS, *Sound*.

RADIO-MUSCULAR, *Radio-muscula'ris*. A name given to the branches sent off from the radial artery to the muscles of the forearm in the first part of its course; as well as to the nervous filaments which the radial nerve sends to the same muscles.

RADIS, *Raphanus hortensis*—*r. de Cheval*, *Cochlearia armoracia*.

RADISH, *Raphanus hortensis*—*r. Water*, *Nasturtium amphibium*.

RADIUS, 'a spoke'—so called from its shape. *Cervicis*, *Foc'ile minus seu superior*, *Canna minor*, *Os ad cubit'u'm*, *Additamen'tum ulnæ*, *Manu'brium manus*, *Parape'chyon*, *Arund'is brachii minor*, (F.) *Os du Rayon*. A long, prismatic bone, the upper and lesser extremity of which is called the *head*. This is supported by a *cervix* or *neck*. At the part where the neck is conjoined with the body of the bone is the *tubercle* or *bicipital*

tuberosity or *eminence* for the insertion of the biceps. The radius is articulated, above, with the os humeri and with the lesser sigmoid cavity of the ulna: below, with the scaphoides, semilunare, and the head of the ulna. Its inferior extremity, which is larger than the superior, is flattened before and behind: it is irregularly quadrilateral; and has, below, a double facette to be articulated with the first two bones of the carpus. On the outer side is the styloid process; and, on the inner, a concave facette, which joins the ulna. Behind, are grooves for the passage of the extensor tendons. The radius is developed from three points of ossification; one for the body, and one for each extremity.

RADIUS, COL DU, *Collum radii*.

RA'DIX, genitive, *radicis*, *Rhiza*. A root: also, the male organ.

Radix, *Root* or *fang* of a tooth, is the part contained in the alveolus. The *radix* or *root* of the nail is the portion hidden under the skin, &c. Surgeons give the name to prolongations, sent by scirrhus, cancerous, or other tumours into the neighbouring parts.

Five Aperient Roots, *Quinque radi'ces aperient'es*, were, of old, asparagus, butchers' broom, fennel, parsley, and smallage.

Five Lesser Aperient Roots, *Quinque radi'ces aperient'es mino'res*, were caper, dandelion, eryngo, madder, and rest-barrow.

RADIX ASCLEPIADIS CRISPA, see *Gomphocarpus crispus*—*r. Braziliensis*, *Ipecacuanha*.

RADIX CORDIS, *Basis Cordis*. The base of the heart.

RADIX GINSENG, see *Panax quinquefolium*—*r. Indica Lopeziana*, *Lopez radix*—*r. Lingue*, see *Tongue*—*r. Lopeziana*, *Lopez radix*—*r. Ninsi*, *Panax schinseng*, *Sium ninsi*—*r. Rubra*, *Rubia*—*r. Serpentinum*, see *Ophiorrhiza mungos*—*r. Unguis*, *Nail*, *root* of—*r. Ventris*, *Umbilicus*.

RADULA, *Raspatorium*.

RADZYGE, *Radzygin*, *Radesyge*, *Radezyge*, *Tha'ria*, (properly *Theria*, from *θηρion*, ('*thos*'), 'a malignant ulcer.') *Lepra borea'lis seu Norveg'ica*, *Norwegian Lep'rosy*. A name given, in Norway, to a disease, bearing considerable analogy to the yaws. Some have esteemed it a species of lepra or elephantiasis.

Another form—the *Spedalske*, or *Spedalskhd*—of Norway, appears to be a variety of tubercular lepra. Under the epithelial scales, numerous acari have been found; and the affection has been called *Scab'ies crusto'sa*, and *Scab'ies Norveg'ica Boeck'ii*, after Boeck, who well described a case of it.

RAFFLE'SIA ARNOLD'I. A plant of Java. *Order*, *Rafflesiaceæ*, a decoction of which is employed there as an astringent in relaxed conditions of the genito-urinary apparatus.

RAFRAICHISANT, *Refrigerant*.

RAGE, *Ira*, *Furor brevis*, *Orgé*, *Thymos*, (F.) *Colère*, *Rage*. Violent passion, characterized by contraction of the muscles of the face, violence in every movement, extreme irritation of the nervous system, acceleration of the blood's motion, and, generally, redness and swelling of the face.

RAGE, *Rhage*.

RAGE, *Hydrophobia*, *Rage*.

RAGE DE TÊTE. An epidemic and fatal malady, which prevailed in France in 1481. It was presumed to be epidemic meningitis.

RAGWEED, *Ambrosia elatior*—*r. Golden*, *Senecio aureus*—*r. Great*, *Ambrosia trifida*.

RAGWORT, *Senecio Jacobææ*.

RAIA BATIS, and **R. CLAVATA**, see *Oleum jecoris aselli*.

RAIDEUR, *Rigidity*—*r. Cadavérique*, *Rigor mortis*.

RAIFORT, *Raphanus hortensis*—*r. Sauvage*, *Cochlearia armoracia*.

RAINBOW WORM, *Herpes iris*.

RAINURE, Groove—*r. Digastrique*, Mastoid groove.

RAISIN, see *Vitis vinifera*—*r. d'Amérique*, *Phytolacca decandra*—*r. de Bois*, *Vaccinium myrtillus*—*r. d'Oure*, *Arbutus uva ursi*—*r. de Renard*, Paris.

RAISING, Expectoration.

RAISINÈRE (F.), from *raisin*, 'a grape.' A name given to a small granular, brownish or blackish tumour, which forms occasionally at the surface of the cornea, in consequence of ulcers or accidental wounds of that membrane.—Nysten.

RAISINS, *Uvæ passæ*, see *Vitis vinifera*—*r. de Corinthe*, see *Vitis Corinthiaca*.

RAISON, Reason.

RAKASIRA. An American balsam; of a brownish or brownish-red colour; semi-transparent; brittle; softening and becoming tenacious by heat; inodorous when cold, but exhaling a very agreeable smell when heated; and possessing a balsamic and slightly bitter taste. The precise vegetable that furnishes it is not known. It has been used chiefly as a balsam in gonorrhœa and urinary affections.

RAKE-TEETH. A term, used in the North of England for teeth which are wide apart, and similar to those of a rake.

RAKIA, see Spirit.

RAKOSIS, *Rachosis*.

RÂLE (F.), *Rattle*, *Rhonchus*, *Rhenchus*, *Rhenis*, from (D.) *Ratel*; *Rhoncus*, *Cerchnus*, *Rhogmos*: (probably an example of onomatopœia.) Noise produced by the air in passing through mucus, of which the lungs are unable to free themselves. This condition is chiefly observed at the approach of death, and is commonly called, "the rattles."

The term *Râle* has been given to different sounds during respiration, caused by the air in connection with fluid contained in the bronchi, or areolæ of the pulmonary tissue; which are perceived by the stethoscope.

RÂLE BRONCHIQUE SEC, *Râle sonore*—*r. Bronchique humide*, *R. muqueux*—*r. Caverneux*, *Gurgling*—*r. Cavernuleux*, see *Gurgling*—*r. Crépitant*, *Rhonchus crepitans*, see *Crepitant*—*r. c. Reduz*, see *Crepitant*.

RÂLE MUQUEUX, *R. bronchique humide*, *Rhonchus mucosus*, *Hygrobronchiorrhonchus*, *Mucous Rattle*. The sound produced by the passage of air through mucus accumulated in the larger bronchi or trachea, or through softened tubercular matter. This *râle* occurs in catarrh, and in softened tubercle. When carried to a very high degree, it constitutes *gurgling*, (F.) *Gargouillement*. The *subcrepitant rhonchus*, (F.) *Râle sous-crépitant*, is produced by the bubbling of air through liquid of variable consistence in the minute bronchial tubes. When heard at the base of both lungs, it is a sign of capillary bronchitis. When heard at one base only, it is generally connected with tuberculosis higher up.

RÂLE SIBILANT, *Rhonchus sibilus* seu *sib'illans*, *Sib'illant Rattle*. A slight, though prolonged, whistling sound, occurring either at the commencement of inspiration, or of expiration, owing to the presence of mucus, thin and viscid, but not abundant, which obstructs, more or less completely, the smaller bronchial ramifications. It is seated in the small tubes, and occurs in the first stage of bronchitis. The *clicking* and *whistling* varieties differ, in the former being short, the latter prolonged. The former is a quick, sharp sibilus or whistle, which ceases almost in-

stantaneously; the latter a prolonged sibilus, of less sharpness, lasting almost the whole time of the movement which it accompanies. To these may be added the *hissing* variety, the name sufficiently indicating its character.

RÂLE SONORE, *R. bronchique sec* ou *ronflant*, *Rhonchus sonorus*, *Sonorous Rattle*. A sound resembling at times the snoring—(F.) *Ronflement*—of a person asleep; at others the sound of the bass string of an instrument when rubbed with the finger, and not unfrequently the cooing—(F.) *Roucoulement*—of a dove. It seems to be caused by a contraction of the bronchial tubes, and is characteristic of chronic catarrh.

RÂLE SOUS-CRÉPITANT, see *Râle muqueux*—*r. Vésiculaire*, see *Crepitation*.

RAMEAU, Ramus.

RAMBERGE, *Mercurialis*.

RAMEN'TA INTESTINO'RUM, *Scrapings of the Intestines*, (F.) *Rachures des Boyaux*. The shreds or scrapings, as it were, of the mucous membrane of the bowels, often discharged in malignant dysentery. The evacuation in which these are contained has been termed *Diachorea mæcysmatodes*.

RAMENTUM, Fragment, Scobis.

RAMEX, Hernia—*r. Varicosus*, *Varicocele*.

RAMIFICA'TION, *Ramifica'tio*, from *ramus*, 'a branch,' and *fic*, 'to become.' A division of arteries, veins, or nerves into branches. Also, a branch itself.

RAMILLE, *Ramusculus*.

RAMOLLISSEMENT, *Mollities*—*r. du Cerveau*, *Mollities cerebri*—*r. du Cœur*, *Cardiomalacia*—*r. de l'Estomac*, *Gastromalacia*—*r. du Foie*, *Hepatomalacia*—*r. Gris*, see *Hepaticization*—*r. de la Moelle épinière*, *Mollities medullæ spinalis*—*r. Rouge*, see *Hepaticization*—*r. de l'Intestin*, *Enteromalacia*—*r. des Os*, *Mollities ossium*.

RAMPANT (F.), from *rampier*, 'to climb.' *Repens*, *Reptans*. That which creeps or crawls. The French *Bandage rampant* has the folds arranged spirally around the part, leaving intervals between them. It is a retentive bandage, but is easily displaced, and, therefore, not employed.

RAMPES DU LIMAÇON, *Scalæ* of the cochlea.

RAMPHOS, *Rostrum*.

RAMULUS, *Ramusculus*.

RAMUNCULUS, *Ramusculus*.

RAMUS, *Clados*, a branch, (F.) *Rameau*. A division of an artery, vein, or nerve. Also, the male organ.

RAMUS DURIOR SEPTIMÆ CONJUGATIONIS, *Facial nerve*.

RAMUS'ULUS, *Ram'ulus*, *Ramun'culus*, *Sur'culus*, *Cladis'cos*, diminutive of *ramus*, 'a branch.' A division of a ramus. (F.) *Ramuscule*, *Ramille*.

RANA ESCULEN'TA, 'esculent frog,' (F.) *Grenouille*. This frog and the bull-frog, *Rana Taurina*—are eaten as delicacies. They are nutritious, and easily digested.

RANA, *Ranula*.

RANCE, *Rancid*.

RANCID, *Ran'cidus*, *Sapros*, (F.) *Rance*, from *rancere*, 'to be stale.' An epithet given to fatty bodies, which have become acrid from age or the contact of air. Mild ointments may thus become very irritating, and unfit for the purposes for which they were originally intended.

RAND'IA DUMETO'RUM, *R. Spino'sa*, *Can'thinum*, *Coronatum*, *Garden'ia* seu *Posoque'ria*, *Dumeto'rum* seu *Spino'sa*. A Coromandel shrub; *Ord. Cinchonacæ*; the fruit of which, bruised and thrown on the water, intoxicates, or even

kills fish. In the form of powder, it is a powerful emetic.

RANDIA SPINOSA, R. dumetorum.

RANINE, *Rani'na*; from *Rana*, 'a frog.' The termination of the lingual artery is so called; that is, the portion of the artery which advances horizontally between the genio-glossus and lingualis muscles as far as the point of the tongue. The ranine vein follows the same course as the artery, and opens into the internal jugular or into the superior thyroid.

RANTREE, *Sorbus aucuparia*.

RAN'ULA, diminutive of *rana*, 'a frog,' so called from its resemblance; *Dysphag'ia Ranula*, *Emphrag'ma saliva'rē*, *Ba'trachos*, *Batra'chium*, *Hydroglos'sa*, *Hypoglossis*, *Hypoglossum*, *Hypoglot'tis*, *Hypogloss'ium*, *Rana*, *Frog Tongue*, (F.) *Grenouillette*. A small, soft, fluctuating, and semi-transparent tumour, which forms under the tongue, owing to the accumulation of saliva in Wharton's duct. In order to give exit to the accumulated fluid, and to establish a permanent opening for the discharge of the saliva, the cyst may be punctured, and the opening maintained by a minute cylinder, with a button at each extremity, as has been recommended by Dupuytren; or a portion of the cyst may be removed by the scissors, and the mouth be frequently washed by any astringent lotion. Sometimes, the salivary duct is found plugged by a small calculus.

RANULA LAPIDEA, see *Calculi*, salivary.

RANUNCULUS, (F.) *Renoncule*, (Sc.) *Yellow gowan*, *Ord. Ranunculacæ*; *Sex. Syst. Polyandria Polygynia*. Diminutive of *rana*, 'a frog,' because found in marshy places where frogs abound. A genus of plants, most of the species of which, when taken internally, act as acrid poisons. See *Poison*. They are, also, acrid and vesicant, when applied to the skin, and have accordingly been often used as counter-irritants. The following species answer to this description.

RANUNCULUS ABORTIVUS, *Small-flowered Crow-foot*, *Chicken-pepper*.

RANUNCULUS ACRIS, *R. pratensis* seu *Steven'ii* seu *rufulus* seu *sylvaticus* seu *Sic'ulus*, *Upright Meadow* or *Tall Crow-foot*, *Butter Cups*, (Prov.) *Craissey*, *Guilty Cups*, *Yellow-weed*, *Blisters-weed*, *Pilewort*, *Burwort*, *Meadow-bloom*, *Yellows*, (F.), *Bouton d'or*.

RANUNCULUS ALBUS, *Anemone nemorosa*.

RANUNCULUS AQUATICUS, *R. sceleratus*.

RANUNCULUS BULBO'SUS, *R. lætus*, *Ranunculus* (Ph. U. S.), *Bulbous* or *Bulbous-rooted Crow-foot*, *Butter Cups*, (F.) *Renoncule bulbeuse*, *Bassinet*.

RANUNCULUS DIGITATUS, *R. sceleratus*.

RANUNCULUS DECLINATUS, *R. flammula*.

RANUNCULUS FICARIA, *R. vernus* seu *præcox*, *Chelido'nium minus*, *Scrophula'ria minor*, *Ficaria ranunculoides* seu *verna* seu *communis*, *Chelido'nia rotundifolia minor*, *Cuscut'ia hamorrhoidalis herba*, *Ægole'thron*, *Lesser celandine*, *Pilewort*, (F.) *Petite Chelidoine*, *Petite Éclairé*, *Ficaire*, *Herbe aux hémorrhoides*. The leaves and root have been used medicinally;—the leaves as an antiscorbutic; the root as a cataplasm in piles.

RANUNCULUS FLAM'MULA, *R. declinatus*, *Flam'mula*, *Surre'ta alba*, *Smaller Water Crow-foot*, *Lesser Spearwort*, (F.) *Petite Douve*.

RANUNCULUS ILLYRICUS.

RANUNCULUS LINGUA.

RANUNCULUS LÆTUS, *R. bulbosus* — *r. Nemorosus*, *Anemone nemorosa* — *r. Palustris*, *R. sceleratus* — *r. Pratensis*, *R. acris*.

RANUNCULUS PENNSYLVANICUS, *Bristly Crow-foot*.

RANUNCULUS PRÆCOX, *R. ficaria*.

RANUNCULUS PUBES'CENS. Grows in ditches and marshes in South Africa. The expressed

juice is used by the Cape Colonists in cancerous ulcers.

RANUNCULUS REPENS, *Creeping Crow-foot* — *r. Rufulus*, *R. acris*.

RANUNCULUS SCELERA'TUS, *R. Palus'tris* seu *umbellatus* seu *digitatus* seu *aquat'icus*, *Risus Ap'ium*, *Herba Sarado'nia*, *Herba Scelera'ta*, *Hecato'nia palus'tris*, *Marsh* or *Crushed* or *Celery-leaved Crow-foot*.

RANUNCULUS SICULUS, *R. acris* — *r. Stevenii*, *R. acris* — *r. Umbellatus*, *R. sceleratus* — *r. Vernus*, *R. ficaria* — *r. Virginianus*, *Lobelia syphilitica*.

RAPA NAPUS, *Brassica rapa* — *r. Rotunda*, *Brassica rapa*.

RAPE, *Brassica napus*, *Stuprum*.

RAPHANE'DON, from *ραφανισ*, 'a radish, like a radish.' *Raphanodon*. A transverse fracture of a long bone, leaving no splinters; — what has been called, in France, *Fracture en rave*.

RAPHA'NIA, *Convulsio Raphania*, *Rapha'nia*, *Morbus spasmod'icus malignus* seu *popula'ris* seu *Sile'siacus*, *Eclamps'ia typho'des*, *Synclonus Raphania*. Linnæus gives this name to a convulsive disease, not uncommon in Germany and Sweden, and which has been attributed to the seeds of the *Raph'anus Raphanis'trum* seu *sylvest'ris*, *Raphanis'trum seg'etum* seu *arvensē*, *Rapistrum*, being mixed with the corn. The convulsions are seated in the limbs, and are attended with acute pain. See *Convulsio cerealis*.

RAPHANIA MAÏZITICA, *Pellagra*.

RAPHANIS, *Raphanus hortensis* — *r. Magna*, *Cochlearia armoracia*.

RHAPHANISTRUM ARVENSE, see *Raphania*.

RAPHANUS, *R. hortensis*.

RAPH'ANUS HORTEN'SIS, *Raphanus*, *R. sativus* seu *niger*, *Raph'anis*, *Raph'anus*, *Radic'ula*, *Rhaphys*, *Raphys*, *Rhops*, *Rapys*, *Radish*, (F.) *Raifort*, *Radis*. *Ord. Cruciferae*. *Sex. Syst. Tetradynamia Siliquosa*. The root of this plant is a common esculent. It has, also, been given as a diuretic in calculous affections. The juice, made into a syrup, is exhibited to relieve hoarseness. Radishes have long been celebrated as anti-scorbutics.

RAPHANUS MARINUS, *Cochlearia armoracia* — *r. Raphanistrum*, see *Raphania* — *r. Rusticanus*, *Cochlearia armoracia* — *r. Sylvestris*, see *Raphania*.

RAPHĒ, *Raphē*, *ραφν*, 'a suture.' A name given to prominent lines resembling a raised stitch. Such is the raphe or tramis which divides the scrotum into two parts, and which extends from the anterior part of the anus to the extremity of the penis. The name *Raphē* seu *Raphē superior* seu *Linea Media'na* seu *Sutur'a corporis callo'si exter'na* seu *chorda longitudinalis Lancis'ii* is, also, given to two prominent lines which are longitudinal and parallel, and are seen at the upper surface of the corpus callosum, extending from the anterior to the posterior extremity of that commissure.

RAPHEX, *Chenopodium vulvaria*.

RAPHIANKIS'TRON, from *ραφιον*, 'a needle,' and *αγκιστρον*, 'a hook.' An instrument employed in the formation of artificial pupil.

RAPHION, *Needle*.

RAPHIS, *Needle*.

RAPHYS, *Raphanus hortensis*.

RAPISTRUM, see *Raphania*.

RAPONTIC, *Rheum rhaponticum*.

RAPPORT (F.), *Relat'io*, *Renuncia'tio*, 'a report.' A deposition made by a physician or surgeon, to prove the condition of a person, the nature of a disease, the existence or non-existence of pregnancy, sudden or violent death, &c

Formerly, there were the *Rapports dénonciatifs*, made at the desire of the injured or their friends, and intended to point out to the judges the details of the crime or misdemeanor,—the *Rapports provisoires*, the object of which was to obtain provisions for the injured individual, both as regarded food and medicine, and to defray the expenses of prosecution;—and *Rapports mixtes*, or those which were both *dénonciatifs* and *provisoires*. At present, reports or depositions are generally divided into *judiciary*, or those which serve to throw light on the subject matter of any civil or criminal cause; and *administrative*, which afford information on subjects relating to any branch of the public administration, as on the nuisance, or otherwise, of certain establishments, the character of any epidemic, &c.

RAPPORT, Eructation, Regurgitation.

RAPTUS POST-TERGANEUS, Opisthotonos.

RAPUM MAJUS, *Brassica rapa*.

RAPUNTUM SYPHILITICUM, *Lobelia siphilitica*.

RÁPURE, *Rasura*.

RAPYS, *Raphanus hortensis*.

RAQUETTE, *Cactus opuntia*.

RAREFA'CIENS, *Rarifa'ciens*, *Aræol'icus*, (F.) *Raréfiant*, from *rarus*, 'rare,' and *facere*, 'to make.' Medicines were so termed, which were supposed to be possessed of the power of giving more bulk or extension to the blood or other humours.

RARÉFIANT, *Rarefaciens*.

RARIFACIENS, *Rarefaciens*.

RA'RITAS, (F.) *Rareté*. Rarity; scantiness; fewness.

RARITAS DEN'TIUM. A large space between the teeth, which appear to be, and may be, too few.

RASCATIO, Excretion.

RASCETA, *Carpus*, *Malleolus*—*r. Pedis*, *Tarsus*.

RASCHA, *Carpus*, *Malleolus*.

RASETA, *Carpus*, *Malleolus*.

RASETTA, *Carpus*, *Malleolus*.

RASH, Exanthem—*r. Fever*, *Scarlatina*—*r. Canker*, see *Cynanche maligna*—*r. Lichenous*, *Lichen*—*r. Mercurial*, *Eczema mercuriale*—*r. Mulberry*, see *Typhus*—*r. Nettle*, *Urticaria*—*r. Rose*, *Roseola*—*r. Summer*, *Lichen tropicus*—*r. Tooth*, *Strophulus*—*r. Wildfire*, *Ignis sylvaticus*.

RASOIR, Razor.

RASO'RISM. The Italian medical doctrine of contro-stimulus, of which Rasori was the founder.

RASORIUM, *Rasporium*.

RASPATORIUM, *Rasor'ium*, *Rat'ula*, *Scalprum rasor'ium*, *Xyster*, *Xyste'rium*, *Xystrum*, *Rad'ula*, *Runci'na*, *Runci'n'ula*, *Scalprum*, (F.) *Rugine*, *Grattoir*. An instrument used for rasping bones, to detach the periosteum,—either in certain surgical operations, or for anatomical purposes.

RASPBERRY, *Rubus idæus*—*r. Black*, see *Rubus strigosus*—*r. Ground*, *Hydrastis Canadensis*—*r. Wild*, *Rubus strigosus*.

RASPING, Eructation.

RASPS, *Rubus fruticosus*.

RASSETA PEDIS, *Tarsus*.

RASTETA, *Carpus*, *Malleolus*.

RASU'RA, *Xyema*, *Xysmos*, (F.) *Râpure*, from *radere*, *rasum*, 'to scrape.' A rasuro, or scratch, or erosion. Also, the rasping or shaving of any substance; as *Rasu'ra Cornu Cervi*, *Hartshorn shavings*.

RASURA, Scabs.

RATAFI'A, (F.) *Alcoolé sucré*, *Saccharo-alcoolé*. A liquor prepared by imparting to ardent

spirits the flavour of various kinds of fruits, adding sugar.

RATANHY, *Krameria ratanhia*—*r. of the Antilles*, *Krameria ixiina*.

RATE, Spleen.

RATELEUX, *Spleneticus*.

RÂTELIER, see *Dentier*.

RATIO, *Rationalitas*, from *reor*, *ratus*, 'to think.' A reason, a cause, an explanation, a mode or manner; as *Ratio symptom'atum*, an explanation of the symptoms,—*Ratio meden'di*, *Genus curatio'nis*, *Via curandi*; the reason, mode or manner of cure.

RATIO, Reason—*r. Medendi*, see *Ratio*—*r. Symptomatum*, see *Ratio*.

RAT'IONAL, *Rationalis*, (F.) *Rationnel*. Same etymon as *Ratio*. That which is conformable to reason. This epithet is often opposed to *empirical*; as *rational treatment*. The French, also, say *Signes rationales*, in opposition to *Signes sensibles*. See *Organic*.

RATIONALITAS, *Ratio*.

RATSBANE, *Arsenious acid*.

RATTLE, *Râle*.

RATTLES, *THE*, see *Râle*.

RATTLEBUSH, *Sophora tinctoria*.

RATTLESNAKE, *Crotalus horridus*—*r. Leaf*, *Goodyera pubescens*—*r. Plantain*, *Goodyera pubescens*—*r. Root*, *Nabalus albus*, *Polygala senega*, *Trillium latifolium*—*r. Weed*, *Hieracium venosum*.

RATTLESNAKE'S MASTER, *Agave Virginica*, *Liatris scariosa*, *Nabalus albus*.

RATTLEWEED, *Actæa racemosa*.

RATULA, *Rasporium*.

RAUCEDO, *Raucitas*, *Parapho'nia rauca*, *Asper'itas Arte'ria as'pera*, *Vox rauca*, *Branchus*, *Hoarseness*, (Sc.) *Roup*, (Prov.) *Rawp*, (F.) *Enrouement*, *Raucité*, from *raucus*, 'hoarse.' A change in the voice, which loses its smoothness and becomes low and obscure. It is a common symptom in diseases of the larynx and trachea.

RAUCEDO CATARRH'ALIS. Hoarseness from cold.

RAUCITAS, *Raucedo*.

RAUCITÉ, *Raucedo*.

RAUCOUS, *Raucus*. 'Hoarse,' as a raucous voice—*Vox rauca seu raucis'ona*.

RAVE, *Brassica rapa*.

RAVERY, *Delirium*.

RAWP, *Raucedo*.

RAY, *Diarrhoea*.

RAYONNÉ, *Radiated*.

RAYONS SOUS-IRIENS, *Ciliary processes*.

RAZOR, *Culter Tonsor'ius*, *Machæ'ris*, *Novac'ula*, *Xyrum*, *Xyrus*, (F.) *Rasoir*, from *radere*, *rasum*, 'to shave.' An instrument for shaving. Used in surgery to remove hairs, where they would interfere with any operation.

REACHING, *Vomiturition*.

REACTION, *Reactio*, from *re*, 'again,' and *agere*, *actum*, 'to act.' An action of resistance to a power applied. A species of vital movement or excitement, tending to prevent or destroy the effects of any noxious power applied to the animal economy, and which, by some, has been attributed to the *vis medicatrix naturæ*. It is certainly dependent on the organic instinctive actions of the frame. It is the state of activity which succeeds the action on the nervous system, of certain morbid influences.

READ, *Rennet*.

REAL'GAR, *Realgal*, *Arlada*, *Arladar*, *Auripigmentum rubrum*, *Arsenicum rubrum facti'tium*, *Abessi*, *Sindara'cha Græcorum*. Red Sulphuret of Mercury. A violent poison. See *Poison*.

REAM, Cream.

REASON, *Ratio*, *Logos*, (F.) *Raison*. Same stymon as *Ratio*. The faculty or property of the mind by means of which man perceives the distinction between right and wrong, in physics as well as in morals. *Raison* is a just conclusion resulting from the comparison of two ideas, by virtue of which we form a judgment.

REBOLEA, Mummy.

REBONA, Mummy.

REBOUTEUR, *Renoueur*.

RECEIVER, *Ampulla*, (F.) *Ballon*, *Récipient*, from *recipere*, *receptum*, (*re*, and *capere*, 'to take,') 'to take, to receive.' In pharmacy, a wide-necked globular vessel, the neck of which widens gradually outwards, to receive the tapering beak of the retort in distillation. Like the retort, the receiver may be plain or tubulated.

A quilled receiver is one that has a tapering tube from its lower surface, which is inserted into a graduated bottle through a cork joint, when it is desirable to know accurately the amount of fluid that has passed over.

RECEPTACULA DURÆ MATRIS, Sinuses of the dura mater.

RECEPTACULUM CHYLI, 'receptacle of the chyle;' *R. Pecqueti* seu *Pecquetianum*, *Diversorium* seu *Stagnum chyli*, *Sacculus chyli* seu *ro'ifer* seu *lacteus*, *Alveus ampullo'sus* seu *ampullæ'cena*, *Utriculus lacteus*, *Vesicula chyli*, *Ampulla chylifera* seu *chyli*, *Cisterna lumbaris* seu *chyli*, *Chylodochia*, *Chylodochium*, *Lat'ices lactei*, *Lumbar cistern*, *Reservoir of Pecquet*, (F.) *Réservoir du chyle*, *R. de Pecquet*, *Cisterna lumbaire*. A considerable dilatation of the thoracic duct, in front of the lumbar region of the vertebral column. It is the commencement of the duct.

RECEPTACULUM COTURNI. A triangular space, formed by a separation of the two laminae of the dura mater of the internal ear, near the middle of the posterior surface of the pars petrosa of the temporal bone, where the aqueductus vestibuli terminates. It is always filled with the *Fluid of Cotugno*.

RECEPTACULUM PECQUETI, *R. Chyli*—*r. Sellæ equinae lateribus appositum*, Cavernous sinus.

RECEPTIVITY, *Receptivitas*, (F.) *Réceptivité*. Same etymon as *Receiver*. A word used by Tissot to express the susceptibility of certain organs to receive morbid impressions.

RECEPTUM, Prescription.

RECESSIO, Paroxysm.

RECESSUS, Abscess, Sinus—*r. Sulciformis*, see *Fovea vestibuli*.

RECHAMALCA, *Rachamelca*.

RECHUTE, see *Relapse*.

RECIDIVA MORBI, *Relapse*.

RÉCIDIVE, see *Relapse*.

REC'IPE, (L.) (*re*, and *capere*, 'to take,') A Latin word which signifies 'take.' (F.) *Prenez*, and with which prescriptions are commenced. It is, often, represented by *R*, *R*, and was formerly by γ , the symbol of Jove. See *Abbreviation*.

RÉCIPIENT, *Receiver*.

RECIPROCATIÖN, VIBRATIONS OF, see *Sound*.

RECLINATIO, Reclination—*r. Palpebrarum*, *Ectropion*.

RECLINATIO, *Reclina'tio*, from *reclinare*, *reclinatum*, (*re*, and *clinare*, 'to bend,') 'to bend back.' A mode of operating for the cataract, which consists in applying the needle in a certain manner to the anterior surface of the cataract and depressing it into the vitreous humour, in such a way, that the front surface of the cataract is the upper one and its back surface the lower one.

RECLUSOR PALPEBRARUM, *Levator palpebrae superioris*.

RECOMPOSITION, *Metasynchrisis*.

RECONVALESCENTIA, *Convalescence*.

RECORDATIO, *Memory*.

RECORDATUS, *Memory*.

RECORPORATIVUS, *Metasynchritic*.

RECOVERY, *Restauratio*.

RECREATIO, *Restauratio*.

REC'REMENT, *Recrementum*, from *re*, and *cernere*, *cretum*, 'to sift.' A humour, which, after having been separated from the blood, is again returned to it; such as the saliva, the secretion of serous membranes, &c. Such fluids are said to be *recrementit'ial*, (F.) *Récrémenteux*, *Récrémentiel*; *Humores inquilini*.

RECREMENTA VESICÆ, *Urine*.

RECREMENTITIAL FLUIDS, see *Recrement*.

RECREMENTO-EXCREMENTIT'IAL. Applied to secreted fluids, as the saliva, bile, &c., which are in part reabsorbed and in part excreted.

RECREMENTUM, *Recrement*.

RECRUDESCENTIA, *Polincot'esia*, *Recrudescence*, from *re*, 'again,' and *crudus*, 'raw.' An increase of a disease after a temporary remission.

RECTAL, *Rectalis*. Appertaining, or relating to, the rectum,—as '*rectal tube*.' See *Tube*, *rectal*.

RECTIFICA'TION, *Rectifica'tio*, from *rectum* *facere*, 'to make right.' An operation by which certain liquids are purified by distilling them, either singly or after admixture with other substances.

RECTITIS. A hybrid word, from *rectum*, and *itis*, denoting inflammation; properly *architis*, or *proctitis*. Inflammation of the rectum. *Proctalgia inflammatoria*.

RECTO, in composition, signifies relating or belonging to the rectum.

RECTOCELE, *Cystocele*.

RECTOSTENOSIS, Stricture of the rectum.

RECTO-UTERINE, *Recto-uterinus*, from *rectum*, and *uterus*. Appertaining, or relating, to the rectum and uterus.

RECTO-UTERINE FOLDS, see *Uterus*.

RECTO-VAG'INAL, *Recto-vagina'lis*. That which relates or belongs to the rectum and vagina.

RECTO-VAGINAL FISTULA is owing to a preternatural communication between the rectum and vagina through the above septum.

RECTO-VAGINAL SEPTUM, (F.) *Cloison recto-vaginale*, is formed by the junction—back to back—(or by what the French term *adossement*) of the rectum and vagina. It separates these two canals from each other.

RECTO-VESICAL, *Recto-vesical'is*. Having relation to the rectum and bladder; as the "*recto-vesical fold or pouch*" of the peritoneum, between the rectum and bladder. See *Urinary Bladder*.

RECTUM, from *rectus*, so called from a notion that it is straight. It is not so. *Intestinum Rectum*, *Apeuthy'smenos*, *Euthy'enteron*, *Longaon*, *Longas*, *Longanon*, *Archos*, *Cys'saron*, *Princeps*, *Monocolon* [?], *Bum-gut*, (Prov.) *Jot-gut*. The third and last portion of the great intestine. It forms the continuation of the sigmoid flexure of the colon, occupies the posterior part of the pelvis, and extends from the left side of the sacro-vertebral articulation to the coccyx, before which it opens outwards by the orifice called the *anus*. It is cylindrical, but *sacculated*, as it were, at its inferior portion, where it is consequently larger, and is attached to the sacrum by the mesorectum.

Like the other portions of the canal it is formed of three membranes. Towards the lower extremity, some parallel and longitudinal wrinkles are observable on its inner surface: these are the *Columnæ, Columnæ carnea* of the *Rectum* or of *Morgagni*, (F.) *Colonnes du rectum* ou de *Morgagni*. They are merely folds of the mucous membrane, between which is a number of *Lacunæ*. The arteries of the rectum are numerous. They proceed from the inferior mesenteric, hypogastric, and internal pudic; and are called *hemorrhoidal*. The veins open, partly into the hypogastric vein, partly into the inferior mesenteric. The nerves proceed from the sciatic and hypogastric plexuses. The rectum receives the faecal matters, as they pass from the colon, which serves as a reservoir for them.

RECTUS, 'straight' (F.) Droit. A term, used in anatomy for parts that pursue a straight course; — as

RECTUS ABDOMINIS, (F.) Pubio-sternal, Sternopubii (Ch.), Droit de l'abdomen. A long flat muscle, situate at the anterior part of the abdomen, and separated from that of the opposite side by the *linea alba*. It is attached, above, to the cartilages of the last three true ribs; and, below, to the pubis by a very strong tendon, the outer edge of which is continuous with the fascia transversalis. This muscle is enclosed in an aponeurotic sheath; and, in its course, has three or four transverse aponeurotic intersections. It bends the chest on the pelvis, and conversely.

RECTUS ANTERIOR BREVIS, Rectus capitis internus minor — r. *Anterior longus, R. capitis internus major* — r. *Capitis anticus longus, R. c. internus major*.

RECTUS CAPITIS INTERNUS MAJOR, R. internus major seu *anterior longus seu capitis anticus longus, (F.) Trachélo-basilaire, Grand droit antérieur de la tête*. This muscle is long in proportion to its breadth, flat, and broader above than below. It is situate at the anterior and lateral part of the neck, and arises from the transverse processes of the third, fourth, fifth, and sixth cervical vertebrae. It is inserted into the basiliary process of the occipital bone.

RECTUS CAPITIS INTERNUS MINOR, R. internus minor seu *anterior brevis, Renuens, Annuens, (F.) Petit trachélo-basilaire, Petit droit antérieur de la tête*. This is situate deeper than the last. It is thin, flattened, irregularly quadrilateral, and passes from the anterior surface of the body of the first vertebra, near the origin of the transverse process, to be inserted into the basiliary process.

RECTUS CAPITIS LATERALIS, R. lateralis seu *lateralis Fallopii, Transversalis anticus primus, (F.) Trachélo-atloïdo-basilaire, Atloïdo-cous-occipital (Ch.), Droit latéral de la tête*. There is a muscle of this name on each side of the head. It is flat, quadrilateral, and passes from the transverse process of the atlas to the occiput, behind the jugular fossa. It inclines the head to one side.

RECTUS CAPITIS POSTICUS MAJOR, R. major seu *Capitis posticus minor (Albinus), Spino-axoïdo-occipitalis, (F.) Axoïdo-occipitalis (Ch.), Grand droit postérieur de la tête*. A flat, triangular muscle, situate at the posterior and superior part of the neck. It extends from the spinous process of the second cervical vertebra to the external surface of the lower semicircular ridge of the occipital bone. This muscle extends the head, inclines it to one side, and causes it to experience a rotary motion, which directs the face to its side.

RECTUS CAPITIS POSTICUS MINOR, R. minor, (F.) Tuber-atloïdo-occipital, Atloïdo-occipital (Ch.), Petit droit postérieur de la tête. A flat,

triangular muscle, situate beneath the last, which passes from the posterior arch of the atlas to the occipital bone. It extends the head.

RECTUS CAPITIS POSTICUS MINOR, R. c. p. major — r. *Cruris, R. femoris*.

RECTUS EXTERNUS OCULI, Abductor oculi, Indignabundus, Indignatorius, Iracundus, Oculum movens secundus, Orbito-ectus-scleroticus. All the recti muscles of the eye, four in number, arise from the bottom of the orbit, and are inserted into the sclerotic coat of the eye. The *Rectus externus, (F.) Droit externe, Irascible ou Abducteur*, is divided, at its posterior extremity, into two portions, one of which is attached to the outer part of the foramen opticum, and the other to the sphenoid bone.

RECTUS FEMORIS, Rectus seu Gracilis anterior seu cruris, (F.) Ilio-rotulien, Droit antérieur de la cuisse. A straight muscle, seated at the anterior part of the thigh. It extends from the anterior and inferior spine of the ilium and the cotyloid cavity of the patella. This muscle extends the leg upon the thigh, and conversely. It can, also, bend the thigh upon the pelvis or carry the pelvis forward.

RECTUS INFERIOR OCULI, Depressor oculi, Timidus, Deprimens, Humilis, Oculum movens quartus, (F.) Sous-optico-sphéro-scleroticien, Droit inférieur ou abaisseur de l'œil, Muscle humble, arises below the optic foramen, from the sphenoid bone. It depresses the eye.

RECTUS INTERNUS OCULI, Adducens oculi, Bibitorius, Oculi adductor, Oculum movens primus, (F.) Orbito-intus-scleroticien, Droit interne de l'œil ou adducteur ou buveur. It arises from the sphenoid at the inferior part of the foramen opticum. It turns the eye towards the nose.

RECTUS SUPERIOR OCULI, Attollens oculi seu oculum, Levator seu Elevator oculi, Superbus, Oculum movens tertius, (F.) Sus-optico-sphéro-scleroticien, Droit supérieur ou Élévateur de l'œil, Dédaigneux, is attached to the circumference of the foramen opticum.

All the recti muscles are concerned in the voluntary motions of the eye. The oblique muscles have been supposed by Sir C. Bell to execute the involuntary.[?]

RECTUS INFERIOR FEMORIS, Gracilis — r. *Internus major, R. capitis internus major* — r. *Internus minor, R. capitis internus minor* — r. *Lateralis, R. capitis lateralis* — r. *Lateralis Fallopii, R. capitis lateralis* — r. *Major, R. capitis posticus major* — r. *Minor, R. capitis posticus minor*.

RECUBITORIUM, Anacinterium.

RECUPERATION, Restauratio.

RECUPERATIVE, Recup'peratory, from recuperatio, 'recovery.' Restorative. Recovering. Belonging to recovery.

RECURRENT, Recur'rens, from recurrere, (re, and currere), 'to run back.' A name given to several arterial and nervous branches, which seem to re-ascend towards the origin of the trunk whence they emanate.

RECURRENT ARTERIES. This name has been given to several arteries of the forearm, and to one of the leg. In the forearm, there are, 1. The *Radial recurrent, Artère récurrente de l'épicondyle (Ch.)*, which arises from the superior part of the radial, and ascends between the supinator longus, s. brevis, and the brachialis internus. 2. The *anterior cubital recurrent, A. récurrente de l'épitrochlée (Ch.)*, which arises from the upper and inner part of the cubital, and ascends between the pronator teres and brachialis internus. 3. The *posterior cubital recurrent*, which arises from the preceding artery, or from the cubital itself, and ascends between the olecranon and internal tuberosity of the humerus. 4. The *posterior radial recurrent, A. récurrente olécranienne*

(Ch.). It arises from the posterior interosseous, and ascends between the posterior cubital and anconeus muscles. In the *leg*,—the *Arteria recurrens tibialis*, *A. recurrens du genou* (Ch.), is furnished by the anterior tibial, and ascends to the tibialis anticus muscle.

RECURRENT BANDAGES are formed by convolutions of bandages, which make a kind of cap for the part—as the head—to which they are applied.

RECURVATIO, see Hump.

RECUTITUS, Apella.

REDBEAN, *Abrus precatorius*.

REDBERRY, *Arbutus uva ursi*, *Gaultheria*, *Panax quinquefolium*—*r. Bitter*, *Cornus Florida*—*r. Swamp*, *Vaccinium oxycoccos*.

REDGOWN, *Strophulus*.

RED HEAD, *Asclepias Curassavica*.

REDMAILKES, *Papaver rhoeas*.

REDROD, *Cornus sericea*.

RED ROOT, *Celastrus*, *Lachnanthes tinctoria*, *Sanguinaria Canadensis*.

REDSHANKS, *Polygonum persicaria*.

RED WATER TREE, see *Sassy Bark*.

REDIMICULUM, *Diadema*.

REDINGOTES ANGLAISES, see *Condom*.

REDINTEGRATIO, *Regeneration*, *Restauratio*.

REDIVIA, *Paronychia*.

RÉDONDANCE, *Plethora*.

REDOUBLEMENT, *Exacerbation*, *Paroxysm*.

REDRESSEUR UTÉRIN, *Pessary*, *intra-uterine*.

REDS, *Menses*.

REDUCTIO, *Reduction*, *Taxis*—*r. Præputii Impedita*, *Phimosis*.

REDUCTION, *Reductio*, *Repositio*, *Restitutio*, *Diapylasis*, *Diaplasmus*, *Syntaxis*, from *reducere*, *reductum*, (*re*, and *ducere*, *ductum*, 'to lead,') 'to bring back.' A surgical operation, the object of which is to restore displaced parts to their original situation. Thus, we say the reduction of a luxation, when the displaced parts are again brought to their proper relative situation. The reduction of a hernia is the restoration of the protruded parts to the cavity of the abdomen. This is, also, called *Taxis*.

In pharmacy, the process of restoring oxides, chlorides, sulphurets, &c., to the metallic state.

REDUNDANTIA, *Plethora*.

REDUVIA, *Paronychia*.

REED MACE, *Typha latifolia*.

REFECTIO, *Restauratio*.

REFLECTIVA, *Analeptica*.

REFLECTUS, *Restauratio*.

REFICIENTIA, *Analeptica*.

REFLECTION, *Reflexio*, (*re*, and *flecto*, *flectum*, 'to bend,') A bending or turning back.

Duplicature.

REFLECTOR EPIGLOTTIDIS. A muscle, which arises from the arytenoid cartilage and inner part of the thyroid, and is inserted into the lateral edges of the epiglottis.

REFLEX MOVEMENTS, see *Motion*—*r. System of nerves*, see *Excito-motory*, and *Nerves*.

REFLEXIO, *Reflexion*—*r. Palpebrarum*, *Ectropion*.

RÉFORME, *Invaliding*.

REFRACT'ION, *Refractio*, *Photocampsis*, *Dioclasia*, from *re*, and *frangere*, *fractum*, 'to break.' The change of direction which rays of light experience when they pass obliquely from one medium into another of different density. If the light passes from a rarer to a denser medium, it approaches the perpendicular raised at the point of immersion. The contrary occurs, when it goes from a denser to a rarer medium. The theory of lenses, telescopes, and of the eye, rests entirely on this property of light.

RÉFRIGÉRANT, see *Alembic*.

REFRIGÉRANTS, *Refrigerantia*, *Psychica*, *Antitherma*, *Frigeucentia* (F.) *Rafratchiansants*, *Réfrigératifs*, *Desaltérants*, from *re*, and *frigere*, 'to cool.' Medicines which depress the morbid temperature of the body. The chief reputed refrigerants are the *Potasse Nitras* [?], the *Sodæ Boras* [?], but chiefly the abstraction of caloric by ice and ice-drinks, cold water, cold lemonades, soda-water, and effervescing draughts internally; and externally, cool air, cold water, evaporating lotions, and frigorific mixtures.

RÉFRIGÉRATIFS, *Refrigerants*.

REFRIGERATOR, see *Alembic*.

REGENERATIO, *Reproduction*.

REGENERATION, *Regeneratio*, *Palingenesis*, *Redintegratio*, from *re*, and *generare*, *generatum*, 'to beget.' Reproduction of a part of the body, which has been destroyed.

REG'IMEN, (F.) *Régime*, from *regere*, 'to govern.' The rational and methodical use of food, and of everything essential to life; both in a state of health and disease. It is often restricted, in its meaning, to *Diet*. It is sometimes used synonymously with *hygiene*.

RÉGIME SANITAIRE and SYSTÈME SANITAIRE, mean the measures and regulations adopted to prevent the development and propagation of diseases reputed to be pestilential, — especially plague, yellow fever, and cholera.

REGINA PRATI, *Spirea ulmaria*.

REGIO, *Region*.

REGIO AURICULARIS. The region of the ear.

REGIO BUCCALIS. The region of the cheek.

REGIO CARDIACA, *Epigastric region*.

REGIO CERVICALIS. The region of the neck. The cervical region.

REGIO EPIGASTRICA, *Epigastrium*.

REGIO FACIALIS. The facial region.

REGIO FRONTALIS. The frontal region.

REGIO GASTRICA, *Umbilical region*—*r. Hypogastrica*, see *Hypogastrium*—*r. Hypochondriaca*, *Hypochondre*—*r. Iliaca*, *Iliac region*—*r. Inguinalis*, *Inguinal region*—*r. Ischiadica*, *Ischiadic region*—*r. Jugalis*, *Jugal region*—*r. Lumbalis*, *Lumbi*—*r. Mentalis*, *Mental region*—*r. Mesogastrica*, *Umbilical region*—*r. Nasalis*, *Nasal region*—*r. Nuchæ*, *Nuchal region*—*r. Nuchalis*, *Nuchal region*—*r. Occipitalis*, *Occiput*—*r. Occipitalis inferior*, *Nuchal region*—*r. Perinæi*, *Perinæum*—*r. Præcordiaca*, *Præcordial region*—*r. Pubis*, *Pubic region*—*r. Stomachalis*, *Epigastric region*—*r. Stomachica*, *Epigastric region*—*r. Temporalis*, *Temple*—*r. Umbilicalis*, *Umbilical region*.

RE'GION, *Re'gio*. Same etymon. Anatomists have called *regions*, determinate spaces on the surface of the body or of different organs. Thus, the abdomen has been divided into several; to which different names have been given. See *Abdomen*, *Epigastrium*, *Hypochondrium*, *Hypogastrium*.

RÉGION OMBILICALE, *Umbilical region*.

REGIONAL ANATOMY, see *Anatomy*.

RÈGLES, *Menses*—*r. Déviés*, *Menstruation*, *vicarious*—*r. Difficiles*, *Dysmenorrhœa*.

RÉGLISSE, *Glycyrrhiza*.

RÈGNE ANIMAL, see *Animal*.

REG'ULAR, *Regularis*, from *regula*, (*regere*, 'to govern,') 'a rule.' According to rule. Applied to the types of a disease, and to the pulse, when according to rule.

REGULAR PRACTITIONER. One who practises his profession according to the rules established by law or custom.

REGULUS ANTIMONII, *Antimonium*.

REGURGITA'TION, *Regurgitatio*. The act by which a canal or reservoir throws back sub.

stances accumulated in it. *Mitral regurgitation*, for example, means the reflux of blood through the left auriculo-ventricular opening, during the contraction of the left ventricle. Regurgitation is also applied to the *puking* or *possetting* of infants; and to the *rising* of solids or fluids into the mouth in the adult,—from *re*, and *gorge*, (*gurgies*), 'the throat.' (F.) *Rapport*. What is called "vomiting at pleasure" is regurgitation.

The substances brought into the mouth by regurgitation, the French call *Renvois*.

REIF, Psora.

REIN, Kidney.

REINE DES PRES, *Spiræa ulmaria*.

REINERZ, MINERAL WATERS OF. Rein-ort is a small town in Silesia, three German miles from Glatz. The waters are acidulous chalybeates. The principal springs are the tepid drinking spring, the cold spring, and three bathing springs.

REINFORCEMENT, FASCICULUS OF, *F. Cuneatus*. A band of fibres from the *corpus innominatum*, or lateral part of the antero-lateral column of the medulla, which is continued up behind and around the corpus olivare, is lined on its inner or central surface with cineritious substance, enlarges as it ascends, passes over the upper surface of the pons Varolii, and is expanded into the optic thalamus. It thus forms, in its course, the anterior paries of the fourth ventricle, and is brought into view by brushing away the *tuberculum cinereum* or gray matter of the ventricle. The band has received these names owing to its constantly increasing as it ascends, and being mixed up with gray matter on its inner face, from which it seems to derive new fibres.

REINS, Kidneys, Lumbi.

REJECTIO, *Anagôgê*, from *rejicere*, *rejectum*, (*re*, and *jacere*, 'to cast,') 'to cast back.' Every excretion by the mouth,—spitting, expectoration, regurgitation, or vomiting.

REJECTIO SANGUINIS E PULMONIBUS, *Hæmoptysis*.

REJUVENESCENCE, *Rejuvenescentia*, from *re*, 'again,' and *juvenescens*, 'growing young:' (*juvenis*, 'a youth'.) A renewal of youth. The state of being young again. The assumption of the characteristics of youth by the aged.

RELACHANT, Relaxant.

RELACHEMENT, Relaxation.

RELAPSE, *Hypotrophê*, *Epiotrophê*, *Hypotrophê*, *Morbi recidi'va*, *Hypotropias'mus*, from *re*, 'again,' and *labor, lapsus*, 'I fall down.' The return of a disease, during, or shortly after, convalescence. The French use *Rechute* in this sense; and *Récidive* (*Morbus recidi'vus*) for the return of a complaint, at a greater or less period after the complete recovery of health.

The terms *Relapsing fever*, *Short fever*, *Five-day fever*, *Seven-day fever*, *Bilious Remittent fever*, *Remitting Icteric fever*, and *Mild Yellow fever*, (F.) *Fièvre à rechute*, have been given to a form of continued fever, which has occasionally appeared epidemically, as in Edinburgh, in 1843 and 1847, and was generally attended with icteric symptoms. The main phenomena disappeared at about the end of a week; but in many cases a relapse occurred about the end of a fortnight. Few died of the affection.

RELAPSING FEVER, see Relapse.

RELATIO, *Rapport*.

RELATION, FUNCTIONS OF, see Function.

RELAX'ANT, *Laxans*, *Malac'ticum*, *Chalas'ticum*, *Parac'ticum*, (F.) *Relâchant*, from *re*, and *lazare*, 'to loosen.' A medicine which diminishes

the tension of organs. Mucilaginous and fatty substances are relaxants.

RELAXANTIA, Emollients.

RELAXATIO CORDIS ET ARTERIARUM, Diastole.

RELAX'ATION, *Relaxa'tio*. Same etymon. *Chala'sis*, *Chalas'mos*, (F.) *Relâchement*. In *Physiology*, relaxation is opposed to contraction. In *Pathology*, it means great looseness of a part.

RELÈVE-PAUPIÈRE. An instrument for raising the eyelid.

RELEVEUR DE L'ANUS, Levator ani—*r. de la Lèvre inférieure*, Levator labii inferioris—*r. du Menton*, Levator labii inferioris—*r. de l'Omoplate*, Levator scapulæ—*r. de la Prostata*, Levator prostatae—*r. de l'Urèthre*, Levator urethræ.

RELIG'IOUS DISEASES, (F.) *Maladies religieuses*. Diseases affecting the nervous system, and arising from the excitement of religious emotions. See *Convulsionnaire*.

RELIQUAT, (F.) The remnant or remains of a disease.

REMAK FIBRES, see Fibres, Remak.

REMANISIO MENSIIUM, Amenorrhœa.

REMÈDE, Clyster, Medicament—*r. du Capuchin*, Liquor hydrargyri nitrici—*r. de Durande*, Liquor anodynus terebinthinatus—*r. du Duo d'Antin*, Liquor hydrargyri nitrici.

REMEDIAL, Medicinal.

REMEDiate, Medicinal.

REMEDILESS, Incurable.

REMEDIIUM, Juvans, Medicament—*r. Dentifricium*, Dentifrice.

REMEDY, Medicament—*r. English*, Cinchona.

REMISSIO, Decline, Remission—*r. Cordis et Arteriarum*, Diastole—*r. Morbi*, Remission—*r. Trismus*, Atony.

REMIS'SION, *Remis'sio*, *Epan'esis*, *An'esis*, *Submis'sio*, *Aph'esis*, *En'dosis*, *Meio'sis*, from *remittere*, (*re*, and *mittere*, *missum*, 'to send'), 'to relax.' In a more extensive signification, a temporary diminution of the symptoms of a disease, either acute or chronic, *Remis'sio morbi*. Diminution in the febrile symptoms, such as occurs in a remittent fever between the exacerbations.

REMETIC, Medicament.

REMITTENS ICTERODES, Fever, yellow.

REMITTENT, *Remit'tens*. Same etymon. Any disease which presents remissions.

REMITTENT OF CHILDREN, Fever, infantile remittent.

REMITTENT FEVER, *Febris remit'tens*, *Epan'e'tus*, *Erac'erbating*, *paroxysmal*, *sub-continual*, *endem'ic*, *endem'ial* or *endem'ical fever*, (F.) *Fièvre rémittente*, is one which strikingly exacerbates and remits, but without intermission. The ordinary bilious fever of the United States is a simple remittent—*Epanetus mitis*, (F.) *Fièvre rémittente simple*.

REMITTENT FEVER, BILIOUS, Remittent fever, see Relapse.

REMITTING ICTERIC FEVER, see Relapse.

REM'ORA, from *remorari*, (*re*, and *mora*, 'delay,') 'to arrest.' Stoppage, stagnation,—as of blood. Also, the name of two surgical instruments, intended to retain parts *in situ*. The one was used, formerly, in castration, to prevent the intestines from protruding at the inguinal ring; the other, called *Remora Hilda'ni*, (F.) *Arrêt d'Hildan*, *Echene'is*, was employed to maintain fractures and luxations reduced. It is not now used.

REMORA ARATRI, *Ononis spinosa*.

REN, Kidney.

RENAL, *Rena'lis*, from *Ren*, 'the kidney.' That which relates to the kidney.

RENAL ARTERIES, *Emulg'ent Arteries*, are very large and short, and commonly two in number—one to each kidney. They arise from the sides of the abdominal aorta, forming with it a right angle. The left is commonly a little more anterior and elevated than the right. After a short course, they arrive at the fissure of the kidney, where they divide, before entering it, into three or four very considerable branches.

RENAL CAPSULE, Capsule, renal—*r.* Nerve, see Splanchnic Nerves.

RENAL PLEX'USES, *Emulg'ent Plexuses*. These are two:—one on each side. They proceed from fasciuli of the solar and coeliac plexuses; from the outer side of the semilunar ganglion and the final expansion of the small splanchnic nerves. These plexuses begin by three or four ganglions, situate at the commencement of the renal artery; and they give off from their circumference many fine, straight filaments, which do not anastomose, and which enter the kidney with the divisions of the renal artery. These divisions they accompany.

RENAL VEINS, *Emulgent Veins*. These are very large. Their roots follow the same course as the arteries in the substance of the kidney. In the fissure or pelvic portion of the kidney, they unite into a single trunk, which passes transversely inwards, and opens at the side of the vena cava descendens.

RENCHUS, Snoring, Stertor.

REN'ULUS, *Renid'ulus*, *Ren'ulus*, diminutive of *Ren*, 'a kidney.' A small kidney. A lobe of the kidney.

RENDLES, Rennet.

RENEAL'MIA EXALTA'TA, *Alpin'ia exalta'ta* seu *tubula'ta*. A plant of British Guiana, which is possessed of diaphoretic and diuretic properties: and, in large doses, is emetic. It is much used there in dropsy, rheumatism, dysentery, &c. The bruised rhizoma is the part employed.

RENES SUCCENTURIATI, Capsules, renal.

RENICULUS, Renculus.

RENIFORM. Nephroid.

RENNES. MINERAL WATERS OF. Thermal salines in the department of Aube, France, containing carbonic acid and chloride of magnesium. Temperature 103 to 121°, of Fahr.

RENNET, *Rennet*, *Coag'ulum*, *pytia*, *pytia*, *Pytia*, *Pyet'ia*, from (G.) *gerinnen*, 'to coagulate' [?], (F.) *Caille-lait*. When the fourth stomach or *Abomasus* of the calf is salted and dried, it possesses the property of coagulating milk, when a portion of it is soaked in water, and the infusion—*Rennet*, *Rendles*, *Rouning*, (Prov.) *Earning*, *Preaser*, *Read*, *Running*, *Steep*—is added to milk.

RENNET WHEY, see Serum lactis.

RENONCULE, *Ranunculus*—*r.* *Bulbeuse*, *Ranunculus bulbosus*.

RENOUÉE, *Polygonum aviculare*—*r.* *Acre*, *Polygonum hydropiper*.

RENOUEUR, (F.) *Rebouteur*, *Rhabilleur*, from *re*, and *nouer*, (*nodus*, 'a knot,') 'to tie or knit again.' A bone-doctor:—a bone-setter. One who makes a trade of reducing fractures and dislocations.

RENOVATIO, Ananæosis.

RENSSELAERIA VIRGINICA, *Peltandra Virginica*.

RENUENS, Rectus capitis internus minor.

RENULUS, Renculus.

RENUM ATONIA, Nephrotonia—*r.* Paralysis, Nephrotonia.

RENUNCIATIO, Rapport.

RENVERSEMENT DE LA LANGUE, *Paraglossa*—*r.* de la *Matrice*, Uterus, inversion of the—*r.* des *Paupières*, Ectropion—*r.* du *Rectum*, Proctocèle—*r.* de la *Vessie*, Exocyste, Exstrophia of the bladder.

RENVOIS, see Regurgitation,

REPANDATIO, Lordosis.

REPARATION, Restauratio.

REPARATIVE, *Repar'atory*. Having the power of restoring or repairing.

REPELLENTIA, Repellents.

REPEL'LENTS, *Repellent'ia*, *Repercution'ia*, *Apocrus'tica*, *Repercus'sives*, (F.) *Répercu'sifs*, from *re*, and *pellere*, 'to drive.' Medicines which, when applied to a tumefied part, cause the fluids, that render it tumid, to recede, as it were, from it. Astringents, cold-water, ice, &c., are repellents.

RÉPERCUSSIONS, Repellents.

REPERCUSSION, *Contra-fissura*, *Repercussion*.

REPERCUS'SION, *Repercus'sio*, *Anac'lasis*, from *re*, and *percutio*, *percussum*, (*per*, and *quater*, 'to shake or batter,') 'I strike back.' Disappearance of a tumour or cutaneous eruption, in consequence of the application of a repellent. The action of a repellent remedy. The French writers use the term *Delitescence*, *Delitescencia*, for the sudden disappearance of a tumour, inflammation, eruptive disease, or purulent collection, without the disease occurring in any other part of the body: in this respect delitescence differs from metastasis: *Delitescence* of an inflammation or 'sudden disappearance' is distinguished by certain French pathologists from *Resolution*. *Repercussion* also means *Ballotement*.

REPERCUSSIONS, Repellents.

REPERCUTIENTIA, Repellents.

REPLETIO, Fulness, Plethora.

REPLI LONGITUDINAL DE LA MÉNINGE, Falx cerebri.

REPOSITIO, Anaplasia, Reduction, Taxis.

REPOUSSOIR, Punch—*r.* d'Arêtes, Probang.

REPRIMENS, Epischetius.

REPRODUCTION, *Reproduc'tio*, *Regenera'tio*, from *re*, and *producere*, *productum*, (*pro*, and *ducere*.) The function by which living bodies produce bodies similar to themselves. See Generation. As a general remark, it will be found true, that the larger animals are uniparous, and the smaller, which are more exposed to destruction, multiparous. The mammalia being of the same natural class as man, it may be useful to compare them in this respect.

COMPARATIVE TABLE OF REPRODUCTION IN THE CLASS MAMMALIA.

Species.	Age capable of en- gendering.	Duration of gesta- tion.	Number of young at each birth.	Period of cessation of the function.
<i>Primates.</i>				
Man	14 years	9 months	1 to 3	At a somewhat ad- vanced age.
Monkeys (large).....	3 years	7 months	Do.	
Do. (long tailed)	2 years	6 months	Do.	
<i>Carnivora.</i>				
Bear	2 years	112 days	1 to 4	25 to 30 years.
Badger	3 to 4	20 to 25 years.
Hedgehog	1 year	1 mo. and 10 days	3 to 5	Produces during life
Ferret.....	11 months	Do.	5 to 9 twice a year	
Otter.....	3 months	3 to 4	
Pole-cat, } Marten, } Weasel, and } Ermine, } Didelphus, and } Opossum, and } Phalander, }	before 1 year	56 days	3 to 6	8 to 10 years.
Lion	2 years	3 to 4 months	3 to 4	20 to 25 years.
Tiger, } Leopard }	Do.	3 months	4 to 5	Do.
Lynx	9 weeks	3 to 4	9 years.
Wild Cat	before 1 year	56 days	4 to 6	
Wolf	2 years	73 days	5 to 9	15 to 20 years.
Dog, in the most } natural state, }	10 months	63 days	3 to 6	15 years.
Fox	1 year	2 months	Do.	10 or 12 years.
Jackall	Do.	5 to 8	
Isatis	63 days	6 to 7	
Phoca	3 months (?)	2 to 3	
<i>Rodentia.</i>				
Squirrel.....	1 year	45 days	3 to 4 twice a year	Produces during life
Flying Squirrel.....	3 to 4	Do.
Ondatra	3 to 5	Lives 6 years.
Beaver	4 months	2 to 3	Lives 8 years.
Hare	1 year	30 days	{ 2 to 4 several } times a year }	
Rabbit	6 months	Do.	4 to 8 do.	
Rat	Do.	5 or 6 weeks	5 to 6 do.	Lives 8 or 9 years.
Mouse	Do.	1 month	Do.	Produces during life.
Marmot	6 weeks	2 to 4	Do.
Guinea Pig.....	5 to 6 weeks	3 weeks	5 to 8 times a year	Do.
Dormouse	1 year	3 to 5	
<i>Edentata.</i>				
Armadilla	4 times a year	7 to 8 years. (?)
<i>Ruminantia.</i>				
Camel	4 years	11 months or 1 y'r	1	40 to 50 years.
Dromedary	Do.	Do.	1	Do.
Buffalo	3 years	9 months	1	15 to 18 years.
Beef	2 years	Do.	1 to 2	9 years.
Lama	3 years	Do.	12 years.
Reindeer	2 years	8 months	1	Lives 16 years.
Buck and Doe	1½ year	Do.	1 to 2	Lives 25 or 30 years.
Roebuck	Do.	5 months	Do.	Lives 12 or 15 years.
Saiga (Antelope).....	1 year	Do.	Do.	Lives 15 or 20 years.
Chamois and Wild Goat	Do.	Do.	1, 2, and 3	Lives 18 or 20 years.
Goat	Do.	Do.	Do.	7 years.
Musimon	1½ year	Do.	1 to 2	8 or 10 years.
Sheep	1 year	Do.	Do.	8 years.
<i>Solipedes.</i>				
Horse	2½ years	290 days	1 to 2	25 or 30 years.
Ass	Do.	Do.	Do.	Do.
Zebra	Do.	Do.	Do.	Do.
<i>Pachydermata.</i>				
Hog	9 months or 1 year	4 months	10 to 20	15 years.
Elephant.....	16 years (?)	9 to 11 months	1 or 2	Lives 70 years. (?)
Rhinoceros	5 years (?)	1 or 2 (?)	Lives 60 years. (?)
Hippopotamus	1	
Morse	9 months	1	

REPRODUCTION MÉRISMATIQUE, see Generation.

REPULSORIUM, Punch,
REPURGATIO, *Anacatharsis*, from *re*, and
purgare, 'to cleanse.' According to Castelli—
purgation by transpiration or by expectoration.
Medicines producing this effect are called *Repur-*
gantia.

REQUIES, *Acinesia*.
REQUIETO, *Acinesia*

RES CONTRA-NATURAM. 'Things against
nature,' (F.) *Choses contre-nature*, are those things
which tend to destroy man;—as disease, and
every thing connected with it.

RES CULINARIA, Culinary art—*r. Medica*, Me-
dicina.

RES NATURALES, (F.) *Choses naturelles*. Those
things, which, by their union, were conceived to
constitute the nature of man: such as the ele-
ments, temperaments, humours, spirits, &c.

RES NON NATURALES, (F.) *Choses non-naturelles*. Those things which, anciently, formed the matter of hygiene. See Non-naturals.

RES VENEREA, Coition—r. Vestiaria, Vestitus.

RÉSEAU, Rete, Reticulum—r. *Admirable*, Rete mirabile—r. *Intermédiaire*, see Corpus spongiosum urethræ—r. *Merveilleux*, Rete mirabile.

RESECTIO, Resection—r. Articulorum, see Resection.

RESECTION, *Resec'tio*, from *resecare*, *resectum*, (re, and *secare*), 'to cut off.' A name, given, especially by the French surgeons, to operations in which the carious extremities of long bones, or the unconsolidated extremities of fractured bones forming irregular joints, are removed with the saw.—*Resectio articulo'rum*, *Decapita'tio articulo'rum*.

RESECTION, see Amputation.

RESEDA CRISPATA—r. Luteola—r. Lanceolata—r. Luteola.

RESÉDA LUTEOLA, *R. lanceolata* see *cripata*, *Lute'ola*, *Weld*, *Dyer's Weed*. A European plant naturalized in the United States, which was at one time used as a diaphoretic and diuretic.

RESERANS, Aperient.

RESERVE AIR, see Respiration.

RÉSERVOIR DE LA BILE, Gall-bladder—r. *du Chyle*, *Receptaculum chyli*—r. *des Larmes*, *Lachrymal sac*—r. of *Pecquet*, *Receptaculum chyli*—r. *de Pecquet*, *Receptaculum chyli*—r. of the Thymus, see Thymus.

RESIDENTIA, Sediment.

RESIDUAL AIR, see Respiration.

RESIMUS, Camus.

RESIN, *Resi'na*, *Rheti'nē*, (F.) *Résine*. A vegetable product, commonly dry and concrete, more or less brittle, inodorous or slightly odorous, insipid, or of an acrid warm taste; of a smooth, glassy fracture, heavier than water, inflammable, insoluble in water, soluble in alcohol, ether, and yolk of egg, and negatively electrifiable by friction. Many resins are used in medicine; the greater part are purgative and irritating. Some act like acrid poisons.

RESIN, ACAROID, see Xanthorrhœa—r. Botany Bay, see Xanthorrhœa—r. Cayenne, Caoutchouc—r. of New Holland, see Xanthorrhœa—r. of the Spruce fir, see *Pinus abies*—r. of Xanthorrhœa, see Xanthorrhœa.

RESINA, Resin—r. *Abietis*, see *Pinus abies*, *Pinus picea*—r. *Abietis humida*, see *Pinus abies*—r. *Abietis sicca*, see *Pinus abies*—r. *Acaroides*, see Xanthorrhœa—r. *Alba*, see *Pinus sylvestris*—r. *Alba humida*, see *Pinus Abies*—r. *Cayennensis*, *Caoutchouc*—r. *Copal*, *Copal*—r. *Elastica*, *Caoutchouc*—r. *Flava*, see *Pinus sylvestris*—r. *Fluida Canadensis*, see *Pinus balsamea*—r. *Jalapæ*, *Extractum jalapæ resinosum*—r. *Juniperi*, *Sandarac*—r. *Laricis*, see *Pinus larix*—r. *Lentiscina*, see *Pistachia lentiscus*—r. *Liquida pini balsameæ*, see *Pinus balsamea*—r. *Liquida pini laricis*, see *Pinus larix*—r. *Lutea Novi Belgii*, see Xanthorrhœa—r. *Nigra*, *Colophonia*—r. *Pini*, see *Pinus sylvestris*—r. *Pini empyreumatica liquida*, see *Pinus sylvestris*—r. *Pini abietis sponte concreta*, see *Pinus abies*—r. *Strobilina*, see *Pinus balsamea*, and *Pinus mughos*—r. *Terebinthi*, see *Pistacia terebinthus*.

RESINATUM, *Rhetino'ton*. An epithet for wine impregnated with resin, and used by the ancients as a stomachic.

RÉSINE, Resin—r. *de Cayenne*, *Caoutchouc*—r. *Élastique*, *Caoutchouc*—r. *de Guaiac*, see *Guaiacum*—r. *Gomari*, see *Bursera gummifera*—r. *de Lierre*, see *Hedera helix*.

RESINOCE'NUM, *Rhetinoce'ron*, from *ῥητιν*, 'resin,' and *κενος*, 'wax.' A mixture of resin and wax.—r. *Galen*.

RESINOSA, in Pharmacy, resinous substances.

RÉSOLUTIFS, Discutients.

RESOLUTIO, Resolution—r. *Membrorum*, *Paralysis*.

RESOLUTIO MORBI. The resolution or disappearance of a disease; especially when without any critical evacuation.

RESOLUTIO ET DIABROSIS VENTRICULI, *Gastromalacia*—r. *Nervorum*, *Paralysis*—r. *Ventriculi autopeptica*, *Gastromalacia*—r. *Virium*, *Dialysis*.

RESOLUTION, *Resolu'tio*, *Discus'sio*, from *resolvere*, (re and *solvere*, *solutum*, 'to loosen.') Removal or disappearance, as of a disease. Resolution is one of the terminations of inflammation, in which the inflamed part returns gradually, and without suppuration, to the healthy condition.

RÉSOLUTION DES MEMBRES, Paralysis.

RESOLVENTIA, Discutients.

RES'ONANCE, (F.) *Retentissement*, *Résonance*, *Résonnement*, from *re*, 'again,' and *sono*, 'I sound,' 'A return of sound.' A resounding; a reverberation of sounds. A thrilling of the voice more loud than natural; or its existence in a part where it is not heard in health,—as detected by auscultation. A thickened and hardened state of the lungs, caused either by a mass of crude tubercles, or by inflammation, is generally considered to produce this phenomenon, by rendering the lung a better conductor of the murmur of the voice in the bronchi. Hence the origin of the term *Bronchoph'ony*, from *βρογχος*, 'bronchus,' and *φωνη*, 'the voice;' (F.) *Voix bronchique*, *Voix tubaire*, *V. bourdonnante*.

Resonance is also used in the sense of resounding; as when the chest is said to be resonant on percussion.

RESONANCE, VESIC'ULO-TYMPANIT'IC, see Vesiculo-tympanitic—r. Vibrations of, see Sound.

RESONITUS, Contra-fissura.

RES'ONANT, *Res'onans*, *Res'onus*, (F.) *Retentissant*, *Résonnant*. Resounding. See Resonance.

RÉSONNANCE, Resonance.

RESONNEMENT, Resonance.

RESORBENS, Absorbent.

RESORPTIO, Absorption, Resorption—r. *Cutanea*, see Absorption—r. *Cutis*, see Absorption.

RESORPTION, *Resorp'tio*, from *resorbere*, *resorptum*, (re, 'again,' and *sorbere*), 'to absorb again.' Absorption of a fluid which has been previously deposited in a part by the exhalants, as of pus, blood, serum, &c.

RES'PIRABLE, *Respirab'ilis*. Same etymology as Respiration. Capable of being breathed. Adapted for respiration.

RESPIRAMEN, Respiration.

RESPIRAMENTUM, Respiration.

RESPIRATEUR, Respirator, Respiratory.

RESPIRATIO, Respiration—r. *Abolita*, *Apnoea*—r. *Brevis*, *Dyspnoea*—r. *Difficilis*, *Dyspnoea*—r. *Stertens*, *Stertor*—r. *Stertuosa*, *Stertor*.

RESPIRA'TION, *Respira'tio*, *Anap'nōē*, from *respiro*, (re, and *spiro*), 'I take breath.' *Anapneu'sis*, *Pneu'sis spiratio*, *Respira'men*, *Respiramentum*, *Respira'tus*, *Respir'itus*, *Spir'itus*. A function proper to animals, the object of which is, to place the materials of the blood—the mixture of the venous blood with lymph and chyle—in contact with atmospheric air, in order that it may acquire the vivifying qualities which belong to arterial blood. The organs for executing this function are, in the mammalia, birds, and reptiles, the lungs. In man, the respiration consists of mechanical and chymical phenomena. The mechanical are *Inspiration* and *Expiration*. The evident chymical phenomena consist in the formation of a certain quantity of carbonic acid, the absorption of a part of the oxygen of the air, and

the disengagement of a quantity of water in the state of vapour. In the healthy condition the respiration is easy, gentle, regular, and without noise. In man, the respirations are generally about 35 per minute in the first year of life; 25 during the second; 20 at puberty; and 18 in the adult age.

The air of respiration has been divided into *first*, the *residual air*, or that which cannot be expelled from the lungs, but remains after a full and forcible expiration, estimated at 120 cubic inches: *secondly*, the *supplementary or reserve air* or that which can be expelled by a forcible expiration, after an ordinary outbreathing, valued at 130 cubic inches: *thirdly*, the *breath, tidal or breathing air*, valued at 26 cubic inches: and *fourthly*, the *complementary or complementary air*, or that which can be inhaled after an ordinary inspiration, which amounts to 100 cubic inches. This estimate gives 250 cubic inches as the average volume of air which the chest contains after an ordinary expiration.

RESPIRATION BRONCHIQUE, see *Murmur, respiratory*.

RESPIRATION, BRONCHO-VESICULAR. A morbid modification of the respiratory murmur, having more or less of the characters of the bronchial and the vesicular respiration combined. It denotes incomplete solidification of the lung, as from the presence of tubercle. — Flint.

RESPIRATION OF THE CELLS, *Murmur, respiratory* — *r. Cogged-wheel*, *Respiration, jerking* — *r. Entrecoupée*, *Respiration, jerking* — *r. Exagérée*, see *Puerile* — *r. Hypervésiculaire*, see *Puerile*.

RESPIRATION, JERKING, Cogged-wheel, or Wavy respiration, (F.) *R. Saccadée ou Entrecoupée*. When the murmur of inspiration, in place of being continuous, is interrupted as by starts, it is called 'jerking.' It is a concomitant of incipient pleurisy, pleurodynia, spasmodic asthma, and tuberculosis of the lung with corresponding pleuritic adhesion. — Walshe.

RESPIRATION, LUCTUEUSE, see *Luctueux* — *r. Metallic*, *Tintement métallique* — *r. Nulle*, see *Murmur, respiratory*.

RESPIRATION PROOF, *Docimasia pulmonum*.

RESPIRATION, PUERILE, see *Puerile* — *r. Répense*, see *Murmur, respiratory* — *r. Rude*, see *Murmur, respiratory* — *r. Silencieuse*, see *Murmur, respiratory* — *r. Saccadée*, *R. jerking* — *r. Soufflante*, *Murmur, respiratory* — *r. Supplémentaire*, see *Puerile* — *r. Tubal*, see *Murmur, respiratory* — *r. Wavy*, *Respiration, jerking*.

RESPIRATOIRE, *Respiratory*.

RESPIRATOR, (F.) *Respirateur*. Same etymon. Various instruments, under this name, have been devised for the purpose of tempering the air before it reaches the lungs in winter. One of the best is that of Jeffrey, which is made of several layers of fine wire tissue.

RESPIRATORIUM VENTRIS, *Diaphragm*.

RESPIRATORIUS, *Respiratory*.

RESPIRATORY, *Respiratorius*, (F.) *Respirateur*, *Respiratoire*. Appertaining to respiration. An epithet applied to the murmur heard by auscultation in a healthy adult, during inspiration and expiration. It marks the entrance of the air into the cells, and its passage out of them. See *Murmur*.

RESPIRATORY, EXTERNAL, NERVE, see *Thoracic nerves*.

RESPIRATORY GANGLIA, see *Encephalon*.

RESPIRATORY, INTERNAL, NERVES, *Diaphragmatic Nerves* — *r. Nerve of the face*, *Facial Nerve* — *r. Superior, nerve*, *Spinal Nerve*.

RESPIRATORY TRACT, *Tractus respiratorius*. The middle column of the spinal marrow, described by Sir Charles Bell as that whence the respiratory nerves originate. See *Nerves*.

RESPIRATUS, *Respiration*.

RESPIRATUS, *Respiration*.

RESSERRÉ, (*VENTRE*), *Constipation*.

RESTA BOVIS, *Ononis spinosa*.

RESTAGNATIO, *Erection*.

RESTAURANS, *Analeptic, Corroborant*.

RESTAURATIO, *Réfectio, Refectus, Analépsis, Anatrepsis, Anacon'idé, Analépsis, Anacatépsis, Apocatas'tasis*. Restoration of strength, after disease or great fatigue; *Epanortho'sis, Recreatio, Redintegratio, Restauratio virium*. Return of the organs to their normal condition. *Recovery, Recuperation, Reparation*.

RESTAURATIO VIRIUM, see *Restauratio*.

REST HARROW, *Ononis spinosa*.

RESTIFORMIS, from *restis*, 'a rope,' and *forma*, 'shape.' Shaped like a rope, as the 'Corpora restiformia.'

RESTIFORMIS PROCESSUS, *Coracoid*.

RESTITUTIO, *Reduction* — *r. Organica*, *Morioplastice* — *r. Partium deperditarum*, *Morioplastice*.

RESTLESSNESS, *Agitation*.

RESTORATION, *Restauratio*.

RESTORATIVE, *Analeptic, Corroborant*.

RESTRICITIO INTESTINI RECTI seu *ANI*, *Stricture of the rectum*.

RESUMPTIVUS, *Resumptorius*; from *re*, 'again,' and *umere, sumptum*, 'to take.' Causing to resume. *Resumptivum*. Any agent calculated to restore flesh and strength to one who has been reduced.

RESUMPTORIUS, *Resumptivus*.

RESURRECTION-MAN, *Resurrectionist*.

RESURRECTIONIST, *Resurrection-man*, *Body snatcher*: from *re*, and *urgere, surrectum*, 'to arise.' An individual who violates the sanctuary of the grave to obtain subjects for dissection.

RESUSCITATION, *Resuscitatio*, from *resuscitare*, (*re*, and *suscitare, sursum citare seu eiere*), 'to move up,' 'to stir up anew.' *Revivification*. The restoration of persons, apparently dead, to life. See *Asphyxia* and *Suffocation*.

RETCHING, *Vomitition*.

RETE, *Plexus* — *r. Malpighii*, *Corpus mucosum*.

RETÉ, 'a net;' (*Reticulum*, 'a small net,') *Dictyon*, (F.) *Réseau*. A name given to the interlacing and decussation of blood-vessels, lymphatics, fibres, and nerves when they form a kind of network.

RETÉ has also been applied to the *epiploon*. — *Castelli*.

RETÉ MAJUS, *Epiploon, gastro-colic*.

RETÉ MIRAB'LE, (F.) *Réseau admirable, R. merveilleux*. A name given to the ramifications and anastomoses of the internal carotid and vertebral arteries at the base of the brain.

RETÉ MIRABILE seu **RETÉ VASCULOSUM MALP'GHII** is the plexus formed on the cells of the lungs by the minute ramifications of the pulmonary artery.

RETÉ MUCCOSUM, *Corpus mucosum*.

RETÉ TESTIS, *Reté Vasculosum Testis* seu *Halleri*. An irregular plexus or network, formed by the communication of the *Vasa recta*, *Tubuli* seu *Ductuli recti*, at the upper and back part of the testicle. These *Vasa recta* are sent out at the back part of the testicle, from the convoluted seminal tubes. The *Rete Vasculosum testis* sends out from twelve to eighteen straight tubes, called *Vasa efferentia*, which convey the semen from the testes to the epididymus.

RETENTIO, *Retention* — *r. Aerea*, *Dyspnœa* — *r. Alvina*, *Constipation* — *r. Bilis*, *Ischocolia*.

RETENTIO FLUXUS HÆMORRHOÏDALIS. Suppression of the hæmorrhoidal flux.

RETENTIO FŒTUS, *Dystocia* — *r. Lactis*, *Ischogalactia* — *r. Lochiorum*, *Ischiolochia* — *r. Mensium*,

see Amenorrhœa — *r.* Spermatidis, IschospERMIA — *r.* SUDORIS, Hydroschesis.

RETENTION, *Retentio*, *Apoplexia*, *Apoplexis*, from *retinere*, *retentum*, (*re*, and *tenece*), 'to hold back.' Accumulation of a solid or liquid substance, in canals intended for its excretion, or in a reservoir, intended to contain it for a short period.

RETENTION OF URINE: *Paruria Retentio*, *nis vesicalis*, *Ischuria vesicalis*, *Retentio Uri-næ*, (*F.*, *Rétention d'urine*). A disease in which the urine accumulated in the bladder cannot be evacuated; or, at least, cannot be passed without extreme difficulty. In the former case, the retention is said to be *complete*; in the latter, *incomplete*. Many writers have distinguished three degrees of this affection, to which they have given the names *Dysury*, *Strangury*, and *Ischury*. Retention of urine may depend upon loss of contractility, paralysis of the bladder, &c., or on some obstacle to the passing of the urine, as in cases of pressure of the womb on the bladder, of tumours in its vicinity, foreign bodies in its cavity, inflammation of the urethra, swelling of the prostate, stricture, &c. The pain is extremely violent, and the bladder may be found distended above the pubis. The treatment consists in introducing the catheter, or puncturing the bladder, and in combating the cause, which has produced the retention, by particular means adapted to each case.

RENTISSANT, Resonant.

RENTISSEMENT, Resonance — *r.* *Autophonique*, see *Autophonia*.

RETIA UTERI, see *Uterus*.

RETICULAR, *Retiform*, *Reticularis*, *Retinus*, *Amphiblastroïdes*, *Ductoïdes*, *Retiformis*, from *rete*, (diminutive *reticulum*), 'a net.' That which resembles a net. An epithet applied to many structures in the body.

RETICULAR BODY, *Corpus mucosum* — *r.* Substance, Cellular tissue.

RETICULARIA OFFICINALIS, *Lichen pulmonarius*.

RETICULUM: diminutive of *rete*, 'a net.' The *Bonnet* or *Honeycomb bag* or *stomach*, (*Sc.*) *Kingshood*, (*F.*) *Bonnet*, *Réseau* — the second stomach of the ruminant animal. See also, *Epiploon*, and *Rete*.

RETICULUM CUTANEUM, *Rete mucosum* — *r.* *Mucosum*, *Rete mucosum*.

RETIFORM, *Reticular*.

RETINA, from *rete*, 'a net'; *Tunica retinalis* seu *reticularis* seu *retiformis* seu *intima* seu *nervea oculi*, *Involucrum reti comparatum*, *Vericularis* membra'na, *Amphiblastroïdes membrana*, (*F.*) *Rétine*. A soft, pulpy, grayish, semi-transparent, very thin membrane; extending from the optic nerve to the crystalline, embracing the vitreous humour, and lining the choroid; without, however, adhering to either of those parts. It terminates by a defined edge — *margo dentatus* — at the posterior extremities of the ciliary processes. It is constituted, according to most anatomists, by the expansion of the optic nerve. The retina appears to be formed of several laminae; so joined together, that it is difficult to discriminate them. The one — the innermost — is medullary, and pulpy; the other — the outer — is stronger, and fibro-vascular. The retina is the essential organ of vision; on it the images of objects are impressed. Both it and the optic nerve are devoid of general sensibility. They may be punctured or lacerated without pain being experienced. The nerve of general sensibility distributed to the eye is the fifth pair.

RETINACULA, see *Retinaculum* — *r.* *Morgagnii*, see *Fræna* of the valve of *Bauhin*.

RETINACULUM, from *retinere*, (*re*, and *tenece*), 'to hold back.' A surgical instrument,

formerly used in the operation of castration; and in that of hernia, to prevent the intestines from falling into the scrotum, — *Scultet*. *Retinacula* are also band-like extensions of a cellulose-membranous structure, which seem to suspend the ovum in the Follicle of *De Graaf*. They appear to correspond to the chalazæ of the egg of the bird.

RETINACULUM, *Bride*.

RETINACULUM TEN'DINUM PERONEORUM, *Vagina malleolaris externa*. The ligamentous sheaths that retain in situ at the outer ankle the tendons of the peroneal muscles.

RETINAL, *Retinalis*. Relating or appertaining to the retina.

RÉTINE, *Retina*.

RETINITIS, *Inflammatio Retinalis*, *Amphiblastroïdis*, *Dictyitis*, (*F.*) *Rétinite*, from *retina* and *itis*, a suffix denoting inflammation, — a term of hybrid formation. Inflammation of the retina; an uncommon affection, except as an accompaniment of other forms of ophthalmia.

RÉTINOÏDÊ, see *Rhétinolê*.

RÉTINOLÊ, see *Rhétinolê*.

RETINOSCOPY, *Retinoscopiæ*, from *retina*, and *σκοπεω*, 'I view.' Examination of the retina.

RETINUS, *Reticular*.

RETORT, *Retorta*, *Battia*, *Cornu*, *Cornuta*, *Cornes*, *ta*, *Cornamusa*, (*F.*) *Corne*, from *retorqueo*, (*re*, and *torqueo*, *tortum*), 'I bend back.' A vessel, commonly of glass or earthenware, employed in distillation. It may be *plain*, or have a *tubulure* for admitting substances whilst the distillation is going on. In the latter case, it is said to be *tubulated*, (*F.*) *Cornue tubulée*.

RETRAC'TION, *Retractio*, from *retrahere*, (*re*, and *trahere*, *tractum*), 'to draw back.' State of a part, when drawn towards the centre of the body or backwards; — as retraction of the testicles. Sometimes it is used synonymously with *shortening* or *drawing up*; as in retraction of the thigh.

RÉTRACTION DE LA MATRICE, *Retroversio uteri*.

RETRACTOR ANGULI ORIS, *Buccinator*.

RETRAHENS AURIS, *Posterior* seu *Triceps Auris*, *Tricaudalis*, *Retrahens Auriculæ* seu *Auriculam*, *Secundus proprius Auriculæ*, *Bicaudalis*, *Deprimens Auriculæ*, *Proprius Auris externa*, (*F.*) *Mastoido-conchiformis*, *Mastoido-oribulair* (*Ch.*), *Auriculaire postérieur*. It consists of one or more small fasciculi or fleshy, thin, somewhat irregular fibres, situated behind the ear. These are attached to the mastoid process, and pass horizontally forwards, to terminate at the convexity formed by the concha of the ear. If this muscle had any use, it would carry the ear backwards.

RETRAIT VENTRICULAIRE, *Heart*, hypertrophy of the.

RÉTRÉCISSEMENT, *Arctatio*, *Stricture* — *r.* *de l'Urèthre*, *Urethrostenosis*.

RETRIMENTA VESICÆ, *Urine*.

RETRIMENTUM, *Excrement*.

RETROCESSION, *Retrocesio*, *Retrocesus*, *Retrogressus*, from *retrocedere*, *retrocessum*, (*re*, *tro*, 'backwards,' and *cedere*, 'to give place.') The act of going back. A disappearance or metastasis of a tumour, eruption, &c., from the outer part of the body to the inner.

RETROCESSUS, *Retrocession*.

RETROFLEXIO UTERI, see *Retroversio uteri*.

RETROGRESSUS, *Retrocession*.

RETROPHARYNGEAL, *Retropharyngeus*, from *retro*, 'backwards,' and *pharynx*, 'the pharynx.' Relating to parts behind the pharynx, —

as 'retropharyngeal abscess,'—an abscess behind the pharynx.

RÉTROPULSION DE LA MATRICE, Retroversio uteri

RETRO-UTERINE, *Retro-uteri'nus*, (F.) *Rétro-utérin*, from *retro*, and *uterus*. Behind the uterus, as *retro-uterine abscess*, (F.) *Abscès rétro-utérin*:—an abscess seated behind the uterus.

RETROVACCINATION, from *retro*, 'backwards,' and *vaccination*. Vaccination with matter obtained from the cow, after inoculating the animal with vaccine matter from the human subject.

RETROVERSIO UTERI, (*retro*, 'backwards,' and *vertere*, *versum*, 'to turn.') *Retroversion of the Uterus*, *Edopto'sis Uteri retrover'sa*, *Anapod'sis* seu *Anapodis'mus Uteri*, (F.) *Rétroversion*, *Rétraction*, *Obliquité* ou *Rétropulsion de la Matrice*. A change in the position of the uterus, so that the fundus of the organ is turned towards the concavity of the sacrum, whilst the neck is directed towards the symphysis pubis. This displacement is commonly attended with constipation, tenesmus, and retention of urine. Its usual period of occurrence is between the third and fourth months of pregnancy; before the uterus has escaped above the superior aperture of the pelvis. In the treatment, the catheter must be regularly used twice a day, until the uterus, by its growth, rises above the pelvis. The catheter should be small, flat, and curved considerably more than common, in consequence of the distorted course of the urethra. The bowels must be kept open; and absolute rest, in a recumbent posture, be enjoined. Should it be impracticable to draw off the urine, attempts must be made to replace the uterus. The woman being on her hands and knees, the fore and middle fingers of the accoucheur's left hand are to be passed up the rectum to the fundus uteri; which they must elevate, whilst the cervix uteri is carefully depressed by two fingers of the right hand in the vagina. Should the fingers employed to elevate the fundus not be long enough to effect this object, a piece of whalebone may be substituted, having a small piece of sponge attached to one extremity as a pad.

A simple inclination, or bending of the uterus, backwards, is termed *Retroflex'ion of the Uterus*, *Retroflex'io Uteri*: not unfrequently, however, this is used synonymously with Retroversion of the Uterus.

RÉTROVERSION DE LA MATRICE, Retroversio uteri.

REUNIO, Synthesis.—*r. Partium separatarum*, see Synthesis.—*r. per Primam intentionem*, see Intention.—*r. per Secundam intentionem*, see Intention.—*r. Vasorum*, Anastomosis.

REUNION PRIMITIVE, see Intention.—*r. Secondaire*, see Intention.

REVACCINATION, see Vaccina.

REVALENTA ARABICA, see Evrum.

RÉVASSERIE, Incoherence; see Somnium.

RÊVE, Somnium.

RÊVEIL, see Sleep.—*r. en Sursaut*, see Sleep.

REVELLENT, Derivative.

RÉVERIE, Aphelxia.

REVERY, Aphelxia.

REVIVIFICATION, Resuscitation.

RÉVULSIF, Derivative.

REVULSION, *Revul'sio*, from *re*, and *vellere*, *enlum*, 'to pluck'; *Antis'pasis*. The act of turning the principle of a disease from the organ in which it seems to have taken its seat. Rubefacients, vesicatories, bleeding in the foot, &c., are often used for this purpose. See Derivation.

REVULSIVE, Derivative.—*r. Bleeding*, see Bloodletting.

REVULSORIUS, Derivative.

REX AMARORUM, Soulaamen amara.—*r. Metallorum*, Aurum.

REYNOLDS'S SPECIFIC, see Vinum colchici.

REZZLE, Wheeze.

RHA, Rheum.

RHABBARUM, Rheum.—*r. Album*, Convolvulus mechoacan.—*r. Antiquorum*, Rheum Rhaponticum.—*r. Dioscoridis*, Rheum Rhaponticum.—*r. Monachorum*, Rumex patientia.—*r. Nigrum*, Convolvulus jalapa.—*r. Pauperum*, Thalictum.—*r. Sibericum*, Rheum.—*r. Tartaricum*, Rheum.—*r. Turcicum*, Rheum.

RHABARBE, Rheum.—*r. Blanche*, Convolvulus mechoacan.—*r. Fausse*, Thalictum.—*r. des Indes*, Convolvulus mechoacan.—*r. des Moines*, Rumex Alpinus.—*r. Sauvage*, Rumex patientia.

RHABDOIDES (SUTURA), Sagittal suture.

RHABILLEUR, Renover.

RHACHETÉ, Rachitæ.

RHACHETRUM, Medulla spinalis.

RHACHIEI, Rachitæ.

RHACHIAGRA, Rachisagra.

RHACHIALGIA, Rachialgia, Rhachiodynia, Spinal irritation.

RHACHIALGITIS, Spinal irritation, Myelitis.

RHACHIDAGRA, Rachisagra, Rachialgia, Rhachiodynia.

RHACHIOCYPHOSIS, Gibbositas.

RHACHIODYN'IA, from *'paxis*, 'the vertebral column,' and *odyn'*, 'pain,' *Rhachial'gia*. Pain in the vertebral column.

RHACHIOMYELITIS, Myelitis.

RHACHIOPLEGIA, Paraplegia.

RHACHIOSCOLIOMA, Scoliosis.

RHACHIOSCOLIOTIS, Scoliosis.

RHACHIOSTROPH'OSIS, from *'paxis*, 'the vertebral column,' and *στροφω*, 'I turn'; *Contortio colum'næ vertebral'is*. Deviation of the vertebral column.

RHACHIOTOMUS, *Rhachiot'omum*, from *'paxis*, 'the vertebral column,' and *τομή*, 'incision.' An instrument for opening the vertebral column.

RHACHIPA'GIA, *Rachipa'gia*, from *'paxis*, 'the spine,' and *παγω*, 'I fasten.' A monstrous union of two fetuses, in which they are connected at the spine.

RHACHIPARALYSIS, Paraplegia.

RHACHIPHYMA, Rachiphyma.

RHACHIRRHEUMA, Lumbago.

RHACHIS, Vertebral column.—*r. Nasi*, see Nasus.

RHACHISAGRA, Rachisagra.

RHACHITÆ, from *'paxins*, appertaining or relating to *'paxis*, 'the vertebral column'; *Rhachia'i*, *Rachi'tæ*, *Rhache'tæ*, or *Rachia'i*. The muscles of the spine.

RHACHITES, Medulla spinalis.

RHACHITICUS, Rachiticus.

RHACHITIS, Rachitis.—*r. Adultorum*, Mollities ossium.—*r. Spinalis*, Rachisagra.

RHACHITISMUS, Rachitis.

RHACHOSIS, Rachosis.

RHACHUS, *Rhacus*, *Rhagus*, from *'payw*, or *'pnyvui*, 'to break.' A wrinkle. A laceration of the skin. A lacerated ulcer—*ulcus lac'erum*.

RHACUS, Rhachus, Wrinkle.

RHÆBOCRANIA, Torticollis.

RHAG'ADES, *'payades*, *Rhaga'dia*, from *'payw* or *'pnyvui*, 'to break or rend'; *Scissu'ra*, *Fissu'ra*, *Fina*. Ruptures, chaps, or narrow and long ulcers, which form near the origin of mucous membranes, especially around the anus, and which are generally owing to the syphilitic virus.

RHAGADES, see Fissure.

RHAGÉ, *Ragé*, *'payn*. Same etymon. A suffix denoting violent rupture or discharge, as in Hemorrhage, Menorrhagia, &c.

RHAGE, Fissure.

RHAGES. *Uvæ passæ.*

RHAGI'US, Rhagoides.

RHAGODETIS, Uveitis.

RHAGOÏ'DES, *Rhagoï'des, Rhagoï'deus, Rha'gicus*, from *ῥαγή, ῥαγος*, 'a berry,' 'a grape.' Grape-like; as *Tunica Rhagoides*: the uvea. Also, the choroid coat of the eye.

RHAGUS, Rhachus, Wrinkle.

RHAMNI BACCÆ, see *Rhamnus*—*r. Succus*, see *Rhamnus*.

RHAMNUS, *ῥαμνος, R. catharticus, Spina cerci'na* seu *domestica* seu *infecto'ria, Rhamnus solutivus, Cerviepi'na, Buckthorn, Purging buckthorn, (F.) Nerprun purgatif, Bourguépine, Épine de cerf. Family. Rhamnaceæ. Ser. Syst. Pentandria Monogynia. The berries—*Rhamni Baccæ*—have a faint and disagreeable odour; and bitterish, nauseous taste. Their juice—*Rhamni Succus*—is cathartic, and apt to excite griping, which must be obviated by the use of diluents.*

RHAMNUS ALATER'NUS, *R. Hispan'icus, Alater'nus latifo'lius, Common alaternus, (F.) Alaternæ*. The leaves have been used as detergents and astringents, in the form of gargle especially.

RHAMNUS FRAN'GULA, *Black Alder, Frangula Alnus, Alnus Nigra, (F.) Aune noir, Bourdane, Bourguène*. All the parts of this tree, as well as of the common alder, are astringent and bitter. The bark is most so. The inner, yellow bark of the trunk or root acts as a cathartic. It has also been employed as an anthelmintic.

RHAMNUS HISPANICUS, *R. alaternus*—*r. Jujuba*, see *Jujube*—*r. Lotus*, see *Jujube*—*r. Ziziphus*, see *Jujube*.

RHANTER, *Canthus, (greater.)*

RHAPHANEDON, *Raphanodon.*

RHAPHANIA, *Raphania.*

RHAPHANIS, *Raphanus hortensis.*

RHAPHANUS, *Raphanus hortensis.*

RHAPHE, *Raphe, Suture.*

RHAPHOSYMPH'YSIS, from *ραφή, 'suture,'* and *συνήναι, 'union.'* Union by suture,—as of the bones of the cranium.*

RHAPHYS, *Raphanus hortensis.*

RHAPONTICUM, *Rheum Rhaponticum*—*r. Vulgare, Centaurea centaurium.*

RHAPONTIQUE, *Rheum rhaponticum.*

RHAPS, *Raphanus hortensis.*

RHAPYS, *Raphanus hortensis.*

RHATANIA, *Krameria ratanhia.*

RHAX, see *Vitis vinifera.*

RHEGE, *Cramp, Contusion, Laceration.*

RHEGMA, *Cramp, Contusion, Laceration,*

Rhexis—*r. Ligamentare, Apospasma.*

RHEGMUS, *Cramp, Contusion, Laceration.*

RHEMBASMUS, *Somnambulism.*

RHEMBE, *Rhembus, 'a wandering about,'* from *ῥεμβειν, 'to wander about.'* Irregularity in the return of febrile paroxysms.

RHEMBUS, *Rhembe.*

RHENCHUS, *Rattle, Snoring.*

RHENXIS, *Rattle, Snoring.*

RHEON, *Rheum.*

RHETINE, *Resin.*

RHETINOCERON, *Resinatum.*

RHÉTINOLÉ. A name given by M. Béral to a medicine, which results from the union of a resin with other medicinal substances. The *Rhétinolés* of Béral differ from the *Rhétinoïdes* in containing but one resin, whilst in the latter there are several.

RHÉTINOÏDES, see *Rhétinolé.*

RHETINOTON, *Resinatum.*

RHEUM, (pronounced *room*), *Rheu'ma, ρευμα, (F.) Rhume*; from *ῥεω, 'to flow.'* Any thin watery discharge from mucous membranes or skin; as the thin discharge from the air passages arising from cold.

RHEUM. SALT. A popular name in the United States, like 'tetter,' for various cutaneous affections of the eczematous and herpetic forms more especially. It is at times applied to a kind of chronic psoriasis; and likewise to cutaneous affections in those who immerse the hands in metallic or acid solutions. Webster defines it "a vague and indefinite popular name, applied to almost all the non-febrile cutaneous eruptions which are common among adults, except ring-worm and itch."

RHEUM, *Rhabar'barum, Rheon, Rha, Rheum, Lap'athum orienta'le* seu *Chinen'se, Rhabar'barum ve'rum* seu *Tartar'icum* seu *Siber'icum* seu *Tur'cicum, Rhubarb, (F.) Rhabarbe, Rhubarbe*. From *Rha*, now called *Volga*, a river in Russia from which it was first brought. *Ord. Polygonaceæ. Ser. Syst. Enneandria Trigynia.* The root—*Rheum*—is the part used in medicine. Its odour is aromatic, peculiar, and rather nauseous; taste, somewhat aromatic, subacid, bitterish, and astringent. It colours the saliva and the urine yellow. It is easily pulverized, and forms a powder of a bright buff-yellow colour. Both water and spirit extract its virtues. *Rhubarb root* is purgative, stomachic and astringent. [?] It is chiefly used for the properties first mentioned. Externally, the powder is sometimes sprinkled over ulcers, to assist their granulation. Dose, ℥j to ʒss of the powder. In a smaller dose, it is stomachic.

Numerous species of *Rheum, Rh. palmotum, Rh. undulatum, and Rh. compactum*, have purgative roots.

The rhubarbs most used in the United States are the *Chinese or India Rhubarb*—*Rheum Sinen'se* seu *Indicum*; the *Russian, Turkey or Bucharian Rhubarb*—*Rheum Rus'sicum* seu *Tur'cicum*; and *European Rhubarb*.

RHEUM COMPACTUM, *Rheum*—*r. Indicum*, see *Rheum*—*r. Palmatum, Rheum*.

RHEUM RHAPONTICUM, *Rhapontic Rhubarb; Rhaponticum, Rhabar'barum Diosco'ridis* seu *Antiquo'rum, (F.) Rhapontique, Rhapontic, Rapontic*. The root of this seems to have been the rhubarb of the ancients. It is more astringent than that of the *rheum palmatum*, and less purgative; for the latter purpose, two or three drachms being required for a dose. *French, Rhapontic, or Crimea Rhubarb*, is obtained from this species, as well as from *Rh. undulatum, and Rh. compactum*.

RHEUM RUSSICUM, see *Rheum*—*r. Sinense*, see *Rheum*—*r. Turcicum*, see *Rheum*—*r. Undulatum*, see *Rheum*.

RHEUMA, *Catarrh, Diarrhœa, Rheum, Rheumatism*. Also, inflammation of a fibrous tissue, as in rheumatism and gout.—*Fuchs*.

RHEUMA CATARRHALE, *Bronchitis, (chronic)*—*r. Epidemicum, Bronchitis, chronic, Catarrh, epidemic*—*r. Pectoris, Catarrh*—*r. Uteri, Leucorrhœa*.

RHEUMAPYRA, *Rheumatism, acute.*

RHEUMARTHITIS, see *Rheumatism, acute.*

RHEUMARTHROSIS, see *Rheumatism, acute.*

RHEUMATALGIA, *Rheumatism, chronic*—*r. Arthritica*, see *Rheumatism, acute.*

RHEUMATIC, *Rheumat'icus, Rheumatis'mal, (F.) Rhumatismal, Rhumatique*. Belonging to rheumatism; as rheumatic fever. Also, applied to one affected with rheumatism; (F.) *Rhumatisant*.

RHEUMATIC DIATHESIS, *Diathesis rheumat'ica*. The special condition of the body that gives occasion to rheumatism. It may exist without presenting the usual phenomena in the joints; whilst the heart may be seriously affected.

RHEUMATICS, *Rheumatism.*

RHEUMATISM, *Catarrh'a* from *ῥευμα, 'peuparos, 'a defluxion, catarrh.'* *Dolor Rheumat-*

icus et arthriticus. (Hoffman.) *Rheumatosis Myosotis*, *Rheuma*, *Myodyn'ia*, (vulg.) *Rheumatiz*, *Rheumatic*, (F.) *Rhumatisme*. A kind of shifting phlegmasia or neuralgia, sometimes seated in the muscles, sometimes in the parts surrounding the joints; and at others, within them. Hence the names *Muscular*, *Articular*, and *Synovial*, which have been applied to it. The disease may be *acute*, or *chronic*.

RHEUMATISM OF THE ABDOMEN, Coeliorrheuma.

RHEUMATISM, ACUTE, *Rheumatismus acutus* seu *articulorum acutus* seu *universalis febrilis* seu *inflammatorius* seu *hypersthenicus*, *Arthrosia acuta*, *Rheumatop'gra*, *Rheumap'gra*, *Febris rheumatica inflammatoria*, *Synocha rheumatica*, *Myosotis*, *Myitis*, *Cama* seu *Arthritis rheumatismus*, *Arthro-rheumatismus*, *Synovitis rheumatismalis*, *Hæmo-arthritis*, *Rheumatic Fever*, (F.) *Rhumatisme aigu*, *Rh. articulaire aigu*, *Fièvre rhumatismale*, usually comes on with the ordinary symptoms of fever: soon after which, or simultaneously, or even before the appearance of febrile signs, excruciating pains are felt in different parts of the body, particularly in the larger joints, which are more or less red and swollen; the pain shifting from one to the other, at times with great rapidity. The disease rarely terminates in less than six weeks, during the greater part of which period the febrile symptoms remain severe; and what is peculiar to the disease, the skin may be covered daily with a profuse perspiration, although it feels extremely hot, and the pulse appears in no way modified by it. It is one of the essential symptoms of the affection, and consequently affords no relief. The only danger to be apprehended from acute rheumatism, notwithstanding the apparent severity of the symptoms, is the translation or extension of the disease (*metastatic rheumatism*) to some internal part, especially to the heart. This tendency must always be borne in mind. *Acute* rheumatism seldom terminates in *chronic*; they who are liable to the former are rarely so to the latter, and conversely. The disease will generally run its course in spite of treatment. The usual antiphlogistic plan had better be pursued during the first days of the disease. Blood-letting may be employed, if the vigour of the patient and of the disease seem to require it; but, although the blood is generally buffy, bleeding does not usually seem to afford much relief. The bowels must be kept open; effervescent draughts be administered, and opium be given in full doses to allay pain. This is, generally, all that is necessary during the first week; after which, advantage will usually be obtained from the exhibition of the sulphate of quinia with sulphuric acid. It does not augment the excitement, and will be found proper in almost all cases. Rheumatic inflammation is very different from other varieties, and this may account for the anomaly. After a few weeks, the disease usually goes off, but leaves the patient very liable to a recurrence on slight exposures or errors of diet. At times, acute rheumatism of the joints puts on the appearance of gout, and seems to be a complication of the two affections. It is then called *Gouty* or *Arthritic rheumatism* or *Rheumatic gout*, *Rheumat'gia arthritica*, *Rheumarthrosis*, *Rheumarthrit'is*, *Arthritis rheumatica* seu *juvenilis*.

RHEUMATISM OF THE ARM, *Brachiorrheuma*—*r. Arthritic*, see *Rheumatism*, acute—*r. of the Bladder*, *Cystorrhæma*—*r. of the Bowels*, *Enterorrheuma*.

RHEUMATISM, CAP'SULAR, *Rheumatismus capsularis*. Rheumatism seated in the living membrane of the joints and bursæ of the tendons. The parts most liable to its attacks are the feet and hands, where it is generally easily recognised

by the enlargement of the joints; but the peculiar characters of the disease are, perhaps, most strikingly seen when it attacks the knee.

RHEUMATISM OF THE CHEST, Pleurodynia.

RHEUMATISM, CHRONIC, *Arthrosia chronica*, *Rheumatismus chronicus* seu *nonfebrilis* seu *vulgaris* seu *inveteratus* seu *habitualis* seu *frig'idus*, *Rheumat'gia*, *Rhumat'gia*, *Rheumatodyn'ia*, *Arthrodyn'ia*, *Arthritis arthrodyn'ia*, (F.) *Rhumatisme chronique*, *Rhumat'gie*. is attended with pains in the hips, shoulders, knees, and other large joints. These are at times confined to one joint; at others, shift from one to another, without occasioning inflammation, or fever. In this manner, the complaint often continues for a great length of time, and then goes off. There is no danger attendant upon it; but the patient may become lame, and is always liable to painful recurrences. Neither variety generally terminates in suppuration; but effusion of coagulable lymph is apt to occur, so as to occasion permanent thickening of the parts. The great preventive of chronic rheumatism, and one of the most valuable curative agents, is flannel, worn next the skin. The whole class of rubefacients may, also, be used with advantage. The warm bath, especially the natural warm bath, the temperature of which does not vary, is perhaps the most successful remedy of all.

RHEUMATISM OF THE EXTREMITIES, *Acrorrheuma*—*r. of the Face*, *Prosoporrheuma*—*r. Gouty*, see *Rheumatism*, acute—*r. of the Heart*, *Cardiorrheuma*—*r. Lead*, see *Lead rheumatism*—*r. of the Leg*, *Scelorrheuma*—*r. of a Limb*, *Merorrheuma*—*r. Local*, *Merorrheuma*—*r. of the Loins*, *Lumbago*—*r. Metastatic*, see *Rh. acute*—*r. of the Neck*, *Torticollis*—*r. Partial*, *Merorrheuma*.

RHEUMATISM, PREABDOMINAL, (F.) *Rhumatisme pré-abdominal*. Rheumatism of the muscles of the anterior and lateral portions of the abdomen; supposed by MM. Gênést and Requin to be the same as the *nervous affection of the psoas* of Gooch.

RHEUMATISM ROOT, Jeffersonia Bartoni.

RHEUMATISM OF THE SKIN, see *Dermat'gia*—*r. Topical*, *Merorrheuma*—*r. of the Uterus*, *Metorrheuma*.

RHEUMATISM, VISCERAL. Rheumatism affecting the muscular or fibrous tissues of the viscera.

RHEUMATISM WEED, *Pyrola maculata*, *P. umbellata*.

RHEUMATISMAL, Rheumatic.

RHEUMATISMUS ABDOMINIS, *Coelorrheuma*—*r. Acutus*, *Rheumatism*, acute—*r. Articulorum acutus*, *Rheumatism*, acute—*r. Brachii*, *Brachiorrheuma*—*r. Cancerosus*, *Neuralgia faciei*—*r. Cervicis*, *Torticollis*—*r. Colli*, *Torticollis*—*r. Cordis*, *Cardiorrheuma*—*r. Cruris*, *Scelorrheuma*—*r. Dorsalis*, *Lumbago*—*r. Extremitatum*, *Acrorrheuma*—*r. Faciei*, *Prosoporrheuma*—*r. Febrilis*, *Dengue*—*r. Inflammatorius* seu *hypersthenicus*, *Rheumatismus acutus*—*r. Intestinorum*, *Enterorrheuma*—*r. Intestinorum cum ulcere*, *Dysentery*—*r. Larvatus*, *Neuralgia faciei*—*r. Localis*, *Merorrheuma*—*r. Membrorum singulorum*, *Merorrheuma*—*r. Non febrilis*, *Rheumatism*, acute—*r. Partialis*, *Merorrheuma*—*r. Pectoris*, *Pleurodynia*—*r. Phlegmonodes*, *Myositis*—*r. Spurius nervosus*, *Neuralgia*—*r. Universalis febrilis*, *Rheumatism*, acute—*r. Uteri*, *Metorrheuma*—*r. Vesicæ urinariæ*, *Cystorrhæma*—*r. Vulgaris* seu *inveteratus* seu *habitualis* seu *frigidus*, *Rheumatism*, chronic.

RHEUMATIZ, Rheumatism.

RHEUMATODYNIA, *Rheumatism*, chronic.

RHEUMATOPHTHALMIA, *Rheumophthalmia*; from *ρευμα*, *ρευματος*, 'rheumatism,' and *οφθαλμία*, 'inflammation of the eyes.' Rheumatic ophthalmia.

RHEUMATOPYRA, Rheumatism, *acuta*

RHEUMATOSIS, Rheumatism.

RHEUMOPHTHALMIA, Rheumatophthalmia. Scleritis.

RHEXIS, *ῥῆξις*, *Rhegma*, *Diruptio*, *Ruptio*, *Ruptura*, *Rupture*. From *ῥήνυμι*, 'to break.' Rupture of a vessel or organ. Spontaneous opening of an abscess.

RHEXIS OC'ULI. Bursting of the eye from any cause; spontaneous or excited.

RHINO'SIS, from *ῥινοσσις*, 'to become wrinkled;' *Cutis corrugatio*. Wrinkling of the skin from extenuation of the body; opposed to *εκασις*, or distension from repletion.

RHIGOS, Rigor.

RHIN, Nasus.

RHINAL'GIA, *Rhinodyn'ia*, from *ῥιν*, 'the nose,' and *αλγος*, 'pain.' Pain in the nose.

RHINAL'GIA AB INTRU'SIS. Pain in the nose from foreign bodies.

RHINAL'GIA VERMINO'SA. Pain in the nose from worms.

RHINAN'CHONĒ, *Rhinan'gia*, *Rhinostegno'sis*; from *ῥιν*, 'the nose,' and *αἷον*, 'narrowness.' Contraction of the nose. Snuffles.

RHINANGIA, Rhinanchone, Snuffles.

RHINCHOS, Rostrum.

RHINENCEPHALIA, Cyclopia.

RHINENCHYSIS, *Rhinenchys'ia*, from *ῥιν*, 'the nose,' *εν*, 'in,' and *χύνω*, 'I pour.' An infusion or injection, made into the nostrils, with a syringe called *Rhinen'chites*.

RHINES, Nares.

RHINIT'IS, *Nasi'tis*, from *ῥιν*, *ῥίνας*, 'the nose,' and *ιτις*, denoting inflammation. Inflammation of the nose. Coryza.

RHINITIS ULCEROSA, Ozæna.

RHINOBLENNORRHŒA, Rhinorrhœa.

RHINOC'ACE, *Cory'zu scarlatino'sa*; from *ῥιν*, *ῥίνας*, 'the nose,' and *κακος*, 'evil.' The offensive discharge from the nose, and other symptoms indicating a morbid condition of the Schneiderian membrane, that accompany the worst forms of scarlatina.

RHINOCATARRHUS, Coryza.

RHINOCNESMUS, from *ῥιν*, *ῥίνας*, 'the nose,' and *κνέσσω*, 'itching.' Itching of the nose.

RHINODYNIA, Rhinalgia.

RHINOPHO'NIA, from *ῥιν*, *ῥίνας*, 'the nose,' and *φωνή*, 'the voice.' A nasal voice, called, also, *Dyspho'nia immodulata nasalis*, *Porapho'nia nasalis* seu *res'onans*, *Psellis'mus na'sitas*, *Myceteropho'nia*, *Na'sitas*, *Vox* seu *Parala'lia nasalis*. Speaking through the nose.

RHINOPLAS'TIC, *Rhinoplas'tica* (ars), *Rhinoplasty*, from *ῥιν*, *ῥίνας*, 'the nose,' and *πλασσω*, 'I form.' An epithet applied to the operation for forming a new nose. The *Tuliacotian operation*, so called because described by Tagliacozzi. It consists in bringing down a portion of flesh from the forehead, and causing it to adhere to the anterior part of the remains of the nose.

RHINOP'TIA, from *ῥιν*, 'the nose,' and *οπτάνω*, 'I see.' The act of seeing over the nose. Squinting over the nose.

RHINORRHAGIA, Epistaxis.

RHINORRHAPHY, *Rhinorrhaph'ia*, from *ῥιν*, *ῥίνας*, 'nose,' and *ῥάφω*, 'a suture.' An operation for the removal of epicanthis, which consists in pinching up a longitudinal fold of the skin, including this fold in two elliptical incisions, removing it, and bringing together the edges of the wound by harelip suture.

RHINORRHŒA, *Rhinoblenorrhœa*, from *ῥιν*, *ῥίνας*, 'the nose,' and *ῥέω*, 'I flow.' A discharge of limpid mucus from the nose without any inflammatory symptom. A gleet from the nose.

RHINO'SIS, from *ῥίνας*, 'a skin, leather,' or from *ῥίνας*, 'rugged.' The state of looseness and excess of skin observed in phthisis.—Vogel.

RHINOSTEGNOSIS, Rhinanchone.

RHIPTASMOS, Jactation.

RHIPTOLUSIA, Affusion.

RHIS, Nasus.

RHIZA, Radix.

RHIZAGRA, Punch.

RHIZO'MA, *Rhizome*, from *ῥίζα*, 'a root.' The part of the root of a plant, which consists of wood or flesh, as in the carrot.

RHIZOPH'AGUS, from *ῥίζα*, 'a root,' and *φαγώ*, 'I eat.' One who lives on roots.

RHIZOTOMIST, see Simples.

RHIZOTOMUS, Apothecary.

RHOA, Punica granatum.

RHODELŒ'ON, *Oleum rosa'rum*. Oil of roses, or oil impregnated with roses.

RHO'DIA, *Rhodi'ola ro'sea*, *Rosea*, *Rosewort*, (F.) *Rhodiola*, from *ῥόδον*, 'a rose,' owing to its odour. *Sex. Syst.* Diercia Octandria. The root, *Roseroot*, when dry, has a very pleasant smell, and is slightly astringent. It is not used.

RHO'DINUS, *ῥοδίνος*, *Rho'dinus*, *Rosa'ceus*, *Ro'seus*, *Rosa'tus*, (F.) *Rosat*, *Rosy*. Formed of, or containing, roses.

RHODIOLA ROSEA, Rhodia.

RHODIOLE, Rhodia.

RHODITES VINUM. Wine in which roses have been macerated.

RHO'DIUM LIGNUM, *Rhodium* or *Rosewood*, from *ῥόδον*, 'a rose.' The wood or root of a tree supposed to be *Geni'sta Canarien'sis* of Linnæus. (F.) *Bois de rose*, *Bois de Chypre*. *Ord.* Leguminosæ. *Sex. Syst.* Diadelphia Decandria. The essential oil is a perfume, and possesses cordial and tonic virtues. Its smell is attractive to fish, rats, &c.

RHODIUS, Rhodinus.

RHODODAPHNE, Nerium oleander.

RHODODAPHNIS, Nerium oleander.

RHODODENDRON CHRYSANTHEMUM, *R. chrysanthum* seu *aur'eum* seu *officina'le*, *Oleander*, *Rose-bay*, *Yellow rhododendrum*, *Mountain Laurel*, (F.) *Rosage*, *R. chrysanthe*, *Rose de Sibérie*, from *ῥόδον*, 'a rose,' and *ένδρουν*, 'a tree.' *Nat. Ord.* Ericacææ. *Sex. Syst.* Decandria Monogynia. This plant has been recommended in rheumatism, gout, and syphilis. In an overdose, it produces acro-narcotic symptoms.

RHODODENDRON MAX'IMUM, *Great Laurel*, *American Rose-bay*. An indigenous shrub, *Ord.* Ericacææ, which flowers in July. It is stimulant and astringent, and has been used by the Indians in gout and rheumatism.

RHODODENDRON PUNCTATUM. Indigenous; has similar properties.

RHODOM'ELA PALMA'TA, (*ῥόδον*, 'a rose,' and *μελί*, 'honey,') *Dulse*, *Saccharine Fucus*. One of the Algæ, eaten in large quantities throughout the maritime countries of the north of Europe, and in the Grecian Archipelago.

RHODOMELI, Mel rosæ.

RHODON, Rosa, Rosa centifolia.

RHODOSACCHARUM, *Confectio rosæ galicæ*.

RHODOSTACTON, Mel rosæ.

RHODOSTAGMA, Aqua rosæ.

RHŒA, Punica granatum.

RHŒAS, see Papaver rhœas.

RHOEBDESIS, Absorption.

RHŒUM, Rheum.

RHOGMOS, Rattle.

RHOI'TES, *ῥοῖντες*, a wine impregnated with the pomegranate; from *ῥοα*, the *Pomegranate*, — Dioscorides. A confection, made by boiling the

juice of the pomegranate in honey. — Paulus of Ægina.

RHOMBOÏDE'US, from *ῥομβος*, 'a rhombus,' whose sides are equal, with two obtuse and two acute angles. *Rhomboideus major* and *minor*, *Rhomboïdes*, *Cervici-dorso-scapulaire*. A muscle situate at the posterior inferior part of the neck, and at the posterior part of the back. It has a rhomboidal shape, and is attached, by its inner edge, to the posterior cervical ligament, and to the spinous processes of the last cervical vertebrae, and of the first four or five dorsal. By its outer edge, it is attached to the spinal or posterior edge of the scapula. It is divided into two fasciuli, constituting the *Rhomboideus major* and *Rh. minor*, the latter being situate higher than the other. This muscle draws the scapula backwards and upwards, and impresses upon it a rotatory motion, which brings its inferior angle nearer to the spine; and, consequently, depresses the anterior angle and the shoulder.

RHOMBOS, *ῥομβος*. A bandage mentioned by Galen, so called on account of its rhomboidal shape.

RHONCHAL, *Rhonchal'is*, from *rhonchus*, 'rattle.' Relating or appertaining to rhonchus; — as 'rhonchal fluctuation,' the fremitus that occurs through the influence of respiration as an attendant on cavernous rhonchi. — Walshe.

RHONCHUS, Rattle, Snoring, Stertor — *r. Cavernosus*, see Gurgling — *r. Cavernulosus*, see Gurgling.

RHONCHUS, CLICKING. A rhonchus which has been considered positively indicative of the existence of tubercles in the lungs. [?]

RHONCHUS, CREPITANS, *Râle crépissant* — *r. Crepitans Redux*, see Crepitant — *r. Mucous*, see *Râle muqueux* — *r. Sibilans*, see *Râle crépissant* — *r. Sibilus*, *Râle sibilant* — *r. Sonorus*, *Râle sonore* — *r. Subcrepitant*, see *Râle muqueux*.

RHOPALISMUS, Priapismus.

RHOPALOSIS, Plica, Pupil, Uvea.

RHOPE, Congestion.

RHOTACISMUS, Rotacism.

RHUBARB, Rheum — *r. Chinese*, see Rheum — *r. French*, Rheum Rhaponticum — *r. India*, see Rheum — *r. Krimea*, Rheum Rhaponticum — *r. Monk's*, Rumex Alpinus, Rumex patientia — *r. Poor man's*, Thalictron — *r. Rhapontic*, Rheum Rhaponticum — *r. Russia*, see Rheum — *r. Turkey*, see Rheum — *r. Wild*, Convolvulus panduratus.

RHUBARBE, Rheum — *r. Fausse*, Thalictron — *r. des Indes*, Convolvulus mechoacan — *r. des Pauvres*, Thalictron — *r. Sauvage*, Rumex patientia.

RHUE, Ruta.

RHUM, Rum.

RHUMAPYRA, Rheumatism, acute.

RHUMATALGIA, Rheumatism, chronic.

RHUMATALGIE, Rheumatism, chronic.

RHUMATIQUE, Rheumatic.

RHUMATISANT, Rheumatic.

RHUMATISMAL, Rheumatic.

RHUMATISME, Rheumatism — *r. Aigu*, Rheumatism, acute — *r. Articulaire aigu*, Rheumatism, acute — *r. Chronique*, Rheumatism, chronic — *r. Préabdominale*, Rheumatism, præabdominal — *r. d. l'Utérus*, Metrorrhœmia.

RHUME, Rheum — *r. de Cerveau*, Coryza — *r. de Poitrine*, Catarrh.

RHUS COPALLINA, see Rhus glabra.

RHUS CORIARIA, *Adu'rin*, Sumach, Shumach, Elm-leaved Sumach, (F.) *Sumac des Corroyeurs*, *Roure*. Ord. Anacardiaceæ. Sex. Syst. Pentandria Monogynia. This is the only species of the genus rhus that is innocent. Both the leaves and berries have been used as astringents and tonics.

RHUS GLABRA, 'Smooth Sumach.' Indige-

nous, and official in the Pharmacopœia of the United States. An infusion of the fruit — *Rhus glabra* [?] — which is acid and astringent, has been used as a cooling drink, and as a gargle. An infusion of the inner bark of the root has been recommended in mercurial ptyalism.

RHUS RADICANS, *R. vernia*, *Poison vine* (America). This plant is poisonous (see Poisons, Table of), and is extremely acrid, when applied to the skin, producing erysipelas and vesications, and has been used, like the next variety, in paralysis, and other nervous affections.

RHUS SYLVESTRIS, Myrica gale.

RHUS TYPHINA, *Stag's horn Sumach*, and **RHUS COPALLINA**, *Mountain Sumach*, have similar properties.

RHUS TOXICODENDRON, *Toxicodendron*, *Poison Oak* (America), (F.) *Sumac vénéneux*. This plant has been used in paralytic affections and cutaneous eruptions. It is poisonous, like the last. Dose, gr. ss. to gr. iv.

RHUS TYPHINA, see Rhus glabra.

RHUS VENENATA, *Poison Sumach*, and **RHUS TYPHINA**, have similar properties.

RHUS VERNIX, *Rh. radicans*.

RHYAS, from *ῥέω*, 'I flow.' An ulceration of the lachrymal caruncle, with a constant flow of tears. — Blancard.

RHYPARIA, Rupia, Sordes primarum viarum.

RHYPIA, Rupia.

RHYPOS, Sordes.

RHYPTICA, Detergents.

RHYPTICUS, Abstergent.

RHYSIS, Flux.

RHYSMUS, Rhythm.

RHYSEMA, Wrinkle.

RHYTHM, *Rhythmus*, *ῥυθμος*, *Rhythmus*, (F.) *Rhythme*, 'measured movement.' The order of proportion, as regards time, which reigns between the different movements of an organ — as of the organism in health and disease. Thus, rhythm is applied to the diurnal variations of the pulse; and to the paroxysmal movements of an intermittent.

RHYTHMICAL, Periodical.

RHYTHMICAL CONTRACTIONS OF MUSCLES are those that succeed one another after regular intervals.

RHYTHMUS, Rhythm.

RHYTIDOSIS, Rutilosis.

RIB, Costa — *r. False*, see Costa — *r. Floating*, see Costa — *r. Sternal*, see Costa — *r. True*, see Costa.

RIBES GROSSULARIA, *R. uva crispa*.

RIBES NIGRUM, *R. olidum*, *Grossularia nigra*, *Caseie*, *Black Currant*, *Gazel*, (F.) *Groseillier noir*. Ord. Grossulacææ. Sex. Syst. Pentandria Monogynia. The berries — *Black currants* — have been recommended in sore-throat, as diuretics. The leaves have been advised for the same purpose. They are chiefly used as preserves.

RIBES OLIDUM, *R. nigrum* — *r. Officinale*, *R. rubrum*.

RIBES RUBRUM, *R. officinale* seu *vulgarè*, *Grossularia non spinosa* seu *rubra*, *Red currant*, (F.) *Groseillier rouge*; of which the *White* is only a variety. It is cultivated, and affords an agreeable fruit — *Red currants*, (F.) *Groseilles*. It possesses the qualities of the subacid fruits in general, and makes an excellent preserve.

RIBES UVA CRISPA, *R. grossularia*, *Common gooseberry*. The fruit — *Grossularia*, (F.) *Groseille à maquereau*, *Gooseberry*, *Feaberry* — is agreeable, and makes a good preserve.

RIBES VULGARE, *R. rubrum*.

RIBGRASS, *Plantago lanceolata*.

RIBS, NECK OF THE, *Collum costarum* — *r. Short*, see Costa.

RIBWORT, *Plantago Virginica*.

RICE, *Oryza*.—*r.* Disease, Cholera.—*r.* Ground, see *Oryza*.—*r.* Jelly, see *Oryza*.

RICE WATER, see *Oryza*. The evacuations of cholera have been termed — from their resemblance to rice-water — ‘rice-water discharges;’ and in India, ‘congee discharges.’ — congee in Hindoostanee meaning ‘rice-water.’

RICE, WILD, *Zizania aquatica*.

RICHARDSONIA, see *Ipeacacuanha*.

RICHLEAF, *Collinsonia Canadensis*.

RICHWEED, *Actæa racemosa*, *Ambrosia trifida*, *Collinsonia Canadensis*, *Pilea pumila*.

RICIN, *Ricinus communis*.

RICINOIDES, *Jatropha curcas*.

RICINUS COMMUNIS, from *ricinus*, ‘a tick,’ which its seed resembles; *Ricinus vulgaris* see *viridis*, *Palma Christi*, *Cataputia major*, *Faba purgatrix*, *Cerna*, *Kerna*, *Cici*, *Castor oil plant*, (F.) *Ricin*, *Fève purgatif*. Ord. Euphorbiaceæ. Ser. Syst. Monœcia Monadelphia. Castor-oil seeds, whence the oil is obtained, are inodorous; taste acrid and slightly sweetish. One or two seeds — *Semina Cataputia majoris* seu *Ricini vulgaris* — will act as a cathartic; but the oil is alone used. It is obtained by bruising the castor-seeds, previously decorticated, and then expressing the oil without the application of heat. Recently-drawn castor oil, *Oleum de kereu* seu *kervinam*. *Alker'na*, O. *Ricini* seu *Cicini* seu *Palma liquidum* seu *de Palmâ Christi*. — in the United States often called simply *Oil*, (F.) *Huile de Ricin* — is inodorous and nearly insipid. It is cathartic, operating speedily, and may be used in all cases where powerfully stimulating purgatives would be injurious. Dose, f ʒi to f ʒiss. Hufeland recommends that a drop of croton oil shall be added to an ounce of the oil of poppy, and that the mixture shall be called *Oleum Ricini Officinale*.

At the Cape of Good Hope the oil is obtained from *Ricinus lividus*.

A variety of *Ricinus* at Cape Verd is employed by the inhabitants to increase the secretion of milk. It is applied in the form of poultice, made with the green leaves, to the mammæ, or of a strong decoction, with which the mammæ and the external parts of generation are washed.

RICINUS LIVIDUS, see *Ricinus communis* — *r.* Major, *Jatropha curcas*.—*r.* Minor, *Jatropha manihot* — *r.* *Viridis*, *R. communis* — *r.* *Vulgaris*, *R. communis*.

RICK, *Malleolus*.

RICKETS, *Rachitis*.

RICKETY, *Rachitosis*.

RICTUS LUPINUS, see *Harelip*.

RIDE, *Wrinkle*.

RIDÉ, *Wrinkled*.

RIDING (of Bones). *Ossium superpositio*, (F.) *Chevauchement*. Displacement of the fragments of a bone, chiefly produced by the contraction of muscles, which occasions shortening of the limb; the fractured extremities riding over each other instead of being end to end.

RIDING-BAG, *Incubus*.

RIFE, (Sax.) *ryfe*, (Prov.) *Brief*. Prevailing; as applied to diseases.

RIFT, *Erectation*.

RIGIDITAS, *Rigidity*.

RIGIDITAS ARTICULORUM, *Ancylosis epu'ria*. Stiffness of the joints. False ancylosis.

RIGIDITÉ, *Rigidity* — *r.* *Cadavérique*, see *Rigidity*.

RIGIDITY, *Rigiditas*, from *rigere*, ‘to stiffen,’ (F.) *Rigidité*, *Roideur*, *Roideur*. Great stiffness of fibre, or want of suppleness. The stiffness of the dead body, *Cadavérique rigidity*, (F.) *Roideur* ou *Rigidité cadavérique*, is one of the signs of the cessation of life. It may be removed, how-

ever, for a time, by the injection into the arteries of oxygenated detribinated blood. See *Rigor mortis*.

RIGOR, *Rhigos*, *Algor*, *Coldness*, *Cheim'ia*, (F.) *Frisson*. Same etymon. Sensation of cold, with involuntary shivering or shaking of the whole body: — a symptom of fever. A slighter degree, and at times full rigor, is occasionally termed a *chill*, (Prov.) *cream*; *frigus ten'ue*, *perfrigeratio*.

RIGOR ARTUUM, *Contractura* — *r.* *Cadaverous*, *Rigor mortis* — *r.* *Corporis mortui*, *Rigor mortis* — *r.* *Emortualis*, *Rigor mortis* — *r.* *Maxillæ inferioris*, *Trismus*.

RIGOR MORTIS, *R. corporis mortui* seu *cadaverous* seu *emortualis*, *Neconarce'ma*, (F.) *Roideur* ou *Roideur* ou *Rigidité cadavérique*. ‘Stiffness of death.’ Death stiffening. The rigidity of limbs that occurs on dissolution. See *Rigidity*.

RIGOR NERVORUM, *Tetanus* — *r.* *Nervosus*, *Tetanus*.

RIMA, *Schisma*, *Schiza*; ‘a fissure, a cleft;’ as *Rima* seu *Rimula Glottidis* seu *Laryngis*, the opening of the glottis. *Rima* seu *R. magna vulvæ*, *Introitus*, *Sinus vulvæ*, *Scapha*, *Scaphus* seu *Fissura magna* seu *Scissura vulvæ*, *Schiza*, the opening of the vulva. See *Vulva*. *Rima oris*, the opening of the mouth.

RIMA CANALIS ORBITARII, *Suborbital fissure* — *r.* *ad Infundibulum*, *Vulva cerebri* — *r.* *Laryngis*, *Rima glottidis* — *r.* *Magna Vulvæ*, see *Rima* — *r.* *Oris*, see *Rima* — *r.* *Transversa Magna Cerebri*, *Fissure*, *transverse*, of the brain — *r.* *Vulvæ*, see *Rima*.

RIMÆ, *Rhagades*.

RIMULA GLOTTIDIS, *Rima glottidis* — *r.* *Laryngis*, *Rima glottidis*.

RINÆUS, *Compressor naris*.

RINCHOS, *Rostrum*.

RING, (Sax.) *hping*, *ping*, (Germ.) *Ring*, *Cricos*, (F.) *Anneau*. A name given to natural, circular, or roundish apertures, with muscular or aponeurotic parietes, which serve for the passage of some vessel or canal: — as the *umbilical ring*, *inguinal ring*, &c.

RINGS, FESTOONED, *Annuli fibro-cartilaginei* — *r.* *Fibro-cartilaginous*, *Annuli fibro-cartilaginei*.

RING-FINGER, *Annular finger*.

RINGMAN, *Annular finger*.

RINGWORM, *Herpes circinatus* — *r.* of the Scalp, *Porrigi scutulata* — *r.* *Ulcerative*, *Herpes exedens* — *r.* *Vesicular*, *Herpes præputii*.

RIO VINAIGRO, *Passambio*. A stream of some size in New Grenada, which is strongly impregnated with sulphuric acid.

RIRE or RIS, *Risus* — *r.* *Canin*, *Canine laugh* — *r.* *Moqueur*, *Canine laugh* — *r.* *Sardonien*, *Canine laugh*, *Risus sardonius* — *r.* *Sardonique*, *Canine laugh*.

RISAGON, *Cassumunar*.

RISIGALLUM, *Orpiment*.

RISING, see *Regurgitation* — *r.* of the Lights, *Cynanche trachealis*.

RISING OF ELEVATION OF THE PULSE, *Pulsus elevatio*. The pulse is said to rise, when it becomes more full and frequent, as occurs in the exacerbation of acute diseases.

RISORNIUS NOVUS. Santorini gives this name to a portion of the platysma myoides, which passes from the cheek towards the commisure of the lips.

RISIGALLUM, *Orpiment*.

RISUS, from *ridere*, *risum*, ‘to laugh.’ *Gelus*, *Laughing*, *Laughter*, *Laugh*, (F.) *Rire* ou *Ris*. An involuntary movement of the muscles of the face, and of the lips in particular, accompanied with a sonorous and interrupted respiration, commonly indicating mirth and satisfaction.

RISUS APIUM, *Ranunculus sceleratus* — *r.* *Caninus*, *Canine laugh* — *r.* *Involuntarius*, *Canine*

laugh—*r. Sardonicus*, Canine laugh—*r. Spasticus*, Canine laugh.

RITA-CRISTINA. A celebrated double monster, born in Sardinia, in 1829. It lived to be nearly nine months old. Two distinct bodies were united at the pelvis. Below, the monster was single.

RIVER WEED, *Conserva rivalis*.

RIVING, Eructation.

RIZ, *Oryza*.

ROAN TREE, *Sorbus acuparia*.

ROB, *Roob, Robub, Rohob, Rohab*, (F.) *Extrait des fruits*. This word, of Arabic extraction, means the juice of any fruit thickened to the consistence of honey by evaporation, before it has fermented. The juice of the grape, boiled to this consistence, was called *Sapa*, and, when not quite so thick, *Defrutum*.

ROB CYNOSBATOS, Confectio rosæ caninæ—*r. ex Moris*, Diamorion—*r. Nucum*, Diacaryon.

ROBIN-RUN-AWAY, *Glechoma hederacea*.

ROBIN-RUN-THE-HEDGE, *Galium aparine*.

ROBINIA AMA'RA, *Ord. Leguminosæ. Ser. Syst. Diadelphica Decandria*. The roots of the *Robinia amara* of Cochinchina are bitter, and have been recommended, especially in that country, in diarrhoea and dyspepsia.

ROBINIA PSEUDO-ACA'CIA, *Pseudacacia odorata*, *False Aca'cia*, *Black Locust*, *Locust-tree*; native of America. The flowers are said to possess antispasmodic properties, and the bark of the root to be emetic and cathartic.

ROBORANS, Corroborant.

ROBORANTIA, Tonics.

ROBUB, Rob.

ROCELLA, Lichen rocella—*r. Tinctoria*, Lichen rocella.

ROCHE'S EMBROCATION FOR HOOPING-COUGH. An empirical preparation, composed of *ol. oliv.* 3xvj, *ol. succin.* 3vij, *ol. caryoph.*, q. s. to scent it strongly.

ROCHER, see Temporal bone.

ROCHETTA ALEXANDRINA, Soda.

ROCKBRIDGE ALUM SPRINGS, see Virginia, mineral waters of.

ROCKET, GARDEN, *Brassica eruca*—*r. Roman*, *Brassica eruca*.

ROCK ROSE, *Helianthemum Canadense*, *H. Corymbosum*.

ROCOU, Terra Orleana.

RODDIKIN, *Abomasus*.

RODENS, see *Sorbus acuparia*.

RODEN-TREE, *Sorbus acuparia*.

RODSCHIEDIA BURSA PASTORIS, *Thlaspi bursa*.

ROGNE, Psora.

ROHAB, Rob.

ROHENA BARK, *Swietenia*.

ROHOB, Rob.

ROHUN, see *Strychnos nux vomica*.

ROIDEUR, Rigidity—*r. Cadaverique*, Rigor mortis.

ROISDORFF, MINERAL WATERS OF. Alkaline waters in Germany, which contain carbonic acid and carbonate of soda.

ROLLER, Bandage, Fascia.

ROMARIN, *Rosmarinus*—*r. des Marais*, *Statice limonium*—*r. Sauvage*, *Ledum sylvestre*.

ROME, (CLIMATE OF.) The climate of Rome, as regards its physical qualities, is considered to be one of the best in Italy. It is peculiarly still; and, hence, has been esteemed favourable as a winter climate in pulmonary diseases, and for invalids generally, as it admits of their taking exercise in the open air at a lower temperature than they otherwise could do. In the early stages of consumption, Sir James Clark generally found the climate favourable.

ROMPEURE, Hernia.

ROMPT-PIERRE, *Pimpinella saxifraga*.

RONCE, *Rubus idæus*—*r. Bleue*, *Rubus cæsius*—*r. de Montagne*, *Rubus chamaemorus*—*r. Noire*, *Rubus fruticosus*.

RONCHUS, Snoring, Stertor.

ROND, Round—*r. Grand*, *Teres major*—*r. Petit*, *Teres minor*.

RONE, *Sorbus acuparia*.

RONFLEMENT, Snoring, Stertor, see Rattle—*r. de Diable*, *Bruit de Diabls*.

RONNACHS, *Triticum repens*.

RONNING, Rennet.

ROOB, Rob.

ROOF OF THE MOUTH, Palate.

ROOM, Pityriasis.

ROOT, Radix—*r. Ague*, *Aletris farinosa*—*r. Bitter*, *Apocynum androsæmifolium*—*r. Canada*, *Asclepias tuberosa*—*r. Colic*, *Aletris*—*r. Red*, *Celastrus*, *Lachnanthes tinctoria*—*r. Rough*, *Liatriis*.

ROOT OF SCARCITY, *Beta hybrida*.

ROOT, SQUAW, *Actæa racemosa*—*r. Sweet*, *Acorus calamus*—*r. Unicorn*, *Aletris farinosa*—*r. White*, *Ligusticum actææfolium*—*r. Yellow*, *Xanthoriza apifolia*, *Jeffersonia Bartoni*.

ROPE, Nanus.

ROPE BARK, *Dirca palustris*.

ROQUETTE, *Brassica eruca*, Soda.

RORELLA, *Drosera rotundifolia*.

RORIFEROUS, *Rorifer*, (F.) *Rorifère*, from *ros*, *roris*, 'dew,' and *ferre*, 'to carry.' An epithet given to vessels which pour exhaled fluids on the surface of organs.

RORIFEROUS DUCT, Thoracic duct.

ROS, *Sperm*—*r. Calabrinus*, *Fraxinus ornus*—*r. Solis*, *Drosera rotundifolia*.

ROSA, (*poëv*), *R. centifolia*—*r. Afzeliana*, *R. canina*.

ROSA ALBA, *R. usitatissima* seu *leucantha*. The *White Rose*, (F.) *Rosier blanc*. *Ord. Rosaceæ. Ser. Syst. Icosandria Polygynia*. The flowers of this species possess similar but inferior qualities to the damask.

ROSA ARMATA, *R. canina*—*r. Asturica*, see *Pelagragra*—*r. Austriaca*, *R. Gallica*—*r. Calycina*, *R. canina*.

ROSA CANINA, *R. sylvestris* seu *senticosa* seu *Swartziana* seu *Afzeliana* seu *armata* seu *calycina* seu *rubra vulgaris*, *Cynorhodon*, *Canirubus*, *Caninus Sentic*, *Cynocytinus*, *Cynospasium*, *Cynosbatus*, *Dog rose*, *Wild Briar*, *Hip Tree*, (F.) *Rose du chien*, *Eglantier de chien*, *E. sauvage*. The fruit of this variety, called *Heps* or *Hips*, (*Hops*, (Sc.) *Doghip*, has a sourish taste, and is formed into a conserve. See *Confectio cynosbati*. It is seldom employed, except to give form to more active remedies; as to pills, boluses, linctuses, &c.

ROSA CENTIFOLIA, *Rosa*, *R. Damascena* seu *Pallida*, *Hecatomphylita*, *Damask Rose*, *Hundred leaved* or *Cabbage rose*, (F.) *Rosier à cent feuilles*, *Rhodon*. The petals of this rose have an extremely fragrant odour, and sub-acidulous taste. They are laxative; but are scarcely used for any purposes except for the distillation of rose water, *Oil of Roses*, *Oleum Rosa*, *Attar*, *Otto* or *Essence of Roses*, and the formation of a syrup.

ROSA DAMASCENA, *R. centifolia*.

ROSA GALICA, *R. Austriaca* seu *rubra*, *Red Rose*, (F.) *Rosier de Provins*. The petals of this variety are astringent; and are used in infusion, conserve, &c.

ROSA LEUCANTHA, *R. alba*—*r. Pallida*, *R. centifolia*—*r. Rubra*, *R. Gallica*—*r. Rubra vulgaris*, *R. canina*—*r. Saltans*, *Essera*—*r. Senticosa*, *R. canina*—*r. Swartziana*, *R. canina*—*r. Sylvestris*, *R. canina*—*r. Usatissima*, *R. alba*.

ROSACEA, *Roseola*.

ROSACEUS, *Rhodinus*.

ROSAGE, *Nerium oleander*, *Rhododendron chrysanthemum*.

ROSAGO. Nerium oleander.

ROSALIA. Scarlatina.

ROSALIE. Scarlatina.

ROSAT, Rhodinus.

ROSATUS, Rhodinus.

ROSE, Erysipelas — r. Cabbage, Rosa centifolia — r. Campion, Lychnis githago — r. Christmas, Heliborus niger — r. Corn, Papaver rhoeas — r. Damask, Rosa centifolia — r. Dog, Rosa canina — r. Dog, conserve or confection of, Confectio rosæ caninae.

ROSE DU CHIEN, Rosa canina.

ROSE, HUNDRED-LEAVED, Rosa centifolia — r. Marine, Rosmarinus.

ROSE RASH, Roseola.

ROSE, RED, Rosa Gallica — r. Rock, Helianthemum Corymbosum.

ROSE DE SIBÉRIE, Rhododendron corymbosum — r. Spots, Spots, red.

ROSE, WHITE, Rosa alba.

ROSEA, Rhodia.

ROSEBALM, SCARLET, Monarda coccinea.

ROSEBAY, AMERICAN, Rhododendron maximum.

ROSE DROP, Gutta rosea.

ROSÉE DU SOLEIL, Drosera rotundifolia.

ROSELLINA, Roseolæ.

ROSEMARY, Rosmarinus — r. Marsh, Statice limonium, and S. Caroliniana — r. Wild, Eriophyllum umbellatum.

ROSENHEIM, MINERAL WATERS OF. Rosenheim is 34 miles S. E. from Munich, in Bavaria. The waters contain sulphuretted hydrogen, carbonic acid, carbonate of lime, chlorides of calcium and magnesium, carbonate of soda, and oxide of iron. They are used as tonics.

ROSEOLA, *Exanthēsis Roseola*, *Rose Rash*, (F.) *Rosole*, *Eruption anomale*, *Rosace*, *Fievre rouge*, *Efflorescence érysipélateuse*, from *rosa*, 'the rose,' on account of the colour. An efflorescence in blushing patches, gradually deepening to a rose-colour, mostly circular or oval; often alternately fading and reviving; sometimes, with a colourless nucleus; and chiefly on the cheeks, neck, or arms. It is frequently found, as a symptom, in dentition, dyspepsia, rheumatism, &c., and is of no consequence. See *Roseolæ*, *Rubeola*, and *Scarlatina*.

ROSEOLA ACNOSA, Gutta rosea.

ROSEOLA ÆSTIVA. The affection which prevails in summer. The eruption is of a darker colour, and is attended with more itching.

ROSEOLA ANNULATA. Characterized by patches in the form of rose-coloured rays, with central areas of the colour of the skin.

ROSEOLA FICOSA, Sycosis — r. Typhosa, Spots, red — r. Variolosa, Roseolæ.

ROSE'OLÆ, *Roseola*, *Roselli'na*, *Rube'ola epuria*, *False or French measles*, (F.) *Roséole*, *Fausse rougeole*. This name is also given by some to an acute exanthem, midway between measles and scarlatina, and which belongs to neither one nor the other, as it affords no protection against either. It is, also, termed *Rubeolæ* by some, and *Rube'oloid*, a term, which is applicable to any eruption resembling rubeola. An eruption which often precedes that of small-pox is termed *Roseola Variolosa*.

ROSÉOLE, Roseola, Roseolæ.

ROSE PINK, Chironia angularis.

ROSES, MILK OF. A cosmetic wash, formed, according to one method, of subcarbonate of potassa, gr. xj; oil of almonds, ℥ij; essence of bergamot, ℥ij; aquæ rosæ, ℥ij; orange flower water, ℥ij. Mix.

ROSEUS, Rhodinus.

ROSEWOOD, Rhodium lignum.

ROSEWORT, Rhodia.

ROSIER BLANC, Rosa alba — r. à Cent feuilles, R. centifolia — r. de Provins, Rosa Gallica.

ROSIN, BLACK, Colophonia — r. Brown, Colophonia.

ROSIN WEED, Silphium laciniatum.

ROSIN, WHITE, see Pinus sylvestris — r. Yellow, see Pinus sylvestris.

ROSIO STOMACHI, Cardialgia — r. Ventriculi, Cardialgia.

ROSMARINUM, Rosmarinus.

ROSMARINUS, *R. officinalis* seu *latifolius* seu *hortensis*, *Alchuchil*, *Libano'tis coronaria*, *Dendrob'anus*, *Rosemary*, (F.) *Romarin*, *Rose marine*, *Encensier*; Ord. Labiatae. Ser. Syst. Diandria Monogynia. The tops of this plant, *Rosmarinus* (Ph. U. S.), *Herba Anthos*, have a fragrant, grateful smell; and an aromatic, warm, bitterish taste, which is dependent upon an essential oil, combined with camphor. Rosemary has been recommended in nervous headaches, in the form of infusion. It is now rarely used, except for its odour. The flower is called *Anthos*, ανθος. Its oil — *Oleum Rosmarini* — is official. From two to six or ten drops are given in nervous complaints, rubbed up with sugar.

ROSMARINUS HORTENSIS, Rosmarinus — r. Latifolius, Rosmarinus — r. Officialis, Rosmarinus — r. Stœchadis facie, Teucrium creticum — r. Sylvestris, Ledum sylvestre.

ROSSALIA, Scarlatina — r. Squamosa, Scarlatina.

ROSSIGNOL. A name given, in France, to an affection to which pressmen are subject. It is a sprain, followed by tumefaction of the radio-carpal joint.

ROSSOLIO, see Spirit.

ROSTRUM, *Rhinchos*, *Rinchos*, *Ramphos*, 'a beak.' (F.) *Bec*. A name given to several old forceps, on account of their resemblance to the beaks of different birds. The principal were the *Rostrum corv'num*, *anati'num*, *psittaci'num*, *vultur'num*, *cyg'neum* seu *olori'num*, *grui'num*, *lacerati'num*, &c.

ROSTRUM EXTERNUM, Olecranon — r. Porcinum, Acromion — r. Posterius, Olecranon — r. Spheoideale, Azygos processus.

ROSY BETTY, Erigeron bellidifolium — r. Bush, Spirea tomentosa — r. Drop, Gutta rosea — r. Wheel, Gutta rosea.

ROT, Erection.

ROT, Grinders', Asthma, grinders'.

ROTACISMUS, *Rotacism'us*, *Rottacismus*, *Rhotacism'us*, *Paellis'mus rhotacis'mus*, (F.) *Grassegement*. A vicious pronunciation of the Greek, Ρ, Ρau, common in the northern parts of England; especially near Newcastle. Called, there, the *Burr* or *Burrh*.

ROTACISMUS, Grassegement, Rotacism.

ROTATOR, from *rotare*, *rotatum*, (*rota*, 'a wheel,') 'to turn round,' (F.) *Rotateur*. A name given to several muscles that turn the parts to which they are attached upon their axes; — such as the oblique muscles of the eyes, the pronators, supinators, &c.

ROTATOR, Trochanter.

ROTATO'RES DORSI. Eleven small muscles on each side of the dorsal vertebrae, which arise from the point or upper edge of each transverse process, and pass to the lower edge of the arch of the vertebra above, as far as the base of the transverse process. Their action is to rotate the individual vertebrae on each other.

ROTATO'RES FEMORIS, Obturatores — r. Oculi, Oblique muscles of the eye.

ROTCOLL, Cochlearia armoracia.

ROTTACISMUS, Rotacism.

ROTLERA TINCTORIA. A euphorbiaceous plant of India. A powder is attached to the fruit capsule of this plant, which is brushed off and col-

lected when the fruit is dry. It is called *Kameyla*, is used in Northern India as a vermifuge for dogs; and has proved effectual against tapeworm in man. The dose is two or three drachms, which purges.

ROTULA, Patella, Trochiscus—r. Genu, Patella.

ROTULAD, see Rotular aspect.

ROT'ULAR; from *rotula*, 'the patella.' Relating or appertaining to the patella.

ROT'ULAR ASPECT. An aspect towards the side on which the rotula is situated.—Barclay. *Rot'ulad* is used by the same writer, adverbially, to signify 'towards the rotular aspect.'

ROTULE, Patella.

ROTULIAN, *Rotulia'nus*, (F.) *Rotulien*. Relating or belonging to the rotula or patella.

ROTUNDUS, Round.

ROUCOU, Terra Orleansa.

ROUEN, MINERAL WATERS OF. This water, known under the name, *Eau des fontaines de la Marequerie*, is cold, and contains carbonate of iron, carbonic acid, carbonate of lime, chloride of calcium, &c. It is tonic.

ROUGE, Colcothar—r. d'Angleterre, Colcothar—r. de Prusse, Colcothar—r. du Sang, Hæmatin.

ROUGEOLE, Rubéola—r. Fausse, Roseolæ.

ROUGEURS, Flush.

ROUGH ROOT, Liatris.

ROUILLE, Rubiginous.

ROULEMENT DU LAMBEAU, see Flap.

ROUND, *Teres*, *Rotun'dus*, (F.) *Rond*. Anatomists have given this name to many organs whose fibres are collected in round fasciculi. See *Teres*.

ROUND FORAMEN, *Fora'men rotund'um*. A foramen of the sphenoid bone, a little behind the *foramen lacerum superius*, which gives passage to the second branch of the fifth pair of nerves, called *superior maxillary*.

ROUND HEART, Thapsium atropurpureum.

ROUND LIGAMENTS OF THE UTERUS, *Ligament'a rotund'a Uteri*, (F.) *Cordons sus-pubiens* (Ch.), *Cordons vasculaires*, *Ligaments ronds de l'utérus*. Two cords, which arise from the lateral and superior parts of the uterus, whence they proceed towards the abdominal rings, which they pass through, and terminate by vanishing in the areolar tissue of the groins, mons veneris, and labia majora. These cords are whitish, flattened, and narrower in the middle than at the extremities. According to Mr. Rainey, they are composed chiefly of the striped muscular fibre.

ROUP, Cynanche trachealis, Raucedo.

ROURE, Rhus coriaria.

ROUTINE PRACTITIONER, Routinist.

ROUTINIST, pronounced *routeen'ist*; (F.) *Continier*; from (F.) *routine*, 'a regular habit or practice, unvaried by circumstances.' One who practises in such an unvaried manner:—a *routine practitioner*.

ROVE, Scab.

ROVING, Delirium.

ROWLAND'S SPRINGS, see Georgia, mineral waters of.

ROWTING, Snoring.

RUBBER, INDIAN, Caoutchouc.

RUBBING SOUND, *Bruit de Frottement*.

RUBE'DO, *Rubor*, 'redness'; from *rubere*, 'to be red.' A diffused, but not spotted, redness on any part of the skin; like that which arises from blushing. Blushing.

RUBEFA'CIENT, *Erethis'ma*, *Erethis'ticum*, *Rubefa'ciens*, from *rubere*, 'red,' and *facio*, 'I make.' (F.) *Rubéfiant*. That which produces redness. A medicine which causes redness of the skin. The action is called *Rubefaction*.

RUBÉFIANT, Rubefacient.

RUBE'OLA, *Rubeola vulga'ris*, from *rubeo*, 'I am red.' *Rose'ola*, *Exanthe'sis Rubeola*, *Vetera'na*, *Blac'cia*, *Fersæ*, *Blactiæ*, *Bovillæ*, *Morbill'i*, *Scarlati'na pustulo'sa seu milia'ris*, *Febbris morbill'o'sa*, *Typhus morbill'osus*, *Phenicia'mus*, *Ephy'mata*, *Measles*, (Sc.) *Mirles*, (Prov.) *Creas*, *Crewds*, *Malls*, *Maules*, *Measlings*, (F.) *Rougeole*, *Fièvre morbillense*. One of the major exanthemata; generally affecting individuals but once, and produced by specific contagion. The rash usually appears on the fourth, but sometimes on the third, fifth, or sixth day of a febrile disorder; and, after a continuance of four days, gradually declines with the fever. The disease generally commences from ten to fourteen days after the contagion has been received. The eruption first shows itself in distinct, red, and nearly circular spots, somewhat less than the ordinary areolæ of flea-bites. As these increase in number, they coalesce; forming small patches of an irregular figure, but approaching nearest to that of semi-circles or crescents. These patches are intermixed with single, circular dots and with interstices of the natural colour of the skin. On the face they are slightly raised, so as to give the sensation of inequality of surface to the finger passed over the cuticle. The disappearance of the eruption is followed by desquamation of the cuticle. Measles is not dangerous of itself, but it is liable to induce pneumonia in winter, and dysentery in summer, which are, at times, very fatal. These are apt to come on at the time of, or soon after, the disappearance of the eruption. When they supervene, they must be treated as idiopathic affections. It demands a general antiphlogistic treatment. Willan has pointed out a kind of RUBÉOLA SPURIA, which he calls RUBÉOLA sine CATARRHO. In this the rash runs its regular course, with little fever or catarrhal affection; affording no certain security against the common or regular disease. He also gives the name, RUBÉOLA NIGRA or BLACK MEASLES, to an unusual appearance of the measles about the seventh or eighth day, when the rash becomes suddenly livid, with a mixture of yellow. It is devoid of inconvenience or danger; and is removed in a week or ten days by the mineral acids and tonics.

RUBÉOLA SPURIA, Roseolæ.

RUBÉOLE, Roseolæ.

RUBÉOLOID, see Roseolæ.

RUB'IA, from *rubere*, 'red.' *R. Tincto'rum seu peregrin'a seu sylvest'ris seu major*, *Erythrodianum*, *Radix rubra*, *Dyer's Madder*, *Madder*, (F.) *Garance*. Family, Rubiaceæ. Sec. Syst. Tetrandria Monogynia. The root of this plant—*Rubia*, (Ph. U. S.)—has a bitterish, somewhat austere, taste, and a slight, not agreeable, smell. They were formerly considered deobstruent, detergent, and diuretic.

RUBIG'INOUS, *Rubigin'o'sus*, from *rubigo*, *rubigine*, 'rust.' (F.) *Rouillé*. Rusty. Having the colour of rust. An epithet given to the sputa in certain cases of pneumonia.

RUBINUS VERUS, Anthrax.

RUBOR, Rubedo.

RUBRICA FABRI'LIS, from *rubere*, 'red.' A red, heavy, earthy matter, formerly employed for making drying plasters; *Hard Ruddle*, *Red Chalk*.

RUBRICA SINOP'ICA. A heavy, compact, reddish earth: formerly used in diarrhoea.

RUBRIN, Hæmatin.

RUBUS AFFINIS, *R. fruticosus*; allied to *rubus*, 'red.'

RUBUS ARC'TICUS, *Shrubby Strawberry*, *Bacca Norland'icæ*. Ord. Rosaceæ. Sec. Syst. Icosandria Polygynia. The fruit is recommended by Linnæus as possessing antiseptic, refrigerant, and antiscorbutic qualities.

RUBUS BATUS, R. cæsius.

RUBUS CÆSIUS, R. batus, Dewberry plant, (F.) Ronce bleue. The fruit resembles the blackberry in appearance and properties. The bark of the root of *R. Trivialis* seu *Canadensis* seu *procumbens* seu *flagellaria*, American Dewberry, Low Blackberry, is astringent.

RUBUS CANADENSIS, R. cæsius.

RUBUS CHAMÆMORUS, Cloudberry Tree, Chamæmorus, C. Norvegica, Chamædrya, Cloudberries, Knot-berries, (F.) Ronce de Montagne. The ripe fruit is sometimes prepared like a jam, and used in fevers, &c., to allay thirst. It has been much extolled as an antiscorbutic.

RUBUS FLAGELLARIS, R. cæsius—r. Framboesianus, R. idæus—r. Frondosus, R. fruticosus.

RUBUS FRUTICOSUS, R. affinis seu paniculatus seu niger seu frondosus seu vulgaris; Common Bramble, which affords Blackberries, (Prov.) Brambleberries, Bumblekites, (F.) Ronce noire. The berries are eaten as a summer fruit, of which they form a poor variety. The bark of the root of the *Rubus villosus*, American Blackberry, High or standing blackberry, Hairy American Bramble, is astringent and has been used in the last stages of dysentery, and in cholera infantum.

RUBUS IDÆUS, R. frambœsæna, Batos; Raspberries, (F.) Framboise, Ronce. The fruit of the best varieties—(Sc.) *Rasps*, (Prov.) *Hindberries*—is extremely agreeable; and is much used. It was, formerly, called *Batton moron*.

RUBUS NIGER, R. fruticosus—r. Occidentalis, R. strigosus—r. paniculatus, R. fruticosus—r. Procumbens, R. cæsius.

RUBUS STRIGOSUS, Wild red raspberry, of the United States. The leaves are often used as a mild astringent, as well as those of *Rubus occidentalis*, Black Raspberry.

RUBUS TRIVIALIS, R. cæsius—r. Villosus, R. fruticosus—r. vulgaris, R. fruticosus.

RUCKING, Eructation.**RUCTAMEN, Eructation.****RUCTATIO, Eructation.****RUCTUATIO, Eructation.****RUCTUOSITAS, R. cæsius.**

RUCTUS, Eructation—r. Acidus, Oxyregmia—r. Nidosus, Cnissoregmia.

RUDBECKIA LACINIATA, Cone flower, Thimble weed; indigenous: Order, Compositæ; is diuretic, and has been given in nephritic affections.

RUDDLE, HARD, Rubrica fabrillis.

RUDIMENTAL, Rudimentalis, (F.) Rudimentaire, from rudimentum, 'the first beginning.' That which is but slightly developed.

RUDOLPHIA FRONDOSA, Butea frondosa.

RUE, Ruta—r. Common, Ruta—r. de Chèvre, Galega officinalis—r. Goat's, Galega officinalis—r. Goat's, Virginia, Galega Virginiana—r. Meadow, Thalictron—r. des Miralles, Asplenium ruta muraria—r. des Prés, Thalictron—r. Sauvage, Ruta.

RUELLES, Wrinkles.**RUFT, Eructation.****RUGA, Wrinkle.****RUGINE, Raspatorium.****RUGISEMENT, Rugitus.**

RUGITUS, Brycheth'mos, from rugire, rugitum, 'to roar.' (F.) Rugissement. A Latin word, which ordinarily expresses the roaring of the lion: and which is used, by some medical writers, to designate a particular species of borborygmus compared to it.

RUGOSE, Wrinkled.**RUGOUS, Wrinkled.****RUM, Tafia, Spiritus Jamaicensis, Molasses**

Spirit, (F.) Rhum. A spirit distilled from sugar, or molasses. See Spirit.

RUMA, Mamma.**RUMEN, Hypogastrium, Ingluvies.**

RUMEX ACETOSELLA, Sheep Sorrel, Field Sorrel; indigenous; has similar virtues.

RUMEX ACETOSUS, R. tuberosus seu intermedium, Lap'athum pratense, Oxalis Acetosæ, Acetosa vulgaris seu pratensis seu nostras, Anax'gris, A. arvensis, Sour dock, Sorrel, Common Sorrel, (Sc.) Docken, (F.) Oseille, Vinette, Aigrette, Surelle. Ord. Polygonacæ. Sex. Syst. Hexandria Trigynia. The leaves are sour, and are used in cookery, and also as a refrigerant.

RUMEX ACUTUS, Sharp-pointed Wild dock, Oxyp'athum, Lap'athum, L. sylvestre seu acutum, Rumex, R. paludosus seu pusillus, Lap'athos, (F.) Patience sauvage. A decoction of the root has long been used in cutaneous affections. It is, sometimes, employed for the cure of the itch.

RUMEX ÆTENSIS, R. scutatus.

RUMEX ALPINUS, Acetosa Alpina. The systematic name of the plant that affords the Monk's Rhubarb, (F.) *Rhabarbe des Moines*. See Rumex patientia.

RUMEX AQUATICUS, Rumex hydrolapathum—r. Britannica, Rumex hydrolapathum—r. Confer'tus, R. patientia—r. Dignus, Oxyria reniformis—r. Glaucus, R. scutatus—r. Hastatus, R. scutatus—r. Hastifolius, R. Scutatus.

RUMEX HYDROLAPATHUM, Water-Dock, (Old Eng.) Brokelack, Hydrolapathum, Rumex aquaticus seu Britannica, Herba Britannica, Lap'athum aquaticum, (F.) Parelle, Patience d'eau. The leaves of this plant are subacid, and said to be laxative. The root has been used as a tonic, astringent, and antiscorbutic. It has also been employed externally in some cutaneous affections.

RUMEX INTERMEDIUS, R. acetosus.

RUMEX OBTUSIFOLIUS, Lap'athum sylvestre, Blunt-leaved Dock, Bitter dock. The root of this, as well as of *R. crispus*, Curled, or Sour, or Yellow, has similar virtues to those of the other species. Both have been used in decoction in the same cases as sarsaparilla.

RUMEX OXYOTUS, R. scutatus—r. Paludosus, R. acutus.

RUMEX PATIENTIA, R. confer'tus, Lap'athum hortense, Rhubarbarum monachorum, Hippolap'athum, Patientia, Garden Patience, Monk's Rhubarb, (F.) Patience, Rhubarbe sauvage, Parelle. This root, which is supposed to possess the virtues of rhubarb, but in an inferior degree, is, according to Linnæus, obtained from the *Rumex patientia*; according to Murray, from the *Rumex alpinus*.

RUMEX PUSILLUS, R. acutus.

RUMEX SANGUINEUS. The root of the bloody dock or bloody veined Dock, has an austere and astringent taste; and has been employed in dysentery. It is called, also, *Lap'athum sanguineum*, (F.) *Oseille on Patience Rouge, Sangdragon*.

RUMEX SCUTATUS, R. glaucus seu hastifolius seu Ætensis seu oxyotus seu hastatus, Acetosæ rotundifolia seu Romana seu Rotundifolia hortensis seu scutata, Lap'athum scutum seu rotundifolia, Roman or Garden Sorrel, Green Sauce, French Sorrel, (F.) Oseille boucher, Oseille ronde, Petite O. Virtues the same as those of the common sorrel.

RUMEX TUBEROSUS, R. acetosus.

RUMINATION, Rumination, from ruma or rumen, 'the gullet.' Mercy's mus. A function peculiar to ruminating animals, by which they chew a second time the food they have swallowed. An analogous phenomenon is sometimes seen in man. One who ruminates is in French *mérycole*.

RUMMET, Pityriasis.
RUMP, *Croupon*—*r.* Bone, Coccyx.
RUNCHES, *Sinapis arvensis*.
RUNCINA, *Raspatorium*.
RUNCINULA, *Raspatorium*.
RUNKILL, Wrinkle.
RUNKLE, Wrinkle.
RUNNET, Rennet.
RUNNING, *Cursus*, (*F.*) *Course*. A succession of leaps, executed with rapidity, to transport the body quickly from one place to another.
RUNNING, Rennet.
RUNSH, *Sinapis arvensis*.
RUNUNCULA, *Raspatorium*.
RUPEs, Lips of the Vulva. See *Lip*.
RUPIA, *Rhyphia*, *Ulcus atonicum*, *Ecchly'sis Rhy'pia*, *Rhypha'ria*, from *'purus*, 'filth.' An eruption of large flattish blebs, which contain a fluid—at first serous, afterwards puriform, and often bloody, which rapidly concretes into crusts, at the base of which are ulcers of variable depths. It requires the use of tonics internally, and of stimulating ointments externally.
RUPIA ESCHAROTICA, *Pem'phigus infantilis* seu *gangræno'sus*; known in Ireland under the names—*white blisters*, *eating hiye* and *burnt holes*. An affection, which bears a close similarity to pemphigus, particularly in the absence of a thick rugous crust, whilst in its chief feature, that of ulceration, it evidently belongs to *Rupia*.
RUPTIBILITAS, *Fragilitas*.
RUPTIO, *Rhexis*.
RUPTORIUM, *Cauterium*—*r.* *Commune*, *Potassa fusa*.
RUPTURA, *Erexis*, *Hernia*, *Laceration*, *Rhexis*—*r.* *Cornææ*, *Ceratorrhexis*—*r.* *Herniosa*, *Hernia*—*r.* *Ovarii*, *Oariorrhexis*—*r.* *Uteri*, *Uterus*, rupture of the—*r.* *Vaginæ*, *Colporrhexis*.
RUPTURE, *Hernia*, *Rhexis*—*r.* *du Cœur*, *Cardiorrhexis*—*r.* of the Groin, *Bubonocœle*—*r.* *de la Matrice*, *Uterus*, rupture of the—*r.* *Wort*, *Herniaria glabra*.
RUPTURED, see *Hernial*.
RUSCUS, *R. aculeatus*, *Bruscus*, *Oxymyr'rhinê*, *Oxymyr'sinê*, *Myrtacantha*, *Catan'gelos*, *Myacantha*, *Scopia re'gia*, *Wild Myrtle*, *Centromyr'inê*, *Butcher's Broom*, (*Prov.*) *Knee-holly*, (*F.*) *Petit houx*, *Fragon*. *Ord.* *Smilacææ*. *Sex. Syst.* *Dioecia Monadelphia*. The root has been recommended as an aperient and diuretic.
RUSCUS ACULEATUS, *Ruscus*.

RUSCUS HYPGLOS'SUM, *Ucula'ria*, *Hypoglossum*, *Hippoglossum*, *Bistlin'gua*, *Bonija'cia*, *Laurus Alexandri'na angustifolia*, (*F.*) *Laurier Alexandrin*, *Langue de cheval*. Formerly used against relaxed uvula.

RUSH, *SWEET*, *Acorus calamus*, *Juncus odoratus*.

RUSMA, see *Orpiment*.

RUSPIN'S STYPTIC, see *Styptic*, *Ruspini's*.

RUSPIN'S TINCTURE FOR THE TEETH. An empirical preparation, composed of *Florentine orris* \mathfrak{z} viij, cloves \mathfrak{z} j, rectified spirit Oij, ambergris \mathfrak{z} j. *RUT*, Heat.

RUTA, *'purn*, *R. graveolens* seu *horten'sis* seu *altera*, *Peg'anum*, *Besa'sa*, *Common Rue*, (*F.*) *Rue*, *Rhue*, *Rue sauvage*, *Herbe de grace*; *Ord.* *Rutacææ*. *Sex. Syst.* *Decandria Monogynia*. The plant has a strong, ungrateful odour, and bitter pungent taste. It is acrid, so as to blister the skin. Its properties are reputed to be tonic, stimulant, antispasmodic, and emmenagogue. Dose, gr. xv to \mathfrak{z} ij. The leaves are officinal—*Ruta*. (*Ph. U. S.*)

Oleum Ruta, distilled Oil of Rue, is antispasmodic; and, externally, rubefacient. Dose, gtt. ij to vj.

RUTA ALTERA, *Ruta*—*r.* *Capraria*, *Galega officinalis*—*r.* *Graveolens*, *Ruta*—*r.* *Hortensis*, *Ruta*—*r.* *Muraria*, *Asplenium ruta muraria*.

RUTABULUM, *Penis*.

RUTA'CEUM, from *Ruta*, 'rue.' Vinegar of rue.

RUTIDOMA, see *Rutidosis*.

RUTIDO'SIS, *Rytido'sis*, *Rhytido'sis*, from *ρυτιδω*, 'I wrinkle.' The act of wrinkling; as *Rutido'ma* means a shrivelled or wrinkled body. A destruction or atrophy of the eye. Corrugation and subsidence of the cornea.

RUYSCHIA'NA TUNICA, (*F.*) *Lame ou Membrane Ruyschienne*. According to Ruysch, and the greater part of his disciples, the choroid coat of the eye is formed of two laminae. His son gave to the inner the name *Tunica Ruyschia'na* or *Membra'na Ruyschia'na* seu *Ruysch'ii*. *Tu'nica Cellulo'sa Ruyschia'na* is the areolar substance under the peritoneal covering of the stomach. It is not numbered among the coats of that organ.

RYE, *Secale*—*r.* *Horned*, *Ergot*—*r.* *Robbin's*, *Polytrichum juniperinum*—*r.* *Spurred*, *Ergot*.

RYTIDOSIS, *Rutidosis*.

RYTIS, *Wrinkle*.

S.

S. A. &c., see *Abbreviation*.
SAAG, *Urine*.
SABADILLA, *Veratrum sabadilla*, see *Veratrina*.
SABADILLIN, see *Veratrina*.
SABATILLA, *Veratrum sabadilla*.
SABBATIA, *Chironia angularis*—*s.* *Angularis*, *Chironia angularis*.
SABINA, *Juniperus sabina*.
SABINE, *Juniperus sabina*.
SABLÊ, *Maculo'sus*, 'sanded,' from *sabulum*, 'sand.' Having the appearance as if strewed with sand. The terms *sablê* and *piquetê* are sometimes applied to the brain, when a slice of it exhibits red points as if it had been strewed with red sand. Such is the case in cerebral hyperæmia.

SABOT DE VÉNUS JAUNE, *Cypripedium luteum*.

SABUCUS, *Sambucus*.

SAB'ULOUS, *Sabulo'sus*, *Areno'sus*, *Psammo'des*, from *sabulum*, 'sand.' The sandy deposit often seen in the urine.

SABULUM CONARII, see *Pineal gland*.

SABURA, *Saburra*.

SABUR'RA, *Sabu'ra*, *Sordes*. This word signifies, by derivation, 'coarse sand, or ballast.' In its common acceptation, it means *foulness of the stomach*, as it is sometimes called;—vitiated matters supposed to be retained and accumulated in the stomach—the result of imperfect digestion—and which have been sometimes considered as a morbid product of the mucous secretion of that organ, or of the biliary secretion; at others, as a residuum from alimentary substances badly di-

gested. The *Saburral state*, *Colluvies gastrica*, *Cruditates ventriculi*, *Gastrismus*, (F.) *Embaras gastrique*, *État saburral*, is an accumulation of *saburræ* in the stomach, (F.) *Saburres gastriques*, which the Humourists considered to be the cause of a number of diseases.

SABURRA VERMINOSA, Helminthiasis.

SABURRAL, *Saburralia*. Relating or appertaining to *saburræ*,—as the '*saburral state*.'

SABURRATIO, Arenatio.

SABURRES GASTRIQUES, see *Saburra*.

SAC, see Sack—*s. Herniaire*, Hernial sac—*s. of the Omentum*, see Epiploon, gastro-hepatic.

SACCADE, see Respiration, jerking.

SACCATUS, Encysted.

SACCHAR, Saccharum.

SACCHARHYDROLÉ, Syrup.

SACCHARI FÆX, Melasses.

SACCHARINA, *Saccharolés*.

SACCHARO-ALCOOLÉ, Ratafia.

SACCHAROLÉ LIQUIDE, Mellitum, Syrup—*s. Mou*, Conserva, Electuarius—*s. Solide*, see Pastel, and Tabella.

SACCHAROLÉS (F.), *Sacchari'na*, from *saccharum*, 'sugar.' A name given by MM. Henry and Guibourt to medicines which have sugar or honey for excipient, or for predominant principle. M. Béal so calls an exact mixture of pulverized sugar with other substances also in powder, but in less quantity.

SACCHAROMETER, (F.) *Saccharomètre*, *Pèse-sirop*, from *saccharum*, 'sugar,' and μέτρον, 'measure.' A hydrometer adapted for determining the density of syrups. The marginal figure is graduated to Baumé's scale. See Areometer.

SACCHAROMYCES, *Torula cerevisiæ*.

SACCHARORRHŒA, Glycyrrhœa—*s. Cutanea*, *Sudor dulcis*.

SACCHARORRHŒA LACTEA, *Galactorrhœa saccharata*. A secretion of unusually sweet milk.

SACCHARORRHŒA PULMONALIS, *Sputum dulce seu saccharatum*. The copious expectoration of unusually sweet matter from the lungs.

SACCHARORRHŒA URINOSA, *Diabetes mellitus*.

SACCHARUM, *Sacchar*, *сахар*, from Arabic, or rather Sanscrit, *Sakar*, 'white earth'; *Suchar*, *Succhar*, *Sutter*, *Zuchar*, *Zu'caro*, *As'sakur*, *Tigala*, *Zaccharum*, *Zozar*, *Zu'charum*, *Zu'carum*, *Zuchra*, *Mel arundinaceum*, *Sugar*, (F.) *Sucre*. The produce of the juice of *Saccharum officinarum* seu *officinale* of Linnaeus, *Calamus Indicus* seu *Saccharinus*, *Sugar Cane*, (F.) *Canne à Sucre*. *Cannamelle*; *Arun'do saccharifera* of Sloane; *Ord. Graminæ*; *Sex. Syst. Triandria Digynia*, and also of the beet, maple, &c.

True Sugars are those with which alcoholic fermentation may be effected, by the addition of yeast and water. They are:—

Species.

Varieties.

- | | | |
|--------|---|--|
| 1..... | { | Hard sugar of the cane, maple, beet, chestnut, &c. |
| 2..... | | Liquid sugar, sugar of malt, of the sweet potato, melasses, honey, &c. |
| 3..... | | Hard sugar of the grape, ripe fruits, starch, diabetes, &c. |
| 4..... | | Hard sugar of the mushroom. |

Imperfect sugars are those in which alcoholic fermentation cannot be effected by means of yeast and water.

Species.

Varieties.

- | | |
|--------|-------------------------|
| 1..... | Manna. |
| 2..... | Sugar of Milk. |
| 3..... | Sugar of jelly or glue. |
| 4..... | Liquorice. |
| 5..... | Pieromel. |

Sugar is prepared from the expressed juice of the sugar-cane, boiled with the addition of quicklime or common vegetable alkali. It is used, in pharmacy, for the preparation of syrups, conserves, lozenges, &c. It is nutritious, and is employed as an aliment, and as a eutrophic demulcent and antiseptic. Dissolved in small quantities in water, as in tea, it is apt to disagree with dyspeptics; an evil which does not always occur when the same substance is taken more largely.

Liver or Hepatic Sugar, (F.) *Sucre du Foie*, has been recently found to be a normal secretion of the liver. It is more readily assimilated than other varieties of sugar. See Glucose.

SACCHARUM ACER'NUM, *S. Canadensē*, *Maple Sugar*, is often used as a demulcent.

SACCHARUM ALBUM, *S. purificatum*, *Saccharum* (Ph. U. S.), *Sal Indus*, *White Sugar*, *Refined Sugar* is the *Loaf Sugar*. *Double-refined Sugar* has been called *Tabarzet*.

SACCHARUM ALUMINATUM. Equal parts of white sugar and alum, powdered. Used as a styptic.

SACCHARUM CANDIDUM, *S. candum seu cantum seu canthum*, *Al'phenic*, *Nabot*, *Sugar candy*, *Sugar candian*, (F.) *Sucre Candi*, is the crystallized juice of the cane.

SACCHARUM CANDUM, *S. Candidum*—*s. Cantum*, *S. Candidum*—*s. Canthum*, *S. Candidum*—*s. Commune*, *Saccharum non-purificatum*.

SACCHARUM HORDEA'TUM, *Barley Sugar*, (Prov.) *Sugar barley*, *Penides*, *Penid'ium*, *Saccharum penid'ium*, is a variety of sugar candy.

SACCHARUM LACTIS, *Sugar of milk*.

SACCHARUM NON-PURIFICA'TUM, *Muscovado*, *raw*, *brown* or *moist sugar*, is the *S. communē*, *S. rubrum*, *Mel canna*, or *coarse sugar*.

SACCHARUM OFFICINARUM, see *Saccharum*—*s. Penid'ium*, *Hordeatum*—*s. Rosaceum*, *Confectio rosæ gallicæ*—*s. Saturni*, *Plumbi superacetas*.

SACCHARURE, from *saccharum*, 'sugar.' A medicine resulting from the intimate union of sugar with the medicinal principles of alcohol or ethereal tinctures.—Béal.

SACCULATED, Encysted.

SACCULATUS, Encysted.

SACCULI ADIPOSII, *Fatty vesicles*—*s. Chalicophori*, see *Lacunæ of Bone*—*s. Laryngei*, *Ventricles of the larynx*.

SACCULI MEDICINAL'ES. Bags containing medicinal substances.

SACCULI MEDULLARES, see *Medullary membrane*.

SACCULUS, diminutive of *saccus*, 'a little sac.' This name has been given to two sacs formed by the membranous vestibule, which open into each other, and contain the *otolithes* and *otoconies*: one of these is termed the *sacculus vestibuli*, the other *sacculus*. See *Semicircular canals*.

SACCULUS, Marsupium—*s. Cephalicus*, *Cucupha*—*s. Chylifer seu roifer*, *Receptaculum chyli*—*s. Cordis*, *Pericardium*—*s. Ellipticus*, see *Semicircular canals*—*s. Lactiferus*, see *Lactiferous vessels*.

SACCULUS LARYN'GIS. A cæcal pouch, of variable size, formed by a mucous membrane in the ventricles of the larynx.—Hilton.



Saccharometer.

SACCULUS MEDICINALIS, Pulvinar medicatum —s. Sphericus, see Semicircular canals—s. Vesical, see Bladder, sacculated —s. Vestibuli, Semicircular canals, see Sacculus.

SACCUS, Sac —s. Coli, Cæcum —s. Cordis, Pericardium—s. Herniosus, Hernial sac—s. Epiploicus, Epiploon —s. Intestini crassi, Cæcum —s. Lacrymalis, Lacrymal sac —s. Lacrumalis, Lacrymal sac —s. Lacteus, Receptaculum chyli —s. Mucosus, see Bursæ mucosæ —s. Peritonæi retro-ventricularis, see Epiploon, gastro-hepatic —s. Pleuræ, Pleura—s. Venarum cavarum, Sinus dexter cordis —s. Venarum pulmonalium, Sinus pulmonalis.

SACER, Sacral —s. Ignis, Herpes zoster —s. Morbus, Epilepsy —s. Musculus, Transversalis dorsii.

SACERDOTIS PENIS, Arum maculatum.

SACHET (F.), *Sacculus*, 'a little sac.' A small bag containing perfumes to be placed among clothes to communicate to them an agreeable odour. The following is a formula for the powder they contain. *Powdered cloves and powdered cassia*, each one ounce; *powdered orris root*, an ounce and a half; *powdered yellow sandal wood*, one drachm; *otto of roses*, twenty-four drops; *oil of lavender*, *oil of bergamot*, each one drachm; *musk*, six grains. A scent-bag.

SACK, (F.) *Sec*, 'dry.' A dry wine used by our ancestors; *Vin sec*. Probably Sherry.

SACK, *Saccus*, *Seclusorium*; *Sax. fac*. A bag or pouch.

SACK, *HER'NIAL*, *Saccus herniosus*, (F.) *Sac herniaire*. An envelope or pouch, formed by the serous membrane of the cavity, whence a viscus has escaped. We distinguish, in a hernial sac, its *orifice* or the opening by which it communicates with the great cavity lined by the serous membrane: its *cervix* or neck, (F.) *Collet*, the narrow portion on the outside of the orifice, where strangulation sometimes occurs: the *body*, (F.) *Corps*, which is the round prominent part; and the *fundus* or bottom,—the part opposite to the orifice.

SACK POSSET; from *sack*, 'the wine,' and *posset*. A posset made of sack, milk and other ingredients.

SACK, SEROUS. The shut cavity formed by any serous membrane, as by the pleura or peritoneum.

SACK WHEY, Wine Whey.

SACRAD, see Sacral aspect.

SACRA FISTULA, Vertebral column.

SACRAL, *Sacer*, (F.) *Sacré*. That which belongs or relates to the sacrum. A name given to several parts.

SACRAL ARTERIES are distinguished into, 1. The *anterior or middle sacral*, *Median Artery of the Sacrum* of Chaussier, which arises from the posterior part of the abdominal aorta, in front of the 4th lumbar vertebra. It descends, in a serpentine manner, upon the sacro-vertebral articulation, and on the anterior surface of the sacrum. It furnishes many lateral branches, which unite with the lateral sacral arteries, and enter the anterior sacral foramina, to be distributed to the membranes of the marrow, &c. 2. The *lateral sacral arteries*, which are two in number; one on each side. They arise from the hypogastric, from the ilio-lumbar, or from the gluteal; and descend before the anterior sacral foramina, anastomosing in an arched manner, towards the coccyx, with the middle sacral. They give off *external and posterior branches*, which enter the sacral foramina; and *internal branches*, which are distributed to the pyramidalis, sacrum, &c.

SACRAL ASPECT. An aspect towards the region where the sacrum is situated.—Barclay. *Sacrad*

is used by the same writer adverbially to signify 'towards the sacral aspect.'

SACRAL EXTREMITIES. The lower limbs.

SACRAL GROOVES, (F.) *Gouttières sacrées*. Two longitudinal excavations, slightly marked, at the posterior surface of the sacrum; which appear to terminate the vertebral grooves or gutters.

SACRAL NERVES, (F.) *Nerfs sacrés*, are generally six in number. They arise from the termination of the spinal marrow. The first, united with the great cord which descends from the lumbar plexus, concurs with the three others in the formation of the sacral or sciatic plexus. The third and fourth, with branches of the great sympathetic, form the hypogastric plexus. The fifth and sixth (*coccygeal nerve*) are distributed to the ischio-coccygeus muscle, and to the sphincter and levator ani.

SACRAL PLEXUS, *Sciat'ic plexus*,—the *sacral portion of the crural plexus* of Chaussier, constituted as just mentioned, is situate in front of the pyramidalis. It gives off, backwards, two branches, called the *pudic*, (F.) *honteux*, and the *lesser sciatic*. It terminates in the great sciatic nerve.

SACRÉ, Sacral.

SACRO, in composition, relating or belonging to the sacrum.

SACROCOCCYGE'AL, *Sacro-coccygeus*. Relating to the sacrum and coccyx.

SACROCOCCYGE'AL ARTICULA'TION is that of the lower extremity of the sacrum with the superior *facet* of the coccyx. It is closed by two ligaments;—an *anterior* and a *posterior sacro-coccygeal*.

SACROCASTALIS, *Sacro-lumbalis*—s. *Fémoral*, *Glutæus major*.

SACROCOXAL'GIA, from *sacro*, *coxa*, 'the hip,' and *algos*, 'pain,' a hybrid term. Some authors have called thus a rheumatic affection of the sacro-iliac symphysis.

SACROÏL'IAC, *Sacro-ili'acus*. That which relates to the sacrum and ilium.

SACROÏLIAO ARTICULA'TION, *SYMPHYSIS* or *SYNCHONDRO'SIS*, *Ilio-sacral articulation*, *symphysis* or *synchondrosis*, is the union of two surfaces;—two inclined planes, broader above than below; one formed by the sacrum, the other by the ilium. Both surfaces are incrustated by an articular cartilage. The bands which strengthen them are the two *sacro-sciatic ligaments*, the *sacro-spinal*, a *sacro-iliac* ligament, and some irregular fibres. The *sacro-iliac* ligament is a very thick assemblage of short, dense, irregular fibres, occupying the space which the sacrum and ilium leave between them, behind their articular surface. Some authors have termed some irregular fibres, situate in front of the sacro-iliac articulation, *anterior sacro-iliac ligaments*, in opposition to the preceding fasciæ, which they call *posterior sacro-iliac*.

SACROÏSCHIATICUS, *Sacro-sciatic*.

SACROLOMBAIRE, *Sacro-lumbalis*.

SACROLUM'BAR, *Sacro-lumba'lis*, *Sacro-lombaire*. That which belongs to the sacrum and loins.

SACROLUMBAR, *Sacro-lumba'lis*, *Sacro-lumba'ris Muscle*, *Sacro-costa'lis*, *Dorso-trache'lian portion* of the *Sacro-spinal* of Chaussier, (F.) *Lombocosto-trachélien*, *Sacro-lombaire*, is seated at the posterior part of the trunk; is thick and triangular below; flat and thin in the remainder, and terminates in a point at its upper extremity. It is attached to the posterior surface of the sacrum; to the corresponding portion of the crista ili; to the extremities of the transverse processes of the lumbar vertebræ; to the angles of the eleven inferior ribs; to the tuberosity of the first, and the

posterior tubercle of the transverse processes of the last five cervical vertebrae. The sacro-lumbalis straightens the vertebral column when bent forwards, and maintains it straight. By means of its insertion into the ribs, it can elevate or depress these bones, according as it contracts from above to below, or from below to above. It acts, consequently, sometimes as an inspiratory, at others as an expiratory muscle.

From six or eight of the lower ribs arises an equal number of fleshy portions, which terminate at the inner side of this muscle, and are called *Musculi Accessorii seu Additamentum ad Sacro-lumbalem*, (F.) *Accessoires du Sacro-lombaire*.

From the upper part of the sacro-lumbalis, a fleshy slip, called *Cervicalis descendens*, *Cervicalis descendens Dienerbroeckii*, *Transversalis collateralis colli*, *Accessorius ad Sacro-lumbalem*, (F.) *Transversaire grêle*, runs up, to be fixed to the transverse processes of the fourth, fifth, and sixth cervical vertebrae, by three distinct tendons. These last parts, though separated by some anatomists, belong to the sacro-lumbalis.

SACROSCIATIC, *Sacro-ischiat'icus*. That which belongs to the sacrum and ischium.

SACROSCIATIC LIGAMENTS are two ligaments, which assist in securing the sacro-iliac articulation. The *great or posterior sacro-sciatic ligament* extends from the crista ilii and the sides of the sacrum and coccyx to the tuberosity of the ischium. The *less or anterior sacro-sciatic ligament* arises from the same point, and terminates in the spine of the ischium.

SACROSPINAL, *Dorso-trachelian* portion of the sacro-lumbalis.

SACROSPINALIS. That which relates to the sacrum and spine. Under this name, Chaussier describes the sacro-lumbalis, longissimus dorsi, transversalis, transverso-spinalis, and intertransversalis muscles. He divides it into three portions.

SACROTROCHANTÉRIEN, *Pyramidalis*.

SACROVERTÉBRAL, *Sacro-vertebralis*. That which relates to the sacrum and the vertebrae.

SACROVERTÉBRAL ARTICULATION is the junction of the last vertebrae of the loins with the sacrum. The angle which these two bones form at the anterior part has been called the *Promontory of the Sacrum*, or the *Sacro-vertebral Angle*.

SACROVERTÉBRAL LIGAMENT is a strong ligament, which passes from the anterior part of each of the transverse processes of the last lumbar vertebra to be attached to the base of the sacrum. Its use is to maintain the articulation of those two bones *in situ*.

SACRUM, *Os sacrum seu basilarē seu Alagae seu latum seu clun'ium*, *Lus*, *Luz*, *Vertebra magna*, *Albagiazi*, *Hi'eron ost'con*, *Cluniv*, *Subvertebra*, *Subvertebrum*, *Vetula*; from *sacer*, 'sacred'; because it contributes to protect the genital organs, which were considered sacred, or because it was offered in sacrifice. The bone, which forms the posterior part of the pelvis, and is a continuation of the vertebral column. The sacrum is symmetrical and triangular, situate at the posterior part of the pelvis, and concave anteriorly. It has,—1. A *pelvic or anterior surface*, which corresponds to the pelvic cavity, and has four transverse furrows, that point out the parts where the primitive portions of the bone were separated from each other. On each side there are four foramina, called *anterior sacral*; these are oblique, and decrease in size, from above downwards. Through them pass the anterior branches of the sacral nerves. 2. A *spinal or posterior surface*, which is convex, rough, and covered by muscles. Upon the median line are four horizontal, tubercular eminences, forming a

continuation of the spinous processes of the vertebrae; and below, a triangular notch, where the *sacral canal* terminates. On each side of the surface are four foramina, called *posterior sacral*, which are smaller than the anterior, and give passage to the posterior branches of the sacral nerves. 3. The *vertebral surface or base*. Its greatest extent is transversely. It is articulated with the last lumbar vertebra. On it is seen the orifice of the *sacral canal*, of a triangular shape, which decreases in width from above downward; terminates the vertebral canal; is lined by a prolongation of the cerebral membranes; contains the fasciculi of the sacral nerves; communicates, externally, by the sacral foramina; and terminates at the triangular notch on the posterior surface of the bone. At the sides of the upper orifice of the sacral canal are two articular processes, which receive the articular processes of the last lumbar vertebra. The projection, formed by the union of the base of the sacrum with the last lumbar vertebra, is called the *Promontory*, *Promontorium Ossis Sacri*, *Great Angle of the Sacrum* or *Sacro-vertebral Angle*. 4. A *coccygeal face or apex*, (F.) *Face coccygienne ou Sommet*, which is very narrow, and is articulated with the coccyx. 5. Two *lateral margins*, which have, above, a broad, oblique surface, articulated with the ilium.

The sacrum is developed by 30 to 36 points of ossification.

SADDLE, TURKISH, *Sella Turcica*.

SÆPES, *Sepes*.

SAFFLOW, *Carthamus tinctorius*.

SAFFLOWER, *Carthamus tinctorius*.

SAFFRON, *Crocus*—s. *Bastard*, *Carthamus tinctorius*—s. *Dyers'*, *Carthamus tinctorius*.

SAFFRON FLOWER, *Carthamus tinctorius*.

SAFFRON, MEADOW, *Colchicum autumnale*.

SAFRAN BÂTARD, *Carthamus tinctorius*, *Colchicum autumnale*—s. *des Indes*, *Curcuma longa*—s. *de Mars apéritif*, *Ferri subcarbonas*—s. *de Mars astrigent*, *Colcothar*—s. *des Prés*, *Colchicum autumnale*—s. *Racine de*, *Curcuma longa*.

SAFRANUM, *Carthamus tinctorius*.

SAFRE, *Smalt*.

SAGA, *Sago*.

SAGAPE'NUM, *Sag'apin*, *Serapi'non*, (F.) *Gomme staphique*. A gum-resin, supposed to be obtained from the *Ferula Per'sica*. *Ord. Umbelliferae*. *Sec. Syst. Pentandria Digynia*. Its odour is fetid and alliaceous; taste pungent, bitterish, nauseous. It is in small, agglutinated masses of a yellow colour; tenacious; breaking with a horny fracture. It is a reputed antispasmodic and emmenagogue. Dose, gr. x to ʒss.

SAGAPIN, *Sagapenum*.

SAGE, *Salvia*—s. *Indian*, *Eupatorium perfoliatum*—s. *Jerusalem*, *Pulmonaria*.

SAGE TREE, *Lantana*.

SAGE, WILD, *Salvia Africana*, *S. Lyrata*.

SAGE-FEMME, *Midwife*.

SAGENA, *Epiploon*.

SAGESSE DES CHIRURGIENS, *Sisymbrium Sophia*.

SAGIMEN VITRI, *Soda*, subcarbonate of.

SAGIT'TAL, *Sagitta'lis*, *Sagitta'tus*, from *sagitta*, 'an arrow.'

SAGITTAL FURROW, *Sagittal Groove*, (F.) *Gouttière sagittale*, is a channel on the median line of the inner surface of the skull-cap. It is excavated in the os frontis, in the two parietal and occipital bones, and extends from the crista galli to the internal occipital protuberance. It lodges the longitudinal sinus, which has been called the *sagittal sinus*.

SAGITTAL SUTURE, *Obelæ'a sutur'a seu Raphē*, *Sutura Virga'ta seu jugu'lis seu Rabdo'des*,

Rhabdoides, *Rhabdoides*, *Rhabdoidens* seu *Biparietalis* seu *Interparietalis*. The suture which unites the two parietal bones, and which extends, from before to behind, on the median line, from the coronal suture to the lambdoidal. It is so called, from *sagitta*, 'an arrow,' because it seems to meet the coronal suture as an arrow meets the string of a bow.

SAGITTARIA VARIABILIS. From *sagitta*, 'an arrow,' owing to the prevalent form of the leaves. *Arrowhead*, *Arrowleaf*, *Arrow weed*, *Wapata* of Oregon; *Nat. Ord.* Alismaceæ. Esculent, yielding fecula like arrow-root. The roots are cooling and sub-astringent, and are at times applied to the feet in cases of jaws and dropsical swellings of the legs. The leaves are applied to the breasts to dispel milk.

SAGITTARIUM ALEXIPHARMACUM, *Malacæ radix*, *Canna Indica*, *Arundo Indica*. A root cultivated in the West Indies, and supposed to be a remedy for wounds by poisoned arrows.

SAGITTATUS, Sagittal.

SAGO, *Sagua*, *Saga*, *Sagu*, *Segu*, *Amylum palmaceum*, (F.) *Sagou*, *Sague*. A fecula, obtained from the pith of the *Sagua rumphii*, *Cycas circinalis*, *C. revoluta*, &c., growing in the Moluccas, Philippine Isles, &c., and which is brought to us in small grains. The same substance is, also, obtained from the West Indies, but it is inferior to that from the East. By boiling in water or milk, sago becomes soft and transparent, and forms an agreeable and nutritious food in febrile, calculous, and other disorders. It is made palatable by sugar, lemon-juice, or wine, where the last is not contra-indicated.

To make sago into a proper mucilage for the sick, a tablespoonful of it should be macerated in a pint of water for two hours, and then boiled for fifteen minutes, stirring assiduously. Care must be taken, that the grains are perfectly dissolved by long boiling and constant stirring.

Sago posset is made by putting *sago*, $\frac{3}{4}$ ij, into water, Oij; boiling until the mucilage is formed: then rubbing *sugar*, $\frac{3}{4}$ ss, on the rind of a lemon, and putting it with *tincture of ginger*, $\frac{1}{2}$ ij, into *cherry*, Oss; adding this mixture to the mucilage, and boiling for five minutes.—It is a good restorative.

SAGO, FRENCH, see *Solanum tuberosum*, and *Arrow-root*, common.—s. Portland Island, see *Arum maculatum*.

SAGOU, Sago.

SAGU, Sago.

SAGUE, Sago—s. *Petit*, see *Solanum tuberosum*.

SAGUERUS GOMUTO, see *Toddy*—s. *Rumphii*, see *Toddy*—s. *Saccharifer*, see *Toddy*.

SAGUS, Sago—s. *Rumphii*, see *Sago*.

SAHARA, Insomnia.

SAIDSCHÜTZ WATER, Seidschütz water.

SAIGNÉE, Blood-letting—s. *Spoliative*, *Spoliativa phlebotomia*.

SAIGNEMENT DU NEZ, Epistaxis.

SAILORS' DISEASES, &c., OF. Sailors, by nature of their occupation, are liable to many diseases. Several of these may, however, be ward off by proper attention to cleanliness and ventilation.

SAIN BOIS, *Daphne gnidium*—s. *Doux*, *Adeps præparatus*.

SAINT AMAND, MINERAL WATERS OF. Sulphuretted waters, in the department of Du Nord, France. They contain carbonic and sulphohydric acids. The *Boue* or deposit contains sulphohydric acid, and salts of iron and magnesia. Temperature, 64° to 82° Fahr.

SAINT ANTHONY'S FIRE, Erysipelas.

SAINT AUGUSTINE, CLIMATE OF. This

sea-port of Florida is a frequent retreat during the winter season for the valetudinarian from the more northern parts of the United States. The climate is certainly comparatively mild and equable; but it is liable to frequent north-east storms. There are only a few days, however, in winter, in which the invalid cannot take exercise in the open air. Tampa Bay, and Pensacola, in Florida, and Mobile, in Alabama, which are sometimes visited by the consumptive, are less liable to the storms in question.

SAINT CATHERINE'S SPRINGS, see Canada, Mineral waters of.

SAINT CHARLES'S ROOT, *Carlo Sancto* (Radix).

SAINT CROIX, Santa Cruz.

SAINT JAMES'S WORT, *Senecio Jacobæa*, *Thlaspi bursa*.

SAINT JOHN'S DANCE, see *Mania*, dancing.—*St. John's Wort*, *Hypericum perforatum*.

SAINT KITTS, see *West Indies*.

SAINT LEONARD'S, CLIMATE OF. This place, which is situate about a mile to the westward of Hastings, in Sussex, England, is less sheltered than it, and is therefore a less favourable retreat, during the winter months, for the phthisical valetudinarian.

SAINT MYON, MINERAL WATERS OF. Some cold, acidulous waters at Saint Myon, near Riom, in the department of Puy-de-Dôme. They contain free carbonic acid, subcarbonate of soda, chloride of sodium, and carbonate and sulphate of lime.

SAINT NECTAIRE, MINERAL WATERS OF. Springs in the department of Puy-de-Dôme, France. They contain carbonic acid, bicarbonate of soda, and chloride of sodium. Temperature, 75° to 103° Fahr.

SAINT PARDAUX, MINERAL WATERS OF. Chalybeate springs in the department of Allier, France. They contain carbonic acid and iron.

SAINT SAUVEUR, MINERAL WATERS OF. Sulphurous waters in the department of Hautes-Pyrénées, France. Temperature, 76° to 93° Fahr.

SAINT VINCENT'S, see *West Indies*.

SAINT VITUS'S DANCE, Chorea, see *Mania*, dancing—s. *Vitus's Dance of the Voice*, *Balbuties*.

SAIR, Ulcer.

SAIRE, Essera.

SAL, Salt—s. *Absinthii*, *Potassæ subcarbonas*, see *Potash*—s. *Acetosellæ*, *Potassæ oxalate*—s. *Acetosus ammoniacalis*, *Liquor ammoniæ acetatis*—s. *Acidum benzoës*, *Benjamin*—s. *Aëratum*, see *Potassæ carbonas*—s. *Albus*, *Soda*, *muriate of*—s. *Alembroth*, *Alembroth salt*—s. *Alkali*, *Soda*, *subcarbonate of*—s. *Alkalinus fixus fossilis*, *Soda*—s. *Amarum*, *Magnesiæ sulphas*—s. *Ammoniac*, *volatile*, *Ammoniac carbonas*—s. *Ammoniacum*, *Ammoniac murias*—s. *Ammoniacum martiale*, *Ferrum ammoniatum*—s. *Ammoniacum secretum* *Glauberi*, *Ammoniac sulphas*—s. *Ammoniacum tartareum*, *Ammoniac tartaras*—s. *Ammoniacum vegetabile*, *Liquor ammoniæ acetatis*—s. *Ammoniacus nitrosus*, *Ammoniac nitras*—s. *Ammoniac*, *Ammoniac murias*—s. *Anglicanum*, *Magnesiæ sulphas*—s. *Antiepilepticus of Weissman*, *Cuprum ammoniatum*—s. *Antimonii*, *Antimonium tartarizatum*—s. *Ammoniacum*, *Ammoniac murias*—s. *Armoniacum*, *Ammoniac murias*—s. *Artis*, *Alembroth salt*—s. *Asphaltitis*, *Bit noben*—s. *Catharticus amarus*, *Magnesiæ sulphas*—s. *Catharticus Glauberi*, *Soda*, *sulphate of*—s. *Communis*, *Soda*, *muriate of*—s. *Cornu cervi volatile*, *Ammoniac carbonas*—s. *Culinaris*, *Soda*, *muriate of*—s. *Digestivus Sylvii*, *Potassæ murias*—s. *Diureticus*, *Potassæ acetas*—s. *de Duobus*, *Potassæ sulphas*—s. *Ebsdamense*, *Magnesiæ sulphas*—s. *Enix-*

um. Potassæ supersulphas.—s. Epsomensis, Magnesie sulphas.—s. Essentielle Benzoes, Benjamin.—s. Essentielle corticis Peruviani, Quinine.—s. Febrifugus of Lemery, Potassæ supersulphas.—s. Febrifugus Sylvii, Potassæ murias.—s. Pontium, Soda, muriate of.—s. Fossilis, Soda, muriate of.—s. Fuliginis, Ammonie carbonas.—s. Gemmæ, Soda, muriate of.—s. Herbarum, Potassæ subcarbonas.—s. Indus, Saccharum album.—s. Lixiviosum, see Lixivium.—s. Lucidum, Soda, muriate of.—s. Marinus, Soda, muriate of.—s. Martis, Ferri sulphas.—s. Martis Muriatricum sublimatum, Ferrum ammoniatum.—s. Mirabile perlatum, Soda, phosphate of.—s. Mirabilis Glauberi, Soda, sulphate of.—s. Muriatricum baroticum, Baryta, muriate of.—s. Nitrum, Potassæ nitras.—s. Ossium, Ammonie carbonas.—s. Panchrestum, Potassæ tartras.—s. Petræ, Potassæ nitras.—s. Tartarum, Potassæ subcarbonas.—s. Plumbi, Plumbi superacetas.—s. Polychrestus, Potassæ sulphas.—s. Polychrestus Glaseri, Potassæ sulphas cum sulphure.—s. Polychrestum Seignetti, Soda, tartrate of.—s. Prunellæ, Potassæ nitras fusus sulphatis paucillo mixtus.—s. Rupellensis, Soda, tartrate of.—s. Sapientie, Alembroth salt, Potassæ sulphas.—s. Saturni, Plumbi superacetas.—s. Scientie, Alembroth salt.—s. Secretus Glauberi, Ammonie sulphas.—s. Sedativus Hombergi, Boracic acid.—s. Sedlicensis, Magnesie sulphas.—s. Seignetti, Soda, tartrate of.—s. Sennetti, Potassæ acetas.—s. Seydschützense, Magnesie sulphas.—s. Sodæ, Soda, subcarbonate of.—s. Sodomæus, Bit noben.—s. Succini, Succinic acid.—s. Tartari, Potassæ subcarbonas, see Potash.—s. Tartari essentielle, Tartaric acid.—s. Tartari sulphuratum, Potassæ sulphas.—s. Urinæ, Microcosmicum Sal.—s. Urinosum, Urea.—s. Vegetabilis, Potassæ tartras.—s. Vitæ, Alembroth salt.—s. Vitrioli, Zinci sulphas.—s. Vitrioli narcoticum, Boracic acid.—s. Volatile, Ammonie carbonas, Spiritus ammonie aromaticus.—s. Volatile Benzoes, Benjamin.—s. Volatile boracis, Boracic acid.—s. Volatile oleosum Sylvii, Spiritus ammonie aromaticus.—s. Volatile succini, Succinic acid.—s. Volatilis salis ammoniaci, Ammonie carbonas.—s. Vomitorii vitrioli, Zinci sulphas.

SALAAM CONVULSION, see Convulsion, Salaam.

SALAB, Salep.

SALACITAS, Satyriasis.—s. Vulvæ, Nymphomania.

SALAD, CORN, Valeriana dentata.

SALAP, Salep.

SALCOMBE, (CLIMATE OF.) A small place—the Montpelier of Huxham—which is one of the warmest spots on the S. W. coast of England, and, therefore, in this respect adapted for the phthisical invalid as a winter retreat.

SALEP, *Saloop'* (so pronounced.) *Salab*, *Salap*. The prepared bulb of the *Orchis mas'cula* and other species of *Orchis*. The salep of Cachmere is from a species of *Eulophia*. Salep is in small, irregular, hard masses. In its composition in relation to water, it is closely analogous to tragacanth. It is possessed of similar virtues with sago, tapioca, &c. The mucilage is prepared by dissolving the powdered salep in hot water, with assiduous stirring, and adding to the solution sugar and milk.

SALEP, Otaheite, Arrow-root, (Tabiti).

SALER'NUM, (SCHOOL OF.) An ancient school of medicine at Salerno in Italy. It was established by Charlemagne, in 802; and was the first Christian university at which medicine was taught. It was much celebrated, especially in the 12th century. The work on hygiene, "*De Conservandâ Valetudine*," which bears its name, has passed through several editions in various

languages. It is supposed to have been written by John of Milan, in 1100, and is in Leonine verses.

SALICARIA, *Lythrum salicaria*.

SALICINE, see *Salix*.

SALICORNIA ARABICA, &c., see *Soda*.

SALINE', *Salinus*, *Salina'cius*, *Salinae'idus*, *Salitus*, *Salsus*, *Hali'nus*, *Halm'y'rus*, *'adypnos*, *Halico'des*, *Sal'i'nous*, from *sal*. That which contains a salt, or has the properties of a salt.

SALINE DRAUGHT or MIXTURE. This may be made as follows:—*Potass. carbon.* ℥j, *acid. citric.* seu *tartar.* gr. xv seu succ. limon. recent. ℥ss, *aque* f℥viij, *syrup. cujusvis* ℥j. M. Given in divided doses as a febrifuge.

SALINE EFFERVESCING DRAUGHT or MIXTURE is made by adding a due proportion of either of the acids above mentioned, or of lemon-juice, to the proper proportion of the carbonate of soda or potassa. About 25 grains of the acid saturate 30 grains of the carbonate. See *Powders*, *soda*.

SALINES, see Kentucky, mineral waters of.

SALINOUS, *Saline*.

SALIS AMMONIACI FLORES, *Ammonie carbonas*.

SALITIO, Leap.

SALITUS, *Saline*.

SALIUM, *Saliva*.

SALIUNCA, *Valeriana Celtica*.

SALIVA, *Si'alon*, *Si'elon*, *Sialo'ma*, *Pty'sma*, *Pty'alon*, *Pty'elon*, *Sal'ium*, *Sputa'men*, *Sputa'mentum*, *Sputum*, *Spittle*, (Old. Eng.) *Spaul*, (Prov.) *Spoat*. An odorless, insipid, transparent, slightly viscid fluid; secreted by the parotid, submaxillary and sublingual glands, and poured into the mouth by the ducts of Steno, Wharton, and Rivinus; the use of which is to mix with the alimentary bolus, and to serve in the process of digestion. It is composed, according to Berzelius, of 992.2 parts of water, 2.9 of a particular animal matter, soluble in water, and insoluble in alcohol, *pty'alín*, *principium saliva'le*, *mat'ria sial'i'na*, *Ca'sein* of the *sal'iva*, *Sal'ivary abut'men*, of 1.4 of mucus, 1.7 of chlorides of potassium and sodium, 0.9 of lactate of soda and animal matter, and 0.2 of soda.

SALIVÆ FLUXUS CIBO VISO, see *Salivatio*.

SALIVAIRE, *Anthemis pyrethrum*, *Salivary*.

SALIVAL, *Salivary*.

SALIVANS, *Sialogogue*.

SALIVANT, *Sialogogue*.

SALIVARIS, *Salivary*—s. *Herba*, *Anthemis pyrethrum*.

SALIVARY, *Sal'ivarius*, *Saliva'ris*, *Salivo'rus*, *Sal'i'val*, *Sal'i'vous*, from *saliva*, (F.) *Salivaire*. That which belongs to the saliva.

SALIVARY ALBUMEN, see *Saliva*—s. *Calculi*, see *Calculi*, *salivary*.

SALIVARY FIS'TULE, *Syalosyrin'ges*, are such as depend upon an accidental opening into the excretory ducts of the salivary glands. *Fistulæ* are most commonly met with in the duct of Steno.

SALIVARY GLANDS, *Secreting organs of the saliva*, *Salival glands*, are situate symmetrically, to the number of four, on each side of the face. Their form is very irregular, and their extent varies according to individuals. Their excretory ducts—*Ductus Saliva'rii*, (F.) *Canaux ou Conduits salivaires*—carry the saliva directly into the mouth, without pouring it into any intermediate reservoir, as happens in the case of some of the other glands. The salivary glands have been distinguished into,—1. The *parotid*; 2. The *submaxillary*; 3. The *sublingual*; and 4. The *intra-lingual* or *lingual*, seated at the inferior surface of the tongue, where the mucous membrane forms a fringed fold.

SALIVATIO, Salivation—s. *Mercurialis*, Salivation, mercurial.

SALIVATION, *Saliva'tio*, *S. cal'ida*, *Epiph'ora ptyalis'mus*, *Sialis'mus*, *Sputum Oris*, *Sial'achus*, *Sielismos*, *Sialorrh'ea*, *Sialozoe'mia*, *Ptyalis'mus*, *Pty'alism*, *Fluxus sialiv'æ*, (F.) *Flux de Bouche*, *Flux salivaire*, *Ptyalisme*; from *Saliva*. A superabundant secretion of saliva; occasioned either locally, by the use of irritating masticatories, or under the influence of some cause which acts on the whole economy, and especially of mercurial preparations. In this last case it is accompanied by a coppery taste in the mouth, by swelling of the gums, and sometimes by looseness of the teeth. Salivation may usually be diminished by the use of astringents, laxatives, &c.

Ptyalismus acut'us sympathet'icus, *Salivæ fluxus cibo viso*, *Mouth watering*, the increased flow of saliva produced by the sight, smell, or thought of agreeable food, is a species of ptyalism, as well as the *Ptyalismus acutus mellit'us*, *Apoceno'sis ptyalis'mus mellitus*, *Dulce'do Sputato'rum*, or *Sweet spittle*. In this, the saliva has a sweet or mucilaginous taste.

SALIVATION, *MERCURIAL*, *Saliva'tio mercurialis*, *Sialozoe'mia* seu *Ptyalis'mus mercurialis*, *P. hydrargyratus*, (F.) *Ptyalismus mercuriel*, *Salivation mercurielle*. Salivation produced by mercury. The stomatitis that accompanies it is termed mercurial, *Hydrargyro-stomatit'is*, (F.) *Stomatite mercurielle*.

SALIVATION MERCURIELLE, Salivation, mercurial.

SALIVATUM, *Sialagogue*.

SALIVOSUS, *Salivary*.

SALIVOUS, *Salivary*.

SALIX. *The Willow*, *I'tea*, (Prov.) *Sally*. *Sally-withy*, (Sc.) *Saich*, *Sough*, *Willan*, (F.) *Saule*. A genus of plants in the Linnæan system. *Ord. Salicaceæ*.

SALIX FRAG'ILIS, *S. pen'dula* seu *decipiens* seu *amgdali'na* seu *persicifo'lia* seu *Russelia'na* seu *pippoph'æfolia* seu *Trevira'nii*, or *Crack willow* (Ph. D.) **SALIX ALBA**, *S. cæru'lea* seu *vitel'li'na*, or *White willow* (Ph. D. and Ph. U. S.), (F.) *Osier*. **SALIX LATIFO'LIA**, or *broad-leaved willow*. **SALIX CA'PREA**, *S. Sphacela'ta* seu *toment'osa*, or *Great round-leaved willow* (Ph. L.) **SALIX PENTAN'DRIA** seu *HERMAPHRODIT'ICA*. **SALIX HU'MILIS**, *Low bush willow*, *Speckled willow*. **SALIX VITULI'NA**, **SALIX ERIOCEPH'ALA**. The bark of each of these has a slightly aromatic odour, and a bitter astringent taste, which is nearly the same in all the species. It is tonic and astringent, and has been used in intermittents, and in cases of dyspepsia, worms, &c. The active principle has been separated, and called *Sal'icine*. It has been found in about fourteen species of salix and eight species of *Populus*; is like the quinia in medicinal efficacy. Dose, ʒj to ʒj of the powder. Dose of *Salicine* from 4 to 6 grains.

SALIX EGYPTIACA, *Calaf*.

SALLENDERS, *Malandria*.

SALLIS, *Adeps*.

SALLOW, *Lurid*.

SALLY, *Salix*.

SALLY-WITHY, *Salix*.

SALMIAC, *Ammonie murias*.

SALOOP, *Salep*. Also, *sassafras tea* flavored with milk and sugar. Under this name a tea is sold in the streets of London which consists of an infusion of the wood of *Laurus Sassafras* with the addition of milk and sugar—*Sassafras Tea*.

SALPETRA, *Potassæ nitras*, *Soda*, *nitrate of*.

SALPINGEMPHRAX'IS, from *σαλπιγξ*, *σαλπιγγος*, 'a tube,' and *εμφραξις*, 'obstruction.' Ob-

struction of the Eustachian tube;—a cause of deafness.

SALPINGITIS, from *σαλπιγξ*, *σαλπιγγος*, 'a tube,' and *itis*, 'denoting inflammation.' Inflammation of the Eustachian tube.

SALPINGOCYCE'ISIS, *Paracyc'esis tuba'ria*, *Gravid'itas tuba'ria*, from *σαλπιγξ*, *σαλπιγγος*, 'a tube or trumpet,' and *κυσις*, 'pregnancy.' Tubal pregnancy.—See *Pregnancy*, tubal.

SALPINGOMALL'ÉEN, *Tensor tympani*.

SALPINGOPHARYNG'ÆUS, from *σαλπιγξ*, *σαλπιγγος*, 'a trumpet,' and *φάρυγξ*, 'the pharynx.' Valsalva and Douglas have given this name to a bundle of fibres, which passes from the Eustachian tube to the pharynx.

SALPINGOSTAPHYLINUS, *Circumflex muscles*—s. *Staphylinus internus*, *Levator palati*.

SALPINGOSTENOCH'ORIA, from *σαλπιγξ*, *σαλπιγγος*, 'a tube,' *στενός*, 'narrow,' and *χωρα*, 'space.' Narrowness or stricture of the Eustachian tube.

SALPINX, *Tuba*.

SALSAMENTUM CRUDUM, *Omotarichos*.

SALSAPARILLA GERMANICA, *Sarsaparilla Germanica*.

SALSEDO MUCRUM, *Potassæ nitras*.

SALSEPAREILLE, *Smilax sarsaparilla*—s. *d'Allemagne*, *Sarsaparilla Germanica*—s. *du Brésil*, s. *Coraque*, s. *de Honduras*, s. *de Pérou*, and s. *de Portugal*, see *Smilax sarsaparilla*.

SALSIFI, *Tragopogon porrifolium*.

SALSILAGO, *Muria*.

SALSOLA DECUMBENS, *S. Kali*.

SALSO'LA KALI, *S. Decumb'bens*. *Kali soda* seu *spino'sum cochleat'um*, *Prickly Saltwort*, *Tragus* sive *Tragum Matthi'oli*. *Ord. Chenopodiaceæ*; from *salt*, 'salt,' owing to its saline properties. This plant, as well as *Salsola sati'va* and *Salsola soda* seu *longifo'lia*, *Kali iner'mis*, *Herba vitri*, produces the alkaline salt, commonly called *Barilla*, *Soda*, or *Kelp*.

SALSOLA LONGIFOLIA, *S. Kali*—s. *Sativa*, &c., see *S. Kali* and *Soda*—s. *Soda*, see *S. Kali*.

SALSUGINOSUS, *Halmyrodes*.

SALSUS, *Saline*.

SALT, *Sal*, *Hal*, *Hals*, 'αλς', (F.) *Sel*. A name formerly given to a considerable number of substances of very different nature. At present, chymists apply the term *Salts* exclusively to the combination of an acid with one or more bases. *Neutral salts* are those in which the acid and base reciprocally saturate each other. *Acid salts* or *supersalts*, (F.) *Sur-sels*, are those in which the acid predominates over the base. *Alkaline salts* or *subsalts*, (F.) *Sous-sels*, those in which the base is, on the contrary, predominant. The different denominations of the salts are compounded, 1. Of the name of the acid; the termination of which is changed into *ite*, when the name of the acid is in *ous*;—*ate*, when it is in *ic*. Thus the salts formed by the sulphurous acid are *sulphites*; those by the sulphuric, *sulphates*. 2. This name is followed up by that of the oxyd, which enters into the composition of the salt. Thus, the *sulphate of protoxyd of iron* is a combination of sulphuric acid with the protoxyd of that metal. Frequently, to shorten the name, the word *oxyd* is suppressed; and *proto*, *deuto*, *trito*, &c., put before the acid:—as *protosulphate of iron*. *Simple salts* are those resulting from the combination of an acid with a single base; *double salts* with two different bases: *triple* with three, &c. A *haloid salt* is one constituted after the type of common salt, or which contains a metal and a *salt-radical*, as chlorine, iodine, &c. An *oxy salt* is formed from the combination of an acid with an oxide.

When *salt* is used in an abstract sense, it means the chloride of sodium.

SALT. APERIENT, OF FREDERICK, Soda, sulphate of—s. Baker's, Ammoniac carbonas—s. of Barilla. Soda, subcarbonate of—s. Bay, Soda, muriate of—s. Bitter purging, Magnesie sulphas—s. Common, Soda, muriate of—s. Epsom, Magnesie sulphas—s. Glauber's, Soda, sulphate of—s. (Glauber's, Lymington, see Soda, sulphate of—s. Preston, see Ammoniac carbonas—s. Rochelle, Soda, tartrate of—s. Rock, Soda, muriate of—s. Sea, Soda, muriate of—s. Secondary, Neutral salt—s. Seignette's, Soda, tartrate of—s. Smelling, Ammoniac carbonas—s. of Soda, Soda, subcarbonate of—s. Sore-throat, Potassæ nitras fusus sulphatis paucillo mixtus.

SALT OF TARTAR, see Potassæ subcarbonas, and Potash.

SALT, TASTELESS PURGING, Soda, phosphate of.

SALT OF URINE, Ammoniac carbonas.

SALT, WHITE, Soda, muriate of.

SALT OF WOODSOOT, Ammoniac carbonas—s. of Wormwood, see Potash.

SALTANS ROSA, Urticaria.

SALTATIO, Dancing—s. Sancti Viti, Choreia.

SALT'ERTON. A village about four miles to the eastward of Exmouth, England, frequented as a winter abode by invalids. It is in a small, open valley on the seashore, and is well protected from winds, particularly the northerly.

SALTISH, Halmyrodes.

SALTPETRE, Potassæ nitras.

SALTUS, Leap, Vulva—s. Tendinum, Subul-tus tendinum.

SALTWORT, PRICKLY, Salsola kali.

SALUBRIOUS, Salutory.

SALUBRIOUSNESS, Salubrity.

SALUBRIS, Salutory.

SALUBRITY, *Salu'britas*, *Salu'briousness*, (F.) *Silubrité*, from *salus*, 'health.' Any thing which contributes to the health of individuals or of communities. Situations differ materially from each other in this respect; and the cause of the difference is often an interesting topic of inquiry.

SALUS, Sanitas.

SALUTAIRE, Salutory.

SALUTARY, *Peries'ticos*, *Saluta'ris*, *Salu'ber*, *Salu'bris*, *Hygie'nis*, *Hygias'ticus*, *Hygie'rus*, *Salu'brius*, (F.) *Salutaire*, *Salubre*. Same etymon. Healthy. That which is favourable to health. Some diseases have been considered salutary; but this is doubtful.

SALUTIFEROUS, *Salu'tifer*; from *salus*, *salutis*, 'health,' and *fero*, 'I bear.' Bringing health; health-bearing.

SALVATELLA; same etymon, [?] *Vena salvatella*, is situate on the back of the hand, near its inner margin. It begins on the posterior surface of the fingers and dorsal surface of the hand, by a plexus, formed of a number of radicles, frequently anastomosing with each other; it then ascends to the inner part of the forearm, where it takes the name—*posterior cubital*. The ancients recommended this vein to be opened in certain diseases, as in melancholic and hypochondriacal affections; and they attributed to such abstraction of blood considerable efficacy in the cure of disease: hence its name.

SALVE, BLISTERING, Unguentum lyttæ—s. Deslier's, Ceratum resinæ compositum—s. Eye, Unguentum oxidi zinci impuri—s. Eye, Single-ton's, Unguentum hydrargyri nitrico-oxydi—s. Healing, Cerate of calamine.

SALVIA, (from *salveo*, 'to be sound,' owing to its reputed healing qualities;) *S. officinalis* seu *horten'sis* seu *vulga'ris* seu *minor*, *Eleis'tiaca*, *Sage*, *Sage*, (F.) *Sauge*, *Petite Sauge*, *Thé d'Europe* ou *de France*. Ord. Labiata. Scr. Syst. Diandria Monogynia. Sage leaves, *Salvia* (Ph. U. S.), have a peculiar, aromatic odour, and warm, aromatic, bitterish taste. Their virtues

depend upon an essential oil. They are tonic, stimulant, and carminative. In the form of a weak infusion, Sage is occasionally used as a diluent, and as a wash for sore mouth. Dose, gr. xv to xxx or more.

SALVIA AFRICA'NA, *Wild Sage*, a South African species, has the same medical properties as *Salvia*.

SALVIA COLORATA, *Salvia*.

SALVIA HORMI'NUM, *S. colorata*, *Horminum*, *H. coloratum* seu *sati'vum*, (F.) *Sauge-hormin*, of the Pharmacopœias, is used in similar cases.

SALVIA HORTENSIS, *Salvia*.

SALVIA LYRA'TA, *Wild Sage*, *Cancer Weed*; indigenous. The fresh leaves, bruised, are used to destroy warts. It has been supposed to be of use in cancer.

SALVIA MINOR, *Salvia*—s. *Officinalis*, *Salvia*.

SALVIA SCLA'REA, *Sclarea*, *Gallitrichum*, *Æthio'pis*, *Biser'mas*, *Colus Jovia*, *Common Clary*, (F.) *Sauge esclée*, *Grand sauge*, *Orvale* ou *Tout-bonne*. The leaves have been recommended as corroborants and anti-spasmodics.

SALVIA VITÆ, *Asplenium ruta muraria*—s. *Vulgaris*, *Salvia*.

SAMADERA IN'DICA, Ord. Simarubæ, furnishes the *Niepa bark*, which is possessed of antiperiodic properties.

SAMBU'CUS, from *σαμβυχη*, an ancient musical instrument, said to have been made of its wood. *Sabu'cus*, *Sambucus nigra* seu *vulga'ris* seu *arbo'rea*, *Acté*, *Infelix lignum*; *Elder*, *Ellorne*, *Ellen-tree*, *Ellarne*, (Sc.) *Boartree*, *Bortree*, *Bountree*, (Prov.) *Bottry tree*, *Burtree*, *Ellet*, *Elren*, *Seavo*, (F.) *Sureau commun*. Ord. Caprifoliaceæ. Scr. Syst. Pentandria Digynia—(the flower and bark were formerly called *Cano'pum*). The flowers, berries, and bark of the elder have been used in medicine. The flowers have been reputed diaphoretic and discutient; the berries, aperient and sudorific; the bark, purgative, and, in small doses, deobstruent. The flowers are sometimes used in fomentations, and to form a cooling ointment.

SAMBUCUS ARBorea, *Sambucus*—s. *Chamæ-acte*, *S. ebulus*.

SAMBUCUS EB'ULUS, *S. Chamæacté* seu *he'lion* seu *hu'milis* seu *herba'cea*, *Ebulus*, *Chamæan'te*, *Od'occos*, *Dwarf-elder*, *Danewort*, (F.) *Hèble*, *Yèble*. The root, interior bark, leaves, flowers, *Sambucus* (Ph. U. S.), berries, (*Bacca* seu *Grana Actes*), and seeds, have all been given in moderate doses as resolvents and deobstruents; and in large doses as hydragogues. The berries of *Sambucus Canadensis*, the Common Elder of America, are made official in the Pharmacopœia of the United States. A distilled water of the flowers, *Aqua Sambuci*, *Elder Water*, is official in the London and Edinburgh Pharmacopœias.

SAMBUCUS HELION, *Sambucus ebulus*—s. *Herbacea*, *S. ebulus*—s. *Humilis*, *S. ebulus*—s. *Nigra*, *Sambucus*.

SAMBUCUS PU'BENS, *S. pubescens* seu *race-mo'sa*, *Red-berried Elder*; indigenous. The bark, in decoction, is said to be a valuable hydragogue cathartic.

SAMBUCUS RACEMOSA, *S. pubens*—s. *Vulgaris*, *Sambucus*.

SAMI TERRA, *Samian Earth*. An argillaceous earth, found in the island of Samos, to which were ascribed virtues similar to those of the Terra Lemni.—Galen, Dioscorides.

SA'MIUS LAPIS. A stone found in the Isle of Samos, which the ancients believed to be astrigent and refrigerant.

SAMPHIRE, *Crithum maritimum*.

SAMPSON, BLACK, *Echinacea purpurea*.

SAMPUSCUS, *Thymus mastichina*.

SAMPUSCUS, *Origanum majorana*.

SAMPSYCHINON, *Sampsychinum*. An ointment composed of *σαμψυχον*, which was perhaps marjoram. — Dioscorides.

SANABILIS, Curable.

SANABLE, Curable.

SANAMUNDA, *Geum urbanum*.

SANATIO, Curation, Cure.

SANATIVE, Curative.

SANATORIUM, Sanitarium.

SANATORY, Curative.

SAN GIULIANO, MINERAL WATERS OF.

At the foot of the mountain San Giuliano, a league and a half from Pisa, are several mineral springs; temperature from 84° to 107° Fahr.

The spring, Pozzetto, contains carbonic acid, sulphates of soda, lime, and magnesia; chlorides of sodium and magnesium; carbonates of lime and magnesia; argil and silica. — Santi.

SAND, see Gravel — s. Brain, see Pineal gland — s. Pineal, see Pineal gland.

SANDALUM CÉRULEUM, *Guilandina moringa* — s. Rubrum, *Pterocarpus santalinus*.

SANDARAC, *Sandara'cha*, *Gum Juniper* or *Sanderach*, *Pounce*, *Gummi Juniperi*, *Sandara'ca*, *Vernix sicca*, *Resina Juniperi*, (F.) *Sandaracque*, *Gomme de Genève*. A name given, by the Arabs, to an odoriferous resin, that flows from the *Thuya aphylla*. Ord. Coniferae. *See. Syst.* Monœcia Monadelphia. It is stimulant, like all the resins; but is very little used.

SANDARACHA GRÆCORUM, Realgar.

SAND-BLIND, nearly blind. A defect of vision, in which small particles seem to fly before the eyes. *See* Metamorphopsia.

SANDERS, see Saunders.

SANDIVER, see Vitrum.

SANDIX, *Plumbi oxydum rubrum*.

SANDYX, *Plumbi oxydum rubrum*.

SANG, Blood — s. *Artériel*, Arterial blood — s. *Dragon*, *Rumex sanguineus* — s. *Dragon*, see *Calamus rotang* — s. *Globules du*, Globules of the blood — s. *Perte de*, Hæmorrhagia.

SANGAREE', Wine and water, sweetened and spiced.

SANGLOT, Sob.

SANGSUE, Hirudo.

SANGUIDUCTUS DURÆ MATRIS, Sinuses of the dura mater.

SANGUIIFICATIO, Hæmatosis — s. Debilior, Hyphæmatosis.

SANGUIFLUXUS, Hæmorrhagia — s. Uteri-nus, Metrorrhagia.

SANGUIMOTION, Circulation.

SANGUIMOTORY, Circulatory.

SANGUIN, (*Vaisseau*), Blood-vessel.

SANGUINALIS CORRIGIOLA, *Polygonum aviculare*.

SANGUINA'RIA, from *sanguis*, *sanguinis*, 'blood,' owing to the colour of its juice. *S. Canadensis* seu *acaulis*, Bloodroot, Bloodwort, Puccoon, Red Puccoon, Red Root, Indian Paint, Turmeric. *Nat. Ord.* Papaveraceæ. *See. Syst.* Polyandria Monogynia. A North American plant, the rhizoma, *Sanguinaria* (Ph. U. S.), and seeds of which are used in medicine. They are emetic. Dose of the powder, 8 to 20 grains. They are said to reduce the velocity of the pulse, like digitalis.

SANGUINARIA, *Polygonum aviculare*.

SANGUINE, *Sanguineus*, *Sanguineus*, *Hæmatæus*, *Hæmato'des*, *Hæmato'rus*, *Hæmati'nus*, *Hæmo'des*, *Hæmo'i'des*, *Sanguineus*, *Hæmat'ic*, (F.) *Sanguin*. From *sanguis*, *sanguinis*, 'blood,' Bloody, blood-like; having the colour of blood, as 'a sanguine countenance.' Relating to, or containing blood — as 'sanguine temperament.'

Plethoric.

SANGUINEA, *Potassæ nitras*.

SANGUINENESS, Plethora.

SANGUINEOUS, Plethoric, Sanguine.

SANGUINEUS, Sanguine.

SANGUINIS ABSORPTIO, Hæmorrhophesia — s. Congestio, Congestion — s. Dissolutio, Hæmæteclysis — s. Fluor, Hæmoptysis — s. Fluor Vesicæ, Cystorrhagia — s. Ictus, Apoplexy — s. Missio. Blood-letting — s. Profluvium copiosum, Hæmorrhagia — s. Profluvium ex Ore, Stomatorrhagia — s. Stagnatio, Hæmostasis.

SANGUIN'OLENT, *Sanguinolentus*, from *sanguis*, *sanguinis*, 'blood.' Tinged with blood. An epithet applied, particularly, to the sputa when so tinged. Bloody.

SANGUINOSUS, Sanguine.

SANGUIPURGIUM, from *sanguis*, 'blood,' and *purgare*, 'to cleanse.' Purgation of the blood. A name given by some authors to mild fevers, from an idea that they purged the blood of noxious matters.

SANGUIS, Blood, Sperm — s. Concretus, Coagulum — s. Draconis, see *Calamus rotang* — s. Hirci siccatus, see Blood — s. Menstruus, see Menses — s. in Urinâ, Hæmaturia — s. Virginis primum menstruatum, see Menses.

SANGUISOR'BA CANADENSIS, *Canada Burnett*. From *sanguis*, 'blood,' and *sorbere*, 'to absorb.' A tall indigenous plant — Ord. Rosaceæ — with white, sometimes purple, flowers, which appear from August to October. It has the same virtues as

SANGUISOR'BA OFFICINA' LIS, *S. carnea* seu *rubra*, Italian *Pimpinell*, *Pimpinella Italica* seu *officina'lis*, (F.) *Pimpinella noire*. It was formerly used as an astringent and tonic, in hemorrhages, dysentery, &c.

SANGUISUGA, Hirudo.

SANGUISUGUM, from *sanguis*, 'blood,' and *sugere*, 'to suck.' A barbarous term, to denote too great an accumulation of blood in the heart. — Bonetus.

SANICLE, *Sanicula*, *S. Marilandica* — s. American, *Heuchera cortusa* — s. de *Mariland*, *Sanicula Marilandica* — s. Yorkshire, *Pinguicula vulgaris*.

SANICULA, (from *sano*, 'I heal,') *Saxifraga granulata*.

SANIC'ULA, *S. Europæa* seu *officina'rum* seu *mas*, *Astrantia diaph'nia*, *Cucull'ia Sanicula*, *Cucull'ia*, *Dodeca'theon*, *Sym'phytum petra'um*, *Diaph'nia*, *D. cortusa*, *San'icle*, *Wood March*, (F.) *Toute-saine*. Ord. Umbelliferae. *See. Syst.* Pentandria Digynia. This herb was formerly used as a mild astringent. It is bitterish, and somewhat austere.

SANICULA Eboracensis, *Pinguicula vulgaris* — s. *Europæa*, *Sanicula*.

SANIC'ULA MARILAN'DICA, *San'icle*, *Ma'ryland Sanicle*, *Black Snakeroot*, (F.) *Sanicle de Mariland*. This indigenous plant grows in marshes, or low and shaded ground. It is possessed of tonic virtues; and the powdered root is employed in popular practice as a favourite remedy in intermittent fever. It has been recommended, also, in chorea.

SANICULA MONTANA, *Pinguicula vulgaris* — s. *Officinarum*, *Sanicula*.

SANIDO'DES, *Sanio'des*, 'resembling a board,' from *sans*, gen. *sansûs*, ('a board,' and *idos*, 'resemblance.' One whose thorax is narrow, and, as it were, contracted.

SANIE, Sanies.

SA'NIES, *Ichor*, *Tabum*, *Virus*, *Pus malignum* seu *corrosivum*, (F.) *Sanie*, corrupted from *sanguis*, 'blood.' A thin, serous fluid, presenting some of the qualities of pus and blood, and commonly exhaled at the surface of ulcers.

SANIEUX, Ichorous.

SANIODES, Sanidodes.

SANTOSUS, Ichorous.

SANTOUS, Ichorous.

SANITA'RIMUM, *Sanato'rium*. Same etymology as the next. A retreat selected for valetudina-

rians, in consequence of its salubrity. Also, an establishment for the treatment of the sick.

SANITARY, *San'atory*, from *sanitas*, 'health,' (F.) *Sanitaire*. That which relates to health, and especially to public health; as *établissement sanitaire*, *police sanitaire*.

SANITAS, from *sanus*, 'healthy,' *Hygie'a*, *Hyge'a*, *Hygian'sis*, *Hygi'asis*, *Integ'ritas*, *Prosper'itas*, *Valetu'dinis*, *Salus*, *Hygie'i'a*, *Valetu'do*, *Valitu'do*, *Valetu'do bona seu secu'da seu com'moda*, *Hygie'i'a*, *Euzoödyna'mia*, *Health*, (Sc.) *Heil*, *Heyle*, *Heal*, (F.) *Santé*, from *sanus*, 'sound.' Health has been defined,—a state, in which all the functions are exerted with regularity and harmony. A person may, however, be in health, although certain functions are not performed. A blind man, for example, has health when all his functions, except sight, proceed with regularity. *Sanity* is often used for soundness of mind.

SANITAS, Cure.

SANITY, *Sanitas*.

SANKIRA, *Smilax China*.

SANS PAIRE, (*Veine*), *Azygos vein*.

SANTA CRUZ, (CLIMATE OF.) This West India island is frequently selected as a winter retreat for the phthisical valetudinarian from the United States; but it does not seem entitled to any preference, in such cases, over the other islands of the West Indies, whilst its general salubrity is said to be less than that of many.

SANTAL BLANC, *Santalum album*—*s. Rouge*, *Pterocarpus santalinus*.

SANTALUM, see *Pterocarpus santalinus*.

SANTALUM ALBUM, *S. citrinum seu pallidum seu myrtifolium*, *Sir'ium myrtifolium*, *Yellow Saunders*, (F.) *Santal blanc*. Ord. *Santalaceæ*. Ser. *Syst.* *Tetrandria Monogynia*. The virtues of the wood of the white saunders reside in a volatile oil and resinous principle. The odour is fragrant, and resembles a mixture of ambergris and roses. It has been used as a stimulant and sudorific, but it is not now employed. The saunders woods are called, collectively, in French, *Santaux*.

SANTALUM MYRTIFOLIUM, *S. album*—*s. Rubrum*, *Pterocarpus santalinus*.

SANTÉ, *Sanitas*.

SANTONICUM, *Artemisia santonicum*.

SANTONINE, *San'tonin*, *Cinin*. A peculiar crystalline substance obtained from *Artemisia Santonica*. It possesses the anthelmintic properties of the plant. Dose, five or six grains.

SANTORINI, **FISSURES OF**, (F.) *Incisures ou Scissures de Santorini*. Clefts at the upper and back part of the fibro-cartilaginous portion of the external ear.

SANTORINI, **TUBERCLES OF**. Small cartilaginous projections at the top of the arytenoid cartilages, which support the ligaments of the glottis. See *Corniculum Laryngis*.

SANVE, *Sinapis*.

SAO'RIA, *Saurja*. The ripe and dried fruit of *Maesa seu Bacob'otrys pici*, an Abyssinian tree, which is administered in its native country, in powder, in doses of from 32 to 44 grammes, as a tæniifuge.

SAP, *Sax. sæpe*. (G.) *Soft*, *Ar'borum Succus seu Lympha*, (F.) *Sève*. The circulating fluid of the vegetable, which corresponds to the blood of the animal.

SAPA. see *Rob*.

SAPHA'DA. Paracelsus has given this name to reddish scales adherent to the hair in some eruptions of the hairy scalp.

SAPHENA, *Saphena*.

SAPHENA, *Saphæ'na*, from *σαφης*, 'manifest,' 'evident,' (F.) *Saphène*. A name given to two subcutaneous veins of the lower limb. These

are: 1. The **GREAT SAPHENA VEIN**, *S. interna*, *Tibio-malleolar* (Ch.), (F.) *Grande Veine Saphène ou Saphène interne*. Its radicles occupy the inner part of the great toe, and form a transverse arch on the back of the foot, which unites with the saphena externa, and receives at its convex part, which is turned forwards, a number of branches coming from the toes. These radicles, when united, ascend before the inner ankle, and form a trunk, the size of which is farther increased by several branches that proceed from the region of the tarsus, and that of the metatarsus. This trunk ascends vertically at the inner part of the leg; passes behind the inner condyle of the femur; proceeds along the inner part of the thigh; receives other subcutaneous veins, such as the abdominal subcutaneous, the external pudic, &c., and passes through the inferior opening of the crural canal to empty itself into the femoral vein. 2. The **LESSER VENA SAPHENA** or *Saphena externa*, *Perone'o-malleolar Vein* of Chaussier, (F.) *Petite Veine Saphène ou Saphène externe*, commences by several roots, which are spread over the dorsum and outer side of the foot, and embrace the outer ankle. Behind this, they unite into a single trunk, which ascends obliquely, approaching the tendo Achillis, and proceeds vertically towards the ham to open into the popliteal vein.

SAPHÈNE, *Saphena*.

SAPHE'NOUS, *Sapheno'sus*; from *Saphena*, Of or belonging to the saphena.

SAPHE'NOUS NERVES are two in number, and are branches of the crural or femoral nerve. They are the short *saphenous*—*nervus cutaneus internus*; and the long *saphenous*—*nervus cutaneus internus longus*.

The external *saphenous*, formed by the union of the communicans poplitei and communicans peronei, passes down the leg by the side of the external saphenous vein of the foot.

SAPID, *Sa'porous*, *Sap'idus*, (F.) *Savoureux*, from *sapere*, 'to taste.' Possessed of sapidity. Having the power of affecting the organs of taste.

SAPID'ITY, *Sap'idness*, *Saporos'ity*, *Sapor*, (F.) *Sapidité*. The quality of bodies that gives them taste.

SAPIN COMMUN, *Pinus picea*.

SAPIN'DUS, (*sapo*, and *Indus*, 'Indian soap;') *S. Saponaria*, *Saponaria*, *Soap Tree*, (F.) *Savonnier commun*. Ord. *Sapindaceæ*. The name of the plant that affords the *Soap Nute*, *Saponaria Nu'cula*, *Bac'ce Bermuden'ses*, *Soap-berries*. A spherical fruit about the size of a cherry, the cortical part of which is yellow, glossy, and so transparent as to show the spherical black nut that rattles within, and includes a white kernel. The cortical part of this fruit, which has similar effects with soap in washing, has been recommended in chlorosis, and as a stomachic and deobstruent.

SAPO, *Soap*, *Sax. Tape*, *Smegma*, (F.) *Savon*. A product obtained by treating fatty bodies with caustic alkalies dissolved in water. In the Pharmacopœia of the United States, it means Soap made with soda and olive oil. See *Sapo medicinalis*.

YELLOW SOAP is formed of turpentine, oil, and alkali.

Castile Soap, being made of fine olive oil, is generally used internally. The yellow and soft soaps are the most stimulating; and are, consequently, best adapted for particular cutaneous eruptions.

Internally, soap is purgative and diuretic; externally, it is detergent and stimulant. Soapy water is used with the greatest success, in cases

of poisoning by acids. It is mostly applied externally to bruises and sprains. In *Pharmacy*, it is used to favour the solution of resins.

SAPo AC'IDUS, *Acid Soap*, is a combination of a fixed oil with an acid.

SAPo AMMONIACALIS, *Linimentum ammoniæ fortius*—s. *Ammoniæ oleaceus*, *Linimentum ammoniæ fortius*—s. *Amygdalinus*, *Sapo medicinalis*—s. *Cacaotinus*, *Sapo coconus*—s. *Calcarius*, *Linimentum aquæ calcis*—s. *Castiliensis*, see *sapo medicinalis*.

SAPo COCO'NEUS, *Sapo cacaoti'nis*, *Soap of Cocoa-Nut Oil*, (F.) *Savon de cacao*, has been much used in Germany in herpetic affections.

SAPo DURUS, *S. medicinalis*—s. *Hispanicus*, see *S. medicinalis*.

SAPo KALICUS, *S. Mollis*.

SAPo LIQ'UIDUS, *Liquid Soap*, *Lotio saponacea*, is a cosmetic which is composed of olive oil $\frac{3iv}$, *ol. tartari per deliquium* $\frac{3ss}$. Rub together, and add *aquæ rosarum* $\frac{3xij}$.

SAPo MEDICINALIS, *Sapo amygdali'nis*, *Medicinal Soap*, (F.) *Savon médicinal*, is made with oil of sweet almonds and half its weight of caustic alkali, or, like the *Sapo Hispan'icus* seu *Castilien'sis*, *Spanish or Castile soap*, *Sapo durus*, *Sapo* (Ph. U. S.), (F.) *Savon dur*, of olive oil and soda.

SAPo MOLLIS, *S. viridis* seu *nig'er* seu *kal'licus*, *Common or Soft Soap*, (F.) *Savon mou ou noir*, is made of potass and oil or tallow. The *Sapo vulgaris* or 'common soap' of the Pharm. U. S. is made with soda and animal oil.

SAPo NIGER, *Black Soap*, *Melanosmeg'ma*, is a composition of train oil and an alkali: and **GREEN SOAP**, *Sapo viridis*, of hempseed, linseed, or rape oil, with an alkali. See *Sapo mollis*.

SAPo VEGETABILIS, *Pulvis gummosus alkalinus*—r. *Viridis*, *Sapo mollis*—r. *Vulgaris*, see *Sapo mollis*.

SAPONA'CEOUS, *Sapona'ceus*, (F.) *Saponacé*, from *sapo*, *saponis*, 'soap.' Soapy. Containing soap, as *Linimentum saponaceum*, *Soap liniment*.

SAPONAIRE, *Saponaria*.

SAPONA'RIA, from *Sapo*, 'soap,' the juice forming a lather with water. *Soap-wort*, *Bruise-wort*, *Bouncing Bet*, *Struthium*, *Lana'ria*, *Lychnis officinalis* seu *elysæ'tris*, *Ibici'uma*, *Booti'a vulgaris*, (F.) *Saponaire*, *Savonnère*. Ord. Caryophyllaceæ. *Sex. Syst.* Decandria Digynia. A decoction of the leaves of this plant has been recommended in scrofula, cancer, old syphilis, cutaneous eruptions, jaundice, visceral obstructions, &c. It is hardly ever used.

SAPONARIA, *Sapindus saponaria*.

SAPONARIE NUCULÆ, see *Sapindus saponaria*.

SAPONÆ. Soap, to which are added substances capable of communicating to it new properties, without affecting its own.—Béral.

SAPONEA. A pectoral medicine, prepared with sweetened syrup of violets, and oil of sweet almonds.

SAPONULES, *Sapon'uli*, (F.) *Savonules*. Same etymon. Combinations of volatile or essential oils with different bases.

ACID SAPONULES are combinations of such oils with different acids.

SAPONULÆ. A tincture (*alcoolé*), sufficiently charged with soap to cause it to assume a gelatinous shape.—Béral.

SAPONURE. A medicine, formed of soap, in powder, and of resinous or extractive matters. **SAPOR**, *Sapidity*, *Savour*.

SAPOROSITY, *Sapidity*.

SAPOROUS, *Sapids*.

SAPOTA, *S. Alchras*, *Achras Sapota* seu *australis* seu *Zapota*, *El nispero*, *Oval-fruited Sapota*, (F.) *Sapotier*, *Sapotillier*, *Sapotilla*. Fam.

Sapotacæ. *Sex. Syst.* Hexandria Digynia. A tree which grows in the Antilles, and particularly at St. Domingo. Its fruit is much esteemed. It resembles the marmalade of quinces; and hence has been called *natural marmalade*. The seeds make an emulsion, which has been given in nephritic affections. The bark has been recommended as a febrifuge.

SAPOTA ALCHRAS, *Sapota*.

SAPOTIER, *Sapota*.

SAPOTILLA, *Sapota*.

SAPOTILLIER, *Sapota*.

SAPROPYRA, *Typhus gravior*.

SAPROS, *Putrid*, *Rancid*.

SAPROS/TOMUS, from *sapros*, 'foul, of a bad odour,' and *stoma*, 'mouth.' One who has an offensive breath.

SAPROTES, *Putrefaction*.

SARAPUS, *Sarapous*. One who has a large foot; whose toes are largely separated, or whose feet are much asunder in walking; from *sarpos*, 'I grin,' and *rous*, 'the foot.' See *Kyllosis*.

SARATOGA, MINERAL WATERS OF. The Saratoga springs in New York are ten miles north-east from Ballston. They contain the same ingredients as those of the latter place; and are much frequented by visitors from every part of the United States. The following analysis has been given of the *Congress spring*. In a wine gallon—*gaseous contents*: carbonic acid, 311 cubic inches; atmospheric air, 7. *Solid contents*: chloride of sodium, gr. 385; iodide of sodium, gr. 3.5; bicarbonate of soda, gr. 8.982; bicarbonate of magnesia, gr. 95.788; carbonate of lime, gr. 98.098; carbonate of iron, gr. 5.075; silica, gr. 1.5; bromide of potassium, a trace. Total, gr. 597.943.—Steel.

The *Walton* or *Iodine spring* contains chloride of sodium, carbonate of magnesia, and iodide of sodium.

SARCEPIPOCELE, *Sarcoëpiplocene*.

SARCEPIPLOMPHALOCELE, *Sarcoëpiplomphalocèle*.

SARCEPIPLOMPHALUS, *Sarcoëpiplomphalus*.

SARCIDIUM, *Caruncle*.

SARCINA, *Fœtus*.

SARCINA VENTRIC'ULI, *S. Goodsir'ii*, *L. Sarcina*, 'a pack, a woolpack;' *Merimopœdia ventriculi* (Robin). A microscopic cryptogamous plant found in the fluid of waterbrash.

SARCION, *Caruncle*.

SARCITES, *Anasarca*—s. *Flatusus*, *Emphysema*.

SARCITIS, *Myositis*.

SARCOCE'LE, *Hernia carno'ea*, *Scirrhus Testic'uli*, from *sarx*, *sarkos*, 'flesh,' and *κηλη*, 'a tumour;' *Sarco'ma scroti*, (F.) *Hernie charnue*. *Scirrhus* or cancer of the testicle. This disease affects adults particularly; and appears most commonly after an inflammatory swelling of the testicle. Sometimes it is dependent upon a blow; at others, it makes its appearance without any appreciable cause. It is well known by a hard, heavy, ovoid or spherical swelling of the testicle, which is, at first, slightly, or not at all painful, and merely causes an unpleasant traction on the spermatic cord. There is no heat or change of colour of the skin; the spermatic cord is swollen, and participates in the affection; very painful shootings occur; the lymphatic glands of the abdomen become swollen, and form a tumour, which may, at times, be felt through the abdomen; and the patient, at length, dies with every sign of the cancerous diathesis. The prognosis is very unfavourable. The only means, indeed, that can save life, is the extirpation of the testicle.

SARCOCOL'LA, (F.) *Colle-chair*, from *sarx*,

'flesh,' and *κόλλα*, 'glue.' A resinous matter, obtained from *Penæa Sarcocolla*, *P. micantha*, and other species of *Penæa*; an African shrub, which has been so called from a belief that it facilitates the consolidation of flesh. It is in the form of small, oblong, semitransparent globules, of a yellowish or reddish hue, and smells like aniseed. It was once employed as an astringent and detergent.

SARCODES, Carneous.

SARCOËPIPOCE'LE, *Sarcepiocelle*, from *σαρξ*, *sarkos*, 'flesh,' *επιπλοον*, 'the epiploon,' and *κηλη*, 'a tumour.' Epiplocele, complicated with sarcoma or sarcocele.

SARCOËPIPLOM'PHALUS, *Sarcepiplomphalus*, *Sarcepiplomphalocelle*, from *σαρξ*, *sarkos*, 'flesh,' *επιπλοον*, 'the epiploon,' and *ομφαλος*, 'the navel.' Umbilical hernia, formed by scirrhus epiploon or complicated with sarcoma.

SARCOHYDROCELE, Hydro-sarcocele.

SARCOLEM'MA, *Sarcolemma*, from *σαρξ*, *sarkos*, 'flesh,' and *λεμμα*, 'a coat.' The sheath which surrounds the fibrils of muscle that form a fibre. It is quite distinct from the areolar membrane that binds the fibres into fasciculi.—Bowman. See Perymysium.

SARCOLOGIA, Myology, Sarcology.

SARCOLOGY, *Sarcologia*, from *σαρξ*, *sarkos*, 'flesh,' and *λογος*, 'a discourse.' The part of anatomy which treats of the soft parts. It comprises myology, angiology, neurology, and splanchnology.

SARCO'MA, *Emphy'ma Sarcoma*, *Sarcomatous tumour*, *Sarcosis*, *Porrus*, *Sarcophy'ia*, *Exsarcoma*, *Exsarcoma*, *Tumor carneus*, *Excrescentia carnea*, *Polypus carnosus*, from *σαρξ*, *sarkos*, 'flesh.' Any species of excrescence having a fleshy consistence.

SARCOMA, CARCINOMATOUS, Scirrhus.—s. Cercosis, see Cercosis.—s. Epulis, Epulis.—s. Gelatinous, see Tumour, fibro-cellular.—s. Mastoid, Mammary sarcoma.—s. Medullare, see Encephaloid.—s. Medullary, Encephaloid, Hematodes fungus.—s. Pulp, Hematodes fungus.—s. Scroti, Sarcocele.

SARCOMA, COMMON VAS'CLULAR or OR'GANIZED of Abernethy, *Emphy'ma Sarcoma Vasculosum*, is vascular throughout: texture simple: when bulky, napped on the surface with arborescent veins. Found over the body and limbs generally.

SARCOMATEUX, Sarcomatous.

SARCOMATODES, Sarcomatous.

SARCOMATOSUS, Sarcomatous.

SARCOM'ATOUS, *Sarcomatous*, *Sarcomatoides*, (F.) *Sarcomateux*. Belonging to, or having the characters of sarcoma.

SARCOMPHALUM, Sarcomphalus.

SARCOM'PHALUS, *Sarcophthalmus*, from *σαρξ*, *sarkos*, 'flesh,' and *ομφαλος*, 'the navel.' A scirrhus or fleshy tumour, developed at the navel.

SARCOMYCES, Sarcosponges.

SARCOPHAGUS, Carnivorous, Catheretic.

SARCOPHYIA, Excrescence, Sarcoma.

SARCOPSYLLA PENETRANS, *Chique*.

SARCOPTES, Acarus.

SARCOPTES, SCABIEL, see Psora.

SARCOPYO'DES, from *σαρξ*, *sarkos*, 'flesh,' *πυρ*, 'pus,' and *ειδος*, 'resemblance.' Having the appearance of pus mixed with flesh. An epithet given to certain excreted matters, and particularly to the sputa of those labouring under consumption.

SARCO'SIS, from *σαρξ*, *sarkos*, 'flesh.' The formation of flesh. Also, sarcoma.

SARCO'SIS BULBI, *Exophthal'mia fungosa* seu *sarcomat'ica*. A fleshy growth from the globe of the eye.

SARCOSIS UTERI, Metrauxis.

SARCOSPONG'US, *Sarcosponges*, from *σαρξ*, *sarkos*, 'flesh,' and *σπογγος*, 'fungus.' A fleshy fungus.

SARCOSTOSIS, Osteo-sarcoma.

SARCOTIC, Incarnans.

SARCOUS, from *σαρξ*, *sarkos*, 'flesh.' Of, or belonging to, flesh or muscle—as '*sarcous tissue*.'

SARCOUS EL'EMENTS, *Primitive particles*. The elementary particles, which, by their union, form the mass of muscular fibre.—Bowman.

SARDIASIS, Canine laugh, Rius sardonius.

SARDONIA HERBA, *Ranunculus sceleratus*.

SARDONIASIS, Canine laugh, Rius sardonius.

SARDONIUS RISUS, Canine laugh.

SARE, Essera.

SARACE'NIA, *Side-saddle plant*, *Flytrap*.

The root of *Saracenia flava* and *S. variolaris* is bitter and astringent, and has been used in South Carolina, as a domestic remedy in dyspepsia.

SARRASIN, *Polygonum fagopyrum*.

SARRÊTE, Trismus infantum.

SARRIETTE, *Satureia hortensis*—s. de Crête, *Satureia capitata*.

SARSA, *Smilax sarsaparilla*.

SARSAPARILLA, (S.) Zarza, 'a briar,' and *parilla*, 'a little vine.' *Menispermum Canadense*, *Smilax S.*

SARSAPARILLA, FALSE, *Aralia nudicaulis*.

SARSAPARIL'LA GERMAN'ICA, *Gramen major seu rubrum*, *Salsaparilla Germanica*, *Carex Arenaria*, (F.) *Laiche*, *L. des Sables*, *Salseparille d'Allemagne*, *Chiendent rouge*. Ord. Cyperaceæ. *Sex. Syst.* Monocelia Triandria. A plant, which grows plentifully on the sea-coasts of Europe. It has been recommended in some mucous affections of the trachea, in rheumatism, gout, &c.

SARSAPARILLA, INDIAN, *Hemidesmus Indicus*—s. Wild, *Aralia nudicaulis*.

SARTAPARILLA, *Smilax sarsaparilla*.

SARTO'RIOUS, from *sartor*, 'a tailor,' because used in crossing the legs. *Suto'rius*, *Longissimus Fem'oris*, *Fascia'tilis*, *F. Longus* seu *suto'rius*, (F.) *Rio-cresti-tibial*, *Rio-prætibial* (Ch.), *Couturier*. A muscle situate at the anterior part of the thigh. It is long, small, and flattened, and extends, obliquely, from the anterior and superior spine of the ilium to the superior and inner part of the tibia. This muscle can, also, bend the pelvis on the thigh, and conversely.

SARX, Flesh, Pulp.

SARZA, *Smilax sarsaparilla*.

SASSA GUM. A gum, occasionally carried into the European markets from the East, and answering exactly to Bruce's description of the product of a tree, which he calls *Sassa*. It has considerable resemblance to Gum Tragacanth.

SASSAFRAS, *Laurus sassafras*, *Nectandra cymbarum*—s. Medulla, see *Laurus sassafras*—s. Nut, see *Pichurim* beans—s. Radicis cortex, see *Laurus sassafras*.

SASSAFRAS SWAMP, *Magnolia glauca*.

SASSAFRAS TEA, Saloop.

SASSY BARK, A poisonous bark of a large tree—*Genus*, *Erythrophleum*, *E. Judicia'te*; *Nat. Ord.* Fabaceæ; *Red water tree* of Sierra Leone—growing in Western and Central Africa, used in an ordeal to which persons, suspected of witchcraft, secret murder, and other crimes, are subjected, to test their innocence or guilt. Hence it is called, in Congo, *ordeal bark*, and in Ashantee and Liberia, *doom bark*.

SAT'ELLITE, *Satelles*, genitive *Satellit'is*. That which guards. That which is placed near.

SATELLITE VEINS are those that accompany arteries.

SATHÉ, Penis.

SATIES, Satiety.

SATIETAS, Plesmone, Satiety.

SATIETY, *Sati'etas, Satu'r'itas, Sa'ties, Satura'tio, Plethos, Plesmonê, from satiare* (itself from *satis*, 'enough'), 'to give enough.' Disgust for food; commonly produced by repletion.

SATIN, Lunaria rediviva.

SATURANS, Absorbent (medicine.)

SATURATIO, Satiety, Saturation.

SATURA'TION, *Satura'tio, from saturare, saturatum, (satis, 'enough'), 'to satiate.'* The state of a compound in which its elements are combined in such proportions that a fresh quantity of either cannot be added without producing excess.

SATUREIA CAPITA'TA, from *Sarpos*, 'the Satyrs,' owing to its presumed power of exciting to venery. *Cil'iated Sa'vory, Thymus, T. sylvestris seu capita'tus seu Oret'icus, Cuni'la, (F.) Sarriette de Crète. Ord. Labiatae. Sex. Syst. Didynamia Gymnospermia. Possesses similar properties to thyme.*

SATUREIA HORTEN'SIS, *Satureia, S. sati'va, Cuni'la sativa Plin'ii, Thymbra, Summer Sa'vory, (F.) Sarriette.* An aromatic plant, cultivated for culinary purposes. It has been used, as an excitant, in anorexia, anaphrodisia, &c.

SATUREIA ORIGANOIDES, *Cunila mariana*—s. *Sativa*, s. *hortensis*.

SATURITAS, Satiety.

SATURNI DULCEDO, *Plumbi superacetas.*

SATURNINE, *Saturni'nus, from saturnus, lead.* Containing, or caused by, lead.

SATURNINE BREATH. The peculiar odour of the breath in one labouring under Saturnism.

SATURNISMUS, *Intoxica'tio seu Cachex'ia seu Dyscrasia saturni'na, Molybdocachex'ia, from saturnus, 'lead.'* Poisoning by lead; lead-poisoning; saturnine cachexy.

SATURNUS, Plumbum.

SATYRIASIS, from *Sarpos*, 'a Satyr,' because the Satyrs were greatly addicted to venery. *S. furens, Satyrias'mus, Satyris'mus, Priapismus, Salac'itas, Brachi'na, Araf'con, Lagne'sis Salac'itas, L. furor masculi'nus, Lagne'a, Lagni'a, Lagno'sis, Lasciv'ia, Lasciv'itas, Tenti'go vere'tri.* An irresistible desire, in man, to have frequent connexion with females, accompanied by the power of doing so without exhaustion. The causes are commonly obscure. Sometimes, the abuse of aphrodisiacs has occasioned it. The principal symptoms are:—almost constant erection; irresistible and almost insatiable desire for venery; frequent nocturnal pollutions. Cold lotions; the cold bath; a mild diet; active exercise, &c., are the only means that can be adopted for its removal.

SATYRIASIS FURENS, Satyriasis.

SATYRIASMUS, Satyriasis.

SATYRION, Orchis mascula.

SATYRISMUS, Satyriasis.

SATYRIUM, *Goodyera pubescens.*

SAURJA, Saoria.

SAUCE-ALONE, Alliaria.

SAUCE, GREEN, *Rumex scutatus.*

SAUCH, *Salix.*

SAUER KRAUT, (G.), 'sour cabbage,' (F.) *Chou-crôte.* A culinary preparation, made with chopped cabbage, left to attain an incipient state of acetous fermentation. It is reputed by the Germans to be salubrious, easy of digestion, and eminently antiscorbutic.

SAUGE, *Salvia*—s. *des Bois, Teucrium scor-dium*—s. *Grande, Salvia sclarea*—s. *Hormin, Salvia borminum*—s. *de Jérusalem, Pulmonaria*—s. *Petite, Salvia*—s. *Sclurée, Salvia sclarea.*

SAUGH, *Salix.*

SAULE, *Salix.*

SAUMURE, Muria.

SAUNDERS, RED, *Pterocarpus santalinus*—s. Tree, *Pterocarpus santalinus*—s. Yellow, *Santalum album.*

SAURA, Lizard.

SAURURUS CERNUUS, from *σαυρος*, 'a lizard,' and *οψα*, 'a tail.' *Lizard's Tail, Breast weed.* Indigenous; Order, *Saururacæ*; flowering in June. It has been used in lumbago. The bruised root is emollient, and is a popular application to inflamed breasts.

SAUSAGE POISON, Allantotoxicum.

SAUT, Leap.

SAUTEUSES, see *Convulsionnaire.*

SAUVE-VIE, *Asplenium ruta muraria.*

SAVE, Sage.

SAVEUR, Savour.

SAVINE, *Juniperus sabina.*

SAVINIE, *Juniperus sabina.*

SAVON, Sapo—s. *Ammoniacal, Linimentum ammoniac fortius*—s. *Cucac, Sapo, saponaceus*—s. *Calcaire, Linimentum aquæ calcis*—*Dur, Sapo durus*—s. *Médicinal, Sapo medicinalis*—s. *Mou, Sapo mollis*—s. *Noir, Sapo mollis.*

SAVONNIER COMMUN, *Sapindus saponaria.*

SAVONNIÈRE, Saponaria.

SAVONULE, Saponulus.

SAVONULUS, Saponulus.

SAVORY, CILIATED, *Satureia capitata*—s. Summer, *Satureia hortensis.*

SA'VOUR, *Sapor, Taste, (F.) Saveur.* Quality of bodies, by which they act upon the sense of taste. Chymists, at times, avail themselves of this quality to analyse an unknown substance.

SAVOUREUX, Savoury.

SAVOURY, *Sap'idus, (F.) Savoureux.* An epithet given to bodies that have taste; and particularly to those that excite a very agreeable impression on the organ of taste.

SAVOYAN, *Galium verum.*

SAW, *Prion, Serra, (Da.) Saxe, (Sax.) raga, (F.) Scie.* A surgical instrument, made like a common saw; and used for sawing the bones in amputations, or for removing exostoses, &c. Saws of different sizes and shapes are used in surgery. The crown of the trepan is nothing more than a species of circular saw.

SAW, AMPUTA'TION, *Serra amputato'ria.* The saw used in amputation.

SAW, HEY'S. An instrument used by Mr. Hey, of Leeds, in fractures of the cranium. It consists of a long handle, to which a small saw with a convex or straight edge is attached, and by which a piece of bone of any shape may be removed. With the trephine, the saw must always be circular.

SAW PALMETTO, *Chamærops serratula*—s. Wort, *Liatris.*

SAXIFRAGA, from *saxum*, 'a stone,' and *frango*, 'I break.' *Pimpinella saxifraga*—s. Anglica, *Peucedanum silaus.*

SAXIFRAGA CRASSIFO'LIA. *Ord. Saxifragacæ. Sex. Syst. Decandria Digynia.* The root of this species of saxifrage has been extolled by Pallas, as pre-eminently antiseptic.

SAXIFRAGA GRANULA'TA, *S. alba, Sanic'ula, Sedum, White sax'ifrage, (F.) Saxifrage granulée.* Recommended as diuretic and lithontriptic; but not used.

SAXIFRAGA RUBRA, *Spiræa filipendula*—s. *Vulgaris, Peucedanum silaus.*

SAXIFRAGE, BURNET, *Pimpinella saxifraga*—s. *Granulée, Saxifraga granulata*—s. Meadow, *Peucedanum silaus*—s. White, *Saxifraga granulata.*

SAXIFRAGUS, Lithontriptic.

SAXON'ICUS PULVIS. A powder, prepared with the roots of the wild and cultivated ange-

lica: the asclepias vincetoxicum, &c. It was considered formerly as an antidote.

SCAB, *Scab'ies*, from *scabere*, 'to dig.' *Scall*, (Prov.) *Rove*. An incrustation, which forms upon a sore, owing to the concretion of the fluid discharged from it. An eschar.

SCABBY or SCABBED. Covered or diseased with scabs.

SCABIES, Psora.—s. Agria, Lichen—s. Capitis, Porrigo lupinosa, Porrigo scutulata—s. Capitis favosa, Porrigo favosa—s. Crustosa, Radzyge—s. Fera, Ecthyma.

SCABIES FERINA, *Itch of animals, Mange*, (pronounced *Maynge*.) A cutaneous disease, which affects almost all domestic animals, but especially the horse, sheep, dog, and cow. It is said to have been transmitted to man; but this is questionable.

SCABIES FERINA, Psoriasis—s. Norvegica Boeckii, Radzyge—s. Papuliformis, Prurigo—s. Sicca, Lichen, Psoriasis—s. Venerea contagiosa, Scherlievo.

SCABIEUSE ORDINAIRE, Scabiosa.

SCABIOLA, Scabies.

SCABIOSA, *S. arven'sis*, *Tri'chera arven'sis*, *Field Scabious*, (Sc.) *Eastning wort*, (F.) *Scabieuse ordinaire*, Ord. Dipsacæ. Ser. Syst. Tetrandria Monogynia. From *scaber*, 'rough,' owing to its rough hairy surface.' The plant is bitter and sub-astringent, and was formerly used in the cure of leprous affections and diseased lungs.

SCABIOSA ARVENSIS, Scabiosa—s. Carduifolia, Echinos.

SCABIOSA SUCCI'SA, *Succisa*, *S. pratensis*, *Asteroccephalus succisa*, *Morsus Diab'oli*, *Devil's bit*, (F.) *Mors du Diable*, has similar properties.

SCABIOUS, Erigeron Philadelphicum—s. Field, Scabiosa—s. Sweet, Erigeron heterophyllum.

SCABISH, *Oenothera biennis*.

SCABRITIES, Lichen, Trachoma.

SCALA, 'a ladder.' A machine formerly used for reducing dislocations of the humerus.

SCALE OF THE COCH'LEA, *Gyri seu Ductus spirales seu Canales cochleæ*, (F.) *Rampes ou Échelles du Limaçon*—are the cavities in the cochlea, separated from each other by the spiral septum. The one—*Scala interna seu posterior seu inferior cochleæ*, *S. tym'pani*—would communicate with the cavitas tympani by the fenestra rotunda, were it not for the membrane that closes it. The other—*Scala externa seu anterior seu superior cochleæ*, *S. vestibuli*—opens into the cavity of the vestibule. The scales communicate with each other by an opening at the top of the partition.

SCALE ANTERIOR COCHLEÆ, see *Scala*—s. Clausa, see *Scala*—s. Externa cochleæ, see *Scala*—s. Inferior cochleæ, see *Scala*—s. Interior cochleæ, see *Scala*—s. Posterior cochleæ, see *Scala*—s. Superior cochleæ, see *Scala*—s. Tympani, see *Cochlea*, and *Scala*—s. Vestibuli, see *Cochlea*, and *Scala*.

SCALDRICKS, *Sinapis arvensis*.

SCALE, Sax. *rcale*, from *reylan*, 'to divide, separate'; (G.) *Schale*, 'shell, peel'; *schälen*, 'to shell, to peel.' *Squama*, *Lepis*, (F.) *Escaille*, *Écaille*. An opaque and thickened lamina of cuticle, commonly produced by some degree of inflammation of the true skin, over which it is formed.

SCALE, DRY, Psoriasis.

SCALÈNE ANTÉRIEUR, *Scalenus anticus*—s. *Postérieur*, *Scalenus posticus*.

SCALE'NUS, *σκαληνός*, 'irregular or unequal.' Geometers employ this word to designate a triangle whose three sides are unequal. Anatomists

have given the name to two muscles. 1. *SCALENUS ANTI'CUS*, *Portion of the Costo-trachelian* (Ch.), (F.) *Scalène antérieur*, is situate at the internal and inferior parts of the neck. It is long and triangular; and is inserted, below, at the upper surface of the first rib; and, above, at the anterior tubercle of the transverse processes of the 3d, 4th, 5th, and 6th cervical vertebæ. This muscle bends the neck, and draws it to its side. It can also raise the first rib. 2. *SCALENUS POSTI'CUS*, (F.) *Scalène postérieur*, *Portion of the Costo-trachelian* (Ch.), is situate behind the last. It is, also, long and triangular; but more bulky than the *S. anticus*. It is attached, below, to the outer surface of the first two ribs; and terminates, above, at the summit of the transverse processes of the last six vertebæ of the neck. This muscle bends the neck laterally, and can elevate the first two ribs. Some anatomists, as Albinus and Sömmering, have described five scaleni muscles on each side, and have called them *Scalenus prior*, *S. minimus*, *S. lateralis*, *S. medius*, *S. posticus*. Riolan and Chaussier only describe one scalenus muscle on each side. Cowper and Douglas, and the generality of the English anatomists, admit three, *S. anticus*, *medius*, and *posticus*; or, *primus*, *secundus*, and *tertius*; Winslow, Boyer, and many of the French anatomists,—two, as above described.

SCALING—same etymon as *Scale*—in dentistry, is an operation, which consists in removing the tartar from the teeth.

SCALL, ((Sc.) *Skaw*.) *Impetigo*, *Scab*—s. *Asbestos*, see *Eczema* of the hairy scalp—s. *Crusted*, *Impetigo*—s. *Dry*, *Psoriasis*—s. *Honeycomb*, *Porrigo favosa*—s. *Humid*, *Eczema*—s. *Milky*, *Porrigo larvalis*—s. *Papulous*, *Ecthyma*—s. *Running*, *Eczema*, *Impetigo*—s. *Vesicular*, *Eczema capitis*—s. *Washerwoman's*, *Psoriasis diffusa*.

SCALLARD, *Porrigo*.

SCALLED HEAD, *Porrigo*, *P. scutulata*.

SCALP, Teut. *Schelp*, 'a shell.' (Old Eng.) *Scop*. The integuments covering the head. The hairy scalp is called, also, *Capillit'ium*, *Trichot'on*, *Cæsarices*, (F.) *Chevelure*, *Cuir chevelu*.

SCALPEL, *Scalpel'um*, *Scalpel'lus*, *Culter*, *Incis'orium*, *Smi'leon*; from *scalpo*, 'I cut.' A cutting instrument, formed of a blade of well-tempered steel, very sharp, of variable shape and size, and fixed solidly in a handle. Its use is to divide the soft parts in operations, dissection, &c.

SCALPELLUM, *Scalpel*—s. *Lenticulare*, *Phacopis*.

SCALPELLUS, *Bistouri*, *Scalpel*.

SCALPEUM, *Bistouri*.

SCALPRUM, *Raspatorium*—s. *Chirurgicum*, *Lancet*.

SCALPRUM DENTA'RIUM, *Lima denta'ria*. A tooth-file. A dental file or file used by the dentist.

SCALPRUM RASORIUM, *Raspatorium*.

SCALY, *Squamous*.

SCALY DISEASES OF SKIN, *Lepido'ses*, form the second order in Willan and Bateman's system of cutaneous diseases.

SCAMMA, *Fossa*.

SCAMMONÉE D'ALEP, *Convolvulus scammonia*.

SCAMMONIUM, see *Convolvulus scammonia*—s. *Syriacum*, *Convolvulus scammonia*.

SCAMMONY, *Convolvulus scammonia*.

SCAMMONT, *Bourbon*. The concrete juice of *Periploca Mauritiana*.

SCAMMONT, *GERMANY*. The concrete juice of *Convolvulus sepium*.

SCAMMONT OF MONTPELLIER, *Cynanchum Monspeliacum*.

SCAMNUM HIPPOCRATICUM, (*scamnum*).

'a bench.') *Bathron Hippocrati* on of Galen, (F.) *Banc d'Hippocrate*. A machine invented by Hippocrates for reducing fractures. It was a sort of bed, six feet long, on which the patient was fixed. Straps were attached above and below the fracture or luxation, and extension and counter-extension were effected by a winch. It is described in Galen, Oribasius, Scultetus, &c. See *Bathron*.

SCANDIX, *S. cerefolium*.

SCANDIX BULBOCASTANUM, *Bunium bulbocastanum*.

SCANDIX CEREOFOLIUM, *Scandix*, *Scandyx*, *Pecten Ven'eri*s, *Cerefolium*, *Daucus Seprin'ius*, *Chærophyllum*, *Ch. Sati'vum* seu *Cerefo'lium*, *Anthriscus cerefolium*, *Chærefo'lium*, *Chervil*, (F.) *Cerfeuil*. *Ord. Umbelliferae*. *Sex. Syst.* *Pentandria Dignia*. A culinary herb, which is slightly aromatic and grateful. It is said to be gently aperient and diuretic.

SCANDIX ODORATA, *Sweet Cic'ely*. Properties the same as the other. Also, *Chærophyllum odoratum*.

SCANDULARIUS MUSCULUS, *Parathenar*, *Transversus pedis*.

SCANDYX, *Scandix cerefolium*.

SCAPHA, *Scaphus*, 'a skiff or cock-boat,' from σκαπτόν, 'I make hollow.' The excavation or cavity of the external ear, between the helix and anthelix. The meatus auditorius externus. The rima vulvæ. Also the name of a bandage for the head, mentioned by Galen, and called, likewise, *Tholus Dinele'us*.

SCAPHIA, *Nates*.

SCAPHION, *Cranium*, *Papaver* (Capsule).

SCAPHIUM OCULARE, *Pelvis Ocularis*, *Eye-glass*, (F.) *Bassin ou Gondole oculaire*, *Œillière*. A small porcelain, glass or metallic vessel, used for applying lotions to the eye.

SCAPHO-CARPO-SUPER-PHALANGIUS POLLICIS, *Abductor pollicis brevis*.

SCAPHOID, *Scapho'i'des*, from σκαφή, 'a skiff,' and εἶδος, 'form.' A name given to several parts.

SCAPHOID FOSSA, *Fossa Scapho'i'des*, (F.) *Enfoncement Scaphoïde*. A small cavity at the upper part of the internal ala of the pterygoid process, in which the peristaphylinus internus is attached.

SCAPHOID BONE OF THE HAND, *Os Scaphoïdes Manûs* seu *navicula'rè* seu *cymbifor'mè*, *Boat-like bone*, (F.) *Scaphoïde de la main*. The first bone of the first row of the carpus. The upper surface is convex, and articulated with the radius. The under and outer surfaces are also convex,—to be articulated with the trapezium and trapezoids. Between the upper and under cartilaginous surfaces there is a rough fossa for the insertion of the capsular ligament. The anterior and inner surface has an oval cavity, where it is articulated with the os magnum, which gives name to the bone. There is a process on the outer end of the bone, for the attachment of part of the anterior transverse ligament of the wrist.

SCAPHOID BONE OF THE FOOT, *Os Scaphoïdes Pedis* seu *navicula'rè* seu *cymbifor'mè*, *Boat-like bone*, (F.) *Scaphoïde du Pied*. This bone is situate at the forepart of the astragalus and inner part of the foot. The posterior surface forms a cavity, somewhat like that of a boat, for receiving the head of the astragalus. There is a prominence at the inner side of the bone, for the insertion of tendons, muscles, and strong ligaments. The forepart of the bone is convex, and divided into three articular surfaces for articulation with the ossa cuneiformia. Between the os naviculare and astragalus, the foot has its principal lateral and rotary motions; although each of the other joints of the tarsus contributes a little. Part of the tibialis posticus is inserted into it.

SCAPHOÏDE DE LA MAIN, *Os scaphoïdes manûs*—*s. du Pied*, *Os scaphoïdes pedis*.

SCAPHOÏDO-ASTRAG'ALAN, *Scaphoïdo-astragala'nus*. That which relates to the scaphoid bone and astragalus. The articulation of these bones is so called; and the ligament which strengthens it is termed the *Scaphoïdo-astragolan ligament*.

SCAPHOÏDO-CUBOID, *Scaphoïdo-cuboidæ'us*. That which relates to the scaphoid and cuboid bones. The articulation of those bones is so called.

SCAPHULA, *Navicularis fossa*.

SCAPHUS, *Auditory canal*, *external*, *Rima vulvæ*, *Scapha*.

SCAP'ULA, *Scap'ulum*, *Plata*, *Omopla'ta*, *Homopla'ta*, *Scop'ula*, *Epino'tion*, *Latit'udo hu'meri*, *Omop'latè*, *Os latum hu'meri*, *Scutum thorac'is* seu *cordis*, *Spath'ula*, *Pterygium*, *Chelonium*, *Spat'ula*; *Blade-bone*, *Shoulder-blade*, (Prov.) *Shield-bone*, *Plate-bone*, *Spade-bone*, (F.) *Omo-pla'te*. An irregular, broad, flat bone, of a triangular shape, situate at the posterior part of the shoulder. It has, 1. A posterior or dorsal surface or *Dorsum*, *Testu'do scap'u'læ*, divided transversely into two parts, by a triangular process, called *Spine of the Scapula*; which terminates by a considerable eminence, called *Acromion*. Above the spine, is the *Fossa supra-spinata*; below it, the *F. infra-spinata*. 2. An anterior or costal surface or *venter*, forming the *Fossa subscapularis*. 3. A superior or cervical edge, which is thin, and terminates, anteriorly, by a strong, curved apophysis, called the *coracoid process*. 4. A posterior or vertebral edge, called, also, the *base*. 5. An outer or axillary or inferior edge, having a thick, truncated angle, in which is the *glenoid cavity*. With this is articulated the head of the humerus. The glenoid cavity is separated from the body of the bone by a narrow part called the *cervix* or *neck*. The edges or margins are, also, called *costæ*. The scapula has three angles—an inferior, superior, and anterior.

SCAPULA, *Humerus*.

SCAPULÆ NASI, *Partes laterales nasi*. The lateral parts of the nose.

SCAPULAL'GIA; from *scapula*, 'the shoulder-blade,' and *algos*, 'pain.' Arthralgia of the shoulder-joint.

SCAP'ULAR, *Scap'ulary*, *Scapula'ris*, (F.) *Scapulaire*, from *scapula*, 'the shoulder-blade.' That which relates or belongs to the scapula.

SCAPULAR APONEURO'SIS. A broad, thin aponeurosis, with decussating fibres; which is attached, above, to the spine of the scapula, below, to a crest between the *teres major* and *infra-spinatus*; within, to the spinal edge of the scapula; and which, at its middle, unites with the thin aponeurosis that covers a part of the external surface of the deltoid.

SCAPULAR ARTERIES. These are several. a. The *Superior scapular*, *Superficial scapular* of Sömmering, *Dorsal'is scapulae superior*, *Transversa'lis scapulae*, *T. Hu'meri*. This arises from the subclavian; and often from the inferior thyroid, or from a trunk common to it and the posterior scapular or transverse scapular or cervical. It passes behind the clavical, above the superior edge of the scapula, and buries itself between the dorsal surface of that bone and the muscles inserted into it. b. The *inferior scapular* or *common scapular* or *internal scapular* or *subscapular artery*, *A. Thorac'ica axilla'ris* seu *ala'ris* is of considerable size. It arises from the inferior part of the axillary artery, behind the brachial plexus—gives off several branches to the armpit, and divides into two branches:—the one, *Scapula'ris*

interna, and the smaller of the two, is distributed, almost entirely, to the serratus major anticus, and latissimus dorsi; the other, the larger of the two, the *Dorsalis Scapulae inferior* seu *Circumflexa Scapulae*, which distributes its branches to the subscapular and supra-spinal, and infra-spinal regions.

SCAPULAR NOTCH, see Notch, scapular.

SCAPULAR VEINS follow the same course as the arteries, which they accompany.

SCAPULARE, Scapulary.

SCAP'ULARY, *Scapula'rē*, *Fascia Scapula'ris*, from *scapula*. A broad, linen bandage, divided into two tails for three quarters of its length, the undivided extremity of which is attached to the posterior and middle part of a bandage carried round the body; the tails passing over the shoulders; and being attached to the anterior part of the body bandage, to prevent it from slipping down.

SCAPULODY'NIA, from *scapula*, and *ὄνν*, 'pain.' A hybrid word, signifying pain or rheumatism of the shoulder.

SCAP'ULO-HU'MERAL. An epithet given to the shoulder-joint. It is an orbicular articulation, resulting from the reception of the head of the humerus into the glenoid cavity of the scapula. To keep the bones in connexion, there are: 1. A very strong, conoidal capsular ligament; passing from the circumference of the glenoid cavity around the neck of the humerus. 2. A *fibro-cartilaginous ring*, called the glenoid ligament, and a synovial capsule.

SCAPULO-HUMÉRAL, *Teres major*—*s. Huméro-olécrani*, *Triceps extensor cubiti*—*s. Hyoidien*, *Omo-hyoideus*—*s. Radial*, *Biceps flexor cubiti*—*s. Trochitérien*, *grand*, *Infra-spinatus*.

SCAPUS, Penis, see Hair.

SCAR, Cicatrix.

SCAR, *Seam*. To mark with a cicatrix or seam;—as *scarred*, (*Sc.* *arred*, means marked with a cicatrix or seam.

SCARABÆOLUS HEMISPHERICUS, *Coccus* cacti.

SCAR'BOROUGH, MINERAL WATERS OF. *Scarborough Spa*. A celebrated English chalybeate, at Scarborough, in Yorkshire. One of the springs is a simple, carbonated chalybeate, like that of Tunbridge: the other has an admixture of purging salt. Scarborough, having the additional advantage of sea-bathing, is much frequented by invalids.

SCARCHT, Hermaphrodite.

SCARDAMYG'MUS, *Scardamys's*, (*F.*) *Clignement*, from *σκαρδαμυσσιν*, 'to wink.' Winking.

SCARFSKIN, Epidermis.

SCARIFICA'TION, *Scarifica'tio*, *Amyx'is*, *Schasis*, *Catacas'mus*, *Catuchas'mus*, *Amyg'mus*, *Schasmus*, *Encharas'in*, from *σκαρφαομαι*, 'to make a slight scratch.' The act of scarifying. Also, a small incision, *Amyg'hē*, made into the skin with a lancet, bistoury, or scarificator, for different therapeutical purposes,—as to draw blood, or to discharge some effused fluid. When the scarifications are very superficial, and do not go farther than the skin, they are termed by the French, *Mouchetures*.

SCARIFICA'TOR, *Scarificato'rium*, *Schaste'rium*. Same etymon. An instrument for making scarifications. It consists of a small brass box, of a cubical shape, in which 10 or 12 lancet points turn upon a common pivot. By means of a spring, all these blades can be made to issue at once, and make as many scarifications.

SCARIFICATORIUM, Scarificator.

SCARIOLA, *Cichorium endivia*, *Lactuca scariola*.

SCARIOLE, *Cichorum endivia*.

SCARLATA, *Scarlatina*.

SCARLATINA, from (*I.*) *scarlatte*, 'a deep red.' *Scarlatia*, *Eranthis'sia Rosalia*, *Rosalia*, *Rossalia*, *R. squamo'sa*, *Rosalia*, *Rube'ola*, of some, *Morbilli ignei* seu *confluentes*, *Morbis scarlatinosus*, *Febris scarlatinos'a seu rubri*, *Gutturis morbus epidemicus Foresti*, *Pur'pura*, *Typhus scarlatinus*, *Porphyrismus*, *Porphyrisma*, *Scarlet* or *Rash-fever*, (*F.*) *Fievre rouge*, *F. pourprée*. The characteristic symptoms of scarlatina are:—a scarlet flush, appearing about the second day of fever on the face, neck, and fauces; and progressively spreading over the body; terminating about the 7th day. Two great varieties may be reckoned:—the *S. simplex* seu *febris seu benigna* seu *sine angina*, in which the fever is moderate and terminates with the efflorescence; prostration of strength being trifling, and the contagious property slight;—and the *S. Angino'sa* seu *Parithmitica* seu *Cynanchica* seu *mitior*, in which the fever is severe; the throat ulcerated; the eruption later in its appearance, and less extensive; often changing to a livid hue; the fever being highly contagious. *Scarlatina maligna* seu *gravior*, has been reckoned a variety of this, in its worst degree. It agrees with the *Cynanche maligna* of Cullen.

Scarlatina belongs to the *Major exanthemata*, and is a disease, chiefly, of children. The eruption differs from that of measles, in being an efflorescence not raised above the cuticle. Measles, too, is attended with catarrhal symptoms, whilst the complication, in scarlatina, is *cynanche*. The treatment of simple scarlatina need not be much. It must be antiphlogistic. If the throat be very much ulcerated, acid gargles and counter-irritants must be employed; and if the affection become manifestly typhoid, and the sore throat of a malignant character, the case must be treated like typhus gravior; with antiseptic gargles of bark, acid, &c. Anasarca sometimes supervenes on scarlatina, and requires attention. Purgatives are here demanded as well as the use of sorbefeccients, such as mercury and squill, &c.

SCARLATINA ANGINOSA, see *Scarlatina*—*s. Benigna*, see *Scarlatina*—*s. Cynanchica*, see *Scarlatina*—*s. Febris*, see *Scarlatina*—*s. Gravior*, see *Scarlatina*—*s. Miliaris*, *Rubeola*—*s. Mitior*, see *Scarlatina*—*s. Parithmitica*, see *Scarlatina*—*s. Pustulosa*, *Rubeola*—*s. Rheumatica*, *Dengue*—*s. Sine Angina*, see *Scarlatina*—*s. Urticata*, *Urticaria*.

SCARLATINOUS, *Scarlati'nal*, *Scarlatino'sus*. Relating or appertaining to scarlatina.

SCARLET BERRY, *Solanum dulcamara*.

SCARLET FEVER, *Scarlatina*.

SCATACRASIA, *Scoracrasia*.

SCATACRATIA, *Scoracrasia*.

SCAW, *Sambucus*.

SCEAU DE SALOMON, *Convallaria polygonatum*.

SCELAL'GIA, from *σκελος*, 'the leg,' and *αλγος*, 'pain.' Pain of the leg.

SCELALGIA ANTICA, *Neuralgia femoro-pretibialis*—*s. Postica*, *Neuralgia femoro-poplitea*—*s. Puerperarum*, *Phlegmatia dolens*.

SCELETEU'SIS. Mummification. Also, the mode of making a skeleton, *Skeletope'a*, *Syntar'is*.

SCELETOLOGIA, *Skeletology*.

SCELETON, *Skeleton*.

SCELETOPEA, *Sceleteusis*.

SCELETUM, *Skeleton*—*s. Artificiale*, see *Skeleton*—*s. Naturale*, see *Skeleton*.

SCELONCUS, *Phlegmatia dolens*.

SCELORRHEUMA, *Rheumatismus cruris*, from *σκελος*, 'the leg,' and *ρευμα*, 'defluxion, rheumatism.' Rheumatism of the leg.

SCELOS, Leg.

SCELOTYRBE, Chorea, Porphyræ nautica —
 Febrilis, see Subsaltus tendinum — s. Festinans,
 Paralysis agitans — s. Tarantismus, Chorea.

SCEPARNON, Amnios, Ascia.

SCEPASTERIUM, Couvreechef.

SCIPASTRUM, Couvreechef.

SCIPTUM, Penis.

SCHASIS, Scarification.

SCHASMUS, Scarification.

SCHASTERION, Fleam, Lancet, Scarificator.

SCHIBA ARABUM, Artemisia santonica.

SCHEDULA, Prescription.

SCHERBET', *Sherbet', Serbet', Tacherbet', Sorbet'tum, Sorbet'tum, Sorbet'tum*. A drink prepared by the Turks, Persians, Egyptians, &c., with sugar and the juice of acid fruits.

SCHERLIEVO, *Skerljevo, Male di Scherlievo, Male di Breno, Male di Fiume, Male di Scarlievo, Male scabioso venereo, Margaretizza, Morbus Croatus seu fluminien'sis, Frambe'cia Illy'ica, Scabitis venerea contagio'sa, Venereal Itch*. A name given to a particular form of syphilis, observed in Illyria. It is said to be capable of being communicated without immediate contact; and, that its principal symptoms are: — pains in the bones, ulceration of the fauces, pustules, and fungous growths in different parts of the body.

Affections, known by this name, as well as by those of *Male di Breno, Falcadi'na, Boala and Frenga*, prevail in the frontier countries of the Austrian empire, and are generally regarded as contagious, of syphilitic origin, and as having been introduced from a surrounding country. The terms appear, however, in the course of time, to have been applied to a variety of ulcers, eruptions, and other morbid conditions in no way connected with syphilis, so that they lost their peculiar signification. — Sigmund.

SCHERO'MA. A dryness of the eye from the want of lachrymal secretion. — Dictionaries.

SCHIAS, Neuralgia femoro-poplitea.

SCHIATICUS, Sciatic.

SCHIDAKE'DON, *Schidace'dum, Schindae'ladum, Schindal'amus, Schindal'mus*, from *σχιζω*, 'I cleave.' A longitudinal fracture of a bone. — Galen.

SCHINDACE'DUM, Schidakedon.

SCHINDALAMUS, Schidakedon, Schindylesis.

SCHINDALESIS, Schindylesis.

SCHINDALMUS, Schidakedon, Schindylesis.

SCHINDYLE'SIS, *Schindale'is, Schindal'amus, Schindal'mus*. An articulation of bones — by furrowing, as it were; as in that of the vomer, which constitutes, in part, the septum narium. Also, the act of splitting into small pieces; from *σχιζω*, 'I split into small pieces.' — Hippocrates.

SCHINELÆ'ON, from *σχινος*, 'the mastic tree,' and *ελαιον*, 'oil,' *Oleum lentis'cinum*. Oil of mastic. — Dioscorides.

SCHINOCEPHALUS, Scillocephalus.

SCHINOS, Scilla.

SCHINZNACH or HAPSBURG, MINERAL WATERS OF. These springs are situate three leagues from Baden, and are amongst the most celebrated bathing-places in Switzerland. The waters are of the saline sulphurous kind; and have a high reputation in cutaneous and rheumatic affections, visceral obstructions, and glandular enlargements. They contain sulphate of lime, sulphate of soda, chloride of sodium, chloride of magnesium, sulphate of magnesia, sulphate of lime, carbonate of magnesia, and oxide of iron; sulphohydric acid, and carbonic acid gases.

SCHISMA, Fissure, Rima.

SCHISTOCEPHALUS, from *σχιστος*, 'cleft,

separated,' and *κεφαλη*, 'head.' A monster having some part of the head cleft or fissured. — Gurlt.

SCHISTOCOR'MUS, from *σχιστος*, 'cleft, separated,' and *κορπος*, 'trunk.' A monster having some part of the trunk cleft or fissured. — Gurlt.

SCHISTOM'ELUS, from *σχιστος*, 'cleft, separated,' and *μελος*, 'a limb.' A monster whose limbs are fissured.

SCHISTORRHACHIS, Hydrorachis.

SCHISTOSO'MUS, from *σχιστος*, 'cleft,' and *σωμα*, 'body.' A malformation of the abdomen, in which the fissure and evagination extend over the entire length of the abdomen; the lower extremities being absent, or very little developed, so that the body appears as if truncated inferiorly. — Vogel.

SCHIZA, Rima, Rima vulvæ.

SCHIZATRICH'IA, *Trichos'chia, Schizotrich'ia*, from *σχιζω*, 'I split,' and *τριχis, τριχος*, 'a hair.' The splitting of the hairs at their extremities. — J. P. Frank.

SCHLANGENBAD, MINERAL WATERS OF. Schlangenbad is a much-frequented watering-place in Germany. The waters contain a little carbonic acid, and carbonate of soda. Temperature 86° Fahrenheit.

SCHNAPPS, SCHIEDAM, see Gin.

SCHNEIDERIAN MEMBRANE, Pituitary membrane.

SCHENANTHUS, *Juncus odoratus*.

SCHOLA, School.

SCHOLLERA OXYCOCCOS, *Vaccinium oxycoccos*.

SCHOOL, *Schola*, (F.) *École*. A place of education. In Europe the large Medical Schools are usually attached to the universities. In England, however, there has been no medical school of celebrity at either of the universities of Oxford or Cambridge; owing greatly to their provincial situation. Of late, two schools have existed in London, attached to the London University — University College and King's College. Excellent private schools have, however, long existed in that metropolis. The medical schools of Europe which have been most celebrated, are those of Edinburgh, Leyden, Berlin, Halle, Tübingen, Paris, Montpellier, Bologna, Padua, Pavia, and Pisa. In the United States, the medical schools are numerous; at this time not fewer, perhaps, than 40. Those most numerously attended, are Jefferson Medical College of Philadelphia, the medical department of the University of Pennsylvania, and the University of Nashville.

SCHOOLEY'S MOUNTAIN, MINERAL WATERS OF. The waters of this New Jersey Spring are a carbonated chalybeate. According to Dr. McNevin, they contain extractive, chloride of sodium, chloride of calcium, chloride of magnesium, carbonate of lime, sulphate of lime, carbonate of magnesia, silica, and carbonate, and oxide of iron.

SCHORBUS, Purpura.

SCHOUSBÆA COCCINEA, *Cacocucia coccinea*.

SCHWALBACH or LANGEN-SCHWALBACH, MINERAL WATERS OF. Schwalbach is a town of Germany, seated on the Aa, 32 miles west of Frankfurt. The waters contain chloride of sodium, carbonates of lime and magnesia, iron, sulphate of lime, carbonic acid, oxygen, &c. Employed as a tonic.

SCHWANN, WHITE SUBSTANCE OF, see Substance, white, of Schwann.

SCHWENDECK, MINERAL WATERS OF. These waters are about 5 leagues from Munich, in Bavaria. They contain carbonic acid; carbonate and sulphate of lime, chloride of calcium, chloride of magnesium; carbonate of soda, and

oxide of iron, and are much used for cutaneous affections.

SCIATIC, *Sciaticus*, *Ischiatic*, *Ischiaticus*, *Schiaticus*, (F.) *Sciaticque*, from *ισχίον*, 'the haunch.' A word formed by contraction from *ischiatic*, which latter is still used. See *Ischiatic*.

SCIATIC NERVE, *Great*, *Nervus ischiadicus*, *Great fem'oro-poplite'al* (Ch.), (F.) *Grand nerf sciaticque*, ought to be esteemed a continuation of the sacral plexus. It issues from the pelvis between the pyramidalis and superior gemellus; descends along the posterior part of the thigh; and after having given branches to the gemini, quadratus, obturator externus, glutæus maximus, semi-membranosus, semi-tendinosus, biceps and third abductor muscles, it divides, about the inferior third of the thigh, into two branches; the one, named the *external popliteal*; the other, the *tuternal*.

SCIATIC NERVE, *Lesser*, *Inferior glut'eal branch of the sacral plexus*, *Lesser fem'oro-poplite'al nerve* (Ch.), (F.) *Nerf petit sciaticque*, which seems especially formed by the 2d and 3d sacral, is given off from the lower part of the plexus, and escapes from the pelvis beneath the pyramidalis muscle. It gives branches — *nervi cut'nei clunium inferio'ies* — to the glutæus maximus; a *sciatic branch or infra-pelvic cutaneous or perineal cutaneous*, *Pudenda'lis longus infe'rior*, which passes under the tuberosity of the ischium, and furnishes filaments to the glutæus maximus, and to the muscles of the perineum, urethra, penis, scrotum, &c.; and a *posterior femoral cutaneous branch* — *middle posterior cutaneous nerve* — the filaments of which traverse the femoral aponeurosis, to be distributed on the skin of the posterior part of the thigh. The principal branch of the lesser sciatic nerve divides, near the ham, into two or three filaments, which descend under the integuments almost as far as the inferior part of the leg.

SCIATIC NOTCH, *Great*, *Sciatic Fora'men*, *Incisura ischiad'ica major*, (F.) *Grande échancrure sciaticque ou grand trou sciaticque*, is a large notch at the posterior edge of each os innominatum, below the posterior and inferior iliac spine, which is converted into a foramen by the sacrum and sacro-sciatic ligaments. The *lesser sciatic notch*, *Incisura ischiad'ica minor*, *Luna Albi'ni*, (F.) *Petite échancrure ischiaticque*, is much smaller than the last, from which it is separated by the sciatic spine. It gives passage to the tendon of the obturator internus, and to the internal pudic vessels and nerves.

SCIATIC REGION, *Ischiadic region*.

SCIATIC SPINE, *Spine of the Ischium*, *Spinous Proc'ess of the Ischium*, is a short, flat, pyramidal eminence on the os innominatum, above the preceding foramen; which gives insertion to the lesser sacro-sciatic ligament.

SCIATICA, *Neuralgia femoro-poplitea*.

SCIATIQUE, *Neuralgia femoro-poplitea*, *Sciatic*.

SCIE, *Saw*.

SCIENTIA HERMETICA, *Alchymy* — s. *Medendi*, *Medicina* — s. *Medica*, *Medicina*.

SCIEROPIA, from *σκῆπος*, 'shady,' and *ὤψ*, 'the eye.' A defect of vision, in which all objects have a darker hue than natural.

SCILLA, *σκίλλα*, from *σχιζω*, 'I cleave, I split' [?]. *Scilla seu Squilla marit'ima*, *Scylla*, *Aschil*, *Schinos*, *Oculus Typho'nis*, *Ornithog'olum marit'imum seu squill'a*, *Stella'ris seu Urgan'ea scilla*, *Cepa mari'na scilla*, *Squilla*, *Squill* or *sea onion*, *Eye of Typhon*, (F.) *Ognon marin*. *Ord. Liliacæ*. *Sex. Syst.* *Hexandria Monogynia*. Native of Spain, Austria, &c. The bulb or root of the squill has a bitter, nauseous taste, and is extremely acrid; inflaming the skin when

rubbed on it. Its acrimony, on which its virtues depend, is destroyed by heat, drying, and keeping. It is extracted by vinegar, spirit, and water. In large doses, squill is emetic and purgative; in small doses, diuretic and expectorant. Its active principle has been called *scillitine*. Dose, gr. j to v of the dried root, united or not with mercury.

SCILLA ESCULENTA, *Phalan'gium esculentum*, *Eastern Quamash*, *Wild Hy'acinth*; indigenous; flowering in May. An article of diet amongst the Western Indians. It is made into bread and poultices for inflamed breasts.

SCILLITIC, *Scillit'icus*, (F.) *Scillitique*, from *scilla*, 'squill.' Containing squill.

SCILLOCEPHALUS, *Schinoceph'alus*, from *σκίλλα*, 'the squill,' and *κεφαλή*, 'head.' One who has a large bulbiform head.

SCINCUS, *σκινκος* or *σκινγος*. A species of *Saurus* or *Lacer'ta*, common in Egypt, and much extolled by the ancients as alexipharmic and aphrodisiac.

SCINTILLA VEN'ERIS, 'a spark, a relic of Venus.' A name given, by Paracelsus, to the impotence and loss of power in the limbs, produced by syphilis.

SCINTILLEMENT DE L'CEIL, *Confusio*.

SCIOMACHY, *Skio'machia*, *Skiamachia*; from *σκια*, 'a shade,' and *μαχη*, 'a fight.' A gymnastic exercise with the ancients, which consisted in a mock encounter at boxing and jumping with one's own shadow.

SCIRHUS, *Scirrhus*.

SCIRPUS PALUSTRIS, *Eleocharis palustris*.

SCIRRHE, *Scirrhus*.

SCIRRHOBLEPHARONCUS, see *Scleriasis*.

SCIRRHOCLE, *Orcheocle scirrhus*.

SCIRRHOGASTRIA, *Scirrhus ventric'uli*, (F.) *Squirrogastrie*; from *scirrhus*, and *γαστήρ*, 'stomach.' Scirrhus of the stomach.

SCIRRHOID, *Scirrho'id*, *Scirrho'des*, from *σκῆπος*, 'scirrhus,' and *εἶδος*, 'resemblance.' Resembling scirrhus.

SCIRRHOMA, *Scirrhus*.

SCIRRHOPHTHALMIA, *Sclerophthal'mia*, from *scirrhus*, and *ὀφθαλμος*, 'the eye.' *Cancer oculi*. Cancer of the eye.

SCIRRHOPHTHALMIA PALPEBRARUM, see *Scleriasis*.

SCIRRHOSARCA NEONATORUM, *Induration of the cellular tissue*.

SCIRRHOSIS, *Scirrhus*.

SCIRRHUS, *Scirrho'us*, (F.) *Squirrheus*, from *scirrhus*. Of or belonging to scirrhus.

SCIRRHUS SARCOMA OF AB'ERNETHY, *Emphy'ma sarco'ma scirrho'sum*. Hard, rigid, vascular infarction of glandular follicles; indolent, insentient, glabrous; sometimes shrinking and becoming more indurated. Found in glandulous structures.

SCIRRHUS, *Scirhus*, *Squirrhus*, *Skirrhus*, *Scirrho'ma*, *Scirrho'sis*, *Cancer scirrho'us*, *Carcino'ma simplex seu fibro'sum*, *Induratio maligna*, *Sclerus*, *Carcinom'atous sarco'ma*, *Hard or Fibrous cancer*; from *σκῆπος*, 'hard,' 'indurated,' (F.) *Squirrhe*, *Squirre*, *Schirre*, *Skirre*, *Cancer dur* ou *squirrheux* ou *fibreux*. A disease, so called from the hardness that characterizes it. It is a state of induration, of a peculiar kind, affecting glandular structures generally, but capable of occurring in other textures. It usually precedes carcinoma, of which it may, indeed, be considered as the first stage. Scirrhus must not be confounded with the induration which succeeds inflammation, although the two states are often complicated, and may stand, occasionally, in the relation of cause and effect to each other. Scirrhus is ordinarily accompanied by violent shooting pains. It is also irregular on its surface;

and when cut into has a bluish or grayish white colour. When the surgeon is satisfied of the existence of scirrhus, he had better remove it at once. No other treatment seems to possess much advantage.

Galen uses the term *Scirrhus* for a livid excrescence caused by inflammation.

SCIRRUS ET CARCINOMA VENTRICULI, see Gastrostenosis cardiaca et pylorica—s. Hepatis, Hepatoscirrhosis—s. Intestinorum, Enteropathia cancerosa—s. Palpebrarum, see Scleriosis—s. Testiculii, Sarcocoele—s. Uteri, Metroscirrhosis—s. Ventriculi, Gastroscirrhus.

SCISSION, see Generation.

SCISSIPARITÉ, see Generation.

SCISSORS, *Scissors*, 'a carver,' from *scindere*, *scisum*, 'to cut.' *Forfex*, *Forpex*, *Palis*, *Tomis*, (F.) *Ciseaux*. A well-known instrument in common use, and often employed in surgery. Scissors are composed of two cutting blades, crossing each other and moving upon an axis. They are straight or curved. Sometimes the curve is on the flat side, when they are called, in France, *Ciseaux à cuiller*; at others, on their edges. The scissors, bent so that the blades make an obtuse angle with the handles are also much used. They are called in France *Ciseaux coudés*. Scissors are used for dividing soft, loose, isolated parts, which it would be difficult to reach and to cut with the bistoury. They are employed, likewise, in the operation for harelip, and for removing warts, fungous excrescences, &c.

SCISSURA, Fissure—s. Vulvæ, see Rima.

SCISSURÆ, Rhagades.

SCISSURE DE GLASER, Fissure, glenoid—s. *Grand médiane du Cervelet*, Valley—s. *de Santorini*, Santorini, fissure of.

SKERLJEVO, Scherlievo.

SCLOAREA, Salvia sclarea.

SCLEREMA, Induration of the cellular tissue.

SCLÉRÈME, Induration of the cellular tissue.

SCLEREMIA, Induration of the cellular tissue.

SCLEREMUS, Induration of the cellular tissue.

SCLERENCEPHALIA, *Induratio cerebri*, from *σκληρος*, 'hard,' and *εγκεφαλον*, 'the brain.' (F.) *Endurcissement du cerveau*. Induration of the brain.

SCLERIA, Scleriosis.

SCLERIASIS, *Sclero'ma*, *Sclero'sis*, *Scleria*, *Sclerysmus*, *Seph'irus*, from *σκληρος*, 'hard.' Any kind of hardness or induration. Sometimes hardness of the eyelids, *Collo'itis* seu *Scirrhophtalmia* seu *Scirrhus palpebrarum*, *Scirrhophtalaron'cus*; at others, hardness of the genital organs of the female; probably synonymous with scirrhus.—Galen. Paulus. Hardness of bones, *Fragilitas Os'ium*.

SCLERITIS, Sclerotitis.

SCLERODERMA, Induration of the cellular tissue.

SCLERODES, Sclerotic.

SCLERO-IRITIS, Inflammation of the sclerotic and iris.

SCLEROMA, Induration of the cellular tissue, Scleriosis.

SCLEROMENINX, Dura mater.

SCLERONYXIS, Scleroticonyxis.

SCLEROPHTHALMIA, *Hordeolum*, *Scirrophthalmia*.

SCLEROS, Hard.

SCLEROSARCO'MA, from *σκληρος*, 'hard,' and *σάρκωμα*, 'a fleshy tumour.' A hard, fleshy tumour, seated on the gums, and resembling a cock's comb.

SCLEROSIS, Induration of the cellular tissue, Scleriosis.

SCLEROSTENOSIS CUTANEA, Induration of the cellular tissue.

SCLEROT'IC, *Sclerot'ica*, from *σκληρος*, 'I harden.' *Cornea opa'ca*, *Sclero'des*, *Tunica dura* seu *albugin'ea* seu *du'rrior* et *crass'ior* seu *ex'tima* seu *alba* seu *can'dida* seu *innominat'a ocul'i*, *Sclero'tis*, (F.) *Scleroticque*. A hard, resisting, opaque membrane, of a pearly white colour and fibrous nature, which covers nearly the posterior four-fifths of the globe of the eye, and has the form of a sphere truncated before. It is strengthened by the expansion of the muscles of the eye, to which expansion many anatomists have given the name *Tu'nica albugin'ea*. It forms the *white of the eye*, *Albu'men* seu *Album ocul'i*, *Lon'chades*, *Log'ades*, *Lou'ades*, *Lew'cades*. The sclerotic is penetrated posteriorly by a round opening, for the passage of the optic nerve; and, anteriorly, it has a much larger opening, into which the transparent cornea is received. There are, besides, many small oblong apertures in it for the passage of nerves and vessels, proceeding to the interior of the eye. Its inner surface is of a light brown colour, owing to a delicate areolar tissue containing brown pigment cells; and hence called *Membrana seu Lam'ina fusca*. In the fœtus, the sclerotic may be separated into two very distinct laminae.

SCLEROTICA CERATOIDES, Cornea.

SCLEROTICITIS, Sclerotitis.

SCLEROTICONYX'IS, *Scleronyx'is*, *Scleroticonyx'is*, from *sclerotica*, and *νύξις*, 'puncturing.' Perforation of the sclerotic in the operation for cataract by depression.

SCLEROTICUS, Indurans.

SCLÉROTIQUE, Sclerotic.

SCLEROTIS, Sclerotic.

SCLEROTITIS, *Inflammat'io Sclerot'icæ*, *Leucit'is*, *Lonchadi'tis*, *Logadi'tis*, *Sclerit'is*, *Sclerocit'is*, *Rheumat'ic ophthalm'ia*, *Ophthalm'ia arthrit'ica*, from *sclerotica*, and *itis*, denoting inflammation. Inflammation of the sclerotic coat of the eye.

SCLEROTIUM CLAVUS, see Ergot.

SCLERUNICUS, Indurans.

SCLERUS, Induration, Scirrhus.

SCLERYNTICUS, Indurans.

SCLERYS'MA, from *σκληρός*, 'hard.' Induration; also, induration of the liver.

SCLERYSMUS, Scleriosis.

SCLIRUS, Scirrhus.

SCLOPETA'RUS, from *sclopetum*, 'a gun.' Relating to a gun. Hence, *Aqua sclopetaria*, a water for gunshot wounds. *Vul'nera sclopetaria*, gunshot wounds, &c.

SCLOPETOPLAGÆ, Wounds, gunshot.

SCOBS, *Prisma*, *Xysma*, *Ramen'tum*, *Rasur'a*. A shaving; also, an alkali; and the scoria of any metal.

SCENANTHUM, *Juncus odoratus*.

SCOLECESIS, Helminthiasis.

SCOLECIASIS, Helminthiasis.

SCOLEX, *Ascaris lumbricoides*.

SCOLIOMA, Scoliosis.

SCOLIO'SIS, *Scolio'ma*, *Skolio'sis*, *Rhachio scolio'ma*, from *σκολιος*, 'crooked.' A distortion of the spine to one side. See Hump.

SCOLOPENDRA, *Asplenium scolopendrium*.

SCOLOPENDRIA, *Asplenium ceterach*, and *A. scolopendrium*.

SCOLOPENDRIUM LINGUA, *Asplenium scolopendrium*—s. *Officinarium*, *Asplenium scolopendrium*—s. *Phyllitis*, *Asplenium scolopendrium*—s. *Ruta muraria*, *Asplenium ruta muraria*—s. *Vulgare*, *Asplenium scolopendrium*.

SCOLOPOMACHÆRION, from *σκολομαχ*, 'the woodcock,' and *μαχαίρα*, 'a knife,' so called because it is bent a little at the extremity like a woodcock's bill. A sort of blunt-pointed bistoury.—Scultetus.

SCOLYMUS SATIVUS, *Cynara scolymus*.

SCOOP, Teut. Schoepe, *Cochlea're*, (F.) *Curette*. A surgical instrument of the shape of a spoon, which is used for the extraction of certain foreign bodies. The scoop is often employed to extract balls impacted in the soft parts; to remove calculi from the urinary bladder in lithotomy, &c. A small scoop is sometimes used to extract foreign bodies from the meatus auditorius externus, nasal fossæ, &c.

SCOP. Head, Scalp.

SCOPA REGIA, *Ruscus*.

SCOPARIUS, see *Spartium scoparium*.

SCOPE, *σκοπε*, *Scopy*, *Scop'ia*, *Scopos*, from *σκοπεω*, 'I look around.' A common suffix to words meaning 'view, inspection,' as *cranioscopia*, *stethoscopia*, &c.

SCORPOLINA ATROPOIDES, *Hyoseyamus scopolia*.

SCOPTULA, *Scopula*.

SCOPULA, Brush.

SCOR, Excrement.

SCORACRAT'IA, *Scoracra'sia*, *Scatacra'sia*, *Scatacrati'a*, *Copracrati'a*, *Copracra'sia*, *Incontin'ia Alvi*, from *σκορ*, 'excrement,' and *ακρα*, 'want of control.' Want of power to retain the fæces. Involuntary evacuation of the fæces.

SCORBUT, *Porphyra nautica*, *Purpura hæmorrhagica*—*s. des Alpes*, *Pellagra*—*s. de Terre*, *Purpura hæmorrhagica*.

SCORBU'TIC, *Scorbuticus*, (F.) *Scorbutique*, from *scorbutus*, 'scurvy.' That which belongs to scurvy. One affected with scurvy.

SCORBUS, *Purpura*—*s. Alpinus*, *Pellagra*—*s. Nauticus*, *Porphyra nautica*—*s. Oris*, *Cancer aquaticus*.

SCORDINEMA, *Carebaria*, *Pandiculation*.

SCORDINISMUS, *Carebaria*, *Pandiculation*.

SCORDIUM, *Toucerium scordium*.

SCORDON. Allium.

SCORDOSMINE, Oxyd, cystic.

SCORITH, Sulphur.

SCORIUM, Scrotum.

SCORODON. Allium.

SCORPIODEX'IS, from *σκορπιος*, 'the scorpion,' and *δῆξις*, 'bite'—more properly, *SCORPISIG'MA*; from *σκορπιος*, and *στυγμα*, 'puncture.' The sting of the scorpion.

SCOR'PION, *Scor'pio*, *Scorpius*, from *σκορπιος*, 'I enrage.' A genus of insects having a sting at the extremity of their tail, with which they make dangerous wounds. An Oil of Scorpion was once made from them, which is not now used.

SCORPIOSTIGMA, *Scorpiodexis*.

SCORPIUS, Scorpion.

SCORTUM. Scrotum.

SCORZONE'RA, *S. hu'milis* seu *landa'ta* seu *neruo'sa*, *Escorzone'ra*, from *scurzo*, old Spanish for a viper, *Vipera'ria*, *Serpenta'ria Hispan'ica*, *Offic'inal viper's grass*, *Viper's grass*, *Goat's grass*. Family, *Cichoraceæ*. *Sec. Syst.* *Syngenesia Polygamia*. The roots have been used as alexipharms, and in hypochondriacal disorders and obstructions of the viscera. The root of the *Scorzonera Hispan'ica* is esculent, but inefficacious as a drug.

SCOTAS'MA, *Scoto'ma*, *Scoto'dia*, *Scoto'sia*, *Scotos*; from *σκοτος*, 'darkness.' Obscurity of vision, darkness, *Suffusio rad'ians*. Also *Scotodinia*.

SCOTCH FIDDLE, *Psora*.

SCOTODIA, *Scotasma*, *Scotodinia*.

SCOTODIN'IA, *Scoto'ma*, *Dinus Scoto'ma*, *Scotod'ine*, *Tenebrio'sa verti'go*, *Scotos*, *Scotas'ma*, *Scoto'sia*, *Scot'omy*, from *σκοτος*, 'darkness,' and *δινω*, 'I turn round.' Giddiness, with impaired sight, often succeeded by headach. See *Vertigo*.

SCOTOMA. *Scotasma*. *Scotodinia*.

SCOTOMY, *Scotodinia*.

SCOTOS. *Scotasma*. *Scotodinia*.

SCOTOSIS, *Scotodinia*.

SCOTT'S ACID BATH, *Nitro-muriatic acid bath*. A bath of dilute aqua regia, employed by Dr. Scott, of India, as a remedy in hepatic diseases. Three pints by measure of muriatic acid, and two of nitric acid, are mixed to form the aqua regia. In preparing this for use, a pint of it is mixed with an equal quantity of water. The acid bath consists of three ounces of this dilute acid to every gallon of water.

SCOUR, *Diarrhœa*.

SCOURING, *Diarrhœa*.

SCRAT, *Hermaphrodite*, *Psora*.

SCRATCH, *Psora*.

SCREATUS, *Excretion*.

SCRIPULUM, *Scruple*.

SCRIPTIUM, *Scruple*.

SCRIPTULUM, *Scruple*.

SCRIPULUM, *Scruple*.

SCRIPULUS, *Scruple*.

SCRIVENER'S CRAMP, *Cramp*, *writers'*.

SCROBIC'ULUS, *Scrobic'ulum*, (F.) *Fossette*. A little ditch or furrow; diminutive of *scrobs*, 'a ditch.'

SCROBIC'ULUS CORDIS, *Anticar'dium*, *Procar'dium*, (F.) *Fossette du cœur*, *Avant-cœur*, *Anticœur*. The pit of the stomach. The depression at the anterior part of the abdomen, below the xiphoid cartilage.

SCROBICULUS VARIOLÆ, *Pockmark*.

SCROFFÆ, *Scrofula*.

SCROF'ULA, *Scroph'ula*, *Scroph'ulæ*; from *scrofa*, 'a sow;' as *Chœras* is from *χοιρος*, 'a hog,' because swine were presumed to be subject to a similar complaint. *Scrophulo'sis*, *Morbus scrophulo'sus*, *Vit'ium scrophulo'sum*, *Cacochym'ia* seu *Cachex'ia* seu *Dyscras'ia scrophulo'sa*, *Pada'troph'ia glandulo'sa*, *Struma*, *Chœras*, *Coiræas*, *Tabes Glandula'ris*, *Adeno'sis scrofulo'sa*, *Scroffa*, *King's evil*, *The Evil*, *Quince*, *Cruels* (*Scotticè*), (F.) *Scrophules*, *Scrofulæ*, *Strumes*, *Ecouelles*, *Humeurs froides*, *Tubercules des glandes lymphatiques*, *Mal du Roi*. A state of the system characterized by indolent, glandular tumours, chiefly in the neck; suppurating slowly and imperfectly, and healing with difficulty; the disease ordinarily occurring in those of a sanguine temperament, with thick upper lip, &c. The tumours, after suppuration, degenerate into ulcers; which, in process of time, cicatrize, leaving scars. The internal organs are apt to be attacked in those disposed to scrofula; hence they are often the subjects of phthisis and mesenteric affections. Scrofula is hereditary; and is frequently excited by insufficient or improper regimen, and by a close, confined air during the first years of existence. The best treatment is:—to strengthen the system by animal diet; pure air, and exercise, cold bathing, or sea-water bathing, &c.

SCROFULA ABDOMINALIS INTERNA, *Physconia strumosa*.

SCROFULAIRE, *Scrophularia nodosa*—*s. Aquatique*, *Scrophularia aquatica*.

SCROFULARIA, *Scrophularia*.

SCROFULES, *Scrofula*.

SCROFULEUX, *Scrofulous*.

SCROF'ULOUS, *Scrophulo'sus*, *Scrophula'rius*, *Strumous*, *Strumo'sus*, (F.) *Scrofuloux*, *Strumeux*, Suffering from, or relating to, scrofula.

SCROOBY-GRASS, *Cochlearia officinalis*.

SCROOF, Scarf.

SCROG-LEGGED, *Cnemosciosis*.

SCROPHULA, *Scrofula*—*s. Mesenterica*, *Tabes mesenterica*—*s. Moleucana*, *Frambœsia*.

SCROPHULÆ, *Scrofula*—*s. Gonorrhœicæ*, *Gonorrhœa impura*.

SCROPHULARIA, from *scrofula*, against which it has been employed. *S. Nodosa*.

SCROPHULARIA AQUATICA, *Betonica aquatica*, *Ficaria*, *Ferraria*, *Castranigula*, *Greater Water Figwort*, *Water Betony*, (F.) *Scrophulaire aquatique*. *Ord. Scrophulariaceae*. The leaves are celebrated as correctors of the bad flavour of senna. They were at one time regarded as eminently antiscrofulous, stomachic, and carminative.

SCROPHULARIA FETIDA, *Scrophularia nodosa* — *s. Lanceolata*, *S. Nodosa* — *s. Marilandica*, *S. Nodosa* — *s. Minor*, *Ranunculus ficaria*.

SCROPHULARIA NODOSA, *S. foetida* seu *vulgaris* seu *lanceolata* seu *Marilandica*, *Millenordia*, *Scrophularia*, *Figwort*, *Kennehoort*, *Holmesweed*, *Heal-all*, (F.) *Scrophulaire*, *Herbe aux écrouelles*. The root and leaves of this plant have been celebrated, both as an internal and external remedy, against inflammations, piles, scrofulous tumours, ulcers, &c.

SCROPHULARIA VULGARIS, *Scrophularia nodosa*. **SCROPHULARIUS**, *Scrofulus*.

SCROPHULES, *Scrofula* — *s. Mésentériques*, *Tabes mesenterica*.

SCROPHULOSIS, *Scrofula*.

SCROPHULOSUS, *Scrofulous*.

SCROTAL, *Scrotalis*; from *scrotum*. Relating to the scrotum, — as *Hernia scrotalis*, *scrotal hernia*.

SCROTAL NERVE, see *Genito-crural Nerve*.

SCROTOCELE, *Oscchocele*, *Oscchocele*. A hybrid term; from (L.) *scrotum*, and *κηλη*, 'a tumour.' *Hernia Scrotalis*, *Scrotal hernia*. Inguinal hernia, descending into the scrotum.

SCROTUM, from *scorteus*, *scortum*, 'made of leather' [?]. *Scorium*, *Oschē*, *Os'cheus*, *Os'chus*, *Ocus*, *Os'cheon*, *Och'eus*, *Perin*, *Scortum*, *Marau'pium*, *Bur'eula*, *Bursa testium* seu *virilis*, *Cap'ulus* [?], *Orchas*, *Follic'ulus castralis*, *Her'nia*, the *Purse*, the *Cod*, (F.) *Les bourses*. The integuments which cover the testes. These are nothing more than a prolongation of the skin of the inner part of the thighs, perinæum, and penis. The skin is remarkable for its brown colour, for the numerous rugæ, especially when the scrotum is contracted; for the great number of sebaceous follicles it contains; and for the long and few hairs which issue from it in the adult. The scrotum is divided into two equal parts by a raphe, which extends from the anus to the root of the penis. Its organization is similar to that of the skin of other parts of the body, except that the chorion is thinner.

SCROTUM CORDIS, *Pericardium* — *s. Pendulum*, *Rhachosis*.

SCRUBIE, *Purpura nautica*.

SCRUBIE-GRASS, *Cochlearia officinalis*.

SCRUPLE, *Diodolon*, *Grammarium*, *Scrupulus*, *Scrupulum*, *Scriptulus*, *Scriptulum*, *Scriptium*, *Scrupulum*, *Scriptum*, *Scrupulus*. A weight of twenty grains, Troy; twenty-four grains, Poids de marc.

SCUDAMORE'S MIXTURE, see *Mixture*, *Scudamore's*.

SCULL, *Cranium* — *s. Cap*, *Calvaria*.

SCURF, *Sax. scurf*, *Teut. Schorff*; (Prov.) *Serof*, *Furfura*. Small exfoliations of the cuticle are so called.

SCURFY, *Furfuraceous*.

SCURVY OF THE ALPS, *Pellagra* — *s. Land*, *Purpura hæmorrhagica* — *s. Petechial*, *Purpura simplex* — *s. Sea*, *Porphyra nautica*.

SCURVY GRASS, *Sisyrinchium Bermudianum* — *s. Grass*, *common*, *Cochlearia officinalis* — *s. Grass*, *lemon*, *Cochlearia officinalis* — *s. Grass*, *wild*, *Cochlearia coronopus*.

SCUTELLA, *Cup*.

SCUTELLARIA CAROLINIANA, *S. integrifolia*.

SCUTELLARIA GALERICULATA, from *scutella*, 'a cup,' in allusion to the form of the calyx. *Cassida galericulata*, *Tertianaria*, *Skull-cap*, (F.) *Casside bleue*, *Toque*. *Ord. Labiatae*. *Ser. Syst. Didynamia Angiosperma*. This plant has a bitter taste and a garlic smell. It has been esteemed especially serviceable in tertian ague.

SCUTELLARIA HYSSOPIFOLIA, *S. integrifolia*.

SCUTELLARIA INTEGRIFOLIA, *S. hyssopifolia* seu *Caroliniana*, — another indigenous variety, — is an intense bitter.

SCUTELLARIA LATERIFLORA, *Skull-cap*, *Mad-dog skullcap*, *Mudweed*, *Woodwort*, *Blue Pimpernel*. An indigenous plant, which grows in all parts of the United States in moist places. It was at one time esteemed a preventive of hydrophobia, but is probably possessed of little virtue of any kind. By some, however, it is regarded as a good nervine.

SCUTICULA DIGITORUM MANÛS, *Phalanges of the fingers*.

SCUTTER, *Diarrhoea*.

SCUTULÆ DIGITORUM MANÛS, *Phalanges of the fingers*.

SCUTUM, *Thyroid cartilage* — *s. Cordis*, *Scapula*, *Sternum* — *s. Genu*, *Patella* — *s. Pectoris*, *Sternum*, *Thorax*.

SCUTUM STOMACHICUM. A large plaster, applied to the breast or stomach.

SCUTUM THORACIS, *Scapula*.

SCYBALA, plural of *σκαβλον*, 'dung.' *Enterolithus Scybalum*, *Excrementa Alvi sicci'ora*, *Fæces induratae*. Hard fecal matters discharged in round lumps.

SCYLLA, *Scilla*.

SCYPHOPHORUS PYXIDATUS, *Lichen pyxidatus*.

SCYPHUS AUDITORIUS, *Infundibulum of the cochlea* — *s. Cerebri*, *Infundibulum of the brain* — *s. Vuesseini*, *Infundibulum of the cochlea* *SCYROS*, *Callosity*.

SCYTALIDES DIGITORUM MANÛS, *Phalanges of the fingers*.

SCYTHICA, (*RADIX*), *Glycyrrhiza*.

SCYTHROPASMUS, from *σκυθος*, 'gloomy;' *σκυθρωπαζων*, 'to look gloomy;' *Vultus tet'ricus et mæstus*. A gloomy, depressed countenance, of bad augury in serious diseases.

SCYTITIS, *Cytitis*.

SCYTODEPSIUM, *Tannin*.

SEA AIR. The air in the neighbourhood of the sea contains saline particles commingled with it, which render it a valuable change for scrofulous and debilitated individuals. The sea and sea-coast are usually, also, more equable in their temperature than places inland.

SEA SICKNESS, *Nausea marina*.

SEA-WARE, see *Pila marina* — *s. Water*, see *Water*, *sea*.

SEAL, *GOLDEN*, *Hydrastis Canadensis* — *s. Solomon's*, *Convallaria polygonatum* — *s. Solomon's*, *great*, *Convallaria canaliculata* — *s. Solomon's*, *smaller*, *Convallaria pubescens*.

SEAM, *Cicatrix*.

SEARCHER, *Body searcher*.

SEARCHING, *Sounding*; from (F.) *chercher*, 'to seek.' The operation of ascertaining, by the introduction of a metallic instrument, whether there is a stone in the bladder.

SEASONING, *Acclimation* — *s. Fever*, *Fever*, *stranger's*.

SEAT, *Anus* — *s. Bone*, *Ischion*.

SEBACE, *Sebaceous*.

SEBA'CEOUS, *Seba'ceus*, (F.) *Sébacé*; from *sebum*, 'suet.' That which has the nature of suet.

SEBACEOUS FLUX, Stearrhœa.

SEBACEOUS, SEBIPAROUS or SEBIFEROUS GLANDS, *Sebaceous Follicles* or *Crypts*, *Folliculi seba'ci*, *Cryptæ seba'cæ*, Oil or *Miliary glands*, are small hollow organs, of a rounded or pyriform shape, seated in the substance of the skin, and opening on its surface by a small, excretory duct; furnishing a yellow, unctuous humour, *Smegma cutaneum*, *Sebum cutaneum*, having some analogy with suet, which is destined to lubricate the surface of the body. Follicles of a similar nature exist around the corona glandis of the male, and under the skin of the labia majora and nymphæ of the female. They secrete a sebaceous matter, which emits a peculiar odour; and hence, after the name of those who described them, have been called *Glandulæ odorifera Tyso'ni*, *G. Tyso'ni*, *Glandulæ Littriana'ne* seu *Littrii*, *Tyson's glands*, (*F.*) *Glandes de Littré*; and, also, *Glandulæ odorifera' seu seba'cæ glandis*, *G. cor'ona' penis*, *Folliculi seba'ci corona' glandis*, *Glandulæ odorifera' præputii*.

SEBASTOMANIA, from *σεβαστος*, 'worshiped,' and *mania*. Religious insanity, Demonomania.

SEBESTENA, *Cordia myxa* seu *sebestena* seu *Africa'na* seu *domes'tica* seu *obli'qua*, *Sebestena officina'lis*, *Seb'esten*, *Sebesten*, *Myxa*, *Cornus sanguinea*, *Prunus sebestina*, *Smooth-leaved Cordia*, *Assyr'ian plum*, (*F.*) *Sebestier myxa*. Ord. Cordiaceæ. The dark, black fruit of the *Sebestina* possesses glutinous and aperient qualities, and is exhibited, in decoction, in various diseases of the chest.

SEBESTIER MYXA, *Sebestena*.

SEBIFEROUS, see Sebaceous.

SEBIPAROUS, see Sebaceous.

SEBORRHAGIA, Stearrhœa.

SEBORRHŒA, Stearrhœa.

SEBSTEN, *Sebestina*.

SEBUM, Pinguedo, *Sevum*—s. *Cutaneum*, see Sebaceous glands—s. *Palpebrale*, *Chassie*—s. *Præputiale*, *Smegma præputii*.

SECALE, *S. cereale* seu *barbatum*. The Rye plant. (*F.*) *Seigle*. Ord. Gramineæ. Rye is chiefly used as an article of diet, particularly in the northern parts of Europe, where a spirit is also extracted from it. The grain is disposed to acescency: and hence its internal use at first generally produces a laxative effect.

SECALE BARBATUM, *Secale*—s. *Cornutum*, *Ergot*—s. *Luxurians*, *Ergot*—s. *Maternum turgidum* seu *temulentum*, *Ergot*.

SECALIS MATER, *Ergot*.

SÉCATEUR DES AMYGDALES, *Amygdalotome*.

SECERNENT, Secreting.

SECERNING, Secreting—s. Substance of the kidney, see *Kidney*.

SECESSIO FÆCUM, Defecation.

SECESSUS NIGER, *Melæna*.

SÈCHE, *Sepia*.

SECLUSORIUM, *Sac*.

SEC'ONDARY, from *secundus*, 'the second.' Something that acts subordinately to another; as the secondary symptoms of a disease; that is, those which supervene on the primary. A secondary amputation is one performed after the constitutional effects of an injury have mainly passed away. See Fever, secondary; and Hæmorrhagia, for Secondary Hemorrhage.

SECUNDINES, *Secundines*.

SECRETA. Same etymon as Secretion. Things or matters secreted or separated from the blood.

SECRETING, *Secreto'rius*, *Secer'ning*, *Secern'ent*, *Excer'nent*, (*F.*) *Sécréteur*, *Sécrétoire*. Same etymon as Secretion. That which secretes, — as a secreting organ.

SECRETIO, Secretion—s. *Lactis*, *Galactosis*—s. *Lotii*, *Uropoiesis*—s. *Urinæ*, *Uropoiesis*—s. *Vicaria*, see *Vicarius*.

SECRETION, *Secre'tio*, *Apoc'risis*; from *se-cernere*, *secretum*, 'to separate.' An organic function, which is chiefly executed in the glands, and consists in an elaboration or separation of the materials of the blood, at the very extremities of the arterial system, or rather of the vascular secretory system; and which differs in each organ according to its particular structure; hence the formation of different fluids;—bile, saliva, urine, milk, &c. The secretions are of three kinds:—*exhalant*, *follicular*, and *glandular*. Simple secretions are those which exist ready-formed in the blood, and pass out of the vessels by an act of exosmosis, whilst the more complicated are formed from the liquor sanguinis by cell-agency. The latter, alone, are, by some, called *Secretions*.

SÉCRÉTOIRE, Secreting.

SECRETORIUS, Secreting.

SECT, METHODICAL, Methodists.

SECTIO, *Section*. The act of cutting. A division.

SECTIO ALTA, see *Lithotomy*—s. *Anatomica*, *Dissection*—s. *Cadaveris*, *Autopsia cadaverica*—s. *Cadaveris legalis*, *Autopsia cadaverica*—s. *Cæsarea*, *Cæsarean section*—s. *Franconiana*, see *Lithotomy*—s. *Hypogastrica*, see *Lithotomy*—s. *Lateralis*, see *Lithotomy*—s. *Legalis*, *Autopsia cadaverica legalis*—s. *Mariana*, see *Lithotomy*—s. *Nympharum*, *Nymphotomy*—s. *Perinæalis*, *Perineal section*—s. *Rectovesicalis*, see *Lithotomy*—s. *Renalis*, *Nephrotomy*—s. *Tendinum*, *Tenotomy*—s. *Vagino-vesicalis*, *Colpocystotomia*—s. *Vesicalis*, *Lithotomy*.

SECTION, *Sectio*—s. *Sigaultian*, *Symphysotomy*.

SECUNDÆ, *Secundines*.

SECUNDINÆ, *Secundines*.

SEC'UNDINES, *Hys'tera*, *Deuter'ion*, *Metra*, *Secun'dæ*, *Secundi'na*, *After-birth*, (*F.*) *Arridre-faix*, *Secundines*, *Délivre*, from *secundare*, (*secundus*, 'second,' 'to follow.' All that remains in the uterus after the birth of the child,—viz., the placenta, a portion of the umbilical cord, and the membranes of the ovum. These are commonly not expelled till some time after the birth of the fœtus; hence their name. See *Délivrance*.

SECUNDUS PROPRIORUM AURICULÆ, *Retrahens auris*.

SEDANTIA, Sedatives.

SEDATIVES, *Sedati'va*, *Pauon'tia*, *Sedan'tia*, *Adec'ta*, *Catastaltica*, *Deprimen'tia*, (*F.*) *Sédatifs*, *Calman'ts*, *Tempéran'ts*, from *sedare*, *sedatum*, 'to settle or assuage.' Medicines which directly depress the vital forces, and which are consequently employed whenever it is necessary to diminish preternaturally increased action. The chief reputed sedatives are: *Acidum Hydrocyanicum*, *Acidum Hydro-sulphuricum*, *Tabacum*, *Missio Sanguinis*, and certain gases by inhalation, as azote, carbonic acid [?], carburetted and sulphuretted hydrogen.

SEDEM ATTOLLENS, *Levator ani*.

SEDES, Anus, Excrement—s. *Cruentæ*, *Dysentery*, *Hæmatochezia*—s. *Lactescentes*, *Cœliac flux*—s. *Procidua*, *Proctocele*.

SEDHEE, *Bangue*.

SED'IMENT, *Sedimen'tum*, *Lemma*, *Residen'tia*. Same etymon. (*F.*) *Dépôt*. A deposit formed by the precipitation of some one or more of the substances held in solution or suspension by a liquid. That which is formed in urine, at the time of cooling, has been called *Hypos'tasis*, *Hypostema*, *Residentia*, *Subsidentia*, (*F.*) *Dépôt de l'urine*. This sediment or deposit varies in different states of the system. In calculous affec-

tions, it is one of the most important objects of attention.

SEDIMENTUM, Sediment—*s. Urinæ lateritium*, see Lateritious—*s. Urinæ pityroides*, see Furfuraceous.

SEDITIA, Nates.

SEDLITZ, MINERAL WATERS OF, *Seidlitz water*. Springs in Bohemia, near Prague, which are simple salines. They contain sulphate and carbonate of magnesia, chiefly chloride of magnesium, sulphates of potassa, soda, and lime, carbonic acid and carbonate of lime. They are employed as purgatives in a multitude of diseases.

ARTIFICIAL SEDLITZ POWDERS, in no respect, however, resembling the natural ingredients, are formed of *Sodæ et Potassæ tartras 3ij*, *Sodæ carbon. ʒij*, in one paper; *Acid tartaric. gr. xxxv*, in another. For half a pint of water.

SÉDON BRÛLANT, Sedum—*s. Reprise*, Sedum telephium.

SEDUM, *S. acré seu glaciæ læ seu minus seu vermiculæ rē*, from *sedeo*, 'I sit,' owing to its attachment to rocks and walls. *Illec'ebra*, *Hellec'ebra*, *Illec'ebrium*, *Vermiculæ rē*, *Vermiculæ ria*, *Piper mura'le*, *Sempervivum acré*, *Aci'chryson*, *Æo'nion*, Wall pepper, Jack of the Buttery, Pricket, Bird's bread, Stone crop, Biting Stone crop, Small Houseleek, (*F.*) *Joubarbe acre*, *Orpin brûlant*, *Vermiculaire brûlante*, *Sédon brûlant*. *Ord. Crassulacæ. Sex. Syst. Decandria Pentagynia*. In its recent state it is very acrid, and proves both emetic and cathartic. Externally, in the form of cataplasm, it produces vesications and erosions. It has been recommended in cancerous and malignant ulcers.

SEDUM, *Saxifraga granulata*—*s. Glaciale*, Sedum—*s. Majus*, *Sempervivum tectorum*—*s. Minus*, Sedum.

SEDUM TELEPHIUM, *S. telephoides*, *Faba crassa*, *Cras'sula*, *O. Major*, *Illec'ebra major*, *Telephium*, *Faba'ria cras'sula*, *Faba crassa*, *Anacam'peros*, *A. al'bicans seu triphyll'a seu vulga'ris*, *Orpine*, (*F.*) *Sédon reprise*, *Grand orpin*, *Fève épaisse*, *Joubarbe des vignes*, &c. Formerly used as a cataplasm in cuts, hemorrhoids, corns, whitlows, &c.

SEDUM TELEPHOIDES, *S. Telephium*—*s. Vermiculaire*, Sedum.

SEED, *Sperm.*

SEGMENTA CARTILAGINEA, see Trachea.

SEGMENTA'TION, *Segmenta'tio*, from Lat. *segmentum*, 'a segment,' itself from *seco*, 'to cut.' The act of being divided into segments or portions. (*F.*) *Segmentation*, *Fractionnement*, *Sillonement*. A term applied to the division of the vitellus or yolk into 'segments,' each containing a transparent vesicle, which may be a descendant of the primordial or germ cell.

SEGNIT'IA, *Segnit'ies*, from *segnis*, 'sluggish.' Sluggishness or torpor in the exercise of a function:—as *SEGNITIES ALVI*, Torpor of the bowels; *SEGNITIES VIRIUM*, Languor.

SEGNITIES, *Segnitia*.

SEGRAY, MINERAL WATERS OF. Near Pithiviers, in Gâtinais, in the French department of Loiret, there is a chalybeate spring, which has some reputation.

SEGU, *Sago*.

SEDLITZ, *Sedlitz*.

SEIDSCHÜTZ, **SEYDSCHÜTZ** or **SAID-SCHÜTZ**, MINERAL WATERS OF. Bitter springs, in the neighbourhood of those of Püllna and Sedlitz, in Bohemia, which contain chiefly sulphate, nitrate, muriate, and carbonate of magnesia, sulphates of potassa, soda, and lime; carbonate of lime, iron and manganese. The water is an excellent cathartic, and is well known in Europe.

SEIGLE, *Seca.e*—*s. Cornu*, *Ergot*—*s. Corrompu*, *Ergot*—*s. à Éperon*, *Ergot*—*s. Ergotée*, *Ergot*—*s. Ergotisé*, *Ergot*—*s. Fauz*, *Ergot*.

SEIN, Uterus, see Mamma.

SEIRIASIS, *Coup de Soleil*.

SEIZURES, *HIDDEN*, see Hidden seizures.

SEKISKAYAVODKA, see Spirit.

SEL, *Salt*—*s. Admirable de Lémery*, *Magnesia sulphas*—*s. Admirable perlé*, *Soda*, phosphate of—*s. Ammoniac*, *Ammonia murias*—*s. Arménia*, *Ammonia murias*—*s. Arsenical de Macquer*, *Arseniate of protoxide of potassium*—*s. de Cuisine*, *Soda*, muriate of—*s. de Déroene*, *Narcotine*—*s. d'Égra*, *Magnesia sulphas*—*s. de Lait*, *Sugar of Milk*—*s. Marin*, *Soda*, muriate of—*s. d'Opium*, *Narcotine*—*s. d'Oseille*, *Potassæ oxalas*—*s. de Saturne*, *Plumbi superacetats*—*s. de Verre*, see *Vitrum*—*s. Volatil d'Angleterre*, *Ammonia carbonas*.

SELAGO, *Camphorosma Monspeliaca*, *Lycopodium selago*.

SELENE, Moon, see Nail.

SELENIASIS, *Somnambulism*.

SELENIASMUS, *Somnambulism*.

SELENOBLETUS, *Lunatic*.

SELENOGAMIA, *Somnambulism*.

SELERY, *Apium graveolens*.

SELF-ABUSE, *Masturbation*.

SELF-HEAL, *Prunella*.

SELF-LIMITED. An epithet applied to diseases which appear to run a definite course, but little modified by treatment,—small-pox, for example.

SELF-POLLUTION, *Masturbation*.

SELI'BRA, *Semili'bra*, *Sembella*, 'half a pound.' Six ounces, Troy; eight Avoirdupois.

SELINITES, *σελινιτης*. Wine impregnated with the seeds of the *σελινον* or smallage.—*Dioscorides*.

SELINUM ANGELICA, *Angelica Sylvestris*—*s. Galbanum*, *Bubon galbanum*—*s. Imperatoria*, *Imperatoria*—*s. Lactescens*, *Peucedanum palustre*—*s. Marsh*, *Peucedanum palustre*—*s. Opoponax*, *Pastinaca opoponax*—*s. Oreoselinum*, *Athamanta aureoselinum*—*s. Ostruthium*, *Imperatoria*—*s. Palustre*, *Peucedanum palustre*—*s. Pastinaca*, *Pastinaca sativa*—*s. Peucedanum*, *Peucedanum*—*s. Pubescens*, *Angelica sylvestris*—*s. Sylvestre*, *Angelica sylvestris*, *Peucedanum palustre*—*s. Thysselinum*, *Peucedanum palustre*.

SELLA, *S. Turcica*—*s. Equina*, *S. Turcica*—*s. Familiarica*, *Lasanum*—*s. Obstetricia*, *Diphrys*—*s. Sphenoidalis*, *S. Turcica*.

SELLA TUR'CICA, *Ephipp'ium*, *Fossa Pituitaria*, *Sella*, *S. equina seu sphenoidalis*, *Pituitary fossa*, *Turkish Saddle*, (*F.*) *Selle Turcique ou du Turc*. A depression at the upper surface of the sphenoid bone, which is bounded, anteriorly and posteriorly, by the clinoid processes, and lodges the pituitary gland. It is so called from its resemblance to a Turkish saddle. The broad rough plate, behind, and somewhat overhanging the Sella Turcica is the *Dorsum Ephippii*.

SELLA TURCICA, *Pituitaria fossa*.

SELLE TURCIQUE, *Sella Turcica*—*s. du Turc*, *Sella Turcica*.

SELS NEUTRES, *Neutral salts*.

SELTZ or **SELTZER**, MINERAL WATERS OF. Cold, acidulous springs, the source of which is at Seltz, nine leagues from Strasburg, and in the Department of the Bas-Rhin. They contain carbonates of lime, magnesia, and soda, chloride of sodium, and much carbonic acid. They are refrigerant, tonic, diuretic, and aperient.

ARTIFICIAL SELTZER WATER may be made from *Muriat. acid. gr. xxxv*, *water Oj*, *white marble gr. iij*. Stop up till dissolved. Add *carbonate of Magnesia*, *gr. v*, and, after some time, *subcarbonate of soda*, *gr. xxxij*. Close it till used.

SEMBELLA. Selibra.

SEMECARPUS ANACARDIUM, *Avicennia tomentosa*.

SEMEIOGRAPHY, *Semeiograph'ia*, *Semiography*, *Semiograph'ia*, *Symptomato'graphy*, from *σμεῖον*, 'a symptom,' and *γραφον*, 'a description.' A description of symptoms or signs of disease.

SEMEIOLOGY, *Semeiolog'ice*, *Semiology*, *Semiologic*, *Phænomenolog'ia*, from *σμεῖον*, 'a symptom,' and *λογος*, 'a discourse.' (F.) *Sémiologie*. The branch of pathology whose object is the doctrine of the symptoms or signs of disease.

SEMEION, Sign — s. *Boethematicum*, *Indication*.

SEMEIOSIS, *Semiology*.

SÉMÉIOTIQUE, *Semiology*.

SEMEN, *Sperm* — s. *Badian*, *Illicium anisatum* — s. *Cina*, see *Artemisia santonica* — s. *Contra*, see *Artemisia santonica* — s. *Contra d'Alep*, *Artemisia santonica* — s. *Contra vermes*, *Artemisia santonica* — s. *Masculinum seu virile seu genitale*, *Sperm* — s. *Muliebres*, *Sperm* (of the female) — s. *Sanctum*, see *Artemisia santonica* — s. *Sancti*, see *Artemisia santonica* — s. *Zedoaria*, see *Artemisia santonica*.

SEMPENCE, *Sperm*.

SEMENTINA, *Artemisia santonica*.

SEMI, from *ἡμιος*, 'half.' *Semi* or *semi*, in composition, universally signifies 'half,' both in French and English.

SEMIANIMIS, *Semimortuus*, *Sem'inez*, *Semivivus*; from *semi*, 'half,' and *animus*, 'soul.' Half-living. Half-dead.

SEMI-BULB OF THE FEMALE, *Bulbus vestibuli*.

SEMICANALICULUS, *Sulcus*.

SEMICANALIS, *Sulcus*.

SEMICEPHALUS, *Hemicephalus*.

SEMICIRCULI OSSEI, *Semicircular canals*.

SEMICIRCULAR, *Semicircula'ris*. From *semi*, and *circulus*, 'a circle.' (F.) *Demicirculaire*. That which represents the half of a circle.

SEMICIRCULAR CANALS, *Canal'es seu Ductus semicircula'res seu tuba'form'es seu circula'res*, *Semicirculi ossei*, *Funes seu Canaliculi semicircula'res labyrinthi*, (F.) *Canaux demicirculaires*, are seated in the pars petrosa of the temporal bone, and open into the vestibule, behind which they are situate. They are three in number, of which one is *superior* or *vertical*, *canalis semicircula'ris verticalis super'ior*; another *posterior* or *oblique*, *canalis semicircula'ris verticalis post'erior*, and the third *exterior* or *horizontal*, *canalis semicircula'ris horizontalis*. Their parietes are formed of a bony, compact, hard plate. Their uses in audition are not known: at their termination they have the elliptical arrangements called *ampullae*; and by both extremities they open into the vestibule, in the sac at its superior part—the *Sacculus ellipticus*, *Alveus seu Utriculus* seu *Sinus communis*, *Sinus seu Alveus utriculo'sus*, *Sacculus vestibuli* or *Median sinus*, *Utriculus*. In front of the S. ellipticus, nearer the cochlea, and opposite the foramen ovale, is the *Sacculus sphaer'icus*, or *Sacculus*.

SEMICIRCULUS EXSCULPTUS, *Hiatus diaphragmatis aorticus*.

SEMICONGIUS, Half a gallon, or four pints.

SEMICUBIUM, *Semicutium*, *Inses'cio*, *Inces'cio*, *Encathisma*, *Excathisma*, (F.) *Demibain*, from *semi*, and *cubare*, 'to lie down.' A half-bath, or such as receives only the hips or extremities.

SEMICUPIUM, Bath, half, *Semi-cubium*.

SEMI-FIBULEUS, *Peronæus brevis*.

SEMI-INTEROSSEUS INDICIS, *Abductor indicis*.

SEMI-INTEROSSEUS POLICIS, *Opponens pollicis*.

SEMLIBRA, *Selibra*.

SEMILLA DEL GUACHARO. Various kinds of hard and dry fruits, found in the stomachs of the young Guacharos, a sort of nocturnal bird. The fruits are a celebrated South American remedy in Ague.

SEMILUNAR, *Semiluna'ris*, (F.) *Demilunaire* ou *Semilunaire*, from *semi*, and *luna*, 'the moon.' Having the shape of a half-moon.

SEMILUNAR CARTILAGES, *Semilunar fibrocartilages*, *Sigmoid cartilages*, *Cartilag'ines sigmoideæ seu semiluna'res*, are two fibro-cartilages, which exist between the condyles of the os femoris and the articular surfaces of the tibia. The innermost, which is longer from before to behind than transversely, is nearly semicircular. The outermost forms almost an entire circle; an arrangement, in both cases, corresponding to the different surfaces of the tibia. They are thicker at their outer circumference, which is convex, than at the inner, which is concave and very thin. Both are inserted, before and behind, into the spine of the tibia, by means of fibrous fasciæ.

SEMILUNAR GANGLION, — *Ganglion abdominal'e seu splanchnicum seu transversum seu colic'um*, *G. sürréal*, (Ch.), belongs to the great sympathetic. These ganglia present considerable variety, and are deeply situate in the abdomen, above and behind the supra-renal capsules. They correspond, posteriorly, to the pillars of the diaphragm and the aorta. They have the shape of a crescent reversed, often surrounded by other ganglia of a smaller size. From their periphery numerous filaments proceed, which go to the formation of the solar plexus.

SEMILUNAR NOTCH OF THE STERNUM, *Fourchette* — s. *Valves*, *Sigmoid valves*.

SEMILUNULA UNGUIUM, see *Nail*.

SEMI-MAS, *Eunuch*.

SEMIMASCULATUS, *Eunuch*.

SEMIMASCULUS, *Eunuch*.

SEMIMEMBRANEOUS, *Semimembranosus*.

SEMIMEMBRANOSUS, *Semimembraneus*,

(F.) *Ischio-popliti-fémoral*, *Ischio-popliti-tibial*, (Ch.) *Demimembraneus*. A muscle situate at the posterior part of the thigh. It is flat, thin, narrow, and aponeurotic in its upper third; broader and fleshy in the middle, and tendinous below. It is attached above to the tuberosity of the ischium; and below—by a tendon, which has three divisions—to the outer condyle of the femur, and to the posterior and inner part of the internal tuberosity of the tibia. This muscle is a rotator inwards, and a flexor of the leg. It can also bend the thigh upon the leg. In standing it maintains the pelvis in position, and can even draw it downwards.

SEMIMORTUUS, *Semianimis*.

SEMINA CATAPUTLÆ MAJORIS, see *Ricinus communis*.

SEMINA FRIGIDA MAJO'RA. The *greater cold seeds*. The ancients gave this name to the emulsive seeds of the cucumber, melon, gourd, and water-melon.

SEMINA FRIGIDA MINO'RA. The *lesser cold seeds*. The ancients gave this name to the seeds of the lettuce, purslane, endive, and cichory.

SEMINA RICINI VULGARIS, see *Ricinus communis*.

SEMINAL, *Spermatic* — s. *Filaments*, *Spermatozoa* — s. *Fluid*, *Sperm* — s. *Granules*, see *Granule*.

SEMINALE MEMBRUM, *Penis*.

SEMINALIS, *Spermatic*.

SEMINERVOSUS, *Semi-tendinosus*.

SEMIORBITALIS, *Orbicularis oris*.

SEMIPESTIS, *Typhus*.

SEMINEX, *Semianimis*.

SEMINIA MORBORUM, *Predisposition*.

SEMINIFEROUS, from *semen*, *seminis*, 'sperm,' and *fero*, 'I carry.' *Spermatoph'orous*.

An epithet given to the vessels which secrete and convey the seminal fluid.

SEMINIST, Animalculist.

SEMINIUM, Sperm.

SEMIOPGRAPHY, Semeiography.

SEMIOLGY, Semeiology.

SEMIION, Sign.

SEMIOTICE, Semeiotice — s. Faciei, Physiognomy.

SEMIPLÉGIA, Hemiplegia.

SEMISEXUM, Hemiection.

SEMI SICILICUS. A weight of one drachm or three scruples. — Rhodius.

SEMI SIDERATIO, Hemiplegia.

SEMI SIDERATUS, see Hemiplegia.

SEMI S OMNIS, Coma.

SEMI S OPTUS, Coma.

SEMI S OPORUS, Coma.

SEMI SPECULUM. An instrument used for dilating the incision made into the neck of the bladder in the operation of lithotomy. — Fabricius Hildanus.

SEMI SPINALIS COLLI, *Semispinalis seu transverso-spinalis seu spinalis seu transversalis colli*, *Articulo-spinalis*, *Spinalis cervicis*, *Transverso-spinalis*, (F.) *Demi-épineux ou transversaire-épineux du cou*. This muscle arises from the transverse processes of the six uppermost dorsal vertebrae, by an equal number of distinct tendons, which run obliquely under the complexus, and is inserted in the spinous processes of all the cervical vertebrae, except the first and last. Its action is, to extend the neck obliquely backwards and to one side.

SEMI SPINALIS DORSI, *Semi-spinalis externus*, *Transverso-spinalis dorsii*, *Semi-spinalis*, (Riolan), *Transverso-spinalis*, (F.) *Demi-épineux du dos*, *Transverso-épineux du dos*. This muscle arises from the transverse processes of the seventh, eighth, ninth, and tenth dorsal vertebrae, by as many distinct tendons, which soon grow fleshy; and then again become tendinous, and are inserted into the spinous processes of the sixth or seventh uppermost dorsal, and two lowest cervical vertebrae, by as many tendons. Its action is, to extend the spine obliquely backwards.

SEMI SPINALIS EXTERNUS, *Semi-spinalis dorsii* — s. *Internus*, *Transversalis dorsii* — s. *Spinatus*, *Longissimus dorsii*, *Semi-spinalis dorsii*.

SEMI SPINATUS COLLI, *Semispinalis colli*.

SEMI S SIS, *Semis*. The half of a pound, ounce, drachm, &c.

SEMITENDINOSUS, *Semi-nervosus*, (F.) *Ischio-cræti-tibialis*, *Ischio-prætibialis*, (Ch.) *Demitendineux*, *Deminerveux*. This muscle is situate at the posterior part of the thigh. It extends obliquely, downwards and inwards, from the tuberosity of the ischium — to which it is attached by a flat tendon, common to it and the biceps — as far as the posterior and inner part of the tibia, to which it is attached by means of another round tendon. Its fibres are commonly interrupted by an aponeurosis. Its uses are the same as those of the semimembranosus.

SEMITERTIANA, Hemitritæa.

SEMIUNCIA, Hemioncion.

SEMI VIR, Eunuch.

SEMI VIVUS, Semianimis.

SEMOLA, BULLOCK'S; a farinaceous article of diet, which is recommended as highly nutritious and easily digestible, consequently well adapted for infants and invalids, is said to consist of the gluten of wheat, with a small proportion of starch. Semola and Semolina resemble each other greatly. They appear, indeed, to be identical.

SEMOLINA, *Manna croup*. This may be made by grinding wheat in an apparatus, separating the flour from the middlings, dressing the

latter four different times in a bolting mill, and sifting through parchment sieves, until the product is perfectly freed from bran. In Poland, it is called *Cracow groats*.

Semolina is generally imported into England from the Baltic, and in Russia is said to be manufactured from buckwheat. It has a granular form like some kinds of oatmeal, a dingy white colour, a farinaceous, somewhat insipid, taste, and is well adapted for the diet of children affected with derangement of the bowels. A recent analysis makes it consist of the gluten of wheat with a certain proportion of the starch, part of the latter having been removed.

SEMPERVIVUM ACRE, Sedum.

SEMPERVIVUM TECTORUM, *S. majus seu tectorum*; from *semper*, 'always,' and *vivo*, 'I live,' because always green, *Aizoön*, *Beichalalen*, *Buphthalmus*, *Barba Jovis*, *Agriosin'ara*, *Andrachakara*, *Chrysospermum*, *Houze-Leek*, *Seugreen*, *Ficus*, *Fonets*, (Prov.) *Cypheh*, *Jupiter's Beard*, (F.) *Joubarbe des toits*. The leaves have a mild, subacid austerity, and are often applied to bruises and old ulcers.

SEMUDA, Betula.

SEMU NCIA, Hemioncion.

SENA, Cassia senna.

SÉNÉ, Cassia senna — s. d'Amérique, Cassia Marilandica — s. *Faux*, *Colutea* — s. *Indigène*, *Colutea* — s. *des Prés*, *Gratiola officinalis*.

SENEBIERA CORONOPUS, *Cochlearia coronopus*.

SENECA OIL, Petroleum, (New York.)

SENE'CIO, *S. vulgaris*, from *senex*, 'an old man,' in allusion to the grayish down on many species; *Erigeron*, *Cortalon*, *Erigeron*, *Groundsel*, *Fireweed*, (Sc.) *Groundel*, *Swallow*, (F.) *Senegon commun*. Ord. Compositæ. Ser. Syst. Syngenesia Polygamia Superflua. A common plant in Europe; frequently applied, bruised, to inflammations and ulcers, as refrigerant and antiscorbutic.

SENE'CIO AU'REUS, *Golden ragweed*, *Wild valerian*, *Life root*, *Squaw root*; indigenous. Is an excitant, diaphoretic, and diuretic. The dried plant is used in decoction, — ʒss to ʒj of water.

SENECIO HIERACIFOLIUS, *Erechthites hieracifolius*.

SENECIO JACOBÆA, *Jacobæa*, *St. James's Wort*, *Ragwort*, (Sc.) *Bunwede*, *Weebo*, (F.) *Jacobée*. The leaves have a roughish, bitter, subacid, and extremely nauseous taste. A decoction has been used in dysentery. A poultice of the leaves has been recommended in rheumatic affections; and a decoction of the root has often been employed in cases of wounds and bruises.

SENEÇON COMMUN, Senecio.

SENECTA, Senectus.

SENECTA ANGUIM. The cast skin of a serpent. A decoction of it is said to cure deafness!

SENECTA DECREPITA, *Decrepitude* — s. *Extrema*, *Decrepitude* — s. *Summa*, *Decrepitude* — s. *Ultima*, *Decrepitude*.

SENECTUS, *Senect'us*, from *senex*, 'an old man;' *Geras*, *Sen'ium*, *Ætas seni'is seu prove'cta seu mala*, *Old age*, *Senility*, (F.) *Viellesee*. The last period of life, commencing, according to some, at 60 years of age, — according to M. Flourens at 70, — but varying according to numerous circumstances. It is characterized by progressive diminution of the physical and moral faculties.

SENECTUS ULTIMA, *Decrepitude*.

SENEGA, *Polygala senega*.

SEN'EGAL, GUM, *Gummi Senegalen'se seu Sen'eca seu Sen'ega seu Sen'ica*. The exudation from *Acacia seu Mimosa Senegal*. — Ord. Leguminosæ. Ser. Syst. Polygamia Monœcia — which grows in the country of Africa through

which the river Senegal runs. It is in loose or single drops, much larger than gum Arabic, or the gum which exudes from the cherry tree; and is much more difficult of solution than the former.

SENEGINE, see *Polygala senega*.

SÉNÉGRAIN, *Trigonella fœnum*.

SENEKA, *Polygala senega*.

SÉNEVÉ, *Sinapis*.

SENGREEN, *Sempervivum tectorum*.

SENILE, *Seni'lis*, from *senex*, 'an old man.' Relating or belonging to old age: — as '*senile delirium*;' *dotage*.

SENILITY, *Senectus*.

SENNA, *Cassia senna*—s. *Alexandrina*, *Cassia senna*—s. *American*, *Cassia Marilandica*—s. *Bladder*, *Colutea arborescens*—s. *Essence of*, prepared, *Selway's*, see *Infusum sennæ compositum*—s. *Germanica*, *Colutea*—s. *Italica*, *Cassia senna*—s. *Prairie*, *Cassia chamæcrista*—s. *Wild*, *Cassia chamæcrista*, *Cassia Marilandica*.

SENS, *Sense* — s. *Génital*, *Appetite*, *venereal*.

SENSATION, *Sensa'tio*, from *sentire*, *sensum*, 'to feel'; *Æsthe'ma*. The consciousness or cognisance by the brain of an impression caused by an external body on the organs of the senses. (F.) *Sentiment*. It is not necessarily, however, confined to bodies external to us; for we can receive an impression by touching any part of our own body. The body which communicates the impressions needs but to be external to the part impressed. Sensations are divided into *external*, *internal*, and *morbid*. The *external* are communicated by the five organs of the senses. The *internal* are such as occur within the body, and arise from some alteration in the function of the part, for the time being. Hunger and thirst are internal sensations, as well as all the natural wants. *Morbid* sensations may be either internal or external. *Objective sensations* are produced by impressions on the peripheral nerves; as in vision, audition, &c. *Subjective sensations*, are such as originate centrally, or in the encephalon,—as *tinnitus aurium*.

SENSATIONAL, *Sentient*.

SENSE, *Sensus*, *Sen'sio*, *Æsthe'sis*, (F.) *Sens*. Same etymon as *Sensation*. A faculty, possessed by animals, of appreciating impressions from external objects. The senses are five in number: sight, hearing, smell, taste, and touch. Of these, the first two and the last administer more to the intellect; and, hence, have been called by some, the *intellectual senses*; the other two seem more destined for the nutrition of the body: and, hence, have been termed *corporeal* or *nutritive senses*. Other senses have been suggested — as the intense sensation experienced during the venereal act; a sense of heat and cold; a muscular sense, a common sense or *cœnæsthesis*, &c.

SENSES, NERVOUS SYSTEM OF THE, *Medulla oblongata*.

SENSIBILITAS, *Sensibility*—s. *Anomala*, see *Irritable* — s. *Aucta*, see *Irritable* — s. *Morbosa*, see *Irritable*.

SENSIBILITY, *Sensibil'itas*. Same etymon as *Sensation*; *Æsthe'ma*. The faculty of receiving and transmitting impressions, and having the consciousness of them. It thus includes three secondary properties, *impressibility*, (F.) *impressionabilité*, *transmissibility*, and *perceptibility*. Bichat defines it as the property possessed by living bodies, of receiving impressions, whether the individual be conscious of them or not. In the former case, where conscious, he calls the sensibility *animal*: in the latter, where not, he terms it *organic*. This last is common to vegetables and animals, and presides over nutrition, absorption, exhalation, secretion, &c. The other does

not exist in vegetables: it is the origin of the sensations—olfaction, vision, gustation, audition, thirst, hunger, pain, &c. There are few parts of the animal body, but what are sensible — if not in health—in disease. The free extremities of the hair and nails, and the epidermis, are not so. See *Insensibility*.

SENSIFEROUS, *Sentient*.

SENSIO, *Sense*.

SENSITORIUM, *Sensorium*.

SENSORI-VOLIT'IONAL. A term applied to nervous fibres which pass to and from the cerebro-spinal axis, and are respectively concerned in sensation and volition.

SENSO'RIAL, *Sentient*.

SENSORIAL POWER, according to Dr. Wilson Philip, is composed of the two functions of sensation and volition. Dr. Darwin included in it the power of muscular contraction; but it ought not to be extended to acts in which there is not consciousness.

SENSO'RIMUM, *S. communē*, *Or'ganon sensō'rimum internum*, *Sensō'rimum*, *Empō'rimum spirituum*, *Æsthetē'rion*, *Sen'sory*. Same etymon. The common centre of sensations.

SENSORY, *Sensorium*, *Sentient*.

SEN'SORY GANG'LIA. A name given by Dr. Carpenter to a series of ganglionic masses at the base of the brain, which are in direct communication with the nerves of sensation—as the *olfactory*, *optic*, *auditory*, and *gustatory*.

SENSORY NERVES, *Nerves of sensation*, are so called in contradistinction to *Motor* nerves. There are *general sensory nerves* — as those connected with the posterior part of the spinal marrow, and the fifth pair; and *special sensory*, as those of the senses.

SENSUS ÆGRITUDINIS, *Indisposition* — s. *Osmometricus*, *Olfaction* — s. *Perversus*, *Pseudæsthesia*.

SEN'TIENT, *Sen'tiens*, *Sen'sory*, *Ideōg'enous*, *Sensō'rial*, *Sensa'tional*, *Sensif'erous*. Same etymon. Feeling, causing feeling.

SENTIENT EXTREMITIES OF NERVES are their minute terminations in the organs.

SENTINA, *Pituitary gland*.

SENTIMENT, see *Sensation*.

SEPARATO'RIMUM, from *separare*, *separatum*, 'to separate.' A surgical instrument, for separating the pericranium from the skull.

A. *Separatory*, *Vitrum Hypoclepticum*, is a pharmaceutical vessel for separating fluids of different densities from each other.

SEPEDOGENESIS, *Sepedonogenesis*.

SEPEDON, *Putrefaction*.

SEPEDONODES, *Putrid*.

SEPEDONOGEN'ESIS, *Sepedogen'esis*, from *σπένδω*, 'putrescence'; and *γενναις*, 'generation.' A septic disposition or tendency; such as is met with in typhus gravior.

SEPEs, *Se'pes*, 'a hedge,' 'a row,' as *Sepe* seu *Se'ries* seu *Pecten dentium*: — a row of teeth.

SEPEs DENTium, see *Sepe*s.

SEPHIRUS, *Scleritis*.

SE'PIA, *σπία*, *S. officina'lis*, *Se'pium*, *Cuttle-Fish*, (Sc.) *Hose-Fish*, *Slieve-Fish*, (F.) *Sèche*. *Class* Cephalopoda. The osseous part of the cuttle-fish; (F.) *Oe de Sèche*—*Præcip'itans magnūm*—has been used as an absorbent, and is often added to tooth-powders.

SEPLASIA'RIOUS. A name formerly given to one who sold perfumes, ointments, &c. It is derived from *Seplasia*, the name of a public place at Capua, where such preparations were frequently sold. A druggist.

SEPSICHYMTA, *Septochymia*.

SEPSIS, *Putrefaction*.

SEPTÆMIA; from *σπρος*, 'rotten,' and *αἷμα*,

'blood.' A morbid condition of the blood produced by septic or putrid matters.

SEPTAL, *Septalia*, from *Septum*. Relating, or appertaining to, a septum: as 'septal walls of the heart'—those that separate the cavities from each other.

SEPTANA FEBRIS, Fever, septan.

SEPTENARY, *Septenarius*, (F.) *Septenaire*, from *septem*, 'seven.' 'Consisting of seven.' The septenary years of the old believers in critical periods were times at which, it was supposed, important changes might be expected to take place. The grand climacteric was fixed at 63, and it was considered that if a person passed that age, he had well-founded expectations that his life might be protracted to 90.

SEPTENNIAD, Climacteric, (years.)

SEPTERIUS, Septic.

SEPTFOIL, UPRIGHT, Tormentilla.

SEPTIC, *Septicus*, *Septus*, *Seperius*, from *σπνν*, 'to rot', *σπνντος*, 'rotten.' That which produces putrefaction. A substance which corrodes and disorganizes the soft parts without causing much pain.

SEPTIC POISON, see Poison.

SEPTICÆMIA, Septæmia.

SEPTIMESTRIS FÆTUS, see Fœtus septimæstris.

SEPTINERVIA, Plantago.

SEPTOCHYMIA, *Septichymia*; from *σπνν*, 'putrefaction,' and *χυμος*, 'juice.' Putrefaction or putrescence of the humours.

SEPTON, Azote.

SEPTOPYRA, Typhus gravior.

SEPTULUM; diminutive of *septum*. A division between small spaces or cavities—as the *septula* or fibrous cords given off by the mediastinum testis to be inserted into the inner surface of the tunica albuginea.

SEPTUM, (F.) *Cloison*, 'a partition,' from *sepio*, *septum*, 'I fence in.' A part destined to separate two cavities from each other, or to divide a principal cavity into several secondary cavities.

These septa are numerous in the human body. The chief are the following:—

SEPTUM ATRIORUM, *S. Auricularum*. The septum or partition between the auricles of the heart.

SEPTUM ATRICULARUM, *S. Atriorum*—*s. Cerebelli*, Falx cerebelli—*s. Cerebri*, Falx cerebri—*s. Cochleæ auditoriæ*, Lamina spiralis.

SEPTUM CORDIS, *Septum medii cordis*. The partition which separates the two ventricles of the heart.

SEPTUM ENCEPHALI, Tentorium.

SEPTUM LUCIDUM seu *pellucidum* seu *ten'næ* et *pellucidum* seu *medium*, *Speculum*, *S. lucidum*, *Lapis specularis*, *Mediastinum* seu *Diaphragma ventriculorum lateraliu* *cerebri*, (F.) *Septum median* (Ch.), *Cloison transparente*. The soft portion or medullary substance, which separates the two lateral ventricles of the brain from each other. This septum is composed of two laminæ; between which a small cavity exists, filled by a serous fluid, *ventriculus septi pellucidi*, see Ventricles of the Brain. The brothers Wenzel are of opinion that this cavity is lined by a peculiar membrane.

SEPTUM MÆDIAN, *S. lucidum*—*s. Median du cerelet*, Falx cerebelli—*s. Medium cerebri*, *S. lucidum*.

SEPTUM NARIUM, *Interseptum* seu *Dissepimentum* seu *Interfinitum* seu *Discrimen* seu *Imbrex* seu *Diaphragma narium*, (F.) *Cloison des Fosses nasales*. The partition between the nares. It is formed by the vomer, the perpendicular plate of the ethmoid bone, and a cartilage of a triangular shape. These parts are lined by the pituitary

membrane, which receives a considerable number of nerves and vessels.

SEPTUM NASI, see Nares.

SEPTUM PARVUM OCCIPITALE, Falx cerebelli—*s. Pectiniforme*, see Cavernous bodies—*s. Pellucidum*, *S. lucidum*—*s. Scalæ*, Lamina spiralis—*s. Staphylin*, Velum pendulum palati—*s. Tenue* et *pellucidum*, *S. lucidum*—*s. Thoracis*, *Mediastinum*—*s. Transverse*, Tentorium—*s. Transversum*, *Diaphragm*—*s. Vestibuli nervoso-membranaceum*, see Labyrinth.

SEPTUS, Septicus.

SEPUM, Sevum.

SEQUES'TRUM, "the depositing of a disputed thing in the hands of a third person!" The portion of bone, in necrosis, which is dead, and separated from the living bone; acting, whilst retained, as an extraneous body. When the sequester is superficial and small, it is called *Exfolia'tio*. The portion of bone surrounding it is the *Cap'sula sequestra'lis*.

SERAPEUM, Syrup.

SERAPINON, Sagapenum.

SERAPIUM, Syrup.

SERBET, Scherbet.

SÉREUX, Serous.

SÉRICO-CARPUS TORTIFOLIUS, *Coryza bifolia'ta*, *Aster tortifolius*; indigenous; Order, Compositæ: is said to be diuretic.

SERICUM, Silk; from *Seres*—the Thibetans probably—from whom it was first obtained. (F.) *Soie*. The thread spun by the *Bombyx Mori* or *Silkworm*. Silk is used by surgeons for ligatures, &c.

SERICUM ANGLICUM, see Sparadrapum adhæsivum.

SERIES DENTIIUM, see Sepes.

SÉRIEUX, Serious.

SERINGUE, Syringe.

SERIOLA, Cichorium intybus.

SERIOUS, *Serius*, (F.) *Sérieux*, Grave. A term applied to a disease, *Dyspathia*, *Morbus gravis*, and to a symptom that is attended with danger.

SERIS, Cichorium intybus.

SERMOUNTAIN, Laserpitium album.

SERO-COLITIS, see Colitis.

SERO-CYSTIC, *Sero-cysticus*. An epithet for tumours of the female breast, which consist, in the first stage, of one or more membranous cysts, produced perhaps by dilatation of portions of the lactiferous tubes.

SERO-HEPATITIS, see Hepatitis.

SER'OLIN, from *serum* and *olein*. A peculiar fatty matter detected in the blood by Boudet, Lecanu, Sanson, and others.

SÉROSITÉ, Serum.

SEROSITY, Serum.

SEROUS, *Sero'us*, (F.) *Séreau*; from *serum*. Thin; watery. Relating to the most watery portion of animal fluids, or to membranes that secrete them. See Serum.

SEROUS LAYER, see *Tache embryonnaire*—*s. Membranes*, Membranes, serous.

SERPEDO, Psoriasis.

SERPENT, *Serpens*, *Anguis*, *Ophis*; from *serpo*, 'I creep.' A snake.

SNAKE BROTH was, at one time, recommended as a restorative.

Many serpents are venomous; and the bites of some fatal. See Poisons, table of.

SERPENTAIRES NOIRE, *Actæa racemosa*—*s. Rouge*, Polygonum bistorta.

SERPENTARIA, *Aristolochia serpentaria*—*s. Gallorum*, *Arum dracuncul*—*s. Hispanica*, *Scorzonera*—*s. Minor*, *Arum maculatu*—*s. Ni-gra*, *Actæa racemosa*.

SERPENTARIÆ BRAZILIENSIS RADIX, Caineæ radix.

SERPENTINE, see Alembic.

SERPENTINE, Ophites.

SERPENTINUS, Ophites.

SERPES, Herpes.

SERPIG'INOUS, from *serpere*, 'to creep.'

An epithet given to certain affections, which creep, as it were, from one part to another,—'*Serpiginous erysipelas*,' (F.) *Erysipèle serpiginoux*.

SERPIGO, Herpes circinatus, Lichen psoriasis.

SERPILLUM, Thymus serpyllum.

SERPOLET, Thymus serpyllum.

SERPOLLUM, Thymus serpyllum.

SERPYPILLUM, Thymus serpyllum—s. Citratum, see Thymus serpyllum.

SERRA, Saw—s. Amputatoria, Saw, amputation—s. Versatilis, Trepan.

SERRAT'ULA AMA'RA, from *serra*, 'a saw,' owing to its serrated leaves. Family, Cynaracephaleæ. Ser. Syst. Syngenesia Polygamia æqualis. A species of *Sawwort*, recommended in ague.

SERRATULA BEHEN, Centaurea behen.

SERRATUS, Dentate—s. Anticus Minor, Pectoralis minor.

SERRA'TUS MAGNUS. So called from its serrated appearance; from *serra*, 'a saw.' *Serratus major anticus*, *S. major*, (F.) *Costo-basi-scapulaire*, *Costo-scapulaire* (Ch.), *Grand dentelé*. A very broad (especially anteriorly), thin, flat, irregularly quadrilateral muscle, situate at the sides of the thorax. It is attached, before, to the external surface of the first 8 or 9 ribs, by as many distinct digitations; and, behind, to the spinal edge of the scapula. The superior fibres are almost horizontal; the lower become gradually more and more oblique upwards and backwards. This muscle carries the scapula forwards, and causes it to execute a movement of rotation, which directs its inferior angle forwards, and the anterior upwards. When the shoulder is fixed, it raises the ribs.

SERRATUS MAJOR, *Serratus magnus*—s. Major anticus, *Serratus magnus*.

SERRATUS POSTI'CUS INFERIOR, (F.) *Dorso-lombo-costal*, *Lombo-costal* (Ch.), *Petit dentelé postérieur et inférieur*. A broader and thinner muscle than the following, but nearly of the same shape. It is situate obliquely at the lower part of the back; and extends from the spinous processes of the last two or three dorsal vertebrae, and the first two or three lumbar, to the inferior margin of the last four false ribs. This muscle depresses the ribs, and thus concurs in expiration. It is an antagonist to the next.

SERRATUS POSTI'CUS SUPERIOR, (F.) *Cervico-dorso-costal*, *Dorso-costal*, (Ch.) *Petit dentelé supérieur*. A flat, thin, quadrilateral muscle; situate obliquely at the posterior inferior part of the neck, and the superior part of the back. It passes from the posterior cervical ligament, the spinous process of the last cervical vertebra; from the spinous processes of the first two or three dorsal, to the second, third, fourth, and fifth ribs; into which it is inserted by as many digitations. It raises the ribs, and is, consequently, an inspiratory muscle.

SERRE-ARTÈRE, *Presse-artère*.

SERRE-COU (F.), from *serrer*, 'to squeeze,' and *cou*, 'the neck.' An instrument, invented by Chabert, for compressing the jugular vein, in bleeding from that vessel. It surrounds the neck like a collar, and, by means of a pad, exerts the necessary pressure.

SERRE-FINE, (F.), from *serrer*, 'to squeeze,' and *fin*, 'fine.' A small spring forceps, invented by M. Vidal. It is formed of a wire of metal, turned in the middle into two spirals, one in front of the other. This constitutes the spring. Each branch describes nearly the letter S; and one ex-

tremity forms the spiral, the other lays hold of the parts to be united.

SERRE-NEUD (F.), from *serrer*, 'to squeeze,' and *nœud*, 'a knot.' This name has been given to various instruments, employed by the surgeon for tightening a ligature passed round a pediculated tumour, or any part which has to be destroyed by ligature slowly and gradually. Thus, there is the *serre-nœud* of Roderic, the *constrictor* of Herbiniaux, the *serre-nœud* of Desault, &c.

SERRE NŒUD DE DESCHAMPS, *Presse-artère*.

SERTULA CAMPANA, *Trifolium melilotus*.

SERUM, Sperm.

SERUM, 'whey.' *Orrhos*, *Oros*, *Seros'ity*, (F.) *Sérosité*. The most watery portion of animal fluids, exhaled by serous membranes. It is a constituent part of blood, milk, &c.

SERUM OF THE BLOOD, *Hæmydor*, *Orrhos*, *Serum seu Ichor sanguinis*, (F.) *Sérum du Sang*, is the liquid which separates from the blood, when coagulated at rest. It is of a greenish-yellow colour; viscid, slightly coagulable by heat, acids, and alcohol. It is composed of water; chloride of sodium; certain phosphates; and albumen, constantly united to soda, almost in a saponaceous combination.

The fluid which exudes from the albumen of the serum of the blood, when coagulated by heat, is by some called *Seros'ity*.

SÉRUM DU SANG, S. of the blood.

SERUM LACTIS, *Aqua Lactis*; *Serum of Milk*, *Whey*. (Prov.) *Whig*. (F.) *Sérum du lait*, *Petit lait*. That part of milk from which the butter and caseous matter have been separated. It is a transparent, citrine-coloured liquid, containing sugar of milk, mucilage, acetic acid, phosphate of lime, and some other saline substances. It is used as a slightly nutritious diluent and refrigerant. The French have magistral formulæ, the basis of which is whey. They are called *Petite-laits*, or 'wheys.'

SERUM LACTIS HOFFMAN'NI, (F.) *Petit-lait d'Hoffmann*, is a name given to the liquid resulting from treating with boiling water milk which has been evaporated to an almost solid consistence. It is not much used.

RENETT WHEY is made thus: milk 2 pints; rennet $\frac{3}{4}$ ss, infused in a little hot water; mix and keep in a gentle heat for some hours, then strain.—Gray.

SERUM SANGUINIS, Serum of the blood.

SERUM OF SEROUS MEMBRANES, *Serum membranarum serosarum*, *Water of Dropsy*, (F.) *Sérum des membranes séreuses*, *Eau des hydro-piques*. It resembles the serum of the blood,—especially in cases of hydrocele. The water of hydrothorax and ascites contains much less albumen:—sometimes enough to coagulate on the application of heat: at others, not.

SERVICE, MOUNTAIN, *Sorbus aucuparia*.

SERVICE TREE, *Sorbus domestica*.

SERVETTE EN CARRÉ, *Cowrechef*.

SÉSAME, *Sesamum orientale*.

SESAMOID, *Sesamoi'des*, *Sesamo'des*, *Sesamoïdeus*, from *σάμν*, 'a grain of sesamum,' and *ειδός*, 'resemblance.'

SESAMOID BONES, *Ossa seu Ossic'ula sesamoidæa*, are small bones, situate in the substance of tendons, near certain joints. Their number is various. They are commonly more in the male than female. Two are constantly met with beneath the metatarsal-phalangeal articulations of the great toe. Sometimes, the articulations of other bones of the metatarsus with the phalanges of the toes have them likewise. In the hand, there are two before the metacarpophalangeal joint of the thumb. At times, they are

met with in the other metacarpophalangeal articulations. Occasionally, two exist behind the condyles of the femur, in the tendons of the gastrocnemii muscles. A considerable os sesamoides is also met with in the tendon of the peroneus longus. These bones are commonly round, and flattened only on one surface, which is incrustated with cartilage. Their size is very variable, according to the joints in which they exist. Traces of sesamoid bones are not observed in infants: in them, they are replaced by small, cartilaginous concretions. They are developed by a single point of ossification: and are, like the patella (which is a true sesamoid bone,) composed of much cellular substance, covered by a thin layer of compact tissue. The use of these bones is,—to allow the tendon to be inserted into the lever at a greater angle; and thus enable the power to act more advantageously.

Riolan has given the name *Ossa sesamoides* to two bony points sometimes met with;—the one at the outer side of the carotid canal of the temporal bone; and the other at the edge of the cavernous sinus, at the side of the internal carotid artery.

SESAMUM, see *Sesamum orientale*.

SESAMUM ORIENTALE, *S. edule* seu *oleiferum* seu *Indicum*, *Benné*, *Oily Grain*, (F.) *Sésame*, *Jugoline*. *Ord.* Bignoniaceae. The seeds of this African plant are cultivated in Georgia and South Carolina, where the negroes use them with Indian corn as an article of food. They yield a larger proportion of oil than any other vegetable; one hundred weight producing ninety pounds of oil, which is, at least, equal to Florence oil.

Benne oil, *Oleum Sesami*, is used in the Southern States as a laxative. The leaves—*Sesami folia*, (Ph. U. S.)—afford, by infusion, an excellent mucilage.

SESCUN'CIA, *Sesquinci'a*, *Hemio'ion*. An ounce and a half. See *Sesqui*.

SESELI, *Laserpitium siler*—s. *Ægopodium*, *Ligusticum podagraria*—s. *Annum*, *S. tortuosum*—s. *Carum*, *Carum*—s. *Carvi*, *Carum*—s. *Creticum*, *Tordylium officinale*—s. *Foeniculifolium*, *Sison ammi*—s. *Graveolens*, *Apium graveolens*—s. *Meum*, *Æthusa meum*—s. *Patens*, *S. tortuosum*—s. *Pratense*, *Peucedanum silaus*—s. *des Prés*, *Peucedanum silaus*.

SESELI TORTUOSUM, *S. Massilien'se* seu *animum* seu *patens*, *Enanthë striatâ rigida*, *Hartwort* of *Marseilles*. The seeds have a warm taste; and a greater degree of pungency than those of *Laserpitium siler*, which is the *Seseli* of the Pharmacopœias.

SESELIS, *Laserpitium siler*, *Seseli*.

SESQUI. This word, joined with any number, weight, measure, &c., signifies one and a half; as *sesquigranum*, 'a grain and a half.'

SESIQUINCIA, *Sescuncia*.

SETA EQUI'NA, 'horse-hair.' *Hairworm*, *Helminthus Gordii*. A kind of worm, found in stagnant water, from 4 to 6 inches long, and twisted into various knots and contortions; colour pale-brown, with dark extremities. It is common in the intestines of the Laplanders; causing the *Colica Laponica*, in which the gripings are, at times, exceedingly severe.

SETA'CEUM, *Seto*, *Seton*, from *seta*, 'a bristle.' A long strip of fine linen or cotton twist, passed through the skin and areolar membrane, to keep up an issue; or through suppurating cavities, between the fragments of fractured bones, &c., to fulfil certain indications. The term is, also, applied to the issue so established. The operation is simple. It consists in laying hold of a fold of skin, and passing a seton-needle through it, with the thread attached. Every day a fresh

portion of the thread is drawn through the sore. Setons are established in many diseases;—in obstinate ophthalmia, cephalalgia; epilepsy, thoracic and abdominal affections, &c.; but they are not as much used as formerly.

SETARIA ROCELLA, *Lichen rocella*.

SETFAST, see *Furunculus*.

SETO, *Setaceum*.

SETON, *Setaceum*.

SETTERWORT, *Helleborus foetidus*.

SEVADILLA, *Veratrum sabadilla*.

SEVATIO, *Steatoma*.

SÈVE, *Sap*.

SEVENBARK, *Hydrangea arborescens*.

SÈVRAGE, *Weaning*.

SEVUM, *Pinguedo*.

SEVUM, *Sebum*, *Sepum*, *Stear*, *Adips ovilli*, *Sevum vervecinum*. *Suet*, *mutton suet*; *Pinguedo*.

SEVUM PRÆPARATUM, *Sevum* (Ph. U. S.),—*Ovilli Sevum Præparatum*, *Prepared Suet*, (F.) *Graisse de Mouton, Suif*, is formed by cutting suet in pieces; melting over a slow fire, and straining through linen. It is emollient, and is sometimes boiled in milk (3ij to Oj), and is used in diarrhœa and dysentery. Its principal employment is in the formation of plasters.

SEVUM VERVECINUM, *Sevum*.

SEX, *Sexus*, *Genos*, *Physis*. The physical difference between male and female in animals and vegetables:—as the *Male sex*, *Sexus potior* seu *fortior* seu *virilis*; and the *Female sex*, *Sexus sequior*, *S. muliebris*, *Alter Sexus*, 'The Sex.'

SEXTANS, from *sextus*, 'the sixth'; *Sexta pars libræ*. The sixth part of a pound.—Galen.

SEXTARIUS. Same etymon. An ancient measure for liquid and solid substances; the sixth part of a congius. See *Weights and Measures*.

SEX'TULA. Same etymon. *Hexagium*, 'εξαγων.' The sixth part of an ounce.

SEX'UAL, *Sexualis*, from *sextus*, 'sex.' That which relates to the sex; which characterizes the sex.

SEXUAL DISEASES, *Morbi sexuales*, are the diseases of the genital organs; as *SEXUAL FUNCTIONS* are the genital functions.

SEXUAL INTERCOURSE, *Coition*—s. *Organs*, *Genital organs*.

SEXUS, *Sex*—s. *Alter*, see *Sex*—s. *Fortior*, see *Sex*—s. *Muliebris*, see *Sex*—s. *Potior*, see *Sex*—s. *Sequior*, see *Sex*—s. *Virilis*, see *Sex*.

SEYDSCHÜTZ WATER, *Seidschütz water*.

SHACKLE, *Carpus*.

SHACKLE-HAMMED, see *Cnemosecoliosis*.

SHADDOCK. The fruit of *Citrus decumana*, called after the captain of a vessel who first took it to the West Indies from China. It is large; but inferior to the orange in flavour. It keeps, however, longer; and, hence, is useful on long voyages.

SHAKERS, see *Convulsionnaire*.

SHAKES, *Intermittent*.

SHAKING, *Intermittent*, see *Rigor*.

SHALLOT, *Echalotte*.

SHAMPOO'ING, *Champooing*, *Massing*, (F.) *Massage*, *Massement*. A Hindoo process, which consists in pressing the body of the bather in a peculiar manner; cracking the various joints, and employing blows and friction.

'Kneading' consists in pressing alternately upon a part of the body to arouse the muscles to more energetic action. It is employed at times in dyspepsia.

SHAMROCK, INDIAN, *Trillium latifolium*—s. *Water*, *Menyanthes verna*.

SHANKER. *Chancre*.

SHANNONDALE SALINE SPRINGS. These are within a few miles of Charleston, Jefferson County, Virginia. They are mildly aperient and diuretic.

SHARON SPRINGS, NEW YORK. These springs, which are situate about 10 or 12 miles south of Canajoharie, N. Y., and 45 miles west of Albany, are much frequented. They contain, according to the analysis of Dr. Chilton, sulphate of magnesia, sulphate of lime, chloride of sodium, chloride of magnesium, sulphohydrate of sodium, sulphohydrate of calcium, vegetable extractive matter, and free sulphohydric acid gas. Their temperature is 48° Fahr.

SHEATH, Vagina.

SHEDDING OF THE TEETH, see Dentition.

SHEEP POISON, *Kalmia latifolia*.

SHELLBARK, see Hickory.

SHELL FLOWER, *Trillium latifolium* — s.

Liquid, see Calcei murias — s. Membrane, Membrana testis.

SHEPHERD'S PURSE, *Thlaspi bursa* — s. Sun dial, *Anagallis arvensis*.

SHERBET, Scherbet — s. Orange, see Syrupus aurantii.

SHERMAN'S WORM LOZENGES, see Worm lozenges, Sherman's.

SHERRY, Sack, see Wine.

SHIELD-BONE, Scapula.

SHIELD, WATER, *Nelumbium luteum*.

SHILDER, Humerus.

SHIN, *O'crea*, *Crista tib'ia*, Sax. *reina*. The epine or anterior part of the tibia or leg. It is, also, called *Anticnemion*, and *Crea*.

SHINBONE, Tibia.

SHINGLES, Herpes zoster.

SHINLEAF, *Pyrola*.

SHINLOCK, *Brassica ruca*.

SHIP FEVER, see Typhus.

SHITABED, *Leontodon taraxacum*.

SHIVERING, see Rigor.

SHOCK, see Concussion.

SHOCK, NERVOUS, *Vital Shock*, *Vital Depression*, *Nervous Depression*, *Fatal Sinking*. Sudden or instantaneous depression of organic, nervous, or vital power, often with more or less perturbation of body and mind, passing either into reaction, or into fatal sinking, occasioned by the nature, severity or extent of injury, or by an overwhelming moral calamity. — Copland.

SHOCK, VITAL, Shock, nervous.

SHOOT, Crick-in-the-neck.

SHORT, Sax. *recept*, from *reipan*, 'to cut, to shear,' shored, shor'd, short, (F.) Court. A word used, by anatomists, in opposition to long; and to distinguish parts from each other that have otherwise the same name.

SHORT BONES, for example, are those in which the three dimensions—length, breadth, and thickness—are nearly alike; such are the vertebræ, bones of the carpus, tarsus, &c.

SHORT-OF-PUFF, Short-winded — s. Ribs, see Costa.

SHORTSIGHTED, Myopic.

SHORTSIGHTEDNESS, Myopia.

SHORTWINDED, Pury.

SHOTBUSH, *Aralia spinosa*.

SHOTT, *Pleurodynia*.

SHOULDER, Humerus — s. Blade, Scapula — s. Bone, Humerus.

SHOUP, Haunch.

SHOVEL PICKEREL WEED, *Unisema deltifolia* — s. Pond, *Unisema deltifolia*.

SHOW, see Parturition.

SHOW-CHOO, see Spirit.

SHOWER-BATH, see Bath.

SHRUB, perhaps, from Arabic, *sharab*, 'syrup.' A compound of spirits, lemon-juice, and sugar.

SHRUB, SWEET, *Calycanthus* — s. Sweet-scented, *Calycanthus*.

SHRUMP-SHOULDERED, see Hump.

SHUD'DERING, *Frem'itus*, *Phryg'ma*, from (G.) Schauder, Schauer, (F.) *Frémissement*, *Tressaillement*. A peculiar sensation, felt either externally or internally; and which seems to be the result of a spasmodic movement of the parts in which it occurs. A slight feeling of cold sometimes accompanies it. It is, at times, the result of a moral cause, and is often the precursor of shivering.

SHULL-BANE, Humerus.

SHUMAC, *Rhus coriaria*.

SIAGANTRI'TIS, from *σιγων*, 'the jaw,' *αντρον*, 'a cavity,' and *itis*, denoting inflammation. Inflammation of the lining membrane of the antrum of Highmore.

SIAGON, Maxillary bone.

SIAGO'NAGRA, from *σιγων*, 'the jaw,' and *αγρα*, 'a seizure.' *Arthrit'is maxilla'ria*. A gouty or rheumatic affection of the joint of the lower jaw.

SIALACHUS, Salivation.

SIAL'AGOGUE, *Sialago'gus*, *Sial'ogogue*, *Phyal'agogue*, *Ptyal'magogue*, *Saliv'ans*, *Salivat'um*, *Sialocinet'icus*, *Apophlegmat'icans per-os*, (F.) *Salivant*; from *σιλον*, 'saliva,' and *αγω*, 'I expel.' That which provokes the secretion of saliva. Pyrethrum and mercury are sialogogues. Sialogogues may be of two kinds, those which act through the circulation, and those which act immediately on the salivary organs. Mercury belongs to the former, — pyrethrum to the latter class. The chief sialogogues are armoracia, calamus, mezereum, pyrethrum, tabacum, and zingiber.

SIAL'ICA, from *σιλον*, 'saliva.' Medicines, which affect the salivary glands. — Pereira.

SIALISMUS, Salivation.

SIALOCINETICUS, Sialogogue.

SIALOLITHI, Calculi, salivary.

SIALOLOG'IA, from *σιλον*, 'saliva,' and *λογος*, 'a discourse.' A treatise on saliva.

SIALOMA, Saliva.

SIALON, Saliva.

SIALORRHŒA, Salivation.

SIALOS'CHESIS, from *σιλον*, 'saliva,' and *σχεσις*, 'retention.' Retention or suppression of the saliva.

SIALOSYRINGES, Salivary fistulæ.

SIALOZEMIA, Salivation — s. Mercurialis, Salivation, mercurial.

SIAMESE TWINS, see Twins, Siamese.

SIBARE, Phrenitis.

SIBBENS, *Sivvens*, *Frambæ'sia Scot'tica*. An infectious disease in the mountainous parts of Scotland, viewed to be of the venereal kind. *Sivin*, in the Celtic, means raspberry; and it is said that, in this affection, fungi appear like raspberries. From *sivin*, the inhabitants have made *sievin*, *sibben*, *sibbens*, and *sivvens*. By some, this affection is regarded as a complication of the venereal and itch. In the Orkneys, *sibbens* means itch. — Jamieson.

SIB'ILANT, *Sib'ilus*, *Sib'ilans*, from *sibilo*, 'I hiss. I whistle.' Making a hissing or whistling sound. See *Râle sibilant*.

SIBILISMUS AURIUM, Tinnitus aurium.

SIBILUS AURIUM, Tinnitus aurium.

SICCANT'IA, from *sicare*, 'to dry.' *Xeran'tica*, (F.) *Siccatis*. Drying medicines. See Desiccativa.

SICCATIFS, Siccantia.

SICCATIO, Drying.

SICCHA'SIA, *συχχασια*. 'disgust.' A distressing disgust for food; such as is experienced by women with child.

SICIL'ICUM. A Roman weight of about two brachms.

SICK, Sax. *reoc*, *Æger*, *Ægro'tus*, *Nos'eros*, *Noso'des*, *Jacens*, *Laborio'sus*, *Labo'rans*, (F.) *Malade*. Labouring under disease; *Diseas'ed*, *Distem'pered*, *Ill*, *Ailing*, *Disor'dered*. Such is its acceptance in old English, and generally in the United States. In most parts of the United States, if a patient is affected with a slight indisposition, he is said to be *sick*; if with one more severe, he is said to be *ill*; (Prov.) *bad*, *badly*. In England, it most commonly means,—affected with disorder of the stomach, or nausea.

A sick person, *Æger*, *Ægro'tus*, (F.) *Malade*, who is under the charge of a physician is said to be a *patient*, or the patient of the physician. At times, but rarely, patient is used for a sick person in the abstract.

SICK STOMACH, Milk sickness.

SICKLY, *Morbo'sus*, *Valetudina'rins*, *Adyn'atos*, *Nosac'eros*, *Noso'dea*, *Morbose*, *Mor'bulent*, *Unhealth'y*, (So.) *Hough*, (F.) *Maladif*. Subject to be sick, or in feeble health. Ailing.

SICKNESS, Disease, Plague — *s.* Falling, Epilepsy — *s.* Mountain, Puna — *s.* River, Milk sickness — *s.* Sea, Nausea marina — *s.* of the Stomach; Vomiting — *s.* Swamp, Milk sickness.

SICLIUM, Sicileum.

SICUA, Cupping-glass.

SICULA, Beta, Penis.

SICYEDON, Cauldon.

SIDA ABUTILON, Abutilon Avicennæ.

SIDE, PAIN IN THE, Pleurodynia.

SIDERA'TIO, *Sydera'tio*, *Sidera'tion*, from *sidus*, *sideris*, 'a star'; because the condition was thought to be produced by the influence of the stars. The state of one struck suddenly, without apparent cause, and as if by the influence of the stars or planets — *Planet-struck*. The ancients comprised under this name, different morbid conditions, such as paralysis, apoplexy, and gangrene.

SIDERATIO, Apoplexy, Asphyxia, Phrenitis — *s.* *Ossis*, *Spina ventosa*.

SIDERATION, Astrobolismos, Sideratio.

SIDERION, Ferramentum.

SIDERITES, Magnet.

SIDERITIS, Magnet, Teucrium chamæpitys.

SIDEROKRENÆ, Water, mineral, (chalybeate.)

SIDEROPEGÆ, Water, mineral (chalybeate).

SIDEROS, Ferrum.

SIDEROXYLON, Chrysophyllum cainito.

SIDE-SADDLE-PLANT, Sarracenia.

SIDHEE, see Gunjah.

SIDMOUTH (CLIMATE OF). Sidmouth is in Devonshire, England, on the sea-beach, at the mouth of the Sid. It is much frequented by invalids, both during the summer and winter. The climate is, however, damp, and subject to sea-fogs.

SIEGE, Anus.

SIELISMOS, Salivation.

SIELON, Saliva.

SIEVE, see Cribration — *s.* Drum, see Cribration.

SIFFLEMENT (F.), from *siffler*, 'to whistle.' Whistling.

SIFFLEMENT MODULÉ, *Chant des artères*, *Bruit musical*, 'modulated whistling, or music of the arteries.' A kind of sound heard on auscultation, resembling, at times, the humming of certain insects. Various musical sounds, (F.) *Bruits musicaux*, are described from fancied resemblances:—as *Bruit de pialement*, resembling the clucking of the hen;—*Bruit de jappement*, resembling the barking or yelping of a young dog, &c.

The sound is chiefly met with in those labouring under chlorosis.

SIGAULTIAN SECTION, Symphyseotomy.

SIGE, Taciturnity.

SIGH, Suspirium.

SIGHT, Vision — *s.* Askew, Dysopia lateralis — *s.* False, Metamorphopsia, Pseudoblepsia — *s.* Feebleness of, Amblyopia.

SIGILLUM SALAMONIS, Convallaria polygonatum — *s.* Virginitatis, Hymen.

SIGMATOID, Sigmoid.

SIGMOID, *Sigmoid'al*, *Sig'matoid*, *Sigmoï'des*, *Sigmo'des*, *Sigmoï'deus*; from *sigma* and *eidōs*, 'resemblance.' That which has the form of the Greek S or C.

SIGMOID CARTILAGES, Semilunar cartilages.

SIGMOID CAVITIES OF THE ULNÆ, (F.) *Cavités ou Fosses sigmoïdes*, are two notches at the upper part of that bone,—one for the reception of the humerus; the other for that of the radius.

SIGMOID FLEXURE OF THE COLON, *Flexu'ra sigmoï'dea* seu *ili'aca*, is a turn the colon takes immediately before its termination in the rectum.

SIGMOID FOSSA (of the temporal bone), Fossa sigmoidea.

SIGMOID OR SEMILUNAR VALVES, *Valvula Sigmoï'dea*, *Proces'sus Sigmoï'des*, (F.) *Valvules sigmoïdes* ou *semilunaires*, are three valvular folds, at the commencement of the pulmonary artery and aorta. These valves have the form of a crescent, when applied against the parietes of the vessel. When the blood is sent from the ventricles by their contraction, the valves are applied against the sides of the vessel; but, during the diastole of the ventricle, they are depressed, and prevent the blood from returning to the heart. At the middle of the convexity of each valve is a small, hard, triangular granulum, called *Corpus'culum* seu *Glob'ulus Aran'tii*, which completely closes up the passage. It is, also, called *Corpusculum Morgagnii* and *C. Sesamoï'deum*, (F.) *Globule d'Arantius*.

SIGN, *Signum*, *Semi'on*, (F.) *Signe*. Any present or past circumstance, afforded by the examination of a patient, or of matters concerning him, whence a conclusion may be drawn regarding the nature and seat of his disease. The phenomena which precede the disease are called *anæsthetic* or *comment'orative signs*;—those which accompany it are termed *diagnostic*, if they reveal the nature or seat of the disease,—*prognostic*, when they indicate its probable duration and termination. See Symptom.

SIGNA ASSIDENTIA, see Assidens.

SIG'NATURE, *Signatu'ra*, *Signa'tio*, from *signum*, 'a sign'; *Episphagiam'cs*. This term was applied in the Eastern countries, in the middle ages, to mystic characters of good or bad augury, with which it was pretended that every one was marked by the star under which he was born.

SIGNATURES OF PLANTS meant certain peculiarities in their external formation or colour, which indicated that they were adapted for particular diseases;—as in the case of the Lungwort, Liverwort, &c.

SIGNES AVANT-COUREURS, Precursory signs—*s.* *Præcursæurs*, Precursory signs.

SIGNUM, see Parturition — *s.* Characteristicum, see Pathognomonic — *s.* Conjunctum, see Pathognomonic — *s.* Diagnosticum, see Diacritica signa — *s.* Morbi essentialia, see Pathognomonic.

SILACH, *Silac*. An Arabic word, used to designate thickening of the eyelids, *Crassit'ies Palpebrarum*; and, also, *Alopecia*.—Forestus, Stephanus.

SILENCE, see Murmur, respiratory.

SILENE CRASSIFOLIA, Cucubalus behen—

s. Inflata, Cucubalus behen — *s. Thorei*, Cucubalus behen.

SILENE VIRGINICA, *Ground Pink*, *Wild Pink*, *Catch Fly*. *Ord. Caryophyllaceæ*. The root has been employed in decoction, as an efficacious anthelmintic.

SILENE PENNSYLVANICA probably has the same properties.

SILER LANCIFOLIUM, *Laserpitium siler* — *s. Montanum*, *Laserpitium siler*.

SILTQUA, *Cera'tion*; the carat. A weight of about four grains. A pod.

SILQUA ARABICA, *Tamarindus* — *s. Araci aromatici*, see *Vanilla* — *s. Banillæ*, see *Vanilla* — *s. Dulcis*, *Ceratonium siliqua* — *s. Vanigliæ*, see *Vanilla* — *s. Vanillæ*, see *Vanilla*.

SILQUASTRUM PLINII, *Capsicum annum*. *SILK*, *Sericum*.

SILK GRASS, *Yucca filamentosa* — *s. Weed*, *Asclepias Sullivantii*, *Asclepias tuberosa* — *s. Weed*, common, *Asclepias Syriaca* — *s. Weed*, flesh-colored, *Asclepias incarnata* — *s. Weed*, swamp, *Asclepias incarnata* — *s. Worm*, see *Sericum*.

SILLOX, *Suleus* — *s. Horizontal de la veine ombilicale*, see *Liver* — *s. Transversale*, see *Liver* — *s. de la Veine cave inférieure*, see *Liver* — *s. de la Veine porte*, see *Liver*.

SILLONNEMENT, Segmentation.

SILQ, *Camus*.

SILPHIUM, *Laserpitium*.

SILPHIUM LACINIATUM, *Resin weed*, *Compass plant*, *Polar plant*.

SILPHIUM PERFOLIATUM, *Cup Plant*, *Turpentine Sun Flower*, and

SILPHIUM TEREBINTHINACEUM, *Pra'irie Burdock*; *Nat. Order*, *Compositæ*; indigenous in Ohio and Michigan; and flowering from July to September; yield a fragrant gum which is esteemed stimulant and antispasmodic.

SILURUS GLANIS, see *Ichthyocolla*.

SILUS, *Camus*.

SILVER, *Argentum* — *s. Ammonio-chloride of*, see *Argentum* — *s. Chloride of*, see *Argentum* — *s. Chloruret of*, see *Argentum* — *s. Cyanide of*, see *Argentum* — *s. Cyanuret of*, see *Argentum* — *s. Iodide of*, see *Argentum* — *s. Ioduret of*, see *Argentum*.

SILVER LEAF, *Magnolia macrophylla*.

SILVER, *Muriate of*, see *Argentum* — *s. Nitrate of*, *Argenti nitras* — *s. Oxide of*, see *Argentum*.

SILVER AND AMMONIA, *CHLORIDE OF*, see *Argentum* — *s. and Ammonia*, *chloruret of*, see *Argentum* — *s. and Soda*, *Hyposulphite of*, *Sodæ et argenti hyposulphitis*.

SILYBUM, *Carduus marianus* — *s. Maculatum*, *Carduus marianus* — *s. Marianum*, *Carduus marianus*.

SIMA'BA CEDRON. *Nat. Fam.* *Simarubaceæ*. A South American tree, the seeds of which — *cedron seeds* — have been used as an antidote for snake-bites. They have, also, been proposed in mental diseases and in epilepsy. The scrapings of the wood have been used as a substitute for sulphate of quinia.

SIMABA QUASSIOIDES, *Nima quassioides*.

SIMAROUBA, *Quassia simarouba* — *s. Bitter*, *Quassia simarouba* — *s. Fauri*, *Malpighia mourella*.

SIMARUBA, *Quassia simarouba*.

SIMARUBA OFFICINALIS, *Quassia simarouba*.

SIMILATIO, Assimilation.

SIMITIUM, *Plumbi subcarbonas*.

SIMO, *Camus*.

SIMONEA FOLLICULORUM, *Acarus folliculorum*.

SIMOTRACHELUS, *Bysauchen*.

SIMPLER'S JOY, *Verbena hastata*.

SIMPLES. Medicinal herbs. One skilled in simples is called a *Simplist*, a *Rhizomatist*.

SIMPLIST, see *Simples*.

SIMULATED DISEASES, Feigned diseases.

SIMULATI MOREI, Feigned diseases.

SIMUS, *Camus*.

SINA LEVANTICA, *Artemisia santonica*.

SINAPELE'ON, from *σινάπι*, 'mustard,' and *ελαίου*, 'oil.' Oil impregnated with mustard.

SINAPI, *Sinapis*.

SINAPIS, *S. nigra* seu *cer'ua* seu *Olivaria'na*, *Brassica nigra*, *Napus*, *Eru'ca*, *Sinapi's*, *Sinapi*, *Sine'pi*, *Nap'ium*, *Napy*, *Common Black Mustard*, (F.) *Moutarde noire ou grise*, *Senevé*, *Sauve*. *Ord. Cruciferae*. *Sec. Syst.* *Tetradynamia Siliquosa*. The seeds are inodorous, when entire; but, when bruised, the colour is very pungent; taste bitterish and acrid. They yield their virtues to water, and afford a bland oil by expression. The seeds are stimulant, diuretic, emetic, and rubefacient. They are sometimes recommended to be swallowed whole, in dyspepsia, torpor of the bowels, &c. Externally, they are applied in the form of cataplasm. See *Sinapism*.

SINAPIS ALBA, *S. foliosa*, *Bonannia officinalis*, *Napus Leuco-sinapi's*, *White Mustard*, is also used medicinally. It is somewhat less pungent than the black. The distilled oil — *Oleum Sinapis* — is a powerful stimulant and rubefacient. In the pharmacopœias of the United States and London, *Sinapis* is the seeds of *Sinapis nigra* and *Sinapis alba*.

Whitehead's Essence of Mustard consists of oil of turpentine, camphor, and a portion of spirit of Rosemary: to which is added a small quantity of flour of mustard. It is a celebrated embrocation for rheumatism, &c.

Whitehead's Essence of Mustard Pills consist of yellow resin and balsam of tolu enough to make into pills.

Mustard Whey is made by taking bruised mustard seeds 3ss, and milk Oj. Boil together until the milk is curdled, and strain.

This whey is used in dropsy.

SINAPIS ARVEN'SIS, *S. Timoria'na*, *Charlock*, *Wild Mustard*, *Garlock*, *Chadlock*, *Rush*, (Sc.) *Runches*, *Scauldries*, *Skelloch*, *Skellie*. The seeds are sometimes substituted for black mustard.

SINAPIS CER'NUA, *Sinapis* — *s. Eruca*, *Brassica eruca* — *s. Foliosa*, *S. alba* — *s. Nigra*, see *Sinapis alba* — *s. Oliveriana*, *Sinapis* — *s. Tamoriana*, *S. arvensis* — *s. Tuberosa*, *Brassica rapa*.

SINAPISIS, *Bole Armenian*.

SIN'APISM, *Sinapis'mus*, from *sinapis*, 'mustard.' A cataplasm, of which mustard forms the basis, which is used for exciting redness, and acting as a counter-irritant. It is prepared by mixing flour of mustard and vinegar together to the due consistence. See *Cataplasma sinapis*.

SIN'CIPIUT, *Sincipital Region*, *Syn'ciput*, *Hemiceph'alum*. The French use it synonymously with *vertex*, *Mesocranium*, *Mesocranium*. By some, it is applied to the frontal region or forehead, or forehalf of the cranium, *Hemiceph'alum*, *Hemiceph'alum*. The parietal bones have been called *sincipital bones*, *Ossa sincip'itis*, (F.) *Ossa du sinciput*.

SINDON, *σινδών*, very fine linen. A small piece of rag, or round pledget, supported by a thread at the middle, and introduced into the opening in the cranium made by the trephine.

SINEPI, *Sinapis*.

SINEW, *Tendon*.

SINGLETON'S GOLDEN OINTMENT is formed of sulphuret of arsenic (realgar,) and lard or spermaceti ointment.

SINGULT'OUS, *Lygmo'des*, *Lygo'des*, *Lyngo'des*, *Lynggo'des*, (F.) *Singultæus*, from *singultus*, 'hicough.' Relating to or affected with hicough.

SINGULTUS, *Lynx*, *Lygmus*, *Spasmolygmus*, *Clonus* seu *Pneusis singultus*, *Hiccough*, *Hiccup*, (Prov.) *Heuks*, *Yoke*, (Sc.) *Eskin*, *Eeskin*, (F.) *Hoguet*. A noise made by the sudden and involuntary contraction of the diaphragm, and the simultaneous contraction of the glottis, which arrests the air in the trachea. It is a symptom of many morbid conditions, and especially of gangrene, but occurs, frequently, in persons otherwise in good health.

SINISTRAD, from *sinister*, 'to the left hand.' A term used by Dr. Barclay, adverbially, to signify 'towards the sinistral aspect.' See *Mesial*.

SINISTRAL ASPECT, see *Mesial*.

SINKING, **FATAL**, **Shock**, nervous.

SINUEUX, *Sinaous*.

SINUOSITAS, *Sinus*.

SIN'UOUS, *Sinu'ous*, (F.) *Sinueux*, from *sinus*, 'a bay or curve.' An epithet given to certain ulcers, and especially to certain fistulæ, which are tortuous.

SINUS, *Sinu'ositas*, *Colpos*, *Abcon'sio*, *Eurycho'ria*, *Reces'sus*, (F.) *Golfe*, 'a cavity.' Any cavity, the interior of which is more expanded than the entrance. In this sense, anatomists have applied the term to cavities in certain bones of the head and face. It has been, also, applied to certain venous canals, into which a number of vessels empty themselves; such as the sinuses of the dura mater and of the spine; the sinus of the vena porta; uterine sinuses, &c.;—and to the bosom.

In *surgery*, it means a long, narrow, hollow track, leading from some abscess, diseased bone, &c., (F.) *Clapier*.

Sinuses of bones form part of the nasal cavities. They are the frontal, sphenoidal, maxillary, &c. The name has also been given, occasionally, to the ethmoid and mastoid cells.

SINUS ACUSTICUS, Auditory canal, internal—*s. Ampullaceus*, *Cavitas elliptica*—*s. Aortici*, Sinuses of Valsalva—*s. Arteriosus*, *s. pulmonalis*—*s. Basilar*, *Sinus transversus*—*s. Cavernæus*, *Cavernous sinus*—*s. Cavernosi uteri*, *Sinus uterini*—*s. Cavernosus*, *Cavernous sinus*—*s. Cerebri*, Ventricles of the brain—*s. Choroidien*, *S. quartus*—*s. Circular*, of Ridley, *Sinus coronarius*—*s. Circularis*, *S. coronarius*—*s. Communis*, see *Semicircular canals*—*s. Coronaire du cœur*, see *Cardiac veins*.

SINUS CORONARIUS, *S. circula'ris* seu *circula'ris Rid'leyi*, *Circular sinus* of Ridley, *Coronary sinus*. A venous canal, which surrounds, in a more or less regular manner, the pituitary fossa and gland, emptying itself into the corresponding cavernous sinus by each of its extremities.

SINUS CORONARY, *Sinus coronarius*—*s. Coronary* of the heart, *Sinus coronary*, venous.

SINUS CORONARY, VENOUS, *Coronary sinus* of the heart. A name given, on account of its great size, to that portion of the great coronary vein which embraces the left auriculo-ventricular furrow.

SINUS COXÆ, see *Cotyloid*—*s. Droit*, *S. quartus*.

SINUS DEXTER CORDIS, *S. vena'rum cava'rum* seu *veno'sus*, *Saccus vena'rum cava'rum*. The right auricle of the heart:—*A'trium cordis dextrum*. See *Auricula*.

SINUSES OF THE DURA MATER, *S. seu Fis'tulæ seu Cavita'tes seu Ventriculi seu Tri'buli seu Sanguiductus seu Receptacula Duræ matris*, are formed by a separation of the layers of the dura mater, which acts as a substitute for the outer membrane of veins. Bichat admits only four great sinuses, of which all the others seem to be dependencies;—the two lateral sinuses, the straight sinus, and the longitudinal sinus. To the lateral sinuses belong,—the petrous sinuses, superior and inferior; the transverse sinus; the

cavernous sinuses; the coronary sinus, and the occipital sinuses. To the straight sinus belongs the inferior longitudinal. The longitudinal sinus, properly so called, has no sinus under its dependence.

SINUS FALCIFORMIS INFERIOR, *S. longitudinal*, inferior—*s. Falciformis superior*, *S. longitudinal*, superior—*s. Fourth*, *Torcular Herophili*—*s. Genæ pituitarius*, *Antrum* of Highmore—*s. of the Jugular vein*, see *Jugular veins*—*s. Lacrymalis*, *Lachrymal sac*—*s. Lactiferus*, see *Lactiferous vessels*.

SINUSES OF THE LARYNX, Ventricles of the larynx.

SINUSES, LAT'ERAL, *Sinus transver'si*, *S. magni*, *S. latera'les*, (F.) *Sinus latéraux*. Each of these sinuses, distinguished into right and left, begins at the torcular Herophili or confluence of the sinuses, and terminates at the sinus of the jugular vein.

SINUS LATÉRAUX, Sinuses, lateral.

SINUSES, LONGITUDINAL, *S. Longitudina'les*. These are two in number. 1. **SUPERIOR LONGITUDINAL SINUS**, *S. falcifor'mis superior* seu *triangula'ris*, *Sinus median*, (Ch.) A long, triangular canal; convex above, concave below; which occupies the whole of the upper edge of the falx cerebri. It is broad behind, and much narrower before, where it commences by a *cul-de-sac*, at the crista galli of the ethmoid bone. It passes along the frontal or coronal crest, sagittal suture, and the vertical groove of the occipital bone. It receives several veins from the dura mater; and all those that are spread over the convex and plain surfaces of the two hemispheres of the brain. It discharges itself into the upper part of the torcular Herophili. 2. **INFERIOR LONGITUDINAL SINUS**, *S. falcifor'mis inferior*, *Vena falcis cerebri*. This is much narrower than the last, and occupies the inferior edge of the falx cerebri, from its anterior third to the tentorium cerebelli, where it terminates, commonly by two branches, in the straight sinus.

SINUS MAGNI, Sinuses, lateral—*s. Maxillary*, *Antrum* of Highmore—*s. Median*, *Sinus longitudinal superior*—*s. Median*, see *Semicircular canals*—*s. Muliebris*, *Vagina*.

SINUS OCCIP'ITAL, *Sinus occipita'lis*. These sinuses commence at the sides of the foramen magnum ossis occipitis, and ascend—becoming broader and nearer each other—into the substance of the falx cerebelli, where they frequently unite. Each opens into the inferior part of the torcular Herophili. They receive the veins of the falx cerebelli, of the dura mater which lines the fossæ cerebelli, and of the posterior part of the cerebellum. Haller and Sümmering have given them the name of *posterior occipital sinuses*, to distinguish them from the transverse sinus, which they have called *anterior occipital sinus*.

SINUS OCCIPITALIS ANTERIOR, *Sinus transversus*—*s. Ophthalmic*, see *Cavernous sinus*—*s. Perpendicularis*, *S. quartus*—*s. Ovatus*, see *Crista vestibuli*—*s. Pituitarii frontis*, *Frontal sinuses*—*s. Pocularis*, see *Ejaculatory ducts*—*s. Polymorphus*, *Cavernous sinus*—*s. Portarum*, see *Liver*—*s. Prostatae*, see *Ejaculatory Ducts*—*s. Prædendi*, *Vagina*—*s. Pudoris*, *Vagina*, *Vulva*—*s. Pulmonales*, Sinuses of Valsalva.

SINUS PULMONA'LES, *S. sinis'ter cordis* seu *vena'rum pulmona'lium* seu *arterio'sus*, *Saccus vena'rum pulmona'lium*, *A'trium cordis sinis'trum*. The left auricle of the heart. See *Auricula*.

SINUS OF THE PULMONARY VEINS, Auricle, left.

SINUS QUARTUS, *S. perpendicular'is*, *Straight sinus*, (F.) *Sinus choroidien* (Ch.), *Sinus droit*. This sinus—triangular throughout; wide posteriorly; narrower anteriorly; slightly oblique downwards and backwards—passes along the base of the falx

cerebri, from the termination of the inferior longitudinal sinus to the torcular Herophili. It receives the inferior longitudinal sinus, the veins of the lateral ventricles or *venæ Galeni*, the superior cerebellous veins, &c.

SINUS RENUM, Pelvis of the kidney—s. Rhomboideus, see Ventricles of the brain—s. Rotundus, see Crista vestibuli—s. of the Septum lucidum, Ventricle, fifth—s. Sinister cordis, S. pulmonalis—s. Sphenoidalis, Cavernous sinus—s. Straight, S. quartus—s. Superciliares, Frontal sinuses—s. Terminalis, Circulus venosus—s. Transversi, Sinuses, lateral.

SINUS TRANSVER'SUS, *S. occipitalis anterior*, *Basilar sinus*. This sinus lies transversely at the anterior part of the basilar process of the occipital bone. It forms a communication between the two petrous sinuses, and the cavernous sinus of one side, and those of the other. Its breadth is commonly very great. It is situated between two layers of the dura mater, and presents, internally, a kind of cavernous tissue.

SINUS TRIANGULARIS, S. longitudinal, superior—s. Urogenitalis, Urogenital sinus.

SINUS, UTERINI, *S. cavernosi Uteri*, *Uterine sinuses*. A name given to cavities, formed by the uterine veins in the substance of the uterine parietes. They are very large during pregnancy.

SINUS UTRICULOSUS, see Semicircular canals—s. des *Vaisseaux séminifères*, Corpus Highmori.

SINUSUS OF VALSALVA are three depressions in the aorta—*Sinus aortici*,—and pulmonary artery—*Sinus pulmonales*—immediately opposite the semilunar valves. Their use is to leave the edges of the valves free, so that they can be caught by the reflux blood. Where the aorta is about to send off the first of its large branches, at the top of the thorax, it is of great size, and has been called the *Large sinus of Valsalva*.

SINUS OF THE VENA CAVA, Auricle, right—s. Venarum cavarum, S. dexter cordis—s. Venarum pulmonalium, S. pulmonalis—s. Venosus, S. dexter cordis.

SINUSSES, VERTEBRAL, (F.) *Sinus vertebraux*. Two great venous sinuses, different from those of the dura mater, which extend the whole length of the vertebral canal, from the occipital foramen to the end of the sacrum, behind the bodies of the vertebrae, before the dura mater, and at the sides of the posterior vertebral ligament. At their inner side they communicate with each other by means of branches, which form true *transverse sinuses*. They are situated at the middle of the body of each vertebra, passing under the posterior vertebral ligament. They receive, at their middle part, the veins that arise in the spongy tissue of the vertebrae. By their external side, the vertebral sinuses communicate with the posterior branches of the vertebral, intercostal, and lumbar veins. At their posterior side they receive the veins, which creep on the prolongation of the dura mater that envelops the marrow.

SINUS VULVÆ, Rima vulvæ.

SIPERI, Bebeeru.

SIPERINE, see Bebeeru.

SIPHAR, Peritoneum.

SIPHILIS, Syphilis.

SIPHITA, Somnambulism—s. Prava, Chorea.

SIPHO, Siphon—s. Urinæ, Diabetes.

SIPHON, *Sipho*, σῆφον, 'a tube.' A bent tube, whose limbs are of unequal length, employed in pharmacy for transferring liquids from one vessel to another. With this view, the shorter limb is plunged into the liquid; and air is sucked from the extremity of the lower limb. The pressure of the atmosphere forces the fluid through the tube; and the flow will continue as long as the extremity of the longer limb is below the level

of the fluid in which the shorter limb is immersed.

An exhausting syringe is, at times, attached to the longer extremity of the siphon, by means of which the air can be drawn from it.

SIPHON, PLUNGING. An instrument, used in pharmacy for obtaining a specimen,—at times, a measured specimen,—of any liquid. With this view, the instrument is immersed into liquid, and allowed to remain there until full. The opening at the top is then closed with the thumb; and the contents, thus removed, are allowed to flow out by removing the thumb and permitting the entrance of air. It is a form of the *pipette*.

SIPHONIA CAHUCHU, see Caoutchouc—s. *Elastica*, see Caoutchouc.

SIPHUNCULUS, Syringe.

SIRA, Orpiment.

SIRENS, Symmeles.

SIRIASIS, *Coup de Soleil*, Insolation, Phrenitis—s. *Ægyptiaca*, *Coup de Soleil*.

SIRIUM MYRTIFOLIUM, Santalum album.

SIRIUS, Canicula.

SIROP, Syrup—s. *de Capillaire*, see *Adiantum capillus veneris*—s. *de Coralline*, see *Corallina Corsicana*—s. *Simple*, Mel despumatum.

SIRUP, Syrup.

SISON AMMI, *Ethusa ammi*, *Cnidium tenuifolium*, *Pimpinella laterifolia*, *Seseli fœniculifolium*. Ord. Umbelliferae. Sex. Syst. Pentandria Digynia. The plant that affords the *Ammi verum* of the shops; (F.) *Ammi des Boutiques*. The seeds—*Semen Ammiæ cretici seu veri*—were once used as a carminative.

SISON AMOMUM, (F.) *Amome fauve*, is considered to be excitant and diuretic. It is the *field horewort*.

SISON ANISUM, *Pimpinella anisum*—s. *Podagraria*, *Ligusticum podagraria*.

SISTENS, Epischietus.

SISYMBRIUM ALLIARIA, *Alliaria*, s. *Barbarea*, *Erysimum barbarea*—s. *Erucastrum*, *Brassica eruca*—s. *Menthastrum*, *Mentha aquatica*.

SISYMBRIUM NASTURTIUM, *Nasturtium aquaticum seu officinale*, *Laver odoratum*, *Cratævisium*, *Cressi*, *Cardamine fontana seu nasturtium*, *Water Cress*, (F.) *Cresson de fontaine*. Ord. Cruciferae. Ser. Syst. Tetradymania Siliquosa. Water-cresses are commonly used in salad. They have obtained a place in the materia medica for their antiscorbutic qualities.

SISYMBRIUM OFFICINALE, *Erysimum*—s. *Parviflorum*, S. *sophia*.

SISYMBRIUM SOPHIA, *S. parviflorum*, *Herb Sophia*, *Sophia Chirurgorum*, *Fliz* or *Flux-weed*, (F.) *Sagesse des Chirurgiens*. It was once reputed to be antelmintic, and valuable in hysterical affections. It is not used.

Other species of *Nasturtium*, as N. *PALUSSTRÆ*, *Marsh water-cress*; and N. *AMPHIBIUM*, *Water-radish*, possess similar virtues with *Nasturtium officinale*.

SISYRINCHIUM BERMUDIANUM, *Blue-eyed Lily*, *Physic* or *Scurvy grass*; indigenous; Order, Iridaceae; flowering from June to August. Used as a cathartic by the Thompsonians. A decoction of the root is acrid and cathartic.

SITICULOSUS, Dipsodes, see *Alterative*.

SITIENS, Dipsodes.

SITIS, Thirst—s. *Intensa*, *Polydipsia*—s. *Morbosa*, *Polydipsia*.

SITKA, *MINERAL WATERS OF*. See *Archangel*, New, mineral waters of.

SITOS, Aliment.

SITOS ALIENUS, *Ectopia*—s. *Anomalus seu perversus intestinorum*, *Splanchnectopia*—s. *Mutatus*, *Ectopia*, see *Monster*—s. *Perversus*, *Ectopia*.

SITZBATH, see *Bath*, half.

SIUM, *S. nodiflorum* seu *angustifolium* seu *Ber'ula* seu *erectum* seu *incisum*, *Apium sium*, *Helioscia'dium nodiflorum*, *Ber'ula angustifolia*, *Ber'ula*, *Creeping Water-paranep*, (F.) *Berle nodiflora*. *Ord.* Umbelliferae. *Sex. Syst.* Pentandria Digynia. It is stimulant, and supposed to be diuretic and lithontriptic. It is, sometimes, employed in cutaneous eruptions.

SIUM ANGUSTIFOLIUM, *S. nodiflorum*—s. *Apium*, *Apium graveolens*—s. *Berula*, *S. nodiflorum*—s. *Bulbocastanum*, *Bunium bulbocastanum*—s. *Carvi*, *Carum*—s. *Erectum*, *S. nodiflorum*—s. *Graveolens*, *Apium graveolens*—s. *Incisum*, *S. nodiflorum*.

SIUM LATIFOLIUM is the common *Water-paranep* of the United States. It is said to be poisonous.

SIUM LUNA'RE. The root is eaten by the Chepewyans. It has an agreeable nutty flavour. By the Canadians, it is called *Queue de Rat*. The poisonous roots of *Cicuta virosa*, *C. maculata*, and *C. bulbifera*, are often mistaken for it.—Sir J. Richardson.

SIUM NINSE. The plant whose root is called *Radix Ninsi*, *Ninzin*, *Nindsin*. This root was long supposed to be the same as ginseng, and it possesses similar, though weaker properties. See *Panax schinseng*.

SIUM NODIFLORUM, *Sium*—s. *Silaus*, *Peucedanum silaus*.

SIUM SIS'ARUM, *Skirret*, (F.) *Chervi*, *Girole*, is used as a salad in Europe; and has been thought a useful diet in chest affections.

SIUM VULGARE, *Ligusticum podagraria*.

SIVVEN, *Rubus idaeus*.

SIVVENS, *Sibbens*.

SKAW, *Scall*.

SKEELY, *Strabismus*.

SKELETIA, *Emaciation*.

SKELETOL'OGY, *Sceletolog'ia*, (F.) *Squeletologie*, from *skeleton*, and *logos*, 'a discourse.' A treatise on the solid parts of the body. This part of anatomy comprehends *Osteology* and *Syndesmology*.

SKEL'ETON, *Scel'etum*, *Skeleton*, *Cadaver exsiccatum*; (Old Eng. and Prov.) *At'omy*, *Anat'omy*, (Sc.) *Spaig*, (F.) *Squelette*, *Charpente osseuse*; (*σκελετος*, 'dried up,' from *σκελλω*, 'I dry.') A dried body. The aggregate of the hard parts of the body, or the bones. A skeleton is found in almost all animals, although it may not always be formed in the same manner. In some, as in the crustacea and testacea, it is wholly or partly external, *Exo-skeleton*, *Dermo-skeleton*. In others, as in birds, the mammalia, &c., it is internal, *Endo-skeleton*, *Neuro-skeleton*. It serves as a support for other organs. On it the general shape of the body and of its various parts is greatly dependent. When the bones are united by their natural ligaments, the skeleton is said to be natural, *Scel'etum naturale*; when articulated by means of wires, artificial or articulated, *Scel'etum artificiale*.

SKELETOPE'IA, *Sceletopœ'a*, *Sceletœ'ia*, from *skeleton*, and *ποιω*, 'I make.' The part of practical anatomy which treats of the preparation of bones, and the construction of skeletons. Bones are prepared to exhibit their conformation, structure, chymical composition, mode of development, and changes; the different cavities formed by their union; and to demonstrate their connexions, relations, modes of union, &c.

SKELLIE, *Sinapis arvensis*, *Strabismus*.

SKELLOCH, *Sinapis arvensis*.

SKEVISCH, *Erigeron Philadelphicum*.

SKEVISH, *FLEABANE*, *Erigeron Philadelphicum*.

SKEW SIGHT, *Dysopia lateralis*.

SKIN, *Cutis*—s. *Bronzed*, see *Capsule*, *renal*.

SKIN-BIND, *Induration of the cellular tissue*.

SKIN-BOUND DISEASE, *Induration of the cellular tissue*.

SKIN, *SCARE*, *Epidermis*.

SKIN-TIGHT, *Induration of the cellular tissue*.

SKIOMACHIA, *Sciomachy*.

SKIRRET, *Sium sisarum*.

SKIRRHUS, *Scirrhus*.

SKIRSE, *Scirrhus*.

SKITTER, *Diarrhoea*.

SKEKE, *Phytolacca decandra*.

SKOLEMORA FERNAMBUCENSIS, *Geof. fræa vermifuga*.

SKOLIOSIS, *Scoliosis*.

SKUE-SIGHT, *Dysopia lateralis*.

SKULL, *Cranium*.

SKULL-CAP, *Calvaria*, *Scutellaria galericulata*—s. *Cap*, *maddog*, *Scutellaria laterifolia*.

SKUNK-CABBAGE, *Dracontium foetidum*.

SKUNK-WEED, *Dracontium foetidum*.

SLAB, *Torticollis*.

SLABBERING, *Slavering*.

SLATE, *IRISH*, *Hibernicus lapis*.

SLATERS, *Onisci aselli*.

SLATKAIA TRAVA, see *Spirit*.

SLAVERING, *Slab'bering*, *Slob'bering*, *Driv'elling*, (Old E.) *Pirtling*, from *saliva*. *Ptyalis-mus iners*, (F.) *Bave*. Involuntary flow of saliva, from sluggishness of deglutition, without increased secretion. It is seen in the infant, the aged, and the idiot.

SLEEP, *Sax. rlep*: *Somnus*, *Sompnus*, *Sopor*, *Dormit'io*, *Hypnos*, (F.) *Sommeil*. Temporary interruption of our relations with external objects. A repose of the organs of sense, intellectual faculties, and voluntary motion.

The act of quitting this condition is called *awaking*, (F.) *Éveil*, *Réveil*. Sometimes this is progressive and natural; at others, sudden and starting; (F.) *Réveil en sursaut*,—as in nightmare, affections of the heart, hypochondriasis, &c.

SLEEP, *MAGNET'IC*, *Hypno'sis biomagnet'ica*, is the sleep induced by the so-called animal magnetism. See *Magnetism*, *animal*.

SLEEP, *NERVOUS*, see *Magnetism*, *animal*.

SLEEPLESSNESS, *Insomnia*.

SLEEP-WALKING, *Somnambulism*.

SLIACS, *MINERAL WATERS OF*. *Sliacs* is a favourite bathing-place in Hungary, a few miles from Neusohl. Some of the waters are cold; others, thermal: the former are carbonated chalybeates; the latter are much used in the same cases as thermal springs in general.

SLIDING OF THE FLAP, see *Flap*.

SLIEVE-FISH, *Sepia*.

SLING, (G.) *Schlinge*, *Mittel'la*, *Mit'rimum*, (F.) *Écharpe*. A bandage employed to keep the forearm bent on the arm, in cases of injury done to those parts, in fracture of the clavicle, injury of the hands, &c. A handkerchief is often sufficient for this purpose. It must be folded diagonally, and be tied, by the extremities, round the neck. The French surgeons have used many varieties of *Écharpes* or *slings*; as the *Grande écharpe*, *Écharpe moyenne*, *Petite écharpe*, *Écharpe de J. L. Petit*, &c.

SLIPPERS, *Impatiens*.

SLIT AND TAIL BANDAGE, *Invaginated bandage*.

SLOBBERING, *Slavering*.

SLOE, *Viburnum prunifolium*.

SLOE-TREE, *Prunus spinosa*.

SLOKE, *GREEN*, *Uva latissima*.

SLOUGH, *Eschar*.

SLOVEN WOOD, *Artemisia abrotanum*.

SLOWS, *Milk-sickness*.

SLUG, *Limax*—s. *Sea*, *Biche de Mer*.

SLUMBER, *Somnolency*.

SMALLAGE, *Apium graveolens*.

SMALL-POX, Variola—s. Modified, Varioloid.
SMALT, *Protax'ide of Cobalt, Smautum, Saffre, Zaffre, Azur, Blue Starch.* This oxyd has been given in rheumatism, in the dose of ten to twenty grains. A larger quantity excites vomiting.

SMARAG'DUS, *σμαργδος*, (F.) *Émeraude*. The emerald. It was formerly regarded as a tonic.

SMARTWEED, *Polygonum hydropiper*, P. punctatum.

SMECTICA, Detergents.

SMECTIS, *Cimolia terra*.

SMECTRIS, *Cimolia terra*.

SMEGMA, *σμηγμα*, from *σμηγειν*, 'to wipe, to rub off.' Soap, grease, a salve, a liniment. Also, the secretion from the sebaceous follicles of the skin.

SMEGMA ARTICULARE, Synovia—s. Cutaneum, see Sebaceous glands.

SMEGMA PRÆPUTII, *Sebum præputia'lis*. The fatty secretion from the sebaceous follicles of the glans penis.

SMEGMATORRHŒA, Stearrhœa.

SMEGMORRHŒA, Stearrhœa.

SMELL, SENSE OF, Olfaction.

SMELLING, Olfaction.

SMELLOME'S OINTMENT FOR THE EYES, *Ceratum resinæ*.

SMILACINA RACEMOSA, *False Spikenard*; indigenous; *Ord. Liliacæ*; is said to be alterative, diuretic and diaphoretic.

SMILAX, *σμιλαξ*, *Taxus baccata*—s. *Aspera* Chinenensis, *Smilax China*—s. *Aspera* Peruviana, *Smilax sarsaparilla*—s. *Canellæfolia*, *S. pseudo-China*.

SMILAX CHINA, *S. as'pera Chinen'sis*. *Ord. Smilacæ. Sec. Syst. Dicoica Hexandria*. The root—*China, Ch. orientalis seu ponderosa seu vera, Sankira, Gauquava, China root*, (F.) *Squine, Esquine, Racine de Chine*. Formerly much used in cutaneous and venereal affections.

SMILAX GLAUCA; indigenous; has similar virtues to those of *S. Sarsaparilla*.

SMILAX HEDERÆFOLIA, *S. pseudo-China*—s. *Indica Spinosa*, *Smilax pseudo-China*—s. *Officinalis*, *Smilax sarsaparilla*.

SMILAX PSEUDO-CHINA, *S. In'dica spinosa seu canellæfolia seu hederæfolia, China occidentalis seu spu'ria nodo'sa*, *American or West India China*. In scrofulous disorders, the root has been often preferred to the oriental kind. In other cases, it possesses similar virtues.

SMILAX SARSAPARILLA, *S. officinalis, Sarsaparilla, Sartaparilla, Smilax as'pera Peruviana, Sarza, Sarsa, Carivillan'di, Iva Pecan'ga, Macapathi, Zarsa, Zarsaparilla*, (F.) *Salsepareille*. The root of this and other species of *Smilax* is inodorous; taste bitterish, feculaceous, and fibrous; of a dark colour externally; white within. It is possessed of diuretic and demulcent properties, and has been often employed in the sequelæ of syphilis; when, after a mercurial course, nocturnal pains, ulcerations, &c., continue. It has, also, been advised in scrofula, chronic rheumatism, &c.; but its efficacy is, to say the least of it, a matter of doubt. Dose, ℥j to ʒj of the powder.

It is obtained from Honduras, Brazil, and Peru—the varieties being called respectively, *Honduras*, (F.) *Salsepareille de Honduras*; *Brazil*, (F.) *S. de Brésil ou de Portugal*, and *Peruvian Sarsaparilla*, (F.) *S. de Pérou ou caraque*.

SMILAX TAMNOIDES; indigenous; has similar properties.

SMILE, *σμιλη*. A curved, two-edged bistoury. A knife.

SMILEON, Scalpel.

SMIT, see Contagion.

SMITTLE, see Contagious.

SMYRNA, Myrrha.

SMYRNISMOS, Embalming.

SMYRNium OLUSATrum, *Smyr'nium*, from *σμυρα*, 'myrrh', the odour of the seeds resembling that of myrrh; *Hippocist'num, Mace'rona, Macedonist'ium, Herba Alexandrina, Gri'elium, Agrioselinum, Alexan'ders, Alсандers*. A plant formerly cultivated for culinary purposes. The seeds are bitter and aromatic, and the roots more powerfully bitter. They have been recommended as resolvents, diuretics, emmenagogues, &c.

SNAGGLE-TEETH, *Odontoparallaxis*.

SNAGREL, *Aristolochia serpentaria*.

SNAIL, *Limax*—s. *Shell, Cochlea*.

SNAKEHEAD, *Chelone glabra*.

SNAKE-ROOT, *Garuleum bipinnatum*—s.

Birthwort, Aristolochia serpentaria—s. *Black, Actæa racemosa, Sanicula Marilandica*—s. *Bul-ton, Eryngium aquaticum, Liatriis spicata*—s. *Canada, Asarum Canadense*—s. *Dewitt, Prenanthes*—s. *Evergreen, Polygala paucifolia*—s. *Heart, Asarum Canadense*—s. *Virginiana, Aristolochia serpentaria*—s. *White, Eupatorium ageratoïdes*.

SNAKEWEED, *Aristolochia serpentaria, Helenium autumnale, Polygonum bistorta*—s. *American, Cicuta maculata*.

SNEEZING, from *Sax. nieran*, (G.) *Niesen*, (probably an example of onomatopœia), *Olonus Sternutatio, Sternutatio, Sternutamentum, Sternumentum, Ptarmos*, (Sc.) *Neesing*, (F.) *Éternuement*. A convulsive motion of the expiratory muscles, by which the air is driven rapidly, and rushes sonorously, through the nasal fossæ, carrying with it the mucus and foreign bodies which adhere to the pituitary membrane, and have occasioned the movement.

SNEEZE-WORT, *Achillea ptarmica, Helenium autumnale*.

SNIRELS, Nares.

SNOB, Nasal mucus.

SNORING, *Snore*, (Sax.) *ἰσνora, Rhenchus, Renchus, Rhenxis, Ronchus, Rhonchus, Stertor*, (Sc.) *Ronking* (probably a case of onomatopœia), (F.) *Ronflement*. Noise made in the posterior part of the mouth and nasal fossæ during the movements of inspiration, particularly in sleep, both when the person is in health, and in disease, as in apoplexy. In certain affections of the chest—in some cases of pulmonary catarrh—a kind of snoring may be distinctly heard by applying the ear or the stethoscope to the chest.

SNOT, Nasal mucus.

SNOTH, Nasal mucus.

SNOUT, Nasus.

SNOW, *Sax. ἱναβ, Niz, Chion*, (F.) *Neige*. Water in a solid state, which falls from the atmosphere in white flakes. It is used externally as a tonic and discutient.

SNOWBALL, *Cephalanthus occidentalis*.

SNOWBERRY, *Symphoricarpos racemosus*.

SNOWDROP TREE, *Chionanthus Virginica*.

SNOW-BLINDNESS, Blindness produced by the reflection of the solar light from snow. As a preventive, *Snow Eyes* are used by the Esquimaux. These are made of extremely light wood, with a bridge resting on the nose like spectacles, and a narrow slit for the passage of the light.

SNOW EYES, see Snow Blindness.

SNUFF, *CEPHALIC*, *Pulvis asari composi-tus*.

SNUFFLES, *Teut. Snuffeln* (probably an example of onomatopœia), *Grave-do neonato'rum, Rhinan'gia, Rhinanch'onæ*. Breathing hard through the nose, often owing to accumulation of the mucous secretions,—an affection to which

infants are liable, and which is relieved by oiling the interior of the nostrils.

SNURLE, Coryza.

SNURLES, Nares.

SOAP, Sapo—s. Acid, see Sapo.

SOAP BERRIES, Sapindus saponaria.

SOAP, BLACK, see Sapo—s. Castile, Sapo durus.

SOAP OF THE COCOA-NUT OIL, see Sapo.

SOAP, COMMON OR SOFT, see Sapo—s. Liquid, see Sapo—s. Medicinal, see Sapo—s. Spanish, Sapo durus—s. Yellow, see Sapo.

SOAPWORT, Saponaria.

SOB, from Sax. *reopian*, 'to grieve,' (probably an example of onomatopœia); (F.) *Sanglot*. A spasmodic, sudden, and momentary contraction of the diaphragm, immediately followed by relaxation, by which the little air, that the contraction has caused to enter the chest, is driven out with noise. It is an evidence of corporeal or mental suffering.

SOBBY, Corpulent.

SOBOLES, Epigone.

SOBRIETAS, Temperance.

SOCIA PAROTIDIS, see Parotid.

SOCIETY, *Societas*, (F.) *Société*, from *socius*, 'a companion.' An association for the promotion of some common object.

SOCIETIES, MEDICAL. Associations of medical gentlemen for the promotion of science have been numerous, and productive of valuable results. In the United States, societies exist in many of the States, possessing the power of examining into the qualifications of every one who may be desirous of practising within their particular jurisdiction. In some of the States, however, the qualified and unqualified are permitted to exercise their profession with equal privileges and immunities.

Amongst the medical associations that have issued from time to time useful transactions or publications, are the following: IN AMERICA,—The Massachusetts Medical Society; the Physico-medical, and Medical Societies of the State of New York; the College of Physicians of Philadelphia, and the American Medical Association. IN EUROPE,—The Association of Fellows and Licentiates of the King's and Queen's College of Physicians of Ireland; the Medico-Chirurgical and Phrenological Societies of Edinburgh; the College of Physicians, the Medico-Chirurgical the Medical and the Sydenham Societies of London; the Associated Apothecaries and Surgeon Apothecaries of England and Wales, and the Provincial Medical and Surgical Association; the Royal Academy of Surgery, the Royal Society of Medicine, the Société Médicale d'Émulation, the Société de Médecine, the Royal Academy of Medicine, the Société Médicale d'Observation, and the Société de Biologie, of Paris; the Medico-Chirurgical Academy of Vienna; the Medical Societies of Berlin and Copenhagen, &c., &c. IN ASIA,—The Medical and Physical Society of Calcutta.

Valuable medical essays are likewise contained in the published Transactions of the American Philosophical Society; the Royal Societies of London, Edinburgh, Göttingen, &c.; the Imperial Academy of Sciences of Paris; the Imperial Society of Sciences of St. Petersburg and Vienna; the Royal Academy of Sciences of Turin; the Academia Imperialis Leopoldina Naturæ Curiosorum, &c.

SOCORDIA, Hallucination.

SODA. An Arabic word. Protoxide of sodium, (F.) *Soude*, is obtained by adding caustic lime to a solution of carbonate of soda. The soda in solution may be obtained by evaporation as hydrate of soda. Its action is like that of potassa; but it is not official.

Soda impura, *Barilla*, *Bariglia*, *Barillo*, *lor*,

Soda Hispanica, *Anatron*, *Natron*, *Anatón*, *Nitrum Antiquorum*, *Kelp*, *Aphronitrum*, *Sal alkalinus fixus fossilis*, *Carbonas Sodæ impura*, *Subcarbonas Sodæ impura*, *Anachron*. The mineral alkali is obtained from several sources;—naturally, in Egypt: artificially, from the incineration of marine plants; and from the decomposition of sulphate of soda, &c. It is principally, however, derived from plants growing on the sea-coast;—as the *Salic'la sativa*, *S. soda*, *S. tragus*, *Salicornia herba*, *Salicornia*, *Salicornia Arabica*, &c. The *Barilla* ashes are from the ashes of the *Salicornia Europæa*: the *Kelp*, *Soude de Varec*, from the *Fucus vesiculosus* and other varieties;—the *Turkey Barilla*, *Roquette*, *Cendre du Levant*, *Rochet*, *la Alexandrina*, from the *Mesembryanthemum Copticum*;—the *Allicant Barilla*, *Soda Alicanti* or *seu Alonensis*, from the *Mesembryanthemum nodiflorum*;—and the *Carthage'na Barilla*, from the *Salicornia* and *Salic'la*.

Impure subcarbonate of soda, (F.) *Soude du commerce*, consists of subcarbonate of soda, potash, chloride of sodium, clay and other earthy substances. It is chiefly used for preparing the pure subcarbonate.

SODA, Cardialgia, Cephalalgia, Pyrosis.

SODA ACETATA, *Sodæ Acetas*, *Ac'etate of Soda*, *Terra foliata minerala*, (F.) *Acétate de soude*. This salt is used in similar cases with the acetate of potass.

SODA AÉRATA, *Sodæ carbonas*—s. *Alonensis*, *Soda*—s. *Biborate* of, *Borax*—s. *Bicarbonate* of, *Sodæ carbonas*—s. *Bilate* of, *Picromel*—s. *Borate* of, *Borax*—s. *Boraxata*, *Borax*.

SODA, CARBONATE OF, *Sodæ Carbonas* seu *Bicar'bonas*, *Carbonas Na'tricum*, *Soda Aërata*, *Bicar'bonate of Soda*, *Carbonate of protoxide of Sodium*, (F.) *Carbonate de Soude*, *S. effervescente ou aérée ou crayeuse*. (*Sodæ subcarb. libj, aquæ destill. Oijj*. Dissolve the subcarbonate of soda, and pass carbonic acid through the solution; set it aside to crystallize.) Its use is the same as the subcarbonate, than which it is less caustic, and better adapted for effervescing draughts.

SODA CAUSTICA, Protoxide of Sodium, *Caustic Soda*, (F.) *Soude caustique*, *S. à la Chaux*, *Soude pure*. The impure subcarbonate, treated by quicklime, and consequently deprived of its carbonic acid. It may be used as a caustic like the *Potassa fusa*, and is even preferable to it, as it does not deliquesce and run.

SODA, CHLORIDE OF, *S. Chlorinata*, *Sodæ Chloridum* seu *Chlorure'tum*, *Chlorinated Soda*, *Natrum Oxymuriaticum*, *Oxymuriate of Soda*, (F.) *Chlorure de Soude*, *Chlorure d'oxide de Sodium*. This salt is used as a disinfecting agent; both to remove offensive smells in chambers, and to alter the condition of fetid ulcers. See *Liquor Sodæ Chlorinatæ*.

SODA, CHLORINATED, *Soda*, chloride of—s. *Hispanica*, *Soda*—s. *Hydrochlorate* of, *Soda*, muriate of.

SODA, HYPOSULPHITE OF, *Sodæ hyposulph'is*, (F.) *Hyposulfite de Soude*. This salt has been lately extolled in chronic cutaneous and scrofulous affections. It is given in syrup.

SODA, LITHATE OF, *Urate of soda*—s. *Mephite* of, *Soda*, subcarbonate of.

SODA, MURIATE OF, *S. muriata* seu *muriat'ica*, *Alkali minera'le salinum*, *Sodæ murias*, *Sal communis* seu *culina'ris* seu *fontium* seu *gemmæ* seu *marinum* seu *albus* seu *fossilis* seu *luc'idum*, *Mu'ria*, *Natron muriatum*, *Natrum muriaticum*, *Hydrochlorinas Na'tricus*, *Chlorure'tum So'di*, *So'dii Chlor'idum*, *Common salt*, *Sea salt*, *Rock salt*, *Bay salt*, *White salt*, *Quadrich'i'vret* of *Sodium*, *Hydrochlorate of protoxide of Sodium*, *Hydrochlorate of Soda*, *Chloride of Sodium*, (F.) *Chlorure de Sodium*, *Sel marin*, *Sel de Cuisine*.

In an impure state, this salt is one of the most abundant of natural productions. It is soluble in 3 parts of water, and is tonic, purgative, and an-thelmintic. Externally, stimulant. It is a well-known and useful culinary substance. It has been greatly recommended as an antiperiodic in the dose of $\frac{3}{4}$ ss or $\frac{3}{4}$ i.

SODA, NITRATE OF, Sodæ Nitras, Natron seu Natrum nitricum, Nitras natricus, Al'kali mine-ra'le nitratum, Nitrum Rhomboida'le seu cu'bi-cum, Nitras Natricum, Salpe'tra, Nitrate of pro-tox'ide of So'dium, Protoni'trate of Sodium, Cubic Nitre. Its virtues are similar to those of the ni-trate of potass, for which it has been substituted.

SODA, OXYMURIATE OF, Soda, chloride of.

SODA, PHOSPHATE OF, S. Phosphora'ta, Sodæ phos'phas, Phosphas Natricum, Natrum Oxphos-phoro'des, Sal mirab'ile perlatum, Alkali mine-ra'le phosphatatum, Subphosphate of protox'ide of Sodium, Subprotophosphate of Sodium, Sub-phos'phate of Soda, Triba'ic phosphate of Soda, Tasteless purging salt, (F.) Phosphate de Soude, Sel admirable perle. (Bone burnt to whiteness and powdered, lbx; Sulphuric acid, lbvj; Car-bonate of soda, q. s. Mix the bone with the acid in a covered vessel, add a gallon of water, and stir. Digest for three days, occasionally adding a little water to that lost by evaporation, and frequently stirring the mixture. Pour in now a gallon of boiling water, and strain through linen, gradually adding more boiling water, until the liquor passes nearly tasteless. Set the strained liquor by, that the dregs may subside; from which pour off the clear solution, and boil down to a gal-lon. To this solution, poured off from the dregs and heated in an iron vessel, add by degrees the carbonate of soda previously dissolved in hot water, until effervescence ceases, and the phos-phoric acid is completely neutralized; then filter the liquor, and set it aside to crystallize. Having removed the crystals, add, if necessary, a small quantity of carbonate of soda to the liquor, so as to render it slightly alkaline; then alternately evaporate and crystallize so long as crystals are produced. Lastly, preserve the crystals in a well stopped bottle.—Ph. U. S.) Cathartic, in the dose of from $\frac{3}{4}$ ss to $\frac{3}{4}$ i.

SODA, PHOSPHATE OF, TRIBASIC, S. Phosphate of—s. Powders, see Powders, Soda—s. Springs, see Water, Mineral (gaseous).

SODA, SUBCARBONATE OF, Sodæ Subcarbonas, S. Carbonas (Ph. U. S.), Natron præpara'tum, Sal Sodæ, S. al'kali, Flos Salis, Sag't'men Vitri, Salt of Soda or of Barilla, Subcarbonate of pro-tox'ide of Sodium, Mephite of Soda, (F.) Sous-carbonate de Soude. The impure carbonate, dis-solved in water; the solution filtered and crys-tallized. Soluble in two parts of water at 60°. It is much used as an antacid; and also as a deob-struent in scrofulous and other affections. Dose, gr. x to $\frac{3}{4}$ ss.

The dried Subcar'bonate of Soda, Sodæ Sub-car'bonas exsiccata, Sodæ carbonas exsiccatus, (Ph. U. S.) Carbonas Sodæ siccatum, (F.) Sous-carbonate de Soude desséché, is employed as an antacid and lithontripic. Dose, gr. x to xv.

SODA, SUBPHOSPHATE OF, S. phosphate of.

SODA, SULPHATE OF, Sodæ Sulphas, Natron vitriola'tum, Sal cathar'ticus seu mirab'ilis Glau-be'ri, Al'kali minera'le sulphu'ricum, Natrum sul-phu'ricum, Soda vitriola'ta, Sulphas natricus, Ap'e'riant Salt of Frederick, Glauber's Salts, (F.) Sulfate de Soude. Obtained from the salt that remains after the distillation of muriatic acid; the superabundant acid being saturated with sub-carbonate of soda. It is soluble in three parts of water at 60°; is purgative, and, in small doses, diuretic. Dose, $\frac{3}{4}$ j to $\frac{3}{4}$ x.

The Lynton Glauber's Salts is a Sulphate

of Magnesia and Soda, and is obtained from the mother liquor of sea-water, or by dissolving Ep-som salt in a solution of Glauber's salt.

SODA, TARTRATE OF, Tartrate of potass and Soda, Tar'tarised Soda, Rochelle or Seignette's Salt, Soda Tartariza'ta, Sal Rupellen'sis seu Poly-chres'tum Seigneti seu Seignetti, Natron tartari-za'tum, Tartris potas'sæ et Sodæ, Sodæ et Potas'sæ Tartras (Ph. U. S.), Sodæ potassio-tar-tras, Tartras Potas'sæ et Sodæ, (F.) Tartrate de po-tasse et de Soude. (Sodæ carb. lbj; potass. bitart. in pulv. $\frac{3}{4}$ xvj, Aq. bullient. Ov. Dissolve the carbonate in water; and add, gradually, the bitartrate. Filter the solution and crystallize.—Ph. U. S.) It is a gentle cathartic. Dose, $\frac{3}{4}$ ss to $\frac{3}{4}$ i.

SODA, URATE OF, Urate of Soda—s. Vitrio-lata, S. sulphate of.

SODA WATER, Acidulous water, simple.

SODA AND SILVER, HYPOSULPHITE OF, Sodæ et Argenti Hyposulphitis.

SODÆ ACETAS, Soda, acetate of—s. Arsenias, Arseniate of soda—s. Biboras, Borax—s. Bicar-bonas, Soda, carbonate of—s. Boras, Borax—s. Boras alcalescens, Borax—s. Boras supersaturatus, Borax—s. Carbonas, Soda—s. Carbonas, Soda, subcarbonate of—s. Carbonas exsiccatus, see Soda, subcarbonate of—s. Carbonas impurus, Soda—s. Chloras, Soda, chloride of—s. Hypo-sulphis, Hyposulphite of soda—s. Murias, Soda, muriate of—s. Nitras, Soda, nitrate of—s. Phos-phas, Soda, phosphate of—s. et Potassæ tartras, Soda, tartrate of—s. Potassio-tartras, Soda, tartrate of—s. Subboras, Borax—s. Subcarbonas impura, Soda.

SODÆ ET ARGEN'TI HYPOSULPH'IS, Hyposulph'i-te of soda and silver, (F.) Hyposulphite de Soude et d'Argent. This salt is formed by pouring a solution of hyposulphite of soda over oxide of silver recently precipitated by potassa, until the oxide is entirely dissolved. Crystals are afforded by evaporation. A solution of it is used in the same cases as one of the nitrate of silver.

SODII AURO-TERCHLORIDUM, see Gold—s. Chloridum, Soda, muriate of—s. Chlorure-tum, Soda, muriate of.

SODII IOD'IDUM, I'odide of so'dium. This salt has been prescribed in the same cases as the iodide of potassium.

SO'DIUM, Nat'rium, Natro'nium. The metal-lic basis of soda.

SODIUM, CARBONATE OF PROTOXIDE OF, Sodæ carbonas—s. Chloride of, Soda, muriate of—s. Chlorure de, Soda, muriate of—s. Chlorure d'Ox-ide de, Soda, chloride of—s. Protonitrate of, Soda, nitrate of—s. Protoxide of, Soda—s. Pro-toxide of, hydrochlorate of, Soda, muriate of—s. Protoxide of, nitrate of, Soda, nitrate of—s. Pro-toxide of, phosphate of, Soda, phosphate of—s. Protoxide of, subcarbonate of, Soda, subcarbonate of—s. Quadri-chloruret of, Soda, muriate of—s. Subborate of protoxide of, Borax—s. Subprobo-rate of, Borax—s. Subprotophosphate of, Soda, phosphate of.

SODOMA GALLORUM, Syphilis.

SODOMIA, Buggery.

SODOMY, Buggery.

SOFTENING, Mollities.

SOL-POUR-SOL, Homœozygy.

SOIE, Sericum.

SOLIF, Thirst—s. Excessive, Polydipsia.

SOL, Gold.

SOL-LUNAR, Sol-luna'ris; from sol, 'the sun,' and luna, 'the moon.' An epithet applied to a fancied influence exerted by the sun and moon on man in a state of health and disease.

SOLAMEN INTESTINORUM, Pimpinella anisum—s. Scabiosorum, Fumaria.

SOL'ANOID, Solano'i'des; a hybrid word, from

solanum, 'the potato,' and *aidos*, 'resemblance.' Resembling a potato; potato-like. An epithet applied to a form of cancer, which appears to be intermediate between scirrhus and encephaloid, increasing more rapidly than the former, yet approaching it in firmness.

SOLA'NUM, *S. nigrum* seu *vulga'rè* seu *vulga'tum*, Common or Garden Nightshade, (F.) *Morelle à fruit noir*. Family, Solanaceæ. Sec. Syst. Pentandria Monogynia. A narcotic poison. The *Huile de Morelle* is made, by macerating in oil, for several days, the leaves and fruit of this plant. The oil is anodyne and discutient.

SOLANUM CAROLINEN'SE, *Horse nettle*; indigenous. The juice of the berries has been employed successfully in idiopathic tetanus. [?]

SOLANUM DULCAMA'RA, *S. scandens* seu *ligno'sum*, *Dulcamara*, *D. flexu'osa*, *Glycici'eros*, *Ama'ra dulcis*, *Ερυκθας* of Theophrastus, *Bittersweet*, *Woody Nightshade*, *Bittersweet Nightshade*, *Bittersweet Vine*, *Nightshade Vine*, *Violet bloom*, (F.) *Douce-amère*, *Morelle grimpante*, *Vigne de Judée*. The taste of the stalks is bitter, followed by sweetness; hence its name. It has been used in many diseases, as a diuretic and sudorific, especially in chronic rheumatism, humoral asthma, dropsy, various skin diseases, scrofula, and jaundice. Dose, in powder, ℥i to ℥i.

SOLANUM ESCULENTUM, *S. tuberosum* — s. *Fœtidum*, *Datura stramonium* — s. *Furiosum*, *Atropa belladonna* — s. *Hortense nigrum*, *Atropa belladonna* — s. *Humboldtii*, *S. Lycopersicum* — s. *Le-thale*, *Atropa belladonna* — s. *Lignosum*, *S. dulcamara*.

SOLANUM LYCOPER'SICUM; *Lycoper'sicon* *esculentum*, *L. pomum amo'ris* seu *Humboldtii* seu *pseudolycopersicum*, *Malum lycopersicum*, *Love-apple plant*. The fruit of this, called *Toma'ta*, *Toma'to* or *Love-apple*, (F.) *Pomme d'Amour*, is much eaten in the United States; and, with the French, Spaniards, Portuguese, &c., forms an esteemed sauce.

SOLANUM MAGNUM VIRGINIANUM RUBRUM, *Phytolacca decandra* — s. *Maniacum*, *Atropa belladonna*, *Datura stramonium* — s. *Melanocerasus*, *Atropa belladonna*.

SOLANUM MELON'GENA, *Melongenæ*, *Egg-plant*, *Mud Apple plant*; (F.) *Aubergine*. A native of Asia, Africa, and America. Its oblong, egg-shaped fruit, *Pomum Melon'genæ*, *Malum insa'num*, is prepared in soups and sauces in those countries, like the tomato.

SOLANUM NIGRUM, *Solanum*.

SOLANUM NIV'EUM. A South African plant, the leaves of which, applied to foul ulcers, cleanse them. The fresh juice of the berries and leaves, made into an ointment with lard or fat, is used by the farmers for the same purpose.

SOLANUM PARMENTERII, *S. tuberosum* — s. *Pseudolycopersicum*, *S. Lycopersicum* — s. *Quadrifolium*, Paris — s. *Racemosum Americanum*, *Phytolacca decandra*.

SOLANUM SANCTUM; *Pal'estine Nightshade*. The fruit of this is globular, and is much eaten in the countries where it is a native.

SOLANUM SCANDENS, *S. Dulcamara*.

SOLANUM TUBERO'SUM, *S. esculentum* seu *Parmenterii*, *Lycopersicum tuberosum*, *Papas America'nus*, *Pappus America'nus*, *America'num Tuberosum*, *Batta'ta Virginia'na*, *Bata'tas*; the *Pota'to*, (F.) *Pomme de Terre*. A native of Peru. A nutritious and wholesome vegetable; one of the most useful, indeed, that exists. One hundred pounds of potatoes afford ten pounds of starch, which is equal to arrow-root, as a nutrient and demulcent. It is called *Potato Flour*, *P. Starch*, *French Sago*, *Petit Sague*, *Common Arrow-root*, &c.

SOLANUM URENS, *Capsicum annum* — s. *Vesicarium*, *Physalis*.

SOLANUM VIRGINIA'NUM; indigenous; is said to resemble *Solanum nigrum* in its properties.

SOLANUM VULGARE, *Solanum*.

SOLAR, *Sola'ris*, from *sol*, *solis*, 'the sun.' Relating to the sun; having rays like the sun.

SOLAR PLEXUS, *Plexus sola'ris*, *Centrum communè*, *Cerebrum abdomina'le*. A collection of ganglions and filaments, interweaving and anastomosing with each other repeatedly, and belonging to the system of the great sympathetic. This vast plexus, lying on the vertebral column, the aorta, and the pillars of the diaphragm, has a very irregular shape. The ganglions and filaments composing it are soft, reddish, and almost all issue from the semilunar ganglions. It receives, also, branches from the pneumogastrics. It seems to be chiefly intended for the aorta, and accompanies all its branches; furnishing many secondary plexuses, which are called the *infra-diaphragmatic*, *celiac*, *superior mesenteric*, *inferior mesenteric*, *renal* or *emulgent*, *spermatic*, &c.

SOLDANELLA, *Convulvulus soldanella*.

SOLDIER'S WEED, *Matico*.

SOLE, *Solum*, *S. pedis*, *Solea*, *Basis cor'poris*, *Ichnus*, *Superfici'es planta'ris pedis*, *Peti'na*, *Vestig'ium*, *Pe'dion*, *Peza*, *Pelma*, *Planta'ris superficies pedis*, *Facies conca'va seu inferior* seu *Planum seu Concavum seu Pars inferior* seu *Plant'ies* seu *Planta* seu *Vola pedis*, (F.) *Plante du Pied*. The sole of the foot; the under surface of the foot.

SOLEA, *Sole* — s. *Ipecacuanha*, see *Ipecacuanha*.

SOLÉAIRE, *Soleus*.

SOLEN, *Cradle*, see *Vertebral column*.

SOLENA'RIMUM, from *σωλην*, 'a canal, a pipe.' An instrument of surgery, representing a kind of gutter, in the cavity of which the penis was placed, like a limb in a *cradle*. The word *Solen* itself — *σωλην*, *Canalis* — was used, in antiquity, for a grooved channel or cradle, in which a fractured limb was placed.

SOLENOSTEMMA ARGEL, *Cynanchum oleæfolium*.

SOLE'US, *Gastrocnemius inter'nus*, *Tibio-calcaneus*, (Ch.) (F.) *Soléaire*, from *solea*, 'a sole of a shoe,' owing to its shape. A muscle situate at the posterior part of the leg. It is broad, flat, and almost oval; and is attached, above, to the posterior edge of the fibula, to the oblique line on the posterior surface, and to the inner edge of the tibia. It terminates, below, by a large tendon, which unites to that of the gastrocnemius externus, to form the *tendo Achillis*. This muscle extends the foot on the leg, and conversely.

SOLID, *Solidus*, *St'e'cos*, from *solum*, 'the ground.' A body whose integrant molecules are so united by the force of cohesion, that they will not separate by their own weight. The solids, in the human body, are the bones, cartilages, tendons, muscles, ligaments, arteries, veins, nerves, membranes, skin, &c. The anatomy, or rather study of the solids, is called *Stereol'ogy*, from *στερεος*, 'solid,' and *λογος*, 'a discourse.'

SOLIDA'GO, from *solido*, 'I make firm,' in allusion to its properties as a vulnerary. See *Solidago vigaurea*.

SOLIDAGO SARACENICA, *S. virgaurea*.

SOLIDA'GO RIG'IDA, *Rigid Goldenrod*, *Bones'typtic*; indigenous; of the Composite family, flowering in August and September; is astringent and tonic.

SOLIDA'GO VIRGAU'REA, *S. vulga'ris* seu *Saracen'ica*, *Virga aurea*, *Herba do'rea*, *Cony'za comæ aurea*, *Sym'phytum Petra'm*, *El'ichrysum*, *Consol'ida*, *Saracen'ica*, *Golden rod*, (F.) *Verge*

d'or, Herbe des juifs. The leaves of *Solida'go odo'ra*—*SOLIDA'GO* (Ph. U. S.)—*Fragrant or Sweet-scented Golden-rod*—and the flowers have been recommended as aperients and tonics, in urinary diseases, and in internal hemorrhage.

SOLIDA'GO VULGARIS, *S. virgaurea*.

SOLIDISM, *Patholog'ia solida'ris*. The doctrine of those physicians who refer all diseases to alterations of the solid parts of the body. To the believers in this doctrine, the appellation *Sol'id-iste* has been given. The solidists think that the solids alone are endowed with vital properties; that they alone can receive the impression of morbid agents, and be the seat of pathological phenomena. Solidism has been the prevalent doctrine. It is scarcely necessary to say, that in all our investigations, the condition of both solids and fluids must be regarded.

SOLIUM, *Baignoire*, *Tænia solium*.

SOLUBIL'ITY, *Solubil'itas*, from *solvere*, *solutum*, 'to dissolve.' That property of a body by virtue of which it can be dissolved in a menstruum.

TABLE OF THE SOLUBILITY OF SOME SUBSTANCES IN WATER.

NAMES OF SALTS.	Solubility in 100 parts water.	
	at 60°	at 120°
<i>Acids.</i>		
Benzoic.....	0.208	4.17
Citric.....	133	200
Oxalic.....	50	100
Tartaric.....	very soluble	
<i>Salts.</i>		
Acetate of Potassa.....	100	
— Soda.....	35	
Bitartrate of Potassa.....	1.5	3.3
Borate of Soda.....	8.4	16.8
Carbonate of Ammonia.....	50	
— Magnesia.....	25	
— Potassa.....	50	
— Soda.....	6	40
Chlorate of Potassa.....	35	35
— Soda.....	200	
Chloride of Calcium.....	100	
— Magnesium.....	33	
— Potassium.....	35.42	36.16
— Sodium.....	33	100
Muriate of Ammonia.....	33	100
Nitrate of Potassa.....	14.25	100
— Soda.....	33	100
Phosphate of Soda.....	25	50
Sulphate of Ammonia.....	50	100
— Copper.....	25	50
— Iron.....	50	100
— Magnesia.....	100	133
— Potassa.....	6.25	20
— Soda.....	37	125
Supersulphate of Potassa.....	5.0	100
Tartrate of Potassa.....	25	
— Potassa and Soda.....	20	
— Antimony & Soda.....	6.6	33

SOLUM, *Sole*—*s. Pedis*, *Metatarsus*, *Sole*.

SOLUTÉ, *Solution*.

SOLUTIO, *Solution*—*s. Alexiteria Gaubiana*, see *Disinfection*—*s. Ammonia subcarbonatis*, *Liquor ammonia subcarbonatis*—*s. Arsenicalis*, *Liquor arsenicalis*—*s. Arsenicata*, *Liquor arsenicalis*—*s. Arsenitis kalica*, *Liquor arsenicalis*—*s. Calcis*, *Liquor calcis*—*s. Camphoræ ætheræ*, *Tinctura ætheræ camphorata*—*s. Camphoræ oleosa*, *Linimentum camphoræ*—*s. Continui*, *Solution of continuity*—*s. Cupri ammoniati*, *Liquor Cupri ammoniati*—*s. Ferri nitratis*, *Liquor Ferri nitratis*—*s. Guaiaci gummosa*, *Mistura guaiaci*—*s. Muriatis barytæ*, *Baryta*, *muriate of*—*s. Muriatis calcis*, see *Calcis muriæ*—*s. Muriatis ferri ætheræ*, *Tinctura seu alcohol sulfurico-ætheræ ferri*—*s. Muriatis hydrargyri oxygenati*, *Liquor hydrargyri oxy muriatis*—*s. Potassii iodidi iodureta*, *Liquor iodini compositus*—*s. Resinæ guaiaci*

aquea, *Mistura guaiaci*—*s. Solventis Mineralis*, *Liquor Arsenici chloridi*—*s. Subcarbonatis ammoniaci spirituosus*, *Spiritus ammoniac*.

SOLUTIO SULPHATIS CUPRI COMPOSITA, *Aqua vitriol'ica cæru'lea*, *Sydenham's Styptic Water*, (*Sulph. cupri*, \mathfrak{z} ij, *alum.*, *acid. sulph.*, aa \mathfrak{z} ij, *aqua*, \mathfrak{z} viii; dissolve, and filter.) A styptic, and largely diluted, an astringent collyrium.

SOLUTIO, *Solutio*, *Lysis*, (F.) *Dissolution*, from *solvere*, *solutum*, 'to dissolve.' An operation which consists in dissolving a solid body in a menstruum. Likewise, the product of such operation—*Solutum*, (F.) *Soluté*. It means, also, with many, the termination of a disease:—with others, a termination accompanied by critical signs;—and, with others, again, it is synonymous with crisis.

SOLUTION OF ACETATE OF AMMONIA, *Liquor ammonia acetatis*—*s. of Acetate of morphia*, *Liquor morphiæ acetatis*—*s. Alkaline*, see *Fulgokali*—*s. of Alkaline iron*, *Liquor ferri alkalini*—*s. of Alum*, compound, *Liquor aluminis compositus*—*s. of subcarbonate of Ammonia*, *Liquor ammonia subcarbonatis*—*s. of Ammoniated copper*, *Liquor cupri ammoniati*—*s. Arsenical*, *Liquor arsenicalis*—*s. Arsenical*, *De Valangin's*, *Liquor Arsenici chloridi*—*s. of Carbonate of Soda*, *Liquor sodæ carbonatis*—*s. of Charcoal*, concentrated, see *Tinctura catechu*—*s. of Chlorinated soda*, *Liquor sodæ chlorinatæ*—*s. of Chloride of Arsenic*, *Liquor Arsenici chloridi*—*s. of Chloride of Zinc*, see *Burnett's disinfecting liquid*—*s. of Citrate of Ammonia*, *Liquor ammonia citratis*—*s. of Citrate of Magnesia*, see *Magnesia citras*—*s. of Citrate of Potassa*, *Liquor potassæ citratis*—*s. of Continuity*, see *Continuity*—*s. of Corrosive sublimate*, *Liquor hydrargyri oxy muriatis*—*s. De Valangin's*, *Liquor arsenici chloridi*—*s. Donovan's*, see *Arsenic and Mercury*, *Iodide of*—*s. Fowler's*, *Liquor arsenicalis*—*s. Gannal's*, see *Alumina sulphas*—*s. of Gutta percha in Chloroform*, see *Gutta percha*—*s. Heinecke's*, see *Arseniate of Soda*.

SOLUTION, *GOADBY'S*, called after the proposer, is made of bay salt, corrosive sublimate or arsenious acid, and water. It is employed to preserve animal substances.

SOLUTION OF HYDRIOATE OF POTASS, *Liquor potassæ hydriodatis*—*s. of Iodide of iron*, *Liquor ferri iodidi*—*s. of Iodine*, compound, *Liquor iodinii compositus*—*s. of Iron*, nitrated, *Liquor ferri nitratis*—*s. Labarraque's*, *Eau de Javelle*—*s. of Lime*, *Liquor calcis*—*s. Lugol's*, *Liquor iodinii compositus*—*s. of Magnesia*, condensed, *Magnesia*, fluid—*s. of Monro*, *Liquor of Monro*—*s. of Nitrate of iron*, *Liquor ferri nitratis*—*s. of Oxymuriate of mercury*, *Liquor hydrargyri oxy muriatis*—*s. Par les huiles*, *Olea medicinalia*—*s. Pearson's*, see *Arseniate of soda*—*s. Plenk's*, see *Hydrargyri oxydum cinereum*—*s. de Potasse*, *Liquor potassæ*—*s. of Potassa*, *Liquor potassæ*—*s. of Sesquinate of iron*, *Liquor ferri nitratis*—*s. of Soda*, *Liquor sodæ*—*s. of Subacetate of lead*, *Liquor plumbi subacetatis*—*s. of Subacetate of lead*, diluted, *Liquor plumbi subacetatis dilutus*—*s. of Subcarbonate of potass*, *Liquor potassæ subcarbonatis*—*s. of Sulphate of morphia*, *Liquor morphiæ sulphatis*—*s. of Tartarized antimony*, *Vinum antimonii tartarizati*—*s. of Ternitate of sesquioxide of iron*, *Liquor ferri nitratis*—*s. of Van Swieten*, *Liquor hydrargyri oxy muriatis*.

SOLUTIVUS, *Laxative*.

SOLUTUM, *Solution*.

SOLVENS, *Solvent*—*s. Minerale*, *Liquor arsenicalis*.

SOLVENT, *Solvens*, (F.) *Fondant*, from *solvere*, 'to dissolve.' A medicine supposed to possess the property of dissolving or removing obstruc-

tions and extraneous substances. Thus we have solvents of the stone in the bladder, &c. See Dissolventia.

SOLVENT, MINERAL, Liquor arsenicalis.

SOMA, Body, Corpus.

SOMASCETICS, Gymnastics.

SOMASCÉTIQUE, Gymnastics.

SOMATIC, *Somat'icus*; from *σωμα*, *σωματος*, 'the body.' That which concerns the body;—as *somatic* death, or death of the body, in contradistinction to *molecular* death or the death of a part.

SOMATODYM'IA, from *σωμα*, *σωματος*, 'the body,' and *δωω*, 'I enter into.' A double monstrosity in which the trunks are united. Of this, there are several varieties:—*Vertebrody'm'ia*, where the union is by the vertebrae; *Ischiody'm'ia*, by the ossa ischii; *Sternody'm'ia*, by the sternum; *Supraomphalody'm'ia*, (F.) *Sus-omphalodymie*, by the super-umbilical regions; *Supra et infra-omphalody'm'ia*, by the superior and inferior umbilical regions; and *Sterno-omphalody'm'ia*, where the union is by both the sternal and the umbilical regions.—Cruevilhier.

SOMATOL'OGY, *Somatolog'ia*, from *σωμα*, *σωματος*, 'the body,' and *λογος*, 'a discourse.' A treatise on the human body. Anatomy.

SOMATOTOMY, Anatomy.

SOMMEIL, Sleep.

SOMNAMBULATION, Somnambulism.

SOMNAMBULISM, *Somnambula'tio*, *Hypnotat'is*, *Hypnotat'is*, *Hypnotat'is*, *Hypnotat'is*, *Rhembas'mus*, *Lunatic'mus*, *Hypnothergi'a*, *Selenogamia*, *Nyctobate'is*, *Autonyctobatia*, *Selenias'is*, *Selenias'mus*, *Somnambulis'mus*, *Nyctobas'is*, *Noctambula'tio*, *Noctambulis'mus*, *Nyctobadia*, *Nyctobatia*, *Siph'ita*, *Syph'ita*, *Paron'ria am'bulans*, *Noctur'gium*, *Oνειροdy'n'ia acti'va*, *Cenospu'dia*, *Somno-vigil*, *Somnambula'tion*, *Sleep-walking*, from *somnus*, 'sleep,' and *ambulare*, 'to walk.' A condition of the brain which occasions the individual to execute, during sleep, some of those actions that commonly take place when awake. It has been imagined that the somnambulist is guided solely by his subjective sensations; but there are many facts which induce the belief that the external senses are not always entirely suspended. The causes of somnambulism are very obscure; and the principal indication of treatment is,—so to arrange the bed-chamber of the somnambulist, that he cannot get out of it during sleep; and to avoid suppers, as well as all bodily and mental excitement.

SOMNAMBULISM, MAGNET'IC, MESMERIC, or ARTIFIC'IAL, *Somnambulis'mus artificia'lis*, *Cataph'ora magnet'ica*, *Somniph'athy*, is that peculiar condition, into which individuals are thrown by the *Animal Magnetizer*. It occurs only in those of weak nervous temperament; in hysterical females, &c.

SOMNAMBULISMUS, Somnambulism — s. Artificialis, Somnambulism, magnetic.

SOMNAMBULIST, *Somnam'bulus*, *Hyb'nob'ates*, *Hypnob'ates*, *Noctam'bulus*, *Lunatic'us*, *Noctam'bulus*, *Somnam'bulus*. A sleep-walker.

SOMNAMBULIS'TIC. Relating or appertaining to somnambulism, — as the '*somnambulist'ic state*.'

SOMNAMBULO, Somnambulist.

SOMNIA'TIO, from *somnium*, 'a dream.' The act of dreaming.

SOMNIATIO MORBOSA, *Oνειρολα'zus* — s. in *Statu vigili*, Hallucination.

SOMNIF'EROUS, *Som'nifer*, *Soporifer*, *Hypnot'icus*, *Hypnot'icus*, *Soporat'ivus*, *Soporif'ic*, *Soporiferous*, (F.) *Assoupissant*; an epithet for a medicine which causes sleep, from *somnus*, 'sleep,' and *fero*, 'I carry.'

SOMNIL'OQUIST; same etymon as the next. One who talks in his sleep.

SOMNIL'OQUIUM, *Somniloquis'mus*, *Somnil'oquism*, from *somnus*, 'sleep,' and *loquor*, 'I speak.' Talking in the sleep.

SOMNOPATHY, Somnambulism, magnetic.

SOMNIUM, *Eny'p'nion*, *Insom'nium*, a Dream, (Sc.) *Suevin*, *Sueving*, *Suevynung*, *Suenyng*, (F.) *Rêve*, *Songe*. A confused assemblage, or accidental and involuntary combination of ideas and images, which present themselves to the mind during sleep. Dr. S. Mitchell, of New York, used the word *Somnium* to signify the state between sleeping and waking, in which persons perform acts of which they are unconscious.

For 'waking dreams,' see Hallucination.

Those distressing dreams which occur during a state of half sleep, are called, by the French, *Réasseries*.

SOMNO-VIGIL, Somnambulism.

SOMNOLENCY, *Somnolen'tia*, *Hypnæthe'sis*, *Hypno'dia*, *Dormita'tio*, *Carus*, *Letharg'us*, *Cataph'ora*, *Cataphora*, *Coma*, *Somnolen'tum*, *Caro'sis*, *Slumber*, *Heav'iness*, (F.) *Assoupissement*; from *somnus*, 'sleep.' Sleepiness. Also a state intermediate between sleeping and waking. It is often a symptom of disease.

SOMNUS, Sleep.

SOMPNUS, Sleep.

SON, Furfur, Sound.

SONCHUS, *σῶχος*, *S. Olera'ceus*, *Hiera'cium olera'ceum*, *Oicer'bita*, *Sow Thistle*, *Swine-thistle*, (F.) *Laiteron doux*. Ord. *Cichoraceæ*. Most of the species of *Sonchus* abound with a milky juice, which is very bitter, and said to possess diuretic virtues.

SONCHUS FLORIDANUS, *Mulgedium Florida-num*.

SONDE, Sound, Specillum — s. *Cannelée*, Director.

SONDE D'ANEL. A very fine stylet of silver, awl-shaped at one end. It is used for probing the lachrymal passages through the punctum lacrymale.

SONDE DE BELLOC. An instrument invented by Belloc for plugging the nasal fossæ, in cases of hemorrhage. It consists of a curved silver canula, open at both ends, and furnished with a button. The instrument is passed through the nares to the throat; and a plug is attached to the button, so that, when the instrument is drawn forwards through the nose, the posterior nostril is stopped.

SONDE BRISÉE (F.). *Eyed probe*. A long, straight stylet, composed of two parts, which may be screwed to each other; blunt at one end, and furnished at the other with an eye, so that it may be used either as a probe for punctured wounds, or for a seton needle.

SONDE À CONDUCTEUR (F.). A modification of the ordinary catheter;—to facilitate the changing of catheters, without the fear of making false passages. It is a common catheter, open at its beak. The *Stylet* or *Mandrin* is one half longer than that of the ordinary catheter. To use the instrument, the stylet is pushed into the cavity of the bladder through the catheter. The latter is then withdrawn by slipping it over the stylet, which is kept in its place, and serves as a guide to a fresh catheter.

SONDE À DARD, see Lithotomy (high operation).

SONDE DE LAFOREST. A small, crooked canula, used for probing the nasal duct from below upwards, and for throwing injections into it.

SONDE ou PINCE DE HUNTER. A cylindrical silver canula; of the size of a common catheter; open at both extremities, and contain-

ing a wire of iron, one of the ends of which is split into two parts; each being terminated by a small silver scoop, and separating from each other by virtue of their elasticity, when pushed out of the canula, and again approximating when drawn in. It has been used for laying hold of small calculi in the urethra.

SONDE DE VARECQ. Soda.

SONGE, Somnium.

SONITUS AURIUM, Tinnitus aurium — s. Fluidi, Hygrecema — s. Hepaticus, Hepatechema — s. Intestinalis, Enterechema — s. Stomachicus, Gasterechema.

SONOROUS RHONCHUS, Râle sonore.

SONUS, Sound — s. Vocis, Accent.

SOOJA, see Dolichos.

SOOJIE, BASTER'S, a farinaceous article of diet, is said to consist of wheat flour sweetened with sugar.

SOOL, Gastroperiodynia.

SOOT, Fuligo — s. Tea, see Fuligokali — s. Wart, Cancer, chimney-sweeper's.

SOPHIA, Sisymbrium sophia — s. Chirurgorum, Sisymbrium sophia.

SOPHISTICATION, Falsification.

SOPHO'RA HEPTAPHYLLA, Anticholerica. Ord. Leguminosæ. Sex. Syst. Decandria Monogynia. The root and seeds of this shrub are intensely bitter, and are said to have been found useful in cholera, colic, dysentery, &c.

SOPHORA TINCTO'RIA, Baptisia seu Podalyria Tinctoria, Indigofera, Wild Indigo, Indigo Weed, Broom, Indigo Broom, Horseshy Weed, Yellow Broom, Clover Broom, Rattle Bush, Yellow Indigo. An American plant, the taste of whose root is unpleasant, sub-acrid, and nauseous—very similar to that of ipecacuanha. In a large dose, the bark of the root acts both as an emetic and cathartic. It has been considered antiseptic and febrifuge, and has been used, in the form of fomentation or cataplasm, in phagedenic and gangrenous ulcers. A liniment, prepared by simmering the cortical part of the root in cream, has been found an efficacious application to sore nipples and to ulcerated breasts.

Baptisia Leucantha, Tall white false Indigo, an indigenous plant, has the same properties.

SOPHRONISTE DENTES, see Dentition.

SOPHRONISTERES DENTES, see Dentition.

SOPHROSYNE, Temperance.

SOPIENS, Paregoric.

SOPIO, Opium.

SOPOR. A profound sleep, from which the person can be roused with difficulty. It is a symptom in many of the neuroses.

SOPOR CAROTICUS, Carus.

SOPORARIÆ ARTERIÆ, Carotids.

SOPORATIVUS, Somniferous.

SOPORIFEROUS, Somniferous.

SOPORIFIC, Somniferous.

SO'POROUS, So'porus, (F.) Soporeux, from *sopor*, 'sleep.' Sleepy: causing sleep.

SORA, Essera.

SORB TREE, Sorbus domestica.

SORBASTRELLA, Pimpinella saxifraga.

SORBEFA'CIENT, Sorbefaciens, from *sorbere*, 'to suck in,' and *facere*, 'to make.' A remedy that promotes absorption.

The following are the chief sorbefacients or modifiers of nutritive action — *eutrophics*: — Acida Vegetabilia; Alkalia; Ammoniacum; Brominium; Galbanum; Hydrargyri Præparata; Iodinium; Liquor Calci chloridi; Spongia usta; Compression, (methodical;) Friction; Mental Sorbefacients, (Imagination, Emotions.)

SORBETHUM, Scherbet.

SORBETUM, Scherbet.

SORBETUM, Scherbet.

SORBIER DES OISELEURS, Sorbus acuparia.

SORBITIO, Jus.

SORBITO CARNIS seu Ex CARNE. Broth or soup made of meat.

SORBITUM, Jus.

SORBUM, see Sorbus domestica.

SORBUS, Cratægus aria, S. domestica.

SORBUS ACUPA'RIA seu AUCUPA'RIA, Mes'pilus, M. aucuparia, Mountain Service, Mountain Ash, Quicken tree, Roan tree, Witchen, Witch hazel, Witch wood, (Sc.) Roden tree, Roun tree, Rowan tree, Rone, (Prov.) Keer, Rantree, (F.) Sorbier des Oiseleurs, Frêne sauvage. Ord. Rosacæ. The berries — (Sc.) Rodens — of this tree are employed for similar purposes as the last.

SORBUS ALPINA, Cratægus aria.

SORBUS AMERICA'NA, Pyrus Americana, American Mountain Ash. The bark resembles that of the Wild Cherry, and might be used in the same cases.

SORBUS ARIA, Cratægus aria — s. Acuparia, S. acuparia — s. Cydonia, Pyrus cydonia.

SORBUS DOMES'TICA, Sorbus, Pyrus domestica, Mes'pilus domestica, Oie, Service or Sorb Tree, (F.) Cormier. Family, Rosacæ. Sex. Syst. Icosandria Trigynia. The fruit of this tree, Sorbum, is astringent, and has been recommended in diarrhoea and dysentery. It is given in the form of rob. A kind of cider is made from it, and also brandy.

SORBUS MALUS, Pyrus malus.

SORDES, Rhypos, 'Filth.' A dirty-looking sanies, discharged from ulcers. Also, accumulations of the secretions of the mouth, on the teeth in adynamic fevers more especially, and of foul matters in the stomach — *Sordes prima'rum viarum*, Rhyparia. The French call an ulcer, exhibiting such an appearance, *sordid*. Also dirt of any kind.

SORDES, Ichor, Saburra — s. Acidæ, Acidities — s. Aurium, Cerumen — s. Primarum viarum. Embarras gastrique, see Sordes — s. Ventris, Excrement.

SORDICULÆ AURIUM, Cerumen.

SORE, Ulcer.

SORE, BAY. A disease which Mosely considers to be a true cancer. It is endemic in the Bay of Honduras.

SORE MOUTH, Stomatitis — s. Gangrenous, Cancer aquaticus — s. Throat, Cynanche — s. Throat, Clergyman's, Pharyngitis, follicular — s. Throat, inflammatory, Cynanche tonsillaris — s. Throat, putrid, Cynanche maligna — s. Throat, tubercular, Pharyngitis, follicular — s. Throat, ulcerous, Cynanche maligna.

SORENESS, Sax. *ronp*, 'vexed, molested.' *Parap'sis aeris teneritudo*. Painful uneasiness or tenderness, local or general, on being touched with a pressure that does not usually excite distressing feeling. Often, a febrile symptom. Also, the tenderness of a wound, ulcer, &c.

SORGHUM, Panicum Italicum.

SOROCCO, Puna.

SOROCCHE, Puna.

SORORIA'TIO, from *sorores*, 'sisters,' *sororio*, 'to grow up or swell together.' The period at which the breasts of the female become developed. The act of becoming thus developed. A young maiden, whose mammae begin to show, — *sororiantes mammae* — was formerly called *sororians virgo*. — Plantus.

SORREL, COMMON, Rumex acetosus — s. Field, Rumex acetosella — s. French, Rumex scutatus — s. Garden, Rumex scutatus — s. Mountain, Oxalis acetosella, Oxyria reniformis — s. Salt of, Potass, oxalate of — s. Roman, Rumex scutatus — s. Sheep, Rumex acetosella.

SORREL TREE, *Andromeda arborea*, *A. mariana*.
 SORREL, WELSH, *Oxyria reniformis*—*s.* White,
Oxalis acetosella—*s.* Wood, Yellow, *Oxalis stricta*.
 SOSTRUM, *Sotrum*; from *σωζω*, 'to save,'
σωτηρ, 'a preserver.' A reward given to one who
 saves the life of another. A physician's fee or
honorarium, (F.) *Honoraire*.

SOTERIA DOCTRINA, *Medicina*.

SOTERIE AQUÆ, Waters, mineral.

SOTIREL'LA. Ancient name of a medicine
 composed of opium and other narcotics, *nutmeg*,
saffron, *camphor* and *soot*. It was used in cer-
 tain diseases of the teeth.

SOTRUM, *Sotrum*.

SOUBRESAUT, *Subsultus tendinum*.

SOUCHERLOON, *Bit noben*.

SOUCHET DES INDES, *Curcuma longa*—
s. Odorant, *Cyperus longus*—*s.* Rond, *Cyperus*
rotundus.

SOUCI, *Calendula officinalis*, *Panophobia*—
s. des Alpes, *Arnica montana*—*s.* des Champs,
Calendula arvensis—*s.* Ordinaire, *Calendula* offi-
 cinalis.

SOUDE, Soda—*s.* Acétate de, Soda, acetate
 of—*s.* Aéré, Sodæ carbonas—*s.* Borate de, Bo-
 rax—*s.* Borate saturé de, Borax—*s.* Carbonate
 de, Sodæ carbonas—*s.* Caustique, Soda caustica
—s. à la Chaux, Soda caustica—*s.* Chlorure de,
 Soda, chloride of—*s.* du Commerce, Soda—*s.*
Crayeuse, Sodæ carbonas—*s.* Effervescente, Sodæ
 carbonas—*s.* Hyposulfite de, Soda, hyposulphite
 of—*s.* Phosphate de, Soda, phosphate of—*s.* et de
 Potasse, Tartrate de, Soda, tartrate of—*s.* Pure,
 Soda caustica—*s.* Sousborate de, Borax—*s.* Sous-
 carbonate de, Soda, subcarbonate of—*s.* Sulfate
 de, Soda, sulphate of.

SOUFFLE, see Murmur, respiratory—*s.* Am-
 phorique, see Cavernous respiration—*s.* Bron-
 chique, see Murmur, respiratory—*s.* Métallique,
 see Cavernous respiration—*s.* Placentaire, *Bruit*
placentaire—*s.* Trachéal, see Murmur, respiratory
—s. Tubaire, see Murmur, respiratory—*s.* Utérin,
Bruit placentaire—*s.* Voilé, see Cavernous respi-
 ration.

SOUFFLE, WHISPERING. A bellows' sound heard
 over solidified lung and excavations when the
 patient whispers.—*Flint*.

SOUFRE, Sulphur—*s.* Carburé, Carbonis sul-
 phuretum—*s.* Iodure de, Sulphuris iodidum—*s.*
Sublimé, Sulphur sublimatum—*s.* Végétal, Lycopo-
 dium.

SOUILLEURE, Pollution.

SOULA'MEA AMA'RA, *Rex amaro'rum*. A
 shrub of the Moluccas; *Order*, Polygalacæ; all
 the parts of which are of great bitterness—"hor-
 renda amarities." It is employed by the Malays,
 and, it is said, with success in cholera and pleu-
 risy; and as a febrifuge.

SOUND, *Specill'um*, *Stylus*, *Explorato'rium*,
Rad'olus, (F.) *Sonde*. An instrument used by
 surgeons to discover whether there be a stone in
 the bladder. It is usually made of highly pol-
 ished steel, and is shaped like the catheter. The
 operation is termed sounding.

The French *Sonde* has, however, a more ex-
 tensive signification. It means different instru-
 ments introduced into cavities of certain organs,
 or into wounds, fistulas, &c., to investigate their
 condition, or to fulfil some therapeutical indica-
 tion. See *Sonde*.

SOUND, *Sonus*, *Echos*, *Noise*, (F.) *Son*, *Bruit*.
 The sensation produced on the auditory nerve by
 the vibrations of a sonorous body. Sounds may
 be propagated in three modes. 1. By *reciproca-*
tion or *con'sonance*, as when a sounding body, of
 a definite pitch, produces a musical tone when
 another body of the same pitch is sounded near
 it. 2. By *res'onance*, as when a sounding body
 is placed in connection with another, one or more

of whose parts may be thrown into reciprocal vi-
 bration; and 3. By *conduc'tion*, as where the vi-
 brations are transmitted through fluid, liquid, or
 solid media.

SOUND, Catheterize, Syncope—*s.* Auricular,
Apyromele—*s.* Bellows, friction, rasp, saw, lancet,
 &c., see *Bruit*—*s.* Cracked pot, *Bruit de pot fêlé*
—s. Crumpling, pulmonary, *Froissement pulmo-*
naire—*s.* Laryngeal, *Laryngeche*.

SOUND, U'TERINE, *U'terine bougie*, (F.) *Sonde*
utérine. An instrument, somewhat similar to a
 small male catheter, to be introduced into the
 cavity of the uterus, in the diagnosis of diseases
 of that viscus. When notched on the stem, it
 enables the size of the uterine cavity to be ap-
 preciated, and becomes a good *hysterom'eter*.

SOUNDING, Searching, see Sound.

SOUNDNESS OF MIND, Sanity.

SOUP'IR, Sigh.

SOUR BERRY, see *Oxycantha Galeni*, *Vaccini-*
um oxycoccos—*s.* Dock, boreal, *Oxyria reni-*
formis—*s.* Leaf, *Andromeda arborea*—*s.* Tree
Andromeda arborea—*s.* Wood, *Andromeda* ar-
 borea.

SOURCIL, Supercilium.

SOURCILIER, Corrugator supercilii, Super-
 ciliary.

SOURD, Deaf.

SOURDS-MUETS, see *Mutitas surdorum*.

SOUR-MILK, Buttermilk.

SOUR-SOP, *Bromelia ananas*.

SOURWOOD, *Andromeda mariana*.

SOUS-ACROMIO-CLAVI-HUMÉRAL, Del-
 toïd—*s.* Atloïdien, Subatloïdeus—*s.* Azoidien,
 Subaxoïdeus—*s.* Azotate de Bismuth, Bismuth,
 subnitrate of—*s.* Clavier, Subclavius—*s.* Costaux,
 Intercostal—*s.* Cutané, Subcutaneous—*s.* Épi-
 neux, Infra-spinatus—*s.* Lingual, Sub-lingual—*s.*
Mammaire, Submammary—*s.* Maxillaire, Sub-
 maxillary—*s.* Maxillo-Lobal, Depressor anguli
 oris—*s.* Mental, Submental—*s.* Métacarpo-latéri-
 phalangien, Prior annularis—*s.* Optico-sphéno-
 scléroticé, Rectus inferior oculi—*s.* Orbitaire,
 Suborbital—*s.* Pubio-coccygien, Levator ani—*s.*
Pubio-crêti-tibial, Gracilis—*s.* Pubio-prétibial,
 Gracilis—*s.* Pubio-trochantérien-externe, Obtura-
 tor externus—*s.* Pubio-trochantérien-interne, Ob-
 turator internus—*s.* Scapulaire, Subscapularis—*s.*
Scapulo-trochantérien, Subscapularis—*s.* Sels, see
 Salt.

SOUTH CAROLINA, MINERAL WATERS
 OF. *Glenn's Springs*, in Spartanburg District,
 are highly charged with salts of lime, and, not
 far from them, is a chalybeate. *West's Spring*,
 and *Chick's Springs*, a few miles from Greenville,
 resemble Glenn's Springs.

SOUTHERNWOOD, *Artemisia abrotanum*—
s. Field, *Artemisia campestris*—*s.* Maritime, *Ar-*
temisia maritima—*s.* Tartarian, *Artemisia santon-*
ica.

SOWBREAD, *Arthanita*, *Cyclamen*.

SOWENS, Flummery.

SOY, see *Dolichos soja*.

SOYMIDA FEBRIFUGA, *Swietenia febrifuga*
 SPA, MINERAL WATERS OF. A town
 seven leagues southeast of Liège, where there
 are several springs, possessing a sharp acidulous
 taste; containing much carbonic acid, carbonates
 of iron, lime, and magnesia; carbonate of soda,
 and chloride of sodium. The water is much used
 as a tonic.

SPA WATER, ARTIFICIAL. Sodæ subcarb. gr.
 vij; magnesia carbon. 5j; limat. ferri. gr. iij;
 sollii chlorid. gr. j; aqua. Oijj. Impregnate
 with gas from marble powder and sulph. acid.,
 aa 3x.

SPACE, INTERPEDUNCULAR, see Inter-
 peduncular—*s.* Perforated, anterior, *Locus per-*

oratus anticus —s. Perforated, posterior, see Tarsiui pons.

SPADE-BONE, Scapula.

SPADO, Castratus, Eunuch, Spasm.

SPAGYRIA, Chymistry.

SPAG'YRISTS, from *σπαω*, 'I separate,' and *γυρειν*, 'I assemble,' because they reduced compounds into elements, and formed the latter into compounds [?]. A sect of physicians, who pretended to account for the changes that occur in the human body in health and disease, in the same manner as the chymists of their day explained those of the inorganic kingdom. *Medici'na Paracelsi'stica* seu *Spagy'rica*, was likewise called *Hermet'ica*, because it was believed that the means of cure adopted in it had been found in the books of Hermes.

SPAIG, Skeleton.

SPANÆMIA, from *σπας*, 'poor,' and *αἷμα*, 'blood.' Poverty of the blood. Diminution in the quantity of fibrin and red corpuscles of the blood, — as where bleeding has been carried beyond certain limits. — Simon.

SPANÆMIC, *Spanæmicum* (remedium), *Dysplas'ticum*, *Plastily'ticum* et *erethily'ticum*, *Hæmatoly'ticum*, *Dyscrasi'acum*. Same etymon as the last. Relating to *spanæmia*.

A medicine whose protracted use is said to impoverish the blood, — as iodine, bromine, &c.

SPANDARAPUM, Sparadrapum.

SPANING, Weaning.

SPANISH FLY, *Cantharis* —s. Juice, Extractum glycyrrhizæ.

SPANNADRAPUM, Sparadrapum.

SPANOPO'GON, from *σπας*, 'rare,' and *πογόν*, 'beard.' One who has lost his beard, or has a thin beard.

The Greeks called those who had little beard, or who had lost it, *Spanopogo'nes*.

SPARADRAPA, Sparadrapum.

SPARADRAPIER. The name of a machine for spreading sparadraps. A *Plaster machine*. It consists of a table, with two raised pieces, movable, and furnished with points, by which the cloth can be stretched, and of a lamina or blade of metal to extend the plaster over it.

SPARADRA'PUM, *Sparadra'pa*, *Sparadrapus*, *Tela emplas'tica* seu *emplas'trica*, *Spandara'pum*, *Spannadra'pum*, (F.) *Sparadrap*. Any adhesive plaster spread upon linen or paper. The chief sparadraps are the following:—

SPARADRAPUM ADHÆSI'VUM, *Adhæsive Plaster*. A spread plaster of the *Emplastrum adhæsivum*. It is also called *Strapping*.

English Court Plaster is a Sparadrap, (F.) *Taffetas agglutinatif*, *T. gommé*, *Taffetas d'Angleterre*, *Ser'icum Ang'licum*, *Emplastrum Ang'licum*, *E. Anglica'num*, *E. Adhæsivum Woodstock'i*, *E. Ichthyocol'læ telæ induc'tum*, *E. Glutin'o'sum*, *Tela Ichthyocol'læ glutinans*, *Isinglass Plaster*. It is made by stretching black silk, and brushing it over with a solution of isinglass \mathfrak{z} j, in proof spirit \mathfrak{z} xij, to which tinct. benzoin. \mathfrak{z} j, are added. When dry, this is repeated five times; after this, two coats are given it of a solution of tereb. *Chia* \mathfrak{z} iv, in tinct. benz. \mathfrak{z} vj, which render it less liable to crack.

Liston's Isinglass Plaster is made by spreading several coats of strong solution of isinglass in weak spirit over oiled silk, or, still better, over animal membrane previously prepared for the purpose from the peritoneal coat of the cæcum of the ox.

SPARADRAPUM PRO FONTIC'ULIS. *Issue Plaster*, is sometimes made of simple *diachylon*, \mathfrak{b} j; *Burgundy pitch* and *sarcocolla*, each \mathfrak{z} iv; common turpentine, \mathfrak{z} j. Spread upon linen and polished with a moistened calendering glass rubber.

SPARADRAPUM seu **TELA GALTERI**, *Defensive*

Plaster, (F.) *Toile de Gantier*. This is made of olive oil, *lbss*; suet, \mathfrak{z} iv; wax, \mathfrak{z} x; litharge, common turpentine, *thuss*, and mastic, aa. \mathfrak{z} ij; bole armeniac, flour, aa. \mathfrak{z} j. Pour it, while liquid, upon cloth, and spread. Used for issues and to keep on dressings.

SPARADRA'PUM VESICATO'RIVM. Several substances have been introduced as substitutes for blistering plaster, (see *Emplastrum Lyttæ*, under the names, *Tela vesicato'ria*, *Blistering Tissue*, &c., and other forms of which are the *Taffetas vésicant* ou *épispastique*, *Papier épispastique*, *Charta vesicato'ria*. They are made of an ethereal or alcoholic extract of cantharides, or of cantharidin, mixed with wax, and spread, in a very fine layer, on silk or paper previously oiled or waxed. They are efficient agents.

SPARADRAPUM VIRIDÈ, *Corn Plaster*. This may be made of yellow wax, \mathfrak{b} j; Burgundy pitch, \mathfrak{z} xij; common turpentine, \mathfrak{z} vj; verdigris, \mathfrak{z} ij; spread on cloth and polished.

Kennedy's Corn Plaster is made of yellow wax, \mathfrak{b} j; Venice turpentine, \mathfrak{z} j; verdigris, \mathfrak{z} j.

SPARADRAPUS, Sparadrapum.

SPARAGMA, Laceration.

SPARAGMOS, Spasm.

SPARAGUS, Asparagus.

SPARAL'LIVM, *Clyster uteri'nus*. An injection into the vagina. — Ruland and Johnson.

SPAREIDIA. A ligature covered with the white of egg. — Paracelsus.

SPARGANIUM RAMO'SUM, *Great Bur-reed*, from *Sparganum*, owing to the ribbon-like leaves. Indigenous; Order, Typhacææ; flowering in July and August. The roots are subastrigent, but esculent; yielding a fine fecula, similar to salep. They are sometimes made into a poultice for inflamed mammæ.

SPARG'ANON, *Sparg'anum*, from *σπάργω*, 'I swathe,' 'I wrap.' 'Swathing clothes,' a kind of bandage, with which children were formerly surrounded. — *Foësius*. Also, a fascia.

SPARGANO'SIS, *Spargo'sis*, *Intumescen'tia lactea mamma'rum Mastodyn'ia polyg'ala*, from *σπάργω*, 'I am ready to burst.' Extreme distension of the breasts by milk. Sparganosis also means the wrapping of a child in swathing clothes.

SPARGANOSIS PUEPERARUM, *Phlegmatia dolens*.

SPARGANUM, Sparganon.

SPARROWGRASS, Asparagus.

SPARSUS, Sporadic.

SPARTIVM JUN'CEUM, *Spanish broom*. Ord. Leguminosæ. A small European shrub, cultivated in the gardens of the United States, on account of its yellow flowers. The seeds are diuretic and tonic, in small doses; emetic and cathartic in large. They have been used in dropsy — 10 or 15 grains three times a day.

SPARTIVM SCOPA'RIVM, *Cyl'tisus scopari'us*, *Genis'ta*, *G. scopari'ia* seu *hirsu'ta*, *Cytisogenista*, *Broom*, (F.) *Genêt à balai*. *Broom Tops*, *Spartii Cacumina*, *Scopa'rius*, (Ph. U. S.), have a bitter taste, and are possessed of diuretic properties. They have been used in dropsies. Dose, of the powder, \mathfrak{b} j to \mathfrak{z} j. All the genistæ have similar virtues. A decoction of genista, along with the cauterization of the pustules under the tongue, has been recommended in hydrophobia.

SPARTIVM TINCTORIVM, *Genista tinctoria*.

SPASM, *Spasmus*, *Sparag'mos*, *Synol'cè*, *Spado*, from *σπασω*, 'I draw.' The Greeks gave this name to all kinds of convulsions. It is now usually applied to involuntary muscular contractions; and these, again, have been divided into tonic spasm, *Paraton'ia*, which consists in permanent rigidity and immobility of the muscles that are the seat of it (see *Tetanus*); and clonic spasm,

which consists in alternate contractions and relaxations (see Convulsion).

Cullen has a class, *Spasmi*—the *neurospasmi* of Fuchs.

SPASM, CLONIC, Convulsion—s. Cynic, see Canine laugh.

SPASM OF THE GLOTTIS, Asthma thymicum.

SPASM, HISTRIONIC, OF THE FACE, see Tic.

SPASM OF THE LARYNX, Asthma thymicum.

SPASM, MASTICATORY, OF THE FACE, Trismus.

SPASM WITH RIGIDITY, Tetanus.

SPASM, WRITERS'. Involuntary spasms of the muscles of the thumb and index finger, observed, at times, in those who are much accustomed to writing. See Cramp, writers'.

SPASMATICUS, Spasmodicus.

SPASMATODES, Spasmodicus.

SPASME DE LA GLOTTE ET DU THORAX, Asthma thymicum—s. de la Vessie, Cystosasmus.

SPASMI, see Spasm.

SPASMODIC, Convulsive.

SPASMODIC, Spasmodicus.

SPASMOL'OGY, *Spasmolog'ia*, from *σπασμος*, 'spasm,' and *λογος*, 'a treatise.' A treatise on spasms.

SPASMOLYGMUS, Singultus.

SPASMOTICUS, *Spasmat'icus*, *Spasmat'odes*, *Spas'ticus*, *Spastic*, *Spasmod'ic*, (F.) *Spasmodique*, *Spastique*, from *σπασμος*, 'spasm.' Anything relating to spasm. Also, an antispasmodic.

Morbi constrict'o'rii, *M. spasmo'ici*, &c., are diseases accompanied with spasm.

SPASMUS, Convulsion, Spasm—s. Aurium, Otalgia—s. Bronchialis, Asthma—s. Caninus, see Canine laugh—s. Clonicus, Convulsion—s. Cynicus, see Canine laugh—s. Facialis, Tic—s. Glottidis, Asthma thymicum—s. Linguae, Glossospasmus—s. Maxillae inferioris, Trismus—s. Muscularis, Cramp—s. Musculorum Faciei, Canine laugh—s. Nutans, Convulsion, salaam—s. Periphericus, Horrida cutis—s. Universalis, Syncynus—s. Ventriculi, see Cardialgia—s. Vesicae, Cystosasmus.

SPAS'NIA, same etymon as Spasm. [?] A term used by Mercurialis to designate the lancinating pain produced, at times, in the chest by violent fits of coughing.

SPASTIC, Spasmodicus, Tonic spasm.

SPASTICUM, Tetanic.

SPATHA, Hypaleiptron.

SPATHES'TER, from *σπασω*, 'I draw.' A surgical instrument, used for drawing the prepuce over the glans, when too short.—P. Amman.

SPATHOMELE, Spatula.

SPATHULA, Scapula, Spatula—s. Foetida, Iris foetidissima.

SPATHYEMA FETIDA, Dracontium foetidum.

SPATIA INTERLOBULARES, Interlobular fissures.

SPAT'ILĒ, *σπατ'ιλη*, 'human excrement.' A liquid fecal evacuation. Excrement.—Hippocrates.

SPATIUM ORIS, Mouth—s. Trigonum, Lyra.

SPATTERDOCK, Nuphar advena.

SPAT'ULA, *Spath'ula*, diminutive of *Spatha*, *σπαθη*, 'a broad-sword.' *Spathome'le*, *Specill'um latum*. An instrument used for spreading plasters, stirring ointments, holding down the tongue, &c. Also, the scapula.

SPATULA PRO ORE, Glossospatha.

SPAUD, Humerus.

SPAUL, Saliva.

SPAUT-BONE, Humerus.

SPAW-BONE, Humerus.

SPAYING, see Castration.

SPEANING-BRASH, Brash, weaning.

SPEARWORT, Ranunculus flammula.

SPEAUTER, Zinc.

SPECIAL, see Specific.

SPECIALIST, (F.) *Spécialiste*. One who devotes himself to a specialty; as to diseases of particular parts, as of the eye, ear, chest, &c.

SPECIES, (F.) *Espèces*, *Spécialités*, *Thés composés*. Official mixtures of substances cut or bruised. Sometimes used for compound powders, as it always appears to have been formerly. See Pulvis.

SPECIES AROMATICÆ, Pulvis cinnamomi compositus—s. Diacinnamomi, Pulvis cinnamomi compositus—s. Diacretæ, Pulvis cretæ compos.—s. Diagalapæ, Pulvis julapæ compositus—s. Diambræ sine odoratis, Pulvis cinnamomi compositus—s. Diatrageanthæ frigidæ, Pulvis tragacanthæ compositus—s. Hieræ picræ, Pulvis aloes cum canellâ—s. Lætificantes Rhazis, Pulvis cinnamomi compositus—s. e Scordio cum opio, Pulvis cretæ compositus cum opio—s. e Scordio sine opio, Pulvis cretæ compositus.

SPECIF'IC, *Specif'icus*, *Nosocrat'ic*, (F.) *Spécifique*, from *species*, 'a form or fashion,' and *facere*, 'to make.' A substance to which is attributed the special property of removing some particular disease. Probably no such remedy exists. *Mercury* in syphilis, and *sulphur* in the itch, have been regarded as the strongest examples. The word is also used adjectively in the sense of *special*; as "mercury has a *specific* or *special* action on the salivary glands."

SPECIFIC OF HERSEN'SCHWAND. A once celebrated German vermifuge. It is said to have consisted of 10 grains of *Gamboge*, with 20 of *Carbonate of Potassa*. It is affirmed, that mercury and arsenic have also been found in it.—Paris.

SPECIFICUM PARACELSI, Potassæ sulphas.

SPECIL'LUM, *Melè*, *Stylus*, *Explorato'rium*, from *specio*, 'I examine,' *Specia'rium*, a probe, (F.) *Styilet*, *Sonde*. A surgical instrument, employed for examining wounds, fistulas, and for passing setons, &c. It is usually formed of silver; and is terminated, at one end, by an olive-shaped button. The *eyed probe* has an aperture at the other extremity.

SPECILLUM, Evericulum, Hypaleiptron, Sound—s. Auricularium, Apyromele—s. Cereumi, *Bougie*—s. Excavatam, Stylus excavatus—s. Latum, Spatula—s. Minus, Melotis.

SPÉCIOLÉ, Species.

SPECKS OF THE CORNEA, see Caligo.

SPECLARION, Specillum.

SPEC'TACLES, from *spectare*, 'to behold;' *Conspicill'la*, (Sc.) *Breels*, (F.) *Besicles*, *Lunettes ordinaires*, *Conservees*. Glasses to assist the sight; arranged so as to be adapted to both eyes. These glasses are more or less *concave* or *convex*, according as the sight is more or less short, (*myopic*), or long, (*presbyopic*.) When the glass is adapted to one eye, it is called an *Eyeglass*, *Conspicill'um*, *Perspici'llum*, *Vitrum ocula're*.

SPECULARIS LAPIS. A transparent mineral, but of what nature is not clear, which was formerly employed in epilepsy. In old times it was used for glass.—Pliny.

SPECULUM, *Catop'ter*, *Catop'tron*, *Diop'tron*. In Latin, 'a mirror;' from *specio*, 'I see.' In surgery, it means different instruments for dilating cavities, and facilitating their examination. See Dilator. There are various instruments of this kind,—the *S. Ani*, *S. Auris*, *S. Vaginae*, (*Colpeurynt'er*, *Elytreurynt'er*), *S. Matricis*, *S. Oculi*, *S. Oris* ((F.) *Bâillon*), *S. Gutturis*, *S. Vesicae*.

SPECULUM CITRINUM, Orpiment—s. Indicum, Ferri limatura—s. Lucidum, Septum lucidum—s.

Oris, Glossocatochus—s. Veneris, Achillea millefolium.

SPECUS, Vulva—s. Cordis, Ventricle of the Heart—s. Pro medullâ spinali, see Vertebral column—s. Vertebralis, see Vertebral column.

SPEDALSKE, see Radzyge.

SPEDALSKHED, see Radzyge.

SPEECH, Voice, articulated.

SPEECHLESSNESS, Mutitas.

SPEEDIMAN'S PILLS, see Pilulæ aloes et myrrhæ.

SPEEDWELL, BROOKLIME, Veronica beccabunga—s. Female, Antirrhinum elatine—s. Official, Veronica—s. Purslain, Veronica peregrina—s. Water, Veronica beccabunga.

SPELL-BONE, Fibula.

SPELTRUM, Zincum.

SPERAGE, Asparagus.

SPERAGUS, Asparagus.

SPERM, σπέρμα, σπέρματος, from σπέρω, 'I sow.'

Spermatic fluid or liquor, Seminal fluid, Seed, Semen, S. virile seu musculinum seu genitalis, Seminum, Genitalis, Medullæ, Ros, Sanguis, Serum, Humor genitalis seu seminalis seu venereus, Uri'na genitalis, Genit'ra, Sperma, S. virile, Spermium, Tho'rë, Tho'rus, Lac maris, Lagnæ'a, Lagni'a, Lagnæ'ua, Germen, Vita'le virus, Male's milk, Propagatory or genital liquor, Vital or quickening venom, (F.) Semence, Fluide séminal. A whitish, viscid fluid, of a peculiar smell, secreted by the testicles, whence it is carried by the vasa deferentia to the vesiculæ seminales, to be thrown into the vagina, during coition, through the ejaculatory ducts and the urethra. It is the fecundating fluid, and must come into actual contact with the germ of the female. The *Aura sem'inis, Gonau'ra*, is incapable of effecting fecundation. The semen, at the time of emission, is composed of two different fluids; the one liquid and transparent, which is considered to be the secretion of the prostate,—the other, white, and as thick as mucilage; the product of the testicular secretion. The sperm contains, according to Vauquelin, 900 parts of water, 50 of animal mucilage, 10 of soda, and 30 of calcareous phosphate. The animal matter is peculiar, and by some termed *spermatin*.

Microscopic observations show that it contains *spermatozo'a*, or more properly *spermatozo'idæ*; for their animalcular nature is not demonstrated. They are produced in cells—*sperm-cells*—and have probably no more title to be considered animalcular, than the cilia of the ciliated epithelium. By careful examination, other minute, round, granulated bodies may almost always be detected, which are in all cases much less numerous than the spermatozoa. These bodies are the *seminal granules, gran'ula sem'inis*. Pure sperm, in its most perfect state, consists principally of spermatozoids and seminal granules: both of which are enveloped in a small quantity of fluid, *liquor sem'inis, spermat'ic liquor*.

It has been imagined, but erroneously, that during coition there is a secretion of female sperm—*Semen mulie'brë, Thelygonum*. The increased secretion that takes place is chiefly from the glands of Duverney.

Sperm also means spermæti.

Sperm CELL, see Sperm.

SERMA, Sperm—s. Mercurii, Hydrargyris acetatus.

SERMA RANA'RUM, Sperm'i'ola seu Sperm'iolon seu Sperm'i'ola seu Sperm'olum rana'rum. Frog's spawn. Once used in medicine.

SERMACEII, Cetaceum—s. Whale, see Cetaceum.

SERMACRASIA, Spermatorrhœa.

SERMATACRASIA, Spermatorrhœa.

SERMATACRATIA, Spermatorrhœa.

SPERMAT'IC, *Spermat'icus, Semina'lis, Sem'inal, (F.) Spermatique*, from σπέρμα, σπέρματος. That which relates to the sperm. A name given to different parts connected with the organs of generation.

SPERMATIC ARTERIES, *Arteriæ Spermat'icæ seu præparan'tes, (F.) Artères testiculaires, A. de l'ovaire* (Ch.), are two in number—one on each side—and arise from the sides of the aorta, sometimes from the renal arteries. They descend, almost vertically, at the sides of the vertebral column, and are distributed differently in the two sexes. In man, the spermatic artery, situate at the side of the vas deferens, issues by the abdominal ring; gives numerous ramifications to the spermatic cord, and divides into fasciculi of branches, one of which goes to the testicle,—the other to the epididymis. In the female, the spermatic artery, *ova'rian artery*, dips into the pelvis, and passes to the ovarium, Fallopian tube, and round ligament.

SPERMATIC CORD, or CHORD, *Testic'ular Cord, Funic'ulus Spermaticus, Corpus varicosum, (F.) Cordon spermatique ou testiculaire*. The vascular and nervous chord, by which the testicle is suspended. It is composed of the spermatic artery and veins; of other small vessels, of lymphatics; of nervous filaments from the spermatic plexus, and from the genito-crural branch of the lumbo-abdominal plexus; of the vas deferens, and, very often, of a fibro-cellular cord, which unites the peritoneum to the upper part of the tunica vaginalis, and in which encysted hydrocele of the spermatic cord occurs. All these parts are united together by a very lax, areolar tissue, and surrounded by coats, which, reckoning from without, are:—1. The skin and areolar membrane. 2. A fibro-cellular membrane, formed by the fascia superficialis. 3. A very thin layer, formed by fibres of the cremaster muscle, united archwise before, and often also behind, the cord. 4. The proper sheath of the spermatic vessels, or the tubular prolongation furnished by the fascia transversalis to the chord, on a level with the superior orifice of the inguinal canal. The spermatic cord is commonly shorter on the right side than on the left; and of a different size in different individuals. It ascends, almost vertically, from the superior margin of the testicle as far as the lower orifice of the inguinal canal; passes through this canal and enters the abdomen, crossing the epigastric artery. Here it forms an evident elbow, directing its course backward. At this part, also, the organs composing it separate from each other:—the vas deferens descending into the pelvis to pass behind the bladder;—the blood-vessels and lymphatics ascending towards the lumbar region, &c.

SPERMATIC FILAMENTS, Spermatozoa—s. Fluid, Sperm.

SPERMATIC GANGLION. A large ganglion, formed by branches from the hypogastric ganglion, and from the spermatic plexus. It supplies the fundus uteri. Besides these ganglia, Dr. Robt. Lee describes *ves'ical and vag'inal ganglia*, and anterior and posterior *subperitone'al ganglia and plexuses*, which communicate with the preceding, and constitute an extensive nervous rete over the entire uterus.

SPERMATIC LIQUOR, Liquor seminis, see Sperm. SPERMATIC PASSAGES or WAYS, *Via Spermat'icæ*, are the canals concerned in the excretion of semen. See Testicle.

SPERMATIC PLEXUSES of nerves, *Plexus testiculares*, are two in number, and are furnished by the renal plexuses. Their filaments, called *Spermatic nerves*, follow the arteries of the same name to the testicle in man; and to the ovary and Fallopian tube in the female,—*ova'rian nerves*. They

SPERMATOZOÏA, Zoosper'mata, from σπέρμα, σπέρματος, 'sperm,' and ζῷον, 'animal.' Animal'cula semin'alia seu spermat'ica, Fila spermat'ica, Corpuscula sem'inis, spermatococ'ci, Ver-

SPHENOID BONE, *Sphenoïdes os*, *Os basila'rē*
 seu cuneifor'mē seu cu'neo compara'tum seu sphenoid.

wida'lē seu multiformē seu az'ygus seu papilla'rē seu polymorphon seu papilla'rē seu bazilla'rē seu alafur'mē seu sphenoid'um seu respic'or'mē seu inconjugat'um seu pterygoideum seu ala'tum seu colato'rū, Pterygoid bone. An azygous bone, situate on the median line, and at the base of the cranium. It articulates with all the other bones of that cavity; supports them, and strengthens their union. Its form is singular, and resembles a bat with its wings extended. It has, 1. An *inferior or guttural surface*, on which is situate the crista, that joins the vomer; a channel, which concurs in forming the pterygo-palatine foramen; the pterygoid process; the pterygoid fossa; the scaphoid depression; the Vidian or pterygoid canal; the foramina—ovale, spinale, &c. 2. A *superior or cerebral surface*, on which are:—the clinoid processes; the pituitary fossa; the foramina (ovale, rotundum, and spinale); the *Apophysis of Ingrassias or lesser wing*; the foramen opticum, &c. 3. An *occipital or posterior surface*, which is articulated with the basillary process of the occipital bone. 4. An *anterior or orbital nasal surface*; having, anteriorly, a *crista* to unite with the ethmoid bone; and, on each side, a round aperture, which leads into two cavities in the substance of the bone, separated by a middle septum, and called the *sphenoidal sinuses*. 5. Two *zygomato-temporal or external surfaces*, which correspond to the temporal and zygomatic fossæ.

Some divide the sphenoid into *body or middle portion*; and *ala*, which are four in number, and are subdivided into *great (Temporal Plates or Wings, Ala magna)* and *little (Apophyses of Ingrassias, Ala minores, Processus eniformes)*.

The *Sphenoid suture* surrounds the bone.

SPHENOID, SPINOUS PROCESS OF THE, Sphenoid spine.

SPHENOID'AL, Sphenoid'al. That which belongs or relates to the sphenoid bone.

SPHENOIDAL COR'NUA, Cor'ua Sphenoid'al. (F.) *Cornets sphénoïdaux ou de Bertin, Osselets de Bertin, Ossicula Bertin, Ossa triangula'ria, Pyr'amids of Wistar.* Two small, thin, and curved bones, situate between the sphenoid and ethmoid, with which they are confounded in the adult. They have the shape of a hollow pyramid, with the base turned backwards; and are developed by a single point of ossification. They are articulated with the sphenoid, ethmoid, palate bone, and vomer.

SPHENOIDAL OR SUPERIOR ORBITARY FISSURE, Foramen lac'rum superius, (F.) Fente sphénoïdale, is a large fissure, situate between the great and little ala of the sphenoid. It is seen at the upper and back part of the orbit between which and the cranium it is a means of communication, and transmits the third, fourth, and sixth pair of nerves, and the first branch of the fifth pair, to the eye and its appendages.

SPHENOIDAL SPINE, Spina seu Crista sphenoid'al. (F.) *Épine sphénoïdale, Spinous Process of the Sphenoid*;—1. A projecting crista at the inferior surface of the sphenoid-bone, for articulation with the vomer. 2. A triangular process, *Apophyse sous-temporale* (Ch.), met with near the posterior margin of the same bone, behind the foramen spinale. At the point of the spinous process, a *styloid process* is frequently met with.

SPHENOIDES, Cuneiform, Sphenoid—s. Os, Sphenoid bone.

SPHENOMAX'ILLARY, Spheno-maxilla'ris. That which relates to the sphenoid and maxillary bones.

SPHENOMAXILLARY FISSURE, Inferior orbitar Fissure, (F.) Fente sphéno-maxillaire ou orbitaire inférieure; called, also, *Foramen lac'rum inferius seu spheno-maxilla'rē*, is situate at the

posterior part of the angle formed by the union of the internal and inferior parietes of the orbit. It is constituted, above, by the sphenoid bone; below, by the superior maxillary and palate bones; and, before, by the malar bone. It is narrower at the middle than at the extremities, and forms a communication between the orbital cavity and the zygomatic fossa.

SPHENOMAXILLARY FOSSA is a name given by some anatomists, to a depression at the union of the sphenomaxillary and pterygomaxillary fissures.

SPHENO-ORBITAR. A name given by Bérclard to the anterior part of the body of the sphenoid bone, which is developed by a variable number of points of ossification.

SPHENOPALATINE, Sphenopalati'nus. That which relates to the sphenoid and palate bones.

SPHENOPALATINE ARTERY, Large lateral nasal A., is the termination of the internal maxillary. It enters at the posterior part of the superior meatus of the nose, through the sphenopalatine foramen, and spreads its numerous branches on the pituitary membrane covering the septum, the cornua, and the meatus.

SPHENOPALATINE FORAMEN is a round aperture, formed by the vertical portion of the os palati and the sphenoid. It establishes a communication between the nasal fossæ and the zygomatic fossa.

SPHENOPALATINE GANGLION, Ganglion of Meckel, Sphenoid'al G. (Ch.) A small, nervous, cordiform, or triangular ganglion, of variable size, situate without the foramen sphenopalatinum, in the pterygomaxillary fissure. It seems suspended by several nervous filaments to the trunk of the superior maxillary nerve, and gives off *internal or sphenopalatine filaments, inferior or palatine filaments, and a posterior filament*, which is the *Vidian or pterygoid nerve*.

SPHENOPALATINE NERVES, Lat'eral nasal nerves, arise from the ganglion—just described—at its inner part, and enter the nasal fossæ by the sphenopalatine foramen. They are five or six in number, and distribute their filaments to the outer and inner parietes of the nasal fossæ. One of the most remarkable branches is the *Naso-palatine*.

SPHENOPALATINUS, Levator palati.

SPHENOPARI'ETAL, Spheno-parieta'lis. That which belongs or relates to the sphenoid and parietal bones.

SPHENOPARIETAL SUTURE is formed by the articulation of the extremity of the greater ala of the sphenoid with the anterior and inferior angle of the parietal bone.

SPHENO-PTERYGO-PALATINUS, Circumflexus—s. Salpingostaphylinus, Circumflexus—s. Salpingo-mallens, Laxator tympani.

SPHENOSIS, see Wedged.

SPHENOSTAPHYLINUS, Levator palati.

SPHENOTEMP'ORAL, Spheno-tempora'lis. That which belongs to the sphenoid and temporal bones.

SPHENOTEMPORAL SUTURE is the suture at the articulation of the great ala of the sphenoid bone with the squamous portion of the temporal. Bérclard gives the name *sphenotemporal* to the posterior part of the body of the sphenoid, which is developed by distinct points of ossification.

SPHEX, Wasp.

SPHINCTER, σφιγκτήρ, Constrict'or, Musc'ulus constrict'orius, from σφίγω, 'I constrict.' A name given to several annular muscles, which constrict or close certain natural openings.

SPHINCTER ANI, Annula'ris Ani. Many anatomists have described two sphincter muscles of the anus:—1. The *S. exter'ius seu cuta'neus, An-*

pidis'cos, *Coccygio-cutant-sphincter*, *Orbicularis Recti*, *Constrictor Ani*, *Coccygio-anal*, (Ch.) It is situate around the anus; is elliptical from before to behind; flattened, and pierced at its middle. Its fibres describe concentric arcs, which are attached, behind, to the extremity of the coccyx, by a dense, areolar substance; and are confounded, anteriorly, with the bulbo-cavernosi and transversi perinei muscles. This muscle contracts and closes the anus. 2. The inner or internal *Sphincter Ani*, *Sphincter intestinalis* of Winslow, is by many anatomists considered as the termination of the circular fibres of the rectum. It is annular, and situate around the inferior extremity of the rectum, to the extent of about a finger's breadth. It has the same uses as the other.

SPHINCTER GULÆ, *Constrictores pharyngis* — s. *Ilei*, *Bauhin*, valve of — s. *Intestinalis*, *Sphincter ani internus* — s. *Labiorum*, *Orbicularis oris* — s. *Oculi*, *Orbicularis palpebrarum* — s. *Palpebrarum*, *Orbicularis palpebrarum* — s. *Pylori*, see *Pylorus* — s. *Vaginæ*, *Constrictores cunni* — s. *Ventriculi*, *Pylorus*.

SPHINCTER VESICÆ, *Sphincter of the bladder*. Some anatomists have given this name to whitish, elastic, and circular fibres, which surround the neck of the bladder, but do not constitute a particular muscle. Morgagni has given the name *PSEUDO-SPHINCTER* to the anterior fibres of the levator ani, which pass beneath the neck of the bladder, and, by their contraction, close that opening. See *Compressor urethræ*.

SPHONDYLIIUM, *Heracleum spondylium*.

SPHONGUS, *Fungus*, *Spongia*.

SPHRAGIDONYCHARGOCOMETE'A, from *σφραγίς*, 'a seal,' *οὐρά*, *οὐρύς*, 'a nail,' *αργός*, 'white,' or perhaps *αργυρός*, 'silver,' and *κομῆα*, 'I adorn.' A charlatan who adorned his fingers to the very nails with rings. — *Aristophanes*, *Hippocrates*.

SPHYGMA, *Pulse*.

SPHYGMIC, *Sphygmicus*, (F.) *Sphygmique*, from *σφύγμος*, 'pulse.' Relating to the pulse: — pulsatory.

SPHYGMIC ART, *Ars Sphygmica*. The art of judging by the pulse, in health or disease.

SPHYGMICA DOCTRINA, *Sphygmologia*.

SPHYGMICUS, *Throbbing*.

SPHYGMOCEPHALUS, *Crotaphe*.

SPHYGMODES, *Throbbing*.

SPHYGMOLEGIUM, *Pulsilegium*.

SPHYGMOGRAPH, from *σφύγμος*, 'the pulse,' and *γραφω*, 'I describe.' An instrument which, when applied over an artery, indicates the character of the pulse as to force and extent of undulations. — *Vierordt*.

SPHYGMOLOG'IA, *Sphygmology*, *Sphygmica Doctrina*, *Ars Sphygmica*, from *σφύγμος*, 'the pulse,' and *λογος*, 'a description.' The doctrine of the pulse.

SPHYGMOMETER, *Pulsilegium*.

SPHYGMOS, *Pulsation*, *Pulse*.

SPHYGMOSCOPIUM, *Pulsilegium*.

SPHYRA, *Malleolus*.

SPHYXIS, *Pulsation*.

SPIC, *Lavandula Spica*.

SPICA, *Fascia repens*, *Spica bandage*, (F.) *Épi*. A bandage so called in consequence of its somewhat resembling a spike of barley. The turns of the bandage cross like the letter V; — each leaving a third of the roller uncovered. It is distinguished into *ascending* and *descending*. It may be applied over various parts of the body, and in a different manner in each case: thus, there is the *Spica seu Fascia inguinalis*. *Spica inguinalis duplex*, the spica for the shoulder, and another for the thumb.

SPICA, *Lavandula Spica* — s. *Alpina*, *Valeriana*

Celtica — s. *Celtica*, *Valeriana Celtica* — s. *Indica*, *Nardus Indica* — s. *Nardi*, *Nardus Indica*.

SPICEBERRY, *Gaultheria*, *Laurus Benzoin*.

SPICEBUSH, *Laurus Benzoin*.

SPICES, *FOUR*, see *Myrtus pimenta*.

SPICEWOOD, *Laurus Benzoin*.

SPICILLUM, *Specillum*.

SPIDER, see *Aranæa tela*.

SPIDERWORT, *Liliago*, *Tradescantia Virginica*.

SPIGELIA, *S. Marilandica*.

SPIGE'LIA ANTHE'L'MIA, *Demera'ra Pinkroot*, a native of South America and the West India Islands, has properties like the

SPIGE'LIA MARILAN'DICA, *S. Lonic'era*, *Anthe'l'mia*, *Lonic'era Marilan'dica*, *Perennial Wormgrass* or *Indian Pink*, *Caroli'na Pink*, *Starbloom*, *Wormroot*. *Nat. Ord. Gentianacææ*. *Class.* *Pentandria*. *Order*, *Monogynia*. *Indigenous*. The root — *Spigelia* (Ph. U. S.) *Pinkroot* — is celebrated as an anthelmintic, particularly in cases of lumbrici.

It is, also, asserted to have been found serviceable in remittent fever. It is a narcotico-acrid.

Dose, gr. x to ʒss.

SPIGNEL, *Æthusa meum*.

SPIKENARD, *Conyza squarrosa*, *Nardus Indica* — s. *American*, *Aralia racemosa* — s. *False*, *Smalacina racemosa* — s. *Small*, *Aralia radicaulis* — s. *Tree*, *Aralia spinosa*.

SPILANTHES ACME'LLA, *Spilanthus acmella*.

SPILANTHUS ACME'LLA, *S. ciliata* seu *fimbriata*, *Spilanthus acmella*, from *σπίλος*, 'a spot,' and *ανθος*, 'a flower,' owing to its dotted or speckled flower, *Bidens acmella*, *Achmella*, *Acemella*, *A. Mauritanica*, *Verbes'ina acmella*, *Balm-leaved Spilanthus*. *Ord. Compositæ*. *Sez. Syst.* *Syngenesia Polygamia equalis*. This plant possesses a glutinous, bitter taste, and fragrant smell. The herb and seed are said to be diuretic and emmenagogue. They have been used in dropsies, jaundice, fluor albus, and calculous complaints; given in infusion.

SPILANTHUS BALM-LEAVED, *Spilanthus acmella* — s. *Ciliata*, *S. acmella* — s. *Fimbriata*, *S. acmella*.

SPILANTH'US OLERA'CEA, *Spear-leaved Spilanthus*, (F.) *Cresson de Para*. A tincture of the plant has been recommended in toothach.

SPILL, see *Nævus*.

SPILOMA, see *Nævus*.

SPILOSIS, *Epicchrosis* — s. *Ephelis*, *Ephelides* — s. *Poliosis*, *Poliosis*.

SPILSBURY'S ANTISCORBUTIC DROPS. An empirical preparation, formed of *hydrarg. oxymercur.*, *rad. gentian.*, *cort. aurant.*, *sicc. aa ʒij*; *antimon. crud.*, *santal. rubr. aa ʒi*; *spiritus vini rectific.*, *aqua, aa ʒviij*.

SPINA, 'a thorn,' *Spine*, (F.) *Épine*. A process on the surface of a bone, which has been compared to the spines or thorns on certain vegetables. The chief processes of this name are: — the *nasal spine*, the *spine of the scapula*, the *spine of the ischium*, the *four iliac spines*, the *palatine spine*, the *maxillary*, the *sphenoid*, &c. The *spine of the back* is the collection of vertebræ constituting the *vertebral column*.

SPINA, *Penis* — s. *Acida*, *Oxyacantha Galeni* — s. *Egyptiaca*, see *Acaciæ gummi* — s. *Alba*, *Carduus marianus*, *Mespilus oxyacantha*, *Onopordum acanthium* — s. *Bifida*, *Hydrorachis* — s. *Cervina*, *Rhamnus* — s. *Domestica*, *Rhamnus* — s. *Dorsi*, *Vertebral column*, see *Nasus* — s. *Dorsi introrsum Flexa*, *Lordosis* — s. *Ferrea*, *Pin* — s. *Helmontii*, *Aguillon* — s. *Hirci*, *Astragalus verus* — s. *Infectoria*, *Rhamnus* — s. *Nodosa*, *Rachitis* — s. *Sphenoidalis*, *Sphenoidal spine*.

SPINA VENTO'SA, *Spina ventositas*, *Tere'do*,

Osteothrocacé, Fungus seu Tumor fungosus articuli, *Lupia junctura*, *Hyperspongia*, *Flatus spinæ*, *Athrocacé*, *Pædathrocacé*, *Sideratio Ossis*, Cancer seu Gangræna Ossis, *Exosto'is*, *White Swelling* (of some). A term of no definite meaning, as is obvious from these various words having been considered its synonyms. By some, it is defined to be—a disease of the osseous system, in which the texture of the bone dilates, seeming to be distended with air, and constituting a variety of osteo-sarcoma. By others, it is considered to be a tumour arising from an internal caries of a bone; occurring most frequently in the carpus or tarsus. The term itself is a translation from the Arabic of Rhazes. See, also, *Hydrarthrus*, and *Mollities Ossium*.

SPINA VERTEBRALIS, Vertebral column.

SPINACH, Spinacia.

SPINACHIA, Spinacia.

SPINA'CIA, *Spina'chia*, *Spina'cia olera'cea*, *Spin'age*, *Spinach*, (F.) *Épinard*. Ord. Chenopodiaceæ. Sex. Syst. Dicaia Pentandria. A plant which resembles the cabbage in its dietetic powers. The leaves boiled, with the addition of oil, form a good emollient cataplasm. It has been used in phthisical complaints; but its medicinal properties, if it have any, are not now regarded.

SPINACIA OLIERACEA, Spinacia.

SPINÆ, Spinous processes—s. *Ventositas*, *Spina ventosa*.

SPINAGE, Spinacia.

SPINAL, *Spina'lis*, *Spino'eus*, *Spina'tus*, *Rachid'ian*, (F.) *Épinière*, *Rachidien*, from *spina*, 'the spine.' That which relates to the vertebral column.

SPINAL ARTERIES, *Arteriæ spinales*, are two in number, viz: 1. The posterior spinal, *Artère médiane postérieure du Rachis*, (Ch.) It arises from the vertebral, near the corpora pyramidalia, and descends on the posterior surface of the spinal marrow, distributing its ramifications to it. 2. The anterior spinal artery, *A. médiane antérieure*, (Ch.) is larger than the last, and arises, also, from the vertebral. It descends, in a serpentine manner, upon the anterior surface of the marrow; furnishes ramusculi to it, and unites with that of the opposite side, opposite the foramen magnum occipitis. A very tortuous branch arises from this union, which descends as far as the inferior extremity of the marrow, to which it sends numerous divisions.

The term *Spinal Arteries* or *Rachidian Arteries* is also given, in the abstract, to all the arteries of the spinal marrow and vertebral canal. The same may be said of the veins and nerves.

SPINAL CORD, *Medulla spinalis*.

SPINAL FORAM'INA, (F.) *Trous rachidiens*, in the abstract, are the foramina formed by every two contiguous vertebrae, through which the spinal nerves issue. See *Vertebral*. The term *Foramen Spinal'e* seu *spino'eum* is especially applied, however, to a small foramen, in front of the spinous process of the sphenoid bone, through which the middle artery of the dura mater enters the cranium. It is, likewise, called *Foramen epino-spino'eum*, (F.) *Trou Spéno-épineux* ou *petit rond*, *Trou épineux*.

SPINAL GANGLION, see *Encephalon*.

SPINAL IRRITA'TION, *Rhachialgi'tis*, *Rhachial'gia*, *Neural'gia spin'al'is*, *Notal'gia*. A modern pathological view, which refers most nervous diseases to irritation of the spinal cord. This irritation is presumed to be indicated by tenderness on pressure over the spinous process of one or more vertebrae, or over the nerves proceeding from the cord and distributed to the parts at the sides of the spine. Such tenderness, however,

by no means indicates the pathological condition in question, as it is often met with in those enjoying perfect health. The treatment advised is cupping and counter-irritation on each side of the spine, which may be beneficial in such diseases, no matter what part of the frame may be in a morbid state, by exciting a new and revelent impression on a very sensible portion of the cutaneous surface.

SPINAL NERVE, *Accessory of the Par vagum* or *8th pair*, *Accessory nerve of Willis*, *Spinal Accessory*, *Superior res'piratory N.*, *Eleventh pair of encephalic nerves*, (F.) *Spino-cranio-trapézien*, *Trachélo-dorsal* (Ch.), arises from the medulla spinalis, within the vertebral canal, between the anterior and posterior roots of the cervical nerves, at a greater or less distance from the cranium. The roots unite to form the nerve, which ascends into the cranium through the foramen magnum of the occipital bone, and issues by the foramen lacerum posterius, crossing the sterno-cleido-mastoides, to which it gives filaments, and losing itself entirely on the trapezius muscle. The pneumogastric and spinal accessory nerves together—*nervus vagus cum accessorio*—resemble the spinal nerves; the former, with its ganglion, being the posterior root; the latter, the anterior.

SPINAL NERVES, Vertebral nerves—s. *Prolongation*, *Medulla spinalis*.

SPINAL SYSTEM OF NERVES, see *Nerves*.

SPINALES COLLI MINORES, *Interspinales colli*—s. et *Transversales lumborum*, *Transversalis dorsi*.

SPINALIS CERVICIS, *Semi-spinalis colli*—s. *Colli*, *Semi-spinalis colli*.

SPINALIS DORSI, *Spinalis Dorsi major*, (F.) *Grand épineux du dos*. Winslow calls thus some fleshy fasciculi, which are situate on the lateral surfaces of the spinous processes; from the third dorsal vertebra to the first or second lumbar; and which form part of the transverso-spinalis of most authors. The same anatomist calls—*Spinalis Dorsi minor*, *Petit épineux du dos*—some small, fleshy fibres, situate on each side of the interspinal ligament. They are short, flat, and pass from one spinous process to the other. Like the preceding, they form part of the transverso-spinalis. All these fleshy fasciculi strengthen the vertebral column.

SPINATI, *Interspinales colli*.

SPINATUS, *Spinal*.

SPINDLE TREE, *Eunonymus Americanus*.

SPINE, Vertebral column—s. *Curvature* of the, *Gibbositas*—s. *Hæmal*, *Sternum*—s. *Neural*, *Spinous process*.

SPINGEL, *Anethum*.

SPINA-AOXOID-OCCIPITALIS, *Rectus capitis posticus major*—s. *Axoido-trachéli-atloldien*, *Obliquus inferior capitis*.

SPINITIS, *Myelitis*.

SPINNICK, *Nanus*.

SPINO-DORSITIS, *Myelitis*.

SPINOLA, see *Hydrorachis*.

SPINOSUS, *Spinal*.

SPINOUS, *Spino'eus*, (F.) *Épineux*, from *spina*, 'a thorn.' Having the shape of a spine or thorn.

SPINOUS PROC'ESSES or APOPH'YSES of the *Vertebrae*, *Acan'thae*, *Spinæ*, *Cynol'ophoi*, *Neural Spines* of Mr. Owen, (F.) *Apophyses épineuses*, are situate at the posterior part of each vertebra, and afford attachment to the muscles, whose office it is to extend the spine. See *Vertebra*.

SPINTHEROPIA, *Confusio*.

SPINULA, *Pin*.

SPIRAC'ULA, (F.) *Spiracules*, from *spiro*, 'I breathe,' 'Breathing-holes.' Respiratory pores of the skin.

SPIRÆ CEREBRI, *Convolutions*, *cerebral*.

SPIRÆA DENU DATA, *S. ulmaria*.

SPIRÆA FILIPEN'DULA, *Filipendula*, *Saxifraga rubra*, *Dropwort*, (F.) *Filipendula*. Ord. Rosaceæ. Sex. Syst. Icosandria Pentagynia. The root of this plant is said to possess astringent and lithontriptic virtues.

SPIRÆA LOBATA, *S. Palma'ta*, *Queen of the Prairie*, *American Meadowsweet*. An oil, resembling oleum gaultherie, is obtained from the roots and stems.

SPIRÆA OPULIFO'LIA, *Opulus-leaved Spiræa*, *Nine Bark*, has similar properties.

SPIRÆA PALMATA, *S. lobata*.

SPIRÆA TOMENTOSA, *Hardhack*, *Red meadowsweet*, *Steeple bush*, *Rosy bush*, *White leaf*. The root of this indigenous species, *Spiræa* (Ph. U. S.), which is abundant in the northern States of the Union, is tonic and astringent; and is usually administered either in the form of extract or decoction, (*Spirææ* 3ss; *aque* Oj.)

SPIRÆA TRIFOLIATA, *Gillenia trifoliata*.

SPIRÆA ULMA'RIA, *S. denuda'ta*, *Ulmara*, *U. palustris*, *Regi'na Prati*, *Barba capræ*, *Bar'bula capri'na*, *Meadow Sweet*, *Queen of the Meadows*, (F.) *Ulmaire*, *Spirée ulmaire*, *Reine des Prés*, *Vignette*, *Herbe aux Abeilles*. The leaves have been recommended as mild astringents and diuretics—the flowers as antispasmodics and diaphoretics.

SPIRAMENTA PULMONUM, see Pulmo.

SPIRAMENTUM ANIMÆ, Pulmo.

SPIRAMINA PALPEBRARUM, *Lachrymal puncta*—s. Pulmonum, see Pulmo.

SPIRATIO, Respiration.

SPIRÉE ULMAIRE, *Spiræa ulmaria*.

SPIRIT, *Spir'itus*; from *spirare*, 'to breathe, to exhale' (F.) *Espirit*, *Alcolcoat*. A name given to every liquid product of distillation. Spirits were formerly distinguished into—*inflammable*, *acid*, and *alkaline*; and consequently a number of substances were crowded together, which often resembled each other in no other property than in being volatile. The term is now confined to alcoholic liquors, of which the following are the chief:

Arrack. Distilled from coarse palm sugar, named *jaggery*, fermented with the bark of the *Mimosa leucophlea*; also from rice and the fermented juice of the Palm; made in India. Its varieties are:—

1. *Mahwah Arrack*. Made in India from the flowers of the Madhuca tree, *Bassia butyracea*.

2. *Tuba*. Made from palm wine, in the Philippine Islands.

Araka. Distilled, by the Tartars, from koumis, fermented mares' milk.

Araki. Distilled from dates, by the Egyptians.

Arika. A variety of koumis, distilled from fermented mares' milk, in Tartary and Iceland.

Brandy. Distilled from wine, figs, peaches, persimmon, apple, mulberries, and sometimes other fruits, in Europe, Asia, North and South America, wherever wine is made. The best brandy is that of Cognac; the next, that of Bordeaux and Rochelle. The varieties are—

1. *Aguardiente*, (S.) In Peru, the common brandy obtained from grapes is the *Aguardiente de Pisco*, so called, because shipped at the port of Pisco. Another kind, much dearer, and of excellent flavour, is made from Muscatel grapes, and is called *Aguardiente de Italia*. It is sometimes seen in the United States.

2. *Lau*. Made from rice, in Siam.

3. *Rakia*. Made in Dalmatia from the husks of grapes mixed with aromatics.

4. *Rossolio*. Made at Dantzic, from a compound of brandy, rossolis, and other plants.

5. *Troster*. Made on the Rhine, from the husks of grapes, fermented with barley and rye.

6. *Seki-kayavodka*. Made from the lees of wine and fruit at Scio.

Geneva, Hollanda, (*genièvre*, French for juniper.) Distilled from malted barley and rye, rectified on juniper berries, in Holland. Its variety is—

Gin. Made in England, from malted barley, rye, potatoes; rectified with turpentine.

Goldwasser. Distilled at Dantzic, from wheat, barley, and rye, rectified with aniseed, cinnamon, and other spices.

Kirschwasser. Distilled from the Malachet cherry, in Switzerland.

Maraschino. Distilled from the Macarska cherry, at Zara, the capital of Dalmatia.

Rum, (supposed to be derived from the terminal syllable of the word *saccharum*: but the aboriginal Americans call this liquor Rum.) Distilled, in the West Indies and South America, from cane-sugar and molasses, and in North America from maple-sugar. Its variety is—

Slatakaia trava. Made in Kamtschatka, from a sweet grass.

Show-choo. Distilled, in China, from the lees of mandarin, a wine made from boiled rice.

Whisky, (supposed to be derived from *usque*, the first two syllables of *usquebagh*, the original name in Ireland.) Distilled, in Scotland and Ireland, from malted and raw barley, rye, oats, and potatoes; and in the south of France, from sloes. In Ireland it was called *buil-ceann*, or madness of the head. The best Scotch whisky is *Glenlivet*, the best Irish, *Enniskoven*.

Y-wer-a. Distilled, in the Sandwich Islands, from the root of the Tee-root, baked, pounded, and fermented.

To these may be added an intoxicating liquor made by the Afghaniastans, from ewes' milk; and that made in Kamtschatka, from a species of mushroom, named *muchumer*.

SPIRIT OF AMMONIA, *FETID*, *Spiritus ammoniæ foetidus*—s. of Ammonia, succinated, *Spiritus ammoniæ succinatus*—s. of Aniseed, *Spiritus anisi*.

SPIRIT, BATHING, FREEMAN'S, see *Linimentum saponis compos.*—s. Bathing, Jackson's, see *Linimentum saponis compos.*—s. Bezoardic of Bussius, see *Bussii spiritus Bezoardicus*—s. Bone, *Liquor volatilis cornu cervi*.

SPIRIT OF BURRHUS, *Burri spiritus matricialis*—s. of Camphor, *Spiritus Camphoræ*—s. of Carraway, *Spiritus carui*—s. of Colchicum, ammoniated, *Spiritus colchici ammoniatus*—s. of Cinnamon, *Spiritus cinnamomi*—s. of Ether, aromatic, *Spiritus ætheris aromaticus*.

SPIRIT, FUMING, OF BEGUIN, *Ammonia sulphuretum*—s. Fuming, of Boyle, *Ammonia sulphuretum*.

SPIRIT OF HARTSHORN, *Liquor volatilis cornu cervi*—s. of Horseradish, compound, *Spiritus armoracæ compositus*—s. of Juniper, compound, *Spiritus juniperi compositus*—s. of Lavender, *Spiritus lavandulæ*—s. of Lavender, compound, *Spiritus lavandulæ compositus*—s. of Mindererus, *Liquor ammoniæ acetatis*—s. of Nitre, sweet, *Spiritus ætheris nitrici*—s. of Nitric ether, *Spiritus ætheris nitrici*—s. of Pennyroyal, *Spiritus pulegii*—s. of Peppermint, *Spiritus menthæ piperitæ*—s. of Pimento, *Spiritus pimentæ*.

SPIRIT, PROOF, *Spiritus tenuior*—s. Pyroxylic, *Spiritus Pyroxylicus*.

SPIRIT OF ROSEMARY, *Spiritus rosmarini*—s. of Salt, *Muriaticum acidum*—s. of Scurvygrass, golden, see *Spiritus armoracæ compositus*—s. of Spearmint, *Spiritus menthæ viridis*—s. of Sulphuric ether, *Spiritus ætheris sulphurici*—s. of Sulphuric ether, compound, *Spiritus ætheris sulphurici compositus*—s. of Turpentine, *Oleum terebinthinæ rectificatum*.

SPIRIT, VERDIGRIS, Aceticum acidum.

SPIRIT OF VITRIOL, Sulphuric acid—s. of Wine, Spiritus rectificatus—s. of Wine and camphor, Spiritus camphoræ.

SPIRIT, WOOD, Spiritus Pyroxylicus.

SPIRITS, ANIMAL, Nervous fluid.

SPIRITUOUS, Alcoholic.

SPIRITUS, Breath, Life, Respiration, Spirit—s. Æthereus nitrosus, Spiritus ætheris nitrici—s. Æthereus vitriolicus, Spiritus ætheris sulphurici.

SPIRITUS ÆTHERIS AROMATICUS, *Æther sulphuricus cum alcoholæ aromatizatus*, *Eliz'ir vitrioli dulcè*, Sweet *Eliz'ir of Vitriol*, *Viyā'n's Eliz'ir of Vitriol*, *Aromatiz' Spirit of Æther*. (Cinnamon, cort. contus. ʒij, cardam. sem. contus. ʒiiss, piperis longi fruct. cont., zingib. rad. concis., sing. ʒj, spiritus ætheris sulphurici. Oj. Macerate, for 14 days, in a stopped glass vessel and strain. Ph. L.) It is used as a stimulant in nervous affections. Dose, fʒss to fʒj.

SPIRITUS ÆTHERIS COMPOSITUS, Spiritus Ætheris sulphurici compositus—s. Ætheris hydrochlorici, Spiritus Ætheris muriatici.

SPIRITUS ÆTHERIS MURIATICI, *Sp. Æ. Hydrochlorici*, *Sp. Muriaticus æthereus*, *Sp. Salis dulcis*, *Æther Muriaticus alcoholicus*, Spirit of hydrochloric or muriatic ether, *Dulcified marine acid*. An old preparation, made by adding one part of muriatic acid to three parts of rectified spirit, and distilling. Its action is like that of the spiritus ætheris nitrici.

SPIRITUS ÆTHERIS NITRICI, *Sp. Ætheris nitrosi seu nitri dulcis seu Æthereus nitrosus*, *Æther nitricus alcoholizatus*, *A'cidum Nitri dulcificatum*, Sweet Spirit of Nitre, Nitre drops, Spirit of Nitric Ether, (F.) *Æther azoteux alcoolisé*, *Liqueur anodine nitreuse*. (Nitrate of Potassa, in coarse powder, lbij, Sulphuric acid, lbiss; Alcohol, Oixss; Diluted Alcohol, Oj; Carbonate of Potassa, ʒj. Mix the nitrate of potassa and the alcohol in a large glass retort, and having gradually poured in the acid, digest with a gentle heat for two hours; then raise the heat, and distil a gallon. To the distilled liquor add the diluted alcohol and carbonate of potassa, and again distil a gallon.—Ph. U. S.) It is refrigerant, diuretic, antispasmodic, and diaphoretic. Dose, gtt. xxv to fʒj. Specific gravity, 0.834.

SPIRITUS ÆTHERIS OLEOSUS, Spiritus Ætheris sulphurici compositus, Oleum æthereum.

SPIRITUS ÆTHERIS SULPHURICI, *Æther Sulphuricus cum Alcoholæ*, *Liquor Æthereus Sulphuricus*, Spiritus Vitrioli dulcis, S. Ætheris vitrioli, *Liquor Sulphuricus alcoholizatus*, Spirit of Sulphuric Ether. (Æther. sulphuric. Oss; ep. rect. Oj. Mix. Ph. L.) Stimulant, diaphoretic, diuretic, and antispasmodic. Dose, fʒss to fʒij.

SPIRITUS ÆTHERIS SULPHURICI COMPOSITUS, (Ph. U. S. 1842), S. ætheris vitrioli compositus seu Ætheris compositus (Ph. U. S. 1851) seu Spiritus æthereus oleosus, *Liquor anodynus seu anodynus mineralis Hoffmanni*, Compound Spirit of Sulphuric Ether, Hoffmann's Anodyne Liquor, (F.) *Æther hydrique alcoolisé*, *Æther sulfurique alcoolisé*, *Esprit d'Æther sulfurique*, *Liqueur anodine d'Hoffman*. (Æther Sulphuric. Oss; Alcohol, Oj; Ol. Æther. fʒij. Ph. U. S. Ph. L.) Mix. A stimulant and antispasmodic. Specific gravity, 0.816.

SPIRITUS ÆTHERIS VITRIOLICI COMPOSITUS, Spiritus ætheris sulphuric. comp.

SPIRITUS AMMONIÆ, *Alkali ammoniacum spirituosum*, *Alcohol ammoniatum*, *Sp. salis ammoniaci*, *Sp. salis ammoniaci vini seu dulcis seu dulcificatus*, *Ammoniacum alcoholizatum*, *Liquor ammoniaci vinosus*, L. ammoniæ spirituosus,

Lixirium ammoniacale vinosum, *Solutio carbonatis ammoniaci spirituosus*, (F.) *Esprit d'ammoniaque*. (Ammon. muriat. in pulv. subtilissim., calcis, aa lbj; alcohol. fʒxx; aquæ, fʒix. The water is employed to slake the lime; the muriate of ammonia is then well mixed with it, and by means of heat, the ammonia is made to pass into the distilled water contained in a quart bottle. Ph. U. S.) Stimulant and antispasmodic. Dose, fʒss to fʒj.

SPIRITUS AMMONIÆ AROMATICUS, *Alcohol ammoniatum aromaticum*, Spiritus salis volatilis oleosus seu volatilis aromaticus seu ammoniæ compositus, *Sal volatile oleum Sylvestri*, *Tinctura aromatica ammoniata*, *Liquor oleosus Sylvestri*, *Lixirium ammoniacale aromaticum*, *Aromatic ammoniated Alcohol*, *Sal volatile Drops*. (Ammon. muriat. ʒv; Potass. carb. ʒviij; cinnamon. cont., caryophyll. cont. aa ʒij; Limon. cort. ʒiv; alcohol., aquæ aa, Ov. M. Distil Ovijss. Stimulant. Dose, fʒss to fʒj.

SPIRITUS AMMONIÆ COMPOSITUS, Spiritus aromaticus.

SPIRITUS AMMONIÆ FÆTIDUS, *Sp. volatilis fætidus*, *Alcohol ammoniatum fætidum*, *Tinctura asafetide ammoniata*, *Fetid Spirit of Ammonia*, *Fit Drops*. (Ammon. muriat. ʒx, Potass. carb. ʒxvj, *Sp. rectificat.*, *Aquæ aa Oij*, (Imp. meas.) *Asafetide*, ʒv, *Distil Oij*, (Imp. meas.) Macerate for 12 hours, and distil Oiss. Ph. L.) Stimulant and antispasmodic. Dose, fʒss. to fʒj.

SPIRITUS AMMONIÆ SUCCINATUS, *Sp. Ammoniæ cum Succino*, *Sp. Salis Ammoniaci lactescens seu lacteus*, *Tinctura ammoniæ composita*, *Ammoniacum succinatum*, *Aqua Luvæia*, *Liquor ex Ammoniâ et Oleo Succini*, L. ex'citans, *Succinated Spirit of Ammonia*, *Common Eau de Luce*. (Mastich. ʒij, ep. rect. fʒix, ol. Lavand. gtt. xiv, ol. succin. gtt. iv, liq. ammon. fʒx. Macerate the mastich in the alcohol, and pour off the clear tincture; add the rest and shake. Ph. L.) Stimulant and antispasmodic. Dose, gtt. x to fʒss, or fʒj.

SPIRITUS ANIMALES, see Nervous fluid.

SPIRITUS ANISI, *Sp. Anisi compositus*, *Aqua anisi fortis*, A. Sem'inum Anisi composita, Spirit of Aniseed, (F.) *Esprit d'anis*. (Anisi sem. cont. lbss, spirit tenuior. cong., aquæ quod satis sit ad prohibendum empyreuma. Ph. L.) Carminative. Dose, fʒss to fʒss. In the compound spirit, angelica is usually an ingredient.

SPIRITUS ANTHOS, Spiritus rosmarini.

SPIRITUS ARMORACIÆ COMPOSITUS, *Sp. Raphani compositus*, *Aqua Raphani composita*, Compound Spirit of Horseradish, (F.) *Esprit de raifort composé*. (Armorac. rad. recent. concis., quant. cort. exsic., sing. lbj, myrist. nuc. contus. ʒss, spirit. ten. cong., aquæ q. s. ad prohibendum empyreuma. Ph. L.) Stimulant and antiscorbutic. [?] Dose, fʒj to fʒiv.

The Golden Spirit of Scurvy grass—is an empirical preparation—is formed by adding gamboge ʒviij, to one gallon of the sp. armoraciæ compositus.

Greenough's Tincture for the teeth is formed of amygd. amar. ʒij, liq. Brestl., cassiæ bacc., aa ʒiv, irid. Florent. ʒij, coccinell., sal. acetosellæ, alum. aa ʒj, ep. vin. rect. Oij, sp. armoraciæ comp. fʒss.

SPIRITUS BEGUINIS, Ammoniæ sulphuretum—s. Bellidis, see Osmittopsis asteriscoides.

SPIRITUS CAMPHORÆ, *Sp. Camphoratus seu vinosus camphoratus*, *Tinctura Camphoræ* (Ph. U. S.), Spirit of Wine and Camphor, Spirit of Camphor, (F.) *Esprit de Camphre*, *Alcool camphré*, *Eau de Vie camphrée*. (Camphor. ʒiv, alcohol. Oij. Dissolve.) Stimulant, anodyne and discutient. Used only externally.

SPIRITUS CARUI, *Sp. Cari Carui, Aqua Sem-inum Carui* fortis seu *Sem-inum Carui, Spirit of Caraway, Strong Carui Water*, (F.) *Esprit de Carvi*. (*Carui* sem. cont. ℥ss, *Sp. tenuior* cong., *aquæ* q. s. Distil a gallon. Ph. L.) Carminative. Dose, fʒj to fʒss.

SPIRITUS CINNAMOMI, *Sp. Lauri Cinnamomi, Aqua Cinnamomi fortis* seu *Cinnam. spirituosæ, Spirit of Cin-namon, Strong Cinnamon Water*, (F.) *Esprit de Cannelle*. (Cort. cinnam. ʒv, sp. rect. Oivss.) Stimulant and carminative. Dose, fʒj to fʒss.

SPIRITUS COLCHICI AMMONIATVS, *Tinctura Colchici Composita*, *Ammoniated Spirit of Colchicum, Compound Tincture of Colchicum*. (Sem. colchic. cont. ʒij, sp. ammon. aromat. Oj. Ph. L.) Narcotic, cathartic, and diuretic. Used in gout, rheumatism, and dropsy. Dose, from gtt. xxx to fʒi.

SPIRITUS CORNU CERVI, *Liquor volatilis C.C.*—s. Fortior, see Alcohol.—s. *Fumans* Beugini, *Ammonie sulphuretum*—s. *Genitalis*, *Aura seminis*—s. *Jamaicensis*, Rum.

SPIRITUS JUNIPERI COMPOSITVS, *Aqua Juniperi composita*, *Compound Spirit of Juniper*, (F.) *Esprit de genivère composé*. (Ol. junip. ʒis, Ol. carui, Ol. fœnicul., aa ʒx; Alcohol dilut. cong. Solve. Ph. U. S.) Stimulant and diuretic. Dose, fʒj to fʒss.

SPIRITUS LAURI CINNAMOMI, *Spiritus cinnamomi*.

SPIRITUS LAVANDULÆ, *Sp. Lavandulæ spicæ* seu *Lavand. simplex, Lavender Water*, (F.) *Esprit de Lavande*. (*Lavand.* recent. ℞ij. Alcohol. cong., *aquæ* Oij. Distil a gallon. Ph. U. S.) Usually made by adding the oil to rectified spirit. Used as a perfume chiefly.

SPIRITUS LAVANDULÆ COMPOSITVS, *Tinctura Lavandulæ composita*, *Red Hartshorn, Lavender drops, Compound Spirit of Lavender*, (F.) *Esprit de Lavande composé*. (Sp. Lavand. Oijj., sp. ros-marini. Oj, cinnam. contus. ʒj, myristic. cont. ʒss, earyophyll. cont. ʒij, santali concis. ʒijj. Mace-rate for 14 days, and filter through paper. Ph. U. S.) Stimulant. Dose, gtt. xl to fʒij.

SPIRITUS LETHALIS, *Carbonic acid*—s. *Lumbrocorum, Liquor volatilis cornu cervi*.

SPIRITUS MENTHÆ PIPERITÆ, *Sp. Menthæ piperitidis, Aqua Menthæ piperitidis spirituosæ, Spirit of Peppermint*, (F.) *Esprit de Menthe poivrée*. (Olei. menth. pip. ʒvss, sp. rectific. Oivss, *aquæ*, q. s. Distil a gallon. Ph. L.) Carminative and stimulant. Dose, fʒss to fʒij.

SPIRITUS MENTHÆ SATIVÆ, *Spiritus menthæ viridis*.

SPIRITUS MENTHÆ VIRIDIS, *Spiritus menthæ sativæ, Aqua Menthæ vulgæ spirituosæ, Spirit of Spearmint*, (F.) *Esprit de Menthe vert*. (Olei. Menth. virid. ʒvss, sp. rectific. Oivss, aq. q. s. Distil a gallon. Ph. L.) Like the last in properties and dose.

SPIRITUS MILLEPEDARUM, *Liquor cornu cervi*—s. *Muriaticus æthereus, Spiritus ætheris muriatici*.

SPIRITUS MYRISTICÆ, *Sp. Myristicæ Moschatæ* seu *Nucis Moschatæ, Aqua Nucis Moschatæ seu Nephritica, Nutmeg Water*, (F.) *Esprit de Muscade*. (Myrist. cont. ʒij, Alcohol. dil. cong., *aquæ* Oj. Distil a gallon.) Cordial and carminative. Dose, fʒj to fʒiv.

SPIRITUS MYRTI PIMENTÆ, *Spiritus pimentæ*—s. *Nitri acidus, Nitric acid*—s. *Nitri dulcis, Spiritus ætheris nitrici*—s. *Nitri duplex, Nitric acid*—s. *Nitri fumans, Nitric acid*—s. *Nitri Glauberi, Nitric acid*—s. *Nitri simplex*, see *Nitric acid*—s. *Nitri vulgaris*, see *Nitric acid*—s. *Nucis moschatæ, Spiritus myristicæ*—s. *Ophthalmicus Mindereri, Liquor ammoniæ acetatis*.

SPIRITUS PIMENTÆ, *Sp. Myrti Pimentæ seu Pimento, Aqua Mirabilis, Spirit of Pimento*. (Ol. Piment. f. ʒij, Alcohol. dilut. cong. Solve. Ph. U. S.) Cordial and carminative. Dose, fʒj to fʒiv.

SPIRITUS PULEGII, *Aqua Pulegii spirituosæ, Spirit of Pennyroyal, Spirituous Pennyroyal Water*, (F.) *Esprit de Pouliot*. (Olei pulegii ʒvij, sp. rect. Oivss, *aquæ*, q. s. Distil a gallon. Ph. L.) Same as spirit of spearmint in properties and dose.

SPIRITUS PYRO-ACETICUS, *Acetone*.

SPIRITUS PYROXYLICUS, *Pyroxylic spirit, Wood spirit, Wood alcohol, Pyroxylic alcohol, Methylic alcohol, Hydrated oxide of methyle, Bihydrate of Methylene*, (F.) *Esprit pyroxylique, Esprit de bois*. This substance is by some supposed to have been the article given in phthisis, by Dr. Hastings, under the names *Naphtha* and *Wood Naphtha*. It was probably, however, the *Pyro-acetic spirit*. See *Acetone*. When wood is subjected to destructive distillation, about one per cent. of an inflammable volatile liquid, along with other products, passes over, which, when separated and purified, constitutes pyroxylic spirit. It is given in the same cases and doses as *Acetone*.

SPIRITUS RECTIFICATUS, *Sp. Vini rectificatus seu vinosus rectificatus, Cœnostagma, Cœnelæum, Pyroæus, Alcohol, A. Vini, Spirit of Wine*, (F.) *Esprit de Vin rectifié*. Rectified spirit is much used in Pharmacy to dissolve resinous and other substances. It is seldom exhibited internally; although it exists in the diluted state in all vinous and spirituous liquors. Externally, it is employed as a discutient.—S. g., according to the London Pharmacopœia, 0.838; according to that of Dublin, 0.840; of Edinburgh and United States, 0.835, (see Alcohol,) and of Paris, 0.819 to 0.827.

SPIRITUS RECTOR, *Aroma*—s. *Raphani compositus, Spiritus armoraciæ compositus*—s. *Roriferus, Gaz sanguinis*.

SPIRITUS ROSMARINI, *Sp. Rosmarini officinalis, Sp. Anthos, Spirit of Rosemary, Hungary Water*, (F.) *Esprit de Romarin*. (Ol. rosmarini fʒiv, Alcohol. cong. Solve. Ph. U. S.) Stimulant. Dose, fʒj to fʒiv.

SPIRITUS SALIS ACIDUS seu *FUMANS*, *Muriaticum acidum*—s. *Salis ammoniaci, Spiritus ammoniæ*—s. *Salis ammoniaci aquosus, Liquor ammoniæ*—s. *Salis ammoniaci lactescens seu lacteus, Spiritus ammoniæ succinatus*—s. *Salis ammoniaci sulphuratus, Ammonia sulphuretum*—s. *Salis dulcis, Spiritus ætheris muriatici*—s. *Salis Glauberi, Muriaticum acidum*—s. *Salis marini, Muriaticum acidum*—s. *Salis marini coagulatus, Potassæ murias*—s. *Salis volatilis oleosus, Spiritus ammoniæ aromaticus*—s. *Sulphurico-æthereus camphoratus, Tinctura ætheræ composita*—s. *Sulphurico-æthereus martialis, Tinctura seu Alcohol sulphurico-æthereus ferri*—s. *Sulphuris, Sulphurous acid*—s. *Sulphuris per campanam, Sulphurous acid*—s. *Sulphuris volatilis, Ammonia sulphuretum*—s. *Sulphurosus volatilis, Sulphurous acid*.

SPIRITUS TENUIOR, *Alcohol dilutum, Spiritus vinosus tenuior, Proof Spirit*, (F.) *Esprit de vin délayé*. Generally made for pharmaceutical purposes, by mixing equal parts of rectified spirit and water. S. g. Lond. 0.920; Edinb. and United States, 0.935. It is employed in the same cases as those in which the alcohol is used:—chiefly, as a solvent of vegetable matters, in the form of tinctures.

SPIRITUS VENERIS, *Aceticum acidum*—s. *Vini Gallici, Brandy*—s. *Vini rectificatus, Spiritus rectificatus*—s. *Vinosus camphoratus, Spiritus camphoræ*—s. *Vinosus tenuior, Spiritus tenuior*—s.

Vitalis, Nervous fluid — s. Vitrioli, Sulphuric acid — s. Vitrioli acidus Vogleri, Sulphuricum acidum dilutum — s. Vitrioli coagulatus Mynsichti, Potassæ sulphas — s. Vitrioli dulcis, Spiritus ætheris sulphurici — s. Vitrioli phlogisticatus, Sulphurous acid — s. Volatilis aromaticus, Spiritus ammoniæ aromaticus — s. Volatilis foetidus, Spiritus ammoniæ foetidus.

SPIROID CANAL, Aquæductus Fallopii.

SPIROMETER; a hybrid term, from *spiro*, 'I breathe,' and *μετρον*, 'a measure': — properly *Anapnom'eter*, *Pneusom'eter*, *Pneumom'eter*, *Pneumatom'eter*, *Pulmom'eter*, *Pneumonon'eter*. Any instrument for measuring the quantity of air concerned in respiration, and, consequently, the capacity of the lungs. This mode of measurement has been called *Spirom'etry*, as it was formerly called *Pulmon'etry*. The term spirometer is now usually applied to a kind of gasometer introduced by Dr. Hutchinson to estimate the quantity of air contained in, or, rather, expelled from the lungs under different circumstances.

SPIROPTERA HOMINIS, see Worms.

SPISSANTIA, Incrassautia.

SPIT, Sputum.

SPITAL, Hospital.

SPITAL, MINERAL WATERS OF. The village of Spital is situate on the south side of the Tweed, near Tweedmouth. The springs there contain chloride of calcium, sulphate of soda, sulphate and carbonate of lime. They act as saline cathartics.

SPITH'AMA, σπιθαμή, *Dodrans*. A measure of twelve fingers' breadth, or about nine inches.

SPITTING, Exspuition.

SPITTLE, Saliva — s. Sweet, see Salivation.

SPLANCHNA, Entrails.

SPLANCHNECTOP'IA, *Splanchnodias'tasis*, *Situs perver'sus seu anom'alus intestin'o'rum*; from *σπλᾶγχνον*, 'an intestine,' *εκ*, 'out of,' and *τοπος*, 'place.' Misplacement or ectopia of the intestines.

SPLANCHNELMINTHA, Worms,

SPLANCHNEURYS'MA, from *σπλᾶγχνον*, 'an intestine,' and *ευρυνω*, 'I dilate.' Morbid dilatation of the intestines.

SPLANCHNIC, *Splanchn'icus*, *Vis'ceral*, (F.) *Splanchnique*, from *σπλᾶγχνον*, 'a viscus.' Relating or belonging to the viscera.

SPLANCHNIC CAVITIES are the three great cavities of the body;—those of the cranium, chest, and abdomen.

SPLANCHNIC NERVES belong to the great sympathetic, and are two in number on each side. They are distinguished into:—1. The great splanchnic nerve, (F.) *Grand splanchn'ic* (Ch.), Its roots—four or five in number—come from the 6th, 7th, 8th, 9th, and sometimes the 10th, thoracic ganglia. They descend inwards, on the sides of the vertebral column, beneath the pleura, and unite into a single trunk, which enters the abdomen through a separation of the fleshy fibres of the pillars of the diaphragm; passes behind the stomach, and divides into several branches, which pass into the semilunar ganglion. 2. The lesser splanchnic nerve, *Renal nerve*, (F.) *Splanchn'ique accessoire* of Walther, *Petit splanchn'ic*, (Ch.,) is composed of two distinct branches, proceeding from the 10th and 11th thoracic ganglia, which unite on the 12th dorsal vertebra into a cord, that pierces the diaphragm singly; enters the abdomen, and divides into two branches, one of which anastomoses with the greater splanchnic nerve, whilst the other sends its filaments to the renal and solar plexuses.

SPLANCHNICA. Medicines adapted to diseases of the bowels. Also, the 2d order of the class

Calicæ of Good; defined,—diseases affecting the collatitious viscera, and characterized by disquiet or diseased action in the organs auxiliary to the digestive process, without primary inflammation.

SPLANCHNODIATASIS, Splanchnectopia.

SPLANCHNOD'YNĒ, from *σπλᾶγχνον*, 'a viscus,' and *ὀδυνη*, 'pain.' Pain in the bowels.

SPLANCHNOGRAPHIA, Splanchnography.

SPLANCHNOG'RAPHY, *Splanchnog'ra'phia*, from *σπλᾶγχνον*, 'a viscus,' and *γραφη*, 'a description.' The part of anatomy which treats of the viscera.

SPLANCHNOLITHI'ASIS; from *σπλᾶγχνον*, 'a viscus,' and *λίθος*, 'a stone.' The formation of a calculous concretion in any viscus.

SPLANCHNOL'OGY, *Splanchnolog'ia*, from *σπλᾶγχνον*, 'a viscus,' and *λογος*, 'a discourse.' A treatise on the viscera:—also, Splanchnography.

SPLANCHNOPATHI'A, from *σπλᾶγχνον*, 'a viscus,' and *παθος*, 'suffering.' A disease of the intestines.

SPLANCHNOSCLERO'SIS, from *σπλᾶγχνον*, 'a viscus,' and *σκληρος*, 'hard.' Induration of a viscus.

SPLANCHNOT'OMY, *Splanchnotom'ia*, *Splanchno'tomē*, from *σπλᾶγχνον*, 'a viscus,' and *τεμνω*, 'I cut.' Dissection or anatomy of the viscera.

SPLAT-FOOT, see Kylosis.

SPLAYFOOT, see Kylosis.

SPLEEN, σπλην, *Lien*, *Hepar sinist'rum seu adulteri'num*, *Fomes ventriculi*, *Milt*, (Sc.) *Melt*, (F.) *Rate*. A soft, spongy, parenchymatous organ; of a more or less deep violet red, situate deeply in the left hypochondrium, below the diaphragm, above the colon, between the great tuberosity of the stomach and the cartilages of the false ribs, and above and anterior to the kidney. Its ordinary length is $4\frac{1}{2}$ inches; its thickness $2\frac{1}{2}$; and its weight 8 ounces. Its parenchyma, which is bathed in blood, contains a multitude of grayish, soft, and semi-transparent granulations—*splenic corpuscles* or *Malpighian bodies* of the spleen. It is covered by a serous membrane, furnished by the peritoneum; and by a proper coat, of a fibrous nature, which is intimately united with it, and transmits into its interior delicate, solid, and very elastic prolongations—*trabeculae*. At its inner edge is a fissure, *Hilus* seu *Porta lie'nis*, by which the vessels and nerves enter the organ. The functions of the spleen are unknown. The best opinion is, that it appertains to hæmaturia, and acts as a diverticulum to the vascular system.

SPLEEN, Hypochondriasis—s. Malpighian bodies of the, see Spleen—s. Supernumerary, *Lienculus*.

SPLEENWORT, Asplenium, A. filix femina, Asplenium scolopendrium—s. Bush, *Comptonia asplenifolia*.

SPLEN, Spleen.

SPLENAL'GIA, *Neural'gia lie'nis*, *Sple'nica*, *Splenodyn'ia*, from *splen*, and *algos*, 'pain.' Pain in the spleen.

SPLENALGIA BENGALENSIS, see Cachexia, spleen—s. Phlegmonodea, Splenitis—s. Subinflammatoria chronica, Splenoncus.

SPLENATROPH'IA, *Atroph'ia lie'nis*, from *splen*, and *atrophia*, 'atrophy.' Wasting or atrophy of the spleen.

SPLENAUXE, Splenoncus.

SPLENECTAMA, Splenoparectama.

SPLENECTASIS, Splenoparectama.

SPLENECTOM'IA, *Splene'ctomē*, from *splen*, *εκ*, 'out of,' and *τεμνω*, 'I cut.' Extirpation of the spleen.

SPLENECTOP'IA, *Disloc'io seu Ectop'ia lie'nis* seu *splenitis*, *Anomalsplenotopie*—Piorry; from *splen*, and *εκτοπος*, (*εκ*, and *τοπος*), 'out of

place.' Dislocation of the spleen. Not a common occurrence.

SPLENEMPHRAX'IS, *Splenophrax'ia*, from *splen*, and *σπασσω*, 'I obstruct.' Obstruction of the spleen.—Vogel. See Splenoncus.

SPLENES, Hypochondriasis.

SPLENET'ICUS, *Sple'nicus*, *Splenit'icus*, *Lien'ous*, (F.) *Rateleux*. That which relates to the spleen: one labouring under diseased spleen. A medicine adapted for diseases of the spleen.

SPLENIC, *Sple'nicus*, *Sple'nus*, (F.) *Splénique*, from *splen*. That which relates to the spleen.

SPLENIC ARTERY. It arises from the cœliac; and passes from right to left, forming many windings along the upper edge of the pancreas, which lodges it in a special groove. It reaches the fissure of the spleen, and divides into several branches that enter the organ. In its course it furnishes the pancreatic, left gastro-epiploic, and the *vasa brevia*.

SPLENIC CACHEXIA, see Cachexia splenica—s. *Corpuscles*, see Spleen.

SPLENIC PLEXUS is a nervous network, which proceeds from the cœliac plexus, and accompanies the splenic artery, sending secondary plexuses to each division.

SPLENIC VEIN, *Vena sple'nica*, *V. lienal'is*, arises from the spleen, and accompanies the splenic artery; proceeding from left to right, to unite with the superior mesenteric vein, and to form, with it, the abdominal vena porta. In its course, it receives veins corresponding to the *vasa brevia*, the right and left gastro-epiploic, the duodenal, pancreatic, coronaria ventriculi, and inferior mesenteric veins.

SPLENICA, Splenalgia. Also, medicines which affect the spleen.—Pereira.

SPLENICUS, Spleneticus, Splenic.

SPLENIOLA, Compress.

SPLENIS TUMOR, Ague cake.

SPLENISATION, Splenization.

SPLENISCUS, Compress.

SPLENITICUS, Splenic.

SPLENIT'IS, from *splen*, and *itis*, denoting inflammation. Inflammation of the spleen. *Empres'ma splenitis*, *Inflammat'io lie'nis*, *I. splenis*, *Splenal'gia phlegmono'dea*, *Lienit'is*, (F.) *Splénite*, *Inflammation de la Rate*. A rare disease, characterized by local pain, swelling, and heat, which sometimes terminates by suppuration. It generally affects the capsule—*Epi-splenit'is*, *Peri-splenit'is*—and must be treated like other inflammations.

SPLENIUM, Compress—s. *Corporis Callosi*, see *Corpus callosum*—s. *Cruciatum*, *Croix de Malte*.

SPLE'NIUS, *S. Cap'itis* seu *Colli*, (F.) *Cervico-dorso-mastoidien et dorso-trachélien*, *Cervico-mastoidien* (Ch.), *Posterior mastoid muscle*. So called from its resemblance to the spleen of certain animals. A muscle situate at the posterior part of the neck, and upper part of the back. It is much broader above than below, where it terminates in a point. It is attached, by its inner edge, to the spinous processes of the first five dorsal vertebrae; to that of the 7th cervical, and to the inferior part of the posterior cervical ligament. By its upper extremity, it is attached to the transverse processes of the first two cervical vertebrae; to the mastoid process, and to the posterior surface of the occipital bone. Some have considered this muscle to be formed of two portions, which they have called *Splenius colli*, and *Splenius cap'itis*. The splenius extends the head, inclines it, and rotates it. If the splenii of each side act together, they extend the head.

SPLENIZA'TION, *Splenizatio*, *Splenifica'tion*, *Splenifica'tio*, (F.) *Splénisation*, from *splen*. A

term applied to the state of the lung in the first or second stage of pneumonia, in which its tissue resembles that of the spleen.

SPLENOCE'LE, *Her'nia lienal'is*, from *splen*, and *κηλη*, 'a tumour.' Hernia formed by the spleen.

SPLENODYNIA, Splenalgia.

SPLENOG'RAPHY, *Splenograph'ia*, from *splen*, and *γραφη*, 'a description.' A description of the spleen.

SPLENOHÆ'MIA, (F.) *Splénohémie*, from *splen*, and *αἷμα*, 'blood.' Congestion or hyperæmia of the spleen, a common phenomenon in many diseases, especially in typhoid and intermittent fevers.

SPLENOID, *Spleno'i'des*, *Spleno'des*, from *splen*, and *εἶδος*, 'resemblance.' Spleen-like. Having the appearance of the spleen.

SPLENOL'OGY, *Splenolog'ia*, from *splen*, and *λογος*, 'a discourse.' A treatise on the spleen.

SPLENOMALA'CIA, *Lienomala'cia*, from *splen*, and *μαλακος*, 'soft.' Morbid softening of the spleen.

SPLENONCIE, Splenoncus.

SPLENON'CUS, *Megalosple'nia*, *Hypertroph'ia* seu *Supernutrit'io splenis* seu *lie'nis*, *Intumescen'tia* seu *Infarctus* seu *Physco'nia lienis*, *Splenal'gia subinflammato'ria chron'ica*, *Splenemphrax'is*, *Splenaux'is*, *Hypersplenotroph'ia*, *Lien ingens*, (F.) *Splénoncie*, *Hypertrophie de la Rate*; from *splen*, and *ογκος*, 'a tumour.' Tumefaction of the spleen. See Ague cake.

SPLENOPAREC'TAMA, *Splene'ctama*, *Splenoparec'tasis*, *Splene'ctasis*, from *splen*, and *παρεκταμα*, 'excessive volume.' Excessive enlargement of the spleen.—Plouquet.

SPLENOPARECTASIS, Splenoparectama.

SPLENOPHRAXIA, Splenemphraxis.

SPLENORRHAG'IA, from *splen*, and *ρην*, 'rupture.' Hemorrhage from the spleen.

SPLENOSCIR'RHUS, from *splen*, and *σκιρρος*, 'hardness.' Scirrhus or cancer of the spleen.

SPLENOT'OMY, *Splenotom'ia*, *Splenot'omie*, from *splen*, and *τεμνω*, 'I cut.' Dissection of the spleen.

SPLINT, Tent. Splintes, from *splijten*, 'to split.' *Hastel'la*, *Ass'ula*, *Asser'culum*, *Plag'ula*, *Narthex*, *Fer'ula*, (F.) *Atelle*, *Éclisse*. A flexible and resisting lamina of wood, used in the treatment of fractures, to keep the fragments in apposition, and prevent their displacement. Splints are made of iron, wood, bark of trees, leather, gutta percha, pasteboard, &c. They are generally padded, and are fixed by a roller or by tapes, according to circumstances.

SPLINTER, (F.) *Esquille*. Same etymon. This term is sometimes applied to a small portion of wood, which occasionally enters the skin, and gives occasion to much irritation, until it is removed,—*Aculeus lig'neus*, (F.) *Écharde*.

It is also applied to fragments that separate from a fractured or diseased bone.

SPLITCLOTH, *Lin'teum scissum*. A bandage for the head, consisting of a central part, and six or eight tails. The single split cloth has four tails. It is regarded as the most convenient bandage for the head, face and jaws.

SPLITROCK, *Heuchera cortusa*.

SPOAT, *Saliva*.

SPODIUM GRÆCORUM, *Album græcum*.

SPOLIARIUM, *Apodyterium*.

SPO'LIATIVE, *Spoliat'iva*, from *spoliare*, *spoliatum*, 'to take away.' (F.) *Spoliatif*. *Phlebotom'ia spoliati'va*, (F.) *Saignée spoliative*. Blood-letting, used for the purpose of diminishing the mass of blood.

SPOLIATORIUM, *Apodyterium*.

SPONDLES, *Vertebrae*.

SPONDYLAL'GIA, *Spondylodyn'ia*, from *σπονδύλος*, 'a vertebra,' and *αλγος*, 'pain.' Pain in the vertebrae. *Cacorrhachitis*. See *Vertebral Disease*.

SPONDYLARTHRI'TIS, from *σπονδύλος*, 'a vertebra,' *αρθρον*, 'a joint,' and *itis*, denoting inflammation. Gouty or rheumatic inflammation of the spinal column.

SPONDYLARTHROCAE, *Vertebral disease*.

SPONDYLI, *Vertebrae*.

SPONDYLIS, *Coccyx*.

SPONDYLI'TIS, *Inflamma'tio ver'tebrae*, from *σπονδύλος*, and *itis*, denoting inflammation. Inflammation of the vertebrae.

SPONDYLIUM, *Coccyx*, *Heracleum spondylium*.

SPONDYLOCAE, *Vertebral disease*.

SPONDYLODYNIA, *Spondylalgia*.

SPONDYLOPATHI'A, from *σπονδύλος*, 'a vertebra,' and *παθος*, 'a disease.' A disease of the vertebral column.

SPONDYLOPYOSIS, *Vertebral disease*.

SPONDYLUS, *σπονδύλος*. A vertebra. Also, the vertex.

SPONGE, *BASTARD*, *Aleyonium*.

SPONGI. The tumefied glands of the neck.

SPONG'IA, *S. officina'lis* seu *Mari'na*, *Spongos*, *Sphongus*, *Sponge*, (F.) *Éponge*. An organized marine substance, of the *Class* *Porifera*, met with in many seas, but especially in the Mediterranean and Arabian. It is supple, elastic, porous, and capable of absorbing fluids in which it may be placed, and of thus increasing in size. It is employed in *surgery*, for cleansing or washing; for absorbing acrid discharges from ulcers; suppressing hemorrhage when the mouth of the vessel can be compressed by it, and for forming tents for dilating wounds; in which case, the sponge is immersed in melting wax, and cooled before being used. It is then called *Spongia cera'ta*, *Waxed*, *Cera'ted* or *Prepared Sponge* or *Sponge Tent*.

SPONGIA ASSA, *S. Usta*—*s. Cerata*, see *Spongia*—*s. Cynosbati*, *Bedegar*.

SPONGIA USTA, *S. Assa*, *Carbo spon'giae*, *Burnt Sponge*, contains carbonate and phosphate of lime, carbonate of soda, charcoal, and iodine. It is used in bronchocoele, scrofulous affections, and herpetic eruptions; and its good effects seem to be dependent upon iodine. It is made by cutting sponge into pieces, and heating it, that any extraneous matter may be separated; burning it in a close iron vessel until it becomes black and friable, and, lastly, rubbing it into very fine powder.—Ph. U. S. Dose, ʒj to ʒiij.

SPONGIÆ LAPIS. A name given to small friable stones found in sponge. They were formerly esteemed lithontriptic.

SPONGIOIDES, *Spongoid*.

SPONGIOLE, *Spon'giolum*, *Spon'giola*, *Sponge'let*; from *σπγγιον*, 'a small sponge.' The soft succulent extremity of the fibrils or true roots of a plant, by which it absorbs or sucks up fluid. Similar spongioles have been supposed to exist at the terminations of the absorbents in animals.

SPONGION. Ancient name of an epithem made of goats' dung; and also of a malagma, which was considered capable of absorbing the serum in dropsies, like a sponge.—Paulus, *Goræus*.

SPONGIO-PILINE, from *spongia*, 'sponge,' and *pilus*, 'a hair.' A fabric, composed of sponge and wool, felted together in three layers, and coated, on one of its surfaces, with caoutchouc, which renders it impermeable. It is of great service as a cataplasm, fomentation, vehicle for lotions, liniments, &c.

SPONGIOSUM (OS), *Ethmoid bone*.

SPONGIOSUS, *Spongoid*.

SPONGOID, *Spongoi'des*, *Spongioides*, *Spongioides*, *Spongy*, from *σπγγια*, 'sponge,' and *ειδος*, 'resemblance.' Resembling sponge. Of the nature of sponge.

SPONGOID INFLAMMATION, *Hæmatodes fungus*.

SPONGOID TISSUE, (F.) *Tissu spongoïde*. An adventitious spongy tissue, found in cases of rickets.—Guérin.

SPONGOS, *Tonsil*.

SPONGY, *Spongoid*.

SPONGY TISSUE, see *Bone*.

SPONSA SOLIS, *Calendula officinalis*.

SPONTA'NEOUS, *Sponta'neus*, (F.) *Spontanée*, from *sponte*, 'voluntarily.' That which occurs of itself, or without any manifest external cause. Evacuations are termed *spontaneous*, which are not produced by medicine. *Sponta'neous las'situde* is that which is not the result of fatigue. Diseases are also *spontaneous* which supervene without apparent cause. *Sponta'neous Amputa'tion* is the separation of a limb from the fetus in utero.

SPONTANEOUS EVOLUTION, see *Evolution*—*s. Version*, *Evolution*, *spontaneous*.

SPONTÉPARITÉ, *Generation*, *spontaneous*.

SPOON, *Umbilicus*.

SPOONWOOD, *Kalmia latifolia*, *Tilia Americana*.

SPORAD'IC, *Spor'ades*, *Sporad'ici*, *Morbi disper'si* seu *sparsi*, (F.) *Maladies sporadiques*, from *σπειρω*, 'I spread.' Diseases which supervene indifferently in every season and situation, from accidental causes, and independently of any epidemic or contagious influence.

SPORE, *Spor'ule*, *Sporum*, *Spor'ulum*, from *σπειρω*, 'I sow.' The reproductive body in cryptogamous plants, which is analogous to the seed of phanerogamous plants.

SPORID'IUM; same etymon. The reproductive body in algaceous plants, which resembles the spore of other plants.

SPORULE, *Spore*.

SPOBUM, *Spore*.

SPOTS, *CEREBRAL*, *Taches cérébrales*—*s. Germinal*, see *Molecule*—*s. Meningitic*, *Taches méningitiques*.

SPOTS, *RED*, (F.) *Taches rouges*, *Rose spots*, *Rose'ola* seu *Pap'ula typh'o'ea*. Small red spots which appear in typhoid fever, from the 8th to the 15th day of the disease. They are most frequently seen on the lower and middle portions of the chest, and upper part of the abdomen.

SPOTS, *ROSE*, *Spots*, *red*.

SPRAIN, corrupted from *strain* [?]*—Subluxa'tio*, *Distort'io*, *Dias'trophé*, *Dia'strem'ma*, *Luxa'tio imperfec'ta*, *Ezarthre'ma*, *Ezarthro'sis*, *Strem'ma*, *Thlasma Stremma*, *Wrench*, (Sc.) *Wramp*, (Prov.) *Kenck*, *Keak*, (F.) *Détorse*, *Entorse*, *Foulure*. A violent straining or twisting of the soft parts surrounding the joints. According to the degree of the sprain, the fibrous parts of the joint may be simply stretched, or they may be ruptured. This affection, which has been regarded as incomplete luxation, occurs, particularly, in the ginglymoid joints;—as in the foot, wrist, knee, &c. The violence causes an afflux of blood; the serum of the blood is poured out, and often the blood itself, owing to the rupture of vessels; and inflammation arises, which ends in resolution or induration. Local, antiphlogistic applications are required, with perfect rest; and, when the inflammatory action has subsided, cold pumping and friction, with the application of a plaster and bandage, will afford support till the parts affected recover their due tone.

SPRAT, *YELLOW-BILLED*, *Clupea thyrssa*.

SPRUCE BEER, *White Spruce Beer*. A beer made as follows: To 10 gallons of water, add 6 pounds of sugar, and 4 oz. of essence of spruce.

Add yeast, and bottle, when ready. It is grateful and antiscorbutic.

SPRUCE BEER POWDERS may be formed of white sugar, 3j, and ʒij; soda subcarb., gr. xxvj; essence of spruce, gr. x, in each blue paper; and acid of tartar, ʒss in each white paper. For half a pint of water.

SPUITIO, Exspuition.

SPUMA ARGENTI, Plumbi oxydum semivitreum—s. Cerevisiæ, Yest—s. Trium draconum, Antimonium muriatum.

SPUMEUX, Frothy.

SPUMOSUS, Frothy.

SPUNK, Boletus ignarius.

SPUR, Sax. *spur*, (F.) *Éperon des Artères*. The angle at which the arteries leave a cavity or trunk.

SPUR, Ergot.

SPURBLIND, Myopic.

SPURGE, CAPER, Euphorbia lathyris—s. Cypress, Euphorbia Cyparissias—s. Flax, Daphne gnidium—s. Flowering, large, Euphorbia corollata—s. Garden, Euphorbia lathyris—s. Greater, Euphorbia palustris—s. Ipecacuanha, Euphorbia ipecacuanha.

SPURGE LAUREL, Daphne laureola.

SPURGE, MARSH, Euphorbia palustris.

SPURGE OLIVE, Cneorum tricoccon, Daphne mezereum.

SPURGE, PETTY, Euphorbia peplus—s. Sea, Euphorbia paralias—s. Spotted, Euphorbia maculata—s. Sun, Euphorbia helioscopia—s. Wart-wort, Euphorbia helioscopia.

SPUTA, PRUNE JUICE, see Prune Juice.

SPUTAMEN, Saliva, Sputum.

SPUTAMENTUM, Saliva, Sputum.

SPUTATIO, Exspuition.

SPUTATION, Exspuition.

SPUTATORIUS, Expectoant.

SPUTUM, plural Sputa, *Apochrem'ma*, *Chrem'ma*, *Empty's'ma*, *Sputa'men*, *Pty'elon*, *Pytis'ma*, *Spit*, from *spuere*, *sputum*, 'to spit.' (F.) *Crachut*.

The secretions ejected from the mouth in the act of spitting. It is composed of saliva, and of the mucus secreted by the mucous membrane of the nasal fossæ and fauces, and often by the membrane of the larynx and bronchia. The sputa are, sometimes, called *eru'ginous*, when of a greenish-yellow colour; *gummy*, when they resemble a thick solution of gum; *frothy*, (F.) *mousseux*, when covered with froth; *rubig'inous*, (F.) *rubigineux* ou *rouillés*, when of a reddish-yellow; and *streaked*, (F.) *striés*, when mixed with lines of blood. Also, saliva.

SPUTUM CRUENTUM, Hæmoptysis—s. Dulce, Saccharorrhœa pulmonalis—s. Oris, Salivation—s. Saccharatum, Saccharorrhœa pulmonalis—s. Sanguinis, Hæmoptysis.

SQUALLING, Vagitus.

SQUAMA, Scale, Squamous portion, Table.

SQUAMÆ, Scaly diseases.

SQUAMARIA, Plumbago Europæa.

SQUAMEUX, Squamous.

SQUAMIFORMIS, Squamous.

SQUAMOUS, *Squamo'sus*, *Lepidoi'des*, *Lepido'des*, *Squamifor'mis*, *Caly*; from *squama*, 'a scale.' That which relates to or resembles a scale.

SQUAMOUS PORTION of the temporal bone, (F.) *Portion squameuse* ou *écailleuse du Temporal*, is the upper part of the temporal, so called from a fancied resemblance to a scale. See Temporal Bone.

SQUAMOUS SUTURE, *Sutu'ra squamo'sa* seu *mendo'sa* seu *lepidoi'des* seu *lepidodes* seu *crani'i squamo'sa*, (F.) *Suture squameuse* ou *écailleuse*. The suture which unites the squamous portion of the temporal bone to the parietal.

Certain scaly cutaneous affections are also called *squamous*, (F.) *squameux*. See Scale.

SQUARE STALK, *Monarda coccinea*.

SQUARRA TONDENS, *Porrigo decalvans*.

SQUATTING PILLS, *Pilulæ opiatæ*.

SQUAW-MINT, *Hedeoma*.

SQUAW-ROOT, *Actæa racemosa*, *Caulophyllum thalictroides*, *Senecio aureus*.

SQUAW-WEED, *Erigeron Philadelphicus*.

SQUEAMISHNESS, *Fastidium cibi*.

SQUEASINESS, Nausea.

SQUELETTE, Skeleton.

SQUELETTOTOLOGIE, Skeletology.

SQUILL, Scilla—s. Vinegar of, *Acetum scillæ*.

SQUILLA, Scilla.

SQUINANCIE, Cynanche, Cynanche tonsillaris.

SQUINANCY, COMMON, Cynanche tonsillaris.

SQUINANTHIA, Cynanche.

SQUINANTHUS, *Juncus odoratus*.

SQUINE, *Smilax China*.

SQUINSY, Cynanche tonsillaris.

SQUINT-EYED, see Strabismus.

SQUINTER, see Strabismus.

SQUINTING, Strabismus.

SQUINZEY, Cynanche tonsillaris.

SQUIRRE, Scirrhus—s. en *Cuirasse*, see Lig-neous.

SQUIRRHE, Scirrhus.

SQUIRREL CORN, *Dicentra Canadensis*.

SQUIRREUX, Scirrhous.

SQUIRROGASTRIE, Scirrhogastria.

SQUIRRUS, Scirrhus.

SQUIRT, Syringe.

STACHELBERG, MINERAL WATERS OF. Stachelberg Springs are far down in the valley of the Linth, in Switzerland. The waters are sulphureous, and are much visited by gouty and rheumatic invalids.

STACHYS FÆTIDA, Galeopsis.

STACE, Myrrha.

STADE, Stage.

STADIUM, Period, Stage.

STA'DIUM ACME SEU STA'SEOS. The period of the height of a disease.

STA'DIUM ANNIHILATIO'NIS SEU CONVALESCENTIE SEU RECONVALESCENTIE. The period of recovery from disease.

STADIUM AUGMENTI SEU INCREMENTI MORBI. The period or stage of increase of a disease.

STADIUM CALORIS, Stage, (hot)—s. Convalescentiæ, S. annihilationis.

STADIUM DECREMENTI. The period or stage of decrease of a disease.

STADIUM FRIGORIS, Stage, (cold)—s. Incrementi Morbi, S. Augmenti morbi—s. Latentis Contagii, Prodromus—s. Opportunitatis, Prodromus—s. Prodromorum, Prodromus—s. Reconvalescentiæ, S. Annihilationis.

STAFF, (D.) *Staf*. An instrument of great importance in lithotomy,—being the director for the gorget or knife. It is made of steel; and the handle is flat and rough, in order that it may be more securely held. The groove is on its convex surface.

STAFF TREE, CLIMBING, *Celastrus scandens*.

STAFFA, Stapes.

STAG'S PIZZLE, see Cervus.

STAGE, *Sta'dium*, (F.) *Estage*, *Étage*. *Stade*. The period or degree of a disease; especially the period of an intermittent:—as the cold stage,—*sta'dium fri'goris*—hot stage, &c

STAGGERBUSH, *Andromeda mariana*

STAGGERWEED, *Delphinium*.

STA'GIUM. The sixth part of an ounce.

STAGMA, from *σταῶ*, 'I distil,' *Stalag'ma*. Any distilled liquid or liquor.

STAGMA, Gutta.

STAGNA'TION, *Stagna'tio*, from *stagnare*, *stagnatum*, 'to form a pond,' (*stagnum*, 'a pond.') Accumulation and retardation of fluids in any part of the body.

STAGNUM CHYLI, *Receptaculum chyli*.

STAHLIAN. One who follows the doctrine of Stahl.

STAHLIANISM. The doctrine of Stahl, which considered every vital action under the direction and presidency of the soul. See *Animist*.

STALAGMA, *Stagma*.

STALAGMITIS CAMBOGIOÏ'DES, *S. Cambogia*, from *σταλαγμος*, 'a drop'—the gum exuding by drops. Gamboge was long ascribed to this fancied Ceylonese plant. It is now known, that the name was given by Murray to specimens of two distinct trees of different genera. See *Cambogia*.

STALAGMOS, *Distillation*.

STALE, *Urine*.

STALTICUS, *Incarnans*.

STAMMERING, *Balbuties*—s. of the Fingers, Cramp, writers'.

STAMP, *Cachet*—s. Medicine, *Cachet*—s. Oculist, *Cachet*.

STAMPING MILL, see *Pulverization*.

STANG, see *Throbbing*.

STANGOS, *Tin*.

STANNI MURIAS, *Tin*, *muriate* of—s. *Oxydali murias*, *Tin*, *muriate* of—s. *Pulvis*, see *Tin*.

STANNIOLUM, *Tin foil*.

STANNUM, *Tin*—s. *Cinereum*, *Bismuth*—s. *Foliatum*, *Tin foil*—s. *Glaciale*, *Bismuth*—s. *Granulatum*, see *Tin*—s. *Indicum*, *Zincum*.

STAPÉDIEN, *Stapedius*.

STAPÉDIUS, *Stapes*, (F.) *Stapédien*, *Pyramidal-stapédien*. A muscle of the middle ear, which draws the stapes obliquely upwards.

STAPES, genitive *Stap'edis*, (*sto* and *pes*.) *Stapha*, *Staffa*, 'a stirrup.' (F.) *Étrier*. The innermost of the small bones of the ear, so called because it resembles a *stirrup*. It is situate horizontally in the cavity of the tympanum. Its head, which is directed outwards, is articulated with the os orbiculare. Its base faces inwards, and is applied to the fenestra ovalis, which it closes imperfectly, and to the membrane of which it is united. Its branches are one anterior, and the other posterior. They are curved; and the space between them is parabolic. Surgeons have given the name *Stapes* to a bandage used in bleeding from the foot; which forms a figure of 8 around the ankles.

STAPHA, *Stapes*.

STAPHIS, *Delphinium staphisagria*.

STAPHISAGRIA, *Delphinium staphisagria*.

STAPHISAIGRE, *Delphinium staphisagria*.

STAPHYLEMATOMA, from *σταφυλη*, 'the uvula,' and *hæmatoma*. A tumour formed by effusion of blood beneath the mucous membrane of the uvula.

STAPHYLAGRA, *Staphylepartes*.

STAPHYLE, *Uvula*, see *Vitis vinifera*.

STAPHYLEPAR'ES, from *σταφυλη*, 'the uvula,' and *εραϊσος*, 'I lift up.' *Staph'ylogra*. An instrument for laying hold of, and removing the uvula.—Paulus.

STAPHYLINI, see *Azygos muscle*—s. *Medii*, see *Azygos muscle*.

STAPHYLINO-PHARYNGEUS, *Palato-pharyngeus*.

STAPHYLINUS EXTERNUS, *Circumflexus*.

STAPHYLIS, *Nipple*.

STAPHYLITIS, *Uvulitis*.

STAPHYLUM, *Nipple*.

STAPHYLODIALYSIS, *Cionis*, *Staphylodema*.

STAPHYLŒDE'MA, from *σταφυλη*, 'the uvula,' and *dema*. *Angi'na uvula'ris*, *Dyspha'gia uvula'ris*, *Hypostaph'yli'tis*, *Hypostaph'yli'tis*, *Staphylo'dial'ysis*, *Prolapsus seu Propto'is* seu *Casus seu* (*Edema u'vulae*, *Himias*, *Himanto'ma*, *Himanto'is*, *Staphylo'n'cus*, *Staphylo'n'cia*, *Productio uvulae à pituità*, *U'vula decidua seu relaxa'ta*, *Cras'pedon*. Relaxation and tumefaction of the uvula, which, either from inflammation or infiltration, hangs down so as to occasion difficult deglutition, cough, &c. Scarification, astringent gargles, and, if necessary, excision, constitute the treatment.

STAPHYLO'MA, from *σταφυλη*, 'a grape.' *Uva'tio*, *Myoceph'alon*, *Myoceph'alum*, *Parop'sis*, *Staphylo'ma*, *Staphylo'is*, *Polygro'ma*. A name given to different tumours of the anterior surface of the globe of the eye.

STAPHYLOMA CONICUM, *Staphyloma* of the cornea—s. *Pellucidum conicum*, *Staphyloma* of the cornea.

STAPHYLOMA OF THE CORNEA, *Sugar-loaf Cornea* or *Conical Cornea*, *Conical hydrophthal'mia*, *Staphylo'ma con'icum* seu *pellu'idum con'icum*, *Conophthal'mia*, *Prolap'sis cor'neae*, *Ceraton'cus*, *Hypercerato'sis*, *Hypercer'asis*, is an unequal, embossed, bluish or whitish, round or conical tumour; of variable size; formed by the projection of the transparent cornea, which is sometimes attenuated and distended, at others thickened.

STAPHYLOMA OF THE IRIS is the disease described by Scarpa under the name *Procidencia Iridis*. It is hernia of the iris, through an accidental opening of the cornea. Also, *iridauexesis*.

STAPHYLOMA OF THE SCLEROT'IC, *Staphyloma sclerot'ice*, differs from the last, in the prominence existing on a part of the eye covered by the sclerotic, which is much thinner.

STAPHYLOMA VUEÆ, *Iridauexesis*.

STAPHYLONCIA, *Staphylodema*.

STAPHYLONCUS, *Staphylodema*.

STAPHYLOPLAS'TIC, *Staphyloplas'ticus*, from *σταφυλη*, 'the uvula,' and *πλασσω*, 'I form.' An epithet applied to the operation for replacing the soft palate, when it has been lost. *Staph'yloplasty*, *Pal'atoplasty*, a word of hybrid formation, is the operation for closing the palatine vault.

STAPHYLORAPHY, *Velosynth'esis*, *Staphylorrrha'phia*, *Kionorrrha'phia*, *Cionorrrha'phia*, *Uranor'rhap'hè*, *Uranorrrha'phia*, *Uraniscorrrha'phia*, *Uraniscor'rhap'hè*, from *σταφυλη*, 'the uvula,' and *ραφή*, 'a suture.' The operation for uniting a cleft palate. It consists in paring the edges of the cleft; passing ligatures through them, and bringing them together.

STAPHYLOSIS, *Staphyloma*.

STAPHYLOT'OMY, *Staphylotom'ia*, from *σταφυλη*, 'the uvula,' and *τομή*, 'incision.' Amputation of the uvula.

STAR APPLE, **BROAD-LEAVED**, *Chrysophyllum cainito*—s. *Blazing*, *Aletris farinosa*—s. *Bloom*, *Spigelia Marilandica*—s. *Grass*, *Aletris farinosa*, *Hyopoxys erecta*.

STAR, *Knap-weed*, *Centaurea calcitrapa*.

STAR OF THE NORTH, *Geum urbanum*.

STAR STROKE, *Coup-de-soleil*—s. *Wort*, *Aletris farinosa*—s. *Wort*, *golden*, *Bubonium*—s. *Wort*, *water*, *Callitriche verna*.

STARBECK SPA. A chalybeate spring at Harrogate.

STARCH, *Sax. r'tape*, 'rigid, stiff,' *Amylum*.

STARCH, *ANIMAL*, *Corpora amylacea*—s. *Blue*, *Smalt*—s. *Canna*, *Tous-les-mois*.

STARCH CORPUSCLES, *Corpora amylacea*.

STARCH, *IODIDE OF*, *Iod'idium* seu *Iodure'tum Am'gli*, *Am'ylum ioda'tum*, (F.) *Iodure d'Amidon*; prepared by triturating iodine and starch, is said to have the eutrophic effects of iodine; but, by some, is esteemed inert. Dose, four to six grains.

STARCH, POTATO, see *Solanum tuberosum*.

STARCH SUGAR, Glucose.

STARCH, TACCA, Arrowroot, Tahiti.

STARKEY'S PILLS, see *Pilulae ex helleboro et myrrha*.

STASE, Stasis.

STASIS, *stasis*, from *stao*, 'I stop,' (F.) *Stase*. This word has nearly the same signification as stagnation. Stagnation, however, sometimes includes a morbid condition of the fluids, — Stasis never.

STASIS, Station.

STATE MEDICINE, Medicine, state.

STATIC, *Statical*, *Staticus*, (F.) *Statique*.

Same etymon. An epithet applied to the physical phenomena presented by organized bodies in contradistinction to the organic or vital.

STATICA SEPTI NARIUM, *Appendix cutanea septi narium*. The cutaneous appendage to the septum narium.

STATICAL, Static.

STATICE CAROLINIANA, of the United States, (*στατικός*, 'restraining,') *American* or *Seaside Thrift*, *Marsh Rosemary*, *Marsh Root*, *Ink-root*, *Sea Lavender*, *Ord.* *Plumbaginaceae*, is considered to be a mere variety of the *Statice limonium*. The root is the *Statice* — (Ph. U. S.).

STATICE LIMONIUM, *S. Maritimum*, *Behen rādrum*, *Limonium*, *L. majus seu vulgare*, *Behen*, *Behnen Ackmar*, *Sea Lavender*, *Red Behen*, *Marsh Rosemary*, *Lavender Thrift*, (F.) *Behen rouge*, *Romarin des marais*, *Lavande triste ou de mer*. The roots have been esteemed astringent, tonic, and eminently antiseptic.

STATICE MARITIMUM, *S. limonium* — *s. Vulgare*, *S. limonium*.

STATION, *Sta'tio*, *Stasis*, *Status*, from *stare*, *statum*, 'to stand.' The act of standing.

STATIONARY, *Stationarius*, (F.) *Stationnaire*. Same etymon. A name given by Sydenham and Stoll to certain diseases, which depend upon a particular state of the atmosphere; and which prevail in a district for a certain number of years, and then give way to others.

STATISTICS, MEDICAL, *Vital Statistics*, *Biostatistics*, (*Status*, 'a state,') (F.) *Statistique médicale*. That part of medicine which relates to details — *medico-statistical* — connected with the mortality, salubrity, &c., of different countries and situations.

It has, also, been used in a wider sense to signify the same as *Numerical method*, (F.) *Méthode numérique*; which is applied to the systemization of medical facts, or to the use of numbers in the investigation of disease.

STATISTICS, VITAL, Statistics, medical.

STATURE, *Statu'ra*, *Meg'ethos*, *Magnitudo corporis*, from *sto*, *statum*, 'to stand.' The height of the human figure. The following table exhibits the results of several admeasurements, with the heights of some of the monkey species, on the authority of Mr. Lawrence.

	STATURE.	
	Feet.	In.
An Englishman	6	4½
	5	00
	5	4
An Englishwoman	5	00
	5	8
	5	5
European male skeleton	5	10½
Do.	5	5½
A Negro at the Lunatic Hospital, Liver- pool	5	8
Another from Virginia	5	00
Another from the Gold Coast		
Another		

	STATURE.	
	Feet.	In.
Negro skeleton	4	11
Another	5	7½
A Lascar	5	4
Venus de Medici	5	00
Tyson's Chimpanzé (<i>Simia Troglodytes</i>) ..	2	2
Mr. Abel's Orang Utang	2	7
Camper's Do.	less than	30
Mr. White's Monkey	2	2

STATUS, *Acme* — *s. Cruditatis*, *Crudity* — *a. Erethiticus*, *Nervous diathesis* — *s. Gastricus*, *Embaras gastricus* — *s. Nervosus*, *Nervous diathesis* — *s. Parturiendi*, *Parturition* — *s. Sthenicus*, *Sthenia* — *s. Verminosus*, *Helminthiasis*.

STAVESACRE, *Delphinium staphisagria*.

STAXIS, *staxis*, *Apostax'is*. A slight epistaxis or nasal hemorrhage. — *Hippocrates*.

STEAM-DOCTOR. A term applied to one who treats all or most diseases by steam.

STEAR, see *Fat*, *Sevum*.

STEARATE, *Emplastrum*.

STEARIN, see *Fat*.

STEARINUM CETACEUM, *Cetaceum*.

STEAROLÉ, *Pomatum* — *a. Solide*, *Emplastrum*.

STEAROSIS, *Steatosis*.

STEARRHŒA, *Stearorrhœ'a*, *Seborrhœ'a*, *Seborrhag'ia*, *Smegmatorrhœ'a*, *Smegmorrhœ'a*, *Fluxus sebaceus*, *Sebaceous Flux*, from *stear*, 'suet,' and *rho*, 'I flow.' Augmentation of secretion from the sebaceous follicles of the skin.

STEARRHŒA NIGRICANS. A term used by Dr. Neligan, of Dublin, for a black discoloration of the face, observed in young females affected with derangement of the catamenia, and which, in his opinion, is owing to a dark secretion from the cutaneous sebaceous glands.

STEATITES, *Polysarcia*.

STEATOCELE, from *stear*, *e caros*, 'suet,' and *κηλη*, 'a tumour.' A tumour formed by a collection of steatomatous matter in the scrotum.

STEATO'MA, *Tumor lar'vus*, from *stear*, *steatos*, 'suet.' *Sena'tio*, *Emphy'ma encystis steatomata*. An encysted tumour, whose contents are similar to fat.

STEATOPYGA, from *stear*, *steatos*, 'fat,' and *πυγ*, 'buttocks.' *Buttock-hump*. Fat buttocks; such as exist in the Hottentot female.

STEATORRHŒA, *Stearrhœa*.

STEATO'SIS, *Stearo'sis*. Fatty degeneration of a part, or unusual deposition and accumulation of fat in its adipose tissue.

STEATO'SIS CORDIS, *Degenera'tio adipo'sa cordis*, *Cardiethmolipo'sis* and *Cardiomyolipo'sis* (*Piorry*), *Fatty heart*. Unusual deposition of fat on the heart: — by some presumed to be occasionally a true adipous degeneration of the substance of the heart.

STEATOSIS HEPATICA, *Adiposis hepatica*.

STEATOZOON FOLLICULORUM, *Acarus Folliculorum*.

STEEL, *Chalybs*.

STEEP, *Rennet*.

STEEPLE-BUSH, *Spiræa tomentosa*.

STEER'S OPODELDOCH, see *Opodeldoch*, *Steer's*.

STEGMONTH, see *Parturient*.

STEGANOP'ODES, *στεγανόδες*, *Planip'edes*. They who have flat feet, or are web-footed, from *στεγανος*, 'covered,' and *πους*, 'foot.'

STEGNO'SIS, *Steno'sis*, *Constipa'tio*, *Stipa'tio*, *Condensa'tio*, *Obstruc'tio*, from *στεγνω*, 'I constrict.' Constriction or narrowness of the pores and vessels. Stricture. Constipation. Suppression of the evacuations. See *Arctatio*.

STEGNOTICA, *Astringents*.

STEIROSIS, Sterilitas.

STELENGIS, Stridor dentium.

STELLA, *Fas'cia stella'ta*, (F.) *Étoile*. A bandage, improperly compared to a star, because it is crossed like the letter X. This bandage is applied upon one or both shoulders. In the first case, it is called *simple*; in the second, *double*. It was employed in fractures of the sternum, clavicle, and scapula; in luxations of the humerus, &c.

STELLA TERRÆ, Plantago coronopus.

STELLARIA MEDIA, Alsine media—s. Scilla, Scilla.

STELLATA RUPINA, Centaurea calcitrapa.

STELL'ULÆ VERHEYEN'II, (*stella*, 'a star,' *stel'lula*, 'a little star,') *Substan'tia stel'lata serpenti'na seu radiat'a*, *Ell'gch'nia*. A stelliform plexus of veins at the surface of the kidney, which gives origin to the interlobular veins.

STELOCHITIS, Osteocolla.

STENAGMUS, Suspirium.

STENAXIS, Suspirium.

STENIA, Sthenia.

STENICUS, Active.

STENOBREGMATE, see Cranium.

STENOCARDIA, Angina pectoris, Cardiostenosis.

STENOCHO'RIA, *Stenochori'asis*, from *στενος*, 'narrow,' and *χωρα*, 'space.' Angustia. Employed by some for contraction of the vagina.

STENOCHORIA INTESTINI RECTI, Stricture of the rectum—s. *Œsophagi*. Dysphagia constricta.

STENOMYCTE'RIA, from *στενος*, 'narrow,' and *μυκτρον*, 'nose,' Contraction of the nasal passages.

STENOSIS, Stenosis.

STENOSTEGNOSIS, Stenostenosis.

STENOSTENO'SIS, *Stenostegno'sis*, from *Steno*, and *στενωσις*, 'contraction.' *Obstruc'tio ductus Stenonia'ni*. Narrowness or contraction of the duct of Steno.

STENOSTOM'IA, from *στενος*, 'narrow,' and *στρομα*, 'mouth.' Contraction of the mouth.

STENOTHORAC'ES, from *στενος*, 'narrow,' and *σθώρας*, *σθώρακος*, 'the chest.' They who have narrow chests.

STENTOROPHO'NUS, from *Στεντορ*, *Stentor*, a Greek of powerful voice in the army before Troy; and *φωνη*, 'voice.' One who has a strong voice.

STEPHANE, Crown.

STEPHENS, MRS., REMEDY FOR THE STONE. This consisted of lime, prepared by calcining the shells of eggs and snails, made into pills with soap. A decoction was, also, administered, consisting of chamomile, fennel, parsley, and burdock, with a portion of Alicant soap. Its virtues were dependent upon the lime and the tonic properties of the decoction. Hales, Hartley, Kirkpatrick, Lobb, and others, wrote in favour of this nostrum; and it was ultimately bought by the British parliament for £5000—about \$25,000.

STEPHENSIA ELONGATA, Matico.

STEPMOTHER'S BLESSING, Hangnail.

STERA, Uterus.

STERCORACEOUS, *Stercora'rius*, *Diachor'ricus*. *Ster'coral*, from *stercus*, *stercoris*, 'dung.' *Fœcal*, *Fæcal*, (F.) *Stercoraire*, *Stercoral*.

The French use the term *Fistules stercoraires* for those which are kept up by the constant passage of fecal or *stercora'ceous* matter. The name *Stercoraires* is sometimes given, in derision, to those physicians who are fond of purging their patients: à *posteriori* has been similarly used.

STERCORAIRE, Stercoraceus.

STERCUS, Excrement, Fimus—s. Anseris, Fimus anseris—s. Caninum album, Album Græcum—s. Diaboli, Asafœtida—s. Equi non castrati, Fimus.

STERCUS HUMA'NUM, Homerda. Human excrement.

STERCUS LACERTÆ, Crocodilia.

STEREOCYST'IS, from *στερεος*, 'hard,' and *κυστις*, 'cyst.' A hard encysted tumour.

STEREOS, Solid.

STEREOSCOPE, *Stereoscop'ium*, from *στερεος*, 'solid,' and *σκοπεω*, 'I view.' A physical instrument by means of which plane pictures of an object are made on each retina, and so blended as to appear as one in relief. Also, an instrument for detecting a calculus in the bladder, and foreign bodies in the soft parts. It resembles a common catheter, having an expanded free extremity.

STEREOT'ICA, from *στερεος*, 'hard.' Fortuitous lesions or deformities affecting the hard parts. The 2d order, class *Tychica* of Good.

STER'ILE, *Ster'ilis*, *A'gonos*, *Aph'orus*, *Aph'oricus*, *Ablast'ea*, *Infecun'dus*, *Barren*, (F.) *Stérile*. Same etymon as Sterilitas. Not producing fruit; infecund.

STERILIS, Sterile.

STERILITAS, *Agon'ia*, *Steiro'sis*, *Stiro'sis*, *Agenne'sis*, *Agenne'sis*, *Infecun'ditas*, *Steril'ity*, *Acy'e'sis*, *Bar'renness*, *Infecun'dity*, *Apho'ria*, *Atec'nia*, *Ato'cia*, from *στερεος*, or *στερος*, 'hard, barren.' Quality of that which is barren. Sterility may be *absolute* or *relative*. The former depends upon some vicious conformation of the genital organs, apparent or concealed, and occasionally admits of cure. Sterility is *relative* when a female does not conceive with one individual, but does with another. Sterility in females is often dependent upon a condition of the uterus characterized by membranous menstruation.

STERNAD, see Sternal.

STERNAL, *Sterna'tis*, from *sternum*, 'the breast bone.' Relating or appertaining to the sternum or breast bone.

STERNAL ASPECT. The aspect of the body where the sternum is situate.—Barclay. *Sternad* is used by the same writer adverbially to signify towards the sternal aspect.

STERNALGIA, Angina pectoris.

STERNO, in composition, belonging, or relating to, the sternum or breast bone.

STERNO-CLAVIC'ULAR, *Sterno-clavicula'ris*. That which relates to the sternum and clavicle.

STERNO-CLAVIC'ULAR ARTICULA'TION is the union of the inner extremity of the clavicle with a round facetæ at each side of the superior extremity of the sternum. In this articulation there are, besides the intercostal cartilages, two radiated sterno-clavicular, (F.) *Sterno-claviculaires rayonnées*; one anterior, and one posterior—an *inter-clavicular* ligament extending between the two clavicles, and two synovial capsules, separated by an *inter-articular* fibro-cartilage. The costo-clavicular ligament may also be considered to belong to this articulation.

STERNO-CLAVIO-MASTOÏDIEN, Sterno-cleido-mastoideus—s. Cleido-hyoideus, Sterno-hyoideus.

STERNO-CLEIDO BRACHIALIS, Pectoralis major.

STERNO-CLEIDO-MASTOÏDEUS, *Sterno-mastoïdeus*, *Cleido-mastoïdeus*, *Mastoïdeus*, *M. anterior*, *Nuta'tor cap'itis*; (F.) *Sterno-clavio-mastoïdien*, *Sterno-mastoïdien*; from *στερεος*, 'the sternum,' *κλες*, 'the clavicle,' and *μαστος*, 'the mastoid process.' A muscle situate at the anterior and lateral part of the neck. It is long and flattened: simple, above; and divided into two portions below; where it is inserted into the anterior and superior part of the sternum, and the inner quarter of the clavicle. Above, it terminates at the mastoid process, which it embraces, at the neighbouring part of the temporal bone, and at the outer third of the upper curved line

of the occipital bone. It carries the head forward; inclines, and rotates it to one side. When both contract, they bend the head directly forwards.

STERNO-COSTAL, *Triangularis sterni* — *s. Costo-clavio-huméral*, *Pectoralis major* — *s. Huméral*, *Pectoralis major*.

STERNO-HYOIDEUS, *Sterno-cleido-hyoideus*, *Sterno-hyoïdes*, *Hyoïdes primus*; from *στερον*, 'the sternum,' and *υοιδης*, 'the os hyoides.' A muscle situate at the anterior part of the neck; inserted, above, at the inferior edge of the body of the hyoid bone; below, into the upper part of the posterior surface of the sternum; — at the posterior part of the sterno-clavicular articulation, and sometimes into the cartilage of the first rib. It depresses the os hyoides.

STERNO-MASTOÏDIEN, *Sterno-cleido-mastoïdeus* — *s. Pubien*, *Rectus abdominis*.

STERNO-THYROIDÆUS, *Sterno-thyroïdes*, *Bronchius*. A muscle, situate at the anterior part of the neck. It is attached, above, to the oblique line seen on the outer surface of the thyroid cartilage; and terminates, below, at the upper part of the posterior surface of the sternum, and the cartilage of the second rib. When it contracts, it depresses the thyroid cartilage and os hyoides by the medium of the thyro-hyoid membrane.

STERNODYNIA, see *Stomatodynia* — *s. Syncopalis*, *Angina pectoris* — *s. Syncoptica et Palpitans*, *Angina pectoris*.

STERNO-OMPHALODYMIA, see *Somatodynia*.

STERNOPA'GIA, from *στερον*, 'the breast bone,' and *παγω*, 'I fasten.' A monstrous union of costæ, in which they are attached at the sternum.

STERNUM, *Sternon*, from *στερρος*, 'solid,' *Pectus*, *Os asser seu pectoris seu xiphoides seu gladioli seu ensiforme, Scutum cordis seu pectoris*, *Breast Bone*, *Hæmal Spine* of Owen. An azygous symmetrical bone, situate at the forefront of the chest. It is flat; broad above; narrower in the middle; and terminates, below, by a pointed cartilage — the xiphoid. It has an anterior or cutaneous surface, a posterior or mediastinal — a superior or clavicular extremity, and an inferior or abdominal. It is articulated with the clavicles and the seven upper ribs on each side, by means of their cartilages. According to Bécларd, it is formed of six principal bones, which he names, from their position, 1. *Primi-sternal* or *clavi-sternal*: 2. *Duo-sternal*: 3. *Tri-sternal*: 4. *Quarti-sternal*: 5. *Quinti-sternal*: 6. *Ultimi-sternal* or *ensi-sternal*.

STERNUMENTUM, *Sneezing*.

STERNUTAMENTORIA, *Achillea ptarmica*.

STERNUTAMENTUM, *Sneezing*.

STERNUTATIO, *Sneezing*.

STERNUTATOIRE, *Sternutatory*.

STERNUTATORY, *Sternutatorius*, from *sternutare*, *sternutatum*, 'to sneeze.' *Sternutamentum*, *Errhi'num*, *Ptar'micum*, *Anarrhi'num*, (F.) *Sternutatoire*. A substance which provokes sneezing; as tobacco, &c. — See *Errhine*.

STERTEUR, *Stertor*.

STERTOR, *Ronchus* or *Rhonchus*, *Renchus*, *Rhencus*, (F.) *Sterteur*, *Ronflement*, from *stertere*, 'to snore.' The deep snoring which accompanies inspiration in some diseases, particularly in apoplexy. Such respiration is said to be *ster'torous*, (F.) *Sterto'reux*; *Respira'tio stertuo'sa seu stertens seu stertori'ca*.

STERTOROUS, see *Stertor*.

STETHÆMIA, *Hyperæmia Pulmo'num seu Pectoris*, *Congestio Pulmonum seu Pectoris*, (F.) *Hyperémie ou Congestion des Poumons*; from

σθηος, 'the chest,' and *αἷμα*, 'blood.' *Hyperæmia* of the lungs. Congestion or accumulation of blood in the pulmonary vessels.

STETHOCATHARSIS, *Expectoration*.

STETHOCHYSIS, *Hydrothorax*.

STETHODESMIS, *Corset*.

STETHODESMIUM, *Corset*.

STETHODESMUS, *Corset*.

STETHOMETER, from *σθηος*, 'the chest,' and *μετρον*, 'measure.' An instrument, proposed by Dr. Richard Quain, for measuring the extent of movement of the walls of the chest, as a means of diagnosis in thoracic diseases. It consists of a flat case, not unlike a watch-case, having on its upper surface a graduated dial and index, which is acted on by a simple mechanism. A silk cord, long enough to surround one-half the chest, passes through an aperture in the case. The dial is graduated into fifty equal parts, each of which represents the 1-200th of an inch. In using the instrument, the dial is placed firmly against the spine, and the point of the tape over the sternum, or at any other spot according to the part to be examined. The extension of the string, as indicated by the dial, shows the amount of elasticity of the parietes of the thorax.

Under the name *Thoracometer*, Dr. Sibson has described an instrument of his invention, which is employed with a similar view as the stethometer of Dr. Quain. — *Med. Chir. Transact.*, vol. xxxi.

STETHOPARALYSIS, *Paralysis nervorum thoracis, phrenicorum*, &c.; from *σθηος*, 'the chest,' and *παρὰλυσις*, 'paralysis.' Palsy of the muscles of the chest.

STETHORRHEUMA, *Pleurodynia*.

STETHOSCOPE, *Stethoscopium*, *Stethoscopum*, *Echoscopium*, *Pnephonimeter*, *Thoracoscopium*, from *σθηος*, 'the chest,' and *σκοπεω*, 'I examine.' An instrument invented by Laënnec, of Paris, for exploring the chest. The stethoscope, sometimes called *Pectorilique*, is a cylinder of wood, from four inches to a foot long; pierced through and through by a longitudinal canal about a quarter of an inch in diameter. The longer are generally composed of two portions, that fit together by means of a screw, one of which is hollowed at the extremity, in the shape of a funnel. These two portions being screwed to each other, the physician lays hold of the instrument, as he would of a pen, puts the funnel-shaped extremity on the chest of the patient, and applies his ear to the other. To explore the pulsations of the heart, the funnel is plugged up by a piece of the same kind of wood accurately adapted to it, and pierced by a canal of the same width as that in the body of the instrument. Stethoscopes are sometimes flexible, like the flexible ear-trumpet.

This mode of examining affections of the chest, *Stethoscopy*, *Stethoscopia*, &c. is what Laënnec terms *Auscultation médiate*, *Mediate Auscultation*.

STETHOSCOPIA, see *Auscultation*, and *Stethoscope*.

STETHOSCOPTIC, *Stethoscopicus*, (F.) *Stéthoscopique*. Same etymon as *Stethoscope*. Furnished by stethoscopy or auscultation, — as a 'stethoscopic sign.'

STETHOSCOPY, see *Stethoscope*.

STETHUS, *Thorax*.

STEW, *Stove*.

STHENIA, *Steni'a*, *Diathësis sthenica*, *Status sthenicus*, *Via activa*, from *σθενος*, 'strength,' 'power.' Excess of strength; excitement of the organic actions. A word used, especially by the Brunonians, as well as *Asthenia*, — its antitheton. *Sthenic* or *dynamic diseases* are such as depend upon excessive excitement.

STHENICUS, *Active*.

STHENOPYRA, *Synocha*.

STIBI, Antimonium.

STIBIALIS, Antimonial.

STIBIALISM, *Stibialis'mus*, from *stibium*, 'antimony.' The aggregate phenomena produced by antimonials, when given in large quantities:—antimonial intoxication or poisoning.

STIBIATUS TARTRIS LIXIVIÆ, Antimonium tartarizatum.

STIBI BUTYRUM, Antimonium muriatum—s. Deuto-murias sublimatum, Antimonium muriatum—s. et Potassæ deuto-tartras, Antimonium tartarizatum—s. Hydrosulphuretum rubrum, Antimonii sulphuretum rubrum—s. Oxidum acidumuriatico oxygenato paratum, Algaroth—s. Oxidum album mediante nitro confectum, Antimonium diaphoreticum—s. Oxidum præcipitando paratum, Algaroth—s. Oxodes album, Antimonium diaphoreticum—s. Oxodes sulphuretum, Antimonii sulphuretum præcipitatum—s. Oxydi sulphurati hydro-sulphuretum luteum, Antimonii sulphuretum præcipitatum—s. Oxydulati sulphuretum, Antimonii sulphuretum præcipitatum—s. Oxydulum vitreum, Antimonii vitrum—s. Oxidum semivitreum, Antimonii vitrum—s. Oxidum sulphuratum, Oxidum stibii sulphuratum—s. Subhydrosulphas, Antimonii sulphuretum rubrum—s. Suburias præcipitando paratum, Algaroth—s. Sulphurati hydrosulphuretum rubrum, Antimonii sulphuretum rubrum—s. Sulphuretum nigrum, Antimonium—s. Vitrum, Antimonii vitrum.

STIBO-DERMIC, see Unguentum antimonii.

STIBIOSUM HYDROSULPHURETUM CUM SULPHURE, Antimonii sulphuretum præcipitatum—s. Oxidum, Antimonium diaphoreticum.

STIBIUM, Antimonium.

STICHAS ARABICA, *Lavendula stœchas*.

STICKWORT, Agrimony.

STICTA PULMONACEA, Lichen pulmonarius.

STIFF JOINT, MUSCULAR, Contractura.

STIFF JOINTS, Milk sickness.

STIFFNECK, Torticollis.

STIGMA, Cicatricula.

STIGMATA, see Nævus. The breathing pores of insects are so called.

STIGMATA OVARIO'RUM, *Cicatrices Ovariorum*. The cicatrices left in the ovary after the escape of ova.

STILBOMA, Cosmetic.

STILL, Alembic.

STILLA, Gutta.

STILLATIO SANGUINIS E NARIBUS, Epistaxis.

STILLBORN, (F.) *Mort-né*. Born still or lifeless,—dead born.

STILLICIDIUM, *Substillum*, from *stilla*, 'a drop.' This term is often used synonymously with *Strangury*. It means the flow of any fluid—particularly the urine—drop by drop. Also a kind of embrocation, *Stillicidium supra partem*, effected by dropping a liquid upon a part.

STILLIDIUM LACRYMARUM, Epiphora—s. Narium. Coryza—s. Sanguinis e naribus, Epistaxis—s. Supra partem, see Stillidium.

STILLINGIA, *S. sylvatica*, Queen's Root, Queen's delight, Yaw root, *Mar'cory*, Cockup hat. An indigenous plant, which grows in pine barrens, from Virginia to Florida. *Sex. Syst.* Monœcia Monadelphica. *Nat. Ord.* Euphorbiaceæ. The root—*Stillingia*, Ph. U. S.—is said to be used in syphilis, obstinate cutaneous affections, &c., like sarsaparilla.

STIMATOSIS, *Stymatosis*, *Hæmorrhag'ia Penis*, *Stymatorrhag'ia*, *Phallorrhag'ia*, *Caulorrhag'ia*, from *στυμα*, 'priapism,' itself from *στυμι*, 'to erect.' Hemorrhage from the male organ, accompanied or not with priapism.

STIMMI, Antimonium.

STIMULANT, *Stimulans*, *Excitans*, *Inci'tans*, *Incentivum*, *Irritans*, *Exci'tant*, *Incit'ant*, *Hyperæthenic*, *Pyretogenetic*, from *stimulare*, 'to goad.' A medicine which has the power of exciting the organic action of the different systems of the economy. Some stimulants are *diffusible*,—that is, have a prompt but temporary action; others are *permanent* or *persistent*. The action of stimulants is called *Stimulation*, *Stimula'tio*, *Incitatio*.

The chief stimulants are the following: *Æther Sulphuricus*, *Ammonia*, *Asafœtida*, *Brucia*, *Camphora*, *Capsicum*, *Castoreum*, *Cinnamomum*, *Copaiba*, *Cubeba*, *Guaiacum*, *Hydrargyri Præparata*, *Iodinium*, *Nux Vomica*, *Olea Essentialia*, *Opium* (in small dose), *Piper*, *Sinapis*, *Strychnia*, *Zingiber*,—*Alcohol*, *Caloric*, *Electricity*, *Mental Emotions*, (the *Impetuous*.)

STIMULUS, 'a goad.' *Incitamentum*, *Irritamentum*, *Irritantum*, *Potentia Irritans*. Any thing which excites the animal economy.

STINE, Hordeolum.

STENK POKE, *Dracontium fœtidum*—s. Weed, *Chenopodium anthelminticum*, *Datura stramonium*, *Polanisia graveolens*.

STIPATIO, Stegnosis.

STIPATIO TELÆ CELLULOSÆ INFANT'UM. Induration of the cellular tissue of children.

STIROSIS, Sterilitas.

STITCH, same root as *stick*, Sax. *ritician*, (G.) *stecken*, (Sc.) *Stonnd*, (Prov.) *Dag*, (F.) *Point*, (L.) *Punctum*; (from *pungere*, *punctum*, 'to prick or sting.') An acute lancing pain, like that produced by the puncture of a sharp needle.

STITCH, COMMON, Suture, common—s. Royal, see Suture—s. in the Side, *Pleurodynia*.

STIZOLOBIUM, *Dolichos pruriens*.

STLENGIS, *Stridor dentium*.

STOBÆA RUBRICAULIS. A South African plant, *Nat. Ord.* Compositæ. A tincture of the bruised roots is diuretic, and of great service in gravel.

STOCKING, LACED, (F.) *Bas lissé*. A bandage used by surgeons, which consists of a stocking made of firm cloth or dog-skin, admitting of being tightly laced anteriorly. It is employed in varices and ulcers of the legs.

STECHAS ARABICA, *Lavendula stœchas*.

STECHEIOLOGY, *Stœchiolog'ia*, (F.) *Stœchiologie*, from *σταιχιον*, 'element,' and *loyos*, 'a discourse.' A treatise, or the theory, of elements; hence of the elementary parts of organized substances.

STOICHEION, Element.

STOMA, Mouth.

STOMAC'ACĒ, *Stomatoc'acē*, *Stomol'gia*, *Labrisul'cium*, *Cheiloc'acē*, *Uloc'acē*, *Cheilomal'cia*, *Stomatoma'ia*, *Stomatonecro'sis*, *Cancer Oris*, *Canker*, from *στωμα*, 'the mouth,' and *kakos*, 'evil.' Fœtor of the mouth with a bloody discharge from the gums. (F.) *Cancer aqutique*; see *Cancer Aquaticus*. Also, *Scurvy*.

STOMACACE, *Porphyræ nautica*—s. *Gangrænium*, *Cancer aquaticus*—s. *Universale*, *Purpura hæmorrhagica*.

STOMACAL, Stomachal.

STOMACALGIA, Cardialgia.

STOMACH, *Stom'achus*, *Ventric'ulus*, *Nedys*, *Anocœ'lia*, *Gaster*, *Gluttu'patens*, *Maw*, (of animals), (Sc.) *Kyte*, *Pechan*, (Prov.) *Warison*, (F.) *Estomac*, from *στωμα*, 'a mouth.' One of the principal organs of digestion. It is a musculo-membranous reservoir; continuous, on the one side, with the œsophagus; on the other, with the duodenum. It is situate beneath the diaphragm, between the liver and the spleen; and occupies the epigastrium and a part of the left hypochondrium. In it the food is converted into chyme.

When viewed externally, the stomach has. 1. An *anterior face*, which looks a little upwards. 2. An *inferior face*, directed downwards. 3. An *inferior or colic margin*, which is convex and extensive, and is called the *greater curvature*, (F.) *Grand courbure*. It gives origin to the omentum majus. 4. A *superior or diaphragmatic margin*, which is shorter, concave, and is called the *lesser curvature*, (F.) *Petit courbure*. The lesser omentum is attached to this. 5. A *left or œsophageal orifice*, called, also, the *cardia*, *Os ventriculi* or *upper orifice*. 6. A *right or intestinal, or inferior orifice*, called the *pylorus*. 7. A considerable dilatation, situate to the left of the cardia and greater curvature—the *great tuberosity* or *great cul-de-sac* or *fundus* of the stomach; and, 8. A less extensive dilatation, situate to the right of the greater curvature,—the *lesser tuberosity* or *lesser cul-de-sac*, *Antrum Pylo'ri*. The inner surface of the stomach is of a reddish-white colour, and has a marbled appearance. It is constantly covered by thick mucus, and is lined by a mucous membrane, which presents numerous wrinkles. The parietes of the stomach consist of three membranes in superposition. The outermost is serous, and is an extension of the peritoneum. The middle coat is muscular,—some of its fibres running longitudinally; others, transversely, and others obliquely. The innermost membrane is of a mucous nature, *Crusta villosa ventriculi*, *Gastro-mycod'eria*, but not exactly a continuation of the membrane that lines the œsophagus. The mucous and muscular membranes form, at the pylorus, a valve, called the *Pyloric valve*. These three coats are united by a dense, close, areolar membrane; and, between the mucous and muscular coats, along the two curvatures especially, is a quantity of muciparous glands, called *Glands of Brunner*. The arteries of the stomach are very numerous, and proceed from the coronaria ventriculi, the pyloric, splenic, and right and left gastro-epiploic. The veins have the same name, and pursue the same course as the arteries. They pour their blood into the trunk of the vena porta. Its lymphatic vessels are very numerous, and pass into ganglia, situate along the two curvatures. The nerves of the stomach proceed from the pneumo-gastric, and three divisions of the celiac plexus.

STOMACH CELL, see Favulus—s. Disease, Lismosis.

STOMACH, FIRST, Inguivies.

STOMACH PUMP, *Gastren'chyta*, *Antli'a gastrica*, (F.) *Pompe pour l'estomac*. A useful instrument for conveying fluids to the stomach, when deglutition is impracticable, and for pumping out the contents of that organ. It consists of a forcing syringe, to the bottom and nozzle of which flexible tubes can be attached. When the object is to force fluids into the stomach, the stomach-tube must be attached to the nozzle, and passed down the throat,—the tube connected with the bottom of the syringe being placed in the fluid to be injected. To empty the stomach, the stomach-tube must be attached to the bottom of the instrument. This instrument has been of great service where poisons have been taken. The antidote, when any exists, and is at hand, should be administered immediately prior to its use.

The stomach pump was first employed in America by Dr. Physick, in the year 1812.

STOMACH, SECOND, Proventriculus—s. Sick, Milk sickness—s. Third, Ventriculus callosus.

STOMACH TUBE, see Tube, œsophageal.

STOM'ACHAL, *Stomach'ic*, *Stomach'ical*, *Cardi'acus*, *Stomach'icus*, *Stomacha'lis*, *Viscera'lis*, (F.) *Stomacal*, *Stomachique*. That which belongs to the stomach; that which is good for the sto-

mach; which strengthens the stomach. A medicine that gives tone to the stomach.

STOMACHALGIA, Cardialgia.

STOMACHI RESOLUTIO, Dyspepsia.

STOMACHIC, Stomachal.

STOMACHICAL, Stomachal.

STOMACHIQUE, Stomachal.

STOM'ACHUS, from *στομα*, 'mouth.' This word has been used in several different acceptations; for example, for the œsophagus, for the cardiac orifice of the stomach, (see Cardia,) and for the stomach itself. Hippocrates calls the neck of the bladder *στομαχος κωτος*; and the neck of the uterus *της μήτρας στομαχος*.

STOMAL'GIA, *Stomatal'gia*, *Stomatodyn'ia*, from *στομα*, 'mouth,' and *αλγος*, 'pain.' Pain in the mouth.

STOMAPYRA, Aphthæ—s. Aphtha, Aphthæ.

STOMATALGIA, Stomalgia.

STOMATEL'CIA, *Stomateleo'etis*; from *στομα*, *στοματος*, 'the mouth,' and *ελκος*, 'an ulcer.' Ulceration of the mouth.

STOMAT'IA, *Stomatopathi'a*; from *στομα*, *στοματος*, 'the mouth,' Disease of the mouth.

STOMAT'IC, *Stomat'icus*, from *στομα*, *στοματος*, 'mouth.' A medicine used in diseases, &c., of the mouth. Dentifrices, masticatories, &c., are stomatics.

STOMATITE APHTHEUSE, Stomatitis, aphthous—s. Charbonneuse, Cancer aquaticus—s. Couenneuse; Stomatitis, pseudo-membranous—s. Crèmeuse pultrée, Aphthæ—s. Folliculeuse, Stomatitis, aphthous—s. Gangréneuse, Cancer aquaticus—s. Mercurielle, see Salivation, mercurial—s. Ulcéreuse, Stomatitis, aphthous.

STOMATIT'IS, from *στομα*, *στοματος*, 'the mouth,' and *itis*, denoting inflammation. (F.) *Inflammation de la Bouche*; Inflammation of the mouth.

STOMATITIS, APHTHOUS, *S. follic'ular* or *ulcéreus*, *Em'phlysis Aphtha*, *A. Adulto'rum*, (F.) *Stomatite*, *S. aphtheuse* ou *folliculeuse* ou *ulcéreuse*. An inflammation of the follicles of the mouth, constituting the aphthæ of the adult, which is generally accompanied by cephalic, gastric, and general disturbance. It may be either discrete or confluent, and requires general and local treatment, adapted to the case.

STOMATITIS, ERYTHEM'ATOUS. Simple inflammation of the mouth.

STOMATITIS, EXSUDATIVA, Aphthæ—s. Follicular, S. aphthous—s. Gangrenous, Cancer aquaticus—s. Materna, Stomatitis of nursing women.

STOMATITIS, MERCU'RIAL, *S. mercuria'lis*, *Hydrargyrostom'atis*, *Hydrargyrostomatit'is*. Inflammation of the mouth induced by mercury.

STOMATITIS OF NURSING-WOMEN, *S. Nutri'cum* seu *Mater'na*, appears to be a variety of S. aphthous.

STOMATITIS NUTRICUM, Stomatitis of nursing-women.

STOMATITIS, PSEUDOMEMBRANOUS, (F.) *Stomatite couenneuse*, *Diphthérie buccale*, *Pseudo-membranous inflammation of the mouth*;—in its worst form, *Stomatyphus*, (F.) *Fégarite*. Inflammation of the mouth accompanied by the exudation of a false membrane,—a disease which is rarely seen except in large foundling establishments.

STOMATITIS, ULCEROUS, S. aphthous—s. Vesiculosa infantum, Aphthæ.

STOMATOCACE, Stomacace.

STOMATODYNIA, Stomalgia.

STOMATO-GASTRIC, *Sto'mato-gas'tricus*, *Stomo-gastric*, *Stomo-gas'tricus*, from *στομα*, *στοματος*, 'the mouth,' and *γαстр*, 'the stomach.' Appertaining to the mouth and stomach,—as the *stomato-gastric* ganglia of the lower animals. See Encephalon and Nerves.

STOMATOMALACIA PUTRIDA, Cancer aquaticus.

STOMATONECROSIS, Cancer aquaticus, Stomacace.

STOMATONOMA, Stomacace.

STOMATOPANUS, from *στομα*, *στοματος*, 'the mouth,' and *πανος*, 'a glandular tumour.' *Stomatophyma glandulosum*, *Pannus Fav'cium*. Tumefaction of the glands of the mouth.

STOMATOPHYMA, from *στομα*, *στοματος*, 'the mouth,' and *φυμα*, 'a swelling.' A swelling in the mouth.

STOMATOPHYMA GLANDULOSUM, Stomatopanus.

STOMATOPLASTIC, *Stomatoplasticus*; from *στομα*, *στοματος*, 'the mouth,' and *πλασσω*, 'I form.' An epithet for the operation of forming a mouth, where the aperture has been contracted from any cause. *Stomatoplastic*.

STOMATORRHAGIA, from *στομα*, *στοματος*, *Hæmorrhag'ia oris* seu *fav'cium*, *Sto'guinis profluvium ex Orë*, *Hæmorrhoi'des Oris*, *Stomato'sis*, *Stomorrhag'ia*, (F.) *Hæmorrhagie buccale*, from *στομα*, *στοματος*, 'the mouth,' and *ρηννυμι*, 'I break out.' Hemorrhage from the mouth.

STOMATORRHAGIA ALVEOLOSUM, Phatnorrhagia—s. *Gingivarum*, *Ulorrhagia*.

STOMATORRHŒA, Stomatorrhagia.

STOMATOSCOPE, from *στομα*, *στοματος*, 'mouth,' and *σκοπεω*, 'I view.' Any instrument for keeping the mouth open, so as to permit the parts within to be inspected; *Speculum oris*.

STOMATOSEPSIS, Cancer aquaticus.

STOMATOSIS, Stomatorrhagia.

STOMATOSPASMUS, Trismus.

STOMATYPHUS, Stomatitis, pseudomembranous.

STOMENCEPHALUS, Stomocephalus.

STOMOCEPH'ALUS, *Stomenceph'alus*, (F.) *Stomocéphale*, *Stomencéphale*; from *στομα*, 'mouth,' and *κεφαλη*, 'head.' A genus of monsters, having two contiguous eyes, or a double eye occupying the median line; the nasal apparatus atrophied and forming a tube (*trompe*); the jaws rudimentary; and the mouth imperfect or wanting.—I. G. St. Hilaire.

STOMOMANICON, *Platysma myoides*.

STOMORRHAGIA, Stomatorrhagia.

STONE IN THE BLADDER, Calculi, vesical—s. *Binder*, *Osteocolla*—s. *Crop*, *Sedum*—s. *Crop*, biting, *Sedum*—s. *Mint*, *Cunila mariana*—s. *Pook*, *Acne*—s. *Root*, *Collinsonia Canadensis*.

STONES, THE, Testes.

STOOL, Dejectio.

STOOL, CALOMEL, see *Calomel*.

STORAX, *Styrax*—s. *Liquida*, *Liquidambar styraciflua*.

STOREY'S WORM CAKES. An empirical preparation, formed of *calomel* ℥j, *jalap* ℥j, *zingib.* ℥ij, *sacch.* ℥j, *cinnab. antim.* q. s. to colour them; *syrup. simpl.* q. s. to make into cakes.

STORKBILL, *Geranium maculatum*.

STOUND, *Stitch*.

STOVE, *Ætna'rium*, *Stupha*. Old French, *Estuve*; a *Stew*, (F.) *Étuve*. A limited space, warmed artificially, and in which the air is slowly renewed. It is used for drying various substances, as plants, extracts, conserves, &c., or for taking vapour baths. In this case, the *stew* or *stove* is said to be *wet* or *humid*; in the opposite case, it is said to be *dry*, (F.) *sèche*. The latter, which is used by the Turks, is the *Lacon'ium* or *Calid'arium* of the Romans. The former is their *Tepida'rium* or *Vapora'rium*. These kinds of baths greatly excite the cutaneous transpiration; and are valuable agents in rheumatic and other affections.

STRAB, see *Strabismus*.

STRABILISMOS, *Strabismus*.

STRABIS'MUS, *Strabilis'mos*, *Parop'sis Stra-*

bismus, from *στραβος*, 'twisted,' 'one who squints,' *Strabos'ites*, *Lazophthal'mus*, *Distor'sio*, *Lus'citas*, *Illo'sis*, *Ilw'sis*, *Helw'sis*, *Squinting*, *Cross eye*, (Prov.) *Doyle*, *Gleeing*, (Sc.) *Skellie*, *Skelly*, (F.) *Strabisme*, *Vue louche*, *Louchement*. Want of concordance of the optic axes. It may be dependent upon natural or acquired inequality in the action of the motor muscles of the eye; on a convulsive state of one of those muscles; a difference in the sensibility of the two eyes; or a cerebral affection. In the treatment, if the disease seem to depend upon an excess of strength in one of the motor muscles, we must endeavour to give tone to that which is too weak, by placing before the eye a mask having a small aperture on the side to which it is necessary to direct the pupil; or by wearing glasses, provided with a piece of looking-glass, on the side to which the eye is vitiously turned; so that the reflection may be inconvenient, and occasion the organ to be turned in an opposite direction; or by putting a black patch on the angle opposite to that to which the eye is turned. If owing to weakness, the organ must be strengthened by being exerted alone, and by leaving the other at rest. If the disease be symptomatic of cerebral affection, attention must, of course, be directed to the latter.

Strabismus may be *single*, or affect one eye only, or *double*, and it may be *convergent* or *divergent*.

Those affected with Strabismus are called *Strabi*, *Strabo'nes*, *Strabo'ni*, *Strebli*, *Il'i*, *Il'ides*, *Squint'ers*, *Squint-eyed*, (Sc.) *Gleyed*, (F.) *Louches*, *Bigles*. A slight deviation is generally called a *Cast* in the *Eye*, and, sometimes, a *Cock-eye*.

STRABISMUS, CONVERGENT, see *Strabismus*—s. *Divergent*, see *Strabismus*—s. *Double*, see *Strabismus*—s. *Single*, see *Strabismus*.

STRABONES, see *Strabismus*.

STRABONI, see *Strabismus*.

STRABOSITAS, *Strabismus*.

STRABOTOMY, *Strabotom'ia*, from *στραβος*, 'one who squints,' and *τομή*, 'incision.' The operation for the removal of squinting by the division of the muscle or muscles that distort the eye-ball.

STRAE-DEATH, see *Death*.

STRAIN, Sprain.

STRAIN, *colare*, (F.) *couler*, from *σπαγγειν*, *stringere*, (F.) *estreindre*, 'to constrict or squeeze.' To pass decoctions, infusions, &c., forcibly through linen; also, to exert an effort. This is accomplished by fixing firmly the parts where the muscles to be exerted originate, in order that their full power may be developed.

STRAINING, *Nisus*.

STRAITS OF THE PELVIS, see *Pelvis*.

STRAMOINE, *Datura stramonium*.

STRAMONIA, *Datura stramonium*.

STRAMONIUM, *Datura stramonium*—s. *Fœtidum*, *Datura stramonium*—s. *Majus album*, *Datura stramonium*—s. *Vulgatum*, *Datura stramonium*.

STRANGALIDES, from *σπαγγω*, 'I compress.' Small, hard tumours, which form, as in the breast when the milk does not find issue.

STRANGALION, *Tubercle*.

STRANGALIS, *Tubercle*.

STRANGALISMUS, *Strangulation*.

STRANGERS' FEVER, see *Fever*, *strangers'*. STRANGULATIO, *Orthopnoea*, *Strangulation*—s. *Uterina*, *Angone*—s. *Vulvæ*, *Hysteria*.

STRANGULA'TION, *Strangula'tio*, *Pnig'mos*, *Strangalis'mus*, (F.) *Étranglement*, *Strangulation*. From *σπαγγω*, 'I constrict or squeeze.' State of a part too closely constricted. Thus we say that there is *strangulation* of an in-

testinal hernia, when the opening that gives passage to the portion of protruded intestine contracts, so as to intercept the continuity of the digestive canal. In *Legal Medicine*, it means the forcible obstruction of the air passages, by a ligature or by the hand, for criminal purposes. See Suffocation.

STRANGULATOR, see Lycoides.

STRANGURIA, Enuresis.—s. Cruenta, Cystorrhagia.

STRANGURIE, Strangury.—s. Menstruelle, Dysmenorrhœa.

STRANGURIOUS; same etymon as the next. Affected with, or of the nature of, strangury.

STRANGURY, *Strangu'ria*, *Paru'ria stilla'ti'tia*, from *σπρᾶγγειν*, 'to squeeze,' and *οὐρον*, 'urine.' *Dysu'ria*, *Uri'na Stillicid'ium* seu *Sub'til'ium*, (F.) *Strangurie*. Extreme difficulty in evacuating the urine, which issues only drop by drop, and is accompanied with heat, pain, tenesmus at the neck of the bladder, &c. See Ischuria, and Retention.

STRAPPING, WATER, see Water-dressing.—s. Wet, see Water-dressing.

STRATHPEFFER, MINERAL WATERS OF. A sulphuretted water, at Strathpeffer, a few miles west from Dingwall, in Ross-shire, Scotland. It contains sulphate of soda, sulphate of lime, chloride of sodium, sulphate of magnesia, and sulpho-hydric acid.

STRATIOTES, *Achillea millefolium*.

STRATUM BACILLATUM, see Tunica Jacobi.—s. Corneum, Epidermis.—s. Malpighii, *Corpus mucosum*—s. Pigmenti, see Choroid.—s. Proligerum, Proligerous disc.

STRAWBERRY, *Fragaria*—s. Bush, *Euonymus Americanus*—s. Shrubby, *Rubus arcticus*—s. Tree, *Euonymus Americanus*—s. Wild, *Fragaria Virginiana*.

STREAK, PRIMITIVE, Groove, primitive.

STREATHAM, MINERAL WATERS OF. Streatham is a village near London. The waters are saline and cathartic.

STRELLI, see Strabismus.

STREBLOSIS, Luxation.

STREMMMA, Luxation, Sprain.

STREPITO'SUS MORBUS. 'Noisy disease.'

A kind of flatulent disease, said by Sorbait to be not uncommon in the Austrian Alps, in which emphysematous tumours form on different parts of the body,—accompanied by sonorous excretion of gas by the mouth and anus.

STREPITUS AURIUM, Tinnitus aurium.

STRETCHING, Pandaliculation.

STRIE, Vibices.—s. Ciliares, Ciliary processes.—s. Corneæ, *Tenia semicircularis*—s. Longitudinales Lancisii, see *Corpus callosum*—s. Medulares, see Processes teretes.—s. Retinæ subjectæ ligamento ciliari, Halo signatus.—s. Semilunares, *Tenia semicircularis*—s. Terminales, *Tenia semicircularis*—s. Transversales Willisii, see *Corpus callosum*.

STRATED MUSCULAR FIBRE, see Muscular fibre.

STRATUS, *Cannelé*, Grooved, see Corpora striata.

STRIBILIGO, Efflorescence.

STRICTURA, Stricture.—s. Ani, Stricture of the rectum.—s. Intestini recti, Stricture of the rectum.—s. Intestini recti spasmodica, Stricture of the rectum, spasmodic.—s. Œsophagi callosa, Dysphagia constricta.—s. Œsophagi spasmodica, Œsophagismus.—s. Pharyngis seu Œsophagi vera, Dysphagia constricta.—s. Præputii, Phimosis.—s. Urethræ, Urethrostenosis.—s. Ventriculi, Stricture of the stomach.

STRICTURE, *Strictu'ra*, from *stringere*, *strictum*, 'to tie hard.' *Coarcta'tio*, *Stegno'sis*, (F.) *Rétrécissement*. A diminution or contracted con-

dition of some tube or duct, as the Œsophagus, rectum, urethra, &c. This must be dilated by appropriate bougies. Strictures may, also, occur in the intestines: they are unmanageable, with the exception of those of the rectum, which admit of topical applications.

STRICTURE OF THE ŒSOPHAGUS, Dysphagia constricta.—s. of the Pharynx, Dysphagia constricta.

STRICTURE OF THE RECTUM, SPASMODIC, *Strictu'ra Intesti'ni Recti spasmod'ica*, *Obstruc'tio Recti spas'tica*. An affection occurring in the nervous especially, which subsides spontaneously, after a longer or shorter continuance.

Stricture of the rectum, *Strictu'ra Ani* seu *Intesti'ni Recti*, *Archostegno'ma*, *Rectostenosis*, *Euthyenterostenoma*, *Archostegno'sis*, *Archostenosis*, *Proctoëcnosis*, *Angusta'tio* et *Restrictio intestini recti* seu *ani*, *Stenochoria* *intestini recti*, *Proctostenosis organ'ica*, occurs also organically, and is a very serious affection; being generally dependent upon scirrhus.

STRICTURE OF THE STOMACH, *Gastrostenosis*, *Coarcta'tio* seu *Strictu'ra ventriculi*. See Gastrostenosis cardiaca, &c.

STRICTURE OF THE URETHRA, Urethrostenosis.—s. of the Urinary bladder, Cystostenochoria.—s. Vesicæ urinariæ, Cystostenochoria.

STRIDOR DENTIIUM, Brygma.

STRIGA CARTILAGINOSA CORDIS, Isthmus of the fossa ovalis.

STRIGIL, *Strig'ilis*, *Stlen'gis*, *Stelen'gis*. A flesh-brush. Also, an instrument, anciently used in the baths, for scraping off the sweat.

STRIPED MUSCULAR FIBRE, see Muscular fibre.

STROBILITYTES, from *στροβίλος*, *stro'bilus*, 'a cone of the pine.' Wine impregnated with the cones of the pine.

STROEMIA, Cadaba.

STROKE, APOPLEC'TIC. An apoplectic seizure.

STROKE, BACK, OF THE HEART, Impulse, diastolic.

STROKE, PARALYT'IC. A sudden attack of encephalo-spinal paralysis.

STROMA, *Strag'ulum*, *Stramen'tum*; *στρομα*, 'a bed or couch,' from *σπρᾶννω*, *σπρᾶννμι*, 'to spread out; to cover.' The bed or foundation-texture of an organ,—as the stroma of the ovary. Applied, also, to the bed or base of any deposit—as the 'amorphous stroma' of serofulous deposits.

STROMBULCUS, Forceps (bullet.)

STRONGLE, Strongylus.—s. Géant, Strongylus gigas.

STRON'GYLUS, from *στρογγυλος*, 'round.' Hippocrates, Chabert, and others, mean the *Ascaris lumbricoides* by this term. The *Strongylus Gigas*, *Eustron'gylus gigas*, *Ascaris viscera'lis* seu *renalis*, *Fusaria viscera'lis* et *renalis*, (F.) *Strongle*, *S. géant*, is, however, distinct. It has an obtuse, flat head; mouth surrounded with six flattish papillæ: the whole bursa of the male truncated: the tail of the female rounded. It is sometimes met with,—five inches, a foot, a foot and a half, and even three feet long,—and from two lines to half an inch in diameter. It is occasionally found in the human kidney; rarely in other viscera, and still more rarely in the intestinal tube.

STRONGYLUS, Teres.

STROPHOS, Tormina.

STROPHULE, Strophulus.

STROPH'ULUS, *Lichen'ia asis stroph'ulus*, *Ebullit'io*, *Eranthe'ma* seu *Eco'mia strophulus*, *Red Gum*, *Red Gown*, *Tooth Rash*, *White Gum*, *Milk-spots*, (F.) *Strophule*, *Feux de dents*. It consists of an eruption of red, or sometimes

whitish pimples, occurring in early infancy, chiefly about the face, neck, and arms, surrounded by a reddish halo; or interrupted by irregular patches of cutaneous blush. All the varieties under this genus arise in consequence of the extreme vascularity and irritability of the skin at this period of life, when the constitution is liable to be disturbed by irritation, either in the alimentary canal, gums or other parts. None of these eruptions are of any importance, and no medical treatment is usually necessary. If the stomach seem at all disordered, a little rhubarb and magnesia may be administered occasionally. Willan and Bateman reckon five species, — *Strophulus intertinctus*, Red gum or gown; *S. albidus*, white gum; *S. confertus*, rank red gum, Tooth rash; *S. volatilis* or *Erythema volaticum*, and *S. candidus*.

STROPHULUS SYLVESTRIS, Ignis sylvaticus—s. Volaticus, Ignis sylvaticus.

STROPHUS, Tormina.

STRUCTIO, Structura.

STRUCTURAL; same etymon as the next. Of, or belonging to, structure.

STRUCTURE, *Structu'ra*, *Struc'tio*, *Conformatio*, *Catasceu'ē*, from *struere*, *structum*, 'to build.' The arrangement of the different tissues or organic elements of which animals and vegetables are composed. See Tissue.

STRUMA, Bronchocele, Scrofula.

STRUMA ADIPO'SA, *Obe'sitas colli*. Fatty neck. Prominence of the neck from obesity.

STRUMA FUNGOSA, Encephaloid.

STRUMES, Scrofula.

STRUMEUX, Scrofulous.

STRUMOSIS CEREBRI, Encephalophymata — s. Pulmonum, Tubercles of the lungs.

STRUMOSUS, Scrofulous.

STRUMOUS, Scrofulous.

STRUTHIOPH'AGUS, from *στροφιον*, 'a young ostrich,' and *φαγος*, 'I eat.' Struthiophagous tribes still exist in some parts of Africa.

STRUTHIUM, Saponaria.

STRUVE'S LOTION FOR HOOPING-COUGH, see Lotion, *Struve's*, for hooping-cough.

STRYCHNIA, *Strychni'na*, *Strych'nine*, *Strychni'num*, *Strych'nium*, *Vauquelina*, *Tet'anine*. An alkaline principle; solid, crystalline, inodorous, bitter, and excessively poisonous, which has been found in *Strychnos nux vomica*, *S. Colubri'na*, *S. Tieuté*, *Ignat'ia ama'ra*, *Cerbera Thevet'ia*, and *C. Aho'vai*. (*Nucis vomicae*. *rasur.* *℥iv*: *calcei pulv.* *℥vj*: *acid. muriat.* *f℥ij*; *alcohol.* *acid. sulphuric.* *dil.* *liquor. ammoniac.* *carbon. animal. purif.* *aqua.* *aa* q. s. Digest the nux vomica in two gallons of water, acidulated with a fluidounce of muriatic acid, for 24 hours: then boil for two hours, and strain with expression through a strong linen bag. Boil the residuum twice successively in the same quantity of acidulated water, each time straining as before. Mix the decoctions, and evaporate to the consistency of thin syrup; then add the lime previously mixed with a pint of water, and boil for ten minutes, frequently stirring. Pour the mixture into a double linen bag, and having washed the precipitate well with water, press, dry, and powder it. Treat the powder repeatedly with boiling alcohol, until deprived of its bitterness; mix the liquors, and distil off the alcohol by means of a water-bath. Mix the residue with water, and having applied heat, drop in sufficient diluted sulphuric acid to neutralize and dissolve the strychnia; then add purified animal charcoal; boil for a few minutes, filter, evaporate, and crystallize. Dissolve the crystals in water, and add sufficient solution of ammonia to precipitate the strychnia. Lastly, dry the precipitate on bibulous paper. Ph. U. S.) The action—therapeutical and toxic—of strychnia on

man and animals is exactly like that of the alcoholic extract of nux vomica. It is, however, more active. An eighth of a grain is sufficient to kill a large dog; and a quarter of a grain produces marked effects on the human body, when in health. It has been given in paralysis and other cases like the nux vomica. Dose, one-sixteenth to one-twelfth of a grain.

STRYCHNIA, ACETATE OF, *Strychniæ acetat* — s. Hydriodate of, *Strychniæ hydriodas* — s. Iodate of, *Strychniæ iodas* — s. Iodide of, *Strychniæ hydriodas*.

STRYCHNIA, I'ODIDE OF IODHYDRATE OF, (F.) *Iodure d'Iodhydrate de Strychnine*, is made by pouring a solution of ioduretted iodide of potassium into a solution of a salt of strychnia; treating the precipitate with boiling alcohol, and crystallizing. Used in the same cases as strychnia. Dose, a tenth to a sixth of a grain.

STRYCHNIA, MURIATE OF, *Strychniæ murias* — s. Nitrate of, *Strychniæ nitras* — s. Sulphate of, *Strychniæ sulphas*.

STRYCHNIA and ZINC, DOUBLE IODIDE OF, Zinc and Strychnia, double iodide of.

STRYCHNIE ACET'AS, *Strych'nium* seu *Strychni'num acet'icum*, *Acet'as strych'nii* seu *strych'nicus*, *Ac'etate of Strychnia* or *Strychnine*. This salt is formed by the direct combination of strychnia with acetic acid. It is given in the same cases as pure strychnia.

STRYCHNIE HYDRIODAS, *S. iodhy'dras*, *Strych'nium hydroi'd'icum*, *Hydri'odas* seu *Iodhy'dras strych'nicus*, *Hydri'odate* or *Iodide of Strychnia*, is prepared by mixing a solution of iodide of potassium with a concentrated solution of acetate of strychnia.

STRYCHNIE I'ODAS, *Strychni'ne I'odas*, *Strychni'na Ioda'ta*, *Strych'nium* seu *Strychni'num Iod'icum*, *I'odas Strych'nii* seu *Strych'nicus*, *I'odate of Strychnia* or *Strychnine*. This salt may be formed by adding a concentrated solution of iodic acid to strychnia; treating with boiling alcohol; filtering, and leaving it to spontaneous evaporation.

STRYCHNIE MU'RIAS, *Mu'riate of Strychnia*. (The product of the union of muriatic acid and strychnia.) Used in the same cases as strychnia.

STRYCHNIE NITRAS, *Strych'nium ni'tricum*, *Nitrate of Strychnia* or *Strychnine*, (F.) *Nitrate* ou *Azotate de Strychnine*. (The product of the union of nitric acid and strychnia.) Used in the same cases as strychnia.

STRYCHNIE SULPHAS, *Strych'nium sulphu'ricum*, *Sulphate of Strychnia* or *Strychnine*. Has the same properties.

STRYCHNINA, Strychnia.

STRYCHNINE, Strychnia — s. Azotate of, *Strychniæ nitras*.

STRYCHNINUM, Strychnia.

STRYCHNISM, see *Strychnos nux vomica*.

STRYCHINIUM, Strychnia — s. Aceticum, *Strychniæ acetat* — s. Nitricum, *Strychniæ nitras* — s. Sulphuricum, *Strychniæ sulphas*.

STRYCHNOS, *Solanum dulcamara* — s. Colubrina, see *Strychnia*, and *S. Nux vomica* — s. Guianensis, see *Curare* — s. Ignatii, *Ignatia amara*.

STRYCHNOS NUX VOM'ICA, *Nux Vomica*, *Metella*, *Caniram*, *Koochla tree*. *Nat. Ord. Loganiaceæ*. *Sex. Syst. Pentandria Monogynia*. (F.) *Vomiquier*. A tree of the family *Strychnoides*, which grows in India, and the seeds of which have been long sold in the shops, under the names, *Nux Vomica*, *Vomic* or *Poison nut*, *Bachelors' buttons*, &c., (F.) *Noix vomique*. For a long time, these seeds were used only for poisoning rats; but subsequently an alcoholic extract was prepared from them, which has been administered in paralytic affections, in small doses. Dose of the *Alcoholic Extract*, *Extractum nucis*

com'icæ spirituo'sum seu nucis vomicæ resino'sum seu nucis vomicæ alcohol'icum, (F.) *Extrait alcoolique de noix vomique*, one or two grains. It produces a kind of tetanic convulsion in the paralyzed parts, when given to the requisite extent. It has, also, been administered in chronic diarrhoea and dysentery. When taken to such an extent as to induce toxicological phenomena—*Strychnism*—it causes, at first, or when given in a smaller dose, greatly augmented excitability of the spinal nervous system, with slight transient paroxysms of spasms of the limbs, and hurried respiration. In larger doses, the excitability is still more developed, with paroxysms of prolonged general tetanic convulsions, laryngismus or contraction of the glottis, violent efforts of respiration, apoplexy, and asphyxia.

The following is the formula for the preparation of the *EXTRACTUM NUCIS VOMICÆ* of the Pharm U. S. Take of *Nux vomica*, ℞j; *Alcohol*, a sufficient quantity. Expose the *Nux Vomica* to steam till it is softened; then, having sliced and dried it, grind it into powder. Introduce it into an apparatus for displacement, and pour alcohol upon it gradually, until the liquid passes without bitterness. Distil off the greater part of the alcohol from the filtered liquor, and evaporate the residue to the proper consistence.

The bark is the *false angustura* or false *cusparia bark*; *Rohun*. See *Brucæa antidiysenterica*.

The wood of *STRYCHNOS COLUMBINA*, *Modira cantram*, an Indian tree, which contains strychnia, was formerly considered a specific against the bites of venomous animals; and was recommended by Boerhaave in intermittents. It is the *Lignum Colubrinum*, (F.) *Bois de Couleuvre*.

Under the names *στυχυν*, and *στυχνος*, different kinds of nightshade are described by Dioscorides.

STRYCHNOS IGNATII, *Ignatia amara*.

STRYCHNOS POTATORUM, *Clearing nut*. The fruit of a large tree of Silhet, which, when ripe, is sold in the markets of India, and has been introduced into those of London, under the name *Nirmullees* or *Nirmillies*. One of them is well rubbed for a minute or two around the inside of a vessel containing water, which it purifies by causing the impurities, which the fluid contains, to be deposited through its albumen and casein it is supposed.

STRYCHNOS PSEUDOQUI'NA, *Quina do Campo*, of Brazil, is used by the natives as a substitute for cinchona.

STRYCHNOS TIEUTÉ, see *Strychnia*, and *Upas*—*s. Toxicaria*, see *Toxifera*, and *Curare*.

STRYPHNA, *Astringents*.

STRYPHNOS, *Acerb*.

STUDIOSUS MEDICINÆ, *Philiatrics*.

STUDIUM INANE, *Aphelxia otiosa*.

STUDY, *BROWN*, *Aphelxia otiosa*.

STUFF, see *Medicament*.

STUFFING, *Cynanche trachealis*.

STULTITIA, *Fatuitas*.

STUMP, (G.) *Stumpf*, (D.) *Stomp*, 'a stump'; also, 'blunt.' (F.) *Moignon*. The part remaining from which a limb or other part has been amputated or removed in any manner. Also, the portion of a tooth remaining in the alveolus, after the corona has been removed, (F.) *Chicot*.

STUMPFoot, *Kyllosis*.

STUNNED. Old French, *Estonné*, (F.) *Étonné*, (L.) *attonitus*, 'astonished.' An epithet, applied to one who, in consequence of a fall or other accident, has received such a concussion of the brain as to deprive him, for a time, of his mental manifestations.

STUPA, *στυπη*, *Stupha*, *Stuppa*, *στυπη*, *Stuppi'on*. Tow, used in certain surgical apparatuses

and dressings, (F.) *Étoupe*. Also, a *Stupe*, that is, cloth or tow used in fomentations. A flannel or other article wrung out of hot water, plain or medicated, applied to a part, is a *stupe*. The act of applying such stupe is called *stuping*.

STUPE, *Stupa*.

STUPEFACIENTIA, *Narcotics*.

STUPEFACTIFS, *Narcotics*.

STUPEFACTIO, *Narcosis*.

STUPEFIANTS, *Narcotics*.

STUPEION, *Stupa*.

STUPEUR, *Stupor*.

STUPHA, *Stove*, *Stupa*.

STUPIA, *Tin*.

STUPIDITAS, *Idiotism*, *Imbecility*.

STUPING, see *Stupe*.

STUPOR, *Eplex'ia*, *Eplex'ia*, *Consternatio*, (Sc.) *Swarf*, *Swerf*, (F.) *Stupeur*, from *stupere*, 'to be stupefied.' Diminished activity of the intellectual faculties, often amounting to lethargy. It occurs in many affections, especially in the neuroses.

STUPOR, *Narcosis*—*s. Artuum*, *Obdormitio*—*s. Dentium*, *Hæmodia*—*s. Formicans*, *Formication*—*s. Mentis*, *Fatuitas*—*s. Vigilans*, *Catalepsy*.

STUPPE, *Stupa*.

STUPRATION, *Stuprum*.

STUPRUM, *Rape*, *Stupr'ation*, (F.) *Défloration*, *Viol*. Forceible and illicit enjoyment of a married or unmarried female. When committed upon the latter, which is most commonly the case, it is also called, *Défloration*, *Defloratio*, *Devirginatio*, *Vitatio*, *Apanthis'mus*, &c. In judging whether rape has been committed under such circumstances, the state of the genital organs; the age of the individual; and the possibility of accidents and diseases of the parts, will have to be considered. It will be necessary, also, to be acquainted with the evidences of virginity; and to determine,—if signs of virginity no longer exist,—whether its loss be attributable to copulation, or to the introduction into the vagina of any other body than the male organ, &c. Recent deflowering can, of course, be much more readily distinguished than that which has taken place some time.

STUTTERING, *Balbuties*.

STYAN, *Hordeolum*.

STYE, *Hordeolum*.

STYGMATES. M. J. Cloquet has proposed this word, or *Stigmates*, from *εργμα*, 'a mark,' to designate the marks, in form of cicatrices, which remain on the peritoneum after the obliteration of the neck of a hernial sac. They are, commonly, radiated folds, of a whitish appearance, more or less thick, and of a fibrous or fibro-cartilaginous nature.

STYLET, *Specillum*.

STYLI, *Styloid processes*.

STYLIFORM, *Styloid*.

STYLO, in composition, relating or belonging to the styloid process of the temporal bone.

STYLO-CERATOIDES, *Stylo-hyoides*—*s. Cerato-hyoides*, *Stylo-hyoides*.

STYLO-CHONDRO-HYOIDEUS. A name given, by Douglas and Albinus, to a fleshy fasciculus which the stylo-hyoid muscle sometimes presents, and which is attached to the lesser cornu of the os hyoides. It is the same fasciculus which Santorini calls *Stylo-hyo'id es novus*, *Stylo-hyoides alter*.

STYLO-GLOSSUS. A muscle situate at the anterior and superior part of the neck. It is narrow behind, much broader before; is attached to the styloid process of the temporal bone, and to the stylo-maxillary ligament; and is inserted into the side of the tongue. It raises the base of the tongue and carries it backwards.

STYLO-HYOÏDEUS, *Stylo-cer'ato-hyoïdeus*, *Stylo-ceratoï'des*, *Stylo-hyoïdes major*, (F.) *Stylo-hyoïdien*. A muscle, situate at the superior, anterior, and lateral part of the neck. It is thin and narrow, — especially posteriorly: anteriorly, it opens to give passage to the tendon of the digastricus. It is attached to the styloid process of the temporal bone, and to the body of the os hyoides. It raises the os hyoides, and carries it backwards.

STYLO-HYOID LIGAMENT is a fibrous, flattened cord, which passes from the styloid process to the lesser cornu of the os hyoides.

STYLO-HYOID NERVE is the second branch given off by the facial nerve.

STYLOID, *Styliform*, *Styloï'des*, *Graphioï'des*, *Graphoï'des*, *Graphoï'des*, *Graphidoï'des*, *Belonoï'des*, from *στυλος*, 'a style,' 'a peg,' 'a pin,' and *ειδος*, 'shape.' Shaped like a peg or pin.

STYLOID CORNUA, see Hyoides os.

STYLOID PROC'ESSES, *Pencil-like Processes*, *Processus Beloïdes* seu *Belemnoïdes* seu *Belonoï'des* seu *Graphioïdes* seu *Graphoï'des*, *Styli*, are, 1. A long and slender process of the temporal bone, to which are attached the stylo-glossus, stylo-pharyngeus, and stylo-hyoïdeus muscles, and the stylo-hyoid and stylo-maxillary ligaments. 2. Two slender and pyramidal processes at the inferior extremities of the radius and ulna.

STYLO-MASTOID, *Stylo-mastoïdeus*. That which relates to the styloid and mastoid processes.

STYLO-MASTOID ARTERY arises from the posterior auricular, and, in some subjects, from the occipital. It enters the stylo-mastoid foramen; passes along the aqueduct of Fallopius, and spreads its ramifications on the mucous membrane of the tympanum, and in the mastoid cells, and semi-circular canals; and terminates by anastomosing with a branch of the *Arteria meningea media*, which enters by the *Hiatus Fallopii*. Murray has given the name—*Stylo-mastoid* — to the posterior auricular artery itself.

STYLO-MASTOID FORAMEN, *Foramen Stylo-mastoïdeum* seu *Aqueductus Fallopii*, (F.) *Trou anonyme*, is situate at the inferior surface of the petrous portion of the temporal bone, between the styloid and mastoid processes. It terminates the aqueduct of Fallopius, and gives passage to the facial nerve.

STYLO-MAX'ILLARY, *Stylo-mazilla'ris*. That which relates to the styloid processes and jaw.

The **STYLO-MAXILLARY LIGAMENT** is a ligamentous, flat cord, extending between the styloid process and the angle of the jaw.

STYLO-PHARYNGEUS, (F.) *Stylo-thyropharyngien*. A muscle, situate at the anterior and lateral part of the neck. It is slender; thin above; attached to the styloid process of the temporal bone, and terminates in the parietes of the pharynx, as well as at the posterior margin of the thyroid cartilage. It raises the pharynx and carries it backwards. Professor Chaussier has united, under the name *Stylo-pharyngien*, the three constrictor muscles of the pharynx, the stylo-pharyngeus, and palato-pharyngeus.

STYLOPH'ORUM DIPHYL'LUM, *Meconopsis diphyl'la*, *Chelidonium diphyl'lum*, *Celandine* or *Horn Poppy*, *Bruse root*. An indigenous plant, of the family *Papaveraceæ*, growing from western Pennsylvania to Wisconsin, which flowers from May to August. The juice is fetid and narcotic. Infused in wine, it has been given internally in gravel: and has been applied externally to contusions, &c.

STYLO-THYRO-PHARYNGIEN, *Stylo-pharyngien*.

STYLUS. Sound.

STYLUS EXCAVATUS, *Specillum excavatum*. A hollow sound.

STYMA, Erection, Priapismus.

STYMATOSIS, Stigmatosis.

STYPMATA, from *στυμμα*, *στυμμαρος*, 'that which constricts or thickens.' The most solid ointments. Also, the ingredients proper for increasing their consistence.

STYPSIS, Astriiction, Constipation.

STYPTERIA, Alumen, Styptic.

STYPTIC, see Styptics.

STYPTIC, BINELLI'S, *Aqua Binellii* — s. Brocchieri's, *Aqua Brocchieri*.

STYPTIC, EATON'S. A name first given to *Helvetius's Styptic*, when introduced into Great Britain. It is now usually made of calcined green vitriol, 3j; proof spirit, tinged with a little oak bark, Oij.

STYPTIC, HELVETIUS'S, see Ferrum tartarizatum, and Styptic, Eaton's.

STYPTIC, PAGLIARI'S, *Hæmostatic*, *Pagliari's*.

STYPTIC, RUSPINI'S, *Liquor Stypticus Ruspini*. A nostrum, said to consist of gallic acid, a small quantity of sulphate of zinc and opium, dissolved in a mixture of alcohol and rose-water, which has had much reputation as an internal and external astringent.

STYPTICS, *Contrahentia*, *Constrictiva Medicamenta*, *Constringentia*, *Anastaltica*, *Catastaltica*, *Stypteria*, (F.) *Styptiques*, from *στυπτω*, 'I constringe.' Sometimes used synonymously with astringent, but generally applied to a substance employed to arrest hemorrhage, — *Enæmon*. A mechanical styptic is one that arrests the flow of blood, by being applied immediately to the mouth of the bleeding vessel, or by inducing a coagulum, mechanically, in it. A chymical styptic is one which coagulates, chymically, the blood around a bleeding orifice.

STYRAC'INUM O'LEUM. Oil obtained by boiling storax in good olive oil.

STYRAX, *στυράξ*. A resin obtained from *Styrax officinalis* and *S. calamita*. It is the solid storax; — official storax. There are two kinds of storax met with: 1. *Red Storax*, *Gum Storax*, *Thus Judaërum*, *Styrax rubra*, *Styracis balsamum*, *Balsamum styracis officinalis*, which is obtained by incision from the *Styrax officinalis*, and perhaps from the *Liquidambar Orientalis*. The purest is in tears; but it has lost some of its smell in drying. 2. *Common Storax*, *Styrax calamita*. That which has been received in reeds or vessels, and has had saw-dust added immediately to thicken it. The odour of storax is fragrant; the taste aromatic. It is considered, like the other balsams, to be stimulant and expectorant, but is rarely used long.

PURIFIED STORAX, *S. cola'ta* seu *purifica'ta*, *S. preparata*, *Extractum styracis*, is prepared by dissolving storax in alcohol; straining the solution; then distilling off the alcohol with a gentle heat, until the storax acquires the proper consistence. — Ph. U. S. Dose, gr. x to xxx.

STYRAX ALBA, see Myroxylon Peruferum — s. Benzoin, see Benjamin — s. Calamita, see Styraz — s. *Liquida*, *Liquidambar styraciflua* — s. *Officinalis*, see Styraz.

STYSIS, Erection, Priapismus.

SUAVEOLENS, Beneolens.

SUB. A common Latin prefix, often signifying 'beneath,' or 'a lower degree.'

SUBACIDUS, Acidulous.

SUBACTIO, *Chiro'sis*, *Cheiro'sis*, from *subigo*, *subactum*, (*sub*, and *ago*), 'to subject, to prepare.' The operation of making plasters, extracts, &c., with the hand, or with the pestle and mortar.

SUBAFFECTIO, Hydropathia.

SUBAGITATRIX, *Tribade*.

SUB-ATLOÏDEUS, *Infra-atloïdeus*, (F.) *Sous-atloïdien*. That which is situate beneath the

atlas or first cervical vertebra. Chaussier has given this name to the second pair of cervical nerves.

SUB-AXOÏDEUS, *Infra-axoïdeus*, (F.) *Sous-axoïdien*. That which is below the *axis* or *vertebra dentata*. Chaussier has given this name to the third pair of cervical nerves.

SUBAZOTAS BISMUTICUS, Bismuth, subnitrate of.

SUBBORAS NATRICUM, Borax.

SUBCARTILAGINEUM, Hypochondre.

SUBCLAVIAN, *Subclavius*, *Subclavicula'ris*, from *sub*, and *clavis*, 'the clavicle.' That which is under the clavicle.

SUBCLAVIAN ARTERIES, (F.) *Artères sous-clavières*, are situate at the upper part of the chest, and the lateral and lower parts of the neck. The right arises at the corresponding side of the trachea, and comes off from the arteria innominata or brachio-cephalic trunk. The left leaves the aorta at the end of the arch. Both extend as far as the superior surface of the first rib, in the space between the scaleni muscles, beyond which they take the name *axillary arteries*. The subclavian arteries run, for some distance, without sending off any branch. In the neighbourhood of the first rib, however, they give off, 1. *Upwards*, the *vertebral* and *inferior thyroïdal*. 2. *Downwards*, the *internal mammary* and *superior intercostal*. 3. *Outwards*, the *transverse cervical* or *posterior scapular*, the *superior scapular*, and the *posterior cervical* or *profunda*.

SUBCLAVIAN or **SUBCLAVIUS MUSCLE**, *Subclavi'us*, (F.) *Costo-claviculaire* (Ch.), *Sous-clavier*, is situate at the superior and anterior part of the chest. It is attached, by its inner extremity, to the superior surface of the cartilage of the first rib; and by its superior margin and outer extremity to the inferior surface of the clavicle. This muscle depresses the clavicle, and carries it forward. When the shoulder is fixed, it can raise the first rib.

SUBCLAVIAN VEINS, (F.) *Veines sous-clavières*, are continuations of the axillary, and extend from the inferior extremity of the scalenus anticus, in front of which they pass, to the vena cava superior, which they form by their union. The right subclavian vein is very short; the left, longer and broader. These veins receive the mouths of the internal and external jugulars, vertebral and superior intercostal. The left subclavian receives two veins more than the right,—the left internal mammary and left inferior thyroid. The thoracic duct opens into the left subclavian:—the corresponding great lymphatic vessel into the right.

SUBCLAVIC'ULA, from *sub*, and *clavicula*, 'the clavicle.' The first rib.

SUBCLAVICULARIS, Subclavian.

SUBCLAVIUS, Subclavian muscle.

SUBCRUEN'TUS, 'somewhat bloody;' from *sub*, and *cruentus*, 'bloody;' *Hyphæ'mos*. That which has the colour and nature of blood. An epithet given to certain excreted substances, which are mixed with blood, or have the appearance of blood.

SUBCRURÆI, see *Cruræus*.

SUBCUTANEOUS, *subcutaneus*, from *sub*, and *cutis*, 'the skin;' *Intercutaneus*, (F.) *Sous-cutané*. That which is placed immediately under the skin.

SUBCUTANEOUS SECTION, Hypodermatomy.

SUBCUTANEUS COLL, *Superficialis colli*—*s. Malæ* (Nervus), Orbital nerve.

SUBDELIRIUM, Typhomania.

SUB'DITA, *Subditi'tia*, from *subdere*, (*sub*, and *dare*), 'to put under;' *Pro'stheta*. Medicines introduced into some one of the natural apertures;—as suppositories, pessaries, &c.

SUBDITI'TIA, Subdita.

SUBDUCTIO, Dejection.

SUBER, *Cork*:—the bark of the *Quercus Suber* or *Cork Tree*, (F.) *Liège*. Family, Amentaceæ. *Sex. Syst.* Monœcia Polyandria. Cork, when burnt, is applied as a mechanical styptic to bleeding piles, &c. Surgeons use it for making tents; and it has been recommended to be worn as an amulet to suppress the secretion of milk!

SUBETH, Coma.

SUBFRONTALIS SUTU'RA, from *sub*, and *frons*, *frontia*, 'the forehead.' A suture resulting from the articulation of the os frontis with the nasal process of the superior maxillary bone, and the proper bones of the nose.

SUBGRONDATION, Depression.

SUBHUMERATIO, Catomismos.

SUBIGITATRIX, Tribas.

SUBINFLAMMATIO, Subinflammation—*s.* Hepatis, Hepatitis, chronic.

SUBINFLAMMATION, *Subinflammatio*, from *sub*, and *inflammatio*. *Inflammatio'cula*, *Hypophlegma'sia*. A mild degree of inflammation, so slight as hardly to deserve the name *inflammation*. Broussais understood by the term, an augmentation in the vital phenomena of the vessels that convey white blood. There are, however, no such vessels. Lymphatic engorgements, scrofula, herpes, and cancer he considered subinflammations.

SUBINTRAN'TES FEBRES, from *sub*, and *intrare*, 'to enter.' *Communican'tes febres*. Fevers primarily intermittent, whose paroxysms approximate, so that one paroxysm begins before that which preceded it has gone through its stages.—L. Bellini.

SUBJECTIVE SENSATIONS, see Sensation.

SUBJEE, Bangué.

SUBLIGACULUM, Truss.

SUBLIGATURA, Truss.

SUBLIMENTUM, Enœorema.

SUBLIMATE, CORROSIVE, Hydrargyri oxyurias.

SUBLIMA'TION, *Sublima'tio*, *Meteoris'mus*, *Hypo'sis*, from *sublimis*, 'exalted.' An operation by which dry and solid matters are volatilized and condensed at the upper part of a subliming apparatus. Calomel, sal ammoniac, &c., are commonly obtained by sublimation. The product of sublimation is a *Sub'imate*, *Sublima'tum*, (F.) *Sublimé*.

SUBLIMATUM CORROSIVUM, Hydrargyri oxyurias.

SUBLIMATUS CORROSIVUS, Hydrargyri oxyurias.

SUBLIME, Sublimis.

SUBLIME, Enœorema, see Sublimatiou—*s.* *Corrosif*, Hydrargyri oxyurias.

SUBLIMIS, (F.) *Sublime*, 'high, elevated, superficial.' Anatomists have given the name *sublimis* to certain muscles, which are situate more superficially than their kindred muscles. These last they have called *profundi*. Of the former, the *flexor sublimis digitorum* is an example.

The French use the term *respiration sublime* for the respiration which is accompanied by considerable elevation of the ribs, and with separation of the alæ nasi at the time of inspiration.

SUBLINGUA, Uvula.

SUBLIN'GUAL, *Sublingua'tis*, (F.) *Souslingual*, from *sub*, and *lingua*, 'the tongue.' That which is seated under the tongue.

SUBLINGUAL ARTERY, with some, is the *Lingual*. With others, it is a branch given off by the lingual opposite the genio-glossus muscle, which is distributed particularly to the sublingual gland, to the mylo-hyoid, and genio-glossus muscles, &c.

SUBLINGUAL GLAND, *Glan'dula sublingua'tis* seu *Bartholinia'na* seu *Rivinia'na*, is seated in the substance of the inferior paries of the mouth,

beneath the anterior part of the tongue. It is smaller than the submaxillary, of which it frequently seems to be only a prolongation. It is oblong, flattened, amygdaloid, and is covered by the mucous membrane of the mouth, beneath which it projects. This gland has 10 or 12 excretory ducts—*Ductus Riviniani* seu *Waltheriani*—some of which open beneath the frænum linguae, joining Wharton's duct, whilst others pierce the mucous membrane of the mouth separately. It has the same structure as the parotid.

SUBLINGUIUM, Uvula.

SUBLINGUUM, Epiglottis.

SUBLOB'ULAR, *Sublobularis*; from *sub*, and *lobus*, 'a lobe,' *lobulus*, 'a little lobe.' That which is placed under a lobe.

SUBLOBULAR VEINS OF THE LIVER. The veins in which the intra-lobular veins terminate. By their union, the sublobular veins form the hepatic veins.

SUBLUXATIO, Sprain.

SUBMAMMARY, *Submammaris*, (F.) *Sous-mammaire*, from *sub*, and *mamma*, 'the breast.' Under the mamma or breast:—hence, 'submammary inflammation,' inflammation of the areolar tissue beneath the mamma.

SUBMAXILLARY, *Submaxillaris*, *Infra-maxillaris*, (F.) *Sous-maxillaire*, from *sub*, and *maxilla*, 'the jaw.' That which is seated beneath the jaw.

SUBMAXILLARY GLANDION is a small nervous ganglion, situate opposite the gland. It seems to be formed by the superior branch of the Vidian nerve, and communicates with the lingual nerve of the inferior maxillary. Its filaments form a plexus which penetrates the submaxillary gland.

SUBMAXILLARY GLAND, *Maxillary Gland*, *Hypognathic aden*, *Hypoglossal aden*, is a salivary gland, less than the parotid, seated at the inner side of the ramus and body of the inferior maxillary bone, in the triangular space between the two bellies of the digastricus. Irregularly ovoid, and flattened on three surfaces, it has the same aspect and organization as the other salivary glands. Its excretory duct is called *Wharton's Duct*, and terminates at the side of the frænum linguae, by a very small orifice. It is also called *Ductus inferior*.

SUBMENTAL, *Submentalis*, (F.) *Sous-mental*, from *sub*, and *mentum*, 'the chin.' Situate under the chin.

SUBMENTAL ARTERY is furnished by the facial, near the base of the jaw. It passes forwards along the attachment of the mylo-hyoideus, to which it furnishes branches that cross it to anastomose with those of the sublingual. Near the median line it bifurcates, to be distributed to the chin and muscles of the infra-hyoid region.

SUBMENTAL VEIN, which accompanies the artery, opens into the labial.

SUBMERSION, *Submersio*, from *sub*, and *mergere*, *mergum*, 'to plunge.' The act of plunging, or being entirely plunged, in a liquid.

Asphyxia by submersion, *A. Immersorum*, *Drowning*, is not owing to a certain quantity of liquid being introduced into the alimentary or air-passages; but simply to the interception of air, and of the respiratory phenomena. It is a case of suffocation, the signs of which, on dissection, are equivocal, without circumstantial evidence.

SUBMISSIO, Remission—s. *Cordis*, *Systole*.

SUBMU'COUS, *Submucoeus*; from *sub*, and *mucus*. Under a mucous membrane,—as the 'submucous areolar tissue.'

SUBMURIAS AMMONIACO-HYDRARGYRICUS, *Hydrargyrum præcipitatum*.

SUBMUSCULAR, *Submuscularis*; from *sub*,

and *musculus*, 'a muscle.' Seated beneath muscles or a muscular layer.

SUBOCCIPITAL, *Suboccipitalis*; from *sub*, and *occiput*. That which is situated under the occiput.

SUBOCCIP'ITAL NERVE, Occipital nerve.

SUBORBITAR, *Suborbitarius*, *Infra-orbital*, *Infra-orbitalis*, *Infra-orbitarius*, (F.) *Sous-orbitaire*, from *sub*, and *orbita*, 'the orbit.' That which is seated beneath the orbital cavity.

SUBORBITAR ARTERY, *Infra-orbital A.*, proceeds from the internal maxillary, from which it separates near the anterior and superior part of the zygomatic fossa. It enters the infra-orbital canal, and passes through it, giving branches to the anterior and superior dental canal; issues by the foramen infra-orbitarium; and divides, in the fossa canina, into a number of branches, which lose themselves in the neighbouring parts. The infra-orbital vein follows the same course.

SUBORBITAR CANAL, *Infra-orbital Canal*, (F.) *Canal ou Conduit sous-orbitaire*, is a small canal, which runs obliquely through the substance of the inferior paries of the orbit. It begins, behind, by a simple gutter, and divides, anteriorly, into two branches, one of which descends into the anterior paries of the maxillary sinus, whilst the other passes out, in the canine fossa, by an orifice, called *Foramen-infra-orbitarium*, (F.) *Trou sous-orbitaire*. The infra-orbital artery and nerve pass along it.

SUBORBITAR FISSURE, *Infra-orbital Fissure*, *Fissura infra-orbitalis*, *Rima canalis orbitalis*. A fissure in the superior maxillary bone; the superior commencement of the suborbitar canal.

SUBORBITAR FOSSA, Canine fossa.

SUBORBITAR NERVES, *Infra-orbital Nerves*, seem to terminate the superior maxillary. They issue from the infra-orbital canal by the foramen of the same name, and spread out in a radiated manner in the fossa canina, to divide into *superior* or *palpebral filaments*, *inferior* or *labial*, *internal* or *nasal*, and *external* or *buccal*.

SUBPERITONEAL, *Subperitonealis*; from *sub*, and *peritoneum*. That which lies under the peritoneum.

SUBPERITONEAL APONEUROSIS, Fascia, subperitoneal—s. Ganglia, see Spermatic ganglion.

SUBPLACENTA, Decidua membrana.

SUBPOPLITEUS, Popliteus muscle.

SUBPUBIC ARCH, *Arcus subpubicus*, from *sub*, and *pubis*. A strong arch of the superior pelvic aponeurosis, which completes the posterior orifice of the obturator or subpubic canal.

SUBPUBIC MEMBRANE, Obturator membrane.

SUBPUBIO-FEMORALIS, Abductor brevis—s. Prostaticus, Compressor prostatæ.

SUBPURATIO, Suppuration.

SUBPUTRIS, Hyposaprus.

SUBSCAP'ULAR, *Subscapularis*, (F.) *Sous-scapulaire*, *Sous-scapulo-trochinnien*, from *sub*, and *scapula*, 'the shoulder-blade.' That which is beneath the scapula.

SUBSCAPULAR FOSSA, *Fossa Subscapularis*. A considerable excavation at the anterior surface or venter of the scapula, in which the subscapularis is seated.

SUBSCAPULAR MUSCLE, *Subscapularis Muscle*, *Immer'sus*, *Infra-scapularis*, (F.) *Sous-scapulo-trochinnien* (Ch.), *Porte-feuille*, *Muscle-sous-scapulaire*. A muscle situate in the above-mentioned fossa. It is flat, broad, and triangular; is attached, by its base, to the subscapular fossa and to the anterior lip of the edge of the spine of the scapula, and is inserted into the lesser tuberosity of the humerus. It carries the arm in rotation inwards. It can also depress it, and bring it towards the body.

SUBSCAP'ULAR NERVES are two in number.

One arises from the brachial plexus above the clavicle; the other from the posterior aspect of the plexus within the axilla. They are distributed to the subscapularis muscle.

SUBSEROUS, *Subsero'us*; from *sub*, and *serous*. Under a serous membrane,—as the 'subserous areolar tissue.'

SUBSIDENTIA, *Sediment*.

SUBSTANCE ÉMAILÉE DES DENTS, Enamel of the teeth — s. Horny, see Tooth.

SUBSTANCE, HYALINE, *Cytoblastema* — s. Intercellular, *Cytoblastema* — s. *Vitrée des Dents*, Enamel of the teeth.

SUBSTANCE, WHITE, OF SCHWANN. A hollow cylinder within the tubular nervous fibre, which differs in composition and refracting power from the matter that occupies the centre of the tube.

SUBSTANTIA ADAMANTINA DENTUM, Enamel.

SUBSTANTIA FERRUGIN'EA, *Locus cine'reus*. An ash coloured substance, produced by the existence of well marked pigmentary matter in the floor of the fourth ventricle or *rhomboid fossa*.

SUBSTANTIA FILAMENTOSA DENTUM, Enamel — s. *Grisea Centralis*, see *Medulla spinalis* — s. *Hyalina*, *Cytoblastema* — s. *Nigra pedunculi*, see *Peduncles of the brain* — s. *Ossea dentium*, see *Tooth* — s. *Ostioidea*, see *Tooth* — s. *Perforata anterior*, *Locus perforatus anticus*.

SUBSTANTIA PERFORATA MEDIA. When the pia mater has been removed from the space between the crura cerebri, termed the *intercrural* or *interpunctular*, the surface appears cribriform from the perforations of the numerous minute vessels which penetrate it. This is the *substantia perforata media* of Vicq d'Azyr.

SUBSTANTIA PERFORATA POSTERIOR, see *Tarini pons* — s. *Pulposa Dentis*, *Dental pulp* — s. *Radiata*, *Stellula Verheyenii* — s. *Rhomboides*, *Corpus dentatum* — s. *Spongiosa Urethra*, *Corpus spongiosum urethrae* — s. *Stellata serpentina*, *Stellula Verheyenii* — s. *Vitrea dentium*, Enamel of the teeth — s. *Vitrea*, *Cytoblastema* — s. *Vitrea dentium*, Enamel.

SUBSTANTIVE, *Substantivus*; from *sub-stante*, (*sub*, and *stante*), 'to stand under or subsist.' An epithet applied by Dr. Paris to aliments which are nutritious, in contradistinction to *adjective* aliments, which are themselves not possessed of nutritive virtues, but impart to the digestive organs greater energy, so as to enable them to extract more nutriment from the substantive aliments. Meat, for example, is a *substantive* aliment: the condiment, as mustard, eaten with it, an *adjective* aliment.

SUBSTILLUM, *Stillicidium*.

SUBSULTIO, *Palpitation*.

SUBSULTUS TEN'DINUM, *Saltus tendinum*, *Clonus subultus*, *Subultus*, *Carpolog'ia spasmod'ica*, *Myopal'mus*, *Twitching of the Tendons*, (F.) *Soubresaut*, from *subsulto*, *subsultum*, (*sub* and *salto*), 'to make short leaps.' Twitching, communicated to the tendons by the involuntary and instantaneous contraction of the muscular fibres. It is more observable at the wrist than elsewhere; and is an evidence of great cerebral irritability, and debility, when it occurs at an advanced period of fever.

The muscular agitations or twitchings observed in febrile diseases, especially of children, have been termed *Convulsibilitas*, *Scleroty' bē febrilis*, and *Irritabilitas morbo'ea*.

SUBSURTITAS, *Deafness*.

SUBU'BERES, from *sub*, and *ubera*, 'the breasts.' *Infan'tes eugen'tes*, *Sucking children*. Infants at the breast, in contradistinction to *Ecu'beres*, *απογαλακτοι*, or those which have been weaned.

SUBUN'GUAL, *Subun'qual*, *Subun'guic'al*, (F.) *Sous-ungéale*, from *sub*, and *unguis*, 'a nail.' Belonging to parts under the nail; — as *subungual exostosis*, (F.) *Exostose sous-ungéale*, an exostosis under the nail of the great toe especially.

SUBVERSIO STOMACHI, *Vomiturition*.

SUBVERTEBRA, *Sacrum*.

SUBVERTEBRUM, *Sacrum*.

SUBVOLA, *Hypothenar*.

SUC, *Succus* — s. *de Citron*, see *Citrus medica* — s. *Gastrique*, *Gastric juice* — s. *de Kino*, *Kino* — s. *Médullaire*, *Marrow* — s. *du Limon*, see *Citrus medica* — s. *Nourricier*, *Liquor sanguinis* — s. *Pancreatique*, see *Pancreas*.

SUCCA'GO, from *succus*, 'juice.' The inspissated juice of plants; robs, jellies, &c.

SUCCÉDANÉ, *Succedaneum*.

SUCCEDA'NEUM, *Succida'neum*, *Substitutum*, *Antibal'lo'enum*, *Surroga'tum*, *Antembal'lo'enum*, *Quid pro quo*, from *succedere*, (*sub*, and *cedere*), 'to go under, to come in place of another.' (F.) *Succédant*. An epithet for medicines that may be substituted for others possessing similar properties. Also, used, in dentistry, for an amalgam for filling teeth.

SUCCENTURIA'TUS, from *succenturiare*, (*sub*, and *centuria*), 'to supply a soldier to fill a vacancy;' and hence, 'to substitute or put in the room of.' That which is a substitute for another; as *Renes succenturiati*, *Ventriculus succenturiatus*, &c.

SUCCHAR, *Saccharum*.

SUCCIDANEUM, *Succedaneum*.

SUCCIN, *Succinum*.

SUCCINCTURA, *Diaphragm*.

SUCCINCTUS, *Diaphragm*.

SUCCINI RESINA, *Musk*, artificial.

SUCCIN'IC ACID, *Ac'idum Suc'cini*, *A. Succin'icum*, *Acor Succin'eus*, *Sal Succini*, *Sal volatile Succini*, has been considered antispasmodic and diaphoretic, in the dose of from five to twenty grains.

SUC'CINUM, *Electrum*, *Ambra*, *Ambra'm*, *Am-par*, *Beren'cē*, *Amber*, *Yellow Amber*, (F.) *Succin*, *Ambre jaune*, from *succus*, 'juice,' because at one time supposed to exude from a tree. This kind of fossil resin, which is found on the shores of the Baltic, is composed of a resinous matter, of essential oil, and an acid *sui generis*. It is inodorous, except when heated or rubbed; insipid; in fragments of a pale golden-yellow colour, transparent, and has a shining lustre. S. g. 1·08; insoluble in water, and slightly acted on by alcohol. It is used for affording its essential oil — the *O'leum seu Balsamum Suc'cini*, (F.) *Huile de Succin*. This oil possesses stimulant, antispasmodic, diuretic, and rubefacient properties.

The *O'leum Suc'cini rectificatum*, *Rectified Oil of Amber*, is made as follows:—*Oil of Amber*, Oj; water, Ovj. Mix in a glass retort, and distil until Oiv of the water have passed with the oil into the receiver; separate the oil from the water, and keep it in well-stopped bottles.—(Ph. U. S.)

The *White amber*, *Leucelectrum*, (F.) *Ambre blanc*, is a variety.

SUCCINUM CINEREUM, *Ambergris* — s. *Griseum*, *Ambergris*.

SUCCION, *Sucking*.

SUCCISA, *Scabiosa succisa* — s. *Pratensis*, *S. succisa*.

SUCCOCOLLATA, *Chocolate*.

SUCCOLATA, *Chocolate*.

SUCCORY, *WILD*, *Cichorium intybus*, *Chironia angularis*.

SUC'COTASH, *Suck'atash*. A term of Indian derivation. A favorite dish in New England, formed of green Indian corn and beans boiled together.

SUC'CUBUS, from *sub*, and *cubare*, 'to lie.' Some authors have used this word synonymously with nightmare. Others mean by it a female phantom, with which a man in his sleep may believe he has intercourse;—as *Inebulus* has been applied to the male phantom, with which a female may dream she is similarly situate.

SUCCUS, *Chylus*, *Opos*, *Opis'ma*, *Chylis'ma*, *Eechylo'ma*, *Apothlim'ma*, *Juice*, (F.) *Suc*, from *eugere*, suctum, 'to suck.' The fluid obtained by pressing plants, flesh, &c.; *Succus expres'sus*, *Opol's* (official), *Opolide* (magistral), according to Chréaen. It is, also, applied to certain animal fluids, as *S. Gastricus*, &c.

SUCCUS ACACIÆ GERMANICÆ INSPISSATUS, see *Prunus spinosa*—*s. Acaciæ nostratis*, see *Prunus spinosa*—*s. Arborum*, *Sap*—*s. Cicutæ spissatus*, *Extractum conii*—*s. Cyrenaiscus*, *Laserpitium*.

SUCCUS ENTER'ICUS, *S. Intestinal'is*, *Liquor intestinal'is seu enter'icus*, *Intes'tinal Juice*. The fluid copiously secreted from the lining membrane of the small intestines.

SUCCUS EXPRESSUS, *Apothlimma*—*s. Gastricus*, *Gastric juice*—*s. Glycyrrhizæ*, *Extractum glycyrrhizæ*—*s. Hypocistidis*, see *Cytinus*—*s. Intestinalis*, *Succus entericus*—*s. Japonicus*, *Catechu*—*s. Limonis*, see *Citrus medica*—*s. Liquiritiæ*, *Extractum glycyrrhizæ*—*s. Nervæus*, *Nervous fluid*—*s. Nutritius*, *Chyle*—*s. Orni concretus*, see *Fraxinus ornus*—*s. Pancreaticus*, see *Pancreas*—*s. Spissatus Aconiti Napelli*, *Extractum aconiti*—*s. Spissatus atropæ belladonnæ*, *Extractum belladonnæ*—*s. Spissatus conii maculati*, *Extractum cicutæ*—*s. Spissatus hyoseyami nigri*, *Extractum hyoseyami*—*s. Spissatus lactucæ sativæ*, *Extractum lactucæ*.

SUCCUS'ION, *Succus'io*, *Hippocrat'ic Succus'ion*, from *succutio*, *succussum*, (*sub*, and *quatio*), 'I shake from beneath.' A mode of ascertaining the existence of a fluid in the thorax, ('fluctuation by succussion,') by giving the body one or more slight shakes.

SUCCUSSIO, *Ballotement*.

SUCHOR, *Saccharum*.

SUCKATASH, *Succotash*.

SUCKING, *Suc'tio*, *Suctus*, *Athel's'is*, *Bdalis*, *Myse'is*, (F.) *Succion*, from *eugere*, suctum, 'to suck.' The act of drawing liquid into the mouth, by forming a vacuum with the tongue acting as a piston, during inspiration.

SUCKING BOTTLE. A bottle so formed, that, when filled with milk, it can be sucked from instead of the breast.

SUCKING CHILD, *Lactens*. A suckling.

SUCKLE, *Lactare*, *γαλακτορροειν*, *τιθησκειν*, (F.) *Allaiter* on *Alaiter*. Same etymon. To give suck. To nurse.

SUCKLING, *Lactation*, *Lonicera periclymenum*, *Sucking child*.

SUCRE, *Saccharum*—*s. Candi*, *Saccharum candidum*—*s. du Foie*, see *Saccharum*—*s. de Lait*, *Sugar of milk*—*s. de Saturne*, *Plumbi superacetas*—*s. Tors*, *Penidium*.

SUCTIO, *Exsuctio*, *Sucking*.

SUCTION POWER. The force exerted on the blood in the veins, by the active dilatation of the heart. See *Derivation*.

SUCTORIA, from *eugere*, suctum, 'to suck.' Animals provided with mouths for sucking. Also, organs for sucking or adhesion,—hence termed *suctorial*—as 'suctorial oscula.'

SUCTORIAL, see *Suctoria*.

SUCTUS, *Sucking*.

SUDAM'INA, *Hydro'ata*, *Hidro'ta*, *Pap'ule sudora'les*, *Sufferm'ra*, *Alha'sef*, *Planta noctis*, *Boa*, from *sudor*, 'sweat.' Small vesicles, which appear upon the skin, especially in the summer time, in hot countries, and in diseases attended

with much sweating. It is a miliary eruption. See *Miliary fever*.

SUDAMINA, *Desudatio*, *Échaubouluures*, *Lichen tropicus*.

SUDARIUM, *Achicolum*.

SUDATIO, *Ephidrosis*.

SUDATIUNCULA, *Ephidrosis*.

SUDATORIA, *Hidrosis*—*s. Maligna*, see *Sudor Anglicus*.

SUDATORIUM, *Achicolum*.

SUDATORY, *Achicolum*.

SUDOR, *ἰδρως*, *Hidros*, (F.) *Sueur*, 'sweat.' The product of cutaneous transpiration, when visible on the surface of the body. It is colourless; of an odour more or less strong and variable; and of a saline taste. It is composed, according to Berzelius, of water, lactic acid, lactate of soda united with a little animal matter, chloride of sodium, and a little chloride of potassium, an atom of earthy phosphate, and some oxyd of iron.

SUDOR, *Miliary fever*.

SUDOR AN'GLICUS, *Hidron'osus*, *Hidrop'yra*, *Hidrop'yretos*, *Febris seu Mili'aris sudato'ria*, *Sudor mili'aris*, *Morbus sudatorius*, *Ephem'era maligna seu sudatoria seu Britan'nica seu Anglica pest'ilens*, *Pestis Britannica*, *Sudato'ria maligna*, *Sweating Sickness*, *Sweating Fever*, (F.) *Suette*, *S. Miliare*, *S. épidémique*, *Sueur Anglaise*. A very severe epidemic disease, characterized by profuse sweating, which appeared in England in 1486; and recurred, at different times, until about the middle of the 16th century. It was accompanied with coldness, excessive prostration of strength, palpitation, frequency and inequality of the pulse, &c., and terminated favourably or unfavourably in the course of 24 hours. The French have given the name *SUETTE DE PICARDIE*, *Suette des Picards*, *Fièvre suante*; *Mili'aris sudatoria*, *Sudor Picard'icus*, to an epidemic disease, which has appeared several times in that province; the principal symptoms of which were profuse sweats and a miliary eruption. The disease recurred in 1821, and has been described, at length, by M. Rayer, who considers it to consist of a simultaneous state of inflammation of various tissues; and proposes to class it with variola, scarlatina, and measles.

A form of it, in which the perspiration was of a black colour, was called *Sudor An'glicus niger*, *Melanephidro'sis*, *Black English Sweating sickness* or *fever*.

SUDOR COLLIQUATIVUS, *S. Profusus*.

SUDOR CRUEN'TUS, *S. sanguin'eus*, *Hæmatidro'sis*, *Hæmathidro'sis*, *Ephidro'sis cruen'ta*, *Hæmor-rhag'ia per Cutem*, (F.) *Hématidrose*, *Sueur de Sang*; *Sweating of blood*, *Bloody sweat*, *Hemorrhage from the skin*. Cutaneous perspiration intermixed with blood:—a very uncommon affection.

SUDOR DULCIS, *Saccharorrhœa cuta'nea*, *Ephidro'sis sacchar'a'ta*. A discharge of saccharine matter by perspiration.

SUDOR LOCALIS, *Meridrosis*—*s. Miliaris*, *Miliary fever*, *Sudor Anglicus*—*s. Partialis*, *Meridrosis*.

SUDOR PROFU'SUS, *S. Colliquati'vus*. A profuse or immoderate sweat.

SUDOR SANGUINEUS, *Sudor cruentus*—*s. Universalis*, *Panhidrosis*.

SUDORIFER, *Sudoriferous*, *Sudorific*.

SUDORIFEROUS, *Sudor'ifer*, *Hydroph'orus*, from *sudor*, and *fero*, 'I carry.' That which carries sweat, as the *sudoriferous* ducts or glands.

SUDORIFIC, *Sudorific'um*, *Sudor'ifer*, *Hidrot'icum*, *Hidrot'erium*, *Hidrotrope'um*, *Hidropoë'ticum*, (F.) *Sudorifique*, from *sudor*, and *facio*, 'I make.' A medicine which provokes sweating. See *Diaphoretic*.

FOUR SUDORIFIC WOODS, *Quatuor ligna sudorifica*, were, formerly, *guaiacum*, perfumed cherry, *sassafras*, and *sassafras*.

SUDORIPAROUS, see Perspiration.

SUDORIS NULLITAS, Anidrosis — *s. Privatio*, Anidrosis — *s. Suppressio*, Ischidrosis.

SUET, Sebum.

SUETTE, Sudor Anglicus — *s. Epidémique*, Sudor Anglicus — *s. Militaire*, Sudor Anglicus — *s. de Picardie*, Sudor picardicus — *s. des Picards*, Sudor picardicus.

SUEUR, Sudor — *s. Anglaise*, Sudor Anglicus — *s. d'Expression*, Expression, sweat of — *s. de Sang*, Diapedesis, Sudor oruentus — *s. Urineuse*, Uridrosis.

SUFFERSURÆ, Sudamina.

SUFFIMEN, Suffimentum.

SUFFIMENTUM, *Hypocapnis'ma*, *Suffi'men*, *Thymia'ma*, *Suffi'tus*, *Suffumiga'tio*, *Suffumiga'tion*, (F.) *Parfum*; from *suffio*, 'I perfume.' A perfume; suffumige; fumigation.

SUFFITIO, Fumigation.

SUFFITUS, Fumigation, Suffimentum.

SUFFLATIO, Puffiness.

SUFFOCATIO, Suffocation, Orthopnoea — *s. Hypochondriaca*, Hypochondriasis — *s. Hysterica*, Angone — *s. Stridula*, Cynanche trachealis — *s. Uterina*, Angone.

SUFFOCATION, *Suffoca'tio*, *Præfoca'tio*, *Apopnæ'is*, *Asphyx'ia*, *Melanæ'ma*, from *eub*, and *fals*, pl. *fauces*, 'the jaws.' Death, or suspended animation from impeded respiration, whether caused by the inhalation of noxious gases, drowning, hanging, strangling, or smothering. The principal morbid appearances in such cases are:—the lungs of a deep-blue colour, with the blood extravasated in the air-cells; right auricle and ventricle filled with dark blood, as well as the neighbouring veins; lividity of the countenance, turgescence, and, perhaps, rupture of the vessels of the brain.

Treatment of suspended animation by suffocation in general. The patient must be conveyed into a room not too warm. Blood-letting must be used, if at all, with caution;—friction must be employed with salt, or warm flannels; stimulating fluids, in a dilute state, be poured into the stomach by means of a tube, and attempts be made to inflate the lungs. Laryngotomy, if necessary.

SUFFOCATION, *Pnigopho'bia*, (F.) *Étouffement*, is used, as *strangling* is, for threatened suffocation. It means the state of dyspnoea and oppression experienced when a sufficient quantity of air does not enter the lungs, or when the chemical phenomena of respiration are imperfectly executed; as in cases of asthma.

SUFFOCATION UTERINE, Angone, Hysteria.

SUFFRAGO, Poples.

SUFFUMIGATION, Suffimentum.

SUFFUMIGE, Suffimentum.

SUFFUMIGIUM, Fumigation.

SUFFUMINATIO, Fumigation.

SUFFUSIO, Pseudoblepsia — *s. Auriginosa*, Icterus — *s. Bilis*, Icterus — *s. Colorans*, Chromopsia — *s. Dimidiata*, Hemipopia — *s. Exclarans*, Oxyopia — *s. Fellis*, Icterus — *s. Lentis crystallinae*, Cataract — *s. Metamorphopsia*, Metamorphopsia — *s. Multiplicans*, see Diplopia — *s. Myodes*, Metamorphopsia — *s. Nigra*, Amaurosis — *s. Oculi*, Cataract, Pseudoblepsia — *s. Oculorum*, Metamorphopsia — *s. Radians*, Scotasma — *s. Sanguinis*, Echymona.

SUGAR, Saccharum — *s. Barley*, Penidium, Saccharum hordeatum.

SUGAR CANDY, Saccharum candidum — *s. Cane*, see Saccharum.

SUGAR, DIABETIC, Glucose — *s. Fruit*, Glucose

— *s. Grape*, Glucose — *s. Hepatic*, see Saccharum — *s. Honey*, Glucose.

SUGAR OF LEAD, Plumbi superacetat.

SUGAR, LIVER, see Saccharum.

SUGAR, MAGNETIC, *Saccharum Magnet'icum*. A quack preparation, said to be composed of sugar and cannabis. It is given in nervous cases.

SUGAR OF MILK, *Saccharum lactis*, *Galac'to-saccharum*, *Lactose*, *Lactine*, (F.) *Sucre de Lait*, *Sel de Lait*, is obtained from whey by evaporation and crystallization. It possesses the demulcent properties of the sugars.

SUGAR. MUSCLE, Inosite.

SUGAR PINE, see Arrowroot — *s. Starch*, Glucose.

SUGAR, URINARY, Glucose.

SUGARS, VARIOUS, see Saccharum.

SUGARBERRY, *Celtis occidentalis*.

SUGGILLATION, *Sugilla'tio*, *Livor*, *Ecde'ma cruentum*, *Sugillation*, *Bruise*; from *sugillare*, *sugillatum*, 'to beat black and blue.' By most authors used synonymously with ecchymosis and ecchymoma. It is, also, applied to the spots, or ecchymoses which occur in consequence of intrinsic causes,—in disease and in incipient putrefaction. It is common in dead bodies; *Livor seu macula mortuaria*. To this last case it is, indeed, restricted by some medico-legal writers; and an interesting question sometimes arises:—Whether the appearance have been induced before death or afterwards? Whether it be a case of ecchymosis or of sugillation? In the former, the extravasated blood will usually be coagulated; not so in the latter. It is often, however, difficult to form a correct judgment without reference to the history of the case.

SUGGRUNDIUM, perhaps from *suggredi*, (*eub*, and *gradior*), 'to go or come up to.' The eave of a house.

SUGGRUNDIUM OCULI, Superciliary arch — *s. Superciliorum*, Frontal protuberance.

SUICIDA, Autochir.

SUICIDAL INSANITY, Suicide.

SUICIDE, *Suici'dium*: from *sui cædes*, 'murder of one's self.' The act of killing one's self. *Autoch'ria*, *Autocton'ia*, *Idiocton'ia*, *Autophon'ia*. Suicide is very frequently the result of disease of the mind: the individual seeming, at times, to be irresistibly impelled to self-destruction. — *Melancho'lia Suici'dium seu autoch'ria*, *Suici'dal Insanity*. Also, one who commits self-murder.

SUIE, Fuligo.

SUIF, Sebum.

SUITES DE COUCHES, Lochia.

SULCATUS, *Cannelé*, Grooved.

SULCI, plural of Sulcus.

SULCI CILIARES, Halo signatus.

SULCULUS LABII SUPERIORIS, Lacuna Labii superioris.

SULCUS, *Semicanal'is*, *Semicanal'culus*. A furrow. A groove. (F.) *Sillon*. Anatomists have given this name to grooves on the surface of bones and other organs. See Anfractuosity.

French writers on anatomy often use the term *Sillons* for the grooves which lodge the arteries and creep on the surface of bones; whilst those that receive veins are called *gouttières*, 'gutters.' Also, the vulva.

SULCUS ANTERO-POSTERIOR JECORIS, see Liver — *s. Labii superioris*, Lacuna Labii superioris.

SULCUS, LAT'ERAL, ANTE'RIOR, OF THE SPINAL MARROW. A mere trace on the marrow, marked only by the attachment of the filaments of the anterior roots of the spinal nerves.

SULCUS, LAT'ERAL, POSTERIOR, OF THE SPINAL MARROW, corresponds with the attachment of the posterior roots of the spinal nerves.

SULCUS RETRO-GLANDULARIS, see Glans — *s. Sinus lateralis*; *Fossa sigmoidea* — *s. Transversalis*, see Liver — *s. Umbilicalis*, see Liver.

SULFATE D'AMMONIAQUE, Ammoniz sulphas.

SULFIDUM CARBONII, Carbonis sulphuretum.

SULFUR, Sulphur.

SULFURE DE CARBON, Carbonis sulphuretum.

SULPHAS ALUMINARIS, Alumen—s. Ammoniacæ Cupratus, Cuprum ammoniatum—s. Caduricus, Cadmii Sulphas—s. Kalico-aluminicum, Alumen—s. Natrius, Soda, sulphate of—s. Zincum, Zinci sulphas.

SULPHUR, *Sulfur*, *Scorith*, *Thei'on*, *Thion*, *Aerich*, *Terra fuliata*, *Fumus citrinus*, *Aquila*, *Brimstone*, (F.) *Soufre*. Native Sulphur is called *Rock* and *Black Sulphur*, *Sulphur vivum* seu *nigrum*. Sulphur, as we meet with it, is in rolls, *Sulphur in rotulis* seu *rotundum*, *Roll Brimstone*. It is a volcanic production, and is obtained, in large quantities, from *Solfatara*, in Italy. *Roll Sulphur* is impure sulphur, melted, and run into moulds. It is unpleasantly inodorous, when heated or rubbed; insipid; solid, and brittle. S. g. 1.99. Fusible at 226°; volatilizable by heat, and condensing unchanged.

SULPHUR ANTIMONIATUM, Antimonii sulphuretum præcipitatum—s. Caballinum, see Sulphur sublimatum—s. Carburet of, Carbonis sulphuretum—s. Chloratum, Sulphur, chloride of.

SULPHUR, CHLORIDE OF, *Sulphuris Chloridum* seu *Dichloridum*, *Sulphur chloratum*, *Hypochloritum sulphuræum*, *Chlorum hypersulphuratum*, *Dichloride* or *Protochloride* or *Hypochloride* or *Subchloride* of Sulphur, *Bismphuret of Chlorine*, is obtained by passing dry chlorine gas over washed and dried flowers of sulphur, until they are for the most part dissolved, and distilling the decanted fluid from the excess of dissolved sulphur. One drachm of the liquid to an ounce of lard has been used in lepra, psoriasis, and other chronic cutaneous diseases. Ten drops, dissolved in ether, have been given in adynamic fever.

SULPHUR, DICHLORIDE OF, Sulphur, chloride of—s. Hypochloride of, Sulphur, chloride of—s. Iodatum, Sulphuris iodidum—s. Iodide of, Sulphuris iodidum—s. Ioduret of, Sulphuris iodidum—s. Liver of, Potassæ sulphuretum.

SULPHUR LOTUM, *Sulphur sublimatum lotum*, *Washed Sulphur*, *Magistrium Sulphuris*, *Sulphuris Flores loti*, is prepared by pouring boiling water on sublimed sulphur, so as to wash away any acid it may contain, and then drying it.

SULPHUR NIGRUM, *Sulphur vivum*, see Sulphur sublimatum.

SULPHUR PRÆCIPITATUM, *Lac Sulphuris*, *Milk of Sulphur*, *Præcipitated Sulphur*, (F.) *Crème de Soufre*, is prepared by boiling sublimed sulphur and lime together, and adding muriatic acid to precipitate the sulphur.

SULPHUR, PROTOCHLORIDE OF, Sulphur chloride of—s. Subchloride of, Sulphur, chloride of.

SULPHUR SUBLIMATUM, *Sulphur*, (Ph. U. S.) *Sublimed Sulphur*, *Flowers of Sulphur* (F.) *Soufre sublimé*, *Flours de Soufre*, is the sulphur of commerce, sublimed in close vessels.

Sulphur is possessed of stimulant, laxative, and diaphoretic properties. As a laxative, it is used in hemorrhoidal and other affections. In the itch, applied externally and taken internally, it has been considered as a specific [?], and it is a valuable remedy in most cutaneous affections that are local. Dose, ʒss to ʒij.

The fæces or dregs, left in the purification or sublimation of sulphur, are called *Horæ Brimstone*, *Dregs of Sulphur vivum*, *Sulphur caballinum* seu *vivum*, *Sulphuris vivi recemen'ta*. They are very impure, and are only used externally.

SULPHUR, VEGETABLE, *Lycopodium*—s. *Wort*, *Pucedanum*.

SULPHURATUS, *Sulphureous*, *Sulphurated*, *SULPHUREOUS*, *Sulphureus*, *Sulphuratus*, *Sulphuratus*, *Sulphuretted*. Possessing the properties of, or relating to, or containing sulphur.

SULPHURETTED, *Sulphureous*—s. Hydrogen, see Hydrogen, sulphuretted.

SULPHURETUM AMMONIACÆ, *Ammoniz sulphuretum*—s. *Ammoniz hydrogenatum*, *Liquor fumans Boylei*—s. *Kalicum*, *Potassæ sulphuretum*—s. *Lixivie*, *Potassæ sulphuretum*.

SULPHURIC ACID, *Ac'idum Sulphuricum* seu *Sulfuricum*, *Oleum Vitrioli*, *Spir'itus Vitrioli* seu *vitrioli fortis*, *Acidum vitriolicum*, *A. Vitrioli dephlogisticatum*, *Oil of Vitriol*, *Spirit of Vitriol*, *Vitriolic Acid*, *Acor Sulphuris*, (F.) *Acide Sulphurique on Sulfurique*. Sulphuric acid is inodorous; has a strong acid taste, and is corrosive. S. g. 1.845 (Ph. U. S.) It is escharotic, stimulant, and rubefacient. Mixed with lard, it is sometimes used in local pains, and certain cutaneous affections.

SULPHURIC ACID, AROMATIC, *Ac'idum Sulphuricum Aromaticum*, *A. vitriolicum aromaticum*, seu *vitriolicum alcohole aromaticum*, *Elizir vitrioli* seu *vitrioli aromaticum* seu *vitrioli Mynsichti* seu *vitrioli Edinburgensium* seu *vitrioli cum tinctura aromatica* seu *vitrioli dulcè seu acidaromaticum*, *Alcohol cum aromatibus sulphuricatus*, *Tinctura acidi sulphurici*, is formed of sulphuric acid, f ʒijss; ginger, bruised, ʒj; cinnamon, bruised, ʒiss; Alcohol, Oij.—Ph. U. S. Dose, gtt. x to xxx. Its properties are like those of the next preparation, than which it is more frequently employed, being more agreeable.

SULPHURIC ACID, DILUTED, *Ac'idum Sulphuricum Dilutum*, *A. vitriolicum dilutum*, *Elizir vitrioli*, *Spir'itus vitrioli ac'idus Vogleri*, *Elizir of Vitriol*, (F.) *Acide sulfurique délayé*, is formed, according to the Pharmacopœia of the United States, by adding aq. destillat. f ʒxij, to acid. sulph. f ʒj. It is tonic, astringent, and refrigerant. Dose, gtt. x to xx, largely diluted. When used as a gargle, half a drachm may be added to ʒvj of water.

Very dilute sulphuric acid was formerly termed *Phlegma vitrioli*.

SULPHURIS CARBURETUM, *Carbonis sulphuretum*.

SULPHURIS CHLORIDUM, *Sulphur*, *chloride of*—s. *Dichloridum*, *Sulphur*, *chloride of*.

SULPHURIS IODIDUM, *S. Ioduretum*, *Sulphur Iodatum*, *Iodure* or *Iodide of Sulphur*, (F.) *Soufre Ioduré*, *Iodure de Soufre*. It is thus directed to be prepared in the Pharmacopœia of the U. S. (*Iodin*. ʒiv; *sulphur*. ʒj.) Rub the iodine and sulphur together in a glass, porcelain, or marble mortar, until they are thoroughly mixed. Put the mixture into a matrass, close the orifice loosely, and apply a gentle heat, so as to darken the mass without melting it. When the colour has become uniformly dark throughout, increase the heat so as to melt the iodide; then incline the matrass in different directions, in order to return into the mass any portions of iodine which may have condensed on the inner surface of the vessel: lastly, allow the matrass to cool, break it, and put the iodide into bottles, which are to be well stopped.

SULPHURIUS, *Sulphureous*.

SULPHUROSUS, *Sulphureous*.

SULPHUROUS, *Sulphureous*.

SULPHUROUS ACID, *Ac'idum sulphurosicum* seu *sulphuris volat'ile* seu *Sulphurosium* seu *Sulfurosium*, *Spir'itus sulphuris* seu *sulphurosus volat'ilis* seu *sulphuris per campanam* seu *vitrioli phlogisticatus*, *Gas sulphuris*, (F.) *Acide Sulfureux*, *Air Acide Vitriolique*. Obtained by treating mercury with concentrated sulphuric acid, or by burning sulphur. It is sometimes

used in the gaseous, — at others, in the liquid, state. Sulphurous acid gas is used to disinfect clothes, confined air, letters coming from infected places, &c. It is also exhibited, with great success, in the form of fumigation, in the treatment of certain chronic cutaneous affections, and of sciatic and rheumatic pains in general. If respired pure, it acts in the manner of violent irritants; and is not long in producing death.

SULTZMALT, MINERAL WATERS OF. Sultzmalt is a village in the department of the Upper Rhine; where are several sources of cold, acidulous waters, which seem to contain carbonic acid, subcarbonate of soda, subcarbonate of lime, sulphate of lime, and a little bitumen. They are refrigerant, aperient, and alterative.

SUMAC DES CORROYEURS. *Rhus coriaria* — *s. Vénéneux*, *Rhus toxicodendron*.

SUMACH, *Rhus coriaria* — *s. Mountain*, *Rhus copallina* — *s. Poison*, *Rhus venenata* — *s. Stag's Horn*, *Rhus typhina*.

SUMBUL. The name of a root, *Sum'buli radix*, *Jatamansi*, *Musk-root*, presumed to belong to the Umbelliferæ family, much employed in Russia and Germany against cholera. Its exact botanical history is not determined. It contains a strongly odorous principle like that of musk; and is regarded as an antispasmodic.

SUMBULI RADIX, Sumbul root.

SUMEN, Hypogastrium.

SUMMER COMPLAINT. A popular name in the United States, for diarrhoea occurring in summer. It is often, also, made to include dysentery and cholera infantum. With some it means cholera infantum only.

SUMMER DISEASES, see Estival — *s. Plant winter worm*, *Sphaeria Sinensis* — *s. Rash*, *Lichen tropicus*.

SUMMITAS CUTIS, Epidermis.

SUMMUS HUMERUS, Acromion.

SUN BURN, Ephelides — *s. Dew*, *Drosera rotundifolia* — *s. Flower, false*, *Helenium autumnale* — *s. Flower, swamp*, *Helenium autumnale* — *s. Flower, tickweed*, *Coreopsis trichosperma* — *s. Flower, turpentine*, *Silphium perfoliatum* — *s. Pain*, see *Hemicrania* — *s. Rash*, *Lichen* — *s. Stroke*, *Coup de Soleil* — *s. Stroke*, *Egyptian*, *Coup de Soleil*.

SUPELLECTILE DIOGENIS, Palm.

SUPELLEX ANATOMICA, Museum anatomicum — *s. Diogenis*, Palm.

SUPER. A common Latin prefix; in composition, signifying 'above.'

SUPERBUS, Rectus superior oculi.

SUPERCHLORIDUM FORMYLICUM. Chloroform.

SUPERCILIARIS, Corrugator supercilii.

SUPERCILIARY, *Superciliaris*, from *super*, and *cilium*, 'the edge of the eyelid.' That which relates to the supercilia or eyebrows. (F.) *Sourcilier* ou *Surcilier*.

SUPERCILIARY ARCHES, *Superciliary Ridges*, *Arcus superciliar' res*, *Suggrun'dia oculo'rum*. (F.) *Arcades sourcilières*, are two slightly projecting apophyses, situate at the anterior surface of the os frontis, above the superior margin of the orbits. They are separated by the nasal *bossé*, and covered by the superciliaris muscle, to which they give attachment.

SUPERCILIARY RIDGES, Superciliary arches.

SUPERCILIUM, *Ophrys*, *Oph'rya*, *Vallum*, (F.) *Sourcil*. Same etymon. The eyebrows (Sc.) *Brees*, (Prov.) *Eye-breen*, are two arched eminences, — convex above, and more or less prominent, according to the individual, — which lie on the superciliary arch of the os frontis, above the eyelids, and extend from the sides of the root of the nose to the temple. Their inner extremity or head is more prominent than the outer or tail. The eyebrows are covered with short and stiff

hairs, which are directed, obliquely, from within outwards, and are, commonly, of the colour of the hair. The skin in which these hairs are implanted, rests on the *orbicularis palpebrarum*, *frontalis*, and *superciliaris* muscles, which move the eyebrows in different directions. The eyebrows serve as an ornament to the countenance; and by covering the eye defend it from the impression of too strong a light. They direct the perspiration also from the eye.

SUPERCILIUM ACETAB'ULLI. The edge of the cotyloid cavity of the os innominatum.

SUPERCONCEPTION, see Superfætation.

SUPEREVACUATIO, Hypercrisis.

SUPERFECUNDATION, see Superfætation.

SUPERFIBRATION, Hyperinosis.

SUPERFICIA' LIS COLLI, *Subcuta'neus Colli*.

A nerve formed by communicating branches from the second and third cervical nerves. It is distributed to the superficial parts of the neck.

SUPERFICIES ORBITALIS, see Maxillary bone (superior) — *s. Plantaris Pedis*, Sole.

SUPERFECUNDATIO, Superfætation.

SUPERFÆTATIO, Superfætation.

SUPERFÆTATION, *Superfætatio*, *Epicye'sis*, *Hypercye'sis*, *Hypercye'ma*, *Superimprægna'tio*, *Epig'one*, *Superfæcunda'tio*, *Superfecunda'tion*, *Superimprægna'tion*, from *super* and *fætare*, 'to breed.' Conception of a foetus — *epicye'ma* — in a uterus which already contains one. The impregnation of a female already pregnant. The possibility of this has been denied; but there is abundant evidence to show, that it is possible at a very early period of pregnancy. Twin cases may be of this kind of *superconception*.

SUPERGEMINALIS, Epididymis.

SUPERGENUALIS, Patella.

SUPERHUMERALE, Epomis.

SUPERIMPRÆGNATIO, Superfætation.

SUPERIOR AURIS, *Attollens aurem* — *s. Rotundus clitoridis*, *Ischio-cavernosus*.

SUPERLABIA, Clitoris.

SUPERLIGAMEN, Epidesmus.

SUPERLIGULA, Epiglottis.

SUPERNUMERY FINGER, *Digitus supernumerarius*.

SUPERNUTRITIO LIENIS, *Splenoncus* — *s. Splenis*, *Splenoncus*.

SUPEROXALAS KALICUM, Potass, oxalate of.

SUPERPURGATIO, Hypercatharsis.

SUPERSCAPULARIS, *Supraspinatus* — *s. Inferior*, *Infra-spinatus*.

SUPINATEUR COURT ou *PETIT*, *Supinator radii brevis* — *s. Grand ou long*, *Supinator radii longus*.

SUPINA'TION, *Supina'tio*, from *supinus*, 'lying on the back.' The movement in which the forearm and hand are carried outwards, so that the anterior surface of the latter becomes superior.

In Pathology, *Supination*, *Supina'tio*, *Hyptias'ma*, *Hyptias'mus*, means the horizontal position on the back, with the head thrown back, and the legs and arms extended. It is often a sign of great weakness in disease.

SUPINATOR. Same etymon. That which produces supination. A name given to two muscles of the forearm.

SUPINATOR BREVIS seu *MINOR*, *Supinator radii brevis* — *s. Longus seu major*, *Supinator radii longus*.

SUPINATOR RADII BREVIS, *S. brevis seu minor*, (F.) *Epicondyl'o-radial* (Ch.), *Supinateur court* ou *petit supinateur*. A muscle, seated at the outer and back part of the forearm. It is flattened, triangular, and curved upon itself, in such a manner that it embraces the superior extremity of the radius. It is attached, on the one hand, to the external tuberosity of the humerus, and to

a small portion of the posterior surface of the ulna, and is inserted into the upper third of the outer and upper surfaces of the radius. It produces, by contracting, the motion of supination.

SUPINATOR RADII LONGUS, *S. longus* sive *major*, *Brachio-radialis*, (F.) *Humero-radialis* (Ch.), *Long ou grand supinateur*, is seated at the anterior and outer part of the forearm. It arises from the inferior part of the outer edge of the os humeri, and from the external intermuscular aponeurosis; and is inserted, by a long tendon, into the outer edge of the radius, near the base of the styloid process. It causes the supination of the forearm, which it can likewise bend on the arm.

SUPPEDANEA, Hypopodia.

SUPPLANTALIA, Hypopodia.

SUPPLEMENTA EPIPLOICA, Appendiculæ pinguedinosæ.

SUPPLEMENTARY AIR, see Respiration.

SUPPORTER, ABDOMINAL, Belt, Russian — *s. Utero-Abdominal*, see Belt, Russian.

SUPPOSITION DE PART, Pregnancy, pre-tended.

SUPPOSITOIRE, Suppositorium.

SUPPOSITORIUM, Suppository — *s. Uterinum*, Pessary.

SUPPOSITORY, *Supposito-rium*, *Hypotheton*, *Bacillus*, *Balanus*, *Balanus*, *Glands*, *G. embiddit'ia*, *Nodulus intestinalis*, from *supponere*, *euppositum*, (*sub* and *ponere*, 'to put'), 'to put under,' (F.) *Suppositoire*. A solid medicine, in the form of a cone or cylinder, intended to be introduced into the rectum; — either for the purpose of favouring intestinal evacuations, or to act as an anodyne. Purgative suppositories are made of soap, aloes, or any other irritating substance.

SUPPRESSIO ALVI, Constipation — *s. Lochiorum*, *Ischolochia* — *s. Lotii*, *Ischuria* — *s. Mensium*, *Amenorrhœa* — *s. Menstruationis*, *Amenorrhœa* — *s. Sudoris*, *Ischidrosis*.

SUPPRESSION DU FLUX MENSTRU-EL, *Amenorrhœa* — *s. de Part*, see Parturition — *s. d'Urine*, *Ischuria*.

SUPPURANS, Suppurative.

SUPPURARE, Suppurate.

SUPPURATE, *Suppurat'è*, (F.) *Suppurer*, (Prov.) *Beal* (vulgarly). To form or secrete pus.

SUPPURATION, *Suppuratio*, *Suppuratio*, *Ecpyema*, *Purulentia*, *Pyosis*, *Pyeis*, *Diapyeima*, *Diapyeis*, *Pyogenia* simplex, *Purulence*, *Purulence*, *Gath'ering*, (F.) *Aboutissement*. Formation or secretion of pus, (see, also *Pyogenia*.) from *sub*, 'under,' and *pus*, *puris*. It is a frequent termination of inflammation, and may occur in almost any of the tissues. This termination is announced by slight chills; by remission of the pain, which, from being lancinating, becomes heavy; by a sense of weight in the part, and, when the collection of pus can be easily felt, by fluctuation. When pus is thus formed in the areolar membrane, and is collected in one or more cavities, it constitutes an *abscess*. If it be formed from a surface exposed to the air, it is an *ulcer*: and such ulcers we are in the habit of establishing artificially in certain cases of disease.

SUPPURATIVE, *Suppurans*, *Suppurativus*, *Diapyeiticus*, *Ecpyoticus*, *Epyiscon'tus*, (F.) *Suppuratif*. Same etymon. That which facilitates suppuration. Suppuratives are, generally, vesicants mixed with lard.

SUPPURER, Suppurate.

SUPRA. A common Latin prefix; in composition, signifying 'above.'

SUPRA-COSTALES, *Levato-res Costa-rum*, (F.) *Sur-costaux*, from *supra*, and *costa*, 'a rib.' A name given to twelve small, fleshy fasciculi, which are flattened, triangular, and with radiating fibres. They pass from the transverse processes of the dorsal vertebrae to the superior mar-

gin of the rib beneath. Spigelius, Cowper, Boyer, Chaussier, and others, consider them to form part of the intercostales externi.

SUPRA-HEPATIC VEINS, see Hepatic veins.

SUPRA-OMPHALODYMIA, see Somatodymia — *s. et Infra-omphalodymia*, see Somatodymia.

SUPRA-ORBITAR, *Supra-orbita'lis*, *Supra-orbitarius*, (F.) *Sus-orbitaire*, from *supra*, and *orbita*, 'the orbit.' That which is situate above the orbit.

SUPRA-ORBITAR FORA'MEN, *F. Supra-orbitarium* seu *orbitarium superius*, (F.) *Trou sus-orbitaire ou orbitaire supérieur*, is a foramen or notch completed by a ligament at the inner third of the orbital arch. It gives passage to the *supra-orbitar* or *superiiliary artery*, a branch of the ophthalmic, which ascends the forehead, and is distributed thereto.

SUPRA-PUBIAN, *Supra-pubic*, *Supra-pubianus*, from *supra*, and *pubes*. That which is seated above the pubis.

SUPRA-PUBIAN NERVE, (F.) *Nerv sus-pubien*, is the internal ramus of the inguino-cutaneous branch of the first lumbar nerve. It is the *genito-crural* branch of Bichat. It descends almost vertically before the *psaos* muscle; and, near the crural arch, divides into two branches. One of these accompanies the spermatic cord, and is distributed to the cremaster, scrotum, and integuments on the inner part of the thigh. The other issues from the pelvis with the femoral filaments; traverses the aponeurosis of the thigh, and is distributed to the integuments of the upper and anterior part of the limb.

SUPRA-RENAL, *Supra-rena'lis*, (F.) *Surrenal*, from *supra*, and *ren*, 'the kidney.' Seated above the kidney — as the *supra-renal* capsule.

SUPRA-SCAPULAR, *Supra-scapula'ris*, from *supra*, and *scapula*. Seated above the scapula; — as

SUPRA-SCAPULAR NERVE. This arises from the fifth cervical; passes through the supra-scapular notch, and is distributed to the supra-spinatus and infra-spinatus muscles.

SUPRA-SCAPULARIS, *Supra-scapular*, and *Supra-spinatus* muscle.

SUPRA-SEMIORBITULARIS, *Orbicularis* oris.

SUPRA-SPINATUS, *Supra-spino'sus*, (F.) *Sur-épineux*, *Sus-épineux*, from *supra*, and *spina*. That which is seated above the spine.

SUPRA-SPINATA FOSSA, (F.) *Fosse sus-épineuse*, is a triangular depression above the spine of the scapula.

SUPRA-SPINATUS MUSCLE, *Super-scapula'ris*, *Supra-scapularis*, (F.) *Sus-spino-scapulo-trochitérien*, *Petit sus-scapulo-trochitérien* (Ch.), is situated in the fossa supra-spinata. It is elongated, thick, and triangular, and is attached, on the one hand, to the two inner thirds of the fossa supra-spinata; and terminates, on the other, at the anterior part of the great tuberosity of the humerus. This muscle raises the arm.

SUPRA-SPINOSA LIGAMEN'TA, *Supra-spinal ligaments*, are two in number. 1. The *Dorso-lumbo-supra-spinal ligament*, (F.) *Ligament sur-épineux dorso-lombaire*, extending above the spinous processes of the dorsal and lumbar vertebrae from the 7th cervical as far as the median crista of the sacrum. 2. *Cervical-supra-spinal ligament*, (F.) *Ligament sur-épineux cervical*, which extends above all the cervical spinous processes, and is attached, above, to the outer occipital protuberance.

SUR, *Supra*. A French prefix, signifying 'above.'

SURA, *Gastrocnem'è*, *Gastrocnem'ia*, *Gastrocnemium*; *Calf of the leg*, (F.) *Mollet*, *Gras de*

Jambe. The projection, formed at the posterior part of the leg by the gastrocnemii muscle. Also, the fibula.

SURCILIER, *Corrugator supercilii*, Superciliary.

SURCOSTAUX, Supracostales.

SURCULATION, see Generation.

SURCULI FELLEI, Pori biliarii.

SURCULUS, Ramusculus.

SURDENT (F.), from *sur*, and *dens*, 'a tooth.'

When a milk-tooth does not fall out at the period of the second dentition, but is merely forced to one side by the new tooth, it is called a *surdent*.

SURDI-MUTITÉ, Mutitas surdorum.

SURDITAS, Cophosis, Deafness.

SURDITÉ, Cophosis, Deafness.

SURDOMUTITAS, Deaf-dumbness.

SURDUS, Deaf.

SUREAU COMMUN, Sambucus.

SURELLE, *Oxalis acetosella*, *Rumex acetosa*.

SURÉPINEUX, Supra-spinatus.

SUR-EXCITATION, see Excitement, and Super-excitation.

SURFEIT, Acrasia, Colica crapulosa.

SURGEON, *Chirurgæon* : *Chirō'nax*, *Chirurgus*, *Vul'nèrum medicus*, *Cheirid'èter*, *Chirid'èter*, *Iatrus*, (F.) *Chirurgien*, old French, *Surgien*, from *χυρ*, 'the hand,' and *εργον*, 'work.' One who practises surgery. The surgeon, says Celsus, ought to be young, or, at all events, not advanced in life. His hand must be steady, expert, and never tremulous : he must use the right or left with equal facility : his sight must be clear and penetrating : his mind intrepid ; and, although unmoved when he proceeds to operate, he must not be in haste, nor cut less than is necessary, but finish his operation, as if the cries of the patient made no impression upon him.

In the middle ages, the Universities of Italy and Spain sent out educated surgeons, as well as physicians, under the name *chirurgi-physici*.

SURGEON-APOTHECARY, One who unites the practice of surgery with that of the apothecary. A general practitioner. He is the ordinary medical man or family medical attendant in England. See Apothecary.

SURGEON-DENTIST, Dentist.

SUR-GEONCY, The office of a surgeon.

SUR'GERY, *Chirurg'ia*, *Chiriatri'a*, *Cheiriz'is*, *Chi'risis*, *Chiriz'is*, *Chirid'mus*, *Metacheiriz'is*, *Metach'risis*, *Metachiriz'mus*, *Tractat'io manu'al'ia*, *Medic'ina operat'iva seu eff'icax*, (F.) *Chirurgie*. Same etymon as surgeon. The part of the healing art which relates to external diseases ; their treatment ; and, especially, to the manual operations adapted for their cure. Also, the office of a surgeon.

SURGERY, ANAPLASTIC, Morioplastice — s. Autoplastic, Morioplastice — s. Conservative, Surgery, salvation — s. Dental, Dentistry.

SURGERY, MILITARY, *Chirurg'ia militaris*, (F.) *Chirurgie militaire*, is that which relates to the operations to be performed on the field of battle, or subsequently ; — with the attention to be paid to the wounded, &c.

SURGERY, OPERATIVE, is termed *Acir'gia*, *Acir'gia*, and *Acidur'gia*, (F.) *Médecine opératoire*.

SURGERY, PLASTIC, Morioplastice.

SURGERY, SALVATION, *Conser'vative Surgery*, is that which looks rather to the preservation of a limb for example than its removal. One, who is addicted to operating in almost all cases, is sometimes called a *knifeman*.

SUR'GICAL, *Chirurg'icus*, (F.) *Chirurgical*, *Chirurgique*. Same etymon. That which relates to surgery ; as surgical anatomy, surgical operations, surgical instruments, &c.

SURGIEN, Surgeon.

SURINAMINE, see *Geoffræa Surinamensis*.

SUR-IRRITATION, see Irritation.

SURNUMÉRAIRE (Doigt), Digitus supernumerarius.

SUROXYGÉNÈSES. From *sur* and *oxygen*. A name given by Baumes to diseases attributed by him to superabundance of oxygen.

SURPEAU, Epidermis.

SURRECTA ALBA, *Ranunculus flammula*.

SUR-RÉNAL, Supra-renal.

SURROGATUM, *Succedaneum*.

SURSELS, see Salt.

SURTOUTS LIGAMENTEUX DE LA COLONNE VERTÉBRALE, Vertebral ligaments.

SURUMPE (S.) A disease to which the traveller in the Cordilleras is liable. It consists of a violent inflammation of the eyes, caused by the reflection of the bright rays of the sun. The conjunctiva is highly inflamed, and the lids swell and bleed. The pain is intense, and frequently induces delirium. — Tschudi.

SURVIE, Survivorship.

SURVIVORSHIP, (F.) *Survie*, from *super-vivere*, (*super*, and *vivere*), (F.) *survivre*, 'to live longer.' In legal medicine, this word means the probability that any individual has survived others in an accident, which has been fatal to all. This is often an important question, connected with the inheritance of property, and on which physiology can rarely afford more than presumptive grounds for conclusion.

SUS, SUSUM, (F.) *Sus*, same as *Sursum*, 'above.' A French prefix, signifying 'above,' as in

SUS-ACROMIO-HUMÉRAL, Deltoid.

SUS-CARPIEN ; *sus*, and *carpe*, 'the wrist.' Above the wrist. An epithet given by Chaussier to the dorsal artery of the wrist — *Artère sus-carpienne* — a branch of the radial.

SUS-ÉPINEUX, Supra-spinatus — s. *Maxillo-labial*, Levator anguli oris — s. *Maxillo-labial*, grand, Levator labii superioris alaeque nasi — s. *Maxillo-labial moyen*, Levator labii superioris proprius — s. *Maxillo-nasal*, Compressor naris — s. *Omphalodymie*, see *Somatodymia* — s. *Optico-phéno-scléroticien*, Rectus superior oculi — s. *Orbitaire*, Supra-orbital — s. *Phalangien du ponce*, Abductor pollicis brevis — s. *Pubio-fémoral*, Pectinalis — s. *Scapulo-trochitérien le plus petit*, Teres minor — s. *Scapulo-trochitérien petit*, Supra-spinatus — s. *Scapulo-scapulo-trochitérien*, Supra-spinatus.

SUSCEPTIBIL'ITY, *facile suscipiens*, (*sus*, and *cipio*, 'I take,') 'easily taking, or undertaking.' Sensibility, in its most enlarged acceptation. Impressibility. This term, with the epithet *nervous*, is sometimes used for that increase in the moral sensibility which is particularly observed in nervous affections.

SUSCEPTIO INTESTINORUM, Intussusception.

SUSPENDICULUM, Cremaster.

SUSPENSÀ, Enæorema.

SUSPENSEUR, Suspensory.

SUSPENSOIRE, Suspensory.

SUSPENSORIUM, Suspensory — s. Hepatic, Suspensory ligament of the liver — s. Testis, Cremaster.

SUSPEN'SORY, *Suspensio'rius*, from *suspenderè*, (*sus*, and *pendere*, 'to hang,') 'to hang up.' (F.) *Suspenseur*. That which sustains or suspends.

SUSPENSORY BANDAGE, *Suspensio'rium*, *Suspensoir* ou *Suspensoire*, is a bandage intended to support the scrotum, in cases of diseases of the testicle or of scrotal hernia. It consists of a kind of bag, fixed above to a cincture, and retained

below by strings passing between the thighs, and fixed to the cincture behind.

SUSPENSORY LIGAMENT OF THE LIVER, *Ligamentum latum, Suspensorium hepatis, Perpendiculum hepatis*, is a triangular duplicature of the peritoneum, between the inferior surface of the diaphragm and the upper surface of the liver, which is continuous with the great falc of the umbilical vein.

SUSPENSORY LIGAMENT OF THE PENIS is a fibro-cellular fascia, of a triangular shape, which extends from the anterior part of the symphysis pubis to the corpus cavernosum, which it sustains. A similar ligament belongs to the clitoris.

SUSPENSUM, Enæorema—s. *Urinæ, Enæorema*.

SUSPIRATION, Suspirium.

SUSPIRATUS, Suspirium.

SUSPIRITUS, Suspirium.

SUSPIRITUM, Suspiratio, Suspiratus, Suspiritus, from *sus*, and *spiro*, 'I breathe'; *Myrmus, Mychmus, Mychthismus, Stenagmus, Stenax's, Sigh, Suspiration, (F.) Soupir*. A short breathing. A slow, voluntary contraction of the diaphragm and intercostal muscles, the effect of which is to restore the due relation between the circulation and respiration, or to free ourselves from an inconvenient sense of weight felt in the chest, when we are much depressed, — a weight, which appears to be chiefly dependent on the functions of the heart being disturbed through the moral influence. The sigh differs from the sob: — the latter being involuntary and spasmodic. Sighing, occurring in fevers to a great extent, indicates considerable depression. When accompanied with a vocal sound, it constitutes a groan.

SUSPIRITUM CARDIACUM, Angina pectoris.

SUSTENTACULUM, Aliment.

SUSTENTATOR CLITORIDIS, Erector clitoridis — s. *Penis, Erector penis.*

SUSURATION, Whispering.

SUSURRUS, Bourdonnement — s. *Aurium, Tinnitus aurium.*

SUTELA, Suture.

SUTERBERRY, Xanthoxylum fraxineum.

SUTORIUS, Sartorius.

SUTTER, Saccharum.

SUTURA ABDOMINALIS, Gastroraphy — s. *Arcaalis, Coronal suture* — s. *Biparietalis, Sagittal suture* — s. *Clavata, Suture, quilled* — s. *Corporis callosi externa, see Raphe* — s. *Cranii squamosa, Squamous suture* — s. *Dentata, see Suture* — s. *Frontoparietalis, Coronal suture* — s. *Interparietalis, Sagittal suture* — s. *Jugalis, Sagittal suture* — s. *Lambdoidalis, Lambdoid suture* — s. *Limbosa, see Suture* — s. *Lepidoides, Squamous suture* — s. *Mendosa, Squamous suture* — s. *Notha, see Suture* — s. *Oblelea, Sagittal suture* — s. *Ossium Spuria, Harmony* — s. *Proræ, Lambdoid suture* — s. *Punctata, Suture, common* — s. *Puppis, Coronal suture* — s. *Rhabdoides, Sagittal suture* — s. *Serrata, see Suture, Sagittal suture* — s. *Spuria, see Suture* — s. *Squamosa, see Suture* — s. *Vera, see Suture.*

SUTURAL, Sutura'tia; same etymon as the next. Relating to a suture.

SUTURE, Sutura, Sute'la, Armè, Rhapha, from suere, sutum, 'to stitch': Dove-tail joint, (F.) Engrenure. A kind of immovable articulation, in which the bones unite by means of serrated edges, which are, as it were, *dove-tailed* into each other. The articulations of the greater part of the bones of the skull are of this kind. The coronal, lambdoidal, and sagittal sutures are called *true sutures, sutura vera*, the two temporal are called *false sutures, sutura notha seu spuria seu squamosa*.

A *dentated suture* — *Sutura dentata* — is one in which the processes are long and dentiform,

as in the interparietal suture. In a *serrated suture* — *sutura seu gyata's serrata* — the indentations and processes are small and fine, like the teeth of a saw, as in the suture between the two portions of the frontal bone. In the *sutura limbo'sa* there is along with the dentated margins a degree of bevelling of one, so that one bone rests on the other, as in the occipito-parietal suture.

SUTURE, in Surgery, Rha'phè. means an operation which consists in stitching the lips of a wound to procure their union. The suture was, at one time, largely employed; but, in modern times, its use has been wisely restricted. There are few wounds in which the edges may not be brought together by adhesive plaster appropriately applied. This kind of union, by adhesive plaster simply, has been absurdly termed *Sutura sicca* or *dry suture*, in opposition to the *Sutura cruenta* or suture properly so called. The suture is generally performed with waxed thread and straight or curved needles, according to the arrangement of parts.

The chief sutures are the *interrupted suture*, the *quilled suture*, the *glover's suture*, and the *twisted suture*. 1. The **INTERRUPTED SUTURE, (F.) Suture à points séparés, Suture entrecoûpée**, is performed by passing a needle, armed with a ligature, through the lips of the wound previously brought into contact; and then tying the extremities of the thread. The other stitches are made in the same manner. They are separate or interrupted. It is the suture most commonly used. 2. The **QUILLED SUTURE, Sutura clavata, (F.) Suture enchevillée ou emplumée ou empenée, Compound Suture**, is merely the interrupted suture, with this difference, that the ligatures are not tied over the face of the wound, but over two quills or rolls of plaster, or bougies, which are laid along the sides of the wound. It is used by some surgeons, on account of there being less danger of the stitches tearing out. It is not much employed. The **CLAMP SUTURE**, used by Dr. Sims, of New York, in the treatment of vesico-vaginal fistula, is a modification of this. It is formed by two cylinders of silver or lead, perforated at several points for the passage of small pieces of fine silver wire, which are to supply the places of thread, and are prevented from slipping by perforated shot carried down upon them, pressed against the cylinders, and kept in situ by being firmly pinched with pliers. 3. The **GLOVER'S SUTURE, Contin'ued or Uninterrupted Suture, (F.) Suture de Pelletier, ou en surget**, is executed by introducing the needle first into one lip of the wound from within outwards, then into the other in the same way; and, in this manner, sewing up the whole extent of the wound. It has been chiefly used in wounds of the intestinal canal. It is now rarely employed except for stitching up dead bodies. The common stitch or suture, *Sutura punctata, (F.) Suture à points passés*, has been used in similar cases. 4. The **ROYAL STITCH or SUTURE** is the name of an old operation for the cure of *lumbocele*. It consisted in putting a ligature under the hernial sac, close to the abdominal ring, and then tying that part of the sac, so as to render it impervious by the adhesive inflammation thus excited. 5. The **TWISTED SUTURE, (F.) Suture entortillée**, is employed, chiefly, to unite wounds of the cheeks and of the free edges of the lips. To perform it, a pin or needle is passed through the edges of the wound, so as to keep them accurately in contact. A piece of thread is then wound accurately around the pin, from one side of the division to the other, in the form of the figure ∞ . The number of pins and sutures must, of course, correspond with the extent of the

wound. This is the suture used in cases of hare-lip. The **BUTTON SUTURE**, of Dr. Bozeman, of Alabama, employed in vesico-vaginal fistula, is a modification of this. 6. The **SUTURE À ANSE**, of Le Dran, has only been employed in cases of wounds of the intestines. For this purpose, the edges of the wound are brought together, and as many needles, with unwaxed threads, used as there are to be stitches. Each stitch is then made by passing a needle through the edges; and all the ligatures which belong to one side of the wound are tied together, and twisted so as to form a thread, which is fixed externally. 7. The **IMPLANTED SUTURE**, (F.) *Suture implantée*, is formed by pins arranged parallel to the edges of the wound, and implanted in their substance. By means of threads, the needles are approximated so as to bring the edges of the wound together. 8. The **PLASTIC SUTURE**, so called by Prof. Pancoast, of Philadelphia, its proposer, is a suture by tongue and groove. The groove is made by an incision in the part, which is to receive the flap. The tongue is made on the flap by bevelling off the two surfaces, cuticular and adipose. The suture is made by passing a loop of thread, by two punctures, deep through the inner lip of the groove. The two needles are passed through the base of the tongue, and again through the outer lip of the groove. The two ends of the thread are then tied over a small roll of adhesive plaster, by which the tongue is, of necessity, forced deep into the groove. Four raw surfaces are thus firmly held in connection, making the union of parts, by this suture, a very certain process.

SUTURE, **BUTTON**, see **SUTURE**, (twisted) — *s.* **CLAMP**, see **SUTURE**, (quilled) — *s.* **COMPOUND SUTURE**, quilled — *s.* **CONTINUED SUTURE**, glover's — *s.* *Écaillée*, Squamous suture — *s.* *Empenée*, Suture, quilled — *s.* *Emplumée*, Suture, quilled — *s.* *Enchevillée*, Suture, quilled — *s.* *Entortillée*, Suture, twisted — *s.* *Entrecoupée*, Suture, interrupted — *s.* **FALSE HARMONY**, see **SUTURE** — *s.* *Implantée*, see **SUTURE**, (implanted) — *s.* *Occipitale*, Lambdoid suture — *s.* *Occipito-parietal*, Lambdoid suture — *s.* *de Pelletier*, Suture, glover's — *s.* *Périnéale*, Perineorrhaphy — *s.* **PLASTIC**, see **SUTURE** — *s.* *à Points passés*, Suture, common — *s.* *à Points séparés*, Suture, interrupted — *s.* **SAGITTAL**, see **SAGITTAL SUTURE** — *s.* *Squamense*, Squamous suture — *s.* *en Surget*, Suture, glover's — *s.* **SUPERFICIAL HARMONY** — *s.* **UNINTERRUPTED**, see **SUTURE**, glover's.

SWAB, from Sax. *ƿrebban*, 'to clean with a mop.' A piece of sponge or rag tied to a rod of whalebone or wood, for cleansing the mouth of the sick, or applying remedial agents to deep-seated parts. The process is termed *Swabbing*.

SWADDLING CLOTHES, Swathing clothes.

SWAGBELLY, Physconia.

SWALLOW, Hirundo — *s.* **WORT**, *Asclepias vincetoxicum* — *s.* **WORT**, orange, *Asclepias tuberosa* — *s.* **WORT**, tuberous-rooted, *Asclepias tuberosa* — *s.* **WORT**, white, *Asclepias vincetoxicum*.

SWALLOW'S NEST, *Nidus hirundinis*.

SWALME, Excrescence, Tumour.

SWAMP CABBAGE, *Dracontium foetidum* — *s.* **DOGWOOD**, *Petala trifoliata* — *s.* **SICKNESS**, Milk sickness — *s.* **WOOD**, *Cephalanthus occidentalis*, *Dicaea palustris*.

SWARFF, Stupor, Syncope.

SWATHING CLOTHES, *Swaddling Clothes*, *Incunabula*, *Fasciæ*, *Panni*, from Sax. *ƿreowan*, 'to bind.' (F.) *Maillet*. The kind of clothing or bandaging formerly applied to children during the first year of life. It is now, properly, laid aside, and every freedom allowed to muscular exertion.

SWEAT, **SUDOR** — *s.* **BLOODY**, *Sudor cruentus* —

s. **GLANDS**, see **PERSPIRATION** — *s.* **ROOT**, *Polemonium reptans* — *s.* **SCAB**, *Crusta genu equinae*.

SWEATING, *Epidrosis* — *s.* **BATH**, *Achilolum* — *s.* **OF BLOOD**, see **DIAPEDESIS**, and *Sudor cruentus* — *s.* **HOUSE**, *Achilolum* — *s.* **PLANT**, *Eupatorium perfoliatum* — *s.* **ROOM**, *Achilolum* — *s.* **SICKNESS**, *Sudor Anglicus*.

SWEATING SICKNESS, **MALWAH**. A disease which appears to be allied to the worst form of cholera, and to bear a close relation to malignant congestive fever. As its name imports, it occurs in Malwah in India.

SWEBBING, Syncope.

SWEE, Vertigo.

SWEEMING, Syncope.

SWEEMISH, see Syncope.

SWEET BITTER, *Triosteum* — *s.* **BREAD**, *Pancreas*, *Thymus* — *s.* **BUSH**, *Comptonia asplenifolia* — *s.* **FERN**, *Comptonia asplenifolia* — *s.* **FLAG**, *Acorus calamus* — *s.* **POD**, *Ceratonium siliqua* — *s.* **ROOT**, *Aralia nudicaulis*.

SWELLING, *Epanastema*, *Exaroma*, *Excedosis*, *Presma*, *Oncus*, *Extumefactio*, *Tumor*, *Tumefactio*, (Prov.) *Blowing*, *Boist*, *Coul*, (Old Eng.) *Boche*, *Bollynge*, *Bolning*, (F.) *Enflure*, *Gonflement*, from Sax. *ƿrellan*, (D.) *swellen* (G.) *schwellen*, 'to swell,' Morbid increase of bulk in the whole, or a part, of the body. See **Tumour**.

SWELLING, **WHITE**, *Hydrarthrus*.

SWENSIE, Cynanche.

SWENYNG, Somnium.

SWERF, Stupor, Syncope.

SWERTIA CHIRAYITA, *Gentiana chirayita* — *s.* **DIFORMIS**, see *Calumba* — *s.* **FRASERA**, see *Calumba* — *s.* **LUTEA**, *Gentiana lutea*.

SWEUIN, Somnium.

SWEVYNG, Somnium.

SWEVYNYNG, Somnium.

SWIETENIA FEBRIFUGA, after Van Swieten, *S. So'nida*, *Soym'ida* seu *Cedre'la febrifuga*, *Febrifuge Swietenia*, (F.) *Mahogan febrifuge*. *Ord.* *Meliaceæ*. *Secz. Syst.* *Decandria Monogynia*. The bark of this tree, which is a native of the East Indies, called there *Rohena Bark*, has an aromatic odour and bitter taste. It is tonic and stomachic, and has been used in intermittents, and in diseases where bitter tonics are indicated. Dose, ʒj to ʒij of the powder.

SWIETENIA MAHOGANI, *Cedrus Mahogani*, *Mahogany Tree*, grows in the West Indies and tropical America, and is possessed of similar properties.

SWIETENIA SENEGALENSIS, *Khaya Senegalen'sis*. A common forest tree on the borders of the Gambia. The bark called *Cail-cedra*, and *Karson Khayi*, is bitter and antiperiodic.

SWIETENIA SOMIDA, *S. Febrifuga*.

SWIM, Air bladder.

SWIMBLADDER, Air bladder.

SWIMMING, *Natation* — *s.* **BLADDER**, Air bladder — *s.* **OF THE HEAD**, Vertigo.

SWINACIE, Cynanche tonsillarlis.

SWINE-POX, see *Varicella*.

SWINE-SAME, *Adeps*.

SWINE-THISTLE, *Sonchus*.

SWINNYING, Vertigo.

SWOON, Syncope.

SWOONING, Syncope.

SWOUNDING, Syncope.

SYALADENTITIS, *Inflammatio glandularum salivarium*, from *σάλων*, 'saliva,' *adn̄*, 'a gland,' and *itis*, denoting inflammation. Inflammation of the salivary glands.

SYCAMINOS, *Morus nigra*.

SYCAMINUM, see *Morus nigra*.

SYCE, *Ficus*, *Ficus carica*.

SYCEA, *Ficus*.

SYCEPHALUS, from *σύν*, 'with,' 'together,'

and κεφαλή, 'head.' A monster having two heads united together.

SY'CION, σικιον, from σικον, 'a fig.' A decoction of dried figs.

SY'CIOS ANGULA'TUS. One-seeded star cucumber, *Wild Bry'ony*. The roots and seeds of this indigenous plant, Order, Cucurbitaceæ, are bitter, purgative and diuretic. It is used in dropsy in the same cases as bryony.

SYCI'TES, from σικον, 'a fig.' Wine, in which figs have been infused.

SYCITES, Catorchites.

SYCOMA, Ficus.

SYCO'SIS. Same etymon. A tumour of the shape of a fig. Also, a fungous ulcer. A horny excrescence about the eyelids. Dr. Bateman defines it:—an eruption of inflamed but not very hard tubercles occurring on the bearded portion of the face and on the scalp, in adults; and usually clustering together in irregular patches;—*Phyma Sycosis, Rose'ola fice'la, Aene men'ta-gra*. Of this he has given two species: 1. *Sycosis Menti, S. barbæ, Men'tagra, Menti'go, Varus Men'tagra, Mentagraph'ya, Chinwelk, Chinwelk, Barber's Itch, (F.) Dartre pustulose mentagre*: and, 2. *Sycosis Capit'li, S. Capit'li'ti*. The treatment, in both cases, resembles that for porrigo favosa.

SYCOSIS, Ficus.

SYCUM, Ficus.

SYDERATIO, Sideratio.

SYLLEPSIS, Conception.

SYLLEPSIOLOG'IA, from συλληψις, 'conception,' and λογος, 'a description.' The doctrine of conception.

SYM, SYN, συν, 'with, together,' like the *cum*, *con*, *col*, *cor*, &c., of the Latins. A common prefix. Before *b*, *p*, *ph*, *ps*, and *m*,—*sym*, or *syn*, is changed into *sym* or *syn*, (*com*;) before *c*, *ch*, *g*, *k*, and *x*, into *syn* or *syn*, *συγ*, *ξυγ*, (*con*;) before *l* into *eyl* or *xyl*, (*col*;) before *r* into *eyr* or *zyr* (*cor*;) and before *s* into *sy* or *xy*, and *sys*, (*co* and *cos*.)

SYMBLEPHARON, Ankyloblepharon.

SYMBLEPHARO'SIS, *Concre'tio palpebra-rum cum bulbo oc'uli*, from *sym*, and βλεφαρον, 'an eyelid.' A preternatural adhesion between the eyelids; and especially between the upper eyelid and globe of the eye.

SYMBLEPHAROSIS, Ankyloblepharon.

SYMBOL, *Sym'bolum, Character, (F.) Carac-tère*, from συμβάλλω (*syn*, and βάλλω, 'I throw'), 'I compare.' A sign or representation of something else.

The following are the symbols used in prescriptions:

R, Recipe. See R.

gr., *Grammum*, a grain.

ʒ, *Scrupulus seu scrupulum*, a scruple.

ʒ, *Drachma*, a drachm.

ʒ, *Uncia*, an ounce.

lb, *Libra*, a pound.

lb, *Libra*, pounds.

ʒ, *Minimam*, a minim.

fʒ, *Fluidrachma*, a fluidrachm.

fʒ, *Fluiduncia*, a fluidounce.

O, *Octarius*, a pint.

C, *Congius*, a gallon. See Abbreviation.

SYMBOLE, Commissure.

SYMBOLEUSIS, Consultation.

SYMBOLISMUS, Sympathy.

SYMBOLIZATIO, Sympathy.

SYMBOLICA, Symptomatology.

SYMBOLUM, Sympathy.

SYMÉTRIE, Symmetry.

SYMÉTRIQUE, Symmetrical.

SYM'MELES, from *sym*, and μέλος, 'an extremity.' *Monoporus*. A genus of monsters whose extremities are joined together: *Sirens*.—I. G. St. Hilaire.

SYMMETRIA, Symmetry.

SYMMETRICAL, *Symmet'ricus, συμμετρικός, (F.) Symétrique*, from *sym*, and μετρεῖν, 'to measure.' An epithet given to those parts of the body, which, if seated on the median line, may be divided into two equal, and perfectly like halves; or which—if situate, the one to the right, the other to the left of this line—have a similar conformation, and a perfectly analogous arrangement.

SYMMETRICAL DISEASES, see Symmetry.

SYMMETRY, *Symmet'ria, Commen'sum, Membro'rum Omnium Competen'tia, Proportio, (F.) Symétrie*. The regularity in form, observed in many parts, situate on the median line; and, also, the resemblance which exists between many organs that are situate on each side of the median line; *Bilatéral Symmetry*. Bichat first noticed that this symmetry was confined to the organs of animal life; as the head, brain, spine, organs of sense, locomotion, &c.

It has been affirmed—and there is evidence in favour of the assertion—to be a law of the animal economy, that when uninfluenced by disturbing agents, all general or constitutional diseases affect equally and similarly the corresponding parts of the two sides of the body. This has been called the *Symmetry of Diseases*; and the resulting diseases are said to be *symmetrical*.

SYMMETRY OF DISEASES, see Symmetry.

SYMPARATERESIS, Observation.

SYMPASMA, Catapasma.

SYMPATHESIS, Sympathy.

SYMPATHETIC, *Sympath'icus, Sympatheti-cus, (F.) Sympathique*, from *sym*, and πασος, 'suffering.' That which depends on sympathy. Sympathetic affections of an organ are those morbid phenomena that supervene without any morbid cause acting directly on it, and by the reaction of another organ primarily affected. Thus, itching of the nose is a *sympathetic affection*, produced by irritation in the intestinal canal.

The epithet *sympathetic* is, also, given to different nerves. See Trisplanchnic, Pneumogastric, and Facial.

SYMPATHETIC, GREAT, Trisplanchnic nerve—s. Middle, Pneumogastric—s. Nerve, Trisplanchnic nerve.

SYMPATHETIC POWDER, *Pulvis sympathet'icus* or *Powder of Sympathy* of Sir Kenelm Digby, was composed of calcined sulphate of iron, prepared in a particular manner. It was long supposed to be able to cure a wound, if applied to the weapon that inflicted it; or even to a portion of the bloody clothes. It is proper to remark, however, that the wound itself was carefully excluded from the air, and from all irritation.

SYMPATHETICUS MINOR, Facial nerve.

SYMPATHETISMUS, Sympathy.

SYMPATHIA, Consent of parts.

SYMPATHICUS, Sympathetic.

SYMPATHIQUE, Sympathetic — s. *Nerf*, Trisplanchnic nerve.

SYMPATHY, *Sympathi'a, Sympathe'is, Compas'sio, Consensus, Sympathetis'mus, Xympathi'a, Conflux'io, Conspira'tio, Commer'cium, Consor'tium, Homoiopathi'a, Homœoth'nia, Adelphix'is, Symbolis'mus, Symboliza'tio, Sym'bolum*. Same etymon. The connexion that exists between the action of two or more organs, more or less distant from each other; so that the affection of the first is transmitted, secondarily, to the others, or to one of the others, by means unknown. A knowledge of the particular sympathies between

different organs throws light on the etiology of diseases, their seat, and the particular organs towards which our therapeutical means should be directed. See Irritation.

SYMPEPSIS, Coction.

SYMPEXIS, Concretion.

SYMPHORA, Congestion.

SYMPHOREMA, Congestion.

SYMPHORESIS, Congestion.

SYMPHORICAR'PUS RACEMOSUS, *Snowberry*; from *συμφορεω*, 'to bear together,' and *καρπος*, 'fruit,' owing to the clustered berries. An indigenous plant, *Order* Caprifoliaceæ, *Tribe* Loniceræ, whose berries are large, bright white, and remain till winter; flowering from June to September. The root is tonic and astringent, and has been used in ague; and by the Indians in syphilis.

SYMPHYIA, Symphysis.

SYMPHYOCEPH'ALUS, *Metop'ages*, *Cephalop'ages*, from *συνφυς*, (*συν*, and *φυω*, 'to grow,') 'united together,' and *κεφαλη*, 'head.' A monstrosity in which twins are united by the head.

SYMPHYSEOT'OMY, *Symphysiot'omy*, *Synchondrot'omy*, *Symphyseotom'ia*, *Symphysiotom'ia*, from *συνφυσις*, and *τομω*, 'I cut.' The operation or section of the *Symphysis*, *Sigaultian operation* or *section*. This operation is performed with the view of augmenting the diameter of the pelvis, in order to facilitate delivery in certain cases of faulty conformation of that cavity, with wedging of the head of the child therein. To perform the operation, (F.) *désymphyser*, the female is placed on her back; the pubes is shaved; and, with a strong scalpel, the soft parts are divided on the median line. The articulation is then cut. As soon as the fibrous parts, which hold the bones together, are divided, they separate to a certain distance from each other; and the delivery may be accomplished. The knife, with which the operation is performed, is called, by the French, *Couteau symphysien*.

SYMPHYS'IA, from *συνφυω*, 'to grow together.' A malformation consisting in the union of parts that are usually divided. — Breschet.

SYMPHYSIOTOMIA, *Symphyseotomy*.

SYMPHYSIS, *Symphy'ia*, *Coalescen'tia*, *Coal'itus*, *Coalit'io*, (F.) *Symphysie*, from *συνφυω*, (*συν*, 'with,' and *φυω*, 'I grow'), 'I unite together.' A union of bones. The bond of such union. The aggregate of means used for retaining bones *in situ* in the articulations. The name *symphysis* has, however, been more particularly appropriated to certain articulations; as to the *symphysis pubis*, *sacro-iliac symphysis*, &c. See Monster.

SYMPHYSIS, Coalescence, Commissure, Insertion — *s.* Cartilaginea, *Synchondrosis* — *s.* Ligamentosa, *Syndesmosis* — *s.* Ossium muscularis, *Syssarcosis* — *s.* Pubis, *Pubic articulation*.

SYMPHYSOTOMY, *Symphyseotomy*.

SYMPHYTOS, Hereditary.

SYMPHYTUM. Same etymon. *Symphytum officinale* seu *album* seu *Bohemicum*, *Consol'ida major*, *Comfrey*, *Alum*, *Alus*, (Prov.) *Knit-back*, (F.) *Grande Consoude*, *Oreilles d'âne*. *Ord.* Boraginaceæ. *Sex. Syst.* Pentandria Monogynia. The roots of this European plant are very mucilaginous, and are employed as emollients. They are used in decoction in hæmoptysis, dysentery, chronic diarrhœa, &c.

SYMPHYTUM, *Solidago virgaurea* — *s.* *Album*, *Symphytum* — *s.* *Bohemicum*, *Symphytum* — *s.* *Minimum*, *Bellis* — *s.* *Minus*, *Prunella* — *s.* *Officinale*, *Symphytum*.

SYMPHYTUM PETRÆ'UM, *Coris Monapeliensis*, *Montpellier Coria*. An intensely bitter and nauseous plant, which is said to have been useful in syphilis. See, also, *Sanicula*.

SYMPLOCARP'US ANGUSTISPATHA, *Dracontium angustispatha* — *s.* *Fœtidus*, *Dracontium fœtidum*.

SYMPOD'IA, *Monopod'ia*, from *sym*, and *πους*, *podos*, 'a foot.' A monstrosity in which there is coalescence of the lower extremities.

SYMPTOM, *Sympto'ma*, *Ac'cidens*, *Casus*, from *sym*, and *πτωω*, 'a fall,' from *πτωω*, 'I fall.' Any change, perceptible to the senses in any organ or function, which is connected with morbid influence. It is by the aggregate and succession of symptoms that a disease is detected. The term *symptoms of symptoms* has been used for the effects which result from the symptoms of a disease; but which effects are not essentially connected with the disease itself. Thus, the debility arising from the frequency of evacuations in dysentery is a symptom of symptoms. *Symptom*, at one time, was generally used in the same sense as *sign*; but with many, perhaps most, of the present day, the former signifies a functional or vital phenomenon of disease, whilst the latter is applied to that which is more directly physical; — and hence the expressions *functional or vital phenomena or symptoms*, in contradistinction to the *physical signs* afforded by auscultation, percussion, &c. See *Sign*.

SYMPTOMA TURPIDUDINIS, *Nymphomania*.

SYMPTOMAT'IC, *Symptomat'icus*, (F.) *Symptomatique*. That which is a symptom of some other affection. Thus we speak of a symptomatic disease, in opposition to one that is idiopathic.

SYMPTOMATOGRAPHY, *Semeiography*.

SYMPTOMATOL'OGY, *Symptomatology'ia*, *Symbolog'ica*, from *symptoma*, and *λογος*, 'a treatise.' The branch of pathology, the object of which is the knowledge of symptoms.

SYMPTO'SIS, from *sym*, and *πτωω*, 'I fall.' Depression or collapse, emaciation or atrophy, either of the whole body or of a part.

SYN, *Sym*.

SYNEREMA, *Corrugation*.

SYNETHION, *Concausa*.

SYNANCHE, *Cynanche*, *Cynanche tonsillar*.

SYNANCIE, *Cynanche*.

SYNAPHYMENTITIS, see *Ophthalmia*.

SYNAPTASE, see *Amygdalin*.

SYNARTHRODIA, *Synarthrosis*.

SYNARTHRO'DIAL, *Synarthrodial'ia*, from *syn*, and *ἄρθρωσις*, 'articulation.' That which belongs to synarthrosis; as *Synarthrodial articulation*.

SYNARTHRO'SIS, *Synarthro'dia*, *Abarticulatio*, *Coarticulatio*. An immovable articulation. *Synarthrosis* includes three species: *Suture*, *Harmony*, and *Gomphosis*.

SYNATHROESIS, *Congestion*.

SYNATHROISMUS, *Congestion*.

SYNCANTHUS, from *syn*, and *κανθος*, 'the corner of the eye.' A morbid adhesion between the globe of the eye and the orbit.

SYNCHONDRO'SIS, *Sym'physis seu Cop'ula seu Connea'io cartilagin'ea*, *Chondrosyndes'mus*, from *syn*, and *χονδρος*, 'a cartilage.' Union or articulation of bones by means of cartilage: — as the articulations of the ribs with the sternum, by means of their cartilages of prolongation.

SYNCHONDROTOMY, *Symphyseotomy*.

SYNCHRONIUS, *Synchronous*.

SYN'CHRONOUS, *Syn'chronus*, *Synchron'ius*, *Isoch'ronous*, from *syn*, and *χρονος*, 'time.' That which occurs at the same time. Thus the pulsations of an artery are said to be synchronous with those of the heart.

SYNCHRONUS, *Synchronous*.

SYNCHYSIS, *Confusio* — *s.* *Sparkling*, *Confusio*.

SYNCIPUT, *Sineiput*.

SYNCLONESIS, Synclonus.

SYNCLONUS, *Synclonus'sis*, from *συγκλονω*, (*συν*, and *κλονω*, 'I shake,') 'I shake together.' Tremulous, simultaneous, and chronic agitation of various muscles, especially when excited by the will; *Clonici universales*, *Clonus*, (Young.) Also, general spasm, *epasmus universalis*. See Tremor, Chorea, &c.

SYNCLONUS BALLISMUS, Paralysis agitans — s. Beriberia, Beriberi — s. Chorea, Chorea — s. Raphania, Raphania — s. Saltans, Chorea — s. Tremor, Tremor.

SYNCOMISTUS, from *συγκομιζω*, (*συν*, and *κομιζω*), 'I bring together.' *Autop'gyros*, *Panis domesticus*, *secundarius*, *cibarius*, *confusus*. Bread of the whole wheat, or of the unsifted meal. *Household bread*. Also a cataplasm made of the same.

SYNCOPAL, *Syncopalis*, *Syncope'sis*, *Syncope'tus*. Same etymon as Syncope; as *Syn'copal Fever*, *Febris Syncope'alis*. A pernicious intermittent, characterized by repeated attacks of syncope.

SYNCOPE, from *συκοπω*, (*συν*, and *κοπω*, 'to fall,') 'I fall down.' *An'imi defectio seu deliquium*, *Leipopsych'ia*, *Lipopsych'ia*, *Ap'eych'ia*, *Ap'eych'ia*, *Lipsis An'imi*, *Asphyx'ia*, *Apopsych'ia*, *Lipothym'ia*, *Leipothym'ia*, *Lipophyx'ia*, *Eclips'is*, *Su'bitus animi casus*, *Anepcephaloh'mia*, *Ap'eychy*, *Swoon*, *Swooning*, *Fainting*, *Fainting fit*, *Faintness*, *Exol'ution*, (Sc.) *Drac*, *Dualm*, *Sicaff*, *Swerf*, *Swarth*, (Prov.) *Fower*, *Swebbing*, *Sweeming*, *Swooning*, *Sound*, (F.) *Evanouissement*, *Défaillance*, *Pamoi'on*. Complete and, commonly, sudden loss of sensation and motion, with considerable diminution, or entire suspension of the pulsations of the heart and the respiratory movements. Syncope is, commonly, an affection of no consequence; but, sometimes, it is an index of diseased heart. Placing the patient in the horizontal posture; sprinkling cold water over the face, and the application of volatile substances to the nose, are all that will be required during the fit. See Asphyxia, and Suffocation.

An approach to this condition is termed *faintness*; and the person is said to be *faint*, (Prov.) *Sweemish*.

SYNCOPE ANGENS, Angina pectoris — s. Anginosa, Angina pectoris.

SYNCOPTICUS, Syncopeal.

SYNCOPTUS, Syncopeal.

SYNCRITICA, Astringens.

SYNDACTYLUS, Aschistodactylus.

SYNDESMITIS, see Ophthalmia. Also, inflammation of articular ligaments: *Inflammatio ligamentorum*, from *syndemos*, 'ligament,' and *itis*, denoting inflammation.

SYNDESMOGRAPHY, *Syndesmograph'ia*, from *syndemos*, 'a ligament,' and *γραφω*, 'a description.' A description of the ligaments. The part of anatomy which describes the ligaments.

SYNDESMOL'OGY, *Syndesmolog'ia*, from *syndemos*, 'a ligament,' and *λογος*, 'a treatise, discourse.' A treatise on the ligaments.

SYNDESMO-PHARYNGE'US. A fasciculus of the constrictor pharyngis superior.

SYNDESMOS, Ligament.

SYNDESMOSIS, *Synneuro'sis*, *Synyme'nesis*, *Connex'io* seu *Syn'physis ligament'ia*, *Commis'sura ner'va*, *Conner'atio*, *Colligatio*, from *syndemos*, 'ligament.' The articulation of two or more bones by means of ligament; — as of the radius with the ulna.

SYNDESMOT'OMY, *Syndesmotom'ia*, from *syndemos*, 'ligament,' and *τομω*, 'I cut or dissect.' Dissection of the ligaments.

SYNDESMUS, Bandage, Ligament.

SYN'DROME, from *syn*, and *δρομος*, 'a course.' *Concur'sus*, *Concourse*. A name given by the Empirical Sect to the union of symptoms which takes place in diseases: — *Syndromē plethor'ica*, for example, meant the union of symptoms produced by plethora.

SYNDYASMUS, Coition.

SYNECHES, see Continued fever.

SYNECHI'A, from *syn*, and *εχειν*, 'to have,' or 'to hold.' Adhesion between the iris and transparent cornea, or between the iris and crystalline lens. The former case is called *Synechia anterior*; the latter, *S. posterio*.

SYNERGIA, Synergy.

SYNERGIA, Synergy.

SYNERGIE, Synergy.

SYNERGY, *Synerg'ia*, *Synergery'ia*, (F.) *Synergie*; from *syn*, and *εργον*, 'work.' A correlation or concurrence of action between different organs in health; and, according to some, in disease.

SYNESIS, Intellect.

SYNEUROSIS, Syndesmosis.

SYNEZISIS, *Synize'sis*, *Consident'ia*, from *syn*, and *συζυναι*, 'to join.'

SYNEZISIS PUPILL'Æ, *Parop'sis Synizesis*, *Caligo* & *Synizesis*, *Caligo seu Imperforatio* seu *Phthisis pupillæ*, *Amyo'sis*, *Closure* or *occlusion* or *obliteration* of the Pupil. This disease is sometimes a faulty conformation, depending upon the continued existence of the membrana pupillaris, or the absence of the pupil. It is, in other words, *congenital*. At other times, it is owing to accident, and supervenes on violent ophthalmia, or on the operation for cataract. They who are affected with synezisis are completely blind: the iris having no central aperture. In certain cases, a remedy may be obtained by making an opening in the iris, which may supply the place of the pupil, and through which the rays of light may reach the retina. This operation is called the operation for artificial pupil.

SYNGENES, Hereditary.

SYNGENICUS, Congenital.

SYNHYMENESIS, Synymenesis.

SYNIMENESIS, Synymenesis.

SYNIZESIS, Considentia, Synyzisis.

SYNEUROSIS, Syndesmosis.

SYN'OCHA, from *συνεχω*, (*συν*, and *εχω*, 'I have or hold,') 'I continue.' *Inflam'matory* or *Dynam'ic Fever*, *Angiop'ya*, *Sthenop'ya*, *Febris caus'odes seu caumat'odes seu deu'rens seu ardens seu contin'ua sanguin'ea seu contin'ua non putris seu amon'stica seu acut'a seu acut'a sanguin'ea seu contin'ens seu continens non pu'trida seu angioten'ica seu sthenica seu sanguin'ea seu inflam'matoria*, *Homot'onos*, *Syn'ochus* (Vogel), *S. impu'tris* (Galen), *Ene'cia cauma seu ton'ica*, *Phlogop'yrus*, *Cauma*, *Calentura continua*, (F.) *Fièvre ardente ou continente inflammatoire ou sanguine ou irritative ou angioténique ou inflammatoire*, *Gastro-entérite intense*; *General inflammation*. A species of continued fever, characterized by increased heat; and by quick, strong, and hard pulse; urine high-coloured; disturbance of mind slight. It requires, of course, the most active treatment.

SYNOCHA RHEUMATICA, Rheumatism, acute — s. *Vaccina*, *Vaccina*.

SYN'OCHAL, *Synocha'lis*; same etymon. Relating to, or having the characters of, synocha.

SYN'OCHUS, *Ene'cia syn'ochus*, *Febris lenta*, (Linnaeus) same etymon. Continued fever, compounded of synocha and typhus; — in its commencement often resembling the former; in its progress, the latter: *Mixed fever*, *Common continued fever*, *Syn'ochoid fever*. See Continued fever, Fever, and Typhus.

SYNOCHUS, Synocha — s. *Biliosa*. Fever, gastric, Fever, bilious — s. *Catarrhalis*, *Influenza* — s. *Icte-*

rodes, Fever, yellow — s. Imputris, Synocha — s. Miliaria, Miliary fever — s. Varicella, Varicella.

SYNOLCE, Spasm.

SYNONIA, Synovia.

SYNSTEOGRAPHY, *Synosteographia*, from *syn*, σύν, a 'bone,' and *γραφω*, 'I describe.'

The part of anatomy which describes the joints.

SYNSTEOL'OGY, *Synosteologia*, from *syn*, σύν, 'a bone,' and *λογος*, 'a treatise.' A treatise on the joints.

SYNSTEO'SIS, *Synostosis*, from *syn*, σύν, and *στέον*, 'a bone.' Union by means of bone.

SYNSTEOT'OMY, *Synosteotomia*, from *syn*, σύν, 'with,' *στέον*, 'a bone,' and *τεμνω*, 'to cut.' Dissection of the joints.

SYNSTOSIS, Synostosis.

SYNOVIA, from *syn*, σύν, and *ων*, 'an egg.'

Mucila'go, *Un'guen* seu *Unguen'tum* seu *Smegma articula'rè*, *Syno'nia*, *Azun'gia articula'ris*, *Mucila'go articulo'rum* seu *junctura'rum*, *Humor articula'ris*, *Hydrarthrus*, *Humor* seu *Hu'midum nati'vum* seu *Gluten articulo'rum*, *Arthrohydrine*, (F.) *Synovie*. A fluid resembling the white of egg, which is exhaled by the synovial membranes surrounding the moveable articulations. The synovia has been found, on analysis, to consist of water, albumen, soda, chloride of sodium, phosphate of lime, and — Fourcroy fancied — uric acid. Its use is to lubricate the joints, and the sheaths in which tendons play.

SYNOVIAL, *Synovia'lis*. That which relates to the synovia.

SYNOVIAL GLANDS, *Havers's Glands*, *Glandulæ articula'res* seu *Synovia'les* seu *Haversia'næ*, are the secretory fringes, which float in the interior of the synovial capsules (see Capsules.) They do not seem to be glandular. They have been called *Synovial sim'briae*, (F.) *Franges synoviales*.

SYNOVITIS, from *Synovia*, and *itis*, a termination denoting inflammation. A term, badly compounded, applied at times to inflammation of the synovial membrane, *arthrophlogo'sis synovia'lis*. When it affects the bursæ, it is, sometimes called *Bursitis*.

SYNOVITIS RHEUMATISMALIS, Rheumatism, (acute.)

SYNTASIS, *συντασις*, from *συνταίω*, (*syn*, and *ταίω*, 'I stretch,') 'I strain.' Tension of parts: hence *συνταξικά*, are medicines which render parts tense. *Syntex'is*, on the other hand, means *Colliqua'tio*, or wasting of the body; from *συντηνω*, 'I melt together.'

SYNTAXIS, Articulation, Reduction, Taxis — s. Serrata, see Suture.

SYNTECOPYRA, Hectic fever.

SYNTECTYCOPYRA, Hectic fever.

SYNTEO'SIS, from *syn*, σύν, and *τενω*, 'a tendon.' A species of articulation in which two bones are joined by means of tendon; as the sesamoid bones with the toes, the rotula with the tibia, &c.

SYNTERESIS, Prophylaxis.

SYNTERETICUS, Prophylactic.

SYNTEXIS, Colliquation, Consumption, see Syntasis.

SYNTHENA. A term employed by Paracelsus to designate a species of epilepsy, accompanied by cardialgia and tormina.

SYNTHESIS, *Ren'io*, 'composition or putting together,' from *syn*, σύν, and *τιθημι*, 'I place.' A surgical operation, which consists in uniting parts that are divided; and approximating such as are separated or distant from each other. *Ren'io par'tium separata'rum*, *Syn'thesis of continuity*, means the union of the edges of a wound, or the approximation of the extremities of a fractured bone. *Syn'thesis of contigu'ity* is the reduction of displaced organs; as in cases of hernia and luxations.

SYNTHESIS, Composition.

SYNTHETIS'MUS, same etymon. The aggregate of operations and means for reducing a fracture, and maintaining it reduced: — comprising extension, counter-extension, coaptation, and bandaging.

SYNTONINE, see Fibrin.

SYNTRIMMA, Conitrio.

SYNTRIPSIS, Conitrio.

SYNTROPHOS, Connutritus.

SYNULOSIS, Cicatrization.

SYNULOTICA, Cicatrissantia.

SYNUSIA, Coition.

SYNYME'NESIS, *Synhyme'nesis*, *Synime'ne'sis*, from *syn*, σύν, and *υμνη*, 'a membrane.' Union of bones by membrane, as in the case of the bones of the head in the fœtus.

SYNYMENESIS, Syndesmosis.

SYPHAR, Peritonium.

SYPHILIDES, *Syphili'dæ*, (F.) *Dermo-syphili'des*, *Dermatoses véroleuses*. Same etymon as Syphilis. The cutaneous eruptions that accompany constitutional syphilis are so denominated by the French writers.

SYPHILIDIA'TER, *Syphilidia'trus*, from *Syphilis*, and *ιατρος*, 'a physician.' A pox doctor. One who occupies himself in treating syphilis.

SYPHILIDOMANIA, Syphilomania.

SYPHILIPHO'BIA, from *Syphilis* and *φοβος*, 'dread.' A morbid dread of syphilis, giving rise to fancied symptoms of the disease: a form of hypochondriasis.

SYPH'ILIS, *S. vene'rea* seu *malig'na*, *Siphilis*, *Syphili's*, *Cache'ria* seu *Cacochy'm'ia Vene'rea*, *Malum vene'reum*, *Puden'dagra*, *Me'vium*, *Lues*, *L. vene'rea* seu *syphilis*, *Morbus Gal'licus* seu *Ital'icus* seu *Hispan'icus* seu *Neapolita'nus* seu *Ind'icus* seu *aphrodis'ius*, *Paturæ*, *Basilis'cus*, *Boæ*, *Sod'oma Gallo'rum*; *Veneræal*, *Vene'real Disease*, *Poz*, *French Poz*, *French Crust*, *French Distemper*, *Foul Disease*, old cant term, *Grincomes*, (Prov.) *Pip*, (Sc.) *Glengore*, *Glengour*, *Grandgore*, (F.) *Vérole*, *Maladie vénérienne*, *Mal de Naples*, *Mal Français*, *Maladie de Vénus*. The etymology of syphilis is unknown. Some consider it to proceed from *συσ*, 'a hog,' and *φίλω*, 'I love,' others, from *συν*, 'with,' and *φίλω*, 'I love,' and others, with more probability, from *σφίλω*, — formed, by contraction, from *σφαλεος*, — 'a reproach,' &c. It is an infectious disease, communicable by coition, or by the contact of parts that are only lined with a thin epidermis; as the lips, nipple, &c. It generally makes its first appearance by a chancre, followed sooner or later by a bubo. The chancre is a primary symptom or sore; but there are others which are secondary or constitutional. These are: — ulcers in the throat; copper-coloured blotches on the skin; pains in the bones, nodes, &c.: the last are sometimes termed tertiary phenomena. The origin of syphilis is not known. It is believed to have been introduced into Europe from America, at the end of the 15th century [?]. Its progress is extremely variable, as well as its duration, which is often unlimited, if left to itself. Mercury may be esteemed the best remedy, but the disease admits of cure without it. Almost every antisiphilitic composition, whose use is attended with success, probably contains mercury united with diaphoretics or other medicines. It is generally sufficient to keep the mouth sore for a fortnight for the cure of chancre: for venereal sore throat, and other recent constitutional symptoms, eight or ten weeks will be required; and, of course, a longer time, if these symptoms have persisted longer. Sometimes, after mercury has been given for a time, the symptoms continue stationary; and do not yield until the mercury has been discontinued, and

tonics and eutrophics have been administered. The preparations of iodine have been much used, of late, in the revellent treatment of this disease.

SYPHILIS INDICA, Framboesia.

SYPHILIS PSEUDO-SYPHILIS, *Lues Syphilo'des, Pseudo-syphilis*. A term given to ulcers or other affections that resemble syphilis, but are indeterminate in their character,—the symptoms being irregular in their appearance, and usually yielding spontaneously, or under the use of tonics, change of air, &c. The condition has not been at all well defined. The *Sivvens, Radesyge, &c.*, have been referred to this head.

SYPHILISATION, Syphilization.

SYPHILISMUS, Syphilis.

SYPHILITIC, *Syphiliticus*, (F.) *Syphilitique, Vérolique, Vérolé*. Belonging, or relating to syphilis—as 'a syphilitic ulcer.'

SYPHILIZATION, *Syphilisation*, (F.) *Syphilisation*. Saturation of the system by inoculation with the matter of syphilis. It has been used as a curative procedure in syphilis, as well as a preventive of that disease; and for the cure of other cachexiæ, as cancer.

SYPHILODERMA, from *syphilis*, and *δερμα*, 'skin.' A cutaneous eruption of a syphilitic character.—E. Wilson.

SYPHILOG'RAPHER, from *syphilis*, and *γραφω*, 'I describe.' A writer on syphilis.

SYPHILOID, *Syphiloïdes*; from *syphilis*, and *ειδος*, 'resemblance.' Resembling syphilis,—as 'a syphiloid affection.'

SYPHILOMANIA, *Syphilidomania*,—vulgarly, *Noddle Pox*. A mania, with which some persons are affected, so that they subject themselves to antiveneral treatment, under the erroneous impression that they are affected with syphilis.

SYPHITA, Somnambulism.

SYPHO AURICULARIS, Otoschyttes.

SYPHON, see Siphon.

SYRIGMOPHONIA, from *συσπυμος*, 'piping,' and *φωνη*, 'voice.' A piping state of the voice.

SYRIGMUS, Tinnitus aurium.

SYRINGA, Eustachian Tube, Fistula, Syringe, Trachea—s. Clysmatica, Syringe, (Clyster.)

SYRINGA VULGARIS, *Lilæe vulgaris*, Common Lilac. Ord. Oleaceæ. The leaves and fruit are bitter and somewhat acid, and have been used as tonics and antiperiodics.

SYRINGE, *Syr'ina*, *Syrin'ga*, *Sipho Siphunculus*, *Squirt*, (F.) *Seringue*, from *συσπυγξ*, *σπυγγος*, 'a pipe.' A familiar apparatus through which any liquor is injected.

Various forms of syringe are used in medicine and surgery.

SYRINGE, CLYSTER, *Syrin'ga clysmatica*, *Clystanti'um*, (F.) *Clysoponge*. A syringe for administering clysters.

SYRINGE, CUPPING, *Antlia sanguisuga*.

SYRINGES AERIFERI, see Bronchia.

SYRINGICUS, Fistulous.

SYRINGITIS EUSTA'CHII, Inflammation of the Eustachian tube.

SYRINGMOS, Tinnitus aurium.

SYRINGODES, Fistulous.

SYRINGOTOME, *Syringot'omus*, *Syringot'omum*, from *συσπυγξ*, *σπυγγος*, 'a pipe,' 'a fistula,' and *τεμνειν*, 'to cut.' A surgical instrument, formerly used in the operation for *fistula in ano*. It is a bistouri, concave at its edge, and terminated by a long, flexible, probe-pointed stylet. This stylet or probe was first introduced by the external opening of the fistula into the rectum, whence it was drawn externally through the anus, carrying with it the cutting edge of the instrument, which divided the intermediate parts.

SYRINGOT'OMY, *Syringotom'ia*. Same etymon as the last. The operation for fistula in ano by incision.

SYRINX, Eustachian tube, Fistula, Syringe, Trachea—s. Hiera, Vertebral column.

SYRMAIS'MOS, from *συναπαινω*, 'I evacuate.' A gentle evacuation by vomiting or stool.

SYROP, Syrup—s. d'*Ail*, Syrupus allii—s. d'*Armoisee composee*, Syrupus aromaticus—s. de *Chicorée composee*, Syrupus de cichorio compositus—s. des *Cinq racines*, Syrupus e quinque radicibus—s. de *Coquelicot*, Syrupus rheados—s. de *Cuisinier*, Syrupus de sarsaparilla et sennâ compositus—s. d'*Ether*, Syrupus cum æthere sulphurico paratus—s. de *Gingembre*, Syrupus zingiberis—s. de *Guinauwe*, Syrupus althææ—s. de *Longue vie*, Mellitum de mercuriali compositum—s. de *Mercur*, Syrupus de mercurio mediante gummi—s. de *Mou de veau*, see Syrupus zingiberis—s. de *Mûres*, Syrupus mori—s. de *Nerprun*, Syrupus rhamnii—s. de *Pavots blancs*, Syrupus papaveris—s. de *Pommes composee*, see Syrupus sennæ—s. de *Quinquina avec le Vin*, Syrupus de kinâ kinâ cum vino—s. de *Roses pâles*, Syrupus rosæ—s. de *Roses rouges*, Syrupus rosæ—s. de *Vinaigre*, S. aceti.

SYRUP, *Sirup*, *Syr'upus*, *Serapi'um*, *Serape'um*, (F.) *Syrop*, *Sirop*, *Saccharolé liquide*, *Saccharhydrolé*; derived, according to some, from *συρω*, 'I draw,' and *σμος*, 'juice'; but more probably, from the Arabic *Siruph*, *Sirab*, or *Scharab*, which signifies 'a potion.' A saturated solution of sugar in water, either simply, or united with some vegetable or other principle. Syrups, in general, require two pounds of sugar to the pint, and they are judged to be sufficiently boiled when a portion, taken up in a spoon, flows like oil. In the Pharmacopœia of the United States, it is directed, that syrups, whose density is not precisely determined by the process, should have the s. g. 1.261 when boiling, and about 1.319 at ordinary temperatures. Syrups should be kept in small bottles in a cool place. They are chiefly used to render medicines palatable. See Syrupus simplex.

SYRUP OF ALMONDS, Syrupus amygdalæ.

SYRUP, ANTISCORBUTIC, Syrupus antiscorbuticus—s. Balsamic, Syrupus toltanus.

SYRUP OF BUCKTHORN, Syrupus rhamnii—s. of Calf's lungs, see Syrupus zingiberis—s. of Capillaire, common, Syrupus simplex—s. of Cinchona with wine, Syrupus de kinâ kinâ cum vino—s. of Cinchona, Syrupus cinchonæ—s. of Citric acid, Syrupus acidi citrici—s. of Clove July flower, Syrupus caryophylli rubri—s. of Clove pinks, Syrupus Caryophylli rubri—s. of Cochineal, Syrupus coccii—s. of Emetine, Syrupus emetinæ—s. of Ether, Syrupus cum æthere sulphurico paratus—s. of Five roots, Syrupus e quinque radicibus—s. of Garlic, Syrupus allii—s. of Ginger, Syrupus zingiberis—s. of Gum Arabic, Syrupus acaciæ,

SYRUP, HIVE, Syrupus scillæ compositus.

SYRUP OF HONEY, Mellitum simplex—s. of Iodide of iron, Liqueur Ferri iodidi—s. of Ipecacuanha, Syrupus ipecacuanhæ—s. of Jalap, Syrupus jalapæ—s. of Lemon, Syrupus limonis—s. of Marshmallow, Syrupus althææ—s. of Meadow saffron, Syrupus colchici autumnalis—s. of acetate of Morphia, Syrupus morphiæ aceticæ—s. of muriate of Morphia, Syrupus morphiæ muriatis—s. of Mugwort, compound, Syrupus aromaticus—s. of Mulberries, Syrupus mori—s. of Orange juice, see Syrupus aurantii—s. of Orange peel, Syrupus aurantii—s. of Orgeat, Syrupus amygdalæ—s. of Poppies, Syrupus papaveris—s. of Poppy, red, Syrupus rheados—s. of Quinia, Syrupus quiniæ—s. of Rha-

tany, Syrupus krameriae—s. of Rhubarb, Syrupus rhei—s. of Rhubarb, aromatic, Syrupus rhei aromaticus—s. of Rhubarb, spiced, Syrupus rhei aromaticus—s. of Roses, Syrupus rosæ—s. of Roses, red, Syrupus rosæ gallica—s. of Saffron, Syrupus croci—s. of Sarsaparilla, Syrupus sarsaparillæ—s. of Sarsaparilla and senna, compound, Syrupus de sarsaparillâ et sennâ compositus—s. of Sarsaparilla, compound, see Syrupus de sarsaparillâ et sennâ compositus—s. of Sarsaparilla, Indian, Syrupus hemidesmi—s. of Scammony, Syrupus de scammonio—s. of Seneka, Syrupus senegæ—s. of Senna, Syrupus sennæ—s. Simple, Syrupus simplex—s. of Squill, Syrupus scillæ—s. of Squill, compound, Syrupus scillæ compositus—s. of Suceory, compound, Syrupus de cichorio compositus—s. of Tolu, Syrupus tolu-tanæ—s. of Vinegar, Syrupus acetii—s. of Violets, Syrupus violæ—s. of Wild cherry bark, Syrupus Pruni Virginianæ.

SYRUPUS, S. simplex.

SYRUPUS ACACIÆ, Syrup of Gum Arabic, (Acacia 3ij, Sacchar. 3xv, Ag. f 3viiij. Dissolve the gum in the water; add the sugar, and strain. Ph. U. S.) A good pectoral syrup. Dose, f 3j to f 3ij.

SYRUPUS ACETI, S. Ac'idi Aceto'si, Syrup of Vinegar, (F.) Syrup de Vinaigre. (Aceti Oïiss, sacch. purif. lbiiiss. Boil to form a syrup. Ph. E.) It is refrigerant and antiseptic. Dose, f 3j to f 3ij.

SYRUPUS ACIDI ACETOSI, S. acetii.

SYRUPUS AC'IDI CITRICI, Syrup of Citric Acid. (Acid. citric. pulv. 3ij; Ol. Limon. Mij; Syrup. Oij. Rub the acid and oil with an ounce of the syrup; add the rest of the syrup, and dissolve with a gentle heat. Ph. U. S.) Virtues the same as those of the Syrupus Limonis.

SYRUPUS CUM ÆTHERE SULPHURICO PARATUS, Syrup of Ether, (F.) Syrup d'ether. (Very pure sugar, 1000 p.; distilled water, 500 p.; dissolve in the cold, and strain; add sulphuric ether, 48 p. Shake for five or six days, and draw off the limpid syrup. Ph. L.) Properties like those of ether.

SYRUPUS ALII, Syrup of Garlic, (F.) Syrup d'ail. (Allii recent. concis. 3vj, acet. destillat. Oj, sacchar. lbij. Macerate the garlic in the vinegar, in a glass vessel, for four days; then express the liquor and set it by, that the dregs may subside; lastly, add the sugar to the clear liquor, and proceed in the manner directed for syrup. Ph. U. S.) Use, same as that of garlic. Dose, f 3j to f 3ij.

SYRUPUS ALTHÆÆ, S. ex Althæâ seu de Althæâ seu Althææ officinalis, Syrup of Marsh-mallow, (F.) Syrup de Guimauve. (Althææ rad. recent. cont. lbss, sacch. pur. lbij, aquæ Oiv. Boil the root in the water to one-half, and press out the liquor; when cold, defecate; add the sugar, and boil. Ph. L.) Emollient and demulcent. Dose, f 3j to f 3ij.

SYRUPUS ANOMI ZINGIBERIS, S. zingiberis.

SYRUPUS AMYGDALÆ, Syrup of Almonds, S. of Orgeat, Orgeat. (Amygdal. dulc. lbj, amygdal. amar. 3iv, aquæ, Oij, sacchar. lbvj. Blanch the almonds, and rub them in a mortar to a very fine paste, adding, during the trituration, three fluid-ounces of the water, and a pound of the sugar. Mix the paste thoroughly with the remainder of the water; strain with strong expression; add the remainder of the sugar to the strained liquor, and dissolve it with the aid of a gentle heat. Strain the syrup through fine linen, and having allowed it to cool, put it into bottles, which must be well stopped, and kept in a cool place. Ph. U. S.)

SYRUPUS ANTICAUSTICUS, see Anticaustodic.

SYRUPUS ANTISCORBUTICUS, S. de Armora'ciâ

compositus, Antiscorbutic Syrup. (Leaves of cochlearia, buckbean, water-cresses, horse-radish root and bitter oranges, each lbj, cinnamon, 3iss. Cut, and pour on white wine lbvj. Macerate for two days; distil over Oj in a sand-bath; add to this sugar, lbij, and form a syrup. Again, take what remains in the cucurbit: pass it through cloth; let it rest; decant, and add sugar lbij; make into a syrup, and clarify with white of egg. Mix it with the first while yet warm. Ph. P.) Antiscorbutic, alterative, tonic, stomachic. &c. Dose, f 3ij to f 3ij.

SYRUPUS AROMATICUS, S. de Artemisiâ compositus, Compound Syrup of Mugwort, (F.) Syrup d'armoise composé. (Tops of artemisia vulgaris, in flower, 3vj, inula root, ligusticum levisticum, and fennel, each, 3ss, pennyroyal, catmint, savine leaves, each 3vj, marjoram, hyssop, pyrethrum, rue, basil, each 3ij, aniseed and cinnamon, each 3ix; bruise and macerate in hydromel lbxviij; distil over Oss; add to this a pound of sugar. Again, take the liquid that remains in the cucurbit, strain it gently; and add sugar lbvj. Then mix the two. Ph. P.) Tonic, emmenagogue, &c.

SYRUPUS DE ARMORACIÂ COMPOSITUS, S. antiscorbuticus—s. de Artemisiâ compositus, S. antiscorbuticus—s. Asparagi, see Asparagus.

SYRUPUS AURANTII, S. Aurantio'rum seu Citri Aurantii seu Corticis Aurantii seu Aurantii Corticis (Ph. U. S.) seu e Cortic'ibus Aurantio'rum seu de Cortic'ibus Aurantio'rum, Syrup of Orange peel. (Aurant. cort. cont. 3ij, ag. ferv. Oj, sacch. lbijss. Macerate the peel in the water for twelve hours. Add the sugar. Ph. U. S.) Stomachic. Dose, f 3j to f 3ij.

A Syrup of Orange Juice, Orange Sherbet, Syrupus e succo Aurantiorum, may be made by adding white sugar lbj to lbj of the juice of oranges, strained and clarified.

SYRUPUS BALSAMICUS, S. toluatanus.

SYRUPUS CARYOPHYLLI RUBRI, S. Infusio'nis florum caryophylli'rum, S. Dianthi Caryophylli, Syrup of Clove July flower or of Clove Pinks. (Petalorum dianth., caryoph. recent., unguibus resectis, lbj, aquæ bull. Oiv, sacch. purif. lbvj. Ph. E.) Aromatic.

SYRUPUS CASSIÆ SENNÆ, S. sennæ.

SYRUPUS DE CICHORIO COMPOSITUS, S. de Rheo, Compound Syrup of Succory, (F.) Syrup de chicorée composé. (Root of succory 3vj, leaves of succory 3ix, leaves of fumitory and scolopendrium, each 3ij, Alkekengi berries 3ij. Boil in water Oxiv to Oxij: pass through cloth, and dissolve in white sugar lbvj. Boil till it has the consistency of syrup. Again, take water Oviij, rhubarb 3vj, yellow sanders and cinnamon, each 3ss. Infuse for twenty-four hours; pass through linen, and add the preceding syrup. Ph. P.) It is laxative and stomachic. Dose, f 3j to f 3ss.

SYRUPUS CINCHONÆ CUM VINO, S. de kinâ kinâ cum vino.

SYRUPUS CINCHO'NIÆ, Syrup of Cincho'nia. (Simple syrup one pound; sulphate of cinchonia nine grains. Mix.) Six spoonfuls are commonly sufficient to arrest an intermittent.

SYRUPUS CITRI AURANTII, S. aurantii.

SYRUPUS COCCI, Syrup of Cochineal. (Cocci. cont. Div. Ag. destillat. bullient. Oj (Imp. meas.), sacchar. lbij, vel q. s. Sp. rectificat. f 3iiss, vel q. s. Ph. L.) Chiefly used as a coloring agent.

SYRUPUS COL'CHICI AUTUMNA' LIS, S. Colechici, Syrup of Meadow Saffron. (Rad. colch. autumn. recent. 3j, acidi acetosi, f 3xvi, sacch. pur. 3xxvj. Macerate the root in the vinegar for two days, occasionally shaking the vessel; then strain with gentle expression. To the strained fluid add the sugar, and boil to a syrup. Ph. E.) Diuretic, antiarthritic. Dose, f 3j to f 3ss, or more.

SYRUPUS CORTICIS AURANTII, *S. aurantii* — s. Citri medicæ, *S. limonum* — s. e Succo citrorum, *S. limonum*.

SYRUPUS CROCI, *Syrup of Saffron*. (*Croc. stigmat.* 3j, *aq.æ ferrent.* Oj, *sacch. purif.* lbiss. *Ph. L.*) Cardiac. Chiefly used for colouring.

SYRUPUS DIACODION, *Diacodium* — s. *Dianthi Caryophylli*, *S. caryophylli rubri*.

SYRUPUS EMETICÆ, *Syrup of Emet.etine.* (*Simple syrup* lbj, *emetine* 16 gr.) In small doses, expectorant; and in large, emetic.

SYRUPUS EMPYREUMATICUS, *Melasses* — s. *Ferri iodidi*, *Liquor Ferri iodidi*.

SYRUPUS HEMIDESMI, *Syrup of Indian Sarsaparilla*. (*Hemidesm. cont.* 3iv (*avoidup.*), *Aq. destillat. bullient.* Oj (*Imp. meas.*), *sacchar. q. s.* *Ph. D.*) A weak preparation of hemidesmus.

SYRUPUS HYDRARGYRI, *Syrupus de mercurio mediante gummi* — s. *Infusionis fluorum caryophyllorum*, *Syrupus caryophylli rubri*.

SYRUPUS IPECACUANÆ, *Syrup of Ipecacuanha*. (*Ipecac.* in pulv. crass. 3j, *alcohol. dilut.* Oj, *sacchar.* lbiss, *aq.æ q. s.* Macerate the ipecacuanha in the alcohol for 14 days, and filter. Evaporate the filtered liquor to 6 fluidounces, again filter, and add sufficient water to make the liquid measure a pint; add the sugar, and evaporate by a water-bath to a proper consistence. *Ph. U. S.*) It may also be prepared by displacement.

SYRUPUS DE JALAP'Æ, *Syrup of Jalap.* (*Pulv. Jalapæ* 3x, *sem. coriandræ*, *feniculi. dulc.* aa 3ss; *aq.æ f3xijss, cocch.* 3xxv. *Ph. P.*) Cathartic. Dose, f3j to f3ss for children.

SYRUPUS DE KINÀ KINÀ CUM VINO, *S. Cincho'ne cum vino*, *Syrup of Cinchona with wine*, (*F.*) *Syrup de quinquina avec le vin.* (*Cort. cinchonæ officinalis* 3ij, *ext. cinchonæ* 3vi, *vini albi* Vin. Linel. seu *conjuvis ei similis*, Oj, *alcohol* (s. g. '955 to '915.) f3j, *sacchar. alb.* lbiss. Bruise the cinchona; gradually add the alcohol, so as to form a soft mass; pour on the wine; macerate for two days, and strain. Dissolve the extract, add the sugar, and make a syrup. *Ph. P.*) Dose, f3ss to f3j.

SYRUPUS KRAMERIE, *Syrup of Rhat'any.* (*Ext. Kramer.* 3ij, *aq.æ* Oj, *sacchar.* lbjss. Dissolve the extract in the water, and filter; then add the sugar, and form into syrup. *Ph. U. S.* It may also be made by displacement from *kramer.* in pulv. crass. lbj, *sacchar.* lbiss; *aq. q. s.*) An adjunct to astringent mixtures; or given in diarrhœa, a teaspoonful to a child.

SYRUPUS LIMONIS, *S. Limonum* — s. *Succi limonis*, *S. limonum*.

SYRUPUS LIMO'NUM, *S. limonis* (*Ph. U. S.*) seu *citri medicæ* seu *e succo citro'num* seu *e succo limo'num* seu *succi limo'nis*, *Syrup of Lemon*. *Lemon Syrup.* (*Limon. succ. colat.* Oj, *sacchar.* lbij.) Cooling, antiseptic.

SYRUPUS E MECONIO, *S. papaveris* — s. *de Melle*, *Mellitum simplex*.

SYRUPUS DE MERCURIO MEDIAN'TE GUMMI, *S. Hydrargyri*, *Mercurius gummosus Plenck'ii*, (*F.*) *Syrup de mercure*, *Mercur gommeux de Plenck.* (*Hydrargyri pur.* 3j. p. gum. acaciæ, 3ij, *syrup diacod.* 3ss. Rub together till the mercury is extinguished. *Ph. P.*) Possesses the properties of blue pill. Dose, gr. ij to viij, or more.

SYRUPUS MORI, *S. e succo moro'rum* seu *succi mori*, *Syrup of Mul'berries*, (*F.*) *Syrup de mûres.* (*Mori succ. colat.* Oj, *sacchar. purif.* lbij. *Ph. L.*) Gratefully refrigerant.

SYRUPUS E SUCCO MORORUM, *S. mori*.

SYRUPUS MORPH'Æ ACETAT'IS, *Syrup of acetate of morphia.* (*Liq. morphicæ acet.* f3j, *syrup f3xv.* *Ph. D.*) Dose, f3ss to f3j, as an opiate.

SYRUPUS MORPH'Æ MURIAT'IS, *Syrup of muriate of morphia.* (*Liq. morph. muriat.* f3j;

syrup f3xviij. *M. Ph. D.*) Dose, f3ss to f3j, as an opiate.

SYRUPUS MORPH'Æ SULPHAT'IS may be made in the same way.

SYRUPUS NIGER, *Melasses*.

SYRUPUS PAPAVERIS, *S. papaveris somniferi* seu *papaveris albi* seu *emecol'io* seu *de meconio*, *Diacod'ium*, *Syrup of poppies*, (*F.*) *Syrup de pavots blancs ou Diacode.* (*Papav. capent.* exsiccet. et contus. exemptis seminibus, 3xiv, *sacch. pur.* lbij, *aq. ferrent.* cong. ijss. Macerate the capsules in liquor for 12 hours; boil to a gallon, and express strongly. Boil the liquor to Oj; and strain while hot. Defecate by rest, and boil to Oj, adding the sugar so as to form a syrup. *Ph. L.*) One ounce contains about one grain of opium. Anodyne and demulcent.

The Dublin college has a SYRUPUS OPII, called *Common Syrup of Poppies*, which is made by dissolving 18 grains of the *Watery Extract of Opium* in 18 ounces of boiling water. Properties, those of opium and syrup.

SYRUPUS DE PAPAVERE ERRATICO, *S. rhæados* — s. *de Pomis compositus*, see *S. sennæ*.

SYRUPUS PROTEÆ, see *Protea mellifera*.

SYRUPUS PRUNI VIRGINIANÆ, *Syrup of Wild cherry bark.* (*Pruni Virginian.* in pulv. crass. 3v, *sacchar.* lbij, *Aquæ*, q. s. Moisten the bark with water; let it stand for 24 hours; transfer it to a percolator; pour water gradually upon it, until a pint of filtered liquor has passed. Add the sugar in a bottle, and dissolve by agitation. *Ph. U. S.*, 1851.) Dose, as a demulcent expectorant, f3j to f3ij.

SYRUPUS E PULMONIBUS VITULINIS, see *Syrupus Zingiberis*.

SYRUPUS QUINIE, *Syrup of Quinia.* (*Simple syrup*, 2 pints, *sulphate of quinia*, 64 grains.) Six spoonfuls are commonly sufficient to arrest an intermittent.

SYRUPUS E QUINQUE RADICIBUS, *Syrup of five roots*, (*F.*) *Syrup des cinq racines.* (*Rad. apii, feniculi, apii petroselinii.* aa 3v. Infuse for 24 hours in boiling water Oiss, and strain. *Rad. asparag.*, *rusc. aculeati.* aa 3v, *aq.æ*, Oviij. Boil and reduce to half; adding, towards the end, the residue of the preceding infusion. Boil again for some minutes; strain, and add *white sugar*, lbvj. *Ph. P.*) Aperient and diuretic. Used in obstructions of the liver, spleen, and mesentery; and in dropsy, &c. Dose, f3ss to f3j, or more.

SYRUPUS RHAMNI, *S. Rhamni cathartici* seu *de Spinâ cervinâ* seu *spinâ cervinâ*, *Syrup of Buckthorn*, (*F.*) *Syrup de nerprun.* (*Rhamn. baccar. succ. recent.* Oiv, *zingib. rad. concis.*, *pimento bacc. contrit.*, aa 3ss, *sacch. purif.* lbiss. Defecate the juice for 3 days, and strain. To a pint of this, add the ginger and pimento; macerate for 4 hours, and strain; boil what remains to Oiss. Mix the liquors and add sugar, so as to form a syrup. *Ph. L.*) Cathartic, but apt to gripe. Dose, f3iv to f3ij.

SYRUPUS RHEI, *Syrup of Rhubarb.* (*Rhei*, in pulv. crass. 3ij; *Alcohol*, Oss; *Aquæ*, Oiss; *sacchar.* lbij. Mix the alcohol and water; pour f3iv on the rhubarb, mixed with an equal bulk of sand, and allow the whole to stand four hours; put the mass into a percolator, and gradually pour on the alcohol and water. By means of a water-bath, evaporate to f3xiiij; add the sugar, and make a syrup. *Ph. U. S.*) Dose to a child, as a laxative, one or two teaspoonfuls.

SYRUPUS RHEI AROMATICUS, *Aromatic or Spiced Syrup of Rhubarb.* (*Rhei cont.* 3iiss; *Caryophyll.* cont., *Cinnam.* cont. aa 3ss; *Myrist.* cont. 3ij; *Alcohol. dilut.* Oij; *Syrup* Ovj. Macerate the rhubarb and aromatics in the diluted alcohol for 14 days, and strain; then, by means of a water-bath, evaporate the liquor to a pint,

and while still hot, mix with the syrup previously heated. Ph. U. S.) It may also be made by the process of displacement.

The aromatic syrup of rhubarb is a gentle cathartic. Dose to a child, fʒj; to an adult, fʒj.

SYRUPUS DE RHEO, Syrupus de Cichorio compositus.

SYRUPUS RHEO'DOS, *S. de papaverē errat'ico seu papaveris erratici*, Syrup of Red Poppy, (F.) Syrup de coquelicot. (Rhead. petal. recent. lbj, aq. fervent. fʒxvij, sacch. purif. lbiss. To the water, heated in a water-bath, add the petals. Remove the vessel; macerate for 12 hours; express; defecate, and add sugar to form a syrup. Ph. L.) Chiefly used to give colour.

A factitious syrup is often made of aloes, dissolved in simple syrup.

SYRUPUS ROSÆ, *S. rosa'rum soluti'vus seu e rosæ ciccis seu rosæ centifolia seu rosæ'ceus soluti'vus*, Syrup of Roses, (F.) Syrup de roses pâles. (Rosæ centif. petal. exsic. ʒvij, sacch. purif. lbvj, aq. fervent. Oiv. Macerate the petals for 12 hours. Evaporate the strained liquor to lbiss. Add the sugar. Ph. L.) Gently laxative. Dose, fʒij to fʒiss, or more.

SYRUPUS ROSÆ GAL'ICÆ, Syrup of Red Roses, (F.) Syrup de roses rouges. (Rosæ Gallic. petal. exsic. ʒvij, aquæ bull. Ov, sacch. purif. lbvj; prepared like the last. Ph. E.) Mildly astringent. Added to stomachic infusions.

SYRUPUS SACCHARI, S. Simplex.

SYRUPUS SARSAPARIL'Æ, *S. Sarsæ*, S. Sarsæ, Syrup of Sarsaparilla. (Rad. Sarsap. cont. lbj, aquæ bullient. cong., sacch. purif. lbj. Macerate the root for 24 hours; boil to Ovj. Strain while hot; add the sugar, and boil. Ph. L.) Used in the same cases as the root. Dose, fʒj to fʒiv, and more.

SYRUPUS DE SARSAPARIL'Â ET SENNÂ Compositus, Compound Syrup of Sarsaparilla and Senna, (F.) Syrup de Cuisinier. (Rad. Sarsap. concis. lbj. Infuse for 24 hours in warm water Ovj; boil for a quarter of an hour; pass with expression, and boil again the residuum in O of water. Reduce this to Ovj, and repeat the operation 2 or 3 times. Mix all the liquors and boil them together with flowers of borage, white roses, senna leaves and aniseed, each ʒij; reduce to half; pass, and add honey lbj. Boil to a syrup. Ph. P.) It is considered to be alterant, depurative, and sudorific. Dose, fʒss to fʒij.

A similar syrup, under the name **SYRUPUS SARSAPARIL'Æ COMPOSITUS**, Compound Syrup of Sarsaparilla, is directed in the Pharmacopœia of the United States. (Sarsaparill. cont. lbj; Lign. guaiac. rasur. ʒij; Rosæ centifol., Sennæ, Glycyrrhiz. cont. ʒā ʒij; Ol. Sassafras. Ol. Anisi. ʒā ʒvj; Ol. Gaultheria m. iij; Alcohol. dilut. Ox.; Sacchar. lbvij. Macerate the sarsaparilla, guaiacum wood, roses, senna, and liquorice root in the diluted alcohol for 14 days; express and filter. Evaporate the tincture in a warm bath to four pints; filter, add the sugar, and form into a syrup. Lastly, having rubbed the oils with a small quantity of the syrup, mix them thoroughly with the remainder. Ph. U. S.)

It may also be made by the process of displacement.

SYRUPUS SARSÆ, S. Sarsaparillæ — s. Sarzæ, S. Sarsaparillæ.

SYRUPUS DE SCAMMO'NIO, Syrup of Scammony. (Pulv. Scammon. ʒss, sacch. alb. ʒiv, alcohol. (s. g. 985 to 915) fʒvij. Heat the mixture moderately: then set fire to the alcohol with a lighted taper; agitate constantly, until the flame is extinguished. Suffer it to cool: pass through cloth, and add Syrup of Violets, fʒiv. Ph. P.)

ʒj contains 18 grains of scammony. Dose, fʒj to fʒss.

SYRUPUS SCILLÆ, *S. scillit'icus seu scillæ marit'ime*, Syrup of Squilla. (Acet. scillæ. Oj, sacch. lbj. Ph. U. S.) Diuretic, expectorant, and emetic. Used in the same cases as the ozymel.

SYRUPUS SCILLÆ COMPOSITUS, Mel Scillæ Compositum, Compound Honey of Squill, Compound Syrup of Squill, Hicc Syrup. (Scillæ cont.; Senegæ cont. ʒā ʒiv; Antim. et Potassæ Tart. gr. xlvij; Aquæ Oiv; Sacchar. lbiss. Pour the water upon the squills and seneka, and, having boiled to one-half, strain and add the sugar. Then evaporate to three pints, and while the syrup is still hot, dissolve in it the tartrate of antimony and potassa. It may also be made by the process of displacement. Ph. U. S.)

It is often given as an emetic in cases of croup, and in catarrh, especially in children. Dose, to infants, 10 drops to a drachm, repeated every 15 minutes, if vomiting be desirable.

SYRUPUS SENEGÆ, Syrup of Seneka. (Senegæ cont. ʒiv; Aquæ Oj; Sacchar. lbj. Boil the water with the seneka to one half, and strain; then add the sugar, and form into syrup. Ph. U. S.)

It may also be formed by the process of displacement.

This is a favourite stimulant expectorant [?] with many practitioners. Dose, fʒj or fʒij.

SYRUPUS SENNÆ, *S. cassiæ sennæ*, Syrup of Senna. (Sennæ ʒij, sem. fœnicul. cont. ʒj, sacch. pur. ʒxv, aquæ bullient. Oj. Macerate the senna leaves and fennel-seeds with a gentle heat for an hour. Strain; add the sugar, and make into a syrup. Ph. U. S.) Cathartic. Dose, fʒij to fʒiss, or more.

The Parisian Codex has a Syrupus de Sennâ of a compound character, called Syrupus de pomis compositus, (F.) Syrup de pommes composé. It contains, also, borage, bugloss, &c. Dose, fʒss to fʒij.

SYRUPUS SIMPLEX, Syrupus (Ph. U. S.), S. Sacchari, Saccharum clarificatum, Liqueurum, Simple Syrup, Syrup, Common syrup of capillaire. (Sacch. purif. lbiss, aquæ Oj. Dissolve the sugar; set aside for 24 hours; remove the scum, and pour off the clear liquid.) Used to cover nauseous tastes, &c.

SYRUPUS DE SPINÂ CERVINÂ, Syrupus rhamni — s. de Stœchade communis, see Lavendula.

SYRUPUS TOLUT'NUS, S. Tolut'ni seu toluifera balsami seu balsam'icus, Balsam'ic syrup, Syrup of tolu. (Tinct. Tolut. fʒiss; Aquæ Oj; Sacchar. lbiss. Mix the tincture with the sugar in coarse powder; expose it on a shallow dish to a gentle heat, until the alcohol has evaporated; pour the water upon it; heat gradually until the sugar is dissolved, and strain. Ph. U. S.) Use— to give flavour.

SYRUPUS VIOLÆ, *S. violæ odorat'a*, S. violarum, S. e succo violarum, Syrup of Violets. (Flor. recent. viol. odorat. lbj, aq. bull. Oviij, sacch. pur. lbxv. Macerate, in a covered vessel, for 24 hours; pass, without expression, through linen; add the sugar so as to form a syrup. Ph. E.) Very gently laxative. Dose, fʒj to fʒij.

SYRUPUS E SUCCO VIGLARUM, S. violæ.

SYRUPUS ZINGIBERIS, *S. amo'mi zingiberis*, Syrup of Ginger. (F.) Syrup de gingembre. (Tinct. Zingib. fʒiv; Aquæ Oiv; Sacchar. lbx. Mix the tincture with four pounds of the sugar in coarse powder; and expose it in a shallow dish to a gentle heat until the alcohol has evaporated. Add the rest of the sugar, and then the water in a covered vessel; heat gradually until the sugar is dissolved, and strain. Ph. U. S.) To the strained liquor add the sugar, and form a syrup.

N. B. The Parisian Codex contained a few other compound syrups besides those enume-

rated: amongst these is a *Syrup of calf's lungs*. *Syrupus e pulmonibus vitulinis*, (F.) *Syrup de mon de veau*, as an expectorant.

These syrups are made according to the usual rule, and possess the virtues of the substances after which they are named.

SYS, Sym.

SYSPA'SIA, from *συσπαι*, (*συν*, and *σπαι*, 'I draw,') 'I contract.' Clonic spasm, with diminished sensibility and inability of utterance. A genus in Good's Nosophy, which includes convulsion, epilepsy, and hysteria.

SYSPASIA CONVULSIO, Convulsion—s. Epilepsia, Epilepsy—s. Hysteria, Hysteria.

SYSSARCO'SIS, from *συν*, 'with,' and *σάρξ*, *σάρκος*, 'flesh.' *Commissu'ra ossium car'nea*, *Symph'ysis ossium muscula'ris*, *Connex'io ossium carno'sa*, *Car'na'tio*, *Concar'na'tio*. Union of bones by means of muscles; such as the union of the os hyoides with the lower jaw, of the scapula with the ribs, &c.

SYSTALCIS, Systole.

SYSTAL'TIC, *Systal'ticus*, *Peristal'tic*; from *συστέλλω*, (*συν*, and *στέλλω*,) 'I contract.' A systaltic motion in a part is one in which there is alternate contraction and dilatation. See Systole.

SYSTATICA, from *συνιστημι*, (*συν*, and *ιστημι*, 'I place,') 'I associate.' Diseases affecting several or all of the sensorial powers simultaneously; irritation or inertness of mind, extending to the muscles, or external senses; or of the muscles, or external senses, extending to the mind;—the 4th order, class *Neurotica*, of Good.

SYSTEM, *Syste'ma*, from *συν*, 'together,' and *ιστημι*, 'I place.' This word, taken in a good sense, means the collection and arrangement of different matters having an analogy to each other, or working towards the same end. In natural

history, it means a methodical arrangement of objects for the purpose of facilitating their study. System is, however, sometimes used in an unfavourable sense in physical science, and often means,—a purely gratuitous supposition, by which the march of nature is attempted to be universally explained, and to which every thing is made to bend. In anatomy, system is an assemblage of organs, composed of the same tissues, and intended for similar functions; for instance, the nervous system, muscular system, &c.

SYSTEMA, System—s. Vasorum, Vascular system.

SYSTEMATISM. A method, according to which every fact of a science is collected around an opinion, true or false. One who reduces things to such a method is called a *Sys'tematist*.

SYSTEMATIST, see Systematism.

SYSTEMATOLOGY, *Systematolog'ia*, from *συστημα*, 'system,' and *λογος*, 'a discourse.' The doctrine or a treatise of systems.

SYSTÈME VASCULAIRE À SANG NOIR, see Vascular—s. *Vasculaire à sang rouge*, see Vascular.

SYSTEMIC. Belonging to the general system—as 'systemic circulation' in contradistinction to the 'pulmonic.'

SYS'TOLE. Same etymon as Systaltic. *Systal'sis*, *Constric'tio*, *Contractio* seu *Angusta'tio* seu *Submissio* seu *Mica'tio cordis*. The movement of constriction or contraction of the heart, to give impulse to the blood, or to cause its progression. It is opposed to the *Diastole* of the organ.

SYS'TOLIC, *Systol'icus*, (F.) *Systolique*; same etymon. Relating to the systole of the heart,—as 'systolic murmur.'

SZYGIUM NERVORUM, *Par nervorum*.

T.

T BANDAGE, *T fas'cia*, *Fas'cia T formis* seu *Heliodo'ri*, *Cru'cial Bandage*. A bandage shaped like the letter T. It consists of a strip of linen, of variable length and breadth, the middle of which is attached, at right angles, to the extremity of another strip, also of variable length and breadth. When two such strips are attached, at a distance from each other, the bandage is called a *double T*. This bandage is used, principally, for supporting the dressings after the operation for fistula in ano, in diseases of the perinæum, groin, anus, &c.

T FASCIA, T Bandage.

TABAC, *Nicotiana tabacum*—*t. des Savoyards*, *Arnica montana*—*t. des Montagnes*, *Arnica montana*—*t. des Vosges*, *Arnica montana*.

TABACUM, *Nicotiana tabacum*.

TAB'ANUS, called, by the *voyageurs*, *Bulldog*. A troublesome fly in the Hudson's Bay territory, which makes a perforation like a leechbite, from which it sucks the blood.

TABARZET, *Saccharum album*.

TAB'ASHEER, *Tabaschir*, *Tabazir*. A substance found in the knots of the bamboo, in the East Indies, as a consequence of a morbid condition of the part. It is distinguished for its great proportion of silicic acid, in combination with a small quantity of mineral and vegetable matters; as well as for its slight refractive property. It has been used in obstruction of the bowels. The

sweet juice of the bamboo stalks has also been called *Tabasheer*.

TABEFACIO, Emaciation.

TABEFFY, Emaciate.

TABEL'LA, *Tab'ula*, *Tabula'tum*, *Lozan'ga*, *Lozan'gia*, *Mor'sulus*, *Morsel'lus*, *Lozenge*, *Tes'sel'la*, (F.) *Tablette*, 'a small table.' A solid medicine, composed of a powder, sugar, and mucilage, *Electua'rium sol'idum*; and generally shaped like a disk. See *Pastil*, and *Troch*. The three comprise the *Saccharol'es solid'es*.

TABELLA, Table.

TABELLÆ ABSORBENTES, T. de Mag-nesiâ.

TABELLÆ AC'IDI OXAL'ICI, T. ad sitem, *Lozenges of Oxalic Acid*. (*Acid. oxalic. 3j*, *sacch. alb. 3ss*, *ol. limonis*, *gtt. xvij* seu *gtt. xij*. Make into lozenges with mucilage of gum tragacanth. *Ph. P.*) Lozenges are made in the same way with the citric and tartaric acids. They are refrigerant.

TABELLÆ DE ALTHE'Â, *Lozenges of Marsh-mallows*. (*Rad. althææ pulv. 3jss*, *sacchar. alb. pulv. 3jss*; f. massa. *Ph. P.*) Demulcent; expectorant.

TABELLÆ ANTIMONIA'LES KUNCKE'LII, T. de stib'i sulphure'to, *Lozenges of Sulph'uret of Ant'i-mony*, *Mor'suli antimonial'es seu stibi'i seu stibi-a'ti*, *Trochisci stibiati*. (*Amygdal. dulc. 3j*, *sacch. Ross. sem. cardam. cinnam. pulv., sulph.*

antim. præpar., āā 3j; flant tabellæ. *Ph. P.* Each to weigh 10 grains.) Recommended in cutaneous diseases, blennorrhagia, arthritic pains, &c. Dose, 4 or more.

TABELLE AD ARDOREM VENTRICULI, Trochisci carbonatis calcis — t. Cardialgiæ, Trochisci carbonatis calcis.

TABELLE DE CAT'CHU ET MAGNE'SIÂ, Lozenges of Catechu and Magnesia, (F.) *Tablettes de catehou et de magnésie.* (P. catech. 3vj, p. magnes. 3iv, p. cinnam. 3iij, sacch. alb. lss, g. tragac. gr. xij, aq. cinnam. q. s. ut f. massa. *Ph. P.* Each to weigh 12 gr.) Antacid, absorbent, and stomachic. Dose, 5 to 10 lozenges.

TABELLE CINCHO'NÆ, T. de kinâ kinâ, *Bark Lozenges.* (Ext. cinchon. sicc. 3ss, sacch. 3iv, cinnam. 3ss, mucilag. g. trag. q. s. ut f. massa. *Ph. P.*) Tonic.

TABELLE DE CITRO ET DIACARTHAMO, T. de Scammonio et Sennâ.

TABELLE DE FERRO, *Steel lozenges, Tabulæ martia'les seu instant'les seu antichlorot'icæ.* (Ferri limat. porphyria. 3ss, p. cinnam. 3j, p. sacch. 3v, g. trag. et aq. cinnam. q. s., ut fiat massa. Each lozenge to weigh 12 gr. *Ph. P.*) Tonic, stomachic. Dose, v to x.

Aromatic lozenges of steel. These consist of sulphate of iron, with a small proportion of the Tincture of Cantharides.

TABELLE DE KINÂ KINÂ, T. cinchonæ.

TABELLE DE MAGNESIÂ, T. absorbent'les, *Magnesia lozenges.* (Magnes. 3j, sacchar. alb. 3iv, mucil. g. tragac. q. s. *Ph. P.*) Absorbent and antacid.

TABELLE DE RHEO, *Rhubarb lozenges.* (P. rhej. 3ss, sacch. 3v, mucilag. gum. tragac. q. s.; f. massa. Each to contain 12 grains. *Ph. P.*) Stomachic and laxative.

TABELLE DE SCAMMONIO ET SENNÂ COMPOS'ITE, T. de citro et diacarthamo, *Compound lozenges of scammony and senna.* (Scammon. 3iij, Sennâ 3ivss, rhej. 3iss, caryoph. 3j, corticis limon. confect. 3j, sacch. 3vj; f. massa. *Ph. P.*) Purgative. Dose, 3ij to 3vj, or more.

TABELLE AD SITEM, T. acidi oxalici — t. ad Sodam, Trochisci carbonatis calcis — t. de Stibii sulphureto, T. antimonialis Kunckelii.

TABELLE DE SULPHURE SIM'PLICES, *Simple sulphur lozenges.* (Sulphur lot. 3ss, sacch. 3iv, muc. g. tragac. cum aquâ rosâ præpar. q. s. *Ph. P.*) Expectorant, demulcent, &c. Used in diseases of the chest, and in cutaneous affections. Dose, a lozenge.

TABELLE DE SULPHURÊ COMPOS'ITÆ, *Compound sulphur lozenges.* (Sulph. lot. 3ij, acid. benzoic. gr. xij, rad. irid. p. 3ss, ol. anisi, gtt. viij, sacch. 3vss, g. trag. q. s. *Ph. P.*) Aperient, expectorant, and stimulant [?]. Dose, 4 a day.

TABERNA MEDICA. Pharmacopolium.

TABERNÆMONTANA U'TILIS, (after Tabernæmontanus, a German botanist.) A tree of British Guiana, of the family Apocynaceæ, called, by the natives, *Hya-Hya*, the bark and pith of which yield a milk, which is used by the savages as a refreshing drink.

TABES, *Marasmus Tabes, Tabum*, from *tabere*, 'to decay'; *Te'kedon*, *Atroph'ia*, *Tab'idness*, *Decline*, *Wasting*. Emaciation of the whole body, with general languor, hectic fever, and, for the most part, depressed spirits.

TABES, Atrophy, Phthisis — t. Coxaria, Ischiophthisis — t. Diabetica, Diabetes mellitus — t. Diuretica, Diabetes mellitus.

TABES DORSALIS, T. Dorsud'lis, *Marasmus tabes dorsa'lis*, *Phthisis no'tias seu dorsa'lis seu sicca*, *Myelophthi'sis*, *Myelatrophi'ia*, *Myelanol'is*, *Tabes ossis sacri*. A wasting of the body, attended at first with a pain in the back or loins, and afterwards in the neck or head, — caused, it

has been conceived, by too early or too frequent addition to venery.

TABES GLANDULARIS, T. mesenterica, Scrofula — t. Ischiadica, Ischiophthisis — t. Lactea, Atrophia lactantium, see Galactia.

TABES MESENTER'ICA, T. glandula'ris seu scrofulosa, *Atroph'ia mesenter'ica seu infant'um seu glandula'ris seu infant'is seu dorsi*, *Parabys'ma mesenter'icum*, *Ma'cies Infant'um*, *Pædatrophi'ia*, *Scrofula seu Rachial'gia seu Physco'nia mesenterica*, *Myelophthi'sis sicca*, *Febria hec'tica infant'um*, *Phthisis mesara'ica*, *Mesenterophthi'sis*, *Mesenter'ic Disease*, (F.) *Carreau*, *Chartre*, *Scrofules* ou *Écrouelles mésentériques*, *Étiëie* ou *Rachialgie* ou *Atrophie* ou *Physconie mésentérique*, *Entéro-mésentérite*, *Tubercules des ganglions mésentériques*, *Phthisis mésentérique*. An engorgement and tubercular degeneration of the mesenteric glands, followed by emaciation and general disorder of the nutritive functions. The disease is often owing to inflammation or irritation of the lining membrane of the intestines, giving occasion to enlargement of the glands of the mesentery, — as injury to the upper or lower extremities causes inflammation, and enlargement of the lymphatic glands of the axilla and groin. It occurs, particularly, in children of a scrofulous diathesis; and in those who are weaned too early, or fed on indigestible substances. Diarrhoea, emaciation, anorexia, or immoderate appetite; hardness and swelling of the abdomen; and, towards the end, hectic fever, are the chief symptoms of this affection; the prognosis of which is unfavourable. On opening the bodies of those who die of the disease, the mesenteric glands are found disorganized, tubercular, or in a more or less advanced state of suppuration. The treatment is the same as in scrophulosis. Frictions may be added over the abdomen and spine; and mercury, in small doses, so as to excite a new action in the system. The food must be easy of digestion, and the bowels be kept soluble.

TABES NUTRICUM, Atrophia lactantium, see Galactia — t. Ossis sacri, *Tabes dorsalis* — t. Pectoræa, *Rachitis* — t. Pulmonalis, *Phthisis pulmonalis* — t. Scrofulosa, T. mesenterica — t. Senum, *Marasmus senilis*.

TABESCERE, Emaciate.

TABID, *Tab'idus*, (F.) *Tabide*. Same etymon as *Tabes*. Relating or appertaining to *tabes*. Wasted by disease.

TABIDNESS, *Tabes*.

TABIFIC, *Tabif'icus*, (F.) *Tabifique*, from *tabes*, 'consumption,' and *facere*, 'to make.' That which causes consumption or phthisis.

TABLE, *Tab'ula*, *Tabell'a*, *Tabula'tum*, *Lam'ina*, *Paries*, *Squama*. A name given to the plates of compact tissue, which form the bones of the cranium. Of these, one is external; the other internal, and called *Tab'ula vi'trea*, on account of its brittleness.

TABLES DE MORTALITÉ, Mortality, bills of.

TABLET, Trochiscus — t. Votive, *Tabula votiva*.

TABLETTE, *Tabella*.

TABLETTES DE CACHOU ET DE MAGNÉSIE, *Tabellæ de catechu et magnesiâ*.

TABLIER, *Ventrale cutaneum*.

TABOURET, *Thlaspi bursa*.

TABULA, *Tabella*, *Table*.

TAB'ULA VOTI'VA, *Votive tablet*. A tablet, hung up as an offering — *anathe'ma* — in the ancient temples by the sick, descriptive of their diseases, and mode of cure; or by those who had been preserved from some calamity.

TABULÆ ANTICHLOROTICÆ, *Tabellæ de ferro* —

t. *Instaurantes*, *Tabellæ de ferro*—t. *Martiales*, *Tabellæ de ferro*.

TABULATUM, *Tabella*, *Table*.

TABUM, *Ichor*, *Sanies*, *Tabes*.

TAC, *Influenza*.

TACAMAHACA, *Fagara octandra*—t. *Caragna*, *Caranua*.

TACCA OCEANICA, see *Arrowroot*, (*Tahiti*.)

TACHE, *Macula*.

TACHE CÉRÉBRALE, *Tache méningitique*.

A red mark or stain, observable in a large number of hydrocephalic children, when pressure is made on the skin.—*Trousseau*.

TACHE EMBRYONNAIRE, 'Embryonic spot,' *Cu'mulus*, *C. germinativus* seu *prolig'erus*, *A'rea germinativa*, *Nu'cleus cicatriculæ* seu *blastoderm'atis*, *Ger'minal cu'mulus*. A name given by M. Coste to a spot seen on the blastodermata about the end of the first week, from which the development of the embryo proceeds.

According to Bischoff, the blastodermic vesicle consists here of a double layer, (F.) *Feuillet*—the upper one the *serous* or *animal*; the lower, the *mucous* or *vegetative*; and between these is afterwards formed the *vascular layer*.

TACHE GERMINATIVE, see *Molecule*—t.

Mélanienne, see *Nævus*—t. *Méningitique*. *Tache cérébrale*—t. *de Naissance*, *Nævus*—t. *Sanguine*, see *Nævus*.

TACHENCAUSIS, *Combustion*, *human*.

TACHES HÉPATIQUES, *Chloasma*.

TACHES LAITEUSES, *Plaques lacteuses*, 'milky spots or patches.' White spots occasionally seen on the heart, and regarded by many pathologists as the results of previous pericarditis.

TACHES DE ROUSSEUR, *Ephelides*—t. *Rouges*, *Spots*, *red*.

TACHÊTE DE ROUSSEURS, see *Ephelides*.

TACHYIA'TER, *Tachyia'trus*. Same etymon as the next. One who cures speedily.

TACHYIATRIA, from *ταχως*, 'quick,' and *ιατρεία*, 'healing.' The art of curing quickly.

TACHY'TOMY, *Tachytom'ia*, from *ταχως*, 'quick,' and *τομή*, 'incision.' The art of cutting or operating rapidly.

TACITURNITY, *Taciturn'itas*, *Sigê*, 'silence,' from *tacere*, *tacitum*, 'to be silent.' 'State of a person who does not speak.' Morbid taciturnity sometimes exists as a symptom of the neuroses, especially of melancholy.

TACT, *Tactus*, from *tangere*, *tactum*, 'to touch.' A modification of the sense of touch. Passive judgment by the cutaneous nerves regarding bodies exterior to the skin. Tact is passive: Touch, active.

TACTILE, *Tact'ilis*; same etymon. Relating or appertaining to touch.

TACTION, Touch.

TACTUS, *Tact*, *Touch*—t. *Dolorificus*, *Haptodysp'horia*—t. *Eruditus*, see *Palpation*.

TAE, *Digitus pedis*.

TEDA, *Teda*, *Dæa*, *daïs*, 'a torch.' A term used for a medicine employed under the form of fumigation, pessary, or troch.

TÆ'NIA, *Te'nia*, *Fas'cia*, from *ταΐα*, 'a fillet, a riband.' See *Bandage*. Also a genus of intestinal worms, characterized by an elongated, flattened, and articulated body, with four suckorial oscula on the head. Of this genus, two only inhabit the human body—the *Tænia lata*, (see *Bothriocephalus latus*), and *Tænia solium*.

TÆNIA ACEPHALA, *Bothriocephalus latus*—t. *à Anneaux courts*, *Bothriocephalus latus*—t. *à Anneaux longs*, *Tænia solium*—t. *Articulos demittens*, *Tænia solium*—t. *Cucurbitina*, *Tænia solium*—t. *Dentata*, *Bothriocephalus latus*—t. *Grisea*,

Bothriocephalus latus—t. *Hippocampi*, *Corpora fibriata*—t. *Humana armata*, *Tænia solium*—t. *Humana inermis*, *Bothriocephalus latus*—t. *Hydatigena*, *Hydatid*—t. *Lata*, *Bothriocephalus latus*—t. *à Longues articulations*, *Tænia solium*—t. *Membranacea*, *Bothriocephalus latus*—t. *Nervosa Halleri*, *Anglion* of *Gasser*—t. *Non arme*. *Bothriocephalus latus*—t. *Osculis lateralibus geminis*, *Bothriocephalus latus*—t. *Osculis lateralibus solitariis*, *Bothriocephalus latus*—t. *Osculis marginalibus solitariis*, *Tænia solium*—t. *Osculis superficialibus*, *Bothriocephalus latus*—t. *Prima*, *Bothriocephalus latus*—t. *Sans épine*, *Tænia solium*—t. *Secunda*, *Tænia solium*.

TÆNIA SEMICIRCULARIS, *T. stri'ata*, *Gem'inum centrum semicircula're*, *Limbus post'rior cor'poris stri'cti*, *Stria cor'neæ seu semihum'eres seu termina'les*, *Fra'nulum novum*, *Fascia semicircula'ris*, (F.) *Bandelette des éminences pyriformes* (Ch.), *Bandelette*, *B. demicirculaire*, *Lame cor'née*. A medullary, grayish, semitransparent layer, situated in the groove that separates the thalamus opticus from the corpus striatum in the lateral ventricle.

TÆNIA SOLIUM, *T. os'culis marginal'ibus solitariis* seu *cucurbiti'na seu huma'na arma'ta seu artic'ulos demitt'ens seu secun'da*, *Hal'ysis solium*, *Vermis cucurbiti'nus*, *Lumbri'cus latus*, *Solium*, *Tænia sans épine* ou *à longues articulations* ou *à anneaux longs*, *Ver solitaria* ou *cucurbitin*, *Long Tape Worm*, has an almost hemispherical, discrete head; an obtuse beak; the neck full anteriorly; all the joints slightly obtuse, the anterior very short; the next almost square, and the rest oblong; the marginal foramina vaguely alternated. The *tænia solium* is commonly a few feet long; sometimes its length is enormous. It is said to have been met with 600 feet long. It principally inhabits the small intestines, and is not common. The oil of turpentine, in some form, largely exhibited, is perhaps the best remedy.

TÆNIA STRIATA, *T. semicircularis*—t. *Tarini*, *Lamina cornea*—t. *Tenella*, *Bothriocephalus latus*—t. *Vulgaris*, *Bothriocephalus latus*.

TÆNIACIDE, *Tæniafuge*.

TÆNIE COLI, see *Colon*—t. *Valsalvæ Coli*, see *Colon*.

TÆNIAFUGE, *Tæ'nifuge*, *Tæ'niacide*, *Tæ'nicide*, *Tæni'afugum*, from *tænia*, and *fugo*, 'I drive off.' A remedy calculated to expel or destroy *tænia*.

TÆNICIDE, from *tænia*, and *cædo*, 'I kill.' A destroyer of *tænia* or tapeworm.

TÆNIFUGE, *Tæniafuge*.

TÆNIOLA, *Bandelette*; diminutive of *Tænia*. Used by Gerber for a simple, band-like, or flat fibre. *Filum tæniaform'e*, seen under the microscope.

TAFFETAS AGGLUTINATIF, *Sparadrapum adhæsivum*—t. *d'Angleterre*, see *Sparadrapum adhæsivum*—t. *Épispastique*, *Sparadrapum vesicatorium*—t. *Gommé*, see *Sparadrapum adhæsivum*—t. *Vésicant*, *Sparadrapum vesicatorium*.

TAFIA, *Rum*.

TAIE, see *Caligo*.

TAILLADE, *Incision*, see *Entaille*.

TAILLE, *Lithotomy*—t. *Hypogastrique*, *Hypogastric operation* of *lithotomy*, see *Lithotomy*—t. *Postérieure*, see *Lithotomy*—t. *Recto-vésicale*, see *Lithotomy*—t. *par le Rectum*, see *Lithotomy*—t. *Sus-pubienne*, see *Lithotomy*.

TAKAMAHAKA, *Fagara octandra*.

TAKE, *Attack*.

TAKEN DOWN, see *Attack*.

TAKING, see *Contagious*.

TAL. A word formerly employed for *alkali*. It also meant the dung of the peacock.

TALC EARTH, Magnesia.

TALCUM NEPHRITICUM, Nephreticus lapis.

TALENTUM, *ταλαντον*. An ancient weight equal to 54 pounds, 2 oz., 5 dr., and 24 grains. *Poids de Marc*. See Weights and Measures.

TALEOLUS, Malleolus.

TALIACOTIAN OPERATION, Rhinoplastic.

TALINUM UMBELLATUM, see *Mistura*.

TALIPES, Kyllosis (Clubfoot) — t. Calcanens, see Kyllosis — t. Equinus, see Kyllosis — t. Valgus, see Kyllosis — t. Varus, see Kyllosis.

TALLAHATTA SPRINGS, see Alabama, mineral waters of.

TALON, see Calcanem.

TALPA, 'a mole.' *Topina'ria*, *Tupina'ria*, *Talpa'ria*, *Testudo*, (F.) *Taupe*. A tumour situated on the head, which has been supposed to burrow like a mole.

TALPARIA, Talpa.

TALUS, Astragalus, Malleolus, see Calcanem. Pessary.

TAMARA CONGA, *Averrhoa carambola*.

TAMARÆA ZECLA, Tamarindus.

TAMARIN, Tamarindus.

TAMARIN'DI PULPA, *Pulp of Tamarinds*, (*Tamarinda*, q. s. Digest them with a small quantity of water, until they become of a uniform consistence; then separate the seeds and filaments, by pressing through a hair-sieve. Ph. U. S.)

TAMARINDS, PULP OF, Tamarindi pulpa.

TAMARINDUS, *T. Indica*, *Tamarindus*, *Orypha'nicon*, *Sil'igna Arab'ica*, *Balampul'i*, *Tamaræa zecla*, *Oxyphen'icia*, *Aca'cia Indica*, *Tamarind tree*, (F.) *Tamarin*. Ord. Leguminosæ. *Sex. Syst.* Triandria Digynia. The pulp of the tamarind with the seeds connected by numerous tough strings, *Tamarindus* (Ph. U. S.), is exported from the West Indies, preserved in syrup. It is refrigerant and laxative. *Tamarind whey* is made by boiling 3ij of the pulp with Hiss of milk. When strained, it is used as a diluent in fever.

TAMARISCUS, Tamarix Gallica.

TAMARIX, *T. Gal'lica* seu *Pentan'dra*, *Tamarisc'us*. The *Tamarisk tree*. Ord. Tamariscinæ. *Sex. Syst.* Pentandria Pentagynia. The bark, wood, and leaves were formerly employed medicinally; — the former for its aperient and corroborant virtues, in obstructions of the liver; the latter in icterus, hæmoptysis, and some cutaneous affections.

TAMBOUR, *CAVITÉ DU*, Tympanum.

TAMPA BAY, see Saint Augustine.

TAMPON, see Plugging.

TAMPONNEMENT, Plugging.

TAMUS COMMUNIS, *An'pelos melæ'na* (Disco-rides), *Chiro'nia*, *Gynæcon'thê* seu *Apro'nia* (Pliny), *Bryo'nia nigra*, *Common Black Bryony*. The root is diuretic, and has been given as an antilithic. In the herb-shops of Europe, it is sold as a topical application for removing bruise marks, and in France is called *Herbe aux femmes battues*; 'herb for beaten or bruised women.'

TANACE'TUM, *T. vulga'rê*, *Tana'sia*, *Athana'sia*, *Parthe'nium mas*, *Hind-heel*, *Tansy*, (F.) *Tanaisie vulgaire*, *Herbe aux vers*, *Coq des Jardins*. Ord. Compositæ. *Sex. Syst.* Syngenesia Polygamia. The leaves and flowers of tansy have a peculiar, strong odour, and warm bitter taste. They are tonic, deobstruent [?], and anthelmintic. Dose, ʒss to ʒj. The infusion is usually drunk as tea.

TANACE'TUM BALSAM'ITA, *T. horten'sê*, *Balsam'ita mas* seu *vulga'rê*, *seu smar'colens* seu *majus*, *Costus horto'rum*, *Mentha Roma'na* seu *Saracen'ica*, *Costmary*, *Alecost*, *Cock-mint*, (F.) *Menthe coq*, *Herbe au coq*, *Grand Bruime*. A fragrant-smelling herb, somewhat resembling mint, and

formerly much esteemed as a corroborant, carminative, and emmenagogue.

TANACE'TUM HORTENSE, T. balsamita.

TANACE'TUM MULTIFLO'RUM. A South African plant, which is used as a tonic, antispasmodic and anthelmintic. It is given in flatulency, gout, amenorrhœa and dropsy: but is especially prescribed in cases of ascariæ lumbricoïdes and other entozoa. It is administered in the form of powder or infusion; the latter acting also as a diaphoretic and diuretic. It is used likewise as a resolvent and anodyne fomentation.

TANASIE VULGAIRE, Tanacetum.

TANASIA, Tanacetum.

TANGÊ, *ταγγη*, 'rancidity.' A putrescent tumour.

TANGHINIA, *Cerbera tanghin*.

TANGLES, BIG, *Pila marina* — t. Blue, *Gaylussacia dumosa*.

TANNINUM, Tanninum.

TANNIC ACID, Tannin.

TANNICUM PURUM, Tannin.

TANNIN, from (F.) *tanner*, 'to tan.' *Tan'nicum purum*, *Tanni'num*, *Tan'num*, *Princip'ium Adstringens* seu *Scytodeph'eicum* seu *Byrodeph'eicum* seu *Coria'ceum*, *Scytodeph'eicum*, *Mate'ria Scytodeph'ica*, *Acidum Tan'nicum* seu *Scytodeph'icum* seu *Quercitan'icum*, *Tannic acid*, (F.) *Acide tannique*. A principle obtained from oak bark and other astringent vegetables, so called from its forming the principal agent in the process of tanning. Pure tannin, obtained from galls, by the action of ether. (see Ph. U. S.) has been employed as an excellent and powerful astringent. Dose, two grains, repeated several times a day.

TANSY, Tanacetum — t. Wild, *Potentilla anserina*.

TAPER, HIGH, *Verbasum nigrum*.

TAPE'TUM, *Tapes*, *Tape'te*, 'tapestry, a carpet.' *Tape'tum choroidæ*. A shining spot, on the outside of the optic nerve, in the eyes of certain animals, which is owing to the absence of the pigmentum nigrum, occasioning the reflection of a portion of the rays from the membrana Ruyschiana. Its use appears to be, by reflecting the rays, to cause a double impression on the retina, and thus add to the intensity of vision.

TAPETUM, see Corpus Callosum.

TAPETUM ALVE'OLI, *Membr'na exter'na den'tium*. The outer membrane of the teeth. The lining membrane or periosteum of the alveoli.

TAPEWORM, BROAD, *Bothriocephalus latus* — t. Long, *Tænia solium*.

TAPIOCA, see *Jatropha manihot* — t. Pearl, see *Jatropha manihot*.

TAPSUS BARBATUS, *Verbasum nigrum*.

TAPSARIA, *Decoctum hordei*.

TAR, see *Pinus sylvestris* — t. Barbadoes, *Petroleum*.

TAR BEER, see *Pinus sylvestris*.

TAR, MINERAL, *Pissaspphaltum*.

TAR WATER, see *Pinus sylvestris* — t. Wine, see *Pinus sylvestris*.

TAR'ACME, *Tarax'is*, *Tarag'ma*, *Tarag'mus*, 'disorder.' A disorder of sight which arises from a blow, compression of the eye, or the action of dust, smoke or other external cause. By some, it is understood to signify mild or incipient ophthalmia.

Tarax'is της κοιλίας, 'disturbance of the belly.' Diarrhœa.

TARAGMA, Tarache.

TARAGMUS, Tarache.

TARAGON, *Artemisia dracunculæ*.

TARAHU, *Peltandra Virginica*.

TARANTISM'US, *Tarentis'mus*, *Tarantalis'mus*, *Diomæ'nium*, *Choræ'a Sancti Valentini*, *Tarentulism*, *Tar'antism*, from *Tarentum*, or *Taranto*, where it was observed. A feigned or

imaginary disease in Apulia, characterized by excessive avidity for dancing at the sound of instruments, and which was ascribed, by the vulgar, to the bite of the tarantula. A sort of tarantismus, considered peculiar to Africa, and believed to be spontaneous, has been called *Tarentismus Tangitanius*, *Janon-tarentime*.

TARANTULA. Same etymon. *Taren'tula*, (F.) *Tarentule*. A species of spider—the *Aranea Tarentula* of Linnaeus, *Lycosa Tarentula* of Latreille—whose bite has been looked upon as dangerous. Music has been thought to cure it.

TARASCON, MINERAL WATERS OF. Chalybeate springs in the department of Arriège, France. They contain iron and carbonic acid.

TARAXACUM, *Leontodon taraxacum* — t. *Dens leonis*, *Leontodon taraxacum*.

TARAXIS, *Tarache*, see *Ophthalmia*.

TARCHON SYLVESTRIS, *Achillea ptarmica*.

TARENTISMUS, *Tarantismus* — t. *Tangitanus*, *Tarantismus*.

TARENTULA, *Tarantula*.

TARENTULISM, *Tarantismus*.

TARGAR, *Juniperus communis*.

TARI. Palm or cocoa wine, from which sugar was obtained, called *Jagre*. It was once used as a tonic.

TARI'NI PONS, (F.) *Pont de Tarin*. A cineritious substance, between the corpora albicantia, which unites the crura and assists in forming the bottom of the third ventricle. The fossa, at the bottom of which it is placed, is called *Locus perforatus posterior*, *Posterior perforated space*, *Substantia seu Lamina perforata posterior*.

TARO. *Arm esculentum*.

TARSAL, *Tar'seus*, from *tarsus*, (F.) *Tarsien*. That which relates to the tarsus.

TARSAL ARTICULATIONS result from the union of the bones of the tarsus with each other.

TAR'SEA LATA (LIGAMENTA). Three ligaments, described by Winslow as stretching from the edge of the orbit to that of the tarsi, with which, he thought, they formed a complete layer of the eyelids. Zinn could not trace the ligaments all the way to the tarsi; whilst Haller doubted altogether their ligamentous texture, and Zeis regards them as merely areolar substance.

TARSEUS, *Tarsal*.

TARSO, in composition, belonging or relating to the tarsus.

TARSO-METATAR'SAL, *Tarso-metatar'seus*. That which relates to tarsus and metatarsus.

TARSO-METATARSAL ARTICULATIONS are those between the bones of the second row of the tarsus and the metatarsal bones. The term *Tarso-metatarsal* is likewise applied to ligaments connected with these joints, some of which are *dorsal*, and others *plantar*.

TARSO-MÉTATARSIPHALANGIEN DU POUCE, *Adductor pollicis pedis* — t. *Phalangiën du pouce*, *Flexor brevis pollicis pedis* — t. *Sous-phalangiën du petit orteil*, *Flexor brevis minimi digiti pedis* — t. *Sous-phalangiëtien du premier orteil*, *Flexor brevis pollicis pedis*.

TARSOPHY'MA, from *Tarsus*, and *φύμα*, 'a tumour.' A swelling of the tarsus.

TARSOR/RHAPHY, *Tarsorrhaph'ia*, from *tarsus*, 'the tarsal cartilage,' and *ρᾶφη*, 'suture.' An operation for diminishing the size of the opening between the eyelids, when it is enlarged by surrounding cicatrices.

TARSOT'OMY, *Tarsotom'ia*, from *tarsus*, and *τομή*, 'incision.' The section or removal of the tarsal cartilages.

TARSUS, *ρᾶσος*, *Planta prima seu Ras'se'ta seu Ras'se'ta pedis*, *Pe'dium*, *Claviculæ pedis nodus*. 1. The posterior part of the foot. The tarsus consists of 7 bones, arranged transversely in two rows, between which amputation is sometimes

performed. The *first or Tibial row*, (F.) *Première rangée*, *R. jambière*, consists of the astragalus and os calcis; the *second or metatarsal*, (F.) *Seconde rangée*, *R. métatarsienne*, of the scaphoides, cuboides, and three ossa cuneiformia. 2. The *Tarsi*, *Tarsal cartilages*, *Tarsal fibro-cartilages*, *Orchoi*, *Crepid'ines seu Coro'næ seu Chelæ seu Un'gulæ Palpebra'rum*, *Tharsi*, *Oil'ia*, are two small cartilaginous layers, seated in the substance of the free edge of each eyelid. The upper is larger and broader than the lower. Both commence at the bifurcated extremity of the tendon of the orbicularis palpebrarum, and pass to the opposite commissure of the eyelids. They are broader in the middle than at their extremities.

TARTAR, *Antimonium tartarizatum* (improperly), *Potassæ supertartas impurus* — t. *Calcareous*, *Potassæ subcarbonas* — t. *Crystals of*, *Potassæ supertartas*.

TARTAR EMETIC, *Antimonium tartarizatum*.

TARTAR, SALT OF, see *Potash*, and *Potassæ subcarbonas* — t. *Soluble*, *Potassæ tartas*, see *Potash*.

TARTAR OF THE TEETH, *Odontolithos*.

TARTAR'EI MORBI. A term employed by Paracelsus to designate certain diseases which he attributed to tartar.

TARTARI CRYSTALLI, *Potassæ supertartas*.

TARTAR'IC ACID, *Ac'idum tartar'icum seu tartar'um seu tar'tari essentia'lē*, *Acor tartar'icus*, *Sal tartari essentia'lē*, *Tarta'reous acid*, (F.) *Acide tartrique*. (*Potassæ bitar't. Biiss, ag. destillat. ferr.*, cong. iij. cretæ ppt. lbj, *acidi sulph. lbj*.) Boil the bitartrate with two gallons of the water, and gradually add the prepared chalk, until no more bubbles appear; let the tartrate of lime subside; then pour off the fluid and wash the tartrate of lime with distilled water, until it is tasteless. Pour upon it the sulphuric acid, diluted with a gallon of boiling distilled water; and set the mixture aside for 24 hours, occasionally stirring it. Filter and evaporate it to crystallize. (*Ph. L.*) It is refrigerant and antiseptic, but is chiefly used for making effervescent draughts with soda.

TARTARUM, *Potassæ supertartas impurus* — t. *Soluble*, *Potassæ tartas* — t. *Vitriolatum*, *Potassæ sulphas*.

TARTARUS AMMONIÆ, *Ammonie tartas* — t. *Chalybeatus*, *Ferrum tartarizatum* — t. *Crudus*, *Potassæ supertartas impurus* — t. *Dentium*, *Odontolithos* — t. *Emeticus*, *Antimonium tartarizatum* — t. *Kalioferrieus*, *Ferrum tartarizatum* — t. *Kalinus*, *Potassæ tartas* — t. *Lixivæ*, *Potassæ tartas* — t. *Martialis*, *Ferrum tartarizatum* — t. *Potassæ ferruginosus*, *Ferrum tartarizatum* — t. *Stibiatus*, *Antimonium tartarizatum* — t. *Subpotassicus*, *Potassæ supertartas* — t. *Superpotassicus*, *Potassæ supertartas* — t. *Tartarizatus*, *Potassæ tartas*.

TARTRATE D'AMMONIAQUE, *Ammonie tartas*.

TARTRE BRUT, *Potassæ supertartas impurus* — t. *Crû*, *Potassæ supertartas impurus* — t. *des Dents*, *Odontolithos* — t. *Émétique*, *Antimonium tartarizatum* — t. *Stibié*, *Antimonium tartarizatum*.

TASIS, *Extension*, *Tension*.

TASTE, *Gustus*, *Geusis*, *Geuma*, *Geuthmos*, (F.) *Goût*. One of the five external senses, which instructs us of the savours of bodies, and of which the tongue is the principal organ.

Gustation, *Gustatio*, is the exercise of this faculty. In disease, the taste frequently becomes variously vitiated. The nerves of taste are probably the glosso-pharyngeal, and, perhaps also, the lingual branch of the fifth pair.

TASTE, *Savour*.

TATZÉ. The fruit of *Myrsi'na Africa'na*, a native of Abyssinia, the Cape of Good Hope, the

Azores, and Algeria, which is used as a tænia-fuge.

TATTOO'ING. A practice, with the South Sea Islanders, of puncturing the skin, and forcing coloring matters into the punctures. Sailors in all countries are in the custom of practising it by rubbing or forcing gunpowder into the punctures. The figures, thus formed, remain through life.

TAUPE, Talpa.

TAUROCOL'LA, from *ταυρος*, 'a bull,' and *κόλλα*, 'glue.' Glue made of the ears and genitals of the bull. It was formerly much esteemed.

TAURUS, Penis, Perineum.

TAXIS, 'arrangement' from *τάσσω*, 'I order,' 'I arrange'; *Reposit'io*, *Reduct'io*, *Syntax'is*. A methodical pressure exerted by the hand on a hernial tumour, for the purpose of reducing it. This operation is generally easy in herniæ that are free from adhesions, of small size, and with a large opening into the sac; but it is very difficult and frequently impracticable in those that are adherent or strangulated. To perform the taxis, the patient should be put in the horizontal posture, and in one in which the parietes of the abdomen are relaxed. The manual procedure varies in each species of hernia. In all cases, too strong pressure of the parts must be avoided; as inflammation and gangrene might be induced. The displaced organs must be pushed into the abdomen, according to the direction of the axis of the hernial sac. The reduction is sometimes favoured by blood-letting *ad deliquium*; the warm bath; warm fomentations; tobacco; emollient glysters; and, at times, by cold applications, as ice, &c. In a case of strangulated hernia, should the taxis fail, an operation becomes necessary.

TAXO'DIUM DIS'TICHUM, *Cupressus disticha*, *Schuber'tia disticha*, *Cypress*, *Bald Cypress*; indigenous. *Nat. Ord.* Coniferae. A balsam, obtained from it, is applied to cuts and wounds. The resin from the cones is diuretic and carminative.

TAXONOMY, Biotaxy.

TAXUS BACCA'TA, *ραῖος*. The *Yew tree*, *Smilac*, *Smilax*, *Thym'alos*, (F.) *If.* Family, Coniferae. *Sex. Syst.* Dicoecia Monadelphica. The ancients believed the yew to be poisonous; but the fruit may be eaten without inconvenience. The leaves have been often recommended in epilepsy. The resin, obtained from the bark, has been called *albir*.

TAXUS CANADEN'SIS, *American Yew*, *Ground Hemlock*; indigenous. The leaves are said to be sedative.[?]

TAXWAX, see Nucha.

TAYE, Caligo.

TAYLOR'S REMEDY FOR DEAFNESS, see Allium.

TEA, Thea—t. Apple, see Apple tea—t. Bahama, Lantana camara—t. Beef, see Beef tea—t. Berried, Gaultheria.

TEA BERRY, Gaultheria.

TEA, BLANKENHEIM, see Galeopsis grandiflora—t. Brazil, Lantana pseudothea—t. Bushman's, Methescophyllum glaucum—t. Caffre, Helichrysum nudifolium—t. Chicken, see Chicken tea—t. Flaxseed, Infusum Lini compositum—t. Ginger, Infusum Zingiberis—t. Hottentots', Helichrysum serpyllifolium—t. Labrador, Ledum latifolium—t. Lemon-peel, see Citrus medica—t. Lye, Physick's, see Fuligokali—t. Marsh, Ledum palustre—t. Mexico, Chenopodium ambrosioides—t. Mint, Infusum menthae—t. Mountain, Gaultheria—t. Mutton, see Mutton tea—t. New Jersey, Celastrus—t. Oswego, Monarda coccinea—t. Red-berry, Gaultheria—t. Sassafras, Saloop—t. Slippery elm, Infusum Lini compositum—t. Soot,

see Fuligokali—t. South sea, Ilex vomitoria—t. Spanish, Chenopodium ambrosioides—t. Veal, see Veal tea.

TEAK, Paronychia.

TEAR, Sax. *tear*, *Lac'ryma*, *Lach'ryma*, *Lac'rima*, *Lac'rimum*, *Dacry*, *Dac'ryma*, *Dac'ryon*, *Humor lacryma'lis*, *Eyedrop*, *Weeping-tear*, (F.) *Larpe*. An excrementitious humour, secreted by the lachrymal gland, and poured between the globe of the eye, and the eyelids, to facilitate the motions of those parts. The tears are saline, inodorous, and colourless. They turn the syrup of violets green; and contain chloride of sodium, with excess of base; phosphates of lime and soda, and albumen, and a special organic matter called *Lac'rymine* and *Thyranine*.

TEARS, DEERS', Bezoar of the deer.

TEASEL, CULTIVATED, *Dipsacus sylvestris*.

TEASICK, Phthisis pulmonalis.

TEAT, Nipple.

TEAZEL, *Eupatorium perfoliatum*.

TECEDON, Phthisis.

TECNOCTONIA, Infanticide.

TECNOL'OGY, *Tecol'ogy*, *Tecnolog'ia*, *Tecollog'ia*, from *τεκνος* and *τεκος*, 'a child,' and *λογος*, 'a description.' A treatise on children.

TECOLITHOS, Judeus (lapis.)

TECO'MA RADICANS, *Bigno'nia radicans*, *Trumpet flower*, *Trumpet creeper*, *Virgin'ia creeper*. Indigenous; the orange and scarlet flowers appearing in July. Order, Bignoniaceæ. The leaves are acrid.

TECTARIA CALAHUALA, see Calagualæ radix—t. Ferruginea, see Calagualæ radix.

TEDA, Tæda.

TEETH, see Tooth—t. Barred, *Barrées* (*Dents*)—t. Eye, Canine teeth—t. Jaw, Molar teeth—t. Milk, see Dentition—t. Neck of the, Collum dentium—t. Shedding of the, see Dentition—t. Vitreous substance of the, Enamel of the teeth—t. Wisdom, see Dentition—t. Wit, see Dentition.

TEETHING, Dentition.

TEGIMEN, Integument.

TEGMEN, Integument.

TEGULA HIBERNICA, *Hibernicus lapis*.

TEGUMEN, Integument.

TÉGUMENT, Integument.

TEGUMENT'ARY, *Tegumenta'rius*, (F.) *Tegumentaire*. Relating or appertaining to the integument.

TEGUMEN'TUM, Integument—t. Auris, see Tympanum—t. Ventriculorum cerebri, Centre, oval, of the brain.

TEIGNE, Pityriasis, Porrigo, *Porrigo lupinosa*—t. Annulaire, *Porrigo scutulata*—t. *en Anneau*, *Porrigo scutulata*—t. Favose, *Porrigo favosa*—t. Furfuracæ, *Porrigo furfurans*—t. Nummulaire, *Porrigo scutulata*.

TEINESMOS, Tenesmus.

TEINODYN'IA; from *τείνω*, 'to extend,' and *ὄδυν*, 'pain.' Pain of the sinews or tendons.

TEINOPHLOGO'SIS; from *τείνω*, 'to extend,' and *φλογωσις*, 'inflammation.' Inflammation of the sinews or tendons.

TEINTURE, Tincture—t. *Éthérée*, see *Æthe'*. real—t. *Rubéfiante*, *Tinctura cantharidis*.

TEKEDON, Tabes.

TELA, Texture, Tissue—t. Accidentalalis Medullaris, *Encephaloid*—t. Cavernosa, *Cavernous texture*—t. Cellulosa, *Cellular tissue*—t. Cornea, *Corneous tissue*—t. Choroida, *Choroidea tela*—t. Emplastica, *Sparadrapum*—t. Ereclitis, *Erectile tissue*—t. Fibrosa, see *Fibrous*—t. Galteri, *Sparadrapum Galteri*—t. Hippocratis cribrrosa, *Cellular tissue*—t. Hyemenochondroides, *Tissue, cartilaginous*—t. Hyemenochondroides, *Tissue, cartilaginous*—t. Ichthyocollæ glutinans, see *Sparadra*

pum adhæresivum—t. Inodularis, Tissue, inodular—t. Organica, Histos—t. Vesicatoria, Sparadrapum vesicatorium.

TELAMON, Habena.

TELAMO'NES, τελαμώνες, 'bandages.' Charpie, used for wounds; or linen employed for bandages.

TELANGECTASIS, Telangiectasia.

TELANGIECTA'SIA, *Telangiect'asis*, *Telangiect'asis*, *Angiectelecta'sia*, *Hæmatotelangi'o'sis*, *Hæmotelangi'o'sis*, *Angiect'a'sia*, *Anastomo'sis aneurysmat'ica*; from *telos*, 'far,' *αγγειον*, 'vessel,' and *ectasis*, 'dilatation.' Dilatation of far or capillary vessels. Nævus maternus. Aneurism by anastomosis. According to some, fungus hæmatodes. See *Hæmatodes fungus*.

TELANGIECTASIA OCULI, Cirsophthalmus.

TELEPH'IUM, *Ulcus telephium*, *Achille'um*. The ancients gave this name to any ulcer of a very bad kind, and difficult to cure, because the wound, which Achilles inflicted on Telephus, degenerated into a similar ulcer.

TELEPHIUM, Sedum telephium.

TELLURISMUS, Magnetism, animal.

TELLURIUM, (F.) *Tellure*; from *tellus*, *telluris*, 'the earth.' A metal first met with in the gold mines of Transylvania. It is of a bluish white colour, very volatilizable; specific gravity 6.115; and oxidizable in the air and by heat.

The salts of tellurium occasion the exhalation of so disagreeable an alliaceous odour from the body, that they have not been much experimented with as therapeutical agents.

TELUM, Penis.

TEMPE, Temple.

TEMPERAMENT, *Temperamen'tum*, *Temperatura*, *Tempera'tio*, *Tempe'ries*; from *temperare*, 'to moderate,' originally from *tempus*, *temporis*, 'time,' the great temperer; *Complex'io*, *Crisis*, *Eucra'sia*. A name given to the remarkable differences that exist between individuals, in consequence of the variety of relations and proportions between the constituent parts of the body. Thus, of old, it was supposed that, according to the predominance of any one of the four humours then generally admitted,—blood, lymph, bile, and atrabillis or black bile,—the manifestations of the functions were tempered or modified so as to give rise to a sanguine, lymphatic, choleric, or atrabillious or melancholic predominance or temperament. The idea of the existence of certain of these humours has been banished; yet the influence of some, as of the blood, lymph, and bile, is still admitted by many. Too much influence has, doubtless, been assigned to temperament in the sense in which it was formerly and still is generally understood, although there can be no question that difference of organization occasions a difference in the activity of organs.

TEMPERAMENT, BILIOUS, Choleric temperament.

TEMPERANCE, *Sophros'ynê*, *Nephal'iotês*, *Sobri'etas*, *Sobri'ety*. Same etymon. Moderation, opposed to gluttony and drunkenness. The advantage of temperance, in a hygienic point of view, requires no comment.

TEMPÉRANTS, Sedatives.

TEMPERATIO, Temperament.

TEMPERATURA, Temperament.

TEMPERIES, Temperament.

TEMPLE, *Tempus*, *Re'gio Tempora'lis*, *Crotaphus*, (F.) *Tempe*. A depression, on each side of the head, between the forehead and eye anteriorly, and the ear posteriorly. The temples, distinguished into *right* and *left*, correspond to the temporal fossa on each side. The word is said to be from the Latin *tempus*, 'time,' because in this

part the hair first begins to turn white, and to indicate age.

TEMPORAL, *Tempora'lis*. That which relates to the temples.

TEMPORAL APONEUROSIS is a very strong aponeurosis, of a bluish, shining colour, which is attached to the whole of the curved line of the temporal bone; to the posterior edge of the malar bone; to the superior margin of the zygomatic arch; and forms an envelope for the temporal muscle.

TEMPORAL ARTERIES, *Crotaphite Arteries*, are several:—1. The *Temporal Artery*, properly so called, arises from the outer side of the superior part of the external carotid. It ascends between the ramus of the jaw, the meatus auditorius externus, and the parotid gland, which covers it as far as the zygomatic arch. Above this it becomes subcutaneous; and, when it has attained the middle of the temporal region, it divides into two branches; the one anterior—A. *tempora'lis fronta'lis* seu *inter'na* seu *ante'rior*; the other posterior—the A. *tempora'lis occipita'lis* seu *exter'na* seu *supe'rior*. The temporal artery, before its division, gives off, 1. Branches to the parotid; to the joint of the lower jaw; to the meatus auditorius, and to the masseter. 2. The *transverse artery of the face*. 3. The *middle temporal*, which arises beneath the cheek-bones, and traverses the temporal aponeurosis, immediately beneath that bone. II. The *Deep-seated Temporals*—A. *tempora'les profun'dæ*. These are two in number. They arise from the internal maxillary, and have been distinguished, from their position, into *anterior* and *posterior*. They send their principal divisions to the temporal muscle.

TEMPORAL BONE, *Os tem'poris* seu *tempora'le* seu *arcua'te* seu *arma'te* seu *parieta'le inferius* seu *crotaphum* seu *memen'to mori* seu *mendo'sum* seu *crotaphit'icum* seu *lapido'sum*, is seated at the lateral and inferior part of the cranium, of which it forms part; and contains, within it, the special organs of audition. It is usually divided into three portions:—1. The *squamous*, *Pars squamo'sa*, *Squama*, *Lepi'sma*, *Os tepido'i'des* seu *squamo'sum* seu *squm'i'for'me*, (F.) *Portion écailleuse* ou *squameuse*. 2. The *mastoid portion*, and, 3. The *Petrous portion*, *Petrus Proc'ess*, *Pyramis trig'ona*, *Os lapide'um* seu *lapido'sum* seu *sax'eum* seu *petro'sum* seu *lithoi'des* seu *durum* seu *prærupte rupi assimila'tum* seu *nervale*, *Pars petro'sa*, *P. pyramida'lis ossis tem'poris*, (F.) *Portion pierreuse*, *Rocher*, *Os pétreux*, *Apophyse pyramidale*, *A. pétée*.

It has, first, an *auricular* or *external face*, on which is a surface that makes part of the temporal fossa; the *zygomatic process*; the *glenoid cavity*; the *fissure of Glaser*; the *meatus auditorius externus*; the *mastoid process*; the *digastric groove*; the *canalis caroticus*; the *jugular fossa*; the *styloid process*; the *stylo-mastoid foramen*, &c. 2. A *cerebral* or *internal surface*, which corresponds to the cavity of the cranium, and on which is the *Pars petrosa*, the *hiatus Fallopi*, the *meatus auditorius internus*, the *fissure of the aquæductus vestibuli*, a *portion of the gutter for the lateral sinus*, &c. 3. A *circumference*, on which is observed the orifice of the *bony portion of the Eustachian tube*, the *aquæductus cochleæ*, &c. The temporal bone is articulated with the sphenoid, occipital, parietal, malar, and inferior maxillary bones. It is developed from five points of ossification.

TEMPORAL FOSSA, *Fossa tempora'lis*, is an excavation, observed at each side of the head. It is filled by the muscle whose name it bears, and is formed by the temporal and sphenoid, below; by the parietal and frontal bones, above; is separated, by a transverse crista, from the

zygomatic fossa; and completed, before, by the cheek-bone.

TEMPORAL MUSCLE, *Temporalis*, *Crotaphites*, (F.) *Arcadi-temporo maxillaire*, *Temporo maxillaire* (Ch.), is a broad, flattened, triangular muscle, which fills the temporal fossa. Its fibres are attached to the temporal aponeurosis and fossa; and all converge into a strong tendon, which passes under the zygomatic arch, and is inserted into the coronoid process of the lower jaw. This muscle raises the lower jaw, and applies the teeth against each other.

TEMPORAL NERVES. These are distinguished into the *superficial temporal*, furnished by the inferior maxillary branch of the fifth pair, behind the condyle of the jaw; and the *temporales profundi*, — anterior and posterior, — furnished by the same nerve at its exit from the cranium. Sömmering has given the name *temporal* to the divisions sent by the facial nerve to the temporal regions.

TEMPORAL VEINS were formerly called *Ætoi* or *Ætoi Phlebes*, *Aq'uila Venæ*.

TEMPORO-AURICULAIRE, *Attollens* aumen.

TEM'PORO-FA'CIAL, *Tem'poro-facia'lis*. Belonging to the temple and face.

TEMPORO-FACIAL NERVE, *Nervus temporo-facia'lis*. A branch of the facial nerve distributed to the face and temple.

TEMPORO-MAXILLAIRE, *Temporo-maxillary*, see Temporal muscle.

TEMPORO-MAX'ILLARY, *Tem'poro-maxilla'ris*, (F.) *Temporo-maxillaire*. That which belongs to the temporal bone and lower jaw.

TEMPORO-MAXILLARY ARTICULATION or joint of the lower jaw, is seated between the glenoid cavity and transverse process of the temporal bone on the one hand, and the condyle of the lower jaw on the other. Two thin cartilages cover the surfaces of these bones. The one is common to the transverse process, and the portion of the glenoid cavity anterior to the fissure of Glaser; the other belongs to the condyle. This articulation is furnished with an inter-articular fibro-cartilage, two synovial capsules, and two lateral ligaments: — the one internal, and the other external.

TEMPORO-MAXILLARY NERVES of Bichat are the divisions of the facial nerve, distributed to the temporal and maxillary regions.

TEMPUS, Temple — t. Intercalare, Apyrexia, Intermission.

TEM'ULENCE, *Ebri'etas*, *Ebrios'itas*, *Methys'mus*, *Meth'ysis*, *Temulen'tia*, *Tem'ulency*, *Inebria'tion*, *Intoxica'tion*, *Drunk'ness*: — from wine, *Enophyl'gia*, *Enophyl'xis*, *Vinolen'tia*, *V'no-lence*, (F.) *Ivresse*; from *temetum*, μέθυ, 'wine, any intoxicating drink.' Plater gave this name to delirium; Ettmüller, to an apoplectic condition, depending upon drunkenness; the *Apoplex'ia Temulen'tia* or dead drunkenness. Commonly, *Temulentia* is used synonymously with drunkenness, *Methē*; and is often employed in the description of diseases to indicate a state resembling drunkenness. *Mania e Temulentia* is the same as *Delirium Tremens*. The French use the term *Ivresse* for drunkenness, *Ivrognerie* for habitual drunkenness.

TEMULENTIA, *Temulency*.

TENACULA, Bone-nippers — t. Volsella, Forceps (lithotomy).

TENAC'ULUM, (*tenax*, *tenacia*, 'holding fast.') from *teneo*, 'I hold.' A kind of fine hook attached to a handle which is thrust through the parietes of a blood-vessel to draw it out and enable it to be tied.

TENACULUM, ASSALINI'S. An ingenious instrument, invented by Assalini. It consists of a for-

ceps, or double tenaculum, the points of which are fine, sharp, and bifurcated, so as to be readily received into each other when the instrument is closed on the artery, which it is by a spring. It is of great use when the surgeon has no assistant at hand. The instrument is figured in S. Cooper's "First Lines."

TENAILLE INCISIVE, Bone-nippers.

TENASMUS, *Tenesmus*.

TENDINES CORONARIE, Coronary tendons.

TEN'DINOUS, *Tendino'sus*, *Tendin'ous*, *Sin'evey*, (F.) *Tendineux*, from *tendo*, *tendinis*, 'a tendon.' Relating or appertaining to, or having the nature of tendons.

TENDO, Tendon — t. Cordis venosus, Circulus callosus Halleri.

TENDON, *Tendo*, *Pronerva'tio*, *Tenon*, from *τενω*, *tendo*, 'I stretch.' *Sinew*, *Thew*, *Leader*, (Prov.) *Guider*. A fibrous cord, more or less round, long, or flattened; of a white, pearl colour, and composed of very close parallel fibres. The tendons are attached to the bones, by one extremity; by the other, they receive the insertions of the fleshy fibres: sometimes they afford attachment to fleshy fibres at both ends; — as in the digastric muscle. The tendons must be considered as so many cords, for transmitting the motion of muscles to the bones or levers.

TENDON D'ACHILLE, Achillis tendo.

TENDON, CORDIFORM, OF THE DIAPHRAGM, Center, phrenic.

TENDONS, TWITCHING OF THE, Sub-sultus tendinum.

TENEOTOMIA, Tenotomy.

TÉNESME, *Tenesmus* — t. de l'Œsophage, Œsophagismus — t. Vésical, Bradysuria.

TENES'MUS, *Teines'mos*, *Tenas'mus*, from *τενω*, 'I stretch.' *Proct'ica* seu *Obstipa'tio* *Tenesmus*, *Proctospas'mus*, *Tinag'mus*, (F.) *Ténesme*, *Épreintes*. Frequent, vain and painful desires to go to stool; — one of the chief symptoms of inflammation of the lining membrane of the digestive tube, as of dysentery.

TENESMUS VESICÆ, Bradysuria — t. Vesicæ mucosus, Cystirrhœa.

TENETTES, Forceps (craniotomy)

TÉNIA, Tænia.

TENNESSEE, MINERAL WATERS OF. On the waters of the French Broad River, in this State, is a large, clear, thermal spring. The heat is so great that, on first going into the water, it is almost insupportable.

TENON, Tendon.

TENONT'AGRA, from *τενω*, *τενοντος*, 'tendon,' and *αγρα*, 'seizure.' A variety of gout, which affects the tendinous parts of muscles, or the tendinous ligaments which strengthen certain articulations.

TENONTOTOMIA, Tenotomy.

TENONTOTRO'TI, from *τενω*, *τενοντος*, 'tendon,' and *τροπος*, 'I wound.' They who are wounded in the tendons. — Galen.

TENOSYNI'TIS, *Tenosynovi'tis* [?], (F.) *Ténosynite*. An imperfectly formed term, from *τενω*, 'a tendon,' *syn*, and *itis*, denoting inflammation. Inflammation of the synovial sheath of a tendon.

TÉNOTOME, *Tenot'omus*; same etymon as the next. An instrument for dividing tendons, especially by the subcutaneous method.

TENOT'OMY, *Tenotom'ia*, *Teneotom'ia*, *Tenontotom'ia*, *Sec'tio* seu *Dissec'tio ten'dinum*, (F.) *Ténotomie*, from *τενω*, 'a tendon,' and *τομή*, 'incision.' The operation of dividing a tendon, often practised for the removal of luxathrus, or deviation of the joints — as in club-foot.

TENSEUR DE L'APONÉVROSE FÉMO-RALE, Fascia lata muscle.

TEN'SION, *Ten'sio*, *Tensu'ra*, *Tasis*, *Diat'asis*, *Enton'ia*, (*Entonia* is, also, used for over-tension, *Conten'sio*, *Disten'sio*, from *tendere*, *tensum*, 'to stretch.' State of a part, which is stretched, when its textures are distended by the afflux of fluids, or the accumulation of gas; or when its fibres are drawn, as it were, in an opposite direction, and are, consequently, more resisting than usual.

TENSIVE, *Tensi'vus*, *Tono'des*, (F.) *Tensif*. Same etymon. Accompanied by tension or by a feeling of tension—as a 'tensive pain.'

TENSOR CHOROIDEÆ, Ciliary muscle—*t. Palati*, *Circumflexus musculus*.

TENSOR TARSII. A small muscle, belonging to the inner commissure of the eyelids, well described by Professor Horner, of Philadelphia, and hence termed *Muscle of Horner*. It is about three lines broad and six long; arises from the posterior flat surface of the os unguis, near its junction with the ethmoid bone, and passes forwards and outwards, lying on the posterior face of the lachrymal ducts. As it approaches the commissure of the lids, it splits into two nearly equal parts, each of which is appropriated to a duct, and inserted along its course, almost to the punctum lacrymale. Its chief office seems to be, to influence the position of the punctum lacrymale, and to regulate the course of the tears through the lachrymal ducts.

TENSOR TYMPANI, *Inter'nua Auris*, *Internus Mal'lei*, (F.) *Salpingo-malleus*. A small muscle, which arises from the petrous portion of the temporal bone, and the cartilage of the Eustachian tube, and terminates by a tendon, which is reflected over the processus cochleariformis, and is inserted into the apophysis of the handle of the malleus.

TENSOR VAGINÆ FEMORIS, *Fascia lata muscle*.

TENSURA, *Tension*.

TENSUS, *Penis*.

TENT, *Tenta*, *Turun'da*, and its diminutive, *Turun'dula*, *Penic'ulus*, *Penicil'lus*, *Penicil'lum*, *Motos*, *Lemni'cus*, *Licin'ium*, (F.) *Tente*, *Mèche*, from *tendere*, *tensum* or *tentum*, 'to stretch.' In surgery, tents are small rolls of lint, of a cylindrical or pyramidal shape, which are introduced into wounds and deep ulcers, to prevent them from closing before they are filled up from the bottom. Tents are sometimes, also, made of prepared sponge, gentian root, &c. They are not so much used as they formerly were.

TENTACULARIA, *Filaria hominis bronchialis*.

TENTACULUM CEREBELLI, *Tentorium*.

TENTA'MEN MED'ICUM. 'A medical trial.' From *tentare*, 'to try.' An examination in medicine, in the Prussian universities, which is conducted both in writing and *vivâ voce* before the Dean of the Medical Faculty.

TENTE, *Tent*—*t. du Cervelet*, *Tentorium*.

TENTIGO VENEREA, *Nymphomania*—*t. Veretri*, *Satyrasis*.

TENTIPEL'LUM, from *tendere*, *tensum* or *tentum*, 'to stretch,' and *pellis*, 'the skin.' Ancient name of a cosmetic, with which, it was pretended, wrinkles could be effaced.

TENTORIUM, *T. cerebelli* *lo super-externum*, *T. cerebelli*, *Tentac'ulum cerebelli*, *Septum enceph'ali*, *Diaphrag'ma cer'ebri*, *Intersep'ta horizonta'lia Pacchioni*, *Processus transversus duræ matris*, *Lateral Processes of the Dura Mater*, *Transverse septum*, (F.) *Tente du cervelet*, *Plancher du Cerveau*, *Septum transversæ* (Ch.), from *tendere*, *tensum* or *tentum*, 'to stretch.' A process of the dura mater, which separates the cerebrum from the cerebellum. It extends from the internal horizontal spine of the occipital bone, directly forwards to the sella Turcica of the sphenoid.

TENTORIUM CEREBELLI, *Tentorium*.

TENTUM, *Penis*.

TENTWORT, *Asplenium ruta muraria*.

TENUIS MATER, *Pia mater*.

TENUITAS AQUOSA, *Prædominium aquæ*.

TEPHRO'SIA APOLLINE'A, *Gale'ga Apollin'e'a* of Egypt and Nubia. Order, *Leguminosæ*. The leaves are often found mixed with senna.

TEPHROSIA PURPU'REA, *Gale'ga purpu'rea* of the coast of Coromandel. The root is bitter, and a decoction of it is prescribed by the native doctors in dyspepsia, lientery, and tympanites.

TEPHROSIA SENNA. The leaves are used for senna by the people of Popayan.

TEPHROSIA TOXICA'RIA, *Gale'ga toxica'ria* of Cayenne. The leaves are used for poisoning fish. It has been suggested as a substitute for digitalis.

TEPHROSIA VIRGINIANA, *Galega Virginiana*.

TEPIDARIUM, see *Stove*.

TEPLITZ, *Töplitz*.

TERAS, *Monster*.

TERATOG'ENY, *Teratogen'ia*, from *teras*, *teras*, 'a monster,' and *genesis*, 'generation.' The formation of monsters.

TERATOL'OGY, *Teratolog'ia*, from *teras*, *teras*, 'a monster,' and *logos*, 'a discourse.' A treatise on monsters.

TERATOSIS, see *Monster*.

TERCERON, see *Mulatto*—*t. Black*, see *Mulatto*.

TEREBELLA, *Trepan*.

TEREBELLUM. A name given by Dugès to a perforator, *Perce crâne*, of his invention.

TÉRÉBENTHINE, *Terebinthina*—*t. de Bordeaux*, see *Pinus sylvestris*—*t. de Chio*, see *Pistacia terebinthus*—*t. de Copahu*, *Copaiba*—*t. Commune*, see *Pinus sylvestris*—*t. d'Égypte*, see *Amyris opobalsamum*—*t. de Giléad*, see *Amyris opobalsamum*—*t. du Grand Caire*, see *Amyris opobalsamum*—*t. de Judée*, see *Amyris opobalsamum*—*t. de Mèlèze*, see *Pinus larix*.

TÉRÉBENTHINE, *Terebinthinate*.

TEREBIN'THINA, from *terpivdos*, or *terpeivdos*, 'the turpentine tree.' *Tur'pentine*, *Botin*, *Albotin*, *Altibibat*, *Brutino*, (F.) *Térébenthine*. A substance, of the consistence of honey, which flows from many trees of the terebinthaceæ and coniferæ families. It is viscid; shining; more or less transparent; inflammable; of a warm, pungent taste; strong smell; and is entirely composed of resin and essential oil, without benzoic acid. It is soluble in alcohol. See *Turpentine*, *Pinus palustris* and *Pinus sylvestris*. All the turpentine are stimulant, diuretic, and cathartic.

TEREBIN'THINA ARGENTORATENSIS, see *Pinus picea*—*t. Balsamea*, see *Pinus balsamea*—*t. Canadensis*, see *Pinus balsamea*—*t. Chia*, see *Pistacia terebinthus*—*t. Communis*, see *Pinus sylvestris*—*t. Cypria*, see *Pistacia terebinthus*—*t. Empyreumatica*, see *Pinus sylvestris*—*t. Laricina*, see *Pinus larix*—*t. Veneta*, see *Pinus larix*—*t. Vera*, see *Pistacia terebinthus*—*t. Vulgaris*, see *Pinus sylvestris*—*t. de Vénise*, see *Pinus larix*.

TEREBIN'THINATE, *Terebinthina'tus*, (F.) *Térébenthiné*. Having the qualities of turpentine.

TEREBINTHUS, *Terminthus*—*t. Gummifera*, *Bursera gummifera*—*t. Lentiscus*, *Pistacia lentiscus*—*t. Vulgaris*, *Pistacia terebinthus*.

TEREBRA, *Trepan*.

TEREBRATIO, *Trepanning*.

TEREDO OSSIIUM, *Caries*, *Spina ventosa*.

TERES, *Stron'gylus*, *Cylindroi'des* 'long and round,' from *tero*, 'to rub,' quasi 'rubbed smooth or round.' An epithet given to many organs, the fibres of which are collected into round fasciculi.

TERES LIGAMENTUM, (F.) *Ligamentum rond*.

This name is given to the round ligament in the cotyloid cavity of the os innominatum, which is attached to the head of the bone, and to the bottom of the cavity. It is, also, given to a small fibrous fascia, which extends from the osseous process of the ulna to below the bicipital tubercle of the radius.

TERES MAJOR, (F.) *Anguli-scapulo-humeral*, *Scapulo-humeral* (Ch.), *Grand rond*, is situate at the inferior and posterior part of the shoulder. It is attached, on the one hand, to the outer surface of the scapula; to the corresponding part of its axillary margin; and, on the other, to the posterior margin of the bicipital groove of the humerus. It carries the arm backwards and inwards, and carries it, also, inwards in rotation. When it acts with the longissimus dorsi and pectoralis major, it applies the arm against the lateral part of the chest, and keeps it forcibly there.

TERES MINOR, *Nonus humeri Placentini*, (F.) *Le plus petit sus-scapulo-trochitérien* (Ch.), *Petit rond*, *Margini-sus-scapulo-trochitérien*, is situate at the posterior and inferior part of the shoulder; it is narrow and flattened, from above to below, in its inner half; and, from before to behind, in the outer. It is attached, on the one part, to the outer surface of the scapula, near its inferior angle; and, on the other, to the inferior part of the great tuberosity of the humerus. It depresses the arm, and causes it to turn on its axis from within outwards. It also carries it backwards.

TERESIS, Observation.

TERETRIUM, Trepan.

TERETRON, Trepan.

TERGAL, Dorsal.

TERGEMINI, Trigemini.

TERGUM, Dorsum, Vertebral column.

TERMINALIA, see Myrobalanus.

TERMINOLOGY, Nomenclature.

TERMIN'THUS, *Terebin'thus*, from *τερεβινθος*, 'the turpentine tree.' *Phyma Anthrax*, *Termin'thus*, *Berry* or *Fungous carbuncle*. The ancients gave this name to a tumour, surmounted by a black pustule, and resembling the fruit of the turpentine tree.

TERMIN'THUS, *Pistacia terebinthus*.

TERMINUS SUPERIOR ASPERÆ ARTERIÆ, Larynx.

TERMINOLOGY, Nomenclature.

TERMS, Menses.

TERRA ABSORBENS MINERALIS, *Magnesia carbonas*—t. *Aluminis*, *Argilla pura*—t. *Amaræ atrata*, *Magnesia carbonas*—t. *Amaræ sulphurica*, *Magnesia sulphas*—t. *Argillacea hydrata*, *Argilla pura*—t. *Argillacea pura*, *Argilla pura*—t. *Bolaris*, *Argilla pura*—t. *Foliata*, *Sulphur*—t. *Foliata mineralis*, *Soda acetata*—t. *Foliata tartari*, *Potassæ acetat*—t. *Fornacum*, see *Bricks*—t. *Fullonica*, *Cimolia purpurescens*—t. *Japonica*, *Catechu*, *Nauclea gambir*.

TERRA LEM'NIA, *Lemp'nias*, *Argilla bolus flava*, *Lem'nian Earth*. A name given by the ancients to a solid, reddish, astringent, substance, prepared with the pulp of the fruit of the Baobab, an African tree, according to Prosper Alpinus. It seems, however, to be an argillaceous earth. It has been employed as an astringent.

TERRA LIVON'ICA. A *terra sigillata* or sealed earth from Livonia, redder than that from Silesia, and very astringent.

TERRA MERITA, *Curcuma longa*.

TERRA NOCERIA'NA. A whitish, soft, and astringent earth, found in the environs of Nocera, in Italy.

TERRA ORLEA'NA, *Pigmentum urucu*, *Urucu*, (F.) *Rocou*, *Roucou*, *Bichet*. A pigmental matter obtained from the seeds of *Bixa Orleana* seu *Orellana* seu *Americana*, *Orellana*, *Orleana*.

In Jamaica, it is recommended in dysentery, and is considered to possess astringent and stomachic qualities. It is called, according to its shape, *Flag*, *Roll*, or *Egg Annotto*.

Spanish Annotto, *Bixa*, is in small, oblong cakes. It is chiefly employed as a colouring matter.

TERRA OS'SEA, *Materia ossea*. Bony matter. The earthy portion of bones.

TERRA PONDEROSA, *Baryta*—t. *Ponderosa murata*, *Baryta*, *muriate* of—t. *Ponderosa salita*, *Baryta*, *muriate* of—t. *Saponaria*, *Cimolia purpurescens*.

TERRA PORTUGAL'LICA, *Barros*, *Bucaros*. A reddish, astringent, styptic earth, obtained from Portugal.

TERRA SIGILLA'TA, *Sealed Earth*. A bole or earthy matter, made into little cakes or flat masses, and stamped with certain impressions, as with the head of the Grand Seigneur. To these belong the *Terra Silesiaca*, *Terra Lemnia*, *T. Turcica*. See *Bole* and *Bolus alba*.

TERRA SILESIACA, see *Terra sigillata*—t. *Talcosa oxyanthracodes*, *Magnesia carbonas*—t. *Turcica*, see *Terra sigillata*.

TERRA URI'NÆ. The earthy deposit in the urine.

TERRA VITRIOLI DULCIS, *Colcothar*.

TERRÆ SIGILLATÆ, see *Bole*, *Bolus alba*, and *Terra sigillata*.

TERRAPIN, *Emys palustris*.

TERRE BOLAIRE, *Bolus*—t. *Magnésienne*, *Magnesia carbonas*—t. *Noix*, *Bunium bulbocastanum*—t. *Pesante*, *Baryta*—t. *Sigillée*, *Bolus alba*.

TERRETTE, *Glechoma hederacea*.

TERTIAN FEVER, *Febris tertiana*, *An'etus tertianus*, *Tritæ'us*, *Tertian* or *Third Day Ague*, *Third Day Fever*, (F.) *Fièvre tierce*, from *tertius*, 'the third.' An intermittent, whose paroxysms recur every third day, or every 48 hours. The mildest, and the most pernicious, intermittents belong to this head. As a general rule, it is the most manageable form of ague.

A *double tertian*, (F.) *Double tierce* (*Fièvre*), has a paroxysm every day,—the paroxysms of alternate days being associated.

TERTIANARIA, *Scutellaria galericulata*.

TESSARA, Cuboid.

TESSELLA, *Tabella*.

TEST, **DANIEL'S**, see *Docimasia pulmonum*—t. *Hydrostatic*, see *Docimasia pulmonum*—t. *Plouquet's*, see *Docimasia pulmonum*—t. *Static*, see *Docimasia pulmonum*.

TESTA, ('a shell,') see *Ostrea*—t. *Ovi*, see *Membrana putaminis*.

TESTA PRÆPARATA, *Prepared Oyster-shell*. Oyster-shell, freed from extraneous matter, washed with boiling water, reduced to powder, and prepared in the same manner as *Creta præparata*.

TESTA'CEOUS, *Testa'ceus*, (F.) *Testacé*, from *testa*, 'a shell.' A powder, consisting of burnt shells. These contain carbonate of lime chiefly, and hence the term has been applied to cretaceous substances. See *Creta*.

TESTÆ FORNACEÆ, *Bricks*—t. *Ostreæ*, see *Ostrea*.

TESTES. The Testicles, see *Testicle*. *Vulgarly*, the *Stones*. A name, also, given to the inferior tubercles of the corpora quadrigemina, to distinguish them from the superior, called *Nates*.

TESTES CEREBRI, see *Quadrigemina corpora*.

TESTICLE, *Testis*, *T. viri'lis*, *Testic'ulus*, *Orchis*, *Pomum amo'ris*, *Didymus*, *Her'nia*, *Gem'ini* (pl.), *Gemelli* (pl.), *Ovum*, (pl.) *Testic'ule*, from *testis*, 'a witness,' because the testicles are evidences of virility. The name of two glandular organs, contained in the scrotum, whose office is to secrete sperm. The substance of the testicle

consists of numerous conical, flattened lobules — *lobuli testis* — whose bases are directed towards the surface of the organ, and the apices towards the corpus Highmorianum. They are formed of a gray and softish substance, composed of tortuous canals, called *Tubuli seu Canaliculi semiferi seu seminales, Vasa seminalia, Vascula serpentina*, folded on each other, and of extreme fineness; the number of which, according to the estimate of Monro, is 62,500, and the total length 5,208 feet. These canals, uniting in their course, pass through the corpus Highmorianum, concur in forming the epididymis, and give origin to the vas deferens. The testicles are covered, immediately, by a fibrous membrane, of an opaque white colour, and very tough, called *Tunica Albuginea*, or, simply, *Albuginea, Peritesticis*, (F.) *Pérididyme*, of which the corpus Highmorianum is only an enlargement. Over this, again, is the tunica vaginalis, the second envelope of the testis.

TESTICLE, SWELLED, Hernia humoralis.

TESTICONDUS, Cryptorchis.

TESTICULAR, *Testicularis*, (F.) *Testiculaire*. Belonging or relating to the testicle.

TESTICULE, Testicle.

TESTICULUS, Testicle — t. Accessorius, Epididymis — t. Caninus, Orchis mascula.

TESTIMONIUM MATURITATIS. 'Evidence of fitness.' A certificate of complete education, required, in the Prussian universities, of a candidate, before he can be admitted to examination for the degree of Doctor of Medicine.

TESTIS FEMINEUS, Ovary — t. Irritable, Orchialgia — t. Minor, Prostate — t. Muliebris, Ovary — t. Virilis, Testicle.

TESTUDINATIO CRANII, Camarosis.

TESTUDO, *Chelone*, (F.) *Tortue*. A genus of reptiles, including the turtle, the flesh of which is much esteemed as an article of diet.

Also, an encysted tumour, which has been supposed to resemble the shell of a turtle; *Emphyma encystis ganglion*. See Talpa.

TESTUDO CEREBRI, Fornix — t. Scapulae, see Scapula.

TET, Nipple.

TET'ANIC, *Tetan'icum, Spas'ticum, Convulsivum, Convulsivae*, (F.) *Tétanique*, from *teino*, 'I stretch.' A remedy, which acts on the nerves, and, through them, on the muscles, occasioning, in large doses, convulsions. The chief agents of this class are *Nux Vomica, Strychnia, Brucia, Arnica*, and *Toxicodendron*.

Also, an epithet for a tonic convulsion.

TETANINE, Strychnia.

TETANOMATA, Tetanothra.

TETANO'THRA, *Tetanom'ata, Erugato'ria*, from *teravos*, 'free from wrinkles.' Medicines which remove wrinkles. — *Gorræus*.

TÉTANOS, Tetanus.

TET'ANUS, from *teino*, 'I stretch.' *Spasmodic rigidity, Convulsio In'dica, Holoton'ia, Holoton'icus, Holotet'anicus, Tet'anus universalis, Rigor nervosus seu nervorum, Extensio seu Distensio, Entesia tetanus, Cat'ochus cervinus seu holoton'icus*, (F.) *Tétanos, Mal de cerf*. A disease which consists in a permanent contraction of all the muscles or merely of some, without alternations of relaxation. It is characterized by closure of the jaws; difficulty or impracticability of deglutition; rigidity and immobility of the limbs and trunk, which is sometimes curved forwards (*Emprosthotonos*), sometimes backwards (*Opiethotonos*), and sometimes to one side (*Pleurothotonos*). When tetanus is confined to the muscles of the jaws, it is called *Trismus*. It is a most formidable affection. The means of treatment are. — copious and repeated blood-letting;

bathing, cold and warm; powerful doses of opium, and other narcotics. In *Traumatic Tetanus*, or that which supervenes on a wound, every extraneous matter must, of course, be removed from the wound, and soothing applications be made to it.

TETANUS ANTICUS, *Emprosthotonos* — t. Dolorificus, Cramp — t. Dorsalis, *Opisthotonos* — t. Lateralis, *Pleurothotonos* — t. Maxillae inferioris, *Trismus* — t. Posterganeus, *Opisthotonos* — t. Posticus, *Opisthotonos* — t. Universalis, see *Tetanus*.

TETARTÆUS, Quartan.

TETARTOPH'IA, from *terapros*, 'the fourth,' and *φω*, 'I arise.' *Quarta'nus remit'tens*. A quartan, in which the intermission is inordinately short or imperfect.

TÊTE, Caput, Head — t. de Veau, Elephantiasis of the Greeks.

TETRADRACH'MON, from *τετρας*, 'four,' and *δραχμη*, 'a drachm.' A weight of four drachms or half an ounce.

TETRAGONUS, *Platysma myoides*.

TETRAHIT LONGIFLORUM, *Galeopsis grandiflora*.

TETRAMY'RON, from *τετρας*, 'four,' and *μυρον*, 'an ointment.' An ointment of four ingredients. — *Galen*.

TETRANGURIA, *Cucurbita citrullus*.

TETRANTHERA PICHURIM, *Pichurim* beans.

TETRA'O COTUR'NIX, *Coturnix*, the Quail, (F.) *Caille*; *Class*, Aves; *Ord*, Gallinaceæ. The quail is an esteemed article of diet. The Greek name is *ορνις, ορνυος*; and the places, called *Ortygia*, are named after it. The excrements of the quail were formerly extolled in epilepsy, and the fat in specks of the eye.

TETRAPHAR'MACUM, from *τετρας*, 'four,' and *φαρμακον*, 'a medicine.' A medicine consisting of four ingredients.

TETRAS'CEIUS, from *τετρα*, 'four,' and *σκελος*, 'leg.' A monster having four legs. — *Gurli*.

TETROB'OLON. The weight of four oboli, or two scruples. — *Gorræus*.

TETOROS, *Astragalus*.

TETER, Herpes — t. Crusted, *Impetigo* — t. Honeycomb, *Porrigio favosa* — t. Humid, *Eczema impetigo* — t. Milky, *Porrigio larvalis* — t. Pustular, *Impetigo* — t. Running, *Impetigo* — t. Scaly, *Psoriasis*.

TETTERWORT, *Chelidonium majus*.

TEUCRIUM ÆGYPTIACUM, *T. capitatum* — t. Belium, *T. capitatum*.

TEUCRIUM CAPITA'TUM, *T. Bel'ium seu Ægyptiacum, Po'lium Capitatum, Poley Mountain of Montpel'lier, Ord. Labiatae*, is, generally, substituted for *T. chamædrys*. The common *Poley Mountain* is the *Teucrium montanum*; and the *Golden Poley Mountain*, the *Teucrium Po'lium*.

TEU'CRIMUM CHAMÆ'DRYS, *T. officinale, Chamædrys, C. minor repens seu vulgare, Quercula minor seu Calamandrina, Trisa'go, Chamædrops, Triza'go, Common, Creeping or Small German, English Treacle*, (F.) *Petit chène, Chénette, Germandrée officinale*. This is possessed of aromatic and bitter properties. The dose of the dried powder is from ʒss to ʒj.

TEUCRIUM CHAMÆ'PITYS, *Chamæpitys, Arthetica, Arthetica, Aju'ga, A. seu Bu'gula Chamæpitys, Abiga, Chamæmor'is, Iva arthritica, Holoc'yon, Io'nia, Sideritis, Common Groundpine*, (F.) *Petite Ivette*. Properties like the last. The tops and leaves have been recommended as aperients and tonics; especially in female obstructions, and in paralytic disorders.

TEUCRIUM CRE'TICUM, *T. hyssopifolium seu Rosmarinifolium, Rosmarinus ste'chadis facie*,

P. Cay-mountain of Candy, Po'tium Cret'icum. The tops and whole herb enter into the old compounds, *Mithridate* and *Theriaca*. It has a moderately aromatic smell; and a nauseous, bitter taste, and is placed among the aperients and corroborants.

TEUCRIUM FLAVUM has similar properties to *T. chamædrys*.

TEUCRIUM HYSSOPIFOLIUM, *T. Creticum*.

TEUCRIUM IVA, *Chamæpitys moscha'ta, Iva, I. moscha'ta Mompelien'sium, Chamæpitys anthyllus*, French Groundpine, (F.) *Ivette musquée*. It has similar virtues to *T. Chamæpitys*, but is weaker.

TEUCRIUM MARITIMUM, *T. Marum*.

TEUCRIUM MARUM, *T. marit'imum, Marum Syriacum* seu *Cret'icum* seu *verum* seu *Cortu'si, Majorana Syri'aca, Chamædry's inca'na marit'ima* seu *marum, Orig'anum Syriacum, Marum germanicum, Syrian herb mastich, Cat thyme*, (F.) *Marum* ou *Germandrée maritime, Herbe aux chats*. A very aromatic plant, of a camphorated smell; formerly much used in medicine as a tonic, antispasmodic, emmenagogue, &c.

TEUCRIUM OFFICINALE, *T. chamædry's*—*t. Palustre, T. scordium*—*t. Pyramidale, Ajuga*—*t. Rosmarinifolium, T. Creticum*.

TEUCRIUM SCORDIUM, *T. palust're, Scordium, Trisa'go palust'ris, Chamædry's palust'ris* seu *scord'ium, Al'hum red'olens, Water Germander*, (F.) *Germandrée scorodone, Sauge des bois, Germandrée aquatique*. The leaves have a garlicky smell, and bitterish, slightly pungent taste. It has the tonic properties of the other *Teucra*.

TEXTUM INTERLOBULARE, *Interlobular tissue*.

TEXTURA, *Texture*—*t. Organica, Histos*.

TEXTURAL Same etymon as *Texture*. Relating or appertaining to a texture.

TEXTURE, *Textu'ra, Textus*, from *texere, texum*, 'to weave.' *Tela, Hypha, Hyphæ*. The particular arrangement of the tissues that constitute an organ.

TEXTUS, *Texture, Tissue*—*t. Cellulosus, Cellular tissue*—*t. Desmosus, Desmoid tissue*—*t. Erectilis, Erectile texture*—*t. Nervorum, Plexus nervorum*—*t. Organicus, Cellular tissue*—*t. Papillaris, Corpus papillare*—*t. Parenchymalis, Cellular tissue*.

THALIA, *Radzyge*.

THALAMUS, *Σάλας*, (F.) *Couche*. 'A room or chamber; a bed.' The place at which a nerve originates, or has been considered to originate.

THALAMI NERVORUM OPTICORUM seu OPTICI, *Eminent'ie magnæ cer'ebr'i, Ganglia cerebri post'ica, Crura medull'æ oblongat'æ* (of some), *Striat'a corpora super'na posterio'ra, Colliculi nervorum optico'rum, Optic thalami, Posterior cerebral ganglion*, (F.) *Couches des nerfs oculaires* (Ch.), *C. des nerfs optiques, C. optiques*. Two rounded and irregular surfaces, which are seen exposed in the lateral ventricles of the brain, and in the third ventricle, the inferior surface of which presents two projections, called *Corpora geniculat'a*, that furnish several filaments to the optic nerves. The name was given from a belief that the optic nerves originate from them. They, however, arise more posteriorly, and adhere merely to the inner margin of those bodies. Gall considers, that the thalami act as ganglions to the nerves; and, hence, he calls them *Grand ganglion cérébral inférieur*. They are, also, called *Posterior ganglion of the cerebrum*. From the thalami and corpora striata fibres proceed upwards to constitute the convolutions of the brain, and the various bands that connect the different parts of the brain together. The upper and inner parts of the thalami are so closely connected as to form one continued surface, called *Commissu'ra mollis*.

The posterior parts turn downwards and outwards; after which they are elongated to form the two white cords, termed *Tractus opt'ici*.

THALAMUS REGALIS, *Pericardium*.

THALICTRON, *Σάλιτρον, Thalictrum, T. magnum* seu *flavum* seu *flaves'cens* seu *pratens* seu *ni'gricans* seu *rugo'sum* seu *vagina'tum, Rhabar'barum pau'perum, Meadow-rue, Poor Man's Rhubarb*, (F.) *Pigamon jaunâtre, Rue des prés, Fausse-Rhabarbe, Rhubarbe des pauvres*. Ord. Ranunculacæ. *Sex. Syst. Polyandria Polygynia*. The root resembles rhubarb in its properties.

THALICTRUM ANEMONIDES, *Meadow Rue, Rue-leaved Anem'ony*; an indigenous plant, flowering in April and May.

THAMARINDUS, *Tamarindus*.

THAME, *Pollex*.

THANASIMUS, *Mortal*.

THANATODES, *Mortal*.

THANATOID, *Thanatoï'des*; from *θάνατος*, 'death,' and *ειδός*, 'resemblance.' Resembling death. Apparently dead.

THANATOLOGY, *Thanatolog'ia, Thnescolog'ia*, from *θάνατος*, 'death,' and *λογος*, 'a discourse.' A description, or the doctrine, of death.

THANATOS, *Death*.

THAPSIA, from the island Thapsus. The deadly carrot, *thapsia ascle'pias*. *Family, Umbelliferae. Sex. Syst. Pentandria Dignia*. The root operates violently, both upwards and downwards. It is not used.

THAPSUS BARBATUS, *Verbascum nigrum*.

THARSI, see *Tarsus*.

THASPIUM ACTÆIFOLIUM, *Ligusticum actæifolium*.

THASPIUM ATROPURPUREUM, *Th. cordat'm, Round heart. Order, Umbelliferae*. The flowers, which appear in June, are of a dark purple.

THASPIUM BARBINO'DE, *Meadow Parsnep*. An indigenous plant, of the Northern and Western States, which flowers in June. Flowers, yellow.

Both plants have been esteemed vulnerary, antispasmodic, and diaphoretic; and as antidotes to the bite of a rattlesnake.

THASPIUM CORDATUM, *Th. atropurpureum*.

THE, *Thigh*.

THÉ, *Thea*—*t. des Apalaches, Ilex vomitoria*—*t. du Canada, Gaultheria*—*t. Composé, Species*—*t. d'Europe, Salvia veronica*—*t. de France, Salvia*—*t. de Jersey, Celastrus*—*t. de la Mer du Sud, Ilex vomitoria*—*t. du Mexique, Chenopodium ambrosioides*—*t. de Simon Pauli, Myrica gale*—*t. Suisse, Falltranck*.

THEA, *Chaa, Tea*, (F.) *Thé*. There are two principal species of tea-plant; the *Thea Bohe'a*, and *Thea vir'idis*; the black tea, and the green. Ord. Camelliæ. *Sex. Syst. Polyandria Monogynia*. In commerce, many kinds of tea are met with. Several of the differences between these appear to result from the age at which the leaves have been collected, and the mode of their desiccation.

Tea-drinking was introduced into Europe about the year 1666; since which time its use has become almost universal; and, in spite of the assertions of medical terrorists, it is, except in particular idiosyncrasies, entirely harmless.

By an analogical transference, very common in language, the word *Tea* has been used almost synonymously with infusion, as *Beef tea, Mint tea*, &c.

THEA GERMANICA, *Veronica*.

THEBESIIUS, VEINS OF. A name given to supposititious branches of the coronary veins, which Christopher Adam Thebesius, a German anatomist, described as opening into the heart by small depressions observable in the right auricle; and which have been called *Foramina*

Thebesii. No such veins can be demonstrated. The valve, at the orifice of the coronary vein, in the right ventricle, is called *Valvula Thebesii*, *V. Guifartiana*, *V. venæ magnæ*.

THECA, Case, Vagina, Sheath—t. Cerebri, Cranium—t. Cordis, Pericardium—t. Folliculi, see Folliculi Graafiani—t. Vertebralis, see Vertebral canal.

THECITIS, from *theca*, and *itis*, denoting inflammation. A term, of hybrid formation, signifying inflammation of the sheaths of tendons.

THEIN, see *Coffea Arabica*.

THEION, Sulphur.

THEIOPEGÆ, Waters, mineral (sulphureous.)

THEIOTHERMÆ, Waters, mineral, sulphureous.

THELASIS, Lactation.

THELASMUS, Lactation.

THELASTRIA, Nurse.

THELE, Nipple.

THELITIS, (F.) *Thélite*; from *θηλη*, 'the female nipple,' 'the female breast,' and *itis*, denoting inflammation. Inflammation of the nipple.

THELONCUS, Mastoncus.

THELORRHAGIA, from *θηλη*, 'nipple,' and *ρρῆξις*, 'a rupture.' Hemorrhage from the nipple.

THELOS, Mamma.

THELYGONIA, from *θηλος*, 'pertaining to the female sex,' and *γενν*, 'generation.' The part taken by the female in the act of generation. Also, nymphomania.

THELYGONUM, Sperm (of the female.)

THELYPTERIS, Pteris aquilina.

THENAD, Thenal.

THENAL, *Thenar*; same etymon as *thenar*. Relating or appertaining to the thenar.

THENAL ASPECT. An aspect towards the side on which the thenar is situated.—Barclay. *Thenad* is used, adverbially, by the same writer, to signify 'towards the thenal aspect.'

THENAR, *Thenar*, from *τενω*, 'I strike,' 'I push.' The palm of the hand, or sole of the foot.

THENAR, Flexor brevis pollicis manus, Palm.

THENAR or THENAL EMINENCE is the projection at the anterior and outer part of the hand, formed by the abductor brevis, opponens, and flexor brevis pollicis.

THENAR or THENAL MUSCLE. Riolan and Winslow give this name to the fleshy mass, formed of the abductor brevis, opponens pollicis, and the interior part of the flexor brevis pollicis. In the foot, Winslow gives the name *Thenal muscle* to the abductor and flexor brevis pollicis pedis.

THEOBROMA CACAO, Cacao.

THEOMANIA, Demonomania.

THEOPLEGIA, Apoplexy.

THEOPLEXIA, Apoplexy.

THEORETICAL, *Theoreticus*, *Theoricus*, from *θεωρεω*, 'I contemplate.' (F.) *Théorique*, *Théorique*. That which is confined to theory, or is connected with it. An epithet also applied to a set of physicians, who founded their doctrine chiefly on reasoning.

THEORIA, Theory.

THEORIQUE, Theoretical.

THEORY, *Theoria*, (F.) *Théorie*. Same etymon. The speculative part of a science. The connexion established in the mind between a general fact, or the least possible number of general facts, and all the particular facts dependent thereon: for example, the motions of the heavenly bodies, and the most important natural phenomena are connected with a single fact, previously known by observation; viz. that the force of gravity acts inversely according to the square of the distance. This constitutes the theory of universal gravitation. *Theory* must not be confounded with *system*. *Theory* regards nature as it is, and is a rigid deduction from facts. *System* is too

often the creature of the imagination, to which nature is made to bend.

THEORY OF MEDICINE, *Institutes of Medicine*, *Theoretical Medicine*, is that part of the science which attempts philosophically to account for the various phenomena that present themselves during health as well as in disease. It is the philosophy of Medicine. The *Institutes of Medicine* are generally considered to comprise Physiology and its applications to Pathology, Hygiene, and Therapeutics. By some they are considered to include General Pathology and General Therapeutics.

THEIOTHERMÆ, Waters, mineral, sulphureous.

THERAPEIA, Curation, Therapeutics.

THERAPEUSIS, Therapeutics.

THERAPEUTA, Therapist.

THERAPEUTICE, Therapeutics.

THERAPEUTICS, *Therapeutiçæ*, *Therapeutica*, *Therapeut'is*, *Isotroch'nicæ*, *Therapeut'a*, *Therapi'a*, *Curato'ria* *Meth'odus*, *Methodus* *Meden'di*, *Practice of Physic*, (F.) *Thérapeutique*, *Thérapie*, from *θεραπεω*, 'I wait upon, I alleviate, I attend upon the sick.' That part of medicine, the object of which is the treatment of disease. With others, the department which comprises an explanation of the *modus operandi* of medicines.

THERAPEUTIQUE, Therapeutics.

THERAPEUTIST, *Therapeut'a*. Same etymon. One who practices therapeutics. A practitioner of medicine. To be a good therapist, a man must be well versed in every department of medicine, and be capable of observing and reasoning well. He may be a good observer, and yet a bad reasoner. He cannot practice well unless he is both. Hence, the comparatively small number of good therapists.

THERAPIA, Therapeutics.

THERAPIE, Therapeutics.

THERIA, Radzyge.

THERIAC, Theriaca—t. of Antipater, Antipatri theriaca.

THERIACA, *Theriaca*, *Ther'iac*, from *θηρ*, 'a ferocious or venomous animal.' *Treacle*, *Melasses*, (F.) *Thériaque*. Also, a medicine, believed to be capable of curing or preventing the effects of the bite of a venomous animal. In this sense it is chiefly used by writers. *Theriac* and *Theriacal* have been used adjectively for medicinal.

THERIACA ANDROMACHI, *Venice Treacle*, is an ancient alexipharmic electuary; consisting of a farrago of 61 different ingredients, which possessed the most opposite properties. It was invented by Andromachus of Crete, and prepared by order of Nero. It has received various modifications; and, to the discredit of the *Faculté de Médecine* of Paris, long held its place in their Codex, with even an additional number of ingredients. The *Electuarium Opium polypharmacum* of the Codex has contained acid ingredients, 5; astringent, 5; bitter, 22; indigenous aromatics, 10; umbelliferous aromatics, 7; balsams and resinous substances, 8; fetid ingredients, 6; narcotics, 1; earthy substances, 1; gummy or amylaceous, 4; saccharine, 3; total, 72! and one of these the flesh of the viper! A little more than a grain of opium is contained in each drachm of the compound.

THERIACA ARTICULORUM, Colchicum—t. Celestis, Tinctura opii—t. Communis, Melasses—t. Edinensis, Confectio opii.

THERIACA GERMANORUM. An extract or rob, prepared from juniper berries. Used as a stomachic.

THERIACA LONDINENSIS, *Cataplasma Cymini*. A cataplasm of cumin seed, bay berries, german-

der, snake-root, cloves, honey, and sometimes opium, or syrup of poppies.

THERIACA RUSTICORUM, Allium.

THERIAQUE, Theriaca.

THERIAKI. In Turkey, *opium-eaters*, or they who indulge largely in the use of opium, are so called.

THERIATRICA, Veterinary art.

THERIODES, Ferine.

THERIO'MA, Therion, from *θηρ*, 'a venomous animal.' Any extremely malignant ulcer. Sometimes confined to ulcers of the lungs.

THERION, Therioma.

THERIO'TOMY, Zootomy.

THERMA, Heat—*t. Emphytum, Animal heat.*

THERMÆ, Σεππας, Ther'mata. Warm baths or springs. See *Waters (mineral).*

THERMÆ FABARIÆ, Pfeffer's, Mineral Waters of—*t. Piperinæ, Pfeffer's, Mineral Waters of*—*t. Plumbariæ, Plombières, Mineral Waters of.*

THERMAL, Therma'lis; from *Seppas*, 'heat,' *Seppos*, 'hot.' Relating or appertaining to heat. Hot, warm. As 'a thermal mineral water.'

THERMANTICA, Calefacients.

THERMASMA, Fomentation.

THERMATA, Thermæ.

THERMOMETER, see Heat.

THERMUM EMPHYTUM, Animal heat, Bio-lychium.

THESIS, thesis, from *τιθημι*, 'to put.' 'A position or proposition.' *Disputa'tio inaugura'lia, Inaug'ural Dissertation.* The name usually given to the essay composed by a candidate for graduation in medicine, which he is at times required to defend publicly. Often, however, it is a mere form, giving useless labour and trouble to the student, inasmuch as it is executed as a task, and never afterwards regarded by the preceptor or by others. Sandifort, Haller, Smellie, and Stoll, have published collections of these.

Also, a suffix denoting 'arrangement;' hence *Diathesis*.

THEVE'TIA A'HOUI, Ahouai. The kernels of the nut of this Brazilian tree are said to be a violent poison. At the Antilles, its nuts are called *Noix de Serpent*; and they are used against the bites of serpents.

THEW, Tendon.

THEY, Thigh.

THICK-LISTED, Ptery.

THIGH, Sax. ðeoð, Femur, Femem, Crus, Me-rus, Me'rium, (Sc.) The, They, (F.) Cuissæ. (Of old, *Femur* meant the thigh generally, and especially the outer part;—*Femem*, the inner). The part of the lower limb which extends from the pelvis to the leg. The thigh is larger at its superior than inferior part, and has the form of an inverted and truncated cone, slightly depressed from within outwards. Above, it is bounded, anteriorly, by the groin; externally, by the hip; behind, by the fold of the nates; and within, by the perineal region. Below, it is bounded, anteriorly, by the prominence of the knee; posteriorly, by the ham. It is formed of a considerable number of muscles, blood-vessels, lymphatics, nerves, &c.; and is covered by a strong aponeurosis.

THIGHBONE, Femur—*t. Neck of the, Collum femoris.*

THIMBLE WEED, Rudbeckia laciniata.

THION, Sulphur.

THIRST, Sax. ðýrr, (D.) Dorst, Sitie, Dipsa, Potio'nis desid'rium, (F.) Soif. A simple desire or an absolute want for liquids. Physiologists are not entirely agreed regarding the seat of thirst; some place it in the fauces; others in the stomach. Its immediate cause is not known. It has been attributed to a dry condition of the nervous papillæ of the pharynx, produced by suppression of the salivary and mucous secre-

tions. This is probably true; but, again, it is owing to the wants of the system,—a supply of fluid being required to compensate the numerous losses that are constantly taking place. Thirst is an internal sensation,—an instinctive want—arising from organization, and inexplicable. It is an urgent symptom in many diseases, particularly in those of vascular excitement.

THIRST, EXCESSIVE, Polydipsia—*t. Morbid Dipsosis.*

THIRSTY, Dipsodes.

THISTLE, BLESSED, Centaurea benedicta—*t. Cotton, Onopordium acanthium*—*t. Globe, Echinops*—*t. Holy, Centaurea benedicta*—*t. Ladies', Carduus Marianus*—*t. Milk, common, Carduus Marianus*—*t. Pine, Atractylus gummifera*—*t. Sow, Sonchus oleraceus*—*t. Star, Centaurea calcitrapa*—*t. Yellow, Argemone Mexicana.*

THLADIAS, Eunuch.

THLASIAS, Eunuch.

THLASIS, Contusion—*t. Depressio, Depression.*

THLASMA, Contusion—*t. Concussio, Concussion*—*t. Stremma, Sprain.*

THLASPI, Pennycreese. Ord. Cruciferae. Sex. Syst. Tetradynamia Siliculosa. Two species of thlaspi are directed, in some pharmacopœias, for medical use:—*Thlaspi arven'se* or *Treacle mustard*, and *Thlaspi campest're* seu *hiru'tum* seu *vulga'tius, Lepid'ium campest're, Lep'ia* seu *Ibe'ris campest'ris, Mith'ridate mustard.* The pharmaceutical properties of both kinds resemble those of common mustard.

THLASPI BURSA, Th. bursa'tum, Thlaspi seu *Capsell'a* seu *Ibe'ris* seu *Nastur'tium* seu *Rod-schied'ia bursa pasto'ris, Bursa pasto'ris, Shepherd's purse, Coccart, St. James's wort, Poor man's Pharmacy, (F.) Fleur de St. Jacques, Bonne à berger ou à pasteur, Tabouret.* A common European plant, introduced into this country. It is slightly astringent, but is little used.

THLASPI BURSATUM, Thlaspi bursa—*th. Nasturtium, Lepidium sativum*—*th. Sativum, Lepidium sativum.*

THLBIAS, Eunuch.

THLIPSENCEPHALIA, Derencephalia.

THLIPSENCEPH'ALUS, from θλιψις, 'compression,' and κεφαλος, 'the encephalon.' A monster in whom the skull is open, not merely in the frontal and parietal, but also in the occipital regions—a distinct fontanelle not existing.

THLIP'SIS, θλιψις, Compres'sio. Compression, and especially constriction of vessels by an external cause. Oppression.

THNESCOLOGIA, Thanatology.

THOLUS, Achiculum—*t. Dioecus, Scapha.*

THOMPSON'IAN, One who practises or believes in Thompsonianism.

THOMPSON'IANISM, Thompson'sonism. A fanciful doctrine, of which a person of the name of Thompson, of New York, is the founder. One of its leading principles is, that the human body is composed of four elements [?], earth, air, fire and water; and one of its apothegms,—that metals and minerals are in the earth, and, being extracted from the depths of the earth, have a tendency to carry all down into the earth who use them; that the tendency of all vegetables is to spring up from the earth, and therefore to uphold mankind from the grave.

The Thompsonians are Botanical Doctors. See *Herb doctor.*

THORACENTESIS, Thoracocentesis.

THORACHIQUE, Thoracic.

THORAC'IC, Thorac'icus, from *thorax, thoracis, 'the chest,' (F.) Thoracique* ou *Thorachique.* That which relates or belongs to the chest; as *Thoracic vis'cera, &c.*

THORACIC INFERIOR ARTERY, Arte'ria Thorac'ica Exter'na Inferior seu *Longa* seu *mam-*

ma'ria externa seu Thoracica inferior, (F.) *Deuxième des Thoraciques* (Ch.), arises from the axillary, a little below the preceding, and descends vertically over the latter part of the thorax, and the serratus major anticus. It afterwards bends inwards; becomes subcutaneous, and divides into several branches, which embrace the breast. It gives numerous ramifications to the pectoralis major, serratus anticus, intercostals, ganglions of the axilla, mamma, &c.

THORACIC SUPERIOR ARTERY, *Arte'ria Thoracica Exter'na Superior*, (F.) *Artère thoracique supérieure, Première des Thoraciques* (Ch.), arises from the axillary artery or from the acromial; and descends forwards between the Pectoralis major and P. minor, to which it distributes itself by a great number of branches. In some subjects, there are two or three *Arteriæ thoracicae externæ superiores*.

THORACIC DUCT, *Ductus thorac'icus seu thoracicus post'e'rior seu vertebra'lis seu chy'lifer seu chyli seu lact'eus seu ro'rifer seu Pecqueti seu Pecqueti'anus, Vena alba thorac'is, Alveus ampullosus, Distributo'ria lactea thoracica, Galax'ia, Duct of Pecquet, Alimentary Duct*, (F.) *Canal ou Conduit Thoracique*, is the duct in which the lymphatics of the lower limbs, abdomen, left superior extremity, left side of the head, neck, and thorax terminate. It begins at the *receptaculum chyli*, which is formed by the union of five or six large lymphatic trunks,—themselves formed from the union of all the absorbent plexuses of the abdomen. The duct ascends into the chest through the pillars of the diaphragm, and by the side of the aorta and vena azygos. It contracts in dimension, as far as the 6th dorsal vertebra, when it inclines towards the left hand; ascends behind the arch of the aorta; passes behind the left internal jugular vein, and opens at the posterior part of the subclavian vein of the same side. Its embouchure is furnished with two valves, which prevent the blood from passing from the vein into the duct.

THORACIC LIMBS are the upper limbs; so called because they are articulated with the lateral and upper parts of the chest.

THORACIC, LONG or INFERIOR, Mammary inferior external artery.

THORACIC NERVES. The *short or anterior thoracic nerves* are two in number. They arise from the brachial plexus, and divide into an anterior and a posterior branch:—the former distributed to the pectoralis major muscle; the latter uniting with a branch of the other to form a loop, from which numerous branches are given off to the pectoralis major and p. minor.

The *long thorac'ic nerve, Post'e'rior thorac'ic, External res'piratory* of Sir Charles Bell, is a long branch, which arises from the fourth and fifth cervical nerves, immediately after their escape from the intervertebral foramina, and passes downwards to be distributed upon the serratus magnus muscle.

THORACIC REGIONS are the different regions of the chest. Thus we say, *anterior, lateral, and superior thoracic, &c.*

THORACICA, Pectorals:

THORACICS, FIRST OF THE, Mammary superior external artery.

THORACIQUE, Thoracic.

THORACOCENTE'SIS, Thoracente'sis, Thoracot'omy, from *ῥωπαζ, ῥωπακος*, 'the chest,' and *κέντρος*, 'puncture.' *Paracente'sis thorac'is*. Perforation of the chest to evacuate contained fluid—as in empyema.

THORACOCYSTIS, Thoracystis.

THORACODID'YMUS, from *ῥωπαζ, ῥωπακος*, 'the chest,' and *ἰδύμος*, 'a twin.' A monstrosity in which twins are united by the thorax.—Gurli.

THORACODYNE, Pleurodynia.

THORACO-FACIAL, *Platysma myoides*.

THORACO-GASTRODID'YMUS, *Did'ymus sym'phyo-thoracogast'rius, Xiphodid'ymus*: from *ῥωπαζ, ῥωπακος*, 'the chest,' *γαστήρ*, 'the belly,' and *ἰδύμος*, 'a twin.' A monstrosity in which twins are united by the chest and abdomen.—Gurli.

THORACO-MAXILLO-FACIAL, Platysma myoides.

THORACOMETER, Stethometer.

THORACOPATH'IA, from *ῥωπαζ, ῥωπακος*, 'the chest,' and *πάθος*, 'disease.' Disease or suffering in the chest.

THORACOSCOPIA, see Auscultation.

THORACOSCOPIUM, Stethoscope.

THORACOTOMY, Thoracocentesis.

THORACYST'IS, Thoracocyst'is, from *ῥωπαζ, ῥωπακος*, 'the chest,' and *κύστις*, 'a bladder.' Encysted dropsy of the chest. Hydatids in the chest.

THORAX, ῥωπαζ, 'a cuirass, a coat of mail.' *Cith'arus, Venter me'dius, Pectus, Stethus, Scutum pect'oris, Bir, Chelys, Cassa, the Breast, the Chest*, (F.) *Poirine*. One of the splanchnic cavities, *Cavum thorac'is*, bounded, posteriorly by the vertebrae; laterally, by the ribs and scapula; anteriorly, by the sternum; above, by the clavicle; and below, by the diaphragm. It is destined to lodge and protect the chief organs of respiration and circulation—the lungs and the heart.

THORAX, Corset.

THORE, Sperm.

THORN APPLE, Datura stramonium—t. Red, Datura Sanguinea.

THOROUGHSTEM, Eupatorium perfoliatum.

THOROUGHWAX, Eupatorium perfoliatum.

THOROUGHWORT, Eupatorium perfoliatum.

THOROW-WAX, Bupleurum rotundifolium.

THORULUS STRAMINEUS, Fanon.

THRENNINE, see Tears.

THREADWORM, Dracunculus—t. Long, Tri chocephalus.

THREPSIS, Assimilation, Nutrition.

THREPSOL'OGY, Threpsolog'ia, from *θρεψις*, 'nutrition,' and *λογος*, 'a description.' The doctrine of, or a treatise on, the nutrition of organized bodies.

THREPTICE, Assimilation.

THRIDACE, Lactucarium.

THRIDAX, Lettuce.

THRIFT, LAVENDER, Statice limonium—t. American, Statice Caroliniana—t. Sea-side, Statice Caroliniana.

THRIX, Capillus, Hair.

THROAT, ῥωπα, ῥωπα, Jugulum, Interstit'ium Jugula'rē, Guttur, Sphagē, (Prov.) *Chokes, Droat, Haise, Hose*, (Sc.) *Halse, Hawse, Witter*. The anterior part of the neck. (F.) *Gorge*. Also, the pharynx.

THROATROOT, Geum Virginianum, Liatris.

THROATWORT, Campanula trachelium.

THROBBING, Pulsat'rus, Sphygm'icus, Sphygm'o'es, Pul'satory, (F.) Pulsatif. A throbbing pain, (Prov.) *Staug*, (F.) *Douleur pulsative*, is a kind of pain which is, or seems to be caused or augmented by the pulsation of arteries.

THROE, Agony, Pains, labour.

THROMBI LACTEI, Infarctus mammae lacteus.

THROMBOCYS'TIS, from *ῥωπακος*, 'a clot,' and *κύστις*, 'a cyst.' The cyst occasionally surrounding a clot of blood—as in encephalic hemorrhage.

THROMBOSIS, Coagulation, Thrombus.

THROMBUS, ῥωπακος, 'a clot,' from *ῥωπαζω*, 'to coagulate,' *Thromb'osis, Thrombus, Hæmatoma*. A small, hard, round, bluish tumour; formed by an effusion of blood in the vicinity of a vein which has been opened in the operation of blood-

letting. The thrombus is most commonly owing to the opening in the vein and that of the skin not corresponding; to excessive smallness of the cutaneous orifice; or to small, fatty granules, which prevent the discharge of the blood. Compresses, dipped in salt water; camphorated spirit, and slight compression, usually disperse it. See Blood.

THROMBUS NEONATORUM, Cephalæmatoma.

THROPILL, Trachea.

THROPPLE, Trachea.

THROTTLE, Trachea.

THROW, Agony, Pains, labour.

THROWORT, Leonurus cardiaca.

THRUSH, Aphthæ — t. Milk, Aphthæ — t. White, Aphthæ.

THRUSTY, Dipsodes.

THRYPSIS, Communion.

THUYA OCCIDENTALIS, Thuya occidentalis.

THUMB, Pollex.

THUREA, Juniperus lycia — t. Virga, Juniperus lycia.

THUS, see Pinus abies — t. Fœmininum, see Pinus abies — t. Judæorum, Croton cascarilla, Styra, Thymiana — t. Libanotos, Juniperus lycia — t. Masculum, Juniperus lycia — t. Verum, Juniperus lycia — t. Vulgaris, see Pinus abies.

THUYA APHYLLA, see Sandarac.

THUYA seu THUYA OCCIDENTALIS, *T. obtusa*, *Cupressus Arbor Vita*, *American Arbor Vita*, *Tree of Life*. Ord. Conifere. The leaves and wood were formerly in high repute as resolvents, sudorifics, and expectorants, and were given in phthisical affections, intermittent fevers, and dropsies. The expressed juice has been applied to condylomata. The arrangement of the medullary matter of the cerebellum, termed *Arbor Vita*, is also called *Thuya*.

THYLACIITIS, Gutta rosea.

THYM, Thymus.

THYMA, Thymion.

THYMALOS, Taxus baccata.

THYMASTHMA, Asthma thymicum.

THYMBRA, Satureia hortensis — t. Hispanica, Thymus mastichina.

THYME, CAT, Teucrium Marum — t. Common, Thymus — t. Lemon, see Thymus serpyllum — t. Mother of, Thymus serpyllum — t. Virginia, Pycnanthemum linifolium — t. Wild, Thymus serpyllum.

THYMELEA, Daphne gnidium — t. Laureola, Daphne laureola — t. Mezereum, Daphne mezereum — t. Monspelica, Daphne gnidium.

THYMELCO'SIS, from *Thymos*, 'thymus,' and 'ekros,' 'an ulcer.' Ulceration of the thymus gland.

THYMIAMA, *Thymia*, (*Thym*, 'I burn perfumes') 'a perfume'; *Thus Judæorum*, *Muskwood*, (F.) *Narcaphte*. A bark from Syria, Cilicia, &c., supposed to be the product of the liquid storax tree. It has an agreeable, balsamic smell; approaching that of liquid storax.

THYMIAMA, Fumigation, Suffimentum.

THYMIASIOTECHNIA, Thymiatechny.

THYMIASIS, Fumigation.

THYMIATECH'NIA, *Thymiatech'nia*, *Thymiasiotech'nia*, *Cura fignigato'ria*, from *Thymia*, 'an odour,' and *teyxn*, 'art.' The art of employing perfumes in medicine.

THYMIC, *Thymicus*, (F.) *Thymique*. Having relation to the thymus, as "thymic asthma."

THYM'ION, *Thyma*, *Thymus*, *Porrus*, *Verruca rhagoidea Verruca minor*, from *Thymos*, 'thyme.' A small wart on the skin, resembling a bud of thyme.

THYMIOSIS, Framboesia — t. Indica, Framboesia.

THYMIQUE, Thymic.

THYMITES. Wine impregnated with thyme THYMITIS, from *Thymos*, 'thymus,' and *itis*, denoting inflammation. Inflammation of the thymus gland.

THYMOPATHIA, *Psychopathia*, from *Thymos*, 'the mind,' and *pathos*, 'affection.' A disease of the mind.

THYMOS, Rage.

THYMOXAL'ME, from *Thymos*, 'thyme,' *oxys*, 'acid,' and *als*, 'salt.' A compound of thyme, vinegar and salt.

THYMUS, *Thymos*, *Glandula Thymus*, *Corpus Thymicum* seu *Thymicum*, *Glandium*, *Thymus gland*, *Corpus incomprehensibile* of Jos. Frank. Sweetbread of the butcher. An organ, concerned in hæmatosis, which is seated in the upper separation of the anterior mediastinum. The thymus has the appearance of a glandular body. It is oblong; bilobate; soft, and very variable in size and colour. In the fœtus, it is very large, and contains in a central cavity — reservoir of the thymus — a milky fluid; but it gradually disappears, and in old age is scarcely discernible. The arteries, called *thymicæ*, are from the inferior thyroid, internal mammary, bronchial, and mediastinal. The veins have the same arrangement. It receives some nervous filaments from the pneumogastric nerves, the phrenic, and the inferior cervical ganglia.

THYMUS, *T. vulgaris* seu *tenuifolius* seu *hortensis*, *Common Thyme*, (F.) *Thym*, *T. ordinaire*, *Farigoule*. Ord. Labiata. *Ses. Syst.* Didymia *Gymnospermia*. This herb has an agreeable, aromatic smell; and a warm, pungent taste. It is reputed to be resolvent, emmenagogue, tonic, and stomachic. It is not much used.

THYMUS, *Satureia capitata*, *Thymion* — t. *Calaminthus*, *Melissa calamintha* — t. *Capitatus*, *Satureia capitata* — t. *Ciliatus*, *T. mastichina* — t. *Creteus*, *Satureia capitata* — t. *Hortensis*, *Thymus* — t. *Includens*, *T. serpyllum*.

THYMUS MASTICH'NA, *T. ciliatus*, *Marum vulgare*, *Samp'euchus*, *Clinopodium*, *Mastich'ina Gallo'rum*, *Thymbra Hispanica*, *Jaca Indica*, *Common herb Mastich*. A low, shrubby, Spanish plant, used as an errhine. It has a smell like mastich. Its virtues resemble those of the *Marum Syriacum*; but it is said to be less powerful.

THYMUS MULTIFLORUS, *Melissa nepeta* — t. *Nepeta*, *Melissa nepeta*.

THYMUS, RESERVOIR OF THE, see Thymus.

THYMUS SERPYLLUM, *T. includens*, *Herpyllus*, *Herpyllus*, *Serpyllum*, *Serpyllum*, *Serpyllum*, *Gila'rum*, *Serpyllum vulgare minus*, *Mother of Thyme*, *Wild Thyme*, (F.) *Serpolet*. This plant has the same sensible properties as the garden thyme; but has a milder and rather more grateful flavour. *Lemon Thyme*, the *Serpyllum citratum*, is merely a variety of the *Thymus Serpyllum*. It is very pungent; and has a particularly grateful odour, approaching that of lemons.

THYMUS SYLVATICUS, *Clinopodium vulgare* — th. *Sylvestris*, *Satureia capitata* — th. *Tenuifolius*, *Thymus* — th. *Vulgaris*, *Thymus*.

THYRA, *Thyra*, 'a gate, folding-door.' In composition, *Thyreos* and *Thyro*, *Thyreos*, 'a shield resembling a folding-door,' mean the thyroid cartilage.

THYREMPHRAXIS, Bronchocele.

THYREO, *Thyro*, in composition, relating to the thyroid cartilage or gland.

THYREOADENITIS, Thyreoiditis.

THYREO-ARYT'ENOID, *Thyro-arytenoides* or *Thyro-arytenoides*. That which relates to the thyroid and arytenoid cartilages.

THYREO-ARTENOID LIG'AMENTS, *Inferior Ligaments of the Larynx*, *Lips of the Glottis*, *Vocal Cords*. are two ligaments about two lines broad, formed of elastic and parallel fibres, which are

contained in a duplicature of the mucous membrane of the larynx. They extend horizontally from the anterior projection at the base of each arytenoid cartilage to the middle of the angle of union of the thyroid cartilage. They are the essential organs of voice. See Glottis.

THYREO-ARYTENOID MUSCLES, THYREO-ARYTENOIDÆ, are thin muscles, which arise from the middle and inferior part of the posterior surface of the thyroid cartilage; whence they proceed backwards and outwards, to be inserted into the outer part of the base of the arytenoid cartilage. They give the necessary tension to the ligaments of the glottis, in the production of the voice. Santorini has given the name *Thyre-arytenoideus obliquus* to a portion of the *arytenoideus* muscle.

THYREO-EPIGLOT'TIC, Thyreo-epiglottides, Thyrepiglott'icus. Sabatier and Santorini have given this name to the outer portion of the thyro-arytenoid muscle; because it passes from the thyroid cartilage to the anterior part of the epiglottis.

THYREO-HYOID, Thyro-hyoides, Thyrohyoides, Hyodeothyroides. That which belongs or relates to the thyroid cartilage and os hyoides.

THYREO-HYOID or HYO-THYROID MEMBRANE is a very broad, yellowish, fibrous membrane, thicker at the middle than at the extremities, which is attached, above, to the posterior surface of the body and great cornu of the os hyoides; and, below, to the whole superior edge of the thyroid cartilage.

THYREO-HYOÏDEUS or Hyo-thyreoides Muscle is situate at the anterior and superior part of the neck. It is attached to the oblique line on the anterior surface of the thyroid cartilage; to the inferior edge of the body of the os hyoides, and to the anterior portion of its great cornu. It approximates the thyroid cartilage and os hyoides to each other, by raising the cartilage, and with it the whole larynx: or, it can depress the os hyoides.

THYREO-PHARYNGEUS, Thyro-Pharyngeus.

THYREOID, Thyroid.

THYREOÏDIEN, Thyroideal:

THYREOÏDEUS, Thyroideal.

THYREOÏTIS, Thyreoiden'tis, Thyreoidi'tis, Angi'na thyreoides, from *Suppos*, 'a shield,' and *tis*, denoting inflammation. Inflammation of the thyroid gland.

THYREONCUS, Bronchocele.

THYREOPHRAXIA, Bronchocele.

THYREPIGLOTTICUS, Thyreoëpiglottic.

THYRO, see Thyreo.

THYRO-ARYTENOID, Thyreo-arytenoid.

THYRO-CRICO-PHARYNGEUS, see Crico-pharyngeal.

THYRO-HYOIDES, Thyreo-hyoid.

THYRO-PHARYNGEUS, Thyreo-pharyngeus.

Valsalva, Morgagni, Santorini, and Winslow have given this name to the portion of the constrictor pharyngis inferior, which is attached to the thyroid cartilages.

THYRO-PHARYNGEUS, Constrictor pharyngis, see Crico-pharyngeal.

THYRO-PHARYNGO-STAPHYLINUS, Palato-pharyngeus.

THYROCELE, Bronchocele.

THYROID, Thy'reoid, Thyroï'des, from *Thupa*, 'a gate or folding-door,' or from *Suppos*, 'a shield,' and *eidos*, 'form.' That which has the shape of a folding-door.

THYROID CARTILAGE, Cartila'go Scutifor'mis seu *Scutalis* seu *Clypealis*, *Coëperto'rium*, *C. Peltalis* seu *peltatus*, *Scutum*, *Adami Moræus* *Os*, is the largest of the cartilages of the larynx, at

the anterior part of which it is situate. It is larger transversely than vertically; broader above than below; and seems formed by the junction of the two quadrilateral plates, which produce, by their union, an angle that projects forwards. Its two posterior margins terminate above, and on each side, by an ensiform prolongation, called the *greater cornu of the thyroid cartilage, cornu superius* seu *longum*; and, below, by a less prominent eminence, the *lesser cornu, Cornu inferius* seu *brevè*, which is articulated with the cricoid.

THYROID GLAND, Corpus Thyreoidæum, Glan'dula Thyreoidæa, (F.) Gland ou Corps Thyroïde, is an organ, the uses of which are not known. It covers the anterior and inferior part of the larynx, as well as the first rings of the trachea, and seems to be composed of two lobes, flattened from before to behind, which are united by a transverse prolongation of the same nature as themselves, called the *Isthmus*. The tissue of the thyroid is soft, spongy, and of a brownish colour; but its intimate structure is unknown. It is formed of several distinct lobules; collected in lobes of greater or less size. These are composed of granulations, some of which contain a yellowish or milky fluid. The thyroid gland receives four large arteries, called *Thyroideal*, as well as corresponding veins. Its nerves proceed from the pneumogastric, and from the cervical ganglia. No excretory duct has ever been found in it. Its uses are not known.

THYROIDÆAL, Thyroideus, Thyreoides, (F.) Thyroïdien, Thyroëdien. That which concerns the thyroid gland or cartilage.

THYROIDÆAL ARTERIES are two on each side.

1. The *superior Thyroideal, superior laryngeal, superior guttural*, arises from the anterior part of the external carotid, and proceeds to the upper part of the thyroid gland, after having given off a *laryngeal branch* and a *crico-thyroid*. 2. The *inferior Thyroideal, inferior guttural*, much larger than the last, arises from the upper part of the subclavian. It passes, in a serpentine manner, to the inferior part of the thyroid gland, to which it is distributed, after having given a number of branches to the neighbouring parts, among which is the *cervicalis ascendens*.

THYROIDÆAL VEINS are, 1. A *superior thyroideal*, and several *middle thyroideal*, which open into the internal jugular vein. 2. An *inferior thyroideal*, much larger than the preceding, whose branches form — by anastomosing with those of the opposite side, in front of the trachea — a very remarkable venous plexus, which J. Cloquet calls the *infra-thyroideal plexus*. The inferior thyroideal veins open, — the left into the corresponding subclavian; — the right, into the vena cava superior.

THYROIDÏEN, Thyroideal.

THYROPHRAXIA, Bronchocele.

THYRSUS, Penis.

THYSSELINUM, Peucedanum palustre.

THYSSELUM PALUSTRE, Peucedanum palustre.

TIBI-PÉRONÉO-TARSIEN, Peronæus longus.

TIB'IA. A Latin word, which signifies a flute or pipe. *Cnemè, Procnè'mè, Procnè'mium, Fo'cièz majus, Arun'do* seu *Canna major, Canna major* seu *dome'stica eruris, Shinbone*. The largest bone of the leg. A long, irregular, and triangular bone, situate on the inner side of the fibula. It has, 1. A *Superior or femoral extremity*, which is very large, rounded, and has two eminences at the sides, called *Tuberosities of the Tibia, or Tibial tuberosities*. The outermost is articulated with the fibula. On its head are two articular, oval, concave surfaces, separated by a process, called

Spine of the tibia, which are articulated with the condyles of the *Os femoris*. 2. An *inferior or tarsal extremity*, having an articular surface beneath, which joins the astragalus; within, a triangular eminence, which forms the malleolus internus; and, on the outside, a triangular surface, which articulates with the fibula. 3. The body or shaft of the tibia has three faces, separated by three angles, the anterior of which is the most prominent, and is called the *spine or crista of the tibia or tibial spine*. It is the *shin*. The tibia is articulated with the femur, fibula, and astragalus. It is developed by three points of ossification, one for the body, and one for each extremity.

TIBIA MINIMA, Fibula.

TIBIÆUS, Tibial.

TIBIAL, *Tibia'lis, Tibiæ'us*, (F.) *Tibial, Jambier*. From *tibia*. That which relates to the tibia or to the leg.

TIBIAL APONEUROSIS, (F.) *Aponévrosé Jambière*, surrounds the muscles of the leg. It is continuous, above, with the femoral aponeurosis, and arises, also, from the head of the fibula, and from several fibrous expansions which detach themselves from the tendons of the triceps, sartorius, gracilis, and semitendinosus. Thence it descends around the leg, attaching itself to the whole extent of the anterior and inner edges of the tibia. It sends, below, an expansion, which passes before the tendo Achillis, and is continuous with the anterior annular ligament of the tarsus. It is attached, also, to the sheath of the peronei muscles.

TIBIAL ARTERIES are two in number. 1. *Arteria Tibia'lis Anti'ca*, which arises from the popliteal, before its bifurcation: passes immediately through the opening in the upper part of the interosseous ligament, and reaches the anterior part of the leg; when it turns downwards, and descends obliquely, between the extensor communis digitorum pedis and the tibialis anticus, anterior to the interosseous ligament; glides under the anterior annular ligament of the tarsus, and takes the name *Dorsalis tarsi*. Amongst the numerous branches, given off by the tibialis antica, may be distinguished—the *Tibialis recurvens*, *A. malleolaris interna*, and *A. malleolaris externa*. 2. *A. Tibialis post'ca*, arises from the bifurcation of the popliteal artery; and, slightly tortuous, descends between the two planes of the posterior muscles of the leg; until it arrives under the arch of the calcaneum, where it divides into two branches—the *Plantar arteries*.

TIBIAD, see Tibial aspect.

TIBIAL ASPECT. An aspect towards the side on which the tibia is situated.—Barclay. *Tibiad* is used adverbially by the same writer, to signify, 'towards the tibial aspect.'

TIBIAL NERVES are two in number. 1. The *anterior tibial nerve, Prétibio-supplantaire*, (Ch.) One of the two branches in which the external popliteal terminates. It accompanies the *arteria tibialis antica*. 2. The *posterior tibial nerve, Branche tibiale du nerf fémoro-poplitée* (Ch.), is the *internal popliteal*.

TIBIAL VEINS, ANTERIOR AND POSTERIOR, have the same arrangement as the arteries which they accompany.

TIBIALIS ANTI'CUS, *Catænæ musculus*, (F.) *Tibio-sus-métatarsien, Tibio-austarsien* (Ch.), *Jambier antérieur*. A muscle situate at the anterior part of the leg. Above, it has the shape of a triangular prism; below, it is slender and tendinous. It is attached, above, to the anterior part of the external tuberosity of the tibia; to the superior half of the outer surface of that bone; and to the anterior surface of the interosseous ligament. Below, its tendon terminates at the base of the first cuneiform bone, and at the

posterior extremity of the first metatarsal bone. This muscle bends the foot on the leg, and directs its point inwards, at the same time that it raises its inner edge. It can, also, bend the leg on the foot, and prevent it from falling backwards in standing.

TIBIALIS GRACILIS, Plantar muscle.

TIBIALIS POSTI'CUS, (F.) *Tibio-tarsien, Tibio-sous-tarsien* (Ch.), *Jambier postérieur*. This muscle is situate at the posterior and deep part of the leg. It is thicker above than below; and is attached, above, to the posterior surface of the fibula; to the posterior surface of the tibia, and to the interosseous ligament. Its tendon terminates, below, at the tuberosity on the inferior extremity of the os scaphoides. This muscle extends the foot on the leg; at the same time raising its inner edge. It, likewise, extends the leg on the foot.

TIBIO-CALCANIEN, Soleus—*t. Phalangetien*, *Flexor longus digitorum pedis profundus perforans*—*t. Sous-tarsien*, *Tibialis posticus*—*t. Sus-métatarsien*, *Tibialis anticus*—*t. Sus-tarsien*, *Tibialis anticus*.

TIBIO-TARSAL, *Tibio-tar'seus*, (F.) *Tibio-tarsienne*. What relates to the tibia and tarsus.

TIBIO-TARSAL ARTICULATION is the articulation of the foot with the leg. It is formed by the tibia and the astragalus; and is strengthened by lateral, and by anterior and posterior ligaments.

TIBIO-TARSIEN, *Tibialis posticus*.

TIC, *Twitching, Vellication*. A local and habitual convulsive motion of certain muscles; and, especially, of some of those of the face. It is, sometimes, called *Tic convulsif* ou *Tic non douloureux*, *Histric'ic Spasm of the Face*, to distinguish it from *Tic douloureux* or *neuralgia faciei*, and has been termed a *partial chorea* or form of *chronic chorea*, *Spasmus facia'lis*, *Chorea faciei*.

TIC DOULEUREUX, Neuralgia, facial—*A non douloureux*, see Tic.

TICKLE WEED, *Veratrum viride*.

TICKLING, (Sc.) *Kitling, Kitilling*, (F.) *Chatouillement*. This word, sometimes, means the action of tickling (*titilla'tio, titilla'tion*); at others, the sensation produced by this action (*pruri'tus*). A vivid sensation, which commonly causes laughter, and a state of general spasm that may be dangerous if too long protracted. There are some parts of the body, where we are easier tickled than others; for example, the sole of the feet, and the hypochondriac regions.

TICKSEED SUNFLOWER, *Coreopsis trichosperma*.

TICKWEED, *Hedeoma*—*t. Sunflower*, *Coreopsis trichosperma*.

TICO'REA FEBRIFUGA; a Brazilian plant. Order, Rutaceæ, the bark of which is intensely bitter, astringent and antiperiodic.

TICOREA JASMINIFLO'RA. A decoction of the leaves is said to be drunk by the Brazilians as a cure for frambæsia.

TIDAL AIR, see Respiration.

TIERCE, see Tertian.

TIGE PITUITAIRE, Infundibulum of the brain—*t. Sus-sphénoïdale*, Infundibulum of the brain.

TIGILLUM, Crucible.

TIGLIA SEU TIGLII GRANA, see Croton Tigium.

TIGRETIER, see Mania, dancing.

TIGULA, Saccharum.

TIKIMMA, *Cacoucia coccinea*.

TILBURY, MINERAL WATERS OF. A saline chalybeate at West Tilbury, in Essex, England. It is not much frequented.

TIL'IA, *T. Europæa seu grandiflo'ra seu co-*

disfolia seu *pauciflora* seu *pataphylla*, (akin to *arbuta*, 'the elm tree'). *Phil'gra*. The *Lime* or *Linden tree*, (F.) *Tilleul*. Ord. *Tiliaceae*. The flowers have been supposed to possess anodyne and antispasmodic virtues.

TILIA AMERICANA. *T. glabra* seu *Canadensis*, *Linden*, *Linn*, *Basewood*, *Whitewood*, *Spoonwood*. A mucilage prepared by macerating the inner bark in cold water, has been applied to burns. It may be made into a cataplasm by thickening it with bran.

TILIA GLABRA, T. Americana—*t. Canadensis*, *T. Americana*.

TILLEUL, *Tilia*.

TILlicherry CORTEX, *Nerium* antidy-sentericum.

TILMA, *Lintum*.

TILMOS, *Eulsion*.

TILMUS, *Carphologia*.

TILY, OF THE INDIANS, *Viburnum dentatum*.

TIMAC. The name of an East India root, but of what vegetable is not clear. It is said to possess diuretic properties; and hence has been given in dropsies.

TIMBRE (F.), 'the sound of a bell.' The quality of sound by which one voice, or one instrument, can be distinguished from another when producing the same note.

TIMIDUS, *Rectus inferior oculi*.

TIN, (D.) *Ten*. *Stannum*, *Cassiteros*, *Cattiteros*, *Stangos*, *Jupiter*, *Stu'pia*, *Laos*, *Plumbum album* seu *can'didum*, (F.) *Étain*. A solid metal; of a peculiar odour when rubbed; insipid; colour whitish; s. g. 7.291; giving a peculiar tinkling sound when struck. It is used, in medicine, only as a mechanical antihelmintic. Dose, $\mathfrak{z}\mathfrak{j}$ of the *Pulex Stanni*, *Stannum granulatum* or *Granular tin*, in molasses. This is made by melting tin in an iron vessel over the fire, and, while it is cooling, stirring until it is reduced to a powder, which is passed through a sieve.—Ph. U. S.

TIN, BUTTER OR, *Tin*, muriate of—*t. Chloruret* of *Tin*, muriate of—*t. Deutohydrochlorate* of *T. muriate* of.

TIN-FOIL, *Stannum foliatum*, *Stanni'olum*, is used for plugging carious teeth, &c.

TIN-GLASS, *Bismuth*—*t. Granular*, see *Tin*—*t. Muriate* of, superoxygenated, *T. muriate* of.

TIN, MURIATE OF, *Fuming liquor of Libavius*, *Murius Stanni* seu *Oxyd'uli Stanni*, *Butter* or *Chlor'uret* or *Deu'to-hydro-chlorate* or *Superoxygenated Muriate* of *Tin*. This is formed of *tin*, one part; concentrated *muriatic acid*, three parts. To be crystallized by the aid of heat. A violent cathartic. Dose, gr. ij or iij.

TIN, SULPHURET OF, *Aurum musivum*.

TINAGMUS, *Concussion*.

TINASMUS, *Tenismus*.

TINCTU'RA, from *tingere*, *tinctum*, 'to dye.' *Tincture*, *Essentia*, (F.) *Teinture Alcoolique*, *Alcoolé*, *Alcohol*, *Alcoholature*. The term *tincture* is generally restricted to spirituous solutions of vegetable, animal, and some saline substances. It corresponds, therefore, with the word *Quintessence*, in one of its old significations; and with the *Alcoholatum* of the *Codex* of Paris. It is not unusual, however, to speak of *aqueous tincture*, *ethereal tincture* (*Éthérolature*; *Béral*), &c. *Tinctures* are usually made either with pure alcohol or proof spirit. The former are precipitated by water; and, therefore, are seldom used internally; the latter are common additions to infusions, decoctions, &c. They ought not, of course, to be united with any vehicle which can decompose them, or separate any thing from them in a palpable form; unless such decomposition is desired by the prescriber. In making *tinctures*, the ingredients should be reduced to a coarse

powder; and the maceration be made in close vessels, exposed to a heat of about 80°, and frequently shaken. When completed, they must be filtered, and put away for use in close bottles. When the process of displacement is employed, great care must be taken, so that the substances treated may be, as far as possible, exhausted of their soluble principles, and a perfectly clear tincture be obtained. To those who are not familiar with the process, the plan of maceration is preferable.—Ph. U. S.

TINCTURA ABSIN'THII COMPOSITA, *Essentia absin'thii composita* seu *amara*, *Compound Tincture* of wormwood. (*Absinth*, *artem. pontic*, *caryoph. aa* $\mathfrak{z}\mathfrak{ss}$, each. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, alcohol. Oss. Macerate for fifteen days. Ph. P.) Tonic, stomachic, vermifuge, and carminative. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$.

TINCTURA ACACIÆ CATECHU, *T. catechu*—*t. Acetatis ferri cum alcohole*, see *T. ferri acetatis*—*t. Acidi sulphurici*, *Sulphuricum acidum aromaticum*.

TINCTURA ACONIT'II, (Ph. U. S. 1842.) *Tinctura Aconiti folio'rum*, (Ph. U. S. 1851.) *Tincture* of *Ac'onite*, (*Aconit. folior.* $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, alcohol dilut. Oij. Macerate for 14 days, express, and filter through paper. It may also be made by displacement. Ph. U. S.) Dose, gtt. xx or xxx. It is rubbed on the seat of neuralgia.

TINCTURA ACONIT'II RADICIS, *Tincture* of *Aconite root*, (*Aconit. radices contus.* $\mathfrak{h}\mathfrak{j}$, Alcohol. Oij. Macerate for 14 days; express and filter. It may also be prepared by displacement. Ph. U. S.) Uses same as the last.

TINCTURA ÆTHEREÆ CAMPHORATÆ, *Solutio camphora æthereæ*, *Liquor nervinus Bangii*, *Spiritus sulphurico-æthereus camphoratus*, *Naphtha vitrioli camphorata*. (*Camphor. p. i. Æther sulphuric. p. ii.*) Stimulant in atonic diseases, arthritic cardiacal, and spasm. Dose, 20 to 30 drops in white wine.

TINCTURA ÆTHEREÆ FERRI, *Alcohol* (seu *Tinctura*) *sulphurico-æthereus ferri*—*t. Alcoholica Chinæ composita*, *T. cinchonæ composita*—*t. Alcoholica corticum aurantium Whyttii*, *Tinctura cinchonæ amara*—*t. Alexipharmaca Huxhami*, *Tinctura cinchonæ composita*.

TINCTURA AL'OEÆ, *T. Al'oes Socotri'næ*, *Essentia Aloes*, *Al'cohol cum Aloë perfoliatâ*, *Tincture* of *Aloes*. (*Aloes*, pulv. $\mathfrak{z}\mathfrak{j}$, ext. glycyrrh. $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, aquæ Oiss, alcohol. Oss. Macerate for 14 days, and filter. Ph. U. S.) Properties, those of aloes. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$.

TINCTURA AL'OEÆ ÆTHEREÆ, *T. Aloës vitriolatæ*, *Æthereal Tincture* of *Aloes*. (*Aloes Socot.*, *Myrrh.*, aa $\mathfrak{z}\mathfrak{ss}$, croci $\mathfrak{z}\mathfrak{j}$, sp. ætheris sulph. Oj. Digest the myrrh in the ether for 4 days; then add the aloes and saffron. Ph. E.) Stimulant and cathartic. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{j}$.

TINCTURA AL'OEÆ COMPOSITA, *Elixir Proprietatis* seu *aperitivum* seu *Aloes*, (*Tinctura Aloes cum Myrrhâ seu Aloës et Myrrhæ* (Ph. U. S.), seu *Aloes et Myrrhæ crocata*, *Elixir of long life*, *Compound Tincture of Aloes*, (F.) *Elixir de longue vie*, *Baume de vie de Lelièvre*, (*Aloes*, in pulv., $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, croci, $\mathfrak{z}\mathfrak{j}$, *Tinct. Myrrhæ*, Oij.) Macerate for 14 days, and filter. Purgative and stomachic. Dose, $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ to $\mathfrak{f}\mathfrak{z}\mathfrak{j}$.

Boerhaave's Elixir consisted of aloes, myrrh, saffron, and tartrate of potassa, digested in alcohol and water.

Radcliffe's Elixir is formed of *Aloes socotr.* $\mathfrak{z}\mathfrak{j}$, cort. cinnam., rad. zedoar., aa $\mathfrak{z}\mathfrak{ss}$, rad. rhei $\mathfrak{z}\mathfrak{j}$, coccinell. $\mathfrak{z}\mathfrak{ss}$, syrup. rhamni $\mathfrak{z}\mathfrak{i}\mathfrak{j}$, sp. tenuior. Oj, aquæ puræ $\mathfrak{f}\mathfrak{z}\mathfrak{v}$.

TINCTURA AL'OEÆ ET MYRRHÆ, *T. aloës composita*—*t. Aloes et myrrhæ crocata*, *T. A. composita*—*t. Aloes vitriolata*, *T. A. ætherea*—*t. Amara*, *T. gentianæ composita*—*t. Ammoniac composita*, *Spiritus ammoniac succinatus*—*t.*

Amomi repentis, *T. cardamomi* — *t. Amomi zingiberis*, *T. zingiberis*.

TINCTURA ANGUSTURÆ, *T. Cuspariæ*, *Tincture of Angustura*. (*Cort. cuspar. in pulv. crass. redact.* ʒij, *sp. vin. ten. Oij.* Digest. *Ph. D.*) Dose, fʒj to fʒij.

TINCTURA ANTIFEBRILIS WARBURGI, see *Beberu*.

TINCTURA ANTIDONTALGICA, *Toothach Tincture*. (*Tamini. ʒj, Mastich. gr. v, æther. ʒij. M.*) To be applied on cotton wool to the tooth previously dried.

TINCTURA ARISTOLOCHIE SERPENTARIÆ, *T. serpentariæ* — *t. Aromaticæ*, *T. cinnamomi composita* — *t. Aromaticæ ammoniata*, *Spiritus ammoniæ aromaticus*.

TINCTURA ASSAFÆTIDÆ, *T. Ferulæ Assafætidæ*, *T. fœtida seu assafætidæ*, *Tincture of Assafætidæ*. (*Assafætid. ʒiv, alcohol. Oij.* Macerate for 14 days and filter. *Ph. D.*) Dose, gtt. xv to fʒj.

TINCTURA ASSAFÆTIDÆ AMMONIATA, *Spiritus Ammoniæ fœtidus*.

TINCTURA AURANTII, *T. Corticis Aurantii*, *Tincture of Orange Peel*. (*Aurant. cort. recent. ʒij, sp. rect. Oij.* Digest for 14 days.) Stomachic. Used as an adjunct to stomachic draughts. Dose, fʒss to fʒij.

TINCTURA AUREA, *Alcohol seu tinctura sulphurico-æthereus ferri*.

TINCTURA BELLADONNÆ, *Tincture of Belladonna*. (*Belladonnæ. ʒiv, alcohol. dilut. Oij.* Macerate for 14 days; express, and filter through paper. It may, also, be prepared by displacement. *Ph. U. S.*)

TINCTURA BENZOÏNI COMPOSITA, *T. Benzoës composita*, *Balsamum Catholicum seu Persicum seu traumaticum*, *Elizir traumaticum*, *Compound Tincture of Benzoïn, Friar's or Vervain's Balm*, *Wade's or Jesuit's drops*, *Commander's or Wound Balm*, *Balm for Cuts*. (*Benzoïn. ʒij, styrac. purif. ʒij, Balsam. tolu. ʒj, aloes, in pulv. ʒss, alcohol. Oij.* Macerate for 14 days and filter. *Ph. U. S.*) Stimulant. Used chiefly to wounds and ulcers.

The basis of *Turlington's Balm of Life*, *Turlington's Balm*, is the Compound Tincture of Benzoïn. The following form is recommended by a committee of the Philadelphia College of Pharmacy:—*Alcohol. Oviij, Benzoïn. ʒxij, Styrac. liquid. ʒiv, Aloes Socotrin. ʒj, Balsam. Peruvian. ʒij, Myrrha ʒj, Rad. Angelic. ʒss, Balsam. Tolu. Ext. Glycyrrh. ʒā ʒj.* Digest for 10 days, and strain.

The *Essence of Coltsfoot* consists of equal parts of the Compound Tincture of Benzoïn and *Balsam of Tolu*, to which is added double the quantity of *Rectified Spirit of Wine*. It is sold as a pectoral.

TINCTURA BUCHU, *T. Buchu*, *Tincture of Buchu*. (*Buch. ʒv, Alcohol. dilut. Oij* (*Imp. meas.*). Digest for seven days and filter. It may be made, also, by percolation. *Ph. E.*) Dose, fʒj to fʒij.

TINCTURA BUCKU, *Tinctura Buchu*.

TINCTURA CALUMBÆ, *T. Columbae seu Columbae* (*Ph. U. S.*), *Tincture of Columbo*, *Columbo Bitters*, (*Columb. rad. cont. ʒiv, alcohol. dilut. Oij.* Macerate for 14 days; express, and filter. *Ph. U. S.*) Dose, fʒss to fʒiv. It may, also, be made by displacement.

TINCTURA CAMPHORÆ, *Spiritus camphoræ*.

TINCTURA CAMPHORÆ COMPOSITA, *T. Opii Camphorata* (*Ph. U. S.*), *Compound Tincture or Camphorated Tincture of Opium*, *Elizir Paregoricum*, *Paregoric Elizir*. (*Opii, in pulv. Acid Benzoic. ʒā ʒj, ol. anisi, fʒj, mellis despumat. ʒij, Camphor. ʒij, alcohol. dilut. Oij.* Macerate for 14 days, and filter through paper. *Ph.*

U. S.) One fluidounce contains nearly gr. ij of opium. Anodyne. Dose, fʒj to fʒij.

Squire's Elizir, an empirical carminative preparation, is thus made, according to one formula. (*Opii ʒiv, camphor. ʒj, cocci. ʒj, fœnicul. dulc. ʒij, tinct. serpent. Oj, sp. anisi. cong. ij, aqua Oij, auri musci, ʒvj.*)

TINCTURA CAN'NABIS INDICÆ, *Tincture of Indian Hemp*. (*Ext. Cannab. purif. ʒss* (*avoidsu- pois*), *sp. rectificat. Oss* (*Imp. meas.*) Dissolve. *Ph. D.*) Dose, about 40 drops.

TINCTURA CANTHARIDIS, *T. O. Vesicato'riæ*, *T. Lyttæ seu Melœs vesicato'rii*, *Tincture of Cantharides*, *Tincture of Blistering Flies*, *T. of Spanish Flies*, (*F.*) *Teinture rubéfiante*. (*Cantharid. cont. ʒij, alcohol. dilut. Oij.* Macerate for 14 days; express, and filter. It may, also, be prepared by displacement. *Ph. U. S.*) Stimulant and diuretic, but not often used internally. Externally, stimulant. Dose, gtt. x.

Matthew's Injection, a once celebrated nostrum for fistula in ano, consisted wholly of a dilute Tincture of Cantharides.

TINCTURA CAPSICI, *Tincture of Capsicum or Cayenne Pepper*. (*Capsic. ʒj, alcohol. dil. Oij.* Macerate for 14 days, and filter. It may, also, be prepared by displacement. *Ph. U. S.*) Stimulant. Dose, fʒss to fʒj.

TINCTURA CARDAMOMI, *T. amo'mi repentis*, *Tincture of Cardamoms*. (*Cardam. cont. ʒiv, alcohol. dilut. Oij.* Macerate for 14 days, strain, and filter. It may, also, be made by displacement. *Ph. U. S.*) Use, the same as the seeds. Dose, fʒj to fʒiv.

Solomon's Balm of Gilead, an empirical preparation, seems to be an aromatic tincture, — of which Cardamoms form a leading ingredient, — made with brandy. Some affirm that it contains cantharides.

TINCTURA CARDAMOMI COMPOSITA, *Tinctura Stomachica*, *Compound Tincture of Cardamoms*, *Stomach Tincture*. (*Cardamom. cont. ʒvj, Carui. cont. ʒij, cinnam. cont. ʒv, Uva passa demptic acinis, ʒv, cocci. cont. ʒj, alcohol. dilut. Oiss* Macerate for 14 days; express and filter. *Ph. U. S.*) Stomachic and carminative. Dose, fʒij to fʒiv.

TINCTURA CARMINATI'VA SYLVII, *Alcoolatum Carminativum Sylvii*, *Carminative Elizir of Sylvius*. (*Rad. angelic. 4 p, rad. imperator., galang. ʒā 6 p, ol. rosamarin., majoran., rutæ, basilic. ʒā 48 p, baccar. laur. nobil. 12 p. sem. angelic., Uguet. levist., anis. ʒā 16 p., singib., nuc. moschat. ʒā 6 p., cinnam. 12 p., caryoph., cort. limon. ʒā 4 p.* Add alcohol 1500 p. Macerate and distil, 1000 p. *Ph. P.*) A warm stomachic, carminative, &c. Dose, fʒss to fʒss.

TINCTURA CASCARILLÆ, *T. Crotonis Eleutheriæ*, *Alcohol cum Crotonis Cascarillæ*, *Tincture of Cascarilla*. (*Cascarill. cort. cont. ʒiv, sp. ten. Oij. Ph. L.*) Dose, fʒj to fʒiv.

TINCTURA CASTOREI, *T. Castorei Ros'ici seu Castorei Canadensis*, *Essentia Castorei*, *Alcohol castoria'um*, *Tincture of Castor*. (*Castor cont. ʒij, alcohol. Oij.* Macerate for 7 days, express, and filter. *Ph. U. S.*) Tonic and antispasmodic. Dose, gtt. xx to fʒij, or more.

TINCTURA CASTOREI COMPOSITA, *T. C. ammoniata*, *Compound Tincture of Castor*. (*Cast. ʒj, g. asafætid. ʒss, alcohol. ammoniat. Oj. Ph. E.*) Antispasmodic. Dose, fʒj to fʒiv. Called, also, *Elizir fœtidum*, *Tinctura castorei fœtida seu fœtida ammoniata*.

TINCTURA CAT'EBU, *Tinctura Japonica seu Mino'sæ Catechu seu Aca'cia Catechu*, *Tincture of Catechu*. (*Catech. ʒij, cinnam. contus. ʒij, alcohol. dilut. Oij.* Macerate for 14 days, express and filter. *Ph. U. S.*) Astringent. Dose, fʒj to fʒij.

The Concentrated Solution of Charcoal, sold for cleaning the teeth, seems to be little more than the tincture of catechu.

TINCTURA CHIRETTÆ, *Tincture of Chiretta*. (Chirett. cont. $\mathfrak{Z}\text{v}$, (avoid.) alcohol. dilut. Oij. (Imp. meas.) Macerate for 14 days. Ph. D.) Dose, f $\mathfrak{Z}\text{j}$, or f $\mathfrak{Z}\text{ij}$, as a tonic.

TINCTURA CICUTÆ, *T. conii maculati*.

TINCTURA CINCHONÆ, *T. Corticis Peruvia'ni Simplex seu Cort. Peruvia'ni*, *Tincture of Cinchona or of Bark*. (Cinchon. flav. in pulv. $\mathfrak{Z}\text{vj}$, alcohol. dilut. Oij. Macerate for 14 days, express, and filter through paper. It may, also, be made by displacement. Ph. U. S.) Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{iv}$, or more.

TINCTURA CINCHONÆ AMA'RA seu alcohol'ica corticum aurantiorum Whyttii seu Kinki'næ ama'ra, *Elizir antihypochondriacum seu rob'orans Whyttii seu Whyttii seu stomach'icum spirituosum, Essent'ia Corticis Peruvia'ni Composita*. (Cinch. p. viij, Gentian. Rad., Cort. Aurant. p. iij. Alcohol. ('900) 96 p. Digest. Dose, one or two teaspoonfuls.)

TINCTURA CINCHONÆ AMMONIA'TA, *T. Corticis Peruvia'ni volatilis*, *Ammoniated or Volatile Tincture of Bark*. (Cinchon. lancifol. cort. cont. $\mathfrak{Z}\text{iv}$, sp. ammon. aromat. Oij. Ph. L.) In dyspepsia, combined with acidity and languor.

TINCTURA CINCHONÆ ÆTHE'REA COMPOS'ITA, *T. Kinæ Kinæ æthe'rea compos'ita, Elizir'ium Antisepticum Docto'ris Chausier, Chausier's Antiseptic Elizir*. (Cinchon. officin. $\mathfrak{Z}\text{ij}$, cascarill. $\mathfrak{Z}\text{ss}$, cinnam. $\mathfrak{Z}\text{ij}$, croci $\mathfrak{Z}\text{ss}$, sacch. alb. $\mathfrak{Z}\text{xxxv}\text{ij}$. Put these bruised into a mattress, and add vin. alb. Hispanic. vel vin. muscat. alcohol. $\mathfrak{a}\mathfrak{a}$ Oj. Macerate for two days, and add sulphuric ether f $\mathfrak{Z}\text{iss}$. Ph. P.) Tonic, stimulant, and antiseptic. Dose, f $\mathfrak{Z}\text{ss}$ to f $\mathfrak{Z}\text{j}$.

TINCTURA CINCHONÆ COMPOS'ITA, *T. Corticis Peruvia'ni Composita seu febrifuga Docto'ris Huxhami seu Alexiphar'maca Huxhami seu Alcohol'ica Chinæ Compos'ita, Essent'ia Chinæ seu Corticis Peruvia'ni Alexiphar'maca Huxhami seu antiseptic'a Huxhami, Compound Tincture of Cinchona or Bark, Huxham's Tincture of Bark*. (Cinchon. rubr. pulv. $\mathfrak{Z}\text{ij}$, aurant. cort. cont. $\mathfrak{Z}\text{iss}$, serpent. cont. $\mathfrak{Z}\text{ij}$, croci, santal. $\mathfrak{a}\mathfrak{a}$ $\mathfrak{Z}\text{j}$, alcohol. dilut. f $\mathfrak{Z}\text{xx}$. Macerate for 14 days, express, and filter. Ph. U. S.) It may, also, be made by displacement. More grateful than the simple tincture. Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{ij}$.

TINCTURA CINNAMOMI, *T. Lauri Cinnamomi, Essent'ia Cinnamomi, Tincture of Cinnamon*. (Cinnam. cont. $\mathfrak{Z}\text{ij}$, alcohol. dilut. Oij. Macerate for 14 days, express, and filter. Ph. U. S.) It may, also, be prepared by displacement. Stomachic. Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{ij}$.

TINCTURA CINNAMOMI COMPOS'ITA, *T. seu Essent'ia Aromat'ica, Alcoöl cum Aromat'ibus Compos'itus, Compound Tincture of Cinnamon*. (F. Eau de Bonferme ou d'Armagnac. (Cinnam. cont. $\mathfrak{Z}\text{j}$, cardam. cont. $\mathfrak{Z}\text{ss}$, zingib. cont. $\mathfrak{Z}\text{ij}$, alcohol. dilut. Oij. Macerate for 14 days, express, and filter. Ph. U. S.) It may, also, be prepared by displacement. Use, same as the last. Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{ij}$.

TINCTURA COCCI CACTI, *Tincture of Coch'ineal*. (Cocci pulv. $\mathfrak{Z}\text{j}$ (avoid.), alcohol. dilut. Oj. (Imp. meas. Ph. D.) Chiefly a colouring ingredient. At times, given in nervous affections. Dose, gtt. xij to f $\mathfrak{Z}\text{j}$.

TINCTURA DE COCHLEA'RIS, *Alcoola'tum de Cochle'a'riis seu antiscorb'uticum*. (Fol. cochlear. recent. 2500 p., rad. armoracæ, 230 p., alcohol. (22° to 32° Beaumé) 3000 p., distil off 2000 p. Ph. P.) Antiscorbatic. Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{j}$.

TINCTURA COL'CHICI, *T. C. Sem'inis*, (Ph. U. S.), *Tincture of Colchicum Seed*. (Colchic. sem.

cont. $\mathfrak{Z}\text{iv}$, alcohol. dilut. Oij. Macerate for 14 days, express, and filter. Ph. U. S.) It may, also, be prepared by displacement. Dose, ten to sixty drops.

TINCTURA COLCHICI COMPOSITA, *Spiritus colchici ammoniatus*—*t. Colchici seminis*, *T. colchici*—*t. Colombæ*, *T. Calumbæ*.

TINCTURA CONI'Æ, *T. C. Macula'ti*, *T. Cicutæ*, *Tincture of Hemlock*. (Conti fol. $\mathfrak{Z}\text{iv}$, alcohol. dil. Oij. Macerate for 14 days, express, and filter. Ph. U. S.) It may also be made by displacement. Use;—the same as that of the leaves.

TINCTURA CONII MACULATI, *T. conii*—*t. Convolvuli Jalapæ*, *T. jalapii*—*t. Corticis aurantii*, *T. aurantii*—*t. Corticis Peruviani composita*, *T. cinchonæ comp.*—*t. Corticis Peruviani simplex*, *T. cinchonæ*—*t. Corticis Peruviani volatilis*, *T. cinchonæ ammoniata*.

TINCTURA CROCI SATI'VI, *T. Croci, Tincture of Saffron*. (Croc. Angl. cont. $\mathfrak{Z}\text{j}$, alcohol. dilut. $\mathfrak{Z}\text{xv}$. Ph. E.) Slightly stomachic [?]. Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{ij}$.

TINCTURA DE CROCO COMPOS'ITA, *Elizir'ium de Garus, Alcoola'tum de Croco Compos'itum, Compound Tincture of Saffron, Elizir of Garus*. (Aloes Socotr. 320 p., myrrh. 64 p., croci. 32 p., cinnam., nuc. moschat., caryoph. $\mathfrak{a}\mathfrak{a}$, 16 p., alcohol. 1000 p. ag. flor. aurant. 5000 p. Digest for two days; distil 4000 p. and add syrup of capillaire, 5000 p. Ph. P.) Aromatic, cordial, stomachic. It is also called *Elizir cordia'le* et *stomach'icum*, and *E. anticoll'icum crocata'tum*.

TINCTURA CROTONIS, *Tincture of Croton*. Under this name Mr. E. Wilson recommends 'a most valuable cutaneous stimulant,' made by adding four fluidounces of alcohol to one ounce of the bruised seeds of *Croton tiglium*.

TINCTURA CROTONIS ELEUTHERIÆ, *T. cascarillæ*.

TINCTURA CUBE'Æ, *Tincture of Cubebs*. (Cubeb. cont. $\mathfrak{Z}\text{iv}$, alcohol. dilut. Oij. Macerate for 14 days, express, and filter. Ph. U. S.) It may, also, be prepared by displacement. Dose, one or two fluidrachms.

TINCTURA CUSPARIÆ, *Tinctura Augusturæ*.

TINCTURA DIGITA'LIS, *T. Digitalis purpu'reæ, Tincture of Digitalis*. (Digital. $\mathfrak{Z}\text{iv}$, alcohol. dilut. Oij. Macerate for 14 days, express, and filter.) It may, also, be prepared by displacement. Use, same as that of the leaves. Dose, gtt. x.

TINCTURA ER'GOTÆ, *Tincture of Ergot*. (Ergot. in pulv. crass. $\mathfrak{Z}\text{vij}$ (avoid.), Sp. tenuior. Oij. (Imp. meas.) Macerate for 14 days, express, and filter. Ph. D.) Dose, f $\mathfrak{Z}\text{j}$ to f $\mathfrak{Z}\text{ij}$.

TINCTURA ER'GOTÆ ÆTHE'REA, *Ethe'real Tincture of Ergot*. (Ergot. contus. $\mathfrak{Z}\text{xv}$, Æther. Oij. (Imp. meas.) Macerate for 7 days, express, and filter. Ph. L.) Dose, 60 to 90 minims.

TINCTURA FERRIFUGA DOCTORIS HUXHAMI, *T. cinchonæ composita*.

TINCTURA FERRI ACETA'TIS, *Tincture of Acetate of Iron*. (Potass. acetat. $\mathfrak{Z}\text{ij}$, sulph. ferri. $\mathfrak{Z}\text{j}$, sp. rectif. Oij. Rub the acetate and sulphate into a soft mass. Dry and rub with the spirit. Digest for 7 days, and pour off. Ph. D.) Tonic and astringent. Dose, gtt. xx to f $\mathfrak{Z}\text{j}$. The Dublin college directs a *Tinctura Acetatis Ferri cum Alcoho'le*, which differs from the other, in having half the quantity, only, of the Acetate of Potass.

TINCTURA FERRI AMMONIA'TA, *T. Ferri Ammoniacalis seu Florum martia'lium seu Martis Mynsich'ti, Tincture of Ammoniated Iron*. (Ferri ammon. $\mathfrak{Z}\text{iv}$, sp. ten. Oj. Ph. L.) Use—same as that of the ammoniated iron. Dose, f $\mathfrak{Z}\text{ss}$ to f $\mathfrak{Z}\text{j}$.

TINCTURA FERRI AMMO'NIO-CHLO'RIDI, *Tincture of Ammonio-chloride of Iron*. (Ferri ammonio-chlorid. $\mathfrak{Z}\text{iv}$, alcohol. dilut., aquæ destillat.

33 Oj. (Imp. meas.) Dissolve and filter. *Ph. L.*)
Dose, gtt. x to xx.

TINCTURA FERRI CHLORIDI, *T. ferri muriatis.*

TINCTURA FERRI MURIATIS, *T. Ferri chloridi* (*Ph. U. S.*) seu *seuqui-chloridi* seu *Murtis in Spiritu Salis* seu *Murtis cum Spiritu Salis* seu *Ferri Muriat'is*, *Liquor Ferri Muriat'is*, *Tincture of Muriate of Iron*, *Tincture of Steel*, *Alcohol ferrat'us*. (*Ferri subcarb. ibss, acid. muriat. Oj, alcohol. Oij.*) Add the acid to the subcarbonate in a glass vessel, and shake during three days. Pour off and add the spirit. *Ph. U. S.*) Tonic and styptic. It has been given, also, with advantage in erysipelas and scarlatina in the dose of \mathcal{M} x to \mathcal{M} xx, every two or three hours. Dose, gtt. x to xx.

Bestucheff's Tincture or *Klaproth's Tincture* is an ethereal tincture of chloride of iron. It is composed of dry perchloride of iron, one part; *Hoffmann's anodyne liquor*, seven parts.

The *Antivenereal Drops*, at one time so celebrated at Amsterdam, were examined by Scheele, and found to be a tincture of chloride of iron and corrosive sublimate.

TINCTURA FERRI SESQUICHLORIDI, *T. Ferri muriatis*—*t. Ferulæ Assafoetidæ*, *T. Assafoetidæ*—*t. Florum martialium*, *T. ferri ammoniata*—*t. Foetida*, *T. assafoetidæ*—*t. Foetida ammoniata*, *T. castorei composita*—*t. Fuliginis*, see *Fuligo*.

TINCTURA GAL'BANI, *Tincture of Galbanum*. (*Galb. 3ij, sp. tenuior. Oij. Ph. D.*) Stimulant and antispasmodic. Dose, f3j to f3ij.

TINCTURA GALLA'RUM, *T. Gallæ*. (*Ph. U. S.*) *Tincture of Galls*. (*Gall. contus. 3iv, alcohol. dilut. Oij.*) Macerate for 14 days, express and filter. *Ph. U. S.*) It may, also, be prepared by displacement. Astringent. Dose, f3j to f3ij.

TINCTURA GENTIANÆ COMPOSITA, *Tinct. Ama'ra*, *Elizir Stomach'icum*, *Compound Tincture of Gentian*, *Spirit Bitters*. (*Gent. concis. 3ij, aurant. cort. 3j, cardam. cont. 3ss, alcohol. dilut. Oij.*) Macerate for 14 days, express, and filter. It may, likewise, be prepared by displacement. *Ph. U. S.*) Tonic and stomachic. Dose, f3j to f3ij.

Brodum's Nervous Cordial consists of the *Tinctures of Gentian*, *Columba*, *Cardamom*, and *Bark*, with the *Compound Spirit of Lavender* and *Wine of Iron*.

Stoughton's Elizir is a compound tincture of gentian. The *Elizir'ium Docto'ris Stoughton* seu *Tinctu'ra Ama'ra* of the *Parisian Codex* is prepared as follows:—*Absinth. summit. exsiccata*, *Teucri chamædrys*. *Rad. gentian.*, *cort. aurant.*, 33 3vj, *casca'illa 3j, rhei 3ss, aloe socotr. 3j, alcohol. Oij.*) Digest.

TINCTURA GUAIA'CI, *T. Guaiaci Officin'alis*, *Tincture of Guaiacum*. (*Guaiac. ibss, alcohol. Oij.*) Macerate for 14 days, and filter. *Ph. U. S.*) Stimulant and sudorific. Used especially in rheumatic and arthritic cases. Nearly the same as *Hill's Essence of Bardana* or *Burdock*.

TINCTURA GUAIA'CI AMMONIATA, *T. Guaiaci'na Volat'ilis*, *T. Guaiaci*, (*Ph. L. 1788.*) *Elizir Guaiaci Volatilis*, *Alcohol cum Guaiaco officin'ale ammoniat'us*, *A. ammoniæ et Guaiaci*, *Ammoniated Tincture of Guaiacum*, *Volatile Tincture of Guaiacum*. (*Guaiac. pulv. 3iv, sp. ammon. arom. Oiss.*) Macerate for 14 days, and filter. *Ph. U. S.*) Properties same as the last. Dose, f3j to f3ij.

Hatfield's Tincture—a nostrum—consists of *Guaiacum* and *Soap*, 33 3ij. rectified spirit, *Oiss.*

TINCTURA HELLEB'ORI, *T. Helleb'ori Nigri*, *T. Melamp'odii*, *Tincture of Black Hellebore*. (*Helleb. nigr. cont. 3iv, alcohol. dilut. Oij.*) Macerate for 14 days, express and filter. It may also be prepared by displacement. *Ph. U. S.*) Dose, f3ss to f3j.

TINCTURA HIERA, *Vinum aloes.*

TINCTURA HU'MULI, *T. Lu'puli*, *Tincture of Hops*. (*Humuli, 3v, alcohol. dilut. Oij.*) Tonic and sedative. Dose, f3ss to f3ij.

TINCTURA HYOSCY'AMI, *T. Hyoscyami nigri*, *Tincture of Henbane*. (*Hyoscyam. fol. 3iv, alcohol. dilut. Oij.*) Macerate for 14 days, express and filter. It may also be prepared by displacement. *Ph. U. S.*) Narcotic, anodyne. Dose, gtt. xx. to f3ij.

TINCTURA IO'DINI, (*Ph. U. S.*, 1842,) *T. Iodin'ii*, (*Ph. U. S.*, 1851,) *Alcohol Io'dii*, *Tincture of Iodine*. (*Iodin. 3j, alcohol. Oj.*) Dissolve. *Ph. U. S.*) The tincture spoils by keeping, or at least, deposits the iodine. It has been much used in goitre, &c. Dose, ten drops three times a day.

An ethereal and a chloroform tincture of iodine have been used, to affect the system with iodine by inhalation.

TINCTURA IODINI COMPOSITA, (*Ph. U. S.*, 1842,) *T. Iodin'ii Composita*, (*Ph. U. S.*, 1851,) *Compound Tincture of Iodine*. (*Iodin. 3ss, Potassii iodid. 3j, Alcohol. Oj.*) Dissolve. *Ph. U. S.*) Dose, 10 to 30 drops.

TINCTURA JALA'PII, *T. Jalapæ*, (*Ph. U. S.*) seu *Convolvuli Jalapæ*, *Vali'ga*, *Tincture of Jalap*. (*Jalap. pulv. 3vi, Alcohol. dilut. Oij.*) Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be made by displacement. Cathartic. Dose, f3j to f3iv.

TINCTURA JALA'PII COMPOSITA, *Elizir Jalapæ* *Compositum* seu *purgans seu anthelmin'ticum Suco'rum*, *Essentia cathol'ica purgans Roth'ii*, (*F.*) *Eau de vie Allemande*. (*Jalap. ibss, Scammon. 3j, Alcohol. (22°) Ovj.*) Macerate for eight days. *Ph. P.*) Dose, 60 to 100 drops.

TINCTURA JAPONICA, *Tinctura catechu*—*t. Kinæ kinæ ætherea composita*, *Tinctura cinchonæ ætherea composita*—*t. Kinkinæ amara*, *Tinctura cinchonæ amara*.

TINCTURA KINO, *Tincture of Kino*. (*Kino*, in pulv., 3vj; *Alcohol. dilut. q. s.*) Put the kino, mixed with an equal bulk of sand, into a percolator, and gradually pour on diluted alcohol until f3viij of filtered liquor pass. The tincture should be renewed frequently, and kept in closely stopped bottles. *Ph. U. S.*) Astringent. Dose, f3j to f3ij.

TINCTURA KRAMER'IE, *Tincture of Rhatany*. (*Kramer. in pulv. 3vj, alcohol. dilut. Oij.*) Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Dose, f3j to f3ij.

TINCTURA LACTUCA'RIL, *Tincture of Lactucarium*. (*Lactucar. in pulv. 3iv, Sp. tenuior. Oij.*) (Imp. meas.) Best prepared by percolation. *Ph. E.*) Dose, 30 minims to f3j.

TINCTURA LAURI CINNAMOMI, *Tinctura cinnamomi*—*t. Lavandulæ composita*, *Spiritus lavand. comp.*

TINCTURA LIMO'NIS, *Tincture of Lemon peel*. (*Limon. cort. 3iiiss, Sp. tenuior. Oij.*) (Imp. meas.) Macerate for 7 days, express and filter. *Ph. D.*) Dose, f3j to f3ij.

TINCTURA LOBE'LIÆ, *Tincture of Lobelia*. (*Lobel. 3iv, alcohol. dilut. Oij.*) Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Dose, f3j to f3ij.

TINCTURA LOBE'LIÆ ÆTHE'REA, *Ethe'real Tincture of Lobelia*. (*Lobel. pulv. 3v, Sp. æther. sulph. Oij.*) (Imp. meas.) *Ph. E. and L.*) Dose, same as the last.

TINCTURA LUPULI, *Tinctura Humuli*, *T. Lupulinæ*.

TINCTURA LUPULINÆ, *T. Lu'puli*, (*Ph. E.*) *Tincture of Lu'puline*. (*Lupulin. 3iv, alcohol.*

Oij. Macerate for 14 days, and filter. *Ph. U. S.* Dose, f ʒi to f ʒij.

TINCTURA LYTTE, Tinctura cantharidis—t. Martis Glauberi, see Ferrum tartarizatum—t. Martis in spiritu salis, Tinctura ferri muriatis—t. Martis Mynsichti, Tinctura ferri ammoniata—t. Martis vinosa, Vinum ferri.

TINCTURA MATICO, Tincture of Matico. (*Matico* in pulv. crass. ʒviij, (avoid.) *Sp. tenuior*. Oij. (Imp. meas.) Macerate for 14 days, express, and filter. *Ph. D.*) Dose, f ʒi to f ʒij.

TINCTURA MELAMPODII, Tinctura hellebori nigri—t. Meleses vesicatorii, Tinctura cantharidis—t. Metallorum, Liliun Paracelsi—t. Mimosa catechu, Tinctura catechu.

TINCTURA MOSCHI, Tincture or Essence of Musk. (*Mosch.* in pulv. redact. ʒij. *sp. vini rect.* Oj. Dose, f ʒi to f ʒiv.

TINCTURA MYRRHÆ, Tincture of Myrrh, Simple Tincture of Myrrh. (*Myrrhæ contusæ* ʒiv, alcohol Oij. Macerate for 14 days, and filter. *Ph. U. S.*) Tonic, deobstruent, antiseptic, and detergent. It is chiefly used in gargles, and applied to foul ulcers, spongy gums, &c.

Hudson's Preservative for the Teeth and Gums consists of *Tinct. myrrh.*, *tinct. cinch.*, *aq. cinnam.* aa ʒijij, *eau d'arquebuse* ʒj, *pulv. gum. Arab.* ʒss. *M.*

TINCTURA NERVINA BESTUCHEFFII, Tinctura seu alcohol sulphurico-æthereus ferri—t. Nervina Halensis, Tinctura seu alcohol sulphurico-æthereus ferri—t. Nervino-tonica, Lamotte's, Alcohol seu Tinctura sulphurico-æthereus ferri.

TINCTURA NUCIS VOMICÆ, Tincture of Nuc Vomica. (*Alcohol. s. g.* 837, f ʒj; *Dry Extract of Nuc Vomica*, 3 gr.) Dose, gtt. v to xxx, in cases where the nuc vomica is indicated.

It is directed, in the last edition of the *Ph. U. S.* (1851), to be prepared as follows:—*Nucis vomicæ rasur.*, ʒviij; *alcohol*, Oij. Macerate for 14 days, express, and filter. It may also be prepared by the process of displacement. Dose, gtt. v to xx.

TINCTURA OLEI MENTHÆ PIPERITÆ, Tincture of Oil of Peppermint, Essence of Peppermint. (*Ol. menthæ piperitæ*, f ʒij, *alcohol*, Oj. Dissolve. *Ph. U. S.*) Dose, 5 drops, as a carminative.

TINCTURA OLEI MENTHÆ VIRIDIS, Tincture of Spearmint, Essence of Spearmint, is made in the same manner. *Ph. U. S.*

TINCTURA OPII, T. Thebæica seu Sedati'va, Theriaca cœlestis, Alcohol cum Opio, Tincture of Opium, Liquid Laudanum. (*Opii pulv.* ʒijss, *alcohol. dilut.* Oij. Macerate for 14 days, express, and filter. *Ph. U. S.*) Nineteen minims contain about one grain of opium. Properties, those of opium. Dose, gtt. xx to lx and more. By macerating the dregs remaining after making the Tinctura Opii in a solution of Tartaric Acid, a solution is obtained, which is said to be devoid of the exciting and constipating properties of the tincture of opium. Gray calls it *Haden's Liquor Opii Sedati'vus*.

TINCTURA OPII ACETATA, Acetated Tincture of Opium. (*Opii ʒij*; *aceti*, f ʒxij; *alcohol*. Oss. Rub the opium with the vinegar; add the alcohol, and having macerated for 14 days, express and filter. *Ph. U. S.*) Twenty drops are equivalent to a grain of opium.

TINCTURA OPII AMMONIATA, Ammoniated Tincture of Opium, Edinburgh Pargoric Elixir. (*Flor. benz.*, *croci*, aa ʒijij, *opii ʒij*, *ol. anisi* ʒss, *alcohol. ammoniat.* f ʒxvj. Digest. *Ph. E.*) f ʒj contains one grain of opium. Used like the Tinctura Camphoræ Composita.

TINCTURA OPII CAMPHORATA, Tinctura camphoræ composita—t. Plumbosa, Liquor Plumbi subacetatis dilutus.

TINCTURA QUASSIÆ, T. Quassiæ excel'sæ,

Tincture of Quassia. (*Quassiæ rasur.* ʒij, *alcohol. dilut.* Oij. Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Tonic. Dose, f ʒi to f ʒij.

TINCTURA QUASSIÆ COMPOSITA, Compound Tincture of Quassia (*Cardam. sem. contus.*, *Cocci contus.* aa ʒss; *Cinnam.*, *Quassiæ*, aa ʒvj; *Uvar. pasar.* ʒviij; *Sp. tenuior*. Oij (Imp. meas.). Digest for 7 days; express and filter. May be prepared also by percolation. *Ph. E.*) Dose, as a tonic, f ʒi or f ʒij.

TINCTURA QUINÆ COMPOSITA, Compound Tincture of Quina. (*Quinæ sulph.* ʒv and ʒij; *Tinct. aurantii*, Oij, (Imp. meas.) Oij. Digest for 7 days and filter. *Ph. L.*) f ʒj, contains a grain of the sulphate.

TINCTURA QUINÆ SULPHATIS, Tincture of Sulphate of Quina. (*Sulphate of Quina*, gr. vj; *alcohol.* (847) f ʒj. *M.*) Dose, f ʒi to f ʒiv, in the day.

Tinctura Cincho'niæ Sulpha'tis may be made in a similar manner.

TINCTURA RHABARBARI, Tinctura rhei—t. Rhabarbari composita, Tinctura rhei composita—t. Rhabarbari vinosa, Vinum rhei palmati.

TINCTURA RHEI, T. Rhabarb'ari seu Rhabarbari spirituo'sa seu Rhei seu Rhei palma'ti, Tincture of Rhubarb. (*Rhei contus.* ʒij, *cardam. cont.* ʒss, *alcohol. dilut.* Oij. Macerate for 14 days, express, and filter. It may also be prepared by displacement. *Ph. U. S.*) Stomachic or purgative, according to the dose employed: f ʒi to ʒij, as a stomachic; f ʒiv to ʒj, as a purgative.

TINCTURA RHEI COMPOSITA, T. Rhabarbari composita, Compound Tincture of Rhubarb. (*Rhei rad. concis.* ʒij, *glycyrrhiz. rad.* cont. ʒss, *zingib. rad. concis.*, *croci stigmat.* aa ʒij, *aqvæ* f ʒxij, *sp. ten.* Oj. *Ph. L.*) Properties same as the last.

TINCTURA RHEI ET AL'OE, Elixir sacrum, Tincture of Rhubarb and Aloes. (*Rhei concis.* ʒx, *aloes pulv.* ʒvj, *cardam. cont.* ʒss, *alcohol. dilut.* Oij. Macerate for 14 days, express, and filter. *Ph. U. S.*) Properties like the last.

Rymer's Car'diac Tincture consists of an infusion of Capsicum, Camphor, Cardamom seeds, Rhubarb, Aloes and Castor in Proof Spirit, with a very small quantity of Sulphuric Acid.

TINCTURA RHEI ET GENTIANÆ, Tinctura Rhei Amara, Tincture of Rhubarb with Gentian. (*Rhei cont.* ʒij, *gentian. cont.* ʒss, *alcohol. dilut.* Oij. Macerate for 14 days, express and filter. *Ph. U. S.*) It may also be prepared by displacement. Properties same as the last.

TINCTURA RHEI ET SENNÆ, Tincture of Rhubarb and Senna, Warner's Gout Cordial. (*Rhei cont.* ʒj, *sennæ*, ʒij, *coriandr. cont.*, *fœnicul. cont.* aa ʒj, *santal. rasur.* ʒij, *croci*, *ext. glycyrrhiz. rad.* aa ʒss, *urar. pasar.* acinis exemptis, *ibss*, *alcohol. dilut.* Oij. Macerate for 14 days; express and filter. *Ph. U. S.*) Dose, f ʒss to f ʒij.

TINCTURA RHEI, Tinctura rhei—t. Rhei amara, Tinctura rhei et gentianæ—t. Sacra, Vinum aloes.

TINCTURA SANGUINARIÆ, Tincture of Blood-root. (*Sanguinar. cont.* ʒiv, *alcohol. dilut.* Oij. Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Dose, 30 to 60 drops.

TINCTURA SAPONIS CAMPHORATA, Linimentum saponis—t. Saponis et opii, Linimentum saponis et opii.

TINCTURA SCILLÆ, Tincture of Squill. (*Scillæ ʒiv*, *alcohol. dilut.* Oij. Macerate for 14 days, express, and filter. It may also be made by displacement. *Ph. U. S.*) Dose, gtt. xv to lx.

TINCTURA SEDATIVÆ, Tinctura opii.

TINCTURA SENNÆ, *T. Sennæ Composita*, *Elixir Salutis*, *Daffy's Elixir*, *Tincture of Senna*. (*Senna fol.* $\mathfrak{z}\text{ij}$, *carui sem.* cont. $\mathfrak{z}\text{ij}$, *cardam.* sem. cont. $\mathfrak{z}\text{j}$, *urarum passarum demptis acinis*, $\mathfrak{z}\text{iv}$, *sp. ten.* Oij . *Ph. L. & D.*) Stomachic, carminative and cathartic. Dose, $\mathfrak{f}\mathfrak{z}\text{ij}$ to $\mathfrak{f}\mathfrak{z}\text{j}$.

TINCTURÆ SENNÆ ET JALAPÆ, *Tincture of Senna and Jalap*. (*Sennæ* $\mathfrak{z}\text{ij}$, *jalap. pulv.* $\mathfrak{z}\text{j}$, *coriandr.* cont., *carui* cont., $\mathfrak{a}\mathfrak{a}$ $\mathfrak{z}\text{ss}$, *cardam.* cont. $\mathfrak{z}\text{ij}$, *sacch.* $\mathfrak{z}\text{iv}$, *alcohol. dilut.* Oij . Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Dose, $\mathfrak{f}\mathfrak{z}\text{ij}$ to $\mathfrak{f}\mathfrak{z}\text{j}$.

TINCTURA SERPENTARIÆ, *T. Serpentariæ* seu *Virginianæ* seu *Aristolochiæ Serpentaria*, *Tincture of Snakeroot*. (*Rad. serpent.* $\mathfrak{z}\text{ij}$, *alcohol. dilut.* Oij .* Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Tonic. Dose, $\mathfrak{f}\mathfrak{z}\text{ss}$ to $\mathfrak{f}\mathfrak{z}\text{ij}$.

TINCTURA STOMACHICA, *Tinctura cardamomi composita*.

TINCTURA STRAMONII, *Tincture of Stramonium*. (*Stramon. sem.* cont. $\mathfrak{z}\text{iv}$, *alcohol. dilut.* Oij . Macerate for 14 days, express, and filter. *Ph. U. S.*) It may also be prepared by displacement. Dose, ten to twenty minims.

TINCTURA STRYCHNINÆ, *Tincture of Strychnia*. (*Alcohol. s. g.* $\mathfrak{z}\mathfrak{z}\mathfrak{v}$, Oj , *Strychnia*, *gr. iij.*) Dose, 6 drops to 24. Properties, those of strychnia.

TINCTURA (seu ALCOHOL) SULPHURICO-ÆTHEREUS FERRI, *Gutta nervinæ*, *Liquor Anodynus Martialis*, *Alcohol æthe rei ferri*, *tus*, *Naphtha Vitrioli Martialis*, *Solutio Muriatis ferrici æthereæ*, *Spiritus sulphurico-æthereus martialis*, *Tinctura nervinæ Halen'sis* seu *Tonic-nervinæ Halen'sis* seu *auræa* seu *nervina Bestucheffii* seu *nervino-tonica* (Lamotte's) seu *ætherea ferri*, *Æther Martialis*, *De La Motte's Golden Drops*, *Bestucheff's Nervous Tincture*, (*F.*) *Elixir d'Or de M. le Général De La Motte*. An ethereal solution of muriate of iron. It is much used in gout, hypochondriasis, &c.

TINCTURA SULPHURIS VOLATILIS, *Liquor fumans Boylei*—*t. Thebaica*, *Tinctura opii*, *Vinum opii*.

TINCTURA TOLUIFERÆ BAL'SAMI, *T. Balsami Tolutani*, *T. Tolutani* (*Ph. U. S.*, 1842.), *T. Tolutana* (*Ph. U. S.*, 1851), *Tincture of Balsam of Tolu*, (*Bals. Tolutan.* $\mathfrak{z}\text{ij}$, *alcohol* Oij .) Reputed to be expectorant and corroborant. Dose, $\mathfrak{f}\mathfrak{z}\text{ss}$ to $\mathfrak{f}\mathfrak{z}\text{j}$, or more.

TINCTURA TONICA NERVINA HALENSIS, *Tinctura (seu Alcohol) sulphurico-æthereæ ferri*.

TINCTURA VALERIANÆ, *Tincture of Valerian*. (*Valerian.* cont. $\mathfrak{z}\text{iv}$, *alcohol. dilut.* Oij . Macerate for 14 days, express, and strain. It may also be prepared by displacement. *Ph. U. S.*) Stimulant and antispasmodic. Dose, $\mathfrak{f}\mathfrak{z}\text{ss}$ to $\mathfrak{f}\mathfrak{z}\text{ij}$.

TINCTURA VALERIANÆ AMMONIATA, *T. Valerianæ Volatilis*, *Ammoniated or Volatile Tincture of Valerian*. (*Valerianæ* cont. $\mathfrak{z}\text{iv}$, *sp. ammon. arom.* Oij . Macerate for 14 days, express, and filter. It may also be made by displacement. *Ph. U. S.*) Like the former. Dose, $\mathfrak{f}\mathfrak{z}\text{ss}$ to $\mathfrak{f}\mathfrak{z}\text{ij}$.

TINCTURA VERATRIS, *T. Veratri albi*, *Tincture of White Hellebore*. (*Rad. veratri alb.* cont. $\mathfrak{z}\text{vij}$, *alcohol. dilut.* Oijss . *Ph. E.*) Emetic and cathartic; but not very manageable. Dose, gtt. v to x .

TINCTURA ZINGIBERIS, *T. Amomi Zingiberis*, *Tincture of Ginger*. (*Zingib. cont.* $\mathfrak{z}\text{vij}$, *alcohol.* Oij . Macerate for 14 days, express, and filter. It may also be made by displacement. *Ph. U. S.*) Stimulant and carminative. Dose, $\mathfrak{f}\mathfrak{z}\text{ss}$ to $\mathfrak{f}\mathfrak{z}\text{j}$.

TINCTURE, *Tinctura*—*t. of Aconite*, *Tinctura aconiti*—*t. of Aconite root*, *Tinctura aconiti radicis*—*t. of Aloes*, *Tinctura aloes*—*t. of Aloes*, compound, *Tinctura aloes composita*—*t. of Aloes*, ethereal, *Tinctura aloes æthereæ*—*t. of Ammonia*,

compound, *Spiritus ammoniæ succinatus*—*t. of Angustura*, *Tinctura angusturæ*—*t. of Assafetida*, *Tinctura assafetidæ*—*t. of Balsam of Tolu*, *Tinctura toluiferæ balsami*—*t. of Bark*, *Tinctura cinchonæ*—*t. of Bark*, ammoniated, *Tinctura cinchonæ ammoniata*—*t. of Bark*, compound, *Tinctura cinchonæ composita*—*t. of Bark*, *Huxham's*, *Tinctura cinchonæ composita*—*t. of Bark*, volatile, *Tinctura cinchonæ ammoniata*—*t. of Belladonna*, *Tinctura belladonnæ*—*t. of Benzoïn*, compound, *Tinctura benzoïn composita*—*t. of Blistering fly*, *Tinctura cantharidis*—*t. of Bloodroot*, *Tinctura sanguinariæ*—*t. of Buchu*, *Tinctura buchu*—*t. of Camphor*, compound, *Tinctura camphoræ composita*—*t. of Cantharides*, *Tinctura cantharidis*—*t. of Capsicum*, *Tinctura capsici*—*t. of Cardamoms*, *Tinctura cardamomi*—*t. of Cardamoms*, compound, *Tinctura cardamomi composita*—*t. of Cardiac*, *Rymer's*, *Tinctura rhei et aloës*—*t. of Cascarella*, *Tinctura cascariillæ*—*t. of Castor*, *Tinctura castorei*—*t. of Castor*, compound, *Tinctura castorei composita*—*t. of Catechu*, *Tinctura catechu*—*t. of Cayenne pepper*, *Tinctura capsici*—*t. of Chiretta*, *Tinctura chirettæ*—*t. of Cinchona*, *Tinctura cinchonæ*—*t. of Cinchona*, compound, *Tinctura cinchonæ composita*—*t. of Cinnaon*, *Tinctura cinnaonmi*—*t. of Cinnamon*, compound, *Tinctura cinnamomi composita*—*t. of Cochineal*, *Tinctura cocci cacti*—*t. of Colchicum*, compound, *Spiritus colchici ammoniatus*—*t. of Colchicum seeds*, *Tinctura colchici*—*t. of Columbo*, *Tinctura calumbæ*—*t. of Croton*, *Tinctura crotonis*—*t. of Cubebs*, *Tinctura cubebæ*—*t. of Digitalis*, *Tinctura digitalis*—*t. of Ergot*, *Tinctura ergotæ*—*t. of Ergot*, ethereal, *Tinctura ergotæ æthereæ*—*t. of Galbanum*, *Tinctura galbani*—*t. of Galls*, *Tinctura gallarum*—*t. of Gentian*, compound, *Tinctura gentianæ composita*—*t. of Ginger*, *Tinctura zingiberis*—*t. of Gout*, *D. Wilson's*, see *Colchicum autumnale*—*t. of Guaiacum*, *Tinctura guaiaci*—*t. of Guaiacum*, ammoniated, *Tinctura guaiaci ammoniata*—*t. of Guaiacum*, volatile, *Tinctura guaiaci ammoniata*—*t. Hatfield's*, see *Tinctura guaiaci ammoniata*—*t. of Hellebore*, black, *Tinctura hellebori*—*t. of Hellebore*, white, *Tinctura veratri*—*t. of Hemlock*, *Tinctura conii*—*t. of Henbane*, *Tinctura hyoscyami*—*t. of Hops*, *Tinctura humuli*—*t. of Indian hemp*, *Tinctura cannabis Indicæ*—*t. of Iodine*, *Tinctura iodini*—*t. of Iodine*, compound, *Tinctura iodini composita*—*t. of Iron*, acetate of, *Tinctura ferri acetatis*—*t. of Iron*, ammoniated, *Tinctura ferri ammoniata*—*t. of Iron*, ammonio-chloride of, *Tinctura ferri ammonio-chloridi*—*t. of Iron*, muriated, *Tinctura ferri muriatis*—*t. of Jalap*, *Tinctura jalapii*—*t. of Kino*, *Tinctura kino*—*t. Klaproth's*, see *Tinctura ferri muriatis*—*t. of Lactucarium*, *Tinctura lactucarii*—*t. of Lemon peel*, *Tinctura limonis*—*t. of Lobelia*, *Tinctura lobeliæ*—*t. of Lobelia*, ethereal, *Tinctura lobeliæ æthereæ*—*t. of Ludwig*, see *Ferrum tartarizatum*—*t. of Lupuline*, *Tinctura lupulinæ*—*t. of Matico*, *Tinctura matico*—*t. of Musk*, *Tinctura moschi*—*t. of Myrrh*, *Tinctura myrrhæ*—*t. Nervous*, *Bestucheff's*, *Tinctura seu alcohol sulphurico-æthereus ferri*—*t. of Nux vomica*, *Tinctura nucis vomicæ*—*t. of Oil of Peppermint*, *Tinctura olei menthæ piperitæ*—*t. of Oil of Spearmint*, *Tinctura olei menthæ viridis*—*t. of Opium*, *Tinctura opii*—*t. of Opium*, acetated, *Tinctura opii acetata*—*t. of Opium*, ammoniated, *Tinctura opii ammoniata*—*t. of Opium*, camphorated, *Tinctura camphoræ composita*—*t. of Orange peel*, *Tinctura aurantii*—*t. of Quassia*, *Tinctura quassiæ*—*t. of Quassia*, compound, *Tinctura quassiæ composita*—*t. of Quinia*, sulphate of, *Tinctura quiniæ sulphatis*—*t. of Quinia*, compound, *Tinctura quiniæ composita*—*t. of Rhatany*, *Tinctura kramerisæ*—

t. of Rhubarb, *Tinctura rhei*—t. of Rhubarb and aloes, *Tinctura rhei et aloes*—t. of Rhubarb, compound, *Tinctura rhei composita*—t. of Rhubarb and gentian, *Tinctura rhei et gentianæ*—t. of Rhubarb and senna, *Tinctura rhei et sennæ*—t. of Saffron, *Tinctura croci sativi*—t. of Saffron, compound, *Tinctura de croco composita*—t. of Senna, *Tinctura sennæ*—t. of Senna and jalap, *Tinctura sennæ et jalapæ*—t. of Snake root, *Tinctura serpentaria*—t. of Soap, camphorated, *Limentum saponis*—t. of Spanish fly, *Tinctura cantharidis*—t. of Spearmint, *Tinctura olei menthæ viridis*—t. of Squill, *Tinctura scillæ*—t. of Steel, *Tinctura ferri murialis*—t. Stomach, *Tinctura cardamomi composita*—t. of Stramonium, *Tinctura stramonii*—t. of Strychnia, *Tinctura strychniæ*—t. for the Teeth, Greenough's, see *Spiritus armoraciæ compositus*—t. Toothache, *Tinctura antidontalgica*—t. of Valerian, *Tinctura valerianæ*—t. of Valerian, ammoniated, *Tinctura valerianæ ammoniata*—t. of Valerian, volatile, *Tinctura valerianæ ammoniata*—t. of Wormwood, compound, *Tinctura absinthii composita*.

TINDER, *Boletus ignarius*.

TINEA, Porri—t. Faciei, *Porrigo larvalis*—t. Favosa, *Porrigo favosa*—t. Ficosa, *Porrigo scutulata*—t. Furfuracea, *Porrigo furfurans*—t. Granulata, *Porrigo scutulata*, *Porrigo lupinosa*—t. Lactea, *Porrigo larvalis*—t. Lupina, *Porrigo lupinosa*—t. Mucosa, *Porrigo lupinosa*—t. Porriginosa, *Porrigo furfurans*—t. Tarsi, *Ophthalmia tarsi*—t. Tondens, *Porrigo decalvans*.

TINGLING, from *tinnire*, 'to jingle.' (Sc.) *Dirling*. A sharp, purring or thrilling sensation, not always painful.

TINKER'S WEED, *Triosteum*.

TINKLING, METALLIC, *Tintement métallique*.

TINNIMENTUM METALLICUM, *Tintement métallique*.

TINNITUS AU'RIMUM, (*tinnio*, *tinnitum*, 'to tinkle,' 'ringing of the ears.' *Fluctuatio seu Sonitus seu Susur'rus seu Sib'itus seu Strep'itus seu Sibilismus au'rium*, *Parac'usis imaginaria*, *Ote'chus*, *Syrig'mus*, *Syring'mos*, *Sibilus clango'nis*, *Echos*, *Eneche'ma*, *Bombus*, *Bombitatio*, *Murmur*, (F.) *Tintement*, T. d'Oreille, *Tintouin*, *Bourdonnement*, *Bruissement*. An imaginary sound, like the ringing of a bell, the noise of wind, the murmur of water, &c., heard in health and in sickness. It is, often, an accompaniment of cerebral disorder.

TINTEMENT, *Tinnitus aurium*—t. *Bullaire*, T. *métallique*.

TINTEMENT MÉTALLIQUE (F.), *Cliquetis métallique*, *Tintement bullaire*, *Tinnimentum metallicum*, *Canor stethoscop'icus*, *Metallic Respiration*, *Metallic Voice*, *Metallic Tinkling*. A particular noise, heard by the stethoscope when applied to the chest, and which, according to Laënnec, resembles that caused by striking glass, metal, or porcelain, with a pin; but is, perhaps, more like the sound of the keys of a musical snuff-box. The *tintement métallique* or *metallic tinkling* is heard on causing the patient to speak or breathe; but is more distinct when he coughs. When the phenomenon is not so strongly marked, it produces only the *metallic resonance*. The 'metallic tinkling' is a pathognomonic symptom of a communication between the bronchia and cavity of the chest; in other words, of *Pneumothorax*.

TINTEMENT D'OREILLE, *Tinnitus aurium*.

TINTINNABULUM, *Uvula*.

TINTOUIN, *Tinnitus aurium*.

TIPFOOT, see *Kylosia*.

TIPSARIA, *Decoctum hordei*.

TIRE-BALLE, *Forceps* (bullet.)

TIRE-FOND (F.). A surgical instrument, formerly used to elevate the piece of bone sawed off by the trephine.

TIRE-PUS, *Pyulcon*.

TIRE-TÊTE (F.). A name given to different instruments used for extracting the head of the child when left in the uterus, and in certain cases of difficult labour. The crotchet is one of those, as well as the *Tire-tête à bascule* of Levret, &c.

TIRES, Milk sickness.

TISANE, *Pti'sana* or *Ptis'sana*, *Ptisane*, from *πτισάνη*, 'pearl barley,' itself from *πτισσω*, 'I decorticate.' (F.) *Hydrolé*. A name given to aqueous medicines, containing but little, if any, medicinal agent. The ancients gave the name particularly to the decoction of barley.

TISANE COMMUNE, *Decoctum hordei*.

TISANE DE FELTZ. A celebrated antisyphilitic, formed of *sarsaparilla*, *isinglass*, *sulphuret of antimony* and *water*, boiled together. According to some, bichloride of mercury; and, to others, arsenic enters into its composition. M. Rayer, supposing that it acts by the arsenic it contains, proposes to replace it by arseniate of potassa; and Guibourt, who is of the same opinion, by arsenious acid, in order to ensure certainty of action.

TISIC, *Phthisis*.

TISICAL, *Phthisicus*.

TISSICK, see *Phthisis*.

TISSU, *Tissue*—t. *Accidental*, see *Accidental*, and *Tissue*, *accidental*—t. *Caverneux*, *Cavernous texture*—t. *Celluleux des os*, *Cancelli*—t. *Conjonctif*, *Cellular tissue*—t. *Inodulaire*, *Tissue*, *inodular*—t. *Jaune*, see *Fibrous*—t. *Lardacé*, see *Lardaceous*—t. *Spongoïde*, *Spongioid tissue*.

TISSUE, *Textus*, *Tela*, (F.) *Tissu*. By this term, in anatomy, is meant the various parts, which, by their union, form the organs; and are, as it were, their anatomical elements. Histological anatomy is the anatomy of the tissues, which are the seat of the investigations of the pathological anatomist. The best division, indeed, of diseases would be according to the tissues mainly implicated. For the elementary tissues, see *Fibre*. The compound tissues, which, by associating or combining variously, form every organ of the body, have been variously classed: but every division is imperfect and liable to objection. The following, by Messrs. Todd and Bowman, is one of the most recent.

TABULAR VIEW OF THE TISSUES OF THE HUMAN BODY.

Examples.	
1. Simple membrane, homogeneous, or nearly so, employed alone, or in the formation of compound membranes.	Posterior layer of the cornea.—Capsule of the lens.—Sarcolemma of muscle, &c.
2. Filamentous Tissues, the elements of which are real or apparent filaments.	White and yellow fibrous tissues.—Areolar tissue.
3. Compound membranes, composed of simple membrane and a layer of cells of various forms, (epithelium or epidermis,) or of areolar tissue and epithelium.	Mucous membrane.—Skin.—True or secreting glands.—Serous and synovial membranes.
4. Tissues which retain the primitive areolar structure as their permanent character.	Adipose tissue.—Cartilage.—Gray nervous matter.
5. Sclerous or hard tissue.	Bone.—Teeth.
6. Compound tissues.	
a. Composed of tubes of homogeneous membrane, containing a peculiar substance.	Muscle.—Nerve.
b. Composed of white fibrous tissues and cartilage.	Fibro-cartilage.

TISSUE, ACCIDENTAL or ADVENTITIOUS, consists of every substance—foreign to the primitive organization of the economy, but yet organized and living—which may be developed in the anterior or at the surface of organs. Laënnec divides the accidental tissues into two sections: 1. Those

that resemble others in the animal economy, or *analogous accidental tissues*; and, 2. Those that have nothing analogous in the body, and which are always the result of a morbid process—the *heterologous* or *heteroclitie accidental tissues*. To the first class belong, ossifications; accidental fibrous, fibro-cartilaginous, cartilaginous, areolar, and corneous tissues; hairs; the serous membranes of certain encysted tumours; the mucous membranes of fistulous canals, and accidental synovial membranes. To the second belong tubercles, scirrhi, melanosis, &c.

TISSUE, AREOLAR, Cellular tissue.

TISSUE, CARTILAG'INOUS, *Hymenochondroïdes* seu *Hymenochondroïdes* (*Textura seu Tela*.)

TISSUE, CONNECTIVE, Cellular tissue—t. Cribiform, Cellular tissue—t. Fibrous, see Fibrous—t. Filamentous, Cellular tissue.

TISSUE FORMATION, Histogeny.

TISSUE, INOD'ULAR, *Tela inodula'ris*, (F.) *Inodula*, *Tissu inodulaire*. A name given by Delpech to an accidental fibrous tissue developed in suppurating wounds, which is the principal agent in cicatrization. It has, at first, the appearance of a reddish cellulose-fibrous layer; but soon loses its vascularity; and its fibres, which pass in all directions, become of a dull white colour, and of a consistence and hardness, that may be compared to those of the strongest articular ligaments. It is well seen in the cicatrix left after burns.

TISSUE, LAMINATED, Cellular tissue—t. Ligamentous, Desmoid tissue—t. Mucous, Cellular tissue—t. Porous, Cellular tissue—t. Reticulated, Cellular tissue.

TIT, Nipple.

TITHENE, Nurse.

TITHYMALUS CYPARISSIAS, *Euphorbia cyparissias*—t. Lathyris, *Euphorbia lathyris*—t. Latifolius, *Euphorbia lathyris*—t. Palustris, *Euphorbia palustris*—t. Paralias, *Euphorbia paralias*.

TITILLAMENTUM, Gargarism.

TITILLATION, Tickling.

TITTHE, Nipple.

TITTHION, Nipple.

TITTHIS, Nipple.

TITTHOS, Mamma, Nipple.

TITTY, Mamma, Nipple.

TITUBANTIA, *Bredouillement*.

TITUBATIO, Fidgets, Vacillatio—t. Linguae, Balbuties.

TJETTEK, Upas tiénté.

TO-AND-FRO-SOUND, *Bruit de frottement*.

TOAD-FLAX, *Antirrhinum linaria*—t. f. Bastard, *Comandra umbellata*.

TOAST-WATER, *Aqua tostæ panis*. Toast well half a slice of a stale quartern loaf; put it into a pitcher, and pour over it a quart of water. After two hours, decant the water from the bread. It is a common drink in febrile affections.

TOBACCO, *Nicotiana tabacum*—t. English, *Nicotiana rustica*—t. Indian, *Lobelia inflata*—t. Poison, *Hyoscyamus niger*—t. Wild, *Lobelia inflata*, *Nicotiana rustica*.

TOCETOS, Parturition.

TOCODYMYCODYRITIS MALIGNA VAGINALIS, Colpocæ puerperarum.

TOCOLOGY, Obstetrics.

TOCOS, Parturition.

TODDY. A name given, by the English, to sweet, delicious and refreshing liquors procured in India by wounding the *spathæ* of certain palms. The best is obtained from the *Cocos nucifera*.—Ainslie.

It is obtained in large quantities from *Saguë-rus saccharifer* seu *Rumphi*, *Palma Indica virna'ria* secundâ, *Saguë-rus* seu *Gomutus Gomuto*,

Aren'ga saccharif'era, *Gomuto palm*. It is called, also, *Neva*, and, by boiling, yields a coarse kind of sugar, called *Jaggery*. The inhabitants of the Moluccas used, in their wars, a liquor obtained by macerating the fruit, which the Dutch called *Hell water*, *Aqua infernalis*.

Toddy is also the common name for a mixture of spirit and water sweetened.

TODDY TREE, Mamei.

TOE, Digitus pedis.

TOEPLITZ, see Töplitz.

TOFUS, Tophus.

TOILE D'ARAIGNÉE, Aranæ tela—t. Choroidienne, see Choroid—t. de Gautier, Sparadrapum Galteri.

TOKAS, Puerpera.

TOKEN, PLAGUE, Plague token.

TOKOLOGY, Obstetrics.

TOLA, Tonsil.

TOLERANCE, *Tolerantia*; from *tolerare*, 'to bear.' The power of bearing. A word used by the Italian school of Rasori, to signify the power of bearing large doses of certain potent remedies, as tartrate of antimony and potassa. See *Contra-stimulus*.

TOLES, Tonsil.

TOLIUM, Tonsil.

TOLLES, Tonsil.

TOLLUM, Tonsil.

TOLUIFERA BAL'SAMUM, *Myroxylon* seu *Myrospermum Toluiferum*. Ord. Leguminosæ. *Sex. Syst.* Decandria Monogynia. The name of the tree which affords the Balsam of Tolu, *Balsamum Tolutanum* (Ph. U. S.), *Tolutanum*, *Toluif'era Balsami Balsamum*, *Guac'onaz*, *Red Balsam of Peru*, (F.) *Baume de Tolu*, *B. de Carthagène* ou d'*Amérique* ou de *Saint Thomas*. It has a very fragrant odour, and warm, sweetish taste; is thick, and of a reddish-yellow colour. It is stimulant and expectorant [?], and is used in coughs. It is chiefly, however, employed on account of its flavour.

TOLUTANUM, see *Toluifera balsamum*.

TOMACINUM, Iscium.

TOMACULUM, Iscium.

TOMATA, *Solanum lycopersicum*.

TOMATO, *Solanum lycopersicum*.

TOME, *Tomus*, *τὸν, τομῶς*, 'incision.' A common suffix, as in *Lithotomy*, *Bronchotomy*, &c.

TOME. Cut. Incision.

TOMETON, Knife.

TOMENTUM CER'EBRI, from *to mentum*, 'a flock of wool.' The infinite number of small vessels which pass between the pia mater and convolutions of the brain. They enter the substance of the brain at right angles.

TOMEUS, Knife.

TOMIS, Scissors.

TOMOTOCIA, Cæsarean section.

TOM-TOE, Pollex, see *Digitus pedis*.

TONE, *Tonus*, *τονος*, from *roveo*, 'I brace or give vigour.' The state of tension or firmness proper to each organic tissue. It is the effect of tonicity.

TONE, see Elasticity.

TONGA, see *Datura sanguinea*.

TONGRES, MINERAL WATERS OF. The town of Tongres is about a league from Maestricht in the Low Countries. The springs, of which there are several, are chalybeate, containing carbonates of iron and magnesia.

TONGUE, Sax. *tung*, (D.) *Tong*, *Lingua*, *Glossa*, *Glotta*, (F.) *Langue*. The tongue is a muscular, symmetrical, and very mobile organ; situate in the mouth, from the os hyoides and epiglottis to the incisor teeth. Its particular use is, to procure the sensation of savours; and it aids in the acts of sucking, mastication, deglutition, pronunciation, and expuition. It has the

shape of a pyramid, flattened above and below; rounded at its angles, and terminated, anteriorly, by a blunt point. Its *upper surface*, called the *dorsum*, is free, flat, and divided into two portions by a median furrow, *lin'ea media'na*; at the posterior extremity of which is the *foramen cœcum* or *Foramen of Morgagni*. The *inferior surface* is free, and covered by the mucous membrane on its anterior third and sides: but behind and in the middle it is attached to the lower jaw by means of the *genio-glossus*; and to the *hyoides* by the *hyoglossus*. Its *margins* are rounded; thick behind, thinner anteriorly. Its *tip*, or *point*, or *dental extremity* is rounded and free:—the *root*, *base*, or *hyoid extremity*, *Radix linguae*, is continuous with the *epiglottis* and *velum palati*. It is very thick at the *foramen cœcum*; but thinner as it approaches the *hyoid bone*. The tongue is composed of muscles, and covered by a mucous membrane—*Periglottis*, *Involucrum linguae*, *Membrana seu Tunica vaginalis seu Cutis seu Epidermis seu Tunica seu Involucrum mucosum seu Crusta villosa linguae*—which forms also the *frœnum* beneath the tongue. On this membrane are seen the different *papillæ*. The nerves of the organ proceed from the *inferior maxillary, glosso-pharyngeal* and *hypoglossal*. The first and second are connected with taste; the last with the motions of the organ. Some recent observers, however, consider the *glosso-pharyngeal* to be the special nerve of gustation. The arteries are the *lingual*, from the external carotid; and the *palatine* and *tonsillar*, from the labial. The veins are the *superficialis linguae, ramina, lingualis*, and *submental*. They open into those of the pharynx and larynx. Its lymphatic vessels pass into *ganglia*, situate at the edge of the *hyoglossi muscles*.

TONGUE, BLACK. A fever which prevailed in the Western States in the winter of 1842-3; probably typhoid fever. According to some, an epidemic erysipelas.

TONGUE, HERNIA OF THE, Glossocœle—t. Inflammation of the, *Glossitis*—t. Red, fever, see *Typhus*.

TONGUE SCRAPER, Cure-langue—t. Tie, see *Ankyloglossum*.

TON'IC, Ton'icus, (F.) Tonique. Same etymon as *Tone*. Relating to *tone*.

Tonic, Ton'icus, Tono'nicus, Antasthen'icus, (F.) Tonique, Fortifiant, in Therapeutics, means a medicine, which has the power of exciting slowly, and by insensible degrees, the organic actions of the different systems of the animal economy, and of augmenting their strength in a durable manner. Bitter, vegetable substances which are not associated with an acrid or narcotic principle; preparations of iron; cold water, &c., act as tonics. They are also called *Comfortant'ia, Confortati'va, Corroborant'ia, Confirmant'ia, Roborant'ia*, &c.

The following are the chief Tonics:—*Acidum Arseniosum, Acidum Nitricum, Acidum Sulphuricum, Argenti Nitras, Bismuthi Subnitrates, Carbo Ligni, Cascarella, Cinchona, Cinchonina, Columba, Cornus Florida, Cupri Acetas, Cupri Sulphas, Cuprum ammoniatum, Cusparia, Eupatorium perfoliatum, Ferri Præparata, Gentiana, Liriodendron, Lupulina, Piperina, Prunus Virginiana, Quassia, Quinia, Salix, Salicina, Serpentina, Zinci præparata; Aquæ minerales acidulæ et ferruginæ; Mental Tonics, (Confidence, Hope, Amusement.)*

TONIC POWER, Irritability.

TONIC SPASM, Spasmus ton'icus, Convulsio tonica, Conductio, Spas'ticus, Tonos, Ento'nia, Entasia, Enta'sia. A rigid contraction of muscles without relaxation, as in *trismus, tetanus*, &c. See *Tetanus* and *Spasmus*.

TONIC'ITY, Tonic'itas. Same etymon as

Tone. The faculty that determines the general tone of the solids. Excessive tonicities causes *er'ethism* or *crispness*; deficient tonicities, *atony* or *weakness*. Tonicity has been used in the same sense as irritability.

TONIQUE, Tonic.

TONKA BEAN, Diplexer odorata.

TONODES, Tensive.

TONOS, Tonic spasm—t. *Trismus, Trismus.*

TONOTICUS, Tonic.

TONSIL, Tonsill'a, Amygdala, Parist'h'mia, Parist'h'mium, Amphibranchium, Glan'dula colli, Tola, Tolles, Toles, To'l'ium, Tollum, Spongos, An'tias, Pl. Anti'ades, Almond of the ear, Almond of the throat, (F.) Amygdale, Tonsille, (Prov.) Ear-kecker. An ovoid collection of follicles, about half an inch in length, similar in shape to an almond, and situate, on each side, between the pillars of the *velum palati*. The inner surface, projecting into the isthmus of the throat, is covered by the mucous membrane, in which are the orifices of a dozen cells, whence a transparent, viscid mucus oozes on pressure. This mucus is destined to facilitate deglutition, by lubricating the isthmus faucium; and is forced out during the passage of the alimentary bolus.

TONSIL-GUILLOTINE, Amygdalotome.

TONSILLÆ PESTILENTES, Cynanche maligna.

TONSILLARUM GANGRÆNA, Cynanche maligna.

TONSILLE, Tonsil.

TONSILLIT'IC, Tonsillit'icus. Belonging or relating to the tonsils; as the '*tonsillitic branches of the glosso-pharyngeal nerve*.'

TONSILLITIS, Cynanche tonsillaris.

TONSILLOTOME, Amygdalotome.

TONSORIUS CULTER, Razor.

TOODLE, Tooth.

TOOTH, Sax. toð, Dens, Odous, (Prov.) Toode, (F.) Dent. The teeth, *Dentes, Mord'ices*, are small, hard bones, fixed in the alveoli of each jaw, and serving to lay hold of, and to cut, tear, and triturate alimentary substances. True teeth are found only in man, the mammalia, reptiles, and fish. The teeth have, in general, the shape of an irregular cone, the base of which is towards the interior of the mouth; and the apex, single or divided, towards the bottom of the alveoli. Each tooth has a part situate out of the alveolus, and covered by a vitriform substance. This part is called the *crown* or *coro'na*. It is bounded by a narrower portion, called the *cervix, (F.) collet*, which separates it from the portion in the alveolus—the *fang* or *root, (F.) racine*. In the adult, the teeth are 32 in number; 16 to each jaw: these consist of 4 *incisors*, 2 *canine*, 4 *bicuspidati*, and 6 *molares*. (See these various terms.) The teeth enclose a vascular and nervous pulp, in a central cavity, *Cav'itas Pulpæ, Cavum Dentis, Antrum dentale*. They are formed of a very hard, compact tissue; and the corona is covered by a vitreous substance called *Enam'el*. The incisor and canine teeth are developed by a single point of ossification; the lesser molares, by two; and the larger, by four or five. See *Dentition*. Researches by Retzius, Müller and others, have led them to believe that the ivory of the tooth or *Dentine, Ebur, Substantia os'sea* of Malpighi, proper tooth substance, bone of the tooth, osseous substance of tooth, tooth bone, consists of tubular fibres, which contain organic deposits of calcareous salts; and of intertubular substance, in which are observed corpuscles or osseous cells of the tooth, called, by Professor Owen, *calic'iferous cells*, which also contain earthy salts. At the part where the enamel terminates at the cervix of the tooth, the *crusta petro'ea, cementum* or *cortical substance*, commences in an extremely

thin stratum, passes down and gradually increases in thickness towards the extremity of the root, where it is generally the thickest. It is said by Nasmyth to have been traced on the enamel, and Purkinje and Fränkel thought that it lined the *cavitas pulpæ*. They gave it the name *Substantia ostioides*. It is the *material tartarea* of Malpighi. It has been found, however, that the hard substance deposited on the inner surface of the dentine, at the age of twenty years or later, and which encroaches on the cavity, has a distinct structure from the cement, and resembles in part dentine, and in part bone. It is the *horny substance* of Blumenbach, the *osteodentine* of Owen, and the *secondary dentine* of Tomes. In the compound teeth of the herbivorous animal, a deposition is found on the surface of the enamel, which has been called *cementum*. Müller, however, regards it to be a deposit from the salts and the saliva, and to be essentially the same as what is called tartar in the human subject.

TOOTH-ACH, Odontalgia.—t. Bone, see Tooth — t. Bush, *Xanthoxylum fraxineum* — t. Tree, *Aralia spinosa*, *Xanthoxylum clava Herculis*.

TOOTH EDGE, *Agacement des dents*, *Hæmodia* — t. Paste, *Dentifricium* — t. Powder, *Dentifrice* — t. Rash, *Strophulus* — t. Shell, dog-like, *Dentalium*.

TOPHAŒCEOUS, *Tophæceus*, (F.) *Tophacé*; from *tophus*. Relating or belonging to tophus, — as 'a *tophæceous* concretion.'

TOPHACEOUS CONCRETION, *Tophus*.

TOPHUS, *τοφος*, *Tofus*, 'a porous or friable stone.' *Osteo'tophus*, *Tophæceus* Concretion. A collection of hard, calcareous matter, which forms, particularly, at the surface of joints affected with gout; (see *Calculi*, *arthritic*), and occasionally, in the interior of organs, around the teeth, &c. It likewise meant gravel, and calchala.

TOPIC, *Topical*, *Topicus*. From *τοπος*, *locus*, 'a place.' A local application; (F.) *Topique*. See Local.

TOPICUS, Local.

TOPINARIA, Talpa.

TOPIQUE, Topic.

TÖPLITZ or TEPLITZ, MINERAL WATERS OF. Töplitz is a town of Bohemia, celebrated for its numerous thermal springs. The water contains chloride of sodium; carbonate, and sulphate of soda; and carbonates of lime and iron, &c. Temp. 114° to 122° Fahr.

TOPOGRAPHICAL ANATOMY, see Anatomy.

TOQUE, *Scutellaria galeuriculata*.

TORCULAR, *Tourniquet*. A press for wine, oil, or cider. From *torqueo*, 'I twist.'

TORCULAR HEROPH'ILI, *Press of Heroph'ilus*, *Lechene'ion*, *Lenos*, *Lenum*, *Fourth Sinus*, (F.) *Confluent des Sinus*, *Pressoir d'Hérophile*. A smooth and polished cavity, of irregular shape, in which several sinuses of the dura mater meet. It is situate before the internal occipital protuberance, at the union of the three great duplicatures of the dura mater. It has six openings into it; — one above, that of the superior longitudinal sinus; two below, answering to the occipital sinuses; one anterior, belonging to the straight sinus; and two lateral, and very broad, which lead into the lateral sinuses; and convey from the cavity the blood that has been poured into it by the others.

TORCULUM, *Tourniquet*.

TORDYL'IUM, *T. Officina'le*, *Ses'eli*, *Cret'icum* seu *Montanum*, (F.) *Séseli de Crète*. Family, Umbelliferae. Sex. Syst. Pentandria Digynia. The seeds and roots of this Southern European plant are considered carminative and diuretic. They enter into the theriaca.

TOR'NIA ASIATICA, *Cela-dolo*. A small Malabar plant, the juice of whose leaves, sweetened with sugar, is said to cure gonorrhœa.

TORMENTIL, *Geranium maculatum*, *Tormentilla*.

TORMENTIL'LA, *T. erecta* seu *sylvestris* seu *officina'lis* seu *tuberosa*, *Fraga'ria tormentil'la officina'lis*, *Potentilla tormentilla*, *Heptaphyll'um*, *Consolida rubra*, Common *Tormentil*, *Upright Septfoil*, (F.) *Tormentille*. Ord. Rosaceae. Sex. Syst. Icosandria Polygynia. The root *Tormentilla* (Ph. U. S.) has a slightly aromatic odour; and austere, styptic taste. It is astringent. Dose, of the powder, gr. x to ʒi.

TORMENTUM, Ileus, Intussusceptio.

TOR'MINA, *Strophos*, *Calial'gia*, *Encæst'al'gia*, *Anile'ma*, *Anile'sia*, *Gripes*, *Multigrubs*, *Fretting*, (F.) *Tranchées*. From *torqueo*, 'I twist.' Acute colicky pains. Dysentery.

TORMINA CELSI, Dysentery — t. *Parienturium*, Pains, labour — t. *Post partum*, see Pains, labour.

TORNACULUM, *Tourniquet*.

TOROSITAS, *Corpulent'ia carnos'a*, *Hab'itus athlet'icus*. Muscular strength; from *torosus*, 'brawny,' — *torus*, 'brawn.'

TOROSUS, Muscular.

TORPEDO, Narcosis.

TORPEFACTIO UNIVERSALIS, *Holonan-cosis*.

TORPID, *Torpidus*, (F.) *Engourdi*; from *torpeo*, 'I am benumbed.' Numb. Incapable of exertion and of feeling. Dull, stupid, inactive.

TORPIDITAS, *Torpor*.

TORPIDUS, *Narcotized*, *Torpid*.

TORPOR, *Torpid'itas*, *Noth'rites*, *Parap'sis experts*, (F.) *Engourdissement*, from *torpeo*, 'I am benumbed.' State of a part which becomes numb, and almost incapable of motion. *Torpor*, *Torpid'ity*, *Torpid'ness*, *Torpid'itude* are also applied to a debilitated, sluggish condition of a part, or of the whole of the body; *Aër'gia*, (F.) *Accablement*.

TORPOR, Narcosis — t. of the Colon, see Colon, torpor of the — t. *Intestinorum*, Constipation — t. *Soporificus*, *Nocar*.

TORQUAY. A town in Devonshire, England, which, on account of the mildness of its climate, and its sheltered situation from the bleaker winds, is a frequent retreat for the phthisical invalid.

TOR'SION, *Tor'sio*. From *torquere*, *torum*, 'to twist.' The act of twisting. Torsion of the extremities of bleeding vessels — *Angios'trophè* — until the blood ceases to flow, is sometimes used to arrest hemorrhage.

TORSK, see *Oleum jecoris aselli*.

TORTELE, *Erysimum*.

TORTICOL'IS, *Caput seu Collum obsti'pum*, *Obsti'pa cervix*, *Obsti'p'itas cap'itis* seu *Colli*, *Trachel'agra*, *Enta'sia* *Loxia*, *Lox'ia*, *Cephalox'ia*, *Ancylode'rè*, *Ancylode'ris*, *Rhæbra'nia*, *Auchenorrhæ'ma*, *Rheumatism'us colli* seu *cervi'cis*, *Stiffneck*, *Wryneck*, (Prov.) *Slab*; from *torquere*, *torum*, 'to twist,' and *collum*, 'the neck.' A variety of rheumatism, seated in the muscles of the neck, which prevents the motion of the head, and causes the patient to hold it inclined to the side affected. It is, commonly, of short duration; usually disappearing in a few days. Rubefacients are the best topical applications. The term is also applied to permanent contraction of the muscles of the neck, which causes the head to be held to one side. The remedy is the division of the contracted muscles.

TORTOISE, FRESHWATER, GREAT, *Hydraspis expansa*.

TORTUE, *Testudo*.

TORTURA ORIS, Canine laugh, *Neuralgia faciei* — t. Oris paralytica *Linnaei*, Paralysis. Bell's.

TOR'ULA CEREVISIÆ, *Saccharom'ycetes* seu

Mycoderma cerevisia, *Cryptococcus fermentum* seu *cerevisia*, (F.) *Champignon du ferment*, *Yeast plant*. One of the simplest forms of vegetation, capable, under favourable circumstances, of vegetating rapidly, and assisting in producing the fermentation of saccharine substances. The plant has been found in vomited fluids, and in feces.

TORULUS GLANDIS, Crown of the glands.

TORUS, Muscle — *t. Manûs*, Metacarpus.

TOTA BONA, *Chenopodium bonus Henricus*.

TOTSU KASO, *Sphæria Sinensis*.

TOTTY, Vertiginous.

TOTUM CARNOSUM, Pancreas.

TOUCH, *Tactus*, *Tactio*, *Hapê*, *Hapsis*, *Tactio*, *Palpation*, (F.) *Toucher*. One of the five senses, and that which makes known to us the palpable qualities of bodies, such as their consistence, quality, &c. It is seated, particularly, in the hands; and differs from tact in being active. See *Esaphe*, and *Palpation*.

'To touch the mouth with mercury' is to produce the constitutional effects of that agent, as evinced by the falling away of the gums, &c.

TOUCH-ME-NOT, *Impatiens fulva*.

TOUCH, ROYAL. It was at one time almost universally believed, in England, that the royal touch would remove *Serofula* or *King's Evil* (as it was thence called). This superstition is now entirely exploded. Edward, the Confessor, was the first English king who touched for the Evil.

TOUCHER, *Esaphe*, *Touch* — *t. Vaginal*, *Esaphe*.

TOUCHWOOD, *Boletus igniarius*.

TOUR DE MAÎTRE, *Coup de Maître*.

TOURBILLON VASCULAIRE, *Vasa vasorum*.

TOURNESOL, *Heliotropium Europæum*.

TOURNOIE, *Paronychia*.

TOURNIQUET, *Torcular*, *Torculum*, *Tornaculum*, *Prælium*, *Prælium seu Prelium arterialem*, *Verticillum*, *Fascia tortilis*, from (F.) *tourner*, 'to turn.' A surgical instrument for stopping the course of blood in the limbs, by exerting a strong compression upon the principal artery. It was invented by a French surgeon, Morel, in the middle of the 17th century; and was modified subsequently, by Nuck, Verduc, Monro, Petit, and Louis. It is used to suspend, for a time, the circulation in a limb, during great operations; to arrest arterial hemorrhage in cases of wounds; to compress certain aneurismal and other tumours, &c.

TOUS-LES-MOIS, *Amylum cannaecum*, *Canna Starch*. A fecula obtained from the root of *Canna edulis*. It resembles potato starch, and may be substituted for arrow-root. It is obtained from St. Kitts.

TOUT, *Nates*.

TOUTE BONNE, *Salvia sclarea*.

TOUTE ÉPICE, see *Myrtus pimenta* — *t. Saine*, *Sanicula*.

TOUX, *Tussis* — *t. Bleue*, *Pertussis* — *t. Bronchique*, *Bronchial cough*, *Cough*, *tubal* — *t. Caverneuse*, see *Cavernous respiration* — *t. Convulsive*, *Pertussis* — *t. Quinteuse*, *Pertussis* — *t. Tubaire*, *Bronchial cough*, *Cough*, *tubal*.

TOXICÆMIA, *Toxicohæmia*.

TOXICOL, *Poisonous*.

TOXICATIO, *Poisoning*.

TOXICODENDRON, *Rhus toxicodendron*.

TOXICOHÆMIA, *Toxicæmia*, from *toxicum*, 'a poison,' and *αἷμα*, 'blood.' *Poisoning of the blood*. — *Piorry*.

TOXICOL'OGY, *Toxicolog'ia*, from *toxicum*, (itself from *τοξον*, 'a bow,') 'a poison,' and *λογος*, 'a description.' A treatise on poisons.

TOXICOSES, (G.) *Toxikosen*, from *toxi-*

cum, 'a poison.' A family of diseases, according to the classification of Fuchs, caused by the reception of poisons into the system.

TOXICUM, *Poison*, *Venom*.

TOXIQUE, *Poisonous*.

TRABECULA CEREBRI, *Corpus callosum* — *t. Cordis*, *Columnæ carneæ*.

TRABECULÆ CORPORUM CAVERNOSORUM, see *Cavernous bodies*.

TRABECULÆ SPLENIS, see *Spleen* — *t. Willisii*, *Chordæ Willisii*.

TRABES CORDIS, *Columnæ carneæ*.

TRABS, *Penis* — *t. Cerebri*, *Corpus callosum* — *t. Medullaris*, *Corpus callosum*.

TRACE, *PRIMITIVE*, *Groove*, *primitive*.

TRACHEA, (generally, however, accented *Trachea*), *Trachæa* seu *Trachei'a arte'ria*, *Bronchus*, *Aspera arteria*, *Fistula spiritalis* seu *pulmonalis*, *Canna*, *Syring'a*, *Syrinx*; the *Windpipe*, *Wosen*, *Weasand*, *Weesel*, *Throttle*, (Sc.) *Throppill*, (Prov.) *Keekcorn*, *Kecker*, *Thropple*, (F.) *Trachée artère*; from *τραχῦς*, 'rough,' and *ἀρτηρία*, 'artery,' composed of *ἀν*, 'air,' and *ῥησις*, 'to keep.' The trachea is a cylindrical, fibro-cartilaginous and membranous tube, flattened behind, and situate on the median line, before the vertebral column, from the inferior part of the larynx, as far as the third dorsal vertebra. There it divides into two branches, the *bronchia*, which separate to go to each lung. The trachea, the function of which is to convey air to the lungs during respiration, is composed, 1. Of cartilaginous rings, *An'uli cartilagin'e'i*, *Segmen'ta cartilagin'ea*, *Orbes cartilagin'o'si*, incomplete behind; situate one above the other, and kept in situ by a fibrous membrane. 2. Of a mucous membrane, which lines it internally, and contains numerous mucous follicles. 3. Of transverse muscular fibres, not well-marked, which occupy its posterior surface. 4. Of vessels, which proceed from the superior and inferior thyroideal, and 5. Of nerves, that are given off by the pneumogastric and the cervical ganglia.

The breathing tubes of insects are termed *Tracheæ*.

TRACHEA ARTERIA, *Trachea*.

TRACHEÆ, see *Trachea*.

TRACHEAL, *Trachealis*. Appertaining to the trachea. An epithet applied to respiration as heard through the stethoscope opposite the trachea, larynx, and root of the bronchia; the air appearing as if sucked in from the cylinder during inspiration, and expelled during expiration.

TRACHEAL GLANDS. Mucous follicles on the posterior surface of the trachea. They are small, flattened ovoid bodies.

TRACHEALIS. The fibres of unstriated muscle, which pass from one portion of each of the tracheal cartilages to the other, at the posterior part of the trachea, have been so called by some anatomists. When they contract, they approximate the extremities of the cartilage; diminish the calibre of the trachea; and thus increase the velocity of the air forced through it, as in coughing, and facilitate the removal of any irritating substance in contact with the lining membrane.

TRACHÉE ARTÈRE, *Trachea*.

TRACHEITIS, *Cynanche trachealis*.

TRACHELAGRA, *Torticollis*. Also, *gout in the neck*.

TRACHELIAN, *Cervical*.

TRACHELISMUS, from *τραχηλος*, 'the neck.'

A term proposed by Dr. Marshall Hall, for a fancied spasmodic action of the muscles of the neck, which he esteems to be the cause of many morbid phenomena, by inducing compression of the veins of the neck and an impeded return of blood from the head.

TRACHELITIS, Cynanche trachealis.
TRACHELIUM AMERICANUM, Lobelia cardinalis.

TRACHÉLO-ATLOÏDO-OCCIPITAL, Obliquus superior capitis—*t. Atloïdo-basilaire*, Rectus capitis lateralis—*t. Basilaire*, Rectus capitis internus major—*t. Basilaire, petit*, Rectus capitis internus minor.

TRACHÉLO-DIAPHRAGMATIC, *Trachelo-diaphragmaticus*. The fourth pair of cervical nerves, whence the phrenic nerve chiefly arises.—Chaussier.

TRACHÉLO-MASTOÏDIEN, Complexus minor—*t. Occipital*, Complexus—*t. Scapulaire*, Levator scapulae.

TRACHELOS, Collum.

TRACHE'O-BRONCHITIS. Inflammation of the trachea and bronchia.

TRACHEOCELE, Bronchocele.

TRACHEOPHONY, Laryngophony.

TRACHEOPHTHISIS, see Pthisis laryngea.

TRACHEOPHYMA, Bronchocele.

TRACHEORRHAG'IA, *Hæmoptœ* seu *Hæmoptysis laryngea et trachealis*, from *τραχεια*, 'the trachea,' and *ρηνυμι*, 'to break forth.' Hæmorrhage from the trachea.

TRACHEOSTENOSIS, from *τραχεια*, 'the trachea,' and *στενωσις*, 'contraction.' Contraction or narrowness of the trachea.

TRACHEOTOME, from *trachea*, and *τομή*, 'incision.' An instrument of the trocar kind, for opening the trachea.

TRACHEOTOMY, see Bronchotomy.

TRACHITIS, Cynanche trachealis.

TRACHOMA, from *τραχυς*, 'rough.' *Aspre'do*, *Aspritu'do*, *Pladar'otes*, *Asperitu'do*, *Scabrit'ies*, *Gran'ular conjunct'iva*, *Gran'ular eyelid*, (F.) *Asperité des Paupières*. A roughness of the inner surface of the eyelids. A variety of ophthalmia, of which three kinds have been designated:—*T. scabulo'sum*, from sand, &c., getting between the eyelids; *T. carunculo'sum*, *Pladar'otes*, *Pladar'osis*, *Morum palpebræ internæ*, from fleshy excrescences; and *T. herpet'icum*, from hard pustules on the inner surface of the eyelids. This last is, also, called *Fico'sis*, and *Palpebra fico'sa*.

TRACT, OPTIC, see Optic nerves—*t. Respiratory*, see Respiratory tract.

TRACTATIO MANUARIA, Surgery.

TRACTORATION, Perkinism, see Tractors, metallic.

TRACTORS, METALLIC. Metallic agents, introduced by Dr. Elisha Perkins, of Connecticut, about the end of the last century, to cure diseases. The parts were rubbed with these tractors, and hundreds of thousands of cases were reported to have been cured by the *Tractor'ation*. The affections, in which they were chiefly used, were local inflammations, and pains in different parts. The good effects were, doubtless, owing to the influence exerted by the mind over the body. The bubble did not exist long. See Perkinism.

TRACTUS INTESTINORUM, Intestinal Tube—*t. Opticus*, see Optic nerves—*t. Respiratorius*, Respiratory tract.

TRACTUS SPIRALIS FORAMINULENTUS. Minute openings at the base of the modiolus, arranged in a spiral manner, which transmit the filaments of the cochlear nerve. The central canal of the Tractus is longer than the rest, and is called *Tubulus centralis modioloti*.

TRADESCANTIA VIRGINICA, *Spiderwort*; indigenous; Order, Commelynacæ; flowering from May to August. The roots are demulcent.

TRAGACAN'THA, from *τραγος*, 'a goat,' and *ακανθα*, 'a thorn,' owing to its hirsute boughs.

Gummi Tragacantha seu *Astragal'is Tragacanthæ*, *Gum Tragacanthæ*, *Gum Dragon*. The concrete juice of the *Astragalus Tragacantha* or *Astragalus Verus*, a native of Persia. Ord. Leguminosæ. *Sec. Syst.* Diadelphia Decandria. (F.) *Gomme Adragant*, *Adragant*, *Adragante*, *Adraganthe*. This gum is inodorous; nearly insipid; of a whitish colour; semi-transparent, and striated. It is in thin, vermiform pieces; and does not form a smooth, uniform mucilage with water. It is demulcent, but is rarely used alone.

TRAGÆA AROMATICA, Pulvis cinnamomi compositus.

TRAG'ICUS, *Tragien* (Ch.), from *Tragus*. A small, flat, triangular muscle which almost wholly covers the outer surface of the tragus, arising from its base, and terminating at the top of the eminence.

TRAGIEN, Tragicus.

TRAGOMASCHALIA, *Tragomas'chalæ*, from *τραγος*, 'a goat,' and *μασχάλν*, 'axilla.' The strong unpleasant odour of the armpits.

TRAGOPHONIA, Egophony.

TRAGOPO'GON, *Barba hirci*; from *τραγος*, 'a goat,' and *πογων*, 'beard.' A genus of plants; Ord. Cichoracæ. *Sec. Syst.* Syngenesia Polygamia Æqualis. (F.) *Barbe de Bouc*; of which the roots of the *Tragopogon porrifolium*, *Salsifi*, *Oyster plant*, *Oyster-root*, (F.) *Salsifis* ou *Cer-cifis*, and the young shoots of the *T. pratensis*, *Meadow Salsifi*, *Common Goat's Beard*, (F.) *Salsifs des prés*, are eaten as food. The root of the latter has been used as a diuretic.

TRAGOSELINUM, *Pimpinella saxifraga*—*t. Angelica*, *Ligusticum podagraria*—*t. Magnum*, *Pimpinella magna*—*t. Majus*, *Pimpinella magna*—*t. Saxifraga*, *Pimpinella magna*.

TRAGUS; *τραγος*, 'a goat,' from its being furnished, in some persons, with hair, like the beard of a goat. *Hircus*, *Hirquus*. A small, flattened, triangular nipple, situate before the meatus auditorius externus, which it appears to conceal. It is covered with hair in old people, whence its name. Also, the peculiar goat-like smell of the axillæ.

TRAGUS MATTHIOLI, *Salsola kali*.

TRAINASSE, Polygonum aviculare.

TRAITEMENT, Curatio—*t. Palliatif*, see Palliation.

TRAMIS, Perinæum.

TRANCE, Catalepsy, Ecstasis.

TRANCHÉES, Tormina—*t. Utérines*, Pains, after.

TRANS, in composition, 'through, over, or beyond.'

TRANSFIGURATIO, Transformation.

TRANSFORMA'TION, *Transforma'tio*, *Transfigura'tio*, *Transmuta'tio*, *Metab'olê*, from *trans*, and *forma*, 'form.' Change of form.

TRANSFORMATIONS DE TISSU, (F.) The French pathologists designate thus the change that occurs in an organ, the tissue of which becomes similar to that of another organ. The chief transformations are cartilaginous, osseous, and adipous. See Tissue.

TRANSFUSION, *Transfusio*, *Metach'ysis*, from *transfundere* (*trans*, and *fundere*, *fundum*, 'to pour') 'to pour from one vessel into another.'

TRANSFUSION OF BLOOD, *Transfusio Sanguinis*, *Cura Medea'na*, *Methamachym'ia*, *Hæmatometach'ysis*, *Hæmometach'ysis*, *Diach'ysis*, *Transplanta'tio med'ica nova*, *Chirur'gia transfusio'ria*, *Exotichæmato'sis*. The act of passing the blood of one animal into the veins of another; an operation which was formerly used to fulfil various therapeutical intentions. It has been revived in cases of uterine hemorrhage; and, apparently, with some success.

The operation can only be performed safely

on animals having like kinds of blood. See Globule.

TRANSLATION, Metastasis.

TRANSMISSIBILITY, see Sensibility.

TRANSMUTATIO, Transmutation.

TRANSPIRATIO, Perspiration—t. Unilateral, Hemidiaphoresis.

TRANSPIRATOIRE, Perspiratory.

TRANSPANTA'TIO, (trans, and planta,) Metaphyte'a. A pretended method of curing diseases, devised by Paracelsus—by making them pass from one individual to another, either animal or vegetable.

TRANSPANTATIO, Morioplastice—t. Medica nova, Transfusion.

TRANSPANTA'TIO DENTIS, *Insit'io dentis*. The act of transplanting a tooth from one person to another.

TRANSPORT, Delirium.

TRANSPPOSITIO, Metathesis.

TRANSPPOSIT'ION, *Transposit'io*, from trans, 'over,' and ponere, positum, 'to put.' *Metath'esis*. Change of situation.

TRANSPPOSITION OF THE VIS'CERA, *Polype'ria promisc'ua translati'va*, *Intus inver'sus*, is a congenital vice of conformation, which consists in the viscera being found out of the situations they ordinarily occupy: the heart, for example, being on the right side; the liver on the left, &c.

TRANSUDATIO, Diapedesis.

TRANSUDA'TION, *Transuda'tio*, (F.) *Transudation*, from trans, 'through,' and sudare, sudatum, 'to sweat.' The passage of a fluid through the tissue of any organ, which may collect in small drops on the opposite surface, or evaporate from it.

TRANSUDATION, Exosmose.

TRANSVERSAIRE ÉPINEUX, *Transversalis dorsii*—t. *Épineux du cou*, *Semispinalis colli*—t. *Grêle*, see *Sacro-lumbalis*—t. *Épineux du dos*, *Semi-spinatus dorsii*.

TRANSVERSAL DE LA CONQUE, *Transversalis auris*—t. *du Nez*, *Compressor nasi*—t. *des Orteils*, *Transversus pedis*.

TRANSVERSALIS, Transverse.

TRANSVERSA' LIS ABDOM'INIS, (F.) *Lombo-abdominal* (Ch.), *Transverse de l'Abdomen*. This muscle is seated, deeply, on the lateral parts of the abdomen. It is flat, thin, and broader before than behind. It is attached, above, to the cartilage of the last true rib, and to those of every false rib; and, below, to the inner lip of the crista of the ilium; and to the two outer thirds of the crural arch, and to the upper part of the pubis. Its fibres, moreover, are inserted, behind, by means of broad aponeuroses, into the top of the transverse and spinous processes of the first four lumbar vertebrae; and, before, into the linea alba, and the ensiform cartilage. This muscle, all of whose fibres have a transverse direction, constricts the belly, and diminishes the base of the chest, by drawing inwards the ribs to which it is attached.

TRANSVERSALIS ANTICUS PRIMUS, *Rectus capitis lateralis*—t. *Collateralis colli*, see *Sacro-lumbalis*.

TRANSVERSA' LIS COLLI, *Transversalis Major colli*, *Portion of the Sacro-spinal* (Ch.), (F.) *Transversaire*. A muscle, seated at the posterior and lateral part of the neck and upper part of the back. It is attached to the transverse processes of the five or six lowest vertebrae of the neck; and to those of the four or five first dorsal. It extends the vertebrae of the neck, and inclines them to its side.

TRANSVERSALIS COLLI, *Semi-spinalis colli*.

TRANSVERSALIS DORSI, *Transverso-spinalis*, *Multi'fidus Spinae*, *Lumbo-cervical portion of the*

Spinal (Ch.), (*Transverso-spinalis Lumbo'rurum*, *Transverso-spinalis Dorsi*, *Transverso-spinalis Colli*), *Musculus sacer*, *Spinales et transversales lumborum*, *Semi-spinalis internus* sive *Transverso-spinalis colli pars interna* (Winslow), *Lombodorsi-spinal*, &c., (F.) *Transversaire épineux*. This muscle, usually called, by English anatomists, *Multi'fidus spinæ*, and including all the *transverso-spinales*, is situate in the vertebral gutters, extending from the posterior part of the sacrum to the second cervical vertebra. It is thicker in the neck and loins than in the back and behind the sacrum. Its use is—to straighten the vertebral column, and to impress upon it a slight rotary motion.

TRANSVERSALIS NASI, *Compressor nasus*.

TRANSVERSARIUS, Transverse.

TRANSVERSE, *Transver'sus*, *Transversa'lis*, *Transversa'rius*, from trans, 'over,' and vertere, versum, 'to turn.' That which runs across; also, that which relates to the transverse processes of the vertebrae.

TRANSVERSE ARTERY OF THE FACE, arises from the temporal; passes transversely across the face, in front of the condyle of the lower jaw, and gives its branches to the different muscles of the cheek.

TRANSVERSE FISSURE, see Liver—t. *Fossa*, see Liver—t. *Ligament of the Atlas*, *Annular ligament*.

TRANSVERSE PERINEAL ARTERY, *Urethro-bulbar* (Ch.), is given off from the upper branch of the internal pudic; and passes inwards and forwards above the transversus perinaei muscles, as far as the bulb of the urethra, into which it dips, subdividing into numerous branches.

TRANSVERSE PROC'ESSES OF THE VER'TEBRÆ, *Diapoph'yses* of Owen, (F.) *Apophyses transverses des vertèbres*, are the bony eminences that jut out transversely and posteriorly from the sides of the vertebrae.

TRANSVERSE SUTURE, *Suti'ra transversa'lis*, runs across the face, and sinks into the orbit; joining the bones of the skull to those of the face, but with so many irregularities and interruptions, that it can scarcely be recognised as a suture.

TRANSVERSE DE L'ABDOMEN, *Transversalis abdominis*—t. *de l'Oricule*, *Transversus auris*—t. *du Périnée*, *Transversus perinaei*.

TRANSVERSO-SPINAL, *Semi-spinalis colli*, *Semi-spinalis dorsii*—t. *Spinalis*, *Transversalis dorsii*—t. *Spinalis colli*, *Semi-spinalis colli*—t. *Spinalis dorsii*, *Semi-spinalis dorsii*.

TRANSVER/SUS AURIS, *Transver'sus Auriculæ*, (F.) *Transverse de l'Oricule* (Ch.), *Transversal de la Conque*, is often wanting. It is formed of some transverse fibres, situate transversely, behind the projection of the helix which divides the concha into two portions.

TRANSVERSUS PEDIS, *Scandula'rius*, (F.) *Transversal des Orteils*, *Métatarso-sous-phalangien transversal du premier orteil* (Chaussier). A muscle, which arises by fleshy slips from the heads of the metatarsal bones of the three outer toes. Its tendon is inserted into the base of the first phalanx of the great toe, being blended with that of the adductor pollicis.

TRANSVERSUS PERINÆI, *Leva'tor Ani parvus*, (F.) *Ischio-périneal* (Ch.), *Transverse du Périnée*, is placed at the posterior part of the perinæum. It is thin, triangular, and situate transversely. Its external extremity is attached to the ramus and tuberosity of the ischium: its inner extremity is confounded with its fellow of the opposite side, and with the anterior part of the sphincter ani and posterior part of the bulbo-cavernosus. The use of this muscle is,—to compress the urethra, and to support the rectum and bladder. To a

fasciculus of this muscle, Santorini has given the name *Urethra Elevator* seu *Ejaculator*.

TRANSVERSUS PERINÆI ALTER, *Prostaticus inferior*. A small muscle which occasionally accompanies the last.

TRAPA NATANS, (F.) *Macre flottante*, Noix ou *Châtaigne d'Eau*. Ord. Onagraceæ. Ser. Syst. Tetrandria Monogynia. The plant which affords the *Nuces Aquaticæ*, *Tribuli aquatici*, *Caltropæ*. The nut is considered nutrient and demulcent, and is said to be useful in diarrhoea from abrasion of the bowels, and in calculus. A poultice is sometimes made of it to discuss hard and indolent tumours.

TRAPEZA, Mensa.

TRAPÈZE, Trapezium, Trapezius.

TRAPEZIUM OS, *τραπεζιον*, 'a trapezium.' So called from its shape. Os multangulum majus, (F.) Os trapèze. The first bone of the second row of the carpus. It is articulated, above, with the scaphoides; below, with the first bone of the metacarpus; within, with the trapezoides, and second metacarpal bone. Anteriorly, posteriorly, and externally, it gives attachment to ligaments.

TRAPEZIUS, *Cucullaris*, (F.) *Dorso-scapularis* (Ch.), *Trapèze*, *Capuchon*. A muscle, seated at the posterior part of the neck and shoulder, and at the upper part of the back. It has the shape of a trapezium; is attached, on the one hand, to the inner third of the upper curved line of the occipital bone; to the posterior cervical ligament; to the spinous process of the 7th cervical vertebra; and to those of all the dorsal vertebrae; and, on the other hand, to the spine of the scapula; the acromion and the outer third of the posterior margin of the clavicle. Its upper fibres are situate obliquely downwards and outwards; the transverse and inferior, obliquely outwards and inwards. This muscle elevates the shoulder; carries it backwards, or depresses it, according as its upper, middle, or lower fibres contract separately. It straightens the head, also, and inclines it to one side.

TRAPÉZOID LIGAMENT, *Ligamentum trapezoides*. The anterior part of the coraco-clavicular ligament. It is a very strong, fibrous fasciculus, which has the shape of a trapezium, and is situate obliquely, between the acromion process and clavicle.

TRAPEZOÏDES, OS, Os multangulum minus seu *pyramida le carpi*, from *τραπεζιον*, 'a trapezium,' and *ειδος*, 'shape or resemblance.' The second bone of the second row of the carpus,—so called from its shape. It is smaller than the trapezium, on the inside of which it is situate. It is articulated, above, with the os naviculare; below, with the second metacarpal bone; on the outside, with the trapezium; and internally, with the magnum. Anteriorly and posteriorly, it affords attachment to ligaments.

TRAUBENKUR, Grape cure.

TRAULISMUS, see Balbuties and Blæsitas.

TRAULOTES, see Blæsitas.

TRAUMA, Wound.

TRAUMATIC; *Traumaticus*, *Vulnerarius*, *Vulnerary*, (F.) *Traumatique*, from *τραυμα*, *traumas*, 'a wound.' Anything relating to a wound.

TRAUMATICA, Detergents, Vulneraries.

TRAUMATICINE; same etymon as Traumatic. A name given by Eulenberg to a solution of gutta percha in chloroform, which is applied externally in various chronic cutaneous diseases.

TRAUMATICUS, Traumatic, Vulnerary.

TRAUMATISM, *Traumaticus*. The condition of the organism occasioned by a grave wound.

TRAUMATO-COM'UM, from *τραυμα*, *traumas*, 'a wound,' and *κομω*, 'I take care of.' An hospital for the reception of the wounded.

TRAUMATOP'YRA, *Febris traumatica*; from *τραυμα*, *traumas*, 'a wound,' and *πυρ*, 'fire.' A wound fever, or fever consecutive to a wound.

TRAVAIL, Parturition.

TRAVAIL D'ENFANT, Parturition.

TRAVELLE'S JOY, Clematis vitalba.

TREACLE, Melasses, Theriaca—t. English, *Teucrium chamædrys*—t. Venice, Theriaca.

TREAD, Cicatrícula—t. of the Cock, Molecule.

TREATMENT, Curatio.

TREATMENT, DRY. One in which the patient is almost wholly, if not wholly, restricted from the alimentary use of fluids; a valuable *modus medendi* in inflammatory and hemorrhagic conditions.

TREE, ELK, *Andromeda arborea*—t. of Life, *Thuya occidentalis*—t. Sour, *Andromeda arborea*.

TRÈFLE DE CASTOR, *Menyanthes trifoliata*—t. D'Eau, *Menyanthes trifoliata*—t. des Marais, *Menyanthes trifoliata*.

TREFOIL, *Hepatica triloba*—t. Bean, stinking, *Anagris*—t. Marsh, *Menyanthes verna*—t. Sour, *Oxalis acetosella*—t. Shrubby, *Ptelea trifoliata*—t. Water, *Menyanthes trifoliata*.

TREMA, Foramen.

TREMBLEMENT, Tremor—t. Métallique, see Tremor.

TREMBLES, Milk-sickness.

TREMBLING, Tremor—t. Fevers, Intermittent fever.

TREMELLA AURICULA JUDÆ, *Peziza auricula*.

TREMOR, *Trepida'tio*, *Syn'clonus Tremor*, *Tromo-spasmus*, *Tromus*, *Trembling*, (F.) *Tremblement*, from *tremere*, 'to tremble.' Involuntary agitation of the body, or of some part of it, without any obstacle to voluntary motion. It depends, generally, upon debility of the muscular, or of the nervous system; and, hence, is observed in convalescence, and in typhoid affections. It occurs, also, in old people; in hard drinkers; workers in mercury, lead, &c.: in the last case, it is called by the French, *Tremblement métallique*, *Tremor metallicus*. Tremor seems to resemble paralysis more than convulsion. See Paralysis agitans and Pavitatio.

TREMOR CORDIS, Hippus, Palpitation—t. Metallicus, see Tremor—t. Potatorium, see Delirium tremens—t. Purring, *Frémissement catinaire*.

TREPAN, *Trep'anum*, *Trip'anon*, *τρειπανον*, *Terebel'ta*, *Pereter'ium*, *Peretor'ium*, *Serra versatilis*, *Chæn'icis*, *Chænic'ion*, *Ter'ebra*, *Ter'etron*, *Ter'etrium*, *Ter'ibulum*, (F.) *Trep'an*, from *τρειπαιω*, 'I perforate.' A surgical instrument resembling a winble, and worked in the same manner. It is used for removing portions of bone, and particularly of the bones of the skull. The term *trep'an* is given more particularly to the part of the instrument that makes the perforation. The handle is so constructed as to receive different bits,—as the crown, the *Trep'an perforatif* or *Perite'ri'on*, the *T. exfoliatif* or *Desquamato'rium trep'anum*, and the *T. abaptis'ta*.

In Great Britain and the United States, this instrument has been superseded by the trephine, which is more manageable.

TREPANATIO, Trepanning.

TRÉPANATION, Trepanning.

TREPAN'ING, *Terebra'tio*, *Anatre'sis*, *Cephalotripe'sis*, *Trepanda'tio*, *Trype'sis*, *Treis*, *Pris'sis*, *Prismus*, *Forda'tio*, (F.) *Trépanation*. Same etymon as Trepan. The methodical application of the trepan.

TREPANUM, Trepan.

TREPANUM, Trepan.

TREPHINE. The instrument which has replaced the trepan in some countries. It consists of a simple, cylindrical saw; with a handle placed

transversely like that of a gimlet, and from the centre of the circle, described by the teeth of the saw, a sharp little perforator projects, called the *Centre-pin*. The centre-pin is capable of being removed, at the surgeon's option, by means of a key for the purpose. It is used to fix the instrument until the teeth of the saw have made a circular groove, sufficiently deep for it to work steadily. The pin must then be removed. Sometimes the pin is made to slide up and down, and to be fixed in any position, by means of a screw.

TREPHOUSA, Nurse.

TREPIDATIO, Tremor—t. Cordis, Cardiotromus.

TRÉPIED VITAL, Tripod, vital.

TREPON'DO. A weight of three pounds.—Scribonius.

TRE'SIS, *τρῆσις*, from *τρεω*, 'I bore.' Properly, a perforation. A wound. A forcible solution of continuity in a soft part, commencing externally. A genus in the nosology of Good.

TRESIS, Trepanning, Wound—t. Causis, Burn—t. Punctura, Puncture—t. Vulus, Wound—t. Vulus laceratum, see Laceration—t. Vulus simplex, Cut.

TRESSAILLEMENT, Shuddering.

TRESSO'RIA, low Latin *Episi'on*, *ἐπισιόν*. A term formerly applied to the hair covering the mons veneris, the absence of which has, by the vulgar, been esteemed a matter of reproach.

TRI, *τρι*, in composition, 'three.'

TRIACLE, Melasses.

TRIANGLE MÉDULLAIRE, Fornix.

TRIANGLE, VESICAL, see Urinary Bladder.

TRIANGULAR, *Trigo'nos*, *Triangula'ris*, from *tri*, and *angulus*, 'an angle.' That which has three angles.

TRIANGULAR OF INFRA-PUBIAN LIGAMENT, *Ligamentum triangula're* seu *infra-pubianum*, is a ligamentous fascia with short fibres, situate transversely beneath the symphysis pubis, which it strengthens. Also, a ligament extending from the coracoid to the acromial process of the scapula.

TRIANGULARIS, Depressor anguli oris—t. Coccygis, Coccygeus—t. Nasi, Compressor naris.

TRIANGULA'RIS STERNI, *Sterno-costal* (Ch.), *Pectora'lis inter'nus*. A muscle situate at the anterior and inner part of the chest, behind the cartilages of the ribs. It has the shape of a lengthened triangle, the base of which is directed downwards. It is attached to the posterior, lateral, and inferior part of the sternum, whence its fibres pass upwards and outwards, and terminate by as many distinct digitations at the cartilages of the 3d, 4th, 5th, and 6th true ribs. This muscle depresses the ribs and contributes to expiration.

TRIBAS, (F.) *Tribade*, from *τριβω*, 'I rub.' *Fricatrix*, *Frica'trix*, *Futu'trix*, *Mas'cula*, *Subagitat'rix*, *Subgita'trix*. A female whose clitoris is so large as to cause her to be regarded as a hermaphrodite. One who acts as a male with another female.

TRIBULI AQUATICI, see *Trapa natans*.

TRIBUTUM LUNARE, Menses—t. Menstruum, Menses.

TRICÆ, Plica—t. Incuborum, Plica—t. Scroforum, Plica.

TRICAGO, *Teucrium chamædrys*.

TRICAUDALIS, *Retrahens auris*.

TRICEPH'ALUS, from *tri*, and *κεφαλη*, 'head.' A monster with three heads.

TRICEPS, from *tri*, and *caput*, 'head.' A name given to muscles that have three fleshy bundles at one of their extremities.

TRICEPS ADDUC'TOR FEM'ORIS. Under this

appellation are comprehended three distinct muscles: *Adductor brevis*, *A. longus*, and *A. magnus*.

TRICEPS AURIS, *Retrahens auris*—t. *Brachialis*, *Triceps extensor cubiti*.

TRICEPS CRURIS, *Femora'lis*, (F.) *Trifémoro-rotulien* (Ch.), *Crural*, *Trifémoro-tibi-rotulien*, *Triceps crural* ou *de la cuisse*, is situate at the anterior, inner, and outer part of the thigh. It is formed of three bundles at its superior extremity, viz. 1. A middle fasciculus (the *cruralis* of most anatomists). 2. An external fasciculus (the *vastus externus*) and 3. An inner fasciculus (the *vastus internus*). It is attached, above, to the anterior, inner, and outer surfaces of the femur, and to the two edges of the linea aspera, from the base of the trochanter to within four fingers' breadth of the knee. Below, it terminates by a large tendon, which is attached to the base and edges of the patella, as well as to the inner and outer tuberosities of the tibia. The triceps extends the leg on the thigh, and conversely.

TRICEPS DE LA CUISSE, *Triceps cruris*—t. *Crural*, *Triceps cruris*.

TRICEPS EXTEN'SOR CU'BITI, *Triceps brachia'lis*, (F.) *Scapulo-huméro-olecrani* (Ch.), *Triceps brachial*. This muscle, which occupies all the posterior part of the os humeri, is described as two muscles by Douglas, and as three by Winslow. It arises by three heads. The first or long head,—long head of the *biceps externus* of Douglas, *Anconeus major* of Winslow,—arises from the anterior extremity of the inferior costa of the scapula, near its neck, and below the origin of the *teres minor*. The second head, or short head of the *biceps externus* of Douglas, *Anconeus externus* of Winslow, arises from the upper and outer part of the os humeri, at the base of the great tuberosity; and the third head,—*Brachialis externus* of Douglas, *Anconeus internus* of Winslow,—the shortest of the three, arises from the back part of the os humeri, behind the flat tendon of the *latissimus dorsi*. These three portions unite about the middle of the arm, so as to form one thick and powerful muscle, which is inserted at the upper part of the olecranon. It extends the forearm on the arm, and conversely.

TRICEPS MAGNUS, *Adductor magnus*—t. Minor, *Adductor longus femoris*—t. Secundus, *Adductor brevis*.

TRICHÆ INCUBORUM, Plica.

TRICHANGIA, Capillary vessels

TRICHANGIECTA'SIA, *Angiecta'sia*, *Trichocirsus*, from *τριχ*, *τριχος*, 'a hair,' *αγγειον*, 'a vessel,' and *εκτασις*, 'dilatation.' Morbid dilatation of the capillary vessels.

TRICHAUX'È, from *τριχ*, *τριχος*, 'hair,' and *αύξη*, 'increase.' Increase in the quantity and size of the hair, without change of texture.

TRICHERA ARVENSIS, *Scabiosa*.

TRICHIA, Entropion.

TRICHIA'ASIS, *Trichio'sis*, *Tricho'sis*, from *τριχ*, *τριχος*, 'hair.' *Morbus pila'ris*. This name has been given to several affections. 1. To a disease of the kidneys or bladder, in which filamentous substances, resembling hairs, are passed in the urine. This is also called *Pilimictio*. It must be regarded as a variety of *Catar'rhus vesicæ*. 2. To a painful swelling of the breasts, in child-bed women, when the milk is excreted with difficulty:—a disease attributed, by Aristotle, to a hair swallowed by accident and carried to the breasts through the circulation; and 3dly, To inversion of the eye-lashes. See Entropion.

TRICHIASIS LACTEA, *Infarctus mammae lacteus*.

TRICHILIA, Cathartics.

TRICHIL'IA EMET'ICA, *Elcaja*. A tree, Order, Meliaceæ, common on the mountains of Yemen

and in Senegal. The fruit is used, by the Arabs, as an emetic. The ripe seeds are formed with sesamum oil into an ointment for the itch.

TRICHILIA TRIFOLIA'TA, a bush of Curaçoa, a decoction of the roots of which is used to promote abortion.

TRICHINA SPIRALIS. Same etymon. A small species of entozoa discovered in the muscles of voluntary motion. Trichinae consist of very minute cysts, of an oblong figure, having considerable resemblance, in size and colour, to young pediculi, and are somewhat allied to the eels found in paste and vinegar. No symptoms have as yet been observed to be pathognomonic of their presence.

TRICHIOSIS, Trichiasis—t. Distrix, Distrix.

TRICHIS'MUS. Same etymon, *Capilla'tio*, *Fissu'ra pila'ria*. A capillary fracture, fissure, or crack.

TRICHIURIA, Trichocephalus.

TRICHOÉPHALE DE L'HOMME, Trichocephalus.

TRICHOCEPH'ALUS, *Trichoceph'alus dispar*; from *σπῆξ*, *τρίχος*, 'a hair,' and *κεφαλή*, 'head.' *Trichu'ris*, *T. vulgaris* seu *intestinalis*, *Ascaris trichu'ria*, *Trichiu'ria*, *Trichocephalus* seu *Mastigo'des hominis*, *Long thread-worm*, (F.) *Trichocephale de l'homme*. A worm, from an inch and a half to two inches long; the head acute; the body spirally involutioned in the male, almost straight in the female. The *trichocephalus dispar* generally inhabits the cæcum and colon, and is rarely met with in the small intestines. It gives rise to no peculiar symptoms, and requires the usual anthelmintic treatment.

TRICHOCIRSUS, Trichiogectasia.

TRICHOG'ENOUS, from *σπῆξ*, *τρίχος*, 'a hair,' and *γενῶναι*, 'I generate.' Hair-forming; as a 'trichogenous remedy,'—one that promotes the growth of hair.

TRICHO LABIS, Madisterium.

TRICHO LABIUM, Madisterium.

TRICHOLOGIA, Carphologia.

TRICHOMA, Capillamentum, Plica.

TRICHOMANES, Asplenium trichomanoides—t. Crenata, Asplenium trichomanoides.

TRICHOMAPHYTE, from *trichoma*, 'plica,' and *φυῶν*, 'a vegetable.' A cryptogamous growth, found in plica, and said, by some, to be its cause.

TRICHOMAPHYTON, Mycoderma plicæ.

TRICHOMATION, Capillamentum.

TRICHOMATOSE HAIR, Plica.

TRICHOM'ONAS, *Trichom'onad*; from *σπῆξ*, *τρίχος*, 'a hair,' and *μονάδ*. An animalcule found by M. Donnè in the mucus of the vagina, where cleanliness has not been attended to, and to which he has given the name *Trichom'onas vagina'lis*. By some it is considered to be nothing more than ciliated epithelium separated from the uterus.

TRICHOMYCES TONSURANS, Porrigio decalvans.

TRICHONOSIS, Porrigio—t. *Furfuracea*, Porrigio decalvans—t. *Plica*, *Plica*—t. *Poliosis*, *Poliosis*.

TRICHON'OSUS, from *σπῆξ*, *τρίχος*, 'hair,' and *ὄσος*, 'disease.' A disease of the hair.

TRICHOPHY'IA, *Trichophy'tica*, from *σπῆξ*, *τρίχος*, 'hair,' and *φυῶν*, 'to grow.' Remedies that promote the growth of the hair.

TRICHOPHYTON TONSURANS, see Porrigio decalvans.

TRICHOSAN'THES, AMA'RA, (*σπῆξ*, *τρίχος*, 'hair,' and *ἀνθή*, 'a flower.') Order, Cucurbitaceæ; a plant of India; the seeds are bitter and astringent, sometimes emetic. In Jamaica, it is considered to be a poison, and is used for killing rats.

TRICHOSANTHES, CORDA'TA. The root is used

by the natives of India as a substitute for Calamba root.

TRICHOSANTHES CUCUMER'INA. The fruit is considered in India to be anthelmintic.

TRICHOSANTHES INC'I'SA, *Gashed Trichosanthes*, of the forests of India. The fruit is esteemed to be poisonous. Pounded and intimately blended with cocoa nut oil, it is used in India as an application to offensive sores of the ears, and to ozena.

TRICHOSANTHES VILLO'SA, of Java. The fruit acts like colocynth.

TRICHOSCHISIS, Schizotrichia.

TRICHO'SIS, *Pila'rē malum*; from *σπῆξ*, *τρίχος*, 'hair.' Morbid organization or deficiency of hair. A genus in the system of Good.

TRICHOSIS, Entropion, Trichiasis—t. Area, Porrigio decalvans—t. Athrix, Alopecia—t. Distrix, Distrix—t. Furfuracea, Porrigio furfurans—t. Hirsuties, Hirsuties—t. Plica, Plica—t. Poliosis, Poliosis—t. Setosa, Hystriacis.

TRICHOTON, Scalp.

TRICHURIS, Trichocephalus.

TRICLISPERMA GRANDIFLORA, Polygala paucifolia.

TRICUS'PID, *Tricus'pis*, *Tricus'pidalis*, *Tricus'pidatus*, from *tri*, and *cuspid*, *cuspidis*, 'a point.' That which has three points.

TRICUSPID VALVES, *Valvula Tricus'pides* seu *triglo'chines* seu *triul'ca*, are three triangular, valvular duplicatures, formed by the inner membrane of the right cavities of the heart, around the orifice by which the auricle communicates with the ventricle. The elongated and pointed top of each of these valves is continuous with the chordæ tendineæ, which set out from the columnæ carneæ of the heart. These valves are depressed to allow the blood to flow from the auricle into the ventricle; and are raised to prevent its reflux into the auricle.

TRIENS, *Un'cia qua'tuor*. From *tri*, 'three.' The third part of a pound. Four ounces, Troy.—Galen.

TRIFACIAL, Trigemini.

TRIFÉMORO-ROTULIEN, *Triceps cruris*—t. *Tibi-rotulien*, *Triceps cruris*.

TRIFOLIUM ACETOSUM, *Oxalis acetosella*—t. *Aquaticum*, *Menyanthes trifoliata*—t. *Aureum*, *Hepatica triloba*—t. *Cervinum*, *Eupatorium cannabinum*—t. *Fibrinum*, *Menyanthes trifoliata*—t. *Hepaticum*, *Hepatica triloba*.

TRIFOLIUM MELILO'TUS OFFICINA' LIS, *T. offic'ina'le* seu *odora'tum* seu *caballi'num*; from *tri*, and *folium*, 'a leaf.' *Melilotus*, *Melilotus cit'rina*, *Lotus sylves'tris*, *Ser'tula Campan'a*, *Coro'na reg'ia*, *Common mel'ilot*, (F.) *Mélot*. Ord. Leguminosæ. See *Syst. Diadelpia Decandria*. The infusion or distilled water of this plant has been employed in cases of ophthalmia. It has also been used in fomentations, glisters, &c.

TRIFOLIUM OFFICINALE, *T. melilotus officinalis*—t. *Paludosum*, *Menyanthes trifoliata*—t. *Palustre*, *Menyanthes trifoliata*.

TRIGAS'TRIC, *Trigas'tricus*, *Triven'ter*; from *tri*, and *γαστρον*, 'belly.' That which has three bellies. An epithet for muscles so circumstanced.

TRIGEM'INI, *Tergem'ini*, from *tri*, and *gem'inus*, 'double,' 'threefold,' 'triplets.' *The fifth pair of nerves*, *Nervi divi'si seu gustato'rii*, *Nervus quintus seu tremellus seu mixtus seu synpathet'icus me'dius seu synpath'icus medius seu cran'gius seu innomin'atus*, *Par trigem'inum seu quintum nervorum cerebr'alium seu trium funiculo'rum*, *Trifac'ial*, (F.) *Nerv gustatif innomin'at*, *Nerv à trois cordes*, *Nerv trijumeau*. This nerve arises below the tuber annulare by one large posterior root from the corpora restiformia, and by two small anterior roots from the corpora olivaria, whence, pro-

ceeding forwards to the side of the sella Turcica, the filaments composing the large root form a ganglion—the Gasserian. Emerging from this ganglion, the filaments, of which the large root consists, form, with those constituting the small roots, and which had not passed through the ganglion, two flattened trunks, each of which is soon divided into three branches, in such a manner, however, that the *first*, or *ophthalmic branch*, and the *second* or *upper maxillary*, are composed exclusively of the filaments that arose from the corpora restiformia and passed through the Gasserian ganglion; whilst the *third* or *lower maxillary branch* is composed in part of those filaments, and in part of those that arose from the corpora olivaria, and passed on one side of the ganglion without going through it. The whole, therefore, of the first and second branches, and a part of the third are sensoriferous nerves, whilst the remaining part of the third—the *manducatory* or *masticatory*—is wholly motoriferous, and passes to the pterygoid and other muscles concerned in mastication.

TRIGONE CÉRÉBRAL, Fornix—*t. Vésical*, see Urinary Bladder.

TRIGONELLA FENUM, *T. Fenum Græcum*, diminutive of *trigona*, three sided—from the shape of its flower; *Fen'ugreek, Fe'nugreek, Fœnum Græcum, B'ucerus, B. Fœnum Græcum, Carphos, Ægoc'eras, (F.) Fœnugrec, Sântegrain, Ord. Leguminosæ. Sex. Syst. Diadelphia Decandria*. A native of the south of France. The seeds have a strong, disagreeable smell, and an unctuous, farinaceous taste; accompanied by a slight bitterness. They are chiefly used in poultices.

TRIGONIUM OLFACTORIUM, see Mamillary encephalon.

TRIGONOS, Triangular.

TRIGONUS VESICÆ, see Urinary bladder—

t. Lieutaudi, see Vesical bladder.

TRIJUMEAUX, Trigemini.

TRILLIUM LATIFOLIUM, from *trilix*, 'triple,'—all the parts being in threes. *Broad-leaf Bethroot, Bethroot, Rattlesnake root, Wake-robin, Coughroot, Indian Balm, Ground Lily, Jew's Harp, Indian Shamrock, Pariswort, True-love. Ord. Trilliaceæ. Sex. Syst. Hexandria Trigynia*. A plant peculiar to North America, which blossoms in the spring. The root is astringent.

There are many species of *Trillium*, all of which possess the same properties.

TRIOCEPHALIA, Aprosopia.

TRIORCHIS, from *tri*, and *orchis*, 'a testicle.' One who has three testicles.

• **TRIOSTOSPERMUM**, *Triosteum*.

TRIOSTEUM, *T. Perfoliatum* seu *majus*, from *tri*, and *osteon*, 'a bone,' in allusion to the three bony nuts; *Triosteospermum, Bastard Ipecacuan'ha, Fever Root, Feverwort, Tinker's weed, Horse ginseng or gentian, Ipecac., Wild Coffee, White ginseng, Gent'ian, White Gentian, Sweet Bitter. Indigenous; Ord. Caprifoliaceæ*. In very large doses, the bark of the root proves emetic. It is a good cathartic, in doses of twenty or thirty grains. It sometimes operates as a diuretic.

TRIOSTEUM AUGUSTIFOLIUM, *T. minus*, has similar virtues.

TRIPÉ DE ROCHE. Different lichens of the species *Gyrophora*, inhabiting the Arctic regions of America, bear this name. They are eaten as food by the hunters.

TRIPHARMACUM, τριφάρμακον. From *tri*, and *pharmakon*, 'a medicine.' A medicine composed of three ingredients.

TRIPLET, from *triplex* 'triple.' One of three

children born at the same accouchement or gestation. Also, relating to triplets—as a 'triplet conception.'

TRIPLET MONSTERS, *Monstra trip'lica* seu *trigem'ina*, are such as have parts of the body tripled. They are rare.

TRIPOD, VITAL, (F.) *Trépied vital*. The heart, lungs, and brain have been so called, as their united action is necessary for the maintenance of life.

TRIPSIS, Contusio, Trituration.

TRIPUDIATIO SPASTICA, Chorea.

TRIQUETRA OSSA, Wormiana ossa.

TRIQUETRUM, (OS.) Cuneiform (bone.)

TRISIS, Brygmus.

TRISMUS, from τριζω, 'I gnash.' *Entia'seu Tenos Triemus, Capis'trum, Masticatory spasm of the face, Locked Jaw, Spasmus seu Rigor seu Tet'anus maxill'æ inferio'ris, Gnathospasmus, Stomatospasmus, (F.) Mal de Machoire*. Spastic closure of the under jaw; a partial tetanus. Cullen makes two varieties—*T. nascentium* seu *neonatorum*, *Nineday fits, (F.) Sarrête*, attacking infants during two weeks from birth; and *T. traumaticus*, originating from a wound.

TRISMUS CAPISTRATUS, Brédissure—*t. Catarrhalis maxillaris, Neuralgia maxillaris*—*t. Clonicus, Neuralgia faciei*—*t. Cynicus*, see Canine laugh—*t. Diastrophe Sauvagesii, Paralytic, Bell's*—*t. Dolorificus, Neuralgia faciei*—*t. Maxillaris, Neuralgia faciei*—*t. Nascentium*, see *Trismus*—*t. Neonatorum*, see *Trismus*—*t. Sardonicus, Risus sardonicus*.

TRISPERMUM, from *tri*, and σπέρμα, 'a grain or seed.' A cataplasma, consisting of three seeds,—*cumin, bay, and smallage*. *Gorraus*.

TRISPLANCHNIA, Cholera.

TRISPLANCHNIC, *Trisplanch'nicus, (F.)* *Trisplanchnique*; from *tri*, and σπλαγχνον, 'viscus.' That which relates to three orders of viscera.

TRISPLANCHNIC NERVE, *Nervus trisplanch'nicus* seu *sympathet'icus magnus* seu *sympath'icus* seu *ganglion'icus* seu *vaso-moto'rius, (F.) Nerf sympathique ou grand sympathique, Great sympath'ic, Intercos'tal, Ganglion'ic or ver'tebral nerve, Nerve of organ'ic life*;—named by Bichat the *organic nervous system*, in opposition to the other nervous parts, called the *animal nervous system*. Gall terms it, the *nervous system of the automatic functions*. It has also been called the *ganglionic nervous system*. It is termed *Trisplanchie*, because it distributes its branches to the organs in the three great splanchnic cavities—the head, chest and abdomen. It is composed of a series of ganglia, united by intermediate branches, and passing along the side of the spine to the lowest part of the trunk; communicating, by a branch, with each of the thirty pairs of spinal nerves and several of the cerebral nerves; and detaching nerves from its several ganglia, which accompany the arteries, and are distributed, particularly, to the organs of involuntary functions. At its upper part it is concealed in the canalis caroticus and cavernous sinus, and is joined there by a branch of the 6th pair of nerves, and by a twig from the Vidian nerve of the 5th pair. In passing down by the side of the spine, it forms three ganglia in the neck:—the *superior, middle, (thyroid ganglion,) and inferior, (ver'tebral ganglion)*; 12 in the back—the *thoracic*; 5 in the loins—the *lumbar*; and three or four *sacral*. When it reaches the coccyx it unites with its fellow from the opposite side, forming a small ganglion—the *coccygeal, Gan'glion impar, az'ygous gan'glion*. Meckel divides it into two portions—1. The *central ganglions* or *central part*, composed of the numerous ganglionic plexuses in the abdomen; the centre of which

is the *semilunar* and *solar plexuses*. 2. The *external ganglions*, *Limitrophes* or *external part*, which comprise the series of ganglions situate at the sides of the spine, from the base of the cranium to the sacrum. The great sympathetic seems to form a distinct nervous system, destined for the organs of involuntary motion; for although communicating with both brain and spinal marrow, it does not seem to be immediately under the influence of either. Its special functions are not yet, however, well understood, and are the subject—as they have always been—of much interesting investigation.

TRISPLANCHNITIS, Cholera, (spasmodic.)

TRISSAGO, Teucrium chamædrys—t. Palustris, Teucrium scordium.

TRISTEMANIA, Melancholy.

TRISTERN'AL. Bêclard has given this name to the third bony portion of the sternum;—that which corresponds to the third intercostal space.

TRISTITIA, Athymia, Moror.

TRITEOPHYA AMERICANA, Fever, yellow.

TRITEOPHYES, *Tritæoph'ya*, from *τρίτατος*, 'tertian,' and *φωσ*, 'I become.' A fever, whose type approaches the tertian. Sauvages gave the name *Tritæophya Vratislavien'sis*, to an epidemic, which broke out in the Prussian army at Breslaw, in the middle of the last century.

TRITEUS, Tertian fever.

TRITICUM ÆSTIVUM, from *terere*, tritum, 'to thrash.' *T. cereal'è* seu *Ceuallos* seu *compactum* seu *erina'ceum* seu *Gaertnerianum* seu *Linnæ'um* seu *pruin'osum*, *Summer wheat*; and *TRITICUM HYBERNUM* seu *glau'cum* seu *Muticum* seu *Sic'ulum*, *Winter wheat*, (F.) *Froment*. Ord. Gramineæ. *Sex. Syst.* Triandria Digynia. From these plants our wheaten flour is obtained, by grinding the seeds—*Wheat*, *Pyrus*, *Fruentulum*—and to them we are indebted for our bread and pastry. The nutritive properties of wheaten flour are so well known as not to need comment.

Bread, (F.) *Pain*, mixed with milk, constitutes the ordinary emollient poultice;—and the *crumbs of bread*, *Micæ panis*, are a common excipient for pills and boluses.

TRITICUM ARVENSE, *T. repens*—t. *Cereale*, *T. æstivum*—t. *Ceuallos*, *T. æstivum*—t. *Compactum*, *T. æstivum*—t. *Faginum*, *Polygonum*, *Polygonum* *fagopyrum*—t. *Gaertnerianum*, *T. æstivum*—t. *Glau'cum*, *T. hybernium*, and *T. repens*—t. *Hybernium*, see *T. æstivum*—t. *Muticum*, *T. hybernium*—t. *Linnæ'um*, *T. æstivum*—t. *Pruinosum*, *T. æstivum*.

TRITICUM REPENS, *T. arven'se* seu *glau'cum*, *Agropyrum lævis'simum*, *Bromus glaber*, *Gramen caninum* seu *Dioscor'idis* seu *repens*, *Loliat'ceum* *radic'è repen'te*, *Dog-grass*, *Couch-grass*, *Quickens*, *Twiteh-grass*, (Sc.) *Romnachs*, (F.) *Chicendent*. The roots are sweet, and possess aperient properties. They are said, also, to be diuretic.

TRITICUM SICULUM, *T. hybernium*.

TRITIO, Contritio, Trituration.

TRITON PALUSTRIS, see Ectozoa.

TRITORIUM, Infundibulum.

TRITURA, Contritio.

TRITURAT'ION, *Trituratio*, *Tritura*, *Tritus*, *Trit'io*, *Tripis*, from *terere*, *tritum*, 'to bruise.' The act of reducing a substance to powder.

TRITURATION, Contritio.

TRITUS, Contritio, Trituration.

TRIVELIN, see Liver.

TRIVENTER, Trigastrie.

TROCAR, *Trochar*, *Acus paracent'ica* seu *paracent'ica* seu *tri'quetra vulg'aris*, *Paracent'ion*, *Paracentet'ion*. Corrupted from the (F.) *Troisquarts* or *Trocart*, 'three quarters;' so called from the triangular shape of the point—*Tri'quetrum*. An instrument used for evacuating fluids

from cavities, particularly in ascites, hydrocele, &c. A trocar consists of a perforator or stylet and a canula, which is so adapted to the perforator, that, when the puncture is made, both enter the wound with facility; after which, the stylet being withdrawn, the canula remains in the wound and affords the fluid a ready passage outwards.

TROCART, Trocar.

TROCH, Trochiscos.

TROCHANTER, *Trokanter*, *Rota'tor*, from *τροχάω*, 'I turn.' Anatomists have given the names *great* and *little trochanter* to two processes at the proper extremity of the femur. The *great trochan'ter*, *glutius*, is the one on the outside; the *less* is lower down and more internally. Both afford attachment to rotator muscles of the thigh, and hence their name. Chaussier, by the word *trochanter*, means the larger process; the smaller he calls *trochantin*.

TROCHANTER'IAN, *Trochanteria'nus*, *Trokanter'ian*. That which belongs or relates to the great trochanter.—Chaussier.

TROCHANTIN'IAN, *Trokan'tin'ian*, *Trochantin'ius*. That which belongs or relates to the trochantin or less trochanter.

TROCHAR, Trocar.

TROCHES OF CARBONATE OF LIME,

Trochisci carbonatis calcis—t. of Chalk, *Trochisci carbonatis calcis*—t. Escharotic, *Trochisci escharotici*—t. Gum, *Trochisci gummosi*—t. of Gum Arabic, *Trochisci aacicie*—t. of *Ipeacuanha*, *Trochisci ipeacuanhæ*—t. of *Lactucarium*, *Trochisci lactucarii*—t. *Liquorice*, *Trochisci glycyrrhizæ*—t. *Liquorice*, with opium, *Trochisci glycyrrhizæ cum opio*—t. of *Magnesia*, *Trochisci magnesiæ*—t. of *Morphia*, *Trochisci morphiæ*—t. of *Morphia* and *ipeacuanha*, *Trochisci morphiæ et ipeacuanhæ*—t. *Nitre*, *Trochisci nitratis potassæ*—t. of *Peppermint*, *Trochisci menthæ piperitæ*—t. of *Tartaric acid*, *Trochisci acidi tartarici*.

TROCHIA, Orbit.

TROCHILIA, Trochlea.

TROCHIN, from *τροχάω*, 'to turn.' Chaussier has given this name to the smaller of the tuberosities at the upper extremity of the os humeri; because it gives attachment to one of the rotator muscles of the arm,—the subcapularis.

TROCHIN'IAN, *Trochin'ianus*. That which belongs or relates to the trochin.

TROCHISCUS, *Phthois*, *Artis'cus*, *Collis*, *Colis*, *Phthois'cos*, *Rot'ula*, *Cyclis'mos*, diminutive of *τροχός*, 'a wheel.' A troch or round table, (F.) *Trochisque*. See *Pastil*, and *Tabella*. A solid medicine, prepared of powders, incorporated by means of mucilage, crumb of bread, juices, of plants, &c. In French nomenclature, the *trochisque* differs from the *tablette*,—the former containing no sugar. This form of preparation is adapted for the purpose of allowing the medicines of which the troch is composed to dissolve slowly in the mouth, and to pass gradually into the stomach.

The Parisian Codex has two formulæ for troches for external use.

TROCHISCI ACA'CLÆ, *Troches of Gum Arabic*. (*Acac. 3iv*; *Amyli 3j*; *Sacchar. lbj*. Make into a mass with rose water, and form lozenges.—Ph. E.) To allay cough.

TROCHISCI AC'IDI TARTAR'ICI, *Troches of Tartaric Acid*. (*Acid. Tartar. 3ij*; *Sacchar. 3viii*; *Ol. Limon. Mx*. Form into troches with mucilage.—Ph. E.) Refrigerant and demulcent.

TROCHISCI BECHICI NIGRI, *T. glycyrrhizæ glabræ*.

TROCHISCI CARBONA'TIS CALCIS, *T. e Cretâ*, *T. Cretæ* (Ph. U. S.), *Troches of Carbonate of lime* or *chalk*, *Lozenges for the heartburn*, *Tabellæ ad*

ardo' rem ventriculi seu cardiac' gicæ seu ad sodan. (Cret. ppt. $\frac{3}{4}$ iv; acaciæ. in pulv. $\frac{3}{4}$ j; myrist. in pulv. $\frac{3}{4}$ j; sacch. in pulv. $\frac{3}{4}$ vj. M. Form into troches with water.) Antacid and absorbent.

TROCHISCI E CRETÂ, T. carbonatis calcis—t. Cretæ, T. carbonatis calcis.

TROCHISCI seu PASTILLI EMETINÆ PECTORALES, *Pectoral Lozenges of Emetine.* (Sugar, $\frac{3}{4}$ iv; emetine, 32 grains. Form into lozenges of nine grains each.) One is a dose.

TROCHISCI ESCHAROTICI, *Escharot'ic Troches.* (*Hydrarg. oxymerc.* p. 8; amyli, p. 16; mucilag. g. tragac. q. s. Make into troches, to which a few drops of laudanum may be added.—Ph. P.) Used in foul ulcers, to remove excrescences, &c.

TROCHISCI ESCHAROTICI DE MIN'IO. (*Oxyd. plumbi. rubr.* p. 16; *hydrarg. oxymerc.* p. 32; *mic. panis.* p. 128; *aq. rosæ.* q. s.—Ph. P.) Used like the last.

TROCHISCI GLYCYRRHIZÆ, T. G. *Glabræ seu Bech'ici nigri, Liquorice troches, Black pectoral lozenges.* (Ext. glyc. glabr., gum. acac. aa p. j; sacch. purif. p. ij. Dissolve in warm water; strain; evaporate, and form into troches.—Ph. E.) Demulcent. To allay cough.

TROCHISCI GLYCYRRHIZÆ CUM O'PIO, T. G. et opii (Ph. U. S.) seu *Theba'ici seu sedati'vo-balsam'ici seu pectorales regis Danorum, Liquorice Troches with Opium, Wistar's Lozenges.* (Opil, in pulv. $\frac{3}{4}$ ss; ext. glycyrrhiz. in pulv., sacchar. in pulv., acaciæ, in pulv., aa $\frac{3}{4}$ xx; ol. anisi, f $\frac{3}{4}$ j. Make into troches.—Ph. U. S.) Demulcent and anodyne.

Spitta's Lozenges for coryza, hoarseness, and sore-throat, are composed as follows:—(Cubeb. recent. in pulv. $\frac{3}{4}$ j; extract. glycyrrhiz. $\frac{3}{4}$ j; myrrocyl. $\frac{3}{4}$ j; acaciæ, $\frac{3}{4}$ iv; syrup, q. s. Make into troches of ten grains each.

TROCHISCI GUMMO'SI, *Gum Troches.* (G. acac. p. v; amyli, p. i; sacch. pur. p. xij. Make up the troches with rose-water.—Ph. E.) Demulcent.

TROCHISCI IPECACUAN'HÆ, *Troches of Ipecacuanha.* (Ipecacuanh. in pulv. $\frac{3}{4}$ ss; sacchar. in pulv. $\frac{3}{4}$ xiv; marantæ, in pulv. $\frac{3}{4}$ iv; mucilag. tragacanth. q. s. Make into troches.—Ph. U. S.)

TROCHISCI LACTUCARI, *Troches of Lactucarium.* (Prepared with Lactucarium in the same proportion and manner as the opium lozenges.—Ph. E.) Each lozenge contains between the fifth and sixth of a grain of lactucarium.

TROCHISCI MAGNE'SIÆ, *Troches of Magnesia.* (Magnes. $\frac{3}{4}$ iv; sacchar. $\frac{1}{2}$ j; myrist. pulv. $\frac{3}{4}$ j; mucilag. tragacanth. q. s. Make into troches.—Ph. U. S.) Useful in acidity of the stomach, with constipation.

TROCHISCI MENTHÆ PIPERITÆ, *Troches of Peppermint.* (Ol. menth. piperit. f $\frac{3}{4}$ j; sacchar. in pulv. $\frac{1}{2}$ j; mucilag. tragacanthæ, q. s. Make into troches.—Ph. U. S.) Used as a carminative.

TROCHISCI MORPH'Æ, *Troches of Morphia.* (Morph. Muriat. $\frac{1}{2}$ j; Tinct. Tolut. $\frac{3}{4}$ ss; Sacchar. $\frac{3}{4}$ xv. Make into lozenges, with mucilage, weighing 15 grains each.—Ph. E.) Each lozenge contains about one-fortieth of a grain of muriate of morphia. Used to allay cough.

TROCHISCI MORPH'Æ ET IPECACUAN'HÆ, *Troches of Morphia and Ipecacuanha.* (Morph. mur. $\frac{1}{2}$ j; Ipecac. pulv. $\frac{3}{4}$ j; Tinct. Tolut. f $\frac{3}{4}$ ss; Sacchar. $\frac{3}{4}$ xv. Make into a mass with mucilage, and form lozenges, 15 grains each.—Ph. E.) Each lozenge contains about one-fortieth of a grain of the muriate. Used to allay cough.

TROCHISCI NITRA'TIS POTAS'SÆ, *Nitre Troches.* (Potassæ nitrat. p. j; sacch. pur. p. iii. Make into troches with mucilage of tragacanth.—Ph. E.) In slight inflammatory sore-throat and fever.

TROCHISCI PECTORALES REGIS DANORUM, T. glycyrrhizæ cum opio—t. Sedativo-balsamice, T. glycyrrhizæ cum opio.

TROCHISCI SODÆ BICARBONATIS, *Troches of Bicarbonate of Soda.* (Sodæ bicarb. $\frac{3}{4}$ iv; sacchar. in pulv. $\frac{1}{2}$ j; mucilag. tragacanth. q. s. To be made into a mass, and divided into troches, each weighing ten grains.—Ph. U. S.) Antacid.

The *Pastilles de Vichy, Pastilles digestives d'Hauterive-Vichy, Pastilli digestivi Dorcei'i*, composed of bicarbonate of soda, oil of peppermint, and sugar, are much used in France, like the Vichy waters, in cardialgia, gout, and lithiasis.

TROCHISCI STIBIATI, *Tabellæ antimoniales Kunkeli*—t. Thebaici, T. glycyrrhizæ cum opio.

TROCHISQUE, Trochiscos.

TROCH'ITER, from τροχαιον, 'to turn.' The larger of the two tuberosities at the upper extremity of the os humeri; so called because it affords insertion to rotator muscles.—Chaussier.

TROCHITER'IAN, in the language of Chaussier, means any thing belonging or relating to the trochiter.

TROCH'LEA, *Trochil'ia*, (F.) *Trochlée* ou *Troklée*. A pulley; from τροχαιον, 'to turn,'—for example, the articular surface at the lower extremity of the os humeri; so called from its forming a kind of pulley on which the ulna moves, in flexion and extension of the forearm. Also, the cartilaginous pulley over which the tendon of the trochlearis muscle passes, at the upper and inner part of the orbit.

TROCHLEA LABYRINTHI, Cochlea.

TROCHLEARIS, Obliquus superior oculi.

TROCHLEATEUR, GRAND, Obliquus superior oculi—t. Petit, Obliquus superior oculi.

TROCHLÉE, Trochlea.

TRO'CHOID, from τροχος, 'a wheel,' and εδος, 'resemblance.' *Trochoi'des, Ax'ea commissu'ra*. An articulation, in which one bone turns upon another, like a wheel upon its axle; as the *Atlo-ido-azoid articulation*.

TROCHOIDES, Axæa commissura.

TROENE, Ligustrum vulgare.

TROISQUARTS, Trocar.

TROKANTER, Trochanter.

TROKANTERIAN, Trochanterian.

TROKANTINIAN, Trochantinian.

TROKLÉE, Trochlea.

TROLIUS LAXUS, T. *America'nus, Globe flower; Order, Ranunculacæ; indigenous; has the same medical properties as the Ranunculus.*

TROMOMANIA, Delirium tremens.

TROMOPARANÆA, Delirium tremens.

TROMOS, see Paralysis agitans.

TROMOSPASMUS, Tremor.

TROMPE, Tuba—t. d' *Eustache, Eustachian tube—t. de Fallope, Tuba Fallopiana—t. Utrérine, Tuba Fallopiana.*

TROMUS, Tremor.

TRONC, Trunk—t. *Brachial, Brachial artery—t. Basilaire, Basilar artery.*

TROPÆ'OLUM MAJUS, *Nasturtium Ind'icum seu Peruvia'num, Acrivi'ola, Flos sanguin'eus monard'i, Cardamin'dum majus, Card'amum majus, Indian Cress. Ord. Geraniacæ. Ser. Syst. Octandria Monogynia.* (F.) *Cresson des Indes, Cresson du Mexique, Capucine.* An antiscorbutic.

TROPÆ'OLUM TUBERO'SUM, *Ulluco*. A tuberous plant, cultivated in the Sierra, Peru. It is smaller than the potato.

TROPHALIS, Colostrum.

TROPHE, Aliment, Cibatio, Pabulum.

TROPHI. Same etymon as Trophy. In in-

sects, the parts of the mouth that are employed in receiving and preparing the food.

TROPHICAL, *Trophicus*, (F.) *Trophique*. Same etymon as *Trophy*. Relating to nourishment or nutrition.

TROPHICAL NERVES. The organic nerves, or nerves of the sympathetic system, have been so termed, from the belief that they are concerned in nutrition.

TROPHIMOS, Nutrition.

TROPHONEUROSES, from *τροφή*, 'nourishment,' and *neurosis*. Morbid conditions of the process of nutrition, owing to modified nervous influence. — Romberg.

TROPONOUSI, *Trophonousi*, from *τροφή*, 'nourishment,' and *nosos*, 'a disease.' Diseases of nutrition.

TROPHY, *Trophia*, *τροφή*, a suffix denoting 'nourishment or nutrition:' as in *hypertrophy*, 'excess of nourishment or nutrition.'

TROPONU'SI, *Morbi tropici*, from *τροπή*, the solstitial or tropical point where the sun turns or alters its course, and *nosos*, 'a disease.' Diseases that prevail in the tropics.

TROSTER, see Spirit.

TROU, Foramen.—*t. Anonyme*, Foramen stylo-mastoideum.—*t. Aveugle ou Borge*, Cæcum foramen.—*t. Borge* of the Medulla oblongata, Foramen cæcum of the medulla oblongata.—*t. de Botal*, Botal foramen. *Ovale foramen*—*t. Déchiré antérieur*, Lacerum anterior foramen.—*t. Déchiré postérieur*, Lacerum posterior foramen.—*t. Épineux*, Spinale foramen.—*t. Mentonnier*, Mental foramen.—*t. Optique*, Foramen opticum.—*t. Orbitaire supérieur*, Foramen supra-orbitarium.—*t. Orbitaire supérieur*, Orbital foramen superior.—*t. Ovale*, Ovale foramen.—*t. Rond, petit*, Spinale foramen.—*t. Sous-orbitaire*, see Suborbital canal.—*t. Sous-pubien*, Obturatorium foramen.—*t. Sphéno-épineux*, Spinale foramen.—*t. Sus-orbitaire*, Foramen supra-orbitarium.

TROUS CAROTIDIENS, Carotica foramina.—*t. Condyloïdiens*, Condyloidea foramina.—*t. de Conjugaison*, Conjugationis foramina, see Vertebra.—*t. Olfactifs*, Olfactory foramina.—*t. Rachidiens*, Spinal foramina.

TROUSSE, Case (pocket).

TROUSSE-GALANT, Cholera morbus.

TROUSSEAU, Fasciculus.

TROUSSEAUX ARRIÈRE-MÉSENTÉRIQUES (F.). A name given by Winslow to the nervous plexus situate between the two mesenteric plexuses, of which it is a dependence, and which descends into the pelvis to concur in the formation of the hypogastric plexus.

TRUBS, Lycoperdon tuber.

TRUCK, see Medicament.

TRUCULENTUS, Homicidal.

TRUDGEBACKED, see Hump.

TRUELOVE, Paris, Trillium latifolium.

TRUFFE, Lycoperdon tuber.

TRUFFLE, Lycoperdon tuber.—*t. Hart's*, Elaphomyces granulatus,

TRULLA, Fart.

TRUMBUS, Thrombus.

TRUMPET CREEPER, Tecoma radicans.—*t. Flower*, Tecoma radicans.—*t. Weed*, Eupatorium purpureum.

TRUNCUS, Trunk.

TRUNK, *Truncus*, (F.) *Tronc*. The principal part of the body, to which the limbs are articulated. The trunk has been divided into three parts:—*a superior*, the head; *a middle*, the thorax, and *an inferior*, the pelvis. These three regions contain the great splanchnic cavities. They are united by a common stalk—the vertebral column. To the middle are articulated the

superior extremities; and, to the inferior, the lower extremities.

The trunk of an artery, vein, nerve, or lymphatic, is its largest part, which has not yet given off branches. It is, also, applied to the parent vessel, after branches have been sent off.

TRUSS, from (F.) *trousser*, 'to tie up.' *Brachierium*, *Brachierium*, *Brachilè*, *Brachiorolum*, *Brachiorolum*, *Amma*, *Hamma*, *Subligaculum*, *Subligatura*, (F.) *Brayer*. A hernial bandage for keeping hernia reduced. The ancient trusses were inelastic. They consisted simply of a pad and strap. At the present day, elastic bandages are alone employed. They possess the advantage of making an equable and continuous pressure on the aperture of the sac, and of yielding to the changes of form in the abdominal parietes. The most important part of these bandages is a piece of narrow steel, flat, and adapted to the shape of the body. This is the *spring*, (F.) *Ressort*. It embraces the body on the side affected; extends from beyond the spine, and terminates opposite the ruptured part, by an iron plate to which it is riveted. The posterior surface of this plate is furnished with a convex pad, (F.) *Pelote*, adapted in shape and size to the aperture which it has to close. The spring is covered with leather, and its inner surface is padded; a strap extends from its posterior extremity; passes round the sound side, and is attached to the plate of the pad. This strap is pierced with numerous holes, which allow the patient to tighten or slacken it at pleasure. Trusses have been variously constructed; and different patents obtained for the modifications. They all act upon the above principle. Sometimes they have a double pad.

TRYGODES, Feculent.

TRYPANON, Trepan.

TRYPSIS, Trepanning.

TRYPSIS, Friction.

TSCHERBET, Sherbet.

TSHETTIK, Upas tieuté.

TSIANA, Costus.

TSJAMPACA, Michelia champaca.

TUBA, Tube.—*t. Aristotelica*, Eustachian tube.—*t. Cæca*, Tuba Fallopiana.—*t. Uterina*, Tuba Fallopiana.—*t. Vocalis*, Tube, vocal.

TUBAL PREGNANCY, see Pregnancy, tubal.—*t. Respiration*, see Murmur, respiratory.

TUBE, *Tuba*, *Trumpet*, *Salpinx*, (F.) *Trompe*. A name given to some parts which are hollow and have the shape of a trumpet.

TUBE, ALIMENTARY, Canal, alimentary.—*t. Eustachian*, Eustachian tube.

TUBE, FALLOPIAN, *Tuba Fallopiana* seu *Fallopia* seu *uteri* seu *cæca*, *Metrosalpinx*, *Hysterosalpinx*, *Oviductus muliebris*, *Vas deferens muliebris*, *Ductus varius* seu *Processus lateralis* seu *Meatus semina* seu *Meatus semina* seu *uteri*, *Oviduct*, *Vector canal*, (F.) *Trompe de Fallope* ou *utérine*, is a canal, floating in the abdomen, and extending on each side from the superior angles of the uterus as far as the sides of the brim of the pelvis. They are 4 or 5 inches long. Near the uterus, they are straight and very narrow; but, in their course, they widen, become tortuous, and terminate by a free, expanded, and *fimbriated extremity*; called, also, *Folia* seu *ornamentum*, *Morsus diaboli*, *Infundibulum* seu *Digitatio* seu *Fimbria* seu *Vexille* seu *Lacinia tubarum Fallopia*, (F.) *Morceau frangé*, *Pavillon de la trompe*, *Morceau du diable*. Within, the tubes are hollow and lined by a mucous membrane: the tissue exterior to which is spongy and erectile. The use of the Fallopian tubes is to conduct the sperm to the ovarium, and the fecundated ovum back to the uterus. They establish a communication between the cavity of the peritoneum and that of the

uterus, the abdominal termination being termed *Ostium abdominale*; the uterine *Ostium uterinum*. When the tube embraces the ovary, a funnel-shaped cavity—*Corolla infundibuliformis*—is formed at its extremity, into which the ovule passes.

TUBE, INTESTINAL, Canal, alimentary.

TUBE, (ŒSOPHAGEAL, Stomach Tube. A long elastic gum tube, capable of being passed into the œsophagus or stomach.

TUBE, POLLEN, see Pollen.

TUBE, RECTAL, *Defecation tube*. An elastic gum tube, similar to the last, which is sometimes passed into the colon, to remove air from that intestine, or to enable enemas to be thrown up into it.

TUBE, STOMACH, see Tube, œsophageal.

TUBE, VOCAL, *Tuba vocalis*. The part of the air passages above the inferior ligaments of the larynx, including the passages through the nose and mouth.

TUBER, Hump, Lycoperdon tuber, Tubercle —t. Annulare, Pons varolii—t. *Atloïdo-occipital*. Rectus capitis posticus minor—t. Cervinum, Elaphomyces granulatus—t. Cibarium, Lycoperdon tuber.

TUBER CINE'REUM, *T. C. cum Infundibulo*. A grayish tubercle, seen at the base of the brain behind the commissure of the optic nerves, which is continuous with the infundibulum.

TUBER CORPORIS CALLOSI, see Corpus callosum.

TUBER FRONTALIS, Frontal protuberance —t. Gulosorum, Lycoperdon tuber —t. Gutturorum, Bronchocele —t. *Ischio-trochantérien*, Quadratus femoris—t. Ischii, see Ischiatic—t. Nigrum, Lycoperdon tuber —t. Parietale, see Parietal Bones —t. Pellagra, Pellagra—t. Tympani, Promontory of the tympanum.

TUBERA TERRÆ, Lycoperdon tuber.

TUBERCLE, *Tuber culum*, *Tuber*, *Strangalion*, *Strangalis*, *Tuber culous deposit*, (vulg.) a *Kernel*, a *Knot*, (F.) *Tubercule*. A tumour in the substance of organs, from the production of new matter;—sensation null; growth sluggish. In *Pathological Anatomy*, the term is generally given to a species of degeneration which consists of an opaque matter, of a pale yellow colour: this, in its crude condition, has a consistence analogous to that of concrete albumen: it subsequently becomes soft and friable, and gradually acquires a consistence and appearance analogous to those of pus. Tubercles may be developed in different parts of the body: but they are most frequently observed in the lungs and mesentery. Pulmonary tubercles, *Pneumophymata*, *Tuber cula Pulmonum*, (F.) *Tubercules pulmonaires*, are the cause of *Tubercular Phthisis*. *Mesenteric tubercles* are often met with in the same affection, but particularly in *Tabes Mesenterica*. Laënnec classes tubercles among the accidental tissues, which have no resemblance to the natural tissues, and which never exist except in consequence of morbid action. Others consider them as a scrofulous degeneration. Broussais conceives that they are produced exclusively by inflammation of the white vessels, whose existence, however, is more than questionable. When tubercles in any organ are few in number, they may pass to the state of permanent induration without danger to the patient; but when they are very numerous, they usually cause serious mischief. Dr. Baron considers that they are produced by hydatids. The morbid condition which gives rise to the development of tubercles, *Tuberculization*, is termed *Tuberculo'sis*, *Morbus tuberculo'sus*, *Strumosis* seu *Phymato'sis pulmonum*, and *Dyscrasia Tuberculo'sa*. Tubercles are seen of various forms,—to wit, in fine points, (F.) *Poussière tuberculeuse*, gray and yellow granu-

lations, miliary tubercles; and gray and yellow tubercular masses, softened, and cretaceous.

TUBERCLE, BLACK, Melanosis—t. Painful subcutaneous, see Neuroma.

TUBERCLES OF THE AREOLA, see Mamma.

TUBERCLES, PEARLY, *T. sebaceus miliary*, *Follicular elevations*. Small collections of sebaceous matter in the sebaceous glands or ducts, often seen in the thin skin of the lower eyelids, where they sometimes attain an inconvenient size.

TUBERCLES, SEBACEOUS MILIARY, T. pearly.

TUBER'CU'LA ARAN'TII, *Corpora* seu *puscula Arantii*; *Tubercles of Arantius or Aranzi*. Small eminences at the middle part of the convex edge of the semilunar valves of the pulmonary artery and aorta.

TUBER'CU'LA ARTHRITICA, Calculi, arthritic —t. Cerebri, Encephalophymata—t. Gonorrhœica, Gonorrhœa impura—t. Hepatis, *Hépatostromosis*—t. Intestinorum, Enterophymata —t. Nervorum nodosa, Ganglions, nervous —t. Parva duræ matris, Glandula Pacchioni —t. Peritonæi, Cœliophymata —t. Pulmonum, see Tubercles (of the lungs) —t. Quadrigemina, Quadrigemina Tubercula.

TUBER'CU'LAR, *Tuber culous*, *Tuberculo'sus*, *Tuber culate*, *Tuber culated*, (F.) *Tuberculeux*, *Tuberculé*. That which relates to tubercles, or which is formed by tubercles.

TUBERCULAR OR TUBERCULOUS CONSTITUTION or DIATHESIS, *Phthisio'sis*, is the habit of body which predisposes to tubercular phthisis; as *Tuberculous Cachexia*, *Tuberculo'sis*, *Morbis tuberculo'sus*, *Dyscrasia tuberculo'sa*, is the condition of one labouring under tuberculosis.

TUBERCULAR MATTER. A morbid production, contained in cysts, or loose in the tissue of organs. It varies, in its consistence, at the different periods of the tubercular affection: being, at first, compact, and yellowish; at times, calcareous; afterwards pulaceous, semi-fluid, and curdy. See Tubercle.

TUBERCULAR PHTHISIS is that produced by the development of tubercles in the lungs.

TUBERCULAR SORE THROAT, Pharyngitis, follicular.

TUBERCULATE, Tubercular.

TUBERCULATE SARCO'MA of ABERNETHY, *Emphy'ma Sarcoma tuberculo'sum*, *Tumor tuberculo'sus*. Tubercles, firm, round, and clustering; pea-sized or bean-sized; yellowish or brownish-red; when large, disposed to ulcerate, and produce a painful, malignant, and often fatal sore. Found chiefly in the lymphatic glands of the neck; often, simultaneously, in other glands and organs.

TUBERCULATED, Tubercular —t. Liver, Cirrhosis.

TUBERCULE, Tubercle.

TUBERCULÉ, Tubercular.

TUBERCULES BIJUMEAUX, Quadrigemina tubercula —t. des Glandes lymphatiques, Scrofula —t. du Foie, *Hépatostromosis* —t. des Ganglions Mésentériques, *Tabes mesenterica* —t. of the Larynx and Fauces, Pharyngitis, follicular —t. Miliary, see Granulation —t. *Pisiformes*, Mamillary tubercles—t. *Pulmonaires*, see Tubercle —t. *Quadrijumeaux*, Quadrigemina corpora.

TUBERCULEUX, Tubercular.

TUBERCULI'TIS. A word of hybrid formation: from *tuberculum*, 'a tubercle,' and *itis*, denoting inflammation. Inflammation of a tubercle or of tubercular masses. —Jos. Frank.

TUBERCULIZATION, see Tubercle.

TUBERCULOSIS, see Tubercle, and Tubercule. Cachexia — *t.* Laryngis et Tracheæ, Phthisis laryngea — *t.* Pulmonum, Phthisis pulmonalis — *t.* Vertebrae, Vertebral disease.

TUBERCULOSUS, Tubercular.

TUBERCULOUS, Tubercular — *t.* Deposit, Tubercle — *t.* Disease of the Lungs, Phthisis pulmonalis — *t.* Dust, *Poussière tuberculeuse*.

TUBERCULUM, diminutive of *tuber*, Tubercle.

TUBERCULUM CINE'REUM, 'Ash coloured tubercle,' *Fasci'ola cin'erea*. A mass of cineritious substance at the top of the calamus scriptorius, which is continuous below with the posterior horn of cineritious substance in the cord, and upon the sides with the corpus testiforme.

TUBERCULUM LACRYMALE, see Lachrymal puncta — *t.* Lower, Lower, tubercle of.

TUBEROSITAS, Tuberosity — *t.* Tympani, Promontory of the tympanum.

TUBÉROSITÉ BICIPITALE, Bicipital tubercle — *t.* Sciatique, see Ischiatic.

TUBEROSITY, *Tuberos'itas*, (F.) *Tubérosité*, from *tuber*, 'a bump.' Anatomists have given this name to an eminence or process, the surface of which is unequal and rough; and which gives attachment to muscles or ligaments.

TUBES OF FERREIN, see Uriniferous Tubes.

TUBI MEMBRANACEI, see Villous membranes.

TUBULAR, *Tubula'ris*, *Tubulo'sus*, (F.) *Tubuleux*, *Tubulaire*. Same etymon as the next. Relating to or having the form of a small tube. An epithet applied to glands — *Glandulæ tubulo'sæ*, which, like those of the stomach, have the shape of small tubes.

TUBULAR BLOWING SOUND, see Murmur, respiratory — *t.* Neurine, see Neurine — *t.* Substance of the Kidney, see Kidney.

TUBULE, *Tubulus*; diminutive of *Tuba*, 'a tube.' A small tube.

TUBULEUX, Tubular.

TUBULI BELLINI, Uriniferous tubes — *t.* Contorti, see Uriniferous tubes — *t.* Duræ matris, Sinuses of the dura mater — *t.* Ferreini, see Uriniferous tubes — *t.* Nervi, Nerve fibres — *t.* Pelvis renum, see Calix — *t.* Recti, see Rete testis and Uriniferous tubes — *t.* Seminiferi, see Testicle — *t.* Seminales, see Testicle — *t.* Uriniferi Bellini, Uriniferous tubes.

TUBULOSUS, Tubular.

TUBULUS, Cannula — *t.* Centralis Modiol, see Tractus spiralis foraminulentus.

TUBUS ACUSTICUS, Ear-trumpet — *t.* Alimentarius, Canal, alimentary — *t.* Intestinorum, Canal, alimentary — *t.* Nervorum, Neurilemma — *t.* Pollinicus, see Pollen — *t.* Vermicularis cæci, Appendix vermiformis cæci.

TUCKAHO, Lycoperdon tuber.

TUE-CHIEN, Colechium autumnale.

TULBAGHIA ALLIA'CEA. A South African plant, *Nat. Ord.* Asphodeleæ, the bulbs of which, boiled in milk, are recommended in phthisis and as an anthelmintic.

Tulbaghia cepa'cea and *T. viola'cea*, serve the same purpose.

TULIP, CAPE, *Homeria collina* — *t.* Tree, *Liriodendron*.

TULPIER, *Liriodendron*.

TULIPIFERA LIRIODENDRON, *Liriodendron tulipifera*.

TUMEFACATIO, Swelling — *t.* Mollis, Puffiness. **TUMESCENTIA PITUITOSA**, Leucophlegmatia.

TUMEUR, Tumour — *t.* Anomale, see Hæmatodes fungus — *t.* Blanche, Hydrarthrus — *t.* Caverneuse, see Hæmatodes fungus — *t.* Encéphaloïde, Encephaloid — *t.* Érectile, see Hæmatodes

fungus — *t.* Fongôide, Encephaloid — *t.* Fongueuse, Encephaloid — *t.* Hémorrhagiale circonscrite, see Aneurism — *t.* Lymphatique des articulations, Hydrarthrus — *t.* Ossivoire, Encephaloid — *t.* Pulpeuse, Encephaloid — *t.* Spléniforme, Encephaloid — *t.* Spongieuse, Encephaloid — *t.* Variqueuse, see Hæmatodes fungus.

TUMOR, Swelling, Tumour — *t.* Albus, Hydrarthrus — *t.* Anomalous, Hæmatodes fungus — *t.* Artium, Obdormitio — *t.* Capitis Sanguineus, Cephalæmatoma — *t.* Carneus, Sarcoma — *t.* Cysticus, Encystis — *t.* Cysticus serosus, Hygroma — *t.* Desmoids, Tumour, fibrous — *t.* Fibrosus, Tumour, fibrous — *t.* Ficosus, Ficus — *t.* Flatulentus, Emphysema — *t.* Fugax, see Fugacious — *t.* Fungosus articuli, Spina ventosa — *t.* Fungosus sanguineus, Hæmatodes fungus — *t.* Gangliiformis, Ganglion, nervous — *t.* Genarum, Meloncus — *t.* Geni albus, Genocæce — *t.* Glandulæ parotidæ, Parotidoneus — *t.* Lacteus, Galactocœle — *t.* Lardeus, Steatoma — *t.* Malarum, Meloncus — *t.* Ovarii, Oarioncus — *t.* Parotidis, Parotidoneus — *t.* Prostate, Prostatoncus — *t.* Renalis, Nephroncus — *t.* Sanguineus, Hæmatoma — *t.* Squamiformis carnosus, Lepidosarcoma — *t.* Tuberculosus, Tuberculate sarcoma — *t.* Tunicatus, Encystis — *t.* Uteri, Hysteroncus.

TUMOUR, from *tumeo*, 'I swell;' *Tumor*, *Oncos*, *Oncus*, *Epar'ma*, *Epar'sis*, *Emphy'ma*, *Excrecen'tia* (Sauvages); *Ephy'ma*, (Sc.) *Suadme*, (F.) *Tumour*. A rising or prominence, of greater or less size, developed by a morbid cause in some part of the body. Cullen defines a tumour to be "a partial swelling without inflammation." Sauvages restricts the term to a rising formed by the congestion of a fluid; and he calls *excrecences* those which have a fleshy or osseous consistence. Boyer defines it "any preternatural eminence developed on any part of the body." Good uses it synonymously with tubercle. Boyer's is the best general definition. Tumours differ greatly from each other; according to their seat, the organs interested, their nature, &c.

The French use the term *Tumeur blanche* synonymously with *White Swelling*; — see Hydrarthrus. The term is, however, applied by the generality of surgeons to swellings of the greater joints, without change of colour of the skin, and of a more or less firm consistence, which are dependent upon disease of the osseous or of the soft parts about the joint.

Scrofula is the most common cause of these affections.

TUMOUR, CELLULO-FIBROUS, Tumour fibré-cellular — *t.* Epithelial, see Epithelial.

TUMOUR, FIBRO-CELLULAR, *Cellulo-fibrous tumour*. A soft and elastic form of fibrous tumour, a variety of the *gelatinous sarcoma* of Rokitsky, of which the softer kind of polypus of the nose is an example.

TUMOUR, ERECTILE. A tumour produced by the development of a soft, vascular tissue, which is susceptible of elevation and depression. See Erectile.

TUMOUR, FIBRO-PLASTIC, Sarcoma.

TUMOUR, FIBROUS, *Fibrous growth*, *Fibron'a*, *Fibroid*, *Ino'ma*, *Tumor fibro'sus* seu *desmoids*, *Inchylo'ma*. A tumour or growth formed of fibrous tissue.

TUMOUR, LAMINATED NACREOUS FATTY, Cholesteatoma — *t.* Milk-like, Encephaloid — *t.* Phantom, see Fantom.

TUMOUR, VARICOSE, *Tumor varicosus*. A circumscribed, bluish tumour, formed by the dilatation and turgescence of the capillary vessels of a part.

TUMOUR, VASCULAR, see Hæmorrhoids and Nævus.

TUNAS, see Cactus opuntia.

TUNBRIDGE, MINERAL WATERS OF. A celebrated acidulous chalybeate in Kent, England, a few miles distant from the village of Tunbridge, and thirty-six miles south of London, which is much frequented. It contains carbonic acid, carbonate of iron, and sulphate of magnesia.

TUNDA OCULI, Conjunctiva membrana.

TUNDUS CORDIS, Mucro cordis.

TUNIC, *Tunica, Chiton, a coat.* An envelope. A name given to different membranes, which envelop organs; as the tunics or coats of the eye, stomach, bladder, &c.

TUNICA ACINALIS, Uvea — *t. Aciniformis, Uvea — t. Acinosa, Uvea — t. Adnata, Conjunctiva — t. Adventitia, see Artery — t. Agnata, Conjunctiva — t. Alba oculi, Sclerotic — t. Albuginea, Albuginea, see Kidney — t. Albuginea oculi, Sclerotic — t. Allantoides, Allantois — t. Arachnoides, Arachnoid membrane — t. Aranea, Arachnoid membrane — t. Caduca, Decidua — t. Candida oculi, Sclerotic — t. Carnosa, Panniculus carnosus — t. Cellularis, see Artery — t. Conjunctiva, Conjunctiva — t. Costalis, Pleura — t. Crassa, Decidua — t. Crystallina, Arachnoid membrane — t. Dartos, see Dartos — t. Dura oculi, Sclerotic — t. Durior et crassa, Sclerotic — t. Elastica, see Artery — t. Elytroides, Vaginal coat — t. Erythroides, Cremaster — t. Exterior ovi, Decidua — t. Externa, see Artery — t. Extima oculi, Sclerotic — t. Farciminialis, Allantois — t. Fibrosa, see Folliculi Graafiani — t. Filamentosa, Decidua — t. Glabra, see Artery.*

TUNICA GRANULO'SA. An investment of the ovum exterior to the zona pellucida, described by Dr. Barry, but the existence of which, as a structure, distinct and independent from the membrana granulosa, is denied.

TUNICA HORTENSIS, *Dianthus caryophyllus* — *t. Innominata oculi, Sclerotic — t. Interna oculi, Retina — t. Intima, see Artery.*

TUNICA JACO'BI, Membra'na Jacob'bi, Jacob's coat or membrane. A highly delicate serous membrane, interposed between the retina and the choroid coat, discovered by Dr. Jacob, of Dublin. Its structure is curious, being composed of cylindrical, transparent, highly refractive staff-shaped bodies, and hence called *Stratum bacillo'rum* seu *bacilla'tum, Bacillar layer of the retina*, which are arranged perpendicularly to the surface of the retina, their outer extremities being imbedded, to a greater or less depth, in a layer of the pigmentum nigrum.

TUNICA MEDIA, see Artery — *t. Muscularis, Dartos — t. Nervea oculi, Retina — t. Nervorum, Neurilemma — t. Perforata, Uvea — t. Prætensa abdominis, Peritonæum — t. Propria, see Kidney — t. Reticularis, Retina — t. Retiformis, Retina — t. Rhagoides, Choroidæ tunica, Uvea — t. Rubicunda scroti, Dartos — t. Rubra, Dianthus caryophyllus — t. Ruyschiana, see Choroid — t. Subcostalis, Pleura — t. Thoracis, Corset — t. Uvalis, Uvea — t. Uvea, Uvea — t. Uviformis, Uvea — t. Vaginalis, see Vaginal — t. Vaginalis Linguae, see Tongue — t. Vaginalis Propria, Vaginal coat of the Testicle — t. Vaginalis reflexa, Vaginal coat of the Testicle — t. Vaginalis testis, see Testicle — t. Vasculosa oculi, Choroid, see Eye.*

TUNICA VASCULO'SA TESTIS, Pia mater testis. An extremely delicate membrane, consisting of minute ramifications of the spermatic vessels united by areolar tissue, which separates the lobules of the testis from each other. It is situated immediately within the tunica albuginea, and encloses the substance of the gland. It sends processes inward that separate the lobules, in the same manner as the pia mater is reflected between the convolutions of the brain.

TUNICA VITREA, Hyaloid membrane.

TUNICULA, Clitoris.

TUNIQUE ALBUGINÉE, Albuginea — t. Commune de Bichat, Endangium.

TUN'NELLING. A term, sometimes employed for the treatment of stricture of the urethra, by making pressure upon it from within so as to promote its removal by absorption.

TUNNYFISH, SALTED, Omotrichos.

TUPHLO-ENTERITIS, Typhlo-enteritis.

TUPHUS, Typhus.

TUPINARIA, Talpa.

TURBA HÆMORRHOIDALIS, see Hæmorrhoids.

TURBATIONES ANIMI, Affections of the mind.

TURBINATED BONES, Ossa turbina'ta. Bones shaped like a top; from *turbo, turbinis* or *turben*, 'a top'; *Ossa spongio'sa, O. ter'tia maxilla' superioris, Cornua, Concha na'rium seu vent'ris, Tur'bines nasi seu na'rium, Conchyl'ia, Buccina, Lam'ina spongio'sæ nasi, (F.) Cornets.* Very thin, bony plates, rolled up in the form of horns, and situate in the nasal fossa. 1. *Ossa spongio'sum supe'rius* or *Cornet de Morgagni*; — the highest. 2. *Ossa spongiosum me'dium*, the middle cornu of French anatomists, — the *ethmoid'al cornu* or *Ossa spongiosa superioris seu turbinatum*, of English anatomists, *Ossa convolutum superioris, Concha, C. na'rium superior seu Morgagnia'na, (F.) Cornet moyen.* This bone projects inwards and downwards into the cavity of the nose, from the ethmoid cells at the side of the nasal lamella, and serves to enlarge the organ of smell. It is convex towards the septum, and concave outwards. 3. *Ossa spongiosum seu turbinatum inferius, Concha inferior, Convolutum inferius oss, Os sous-ethmoid'al (Ch.)*, projects at the under part of the side of the nose, is convex towards the septum, and concave outwards. It is connected to the os maxillare, os palati, and os unguis. 4. *Sphenoid'al cornu.*

All the turbinated bones extend the olfactory surface.

TURBINES NASI seu NARIUM, Turbinated bones.

TURBITH BLANC, Globularia alypum — t. Mineral, Hydrargyrum vitriolatus — t. Montpellier, Globularia alypum — t. Noir, see Hydrargyrum oxydum cinereum — t. Plant, Convolvulus turpethum.

TURBO CEREBRI, Pineal gland.

TURGENTIA, Turgescence.

TURGES'CENCE, Turges'cency, Turgid'ity, Tur'gidness, Turgescent'ia, Turgen'tia, Orgas'mus, from *turgescere*, 'to begin to swell', *turgere*, 'to swell.' Superabundance of humours in a part. The term *Turgescence of Bile* was formerly used to denote the passage of that fluid into the stomach, and its discharge by vomiting. A vital action of turgescence or expansibility — *Turgor vita'lis* — has been presumed to exist in certain organs, as in the capillaries; but the generality of physiologists do not admit it.

TURGESCENTIA, Turgescence.

TURGESCENT'IA VESIC'ULE FELL'Æ, Ascites Hep'ato-cyst'icus, Physco'nia seu Cystoc'le bili'o'sa, Hepat'al'gia Petiti'na, Hydrocholecys'tis, Hydrops vesicæ fell'æ, (F.) Hydropisie de la Vésicule du Fiel, Distension de la Vésicule du Fiel, Cholecystiectasie. Distension of the gall-bladder by bile.

TURGIDITY, Turgescence.

TURGINESS, Turgescence.

TURGOR VITALIS, see Turgescence.

TURMERIC, Curuma longa, Sanguinaria Canadensis.

TURN OF LIFE, see Menses.

TURNING, Versio from (F.) *turner*, 'to turn.' (F.) *Versio.*

VERSIO FETUS. The operation of bringing

down the feet, or some part of the lower extremity, when the presentation of the child is such that it cannot be delivered by the natural efforts.

TURNIP, *Brassica rapa* — t. Dragon, *Arum triphyllum* — t. Indian, *Arum triphyllum* — t. Pepper, *Arum triphyllum*.

TURNS, Menses.

TURPENTINE, *Terebinthina* — t. Bordeaux, see *Pinus sylvestris* — t. Briangon, see *Pinus cembra* — t. Chio, see *Pistacia terebinthus* — t. Common, see *Pinus sylvestris* — t. Common, American, see *Pinus sylvestris* — t. Cyprus, see *Pistacia terebinthus* — t. Damarra, see *Pinus damarra* — t. Dombeya, see *Dombeya excelsa* — t. Horse, see *Pinus sylvestris* — t. Strasburg, see *Pinus picea*.

TURPENTINE TREE, *Pistacia terebinthus*.

TURPENTINE, VENICE, see *Pinus larix* — t. Venice, true, see *Pistacia terebinthus* — t. White, see *Pinus sylvestris*.

TURPETHUM, *Convolvulus turpethum* — t. Minerale, *Hydrargyrum vitriolatus* — t. Nigrum, *Hydrargyri oxydum cinereum*.

TURQUETTE, *Herniaria glabra*.

TURQUOISE. This stone has been long imagined to change its colour, according as the wearer is at the time in good or bad health: — a superstitious notion.

TURTLE BLOOM, *Chelone glabra* — t. Green, *Chelonia mydas* — t. Head, *Chelone glabra*.

TURUNDA, Tent.

TURUN'DÆ ITALICÆ. Pastes made of the finest and most glutinous wheat — known under the names of *Maccaroni*, *Vermicelli* and *Italian* or *Cagliari* paste.

TURUNDULA, Tent.

TUSCARORA ACID SPRING, see Canada, mineral waters of.

TUSSACA RETICULATA, *Goodyera pubescens*.

TUSSEDO, Tussis.

TUSSICULA, Tussis.

TUSSICULAR, *Tussicula'ris*, *Tussiculosa*, from diminutive of *tussis*, 'a cough.' Relating to a cough, or to a slight cough.

TUSSICULOSUS, Tussicular.

TUSSILAGO, *T. Farfara seu vulga'ris*, from *tussis*, 'a cough,' which it relieves; *Be'chium*, *Be'chion*, *Cal'ceum equi'num*, *Ohama'leu'cè*, *Chama'gei'ron*, *Fil'i'us ante patrem*, *Farfarel'la*, *Gallio'marchus*, *Farfara*, *F. Be'chium*, *Un'gula caballi'na*, *Coltsfoot*, *Foal-foot*, (F.) *Pas d'âne*, *Herb de Saint-Quirin*. Ord. Compositæ. Sex. Syst. Syngenesia Polygamia Superflua. It is demulcent and expectorant, and has been used in coughs, and pulmonary affections in general, and in some cutaneous diseases.

TUSSILAGO FARFARA, Tussilago — t. Hybrida, T. petasites.

TUSSILAGO PETASITES, *T. hybrida seu petasites hermaphrodita*, *Petasites*, *P. major seu rubens seu officinalis seu hybrida seu vulga'ris*, *Butterbur*, *Pestilent-wort*. The roots of this plant have been recommended as aperient and alexipharmic. They have a strong smell; and a bitterish, acrid taste, of the aromatic kind, but not agreeable.

TUSSILAGO PETASITES HERMAPHRODITA, T. petasites — t. Vulgaris, Tussilago.

TUSSIS — diminutive, *Tussicula*; *Begma*, *Bex*, *Tusse'do*, *Bexis*, *Tussis nervosa*, *Catar'rhus*, *Pneusis Tussis*, *Cough*, *Coughing*, (Prov.) *Couf*, (Sc.) *Hoist*, *Hoast*, *Hoist*, (F.) *Toux*. A violent, sonorous, frequent, and short expiration, by means of which the air, in passing rapidly through the bronchia and trachea, carries along with it the mucus of the parts, which forms the sputa. The cough is said to be dry, when unaccompanied by expectoration. It is symptomatic of many diseases. See *Bronchitis*.

TUSSIS AMPHEMERINA, *Pertussis* — t. *Asinica*, *Pertussis* — t. *Bronchica*, *Cough*, tubal, see *Bronchitis* — t. *Canina*, *Pertussis* — t. *Catarrhalis Simplex*, *Catarrh* — t. *Clamosa*, *Pertussis* — t. *Clangosa*, *Pertussis* — t. *Convulsiva*, *Pertussis* — t. *Delassans*, *Pertussis* — t. *Ferina*, *Pertussis* — t. *Pueros strangulans*, *Pertussis* — t. *Quinta*, *Pertussis* — t. *Senilis*, *Bronchitis* (chronic) — t. *Spasmodica*, *Pertussis* — t. *Stomachalis*, *Pertussis* — t. *Suffocans*, *Pertussis* — t. *Suffocativa*, *Pertussis* — t. *Tussiculosa*, *Pertussis*.

TUSSIVE; *Tussivus*; from *tussis*, 'cough.' Belonging or relating to cough: — as *tussive vibration*; the vibration of the parietes of the chest, caused by coughing.

TUTAM'INA, pl. of *Tutamen*, 'a defence,' 'a protection,' from *tutare*, 'to defend.' Parts are so called which defend or protect each other.

TUTAMINA CEREBRI are the scalp, pericranium, bones of the skull, and the meninges.

TUTAMINA OCULI are the eyebrows, eyelids, and lachrymal apparatus.

TUTENAG, Zincum.

TUTHIA, Tutia.

TUT'IA, *Pom'pholyx*, *Cadmia*, *Alfusa*, *Mesera*, *Cupri'tis*, *Tuthia*, *Oxydum zinci impurum*. The oxyd of zinc that attaches itself to the chimneys of furnaces in which ores of zinc are smelted. It is in the form of gray incrustations; and is sometimes used for making an eye-salve. It is prepared, for this purpose, by levigation.

TUTTERING, Stuttering.

TWIGG'S HAIR DYE, see Hair dye.

TWIN, Gemellus.

TWINGE, see Pain.

TWINKLING OF THE EYE, Nictation.

TWINLEAF, COMMON, *Jeffersonia Bartoni*.

TWINROOT, *Orehis*.

TWINS, SIAMESE. Two brothers, Chang and Eng, connected by means of a ligamentous band passing from the epigastrium of one to that of the other, who were exhibited in various parts of Europe and America, about the year 1830. They have lived in North Carolina, are married, and have children. See *Xiphopages*.

TWITCH-GRASS, *Triticum repens*.

TWITCHING, Tic.

TYCH'ICA, from *τυχη*, 'accident.' Fortuitous lesions or deformities. The 7th class of Good's *Nosology*.

TYLE, Callosity.

TYLOMA, Callosity.

TYLOPH'ORA ASTHMATICA, *Asclepias asthmatica*, *Cynan'chum Ipecacuan'ha seu vomito'rium seu viridifo'rum seu tomento'sum*, *Ipecacuan'ha* of the *Isle of France*, (F.) *Ipecacuan'ha blanc de l'île de France*. Order, *Asclepiadaceæ*. A native of various parts of the East Indies, both of the continent and islands. The root is acrid, and is used on the coast of Coromandel as a substitute for *ipecacuan'ha*. In large doses, it is emetic; in smaller, often repeated, it is cathartic. It is recommended, also, as a diaphoretic, and especially in humeral asthma.

TYLOSIS, Callosity.

TYLUS, Callosity.

TYMMA, Wound.

TYMPAN, Tympanum.

TYMPANAL, Tympanic.

TYMPANAL, *Cadre du tympan*.

TYMPANIA, Tympanites.

TYMPANIAS, Tympanites.

TYMPANIC, *Tympanicus*, *Tym'panal*, (F.) *Tympanique*; same etymon as *Typanum*. Relating to the tympanum, as 'tympanic canal.'

TYMPANIC NERVE, see *Petrus ganglion*.

TYMPANIQUE, Tympanic.

TYMPANITA, Tympanites.

TYMPANITE, Tympanites.

TYMPANITES. *Emphysema seu Pneumato-*
sis Abdominis, Hydrops Tympanites, Affec-
tio tympanitica, Ectympanosis, Hydrops abdominis
aëreus, Aëro-sis, Aëro-enterectasia, Aërenter-
ectasia, Enterærectasia, Physocœlia, Tympano-
sis, Physema, Physes, Hydrops siccus et flut-
ulentus, Tympania, Tympanias, Tympanita, Tym-
punitis, Emphysema Tympanites, E. Tympanit-
icum, Meteorismus, Tympany, Meteorism, Wind
dropsy, (F.) Tympanite, Pneumatose péritonéale,
Balonnement, from τυμπανον, 'a drum,' so called
because the abdomen is swelled with wind, and
sounds like a drum when struck. A swelling of
the abdomen, caused by accumulation of air in
the intestinal tube or in the peritoneum. Tym-
panites may be idiopathic or symptomatic. The
former depends upon exhalation of air from the
inner surface of the intestine, or from the decom-
position of substances contained in it; the latter
is the result of some organic affection, and often
depends upon obliteration of the digestive tube,
which prevents the gas from escaping. Idiopa-
thic tympanites may be more readily cured. The
symptomatic is usually fatal. On dissection in
such cases, the stomach and intestines are found
enormously distended; with strangulation or
disorganization of some part of the mucous mem-
brane of the intestine. Carminatives may be
proper when tympanites depends upon atony of
the digestive organs: but, in cases of the symp-
tomatic kind, they cannot be productive of ad-
vantage, and may do harm.

Sometimes the air is contained in the cavity
of the peritoneum—constituting *Tympanites ab-*
dominalis, Aëropéritonie, of some:—the accumu-
lation in the intestines being called *Tympanites*
intestinalis.

TYMPANITES, UTERINE, Physometra.

TYMPANITIS, *Inflammatio tympani*; from
τυμπανον, 'a drum,' and itis, denoting inflamma-
tion. Inflammation of the lining membrane of
the middle ear. Also, tympanites.

TYMPANOSIS, Tympanites.

TYMPANUM, *τυμπανον, 'a drum,' The Drum*
or barrel of the ear. The names Cavity of the
Tympanum or of the Drum, Cavitas seu Cavum
Tympani, Antrum seu cavitas antro-a Auris,
(F.) Tympan, Cavité du tympan, Caisse du tam-
bour ou du Tympan—have been given to a cavity
of an irregular shape, which constitutes the mid-
dle ear, and has been compared to a drum. It is
situate in the pars petrosa of the temporal bone,
between the meatus auditorius externus and the
labyrinth. This cavity is lined by a mucous
membrane; and communicates, externally, by
means of the Eustachian tube, which is situate
between it and the pharynx. The tympanum
has six parietes. 1. An external, which is almost
entirely formed of the Membrane of the Drum,
Membrana tympani, Diaphragma seu Mediasti-
num seu Tegumentum auris, Myringa, Myrina,
(F.) Membrane du tympan—a species of thin,
transparent, circular septum; convex within, and
concave without, which closes the inner extremity
of the meatus auditorius. 2. An internal, on
which are the Fenes'tra ovalis, Promontory; and
Foramen rotundum. 3. A superior. 4. An in-
ferior, on which is the Glenoid fissure. 5. A pos-
terior, on which are the opening of the mastoid
cells, and the pyramid. 6. An anterior, on which
are the Processus cochleaformis and the inner
orifice of the Eustachian tube. The cavity of the
tympanum is crossed by a series of four small
bones, articulated with each other by diarthrosis,
moved by certain muscles, and representing a
kind of bent lever extending from the membrana
tympani to the fenestra ovalis. They are called

the *Bones or Ossicles of the Ear*, and are the
malleus, incus, os orbiculare, and stapes.

For the nervous plexus of the tympanum, see
Petrosal Ganglion, and Otic Ganglion.

TYMPANUM MINUS, see Fenestra—t. Secunda-
rium, see Fenestra.

TYMPANY, Tympanites—t. of the Womb,
Physometra.

TYPE, *Τύπος, Periodus morbi*, from τυπω, 'a
stamp,' itself from τυττω, 'I strike.' The type is
the order in which the symptoms of a disease ex-
hibit themselves, and succeed each other. It
may be continued, intermittent, or remittent.

TYPHA ANGUSTIFOLIA has the same prop-
erties as

TYPHA AROMATICA, Acorus calamus, and

TYPHA LATIFOLIA, (τύπος, 'a marsh,' [?]). Cat-
tail Reed mace, Bulls-egg, (F.) Massette, Masse
ou Quenouille d'Eau. A plant common in
marshes and ponds, in this country and Europe.
Ord. Typhaceæ. Sex. Syst. Monœcia triandra.
Its roots have been used in dysentery and blen-
norrhagia. The young shoots, before they reach
the surface of the water, when boiled, taste like
asparagus.

The down, (F.) Aigrette, Poils, which accom-
panies the flowers, has been applied to burns like
cotton.

TYPHICUS, Typhoid.

TYPHIQUE, Typhoid, Typhus.

TYPHLITIS, Typhlo-enteritis.

TYPHLO-ENTERITIS, *Typhlo-enteritis, Ty-*
phlitia, Typhloteritis, Inflammatio Cæci, Phleg-
monous Tumour of the Cæcum, (F.) Inflammation
du Cæcum, Cécite (Piorry); from τυφλος, cæcus,
and enteritis, 'inflammation of the intestines.'
Inflammation of the cæcum, occasionally leading
to perforative ulceration.

TYPHLOENTERUM, Cæcum.

TYPHLOPS, Cæcus.

TYPHLOSIS, Cæcitas.

TYPHLOTERITIS, Typhlo-enteritis.

TYPHLOTERON MONOCOLON, Cæcum.

TYPHLOTERUM, Cæcum.

TYPHLOTES, Cæcitas.

TYPHODES, Typhoid.

TYPHOËMIE ENTÉRIQUE, see Typhus.

TYPHOHÆMIA, see Putrefaction.

TYPHOID, *Typhoides, Typhodes, (τυφώδης,*
'stupefied,') Typhicus, Typhosus, Lac'tica, (F.)
Typhique, Typhoide, Typhode. From typhus,
and αἶδος, 'resemblance.' Appertaining to or re-
sembling typhus; as Typhoid Fever. Also, ac-
cording to some, Typhous Fever, typhus. Fuchs
has employed the word Typhoide for a family of
diseases, to distinguish them from Phlogoses: of
this family, Stomacæ, Pharyngocacæ, and Pneu-
mocacæ are examples.

TYPHOID AFFECTION, see Typhus—t. Fever of
India. Cholera.

TYPHOMANIA, from τυφος, 'stupor,' and
μανία, 'delirium.' Delirium with stupor, Subde-
lirium. The kind of delirium common in
typhus.

TYPHOMANIA, Coma vigil, Insomnia.

TYPHONIA, Coma vigil.

TYPHOSEPSES, Typhoseptoses.

TYPHOSEPTOSES, *Typhosep'ses*, from τυφος,
and επις, 'putrefaction.' Typhous affections.

TYPHOSUS, Typhoid.

TYPHOUS, *Typhosus, (F.) Typhique.* Same
etymon as Typhus. Belonging or relating to
typhus.

TYPHOUS DEPOS'IT. A peculiar substance of
new formation found in the areolar membrane,
between the mucous and muscular coats of the
patches of Peyer in typhoid fever.

TYPHOUS FEVER, see Typhoid.

TYPHUS. *Semipet'is*, *Typhus*, *Febris Typho'des* seu *asthen'ica* seu *adynam'ica*, *Asthenop'gra*, (F.) *Fièvre adynamique ou ataxique ou adynamique ataxique*, *Typhus d'Europe*; from τυφος, 'stupor.' A fever characterized by small, weak, and unequal, but usually frequent pulse, with great prostration of strength, and much cerebral disturbance; its duration being usually from a fortnight to three weeks or longer. It is continued fever, accompanied with great cerebral irritation and prostration. (See Fever.) By most writers, this disease has been divided into two varieties—the *Typhus mitior* and *Typhus gravior*. **TYPHUS MITIOR** seu *simplex* seu *nervosus* seu *comatosus*, *Febris lenta nervosa seu putrida nervosa seu hec'tica maligna nervosa*, *Blethrop'gyrus*, *Neurop'gra*, *Neuropyr'etus*, *Nervous Fever*, (F.) *Fièvre nerveuse*, is characterized by slight shiverings, heavy, vertiginous headach; great oppression, peculiar expression of anxiety, nausea, sighing, despondency, and coma or quiet delirium. **TYPHUS GRAVIOR** seu *car'cerum* seu *castr'ens* seu *anthropophthor'acus* seu *contagiosus exanthemat'icus* seu *nostratus* seu *Europæ'us* seu *communis* seu *bell'icus* seu *exanthemat'icus* seu *Hiber'nicus* seu *petechia'lis*, *Febris nervosa epidem'ica* seu *nervosa exanthemat'ica* seu *maligna cum sopor'e* seu *nervosa petechia'lis* seu *pestilential'is* *Europæ'us* seu *puncticula'ris* seu *saprop'gra* seu *car'cerum* et *nosocomio'rum* seu *putrida* seu *continens putrida* seu *continua putrida* seu *pestilens maligna* seu *navitica pestilential'is* seu *petechia'lis*, *Septop'gra*, *Porphryotyphus*, *Pulicariis morbus*, *Pestis bell'ica*, *True* or *Exanthemat'ic* or *contagious typhus*, *Nervous fever* with *exanthematous eruption*, *Putrid*, *Jail*, *Hospital* or *Ship Fever*, (F.) *Fièvre des Camps*, *F. nautique*, *F. d'hôpital*, *F. nosocomiale*, *F. des Prisons*, *F. lenticulaire*, *Typhus contagieux* is attended with rigors and heat, alternating; little or no perspiration; pulse, tense and hard, usually quick, but fluttering; pain over the forehead and vertex; a 'mulberry rash,' appearing on the fifth to the seventh day of the disease, and lasting till the end; delirium succeeded by stupor, signs of incipient putrescence,—as true petechiæ, vibices, hemorrhages, &c. Typhus, under particular circumstances, seems to be communicable by contagion; but if proper attention be paid to cleanliness and ventilation, there is not much danger of communication. The general principles of treatment, laid down under the head of Fever, are applicable here. All the varieties of continued fever resemble each other in the cardinal points, although they may differ in many of their characters. Ventilation and cleanliness; warm or cold ablation or sponging; keeping the bowels free; and attending to symptoms as they may rise, with the avoiding of all irritation of every kind, are the main indications. When symptoms of great prostration occur, the strength must be supported by wine; and tonics be exhibited, with the mineral acids. Of the tonics, the sulphate of quinia is perhaps as good as any, whilst it is free from the objection which applies to most others,—of being too bulky. In the whole management of this affection, however, the physician has to discriminate carefully between oppression and prostration. The former may require evacuations; the latter cannot.

Of late years, it has been maintained by many, that there is an essential difference between typhoid fever and typhus; that the former is an abdominal affection, and dependent upon, or connected with, an inflamed or ulcerated condition of the mucous follicles of the intestines; hence it has received the names *Fièvre mésentérique*, *F. typhoïde*, *Maladie ou Affection typhoïde*, *F. entérotyphoïde*, *Entérite typho-hémique*, *Typhoémie*

entérique, *Entéro-mésentérique typhoïde*, *Exanthéma intestinal*, *Dolich'entérie*, *Dolichentérite*, &c., *Hæm'diclidite*, *Entérite folliculeuse*, *Gastro-entérite* with nervous affection of the brain, *Follicular gastroenteritis*, *F. enteritis*, *Enteritis folliculo'sa*, *Typhus abdomina'lis*, *T. intestinal'is*, *T. Sporadicus*, *Enterotyphus*, *T. ganglionar'is abdominal'is*, *Febris intestinal'is ulcerosa*; *F. nervosa gastrica*, *F. nervosa enterica*, *F. enterica*, *F. nervosa mésentérica*, *Enterhelio'sis nervosa*, *Typhus enter'icus*, *Ileop'gra*, *Ileotyphus*, *Typhoid affection*, *Enteric fever*. Accompanying this form of fever there is generally a typhoid eruption, consisting of rose-red papulæ, (see Spots, red,) which appear mostly on the abdomen. They are distinct from true petechiæ, as they can be removed by pressure; and distinct also from sudamina, which are perhaps the universal accompaniments of sweating, and are consequently a variety of miliaria. The common continued fever or synochus of Great Britain, and of this country, has been presumed by some to be this variety: but farther observations are needed to establish the distinctions which have been attempted; and there is reason to believe, that the abdominal affection is a complication, existing in certain cases and epidemics of adynamic fever, and not in others. The *Red Tongue Fever* of Kentucky is considered by Professor Bartlett to be typhoid fever.

Certain modern French writers have considered the Typhus of the European continent, *Typhus d'Europe*, *pestilential fever*, *contagious typhus*, *Hungary fever*, *Hospital fever*, *camp fever*, *jail fever*, *lenticular petechial fever*, (F.) *Peste d'Europe*, &c., to be identical with the typhoid affection. They admit but one fever on the European continent—the typhoid. Typhus, they maintain—but by no means establish the position—to be peculiar to England.

TYPHUS ABDOMINALIS, see Typhus—*t. d'Amérique*, Fever yellow—*t. Anthracicus*, Plague—*t. Anthropophthoracus*, Typhus—*t. Aphthodes*, *Aphthæ*—*t. Bellicus*, Typhus—*t. Bengalensis*, *Cholera*—*t. Bubonicus*, Plague—*t. Carcerum* Typhus gravior—*t. Castr'ensis*, Typhus gravior—*t. Cerebral*, *Meningitis*, *cerebro-spinal*—*t. Cerebro-spinal*, *Meningitis*, *cerebro-spinal*—*t. Comatosus*, Typhus mitior—*t. Communis*, Typhus—*t. Contagious*, Typhus—*t. Contagious*, Typhus—*t. Contagious exanthematicus*, Typhus—*t. d'Europe*, Typhus—*t. d'Europe*, see Typhus—*t. Exanthematicus*, Typhus—*t. Ganglionaris abdominalis*, see Typhus—*t. Gravissimus*, Plague—*t. Hibernicus*, *T. gravior*—*t. Icterodes*, Fever, yellow—*t. Jaune*, Fever, yellow—*t. Intestinalis*, see Typhus—*t. Miasmatica ataxique putrida jaune*, Fever, yellow—*t. Morbillosus*, *Rubeola*—*t. Nervosus*, Typhus mitior—*t. Nostratus*, Typhus—*t. d'Orient*, Plague—*t. Pestilentialis*, Plague—*t. Pestis*, Plague—*t. Petechialis*, Typhus gravior—*t. Puerpéral*, see Peritonitis—*t. Scarlatinus*, *Scarlatina*—*t. Simplex*, Typhus mitior—*t. Sporadicus*, see Typhus—*t. Tropicus*, Fever, yellow—*t. True*, Typhus—*t. Vesicularis*, Pemphigus.

TYPIC, *Typical*, *Typicus*, (F.) *Typique* Same etymon as *Type*. Characterized by periodicity,—as a 'typical fever;' or one which observes a particular type.

TYPOSES, see Periodicity.

TYPUS, Type.

TYRANNUS, see Critical days.

TYRBASIA, Agitation.

TYRBE, Agitation.

TYREM'ESIS, *Tyromet'esis*, from τυρος, 'cheese,' and εμεσις, 'vomiting,' *Tyros'is*, *Tyren'is*. Vomiting of curdy matter, in infants especially.

TYREUSIS, Tyremesis, Tyrosis.

TYRIA, Porrigo decalvans.

TYRIASIS, Elephantiasis Arabica.

TYRINE, Casein.

TYRODES, Cheesy.

TYROMA, Porridge decalvans.

TYROS, Cheese.

TYROSIS, Tyremesis. Also, the curdling of milk in the stomach; *Tyreusis*.

TYROTEMESIS, Tyremesis.

U.

UBERIS APEX, Nipple, Papilla.

ULA, Gingivæ.

ULATROPH'IA, from *ουλον*, 'the gum,' and *atrophia*, atrophy. Shrinking of the gums; falling away of the gums.

ULCER, *Ulcus*, genitive *Ulcera*, diminutive *Ulcusculum*; *Hulcus*, *ἑλκος*, *Helcos*, *Elco'sis*, *Elco'ma*, *Helcus*, *Helco'ma*, a Sore. (Sc.) *Sair*, (Prov.) *Kile*, (F.) *Ulcère*. A solution of continuity in the soft parts, of longer or shorter standing, and kept up by some local disease or constitutional cause. Richerand makes four great distinctions between a wound and an ulcer. 1. A wound arises from the action of an extraneous body:—the cause of an ulcer is inherent in the economy. 2. A wound is always idiopathic:—an ulcer is always symptomatic. 3. A wound has essentially a tendency to heal, because the action of its cause has been momentary:—an ulcer, on the contrary, has a tendency to enlarge, because its cause persists. 4. The treatment of a wound is purely surgical; that of an ulcer is medical as well. The immediate cause of an ulcer is perverted nutrition; the formative cells losing their plastic properties. Ulcers have been variously divided, according to their nature;—into *simple*, *sinuous*, *fistulous*, *fungous*, *gangrenous*, *scorbutic*, *syphilitic*, *cancerous*, *incontinent*, *scrofulous*, *phagedenic*, *virulent*, *cacœthic*, *sordid*, *carious*, *varicose*, &c. The treatment consists in removing, by appropriate means, the internal and the local causes which keep up the ulceration, and in exciting the vessels of the part, where necessary, by appropriate bandaging, &c.

ULCER, JACOB'S, *Lupus*—u. Malignant, Hospital gangrene—u. Putrid, Hospital gangrene—u. Warty of Marjolin, see Warty Tumour of Cicatrices.

ULCERA INTESTINALIA, Gastrectosis—u. Laryngis, Phthisis laryngea—u. Serpentina oris, Aphthæ—u. Uteri, see Metrophthisis—u. Ventriculi, Gastrectosis—u. Vesicæ, Cystophthisis.

ULCERATED, *Ulcera'tus*, *Emulcera'tus*, *Helco'des*, (F.) *Ulcéré*; from *ulcus*, *ulceris*, 'an ulcer.' In the state of an ulcer. Affected with ulcers.

ULCERATIO INTESTINALIS, Enterelcosis—u. Ventriculi, Gastrectosis.

ULCERATION, *Ulcera'tio*, *Exulceratio*, *Pyogen'ia corrosi'va*. Same etymon. A superficial ulcer. The formation of an ulcer—*Exelco'sis*, *Elco'sis*, and *Helco'sis*.

ULCERATIVE, *Ulcera'tivus*, (F.) *Ulcératif*. Relating to ulceration,—as an 'ulcerative process.'

ULCÈRE, Ulcer—u. *du Larynx*, Phthisis laryngea.

ULCÉREUX, Ulcerous.

ULCEROUS, *Ulcero'sus*, *Helco'des*, *Helcoïdes*, (F.) *Ulcéreux*. Same etymon. Having the character of an ulcer. Affected with ulcers.

ULCERS, EGYPTIAN, *Ægyptia* ulcera.

ULCUS, Ulcer—u. *Atonicum*, *Rupia*—u. *Cacœthes*, see *Cacœthes*—u. *Cancerosum*, *Cancer*, *Chancre*—u. *Dacryodes*, see *Dacryodes*—u. *Depascens*, *Phagedenic* ulcer—u. *Ferinum*, *Phage-*

denic ulcer—u. *Fistulosum*, *Hypophora*—u. *Hypulum*, see *Hypulus*—u. *Narium* *fœtens*, *Ozæna*—u. *Obambulans*, *Phagedenic* ulcer—u. *Pulmonum*, *Phthisis pulmonalis*—u. *Sinuosum*, *Hypophora*—u. *Syriacum*, *Cynanche maligna*—u. *Telephium*, *Telephium*—u. *Tuberculosis*, *Lupus*—u. *Uteri*, *Hysterelcosis*, *Leucorrhœa*.

ULCUSCULA PRÆPUTII, Herpes præputii.

ULCUSCULUM, Ulcer—u. *Cancerosum*, *Chancre*.

ULE, Cicatrix.

ULETICUS, Gingivalis.

ULICUS, Gingivalis.

ULITIS, from *ουλος*, 'solid,' (*σπερ ουλη*, 'solid flesh:') hence, *ουλον*, 'the gum,' and *itis*, denoting inflammation. *Ouli'tis*, *Inflammat'io gingivæ*, *Periodonti'tis gingivarum*, (F.) *Géngivite*, *Inflammation des gencives*, from *ουλον*, 'the gum,' and *itis*, denoting inflammation. Inflammation of the gums.

ULITIS SEPTICA, Cancer aquaticus.

ULLUCO, *Tropæolum tuberosum*.

ULMAIRE, *Spiræa ulmaria*.

ULMARIA, *Spiræa ulmaria*—u. *Palustris*, *Spiræa ulmaria*.

ULMUS, *U. campestris* seu *glabra* seu *montana* seu *nuda* seu *sativa* seu *scabra*, *Ptelea*, *Common Elm*, (F.) *Orme*, *Ormeau*, *Orme pyramidale* ou *châmpêtre*. *Ord. Ulmaceæ*. *Sez. Syst.* *Pentandria Digynia*. The inner, rough bark—*Ulmus*, (Ph. L.)—of this tree is inodorous; and has a bitter, austere, and mucilaginous taste. It is considered tonic, alterative, and diuretic; and has been used, chiefly, in lepra and other cutaneous affections.

ULMUS ALA'TA, *Lynn Wahoo*. The bark of the root has been used successfully as a cataplasm in phlegmonous inflammation.

ULMUS AMERICA'NA, *U. fulva* seu *rubra*, *Rough-leaved*, *Red*, or *Slippery Elm*. The inner bark—*Ulmus*, (Ph. U. S.)—is highly mucilaginous, and is used in coughs, diarrhoeas, and dysenteries; also, as a poultice for tumours, lacerated and contused wounds, &c.

ULMUS CAMPESTRIS, *Ulmus*—u. *Fulva*, *U. Americana*, *U. rubro*—u. *Glabra*, *Ulmus*—u. *Montana*, *Ulmus*—u. *Nuda*, *Ulmus*—u. *Sativa*, *Ulmus*—u. *Scabra*, *Ulmus*.

ULNA, *O'lenê*, *Foc'ilê inferius* seu *majus*, *Canna Bra'chii*, *Os cubiti inferius*, *Os procubitalê*, *Cu'bitus*, *A'un'do bra'chii major*, *Cubit*. Anatomists have given this name to the bone of the forearm which forms the prominence of the elbow, during the flexion of that joint. It is the longer and larger of the two bones, and is inserted at the inner side. Its upper and larger extremity has, behind, a considerable process, called *olecranon*; and, before, a smaller one, called *coronoid*. These two eminences are separated from each other by the greater sigmoid or semilunar fossa or olecranonoid cavity, which receives the articular trochlea of the humerus. The tubercle of the ulna is a small, rough spot, under the root of the coronoid process, for the

insertion of the *Brachialis internus*. On the outside of this extremity of the bone is the *lesser sigmoid cavity*. It is articulated with the head of the radius. The inferior extremity of the cubitus, which is much smaller than the superior, and articulated with the radius, has a surface that corresponds to the fibro-cartilage which separates it from the os cuneiforme; and a pointed process, called *styloid*. The body of the ulna is manifestly triangular. It is articulated with the os humeri and radius, and, mediately, with the os cuneiforme. It is developed by three points of ossification.

Also, the elbow.

ULNAD, see Ulnar aspect.

ULNAR, Cubital.

ULNAR ASPECT. An aspect towards the side on which the ulna is situated.—Barclay. Ulnad is used adverbially by the same writer, to signify 'towards the ulnar aspect.'

ULNARIS, Cubital—u. Externus, see Cubital muscles—u. Gracilis, Palmaris longus—u. Internus, see Cubital muscles.

ULOCACE, Cancer aquaticus, Stomacacæ.

ULON'CUS, from *ουλον*, 'the gum,' and *ογκος*, 'tumour.' Swelling of the gums; epulis.

ULORRHAG'IA, *Oulorrhag'ia*, *Stomatorrhag'ia*, *Hæmorrhag'ia gingiva'rum*, *Ulorrhæ'a*, from *ουλον*, 'the gum,' and *ρηννμι*, 'I break forth.' Bleeding from the gums.

ULORRHŒA, Ulorrhagia.

ULOTICA, Cicatrissantia.

UL'TIMI-STERNAL. Professor Bécлар gave this appellation to the sixth or last bony portion of the sternum. He calls it also *Ensi-ster-nal*.

UL'TIMUM MO'RIENS, 'last dying.' A term given to the part of the economy which dies last.

ULVA CRISPA, *Fucus crispus*.

ULVA LATIS'SIMA, *Broad Green Laver*, *Green Sloke*, *Oyster green*; *Order*, *Algæ*; is used at table cooked in the same way as the next, to which it is greatly inferior.

ULVA UMBILICA'LIS, *Por'phyra lacinia'ta* seu *umbilica'lis*, *Lacinia'ted Purple Laver*, abundant on the shores of Great Britain, is pickled with salt and sold in London as *Laver*. It is eaten at table, stewed with pepper, butter or oil and lemon-juice or vinegar. It has been suggested, also, as an antiscorbutic.

UMBIL, YELLOW, *Cypripedium luteum*.

UMBILIC, Umbilical, Umbilicus.

UMBILICAL, *Um'bilic*, *Umbilica'lis*, *Umbilica'tus*, (F.) *Ombilical*, from *umbilicus*, 'the navel.' That which belongs or relates to the navel.

UMBILICAL ARTERIES, which exist only in the fœtus, seem, as it were, continuations of the primitive iliacs. They clear the umbilical ring, and proceed to the placenta, to which they carry the residuum of the blood sent to the fœtus by the umbilical vein. As soon as respiration is established, the blood ceases to pass by the umbilical arteries, which become obliterated, as well as the vein, and are transformed into two very strong ligamentous cords,—the vein forming the *Ligamentum rotundum* of the liver.

UMBILICAL CORD, *Funiculus umbilicalis*.

UMBILICAL REGION, *Re'gio umbilica'lis* seu *ga'strica* seu *Mesogastrica*, *Mesogas'trium*, (F.) *Région ombilicale*, is the middle region of the abdomen, in which the umbilicus is placed. The sides of this are called the *flanks* or *lumbar regions*. In the umbilical region are the omentum majus; the inferior extremity of the duodenum; the jejunum; and the greater part of the mesentery; the aorta; vena cava; the trunks of the renal arteries and veins; the origin of the spermatic arteries, &c.

UMBILICAL RING, *An'nulus umbilica'lis*, (F.) *Anneau ombilical*, is a fibrous ring which surrounds the aperture of the umbilicus, and through which umbilical hernia occurs in children.

UMBILICAL VESICLE, *Vesicula umbilicalis*.

UMBILICAL VESSELS, (F.) *Vaisseaux ombilicaux*, include the two arteries, and umbilical vein. The umbilical vein arises from the placenta, and terminates at the fissure on the inferior surface of the liver of the fœtus, to which it conveys the blood necessary for its nutrition.

UMBILICALIS, Umbilical.

UMBILICATUS, Umbilical.

UMBILICUS, *Umbili'cum*, *Umbo*, *Om'phalos*, *Radix* seu *Medium ventris*, *Mesompha'lium*, *Meson'phalum*, *Protne'sis* (novel of a new-born child); the *umbilic*, *navel*, (Prov.) *Spoon*; from *umbo*, 'the button or prominence in the midst of a buckler,' or from *υμφαλος*, or rather *ομφαλος*, which signifies the same, as well as the navel; (F.) *Ombilic*, *Nombril*. A round cicatrix, about the median line of the abdomen. It is in the situation of an aperture, which, in the fœtus, affords passage to the vessels of the umbilical cord.

UMBILICUS MARI'NUS, *Cotyle'don marina*, *Andro'sacæ*, *Acetabulum marinum*, *Andro'sacæ Matthi'oli*, *Fungus petræus marinus*. A submarine production, found on the rocks and the shells of fishes, in the south of France. It is reputed to be anthelmintic and diuretic.

UMBO, Elbow, Umbilicus.

UMBRELLA TREE, *Magnolia tripetala*.

UMSCHLAG, see Compress.

UNBOWELING, *Exenterismus*.

UNCARIA GAMBIR, *Nauclea Gambir*.

UNCAS, INDIAN, *Veratrum viride*.

UNCIA, Ounce.

UNCIFORM, *Uncifor'mis*, *Uncina'tus*, from *uncus*, 'a hook,' and *forma*, 'shape.' That which has the shape of a hook.

UNCIFORM BONE, *Os unci'form'e* seu *hamatum* seu *uncina'tum*, (F.) *Os crochu*. The fourth bone of the second row of the carpus. Its shape is very irregular. Inwards and forwards, it has a considerable eminence, which is curved upon itself, and gives attachment to the annular ligament of the carpus. It is articulated with the os semilunare, os magnum, os cuneiforme, and the fourth and fifth metacarpal bones. It is developed from a single point of ossification.

UNCIFORM EMINENCE, *Hippocampus minor*.

UNCIFORM PROCESS, *Proces'sus unci'na'tus* seu *Blumenbach'i*, *Great Process of the Ethmoid*. The most considerable of the thin irregularly twisted laminæ on the inferior surface of the ethmoid bone, which narrow the opening of the maxillary sinuses.

UNCINATUS, Unciform.

UNCINUS, Hook.

UNCTION, *Inunction*, *Unguentum*, *Unguentum hydrargyri*.

UNCUS, Hook.

UNDERCLIFF, ISLE OF WIGHT, ENGLAND, (CLIMATE OF.) The Undercliff comprehends a tract of country from Dunnose to St. Catherine's Hill, on the southeast coast, about six miles in length, and from a quarter to half a mile in breadth. It is well protected from the colder winds; and the climate is remarkably equable, as well as mild and dry, so that there are not many days during the winter in which the invalid cannot take exercise in the open air. It is an excellent climate for the pulmonary valetudinarian.

UNDIM'IA. A name given, by Paracelsus, to a tumour, filled with a gelatinous matter, like the white of an egg.

UNDOSUS, *Cymatodes*.

UNDULATIO. Fluctuation.

UNEDO, *Arbutus unedo* — u. *Papyracea*, *Arbutus unedo*.

UN'GEAL, *Ungea'lis*, from *unguis*, 'a nail.' Belonging, or relating, to the nails of the hands or feet.

UNGUEAL MATRIX, see Nail.

UNGUEN, *Unguentum* — u. *Articulare*, *Synovia*.

UNGUENT, ARMATORY, *Hoplochryisma*.

UNGUENTARIA, see *Myristica moschata*.

• UNGUENTA'RIOUS, *Myropo'us*, *Myrep'ous*, *Myropo'les*, *Myropo'lus*, *Pigmenta'rius*; from *unguentum*, 'an ointment.' One who makes and sells ointments. A perfumer.

UNGUEN'TUM, *Unguen*, *Myron*, *Uction*, (F.) *Onguent*, from *ungere*, *unctum*, 'to anoint.' Ointment. Ointments are topical applications, of a consistence more or less analogous to that of lard. They are used, chiefly, as local applications to ulcers and wounds; but are sometimes rubbed upon a part in cutaneous affections; and, especially, where it is desired that the constituents of the ointment should be absorbed. Ointments are of less consistence than cerates, although the terms are often used indiscriminately.

UNGUEN'TUM AC'IDI NITRO'SI, *Ointment of Nitrous Acid*, *Poma'tum seu Unguen'tum Ni'tricum seu Oxygena'tum*, (F.) *Pommade oxygénée*, *Graisse oxygénée*, *Pommade d'Alyon*. (*Adip. suilli*, lbj; *acid. nitros*. ʒvj. Melt the fat, and rub in the acid gradually, till cold. *Ph. E.*) Stimulant. Used in itch and other cutaneous eruptions, and in foul ulcers.

• UNGUENTUM ADIPOCERÆ CETORUM, *Ceratum cetacei* — u. *Ægyptiacum*, *Linimentum æruginis* — u. *Ægyptium album*, *Crinomyron* — u. *Æruginis*, *U. subacetatis cupri* — u. *Album*, *Unguentum cetacei* — u. *Album Resolvens*, *Linimentum ammoniæ fortius*.

UNGUENTUM DE ALTHEÆ, *U. de Terebin'thinâ et cerâ seu citr'inum seu res'inæ pini sylvestris compositum seu resumptivum seu lenitivum*, *Cera'tum de Althæâ*, *Emplast'rum mucilaginosum seu emolliens seu citr'inum seu flavum seu de Althæâ*, *Unguentum flavum*, *Morsh mal'low Ointment*, (F.) *Onguent de guimauve*, *O. d'althæa*. (*Olæi de mucilaginis*, 1000 p., *cerâ flavæ*, 250 p., *resinæ terebinthin.*, 3â 125 p. Liquefy; strain and stir till cold. *Ph. P.*) Common Olive Oil might be substituted for the Oil of Mucilages.

UNGUENTUM AMYGDALINUM, *Ceratum Galeni*.

UNGUENTUM ANTIMONII, *U. Antimonia'le*, *U. A. Tartarizati*, *U. A. Potassio-tar'tratiss*, *U. tar'tari emel'ici*, *Ointment of tar'tarized antimony*, *Antimonia'l Ointment*, *Tartar emetic ointment*. (*Antimon. et potass. tartrat. in pulv. subtiliss*. ʒij, *adipis*. ʒj. *Ph. U. S.*) If a drachm of this ointment be rubbed upon any part, night and morning, it will excite, in a few days, a painful pustular eruption. Hence, it is employed as a counter-irritant. This preparation is also called *Ung. Tar'tari stibiati seu emel'ici*, *Ung. e Tar'taro stibiato*, *Adeps Tartaræ stibii medicatus*, *Ung. Tar'tratis Potass'æ stibiati*, (F.) *Pommade d'Autenrieth*, *Pommade stibiée*. This method of employing the tartrate of antimony and potassa, as a revellent, is, at times, called *stibio-dermie*, (F.) *La méthode stibio-dermique*.

UNGUENTUM ANTIMONII POTASSIO-TARTRATIS, *Unguentum antimonii* — u. *Antimonii tartarizati*, *U. antimonii* — u. *Antipernius*, see *Antipernius*.

UNGUENTUM AQUÆ ROSÆ, *Ointment of Rose water*, *Cold Cream*. (*Aq. Rosæ*, fʒj; *ol. amygdal.* fʒij; *cetacei*, ʒss; *cerâ albæ*. ʒj. Melt together, in a water-bath, the oil, spermaceti, and wax; then add the rose-water, and stir until it is cold. *Ph. U. S.*) Cooling to irritated surfaces.

UNGUENTUM ARMARIUM, *Hoplochryisma* — u. *Arthanita*, see *Arthanita* — u. *Articulare*, *Synovia* — u. *Basilicon flavum*, *Ceratum resinæ* — u. *Basilicon nigrum*, *Unguentum resinæ nigre* — u. *Basilicon viride*, *Unguentum subacetatis cupri*.

UNGUENTUM BELLADONNÆ, *Ointment of Belladonna*. (*Ext. belladonnæ* ʒi; *adipis* ʒj; *M. Ph. U. S.*) An anodyne application to irritable ulcers, &c.

UNGUENTUM CANTHARIDIS, *Unguentum lyttæ*.

UNGUENTUM CERÆ (*albæ seu flavæ*), *Unguentum simplex* (*Ph. U. S.*), *Linimentum simplex*, *Wax Ointment*, *Oil and Bees' Wax*, (F.) *Onguent de cire*. (*Cerâ lbj*, *adipis lbiv* (*Ph. U. S.*), or — *cerâ ʒij*, *ol. oliv.* ʒv.) Emollient. A mild dressing. Also, the basis of most of the compound ointments.

UNGUENTUM CERATUM, *Ceratum Galeni* — u. *Cerussæ*, *Unguentum oxidi plumbi albi*, *Unguentum plumbi subcarbonatis* — u. *Cerussæ acetatæ*, *Ceratum plumbi superacetatis*, *U. plumbi superacetatis* — u. *Citrinum*, *U. de althæâ*, *U. hydrargyri nitratis*.

UNGUENTUM CETA'CEI, *U. Sper'matis Ceti seu album*, *Linimentum album*, *Spermaceti ointment*, (F.) *Onguent de blanc de baleine*. (*Cetacei ʒvj*, *cerâ albæ ʒiv*, *ol. oliv.* fʒijj. *Ph. L.*) The ordinary dressing for blisters and excoriations. See *Ceratum cetacei*.

UNGUENTUM COCCULI, *Ointment of Cocculus Indicus*. (Take any quantity of *Cocculus Indicus*; separate and preserve the kernels; beat them well in a mortar, first alone, and then with a little lard; then add lard till it amounts altogether to five times the weight of the kernels. *Ph. E.*) Used to destroy vermin, and to cure scabies and ringworm of the scalp.

UNGUENTUM CONII, *Ointment of Hemlock*. (*Conii fol. recent.*; *adipis* 3â lbj. Boil the hemlock in the lard till it becomes crisp, and express through linen. *Ph. L.*) Used to irritable tumours and ulcers.

UNGUENTUM CREASOTI, *Ointment of Creasote*. (*Creasot.* fʒss; *adipis*, ʒj. Add the creasote to the lard previously melted with a moderate heat, and stir constantly till cold. *Ph. U. S.*) Used in chronic cutaneous affections.

UNGUENTUM CUPRI SUBACETATIS, *Unguentum subacetatis cupri* — u. *Diapompholygos*, *Unguentum oxidi zinci impuri*.

UNGUENTUM DIGESTIVUM SIMPLEX, *U. Terebin'thinæ et ovo'rum vitell.*, *Simple digestive ointment*, (F.) *Onguent digestif simple*. (*Terebinthin. pur.* 64 p. *Vitell. ovor.* No. 2, vel 32 p. Mix, for a long time, in a glass mortar, and add, gradually, enough of the Oil of Hypericum (*Millepertuis*) to make a soft ointment. *Ph. P.*) Detergent. It may be rendered more so by adding a little of the Acetate of copper.

UNGUENTUM ELEMII, *Balsam of Arcæus*, *Unguentum elemi compositum*.

UNGUENTUM EL'EMI COMPOSITUM, *Unguentum e gummi elemi seu de terebin'thinâ et adip'ibus seu elemi*, *Linimentum Arcæ'i*, *Ointment of Elemi*, (F.) *Baume d'Arcæus*. (*Elemi lbj*, *terebinthinæ vulgaris ʒx*, *sevi præparati lbij*, *olivæ olei fʒijj*. Melt the elemi with the suet; remove it from the fire; and mix in the turpentine and oil. Strain the mixture through a linen cloth. *Ph. L.*) Stimulant and digestive. To keep open issues, &c.

UNGUENTUM EPISPASTICUM DE DAPHNE GNIPIO, *Adeps Corticis daphnæ gnidi medicatus*, (F.) *Pommade de Garou*. (*Adip. præpar.* 320 p., *cerâ*, 32 p., *cort. daphn. gnid.* 128 p. Melt the fat and wax; and add the bark softened in water. Boil until the water is evaporated; then pass through cloth. *Ph. P.*) To keep open blisters, issues, &c. See *Unguentum Mezæci*.

UNGUENTUM EPISPASTICUM FORTIUS, Cerate of cantharides—u. Epispasticum mitius, Unguentum lyttæ—u. Epispasticum viride, Unguentum lyttæ medicatum—u. Flavum, Unguentum de althæa—u. e Gummi eleui, Unguentum compositum—u. Hellebori albi, Unguentum veratri.

UNGUENTUM GALLÆ, Ointment of Galls, (Gallæ in pulv. ʒj; adipis, ʒviij. M. Ph. U. S.) As an astringent in piles, indolent ulcers, &c.

UNGUENTUM GALLÆ COMPOSITUM, U. Gallæ et Opii, Compound Ointment of Galls. (Gall. in pulv. subtil. ʒvj, adipis. ʒvj, opii pulv. ʒiss. M. Ph. L.) Used in painful hemorrhoids. ʒss or ʒj, of camphor is sometimes added.

UNGUENTUM HYDRARGYRI, U. cæruleum seu Neapolitanum, Cera'tum mercurial'e, Adeps Hydrargyri medicatus, Mercurial Ointment, Blue ointment, Unction, (F.) Onguent mercuriel ou Napolitain.

The Strong Mercurial Ointment, Unguentum Hydrargyri fortius of the London Pharmacopœia, U. Hydrargyri (Ph. U. S.), is thus made—Hydrargyri lbij, adipis ʒxxij, sevi ʒj. Rub till the globules entirely disappear.

The Milder Mercurial Ointment,—U. Hydrargyri Mitius,—is made by adding, to one pound of the strong, two pounds of lard.

Mercurial ointment is antisyphilitic and discutient. It is rubbed on the inside of the thigh, in venereal affections; especially in those cases where mercury, given internally, runs off by the bowels. The stronger ointment is used for this purpose. The weaker is employed as a dressing, or as an application to cutaneous eruptions.

UNGUENTUM HYDRARGYRI AMMONIATI, Unguentum hydrargyri præcipitati albi—u. Hydrargyri calcis albæ, Unguentum hydrargyri præcipitati albi—u. Hydrargyri compositum, Unguentum hydrargyri nitratis—u. Hydrargyri deutoxydi nitrati, Unguentum hydrargyri nitrico-oxydi.

UNGUENTUM HYDRARGYRI IODIDI, Ointment of Iodide of Mer'cury. (Hydrarg. Iodid. ʒj, ceræ alb. ʒij, adipis ʒvj. Ph. L.)

UNGUENTUM HYDRARGYRI IODIDI RUBRI, Ointment of Red Iodide of Mer'cury. (Hydrarg. Iodid. Rubr. ʒj, Ung. ceræ albæ ʒvij. Ph. D.) Both these ointments may be applied to scrofulous and indolent ulcers.

UNGUENTUM EX HYDRARGYRO MITIUS DICTUM CINEREUM, Unguentum oxydi hydrargyri cinereum.

UNGUENTUM HYDRARGYRI NITRATIS, U. hydrargyri nitrati seu supernitratis hydrargyri seu hydrargyri compositum seu mercuriale citrinum seu citrinum ad scabiem, Balsamum mercurial'e, Adeps nitratis hydrargyri medicatus, Ointment of Nitrate of Mer'cury, Citrine Ointment, (F.) Onguent de nitrate de mercure, O. citrin contre la gale. (Hydrarg. ʒj, acid. nitric. f ʒxiv, olei bubuli f ʒix, adipis ʒij. Dissolve the mercury in the acid; and, while hot, add the oil and melted lard. Ph. U. S.)

A milder ointment,—Ung. nitratis Hydrargyri mitius,—is made with triple the quantity of oil and lard. It is stimulant and detergent, and is much used as an application to herpes, porrigo, and cutaneous eruptions. The weaker ointment is sometimes used in ophthalmia tarsi, &c.

UNGUENTUM HYDRARGYRI NITRICO-OXYDI, U. subnitratis hydrargyri seu Oxydi Hydrargyri rubri, U. H. oxydi rubri (Ph. U. S.), U. deutoxydi hydrarg. nitrati seu Hydrargyri rubri seu ex Hydrargyro oxydato rubro seu mercurii rubrum seu ophthalmicum rubrum seu subnitratis hydrargyri, Balsamum ophthalmicum rubrum, Ointment of Nitric Oxyd of Mercury, Red Precipitate Ointment, Golden Ointment, Singleton's Eye Salve. (Hydrarg. oxyd. rubr. in pulv. subtiliss. ʒj; Ung. simpl. ʒj. Add the oxyd, re-

duced to a fine powder, to the melted ointment. Ph. U. S.) Stimulant and escharotic. Applied to foul ulcers, to inflammation of the tarsi, &c. &c.

Belleville's Cerate, which has been long used in cases of scalded head, and of chronic cutaneous diseases in general, is made as follows:—Plumbi acet. ʒj; hydrarg. oxid. rubr. ʒss; hyd. chlorid. mit. ʒij; ceræ albæ ʒiv; ol. oliv. ʒvj. Melt the last two, and add the others in fine powder.

UNGUENTUM HYDRARGYRI OXYDI RUBRI, Unguentum hydrargyri nitrico-oxydi—u. ex Hydrargyro oxydato rubro, Unguentum hydrargyri nitrico-oxydi—u. Hydrargyri oxydi rubri, Unguentum hydrargyri nitrico-oxydi.

UNGUENTUM HYDRARGYRI PRÆCIPITATI ALBI, U. Hydrargyri ammoniat'i, (Ph. U. S.) seu submuriat'is hydrargyri ammoniat'is seu ad scabiem Zelleri seu mercurial'e album seu mundificans Zelleri seu e mercurio præcipitato seu calcis hydrargyri albæ, Ointment of white precipitate. (Hyd. ammon. ʒj, ung. simpl. ʒiss. Melt the fat, and add the precipitate.) Detergent. Used in cutaneous eruptions, to destroy vermin, &c.

UNGUENTUM HYDRARGYRI RUBRI, U. Hydrargyri nitrico-oxydi—u. Hydrargyri submuriat'is ammoniat'i, U. Hydrargyri præcipitati albi—u. Hydrargyri subnitratis, U. hydrargyri nitrico-oxydi—u. Hydrargyri subnitratis, U. hydrargyri nitrico-oxydi—u. Hydrargyri supernitratis, U. hydrargyri nitratis—u. Infusi meloes vesicatorii, U. lyttæ—u. Irritans, U. lyttæ.

UNGUENTUM IODINI, U. Iodin'ii, Ointment of Iodine. (Iodin. ʒj; Potass. iodid. gr. iv; Aquæ Mvi; Adipis. ʒj. Rub the iodine and iodide first with the water until liquefied, and then with the lard until mixed. Ph. U. S.) Used as a local application to goitrous and other swellings.

UNGUENTUM IODINI COMPOSITUM, U. Iodin'ii compositum, Compound Ointment of Iodine. (Iodin ʒss; Potassii iodid. ʒj; Alcohol, f ʒj; adipis ʒij. Rub the iodine and iodide first with the alcohol and then with the lard until they are well mixed. Ph. U. S.)

UNGUENTUM LAURINUM, O'cum laurinum, Adeps lauro-medicatus, Bay Ointment, (F.) Onguent ou Huile de laurier. (Fol. lauri, 500 p., adip. suillæ, 1000 p. Bruise in a marble mortar; evaporate all humidity by boiling; and add Bay berries, in powder, 500 p. Digest and pass through linen. Ph. P.) Reputed to be corroborant and nervine. It is simply emollient.

UNGUENTUM LENITIVUM, U. de althæa—u. Lilaceum, Crinomyron.

UNGUENTUM LYTTÆ, Ung. cantharidis seu vesicatorium seu irritans seu ad vesicata seu infusio meloes vesicatorii seu epispasticum mitius, Blistering Salve, Ointment of Spanish Flies, Ointment of Infusion of Spanish Flies, Milder blistering Ointment. (Cantharid. in pulv. ʒij; aquæ destillat. Oss; cerat. resin. ʒviij. Boil down the water with the Spanish flies to one-half, and strain; then mix the cerate with the strained liquor, and evaporate to the proper consistence. Ph. U. S.) It is stimulant and epispastic; and is chiefly used for keeping up the discharge from a blistered surface.

UNGUENTUM LYTTÆ MEDICATUM, Adeps cantharid'ibus medicatus, Pomatum seu Unguentum epispasticum viride stimulans ac'ribus, Green blistering Ointment, (F.) Pomade épispastique verte. (Pomati seu unguenti cantharid. 64 p., unguent. populei 168 p., ceræ albæ 256 p., cupri acetat., ext. opii, aa 24 p. Melt the unguentum populeum, and wax, and add, as they cool, the oxyd of copper, cantharides, and opium. Ph. P.) Use. same as the last.

UNGUENTUM LYTTÆ MEDICATUM ALIUD, Adeps cantharid'ibus medicatus alius, Pomatum epispasticum flavum et mitius, Yellow Blister.

ing Ointment, (F.) Pomme épispastique jeune. (Pole. cantharid. 120 p., adip. præp. 1680 p., aquæ 250 p. Melt the lard; mix the cantharides, and add the water at the same time; place the whole on a gentle fire; stir continually for two hours, and add a little water to replace that which has been evaporated. Pass through linen and strain. Keep it in a water-bath, and add powdered turmeric, 8 p. Let it cool till the superfluous liquid separates; pour off and add yellow wax, 250 p., Oil of Lemon, 8 p. Ph. P.) A complex preparation. Use, same as the last.

An ointment, not very philosophically combined, was used by Dr. Physick, which is known by the name of "*Dr. Physick's Issue Ointment.*" It is made by boiling half an ounce of powdered cantharides in two ounces of rose-water, and evaporating until one-half of the fluid is driven off, having previously added 15 grains of tartar emetic. The decoction is to be strained, and three ounces of olive oil, an ounce and a half of white wax, and one ounce of spermaceti added to it, and the mixture then boiled till all the water is evaporated.

UNGUENTUM MELANICUM CAUSTICUM, Causticum Æthiopicum.

UNGUENTUM MERCURIALE ALBUM, Unguentum hydrargyri præcipitati albi—u. Mercuriale citrinum, Unguentum hydrargyri nitratis—u. e Mercurio præcipitato, Unguentum hydrargyri præcipitati albi—u. Mercurii rubrum, Unguentum hydrargyri nitrico-oxidi—u. Mundificans Zelleri, Unguentum hydrargyri præcipitati albi.

UNGUENTUM MEZE'REI, *Ointment of Mezereon. (Mezer. incis. transversum, 3iv; adipis, 3xiv; ceræ albæ, 3ij; moisten the mezereon with a little alcohol, and beat in an iron mortar, until reduced to a fibrous mass; then digest it with the lard, in a salt-water bath, for 12 hours; strain forcibly, and allow the strained liquor to cool slowly, so that any undissolved matters may subside; from these separate the medicated lard; melt it with the wax at a moderate heat, and stir constantly till they are cold. Ph. U. S.) To keep open blisters, issues, &c.*

UNGUENTUM MURIA'TIS HYDRARGYRI OXYGENA'TO MEDICA'TUM, *Adeps muriatæ hydrargyri oxygenatæ medicatæ, Cyrillo's Ointment, (F.) Pomme de Cyrillo. (Hydrar. oxyzmur. 4 p.; adipis suilli præp. 32 p.—Ph. P.) Antivenereal and alterative. Used in friction, 3ss at a time.*

UNGUENTUM NARDINUM, see Nardus Indica—u. Neapolitanum, Unguentum hydrargyri—u. Nervinum, Unguentum Sambuci—u. Nitricum seu oxygenatum, Unguentum acidi nitrosi.

UNGUENTUM OPHTHALMICUM, *Adeps oxido hydrargyri rubro et plumbi acetatæ medicatus, Regent's Pomme ophthalmique, (F.) Pomme ophthalmique. (Butyr. recent., aq. rosar. lavat. aa 4½ p.; camphor., oxyd. hydrarg. rubr., acetatis plumbi, aa 0.25 p. Mix intimately.—Ph. P.) In chronic inflammation of the conjunctiva or tarsi.*

UNGUENTUM OPHTHALMICUM, Unguentum oxidi zinci impuri—u. Ophthalmicum rubrum, Unguentum hydrargyri nitrico-oxidi.

UNGUENTUM OPII, *Ointment of Opium. (Opii pulv. 3j; adipis 3i; M.—Ph. L.) To be used where the local application of opium by inunction is needed.*

UNGUENTUM OX'IDI HYDRARGYRI CINE'REI, *Adeps seu Unguentum ex hydrargyro mitius dictum cinereum, Ointment of gray oxide of mercury, (F.) Onguent gris. (Oxyd. hydrarg. ciner. p. j; adipis suilli, p. iij.—Ph. E.) Properties like the mercurial ointment. It is not much used.*

UNGUENTUM OX'IDI PLUMBI ALBI, *U. cerusæ seu album seu subacetatis plumbi, Ointment of white oxide of lead, White ointment. (Ung. simplicis, p. v; oxyd. plumb. alb. p. j.—Ph. E. and*

P.) Cooling and desiccative. Applied to excoriated and burnt surfaces.

UNGUENTUM OXIDI PLUMBI CARBONATI, U. Plumbi subcarbonatis—u. Oxidi zinci, Unguentum zinci.

UNGUENTUM OX'IDI ZINCI IMP'URI, *U. tinctæ seu ophthalmicum seu diopompholygus, Adeps oxidi zinci medicatus, Eye-salve, Tatty ointment. Ointment of impure oxide of zinc. (Linn. simpl. p. v; oxyd. zinci. imp. præp. p. j.—Ph. E.) Generally used in ophthalmia tarsi and atonic ophthalmia.*

UNGUENTUM PICIS LIQ'UIDÆ, U. Picis seu e Picæ, Tar Ointment. (Picis liquid., sevi, sing. ʒij. Melt, and stir till cold.—Ph. U. S.) Stimulant and detergent. Used in tinea capitis, and other cutaneous eruptions.

UNGUENTUM PICIS NIGRÆ, Unguentum resinæ nigre—u. Plumbi acetatis, Ceratum plumbi superacetatis—u. Plumbi carbonatis, Unguentum plumbi subcarbonatis.

UNGUENTUM PLUMBI COMPOS'ITUM, Compound Ointment of Lead. (Emp. Plumbi, ʒij; ol. oliv. f 3xvij; cret. præpar. 3vj; acid. acet. dil. f 3vj. Dissolve the plaster in the oil with a slow fire; then add the chalk, and afterwards the acid, constantly stirring until they become cold.—Ph. L.) Used as a dressing to flabby, indolent ulcers.

UNGUENTUM PLUMBI IOD'IDI, *Ointment of Iodide of Lead. (Plumb. iodid. 3j; adipis, 3vij; M.—Ph. L.) Used as a eutrophic in chronic, glandular, and other swellings.*

UNGUENTUM PLUMBI OXYDI CARBONATI, Unguentum plumbi subcarbonatis—u. Plumbi subacetatis, Unguentum oxidi plumbi albi—u. Plumbici carbonatis, Unguentum plumbi subcarbonatis.

UNGUENTUM PLUMBI SUBCARBONA'TIS, U. plumbi carbonat'is (Ph. U. S.) seu album seu album simplex seu carbonat'is plumbici seu cerusæ seu oxidi plumbi carbonat'is, *Cera'tum de cerusæ, (F.) Onguent blanc de Rhazes, O. de cêruse, O. de Tornamira. (Plumbi carbonat. in pulv. subtiliss. 3ij; Unguent. simpl. ʒij. Add the carbonate of lead to the ointment previously softened over a gentle fire, and mix.—Ph. U. S.) Chiefly used as a dressing for burns.*

UNGUENTUM PLUMBI SUPERACETA'TIS, U. acetat'is plumbi seu saturni'num seu cerusæ acetatæ, *Balsamum universa'le, Ointment of sugar of lead. (Olive oil. ʒss; white wax, 3iss; superacetate of lead, 3ij.) Cooling, desiccative. Called, also, Balsamum saturni'num, Butyrum saturni, Linimentum plumba'tum, Poma'tum saturni, Ung. triphar'macum.*

UNGUENTUM POPULE'UM, *Adeps papav'esh, hyoscy'amo, et belladon'næ medicatus.* Fresh buds of the populus nigra, 500 p., macerated in adeps suillus præp. melted, 1500 p., for 24 hours. Preserve this. Take black poppy leaves, belladonna, hyoscyamus, black nightshade, aa 128 p.; bruise in a mortar with the former preparation; boil until all the humidity is evaporated, and strain.—Ph. P.) Anodyne. Applied to local inflammations, hemorrhoids, sore nipples, cancer, &c. Green elder ointment is often sold for this.

UNGUENTUM POPEL'EUM, see Populus.

UNGUENTUM POTAS'SÆ HYDRIODA'TIS, U. potas'sii iod'idi (Ph. U. S.), *Ointment of hydriodate of potass. (Potass. iodid. in pulv. subtil. 3i; aq. bullient. f 3i; adipis, 3j. Dissolve the iodide in the water, and mix the solution with the lard.—Ph. U. S.) Half a drachm of this ointment is rubbed on scrofulous glands, goitre, &c.*

UNGUENTUM POTASSII IODIDI, Unguentum Potassæ Hydriodatis—u. Pulveris melocæ vesicatorii, Cerate of cantharides—u. Resinæ flavæ, Ceratum resinæ flavæ.

UNGUENTUM RESINÆ NIGRÆ, U. resinæ nigre seu basilicon nigrum seu Picis nigre seu Picis

tetrapharm'acum, Black basilicon, Black resin ointment, Ointment of pitch, Ointment of black pitch. (Resin nigr., ceræ flavæ, resinæ flav. 3i; 3i; olei olivæ, Oj.—Ph. L.) Uses, the same as the Ceratum resinæ or Yellow basilicon.

UNGUENTUM RESINÆ PINI SYLVESTRIUS COMPOSITUM, Unguentum de Althæa—u. Resinosum, Ceratum resinæ flavæ—u. Resumptivum, Unguentum de althæa.

UNGUENTUM SABINÆ. Ceratum Sabinæ.

UNGUENTUM SAMBU'CI, U. sambucinum seu nervi'num, Green ointment, Elder ointment, (F.) Onguent de sureau. (Sambuci flor. ʒij; adipis præp. ʒij. Boil.—Ph. L.) Emollient.

UNGUENTUM SATURNINUM, Unguentum plumbi superacetatis—u. ad Scabiem Zelleri, Unguentum hydrargyri præcipitati albi—u. Simplex, Ceratum Galeni, Unguentum ceræ—u. Solidum de cicuta, Emplastrum cicutæ—u. Spermatidis ceti, Unguentum cetacei.

UNGUENTUM STRAMO'NII, Ointment of Stramonium. (Ext. stramon. fol. 3j; adipis, 3i. Rub the extract with a little water until uniformly soft, and then with the lard.—Ph. U. S.) An anodyne ointment to irritable ulcers, hemorrhoids, &c.

UNGUENTUM SUBACETA'TIS CUPRI, U. cupri subaceta'tis (Ph. U. S.) seu deter'gens seu basilicum vir'ide seu æru'ginis, Balsamum viridæ, Ointment of subacetate of copper, Ver'digris ointment. (Cupri subacet. in pulv. subtiliss. 3j; ung. simpl. 3xv. Add the subacetate to the ointment previously melted with a moderate heat, and stir till cold.—Ph. U. S.) Detergent and escharotic.

UNGUENTUM SULPHURATUM AD SCA'BIE' M, Adeps sulphuræ et ammo'niæ muria'tæ medica'tus. (Adipis, 120 p.; sulph. lot. 60 p.; ammoniæ muriat., aluminæ sulphat. 3a 4 p.; fiat unguentum.—Ph. P.) Antipsoric.

Bailey's Itch Ointment is composed of olive oil, lard, nitre, alum, sulphate of zinc, and cinnabar; scented with oil of aniseed, oil of origanum, and oil of spike, and coloured with alkane-root.

UNGUENTUM SULPHURATUM ALCALI'NUM AD SCA'BIE' M, Adeps sulphuræ et carbona'tæ potassæ medica'tus, Hel'merick's Alkaline Sulphurated Itch Ointment. (Adip. præp. 800 p.; sulphur. lot. 200 p.; carbonat. potass. pur. 100 p.; fiat ung.—Ph. P.) Use, same as the last.

Bateman's Itch Ointment resembles this. It is coloured with cinnabar, and scented with essence of bergamot.

UNGUENTUM SUL'PHURIS, U. e sulphuræ, Sulphur ointment. (Sulph. ʒij; adip. ʒij.—Ph. U. S.) Used chiefly in itch, and some herpetic affections.

UNGUENTUM SULPHURIS COMPOSITUM, U. Antipsoricum, Compound Sulphur Ointment, Itch Ointment, Jackson's Itch Ointment, Pommade Antipsorique de Jasser. (Sulph. sublim. lbss; veratri rad. contus. 3ij; potassæ nitrat. 3j; saponis molliis, lbss; adipis, lbss; ol. bergamot. Mxxx.—Ph. L.) The UNGUENTUM SULPHURIS COMPOSITUM of the Pharm. U. S. is made as follows:—Sulphur. 3j; hydrargyri ammoniat., acid. benzoic. 3a 3j; ol. bergamot., acid. sulphuric. 3a f3j; potassæ nitrat. 3ij; adipis, lbss. To the melted lard add the other ingredients, and stir till cold. Use, the same as the last; but it is more stimulating.

UNGUENTUM SULPHURIS IOD'IDI, Ointment of Podide of Sulphur. (Sulphur. iodic. 3ss; adipis, 3j. Rub the iodide with a little of the lard; add the remainder, and mix.—Ph. U. S.) An excellent application in chronic cutaneous affections.

UNGUENTUM SUSINUM, Crinomyron.

UNGUENTUM TAB'ACI, Tobacco Ointment. (Tabac. recent. concis. 3j; adipis, ʒij. Boil the to-

bacco in the lard, over a gentle fire, till it becomes friable; then strain through linen.—Ph. U. S.) Applied to irritable ulcers, tinea capitis, &c.

UNGUENTUM TARTARI EMETICI, U. antimoni tartarizati—u. Tartari stibiat, U. antimoni tartarizati—u. e Tartaro potassæ stibiat, U. antimoni tartarizati—u. Tartratis potassæ stibiat, U. antimoni tartarizati—u. de Terebinthinæ et adipibus, U. elemi compositum—u. de Terebinthinæ et ceræ, U. de althæa—u. Terebinthinæ et ovorum vitelli, U. digestivum simplex—u. Tetrapharmacum, Basilicon, U. resinæ nigræ—u. Tripharmacum, U. plumbi superacetatis—u. Tutia, U. oxidæ zinci impuri.

UNGUENTUM VERA'TRI, U. Veratri albi (Ph. U. S.) seu helleb'ori albi, White hellebore ointment. (Veratri. pulv. 3ij; adipis, 3viij; olei limon. Mxx.) Used in cases of itch, where sulphur is objected to. It is not equally efficacious.

The principal ingredients in the EDINBURGH OINTMENT are white hellebore and muriate of ammonia.

UNGUENTUM AD VESICATA, U. lyttæ—u. ad Vesicatoria, Cerate of cantharides—u. Vesicatorium, Unguentum lyttæ.

UNGUENTUM ZINCI, U. ox'idi zinci seu zinci oxidi (Ph. U. S.), Zinc ointment. (Zinci oxid. 3j; adipis, 3vj.) Use, the same as that of the Unguentum tutiæ.

UNGUENTUM ZINCI OXIDI, U. zinci.

UNGUIS, Hippocampus minor, Nail, Onyx, Pterygion—u. Avis, Hippocampus minor—u. Halleri, Hippocampus minor—u. Odoratus, Blatta Byzantia.

UNGUIS OS, Os lacryma'le. A small, quadrilateral, very thin, and semi-transparent bone, which has been compared to a human nail, and is situate at the anterior and inner part of the orbit. It aids in the formation of the lachrymal gutter and the nasal duct. It is articulated with the frontal, superior maxillary, ethmoid, and inferior spongy bones, and is developed by a single punctum ossificationis.

UNGULA CABALLINA, Tussilago.

UNGULÆ PALPEBRARUM, see Tarsus.

UNHEALTHY, Sickly.

UNICORN ROOT, Chamælerium luteum.

UNILOC'ULAR, Unilocularis, from unus, 'one,' and locus, (diminutive of locus,) 'a cell.' Having one cell or cavity, as biloc'ular means having two cells or cavities; multiloc'ular, having many.

UNIO. Pearl.

UNIOCLUSUS, Cyclops, Monoculus.

UNION BY THE FIRST AND SECOND INTENTIONS, see Intention.

UNIPARIENS, Uniparous.

UNIPARIENT, Uniparous.

UNIP'AROUS, Unipa'rient, Unipariens, from unus, 'one,' and pario, 'I bring forth.' Producing one at a birth.

UNISE'MA DELTIFO'LIA, Pontede'ria corda'ta, Shovel Pickereaved, Pond Shovel, Shovel-leaf, Water Plantain. An indigenous plant, Ord. Pontederiaceæ, Sex. Syst. Hexandria Monogynia, which grows in water, and flowers from June to August. The root is emollient and astringent.

UNITAIRES (MONSTRES) (F.), from unitas, 'oneness.' A class of monsters in which are found only the complete or incomplete elements of a single individual. In contradistinction to Compound monsters, (F.) Monstres composés, in which there are the complete or incomplete elements of more than one.

UNITING BANDAGE, (F.) Bandage unissant. A bandage for bringing the edges of a wound into approximation. The arrangement

of such a bandage differs according as the wound is transverse or longitudinal.

UNIVOCAL GENERATION, see Generation.

UNOCULUS, *Borgne*. Cyclops, Monoculus.

UNORGANIZED, Inorganic.

UNSTRIPED MUSCULAR FIBRE, see Muscular fibre.

UNWELL, see Menstruation.

UPAS. A word used in the Archipelagos of Molucca and Sunda, and signifying *Vegetable Poison*, *Ipo*, *Bohon*, *Pohon*, or *Boa* or *Boon Upas*, *Upas*, *Pohon Antiar*, *Antjar*. A substance used by the natives of those places for poisoning their arrows. The smallest quantity is capable of destroying the largest animal. The tree which produces it is said to be the *Antiaris Toxicaria*. Another species of upas, *U. Tieuté*, *Tshettik* or *Tjatch*, is the produce of a *Stychnos*, *S. Tieuté*. Many idle stories are told regarding the upas, most of which are fabulous.

URACHUS, *Uracus*, *Urinaculum*, from *ουρον*, 'urine,' and *ἔχω*, 'I contain.' (F.) *Ouarque*. The urachus, in certain animals, is a long membranous canal, which arises from the bladder, makes its exit from the abdomen by the umbilicus, and terminates in the pouch called *allantois*. According to Sabatier, this is the arrangement of the urachus in the human foetus; but Bichat and other anatomists consider, that when the canal exists in it, it is a malformation. They regard it as a kind of suspensory ligament of the bladder, extending from the top of that organ to the umbilicus; and, at the umbilicus, conjoined with the abdominal aponeurosis.

URACRATIA, Enuresis.

URACUS, Urachus.

URÆMIA, *Urhæmia*, (F.) *Urémie*, from *ουρον*, 'urine,' and *αἷμα*, 'blood.' A condition of the blood in which it contains urine or urea.

URALI, Curare.

URANA, Ureter.

URANE, Urinal.

URANISCONTITIS, from *ουρανισκος*, 'the palate,' and *itis*, denoting inflammation. *Palatit'is*. Inflammation of the palate.

URANISCOPLAS'TY, *Uraniscoplas'ticē*, from *ουρανισκος*, 'the palate,' and *πλαστικός*, 'forming.' The operation of engrafting in case of deficiency of the soft palate.

URANISCORRHAPHIA, Staphylorrhaphy.

URANISCOS, Palate.

URANORRHAPHIA, Staphylorrhaphy.

URAPOSTEMA, *Abscessus urinae*, from *ὑράδεμα*, from *ουρον*, 'urine,' and *αποστῆμα*, 'an abscess.' A tumefaction produced by the escape of urine. An abscess containing urine.

URARI, Curare.

URAS, Urate.

URATE, *Uras*, *Lithate*, *Lithas*; from *ουρον*, 'urine.' A generic name for salts, formed by the combination of uric or lithic acid with different bases. *Urate* or *Lithate of Soda*, for example, is found in arthritic and certain urinary concretions:—*Urate* or *Lithate of Ammonia* in certain urinary deposits.

URÆA, *Ureu*, *Urenium*, *Uricum*, *Materia urinae*, *Principium seu Sal urinae*, *Nephrie*, (F.) *Urée*. Same etymon. A white, pearly, brilliant substance, in transparent plates, inodorous, and of a cool taste; very soluble in water; deliquescent, and soluble in alcohol. It exists, in great quantity, in the urine; and is an essential part of it. It has been used as a diuretic. Dose, gr. xv to ʒj.

UREC'CHYSIS, *Urech'ysis*, from *ουρον*, 'urine,' and *ἔκχυσις*, 'effusion.' Effusion of urine into the cellular membrane.

UREDIO, Urticaria.

URÉE, Urea.

UREMA. Urine.

URÆMIE, Uræmia.

UREN, Urea.

URENIUM, Urea.

URENS, Caustic.

UREORRHŒA, Diabetes.

URES-ESTHESIS, Diuresæsthesis.

URESIS, Micturition.

URE'TER, *ουρητηρ*. Same etymon as *Urate*, *U'raua*, *Vena alba renium*, *Canalis nervus fistulæ*, *urenum*, *Ductus seu Ductor Uri'nae*, *Vas urinarium*, (F.) *Uretère*. A long, excretory canal—membranous and cylindrical—which conveys the urine from the kidney to the bladder. It extends, obliquely, between the pelvis of the kidney, with which it is continuous, and the *bas fond* of the bladder, into which it opens. It enters at the posterior and inferior part, for some distance between the coats; and opens into the cavity of the bladder, at the posterior angle of the *Trigone vésical*. The ureter is formed of an exterior, white, opaque, and fibrous membrane; and of an inner, which is mucous.

URETERAL'GIA, from *ουρητηρ*, 'the ureter,' and *αλγος*, 'pain.' Pain in the course of the ureter,—as from calculus, *Ureteral'gia calculosa*.

URETERALGIA CALCULOSA, see Ureteralgia—u. Inflammatoria, Ureteritis.

URETÈRE, Ureter.

URETER'ICUS. An epithet for ischury, &c., arising from an affection of the ureter.

URETERITIS, *Ureteral'gia inflammatoria*, *Ureterophlegmasia*, from *ureter*, and *itis*, denoting inflammation. Inflammation of the ureter; generally caused by calculus.

URETERO-LITH'ICUS, from *ureter*, and *λίθος*, 'a stone.' An epithet applied to ischury or other affection caused by a stone in the ureter.

URETEROL'ITHUS. Same etymon. Calculus in the ureter.

URETEROPHLEGMASIA, Ureteritis.

URETERO-PHLEGMAT'ICUS, from *ureter*, and *φlegμα*, 'phlegm or mucus.' Any affection caused by mucus accumulated in the ureter.

URETERO-PY'ICUS, from *ureter*, and *πυον*, 'pus.' An epithet applied to affections caused by the presence of pus in the ureter.

URETERORRHAG'IA, from *ureter*, and *ρῆμα*, 'rupture.' Hemorrhage from the ureter.

URETERO-STOMAT'ICUS, from *ureter*, and *στομα*, 'orifice.' That which is dependent upon obstruction of the orifice of the ureter.

URETERO-THROMBOÏ'DES, from *ureter*, *θρομβος*, 'grumous blood,' and *εἶδος*, 'resemblance.' Ischury from grumous blood in the ureter.

URE'THRA, *ουρηθρα*, *Canalis seu Ductus urinae*, *Urias*, *Ure'tra*, *Fistula urinaria*, *Iter urinarium* seu *Uri'nae*; (F.) *Urèthre*, *Urètre*, from *ουρον*, 'urine.' The excretory ducts for the urine in both sexes; and for the sperm in man. In the latter, it is 8 or 10 inches long; and extends from the neck of the bladder to the extremity of the penis, in which its external aperture is situate. In the male urethra, three portions are distinguished:—1. A *Prostatic portion*, near the bladder, about an inch and a quarter in length, which traverses the prostate gland:—2. A *membranous portion*, *Isthmus ure'thræ*, *Para ure'thræ membranacea*, from 8 to 10 lines long, which is separated from the rectum by the areolar tissue only:—3. A *spongy portion*, which expands, anteriorly, to form the glans, and which commences at the *bulb of the urethra*. Before opening externally, the canal has a dilatation, called the *Fossa* or *Fossicula navicularis*. On the interior of the urethra two white lines are perceptible—the one above, the other below. The latter commences at the *verumontanum*; and

at the surface of this, the *ejaculatory ducts*, and *prostatic ducts* open. The urethra is lined, in its whole extent, by a mucous membrane; on which are seen, particularly at the bulb, the orifices of a number of follicular ducts, *Lacunæ* or *Sinuses of Morgagni*. At about an inch and a half from the opening of the meatus, one of these lacunæ is generally found, much larger than the rest, and is named *Lacuna Magna*. The arteries of the urethra are numerous, and proceed from the internal pudic;—the largest branches entering the bulb. The veins follow the course of the arteries. The lymphatics pass into the inguinal and hypogastric plexuses. The nerves come from the pudic and gluteal. In the *female*, the urethra is only about an inch long. It is wider than in man, and susceptible of great dilatation. It is very wide at its commencement; and descends obliquely forwards, to terminate at the top of the external aperture of the vagina by the *meatus urinarius*. In its course, it is slightly curved, the concavity being forwards.

URETHRA, BLIND DUCTS OF THE, see Cæcus—u. Mucous lacunæ of the, see Cæcus.

URETHRAL, *Urethral'is*, *Ure'thricus*. Relating to the urethra.

URETHRAL'GIA, from *urethra*, and *algos*, 'pain.' Pain in the urethra. Gonorrhœa.

URÊTHRE, Urethra.

URETHREMPHRAXIS, Urethrophraxia.

URETHRITIS, Gonorrhœa.

URETHROBLENNORRHEA, from *urethra*, *βλένω*, 'mucus,' and *ρῆω*, 'I flow.' Gonorrhœa; gleet.

URETHROCYSTAUCHENOTOMIA, see Lithotomy.

URETHROCYSTEOANEURYSMATOTOMIA, see Lithotomy.

URETHROCYSTEOTRACHELOTOMIA, see Lithotomy.

URETHROHÆMORRHAGIA, Urethrorrhagia.

URETHROPHRAX'IS, *Urethremphrax'is*, from *urethra*, and *φρασσαω*, 'I obstruct.' Obstruction of the urethra.

URETHROPLAS'TIC, *Urethroplas'ticus*; from *urethra*, and *πλασσαω*, 'I form.' An epithet given to the operation, *U'rethroplasty*, which supplies defects in the canal of the urethra through which the urine escapes.

URETHRORRHA'GIA, *Urethrorhamorrh'gia*, *Caulorrh'gia* seu *Hæmatu'ria stillatit'ia*, from *urethra*, and *ρῆω*, 'I break out.' Hemorrhage from the urethra.—Frank.

URETHRORRHŒA, from *urethra*, and *ρῆω*, 'I flow.' A discharge of any kind from the urethra.

URETHROSTENO'SIS, *Strictu'ra ure'thræ*, *Urostenosis*, *Urethrosten'ia*, (F.) *Rétrécissement de l'Ure'thre*, from *urethra*, and *στενωω*, 'narrowness.' Stricture of the urethra.

URETHROTOME, *Urethrot'omus*, (F.) *Uréthrotome*, from *urethra*, and *τεμνω*, 'I cut.' An instrument so called by Le Cat, with which he divided the skin, and laid open the urethra, in the operation for lithotomy. Also, an instrument for dividing strictures of the urethra.

URETHROTOMY, *Urethrotom'ia*; same etymon. An operation, which consists in opening the urethra with the *Urethrotome*, for the removal of stricture. See *Boutonnière*.

URETHRYMENO'DES, from *urethra*, and *μῆνις*, 'a membrane.' That which is caused by a membrane formed in the urethra.

URETICUS, Diuretic, Urinary.

URETRA, Urethra.

URÊTRE, Urethra.

URETRIS, Urinal.

URGINEA SCILLA, Scilla.

URHEMIA, Uremia.

URIAS, Fistula (urinary), Urethra.

URIASIS, Lithia, Uresis.

URIC ACID, *Ac'idum ur'icum* seu *uroli'ticum* seu *lith'icum*, *Lith'ic*, *Ury'lic*, *Cal'cidous*, *Bezoär'dic*, *Lithia'sic* or *Lithen'ic Acid*, (F.) *Acide urique*. An acid which exists in human urine, chiefly, if not wholly, in the form of urate of soda or urate of ammonia, and which in combination constitutes the greater part of urinary calculi. It is often deposited from the urine in health. See Calculi, urinary.

URIC OXIDE, *Ox'idum ur'icum*, *Xanthic oxide*, *Xanthine*, *Urous acid*. A substance not discovered in healthy urine, although probably bearing some relation to the yellow colouring matter. It is a very rare ingredient of calculous concretions, and little is known either of its chemical or pathological history.

URICUM, Urea.

URICUS, Urinary.

URIDRO'SIS, from *ουρον*, 'urine,' and *ιδρωσ*, 'sweat.' Urinous sweating. (F.) *Sueur urineux*.

URINA CEREBELLA, see Cerebella—u. Chyli, see Urine—u. Cibi, see Urine—u. Diabetica, Urine, diabetic—u. Febrilis, Urine, febrile—u. Furfuracea, see Furfuraceous—u. Galactodes, see Galactodes—u. Genitalis, Sperm—u. Jumentosa seu Jumentaria, Urine, jumentous—u. Mucosa, Cystirrhœa—u. Nephelodes, see Nepheloid—u. Oroboides seu Orobodes, see Oroboides—u. Pericardii, see Pericardium—u. Potius, see Urine—u. Sanguinis, see Urine—u. Vaccæ, Urine, cows'.

URINACULUM, Urachus, Urinal.

URINÆ PROFLUVIUM, Diabets—u. Suppressio, Ischuria—u. Stillicidium, Strangury—u. Substillum, Strangury.

URINAIRE, Urinary.

URINAL, *Urinato'rium*, *Chernib'ium*, *Uranê*, *Ure'tris*, *Urinac'ulum*, *Urodoch'um*, *Urod'ochê*, *Urodoch'um*, *Uratra'cium*, *Matra'cium*, *Mat'ula*, *Matel'la*; from *urina*, 'urine.' A name given, 1. To certain vessels, used to pass the urine into. 2. To reservoirs, of different shapes and character, which are adapted to the penis, in cases of incontinence of urine, and which are intended to receive the urine as it flows off.

URINALIS, Diuretic, Urinary.

URINARIA, Antirrhinum linaria, Leontodon taraxacum.

URINARY, *Urina'rius*, *Urina'lis*, *Urino'sus*, *U'ricus*, *Ure'ticus*, (F.) *Urinaire*. Same etymon. That which relates to the urine.

URINARY BLADDER, *Bladder*, *Ves'ica Urina'ria*, *Cystis*, *Uracys'tis*, (F.) *Vessie*. A musculo-membranous reservoir, intended for the reception of the urine, until the accumulation of a certain quantity solicits its excretion. The bladder is situate in the hypogastric region, between the rectum and pubis in man; and between the pubis and vagina in the female. Its upper region is not covered, except by the peritoneum, posteriorly. From its centre arises the *urachus*. Its lower region is bounded, anteriorly, by the prostate, in man; and rests on the vesicula seminales and rectum. In the female, it corresponds to the vagina. The lowest portion of this region bears the name *Bas-fond*. The anterior surface of the organ is not covered by peritoneum. It is attached to the symphysis pubis by a fibro-cellular expansion called the *anterior ligament*. The *posterior ligaments* or *recto-vesical folds—plicæ semilunares*—are duplicatures of the peritoneum on its posterior surface, which extend from this surface to the rectum in the male, and to the uterus in the female. The bladder is composed of a serous membrane, formed by the peri-

toneum;—of a muscular membrane; of a somewhat thick layer of areolar tissue; and is lined on its inner surface by a mucous membrane, which is sometimes raised up, here and there, in long prominences or *columns* by the muscular fibres beneath. To this condition, the French anatomists give the term—*vessie à colonnes*. In the *bas-fond* the two ureters open, and the urethra commences; and the triangular space, which the apertures of these three canals leave between them, is called, by the French, the *Trigone vésical* or *Vesical Triangle*, *Trigo'nus* seu *Trigo'num vesi'cæ* seu *Lieutau'di*, *Corpus trigo'num*. In the orifice of the urethra, which is also called the neck of the bladder, there is, at its lower part, a more or less prominent tubercle, which is formed by the anterior angle of the *Trigone vésical*, and is called the *Vesical uvula*, *U'vula vesi'cæ*, (F.) *Luette vésicale*.

The arteries of the bladder proceed from the hypogastric, umbilical, sciatic, middle hemorrhoidal, and internal pudic. Its veins, which are more numerous than the arteries, open into the hypogastric venous plexus. Its nerves emanate from the sciatic and hypogastric plexuses; and its lymphatic vessels pass into the hypogastric ganglia.

URINARY CALCULI. The calculi which form in the urinary passages. See Calculi, urinary.

URINARY PASSAGES, (F.) *Voces urinariæ*. The aggregate of canals and cavities intended to contain the urine and convey it externally. These *ways* are composed of excretory ducts, which form the tubular portion of the kidney; of calices; pelvis; ureters; bladder; and urethra.

URINATION, Micturition. M. Béraud elaborately defines it—a function, the result of which is the expulsion of matters principally solid, but held in solution, which have become improper for nutrition.

URINATORIUM, Urinal.

URINE, *Uri'na*, *ουρον*, *Uron*, *Ure'ma*, *Lot'ium*, *Lot*, *Lapis aureus*, *Omic'ma*, *Recremen'ta* seu *Retrimen'ta vesi'cæ*, *Aqua pedum*, *Aqua*, (Old Eng.) *Stale*, *Lee*, (Prov.) *Herrin*, *Netting*, *Saag*. An excremental fluid; secreted by the cortical part of the kidney; filtered through the tubular portion; poured *guttatim* from the apices of the tubular papillæ into the pelvis of the kidney; and transmitted by it to the ureters; which convey it slowly, but in a continuous manner, into the bladder, where it remains deposited, until its accumulation excites a desire to void it. The excretion of the fluid takes place through the urethra; and is caused by the action of the abdominal muscles and diaphragm and the contraction of the fibrous coat of the bladder. Urine is transparent; of a citron-yellow colour; of a peculiar odour, and of an acid, saline, and slightly bitter taste. That which is passed some time after taking fluid, is less coloured, and less odorous and dense than that which is voided 7 or 8 hours after eating. The first is the *Uri'na potûs*, (F.) *Urine de la Boisson*, *Urine of drink*;—the latter *Uri'na san'guinis*, (F.) *Urine de la digestion*, *Urine de la coction*, *Urine cuite*, *Cocted urine*, *Urine of digestion*. Three distinct varieties of urine may, according to Dr. Golding Bird, be recognised: *First*, That passed some little time after drinking freely of fluids, which is generally pale, and of low specific gravity—1.003 to 1.009—*Urina potûs*. *Secondly*, That secreted after the digestion of a full meal; s. g. 1.020 to 1.028, or even 1.030—*Urina chyli* seu *cibi*. *Thirdly*, That secreted independently of the immediate stimulus of food and drink, as after a night's rest, which is usually of average density—1.015 to 1.025, and presents the essential characters of urine—*Urina sanguinea*.

The following are the ordinary constituents of healthy human urine:—urea, uric acid, [hippuric acid;] extractive matters, embracing alcohol extract, spirit extract, and water extract, with their respective constituents; mucus; brown colouring matter of the urine, (hæmaphæin;) red colouring matter of the urine, (uroerythrin;) carbonic, lactic, hydrochloric, sulphuric, phosphoric, silicic, and hydrofluoric acids; soda; potassa; ammonia; lime; magnesia; and peroxide of iron. The proportion of solid matters in 1000 parts of urine may vary from 20 to 70. The following table expresses the relative amounts of the different components in every 100 parts of the solid matter, according to the analysis of different observers:

	Berzelius.	Lehmann.	Simon.	Marchand.
Urea,	45.10	49.68	33.80	48.91
Uric Acid,	1.50	1.61	1.40	1.59
Extractive matter.				
Ammonia. Salts,				
and Chloride of				
Sodium,	36.30	28.95	42.60	32.49
Alkaline Sulphates,	10.30	11.58	8.14	10.18
Alkaline Phosphates,	6.88	5.96	6.50	4.57
Phosphates of lime				
and magnesia,	1.50	1.97	1.59	1.81

Its average specific gravity is about 1.015. The quantity passed by the adult in the 24 hours varies. On the average it may amount to two pounds and a half. Sometimes, the urine, in health and disease, exhibits appearances that have received special names. Thus, that which is characterized by a deficiency of solid matters generally, has been called *Hydrû'ria*, ('vôap, 'water;') by a deficiency of urea, *Anazotû'ria*, (av, privative, and azote;) by a superabundance of urea, *Azotû'ria*. Urine in which the deposit consists of lithic acid, and the lithates, has been termed *Lithû'ria*; and that in which the deposit consists of the earthy and earthy-alkaline phosphates, *Phosphû'ria*, *Ceramû'ria*, (κεραμος, 'potters' earth,') &c. &c. At times, a *pellicle* forms on the surface (*Cremor urinæ*) which consists of mucus and some salts: at others, there is a *cloud*, (*Nubec'ula*, *nubes*,) towards its upper part, which is called *Enaore'ma* when it is suspended, as it were, about the middle of the liquid. Occasionally, there is a sediment, to which the names *Hypotasia*, *Sedimen'tum*, &c., have been given. (See these various words.) The urine is often deserving of attention in disease; but it has been considered a matter of more importance than it merits. Some empirics have presumed to pronounce on the nature of disease by simply inspecting it. See Uroscopy.

Human urine was, at one time, considered aperient; and was given in jaundice, in the dose of one or two ounces. Cows' urine, *Urina Vaccæ*, *Allflower water*, was once used, warm from the cow, as a purge, in the dose of Oss.

Urine has been variously designated by pathologists, according to its appearance, &c., for example:—

URINE, ANÆMIC, *Uri'na anæ'mica*. A term used, but not felicitously, for the form of urine that occurs in anæmia, chlorosis, &c.—*Decquerel*.

URINE ARDENTE. Ardent urine—u. Bloody. Hæmaturia—u. de la Boisson, Urine, hysterical, see Urine—u. *Briqueté*, see Lateritious—u. de la Coction, see Urine.

URINE, CRUDE, (F.) *Urine crue*. Urine that is thin, transparent, and but little coloured, affording neither cloud nor deposit.

URINE CRUE, Urine, crude—u. *Cuite*, see Urine.

URINE, CYSTIN'IC, *Cystinu'ria*. Urine containing cystine or cystic oxide in solution, or as a deposit.

URINE, DIABET'IC, *Uri'na diabética*. Urine passed in *Diabetes mellitus*, *Melitu'ria*. It contains a large quantity of sugar analogous to that of the grape, which may be separated in the form of crystals. In other respects, the urine is very abundant and very limpid in that affection.

URINE DE LA DIGESTION, see Urine.

URINE, DROP'SICAL, is an ammoniacal urine, often containing much albumen and scarcely any urea.

URINE, DYSPEP'TIC, has been found to precipitate tannin abundantly, and to soon putrefy.

URINE ÉPAISSE, Urine, mucilaginous.

URINE, FEBRILE, *Uri'na febrilis*. Urine discharged in inflammation and inflammatory fever.

URINE, FILAMENTOUS, see Filamentous.

URINE, FLOCCULENT, (F.) *Urine floconneuse*, is when the fluid is troubled by flocculi, suspended in it.

URINE, FURFURA'CEOUS, see Furfuraceous—*U. floconneuse*, Urine, flocculent—*u. des Goutteux*, Urine, gouty.

URINE, GOUTY, (F.) *Urine des goutteux*, contains much phosphate of lime, and, after severe attacks of gout, precipitates a mixture of the uric and rosacic acids.

URINE HUILEUSE, Urine, oily.

URINE, HYSTER'ICAL, is clear, colourless, and similar to the *Urine de la Boisson*.

URINE, ICTER'ICAL, contains bile.

URINE, INCONTINENCE OF, Enuresis.

URINE OF INFLAMMATORY DISEASES. This is, generally, small in quantity, and very high-coloured.

URINE, INVOLUNTARY DISCHARGE OF, Enuresis.

URINE, JUMENTOUS, *Uri'na jumento'sa* seu *jumenta'ria*, (F.) *Urine Jumentouse*. A term used, by the French, for ammoniacal urine, rendered turbid by a substance similar to small grains of dust, which causes it to resemble the urine of herbivorous animals.

URINE LACTESCENTE, U. milky—*u. Lait-euse*. Urine, milky.

URINE, MILKY, (F.) *Urine lactescente ou lait-euse*. White and troubled urine.

URINE, MUCILAG'INOUS, (F.) *Urine épaisse*. Urine containing much mucus.

URINE, NERVOUS. Thin and very liquid urine; such as is passed shortly after the attack of nervous diseases.

URINE OF NERVOUS FEVERS is generally very thin; and often deposits a mixture of uric and rosacic acids.

URINE, OILY, (F.) *Urine huileuse*, is that which pours like oil, or which has an oily pellicle at its surface.

URINE, OXAL'IC, *Oxalu'ria*. Urine containing a salt of the oxalic acid.

URINE, PHOSPHORES'CENT, *Phosphoru'ria*, *Phosphu'ria*. Urine which is luminous or phosphorescent.

URINE, PURPURIC, see Porphyruria.

URINE, RICK'ETY, contains much phosphate of lime.

URINE, SANDY, Arenosa *urina*—u. Stoppage of, *Ischuria*—*u. Ténue*, Urine, thin.

URINE, THIN, (F.) *Urine ténue*. Urine that is transparent, slightly coloured, and but little dense.

URINEUX, Urinous.

URINIFEROUS, *Urinifer*, (F.) *Urinifère*, from *ovov*, 'urine,' and *fero*, 'I carry.' That which carries urine.

URINIFEROUS SUBSTANCE OF THE KIDNEY, see Kidney.

URINIFEROUS TUBES OR DUCTS OF BELLINI, *Tu'buli* seu *duc'tus uriniferi Bellini*, *Tu'buli Bellini* seu *recti*, *Bellinia'ni Ductus*, *Bellinia'na*

Fistulæ. The collection of small tubes, converging from the cortical part of the kidney to the apices of the papillæ. The tortuous uriniferous ducts are called *Tubes of Ferrein*, *Tu'buli contort'i* seu *Ferrein'ni*.

URINIP'AROUS, (F.) *Urinipare*, from *urina*, 'urine,' and *parere*, 'to produce.' An epithet for tubes in the cortical portion of the kidney, which prepare the urine.

URINODES, Urinous.

URINOM'ETER, from *ovov*, 'urine,' and *μετρον*, 'measure.' A small hydrometer for taking the specific gravity of urine. The urinometer suggested by Dr. Prout is divided into 60 degrees, the zero being the point at which it floats in distilled water. The numbers on the scale added to 1000, the specific gravity of water, give the particular specific gravity: thus, supposing the number cut by the surface of the liquid to be 30, it would indicate a s. g. of 1.030. The letters H. S. on the back of the scale mean "healthy standard," which ranges from 10° to 20°.

URINOUS, *Urino'sus*; *Urino'sus*, (F.) *Urineux*. Having the character of urine. Similar to urine.

UROCE'LE, *Cede'ma scroti* *urino'sum*, *Uroscheoce'le*, *Oucheoce'le* *urina'lis*, from *ovov*, and *κελη*, 'a rupture.' A tumour of the scrotum, owing to the effusion of urine into the same, and into the areolar tissue.

UROCHES'IA, *Urochez'ia*, from *ovov*, and *χεζειν*, 'to go to stool.' A discharge of urine by the bowels.

UROCRASIA, Enuresis.

UROCRIS'IA, from *ovov*, and *κρινω*, 'I judge.' *Urocrite'ria*. A judgment formed either of the urine, or of disease after inspecting the urine.

UROCRITERION, Urocrisia.

UROCYSTIS, Urinary bladder.

UROCYSTITIS, Cystitis.

UROCYSTOCATARRHUS, Cystirrhœa.

URODIAL'YSIS, from *ovov*, and *διαλυσις*, 'dissolution, loss of strength.' A partial and temporary suspension of the function of the kidney, such as occurs in the course of most acute diseases. A condition of the function of the kidney and of the urine, similar to that observed in acute diseases, but without local mischief or preceding inflammatory fever, has been termed, when occurring in children, *Urodial'ysis Neonato'rum*; in the aged, *Urodialysis Senum*.—Schönlein.

URODOCHIUM, Urinal.

URODYN'IA, from *ovov*, and *δύνη*, 'pain.' Pain during the excretion of the urine.

URÆDEMA, Uræpostema.

UROERYTHRIN, see Porphyruria.

UROGENITAL, *Urogenita'lis*, *Gen'ito-u'ri-nary*, *Gen'ito-urina'rius*. Relating or appertaining to the urinary and genital apparatus—as the 'urogenital organs'—*Organa urogenita'lia*.

UROGENITAL SINUS, *Sinus uro-genita'lis*. A cavity or canal in the embryo of the mammalia opening externally, in which the excretory ducts of the Wolffian bodies, the ureters and the efferent parts of the generative apparatus, terminate internally. It is prolonged into the urachus. Subsequently it becomes separated by a process of division into a *pars urina'ria* and a *pars genita'lis*. The former, extending towards the urachus, is converted into the urinary bladder;



Urinometer.

whilst from the latter are formed the *resicula seminules* in the male, and the middle portion of the uterus in the female.—J. Müller.

UROLITHI, Calculi, urinary.

UROLITHIASIS, Lithia.

UROLITHOLOG'IA, from *ουρον*, *λιθος*, 'a stone,' and *λογος*, 'a description.' A description of urinary calculi.

UROLITHOTOMIA, Lithotomy.

UROLOGY, Urology.

UROMANT'IA, from *ουρον*, and *μαντεία*, 'divination.' The art of divining diseases by simple inspection of the urine; *Uroscopia*. One professing to be able to do this is called *Uromantes*. Vulgarly, a *water-doctor*, *water-caster*.

UROM'ILES, (F.) *Uromèle*, from *ουρα*, 'a tail,' and *μελος*, 'a limb.' A genus of monsters in which the two lower limbs are united, very incomplete, and terminated by a single foot, which is almost always imperfect, and the sole of which is turned forwards.—I. G. St. Hilaire.

UROMPH'ALUS, *Her'nia U'rachi*. A monstrosity, which consists in the protrusion of the urachus at the umbilicus.

URON, Urine.

URON'CUS, from *ουρον*, and *ογκος*, 'a swelling.' A urinary swelling.

URONOL'OGY, *Uronolog'ia*, *Urol'ogy*, *Urol'og'ia*, from *ουρον*, and *λογος*, 'a description.' The part of medicine which treats of the urine.

UROPH'THISIS, Diabetes mellitus.

UROPLA'NIA, from *ουρον*, and *πλάνη*, 'wandering.' The presence of urine in other parts than the urinary organs.

UROPOE'ISIS, *Uropoe'isis*, from *ουρον*, and *ποιω*, 'I make.' *Secre'tio urinae* seu *lot'il*. The secretion of urine by the kidney.

UROPOIESIS, Uropoesis.

UROPYGION, Coccyx.

UROPYGIUM, *Coccyon*.

URORRHODINE, see *Porphyruria*.

URORRHAGIA, Diabetes.

URORRHŒA, Enuresis, Diabetes.

UROSCHEOCELE, Urocele.

UROSCOPIA, see *Uromantia*.

URO'SES, from *ουρον*, 'urine.' Diseases of the urinary organs.—Alibert.

UROSTEALITH, *Urosteal'ithos*, from *ουρον*, *στεα*, 'suet,' and *λιθος*, 'stone.' A peculiar fatty ingredient of urinary calculi, discovered by Heller.

UROSTENOSIS, Urethrostenosis.

UROUS ACID, Uric oxide.

UROZEMIA, Diabetes—u. Albuminosa, Kidney, Bright's disease of the—u. Mellita, Diabetes mellitus.

URRHODINE, see *Porphyruria*.

URROSACINE, see *Porphyruria*.

URTICA, U. *Dioi'ca* seu *major* seu *his'pida*, *Pyr'ethrum*, *Ad'icè*, Common stinging nettle, (Prov.) *Etelle*, (F.) *Ortie*. Ord. *Urticacæ*. *Sec. Syst.* Monoclea Tetrandia. This plant has been long used for medical and culinary purposes. The young shoots, in the spring, are considered to possess diuretic and antiscorbutic properties. The pain produced by its sting is well known, but it is not to be compared to the torture occasioned by *Urtica stim'ulans*, and *U. crenulata* of India; and, still less, to *Urtica urentiss'ima* of Timor, which is called by the natives *Daoun Setan*, 'Devil's leaf,' on account of the persistence and severity of the pain.

URTICA BALEARICA, U. *pilulifera*—u. *Crenulata*, see *Urtica*—u. *Dioica*, *Urtica*—u. *Hispida*, *Urtica*—u. *Iners magna fetidissima*, *Galeopsis*—u. *Major*, U. *dioica*—u. *Minor*, U. *urens*—u. *Mortua*, *Lamium album*.

URTICA PILULIF'ERA, U. *Pilulo'sa* seu *Roma'na* seu *Balea'rica*, *Pill-bearing Nettle*, (F.) *Ortie*

Romaine. The seed was formerly given in diseases of the chest.

URTICA PILULOSA, U. *pilulifera*—u. *Romana*, U. *pilulifera*—u. *Stimulans*, see *Urtica*.

URTICA URENS, U. *minor*, *Dwarf nettle*, (F.) *Ortie brulante*. A lesser variety of the nettle, possessing similar virtues.

URTICA URENTISSIMA, see *Urtica*.

Nettles have been used as stimulants to paralytic limbs.

URTICAIRE, *Urticaria*.

URTICA'RIA, from *urtica*, 'a nettle,' (uro, 'I burn,') *Enanthe'sis urtica'ria*, *Ure'do*, *Searlati'na* seu *Febris* seu *Pur'pura urtica'ta*, *Onido'sis*, *Urtica'ria porcel'iana*, *Exanthe'ma urtica'tum*, *Epinye'tis pruriginosa*, *Febris rubra pruriginosa*, *Saltus rosa*, (F.) *Urticaire*, *Fievre Ortie*, *Porcelaine*, *Nettle-rash*, *Nettle-springs*; vulgarly *Hives*, *Bold Hives*; from *urtica*, 'a nettle,' because the eruption resembles that produced by the stings of a nettle. A disease distinguished by those elevations of the cuticle called *wheals*. They have a white top, and are often surrounded by a diffuse redness. They excite intolerable itching, especially when the person is warm in bed. The eruption frequently goes and comes once or more in the 24 hours, and is generally dependent upon gastric derangement. In some people, nettle-rash is induced by shell-fish, eggs, nuts, &c. It is not usually a disease of any consequence, and its treatment is limited to the use of absorbent laxatives, and simple general means. Willan has described six varieties of the complaint:—*Urtica'ria febrilis*, U. *evan'ida*, U. *perstans*, U. *confer'ta*, U. *subcutanea*, and U. *tuberosa*.

URTICARIA PORCELLANA, *Essera*, *Urticaria*.

URTICA'TIO, *Onido'sis*, *Urtica'tion*, from *urtica*, 'a nettle.' A sort of flagellation, with nettles;—used with the intention of exciting the skin.

URUCU, Terra Orleana.

URYLIC ACID, Uric acid.

USNEA, Lichen saxatilis—u. Florida hirta, Lichen plicatus—u. Hirta, Lichen plicatus—u. Plicata, Lichen plicatus.

USSACU, Hura Brasiliensis.

USTILAGO, Ergot.

USTIO, Burn, Ustion.

USTION, *Us'tio*, *Causis*, from *urere*, *urum*, 'to burn.' The act of burning or of applying the actual cautery. Also, the effect of cauterization; a burn.

USTULA'TIO, *Ustulo'tion*. Same etymon. A pharmaceutical operation by which a substance is dried by heat. It is often restricted to the roasting of ores to drive off the volatile matters, as in the case of arsenic.

USUALIA, Officials.

USUS, Habit.

UTER, Uterus.

UTERI ADSCENSUS, Hysteria—u. Antever-sio, Hysteroloxia anterior—u. Carcinoma, Metrocarcinoma—u. Convulsivus morbus, Lorind matricis—u. Coryza, Leucorrhœa—u. Exulceratio, Hysteroleosis—u. Flexio, Hysteroloxia—u. Inclination, Hysteroloxia—u. Obliquitas, Hysteroloxia—u. Pronatio, Hysteroloxia anterior—u. Pruritus, Nymphomania—u. Reflexio completa, Hysteroloxia posterior—u. Retroversio, Hysteroloxia posterior—u. Rheuma, Leucorrhœa—u. Ulcus, Hysteroleosis—u. Versio incompleta, Hysteroloxia.

UTERINE, *Uteri'nus*, from *uterus*, 'the womb.' That which belongs or relates to the womb.

UTERINE ARTERY, *Arte'ria uteri'na hypogas'trica*. Its size is always in a ratio with the greater or less state of development of the uterus. It arises from the hypogastric, either separately

or with the umbilical, or from the internal pudic: passes upon the lateral parts of the vagina, to which it gives branches; and afterwards ascends on the sides of the uterus, and divides into a considerable number of very tortuous branches, which enter the tissue of the organ.

UTERINE BOUGIE, Sound, uterine.

UTERINE BROTHER. One born of the same mother, but of a different father.

UTERINE GLANDS, Utricular glands.

UTERINE MUSCLE OF RUYSCB consists of some fleshy fibres at the fundus of the uterus, which he regarded as forming a particular muscle.

UTERINE SOUND, Sound, uterine.

UTERINUS FRATER. In law, a brother by the mother's side.

UTERO-ABDOMINAL SUPPORTER, see Belt, Russian.

UTERO-GESTATION, Pregnancy.

UTEROMANIA, Nymphomania.

UTERUS, *ov&ap, Uter, Utriculus, Vulva, Loci, L. mulie'bres, Vulva, Venter, Conceptaculum, Matrice, Arcum seu Ager natu'rae, Hysteria, Metra, Stera, An'gus, Delphys, Alvus, Epomphalion, Guster, Mater, Nedys, Postreua, the Womb, (Sc.) Weam, (Prov.) Wem, (F.) Matrice, Sein.* The uterus, destined to lodge the fetus from the commencement of conception till birth, is a hollow symmetrical organ, having the shape of a truncated conoid: it is situate in the pelvis, between the bladder and the rectum, above the vagina, and below the convolutions of the small intestine. The uterus is flattened from before to behind, and is nearly an inch in thickness. It is two inches broad at its highest part, and becomes narrower towards the vagina; terminating in a contracted portion, called the *cervix* or *neck*, to distinguish it from the rest of the organ, called the *body*. From its *fundus* or upper portion, the Fallopian tubes pass off. The mouth is called *Os Uteri*, *Os Tinca*, *Os uteri externum*, *Orific'ium vagina'le*. At the point where the body of the uterus is continuous, below, with the neck, the cavity is slightly constricted, forming what is sometimes named the *internal orifice*, *Os uteri internum*, *Os internum*, *Isthmus seu ostium uteri*, *Orific'ium uterinum*. Between these is the *Canalis cervicis*.

The cavity of the uterus is very small, and its parietes are thick. The portion of the cavity corresponding with the body is triangular and flattened. Its superior angles present the extremely fine orifices of the Fallopian tubes. The cavity of the neck is slightly dilated before opening into the vagina. The uterus is composed, 1. Of a *serous membrane*. This is external, and furnished by the peritoneum; which is reflected upon the uterus from the posterior surface of the bladder and the anterior surface of the rectum. These folds of the peritoneum are sometimes called, respectively, the *anterior* and the *posterior ligaments*. The former are also named the *vesico-uterine* and the latter the *recto-uterine folds* or *plicae semilunares*. It forms, also, the broad ligaments of the uterus, *Ligamentum lata*, *Ala vespertilio'nis*, *Re'tia Uteri*. These contain, above, the Fallopian tube; beneath and anteriorly, the round ligaments; and, behind, the ovary. They are two membranous productions or duplicatures of the peritoneum, sent from the edges of the uterus and posterior extremity of the vagina, in a transverse direction, to be fixed to, and line, the side of the pelvis. 2. Of a *mucous membrane*, which is a prolongation of that of the vagina, and extends into the tubes. 3. Of a *proper tissue*, whose thickness is considerable. It is of a close texture; is grayish and elastic; and resists the instrument that divides it. Its intimate nature is but little known. During gestation it is truly muscular. The arteries of the uterus proceed

from the spermatic and hypogastric. Its veins follow the same course, and bear the same name as the arteries. They are very tortuous, and form large sinuses in the parietes of the organ, called *Uterine Sinuses*. Its nerves proceed from the sciatic and hypogastric plexuses. Its lymphatic vessels are very multiplied; and during pregnancy acquire enormous dimensions. The changes experienced by the uterus during gestation, and after delivery, are very numerous and important to be known. They regard its situation, direction, size, shape, texture, connexions vital properties, &c.

UTERUS, Abdomen—u. Anteversion of the, Hysteroloxia anterior.

UTERUS BIFIDUS. A double uterus. One separated into two cavities.

UTERUS DUPLEX, Dihysteria—u. Hour-glass contraction of the, *Chaton*—u. Inclinator, Hysteroloxia.

UTERUS, IR'BITABLE, Neural'gia of the Uterus. A disease characterized by deep-seated pain in the lower part of the abdomen, and in the back and loins; generally diminished by lying down, and increased by exercise, and more severe for a few days preceding and during menstruation. On examination *per vaginam*, the uterus is found to be tender on pressure, *Metriyperaesthesia*; and the cervix and body slightly swollen, but not usually indurated. It is an obstinate affection, but frequently ends of itself after years of suffering. The treatment must be,—the horizontal posture; local blood-letting to the uterus; emollients; soothing injections; narcotics, warm bath, &c. See Hysteralgia.

UTERUS, INVERSION OF THE, *Inversio Uteri, Metraastrophie, Edoptosis Uteri inversa*, (F.) *Inversio* or *Renversement de la Matrice*. The uterus, displaced and turned inside out. This can only happen through want of caution on the part of the medical practitioner. It is a fatal accident, unless speedily reduced. This must be done by gradually returning the superior part, by grasping it with the hand, and making the rest follow.

UTERUS MASCULINUS, see Ejaculatory ducts—u. Obliquus, Hysteroloxia—u. Retroversion of the, Hysteroloxia posterior.

UTERUS, RUPTURE OF THE, *Metrorrhæ'is, Hysterorrhæ'is, Ruptura seu Perforatio Uteri, Laceration of the Womb*, (F.) *Rupture or Déchirure de la Matrice*. This dreadful accident occurs during labour, and is known to have taken place, when the child's head suddenly recedes during a pain, with vomiting, sinking of the pulse, cold sweats, &c. When it is evident that this has occurred, the hand must be passed into the uterus, and the feet of the child be brought down.

Cases are on record, in which partial lacerations of the uterus have healed, and the patient has done well. This is a rare event, and not to be anticipated.

UTRICLE, see Semicircular canals.

UTRICLE, PRIMORDIAL, *Utriculus primordia'lis*. A delicate membrane, of protein composition, which lines the cell wall of vegetable structures, and from the exterior of which the layer of cellulose is secreted.

UTRICULAR, *Utricula'ris*; from *utriculus*, (diminutive of *uter, utris*), 'a small bag or bottle.' Relating to or resembling a small bag or bottle.

UTRICULAR GLANDS, *Glandulae utricular'es seu uteri'næ*. The glands or follicles in the interior of the uterus, which are concerned in the formation of the decidua.

UTRICULUS, see Ejaculatory ducts, Uterus—u. Communis, see Semicircular canals—u. Lac-teus, Receptaculum chyli—u. Primordialis, Utri-

cle, primordial—u. Prostaticus, see Ejaculatory ducts.

UVA, Uvula, see Vitis vinifera—u. Inversa, Paris—u. Lupina, Paris—u. Passa, see Vitis vinifera—u. Ursi, see Arbutus uva ursi—u. Vulpina, Paris.

UVÆ CORINTHIACÆ, see Vitis Corinthiaca.

UVÆ PASSÆ, Rhages, Dried Grapes, Raisins, see Vitis vinifera.

UVÆ PASSÆ MAJORES, see Vitis vinifera—u. Passæ minores, see Vitis Corinthiaca—u. Passæ sole siccatae, see Vitis vinifera.

UVARIA TRILOBA, see Carica papaya.

UVATIO, Ceratocelle, Staphyloma.

U'VEA, from uva, 'an unripe grape.' *Tu'nica acino'na seu acinifor'mis seu uua'lis seu acina'lis seu uvisfor'mis seu rhagoides seu uvea seu perfor'ata, Membrana uvea, Rhagoi'des, Rhox, (F.) Uvée.* Some anatomists have given this name to the choroid coat of the eye; others, to the posterior layer of the iris, on account of the black and very thick varnish that covers it. See Iris.

UVEA, COMMISSURE OF THE, Ciliary ligament.

UVÉE, Uvea.

UVEITIS; a word of hybrid formation; from uvea, 'the uvea,' and itis, denoting inflammation; properly, *Rhagodei'tis*. Inflammation of the uvea. See Iritis.

UVIGENA, Uvula.

UVIGERA, Uvula.

U'VULA, diminutive of uva, 'a grape.' *Pinnaculum for'nice guttura'lis, Uva, Staph'ylē, Tintinnabulum, Columel'la, Cion, Cionis, Gargareon, Colum'na oris, Gurgu'lio, Interseptum, Processus seu Cartila'go U'vifer, Uvig'ena, Uvig'era, Pap of the Throat, Himas, Plectrum, Sublin'gua, Sublin'guum, (Sc.) Pap of the Bass, (F.)*

Luette. A fleshy appendix or prolongation, which hangs from the middle or free edge of the velum palati. The uvula has a conical shape, and is of a greater or less size in individuals. It is formed, especially, by the mucous membrane of the velum palati; and contains, within it, the *azygos uvulae* muscle. Its use is not clear.

Also, a small prominence or lobule in the portion of the cerebellum that forms the posterior boundary of the fourth ventricle, *U'vula cerebell'i*. It is on the median line, and behind the nodulus.

UVULA, see Vermiform process inferior—u. Cerebelli, see Uvula—u. Decidua, Staphylœdema—u. Relaxata, Staphylœdema—u. Vesicæ, see Urinary bladder—u. Vesical, see Urinary bladder.

U'VULAR, *Uvula'ris*, from uvula. That which belongs to the uvula.

UVULAR GLANDS are small follicles, belonging to the mucous membrane covering the uvula.

UVULARIA, so called from the flowers hanging like the uvula [?], *Ruscus hypoglossum*.

UVULA'RIA PERFOLIA'TA, *Smaller Bellheart*; indigenous; Order, Melanthaceæ; flowering in May and June. The root, when fresh, is subacrid and mucilaginous. A decoction of the plant is used in sore mouth. It is said to cure the bites of rattlesnakes.

UVULARIA GRANDIFLO'RA and *U. Sessilifolia* have similar properties.

UVULITIS, a word of hybrid formation, from uvula, and itis, denoting inflammation. *Angi'na uvula'ris, Staphyli'tis, Cioni'tis, Inflamma'tio u'vulae, (F.) Inflammation de la Luette, Falling down of the Palate.* Inflammation of the uvula. When to a slight degree, *Hypostaphyli'tis*.

V.

VACCIN, see Vaccina.

VACCINA, *V. primig'ena seu vaccinato'ria seu antivariolo'sa, Vaccin'ia, Vari'ola vaccina, V. vaccinia seu tuto'ria seu tutrix, Exanthema vaccina seu antivariolo'sum, Syn'ocha vaccina, Em'phlysis vaccina inser'ta, Vacci'ola, V. scutell'ata, V. vera, Vari'ola tuto'ria, Cow-pox, Kine Pock, Inoculated Cow-pox, from vacca, 'a cow'; (F.) Vaccine.* The cow-pox is a disease of the cow, arising spontaneously, which, if transmitted to man by means of inoculation, may preserve him from small-pox contagion. The promulgation of this valuable property of the vaccine virus is due to Dr. Jenner. When inoculating for small-pox in Gloucestershire, in 1775, he was surprised to meet with a number of persons in the same district, on whom the insertion of the virus produced no effect, although they had never passed through small-pox. On investigation, he found that all these persons had previously been affected by a pustular eruption of the cow, which they had taken by milking the animal; and which eruption they called *Cow-pox*. This discovery induced him to institute experiments, and he found that such preservative effect actually existed. The *Vaccine Virus, Virus Vaccinum, (F.) Vaccine* or the fluid of the eruption, is not now usually taken immediately from the cow: experience having demonstrated that it is nearly as efficacious when obtained from the human vesicle. The characteristic of cow-pox eruption is:—a semf-transparent, pearl-coloured vesicle,

with a circular or somewhat oval base; its upper surface, until the end of the 8th day, being more elevated at the margin than in the centre; and the margin itself being turgid, shining, and rounded, so as often to extend a little over the line of the base. This vesicle is filled with clear lymph, contained in numerous little cells, that communicate with each other. After the 8th or 9th day from the insertion of the virus, it is surrounded by a bright red, circumscribed areola, which varies in its diameter, in different cases, from a quarter of an inch to two inches; and is usually attended with a considerable tumour, and hardness of the adjoining areolar membrane. This areola declines on the 11th or 12th day; the surface of the vesicle then becomes brown in the centre; and the fluid in the cells gradually concretes into a hard, rounded scab or crust, of a reddish-brown colour, which at length becomes darker, contracted, and dry, but is not detached till three weeks after vaccination. It leaves a permanent, circular cicatrix, about five lines in diameter, and a little depressed; the surface being marked with very minute pits or indentations, denoting the number of cells of which the vesicle had been composed. This is the progress of the genuine cow-pox. At times, fever occurs about the 9th or 11th day; but, at others, not the slightest indisposition is felt. Aberrations from this progress occur, which must be attended to; as the same preservative influence is not exerted by irregular or spurious cow-pox. If the erup-

tion runs speedily to a height, has an areola too early, is irregular at its edges, and contains pus instead of a limpid fluid,—it cannot at all be depended upon.

Small-pox occurs, occasionally, as an epidemic after vaccination. For a while, the enthusiastic advocates of vaccination would not admit the affection to be small-pox; and, accordingly, they gave it the name Varioloid. It is unquestionably small-pox, but modified, and greatly divested of its terrors by previous vaccination; and it is said to happen less frequently after vaccination than after small-pox inoculation. Vaccination is now practised everywhere, except among those in whom ignorance and prejudice exclude the lights of reason and philanthropy. There is ground, however, for the belief, that for full protection revaccination may be occasionally necessary.

VACCINA ANTIVARIOLOSA, Vaccina—v. Primi-gena, Vaccina—v. Tutoria, Vaccina—v. Tutrix. Vaccina—v. Vaccinatoria, Vaccina.

VACCINAL, Vaccine.

VACCINATION, Vaccinat'io, from vacca, 'a cow'; Cow-pox inoculation, Jennerian inoculation. An operation which consists in inserting the vaccine virus under the cuticle, so that it may come in contact with the absorbents. It is extremely simple, and requires merely that the point of the lancet should be held under the cuticle for a short time. The best time for taking the matter from the Cow-pox vesicle, for vaccination, is about the 7th or 8th day, before the appearance of the areola, when it ought to be limpid and transparent. The most certain method is, to pass it from arm to arm; but this is not always convenient; and, hence it is taken upon points or pieces of quill, or between glasses. It must, of course, be softened for use. This is done by means of a little water, which does not destroy its efficiency. The scab, too, which falls off about the end of three weeks—sooner or later—is capable of communicating the affection, if dissolved in water; and this is the most common method of transmitting the vaccine virus in some countries.

VACCINATOR, Vac'cinist. One who inoculates for the cow-pox.

VACCINE, Vaccina.

VACCINE, Vaccin'us, Vaccina'lis, (F.) Vaccinal; same etymon. Relating or appertaining to vaccination, as the 'vaccine vesicle,' 'a vaccine physician.'

VACCINELLA. A term employed by some to designate spurious cow-pox. Vaccin'ois, Vacci'ola spu'ria seu lepro'sa, (F.) Vaccinelle, Vaccinoïde.

VACCINIA, Vaccina—v. Palustris, Vaccinium oxycoccos.

VACCINIST, Vaccinator.

VACCINIUM MACROCARPON, see V. oxycoccos.

VACCINIUM MYRTILLUS, either corrupted from *hyacinthus*, or, and more probably, *baccinum*; from *baccæ*, 'berries,' in allusion to its fruit. *Vitis idæa myrtillus*, Vaccin'ia myrtillus, Vaccin'ium nigrum, Myrtle Berry, Bilberry, Bleaberry, Whortleberry, Black Whortle, (Sc.) Blueberry, (Prov.) Hurtleberry, Whimberry, (F.) Airelle Anguleuse, Raisin de Bois. Ord. Vaccinieæ. Ser. Syst. Octandria Monogynia. The berries—*Baccæ Myrtillorum*—are esteemed to be antiscorbutic, when prepared with vinegar. They are also said to possess antiseptic and sub-astringent properties when dried. They are not used in medicine; but are frequently eaten with milk and sugar.

VACCINIUM NIGRUM, V. myrtillus.

VACCINIUM OXYCOCOS, *Oxycoc'cos*, *O. palus-*

tris, *Schollera oxycoccos*, *Vaccinia palustris*, *Vitis idæa palustris*, Moorberry, Cranberry, Mossberry, Swamp Redberry, Sourberry, (F.) Canneberge. The berries are pleasantly acid and cooling. They make an agreeable preserve. The common American Cranberry is from *Vaccinium macrocarpon*.

VACCINIUM PENNSYLVANICUM, Low Blueberry. The leaves and bark of the root are astringent.

VACCINIUM PUNCTATUM, V. vitis idæa—v. Resinosum, Gaylussacia resinosa.

VACCINIUM STAMINEUM, Buckberry, Squaw Huckleberry, Deerberry. The leaves of these plants are astringent.

VACCINIUM VITIS IDÆA, V. punctatum, *Vitis idæa punctata*, *Am'pelos idæa*, Red Bilberry, (Sc.) Brawline, Alpine Cranberry, Whortleberry, Cowberry, (F.) Airelle ou Canneberge punctée. A decoction of the leaves has been used in calculous affections. They are astringent. The ripe berries are pleasantly acid, and refrigerant. Whortleberries are furnished by different species of *Vaccinium*.

VACCINOÏDE, Vaccinella.

VACCINOIS, Vaccinella.

VACCINUS, Vaccine.

VACCIOLA, Vaccina—v. Leprosa, Vaccinella—v. Scutellata, Vaccina—v. Spuria, Vaccinella—v. Vera, Vaccina.

VACILLATIO, Tituba'tio, from vacillare, 'to waver.' Staggering; waddling—*claudica'tio anati'ca*.

VACILLATIO DENTUM, Odontoseisis.

VAGIN, Vagina.

VAGINA, 'a sheath,' V. mulie'bris seu uteri'na seu Uteri seu Penis, Sinus mulie'bris seu Pud'oris seu Pudendi, Col'eus, Peris, Perin, Colpos, Vulvo-uterine canal, (F.) Vagin, Conduit de la pudeur. A cylindrical canal, five or six inches long; situate within the pelvis, between the bladder and rectum. It communicates by one extremity, Aulos, with the vulva; by the other, Fornix seu Fundus, with the womb, Pars seu Portio vagina'lis u'teri, the neck of which it embraces. The vagina is lined, internally, by a mucous membrane; and is partly closed, in young females, by the hymen, and carunculae myrtiformes. Around the mucous membrane is a layer of spongy, erectile tissue; and a celluloso-vascular membrane. A constrictor muscle; numerous vessels furnished by the vaginal artery, particularly; veins and nerves, proceeding from the sciatic plexus, also, enter into the composition of the organ.

The term Vagina or sheath, Theca, is extended to many parts, which serve as envelopes to others.

Sheath, (F.) Gaine, is sometimes applied to the fascia enveloping the limbs; to areolar membrane, enveloping the muscles, and entering between their fibres; to that which surrounds the blood-vessels; to the synovial membranes surrounding tendons, and lining the gutters of bones, &c.

VAGINA CORDIS, Pericardium—v. Femoris, Fascia lata aponeurosis—v. Funiculi Umbilicalis, see Funiculus umbilicalis—v. Glissonii, Capsule of Glisson—v. Malleolaris externa, Retinaculum tendinum perinæorum—v. Muliebris, Vagina—v. Muscularis, Perimysium—v. Nervorum, Neurilemma—v. Penis, Vagina—v. Pili, see Hair—v. Portæ, Capsule of Glisson—v. Uterina, Vagina.

VAGINA OF THE VENA PORTA. Sheath of the Vena Porta, (F.) Gaine de la Veine-porte, is the areolar membrane that accompanies the divisions of the vena porta into the substance of the liver.

VAGINÆ SYNOVIALES, Bursæ mucosæ.

VAG'INAL, Vagina'lis, from vagina, 'a sheath. Relating or pertaining to a sheath.

VAGINAL ARTERY. This exists, as a matter of

course, only in the female. It arises from the hypogastric or from the uterine, vesical, internal pudic, umbilical, &c., and sends its branches, first to the lateral parts of the vagina, and afterwards to the anterior and posterior surfaces.

VAGINAL BILIARY PLEXUS. A plexus of ramifications of the hepatic duct through the capsule of Glisson.

VAGINAL BRANCHES OF THE VENA PORTA are those which have to pass through the sheath of the capsule of Glisson, previous to entering the interlobular spaces. In this course, they form the *vaginal plexus*. There are, also, *vaginal branches*, and a *vaginal plexus* formed by the hepatic artery.

VAGINAL COAT OF THE EYE, see Eye.

VAGINAL COAT OF THE TESTICLE, *Tunica Vaginalis* seu *Elytroides* seu *Elytrodes*, is the serous membrane which envelops the testicle. Before the descent of the testicle, the tunica vaginalis does not exist. It is formed by a prolongation, furnished by the peritoneum to the testicle when it issues from the abdomen. The tunica vaginalis forms a close sac, investing the testicle and epididymis, without containing them in its cavity, and reflected so as to form a bag around the circumference of the testis; hence it is divided into *Tunica vaginalis propria* and *Tunica vaginalis reflexa*. Between the two layers, the inner surface is smooth, and lubricated by a serous fluid.

VAGINAL GANGLIA, see Spermatic ganglion.

VAGINAL PLEXUS, see Vaginal branches of the vena porta.

VAGINAL PROCESS OF THE TEMPORAL BONE, (F.) *Apophyse vaginale* ou *engainante*, *Gaine de l'apophyse styloïde*, Chaton, is a bony crista, which embraces the base of the styloid process.

VAGINAL PULSE, Pulse, vaginal.

VAGINITIS, Leucorrhœa (vaginal.)

VAGINO-VESICAL, Vesico-vaginal.

VAGISSEMENT, Cry, Vagitus.

VAGITUS, (F.) *Vagissement*. The cry of the new-born child. *Squalling*.

VAGITUS UTERINUS. The cry of the child in utero.

VAGUS NERVUS, Pneumogastric.

VAIRON (F.), *Dispar oculus*. One who has eyes of different colours. It also means a *walleyed* animal, or one whose iris is surrounded by a whitish circle.

VAISSEAU, Vessel—*v. Sanguin*, Blood-vessel.

VAISSEAUX CAPILLAIRES, Capillary vessels—*v. Chylifères*, Chyliferous vessels—*v. Lactées*, Chyliferous vessels—*v. Lactifères*, Lactiferous vessels—*v. Omphaliques*, Umbilical vessels—*v. Tourbillonnés*, Vasa vorticosa.

VALANIDA, Fagus sylvatica.

VALANTIA APARINE, Galium aparine.

VALERIAN, AMERICAN, Cyripedium luteum, Valeriana pauciflora—*v. Garden*, Valeriana phu—*v. Greek*, Polemonium reptans—*v. Official*, Valeriana—*v. Wild*, Senecio aureus, Valeriana, *V. pauciflora*.

VALERIA'NA, *V. officinalis* seu *minor*, *Leucolach'anum*, Wild or *Officinal Valerian*. *Ord. Valerianaceæ. Sex. Syst. Triandria Monogynia*. The root, *Valeria'na* (Ph. U. S.), has been long extolled as an efficacious remedy in epilepsy, and in a variety of nervous complaints. It has been also advised as a tonic, antispasmodic, and emmenagogue. It has a strong, fetid odour; bitterish, subacid, and warm taste; and its virtues are extracted by water and alcohol. The distilled oil is official in the Pharmacopœia of the United States.

VALERIANA CAPEN'SIS, a South African species, has similar virtues.

VALERIANA CELTICA, *V. Saxatilis*, *Spica Celtica* seu *Celtica Dioscoridis* seu *Alpina*, *Nardus Celtica*, *Salina*, *Nardum Gallicum*, *Celtic Nard*, (F.) *Nard Celtique*. The roots of this Alpine plant have been recommended as a stomachic, carminative and diuretic. It has a moderately strong smell, and a warm, bitterish, sub-acrid taste.

VALERIANA DENTA'TA, *V. locusta*, *Corn Salad*. A wholesome, succulent plant, cultivated in Europe as a salad. It is antiscorbutic and gently aperient.

VALERIANA MAJOR, *V. phu*—*v. Locusta*, *V. dentata*—*v. Minor*, Valeriana—*v. Officialis*, Valeriana.

VALERIANA PAUCIFLO'RA, *American* or *Wild Valerian*, and *V. sylvatica*, have the properties of the other Valerians.

VALERIANA PHU, *V. major*, *Garden Valerian*, (F.) *Grande Valériane*. The root has been recommended in rheumatism; especially in sciatica, and in epilepsy.

VALERIANA SAXATILIS, *V. celtica*—*v. Spica*, *Nardus Indica*—*v. Sylvatica*, see *V. pauciflora*.

VALÉRIANE, GRANDE, Valeriana phu.

VALET À PATIN (F.), *Volse'lla Patini*. The common ligature forceps is a modification of this instrument invented by one Patin. It consists of two branches, united at the middle by a hinge, which may be separated or approximated by means of a screw or running ring. It is used for seizing hold of vessels that are to be tied.

VALETUDINAIRE, Valetudinary.

VALETUDINARIUM, Hospital.

VALETUDINARIUS, Sickly, Valetudinary.

VALETU'DINARY, *Valetudina'rian*, *Valetudina'rius*, *Inval'idus*, (F.) *Valetudinaire*, from *valetudo*, 'health,' *valet*, 'to be well.' One of delicate health. One subject to frequent diseases. *An Invalid*.

VALETUDO, Sanitas—*v. Adversa*, Disease—*v. Bona*, Sanitas—*v. Commoda*, Sanitas—*v. Secunda*, Sanitas.

VALGUS, see *Cnemocoliosis* and *Kyllosis*.

VALIGA, Tinctura jalapii.

VALITUDO, Sanitas.

VALLECULA, Valley—*v. Unguis*, see Nail.

VALLEY, *Vallee'ula*, *V. Reil'ii*, (diminutive of *vallis*, 'a valley,') (F.) *Grand Seigneurs Médiane du Cervelet*. The deep fissure on the inferior part of the cerebellum, which divides it into two symmetrical portions.

VALLUM. The eyebrow or supercilium. Also, a species of bandage.—Galen.

VALS, MINERAL WATERS OF. A French acidulous chalybeate at Vals, six leagues from Viviers, in the department of Ardèche. There are five springs, containing carbonic acid, sulphate of iron, and alum. They are employed in passive hemorrhages, leucorrhœa, &c.

VALVE, *Val'vula*, *Dielis*, (F.) *Valvule*, from *valva*, 'folding-doors,' itself from *volvo*, 'I fold up.' A small door. Any membrane or duplication of a membrane, which prevents a reflux of humours or other matters in the vessels and canals of the animal body. There are some valves, whose functions appear to be,—to retard or to modify the course of substances along canals, &c. See *mitral*, *sigmoid*, *tricuspid*, *ileo-cæcal valve*, &c.

VALVE OF FALLOPIUS, Bauhin, valve of—*v. Ileo-cæcal*, Bauhin, valve of—*v. Ileo-colic*, Bauhin, valve of—*v. of Tulpius*, Bauhin, valve of—*v. of Varolius*, Bauhin, valve of.

VALVES, KERKRINGIAN, *Valvulæ conniventes*—*v. Semilunar*, Sigmoid valves.

VALVULA, Valve—*v. Cæci*, Bauhin, valve of—*v. Cerebelli*, *Valvula Vieussensii*—*v. Coli*, Bau-

bin. valve of — v. Foraminis ovalis, Ovalis fossa — v. Guiffartiana, see Thebesius, veins of — v. Hei. Baubin, valve of — v. Magna Cerebri, V. Vieussenii.

VALVULA SEMILUNARIS, *Plica Semilunaris* seu *lunata*, *Membrana semilunaris conjunctiva*. A small doubling of the tunica conjunctiva, which lies between the caruncula lacrymalis and the ball of the eye. In birds, this is large, and called *Membrana nictitans* seu *Palpebra tertia*.

VALVULA TARNI, Velum medullare posterius — v. Thebesii, see Thebesius, veins of — v. Vaginata, Hymen — v. Venæ magnæ, see Thebesius, valve of.

VALVULA VIEUSSENI, *V. cerebelli* seu *Willisia* seu *magna cerebri*, Velum Vieussenii seu *interjectum cerebelli*, V. apophysi vermiformi obtentum, V. medullaræ seu medullaræ anticum seu *anterioris*, *Lacunar ventriculi quarti superior*, (F.) *Valvule de Vieussens*, *Voile médullaire supérieure*. A thin, medullary lamina, over the under end of the Aquæductus Sylvii and upper part of the fourth ventricle. At the sides of the valves are two medullary tracts, called *Processus ad Testes*, *Columina Valvulæ Vieussenii* or *Oblique inter-cerebral Commissure*.

VALVULA WILLISIANA, Valvula Vieussenii.

VALVULÆ BICUSPIDALES, Mitral valves.

VALVULÆ CONNIVENTES, *V. C. Kerkringii*, *Plicæ conniventes*, *Kerkringian valves*. Numerous circular folds in the intestinal canal from the pyloric orifice through the greater part of the small intestine. They are folds of the mucous membrane; and their chief use appears to be, to retard the course of the alimentary mass, and to afford a larger surface for the origin of chyloferous vessels.

VALVULÆ CORDIS, Ostiola cordis — v. Cordis mitrales, Mitral valves — v. Tricuschines, Tricuspid valves — v. Trisulcæ, Trisulcid valves.

VALVULE, Valve — v. d'Eustache, Eustachian valve — v. de Pyloro, see Pylorus — v. de Vieussens, Valvula Vieussenii.

VALVULES ÉPISCOPALES, Mitral valves — v. Sigmoides ou *Semilunaires*, Sigmoid valves.

VANDELIA DIFFUSA. A native of Brazil, Guiana and Isle of France; Order, Scrophulariaceæ; is used there as an antibilious emetic and antiperiodic; and is regarded as an efficacious remedy in malignant fever and dysentery, especially in cases connected with a disordered state of the liver.

VANELLOE, Vanilla.

VANIL'LA, *Banil'ia*, *Banilas*, *Banilloës*, *Banilla*, *Benzo'ënil*, *Aracus aromaticus*, (F.) *Vanille*, *Baume de Vanille*. The *Vanellæ*—*Sil'iqua vanill'æ* seu *Banill'æ* seu *Vanigliæ* seu *vanillæ* seu *Ar'aci aromatici* — is a long, flattish pod; containing, under a wrinkled, brittle shell, a reddish-brown pulp, with small, shining, black seeds. The parasitic plant which affords this pod is the *Epidendrum Vanilla*, *Vanilla aromatica* seu *Mexicana*, and other species. Family, Orchidaceæ. *Sex. Syst.* Gynandria Monandria. Vanelloes have an unctuous, aromatic taste, and a fragrant smell, like that of some of the finer balsams, heightened with musk. Although chiefly used as perfumes, they are reputed to possess aphrodisiac virtues.

VANILLA AROMATICA, see Vanilla.

VAPEURS, Hypochondriasis, Hysteria.

VAPOR SANGUINIS, Gaz sanguinis.

VAPORARIUM, *Amist'rium*, *Lacunculæ*, *Pyria*, *Pyriate'rium*, *Vapour bath*, from *vapor*, 'steam.' A useful remedy in rheumatic and cutaneous affections; and wherever it is important to relax the cutaneous capillaries. In the *Russian Vapour Bath*, the vapour is produced by

throwing water over hot stones. Its temperature is from 122° to 145°.

VAPORARIUM, Hypocaustum, see Stove.

VAPORATIO, Evaporation.

VAPORES UTERINI, Hysteria.

VAPOUR, Flatulence — v. Bath, Vaporarium.

VAPOURISH, Hypochondriac, Hysterical.

VAPOUROUS, Windy.

VAPOURS, Hypochondriasis, Hysteria.

VAPOURY, Hypochondriacal, Hysterical.

VARA, Barre.

VAREC, Kelp — v. *Vésiculeux*, *Fucus vesiculosus*.

VARENI, Ambulo-flatulentus, Gout (wandering).

VARICE, Varix.

VARICELLA, (diminutive of *Variola*, Small-pox,) *Em'phlysis* seu *Exanthema* seu *Syn'ochus varicella*, *Variola lymphatica* seu *spuria* seu *sylist'ris*, *Variola notha* seu *illegitima* seu *vulgaris* seu *puella*, *Pseudo-variola*, *Æol'ion*, *Æollium*, *Chicken pox*, (F.) *Varicelle*, *Pétite érole volante*, *Vérolette*, *Fausse variole*. A disease characterized by vesicles scattered over the body, which are glabrous, transparent, and about the size of peas. They appear in successive crops, are covered by a thin pellicle, and, about the third, fourth, or fifth day from their appearance, burst at the top, and concreate into small, puckered scabs, which rarely leave a pit in the skin. Three varieties have, sometimes, been designated. 1. The *V. lentiformes*, or *lenticular chicken pox*, which are irregularly circular, and flattened on the top; the fluid being at first whitish, and afterwards straw-coloured. This is the common *chicken pox*. 2. The *Varicella coniformis*, *Pent'phigus varioloides*, *Variola lymphatica*, *Hydrach'nia*, *Conoid'al Chicken Pox*, *Swine Pox*, *Water Pox*, *Water Jags*, in which the vesicles are generally acuminate, and the fluid pellucid throughout; and, 3dly, the *Varicella globularis* or *Hæes*, in which the vesicles are globular and larger; the fluid being, at first, whey-coloured, and afterwards yellowish. These divisions are, however, very arbitrary, for they are generally confounded.

Hornpox is a popular variety of varicella.

Variola has been, sometimes, mistaken for this disease, and conversely. A little attention to the history of the eruption in the two affections will prevent error. Varicella is not contagious. At times, it prevails epidemically. It cannot be propagated by inoculation. The treatment is extremely simple; rest, abstinence, and the antiphlogistic regimen, being all that is generally required. See Varioloid.

VARICELLA VARIOLOIDA Varioloid.

VARICELLAR FEVER, see Varioloid.

VARICIFORMES PARASTATÆ. A name given, by some authors, to the ducts of the epididymis, because they seem knotty and varicose.

VARICOBLEPH'ARON, from *varix*, *varicis*, and *βλεφαρον*, 'eyelid.' A varicose tumour of the eyelid.

VARICOCE'LE, from *varix*, *varicis*, and *κλην*, 'a tumour' *Osciocele* seu *varico'sa*, *Her'nia varico'sa*, *Rameæ varico'sus*, *Spermatocele*, *Circocele*. A varicose dilatation of the veins of the scrotum and spermatic cord. This latter variety of the disease is more common on the left side than the right. It appears under the form of a soft, doughy, unequal, knotty, compressible, and indolent tumour, situate in the course of the cord, and increasing from below upwards. It is less when the individual is in the recumbent posture, and increases when he is erect. It is, ordinarily, an affection of no moment, and is very common, requiring merely the use of a proper suspensory bandage. It has been advised, in bad cases, to

tie the enlarged vein, but the operation is not free from danger induced by phlebitis. Sir A. Cooper has proposed the removal of a portion of the scrotum, so as to render the remainder of the scrotum a natural bandage. This plan has been found effectual.

VARICOMPHALUS, from *varix*, *varicis*, and *omphalos*, 'the umbilicus.' Varicose tumour of the umbilicus; cirsiomphalus.

VARICOSE, *Varico'sus*, (F.) *Variqueux*. From *varix*, *varicis*. Affected with varix; or belonging to, or connected with, varix;—as a *varicose vein*, a *varicose ulcer*, that is, one kept up by varices.

VARICOSITAS, see *Varix*—v. *Conjunctivæ*, *Cirsiophthalmia*.

VARICOSITY, see *Varix*.

VARICULA, *Varix parva*. Diminutive of *varix*. Also, a varicose swelling of the veins of the tunica conjunctiva.—M. A. Severinus.

VARIOLA, *Variolæ*, *V. veræ*, *Varioli*, *Empy-eis Variola*, *Euphlog'ia*, *Chaspè*, *Pestis seu Febris variolosa*, *Æolec'thyma*, *Small-Pox*, *Pox* (N. of England); from *varius*, 'spotted,' or from *vari*, 'pimples;' (F.) *Variolæ*, *Petite Vérole*, *Picote*. A disease, now of somewhat less interest than before the discovery of vaccination. It is of a very contagious nature, and is supposed to have been introduced into Europe from Asia, at an early period of the middle ages. It is characterized by fever, with pustules appearing from the third to the fifth day, and suppurating from the eighth to the tenth, and it possesses all the distinctive properties of the major exanthemata. It is capable of being produced by inoculation, but this inoculated small-pox—*Variola inserta*—communicates the disease as readily through the air as the natural small-pox, or that received without inoculation.

Small-pox is distinguished into two classes, according to the character of the eruption, the *discrete* or *distinct*, and the *confluent*. 1. In **VARIOLA DISCRETA**, *Empyysis variola discreta*, *Variola discreta benigna*, *Variolæ regulares seu discretæ*, *Distinct Small-pox*, the pustules are usually of the size of peas; distinct, distended, and circular; the intervening spaces being red; the fever inflammatory throughout, and ceasing when the eruption is complete. 2. In **VARIOLA CONFLUENS**, *Empyysis variola confluenta*, *Variolæ regulares confluentes*, *Venic'ula Divæ Barbaræ*, *Vésicules de Sainte-Barbe*, *Confluent Small-pox*, the pustules are confluent, or run together; flaccid and irregularly circumscribed; the intervening spaces being pale, the accompanying fever typhoid. In children, diarrhoea, and, in adults, pyralism, with swelling of the hands and feet, generally appear, towards the period of the *secondary fever*, which occurs from the tenth to the thirteenth day. The fever that precedes and accompanies the eruption is called the *Eruptive fever*. The prognosis is favourable in the distinct kind, if properly managed. The confluent is always dangerous,—the unfavourable symptoms being—flattening of the pustules, or subsidence of the eruption; the breathing becoming much obstructed or oppressed; or marks of inflammatory and congestive affections occurring in the different viscera. The treatment of the distinct kind is very simple: the antiphlogistic regimen being all that is necessary. The confluent kind requires the same management as typhus fever.

When the pimples are confluent in patches, the patches, being, however, separated by intervals of unaffected skin; it constitutes the clustered, coherent, or corymbose variety—*Variola corym-bosa*.

VARIOLA AMBOINENSIS, *Frambesia*—v. *Corymbosa*, see *Variola*—v. *Inserta*, *Variola* (by inoculation)—v. *Lymphatica*, *Varicella*—v. *Mitigata*, *Varioloid*.

VARIOLA sine VARIOLIS, *Variolous Fever*. These terms are applied to cases occasionally observed during the prevalence of variola, in which the constitutional phenomena are present without the cutaneous eruption.

VARIOLA SPURIA, *Varicella*—v. *Vaccina*, *Vaccina*—v. *Vaccinatorum*, *Varioloid*.

VARIOLÆ, *Variola*—v. *Equine*, see *Grease*—v. *Illegitimæ*, *Varicella*—v. *Modificatæ*, *Varioloid*—v. *Nothæ*, *Varicella*—v. *Pucellæ*, *Varicella*—v. *Sylvestris*, *Varicella*—v. *Tutoriæ*, *Vaccina*—v. *Veræ*, *Variola*—v. *Volaticæ*, *Varicella*.

VARIOLATION, *Inoculation* (Small-pox.)

VARIOLE, *Variola*.

VARIOLI, *Variola*.

VARIOLIQUE, *Variolous*.

VARIOLOID, *Variolois*, *Varioloi'des*, *Vari'ola modificata seu mitigata seu vaccinatum*, *Varicella* (of some), *Varicella Varioloidea*, from *variola*, small-pox, and *eidos*, 'form,'—resembling small-pox. This is, really, small-pox, modified by previous inoculation or vaccination; and hence it has been properly called *modified small-pox*. That it is small-pox is proved by the fact, that matter, taken from a varioloid pustule, will communicate small-pox to one who has never been vaccinated. It is, almost always, a milder disease than small-pox; and this circumstance, with its shorter duration, exhibits the salutary effects of previous vaccination or inoculation. It has appeared epidemically.

Occasionally, when varioloid is epidemic, *varioloid* or *varicellar fever* occurs without the eruption. This is the *varicellæ sine varicellis* of Mr. Erasmus Wilson, who calls varioloid, *varicella*.

VARIOLOID FEVER, see *Varioloid*.

VARIOLOIS, *Varioloid*.

VARIOLOUS, *Variolo'sus*, (F.) *Variolique*. Relating or belonging to variola.

VARIQUEUX, *Varicose*.

VARIX, genitive *varicis*, *Phlebeury'sma*, *Phlebec'tasis*, *Her'nia seu Ectasia vena'rum*, *Angiectasia vena'sa*, *Ix'ia*, *Circus*, *Cedma*, *Var'icose Vein*, (F.) *Varice*, is considered, by some, to come from the verb *variare*, 'to turn,' 'to twist,'—on account of the sinuosities of varicose veins. Dilatation of a vein; *Varicos'itas*, *Varicosity*. Varices are owing to local retardation of the venous circulation; and, in some cases, to relaxation of the parietes of the veins. They are very common in the superficial veins of the lower limbs, especially of pregnant females. The tumour or tumours, formed by varices, are soft, knotty, unequal, indolent, and livid; without pulsation, and yielding readily to the impression of the finger; but returning as soon as the compression is discontinued. Sometimes, the vein bursts and gives rise to hemorrhage. The treatment is usually palliative, and consists in exerting a uniform and constant pressure upon the part, by means of an appropriate bandage. The radical cure has, also, been obtained by extirpation, division of the vein, or ligature. The second course is apt to be followed by phlebitis.

VAROLI, *BRIDGE OF*, *Pons Varolii*.

VARUS, *Aene*, *Ionthus*, see *Entogonyancon* and *Kyllosis*—v. *Mentagra*, *Sycosis*.

VAS, *Vessel*, *Penis*—v. *Deferens mulieris*, *Tuba Fallopiana*—v. *Urinarium*, *Ureter*.

VASA ABERRANTIA. Peculiar ramifications of the biliary duct in the liver are so called by E. H. Weber.—Kölliker.

VASA ABSORBENTIA, Absorbent vessels—*v.* **Afferentia**, see Afferent and Efferent—*v.* **Capillaria**, Capillary vessels—*v.* **Chylifera** seu **Chylofera**, Chyliferous vessels.

VASA CHYLOPOËTICA URINIFERA, Blood-vessels, which, according to Lippi, convey drinks from the chyliferous vessels to the renal veins.

VASA EFFERENTIA, see Efferent—*v.* **Exhalantia**, Exhalant vessels—*v.* **Hydragoga**, Lymphatic vessels—*v.* **Inferentia**, see Afferent and Efferent—*v.* **Lactea**, Chyliferous vessels, Lactiferous vessels—*v.* **Meningea**, Meningeal vessels—*v.* **Sedalia**, Hæmorrhoidal vessels—*v.* **Lymphatica**, Lymphatic vessels—*v.* **Nutrientia**, **Vasa vasorum**—*v.* **Nutritia**, **Vasa vasorum**—*v.* **Recta**, see **Rete testis**—*v.* **Resorbentia**, Absorbent vessels, Lymphatic vessels—*v.* **Serosa**, see Capillary vessels.

VASA VASORUM, *V nutritiva* seu *nutrientia*. Small vessels that supply larger. However minutely a vessel is traced, its parietes will be found supplied with blood for its nutrition by a smaller vessel.

VASA VORTICOSA, (F.) *Vaisseaux tourbillonnés*, *Tourbillon vasculaire*. The contorted vessels which creep on the choroid coat of the eye; ciliary veins.

VASCOA AMPLEXICAULIS. A South African shrub, *Nat. Ord.* Leguminosæ, the roots of which taste like liquorice. They are used, in decoction, in catarrh and phthisis, and are a good substitute for liquorice.

VASCULA SERPENTINA, see Testicle.

VASCULAR, *Vascularis*, *Vasculous*, *Angéal*, (F.) *Vasculaire*, from *vas*, *vasis*, 'a vessel,' *vasculum*, 'a little vessel.' That which belongs or relates to vessels;—arterial, venous, or lymphatic; but generally restricted to blood-vessels only. Full of vessels. Bichat gave the name *Vascular system*, *Système vasorum*, *Hæmatanlica*, to the blood-vessels, and of this he made two divisions. 1. The *Arterial system*, or *System of red blood*, (F.) *Système vasculaire à sang rouge*, which commences with the radicles of the pulmonary veins; and includes the pulmonary veins, the left cavities of the heart, and the aorta with all its ramifications. The blood, in this system, is distributed to the general capillary system, where it loses its arterial quality and becomes venous. 2. The *Venous system* or *Vascular system of black blood*, (F.) *Système vasculaire à sang noir*, carries the blood from every part of the system to the lungs. It takes its origin in the general capillary system; continues in the veins, right auricle, and ventricle of the heart and the pulmonary artery and its branches, and is finally lost in the capillary system of the lungs. See Circulation.

VASCULAR LAYER, see *Tache embryonnaire*—*v.* **Radicals**, Radical vessels—*v.* **Substance of the Kidney**, see Kidney—*v.* **System**, see *Vascular*—*v.* **System**, intermediate, Capillary system—*v.* **System**, peripheral, Capillary system—*v.* **Tumour**, see Hæmorrhoids.

VASCULARITY, *Vascularitas*, (F.) *Vascularité*. The state of being vascular, or largely supplied with vessels.

VASCULUM, Penis, Vessel.

VASCULUM ABERRANS, *Appendix to the Epididymis*. A small convoluted duct, of variable length, generally connected with the duct of the epididymis, immediately before the commencement of the *vas deferens*.

VASE À SAIGNER, Cup.

VASTUS. That which is *vast* or has a great extent.

VASTUS EXTERNUS and **VASTUS INTERNUS** are

two considerable, fleshy masses, which form part of the *Triceps cruris*.

VASUM, Vessel.

VATER, AMPULLA OF, (F.) *Ampoule de Vater*. An ampulla formed by the orifice of the pancreatic duct in the duodenum.

VATER, CORPUSCLES OF, Corpuscles, Pacinian.

VATERIA INDICA, see Copal.

VATICINIUM CHIROMANTICUM, Chiromancy.

VAULT, (old F. *Voulte*), 'a round or turn;' from *volvere*, *volutum*, 'to turn;' *Fornix*, *Cam'era*, (F.) *Voûte*. A name given by anatomists to several parts which are rounded superiorly; concave and arched inferiorly,—as the *Vault of the cranium* or the upper part of the skull,—the *Palatine vault* or *roof of the mouth*, &c., &c.

VAUQUELINE, *Strychnia*.

VEAL TEA. This may be made in the same manner as beef tea, by using a pound of fillet of *veal*, free from fat and sliced, and a pint and a half of *boiling water*; and boiling for half an hour. It may also be made with the same quantity of the fleshy part of a knuckle of veal.

VECORDIA, Idiotism.

VECTIS, Lever—*v.* **Elevatorius**, Elevator—*v.* **Triploides**, Elevator, triploid.

VECTOR CANAL, Tube, Fallopiun.

VEGETALITY, *Vegetalitas*, (F.) *Végétalité*. The aggregate of physiological phenomena, which exist alone in the vegetable, but are common to both plants and animals. See Animality.

VEGETARIAN, see Vegetarianism.

VEGETARIANISM. Same etymon as Vegetation. A modern term, employed to designate the view, that man, for his full mental and corporeal development, ought to subsist on the direct productions of the vegetable kingdom, and totally abstain from flesh and blood. An embracer and practiser of this doctrine is called a *vegetarian*.

VÉGÉTATIF, Vegetative.

VEGETATION, *Vegetatio*, 'growing like a plant,' from *vegeo* or *vigeo*, 'I flourish.' A morbid production, which rises as an excrescence on the valves of the heart, in syphilis, &c.—as the *Cauliflower excrescence*, &c. The term is also applied to the fleshy granulations or *Hypersarcoses*, which sometimes arise at the surfaces of wounds or ulcers.

VEGETATIVE, *Vegetativus*, (F.) *Végétatif*, same etymon as the last. Having relation to growth or nutrition; hence, the *vegetative* or *nutritive* functions.

VEGETATIVE LAYER, see *Tache embryonnaire*.

VEGETATIVUS, Vegetative.

VEHICLE, *Vehiculum*, *Oche'na*, *Constit'uena*, (F.) *Véhicule*, from *vehere*, 'to carry.' Anything which serves to carry; thus air is the *vehicle* of sound; the arteries are the *vehicles* of the blood, &c. In *Pharmacy*, the name *vehicle* is given to any excipient; or to substances which serve as the media of administration for any medicine.

VÉHICULE, Vehicle.

VEILED PUFF, see *Cavernous respiration*.

VEIN, *Vena*, diminutive *ven'ula*, *Phlebs*, *Phleps*, (F.) *Veine*. The veins are vessels for the conveyance of black blood from every part of the body to the heart. They are found wherever there are arteries, and, altogether, form the *venous system*, which may be subdivided into two distinct secondary systems—1. The *General venous system*, which commences in all the organs, by very minute radicles; and terminates in the heart by the cava and the coronary vein. 2. The *Abdominal*

venous system, which is limited to the abdomen : cavity; commences, also, by a great number of branches, and terminates in the liver by a single trunk, which subdivides in that organ. It is called, also, the *System of the Vena Porta* or the *Portal System*. See *Porta*. The veins form two sets, one *deep-seated*, *Venæ satel'ites*, *V. concom'ites*, which accompany the arteries; the other *superficial* or *subcutaneous*. Their parietes, which are thinner than those of the arteries, are composed of two coats; the one, external, loose, extensible, and with parallel, longitudinal fibres—the *proper membrane*; the other, the *common* or *inner membrane*, which is thin, polished, and has numerous duplicatures or folds forming parabolic valves. These valves are most numerous in the veins of the lower extremities. The coats of the veins are supplied with small arteries or *vasa vasorum*, veins, and lymphatics. They have not as many nervous filaments as the arteries. These filaments are from the great sympathetic.

TABLE OF THE VEINS.

1. VEINS WHICH FORM THE SUPERIOR VENA CAVA.

1. Subclavians.	Receive the	1. Axillary which	Receives the	1. Posterior ulnar.
	1. Axillary which	1. Axillary which	1. Basilic, formed of the	2. Anterior ulnar.
	2. Cephalic, forming the	2. Cephalic, forming the	3. Circumflex veins.	3. Median basilic.
2. External Jugular.	1. Axillary which	1. Axillary which	4. Inferior scapular.	4. Inferior scapular.
	2. Cephalic, forming the	2. Cephalic, forming the	5. Long thoracic.	5. Long thoracic.
	3. Circumflex veins.	3. Circumflex veins.	6. Superior thoracic.	6. Superior thoracic.
3. Internal Jugular.	1. Axillary which	1. Axillary which	7. Acromial veins.	7. Acromial veins.
	2. Cephalic, forming the	2. Cephalic, forming the	8. Pterygoid.	8. Pterygoid.
	3. Circumflex veins.	3. Circumflex veins.	9. Spheno-palatine.	9. Spheno-palatine.
4. Inferior Jugular.	1. Axillary which	1. Axillary which	10. Alveolar.	10. Alveolar.
	2. Cephalic, forming the	2. Cephalic, forming the	11. Infra-orbital.	11. Infra-orbital.
	3. Circumflex veins.	3. Circumflex veins.	12. Mental.	12. Mental.
5. Inferior Jugular.	1. Axillary which	1. Axillary which	13. Inferior dental.	13. Inferior dental.
	2. Cephalic, forming the	2. Cephalic, forming the	14. Deep temporal.	14. Deep temporal.
	3. Circumflex veins.	3. Circumflex veins.	15. Middle temporal.	15. Middle temporal.
6. Inferior Jugular.	1. Axillary which	1. Axillary which	16. Anterior auricular.	16. Anterior auricular.
	2. Cephalic, forming the	2. Cephalic, forming the	17. Transverse of the face.	17. Transverse of the face.
	3. Circumflex veins.	3. Circumflex veins.	18. Cervical cutaneous.	18. Cervical cutaneous.
7. Inferior Jugular.	1. Axillary which	1. Axillary which	19. Trachelo-scapular, &c.	19. Trachelo-scapular, &c.
	2. Cephalic, forming the	2. Cephalic, forming the	20. Superior Cerebral Veins.	20. Superior Cerebral Veins.
	3. Circumflex veins.	3. Circumflex veins.	21. Veins of the Corpus Striatum.	21. Veins of the Corpus Striatum.
8. Inferior Jugular.	1. Axillary which	1. Axillary which	22. Veins of the Choroid plexus.	22. Veins of the Choroid plexus.
	2. Cephalic, forming the	2. Cephalic, forming the	23. Superior Cerebellar Veins.	23. Superior Cerebellar Veins.
	3. Circumflex veins.	3. Circumflex veins.	24. Inferior Cerebellar Veins.	24. Inferior Cerebellar Veins.
9. Inferior Jugular.	1. Axillary which	1. Axillary which	25. Lateral and inferior Cerebral Veins.	25. Lateral and inferior Cerebral Veins.
	2. Cephalic, forming the	2. Cephalic, forming the	26. Lachrymal.	26. Lachrymal.
	3. Circumflex veins.	3. Circumflex veins.	27. Central of the retina.	27. Central of the retina.
10. Inferior Jugular.	1. Axillary which	1. Axillary which	28. Infra-orbital.	28. Infra-orbital.
	2. Cephalic, forming the	2. Cephalic, forming the	29. Ciliary.	29. Ciliary.
	3. Circumflex veins.	3. Circumflex veins.	30. Ethmoidal.	30. Ethmoidal.
11. Inferior Jugular.	1. Axillary which	1. Axillary which	31. Palpebral, and	31. Palpebral, and
	2. Cephalic, forming the	2. Cephalic, forming the	32. Nasal Veins.	32. Nasal Veins.
	3. Circumflex veins.	3. Circumflex veins.	33. Palpebral, and	33. Palpebral, and
12. Inferior Jugular.	1. Axillary which	1. Axillary which	34. Superciliary Veins.	34. Superciliary Veins.
	2. Cephalic, forming the	2. Cephalic, forming the	35. Dorsals of the nose.	35. Dorsals of the nose.
	3. Circumflex veins.	3. Circumflex veins.	36. Superior Coronaries	36. Superior Coronaries
13. Inferior Jugular.	1. Axillary which	1. Axillary which	37. Inferior of the lip.	37. Inferior of the lip.
	2. Cephalic, forming the	2. Cephalic, forming the	38. Several Buccal and	38. Several Buccal and
	3. Circumflex veins.	3. Circumflex veins.	39. Masseteric Veins.	39. Masseteric Veins.
14. Inferior Jugular.	1. Axillary which	1. Axillary which	40. Ranine.	40. Ranine.
	2. Cephalic, forming the	2. Cephalic, forming the	41. Submental, and	41. Submental, and
	3. Circumflex veins.	3. Circumflex veins.	42. Inferior Palatine.	42. Inferior Palatine.
15. Inferior Jugular.	1. Axillary which	1. Axillary which	43. Lingual and Pharyngeal Veins.	43. Lingual and Pharyngeal Veins.
	2. Cephalic, forming the	2. Cephalic, forming the	44. Superior thyroid.	44. Superior thyroid.
	3. Circumflex veins.	3. Circumflex veins.	45. Occipital, and	45. Occipital, and
16. Inferior Jugular.	1. Axillary which	1. Axillary which	46. Veins of the Diploë.	46. Veins of the Diploë.
	2. Cephalic, forming the	2. Cephalic, forming the		
	3. Circumflex veins.	3. Circumflex veins.		

2. Right Internal Mammary Vein.
3. Inferior Thyroid Vein, opening into the Vena Cava, between the two Subclavians.
4. Vena Azygos, which receives the
 1. Right Bronchial.
 2. Intercoastal Veins.
 3. Semi-azygos.

2. VEINS WHICH FORM THE INFERIOR VENA CAVA.

1. Common Iliac.	Receive the	1. External Iliac, formed by the—	2. Popliteal, commencing by 3. veins which accompany the fibular arteries, and receiving the—	1. External Saphena and 2. Internal Saphena, which receive—	1. Several abdominal veins. 2. Circumflex Iliac, and 3. External pudic veins.
	2. Internal Iliac which receive the—	2. Internal Iliac which receive the—	3. Vesical Veins, commencing with the 2. Sacro-lateral Veins.	1. Dorsal veins of the penis of the male. 2. Veins of the clitoris, in the female.	
	3. Middle Sacral Vein.	3. Middle Sacral Vein.	4. Lumbar Veins—four in number on each side, commencing by an—	1. Abdominal branch. 2. Dorsal branch.	
2. Common Iliac.	5. Spermatc Veins, commencing with the	5. Spermatc Veins, commencing with the	6. Renal Veins.	1. Spermatc plexus in the male. 2. Those of the Ovarium, Fallopian tube, &c. in the female.	
	6. Renal Veins.	6. Renal Veins.	7. Capsular and Adipose Veins.		
	7. Capsular and Adipose Veins.	7. Capsular and Adipose Veins.	8. Hepatic Veins.		
3. Common Iliac.	9. Middle, 10. Left, and 11. Right	9. Middle, 10. Left, and 11. Right	12. Inferior Diaphragmatic Veins, two in number.		
	12. Inferior Diaphragmatic Veins, two in number.	12. Inferior Diaphragmatic Veins, two in number.			

3. VEINS OF THE HEART.

1. Great right Coronary.
2. Small right Coronary.
3. Left Coronary Veins.

4. VEINS WHICH FORM THE VENA PORTA.

1. Splenic Vein.
 - Receives the
 1. Veins which correspond to the vasa brevia.
 2. Right and left gastro-epiploic.
 3. Duodenal, and
 4. Pancreatic veins.
 5. Coronary Vein of the Stomach, and
 6. Small mesenteric vein.
2. Superior Mesenteric Vein.

VEIN, BRONCHIAL, LEFT, see Azygos—v. Demi-azygos, see Azygos—v. Frontal, see Facial vein—v. Semi-azygos, see Azygos (vena)—v. Superior intercostal, left, see Azygos—v. Tibio-malleolar, Saphena, great—v. Palato-labial, Facial vein—v. Peroneo-malleolar, Saphena minor—v. Varicose, Varix.

VEINE, Vein—v. Cave, Cava vena—v. Céphalique, Jugular vein, internal—v. Céphalique du ponce, Cephalic vein of the thumb—v. Cubitale cutanée, Basilic vein—v. Fessière, Gluteal vein—v. de Médine, Dracunculus—v. Porte, Porta vena—v. Prélombo-thoracique, Azygos, (vena)—v. Préparate, Facial vein—v. Radiale cutanée, Cephalic vein—v. Sans paire, Azygos, vein—v. Trachélo-sous-cutanée, Jugular vein, external.

VEINES ARTÉRIELLES, Arterial veins—v. Ciliaires, Ciliary veins—v. Sous-clavières, Subclavian veins—v. Sus-hépatiques, Hepatic veins. VEINEUX, Venous.

VEINS, CORONARY, Cardiac veins—v. Intralobular, Hepatic veins—v. Supra-hepatic, Hepatic veins.

VEINSTONE, Phlebolite.

VEINULE, Venula.

VEJUCO DE MAVACURE, see Curare.

VELAMEN NATIVUM, Integument.

VELAMENTA BOMBYCINA, Villous membranes—v. Cerebralia, Meninges—v. Infantis, Membranes of the foetus.

VELAMENTUM ABDOMINALE, Peritoneum—v. Corporis commune, Integument—v. Linguae, Glosso-epiglottic ligament—v. Nativum, Integument.

VÉLAR, Erysimum.

VELLICATION, Itc.

VELOSYNTHESIS, Staphylorraphy.

VELUM, 'a veil, a cover,' Velum pendulum palati — v. Apophysi vermiformi obtentum, Valvula Vieussenii — v. Interjectum cerebelli, Valvula Vieussenii — v. Medullare anticum, Valvula Vieussenii.

VELUM MEDULLA'RE INFERIUS, (F.) *Voile médullaire inférieur*. A name given, by Reil, to the medullary band or strip, by which the cerebellum communicates with the medulla.

VELUM MEDULLA'RE POSTE'RIOUS seu POSTICUS seu SEMILUNA'RE, *Valvula Tari'ni*. The small posterior valve of the cerebellum; a thin medullary plate, situate between the medulla oblongata and vermis inferior, in front of the nodulus.

VELUM PALATI, Velum pendulum palati — v. Palatinum, Velum pendulum palati.

VELUM PENDULUM PALA'TI, 'Pendulous veil of the palate,' *Velum*, *V. Palati*'num seu *Palati* seu *Pen'dulum* seu *Staphyli'num*, *Pala'tum pen'dulum* seu *mollè* seu *mob'ile*, *Pen'dulum* seu *Clastrum Palati*, *Palatum Mollè*, *Soft Palate*, (F.) *Septum Staphylin* (Ch.), *Voile du Palais*. A movable, soft, and broad septum, attached to the posterior extremity of the palatine arch, and separating the mouth from the pharynx. Its form is nearly quadrilateral. Its anterior surface corresponds to the mouth; the posterior to the pharynx. Its superior edge is fixed to the palatine arch; the inferior is free, floating, and has an appendix at its middle — the *uvula*. The lateral margins of the velum are held down by two folds, separated from each other by a triangular space; these are called the *pillars*; the anterior passes to the base of the tongue, the posterior to the sides of the pharynx. The velum palati is formed of a mucous coat, which contains numerous follicles, and of a muscular stratum, in which are the peristaphylini interni and p. externi muscles, the glosso-staphylini, the pharyngo-staphylini and the palato-staphylini. The arteries of the velum palati are furnished by the internal maxillary, the labial and the superior pharyngeal. Its veins open into the internal jugular. Its nerves are sent off from the ganglion of Meckel, and come from the palatine branches. The glosso-pharyngeal nerve, likewise, sends it some filaments.

VELUM STAPHYLINUM, Velum pendulum palati — v. Vieussenii, Valvula Vieussenii.

VELVET LEAF, Abutilon Avicennæ, Pareira.

VENA, Vein, Penis—v. Alba renum, Ureter—v. Alba thoracis, Thoracic duct—v. Apoplectica, Jugular vein, internal—v. Arteriosa, Porta vena, Pulmonary artery—v. Capitis, Cephalic vein—v. Cava, Cava vena—v. Cubiti interior, Basilic vein—v. Demiazgygos, see Azygos—v. Falcis cerebri, Sinus longitudinalis inferior—v. Hemiazgyga, see Azygos—v. Lactea, Porta vena—v. Lienalis, Splenic vein—v. Magna, Porta vena—v. Medina Arabum, Dracunculus—v. Medinensis, Dracunculus—v. Ostiaria, Porta vena—v. Pari carens, Azygos, (vena)—v. Portalis, Porta vena—v. Portarum, Porta vena—v. ad Portas, Porta vena—v. quæ ad Portas est, Porta vena—v. Ramalis, Porta vena—v. sine Pari, Azygos, (vena)—v. Splenica, Splenic vein—v. Stelechiza, Porta vena—v. Subalaris, Vena axillaris—v. Terminalis, Circulus venosus.

VENÆ APOPLECTICÆ, Jugular veins — v. Cavæ hepatis, Hepatic veins—v. Centrales, Intralobular veins—v. Concomites, see Vein—v. Galeni, Choroid, (veins)—v. Innominatæ, see Innominatum—v. Intralobulares, Intralobular veins—v. Lactææ, Chyliferous vessels—v. Lymphaticæ, Lymphatic vessels—v. Micantes, (arteries,) see Artery—v. Pulsatiles, (arteries,) see Artery—

v. Satellites, see Vein—v. Sororales, Jugular veins—v. Sphagitides, Jugular veins.

VENÆSECTIO, see Blood-letting—v. Revulsoria, see Blood-letting.

VENÆ VORTICO'SÆ. The veins which principally form the external or venous layer of the choroid coat of the eye; so called from their peculiar arrangement; from *vertex* seu *vortex*, 'a whirlpool.'

VENAL, Venous.

VENEFICIUM, Poisoning.

VENENATION, Poison, Poisoning, Venom.

VENENATUS, Venomous.

VÉNÉNEUX, Poisonous.

VENENOSE, Venomous.

VENENUM, Poison, Venom.

VENEREAL, *Vene'reus*, *Aphrodisiacus*, *Vene'rean*, (F.) *Vénérien*, from *Venus*, *veneria*, 'the goddess of pleasure.' That which relates to the pleasures of love. In a more restricted sense, the word *venereal*—as well as *aphroditic*—is synonymous with *syphilitic*, and also with *syphilis*. Some authors have proposed that those diseases should be called *venereal* which are produced by excess in venery; and that *syphilitic* should be limited to those which are the result of impure connexion. It is, also, used in the same sense as aphrodisiac,—adapted to excite venereal desire.

VENEREAL DESIRE, Libido — v. Disease, Syphilis.

VÉNÉRIEN, Venereal. This term, also, means one labouring under syphilis; as *Hôpital des vénériens*.

VENERIVAGUS, Libidinous.

VENERY. Same etymon. The pleasures of love. Sexual communication. Coition.

VENESECTIO, see Blood-letting.

VENIMEUX, Venomous.

VENIN, Venom.

VENOM, *Vene'num*, *Tox'icum*, *Venenat'ion*, (F.) *Venin*. A poisonous fluid secreted by certain animals, as the viper, in a state of health, and which they preserve in a particular reservoir, to use as a means of attack or defence. It differs, according to some, from *virus*, as the latter is always the result of a morbid process, which may be produced by the individuals who receive it, in their turn, and be transmitted to others.

VENOM, QUICKENING, Sperm—v. Vital, Sperm.

VENOMOUS, *Venenat'us*, *Ven'enosus*, (F.) *Venimeux*. An epithet applied to animals which have a secretion of venom, as the viper, rattlesnake, &c., as well as to the venom itself; and, by some, to liquids in the animal body, which have been so perverted by previous disease, that their contact occasions serious mischief in sound individuals; as happens in hydrophobia. *Virus* is more commonly used in this case.

VENOSITAS, Venosity — v. Sanguinis, Prædominium sanguinis venosus.

VENOSITY, *Venos'itas*, *Ve'nousness*, *Venous state* or *cachex'ia*, *Melano'mia*, *Melano'sis universalis*, from *vena*, 'a vein.' A condition in which, it has been supposed, the blood is moved slowly; is more venous; and the venous blood itself in greater quantity. This state is said to be chiefly observed in hemorrhoids, gout, hypochondriasis, melancholia, &c.—Puchelt.

VENOUS, *Venal*, *Ven'ous*, (F.) *Veineux*. Same etymon. Appertaining to a vein, or to the blood of the veins; as *venous cachexia*, *venous pulse*.

VENOUS CACHEXIA, Venosity — v. Hum, *Bruit de Diable*—v. State, Venosity—v. System, see Vascular—v. System, abdominal, Porta vena.

VENOUS SYSTEM OF JACOBSON. A system of vessels in the three classes of vertebrated animals other than the mammalia, which forms a

communication between the vena porta and the vena cava.

VENOUSNESS, Venosity.

VENT, Wind.

VENTER, Abdomen, Belly, Uterus. In law, venter designates the maternal parentage of children. Thus, if A were B's child by his first wife, he would be described in law as "by the first venter." "A died seized, leaving two infant daughters by different venters."

VENTER IMUS, Abdomen, Hypogastrium, Neïsera—v. Inferius, Abdomen—v. Infimus, Abdomen—v. Magnus, Inguivies—v. Medius, Thorax—v. Parvus, Hypogastrium—v. Propendens, Hysteroloxia anterior—v. Renum, Pelvis of the kidney—v. Supremus, Cavum cranii.

VENTEUX, Windy.

VENTILATION, *Ventila'tio*, from *ventus*, 'wind,' *ventulus*, 'a little wind.' The act of renewing the air of vessels, hospitals, and apartments, or of situations where many people are collected together. The utility of such renewal must be evident, when it is considered that atmospheric air loses its oxygen during respiration, and that carbonic acid supplies its place. Stagnant air, also, becomes loaded with numerous impurities; and hence one of the greatest hygienic improvements, of modern times, has been a proper attention to circulation of air. It is, moreover, one of the most valuable additions to our stock of therapeutical agencies in many diseases.

Instruments used for the purpose of renewing the air, in ships, apartments, &c., have been called ventilators.

VENTOSITÉ, Flatulence.

VENTOSITY, Flatulence.

VENTOUSES SCARIFIÉES, Cupping with the scarificator.

VENTOUSES SÈCHES, Cupping, dry.

VENTRAL, from *venter*, 'the belly.' Abdominal.

VENTRAL HERNIA, *Her'nia ventra'lis*. Direct hernia occurring in any part of the abdomen. It is most frequently found between the recti muscles.

VENTRALE, 'an apron,' Belt (Russian.)

VENTRALE CUTANEUM, 'Cutaneous apron.' (F.) *Tablier*. An elongation of the nymphæ in the Bosjesman female, which hangs down before the vulva like an apron.

VENTRE, Abdomen—v. *Inférieur*, Abdomen—v. *Resserré*, Constipation.

VENTRE INSPICIENDO DE, see *De ventre inspicendo*.

VENTRICLE, *Ventric'ulus*, diminutive of *venter*, 'belly.' (F.) *Ventricale*. Literally, a little belly. A name given, in anatomy, to various parts.

VENTRICLE OF ARAN'TIUS. A small cavity, lined with gray substance, at the point of the calamus scriptorius.

VENTRICLES OF THE BRAIN, *Ventric'uli seu Cavita'tes seu Sinus Cer'ebr'i*, (F.) *Ventricules du cerveau*, are five cavities in the interior of that organ, which are distinguished into the lateral ventricles, middle ventricle, fourth ventricle or ventricle of the cerebellum, and fifth ventricle. The middle ventricle or third ventricle, (F.) *Ventricule moyen du cerveau*, is situate in front of the pineal gland. It is bounded, above, by the tela choroidea and the fornix; below, by a portion of cerebral substance, which separates it from the base of the cranium; laterally, by the thalami nervorum opticozum; anteriorly, by a medullary cord, called the anterior commissure; and, posteriorly, by another medullary cord, called the posterior commissure. It communicates with the

lateral ventricles by two openings, situate behind the anterior pillar or crus of the fornix. The lateral ventricles, *Ventric'uli latera'les seu magni seu majo'res seu prio'res seu superio'res seu trico'rne*, (F.) *Ventricules latéraux*, are two in number. They are large, and, owing to their considerable curvature, occupy a great part of the cerebral centre. They are bounded, above, by the inferior surface of the corpus callosum; anteriorly, by the curved portion of the same body; posteriorly, by a considerable mass of cerebral substance; and, internally, they are separated from each other by the septum of the ventricles, — a medullary plate, which is given off from the lower part of the corpus callosum. The lateral ventricles have, in their upper and anterior portion, the *Corpora striata*, *Thalami optici*, and *Tania semicircularis*: — their inferior portion contains the *corpus fimbriatum*, *cornu Ammonis*, &c. Each of the ventricles has a body or centre — *cella me'dia* — and horns, *cornua*. The Fourth ventricle, *Ventriculus cerebelli seu quartus seu Aran'tii*, *Sinus rhomboï'deus*, (F.) *Quatrième ventricule*, is situate beneath the aqueduct of Sylvius, by which it communicates with the third ventricle. It extends from this aqueduct to the upper part of the medulla; and is seated in the substance of the tuber annulare. The Fifth ventricle, *Ventric'ulus septi pellu'cidi*, *Inciue'ra septi*, (F.) *Cinquième ventricule*, is the small serous cavity between the laminae of the septum lucidum; called, also, *Sinus of the septum lu'cidum*, *S. of the median septum* (Ch.).

VENTRICLES OF THE HEART, *Ventric'uli cordis*, *Specus cordis*, (F.) *Ventricules du cœur*, are two in number. The Right ventricle, *Pulmonary ventricle*, *Ventric'ulus anterior seu primus seu pulmona'lis*, (F.) *Ventricule droit ou pulmonaire*, is situate at the fore-part of the heart, the base and apex corresponding to those of the heart. In it are seen the *Columnæ carneæ* and *Cordæ tendineæ*. The right ventricle communicates with the right auricle by the *Ostium venosum*, and from it the *Pulmonary artery* arises. Around the auricular orifice is a tendinous margin or ring, from which the tricuspid valve is given off. The use of the ventricle is to send the blood, by its contraction, along the pulmonary artery to the lungs. The Left ventricle, *Ventric'ulus aorticus seu posterior seu secundus*, (F.) *Ventricule gauche ou aortique*, is situate at the posterior and left part of the heart. Its parietes are much thicker and stronger than those of the right, owing to its having to send the blood over the whole system. The inner surface has the same general appearance as that of the right. The *Columnæ carneæ* and *Cordæ tendineæ* are, however, much stronger. It communicates with the left auricle by means of the *Mitral valve*; and from it proceeds the largest blood-vessel of the body — the aorta.

VENTRICLES OF THE LARYNX, *Ventric'uli seu Sac'culi larynge'i*, *Ventric'uli Morgagn'i*, *Sinu'ses of the larynx*, (F.) *Ventricules ou Sinus du larynx*, are two depressions in the larynx, comprised between the superior and inferior ligaments.

VENTRICLE, PULMONARY, V. right.

VENTRICULAR, *Ventricula'ris*, (F.) *Ventriculaire*. Same etymon as Ventricle. Relating to a ventricle.

VENTRIGULE, Ventricle.

VENTRICULE AORTIQUE, Ventricle, left — v. *Cinquième*, Ventricle, fifth — v. *Droit*, Ventricle, right — v. *Gauche*, Ventricle, left — v. *Moyen du Cerveau*, Ventricle, middle, of the brain — v. *Pulmonaire*, Ventricle, right — v. *Quatrième*, Ventricle, fourth.

VENTRICULES DU CERVEAU, Ventricles of the brain — v. *du Cœur*, Ventricles of the heart — v. *Latéraux*, Ventricles, lateral.

VENTRICULI CORDIS, Ventricles of the heart—*v. Dilatatio*, Gastrectasis—*v. Dissolutio*, Gastro-malacia—*v. Duræ matris*, Sinuses of the dura mater—*v. Emolliatio*, Gastro-malacia—*v. Exulceratio*, Gastrectosis—*v. Fluxus*, Gastrorrhœa—*v. Inflammatio*, Gastritis—*v. Laryngei*, Ventricles of the larynx—*v. Laterales*, see Ventricles, lateral, of the brain—*v. Magni*, Ventricles, lateral, of the brain—*v. Majores*, Ventricles, lateral—*v. Metamorphosis gelatiniformis*, Gastro-malacia—*v. Morgagni*, Ventricles of the larynx—*v. Perforatio*, Gastro-brosis—*v. Priores*, Ventricles, lateral—*v. Pseudophlogosis resolutiva et colliquativa*, Gastro-malacia—*v. Superiores*, Ventricles, lateral—*v. Tricornes*, Ventricles, lateral—*v. Uleeratio*, Gastrectosis.

VENTRICULUS, Stomach, Ventricle—*v. Anterior*, Ventricle, right—*v. Aorticus*, Ventricle, left—*v. Arantii*, see Ventricles of the brain.

VENTRICULUS CALLOSUS, 'Callous stomach.' The gizzard or third stomach of the gallinaceous bird. (F.) *Gésier*. It consists of a double-bellied or *digestive* muscle; is lined by a firm cuticle, and forms a kind of pouch, depending from the alimentary tube, in which the second stomach terminates, and from which the duodenum or *Ventriculus succenturiatus* issues. In the interior of the gizzard there are numerous pebbles, which seem to be destined to aid in the trituration of the food. The gizzard is an organ of mastication, and is possessed of great muscular power.

VENTRICULUS CEREBELLI, see Ventricles of the brain—*v. Intestinalis*, Abomasus—*v. Posterior*, Ventricle, left—*v. Primus*, Ventricle, right—*v. Pulmonalis*, Ventricle, right—*v. Quartus*, see Ventricles of the brain—*v. Secundus*, Ventricle, left—*v. Septi Pellucidi*, see Ventricles of the brain—*v. Succenturiatus*, Duodenum.

VENTRILOQUE, Engastrimythos.

VENTRILOQUISM, see Engastrimyth.

VENTRILOQUIST, Engastrimyth.

VENTRILOQUY, see Engastrimyth.

VENTRIOSUS, Abdominous.

VENTRIPOIENT, Abdominous.

VENTROSITAS, Physcinia.

VENTROSUS, Abdominous.

VENTS, Flatulence.

VENTUS, Wind.

VENULA, *Phlebon*, diminutive of *Vena*. See Vein. (F.) *Veinule*. A small vein. Hippocrates seems to use it in the same sense as *Arteria*.

VENULÆ CENTRALES, Intralobular veins—*v. Intralobulares*, Intralobular veins.

VENUS, Clitoris, Coition, Cuprum—*v. Crystals of*, Cupri subacetat—*v. Crystaux de*, Cupri subacetat.

VER CUCURBITIN, *Tænia solium*—*v. Cutané*, Dracunculus—*v. Filaire*, Dracunculus—*v. de Guinée*, Dracunculus—*v. Luisant*, Cicindela—*v. de Médecine*, Dracunculus—*v. Solitaire*, Bothrioccephalus latus, *Tænia solium*.

VERATRIA, Veratrina.

VERATRINA, *Verat'ria*, *Veratrine*; from *veratrum*. A vegetable alkali, found in the seeds of *Veratrum sabadilla*, *V. officinale*, *Helonias officinale*, and *Asagæa officinale*, which are known in commerce under the names *Cevadilla*, *Cevadilla*, and *Sabadilla*, and are imported from Mexico. They usually occur, in commerce, mixed with the fruit of the plant, are two or three lines long, of a black colour, and are shining, flat, shrivelled, winged, and elastic seeds. *Veratrine* is solid, white, pulverulent, and inodorous; of a very acrid taste; decomposable by heat; very soluble in water, and soluble in alcohol. When prepared in the usual manner, it still contains other substances,—*Sabadillin*, *Veratrin*, gum resin of the *sabadilla*, and black greasy matter.

It is extremely poisonous, and but little used internally in medicine. Magendie gave it as an emetic and drastic in very small doses. It has been recommended to be applied in the way of friction, or endermically, (from gr. v to xx to 3j of lard or cerate,) in various nervous affections. The following is the formula for its preparation in the Pharmacopœia of the United States:—Take of *cevadilla*, lbij; alcohol, 2 gallons; sulphuric acid, solution of ammonia, purified animal charcoal, and magnesia, each a sufficient quantity. Boil the *cevadilla* in a gallon of the alcohol, in a retort with a receiver attached, for an hour, and pour off the liquor. To the residue add another gallon of the alcohol, together with the portion recently distilled; again boil for an hour, and pour off the liquor. Repeat the boiling a third time with the remaining alcohol, and with that distilled in the previous operation. Press the *cevadilla*, mix and strain the liquors, and, by means of a water-bath, distil off the alcohol. Boil the residue three or four times in water acidulated with sulphuric acid, mix and strain the liquors, and evaporate to the consistence of syrup. Add magnesia in slight excess, shake the mixture frequently, then express and wash what remains. Repeat the expression and washing two or three times, and having dried the residue, digest it with a gentle heat several times in alcohol, and strain after each digestion. Distil off the alcohol from the mixed liquor, boil the residue for fifteen minutes in water with a little sulphuric acid and purified animal charcoal, and strain. Having thoroughly washed what remains, mix the washings with the strained liquor, evaporate with a moderate heat to the consistence of syrup, and then drop in as much solution of ammonia as may be necessary to precipitate the *veratrin*. Lastly, separate and dry the precipitate. See *Colchicum autumnale*.

VERATRINE, Veratrina.

VERATRINE, SULPHATE OF, *Verat'rie sulphas*, (F.) *Sulfate de Veratrine*, possesses the same virtues as veratrine. A solution of one grain in two ounces of distilled water has been suggested as a substitute for the *Eau médicinale d'Huon*—a celebrated gout remedy—in the dose of f3j to f3ij.

VERATRUM ALBUM, (*verè atrum*, 'truly black,' from the colour of the roots [?]). *Helleb'orus albus*, *Elleb'orum album*, *Elleborus albus*, *Cherbachem*, *White hellebore*, *Langwoort*, (F.) *Helébore blanc*. *Sex. Syst.* Polygamia Monœcia. *Nat. Ord.* Melanthaceæ (Lindley). The root or rhizoma of this common European plant is indebted to the veratrine for its properties. It is inodorous; has a bitterish, acrid, nauseous taste, and is violently emetic and purgative, even when applied to an ulcerated surface. It is, also, possessed of errhine properties, and acts, externally, as a stimulant. It is rarely exhibited internally. Sometimes it is given as a *sneezing powder*; but it operates with great violence. It forms, also, an itch ointment. See *Ung. Veratri*.

VERATRUM LUTEUM, *Chamaelirium luteum*.

VERATRUM SABADILLA, *Sabadilla*, *Subatilla*, *Causis Intersector*, *Causiticum Americanum*, *Cevadilla Hispanorum*, *Sevadilla*, *Hor'deum causiticum*, (F.) *Cevadille*. The seeds of this plant, *Sabadilla*, *Cevadilla* (Ph. U. S.), have been used, mostly as anthelmintics. Dose, gr. ij to a child four years old. It possesses the same general properties as the *veratrum album*.

VERATRUM VIRIDE, *American Hellebore*, *Swampy hellebore*, *Indian poke*, *Indian Uncas*, *Poke weed*, *Bear weed*, *Itch weed*, *Tickle weed*, *Earth gall*, *Devilbit*, *Wolfbane*, *Puppet root*, is indigenous in the swampy districts of almost every part of the United States. Its properties are like those of

the genus. It has been brought forward, however, in recent times, in the Southern States, as a powerful sedative in inflammatory diseases. Dose, eight drops of a saturated tincture.

VERBASCULUM, *Primula veris*.

VERBASCUM ALBUM, *V. thapsus*—*v. Alatum*, *V. thapsus*—*v. Collinum*, *V. thapsus*—*v. Crassifolium*, *V. thapsus*—*v. Elongatum*, *V. thapsus*—*v. Lychinitis*, *V. thapsus*—*v. Montanum*, *V. thapsus*.

VERBASCUM NIGRUM, (supposed to be a corruption from *Barbasum*, owing to its hairy coat; *barba*, 'a beard.') *Candela regia*, *Thapsus barbatus*, *Thapsus barbatus*, *Candalaria*, *Lanaria*, *Black Mullein*; and VERBASCUM THAPSUS, *V. thapsus*—*v. Alatum* seu *collinum* seu *album* seu *crassifolium* seu *elongatum* seu *lychnitis* seu *montanum* seu *simplex*, *Phlogmus*, *Phlomis*, *Phlox*, great broad-leaved or yellow Mullein, *High Taper*, *Cow's Lungwort*, (F.) *Molène*, *Bouillon blanc*, *Bonhomme*, *Cierge de notre dame*, *Herbe Saint-Fiacre*. *Ord. Scrophulariaceæ*, *Sex. Syst. Pentandria Monogynia*,—especially the latter,—have been used as emollients, applied externally in the form of fomentation. The large leaves have been chosen for this purpose. Occasionally, the mucilage which they contain has been extracted by decoction or infusion, and administered in pectoral affections.

VERBASCUM SIMPLEX, *V. thapsus*—*v. Thapsoides*, *V. thapsus*—*v. Thapsus*, see *V. nigrum*.

VERBENA HASTATA, *V. paniculata*, *Blue vervain*, *Simpler's Joy*, an American species, is more bitter than the European, and said to be emetic.

VERBENA OFFICINALIS, *Verbena*, *Dichromos*, *Verbena*, *Peristephium*, *Hierobotanè*, *Herba sacra*, *Vervain*, *Purvain*, (Prov.) *Berbine*, *Juno's Tears*, (F.) *Verveine*, *Herbe à tous les maux*; *Ord. Verbenacæ*. *Sex. Syst. Diandria Monogynia*. A plant formerly held sacred; used in sacrifices, and suspended round the neck as an amulet. In the form of cataplasm, it is said to have relieved obstinate headaches, and to have been of service in scrofula. It is not used.

VERBENA PANICULATA, *V. hastata*.

VERBENA URticO'RIA, an indigenous species, has been used in poisoning from *Rhus Toxicodendron*.

VERBENACA, *Verbena officinalis*.

VERBESINA ACNELLA, *Spilanthes Acmella*.

VERBESINA VIRGINICA, *Crown Beard*. Indigenous. *Order*, *Compositæ*. A decoction of the roots is used by the Indians as diaphoretic and depurative.

VERBUS, *Penis*.

VERCOQUIN, *Phrenitis verminosa*. A form of phrenitis, attributed to the presence of a worm in the brain!

VERDET, *Cupri subacetat*.

VERDIGRIS, *Cupri subacetat*.

VERENDA, *Genital organs*.

VERETRUM, *Penis*.

VERGE, *Penis*—*v. d'Or*, *Solidago virgaurea*.

VERGETTE DE PHILADELPHIE, *Erigeron Philadelphicum*.

VERGETURES, *Vibices*.

VERJUICE, *Ompha'cium*, (F.) *Verjus*, *Verd-jus*, (*verd*, 'green,' and *jus*, 'juice.') "The juice of green fruits." A sour kind of grape. Also, an acid liquor, prepared from very sour grapes or crab-apples. It is principally used in culinary preparations; although, occasionally, an ingredient in medicinal compounds. See *Vitis vinifera*.

VERJUS, *Verjuice*.

VERMES, *Worms*—*v. Cerebri*, *Choroid plexus*.

VERMICELLI, see *Turundæ Italicæ*.

VERMICULAIRE BRÛLANTE, *Sedum*.

VERMICULAR, *Vermiform*.

VERMICULARIS, *Lumbricalis*, *Sedum*.

VERMICULI SPERMATICI, *Spermatozoa*.

VERMICULUM, *Vermiculum*. An elixir; a tincture.—*Ruland and Johnson*.

VERMICULUS CAPILLARIS, *Dracunculus*.

VERMIFORM, *Vermicular*, *Vermicularis*, *Vermiformis*, (F.) *Vermiculaire*, *Vermiforme*, from *vermis*, 'a worm,' and *forma*, 'form.' That which resembles a worm.

VERMIFORM PROC'ESSES, *Processus seu Pro-tuberantia seu appendices vermiformes*, *Vermes*, *Monticuli cerebelli*, *Worms*, (F.) *Éminences vermiformes du cervelet*, are two medullary projections, at the surface of the cerebellum. The superior vermiform process, *Vermis superior cerebelli*, is an elongated projection, at the anterior and middle part of the superior surface of the cerebellum;—and the inferior vermiform process, *Vermis inferior cerebelli*, *Pyramid of Malacarne*, is a somewhat larger projection, situate in the depression at the inferior surface of the same organ. The inferior vermiform process is usually described as consisting of three portions—the pyramid, the *uvula*, and the *nodulus*. The anterior portion of the superior vermiform process is elevated, and termed *Monticulus*.

VERMIFUGE, *Anthelmintic*.

VERMIFUGE, *MATHIEU'S*. This consisted of two distinct electuaries; the one for killing, the other for expelling tape-worm. The former is made of an ounce of tin filings, and six drachms of *ferri root*.

VERMIFUGUS, *Anthelmintic*.

VERMILION, *Hydrargyri sulphuretum rubrum*.

VERMILION DE PROVENCE, *Antimoni sulphuretum rubrum*.

VERMINATIO, *Helminthiasis*.

VERMINATION, *CUTANEOUS*, *Malis*.

VERMINEUX, *Vermineous*.

VERMINOUS, *Vermineous*, (F.) *Vermineux*. From *vermis*, 'a worm.' That which is caused by worms; as 'a verminous affection.'

VERMIS, see *Vermiform processes*—*v. Bombycinus*, *Cornu ammonis*—*v. Cerebelli*, *Vermiform process*—*v. Cerebri*, *Fever*, *Hungary*—*v. Cucurbitinus*, *Tænia solium*—*v. Digni*, *Paronychia*.

VERMITIO, *Helminthiasis*.

VERNIX CASEO'SA, 'cheesy varnish.' The fatty varnish, or deposit, often found on the fœtus, which is secreted by the sebaceous follicles of the skin.

VERNIX SICCA, *Sandarac*.

VERONICA ANTHELMINTICA, *Ascaricida anthelmintica*, *Calagira*, *Calageri*. A plant which is indigenous in India. *Ord. Compositæ*. *Sex. Syst. Syngenesia Polygamia superflua*. It has great celebrity in the East Indies, and is a bitter tonic; but the seeds are chiefly used as an anthelmintic, in cases of ascarides and lumbrici in children.

VERONICA NOVEBORACENSIS, *Iron weed*, *Flat top*; an indigenous plant; whose purple flowers appear in August and September, is reported to possess cathartic properties.

VÉROLE, *Syphilis*—*v. Petite*, *Varicella*—*v. Petite volante*, *Varicella*.

VÉROLÉ, *Syphilitic*.

VÉROLETTE, *Varicella*.

VÉROLIQUE, *Syphilitic*.

VERONICA, *V. officinalis* seu *Tournefortii* seu *mas*, *Thea Germanica*, *Betonica Pauli*, *Chamaedrys*, *Male Veronica*, *Fluellin*, *Official Speedwell*, *Eighbright*, *Paul's Betony*, (F.) *Véronique*, *Thé d'Europe*, *Herbe aux ladres*. *Ord. Siro-*

phulariaceæ. *Sec. Syst. Diandra Monogynia.* This plant was once used as a pectoral.

VERONICA AQUATICA, V. beccabunga.

VERONICA BECCABUN'GA, *Beccabunga, Anagallis seu veronica aquatica, Laver Germanicum, Cypa'ia, Water Pimpernel, Brooklime, Brooklime Speedwell, Water Speedwell, Neckweed, Water Purslain, (Sc.) Water-purpie, (F.) Véronique cressonée.* This plant was formerly used in a variety of diseases. It is, however, almost insipid and inodorous; and has been long banished from practice.

VERONICA MAS, Veronica — v. *Officialis*, Veronica.

VERONICA PEREGRINA, *Neckweed, Purslain, Speedwell*; indigenous; flowering from April to June, is given internally in scrofulosis, and is applied as a wash to scrofulous tumours.

VERONICA PURPUREA, *Betonica officinalis* — v. *Tournefortii*, Veronica — v. *Virginica*, *Leptandra Virginica*.

VÉRONIQUE, Veronica — v. *Cressonée*, Veronica beccabunga.

VERPA, Penis.

VERRE, Vitrum — v. *d'Antimoine*, Antimonii vitrum.

VERRUCA, diminutive *Verrucula, Ecphy'ma verruca, Phymato'sis verruco'sa, a Wart, (Prov.) Wret, (F.) Verruc, Poireau.* A small, hard, mamillary tumour, which forms at the surface of the skin, and particularly on that of the hands and face. Warts are generally rugous at the surface, and broad at the base; their roots being implanted in the derma by whitish filaments, which are dense, semifibrous, and very numerous.

Warts may be destroyed by caustic, or removed by excision. Their vitality is, generally, not great.

VERRUCA CARNOsa, Condyloma — v. *Formicaria*. Formica — v. Minor, Thyunion — v. *Palpebrarum*, see Hydatid — v. *Pensilis, Acrochordon* — v. *Rhagoidea*, Thyunion.

VERRUCARIA, *Calendula officinalis, Heliotropium Europæum*.

VERRUCOSE, Warty.

VERRUCOUS, Warty.

VERRUCULA, Verruca.

VERRUE, Verruca.

VERS, Worms.

VERSE-CHARMS, see Charm.

VERSIO, Turning — v. *Fœtis*, Turning — v. *Spontanea*, Evolution, spontaneous — v. *Uteri*, Hysteroloxia.

VERSION, Turning.

VERT-DE-GRIS, Cupri subacetat.

VERTEBRA DENTATA, Axis — v. *Dorsi Prima*, Lophia — v. *Lumborum Prima*, Asphalites — v. *Magna*, Sacrum.

VERTEBRÆ, *Spondyli, Spondles, Whetle-bones, (F.) Vertèbres*; from *vertere*, 'to turn.' This name has been given to the bones, which by their union form the vertebral or spinal column. The vertebræ are short, thick, very angular bones; twenty-four in number, and situate one above another. The body—centrum, of Owen—of the vertebra is the middle and anterior portion. Above and below it is articulated, by means of a fibro-cartilage, with the contiguous vertebræ. The Processes, (F.) *Masses apophysaires*, of each vertebra are:—1. The spinous processes, situate behind, and in the median line. 2. The two transverse processes, standing off from each side. 3. The four articular, or oblique processes — *zygapophyses*, of Owen — two superior, and two inferior — which serve to unite the vertebra with each other. These transverse and articulating processes are made continuous with the lateral and posterior parts of the body of the vertebra by narrow bony portions or pedicles, on

each of which are notches. These, by their union with similar notches in the contiguous vertebra, form the lateral foramina, (F.) *Trous de conjugaison*, through which the spinal nerves leave the vertebral canal. The different processes are united with each other, so as to form a kind of ring of the lateral and posterior parts of the vertebra; and the foramen, within this, — that is, between the body and processes, *foramen vertebra'le, rachid'ium*, — concurs in forming the vertebral canal. The arch or lamina of a vertebra, *neurapoph'ysis*, is the part from which the several processes project. The vertebræ have particular characters, according to the regions they occupy. They are divided into seven cervical, twelve dorsal, and five lumbar. The first cervical is called atlas; the second, axis or vertebra dentata; and the seventh, vertebra prom'ens. The vertebræ are articulated with each other. Those of the back are joined, moreover, to the ribs by their articulating surfaces — *parapophyses*. The first or atlas is articulated with the occipital bone; and the last lumbar vertebra with the sacrum. The cervical vertebræ are developed by nine points of ossification; the dorsal and lumbar by eight.

VERTEBRAL, *Vertebra'lis*. That which relates or belongs to the vertebræ.

VERTEBRAL ARTERY, *Posterior cerebral artery*, (Ch.), is the largest of the branches of the subclavian. It arises from the posterior part of that vessel; is lodged in a canal, (F.) *Petit canal vertébral*, formed by the foramina in the transverse processes; enters the cranium, and proceeds to anastomose with its fellow of the opposite side, and to form the Basilar artery. Besides the branches sent to the neck, the vertebral artery gives off the anterior and posterior spinal arteries, and the inferior cerebellous, in the cavity of the cranium.

VERTEBRAL COLUMN, *Rachis, Rhachis, Notoz, Acan'tis, Carina, C. hom'ina, Spina, S. dors' seu vertebra'lis, Colum'na spinal'is seu vertebra'lis seu dors', Dorsum, Tergum, Comp'ges Vertebra'rum, Acan'tia, Sacra fistula, H'era Syrinx, Spine, Back bone, (Sc.) Backsprent, (F.) Colonne vertébrale, Épine du dos, Échine.* A kind of bony column, situate at the posterior and central part of the trunk; extending from the head to the sacrum; flexible in every direction, yet very solid and hollowed by a canal called *vertebral*, which gives it lightness without diminishing its strength, and runs through its whole length. The vertebral column is formed by the superposition of the vertebræ. It represents a pyramid, whose base is below. Anteriorly, it is convex in the neck; concave in the back, and convex, again, in the loins. Behind, the curvatures are of course the reverse of these. There is also, in the dorsal region, a lateral curvature, the concavity of which is towards the left side. Considered as a whole, the vertebral column has an anterior or prepsinal surface; a posterior or spinal; two lateral; a base; and an apex or top. In the nomenclature of Chaussier, the anterior surface, in the neck, is called *trachelian*; in the back, *predorsal*; and in the loins, *prelumbur*. The dorsal surface has, at its middle, the spinous processes; and, at its sides, the vertebral gutters, (F.) *Gouttières vertébrales*, formed by the succession of the vertebral plates, (F.) *Lames vertébrales*, which, by their junction, constitute the spinous processes. These are filled by the sacro-spinalis muscle. The sides or lateral surfaces present the foramina resulting from the union of the vertebral notches, (F.) *Échancrures vertébrales*. These are called *vertebral foramina*, (F.) *Trous de conjugaison*, and give passage to the vertebral nerves. The base

of the vertebral column is oblique, to unite with the sacrum; and forms with it an angle, called *sacro-vertebral angle* or *promontory*. The apex or top of the vertebral column is articulated with the occipital bone. Sometimes, the term *vertebral column* includes the whole of the column from the occipital bone to the extremity of the coccyx. The *Vertebral* or *Spinal* or *Rachidian Canal*, *Solen*, *Canal'is medullæ Spinal'is*, *Spæcis pro medull'a Spinal'i* seu *vertebra'lie*, exists through the whole of the vertebral column, whose curvatures it follows. It is continuous, above, with the cranium; below, with the sacral canal, if we consider the vertebræ to terminate at the sacrum. It is lined by a prolongation of the meninges, forming the *Theca vertebra'lis*, and lodges the spinal marrow.

The vertebral column, which unites lightness with solidity, serves to support the head and chest. It is the seat of all the movements of the trunk, the weight of which it transmits to the pelvis. It lodges and protects the spinal marrow, and gives passage to the vertebral nerves, and to numerous vessels.

VERTEBRAL DISEASE, *Rachial'gia* (of some,) *Caries of the Vertebrae*, *Spondylarthrocacæ*, *Spondylolocacæ*, *Kyp'hosis inflammato'ria* seu *paralytica*, *Gibbositas cariosa*, *Cal'ries vertebra'rum*, *Spondylal'gia*, *Spondylomyel'itis*, *Tuberculo'sis Vertebra'rum*. The spinal column is liable to distortion, (see *Hump*), at times, without the vertebræ being much, if at all, diseased. At other times, a more formidable affection is induced; the vertebræ becoming carious, and the intervertebral cartilages ulcerated; followed by a more or less complete loss of power over the lower extremities. In such case, on examining the spine, one or more of the spinous processes is found to project beyond the rest. The disease was first well described by Mr. Percival Pott; and, hence, it has been called, by the French, *Maladie de Pott*, *Malum* seu *Gibbus Pot'tii*, *Pott's disease* or *curvature*. It usually occurs in scrofulous subjects, and the treatment consists in establishing a counter-irritation by blisters, issues, or the moxa, opposite the transverse processes of the diseased vertebræ. The sore must be suffered to remain open and the patient kept in the horizontal posture until the use of the limbs is recovered; employing tonics, the free admission of fresh air, &c., at the same time.

VERTEBRAL LIG'AMENTS, (F.) *Ligaments vertébraux*, *Surtoits ligamenteux de la colonne vertébrale*, are distinguished into *anterior* and *posterior*. They consist of long, ligamentous coverings; the *anterior* occupying the anterior part of the vertebral column; and the *posterior* seeming to arise from the occipito-axoid ligament, and seated behind the bodies of the vertebræ, from the axis to the sacrum.

VERTEBRAL NERVE, *Trisplanchnic*.

VERTEBRAL NERVES, *Spinal Nerves*, (F.) *Nerfs rachidiens*, spring successively from the spinal marrow; from its origin to its termination. These nerves—30 pairs in all—arise by two roots; one *anterior*, from the *anterior* or *an'tero-lat'eral column* of the spinal marrow, and the other *posterior*, from the *posterior* or *pos'tero-lat'eral column*, and afterwards issue by the vertebral and sacral foramina. Each of these roots seems to have a distinct function; the *posterior* being destined to convey sensation to parts; the *anterior*, motion. On the posterior root, before its junction with the anterior, there is a ganglion. Of the 30 pairs, 7 are *cervical nerves*, 12 *dorsal*, 5 *lumbar*, and 6 *sacral*. See *Nerves*.

VERTEBRATED, *Vertebra'tus*, (F.) *Vertébré*. Provided with vertebræ.

VERTÈBRES, *Vertebrae*.

VERTÉBRODYMIA, see *Somatodymia*.

VERTEBRO-ILIAC LIGAMENT, *Ileo-lum-bar ligament*.

VERTEX, *Cor'yp'hæ*, *Acro'ria*, *Spon'dylus*. The top or highest part or crown of the head (Sc.) *Centel*, *Nom*, *Pallat*, *Palad*. See *Sinciput*.

VERTEX CORDIS, *Mucro Cordis*—v. *Cubiti*, *Olecranon*—v. *Palpitans*, *Fontanella*.

VERTIBULUM, *Trepan*.

VERTICULA, *Articulation*.

VERTICULUM, *Articulation*.

VERTICULUS, *Articulation*.

VERTIGE, *Vertigo*.

VERTIGINOUS, *Vertiginosus*, (Old Eng. and Prov.) *Totty*, (F.) *Vertigineux*. Relating or pertaining to vertigo; *Giddy*, *Dizzy*.

VERTIGO, from *vertere*, 'to turn;' *Dinos*, *Rin'gos*, *Dinus* seu *Hallucina'tio* seu *Aural'gia vertigo*, *Circumgyrat'io*, *Gid'diness*, *Diz'ziness*, *Swimming of the head*, (Prov.) *Swoe*, *Swinnying*, (F.) *Vertige*, *Étourdissement*. A state in which it seems that all objects are turning round; or that the individual himself is performing a movement of gyration. Two varieties have been designated;—*simple vertigo*, where the objects are seen; and *scotodynia*, in which the sight is obscured, and the individual often falls. Vertigo is dependent upon the condition of the brain; and often announces an attack of apoplexy or epilepsy.

VERTIGO TENEBRICOZA, *Scotodynia*.

VERUGA (S.), from *verruca*. A peculiar disease affecting certain localities of Peru, which manifests itself by sore-throat, pains in the bones, and other febrile symptoms. In the course of a few days, an eruption of red pimples or boils appears. These sometimes increase in magnitude till they become nearly as large as an egg, and discharge blood copiously. It is ascribed to the water of certain springs in the valleys, and especially in that of *Sura*: called *Ayua de Veruga*.—*Tschudi*.

VERU MONTANUM, *Gallinaginis caput*.

VERVAIN, *Verbena officinalis*—v. *Blue*, *Verbena hastata*.

VERVEINE, *Verbena officinalis*.

VESA'NIA, *Vesania*, 'madness.' Derangement of the intellectual and moral faculties, without coma or fever. Many nosologists have used this as a generic term; under which they have included the different species of mental alienation, hypochondriasis, somnambulism, and even hydrophobia. See *Nosology*.

VESANIA MANIA, *Mania*.

VESANIE, *Insanity*.

VESICA BILIARIA, *Gall-bladder*—v. *Cordis*, see *Pericardium*—v. *Fellea*, *Gall-bladder*—v. *Intestinalis*, *Vesicula umbilicalis*—v. *Prostatica*, see *Ejaculatory ducts*—v. *Saccata*, *Cystostenochoria*—v. *Urinaria*, *Urinary bladder*.

VESICÆ IRRITABILITAS, *Cysterethismus*—v. *Unguinosæ tendinum*, *Bursæ mucosæ*.

VESICAL, *Vesica'lis*, from *vesica*, 'the bladder.' That which belongs or relates to the bladder.

VESICAL ARTERIES vary in number and origin. The umbilical artery always sends off two or three: the middle hemorrhoidal, internal pudic, and obturator furnish others, which ramify and anastomose in the parietes of the bladder; the hypogastric furnishes one which is somewhat larger, the *Vesico-prostatic* (Ch.), and proceeds to the *bas-fond* of the bladder, sending numerous ramifications to it as well as to the surrounding parts.

VESICAL GLANDS, see *Spermatic Glandion*—

v. Ligaments, see Urinary bladder — v. Oxyd, Oxyd, cystic — v. Triangle, see Urinary bladder.

VESICAL VEINS, much more numerous than the arteries, open into the hypogastric plexus.

VESICANT, Blister, Epispastic.

VESICARIUM, Ovarium.

VESICATION, *Vesicatio*; from *vesica*, 'a bladder.' The action of a vesicant. The formation of blisters.

VÉSICATOIRE, Blister — v. *Magistral*, Magistral.

VESICATORIUM, Blister.

VESICLE, Vesicula — v. Allantoid, Allantois — v. Blastodermic, see Molecule — v. Cowpox, see Vaccination.

VESICLE OF DEVELOPMENT, *V. of Evolution*. A vesicle or cyst concerned in histogeny.

VESICLE OF EVOLUTION, *V. of Development* — v. Germinal, see Molecule — v. Purkinjean, see Molecule — v. Umbilical, Vesicula umbilicalis.

VESICLES, ASCHERSONIAN. When fluid fat and fluid albumen are shaken together, the resulting fat globules become surrounded by an albuminous layer. The vesicles, thus formed, are called after Ascherson, their discoverer.

VESICLES OF DE GRAAF. Folliculi Graafiani — v. Seminal, Vesiculæ seminales.

VESICOCE'LE. Badly compounded for Cystocele.

VESICO-UTERINE, *Vesi'co-uteri'nis*, from *vesica*, 'the bladder,' and 'uterus.' Appertaining, or relating to, the bladder and uterus.

VESICO-UTERINE FOLDS, see Uterus.

VESICO-VAG'INAL. *Vesi'co-vaginal'is*. *Vag'ino-ves'ical*. Relating to the bladder and the vagina; as *vesico-vaginal fistula*.

VESIC'ULA, *Cystinz*, *Ves'icle*, *Blister*, (F.) *Vésicule*; diminutive of *vesica*, 'a bladder.' A small bladder or cyst.

VESICULA, *Ves'icle*, *Ephly'sis*, *Cystinz*, in pathology, means a small orbicular elevation of the cuticle containing lymph, which is sometimes clear and colourless, but often opaque and whitish or pearl-coloured. It is succeeded, either by a scurf, or by a laminated scab.

VESICULA ÆNIGMATOSA, Vesicula umbilicalis — v. Alba, Vesicula umbilicalis — v. Bilis, Gall-bladder — v. Blastodermatica, see Molecule — v. Blastodermica, see Molecule — v. Chyli, Receptaculum chyli — v. Erythroidea, Erythroid vesicle — v. Fellis, Gall-bladder — v. Gangranescens, see Anthrax — v. Germinativa, see Molecule — v. Omphalo-mesenterica, Vesicula umbilicalis — v. Ovalis, Vesicula umbilicalis — v. Prolifera, see Molecule — v. Prostatica, see Ejaculatory ducts.

VESIC'ULA UMBILICAL'IS, *V. alba* seu *omphalo-mesenter'ica* seu *anigmatosa* seu *oval'is*, *Vesi'ca intestinal'is*, *Vitell'ic'us*, *Umbil'ical* or *Intestinal Vesicle*, *Vitelline Pouch*, *Vitell'icle*. A small, pyriform, round, or spheroidal sac, which, about the fifteenth or twentieth day after fecundation, is of the size of a common pea, and probably acquires its greatest dimensions in the course of the third or fourth week. After the seventh week, it becomes shrivelled and disappears insensibly. It seems to be situate between the chorion and amnion, and is commonly adherent either to the outer surface of the chorion, or to the inner surface of the chorion, but at times is loose between them. The *vitelline ped'icle*, which attaches the vesicle to the embryo, is continuous with the intestinal canal; and, up to twenty or thirty days of embryonic life, is hollow, — *Ductus omphalo-mesenter'icus* seu *vitello-intestinal'is* seu *vitell'ic'us*, *vitelline Duct*. Its arteries are the omphalo-mesenteric. The *vitelline fluid*, contained in the vesicle — in the *Vitelline* or *Yolk membrane* — has been compared, from analogy, to the *vitellus* or

yolk of birds: and, like it, is an oleaginous substance, and presumed to be inservient to the nutrition of the embryo.

VESICULÆ constitute the 6th order of Bateman's classification of *Cutaneous Diseases*.

VESICULÆ, see Villous membranes — v. Divæ Barbaræ, Variola confluens — v. Gingivarum, Aphthæ — v. Pulmonales, see Cellule — v. Rotundæ cervicis uteri, Nabothi glandulæ — v. Sanguinis, Globules of the blood — v. Seminales mulierum, Nabothi glandulæ — v. Spermaticæ, Vesiculæ seminales.

VESICULÆ SEMINAL'ES, *V. Semina'ria* seu *spermatic'æ*, *Cap'sulæ semina'les*, *Conceptac'ula semina'ria*, *Gonecys'tides*, *Sem'inal ves'icles*, (F.) *Vésicules séminales*, *Gardouches*, *Gréniers*, are two small membranous sacs, which serve as reservoirs for the sperm, and probably secrete a fluid to be mixed with the sperm. They are situate beneath the bladder, and above the rectum, behind the prostate and to the outer side of the vasa deferentia. They are of an irregular conoidal shape; flattened from above to below; embossed on their surface, and of a grayish tint. Their *posterior extremity* or *fundus* terminates by a round cul-de-sac. Their *anterior extremity* or *neck* is narrow and elongated, and is continuous with the excretory duct of the vesicula. This duct joins the vas deferens, and forms with it the ejaculatory duct. The interior of the vesiculæ exhibits several cells, which communicate with each other, and are lined by a very thin, mucous membrane.

VESIC'ULAR, *Vesicula'ris*, (F.) *Vésiculaire*. Of or belonging to a vesicle:—full of or containing small vesicles or cells. The cortical or gray matter of the brain is sometimes termed *vesicular*. See Cortex cerebri and Neurine.

VÉSICULE, Vesicula — v. *Bilia' e*, Gall-bladder — v. *du Fiel*, Gall-bladder — v. *Germinative*, see Molecule — v. *Ovulifère*, see Folliculi Graafiani.

VÉSICULES DE GRAAF, Folliculi Graafiani — v. *de Sainte Barbe*, Variola confluens — v. *Séminales*, Vesiculæ seminales.

VESIC'ULO-TYMPANIT'IC, *Vesic'ulo-tympanit'icus*. An epithet for a sound or resonance heard on percussion, which consists of the vesicular and tympanitic sounds combined. It occurs when the parenchyma of the lungs is incompletely solidified. — Flint.

VESPA, Wasp.

VESSEL, *Vas*, *Vasum*, diminutive *Vasculum*, *Angel'on*, *Angos*, *Conceptac'ulum*, from (L.) *vas*, *vasculum*, (F.) *Vaisseau*. A canal, more or less elastic, formed by the superposition of membranes, and distinguished, according to its uses and general arrangement, into *Artery*, *Vein*, and *Lymphatic*. See, also, *Vascular*.

VESSELOUP, Lycoperdon.

VESSE, Urinary bladder — v. *à Colonnes*, see Urinary bladder — v. *Col de la Collum vesicæ*.

VESSIE NATATOIRE, Air-bladder.

VESTIBULAR NERVE, see Auditory nerve.

VES'TIBULE, *Vestib'ulum*, from *vestio*, 'to cover.' The porch or entrance to a house.

VESTIBULE, *Vestib'ulum*, *V. Vagi'næ*, *Prona'us*, *Atrium Vagi'næ*, *Amb'itus genit'alis mulie'bris*, is a smooth cavity between the perinaeum and nymphæ in the female, which leads to two passages — to the urethra above, and to the vagina below.

VES'TIBULE OF THE EAR, *Vestib'ulum*, *V. labyrinthi*, (F.) *Vestibule*. A cavity of an irregular shape, which forms part of the labyrinth or internal ear. The vestibule is situate on the inner side of the tympanum; on the outer side of the *mentus auditorius internus*; before the semicircular canals, and behind the cochlea. It is lined

by a particular membrane, and contains the liquor of Cotunnus and branches of the auditory nerve. There is also another membrane, constituting the *membranous vestibule*, but it is not an exact imitation of the osseous cavity. It is composed of two distinct sacs, which open into each other—the *sacculus vestibuli*, and the *sacculus*. There are three contiguous cavities in the vestibule; one of which, the *semi-oval*, is situate above; another, the *hemispherical*, below; and the third, of *culiform*, which is the orifice of the aquæductus vestibuli, behind.

VESTIBULUM LABYRINTHI, Vestibule of the ear—*v. Vaginæ*, Vestibule.

VESTIGE (F.) *Vestig'ium*, 'a vestige,' 'a trace.' A kind of fracture of flat bones, consisting of a simple incision having the mark of the instrument that caused it. The term is also applied to a part of the body, which may have no evident use in man, and yet may be greatly employed in animals. For example, the muscles of the external ear in man are almost devoid of use, or are *vestiges* of what they are in certain animals.

VESTIGIUM, Sole, *Vestige*—*v. Foraminis ovalis*, *Ovalis fossa*—*v. Pedis*, *Metatarsus*.

VESTITUS, *Esthe'ma, Esthe'sia, Res vestia'ria*, *Clothing, Dress, Vesture*, (F.) *Vêtement*, from *vestire, vestitum*, 'to clothe.' Any substance used for the purpose of covering nakedness, or preserving from atmospheric temperature by being worn on the body. The best clothing in winter is composed of articles that are bad conductors of caloric, and do not admit of ready evaporation of the matter of perspiration. In this respect, woollen vestments deserve the first place, and cotton the second. Colour has likewise to be regarded—the darkest hues transmitting the heat most readily through them. It has been found that the same rule applies to odours—the darker colours retaining a greater amount of odorous particles than the lighter. Hence the necessity of white walls, white bed-clothes, &c., in the wards of hospitals; for it is probable that the like law applies to contagious miasms.

VESTURE, *Vestitus*.

VETA, Puna.

VÊTEMENT, *Vestitus*.

VETERANA, *Rubeola*.

VETERINARIUS, see *Hippiater* and *Veterinary*.

VETERINARY, *Veterina'rius*, (F.) *Vétérinaire*, from *veterinus*, 'that which bears burdens;' itself, perhaps, from *veheterinus*, from *vehere*, 'to carry.' That which appertains to beasts of burden: hence, *veterinary surgeon*, (see *Farrier*), *veterinary college*, and *veterinary medicine*.

VETERINARY ANATOMY, see *Anatomy*.

VETERINARY ART, *Ars seu Medic'ina veterina'ria*, *Zoô'i'asis, Zoia'tri'a, Theria'trica, Mulomedici'na, Zoôtherapi'a, Zoôtherapeu'tica, Ars zoia'trica, Far'riery*. The art of treating the diseases, &c., of cattle.

VETERNOSITAS, *Coma vigil*.

VETERNUS, *Coma vigil*.

VETIVERIA ODORATA. *Andropogon muricatus*.

VETTONICA, *Betonica officinalis*.

VETULA, *Sacrum*.

VEXILLA TUBARUM FALLOPII, see *Tube*, *Fallopian*.

VEXILLUM, *Drapeau*.

VIA, Way—*v. Curandi*, *Ratio medendi*—*v. Stomachi et ventris*, *Œsophagus*.

VIABILITY, see *Viable*.

VIABLE, *Viab'ilis*, from *via*, 'way;' or from (F.) *vie*, 'life,' (Orfila.) A word, which may be introduced with great advantage from the French into our language. It is an epithet applied to a *status* whose organs are properly formed, and so

developed as to permit its continued existence. Every *fœtus* is not *viable*. *Anencephali* have lived ten or twelve days, yet they are not viable. The older the *fœtus*, the more viable is it. It is an idle prejudice that a child, born at the end of eight months, is less likely to live than one born at the end of seven. The signs of *viability* are drawn from the weight, length, external conformation, &c., of the *fœtus*. It may be laid down as a general rule, that no *fœtus* born before the end of the seventh month is viable.

VIE CHYLIFERÆ, *Chyliferous vessels*.

VIE CLANDESTINÆ, *Clandestine Passages*. Supposititious secret ways or passages or 'royal roads,' to account for the rapid transmission of substances taken into the stomach to the kidneys or bladder.

VIE EXTRAORDINARIÆ LACTIS, *Galactoplasia*—*v. Lacrymales*, *Lachrymal passages*—*v. Primæ*, *Ways, digestive*—*v. Secundæ*, *Ways, second*.

VIAL, *Phiala*.

VIANDES FAISANDÉES, see *Hyposaprus*.

VIBICES, *Mol'opes, Striæ*, (F.) *Vergetures*, 'Stripes.' Large purple spots, like the marks produced by the strokes of a whip, which appear under the skin in certain malignant fevers. They are unfavourable indications, as they denote great prostration.

VIBRAMEN, *Oscillation*.

VIBRANS, *Vibratory*.

VIBRATILE, *Vibratory*.

VIBRATILITY, *Vibrat'il'itas*, (F.) *Vibratilité*, from *vibrare, vibratum*, 'to quiver.' Capability of being thrown into vibration.

VIBRATIO, *Oscillation*—*v. Cardiaca*, *Palpitation*—*v. Cordis*, *Palpitation*.

VIBRATION, *PECTORAL*, see *Frenitus*—*v. Tactile*, see *Frenitus*—*v. Tussive*, see *Tussive*.

VIBRATIONS OF SOUND, see *Sound*.

VIBRATOR, *Deferens vas*.

VIBRATORY, *Vibrans, Vi'brative*, (F.) *Vibratile, Vibratoire*. An epithet applied to the pain that accompanies some neuralgiæ, which seems to the patient to vibrate among the nerves.

VIBRATORY MOTION, see *Cilia*.

VIBRATUS, *Oscillation*.

VIBRIO: from *vibrare*, 'to quiver.' A genus of infusory animalcules, of which different species are met with,—for example, *Vibrio prolifer* in almost all putrescent fluids containing protein, and in the pus of chancre; in which last the *V. lineola* has likewise been found.

VIBRIS'Æ. Same etymon. The hairs which grow at the entrance of the nostrils and of other outlets. In the feline tribe, the whiskers.

VIBURNUM CASSINOIDES, *Ilex Paraguensis*.

VIBURNUM DENTA'TUM, *Arrow Wood, Mealy Tree, Tily* of the Indians. *Ord. Caprifoliaceæ*. A high indigenous shrub, of the same family as *V. lentago*; the bark of which has been used by the Indians as a diuretic.

VIBURNUM EDU'LE flourishes in the Hudson's Bay territory. The fruit has an orange colour, and is less acid, more fleshy, and more agreeable to the taste than the cranberry.

VIBURNUM LEVIGATUM, *Ilex Paraguensis*.

VIBURNUM LENTA'GO, *Sweet Viburnum, Nannyberry*. A handsome indigenous tree, of the honeysuckle family—*Caprifoliaceæ*—which flowers in May and June. It has been used as an antiperiodic.

VIBURNUM PRUNIFOLIUM, *Black Haw, Sloe-leaved Viburnum, Sloe*. Indigenous: flowering in May. The leaves have been used as tea.

VIBURNUM SLOW-LEAVED, *V. prunifolium*—*v. Sweet, V. lentago*.

VIC-LE-COMTE, *MINERAL WATERS OF*. *Vic-le-Compte* is a small village in the de-

partment of Puy de Dôme, three leagues from Issoire: where there is an acidulous chalybeate, which is aperient.

VICARIOUS, *Vicarius*, from *vicis*, 'change, place.' Taking the place of another:—as 'vicarious secretion,' *Secretio vicaria*; a secretion which takes place in one part instead of another.

VICAR'S BRAY, MINERAL WATERS OF. This spring is situate in Perthshire, Scotland, about two miles from Dollor, and near to Blarngone. The water is one of the strongest chalybeates known.

VICE, Cachosis.

VICE DE CONFORMATION. Faulty formation of a part of the body.

VICHY, MINERAL WATERS OF. Carbonated alkaline chalybeates, the sources of which are at Vichy, six leagues from Moulins, in the department of Allier. The springs are seven in number. Almost all are thermal; give off bubbles of carbonic acid gas, and have a slightly sulphureous odour. They deposit carbonates of lime and magnesia, and a little oxide of iron; and afford, on analysis, carbonates of lime, soda, iron, and magnesia, carbonic acid, sulphate of soda, and chloride of sodium. They are employed as tonics, particularly in chronic affections of the abdominal viscera, in old catarrhs, diseases of the joints, &c.

VICHY WATER, FACTIVIOUS, Aqua Vicien'sis, (F.) *Eau de Vichy*, is made of simple acidulous water, impregnated with twice its bulk of carbonic acid, 3xxss ; subcarbonate of soda, gr. xxxij; sulphate of soda, gr. xvj; chloride of sodium, gr. iv; subcarbonate of magnesia, gr. ss; chloride of iron, gr. $\frac{1}{2}$. Mix. Ph. P.)

VICIA ERVILIA, Ervum.

VIC'IA FABA, *Faba*, *F. major* seu *vulgaris*, *Orobus faba*, Common Garden Bean, (F.) *Fève* ou *Fève des marais*. Ord. Leguminosæ. Sex. Syst. Diadelphia Decandria. A native of Egypt, of which there are many varieties. Beans are not easy of digestion. They are proverbially flatulent, especially when old. When young, they are more digestible and wholesome. Bean meal was one of the four resolvent meals of the ancients.

VICINTRACTUS, Erysipelas.

VICINITRAHA, Erysipelas.

VICTORIALIS FEMINEA, *Gladiolus vulgaris*—v. *Longa*, *Allium victorale*—v. *Rotunda*, *Gladiolus vulgaris*.

VICTUS RATIO, Diet.

VIDANGES, Lochia.

VIDIAN CANAL, Pterygoid canal—v. Nerve, Pterygoid nerve.

VIE, Life—v. *Durée de la*, see Longevity—v.

Moyenne, see Longevity.

VIEILLESSE, Senectus.

VIELLOTE, Colchicum autumnale.

VIERGE, Virgin.

VIF ARGENT, Hydrargyrum.

VIGILANCE, Insomnia.

VIGILANTIA, Egregorsis.

VIGILATIO, Egregorsis.

VIGILIA, Egregorsis.

VIGILIE, Egregorsis—v. *Nimix*, Coma vigil.

VIGILIUM, Egregorsis.

VIGNE, *Vitis vinifera*—v. *Blanche*, *Bryonia*

alba—v. de *Judée*, *Solanum dulcamara*—v. du

Nord, *Humulus lupulus*—v. *Vierge*, *Bryonia alba*.

VIGNETTE, *Spiraea ulmaria*.

VIGNOBLE, *Mercurialis*.

VIGOR, Acmé.

VILLA FRANCA, (CLIMATE OF.) This

little town is situate immediately to the eastward of Montalbano, which separates the bay of Villa Franca from that of Nice. It is on the southern declivity of a steep and lofty range of mountains;

and is more effectually sheltered from the north and north-west winds than Nice, with the climate of which that of Villa Franca closely corresponds.

VILLI, see Villous membranes.

VILLOUS, *Villosus*, (F.) *Villeux*, froz. *villus*, 'a hair.'

VILLOUS MEMBRANES or COATS, Velamenta Bombycina, (F.) *Membranes villieuses*, *Membranes veloutées*, are such as are covered with soft papillæ or *Villi*, *Flocci*, *Floculi*, *Vesiculae*, *Cylindri*, *Tabi Membranae*, constituted of blood-vessels, nerves, and absorbents; or such as are covered with fine down, so as to cause them to feel soft and velvety. The term is, however, chiefly restricted to the former. Chaussier gives the name *simple villous membranes* to the serous membranes; and that of *complicated or follicular villous membranes* to the mucous membranes.

VIN, Wine—v. de *Bordeaux*, Claret—v. de *Poulet*, see *Poulet*—v. de *Quinginaux* composé. Vinum de kinâ kinâ compositum—v. *Rouge d'Oporto*, Vinum Portugalicum—v. *Sec*, Sack.

VINAIGRE, Acetum, see Aceticum acidum—v. *Scillitique*, Acetum scillæ—v. de *Bois*, Pyroligneous acid.

VINAIGRES MÉDICINAUX, Acetica.

VINCA MINOR, *V. pervinca*, from *vincio*, 'I bind,' because of its usefulness in making bands. *Clematis daphnoides major*, *Lesser periwinkle*, (F.) *Perenche*, *Violette des Sorciers*. It has bitter and astringent properties, and has been used in gargles and diseases of the chest.

VINCETOXICUM, *Asclepias vincetoxicum*.

VINCTURA, Fascia.

VINCULA, Fascia.

VINCULUM, Bandage, Lien—v. Caninum, Frænum penis—v. Linguae, Frænum linguae—v. Præputii, Frænum penis—v. Umbilicale, Funiculus umbilicalis.

VINE, *Vitis vinifera*—v. Grape, *Vitis vinifera*.

VINE POISON, Rhubarb radicans.

VINE, Wild, *Paireira*.

VINEGAR, AROMATIC SPIRIT OF, see Aceticum acidum.

VINEGAR OF CANTHARIDES, Acetum cantharidis.

VINEGAR, COMMON, Acetum—v. Distilled, see Aceticum acidum—v. Estragon, *Artemisia dracunculula*—v. *Marseilles*, Acetum aromaticum.

VINEGAR OF MEADOW SAFFRON, Acetum colchici—v. of Opium, *Guttæ nigræ*.

VINEGAR, RADICAL, Aceticum acidum.

VINEGAR OF SQUILLS, Acetum scillæ—v. of the Four Thieves, Acetum aromaticum.

VINEGAR, THIEVES', Acetum aromaticum—v. Whey, see Acetum.

VINEGAR OF WOOD, see Aceticum acidum.

VINETTE, *Rumex acetosa*.

VINETTIER, *Oxyacantha Galeni*.

VINEUX, Vinous.

VINOLENTA, see Temulentia.

VINOLENTIA, Temulentia.

VINOSUS, Vinous.

VINOUS, *Vinosus*, *Enoëdes*, *Enoëdes*, (F.)

Vineux, from *vinum*, 'wine.' Relating to wine. Having the character of, or resembling, wine. Made of wine.

VINUM, Wine, Sherry wine (Ph. U. S. 1842). See Wine. A medicinal wine, (F.) *Vin medicinal*, *Onalé*, *Onolature*.

VINUM ABSINTHITES, Wine of Wormwood. (*Fot. absinth. exsic.*, *absinth. Pontic.*, αἷν ζῳ, *vinum albi* Oir. Macerate for 24 hours, and strain. Ph. P.) Stimulant, tonic, febrifuge, vermifuge.

VINUM ADUSTUM, Brandy—v. Album, Sherry wine (Ph. U. S. 1851), see Wine—v. Album Hispanicum, see Wine—v. Album Montanum, see Wine.

VINUM AL'OEËS, *V. aloës Sacotorina* seu *aloëticum* seu *aro'mo-aloëticum*, *Tinctura hiëra*, *Tinct-*

u'ra sacra, Wine of Aloes. (Aloes in pulv. $\mathfrak{z}\text{ij}$. cardam. cont., Zingib. cont., aa $\mathfrak{z}\text{ij}$; vini alb. Oj. Macerate for 14 days, and filter. Ph. U. S.) Purgative and stomachic, according to the dose. Dose, f $\mathfrak{z}\text{ij}$ to f $\mathfrak{z}\text{ij}$.

VINUM AMARUM, V. Gentianæ comp.—v. Amarum scilliticum, &c., V. diureticum amarum—v. Amarum cum spiritu vini, V. gentianæ compositum—v. Antimoniale, V. antimonii tartarizati—v. Antimoniatum Huxhami, V. antimonii tartarizati—v. Antimonii potassio-tartratis, V. antimonii tartarizati.

VINUM ANTIMONII, V. antimo'nii tartariza'ti seu tartra'tis antimonii seu antimo'nii potas'sio-tartratis seu benedictum seu antimoniatum Huxhami seu antimoniale seu emet'icum, Liquor tartari emet'ici, Essent'ia antimonii seu stil'bi, Aqua benedicta Rulandii, Solution of tartarized antimony, Antimonial wine. (Ant. et Potass. tart. $\mathfrak{z}\text{ij}$, vini albi f $\mathfrak{z}\text{x}$. Dissolve the tartarized antimony in the wine. Ph. U. S.) Each fluidounce contains two grains of the tartrate.

The ordinary Antimonial wine is made with glass of antimony $\mathfrak{z}\text{ij}$, sherry Oiss.

Ward's Red Drop consists of a strong vinous solution of tartarized antimony.

VINUM ANTISCORBU'TICUM, Antiscorbutic wine. (Rad. armaracia recent. $\mathfrak{z}\text{ij}$, fol. cochlearia recent., nasturtii aquatici, trifol. aquatici, semin. sinapi, aa $\mathfrak{z}\text{ss}$, ammonia mural. $\mathfrak{z}\text{ij}$, vin. albi Oj. Macerate for 36 hours: filter, and add tinctura cochlear. $\mathfrak{z}\text{ss}$. Ph. P.) Stimulant and tonic. Dose, f $\mathfrak{z}\text{ss}$ to f $\mathfrak{z}\text{ij}$.

VINUM AROMATICUM, Aromatic wine. (Species aromatic. $\mathfrak{z}\text{iv}$, vin. rubr. Oj. Macerate for 6 days and filter. Ph. P.) Stimulant and tonic. Used, externally, in fomentations, embrocations, &c.

VINUM AROMO-ALOETICUM, V. aloes—v. Benedictum, V. antimonii tartarizati.

VINUM CAMPA'NUM, V. Campanaceum. Champagne wine. A well-known sparkling wine of France. See Wine.

VINUM CANARINUM, see Wine—v. Chalybeatum, V. ferri—v. Cinchonæ compositum, V. de kinâ kinâ comp.

VINUM COL'CHICI, V. Colchici Rad'cis, Wine of Colchicum, Wint's Eau d'Huison. (Colchic. rad. bene contus. lbj, vini alb. Oj. Macerate for 14 days, with occasional agitation, and then express strongly, and filter. Ph. U. S.) It may also be prepared by displacement. Diuretic, sedative, and purgative. Chiefly used in gout. Dose, gtt. 40 to f $\mathfrak{z}\text{j}$.

Reynolds's Specific for gout and rheumatism consisted of the fresh bulb of Colchicum $\mathfrak{z}\text{viij}$, Sherry wine, f $\mathfrak{z}\text{xvj}$. The colchicum was macerated for eight or ten days in the wine, at a gentle heat; coloured with syrup of poppies and flavoured with rum. Reynolds is said to have killed himself by taking an over-dose of it.

VINUM COLCHICI RADICIS, V. colchici.

VINUM COLCHICI SEM'INIS, Wine of Colchicum seed. (Colchic. sem. cont. $\mathfrak{z}\text{iv}$; vini alb. Oj. Macerate for 14 days, with occasional agitation, then express and filter. Ph. U. S.) Dose, f $\mathfrak{z}\text{j}$ to f $\mathfrak{z}\text{ij}$.

VINUM CREMATUM, Brandy.

VINUM DIURETICUM AMARUM, V. amarum scilliticum compositum in Charitatis Nosocomio usurpatum, Bitter diuretic wine. (Cort. cinchon., Winteran., limon. aa $\mathfrak{z}\text{ij}$, rhiz. toxicodendr., scillitæ, cucumini. angelicæ, aa $\mathfrak{z}\text{ss}$, fol. absinth., melissa, aa $\mathfrak{z}\text{ij}$, buccar. junip., macis, aa $\mathfrak{z}\text{ss}$, vin. alb. Ovij. Macerate and filter. Ph. P.) Stimulant, tonic, and diuretic. Dose, f $\mathfrak{z}\text{ss}$ to f $\mathfrak{z}\text{ss}$.

VINUM EMETICUM, V. antimonii tartarizati.

VINUM ER'GOTE, Wine of Ergot. (Ergotæ, cont., $\mathfrak{z}\text{ij}$; vini alb. Oj. Macerate for 14 days,

with occasional agitation; then express and filter. Ph. U. S.) Dose, $\mathfrak{M}\text{xx}$ to f $\mathfrak{z}\text{j}$ to produce the peculiar effects of ergot.

VINUM FERRATUM, V. Ferri.

VINUM FERRI, V. Chalybeatum seu ferratum seu Martiatum seu Martis, Tinctura Martis cinchosa seu Martis cum vino, Steel wine, Wine of Iron. (Iron wire, $\mathfrak{z}\text{ij}$; Sherry wine, Oij (Imp. meas.). Digest for thirty days and filter. Ph. L.) Tonic. Dose, f $\mathfrak{z}\text{ij}$ to f $\mathfrak{z}\text{iv}$.

It is sometimes made by adding two pints of wine to two ounces of iron filings; macerating and filtering.

VINUM GENTIANÆ, V. G. Compositum seu Amarum seu Amarum cum spiritu Vini, Infusum amarum vino'sum, Compound wine of gentian, Wine bitters. (Rad. gentian. luteæ $\mathfrak{z}\text{ss}$, cort. cinchon. offic. $\mathfrak{z}\text{ij}$, cort. aurant. $\mathfrak{z}\text{ij}$, canella alb. $\mathfrak{z}\text{ij}$, alcohol. dilut. f $\mathfrak{z}\text{iv}$, vini alb. Hispanic. Oijss. Ph. E.) Dose, f $\mathfrak{z}\text{j}$ to f $\mathfrak{z}\text{vj}$.

VINUM HIPPOCRATICUM, Claret—v. Hispanicum, Vinum, see Wine—v. Hordeaceum, Cerevisia—v. Hydrotodes, see Hydrotide.

VINUM IPECACUAN'HE, Ipecacuan'ha wine. (Ipecacuan. cont. $\mathfrak{z}\text{ij}$, vini alb. Oij.) Macerate for 14 days, express, and filter. Emetic and diaphoretic. Dose, as an emetic, f $\mathfrak{z}\text{iv}$ to f $\mathfrak{z}\text{x}$.

VINUM DE KINâ KINâ COMPOSITUM, V. cincho'næ compositum, Compound bark wine, (F.) Vin de quinquina composé. (Cort. cinchon. pulv. lbss, quassia, cort. Winteran., cort. aurant. sicc. aa $\mathfrak{z}\text{ss}$, alcohol. (12° to 22° Baumé) Oj. Macerate for 24 hours; and add vini albi Ovi. Macerate for four days, and filter. Ph. P.) Tonic, febrifuge, and stomachic. Dose, f $\mathfrak{z}\text{ss}$ to f $\mathfrak{z}\text{ij}$.

VINUM MARTIATUM, V. ferri—v. Martis, V. ferri—v. Medicatum, see Claret.

VINUM MEDICINALE, Medicated Wine. Wine, holding in solution one or more medicines. Medicinal wines are obtained by macerating or digesting medicinal substances in wine. They differ, of course, according to the nature of the medicine employed, the strength of the wine, &c. In a late Pharmacopœia of the London College, all the wines were changed into weak tinctures; but the old names were left unaltered. Sherry is the wine usually employed. The following are medicated wines.

VINUM MERUM, Acratum—v. Nicotianæ tabaci, V. tabaci—v. Opium fermentatione paratum, Laudanum abbatris Rousseau.

VINUM OPII, V. de opio compositum, Laudanum liquidum Sydenhami seu Hoffmanni, Tinctura Thebæica, Wine of Opium, Sydenham's Laudanum, Ford's laudanum. (Opium in pulv. $\mathfrak{z}\text{ij}$; cinnam. cont., caryophyll. cont. aa $\mathfrak{z}\text{ij}$, vini albi Oj. Macerate for 14 days; express and filter.—Ph. U. S.) Narcotic. Dose, gtt. xv to f $\mathfrak{z}\text{j}$.

VINUM DE OPIO COMPOSITUM, V. opii—v. Picatum, Pissites.

VINUM PORTUGALICUM, V. rubrum, (Ph. U. S. 1851.) (F.) Vin rouge d'Oporto. Wine of Portugal or of Oporto. Port wine. A strong astringent wine, not much drunk in the United States; but greatly used in England.

VINUM QUINIE, Wine of Quinine. (Wine lbj, sulphate of quinia gr. x. Mix.) Dose, f $\mathfrak{z}\text{ss}$ or more; according as it is used as a stomachic, or to arrest ague. A Vinum Cincho'nice may be made in the same way.

VINUM RHEI, V. R. palma'ti, V. Rhabarbari, Tinctura Rhabarbari Vinosæ, Elizir Rhei dulcæ, E. R. Compositum, Rhabarb Wine. (Rhei cont. $\mathfrak{z}\text{ij}$, canella cont. $\mathfrak{z}\text{ij}$, alcohol. dilut. f $\mathfrak{z}\text{ij}$, vini albi, Oj. Macerate for 14 days; express and filter.—Ph. U. S.) Laxative and stimulant. Dose, f $\mathfrak{z}\text{iv}$ to f $\mathfrak{z}\text{ss}$.

VINUM RHEI PALMATI, V. Rhei—v. Rhenanum,

see Wine — *v. Roabbarbari*, *V. rhei palmati* — *v. Rubrum*, *V. Portugallicum* — *v. Rubrum Portugallicum*, see Wine.

VINUM TABACI, *Vinum Nicotianæ Tabaci*, *Tabacco Wine*. (*Tabac*, incis. $\bar{3}$), *vini albi*, Oj. Macerate for 14 days; express and strain. *Ph. U. S.*) Narcotic, diuretic, and antispasmodic. In dropsical cases, colica pictionum, and ileus. Not used. Dose, grt. xx to xxxvj.

VINUM VERATRI, *V. V. albi*, *Wine of White Hellebore*. (*Veratri cont.* $\bar{3}$ iv; *Vini albi*, Oj. Macerate for 14 days, express and filter. — *Ph. U. S.*) In cutaneous affections, and in gout. Dose, ten minims.

VINUM VERATRI ALBI, *V. Veratri* — *v. Xericum*, see *Vinum*.

VIOL, *Stuprum*.

VIOLA, *Viola odorata seu maritima*, *Ion*, *Matronalis*, *Viola'ria*, *Sweet Violet*, (F.) *Violette à bouquets*, *V. odorante*. *Ord. Violaceæ*. The flowers of the sweet violet, have a pleasant, peculiar odour, and scarcely any taste. They are slightly laxative.

VIOLA CANINA, *V. sylvestris*, *Dog Violet*. The root of this plant is emetic and cathartic. Dose, $\bar{3}$ j.

VIOLA IPECACUANHA, see *Ipecacuanha* — *v. Lutea*, *Cheiranthus cheiri* — *v. Martia*, *Viola* — *v. Odorata*, *Viola* — *v. Palustris*, *Pinguicula vulgaris*.

VIOLA PEDA'TA, *Viola*, (*Ph. U. S.*) This indigenous violet, which grows from New England to Carolina, is regarded by some as a useful demulcent in pectoral affections.

VIOLA SYLVESTRIS, *V. Canina*.

VIOLA TRI'COLOR, *Jacq's tricolor*, *Herba seu Flos Trinitatis*, *Heart's ease*, *Pansy*, *Peney*, *Pansies*, (Prov.) *Beedy's eyes*, *Bidly's eyes*, (F.) *Pensée*, *P. sawage*. The decoction of this beautiful little plant has been recommended in diseases of the skin, &c. Its roots are, also, emetic and cathartic.

VIOLARIA, *Viola*.

VIOLET, DOG, *Viola canina* — *v. Dog's Tooth*, *Erythronium Americanum*.

VIOLET BLOOM, *Solanum dulcamara*.

VIOLET, SWEET, *Viola*.

VIOLETTE À BOUQUETS, *Viola* — *v. Odorante*, *Viola* — *v. des Sorciers*, *Vinca minor*.

VIOLIER JAUNE, *Cheiranthus cheiri*.

VIORNA, *Clematis vitalba*.

VIPER, EGYPTIAN, *Aspis* — *v. Ten-inch long*, *Echidna ocellata*.

VIPER GRASS, *Scorzonera*.

VIPERA, *Coluber berus*.

VIPERARIA, *Aristolochia serpentaria*, *Scorzonera*.

VIPERINA VIRGINIANA, *Aristolochia serpentaria*.

VIPÉRINE DE VIRGINIE, *Aristolochia serpentaria*.

VIPER'S DANCE, *Chorea*.

VIR, *Aner*, *Penis* — *v. Effeminatus*, *Gynæcaner*.

VIRAGO, see *Pogoniasis*.

VIREUX, *Virulent*.

VIRGA, *Penis* — *v. Cereæ*, *Bougie* — *v. Cerebri*, *Pineal gland*.

VIRGILLA, *Cladrastis tinctoria*.

VIRGIN, *Virgo*, genitive *virginis*, *Parthenus*, *Expers nuptiarum*, (F.) *Vierge*, *Pucelle*. A female who has never had sexual intercourse. Applied, also, occasionally to the male so circumstanced.

VIRGIN-MARY THISTLE, *Centaurea benedicta*.

VIR'GINAL, *Virginalis*, from *virgo*, *virginis*, 'a virgin.' The external genital organs of the virgin.

VIRGINAL MEMBRANE, *Hymen*.

VIRGINIA, MINERAL WATERS OF. There are several celebrated mineral springs in Virginia,

on the western side of the Blue Ridge. Two Thermal springs exist in Bath county. The Warm Spring, as it is called, issues with a bold stream, sufficient to turn a mill, and to keep the water of its basin at the temperature of 97°. The Hot Spring is a few miles from the warm. Its temperature is 107°. They are both slightly sulphureous. The Sweet Springs are at the eastern foot of the Allegheny, in Botetourt county; 42 miles from the Warm Springs. The water has a temperature of 70°, and is highly impregnated with carbonic acid. The White Sulphur, as well as the Red Sulphur, and Salt Sulphur Springs, are also much celebrated. The Red Spring is affirmed to exert a sedative influence on the pulse, diminishing the number of its beats. Prof. W. B. Rogers analyzed the water of the White Sulphur, and found it to be composed as follows:—Gaseous contents in a wine gallon. Sulphuretted hydrogen, 2.5 cubic inches. carbonic acid, 2; oxygen, 1.448; nitrogen, 3.552.—Total, 9.5. Solid contents in a pint: sulphate of magnesia, 5.588 grains; sulphate of lime, 7.744; carbonate of lime, 1.150; chloride of calcium, 0.204; chloride of sodium, 0.180; oxide of iron, a trace; loss, 0.410. Total, 15.276 grains.

The Rockbridge Alum Springs are situated 17 miles west of Lexington, and are in much favour. Their solid constituents are said to be,—sulphate of potassa, 1.765 gr.; sulphate of lime, 3.263; sulphate of magnesia, 1.763; protoxide of iron, 4.863; alumina, 17.905; erenate of ammonia, 0.700; chloride of sodium, 1.008; silicic acid, 2.840; free sulphuric acid, 15.224; free carbonic acid, 7.536 — water 58315.133. They are used in dyspepsia, scrofula, chronic diarrhoea, and cutaneous affections. The Bath Alum Springs are about 17 miles west of the Rockbridge Springs, in Bath county. They consist of free sulphuric acid, 7.878 grs; carbonic acid, 3.846; sulphate of potassa, 0.258; magnesia, 1.282; lime, 2.539; protoxide of iron, 21.776; alumina, 12.293; erenate of ammonia, 1.776; silicate of soda, 3.150; water, 58317.202. They are more chalybeate than the last.

About 15 miles from Staunton are Stribling's Springs—a sulphuretted water; and near Warrenton, in Fauquier county, there is a sulphureous water, called Lee's Springs, which is much frequented. Berkley or Bath, and Capon Springs are likewise celebrated.

VIRGINIA, *Hymen*.

VIRGINIS VIITIATIO, *Defloration*.

VIRGIN'ITY, *Virgin'itas*, *Parthen'e'ia*, *Integritas*, *Maid'enhoo*, *Maid'enhead*, (F.) *Virginité*, *Pucelage*; from *virgo*, *virginis*, 'a virgin.' The state of one who has never had carnal commerce.

VIRGIN'S BOWER, *COMMON*, *Clematis vitalba*, *Clematis Virginica* — *v. Bower*, upright, *Clematis recta* — *v. Sweet-scented*, *Clematis flammula*.

VIRGIN'S MILK, *Lac Virg'inis* seu *Virginalis*, (F.) *Lait virginal*. A tincture of benjamin or of any other balsam, precipitated by the addition of water, which gives it a milky appearance. Also, sub-acetate of lead precipitated by water. It has been used as a cosmetic to remove freckles, &c.

VIRGINITAS DEFLORATA, *Defloration*.

VIRGO, *Virgin* — *v. Aurea*, *Solidago virgaurea*.

VIRGULA, *Penis*.

VIRIDEÆRIS, *Cupri subacetat* — *v. Montana*, *Cuprum*.

VIRILE, *Penis*.

VIRILES'CENTE: from *virilis*, (*vir*, 'a man,') 'manly.' The state of the aged female in which she assumes certain of the characteristics of the male. — *Mehlis*.

VIRILIA, *Pubidibia*.

VIRILITY, *Adult age*.

VIROSUS, *Virulent*.

VIRTUE, Faculty.

VIRTUS FORMATRIX, Plastic force — *v.* 1 *astica*, Plastic force.

VIRULENT, *Virulentus*, *Virosus*, (F.) *Vireux*, from *virus*. That which is of the nature of a virus or which is caused by a virus. Highly noxious.

VIRUS, from *vis*, 'force,' plural *vires* [?]. A Latin word, which signifies poison; but which, in medicine, has a somewhat different acceptation. By it is understood a principle, unknown in its nature and inappreciable by the senses, which is the agent for the transmission of infectious diseases. Thus we speak of the *variolic*, *vaccinæ*, and *syphilitic viruses*. *Virus* differs from *venom* in the latter being a secretion natural to certain animals, whilst the former is always the result of a morbid process, — a *morbid poison*.

VIRUS, Ichor, Poison, Sanies — *v.* *Vaccine*, see *Vaccina* — *v.* *Vitale*, *Sperm*.

VIS, Force — *v.* *Adhæsis*, *Cohesion* (force of) — *v.* *Attractionis*, *Cohesion* (force of) — *v.* *Aucta*, *Sthenia* — *v.* *Cohæsis*, *Cohesion* (force of) — *v.* *Conservatrix*, *V. medicatrix naturæ* — *v.* *Elastica*, *Elasticity* — *v.* *Essentialis*, *Plastic force*.

VIS EXPULTRIX. The organic force by which, it was supposed, the useless was removed from the body.

VIS FORMATIVA, Plastic force — *v.* *Generatrix*, *Potentia generandi* — *v.* *Imminuta*, *Asthenia* — *v.* *Insita*, *Irritability* — *v.* *Irritabilitatis*, *Irritability*.

VIS MEDICATRIX NATURÆ, *V. conservatrix*, *Autoerati'a*, *Autonom'ia*, *Autoerato'ria*, *Authy-gian'sis*, *Autotherapei'a*, *Physiat'ricæ*, *Physio-autocati'a*, *Physiautoerati'a*, (F.) *Force médicale*. A term, employed by some, to express that instinctive healing power, in an animal or vegetable, by virtue of which it can repair injuries inflicted upon it, or remove disease.

VIS METABOLICA, Metabolic force.

VIS MOR'TUA, Under this name Haller included the mechanical properties of membrane, — flexibility, extensibility, elasticity, &c. — which are totally independent of the vital properties. They remain some time after the complete extinction of life in all its functions; seem to be connected with the mechanical arrangement of particles and the chymical composition of the substance in which they reside, and not to be affected until the progress of decomposition has become sensible.

VIS MUSCULARIS, Myodynamia — *v.* *Nervorum*, *Nervous power* — *v.* *Nervosa*, *Nervous power* — *v.* *Plastica*, *Plastic force* — *v.* *Reproductiva*, *Plastic force* — *v.* *Vitæ*, *Irritability*.

VIS VITA'LIS, *Vis seu Fac'ultas Zo'tica*, (F.) *Force vitale*. The vital force — *Biod* of Baron Von Reichenbach — which distinguishes living matter from the dead. Also, *Irritability*.

VIS ZOTICA, *Vis vitalis*.

VISAGE, Countenance.

VISCAGO BEHEN, *Cucubalus behen*.

VISCERA, Entrails.

VISCERAL, Splanchnic.

VISCERALIS, Stomachal.

VISCERATION, Exenterismus.

VISCÈRES, Entrails.

VISCONIA, *Physconia*.

VISCUM ALBUM, *Ixos*, *Ix'ia*, *Dabach*, *Mis'tetoe*, (F.) *Gui*, *Gui de Chêne*, *Gillon*. A parasitic plant, which grows on a number of trees. *Order*, *Loranthacæ*. *Sec. Syst.* *Dioecia Tetrandria*. That of the oak — *Viscum quercinum* — has been most celebrated, but it does not differ from the others. It was, at one time, much used in epilepsy and other neuroses; the wood or leaves

being given in powder, in the dose of \mathfrak{zss} or \mathfrak{zj} . The berries, also called *Pommes hémorrhoidales*, from their supposed virtues, are drastic; and, when applied externally, are maturative.

VISCUM CAPENSE. A South African parasitic shrub, which grows on the stems of several species of *Rhus* and *Euclea*.

VISCUM FLAVESCENS, *V. verticillatum*, *Yellow* or *Yellowish Mistletoe*, indigenous, has been given in similar cases as the *Viscum album*.

VISCUM VERTICILLATUM, *V. flavescens*.

VISIO, Pupil, Vision.

VISION, *Sight*, *Vis'io*, *Vius*, from *videre*, *visum*, 'to see,' *Opsis*, *Omma*, (F.) *Vue*. The action of seeing. The sight is one of the five senses, and the eye is its organ. By it we distinguish colours, and appreciate most of the physical qualities of external objects. The eye is composed of different transparent media, whose curvatures and refractive powers are so combined as to render insensible the aberrations of sphericity and refrangibility, and to concentrate the numerous rays proceeding from more or less distant objects. These rays traverse, successively, the transparent cornea, the aqueous humour, the crystalline, the vitreous humour, and strike upon the retina, on which they impress the figure of the object whence they have proceeded; and, in order that no obscurity may arise from the reflections that might otherwise occur, the anterior of the eye is lined by a membrane — the choroid — which is itself covered internally by a dark pigment. This absorbs the light, like the dark pigment in the interior of telescopes, and thus obviates confusion. As the rays proceeding from an object must cross each other in their course to the retina, it follows that those proceeding from the upper part of an object must fall upon the lower part of the retina, and conversely; so that the object must be impressed upon the retina in an inverted position; yet experience teaches us to appreciate the position accurately.

VISION, BINOCULAR, see *Binocular* — *v.* *Coloured*, *Chromopsia* — *v.* *Double*, *Diplopia* — *v.* *Iridescent*, *Chromopsia* — *v.* *Manifold*, *Polyopsia* — *v.* *Multiple*, *Polyopsia* — *v.* *Phantasmatum*, *Metamorphopsia* — *v.* *Single*, see *Binocular*.

VISNOMY, Physiognomy.

VISORIUM ORGANUM, Eye.

VISORIUS, Visual.

VIS'UAL, *Visua'lis*, *Viso'rius*, (F.) *Visuel*. Same etymon as *Vision*. That which concerns the sight or belongs to vision.

VISUAL ANGLE is the angle formed by the crossing of two rays, proceeding from opposite points of any body, in their passage through the pupil. It is by the size of this angle that we judge, to a certain extent, of the dimensions of objects.

VISUEL, Visual.

VISUS, Vision — *v.* *Aceris*, *Oxydercia* — *v.* *Acrior*, *Nyctalopia* — *v.* *Coloratus*, *Chromopsia* — *v.* *Debilitas*, *Asthenopia* — *v.* *Decolor*, *Achromatopsia* — *v.* *Defiguratus*, *Metamorphopsia* — *v.* *Dimidiatus*, *Hemipopia* — *v.* *Diurnus*, *Hemeralopia* — *v.* *Duplicatus*, *Diplopia* — *v.* *Hebetudo*, *Amblyopia* — *v.* *Juvenum*, *Myopia* — *v.* *Lucidus*, *Photopsia* — *v.* *Muscarum*, see *Metamorphopsia* — *v.* *Nebulosus*, *Nebula* — *v.* *Nocturnus*, *Nyctalopia* — *v.* *Senilis*, *Presbytia*.

VITA, Life — *v.* *Propria*, *Irritability*.

VITAL, *Vita'lis*, *Biot'ic*, *Biot'icus*, (F.) *Vital*, from *vita*, 'life.' Relating or appertaining to life.

VITAL CAPAC'ITY, *Pulmon'ic* or *Lung capacity*. A term employed by Dr. Hutchinson to designate the number of cubic inches of air, which a person can forcibly expire after a full inspiration.

VITAL DYNAMICS, see Dynamic—v. Force, Vis vitalis.

VITAL PRINCIPLE, *Flam'mula seu Aura vitalis*, *Microcosme'tor*, (F.) *Plamme vitale*, *Vital Flame*. The principle which, in association with matter, as in organized bodies, controls its manifestations and properties, and distinguishes organized matter from the inorganic. We know nothing of this principle, except in its connexion with organization.

VITAL PROPERTIES, see Properties, vital—v. Statistics, Statistics, medical.

VITALBA, *Clematis vitalba*.

VITALISM, *Vitalism'us*. The doctrine of the *vitalist*—*vitalis'ta*—who ascribes every function of the organism to the direct agency of the vital principle.

VITALIST, see Vitalism.

VITALITY, *Vital'itas*, (F.) *Vitalité*, from *vita*, 'life.' The vital principle: also, used in the sense of vital action or movement.

VITALIZE. To endow with life.

VITALS, *Vital'ia*. Parts that are necessary for life—the heart, lungs, and nervous centres more especially.

VITARO'BORANT, PALMER'S. A farinaceous preparation, which appears to consist of a mixture of the red or Arabian lentil and wheat flour, the latter in considerable amount, sweetened with sugar. It bears considerable resemblance to Revalenta.

VITELLARY, Vitelline—v. Membrane, Zona pellucida.

VITELLINE, Zona pellucida.

VITELLINE, *Vitell'inus*, *Vit'ellary*, *Lecith'les*, from *vitellus*, 'the yolk of an egg.' Appertaining to or resembling the vitellus or yolk. Also, the yolk membrane. See *Vesicula umbilicalis*.

VITELLINE DISC, Proligerous Disc—v. Fluid, see *Vesicula umbilicalis*.

VITELLINE PED'ICLE or APOPH'YSIS is the pedicle which attaches the umbilical vesicle to the embryo. See *Vesicula umbilicalis*.

VITELLINE POUCH, *Vesicula umbilicalis*—v. Vessels, Omphalo-mesenteric.

VITELLO-MESENTERIC, Omphalo-mesenteric.

VITELLUM OVI, see Ovum.

VITELLUS OVI, see Ovum.

VITEX, from *vicio*, 'I bind' [?]; *V. Agnus Castus* seu *verticilla'ta*, (F.) *Gatillier*, *Petit Poivre*, *Poivre de moine* ou *sauvage*. Ord. *Verbenaceæ*. *Sex. Syst.* *Didynamia Angiospermia*. The *Chaste Tree*. The seeds, when fresh, have a fragrant smell, and an acrid, aromatic taste. Formerly, they were celebrated as antaphrodisiacs.

VITI SALTUS, *Chorea*.

VITIA PRIMÆ CONFORMATIONIS, see Monster.

VITIATIO, *Stuprum*.

VITILIGINES, *Ephelides*.

VITILIGO, *Lepra*—v. *Alphus*, *Lepra alphoides*—v. *Melas*, *Lepra nigricans*—v. *Nigra*, *Lepra nigricans*.

VITILIGO'DEA, from *vitiligo*, and *eidōs*, 'resemblance.' A disease of the skin, so called from its resembling the vitiligo of Willan. It appears under two forms; either as tubercles—*V. tuberosa*—varying from the size of a pin's head to that of a large pea, isolated or confluent; or as yellow patches of irregular outline, slightly elevated, and with but little hardness—*V. plana*.

VITIS ALBA SYLVESTRIS, *Bryonia alba*—v. *Apyrena*, see *V. Corinthiaca*.

VITIS CORINTHIACA. Ord. *Vitaceæ*. *Sex. Syst.* *Pentandria Monogynia*. This plant affords the *Uvæ Corinthiacæ passæ mino'res*, *Passæ Corinthiacæ*, *Pas'sulæ mino'res*, *Corinthiacæ*, *Currants*,

(F.) *Raisins de Corinthe*; which are the dried fruit of the *Vitis Corinthiaca* or *Vitis apyrena*; and are similar, in their properties, to the raisin.

VITIS IDÆA MYRTILLUS, *Vaccinium myrtillus*—v. *Idæa Palustris*, *Vaccinium oxycoccos*—v. *Idæa punctata*, *Vaccinium vitis idæa*—v. *Sativa*, *V. vinifera*.

VITIS VINIF'ERA, *V. sat'iva*, *Am'pelos*, *A. Enoph'oros*, *Grape vine*, *Vine*, (F.) *Vigne*. Vine leaves and the tendrils have an astringent taste, and were formerly used in diarrhoea, hemorrhage, &c. The juice or sap, called *Lacryma*, has been recommended in calculous disorders and in cases of chronic ophthalmia. The juice of the unripe fruit is called *Verjuice*. The *Grape*, *Uva*, *Botrys*, *Staph'yle*, *Rhas*, (F.) *Raisin*, when ripe, is a wholesome delicious fruit; and, when dried, forms the *Uvæ passæ majo'res*, *Pass'ulæ majo'res*, *Uvæ passæ solè sicca'tæ*, *Uvæ passa* (Ph. U. S.), *Zibebæ*, *Raisins*. These are inodorous; and to the taste, sub-acidulous, sweet, and mucilaginous. They are demulcent and nutritive.

VITIUM, Disease—v. *Caducum*, *Epilepsy*.

VITIUM CONFORMATIONIS. A disease or fault of conformation.

VITIUM DIVINUM, *Epilepsy*—v. *Herculeum*, *Epilepsy*—v. *Scrophulosum*, *Scrofula*.

VITRARIA, *Parietaria*.

VITRÉ, Hyaline.

VITREO-CAPSULITIS, *Hyalitis*.

VITREOUS, Hyaline—v. Humour, Corpus vitreum—v. Humour of the ear, *Vitrine auditive*.

VITREUS, Hyaline.

VITREUX, Hyaline.

VITRINA OCULARIS, *Corpus vitreum*.

VITRINE AUDITIVE (F.), from *vitrum*, 'glass'; *A'quila vit'rea audit'iva*, *End'olymph*, *Liquor* or *Fluid* of *Scarpa*, *Vitreous Humour of the Ear*. The fluid contained in the membranous labyrinth of the ear, so called from its resemblance to the vitreous humour of the eye.

VITRIOL, Ferri sulphas—v. Blue, Cupri sulphas—v. Green, Ferri sulphas—v. Roman, Cupri sulphas—v. White, Zinci sulphas.

VITRIOLUM ALBUM, Zinci sulphas—v. Ammoniacale, Ammonia sulphas—v. *Ceruleum*, Cupri sulphas—v. Cupri, Cupri sulphas—v. Cyprum, Cupri sulphas—v. Ferri, Ferri sulphas—v. Goslariense, Zinci sulphas—v. Martis, Ferri sulphas—v. Romanum, Cupri sulphas—v. Veneris, Cupri sulphas—v. Viride, Ferri sulphas—v. Zinci, Zinci sulphas.

VITRUM, *Hy'alos*, *Glass*, *Brancia*, (F.) *Verre*. Glass, coarsely powdered, has been sometimes used to remove specks of the cornea.

Powdered glass has been regarded, but erroneously, as a poison. It acts only mechanically.

The saline scum, which swims on glass when first made, has been used in tooth-powders. It is called *Sandiver*, *Sel de verre*, *Glass Gall*, *Fel vitri*, &c.

VITRUM ANTIMONII, see Antimonii vitrum—v. Hypocleptium, Separatorium—v. Oculare, see Spectacles—v. Sublimatorium, Aludel.

VITTA, *Bandelette*, *Caul*.

VITTARIA CETERACH, *Asplenium ceterach*.

VITTIE VAYR, *Andropogon muricatus*.

VIVACE, from *vivax*, *vivaci*, (*vivere*, 'to live'), 'living long, tenacious of life.' An epithet given by Levret to uterine polypi, which he considers to be, commonly, vegetations of some ulcer in the interior of the uterus, and incurable.

VIVASECTIO, *Vivisection*.

VIVERRA CIVETTA, see Civetta.

VIVIPARE, Viviparous.

VIVIPARITÉ, see Viviparous.

VIVIP'AROUS, (F.) *Vivipare*, from *vivus*, 'alive,' and *pario*, 'I bring forth.' An epithet applied to animals whose young are brought forth detached, without requiring to be hatched, as in the *Oviparous*. The condition may be termed *Vivip'arousness*, (F.) *Viviparité*. See Generation.

VIVIPAROUSNESS, see *Viviparous*.

VIVISECT'ION, *Vivisectio*, *Vivæsectio*, *Biotom'ia*, from *vivus*, 'alive,' and *secare*, *sectum*, 'to cut.' The act of opening or dissecting living animals. One who performs vivisections is a vivisector.

VOCAL, Oral — *v.* Cords, see Cords, vocal — *v.* Ligaments, see Cords, Vocal — *v.* Tube, see Tube, vocal.

VOCALIS, Vowel.

VOCE DI PETTO, see Voice — *v.* di Testa, see Voice.

VOICE, *Vox*, *Audē*, *Phonē*, (F.) *Voix*, from *vocare*, 'to name, to call.' An appreciable sound produced by the air, when driven from the lungs, throwing the inferior ligaments of the glottis into vibration. The function by which this is effected has been called *Phona'tion*. It is a function of animal life; and, in animals, is limited to the production of the *simple or instinctive voice*, as well as in the idiot; but, in intellectual man, becomes much more complicated and articulated, — an evidence of his intellectual superiority.

VOICE, ARTICULATED, *Speech*, *Loque'la*, *Glossa*, *Glotta*, *La'tia*, *Phrasis*, (F.) *Parole*, *Voix articulée*, is the voice as produced by the glottis, modified by the motions of the tongue, lip, and other parts of the vocal tube. Most physiologists agree in regarding the glottis, or the space between the thyro-arytenoid ligaments plus the ligaments themselves, as the part where vocal sounds are formed; but the mechanism of this formation is a matter of dispute. The greater part of individuals regard the glottis and its ligaments, with the *vocal tube*, or the space which the sound has to traverse after it is produced in the glottis, as a reed instrument, like the clarinet, the inferior ligaments of the glottis constituting the reed. In speech, as ordinarily effected, the tongue, lips, teeth, &c., are concerned, but there are some authentic cases on record, where it has been accomplished after the tongue has been removed. In such cases, the articulated voice must have been produced in the glottis alone, or in it aided by an obscure action of the parts above it.

The singing voice is not limited to the larynx; the pharynx would seem to be concerned also. The notes that are produced in the latter have been termed *supra-laryngeal*, or *notes of the second register*. They constitute the *pharyngeal voice*, *falsetto* or *fauvette voice* or *voce di testa* (I.), *Voix de tête*, *Voix de fausset* (F.). The *laryngeal voice* or *voice of the first register*, *Chest voice*, is the *Voce di Petto* (I.), *Voix de Poitrine* (F.). The *laryngeal voice*, which is always more elevated, by an octave, in the female than in the male, furnishes the types called *Soprano*, *Alto* or *Contralto*, *Tenor* and *Bass*. The first two belong to the female voice; the last two to the male. The *Baritone* is between the bass and tenor.

VOICE, BLEATING, Egophony — *v.* Cavernous, Pectoriloquy — *v.* Chest, see Voice, articulated.

VOICE, CHOL'ERIC, *Vox Chol'er'ica*. The peculiar husky voice of those affected with cholera.

VOICE, CONVUL'SIVE, *Vox convulsi'va*, (F.) *Voix convulsive*. A neurosis of the voice, consisting in the production of discordant sounds — acute and grave — which cannot be brought to the natural tones by any effort of volition. It seems to be caused by disordered contraction of the muscles of the larynx.

VOICE, FALSETTO, see Voice — *v.* Goat's, Egophony — *v.* Laryngeal, Laryngophony, see Voice

— *v.* Pharyngeal, see Voice — *v.* Tracheal. Laryngophony.

VOIE, Way.

VOIES DIGESTIVES, Ways, digestive — *v.* *Lacrymales*, Lachrymal passages — *v.* *Premières*, Ways, digestive — *v.* *Secondes*, Ways, second — *v.* *Urinaires*, Urinary passages.

VOILE DU PALAIS, Velum pendulum palati — *v.* *Médullaire inférieure*, Velum medullare inferius — *v.* *Médullaire supérieure*, Valvula Vieussenii.

VOIX, Voice — *v.* *Aigre*, see Acid — *v.* *Articulée*, Voice, articulated, Pectoriloquy — *v.* *Bourdonnante*, see Resonance — *v.* *Bronchique*, Resonance — *v.* *Caverneuse*, Pectoriloquy — *v.* *Chérotante*, Egophony — *v.* *Égophonique*, Egophony — *v.* *de l'ausset*, see Voice, articulated — *v.* *de Poitrine*, see Voice, articulated — *v.* *de Polichinelle*, Egophony — *v.* *Sénile*, Egophony — *v.* *de Tête*, see Voice, articulated — *v.* *Tubaire*, see Resonance.

VOLA, Palm — *v.* Manus. Palm — *v.* Pedis, Sole.

VOLANT D'EAU, Nymphæa alba.

VOLATICA, Lichen.

VOLATILIZA'TION, *Volatilisa'tio*, *Volatilisatio*, from *volare*, 'to fly.' The operation of reducing volatilizable substances into gas or vapour.

VOLATISATIO, Volatilization.

VOLIT'ION, *Volit'io*, from *volō*, 'to will.' The act of willing. Hence:

VOLIT'IONAL, relating to volition. Thus the muscles may be moved by a *volitional* or by an *emotional* impulse.

VOLNUS, Wound.

VOLONTAIRE, Voluntary.

VOLONTÉ, Voluntas.

VOLSELLA, Acanthobolos, Forceps, Madistarium — *v.* Patini, *Valet à Patin*.

VOLTAISM, Galvanism.

VOLTUS, Countenance, Face.

VOL'UNTARY, *Hec'u'sius*, *Volunta'rius*, (F.) *Volontaire*, from *voluntas*, 'will.' Relating to the will; as 'voluntary muscles,' muscles which act in obedience to the will.

VOLUNTAS, *Volent'ia*, *Arbit'rarium*, *Cupid'itas*, *Boule'sis*, *Bule'sis*, *Thel'e'ma*, (F.) *Volonté*, from *volō*, 'I will.' Will or desire.

VOLUNTAS seu CUPID'ITAS DESEEND'I, *Lubi'do intest'i'ni*. The desire to go to stool.

VOLVA, Vulva.

VOLVULUS, Ileus — *v.* Intestinorum, Intussusceptio.

VOMENDI CONAMEN INANE, Vomiturition.

VOMER, 'a ploughshare,' *Os vo'meris*, *Os maxill'æ superio'ris undec'imum*, *Ara'trum*. One of the bones of the face, so called from its resemblance. It is an azygous bone, forming the posterior part of the septum narium. It is thin, flat, and quadrilateral; and is articulated, below, to the superior maxillary and palatine bones; and above, to the sphenoid, ethmoid, and cornets of Bertin. It is developed by a single point of ossification. Also, the penis.

VOMICA, *Pyocys'tis*, from *vomere*, 'to vomit,' 'to cast up.' (F.) *Vomique*, *Crachement de pus*. A term used by some authors for any encysted collection of purulent matter in the interior of a viscus. The acceptation is commonly, however, restricted to purulent collections in the lungs, encysted or not, which may be discharged by the mouth, by breaking into the bronchial tubes. Two kinds of vomicae are distinguished; the one caused by the softening or breaking down of pulmonary tubercles; the other, of much more rare occurrence, formed by a circumscribed abscess of the lungs. Instead of opening into the bronchia, the abscess is sometimes discharged into the cavity of the pleura, causing empyema.

VOMICINE, Brucia.

VOMIQUE, Vomica.

VOMIQUEER, *Strychnos nux vomica*.

VOMISSEMENT, Vomiting—*v. de Sang*, Hæmatemesis.

VOMIT, Emetic—*v. Black*, Fever, yellow, Melæna.

VOMIT, DRY, OF MARRIOTT. A once celebrated emetic, called 'dry,' from its having been given without solution. It consisted of equal portions of tartrate of antimony and potassa, and sulphate of copper.

VOMITIF, Emetic.

VOMITINE, Emetice.

VOMITING, Vomitus, Vomit'io, Vomit'ium, Vom'itum, Evomit'io, An'atropê, Em'esis, Palnus vom'itus, *Ægritudo ventriculi*, Spewing, Puking, Sickness of the Stomach, (Old Eng.) Parbreaking, (Sc.) Bokking, Bocking, Braking, Broacht, (Prov.) Hull'pung, (vulg.) Casca'ding, Casting, (F.) Vomissement, from vomere, vomitum, 'to throw up.' The act by which solids and liquids, contained in the stomach, are ejected by the mouth. Vomiting is the effect of some special irritation of the nerves of the stomach, calling, by reflex action, on the appropriate muscles to expel the contents of the viscus. It is not accomplished solely by the contraction of the stomach, as was for a long time believed. That organ, indeed, assists in the process; but the main cause is the compression exerted upon the parietes of the viscus by the diaphragm and abdominal muscles. Vomiting is a symptom common to numerous diseases. At times, it is sympathetic, as in affections of the kidney, uterus, brain, &c. At others, it is symptomatic, as in gastritis, peritonitis, &c. When very troublesome, it may often be palliated by effervescent draughts, aromatics, sinapisms, or blisters to the epigastric region, &c.

VOMITING AT PLEASURE, Regurgitation—*v. of Blood*, Hæmatemesis—*v. Stercoraceous*, Copremesis.

VOMITIO, Vomiting—*v. Sanguinis*, Hæmatemesis.

VOMITIUM, Vomiting.

VOMITO NEGRO, Fever, yellow—*v. Prieto*, Fever, yellow.

VOMITORIUM, Emetic.

VOMITORY, Emetic.

VOMITUM, Vomiting.

VOMITURIT'ION, Vomiturit'io, Emes'ia, Dysemes'ia, Subver'sio stom'achi. Same etymon as Vomiting. Ineffectual efforts to vomit; Vom'en'di con'd'men in'a'nê, Retching, Reaching, Herv'ing, (Old Eng.) Bolke, (Sc.) Bok, Bock, Braking. Some authors mean, by this term, a vomiting of but little matter, or that which is effected almost without effort. The first acceptance is the most general.

VOMITUS, Vomiting—*v. Cruentus*, Hæmatemesis—*v. Fæculentus*, Copremesis—*v. Marinus*, Nausea marina—*v. Navigantium*, Nausea marina—*v. Niger*, Fever, yellow—*v. Pituitosus*, Blennemesis—*v. Profusus*, Hyperemesis—*v. Purulentus*, Pyemesis—*v. Sanguinis*, Hæmatemesis—*v. Stercoris*, Copremesis.

VOORARA, Curare.

VOTIVE TABLET, Tabula votiva.

VOUACAPUA AMERICANA, Geoffræa inermis.

VOUSSURE PRÉCORDIALE, 'Præcordial arching or prominence.' The elevation or arching of the parietes of the thorax, witnessed in certain thoracic affections, as in hypertrophy of the heart.

VOÛTE, Vault—*v. Médullaire*, Corpus callosum—*v. à Trois Piliers*, Fornix.

VOWEL, Voca'tis, from Latin vox, vocis, 'the

voice,' and roco, 'to call.' (F.) Voyelle. Physiologically, a continuous breath or sound, produced in the glottis; but more or less modified by the form of the vocal tube, through which it has to pass. In the English language, *a, e, i, o, u, w*, and *y*, are vowels.

VOX, Voice—*v. Cholericæ*, Voice, choleric—*v. Clangosa*, Oxyphonia—*v. Convulsiva*, Voix Convulsive—*v. Nasalis*, Rhinophonia—*v. Rauca*, Raucedo, Raucous voice—*v. Raucisona*, Raucous voice.

VOYELLE, Vowel.

VUE, Vision—*v. Basse*, Myopia—*v. Courte*, Myopia—*v. Diurne*, Hemeralopia—*v. Faible*, Amblyopia—*v. Longue*, Presbytia—*v. Louche*, Strabismus—*v. Nocturne*, Nyctalopia.

VULGA, Vulva.

VULNERA SCLOPETARIA, see Wound—*v. à Sclopetis*, see Wound.

VULNÉRAIRE, Vulnerary—*v. Suisse*, see Falltranck.

VULNERARIUS, Traumatic.

VULNERARY, Vulnerarius, Traumaticus, Traumat'ic, from vulnus, vulneris, 'a wound.' (F.) Vulnéraire. A name applied by the ancients to medicines, which they considered capable of favouring the consolidation of wounds. The property has been ascribed to various plants.

VULNERATIO NERVI, Neurotrosis.

VULNERUM VINCTURA, Deligation.

VULNUS, Wound—*v. Laceratum*, see Laceration—*v. Sclopetarium*, Wound, gun-shot—*v. Simplex*, Cut.

VULNUSCULUM, Wound.

VULPIS MORBUS, Alopecia.

VULTUEUX (F.), Vultuos'ous. Corvisart has used the expression, *Face vultueuse*, *Face vultuos'a*, for the face when it is more florid and swollen than in the natural state. This appearance, which is observed in active aneurism of the heart, according to Corvisart, is owing to the afflux of blood into the arterial capillaries, and hence differs from the livid and swollen appearance caused by the accumulation of blood in the venous capillaries, constituting the *Face injectée*.

VULTUS, Countenance, Face—*v. Tetricus* et Mœstus, Scythrospasmus.

VULVA, Uterus, see Vulva.

VULVA, volva (formerly, the uterus, especially of animals). *Hortus*, *Cunus*, *Puden'dum mulie'brê*, *Sinus pud'o'ris*, *Ictar*, *Choiras*, *Aræum*, *Bacca*, *Bulga*, *Cador'eus*, *Custos*, *Femen*, *An'ulus*, *Femur summum*, *Follic'ulus*, *Fundus*, *Gre'mium*, *Hiat'us*, *Lam'vium*, *Navis*, *Oppid'ulum*, *Specus*, *Vulga*, *Cymba*, *Cythus*, *Saltus*, *Clitor'ium*, *Ost'ium*, *Sulcus*, *Amphicaut'is*, *Concha*, *Porcus*, *Porta*, *Interfemin'eum*, *Mulie'bria*, *Fossa magna mulie'bris*, *Epis'e'on*, *Es'chara*, *Hor'tulus cupid'inis*, *Boubol'ion*, *Byssos*, *Cava*, *Caver'na*, *Del'phys*, *Della*, *For'ea*, *Gynæce'um*, *Loci mulie'bres*, *Female puden'dum*, *Female organs of generation*, *Female parts*, (F.) Vulve. A longitudinal opening between the projecting parts of the external organs of generation in the female, extending from the mons veneris to the perineum, and which is called, also, *Fossa magna*. Some anatomists mean, by this term, the whole of the external genital organs of the female:—mons veneris; labia majora, and the cleft or vestibule separating them; the clitoris; nymphæ; meatus urinarius; entrance of the vagina; with the hymen or the caruncula myrtiformes, fossa navicularis and fourchette.

VULVA, *Fora'men commu'nê ante'rius*, *Iter seu Ad'itus ad infundib'ulum*, *I. seu Ad'itus ad ter'tium ventric'ulum*, *Rima ad infundib'ulum*, *Aper'tura ante'rior ventric'uli ter'tii cer'e'brî*. *Ad*

opening situate at the junction of the optic thalami, in the third ventricle of the brain, and immediately beneath the anterior crus of the fornix.

VULVAIRE, *Chenopodium vulvaria*.

VULVAR, *Vulva'ris*, from *vulva*. That which relates to the vulva. Chaussier gives this name

to the external pudic arteries, because they are distributed to the vulva.

VULVARIA, *Chenopodium vulvaria*.

VULVE, *Vulva*.

VULVITIS, a hybrid word, from *vulva*, and *itis*, denoting inflammation. Inflammation of the vulva.

VULVO-UTERINE CANAL, *Vagina*.

W.

WABRAN LEAVES, *Plantago*.

WAD, Graphites.

WAHOO, see *Eunonymus*.

WAIR, see *Pila marina*.

WAISTCOAT, STRAIT, *Indu'cula*, (F.) *Camisole*, *Gilet de force*. A dress, used for restraining maniacs, or those labouring under violent delirium. It has long sleeves, which are tied behind the body, so that the arms cannot be extricated from them. It ought, of course, to be made of very resisting materials.

WAKE-ROBIN, *Arum maculatum*, *Trillium latifolium*.

WALE, *Wheat*.

WALKING, *Gressus*, *Inces'sus*, *Deambulatio*, *Ambulatio*, *Ead'itis*, (F.) *La Marche*, *Progression*. Sax. *pealcan*, 'to walk.' The action by which we change place, by means of a succession of steps in the same direction. In walking forwards, the centre of gravity is constantly altered; and a series of small, parabolic curves is described on the heads of the thigh-bones, the extent of which is regulated by volition, under the guidance of vision.

WALLA'CHIA, **MINERAL WATERS OF**. There are many excellent sulphureous springs in this country, especially those at Bobotsch, Fincschti, Sibitschiudi Suz, Brása, Otschin, Serbo-neschte, Kimpalungi, Kosia, and Oloneschti, Chalybeate and saline springs have, also, been found at Sibitschiudi Suz.—Siller.

WALLFLOWER, *Cheiranthus cheiri*.

WALLRUE, *Asplenium ruta muraria*.

WALL-TEETH, *Molar teeth*.

WALNUT, *Juglans regia* — w. Black, *Juglans regia* — w. White, *Juglans cinerea*.

WALTHERIA DOURADINHA. A plant of Brazil, *Order*, *Duruliaceæ*, which is used there in diseases of the chest, and in venereal affections. It is a demulcent.

WALTON, **MINERAL WATERS OF**. A mineral spring near Tewkesbury, in Gloucestershire, England. It is an acidulous chalybeate.

WAMPEE, *Peltandra Virginia*.

WANGTEETH, *Molar teeth*.

WANT, from past participle of Sax. *panian*, 'to wane,' 'to fall away.' 'Need of that which is necessary.' (F.) *Besoin*. This word is sometimes used to express the imperious necessity which impels us to take food, drink, &c. (See *Hunger* and *Thirst*.) In a more general sense, it means the impressions transmitted by the organs to the brain, when it is necessary that they should act: these wants are called *instinctive desires* and *internal sensations* or *stimuli*. Such are the desires or wants to eat, drink, void the urine, breathe (*besoin de respirer*), &c.

WANT, in the sense of indigence, (L.) *Egestas*, *Indigentia*, is the cause of a number of diseases.

WARD'S FEVER-DROPS, see *Bebeeru*.

WARD'S ESSENCE FOR THE HEADACH, *Linimentum camphoræ compositum* — w. White Drops, see *Hydrargyri nitras*.

WARE, see *Pila marina*.

WARISON, *Stomach*.

WARK, *Pain*.

WARMBRUNN, **MINERAL WATERS OF**. Much frequented springs in Silesia, about a league from Hirschberg. They are thermal, 95° to 100° Fah., and are chiefly used as baths. The impregnating materials are carbonate of soda, sulphate of soda, sulphate of lime, chloride of sodium, carbonate of magnesia, and sulphohydric acid gas.

WARM SPRINGS, see *Georgia*, *Mineral waters of*, and *Virginia*, *Mineral waters of*.

WARNERA CANADENSIS, *Hydrastis Canadensis*.

WARRENTON SPRINGS, see *Virginia*, *mineral waters of*.

WART, *Verruca* — w. *Corneous*, *Cornu* — w. *Moist*, *Acrothymion*.

WARTY, *Ver'rucose*, *Ver'rucous*, *Verruco'sus*, from Sax. *peapw*. Full of warts. Resembling, relating, or belonging to warts.

WARTY TUMOUR OF CICATRICES. A name given by Mr. Howship to the tumour, which sometimes appears on an old scar, many years, perhaps, after the injury that caused it. The ulcer, formed under such circumstances, has been called the *warty ulcer of Marjolin* — having been well described by that surgeon.

WASH, BLACK, see *Hydrargyri submurias* — w. *Phagedenic*, see *Hydrargyri submurias* — w. *Preventive*, *Lotion*, *Hannay's* — w. *Red*, *Lotie rubra*, and *Lotion*, camphorated, of *Bates* — w. *White*, *Liquor plumbi subacetatis dilutus* — w. *Yellow*, see *Hydrargyri oxyurias*.

WASHER WOMAN'S SCALL, *Psoriasis diffusa*.

WASHITAW SPRINGS, see *Arkansas*, *mineral waters of*.

WASP, *Vespa*, *Sphex*, (F.) *Guêpe*. A genus of gregarious insects, like the bee and the ant. They are armed with a sting, which pours a poisonous fluid into the puncture made by it. (See *Poisons*, *Table of*.) The best application is the sp. *ammoniac aromaticus*, or some preparation of ammonia.

WASSERBURG, **MINERAL WATERS OF**. Wasserburg is a town of Bavaria, seated on the Inn, 28 miles east of Munich. The water, called, also, *Aqua Aga'tii*, contains carbonic acid; carbonates of lime and magnesia; sulphates of lime and magnesia; chloride of sodium, carbonate of soda and oxide of iron.

WASSERKUR, *Hydrosudotherapeia*.

WASTING, *Consumption*, *Tubes*.

WATER, Sax. *wæter*, (G.) *Wasser*, *Aqua*, *Hydor*, *ὑδωρ*, *Alma*, (F.) *Eau*, *Protoxide of hydrogen*. Water was at one time regarded as a simple element. It is now known to consist of eight parts by weight of oxygen, and one of hydrogen. It freezes at 32°, and boils at 212° Fahrenheit: and is taken as the unit in all tables of specific gravity of solid and liquid sub-

stances. Water, as we meet with it, is very different in its state of purity; generally, it contains small quantities of saline matter; at other times, as in sea-water and mineral waters, it is largely impregnated. The characters of water, as met with under ordinary circumstances, are well compared by Celsus, "*Aqua levissima pluvialis est; deinde fontana; tum ex flumine; tum ex puteo; post hæc ex nive, aut glacie; gravior his ex lacu; gravissima ex palude.*" Pure water is one of the best diluents that can be used. When cold, and especially when iced, it is one of the most useful refrigerants; and, in the form of the cold, tepid, and warm bath, it is serviceable in numerous diseases.

WATER OF AIX-LA-CHAPELLE, see Aix-la-Chapelle — w. Allflower, Urine, cows' — w. Alum, Bates's, Liquor aluminis compositus — w. of carbonate of Ammonia, Liquor ammoniæ subcarbonatis — w. Anise, see Pimpinella anisum — w. Apple, Apple-tree — w. Barley, Decoctum hordei.

WATER BED, *Hydrostatic bed*. A bed, contrived by Dr. Arnott, which, in consequence of its being placed on water, is well adapted for the bedridden.

WATER, BINELLI, Aqua Binellii.

WATER OF BITTER ALMONDS, Aqua Amygdalarum concentrata.

WATER BLEBS, Pemphigus — w. Brash, Pyrosis.

WATER, BROCCHERI, Aqua Brocccherii — w. Camphor, Mistura Camphoræ.

WATER CANKER, Cancer aquaticus.

WATER, CARAWAY, see Carum carui — w. Caraway, strong, Spiritus carui — w. of Cassia, see Laurus Cassia.

WATER CASTER, see Uromantia.

WATER, CHERRY LAUREL, see Prunus lauro-cerasus — w. Chicken, Chicken-tea — w. Chlorine, see Chlorine — w. Cinnamon, Aqua cinnamomi — w. Cinnamon, strong, Spiritus cinnamomi — w. Cologne, *Eau de Cologne*.

WATER CURE, Hydrosudotherapia.

WATER, DILL, see Anethum graveolens.

WATER, DISTILLED, *Aqua destillata*, (F.) *Eau distillée*, *Hydrolat*, *Holy Water*, is generally ordered in extemporaneous prescriptions, but is rarely used. In some cases, however, it is absolutely necessary; — the water, containing any saline impregnation — hard water, for example — decomposes certain substances. Distilled water has been recommended as a diet-drink in cancerous affections; and it is used in the religious ceremonies of the Catholic church. In the Pharmacopœia of the United States, it is directed to be made as follows: — Take of water, ten gallons. First distil two pints, and throw them away; then distil eight gallons. Keep the distilled water in glass bottles.

Distilled waters, (F.) *Hydrolats*, consist of common water, charged by distillation with the volatile principles of medicinal agents. See *Aquæ destillatæ*.

WATER-DOCTOR, see Uromantia.

WATER-DRESSING. The treatment of wounds and ulcers by the application of water. It generally consists in dipping folds of lint in water, and placing them on the part: — the whole being covered with oiled silk, elastic gum, or spongiopiline. *Water strapping* or *wet strapping* means the treatment of ulcers, and certain cutaneous affections of the extremities, &c., with strips of linen or cotton saturated with water.

WATER, ELDER, see Sambucus — w. Epileptic of Langius, Langii aqua epileptica — w. Fennel Aqua fœniculi.

WATER FLAG, YELLOW, Iris pseudacorus.

WATER, FLORIDA, — a celebrated perfume, — may be made of *Ol. Bergami* f3iv; *Tinct. benzoin. c.* f3ij; *Alcohol.* cong.

WATERFRIGHT, Hydrophobia.

WATER, GOULARD, Liquor plumbi subacetatis dilutus.

WATER, HARD. Common water, which decomposes and curdles soap; in contradistinction to *soft water*, which is readily miscible with soap. The hardness of water depends upon the presence in it of earthy salts, the most common of which is sulphate of lime.

WATER, HELL, see Toddy.

WATER, HOLY, Water, distilled — w. Honey, see Mel.

WATER IN THE HEAD, Hydrocephalus.

WATER, HUNGARY, Spiritus rosmarini.

WATER JAGS, see Varicella.

WATER, LAKE, *Aqua ex lacu*, (F.) *Eau de lac*, is apt to be impure, from decayed animal and vegetable matters. A great deal will, however, depend upon the magnitude of the collection, and the degree of stagnation.

WATER, LAUREL, see Prunus lauro-cerasus — w. Lavender, Spiritus lavandulæ — w. Lead, Liquor plumbi subacetatis dilutus — w. Lemonpeel, see Lemonpeel Tea — w. Lime, Liquor calcis — w. Lime, compound, Liquor calcis compositus — w. Magnesia, aerated, Magnesia, fluid — w. Magnesia, carbonated, Magnesia, fluid.

WATER, MARSH, *Aqua paludosa*, (F.) *Eau de Marais*, is the most impure of all. It is generally more stagnant, and of course more loaded with decomposing animal and vegetable matters.

WATER, MEDICATED, see *Aquæ medicatæ*.

WATER, MINERAL, *Aqua mineralis seu salubris seu medica* seu *Sote'ria*, *Fons salutaris seu medicatus seu sote'rius*, (F.) *Eau minérale ou médicinale naturelle*. Water, holding in solution different saline and gaseous substances in sufficient quantity to be possessed of medicinal properties, or of a temperature different from that of the ordinary springs of the country.

The following Table by Dr. Perrier exhibits the composition of some of the principal mineral waters.

FIXED CONSTITUENTS

ENTERING INTO THE COMPOSITION OF SOME OF THE MORE CELEBRATED MINERAL SPRINGS.

PROPORTIONS IN 10,000 PARTS OF WATER.

NOTE.—In reducing the analyses contained in this Table to a uniform measure, in order to render them susceptible of direct comparison with each other, the old English gallon is assumed to be as = 58,338 grains; the wine pint = 7,305 grains; the imperial gallon = 70,000 grains; and the German 16-ounce measure = 7,168 grains.

The different salts have been reduced to their elementary constituents by Wollaston's scale of chemical equivalents.

THERMAL.

NAME.	COUNTRY.	ACIDS.			BASES.			Oxide of Iron.	Silica.	SUM.	AUTHORITY AND DATE.	REMARKS.	
		Carbo- nic.	Sulphu- ric.	Muri- atic.	Soda.	Lime.	Mag- nesia.						
Ischia...	Ischia...	0	19.30	29.05	34.50	2.08	2.35	3.19	0.40	94.44	Giudice.....	Iron in the state of ferruginous alumina ; sub-borate of soda 2.79. Iron as in San Restituta. Free carbonic acid 18.9 ; azote ; pot-ash 0.83 ; alumina 0.56 ; organic extract 2.37.	
Gurgitello.....	Ditto.....	14.55	9.05	11.18	31.45	2.95	2.38	1.39	0.56	74.03	Ditto		
Wiesbaden.....	Nassau..	1.38	0.87	33.09	31.99	5.29	0.92	0.05	0.26	57.63	Kastner, 1823.		
Carlsbad Sprudel.....	Bohemia	7.45	14.50	6.40	24.55	1.75	0.85	0.02	0.75	54.59	Berzelius, 1822	Minute traces of phosphoric and fluor- acids, strontian, alumina, and man- ganese.	
St. Nectaire.....	France..	15.13	0.87	13.00	23.90	2.45	1.14	0.14	1.00	53.94	Berthier		
Vichy.....	Ditto....	15.81	1.55	0.72	24.17	0.27	0	0.01	0	42.75	Ditto		
Ems (Kräichensquelle).....	Nassau..	20.32	0.76	2.43	14.87	2.65	1.45	0.65	trace	42.74	Kastner, 1830 ..	Alumina, a trace ; oxide of manga- Potash 2.73. Magnesia, by Seudamore.	
Olmitello.....	Ischia..	0	6.91	5.95	5.98	0	1.80	0	0.42	24.43	Giudice.....		
Bath (King's Bath).....	England.	2.38	8.71	0	2.79	5.29	0.77	0.03	0.37	20.53	Phillips.....		
Mont d'Or.....	France..	2.86	0.36	2.05	4.74	0.90	0.28	0.01	0.10	13.39	Berthie.....	Alumina 0.48. Alumina 0.05. Free carbonic acid 12.99. Phosphoric acid 6.0. Azote 2.01 by Pearson.	
Geyser.....	Iceland..	0	0.81	1.32	2.74	0	0	0	5.40	10.75	Black, 1791.....		
Chaudes Aigues.....	France..	3.07	0	0.86	3.12	1.01	0.59	0.15	1.16	9.96	Berthier		
Ryeum (Reikum).....	Iceland..	0	0.71	1.55	2.43	0	0	0	3.73	8.47	Black, 1791		
Bristol hot well.....	England.	1.02	2.25	1.17	1.17	0.82	0.43	0	0	8.19	Garrick, 1797		
Schlagentbad.....	Nassau..	3.50	0	0.85	2.42	0.90	0.53	0	0	6.96	Kastner, 1823 ..		
Teplitz.....	Bohemia	1.89	0.40	0.29	2.62	0.36	0.18	0.03	0.42	6.24	Berzelius, 1822..		
Buxton.....	England	0.78	0.06	0.28	0.19	1.04	0.04	0	0	2.70	Seudamore, 1820		

WATER, MINERAL

977

WATER, MINERAL

Vicar's Bridge.....	Scotland	0	203.00	0.18	13.20	563.10	Connell, 1831..	Potash, a trace.
Püllna	Bohemia	4.63	182.83	17.20	45.45	0	Struve.....	Potash 3.55.
Seidenschütz	Ditto.....	4.81	97.43	1.43	14.20	0.16	Ditto	{ Nitric acid 7.75; phosphoric acid; pot- ash 3.61; strontian 0.03; alumina; oxide of manganese.
Leamington (Royal Pump)...	England.	0	20.80	71.90	12.00	trace	Thomson, 1830..	Traces of iodine and bromine by Daubeny.
Harrowgate (old sulphur well)	Ditto....	1.20	0.80	76.83	4.95	1.09	Seudamore, 1819	
Airtheigh (first spring)	Scotland	0	4.25	32.10	0.31	0	Thomson, 1828..	
Cheltenham (old well).....	England.	0	11.05	50.65	32.50	0	Seudamore, 1819	Trace of iodine by Daubeny.
Hartfell aluminous chalybeate	Scotland	0	63.23	0	4.26	trace	Thomson, 1828..	Alumina 5.10.
Isle of Wight.....	England.	0	48.58	2.95	0	40.00	Marcet	Alumina 7.77.
Marienbad (Ferdinandsquelle)	Bohemia	9.52	27.60	9.50	5.68	0.96	Steinmann, 1820	{ Phosphoric acid, lithion, strontian, alu- mina, manganese, by Berzelius.
Dunblane (north spring).....	Scotland	0.30	2.23	33.55	2.87	0.12	Murray, 1814...	
Vals	France..	22.48	0.29	15.30	14.04	0	Berthier	
Bilin	Bohemia	20.51	3.40	33.34	0.22	0.45	Reuss, 1788....	
Franzenbad (Franzensbrunn).	Ditto....	5.44	18.50	6.21	2.47	0	Tromsdorf, 1820	{ Phosphoric acid, lithion, strontian, alu- mina, manganese, by Berzelius.
Pitcaithly	Scotland	0.30	0.73	26.70	1.26	0.48	Murray, 1814...	
Reisdorf	Rhenish	6.76	2.50	8.50	13.99	trace	Bischoff, 1826..	Potash.
Epsom	Prussia	2.50	14.80	16.00	1.78	0.21	Daubeny, 1830..	Trace of bromine.
Selters (Nieder).....	Nassau..	5.37	0.18	5.64	0	0	Bischoff, 1826..	Phosphoric acid 0.19.
Fachingen.....	Ditto....	11.49	0.12	16.06	1.37	0.38	Ditto	Phosphoric acid 000.5.
Soden	Ditto....	3.35	0.07	2.63	1.83	0.11	Meyer, 1820....	
Moffat	Scotland	0	2.80	10.72	4.08	0.06	Thomson, 1828..	
Pymont	Germany	4.03	9.83	12.75	0.68	0	Struve.....	{ Phosphoric acid, potash, strontian, man- ganese.
Marienfels.....	Nassau..	4.83	0.18	1.29	85.6	0.68	Kastner.....	{ Potash 1.19; strontian, manganese, phos- phoric acid.
Strathpeffer (pump-room)...	Scotland	0	9.32	3.42	2.20	trace	Thomson, 1828..	Potash.
Gellnau	Nassau..	6.00	0.07	6.18	2.33	0	Bischoff, 1826..	Phosphoric acid .019.
Wellbach	Ditto....	3.61	0.81	5.08	1.46	0.14	Creve, 1810....	Sulphur-resin 0.48.
Hartell Spa	Scotland	0	2.76	4.60	2.05	0	Thomson, 1828..	Alumina, a trace.
Langenschwalbach (Wein- brunn)	Nassau..	3.84	0.12	3.05	2.30	0	Kastner, 1829...	{ Potash, lithion, iodine, strontian, alu- mina, manganese, phosphoric acid.
Spa	Belgium	6.63	0.51	0.12	1.55	0.67	Struve	{ Phos. acid; potash 0.58; manganese.
Carlsbad (Sauerling)	Bohemia	0.28	0.23	0.07	0.74	0.68	Berzelius, 1822.	{ Phosphoric acid, fluato of lime, alumina, oxide of manganese.
Tunbridge	England.	0.20	0.14	0.29	0.39	0.37	Seudamore, 1816	Oxide of manganese.
Malvern.....	Ditto....	0.32	0.14	0.08	0.02	0	Phillip, 1805....	

Mineral waters may be divided into four classes:—1. *Gaseous* or *Acidulous*. 2. *Chalybeate*. 3. *Saline*. 4. *Sulphureous*. These may be *thermal*, or *cold*,—*natural*, or *artificial*. Many of these divisions, however, run into each other; some of the substances, which they hold dissolved, belonging to one class as well as to another.

1. *Gaseous, Acidulous* or *Carbonated Mineral Waters*, *Soda* or *Beer Springs*, *Aqua mineralis acidula*, (F.) *Eaux minérales gazeuses ou acides*. The waters referred to this class are those that contain carbonic acid gas in such quantity as to communicate to it certain sensible qualities. Waters impregnated with free carbonic acid, sparkle when drawn from the spring, or when poured into a glass. They have a sharp, acidulous taste; but become vapid from exposure to the air. Along with the carbonic acid, there are generally present portions of saline, earthy or metallic matter, chiefly carbonates of lime, magnesia, and iron. Waters, highly impregnated with carbonic acid gas, are grateful to the stomach; increase the appetite, and are diuretic; hence, their utility in dyspepsia, hypochondriasis, and gout. Their properties are, of course, modified by the saline matter that may be also contained in them. The most celebrated amongst the acidulous waters are those of Bar, Chateaudon, St. Myon, Mont d'Ore, Langeac, Seltzer, Schlangenbad, Sultzmat, Pyrmont, Spa, Carlsbad, Cheltenham, Scarborough, Saratoga, Ballston, and the Sweet Springs of Virginia.

2. *Chalybeate* or *Ferruginous Mineral Waters*, *Aqua mineralis ferruginea* seu *Chalybeata* seu *Martialis* seu *Ferroea*, *Chalybeo-gæ*, *Chalybokre-na*, *Sidero-pe-gæ*, *Siderokre-na*, (F.) *Eaux minérales ferrugineuses*, contain iron—sulphate, chloride or carbonate, generally the latter—held in solution by an excess of acid. Chalybeate waters have a peculiar styptic taste. They are transparent when taken from the spring; but, when exposed for some time to the air, a pellicle forms on the surface, and a deposit of the iron takes place. Chalybeate waters are used as tonics in debility of all kinds; in all cases, in fact, where iron is considered to be indicated. They are the most numerous of all the classes of mineral waters. In this dictionary, those only are specified which have some celebrity; yet there are about one hundred; more than sixty of these being saline, and forty sulphureous. The most celebrated chalybeates are those of Tunbridge, Scarborough, Spa, Bussang, Forges, Vichy, Pyrmont, Passy, Provins, and Vals.

3. *Saline Mineral Waters*, *Aqua minerales salina*, *Anna alcalizata* (Paracelsus), *Hydralma*, (F.) *Eaux minérales salines*. Waters, holding in solution different saline substances in greater quantity than the acidulous waters. They differ in properties, according to the salts contained in them. The salts usually present are sulphates, chlorides, and carbonates: and the bases, with which the acids forming these are combined, are soda, magnesia, and lime. Saline mineral waters are usually aperient; and the most noted are those of Seltzer, Sedlitz, Balaruc, Bourbonne-les-Bains, Baden, Epsom, Cheltenham, &c.

To this class may also be added *Sea water*.

4. *Sulphureous Mineral Waters*, *Aqua minerales sulphureæ*; when warm, termed *Theotherma*, *Theope-gæ*, *Theotherma*, *Fontes sulphurei calidi*, (F.) *Eaux minérales sulfureuses ou sulfurées ou sulfureuses ou hépatiques*, &c. These waters owe their distinguishing character to an impregnation of sulphuretted hydrogen, and are at once recognised by their peculiar fetid smell, resembling that of rotten eggs. They usually contain saline substances, which modify their

powers. From the action of the sulphuretted hydrogen, they are useful in cutaneous affections; and, from the combined action of this and the saline matter, they are valuable agents in diseases of the digestive organs. They are also employed in cutaneous eruptions; and the warm sulphur baths have been especially celebrated in such cases, as well as in rheumatic affections. The most renowned sulphureous waters are those of Baréges, Bagnères-de-Luchon, Cauterets, Bonnes, Aix in Savoy, Aix-la-Chapelle, Enghien, Harrogate, Moffat, Virginia Springs, &c.

Some springs, as those of Bath, Buxton, Dax, Matlock, Warm and Hot Springs of Virginia, &c., are almost pure *Thermal Mineral Waters*. They are valuable agents in rheumatic affections; the warmth being equable during the whole period the individual is immersed in them; which cannot be the case in the artificial thermal bath.

WATER, MINERAL, Acidulous water, simple.

WATERS, MINERAL, ARTIFICIAL, (F.) *Eaux minérales artificielles ou factices*. These are imitations of the natural; and some of them—as the factitious Cheltenham water, and Sedlitz water—answer the purpose of the natural water tolerably well. The acidulous and chalybeate waters are, however, most easily imitated.

WATER, NUTMEG, Spiritus myristicæ.

WATER PANG, Pyrosis—w. Parsnep, creeping, Sium—w. Pennyroyal, spirituous, Spiritus pulegii—w. Peppermint, Aqua menthæ piperitæ—w. Pimento, see Myrtus pimenta.

WATER POX, see Variella.

WATER, PROTOXIDE OF NITROGEN, Aqua nitrogenii protoxydi.

WATER PURPLE, Veronica beccabunga—w. Qualm, Pyrosis.

WATER, RAIN, *Aqua pulvisalis* seu *pluvia* seu *imbrum*, (F.) *Eau de pluie*, when collected at a distance from houses or other elevated objects, is the purest natural water, and has the least specific gravity. The only bodies which it usually holds in solution, are carbonic acid, and minute traces of carbonate of lime and chloride of calcium.

WATER, RICE, see Oryza.

WATER, RIVER, *Aqua fluvialis*, is derived from the conflux of numerous springs and rain water. It is, generally, pretty pure.

WATER, ROSE, Aqua rosæ.

WATER, SEA, *Aqua Marina*, *Humor Doridis*, (F.) *Eau de Mer*, *Eau salée*. This contains chlorides of sodium, magnesium, and calcium, and sulphate of magnesia. It is cathartic, and forms the usual gyster at sea. It makes an excellent tonic bath; superior in most cases—especially in scrofula—to the fresh-water bath.

WATER, SEARLE'S PATENT OXYGENOUS AERATED, Aqua Nitrogenii protoxydi—w. Seydchütz, Sedlitz water.

WATER SHIELD, Brasenias hydropeltis.

WATER, SNOW, *Aqua nivata*, (F.) *Eau de neige*, has usually been deemed unwholesome. It exactly resembles rain water in composition, and is equally salubrious.

WATER, SODA, Acidulous water, simple—w. Soft, see Water, hard—w. Spearmint, Aqua menthæ viridis.

WATER, SPRING, *Aqua fontana*, *Hydrope-gæ*, (F.) *Eau de fontaine*, contains, in addition to the substances detected in rain water, more or less sulphate of lime. When this is to such an extent as to curdle soap, the water is said to be *hard*; if not, *soft*. Hard water is, of course, inferior to soft, for domestic and medicinal purposes.

WATER STRAPPING, see Water dressing.

WATER, STYPTIC, SYDENHAM'S, Solutio sul-

phatis supri composita—w. Tar, see *Pinus sylvestris*—w. Toast, see *Toast-water*.

WATER TARGET, *Brasenia hypopeltis*.

WATER, WELL, *Aqua putealis* seu *puteana* seu *ex puteo*, (F.) *Eau de puit ou de source*, is the same as spring water, but liable to impregnation, owing to the land springs filtering into the wells, and conveying impurities into them.

WATERS, THE, Liqueur Amnii—w. Bag of, see *Liquor Amnii*—w. Discharged or broken, *Profusio aquæ*—w. Distilled, *Aquæ destillatæ*—w. First, *Primitiæ*—w. Green, see *Lochia*.

WATTWEILER, MINERAL WATERS OF. Wattweiler is a small town in the department of Haut-Rhin, at the foot of the Vosges. The waters are acidulous chalybeates, and contain carbonates of iron, lime, and soda, chloride of sodium, and free carbonic acid.

WAX, Cera—w. Ear, *Cerumen*.

WAX, MYRTLE. A wax obtained from the fruit of *Myrica cerifera*. It has been prescribed in dysentery.

WAX, YELLOW AND WHITE, Cera flava et alba.

WAXING KERNELS, Waxen-kernels, Crescentia, (F.) *Croissances, Croisants, Glandes de croissance*. A popular term for small tumours, formed by an enlargement of the lymphatic glands—in the groins of children particularly. They have been looked upon as connected with the growth of the body,—hence their name.

WAY, Saxon *pæz*, *Via, Passage, Odos*, (F.) *Voie*. A name given to different canals.

WAYBREAD, Plantago major.

WAYS, DIGESTIVE, Digestive Passages, Primæ Viæ, (F.) *Voies digestives, Premières voies*, are the series of hollow organs of digestion, composed of mouth, œsophagus, stomach, and small and large intestines. The term is often restricted to the first three of these.

WAYS, SECOND, Second Passages, Secundæ viæ, (F.) *Secondes voies*, comprise the chyliferous, lymphatic and blood vessels.

WEAKSIGHTEDNESS, Asthenopia.

WEAL, Wheal.

WEAM, Abdomen, Uterus.

WEAM-ILL, Colic.

WEANING, from Saxon *penan*, 'to wean,' *Ablactatio, Apogalactismus*, (Prov.) *Spaning*, (F.) *Sévrage*. The act of depriving the infant, permanently, of the breast, in order to feed it on other and more solid nourishment.

WEANING BRASH, see Brash, weaning.

WEARING, Consumption.

WEATHERCOCKS, Impatiens.

WEB, Caligo, Pterygium—w. Muscular, *Panniculus carnosus*.

WEB-EYE, Caligo.

WEBBED FINGERS, Dactylion.

WEBERIAN CORPUSCLE, see Ejaculatory Ducts—w. Organ, see *Ejaculatory Ducts*.

WEDGED, (D.) *wegge*, *Inclusus, Inhaerens*, (F.) *Enclavé*. The head of the fœtus is said to be *wedged, locked or impacted* in the pelvis, when it remains fixed, notwithstanding the uterine efforts.

The condition is called *Incuneatio* seu *Inclavatio fœtus, Sphenosis*, (F.) *Enclavement*. See *Paragomphosis*.

WEEBO, Senecio Jacobææ.

WEED, see Mastitis—w. Arrow, *Sagittaria variabilis*—w. Beetle, *Galax aphylla*—w. Breast, *Saururus cernuus*—w. in the Breast, see *Mastitis*—w. Butter, *Erigeron Canadense*—w. Canker, *Goodyera pubescens*—w. Dyer's, *Reseda luteola*—w. Field, *Anthemis cotula*—w. Horse,

Erigeron Canadense—w. Horsefly, *Sophora tinctoria*—w. Milk, *Apocynum androsaemifolium*—w. Phthisis, *Ludwigia palustris*—w. Pride, *Erigeron Canadense*—w. Silver, *Potentilla anserina*—w. Soldiers', *Matico*—w. Thimble, *Rudbeckia laciniata*—w. Wind, *Asclepias tuberosa*.

WEEPING, Fletus—w. Tear, *Tear*.

WEESEL, Trachea.

WEID, see Mastitis.

WEIGHT, from Saxon, wægan, 'to weigh,' (F.) *Pesanteur*. A sensation of heaviness or pressure over the whole body, or over a part—the stomach or head, for example.

WEIGHTS AND MEASURES, Pondera et Mensuræ, (F.) *Poids et Mesures*. The importance of possessing a uniform system of weights and measures has impressed the scientific of all countries, and numerous endeavours have been made to accomplish the object. It is, however, a matter of considerable difficulty, and one not likely to be attained. The new French measures are upon decidedly the best footing, but they are not adopted out of France. The greatest diversity prevails over Europe in the measures, both of weight and capacity. Some of the following tables will show, that every subdivision of the pound, as well as the pound itself, differs in England and in France.

WEIGHTS.

Troy Weight, as used by the British Apothecari.
—*Pondus pharmaceuticum seu medicinale.*

The pound lb	}	contains	12 ounces.
The ounce oz			8 drachms.
The drachm ℥			3 scruples.
The scruple ℥			20 grains.
The grain gr.			

These, and the signs by which they are denoted, are the same in all the British Pharmacopœias, as well as in the American.

lb.	Oz.	Drachms.	Scruple.	Gr.
1	12	96	288	5760
	1	8	24	480
		1	3	60
			1	20

Poids de Marc.

The pound lb	}	contains	16 ounces.
The ounce oz			8 drachms.
The drachm ℥			3 scruples.
The scruple ℥			24 grains.

Avoirdupois.

Pound.	Ounces.	Drachms.	Troy grains.
lb. 1.	16	256	7000
	1	16	437.5
		1	27.34375

The avoirdupois drachm is sometimes divided into three scruples, and the scruple into ten grains. The pound of 7680 grains avoirdupois = 7000 grains troy, and hence 1 grain troy = .97 grain avoirdupois.

The *Poids de Marc* is that employed by the French Pharmacians, when the new weights are not. The Avoirdupois is now used by the Dublin College.

The following tables exhibit the relative value of the old French and English weights:

Poids de Marc.	Troy Wt.	Avo. d.	Troy grs.
1 pd (livre) =	1.31268 lb =	1.080143 lb =	7561
1 oz. (once) =	.984504 oz =	1.080143 oz =	47.25625
1 dr. (gros.) =	.954504 dr =		59.0703125
1 gr. =			.520421

Troy.	Poids de Marc.	French grains.
1 pound =	0.76180 lb.	= 7561
1 ounce =	1.01574 once	= 585.083
1 drachm =	1.01574 gros	= 73.135
1 grain		= 1.219

Avoirdupois.	Poids de Marc.	French grains.
1 pound =	0.925803 lb.	= 8532.3
1 ounce =	0.925803 once	= 533.27

To convert	French grains into Troy grains, divide by	1.2189
"	Troy grains into French grains, multiply by	
"	French ounces into Troy ounces, divide by	1.015734
"	Troy ounces into French ounces, multiply by	
"	French pounds (poids de marc) into Troy pounds, multiply by	1.31268
"	Troy pounds into French pounds, divide by	

Troy grain.	Poids de Marc or French grain.
1	= 1.219
2	= 2.438
3	= 3.657
4	= 4.876
5	= 6.095
6	= 7.314
7	= 8.533
8	= 9.752
9	= 10.971

French grain.	Troy grain.
1	= 0.8203
2	= 1.6406
3	= 2.4609
4	= 3.2812
5	= 4.1015
6	= 4.9218
7	= 5.7421
8	= 6.5624
9	= 7.3827

New French Decimal or Metrical Weights.

	Troy grains.	lb. oz. dr. gr.
Milligramme =	.0154	
Centigramme =	.1543	
Decigramme =	1.5434	
Gramme =	15.4340	
Decagramme =	154.3402	= 0 3 1 34.4
Hectogramme =	1543.4023	= 0 3 1 44.4
Kilogramme =	15434.0234	= 2 8 1 24
Myriagramme =	154340.2344	= 26 9 6 0

Comparative Value of the French Medicinal Pound and that of other Places.

	lbs. oz. dr. gr.	
Germany.....	0 15 4 48	
Amsterdam.....	1 00 0 42	
Belgium.....	1 00 0 42	
Berne.....	1 00 0 00	
Copenhagen.....	0 15 3 20 1/4	
Florence.....	0 11 0 50	divided into 12 oz.
Genoa.....	0 10 5 60	do. into 12 oz.
Geneva.....	1 00 0 18	
Hamburg.....	0 15 2 15	
Lisbon.....	0 15 7 68	
London.....	0 12 3 12	divided into 12 oz.
Madrid.....	0 15 0 16	
Milan.....	0 9 3 00	
Monaco.....	0 15 2 23	
Naples.....	0 10 7 54	divided into 12 oz.
Paris.....	1 00 0 00	divided into 16 oz.
Rome.....	0 11 0 50	do. into 12 oz.
Stockholm.....	0 13 7 8	
United States.....	0 12 3 12	divided into 12 oz.
Warsaw.....	1 10 4 24	
Verona.....	0 8 6 00	
Vienna.....	1 2 2 32	

MEASURES OF CAPACITY.

English Measure of Fluids.

The gallon (congius).	0	contains {	8 pints.
The pint (octarius) ..	f 3		16 fluidounces.
The fluidounce.....	f 3		8 fluidrachms.
The fluidrachm.....	f 3		60 minims.
The minim.....	℥		

Proportions of the Apothecaries' and Wine Gallon.

Gal. Pints.	Fluidoz.	Fluidr.	Minims.	Cub. inches.
1 = 8	= 128	= 1024	= 61440	= 231
1 = 1	= 16	= 128	= 7680	= 28.875
	1 =	8	= 480	= 1.6047
		1 =	60	= .2256

Imperial Measure, adopted by the London College in their Pharmacopoeia of 1836.

Gallon.	Pints.	Fluidounces.	Fluidr.	Minims.
1 = 8	= 160	= 1280	= 76800	
1 = 1	= 20	= 160	= 9600	
	1 =	8	= 480	
		1 =	60	

Comparative Value of the Proportions of the Wine and Imperial Gallons.

Wine.	Pints.	Fluidoz.	Fluidr.	Minims.
1 gallon =	6	13	2	23
1 pint =		16	5	18
1 fluidounce =		1	0	20
1 fluidrachm =			1	2 1/2

Imperial.	Gallon.	Pint.	Fluidoz.	Fluidr.	Minims.
1 gallon =	1	1	9	5	8
1 pint =		1	3	1	38
1 fluidounce =				7	41
1 fluidrachm =					58

French Measures of Capacity.

	English cubic inches.	Wine Measure.
Millilitre =	.061028	= 16.2318 minims.
Centilitre =	.610280	= 2.7053 fluidrachms.
Decilitre =	6.102800	= 3.3816 fluidounces.
Litre =	61.028000	= 2.1135 pints.
Decalitre =	610.280000	= 2.6419 gallons.
Hectolitre =	6102.800000	
Kilolitre =	61028.000000	
Myrialitre =	610280.000000	

Approximate comparison between the ancient French Measures of Capacity, and the new, and conversely.

	Grammes.	Litres.	Decilitres.	Centilitres.
1 poisson =	125	or 0	1	2
1 demi-setier =	250	or 0	2	5
1 chopine =	500	or 0	5	0
1 pintle =	1000	or 1	0	0

	Grammes.	lbs. oz. dr. gr.
1 centilitre =	10	or 0 0 2 1/2 00
1 decilitre =	100	or 0 3 2 00
1 litre =	1000	or 2 0 3 36 = 2.113 Eng. wine

Measures of Length.

	Inches.
1 line, the 12th part of an inch.	
3 barleycorns	1.000
A palm or hand's breadth (Scripture measure).....	3.648
A hand (horse measure)	4.000
A span (Scripture measure).....	10.944
A foot.....	12.000
A cubit (Scripture measure for common purposes).....	18.000
A cubit (Scripture measure for sacred purposes).....	21.888
A Flemish ell.....	27.000
A yard.....	3 ft. 00
An English ell.....	3 " 09
A fathom or toise.....	6 " 00

New French Measures of Length.

	English inches.
Millimètre	0.039
Centimètre	0.393
Decimètre	3.937
Mètre	39.371 = yd. 1 ft. 0 3/4

VALUES OF THE GRECIAN, ROMAN, AND ARABIAN
WEIGHTS AND MEASURES IN POIDS DE MARC.

1. *Weights of the Ancient Greeks.*

	lbs.	oz.	dr.	gr.
The talent (ταλαντον).....	54	2	5	24
The mina (μνᾶ).....	14	3	40	
The drachm (δραχμη).....	1	11		
The obolus (οβολος).....				13
The ceration (κερατιον).....				4
The chalcus (χαλκος).....				2
The septon (σεπτον).....				1/8

Dr. Milligan, in his edition of Celsus, gives the following table, exhibiting the Troy weight of measures of capacity and of weight in use amongst the Romans.

	Urna.	Libra.	Uncia.	Denarius.	Scrupulus.	Sextans.	Chalcus.	Grana.
Amphora.....	2	= 80	= 960	= 6720	= 20160	= 40320	= 403200	= 420480
Urna.....	1	= 40	= 480	= 3360	= 10080	= 26160	= 210600	= 210240
Congius.....	1/4	= 10	= 120	= 840	= 25440	= 5040	= 50400	= 52920
Sextarius.....	1/8	= 5	= 60	= 420	= 12720	= 2520	= 25200	= 26460
Libra.....	1	= 1	= 12	= 84	= 252	= 504	= 5040	= 5256
Hemina.....			8 4-7	= 60	= 180	= 360	= 3600	= 3759
Acetabulum.....			2 1-7	= 15	= 45	= 90	= 900	= 939
Esquicyathus.....			2 1-7	= 15	= 45	= 90	= 900	= 939
Cyathus.....			13-7	= 10	= 30	= 60	= 600	= 626
Sesuncia.....			1 1/2	= 10 1/2	= 31 1/2	= 63	= 630	= 657
Uncia.....			1	= 7	= 21	= 42	= 420	= 438
Cochleare.....				2 1/2	= 7 1/2	= 15	= 150	= 156
Drachma.....				1	= 3	= 6	= 60	= 62 4-7
Denarius.....				1	= 3	= 6	= 60	= 62 4-7
Scrupulus.....					1	= 2	= 20	= 20 2/3
Scrupulus dimidiatus.....					1/2	= 1	= 10	= 10 1/3
Obolus.....						1	= 10	= 10 2/3
Sextans.....						1	= 10	= 10 2/3
Chalcus.....							1	= 11 30

He gives also the following 'Carmen Mnemonicum,' which exhibits the analogies of the Roman and British imperial weights.

To the congius and gallon, each, ten pounds allow,

On the bushel and amphora eighty bestow;
Rome's pound, as in Troy weight, twelve ounces obtain,

But her ounce is Abordupois, strictly, in grains:

Denarii drams, scruples scrupls define,
Septarius answers to our bottle of wine,
And the mina's a sport pint—fourteen ounces in fine.

3. *Weights said to be of the Arabians, Modern Greeks, and Latins of the barbarous periods of the Middle Ages.*

	oz.	dr.	gr.
The alchemien.....	14	3	40
The manes or ominos.....	10	6	28
Sacros, aughben, adar, assatil.....			7 16
The great or royal nut.....			3 44
The sextarius, staler.....			3 44
The lesser nut.....			2 50
Alivianus.....			2 29
Aureus, Alcobolus.....			2 14
The hazel nut, bendacate, holca, alchi, dar-chimi, adogchilos, ologinat, nabach.....	1	11	
The acorn, lupine, Egyptian or Syrian bean, the bacchid.....			42
The Alexandrian bean or tremessis.....			30
The Greek bean or gramme, the kermel, gonum, harmi, grachus.....			21
The ring, cumulus, seminet, onolotich, onolossat.....			11
The danich.....			8
The kirat, alkilat, kararil.....			4

2. *Weights of the Ancient Romans.*

	oz.	dr.	gr.
The pound (libra).....	10	6	48
The ounce (uncia).....			7 16
The duella.....			2 29
The sicilicus.....			1 58
The sextula.....			1 14
The consular denier, denarius.....			1 2
The imperial denier or drachm, drachma.....			65
The victoriatus.....			37
The scripulum or scruple.....			21
The obolus.....			11
The stilqua.....			4

The Greeks divided their obolus into *chalci* and *lepta*: some divided it into 6 *chalci*, and each *chalcus* into 7 *lepta*; others into 8 *chalci*, and every *chalcus* into 8 *lepta* or *minuta*.

1. *Greek Measures.*

	lbs.	oz.	dr.	gr.
The metretes (μετρητης).....	84	4	3	00
The chus, choa, congius (χαψ).....		7	0	2 66
The xestes (ξεστης).....	1	1	7	44
The cotyla (κοτυλη).....			8	7 58
The tetartion (τεταρτιον).....			4	3 65
The ozybaphon (οζυβαφον).....			2	1 69
The cyathos (κυαθος).....			1	4 00
The concha (κοχηνη).....				6 00
The mystron (μυστρον).....				3 00
The chama minor (χημη).....				2 00
The cochliarion (κοχλιαριον).....				1 00

2. *Roman Measures of Capacity.*

	lbs.	oz.	dr.	gr.
The amphora or cadus.....	56	9	7	24
The urna.....	28	1	3	48
The congius.....			7	0 2 66
The sextarius.....			1	4 7 44
The hemina.....				10 1 18
The quartarius.....				5 0 45
The acetabulum.....				4 2 23
The cyathus or small glass.....				1 5 30
The ligula or spoonful.....				3 21

3. *Measures said to be of the Arabian, Arabist, and Latin Physicians of the Middle Ages.*

	lbs.	oz.	dr.	gr.
The missohacs weighed.....	3	8	1	33
Aben, kirt, ejub, eberia, or Roman mina.....				1 6 0 00
The phid, hassilutis, or hassilutinus.....				10 1 18
The caliz or rejelat.....				5 0 44
The handful, pugillum, cornutum.....				8 2 68
The hassuf, cessas, or anesime.....				2 2 20
The conos or coalus, alcantus or almunet, briate cuabus.....				1 5 54
The lesser bacchales.....				5 56
The largest spoonful.....				4 44
A spoonful.....				1 52
The celones or rechanarium.....				1 62
A small spoonful or flugerina, or cyanes.....				1 11
The smallest spoonful or fuhaiet.....				42

A good section on weights and measures is contained in the edition of "The Seven Books of Paulus Ægineta," by Mr. Francis Adams, published by the Sydenham Society of London, vol. iii. p. 609, London, 1847.

Besides the weights and measures above mentioned, employed by the moderns, there are modes of estimating the quantities of substances by approximation. For example:—

A *glassful* or *cupful*, (F.) *Verre*, is reckoned to contain 4 or 5 fluidounces.

A *wine-glassful*, $\frac{1}{2}$ ounce or two ounces.

A *table-spoonful*, (F.) *Cuiller à bouche*, about half an ounce.

A *coffee* or *dessert-spoonful*, about 3 drachms.

A *tea-spoonful*, a fluidrachm.

A *handful*, *manipulus*, (F.) *Poignée*, as much as can be held in the hand.

The *Pugillus* (F.) *Pincée* is as much as can be held by the three fingers.

These last quantities are occasionally prescribed by the French practitioners. Their weights must, of course, vary, according to the article. The authors of the Parisian codex have valued them, as regards certain substances.

	French.	
	oz. dr.	
A <i>Manipulus</i> of barley weighs.....	3	$2\frac{1}{2}$
" linseed.....	1	4
" linseed meal.....	3	3
" dried mallow leaves.....	1	3
" dried cicory leaves.....	1	0
" flowers of the tilia.....	1	$2\frac{1}{2}$
	dr. scr. gr.	
A <i>Pugillus</i> of chamomile flowers weighs 2		
" arnica.....	1	2
" marsh-mallow.....	1	1
" mallow.....		60
" fennel seeds.....	1	60
" aniseed.....	1	$\frac{1}{2}$ 60

A *hen's egg*, newly laid, weighs about 2 ounces and 2 drachms: when deprived of its shell, 2 ounces. The *white* weighs 1 ounce, 2 drachms, and 57 grains: the *yolk*, 5 drachms and 15 grains.

WEILBACH, MINERAL WATERS OF. The village of Weilbach is in the plain between the Maine and the southern extremity of the Taunus hills. It has in its neighbourhood a cold sulphureous alkaline spring, the water of which is bottled and exported, although not of great efficacy. Its use is at times combined with that of the saline springs in Nassau, and of Baden-Baden, in various chronic abdominal and thoracic affections.

WEINTRAUBENKUR, Grape cure.

WELCOME TO OUR HOUSE, *Euphorbia cyparissias*.

WELD, *Reseda luteola*.

WELK, *Welk*. An inequality; a protuberance; a cutaneous eruption of this character. Acne.

WELTER'S BITTER, Acid, carbazotic.

WEM, Abdomen, Uterus.

WEMDING, MINERAL WATERS OF. These waters are situate about four leagues from Donauwörth in Bavaria. They contain sulphuretted hydrogen; carbonates of lime, magnesia, and soda; sulphates of lime and magnesia; a little chloride of calcium, and oxide of iron; and are much used in asthenic diseases.

WEN, Sax. *pen*, *Lu'pia*, *Tumour*, (F.) *Loup*. A name given to circumscribed, indolent tumours, without inflammation or change of colour of the skin. They may be met with in almost every part of the body. Their magnitude, as well as nature, is very variable. The term is sometimes given to an encysted tumour, and to bronchocele.

WESTERN ISLANDS, Azores.

WEST INDIES, (CLIMATE OF.) The climate of the West Indies has long been regarded as favourable to the consumptive. In confirmed consumption, it appears, however, to be positively injurious. As a prophylactic for those predisposed to consumption, it is generally favourable. The nature of the patient's constitution, in regard to the effects of elevated temperature, has always to be taken into account. In diseases that are benefited by warm climates, the West Indies afford an appropriate residence. The islands in which the invalid can best obtain all the advantages of the climate, are Jamaica, Barbadoes, St. Vincents, Antigua, St. Kitts, and Santa Cruz.

WEST'S SPRING, see South Carolina, mineral waters of.

WET STRAPPING, see Water dressing.

WHAHOO, see *Eunymus*.

WHEAL, *Weal*, *Wale*, *Pomphus*. A ridge, or elevation of the skin, produced by a rod or whip; or as if produced in that manner. Such elevations are seen in urticaria.

WHEAT, *Triticum*—w. Indian, *Zea mays*—w. Turkey, *Zea mays*.

WHEEZE, (Prov.) *Rezzle*, Sax. *hpeoron*, formed probably, like *whiz* and *whisper*, from the sound it expresses. To breathe with difficulty and noise, as in asthma.

WHELK, *Acne*, *Welk*—w. Chin, *Sycosis*—w. Rosy, *Gutta rosea*.

WHETLEBONES, *Vertebræ*.

WHEY, Serum lactis.

WHEY-CURE, (G.) *Molkenkur*. In the Kur-saal of many of the German watering places, a space is set apart for whey drinkers. The whey is a pleasant diluent, and the saccharine matter it contains may act as a eutrophic; but the main effect on the valetudinarian is probably produced by the new impressions made upon him by his visit to the watering place.

WHEY, MUSTARD, see *Sinapis*—w. Rennet, see *Serum lactis*—w. Sack, *Wine-whey*—w. Tamarind, see *Tamarindus*—w. Vinegar, see *Acetum*—w. Wine, *Wine-whey*.

WHIFFING MURMUR, see *Murmur*, respiratory.

WHIG, Buttermilk, Serum lactis.

WHIMBERRY, *Vaccinium myrtillus*.

WHISKY, see Spirit. A spirituous liquor obtained from oats, potatoes, &c., by distillation.

WHISKY LIVER, Liver, nutmeg.

WHISPER, CAVERNOUS, see *Cavernous*.

WHIS'PERING; evidently an example of onomatopoeia (*hoois'pering*), *Susurra'tio*, *Susurra'tion*. Articulation of the air sent through the vocal tube without any action of the glottis.

WHISPERING SOUFFLE, see *Souffle*.

WHISTLING, *Sifflement*, see *Râle sibilant*.

WHITE BALL, *Cephalanthus occidentalis*—w. Blisters, *Rupia escharotica*.

WHITE, FLAKE, *Plumbi subcarbonas*.

WHITE LEAF, *Pyrola maculata*, *Spiræa tomentosa*—w. Leg, *Plegmatia dolens*—w. Matter of the brain, see *Cerebrum*—w. Root, *Angelica lucida*, *Asclepias tuberosa*, *Ligusticum actæifolium*.

WHITE, SPANISH, Bismuth, subnitrate of.

WHITE SUBSTANCE OF SCHWANN, see Nerve fibre—w. Sulphur, see *Virginia*, mineral waters of—w. Swelling, *Hydrarthrus*—w. Swelling of lying-in women, *Plegmatia dolens*—w. Weed, *Chrysanthemum leucanthemum*—w. Wood, *Liriodendron*, *Tilia Americana*.

WHITENESS OF COMPLEXION, Paleness.

WHITES, THE, *Leucorrhæa*.

WHITLEYA STRAMONIFOLIA, *Anisodus luridus*.

WHITLOW, Paronychia.

WHOOPIING-COUGH, Pertussis.

WHORLYWORT, *Leptandria purpurea*.

WHORTLEBERRY, *Vaccinium myrtillus*,
Vaccinium vitis idææ — *w. Beas'*, *Arbutus uva*
ursi.

WHORTS, BLACK, *Vaccinium myrtillus*.

WHYTT'S DISEASE, Hydrocephalus in-
ternus.

WIDOW-WAIL, *Cneorum tricoccum*, *Daphne*
Alpina.

WIESBADEN, MINERAL WATERS OF,
see Wiesbaden.

WIESSAU, MINERAL WATERS OF. These
waters are situate about four leagues from the
convent of Waldsassen in Bavaria. They contain
carbonic acid; carbonates of lime and magnesia;
chlorides of calcium, magnesium, and aluminium;
carbonate of soda, and much oxide of iron. The
Bavarians consider them to resemble the waters
of Pyrmont.

WILD, Sax. *pilo*, (D.) *Wild*. An epithet
given to the countenance, when not in harmony
with the condition of the individual, and indica-
ting strong mental emotion; — a *wild look*, (F.)
Physiognomie égarée.

WILDBAD, MINERAL WATERS OF. Wild-
bad is a small town of Wurtemberg, three Ger-
man miles from Baden-Baden. It is much fre-
quented as a watering-place. The water is ther-
mal, from 95° to 100° of Fahr., and is much em-
ployed in cases in which thermal waters are indi-
cated. It is used internally, a pint not con-
taining more than a grain of solid ingredients,
which consists of salts of soda and lime. The
proportion of carbonic acid gas is very small.

WILDFIRE, *Erysipelas*.

WILDFIRE RASH, *Ignis sylvaticus*.

WILDUNGEN, MINERAL WATERS OF.
Wildungen is situate a few miles from Cassel in
Germany. In its neighbourhood are several mi-
neral springs; the chief of which, according to
Stucke, contain a bituminous matter; chloride
of sodium and sulphate of soda; carbonate and
sulphate of lime, &c. They are used, chiefly, as
refrigerants.

WILL, *Voluntas*.

WILLAN, *Salix*.

WILLOW, *Salix* — *w. Herb. Common*, *Lythrum*
salicaria — *w. Herb. great*, *Epilobium angustifo-*
lium — *w. Herb. purple*, *Lythrum salicaria* — *w.*
Herb. purple veined, *Epilobium coloratum*.

WILLOW, LOW-BUSH, *Salix humilis* — *w. Speck-*
led, *Salix humilis* — *w. Red*, *Cornus sericea* — *w.*
Rose, *Cornus sericea* — *w. Sweet*, *Myrica gale*.

WIND, Sax. *pinæ*, (an example of onomatopoeia — *ooid*) *Ventus*, An'emos, (F.) *Vent*. Winds
exert considerable influence on the animal econo-
my; acting by their temperature, which neces-
sarily modifies that of the circumambient air, as
well as by their moisture or dryness, and by the
emanations of different kinds, which they trans-
port to greater or less distances. The winds
which blow over a considerable extent of country,
are dry and serene: those which come from the
ocean are moist and chilly.

WIND, see Flatulence.

WIND OF A BALL. A term applied to the com-
pression of air, supposed to be produced by the
passage of a ball near a part of the body, so as
to occasion what has been called a *wind con-*
tusion.

WIND CONTUSION, see Wind of a Ball — *w.*
Dropsy, *Emphysema*, *Tympanites* — *w. Flower*,
Anemone — *w. Pipe*, *Trachea* — *w. Root*, *Asclepias*
tuberosa — *w. Wood*, *Asclepias tuberosa*.

WINDINESS, Flatulence.

WINDY, *Flatulentus*, *Flatulent*, *Vaporosus*,
(F.) *Venteux*, *Flatueux*. That which causes wind

or flatulence; also, that which is caused by wind
or flatulence: as *windy food*, *windy colic*, &c.
Also, one affected with flatulence, or who is
'troubled with the wind.'

WINE, Sax. *pin*, (G.) *Wein*, originally, per-
haps, from *ovos*, *Enos*, *Vinum*, *medu*, *Methu*, (F.)
Vin. A name given by chymists to all liquors
that have become spirituous by fermentation.
The term is generally, however, restricted to the
fermented juice of the grape, or of fruits, the pro-
duct of whose fermentation resembles, in almost
all respects, that of the juice of the grape. Wine
is composed of alcohol; saccharine matter; malic
acid; tartaric acid; bitartrate of potassa; acetic
acid; an extractive colouring matter, more or less
bitter, and partly resinous; and sometimes of an
aromatic substance. The extractive colouring
matter is chiefly met with in red wines. All these
constituents, except the alcohol, are found ready
formed in the grape. The alcohol proceeds from
the decomposition of the saccharine matter. A
part of the acetic acid is also formed during fer-
mentation.

The following table, drawn up by Mr. Brande,
exhibits the quantity of spirit in different kinds
of wine and liquors. It is proper, however, to
remark, that many of these wines are prepared
for the London market, and are more brandied, or
"reinforced," than the same varieties sold in the
United States. This is strikingly the case with
port. Dr. Henderson, too, has remarked, that
some of the wines analyzed by Mr. Brande were
mixed with a considerable quantity of adventi-
tious alcohol. Dr. Henderson's additions and
corrections have the letter H. affixed.

PROPORTION OF ALCOHOL, S. G. 0.825, IN ONE HUNDRED
PARTS BY MEASURE OF THE FOLLOWING WINES, AND MALT
AND SPIRITUOUS LIQUORS.

1. Lissa.....	26.47
Do.....	24.35
Average, (a)	25.41
2. Raisin Wine.....	26.40
Do.....	26.77
Do.....	23.20
Average,	25.12
3. Marsala.....	26.03
Do.....	25.05
Average, (b)	25.09
4. Port — average of six kinds.....	23.48
Do. — highest.....	25.83
Do. — lowest.....	21.40
5. Madeira.....	24.42
Do.....	23.93
Do. (Sercial).....	21.45
Do.....	19.24
Average,	22.27
6. Currant Wine.....	20.55
7. Sherry.....	19.51
Do.....	19.83
Do.....	18.79
Do.....	18.25
Average,	19.17
8. Teneriffe.....	19.79
9. Colares.....	19.75
10. Lacryma Christi.....	19.70
11. Constantia — white.....	19.75
Do. — red.....	18.92
Average, (c)	19.24
12. Lisbon.....	18.94
13. Malaga.....	18.94
14. Bucellas.....	18.49
15. Red Madeira.....	22.30
Do.....	18.40
Average,	20.35
16. Cape Muscat.....	18.25
17. Cape Madeira.....	22.94
Do.....	20.50
Do.....	18.11
Average,	20.61
18. Grape Wine.....	18.11
19. Calcevala.....	19.20
Do.....	18.10
Average,	18.65
20. Vidonia.....	19.25
21. Alba Flora.....	17.28

(a) 15.90 H. (b) 18.40 H. (c) 14.50 H.

22. Malaga.....	17-26
23. White Hermitage.....	17-43
24. Roussillon.....	19-00
Do.....	17-26
Average.....	18-13
25. Claret.....	17-11
Do.....	16-32
Do.....	14-08
Do.....	12-91
Average, (d).....	15-10
26. Malmsey Madeira.....	16-40
27. Lunel.....	15-52
28. Scheraz.....	15-52
29. Syracuse.....	15-28
30. Sauterne.....	14-22
31. Burgundy.....	16-60
Do.....	15-22
Do.....	14-53
Do.....	11-95
Average.....	14-57
32. Hock.....	14-37
Do.....	13-00
Do. (old in cask).....	8-88
Average.....	12-08
Rudesheimer (1811).....	H. 10-72
Do. (1800).....	H. 12-22
Average, H.....	11-47
Johannisberger.....	H. 8-71
33. Nier.....	14-63
34. Barsac.....	13-86
35. Tent.....	13-30
36. Champagne (still).....	13-80
Do. (sparkling).....	12-80
Do. (red).....	12-56
Do.....	11-30
Average.....	12-61
37. Red Hermitage.....	12-32
38. Vin de Grave.....	13-94
Do.....	12-80
Average.....	13-37
39. Frontignac.....	12-79
40. Côte Rôtie.....	12-32
41. Gooseberry Wine.....	11-84
42. Orange Wine—average of six samples made by a London manufacturer.....	11-26
43. Tokay.....	9-88
44. Elder Wine.....	9-87
45. Rhenish Wine.....	H. 8-71
46. Cider—highest average.....	9-87
Do. lowest.....	5-21
47. Perry—average of four samples.....	7-26
48. Mead.....	7-32
49. Ale (Burton).....	8-88
Do. (Edinburgh).....	6-20
Do. (Dorchester).....	5-66
Average.....	6-87
50. Brown Stout.....	6-80
51. London Porter—average.....	4-20
Do. Small Beer—average.....	1-28
52. Brandy.....	53-39
53. Rum.....	53-68
54. Gin.....	51-60
55. Scotch Whisky.....	54-32
56. Irish do.....	53-90

The only wine recommended in some pharmacopœias is *Sherry*, *Vinum*, *V. album Hispanicum*, *V. Xericum*, *Leuceænus*. It is a dry wine; the least variable in its properties, and agrees best with the dyspeptic. Other wines used officinally are—*Canary* or *Sack wine*, *Vinum Canarium*; *Mountain wine*, *Vinum album montanum*; *Tent wine*; *Rhenish wine*, *Vinum Rhenanum*; and *Port wine*, *Vinum rubrum Portugallicum*.

The wines habitually drunk are almost innumerable. Those that are sweet, or contain a large portion of free, saccharine matter, are decidedly the least wholesome to the dyspeptic.

When wine is good, and of a proper age, it is tonic and nutritive; when new, flatulent and cathartic, disagreeing with the stomach and bowels. In medicine, it is a valuable tonic in the last stage of typhus, when the skin is not too hot and dry. Its use, however, requires great caution; and when once commenced under proper circumstances, it cannot be suddenly dropped without mischief. It is, perhaps, the best permanent stimulus in the catalogue of the materia medica.

(d) 12-91 H.

Mulled Wine is made as follows:—Take of bruised cinnamon $\frac{3}{4}$; half a nutmeg grated; and ten bruised cloves. Infuse in boiling water Oss for an hour; strain and add sugar $\frac{3}{4}$. Pour the whole into a pint of hot Port or Sherry. Useful in adynamic conditions.

WINE, AMINÆAN, *Aminæum vinum*.

WINE OF ALOES, *Vinum aloes*.

WINE, ANTIMONIAL, *Vinum antimonii*—w. Antiscorbutic, *Vinum antiscorbuticum*—w. Aromatic, *Vinum aromaticum*—w. Bark, compound, *Vinum de kinâ kinâ compositum*—w. Barley, *Cerevisia*—w. Bitter, diuretic, *Vinum diureticum amarum*.

WINE BITTERS, *Vinum gentianæ compositum*. The formulæ for these may be various. The following is as good and as agreeable as any. (*Rad. gentian. lbj*, *cort. aurant. 3x*, *sem. cardam. cont. 3iv*, *cinnam. cort. 3iv*, *wine*, foreign or domestic, three gallons and a half.) Tonic and stomachic.

WINE, CEDAR, *Cedrinum vinum*—w. Champagne, *Vinum campanum*—w. Chian, *Chium vinum*.

WINE OF COLCHICUM, *Vinum colchici*—w. of Colchicum seed, *Vinum colchici seminis*—w. of Ergot, *Vinum ergotæ*—w. of Gentian, compound, *Vinum gentianæ compositum*—w. of Hellebore, white, *Vinum veratri*.

WINE, IPECACUANHA, *Vinum Ipecacuanhæ*.

WINE OF IRON, *Vinum ferri*.

WINE, MEDICATED, *Vinum medicinale*—w. Mulled, see Wine.

WINE OF OPIUM, *Vinum opii*.

WINE, PORT, *Vinum Portugallicum*—w. Pullet, *Poulet, vin de*.

WINE OF QUINIA, *Vinum quiniæ*.

WINE, RHUBARB, *Vinum rhei*—w. Sherry, *Vinum*—w. Steel, *Vinum ferri*—w. Tobacco, *Vinum tabaci*—w. Tar, see *Pinus sylvestris*.

WINE WHEY, *White Wine Whey*, (Prov.) *Sack-whey*. Take of good milk two-thirds of a pint, and add water to make a pint. Take of sherry, or any other good white wine, two glasses, and of sugar a dessert-spoonful. Place the milk and water in a deep pan on the fire; and the moment it boils, pour into it the wine and sugar. Stir assiduously for 12 or 15 minutes, whilst it boils. Lastly, strain through a sieve.

It is a good mode of giving wine in adynamic states.

WINE, WHITE, see Wine.

WINE OF WORMWOOD, *Vinum absinthites*.

WING-SEED, *Ptelea trifoliata*.

WINKERS, *Cilka*.

WINKING, *Scardamylgus*.

WINTER'S BARK TREE, *Wintera aromatica*.

WINTER BERRY, SMOOTH, *Prinos lævigatus*—w. Berry, Virginia, *Prinos*—w. Berry, whorled, *Prinos*.

WINTER BLOOM, *Hamamelis Virginiana*—w. Cough, Bronchitis, chronic—w. Green, *Gaultheria*—w. Green, round-leaved, *Pyrola*—w. Green, spotted, *Pyrola maculata*.

WINTERA, see *Wintera aromatica*.

WINTERA AROMATICA, *Drimys Winteri*, *Wintera'na aromatica*, *Winter's bark Tree*; Ord. Magnoliacæ. The bark, *Wintera* (Ph. U. S.), *Cortex Wintera'nis* seu *Winteranus Magellan'icus*, *C. Magellan'icus*, *Cinnamomum Magellan'icum*, (F.) *Écorce de Winter*, *Cannelle poirée*, is very much allied in its properties to *Canella alba*.

WINTERANA AROMATICA, *Wintera aromatica*.

WIRTSPRINGS, Hangnails.

WISBADEN, MINERAL WATERS OF. Wisbaden is a town of Germany, about 6 miles

north of Mainz, and 22 west of Frankfort. It has been long celebrated for its hot springs; and is much frequented. There are numerous cold and thermal springs,—the former containing sulphuric acid; the latter being saline and acidulous. The temperature of the hot springs varies from 117° to 160° Fahr.

WISMUTHUM, Bismuth.

WITCHEN, *Sorbus acuparia*.

WITCHHAZEL, *Hamamelis Virginiana*, *Sorbus acuparia*.

WITCHMEAL, *Lycopodium*.

WITCHWOOD, *Sorbus acuparia*.

WIT TEETH, see Dentition.

WITTER, Throat.

WIZEN, *Cæophagus*.

WOAD, *Isatis tinctoria*.

WOD, Insane.

WODNES, Insanity, Mania.

WOLFBANE, *Veratrum viride*.

WOLFSBANE, *Aconitum* — w. Wholesome, *Aconitum anthora*.

WOLFSCLAW, *Lycopodium*.

WOLFSJAW, see Harelip.

WOMAN, pl. women, *Gynæ*, (F.) *Femme*, Sax. *pirman*, 'wifeman,' 'woofman.' The adult female of the human race.

WOMANHOOD, *Mulieritas*.

WOMB, Uterus—w. Falling down of the, *Prolapsus uteri*.

WOMB GRAIN, *Ergot*.

WOMB, LACERATION OF THE, Uterus, rupture of the—w. Tympany of the, *Physometra*.

WONGE, Cheek.

WOOD, BRAZIL, *Cæsalpinia echinata* — w. Elk, *Andromeda arborea* — w. Pernambuco, *Cæsalpinia echinata*.

WOOD-LICE, *Onisci aselli*.

WOOD-MARCH, *Sanicle*.

WOOD, NICARAGUA, see *Cæsalpinia* — w. Peach, see *Cæsalpinia* — w. Pernambuco, *Cæsalpinia echinata* — w. Sappan, *Cæsalpinia sappan* — w. Sappan, *Cæsalpinia sappan*.

WOOD SORREL, *Oxalis acetosella*.

WOOD, SOUR, *Andromeda arborea* — w. Waxen, *Genista tinctoria*.

WOODWEX, *Genista tinctoria*.

WOODBINE, COMMON, *Lonicera periclymenum*.

WOODROOF, SWEET-SCENTED, *Asperula odorata*.

WOODS, SUDORIFIC, (F.) *Bois sudorifiques*. This term is applied, collectively, to guaiacum, sassafras, china, and sarsaparilla, which are often used together to form the sudorific decoction.

WOORALI, Curare.

WOORARA, Curare.

WORM, see Alembic, Vermiform process — w. Bark tree, *Geoffræa inermis* — w. Cakes, *Story's*, see Cakes, worm, *Story's* — w. Disease, *Helminthiasis* — w. Goosefoot, *Chenopodium anthelminticum* — w. Grass, perennial, *Spigelia Marilandica*.

WORM, GUINEA, *Dracunculus*.

WORM LOZENGES, CHING'S. Empirical preparations, which consist of yellow and brown lozenges. The former are directed to be taken in the evening; the latter on the following morning. The *Yellow Lozenges*. (*Saffron*, 3ss, water, Oj; boil, and strain; and add of white panacea of mercury, (calomel washed in spirit of wine,) lbj; white sugar, 28 lbs, mucilage of gum tragacanth q. s. to make a mass. Divide, so that each lo-

zenge may contain a grain of the panacea.) The *Brown Lozenges*. (*Panacea* 3vij, resin of jalap lbijss, white sugar lbij, mucilage of gum tragacanth q. s. Each lozenge to contain gr. ss of the panacea.)

WORM LOZENGES, SHERMAN'S, are said to be composed of calomel, gamboge, and sugar.

Peters's Worm Lozenges are said to be composed in a similar manner.

WORM ROOT, *Spigelia Marilandica* — w. Seed, *Artemisia antonica*, *Chenopodium anthelminticum* — w. Seed, goosefoot, *Chenopodium anthelminticum* — w. Weed, Corsican, *Corallina Corsicana*, *Polanisia graveolens* — w. Weed, white, *Corallina* — w. Wood, biennial, *Artemisia biennis* — w. Wood, Canada, *Artemisia canadensis* — w. Wood, common, *Artemisia absinthium* — w. Wood, creeping, *Artemisia rupestris* — w. Wood, lesser, *Artemisia Pontica* — w. Wood, Roman, *Artemisia Pontica* — w. Wood, sea, *Artemisia maritima* — w. Wood, silky, *Artemisia glacialis*.

WORMIAN Ossa seu OSSIC'ULA, O. *epact'lia* seu *raphogeminant'ia* seu *trig'uetra* seu *triangular'ia* *Blasii* seu *sutura'rum*, *Claves calvar'ia*, Wormian bones, (F.) *Oss Wormiens* ou *épactaux* ou *surnuméraires* (Ch.), ou *intercalés* ou *triangulaires* ou *Clefs du crâne*. Small bones, which are sometimes observable in the sutures of the bones of the cranium, of which they form a part. They were called *Wormiana*, from Olaus Wormius, who is said to have first described them. They had been previously, however, mentioned by G. Andernach, a physician at Strasburg. The Ossa Wormiana exist more commonly in the sutures of the vault of the skull; especially in the lambdoidal, sagittal, and squamous. They are uncommon at the base of the skull. Their size is very variable, and shape irregular. Their structure and development are similar to those of the other bones of the cranium; and, like them, they are bounded by sutures.

WORMIT, *Artemisia absinthium*.

WORMS, originally from (L.) *Vermes*; (F.) *Vers*; *Entozo'a*, *Entozo'o'ria*, *Enterozo'o'a*, *Endozo'o'a*, *Entelmin't'a*, *Enthelmin'thes*, *Helmin'thi*, *Entoparasites*, (F.) *Entozoaires*. Animals whose common character is that of existing only in other animals; hence their name *entozoa*, from *entos*, 'within,' and *zoo*, 'animal.' They are met with, not only in the natural cavities, but even in the tissue of the organs, but chiefly in the intestines — *Vermes intest'i'ni*, *Helmin'thia alvi* seu *pod'icis*, *Splanchnelmin't'ia*, *Intes'tinal worms*, (F.) *Vers intestinaux*, *Angibromelmin'thie*. Piorry, Cuvier, Rudolphi, Brera, Bremser, Laënnec, and others, have endeavoured to classify them, — some according to their shape; others, according to their anatomical characters.

The following table exhibits most of the entozoa which have been met with in the human body, and their usual habitats: —

Entozoa.	Where found.
<i>Trichoceph'alus dispar</i> , <i>Oxyu'ris vermicula'ris</i> , <i>As'caris ala'ta</i> , <i>As'caris lumbricoi'des</i> , <i>Ancylos'toma duodenale</i> , <i>Bothrioceph'alus latus</i> , <i>Tæ'nia so'lium</i> , <i>Pentas'toma constrictum</i> , <i>Tæ'nia nana</i> ,	Intestines.
<i>Distoma hæmato'bium</i> ,	
<i>Ditrachyo'eras rudis</i> , (?)	Intestines and Venæ porta.
<i>Diplosoma crena'ta</i> , <i>Spiro'tera hom'inis</i> , <i>Dactyl'ius aculea'tus</i> ,	
	Urinary bladder.

Dis'torta her'aticum,	Gall bladder.
Dis'toma lanceola'tum,	Hepatic duct.
Stron'gylus gigas,	} Kidney.
Tetras'toma rena'lē,	
Fila'ria oc'uli,	Eye.
Monos'toma lentis,	Crystalline.
Distoma oc'uli huma'ni,	Capsule of crystal-
Acephalocyst'is endog'ena,	line.
Echinococ'cus hom'inis,	Liver.
Polys'toma pinguic'ola,	} Liver, Spleen, and
Hexathyrid'ium pinguic'ola,	
Polys'toma vena'rum seu	Omentum.
sanguic'ola,	Ovary.
Hexathyrid'ium vena'rum,	} Veins.
Fila'ria bronchia'lis,	
Trichi'na spira'lis,	} Bronchial glands.
Cysticer'cus cellulō'sæ,	
Acephalocyst'is multifi'da,	Muscles.
Fila'ria medinen'sis,	Brain.
	Areolar texture.

The most common of these are:—the *Oxyuris vermicularis*, *Ascaris lumbricoides*, and *Tænia*, which are found in the intestines. The origin of these worms is extremely singular, and more favourable than any other fact to the hypothesis of spontaneous generation in the lowest tribes of animated nature. They are certainly not identical with any worms out of the body. They are most commonly met with in children improperly fed; but their presence cannot be positively detected by any symptoms, except that of the discharge of the worms themselves in the evacuations. They may, of course, give rise to every kind of irritation in the intestinal tube, and to sympathetic disturbance in most of the functions; but all these signs may be produced by intestinal irritation resulting from other causes.

Anthelmintics or *vermifuges*, employed, as the name imports, for their removal, may be of two kinds—*mechanical* and *true*. To the first class—the *mechanical*—belong *Emetics*, *Purgatives*, *Mucuna*, *powdered Tin*, &c. To the latter—the *true anthelmintics*,—*Turpentine*, *Chenopodium anthelminticum*, *Pink-root*, *Semina Santonici*, &c. Of these, the best is turpentine. See *Oleum terebinthinæ*. The great object, however, is to prevent their generation. This must be done by generous diet, fresh air, and tonics, where necessary.

WORMS. *Aene punctata*.

WORSUM, Pus.

WORT, Sax. *wýrt*, *Mustum*, *Infusum Bynea* seu *Bra'sii* seu *Malti*, (F.) *Mout de la Bière*. An infusion of salt. This has been recommended in scurvy. One measure of ground malt is added to three equal measures of boiling water. The mixture must be well stirred, and left to stand covered three or four hours. Dose, one to four pints, daily. It has, also, been recommended in other cases, where a strong putrescent disposition

appeared to prevail in the fluids, as in cancerous and phagedenic ulcers. It is not now employed

WORTWALE, Hangnail.

WOKEN, Trachea.

WOUND, Sax. *wund*, *Vulnus*, *Volnus*, (diminutive *Vulnusculum*), *Plaga*, *Trauma*, *Troma*, *Trenis*, *Treis* *Vulnus*, *Blabē*, *Tynnma*, (F.) *Plaie*, *Blessure*. A solution of continuity in the soft parts, produced by some mechanical agent. Wounds present innumerable differences, as regards their situation; the parts interested; their direction; size; shape; the nature of the instrument or agent by which they are produced; their more or less simple or complex character, duration, &c. A wound is called *incised*, (F.) *Coupure*, *Incision*, when made by a cutting instrument;—*punctured*, *Nygma*, (F.) *Piqure*, when made by a pointed instrument;—*lacerated*, (F.) *Déchirure*, *Plaie par arrachement*, when the parts are lacerated or torn by the wounded body; *poisoned*, *Cellulitis venenata*, *Neurisia*, (F.) *Plaie envenimée*, when some virulent or venomous substance has been introduced;—and *contused*, (F.) *Plaie contuse*, when produced by a blunt body. *Gunshot wounds*, *Sclopetopla'gæ*, *Vul'nera sclopetaria* seu *à sclopet'is*, (F.) *Plates d'armes à feu* ou *d'arquebuse* ou *d'arquebuse*, *Coup de feu*, belong to the last division.

WOURALI, Curare.

WOURARI, Curare.

WOURARU, Curare.

WOURSUM, Pus.

WRACK, SEA, *Fucus vesiculosus*—*w*. Bladder, yellow, *Fucus vesiculosus*.

WRAMP, Sprain.

WRAPPING UP, (of the hydropathists,) *Emballottage*.

WRATAACK, Nanus.

WRENCH, Sprain.

WRET, Verruca.

WRETWEED, *Euphorbia palustris*.

WRIGHTIA ANTIDYSENTERICA, *Nerium antidysentericum*.

WRINKLE, Sax. *wrincle*, *Ruga*, *Rytis*, *Rhacus*, *Rhachus*, *Rhagus*, *Rhyssē'ma*, *Corrugatio cutis*, *Ruelles*, (Sc.) *Runkle*, *Runkill*, (Prov.) *Crinkle-crackle*, (F.) *Ride*. A furrow or fold in the skin of the face, forehead, vagina, &c. See *Corrugation*.

WRINKLED, *Rugo'sus*, *Rugous*, *Rugose*, (F.) *Ridé*. Full of wrinkles.

WRIST, Carpus.

WRIST-DROP. Paralysis of the muscles of the forearm, induced by the poison of lead.

WRITERS' SPASM, see *Spasm*, *writers'*.

WURALI, Curare.

WURSTGIFT, Allantotoxicum.

WRONG HEIR, *Canellus*.

WRYNECK, *Torticollis*.

X.

X LEG, Entogonyancon.

XALAPPA, Convolvulus jalapa.

XANTHIC OXIDE, Uric oxide.

XANTHINE, Uric oxide.

XANTHIUM, *X. Strumarium* seu *Orientalis* seu *Vulgaris*, *Barda'na* minor, *Lappa* minor, *Lesser Burdock*, *Burweed*, *Burthistle*, *Clobur*, *Cockle-hur*, (F.) *Lampourde*, *Petit Glouteron*, *Petite Bardane*, from *ξανθος*, 'yellow,' because of the colour the plant is said to yield. This herb, as one of its names imports, was once esteemed for the cure of scrofula. The seeds have been administered in cutaneous affections.

XANTHIURIA, *Xanthuria*.

XANTHOP'SIA, from *ξανθος*, 'yellow,' and *ψις*, 'vision.' Yellow vision, — as sometimes occurs in jaundice.

XANTHORRHIZA, (*ξανθος*, 'yellow,' and *ρίζα*, 'root.') *Xanthorrhiza apiifolia*.

XANTHORRHIZA APIIFOLIA, *Xanthorrhiza*, *X. simplicissima* seu *Marbo'ica*, *Zanthorrhiza tinctoria*, *Yellow Root*, *Parsley-leaved Yellow Root*, *Yellow wort*, *Shrub yellow root*. The root of this American plant—*Xanthorrhiza*, (Ph. U. S.) — is a strong and pleasant bitter; and, in the dose of two scruples, sits easy upon the stomach.

XANTHORRHOEA, *Grass Tree*. A genus of trees in Australia, *Nat. Order*, *Liliaceae*, of which there are several species. Two resins are obtained from them; — one the *yellow resin* of *Xanthorrhoea* or of *New Holland*, *Resin lu'tea* *Novi Belgii*, *Botany Bay resin*, *Ac'aroid resin* or *gum*, *resin'na* vel *gummi acaroides*, which probably resembles *tolu* and *storax* in medicinal properties; — the other the *red resin* of *Xanthorrhoea*, *black-bay gum*.

XANTHOS, Yellow.

XANTHOSIS. A term applied to the yellow discolouration often observed in cancerous tumours — especially in encephaloid of the testicle. — Lebert.

XANTHOXYLUM FRAXIN'EUM, (*ξανθος*, 'yellow,' and *ξύλον*, 'wood.') *X. americanum*, *X. fraxinifolium*, *X. ramiflorum*, *X. tricar'pum*, *Prickly ash*, *Shrubby prickly ash*, *Toothache bush*, *Toothache tree*, *Pellitory*, *Yellow wood*, *Suterry*. *Ord.* *Xanthoxylaceae*. A spirituous infusion of the bark — *Xanthoxylum* — Ph. U. S. — has been highly recommended in cases of violent colic. The fresh juice, expressed from the root of *xanthoxylum*, is said to afford relief in the dry bellyache of the West Indies. A decoction of the bark is sometimes used as a wash to foul ulcers.

XANTHOXYLUM CAROLIA'NUM has similar properties; but is more acrid.

XANTHOXYLUM CLAVA HERCULIS of the West Indies, (F.) *Clavaler à feuilles de frêne*, *Frêne épineux* is a very strong stimulant, and powerful sialogogue: and has been given internally in rheumatism.

XANTHOXYLUM OCTANDRA, *Fagara octandra*.

XANTHURIA, *Xanthuria*, from *ξανθος*, 'yellow,' and *ουρον*, 'urine.' A condition of the system and of the urine, in which xanthic oxide is deposited from the urine.

XANTICA, Carminatives.

XANTOLINA, *Artemisia santonica*.

XENODOCEUM, Hospital.

XENODOCEUM, Hospital.

XERANSIS, Arefaction, Drying.

XERANTIA, Siccantia.

XERA'SIA. Same etymon; from *ξηρος*, 'dry.' A disease of the hairs, which become dry, cease to grow, and resemble down covered with dust. — Galen.

XERION, Catapasma.

XERCOLLYRIUM, from *ξηρος*, 'dry,' and *κόλλυριον*, 'a collyrium,' *Collyrium siccum*. A dry collyrium.

XERODER'MA, from *ξηρος*, 'dry,' and *δερμα*, 'skin.' Diminution of secretion of the sebaceous glands.

XEROMA, Xerophthalmia — x. Conjunctival. Xerophthalmia — x. Lachrymal. Xerophthalmia.

XEROMA, LACHRYMAL. Suppression of the lachrymal secretion. See Xerophthalmia.

XEROMYCTER, Xeromycteria.

XEROMYCTERIA, *Xeromycter*, from *ξηρος*, 'dry,' and *μύκτηρ*, 'the nose.' Dryness of the nose.

XEROMYRON, *Unguentum siccum*; from *ξηρος*, 'dry,' and *μυρον*, 'an ointment.' A dry ointment. — Gorræus.

XEROPHAGIA, from *ξηρος*, 'dry,' and *φαγω*, 'I eat.' Excessive use of dry aliment. A kind of regimen to which the ancient athlete subjected themselves, with the view of increasing their strength.

XEROPHTHAL'MIA, *Ophthalmia sicca*, *Ophthalmoxerosis*, *Xero'ma*, from *ξηρος*, 'dry,' and *ὀφθαλμα*, 'inflammation of the eye.' An inflammation of the eye, without discharge. The term — as well as *Conjunctival* and *Lachrymal Xero'ma*, and *Cuticular Conjunctiva* — has been given to cases in which the conjunctiva is so changed, that it presents more of the characters of skin than of mucous membrane. It is incapable of radical cure. Also, *Lippitudo*.

XEROTRIBIA, see Friction.

XEROTRIPSIS, Friction, dry.

XIPHISTERNUM, Xiphoid.

XIPHODES, Xiphoid.

XIPHODYMUS, Thoraco-gastrodidymus.

XIPHOID, *Xiphoi'des*, *Xipho'des Ensiformis*, from *ξίφος*, 'a sword,' and *εἶδος*, 'shape;' *Cartilago Xiphoides* seu *Xiphoides* seu *Scutalis* seu *Ensiformis* seu *Pelta'lis* seu *Pelta'us*, *Furcella inferior*, *Mucronatum os*, *Mucronata Cartilago*, *Chondros*, *Processus xiphosternalis* seu *xiphoides* seu *sterni xiphoides*, *Xiphisternum*, *Ensiform cartilage*, (F.) *Appendix ou cartilage xiphoides*, *C. Mucroné*, *Appendice sous-sternale*, *Palette*. The name of the appendix which terminates the sternum beneath; and which is so called from some resemblance to a sword.

XIPHOID LIG'AMENT, *Costo-xiphoid L.* (F.) *Ligament xiphoidien ou costo-xiphoidien*, is a small, very thin, ligamentous fasciculus, which passes from the cartilage of prolongation of the 7th rib to the anterior surface of the xiphoid cartilage, into which it is inserted by its fibres decussating with those of the opposite ligament.

XIPHOPAGES, *Didymus Symphyo-epigastricus*; from *ξίφος*, 'a sword,' and *πνυμμι*, 'I fix.' A monstrosity, in which twins are united by the epigastrium, as in the case of the Siamese twins.

XYLO-ALOES, Agallochum — x. Balsamum. see Amyris opobalsamum — x. Cassia, Laurus cassia — x. Cinnamomum, Laurus cinnamomum.

XYLUM, Gossipium.

XYMPATHIA, Sympathy.

XYRIS, Iris foetidissima.

XYRIS BULBO'SA, X. *Indica* seu *Caroliniana*, *Yellow-eyed Grass*; indigenous; Order, Xyridaceæ; flowering from July to September. The roots and leaves are said to be used by the Hin-

doos in lepra, and chronic cutaneous diseases in general.

XYRUM, Novacula.

XYRUS, Novacula.

XYSMA, Linteum, Rasura, Scobs.

XYSMOS, Rasura.

XYSTER, Raspatorium.

XYSTOS, Linteum.

XYSTRUM, Raspatorium — x. Ophthalmicum, Ophthalmoxyster.

Y.

YAM. This esculent root is principally obtained from three species of *Dioscorea*, — *alata bulbifera*, and *sativa*. They grow spontaneously in both the Indies, and their roots are eaten promiscuously, as the potato is with us. Their taste is somewhat like that of the potato.

YAM ROOT, WILD, *Dioscorea villosa*.

YARD, Penis.

YARROW, COMMON, *Achillea millefolium*.

YAUPON, *Ilex vomitoria*.

YAVA SKIN, Elephantiasis Arabum.

YAWNING, from Sax. *geonan*, Chasmē, Chasmus, *Hia'tus*, *Oscē'do*, *Oscita'tio*, *Clonus pandicula'tio maxilla'rum*, Gaping, (F.) *Baillement*. Yawning consists of a deep inspiration, with considerable separation of the jaws, executed slowly and in an involuntary manner, and followed by a prolonged and more or less sonorous expiration. This respiratory movement is preceded by a feeling of oppression in the epigastric region, and in the muscles of respiration and mastication. Yawning has been conceived to be owing to torpor in the pulmonary circulation; the causes producing it being commonly ennui, fatigue, sleepiness, monotonous sounds, hunger, sympathy, &c. Yawning often precedes the invasion of certain intermittents, attacks of hysteria, &c., and it may, by its frequency, and obstinacy, become a true disease.

YAW ROOT, *Stillingia*.

YAWS, *Framboesia*.

YEAST, Yest — y. Plant, *Torula cerevisiæ*.

YÈBLE, *Sambucus ebulus*.

YELK, see Ovum.

YELLOW, *Flavus*, *Lu'teus*, *Xanthus*, from past participle of Sax. *geælan*, 'to burn.' One of the seven primitive colours.

YELLOW ASH, *Cladrastis tinctoria* — y. Berry, *Podophyllum montanum* — y. Fever, see Fever, yellow — y. Fever, mild, see Relapse — y. Jack, Fever, yellow.

YELLOW LIG'AMENTS, (F.) *Ligaments jaunes*, occupy the spaces between the vertebral plates, from the interval, which separates the second vertebra from the third, to that which separates the last vertebra from the sacrum. They are so called in consequence of their colour.

YELLOW PAINT, *Hydrastis Canadensis* — y. Root, *Jeffersonia Bartoni* — y. Root, parsley-leaved, *Xanthorrhiza apiifolia* — y. Root, shrub, *Xanthorrhiza apiifolia*.

YELLOW SPRING, see Ohio, Mineral waters of.

YELLOW SPRINGS. These are in Chester county, Pennsylvania. The temperature of the chief spring is 53° Fahr. The water contains no impregnation.

YELLOW STAR, *Helenium autumnale* — y. Weed, *Ranunculus acris* — y. Wood, *Xanthoxylum fraxineum* — y. Wort, Parsley-leaved, *Xanthorrhiza apiifolia*.

YELLOWWS, *Cypripedium luteum*, *Icterus*, *Ranunculus acris*.

YERBA DEL SOLDADA, Matico — y. *de Huaca*, *Datura sanguinea*.

YEST, Yeast, Sax. *ȝerc*, from *ȝerc*, 'a storm,' *ȝercis*, 'stormy;' *Cerevisiæ Fermentum*, *Spuma*, *Cerevisiæ*, *Barm*, (F.) *Levure*, *Levure de la Bière*. The scum which collects on the surface of beer whilst fermenting. Yest is antiseptic and tonic. It has been much extolled in putrid fevers, — a table-spoonful being given as a dose, in porter, wine, &c. It is, also, applied in the form of cataplasm, to foul ulcers. In all these cases, it acts by virtue of the generation of carbonic acid gas.

YEUX ARDENS, Ardent eyes — y. *d'Écrevisse*, *Cancerorum chela*.

YEW TREE, *Taxus baccata* — y. American, *Taxus Canadensis*.

YLECH, *Ilech*.

YLIATER, *Ilech*.

YOKE, *Singultus*.

YOLK, see Ovum, and *Vesicula umbilicalis* — y. Membrane, see *Vesicula umbilicalis*.

YORK SPRINGS. These are in Adams county, Pennsylvania. One of them is a saline, containing sulphates of magnesia and lime, and chloride of sodium, and is a mild cathartic and diuretic; — the other is a strong chalybeate.

YOUK, Itching, *Psora*.

YOUTH, Adolescence.

YOPON, *Ilex vomitoria*.

YPSELOGLOSSUS, *Basiloglossus*.

YRIDES, Orpiment.

YSAMB'RA, *Pedigamb'ra*. An ancient Spanish poison, of which hellebore formed the greatest part.

YUCCA, *Jatropha manihot*.

YUCCA FILAMENTO'SA, *Bear's grass*, *Silk grass*; indigenous. A tincture of the root has been used in gonorrhœa.

YUKE, Itching, *Psora*.

Y-WER-A, see Spirit.

Z.

ZACCHARUM, Saccharum.

ZAEL, Borosail.

ZAFFRAN, Crocus.

ZAFFRE, Smalt.

ZAMBO, see Mulatto.

ZAMIA INTEGRIFOLIA, see Arrowroot—*z.*
Pumila, see Arrowroot.

ZANNA. A kind of Armenian bole, used by the ancients as a desiccative. It was also called *Zarnacha*.

ZANTHORRIZA APIIFOLIA, *Xanthorrhiza apiifolia*.

ZANTHOXYLUM, *Xanthoxylum*.

ZAPANIA NODIFLORA, *Anacoluppa*.

ZARNACHA, *Zanna*.

ZARSA, *Smilax sarsaparilla*.

ZARZAPARILLA, *Smilax sarsaparilla*.

ZEA MAYS, *Z. vulgare*, *Mays*, *M. zea* seu *Americana*, *Ador*, *Indian Corn*, *Corn*, *Turkey* or *Indian Wheat*, *Maize*, *Mayz*, (F.) *Mais*, *Graine de Turquie*, *Blé d'Inde* ou *d'Espagne* ou *d'Italie* ou *de Turquie*. *Ord. Graminæ*. A native of America, and much used over the whole of the continent as an article of diet. It is ground for the same purposes as wheat, and forms an excellent bread.

ζea or *ζεα*, was the ancient name of a sort of corn.

ZÉDOAIRE ROND, *Kæmpferia rotunda*.

ZEDOARIA, *Kæmpferia rotunda*—*z. Longa*, see *Kæmpferia*—*z. Rotunda*, see *Kæmpferia*.

ZEDOARY, *Kæmpferia rotunda*.

ZELOSIS, *Zelotypia*.

ZELOTYP'IA, from *ζηλος*, 'love,' and *τυπος*, 'form,' 'model,' *Zelo'eis*. It has been proposed to apply this name to the violent mental affection produced by the acknowledged or presumed infidelity of a beloved object; and to the consequences of such affection;—melancholy, mania, &c.

ZEMA, Decoction.

ZEN'ICON, *ζενικον*. Ancient name of a Celtic poison, used for poisoning arrows.

ZEOCRITON DISTICHUM, *Hordeum distichon*.

ZEPHACH, *Peritonæum*.

ZERMA, *Zerna*.

ZERNA, *Zerna*. Amongst the barbarous Latin writers it signified ulcerated impetigo, *Impetigo ulcerata*.

ZERUMBET, *Cassumuniar*.

ZESIS, Effervescence, Decoction.

ZEST, see *Citrus medica*.

ZESTOLUSIA, Bath, hot.

ZIBEBÆ, see *Vitis vinifera*.

ZIBETHUM, *Civetta*.

ZIGNITES, *Zingrites*.

ZINC, *Zincum*, *Zinkum*, *Speltrum*, *Speanter*, *Tutenag*, *Zinchum*, *Zint*, *Stannum Indicum*. Perhaps formed etymologically from (G.) *Zin*, 'Tin.' A metal obtained from calamine and blende. Its colour is bluish-white; lustre of the fresh surface considerable, but soon dulled, from the facility with which it oxidates; hard; texture striated. S. g. 7.190. Melts at 700° Fah., burns with a bright flame in a higher temperature, and is vo-

latilized in the form of a white, flocculent oxyd. Its use, in medicine, is in the formation of the following preparations.

ZINC, ACÉTATE DE, *Zinci acetat*—*a.* But-ter of, *Zinci chloridum*—*z.* Chloride of, *Zinci chloridum*—*z.* *Chlorure de*, *Zinci chloridum*—*z.* *Cyanure de*, *Zinci cyanuretum*—*z.* *Ferrohydrocyanate of*, *Zinci cyanuretum*—*z.* *Flowers of*, *Zinci oxydum*—*z.* *Hydrocyanate of*, *Zinci cyanuretum*—*z.* *Iodide of*, *Zinci Iodidum*—*z.* *Ioduret of*, *Zinci Iodidum*.

ZINC AND MORPH'IA, DOUBLE I'ODIDE OF, (F.) *Iodure double de Zinc et de Morphine*. This salt is prepared by boiling iodide of iodhydrate of morphia with water and zinc, after they have acted on each other for some days; filtering and crystallizing. It is considered to combine the properties of morphia and the salts of zinc, and has been prescribed as an anodyne and antispasmodic. Dose, a quarter of a grain.

ZINC, OXIDE DE, *Zinci oxydum*—*z.* *Oxyd of*, *Zinci oxydum*.

ZINC AND STRYCH'NIA, DOUBLE I'ODIDE OF. This salt is prepared by digesting for several days in hot water, iodide of iodhydrate of strychnia and zinc; filtering and crystallizing. It has been proposed in the same cases as strychnia.

ZINC, SULFATE DE, *Zinci sulphas*—*z.* Sulphate of, *Zinci sulphas*—*z.* *Valerianate of*, *Zinci Valerianas*.

ZINCHUM, *Zincum*.

ZINCI ACÉ'TAS, *Ac'etate of Zinc*, (F.) *Acétate de Zinc*. (*Plumbi acet. lbj, zinc. granul. ℥ix, aquæ destillat. Oij.*) Dissolve the acetate of lead in the water, and filter. Add the zinc to the solution, and agitate them occasionally together, in a stopped bottle, for five or six hours, or until the liquor yields no precipitate with a solution of iodide of potassium. Filter the liquor, evaporate it with a moderate heat to one-fifth, and set it aside to crystallize. Pour off the liquid, and dry the crystals on bibulous paper.

Should the crystals be coloured, dissolve them in distilled water, and having heated the solution, drop into it, while hot, a filtered solution of chlorinated lime, until it ceases to let fall sesquioxide of iron; filter the liquor, acidulate it with a few drops of acetic acid, evaporate, and crystallize.—*Ph. U. S.*)

Frequently used to form a gently astringent wash in ophthalmia and other local inflammations; and as an injection in gonorrhœa.

ZINCI CARBONAS IMPURUS, *Calamina*.

ZINCI CARBONAS PRÆCIPITATUS, *Precipitated carbonate of Zinc*. (*Zinci sulphat., sodæ carbonat. āā lbj; aq. bullient. cong.*) Dissolve the sulphate of zinc and carbonate of soda respectively in Oiv of water. Mix the solutions; stir, and set the mixture by, that the powder may subside. Having poured off the liquid, wash the precipitated carbonate of zinc with hot water, until the fluid of the washings is nearly tasteless, and dry with a gentle heat.—*Ph. U. S.*) Used to prepare the *Zinci oxidum*.

ZINCI CARBONAS PRÆPARATUS, *Calamina præparata*—*z.* *Chloridi liquor*, see *Burnett's Disinfecting Liquid*.

ZINCI CHLORIDUM, *Z. chloruretum*, *Zincum chloratum seu muriaticum (oxydatum)*, *Chloride*

or Chloruret of Zinc, (F.) *Chlorure de Zinc*. (Zinc. in frustulis, ʒijss; acid. nitric., creta preparat. āā ʒi; acid. muriat. q. s. To the zinc, in the porcelain vessel, add gradually sufficient muriatic acid to dissolve it; strain, add the nitric acid, and evaporate to dryness. Dissolve the dry mass in water, add the chalk, allow the mixture to stand for 24 hours, filter, and again evaporate to dryness.—Ph. U. S.) It is very deliquescent in the air, forming the *Butter of Zinc*, *Butyrum Zinci*.

Chloride of Zinc is a powerful escharotic, and has been lately employed as an effective antiseptic.

ZINCI CHLORURETUM, *Z. chloridum*.

ZINCI CYANURETUM, *Z. cyan'idum* seu *Hydrocyanas* seu *Prussias*, *Zincum Cyanogenatum* seu *Borus'sicum* seu *Zool'i'cum*, *Hydrocyanate of Zinc*, (F.) *Cyanure de Zinc*. This salt is formed by adding sulphate of zinc to hydrocyanate of potassa so long as any precipitate is thrown down. When dried and calcined at a low heat, the product is a mixture of cyanuret of zinc and of potassium. It has been recommended in nervous cardiacgia. Dose, from one-sixteenth to one-twelfth of a grain.

It is chiefly used as a caustic; one part of the chloride being mixed with four, three, or two parts of flour, according to the desired strength, and a few drops of water being added to form a paste.

ZINCI FERROHYDROCYANAS, *Zinci ferrocyanuretum*, *Cyanuretum Ferrozin'icium*, *Zincum Ferrohocydran'icium*, *Ferrohydrocyanate of Zinc*. Made by the mutual decomposition of boiling-hot solutions of sulphate of zinc and ferrocyanate of potassa. It has been recommended in the same cases as the last preparation.

ZINCI HYDROCYANAS, *Zinci cyanuretum*.

ZINCI IODIDUM, *Z. Ioduretum*, *Zincum Ioda'tum*, *Iodetum Zinci* seu *Zin'icium*, *Protoioduretum Zinci*, *Iodide or Iod'uret of Zinc*, (F.) *Iodure de Zinc*. This salt is made by boiling together iodine and zinc to dryness, and subliming. An ointment, composed of a drachm to an ounce of lard, has been used in cases where the external use of iodide of potassium is indicated.

ZINCI IODURETUM, *Z. iodidum* — *z. Nix*, *Zinci oxydum*.

ZINCI OXYDUM, *Zincum calcinata'tum*, *Nihil Album*, *Lana Philosoph'rum* seu *Philosoph'ica*, *Nix Zinci*, *Nihil Gris'eum*, *Pom'pholyz*, *Oxyd of Zinc*, *Flowers of Zinc*, (F.) *Oxide ou Blanc de Zinc*. (Procured by burning zinc in a long, deep crucible, placed so as to collect the sublimate. Also, by exposing the precipitate thrown down from a solution of sulphate of zinc by carbonate of ammonia to a strong heat, so as to drive off the carbonic acid.—Ph. U. S.) It is tonic and antispasmodic; externally, detergent and desiccative. Employed in epilepsy, chorea, &c. For its external use, see Ung. Zinci.

ZINCI OXYDUM IMPURUM, *Tutia* — *z. Prussias*, *Zinci cyanuretum*.

ZINCI SULPHAS, *Sulphate of Zinc*, *Zincum Vitriola'tum*, *Vitriolum Album* seu *Goslarien'se* seu *Zinci*, *Sulphas Zin'icium*, *Gilla Vitrioli* seu *Theophrasti*, *Sal Vitrioli* seu *Vomit'orium Vitrioli*, *Chalcant'rum Album*, *White Vitriol*, *White Cop'eras*, (F.) *Sulfate de Zinc*. (Zinc. in frustul. ʒiv; acid. sulphuric. ʒvj; aquæ destillat. Oiv. To the zinc and water, previously introduced into a glass vessel, add by degrees the sulphuric acid, and, when the effervescence shall have ceased, filter the solution through paper; then boil it down till a pellicle begins to form, and set aside to crystallize.—Ph. U. S.) The sulphate of zinc is odororous; taste, styptic. It is in white, semi-

transparent, efflorescent crystals; soluble in three parts of water at 60°. It is emetic, tonic, antispasmodic, and, externally, astringent. A nostrum, sold under the name *Antipertus'sis*, and, as its title imports, given in whooping-cough, contains this salt as its chief ingredient. It is emetic. Dose, as an emetic, gr. x to ʒss; as a tonic, gr. j to ij. Used externally as a collyrium, gr. j. to water f ʒj.

ZINCI VALERIANAS, *Zincum valerian'icum*, *Valer'ianate of Zinc*. A salt formed by saturating valerianic acid with freshly prepared carbonate of zinc. It is in pure white, pearly scales, and has been recommended as a tonic in neuralgic diseases, epilepsy, &c. The dose is one or two grains several times a day. It is officinal in the Dublin Pharmacopœia.

ZINCUM, *Zinc* — *z. Borussicum*, *Zinci cyanuretum* — *z. Calcinatum*, *Zinci oxydum* — *z. Chloratum*, *Zinci chloridum* — *z. Cyanogenatum*, *Zinci cyanuretum* — *z. Ferrohocydranicum*, *Zinci ferrohocydranas* — *z. Iodatium*, *Zinci iodidum* — *z. Muriaticum*, *Zinci chloridum* — *z. Valerianicum*, *Zinci valerianas* — *z. Vitriolatium*, *Zinci sulphas* — *z. Zooticum*, *Zinci cyanuretum*.

ZINGIBER, *Amomum zingiber* — *z. Cassumuniar*, see *Cassumuniar* — *z. Cliffordia*, see *Cassumuniar* — *z. Germanicum*, *Arum maculatum* — *z. Officinale*, *Amomum zingiber* — *z. Purpureum*, see *Cassumuniar* — *z. Spurius*, see *Cassumuniar* — *z. Zerumbet*, see *Cassumuniar*.

ZINGIBERIN, *Piperoid*.

ZINGRITES, *Zigni'tes*. A stone, resembling glass; to which the ancients attributed marvelous properties.

ZINKUM, *Zincum*.

ZINT, *Zincum*.

ZINZIBER, *Amomum zingiber*.

ZIPHAC, *Peritonæum*.

ZIRBUS, *Epiploon* — *z. Adipinus*, *Epiploon*, *gastro-colic*.

ZIZANIA AQUATICA, *Wild Rice*; *Ord. Graminæ*; called by the Indians, *Menomene*, grows abundantly on the marshy margins of the northern lakes and waters of the upper branches of the Mississippi. The grain resembles oats. It furnishes the northern savages and the Canadian traders and hunters with their annual supplies of grain. *Zizani'ov* was the ancient name of some wild grain.

ZIZYPHUM, *Jujube*.

ZIZYPHUS JUJUBA, see *Jujube* — *z. Nitida*, see *Jujube*.

ZIZYPHUS CENOP'RIA, *Z. Napeca*; *Ord. Rhamnaceæ*; common all over India. The fruit is eaten by the natives. It is of a pleasant acid taste. A decoction of the bark of the fresh root is used to promote the healing of fresh wounds.

ZIZYPHUS SATIVA, see *Jujube* — *z. Sylvestris*, see *Jujube*.

ZOA'GRRIA, from *ζωη*, 'life,' and *αγρω*, 'I catch up.' The saving and preservation of life. *Zoësis*. A reward for preserving life; *Sostrium*. A votive offering hung up in the temples for a life saved.

ZOANTHRO'PIA, *Melanchol'ia zoanthro'pia*, from *ζωον*, 'an animal,' and *ανθρωπος*, 'a man.' A species of monomania, in which the patient believes himself transformed into an animal. *Lycanthropia* and *Cynanthropia* belong to this species of monomania.

ZOARA, *Insomnia*.

ZOE, *Life*.

ZOGRAPHIA, *Zoographia*.

ZOIATRIA, *Veterinary art*.

ZOIATRUS, Hippiator.

ZOMIDIUM, Jus.

ZOMOS, *Consommé*, Jus.

ZONA, Herpes zoster — z. Castitatis, Hymen — z. Ciliaris, Ciliary zone — z. Ignea, Herpes zoster.

ZONA PELLUCIDA, *Oolemma pellucidum*, Vitellary membrane, Chorion (of some), (F.) Vitelline, Membrane vitelline. The membrane that surrounds the yolk in the mammalia is so called on account of its thickness and peculiar transparency.

ZONA SERPIGINOSA, Herpes zoster.

ZONA TENDINO'SA. The whitish circle around the auriculo-ventricular orifice of the right side of the heart.

ZONA VIRGINITATIS, Hymen.

ZONE, Cingulum.

ZONULA CILIARIS, see Ciliary, and Ciliary zone — z. Cochleæ, see Lamina spiralis — z. Hildani, Cingulum Hildani — z. Zinnii, see Ciliary zone.

ZOOBIOLOGIA, see Physiology.

ZOOCHEMIA, Chymistry, animal.

ZOOCHEMICAL, *Zoöchymical*, *Zoöchemicus*, *Zoöchymicus*. Appertaining or having relation to animal chymistry.

ZOOCHEMY, Chymistry, animal.

ZOÖCHYMY, Chymistry, animal.

ZOÖGENIUM, Azote.

ZOÖG'ENY, *Zoögen'ia*, *Zoögen'esia*, from *ζωον*, 'animal,' and *γενεσις*, 'generation.' The doctrine of animal formation.

ZOÖG'RAPHY, *Zoögra'phia*, *Zogra'phia*, from *ζωον*, and *γραφη*, 'a description.' A description of animals. The natural history of animals.

ZOÖHEMATINE, Hæmatine.

ZOÖHY'LAS, from *ζωον*, and *ὑλη*, 'matter.' Animal matter.

ZOÖHY'LICA CONCREMEN'TA. Concretions formed of animal matter.

ZOÖIASIS, Veterinary art.

ZOÖID, *Zoöides*, *Zoö'des*; from *ζωον*, 'animal,' and *ειδος*, 'resemblance.' Resembling or belonging to an animal. Every living cell may be regarded as a Zooid.

ZOÖLOGIST, *Zoölogie'sta*, *Zoölogus*, from *ζωον*, and *λογος*, 'description.' One skilled in the natural history of animals.

ZOÖLOGY, *Zoölog'ia*. Same etymon. The natural history of animals.

ZOÖMAGNETISMUS, Magnetism, animal.

ZOÖMYCUS, Mucus.

ZOÖMYXA, Mucus.

ZOÖN, Animal.

ZOÖNOM'IA, *Organozoönom'ia*, *Zoö'nomy*, from *ζωον*, and *νομος*, 'law.' The laws which govern the organic actions of animals in general:—*Animal Physiology*, *Zoöphysiolog'ia*.

ZOONOSOLOG'IA, from *ζωον*, *νομος*, 'a disease,' and *λογος*, 'a description.' The nosology of animals.

ZOÖPATHOLOG'IA, from *ζωον*, *παθος*, 'a disease,' and *λογος*, 'a description.' The pathology of animals.

ZOÖPH'AGOUS, *Zoöph'agus*, (F.) *Zoöphage*, from *ζωον*, and *φαγω*, 'I eat.' Feeding on animals. Carnivorous.

ZOÖPTHALMUS, Hydropththalmia.

ZOÖPHYSIOLOGIA, Physiology, animal, Zoönomia.

ZOÖPHYTE, *Zoöph'ytum*, *Phytozoön*, Animal plant, from *ζωον*, and *φυτον*, 'a plant.' A name

given to the lowest class of animals, which seem to form the transition between the animal and the vegetable kingdoms.

ZOÖSIS, Animation, Zoagria.

ZOÖSPERMATA, Spermatozoa.

ZOÖSPERMES, Spermatozoa.

ZOÖTHAP'SIS, from *ζωον*, 'life,' and *θανασις*, 'I inter.' Burial alive. Premature interment.

ZOÖTOMIST, *Zootom'icus*. Same etymon as the next. One who dissects animals.

ZOÖTOMY, *Zoötom'ia*, from *ζωον*, and *τομή*, 'I cut.' *Anatom'ia animalis* seu *comparativa* seu *comparata*, *Theriotomy*, *Comparative Anatomy*. The anatomy of animals. Animal anatomy.

ZOPISSA, see *Pinus sylvestris*.

ZOPYRON, Fomites.

ZOSTER, Herpes zoster.

ZOSTERA MARINA, Pila marina — z. Martima, Pila marina.

ZOSTÈRE, Pila marina.

ZOUZINETTE, Narcissus pseudonarcissus.

ZOZAR, Saccharum.

ZUCARO, Saccharum.

ZUCCARUM, Saccharum.

ZUCHAR, Saccharum.

ZUCHARUM, Saccharum.

ZUCHRA, Saccharum.

ZULAPIUM, Julep.

ZYGAPOPHYSES, see *Vertebræ*.

ZYGOMA, Males os.

ZYGOMATIC, *Zygomat'icus*, *Juga'lis*, *Jugal*, (F.) *Zygomatique*. That which relates to the *zygoma* or cheek-bone.

ZYGOMATIC ARCH, see *Zygomatie process* — z. Bone, Males os.

ZYGOMATIC FOSSA is the space included between the posterior margin of the outer ala of the pterygoid process, and the crista which descends from the malar tuberosity to the superior alveolar margin.

ZYGOMATIC MUSCLES are two in number. 1. *Zygomat'icus major*, *Distort'or Oris*, (F.) *Zygomato-labial* (Ch.), *Muscle grand zygomatique*, which is situate obliquely at the fore-part and sides of the face. It is a small, rounded muscle; arises from the outer surface of the os malar, and is inserted at the commissure of the lips, which it raises and draws outwards. It acts principally in laughing. 2. *Zygomat'icus minor*, (F.) *Petit Zygomato-labial*, (Ch.), *Petit Zygomatique*, does not always exist. It is situate on the inner side of the last; arises from the outer surface of the os malar, and is inserted at the corner of the mouth, or rather into the upper lip, which it raises and draws outwards.

ZYGOMATIC PROCESS, *Processus zygomat'icus*, *Apoph'ysis Zygomat'ica*, *Malar* or *Jugal process*, (F.) *Apophyse Zygomatique*, arises from the outer surface of the temporal bone by two roots, one of which is transverse, (*Condyle du temporal*, Ch.) and the other longitudinal: between them is the glenoid cavity. The process passes forwards, and is articulated with the posterior angle of the os malar, with which it forms a bony bridge, called the *Zygomat'ic arch*, *Arcus* seu *Pons zygomat'icus*, (F.) *Arcade Zygomatique*. The cavity, beneath this bridge, is sometimes called *Zygo'ma*.

ZYGOMATIC SUTURE is placed between the zygomatic parts of the temporal and cheek bones; and slants obliquely downwards and backwards;

ZYGOMATIQUE, *Zygomatie* — z. *Grand*, *Zygomat'icus major* — z. *Petit*, *Zygomat'icus minor*.

ZYGOMATO-LABIAL, *Zygomaticus major*—*z. Labial, petit*, *Zygomaticus major*—*z. Maxillaire*, *Masseter*—*z. Oriculaire*, Anterior auris.

ZYGOPHYLLUM FABA'GO, (F.) *Fabagelle*. Ord. *Zygophyllaceæ*. A Syrian and Mauritanic plant, possessed of an acrid and bitter taste. It has vermifuge properties. The Arabs believe that the recently expressed juice of *Zygophyllum simplex*, is capable of removing specks from the cornea.

ZYMA, Ferment.

ZYMOMA, Ferment

ZYMOSIS, Fermentation, see *Zymotic*.

ZYMOT'IC, *Zymot'icus*, (F.) *Zymotique*, from ζυμω, 'to ferment.' Relating or appertaining to fermentation. A *zymotic disease* is any epidemic, endemic, contagious or sporadic affection, which is produced by some morbid principle acting on the organism similar to a ferment—as the major exanthemata, pyohæmia, &c. *Zymoses* has been used by some to signify such affections.

ZYTHOGALA, Posset.

ZYTHUM, Cerevisia.

ZYTHUS, Cerevisia.

THE END.

BLANCHARD & LEA'S MEDICAL AND SURGICAL PUBLICATIONS.

TO THE MEDICAL PROFESSION.


The prices on the present catalogue are those at which our books can generally be furnished by booksellers throughout the United States, who can readily procure any which they may not have on hand. To physicians who have not convenient access to bookstores, we will, as long as the existing rates of postage remain unchanged, forward them at these prices, *free* by mail, to any post-office in the United States under 1,500 miles. As we open accounts only with booksellers, the amount must in every case, without exception, accompany the order, and we assume no risks of the mail, either on the money or on the books; and as we deal only in our own publications, we can supply no others. Gentlemen desirous of purchasing will, therefore, find it more advantageous to deal with the nearest booksellers whenever practicable.

BLANCHARD & LEA.

PHILADELPHIA, October, 1862.

* * We have now ready a new edition of our **ILLUSTRATED CATALOGUE** of Medical and Scientific Publications, forming an octavo pamphlet of 80 large pages, containing specimens of illustrations, notices of the medical press, &c. &c. It has been prepared without regard to expense, and will be found one of the handsomest specimens of typographical execution as yet presented in this country. Copies will be sent to any address, by mail, free of postage, on receipt of nine cents in stamps.

Catalogues of our numerous publications in miscellaneous and educational literature forwarded on application.

 The attention of physicians is especially solicited to the following important new works and new editions, just issued or nearly ready:—

	See page	
Ashton on the Rectum,	3	
Bumstead on Venereal,	5	
Barclay on Medical Diagnosis,	5	
Barwell on the Joints,	6	
Condie on Diseases of Children,	8	
Churchill's Midwifery,	9	
Druitt's Surgery,	10	
Dalton's Human Physiology, 2d edition,	11	
Dunglison's Medical Dictionary,	12	
Erichsen's System of Surgery,	14	
Flint on the Heart,	14	
Fownes' Manual of Chemistry,	15	
Gross's System of Surgery,	16	
Gray's Anatomy, Descriptive and Surgical, 2d edition,	17	
Hamilton on Fractures and Dislocations,	18	
Hodge on Diseases of Women,	19	
Lyons on Fever,	21	
Meigs on Diseases of Women,	21	
Parrish's Practical Pharmacy,	25	
Stillé's Therapeutics and Materia Medica,	27	
Simpson on Diseases of Women,	27	
Sargent's Minor Surgery, new edition,	28	
Taylor's Medical Jurisprudence,	28	
Toynbee on the Ear,	29	
Watson's Practice of Physic,	30	
Walshe on the Heart,	32	
Winslow on Brain and Mind,	32	
West on Diseases of Women,	32	

TWO MEDICAL PERIODICALS, FREE OF POSTAGE,

Containing over Fifteen Hundred large octavo pages,
FOR FIVE DOLLARS PER ANNUM.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, subject to postage, when not paid for in advance, \$5 00
THE MEDICAL NEWS AND LIBRARY, invariably in advance, 1 00
OR, BOTH PERIODICALS mailed, FREE OF POSTAGE (as long as the existing rates are maintained), to any post-office in the United States, for Five Dollars remitted in advance.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES,

EDITED BY ISAAC HAYS, M. D.,

is published Quarterly, on the first of January, April, July, and October. Each number contains at least two hundred and eighty large octavo pages, handsomely and appropriately illustrated,

wherever necessary. It has now been issued regularly for more than FORTY years, and it has been under the control of the present editor for more than a quarter of a century. Throughout this long period, it has maintained its position in the highest rank of medical periodicals both at home and abroad, and has received the cordial support of the entire profession in this country. Its list of Collaborators will be found to contain a large number of the most distinguished names of the profession in every section of the United States, rendering the department devoted to

ORIGINAL COMMUNICATIONS

full of varied and important matter, of great interest to all practitioners.

As the aim of the Journal, however, is to combine the advantages presented by all the different varieties of periodicals, in its

REVIEW DEPARTMENT

will be found extended and impartial reviews of all important new works, presenting subjects of novelty and interest, together with very numerous

BIBLIOGRAPHICAL NOTICES,

including nearly all the medical publications of the day, both in this country and Great Britain, with a choice selection of the more important continental works. This is followed by the

QUARTERLY SUMMARY,

being a very full and complete abstract, methodically arranged, of the

IMPROVEMENTS AND DISCOVERIES IN THE MEDICAL SCIENCES.

This department of the Journal, so important to the practising physician, is the object of especial care on the part of the editor. It is classified and arranged under different heads, thus facilitating the researches of the reader in pursuit of particular subjects, and will be found to present a very full and accurate digest of all observations, discoveries, and inventions recorded in every branch of medical science. The very extensive arrangements of the publishers are such as to afford to the editor complete materials for this purpose, as he not only regularly receives

ALL THE AMERICAN MEDICAL AND SCIENTIFIC PERIODICALS,

out also twenty or thirty of the more important Journals issued in Great Britain and on the Continent, thus enabling him to present in a convenient compass a thorough and complete abstract of everything interesting or important to the physician occurring in any part of the civilized world.

To their old subscribers, many of whom have been on their list for twenty or thirty years, the publishers feel that no promises for the future are necessary; but those who may desire for the first time to subscribe, can rest assured that no exertion will be spared to maintain the Journal in the high position which it has occupied for so long a period.

By reference to the terms it will be seen that, in addition to this large amount of valuable and practical information on every branch of medical science, the subscriber, by paying in advance, becomes entitled, without further charge, to

THE MEDICAL NEWS AND LIBRARY,

a monthly periodical of thirty-two large octavo pages. Its "NEWS DEPARTMENT" presents the current information of the day, while the "LIBRARY DEPARTMENT" is devoted to presenting standard works on various branches of medicine. Within a few years, subscribers have thus received, without expense, many works of the highest character and practical value, such as "Watson's Practice," "Todd and Bowman's Physiology," "Malgaigne's Surgery," "West on Children," "West on Females, Part I.," "Habershon on the Alimentary Canal," &c.

While the work at present appearing in its columns is

CLINICAL LECTURES ON THE DISEASES OF WOMEN.

By PROFESSOR J. Y. SIMPSON, of Edinburgh.

WITH NUMEROUS HANDSOME ILLUSTRATIONS.

These Lectures, published in England under the supervision of the Author, carry with them all the weight of his wide experience and distinguished reputation. Their eminently practical nature, and the importance of the subject treated, cannot fail to render them in the highest degree satisfactory to subscribers, who can thus secure them without cost. These Lectures are continued in the "News" for 1862.

It will thus be seen that for the small sum of FIVE DOLLARS, paid in advance, the subscriber will obtain a Quarterly and a Monthly periodical,

EMBRACING NEARLY SIXTEEN HUNDRED LARGE OCTAVO PAGES.

Those subscribers who do not pay in advance will bear in mind that their subscription of Five Dollars will entitle them to the Journal only, without the News, and that they will be at the expense of their own postage on the receipt of each number. The advantage of a remittance when ordering the Journal will thus be apparent.

Remittances of subscriptions can be mailed at our risk, when a certificate is taken from the Postmaster that the money is duly inclosed and forwarded.

Address

BLANCHARD & LEA, PHILADELPHIA.

ASHTON (T. J.),

Surgeon to the Blenheim Dispensary, &c.

ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE RECTUM AND ANUS; with remarks on Habitual Constipation. From the third and enlarged London edition. With handsome illustrations. In one very beautifully printed octavo volume, of about 300 pages. (*Just Issued.*) \$2 00.

The most complete one we possess on the subject. *Medico-Chirurgical Review.*

We are satisfied, after a careful examination of the volume, and a comparison of its contents with those of its leading predecessors and contemporaries, that the best way for the reader to avail himself of

the excellent advice given in the concluding paragraph above, would be to provide himself with a copy of the book from which it has been taken, and diligently to con its instructive pages. They may secure to him in any a triumph and fervent blessing.—*Am. Journal Med. Sciences.*

ALLEN (J. M.), M. D.,

Professor of Anatomy in the Pennsylvania Medical College, &c.

THE PRACTICAL ANATOMIST; or, The Student's Guide in the Dissecting-room. With 266 illustrations. In one handsome royal 12mo. volume, of over 600 pages, leather. \$2 25.

We believe it to be one of the most useful works upon the subject ever written. It is handsomely illustrated, well printed, and will be found of convenient size for use in the dissecting-room.—*Med. Examiner.*

However valuable may be the "Dissector's Guides" which we, of late, have had occasion to

notice, we feel confident that the work of Dr. Allen is superior to any of them. We believe with the author, that none is so fully illustrated as this, and the arrangement of the work is such as to facilitate the labors of the student. We most cordially recommend it to their attention.—*Western Lancet.*

ANATOMICAL ATLAS.

By Professors H. H. SMITH and W. E. HORNER, of the University of Pennsylvania. 1 vol. 8vo., extra cloth, with nearly 650 illustrations. See SMITH, p. 331.

ABEL (F. A.), F. C. S. AND C. L. BLOXAM.

HANDBOOK OF CHEMISTRY, Theoretical, Practical, and Technical; with a Recommendatory Preface by Dr. HOFMANN. In one large octavo volume, extra cloth, of 662 pages, with illustrations. \$3 25.

ASHWELL (SAMUEL), M. D.,

Obstetric Physician and Lecturer to Guy's Hospital, London.

A PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN.

Illustrated by Cases derived from Hospital and Private Practice. Third American, from the Third and revised London edition. In one octavo volume, extra cloth, of 528 pages. \$3 00.

The most useful practical work on the subject in the English language.—*Boston Med. and Surg. Journal.*

The most able, and certainly the most standard and practical, work on female diseases that we have yet seen.—*Medico-Chirurgical Review.*

ARNOTT (NEILL), M. D.

ELEMENTS OF PHYSICS; or Natural Philosophy, General and Medical.

Written for universal use, in plain or non-technical language. A new edition, by ISAAC HAYS, M. D. Complete in one octavo volume, leather, of 484 pages, with about two hundred illustrations. \$2 50.

BIRD (GOLDING), A. M., M. D., &c.

URINARY DEPOSITS: THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS. Edited by EDMUND LLOYD BIRKETT, M. D. A new American, from the fifth and enlarged London edition. With eighty illustrations on wood. In one handsome octavo volume, of about 400 pages, extra cloth. \$2 00. (*Just Issued.*)

The death of Dr. Bird has rendered it necessary to entrust the revision of the present edition to other hands, and in his performance of the duty thus devolving on him, Dr. Birkett has sedulously endeavored to carry out the author's plan by introducing such new matter and modifications of the text as the progress of science has called for. Notwithstanding the utmost care to keep the work within a reasonable compass, these additions have resulted in a considerable enlargement. It is, therefore, hoped that it will be found fully up to the present condition of the subject, and that the reputation of the volume as a clear, complete, and compendious manual, will be fully maintained.

BENNETT (J. HUGHES), M. D., F. R. S. E.,

Professor of Clinical Medicine in the University of Edinburgh, &c.

THE PATHOLOGY AND TREATMENT OF PULMONARY TUBERCULOSIS, and on the Local Medication of Pharyngeal and Laryngeal Diseases frequently mistaken for or associated with, Phthisis. One vol. 8vo., extra cloth, with wood-cuts. pp. 130. \$1 25.

BARLOW (GEORGE H.), M. D.

Physician to Guy's Hospital, London, &c.

A MANUAL OF THE PRACTICE OF MEDICINE. With Additions by D. F. CONDIE, M. D., author of "A Practical Treatise on Diseases of Children," &c. In one handsome octavo volume, leather, of over 600 pages. \$2 75.

We recommend Dr. Barlow's Manual in the warmest manner as a most valuable *va-de-mecum*. We have had frequent occasion to consult it, and have

found it clear, concise, practical, and sound.—*Boston Med. and Surg. Journal.*

BUDD (GEORGE), M. D., F. R. S.,
Professor of Medicine in King's College, London.

ON DISEASES OF THE LIVER. Third American, from the third and enlarged London edition. In one very handsome octavo volume, extra cloth, with four beautifully colored plates, and numerous wood-cuts. pp. 500. \$3 00.

Has fairly established for itself a place among the classical medical literature of England.—*British and Foreign Medico-Chir. Review.*

Dr. Budd's Treatise on Diseases of the Liver is now a standard work in Medical literature, and during the intervals which have elapsed between the successive editions, the author has incorporated into the text the most striking novelties which have characterized the recent progress of hepatic physiology and pathology; so that although the size of the book

is not perceptibly changed, the history of liver diseases is made more complete, and is kept upon a level with the progress of modern science. It is the best work on Diseases of the Liver in any language.—*London Med. Times and Gazette.*

This work, now the standard book of reference on the diseases of which it treats, has been carefully revised, and many new illustrations of the views of the learned author added in the present edition.—*Dublin Quarterly Journal.*

BY THE SAME AUTHOR.

ON THE ORGANIC DISEASES AND FUNCTIONAL DISORDERS OF THE STOMACH. In one neat octavo volume, extra cloth. \$1 50.

BUCKNILL (J. C.), M. D., AND **DANIEL H. TUKE, M. D.,**
Medical Superintendent of the Devon Lunatic Asylum. Visiting Medical Officer to the York Retreat.

A MANUAL OF PSYCHOLOGICAL MEDICINE; containing the History, Nosology, Description, Statistics, Diagnosis, Pathology, and Treatment of INSANITY. With a Plate. In one handsome octavo volume, of 536 pages. \$3 00.

The increase of mental disease in its various forms, and the difficult questions to which it is constantly giving rise, render the subject one of daily enhanced interest, requiring on the part of the physician a constantly greater familiarity with this, the most perplexing branch of his profession. At the same time there has been for some years no work accessible in this country, presenting the results of recent investigations in the Diagnosis and Prognosis of Insanity, and the greatly improved methods of treatment which have done so much in alleviating the condition or restoring the health of the insane. To fill this vacancy the publishers present this volume, assured that the distinguished reputation and experience of the authors will entitle it at once to the confidence of both student and practitioner. Its scope may be gathered from the declaration of the authors that "their aim has been to supply a text book which may serve as a guide in the acquisition of such knowledge, sufficiently elementary to be adapted to the wants of the student, and sufficiently modern in its views and explicit in its teaching to suffice for the demands of the practitioner."

BENNETT (HENRY), M. D.

A PRACTICAL TREATISE ON INFLAMMATION OF THE UTERUS, ITS CERVIX AND APPENDAGES, and on its connection with Uterine Disease. To which is added, a Review of the present state of Uterine Pathology. Fifth American, from the third English edition. In one octavo volume, of about 500 pages, extra cloth. \$2 00.

BROWN (ISAAC BAKER),

Surgeon-Accoucheur to St. Mary's Hospital, &c.

ON SOME DISEASES OF WOMEN ADMITTING OF SURGICAL TREATMENT. With handsome illustrations. One vol. 8vo., extra cloth, pp. 276. \$1 60.

Mr. Brown has earned for himself a high reputation in the operative treatment of sundry diseases and injuries to which females are peculiarly subject. We can truly say of his work that it is an important addition to obstetrical literature. The operative suggestions and contrivances which Mr. Brown describes, exhibit much practical sagacity and skill,

and merit the careful attention of every surgeon—accoucheur.—*Association Journal.*

We have no hesitation in recommending this book to the careful attention of all surgeons who make female complaints a part of their study and practice.—*Dublin Quarterly Journal.*

BOWMAN (JOHN E.), M. D.

PRACTICAL HANDBOOK OF MEDICAL CHEMISTRY. Second American, from the third and revised English Edition. In one neat volume, royal 12mo., extra cloth, with numerous illustrations. pp. 288. \$1 25.

BY THE SAME AUTHOR.

INTRODUCTION TO PRACTICAL CHEMISTRY, INCLUDING ANALYSIS. Second American, from the second and revised London edition. With numerous illustrations. In one neat vol., royal 12mo., extra cloth. pp. 350. \$1 25.

BEALE ON THE LAWS OF HEALTH IN RELATION TO MIND AND BODY. A Series of Letters from an old Practitioner to a Patient. In one volume, royal 12mo., extra cloth. pp. 296. 80 cents.

BUSHNAN'S PHYSIOLOGY OF ANIMAL AND VEGETABLE LIFE; a Popular Treatise on the Functions and Phenomena of Organic Life. In one handsome royal 12mo. volume, extra cloth, with over 100 illustrations. pp. 234. 80 cents.

BUCKLER ON THE ETIOLOGY, PATHOLOGY, AND TREATMENT OF FIBRO-BRONCHITIS AND RHEUMATIC PNEUMONIA. In one 8vo. volume, extra cloth. pp. 150. \$1 25.

BLOOD AND URINE (MANUALS ON). BY JOHN WILLIAM GRIFFITH, G. OWEN REESE, AND ALFRED MARKWICK. One thick volume, royal 12mo., extra cloth, with plates. pp. 460. \$1 25.

BRODIE'S CLINICAL LECTURES ON SURGERY. 1 vol. 8vo. cloth. 350 pp. \$1 25.

BUMSTEAD (FREEMAN J.) M. D.,

Lecturer on Venereal Diseases at the College of Physicians and Surgeons, New York, &c.

THE PATHOLOGY AND TREATMENT OF VENEREAL DISEASES,
including the results of recent investigations upon the subject. With illustrations on wood. In one very handsome octavo volume, of nearly 700 pages, extra cloth; \$3 75. (Now Ready.)

By far the most valuable contribution to this particular branch of practice that has seen the light within the last score of years. His clear and accurate descriptions of the various forms of venereal disease, and especially the methods of treatment he proposes, are worthy of the highest encomium. In these respects it is better adapted for the assistance of the every-day practitioner than any other with which we are acquainted. In variety of methods proposed, in minuteness of direction, guided by careful discrimination of varying forms and complications, we write down the book as unsurpassed. It is a work which should be in the possession of every practitioner.—*Chicago Med. Journal*, Nov. 1861.

The foregoing admirable volume comes to us, embracing the whole subject of syphilology, resolving many a doubt, correcting and confirming many an entertained opinion, and in our estimation the best, completest, fullest monograph on this subject in our language. As far as the author's labors themselves are concerned, we feel it a duty to say that he has not only exhausted his subject, but he has presented to us, without the slightest hyperbole, the best digested treatise on these diseases in our language. He has carried its literature down to the present moment, and has achieved his task in a manner which cannot but redound to his credit.—*British American Journal*, Oct. 1861.

We believe this treatise will come to be regarded as high authority in this branch of medical practice, and we cordially commend it to the favorable notice of our brethren in the profession. For our own part, we candidly confess that we have received many new ideas from its perusal, as well as modified many views which we have long, and, as we now think, erroneously entertained on the subject of syphilis.

To sum up all in a few words, this book is one which no practising physician or medical student can very well afford to do without.—*American Med. Times*, Nov. 2, 1861.

The whole work presents a complete history of venereal diseases, comprising much interesting and valuable material that has been spread through medical journals within the last twenty years—the period of many experiments and investigations on the subject—the whole carefully digested by the aid of the author's extensive personal experience, and offered to the profession in an admirable form. Its completeness is secured by good plates, which are especially full in the anatomy of the genital organs. We have examined it with great satisfaction, and congratulate the medical profession in America on the nationality of a work that may fairly be called original.—*Berkshire Med. Journal*, Dec. 1861.

One thing, however, we are impelled to say, that we have met with no other book on syphilis, in the English language, which gave so full, clear, and impartial views of the important subjects on which it treats. We cannot, however, refrain from expressing our satisfaction with the full and perspicuous manner in which the subject has been presented, and the careful attention to minute details, so useful—not to say indispensable—in a practical treatise. In conclusion, if we may be pardoned the use of a phrase now become stereotyped, but which we here employ in all seriousness and sincerity, we do not hesitate to express the opinion that Dr. Bumstead's Treatise on Venereal Diseases is a "work without which no medical library will hereafter be considered complete."—*Boston Med. and Surg. Journal*, Sept. 5, 1861.

BARCLAY (A. W.), M. D.,

Assistant Physician to St. George's Hospital, &c.

A MANUAL OF MEDICAL DIAGNOSIS; being an Analysis of the Signs and Symptoms of Disease. Second American from the second and revised London edition. In one neat octavo volume, extra cloth, of 451 pages. \$2 25. (Now ready.)

The demand for a second edition of this work shows that the vacancy which it attempts to supply has been recognized by the profession, and that the efforts of the author to meet the want have been successful. The revision which it has enjoyed will render it better adapted than before to afford assistance to the learner in the prosecution of his studies, and to the practitioner who requires a convenient and accessible manual for speedy reference in the exigencies of his daily duties. For this latter purpose its complete and extensive Index renders it especially valuable, offering facilities for immediately turning to any class of symptoms, or any variety of disease.

The task of composing such a work is neither an easy nor a light one; but Dr. Barclay has performed it in a manner which meets our most unqualified approbation. He is no mere theorist; he knows his work thoroughly, and in attempting to perform it, has not exceeded his powers.—*British Med. Journal*.

We venture to predict that the work will be deservedly popular, and soon become, like Watson's Practice, an indispensable necessity to the practitioner.—*N. A. Med. Journal*.

An inestimable work of reference for the young practitioner and student.—*Nashville Med. Journal*.

We hope the volume will have an extensive circulation, not among students of medicine only, but practitioners also. They will never regret a faithful study of its pages.—*Cincinnati Lancet*.

An important acquisition to medical literature. It is a work of high merit, both from the vast importance of the subject upon which it treats, and also from the real ability displayed in its elaboration. In conclusion, let us bespeak for this volume that attention of every student of our art which it so richly deserves—that place in every medical library which it can so well adorn.—*Peninsular Medical Journal*.

BARTLETT (ELISHA), M. D.**THE HISTORY, DIAGNOSIS, AND TREATMENT OF THE FEVERS OF THE UNITED STATES.** A new and revised edition. By ALONZO CLARK, M. D., Prof. of Pathology and Practical Medicine in the N. Y. College of Physicians and Surgeons, &c. In one octavo volume, of six hundred pages, extra cloth. Price \$3 00.

It is a work of great practical value and interest, containing much that is new relative to the several diseases of which it treats, and, with the additions of the editor, is fully up to the times. The distinctive features of the different forms of fever are plainly and forcibly portrayed, and the lines of demarcation carefully and accurately drawn, and to the American practitioner is a more valuable and safe guide than any work on fever extant.—*Ohio Med. and Surg. Journal*.

This excellent monograph on febrile disease, has

stood deservedly high since its first publication. It will be seen that it has now reached its fourth edition under the supervision of Prof. A. Clark, a gentleman who, from the nature of his studies and pursuits, is well calculated to appreciate and discuss the many intricate and difficult questions in pathology. His annotations add much to the interest of the work, and have brought it well up to the condition of the science as it exists at the present day in regard to this class of diseases.—*Southern Med. and Surg. Journal*.

BARWELL (RICHARD,) F. R. C. S.,

Assistant Surgeon Charing Cross Hospital, &c.

A TREATISE ON DISEASES OF THE JOINTS. Illustrated with engravings on wood. In one very handsome octavo volume, of about 500 pages, extra cloth; \$5 00. (Now Ready.)

"A treatise on Diseases of the Joints equal to, or rather beyond the current knowledge of the day, has long been required—my professional brethren must judge whether the ensuing pages may supply the deficiency. No author is fit to estimate his own work at the moment of its completion, but it may be permitted me to say that the study of joint diseases has very much occupied my attention, even from my studentship, and that for the last six or eight years my devotion to that subject has been almost unremitting. . . . The real weight of my work has been at the bedside, and the greatest labor devoted to interpreting symptoms and remedying their cause."—AUTHOR'S PREFACE.

At the outset we may state that the work is worthy of much praise, and bears evidence of much thoughtful and careful inquiry, and here and there of no slight originality. We have already carried this notice further than we intended to do, but not to the extent the work deserves. We can only add, that the perusal of it has afforded us great pleasure. The author has evidently worked very hard at his subject, and his investigations into the Physiology and Pathology of Joints have been carried on in a manner which entitles him to be listened to with attention and respect. We must not omit to mention the very admirable plates with which the volume is enriched. We seldom meet with such striking and faithful delineations of disease.—*London Med. Times and Gazette*, Feb. 9, 1861.

We cannot take leave, however, of Mr. Barwell, without congratulating him on the interesting amount of information which he has compressed into his book. The work appears to us calculated

to be of much use to the practising surgeon who may be in want of a treatise on diseases of the joints, and at the same time one which contains the latest information on articular affections and the operations for their cure.—*Dublin Med. Press*, Feb. 27, 1861.

This volume will be welcomed, both by the pathologist and the surgeon, as being the record of much honest research and careful investigation into the nature and treatment of a most important class of disorders. We cannot conclude this notice of a valuable and useful book without calling attention to the amount of *bonâ fide* work it contains. In the present day of universal book-making, it is no slight matter for a volume to show laborious investigation, and at the same time original thought, on the part of its author, whom we may congratulate on the successful completion of his arduous task.—*London Lancet*, March 9, 1861.

CARPENTER (WILLIAM B.), M. D., F. R. S., &c.,

Examiner in Physiology and Comparative Anatomy in the University of London.

PRINCIPLES OF HUMAN PHYSIOLOGY; with their chief applications to Psychology, Pathology, Therapeutics, Hygiene, and Forensic Medicine. A new American, from the last and revised London edition. With nearly three hundred illustrations. Edited, with additions, by FRANCIS GURNEY SMITH, M. D., Professor of the Institutes of Medicine in the Pennsylvania Medical College, &c. In one very large and beautiful octavo volume, of about nine hundred large pages, handsomely printed and strongly bound in leather, with raised bands. \$4 25.

In the preparation of this new edition, the author has spared no labor to render it, as heretofore, a complete and lucid exposition of the most advanced condition of its important subject. The amount of the additions required to effect this object thoroughly, joined to the former large size of the volume, presenting objections arising from the unwieldy bulk of the work, he has omitted all those portions not bearing directly upon HUMAN PHYSIOLOGY, designing to incorporate them in his forthcoming Treatise on GENERAL PHYSIOLOGY. As a full and accurate text-book on the Physiology of Man, the work in its present condition therefore presents even greater claims upon the student and physician than those which have heretofore won for it the very wide and distinguished favor which it has so long enjoyed. The additions of Prof. Smith will be found to supply whatever may have been wanting to the American student, while the introduction of many new illustrations, and the most careful mechanical execution, render the volume one of the most attractive as yet issued.

For upwards of thirteen years Dr. Carpenter's work has been considered by the profession generally, both in this country and England, as the most valuable compendium on the subject of physiology in our language. This distinction it owes to the high attainments and unwearied industry of its accomplished author. The present edition (which, like the last American one, was prepared by the author himself), is the result of such extensive revision, that it may almost be considered a new work. We need hardly say, in concluding this brief notice, that while the work is indispensable to every student of medicine in this country, it will amply repay the practitioner for its perusal by the interest and value of its contents.—*Boston Med. and Surg. Journal*.

This is a standard work—the text-book used by all medical students who read the English language. It has passed through several editions in order to keep pace with the rapidly growing science of Physiology. Nothing need be said in its praise, for its merits are universally known; we have nothing to say of its defects, for they only appear where the science of which it treats is incomplete.—*Western Lancet*.

The most complete exposition of physiology which any language can at present give.—*Brit. and For. Med.-Chirurg. Review*.

The greatest, the most reliable, and the best book on the subject which we know of in the English language.—*Stethoscope*.

To eulogize this great work would be superfluous. We should observe, however, that in this edition the author has remodelled a large portion of the former, and the editor has added much matter of interest, especially in the form of illustrations. We may confidently recommend it as the most complete work on Human Physiology in our language.—*Southern Med. and Surg. Journal*.

The most complete work on the science in our language.—*Am. Med. Journal*.

The most complete work now extant in our language.—*N. O. Med. Register*.

The best text-book in the language on this extensive subject.—*London Med. Times*.

A complete cyclopædia of this branch of science.—*N. Y. Med. Times*.

The profession of this country, and perhaps also of Europe, have anxiously and for some time awaited the announcement of this new edition of Carpenter's Human Physiology. His former editions have for many years been almost the only text-book on Physiology in all our medical schools, and its circulation among the profession has been unsurpassed by any work in any department of medical science.

It is quite unnecessary for us to speak of this work as its merits would justify. The mere announcement of its appearance will afford the highest pleasure to every student of Physiology, while its perusal will be of infinite service in advancing physiological science.—*Ohio Med. and Surg. Journ.*

CARPENTER (WILLIAM B.), M. D., F. R. S.,

Examiner in Physiology and Comparative Anatomy in the University of London.

THE MICROSCOPE AND ITS REVELATIONS. With an Appendix containing the Applications of the Microscope to Clinical Medicine, &c. By F. G. SMITH, M. D. Illustrated by four hundred and thirty-four beautiful engravings on wood. In one large and very handsome octavo volume, of 724 pages, extra cloth, \$4 00; leather, \$4 50.

Dr. Carpenter's position as a microscopist and physiologist, and his great experience as a teacher, eminently qualify him to produce what has long been wanted—a good text-book on the practical use of the microscope. In the present volume his object has been, as stated in his Preface, “to combine, within a moderate compass, that information with regard to the use of his ‘tools,’ which is most essential to the working microscopist, with such an account of the objects best fitted for his study, as might qualify him to comprehend what he observes, and might thus prepare him to benefit science, whilst expanding and refreshing his own mind.” That he has succeeded in accomplishing this, no one acquainted with his previous labors can doubt.

The great importance of the microscope as a means of diagnosis, and the number of microscopists who are also physicians, have induced the American publishers, with the author's approval, to add an Appendix, carefully prepared by Professor Smith, on the applications of the instrument to clinical medicine, together with an account of American Microscopes, their modifications and accessories. This portion of the work is illustrated with nearly one hundred wood-cuts, and, it is hoped, will adapt the volume more particularly to the use of the American student.

Those who are acquainted with Dr. Carpenter's previous writings on Animal and Vegetable Physiology, will fully understand how vast a store of knowledge he is able to bring to bear upon so comprehensive a subject as the revelations of the microscope; and even those who have no previous acquaintance with the construction or uses of this instrument, will find abundance of information conveyed in clear and simple language.—*Med. Times and Gazette.*

Although originally not intended as a strictly

medical work, the additions by Prof. Smith give it a positive claim upon the profession, for which we doubt not he will receive their sincere thanks. Indeed, we know not where the student of medicine will find such a complete and satisfactory collection of microscopic facts bearing upon physiology and practical medicine as is contained in Prof. Smith's appendix; and this of itself, it seems to us, is fully worth the cost of the volume.—*Louisville Medical Review.*

BY THE SAME AUTHOR.

ELEMENTS (OR MANUAL) OF PHYSIOLOGY, INCLUDING PHYSIOLOGICAL ANATOMY. Second American, from a new and revised London edition. With one hundred and ninety illustrations. In one very handsome octavo volume, leather. pp. 566. \$3 00.

In publishing the first edition of this work, its title was altered from that of the London volume, by the substitution of the word “Elements” for that of “Manual,” and with the author's sanction the title of “Elements” is still retained as being more expressive of the scope of the treatise.

To say that it is the best manual of Physiology now before the public, would not do sufficient justice to the author.—*Buffalo Medical Journal.*

In his former works it would seem that he had exhausted the subject of Physiology. In the present, he gives the essence, as it were, of the whole.—*N. Y. Journal of Medicine.*

Those who have occasion for an elementary treatise on Physiology, cannot do better than to possess themselves of the manual of Dr. Carpenter.—*Medical Examiner.*

The best and most complete exposé of modern Physiology, in one volume, extant in the English language.—*St. Louis Medical Journal.*

BY THE SAME AUTHOR.

PRINCIPLES OF COMPARATIVE PHYSIOLOGY. New American, from the Fourth and Revised London edition. In one large and handsome octavo volume, with over three hundred beautiful illustrations. pp. 752. Extra cloth, \$4 80; leather, raised bands, \$5 25.

This book should not only be read but thoroughly studied by every member of the profession. None are too wise or old, to be benefited thereby. But especially to the younger class would we cordially commend it as best fitted of any work in the English language to qualify them for the reception and comprehension of those truths which are daily being developed in physiology.—*Medical Counsellor.*

Without pretending to it, it is an encyclopedia of the subject, accurate and complete in all respects—a truthful reflection of the advanced state at which the science has now arrived.—*Dublin Quarterly Journal of Medical Science.*

A truly magnificent work—in itself a perfect physiological study.—*Ranking's Abstract.*

This work stands without its fellow. It is one few men in Europe could have undertaken; it is one

no man, we believe, could have brought to so successful an issue as Dr. Carpenter. It required for its production a physiologist at once deeply read in the labors of others, capable of taking a general, critical, and unprejudiced view of those labors, and of combining the varied, heterogeneous materials at his disposal, so as to form an harmonious whole. We feel that this abstract can give the reader a very imperfect idea of the fulness of this work, and no idea of its unity, of the admirable manner in which material has been brought, from the most various sources, to conduce to its completeness, of the lucidity of the reasoning it contains, or of the clearness of language in which the whole is clothed. Not the profession only, but the scientific world at large, must feel deeply indebted to Dr. Carpenter for this great work. It must, indeed, add largely even to his high reputation.—*Medical Times.*

BY THE SAME AUTHOR. (*Preparing.*)

PRINCIPLES OF GENERAL PHYSIOLOGY, INCLUDING ORGANIC CHEMISTRY AND HISTOLOGY. With a General Sketch of the Vegetable and Animal Kingdom. In one large and very handsome octavo volume, with several hundred illustrations.

BY THE SAME AUTHOR.

A PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. New edition, with a Preface by D. F. CONDIE, M. D., and explanations of scientific words. In one neat 12mo. volume, extra cloth. pp. 178. 50 cents.

CONDIE (D. F.), M. D., &c.

A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Fifth edition, revised and augmented. In one large volume, 8vo., leather, of over 750 pages. \$3 25. (*Just Issued*, 1859.)

In presenting a new and revised edition of this favorite work, the publishers have only to state that the author has endeavored to render it in every respect "a complete and faithful exposition of the pathology and therapeutics of the maladies incident to the earlier stages of existence—a full and exact account of the diseases of infancy and childhood." To accomplish this he has subjected the whole work to a careful and thorough revision, rewriting a considerable portion, and adding several new chapters. In this manner it is hoped that any deficiencies which may have previously existed have been supplied, that the recent labors of practitioners and observers have been thoroughly incorporated, and that in every point the work will be found to maintain the high reputation it has enjoyed as a complete and thoroughly practical book of reference in infantile affections.

A few notices of previous editions are subjoined.

Dr. Condie's scholarship, acumen, industry, and practical sense are manifested in this, as in all his numerous contributions to science.—*Dr. Holmes's Report to the American Medical Association.*

Taken as a whole, in our judgment, Dr. Condie's Treatise is the one from the perusal of which the practitioner in this country will rise with the greatest satisfaction.—*Western Journal of Medicine and Surgery.*

One of the best works upon the Diseases of Children in the English language.—*Western Lancet.*

We feel assured from actual experience that no physician's library can be complete without a copy of this work.—*N. Y. Journal of Medicine.*

A veritable paediatric encyclopædia, and an honor to American medical literature.—*Ohio Medical and Surgical Journal.*

We feel persuaded that the American medical profession will soon regard it not only as a very good, but as the VERY BEST "Practical Treatise on the Diseases of Children."—*American Medical Journal.*

In the department of infantile therapeutics, the work of Dr. Condie is considered one of the best which has been published in the English language.—*The Stethoscope.*

We pronounced the first edition to be the best work on the diseases of children in the English language, and, notwithstanding all that has been published, we still regard it in that light.—*Medical Examiner.*

The value of works by native authors on the diseases which the physician is called upon to combat, will be appreciated by all; and the work of Dr. Condie has gained for itself the character of a safe guide for students, and a useful work for consultation by those engaged in practice.—*N. Y. Med. Times.*

This is the fourth edition of this deservedly popular treatise. During the interval since the last edition, it has been subjected to a thorough revision by the author; and all new observations in the pathology and therapeutics of children have been included in the present volume. As we said before, we do not know of a better book on diseases of children, and to a large part of its recommendations we yield an unhesitating concurrence.—*Buffalo Med. Journal.*

Perhaps the most full and complete work now before the profession of the United States; indeed, we may say in the English language. It is vastly superior to most of its predecessors.—*Transylvania Med. Journal.*

CHRISTISON (ROBERT), M. D., V. P. R. S. E., &c.

A DISPENSATORY; or, Commentary on the Pharmacopœias of Great Britain and the United States; comprising the Natural History, Description, Chemistry, Pharmacy, Actions, Uses, and Doses of the Articles of the Materia Medica. Second edition, revised and improved, with a Supplement containing the most important New Remedies. With copious Additions, and two hundred and thirteen large wood-engravings. By R. EGLESFELD GRIFFITH, M. D. In one very large and handsome octavo volume, leather, raised bands, of over 1000 pages. \$3 50.

COOPER (BRANSBY B.), F. R. S.

LECTURES ON THE PRINCIPLES AND PRACTICE OF SURGERY.

In one very large octavo volume, extra cloth, of 750 pages. \$3 00.

COOPER ON DISLOCATIONS AND FRACTURES OF THE JOINTS.—Edited by BRANSBY B. COOPER, F. R. S., &c. With additional Observations by Prof. J. C. WARREN. A new American edition. In one handsome octavo volume, extra cloth, of about 500 pages, with numerous illustrations on wood. \$3 25.

COOPER ON THE ANATOMY AND DISEASES OF THE BREAST, with twenty-five Miscellaneous and Surgical Papers. One large volume, imperial 8vo., extra cloth, with 252 figures, on 36 plates. \$2 50.

COOPER ON THE STRUCTURE AND DISEASES OF THE TESTIS, AND ON THE THYMUS GLAND. One vol. imperial 8vo., extra cloth, with 177 figures on 29 plates. \$2 00.

COPLAND ON THE CAUSES, NATURE, AND TREATMENT OF PALSY AND APOPLEXY. In one volume, royal 12mo., extra cloth. pp. 326. 80 cents.

CLYMER ON FEVERS; THEIR DIAGNOSIS, PATHOLOGY, AND TREATMENT In one octavo volume, leather, of 600 pages. \$1 50.

COLOMBAT DE L'ISERE ON THE DISEASES OF FEMALES, and on the special Hygiene of their Sex. Translated, with many Notes and Additions, by C. D. MEIES, M. D. Second edition, revised and improved. In one large volume, octavo, leather, with numerous wood-cuts. pp. 720. \$3 50.

CARSON (JOSEPH), M. D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania.

SYNOPSIS OF THE COURSE OF LECTURES ON MATERIA MEDICA AND PHARMACY, delivered in the University of Pennsylvania. Second and revised edition. In one very neat octavo volume, extra cloth, of 208 pages. \$1 50.

CURLING (T. B.), F. R. S.,

Surgeon to the London Hospital, President of the Hunterian Society, &c.

A PRACTICAL TREATISE ON DISEASES OF THE TESTIS, SPERMATIC CORD, AND SCROTUM. Second American, from the second and enlarged English edition. In one handsome octavo volume, extra cloth, with numerous illustrations. pp. 420. \$2 00.

CHURCHILL (FLEETWOOD), M. D., M. R. I. A.
ON THE THEORY AND PRACTICE OF MIDWIFERY. A new American from the fourth revised and enlarged London edition. With Notes and Additions, by D. FRANCIS CONDIE, M. D., author of a "Practical Treatise on the Diseases of Children," &c. With 104 illustrations. In one very handsome octavo volume, leather, of nearly 700 large pages. \$3 50. (*Just Issued.*)

This work has been so long an established favorite, both as a text-book for the learner and as a reliable aid in consultation for the practitioner, that in presenting a new edition it is only necessary to call attention to the very extended improvements which it has received. Having had the benefit of two revisions by the author since the last American reprint, it has been materially enlarged, and Dr. Churchill's well-known conscientious industry is a guarantee that every portion has been thoroughly brought up with the latest results of European investigation in all departments of the science and art of obstetrics. The recent date of the last Dublin edition has not left much of novelty for the American editor to introduce, but he has endeavored to insert whatever has since appeared, together with such matters as his experience has shown him would be desirable for the American student, including a large number of illustrations. With the sanction of the author he has added in the form of an appendix, some chapters from a little "Manual for Midwives and Nurses," recently issued by Dr. Churchill, believing that the details there presented can hardly fail to prove of advantage to the junior practitioner. The result of all these additions is that the work now contains fully one-half more matter than the last American edition, with nearly one-half more illustrations, so that notwithstanding the use of a smaller type, the volume contains almost two hundred pages more than before.

No effort has been spared to secure an improvement in the mechanical execution of the work equal to that which the text has received, and the volume is confidently presented as one of the handsomest that has thus far been laid before the American profession; while the very low price at which it is offered should secure for it a place in every lecture-room and on every office table.

A better book in which to learn these important points we have not met than Dr. Churchill's. Every page of it is full of instruction; the opinion of all writers of authority is given on questions of difficulty, as well as the directions and advice of the learned author himself, to which he adds the result of statistical inquiry, putting statistics in their proper place and giving them their due weight, and no more. We have never read a book more free from professional jealousy than Dr. Churchill's. It appears to be written with the true design of a book on medicine, viz: to give all that is known on the subject of which he treats, both theoretically and practically, and to advance such opinions of his own as he believes will benefit medical science, and insure the safety of the patient. We have said enough to convey to the profession that this book of Dr. Churchill's is admirably suited for a book of reference for the practitioner, as well as a text-book for the student, and we hope it may be extensively purchased amongst our readers. To them we most strongly recommend it.—*Dublin Medical Press*, June 20, 1860.

To bestow praise on a book that has received such marked approbation would be superfluous. We need only say, therefore, that if the first edition was thought worthy of a favorable reception by the medical public, we can confidently affirm that this will be found much more so. The lecturer, the practitioner, and the student, may all have recourse to its pages, and derive from their perusal much interest and instruction in everything relating to theoretical and practical midwifery.—*Dublin Quarterly Journal of Medical Science*.

A work of very great merit, and such as we can confidently recommend to the study of every obstetric practitioner.—*London Medical Gazette*.

This is certainly the most perfect system extant. It is the best adapted for the purposes of a text-book, and that which he whose necessities confine him to one book, should select in preference to all others.—*Southern Medical and Surgical Journal*.

BY THE SAME AUTHOR. (*Lately Published.*)

ON THE DISEASES OF INFANTS AND CHILDREN. Second American Edition, revised and enlarged by the author. Edited, with Notes, by W. V. KEATING, M. D. In one large and handsome volume, extra cloth, of over 700 pages. \$3 00, or in leather, \$3 25.

In preparing this work a second time for the American profession, the author has spared no labor in giving it a very thorough revision, introducing several new chapters, and rewriting others, while every portion of the volume has been subjected to a severe scrutiny. The efforts of the American editor have been directed to supplying such information relative to matters peculiar to this country as might have escaped the attention of the author, and the whole may, therefore, be safely pronounced one of the most complete works on the subject accessible to the American Profession. By an alteration in the size of the page, these very extensive additions have been accommodated without unduly increasing the size of the work.

BY THE SAME AUTHOR.

ESSAYS ON THE PUERPERAL FEVER, AND OTHER DISEASES PECULIAR TO WOMEN. Selected from the writings of British Authors previous to the close of the Eighteenth Century. In one neat octavo volume, extra cloth, of about 450 pages. \$2 50.

The most popular work on midwifery ever issued from the American press.—*Charleston Med. Journal*.

We were reduced to the necessity of having but one work on midwifery, and permitted to choose, we would unhesitatingly take Churchill.—*Western Med. and Surg. Journal*.

It is impossible to conceive a more useful and elegant manual than Dr. Churchill's Practice of Midwifery.—*Provincial Medical Journal*.

Certainly, in our opinion, the very best work on the subject which exists.—*N. Y. Annalist*.

No work holds a higher position, or is more deserving of being placed in the hands of the tyro, the advanced student, or the practitioner.—*Medical Examiner*.

Previous editions, under the editorial supervision of Prof. R. M. Huston, have been received with marked favor, and they deserved it; but this, reprinted from a very late Dublin edition, carefully revised and brought up by the author to the present time, does present an unusually accurate and able exposition of every important particular embraced in the department of midwifery. * * The clearness, directness, and precision of its teachings, together with the great amount of statistical research which its text exhibits, have served to place it already in the foremost rank of works in this department of medical science.—*N. O. Med. and Surg. Journal*.

In our opinion, it forms one of the best if not the very best text-book and epitome of obstetric science which we at present possess in the English language.—*Monthly Journal of Medical Science*.

The clearness and precision of style in which it is written, and the great amount of statistical research which it contains, have served to place it in the first rank of works in this department of medical science.—*N. Y. Journal of Medicine*.

Few treatises will be found better adapted as a text-book for the student, or as a manual for the frequent consultation of the young practitioner.—*American Medical Journal*.

CHURCHILL (FLEETWOOD), M. D., M. R. I. A., &c.
ON THE DISEASES OF WOMEN; including those of Pregnancy and Child-bed. A new American edition, revised by the Author. With Notes and Additions, by D. FRANCIS CONDIE, M. D., author of "A Practical Treatise on the Diseases of Children." With numerous illustrations. In one large and handsome octavo volume, leather, of 768 pages. \$3 00.

This edition of Dr. Churchill's very popular treatise may almost be termed a new work, so thoroughly has he revised it in every portion. It will be found greatly enlarged, and completely brought up to the most recent condition of the subject, while the very handsome series of illustrations introduced, representing such pathological conditions as can be accurately portrayed, present a novel feature, and afford valuable assistance to the young practitioner. Such additions as appeared desirable for the American student have been made by the editor, Dr. Condie, while a marked improvement in the mechanical execution keeps pace with the advance in all other respects which the volume has undergone, while the price

It comprises, unquestionably, one of the most exact and comprehensive expositions of the present state of medical knowledge in respect to the diseases of women that has yet been published.—*Am. Journ. Med. Sciences.*

This work is the most reliable which we possess on this subject; and is deservedly popular with the profession.—*Charleston Med. Journal*, July, 1857.

We know of no author who deserves that approbation, on "the diseases of females," to the same

extent that Dr. Churchill does. His, indeed, is the only thorough treatise we know of on the subject; and it may be commended to practitioners and students as a masterpiece in its particular department.—*The Western Journal of Medicine and Surgery.*

As a comprehensive manual for students, or a work of reference for practitioners, it surpasses any other that has ever issued on the same subject from the British press.—*Dublin Quart. Journal.*

DICKSON (S. H.), M. D.,
 Professor of Practice of Medicine in the Jefferson Medical College, Philadelphia.

ELEMENTS OF MEDICINE; a Compendious View of Pathology and Therapeutics, or the History and Treatment of Diseases. Second edition, revised. In one large and handsome octavo volume of 750 pages, leather. \$3 75. (Just Issued.)

The steady demand which has so soon exhausted the first edition of this work, sufficiently shows that the author was not mistaken in supposing that a volume of this character was needed—an elementary manual of practice, which should present the leading principles of medicine with the practical results, in a condensed and perspicuous manner. Disencumbered of unnecessary detail and fruitless speculations, it embodies what is most requisite for the student to learn, and at the same time what the active practitioner wants when obliged, in the daily calls of his profession, to refresh his memory on special points. The clear and attractive style of the author renders the whole easy of comprehension, while his long experience gives to his teachings an authority everywhere acknowledged. Few physicians, indeed, have had wider opportunities for observation and experience, and few, perhaps, have used them to better purpose. As the result of a long life devoted to study and practice, the present edition, revised and brought up to the date of publication, will doubtless maintain the reputation already acquired as a condensed and convenient American text-book on the Practice of Medicine.

DRUITT (ROBERT), M. R. C. S., &c.

THE PRINCIPLES AND PRACTICE OF MODERN SURGERY. A new and revised American from the eighth enlarged and improved London edition. Illustrated with four hundred and thirty-two wood-engravings. In one very handsomely printed octavo volume, leather, of nearly 700 large pages. \$3 50. (Just Issued.)

A work which like DRUITT'S SURGERY has for so many years maintained the position of a leading favorite with all classes of the profession, needs no special recommendation to attract attention to a revised edition. It is only necessary to state that the author has spared no pains to keep the work up to its well earned reputation of presenting in a small and convenient compass the latest condition of every department of surgery, considered both as a science and as an art; and that the services of a competent American editor have been employed to introduce whatever novelties may have escaped the author's attention, or may prove of service to the American practitioner. As several editions have appeared in London since the issue of the last American reprint, the volume has had the benefit of repeated revisions by the author, resulting in a very thorough alteration and improvement. The extent of these additions may be estimated from the fact that it now contains about one-third more matter than the previous American edition, and that notwithstanding the adoption of a smaller type, the pages have been increased by about one hundred, while nearly two hundred and fifty wood-cuts have been added to the former list of illustrations.

A marked improvement will also be perceived in the mechanical and artistic execution of the work, which, printed in the best style, on new type, and fine paper, leaves little to be desired as regards external finish; while at the very low price affixed it will be found one of the cheapest volumes accessible to the profession.

This popular volume, now a most comprehensive work on surgery, has undergone many corrections, improvements, and additions, and the principles and the practice of the art have been brought down to the latest record and observation. Of the operations in surgery it is impossible to speak too highly. The descriptions are so clear and concise, and the illustrations so accurate and numerous, that the student can have no difficulty, with instrument in hand, and book by his side, over the dead body, in obtaining a proper knowledge and sufficient tact in this much neglected department of medical education.—*British and Foreign Medico-Chirurg. Review*, Jan. 1860.

In the present edition the author has entirely rewritten many of the chapters, and has incorporated the various improvements and additions in modern surgery. On carefully going over it, we find that

nothing of real practical importance has been omitted; it presents a faithful epitome of everything relating to surgery up to the present hour. It is deservedly a popular manual, both with the student and practitioner.—*London Lancet*, Nov. 19, 1859.

In closing this brief notice, we recommend as cordially as ever this most useful and comprehensive hand-book. It must prove a vast assistance, not only to the student of surgery, but also to the busy practitioner who may not have the leisure to devote himself to the study of more lengthy volumes.—*London Med. Times and Gazette*, Oct. 23, 1859.

In a word, this eighth edition of Dr. Drutt's Manual of Surgery is all that the surgical student or practitioner could desire.—*Dublin Quarterly Journal of Med. Sciences*, Nov. 1859.

DALTON, JR. (J. C.), M. D.

Professor of Physiology in the College of Physicians, New York.

A TREATISE ON HUMAN PHYSIOLOGY, designed for the use of Students and Practitioners of Medicine. Second edition, revised and enlarged, with two hundred and seventy-one illustrations on wood. In one very beautiful octavo volume, of 700 pages, extra cloth, \$4 00; leather, raised bands, \$4 50. (*Just Issued*, 1861.)

The general favor which has so soon exhausted an edition of this work has afforded the author an opportunity in its revision of supplying the deficiencies which existed in the former volume. This has caused the insertion of two new chapters—one on the Special Senses, the other on Imbibition, Exhalation, and the Functions of the Lymphatic System—besides numerous additions of smaller amount scattered through the work, and a general revision designed to bring it thoroughly up to the present condition of the science with regard to all points which may be considered as definitely settled. A number of new illustrations has been introduced, and the work, it is hoped, in its improved form, may continue to command the confidence of those for whose use it is intended.

It will be seen, therefore, that Dr. Dalton's best efforts have been directed towards perfecting his work. The additions are marked by the same features which characterize the remainder of the volume, and render it by far the most desirable textbook on physiology to place in the hands of the student which, so far as we are aware, exists in the English language, or perhaps in any other. We therefore have no hesitation in recommending Dr. Dalton's book for the classes for which it is intended, satisfied as we are that it is better adapted to their use than any other work of the kind to which they have access.—*American Journal of the Medical Sciences*, April, 1861.

It is, therefore, no disparagement to the many books upon physiology, most excellent in their day, to say that Dalton's is the only one that gives us the science as it was known to the best philosophers throughout the world, at the beginning of the current year. It states in comprehensive but concise diction, the facts established by experiment, or other method of demonstration, and details, in an understandable manner, how it is done, but abstains from the discussion of unsettled or theoretical points. Herein it is unique; and these characteristics render it a text-book without a rival, for those who desire to study physiological science as it is known to its most successful cultivators. And it is physiology thus presented that lies at the foundation of correct pathological knowledge; and this in turn is the basis of rational therapeutics; so that pathology, in fact, becomes of prime importance in the proper discharge of our every-day practical duties.—*Cincinnati Lancet*, May, 1861.

Dr. Dalton needs no word of praise from us. He is universally recognized as among the first, if not the very first, of American physiologists now living. The first edition of his admirable work appeared but two years since, and the advance of science, his

own original views and experiments, together with a desire to supply what he considered some deficiencies in the first edition, have already made the present one a necessity, and it will no doubt be even more eagerly sought for than the first. That it is not merely a reprint, will be seen from the author's statement of the following principal additions and alterations which he has made. The present, like the first edition, is printed in the highest style of the printer's art, and the illustrations are truly admirable for their clearness in expressing exactly what their author intended.—*Boston Medical and Surgical Journal*, March 28, 1861.

It is unnecessary to give a detail of the additions; suffice it to say, that they are numerous and important, and such as will render the work still more valuable and acceptable to the profession as a learned and original treatise on this all-important branch of medicine. All that was said in commendation of the getting up of the first edition, and the superior style of the illustrations, apply with equal force to this. No better work on physiology can be placed in the hand of the student.—*St. Louis Medical and Surgical Journal*, May, 1861.

These additions, while testifying to the learning and industry of the author, render the book exceedingly useful, as the most complete exposé of a science, of which Dr. Dalton is doubtless the ablest representative on this side of the Atlantic.—*New Orleans Med. Times*, May, 1861.

A second edition of this deservedly popular work having been called for in the short space of two years, the author has supplied deficiencies, which existed in the former volume, and has thus more completely fulfilled his design of presenting to the profession a reliable and precise text-book, and one which we consider the best outline on the subject of which it treats, in any language.—*N. American Medico-Chirurg. Review*, May, 1861.

DUNGLISON, FORBES, TWEEDIE, AND CONOLLY.

THE CYCLOPÆDIA OF PRACTICAL MEDICINE: comprising Treatises on the Nature and Treatment of Diseases, Materia Medica, and Therapeutics, Diseases of Women and Children, Medical Jurisprudence, &c. &c. In four large super-royal octavo volumes, of 3254 double-columned pages, strongly and handsomely bound, with raised bands. \$12 00.

** This work contains no less than four hundred and eighteen distinct treatises, contributed by sixty-eight distinguished physicians, rendering it a complete library of reference for the country practitioner.

The most complete work on Practical Medicine extant; or, at least, in our language.—*Buffalo Medical and Surgical Journal*.

For reference, it is above all price to every practitioner.—*Western Lancet*.

One of the most valuable medical publications of the day—as a work of reference it is invaluable.—*Western Journal of Medicine and Surgery*.

It has been to us, both as learner and teacher, a work for ready and frequent reference, one in which modern English medicine is exhibited in the most advantageous light.—*Medical Examiner*.

The editors are practitioners of established reputation, and the list of contributors embraces many of the most eminent professors and teachers of London, Edinburgh, Dublin, and Glasgow. It is, indeed, the great merit of this work that the principal articles have been furnished by practitioners who have not only devoted especial attention to the diseases about which they have written, but have also enjoyed opportunities for an extensive practical acquaintance with them and whose reputation carries the assurance of their competency justly to appreciate the opinions of others, while it stamps their own doctrines with high and just authority.—*American Medical Journal*.

DEWEES'S COMPREHENSIVE SYSTEM OF MIDWIFERY. Illustrated by occasional cases and many engravings. Twelfth edition, with the author's last improvements and corrections. In one octavo volume, extra cloth, of 600 pages. \$3 20.

DEWEES'S TREATISE ON THE PHYSICAL

AND MEDICAL TREATMENT OF CHILDREN. The last edition. In one volume, octavo, extra cloth, 548 pages. \$2 50.

DEWEES'S TREATISE ON THE DISEASES OF FEMALES. Tenth edition. In one volume, octavo extra cloth, 532 pages, with plates. \$3 00.

DUNGLISON (ROBLEY), M. D.,

Professor of Institutes of Medicine in the Jefferson Medical College, Philadelphia.

NEW AND ENLARGED EDITION.

MEDICAL LEXICON; a Dictionary of Medical Science, containing a concise

Explanation of the various Subjects and Terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Pharmacy, Obstetrics, Medical Jurisprudence, Dentistry, &c. Notices of Climate and of Mineral Waters; Formule for Official, Empirical, and Dietetic Preparations, &c. With French and other Synonymes. Revised and very greatly enlarged. In one very large and handsome octavo volume, of 992 double-columned pages, in small type; strongly bound in leather, with raised bands. Price \$4 00.

Especial care has been devoted in the preparation of this edition to render it in every respect worthy a continuance of the very remarkable favor which it has hitherto enjoyed. The rapid sale of FIFTEEN large editions, and the constantly increasing demand, show that it is regarded by the profession as the standard authority. Stimulated by this fact, the author has endeavored in the present revision to introduce whatever might be necessary "to make it a satisfactory and desirable—if not indispensable—lexicon, in which the student may search without disappointment for every term that has been legitimated in the nomenclature of the science." To accomplish this, large additions have been found requisite, and the extent of the author's labors may be estimated from the fact that about SIX THOUSAND subjects and terms have been introduced throughout, rendering the whole number of definitions about SIXTY THOUSAND, to accommodate which, the number of pages has been increased by nearly a hundred, notwithstanding an enlargement in the size of the page. The medical press, both in this country and in England, has pronounced the work indispensable to all medical students and practitioners, and the present improved edition will not lose that enviable reputation.

The publishers have endeavored to render the mechanical execution worthy of a volume of such universal use in daily reference. The greatest care has been exercised to obtain the typographical accuracy so necessary in a work of the kind. By the small but exceedingly clear type employed, an immense amount of matter is condensed in its thousand ample pages, while the binding will be found strong and durable. With all these improvements and enlargements, the price has been kept at the former very moderate rate, placing it within the reach of all.

This work, the appearance of the fifteenth edition of which, it has become our duty and pleasure to announce, is perhaps the most stupendous monument of labor and erudition in medical literature. One would hardly suppose after constant use of the preceding editions, where we have never failed to find a sufficiently full explanation of every medical term, that in this edition "*about six thousand subjects and terms have been added*," with a careful revision and correction of the entire work. It is only necessary to announce the advent of this edition to make it occupy the place of the preceding one on the table of every medical man, as it is without doubt the best and most comprehensive work of the kind which has ever appeared.—*Buffalo Med. Journ.*, Jan. 1858.

The work is a monument of patient research, skilful judgment, and vast physical labor, that will perpetuate the name of the author more effectually than any possible device of stone or metal. Dr. Dunglison deserves the thanks not only of the American profession, but of the whole medical world.—*North Am. Medico-Chir. Review*, Jan. 1858.

A Medical Dictionary better adapted for the wants of the profession than any other with which we are acquainted, and of a character which places it far above comparison and competition.—*Am. Journ. Med. Sciences*, Jan. 1858.

We need only say, that the addition of 6,000 new terms, with their accompanying definitions, may be said to constitute a new work, by itself. We have examined the Dictionary attentively, and are most happy to pronounce it unrivalled of its kind. The erudition displayed, and the extraordinary industry which must have been demanded, in its preparation and perfection, redound to the lasting credit of its author, and have furnished us with a volume *indispensable* at the present day, to all who would find themselves *au niveau* with the highest standards of medical information.—*Boston Medical and Surgical Journal*, Dec. 31, 1857.

Good lexicons and encyclopedic works generally, are the most labor-saving contrivances which literary men enjoy; and the labor which is required to produce them in the perfect manner of this example is something appalling to contemplate. The author

tells us in his preface that he has added about six thousand terms and subjects to this edition, which, before, was considered universally as the best work of the kind in any language.—*Silliman's Journal*, March, 1858.

He has razed his gigantic structure to the foundations, and remodelled and reconstructed the entire pile. No less than six thousand additional subjects and terms are illustrated and analyzed in this new edition, swelling the grand aggregate to beyond sixty thousand! Thus is placed before the profession a complete and thorough exponent of medical terminology, without rival or possibility of rivalry.—*Nashville Journ. of Med. and Surg.*, Jan. 1858.

It is universally acknowledged, we believe, that this work is incomparably the best and most complete Medical Lexicon in the English language. The amount of labor which the distinguished author has bestowed upon it is truly wonderful, and the learning and research displayed in its preparation are equally remarkable. Comment and commendation are unnecessary, as no one at the present day thinks of purchasing any other Medical Dictionary than this.—*St. Louis Med. and Surg. Journ.*, Jan. 1858.

It is the foundation stone of a good medical library, and should always be included in the first list of books purchased by the medical student.—*Am. Med. Monthly*, Jan. 1858.

A very perfect work of the kind, undoubtedly the most perfect in the English language.—*Med. and Surg. Reporter*, Jan. 1858.

It is now emphatically the Medical Dictionary of the English language, and for it there is no substitute.—*N. H. Med. Journ.*, Jan. 1858.

It is scarcely necessary to remark that any medical library wanting a copy of Dunglison's Lexicon must be imperfect.—*Cin. Lancet*, Jan. 1858.

We have ever considered it the best authority published, and the present edition we may safely say has no equal in the world.—*Peninsular Med. Journal*, Jan. 1858.

The most complete authority on the subject to be found in any language.—*Va. Med. Journal*, Feb. '56.

BY THE SAME AUTHOR.

THE PRACTICE OF MEDICINE. A Treatise on Special Pathology and Therapeutics. Third Edition. In two large octavo volumes, leather, of 1,500 pages. \$6 25.

DUNGLISON (ROBLEY), M. D.,

Professor of Institutes of Medicine in the Jefferson Medical College, Philadelphia.

HUMAN PHYSIOLOGY. Eighth edition. Thoroughly revised and extensively modified and enlarged, with five hundred and thirty-two illustrations. In two large and handsomely printed octavo volumes, leather, of about 1500 pages. \$7 00.

In revising this work for its eighth appearance, the author has spared no labor to render it worthy a continuance of the very great favor which has been extended to it by the profession. The whole contents have been rearranged, and to a great extent remodelled; the investigations which of late years have been so numerous and so important, have been carefully examined and incorporated, and the work in every respect has been brought up to a level with the present state of the subject. The object of the author has been to render it a concise but comprehensive treatise, containing the whole body of physiological science, to which the student and man of science can at all times refer with the certainty of finding whatever they are in search of, fully presented in all its aspects; and on no former edition has the author bestowed more labor to secure this result.

We believe that it can truly be said, no more complete repertory of facts upon the subject treated, can anywhere be found. The author has, moreover, that enviable tact at description and that facility and ease of expression which render him peculiarly acceptable to the casual, or the studious reader. This facility, so requisite in setting forth many graver and less attractive subjects, lends additional charms to one always fascinating.—*Boston Med. and Surg. Journal*.

The most complete and satisfactory system of Physiology in the English language.—*Amer. Med. Journal*.

The best work of the kind in the English language.—*Silliman's Journal*.

The present edition the author has made a perfect mirror of the science as it is at the present hour. As a work upon physiology proper, the science of the functions performed by the body, the student will find it all he wishes.—*Nashville Journ. of Med.*

That he has succeeded, most admirably succeeded in his purpose, is apparent from the appearance of an eighth edition. It is now the great encyclopædia on the subject, and worthy of a place in every physician's library.—*Western Lancet*.

BY THE SAME AUTHOR. (A new edition.)

GENERAL THERAPEUTICS AND MATERIA MEDICA; adapted for a Medical Text-book. With Indexes of Remedies and of Diseases and their Remedies. SIXTH EDITION, revised and improved. With one hundred and ninety-three illustrations. In two large and handsomely printed octavo vols., leather, of about 1100 pages. \$6 00.

In announcing a new edition of Dr. Dunglison's *General Therapeutics and Materia Medica*, we have no words of commendation to bestow upon a work whose merits have been heretofore so often and so justly extolled. It must not be supposed, however, that the present is a mere reprint of the previous edition; the character of the author for laborious research, judicious analysis, and clearness of expression, is fully sustained by the numerous additions he has made to the work, and the careful revision to which he has subjected the whole.—*N. A. Medico-Chir. Review*, Jan. 1858.

The work will, we have little doubt, be bought and read by the majority of medical students; its size, arrangement, and reliability recommend it to all; no one, we venture to predict, will study it without profit, and there are few to whom it will not be in some measure useful as a work of reference. The young practitioner, more especially, will find the copious indexes appended to this edition of great assistance in the selection and preparation of suitable formulæ.—*Charleston Med. Journ. and Review*, Jan. 1858.

BY THE SAME AUTHOR. (A new Edition.)

NEW REMEDIES, WITH FORMULÆ FOR THEIR PREPARATION AND ADMINISTRATION. Seventh edition, with extensive Additions. In one very large octavo volume, leather, of 770 pages. \$3 75.

Another edition of the "New Remedies" having been called for, the author has endeavored to add everything of moment that has appeared since the publication of the last edition.

The articles treated of in the former editions will be found to have undergone considerable expansion in this, in order that the author might be enabled to introduce, as far as practicable, the results of the subsequent experience of others, as well as of his own observation and reflection; and to make the work still more deserving of the extended circulation with which the preceding editions have been favored by the profession. By an enlargement of the page, the numerous additions have been incorporated without greatly increasing the bulk of the volume.—*Preface*.

One of the most useful of the author's works.—*Southern Medical and Surgical Journal*.

This elaborate and useful volume should be found in every medical library, for as a book of reference, for physicians, it is unsurpassed by any other work in existence, and the double index for diseases and for remedies, will be found greatly to enhance its value.—*New York Med. Gazette*.

The great learning of the author, and his remarkable industry in pushing his researches into every source whence information is derivable, have enabled him to throw together an extensive mass of facts and statements, accompanied by full reference to authorities; which last feature renders the work practically valuable to investigators who desire to examine the original papers.—*The American Journal of Pharmacy*.

ELLIS (BENJAMIN), M. D.

THE MEDICAL FORMULARY; being a Collection of Prescriptions, derived from the writings and practice of many of the most eminent physicians of America and Europe. Together with the usual Dietetic Preparations and Antidotes for Poisons. To which is added an Appendix, on the Endermic use of Medicines, and on the use of Ether and Chloroform. The whole accompanied with a few brief Pharmaceutical and Medical Observations. Eleventh edition, revised and much extended by ROBERT P. THOMAS, M. D., Professor of Materia Medica in the Philadelphia College of Pharmacy. (Preparing.)

ERICHSEN (JOHN),

Professor of Surgery in University College, London, &c.

THE SCIENCE AND ART OF SURGERY; BEING A TREATISE ON SURGICAL INJURIES, DISEASES, AND OPERATIONS. New and improved American, from the second enlarged and carefully revised London edition. Illustrated with over four hundred engravings on wood. In one large and handsome octavo volume, of one thousand closely printed pages. leather, raised bands. \$4 50. (*Just Issued.*)

The very distinguished favor with which this work has been received on both sides of the Atlantic has stimulated the author to render it even more worthy of the position which it has so rapidly attained as a standard authority. Every portion has been carefully revised, numerous additions have been made, and the most watchful care has been exercised to render it a complete exponent of the most advanced condition of surgical science. In this manner the work has been enlarged by about a hundred pages, while the series of engravings has been increased by more than a hundred, rendering it one of the most thoroughly illustrated volumes before the profession. The additions of the author having rendered unnecessary most of the notes of the former American editor, but little has been added in this country; some few notes and occasional illustrations have, however, been introduced to elucidate American modes of practice.

It is, in our humble judgment, decidedly the best book of the kind in the English language. Strange that just such books are no oftener produced by public teachers of surgery in this country and Great Britain. Indeed, it is a matter of great astonishment, but no less true than astonishing, that of the many works on surgery republished in this country within the last fifteen or twenty years as text-books for medical students, this is the only one that even approximates to the fulfilment of the peculiar wants of young men just entering upon the study of this branch of the profession.—*Western Jour. of Med. and Surgery.*

Its value is greatly enhanced by a very copious well-arranged index. We regard this as one of the most valuable contributions to modern surgery. To one entering his novitiate of practice, we regard it the most serviceable guide which he can consult. He will find a fulness of detail leading him through every

step of the operation, and not deserting him until the final issue of the case is decided.—*Sethoscope.*

Embracing, as will be perceived, the whole surgical domain, and each division of itself almost complete and perfect, each chapter full and explicit, each subject faithfully exhibited, we can only express our estimate of it in the aggregate. We consider it an excellent contribution to surgery, as probably the best single volume now extant on the subject, and with great pleasure we add it to our text-books.—*Nashville Journal of Medicine and Surgery.*

Prof. Erichsen's work, for its size, has not been surpassed; his nine hundred and eight pages, profusely illustrated, are rich in physiological, pathological, and operative suggestions, doctrines, details, and processes; and will prove a reliable resource for information, both to physician and surgeon, in the hour of peril.—*N. O. Med. and Surg. Journal.*

FLINT (AUSTIN), M. D.,

Professor of the Theory and Practice of Medicine in the University of Louisville, &c.

PHYSICAL EXPLORATION AND DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS. In one large and handsome octavo volume, extra cloth, 636 pages. \$3 00.

We regard it, in point both of arrangement and of the marked ability of its treatment of the subjects, as destined to take the first rank in works of this class. So far as our information extends, it has at present no equal. To the practitioner, as well as the student, it will be invaluable in clearing up the diagnosis of doubtful cases, and in shedding light upon difficult phenomena.—*Buffalo Med. Journal.*

A work of original observation of the highest merit. We recommend the treatise to every one who wishes to become a correct auscultator. Based to a very large extent upon cases numerically examined, it carries the evidence of careful study and discrimination upon every page. It does credit to the author, and, through him, to the profession in this country. It is, what we cannot call every book upon auscultation, a readable book.—*Am. Jour. Med. Sciences.*

BY THE SAME AUTHOR. (*Now Ready.*)

A PRACTICAL TREATISE ON THE DIAGNOSIS, PATHOLOGY, AND TREATMENT OF DISEASES OF THE HEART. In one neat octavo volume, of about 500 pages, extra cloth. \$2 75.

We do not know that Dr. Flint has written anything which is not first rate; but this, his latest contribution to medical literature, in our opinion, surpasses all the others. The work is most comprehensive in its scope, and most sound in the views it enunciates. The descriptions are clear and methodical; the statements are substantiated by facts, and are made with such simplicity and sincerity, that without them they would carry conviction. The style is admirably clear, direct, and free from dryness. With Dr. Walshe's excellent treatise before us, we have no hesitation in saying that Dr. Flint's book is the best work on the heart in the English language.—*Boston Med. and Surg. Journal.*

We have thus endeavored to present our readers with a fair analysis of this remarkable work. Preferring to employ the very words of the distinguished author, wherever it was possible, we have essayed to condense into the briefest space a general view of his observations and suggestions, and to direct the attention of our brethren to the abounding stores of valuable matter here collected and arranged for their use and instruction. No medical library will hereafter be considered complete without this volume; and we trust it will promptly find its way into the hands of every American student and physician.—*N. Am. Med. Chir. Review.*

This last work of Prof. Flint will add much to his previous well-earned celebrity, as a writer of

great force and beauty, and, with his previous work, places him at the head of American writers upon diseases of the chest. We have adopted his work upon the heart as a text-book, believing it to be more valuable for that purpose than any work of the kind that has yet appeared.—*Nashville Med. Journ.*

With more than pleasure do we hail the advent of this work, for it fills a wide gap on the list of text-books for our schools, and is, for the practitioner, the most valuable practical work of its kind.—*N. O. Med. News.*

In regard to the merits of the work, we have no hesitation in pronouncing it full, accurate, and judicious. Considering the present state of science, such a work was much needed. It should be in the hands of every practitioner.—*Chicago Med. Journal.*

But these are very trivial spots, and in no wise prevent us from declaring our most hearty approval of the author's ability, industry, and conscientiousness.—*Dublin Quarterly Journal of Med. Sciences.*

He has labored on with the same industry and care, and his place among the first authors of our country is becoming fully established. To this end, the work whose title is given above, contributes in no small degree. Our space will not admit of an extended analysis, and we will close this brief notice by commending it without reserve to every class of readers in the profession.—*Peninsular Med. Journ.*

FOWNES (GEORGE), PH. D., &c.

A MANUAL OF ELEMENTARY CHEMISTRY; Theoretical and Practical.

From the seventh revised and corrected London edition. With one hundred and ninety-seven illustrations. Edited by ROBERT BRIDGES, M. D. In one large royal 12mo. volume, of 600 pages. In leather, \$1 65; extra cloth, \$1 50. (*Just Issued.*)

The death of the author having placed the editorial care of this work in the practised hands of Drs. Bence Jones and A. W. Hoffman, everything has been done in its revision which experience could suggest to keep it on a level with the rapid advance of chemical science. The additions requisite to this purpose have necessitated an enlargement of the page, notwithstanding which the work has been increased by about fifty pages. At the same time every care has been used to maintain its distinctive character as a condensed manual for the student, divested of all unnecessary detail or mere theoretical speculation. The additions have, of course, been mainly in the department of Organic Chemistry, which has made such rapid progress within the last few years, but yet equal attention has been bestowed on the other branches of the subject—Chemical Physics and Inorganic Chemistry—to present all investigations and discoveries of importance, and to keep up the reputation of the volume as a complete manual of the whole science, admirably adapted for the learner. By the use of a small but exceedingly clear type the matter of a large octavo is compressed within the convenient and portable limits of a moderate sized duodecimo, and at the very low price affixed, it is offered as one of the cheapest volumes before the profession.

Dr. Fownes' excellent work has been universally recognized everywhere in his own and this country, as the best elementary treatise on chemistry in the English tongue, and is very generally adopted, we believe, as the standard text-book in all our colleges, both literary and scientific.—*Charleston Med. Journ. and Review.*

A standard manual, which has long enjoyed the reputation of embodying much knowledge in a small space. The author has achieved the difficult task of condensation with masterly tact. His book is concise without being dry, and brief without being too dogmatical or general.—*Virginia Med. and Surgical Journal.*

The work of Dr. Fownes has long been before the public, and its merits have been fully appreciated as the best text-book on chemistry now in existence. We do not, of course, place it in a rank superior to the works of Brande, Graham, Turner, Gregory, or Gmelin, but we say that, as a work for students, it is preferable to any of them.—*London Journal of Medicine.*

A work well adapted to the wants of the student. It is an excellent exposition of the chief doctrines and facts of modern chemistry. The size of the work, and still more the condensed yet perspicuous style in which it is written, absolve it from the charges very properly urged against most manuals termed popular.—*Edinburgh Journal of Medical Sciences*

FISKE FUND PRIZE ESSAYS—THE EFFECTS OF CLIMATE ON TUBERCULOUS DISEASE. By EDWIN LEE, M. R. C. S., London, and THE INFLUENCE OF PREGNANCY ON THE DEVELOPMENT OF TUBERCLES By

EDWARD WARREN, M. D., of Edenton, N. C. Together in one neat 8vo. volume, extra cloth, \$1 00. **FRICK ON RENAL AFFECTIONS;** their Diagnosis and Pathology. With illustrations. One volume, royal 12mo., extra cloth. 75 cents

FERGUSON (WILLIAM), F. R. S.,

Professor of Surgery in King's College, London, &c.

A SYSTEM OF PRACTICAL SURGERY. Fourth American, from the third and enlarged London edition. In one large and beautifully printed octavo volume, of about 700 pages, with 393 handsome illustrations, leather. \$3 00.

GRAHAM (THOMAS), F. R. S.

THE ELEMENTS OF INORGANIC CHEMISTRY, including the Applications of the Science in the Arts. New and much enlarged edition, by HENRY WATTS and ROBERT BRIDGES, M. D. Complete in one large and handsome octavo volume, of over 800 very large pages, with two hundred and thirty-two wood-cuts, extra cloth. \$4 00.

*** Part II., completing the work from p. 431 to end, with Index, Title Matter, &c., may be had separate, cloth backs and paper sides. Price \$2 50.

From Prof. E. N. Horsford, *Harvard College.*

It has, in its earlier and less perfect editions, been familiar to me, and the excellence of its plan and the clearness and completeness of its discussions, have long been my admiration.

No reader of English works on this science can

afford to be without this edition of Prof. Graham's Elements.—*Silliman's Journal*, March, 1858.

From Prof. Wolcott Gibbs, *N. Y. Free Academy.*

The work is an admirable one in all respects, and its republication here cannot fail to exert a positive influence upon the progress of science in this country.

GRIFFITH (ROBERT E.), M. D., &c.

A UNIVERSAL FORMULARY, containing the methods of Preparing and Administering Official and other Medicines. The whole adapted to Physicians and Pharmacutists. SECOND EDITION, thoroughly revised, with numerous additions, by ROBERT P. THOMAS, M. D., Professor of Materia Medica in the Philadelphia College of Pharmacy. In one large and handsome octavo volume, extra cloth, of 650 pages, double columns. \$3 00; or in sheep, \$3 25.

It was a work requiring much perseverance, and when published was looked upon as by far the best work of its kind that had issued from the American press. Prof. Thomas has certainly "improved," as press. Prof. Thomas has certainly "improved," as well as added to this Formulary, and has rendered it additionally deserving of the confidence of pharmacutists and physicians.—*Am. Journal of Pharmacy.*

We are happy to announce a new and improved edition of this one of the most valuable and useful works that have emanated from an American pen. It would do credit to practitioners of medicine; it is of daily usefulness to practitioners of medicine; it is better adapted to their purposes than the dispensatories.—*Southern Med. and Surg. Journal.*

It is one of the most useful books a country practitioner can possibly have.—*Medical Chronicle.*

This is a work of six hundred and fifty-one pages, embracing all on the subject of preparing and administering medicines that can be desired by the physician and pharmacist.—*Western Lancet.*

The amount of useful, every-day matter for a practicing physician, is really immense.—*Boston Med. and Surg. Journal.*

This edition has been greatly improved by the revision and ample additions of Dr. Thomas, and is now, we believe, one of the most complete works of its kind in any language. The additions amount to about seventy pages, and no effort has been spared to include in them all the recent improvements. A work of this kind appears to us indispensable to the physician, and there is none we can more cordially recommend. *N Y Journal of Medicine.*

GROSS (SAMUEL D.), M. D.,

Professor of Surgery in the Jefferson Medical College of Philadelphia, &c.

Enlarged Edition—Now Ready, January, 1862.

A SYSTEM OF SURGERY: Pathological, Diagnostic, Therapeutic, and Operative. Illustrated by TWELVE HUNDRED AND TWENTY-SEVEN ENGRAVINGS. Second edition, much enlarged and carefully revised. In two large and beautifully printed octavo volumes, of about twenty-two hundred pages; strongly bound in leather, with raised bands. Price \$12.

The exhaustion in little more than two years of a large edition of so elaborate and comprehensive a work as this is the best evidence that the author was not mistaken in his estimate of the want which existed of a complete American System of Surgery, presenting the science in all its necessary details and in all its branches. That he has succeeded in the attempt to supply this want is shown not only by the rapid sale of the work, but also by the very favorable manner in which it has been received by the organs of the profession in this country and in Europe, and by the fact that a translation is now preparing in Holland—a mark of appreciation not often bestowed on any scientific work so extended in size.

The author has not been insensible to the kindness thus bestowed upon his labors, and in revising the work for a new edition he has spared no pains to render it worthy of the favor with which it has been received. Every portion has been subjected to close examination and revision; any deficiencies apparent have been supplied, and the results of recent progress in the science and art of surgery have been everywhere introduced; while the series of illustrations has been enlarged by the addition of nearly three hundred wood-cuts, rendering it one of the most thoroughly illustrated works ever laid before the profession. To accommodate these very extensive additions, the work has been printed upon a smaller type, so that notwithstanding the very large increase in the matter and value of the book, its size is more convenient and less cumbersome than before. Every care has been taken in the printing to render the typographical execution unexceptionable, and it is confidently presented as a work in every way worthy of a place in even the most limited library of the practitioner or student.

A few testimonials of the value of the former edition are appended.

Has Dr. Gross satisfactorily fulfilled this object?

A careful perusal of his volumes enables us to give an answer in the affirmative. Not only has he given to the reader an elaborate and well-written account of his own vast experience, but he has not failed to embody in his pages the opinions and practice of surgeons in this and other countries of Europe. The result has been a work of such completeness, that it has no superior in the systematic treatises on surgery which have emanated from English or Continental authors. It has been justly objected that these have been far from complete in many essential particulars, many of them having been deficient in some of the most important points which should characterize such works. Some of them have been elaborate—too elaborate—with respect to certain diseases, while they have merely glanced at, or given an unsatisfactory account of, others equally important to the surgeon. Dr. Gross has avoided this error, and has produced the most complete work that has yet issued from the press on the science and practice of surgery. It is not, strictly speaking, a Dictionary of Surgery, but it gives to the reader all the information that he may require for his treatment of surgical diseases. Having said so much, it might appear superfluous to add another word; but it is only due to Dr. Gross to state that he has embraced the opportunity of transferring to his pages a vast number of engravings from English and other authors, illustrative of the pathology and treatment of surgical diseases. To these are added several hundred original wood-cuts. The work altogether commands itself to the attention of British surgeons, from whom it cannot fail to meet with extensive patronage.—*London Lancet*, Sept. 1, 1860.

Of Dr. Gross's treatise on Surgery we can say no more than that it is the most elaborate and complete work on this branch of the healing art which has ever been published in any country. A systematic work, it admits of no analytical review; but, did our space permit, we should gladly give some extracts from it, to enable our readers to judge of the classical style of the author, and the exhausting way in which each subject is treated.—*Dublin Quarterly Journal of Med. Science*.

The work is so superior to its predecessors in matter and extent, as well as in illustrations and style of publication, that we can honestly recommend it as the best work of the kind to be taken home by the young practitioner.—*Am. Med. Journ.*

With pleasure we record the completion of this long-anticipated work. The reputation which the author has for many years sustained, both as a surgeon and as a writer, had prepared us to expect a treatise of great excellence and originality; but we confess we were by no means prepared for the work which is before us—the most complete treatise upon surgery ever published, either in this or any other country, and we might, perhaps, safely say, the most original. There is no subject belonging properly to surgery which has not received from the author a due share of attention. Dr. Gross has supplied a want in surgical literature which has long been felt by practitioners; he has furnished us with a complete practical treatise upon surgery in all its departments. As Americans, we are proud of the achievement; as surgeons, we are most sincerely thankful to him for his extraordinary labors in our behalf.—*N. Y. Monthly Review and Buffalo Med. Journal*.

BY THE SAME AUTHOR.

ELEMENTS OF PATHOLOGICAL ANATOMY. Third edition, thoroughly revised and greatly improved. In one large and very handsome octavo volume, with about three hundred and fifty beautiful illustrations, of which a large number are from original drawings. Price in extra cloth, \$4 75; leather, raised bands, \$5 25. (*Lately Published.*)

The very rapid advances in the Science of Pathological Anatomy during the last few years have rendered essential a thorough modification of this work, with a view of making it a correct exponent of the present state of the subject. The very careful manner in which this task has been executed, and the amount of alteration which it has undergone, have enabled the author to say that "with the many changes and improvements now introduced, the work may be regarded almost as a new treatise," while the efforts of the author have been seconded as regards the mechanical execution of the volume, rendering it one of the handsomest productions of the American press.

We most sincerely congratulate the author on the successful manner in which he has accomplished his proposed object. His book is most admirably calculated to fill up a blank which has long been felt to exist in this department of medical literature, and as such must become very widely circulated amongst all classes of the profession.—*Dublin Quarterly Journ. of Med. Science*, Nov. 1857.

We have been favorably impressed with the general manner in which Dr. Gross has executed his task of affording a comprehensive digest of the present state of the literature of Pathological Anatomy, and have much pleasure in recommending his work to our readers, as we believe one well deserving of diligent perusal and careful study.—*Montreal Med. Chron.*, Sept. 1857.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON FOREIGN BODIES IN THE AIR-PASSES. In one handsome octavo volume, extra cloth, with illustrations.

GROSS (SAMUEL D.), M. D.

Professor of Surgery in the Jefferson Medical College of Philadelphia, &c.

A PRACTICAL TREATISE ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA. Second Edition, revised and much enlarged, with one hundred and eighty-four illustrations. In one large and very handsome octavo volume, of over nine hundred pages. In leather, raised bands, \$5 25; extra cloth, \$4 75.

Philosophical in its design, methodical in its arrangement, ample and sound in its practical details, it may in truth be said to leave scarcely anything to be desired on so important a subject.—*Boston Med. and Surg. Journal*.

Whoever will peruse the vast amount of valuable practical information it contains, will, we think,

agree with us, that there is no work in the English language which can make any just pretensions to be its equal.—*N. Y. Journal of Medicine*.

A volume replete with truths and principles of the utmost value in the investigation of these diseases.—*American Medical Journal*.

GRAY (HENRY), F. R. S.

Lecturer on Anatomy at St. George's Hospital, London, &c.

ANATOMY, DESCRIPTIVE AND SURGICAL. The Drawings by H. V. CARTER, M. D., late Demonstrator on Anatomy at St. George's Hospital; the Dissections jointly by the AUTHOR and Dr. CARTER. Second American, from the second revised and improved London edition. In one magnificent imperial octavo volume, of over 800 pages, with 388 large and elaborate engravings on wood. Price in extra cloth, \$6 25; leather, raised bands, \$7 00. (*Now Ready*, 1862.)

The speedy exhaustion of a large edition of this work is sufficient evidence that its plan and execution have been found to present superior practical advantages in facilitating the study of Anatomy. In presenting it to the profession a second time, the author has availed himself of the opportunity to supply any deficiencies which experience in its use had shown to exist, and to correct any errors of detail, to which the first edition of a scientific work on so extensive and complicated a science is liable. These improvements have resulted in some increase in the size of the volume, while twenty-six new wood-cuts have been added to the beautiful series of illustrations which form so distinctive a feature of the work. The American edition has been passed through the press under the supervision of a competent professional man, who has taken every care to render it in all respects accurate, and it is now presented, without any increase of price, as fitted to maintain and extend the popularity which it has everywhere acquired.

With little trouble, the busy practitioner whose knowledge of anatomy may have become obscured by want of practice, may now resuscitate his former anatomical lore, and be ready for any emergency. It is to this class of individuals, and not to the student alone, that this work will ultimately tend to be of most incalculable advantage, and we feel satisfied that the library of the medical man will soon be considered incomplete in which a copy of this work does not exist.—*Madras Quarterly Journal of Med. Science*, July, 1861.

This edition is much improved and enlarged, and contains several new illustrations by Dr. Westmacott. The volume is a complete companion to the dissecting-room, and saves the necessity of the student possessing a variety of "Manuals."—*The London Lancet*, Feb. 9, 1861.

The work before us is one entitled to the highest praise, and we accordingly welcome it as a valuable addition to medical literature. Intermediate in fulness of detail between the treatises of Saarpay and of Wilson, its characteristic merit lies in the number and excellence of the engravings it contains. Most of these are original, of much larger than ordinary size, and admirably executed. The various parts are also lettered after the plan adopted in Holden's Osteology. It would be difficult to over-estimate the advantages offered by this mode of pictorial illustration. Bones, ligaments, muscles, bloodvessels, and nerves are each in turn figured, and marked with their appropriate names; thus enabling the student to comprehend, at a glance, what would otherwise often be ignored, or at any rate, acquired only by prolonged and irksome application. In conclusion, we heartily commend the work of Mr. Gray to the attention of the medical profession, feeling certain that it should be regarded as one of the most valuable contributions ever made to educational literature.—*N. Y. Monthly Review*. Dec. 1859.

In this view, we regard the work of Mr. Gray as far better adapted to the wants of the profession, and especially of the student, than any treatise on anatomy yet published in this country. It is destined, we believe, to supersede all others, both as a manual of dissections, and a standard of reference to the student of general or relative anatomy.—*N. Y. Journal of Medicine*, Nov. 1859.

For this truly admirable work the profession is indebted to the distinguished author of "Gray on the Spleen." The vacancy it fills has been long felt

to exist in this country. Mr. Gray writes throughout with both branches of his subject in view. His description of each particular part is followed by a notice of its relations to the parts with which it is connected, and this, too, sufficiently ample for all the purposes of the operative surgeon. After describing the bones and muscles, he gives a concise statement of the fractures to which the bones of the extremities are most liable, together with the amount and direction of the displacement to which the fragments are subjected by muscular action. The section on arteries is remarkably full and accurate. Not only is the surgical anatomy given to every important vessel, with directions for its ligation, but at the end of the description of each arterial trunk we have a useful summary of the irregularities which may occur in its origin, course, and termination.—*N. A. Med. Chir. Review*, Mar. 1859.

Mr. Gray's book, in excellency of arrangement and completeness of execution, exceeds any work on anatomy hitherto published in the English language, affording a complete view of the structure of the human body, with especial reference to practical surgery. Thus the volume constitutes a perfect book of reference for the practitioner, demanding a place in even the most limited library of the physician or surgeon, and a work of necessity for the student to fix in his mind what he has learned by the dissecting knife from the book of nature.—*The Dublin Quarterly Journal of Med. Sciences*, Nov. 1858.

In our judgment, the mode of illustration adopted in the present volume cannot but present many advantages to the student of anatomy. To the zealous disciple of Vesalius, earnestly desirous of real improvement, the book will certainly be of immense value; but, at the same time, we must also confess that to those simply desirous of "cramming" it will be an undoubted godsend. The peculiar value of Mr. Gray's mode of illustration is nowhere more markedly evident than in the chapter on osteology, and especially in those portions which treat of the bones of the head and of their development. The study of these parts is thus made one of comparative ease, if not of positive pleasure; and those bugbears of the student, the temporal and sphenoid bones, are shorn of half their terrors. It is, in our estimation, an admirable and complete text-book for the student, and a useful work of reference for the practitioner; its pictorial character forming a novel element, to which we have already sufficiently alluded.—*Am. Journ. Med. Sci.*, July, 1859.

GIBSON'S INSTITUTES AND PRACTICE OF SURGERY. Eighth edition, improved and altered. With thirty-four plates. In two handsome octavo volumes, containing about 1,000 pages, leather, raised bands. \$6 50

GARDNER'S MEDICAL CHEMISTRY, for the use of Students and the Profession. In one royal 12mo. vol., cloth, pp. 396, with wood-cuts. \$1.

GLUGE'S ATLAS OF PATHOLOGICAL HISTOLOGY. Translated, with Notes and Addi-

tions by JOSEPH LEIDY, M. D. In one volume, very large imperial quarto, extra cloth, with 320 copper plate figures, plain and colored, \$5 00.

HUGHES' INTRODUCTION TO THE PRACTICE OF AUSCULTATION AND OTHER MODES OF PHYSICAL DIAGNOSIS IN DISEASES OF THE LUNGS AND HEART. Second edition 1 vol. royal 12mo., ex. cloth, pp. 304. \$1 00.

HAMILTON (FRANK H.), M. D.,

Professor of Surgery in the Long Island College Hospital.

A PRACTICAL TREATISE ON FRACTURES AND DISLOCATIONS. In one large and handsome octavo volume, of over 750 pages, with 289 illustrations. \$4 25. (*New Ready*, January, 1860.)

Among the many good workers at surgery of whom America may now boast not the least is Frank Hastings Hamilton; and the volume before us (we say it with a pang of wounded patriotism) the best and handiest book on the subject in the English language. It is in vain to attempt a review of it; nearly as vain to seek for any sins, either of commission or omission. We have seen no work on practical surgery which we would sooner recommend to our brother surgeons, especially those of "the services," or those whose practice lies in districts where a man has necessarily to rely on his own unaided resources. The practitioner will find in it directions for nearly every possible accident, easily found and comprehended; and much pleasant reading for him to muse over in the after consideration of his cases.—*Edinburgh Med. Journ.* Feb. 1861.

This is a valuable contribution to the surgery of most important affections, and is the more welcome, inasmuch as at the present time we do not possess a single complete treatise on Fractures and Dislocations in the English language. It has remained for our American brother to produce a complete treatise upon the subject, and bring together in a convenient form those alterations and improvements that have been made from time to time in the treatment of these affections. One great and valuable feature in the work before us is the fact that it comprises all the improvements introduced into the practice of both English and American surgery, and though far from omitting mention of our continental neighbors, the author by no means encourages the notion—but too prevalent in some quarters—that nothing is good unless imported from France or Germany. The latter half of the work is devoted to the consideration of the various dislocations and their appropriate treatment, and its merit is fully equal to that of the preceding portion.—*The London Lancet*, May 5, 1860.

It is emphatically the book upon the subjects of which it treats, and we cannot doubt that it will continue so to be for an indefinite period of time. When we say, however, that we believe it will at once take its place as the best book for consultation by the practitioner; and that it will form the most complete, available, and reliable guide in emergencies of every nature connected with its subjects; and also that the student of surgery may make it his textbook with entire confidence, and with pleasure also, from its agreeable and easy style—we think our own

opinion may be gathered as to its value.—*Boston Medical and Surgical Journal*, March 1, 1860.

The work is concise, judicious, and accurate, and adapted to the wants of the student, practitioner, and investigator, honorable to the author and to the profession.—*Chicago Med. Journal*, March, 1860.

We regard this work as an honor not only to its author, but to the profession of our country. Were we to review it thoroughly, we could not convey to the mind of the reader more forcibly our honest opinion expressed in the few words—we think it the best book of its kind extant. Every man interested in surgery will soon have this work on his desk. He who does not, will be the loser.—*New Orleans Medical News*, March, 1860.

Now that it is before us, we feel bound to say that much as was expected from it, and onerous as was the undertaking, it has surpassed expectation, and achieved more than was pledged in its behalf; for its title does not express in full the richness of its contents. On the whole, we are prouder of this work than of any which has for years emanated from the American medical press; its sale will certainly be very large in this country, and we anticipate its eliciting much attention in Europe.—*Nashville Medical Record*, Mar. 1860.

Every surgeon, young and old, should possess himself of it, and give it a careful perusal, in doing which he will be richly repaid.—*St. Louis Med. and Surg. Journal*, March, 1860.

Dr. Hamilton is fortunate in having succeeded in filling the void, so long felt, with what cannot fail to be at once accepted as a model monograph in some respects, and a work of classical authority. We sincerely congratulate the profession of the United States on the appearance of such a publication from one of their number. We have reason to be proud of it as an original work, both in a literary and scientific point of view, and to esteem it as a valuable guide in a most difficult and important branch of study and practice. On every account, therefore, we hope that it may soon be widely known abroad as an evidence of genuine progress on this side of the Atlantic, and further, that it may be still more widely known at home as an authoritative teacher from which every one may profitably learn, and as affording an example of honest, well-directed, and untiring industry in authorship which every surgeon may emulate.—*Am. Med. Journal*, April, 1860.

HOBLYN (RICHARD D.), M. D.

A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. A new American edition. Revised, with numerous Additions, by ISAAC HAYS, M. D., editor of the "American Journal of the Medical Sciences." In one large royal 12mo. volume, leather, of over 500 double columned pages. \$1 50.

To both practitioner and student, we recommend this dictionary as being convenient in size, accurate in definition, and sufficiently full and complete for ordinary consultation.—*Charleston Med. Journ.*

We know of no dictionary better arranged and adapted. It is not encumbered with the obsolete terms of a bygone age, but it contains all that are now in

use; embracing every department of medical science down to the very latest date.—*Western Lancet*.

Hoblyn's Dictionary has long been a favorite with us. It is the best book of definitions we have, and ought always to be upon the student's table.—*Southern Med. and Surg. Journal*.

HOLLAND'S MEDICAL NOTES AND REFLECTIONS. From the third London edition. In one handsome octavo volume, extra cloth. \$3.

HORNER'S SPECIAL ANATOMY AND HIS-

TOLOGY. Eighth edition. Extensively revised and modified. In two large octavo volumes, extra cloth, of more than 1000 pages, with over 300 illustrations. \$6 00.

HODGE (HUGH L.), M. D.,

Professor of Midwifery and the Diseases of Women and Children in the University of Pennsylvania, &c.

ON DISEASES PECULIAR TO WOMEN, including Displacements of the Uterus. With original illustrations. In one beautifully printed octavo volume, of nearly 500 pages, extra cloth. \$3 25. (*Now Ready.*)

We will say at once that the work fulfils its object capably well; and we will moreover venture the assertion that it will inaugurate an improved practice throughout this whole country. The secrets of the author's success are so clearly revealed that the attentive student cannot fail to insure a goodly portion of similar success in his own practice. It is a credit to *all* medical literature; and we add, that the physician who does not place it in his library, and who does not faithfully con its pages, will lose a vast deal of knowledge that would be most useful to himself and beneficial to his patients. *It is a practical work of the highest order of merit; and it will take rank as such immediately.*—*Maryland and Virginia Medical Journal*, Feb. 1861.

This contribution towards the elucidation of the pathology and treatment of some of the diseases peculiar to women, cannot fail to meet with a favorable reception from the medical profession. The character of the particular maladies of which the work before us treats; their frequency, variety, and obscurity; the amount of malaise and even of actual suffering by which they are invariably attended; their obstinacy, the difficulty with which they are overcome, and their disposition again and again to recur—these, taken in connection with the entire competency of the author to render a correct account of their nature, their causes, and their appro-

priate management—his ample experience, his matured judgment, and his perfect conscientiousness—invest this publication with an interest and value to which few of the medical treatises of a recent date can lay a stronger, if, perchance, an equal claim.—*Am. Journ. Med. Sciences*, Jan. 1861.

Indeed, although no part of the volume is not eminently deserving of perusal and study, we think that the nine chapters devoted to this subject, are especially so, and we know of no more valuable monograph upon the symptoms, prognosis, and management of these annoying maladies than is constituted by this part of the work. We cannot but regard it as one of the most original and most practical works of the day; one which every accoucheur and physician should most carefully read; for we are persuaded that he will arise from its perusal with new ideas, which will induce him into a more rational practice in regard to many a suffering female, who may have placed her health in his hands.—*British American Journal*, Feb. 1861.

Of the many excellences of the work we will not speak at length. We advise all who would acquire a knowledge of the proper management of the maladies of which it treats, to study it with care. The second part is of itself a most valuable contribution to the practice of our art.—*Am. Med. Monthly and New York Review*, Feb. 1861.

The illustrations, which are all original, are drawn to a uniform scale of one-half the natural size.

HABERSHON (S. O.), M. D.,

Assistant Physician to and Lecturer on Materia Medica and Therapeutics at Guy's Hospital, &c.

PATHOLOGICAL AND PRACTICAL OBSERVATIONS ON DISEASES OF THE ALIMENTARY CANAL, ŒSOPHAGUS, STOMACH, CÆCUM, AND INTES-TINES. With illustrations on wood. In one handsome octavo volume of 312 pages, extra cloth \$1 75. (*Now Ready.*)

JONES (T. WHARTON), F. R. S.,

Professor of Ophthalmic Medicine and Surgery in University College, London, &c.

THE PRINCIPLES AND PRACTICE OF OPHTHALMIC MEDICINE AND SURGERY. With one hundred and ten illustrations. Second American from the second and revised London edition, with additions by EDWARD HARTSHORNE, M. D., Surgeon to Wills' Hospital, &c. In one large, handsome royal 12mo. volume, extra cloth, of 500 pages. \$1 50.

JONES (C. HANDFIELD), F. R. S., & EDWARD H. SIEVEKING, M. D.,
Assistant Physicians and Lecturers in St. Mary's Hospital, London.

A MANUAL OF PATHOLOGICAL ANATOMY. First American Edition, Revised. With three hundred and ninety-seven handsome wood engravings. In one large and beautiful octavo volume of nearly 750 pages, leather. \$3 75.

As a concise text-book, containing, in a condensed form, a complete outline of what is known in the domain of Pathological Anatomy, it is perhaps the best work in the English language. Its great merit consists in its completeness and brevity, and in this respect it supplies a great desideratum in our literature. Heretofore the student of pathology was

obliged to glean from a great number of monographs, and the field was so extensive that but few cultivated it with any degree of success. As a simple work of reference, therefore, it is of great value to the student of pathological anatomy, and should be in every physician's library.—*Western Lancet*.

KIRKES (WILLIAM SENHOUSE), M. D.,

Demonstrator of Morbid Anatomy at St. Bartholomew's Hospital, &c.

A MANUAL OF PHYSIOLOGY. A new American, from the third and improved London edition. With two hundred illustrations. In one large and handsome royal 12mo. volume, leather. pp. 586. \$2 00. (*Lately Published.*)

This is a new and very much improved edition of Dr. Kirkes' well-known Handbook of Physiology. It combines conciseness with completeness, and is, therefore, admirably adapted for consultation by the busy practitioner.—*Dublin Quarterly Journal*.

One of the very best handbooks of Physiology we possess—presenting just such an outline of the science as the student requires during his attendance upon a course of lectures, or for reference whilst preparing for examination.—*Am. Medical Journal*
Its excellence is in its compactness, its clearness,

and its carefully cited authorities. It is the most convenient of text-books. These gentlemen, Messrs. Kirkes and Paget, have the gift of telling us what we want to know, without thinking it necessary to tell us all they know.—*Boston Med and Surg. Journal*.

For the student beginning this study, and the practitioner who has but leisure to refresh his memory, this book is invaluable, as it contains all that it is important to know.—*Charleston Med. Journal*.

KNAPP'S TECHNOLOGY; or, Chemistry applied to the Arts and to Manufactures. Edited by Dr. RONALDS, Dr. RICHARDSON, and Prof. W. R. JOHNSON. In two handsome 8vo. vols., without 500 wood-engravings. \$6 00.

LAYCOCK'S LECTURES ON THE PRINCIPLES AND METHODS OF MEDICAL OBSERVATION AND RESEARCH. For the Use of Advanced Students and Junior Practitioners. In one royal 12mo. volume, extra cloth. Price \$1.

LALLEMAND AND WILSON.

A PRACTICAL TREATISE ON THE CAUSES, SYMPTOMS, AND TREATMENT OF SPERMATORRHOEA. By M. LALLEMAND. Translated and edited by HENRY J. McDUGALL. Third American edition. To which is added — ON DISEASES OF THE VESICULÆ SEMINALES; AND THEIR ASSOCIATED ORGANS. With special reference to the Morbid Secretions of the Prostatic and Urethral Mucous Membrane. By MARRIS WILSON, M. D. In one neat octavo volume, of about 400 pp., extra cloth. \$2 00. (*Just Issued.*)

LA ROCHE (R.), M. D., &c.

YELLOW FEVER, considered in its Historical, Pathological, Etiological, and Therapeutical Relations. Including a Sketch of the Disease as it has occurred in Philadelphia from 1699 to 1854, with an examination of the connections between it and the fevers known under the same name in other parts of temperate as well as in tropical regions. In two large and handsome octavo volumes of nearly 1500 pages, extra cloth. \$7 00.

From Professor S. H. Dickson, Charleston, S. C.,
September 18, 1855.

A monument of intelligent and well applied research, almost without example. It is, indeed, in itself, a large library, and is destined to constitute the special record as a book of reference, in the subject of which it treats, to all future time.

We have not time at present, engaged as we are, by day and by night, in the work of combating this very disease, now prevailing in our city, to do more than give this cursory notice of what we consider as undoubtedly the most able and erudite medical publication our country has yet produced. But in view of the startling fact, that this, the most malign-

ant and unmanageable disease of modern times, has for several years been prevailing in our country to a greater extent than ever before; that it is no longer confined to either large or small cities, but penetrates country villages, plantations, and farm-houses; that it is treated with scarcely better success now than thirty or forty years ago; that there is vast mischief done by ignorant pretenders to knowledge in regard to the disease, and in view of the probability that a majority of southern physicians will be called upon to treat the disease, we trust that this able and comprehensive treatise will be very generally read in the south.—*Memphis Med. Recorder.*

BY THE SAME AUTHOR.

PNEUMONIA; its Supposed Connection, Pathological and Etiological, with Autumnal Fevers, including an Inquiry into the Existence and Morbid Agency of Malaria. In one handsome octavo volume, extra cloth, of 500 pages. \$3 00.

LAWRENCE (W.), F. R. S., &c.

A TREATISE ON DISEASES OF THE EYE. A new edition, edited, with numerous additions, and 243 illustrations, by ISAAC HAYS, M. D., Surgeon to Will's Hospital, &c. In one very large and handsome octavo volume, of 950 pages, strongly bound in leather with raised bands. \$5 00.

LUDLOW (J. L.), M. D.

A MANUAL OF EXAMINATIONS upon Anatomy, Physiology, Surgery, Practice of Medicine, Obstetrics, Materia Medica, Chemistry, Pharmacy, and Therapeutics. To which is added a Medical Formulary. Third edition, thoroughly revised and greatly extended and enlarged. With 370 illustrations. In one handsome royal 12mo. volume, leather, of 816 large pages. \$2 50.

We know of no better companion for the student during the hours spent in the lecture room, or to refresh, at a glance, his memory of the various topics

crammed into his head by the various professors to whom he is compelled to listen.—*Western Lancet*, May, 1857.

LEHMANN (C. G.)

PHYSIOLOGICAL CHEMISTRY. Translated from the second edition by GEORGE E. DAY, M. D., F. R. S., &c., edited by R. E. ROGERS, M. D., Professor of Chemistry in the Medical Department of the University of Pennsylvania, with illustrations selected from Funke's Atlas of Physiological Chemistry, and an Appendix of plates. Complete in two large and handsome octavo volumes, extra cloth, containing 1200 pages, with nearly two hundred illustrations. \$6 00.

The work of Lehmann stands unrivalled as the most comprehensive book of reference and information extant on every branch of the subject on which it treats.—*Edinburgh Journal of Medical Science.*

The most important contribution as yet made to Physiological Chemistry.—*Am. Journal Med. Sciences*, Jan. 1856.

BY THE SAME AUTHOR. (*Lately Published.*)

MANUAL OF CHEMICAL PHYSIOLOGY. Translated from the German, with Notes and Additions, by J. CHESTON MORRIS, M. D., with an Introductory Essay on Vital Force, by Professor SAMUEL JACKSON, M. D., of the University of Pennsylvania. With illustrations on wood. In one very handsome octavo volume, extra cloth, of 336 pages. \$2 25.

From Prof. Jackson's Introductory Essay.

In adopting the handbook of Dr. Lehmann as a manual of Organic Chemistry for the use of the students of the University, and in recommending his original work of PHYSIOLOGICAL CHEMISTRY for their more mature studies, the high value of his researches, and the great weight of his authority in that important department of medical science are fully recognized.

LYONS (ROBERT D.), K. C. C.,

Late Pathologist in-chief to the British Army in the Crimea, &c.

A TREATISE ON FEVER; or, selections from a course of Lectures on Fever. Being part of a course of Theory and Practice of Medicine. In one neat octavo volume, of 362 pages, extra cloth; \$2 00. (*Now Ready.*)

From the Author's Preface.

"I am induced to publish this work on Fever with a view to bring within the reach of the student and junior practitioner, in a convenient form, the more recent results of inquiries into the Pathology and Therapeutics of this formidable class of diseases.

"The works of the great writers on Fever are so numerous, and in the present day are scattered in so many languages, that they are difficult of access, not only to students but also to practitioners. I shall deem myself fortunate if I can in any measure supply the want which is felt in this respect.

We have great pleasure in recommending Dr. Lyons' work on *Fever* to the attention of the profession. It is a work which cannot fail to enhance the author's previous well-earned reputation, as a diligent, careful, and accurate observer.—*British Med. Journal*, March 2, 1861.

Taken as a whole we can recommend it in the highest terms as well worthy the careful perusal and study of every student and practitioner of medi-

cine. We consider the work a most valuable addition to medical literature, and one destined to wield no little influence over the mind of the profession.—*Med. and Surg. Reporter*, May 4, 1861.

This is an admirable work upon the most remarkable and most important class of diseases to which mankind are liable.—*Med. Journ. of N. Carolina*, May, 1861.

MEIGS (CHARLES D.), M. D.,

Professor of Obstetrics, &c. in the Jefferson Medical College, Philadelphia.

OBSTETRICS: THE SCIENCE AND THE ART. Third edition, revised and improved. With one hundred and twenty-nine illustrations. In one beautifully printed octavo volume, leather, of seven hundred and fifty-two large pages. \$3 75.

Though the work has received only five pages of enlargement, its chapters throughout wear the impress of careful revision. Expunging and rewriting, remodelling its sentences, with occasional new material, all evince a lively desire that it shall deserve to be regarded as improved in manner as well as matter. In the matter, every stroke of the pen has increased the value of the book, both in expungings and additions.—*Western Lancet*, Jan. 1857.

The best American work on Midwifery that is accessible to the student and practitioner.—*N. W. Med. and Surg. Journal*, Jan. 1857.

This is a standard work by a great American Obstetrician. It is the third and last edition, and, in the language of the preface, the author has "brought the subject up to the latest dates of real improvement in our art and Science."—*Nashville Journ. of Med. and Surg.*, May, 1857.

BY THE SAME AUTHOR. (*Just Issued.*)

WOMAN: HER DISEASES AND THEIR REMEDIES. A Series of Lectures to his Class. Fourth and Improved edition. In one large and beautifully printed octavo volume, leather, of over 700 pages. \$3 60.

In other respects, in our estimation, too much cannot be said in praise of this work. It abounds with beautiful passages, and for conciseness, for originality, and for all that is commendable in a work on the diseases of females, it is not excelled, and probably not equalled in the English language. On the whole, we know of no work on the diseases of women which we can so cordially commend to the student and practitioner as the one before us.—*Ohio Med. and Surg. Journal*.

The body of the book is worthy of attentive consideration, and is evidently the production of a clever, thoughtful, and sagacious physician. Dr. Meigs's letters on the diseases of the external organs, contain many interesting and rare cases, and many instructive observations. We take our leave of Dr. Meigs, with a high opinion of his talents and originality.—*The British and Foreign Medico-Chirurgical Review*.

Every chapter is replete with practical instruction, and bears the impress of being the composition of an acute and experienced mind. There is a terseness, and at the same time an accuracy in his description of symptoms, and in the rules for diagnosis,

which cannot fail to recommend the volume to the attention of the reader.—*Ranking's Abstract*.

It contains a vast amount of practical knowledge, by one who has accurately observed and retained the experience of many years.—*Dublin Quarterly Journal*.

Full of important matter, conveyed in a ready and agreeable manner.—*St. Louis Med. and Surg. Jour.*

There is an off-hand fervor, a glow, and a warm-heartedness infecting the effort of Dr. Meigs, which is entirely captivating, and which absolutely hurries the reader through from beginning to end. Besides, the book teems with solid instruction, and it shows the very highest evidence of ability, viz., the clearness with which the information is presented. We know of no better test of one's understanding a subject than the evidence of the power of lucidly explaining it. The most elementary, as well as the obscurest subjects, under the pencil of Prof. Meigs, are isolated and made to stand out in such bold relief, as to produce distinct impressions upon the mind and memory of the reader.—*The Charleston Med. Journal*.

BY THE SAME AUTHOR.

ON THE NATURE, SIGNS, AND TREATMENT OF CHILDBED FEVER. In a Series of Letters addressed to the Students of his Class. In one handsome octavo volume, extra cloth, of 365 pages. \$2 50.

The instructive and interesting author of this work, whose previous labors have placed his countrymen under deep and abiding obligations, again challenges their admiration in the fresh and vigorous, attractive and racy pages before us. It is a de-

lectable book. * * * This treatise upon childbed fevers will have an extensive sale, being destined, as it deserves, to find a place in the library of every practitioner who seems to lag in the rear.—*Nashville Journal of Medicine and Surgery*.

BY THE SAME AUTHOR; WITH COLORED PLATES.

A TREATISE ON ACUTE AND CHRONIC DISEASES OF THE NECK OF THE UTERUS. With numerous plates, drawn and colored from nature in the highest style of art. In one handsome octavo volume, extra cloth. \$4 50.

MACLISE (JOSEPH), SURGEON.

SURGICAL ANATOMY. Forming one volume, very large imperial quarto.

With sixty-eight large and splendid Plates, drawn in the best style and beautifully colored. Containing one hundred and ninety Figures, many of them the size of life. Together with copious and explanatory letter-press. Strongly and handsomely bound in extra cloth, being one of the cheapest and best executed Surgical works as yet issued in this country. \$11 00.

* * The size of this work prevents its transmission through the post-office as a whole, but those who desire to have copies forwarded by mail, can receive them in five parts, done up in stout wrappers. Price \$9 00.

One of the greatest artistic triumphs of the age in Surgical Anatomy.—*British American Medical Journal*.

No practitioner whose means will admit should fail to possess it.—*Ranking's Abstract*.

Too much cannot be said in its praise; indeed, we have not language to do it justice.—*Ohio Medical and Surgical Journal*.

The most accurately engraved and beautifully colored plates we have ever seen in an American book—one of the best and cheapest surgical works ever published.—*Buffalo Medical Journal*.

It is very rare that so elegantly printed, so well illustrated, and so useful a work, is offered at so moderate a price.—*Charleston Medical Journal*.

Its plates can boast a superiority which places them almost beyond the reach of competition.—*Medical Examiner*.

Country practitioners will find these plates of immense value.—*N. Y. Medical Gazette*.

A work which has no parallel in point of accuracy and cheapness in the English language.—*N. Y. Journal of Medicine*.

We are extremely gratified to announce to the profession the completion of this truly magnificent work, which, as a whole, certainly stands unrivalled, both for accuracy of drawing, beauty of coloring, and all the requisite explanations of the subject in hand.—*The New Orleans Medical and Surgical Journal*.

This is by far the ablest work on Surgical Anatomy that has come under our observation. We know of no other work that would justify a student, in any degree, for neglect of actual dissection. In those sudden emergencies that so often arise, and which require the instantaneous command of minute anatomical knowledge, a work of this kind keeps the details of the dissecting-room perpetually fresh in the memory.—*The Western Journal of Medicine and Surgery*.

MILLER (HENRY), M. D.,

Professor of Obstetrics and Diseases of Women and Children in the University of Louisville.

PRINCIPLES AND PRACTICE OF OBSTETRICS, &c.; including the Treatment of Chronic Inflammation of the Cervix and Body of the Uterus considered as a frequent cause of Abortion. With about one hundred illustrations on wood. In one very handsome octavo volume, of over 600 pages. (*Lately Published.*) \$3 75.

We congratulate the author that the task is done. We congratulate him that he has given to the medical public a work which will secure for him a high and permanent position among the standard authorities on the principles and practice of obstetrics. Congratulations are not less due to the medical profession of this country, on the acquisition of a treatise embodying the results of the studies, reflections, and experience of Prof. Miller. Few men, if any, in this country, are more competent than he to write on this department of medicine. Engaged for thirty-five years in an extended practice of obstetrics, for many years a teacher of this branch of instruction in one of the largest of our institutions, a diligent student as well as a careful observer, an original and independent thinker, wedded to no hobbies, ever ready to consider without prejudice new views, and to adopt innovations if they are really improvements, and withal a clear, agreeable writer, a practical treatise from his pen could not fail to possess great value.—*Buffalo Med Journal*.

In fact, this volume must take its place among the standard systematic treatises on obstetrics; a posi-

tion to which its merits justly entitle it. The style is such that the descriptions are clear, and each subject is discussed and elucidated with due regard to its practical bearings, which cannot fail to make it acceptable and valuable to both students and practitioners. We cannot, however, close this brief notice without congratulating the author and the profession on the production of such an excellent treatise. The author is a western man of whom we feel proud, and we cannot but think that his book will find many readers and warm admirers wherever obstetrics is taught and studied as a science and an art.—*The Cincinnati Lancet and Observer*.

A most respectable and valuable addition to our home medical literature, and one reflecting credit alike on the author and the institution to which he is attached. The student will find in this work a most useful guide to his studies; the country practitioner, rusty in his reading, can obtain from its pages a fair résumé of the modern literature of the science; and we hope to see this American production generally consulted by the profession.—*Va. Med. Journal*.

MACKENZIE (W.), M. D.,

Surgeon Oculist in Scotland in ordinary to Her Majesty, &c. &c.

A PRACTICAL TREATISE ON DISEASES AND INJURIES OF THE

EYE. To which is prefixed an Anatomical Introduction explanatory of a Horizontal Section of the Human Eyeball, by THOMAS WHARTON JONES, F. R. S. From the Fourth Revised and Enlarged London Edition. With Notes and Additions by ADDINELL HEWSON, M. D., Surgeon to Wills Hospital, &c. &c. In one very large and handsome octavo volume, leather, raised bands, with plates and numerous wood-cuts. \$5 25.

The treatise of Dr. Mackenzie indisputably holds the first place, and forms, in respect of learning and research, an Encyclopædia unequalled in extent by any other work of the kind, either English or foreign.—*Dixon on Diseases of the Eye*.

Few modern books on any department of medicine or surgery have met with such extended circulation, or have procured for their authors a like amount of European celebrity. The immense research which it displayed, the thorough acquaintance with the subject, practically as well as theoretically, and the

able manner in which the author's stores of learning and experience were rendered available for general use, at once procured for the first edition, as well on the continent as in this country, that high position as a standard work which each successive edition has more firmly established. We consider it the duty of every one who has the love of his profession and the welfare of his patient at heart, to make himself familiar with this the most complete work in the English language upon the diseases of the eye.—*Med. Times and Gazette*.

MAYNE'S DISPENSATORY AND THERAPEUTICAL REMEMBRANCER. With every Practical Formula contained in the three British Pharmacopæias. Edited, with the addition of the Formulae of the U. S. Pharmacopæia, by R. E. GRIFFITH, M. D. 12mo. vol. ex. cl., 300 pp. 75 c.

MALGAIGNE'S OPERATIVE SURGERY, based on Normal and Pathological Anatomy. Translated from the French by FREDERICK BRITTON, A. B., M. D. With numerous illustrations on wood. In one handsome octavo volume, extra cloth, of nearly 800 pages.

MILLER (JAMES), F. R. S. E.,
Professor of Surgery in the University of Edinburgh, &c.

PRINCIPLES OF SURGERY. Fourth American, from the third and revised Edinburgh edition. In one large and very beautiful volume, leather, of 700 pages, with two hundred and forty illustrations on wood. \$3 75.

The work of Mr. Miller is too well and too favorably known among us, as one of our best text-books, to render any further notice of it necessary than the announcement of a new edition, the fourth in our country, a proof of its extensive circulation among us. As a concise and reliable exposition of the science of modern surgery, it stands deservedly high—we know not its superior.—*Boston Med. and Surg. Journal.*

The work takes rank with Watson's Practice of Physic; it certainly does not fall behind that great work in soundness of principle or depth of reasoning and research. No physician who values his reputation, or seeks the interests of his clients, can acquit himself before his God and the world without making himself familiar with the sound and philosophical views developed in the foregoing book.—*New Orleans Med. and Surg. Journal.*

BY THE SAME AUTHOR. (Just Issued.)

THE PRACTICE OF SURGERY. Fourth American from the last Edinburgh edition. Revised by the American editor. Illustrated by three hundred and sixty-four engravings on wood. In one large octavo volume, leather, of nearly 700 pages. \$3 75.

No eulogium of ours could add to the popularity of Miller's Surgery. Its reputation in this country is unsurpassed by that of any other work, and, when taken in connection with the author's *Principles of Surgery*, constitutes a whole, without reference to which no conscientious surgeon would be willing to practice his art.—*Southern Med. and Surg. Journal.*

It is seldom that two volumes have ever made so profound an impression in so short a time as the "Principles" and the "Practice" of Surgery by Mr. Miller—or so richly merited the reputation they have acquired. The author is an eminently sensible, practical, and well-informed man, who knows exactly what he is talking about and exactly how to talk it.—*Kentucky Medical Recorder.*

By the almost unanimous voice of the profession,

his works, both on the principles and practice of surgery have been assigned the highest rank. If we were limited to but one work on surgery, that one should be Miller's, as we regard it as superior to all others.—*St. Louis Med. and Surg. Journal.*

The author has in this and his "Principles," presented to the profession one of the most complete and reliable systems of Surgery extant. His style of writing is original, impressive, and engaging, energetic, concise, and lucid. Few have the faculty of condensing so much in small space, and at the same time so persistently holding the attention. Whether as a text-book for students or a book of reference for practitioners, it cannot be too strongly recommended.—*Southern Journal of Med. and Physical Sciences.*

MORLAND (W. W.), M. D.,
Fellow of the Massachusetts Medical Society, &c.

DISEASES OF THE URINARY ORGANS; a Compendium of their Diagnosis, Pathology, and Treatment. With illustrations. In one large and handsome octavo volume, of about 600 pages, extra cloth. (Just Issued.) \$3 50.

Taken as a whole, we can recommend Dr. Morland's compendium as a very desirable addition to the library of every medical or surgical practitioner.—*Brit. and For. Med.-Chir. Rev.*, April, 1859.

Every medical practitioner whose attention has been to any extent attracted towards the class of diseases to which this treatise relates, must have often and sorely experienced the want of some full, yet concise recent compendium to which he could

refer. This desideratum has been supplied by Dr. Morland, and it has been ably done. He has placed before us a full, judicious, and reliable digest. Each subject is treated with sufficient minuteness, yet in a succinct, narrative style, such as to render the work one of great interest, and one which will prove in the highest degree useful to the general practitioner.—*N. Y. Journ. of Medicine.*

BY THE SAME AUTHOR —(Now Ready.)

THE MORBID EFFECTS OF THE RETENTION IN THE BLOOD OF THE ELEMENTS OF THE URINARY SECRETION. Being the Dissertation to which the Fiske Fund Prize was awarded, July 11, 1861. In one small octavo volume, 83 pages, extra cloth. 75 cents.

MONTGOMERY (W. F.), M. D., M. R. I. A., &c.,
Professor of Midwifery in the King and Queen's College of Physicians in Ireland, &c.

AN EXPOSITION OF THE SIGNS AND SYMPTOMS OF PREGNANCY.

With some other Papers on Subjects connected with Midwifery. From the second and enlarged English edition. With two exquisite colored plates, and numerous wood-cuts. In one very handsome octavo volume, extra cloth, of nearly 600 pages. (Lately Published.) \$3 75.

A book unusually rich in practical suggestions.—*Am. Journal Med. Sciences*, Jan. 1857.

These several subjects so interesting in themselves, and so important, every one of them, to the most delicate and precious of social relations, controlling often the honor and domestic peace of a family, the legitimacy of offspring, or the life of its parent, are all treated with an elegance of diction, fulness of illustrations, acuteness and justice of reasoning, unparalleled in obstetrics, and unsurpassed in medicine. The reader's interest can never flag, so

fresh, and vigorous, and classical is our author's style; and one forgets, in the renewed charm of every page, that it, and every line, and every word has been weighed and reweighed through years of preparation; that this is of all others the book of Obstetric Law, on each of its several topics; on all points connected with pregnancy, to be everywhere received as a manual of special jurisprudence, at once announcing fact, affording argument, establishing precedent, and governing alike the jurymen, advocate, and judge.—*N. A. Med.-Chir. Review.*

MOHR (FRANCIS), PH. D., AND REDWOOD (THEOPHILUS).

PRACTICAL PHARMACY. Comprising the Arrangements, Apparatus, and Manipulations of the Pharmaceutical Shop and Laboratory. Edited, with extensive Additions, by Prof. WILLIAM PROCTER, of the Philadelphia College of Pharmacy. In one handsomely printed octavo volume, extra cloth, of 570 pages, with over 500 engravings on wood. \$2 75.

NEILL (JOHN), M. D.,

Surgeon to the Pennsylvania Hospital, &c.; and

FRANCIS GURNEY SMITH, M. D.,

Professor of Institutes of Medicine in the Pennsylvania Medical College.

AN ANALYTICAL COMPENDIUM OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE; for the Use and Examination of Students. A new edition, revised and improved. In one very large and handsomely printed royal 12mo. volume, of about one thousand pages, with 374 wood-cuts. Strongly bound in leather, with raised bands. \$3 00.

The very flattering reception which has been accorded to this work, and the high estimate placed upon it by the profession, as evinced by the constant and increasing demand which has rapidly exhausted two large editions, have stimulated the authors to render the volume in its present revision more worthy of the success which has attended it. It has accordingly been thoroughly examined, and such errors as had on former occasions escaped observation have been corrected, and whatever additions were necessary to maintain it on a level with the advance of science have been introduced. The extended series of illustrations has been still further increased and much improved, while, by a slight enlargement of the page, these various additions have been incorporated without increasing the bulk of the volume.

The work is, therefore, again presented as eminently worthy of the favor with which it has hitherto been received. As a book for daily reference by the student requiring a guide to his more elaborate text-books, as a manual for preceptors desiring to stimulate their students by frequent and accurate examination, or as a source from which the practitioners of older date may easily and cheaply acquire a knowledge of the changes and improvement in professional science, its reputation is permanently established.

The best work of the kind with which we are acquainted.—*Med. Examiner.*

Having made free use of this volume in our examinations of pupils, we can speak from experience in recommending it as an admirable compend for students, and as especially useful to preceptors who examine their pupils. It will save the teacher much labor by enabling him readily to recall all of the points upon which his pupils should be examined. A work of this sort should be in the hands of every one who takes pupils into his office with a view of examining them; and this is unquestionably the best of its class.—*Transylvania Med. Journal*

In the rapid course of lectures, where work for

the students is heavy, and review necessary for an examination, a compend is not only valuable, but it is almost a *sine qua non*. The one before us is, in most of the divisions, the most unexceptionable of all books of the kind that we know of. The newest and soundest doctrines and the latest improvements and discoveries are explicitly, though concisely, laid before the student. There is a class to whom we very sincerely commend this cheap book as worth its weight in silver—that class is the graduates in medicine of more than ten years' standing, who have not studied medicine since. They will perhaps find out from it that the science is not exactly now what it was when they left it off.—*The Stethoscope.*

NELIGAN (J. MOORE), M. D., M. R. I. A., &c.

ATLAS OF CUTANEOUS DISEASES. In one beautiful quarto volume, extra cloth, with splendid colored plates, presenting nearly one hundred elaborate representations of disease. \$4 50.


This beautiful volume is intended as a complete and accurate representation of all the varieties of Diseases of the Skin. While it can be consulted in conjunction with any work on Practice, it has especial reference to the author's "Treatise on Diseases of the Skin," so favorably received by the profession some years since. The publishers feel justified in saying that few more beautifully executed plates have ever been presented to the profession of this country.

Neligan's Atlas of Cutaneous Diseases supplies a long existent desideratum much felt by the largest class of our profession. It presents, in quarto size, 16 plates, each containing from 3 to 6 figures, and forming in all a total of 90 distinct representations of the different species of skin affections, grouped together in genera or families. The illustrations have been taken from nature, and have been copied with such fidelity that they present a striking picture of life; in which the reduced scale aptly serves to

give, at a *coup d'œil*, the remarkable peculiarities of each individual variety. And while thus the disease is rendered more definable, there is yet no loss of proportion incurred by the necessary concentration. Each figure is highly colored, and so truthful has the artist been that the most fastidious observer could not justly take exception to the correctness of the execution of the pictures under his scrutiny.—*Montreal Med. Chronicle.*

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN. Third American edition. In one neat royal 12mo. volume, extra cloth, of 334 pages. \$1 00.

 The two volumes will be sent by mail on receipt of *Five Dollars.*

OWEN ON THE DIFFERENT FORMS OF THE SKELETON, AND OF THE TEETH.

One vol. royal 12mo., extra cloth with numerous illustrations. \$1 25

PIRRIE (WILLIAM), F. R. S. E.,

Professor of Surgery in the University of Aberdeen.

THE PRINCIPLES AND PRACTICE OF SURGERY. Edited by JOHN NEILL, M. D., Professor of Surgery in the Penna. Medical College, Surgeon to the Pennsylvania Hospital, &c. In one very handsome octavo volume, leather, of 780 pages, with 316 illustrations. \$3 75.

We know of no other surgical work of a reasonable size, wherein there is so much theory and practice, or where subjects are more soundly or clearly taught.—*The Stethoscope.*

Prof. Pirrie, in the work before us, has elaborately

discussed the principles of surgery, and a safe and effectual practice predicated upon them. Perhaps no work upon this subject heretofore issued is so full upon the science of the art of surgery.—*Nashville Journal of Medicine and Surgery.*

PARRISH (EDWARD),

Lecturer on Practical Pharmacy and Materia Medica in the Pennsylvania Academy of Medicine, &c.

AN INTRODUCTION TO PRACTICAL PHARMACY. Designed as a Text-Book for the Student, and as a Guide for the Physician and Pharmacist. With many Formulae and Prescriptions. Second edition, greatly enlarged and improved. In one handsome octavo volume of 720 pages, with several hundred Illustrations, extra cloth. \$3 50. (*Just Issued.*)

During the short time in which this work has been before the profession, it has been received with very great favor, and in assuming the position of a standard authority, it has filled a vacancy which had been severely felt. Stimulated by this encouragement, the author, in availing himself of the opportunity of revision, has spared no pains to render it more worthy of the confidence bestowed upon it, and his assiduous labors have made it rather a new book than a new edition, many portions having been rewritten, and much new and important matter added. These alterations and improvements have been rendered necessary by the rapid progress made by pharmaceutical science during the last few years, and by the additional experience obtained in the practical use of the volume as a text-book and work of reference. To accommodate these improvements, the size of the page has been materially enlarged, and the number of pages considerably increased, presenting in all nearly *one-half more* matter than the last edition. The work is therefore now presented as a complete exponent of the subject in its most advanced condition. From the most ordinary matters in the dispensing office, to the most complicated details of the vegetable alkaloids, it is hoped that everything requisite to the practising physician, and to the apothecary, will be found fully and clearly set forth, and that the new matter alone will be worth more than the very moderate cost of the work to those who have been consulting the previous edition.

That Edward Parrish, in writing a book upon practical Pharmacy some few years ago—one eminently original and unique—did the medical and pharmaceutical professions a great and valuable service, no one, we think, who has had access to its pages will deny; doubly welcome, then, is this new edition, containing the added results of his recent and rich experience as an observer, teacher, and practical operator in the pharmaceutical laboratory. The excellent plan of the first is more thoroughly, and in detail, carried out in this edition.—*Peninsular Med. Journal*, Jan. 1860.

Of course, all apothecaries who have not already a copy of the first edition will procure one of this; it is, therefore, to physicians residing in the country and in small towns, who cannot avail themselves of the skill of an educated pharmacist, that we would especially commend this work. In it they

will find all that they desire to know, and should know, but very little of which they do really know in reference to this important collateral branch of their profession; for it is a well established fact, that, in the education of physicians, while the science of medicine is generally well taught, very little attention is paid to the art of preparing them for use, and we know not how this defect can be so well remedied as by procuring and consulting Dr. Parrish's excellent work.—*St. Louis Med. Journal*, Jan. 1860.

We know of no work on the subject which would be more indispensable to the physician or student desiring information on the subject of which it treats. With Griffith's "Medical Formulary" and this, the practising physician would be supplied with nearly or quite all the most useful information on the subject.—*Charleston Med. Jour. and Review*, Jan. 1860

PEASLEE (E. R.), M. D.,

Professor of Physiology and General Pathology in the New York Medical College.

HUMAN HISTOLOGY, in its relations to Anatomy, Physiology, and Pathology; for the use of Medical Students. With four hundred and thirty-four illustrations. In one handsome octavo volume, of over 600 pages. (*Lately Published.*) \$7 75.

It embraces a library upon the topics discussed within itself, and is just what the teacher and learner need. Another advantage, by no means to be overlooked, everything of real value in the wide range which it embraces, is with great skill compressed into an octavo volume of but little more than six hundred pages. We have not only the whole subject of Histology, interesting in itself, ably and fully discussed, but what is of infinitely greater interest to the student, because of greater practical value, are its relations to Anatomy, Physiology, and Pathology, which are here fully and satisfactorily set forth.—*Nashville Journ. of Med. and Surgery*.

We would recommend it to the medical student and practitioner, as containing a summary of all that is known of the important subjects which it treats; of all that is contained in the great works of Simon and Lehmann, and the organic chemists in general. Master this one volume, we would say to the medical student and practitioner—master this book and you know all that is known of the great fundamental principles of medicine, and we have no hesitation in saying that it is an honor to the American medical profession that one of its members should have produced it.—*St. Louis Med. and Surg. Journal*.

PEREIRA (JONATHAN), M. D., F. R. S., AND L. S.

THE ELEMENTS OF MATERIA MEDICA AND THERAPEUTICS.

Third American edition, enlarged and improved by the author; including Notices of most of the Medicinal Substances in use in the civilized world, and forming an Encyclopædia of Materia Medica. Edited, with Additions, by JOSEPH CARSON, M. D., Professor of Materia Medica and Pharmacy in the University of Pennsylvania. In two very large octavo volumes of 2100 pages, on small type, with about 500 illustrations on stone and wood, strongly bound in leather, with raised bands. \$9 00.

*** Vol. II. will no longer be sold separate.

PARKER (LANGSTON),

Surgeon to the Queen's Hospital, Birmingham.

THE MODERN TREATMENT OF SYPHILITIC DISEASES, BOTH PRIMARY AND SECONDARY; comprising the Treatment of Constitutional and Confirmed Syphilis, by a safe and successful method. With numerous Cases, Formulae, and Clinical Observations. From the Third and entirely rewritten London edition. In one neat octavo volume, extra cloth, of 316 pages. \$1 75.

ROYLE'S MATERIA MEDICA AND THERAPEUTICS; including the Preparations of the Pharmacopœias of London, Edinburgh, Dublin, and of the United States. With many new medicines. Edited by JOSEPH CARSON, M. D. With ninety-eight illustrations. In one octavo volume, extra cloth, of about 700 pages. \$3 00.

RAMSBOTHAM (FRANCIS H.), M. D.

THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY, in reference to the Process of Parturition. A new and enlarged edition, thoroughly revised by the Author. With Additions by W. V. KEATING, M. D., Professor of Obstetrics, &c., in the Jefferson Medical College, Philadelphia. In one large and handsome imperial octavo volume, of 650 pages, strongly bound in leather, with raised bands; with sixty-four beautiful Plates, and numerous Wood-cuts in the text, containing in all nearly 200 large and beautiful figures. \$5 00.

From Prof. Hodge, of the University of Pa.

To the American public, it is most valuable, from its intrinsic undoubted excellence, and as being the best authorized exponent of British Midwifery. Its circulation will, I trust, be extensive throughout our country.

It is unnecessary to say anything in regard to the utility of this work. It is already appreciated in our country for the value of the matter, the clearness of its style, and the fulness of its illustrations. To the physician's library it is indispensable, while to the student as a text-book, from which to extract the material for laying the foundation of an education on obstetrical science, it has no superior.—*Ohio Med and Surg. Journal*.

The publishers have secured its success by the

truly elegant style in which they have brought it out, excelling themselves in its production, especially in its plates. It is dedicated to Prof. Meigs, and has the emphatic endorsement of Prof. Hodge, as the best exponent of British Midwifery. We know of no text-book which deserves in all respects to be more highly recommended to students, and we could wish to see it in the hands of every practitioner, for they will find it invaluable for reference.—*Med. Gazette*.

RICORD (P.), M. D.

A TREATISE ON THE VENEREAL DISEASE. By JOHN HUNTER, F. R. S.

With copious Additions, by PH. RICORD, M. D. Translated and Edited, with Notes, by FREEMAN J. BUMSTEAD, M. D., Lecturer on Venereal at the College of Physicians and Surgeons, New York. Second edition, revised, containing a *résumé* of RICORD'S RECENT LECTURES ON CHANCERE. In one handsome octavo volume, extra cloth, of 550 pages, with eight plates. \$3 25. (Just Issued.)

In revising this work, the editor has endeavored to introduce whatever matter of interest the recent investigations of syphilographers have added to our knowledge of the subject. The principal source from which this has been derived is the volume of "Lectures on Chancre," published a few months since by M. Ricord, which affords a large amount of new and instructive material on many controverted points. In the previous edition, M. Ricord's additions amounted to nearly one-third of the whole, and with the matter now introduced, the work may be considered to present his views and experience more thoroughly and completely than any other.

Every one will recognize the attractiveness and value which this work derives from this presenting the opinions of these two masters side by side. But, it must be admitted, what has made the fortune of the book, is the fact that it contains the "most complete embodiment of the veritable doctrines of the Hôpital du Midi," which has ever been made public. The doctrinal ideas of M. Ricord, ideas which, if not universally adopted, are incontestably dominant, have heretofore only been interpreted by more or less skillful

secretaries, sometimes accredited and sometimes not.

In the notes to Hunter, the master substitutes himself for his interpreters, and gives his original thoughts to the world in a lucid and perfectly intelligible manner. In conclusion we can say that this is incontestably the best treatise on syphilis with which we are acquainted, and, as we do not often employ the phrase, we may be excused for expressing the hope that it may find a place in the library of every physician.—*Virginia Med. and Surg. Journal*.

BY THE SAME AUTHOR.

RICORD'S LETTERS ON SYPHILIS. Translated by W. P. LATTIMORE, M. D.

In one neat octavo volume, of 270 pages, extra cloth. \$2 00.

ROKITANSKY (CARL), M. D.,

Curator of the Imperial Pathological Museum, and Professor at the University of Vienna, &c.

A MANUAL OF PATHOLOGICAL

bound in two, extra cloth, of about 1200 pages. KING, C. H. MOORE, and G. E. DAY. \$5 50.

The profession is too well acquainted with the reputation of Rokitansky's work to need our assurance that this is one of the most profound, thorough, and valuable books ever issued from the medical press. It is *sui generis*, and has no standard of comparison. It is only necessary to announce that it is issued in a form as cheap as is compatible with its size and preservation, and its sale follows as a matter of course. No library can be called complete without it.—*Buffalo Med. Journal*.

An attempt to give our readers any adequate idea of the vast amount of instruction accumulated in these volumes, would be feeble and hopeless. The effort of the distinguished author to concentrate in a small space his great fund of knowledge, has

ANATOMY. Four volumes, octavo, Translated by W. E. SWAINE, EDWARD SIEVE-

so charged his text with valuable truths, that any attempt of a reviewer to epitomize is at once paralyzed, and must end in a failure.—*Western Lancet*.

As this is the highest source of knowledge upon the important subject of which it treats, no real student can afford to be without it. The American publishers have entitled themselves to the thanks of the profession of their country, for this timely and beautiful edition.—*Nashville Journal of Medicine*.

As a book of reference, therefore, this work must prove of inestimable value, and we cannot too highly recommend it to the profession.—*Charleston Med. Journal and Review*.

This book is a necessity to every practitioner.—*Am. Med. Monthly*.

RIGBY (EDWARD), M. D.,

Senior Physician to the General Lying-in Hospital, &c.

A SYSTEM OF MIDWIFERY. With Notes and Additional Illustrations.

Second American Edition. One volume octavo, extra cloth, 422 pages. \$2 50.

BY THE SAME AUTHOR. (Lately Published.)

ON THE CONSTITUTIONAL TREATMENT OF FEMALE DISEASES.

In one neat royal 12mo. volume, extra cloth, of about 250 pages. \$1 00.

STILLE (ALFRED), M. D.

THERAPEUTICS AND MATERIA MEDICA; a Systematic Treatise on the Action and Uses of Medicinal Agents, including their Description and History. In two large and handsome octavo volumes, of 1789 pages. (Just Issued.) \$8 00.

This work is designed especially for the student and practitioner of medicine, and treats the various articles of the Materia Medica from the point of view of the bedside, and not of the shop or of the lecture-room. While thus endeavoring to give all practical information likely to be useful with respect to the employment of special remedies in special affections, and the results to be anticipated from their administration, a copious Index of Diseases and their Remedies renders the work eminently fitted for reference by showing at a glance the different means which have been employed, and enabling the practitioner to extend his resources in difficult cases with all that the experience of the profession has suggested.

Rarely, indeed, have we had submitted to us a work on medicine so ponderous in its dimensions as that now before us, and yet so fascinating in its contents. It is, therefore, with a peculiar gratification that we recognize in Dr. Stille the possession of many of those more distinguished qualifications which entitle him to approbation, and which justify him in coming before his medical brethren as an instructor. A comprehensive knowledge, tested by a sound and penetrating judgment, joined to a love of progress—which a discriminating spirit of inquiry has tempered so as to accept nothing new because it is new, and abandon nothing old because it is old, but which estimates either according to its relations to a just logic and experience—manifests itself everywhere, and gives to the guidance of the author all the assurance of safety which the difficulties of his subject can allow. In conclusion, we earnestly advise our readers to ascertain for themselves, by a study of Dr. Stille's volumes, the great value and interest of the stores of knowledge they present. We have pleasure in referring rather to the ample treasury of undoubted truths, the real and assured conquest of medicine, accumulated by Dr. Stille in his pages; and commend the sum of his labors to the attention of our readers, as alike honorable to our science, and creditable to the zeal, the candor, and the judgment of him who has garnered the whole so carefully.—*Edinburgh Med. Journal*.

Our expectations of the value of this work were based on the well-known reputation and character of the author as a man of scholarly attainments, an elegant writer, a candid inquirer after truth, and a philosophical thinker; we knew that the task would be conscientiously performed, and that few, if any, among the distinguished medical teachers in this country are better qualified than he to prepare a systematic treatise on therapeutics in accordance with the present requirements of medical science. Our preliminary examination of the work has satis-

fied us that we were not mistaken in our anticipations.—*New Orleans Medical News*, March, 1860.

The most recent authority is the one last mentioned, Stille. His great work on "Materia Medica and Therapeutics," published last year, in two octavo volumes, of some sixteen hundred pages, while it embodies the results of the labor of others up to the time of publication, is enriched with a great amount of original observation and research. We would draw attention, by the way, to the very convenient mode in which the *Index* is arranged in this work. There is first an "Index of Remedies;" next an "Index of Diseases and their Remedies;" Such an arrangement of the Indices, in our opinion, greatly enhances the practical value of books of this kind. In tedious, obstinate cases of disease, where we have to try one remedy after another until our stock is pretty nearly exhausted, and we are almost driven to our wit's end, such an index as the second of the two just mentioned, is precisely what we want.—*London Med. Times and Gazette*, April, 1861.

We think this work will do much to obviate the reluctance to a thorough investigation of this branch of scientific study, for in the wide range of medical literature treasured in the English tongue, we shall hardly find a work written in a style more clear and simple, conveying forcibly the facts taught, and yet free from turgidity and redundancy. There is a fascination in its pages that will insure to it a wide popularity and attentive perusal, and a degree of usefulness not often attained through the influence of a single work. The author has much enhanced the practical utility of his book by passing briefly over the physical, botanical, and commercial history of medicines, and directing attention chiefly to their physiological action, and their application for the amelioration or cure of disease. He ignores hypothesis and theory which are so alluring to many medical writers, and so liable to lead them astray, and confines himself to such facts as have been tried in the crucible of experience.—*Chicago Medical Journal*.

SMITH (HENRY H.), M. D. AND HORNER (WILLIAM E.), M. D.
AN ANATOMICAL ATLAS, illustrative of the Structure of the Human Body.

In one volume, large imperial octavo, extra cloth, with about six hundred and fifty beautiful figures. \$3 00.

These figures are well selected, and present a complete and accurate representation of that wonderful fabric, the human body. The plan of this Atlas, which renders it so peculiarly convenient for the student, and its superb artistic execution, have been already pointed out. We must congratulate the student upon the completion of this Atlas as it is the most convenient work of the kind that has yet appeared; and we must add, the very beautiful manner in which it is "got up" is so creditable to the country as to be flattering to our national pride.—*American Medical Journal*.

SHARPEY (WILLIAM), M. D., JONES QUAIN, M. D., AND RICHARD QUAIN, F. R. S., &c.

HUMAN ANATOMY. Revised, with Notes and Additions, by JOSEPH LEIDY, M. D., Professor of Anatomy in the University of Pennsylvania. Complete in two large octavo volumes, leather, of about thirteen hundred pages. Beautifully illustrated with over five hundred engravings on wood. \$6 00.

SIMPSON (J. Y.), M. D.,

Professor of Midwifery, &c., in the University of Edinburgh, &c.

CLINICAL LECTURES ON THE DISEASES OF FEMALES. With numerous illustrations.

This valuable series of practical Lectures is now appearing in the "MEDICAL NEWS AND LIBRARY" for 1860, 1861, and 1862, and can thus be had without cost by subscribers to the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES." See p. 2.

SOLLY ON THE HUMAN BRAIN; its Structure, Physiology, and Diseases. From the Second and much enlarged London edition. In one octavo volume, extra cloth, of 500 pages, with 120 woodcuts. \$2 00.

SKYE'S OPERATIVE SURGERY. In one very

handsome octavo volume, extra cloth, of over 650 pages, with about one hundred woodcuts. \$3 25.

SIMON'S GENERAL PATHOLOGY, as conducive to the Establishment of Rational Principles for the prevention and Cure of Disease. In one octavo volume, extra cloth, of 212 pages. \$1 25.

SARGENT (F. W.), M. D.

ON BANDAGING AND OTHER OPERATIONS OF MINOR SURGERY.

New edition, with an additional chapter on Military Surgery. One handsome royal 12mo. vol., of nearly 400 pages, with 184 wood cuts. Leather, \$1 50. (*Now Ready.*)

The value of this work as a handy and convenient manual for surgeons engaged in active duty in the field and hospital, has induced the publishers to render it more complete for those purposes by the addition of a chapter on gun-shot wounds and other matters peculiar to military surgery. In its present form, therefore, with no increase in price, it will be found a very cheap and convenient vade-mecum for consultation and reference in the daily exigencies of military as well as civil practice.

We have read Bourgerie's Minor Surgery with pleasure and profit, but in many respects the volume now before us immeasurably transcends it. We consider that no better book could be placed in the hands of an hospital dresser, or the young surgeon, whose education in this respect has not been perfected. We most cordially commend this volume as one which the medical student should most closely study, to perfect himself in these minor surgical operations in which neatness and dexterity are so much required, and on which a great portion of his reputation as a future surgeon must evidently rest. And to the surgeon in practice it must prove itself a valuable volume, as instructive on many points which he may have forgotten.—*British American Journal*, May, 1862.

The instruction given upon the subject of *Bandaging*, is alone of great value, and while the author modestly proposes to instruct the students of medicine, and the younger physicians, we will say that experienced physicians will obtain many exceedingly valuable suggestions by its perusal. Without attempting to particularize further, we will conclude our brief notice by saying, that it will be found one of the most satisfactory manuals for reference in the field, or hospital yet published; thoroughly adapted to the wants of Military surgeons, and at the same time equally useful for ready and convenient reference by surgeons everywhere.—*Buffalo Med. and Surg. Journal*, June, 1862.

SMITH (W. TYLER), M. D.,

Physician Accoucheur to St. Mary's Hospital, &c.

ON PARTURITION, AND THE PRINCIPLES AND PRACTICE OF OBSTETRICS. In one royal 12mo. volume, extra cloth, of 400 pages. \$1 25.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON THE PATHOLOGY AND TREATMENT OF LEUCORRHOEA. With numerous illustrations. In one very handsome octavo volume, extra cloth, of about 250 pages. \$1 50.

TANNER (T. H.), M. D.,

Physician to the Hospital for Women, &c.

A MANUAL OF CLINICAL MEDICINE AND PHYSICAL DIAGNOSIS.

To which is added The Code of Ethics of the American Medical Association. Second American Edition. In one neat volume, small 12mo., extra cloth, 87½ cents.

TAYLOR (ALFRED S.), M. D., F. R. S.,

Lecturer on Medical Jurisprudence and Chemistry in Guy's Hospital.

MEDICAL JURISPRUDENCE. Fifth American, from the seventh improved and enlarged London edition. With Notes and References to American Decisions, by EDWARD HARTSHORNE, M. D. In one large 8vo. volume, leather, of over 700 pages. (*Now Ready.*) \$3 25.

This standard work having had the advantage of two revisions at the hands of the author since the appearance of the last American edition, will be found thoroughly revised and brought up completely to the present state of the science. As a work of authority, it must therefore maintain its position, both as a text-book for the student, and a compendious treatise to which the practitioner can at all times refer in cases of doubt or difficulty.

No work upon the subject can be put into the hands of students either of law or medicine which will engage them more closely or profitably; and none could be offered to the busy practitioner of either calling, for the purpose of casual or hasty reference, that would be more likely to afford the aid desired. We therefore recommend it as the best and safest manual for daily use.—*American Journal of Medical Sciences*.

It is not excess of praise to say that the volume before us is the very best treatise extant on Medical Jurisprudence. In saying this, we do not wish to be understood as detracting from the merits of the excellent works of Beck, Ryan, Traill, Guy, and others; but in interest and value we think it must be conceded that Taylor is superior to anything that has preceded it.—*N. W. Medical and Surg. Journal*

American and British legal medicine. It should be in the possession of every physician, as the subject is one of great and increasing importance to the public as well as to the profession.—*St. Louis Med. and Surg. Journal*.

This work of Dr. Taylor's is generally acknowledged to be one of the ablest extant on the subject of medical jurisprudence. It is certainly one of the most attractive books that we have met with; supplying so much both to interest and instruct, that we do not hesitate to affirm that after having once commenced its perusal, few could be prevailed upon to desist before completing it. In the last London edition, all the newly observed and accurately recorded facts have been inserted, including much that is recent of Chemical, Microscopical, and Pathological research, besides papers on numerous subjects never before published.—*Charleston Med. Journal and Review*.

It is at once comprehensive and eminently practical, and by universal consent stands at the head of

BY THE SAME AUTHOR.

ON POISONS, IN RELATION TO MEDICAL JURISPRUDENCE AND MEDICINE. Second American, from a second and revised London edition. In one large octavo volume, of 755 pages, leather. \$3 50.

Mr. Taylor's position as the leading medical jurist of England, has conferred on him extraordinary advantages in acquiring experience on these subjects, nearly all cases of moment being referred to him for examination, as an expert whose testimony is generally accepted as final. The results of his labors, therefore, as gathered together in this volume, carefully weighed and sifted, and presented in the clear and intelligible style for which he is noted, may be received as an acknowledged authority, and as a guide to be followed with implicit confidence.

TODD (ROBERT BENTLEY), M. D., F. R. S.,
Professor of Physiology in King's College, London; and
WILLIAM BOWMAN, F. R. S.,
Demonstrator of Anatomy in King's College, London.

THE PHYSIOLOGICAL ANATOMY AND PHYSIOLOGY OF MAN. With about three hundred large and beautiful illustrations on wood. Complete in one large octavo volume, of 950 pages, leather. Price \$4 50.

Gentlemen who have received portions of this work, as published in the "MEDICAL NEWS AND LIBRARY," can now complete their copies, if immediate application be made. It will be furnished as follows, free by mail, in paper covers, with cloth backs.

PARTS I., II., III. (pp. 25 to 552), \$2 50.

PART IV. (pp. 553 to end, with Title, Preface, Contents, &c.), \$2 00

Or, **PART IV., SECTION II.** (pp. 725 to end, with Title, Preface, Contents, &c.), \$1 25.

A magnificent contribution to British medicine, and the American physician who shall fail to peruse it, will have failed to read one of the most instructive books of the nineteenth century.—*N. O. Med. and Surg. Journal.*

It is more concise than Carpenter's Principles, and more modern than the accessible edition of Müller's Elements; its details are brief, but sufficient; its descriptions vivid; its illustrations exact and copious; and its language terse and perspicuous.—*Charleston Med. Journal.*

We know of no work on the subject of physiology

so well adapted to the wants of the medical student. Its completion has been thus long delayed, that the author might secure accuracy by personal observation.—*St. Louis Med. and Surg. Journal.*

Our notice, though it conveys but a very feeble and imperfect idea of the magnitude and importance of the work now under consideration, already transcends our limits; and, with the indulgence of our readers, and the hope that they will peruse the book for themselves, as we feel we can with confidence recommend it, we leave it in their hands.—*The Northwestern Med. and Surg. Journal.*

TODD (R. B.) M. D., F. R. S., &c.

CLINICAL LECTURES ON CERTAIN DISEASES OF THE URINARY ORGANS AND ON DROPSIES. In one octavo volume, 284 pages. \$1 50.

BY THE SAME AUTHOR. (*Now Ready.*)

CLINICAL LECTURES ON CERTAIN ACUTE DISEASES. In one neat octavo volume, of 320 pages, extra cloth. \$1 75.

TOYNBEE (JOSEPH), F. R. S.,

Aural Surgeon to, and Lecturer on Surgery at, St. Mary's Hospital.

A PRACTICAL TREATISE ON DISEASES OF THE EAR; their Diagnosis, Pathology, and Treatment. Illustrated with one hundred engravings on wood. In one very handsome octavo volume, extra cloth, \$3 00. (*Just Issued.*)

The work, as was stated at the outset of our notice, is a model of its kind, and every page and paragraph of it are worthy of the most thorough study. Considered all in all—as an original work, well written, philosophically elaborated, and happily illustrated with cases and drawings—it is by far the ablest monograph that has ever appeared on the anatomy and diseases of the ear, and one of the most valuable contributions to the art and science of surgery in the nineteenth century.—*N. Amer. Medical-Chirurg. Review*, Sept. 1860.

To recommend such a work, even after the mere hint we have given of its original excellence and value, would be a work of supererogation. We are speaking within the limits of modest acknowledgment,

and with a sincere and unbiassed judgment, when we affirm that as a treatise on Aural Surgery, it is without a rival in our language or any other.—*Charleston Med. Journ. and Review*, Sept. 1860.

The work of Mr. Toynbee is undoubtedly, upon the whole, the most valuable production of the kind in any language. The author has long been known by his numerous monographs upon subjects connected with diseases of the ear, and is now regarded as the highest authority on most points in his department of science. Mr. Toynbee's work, as we have already said, is undoubtedly the most reliable guide for the study of the diseases of the ear in any language, and should be in the library of every physician.—*Chicago Med. Journal*, July, 1860.

WILLIAMS (C. J. B.), M. D., F. R. S.,

Professor of Clinical Medicine in University College, London, &c.

PRINCIPLES OF MEDICINE. An Elementary View of the Causes, Nature, Treatment, Diagnosis, and Prognosis of Disease; with brief remarks on Hygienics, or the preservation of health. A new American, from the third and revised London edition. In one octavo volume, leather, of about 500 pages. \$2 50. (*Just Issued.*)

We find that the deeply-interesting matter and style of this book have so far fascinated us, that we have unconsciously hung upon its pages, not too long, indeed, for our own profit, but longer than reviewers can be permitted to indulge. We leave the further analysis to the student and practitioner. Our judgment of the work has already been sufficiently

expressed. It is a judgment of almost unqualified praise.—*London Lancet.*

A text-book to which no other in our language is comparable.—*Charleston Medical Journal.*

No work has ever achieved or maintained a more deserved reputation.—*Va. Med. and Surg. Journal.*

WHAT TO OBSERVE

AT THE BEDSIDE AND AFTER DEATH, IN MEDICAL CASES.

Published under the authority of the London Society for Medical Observation. A new American, from the second and revised London edition. In one very handsome volume, royal 12mo., extra cloth. \$1 00.

To the observer who prefers accuracy to blunders and precision to carelessness, this little book is invaluable.—*N. H. Journal of Medicine.*

One of the finest aids to a young practitioner we have ever seen.—*Peninsular Journal of Medicine.*

New and much enlarged edition—(Just Issued.)

WATSON (THOMAS), M. D., &c.,

Late Physician to the Middlesex Hospital, &c.

LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC.

Delivered at King's College, London. A new American, from the last revised and enlarged English edition, with Additions, by D. FRANCIS CONNIE, M. D., author of "A Practical Treatise on the Diseases of Children," &c. With one hundred and eighty-five illustrations on wood. In one very large and handsome volume, imperial octavo, of over 1200 closely printed pages in small type; the whole strongly bound in leather, with raised bands. Price \$4 25.

That the high reputation of this work might be fully maintained, the author has subjected it to a thorough revision; every portion has been examined with the aid of the most recent researches in pathology, and the results of modern investigations in both theoretical and practical subjects have been carefully weighed and embodied throughout its pages. The watchful scrutiny of the editor has likewise introduced whatever possesses immediate importance to the American physician in relation to diseases incident to our climate which are little known in England, as well as those points in which experience here has led to different modes of practice; and he has also added largely to the series of illustrations, believing that in this manner valuable assistance may be conveyed to the student in elucidating the text. The work will, therefore, be found thoroughly on a level with the most advanced state of medical science on both sides of the Atlantic.

The additions which the work has received are shown by the fact that notwithstanding an enlargement in the size of the page, more than two hundred additional pages have been necessary to accommodate the two large volumes of the London edition (which sells at ten dollars), within the compass of a single volume, and in its present form it contains the matter of at least three ordinary octavos. Believing it to be a work which should lie on the table of every physician, and be in the hands of every student, the publishers have put it at a price within the reach of all, making it one of the cheapest books as yet presented to the American profession, while at the same time the beauty of its mechanical execution renders it an exceedingly attractive volume.

The fourth edition now appears, so carefully revised, as to add considerably to the value of a book already acknowledged; wherever the English language is read, to be beyond all comparison the best systematic work on the Principles and Practice of Physic in the whole range of medical literature. Every lecture contains proof of the extreme anxiety of the author to keep pace with the advancing knowledge of the day, and to bring the results of the labors, not only of physicians, but of chemists and histologists, before his readers, wherever they can be turned to useful account. One scarcely knows whether to admire most the pure, simple, forcible English—the vast amount of useful practical information condensed into the Lectures—or the manly, kind-hearted, unassuming character of the lecturer shining through his work.—*Lond. Med. Times.*

Thus these admirable volumes come before the profession in their fourth edition, abounding in those distinguished attributes of moderation, judgment, erudite cultivation, clearness, and eloquence, with which they were from the first invested, but yet richer than before in the results of more prolonged observation, and in the able appreciation of the latest advances in pathology and medicine by one of the most profound medical thinkers of the day.—*London Lancet.*

The lecturer's skill, his wisdom, his learning, are equalled by the ease of his graceful diction, his eloquence, and the far higher qualities of candor, of courtesy, of modesty, and of generous appreciation of merit in others.—*N. A. Med.-Chir. Review.*

Watson's unrivalled, perhaps unapproachable work on Practice—the copious additions made to which (the fourth edition) have given it all the novelty and much of the interest of a new book.—*Charleston Med. Journal.*

Lecturers, practitioners, and students of medicine will equally hail the reappearance of the work of Dr. Watson in the form of a new—a fourth—edition. We merely do justice to our own feelings, and, we are sure, of the whole profession, if we thank him for having, in the trouble and turmoil of a large practice, made leisure to supply the hiatus caused by the exhaustion of the publisher's stock of the third edition, which has been severely felt for the last three years. For Dr. Watson has not merely caused the lectures to be reprinted, but scattered through the whole work we find additions or alterations which prove that the author has in every way sought to bring up his teaching to the level of the most recent acquisitions in science.—*Brit. and For. Medico-Chir. Review.*

WALSHE (W. H.), M. D.,

Professor of the Principles and Practice of Medicine in University College, London, &c.

A PRACTICAL TREATISE ON DISEASES OF THE LUNGS; including the Principles of Physical Diagnosis. A new American, from the third revised and much enlarged London edition. In one vol. octavo, of 468 pages. \$2 25.

The present edition has been carefully revised and much enlarged, and may be said in the main to be rewritten. Descriptions of several diseases, previously omitted, are now introduced; the causes and mode of production of the more important affections, so far as they possess direct practical significance, are succinctly inquired into; an effort has been made to bring the description of anatomical characters to the level of the wants of the practical physician; and the diagnosis and prognosis of each complaint are more completely considered. The sections on TREATMENT and the Appendix (concerning the influence of climate on pulmonary disorders), have, especially, been largely extended.—*Author's Preface.*

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON THE DISEASES OF THE HEART AND GREAT VESSELS, including the Principles of Physical Diagnosis. Third American, from the third revised and much enlarged London edition. In one handsome octavo volume of 420 pages, extra cloth. \$2 25. (Just Ready.)

From the Author's Preface.

The present edition has been carefully revised; much new matter has been added, and the entire work in a measure remodelled. Numerous facts and discussions, more or less completely novel, will be found in the description of the principles of physical diagnosis; but the chief additions have been made in the practical portions of the book. Several affections, of which little or no account had been given in the previous editions, are now treated of in detail. Functional disorders of the heart, the frequency of which is almost rivalled by the misery they inflict, have been closely reconsidered; more especially an attempt has been made to render their essential nature clearer, and consequently their treatment more successful, by an analysis of their dynamic elements.

New and much enlarged edition—(Just Issued.)

WILSON (ERASMUS), F. R. S.

A SYSTEM OF HUMAN ANATOMY, General and Special. A new and revised American, from the last and enlarged English Edition. Edited by W. H. GOBRECHT, M. D., Professor of Anatomy in the Pennsylvania Medical College, &c. Illustrated with three hundred and ninety-seven engravings on wood. In one large and exquisitely printed octavo volume, of over 600 large pages; leather. \$3 25.

The publishers trust that the well earned reputation so long enjoyed by this work will be more than maintained by the present edition. Besides a very thorough revision by the author, it has been most carefully examined by the editor, and the efforts of both have been directed to introducing everything which increased experience in its use has suggested as desirable to render it a complete text-book for those seeking to obtain or to renew an acquaintance with Human Anatomy. The amount of additions which it has thus received may be estimated from the fact that the present edition contains over one-fourth more matter than the last, rendering a smaller type and an enlarged page requisite to keep the volume within a convenient size. The editor has exercised the utmost caution to obtain entire accuracy in the text, and has largely increased the number of illustrations, of which there are about one hundred and fifty more in this edition than in the last, thus bringing distinctly before the eye of the student everything of interest or importance.

It may be recommended to the student as no less distinguished by its accuracy and clearness of description than by its typographical elegance. The wood-cuts are exquisite.—*Brit. and For. Medical Review*.

An elegant edition of one of the most useful and accurate systems of anatomical science which has been issued from the press. The illustrations are really beautiful. In its style the work is extremely concise and intelligible. No one can possibly take up this volume without being struck with the great

beauty of its mechanical execution, and the clearness of the descriptions which it contains is equally evident. Let students, by all means examine the claims of this work on their notice, before they purchase a text-book of the vitally important science which this volume so fully and easily unfolds.—*Lancet*.

We regard it as the best system now extant for students.—*Western Lancet*.

It therefore receives our highest commendation.—*Southern Med. and Surg. Journal*.

BY THE SAME AUTHOR. (Just Issued.)

ON DISEASES OF THE SKIN. Fourth and enlarged American, from the last and improved London edition. In one large octavo volume, of 650 pages, extra cloth, \$2 75.

The writings of Wilson, upon diseases of the skin, are by far the most scientific and practical that have ever been presented to the medical world on this subject. The present edition is a great improvement on all its predecessors. To dwell upon all the great merits and high claims of the work before us, *seriatim*, would indeed be an agreeable service; it would be a mental homage which we could freely offer, but we should thus occupy an undue amount of space in this *Journal*. We will, however, look

at some of the more salient points with which it abounds, and which make it incomparably superior in excellence to all other treatises on the subject of dermatology. No mere speculative views are allowed a place in this volume, which, without a doubt, will, for a very long period, be acknowledged as the chief standard work on dermatology. The principles of an enlightened and rational therapeia are introduced on every appropriate occasion.—*Am. Jour. Med. Science*, Oct. 1857.

ALSO, NOW READY,

A SERIES OF PLATES ILLUSTRATING WILSON ON DISEASES OF THE SKIN; consisting of nineteen beautifully executed plates, of which twelve are exquisitely colored, presenting the Normal Anatomy and Pathology of the Skin, and containing accurate representations of about one hundred varieties of disease, most of them the size of nature. Price in cloth \$4 25.

In beauty of drawing and accuracy and finish of coloring these plates will be found equal to anything of the kind as yet issued in this country.

The plates by which this edition is accompanied leave nothing to be desired, so far as excellence of delineation and perfect accuracy of illustration are concerned.—*Medico-Chirurgical Review*.

Of these plates it is impossible to speak too highly. The representations of the various forms of cutaneous disease are singularly accurate, and the coloring exceeds almost anything we have met with in point of delicacy and finish.—*British and Foreign Medical Review*.

We have already expressed our high appreciation of Mr. Wilson's treatise on Diseases of the Skin. The plates are comprised in a separate volume, which we counsel all those who possess the text to purchase. It is a beautiful specimen of color printing, and the representations of the various forms of skin disease are as faithful as is possible in plates of the size.—*Boston Med. and Surg. Journal*, April 8, 1858.

BY THE SAME AUTHOR.

ON CONSTITUTIONAL AND HEREDITARY SYPHILIS, AND ON SYPHILITIC ERUPTIONS. In one small octavo volume, extra cloth, beautifully printed, with four exquisite colored plates, presenting more than thirty varieties of syphilitic eruptions. \$2 25.

BY THE SAME AUTHOR.

HEALTHY SKIN; A Popular Treatise on the Skin and Hair, their Preservation and Management. Second American, from the fourth London edition. One neat volume, royal 12mo., extra cloth, of about 300 pages, with numerous illustrations. \$1 00; paper cover, 75 cents.

BY THE SAME AUTHOR.

THE DISSECTOR'S MANUAL; or, Practical and Surgical Anatomy. Third American, from the last revised and enlarged English edition. Modified and rearranged, by WILLIAM HUNT, M. D., Demonstrator of Anatomy in the University of Pennsylvania. In one large and handsome royal 12mo. volume, leather, of 582 pages, with 154 illustrations. \$2 00.

WINSLOW (FORBES), M. D., D. C. L., &c.

ON OBSCURE DISEASES OF THE BRAIN AND DISORDERS OF THE MIND; their incipient Symptoms, Pathology, Diagnosis, Treatment, and Prophylaxis. In one handsome octavo volume, of nearly 600 pages. (Just Issued.) \$3 00.

We close this brief and necessarily very imperfect notice of Dr. Winslow's great and classical work, by expressing our conviction that it is long since so important and beautifully written a volume has issued from the British medical press.—*Dublin Med. Press*, July 25, 1860.

We honestly believe this to be the best book of the season.—*Ranking's Abstract*, July, 1860.

It called us back to our old days of novel reading, it kept us from our dinner, from our business, and from our slumbers; in short, we laid it down only when we had got to the end of the last paragraph, and even then turned back to the perusal of several passages which we had marked as requiring further study. We have failed entirely in the above notice to give an adequate acknowledgment of the profit and pleasure with which we have perused the above work. We can only say to our readers, study it

yourself; and we extend the invitation to unprofessional as well as professional men, believing that it contains matter deeply interesting not to physicians alone, but to all who appreciate the truth that: "The proper study of mankind is man."—*Nashville Medical Record*, July, 1860.

The latter portion of Dr. Winslow's work is exclusively devoted to the consideration of Cerebral Pathology. It completely exhausts the subject, in the same manner as the previous seventeen chapters relating to morbid psychical phenomena left nothing unnoticed in reference to the mental symptoms premonitory of cerebral disease. It is impossible to overrate the benefits likely to result from a general perusal of Dr. Winslow's valuable and deeply interesting work.—*London Lancet*, June 23, 1860.

It contains an immense mass of information.—*Brit. and For. Med.-Chir. Review*, Oct. 1860.

WEST (CHARLES), M. D.,

Accoucheur to and Lecturer on Midwifery at St. Bartholomew's Hospital, Physician to the Hospital for Sick Children, &c.

LECTURES ON THE DISEASES OF WOMEN. Second American, from the second London edition. In one handsome octavo volume, extra cloth, of about 500 pages; price \$2 50. (Now Ready, July, 1861.)

* * * Gentlemen who received the first portion, as issued in the "Medical News and Library," can now complete their copies by procuring Part II, being page 309 to end, with Index, Title matter, &c., 8vo., cloth, price \$1.

We must now conclude this hastily written sketch with the confident assurance to our readers that the work will well repay perusal. The conscientious, painstaking, practical physician is apparent on every page.—*N. Y. Journal of Medicine*, March, 1858.

We know of no treatise of the kind so complete and yet so compact.—*Chicago Med. Jour.* Jan. 1858.

A fairer, more honest, more earnest, and more reliable investigator of the many diseases of women and children is not to be found in any country.—*Southern Med. and Surg. Journal*, January 1858.

We gladly recommend his Lectures as in the highest degree instructive to all who are interested in obstetric practice.—*London Lancet*.

We have to say of it, briefly and decidedly, that it is the best work on the subject in any language; and that it stamps Dr. West as the *facile princeps* of British obstetric authors.—*Edinb. Med. Jour.*

As a writer, Dr. West stands, in our opinion, second only to Watson, the "Macaulay of Medicine;" he possesses that happy faculty of clothing instruc-

tion in easy garments; combining pleasure with profit, he leads his pupils, in spite of the ancient proverb, along a royal road to learning. His work is one which will not satisfy the extreme on either side, but it is one that will please the great majority who are seeking truth, and one that will convince the student that he has committed himself to a candid, safe, and valuable guide.—*N. A. Med.-Chirurg. Review*, July, 1858.

Happy in his simplicity of manner, and moderate in his expression of opinion, the author is a sound reasoner and a good practitioner, and his book is worthy of the handsome garb in which it has appeared.—*Virginia Med. Journal*.

We must take leave of Dr. West's very useful work, with our commendation of the clearness of its style, and the industry and sobriety of judgment of which it gives evidence.—*London Med Times*.

Sound judgment and good sense pervade every chapter of the book. From its perusal we have derived unmixed satisfaction.—*Dublin Quart. Journ.*

BY THE SAME AUTHOR. (Just Issued.)

LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD.

Third American, from the fourth enlarged and improved London edition. In one handsome octavo volume, extra cloth, of about six hundred and fifty pages. \$2 75.

The three former editions of the work now before us have placed the author in the foremost rank of those physicians who have devoted special attention to the diseases of early life. We attempt no analysis of this edition, but may refer the reader to some of the chapters to which the largest additions have been made—those on Diphtheria, Disorders of the Mind, and Idiocy, for instance—as a proof that the work is really a new edition; not a mere reprint. In its present shape it will be found of the greatest possible service in the every-day practice of nine-tenths of the profession.—*Med. Times and Gazette*, London, Dec. 10, 1859.

All things considered, this book of Dr. West is by far the best treatise in our language upon such modifications of morbid action and disease as are witnessed when we have to deal with infancy and childhood. It is true that it confines itself to such disorders as come within the province of the physician, and even with respect to these it is unequal as regards minuteness of consideration, and some

diseases it omits to notice altogether. But those who know anything of the present condition of pædiatrics will readily admit that it would be next to impossible to effect more, or effect it better, than the accoucheur of St. Bartholomew's has done in a single volume. The lecture (XVI.) upon Disorders of the Mind in children is an admirable specimen of the value of the later information conveyed in the Lectures of Dr. Charles West.—*London Lancet*, Oct. 22, 1859.

Since the appearance of the first edition, about eleven years ago, the experience of the author has doubled; so that, whereas the lectures at first were founded on six hundred observations, and one hundred and eighty dissections made among nearly fourteen thousand children, they now embody the results of nine hundred observations, and two hundred and eighty-eight post-mortem examinations made among nearly thirty thousand children, who, during the past twenty years, have been under his care.—*British Med. Journal*, Oct. 1, 1859.

BY THE SAME AUTHOR.

AN ENQUIRY INTO THE PATHOLOGICAL IMPORTANCE OF ULCERATION OF THE OS UTERI. In one neat octavo volume, extra cloth. \$1 00.

WHITEHEAD ON THE CAUSES AND TREATMENT OF ABORTION AND STERILITY.

Second American Edition. In one volume, octavo extra cloth, pp. 308. \$1 75.

